

Construction of solutions for the critical polyharmonic equation with competing potentials

Wenjing Chen* and Zexi Wang†

School of Mathematics and Statistics, Southwest University, Chongqing, 400715, P.R. China

Abstract

In this paper, we consider the following critical polyharmonic equation

$$(-\Delta)^m u + V(|y'|, y'')u = Q(|y'|, y'')u^{m^*-1}, \quad u > 0, \quad y = (y', y'') \in \mathbb{R}^3 \times \mathbb{R}^{N-3},$$

where $N > 4m + 1$, $m \in \mathbb{N}^+$, $m^* = \frac{2N}{N-2m}$, $V(|y'|, y'')$ and $Q(|y'|, y'')$ are bounded nonnegative functions in $\mathbb{R}^+ \times \mathbb{R}^{N-3}$. By using the reduction argument and local Pohožaev identities, we prove that if $Q(r, y'')$ has a stable critical point (r_0, y''_0) with $r_0 > 0$, $Q(r_0, y''_0) > 0$, $D^\alpha Q(r_0, y''_0) = 0$ for any $|\alpha| \leq 2m - 1$ and $B_1 V(r_0, y''_0) - B_2 \sum_{|\alpha|=2m} D^\alpha Q(r_0, y''_0) \int_{\mathbb{R}^N} y^\alpha U_{0,1}^{m^*} dy > 0$, then the above problem

has a family of solutions concentrated at points lying on the top and the bottom circles of a cylinder, where B_1 and B_2 are positive constants that will be given later.

Keywords: Polyharmonic equation; Competing potentials; Reduction argument; Local Pohožaev identities.

2020 Mathematics Subject Classification: 35J60; 35B38; 35J91.

1 Introduction

In this paper, we concern the following polyharmonic equation with double potentials

$$(-\Delta)^m u + V(y)u = Q(y)u^{m^*-1}, \quad u > 0, \quad u \in H^m(\mathbb{R}^N), \quad (1.1)$$

where $N > 4m + 1$, $m \in \mathbb{N}^+$, $m^* = \frac{2N}{N-2m}$, $V(y)$ and $Q(y)$ are two potential functions, and $H^m(\mathbb{R}^N)$ is the Hilbert space with the scalar product

$$\langle u, v \rangle = \begin{cases} \int_{\mathbb{R}^N} \left((\Delta^{\frac{m}{2}} u)(\Delta^{\frac{m}{2}} v) + V(y)uv \right) dy, & \text{if } m \text{ is even,} \\ \int_{\mathbb{R}^N} \left((\nabla(\Delta^{\frac{m-1}{2}} u)) \cdot (\nabla(\Delta^{\frac{m-1}{2}} v)) + V(y)uv \right) dy, & \text{if } m \text{ is odd.} \end{cases}$$

In the case of $m = 1$, (1.1) reduces to

$$-\Delta u + V(y)u = Q(y)u^{2^*-1}, \quad u > 0, \quad u \in H^1(\mathbb{R}^N), \quad (1.2)$$

*E-mail address: wjchen@swu.edu.cn (W. Chen), zzwangmath@163.com (Z. Wang).

†Corresponding author.

where $2^* = \frac{2N}{N-2}$. If $V(y) = 0$, (1.2) can be written as

$$-\Delta u = Q(y)u^{2^*-1}, \quad u > 0, \quad u \in D^{1,2}(\mathbb{R}^N), \quad (1.3)$$

which comes from the following prescribed curvature problem by using the stereographic projection

$$-\Delta_{\mathbb{S}^N} u + \frac{N(N-2)}{2}u = \tilde{Q}(y)u^{2^*-1}, \quad u > 0 \quad \text{on } \mathbb{S}^N,$$

where $\tilde{Q}(y)$ is a potential function. Recent years, much attention has been paid to study the existence of solutions for (1.3) under suitable assumptions on $Q(y)$. Using the number of the bubbles of solutions as the parameter, Wei and Yan [30] obtained infinitely many solutions on a circle for (1.3) when $Q(y)$ is radially symmetric. Since then, the idea of [30] has been exploited and further developed by many scholars in the search for infinitely many solutions for equations with critical or asymptotic critical growth, see e.g. [7–9, 14, 17, 21, 23–25, 29, 31, 32]. In particular, Peng, Wang and Wei [25] constructed infinitely many solutions for (1.3) under a weaker symmetry condition on $Q(y)$, where the concentration point can be a saddle point of $Q(y) = Q(|\tilde{y}'|, \tilde{y}'')$, $y = (\tilde{y}', \tilde{y}'') \in \mathbb{R}^2 \times \mathbb{R}^{N-2}$. Recently, Duan, Musso and Wei [9] constructed a new type of solutions for (1.3) when $Q(y)$ is radially symmetric, and the solutions concentrate at points lying on the top and the bottom circles of a cylinder. More precisely, these solutions are different from [25, 30] and have the form

$$\sum_{j=1}^k W_{\bar{x}_j, \lambda} + \sum_{j=1}^k W_{\underline{x}_j, \lambda} + \varphi_k,$$

where $W_{x, \lambda}(y) = \left(\frac{\lambda}{1+\lambda^2|y-x|^2}\right)^{\frac{N-2}{2}}$, φ_k is a remainder term,

$$\begin{cases} \bar{x}_j = (\bar{r}\sqrt{1-\bar{h}^2} \cos \frac{2(j-1)\pi}{k}, \bar{r}\sqrt{1-\bar{h}^2} \sin \frac{2(j-1)\pi}{k}, \bar{r}\bar{h}, 0), & j = 1, 2, \dots, k, \\ \underline{x}_j = (\bar{r}\sqrt{1-\bar{h}^2} \cos \frac{2(j-1)\pi}{k}, \bar{r}\sqrt{1-\bar{h}^2} \sin \frac{2(j-1)\pi}{k}, -\bar{r}\bar{h}, 0), & j = 1, 2, \dots, k, \end{cases}$$

with \bar{h} goes to zero, and \bar{r} is close to some $r_0 > 0$.

When $Q(y) = 1$, Benci and Cerami [1] first obtained the existence result for (1.2), and proved that if $\|V\|_{\frac{N}{2}}$ is suitably small, then (1.2) has a solution. After [1], there is no other result for (1.2) until the contribution made by Chen, Wei and Yan [3]. They proved that when $N \geq 5$, $V(y)$ is radially symmetric, and $r^2V(r)$ has a local maximum point or a local minimum point, (1.2) possesses infinitely many solutions. Later, Peng, Wang and Yan [26] generalized the result of [3] from $V(y)$ is radial to partial radial. Inspired by the results of [9] and [26], under a weaker symmetry condition on $V(y)$, Du et al. [6] obtained a new type of solutions for (1.2), which are different from those obtained in [26]. For more related results about (1.2), we refer the readers to [5, 10, 16, 20, 28] and references therein.

Last but not least, we mention that, by combining the finite dimensional reduction argument and local Pohožaev identities, He, Wang and Wang [19] studied (1.2) with double potentials, and constructed infinitely many solutions, where they assumed that:

$$(C_1) \quad V(y) = V(|\tilde{y}'|, \tilde{y}'') \text{ and } Q(y) = Q(|\tilde{y}'|, \tilde{y}'') \text{ are bounded nonnegative functions, where } y = (\tilde{y}', \tilde{y}'') \in \mathbb{R}^2 \times \mathbb{R}^{N-2};$$

(C₂) $Q(\tilde{r}, \tilde{y}'')$ has a stable critical point $(\tilde{r}_0, \tilde{y}_0'')$ in the sense that $Q(\tilde{r}, \tilde{y}'')$ has a critical point $(\tilde{r}_0, \tilde{y}_0'')$ satisfying $\tilde{r}_0 > 0$, $Q(\tilde{r}_0, \tilde{y}_0'') = 1$, and

$$\deg(\nabla Q(\tilde{r}, \tilde{y}''), (\tilde{r}_0, \tilde{y}_0'')) \neq 0;$$

(C₃) $V(\tilde{r}, \tilde{y}'') \in C^1(B_\rho(\tilde{r}_0, \tilde{y}_0''))$, $Q(\tilde{r}, \tilde{y}'') \in C^3(B_\rho(\tilde{r}_0, \tilde{y}_0''))$, and

$$V(\tilde{r}_0, \tilde{y}_0'') \int_{\mathbb{R}^N} U_{0,1}^2 dy - \frac{\Delta Q(\tilde{r}_0, \tilde{y}_0'')}{2^{*N}} \int_{\mathbb{R}^N} y^2 U_{0,1}^{2^*} dy > 0,$$

where $\rho > 0$ is a small constant, and $U_{0,1}$ is the unique positive solution of $-\Delta u = u^{2^*-1}$ in \mathbb{R}^N . More recently, the fractional case was considered by Liu [22].

If $m \geq 2$, (1.1) becomes the equation involving polyharmonic operator. It is worth pointing out that the polyharmonic operator has found considerable interest in the literature due to its geometry roots, and problems involving polyharmonic operator present new and changing features compared with the elliptic operator (namely $m = 1$). To our best knowledge, few results are known for (1.1) besides [2, 11–13, 15]. In particular, Guo and Li [12] first obtained infinitely many solutions for (1.1), where $V(y) = 0$ and $Q(y)$ is radially symmetric. For $Q(y) = 1$ and $V(y)$ is partial radial, we got a new type of solutions in [2], which are concentrated at points lying on the top and the bottom circles of a cylinder.

Motivated by the results mentioned above, especially [2], [9] and [19], in this paper, under a weaker symmetry condition on $V(y)$ and $Q(y)$, namely, $V(y) \neq 0$, $Q(y) \neq 1$, and $V(y)$, $Q(y)$ are partial radial, we intend to construct a new type of solutions for problem (1.1), which concentrate at points lying on the top and the bottom circles of a cylinder. Moreover, we would like to study how the critical points of $V(y)$ and $Q(y)$ affect the existence of solutions.

Before the statement of the main results, let us first introduce some notations. We assume that $V(y)$ and $Q(y)$ satisfy the following assumptions:

(A₁) $V(y) = V(|y'|, y'')$ and $Q(y) = Q(|y'|, y'')$ are bounded nonnegative functions, where $y = (y', y'') \in \mathbb{R}^3 \times \mathbb{R}^{N-3}$;

(A₂) $Q(r, y'')$ has a stable critical point (r_0, y_0'') in the sense that $Q(r, y'')$ has a critical point (r_0, y_0'') satisfying $r_0 > 0$, $Q(r_0, y_0'') = 1$, $D^\alpha Q(r_0, y_0'') = 0$ for any $|\alpha| \leq 2m - 1$, and

$$\deg(\nabla Q(r, y''), (r_0, y_0'')) \neq 0;$$

(A₃) $V(r, y'') \in C^1(B_\rho(r_0, y_0''))$, $Q(r, y'') \in C^{2m+1}(B_\rho(r_0, y_0''))$, and

$$B_1 V(r_0, y_0'') - B_2 \sum_{|\alpha|=2m} D^\alpha Q(r_0, y_0'') \int_{\mathbb{R}^N} y^\alpha U_{0,1}^{m^*} dy > 0,$$

where $\rho > 0$ is a small constant, B_1 and B_2 are positive constants given in Lemma 3.1.

Recall that (see [27])

$$U_{x,\lambda}(y) = P_{m,N}^{\frac{N-2m}{4m}} \left(\frac{\lambda}{1 + \lambda^2 |y - x|^2} \right)^{\frac{N-2m}{2}}, \quad \lambda > 0, \quad x \in \mathbb{R}^N,$$

is the unique radial solution of the equation

$$(-\Delta)^m u = u^{m^*-1}, \quad u > 0, \quad \text{in } \mathbb{R}^N,$$

where $P_{m,N} = \prod_{h=-m}^{m-1} (N+2h)$ is a constant.

Define

$$H_s = \left\{ u : u \in H^m(\mathbb{R}^N), u(y_1, y_2, y_3, y'') = u(y_1, -y_2, -y_3, y''), \right. \\ \left. u\left(r \cos \theta, r \sin \theta, y_3, y''\right) = u\left(r \cos\left(\theta + \frac{2j\pi}{k}\right), r \sin\left(\theta + \frac{2j\pi}{k}\right), y_3, y''\right) \right\},$$

where $r = \sqrt{y_1^2 + y_2^2}$ and $\theta = \arctan \frac{y_2}{y_1}$.

Let

$$\begin{cases} x_j^+ = (\bar{r} \sqrt{1 - \bar{h}^2} \cos \frac{2(j-1)\pi}{k}, \bar{r} \sqrt{1 - \bar{h}^2} \sin \frac{2(j-1)\pi}{k}, \bar{r} \bar{h}, \bar{y}''), & j = 1, 2, \dots, k, \\ x_j^- = (\bar{r} \sqrt{1 - \bar{h}^2} \cos \frac{2(j-1)\pi}{k}, \bar{r} \sqrt{1 - \bar{h}^2} \sin \frac{2(j-1)\pi}{k}, -\bar{r} \bar{h}, \bar{y}''), & j = 1, 2, \dots, k, \end{cases}$$

where \bar{y}'' is a vector in \mathbb{R}^{N-3} , $\bar{h} \in (0, 1)$ and (\bar{r}, \bar{y}'') is close to (r_0, y_0'') .

In this paper, we consider the following three cases of \bar{h} in the process of constructing solutions:

- **Case 1.** \bar{h} goes to 1;
- **Case 2.** \bar{h} is separated from 0 and 1;
- **Case 3.** \bar{h} goes to 0.

We will use $U_{x_j^\pm, \lambda}$ as an approximate solution. To deal with the slow decay of this function when N is not big enough, we introduce the smooth cut-off function $\xi(y) = \xi(|y|, y'')$ satisfying $\xi = 1$ if $|(r, y'') - (r_0, y_0'')| \leq \delta$, $\xi = 0$ if $|(r, y'') - (r_0, y_0'')| \geq 2\delta$, and $0 \leq \xi \leq 1$, where $\delta > 0$ is a small constant such that $Q(r, y'') > 0$ if $|(r, y'') - (r_0, y_0'')| \leq 10\delta$.

Denote

$$Z_{x_j^\pm, \lambda} = \xi U_{x_j^\pm, \lambda}, \quad Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^* = \sum_{j=1}^k U_{x_j^+, \lambda} + \sum_{j=1}^k U_{x_j^-, \lambda}, \quad Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} = \sum_{j=1}^k \xi U_{x_j^+, \lambda} + \sum_{j=1}^k \xi U_{x_j^-, \lambda}.$$

As for the **Case 1**, we assume that $\nu = N - 4m - \iota$, $\iota > 0$ is a small constant, $k > 0$ is a large integer, $\lambda \in [L_0 k^{\frac{N-2m}{N-4m-\nu}}, L_1 k^{\frac{N-2m}{N-4m-\nu}}]$ for some constants $L_1 > L_0 > 0$ and $(\bar{r}, \bar{h}, \bar{y}'')$ satisfies

$$|(\bar{r}, \bar{y}'') - (r_0, y_0'')| \leq \frac{1}{\lambda^{1-\vartheta}}, \quad \sqrt{1 - \bar{h}^2} = M_1 \lambda^{-\frac{\nu}{N-2m}} + o(\lambda^{-\frac{\nu}{N-2m}}), \quad (1.4)$$

where $\vartheta > 0$ is a small constant, M_1 is a positive constant.

Theorem 1.1. *Assume that $N > 4m + 1$, if $V(y)$ and $Q(y)$ satisfy (A_1) , (A_2) and (A_3) , then there exists an integer $k_0 > 0$, such that for any $k > k_0$, problem (1.1) has a solution u_k of the form*

$$u_k = Z_{\bar{r}_k, \bar{h}_k, \bar{y}_k'', \lambda_k} + \phi_k,$$

where $\lambda_k \in [L_0 k^{\frac{N-2m}{N-4m-\nu}}, L_1 k^{\frac{N-2m}{N-4m-\nu}}]$ and $\phi_k \in H_s$. Moreover, as $k \rightarrow \infty$, $|(\bar{r}_k, \bar{y}_k'') - (r_0, y_0'')| \rightarrow 0$, $\sqrt{1 - \bar{h}_k^2} = M_1 \lambda_k^{-\frac{\nu}{N-2m}} + o(\lambda_k^{-\frac{\nu}{N-2m}})$, and $\lambda_k^{-\frac{N-2m}{2}} \|\phi_k\|_\infty \rightarrow 0$.

For the **Case 2** and **Case 3**, we assume that $k > 0$ is a large integer, $\lambda \in [L'_0 k^{\frac{N-2m}{N-4m}}, L'_1 k^{\frac{N-2m}{N-4m}}]$ for some constants $L'_1 > L'_0 > 0$ and $(\bar{r}, \bar{h}, \bar{y}'')$ satisfies

$$|(\bar{r}, \bar{y}'') - (r_0, y_0'')| \leq \frac{1}{\lambda^{1-\vartheta}}, \quad \bar{h} = a + M_2 \lambda^{-\frac{N-4m}{N-2m}} + o(\lambda^{-\frac{N-4m}{N-2m}}), \quad (1.5)$$

where $a \in [0, 1)$, $\vartheta > 0$ is a small constant, M_2 is a positive constant.

Theorem 1.2. *Assume that $N > 4m + 1$, if $V(y)$ and $Q(y)$ satisfy (A_1) , (A_2) and (A_3) , then there exists an integer $k_0 > 0$, such that for any $k > k_0$, problem (1.1) has a solution u_k of the form*

$$u_k = Z_{\bar{r}_k, \bar{h}_k, \bar{y}_k'', \lambda_k} + \phi_k.$$

where $\lambda_k \in [L_0 k^{\frac{N-2m}{N-4m}}, L_1 k^{\frac{N-2m}{N-4m}}]$ and $\phi_k \in H_s$. Moreover, as $k \rightarrow \infty$, $|(\bar{r}_k, \bar{y}_k'') - (r_0, y_0'')| \rightarrow 0$, $\bar{h}_k = a + M_2 \lambda_k^{-\frac{N-4m}{N-2m}} + o(\lambda_k^{-\frac{N-4m}{N-2m}})$, and $\lambda_k^{-\frac{N-2m}{2}} \|\phi_k\|_\infty \rightarrow 0$.

Remark 1.1. The condition $N > 4m + 1$ is used in Lemmas 2.4 and 3.1 to guarantee the existence of a small constant $\iota > 0$ for Theorem 1.1 ($\iota = 0$ in Theorem 1.2). Moreover, it is needed in the estimates of local Pohožaev identities (3.1) and (3.2), we can see this in Lemma 3.2.

Remark 1.2. The condition $D^\alpha Q(r_0, y_0'') = 0$ for any $|\alpha| \leq 2m - 1$ is used in the estimate of the error term, see Lemma 2.4 for more details.

Remark 1.3. Combining Theorems 1.1 and 1.2 with the results obtained in [2], we can see that the role of the stable critical point of $Q(r, y'')$ in constructing solutions is more important than that of $V(r, y'')$. Moreover, (A_3) implies that $V(r_0, y_0'')$ can affect the sign of $B_0 := \sum_{|\alpha|=2m} D^\alpha Q(r_0, y_0'') \int_{\mathbb{R}^N} y^\alpha U_{0,1}^{m*} dy$ and B_0 can be nonnegative.

The paper is organized as follows. In Section 2, we will carry out the reduction procedure. Then we will study the reduced problem and prove Theorem 1.1 in Section 3. Section 4 is devoted to the proof of Theorem 1.2. In Appendix A, we put some basic estimates. Throughout the paper, C denotes positive constant possibly different from line to line, $A = o(B)$ means $A/B \rightarrow 0$ and $A = O(B)$ means that $|A/B| \leq C$.

2 Reduction argument

Let

$$\|u\|_* = \sup_{y \in \mathbb{R}^N} \left(\sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N-2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N-2m}{2} + \tau}} \right) \right)^{-1} \lambda^{-\frac{N-2m}{2}} |u(y)|,$$

and

$$\|f\|_{**} = \sup_{y \in \mathbb{R}^N} \left(\sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N+2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N+2m}{2} + \tau}} \right) \right)^{-1} \lambda^{-\frac{N+2m}{2}} |f(y)|,$$

where $\tau = \frac{N-4m-\nu}{N-2m-\nu}$. For $j = 1, 2, \dots, k$, denote

$$Z_{j,2}^\pm = \frac{\partial Z_{x_j^\pm, \lambda}}{\partial \lambda}, \quad Z_{j,3}^\pm = \frac{\partial Z_{x_j^\pm, \lambda}}{\partial \bar{r}}, \quad Z_{j,l}^\pm = \frac{\partial Z_{x_j^\pm, \lambda}}{\partial \bar{y}_l''}, \quad l = 4, 5, \dots, N.$$

For later calculations, we divide \mathbb{R}^N into k parts, for $j = 1, 2, \dots, k$, define

$$\Omega_j := \left\{ y : y = (y_1, y_2, y_3, y'') \in \mathbb{R}^3 \times \mathbb{R}^{N-3}, \left\langle \frac{(y_1, y_2)}{|(y_1, y_2)|}, \left(\cos \frac{2(j-1)\pi}{k}, \sin \frac{2(j-1)\pi}{k} \right) \right\rangle_{\mathbb{R}^2} \geq \cos \frac{\pi}{k} \right\},$$

where $\langle \cdot, \cdot \rangle_{\mathbb{R}^2}$ denotes the dot product in \mathbb{R}^2 . For Ω_j , we further divide it into two separate parts

$$\Omega_j^+ := \{ y : y = (y_1, y_2, y_3, y'') \in \Omega_j, y_3 \geq 0 \},$$

$$\Omega_j^- := \{ y : y = (y_1, y_2, y_3, y'') \in \Omega_j, y_3 < 0 \}.$$

We also define the constrained space

$$\mathbb{H} := \left\{ v : v \in H_s, \int_{\mathbb{R}^N} Z_{x_j^+, \lambda}^{m^*-2} Z_{j,l}^+ v dy = 0, \int_{\mathbb{R}^N} Z_{x_j^-, \lambda}^{m^*-2} Z_{j,l}^- v dy = 0, \quad j = 1, 2, \dots, k, \quad l = 2, 3, \dots, N \right\}.$$

Consider the following linearized problem

$$\begin{cases} (-\Delta)^m \phi + V(r, y'') \phi - (m^* - 1) Q(r, y'') Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^{m^*-2} \phi \\ = f + \sum_{l=2}^N c_l \sum_{j=1}^k \left(Z_{x_j^+, \lambda}^{m^*-2} Z_{j,l}^+ + Z_{x_j^-, \lambda}^{m^*-2} Z_{j,l}^- \right), \quad \text{in } \mathbb{R}^N, \\ \phi \in \mathbb{H}, \end{cases} \quad (2.1)$$

for some real numbers c_l .

In the sequel of this section, we assume that $(\bar{r}, \bar{h}, \bar{y}'')$ satisfies (1.4). Then, we have

Lemma 2.1. *Assume that ϕ_k solves (2.1) for $f = f_k$. If $\|f_k\|_{**}$ goes to zero as k goes to infinity, so does $\|\phi_k\|_*$.*

Proof. Assume by contradiction that there exist $k \rightarrow \infty$, $\lambda_k \in [L_0 k^{\frac{N-2m}{N-4m-\nu}}, L_1 k^{\frac{N-2m}{N-4m-\nu}}]$, $(\bar{r}_k, \bar{h}_k, \bar{y}_k'')$ satisfying (1.4) and ϕ_k solving (2.1) for $f = f_k$, $\lambda = \lambda_k$, $\bar{r} = \bar{r}_k$, $\bar{h} = \bar{h}_k$, $\bar{y}'' = \bar{y}_k''$ with $\|f_k\|_{**} \rightarrow 0$ and $\|\phi_k\|_* \geq C > 0$. Without loss of generality, we assume that $\|\phi_k\|_* = 1$. For simplicity, we drop the subscript k .

From (2.1), we have

$$\begin{aligned} |\phi(y)| &\leq C \int_{\mathbb{R}^N} \frac{1}{|y-z|^{N-2m}} Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^{m^*-2}(z) |\phi(z)| dz + C \int_{\mathbb{R}^N} \frac{1}{|y-z|^{N-2m}} |f(z)| dz \\ &+ C \int_{\mathbb{R}^N} \frac{1}{|y-z|^{N-2m}} \left| \sum_{l=2}^N c_l \sum_{j=1}^k \left(Z_{x_j^+, \lambda}^{m^*-2} Z_{j,l}^+ + Z_{x_j^-, \lambda}^{m^*-2} Z_{j,l}^- \right) \right| dz. \end{aligned}$$

The rest of the proof is similar to [2, Lemma 2.1], so we omit it. \square

From Lemma 2.1, applying the same argument as the proof of [4, Proposition 4.1], we can prove the following result.

Lemma 2.2. *There exists an integer $k_0 > 0$, such that for any $k \geq k_0$ and $f \in L^\infty(\mathbb{R}^N)$, problem (2.1) has a unique solution $\phi = L_k(f)$. Moreover,*

$$\|L_k(f)\|_* \leq C \|f\|_{**}, \quad |c_l| \leq \frac{C}{\lambda \eta_l} \|f\|_{**},$$

where $\eta_2 = -\beta_1$, $\beta_1 = \frac{\nu}{N-2m}$, $\eta_l = 1$ for $l = 3, 4, \dots, N$.

Now, we consider the following problem

$$\begin{cases} (-\Delta)^m(Z_{\bar{r},\bar{h},\bar{y}'',\lambda} + \phi) + V(r, y'')(Z_{\bar{r},\bar{h},\bar{y}'',\lambda} + \phi) \\ = Q(r, y'')(Z_{\bar{r},\bar{h},\bar{y}'',\lambda} + \phi)_+^{m^*-1} + \sum_{l=2}^N c_l \sum_{j=1}^k \left(Z_{x_j^+, \lambda}^{m^*-2} Z_{j,l}^+ + Z_{x_j^-, \lambda}^{m^*-2} Z_{j,l}^- \right), \quad \text{in } \mathbb{R}^N, \\ \phi \in \mathbb{H}. \end{cases} \quad (2.2)$$

The main result of this section is as follows.

Proposition 2.1. *There exists an integer $k_0 > 0$, such that for any $k \geq k_0$, $\lambda \in [L_0 k^{\frac{N-2m}{N-4m-\nu}}, L_1 k^{\frac{N-2m}{N-4m-\nu}}]$, $(\bar{r}, \bar{h}, \bar{y}'')$ satisfies (1.4), problem (2.2) has a unique solution $\phi = \phi_{\bar{r},\bar{h},\bar{y}'',\lambda}$ satisfying*

$$\|\phi\|_* \leq C \left(\frac{1}{\lambda} \right)^{\frac{2m+1-\beta_1}{2} + \varepsilon}, \quad |c_l| \leq C \left(\frac{1}{\lambda} \right)^{\frac{2m+1-\beta_1}{2} + \eta + \varepsilon},$$

where $\varepsilon > 0$ is a small constant.

Rewrite (2.2) as

$$\begin{cases} (-\Delta)^m \phi + V(r, y'') \phi - (m^* - 1) Q(r, y'') Z_{\bar{r},\bar{h},\bar{y}'',\lambda}^{m^*-2} \phi \\ = N(\phi) + E_k + \sum_{l=2}^N c_l \sum_{j=1}^k \left(Z_{x_j^+, \lambda}^{m^*-2} Z_{j,l}^+ + Z_{x_j^-, \lambda}^{m^*-2} Z_{j,l}^- \right), \quad \text{in } \mathbb{R}^N, \\ \phi \in \mathbb{H}, \end{cases} \quad (2.3)$$

where

$$N(\phi) = Q(r, y'') \left((Z_{\bar{r},\bar{h},\bar{y}'',\lambda} + \phi)_+^{m^*-1} - Z_{\bar{r},\bar{h},\bar{y}'',\lambda}^{m^*-1} - (m^* - 1) Z_{\bar{r},\bar{h},\bar{y}'',\lambda}^{m^*-2} \phi \right),$$

and

$$\begin{aligned} E_k &= \underbrace{Q(r, y'') Z_{\bar{r},\bar{h},\bar{y}'',\lambda}^{m^*-1}}_{:=I_1} - \underbrace{\sum_{j=1}^k \left(\xi U_{x_j^+, \lambda}^{m^*-1} + \xi U_{x_j^-, \lambda}^{m^*-1} \right)}_{:=I_2} - \underbrace{V(r, y'') Z_{\bar{r},\bar{h},\bar{y}'',\lambda}}_{:=I_2} - \underbrace{\sum_{l=0}^{m-1} \binom{m}{l} (-\Delta)^{m-l} \xi (-\Delta)^l Z_{\bar{r},\bar{h},\bar{y}'',\lambda}^*}_{:=I_3} \\ &+ \underbrace{2m \sum_{l=0}^{m-1} \binom{m-1}{l} \nabla((- \Delta)^{m-l-1} \xi) \cdot \nabla((- \Delta)^l Z_{\bar{r},\bar{h},\bar{y}'',\lambda}^*)}_{:=I_4} \\ &+ \underbrace{\sum_{l=1}^{m-1} a_l \sum_{i_1, i_2, \dots, i_{m-l}=1}^N \sum_{s=0}^{l-1} \binom{l-1}{s} \nabla \left(\frac{\partial^{m-l}((- \Delta)^s Z_{\bar{r},\bar{h},\bar{y}'',\lambda}^*)}{\partial y_{i_{m-l}} \cdots \partial y_{i_1}} \right) \cdot \nabla \left(\frac{\partial^{m-l}((- \Delta)^{l-s-1} \xi)}{\partial y_{i_{m-l}} \cdots \partial y_{i_1}} \right)}_{:=I_5}, \end{aligned}$$

where $a_l, l = 1, 2, \dots, m-1$ are some constants.

In the following, we will make use of the contraction mapping theorem to prove that (2.3) is uniquely solvable under the condition that $\|\phi\|_*$ is small enough, so we need to estimate $N(\phi)$ and E_k , respectively.

Lemma 2.3. *If $N > 4m + 1$, then*

$$\|N(\phi)\|_{**} \leq C \|\phi\|_*^{\min\{2, m^*-1\}}.$$

Proof. The proof is similar as that of [2, Lemma 2.3], we omit it here. \square

Lemma 2.4. *If $N > 4m + 1$, then there exists a small constant $\varepsilon > 0$ such that*

$$\|E_k\|_{**} \leq C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}+\varepsilon}.$$

Proof. By symmetry, we assume that $y \in \Omega_1^+$. Then it follows

$$|y - x_j^-| \geq |y - x_j^+| \geq |y - x_1^+|, \quad j = 1, 2, \dots, k, \quad (2.4)$$

$$|y - x_j^+| \geq C|x_j^+ - x_1^+|, \quad |y - x_j^-| \geq C|x_j^- - x_1^+|, \quad j = 1, 2, \dots, k. \quad (2.5)$$

For I_1 , we have

$$\begin{aligned} I_1 &= Q(r, y'') \left(\sum_{j=1}^k (\xi U_{x_j^+, \lambda} + \xi U_{x_j^-, \lambda}) \right)^{m^*-1} - \sum_{j=1}^k \left(\xi U_{x_j^+, \lambda}^{m^*-1} + \xi U_{x_j^-, \lambda}^{m^*-1} \right) \\ &= Q(r, y'') \left[\left(\sum_{j=1}^k (\xi U_{x_j^+, \lambda} + \xi U_{x_j^-, \lambda}) \right)^{m^*-1} - \sum_{j=1}^k \left(\xi U_{x_j^+, \lambda}^{m^*-1} + \xi U_{x_j^-, \lambda}^{m^*-1} \right) \right] \\ &\quad + (Q(r, y'') - 1) \sum_{j=1}^k \left(\xi U_{x_j^+, \lambda}^{m^*-1} + \xi U_{x_j^-, \lambda}^{m^*-1} \right) \\ &:= I_{11} + I_{12}. \end{aligned}$$

From [2, Lemma 2.4], we obtain

$$\|I_{11}\|_{**}, \|I_2\|_{**}, \|I_3\|_{**}, \|I_4\|_{**}, \|I_5\|_{**} \leq C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}+\varepsilon}. \quad (2.6)$$

In the following, we estimate $\|I_{12}\|_{**}$. In the region $|(r, y'') - (r_0, y_0'')| \leq \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{4m}+\varepsilon}$, using the Taylor's expansion, we have

$$\begin{aligned} |I_{12}| &= \left| \left(\sum_{|\alpha|=2m} \frac{1}{(2m)!} D^\alpha Q(r_0, y_0'') (y - y_0)^\alpha + o(|y - y_0|^{2m}) \right) \sum_{j=1}^k \left(\xi U_{x_j^+, \lambda}^{m^*-1} + \xi U_{x_j^-, \lambda}^{m^*-1} \right) \right| \\ &\leq C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}+\varepsilon} \lambda^{\frac{N+2m}{2}} \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{N+2m}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{N+2m}} \right) \\ &\leq C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}+\varepsilon} \lambda^{\frac{N+2m}{2}} \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N+2m}{2}+\tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N+2m}{2}+\tau}} \right). \end{aligned}$$

On the other hand, in the region $\left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{4m}+\varepsilon} \leq |(r, y'') - (r_0, y_0'')| \leq 2\delta$,

$$|y - x_j^\pm| \geq |(r, y'') - (r_0, y_0'')| - |(r_0, y_0'') - (\bar{r}, \bar{y}'')| \geq \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{4m}+\varepsilon} - \frac{1}{\lambda^{1-\vartheta}} \geq \frac{1}{2} \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{4m}+\varepsilon},$$

which leads to

$$\frac{1}{1 + \lambda|y - x_j^\pm|} \leq C \left(\frac{1}{\lambda}\right)^{\frac{2m-1+\beta_1}{4m}-\varepsilon},$$

then

$$\begin{aligned}
|I_{12}| &\leq C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}+\varepsilon} \lambda^{\frac{N+2m}{2}} \sum_{j=1}^k \left(\frac{1}{(1+\lambda|y-x_j^+|)^{\frac{N+2m}{2}+\tau}} \frac{\lambda^{\frac{2m+1-\beta_1}{2}+\varepsilon}}{(1+\lambda|y-x_j^+|)^{\frac{N+2m}{2}-\tau}} \right. \\
&\quad \left. + \frac{1}{(1+\lambda|y-x_j^-|)^{\frac{N+2m}{2}+\tau}} \frac{\lambda^{\frac{2m+1-\beta_1}{2}+\varepsilon}}{(1+\lambda|y-x_j^-|)^{\frac{N+2m}{2}-\tau}} \right) \\
&\leq C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}+\varepsilon} \lambda^{\frac{N+2m}{2}} \lambda^{\frac{2m+1-\beta_1}{2}+\varepsilon} \left(\frac{1}{\lambda}\right)^{(N+2m-\tau)(\frac{2m-1+\beta_1}{4m}-\varepsilon)} \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1+\lambda|y-x_j^+|)^{\frac{N+2m}{2}+\tau}} + \frac{1}{(1+\lambda|y-x_j^-|)^{\frac{N+2m}{2}+\tau}} \right) \\
&\leq C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}+\varepsilon} \lambda^{\frac{N+2m}{2}} \sum_{j=1}^k \left(\frac{1}{(1+\lambda|y-x_j^+|)^{\frac{N+2m}{2}+\tau}} + \frac{1}{(1+\lambda|y-x_j^-|)^{\frac{N+2m}{2}+\tau}} \right),
\end{aligned}$$

where we used the fact that $(\frac{N+2m}{2}-\tau)(\frac{2m-1+\beta_1}{4m}-\varepsilon) \geq \frac{2m+1-\beta_1}{2}+\varepsilon$ if $\varepsilon > 0$ small enough since $N > 4m+1$ and ι is small. Therefore, we obtain

$$\|I_{12}\|_{**} \leq C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}+\varepsilon}. \quad (2.7)$$

Combining (2.6) and (2.7), we obtain the result. \square

Now we come to prove Proposition 2.1.

Proof of Proposition 2.1. Denote

$$\mathbb{E} = \left\{ \phi : \phi \in C(\mathbb{R}^N) \cap \mathbb{H}, \quad \|\phi\|_* \leq C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}} \right\}.$$

By Lemma 2.2, (2.3) is equivalent to find a fixed point for

$$\phi = T(\phi) := L_k(N(\phi) + E_k). \quad (2.8)$$

Hence, it is sufficient to prove that T is a contraction map from \mathbb{E} to \mathbb{E} . In fact, for any $\phi \in \mathbb{E}$, by Lemmas 2.2, 2.3 and 2.4, we have

$$\begin{aligned}
\|T(\phi)\|_* &\leq C \|L_k(N(\phi))\|_* + \|L_k(E_k)\|_* \leq C \|N(\phi)\|_{**} + C \|E_k\|_{**} \\
&\leq C \|\phi\|_*^{\min\{2, m^*-1\}} + C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}+\varepsilon} \leq C \left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2}}.
\end{aligned}$$

This shows that T maps from \mathbb{E} to \mathbb{E} .

On the other hand, for any $\phi_1, \phi_2 \in \mathbb{E}$, we have

$$\|T(\phi_1) - T(\phi_2)\|_* \leq C \|L_k(N(\phi_1)) - L_k(N(\phi_2))\|_* \leq C \|N(\phi_1) - N(\phi_2)\|_{**}.$$

By symmetry, we assume that $y \in \Omega_1^+$. If $N \geq 6m$, by (2.5), the Hölder inequality and Lemmas A.1, A.3, we have

$$|N(\phi_1) - N(\phi_2)| \leq C |N'(\phi_1 + \kappa(\phi_2 - \phi_1))| |\phi_1 - \phi_2| \leq C (|\phi_1|^{m^*-2} + |\phi_2|^{m^*-2}) |\phi_1 - \phi_2|$$

$$\begin{aligned}
&\leq C(\|\phi_1\|_*^{m^*-2} + \|\phi_2\|_*^{m^*-2})\|\phi_1 - \phi_2\|_*\lambda^{\frac{N+2m}{2}} \\
&\quad \times \left(\sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N-2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N-2m}{2} + \tau}} \right) \right)^{m^*-1} \\
&\leq C(\|\phi_1\|_*^{m^*-2} + \|\phi_2\|_*^{m^*-2})\|\phi_1 - \phi_2\|_*\lambda^{\frac{N+2m}{2}} \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N+2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N+2m}{2} + \tau}} \right) \\
&\quad \times \left(\sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^\tau} + \frac{1}{(1 + \lambda|y - x_j^-|)^\tau} \right) \right)^{m^*-2} \\
&\leq C(\|\phi_1\|_*^{m^*-2} + \|\phi_2\|_*^{m^*-2})\|\phi_1 - \phi_2\|_*\lambda^{\frac{N+2m}{2}} \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N+2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N+2m}{2} + \tau}} \right) \\
&\quad \times \left(1 + \sum_{j=2}^k \frac{1}{(\lambda|x_j^+ - x_1^+|)^\tau} + \sum_{j=1}^k \frac{1}{(\lambda|x_j^- - x_1^+|)^\tau} \right)^{m^*-2} \\
&\leq C(\|\phi_1\|_*^{m^*-2} + \|\phi_2\|_*^{m^*-2})\|\phi_1 - \phi_2\|_*\lambda^{\frac{N+2m}{2}} \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N+2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N+2m}{2} + \tau}} \right),
\end{aligned}$$

that is

$$\|T(\phi_1) - T(\phi_2)\|_* \leq C(\|\phi_1\|_*^{m^*-2} + \|\phi_2\|_*^{m^*-2})\|\phi_1 - \phi_2\|_* < \frac{1}{2}\|\phi_1 - \phi_2\|_*.$$

Therefore, T is a contraction map from \mathbb{E} to \mathbb{E} . The case $4m + 1 < N \leq 6m - 1$ can be discussed in a similar way.

By the contraction mapping theorem, there exists a unique $\phi = \phi_{\bar{r}, \bar{h}, \bar{y}'', \lambda}$ such that (2.8) holds. Moreover, by Lemmas 2.2, 2.3 and 2.4, we deduce

$$\|\phi\|_* \leq C\|L_k(N(\phi))\|_* + \|L_k(E_k)\|_* \leq C\|N(\phi)\|_{**} + C\|E_k\|_{**} \leq C\left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2} + \varepsilon},$$

and

$$|c_l| \leq \frac{C}{\lambda^{\eta_l}} (\|N(\phi)\|_{**} + \|E_k\|_{**}) \leq C\left(\frac{1}{\lambda}\right)^{\frac{2m+1-\beta_1}{2} + \eta_l + \varepsilon},$$

for $l = 2, 3, \dots, N$. This completes the proof. \square

3 Proof of Theorem 1.1

Recall that the functional corresponding to (1.1) is

$$I(u) = \begin{cases} \frac{1}{2} \int_{\mathbb{R}^N} \left(|\Delta^{\frac{m}{2}} u|^2 + V(r, y'') u^2 \right) dy - \frac{1}{m^*} \int_{\mathbb{R}^N} Q(r, y'') (u)_+^{m^*} dy, & \text{if } m \text{ is even,} \\ \frac{1}{2} \int_{\mathbb{R}^N} \left(|\nabla(\Delta^{\frac{m-1}{2}} u)|^2 + V(r, y'') u^2 \right) dy - \frac{1}{m^*} \int_{\mathbb{R}^N} Q(r, y'') (u)_+^{m^*} dy, & \text{if } m \text{ is odd.} \end{cases}$$

Let $\phi = \phi_{\bar{r}, \bar{h}, \bar{y}'', \lambda}$ be the function obtained in Proposition 2.1 and $u_k = Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} + \phi$. In this section, we will choose suitable $(\bar{r}, \bar{h}, \bar{y}'', \lambda)$ such that u_k is a solution of problem (1.1). For this purpose, similar to the arguments of [2, Proposition 3.1], we have the following result.

Proposition 3.1. *Assume that $(\bar{r}, \bar{h}, \bar{y}'', \lambda)$ satisfies*

$$\int_{D_\varrho} ((-\Delta)^m u_k + V(r, y'') u_k - Q(r, y'') (u_k)_+^{m^*-1}) \langle y, \nabla u_k \rangle dy = 0, \quad (3.1)$$

$$\int_{D_\varrho} ((-\Delta)^m u_k + V(r, y'') u_k - Q(r, y'') (u_k)_+^{m^*-1}) \frac{\partial u_k}{\partial y_i} dy = 0, \quad i = 4, 5, \dots, N, \quad (3.2)$$

and

$$\int_{\mathbb{R}^N} ((-\Delta)^m u_k + V(r, y'') u_k - Q(r, y'') (u_k)_+^{m^*-1}) \frac{\partial Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}}{\partial \lambda} dy = 0, \quad (3.3)$$

where $u_k = Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} + \phi$ and $D_\varrho = \{y : y = (y', y'') : (|y'|, y'') - (r_0, y''_0) \leq \varrho\}$ with $\varrho \in (2\delta, 5\delta)$, then $c_l = 0$ for $l = 2, 3, \dots, N$.

Lemma 3.1. *We have*

$$\begin{aligned} & \int_{\mathbb{R}^N} ((-\Delta)^m u_k + V(r, y'') u_k - Q(r, y'') (u_k)_+^{m^*-1}) \frac{\partial Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}}{\partial \lambda} dy \\ &= 2k \left(-\frac{\tilde{B}_1}{\lambda^{2m+1}} + \sum_{j=2}^k \frac{\tilde{B}_2}{\lambda^{N-2m+1} |x_j^+ - x_1^+|^{N-2m}} + \sum_{j=1}^k \frac{\tilde{B}_2}{\lambda^{N-2m+1} |x_j^- - x_1^+|^{N-2m}} + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right) \right) \\ &= 2k \left(-\frac{\tilde{B}_1}{\lambda^{2m+1}} + \frac{\tilde{B}_3 k^{N-2m}}{\lambda^{N-2m+1} (\sqrt{1-\bar{h}^2})^{N-2m}} + \frac{\tilde{B}_4 k}{\lambda^{N-2m+1} \bar{h}^{N-2m-1} \sqrt{1-\bar{h}^2}} + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right) \right), \end{aligned}$$

where $\tilde{B}_1, \tilde{B}_2, \tilde{B}_3, \tilde{B}_4$ are some positive constants.

Proof. By symmetry, we have

$$\begin{aligned} & \int_{\mathbb{R}^N} ((-\Delta)^m u_k + V(r, y'') u_k - Q(r, y'') (u_k)_+^{m^*-1}) \frac{\partial Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}}{\partial \lambda} dy \\ &= \left\langle I'(Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}), \frac{\partial Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}}{\partial \lambda} \right\rangle + 2k \left\langle (-\Delta)^m \phi + V(r, y'') \phi - (m^* - 1) Q(r, y'') Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^{m^*-2} \phi, \frac{\partial Z_{x_1^+, \lambda}}{\partial \lambda} \right\rangle \\ & \quad - \int_{\mathbb{R}^N} Q(r, y'') \left((Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} + \phi)_+^{m^*-1} - Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^{m^*-1} - (m^* - 1) Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^{m^*-2} \phi \right) \frac{\partial Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}}{\partial \lambda} dy \\ & := \left\langle I'(Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}), \frac{\partial Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}}{\partial \lambda} \right\rangle + 2k I_1 - I_2. \end{aligned}$$

Moreover,

$$\begin{aligned} \frac{\partial I(Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda})}{\partial \lambda} &= \frac{\partial I(Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^*)}{\partial \lambda} + O\left(\frac{k}{\lambda^{2m+1+\varepsilon}}\right) \\ &= \int_{\mathbb{R}^N} V(r, y'') Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^* \frac{\partial Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^*}{\partial \lambda} dy + \int_{\mathbb{R}^N} (1 - Q(r, y'')) (Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^*)^{m^*-1} \frac{\partial Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^*}{\partial \lambda} dy \\ & \quad - \int_{\mathbb{R}^N} \left((Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^*)^{m^*-1} - \sum_{j=1}^k U_{x_j^+, \lambda}^{m^*-1} - \sum_{j=1}^k U_{x_j^-, \lambda}^{m^*-1} \right) \frac{\partial Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^*}{\partial \lambda} dy + O\left(\frac{k}{\lambda^{2m+1+\varepsilon}}\right) \end{aligned}$$

$$:= I_3 + I_4 - I_5 + O\left(\frac{k}{\lambda^{2m+1+\varepsilon}}\right).$$

From [2, Lemmas 3.1, B.1], we have

$$|I_1| = O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right), \quad |I_2| = O\left(\frac{k}{\lambda^{2m+1+\varepsilon}}\right), \quad I_3 = 2k\left(-\frac{B_1 V(r_0, y_0'')}{\lambda^{2m+1}} + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right)\right),$$

and

$$I_5 = 2k\left(-\sum_{j=2}^k \frac{B_3}{\lambda^{N-2m+1}|x_j^+ - x_1^+|^{N-2m}} - \sum_{j=1}^k \frac{B_3}{\lambda^{N-2m+1}|x_j^- - x_1^+|^{N-2m}} + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right)\right),$$

for some positive constants B_1 and B_3 .

In the following, we estimate I_4 . By symmetry, using Lemmas A.1, A.3, and the Taylor's expansion, we have

$$\begin{aligned} I_4 &= 2k \left[\int_{\mathbb{R}^N} (1 - Q(r, y'')) U_{x_1^+, \lambda}^{m^*-1} \frac{\partial U_{x_1^+, \lambda}}{\partial \lambda} dy + O\left(\frac{1}{\lambda^{\beta_1}} \int_{\mathbb{R}^N} U_{x_1^+, \lambda}^{m^*-1} \left(\sum_{j=2}^k U_{x_j^+, \lambda} + \sum_{j=1}^k U_{x_j^-, \lambda}\right) dy\right) \right] \\ &= 2k \left[\int_{B_{\lambda^{-\frac{1}{2}+\varepsilon}}(x_1^+)} (1 - Q(r, y'')) U_{x_1^+, \lambda}^{m^*-1} \frac{\partial U_{x_1^+, \lambda}}{\partial \lambda} dy + \int_{B_{\lambda^{-\frac{1}{2}+\varepsilon}}(x_1^+)^c} (1 - Q(r, y'')) U_{x_1^+, \lambda}^{m^*-1} \frac{\partial U_{x_1^+, \lambda}}{\partial \lambda} dy \right. \\ &\quad \left. + O\left(\frac{1}{\lambda^{\beta_1}} \int_{\mathbb{R}^N} U_{x_1^+, \lambda}^{m^*-1} \left(\sum_{j=2}^k U_{x_j^+, \lambda} + \sum_{j=1}^k U_{x_j^-, \lambda}\right) dy\right) \right] \\ &= 2k \left[\int_{B_{\lambda^{-\frac{1}{2}+\varepsilon}}(x_1^+)} (1 - Q(r, y'')) U_{x_1^+, \lambda}^{m^*-1} \frac{\partial U_{x_1^+, \lambda}}{\partial \lambda} dy + O\left(\frac{1}{\lambda^{\frac{N}{2}+N\varepsilon+\beta_1}}\right) \right. \\ &\quad \left. + O\left(\frac{1}{\lambda^{\beta_1}} \left(\sum_{j=2}^k \frac{1}{(\lambda|x_j^+ - x_1^+|)^{N-\varepsilon}} + \sum_{j=1}^k \frac{1}{(\lambda|x_j^- - x_1^+|)^{N-\varepsilon}}\right)\right) \right] \\ &= 2k \left[\int_{B_{\lambda^{-\frac{1}{2}+\varepsilon}}(x_1^+)} (1 - Q(r, y'')) U_{x_1^+, \lambda}^{m^*-1} \frac{\partial U_{x_1^+, \lambda}}{\partial \lambda} dy + O\left(\frac{1}{\lambda^{\frac{N}{2}+N\varepsilon+\beta_1}}\right) + O\left(\frac{1}{\lambda^{\beta_1 + \frac{2m}{N-2m}(N-\varepsilon)}}\right) \right] \\ &= 2k \left[\int_{B_{\lambda^{-\frac{1}{2}+\varepsilon}}(x_1^+)} (1 - Q(r, y'')) U_{x_1^+, \lambda}^{m^*-1} \frac{\partial U_{x_1^+, \lambda}}{\partial \lambda} dy + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right) \right] \\ &= 2k \left[-\sum_{|\alpha|=2m} \frac{1}{(2m)!} D^\alpha Q(r_0, y_0'') \int_{B_{\lambda^{-\frac{1}{2}+\varepsilon}}(x_1^+)} (y - y_0)^\alpha U_{x_1^+, \lambda}^{m^*-1} \frac{\partial U_{x_1^+, \lambda}}{\partial \lambda} dy + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right) \right] \\ &= 2k \left[-\sum_{|\alpha|=2m} \frac{1}{(2m)!} D^\alpha Q(r_0, y_0'') \int_{B_{\lambda^{-\frac{1}{2}+\varepsilon}}(0)} (y + x_1^+ - y_0)^\alpha U_{0, \lambda}^{m^*-1} \frac{\partial U_{0, \lambda}}{\partial \lambda} dy + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right) \right] \\ &= 2k \left[-\sum_{|\alpha|=2m} \frac{1}{(2m)!} D^\alpha Q(r_0, y_0'') \int_{B_{\lambda^{-\frac{1}{2}+\varepsilon}}(0)} (y + x_1^+ - y_0)^\alpha \frac{1}{m^*} \frac{\partial U_{0, \lambda}^{m^*}}{\partial \lambda} dy + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right) \right] \\ &= 2k \left[-\sum_{|\alpha|=2m} \frac{1}{(2m)!} \frac{1}{m^*} D^\alpha Q(r_0, y_0'') \frac{\partial}{\partial \lambda} \int_{B_{\lambda^{-\frac{1}{2}+\varepsilon}}(0)} \left(\frac{y}{\lambda} + x_1^+ - y_0\right)^\alpha U_{0, 1}^{m^*} dy + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right) \right] \end{aligned}$$

$$\begin{aligned}
&= 2k \left[\sum_{|\alpha|=2m} \frac{1}{(2m-1)!} \frac{1}{m^*} D^\alpha Q(r_0, y_0'') \frac{1}{\lambda^{2m+1}} \int_{B_{\lambda^{\frac{1}{2}+\varepsilon}(0)}} y^\alpha U_{0,1}^{m^*} dy + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right) \right] \\
&= 2k \left[\sum_{|\alpha|=2m} \frac{1}{(2m-1)!} \frac{1}{m^*} D^\alpha Q(r_0, y_0'') \frac{1}{\lambda^{2m+1}} \int_{\mathbb{R}^N} y^\alpha U_{0,1}^{m^*} dy + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right) \right] \\
&= 2k \left[\frac{B_2}{\lambda^{2m+1}} \sum_{|\alpha|=2m} D^\alpha Q(r_0, y_0'') \int_{\mathbb{R}^N} y^\alpha U_{0,1}^{m^*} dy + O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right) \right],
\end{aligned}$$

where $B_2 = \frac{1}{(2m-1)!} \frac{1}{m^*}$, and we used the facts that $\frac{N}{2} + N\varepsilon + \beta_1 \geq 2m + 1 + \varepsilon$ and $\beta_1 + \frac{2m}{N-2m}(N-\varepsilon) \geq 2m + 1 + \varepsilon$ if $\varepsilon > 0$ small enough since $N > 4m + 1$ and ι is small. Using Lemma A.4 and the condition (A₃), we derive the conclusion with

$$\tilde{B}_1 = B_1 V(r_0, y_0'') - B_2 \sum_{|\alpha|=2m} D^\alpha Q(r_0, y_0'') \int_{\mathbb{R}^N} y^\alpha U_{0,1}^{m^*} dy > 0, \quad \tilde{B}_2 = B_3.$$

□

By Lemma A.5, we have

$$\int_{D_e} (-\Delta)^m u_k \langle y, \nabla u_k \rangle dy = \frac{1}{2} \int_{\partial D_e} f_m(u_k, u_k) d\sigma - \frac{N-2m}{2} \int_{D_e} u_k (-\Delta)^m u_k dy. \quad (3.4)$$

Integrating by parts, we obtain

$$\int_{D_e} V(y) u_k \langle y, \nabla u_k \rangle dy = \frac{1}{2} \int_{\partial D_e} \varrho V(y) u_k^2 d\sigma - \frac{1}{2} \int_{D_e} \langle y, \nabla V(y) \rangle u_k^2 dy - \frac{N}{2} \int_{D_e} V(y) u_k^2 dy, \quad (3.5)$$

and

$$\begin{aligned}
\int_{D_e} Q(y) (u_k)_+^{m^*-1} \langle y, \nabla u_k \rangle dy &= \frac{1}{m^*} \int_{\partial D_e} \varrho Q(y) (u_k)_+^{m^*} d\sigma \\
&\quad - \frac{1}{m^*} \int_{D_e} \langle y, \nabla Q(y) \rangle (u_k)_+^{m^*} dy - \frac{N}{m^*} \int_{D_e} Q(y) (u_k)_+^{m^*} dy.
\end{aligned} \quad (3.6)$$

Combining (3.4), (3.5) and (3.6), we know that (3.1) is equivalent to

$$\begin{aligned}
&\frac{2m-N}{2} \int_{D_e} u_k (-\Delta)^m u_k dy - \frac{1}{2} \int_{D_e} \langle y, \nabla V(y) \rangle u_k^2 dy - \frac{N}{2} \int_{D_e} V(y) u_k^2 dy \\
&\quad + \frac{1}{m^*} \int_{D_e} \langle y, \nabla Q(y) \rangle (u_k)_+^{m^*} dy + \frac{N}{m^*} \int_{D_e} Q(y) (u_k)_+^{m^*} dy \\
&= O\left(\int_{\partial D_e} \left(\phi^2 + |\phi|^{m^*} + \sum_{j=1}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j} \phi| + \sum_{j=0}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j-1} \phi| \right) d\sigma \right),
\end{aligned} \quad (3.7)$$

since $u_k = \phi$ on ∂D_e .

Similarly, for $i = 4, 5, \dots, N$, Lemma A.6 shows

$$\int_{D_e} (-\Delta)^m u_k \frac{\partial u_k}{\partial y_i} dy = \frac{1}{2} \int_{\partial D_e} g_{m,i}(u_k, u_k) d\sigma. \quad (3.8)$$

Integrating by parts, we have

$$\int_{D_e} V(r, y'') u_k \frac{\partial u_k}{\partial y_i} dy = \frac{1}{2} \int_{\partial D_e} V(r, y'') u_k^2 \nu_i d\sigma - \frac{1}{2} \int_{D_e} \frac{\partial V(r, y'')}{\partial y_i} u_k^2 dy, \quad (3.9)$$

and

$$\int_{D_e} Q(r, y'') (u_k)_+^{m^*-1} \frac{\partial u_k}{\partial y_i} dy = \frac{1}{m^*} \int_{\partial D_e} Q(r, y'') (u_k)_+^{m^*} \nu_i d\sigma - \frac{1}{m^*} \int_{D_e} \frac{\partial Q(r, y'')}{\partial y_i} (u_k)_+^{m^*} dy, \quad (3.10)$$

where $\nu = (\nu_1, \nu_2, \dots, \nu_N)$ denotes the outward unit normal vector of ∂D_e . Combining (3.8), (3.9) and (3.10), we know that (3.2) is equivalent to

$$\begin{aligned} & \int_{D_e} \frac{\partial V(r, y'')}{\partial y_i} u_k^2 dy - \frac{2}{m^*} \int_{D_e} \frac{\partial Q(r, y'')}{\partial y_i} (u_k)_+^{m^*} dy \\ &= O\left(\int_{\partial D_e} \left(\phi^2 + |\phi|^{m^*} + \sum_{j=1}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j} \phi| \right) d\sigma \right), \quad i = 4, 5, \dots, N. \end{aligned} \quad (3.11)$$

Multiplying (2.2) by u_k and integrating in D_e , we obtain

$$\begin{aligned} & \int_{D_e} u_k (-\Delta)^m u_k dy + \int_{D_e} V(y) u_k^2 dy \\ &= \int_{D_e} Q(y) (u_k)_+^{m^*} dy + \sum_{l=2}^N c_l \sum_{j=1}^k \int_{D_e} \left(Z_{x_j^+, \lambda}^{m^*-2} Z_{j,l}^+ + Z_{x_j^-, \lambda}^{m^*-2} Z_{j,l}^- \right) Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} dy. \end{aligned}$$

Thus, (3.7) can be reduced to

$$\begin{aligned} & m \int_{D_e} V(y) u_k^2 dy + \frac{1}{2} \int_{D_e} \langle y, \nabla V(y) \rangle u_k^2 dy - \frac{1}{m^*} \int_{D_e} \langle y, \nabla Q(y) \rangle (u_k)_+^{m^*} dy \\ &= \frac{2m-N}{2} \sum_{l=2}^N c_l \sum_{j=1}^k \int_{D_e} \left(Z_{x_j^+, \lambda}^{m^*-2} Z_{j,l}^+ + Z_{x_j^-, \lambda}^{m^*-2} Z_{j,l}^- \right) Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} dy \\ &+ O\left(\int_{\partial D_e} \left(\phi^2 + |\phi|^{m^*} + \sum_{j=1}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j} \phi| + \sum_{j=0}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j-1} \phi| \right) d\sigma \right). \end{aligned} \quad (3.12)$$

Using (3.11), we can rewrite (3.12) as

$$\begin{aligned} & m \int_{D_e} V(y) u_k^2 dy + \frac{1}{2} \int_{D_e} r \frac{\partial V(r, y'')}{\partial r} u_k^2 dy - \frac{1}{m^*} \int_{D_e} r \frac{\partial Q(r, y'')}{\partial r} (u_k)_+^{m^*} dy \\ &= \frac{2m-N}{2} \sum_{l=2}^N c_l \sum_{j=1}^k \int_{D_e} \left(Z_{x_j^+, \lambda}^{m^*-2} Z_{j,l}^+ + Z_{x_j^-, \lambda}^{m^*-2} Z_{j,l}^- \right) Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} dy \\ &+ O\left(\int_{\partial D_e} \left(\phi^2 + |\phi|^{m^*} + \sum_{j=1}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j} \phi| + \sum_{j=0}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j-1} \phi| \right) d\sigma \right). \end{aligned} \quad (3.13)$$

A direct computation gives

$$\sum_{j=1}^k \int_{D_e} \left(Z_{x_j^+, \lambda}^{m^*-2} Z_{j,l}^+ + Z_{x_j^-, \lambda}^{m^*-2} Z_{j,l}^- \right) Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} dy = O\left(\frac{k\lambda^m}{\lambda^{2m}}\right),$$

this with Proposition 2.1 yields

$$\sum_{l=2}^N c_l \sum_{j=1}^k \int_{D_e} \left(Z_{x_j^+, \lambda}^{m^*-2} Z_{j,l}^+ + Z_{x_j^-, \lambda}^{m^*-2} Z_{j,l}^- \right) Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} dy = O\left(\frac{k}{\lambda^{3m + \frac{1-\beta_1}{2} + \varepsilon}}\right) = o\left(\frac{k}{\lambda^{2m}}\right).$$

Therefore, (3.13) is equivalent to

$$\begin{aligned} & \int_{D_e} \frac{1}{2r^{2m-1}} \frac{\partial(r^{2m}V(r, y''))}{\partial r} u_k^2 dy - \frac{1}{m^*} \int_{D_e} r \frac{\partial Q(r, y'')}{\partial r} (u_k)_+^{m^*} dy \\ & = o\left(\frac{k}{\lambda^{2m}}\right) + O\left(\int_{\partial D_e} \left(\phi^2 + |\phi|^{m^*} + \sum_{j=1}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j} \phi| + \sum_{j=0}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j-1} \phi|\right) d\sigma\right). \end{aligned} \quad (3.14)$$

First, we estimate (3.11) and (3.14) from the right hand, and it is sufficient to estimate

$$\int_{D_{4\delta} \setminus D_{3\delta}} \left(\phi^2 + |\phi|^{m^*} + \sum_{j=1}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j} \phi| + \sum_{j=0}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j-1} \phi|\right) dy.$$

We first prove

Lemma 3.2. *It holds*

$$\int_{\mathbb{R}^N} \left(\phi(-\Delta)^m \phi + V(r, y'') \phi^2\right) dy = O\left(\frac{k}{\lambda^{2m+1-\beta_1+\varepsilon}}\right).$$

Proof. Multiplying (2.2) by ϕ and integrating in \mathbb{R}^N , we have

$$\begin{aligned} & \int_{\mathbb{R}^N} \left(\phi(-\Delta)^m \phi + V(r, y'') \phi^2\right) dy \\ & = \int_{\mathbb{R}^N} \left(Q(r, y'')(Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} + \phi)_+^{m^*-1} - V(r, y'') Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} - (-\Delta)^m Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}\right) \phi dy \\ & = \int_{\mathbb{R}^N} Q(r, y'') \left((Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} + \phi)_+^{m^*-1} - Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^{m^*-1}\right) \phi dy + \int_{\mathbb{R}^N} (Q(r, y'') - 1) Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^{m^*-1} \phi dy \\ & \quad - \int_{\mathbb{R}^N} V(r, y'') Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} \phi dy + \int_{\mathbb{R}^N} \left(Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^{m^*-1} - (-\Delta)^m Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}\right) \phi dy \\ & := I_1 + I_2 - I_3 + I_4. \end{aligned}$$

From [2, Lemma 3.2], we obtain

$$|I_1| = O\left(\frac{k}{\lambda^{2m+1-\beta_1+\varepsilon}}\right), \quad |I_3| = O\left(\frac{k}{\lambda^{2m+1-\beta_1+\varepsilon}}\right), \quad |I_4| = O\left(\frac{k}{\lambda^{2m+1-\beta_1+\varepsilon}}\right).$$

In the following, we estimate I_2 . By symmetry, using Lemmas A.1, A.3, and the Taylor's expansion, we have

$$\begin{aligned}
|I_2| &\leq C\|\phi\|_*\lambda^N \int_{\mathbb{R}^N} |Q(r, y'') - 1| \left(\sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{N-2m}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{N-2m}} \right) \right)^{m^*-1} \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N-2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N-2m}{2} + \tau}} \right) dy \\
&\leq C\|\phi\|_*\lambda^N \int_{\mathbb{R}^N} |Q(r, y'') - 1| \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{N+2m}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{N+2m}} \right) \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N-2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N-2m}{2} + \tau}} \right) dy \\
&\leq Ck\|\phi\|_*\lambda^N \left\{ \int_{B_{\lambda^{-\frac{1}{2} + \varepsilon}}(x_1^+)} + \int_{B_{\lambda^{-\frac{1}{2} + \varepsilon}}(x_1^+)^c} \right\} |Q(r, y'') - 1| \frac{1}{(1 + \lambda|y - x_1^+|)^{N+2m}} \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N-2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N-2m}{2} + \tau}} \right) dy \\
&\leq Ck\|\phi\|_*\lambda^N \int_{B_{\lambda^{-\frac{1}{2} + \varepsilon}}(x_1^+)} |Q(r, y'') - 1| \frac{1}{(1 + \lambda|y - x_1^+|)^{N+2m}} \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N-2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N-2m}{2} + \tau}} \right) dy + Ck \left(\frac{1}{\lambda} \right)^{(\frac{1}{2} + \varepsilon)(\frac{N}{2} + m) + \frac{2m+1-\beta_1}{2} + \varepsilon} \\
&\leq Ck\|\phi\|_*\lambda^N \int_{B_{\lambda^{-\frac{1}{2} + \varepsilon}}(x_1^+)} |Q(r, y'') - 1| \frac{1}{(1 + \lambda|y - x_1^+|)^{N+2m}} \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N-2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N-2m}{2} + \tau}} \right) dy + \frac{Ck}{\lambda^{2m+1-\beta_1+\varepsilon}} \\
&\leq Ck\|\phi\|_*\lambda^N \left| \sum_{|\alpha|=2m} \frac{1}{(2m)!} D^\alpha Q(r_0, y_0'') \right| \int_{B_{\lambda^{-\frac{1}{2} + \varepsilon}}(x_1^+)} |y - y_0|^\alpha \frac{1}{(1 + \lambda|y - x_1^+|)^{N+2m}} \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1 + \lambda|y - x_j^+|)^{\frac{N-2m}{2} + \tau}} + \frac{1}{(1 + \lambda|y - x_j^-|)^{\frac{N-2m}{2} + \tau}} \right) dy + \frac{Ck}{\lambda^{2m+1-\beta_1+\varepsilon}} \\
&\leq Ck\|\phi\|_* \left| \sum_{|\alpha|=2m} \frac{1}{(2m)!} D^\alpha Q(r_0, y_0'') \right| \int_{B_{\lambda^{\frac{1}{2} + \varepsilon}}(0)} \left| \frac{y}{\lambda} + x_1^+ - y_0 \right|^\alpha \frac{1}{(1 + |y|)^{N+2m}} \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1 + |y + \lambda(x_1^+ - x_j^+)|)^{\frac{N-2m}{2} + \tau}} + \frac{1}{(1 + |y + \lambda(x_1^+ - x_j^-)|)^{\frac{N-2m}{2} + \tau}} \right) dy + \frac{Ck}{\lambda^{2m+1-\beta_1+\varepsilon}} \\
&\leq C \frac{k\|\phi\|_*}{\lambda^{2m}} + \frac{Ck}{\lambda^{2m+1-\beta_1+\varepsilon}} = O\left(\frac{k}{\lambda^{2m+1-\beta_1+\varepsilon}} \right),
\end{aligned}$$

where we used the fact that $(\frac{1}{2} + \varepsilon)(\frac{N}{2} + m) + \frac{2m+1-\beta_1}{2} + \varepsilon \geq 2m + 1 - \beta_1 + \varepsilon$ if $\varepsilon > 0$ small enough since $N > 4m + 1$. This completes the proof. \square

With a similar argument of [2], we have the following lemmas.

Lemma 3.3. *It holds*

$$\int_{D_{4\delta} \setminus D_{3\delta}} (\phi^2 + |\phi|^{m^*}) dy = O\left(\frac{k}{\lambda^{2m+1-\beta_1+\varepsilon}}\right).$$

Lemma 3.4. *It holds*

$$\|\phi\|_{C^{2m-1}(D_{4\delta} \setminus D_{3\delta})} \leq \frac{C}{\lambda^{m+\tau+\varepsilon}}.$$

From Lemmas 3.2, 3.3 and 3.4, we know

$$\int_{D_{4\delta} \setminus D_{3\delta}} \left(\phi^2 + |\phi|^{m^*} + \sum_{j=1}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j} \phi| + \sum_{j=0}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j-1} \phi| \right) dy = O\left(\frac{k}{\lambda^{2m+1-\beta_1+\varepsilon}}\right) = o\left(\frac{k}{\lambda^{2m}}\right).$$

Thus, there exists $\varrho \in (3\delta, 4\delta)$ such that

$$\int_{\partial D_\varrho} \left(\phi^2 + |\phi|^{m^*} + \sum_{j=1}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j} \phi| + \sum_{j=0}^{2m-1} |\nabla^j \phi| |\nabla^{2m-j-1} \phi| \right) d\sigma = o\left(\frac{k}{\lambda^{2m}}\right). \quad (3.15)$$

Conversely, we need to estimate (3.11) and (3.14) from the left hand, and we have the following lemma.

Lemma 3.5. *[2, Lemma 3.5] For any function $h(r, y'') \in C^1(\mathbb{R}^{N-2}, \mathbb{R})$, there holds*

$$\int_{D_\varrho} h(r, y'') u_k^2 dy = 2k \left(\frac{1}{\lambda^{2m}} h(\bar{r}, \bar{y}'') \int_{\mathbb{R}^N} U_{0,1}^2 dy + o\left(\frac{1}{\lambda^{2m}}\right) \right).$$

Lemma 3.6. *For any function $h(r, y'') \in C^1(\mathbb{R}^{N-2}, \mathbb{R})$, there holds*

$$\int_{D_\varrho} h(r, y'') (u_k)_+^{m^*} dy = 2k \left(h(\bar{r}, \bar{y}'') \int_{\mathbb{R}^N} U_{0,1}^{m^*} dy + o\left(\frac{1}{\lambda^{1-\varepsilon}}\right) \right).$$

Proof. Since $u_k = Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} + \phi$, we have

$$\begin{aligned} \int_{D_\varrho} h(r, y'') (u_k)_+^{m^*} dy &= \int_{D_\varrho} h(r, y'') Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^{m^*} dy + O\left(\int_{D_\varrho} |\phi|^{m^*} dy\right) \\ &\quad + O\left(\int_{D_\varrho} Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda} |\phi|^{m^*-1} dy\right) + O\left(\int_{D_\varrho} Z_{\bar{r}, \bar{h}, \bar{y}'', \lambda}^{m^*-1} |\phi| dy\right) \\ &:= I_1 + I_2 + I_3 + I_4. \end{aligned}$$

For I_1 , a direct computation leads to

$$I_1 = 2k \left(h(\bar{r}, \bar{y}'') \int_{\mathbb{R}^N} U_{0,1}^{m^*} dy + o\left(\frac{1}{\lambda^{1-\varepsilon}}\right) \right).$$

For I_2 , from the proof of Lemma 3.3, we have

$$I_2 = O\left(\frac{k}{\lambda^{2m+1-\beta_1+\varepsilon}}\right) = o\left(\frac{k}{\lambda^{1-\varepsilon}}\right).$$

By symmetry, we obtain

$$\begin{aligned}
I_3 &\leq C\|\phi\|_*^{m^*-1}\lambda^N \int_{\mathbb{R}^N} \sum_{j=1}^k \left(\frac{1}{(1+\lambda|y-x_j^+|)^{N-2m}} + \frac{1}{(1+\lambda|y-x_j^-|)^{N-2m}} \right) \\
&\quad \times \left(\sum_{j=1}^k \left(\frac{1}{(1+\lambda|y-x_j^+|)^{\frac{N-2m}{2}+\tau}} + \frac{1}{(1+\lambda|y-x_j^-|)^{\frac{N-2m}{2}+\tau}} \right) \right)^{m^*-1} dy \\
&\leq C\|\phi\|_*^{m^*-1}\lambda^N \int_{\mathbb{R}^N} \sum_{j=1}^k \left(\frac{1}{(1+\lambda|y-x_j^+|)^{N-2m}} + \frac{1}{(1+\lambda|y-x_j^-|)^{N-2m}} \right) \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1+\lambda|y-x_j^+|)^{\frac{N+2m}{2}+\frac{N+2m}{N-2m}\tau}} + \frac{1}{(1+\lambda|y-x_j^-|)^{\frac{N+2m}{2}+\frac{N+2m}{N-2m}\tau}} \right) dy \\
&\leq Ck\|\phi\|_*^{m^*-1} = o\left(\frac{k}{\lambda^{1-\varepsilon}}\right),
\end{aligned}$$

and

$$\begin{aligned}
I_4 &\leq C\|\phi\|_*\lambda^N \int_{\mathbb{R}^N} \sum_{j=1}^k \left(\frac{1}{(1+\lambda|y-x_j^+|)^{N+2m}} + \frac{1}{(1+\lambda|y-x_j^-|)^{N+2m}} \right) \\
&\quad \times \sum_{j=1}^k \left(\frac{1}{(1+\lambda|y-x_j^+|)^{\frac{N-2m}{2}+\tau}} + \frac{1}{(1+\lambda|y-x_j^-|)^{\frac{N-2m}{2}+\tau}} \right) dy \\
&\leq Ck\|\phi\|_* = o\left(\frac{k}{\lambda^{1-\varepsilon}}\right).
\end{aligned}$$

So we get the result. \square

Now we will prove Theorem 1.1.

Proof of Theorem 1.1. Through the above discussion, applying (3.15) and Lemmas 3.5, 3.6 to (3.11) and (3.14), we can see that (3.1) and (3.2) are equivalent to

$$2k\left(\frac{1}{\lambda^{2m}}\frac{1}{2\bar{r}^{2m-1}}\frac{\partial(\bar{r}^{2m}V(\bar{r},\bar{y}''))}{\partial\bar{r}}\int_{\mathbb{R}^N}U_{0,1}^2dy - \frac{1}{m^*}\bar{r}\frac{\partial Q(\bar{r},\bar{y}'')}{\partial\bar{r}}\int_{\mathbb{R}^N}U_{0,1}^{m^*}dy + o\left(\frac{1}{\lambda^{1-\varepsilon}}\right)\right) = o\left(\frac{k}{\lambda^{2m}}\right),$$

and

$$2k\left(\frac{1}{\lambda^{2m}}\frac{\partial V(\bar{r},\bar{y}'')}{\partial\bar{y}_i}\int_{\mathbb{R}^N}U_{0,1}^2dy - \frac{2}{m^*}\frac{\partial Q(\bar{r},\bar{y}'')}{\partial\bar{y}_i}\int_{\mathbb{R}^N}U_{0,1}^{m^*}dy + o\left(\frac{1}{\lambda^{1-\varepsilon}}\right)\right) = o\left(\frac{k}{\lambda^{2m}}\right), \quad i = 4, 5, \dots, N.$$

Therefore, the equations to determine (\bar{r}, \bar{y}'') are

$$\frac{\partial Q(\bar{r}, \bar{y}'')}{\partial \bar{r}} = o\left(\frac{1}{\lambda^{1-\varepsilon}}\right), \quad (3.16)$$

and

$$\frac{\partial Q(\bar{r}, \bar{y}'')}{\partial \bar{y}_i} = o\left(\frac{1}{\lambda^{1-\varepsilon}}\right), \quad i = 4, 5, \dots, N. \quad (3.17)$$

Moreover, by Lemma 3.1, the equation to determine λ is

$$-\frac{\tilde{B}_1}{\lambda^{2m+1}} + \frac{\tilde{B}_3 k^{N-2m}}{\lambda^{N-2m+1}(\sqrt{1-\bar{h}^2})^{N-2m}} + \frac{\tilde{B}_4 k}{\lambda^{N-2m+1}\bar{h}^{N-2m-1}\sqrt{1-\bar{h}^2}} = O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right), \quad (3.18)$$

where $\tilde{B}_1, \tilde{B}_3, \tilde{B}_4$ are positive constants.

Let $\lambda = tk^{\frac{N-2m}{N-4m-\nu}}$ with $\nu = N - 4M - \iota$, ι is a small constant, then $t \in [L_0, L_1]$. From (3.18), we have

$$-\frac{\tilde{B}_1}{t^{2m+1}} + \frac{\tilde{B}_3 M_1^{N-2m}}{t^{N-2m+1-\nu}} = o(1), \quad t \in [L_0, L_1].$$

Define

$$F(t, \bar{r}, \bar{y}'') = \left(\nabla_{\bar{r}, \bar{y}''} Q(\bar{r}, \bar{y}''), -\frac{\tilde{B}_1}{t^{2m+1}} + \frac{\tilde{B}_3 M_1^{N-2m}}{t^{N-2m+1-\nu}} \right).$$

Then, it holds

$$\deg\left(F(t, \bar{r}, \bar{y}''), [L_0, L_1] \times B_{\lambda^{\frac{1}{1-\nu}}}\left((r_0, y_0'')\right)\right) = -\deg\left(\nabla_{\bar{r}, \bar{y}''} Q(\bar{r}, \bar{y}''), B_{\lambda^{\frac{1}{1-\nu}}}\left((r_0, y_0'')\right)\right) \neq 0.$$

Hence, (3.16), (3.17) and (3.18) has a solution $t_k \in [L_0, L_1]$, $(\bar{r}_k, \bar{y}_k'') \in B_{\lambda^{\frac{1}{1-\nu}}}\left((r_0, y_0'')\right)$. \square

4 Proof of Theorem 1.2

In this section, we give a brief proof of Theorem 1.2. We define $\tau = \frac{N-4m}{N-2m}$.

Proof of Theorem 1.2. We can verify that

$$\frac{k}{\lambda^\tau} = O(1), \quad \frac{k}{\lambda} = O\left(\left(\frac{1}{\lambda}\right)^{\frac{2m}{N-2m}}\right). \quad (4.1)$$

Using (4.1) and Lemmas A.3, A.4, we get the same conclusions for problems arising from the distance between points $\{x_j^+\}_{j=1}^k$ and $\{x_j^-\}_{j=1}^k$.

Moreover, by Lemma A.2, we have

$$|Z_{j,2}^\pm| \leq C\lambda^{-\beta_2} Z_{x_j^\pm, \lambda}, \quad |Z_{j,l}^\pm| \leq C\lambda Z_{x_j^\pm, \lambda}, \quad l = 3, 4, \dots, N, \quad (4.2)$$

where $\beta_2 = \frac{N-4m}{N-2m}$.

Using (4.1) and (4.2), with a similar step in the proof of Theorem 1.1 in Sections 2 and 3, we know that the proof of Theorem 1.2 has the same reduction structure as that of Theorem 1.1 and u_k is a solution of problem (1.1) if the following equalities hold:

$$\frac{\partial Q(\bar{r}, \bar{y}'')}{\partial \bar{r}} = o\left(\frac{1}{\lambda^{1-\varepsilon}}\right), \quad (4.3)$$

$$\frac{\partial Q(\bar{r}, \bar{y}'')}{\partial \bar{y}_i} = o\left(\frac{1}{\lambda^{1-\varepsilon}}\right), \quad i = 4, 5, \dots, N, \quad (4.4)$$

$$-\frac{\tilde{B}_1}{\lambda^{2m+1}} + \frac{\tilde{B}_3 k^{N-2m}}{\lambda^{N-2m+1}(\sqrt{1-\bar{h}^2})^{N-2m}} + \frac{\tilde{B}_4 k}{\lambda^{N-2m+1}\bar{h}^{N-2m-1}\sqrt{1-\bar{h}^2}} = O\left(\frac{1}{\lambda^{2m+1+\varepsilon}}\right). \quad (4.5)$$

Let $\lambda = tk^{\frac{N-2m}{N-4m}}$, then $t \in [L'_0, L'_1]$. Next, we discuss the main items in (4.5).

Case 1. If $\bar{h} \rightarrow A \in (0, 1)$, then $(\lambda^{\frac{N-4m}{N-2m}} \bar{h})^{-1} \rightarrow 0$ as $\lambda \rightarrow \infty$, from (4.5), we have

$$-\frac{\tilde{B}_1}{t^{2m+1}} + \frac{\tilde{B}_3}{t^{N-2m+1}(\sqrt{1-A^2})^{N-2m}} = o(1), \quad t \in [L'_0, L'_1].$$

Define

$$F(t, \bar{r}, \bar{y}'') = \left(\nabla_{\bar{r}, \bar{y}''} Q(\bar{r}, \bar{y}''), -\frac{\tilde{B}_1}{t^{2m+1}} + \frac{\tilde{B}_3}{t^{N-2m+1}(\sqrt{1-A^2})^{N-2m}} \right).$$

Then, it holds

$$\deg\left(F(t, \bar{r}, \bar{y}''), [L_0, L_1] \times B_{\lambda^{\frac{1}{1-\vartheta}}}\left((r_0, y_0'')\right)\right) = -\deg\left(\nabla_{\bar{r}, \bar{y}''} Q(\bar{r}, \bar{y}''), B_{\lambda^{\frac{1}{1-\vartheta}}}\left((r_0, y_0'')\right)\right) \neq 0.$$

Hence, (4.3), (4.4) and (4.5) has a solution $t_k \in [L'_0, L'_1]$, $(\bar{r}_k, \bar{y}_k'') \in B_{\lambda^{\frac{1}{1-\vartheta}}}\left((r_0, y_0'')\right)$.

Case 2. If $\bar{h} \rightarrow 0$ and $(\lambda^{\frac{N-4m}{N-2m}} \bar{h})^{-1} \rightarrow 0$ as $\lambda \rightarrow \infty$, from (4.5), we have

$$-\frac{\tilde{B}_1}{t^{2m+1}} + \frac{\tilde{B}_3}{t^{N-2m+1}} = o(1), \quad t \in [L'_0, L'_1].$$

Define

$$F(t, \bar{r}, \bar{y}'') = \left(\nabla_{\bar{r}, \bar{y}''} Q(\bar{r}, \bar{y}''), -\frac{\tilde{B}_1}{t^{2m+1}} + \frac{\tilde{B}_3}{t^{N-2m+1}} \right).$$

Then, we can find a solution $(t_k, \bar{r}_k, \bar{y}_k'')$ of (4.3), (4.4) and (4.5) as before.

Case 3. If $\bar{h} \rightarrow 0$ and $(\lambda^{\frac{N-4m}{N-2m}} \bar{h})^{-1} \rightarrow A \in (C_1, M_2)$ for some positive constant C_1 as $\lambda \rightarrow \infty$, from (4.5), we have

$$-\frac{\tilde{B}_1}{t^{2m+1}} + \frac{\tilde{B}_3}{t^{N-2m+1}} + \frac{\tilde{B}_4 A^{N-2m-1}}{t^{2m+1+\frac{N-4m}{N-2m}}} = o(1), \quad t \in [L'_0, L'_1].$$

Since $N-2m+1$ and $2m+1+\frac{N-4m}{N-2m}$ are strictly greater than $2m+1$, there exists a solution of (4.3), (4.4) and (4.5) as before. \square

Appendix A Some basic estimates

In this section, we give some basic estimates.

Lemma A.1. [30, Lemma B.1] For $y, x_i, x_j \in \mathbb{R}^N$, $i \neq j$, let

$$g_{ij}(y) = \frac{1}{(1+|y-x_i|)^{\kappa_1}} \frac{1}{(1+|y-x_j|)^{\kappa_2}},$$

where $\kappa_1, \kappa_2 \geq 1$ are constants. Then for any constant $0 < \sigma \leq \min\{\kappa_1, \kappa_2\}$, there exists a constant $C > 0$ such that

$$g_{ij}(y) \leq \frac{C}{|x_i - x_j|^\sigma} \left(\frac{1}{(1+|y-x_i|)^{\kappa_1+\kappa_2-\sigma}} + \frac{1}{(1+|y-x_j|)^{\kappa_1+\kappa_2-\sigma}} \right).$$

Lemma A.2. As $\lambda \rightarrow \infty$, we have

$$\frac{\partial U_{x_j^\pm, \lambda}}{\partial \lambda} = O(\lambda^{-1} U_{x_j^\pm, \lambda}) + O(\lambda U_{x_j^\pm, \lambda}) \frac{\partial \sqrt{1 - \bar{h}^2}}{\partial \lambda} + O(\lambda U_{x_j^\pm, \lambda}) \frac{\partial \bar{h}}{\partial \lambda}.$$

Hence, if $\sqrt{1 - \bar{h}^2} = C\lambda^{-\beta_1}$ with $0 < \beta_1 < 1$, we have

$$\left| \frac{\partial U_{x_j^\pm, \lambda}}{\partial \lambda} \right| \leq C \frac{U_{x_j^\pm, \lambda}}{\lambda^{\beta_1}}.$$

If $\bar{h} = C\lambda^{-\beta_2}$ with $0 < \beta_2 < 1$, then we have

$$\left| \frac{\partial U_{x_j^\pm, \lambda}}{\partial \lambda} \right| \leq C \frac{U_{x_j^\pm, \lambda}}{\lambda^{\beta_2}}.$$

Proof. The proof is standard, we omit it. □

Lemma A.3. [6, Lemma A.2] For any $\gamma > 0$, there exists a constant $C > 0$ such that

$$\sum_{j=2}^k \frac{1}{|x_j^+ - x_1^+|^\gamma} \leq \frac{Ck^\gamma}{(\bar{r}\sqrt{1 - \bar{h}^2})^\gamma} \sum_{j=2}^k \frac{1}{(j-1)^\gamma} \leq \begin{cases} \frac{Ck^\gamma}{(\bar{r}\sqrt{1 - \bar{h}^2})^\gamma}, & \gamma > 1; \\ \frac{Ck^\gamma \log k}{(\bar{r}\sqrt{1 - \bar{h}^2})^\gamma}, & \gamma = 1; \\ \frac{Ck}{(\bar{r}\sqrt{1 - \bar{h}^2})^\gamma}, & \gamma < 1, \end{cases}$$

and

$$\sum_{j=1}^k \frac{1}{|x_j^- - x_1^+|^\gamma} \leq \sum_{j=2}^k \frac{1}{|x_j^+ - x_1^+|^\gamma} + \frac{C}{(\bar{r}\bar{h})^\gamma}.$$

Lemma A.4. [2, Lemma A.6] Assume that $N > 4m + 1$, as $k \rightarrow \infty$, we have

$$\sum_{j=2}^k \frac{1}{|x_j^+ - x_1^+|^{N-2m}} = \frac{A_1 k^{N-2m}}{(\sqrt{1 - \bar{h}^2})^{N-2m}} \left(1 + o\left(\frac{1}{k}\right)\right),$$

and if $\frac{1}{hk} = o(1)$, then

$$\sum_{j=1}^k \frac{1}{|x_j^- - x_1^+|^{N-2m}} = \frac{A_2 k}{\bar{h}^{N-2m-1} (\sqrt{1 - \bar{h}^2})} \left(1 + o\left(\frac{1}{hk}\right)\right) + O\left(\frac{1}{(\sqrt{1 - \bar{h}^2})^{N-2m}}\right),$$

or else, $\frac{1}{hk} = C$, then

$$\sum_{j=1}^k \frac{1}{|x_j^- - x_1^+|^{N-2m}} = \left(\frac{A_3 k}{\bar{h}^{N-2m-1}}, \frac{A_4 k}{\bar{h}^{N-2m-1}} \right),$$

where A_1, A_2, A_3 and A_4 are some positive constants.

Suppose that u and v are two smooth functions in a given bounded domain Ω , define the following two bilinear functionals

$$L_1(u, v) = \int_{\Omega} \left((-\Delta)^m u \langle y, \nabla v \rangle + (-\Delta)^m v \langle y, \nabla u \rangle \right) dy,$$

$$L_{2,i}(u, v) = \int_{\Omega} \left((-\Delta)^m u \frac{\partial v}{\partial y_i} + (-\Delta)^m v \frac{\partial u}{\partial y_i} \right) dy, \quad i = 1, 2, \dots, N.$$

We have

Lemma A.5. [18, Proposition 3.3] For any integer $m > 0$, there exists a function $f_m(u, v)$ such that

$$L_1(u, v) = \int_{\partial\Omega} f_m(u, v) dy - \frac{N-2m}{2} \int_{\Omega} \left(v(-\Delta)^m u + u(-\Delta)^m v \right) dy.$$

Moreover, $f_m(u, v)$ has the form

$$f_m(u, v) = \sum_{j=1}^{2m-1} \bar{l}_j(y, \nabla^j u, \nabla^{2m-j} v) + \sum_{j=0}^{2m-1} \tilde{l}_j(\nabla^j u, \nabla^{2m-j-1} v),$$

where $\bar{l}_j(y, \nabla^j u, \nabla^{2m-j} v)$ and $\tilde{l}_j(\nabla^j u, \nabla^{2m-j-1} v)$ are linear in each component.

Lemma A.6. [18, Proposition 3.1] For any integer $m > 0$, there exists a function $g_{m,i}(u, v)$ such that

$$L_{2,i}(u, v) = \int_{\partial\Omega} g_{m,i}(u, v) dy.$$

Moreover, $g_{m,i}(u, v)$ has the form

$$g_{m,i}(u, v) = \sum_{j=1}^{2m-1} l_{j,i}(\nabla^j u, \nabla^{2m-j} v),$$

where $l_{j,i}(\nabla^j u, \nabla^{2m-j} v)$ is bilinear in $\nabla^j u$ and $\nabla^{2m-j} v$.

Acknowledgments

The research has been supported by Natural Science Foundation of Chongqing, China CSTB2024 NSCQ-LZX0038.

Statements and Declarations

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest. The manuscript has no associated data.

References

- [1] V. Benci, G. Cerami, Existence of positive solutions of the equation $-\Delta u + a(x)u = u^{\frac{N+2}{N-2}}$ in \mathbb{R}^N , *J. Funct. Anal.*, **88** (1990): 90-117.
- [2] W.J. Chen, Z.X. Wang, New type of solutions for the critical polyharmonic equation, *arXiv preprint*, (2024): arXiv.2405.16095.
- [3] W.Y. Chen, J.C. Wei, S.S. Yan, Infinitely many solutions for the Schrödinger equations in \mathbb{R}^N with critical growth, *J. Differential Equations*, **252** (2012): 2425-2447.
- [4] M. del Pino, P. Felmer, M. Musso, Two-bubble solutions in the super-critical Bahri-Coron's problem, *Calc. Var. Partial Differential Equations*, **16** (2003): 113-145.
- [5] F. Du, Q.Q. Hua, C.H. Wang, A new type of bubble solutions for a critical fractional Schrödinger equation, *arXiv preprint*, (2023): arXiv.2307.02272.
- [6] F. Du, Q.Q. Hua, C.H. Wang, Q.F. Wang, Multi-piece of bubble solutions for a nonlinear critical elliptic equation, *J. Differential Equations*, **393** (2024): 102-138.
- [7] L.X. Duan, Q. Guo, Infinitely many bubbling solutions and non-degeneracy results to fractional prescribed curvature problems, *arXiv preprint*, (2022): arXiv.2207.14441.
- [8] L.P. Duan, M. Musso, New type of solutions for the nonlinear Schrödinger equation in \mathbb{R}^N , *J. Differential Equations*, **336** (2022): 479-504.
- [9] L.P. Duan, M. Musso, S.T. Wei, Doubling the equatorial for the prescribed scalar curvature problem on S^N , *NoDEA Nonlinear Differential Equations Appl.*, **30** (2023): 40.
- [10] Y. Gao, Y.X. Guo, New type of solutions for Schrödinger equations with critical growth, *arXiv preprint*, (2024): arXiv.2401.11111.
- [11] Y. Gao, Y.X. Guo, Y.C. Hu, Double-tower solutions for higher-order prescribed curvature problem, *Ann. Mat. Pura Appl.*, **203** (2024): 1327-1360.
- [12] Y.X. Guo, B. Li, Infinitely many solutions for the prescribed curvature problem of polyharmonic operator, *Calc. Var. Partial Differential Equations*, **46** (2013): 809-836.
- [13] Y.X. Guo, T. Liu, The prescribed scalar curvature problem for polyharmonic operator, *Ann. Mat. Pura Appl.*, **200** (2021): 953-982.
- [14] Y.X. Guo, T. Liu, Large energy bubble solutions for Schrödinger equation with supercritical growth, *Adv. Nonlinear Stud.*, **21** (2021): 421-445.
- [15] Y.X. Guo, T. Liu, J.J. Nie, Construction of solutions for the polyharmonic equation via local Pohozaev identities, *Calc. Var. Partial Differential Equations*, **58** (2019): 123.
- [16] Y.X. Guo, T. Liu, J.J. Nie, Solutions for fractional Schrödinger equation involving critical exponent via local Pohozaev identities, *Adv. Nonlinear Stud.*, **20** (2020): 185-211.
- [17] Y.X. Guo, J.J. Nie, Infinitely many non-radial solutions for the prescribed curvature problem of fractional operator, *Discrete Contin. Dyn. Syst.*, **36** (2016): 6873-6898.

- [18] Y.X. Guo, S.J. Peng, S.S. Yan, Local uniqueness and periodicity induced by concentration, *Proc. Lond. Math. Soc.*, **114** (2017): 1005-1043.
- [19] Q.H. He, C.H. Wang, D.B. Wang, Construction of solutions for a critical problem with competing potentials via local Pohozaev identities, *Commun. Contemp. Math.*, **24** (2022): 2050071.
- [20] Q.H. He, C.H. Wang, Q.F. Wang, New type of positive bubble solutions for a critical Schrödinger equation, *J. Geom. Anal.*, **32** (2022): 278.
- [21] Y.Y. Li, J.C. Wei, H.Y. Xu, Multi-bump solutions of $-\Delta u = K(x)u^{\frac{n+2}{n-2}}$ on lattices in \mathbb{R}^n , *J. Reine Angew. Math.*, **743** (2018): 163-211.
- [22] T. Liu, Construction of infinitely many solutions for fractional Schrödinger equation with double potentials, *Z. Angew. Math. Phys.*, **75** (2024): 102.
- [23] C.G. Liu, Q. Ren, Infinitely many non-radial solutions for fractional Nirenberg problem, *Calc. Var. Partial Differential Equations*, **56** (2017): 52.
- [24] M. Musso, J.C. Wei, S.S. Yan, Infinitely many positive solutions for a nonlinear field equation with super-critical growth, *Proc. Lond. Math. Soc.*, **112** (2016): 1-26.
- [25] S.J. Peng, C.H. Wang, S.T. Wei, Constructing solutions for the prescribed scalar curvature problem via local Pohozaev identities, *J. Differential Equations*, **267** (2019): 2503-2530.
- [26] S.J. Peng, C.H. Wang, S.S. Yan, Construction of solutions via local Pohozaev identities, *J. Funct. Anal.*, **274** (2018): 2606-2633.
- [27] C.A. Swanson, The best Sobolev constant, *Appl. Anal.*, **47** (1992): 227-239.
- [28] J. Vétois, S.D. Wang, Infinitely many solutions for cubic nonlinear Schrödinger equations in dimension four, *Adv. Nonlinear Anal.*, **8** (2019): 715-724.
- [29] C.H. Wang, S.T. Wei, Large number of bubble solutions for a fractional elliptic equation with almost critical exponents, *Proc. Roy. Soc. Edinburgh Sect. A*, **151** (2021): 1642-1681.
- [30] J.C. Wei, S.S. Yan, Infinitely many solutions for the prescribed scalar curvature problem on \mathbb{S}^N , *J. Funct. Anal.*, **258** (2010): 3048-3081.
- [31] J.C. Wei, S.S. Yan, Infinitely many positive solutions for the nonlinear Schrödinger equations in \mathbb{R}^N , *Calc. Var. Partial Differential Equations*, **37** (2010): 423-439.
- [32] J.C. Wei, S.S. Yan, Infinitely many nonradial solutions for the Hénon equation with critical growth, *Rev. Mat. Iberoam.*, **29** (2013): 997-1020.