Quantum Computing Universal Thermalization Dynamics in a (2+1)D Lattice Gauge Theory

Niklas Mueller,^{1,*} Tianyi Wang,^{2,3,4} Or Katz,^{3,5,6} Zohreh Davoudi,^{7,8,4,9} and Marko Cetina^{2,3,5,4}

¹InQubator for Quantum Simulation (IQuS), Department of Physics,

University of Washington, Seattle, WA 98195, USA

²Department of Physics, Duke University, Durham, NC 27708

³Duke Quantum Center, Duke University, Durham, NC 27701

⁴ The NSF Institute for Robust Quantum Simulation,

University of Maryland, College Park, Maryland 20742, USA

⁵Department of Electrical and Computer Engineering, Duke University, Durham, NC 27708

⁶School of Applied and Engineering Physics, Cornell University, Ithaca, NY 14853

⁷Department of Physics and Maryland Center for Fundamental Physics,

- University of Maryland, College Park, MD 20742 USA
- ⁸Joint Center for Quantum Information and Computer Science,

NIST and University of Maryland, College Park, MD 20742 USA

⁹National Quantum Laboratory (QLab), University of Maryland, College Park, MD 20742 USA

Simulating non-equilibrium phenomena in strongly-interacting quantum many-body systems, including thermalization, is a promising application of near-term and future quantum computation. By performing experiments on a digital quantum computer consisting of fully-connected optically-controlled trapped ions, we study the role of entanglement in the thermalization dynamics of a Z_2 lattice gauge theory in 2+1 spacetime dimensions. Using randomized-measurement protocols, we efficiently learn a classical approximation of non-equilibrium states that yields the gap-ratio distribution and the spectral form factor of the entanglement Hamiltonian. These observables exhibit universal early-time signals for quantum chaos, a prerequisite for thermalization. Our work, therefore, establishes quantum computers as robust tools for studying universal features of thermalization in complex many-body systems, including in gauge theories.

I. INTRODUCTION

Thermalization of isolated quantum many-body systems in e.g., ultra-cold atomic gases [1-8], trapped ions [9-12], condensed matter physics [13, 14], cosmology [15], and nuclear and high-energy physics [16, 17], remains a vibrant frontier. Most quantum systems thermalize according to the Eigenstate Thermalization Hypothesis (ETH) [18, 19], which posits that observables eventually attain thermal values after long unitary evolution [20, 21]. Yet, probing large quantum many-body systems undergoing thermalization is inherently challenging due to the non-equilibrium nature of the process, which precludes using first-principle classical computational techniques. Recent advancements in quantum information theory and experiment [22-26] have brought this topic within immediate experimental reach, potentially enabling, via quantum simulation [27–29], verification of important thermalization paradigms, and discovery of novel principles grounded in quantum information science.

While it is commonly posited that quantum chaos and ergodicity are prerequisites for thermalization [30, 31], their demonstration in the context of quantum manybody systems remains somewhat elusive. Indicators of chaos and ergodicity involve measures associated with the eigenvalues of a given Hamiltonian [32–34], or the properties of its eigenstates [35]. A connection has been drawn recently between the *entanglement* of quantum states and quantum chaos, via the so-called entanglement Hamiltonians (EHs) [36, 37]. This connection can, in principle, be leveraged to experimentally probe thermalization via quantum simulation. Most studies to date, nonetheless, have stayed in the realm of theoretical exploration [38–43].

Analog quantum simulation allows one to monitor a quantum system continuously in time. However, current analog quantum simulators have limited programmability and are restricted to probing specific physical models and a limited set of observables [9-12]. In contrast, universal digital quantum computers allow, in principle, the probing of the dynamics of *any* physical model. The universal control in digital quantum computers enables the use of tomographic techniques to extract a wide range of observables, including entanglement. Digitized time evolution, via Trotterization [44, 45] or other schemes [46, 47], limits the near-term simulations to times that are short compared to those of thermalization. Nonetheless, there exist universal phenomena that are indicative of quantum chaos and ergodicity at earlier times. Probing these phenomena, therefore, is an attractive near-term opportunity for digital quantum computers.

Gauge theories and their lattice formulations are among prime physical models whose simulations will benefit from quantum-computing technology [29, 48– 51]. Gauge theories are key in high-energy and nuclear physics [52, 53], condensed and synthetic quantum mat-

^{*} niklasmu@uw.edu




FIG. 1. Overview of the experiment and the entanglement-Hamiltonian tomography. (a) Schematic of the trapped-ion experiment: 15 optically-controlled ions in a linear trap realize a universal digital quantum computer. Single-qubit and all-to-all two-qubit gates are implemented by an array of individually-focused laser beams (blue, vertical) and a global laser beam (blue, horizontal). Purple beams indicate the operation of a two-qubit gate that entangles the corresponding ions' internal states. (b) Schematic of the randomized-measurement protocol for entanglement-Hamiltonian tomography. This protocol extracts a classical approximation of a reduced density matrix associated with a subsystem of the quantum state $|\psi(t)\rangle$, from which the presence or absence of quantum chaos is inferred. The protocol consists of measuring observables in a single-qubit randomized basis, then classically learning the entanglement Hamiltonian $H_{\rm E}(\{\beta_i\})$, which parameterizes the reduced quantum state with parameters $\{\beta_i\}$, so as to optimally reproduce all measurements. The statistical behavior of the eigenvalue spectrum of the entanglement Hamiltonian is then analyzed: e.g., eigenvalue repulsion indicates quantum chaos, as detailed in the main text.

ter [54–57], local fermion-to-qubit mappings [58–60], and quantum-error correction [61–65]. Studying thermalization dynamics of gauge theories, e.g., in early universe and in high-energy particle collisions, remains challenging using first-principles simulation methods [17]. As a first step in experimentally probing thermalization dynamics of gauge theories, we study a Z_2 lattice gauge theory (LGT) in 2 + 1 spacetime dimensions [66–68] using a digital trapped-ion quantum computer [67, 69–71]. We use a chain of fifteen ¹⁷¹Yb⁺ ions to realize a generalpurpose fully-connected digital quantum computer with twelve qubits, and use this computer to natively and accurately encode system's initial state, evolve it in time, and measure observables.

Our analysis relies on EH tomography [72–77] in combination with randomized-measurement protocols [78–89] to learn representations approximating nonequilibrium states. The EH is defined as $H_{\rm E} \equiv -\log(\rho_A)$, where ρ_A denotes the reduced density matrix of subsystem A formed by bipartitioning a (pure) quantum state. The utility of EH as a theoretical and experimental tool stems from the observation that, in many cases, it consists of approximately local operators [90, 91]. This resemblance to conventional Hamiltonians, i.e., energy operators governing system dynamics, facilitates simple theoretical analysis: If an EH is (k-)local, it can be described by parameters whose number scales polynomially with the system size, unlike the matrix ρ_A which requires an exponential number of parameters.

Randomized-measurement protocols [89] can be used to learn the EH: after repeatedly preparing and then measuring the quantum state in a randomly chosen basis, one can fit a parameterized EH to the obtained measurement outcomes. Because the spectrum of $H_{\rm E}$, known as the entanglement spectrum, scales logarithmically with the Schmidt eigenvalues of ρ_A , it is difficult to learn an EH and reproduce the *entire* entanglement spectrum with arbitrary accuracy. Randomized-measurement protocols may require resolving exponentially small probabilities even for relatively small systems, which is out of reach of current quantum devices and perhaps fault-tolerant devices, too. The present experiments can, therefore, only aim at approximating $H_{\rm E}$ and accurately reproducing its low-energy spectrum. Applications of this approach include the verification of topological phases [41, 86].

Nonetheless, what if the precise quantitative structure of $H_{\rm E}$ were not crucial, but only its statistical properties were? This viewpoint is that of random matrix theory [92, 93], where metrics like level distribution and spectral form factor differentiate between integrable and chaotic dynamics, while being indifferent to quantitative details, with the notable exception of symmetries. This statistical perspective underpins our study. Guided by physical insights regarding the expected operator content of a non-equilibrium EH, we ask whether one can learn, from experimental data, a classical representation of a state to answer a simpler question: does a quantum state exhibit *universal* signatures of quantum chaos evident in the statistical properties of its EH? Crucially, can we discern this scenario from one where the state lacks chaotic behavior?

To answer these questions, we focus on two observables indicative of quantum ergodic and chaotic behavior: the entanglement(-Hamiltonian) gap-ratio distribution (EGRD) [38–40] and the entanglement spectral form factor (ESFF) [42, 94]. The EGRD is predicted to exhibit level repulsion for chaotic states, in contrast to uncorrelated levels in non-chaotic scenarios. Similarly, the ESFF is predicted to display a plateau-ramp structure in chaotic states [95]. To experimentally test these predictions, we initialize our system in a product state and perform a quantum quench by digitally evolving the system for a variable length of time. At short evolution times, we observe an EGRD that is characteristic of uncorrelated states. With increasing evolution time, we observe the onset of level repulsion and a ramp-plateau structure in the ESFF, indicative of quantum chaos. Detailed analysis, including comparison with emulated data, suggests that the observed behavior primarily arises from the thermalization dynamics of the isolated quantum system under simulation, with experimental inaccuracies and shortcomings of our ansatz playing a small, though nonnegligible, role. Our work, therefore, establishes presenttime quantum computers as robust tools for studying universal feature of thermalization dynamics in complex many-body systems.

This manuscript is organized as follows. In Secs. II and III, we introduce, respectively, the model and the experimental setup, including state preparation, real-time evolution, and measurement. We discuss our tomography scheme and classical post-analysis in Sec. IV. Finally in Sec. V, we present our main experimental results, reporting the observation of a behavior consistent with quantum chaos in the structure of the EH of time-evolved states. Several Appendices offer additional insights: Appendix A provides further details of the quantum circuits employed; Appendix B investigates finite-size, Trotterization, and late-time effects; and Appendix C offers a comprehensive explanation of our EH-tomography protocol.

II. MODEL

We focus on a Z_2 LGT in 2 + 1 D with the Hamiltonian

$$H = \sum_{\Box} W_{\Box} + g \sum_{\ell} \sigma_{\ell}^{z} \,. \tag{1}$$

Here, $W_{\Box} \equiv \prod_{\ell \in \Box} \sigma_{\ell}^{x}$ is a magnetic-field operator, where \Box denotes the elementary (square) plaquettes of the lattice [see Fig. 2(a)]. σ_{ℓ}^{x} and σ_{ℓ}^{z} are Pauli operators representing gauge link and electric fields on edge ℓ , respectively. These act on spin- $\frac{1}{2}$ hardcore bosons residing on the edges of a two-dimensional spatial square lattice. The first term in Eq. (1) corresponds to the magnetic energy, and the second to the electric energy, with coupling g controlling the relative strength of the two non-commuting contributions. In this formulation, the z basis, therefore, represents the electric basis, while the x basis corresponds to the magnetic basis. The Hamiltonian is expressed in dimensionless units and \hbar is set to one throughout.

We consider the joint +1 eigenspace of the Gauss-law operators $G_j \equiv \prod_{\ell \in +_j} \sigma_{\ell}^z$ as the physical Hilbert space, where $\ell \in +_j$ denote the four links adjacent to a lattice site *j*. In the subsequent discussion, we consider a quasi-1D chain composed of L_x plaquettes along the *x* direction with periodic boundary conditions. Fixed boundary conditions are applied along the short side of the chain (with $L_y = 1$ plaquettes) as illustrated in Fig. 2(b). In this configuration, all Gauss laws contain three links at each




FIG. 2. Model overview. (a) Degrees of freedom (DOFs) in the Z_2 LGT (spin- $\frac{1}{2}$ hardcore bosons denoted by orange circles) residing on the edges of a two-dimensional square array. The electric-field (link) operators on each edge are denoted by pink (blue) ellipses. Four electric operators at site j build the Gauss-law operator G_j , and four link operators covering edges of a plaquette build the magnetic operator W_{\Box} . (b) The spatial lattice considered in this work consisting of $L_x = 10$ plaquettes along the x direction with periodic boundary conditions and $L_y = 1$ plaquette along the y direction with fixed boundary conditions. The short-ribbon operator V_{y} , comprised of electric-field operators at the adjacent links α and β , is shown. (c) The lattice in the dual formulation. Gaugedependent operators (e.g., $\sigma^{z}_{\ell_{\bar{A}A}}$) and the corresponding Gauss laws (e.g., $G_{\bar{A}A}^{\text{dual}}$ denoted by the pink ellipsoid) on the boundary between the system A and its complement \overline{A} (indicated by green thick dashed lines) remain unchanged from the original formulation. However, bulk operators within each subsystem are replaced with gauge-independent Pauli operators $\mu_i^{x,z}$ on each plaquette. The 12 quantum DOFs in this system are mapped to the trapped ion quantum computer, with DOFs shown as pink circles mapped to qubits 1 to 4 in the system and qubits 6 to 11 in the complement. Boundary DOFs, shown as yellow circles, are mapped to qubits 0 and 5. (d) Symmetry operators of the reduced density matrix of a state evolving under Gauss-law-respecting dynamics, as explained in the main text. The thin dashed green line represents the path \mathcal{P} along which the Pauli operators in $V_{x,A}$ act on.

site. Consider a short-ribbon operator $V_y \equiv \sigma_{\alpha}^z \sigma_{\beta}^z$, where α and β are the two opposing links along the inner and the outer circumference at an arbitrary position along the closed chain shown in Fig. 2(b) [96]. This operator commutes with the Hamiltonian and defines superselection sectors of the model; we work in the $V_y = 1$ sector (i.e., short-ribbon operators possess eigenvalue one).

As initial conditions, we randomly select Gauss-lawrespecting electric eigenstates (z-basis product states) of the form $|\Psi(t=0)\rangle \equiv |\Psi_0\rangle = \prod_{\ell} |\ell\rangle$ with $|\ell\rangle \in \{|\uparrow\rangle, |\downarrow\rangle\}$. We then define a subsystem A consisting of $L_A \leq L_x/2$ plaquettes. The reduced density matrix $\rho_A(t)$ of A after evolution time t is related to the density matrix $\rho(t)$ of the whole system as

$$\rho_A(t) \equiv \text{Tr}_{\bar{A}}[\rho(t)], \quad \rho(t) \equiv U(t)\rho_0 U(t)^{\dagger}, \qquad (2)$$

where \overline{A} is the complement of subsystem A shown in Fig. 2 (c) and (d), $\rho_0 \equiv |\Psi_0\rangle\langle\Psi_0|$, and $U(t) \equiv e^{-iHt}$ is the time-evolution operator.

To directly encode the degrees of freedom of the model described onto the qubits, one requires one qubit per link degree of freedom, or $N_q = 3L_x$ qubits in total. To adapt the simulation to the available hardware, we consider a dual formulation of the model that requires only $N_q = L_x + 2$ qubits. In the dual formulation, all operators, except those containing operators having support at the boundaries between the subsystems, labeled $\ell_{\bar{A}A}$ and $\ell_{A\bar{A}}$, are entirely represented using gauge-invariant (Ising dual) spin- $\frac{1}{2}$ variables which act solely within the physical Gauss-law subspace; they are denoted by $\mu_i^{x,z}$ Pauli matrices [see purple circles in Fig. 2(c)]. Concretely, $\mu_i^x \to W_i = \prod_{\ell \in \Box_i} \sigma_\ell^x$, while electric variables are represented by $\mu_i^z \mu_{i-1}^z \equiv \sigma_\ell^z$ where ℓ is the link between plaquettes i and i-1. The corresponding "bulk" Hamiltonian terms for the subsystem A then reads

$$H_A^{\text{dual}} = \sum_{i \in A_{\text{bulk}}} \mu_i^x + g \left(\sum_{\langle i,j \rangle \in A} \mu_i^z \mu_j^z + \kappa \sum_{i \in A} \mu_i^z \right), \quad (3)$$

where $\kappa \equiv 1 + V_y$. Here, $i \in A_{\text{bulk}}$ in the sum over μ_i^x indicates that the original plaquette \Box_i is entirely in A(i.e., does not touch the boundaries). $\langle i, j \rangle \in A$ denotes nearest-neighbor bulk-spin pairs i, j in A. Finally, the sum over μ_i^z runs over all bulk-spin indices i. The terms in the Hamiltonian acting on the complement are defined identically. Importantly, at the boundaries between the subsystems, $\ell_{\bar{A}A}$ and $\ell_{A\bar{A}}$ [see the orange circles in Fig. 2(c)], the gauge-variant variables of the LGT, as described by Eq. (1), are retained. Denoting the Ising-spin index at the A side of one of the boundaries to be k, the Hamiltonian terms coupling the subsystems at this boundary are

$$H_{\bar{A}A}^{\text{dual}} \equiv \mu_k^x \sigma_{\ell_{\bar{A}A}}^x + \mu_{k-1}^x \sigma_{\ell_{\bar{A}A}}^x + g \sigma_{\ell_{\bar{A}A}}^z , \qquad (4)$$

with a similar definition for the other boundary. Note that on the plaquette k containing one boundary link, $\mu_k^x \sigma_{\ell\bar{A}A}^x \equiv W_k$, where $\mu_k^x \equiv \prod_{\ell \in \hat{\Box}_k} \sigma_\ell^x$, with $\tilde{\Box}_i$ referring to all but non-boundary links of plaquette k, and similarly for the plaquette k - 1. The Gauss laws at the boundaries are not eliminated by the duality. The two Gauss laws in the dual model that are independent (one at each boundary) are

$$G_{\bar{A}A}^{\text{dual}} \equiv \mu_k^z \sigma_{\ell_{\bar{A}A}}^z \mu_{k-1}^z \,, \tag{5}$$

and similarly for the boundary at $\ell_{A\bar{A}}$. The dual Hamiltonian of the model is the sum of the terms above:

$$H^{\text{dual}} = H_A^{\text{dual}} + H_{\bar{A}}^{\text{dual}} + H_{\bar{A}A}^{\text{dual}} + H_{A\bar{A}}^{\text{dual}} \,. \tag{6}$$

This formulation ensures that all gauge-invariant variables of the LGT and its dual have the same expectation values. Importantly, by maintaining the Gauss laws at the boundaries, the entanglement properties of the dual formulation are identical to those of the LGT [41]. This is a subtle but crucial distinction from the standard Ising duality [97], which does not preserve the entanglement structure.

The entanglement structure depends on the symmetries of the reduced density matrix, which stem from the remaining Gauss laws. Specifically, $[S_j, \rho_A(t)] = 0$, where in LGT variables, $S_j \equiv \prod_{\ell_j \in A} \sigma_{\ell_j}^x$, with ℓ_j being the two links originating from a boundary site with lattice coordinates $j = (j_x, j_y)$ within A. This is a direct consequence of the state being a Gauss-law eigenstate [41]. In addition, $[V_{x,A}, \rho_A(t)] = 0$ where the *long-ribbon* operator within A is defined as $V_{x,A} \equiv \prod_{\ell \in \mathcal{P}} \sigma_{\ell}^x$ where σ_{ℓ}^x act on the original degrees of freedom along the path \mathcal{P} connecting both boundaries within A, see Fig. 2(d). Not all symmetry operators are independent. For the quasi-1D chain of plaquettes shown in Fig. 2, only one of the two S_i is independent at each boundary. We place one of the boundaries at sites (0,0) and (0,1) and the other at sites $(L_A, 0)$ and $(L_A, 1)$. We then choose the two independent S_j operators to be $S_{(0,0)}$ and $S_{L_A,0}$, henceforth referred to as $S_{A\bar{A}}$ and $S_{\bar{A}A}$, respectively. Note that, when acting on a Gauss-law-respecting state, $V_{x,A} = S_{A\bar{A}}S_{\bar{A}A}$. Consequently, there are four independent symmetry blocks $\rho_A(t) \equiv \bigoplus_{s=1,\dots,4} \rho_{A,s}(t)$ that will play a role in our analysis. In the dual formulation, these symmetry operators are simply $S_{A\bar{A}}^{\text{dual}} \equiv \mu_k^z \sigma_{\ell_{\bar{A}A}}^z$ and $S_{A\bar{A}}^{\text{dual}} \equiv \mu_{k'}^z \sigma_{\ell_{\bar{A}A}}^z$ where k and k' are the first and last plaquettes within A [98]. We denote the eigenvalue $\{1, 1\}, \{1, -1\}, \{-1, 1\}, \{-1,$ and $\{-1, -1\}$ sectors of $\{S_{\bar{A}A}^{\text{dual}}, S_{A\bar{A}}^{\text{dual}}\}$ operators with labels s = 1, 2, 3, 4, respectively.

III. EXPERIMENTAL SETUP

We use a universal trapped-ion digital quantum computer, schematically shown in Fig. 1(a), to simulate the time evolution of the dual formulation of the Z_2 LGT. We consider $L_x = 10$ plaquettes, focusing on the entanglement structure of a subsystem consisting of $L_A = 4$ plaquettes, as shown in Fig. 2(c).

Our experimental system consists of a linear chain of fifteen ¹⁷¹Yb⁺ ions trapped along the **x** axis of a microfabricated ion trap [68] and spaced by approximately 3.7 μ m. The twelve LGT spins are mapped to the ions as shown on Fig. 2(c). The $|\downarrow\rangle$ and $|\uparrow\rangle$ spin states are respectively mapped to the first-order magnetic-field-insensitive hyperfine $|F = 1, m_F = 0\rangle$ and $|F = 0, m_F = 0\rangle$ states of the ²S_{1/2} ground electronic manifold of the ions [99]. The spin states are initialized in the $|\uparrow\rangle$ state by optical pumping and are measured by coupling to the excited ²P_{1/2}, $|F = 0, m_F = 0\rangle$ state using a 369-nm laser, wherein the presence (absence) of emitted photons differentiates between the bright state $|\downarrow\rangle$ and the dark state $|\uparrow\rangle$ with an average 0.3% infidelity, limited by off-resonant photon scattering.

The ions are individually addressed by an equispaced array of 1 μ m-diameter, 355-nm, laser beams oriented perpendicular to the trap surface (z axis), together with a global 300 μ m × 30 μ m beam oriented along the y axis, which is parallel to the trap surface (see Fig. 1(a) and Ref. [5]). These beams drive stimulated Raman transition via the ${}^{2}P_{1/2}$ and ${}^{2}P_{3/2}$ excited electronic states, where a photon is absorbed from the global beam and emitted into an individual beam to flip the qubit from $|\downarrow\rangle$ to $|\uparrow\rangle$ and vice versa. All spin-manipulation light is produced by a single pulsed laser, modified to control its repetition rate to null 4-photon Stark shifts. The phase and amplitude of the 355-nm beams is controlled by single-channel and 32-channel acousto-optic modulators [100].

To minimize addressing errors, crosstalk, and stray coupling to axial motion of the ions [101], single-qubit operations are realized using compound SK1 pulses [102] with Gaussian-shaped sub-segments, with typical infidelities of 0.2%. Entangling operations between any two qubits are realized via variable-angle pairwise Mølmer-Sørensen (MS) gates [103] employing laser-induced statedependent forces on the \approx 3-MHz motional modes of the ion chain oriented along the $\mathbf{y}+\mathbf{z}$ axis. Robust decoupling from the ion motion at the end of each entangling gate is accomplished using amplitude-modulated, detuningrobust pulse waveforms [104]. Typical non-unitary errors of fully-entangling MS gates are 1% and consist predominantly of gate-angle errors and Z-flips of the individual qubits [105].

Our ultimate goal is to study thermalization dynamics in real-time non-equilibrium conditions by learning system's entanglement structure. To this end, we first demonstrate the capability of our digital computer to compute the dynamics of the Z_2 LGT model, including initial-state preparation, time evolution, and subsequent measurements. We initialize the system in a randomly chosen electric eigenstates that respects Gauss law $(|\Psi_0\rangle = |\downarrow\downarrow\downarrow\uparrow\uparrow\downarrow\downarrow\uparrow\uparrow\uparrow\rangle$, see Fig. 2(c) for the qubit mapping, indicated in blue numbers, read from left to right) and simulate its time evolution for q = 0.85 using four Trotter steps that separately implement the noncommuting terms of H^{dual} , i.e., those diagonal in electric (z) or magnetic (x) bases. Our choice of coupling is such that magnetic and electric terms are of similar magnitude, making the model sufficiently non-integrable. For each evolution time t, we apply single-qubit gates to measure all qubits in either the Bloch x or the z bases and repeat each experiment $N_{\rm shots} = 500$ times. The measurement results are used to compute single- and twoqubit observables in the Bloch x and z bases and are plotted in Figure 3 for select observables [106]. The statistical errors are determined by a standard bootstrap resampling analysis with $N_{\text{boots}} = 1000$ bootstrap samples. Quantum circuits implementing Trotterized time evolution and measurements are provided in Appendix A.




FIG. 3. Real-time observables in $Z_2 \ LGT$ in (2+1)D. Measured expectation values of single-qubit (a) and two-qubit (b) observables of qubits $i = 0, \ldots, 5$ of the 10-plaquette theory with g = 0.85 following four steps of Trotterized time evolution with an initial state $|\Psi_0\rangle = |\downarrow\downarrow\downarrow\uparrow\uparrow\downarrow\downarrow\uparrow\uparrow\uparrow\downarrow\rangle$. Each point represents data from $N_{\rm shots} = 500$ experiments, with statistical uncertainties obtained using $N_{\rm boots} = 1000$ bootstrap samples, indicated by error bars. Red (blue) points correspond to measurements in the Bloch z-basis (x-basis); light-blue points indicate the expectation values of non-gauge-invariant observables, which are expected to be zero. Solid lines correspond to exact classically-computed results.

Each Trotter step of evolution uses 12 variable-angle MS gates.

For most evolution times, the expectation values of the physical operators (dark blue and red) show good agreement with the zero-parameter theory prediction (solid lines). We ascribe the $\sim 10\%$ deviation in the evolution rate of the system to miscalibrations of the MS-gate angles. Importantly, the expectation values of gauge-invariance-violating operators (i.e., those that do not commute with the Gauss laws at the two boundaries),

shown in light blue, are consistent with zero within measurement errors. Figure 7 of Appendix A provides a closeup of these observables, showing that gauge-invariance violation is about 5% throughout the time evolution.

To establish the connection between these simulations and the signatures of chaos and thermalization, we now introduce a measurement and analysis protocol that allows the determination of the system's entanglement structure, going beyond low-order correlation functions.

IV. ENTANGLEMENT-HAMILTONIAN SPECTRUM AND TOMOGRAPHY

Our main interest lies in the statistical properties of the Schmidt decomposition of the reduced density matrix of the subsystem A, given by $\rho_A(t) \equiv \sum_{\lambda} p_{\lambda}(t) |\lambda(t)\rangle \langle \lambda(t) |$, and its associated EH [36, 37]

$$H_{\rm E}(t) \equiv -\log(\rho_A(t))\,. \tag{7}$$

The EH's eigenspectrum, called the entanglement spectrum, is defined by $\{\xi_{\lambda}(t) \equiv -\log(p_{\lambda}(t))\}$.

The statistical properties of the entanglement spectrum directly inform the thermalization dynamics of the system. One statistical probe is the distribution, P(r), of the gap ratios, r_{λ} , of the entanglement spectrum [38],

$$r_{\lambda} \equiv \frac{\min(\delta_{\lambda}, \delta_{\lambda-1})}{\max(\delta_{\lambda}, \delta_{\lambda-1})}, \qquad (8)$$

where $\delta_{\lambda} \equiv \xi_{\lambda} - \xi_{\lambda-1} \geq 0$ are the gaps between the eigenvalues ξ_{λ} of $H_{\rm E}$ and $r \equiv \{r_{\lambda}\}$. According to random matrix theory of Hermitian matrices, this quantity distinguishes chaotic from nonchaotic behavior depending on whether the distribution is centered away from zero (level repulsion) or centered around zero (uncorrelated levels) [92]. The gap ratios, nonetheless, only capture local correlations of the entanglement spectrum and, owing to their logarithmic nature, treat entanglement eigenvalues $\xi_{\lambda} \equiv -\log(p_{\lambda})$ associated with very small probabilities $p_{\lambda} \ll 1$ no differently than those where $p_{\lambda} \approx O(1)$.

A complementary view of the statistics of the entanglement spectrum that captures global correlations in the level distribution is afforded by the entanglement spectral form factor (ESFF) [40, 92],

$$\mathcal{F}(\theta;t) \equiv \left\langle \frac{1}{\mathcal{R}_{A}^{2}(t)} \sum_{\lambda,\lambda'} e^{i\theta[\xi_{\lambda}(t) - \xi_{\lambda'}(t)]} \right\rangle, \qquad (9)$$

where $\langle \cdot \rangle$ denotes the average over initial states $|\Psi_0\rangle$. Here, $\mathcal{R}_A \equiv \lim_{\alpha \to 0} \exp\{\frac{1}{1-\alpha}\log(\sum_{\lambda} p_{\lambda}^{\alpha})\}$ [40] is the effective rank of $H_{\rm E}$, whose value depends on the state and lies in the range $[1, 2^{L_A}]$ [107]. Starting from an initial flat behavior, a ramp-plateau structure in the ESFF as function of the 'Fourier time' θ is known to indicate ergodic behavior and imply quantum chaos [40].

While in classical computation of a tractably small system, the state and its entanglement are readily available, in a quantum-simulation experiment, these must be inferred from measurements. To tackle this challenge, we constrain the operator content of the EH to obtain an approximate $H_{\rm E}$, and utilize an EH-tomography scheme based on randomized measurements [72–77]. Concretely, we perform a single-layer, single-qubit randomizedmeasurement [108] of the time-evolved state. At the end of the time evolution, this protocol applies one of $N_{\mathcal{U}}$ different gates $\mathcal{U}_j = \bigotimes_{i=0,\cdots,N_q-1} u_{i,j}$ for $j = 1, \cdots, N_{\mathcal{U}}$ consisting of independent single-qubit gates $u_{i,j}$ sampled from a unitary 2-design [109]. For each \mathcal{U}_j , $N_{\rm shots}$ experiments are performed, measuring all qubits in the Bloch z basis.

We compare the relative frequencies of the different bitstrings b to the prediction based on an approximate EH inspired by the Bisognano-Wichmann (BW) theorem [90, 91]. Explicitly, we assume that $H_{\rm E}(t)$ is a linear combination of k-local terms \mathcal{O}_i , i.e.

$$H_{\rm E}(t; \{\beta_i\}) \equiv \sum_i \beta_i(t) \mathcal{O}_i \,. \tag{10}$$

To find a suitable set of operators $\{O_i\}$, we proceed as follows. Starting from the operator content of the physical Hamiltonian in Eq. (1), containing at most 4-local terms (2-local in the dual formulation), new operators are generated iteratively by forming non-trivial commutators from the existing set. This process can be halted after two iterations ('commutators of commutators') resulting in a maximum of 7-local operators within the LGT framework (or 3-local in the dual formulation). We exclude operators that violate the Gauss laws and those that can be related to others upon application of the Gauss-laws. To ensure Hermiticity, we adjust prefactors and create linear combinations with conjugates [110]. To find the parameters β_i , a global optimization routine, based on standard MATLAB libraries [111], is used to determine the parameterization that most closely reproduces all measurements (refer to Appendix C for details).

V. EXPERIMENTAL STUDY OF THERMALIZATION DYNAMICS OF Z_2 LGT

To study the evolution of the entanglement spectrum in the Z_2 LGT, we repeat the time-evolution experiments shown in Fig. 3 starting from 6 randomly chosen initial electric eigenstates that satisfy the Gauss laws [112]. Since these are not the eigenstates of the full Hamiltonian H, the system undergoes non-trivial time evolution when evolved under H. For each evolution time t, symmetry sector, and initial state, we subsequently perform randomized-measurement tomography with g = 0.85, $N_{\mathcal{U}} = 24$ bases, and $N_{\text{shots}} = 750$ bitstring measurements in each basis, and use these measurements to reconstruct the EH as described in the previous section. This procedure yields a set of gap ratios for each initial state, symmetry sector, and evolution time.




FIG. 4. Statistics of the gap ratios of the spectrum of the entanglement Hamiltonian. (a) Time evolution of the average gap ratio averaged over 6 randomly-drawn initial states and all symmetry sectors. The horizontal lines represent the averages for non-repulsive (Poisson, blue dotted) and repulsive (GUE, red dashed) distributions. Error bars indicate standard deviation over initial states and symmetry sectors. (b) Distribution of the entanglement-spectrum gap ratios, combined across 6 randomly-drawn initial states, all symmetry sectors, and all times in each of the regimes (I), (II), and (III). A total of 504, 504 and 336 gaps are quantum-computed, and the average is shown in orange, along with simulated Bisognano-Wichmann results in the limit of infinite measurements in cyan, and the exact distributions in black. Blue-dotted and red-dashed curves represent Poisson and GUE distributions.

In Fig. 4(a), we plot in orange points the average gap ratio $\langle r \rangle \equiv \sum_{r} r \bar{P}(r)$ of the reconstructed EH as a function of the scaled evolution time gt. The plotted gap ratios are averaged over both the symmetry sectors and the 6 randomly chosen initial states. The black lines correspond to the predicted exact distributions following the Trotterized time evolution. The cyan lines correspond to the EH obtained from an optimal BW-inspired ansatz, where we numerically minimize the the relative entropy (Kullback–Leibler divergence) between the exact state and the ansatz, corresponding to the limit of infinitely many measurements. A buildup of the level repulsion is discernible as the observed average gap ratio $\langle r \rangle$ evolves from ≈ 0.4 predicted for a non-repulsive Poisson distribution (blue dashed line) towards ≈ 0.6 characteristic of repulsive level statistics of a Gaussian Unitary Ensemble (GUE) [38]. Three time regimes (I), (II), and (III) are identified that correspond, respectively, to the evolution of the predicted average gap ratio toward the Poisson-distribution value, toward the GUE value, and to saturation at the GUE value.

In Fig. 4(b) we plot, for the three ranges of evolution times, the corresponding normalized distribution of the gap ratios, P(r), combined over 6 random initial states and 4 symmetry sectors. We observe a clear transition from early-time non-repulsion (Poisson distribution, blue dotted line) in regime (I) to level repulsion (GUE, red) in regime (III). At intermediate times, a distribution is observed between the initial absence of level repulsion and the subsequent emergence of level repulsion. Error bars and bands denote the variance resulting from averaging over symmetry sectors and initial states. The predicted exact distribution (black line) exhibits a sharper peak around zero at the earliest times due to the reduced density matrix not being full rank. Conversely, the BW-inspired ansatz (cvan line) generally parametrizes a full-rank matrix unless couplings are very finely tuned,




FIG. 5. Entanglement spectral form factor. The average entanglement spectral form factor across various initial states, symmetry sectors, and three distinct regimes (I), (II), and (III) identified in Fig. 4. The figures display exact results (black curves), infinite-measurement outcomes (cyan), and quantum-computed experimental data (orange). In panel (III), a purple dotted line indicates a fit of the ramp observed in the quantum-computed data. Our normalization ensures $\langle \mathcal{F}(0) \rangle = 1$, with the plateau occurring at $\langle \mathcal{F}(\infty) \rangle = 1/d_s$, where d_s denotes the dimension of a symmetry block, see Appendix B for details. Shaded areas indicate the standard deviation over initial states, symmetry sectors, and times.

resulting in an overestimation of level repulsion in the initial stages. Calculations indicate that the onset of the three regimes remains consistent for any randomly chosen initial states and depends only on the (subsystem) size when measured in coupling-independent units gt, indicating the universality of thermalization dynamics.

In Fig. 5 we plot the ESSF reconstructed from the data used in Fig. 4. The three panels of the plot correspond to the three time regimes discussed in relation to the EGRD evolution. The displayed theory curves are averaged over initial states, symmetry sectors, and each

of the three time ranges. Starting from an initial flat behavior of the ESSR as a function of θ in panel (I), our data in panels (II) and (III) clearly show the buildup of a ramp-plateau structure, indicating ergodic behavior. In panel (III), we show a fit (purple dotted line) to the observed ramp, indicating a power-law behavior θ^{κ} where $\kappa = 0.6 \pm 0.2$, with the fit error determined by changing the fit regime. This exponent is consistent with the results obtained from the numerical analysis of a significantly larger system in Appendix B.

Our results demonstrate that, using observables that reveal universal statistical or global features of entanglement spectrum, the onset of chaotic behavior can be robustly traced in the entanglement dynamics of arbitrary initial states. Importantly, time digitization and experimental infidelities in current systems do not significantly impact qualitative features of thermalization dynamics.

VI. CONCLUSIONS

Using a digital trapped-ion quantum computer, we observe early-stage thermalization dynamics in a \mathbb{Z}_2 lattice gauge theory in 2+1 dimensions by generating nonequilibrium quantum states and measuring their entanglement structure. As performing quantitative state tomography is extremely challenging and resource intensive, we focused on a simpler question: can a randomizedmeasurement-based entanglement-Hamiltonian tomography recover universal properties of the EH that indicate quantum chaos? We experimentally determine the entanglement spectrum of time-dependent quantum states and use two properties of this spectrum, namely the entanglement gap-ratio distribution [Eq. (8)] and the entanglement spectral form factor [Eq. (9)], as indicators of the emergence of quantum chaos. Our data indicate that both quantities behave as expected: the initially non-repulsive level distribution transitions to a repulsive one. Likewise, a ramp-plateau feature of the ESFF is initially absent but builds up with time. Our results indicate that the timescales for these entanglement-based phenomena mainly depend on subsystem size.

The key technique enabling our analysis is a Bisognano-Wichmann-inspired ansatz for the EH, which allows the parameterization of a nonequilibrium state using a polynomial number of parameters. While this classical parameterization optimally reproduces the observed (randomized) measurement results, in Appendix \mathbf{C} , we show that the higher-lying part of the EH spectrum is not quantitatively recovered. Indeed, the measurement cost for *precise* state tomography still scales exponentially with the subsystem size. Fortunately, to detect the presence of quantum chaos, one needs to only distinguish repulsion from non-repulsion in the EGRD or identify a ramp-plateau structure in the ESFF. Our work indicates that selecting a sufficiently small subsystem of a much larger, potentially classically nonsimulable system, can sufficiently constrain these observables.

Our randomized-measurement procedure remains classically simulable because the lattice sizes we considered are manageable with exact diagonalization. Our procedure is also heavily tailored at minimizing the computational load on the quantum computer, avoiding deep quantum circuits and the need for error mitigation, at the cost of classical post-processing. To extend our approach to larger systems, several steps can be implemented in future work:

- The employed single-qubit one-layer randomization strategy is symmetry ignorant, i.e., it randomizes regardless of the known symmetry structure of the subsystem density matrix $\rho_A(t)$. Although our scheme is tomographically complete, and enables state reconstruction along with its symmetries, it is inefficient. Symmetry-conscious protocols, for the LGT in this work and for other models, have been developed [113–115] and could be used in future work.
- A major methodological uncertainty is the BWinspired parameterization of the EH, which is entirely heuristic. While there is research exploring the operator content of the EH of ground, excited, and thermal states [116], further investigations are needed into the applicability and limitations of EH-based schemes for far-from-equilibrium states. Generally, the BW(-inspired) ansätze describe the *low-energy* regime of the EH well, and incorporating progressively more nonlocal terms improves convergence into the bulk [74] which the EGRD, and to a lesser extent the ESFF, predominantly rely on. Notably, our analysis reveals that despite the quantitative discrepancy in the bulk, the statistical distribution of the EH appears to be accurately reproduced. This suggests that it may be unnecessary to perform *precise* state tomography when one's interest is solely in statistical properties.
- Trotterization corresponds to time evolution with an effective, rather than the desired, Hamiltonian. We observe that employing too few Trotter steps leads to poor convergence of our optimization procedure for the BW-inspired ansatz. In this case, more nonlocal operators must be included in the ansatz; constraining these becomes challenging, especially when relying on a limited number of measurements. Since using a large number of Trotter steps increases experimental errors, the optimal Trotterization and its interplay with the BW ansatz should be further investigated.
- Detailed comparison between the emulator and the experimental data (see Appendix C) highlights the influence of device errors, primarily for the time evolution and tomography steps of our algorithm. The dominant errors in our study are Z-flip errors and over- and under-rotations due to mechanical

motion of the ions [101]. Our tomography protocol is especially susceptible to small single-qubit rotation errors during randomization [108]. Recently demonstrated sympathetic cooling during circuit execution [101] would allow the needed measurement circuits to be executed with higher fidelity. The same technique could extend our study to larger systems, later times, and more Trotter steps, e.g., allowing us to directly test the eigenstate thermalization hypothesis in a digital quantumcomputing set-up.

In summary, our results demonstrate that entanglement structure is a measurable quantity in presentday LGT quantum-simulation experiments, and illustrates the potential value of our approach to probe thermalization dynamics and its robust universal features in strongly coupled isolated quantum many-body systems. A compelling future direction is to extend the investigation to later times, to probe aspects such as pre-thermalization [10, 12, 117–119] or fluctuationdissipation relations [120–123], once experimental capabilities permit. This would also allow for probing the applicability of the Eigenstate Thermalization Hypothesis, e.g., in systems with non-Abelian symmetries [124] and other gauge theories [125, 126]. Furthermore, the experimental and theoretical tools of this study can be applied in a number of other applications, including obtaining thermodynamic quantities such as work and heat exchanged during nonequilibrium processes [127], and detecting phases of matter, including topological phases [86], in quantum-simulation experiments.

- J. Eisert, M. Friesdorf, and C. Gogolin, Quantum manybody systems out of equilibrium, Nature Physics 11, 124 (2015).
- [2] M. Schreiber, S. S. Hodgman, P. Bordia, H. P. Lüschen, M. H. Fischer, R. Vosk, E. Altman, U. Schneider, and I. Bloch, Observation of many-body localization of interacting fermions in a quasirandom optical lattice, Science 349, 842 (2015).
- [3] J. Schachenmayer, L. Pollet, M. Troyer, and A. J. Daley, Thermalization of strongly interacting bosons after spontaneous emissions in optical lattices, EPJ Quantum Technology 2, 1 (2015).
- [4] A. M. Kaufman, M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, Quantum thermalization through entanglement in an isolated many-body system, Science **353**, 794 (2016).
- [5] C. Eigen, J. A. Glidden, R. Lopes, E. A. Cornell, R. P. Smith, and Z. Hadzibabic, Universal prethermal dynamics of bose gases quenched to unitarity, Nature 563, 221 (2018).
- [6] M. Ueda, Quantum equilibration, thermalization and prethermalization in ultracold atoms, Nature Reviews Physics 2, 669 (2020).

ACKNOWLEDGMENT

N.M. thanks H. Froland, A. Polkovnikov, M. Savage, M. Srednicki, X. Yao, T. Zache, P. Zoller, and the participants of the InQubator for Quantum Simulation (IQuS) workshop "Thermalization, from Cold Atoms to Hot Quantum Chromodynamics "(https://iqus.uw. edu/events/iqus-workshop-thermalization/) at the University of Washington in September 2023 for many valuable discussions leading to this work. M.C. thanks L. Feng for valuable discussions and help with the experimental setup leading to this work. N.M. acknowledges funding by the Department of Energy (DOE), Office of Science, Office of Nuclear Physics, IQuS (https: //iqus.uw.edu), via the program on Quantum Horizons: QIS Research and Innovation for Nuclear Science under Award DE-SC0020970. Z.D., M.C., and T.W. were supported by the National Science Foundation's Quantum Leap Challenge Institute for Robust Quantum Simulation under Award OMA-2120757. Z.D. further acknowledges support by the DOE, Office of Science, Early Career Award DE-SC0020271. This work is further supported by a collaboration between the US DOE and other Agencies. This material is based upon work supported by the DOE, Office of Science, National Quantum Information Science Research Centers, Quantum Systems Accelerator.

COMPETING INTERESTS

M.C. is a co-inventor on patents that are licensed from the University of Maryland to IonQ, Inc.

- [7] Y. Le, Y. Zhang, S. Gopalakrishnan, M. Rigol, and D. S. Weiss, Observation of hydrodynamization and local prethermalization in 1d bose gases, Nature 618, 494 (2023).
- [8] Z.-Y. Zhou, G.-X. Su, J. C. Halimeh, R. Ott, H. Sun, P. Hauke, B. Yang, Z.-S. Yuan, J. Berges, and J.-W. Pan, Thermalization dynamics of a gauge theory on a quantum simulator, Science **377**, 311 (2022).
- [9] J. Smith, A. Lee, P. Richerme, B. Neyenhuis, P. W. Hess, P. Hauke, M. Heyl, D. A. Huse, and C. Monroe, Many-body localization in a quantum simulator with programmable random disorder, Nat. Phys. 12, 907 (2016).
- [10] B. Neyenhuis, J. Zhang, P. W. Hess, J. Smith, A. C. Lee, P. Richerme, Z.-X. Gong, A. V. Gorshkov, and C. Monroe, Observation of prethermalization in long-range interacting spin chains, Science Advances 3, 10.1126/sciadv.1700672 (2017).
- [11] W. Morong, F. Liu, P. Becker, K. S. Collins, L. Feng, A. Kyprianidis, G. Pagano, T. You, A. V. Gorshkov, and C. Monroe, Observation of stark many-body localization without disorder, Nature **599**, 393 (2021).

- [12] A. Kyprianidis, F. Machado, W. Morong, P. Becker, K. S. Collins, D. V. Else, L. Feng, P. W. Hess, C. Nayak, G. Pagano, N. Y. Yao, and C. Monroe, Observation of a prethermal discrete time crystal, Science **372**, 1192 (2021).
- [13] R. Nandkishore and D. A. Huse, Many-body localization and thermalization in quantum statistical mechanics, Annu. Rev. Condens. Matter Phys. 6, 15 (2015).
- [14] F. Borgonovi, F. M. Izrailev, L. F. Santos, and V. G. Zelevinsky, Quantum chaos and thermalization in isolated systems of interacting particles, Physics Reports 626, 1 (2016).
- [15] R. Micha and I. I. Tkachev, Turbulent thermalization, Physical Review D 70, 043538 (2004).
- [16] R. Baier, A. H. Mueller, D. Schiff, and D. T. Son, "bottom-up" thermalization in heavy ion collisions, Physics Letters B 502, 51 (2001).
- [17] QCD thermalization: Ab initio approaches and interdisciplinary connections, author=Berges, Jürgen and Heller, Michal P and Mazeliauskas, Aleksas and Venugopalan, Raju, Reviews of Modern Physics **93**, 035003 (2021).
- [18] J. M. Deutsch, Quantum statistical mechanics in a closed system, Physical review a 43, 2046 (1991).
- [19] M. Srednicki, Chaos and quantum thermalization, Physical review e 50, 888 (1994).
- [20] M. Rigol, V. Dunjko, and M. Olshanii, Thermalization and its mechanism for generic isolated quantum systems, Nature 452, 854 (2008).
- [21] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics, Advances in Physics 65, 239 (2016).
- [22] E. Grumbling, Quantum computing, (No Title) (2019).
- [23] F. Arute, K. Arya, R. Babbush, D. Bacon, J. C. Bardin, R. Barends, R. Biswas, S. Boixo, F. G. Brandao, D. A. Buell, *et al.*, Quantum supremacy using a programmable superconducting processor, Nature **574**, 505 (2019).
- [24] I. H. Deutsch, Harnessing the power of the second quantum revolution, PRX Quantum 1, 020101 (2020).
- [25] D. Bluvstein, A. Omran, H. Levine, A. Keesling, G. Semeghini, S. Ebadi, T. T. Wang, A. A. Michailidis, N. Maskara, W. W. Ho, *et al.*, Controlling quantum many-body dynamics in driven rydberg atom arrays, Science **371**, 1355 (2021).
- [26] S. J. Evered, D. Bluvstein, M. Kalinowski, S. Ebadi, T. Manovitz, H. Zhou, S. H. Li, A. A. Geim, T. T. Wang, N. Maskara, *et al.*, High-fidelity parallel entangling gates on a neutral atom quantum computer, arXiv preprint arXiv:2304.05420 (2023).
- [27] J. I. Cirac and P. Zoller, Goals and opportunities in quantum simulation, Nature physics 8, 264 (2012).
- [28] A. J. Daley, I. Bloch, C. Kokail, S. Flannigan, N. Pearson, M. Troyer, and P. Zoller, Practical quantum advantage in quantum simulation, Nature 607, 667 (2022).
- [29] C. W. Bauer, Z. Davoudi, A. B. Balantekin, T. Bhattacharya, M. Carena, W. A. De Jong, P. Draper, A. El-Khadra, N. Gemelke, M. Hanada, *et al.*, Quantum simulation for high-energy physics, PRX quantum 4, 027001 (2023).
- [30] F. Haake, *Quantum signatures of chaos* (Springer, 1991).

- [31] S. Goldstein, J. L. Lebowitz, C. Mastrodonato, R. Tumulka, and N. Zanghi, Normal typicality and von neumann's quantum ergodic theorem, Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences 466, 3203 (2010).
- [32] M. V. Berry and M. Tabor, Level clustering in the regular spectrum, Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences **356**, 375 (1977).
- [33] O. Bohigas, M.-J. Giannoni, and C. Schmit, Characterization of chaotic quantum spectra and universality of level fluctuation laws, Physical review letters 52, 1 (1984).
- [34] L. K. Joshi, A. Elben, A. Vikram, B. Vermersch, V. Galitski, and P. Zoller, Probing many-body quantum chaos with quantum simulators, Physical Review X 12, 011018 (2022).
- [35] M. Pandey, P. W. Claeys, D. K. Campbell, A. Polkovnikov, and D. Sels, Adiabatic eigenstate deformations as a sensitive probe for quantum chaos, Physical Review X 10, 041017 (2020).
- [36] H. Li and F. D. M. Haldane, Entanglement spectrum as a generalization of entanglement entropy: Identification of topological order in non-abelian fractional quantum hall effect states, Physical review letters **101**, 010504 (2008).
- [37] M. Dalmonte, V. Eisler, M. Falconi, and B. Vermersch, Entanglement hamiltonians: from field theory to lattice models and experiments, Annalen der Physik 534, 2200064 (2022).
- [38] V. Oganesyan and D. A. Huse, Localization of interacting fermions at high temperature, Physical review b 75, 155111 (2007).
- [39] T. Rakovszky, S. Gopalakrishnan, S. Parameswaran, and F. Pollmann, Signatures of information scrambling in the dynamics of the entanglement spectrum, Physical Review B 100, 125115 (2019).
- [40] P.-Y. Chang, X. Chen, S. Gopalakrishnan, and J. Pixley, Evolution of entanglement spectra under generic quantum dynamics, Physical review letters 123, 190602 (2019).
- [41] N. Mueller, T. V. Zache, and R. Ott, Thermalization of gauge theories from their entanglement spectrum, Physical Review Letters 129, 011601 (2022).
- [42] C.-T. Ma and C.-H. Wu, Quantum entanglement and spectral form factor, International Journal of Theoretical Physics 61, 272 (2022).
- [43] H. Froland, T. V. Zache, R. Ott, and N. Mueller, Entanglement structure of non-gaussian states and how to measure it (2024), arXiv:2407.12083.
- [44] M. Suzuki, General theory of fractal path integrals with applications to many-body theories and statistical physics, Journal of mathematical physics **32**, 400 (1991).
- [45] A. M. Childs, Y. Su, M. C. Tran, N. Wiebe, and S. Zhu, Theory of trotter error with commutator scaling, Physical Review X 11, 011020 (2021).
- [46] L. Lin, Lecture notes on quantum algorithms for scientific computation, arXiv preprint arXiv:2201.08309 (2022).
- [47] A. M. Childs, Lecture notes on quantum algorithms, Lecture notes at University of Maryland 5 (2017).
- [48] M. C. Bañuls *et al.*, Simulating Lattice Gauge Theories within Quantum Technologies, Eur. Phys. J. D 74, 165

(2020), arXiv:1911.00003 [quant-ph].

- [49] N. Klco, A. Roggero, and M. J. Savage, Standard model physics and the digital quantum revolution: thoughts about the interface, Rept. Prog. Phys. 85, 064301 (2022), arXiv:2107.04769 [quant-ph].
- [50] C. W. Bauer, Z. Davoudi, N. Klco, and M. J. Savage, Quantum simulation of fundamental particles and forces, Nature Rev. Phys. 5, 420 (2023), arXiv:2404.06298 [hep-ph].
- [51] A. Di Meglio, K. Jansen, I. Tavernelli, C. Alexandrou, S. Arunachalam, C. W. Bauer, K. Borras, S. Carrazza, A. Crippa, V. Croft, *et al.*, Quantum computing for high-energy physics: State of the art and challenges. summary of the qc4hep working group, arXiv preprint arXiv:2307.03236 (2023).
- [52] I. J. Aitchison and A. J. Hey, Gauge Theories in Particle Physics: A Practical Introduction, -2 Volume set (Taylor & Francis, 2012).
- [53] C. Quigg, Gauge theories of strong, weak, and electromagnetic interactions (CRC Press, 2021).
- [54] E. Fradkin, Field theories of condensed matter physics (Cambridge University Press, 2013).
- [55] H. Kleinert, Gauge Fields in Condensed Matter: Vol.
 1: Superflow and Vortex Lines (Disorder Fields, Phase Transitions) Vol. 2: Stresses and Defects (Differential Geometry, Crystal Melting) (World Scientific, 1989).
- [56] X.-G. Wen, Topological orders in rigid states, International Journal of Modern Physics B 4, 239 (1990).
- [57] M. A. Levin and X.-G. Wen, String-net condensation: A physical mechanism for topological phases, Physical Review B 71, 045110 (2005).
- [58] Y.-A. Chen, A. Kapustin, and D. Radičević, Exact bosonization in two spatial dimensions and a new class of lattice gauge theories, Annals of Physics **393**, 234 (2018).
- [59] Y.-A. Chen, Exact bosonization in arbitrary dimensions, Physical Review Research 2, 033527 (2020).
- [60] Y.-A. Chen and Y. Xu, Equivalence between fermion-toqubit mappings in two spatial dimensions, PRX Quantum 4, 010326 (2023).
- [61] A. Y. Kitaev, Fault-tolerant quantum computation by anyons, Annals of physics 303, 2 (2003).
- [62] A. Kitaev, Anyons in an exactly solved model and beyond, Annals of Physics **321**, 2 (2006).
- [63] S. Das Sarma, M. Freedman, and C. Nayak, Topological quantum computation, Physics Today 59, 32 (2006).
- [64] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Non-abelian anyons and topological quantum computation, Rev. Mod. Phys. 80, 1083 (2008).
- [65] V. Lahtinen and J. K. Pachos, A Short Introduction to Topological Quantum Computation, SciPost Phys. 3, 021 (2017).
- [66] P. L. W. Maunz, *High Optical Access Trap 2.0.*, Tech. Rep. (Sandia National Lab.(SNL-NM), Albuquerque, NM (United States), 2016).
- [67] C. Monroe, W. C. Campbell, L.-M. Duan, Z.-X. Gong, A. V. Gorshkov, P. W. Hess, R. Islam, K. Kim, N. M. Linke, G. Pagano, *et al.*, Programmable quantum simulations of spin systems with trapped ions, Reviews of Modern Physics **93**, 025001 (2021).
- [68] P. L. W. Maunz, High optical access trap 2.0. 10.2172/1237003 (2016).

- [69] I. Pogorelov, T. Feldker, C. D. Marciniak, L. Postler, G. Jacob, O. Krieglsteiner, V. Podlesnic, M. Meth, V. Negnevitsky, M. Stadler, *et al.*, Compact ion-trap quantum computing demonstrator, PRX Quantum 2, 020343 (2021).
- [70] R. Yao, W.-Q. Lian, Y.-K. Wu, G.-X. Wang, B.-W. Li, Q.-X. Mei, B.-X. Qi, L. Yao, Z.-C. Zhou, L. He, *et al.*, Experimental realization of a multiqubit quantum memory in a 218-ion chain, Physical Review A **106**, 062617 (2022).
- [71] J. G. Bohnet, B. C. Sawyer, J. W. Britton, M. L. Wall, A. M. Rey, M. Foss-Feig, and J. J. Bollinger, Quantum spin dynamics and entanglement generation with hundreds of trapped ions, Science **352**, 1297 (2016).
- [72] H. Pichler, G. Zhu, A. Seif, P. Zoller, and M. Hafezi, Measurement protocol for the entanglement spectrum of cold atoms, Phys. Rev. X 6, 041033 (2016).
- [73] M. Dalmonte, B. Vermersch, and P. Zoller, Quantum simulation and spectroscopy of entanglement hamiltonians, Nat. Phys. 14, 827 (2018).
- [74] C. Kokail, R. van Bijnen, A. Elben, B. Vermersch, and P. Zoller, Entanglement hamiltonian tomography in quantum simulation, Nat. Phys. 17, 936 (2021).
- [75] C. Kokail, B. Sundar, T. V. Zache, A. Elben, B. Vermersch, M. Dalmonte, R. van Bijnen, and P. Zoller, Quantum variational learning of the entanglement hamiltonian, Phys. Rev. Lett. **127**, 170501 (2021).
- [76] T. V. Zache, C. Kokail, B. Sundar, and P. Zoller, Entanglement spectroscopy and probing the Li-Haldane conjecture in topological quantum matter, Quantum 6, 702 (2022).
- [77] N. Mueller, J. A. Carolan, A. Connelly, Z. Davoudi, E. F. Dumitrescu, and K. Yeter-Aydeniz, Quantum computation of dynamical quantum phase transitions and entanglement tomography in a lattice gauge theory, PRX Quantum 4, 030323 (2023).
- [78] M. Paini and A. Kalev, An approximate description of quantum states, arXiv preprint arXiv:1910.10543 (2019).
- [79] H.-Y. Huang, R. Kueng, and J. Preskill, Predicting many properties of a quantum system from very few measurements, Nat. Phys. 16, 1050 (2020).
- [80] H.-Y. Huang, R. Kueng, and J. Preskill, Efficient estimation of pauli observables by derandomization, Phys. Rev. Lett. 127, 030503 (2021).
- [81] H.-Y. Hu, S. Choi, and Y.-Z. You, Classical shadow tomography with locally scrambled quantum dynamics, arXiv preprint arXiv:2107.04817 (2021).
- [82] J. Kunjummen, M. C. Tran, D. Carney, and J. M. Taylor, Shadow process tomography of quantum channels, Physical Review A 107, 042403 (2023).
- [83] R. Levy, D. Luo, and B. K. Clark, Classical shadows for quantum process tomography on near-term quantum computers, Physical Review Research 6, 013029 (2024).
- [84] H.-Y. R. Huang, M. B. Broughton, J. Cotler, S. Chen, J. Li, M. Mohseni, H. Neven, R. Babbush, R. Kueng, J. Preskill, *et al.*, Demonstrating quantum advantage in learning from experiments, Science **376**, 1182 (2021).
- [85] H.-Y. Huang, Learning quantum states from their classical shadows, Nat. Rev. Phys. 4 (2022).
- [86] J. Bringewatt, J. Kunjummen, and N. Mueller, Randomized measurement protocols for lattice gauge theories, Quantum 8, 1300 (2024).

- [87] A. Elben, B. Vermersch, C. F. Roos, and P. Zoller, Statistical correlations between locally randomized measurements: A toolbox for probing entanglement in many-body quantum states, Phys. Rev. A 99, 052323 (2019).
- [88] A. Elben, R. Kueng, H.-Y. R. Huang, R. van Bijnen, C. Kokail, M. Dalmonte, P. Calabrese, B. Kraus, J. Preskill, P. Zoller, *et al.*, Mixed-state entanglement from local randomized measurements, Phys. Rev. Lett. **125**, 200501 (2020).
- [89] A. Elben, S. T. Flammia, H.-Y. Huang, R. Kueng, J. Preskill, B. Vermersch, and P. Zoller, The randomized measurement toolbox, Nat. Rev. Phys. 5, 9 (2023).
- [90] J. J. Bisognano and E. H. Wichmann, On the duality condition for a hermitian scalar field, Journal of Mathematical Physics 16, 985 (1975).
- [91] J. J. Bisognano and E. H. Wichmann, On the duality condition for quantum fields, Journal of mathematical physics 17, 303 (1976).
- [92] T. Guhr, A. Müller-Groeling, and H. A. Weidenmüller, Random-matrix theories in quantum physics: common concepts, Physics Reports 299, 189 (1998).
- [93] M. L. Mehta, Random matrices (Elsevier, 2004).
- [94] C. B. Dağ, S. I. Mistakidis, A. Chan, and H. R. Sadeghpour, Many-body quantum chaos in stroboscopicallydriven cold atoms, Communications Physics 6, 136 (2023).
- [95] Strictly speaking, both indicate quantum ergodicity which usually implies quantum chaos.
- [96] Note that V_y is independent of α and β because, for Gauss law eigenstates, the operator can be acted on by Gauss law operators in such a way that it can be moved around the lattice.
- [97] F. J. Wegner, Duality in generalized ising models, Topol. Asp. Condens. Matter Phys.(Oxford University Press, 2017) pp , 219 (2015).
- [98] This discussion is general, independent of the choice of a V_y eigensector.
- [99] Here, we use the quantum-information convention for the spin states, i.e., $|\uparrow\rangle \equiv |0\rangle$ and $|\downarrow\rangle \equiv |1\rangle$.
- [100] Provided by L3 Harris Corporation.
- [101] M. Cetina, L. Egan, C. Noel, M. Goldman, D. Biswas, A. Risinger, D. Zhu, and C. Monroe, Control of transverse motion for quantum gates on individually addressed atomic qubits, PRX Quantum 3 (2022).
- [102] K. R. Brown, A. W. Harrow, and I. L. Chuang, Arbitrarily accurate composite pulse sequences, Phys. Rev. A 70, 052318 (2004).
- [103] A. Sørensen and K. Mølmer, Quantum computation with ions in thermal motion, Physical review letters 82, 1971 (1999).
- [104] P. H. Leung, K. A. Landsman, C. Figgatt, N. M. Linke, C. Monroe, and K. R. Brown, Robust 2-qubit gates in a linear ion crystal using a frequency-modulated driving force, Phys. Rev. Lett. **120**, 020501 (2018).
- [105] S. Huang, K. R. Brown, and M. Cetina, Comparing shor and steane error correction using the bacon-shor code (2023), arXiv:2312.10851.
- [106] For simplicity, the operators whose expectation values are measured are generally noted as Pauli σ operators. When acting on the bulk dual spins, they should be thought of as the Pauli μ operators, while when acting on the boundary spins, they are the Pauli σ operators in the original LGT.

- [107] In practice, both statistical measures are typically used to describe the properties of *physical* Hamiltonians, whose rank equals the Hilbert-space dimension. For EHs, we employ regularization suitable for non-fullrank matrices. To address levels with probabilities close to zero within machine precision, we define an effective rank. Only the \mathcal{R}_A lowest levels of the EH are included in the analysis. However, experimental constraints, such as finite statistics and device errors, typically lead to EHs that are reconstructed as nearly or exactly full rank, rendering any regularization obsolete. Nonetheless, non-full-rank EHs occur for states computed in exact diagonalization, particularly at early times, that we contrast our data with.
- [108] T. Brydges, A. Elben, P. Jurcevic, B. Vermersch, C. Maier, B. P. Lanyon, P. Zoller, R. Blatt, and C. F. Roos, Probing rényi entanglement entropy via randomized measurements, Science **364**, 260 (2019).
- [109] C. Dankert, R. Cleve, J. Emerson, and E. Livine, Exact and approximate unitary 2-designs and their application to fidelity estimation, Physical Review A 80, 012304 (2009).
- [110] If this iteration were continued, eventually one would end up with an exponentially large set of all possible operators. Our approach is based on a purely heuristic observation that, in practice, only polynomial many (quasi)local operators are required to accurately reproduce the state $\rho_A(t)$ with exponential accuracy [74, 75].
- [111] https://www.mathworks.com/help/gads/ globalsearch.html.
- [112] We numerically checked that the results remain valid for other randomly chosen states in this basis.
- [113] J. Bringewatt, J. Kunjummen, and N. Mueller, Randomized measurement protocols for lattice gauge theories, Quantum 8, 1300 (2024).
- [114] S. N. Hearth, M. O. Flynn, A. Chandran, and C. R. Laumann, Unitary k-designs from random number-conserving quantum circuits, arXiv preprint arXiv:2306.01035 (2023).
- [115] S. N. Hearth, M. O. Flynn, A. Chandran, and C. R. Laumann, Efficient local classical shadow tomography with number conservation, arXiv preprint arXiv:2311.09291 (2023).
- [116] M. K. Joshi, C. Kokail, R. van Bijnen, F. Kranzl, T. V. Zache, R. Blatt, C. F. Roos, and P. Zoller, Exploring large-scale entanglement in quantum simulation, Nature , 1 (2023).
- [117] J. Berges, S. Borsányi, and C. Wetterich, Prethermalization, Physical review letters 93, 142002 (2004).
- [118] M. Gring, M. Kuhnert, T. Langen, T. Kitagawa, B. Rauer, M. Schreitl, I. Mazets, D. A. Smith, E. Demler, and J. Schmiedmayer, Relaxation and prethermalization in an isolated quantum system, Science **337**, 1318 (2012).
- [119] T. Mori, T. N. Ikeda, E. Kaminishi, and M. Ueda, Thermalization and prethermalization in isolated quantum systems: a theoretical overview, Journal of Physics B: Atomic, Molecular and Optical Physics 51, 112001 (2018).
- [120] H. B. Callen and T. A. Welton, Irreversibility and generalized noise, Physical Review 83, 34 (1951).
- [121] R. Kubo, The fluctuation-dissipation theorem, Reports on progress in physics 29, 255 (1966).

- [122] A. P. Orioli and J. Berges, Breaking the fluctuationdissipation relation by universal transport processes, Physical review letters 122, 150401 (2019).
- [123] A. Schuckert and M. Knap, Probing eigenstate thermalization in quantum simulators via fluctuationdissipation relations, Physical Review Research 2, 043315 (2020).
- [124] C. Murthy, A. Babakhani, F. Iniguez, M. Srednicki, and N. Y. Halpern, Non-abelian eigenstate thermalization hypothesis, Physical Review Letters 130, 140402 (2023).
- [125] X. Yao, SU(2) gauge theory in 2+1 dimensions on a plaquette chain obeys the eigenstate thermalization hypothesis, Phys. Rev. D 108, L031504 (2023), arXiv:2303.14264 [hep-lat].
- [126] L. Ebner, A. Schäfer, C. Seidl, B. Müller, and X. Yao, Eigenstate thermalization in (2+ 1)-dimensional su (2) lattice gauge theory, Physical Review D 109, 014504 (2024).
- [127] Z. Davoudi, C. Jarzynski, N. Mueller, G. Oruganti, C. Powers, and N. Y. Halpern, Quantum thermodynamics of nonequilibrium processes in lattice gauge theories, arXiv preprint arXiv:2404.02965 (2024).
- [128] S. Ghosh, R. M. Soni, and S. P. Trivedi, On the entanglement entropy for gauge theories, Journal of High Energy Physics **2015**, 1 (2015).
- [129] K. Van Acoleyen, N. Bultinck, J. Haegeman, M. Marien, V. B. Scholz, and F. Verstraete, Entanglement of distillation for lattice gauge theories, Physical review letters 117, 131602 (2016).
- [130] MATLAB, version 23.2.0.2485118 (R2023a) (The MathWorks Inc., Natick, Massachusetts, 2023).
- [131] MathWorks, (2024). Fmincon documentation. Retrieved May 13, 2024 from https://www.mathworks. com/help/optim/ug/fmincon.html,.
- [132] MathWorks, (2024). Globalsearch documentation. Retrieved May 13, 2024 from https://www.mathworks. com/help/gads/globalsearch.html,.

Appendix A: Circuit Representation for State Preparation, Evolution, and Measurement

In this Appendix, we provide circuits employed to experimentally realize state preparation, time evolution, and randomized measurement in the Z_2 LGT in (2+1)D within the dual Isnig formulation.

We label qubits with indices $q = 0, \ldots, L - 1$, with L = 12. Qubit q = 0 corresponds to the spin at $\ell_{\bar{A}A}$ positioned at the interface between subsystem A and its complement, identifiable by an orange circle in Fig. 2(c). Qubits $q = 1, \ldots, 4$ represent the succeeding dual variables arranged counterclockwise at the center of plaquettes denoted by purple circles in Fig. 2(c). Following this sequence, qubit q = 5 resides at the opposite boundary marked as $\ell_{A\bar{A}}$, and so forth.

Working in the electric eigenbasis and starting from an all-up spin state, we first perform single-qubit σ^x bitflip operations to select a randomly chosen product state in the σ^z basis that is consistent with Gauss laws, i.e., $G_i |\psi(0)\rangle = |\psi(0)\rangle$ for all *j*. Recall that in the dual formulation, there are only two Gauss-law operators each located at one of the subsystem's boundaries: $G_{\bar{A}A}^{\text{dual}} = \mu_{11}^z \sigma_0^z \mu_1^z$ and $G_{A\bar{A}}^{\text{dual}} = \mu_4^z \sigma_5^z \mu_6^z$. Importantly, any spin configuration in either bulk, A and \bar{A} , is physical, as in the dual formulation, only gauge-invariant degrees of freedom are kept in the bulk.

For time evolution, we utilize a first-order Trotter scheme with time-evolution operator

$$U(t) \equiv e^{-itH^{\text{dual}}} \equiv \prod_{\delta t} U(\delta t).$$
 (A1)

We take δt to be a variable time step while keeping the Trotter depth fixed. In the simulations conducted in this work, $t/\delta t = 4$, resulting in states and observables closely resembling the exact time-evolved states at early times. However, for late times, Trotter effects become more pronounced.

 $U(\delta t)$ in Eq. (A1) can be written as $U(\delta t) \equiv \prod_{a \in X, Z, XX, ZZ} U_a$ with $U_a \equiv e^{-i\delta t H_a^{\text{dual}}}$; H_a^{dual} are the respective 1- and 2-local operators of the dual Hamiltonian in Eq. (6), sorted in X and Z operations, as well as single- and two-qubit entangling gates. While XX, X, and Z are native operations, ZZ entangling operations are realized via basis transformation and usage of the native MS gate,

$$R_{ij}^{zz}(\alpha) \equiv R_i^y\left(-\frac{\pi}{2}\right)R_j^y\left(-\frac{\pi}{2}\right)R_{ij}^{xx}(\alpha)R_i^y\left(\frac{\pi}{2}\right)R_j^y\left(\frac{\pi}{2}\right).$$
(A2)

Here, $R_i^{x/y/z}(\alpha) \equiv e^{-i\frac{\alpha}{2}\sigma_i^{x/y/z}}$ and $R_{ij}^{xx}(\alpha) \equiv e^{-i\alpha\sigma_i^x\sigma_j^x}$. The experimental errors of our MS gates increase with the absolute value of the gate angle. To minimize the gate error, all MS gates are optimized as follows. First, all angles are mapped to the regime $\alpha \in [-\pi, \pi]$. Within this range, we make the following substitution

$$R_{ij}^{xx}(\alpha) \rightarrow \begin{cases} R_{ij}^{xx}(\alpha) & \text{if } |\alpha| \leq \frac{\pi}{4} \\ R_{ij}^{xx}(\alpha + \frac{\pi}{2})R_{i}^{x}(\pi)R_{j}^{x}(\pi) & \text{if } |\alpha| \in (\frac{\pi}{4}, \frac{3\pi}{4}] \\ & \& \alpha < 0 \\ R_{ij}^{xx}(\alpha - \frac{\pi}{2})R_{i}^{x}(\pi)R_{j}^{x}(\pi) & \text{if } |\alpha| \in (\frac{\pi}{4}, \frac{3\pi}{4}] \\ & \& \alpha > 0 \\ R_{ij}^{xx}(\alpha + \pi) & \text{if } \alpha < -\frac{3\pi}{4} \\ R_{ij}^{xx}(\alpha - \pi) & \text{if } \alpha > \frac{3\pi}{4} \end{cases}$$
(A3)

so that all MS operations are restricted to the range $|\alpha| \leq \frac{\pi}{4}$.

An example study of the initial-state preparation, time evolution, and measurement in a fixed (Bloch x or z) basis is shown in Fig. 3 (b) and (c) of the main text, where we presented measurement of several gauge-invariant and non-gauge-invariant one- and two-qubit observables. In addition, we present the measurement of several other non-gauge-invariant observables in Fig. 7. These observables are expected to be zero, but remain non-vanishing because of finite measurements and device errors. A main




FIG. 6. Overview of the circuits employed in the simulations of this work. (a) Initial-state preparation, which involves setting a Gauss-law-respecting electric eigenstate in the Bloch z basis. The shown circuit corresponds to one such initial state. (b) Trotterized time evolution, consisting of magnetic interactions (in the x basis) comprised of single- and two-qubit rotations in the dual formulation. The electric part of the Hamiltonian evolution is diagonal consisting of Z and ZZ rotations. All MS gates are implemented according to Eq. (A3) depending on the employed time-evolution step δt . (c) Randomized-measurement circuits. $\gamma_i^{1,2,3}$ at qubit $i = 0, 1, \dots, 5$ are drawn from a circular unitary ensemble. The qubits 0 and 5 marking the two boundaries of the periodic lattice are drawn in orange. The qubit index of the rotation gates are dropped to reduce clutter and can be deduced from the qubit(s) they act on.

source of errors are likely coherent errors, related to the over- and under-rotation of single- and two-qubit gates. While the time-evolution circuit only contains gauge-invariant operations for any gate angle employed, the final single-qubit rotations $R_i^y(-\pi/2)$ and $R_i^x(\pi/2)$ (that transform all qubits *i* from the Bloch *z* basis into the Bloch *x* and *y* bases, respectively) can introduce gauge-invariance violation if their rotation angle is set inaccurately. Additionally, errors stemming from initial-state preparation and readout processes can contribute to this violation.

Appendix B: Finite-size Dependence, Trotterization, and Late-time Behavior

In this Appendix, we perform various tests aiming to extend our study toward larger systems, and to provide an outlook on the expected late-time behavior of nonequilibrium states, features that are inaccessible in current quantum computers. We present a detailed *classical computation* of a fairly large system (L = 22, $L_A = 8$ with coupling g = 1), evolving the state continuously, and computing its entanglement structure using exact diagonalization, hence avoiding a Trotterization error. Such an analysis is possible for the model under consideration, owing to its relative simplicity. This simplicity enables us to achieve results that are fairly insensitive to finite-size effects, even with moderately large systems. A similar classical computation will be impossible for more involved models or in higher dimensions.

We compute the same entanglement-related observables indicative of quantum chaos as in the main text, starting from several randomly drawn initial product states. The EGRD, i.e., the (normalized) distribution of the gap ratios P(r) with r defined after Eq. (8), is computed similarly as in the main text: we consider ten randomly chosen initial states, and combine the gap ratios that are separately computed for each symmetry sector of the respective reduced states and for all initial states. Additionally, we combine the distribution of gap ratios in three time regimes (I), (II), and (III). The result is shown Fig. 8(a). Further, the mean of the gap ratio, combining initial states and symmetry sectors but not times, $\langle r \rangle$, is shown in Fig. 8(b) as a function of scaled time *qt*. The distribution in Fig. 8(b) is peaked near zero first, closely resembling a Poisson distribution. In this regime, the rank $\mathcal{R}_A \ll d_A$, where d_A is the Hilbert-space dimension of the subsystem. The effective rank \mathcal{R}_A is defined after Eq. (9) and plotted for the present example in Fig. 8(c)starting from a product state with $\mathcal{R}_A = 1$ at t = 0. The largest ξ_{λ} values in this regime correspond to extremely small probabilities $p_{\lambda} = \exp(-\xi_{\lambda})$, close to or at the level of machine precision. A regularization of the smallest p_{λ} is required: we manually cut off probabilities below 10^{-15} before computing \mathcal{R}_A , varying this limit by two orders of magnitude in each direction to provide the blue bands in panel Fig. 8(c), then using only the \mathcal{R}_A lowest levels in the analysis of the EGRD and the ESFF. Gray bands represent the variance with respect to the randomly cho-




FIG. 7. Estimation of gauge-invariance violation. Expectation values of several single- and two-qubit non-gauge-invariant operators that are expected to be zero at all times. For better visibility, we plot the different data sets slightly shifted in time gt.

sen initial states. In regime (II), $\langle r \rangle$ quickly jumps to a value associated with a Gaussian Orthogonal Ensemble (GOE), until continuing, albeit at much slower pace, towards a Gaussian Unitary Ensemble (GUE) indicating quantum chaotic behavior. At the end of this stage, $\mathcal{R}_A = d_A$ is maximal, and all probabilities p_λ are well above machine precision, eliminating the need for any regularization. Finally, regime (III) sees the saturation of the gap ratio to GUE level statistics (up to minimal finite lattice-size effects). The beginning and end of the time regimes (I-III) depend on the subsystem size and are chosen as follows, (I): $0 \leq gt < 1.8$, (II): $1.8 \leq gt < 5.0$, and (III): $5.0 \leq gt < 10.0$.

An analogous picture is evident in the time evolution of the ESFF defined in Eq. (9), and shown in Figs. 8 (d) and (e). Panel (e) shows the ESFF for various times $0 \leq qt \leq 10$, starting from a flat distribution at earliest time and showing a plateau-ramp structure at late times. Similarly as in the main text, the ESFF is computed separately for every symmetry sector of the EH to avoid contamination from uncorrelated levels in different sectors, and then averaged over these sectors and over the randomly chosen initial states. Panel (d) shows the time average over regime (III). In this regime, the ramp-plateau is evident. A gray band marks the statistical deviation from the initial-state and symmetry-sector averages. We fit the ramp to the form $\sim \theta^{0.6\pm0.1}$ which is consistent with the experimental data shown in the main text for a much smaller system. The fit error is determined by varying the fit range.

Because the late-time behavior, i.e., $gt \gg 1$, is inaccessible to current Trotter-based digital quantum simulation, we extend our analysis towards this regime, aiming to elucidate the potential outcomes that future quantum simulators might uncover. A similar analysis has been performed previously in Ref. [41]. At late times, it is expected that the entanglement entropy of a subsystem becomes equal to the thermal entropy contained in that subsystem, corresponding to a global Gibbs state with a temperature corresponding to the average energy of the initial state. In Fig. 8 (f), we plot the von Neumann entanglement entropy, for a variety of randomly chosen initial states. The individual curves are color coded relative to their initial average energy density $\sim E_0 = \langle \psi(0) | H | \psi(0) \rangle$, normalized relative to the energy bandwith ΔE , i.e., the difference between the highest and lowest eigenvalues of the EH. Displayed are only initial states whose energies lie within the interquartile range, representing the central 50% energy spectrum of states. States highlighted in yellow denote the highest energies, while those in blue represent the lowest. We were not able to extend this study to even later times to observe the expected saturation of the von Neumann entanglement entropy, owing to the fact that eventually finitesize effects become large. These results demonstrate that entanglement-entropy saturation is not a practical measure of thermalization in present experiments, demanding long evolution times and exhibiting lack of initial-state insensitivity (i.e., universality).

Shown in Fig. 8(g) is a separation of the von Neumann entanglement entropy, S_{vN} , into components, $S_{vN,s}$, related to the symmetry structure of $\rho_A = \bigoplus_s \rho_{A,s}$,

$$S_{vN} = -\sum_{s} p_s \log(p_s) + \sum_{s} p_s S_{vN,s}, \qquad (B1)$$

where the first term is the symmetry component, and the second is the distillable entanglement [128, 129]. Here, $p_s \equiv \text{Tr}[\rho_{A,s}] \leq 1$ with $\sum_s p_s = 1$, and $S_{vN,s} \equiv -\text{Tr}[\bar{\rho}_{A,s}\log(\bar{\rho}_{A,s})]$ is the sector-wise entanglement entropy with $\bar{\rho}_{A,s} \equiv \rho_{A,s}/p_s$. Dashed lines in the plot denote the distillable component, while dotted lines denote the symmetry part. It is evident that both components saturate on different time scales: the symmetry component saturates fairly quickly (albeit still later than the build-up of level repulsion) to its maximal value $\log(1/p_s)$ where $p_s = 1/4$ corresponds to equal mixing of the four symmetry sectors of ρ_A . In contrast, the distillable components dominates the late-time behavior of the von Neumann entropy.

While experimental constraints prevent us from accessing late times, Fig. 9 illustrates the von Neumann entanglement entropy derived from our experimental data, utilizing the same dataset as in the main text. In this figure, exact classically computed results are denoted by black lines, while cyan crosses represent the optimal BW-inspired parameterization assuming infinite measurements, and orange circles are the experimental data points. The right-hand side of the figure displays




FIG. 8. Theoretical analysis for L = 22 and $L_A = 10$ plaquettes. (a) Entanglement-spectrum gap-ratio distribution, combining the gap ratios of ten randomly selected initial states and across all symmetry sectors of the reduced state. The distributions are additionally combined in time over all data in each of regimes (I), (II), and (III), and binned over intervals $\Delta r = 1/12$. (b) Mean of the gap ratio distribution, combining initial states and symmetry sectors. (c) Rank of the reduced density matrix as a function of scaled time. (d) Spectral form factor of the EH, averaged over symmetry sectors and over 10 randomly selected initial product states, averaged over time range (III) ($5 < gt \le 10$). (e) Time dependence of the entanglement spectral form factor. (f) The von Neumann entanglement entropy as a function of scaled time, for 10 randomly chosen initial product states. The color encoding represents the energy of the initial state relative to the energy bandwidth ΔE of the physical Hamiltonian. (g) Distillable versus symmetry components of the von Neumann entanglement entropy.




FIG. 9. Entanglement entropy from experiment. Shown is the von Neumann entanglement entropy in the left panel, comparing exact (black lines), infinite-measurement results using a BW-inspired parametrization (cyan crosses), and quantum-computed data (orange circles). The top and bottom side panels show the symmetry and distillable components of the von Neumann entanglement entropy, respectively.

two panels: the top panel depicts the symmetry compo-

nent, while the bottom panel depicts the distillable component. Notably, the observed behavior closely mirrors the (classically computed) findings of the much larger system in Fig. 8(g). Specifically, the symmetry component saturates at its maximum value of log(4), while the distillable entanglement demonstrates continued growth. Generally, the entanglement measured in our experiment via BW-inspired tomography overshoots the exact result. We attribute this discrepancy primarily to over- or underrotations within the single-qubit random-bases changes.

Appendix C: Entanglement-Hamiltonian Tomography

In this section, we outline the randomized-measurement and analysis strategy behind extracting approximate EHs from experiment. We also systematically investigate the performance of the approach with regard to the number of bases sampled, the number of shots performed, and the influence of device errors.

Our tomography protocol consists of two parts: randomizing the state via single-qubit random unitaries drawn from a circular unitary ensemble, followed by classical post-analysis of the extracted measurements to determine an EH that best describes all observables. We




FIG. 10. Operator set for the entanglement-Hamiltonian ansatz. Representation of the operator set in the entanglement-Hamiltonian ansatz used in our study, encompassing operators positioned inside the subsystem. All operators are Hermitian. Electric fields along the x direction are interdependent (since $V_y = 1$), thus not treated as independent operators. For the specific subsystem with $N_A = 4$, our ansatz incorporates a total of 73 distinct operators. While this is close to the subsystem Hilbert-space dimension, for a larger system, the number of operators scales linearly with the degrees of freedom within the subsystem.




FIG. 11. The operator related to the symmetry sectors of the ansatz state. Because of the (remaining) Gauss laws, some operators outside of the subsystem are identical to operators within its complement. The operator shown, and a similar operator placed on the other boundary, commutes with all other operators in the BW-inspired ansatz shown in Fig. 10. Therefore, their (common) eigenspace are the symmetry sectors of the reduced density matrix.

use single-qubit random circuits, $\mathcal{U} \equiv \bigotimes_i u_i$ where u_i is the following single-qubit unitary

and for each qubit, the angles γ_i^1 , γ_i^2 , γ_i^3 are drawn according to a circular unitary ensemble (an overall phase is ignored) following Refs. [77, 108]. A drawback of

this approach is that it does not maintain the symmetry structure of ρ_A , as it randomizes over the whole Hilbert space instead of each symmetry block of ρ_A . Such a symmetry-ignorant randomization mixes different symmetry sectors, resulting in an outcome that is inaccessible to any physical time-evolved quantum state. We choose it, nonetheless, to avoid the larger circuit depth associated with a symmetry-conscious randomization such as that proposed in Ref. [86]. We note that a symmetryconscious scheme would significantly reduce the measurement cost, as the sampling cost would only scale as the size of the symmetry block instead of the Hilbert-space size. It would thus simplify the classical optimization significantly. However, because the single-qubit scheme is tomographically complete, one can still reconstruct, approximately, the symmetry structure from the data.

The ansatz for the EH is a sum of local operators \mathcal{O}_i ,

$$H_A(t; \{\beta_i\}) \equiv \sum_i \beta_i(t) \mathcal{O}_i , \qquad (C1)$$

which are either the physical-Hamiltonian terms or are chosen iteratively from commutators of those terms. β_i are real parameters to be constrained from data. While the BW theorem hints at an optimal choice of operators \mathcal{O}_i for ground states, our ansatz is heuristic, as a systematic ansatz for non-equilibrium states is not known. Our




FIG. 12. *Gap-ratio averages across initial states*. Comparison of the average gap ratio obtained in experiment (yellow circles) versus using an ideal emulator (blue squares), shown individually for six randomly-drawn initial electric eigenstates compatible with the Gauss laws. We show the spin configuration of the initial state in each panel, which is randomly drawn. Error bars and error bands represent the spread over symmetry sectors. Dotted blue and dashed red lines denote the values associated with the Poisson distribution and GUE, respectively.

criteria for the operators are that i) they are local, involving operators with support on at most two neighboring plaquettes, ii) they are compatible with the symmetries of the reduced state ρ_A (which will be discussed below), and iii) they are independent, meaning they cannot be transformed into each other through Gauss laws when acting on a physical state (although they are certainly not independent in the algebraic sense).

The operators we use are pictorially represented in Figs. 10 plus all operators that are generated from those depicted upon translation. Here for the sake of generality, we represent them in terms of Z_2 variables, albeit, in practice, they are represented in the dual formulation in our algorithm. Figure 11 shows a special operator; this operator (and related examples) relates operators within the subsystem to operators in the complement via Gauss laws. Further, it commutes with ρ_A , and is hence connected to the symmetry blocks of ρ_A . All terms in Figs. 10 are obtained by considering the physical-Hamiltonian operators, stopping at two recursions (i.e., commutators of commutators). Other selection strategies are also feasible.

Given the ansatz and following Ref. [75], we then min-




FIG. 13. Entanglement spectrum from tomography. (a) Lower parts of the entanglement spectrum for times $gt = 0, 0.34, 1.7, \text{ and } 2.38, \text{ comparing exact data (black lines), infinite-measurement results (cyan crosses), emulator data (blue squares), and quantum-computed data (orange circles), for one given initial state <math>|\Psi_0\rangle = |\downarrow\downarrow\downarrow\uparrow\uparrow\downarrow\downarrow\uparrow\uparrow\uparrow\downarrow\uparrow\rangle$. (b) Entanglement spectrum separated into the four symmetry sectors for the same data set at gt=1.02. The inset shows a close-up of the lower part of the spectrum for all symmetry sectors.

imize the functional

$$\left\langle \sum_{b} \left[P_{\mathcal{U}}(b) - \operatorname{Tr} \left[\mathcal{U}^{\dagger} | b \rangle \langle b | \mathcal{U} \rho_{A}(t; \{\beta_{i}\}) \right] \right]^{2} \right\rangle_{\mathcal{U}}$$
(C2)

where $P_{\mathcal{U}}(b)$ is the probability to measure a bitstring b in the basis determined by \mathcal{U} and

$$\rho_A(t; \{\beta_i\}) \equiv \frac{e^{-H_A(t; \{\beta_i\})}}{\text{Tr}[e^{-H_A(t; \{\beta_i\})}]}.$$
 (C3)

is the normalized reduced density matrix parameterized by H_A . Here, $\langle \cdot \rangle_{\mathcal{U}}$ is the average over random circuits. Note that optimizing Eq. (C2) effectively weighs more favorably the largest Schmidt eigenvalues of ρ_A because they, on average, contribute the most to any random observable. Because of this, the optimization more accurately reproduces the low-energy part of H_A . The




FIG. 14. Estimation of basis dependence (emulator data). (a) Estimation of dependence on the number of random bases $N_{\mathcal{U}}$, illustrated for fixed $N_{\rm shots} = 750$ per basis for one representative initial state. Error bars and error bands represent the spread over symmetry sectors. (b) Example of a reconstructed entanglement spectrum, varying the number of random bases for a fixed number of shots $N_{\rm shots} = 750$. The inset shows a close-up of the lower part of the spectrum for all symmetry sectors.

optimization is performed using MATLAB's [130] nonlinear FMINCON optimization package with the 'sqp' algorithm [131]. Convergence and uniqueness of the obtained minimum have been cross checked for several data sets using MATLAB's GLOBALSEARCH routine with default parameters [132]. All EH couplings are confined to the range $\beta_i \in [-50, 50]$. We check explicitly that the routine does not come close to the boundary of the parameter regime. The EH that is obtained in this way is projected into symmetry sectors, which we then separately analyze. For this projection, the symmetry sector can be read off directly from the row and column numbers of the EH when they are interpreted as binary. The corresponding subblock $H_{\rm E,s}$ is then selected for further analysis, where s labels one symmetry sector. For the EGRD analysis, we analyze the gap-ratio distribution separately for every sector, then combine the distributions and re-normalize the total distribution. For the ESFF analysis, we average




FIG. 15. Estimation of shot-noise dependence (emulator data). (a) Estimation of dependence on the number of shots, illustrated for fixed number of random bases $N_U = 24$ for one representative initial state. Error bars and error bands represent the spread over symmetry sectors. (b) Example of a reconstructed entanglement spectrum, varying the number of shots for a fixed $N_U = 24$. Red hexagons represent the infinite-shot limit. The inset shows a close-up of the lower part of the spectrum for all symmetry sectors, a black arrow indicates where emulator data (in the infinite-shot limit) diverges from the ideal BW results.

over the sectors. Finally, EGRD and ESFF shown in the main text and appendices involve a combination/average over initial states. The error band and the error bars in the plots are the standard deviation for the symmetry-sector and initial-state averages.

An example of the eigenvalue spectrum of the EH, comparing the exact values with those obtained from our procedure, is shown in Fig. 13. Panel (a) shows the reconstructed entanglement spectrum for several times, comparing exact data (black lines), infinite-measurement results (cyan crosses), emulator data (blue squares), and quantum-computed data (orange circles), for one given initial state. Panel (b) shows the symmetry-resolved spectrum for gt=1.02. Here, s are the symmetry sectors of ρ_A , see the discussion at the end of Sec. II in the main text. While our analysis effectively describes the low-

lying part of the entanglement spectrum, it fails to quantitatively reproduce the higher part associated with very small probabilities. Additionally, the reconstructed spectrum appears more mixed (more entangled) compared to e.g., the emulator data. This discrepancy primarily is due to finite-measurement statistics and device errors. Comparison with the infinite-(ideal-)measurement results also reveals that the BW-inspired parameterization falls short in capturing even quantitatively higherlying components of the spectrum. It is noteworthy that the limitations of the parametrization become more pronounced at later times where the Trotter-step size is large and even the low-lying part deviates. This is because the effective Trotter Hamiltonian at late times deviates from the known target Hamiltonian from which the BWinspired ansatz is derived.

In Fig. 14, we study the accuracy of our tomography scheme concerning the number of random bases. Each basis is probed with $N_{\rm shots} = 750$ samples, employing emulated data to eliminate the effect of device errors. Applying $N_{\mathcal{U}} = 4$ to $N_{\mathcal{U}} = 128$ random bases shows convergence in the lower-lying range of the entanglement spectrum relatively fast. However, the higher-lying part is not reconstructed even with a large sample size. The top panel displays the average EGRD over time, showing consistent behavior across the sample range. Error bars indicate the spread from combining initial states and symmetry sectors, which decrease with increasing $N_{\mathcal{U}}$. Concretely the error bars are obtained by computing the mean of the gap ratios for every sector and initial states separately, and then computing the standard deviation.

Finally in Fig. 15, we investigate the dependence on the number of measurement of probabilities $P_{\mathcal{U}}(s)$ in each basis (i.e., the number of shots), maintaining a constant number of measurement bases, $N_U = 24$. Once more, convergence is evident in the lower segment of the ES in the lower panel, while the higher portion remains beyond reach even with infinite shots (red octagons). The top panel illustrates the EGRD, indicating consistency, within error bars, with the optimal BW-inspired ansatz result (cyan band).

In principle, given that the scheme is tomographically complete, we anticipate the ability to precisely reconstruct the entire ES with exponential resources. However in practice, we were not able to do so because of the significant bias of the cost function Eq. (C2) towards the low-energy portion of the entanglement spectrum. This bias places considerable strain on the numerical minimization routine, pushing it beyond its numericalaccuracy threshold. Despite this, its accuracy is satisfactory for analyzing experimental data, and we abstained from further optimization attempts. Exploring advanced optimization routines, including machine-learning techniques [84], holds promise for EH tomography in future studies.