

Reverberation mapping of high-mass and high-redshift quasars using gravitational time delays

MIRIAM GOLUBCHIK,¹ CHARLES L. STEINHARDT,^{2,3} ADI ZITRIN,¹ ASHISH K. MEENA,¹ LUKAS J. FURTAJ,¹
DORON CHELOUCHE,^{4,5} AND SHAI KASPI⁶

¹*Department of Physics, Ben-Gurion University of the Negev, P.O. Box 653, Be'er-Sheva 84105, Israel*

²*Cosmic Dawn Center (DAWN), Denmark*

³*Niels Bohr Institute, University of Copenhagen, Lyngbyvej 2, Copenhagen Ø 2100, Denmark*

⁴*Department of Physics, Faculty of Natural Sciences, University of Haifa, Haifa 3498838, Israel*

⁵*Haifa Research Center for Theoretical Physics and Astrophysics, University of Haifa, Haifa 3498838, Israel*

⁶*School of Physics & Astronomy and the Wise Observatory, The Raymond and Beverly Sackler Faculty of Exact Sciences Tel-Aviv University, Israel*

(Received; Revised; Accepted)

ABSTRACT

Mass estimates of black holes (BHs) in the centers of Active Galactic Nuclei (AGN) often rely on the radius-luminosity relation. However, this relation, usually probed by reverberation mapping (RM), is poorly constrained in the high-luminosity and high-redshift ends due to the very long expected lag times. Multiply imaged AGN may thus offer a unique opportunity to explore the radius-luminosity relation at these ends. In addition to comprising several magnified images which enable a more efficient light-curve sampling, the time delay between multiple images of strongly lensed quasars can also aid in making such RM measurements feasible on reasonable timescales: If the time delay is, for example, of the order of the expected time lag, changes in the emission lines in the leading image can be observed around the same time as the changes in the continuum in the trailing image. In this work we probe the typical time-delay distribution in galaxy-cluster lenses and estimate the number of both high-mass ($\sim 10^9 - 10^{10} M_{\odot}$), and high-redshift ($z \gtrsim 4 - 12$) quasars that are expected to be strongly lensed by clusters. We find that less than one, very massive (and luminous, $L_{UV} > 10^{46.5} \text{ erg s}^{-1}$) multiply imaged quasar is expected across the sky down to 30 AB mag. Nonetheless, up to several dozen thousand $M_{BH} \sim 10^6 - 10^8 M_{\odot}$ broad-line AGN at $z > 4$ should be multiply imaged by galaxy clusters and detectable with JWST, hundreds with *Euclid* and several thousands with the *Roman* Space Telescope, across the whole sky. These could supply an important calibration for the BH mass scaling in the early Universe.

Keywords: Quasars – galaxies: clusters: general – gravitational lensing: strong

1. INTRODUCTION

Super massive black holes (SMBHs) are found in the centers of most galaxies. The masses of those SMBHs are of much interest as their evolution is not well understood, and since they are believed to play a major role in their host galaxy evolution. In fact, tight relations seem to hold between the mass of an SMBH and the properties of its host galaxy such as the luminosity or the velocity dispersion of the bulge (e.g., Ferrarese &

Merritt 2000; Kormendy & Ho 2013; Reines & Volonteri 2015; Ding et al. 2020). Measuring SMBH masses as a function of redshift can also shed light onto black hole (BH) formation and the physics of accretion processes (e.g., Wang et al. 2021; Volonteri 2012), and their mass evolution through cosmological time might help constrain different cosmological models (e.g., Fanidakis et al. 2012; Habouzit et al. 2022).

SMBHs are believed to be the central engines of Active Galactic Nuclei (AGNs). The rapid accretion onto the BH results in a high luminosity across a broad range of wavelengths and unique spectral features from which one can estimate the virial mass of the BH. Using a single epoch spectrum, one can resolve the velocity of the

material in the broad line region (BLR) surrounding the BH and, through further assumptions of virial motion, obtain an estimate for the BH mass (e.g., [Greene & Ho 2005](#)). However, this technique strongly relies on the ability to determine the distance R from the central engine to the BLR (often referred to as the size of the BLR) which is usually obtained through empirical scaling relations with the AGN’s continuum luminosity L (often referred to as an R - L relation).

R - L relations can be empirically derived using Reverberation Mapping (RM). The RM technique aims to measure the distance to the BLR in AGNs by measuring the time lag between variations in the continuum flux from the accretion disk, and the response of the emission lines in the BLR to those variations, assuming that this response propagates with the speed of light ([Bahcall et al. 1972](#); [Blandford & McKee 1982](#); [Peterson 1993](#); [Netzer & Peterson 1997](#)). Almost a 100 AGNs were studied using RM to determine the sizes of their BLRs, resulting in an empirical relation connecting the continuum luminosity to the BLR size for different emission lines ([Kaspi et al. 2000, 2005](#); [Bentz et al. 2009, 2013](#)). Indeed, those R - L relations are widely used for virial mass estimates of SMBHs in the centers of AGN, varying in luminosity and redshift. However, because those relations were mostly obtained using (relatively) low redshift and lower-luminosity AGN, RM studies of high-luminosity and high-redshift quasars are needed in order to extend and calibrate these relations to those extreme ends where the relation is not well established and where the physical picture may be different (for example, the BLR might not be necessarily virialized (e.g. [King 2024](#); [Lupi et al. 2024](#))).

Unfortunately, RM of high-luminosity and high-redshift quasars faces several difficulties. High-luminosity quasars are presumed to have large BLR sizes, resulting in long time lags between continuum and emission-line variations. Similarly, high-redshift quasars and AGN (e.g., [Wang et al. 2021](#); [Furtak et al. 2024](#); [Larson et al. 2023](#); [Goulding et al. 2023](#)) will also show longer observed time lags compared to their lower-redshift counterparts, due to cosmological time dilation. As a result, observations for several decades are needed for those RM campaigns ([Kaspi 2018, 2021](#)). Nevertheless, several attempts have been recently made to conduct RM for those types of AGN. In one study, [Kaspi et al. \(2021\)](#) observed photometrically 11 quasars at redshifts $\sim 2 < z < 3.4$ and luminosities $\sim 10^{46.9} < \lambda L_{\lambda}(1350\text{\AA}) < 10^{48.0} \text{ erg s}^{-1}$ for almost twenty years while six of them were monitored spectrophotometrically for 13 years. Their study manages to significantly measure time lags of the C IV $\lambda\lambda 1548, 1551\text{\AA}$ doublet in

three objects and a C III] $\lambda\lambda 1907, 1909\text{\AA}$ time lag in one object. Scaling the BLR size with the UV continuum luminosity they conclude that their objects lie on the previously measured relation. [Shen et al. \(2023\)](#) performed RM using 11-year photometric and a 7-year spectroscopic data for 849 broad-line quasars. They report 339 significantly measured time lag detections for the different emission lines. Although they claim that on average the R - L relations follow existing ones, they report an intrinsic scatter of ~ 0.3 dex in the lags for the H β line at 4861 \AA and the Mg II] $\lambda\lambda 2796, 2803\text{\AA}$ doublet, whereas the C IV R - L relation has an even more substantial scatter of ~ 0.5 dex. It is thus becoming important to understand whether these large scatters stem from intrinsic physical processes at high luminosities or due to the fact that there is often a difficulty to significantly measure time lags for such objects given the monitoring times needed.

In this work, we discuss the possibility to probe the R - L relation for the high-luminosity or high-redshift regimes using multiply imaged quasars. If the strong-lensing (SL) time delays between different multiple images of a lensed quasar are of the same order as the time lags, the expected changes in the emission lines can be observed in a leading multiple image around the same time as the changes in the continuum from a trailing multiple image, so that RM of these types of objects can be done with a reasonable and shorter monitoring time-span than for similar, un-lensed quasars (nominally shorter by one lag-time, but as we also discuss here in practice the gain will be larger, as a lensed quasar will be lensed into several magnified images, supplying a more efficient sampling of the light curve).

About a handful of quasars and AGN, mostly around cosmic noon, are known to be multiply imaged by galaxy clusters (e.g., [Inada et al. 2003](#); [Oguri et al. 2013](#); [Sharon et al. 2017](#); [Acebron et al. 2022](#); [Furtak et al. 2023a](#); [Napier et al. 2023a](#)), whereas some have also been used for cosmography (e.g., [Napier et al. 2023b](#)). Notably, a successful RM of a strongly lensed quasar has been recently done by [Williams et al. \(2021\)](#) where they exploit the time delays between the multiple images of the strongly lensed quasar SDSS J2222+2745 to construct a detailed light curve spending over 6 years in the observed frame with 4.5 years of observations with high cadence. They significantly detect a time lag of $\tau_{cen} = 36.5^{+2.9}_{-3.9}$ rest-frame days for the C IV emission line, whereas the time delays from the two other images they use are of ~ 40 and ~ 700 days. The quasar SDSS J2222+2745 ranged in continuum luminosity with the average value of $\log_{10}[\lambda L_{\lambda}(1350\text{\AA})/\text{erg s}^{-1}] = 44.66 \pm 0.18$. In this illustrative case the time lag was short enough so that

the gain from image multiplicity was mainly manifested in additional data points on the light curve, thus importantly allowing to shorten the overall monitoring time. Here, we wish to focus on objects at the high luminosity end, namely continuum UV luminosities $L_{UV} > 10^{46.5}$ erg s $^{-1}$ (BH masses of $\log_{10}M_{BH} \gtrsim 9$), or at high redshifts (i.e., $z \gtrsim 4$), for which we expect time lags of $\sim 300 - \sim 1000$ days, depending on redshift. For example, a redshift $z \sim 7$, $M_{BH} \sim 10^9 - 10^{9.5} M_{\odot}$ quasar (e.g. Wang et al. 2021) would be expected to show an observed time lag of about 3 or 6 years respectively, requiring decades of monitoring to measure it without lensing multiplicity. In this paper we suggest that SL time delays of lensed quasars may be a useful route to constraining the R-L relation at those regimes. This is becoming especially timely now that JWST has been uncovering a new abundant population of Type I AGN at high redshifts (e.g., Kocevski et al. 2023a; Harikane et al. 2023; Matthee et al. 2023; Larson et al. 2023; Greene et al. 2023), including the first example of a cluster-lensed multiply imaged one at $z \sim 7$ (Furtak et al. 2023b, 2024).

The work is constructed as follows. In section 2 we start by estimating the expected time lags for high-luminosity and high-redshift quasars (§2.1), and then estimate the distribution of time delays between the multiple images of lensed systems in typical lensing clusters (§2.2). In section 3 we estimate the expected number of lensed quasars and AGN that could be found with e.g., the JWST, *Euclid*, or with the *Roman* Space Telescope, in the near future. In section 5 we demonstrate the suggested idea using simulated light curves, and in section 6 we conclude the work. Throughout we assume a standard flat Λ CDM cosmology with $H_0 = 70 \frac{\text{km}}{\text{s Mpc}}$, $\Omega_{\Lambda} = 0.7$, and $\Omega_m = 0.3$. Magnitudes are quoted in the AB system (Oke & Gunn 1983) and all quoted uncertainties represent 1σ ranges unless otherwise stated.

In this work we use the terms “quasars” and “AGN” synonymously. In addition, whenever we discuss lensed AGN in this work, we explicitly refer to multiply imaged examples.

2. THE DISTRIBUTIONS OF TIME LAGS AND LENSING TIME DELAYS

2.1. RM of quasars: the distributions of time lags

As was noted above, RM of quasars relies on the ability to detect temporal variations in the continuum emission from a quasar that lead to, after a time lag which depends on the BLR size, similar variations in the emission lines from the BLR. In this section, we estimate the BLR sizes – and thus time lags – of a representative sample of quasars, namely the SDSS quasar catalog, which




Figure 1. Histograms of the expected BLR sizes for SDSS DR7 quasars. The catalog contains two distinct subsamples (see sub-section 2.1 for details). The $H\beta$ time-lag distribution is plotted for the subsample of quasars which have an optical continuum measurement, assuming the R-L relation by Bentz et al. (2013). The C IV time-lag distribution is plotted for the subsample of quasars which have a UV continuum measurement, using the relation from Kaspi et al. (2021). The distributions seem to each fit well to a Gaussian distribution, with $[\mu, \sigma] = [1.86, 0.21]$ and $[\mu, \sigma] = [1.78, 0.16]$ for $H\beta$ and C IV, respectively. For example, $\sim 9\%$ ($\sim 26\%$) of the expected C IV ($H\beta$) rest-frame time lags will be 100 days or longer. This means 500 – 1300 observed days, depending on redshift. The reader should also note that in general, for the same quasar, $H\beta$ time-lags are expected to be longer by a factor of a few compared to C IV time-lags, so that in practice the $H\beta$ distribution may extend to even longer lag times.

is based on Data Release 7 (DR7) (Shen et al. 2011) and is available online. This catalog supplies continuum luminosities for either $\lambda L_{\lambda}(1350\text{\AA})$ or $\lambda L_{\lambda}(5100\text{\AA})$, depending on redshift.

The distributions of BLR sizes are shown in Fig.1. The expected $H\beta$ time-lag distribution is obtained assuming the R-L relation from Bentz et al. (2013), for those quasars that have a measurement of the continuum luminosity at 5100 Å. The expected C IV time-lag distribution is obtained assuming the R-L relation from Kaspi et al. (2021) (see their eq. 1), using the continuum UV luminosity at 1350 Å. Note that the C IV subsample consists of 52,427 objects, which have a higher redshift (median redshift of 1.96) and are typically brighter (median $\lambda L_{\lambda}(1350\text{\AA}) = 10^{46.1}$ erg s $^{-1}$) than the $H\beta$ subsample (23,093 objects, median redshift of 0.59, median $\lambda L_{\lambda}(5100\text{\AA}) = 10^{44.6}$ erg s $^{-1}$). As can be seen in Fig. 1, the lag-time distributions, plotted as $\log_{10}(R)$ where R is in light days, are approximately Gaussian, with a mean and width of $[\mu, \sigma] = [1.86, 0.21]$ and $[\mu, \sigma] = [1.78, 0.16]$ for $H\beta$ and C IV, respectively. About 9%, roughly, of the

quasars show CIV time lags of 100 days or longer in the rest-frame, and about 26% of the quasars show H β time lags of 100 days or longer in the rest-frame. This translates to 500 – 1300 days for redshifts of $z \sim 4 - 12$, respectively, which we define here as the high redshift regime.

2.2. The distribution of strong-lensing time delays

In this subsection we aim to obtain a general distribution of the time delays caused by strong lensing in clusters. Galaxy clusters act as gravitational lenses, causing the bending of light. When the projected mass density is larger than some critical density for lensing, strong lensing will occur comprising multiple images of background sources. Using the thin lens approximation, light that travels from a source to the observer will be delayed by some time t (Schneider 1985):

$$t(\vec{\beta}, \vec{\theta}) = \frac{D_l D_s}{D_{ls}} \left[\frac{1}{2} (\vec{\theta} - \vec{\beta})^2 - \psi(\vec{\theta}) \right] \frac{(1 + z_l)}{c}, \quad (1)$$

with D_l , D_s and D_{ls} being the angular diameter distances to the lens, source and between the lens and the source, respectively; z_l is the redshift of the cluster; $\vec{\theta}$ and $\vec{\beta}$ represent the position of the image and source, respectively; and $\psi(\vec{\theta})$ is the lensing potential in that position in the image plane (see Narayan & Bartelmann (1996) for more details). The relevant lensing quantities can be obtained for different clusters using SL lens modeling (e.g., Richard et al. 2014; Mahler et al. 2019; Bergamini et al. 2022). We use mass models made previously by our group (e.g., Zitrin et al. 2015; Furtak et al. 2023c,b,a) for three typical high Einstein mass galaxy clusters: RXC0018, MACS0035 and Abell 2744, where the Einstein mass is the projected mass enclosed within the Einstein radius, and three typical lower Einstein mass clusters: Abell 383, Abell 611 and RX J2129. While all these clusters have typical masses of a few times $10^{14} M_\odot$ to $\sim 10^{15} M_\odot$, the high Einstein mass clusters have a corresponding Einstein radii of $\sim 20 - 30''$ (and thus higher enclosed mass) and the low Einstein mass clusters have Einstein radii of about $\sim 10 - 15''$ typically. Two of the three high mass clusters also happen to show a multiply lensed AGN or quasar-like system with three multiple images each: MACS0035 contains a Type I AGN at $z = 2.06$ (Furtak et al. 2023a), and a redshift $z = 7.05$ “red-dot” AGN was found behind Abell 2744 (Furtak et al. 2024, 2023b).

In order to create the time delay (TD) distribution, we randomly plant sources in the source plane and re-lens them to the image plane using the deflection fields from the model. Typically we use a 100 multiply imaged sources in each case. We calculate the time delays

between all available pairs using eq. 1 and repeat this process for different source redshifts, in the ranges of $1 < z < 2$, $2 < z < 3$, $3 < z < 5$, $5 < z < 8$, $8 < z < 10$ and $8 < z < 12$. The SL time-delay distributions are plotted in Fig.2.

We parameterize the TD distributions and note that they can be well fit (R-square of $\simeq 0.98$) by the following parametrization, inspired by a log-normal distribution function: $P(x) = \frac{1}{a \cdot (d-x)} \cdot \exp\left\{-\frac{[\ln(d-x)-b]^2}{c^2}\right\}$, with $a \sim 10 - 16$ typically (depending on the redshift bin), $b \sim 0.1$, $c \sim 0.8$ and $d \sim 4.6$ for most redshift bins. Various previous works (e.g., Oguri et al. 2002, 2003; Li et al. 2012) have also looked into the TD statistics. Another useful way to parameterize the TD distribution is in log-log space, where we find a slope of ~ 0.8 , similar to what was found for, e.g., Abell 1689 (Li et al. 2012).

In Fig. 2 we also compare the lensing TD probability distributions to the time lags expected from the BLR sizes, recalculated to match each redshift bin. We see that there is an overlap in the distributions such that time delays by both types of galaxy clusters cover the entire range of expected time lags, with a higher probability of having the relevant time delays, i.e., of order the time lag, for higher-luminosity and higher-redshift quasars. About 20%–40% of the TD pairs in all clusters overlap with the range of time lags, depending on the redshift of the lensed source. We see that the peak of the TD distribution caused by lower mass clusters, which should be more abundant, agrees better with the peak of the time lag distributions, but those clusters would have also less multiple images per cluster in proportion to their smaller critical area size (which is roughly proportional to the square root of the mass), such that both populations should be useful. Also overplotted is a line marking a typical time delay expected by a massive galaxy lens – where we nominally adopt for this calculation a SIS galaxy of $\sigma_{vel} \sim 250 \text{ km s}^{-1}$ placed at redshift 0.5 (Oguri 2019), demonstrating that time delays in the case of lens galaxies are too short for a significant gain in the wait time needed for RM of high-redshift, high-mass quasars, and that galaxy clusters are needed for that purpose.

3. EXPECTED NUMBERS OF LENSED QUASARS

To examine whether the proposed idea for RM of multiply imaged quasars at the high luminosity and high redshift ends might become realistic in the near future, we estimate the expected number of quasars that would be multiply imaged by galaxy clusters. Given the many uncertainties involved (see for example section 5), our goal is merely to obtain an order-of-magnitude estimate. We focus on the observational properties of three ob-




Figure 2. Overlap of the distribution of SL time-delays by lensing clusters, and the distribution of expected time lags from the BLR sizes for different redshifts. We show the distribution for both high and low mass galaxy clusters. The redshift limits in each panel’s legend indicate the redshift range of the sources that are lensed to create the SL time delay distribution. The time lags are corrected for the specified, central redshift, i.e., they are given in the observed frame as well. *As can be seen, there is substantial overlap such that SL time delays by galaxy clusters completely cover the range of expected lag times.* We also mark with a dashed black line a typical time delay caused by galaxy-galaxy strong lensing by a massive galaxy (where we adopt a SIS galaxy model with $\sigma = 250 \text{ km s}^{-1}$ placed at $z = 0.5$), showing that only galaxy clusters supply large enough time delays for our purpose. As can be seen, smaller galaxy cluster lenses show more overlap, but they are also expected to show less multiple images per cluster. For more details see section 2.

servatories: *Euclid*, the *Roman* Space Telescope, and JWST. The basic idea is to calculate for each observatory, i.e., a filter set and depth limit, how many quasars in the relevant BH mass and redshift regimes fall within the caustics of lensing clusters so that they would be multiply imaged, and would be bright enough, folding in the magnification, to be detected. We adopt relevant quasar luminosity functions (QLFs) and generally follow known prescriptions for the lensing optical depth, where we assume an explicit cluster mass function – from which an Einstein radius function is calculated – and then calculate the magnification bias and expected fraction (and total number) of multiply imaged quasars in the relevant cluster mass and redshift ranges.

3.1. The Quasar Luminosity Function

For the calculations below we adopt two complementary QLFs: For UV selected quasars we use the one from Willott et al. (2010), and to account for recent populations of AGN uncovered with JWST for example, we also use the QLF from Harikane et al. (2023). Both QLFs are given in terms of the UV absolute magnitude at 1450 Å, M_{1450} . The former also has a term for evolution with redshift, which we assume applies for the range of redshifts spanned here ($4 < z < 12$), while the latter is divided into four discrete QLFs, one for each of the following redshift bins - $z \sim 4$, $z \sim 5$, $z \sim 6$, and $z \sim 7$, where we assume that it does not change significantly from $z \sim 7 - 12$ – but as seen below, even as such the redshift window from $z \sim 8 - 12$ only negligibly contributes to the result. Note that, given that we are interested in an order of magnitude and since the results heavily rely on the various assumptions made throughout, for the QLFs we adopt only the best-fit parameters, without propagating their uncertainties (which would affect the uncertainty on our estimates, too). Also, while the QLF from Willott et al. (2010) is less constrained at the faint end, the QLF from Harikane et al. (2023) is better constrained at that regime.

3.2. The total expected number of quasars

For each observatory we then do the following. We first divide the relevant redshift range into bins. For each bin, we redshift a quasar’s SED and define the selection according to the band in which the quasar would be brightest. Then, using the QLF, which itself is divided into magnitude bins for summation, we calculate how many quasars are expected in that redshift bin down to the observational depth. The *total* number of quasars expected to be observable across the sky is then obtained by summing this number up over the different redshift bins.

For the *Euclid* survey we focus on the detection of typical “blue” quasars and adopt the spectrum from Vanden Berk et al. (2001). Following the *Euclid* overview paper (Laureijs et al. 2011), we primarily use the QLF from Willott et al. (2010), but also for comparison employ the Harikane et al. (2023) one. For each redshift bin we choose as the selection band the band in which the rest frame 1450Å falls. If the redshifted wavelength is lower (or higher) than the survey’s available bands, we choose the nearest band to it. Since in this work we are particularly interested in lensed high-mass and high-redshift quasars, we perform the calculation for those two regimes specifically.

We follow the same procedure for the *Nancy Grace Roman Space Telescope*. Both observatories cover roughly similar wavelength ranges into the near-infrared. While the nominal FOV for *Roman* is about half of that of *Euclid*, *Roman* has a two times in diameter larger mirror and the observational limits are deeper than the ones for *Euclid*, hence we may expect higher numbers of quasars to be detectable with it. Again our focus remains on the high-redshift and high-luminosity quasars and as above we adopt the typical SED from Vanden Berk et al. (2001).

Last we turn to perform the same calculation for JWST. In addition to typical “blue” quasars that are detectable with JWST at redshifts higher than ~ 4 , given the new population of “red-dot” AGN at high redshift that has been discovered in JWST observations in increasing numbers (e.g., Furtak et al. 2023b; Kokorev et al. 2023; Matthee et al. 2023; Labbe et al. 2023; Kocovski et al. 2023b, 2024), we also consider here an additional quasar SED model. The SED of this type of object is heavily reddened and thus is brighter towards the red, with some UV light from star formation or scattered AGN radiation. As the representative SED for red-dot objects we use the one from Furtak et al. (2023b), and we only consider here the QLF from Harikane et al. (2023) which was based on or commensurate with this new population of quasars (Greene et al. 2023). Given the red SED, for red-dot AGNs we now choose the selection band to be the one in which the rest frame 5500Å falls, shifting all selection to the red-optical end of the spectrum, which indeed, is currently only observable with JWST for these redshifts. We again focus on the high-redshift population for our calculations and estimate the expected numbers of quasars to be detected with JWST for both blue quasars and red quasars. To account for JWSTs extreme depth capabilities we assume a depth of 30 magnitudes, reachable in several hours with JWST.

3.3. The lensed fraction

Various works have calculated the effect of lensing on the luminosity function and the fraction of some population of lensed sources of interest such as galaxies, quasars or supernovae, and the prescriptions for doing so are well established. Here we largely follow the outlined procedure in Yue et al. (2022), where they obtained an estimate for the fraction of lensed quasars mostly by galaxies. i.e. in a different mass regime that is our interest here, and expand it to massive galaxy clusters. Various components of the formalism are also based on e.g., Oguri & Marshall (2010); Wyithe et al. (2011); Mason et al. (2015a). One other important difference is that in this formalism, given it was usually applied to galaxies, a singular isothermal spheres (SIS) mass density profile is typically assumed, where here we consider the Navarro–Frenk–White (NFW) mass density profile Navarro et al. (1996), which is a more realistic approximation for the mass distribution of clusters.

One can define the optical depth for multiple images, which describes how likely it is for a source at z_s to lie within the caustics of a lens (the Einstein radius mapped back to the source plane), and thus be multiply imaged. The optical depth for multiple images is then given by:

$$\tau_m = f_{sp} \int_0^{z_s} dz_l \int dM \phi(M, z_l) \frac{d^2 V_c}{d\Omega dz_l} \pi \theta_E(M, z_l, z_s)^2 D_l^2, \quad (2)$$

where $\phi(M, z_l)$ is the lens density function and taken here to simply be the halo mass function (see eq. 3), z_l is the redshift of the lens, D_l is the angular diameter distance to it, and f_{sp} is the fraction of the critical area ($\pi \theta_E^2$) in which sources would be multiply imaged (for example, for a SIS lens, $f_{sp} = 1$). The expression $\frac{d^2 V_c}{d\Omega dz_d} = (1 + z_d)^3 c \frac{dt}{dz_d}$ is the differential co-moving volume¹, with $\frac{dt}{dz_d} = \frac{1}{H(z)(1+z)}$ where $H(z) = H_0(\Omega_m(1+z)^3 + \Omega_\Lambda)^{1/2}$.

For our calculations we adopt the mass function from Tinker et al. (2008), namely:

$$\frac{dn}{dM} = f(\sigma) \frac{\rho_m}{M} \frac{d \ln \sigma^{-1}}{dM} \quad (3)$$

with $\rho_m = \rho_{crit} \Omega_m$ being the mean matter density, σ is the RMS variance of a spherical top hat containing a mass M and $f(\sigma)$ is the halo multiplicity function given also in Tinker et al. (2008). In practice, to calculate the mass function (and the power spectrum required therein) we use `cluster_toolkit` and `CLASS`. We refer the reader to <https://cluster-toolkit.readthedocs.io/en/latest/source/massfunction.html> for more informa-

tion. The Einstein radius of a NFW lens depends on the mass, concentration and the redshifts of lens and the source. To obtain the Einstein radius for the NFW lens for each combination of parameters we follow the procedure from Broadhurst & Barkana (2008) and Sadegh & Rephaeli (2008, see also Bartelmann 1996), while we assume the concentration-mass (c-M) relation from Meneghetti et al. (2014):

$$c(M, z) = A \left(\frac{1.34}{1+z} \right)^B \left(\frac{M}{8 \times 10^{14} h^{-1} M_\odot} \right)^C \quad (4)$$

with $A = 3.757 \pm 0.054$, $B = 0.288 \pm 0.077$ and $C = -0.058 \pm 0.017$ (although note, we only use here the best fit values to translate the mass to Einstein radius; we do not propagate the scatter in this relation).

Analytically, for a NFW lens multiple images will occur if the source lies within the radius of the radial critical curve (Meneghetti et al. 2003). We however adopt here the more empirical assumption of $f_{sp} = 0.1$, which we obtained from an ensemble of cluster models for known lensing clusters (e.g., Zitrin et al. 2015): We find that in practice, the angular source-plane area enclosed by the caustics is about $\sim 10\%$ of the angular area enclosed by the critical curves in the image plane (see also Broadhurst et al. (2005)).

Next we need to consider that in flux-limited surveys, objects that lie intrinsically below the detection threshold can be detected if they are sufficiently magnified. However, there exists a trade-off with the smaller area probed in the source plane and the actual gain in number count would depend also on the shape of the luminosity function (Broadhurst et al. 1995; Wyithe et al. 2011; Fialkov & Loeb 2015). The magnification bias can be written as:

$$B = \frac{\int_{\mu_{min}}^{+\infty} d\mu p(\mu) N(> L_{lim}/\mu)}{N(> L_{lim})}, \quad (5)$$

where $p(\mu)$ is the magnification distribution of lensed sources, L_{lim} is the survey flux limit, and $N(> L_{lim})$ is the cumulative luminosity function, i.e., the number of background sources brighter than L_{lim} .

We use $\mu_{min} = 2$ for the magnification bias calculation, and, as the magnification probability function we adopt $p(\mu) \propto 1/(\mu)^3$ (truncated at $\mu = 100$). This distribution seems to approximately describe the magnification distribution of lensed images, also in NFW lenses (see for example Figure 9 in Wyithe et al. (2001)). We also verify this by simulating a NFW lens and calculating the magnification distribution of randomly planted sources.

As is also indicated in Yue et al. (2022), out of all sources with apparent luminosity brighter than L_{lim} ,

¹ Note that in some work the differential co-moving volume is defined such that it contains the D_l^2 factor seen in eq. 2.

the lensed fraction is given by (Mason et al. 2015a):

$$F_{multi} = \frac{B\tau_m}{B\tau_m + B'(1 - \tau_m)} \quad (6)$$

where B' is the magnification bias of non lensed sources, and we assume here it is $B' \approx 1$ as in Yue et al. (2022); Mason et al. (2015b); Wyithe et al. (2011).

After calculating the expected number of quasars to be detected as explained in the previous section, the number is multiplied by the lensed fraction F_{multi} for each redshift bin. Eventually, a total lensed fraction is calculated. We perform the calculation for three different regimes - quasars at all redshifts and luminosity ranges, high luminosity quasars (i.e. $M_{1450} < -26.5$ which corresponds to $\lambda L_{UV,1450} > 10^{46.5}$ erg s $^{-1}$) and high redshift quasars, namely $4 < z < 12$. Last, out of the high redshift candidates we aim to obtain the number of quasars for which the notions we highlight here might be particularly useful, i.e, those with long lag times. We follow the same procedure, only this time for each redshift bin we find the lower limit for the luminosity such that the expected time lag of the C IV line is at least, e.g., a 100 days. As can be seen in Fig.1, the expected time lags for the $H\beta$ emission line will be even longer.

4. RESULTS

We summarize the results in Table 1. For all observatories, we see that the chances to detect a multiply imaged, high-luminosity quasar are very small - with < 1 that are expected to be detected across the sky, with both QLFs. Nonetheless, for the high redshift regime, we obtain that many such lensed AGN should be found. Explicitly, we find that $\sim 3 - 400$ multiply imaged AGN may be detected with *Euclid* over the whole sky, with the exact number depending on the choice of QLF. For the *Roman* Space telescope, we similarly estimate $\sim 13 - 9,800$ detectable, multiply imaged quasars, and for JWST we find that $\sim 3 \times 10^4 - 5 \times 10^4$ quasars multiply imaged by galaxy clusters might be detected all over the sky. In general, then, the numbers of multiply imaged AGN above correspond to about $\sim 10^{-4}$ of the total (unlensed) population.

It is interesting to note that JWST has imaged to date about ~ 10 clusters and in one of them, Abell 2744, a conspicuous multiply imaged AGN at $z \simeq 7$ has already been discovered (Furtak et al. 2023b). If we take this 1 in 10 ratio as representative, then many more such multiply imaged AGN should be detected soon with JWST, as it expands its observations to a larger cluster sample. Moreover, with $\sim 100,000$ clusters in the sky one could expect about $\sim 10,000$ multiply-imaged AGN de-

tectable with JWST, which is of the same order of magnitude as the result obtained in our more detailed calculation, implying that the expected number is representative.

While it is encouraging to see that many AGN multiply imaged by galaxy clusters could be detected across the sky, it is also insightful to examine which fraction of this lensed population has a large expected time lag, such that the gain from lensing would be most significant. Columns 5 and 6 in Table 1 list the expected number of multiply imaged AGN that are expected to have lag times of, as an example, > 50 and > 100 days. For *Euclid*, we expect $\sim 3 - 380$ multiply imaged AGN with > 50 days observed time lags and $\sim 2 - 35$ with > 100 days; $\sim 4 - 2150$ and $\sim 2 - 207$ with the *Roman* space telescope; and $\sim 2750 - 3500$ and $\sim 140 - 190$ with JWST, respectively, across the sky. This is about 1-2 orders of magnitude lower than the total number of high-redshift AGN that are multiply imaged by galaxy clusters, but shows that there exists a lensed population in the sky which can gain substantially from RM campaigns to probe the R-L relation at high-redshift, as we also discuss further below (section 5).

In addition to the above, it is seen in Table 1 that for both *Euclid* and the *Roman* space telescope higher numbers of quasars are generally expected assuming the QLF from Harikane et al. (2023) compared to the QLF from Willott et al. (2010). The order of magnitude difference is not surprising given the high number density of broad-line AGN recently detected with JWST to fainter magnitudes (e.g., Furtak et al. 2023b; Labbe et al. 2023; Greene et al. 2023; Kocevski et al. 2023a; Matthee et al. 2023).

In Table 1, for completeness, the first column lists the numbers of quasars *across all redshifts* that are expected to be multiply imaged by galaxy clusters. It should be noted that both the QLFs we adopt for this calculation are, however, limited in redshift. For the Harikane et al. (2023) QLF, which is defined between redshifts $z \sim 4$ and $z \sim 8$, we assume no additional evolution: We use the $z \sim 8$ QLF for all redshifts above $z \sim 8$, which contributes only negligibly to the final numbers, and the $z \sim 4$ QLF to all redshift below $z \sim 4$. Since the QLF is expected to rise down to $z \sim 2$ and then turn over in resemblance to the cosmic star formation density (e.g., Richards et al. 2006; Shen et al. 2020), we note that the exact effect of this assumption is not clear, in particular since the evolution of “little red-dot” and other JWST selected AGN is not well established yet to lower redshifts. Regarding the QLF from Willott et al. (2010), we note that it has a term for a monotonic evolution with redshift, although it was defined in practice over

QLF reference	all sources	high L	high z	C IV >50 days	C IV > 100 days
<i>Euclid</i>					
Willott et al. (2010)	50 (1.3×10^7)	< 1 (19,800)	3 (84,000)	3 (81,500)	2 (41,800)
Harikane et al. (2023)	2,000 (1.5×10^8)	< 1 (< 1)	400 (2×10^6)	376 (1.7×10^6)	36 (45,000)
<i>Roman Space Telescope</i>					
Willott et al. (2010)	200 (5.9×10^7)	< 1 (19,800)	13 (4.2×10^5)	4 (1.2×10^5)	2 (46,000)
Harikane et al. (2023)	32,700 (2.4×10^9)	< 1 (< 1)	9,800 (1.3×10^8)	2,150 (1.7×10^7)	207 (1.1×10^6)
JWST					
Harikane et al. (2023)	1.3×10^5 (8.8×10^9)	< 1 (< 1)	5.3×10^4 (7.2×10^8)	2,750 (2.4×10^7)	140 (1.1×10^6)
Harikane et al. (2023) (red selection)	—	< 1 (< 1)	3.5×10^4 (5.3×10^8)	3,500 (2.4×10^7)	190 (1.1×10^6)

Table 1. The estimated numbers of lensed quasars that might be detected with *Euclid*, *Roman* space telescope and JWST. In every column, we also specify within the brackets the total number of quasars expected to be detected; *Column 1*: The assumed QLF. The detection limit is calculated assuming a blue selection unless otherwise specified; *Column 2*: The estimated number of *lensed* quasars at all redshifts and all luminosity range; *Column 3*: The estimated number of *lensed* quasars at all redshifts and high luminosity range; *Column 4*: The estimated number of *lensed* quasars at high redshifts; *Columns 5-6*: The estimated number of *lensed* quasars with luminosities and redshift that would result in a C IV observed time lag of above 50 and 100 days;

the redshift range $\sim 5.5 - 6.5$ and the redshift evolution term assumes a monotonically increasing number density towards lower redshifts. In that case, assuming this QLF below $z \sim 4$ might cause an over estimate of the number of quasars at the low redshift regime. A detailed examination of the effect of the $z \sim 1 - 4$ range on the total numbers is out of scope for this work, in particular, since here we focus on the high-redshift regime $z \gtrsim 4$.

5. DISCUSSION

To examine the feasibility of the idea proposed in this work, in the first part of this work we have shown the distribution of time delays by typical lensing clusters and compared it to the time lag distribution of quasars across their mass range and at various redshifts. After showing that indeed galaxy clusters supply useful time delays that can easily be as large as the the time lags expected for highest luminosity and highest redshift AGN, in the second part of this work we then estimated the number of AGN that are expected to be multiply imaged by galaxy clusters and observable with different observatories of interest.

Before concluding the work and while perhaps quite intuitive, we now aim to provide a more visual demonstration of the idea, using an idealized case. In particular, we consider an illustrative scenario in which the expected time lag is equal to the time delay between two images of a quasar. We end by referring to a couple of recent lensed examples for which such RM observational campaigns may be feasible using existing facilities.

To simulate a typical continuum light curve of a quasar, we use mock light curves of the quasar HE0435-1223 generated by H0LiCOW collaboration and available online (Wong et al. 2017; Chen et al. 2019). We

simulate the emission-line light curve by replicating the continuum light curve shifted by a certain time lag and add for illustration purposes, a Gaussian noise with a standard deviation = 0.05 in flux.

In Fig. 3 we show the mock light curves for two lensed images A and B for a SL time delay of 1000 days, where it is in this case equal to the observed time lag. An observed BLR size of a 1000 days corresponds to a physical BLR size of $1000/(1+z)$ light days. At a redshift of $z = 2$, for example, this corresponds to a SMBH luminosity of $\lambda L_\lambda \sim 4 \times 10^{47}$ erg s^{-1} , according to the local scaling relations. For each image we show the same time-frame of 1200 days in the observer frame, while the duration was arbitrarily chosen for illustration purposes. For each image separately, similar to the case in which the quasar were unlensed, the pattern seen in the first 1000 days in continuum is seen repeating itself in emission lines a 1000 days later. In practice, to securely identify such an offset it would require at least several times this timeframe, hence order a decade or more of continuous monitoring. However, due to the TD being similar to the time lag, the same changes in the continuum seen in the trailing image are seen *at the same time* in emission lines in the leading image, essentially saving the years-long wait time to observe the lagging emission line changes. Suppose that in order to detect the ~ 3 year time lag a monitoring campaign spanning 15 years were needed for an unlensed source to accumulate enough data points and timespan (the true value depends on a variety of factors that we do not go into here). Investing 15 years of observing time would be looked upon as an uneasy and perhaps risky investment given that the output is uncertain. In contrast, imagine a lensed quasar such that the source is multiply imaged

with at least three images, which - more often than not - would be the case for wide separation lensed quasars as we consider here (e.g., Sharon et al. 2017; Williams et al. 2021). In such a case the investment becomes much more reasonable: each observation triples the number of data-points compared to the unlensed case, and more importantly, assuming that $t_{TD} \sim t_{lag}$ then the corresponding emission line changes can be observed (even if not immediately identified) already in the first few batches of monitoring. Furthermore, each image would then be magnified, which would also ease the observational follow-up.

As another example, in the case of a smaller BLR size, causing a shorter time lag than the SL time delay, the pattern seen in the first days of observations in continuum would already be seen repeated in the emission lines after a short time lag. This is similar to the way it would appear for an unlensed quasar, where as for the lensed case we mainly gain magnification and additional sampling from the different multiple images, making the reverberation mapping more efficient overall. Also in the opposite scenario where the SL time delay is much shorter than the expected time lag, while we do not gain much in terms of overall wait time, we gain in several other aspects as above. First, we obtain additional points in the light curve with the same observation, compared to the unlensed case (an advantage that increases even further when more than two quasars images are seen, as is often the case in cluster lensing). Second, the images are magnified, which facilitates their follow-up. Third, while of singly imaged (or unlensed) objects can only be observed for several months a year, when they are away from the Sun’s direction, a multiply-imaged quasar may allow us to fill gaps in the light curves also in those otherwise down times.

It should however be noted that even if TDs are used to facilitate RM, the monitoring time should still be sufficiently significant so as not to bias the measurements towards the available TDs.

Recently, a successful reverberation mapping was performed on the multiply imaged quasar, SDSS J2222+2745, at $z = 2.805$ (Williams et al. 2021), demonstrating the usefulness of multiply imaged quasars in that respect. A time lag of $\tau_{cen} = 36.5_{-3.9}^{+2.9}$ restframe days was significantly detected for the C IV emission line, whereas the time delays to the two other images of the quasar are about ~ 40 and ~ 700 days (see also Sharon et al. 2017). Hence, one time delay is of the order of the time lag, similar to first case discussed above (but with a shorter time lag than what we consider here, which is aimed for the high-redshift and high luminosity

regimes), and the second time delay is much longer than the time lag, similar to second case discussed above.

Note that throughout we assume that any object detected by a certain observatory could also be followed up spectroscopically (by the same or other observatories). In addition, given that we only wish to demonstrate the basic idea, we only show an idealized case. We also neglect here other sources of noise such as local microlensing for some of the images, which is known to often be present and cause its own variations to the light curve for some of the components (e.g., Eigenbrod et al. 2008; Mediavilla et al. 2015; Vernardos et al. 2024) – depending on the mass of the point mass and size of the emitting source – although even in such cases the effects can be remedied (see e.g. Williams et al. (2021)). Similarly, we do not take into account so-called “changing-look” AGN (e.g. Ricci & Trakhtenbrot 2023), for which some of the images may not show, for example, broad-line emission, while the other images might (in fact, lensing TDs may be a route for detecting lensed AGN of this type).

One should also bear in mind that in reality the TD is not necessarily known from the lens model to the required accuracy to remove it from the fit altogether. However, it might be first determined from the temporal shift between the continua of the two images before determining the time lag. This may improve the error on the TDs compared to the errors from the lens model’s TD, which would otherwise propagate into the R-L relation as well.

It is also worth highlighting a couple of interesting recent objects which may become useful targets for RM of lensed AGN (other examples naturally exist, too). First we note the triply-imaged Little Red Dot AGN reported by Furtak et al. (2023b) at $z = 7.045$. This object is currently the highest-redshift multiply imaged AGN. It has according to their lens model a TD of ~ 3 years, roughly, between the two closest images, and a TD of about 19 years between the first of these two and the third, more distant image, which is first to arrive. The NIRSpc spectrum of the source was published in Furtak et al. (2024). Based on its luminosity, with a few hours per visit we could detect both the continuum and the $H\beta$ line (no C IV lines were detected in the object, despite the very deep integration time). One can thus envision that a NIRSpc/JWST program, for example, targeting the three images simultaneously, could be used to measure both the continuum and emission lines in each epoch. From the width of $H\beta$, Furtak et al. (2024) derived a BH mass of $4 \times 10^7 M_{\odot}$, which, according to the scaling relations by Kaspi et al. (2021), corresponds to an expected or observed time lag of a couple dozen days. This time lag is relatively short (corresponding to the second




Figure 3. Simulated idealized light curve for the case where the time delay equals the time lag. In the top panel of the figure we demonstrate mock light curves of two images of the same source. The time axis represents the observing time of the images and the y axis represents a normalized flux. The SL time delay between the two images is 1000 days, image A is the trailing image and image B is the leading image. The orange and blue curves correspond to the mock continuum and line emission light curves respectively. Since the two images will have different magnification factors, the leading image shown in the top right panel has its flux normalized lower, as it will usually have a lower magnification than the trailing one. Details on the mock data are given in section 5. In the bottom panel we show the full mock light curves with shaded areas emphasizing the data obtained from each image and the overlap. The mock light curves of the leading image B are corrected by the time delay. We emphasize the time lag between the continuum and emission curves at several points with arrows.

case discussed above), and while its measurement could in principle also be done were it not lensed, its measurement would be facilitated by measuring the three images simultaneously. While the expected time lag for this object does not appear to be very long (so that the gain in wait time is not maximized), it could supply a unique example for probing the RM relation at high-redshift.

A second, recent example is the large separation $z = 3.27$ quasar COOL J0335-1927 (Napier et al. 2023a), for which time delays of ~ 500 days and ~ 130 days between two pairs of its images were predicted from the lens model, although with quite substantial uncertainty. To estimate the expected time lag, we adopt the typical g -band luminosity seen in Fig. 4 in their work, and translate it to an absolute UV magnitude of ~ 25.5 AB, which, based on existing scaling relations (Kaspi et al. 2021), corresponds to an observed time lag of ~ 300 days. The object therefore might benefit significantly from such a campaign. For more discussion on the latter object, see also Napier et al. (2023a).

We thus conclude that designated observations of multiply imaged quasars can open the door for RM studies in a much more efficient manner, as was already demonstrated at lower-redshifts Williams et al. (2021). For the most massive and/or high-redshift quasars for which the time lags may be otherwise too long, this might possibly be the only, feasible way.

6. SUMMARY

In this work we discuss the possibility of using lensing time delays to facilitate the RM of high-luminosity and high-redshift quasars, for which the radius-luminosity relation is currently, poorly constrained. The expected time lags for very-massive or high-redshift quasars can reach several years and would likely require much longer monitoring times than can be comfortably accommodated by typical observing campaigns. Multiply imaged AGN may, however, offer a unique opportunity to explore the radius-luminosity relation at these ends. Not only do strongly lensed quasars comprise several magnified images which boost the efficiency of the light curve sampling, but also, the time delay between the quasar's multiple images can save the long wait times that are

typically required. For example, if the time delay is roughly of the order of the expected time lag, changes in the emission lines in a leading image can be observed around the same time as the changes in the continuum in a trailing image, thus allowing a much reduced wait times and overall monitoring times-spans.

We start by showing the typical distribution of BH time lags compared to the lensing TD distribution from cluster lenses and show that a strong overlap exists. We then estimate the number of multiply imaged quasars that may be detectable in the near future with *Euclid*, *Roman*, or JWST. We find that indeed, a fair amount of multiply imaged high-redshift quasars, out of which some would experience a significantly long observed time lag, should be detectable. Using the time delays between the multiple images, RM of these possible future candidates could be obtained in a reasonable few-year monitoring timescale, probing the poorly known R-L relation in the early Universe.

ACKNOWLEDGEMENTS

M.G would like to thank Yuichi Harikane for a useful discussion. M.G is thankful for the hospitality of the SURF program and cosmic DAWN center where much of this work has been carried out. The BGU group acknowledges support by grant No. 2020750 from the United States-Israel Binational Science Foundation (BSF) and grant No. 2109066 from the United States National Science Foundation (NSF); by the Israel Science Foundation Grant No. 864/23; and by the Ministry of Science & Technology, Israel. D.C. acknowledges support by grants from the German Research Foundation (HA3555-14/1, CH71-34-3) and the Israeli Science Foundation (2398/19, 1650/23). This research made use of *Astropy*,² a community-developed core Python package for Astronomy (Astropy Collaboration et al. 2013; Price-Whelan et al. 2018) as well as the packages `cluster_toolkit`, `CLASS` (Blas et al. 2011), `NumPy` (van der Walt et al. 2011), `SciPy` (Virtanen et al. 2020) and the astronomy library for `MATLAB` (Ofek 2014). The `Matplotlib` package (Hunter 2007) was used to create some of the figures in this work.

REFERENCES

- Acebron, A., Grillo, C., Bergamini, P., et al. 2022, *A&A*, 668, A142
- ² <http://www.astropy.org>
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Bahcall, J. N., Kozlovsky, B.-Z., & Salpeter, E. E. 1972, *ApJ*, 171, 467
- Bartelmann, M. 1996, *A&A*, 313, 697

- Bentz, M. C., Peterson, B. M., Netzer, H., Pogge, R. W., & Vestergaard, M. 2009, *ApJ*, 697, 160
- Bentz, M. C., Denney, K. D., Grier, C. J., et al. 2013, *ApJ*, 767, 149
- Bergamini, P., Acebron, A., Grillo, C., et al. 2022, arXiv e-prints, arXiv:2207.09416
- Blandford, R. D., & McKee, C. F. 1982, *ApJ*, 255, 419
- Blas, D., Lesgourgues, J., & Tram, T. 2011, *JCAP*, 2011, 034
- Broadhurst, T., Benítez, N., Coe, D., et al. 2005, *ApJ*, 621, 53
- Broadhurst, T. J., & Barkana, R. 2008, *MNRAS*, 390, 1647
- Broadhurst, T. J., Taylor, A. N., & Peacock, J. A. 1995, *ApJ*, 438, 49
- Chen, G. C. F., Fassnacht, C. D., Suyu, S. H., et al. 2019, *MNRAS*, 490, 1743
- Ding, X., Silverman, J., Treu, T., et al. 2020, *ApJ*, 888, 37
- Eigenbrod, A., Courbin, F., Sluse, D., Meylan, G., & Agol, E. 2008, *A&A*, 480, 647
- Fanidakis, N., Baugh, C. M., Benson, A. J., et al. 2012, *MNRAS*, 419, 2797
- Ferrarese, L., & Merritt, D. 2000, *ApJL*, 539, L9
- Fialkov, A., & Loeb, A. 2015, *ApJ*, 806, 256
- Furtak, L. J., Mainali, R., Zitrin, A., et al. 2023a, *MNRAS*, 522, 5142
- Furtak, L. J., Zitrin, A., Plat, A., et al. 2023b, *ApJ*, 952, 142
- Furtak, L. J., Zitrin, A., Weaver, J. R., et al. 2023c, *MNRAS*, 523, 4568
- Furtak, L. J., Labbé, I., Zitrin, A., et al. 2024, *Nature*, 628, 57
- Goulding, A. D., Greene, J. E., Setton, D. J., et al. 2023, *ApJL*, 955, L24
- Greene, J. E., & Ho, L. C. 2005, *ApJ*, 630, 122
- Greene, J. E., Labbe, I., Goulding, A. D., et al. 2023, arXiv e-prints, arXiv:2309.05714
- Habouzit, M., Onoue, M., Bañados, E., et al. 2022, *MNRAS*, 511, 3751
- Harikane, Y., Zhang, Y., Nakajima, K., et al. 2023, arXiv e-prints, arXiv:2303.11946
- Hunter, J. D. 2007, *Computing in Science & Engineering*, 9, 90
- Inada, N., Oguri, M., Pindor, B., et al. 2003, *Nature*, 426, 810
- Kaspi, S. 2018, *Astrophysics and Space Science*, 363, doi:10.1007/s10509-018-3305-2
- Kaspi, S. 2021, in *Nuclear Activity in Galaxies Across Cosmic Time*, ed. M. Pović, P. Marziani, J. Masegosa, H. Netzer, S. H. Negu, & S. B. Tessema, Vol. 356, 116–121
- Kaspi, S., Brandt, W. N., Maoz, D., et al. 2021, *ApJ*, 915, 129
- Kaspi, S., Maoz, D., Netzer, H., et al. 2005, *ApJ*, 629, 61
- Kaspi, S., Smith, P. S., Netzer, H., et al. 2000, *ApJ*, 533, 631
- King, A. 2024, *MNRAS*, 531, 550
- Kocevski, D. D., Onoue, M., Inayoshi, K., et al. 2023a, *ApJL*, 954, L4
- . 2023b, *ApJL*, 954, L4
- Kocevski, D. D., Finkelstein, S. L., Barro, G., et al. 2024, arXiv e-prints, arXiv:2404.03576
- Kokorev, V., Fujimoto, S., Labbe, I., et al. 2023, *ApJL*, 957, L7
- Kormendy, J., & Ho, L. C. 2013, *ARA&A*, 51, 511
- Labbe, I., Greene, J. E., Bezanson, R., et al. 2023, arXiv e-prints, arXiv:2306.07320
- Larson, R. L., Finkelstein, S. L., Kocevski, D. D., et al. 2023, *ApJL*, 953, L29
- Laureijs, R., Amiaux, J., Arduini, S., et al. 2011, arXiv e-prints, arXiv:1110.3193
- Li, X., Hjorth, J., & Richard, J. 2012, *JCAP*, 2012, 015
- Lupi, A., Trinca, A., Volonteri, M., Dotti, M., & Mazzucchelli, C. 2024, arXiv e-prints, arXiv:2406.17847
- Mahler, G., Sharon, K., Fox, C., et al. 2019, *ApJ*, 873, 96
- Mason, C. A., Trenti, M., & Treu, T. 2015a, *ApJ*, 813, 21
- Mason, C. A., Treu, T., Schmidt, K. B., et al. 2015b, *ApJ*, 805, 79
- Matthee, J., Naidu, R. P., Brammer, G., et al. 2023, arXiv e-prints, arXiv:2306.05448
- Mediavilla, E., Jiménez-vicente, J., Muñoz, J. A., & Mediavilla, T. 2015, *ApJL*, 814, L26
- Meneghetti, M., Bartelmann, M., & Moscardini, L. 2003, *MNRAS*, 346, 67
- Meneghetti, M., Rasia, E., Vega, J., et al. 2014, *ApJ*, 797, 34
- Napier, K., Gladders, M. D., Sharon, K., et al. 2023a, *ApJL*, 954, L38
- . 2023b, *ApJL*, 954, L38
- Narayan, R., & Bartelmann, M. 1996, *ArXiv Astrophysics e-prints*, astro-ph/9606001, arXiv:astro-ph/9606001
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1996, *ApJ*, 462, 563
- Netzer, H., & Peterson, B. M. 1997, in *Astrophysics and Space Science Library*, Vol. 218, *Astronomical Time Series*, ed. D. Maoz, A. Sternberg, & E. M. Leibowitz, 85
- Ofek, E. O. 2014, *MAAT: MATLAB Astronomy and Astrophysics Toolbox*, *Astrophysics Source Code Library*, record ascl:1407.005, , , ascl:1407.005
- Oguri, M. 2019, *Reports on Progress in Physics*, 82, 126901
- Oguri, M., & Marshall, P. J. 2010, *MNRAS*, 405, 2579

- Oguri, M., Suto, Y., & Turner, E. L. 2003, *ApJ*, 583, 584
- Oguri, M., Taruya, A., Suto, Y., & Turner, E. L. 2002, *ApJ*, 568, 488
- Oguri, M., Schrabback, T., Jullo, E., et al. 2013, *MNRAS*, 429, 482
- Oke, J. B., & Gunn, J. E. 1983, *ApJ*, 266, 713
- Peterson, B. M. 1993, *PASP*, 105, 247
- Price-Whelan, A. M., Sipőcz, B. M., Günther, H. M., et al. 2018, *AJ*, 156, 123
- Reines, A. E., & Volonteri, M. 2015, doi:10.1088/0004-637X/813/2/82. <http://blackhole.berkeley.edu>
- Ricci, C., & Trakhtenbrot, B. 2023, *Nature Astronomy*, 7, 1282
- Richard, J., Jauzac, M., Limousin, M., et al. 2014, arXiv, 1405.3303, arXiv:1405.3303
- Richards, G. T., Strauss, M. A., Fan, X., et al. 2006, *AJ*, 131, 2766
- Sadeh, S., & Rephaeli, Y. 2008, *MNRAS*, 388, 1759
- Schneider, P. 1985, *A&A*, 143, 413
- Sharon, K., Bayliss, M. B., Dahle, H., et al. 2017, doi:10.3847/1538-4357/835/1/5
- Shen, X., Hopkins, P. F., Faucher-Giguère, C.-A., et al. 2020, *MNRAS*, 495, 3252
- Shen, Y., Richards, G. T., Strauss, M. A., et al. 2011, *ApJS*, 194, 45
- Shen, Y., Grier, C. J., Horne, K., et al. 2023, arXiv e-prints, arXiv:2305.01014
- Tinker, J., Kravtsov, A. V., Klypin, A., et al. 2008, *ApJ*, 688, 709
- van der Walt, S., Colbert, S. C., & Varoquaux, G. 2011, *Computing in Science Engineering*, 13, 22
- Vanden Berk, D. E., Richards, G. T., Bauer, A., et al. 2001, *AJ*, 122, 549
- Vernardos, G., Sluse, D., Pooley, D., et al. 2024, *SSRv*, 220, 14
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature Methods*, 17, 261
- Volonteri, M. 2012, *Science*, 337, 544
- Wang, F., Yang, J., Fan, X., et al. 2021, *ApJL*, 907, L1
- Wang, F., Yang, J., Fan, X., et al. 2021, *The Astrophysical Journal Letters*, 907, L1. <https://doi.org/10.3847%2F2041-8213%2Fabd8c6>
- Williams, P. R., Treu, T., Dahle, H., et al. 2021, *ApJ*, 911, 64
- Willott, C. J., Delorme, P., Reylé, C., et al. 2010, *AJ*, 139, 906
- Wong, K. C., Suyu, S. H., Auger, M. W., et al. 2017, *MNRAS*, 465, 4895
- Wyithe, J. S. B., Turner, E. L., & Spergel, D. N. 2001, *ApJ*, 555, 504
- Wyithe, J. S. B., Yan, H., Windhorst, R. A., & Mao, S. 2011, *Nature*, 469, 181
- Yue, M., Fan, X., Yang, J., & Wang, F. 2022, *ApJ*, 925, 169
- Zitrin, A., Fabris, A., Merten, J., et al. 2015, *ApJ*, 801, 44