

# THE ORLOV SPECTRA OF ABELIAN CATEGORIES\*

JUNLING ZHENG<sup>1</sup> AND YU-ZHE LIU<sup>2,†</sup>

ABSTRACT. We introduced the notion of Orlov spectra of Abelian categories, and study its some properties. In particular, we give precise result of Orlov spectra of algebras with type  $\mathbb{A}_n$ .

## CONTENTS

1. Introduction	1
2. Preliminaries	2
2.1. The extension dimension of Abelian category	2
2.2. Ghosts and coghosts	3
2.2.1. Contravariantly finite subcategories and covariantly finite subcategories	4
2.2.2. Ghost lemma and Coghost lemma	4
2.3. Coghosts on Artin algebra	7
2.3.1. Some properties of coghost on Artin algebra	7
2.3.2. Some properties of strong generator over Artin algebra	8
3. Loewy lengths	9
3.1. Properties of Loewy lengths	10
3.2. Strong generators	11
3.3. Loewy lengths of some Artin algebras	12
3.3.1. Loewy lengths of SPI Artin algebras	12
3.3.2. Loewy lengths of semi-simple Artin algebras	13
3.3.3. Loewy length of Artin algebras with Orlov spectra $\{0, 1\}$	14
4. Radical Layer lengths	15
4.1. Torsion pairs and radical Layer lengths	15
4.2. Short exact sequences and radical layer length	16
4.2.1. The first main result and some corollaries	16
4.2.2. The second main result and some corollaries	19
5. Application: The Orlov spectrum with type $\mathbb{A}_n$	23
Acknowledgements	27
References	27

## 1. INTRODUCTION

Rouquier introduced in [21, 22] the dimension of a triangulated category under the idea of Bondal and van den Bergh in [8]. Roughly speaking, it is an invariant that measures how quickly the category can be built from one object. This dimension plays an important role in representation theory [5, 7, 10, 12, 19, 21, 22, 26]. For example, it can be used to

\* **2010 Mathematics Subject Classification:** 18G20, 16E10, 18E10.

**Key words:** Extension dimensions, Artin algebras, Orlov spectrums.

<sup>1</sup> Department of Mathematics, China Jiliang University, Hangzhou, 310018, Zhejiang Province, P. R. China. E-mail: zhengjunling@cjl.u.edu.cn / zjlshuxue@163.com.

<sup>2</sup> School of Mathematics and statistics, Guizhou University, Guiyang 550025, Guizhou Province, P. R. China. E-mail: liuyz@gzu.edu.cn / yzliu3@163.com .

† Corresponding author.

compute the representation dimension of artin algebras [18, 21]. Similar to the dimension of triangulated categories, the (extension) dimension of an abelian category was introduced by Beligiannis in [6], also see [11]. The size of the extension dimension reflects how far an artin algebra is from a finite representation type, some relate result can be see [25] and so on.

The spectrum of a triangulated category is an invariant of the category introduced by Orlov in 2009(see [20]). It can be seen as a generalization of the Rouquier dimension for triangulated categories. In 2012, Ballard, Favero, and Katzarkov systematically studied this invariant and named it the Orlov spectrum(see [5]). Similar to the Orlov spectrum of triangulated categories, we will define the Orlov spectrum for abelian categories, which can be regarded as a generalization of the extension dimension for abelian categories. In this paper, we will mainly investigate some properties of the Orlov spectrum of finite representation type algebras. The main theorems are as follows:

**Theorem 1.1.**

(1) (see Theorem 4.5) *Let  $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$  be an exact sequence in  $\text{mod } A$ . Then*

$$\max\{\ell^{t_S}(L), \ell^{t_S}(N)\} \leq \ell^{t_S}(M) \leq \ell^{t_S}(L) + \ell^{t_S}(N),$$

where  $\ell^{t_S}$  is  $t_S$ -radical layer length (see Definition 4.2). In particular, if  $\ell^{t_S}(L) = 0$ , then  $\ell^{t_S}(N) = \ell^{t_S}(M)$ ; if  $\ell^{t_S}(N) = 0$ , then  $\ell^{t_S}(L) = \ell^{t_S}(M)$ .

(2) (see Theorem 4.14) *If  $A$  is representation-finite, then*

$$\left\{ \left\lceil \frac{\ell^{t_S}(A)}{d} \right\rceil - 1 \mid d \in \mathbb{N} \text{ and } 1 \leq d < \ell^{t_S}(A) \right\} \subseteq \text{OSpec}(A),$$

where  $\lceil \alpha \rceil$  is the least integer greater than  $\alpha$ .

Furthermore, we obtain the following result by using Theorem 1.1, cf. Propositions 5.1 and 5.4.

**Theorem 1.2** (Theorem 5.7). *Let  $\vec{\mathbb{A}}_n$  be the linearly oriented quiver with type  $\mathbb{A}_n$ , i.e.,*

$$\vec{\mathbb{A}}_n = 1 \xrightarrow{a_1} 2 \xrightarrow{a_2} \dots \xrightarrow{a_{n-1}} n.$$

*Then the Orlov spectrum  $\text{OSpec}(A_n)$  of  $A_n := \mathbf{k}\vec{\mathbb{A}}_n$  is  $\{0, 1, 2, \dots, n-1\}$ .*

## 2. PRELIMINARIES

Let  $\mathcal{A}$  be an Abelian category. All subcategories of  $\mathcal{A}$  are full, additive and closed under isomorphisms and all functors between categories are additive. For a subclass  $\mathcal{U}$  of  $\mathcal{A}$ , we use  $\text{add}\mathcal{U}$  to denote the subcategory of  $\mathcal{A}$  consisting of direct summands of finite direct sums of objects in  $\mathcal{U}$ . For two objects  $X$  and  $Y$  in  $\mathcal{A}$ , we denote by  $X \leq Y$  if  $X$  is a subobject of  $Y$ , and denote by  $X \leq_{\oplus} Y$  if  $X$  is a direct summand of  $Y$ . Let  $\mathbb{N}$  be the set of natural numbers  $\{0, 1, 2, \dots\}$ , and  $\mathbb{N}^+$  be the set of positive integers  $\{1, 2, 3, \dots\}$ .

**2.1. The extension dimension of Abelian category.** Let  $\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n$  be subcategories of  $\mathcal{A}$ . Define

$$\mathcal{U}_1 \cdot \mathcal{U}_2 := \text{add}\{X \in \mathcal{A} \mid \text{there exists an sequence } 0 \rightarrow U_1 \rightarrow X \rightarrow U_2 \rightarrow 0 \text{ in } \mathcal{A} \text{ with } U_1 \in \mathcal{U}_1 \text{ and } U_2 \in \mathcal{U}_2\}.$$

For any subcategories  $\mathcal{U}, \mathcal{V}$  and  $\mathcal{W}$  of  $\mathcal{A}$ , by [11, Proposition 2.2] we have

$$(\mathcal{U} \cdot \mathcal{V}) \cdot \mathcal{W} = \mathcal{U} \cdot (\mathcal{V} \cdot \mathcal{W}).$$

Inductively, we can define

$$\mathcal{U}_1 \cdot \mathcal{U}_2 \cdot \dots \cdot \mathcal{U}_n := \text{add}\{X \in \mathcal{A} \mid \text{there exists an sequence } 0 \rightarrow U \rightarrow X \rightarrow V \rightarrow 0 \text{ in } \mathcal{A} \text{ with } U \in \mathcal{U}_1 \text{ and } V \in \mathcal{U}_2 \cdot \dots \cdot \mathcal{U}_n\}.$$

For a subcategory  $\mathcal{U}$  of  $\mathcal{A}$ , set  $[\mathcal{U}]_0 = 0$ ,  $[\mathcal{U}]_1 = \text{add}\mathcal{U}$ ,  $[\mathcal{U}]_n = [\mathcal{U}]_1 \cdot [\mathcal{U}]_{n-1}$  for each  $n \geq 2$ , and  $[\mathcal{U}]_{\infty} = \bigcup_{n \geq 0} [\mathcal{U}]_n$ , cf. [6].

The following two lemmas are useful for this paper.

**Lemma 2.1** ([25, Lemma 2.3]). *Let  $\mathcal{A}$  be an abelian category and  $T_1, T_2 \in \mathcal{A}$  and  $m, n \in \mathbb{N}$ . Then*

- (1)  $[T_1]_m \cdot [T_2]_n \subseteq [T_1 \oplus T_2]_{m+n}$ ;
- (2)  $[T_1]_m \oplus [T_2]_n \subseteq [T_1 \oplus T_2]_{\max\{m,n\}}$ .

**Lemma 2.2** ([25, Lemma 2.4]). *Let  $F : \mathcal{A} \rightarrow \mathcal{B}$  be an exact functor of abelian categories. Then  $F([T]_n) \subseteq [F(T)]_n$  for any  $T \in \mathcal{A}$  and positive integer  $n \geq 1$ .*

**Definition 2.3.** Let  $X$  be an object in  $\mathcal{A}$ . The generation time of  $X$ , denoted  $\Theta_{\mathcal{A}}(X)$  is

$$\Theta_{\mathcal{A}}(X) := \begin{cases} \inf\{n \in \mathbb{N} \mid \mathcal{A} = [X]_{n+1}\}, & \text{if } [X]_t = \mathcal{A} \text{ for some } t \in \mathbb{N}; \\ +\infty, & \text{otherwise.} \end{cases}$$

The object  $X$  is called a *strong generator* of  $\mathcal{A}$  if  $\Theta_{\mathcal{A}}(X)$  is finite.

**Lemma 2.4.** *For an object  $X$  in  $\mathcal{A}$  with  $\Theta_{\mathcal{A}}(X) < +\infty$ , we have*

$$\Theta_{\mathcal{A}}(X) \geq \Theta_{\mathcal{A}}(X \oplus Y)$$

*holds for arbitrary object  $Y$  in  $\mathcal{A}$ .*

*Proof.* By add  $X = [X]_1 \subseteq [X \oplus Y]_1 = \text{add}(X \oplus Y)$  and induction, we have

$$(2.1) \quad [X]_{n+1} \subseteq [X \oplus Y]_{n+1}$$

since, for any integer  $m \geq 0$ ,  $[X]_m \subseteq [X \oplus Y]_m$  yields

$$[X]_{m+1} = [X]_1 \cdot [X]_m \subseteq [X \oplus Y]_1 \cdot [X \oplus Y]_m = [X \oplus Y]_{m+1}.$$

Furthermore, (2.1) follows

$$\Theta_{\mathcal{A}}(X) = \inf\{n \in \mathbb{N} \mid [X]_{n+1} = \mathcal{A}\} \geq \inf\{n \in \mathbb{N} \mid [X \oplus Y]_{n+1} = \mathcal{A}\} = \Theta_{\mathcal{A}}(X \oplus Y)$$

by Definition 2.3 in the case for  $\Theta_{\mathcal{A}}(X) < +\infty$ .  $\square$

**Definition 2.5.** The *Orlov spectrum*  $\text{OSpec}(\mathcal{A})$  of an abelian category  $\mathcal{A}$  is defined as

$$\text{OSpec}(\mathcal{A}) = \{\Theta_{\mathcal{A}}(X) \mid X \text{ is a strong generator of } \mathcal{A}\}.$$

The *extension dimension* ([6]) of  $\mathcal{A}$  is defined as the infimum of  $\text{OSpec}(\mathcal{A})$ , that is,

$$\text{ext.dim } \mathcal{A} := \inf \text{OSpec}(\mathcal{A}).$$

Dually, we can define the *ultimate dimension* of  $\mathcal{A}$  is defined as the supremum of  $\text{OSpec}(\mathcal{A})$ , that is,

$$\text{u.dim } \mathcal{A} := \sup \text{OSpec}(\mathcal{A}).$$

**Remark 2.6.** For artin algebra  $\Lambda$ , we set  $\text{OSpec}(\Lambda) := \text{OSpec}(\text{mod } \Lambda)$ , and  $\text{u.dim}(\text{mod } \Lambda) := \text{u.dim}(\text{mod } \Lambda)$ .

**Lemma 2.7.** *If  $[T]_{n+1} = \mathcal{A} \neq [T]_n$ , then  $n \in \text{OSpec}(\mathcal{A})$ .*

*Proof.* Since  $[T]_1 \subseteq [T]_2 \subseteq \cdots \subseteq [T]_n \neq \mathcal{A}$  and  $[T]_{n+1} = \mathcal{A}$ , we have  $\Theta_{\mathcal{A}}(T) = n$ . It follows  $n \in \text{OSpec}(\mathcal{A})$  by the definition of Orlov spectrum (see Definition 2.5).  $\square$

**2.2. Ghosts and coghosts.** This section contains two parts as follows: in the first part we recall contravariantly finite subcategories and covariantly finite subcategories of Abelian categories; and in the second part we recall ghosts introduced in [6], and introduce coghosts in our paper by dual way.

2.2.1. *Contravariantly finite subcategories and covariantly finite subcategories.* Let  $\mathcal{A}$  be an Abelian category and  $\mathcal{C}$  be a subcategory of  $\mathcal{A}$  in this section. A map  $f : C \rightarrow X$  in  $\mathcal{A}$  with  $C \in \mathcal{C}$  is called a *right  $\mathcal{C}$ -approximation* of  $X$  if for every  $f' : C' \rightarrow X$  with  $C' \in \mathcal{C}$  there exists a map  $g : C' \rightarrow C$  such that the following diagram

$$\begin{array}{ccc} & & C' \\ & \swarrow g & \downarrow f' \\ C & \xrightarrow{f} & X \end{array}$$

commutes, that is,  $f' = f \circ g$ . If for every  $X \in \mathcal{A}$  admits a right  $\mathcal{C}$ -approximation, then  $\mathcal{C}$  is called *contravariantly finite*. Dually, a map  $f : X \rightarrow C$  in  $\mathcal{A}$  with  $C \in \mathcal{C}$  is called a *left  $\mathcal{C}$ -approximation* of  $X$  if for every  $f' : X \rightarrow C'$  with  $C' \in \mathcal{C}$  there exists a map  $g : C' \rightarrow C$  such that the following diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & C \\ f' \downarrow & \swarrow g & \\ C' & & \end{array}$$

commutes, that is,  $f' = f \circ g$ . If for every  $X \in \mathcal{A}$  admits a left  $\mathcal{C}$ -approximation, then  $\mathcal{C}$  is called *covariantly finite*.

2.2.2. *Ghost lemma and Coghost lemma.* Let  $T$  be an object in an Abelian category  $\mathcal{A}$ .

**Definition 2.8.** A morphism  $f : Y \rightarrow A$  (resp.  $f : A \rightarrow Y$ ) in  $\mathcal{A}$  is called a  *$T$ -ghost* (resp.  *$T$ -coghost*) if the induced map  $\text{Hom}_{\mathcal{A}}(T', f)$  (resp.  $\text{Hom}_{\mathcal{A}}(f, T')$ ) is zero for each  $T' \in \text{add } T$ . Furthermore, if a map can be written a composition of  $n$   $T$ -coghost maps, then it is called an  *$n$ -fold ghost* (resp.  *$n$ -fold coghost*).

**Notation 2.9.** For a set  $S$  of some objects lying in an Abelian category  $\mathcal{A}$ , we use  $\langle S \rangle$  be the minimal full subcategory containing  $S$ . For any  $n \in \mathbb{N}^+$ , we set

- (1)  $\text{Gh}_T^n(X, Y) := \{f \in \text{Hom}_{\mathcal{A}}(X, Y) \mid f \text{ is an } n\text{-fold } T\text{-coghost map}\}$ ,
- (2)  $\text{Gh}_T^n(-, Y) := \langle \text{Gh}_T^n(X, Y) \mid X \in \text{Obj}(\mathcal{A}) \rangle$ ,
- (3) and  $\text{Gh}_T^n(X, -) := \langle \text{Gh}_T^n(X, Y) \mid Y \in \text{Obj}(\mathcal{A}) \rangle$ .

**Remark 2.10.** It is easy to see that  $\text{Gh}_T^n(X, Y)$  is an ideals of  $\mathcal{A}$ , that is, for any  $T$ -coghost  $f : X \rightarrow Y$  and arbitrary homomorphism  $g : Y \rightarrow Z$  (resp.,  $g : Z \rightarrow X$ ), the composition  $g \circ f$  (resp.,  $f \circ g$ ) is also a  $T$ -coghost.

Recall that the *kernel* of a morphism  $f : X \rightarrow Y$  in an Abelian category  $\mathcal{A}$  is the morphism  $\eta : K \rightarrow X$  from an object  $K$  in  $\mathcal{A}$  to  $X$  such that

- (1)  $f \circ \eta = 0$ ;
- (2) and, for any morphism  $\eta' : K' \rightarrow X$  satisfying  $f \circ \eta' = 0$ , there is a unique morphism  $\tau : K' \rightarrow K$  such that  $\eta' = \eta \circ \tau$ .

One can check that  $K$  is unique up to isomorphism. In most cases,  $K$  is written as  $\text{Ker } f$ .

**Lemma 2.11.** *The kernel of any left  $\mathcal{C}$ -approximation in  $\mathcal{A}$  is a  $\mathcal{C}$ -coghost.*

*Proof.* Assume that  $f : X \rightarrow T$  is a left  $\mathcal{C}$ -approximation and  $\eta : \text{Ker } f \rightarrow X$  its kernel. Here,  $T$  is an object lying in  $\mathcal{C}$ . For each  $h \in \text{Hom}_{\mathcal{A}}(X, T')$ , there exists a map  $f' : T' \rightarrow T$  such that  $T' \in \mathcal{C}$  and  $h = f' \circ f$  since  $f$  is a left  $\mathcal{C}$ -approximation. Thus, we obtain the following diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker } f & \xrightarrow{\eta} & X & \xrightarrow{f} & T \\ & & \downarrow h \circ \eta & \swarrow h & \searrow f' & & \\ & & T' & & & & \end{array}$$

commutes. Then

$$h \circ \eta = (f' \circ f) \circ \eta = f' \circ (f \circ \eta) = f' \circ (f \circ \eta) = f' \circ 0 = 0.$$

It follows that  $\eta : \text{Ker } f \rightarrow X$  is a  $\mathcal{C}$ -coghost by the definition of coghost.  $\square$

Let  $\text{Fac}(T)$  (resp.  $\text{Sub}(T)$ ) be the subcategory of  $\mathcal{A}$  whose object is isomorphic to a quotient (resp. subobject) of some direct sum  $T^{\oplus I}$ , where  $I$  is a finite index set. The following lemma, i.e., Ghost Lemma, is established by Beligiannis in [6].

**Lemma 2.12** (Ghost Lemma [6, Lemma 1.3]).

- (a) If  $X \in [\text{Fac}(T)]_n$ , then  $\text{Gh}_T^n(X, -) = 0$ .
- (b) If  $\text{add } T$  is contravariantly finite in  $\mathcal{A}$ , then the following are equivalent:
  - (1)  $\text{Gh}_T^n(X, -) = 0$ ;
  - (2)  $X \in [\text{Fac}(T)]_n$ .

Next, we provide a dual result, say Coghost Lemma in our paper, for Ghost Lemma.

**Lemma 2.13** (Coghost Lemma).

- (a) If  $Y \in [\text{Sub}(T)]_n$ , then  $\text{Gh}_T^n(-, Y) = 0$ .
- (b) If  $\text{add } T$  is covariantly finite in  $\mathcal{A}$ , then the following are equivalent:
  - (1)  $\text{Gh}_T^n(-, Y) = 0$ ;
  - (2)  $Y \in [\text{Sub}(T)]_n$ .

*Proof.* (a) If  $n = 1$ . Let  $f : X \rightarrow Y$  be a  $T$ -coghost. Since  $Y \in [\text{Sub}(T)]_1$ , there is a monomorphism  $g : Y \rightarrow T'$  such that  $T' \in \text{add } T$ . And, by the definition of coghost and  $T' \in \text{add } T$ , we have  $g \circ f = 0$  since  $f : X \rightarrow Y$  is a  $T$ -coghost. Thus,  $f = 0$  by using  $g$  to be a monomorphism.

Next, assume that, for any  $n \leq k$ ,  $Y \in [\text{Sub}(T)]_k$  yields that  $\text{Gh}_T^k(-, Y) = 0$ . Considering the case of  $n = k + 1$ . we have, for any  $Y \in [\text{Sub}(T)]_{k+1}$ , the following short exact sequence

$$0 \longrightarrow Y_1 \xrightarrow{g_1} Y \xrightarrow{g_2} Y_2 \longrightarrow 0$$

with  $Y_1 \in [\text{Sub}(T)]_1$  and  $Y_2 \in [\text{Sub}(T)]_k$ , and there is a monomorphism  $h_1 : Y_1 \rightarrow T_1$  such that  $T_1 \in \text{add } T$ . Let  $f \in \text{Gh}_T^{k+1}(-, Y)$ . Then we have  $f = f_1 \circ f_2 \circ \cdots \circ f_k \circ f_{k+1}$ , where each  $f_i : X_i \rightarrow X_{i-1}$  is a  $T$ -coghost (we set  $X_0 := Y$  in this proof). By assumption and  $Y_2 \in [\text{Sub}(T)]_k$ , we have  $\text{Gh}_T^k(-, Y_2) = 0$ . Since  $f_1$  is  $T$ -coghost, we obtain that  $g_2 \circ f_1$  is also a  $T$ -coghost by Remark 2.10. Then

$$g_2 \circ (f_1 \circ f_2 \circ \cdots \circ f_k) = (g_2 \circ f_1) \circ f_2 \circ \cdots \circ f_k \in \text{Gh}_T^k(-, Y_2).$$

Moreover,  $g_2 \circ (f_1 \circ f_2 \circ \cdots \circ f_k) = 0$ . Then, since  $g_1$  is the kernel of  $g_2$ , there exists a unique morphism  $h : X_k \rightarrow Y_1$  such that

$$f_1 \circ f_2 \circ \cdots \circ f_k = g_1 \circ h,$$

that is, we obtain the following diagram

$$\begin{array}{ccccccc}
 & & T_1 & & & & \\
 & & \uparrow h_1 & & & & \\
 0 & \longrightarrow & Y_1 & \xrightarrow{g_1} & Y & \xrightarrow{g_2} & Y_2 \longrightarrow 0 \\
 & & \uparrow & & \uparrow f_1 & & \\
 & & & & X_1 & & \\
 & & & & \uparrow f_2 & & \\
 & & & & \vdots & & \\
 & & & & \uparrow f_k & & \\
 & & & & X_k & & \\
 & & & & \uparrow f_{k+1} & & \\
 & & & & X_{k+1} & & \\
 & & \text{---} h \text{---} & & \text{---} f \text{---} & & 
 \end{array}$$

commutes. Thus,  $h_1 \circ (h \circ f_{k+1}) = (h_1 \circ h) \circ f_{k+1} = 0$  by using  $f_{k+1}$  to be a  $T$ -coghost. It follows that  $h \circ f_{k+1} = 0$ . Then  $f = f_1 \circ \cdots \circ f_{k+1} = g_1 \circ h \circ f_{k+1} = g_1 \circ 0 = 0$ , i.e.,  $\text{Gh}_T^{k+1}(-, Y) = 0$ .

(b) It is trivial that (2) implies (1) by (a). Next, we prove (1) implies (2).

First of all, we show that (1) implies (2) in the case of  $n = 1$ . Assume that  $Y$  is an object satisfying  $\text{Gh}_T^1(-, Y) = 0$ , and  $f : Y \rightarrow T_1$  is a left add  $T$ -approximation of  $Y$ . Then for each map  $h : Y \rightarrow T'_1$  with  $T'_1 \in \text{add } T$ , there is a morphism  $g : T_1 \rightarrow T'_1$  such that the following diagram

$$\begin{array}{ccc}
 \text{Ker } f & \xrightarrow{f_1} & Y & \xrightarrow{f} & T_1 \\
 & & \downarrow h & \searrow g & \\
 & & T'_1 & & 
 \end{array}$$

commutes. Thus,  $h \circ f_1 = (g \circ f) \circ f_1 = g \circ (f \circ f_1) = g \circ 0 = 0$ . It follows that  $f_1$  is a  $T$ -coghost. Furthermore,  $\text{Gh}_T^1(-, Y) = 0$  yields that  $f_1$ , as a morphism in  $\text{Gh}_T^1(\text{Ker } f, Y)$ , is zero. Thus,  $\text{Ker } f = 0$ . Therefore,  $f$  is monomorphic, i.e.,  $Y \in [\text{Sub}(T)]_1$  as required.

Next, suppose that  $\text{Gh}_T^k(-, Y) = 0$ . Considering the following family of exact sequences

$$(2.2) \quad (0 \longrightarrow Y_{i+1} \xrightarrow{f_{i+1}} Y_i \xrightarrow{g_i} T_i)_{0 \leq i < k},$$

where, for all  $i$ ,  $g_i$  is a left add  $T$ -approximation of  $Y_i$ ;  $Y_{i+1} := \text{Ker } g_i$ ; and  $Y_0 := Y$ , we know that  $f_1, \dots, f_{k+1}$  are  $T$ -coghosts by Lemma 2.11. Then we obtain

$$f_1 \circ f_2 \circ \cdots \circ f_{k-1} \circ f_k = 0$$

by  $\text{Gh}_T^k(-, Y) = 0$ , and it follows  $f_k = 0$  by using all  $f_i$  ( $0 \leq i \leq k-1$ ) to be monomorphisms. On the other hand, (2.2) induces two family exact sequences

$$\begin{aligned}
 (0 \longrightarrow Y_{i+1} \xrightarrow{f_{i+1}} Y_i \longrightarrow \text{Im } g_i \longrightarrow 0)_{0 \leq i < k-1} \\
 \text{and } (0 \longrightarrow \text{Im } g_i \longrightarrow T_i)_{0 \leq i < k-1}
 \end{aligned}$$

by the canonical decomposition of  $g_i$ . It is easy to see that  $\text{Im } g_i \in [\text{Sub}(T_i)]_1 \subseteq [\text{Sub}(T)]_1$  holds for each  $0 \leq i \leq k-1$ , and then we have

$$\begin{aligned}
 Y &= Y_0 \in [Y_1]_1 \cdot [\text{Im } g_0]_1 \\
 &\subseteq [Y_2]_1 \cdot [\text{Im } g_1]_1 \cdot [\text{Im } g_0]_1 \\
 &\subseteq [Y_3]_1 \cdot [\text{Im } g_2]_1 \cdot [\text{Im } g_1]_1 \cdot [\text{Im } g_0]_1
 \end{aligned}$$

$$\begin{aligned}
& \vdots \\
& \subseteq [Y_n]_1 \cdot [g_{n-1}]_1 \cdots \cdot [\text{Im } g_2]_1 \cdot [\text{Im } g_1]_1 \cdot [\text{Im } g_0]_1 \\
& \subseteq [0]_1 \cdot [\text{Sub}(T)]_1 \cdots \cdot [\text{Sub}(T)]_1 \cdot [\text{Sub}(T)]_1 \cdot [\text{Sub}(T)]_1 \\
& = [\text{Sub}(T)]_n.
\end{aligned}$$

Therefore, (1) implies (2) by induction.  $\square$

**Remark 2.14.** If  $\mathcal{U}$  is a subcategory of  $\mathcal{V}$ , then  $[\mathcal{U}]_n \subseteq [\mathcal{V}]_n$  holds for all  $n \in \mathbb{N}$ , see [25, Proposition 2.2(2)]. Immediately, for each  $T \in \mathcal{A}$ , we have  $[T]_n \subseteq [\text{Fac}(T)]_n$  and  $[T]_n \subseteq [\text{Sub}(T)]_n$ .

**2.3. Coghosts on Artin algebra.** Let  $\Lambda$  be an Artin  $R$ -algebra defined over Artin ring  $R$  in this section. Then the category  $\text{mod } \Lambda$  of finitely generated right  $\Lambda$ -module (say  $\Lambda$ -module for short) is an Abelian category. Now, we provide some properties of coghosts and strong generator in  $\text{mod } \Lambda$  which are dual of the properties of ghost.

**2.3.1. Some properties of coghost on Artin algebra.** Let  $T$  be a finitely generated  $\Lambda$ -module. The following Lemma is well-known.

**Lemma 2.15.** *The subcategory  $\text{add } T$  of  $\text{mod } \Lambda$  is a covariantly finite subcategory.*

We will prove Proposition 2.16 through the above lemma. Thus, for the convenience of readers, we still provide proof of the above lemma.

*Proof.* First of all, for each  $M \in \text{mod } \Lambda$ ,  $\text{Hom}_\Lambda(M, T)$  can be seen as a finitely generated  $R$ -module defined by

$$\text{Hom}_\Lambda(M, T) \times R \rightarrow \text{Hom}_\Lambda(M, T), (h, r) \mapsto h \cdot r,$$

where  $h \cdot r$  is the map  $M \rightarrow T$  sending each  $m \in M$  to  $f(m)r$ , see [4]. Then  $\text{Hom}_\Lambda(M, T)$  containing a family of right  $R$ -homomorphism  $f_1, \dots, f_n$  which form a set of generators for  $\text{Hom}_\Lambda(M, T)$ .

Next, take an arbitrary map  $g : M \rightarrow N$  with  $N \in \text{add } T$ . Notice that there is a positive integer  $m$  such that  $N$  is isomorphic to a direct summand of  $T^{\oplus m}$  (written as  $N \leq_{\oplus} T^{\oplus m}$  for simplification), we can find two  $\Lambda$ -homomorphisms

$$\eta = \begin{pmatrix} \eta_1 \\ \vdots \\ \eta_m \end{pmatrix} : N \rightarrow T^{\oplus m} \text{ and } \theta = (\theta_1 \cdots \theta_m) : T^{\oplus m} \rightarrow N$$

such that  $\theta \circ \eta = \sum_{i=1}^m \theta_i \eta_i = 1_N$ . Here,  $\eta_1, \dots, \eta_m$  are  $R$ -homomorphisms in  $\text{Hom}_\Lambda(N, T)$ , and  $\theta_1, \dots, \theta_m$  are  $R$ -homomorphisms in  $\text{Hom}_\Lambda(T, N)$ . Thus, for the  $\Lambda$ -homomorphism  $\eta_i \circ g \in \text{Hom}_\Lambda(M, T)$ , there are  $n$  elements  $b_{1j}, \dots, b_{nj}$  such that  $\eta_i \circ g = \sum_{j=1}^n b_{ij} f_j$ . It follows that

$$\eta \circ g = \begin{pmatrix} \eta_1 \circ g \\ \vdots \\ \eta_m \circ g \end{pmatrix} = \begin{pmatrix} \sum_{j=1}^n b_{1j} f_j \\ \vdots \\ \sum_{j=1}^n b_{mj} f_j \end{pmatrix} = \begin{pmatrix} b_{11} & \cdots & b_{1n} \\ \vdots & & \vdots \\ b_{m1} & \cdots & b_{mn} \end{pmatrix} \circ \begin{pmatrix} f_1 \\ \vdots \\ f_n \end{pmatrix}.$$

The right of the above equation is written as  $B \circ f$  for simplification in this proof. Then we obtain

$$g = 1_N \circ g = (\theta \circ \eta) \circ g = \theta \circ (\eta \circ g) = \theta \circ (B \circ f) = (\theta \circ B) \circ f.$$

Thus,  $g$  factors through  $f$ , see the following diagram

$$\begin{array}{ccc}
M & \xrightarrow{f} & T^{\oplus n} \\
\downarrow g & \nearrow \theta \circ B & \\
N & & \\
\downarrow \eta & \nearrow B & \\
T^{\oplus m} & & 
\end{array}$$

$\theta$  (curved arrow from  $T^{\oplus m}$  to  $N$ ),  $\leq_{\oplus}$  (arrow from  $N$  to  $T^{\oplus m}$ ),  $\spadesuit$  (triangle with vertices  $M, N, T^{\oplus n}$ ),  $\clubsuit$  (triangle with vertices  $N, T^{\oplus m}, T^{\oplus n}$ )

whose triangles  $\spadesuit$  and  $\clubsuit$  are commute. Therefore, we get a  $\Lambda$ -homomorphism  $f$  which is a left add  $T$ -approximation of  $M$ . Thus, add  $T$  is covariantly finite.  $\square$

**Proposition 2.16.** *For a chain of add  $T$ -coghosts*

$$Y_n \xrightarrow{f_n} Y_{n-1} \xrightarrow{f_{n-1}} \cdots \xrightarrow{f_3} Y_2 \xrightarrow{f_2} Y_1 \xrightarrow{f_1} Y_0$$

in  $\text{mod } \Lambda$  satisfying  $f_1 \circ f_2 \circ \cdots \circ f_n \neq 0$ , we have  $Y_0 \notin [T]_n$ .

*Proof.* Notice that the composition  $f_1 \circ f_2 \circ \cdots \circ f_n \in \text{Gh}_T^n(Y_n, Y_0)$  is a non-zero  $\Lambda$ -homomorphism, we obtain  $\text{Gh}_T^n(Y_n, Y_0) \neq 0$ . Then, by Lemma 2.15 and Lemma 2.13 (b),  $Y_0 \notin [\text{Sub}T]_n$  holds. Thus,  $Y_0 \notin [T]_n$  since  $[T]_n \subseteq [\text{Sub}T]_n$  by Lemma 2.14.  $\square$

**2.3.2. Some properties of strong generator over Artin algebra.** In this subdivision, we provide some properties for strong generator in the category  $\text{mod } \Lambda$  of finitely generated  $\Lambda$ -module over an Artin algebra  $\Lambda$ .

**Proposition 2.17.** *If  $T$  is a strong generator of  $\text{mod } \Lambda$ , then any automorphism  $f : X \rightarrow X$  defined on non-zero  $\Lambda$ -module  $X$  is not a  $T$ -coghost.*

*Proof.* Assume  $\text{mod } \Lambda = [T]_n$ ,  $n \in \mathbb{N}^+$ . If  $f$  is both an isomorphism and a  $T$ -coghost, then  $f^n$  is an isomorphism which is not zero. In these cases,  $X \notin [T]_n = \text{mod } \Lambda$  by Property 2.16, a contradiction.  $\square$

**Proposition 2.18.** *Given a map  $f : M_1 \rightarrow M_2$  in  $\text{mod } \Lambda$ . If for each indecomposable module  $N \in \text{add } T$ , either  $\text{Hom}_{\Lambda}(M_1, N) = 0$  or  $\text{Hom}_{\Lambda}(M_2, N) = 0$ , then  $f$  is a  $T$ -coghost.*

*Proof.* Let  $N' = \bigoplus_{k=1}^t N_k \in \text{add } T$  be an arbitrary module, where all direct summands  $N_k$  of  $N$  are indecomposable. Then for any homomorphism  $g = \begin{pmatrix} g_1 \\ \vdots \\ g_t \end{pmatrix} : M_2 \rightarrow N'$ , where  $g_k \in \text{Hom}_{\Lambda}(M_2, N_k)$  for each  $1 \leq k \leq t$ , we have

$$g \circ f = \begin{pmatrix} g_1 \circ f \\ \vdots \\ g_t \circ f \end{pmatrix} \in \text{Hom}_{\Lambda}(M_1, N'),$$

where  $g_k \circ f \in \text{Hom}_{\Lambda}(M_1, N_k)$ . Since either  $\text{Hom}_{\Lambda}(M_1, N_k) = 0$  or  $\text{Hom}_{\Lambda}(M_2, N_k) = 0$ , we obtain that  $g_k \circ f = 0$  for each  $1 \leq k \leq t$ , and so  $g \circ f = 0$ .  $\square$

**Proposition 2.19.** *Given a map  $f : M_1 \rightarrow M_2$  in  $\text{mod } \Lambda$ . If  $f$  is both  $T_1$ -coghost and  $T_2$ -coghost, then  $f$  is a  $(T_1 \oplus T_2)$ -coghost.*

*Proof.* For any morphism  $g = \begin{pmatrix} g_1 \\ g_2 \end{pmatrix} \in \text{Hom}_{\Lambda}(M_2, T'_1 \oplus T'_2)$ , where  $T'_1 \in \text{add } T_1$  and  $T'_2 \in \text{add } T_2$ , we have

$$g \circ f = \begin{pmatrix} g_1 \\ g_2 \end{pmatrix} \circ f = \begin{pmatrix} g_1 \circ f \\ g_2 \circ f \end{pmatrix}.$$

Since  $f$  is both  $T_1$ -coghost and  $T_2$ -coghost, we obtain  $g_1 \circ f = 0$  and  $g_2 \circ f = 0$ . Then  $g \circ f = 0$ , i.e.,  $f$  is a  $(T_1 \oplus T_2)$ -coghost.  $\square$

The following lemma is well-known. We will use it to prove Proposition 2.21.



**Lemma 2.20.** *Let  $M$  and  $N$  be two  $\Lambda$ -modules in  $\text{mod}\Lambda$ .*

- (1) *If  $N$  is semi-simple, then  $\text{Hom}_\Lambda(N, M) = \text{Hom}_\Lambda(N, \text{soc } M)$ .*
- (2) *If  $M$  is semi-simple, then  $\text{Hom}_\Lambda(N, M) = \text{Hom}_\Lambda(\text{top } N, M)$ .*

**Proposition 2.21.** *If  $T$  is a strong generator, then  $\text{soc } T \cong \text{top } T \cong \Lambda / \text{rad } \Lambda$ .*

*Proof.* We only show that  $\text{soc } T \cong \Lambda / \text{rad } \Lambda$ , the proof of  $\text{top } T \cong \Lambda / \text{rad } \Lambda$  is similar by using Ghost Lemma 2.12.

First of all, we show that each simple  $\Lambda$ -modules is a direct summand of  $\text{soc } T$ . Otherwise, there is a simple module  $S \in \text{mod}\Lambda$  such that  $\text{Hom}_\Lambda(S, T') = 0$  holds for all  $T' \in \text{add } T$ . Then, clearly, any non-zero homomorphism  $f : S \rightarrow S$  is a  $T$ -coghost by Proposition 2.18. It contradicts with Proposition 2.17 because, by Schur's lemma,  $f$  is an automorphism. Thus,  $S \leq_{\oplus} \text{soc } T$  holds for each simple module  $S$ . It follows that  $\Lambda / \text{rad } \Lambda \cong \bigoplus_S S \leq_{\oplus} \text{soc } T$  as required.

On the other hand,  $\text{soc } T \leq_{\oplus} \bigoplus_S S$  is trivial. Thus  $\text{soc } T \cong \Lambda / \text{rad } \Lambda$ .  $\square$

**Lemma 2.22.** *If  $T$  is a strong generator, then  $\text{Hom}_\Lambda(Y, T) \neq 0$  and  $\text{Hom}_\Lambda(T, Y) \neq 0$  hold for all  $0 \neq Y \in \text{mod}\Lambda$ .*

*Proof.* Since  $f_1 : Y \rightarrow \text{top } Y$ ,  $f_2 : \text{top } Y \rightarrow \text{soc } T = \Lambda / \text{rad } \Lambda$ , and  $f_3 : \text{soc } T \rightarrow T$ , we have  $f_3 \circ f_2 \circ f_1 \neq 0$ . Then  $\text{Hom}_\Lambda(Y, T) \neq 0$ .  $\square$

**Proposition 2.23.** *If  $T$  is a strong generator of  $\text{mod}\Lambda$ , then, for arbitrary projective or injection simple modules  $S_1, \dots, S_t$ , the direct sum  $\bigoplus_{i=1}^t S_i$  lies in  $[T]_1$ .*

*Proof.* By Lemma 2.22, there is  $\Lambda$ -homomorphism  $f : S \rightarrow T$  for any simple module  $S$  such that  $f \neq 0$ . Thus,  $f$  is monomorphic by Schur's lemma. If  $S$  is projective, then  $f$  is a section, i.e.,  $f \simeq \begin{pmatrix} 1_S \\ 0 \end{pmatrix} : S \rightarrow S \oplus T' (\cong T)$ . It follows that  $S \leq_{\oplus} T$ . We can show that  $S \leq_{\oplus} T$  holds in the case for  $S$  to be injective by dual method. Thus,  $\bigoplus_{i=1}^t S_i \in \text{add } T = [T]_1$  since  $\text{add } T$  is closed under direct sum.  $\square$

**Proposition 2.24.** *If  $T$  is a strong generator in  $\text{mod}\Lambda$ , then, for any  $T$ -coghost  $f : X \rightarrow Y$  with  $Y$  indecomposable, we have  $f \in \text{rad}_A(X, Y)$ .*

*Proof.* If  $f \notin \text{rad}_A(X, Y) = \{f \in \text{Hom}_A(X, Y) \mid 1_Y - f \circ g \text{ is an isomorphism for all } g \in \text{Hom}_A(Y, X)\}$ , then there exists a morphism  $g \in \text{Hom}_A(Y, X)$  such that  $1_Y - fg$  is not an isomorphism. Since  $f$  is a  $T$ -coghost, we have  $h \circ f = 0$  for all  $h \in \text{Hom}_A(Y, T')$  with  $T' \in \text{add } T$ . Then

$$h \circ (1_Y - f \circ g) = h \circ 1_Y - h \circ f \circ g = h - 0 \circ g = h.$$

Thus, for each  $n \in \mathbb{N}^+$ , the following equation

$$\begin{aligned} h \circ (1_Y - f \circ g)^n &= h \circ (1_Y - f \circ g) \circ (1_Y - f \circ g)^{n-1} \\ &= h \circ (1_Y - f \circ g)^{n-1} \\ &\quad \vdots \\ &= h \circ (1_Y - f \circ g) \\ &= h \end{aligned}$$

holds. Since  $1_Y - f \circ g$  is not an isomorphism and  $Y$  is indecomposable, we get  $(1_Y - f \circ g)^m = 0$  for some  $m \in \mathbb{N}^+$ . And then we have  $h = h \circ (1_Y - f \circ g)^m = h \circ 0 = 0$ . Thus,  $\text{Hom}_A(Y, T') = 0$ . By Lemma 2.22, we have  $\text{Hom}_A(Y, T') \neq 0$ , this is a contradiction.  $\square$

### 3. LOEWY LENGTHS

The *Loewy length*  $\ell\ell(M)$  of a  $\Lambda$ -module  $M$  in  $\text{mod}\Lambda$  is defined as the length of the ascending chain

$$0 = \text{rad}^n M \leq \text{rad}^{n-1} M \leq \dots \leq \text{rad}^1 M \leq \text{rad}^0 M = M$$

where we use “ $X \leq Y$ ” to represent a submodule (resp. subobject)  $X$  of  $Y$  in  $\text{mod } \Lambda$  (resp.  $\mathcal{A}$ ), and, for each  $1 \leq i \leq n$ ,  $\text{rad}^i M$  is the submodule of  $M$  satisfying  $\text{rad}^i M = \text{rad}(\text{rad}^{i-1} M)$ . Obviously, the Loewy length of any semi-simple module is 1, and the Loewy length of a  $\Lambda$ -module  $M$  is zero if and only if  $M = 0$ .

**3.1. Properties of Loewy lengths.** We recall some well-known results for Loewy length in this subsection.

**Lemma 3.1.** *For arbitrary short exact sequence*

$$0 \longrightarrow X \xrightarrow{f} Y \xrightarrow{g} Z \longrightarrow 0$$

in  $\text{mod } \Lambda$ , we have

$$\max\{\ell(Y), \ell(Z)\} \leq \ell(Y) \leq \ell(X) + \ell(Z).$$

**Lemma 3.2.** *For each  $M \in [T]_m$ , we have  $\ell(M) \leq m \ell(T)$ .*

*Proof.* In the case of  $m = 1$ , it is trivial that if  $M \in [T]_1 = \text{add } T$  then  $\ell(M) \leq \ell(T)$ . Consider the induction hypothesis which admit that if  $m \leq t$  then  $\ell(M) \leq t \ell(T)$ . Then for  $m = t + 1$ , there is a short exact sequence

$$0 \longrightarrow A \longrightarrow M \longrightarrow B \longrightarrow 0$$

such that  $A \in [T]_1$  and  $B \in [T]_t$  hold for arbitrary  $M \in [T]_{t+1}$ . Then, by Lemma 3.1, we have

$$\ell(M) \leq \ell(A) + \ell(B) \leq 1 \ell(T) + t \ell(T) = (1 + t) \ell(T).$$

Thus, this lemma holds by induction.  $\square$

Notice that if  $T$  is a strong generator of  $\text{mod } \Lambda$  and the  $m$  is the positive integer satisfying  $[T]_m = \text{mod } \Lambda$ , then by Lemma 3.2, the following two statement hold.

- (1)  $\ell(\Lambda) \leq m \ell(T)$ ;
- (2) for each  $M \in [\Lambda / \text{rad } \Lambda]_m$ , we have  $\ell(M) \leq m \ell(\Lambda / \text{rad } \Lambda) = m$ .

The statement (1) holds by  $\Lambda \in [T]_m$ ; and the statement (2) can be shown in the case for  $T = \Lambda / \text{rad } \Lambda$ .

**Proposition 3.3.**  $\Theta_{\text{mod } \Lambda}(\Lambda / \text{rad } \Lambda) = \ell(\Lambda) - 1$ .

*Proof.* The case  $\ell(\Lambda) = 1$  is trivial. Now consider the case  $\ell(\Lambda) \geq 2$ . For each  $M \in \text{mod } \Lambda$ , the following family of short exact sequence

$$(0 \longrightarrow \text{rad}^t M \longrightarrow \text{rad}^{t-1} M \longrightarrow \text{rad}^{t-1} M / \text{rad}^t M \longrightarrow 0)_{1 \leq t \leq \ell(\Lambda) - 1}$$

satisfies  $\text{rad}^{t-1} M / \text{rad}^t M \in [\Lambda / \text{rad } \Lambda]_1$  for all  $1 \leq t \leq \ell(\Lambda) - 1$  since all  $\text{rad}^{t-1} M / \text{rad}^t M$  are semi-simple. Then  $[\Lambda / \text{rad } \Lambda]_{\ell(\Lambda)} = \text{mod } \Lambda$ . It follows that  $\Theta_{\text{mod } \Lambda}(\Lambda / \text{rad } \Lambda) \leq \ell(\Lambda) - 1$ .

On the other hand, if  $\Theta_{\text{mod } \Lambda}(\Lambda / \text{rad } \Lambda) \leq \ell(\Lambda) - 2$ , then  $\Lambda \in \text{mod } \Lambda = [\Lambda / \text{rad } \Lambda]_{\ell(\Lambda) - 1}$ . It follows  $\ell(\Lambda) \leq (\ell(\Lambda) - 1) \ell(\Lambda / \text{rad } \Lambda)$  by Lemma 3.2, a contradiction.  $\square$

**Corollary 3.4.** *The following two statements hold.*

- (1)  $\ell(\Lambda) - 1 \in \text{OSpec}(\Lambda)$ .
- (2)  $\text{u.dim}(\text{mod } \Lambda) \geq \ell(\Lambda) - 1$ .

*Proof.* By Proposition 3.3, we have  $\Theta_{\text{mod } \Lambda}(\Lambda / \text{rad } \Lambda) = \ell(\Lambda) - 1$  holds for arbitrary Artin algebra  $\Lambda$ . Then the statement (1) holds by Definition 2.5. Furthermore, the statement (2) is obtained by (1) and the definition of ultimate dimension ( see Definition 2.5) as following formula

$$\text{u.dim}(\text{mod } \Lambda) = \sup(\text{OSpec}(\Lambda)) \geq \ell(\Lambda) - 1$$

as required.  $\square$

**3.2. Strong generators.** Recall that the following well-known lemma which will be used to prove Lemma 3.6 in our paper.

**Lemma 3.5.** *For any short exact sequence*

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

in  $\text{mod } \Lambda$ , we have

$$\text{pd } B \leq \text{pd}(A \oplus C) \text{ and } \text{id } B \leq \text{id}(A \oplus C) \quad ^1$$

**Lemma 3.6.** *Let  $M$  and  $T$  be two  $\Lambda$ -modules such that  $M \in [T]_{n+1}$ . Then  $\text{pd } M \leq \text{pd } T$  and  $\text{id } M \leq \text{id } T$ .*

*Proof.* First of all, by  $M \in [T]_{n+1}$ , there is the following family of short exact sequences

$$(0 \longrightarrow X_i \longrightarrow Y_{i-1} \longrightarrow Z_i \longrightarrow 0)_{1 \leq i \leq n}$$

such that  $Z_{i-1} \leq_{\oplus} Y_{i-1}$  (in this case,  $Y_{i-1} \cong Z_{i-1} \oplus Z'_{i-1}$  for some  $Z'_{i-1}$ ), where  $Z_0 = M$ ;  $X_1, \dots, X_n \in [T]_1$ ; and  $Z_i \in [T]_{n-i+1}$ . Then, naturally, we have the following short exact sequences

$$0 \longrightarrow \left( \bigoplus_{t=1}^i X_t \right) \oplus X_{i+1} \longrightarrow \left( \bigoplus_{t=1}^i X_t \right) \oplus Y_i \longrightarrow Z_{i+1} \longrightarrow 0$$

for any  $1 \leq i \leq n-1$ . Then, by Lemma 3.5 and the above two families of short exact sequences, we have

$$\begin{aligned} \text{pd } M = \text{pd } Z_0 &\leq \text{pd } Y_0 \leq \text{pd}(X_1 \oplus Z_1) \\ &\leq \text{pd}(X_1 \oplus Z_1 \oplus Z'_1) = \text{pd}(X_1 \oplus Y_1) \\ &\leq \text{pd}(X_1 \oplus X_2 \oplus Z_2) \\ &\leq \text{pd}(X_1 \oplus X_2 \oplus Z_2 \oplus Z'_2) = \text{pd}(X_1 \oplus X_2 \oplus Y_2) \\ &\leq \text{pd}(X_1 \oplus X_2 \oplus X_3 \oplus Z_3) \\ &\leq \text{pd}(X_1 \oplus X_2 \oplus X_3 \oplus Z_3 \oplus Z'_3) = \text{pd}(X_1 \oplus X_2 \oplus X_3 \oplus Y_3) \\ &\vdots \\ &\leq \text{pd} \left( \left( \bigoplus_{t=1}^n X_t \right) \oplus Y_n \right) \spadesuit \leq \text{pd } T, \end{aligned}$$

where  $\spadesuit$  holds since all  $X_i$  and  $Y_n$  are  $\Lambda$ -modules lying in  $[T]_1 = \text{add } T$ . We can prove  $\text{id } M \leq \text{id } T$  by similar way.  $\square$

**Proposition 3.7.** *If  $T$  is a strong generator of  $\text{mod } \Lambda$ , then the global dimension  $\text{gl.dim } \Lambda$  of  $\Lambda$  and the projective dimension  $\text{pd } T$  of  $T$  coincide.*

*Proof.* Assume  $\text{mod } \Lambda = [T]_{n+1}$ . Then we have  $\text{pd}(\Lambda/\text{rad } \Lambda) \leq \text{pd } T$  by Lemma 3.6 since it is clear that  $\Lambda/\text{rad } \Lambda$  is a  $\Lambda$ -module in  $\text{mod } \Lambda = [T]_{n+1}$ . It follows that  $\text{gl.dim } \Lambda = \text{pd}(\Lambda/\text{rad } \Lambda) \leq \text{pd } T$ . On the other hand,  $\text{pd } T \leq \text{gl.dim } \Lambda$  by the definition of global dimension of the Artin algebra  $\Lambda$ . Thus,  $\text{pd } T = \text{gl.dim } \Lambda$  as required.  $\square$

**Corollary 3.8.** *Keep the notations from Proposition 3.7, then the following two statements hold.*

- (1) *There is an indecomposable module  $M$  with  $\text{pd } M = \text{gl.dim } \Lambda$  such that  $M \in [T]_1$ .*
- (2) *There is an indecomposable module  $N$  with  $\text{id } N = \text{gl.dim } \Lambda$  such that  $N \in [T]_1$ .*

*Proof.* We only prove (1), the proof of (2) is similar. Considering the decomposition  $T = \bigoplus_{i=1}^t T_i$  of  $T$ , where  $T_1, \dots, T_t$  are indecomposable, we obtain

$$\text{gl.dim } \Lambda = \text{pd } T = \max_{1 \leq i \leq t} \text{pd } T_i,$$

<sup>1</sup>For any module  $X$  in  $\text{mod } \Lambda$ , “ $\text{pd } X$ ” and “ $\text{id } X$ ” are projective dimension and injective dimension of  $X$ , respectively.

that is, there is an index  $i$  such that  $\text{pd } T = \text{pd } T_i$  as required.  $\square$

Next we provide an instance for Corollary 3.8.

**Example 3.9.** Let  $\Lambda = \mathbf{k}Q/\mathcal{I}$  be the basic algebra whose quiver and admissible ideal is given by

$$1 \xrightarrow{a} 2 \xrightarrow{b} 3 \quad \text{and} \quad \mathcal{I} = \langle ab \rangle,$$

respectively. One can check that  $T := S(1) \oplus S(2) \oplus S(3)$  is a strong generator of the finitely generated module category  $\mathbf{mod} \Lambda$  of  $\Lambda$  and the projective dimension of the direct summand  $S(1)$  of  $T$  equals to 2. Moreover, we have  $\text{pd } S(2) = 1$  and  $\text{pd } S(3) = 0$ . Therefore,  $\text{pd } T = \text{pd } S(1) = \text{gl.dim } \Lambda = 2$  in this example.

**3.3. Loewy lengths of some Artin algebras.** In this subsection we consider the Loewy lengths of some Artin algebras. This section containing three parts. In the parts one, i.e., 3.3.1, we show that the Loewy length of an SPI Artin algebra, an Artin algebra whose simple modules are either projective or injective, is less than or equal to two. In the parts two, i.e., 3.3.2, we provide a description of semi-simple algebra by using Loewy length. In the last parts, i.e., 3.3.3, we prove that the Loewy length of an Artin algebra with Orlov spectrum  $\{0, 1\}$  equals to two, and compute the Orlov spectrum of any SPI Artin algebra.

3.3.1. *Loewy lengths of SPI Artin algebras.*

**Proposition 3.10.** *If  $\Lambda$  is an SPI Artin algebra, then  $\ell(\Lambda) \leq 2$ .*

*Proof.* Assume that there is an SPI Artin algebra  $\Lambda$  such that all simple modules are either projective or injective and  $\ell(\Lambda) = n \geq 3$ . Then there is a projective module  $P$  with  $\ell(P) = n$ . In particular, we have  $\ell(\text{rad } P) = n - 1$  and  $\ell(\text{rad}^2 P) = n - 2$ . The following short exact sequence

$$0 \longrightarrow \text{rad}^2 P \longrightarrow \text{rad } P \longrightarrow \text{rad } P / \text{rad}^2 P \longrightarrow 0$$

admits  $\text{rad } P / \text{rad}^2 P = Q \oplus E$  such that  $Q$  is the sum of some simple projective modules and  $E$  is the sum of some simple injective modules by our assumption. Then the pullback of

$$\begin{array}{ccc} & & Q \\ & & \downarrow \leq \oplus \\ \text{rad } P & \xrightarrow{\text{canonical}} & \text{rad } P / \text{rad}^2 P \\ & \text{epimorphism} & \end{array}$$

is of the following form.

$$(3.1) \quad \begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \text{rad}^2 P & \longrightarrow & W & \longrightarrow & Q \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \text{rad}^2 P & \longrightarrow & \text{rad } P & \longrightarrow & \text{rad } P / \text{rad}^2 P \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & E & \xlongequal{\quad} & E \\ & & & & \downarrow & & \downarrow \\ & & & & 0 & & 0 \end{array}$$

Since  $Q$  is projective and  $E$  is injective, we have  $W \cong \text{rad}^2 P \oplus Q$  and  $\text{rad } P \cong W \oplus E$ . Thus, we have

$$n - 1 = \ell(\text{rad } P) = \ell(W \oplus E) = \ell(\text{rad}^2 P \oplus Q \oplus E)$$

$$= \max\{\ell(\text{rad}^2 P), \ell(Q), \ell(E)\} = n - 2$$

by  $n \geq 3$ , a contradiction.  $\square$

**Example 3.11.** The inverse of Proposition 3.10 maybe not hold. For example, let the path algebra  $\Lambda = \mathbf{k}Q/\langle \alpha\beta, \beta\alpha \rangle$ , where the quiver  $Q$  as follows

$$1 \begin{array}{c} \xrightarrow{\alpha} \\ \xleftarrow{\beta} \end{array} 2.$$

In this case, we have  $\ell(\Lambda) = 2$ , but  $S(1)$  and  $S(2)$  are neither projective nor injective.

### 3.3.2. Loewy lengths of semi-simple Artin algebras.

**Lemma 3.12.** *Assume that  $\Lambda$  is a representation-finite Artin algebra which is not semi-simple. Then  $\{0, 1\} \subseteq \text{OSpec}(\Lambda)$ .*

*Proof.* Let  $\text{ind}(\text{mod}\Lambda)$  be the set of all indecomposable  $\Lambda$ -modules (up to isomorphism). Since  $\Lambda$  is representation-finite, that is,  $\text{ind}(\text{mod}\Lambda)$  is a finite set, it is easy to see that

$$T = \bigoplus_{X \in \text{ind}(\text{mod}\Lambda)} X$$

is a strong generator of  $\text{mod}\Lambda$ , and, in this case, we obtain  $[T]_1 = \text{mod}\Lambda$ . Thus,  $0 \in \text{OSpec}(\Lambda)$ .

On the other hand, there is an indecomposable projective module  $P$  whose Loewy length  $\ell(P)$  equals to the Loewy length  $\ell(\Lambda)$  of  $\Lambda$ . Let

$$T' = \bigoplus_{X \in \text{ind}(\text{mod}\Lambda) \setminus \{P\}} X,$$

then the following short exact sequence

$$0 \longrightarrow \text{rad} P \longrightarrow P \longrightarrow P/\text{rad} P \longrightarrow 0$$

shows that  $P \in [T']_2$  since  $\text{rad} P, P/\text{rad} P \in [T']$ . On the other hand,  $P \notin [T']_1$ . That is,  $[T']_1 \subsetneq [T']_2 = \text{mod}\Lambda$ . It follows  $1 \in \text{OSpec}(\Lambda)$ .  $\square$

**Theorem 3.13.** *For an Artin algebra  $\Lambda$ , the following statement are equivalent.*

- (1)  $\Lambda$  is semi-simple.
- (2)  $\ell(\Lambda) = 1$ .
- (3)  $\text{OSpec}(\Lambda) = \{0\}$ .

*Proof.* First of all, since  $\ell(\Lambda) = 1$ , we obtain  $\ell(\Lambda/\text{rad}\Lambda) = 0$ . It follows  $\Lambda/\text{rad}\Lambda = 0$ , i.e.,  $\Lambda$  is semi-simple. Thus, (2) admits (1).

Second, if  $\Lambda$  is semi-simple, then all indecomposable  $\Lambda$ -modules are simple modules which are direct summands of  $\Lambda$ . It is easy to see that  $\ell(\Lambda) = 1$ . Thus, (1) and (2) are equivalent.

Third, we show that (1) admits (3). Notice that semi-simple Artin algebra are always representation-finite. Thus, we can assume that  $\Lambda = \bigoplus_{i=1}^n S_i$  is basic up to Morita equivalence, where  $S_1, \dots, S_n$  are simple. Then we obtain that each basic  $\Lambda$ -module  $M$  is isomorphic to

$$\bigoplus_{i=1}^n S_i^{\oplus t_i}, \quad (t_i \in \{0, 1\})$$

which lies in  $[\Lambda]_1 = \text{mod}\Lambda$ . Thus,  $0 \in \text{OSpec}(\Lambda)$ . However, if a  $\Lambda$ -module  $T$  does not contain some simple module  $S_i$  as its direct summand (up to isomorphism), then  $[T]_n \subsetneq \text{mod}\Lambda$  holds for all  $n \in \mathbb{N}$ . Then  $\text{OSpec}(\Lambda) = \{0\}$ .

Finally, it is easy to prove that  $\Lambda$  is semi-simple if  $\text{OSpec}(\Lambda) = \{0\}$  by Lemma 3.12. Therefore, the statements (1), (2) and (3) are equivalent.  $\square$

3.3.3. *Loewy length of Artin algebras with Orlov spectra  $\{0, 1\}$ .* The following result shows that the Loewy length of an Artin algebra with Orlov spectrum  $\{0, 1\}$  equals to two.

**Proposition 3.14.** *If  $\text{OSpec}(\Lambda) = \{0, 1\}$ , then  $\ell(\Lambda) = 2$ .*

*Proof.* If  $\ell(\Lambda) = n \geq 3$ , then  $n \in \text{OSpec}(\Lambda)$  by Proposition 3.3 (or Corollary 3.4 (1)), a contradiction. Thus,  $\ell(\Lambda) \leq 2$ . By Theorem 3.13 and  $\text{OSpec}(\Lambda) = \{0, 1\}$ , we obtain that  $\Lambda$  is not semi-simple, i.e.,  $\ell(\Lambda) \geq 2$ . Thus,  $\ell(\Lambda) = 2$ .  $\square$

**Theorem 3.15.** *Let  $\Lambda$  be an SPI Artin algebra which is not semi-simple.*

- (1) *If  $\Lambda$  is representation-finite, then  $\text{OSpec}(\Lambda) = \{0, 1\}$ .*
- (2) *If  $\Lambda$  is representation-infinite, then  $\text{OSpec}(\Lambda) = \{1\}$ .*

*Proof.* Assume  $\text{mod}\Lambda = [T]_{n+1}$  for some  $n \in \mathbb{N}$ . By Property 2.23, all simple modules lie in  $[T]_1 = \text{add}T$  since every simple module is either a projective module or an injective module. Thus,  $T \cong (\Lambda/\text{rad}\Lambda) \oplus T'$  for some module  $T'$ . Then we obtain

$$\begin{aligned} \Theta_{\text{mod}\Lambda}(T) &= \Theta_{\text{mod}\Lambda}((\Lambda/\text{rad}\Lambda) \oplus T') \\ \text{(by Lemma 2.4)} \quad &\leq \Theta_{\text{mod}\Lambda}(\Lambda/\text{rad}\Lambda) \\ \text{(Proposition 3.3)} \quad &= \ell(\Lambda) - 1 \\ \text{(Proposition 3.10)} \quad &= 2 - 1 = 1. \end{aligned}$$

It follows that  $\text{u.dim}(\text{mod}\Lambda) = \sup(\text{OSpec}(\Lambda)) \leq 1$ . On the other hand,  $\text{u.dim}(\text{mod}\Lambda) \geq \ell(\Lambda) - 1 = 1$  by Corollary 3.4 (2). That is,  $\text{u.dim}(\text{mod}\Lambda) = 1$ . Thus,  $1 \in \text{OSpec}(\Lambda)$ .

If  $\Lambda$  is representation-finite, then, by Lemma 3.12, we have  $\text{OSpec}(\Lambda) = \{0, 1\}$ , that is, the statement (1) holds; otherwise,  $\text{OSpec}(\Lambda) = \{1\}$ , that is, the statement (2) holds.  $\square$

Next, we provide some examples for Theorem 3.15.

**Example 3.16.** Consider the hereditary algebra  $\Lambda = \vec{A}_2 = \mathbf{k}\mathcal{Q}$  with  $\mathcal{Q} = 1 \xrightarrow{\alpha} 2$ . It is representation-finite, to be more precise, the number of isoclasses of indecomposable module equals to 3. Since the simple module  $S(2)$  is projective and the simple module  $S(1)$  is injective, we have  $\text{OSpec}(\Lambda) = \{0, 1\}$  by Theorem 3.15 (1).

**Example 3.17.** (1) Consider the path algebra  $\Lambda = \mathbf{k}\mathcal{Q}$  given by the following quiver shown in Figure 3.1. It is clearly that  $\Lambda$  is a hereditary algebra, and then  $\Lambda$  is representation-




FIGURE 3.1. The path algebra in Example 3.17 (1)

infinite since the underlying graph of  $\mathcal{Q}$  is not a Dynkin type (cf. [1, Chap VII, Section

2]). Moreover, the simple module  $S(0)$  is projective and, for any  $1 \leq i \leq n$ , the simple module  $S(i)$  is injective. Therefore,  $\text{OSpec}(\Lambda) = \{1\}$  by Theorem 3.15 (2).

(2) Consider the path algebra  $\Lambda = \mathbf{k}\mathcal{Q}$  given by the quiver shown in Figure 3.2, where  $t$  is an even number with  $t \geq 4$ . Now we compute its Orlov spectrum. First, it is clear that




FIGURE 3.2. The path algebra in Example 3.17 (2)

$\Lambda$  is a hereditary algebra. Then, it is not representation finite since the underlying graph of  $\mathcal{Q}$  is not a Dynkin type (cf. [1, Chap VII, Section 2]). On the other hand, it is easy to see that the simple modules  $S(2i)$  ( $i \in \mathbb{N}$ ,  $2 \leq 2i \leq t$ ) is indecomposable projective and  $S(2j+1)$  ( $j \in \mathbb{N}$ ,  $1 \leq 2j+1 \leq t$ ) is indecomposable injective. Therefore,  $\text{OSpec}(\Lambda) = \{1\}$  by Theorem 3.15 (2) as required.

#### 4. RADICAL LAYER LENGTHS

Layer length, introduced by Huard, Lanzilotta, and Hernández in [13], is closely related to finitistic dimension.

##### 4.1. Torsion pairs and radical Layer lengths.

**Definition 4.1.** A *torsion pair* (*torsion theory*) for  $\text{mod } \Lambda$  is a pair  $(\mathcal{T}, \mathcal{F})$  of classes of modules in  $\text{mod } \Lambda$  satisfying the following conditions:

- (1)  $\text{Hom}_\Lambda(M, N) = 0$  for each  $M \in \mathcal{T}$  and  $N \in \mathcal{F}$ ;
- (2) for each  $X \in \text{mod } \Lambda$ , if  $\text{Hom}_\Lambda(X, -)|_{\mathcal{F}} = 0$  then  $X \in \mathcal{T}$ ;
- (3) for each  $Y \in \text{mod } \Lambda$ , if  $\text{Hom}_\Lambda(-, Y)|_{\mathcal{T}} = 0$  then  $Y \in \mathcal{F}$ .

For a given class  $\mathcal{X}$  in  $\text{mod } \Lambda$  and a module  $M \in \text{mod } \Lambda$ , the trace of  $\mathcal{X}$  is the submodule  $\text{Trace}_{\mathcal{X}}(M)$  of  $M$  which is the sum of all images  $\text{Im}(f)$  of the morphisms  $f \in \text{Hom}_\Lambda(X, M)$

with  $X \in \mathcal{X}$ . Let  $(\mathcal{T}, \mathcal{F})$  be a torsion theory for  $\text{mod } \Lambda$ . Let us recall that  $t := \text{Trace}_{\mathcal{T}}$  is the so called torsion radical attached to  $(\mathcal{T}, \mathcal{F})$ . In fact,  $t(M)$  is the largest submodule of  $M$  lying in  $\mathcal{T}$ . And we write  $q_t := 1_{\text{mod } \Lambda}/t$ , i.e.,  $q_t(M) := M/t(M)$  for each  $M \in \text{mod } \Lambda$ . Thus, for each  $M \in \text{mod } \Lambda$ , we have the following short exact sequence

$$0 \longrightarrow t(M) \longrightarrow M \longrightarrow q_t(M) \longrightarrow 0$$

where  $t(M) \in \mathcal{T}$  and  $q_t(M) \in \mathcal{F}$ .

**Definition 4.2** ([13]). Given a torsion pair  $(\mathcal{T}, \mathcal{F})$  with the torsion radical  $t$ . The  $t$ -radical layer length is a function  $\ell^t : \text{mod } \Lambda \rightarrow \mathbb{N} \cup \{\infty\}$  via

$$\ell^t(M) = \inf\{i \geq 0 \mid t \circ F_t^i(M) = 0, M \in \text{mod } \Lambda\}$$

where  $F_t = \text{rad} \circ t$ .

Let  $\mathbf{S}$  be a subset of some simple  $\Lambda$ -modules and  $\mathbf{S}'$  the set of all the others simple  $\Lambda$ -modules. A  $\mathbf{S}$ -filtration of a  $\Lambda$ -module  $M$  is a filtration  $0 \subseteq M_0 \subseteq M_1 \cdots \subseteq M_{m-1} \subseteq M_m = M$  such that  $M_i/M_{i-1} \in \mathbf{S}$  holds for all  $1 \leq i \leq m$ . We use  $\mathcal{F}_{\mathbf{S}}$  to represent the subcategory of  $\text{mod } \Lambda$  generated by all  $\Lambda$ -modules  $M$  with some  $\mathbf{S}$ -filtration. We have the following statements:

- it is easy to check that  $\mathcal{F}_{\mathbf{S}}$  is closed under extensions, submodules and quotients modules;
- and there is a subcategory  $\mathcal{T}_{\mathbf{S}}$  of  $\text{mod } \Lambda$  such that  $(\mathcal{T}_{\mathbf{S}}, \mathcal{F}_{\mathbf{S}})$  is a torsion pair (the corresponding torsion radical is denoted by  $t_{\mathbf{S}}$ ),

and have the following results.

**Proposition 4.3** ([26, Proposition 3.2]). *If  $\mathbf{S} = \emptyset$ , then  $\ell^{t_{\mathbf{S}}}(M) = \ell(M)$  for each  $M \in \text{mod } \Lambda$ .*

**Lemma 4.4.** *Let the pair  $(\mathcal{T}_{\mathbf{S}}, \mathcal{F}_{\mathbf{S}})$  be a torsion pair as above.*

- (1) ([15, Proposition 5.9])  $\mathcal{T}_{\mathbf{S}} = \{M \in \text{mod } \Lambda \mid \text{top}(M) \in \text{add}(\mathbf{S}')\}$ ;
- (2) ([15, Lemma 3.3(c)]) *If  $M \in \mathcal{T}_{\mathbf{S}}$  and  $M \neq 0$ , then  $\ell^{t_{\mathbf{S}}}(M) = \ell^{t_{\mathbf{S}}}(\text{rad } M) + 1$ ;*
- (3) ([15, Lemma 3.4(a)])  $\ell^{t_{\mathbf{S}}}(\bigoplus_{i=1}^n M_i) = \sup_{1 \leq i \leq n} \ell^{t_{\mathbf{S}}}(M_i)$ , where  $M_i \in \text{mod } \Lambda$  for all  $1 \leq i \leq n$ .

**4.2. Short exact sequences and radical layer length.** In this section, we show two main results, i.e., Theorems 4.5 and 4.14, of our paper.

**4.2.1. The first main result and some corollaries.** In this subdivision, we show the following theorem which is the first result of our paper.

**Theorem 4.5.** *Let  $0 \longrightarrow L \longrightarrow M \longrightarrow N \longrightarrow 0$  be an exact sequence in  $\text{mod } \Lambda$ . Then*

$$\max\{\ell^{t_{\mathbf{S}}}(L), \ell^{t_{\mathbf{S}}}(N)\} \leq \ell^{t_{\mathbf{S}}}(M) \leq \ell^{t_{\mathbf{S}}}(L) + \ell^{t_{\mathbf{S}}}(N).$$

*In particular, if  $\ell^{t_{\mathbf{S}}}(L) = 0$ , then  $\ell^{t_{\mathbf{S}}}(N) = \ell^{t_{\mathbf{S}}}(M)$ ; if  $\ell^{t_{\mathbf{S}}}(N) = 0$ , then  $\ell^{t_{\mathbf{S}}}(L) = \ell^{t_{\mathbf{S}}}(M)$ .*

We need some lemmas for proving it.

**Lemma 4.6.** *Let  $X$  be a  $\Lambda$ -module.*

- (1) ([13, Proposition 5.9 (c)])  $t_{\mathbf{S}}(\Lambda_{\Lambda})$  is a two side ideal of  $\Lambda$  and  $t_{\mathbf{S}}(X) = X \cdot t_{\mathbf{S}}(\Lambda_{\Lambda})$ .
- (2) ([4, Propostion 3.5])  $\text{rad } X = X \cdot \text{rad}(\Lambda_{\Lambda})$ .
- (3)  $\text{rad}(t_{\mathbf{S}}(\Lambda_{\Lambda})) = t_{\mathbf{S}}(\Lambda_{\Lambda}) \cdot \text{rad}(\Lambda_{\Lambda})$ .
- (4)  $t_{\mathbf{S}}(\text{rad}(t_{\mathbf{S}}(\Lambda_{\Lambda}))) = t_{\mathbf{S}}(\Lambda_{\Lambda}) \cdot \text{rad}(\Lambda_{\Lambda}) \cdot t_{\mathbf{S}}(\Lambda_{\Lambda})$ .
- (5)  $t_{\mathbf{S}}(F_{t_{\mathbf{S}}}^i(\Lambda_{\Lambda}))$  is an ideal of  $\Lambda$  for each  $i \geq 0$ .

*Proof.* The statement (3) can be obtained by (2) in the case of  $X = t_{\mathbf{S}}(\Lambda_{\Lambda})$ . The statement (4) is obtained by (1) and (3), and the statement (5) is obtained by (1)–(4).  $\square$

**Lemma 4.7.** *For any module  $X \in \text{mod } \Lambda$ , we have  $t_{\mathbf{S}}(F_{t_{\mathbf{S}}}^i(X)) = X \cdot t_{\mathbf{S}}(F_{t_{\mathbf{S}}}^i(\Lambda_{\Lambda}))$  for each  $i \geq 0$ .*



*Proof.* Lemma 4.6 (1) admits this statement under the case for  $i = 0$ . Assume that if  $i = n$  then  $t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^n(X)) = X \cdot t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^n(\Lambda_{\Lambda}))$ . For the case  $i = n + 1$ , we have

$$\begin{aligned}
& t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{n+1}(X)) = t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^n(F_{t_{\mathcal{S}}}(X))) \\
(\text{using assumption}) & \quad = F_{t_{\mathcal{S}}}(X) \cdot t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^n(\Lambda_{\Lambda})) \\
(F_{t_{\mathcal{S}}} = \text{rad} \circ t_{\mathcal{S}}) & \quad = \text{rad}(t_{\mathcal{S}}(X)) \cdot t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^n(\Lambda_{\Lambda})) \\
(\text{Lemma 4.6 (2)}) & \quad = t_{\mathcal{S}}(X) \cdot \text{rad}(\Lambda_{\Lambda}) \cdot t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^n(\Lambda_{\Lambda})) \\
(\text{Lemma 4.6 (2)}) & \quad = t_{\mathcal{S}}(X) \cdot \text{rad}(t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^n(\Lambda_{\Lambda}))) \\
(F_{t_{\mathcal{S}}} = \text{rad} \circ t_{\mathcal{S}}) & \quad = t_{\mathcal{S}}(X) \cdot F_{t_{\mathcal{S}}}^{n+1}(\Lambda_{\Lambda}) \\
(\text{Lemma 4.6 (1)}) & \quad = X \cdot t_{\mathcal{S}}(\Lambda_{\Lambda}) \cdot F_{t_{\mathcal{S}}}^{n+1}(\Lambda_{\Lambda}) \\
(\text{Lemma 4.6 (3) (4)}) & \quad = X \cdot t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{n+1}(\Lambda_{\Lambda})),
\end{aligned}$$

which yields that this lemma holds by induction.  $\square$

**Lemma 4.8.**

- (1) *The functor  $t_{\mathcal{S}}$  and  $\text{rad}$  preserve monomorphism and epimorphism.*
- (2) *For each  $i \geq 0$ ,  $F_{t_{\mathcal{S}}}^i$  and  $t_{\mathcal{S}} \circ F_{t_{\mathcal{S}}}^i$  preserve monomorphism and epimorphism.*

*Proof.* (1) The functor  $t_{\mathcal{S}}$  preserving monomorphism and epimorphism is proven in [26, Lemma 3.3]. By [13, Lemma 3.6(a)],  $\text{rad}$  preserves monomorphism, and by [1, Chapter V, Lemma 1.1],  $\text{rad}$  preserves epimorphism.

(2) This statement is a direct corollary of (1).  $\square$

**Lemma 4.9.** *Let  $X$  be a  $\Lambda$ -module. Then we have  $t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(X)}(X)) = 0$ .*

*Proof.* By Definition 4.2, we have this observation.  $\square$

Now, we prove the first main result of our paper.

**The proof of Theorem 4.5.** By Lemma 4.8, we know that  $F_{t_{\mathcal{S}}} = \text{rad} \circ t_{\mathcal{S}}$  preserve monomorphism and epimorphism. Thus, by [13, Lemma 3.4(b)(c)], we obtain  $\ell\ell^{\mathcal{S}}(L) \leq \ell\ell^{\mathcal{S}}(M)$  and  $\ell\ell^{\mathcal{S}}(N) \leq \ell\ell^{\mathcal{S}}(M)$ , i.e.,  $\max\{\ell\ell^{\mathcal{S}}(L), \ell\ell^{\mathcal{S}}(N)\} \leq \ell\ell^{\mathcal{S}}(M)$ .

Next, we prove the second “ $\leq$ ”, it can be obtained by proving

$$(4.1) \quad t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(L)+\ell\ell^{\mathcal{S}}(N)}(M)) \subseteq t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(L)}(L)) = 0,$$

see Lemmas 4.8 (2) and 4.9, and Definition 4.2.

By Lemma 4.6 (5),  $t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^i(\Lambda_{\Lambda}))$  is an ideal of  $\Lambda$  for each  $i \geq 0$ , then we have

$$\begin{aligned}
(\text{Lemma 4.7}) \quad & (t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(N)}(M)) + L)/L = (M \cdot t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(N)}(\Lambda_{\Lambda})) + L)/L \\
& \quad = M/L \cdot t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(N)}(\Lambda_{\Lambda})) \\
(M/L \cong N) & \quad \cong N \cdot t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(N)}(\Lambda_{\Lambda})) \\
(\text{Lemma 4.7}) & \quad = t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(N)}(N)) \\
(\text{Lemma 4.9}) & \quad = 0.
\end{aligned}$$

Thus,  $t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(N)}(M)) + L = L$ . It follows  $t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(N)}(M)) \subseteq L$ . Furthermore, we obtain

$$\begin{aligned}
& t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(L)+\ell\ell^{\mathcal{S}}(N)}(M)) = (t_{\mathcal{S}}F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(L)} \circ F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(N)})(M) \\
(t_{\mathcal{S}}^2 = t_{\mathcal{S}}) & \quad = (t_{\mathcal{S}}F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(L)} \circ t_{\mathcal{S}}F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(N)})(M) \\
& \quad \subseteq t_{\mathcal{S}}(F_{t_{\mathcal{S}}}^{\ell\ell^{\mathcal{S}}(L)}(L)) \\
(\text{Lemma 4.9}) & \quad = 0,
\end{aligned}$$

that is, (4.1) holds.  $\square$

Here are some corollaries about Theorem 4.5.

**Corollary 4.10.** *Let  $M \in [T]_{n+1}$ . Then  $\ell\ell^{ts}(M) \leq (n+1)\ell\ell^{ts}(T)$ .*

*Proof.* A module  $M$  lying in  $[T]_{n+1}$  provides a family of short exact sequences

$$(0 \longrightarrow Y_i \longrightarrow Z_{i-1} \oplus Z'_{i-1} \longrightarrow Z_i \longrightarrow 0)_{1 \leq i \leq n},$$

where  $Z_0 = M$ ,  $Y_1, \dots, Y_n \in [T]_1$ ,  $Z_i \in [T]_{n-i+1}$ . Then we obtain

$$\begin{aligned} \ell\ell^{ts}(M) &= \ell\ell^{ts}(Z_0) \leq \ell\ell^{ts}(Z_0 \oplus Z'_0) \\ (\text{Lemma 3.5}) \quad &\leq \ell\ell^{ts}(Y_1) + \ell\ell^{ts}(Z_1) \\ &\leq \ell\ell^{ts}(T) + \ell\ell^{ts}(Z_1 \oplus Z'_1) \\ &\leq \ell\ell^{ts}(T) + \ell\ell^{ts}(Y_2) + \ell\ell^{ts}(Z_2) \\ (Y_2 \in \text{add } T) \quad &\leq \ell\ell^{ts}(T) + \ell\ell^{ts}(T) + \ell\ell^{ts}(Z_2) \\ &= 2\ell\ell^{ts}(T) + \ell\ell^{ts}(Z_2) \\ &\vdots \\ &\leq n\ell\ell^{ts}(T) + \ell\ell^{ts}(Z_n) \\ (Z_n \in \text{add } T) \quad &\leq n\ell\ell^{ts}(T) + \ell\ell^{ts}(T) \\ &= (n+1)\ell\ell^{ts}(T). \end{aligned}$$

□

In the case for  $\mathbf{S} = \{S \in \text{mod } A \mid \text{pd } S < \infty\}$ , we define  $\ell\ell^\infty(M) := \ell\ell^{ts}(M)$  in this situation, see [15, Definition 4.2]. Then the following corollary holds.

**Corollary 4.11.** *Let  $0 \longrightarrow L \longrightarrow M \longrightarrow N \longrightarrow 0$  be an exact sequence in  $\text{mod } A$ . Then the following statements hold.*

- (1)  $\max\{\ell\ell(L), \ell\ell(N)\} \leq \ell\ell(M) \leq \ell\ell(L) + \ell\ell(N)$ .
- (2)  $\max\{\ell\ell^\infty(L), \ell\ell^\infty(N)\} \leq \ell\ell^\infty(M) \leq \ell\ell^\infty(L) + \ell\ell^\infty(N)$ .
- (3) *If  $\ell\ell^\infty(L) = 0$  (resp.  $\ell\ell^\infty(N) = 0$ ), then  $\ell\ell^\infty(N) = \ell\ell^\infty(M)$  (resp.  $\ell\ell^\infty(L) = \ell\ell^\infty(M)$ ).*

*Proof.* (1) and (2) are particular cases of Theorem 4.5. We need prove (3).

If  $\ell\ell^\infty(L) = 0$ , then we have

$$\ell\ell^\infty(N) = \max\{\ell\ell^\infty(L), \ell\ell^\infty(N)\} \leq \ell\ell^\infty(M) \leq \ell\ell^\infty(L) + \ell\ell^\infty(N) = \ell\ell^\infty(N)$$

by (2). Thus,  $\ell\ell^\infty(M) = \ell\ell^\infty(N)$  in this situation. We can show that  $\ell\ell^\infty(N) = 0$  admits  $\ell\ell^\infty(L) = \ell\ell^\infty(M)$  by similar way. □

**Remark 4.12.** Note that the functions Loewy length  $\ell\ell$  and infinite layer length  $\ell\ell^\infty$  are particular radical layer length, see more details [13, 15]. Corollary 4.11 (1) is a classical result. The first “ $\leq$ ” in Corollary 4.11 (2) is first established in [15, Proposition 4.5 (a) (b)].

**Corollary 4.13.** *Assume  $\ell\ell^{ts}(A) = n$ . Then for each  $M \in \text{mod } A$ , we have  $\ell\ell^{ts}(F_{ts}^i(M)) \leq n - i$  for each  $0 \leq i \leq n$ .*

*Proof.* By [26, Lemma 2.6], if  $M \in \text{mod } A$  is a  $A$ -module  $M$  with  $\ell\ell^{ts}(M) = n$ , then  $\ell\ell^{ts}(F_{ts}^i(M)) = n - i$  for each  $0 \leq i \leq n$ . Obviously, any module  $M$  has a free precover  $f : A^{\oplus I} \rightarrow M$  which induces a short exact sequence

$$0 \longrightarrow \ker f \longrightarrow A^{\oplus I} \longrightarrow M \longrightarrow 0,$$

where  $I$  is a finite index set. It yields  $\ell\ell^{ts}(M) \leq \ell\ell^{ts}(A) \leq n$  by using Theorem 4.14 as required. □

4.2.2. *The second main result and some corollaries.* In this subdivision, we show the following theorem which is the second result of our paper.

**Theorem 4.14.** *If  $\Lambda$  is representation-finite, then*

$$\left\{ \left\lceil \frac{\ell\ell^{ts}(\Lambda)}{d} \right\rceil - 1 \mid d \in \mathbb{N} \text{ and } 1 \leq d < \ell\ell^{ts}(\Lambda) \right\} \subseteq \text{OSpec}(\Lambda),$$

where  $\lceil \alpha \rceil$  is the least integer greater than  $\alpha$ .

We need the following lemma for proving Theorem 4.14

**Lemma 4.15.** *A family*

$$(0 \longrightarrow M_i \xrightarrow{\varphi_i} M_{i-1} \longrightarrow N_i \longrightarrow 0)_{1 \leq i \leq d}$$

of short exact sequences induces a short exact sequence

$$0 \longrightarrow M_d \longrightarrow M_0 \longrightarrow N_{[0,d]} \longrightarrow 0$$

satisfying  $\ell\ell^{ts}(N_{[0,d]}) \leq \sum_{i=1}^d \ell\ell^{ts}(N_i)$ , where  $d \geq 1$ .

*Proof.* The case for  $d = 1$  is trivial. Next, we show this lemma holds for  $d \geq 2$ .

For simplification, we define  $N_{[u,u+i]} := M_u/M_{u+i}$  which is induced by the cokernel of the composition

$$\begin{aligned} \varphi_{[u,u+i]} &:= \varphi_{u+1} \circ \cdots \circ \varphi_{u+i} : M_{u+i} \rightarrow M_u \\ (0 \leq u < u+i \leq d, 1 \leq i \leq d-u). \end{aligned}$$

Since composition  $\varphi_{[u-1,u+i]}$  is always monomorphic,  $\varphi_{[0,d]}$  induces a short exact sequence which is of the form

$$0 \longrightarrow M_d \xrightarrow{\varphi_{[0,d]}} M_0 \longrightarrow N_{[0,d]} \longrightarrow 0,$$

where  $N_{[0,d]} \cong M_0/M_d$ . Moreover, for any  $1 \leq i \leq d$ , the epimorphism  $\phi_{[d-(i+1),d]} : M_{d-(i+1)} \rightarrow N_{[d-(i+1),d]}$  induced by the cokernel of  $\varphi_{[d-(i+1),d]}$  and the monomorphism  $\phi_{d-i} : N_{[d-i,d]} \rightarrow N_{[d-(i+1),d]}$  obtained by the universal property of the cokernel of  $\varphi_{[d-i,d]}$  provide the following diagram

commutes, whose rows and columns are exact, and the commutative square marked by “♠” is both a push-out and pull-back square. Of course, the above diagram can also be induced by using the snake lemma. Then we obtain a family

$$(0 \longrightarrow N_{[d-j,d]} \longrightarrow N_{[d-(j+1),d]} \longrightarrow N_{d-(j+1)} \longrightarrow 0)_{0 \leq j \leq d-2}$$

of short exact sequences (see the shadow of the above diagram), where  $N_{[d,d]} := N_d$ . Then, by Theorem 4.14, we have

$$\ell\ell^{ts}(N_{[1,d]}) \leq \ell\ell^{ts}(N_{[2,d]}) + \ell\ell^{ts}(N_1)$$

$$\begin{aligned}
&\leq \ell\ell^{ts}(N_{[3,d]}) + \ell\ell^{ts}(N_2) + \ell\ell^{ts}(N_1) \\
&\quad \vdots \\
&\leq \ell\ell^{ts}(N_d) + \cdots + \ell\ell^{ts}(N_2) + \ell\ell^{ts}(N_1) = \sum_{i=1}^d \ell\ell^{ts}(N_i).
\end{aligned}$$

□

Now we prove Theorem 4.14.

**The proof of Theorem 4.14.** This proof is divided to two parts (i) and (ii).

(i) In the case of  $\ell\ell^{ts}(\Lambda) = 1$ , we have

$$\left\{ \left\lceil \frac{\ell\ell^{ts}(\Lambda)}{d} \right\rceil - 1 \mid d \in \mathbb{N} \text{ and } 1 \leq d < \ell\ell^{ts}(\Lambda) \right\} \Big|_{\ell\ell^{ts}(\Lambda)=1} = \{0\} \subseteq \text{OSpec}(\Lambda).$$

(ii) Assume that  $\ell\ell^{ts}(\Lambda) = n \geq 2$ , and for any  $1 \leq d < \ell\ell^{ts}(\Lambda)$ , assume  $n = dm + r$ , where  $0 \leq r < d$ . Thus, we need show  $m - 1 \in \text{OSpec}(\Lambda)$  for proving this theorem. To do this, we construct a  $\Lambda$ -module  $T$  such that  $\mathcal{O}_{\text{mod}\Lambda}(T) = m - 1$  in this proof.

First of all, notice that, for arbitrary  $M \in \text{mod}\Lambda$ , we have the following two short exact sequences

$$(4.2) \quad \begin{aligned} &0 \longrightarrow t_S \circ F_{t_S}^{i-1}(M) \xrightarrow{f_{i-1}} F_{t_S}^{i-1}(M) \longrightarrow q_{t_S} \circ F_{t_S}^{i-1}(M) \longrightarrow 0 \\ &\text{and } 0 \longrightarrow F_{t_S}^i(M) \longrightarrow t_S \circ F_{t_S}^{i-1}(M) \xrightarrow{g_{i-1}} \text{top}(t_S \circ F_{t_S}^i(M)) \longrightarrow 0 \end{aligned}$$

for each  $1 \leq i \leq n - 1$ . Here,  $F_{t_S}^i(M) = \text{rad}(t_S \circ F_{t_S}^{i-1}(M))$ , see Definition 4.2. It follows that the following push-out square

$$\begin{array}{ccc}
t_S \circ F_{t_S}^{i-1}(M) & \xrightarrow{f_{i-1}} & F_{t_S}^{i-1}(M) \\
g_{i-1} \downarrow & & \downarrow \\
\text{top}(t_S \circ F_{t_S}^{i-1}(M)) & \longrightarrow & D_i
\end{array}$$

provides the diagram

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
& & F_{t_S}^i(M) & \xlongequal{\quad} & F_{t_S}^i(M) & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & t_S \circ F_{t_S}^{i-1}(M) & \xrightarrow{f_{i-1}} & F_{t_S}^{i-1}(M) & \longrightarrow & q_{t_S} \circ F_{t_S}^{i-1}(M) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \parallel \\
0 & \longrightarrow & \text{top}(t_S \circ F_{t_S}^{i-1}(M)) & \longrightarrow & D_i & \longrightarrow & q_{t_S} \circ F_{t_S}^{i-1}(M) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \\
& & 0 & & 0 & & 
\end{array}$$

commutes, whose rows and columns are exact. Thus, we obtain that

$$(4.3) \quad \ell\ell^{ts}(D_i) \leq \ell\ell^{ts}(\text{top}(t_S \circ F_{t_S}^{i-1}(M))) + \ell\ell^{ts}(q_{t_S} \circ F_{t_S}^{i-1}(M)) \leq 1 + 0 = 1$$

holds for each  $1 \leq i \leq n - 1$ , and there is a family

$$(0 \longrightarrow F_{t_S}^i(M) \xrightarrow{\varphi_i} F_{t_S}^{i-1}(M) \longrightarrow D_i \longrightarrow 0)_{1 \leq i \leq n-1}$$

of short exact sequences.

Second, the composition  $\varphi_{[ud,(u+1)d]} := \varphi_{(u+1)d} \circ \cdots \circ \varphi_{ud+1}$  ( $1 \leq u < m$ ) (resp.  $\varphi_{[md,md+r]} := \varphi_{md+r} \circ \cdots \circ \varphi_{md+1}$ ) is monomorphic since all  $\varphi_i$  are monomorphic, it induces a short exact sequence which is of the form

$$(4.4) \quad 0 \longrightarrow F_{t_S}^{(u+1)d}(M) \longrightarrow F_{t_S}^{ud}(M) \longrightarrow D_{[ud,(u+1)d]} \longrightarrow 0$$

$$(4.5) \quad (\text{resp. } 0 \longrightarrow F_{t_S}^{(m-1)d-r}(M) \longrightarrow F_{t_S}^{(m-2)d}(M) \longrightarrow D_{[(m-2)d,(m-1)d-r]} \longrightarrow 0).$$

where

$$D_{[ud,(u+1)d]} \cong F_{t_S}^{ud}(M)/F_{t_S}^{(u+1)d}(M)$$

$$(\text{resp. } D_{[(m-2)d,(m-1)d-r]} \cong F_{t_S}^{(m-2)d}(M)/F_{t_S}^{(m-1)d-r}(M)).$$

By Lemma 4.15 and (4.3), we have

$$\ell^{\ell^{\text{ts}}}(D_{[ud,(u+1)d]}) \leq \sum_{i=ud+1}^{(u+1)d} \ell^{\ell^{\text{ts}}}(D_i) = d$$

$$\left( \text{resp. } \ell^{\ell^{\text{ts}}}(D_{[(m-2)d,(m-1)d-r]}) \leq \sum_{i=(m-2)d+1}^{(m-1)d-r} \ell^{\ell^{\text{ts}}}(D_i) \leq d \right).$$

Then, by using (4.4) and (4.5), we have

$$\begin{aligned} [M]_1 &= [\text{rad}^0 \circ t_S^0(M)]_1 = [F_{t_S}^0(M)]_1 \\ &\subseteq [F_{t_S}^d(M)]_1 \cdot [D_{[0,d]}]_1 \\ &\subseteq [F_{t_S}^{2d}(M)]_1 \cdot [D_{[d,2d]}]_1 \cdot [D_{[0,d]}]_1 \\ &\quad \vdots \\ &\subseteq [F_{t_S}^{(m-2)d}(M)]_1 \cdot [D_{[(m-3)d,(m-2)d]}]_1 \cdot \cdots \cdot [D_{[0,d]}]_1 \\ (4.6) \quad &\subseteq [F_{t_S}^{(m-1)d-r}(M)]_1 \cdot [D_{[(m-2)d,(m-1)d-r]}]_1 \cdot [D_{[(m-3)d,(m-2)d]}]_1 \cdot \cdots \cdot [D_{[0,d]}]_1. \end{aligned}$$

Finally, we construct the  $\Lambda$ -module  $T$  as follows

$$T = \bigoplus_{Z \in \mathcal{W}_d} Z,$$

where  $\mathcal{W}_d := \{Z \in \text{ind}(\text{mod } \Lambda) \mid \ell^{\ell^{\text{ts}}}(Z) \leq d\}$ . We claim that  $[T]_m = \text{mod } \Lambda$  and  $[T]_{m-1} \subsetneq \text{mod } \Lambda$  hold. Indeed, by Lemma 4.4 (3), we have

$$\ell^{\ell^{\text{ts}}}(T) = \ell^{\ell^{\text{ts}}}\left(\bigoplus_{Z \in \mathcal{W}_d} Z\right) = \sup\{\ell^{\ell^{\text{ts}}}(Z) \mid Z \in \mathcal{W}_d\} \leq d.$$

Then we have  $[M]_1 \subseteq \overbrace{[T]_1 \cdot [T]_1 \cdot \cdots \cdot [T]_1}^m = [T]_m$  by (4.6). In this case,  $\text{mod } \Lambda = [T]_m$ .

On the other hand, if  $[T]_{m-1} = \text{mod } \Lambda$ , then, by Corollary 4.10, we obtain

$$\ell^{\ell^{\text{ts}}}(\Lambda) \leq (m-1)\ell^{\ell^{\text{ts}}}(T) \leq (m-1)d = md - d < md - j = \ell^{\ell^{\text{ts}}}(\Lambda),$$

This is a contradiction.

Therefore,  $\Theta_{\text{mod } \Lambda}(T) = m-1$ , and then  $m-1 \in \text{OSpec}(\Lambda)$  holds.  $\square$

**Example 4.16.** Take

$$\Lambda = \mathbf{k}Q / \langle (\alpha_1\alpha_2\alpha_3\alpha_4)^5 \rangle,$$

where  $Q$  is defined as

$$\begin{array}{ccc} 1 & \xrightarrow{\alpha_1} & 2 \\ \alpha_4 \uparrow & & \downarrow \alpha_2 \\ 4 & \xleftarrow{\alpha_3} & 3. \end{array}$$

We have seven items as follows:

- (a) Let  $\mathbf{S} = \emptyset$ , then  $\ell\ell^{ts}(A)$  by Proposition 4.3. On the other hand,  $\ell\ell(A) = \sup\{\ell\ell(P(i)) \mid 1 \leq i \leq 4\} = 23$ , then by Theorem 4.14, we have

$$\{0, 1, 2, 3, 4, 5, 7, 11, 22\} \subseteq \text{OSpec}(A).$$

- (b) Let  $\mathbf{S} = \{S(1)\}$ , then  $\ell\ell^{ts}(A) = \sup\{\ell\ell^{ts}(P(i)) \mid 1 \leq i \leq 4\} = \ell\ell^{ts}(P(2)) = 18$ . Furthermore, by Theorem 4.14, we have

$$\{0, 1, 2, 3, 4, 5, 8, 17\} \subseteq \text{OSpec}(A).$$

- (c) Let  $\mathbf{S} = \{S(2)\}, \{S(3)\}$ , or  $\{S(4)\}$ , then  $\ell\ell^{ts}(A) = \sup\{\ell\ell^{ts}(P(i)) \mid 1 \leq i \leq 4\} = \ell\ell^{ts}(P(2)) = 17$ . Furthermore, by Theorem 4.14, we have

$$\{0, 1, 2, 3, 4, 5, 8, 16\} \subseteq \text{OSpec}(A).$$

- (d) Let  $\mathbf{S} = \{S(1), S(2)\}, \{S(1), S(3)\}$ , or  $\{S(1), S(4)\}$ , then  $\ell\ell^{ts}(A) = \sup\{\ell\ell^{ts}(P(i)) \mid 1 \leq i \leq 4\} = \ell\ell^{ts}(P(2)) = 12$ . Furthermore, by Theorem 4.14, we have

$$\{0, 1, 2, 3, 5, 11\} \subseteq \text{OSpec}(A).$$

- (e) Let  $\mathbf{S} = \{S(2), S(3)\}, \{S(2), S(4)\}$ , or  $\{S(3), S(4)\}$ , then  $\ell\ell^{ts}(A) = \sup\{\ell\ell^{ts}(P(i)) \mid 1 \leq i \leq 4\} = \ell\ell^{ts}(P(2)) = 11$ . Furthermore, by Theorem 4.14, we have

$$\{0, 1, 2, 3, 5, 10\} \subseteq \text{OSpec}(A).$$

- (f) Let  $\mathbf{S} = \{S(1), S(2), S(3)\}, \{S(1), S(2), S(4)\}$ , or  $\{S(1), S(3), S(4)\}$ , then  $\ell\ell^{ts}(A) = \sup\{\ell\ell^{ts}(P(i)) \mid 1 \leq i \leq 4\} = \ell\ell^{ts}(P(2)) = 6$ . Furthermore, by Theorem 4.14, we have

$$\{0, 1, 2, 3, 5\} \subseteq \text{OSpec}(A).$$

- (g) Let  $\mathbf{S} = \{S(2), S(3), S(4)\}$ , then  $\ell\ell^{ts}(A) = \sup\{\ell\ell^{ts}(P(i)) \mid 1 \leq i \leq 4\} = \ell\ell^{ts}(P(2)) = 5$ . Furthermore, by Theorem 4.14, we have

$$\{0, 1, 2, 4\} \subseteq \text{OSpec}(A).$$

By (a)–(g), we have

$$\{0, 1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 16, 17, 22\} \subseteq \text{OSpec}(A).$$

**Corollary/Proposition 4.17.** *The Orlov spectrum  $\text{OSpec}(A)$  of the finite-dimensional algebra  $A = \mathbf{k}Q/I$  given by the quiver shown in Figure 4.1 and the admissible ideal  $I = \langle \alpha_1\alpha_2 \cdots \alpha_m \rangle$  and  $2 \leq m \leq n - 1$ . satisfies*

$$(4.7) \quad \{0, 1, 2, \dots, m+n-1\} \subseteq \text{OSpec}(A).$$




FIGURE 4.1. The path algebra in Corollary/Proposition 4.17

*Proof.* We have the five items as follows.

- (a) If  $\mathbf{S} = \emptyset$ , then  $\ell\ell^{ts}(A) = \ell\ell(A)m + n$ .
- (b) If  $\mathbf{S} = \{S(2), S(3), \dots, S(i)\}$  (for any  $i$  satisfying  $2 \leq i \leq m$ ), then  $\ell\ell^{ts}(A) = \sup\{P(i) \mid 1 \leq i \leq n\} = \ell\ell^{ts}(P(2)) = m + n + 2 - 2i$ .
- (c) If  $\mathbf{S} = \{S(1), S(2), S(3), \dots, S(i)\}$  (for any  $i$  satisfying  $2 \leq i \leq m$ ), then  $\ell\ell^{ts}(A) = \sup\{P(i) \mid 1 \leq i \leq n\} = \ell\ell^{ts}(P(2)) = m + n + 1 - 2i$ .
- (d) If  $\mathbf{S} = \{S(1), S(2), S(3), \dots, S(m), S(m+1), \dots, S(i)\}$  (for any  $i$  satisfying  $m+2 \leq i \leq n$ ), then  $\ell\ell^{ts}(A) = \sup\{P(i) \mid 1 \leq i \leq n\} = \ell\ell^{ts}(P(2)) = n + 2 - i$ .
- (e) If  $\mathbf{S} = \{S(2), S(3), \dots, S(m), S(m+1), \dots, S(n)\}$ , then  $\ell\ell^{ts}(A) = \sup\{P(i) \mid 1 \leq i \leq n\} = \ell\ell^{ts}(P(2)) = 1$ .

By Theorem 4.14, we obtain  $\ell\ell^{ts}(A) - 1 \in \text{OSpec}(A)$ . Thus, (a)–(e) yields that  $\{1, 2, \dots, m+n-1\} \subseteq \text{OSpec}(A)$ . Moreover,  $A$  is representation-finite, it follows  $0 \in \text{OSpec}(A)$ , see Lemma 3.12. Therefore, (4.7) holds.  $\square$

## 5. APPLICATION: THE ORLOV SPECTRUM WITH TYPE $\mathbb{A}_n$

Let  $\vec{\mathbb{A}}_n$  be the linearly oriented quiver with type  $\mathbb{A}_n$ , i.e.,

$$\vec{\mathbb{A}}_n = 1 \xrightarrow{a_1} 2 \xrightarrow{a_2} \dots \xrightarrow{a_{n-1}} n.$$

In this subsection, we compute the Orlov spectrum  $\text{OSpec}(A_n)$  of  $A_n := \mathbf{k}\vec{\mathbb{A}}_n$ .

**Proposition 5.1.**  $\{0, 1, 2, \dots, n-1\} \subseteq \text{OSpec}(A_n)$ .

*Proof.* For each  $1 \leq i \leq n-1$ , we have the following three items.

- (a) For the case of  $\mathbf{S} = \emptyset$ , we have  $\ell\ell(A_n) - 1 = n - 1 \in \text{OSpec}(A_n)$ .
- (b) For the case of  $\mathbf{S} = \{S(1)\}$ , we have  $\ell\ell^{ts}(A_n) = \sup\{P(i) \mid 1 \leq i \leq n\} = n - 1$ . Then we obtain  $\ell\ell^{ts}(A_n) - 1 = n - 2 \in \text{OSpec}(A_n)$  by Theorem 4.14.
- (c) For the case of  $\mathbf{S} = \{S(1), \dots, S(j)\}$  ( $1 \leq j \leq n-1$ ), we have  $\ell\ell^{ts}(A_n) = \sup\{P(i) \mid 1 \leq i \leq n\} = n - j$ . Then we obtain  $\ell\ell^{ts}(A_n) - 1 = (n - j) - 1 = n - j - 1 \in \text{OSpec}(A_n)$ .

The above three items yield  $\{0, 1, 2, \dots, n-1\} \subseteq \text{OSpec}(A_n)$ .  $\square$

Next, we define some notation to describe the Auslander-Reiten quiver  $\Gamma(A_n)$  of  $A_n$ . Recall that a path  $\wp_{[i,j]} = a_i a_{i+1} \dots a_j$  ( $j \geq i$ ) on  $\vec{\mathbb{A}}_n$  corresponds to an indecomposable representation  $M_{[i,j]} = ((M_{[i,j]})_k, (M_{[i,j]})_\alpha)_{k \in (\vec{\mathbb{A}}_n)_0, \alpha \in (\vec{\mathbb{A}}_n)_1}$  satisfying

$$(M_{[i,j]})_k = \begin{cases} \mathbf{k}, & \text{if } i \leq k \leq j+1; \\ 0, & \text{otherwise,} \end{cases} \quad \text{and } (M_{[i,j]})_\alpha = \begin{cases} \mathbf{1}_{\mathbf{k}}, & \text{if } \alpha \in \{a_i, a_{i+1}, \dots, a_j\}; \\ 0, & \text{otherwise.} \end{cases}$$

Then each irreducible morphisms in  $\text{mod}A_n$  is one of the following form:

- the irreducible monomorphism  $f_{[i,j]}^+ : M_{[i,j]} \rightarrow M_{[i-1,j]}$  ( $2 \leq i \leq n$ ),
- the irreducible epimorphism  $f_{[i,j]}^- : M_{[i,j]} \rightarrow M_{[i,j-1]}$  ( $2 \leq j \leq n$ );

and in this case,  $\Gamma(A_n) = (\Gamma(A_n)_0, \Gamma(A_n)_1, \mathfrak{s}, \mathfrak{t})$ , where


- $\Gamma(A_n)_0 = \{M_{[i,j]} \mid 1 \leq i \leq j \leq n\}$ ;
- $\Gamma(A_n)_1 = \{f_{[i,j]}^+ \mid 2 \leq i \leq n, 1 \leq j \leq n\} \cup \{f_{[i,j]}^- \mid 1 \leq i \leq n, 2 \leq j \leq n\}$ ;
- $\mathfrak{s}$  and  $\mathfrak{t}$  respectively send any arrow  $f_{[-,-]}^\pm$  to its starting and ending points, see Figure 5.1

For any  $1 \leq m < n$ , let

$$T_m = \left( \bigoplus_{1 \leq j \leq m} M_{[1,j]} \right) \oplus \left( \bigoplus_{i \in (\vec{\mathbb{A}}_n)_0} M_{[i,i]} \right).$$

The computing is divided to two steps as follows:

Step 1: describe all irreducible  $T_m$ -coghosts (see Proposition 5.4);

FIGURE 5.1. The Auslander-Reiten quiver of  $A_n$ 

Step 2: show that  $\text{sup OSpec}(A_n) = n - 1$ .

Thus, we can obtain the Orlov spectrum of  $A_n$  is  $\text{OSpec}(A_n) = \{1, 2, \dots, n - 1\}$  by using Proposition 5.1.

**Lemma 5.2.** All  $f_{[i,n]}^+ : M_{[i,n]} \rightarrow M_{[i-1,n]}$  ( $m + 1 \leq i \leq n$ ) are  $T_m$ -coghosts.

*Proof.* First of all, for any  $M_{[j,j]}$ , we have  $M_{[j,j]}$  is isomorphic to the simple module  $S(j)$  corresponded by  $j \in (\vec{\mathbb{A}}_n)_0$ , and so the following statements hold.

- (1) If  $j \leq i - 1$  then  $\text{Hom}_{A_n}(M_{[i,n]}, M_{[j,j]}) = 0$ .
- (2) If  $j \geq i$ , then  $\text{Hom}_{A_n}(M_{[i-1,n]}, M_{[j,j]}) = 0$ .

The proof of (1) is trivial. Next we prove (2). For any  $1 \leq p < q \leq n$  and  $p < r \leq q$ , the morphism  $h = (h_v)_{v \in (\vec{\mathbb{A}}_n)_0} \in \text{rep}(\vec{\mathbb{A}}_n)$  from the representation  $((M_{[p,q]})_v, (M_{[p,q]})_\alpha)_{v \in (\vec{\mathbb{A}}_n)_0, \alpha \in (\vec{\mathbb{A}}_n)_1}$  of  $M_{[p,q]}$  to the simple representation of  $S(r)$  is a family of  $\mathbf{k}$ -linear maps such that the following diagram commutes.

$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & (M_{[p,q]})_{r-1} & \xrightarrow{d_{r-1}} & (M_{[p,q]})_r & \xrightarrow{d_r} & (M_{[p,q]})_{r+1} \longrightarrow \cdots \\
 & & \downarrow & & \downarrow h_r & & \downarrow \\
 \cdots & \longrightarrow & 0 & \longrightarrow & S(j)e_r & \longrightarrow & 0 \longrightarrow \cdots
 \end{array}$$

( $e_r$  is the path of length zero corresponded by the vertex  $r$  of  $(\vec{\mathbb{A}}_n)_0$ ). Then  $r - 1 \geq p$ , and so  $(M_{[p,q]})_{r-1} = \mathbf{k}$ . Since  $p \neq q$ , we obtain  $(M_{[p,q]})_r = \mathbf{k}$ , and so  $d_{r-1} = 1_{\mathbf{k}}$ . By the commutativity of the above diagram, we obtain  $h_r = 0$ . Take  $p = i - 1, q = n$  and  $r = j \geq i$ . Then any  $h \in \text{Hom}_{A_n}(M_{[i-1,n]}, M_{[j,j]})$  is zero holds for all  $p = i - 1 < r = j \leq q = n$ . Thus, the statement (2) holds. Then, by Lemma 2.18, every  $f_{[i,n]}^+$  ( $m + 1 \leq i \leq n$ ) is an  $M_{[j,j]}$ -coghost ( $j \in (\vec{\mathbb{A}}_n)_0$ ). Furthermore, we obtain that it is also a  $(\bigoplus_{j \in (\vec{\mathbb{A}}_n)_0} M_{[j,j]})$ -coghost by Lemma 2.19.

Next, we prove that every  $f_{[i,n]}^+$  ( $m + 1 \leq i \leq n$ ) is a  $(\bigoplus_{1 \leq j \leq n} M_{[1,j]})$ -coghost ( $1 \leq j \leq m$ ). For any  $g \in \text{Hom}_{A_n}(M_{[i-1,n]}, M_{[1,j]})$ , note that  $i > j$ , then it is obvious that  $g$  is zero. In this case,  $gf_{[i,n]}^+ = 0$ , then  $f_{[i,n]}^+$  is an  $M_{[1,j]}$ -coghost by Lemma 2.18. Thus,  $f_{[i,n]}^+$  is also a  $(\bigoplus_{1 \leq j \leq n} M_{[1,j]})$ -coghost by Lemma 2.19.



Finally, using Lemma 2.19 again, we have  $f_{[i,n]}^+$  is a  $T_m$ -coghost.  $\square$

**Lemma 5.3.** *If  $f_{[r,s]}^+$  ( $s \geq 1$ ) is a  $T_m$ -coghost, then so is  $f_{[r,s-1]}^+$ .*

*Proof.* For any morphism  $h : M_{[r-1,s-1]} \rightarrow T'$  with  $T' \in \text{add } T_m$ , we have

$$(h \circ f_{[r-1,s]}^-) \circ f_{[r,s]}^+ = 0$$

since  $h \circ f_{[r-1,s]}^-$  is a morphism in  $\text{Hom}_{A_n}(M_{[r-1,s]}, T')$  and  $f_{[r,s]}^+$  is a  $T_m$ -coghost. Then

$$\begin{array}{ccc} & M_{[r-1,s]} & \\ f_{[r,s]}^+ \nearrow & & \searrow f_{[r-1,s]}^- \\ M_{[r,s]} & & M_{[r-1,s-1]} \\ f_{[r,s]}^- \searrow & & \nearrow f_{[r,s-1]}^+ \\ & M_{[r,s-1]} & \end{array}$$

FIGURE 5.2. Anti-commutative block in  $\Gamma(\text{mod } A_n)$

$0 = h \circ f_{[r-1,s]}^- \circ f_{[r,s]}^+ = -h \circ f_{[r,s-1]}^+ \circ f_{[r,s]}^-$  (see Figure 5.2). Since  $f_{[r,s]}^-$  is an epimorphism, we obtain  $hf_{[r,s-1]}^+ = 0$ .  $\square$

The following result is given by Lemmas 5.2 and 5.3, which provide all irreducible  $T_m$ -coghosts.

**Proposition 5.4.** *The morphisms  $f_{[i,n-t]}^+$  ( $m+1 \leq i \leq n$ ,  $0 \leq t < n$ ) constitute all irreducible  $T_m$ -coghosts in  $\text{mod } A_n$ .*

*Proof.* Let  $\mathfrak{F} = \{f_{[i,n-t]}^+ \mid m+1 \leq i \leq n, 0 \leq t < n\}$ . We show that a homomorphism  $f$  is an irreducible  $T_m$ -coghost if and only if  $f \in \mathfrak{F}$ .

First of all, for any  $1 \leq t < j$  ( $\leq m$ ),  $f_{[t+1,j]}^+ : M_{[t+1,j]} \rightarrow M_{[t,j]}$  (see the irreducible morphisms  $f_{[-,-]}^+$  in the part A of Figure 5.3) is not a  $T_m$ -coghost. Indeed, every  $f_{[i,j]}^+$  ( $2 \leq i \leq j$ ) is monomorphic, then  $h = \prod_{i=2}^t f_{[i,j]}^+ : M_{[t,j]} \rightarrow M_{[1,j]}$  and  $hf_{[t+1,j]}^+$  are monomorphic, and then  $hf_{[t+1,j]}^+ \neq 0$ . Since  $M_{[1,j]} \in \text{add } T$ ,  $f_{[t+1,j]}^+$  is not a  $T_m$ -coghost.

Secondly, for any  $f_{[r,s]}^+$ , if  $f_{[r,s]}^+$  is not a  $T_m$ -coghost, then by Lemma 5.3, we obtain that  $f_{[r,s+1]}^+$  is not a  $T_m$ -coghost. Thus any irreducible morphisms  $f_{[-,-]}^+$  in the part B of Figure 5.3. Therefore,  $f_{[r,s]}^+$  is an irreducible  $T_m$ -coghost if and only if  $f_{[r,s]}^+ \in \mathfrak{F}$ .

Finally, any  $f_{[r,s]}^- : M_{[r,s]} \rightarrow M_{[r,s-1]}$  is not a  $T_m$ -coghost because  $\epsilon_{[r,s-1]} f_{[r,s]}^- \neq 0$ , where  $\epsilon_{[r,s-1]} : M_{[r,s-1]} \rightarrow \text{top}(M_{[r,s-1]})$  is the natural epimorphism from  $M_{[r,s-1]}$  to its top  $\text{top}(M_{[r,s-1]}) \cong M_{[r,r]} \in \text{add } T_m$ . Thus,  $f_{[r,s]}^- \notin \mathfrak{F}$ .

In conclusion,  $f$  is an irreducible  $T_m$ -coghost if and only if  $f \in \mathfrak{F}$ .  $\square$


**Lemma 5.5.** *For any  $n$  homomorphisms  $f_i : X_i \rightarrow X_{i+1}$  ( $1 \leq i \leq n$ ) in  $\text{rad}_{A_n}(X_i, X_{i+1})$ , we have  $f_n \circ \cdots \circ f_2 \circ f_1 = 0$ .*

*Proof.* Consider the Auslander algebra  $A_n^{\text{Aus}} = \mathbf{k}\Gamma(\text{mod } A_n)/\mathcal{I}(\text{mod } A_n)$  of  $A_n$ , where  $\mathcal{I}(\text{mod } A_n)$  is generated by

- (1) the zero relations  $f_{[t-1,t]}^- f_{[t,t]}^+$  ( $1 < t \leq n$ )
- (2) and the commutative relations  $f_{[r-1,s]}^- f_{[r,s]}^+ + f_{[r,s]}^- f_{[r,s-1]}^+$  ( $2 \leq r \leq n-1$ ,  $3 \leq s \leq n$ ).

Any morphism  $f_i$  is an element in  $A_n^{\text{Aus}}$ , then there are  $k_{i1}, \dots, k_{it_i} \in \mathbf{k}$  such that

$$f_i = \sum_{u=1}^{t_i} k_{iu} \partial_{iu},$$

FIGURE 5.3. The Auslander-Reiten quiver of  $A_n$  ( $T_m$ -coghosts)

where every  $\varphi_{iu}$  is a path on the bound quiver  $(\Gamma(\text{mod} A_n), \mathcal{I}(\text{mod} A_n))$  of  $A_n^{\text{Aus}}$  whose length  $\ell(\varphi_{iu})$  satisfies the following conditions:

- $\ell(\varphi_{iu}) \geq 1$  because  $f_i \in \text{rad}_{A_n}(X_i, X_{i+1}) \subseteq \text{rad}(A_n^{\text{Aus}})$ ;
- and  $\ell(\varphi_{iu}) \leq n - 1$  because  $f_{[2,n]}^+ \circ \cdots \circ f_{[n,n]}^+$  and  $f_{[1,2]}^- \circ \cdots \circ f_{[1,n]}^-$  are the longest paths in  $(\Gamma(\text{mod} A_n), \mathcal{I}(\text{mod} A_n))$  whose length are  $n - 1$ .

Therefore,  $f_n \cdots f_1 \in \text{rad}^n(A_n^{\text{Aus}}) = 0$ , to be more precise, we have

$$f_n \circ \cdots \circ f_2 \circ f_1 = \sum_{(u_1, \dots, u_n) \in (\mathbb{N}_{\leq n}^+)^{\times n}} \prod_{i=1}^n k_{iu_i} \varphi_{iu_i}.$$

Since  $\ell(\varphi_{nu_n} \cdots \varphi_{1u_1})$  equals to either  $\sum_{i=1}^n \ell(\varphi_{iu_i}) (\geq n)$  or zero, we have  $\varphi_{nu_n} \cdots \varphi_{1u_1} = 0$ , and so  $f_n \circ \cdots \circ f_1 = 0$ .  $\square$

**Proposition 5.6.**  $\text{sup}(\text{OSpec}(A_n)) = n - 1$ .

*Proof.* We have  $n - 1 \in \text{OSpec}(A_n)$  by Proposition 5.1. Next, assume that there is an integer  $y \geq n$  such that  $y \in \text{OSpec}(A_n)$ , then there is a strong generator  $T$  satisfying  $[T]_y = \text{mod}(A_n)$ , and  $[T]_y \setminus [T]_{n-1}$  contains at least one non-zero  $A_n$ -module  $Y$  such that there exists a homomorphism  $g \in \text{Gh}_T^y(X, Y)$  which is not zero, where  $X$  is a non-zero  $A_n$ -module (see Coghost Lemma 2.13). Therefore, we can find  $y$   $T$ -coghosts

$$f_1 : X = X_1 \rightarrow X_2, f_2 : X_2 \rightarrow X_3, \dots, f_y : X_y \rightarrow X_{y+1} = Y$$

satisfying the following two conditions:

- all  $f_i$  are not isomorphisms (by Lemma 2.17);
- $g = f_y \circ \cdots \circ f_2 \circ f_1 \neq 0$ .

Consider the Auslander algebra

$$A_n^{\text{Aus}} = \text{End}_{A_n} \left( \bigoplus_{M \in \text{ind}(\text{mod}(A_n))} M \right)$$

of  $A_n$  whose quiver  $\Gamma(A_n) = (\Gamma(A_n)_0, \Gamma(A_n)_1, \mathfrak{s}, \mathfrak{t})$  is shown in Figure 5.1. Then each homomorphism  $f_i$  can be seen as an element in  $A_n^{\text{Aus}}$ . In the above sense,  $f_i$  is either a

homomorphism lying in  $\text{rad}(A_n^{\text{Aus}})$  or a homomorphism of the form

$$(f_i = (f_{i1} \ f_{i2}) : X_i = X_{i1} \oplus X_{i2} \longrightarrow X_{i+1}) \in \mathbf{k}\Gamma(A_n)_0 \oplus \text{rad}(A_n^{\text{Aus}}),$$

such that  $f_{i1} \in \mathbf{k}\Gamma(A_n)_0$ ,  $f_{i2} \in \text{rad}(A_n^{\text{Aus}})$ . Then for any  $h : X_{i+1} \rightarrow T'$ ,  $T' \in \text{add } T$ , we have

$$h \circ f_i = (h \circ f_{i1} \ h \circ f_{i2}) = 0,$$

since  $f_i$  is a  $T$ -coghost, and then  $h \circ f_{i1} = 0$  and  $h \circ f_{i2} = 0$ . The above follows that  $f_{i1}$  is a  $T$ -coghost, it contradicts with Lemma 2.17.  $f_{i1}$  is an isomorphism defined on some semi-simple module. Then each homomorphism  $f_i$ , as an element in  $A_n^{\text{Aus}}$ , lies in  $\text{rad}(A_n^{\text{Aus}})$ , that is,  $f_i \in \text{rad}_{A_n}(X_i, X_{i+1})$ . It contradicts with Lemma 5.5.  $\square$

**Theorem 5.7.**  $\text{OSpec}(A_n) = \{0, 1, 2, \dots, n-1\}$ .

*Proof.* This theorem is obtained by Propositions 5.1 and 5.6 immediately.  $\square$

### Acknowledgements.

This work is supported by National Natural Science Foundation of China (Grant No. 12171207), Guizhou Provincial Basic Research Program (Natural Science) (Grant No. ZK[2024]YiBan066), and Scientific Research Foundation of Guizhou University (Grant Nos. [2022]53, [2022]65, [2023]16).

### ACKNOWLEDGEMENTS

### REFERENCES

- [1] I. Assem, D. Simson, and A. Skowronski. *Elements of the Representation Theory of Associative Algebras: Volume 1: Techniques of Representation Theory*. Cambridge University Press, 2006.
- [2] I. Assem, A. Skowroski, and S. Trepode. The representation dimension of a selfinjective algebra of Euclidean type. *Journal of Algebra*, 459:157–188, 2016.
- [3] M. Auslander. Representation dimension of artin algebras. *Selected works of Maurice Auslander*, 1:505–574, 1999.
- [4] M. Auslander, I. Reiten, and S. O. Smalø. *Representation theory of Artin algebras*, volume 36. Cambridge university press, 1997.
- [5] M. Ballard, D. Favero, and L. Katzarkov. Orlov spectra: bounds and gaps. *Inventiones mathematicae*, 189(2):359–430, 2012.
- [6] A. Beligiannis. Some ghost lemmas, survey for ‘the representation dimension of Artin algebras’, bielefeld. 2008.
- [7] P. A. Bergh, S. Oppermann, and D. A. Jorgensen. The Gorenstein defect category. *The Quarterly Journal of Mathematics*, 66(2):459–471, 2015.
- [8] A. I. Bondal and M. Van den Bergh. Generators and representability of functors in commutative and noncommutative geometry. *Moscow Mathematical Journal*, 3(1):1–36, 2003.
- [9] D. Bravo, M. Lanzilotta, and O. Mendoza. Pullback diagrams, syzygy finite classes and Igusa-Todorov algebras. *Journal of Pure and Applied Algebra*, 223(10):4494–4508, 2019.
- [10] X.-W. Chen, Y. Ye, and P. Zhang. Algebras of derived dimension zero. *Communications in Algebra*, 36(1):1–10, 2008.
- [11] H. Dao and R. Takahashi. The radius of a subcategory of modules. *Algebra & Number Theory*, 8(1):141–172, 2014.
- [12] Y. Han. Derived dimensions of representation-finite algebras. *arXiv preprint arXiv:0909.0330*, 2009.
- [13] F. Huard, M. Lanzilotta, and O. M. Hernández. Layer lengths, torsion theories and the finitistic dimension. *Applied Categorical Structures*, 21(4):379–392, 2013.
- [14] F. Huard, M. Lanzilotta, and O. Mendoza. An approach to the finitistic dimension conjecture. *Journal of Algebra*, 319(9):3918–3934, 2008.
- [15] F. Huard, M. Lanzilotta, and O. Mendoza. Finitistic dimension through infinite projective dimension. *Bulletin of the London Mathematical Society*, 41(2):367–376, 2009.
- [16] Z. Huang and J. Sun. Invariant properties of representations under excellent extensions. *Journal of Algebra*, 358:87–101, 2012.
- [17] Linckelmann and Markus. Finite generation of Hochschild cohomology of Hecke algebras of finite classical type in characteristic zero. *Bulletin of the London Mathematical Society*, 2011.
- [18] S. Oppermann. Lower bounds for Auslander’s representation dimension. *Duke Mathematical Journal*, 148(2):211–249, 2009.
- [19] S. Oppermann and J. Št’ovíček. Generating the bounded derived category and perfect ghosts. *Bulletin of the London Mathematical Society*, 44(2):285–298, 2012.

- [20] D. Orlov, Remarks on generators and dimensions of triangulated categories, *Moscow Mathematical Journal*, 9(1), 153159 (2009)
- [21] R. Rouquier. Representation dimension of exterior algebras. *Inventiones mathematicae*, 165(2):357–367, 2006.
- [22] R. Rouquier. Dimensions of triangulated categories. *Journal of K-theory*, 1(2):193–256, 2008.
- [23] J. Wei. Finitistic dimension and Igusa–Todorov algebras. *Advances in Mathematics*, 222(6):2215–2226, 2009.
- [24] J. Zhang and J. Zheng, Extension dimensions of derivd and stable equivalent algebras, *J. Algebra* **646** (2024) 17-48.
- [25] J. Zheng, X. Ma, and Z. Huang. The extension dimension of Abelian categories. *Algebras and Representation Theory*, 23(3):693–713, 2020.
- [26] J. Zheng and Z. Huang. An upper bound for the dimension of bounded derived categories. *Journal of Algebra*, 556:1211–1228, 2020.
- [27] J. Zheng and Z. Huang. The derived and extension dimensions of Abelian categories. *Journal of Algebra*, 606:243–265, 2022.
- [28] J. Zheng and Z. Huang. The finitistic dimension and chain conditions on ideals. *Glasgow Mathematical Journal*, 64(1):3744, 2022.
- [29] J. Zheng, L. Tian, Q. Shu and J. Zhang. The Extension dimension of syzygy module categories. arXiv: 2405.02921, 2024.