DIMENSION FORMULAS FOR MODULAR FORM SPACES OF RATIONAL WEIGHTS, THE CLASSIFICATION OF ETA-QUOTIENT CHARACTERS AND AN EXTENSION OF MARTIN'S THEOREM

XIAO-JIE ZHU

ABSTRACT. We give an explicit formula for dimensions of spaces of rational-weight modular forms whose multiplier systems are induced by eta-quotients of fractional exponents. As the first application, we give series expressions of Fourier coefficients of the *n*-th root of certain infinite *q*products. As the second application, we extend Yves Martin's list of multiplicative holomorphic eta-quotients of integral weights by first extending the meaning of multiplicativity, then identifying one-dimensional spaces, and finally applying Wohlfahrt's extension of Hecke operators. A table containing 2277 of such eta-quotients is presented. As a related result, we completely classify the multiplier systems induced by eta-quotients of integral exponents. For instance, there are totally 384 such multiplier systems on $\Gamma_0(4)$ for any fixed weight. We also provide SageMath programs on checking the theorems and generating the tables.

Contents

2
2
4
5
6
8
12
18
22
25
33
35
37
39
39

2020 Mathematics Subject Classification. Primary 11F12; Secondary 11F20, 11F25, 11F30, 11L05, 30F10.

Key words and phrases. modular form, dimension formula, rational weight, Dedekind eta function, Hecke operator, multiplicative eta-quotient.

This work is supported in part by Science and Technology Commission of Shanghai Municipality (No. 22DZ2229014).

9. Application: an extension of Martin's list of multiplicative eta-quotients	44
9.1. The algorithm for finding admissible eta-quotients	45
9.2. The table of admissible eta-quotients of type I	48
9.3. Generalized double coset operators	63
9.4. Identities involving Fourier coefficients of weakly admissible eta-quotients	72
10. Miscellaneous observations and open questions	82
Appendix A. Usage of SageMath code	83
References	83

1. INTRODUCTION

1.1. **Dimension formulas.** A holomorphic function f defined on the upper half plane \mathfrak{H} is called a modular form if it satisfies two conditions: first $f\left(\frac{a\tau+b}{c\tau+d}\right) = (c\tau+d)^k \chi\left(\begin{smallmatrix}a&b\\c&d\end{smallmatrix}\right) f(\tau), \tau \in \mathfrak{H}$ for all $\begin{pmatrix}a&b\\c&d\end{smallmatrix}$ in some discrete matrix group G, k being an integer or half integer, called the weight, and χ being a multiplier system of G; second f is holomorphic at all cusps of G, the meaning of which will be clarified in Definition 2.1. Let the space of all modular forms with given G, k and χ be denoted by $M_k(G, \chi)$.

Modular forms, spaces of modular forms and their variants play important roles in mathematics. In the study of these topics, having an explicit formula for the dimension of the complex vector space $M_k(G,\chi)$ is crucial to some applications. For instance, $\dim_{\mathbb{C}} M_k(G,\chi) = 1$ would imply that any function f in $M_k(G,\chi)$ is an eigenfunction for a family of operators called the Hecke operators and hence we obtain nontrivial relations among the Fourier coefficients of f.

A natural and elegant way for computing $\dim_{\mathbb{C}} M_k(G, \chi)$ is to use the Riemann-Roch theorem and the Riemann-Hurwitz formula. Petersson [1, p. 194] applied this method and gave a dimension formula which he called the generalized Riemann-Roch theorem. Petersson's formula concerns arbitrary Fuchsian group G of the first kind, arbitrary complex weight k and arbitrary multiplier system χ while this formula is not so explicit in the sense one can not directly compute the dimension using elementary operations and numerical information about G and χ . To our best knowledge, Shimura [2, Section 2.4] gave explicit formulas for G arbitrary, k being an integer and χ trivial first. Another way for computing $\dim_{\mathbb{C}} M_k(G, \chi)$ is to use the Eichler-Selberg trace formula for Hecke operators; c.f. [3, 4] for the case G being the full modular group, χ trivial and [5] for the case $G = \Gamma_0(N)$ and χ induced by a Dirichlet character modulo N, both dealing with integral weights. See also [6]. For the half-integral weights, Cohen and Oesterlé [7] gave an explicit dimension formula for the case $G = \Gamma_0(4N)$, χ being the product of the multiplier system of the theta series $\sum_{n \in \mathbb{Z}} e^{2\pi i n^2 \tau}$ and that of a Dirichlet character modulo 4N. Since then, there appear many excellent works on computing $\dim_{\mathbb{C}} M_k(G, \chi)$ for certain special G or special χ , or computing the dimensions of certain subspaces; c.f. [8], [9], [10] and [11].

Modular forms of rational weights are less attractive to mathematicians than those of integral or half-integral weights. We know little about them. For instance, it seems no Hecke theory has been attempted. (Maybe there is no such theory.) However, there actually exist many such forms and more importantly, there are interesting applications, e.g., to noncongruence modular forms. Explicit dimension formulas for spaces of rational weights have been obtained by Ibukiyama [12, Lemma 1.7], [13, p. 5] in some special cases, using essentially the method of Petersson [1], but no proof is given.

As our main result, we give explicit formulas for $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi)$ where k is any rational number, χ is the multiplier system of any level N eta-quotient $\prod_{n|N} \eta(n\tau)^{r_n}$ of fractional exponents r_n . See Definition 2.1 for the exact meaning of $M_k(\Gamma_0(N), \chi)$ and (6) for that of χ . The proof is based on Petersson's method [1] and we provide the full details. The following theorem is a special case of Theorem 4.2, which is our main theorem.

A special case of the Main Theorem. If $k > 2 - \frac{6}{m}\varepsilon_2 - \frac{8}{m}\varepsilon_3 - \frac{12}{m}\sum_{c|N}\phi(c, N/c) \cdot (1 - \{\frac{x_c}{24}\})$, then

(1)
$$\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) = \frac{k-1}{12}m + \frac{1}{4}\varepsilon_2 + \frac{1}{3}\varepsilon_3 + \sum_{c|N} \phi(c, N/c) \cdot \left(\frac{1}{2} - \left\{\frac{x_c}{24}\right\}\right),$$

where m, ε_2 and ε_3 , depending on N, are given in (9), (10) and (11) respectively and

$$k = \frac{1}{2} \sum_{n \mid N} r_n, \quad x_c = \sum_{n \mid N} \frac{N}{(N, c^2)} \cdot \frac{(n, c)^2}{n} r_n \quad \text{for } c \mid N.$$

Note that the notation $\phi(c, N/c)$ refers to the Euler totient ϕ of the greatest common divisor of c and N/c.

In particular, if $k \geq 2$, then (1) always holds. For any rational k with 0 < k < 2, there are also infinitely many pairs (N, χ) such that $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi)$ can be computed using (1). See Section 7 for complete lists of such pairs (N, χ) for k = 1/2, 1, 3/2 and r_n being integers. The dimensions for k = 1/2 and k = 3/2 in the case $G = \Gamma_0(4N)$ and χ is the product of the multiplier system of the theta series $\sum_{n \in \mathbb{Z}} e^{2\pi i n^2 \tau}$ and that of a Dirichlet character modulo 4N are previously known: Serre and Starks [14] gave explicit bases of the spaces of weight 1/2; Cohen and Oesterlé [7] established relations between spaces of weight 1/2 and those of weight 3/2. The important and interesting problem of computing dimensions in weight 1 is difficult and the most complete result about this was obtained by Deligne and Serre [15]. Using their theory, one can compute $\dim_{\mathbb{C}} M_1(\Gamma_0(N), \chi)$ for any $N \in \mathbb{Z}_{\geq 1}$ and χ induced by any Dirichlet character modulo N although there is no explicit formula.

The difference between our formulas for $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi)$ with $k \in \frac{1}{2}\mathbb{Z}$ and others' is that we deal with those χ induced by eta-quotients while the existing formulas are about those χ induced by Dirichlet characters or the multiplier system of $\sum_{n \in \mathbb{Z}} e^{2\pi i n^2 \tau}$. Moreover, when $k \in \mathbb{Q} \setminus \frac{1}{2}\mathbb{Z}$, our formulas include the one in [12, Lemma 1.7] and [13, p. 5] as special cases.

After preparing, stating and proving the main theorem and establishing some related tools (Proposition 5.1, Theorem 6.8, Tables 1, 2 and Section 7.3), we will give mainly two applications, both of which are based on identifying one-dimensional spaces. The first application concerns rational-weight modular forms and the second concerns ordinary eta-quotients whose exponents r_n are all integers.

1.2. Application I. We give a series expression of the Fourier coefficients of the n-th root of certain infinite q-products. See Corollary 8.8 for the details. To avoid overlapping with the main text, here we give a randomly chosen example which is different from Examples 8.9 and 8.10.

Proposition 1.1. We have

$$\frac{\sqrt[3]{\prod_{n\in\mathbb{Z}_{\geq 1}}(1-q^n)}\cdot\sqrt{\prod_{n\in\mathbb{Z}_{\geq 1}}(1-q^{2n})^{29}}}{\sqrt[3]{\prod_{n\in\mathbb{Z}_{\geq 1}}(1-q^{4n})^{22}}} = 1 + e^{-\frac{15\pi i}{8}}\frac{(2\pi)^{\frac{15}{4}}}{\Gamma(\frac{15}{4})}}{\cdot \sum_{n\in\mathbb{Z}_{\geq 1}}n^{\frac{11}{4}}} \left(\sum_{c\in\mathbb{Z}_{\geq 1}}\frac{1}{(4c)^{\frac{15}{4}}}\sum_{\substack{0\leq d<4c\\(d,4c)=1}}\mathfrak{e}\left(\frac{dn}{4c} - \frac{1}{24}P(4c,d;\frac{29}{2},-\frac{22}{3})\right)\right)\cdot q^n,$$

where $q = e^{2\pi i \tau}$ throughout the paper, $\tau \in \mathfrak{H}$ and

$$P(4c,d;\frac{29}{2},-\frac{22}{3}) = -88(-d,c) + 174(-d,2c) + 4s(-d,4c) - \frac{45}{2},$$
$$s(-d,c) = \sum_{r=1}^{c-1} \frac{r}{c} \left(\left\{\frac{-dr}{c}\right\} - \frac{1}{2}\right).$$

Corollary 8.8 contains infinitely many such identities of level 4, extending a previous result of the author [16, Theorem 7.1] which concerns levels 2 and 3. For the outline of the proof, say of Proposition 1.1, let the infinite product be f, which is an eta-quotient of fractional exponents. Then $f \in M_{15/4}(\Gamma_0(4), \chi)$ where χ is the multiplier system of f. There is as well an Eisenstein series g in $M_{15/4}(\Gamma_0(4), \chi)$; c.f. Definition 8.3. Now we apply Theorem 4.2 and find that $\dim_{\mathbb{C}} M_{15/4}(\Gamma_0(4), \chi) = 1$. Hence f and g are proportional. The identity thus follows by figuring out the Fourier coefficients of g. 1.3. Application II. This is the major application. In 1996, Martin [17] obtained the complete $list^1$ of integral-weight holomorphic eta-quotients f (with integral exponents) satisfying that

- (a) the multiplier system of f is induced by some Dirichlet character,
- (b) if we write $f = \sum_{n \in \mathbb{Z}_{\geq 0}} c_f(n)q^n$, then $c_f(n_1)c_f(n_2) = c_f(n_1n_2)c_f(1)$ for any coprime positive integers.

We will discard condition (a) and seek for arbitrary holomorphic eta-quotients (with integral exponents) satisfying a multiplicativity property similar to the one in condition (b). We will also give many interesting new identities involving Fourier coefficients of these eta-quotients. Let us first present some randomly chosen examples. For the basic knowledge about Dedekind eta function $\eta(\tau)$ and eta-quotients, see the second half of Section 2.

Example. Let $f(\tau) = \eta(\tau)^{-7} \eta(2\tau)^{17} \eta(4\tau)^{-3} = \sum_{n \in \frac{5}{8} + \mathbb{Z}_{\geq 0}} c_f(n) q^n$. Let l_1, l_2 be coprime odd square-free integers. Then

$$c_f\left(\frac{5l_1^2}{8}\right)c_f\left(\frac{5l_2^2}{8}\right) = c_f\left(\frac{5l_1^2l_2^2}{8}\right).$$

See Proposition 9.33 for details and the proof.

Example. Let $f(\tau) = \eta(3\tau)^2 \eta(9\tau)^{-1} \eta(27\tau) = \sum_{n \in \mathbb{Z}_{\geq 1}} c_f(n)q^n$. Then $c_f(l_1)c_f(l_2) = c_f(l_1l_2)$ whenever $l_1, l_2 \equiv 1 \mod 3$ are square-free positive integers with $(l_1, l_2) = 1$. See Example 9.29 for details.

Example. Let
$$f(\tau) = \eta(\tau)^1 \eta(2\tau)^{-1} \eta(3\tau)^{-1} \eta(4\tau)^1 \eta(6\tau)^4 \eta(12\tau)^{-2} = \sum_{n \in \mathbb{Z}_{\geq 0}} c_f(n) q^n$$
. Then
 $c_f(l) = -\sum_{a|l} \left(\frac{a}{6}\right) = -\prod_{p|l} \left(1 + \left(\frac{p}{6}\right)\right)$

where $1 \leq l \equiv 1 \mod 24$ is square-free. Note that $\left(\frac{a}{6}\right)$ refers to the Kronecker-Jacobi symbol. See Example 9.23 for details.

Martin [17] proved his results using Hecke operators in the sense of [18] while we prove our results using Wohlfahrt's extension of Hecke operators [19]. See also [20, Section 3] for a theory of the double coset version of these operators. These operators are denoted by T_l , where l runs through a multiplicative submonoid L of the positive integers; c.f. Corollary 9.13. As a prerequisite, we give an explicit formula for the action on Fourier coefficients; c.f. Theorem 9.15. Then we aim to find holomorphic eta-quotient f such that $T_l f = c_l \cdot f$ with $c_l \in \mathbb{C}$. At this point Theorem 4.2 enters into play. If f, whose level is N and character is χ , lies in a one-dimensional space $M_k(\Gamma_0(N), \chi)$ or $S_k(\Gamma_0(N), \chi)$, the subspace of cusp forms, then we will have $T_l f = c_l \cdot f$ for all $l \in L$. (L depends on f.) This fact, which is stated in Theorem 9.19, is the main theorem of this application.

¹There is a misprint in [17, p. 4853]. The entry $2^{-1} \cdot 4^4 \cdot 6^{-1} \cdot 8^{-1} \cdot 12^4 \cdot 24$ should be corrected to $2^{-1} \cdot 4^4 \cdot 6^{-1} \cdot 8^{-1} \cdot 12^4 \cdot 24^{-1}$.

It seems that there are more than 10000 eta-quotients that are Hecke eigenforms (in the sense of (83)) for infinitely many l. We list 2277 of them in Table 3. They are what we call admissible eta-quotients of type I with levels

$$N = 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 17, 19, 21, 27$$

For the meaning of admissible eta-quotients of type I, see the beginning part of Section 9. For other levels, either there is no admissible eta-quotient of type I, or there are too many so it is not appropriate to list them in the paper. One can find the SageMath code that generates admissible eta-quotients of type I or II with any given level in Appendix A.

Besides Theorem 9.19, another important result is Theorem 9.24, which says that

$$c_{l_1} \cdot c_{l_2} = c_{l_1 l_2}, \quad l_1, l_2 \in L, \quad (l_1, l_2) = 1.$$

Here $c_l = T_l f/f$. This is the genuine multiplicativity property that is satisfied by all admissible eta-quotients and reduces to the relation $c_f(n_1)c_f(n_2) = c_f(n_1n_2)c_f(1)$ for ordinary multiplicative eta-quotients in [17].

All identities in the above three examples are immediate consequences of Theorems 9.19 and 9.24.

1.4. Other results, structure of the paper and notations. As a related result, we give the complete classification of linear characters that are induced by eta-quotients $\prod_{n|N} \eta(n\tau)^{r_n}$ with $r_n \in \mathbb{Z}$ on the double cover of $\Gamma_0(N)$. See Corollary 6.9 for the conclusion and Examples 6.10, 6.11, 6.12, 6.13, 6.14 for examples. For instance, for each $k \in \frac{1}{2}\mathbb{Z}$, there are totally 384 linear characters of the double cover of $\Gamma_0(4)$ that are induced by $\eta(\tau)^{r_1}\eta(2\tau)^{r_2}\eta(4\tau)^{r_n}$ of weight k. As a comparison, there are only two characters of $\Gamma_0(4)$ that are induced by Dirichlet characters modulo 4 and these two characters are also contained in the above 384 characters with any fixed $k \in \mathbb{Z}$.

The structure of the paper is as follows. We review some elements of rational-weight modular forms in Section 2. The second half of this section contains elements of Dedekind eta function and eta-quotients of fractional exponents. In Section 3 we associate a divisor on certain compact Riemann surface with any rational-weight meromorphic modular form and in addition, we give a detailed proof of the valence formula in the case of rational weights. In Section 4 we state and prove the main theorem via Petersson's method [1]. The following three sections contain some tools which are needed by the following two applications: Section 5 contains a formula that relates the orders at cusps and the exponents (or the character) of an eta-quotient. This formula is due to Bhattacharya (cf. [21, eq. (5.13)]). Here we give more details on the proof. In Section 6 we classify all linear characters that are induced by eta-quotients $\prod_{n|N} \eta(n\tau)^{r_n}$ with $r_n \in \mathbb{Z}$ on the double cover of $\Gamma_0(N)$, as is described in the last paragraph. Section 7, which is subdivided into three subsections, contains tables of dimensions of modular form spaces of weight 1/2, 1 and 3/2 that can be computed by our main theorem. In Sections 8 and 9 we carry out the first and second applications, which has been described in Sections 1.2 and 1.3, respectively. Section 10 contains some comments, open problems and conjectures. Finally, many formulas in this paper have been verified and many tables are generated by SageMath [22] programs. Appendix A contains the usage of the code.

We collect some notations. For a set A, the symbols |A| and # A both denote its cardinality. The notation \mathbb{Q}^A denotes the \mathbb{Q} -vector space of functions from A into \mathbb{Q} . A multiset $\{x_i: i \in I\}$ is defined to be the ordinary set $\{(y, n): n = \#\{i \in I: x_i = y\}, n \neq 0\}$. Its underlying (ordinary) set is still denoted by $\{x_i: i \in I\}$. If $f: X \to Y$ is a function, then $f|_A$ is its restriction to A where A is a subset of X. If f, g are functions then $f \circ g$ is their composition: $f \circ g(x) = f(g(x))$. For a family of sets $A_i, i \in I$, the notation $\bigsqcup_{i \in I} A_i$ denotes the union and only when the family A_i are disjoint can one use this notation.

For a group G, $\langle X \rangle$ denotes the subgroup generated by X where X is a subset of G. If H is a subgroup of G then $[G: H] = \# G/H = \# H \setminus G$ is the index. If G acts on the left (right resp.) on a set X, then $G \setminus X$ (X/G resp.) denotes the set of orbits $G \cdot x$ ($x \cdot G$ resp.) where $x \in X$. If X is itself a group, G is a subgroup of X and the action is the group operation, then $G \setminus X$ (X/G resp.) is called the left coset space (right coset space resp.) For $x \in X$, G_x denotes the stabilizer, that is, the subgroup of $g \in G$ such that gx = x (or xg = x). A character χ of G is a complex linear character, that is, a group homomorphism from G to \mathbb{C}^{\times} , the multiplicative group of nonzero complex numbers. If $|\chi(g)| = 1$ for all $g \in G$, then χ is called unitary. If G' is another group and $\pi: G \to G'$ is a group surjection, then we say the character χ descends to a character on G' if there is a character χ' of G' such that $\chi = \chi' \circ \pi$.

The symbol ϕ refers to the Euler totient function. For integers a, b, \ldots , the notation (a, b, \ldots) denotes the greatest common divisor and $\phi(a, b, \ldots)$ denotes the value of ϕ at (a, b, \ldots) . Let $N \in \mathbb{Z}_{\geq 1}$; the summation range of $\sum_{n|N}$ is implicitly understood to be the *positive* divisors of N. Let p be a prime and $\alpha \in \mathbb{Z}_{\geq 0}$; then $p^{\alpha} \parallel N$ means $p^{\alpha} \mid N$ but $p^{\alpha+1} \nmid N$. The p-adic exponential valuation, $v_p(N)$, is the largest α such that $p^{\alpha} \parallel N$. An empty sum is understood to be 0 and an empty product be 1. For a real number x, the notation [x] means the largest integer not exceeding x and $\{x\} = x - [x]$. The functions $\Gamma(s)$ and $\zeta(s)$ are the usual Euler Gamma function and Riemann zeta function respectively.

Let $m, n \in \mathbb{Z}$; the Kronecker-Jacobi symbol $\left(\frac{m}{n}\right)$ is defined as follows:

- $\left(\frac{m}{p}\right)$ is the usual Legendre symbol if p is an odd prime.
- $\left(\frac{m}{2}\right)$ equals 0 if $2 \mid m$, and equals $(-1)^{(m^2-1)/8}$ if $2 \nmid m$.
- $\left(\frac{m}{-1}\right)$ equals 1 if $m \ge 0$, and equals -1 otherwise.
- $\left(\frac{m}{1}\right) = 1$ by convention.
- $\left(\frac{m}{n}\right)$ is defined to make it a complete multiplicative function of $n \in \mathbb{Z} \{0\}$.
- $\left(\frac{m}{0}\right) = 0$ if $m \neq \pm 1$, and $\left(\frac{\pm 1}{0}\right) = 1$.

We shall freely use the following properties, especially in the proof of Theorem 9.15.

- $\left(\frac{m}{n_1 n_2}\right) = \left(\frac{m}{n_1}\right) \left(\frac{m}{n_2}\right)$. When m = -1, it is required that $n_1, n_2 \neq 0$. $\left(\frac{m_1 m_2}{n}\right) = \left(\frac{m_1}{n}\right) \left(\frac{m_2}{n}\right)$. When n = -1, it is required that either $m_1, m_2 \neq 0$, or one of m_1, m_2 is 0 and the other is nonnegative.
- $\left(\frac{m}{n}\right)\left(\frac{n}{m}\right) = \varepsilon(m,n) \cdot (-1)^{\frac{n-1}{2} \cdot \frac{m-1}{2}}$ where m,n are coprime odd integers, $\varepsilon(m,n) = -1$ if n, m < 0 and $\varepsilon(m, n) = 1$ otherwise.
- $\left(\frac{-1}{n}\right) = (-1)^{\frac{n-1}{2}}$ and $\left(\frac{2}{n}\right) = (-1)^{\frac{n^2-1}{8}}$ where *n* is odd.
- The function $n \mapsto \left(\frac{m}{n}\right)$ is |m|-periodic if $m \equiv 0, 1 \mod 4$; it is |4m|-periodic if $m \equiv 2 \mod 4$.
- The function $m \mapsto \left(\frac{m}{n}\right)$ is *n*-periodic if *n* is odd and positive.

For the proofs, see [23, Section 2.2.2].

2. Modular forms of rational weight

In this section, we review some elements of the theory of modular forms of rational weight (cf. [16, 13, 24, 25, 12, 1]). The concept of modular forms of rational weight is a special case of that of generalized modular forms which was initiated by Knopp and Mason [25]. The multiplier system of a generalized modular form is not required to be unitary or to have finite order, while in this paper, the multiplier systems of all modular forms that occur are unitary and have finite orders. In addition, it seems that the modular forms of rational weights that occur in this paper can only be regarded as modular forms on non-congruence subgroups of $SL_2(\mathbb{Z})$ with trivial character. As a comparison, Freitag and Hill [26] have recently constructed modular forms of weight 1/3 on SU(2,1) whose levels are certain congruence subgroups of SU(2,1). Furthermore, see [27, Section 2] for harmonic weak Maass forms of real weight and their relations with weakly holomorphic modular forms.

First some notations. Let $\operatorname{GL}_2^+(\mathbb{R})$ be the group of all 2×2 real matrices with positive determinants and $\mathrm{SL}_2(\mathbb{R})$ be the subgroup of $\mathrm{GL}_2^+(\mathbb{R})$ whose elements have determinant 1. Let D be a positive integer. Then we define the *D*-cover of $\operatorname{GL}_2^+(\mathbb{R})$ by

$$\widetilde{\operatorname{GL}_{2}^{+}(\mathbb{R})^{D}} = \left\{ \left(\left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right), \varepsilon \left(c'\tau + d'\right)^{\frac{1}{D}} \right) : \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right) \in \operatorname{GL}_{2}^{+}(\mathbb{R}), \varepsilon^{D} = 1 \right\},$$

where $\begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}$ is the matrix in $SL_2(\mathbb{R})$ proportional to $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and $(c'\tau + d')^{\frac{1}{D}}$ means a function of $\tau \in \mathfrak{H} = \{z \in \mathbb{C} : \mathfrak{I}(z) > 0\}$. The notation $\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \varepsilon (c'\tau + d')^{\frac{1}{D}} \right)$ is sometimes abbreviated to $\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \varepsilon\right)$ when D is understood. The composition is given by

$$\begin{pmatrix} \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix}, \varepsilon_1 (c_1'\tau + d_1')^{\frac{1}{D}} \end{pmatrix} \cdot \begin{pmatrix} \begin{pmatrix} a_2 & b_2 \\ c_2 & d_2 \end{pmatrix}, \varepsilon_2 (c_2'\tau + d_2')^{\frac{1}{D}} \end{pmatrix}$$

$$= \begin{pmatrix} \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix} \begin{pmatrix} a_2 & b_2 \\ c_2 & d_2 \end{pmatrix}, \varepsilon_1 \varepsilon_2 \begin{pmatrix} c_1' \frac{a_2'\tau + b_2'}{c_2'\tau + d_2'} + d_1' \end{pmatrix}^{\frac{1}{D}} (c_2'\tau + d_2')^{\frac{1}{D}} \end{pmatrix}.$$

It can be verified straightforwardly that $\operatorname{GL}_{2}^{+}(\mathbb{R})^{D}$ with this composition is a group. One should note that different choices of holomorphic branches of $(c'\tau + d')^{\frac{1}{D}}$ lead to different *D*-covers. We choose the following branch throughout:

$$z^r = \exp(r\log z), \quad -\pi < \Im(\log z) \le \pi$$

It is possible to make $\widetilde{\operatorname{GL}_2^+}(\mathbb{R})^D$ a topological covering of $\operatorname{GL}_2^+(\mathbb{R})$ and hence $\widetilde{\operatorname{GL}_2^+}(\mathbb{R})^D$ becomes a Lie group whose smooth structure is transposed from $\operatorname{GL}_2^+(\mathbb{R})$ using the above covering map (cf. [28]). The motivation for using *D*-covers is that when dealing with modular forms of weight $k \in \frac{1}{D}\mathbb{Z}$, the multiplier systems become group characters on *D*-covers.

Let G be a subgroup of $\operatorname{GL}_2^+(\mathbb{R})$; then by $\widetilde{G^D}$ we understand the preimage of G under the natural projection $\operatorname{GL}_2^+(\mathbb{R})^D \to \operatorname{GL}_2^+(\mathbb{R}), \ ((\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}), \varepsilon) \mapsto (\begin{smallmatrix} a & b \\ c & d \end{smallmatrix})$. For $\gamma \in \operatorname{GL}_2^+(\mathbb{R})$, let $\widetilde{\gamma}$ denote $(\gamma, 1) \in \operatorname{GL}_2^+(\mathbb{R})^D$. (The cover index D should be inferred from the context.) In addition, we set $\overline{G} = \{g \cdot \{\pm (\begin{smallmatrix} 1 & 0 \\ 0 & 1 \end{smallmatrix})\} \colon g \in G\}$, which is a subgroup of $\overline{\operatorname{GL}_2^+(\mathbb{R})} = \operatorname{GL}_2^+(\mathbb{R})/\{\pm (\begin{smallmatrix} 1 & 0 \\ 0 & 1 \end{smallmatrix})\}$.

The group $SL_2(\mathbb{Z})$ is the set of 2×2 integral matrices of determinant 1 and is known as the *full* modular group. We also need the congruence subgroup

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{SL}_2(\mathbb{Z}) \colon c \equiv 0 \mod N \right\},\$$

where N is a positive integer called the level. It is well known that the index $[SL_2(\mathbb{Z}): \Gamma_0(N)] = N \prod_{p|N} (1+1/p)$ where p denotes a prime (cf. [29, Coro. 6.2.13]). The matrices $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $\begin{pmatrix} N & 0 \\ 0 & 1 \end{pmatrix}$ are denoted by I, T, S, B_N respectively.

Let $\mathscr{M}(\mathfrak{H})$ denote the field of meromorphic functions on the upper half plane \mathfrak{H} and $k \in \frac{1}{D}\mathbb{Z}$. Define the *slash operator* of weight k by

$$f|_{k}\left(\left(\begin{smallmatrix}a&b\\c&d\end{smallmatrix}\right),\varepsilon\right)(\tau) = \left(\varepsilon\sqrt[D]{c'\tau+d'}\right)^{-Dk} f\left(\frac{a\tau+b}{c\tau+d}\right),$$

where $f \in \mathscr{M}(\mathfrak{H})$, $(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \varepsilon) \in \widetilde{\operatorname{GL}_2^+(\mathbb{R})}^D$ and $\tau \in \mathfrak{H}$. It is immediate that $\widetilde{\operatorname{GL}_2^+(\mathbb{R})}^D$ acts on $\mathscr{M}(\mathfrak{H})$ on the right via the slash operator of weight k. Let G be a subgroup of $\operatorname{SL}_2(\mathbb{Z})$ and χ be a character² on \widetilde{G}^D . We say f transforms like a modular form of weight k and with multiplier system (or with character) χ if $f|_k \gamma = \chi(\gamma)f$ for any $\gamma \in \widetilde{G}^D$. If this is the case, and $[\operatorname{SL}_2(\mathbb{Z}): G] < +\infty$ and $\chi(\gamma)$ has finite order for any $\gamma \in \widetilde{G}^D$, then for $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{SL}_2(\mathbb{Z})$ we have

$$f|_{k}(\widetilde{a \ b \atop c \ d})(\tau) = \sum_{n \in m^{-1}\mathbb{Z}} c_n q^n,$$

where $q = \mathfrak{e}(\tau) = \exp(2\pi i \tau)$ and $m \in \mathbb{Z}_{\geq 1}$ provided that there is $Y_0 > 0$ (depending on $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$) such that $f|_k(\overbrace{a & b}{b})$ has no poles on $\mathfrak{H}_{Y_0} = \{\tau \in \mathfrak{H} : \mathfrak{H}(\tau) > Y_0\}$. Moreover, the series converges

 $^{^{2}}$ A character always means a unitary linear character, that is, a group homomorphism to the group of complex numbers of absolute value 1.

normally on \mathfrak{H}_{Y_0} . To see this, note that there exists $m \in \mathbb{Z}_{\geq 1}$ such that $\chi\left(\widetilde{\binom{a \ b}{c \ d}}\widetilde{T}^m \widetilde{\binom{a \ b}{c \ d}}^{-1}\right) = 1$ and hence $f|_{k}(\widetilde{a\ d\ b})(\tau+m) = f|_{k}(\widetilde{a\ d\ b})(\tau)$. Therefore, the desired expansion follows from Fourier's theorem or Laurent's theorem. Define the order $\operatorname{ord}_{i\infty}(f|_k(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \varepsilon))$ to be the least n such that $c_n \neq 0$. If the expansion does not exist (that is, $f|_k(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \varepsilon)$ has a nonisolated singularity at infinity) then the order is undefined; if the expansion holds but for any $n_0 \in \mathbb{Z}$ there exists $n < n_0$ such that $c_n \neq 0$ then define $\operatorname{ord}_{i\infty}(f|_k(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \varepsilon)) = -\infty$; if $f|_k(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \varepsilon)$ is identically zero then define $\operatorname{ord}_{i\infty}(f|_k(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \varepsilon)) = +\infty.$

Definition 2.1. Let $D \in \mathbb{Z}_{\geq 1}$ and $k \in \frac{1}{D}\mathbb{Z}$. Let G be a subgroup of $SL_2(\mathbb{Z})$ of finite index and $\chi: \overline{G^D} \to \mathbb{C}^{\times}$ be a character of finite order³ (that is, $\chi(\gamma)$ has finite order for any $\gamma \in \overline{G^D}$). Let $f \in \mathscr{M}(\mathfrak{H})$. Then we say f is a meromorphic modular form of weight k for G with character χ if $f|_k \gamma = \chi(\gamma) f$ for any $\gamma \in \widetilde{G^D}$ and $\operatorname{ord}_{i\infty}(f|_k \gamma) > -\infty$ for any $\gamma \in \widetilde{\operatorname{SL}}_2(\mathbb{Z})^D$. Suppose this is the case. Then we say f is a weakly holomorphic modular form if it has no poles on \mathfrak{H} ; it is a modular function if the weight k = 0; it is a modular form (cusp form respectively) if it is weakly holomorphic and $\operatorname{ord}_{i\infty}(f|_k\gamma) \geq 0$ (> 0 respectively) for any $\gamma \in \widetilde{\operatorname{SL}}_2(\mathbb{Z})^D$. The vector spaces (over \mathbb{C}) of modular forms and of cusp forms are denoted by $M_k(G,\chi)$ and $S_k(G,\chi)$ respectively. When χ is trivial (which means $k \in 2\mathbb{Z}$ and χ always takes the value 1), we let $M_k(G) = M_k(G, \chi)$ and $S_k(G) = S_k(G, \chi).$

Note that the cover index D can be recovered from the domain of χ and $M_k(G, \chi)$ is also denoted by $M_k(\widetilde{G^D}, \chi)$ when needed.

Remark 2.2. Suppose there exists a nonzero meromorphic modular form f of weight k for G with character χ . Then we have the following facts:

- χ(I, ε(1/D)) = ε(-k).
 If −I ∈ G, then χ(-I) = ε(-k/2).
- If D' is another positive integer with $k \in \frac{1}{D'}\mathbb{Z}$ and $D' \mid D$, then $\chi \colon \widetilde{G^D} \to \mathbb{C}^{\times}$ descends to a character on $\widetilde{G^{D'}}$, i.e., χ can be factored as $\chi = \chi_1 p_{D,D'}$ where $p_{D,D'} \colon \widetilde{G^D} \to \widetilde{G^{D'}}$ is the natural projection $(\gamma, \varepsilon) \mapsto (\gamma, \varepsilon^{D/D'})$ and χ_1 is a character on $\widetilde{G^{D'}}$ (which is unique).

The proof is straightforward by considering slash operators acting on f of which we omit the details. Motivated by the first two facts, we call a character χ on $\widetilde{G^D}$ with the properties $\chi(I, \mathfrak{e}(1/D)) =$ $\mathfrak{e}(-k)$ and $\chi\left(\widetilde{-I}\right) = \mathfrak{e}(-k/2)$ if $-I \in G$ a multiplier system for G of weight k of cover index D. Note that if $-I \in G$, then the second property implies the first one. Finally, note that the third fact still holds when we require that χ is a multiplier system for G of weight k of cover index D even if we can *not* find a nonzero form f.

³If χ is not of finite order as considered by Knopp and Mason [25], this definition also makes sense.

Remark 2.3. If f is a meromorphic modular form of weight k for G with character χ , and g is a meromorphic modular form of weight k' for G with character χ' with $k, k' \in \frac{1}{D}\mathbb{Z}$, then it is immediate that $f \cdot g$ is a meromorphic modular form of weight k + k' for G with character $\chi \cdot \chi'$. Moreover, if f and g are both modular forms (cusp forms respectively), then so is $f \cdot g$.

One of the aims of this paper is to give explicit formulas concerning $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi)$ and $\dim_{\mathbb{C}} S_k(\Gamma_0(N), \chi)$ where $k \in \mathbb{Q}$ and χ is a character on $\Gamma_0(N)^D$ of some kind which we explain now.

We need the Dedekind eta function and its logarithm, which are defined by

(2)
$$\eta(\tau) = q^{1/24} \prod_{n \in \mathbb{Z}_{\geq 1}} (1 - q^n), \quad \tau \in \mathfrak{H}$$
$$\log \eta(\tau) = \log \eta(\mathbf{i}) + \int_{\mathbf{i}}^{\tau} \frac{\eta'(z)}{\eta(z)} \, \mathrm{d}z,$$

where $\log \eta(i)$ is the real logarithm. The transformation equations of $\log \eta$ under $SL_2(\mathbb{Z})$ are obtained by Dedekind (cf. [30, Equation (12), Section 3.4]):

(3)
$$\log \eta \left(\frac{a\tau + b}{c\tau + d}\right) - \log \eta(\tau) = 2\pi i \left(\frac{a+d}{24c} + \frac{1}{2}s(-d,c) - \frac{1}{8}\right) + \frac{1}{2}\log(c\tau + d),$$

where $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$ with c > 0 and s(-d, c) is the Dedekind sum

(4)
$$s(h,k) = \sum_{r \mod k} \left(\left(\frac{r}{k} \right) \right) \cdot \left(\left(\frac{hr}{k} \right) \right), \quad h \in \mathbb{Z}, \ k \in \mathbb{Z}_{\geq 1}$$

In the above definition, ((x)) = x - [x] - 1/2 if $x \in \mathbb{R} \setminus \mathbb{Z}$ and ((x)) = 0 if $x \in \mathbb{Z}$. Set

(5)
$$\Psi: \operatorname{SL}_{2}(\mathbb{Z}) \to \mathbb{Z}$$

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{cases} \frac{a+d}{c} + 12s(-d,c) - 3, & c > 0; \\ \frac{a+d}{c} + 12s(d,-c) + 3, & c < 0; \\ b, & c = 0, a > 0; \\ -b - 6, & c = 0, a < 0. \end{cases}$$

Fix a positive integer D. Then the transformation equations of η follow from that of $\log \eta$ and can be expressed as $\eta|_{1/2}(\gamma, \varepsilon) = \chi_{\eta}(\gamma, \varepsilon)\eta$, where χ_{η} is the multiplier system for $SL_2(\mathbb{Z})$ of weight 1/2 of cover index 2D defined by

$$\chi_{\eta}(\gamma,\varepsilon) = \varepsilon^{-D} \mathfrak{e}\left(\frac{\Psi(\gamma)}{24}\right), \quad (\gamma,\varepsilon) \in \widetilde{\mathrm{SL}_{2}(\mathbb{Z})^{2D}}$$

The fact that $\Psi(\operatorname{SL}_2(\mathbb{Z})) \subseteq \mathbb{Z}$ (when we give (5) this is tacitly assumed) follows from the above formula with D = 1, the fact $\operatorname{SL}_2(\mathbb{Z})$ is generated by T and S and the fact $\chi_\eta(\widetilde{T})^{24} = \chi_\eta(\widetilde{S})^{24} = 1$.

Let $N, D \in \mathbb{Z}_{\geq 1}$. By an eta-quotient of level N and cover index D we understand a product $f(\tau) = \prod_{0 < n \mid N} \eta(n\tau)^{r_n}$ with $r_n \in \mathbb{Q}$ such that $D \cdot r_n \in 2\mathbb{Z}$ for any $n \mid N$. Thus an eta-quotient of cover index 2 is just an ordinary eta-quotient whose exponents are integers. The fractional powers are defined by $\eta(n\tau)^{r_n} = \exp(r_n \log \eta(n\tau))$ (cf. (2)). Since η has no poles on \mathfrak{H} , the eta-quotient f is a weakly holomorphic modular form of weight $k = \frac{1}{2} \sum_{n \mid N} r_n$ for $\Gamma_0(N)^D$ with character

(6)
$$\chi \colon \widetilde{\Gamma_0(N)^D} \to \mathbb{C}^{\times}$$
$$\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \varepsilon \end{pmatrix} \mapsto \varepsilon^{-Dk} \mathfrak{e} \left(\frac{1}{24} \sum_{n|N} r_n \Psi \begin{pmatrix} a & bn \\ c/n & d \end{pmatrix} \right)$$

The above formula can be proved using (3), Remark 2.3 and the fact $\eta(n\tau) = n^{-1/4}\eta|_{1/2}\widetilde{B_n}(\tau)$. Note that η has no zeros on \mathfrak{H} according to its infinite product expansion. Therefore any eta-quotient has no zeros or poles on \mathfrak{H} . For the order at infinity, we have (cf. [16, Lemma 4.2])

(7)
$$\operatorname{ord}_{i\infty}(f|_{k}(\widetilde{\left(\begin{smallmatrix}a & b \\ c & d\end{smallmatrix}\right)}) = \frac{1}{24} \sum_{n|N} \frac{(n,c)^{2}}{n} r_{n}, \quad \left(\begin{smallmatrix}a & b \\ c & d\end{smallmatrix}\right) \in \operatorname{SL}_{2}(\mathbb{Z}).$$

We recommend the reader to see [31, Section 2] or [21, Section 5] for more details on ordinary eta-quotients. If (7) are always nonnegative for all $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$, we say f is a holomorphic eta-quotient.

We will investigate the dimensions $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi)$ and $\dim_{\mathbb{C}} S_k(\Gamma_0(N), \chi)$ where $k \in \frac{1}{D}\mathbb{Z}$ and χ is the character of an eta-quotient of weight k, level N and cover index D in the next three sections.

Remark 2.4. The reader should be warned that when dealing with half-integral weights, our notation may differ from other authors', e.g. from the notation encountered in [32] and [14]. Let ψ be a Dirichlet character modulo 4N and let $2k \in \mathbb{Z}_{\geq 1}$. The space $M_0(4N, k, \psi)$ in Serre and Stark's notation is the same thing as $M_k(\Gamma_0(4N), \chi_1\psi_1)$ in our notation where χ_1 is the character of $\eta(\tau)^{-4k}\eta(2\tau)^{10k}\eta(4\tau)^{-4k}$ and ψ_1 is the character that maps $\left(\begin{pmatrix}a & b \\ c & d\end{pmatrix}, \varepsilon\right)$ to $\psi(d)$. The character χ_1 is hidden in many authors' notation of spaces and it sometimes appears in their definitions of slash operators or modular transformations or automorphic factors.

3. Divisors of modular forms

Our method relies on the classical Riemann-Roch theorem, so we recall some elements here. By a compact Riemann surface X, we understand a compact connected Hausdorff topological space equipped with a maximal atlas $\{(U_i, \phi_i)\}$ (i.e., each U_i is open in X and ϕ_i is a homeomorphism from U_i onto an open set in \mathbb{C}) such that when (U_i, ϕ_i) and (U_j, ϕ_j) overlap, the transition map $\phi_j \circ \phi_i^{-1} \colon \phi_i(U_i \cap U_j) \to \phi_j(U_i \cap U_j)$ is holomorphic. It can be shown that X is second-countable (cf. [33, p. 88]). Therefore, according to the well known classification of compact connected topological surfaces (cf. eg. [34, Theorem 10.22]) and the fact that Riemann surfaces are oriented, X is homeomorphic to the 2-sphere \mathbb{S}^2 or to the connected sum of g copies of the real projective plane \mathbb{P}^2 with $g \in \mathbb{Z}_{\geq 1}$. The number g is called the *genus* of X. (If X is a sphere, then its genus is defined to be 0.)

A meromorphic differential ω on X is by definition a holomorphic differential on $X \setminus S$ where S is a finite subset of X such that if $(U, \phi: U \to \phi(U) \subseteq \mathbb{C})$ is any chart of X, then ω is of the form f(z) dz on this chart where f(z) is a meromorphic function on $\phi(U)$ whose poles are exactly $\phi(U \cap S)$. The basic feature of a meromorphic differential is the transformation equation between different charts: if (V, ψ) is another chart that overlaps (U, ϕ) and $\omega = g(w) dw$ on (V, ψ) , then we have

$$f(z) = g(\psi \circ \phi^{-1}(z)) \frac{\mathrm{d}\psi \circ \phi^{-1}(z)}{\mathrm{d}z}, \quad z \in \phi(U \cap V).$$

The set of all meromorphic differentials on X is denoted by $\mathscr{K}(X)$ which is obviously a complex vector space. Let $\mathscr{M}(X)$ denote the complex vector space of all meromorphic functions on X as in Section 2. If there exists a nonzero $\omega_0 \in \mathscr{K}(X)$, then the map $\mathscr{M}(X) \to \mathscr{K}(X)$ that sends f to $f \cdot \omega_0$ is a \mathbb{C} -linear isomorphism. Such ω_0 actually exists according to a fundamental result in the theory of Riemann surfaces (cf. [33, Theorem 1.10, Chapter IV]). As a consequence, nonzero meromorphic functions always exist on any compact Riemann surface.

By a (Q-valued) divisor on X, we understand a function $X \to \mathbb{Q}$ with finite support. We often write a divisor D as a formal sum $\sum_{x \in X} d_x \cdot (x)$ where $d_x = D(x)$. The term "divisor" traditionally refers to a Z-valued divisor but in this paper to a Q-valued divisor. The sets of all Z-valued divisors and of all Q-valued divisors are denoted by $\mathscr{D}(X)$ and $\mathscr{D}_{\mathbb{Q}}(X)$ respectively. Equipped with the pointwise addition, they are abelian groups. We define the *degree* of $D \in \mathscr{D}_{\mathbb{Q}}(X)$ by $\deg(D) = \sum_{x \in X} D(x)$, which is well-defined since the sum is actually a finite sum. Moreover, define the floor function as follows: for $D \in \mathscr{D}_{\mathbb{Q}}(X)$, set $[D] = \sum_{x \in X} [d_x] \cdot (x)$, where $[d_x]$ is the greatest integer that does not exceed d_x . Finally, for two divisors D_1 and D_2 , we say $D_1 \ge D_2$ if $D_1(x) \ge D_2(x)$ far any $x \in X$, and we say $D_1 > D_2$ if $D_1 \ge D_2$ but $D_1 \ne D_2$.

Generally speaking, with any meromorphic section of a holomorphic line bundle over X one can associate a Z-valued divisor. Specifically, if $f \in \mathscr{M}(X)$, then define $\operatorname{div}(f) = \sum_{x \in X} d_x \cdot (x)$ with d_x being the least exponent in the Laurent expansion of f at x in any chart. For $\omega \in \mathscr{K}(X)$ we define $\operatorname{div}(\omega)$ in a similar manner.

The compact Riemann surfaces we need are the modular curves X_G where G is a finite index subgroup of $SL_2(\mathbb{Z})$ which we describe now. Let $\mathfrak{H}^* = \mathfrak{H} \cup \mathbb{P}^1(\mathbb{Q})$ where $\mathbb{P}^1(\mathbb{Q})$ is the projective line over \mathbb{Q} which can be identified with \mathbb{Q} together with a point $i\infty = 1/0$. The group $SL_2(\mathbb{Z})$ acts on $\mathbb{P}^1(\mathbb{Q})$ on the left via $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot (p/q) = \frac{ap+bq}{cp+dq}$. We endow \mathfrak{H}^* with the topology generated by the usual topology of \mathfrak{H} and the sets

$$\{\gamma \tau \colon \tau \in \mathfrak{H}_{Y_0} \cup \{\mathrm{i}\infty\}\}, \quad \gamma \in \mathrm{SL}_2(\mathbb{Z}), \, Y_0 > 0,$$

where $\mathfrak{H}_{Y_0} = \{\tau \in \mathfrak{H} : \mathfrak{F}(\tau) > Y_0\}$. It is immediate that \mathfrak{H}^* with this topology is a Hausdorff space⁴, \mathfrak{H} is an open subset of \mathfrak{H}^* and G acts on \mathfrak{H}^* (on the left) by homeomorphisms. Let $G \setminus \mathfrak{H}^*$ be the orbit space (endowed with the quotient topology) and $p : \mathfrak{H}^* \to G \setminus \mathfrak{H}^*$ be the quotient map. It can be shown that $G \setminus \mathfrak{H}^*$ is Hausdorff (cf. eg. [35, Lemma 1.7.7]) and when $[\mathrm{SL}_2(\mathbb{Z}) : G] < +\infty$ $G \setminus \mathfrak{H}^*$ is compact (cf. eg. [33, Proposition 14.6, Chapter IV]). Hereafter we always assume that $[\mathrm{SL}_2(\mathbb{Z}) : G] < +\infty$. Write $G \setminus \mathfrak{H}^* = G \setminus \mathfrak{H} \cup G \setminus \mathbb{P}^1(\mathbb{Q})$, a disjoint union. Then $G \setminus \mathfrak{H}$ is open in $G \setminus \mathfrak{H}^*$ and $G \setminus \mathbb{P}^1(\mathbb{Q})$ is a finite set since $\mathrm{SL}_2(\mathbb{Z})$ acts transitively on $\mathbb{P}^1(\mathbb{Q})$ and hence $|G \setminus \mathbb{P}^1(\mathbb{Q})| \leq$ $[\mathrm{SL}_2(\mathbb{Z}) : G]$. The orbits in $G \setminus \mathbb{P}^1(\mathbb{Q})$ are called *cusps* of $G \setminus \mathfrak{H}^*$ and by abuse of language, the points in $\mathbb{P}^1(\mathbb{Q})$ are also called cusps. The *width* of a cusp $G \cdot x \in G \setminus \mathbb{P}^1(\mathbb{Q})$ is defined by $w_{Gx} =$ $w_x = [\overline{\mathrm{SL}_2(\mathbb{Z})_x} : \overline{G_x}]$ which is independent of the choice of the representative x. We have (cf. [29, Proposition 6.3.8(b)])

$$[\overline{\mathrm{SL}_2(\mathbb{Z})} \colon \overline{G}] = \sum_{Gx \in G \setminus \mathbb{P}^1(\mathbb{Q})} w_{Gx}.$$

We also need the notion of *elliptic points*. It is well known that the stabilizer of each point in $\{\gamma i: \gamma \in \mathrm{SL}_2(\mathbb{Z})\}$ under the action of $\mathrm{SL}_2(\mathbb{Z})/\{\pm I\}$ is a cyclic group of order 2, the stabilizer of each point in $\{\gamma \rho: \gamma \in \mathrm{SL}_2(\mathbb{Z})\}$ is a cyclic group of order 3 where $\rho = \frac{-1+\sqrt{3}i}{2}$ and the stabilizer of any other point in \mathfrak{H} is trivial. Thus for $\tau \in \mathfrak{H}$, $|\overline{G}_{\tau}| = 1$, 2 or 3 since $|\overline{G}_{\tau}|$ divides $|\overline{\mathrm{SL}_2(\mathbb{Z})}_{\tau}|$. If $|\overline{G}_{\tau}| = 2$ (3 respectively) then we call $p(\tau)$ an elliptic point for $G \setminus \mathfrak{H}^*$ of period 2 (3 respectively). By abuse of language, we also call the number τ an elliptic point if $p(\tau)$ is.

The description and verification of the atlas on $G \setminus \mathfrak{H}^*$ that turns it into a compact Riemann surface are rather technical and tedious but it is basic and well known (cf. eg. [35, Section 1.8]). Let X_G denote the resulting compact Riemann surface. We will mainly use $X_{\Gamma_0(N)}$ hereafter so set $X_0(N) = X_{\Gamma_0(N)}$ for simplicity. Let g be the genus of X_G . Then by applying the Riemann-Hurwitz ramification formula to the holomorphic map $X_G \to X_{\mathrm{SL}_2(\mathbb{Z})}, G \cdot z \mapsto \mathrm{SL}_2(\mathbb{Z}) \cdot z$, noticing that the degree of this map equals $m = [\overline{\mathrm{SL}_2(\mathbb{Z})}: \overline{G}]$ and the genus of $X_{\mathrm{SL}_2(\mathbb{Z})}$ is 0, we obtain (cf. e.g. [36, Theorem 3.1.1] for the case G is a congruence subgroup)

(8)
$$g = 1 + \frac{m}{12} - \frac{\varepsilon_2}{4} - \frac{\varepsilon_3}{3} - \frac{\varepsilon_\infty}{2},$$

where ε_2 and ε_3 are the numbers of elliptic points of X_G of period 2 and 3 respectively and ε_{∞} is the number of cusps of X_G . More formally,

$$\varepsilon_2 = \#\{G \cdot z \in G \setminus \mathrm{SL}_2(\mathbb{Z})i \colon |\overline{G}_z| = 2\},$$

$$\varepsilon_3 = \#\{G \cdot z \in G \setminus \mathrm{SL}_2(\mathbb{Z})\rho \colon |\overline{G}_z| = 3\}, \quad \rho = \frac{-1 + \sqrt{3}i}{2},$$

$$\varepsilon_{\infty} = \#G \setminus \mathbb{P}^1(\mathbb{Q}).$$

⁴The space \mathfrak{H}^* is not locally compact and hence can not be a topological manifold.

For $G = \Gamma_0(N)$, it is known that

(9)
$$m = [\overline{\mathrm{SL}_2(\mathbb{Z})} \colon \overline{\Gamma_0(N)}] = [\mathrm{SL}_2(\mathbb{Z}) \colon \Gamma_0(N)] = N \prod_{p|N} \left(1 + \frac{1}{p}\right)$$

(10)
$$\varepsilon_2 = \begin{cases} \prod_{p \mid N} \left(1 + \left(\frac{-4}{p} \right) \right) & \text{if } 4 \nmid N, \\ 0 & \text{if } 4 \mid N, \end{cases}$$

(11)
$$\varepsilon_{3} = \begin{cases} \prod_{p \mid N} \left(1 + \left(\frac{-3}{p} \right) \right) & \text{if } 9 \nmid N, \\ 0 & \text{if } 9 \mid N, \end{cases}$$

(12)
$$\varepsilon_{\infty} = \sum_{d|N} \phi(d, N/d),$$

where p denotes a prime, $\phi(d, N/d) = \phi(\operatorname{gcd}(d, N/d))$, and ϕ is the Euler function. For proofs, see [29, Corollary 6.2.13, Corollary 6.3.24(b)] and [36, Corollary 3.7.2].

Now let us associate a \mathbb{Q} -valued divisor to any meromorphic modular form of rational weight. For integral or half-integral weights, this association is the same as the ordinary one⁵ (cf. [36, eq. (3.2) and (3.3)]). Our treatment has the feature that one need not distinguish between regular and irregular cusps.

Definition 3.1. Let G be a subgroup of $\operatorname{SL}_2(\mathbb{Z})$ of finite index, $D \in \mathbb{Z}_{\geq 1}$ and $k \in \frac{1}{D}\mathbb{Z}$. Let χ be a multiplier system for G of weight k of cover index D (not assumed to be of finite order). Let f be a *nonzero* meromorphic modular form of weight k for G with character χ . We define its divisor $\operatorname{div}(f) = \sum_{x \in X_G} \operatorname{div}_x(f) \cdot (x) \in \mathscr{D}_{\mathbb{Q}}(X_G)$ as follows:

- (a) If $\tau \in \mathfrak{H}$, then set $\operatorname{div}_{G\tau}(f) = n_0/|\overline{G}_{\tau}|$ where n_0 is the integer such that $f(z) = \sum_{n \ge n_0} c_n (z \tau)^n$, $c_{n_0} \ne 0$.
- (b) If $x \in \mathbb{P}^1(\mathbb{Q})$ and $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ satisfying $\gamma(\mathrm{i}\infty) = x$, then set $\operatorname{div}_{Gx}(f) = \operatorname{ord}_{\mathrm{i}\infty}(f|_k \widetilde{\gamma}) \cdot w_{Gx}$.

One can verify that $\operatorname{div}(f)$ is well-defined, which is not totally trivial but straightforward. The usefulness of this notion relies on the following two simple facts:

- If g is another nonzero meromorphic modular form of weight $k' \in \frac{1}{D}\mathbb{Z}$ for G with character χ' , then we have $\operatorname{div}(fg) = \operatorname{div}(f) + \operatorname{div}(g)$ in $\mathscr{D}_{\mathbb{Q}}(X_G)$ (cf. Remark 2.3).
- If k = 0 and χ is the trivial character, then f descends to a meromorphic function \tilde{f} on X_G . We have $\operatorname{div}(f) = \operatorname{div}(\tilde{f})$ where $\operatorname{div}(\tilde{f})$ is the usual divisor of a meromorphic function.

Note that the factors $1/|\overline{G}_{\tau}|$ and w_{Gx} in Definition 3.1 are chosen to let the latter fact hold (to see this one must dive into the atlas of X_G which we have omitted). Also note that the former fact holds because the factors $1/|\overline{G}_{\tau}|$ and w_{Gx} remain unchanged for all k.

,

 $^{{}^{5}}$ But it is different from the one described in [33, p. 299]

Similarly, we have $\operatorname{div}(f/g) = \operatorname{div}(f) - \operatorname{div}(g)$ and in particular $\operatorname{div}(1/g) = -\operatorname{div}(g)$.

A fundamental theorem on modular forms of rational weight is the following one, which in cases of integral or half-integral weights is sometimes called the *valence formula*.

Theorem 3.2. Let us use the notation of Definition 3.1 and set $m = [\overline{\operatorname{SL}_2(\mathbb{Z})}: \overline{G}]$. Then we have

$$\deg(\operatorname{div} f) = \frac{1}{12}mk$$

Proof. Suppose that $\overline{\operatorname{SL}_2(\mathbb{Z})} = \bigcup_{1 \leq j \leq m} \overline{G\gamma_j}$ is a disjoint union where $\gamma_j \in \operatorname{SL}_2(\mathbb{Z})$ and $\overline{\gamma_j} = \gamma_j \cdot \{\pm I\}$. Set $g = \prod_{1 \leq j \leq m} f|_k \widetilde{\gamma_j}$. Then $g|_{mk} \gamma = c_{\gamma} g$ for any $\gamma \in \widetilde{\operatorname{SL}_2(\mathbb{Z})^D}$ where $c_{\gamma} \in \mathbb{C}^{\times}$. Since g is not identically zero, the map $\gamma \mapsto c_{\gamma}$ is a linear character on $\operatorname{SL}_2(\mathbb{Z})^D$. According to a presentation of $\widetilde{\operatorname{SL}_2(\mathbb{Z})^D}$ (cf. [16, Lemma 5.2]) the order of this character divides 12D and hence g^{12D} transforms like a modular form of weight 12Dmk for $\operatorname{SL}_2(\mathbb{Z})$ with trivial character. Since f is meromorphic at cusps, so is g^{12D} . Therefore g^{12D} is a meromorphic modular form of weight 12Dmk for $\operatorname{SL}_2(\mathbb{Z})$ with trivial character. Applying the usual valence formula for even weights (cf. [29, Theorem 5.6.1]) we obtain deg(div g^{12D}) = Dmk. It remains to prove that $12D \operatorname{deg}(\operatorname{div} f) = \operatorname{deg}(\operatorname{div} g^{12D})$, which is equivalent to deg(div f) = deg(div g). We now prove a stronger assertion, i.e.

(13)
$$\operatorname{div}_{x}(g) = \sum_{y \in p^{-1}(x)} \operatorname{div}_{y}(f)$$

for any $x \in X_{\mathrm{SL}_2(\mathbb{Z})}$, where $p: X_G \to X_{\mathrm{SL}_2(\mathbb{Z})}$ is the natural projection $G \cdot \tau \mapsto \mathrm{SL}_2(\mathbb{Z}) \cdot \tau$. In the case that $x \in \mathrm{SL}_2(\mathbb{Z}) \setminus \mathfrak{H}$, we write $x = \mathrm{SL}_2(\mathbb{Z}) \cdot \tau$ and set $w_\tau = [\overline{\mathrm{SL}_2(\mathbb{Z})}_\tau : \overline{G}_\tau]$ (similar to the notion of the width of a cusp in the group-theoretical aspect). Notice that $\mathrm{ord}_\tau f|_k \widetilde{\gamma}_j = \mathrm{ord}_{\gamma_j \tau} f$ where $\mathrm{ord}_\tau f$ is the integer n_0 such that $\lim_{z\to\tau} f(z)(z-\tau)^{-n_0}$ is nonzero. Thus we have

$$\operatorname{div}_{x}(g) = |\overline{\operatorname{SL}_{2}(\mathbb{Z})}_{\tau}|^{-1} \cdot \sum_{1 \leq j \leq m} \operatorname{ord}_{\gamma_{j}\tau} f.$$

The underlying set of the multiset $\{G \cdot \gamma_j \tau \colon 1 \leq j \leq m\}$ equals $p^{-1}(x)$ and the multiplicity of each element $G \cdot \gamma_j \tau$ equals $w_{\gamma_j \tau}$. It follows that

$$\operatorname{div}_{x}(g) = |\overline{\operatorname{SL}_{2}(\mathbb{Z})}_{\tau}|^{-1} \cdot \sum_{y = G \cdot \tau \in p^{-1}(x)} w_{\tau} \cdot \operatorname{ord}_{\tau} f = \sum_{y \in p^{-1}(x)} \operatorname{div}_{y}(f),$$

i.e., the desired assertion (13) holds. In the other case $x \in \mathrm{SL}_2(\mathbb{Z}) \setminus \mathbb{P}^1(\mathbb{Q})$, we have $x = \mathrm{SL}_2(\mathbb{Z}) \cdot i\infty$ and its width is 1. Therefore

$$\operatorname{div}_x(g) = \sum_{1 \le j \le m} \operatorname{ord}_{\mathrm{i}\infty}(f|_k \widetilde{\gamma_j}).$$

The underlying set of the multiset $\{G \cdot \gamma_j i \infty : 1 \leq j \leq m\}$ equals $p^{-1}(x)$ and the multiplicity of each element $G \cdot \gamma_j i \infty$ equals $w_{\gamma_j i \infty}$. It follows that

$$\operatorname{div}_{x}(g) = \sum_{y \in G \cdot \gamma i \infty \in p^{-1}(x)} w_{\gamma i \infty} \operatorname{ord}_{i \infty}(f|_{k} \widetilde{\gamma}) = \sum_{y \in p^{-1}(x)} \operatorname{div}_{y}(f),$$

17

i.e., the desired assertion (13) holds. This concludes the proof.

Remark 3.3. The above theorem holds even when χ is non-unitary and the proof remains unchanged. For instance, one can apply it to the generalized modular forms constructed in [25].

In the rest we need to know information about the divisor of an eta-quotient times a product of Eisenstein series on $SL_2(\mathbb{Z})$. We let

$$E_4(\tau) = 1 + 240 \sum_{n \in \mathbb{Z}_{\ge 1}} \sigma_3(n) q^n,$$

$$E_6(\tau) = 1 - 504 \sum_{n \in \mathbb{Z}_{\ge 1}} \sigma_5(n) q^n,$$

where $\sigma_k(n) = \sum_{d|n} d^k$. It is well known that $E_4 \in M_4(SL_2(\mathbb{Z}))$ and $E_6 \in M_6(SL_2(\mathbb{Z}))$.

Proposition 3.4. Let $N \in \mathbb{Z}_{\geq 1}$, $r_n \in \mathbb{Q}$ for any $n \mid N$ and $t_1, t_2 \in \mathbb{Z}$. Let χ be the character (6) where $D \in \mathbb{Z}_{\geq 1}$ is chosen so that $Dr_n \in 2\mathbb{Z}$ for any $n \mid N$. Set

$$f(\tau) = \prod_{n|N} \eta(n\tau)^{r_n} \cdot E_4(\tau)^{t_1} \cdot E_6(\tau)^{t_2}.$$

Then f is a nonzero meromorphic modular form of weight $k = \frac{1}{2} \sum_{n} r_n + 4t_1 + 6t_2$ for $\Gamma_0(N)$ with character χ and

$$\deg\left([\operatorname{div} f]\right) = \sum_{c|N} \phi(c, N/c) \cdot \left[\frac{N}{24(N, c^2)} \sum_{n|N} \frac{(n, c)^2}{n} r_n\right] + \left(\frac{t_1}{3} + \frac{t_2}{2}\right) m - \left\{\frac{t_2}{2}\right\} \varepsilon_2 - \left\{\frac{t_1}{3}\right\} \varepsilon_3,$$

where m, ε_2 and ε_3 are given in (9), (10) and (11).

Proof. The fact that f is a nonzero meromorphic modular form follows from Remark 2.3. It remains to compute deg ([div f]). The divisors of E_4 and E_6 on $X_{\Gamma_0(N)}$ are known (cf. eg. [29, Proposition 5.6.5]) as follows:

$$\operatorname{div}_{x}(E_{u}) = \begin{cases} 1/e_{\tau} & \text{if } x = \Gamma_{0}(N) \cdot \tau \in \Gamma_{0}(N) \backslash \operatorname{SL}_{2}(\mathbb{Z}) \cdot \tau_{0}, \\ 0 & \text{otherwise}, \end{cases}$$

where u = 4 or 6, $e_{\tau} = |\overline{\Gamma_0(N)}_{\tau}|$ and $\tau_0 = \rho = \frac{-1+\sqrt{3}i}{2}$ if u = 4; $\tau_0 = i$ if u = 6. On the other hand, if we set $f_0(\tau) = \prod_{n|N} \eta(n\tau)^{r_n}$, then $\operatorname{div}_x(f_0) = 0$ for $x \in \Gamma_0(N) \setminus \mathfrak{H}$ since f_0 has no poles or zeros on \mathfrak{H} . For $x \in \Gamma_0(N) \setminus \mathbb{P}^1(\mathbb{Q})$, we can find $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{SL}_2(\mathbb{Z})$ such that $x = \Gamma_0(N) \cdot \begin{pmatrix} a & b \\ c & d \end{pmatrix} i\infty$. It is known that $w_x = N/(N, c^2)$ (cf. [29, Corollary 6.3.24(a)]). It follows from this and (7) that

(14)
$$\operatorname{div}_{x}(f_{0}) = \frac{N}{24(N,c^{2})} \sum_{n|N} \frac{(n,c)^{2}}{n} r_{n}.$$

Therefore

(15)
$$\deg\left([\operatorname{div} f]\right) = \sum_{x \in \Gamma_0(N) \setminus \operatorname{SL}_2(\mathbb{Z})\rho} [t_1 \cdot \operatorname{div}_x(E_4)] + \sum_{x \in \Gamma_0(N) \setminus \operatorname{SL}_2(\mathbb{Z})i} [t_2 \cdot \operatorname{div}_x(E_6)] + \sum_{x \in \Gamma_0(N) \setminus \mathbb{P}^1(\mathbb{Q})} [\operatorname{div}_x(f_0)],$$

where the three sums in the right-hand side are denoted by S_1 , S_2 and S_3 respectively hereafter. To compute S_1 , let $\operatorname{SL}_2(\mathbb{Z}) = \bigcup_{1 \leq j \leq m} \Gamma_0(N) \gamma_j$ be a disjoint union⁶. Then the underlying set of the multiset $\{\Gamma_0(N) \cdot \gamma_j \rho \colon 1 \leq j \leq m\}$ equals $\Gamma_0(N) \setminus \operatorname{SL}_2(\mathbb{Z})\rho$ and the multiplicity of each element $\Gamma_0(N) \cdot \gamma_j \rho$ equals $|\overline{\operatorname{SL}_2(\mathbb{Z})}_{\gamma_j \rho}|/|\overline{\Gamma_0(N)}_{\gamma_j \rho}| = 3/e_{\gamma_j \rho}$. Hence we have

$$S_1 = \frac{1}{3} \sum_{1 \le j \le m: \ e_{\gamma_j \rho} = 1} \left[\frac{t_1}{e_{\gamma_j \rho}} \right] + \sum_{1 \le j \le m: \ e_{\gamma_j \rho} = 3} \left[\frac{t_1}{e_{\gamma_j \rho}} \right] = \frac{mt_1}{3} - \left\{ \frac{t_1}{3} \right\} \varepsilon_3.$$

Similarly, $S_2 = \frac{mt_2}{2} - \left\{\frac{t_2}{2}\right\} \varepsilon_2$. For S_3 , we need a complete set of representatives of $\Gamma_0(N) \setminus \mathbb{P}^1(\mathbb{Q})$, i.e., the set of $a/c \in \mathbb{P}^1(\mathbb{Q})$ where $c \mid N$ and for each a_0 such that $0 \leq a_0 < (c, N/c)$ with $(a_0, c, N/c) = 1$, a is chosen to satisfy $a \equiv a_0 \mod (c, N/c)$ and (a, c) = 1 (cf. [29, Corollary 6.3.23]). It follows from this and (14) that

$$S_3 = \sum_{c|N} \phi(c, N/c) \cdot \left[\frac{N}{24(N, c^2)} \sum_{n|N} \frac{(n, c)^2}{n} r_n \right]$$

Inserting the expressions for S_1 , S_2 and S_3 into (15) gives the desired formula.

4. The main theorem: Dimension formulas for rational weights

We recall the Riemann-Roch theorem here for proving the main theorem. Let X be a compact Riemann surface of genus g and $D \in \mathscr{D}_{\mathbb{Q}}(X)$. Then the *Riemann-Roch space* is defined by $\mathscr{L}(D) = \{f \in \mathscr{M}(X) : \operatorname{div}(f) \geq -D\}$ with the convention that $0 \in \mathscr{L}(D)$. It is immediate that $\mathscr{L}(D)$ is a complex vector space and that $\mathscr{L}(D) = \mathscr{L}([D])$. Similarly we define $\mathscr{K}(D) = \{\omega \in \mathscr{K}(X) : \operatorname{div}(\omega) \geq D\} \cup \{0\}$. Then the Riemann-Roch theorem states that

(16)
$$\dim_{\mathbb{C}} \mathscr{L}(D) = \dim_{\mathbb{C}} \mathscr{K}(D) + \deg(D) - g + 1,$$

provided that D is integral, i.e., $D \in \mathscr{D}(X)$. Note that it is tacitly understood that $\dim_{\mathbb{C}} \mathscr{L}(D) < +\infty$. For a proof based on the existence theorems for harmonic functions on Riemann surfaces, see [33, p. 249]. Set D = 0; we find that $\dim_{\mathbb{C}} \mathscr{K}(0) = g$, that is, the dimension of the space of all holomorphic differentials is g. Let ω_0 be any nonzero meromorphic differential (necessarily exists) and $K = \operatorname{div}(\omega_0)$. Then $\mathscr{L}(K - D) \to \mathscr{K}(D)$, $f \mapsto f \cdot \omega_0$ is a \mathbb{C} -linear isomorphism. Now set D = K in (16); we find that $\operatorname{deg}(K) = 2g - 2$. Finally, if $D \in \mathscr{D}(X)$ and $\operatorname{deg}(D) > 2g - 2$, then

⁶Since $-I \in \Gamma_0(N)$, this is equivalent to $\overline{\operatorname{SL}_2(\mathbb{Z})} = \bigcup_{1 \leq j \leq m} \overline{\Gamma_0(N)} \overline{\gamma_j}$, and as well to $\widetilde{\operatorname{SL}_2(\mathbb{Z})} = \bigcup_{1 \leq j \leq m} \widetilde{\Gamma_0(N)} \widetilde{\gamma_j}$ both of which are disjoint unions.

 $\deg(K-D) < 0$ and hence $\mathscr{L}(K-D) = \mathscr{K}(D) = \{0\}$ since the degree of a nonzero meromorphic function is 0. It follows that

(17)
$$\dim_{\mathbb{C}} \mathscr{L}(D) = \deg(D) - g + 1, \quad \text{if } D \in \mathscr{D}(X), \, \deg(D) > 2g - 2.$$

Our proof of the main theorem rests on the following general lemma, which is the key point for using the Riemann-Roch theorem.

Lemma 4.1. Let G be a subgroup of $SL_2(\mathbb{Z})$ of finite index, $D \in \mathbb{Z}_{\geq 1}$, $k \in \frac{1}{D}\mathbb{Z}$ and $v \colon \widetilde{G}^D \to \mathbb{C}^{\times}$ be a character. Let $p \colon \mathfrak{H} \to G \setminus \mathfrak{H}$ be the natural projection. Suppose there exists a nonzero meromorphic modular form f_0 of weight k for group G with character v. Then the map

$$\mathscr{L}([\operatorname{div} f_0]) \to M_k(G, v)$$
$$f \mapsto (f|_{G \setminus \mathfrak{H}}) \circ p \cdot f_0$$

is a \mathbb{C} -linear isomorphism. Moreover, let

(18)
$$R = \{ x \in G \setminus \mathbb{P}^1(\mathbb{Q}) \colon \operatorname{div}_x(f_0) \in \mathbb{Z} \}$$

Then the map

$$\mathscr{L}\left([\operatorname{div} f_0] - \sum_{x \in R} (x)\right) \to S_k(G, v)$$
$$f \mapsto (f|_{G \setminus \mathfrak{H}}) \circ p \cdot f_0$$

is also a \mathbb{C} -linear isomorphism.

Proof. Let $f \in \mathscr{L}([\operatorname{div} f_0])$ be arbitrary. Since p is holomorphic, $(f|_{G\setminus\mathfrak{H}}) \circ p$ is a modular function for group G with trivial character. Thus $(f|_{G\setminus\mathfrak{H}}) \circ p \cdot f_0$ is a meromorphic modular form of weight k for group G with character v by Remark 2.3. Now assume $f \neq 0$ and let $x \in X_G$ be arbitrary. Since $f \in \mathscr{L}([\operatorname{div} f_0])$ we have $\operatorname{div}_x((f|_{G\setminus\mathfrak{H}}) \circ p) = \operatorname{div}_x(f) \geq -\operatorname{div}_x(f_0)$. It follows that

$$\operatorname{div}_x((f|_{G\setminus\mathfrak{H}})\circ p\cdot f_0) = \operatorname{div}_x((f|_{G\setminus\mathfrak{H}})\circ p) + \operatorname{div}_x(f_0) \ge 0.$$

Hence $(f|_{G\setminus\mathfrak{H}}) \circ p \cdot f_0 \in M_k(G, v)$. It is obvious that the considered map is a \mathbb{C} -linear injection. To prove the surjectivity, let $0 \neq g \in M_k(G, v)$ be arbitrary, then by Remark 2.3 g/f_0 is a modular function for group G with trivial character and hence it descends to some nonzero $f \in \mathcal{M}(X_G)$. For any $x \in X_G$ we have

$$\operatorname{div}_x(f) = \operatorname{div}_x(g/f_0) = \operatorname{div}_x(g) - \operatorname{div}_x(f_0) \ge -\operatorname{div}_x(f_0).$$

Therefore $f \in \mathscr{L}(\operatorname{div} f_0) = \mathscr{L}([\operatorname{div} f_0])$ and the image of f is g from which the surjectivity follows.

The assertion on the second map can be proved in a similar manner and the only thing that need to explain is that $(f|_{G\setminus\mathfrak{H}}) \circ p \cdot f_0 \in S_k(G, v)$ if and only if $f \in \mathscr{L}\left([\operatorname{div} f_0] - \sum_{x \in R}(x)\right)$. If $0 \neq f \in \mathscr{L}\left([\operatorname{div} f_0] - \sum_{x \in R}(x)\right)$ and $x \in R$, then

$$\operatorname{div}_x((f|_{G\setminus\mathfrak{H}})\circ p\cdot f_0) = \operatorname{div}_x(f) + \operatorname{div}_x(f_0) \ge 1 - [\operatorname{div}_x(f_0)] + \operatorname{div}_x(f_0) > 0$$

On the other hand, if $x \notin R$ then $-[\operatorname{div}_x(f_0)] + \operatorname{div}_x(f_0) > 0$ and hence

$$\operatorname{div}_x((f|_{G\setminus\mathfrak{H}})\circ p\cdot f_0) = \operatorname{div}_x(f) + \operatorname{div}_x(f_0) \ge -[\operatorname{div}_x(f_0)] + \operatorname{div}_x(f_0) > 0.$$

It follows that $(f|_{G\setminus\mathfrak{H}}) \circ p \cdot f_0 \in S_k(G, v)$. The converse can be proved similarly.

Now we state and prove the main theorem.

Theorem 4.2. Let $N \in \mathbb{Z}_{\geq 1}$, $r_n \in \mathbb{Q}$ for any $n \mid N$ and $t \in \mathbb{Z}_{\geq 0}$. Let D be a positive integer such that $Dr_n \in 2\mathbb{Z}$ for any $n \mid N$. Let $\chi \colon \widetilde{\Gamma_0(N)^D} \to \mathbb{C}^{\times}$ be the character (6) and m, ε_2 , ε_3 are given by (9), (10), (11) respectively. Set $k = \frac{1}{2} \sum_n r_n + 2t$ and

(19)
$$x_{c} = \sum_{n|N} \frac{N}{(N,c^{2})} \cdot \frac{(n,c)^{2}}{n} r_{n}$$

with $c \mid N$. Suppose either

(20)
$$\sum_{c|N} \phi(c, N/c) \cdot \left(\left[\frac{x_c}{24} \right] + 1 \right) > 0, \ t \ge 1$$

or

(21)
$$k > 2 - \frac{6}{m}\varepsilon_2 - \frac{8}{m}\varepsilon_3 - \frac{12}{m}\sum_{c|N}\phi(c,N/c)\cdot\left(1 - \left\{\frac{x_c}{24}\right\}\right), t = 0.$$

Then we have

(22)
$$\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) = \frac{k-1}{12}m + \left(\frac{1}{4} - \left\{\frac{t}{2}\right\}\right)\varepsilon_2 + \left(\frac{1}{3} - \left\{-\frac{t}{3}\right\}\right)\varepsilon_3 + \sum_{c|N}\phi(c, N/c)\cdot\left(\frac{1}{2} - \left\{\frac{x_c}{24}\right\}\right).$$

Proof. Set

$$f_0(\tau) = \prod_{n|N} \eta(n\tau)^{r_n} \cdot E_4(\tau)^{-t} \cdot E_6(\tau)^t.$$

According to Proposition 3.4 f_0 is a nonzero meromorphic modular form of weight k for group $\Gamma_0(N)$ with character χ . Hence $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) = \dim_{\mathbb{C}} \mathscr{L}([\operatorname{div} f_0])$ by Lemma 4.1. Using Proposition 3.4 again we find that

(23)
$$\deg\left(\left[\operatorname{div} f_{0}\right]\right) = \sum_{c|N} \phi(c, N/c) \cdot \left[\frac{x_{c}}{24}\right] + \frac{t}{6}m - \left\{\frac{t}{2}\right\}\varepsilon_{2} - \left\{-\frac{t}{3}\right\}\varepsilon_{3}.$$

We now prove either (20) or (21) implies that deg ([div f_0]) > 2g - 2 where g is the genus of $X_{\Gamma_0(N)}$ (cf. (8)). First suppose (20) holds. If N = 1, then g = 0, $r_1 \ge 0$ and hence

$$\deg\left([\operatorname{div} f_0]\right) = \left[\frac{r_1}{24}\right] + \frac{t}{6} - \left\{\frac{t}{2}\right\} - \left\{-\frac{t}{3}\right\} > -2 = 2g - 2.$$

If $N \geq 2$, set

$$\gamma(t) = \frac{t}{6}m - \left\{\frac{t}{2}\right\}\varepsilon_2 - \left\{-\frac{t}{3}\right\}\varepsilon_3.$$

We have

$$\gamma(t+1) - \gamma(t) \ge \frac{1}{6}m - \frac{1}{2}\varepsilon_2 - \frac{2}{3}\varepsilon_3 = 2g - 2 + \varepsilon_\infty$$

for $t \ge 1$. Since $N \ge 2$ we have $\varepsilon_{\infty} \ge 2$ and hence $\gamma(t+1) - \gamma(t) \ge 0$. Therefore deg ([div f_0]) as a function of $t \in \mathbb{Z}_{\ge 1}$ is increasing which means we need only to prove

$$\sum_{c|N} \phi(c, N/c) \cdot \left[\frac{x_c}{24}\right] + \frac{1}{6}m - \frac{1}{2}\varepsilon_2 - \frac{2}{3}\varepsilon_3 > 2g - 2.$$

This is actually equivalent to (20) according to (8) and (12). Thereby we have shown that (20) implies deg ([div f_0]) > 2g - 2. Next suppose (21) holds, which is equivalent to

(24)
$$\frac{1}{12}mk > \frac{1}{6}m - \frac{1}{2}\varepsilon_2 - \frac{2}{3}\varepsilon_3 - \varepsilon_\infty + \sum_{c|N}\phi(c, N/c)\left\{\frac{x_c}{24}\right\} = 2g - 2 + \sum_{c|N}\phi(c, N/c)\left\{\frac{x_c}{24}\right\}.$$

Applying Theorem 3.2 and (14) to f_0 (noting t = 0 here) and using the complete set of representatives of $\Gamma_0(N) \setminus \mathbb{P}^1(\mathbb{Q})$ described in the proof of Proposition 3.4 we find that

(25)
$$\frac{1}{12}m \cdot \frac{1}{2}\sum_{n} r_{n} = \deg(\operatorname{div} f_{0}) = \sum_{c|N} \phi(c, N/c) \cdot \frac{x_{c}}{24}$$

Inserting this into (24) we obtain deg ([div f_0]) > 2g - 2 as required.

Thus, in both cases we can apply (17) to $D = [\operatorname{div} f_0], X = X_{\Gamma_0(N)}$. It follows that

$$\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) = \dim_{\mathbb{C}} \mathscr{L}([\operatorname{div} f_0]) = \operatorname{deg}([\operatorname{div} f_0]) - g + 1.$$

Inserting (23), (8) and (25) into the above identity gives the desired formula.

Remark 4.3. If neither (20) nor (21) holds, then the difference between the left-hand side and the right-hand side of (22) equals $\dim_{\mathbb{C}} \mathscr{K}([\operatorname{div} f_0])$ according to (16). It follows that in all cases we have

$$\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) \ge \frac{k-1}{12}m + \left(\frac{1}{4} - \left\{\frac{t}{2}\right\}\right)\varepsilon_2 + \left(\frac{1}{3} - \left\{-\frac{t}{3}\right\}\right)\varepsilon_3 + \sum_{c|N} \phi(c, N/c) \cdot \left(\frac{1}{2} - \left\{\frac{x_c}{24}\right\}\right),$$

where the right-hand side is always an integer. We can derive an upper bound⁷ for dim_C $M_k(\Gamma_0(N), \chi)$ as well. Suppose that $k \ge 0$. (When k < 0 $M_k(\Gamma_0(N), \chi) = \{0\}$ by Theorem 3.2.) Then

$$\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) \le \left[\frac{1}{12}mk\right] + 1,$$

⁷This upper bound is still valid if $\Gamma_0(N)$ is replaced by any subgroup G of finite index in $SL_2(\mathbb{Z})$.

for if $f_1, \ldots, f_j \in M_k(\Gamma_0(N), \chi)$ with $j > \left[\frac{1}{12}mk\right] + 1$, then by linear algebra we can find a nontrivial linear combination $g = \sum_n c_n f_n$ such that $\operatorname{div}_x(g) \ge \left[\frac{1}{12}mk\right] + 1$ where $x \in \Gamma_0(N) \setminus \mathfrak{H}$ is not an elliptic point. If $g \neq 0$ then applying Theorem 3.2 to g we reach a contradiction. Hence g = 0 which means f_1, \ldots, f_j are linearly dependent.

Remark 4.4. We can derive a formula for $\dim_{\mathbb{C}} S_k(\Gamma_0(N), \chi)$ as well, using the second map in Lemma 4.1. However, it may happen that for certain r_n and t (22) holds while the corresponding formula for $\dim_{\mathbb{C}} S_k(\Gamma_0(N), \chi)$ is not applicable. This happens precisely when deg ($[\operatorname{div} f_0]$) > 2g-2 but deg ($[\operatorname{div} f_0]$) - $|R| \leq 2g - 2$. Besides, when k > 2, we always have dim_C $S_k(\Gamma_0(N), \chi) =$ dim_C $M_k(\Gamma_0(N), \chi) - |R|$ which is a well known fact when k is integral and χ is induced by a Dirichlet character. In the rational weight and arbitrary multiplier system case, this holds as well since the Petersson inner product can also be defined. The details are omitted here.

5. Order-character relations

Let us assume (21) holds and consider Theorem 4.2. The sequence $(x_c)_{c|N}$ occurs in the formula for dim_C $M_k(\Gamma_0(N), \chi)$ and is determined by the sequence $(r_n)_{n|N}$ via (19). Note that $(r_n)_{n|N}$ represents an eta-quotient (or its character) and $(x_c)_{c|N}$ represents the orders of this eta-quotient at cusps (or its divisor). Therefore (19) can be regarded as a map that sends the character represented by $(r_n)_{n|N}$ to the orders at cusps $(x_c)_{c|N}$. In some potential applications, the sequence of orders $(x_c)_{c|N}$ is first given, e.g., when one want to known dim_C $M_k(\Gamma_0(N), \chi)$ for fixed N and k, and then one works out $(r_n)_{n|N}$ via (19). Motivated by this, we need an inverse formula of (19). This formula is an enhancement of a statement in [37, p. 129] and is equivalent to [21, eq. (5.13)].

Let us fix some notations in linear algebra. For $N \in \mathbb{Z}_{\geq 1}$, $\mathscr{D}(N)$ denotes the set of positive divisors of N and $\sigma_0(N) = |\mathscr{D}(N)|$. Let V be a Q-vector space of dimension $\sigma_0(N)$, then a $\mathscr{D}(N)$ indexed basis of V is a sequence $(\mathbf{b}_n)_{n \in \mathscr{D}(N)}$ so that $\{\mathbf{b}_n\}$ is a basis of V. Let W be a Q-vector space of dimension $\sigma_0(M)$ with a $\mathscr{D}(M)$ -indexed basis $(\mathbf{b}'_m)_{m|M}$ where M is another positive integer and let $f: V \to W$ be a linear map. The matrix of f with respect to the pair $((\mathbf{b}_n), (\mathbf{b}'_m))$ is the $\mathscr{D}(M) \times \mathscr{D}(N)$ -indexed sequence $(a_{m,n})_{m|M,n|N}$ with the property $f(\mathbf{b}_n) = \sum_{m|M} a_{m,n} \mathbf{b}'_m$. All linear algebra machinery works for such kind of matrices. The difference is just the underlying index sets: we use $\mathscr{D}(N)$ and the ordinary one is $\{1, 2, \ldots, \sigma_0(N)\}$. We call the matrix $(a_{m,n})_{m|M,n|N}$ a $\mathscr{D}(M) \times \mathscr{D}(N)$ -indexed matrix and when M = N a $\mathscr{D}(N)$ -indexed matrix.

Let $N_1, N_2 \in \mathbb{Z}_{\geq 1}$ such that $gcd(N_1, N_2) = 1$ and let $(a_{c_1,n_1})_{c_1,n_1|N_1}$, $(b_{c_2,n_2})_{c_2,n_2|N_2}$ be $\mathscr{D}(N_1)$ indexed and $\mathscr{D}(N_2)$ -indexed matrices respectively. Define

$$(a_{c_1,n_1})_{c_1,n_1|N_1} \otimes (b_{c_2,n_2})_{c_2,n_2|N_2} = (a_{c_1,n_1} \cdot b_{c_2,n_2})_{c_1c_2,n_1n_2|N_1N_2}.$$

It is immediate that this is a well defined \mathbb{Q} -bilinear map from $\mathbb{Q}^{\mathscr{D}(N_1) \times \mathscr{D}(N_1)} \times \mathbb{Q}^{\mathscr{D}(N_2) \times \mathscr{D}(N_2)}$ to $\mathbb{Q}^{\mathscr{D}(N_1N_2) \times \mathscr{D}(N_1N_2)}$. Moreover it is universal among all bilinear maps on $\mathbb{Q}^{\mathscr{D}(N_1) \times \mathscr{D}(N_1)} \times \mathbb{Q}^{\mathscr{D}(N_2) \times \mathscr{D}(N_2)}$ and hence is a tensor product. We call it the *Kronecker product*.

Following Bhattacharya [21] we define $A_N = \left(\frac{N}{(N,c^2)} \cdot \frac{(n,c)^2}{n}\right)_{c,n|N}$ so that (19) becomes $x_c = \sum_{n|N} A_N(c,n)r_n$. It is immediate that $A_{N_1N_2} = A_{N_1} \otimes A_{N_2}$ whenever $gcd(N_1, N_2) = 1$.

Proposition 5.1. Let $N \in \mathbb{Z}_{\geq 1}$, $(x_c)_{c|N}$ and $(r_n)_{n|N}$ be in $\mathbb{Q}^{\mathcal{D}(N)}$. Then the following two relations are equivalent:

(26)
$$x_c = \sum_{n \mid N} \frac{N}{(N, c^2)} \cdot \frac{(n, c)^2}{n} r_n, \quad \forall c \mid N,$$

(27)
$$r_n = \frac{1}{N} \prod_{p|N} \frac{p}{p^2 - 1} \cdot \sum_{c|N} \left(\prod_{p^{\alpha} \parallel N} b_{p,\alpha}(v_p(n), v_p(c)) \right) x_c, \quad \forall n \mid N,$$

where p denotes a prime, $v_p(n)$ is the integer β such that $p^{\beta} \parallel n$ and

$$b_{p,\alpha}(i,j) = \begin{cases} p & \text{if } i = j = 0 \text{ or } \alpha, \\ (p^2 + 1)p^{\min(i,\alpha-i)-1} & \text{if } 0 < i = j < \alpha, \\ -p^{\min(j,\alpha-j)} & \text{if } |i-j| = 1, \\ 0 & \text{otherwise.} \end{cases}$$

In another words, we have

$$A_N^{-1}(n,c) = \frac{1}{N} \prod_{p|N} \frac{p}{p^2 - 1} \cdot \prod_{p^{\alpha} \parallel N} b_{p,\alpha}(v_p(n), v_p(c)).$$

This proposition is due to Bhattacharya (cf. [21, eq. (5.13)]).

Proof. Suppose $N_1, N_2 \in \mathbb{Z}_{\geq 1}$ such that $gcd(N_1, N_2) = 1$. Then A_{N_1} and A_{N_2} are both invertible if and only if $A_{N_1} \otimes A_{N_2}$ is and in this case $(A_{N_1} \otimes A_{N_2})^{-1} = A_{N_1}^{-1} \otimes A_{N_2}^{-1}$ by basic properties of the Kronecker product. Therefore it is sufficient to prove that

$$A_{p^{\alpha}}^{-1}(p^{i}, p^{j}) = \frac{1}{p^{\alpha-1}(p^{2}-1)} \cdot b_{p,\alpha}(i, j),$$

where p is a prime, $\alpha \in \mathbb{Z}_{\geq 1}$ and $0 \leq i, j \leq \alpha$. This is equivalent to

(28)
$$\sum_{\substack{j=0\\\alpha}}^{\alpha} A_{p^{\alpha}}(p^{i}, p^{j}) \cdot b_{p,\alpha}(j, i) = (p^{2} - 1)p^{\alpha - 1}$$

(29)
$$\sum_{j=0}^{\alpha} A_{p^{\alpha}}(p^{i_1}, p^j) \cdot b_{p,\alpha}(j, i_2) = 0 \text{ if } i_1 \neq i_2$$

Note that

$$A_{p^{\alpha}}(p^{i}, p^{j}) = \frac{p^{\alpha}}{(p^{\alpha}, p^{2i})} \cdot \frac{(p^{j}, p^{i})^{2}}{p^{j}} = \begin{cases} p^{\alpha-j} & \text{if } i \leq j \text{ and } i \leq \alpha/2, \\ p^{2i-j} & \text{if } i \leq j \text{ and } i > \alpha/2, \\ p^{\alpha-2i+j} & \text{if } i \geq j \text{ and } i \leq \alpha/2, \\ p^{j} & \text{if } i \geq j \text{ and } i > \alpha/2. \end{cases}$$

Now we begin to prove (28). In the case i = 0, we have

$$\sum_{j=0}^{\alpha} A_{p^{\alpha}}(p^{i}, p^{j}) \cdot b_{p,\alpha}(j, i) = A_{p^{\alpha}}(p^{0}, p^{0}) \cdot b_{p,\alpha}(0, 0) + A_{p^{\alpha}}(p^{0}, p^{1}) \cdot b_{p,\alpha}(1, 0)$$
$$= p^{\alpha} \cdot p + p^{\alpha - 1} \cdot (-1) = (p^{2} - 1)p^{\alpha - 1}$$

as required. In the case $i = \alpha$, we have

$$\sum_{j=0}^{\alpha} A_{p^{\alpha}}(p^{i}, p^{j}) \cdot b_{p,\alpha}(j, i) = A_{p^{\alpha}}(p^{\alpha}, p^{\alpha-1}) \cdot b_{p,\alpha}(\alpha - 1, \alpha) + A_{p^{\alpha}}(p^{\alpha}, p^{\alpha}) \cdot b_{p,\alpha}(\alpha, \alpha)$$
$$= p^{\alpha-1} \cdot (-1) + p^{\alpha} \cdot p = (p^{2} - 1)p^{\alpha-1}$$

as required. In the case $0 < i \le \alpha/2$ we have

$$\sum_{j=0}^{\alpha} A_{p^{\alpha}}(p^{i}, p^{j}) \cdot b_{p,\alpha}(j, i)$$

= $A_{p^{\alpha}}(p^{i}, p^{i-1}) \cdot b_{p,\alpha}(i-1, i) + A_{p^{\alpha}}(p^{i}, p^{i}) \cdot b_{p,\alpha}(i, i) + A_{p^{\alpha}}(p^{i}, p^{i+1}) \cdot b_{p,\alpha}(i+1, i)$
= $p^{\alpha-i-1} \cdot (-p^{\min(i,\alpha-i)}) + p^{\alpha-i} \cdot (p^{2}+1)p^{\min(i,\alpha-i)-1} + p^{\alpha-i-1} \cdot (-p^{\min(i,\alpha-i)})$
= $(p^{2}-1)p^{\alpha-1}$

as required. In the remaining case $\alpha/2 < i < \alpha$ we have

$$\sum_{j=0}^{\alpha} A_{p^{\alpha}}(p^{i}, p^{j}) \cdot b_{p,\alpha}(j, i)$$

= $A_{p^{\alpha}}(p^{i}, p^{i-1}) \cdot b_{p,\alpha}(i-1, i) + A_{p^{\alpha}}(p^{i}, p^{i}) \cdot b_{p,\alpha}(i, i) + A_{p^{\alpha}}(p^{i}, p^{i+1}) \cdot b_{p,\alpha}(i+1, i)$
= $p^{i-1} \cdot (-p^{\min(i,\alpha-i)}) + p^{i} \cdot (p^{2}+1)p^{\min(i,\alpha-i)-1} + p^{i-1} \cdot (-p^{\min(i,\alpha-i)})$
= $(p^{2}-1)p^{\alpha-1}$

which concludes the proof of (28). We can prove (29) in a similar manner which is tedious (there are so many cases) so we omit the details. \Box

Remark 5.2. There is another useful formula concerning the matrix A_N :

$$\sum_{c|N} \phi(c, N/c) \frac{N}{(N, c^2)} \frac{(n, c)^2}{n} = N \prod_{p|N} \left(1 + \frac{1}{p} \right).$$

In another words, the Euclidean inner product of each column of A_N and $(\phi(c, N/c))_{c|N}$ is equal to $N \prod_{p|N} \left(1 + \frac{1}{p}\right)$. This can be proved by applying Theorem 3.2 to $\eta(n\tau)$. As a consequence, $\sum_{c|N} \phi(c, N/c) x_c = N \prod_{p|N} \left(1 + \frac{1}{p}\right) \cdot \sum_{n|N} r_n$.

Example 5.3. We present some examples concerning dimensions of spaces of weight 2 for infinite many unitary characters. Suppose that N is square-free for simplicity. Let $\mathbf{x} = (x_c)_{c|N} \in \mathbb{Q}^{\mathscr{D}(N)}$ such that $\sum_{c|N} x_c = 0$. Then (20) with t = 1 holds since $\phi(c, N/c) = 1$ in the current case. Thus

(22) holds. According to the above remark we have $\sum_{n|N} r_n = m^{-1} \sum_{c|N} x_c = 0$ and hence k = 2. Therefore (22) is equivalent to

(30)
$$\dim_{\mathbb{C}} M_2(\Gamma_0(N), \chi) = g - 1 + \sigma_0(N) + \sum_{c|N} \left[\frac{x_c}{24}\right],$$

where g is the genus of $X_{\Gamma_0(N)}$. The multiplier system χ is the character (6) where $(r_n)_{n|N}$ is determined by **x** via (27) which in the current case is

(31)
$$r_n = \prod_{p|N} (p^2 - 1)^{-1} \cdot \sum_{c|N} \left((-1)^{\#\{p|N: v_p(n) \neq v_p(c)\}} \cdot \prod_{p|N, v_p(n) = v_p(c)} p \right) x_c.$$

More generally, if $N \in \mathbb{Z}_{\geq 1}$ is not divisible by 16 and has no odd square factor except 1, then we have $\phi(c, N/c) = 1$ for any $c \mid N$ as well and hence (30) is still valid. However in this case, (31) should be modified slightly at the prime factor 2.

Now we try to find out one-dimensional spaces among the above spaces. Note that

 $g-1 < \dim_{\mathbb{C}} M_2(\Gamma_0(N), \chi) \le g-1 + \sigma_0(N),$

so it is necessary g = 0 or 1 and N > 1. When g = 0, since $N \in \mathbb{Z}_{\geq 2}$ is not divisible by 16 and has no odd square factor except 1, then $N \in \{2, 3, 4, 5, 6, 7, 8, 10, 12, 13\}$. For such $N \dim_{\mathbb{C}} M_2(\Gamma_0(N), \chi) = 1$ if **x** satisfies

$$\sum_{c|N} x_c = 0, \quad \sum_{c|N} \left[\frac{x_c}{24}\right] = 2 - \sigma_0(N).$$

When g = 1, we have $N \in \{11, 14, 15, 17, 19, 20, 21, 24\}$. For such N, dim_C $M_2(\Gamma_0(N), \chi) = 1$ if **x** satisfies

$$\sum_{c|N} x_c = 0, \quad \sum_{c|N} \left[\frac{x_c}{24} \right] = 1 - \sigma_0(N).$$

Let us look at a very simple case N = 11. Since $\sigma_0(11) = 2$, dim_C $M_2(\Gamma_0(11), \chi) = 1$ if and only if $x_1 \notin 24\mathbb{Z}$ (of course $x_{11} = -x_1$). We have $r_1 = -r_{11} = x_1/10$ so χ is the multiplier system of $(\eta(\tau)/\eta(11\tau))^{x_1/10}$ which descends to a unitary character on $\Gamma_0(11)$. If $x_1 \in 24\mathbb{Z}$, then dim_C $M_2(\Gamma_0(11), \chi) = 2$. There are two completely different subcases in this case: $x_1 \in 120\mathbb{Z}$ and $x_1 \notin 120\mathbb{Z}$. If $x_1 \in 120\mathbb{Z}$, then χ is trivial and the formula reads dim_C $M_2(\Gamma_0(11)) = 2$ which is well known (cf. [29, Corollary 7.4.3]). Otherwise, if $x_1 \notin 120\mathbb{Z}$ then $x_1 \notin 10\mathbb{Z}$, which implies that χ is the character of an eta-quotient of rational exponents. To our best knowledge, the dimension formulas in such situation are new. Moreover, it seems that ker χ is a noncongruence subgroup of $SL_2(\mathbb{Z})$ so that our formulas may be applied to the theory of noncongruence modular forms.

6. The classification of characters induced by eta-quotients

In this section we give a complete classification of the characters of eta-quotients of level N and cover index 2 for any $N \in \mathbb{Z}_{\geq 1}$. The motivation is that, with this classification in hand, we can

know exactly the set of χ for which dim_C $M_k(\Gamma_0(N), \chi)$ can be calculated using Theorem 4.2. Since we only deal with the cover index D = 2, set $\widetilde{\Gamma_0(N)} = \widetilde{\Gamma_0(N)^2}$ in this section.

Now we establish some notations. Each $(r_n)_{n|N} \in \mathbb{Z}^{\mathscr{D}(N)}$ represents an eta-quotient $\prod_{n|N} \eta(n\tau)^{r_n}$ of level N and cover index 2 bijectively and $(r_n)_{n|N}$ with $\sum_n r_n = 2k$ represents an eta-quotient of weight k. Thus, set

$$S(N,k) = \left\{ (r_n)_{n|N} \in \mathbb{Z}^{\mathscr{D}(N)} \colon \sum_{n|N} r_n = 2k \right\}, \quad N \in \mathbb{Z}_{\geq 1}, \ k \in \frac{1}{2}\mathbb{Z},$$

and S(N) = S(N, 0). Then $\mathbb{Z}^{\mathscr{D}(N)} = \bigcup_{2k \in \mathbb{Z}} S(N, k)$ which is a disjoint union. In fact, S(N) is a submodule of $\mathbb{Z}^{\mathscr{D}(N)}$ and each S(N, k) is a coset in $\mathbb{Z}^{\mathscr{D}(N)}/S(N)$. The element $(r_n)_{n|N}$ in $\mathbb{Z}^{\mathscr{D}(N)}$ is customary to be written as $\mathbf{r} = \sum_{n|N} r_n e_n$ where $\{e_n : n \mid N\}$ is the standard basis of $\mathbb{Z}^{\mathscr{D}(N)}$. Moreover, the character (6) with D = 2 is denoted by $\chi_{\mathbf{r}}$ to emphasize its dependence upon $\mathbf{r} \in S(N, k)$.

Two different vectors \mathbf{r} , \mathbf{r}' in S(N, k) may correspond to the same character $\chi_{\mathbf{r}} = \chi_{\mathbf{r}'}$. The following lemma of Newman [38] tells us precisely when this happens.

Lemma 6.1. Let $N \in \mathbb{Z}_{\geq 1}$, $k \in \frac{1}{2}\mathbb{Z}$. Let $\mathbf{r} = (r_n)_{n|N}$ and $\mathbf{r}' = (r'_n)_{n|N}$ be vectors in S(N,k). Then $\chi_{\mathbf{r}} = \chi_{\mathbf{r}'}$ if and only if

$$\sum_{n|N} n \cdot (r_n - r'_n) \equiv 0 \pmod{24},$$
$$\sum_{n|N} \frac{N}{n} \cdot (r_n - r'_n) \equiv 0 \pmod{24},$$
$$\prod_{2 \nmid r_n - r'_n} n \text{ is a perfect square.}$$

Proof. This is a special case of the last assertion of [20, Theorem 3.9]. Setting l = 1 in that assertion and noting that $\chi_{\mathbf{r}} = \chi_{\mathbf{r}'}$ if and only if they are \tilde{I} -compatible give the desired equivalence.

Remark 6.2. It seems that this method does not work in the case $D \ge 3$ since we can not find any generalized double coset operator (cf. [20, Definition 1]) between characters of eta-quotients of cover index $D \ge 3$.

As a corollary, for $\mathbf{r} = (r_n)_{n|N} \in S(N)$, $\chi_{\mathbf{r}}$ is trivial if and only if $\sum_n n \cdot r_n \equiv 0 \mod 24$, $\sum_n \frac{N}{n} \cdot r_n \equiv 0 \mod 24$ and $\prod_{2 \nmid r_n} n$ is a perfect square. The set of all such \mathbf{r} is denoted by Z(N) which is a subgroup of S(N). We define

$$\mathscr{S}(N,k) = S(N,k)/Z(N) = \{\mathbf{r} + Z(N) \colon \mathbf{r} \in S(N,k)\},\$$

and $\mathscr{S}(N) = \mathscr{S}(N,0)$. Thus $\mathbb{Z}^{\mathscr{D}(N)}/Z(N) = \bigcup_{2k \in \mathbb{Z}} \mathscr{S}(N,k)$, which is a disjoint union. Note that elements of $\mathscr{S}(N,k)$ are in one-to-one correspondence with characters of eta-quotients of level N, cover index 2 and weight k according to Lemma 6.1, so to classify such characters we need only to obtain a complete system of representatives of $\mathscr{S}(N,k)$. Moreover, the map $\mathscr{S}(N) \to \mathscr{S}(N,k)$ that sends $\sum_{n|N} r_n e_n + Z(N)$ to $2ke_1 + \sum_{n|N} r_n e_n + Z(N)$ is a bijection. Therefore, we have reduced our task to the study of $\mathscr{S}(N)$. (For the result, see Theorem 6.8.)

Remark 6.3. Let $\chi_{\mathbf{r}} \colon \widetilde{\Gamma_0(N)} \to \mathbb{C}^{\times}$ be a character given by (6) where $\mathbf{r} \in S(N)$. Then $\chi_{\mathbf{r}}$ is the multiplier system of an eta-quotient of weight 0 and cover index 2. According to the third property in Remark 2.2, $\chi_{\mathbf{r}}$ descends to a character on $\Gamma_0(N)$. Thus $\mathscr{S}(N)$ can be regarded as a subgroup of the group of unitary linear characters of $\Gamma_0(N)$.

The sets given in the following definition are subsets of $\mathscr{D}(N)$ that occur in the complete system of representatives of $\mathscr{S}(N)$.

Definition 6.4. Let N be a positive integer and p be a prime divisor of N. If $p \neq 2, 3$ then define $\mathscr{B}_N^p = \{p\}$. For p = 2, we define

$$\mathscr{B}_{N}^{2} = \begin{cases} \{2^{1}\} & \text{if } v_{2}(N) = 1, \\ \{2^{1}, 2^{2}\} & \text{if } v_{2}(N) = 2, \\ \{2^{\alpha-2}, 2^{\alpha-1}, 2^{\alpha}\} & \text{if } v_{2}(N) = \alpha \ge 3 \end{cases}$$

For p = 3, we define

$$\mathscr{B}_{N}^{3} = \begin{cases} \{3^{1}\} & \text{if } v_{3}(N) = 1, \\ \{3^{\alpha-1}, 3^{\alpha}\} & \text{if } v_{3}(N) = \alpha \ge 2. \end{cases}$$

Finally, set $\mathscr{B}_N = \bigcup_{p|N} \mathscr{B}_N^p$. Note that $\mathscr{B}_1 = \emptyset$.

We introduce a useful notation which will be used in the proofs below: if $\mathbf{r}_1, \mathbf{r}_2 \in S(N)$, then $\mathbf{r}_1 \sim_N \mathbf{r}_2$, or simply $\mathbf{r}_1 \sim \mathbf{r}_2$, means $\mathbf{r}_1 - \mathbf{r}_2 \in Z(N)$. This is an equivalence relation compatible with the addition of S(N).

Lemma 6.5. Let $N = 2^{\alpha_0} \cdot 3^{\alpha_1} \cdot N_1$ with $gcd(N_1, 6) = 1$ and $\alpha_0, \alpha_1 \in \mathbb{Z}_{>0}$.

• For any $0 \leq \beta_0 \leq \alpha_0$, $0 \leq \beta_1 \leq \alpha_1$ and $n_1, n_2 \mid N_1$ with $gcd(n_1, n_2) = 1$, there exist $x, y, z \equiv 1 \mod 2$ such that

(32)

$$e_{2^{\beta_0}3^{\beta_1}n_1n_2} \sim x e_{2^{\beta_0}3^{\beta_1}n_1} + y e_{2^{\beta_0}3^{\beta_1}n_2} + z e_{2^{\beta_0}3^{\beta_1}}.$$

• For any $0 \leq \beta_0 \leq \alpha_0$, $0 \leq \beta_1 \leq \alpha_1$ and $n \mid N_1$, there exist $x, y \equiv 1 \mod 2$ and $z \in \mathbb{Z}$ such that

(33)
$$e_{2^{\beta_0}3^{\beta_1}n} \sim x e_{2^{\beta_0}3^{\beta_1}} + y e_n + z e_1.$$

• For any $0 \leq \beta_0 \leq \alpha_0$, $0 \leq \beta_1 \leq \alpha_1$, there exist $x, y, z \in \mathbb{Z}$ such that

(34)
$$e_{2^{\beta_0}3^{\beta_1}} \sim x e_{2^{\beta_0}} + y e_{3^{\beta_1}} + z e_1$$

Proof. First we prove (32). If $n_1 = 1$ or $n_2 = 1$ then (32) is immediate. Thus assume that $n_1, n_2 > 1$. By the definition of Z(N), a solution $(x, y, z) \in \mathbb{Z}^3$ of the system

(35)
$$\begin{cases} 2^{\beta_0} 3^{\beta_1} n_1 x + 2^{\beta_0} 3^{\beta_1} n_2 y + 2^{\beta_0} 3^{\beta_1} z \equiv 2^{\beta_0} 3^{\beta_1} n_1 n_2 \pmod{24} \\ \frac{N}{2^{\beta_0} 3^{\beta_1} n_1} x + \frac{N}{2^{\beta_0} 3^{\beta_1} n_2} y + \frac{N}{2^{\beta_0} 3^{\beta_1}} z \equiv \frac{N}{2^{\beta_0} 3^{\beta_1} n_1 n_2} \pmod{24} \\ x, y, z \equiv 1 \pmod{2} \end{cases}$$

is also a solution of (32). Moreover, a solution of the system

(36)
$$\begin{cases} n_1 x + n_2 y + z \equiv n_1 n_2 \pmod{24} \\ n_2 x + n_1 y + n_1 n_2 z \equiv 1 \pmod{24} \\ x, y, z \equiv 1 \pmod{2} \end{cases}$$

is a solution of (35). Since $2, 3 \nmid n_1, n_2$ we have $n_1^2 \equiv 1 \mod 24$ and $n_2^2 \equiv 1 \mod 24$. Thus the first equation of (36) is equivalent to the second one. Therefore let $x, y \equiv 1 \mod 2$ be arbitrary and set $z \equiv n_1 n_2 - n_1 x - n_2 y$; we obtain a solution of (36) and hence of (32).

The proofs of (33) and (34) are similar of which we omit the details.

Lemma 6.6. Any element in $\mathscr{S}(N)$ can be represented as

(37)
$$\sum_{n \in \mathscr{B}_N} c_n e_n - \left(\sum_{n \in \mathscr{B}_N} c_n\right) e_1 + Z(N),$$

where $c_n \in \mathbb{Z}$ and $0 \leq c_n < 24$ for any $n \in \mathscr{B}_N$.

As a consequence, $\mathscr{S}(N)$ is a finite group and $|\mathscr{S}(N)| \leq 24^{|\mathscr{B}_N|}.$

Proof. Let $\mathbf{r} + Z(N) = \sum r_n e_n + Z(N) \in \mathscr{S}(N)$ be arbitrary. According to Lemma 6.5 we can assume that $r_n = 0$ unless n is a prime power or n = 1. Therefore there exist $\mathbf{r}^{(p)} + Z(N) = \sum_{0 \leq \beta \leq v_p(N)} r_{p^{\beta}}^{(p)} e_{p^{\beta}} + Z(N) \in \mathscr{S}(N)$ such that $\mathbf{r} \sim \sum_{p|N} \mathbf{r}^{(p)}$. (Explicitly, let $r_{p^{\beta}}^{(p)} = r_{p^{\beta}}$ for $\beta > 0$ and $r_{p^0}^{(p)} = -\sum_{\beta > 0} r_{p^{\beta}}$.) It remains to find $c_n \in \mathbb{Z}$ such that

(38)
$$\mathbf{r}^{(p)} \sim \sum_{n \in \mathscr{B}_N^p} c_n e_n - \left(\sum_{n \in \mathscr{B}_N^p} c_n\right) e_1$$

and $0 \le c_n < 24$ for each $p \mid N$. If $p \ne 2, 3$ then (38) is equivalent to

$$\begin{cases} \sum_{1 \le \beta \le v_p(N)} (p^{\beta} - 1) r_{p^{\beta}}^{(p)} \equiv (p - 1) c_p \pmod{24} \\ \sum_{1 \le \beta \le v_p(N)} (N p^{-\beta} - N) r_{p^{\beta}}^{(p)} \equiv (N p^{-1} - N) c_p \pmod{24} \\ c_p \equiv \sum_{\beta \ge 1, r_{p^{\beta}}^{(p)} \equiv 1 \mod 2} \beta \pmod{2}. \end{cases}$$

Since $p^2 \equiv 1 \mod 24$, the solution of this system is $c_p \equiv \sum_{2 \nmid \beta} r_{p^\beta}^{(p)} \mod 24/(12, p-1)$. If p = 2 and $v_2(N) = \alpha \ge 3$, then (38) is equivalent to

$$\begin{cases} \sum_{1 \le \beta \le \alpha} (2^{\beta} - 1)r_{2^{\beta}}^{(2)} \equiv (2^{\alpha} - 1)c_{2^{\alpha}} + (2^{\alpha - 1} - 1)c_{2^{\alpha - 1}} + (2^{\alpha - 2} - 1)c_{2^{\alpha - 2}} \pmod{24} \\ \sum_{1 \le \beta \le \alpha} (N2^{-\beta} - N)r_{2^{\beta}}^{(2)} \equiv (N2^{-\alpha} - N)c_{2^{\alpha}} + (N2^{-\alpha + 1} - N)c_{2^{\alpha - 1}} + (N2^{-\alpha + 2} - N)c_{2^{\alpha - 2}} \pmod{24} \\ \sum_{\substack{1 \le \beta \le \alpha \\ 1 \le \beta \le \alpha}} \beta \equiv \sum_{1 \le \beta < \alpha - 2, r_{2^{\beta}}^{(2)} \equiv 1 \mod 2} \beta \pmod{2}. \end{cases}$$

By considering the subcases $2 \mid \alpha \geq 6$, $2 \nmid \alpha \geq 5$ and $\alpha = 3, 4$ separately and splitting the congruences modulo 24 to congruences modulo 3 and 8, one can verify that this system is solvable. We omit the proof of the case p = 2, $v_2(N) = 1, 2$. If p = 3 and $v_3(N) = \alpha \geq 2$, then (38) is equivalent to

$$\begin{cases} \sum_{1 \le \beta \le \alpha} (3^{\beta} - 1)r_{3^{\beta}}^{(3)} \equiv (3^{\alpha} - 1)c_{3^{\alpha}} + (3^{\alpha - 1} - 1)c_{3^{\alpha - 1}} \pmod{24} \\ \sum_{1 \le \beta \le \alpha} (N3^{-\beta} - N)r_{3^{\beta}}^{(3)} \equiv (N3^{-\alpha} - N)c_{3^{\alpha}} + (N3^{-\alpha + 1} - N)c_{3^{\alpha - 1}} \pmod{24} \\ \sum_{\substack{\beta \in \{\alpha - 1, \alpha\} \\ c_{3^{\beta}} \equiv r_{3^{\beta}}^{(3)} + 1 \mod 2}} \beta \equiv \sum_{1 \le \beta < \alpha - 1, r_{3^{\beta}}^{(3)} \equiv 1 \mod 2} \beta \pmod{2}. \end{cases}$$

As in the above case, by considering the subcases $2 \mid \alpha$ and $2 \nmid \alpha$ separately and splitting the congruences modulo 24 to congruences modulo 3 and 8, one can verify that this system is solvable. The case p = 3, $v_3(N) = 1$ is obvious. Thereby we have shown (38) which concludes the proof. \Box

To state the main theorem of this section, let us introduce the system of congruences

(39)
$$\begin{cases} \sum_{n \in \mathscr{B}_N} (n-1)c_n \equiv 0 \pmod{24} \\ \sum_{n \in \mathscr{B}_N} (Nn^{-1} - N)c_n \equiv 0 \pmod{24} \\ c_p \equiv 0 \pmod{2}, \ \forall p \mid N, p \neq 2, 3 \\ \sum_{2^\beta \in \mathscr{B}_N : \ c_{2\beta} \equiv 1 \mod 2} \beta \equiv 0 \pmod{2}, \quad \sum_{3^\beta \in \mathscr{B}_N : \ c_{3\beta} \equiv 1 \mod 2} \beta \equiv 0 \pmod{2}. \end{cases}$$

Then the element (37) is the zero in $\mathscr{S}(N)$ if and only if $(c_n)_{n \in \mathscr{B}_N}$ satisfies (39) according to the definition of Z(N).

Definition 6.7. Let $N \in \mathbb{Z}_{\geq 2}$. For any ordering b_1, b_2, \ldots, b_t of \mathscr{B}_N (which means $\mathscr{B}_N = \{b_1, b_2, \ldots, b_t\}$ and $b_i \neq b_j$ for $i \neq j$), we define a sequence of positive integers $\Delta_1, \Delta_2, \ldots, \Delta_t$

as follows:

$$\Delta_i = \min\{1 \le m \le 24 : \exists (c_n)_{n \in \mathscr{B}_N} \text{ s.t. } (39) \text{ holds and } c_{b_j} = 0 \ (j < i), \ c_{b_i} = m\}$$

Note that Δ_i is well defined since the set in the right-hand side always contains 24. In addition, be careful that the definition *does not* imply $\sum_{1 \le j \le t} \Delta_j e_j - \left(\sum_{1 \le j \le t} \Delta_j\right) e_1 \in Z(N)$ unless $|\mathscr{B}_N| = 1$.

Theorem 6.8. With the notation of Definition 6.7, a complete system of representatives of $\mathscr{S}(N)$ is given by

(40)
$$\sum_{1 \le j \le t} c_{b_j} e_{b_j} - \left(\sum_{1 \le j \le t} c_{b_j}\right) e_1, \quad 0 \le c_{b_j} < \Delta_j$$

1

Proof. Let $\mathbf{r} + Z(N) \in \mathscr{S}(N)$ be arbitrary. According to Lemma 6.6 we can assume that $\mathbf{r} = \sum_{n \in \mathscr{B}_N} r_n e_n - (\sum_{n \in \mathscr{B}_N} r_n) e_1$ with $r_n \in \mathbb{Z}$. By the definition of Δ_1 , there exists $\mathbf{d}_1 = \sum_{n \in \mathscr{B}_N} d_n e_n - (\sum_{n \in \mathscr{B}_N} d_n) e_1 \in Z(N)$ with $d_{b_1} = \Delta_1$. Hence subtracting a multiple of \mathbf{d}_1 from \mathbf{r} we may assume that $0 \leq r_{b_1} < \Delta_1$. Inductively, suppose we have adjusted \mathbf{r} such that $0 \leq r_{b_j} < \Delta_j$ for $j = 1, 2, \ldots, J$. Let $\mathbf{d}_{J+1} = \sum_{n \in \mathscr{B}_N} d_n e_n - (\sum_{n \in \mathscr{B}_N} d_n) e_1 \in Z(N)$ with $d_{b_j} = 0$ $(j \leq J)$ and $d_{b_{J+1}} = \Delta_{J+1}$. This exists by the definition of Δ_{J+1} . Subtracting a multiple of \mathbf{d}_{J+1} from \mathbf{r} we can assume that $0 \leq r_{b_j} < \Delta_j$ for $j = 1, 2, \ldots, J + 1$. Therefore by induction \mathbf{r} takes the form of (40) modulo Z(N). We have proved that each coset in $\mathscr{S}(N)$ can be represented by an element in (40).

Let $\mathbf{r}_1 = \sum_{1 \leq j \leq t} c_{b_j} e_{b_j} - \left(\sum_{1 \leq j \leq t} c_{b_j}\right) e_1$ and $\mathbf{r}_2 = \sum_{1 \leq j \leq t} d_{b_j} e_{b_j} - \left(\sum_{1 \leq j \leq t} d_{b_j}\right) e_1$ be two different representatives of the form (40). Let $1 \leq J \leq t$ satisfy $c_{b_j} = d_{b_j}$ for j < J and $c_{b_J} \neq d_{b_J}$. Without loss of generality assume that $c_{b_J} < d_{b_J}$. It follows that $0 < d_{b_J} - c_{b_J} < \Delta_J$. Then $\mathbf{r}_2 - \mathbf{r}_1 \notin Z(N)$ by the definition of Δ_J . We have proved that the vectors in (40) are pairwise inequivalent modulo Z(N), which concludes the whole proof.

Corollary 6.9. Let $N \in \mathbb{Z}_{\geq 1}$ and $k \in \frac{1}{2}\mathbb{Z}$. Then the characters (6) of eta-quotients of level N, weight k and cover index 2 are in one-to-one correspondence with the vectors

$$\sum_{1 \le j \le t} r_{b_j} e_{b_j} + \left(2k - \sum_{1 \le j \le t} r_{b_j}\right) e_1, \quad 0 \le r_{b_j} < \Delta_j.$$

Proof. According to the discussion followed by Remark 6.3, the characters of eta-quotients of level N, weight k and cover index 2 can be regarded as the set $\mathscr{S}(N, k)$ and the map $\mathscr{S}(N) \to \mathscr{S}(N, k)$, $\mathbf{r} + Z(N) \mapsto \mathbf{r} + 2ke_1 + Z(N)$ is a bijection. The corollary follows from this and Theorem 6.8. \Box

In the remainder of this section, we give examples of concrete sequences $(\Delta_i)_{1 \le i \le t}$ $(t = |\mathscr{B}_N|)$ for some special N.

Example 6.10. Let $N = p^{\alpha}$ where $p \geq 5$ is a prime and $\alpha \in \mathbb{Z}_{\geq 1}$. Then $\mathscr{B}_N = \{p\}$ and the sequence $(\Delta_i)_{1 \leq i \leq t}$ $(t = |\mathscr{B}_N|)$ contains a single element Δ_1 . The system of congruences (39) is

equivalent to $(p-1)c_p \equiv 0 \mod 24$ and $c_p \equiv 0 \mod 2$ in this case. Thus, $\Delta_1 = \frac{24}{(12,p-1)}$ and a complete system of representatives of $\mathscr{S}(p^{\alpha})$ is given by

$$r_p(e_p - e_1), \quad 0 \le r_p < \frac{24}{(12, p-1)}.$$

As a consequence, for any $k \in \frac{1}{2}\mathbb{Z}$ there are exactly $\frac{24}{(12,p-1)}$ characters $\chi: \widetilde{\Gamma_0(p^{\alpha})} \to \mathbb{C}^{\times}$ of etaquotients of level p^{α} , weight k and cover index 2. It should be noted that, the character of an eta-quotient, say, of cover index 4 may also be regarded as a character on $\widetilde{\Gamma_0(p^{\alpha})}$ (cf. the third fact of Remark 2.2). For instance, the character $\chi: \widetilde{\Gamma_0(p^{\alpha})^8} \to \mathbb{C}^{\times}$ of $\eta(\tau)^{1/4} \eta(p\tau)^{3/4}$ descends to a character on $\widetilde{\Gamma_0(p^{\alpha})} = \widetilde{\Gamma_0(p^{\alpha})^2}$. Such characters are not in the list above.

Example 6.11. Let $N = 2^{\alpha}$ with $\alpha \geq 3$. Then $\mathscr{B}_N = \{2^{\alpha-2}, 2^{\alpha-1}, 2^{\alpha}\}$ which is arranged in the increasing order. We need to calculate the sequence $\Delta_1, \Delta_2, \Delta_3$. The system of congruences (39) is equivalent to

$$\begin{cases} (2^{\alpha-2}-1)c_{2^{\alpha-2}} + (2^{\alpha-1}-1)c_{2^{\alpha-1}} + (2^{\alpha}-1)c_{2^{\alpha}} \equiv 0 \pmod{24} \\ (4-2^{\alpha})c_{2^{\alpha-2}} + (2-2^{\alpha})c_{2^{\alpha-1}} + (1-2^{\alpha})c_{2^{\alpha}} \equiv 0 \pmod{24} \\ \sum_{\substack{\beta \in \{\alpha-2,\alpha-1,\alpha\}\\c_{2^{\beta}} \equiv 1 \mod{2}}} \beta \equiv 0 \pmod{2}. \end{cases}$$

One can verify directly that

$$(\Delta_1, \Delta_2, \Delta_3) = \begin{cases} (2, 24, 8) & \text{if } \alpha \ge 3 \text{ and } 2 \mid \alpha \\ (2, 8, 24) & \text{if } \alpha \ge 3 \text{ and } 2 \nmid \alpha. \end{cases}$$

Therefore, a complete system of representatives of $\mathscr{S}(2^{\alpha})$ with $\alpha \geq 3$ is given by

$$r_{2^{\alpha-2}}e_{2^{\alpha-2}} + r_{2^{\alpha-1}}e_{2^{\alpha-1}} + r_{2^{\alpha}}e_{2^{\alpha}} - (r_{2^{\alpha-2}} + r_{2^{\alpha-1}} + r_{2^{\alpha}})e_1,$$

where $r_{\alpha-2} \in \{0,1\}$ and $0 \leq r_{2^{\alpha-1}} < 24$, $0 \leq r_{2^{\alpha}} < 8$ if $2 \mid \alpha$; $0 \leq r_{2^{\alpha-1}} < 8$, $0 \leq r_{2^{\alpha}} < 24$ if $2 \nmid \alpha$. As a consequence, for any $k \in \frac{1}{2}\mathbb{Z}$ there are exactly 384 characters $\chi \colon \Gamma_0(2^{\alpha}) \to \mathbb{C}^{\times}$ of eta-quotients of level 2^{α} , weight k and cover index 2 when $\alpha \geq 3$. In addition, it is immediate that for $N = 2^2$ we have $\mathscr{B}_4 = \{2^1, 2^2\}$ and $(\Delta_1, \Delta_2) = (24, 8)$ and for $N = 2^1$ we have $\mathscr{B}_2 = \{2^1\}$ and $\Delta_1 = 24$. Thus, there are exactly 192 (24 respectively) characters of eta-quotients of level 4 (2 respectively), cover index 2 and any fixed weight $\in \frac{1}{2}\mathbb{Z}$.

Example 6.12. Let $N = 3^{\alpha}$ with $\alpha \geq 2$. Then $\mathscr{B}_N = \{3^{\alpha-1}, 3^{\alpha}\}$ which is arranged in the increasing order. We need to calculate the sequence Δ_1, Δ_2 . The system of congruences (39) is

equivalent to

$$\begin{cases} (3^{\alpha-1}-1)c_{3^{\alpha-1}} + (3^{\alpha}-1)c_{3^{\alpha}} \equiv 0 \pmod{24} \\ (3-3^{\alpha})c_{3^{\alpha-1}} + (1-3^{\alpha})c_{3^{\alpha}} \equiv 0 \pmod{24} \\ \sum_{\substack{\beta \in \{\alpha-1,\alpha\}\\c_{3^{\beta}} \equiv 1 \mod 2}} \beta \equiv 0 \pmod{2}. \end{cases}$$

One can verify directly that

$$(\Delta_1, \Delta_2) = \begin{cases} (12, 3) & \text{if } \alpha \ge 2 \text{ and } 2 \mid \alpha \\ (3, 12) & \text{if } \alpha \ge 2 \text{ and } 2 \nmid \alpha. \end{cases}$$

Therefore, a complete system of representatives of $\mathscr{S}(3^{\alpha})$ with $\alpha \geq 2$ is given by

$$r_{3^{\alpha-1}}e_{3^{\alpha-1}} + r_{3^{\alpha}}e_{3^{\alpha}} - (r_{3^{\alpha-1}} + r_{3^{\alpha}})e_1,$$

where $0 \leq r_{3^{\alpha-1}} < 12$, $0 \leq r_{3^{\alpha}} < 3$ if $2 \mid \alpha$; $0 \leq r_{3^{\alpha-1}} < 3$, $0 \leq r_{3^{\alpha}} < 12$ if $2 \nmid \alpha$. As a consequence, for any $k \in \frac{1}{2}\mathbb{Z}$ there are exactly 36 characters $\chi \colon \widetilde{\Gamma_0(3^{\alpha})} \to \mathbb{C}^{\times}$ of eta-quotients of level 3^{α} , weight k and cover index 2 when $\alpha \geq 2$. In addition, it is direct that for $N = 3^1$ we have $\mathscr{B}_3 = \{3^1\}$ and $\Delta_1 = 12$. Thus, there are exactly 12 characters of eta-quotients of level 3, cover index 2 and any fixed weight $\in \frac{1}{2}\mathbb{Z}$.

Example 6.13. Let $N = 4p^{\alpha}$ where $p \ge 5$ is a prime and $\alpha \in \mathbb{Z}_{\ge 1}$. Then $\mathscr{B}_N = \{2^1, 2^2, p\}$ which is arranged in the increasing order. We need to calculate the sequence $\Delta_1, \Delta_2, \Delta_3$. The system of congruences (39) is equivalent to

$$\begin{cases} c_2 + 3c_4 + (p-1)c_p \equiv 0 \pmod{24} \\ -2p^{\alpha}c_2 - 3p^{\alpha}c_4 + (4p^{\alpha-1} - 4p^{\alpha})c_p \equiv 0 \pmod{24} \\ c_2, c_p \equiv 0 \pmod{2}. \end{cases}$$

Using elementary number theory one can show that

$$\Delta_1 = 2 \cdot (12, p-1), \quad \Delta_2 = 8, \quad \Delta_3 = \frac{24}{(12, p-1)}.$$

Therefore, a complete system of representatives of $\mathscr{S}(4p^{\alpha})$ with $p \geq 5$ and $\alpha \geq 1$ is given by

$$r_2e_2 + r_4e_4 + r_pe_p - (r_2 + r_4 + r_p)e_1$$

with

$$0 \le r_2 < 2 \cdot (12, p-1), \quad 0 \le r_4 < 8, \quad 0 \le r_p < \frac{24}{(12, p-1)}.$$

As a consequence, for any $k \in \frac{1}{2}\mathbb{Z}$ there are exactly 384 characters $\chi \colon \widetilde{\Gamma_0(4p^{\alpha})} \to \mathbb{C}^{\times}$ of etaquotients of level $4p^{\alpha}$, weight k and cover index 2 when $p \geq 5$ and $\alpha \geq 1$.

32

Example 6.14. Let $N = p_1^{\alpha_1} p_2^{\alpha_2}$ where $p_1, p_2 \ge 5$ are distinct primes and $\alpha_1, \alpha_2 \in \mathbb{Z}_{\ge 1}$. Then $\mathscr{B}_N = \{p_1, p_2\}$. We need to calculate the sequence Δ_1, Δ_2 . The system of congruences (39) is equivalent to

$$\begin{cases} (p_1 - 1)c_{p_1} + (p_2 - 1)c_{p_2} \equiv 0 \pmod{24} \\ (p_1^{\alpha_1 - 1}p_2^{\alpha_2} - p_1^{\alpha_1}p_2^{\alpha_2})c_{p_1} + (p_1^{\alpha_1}p_2^{\alpha_2 - 1} - p_1^{\alpha_1}p_2^{\alpha_2})c_{p_2} \equiv 0 \pmod{24} \\ c_{p_1}, c_{p_2} \equiv 0 \pmod{2}. \end{cases}$$

Using elementary number theory one can show that

$$\Delta_{1} = \frac{24 \cdot (12, p_{2} - 1)}{(12, (p_{1} - 1)(p_{2} - 1))} \cdot \left(\frac{12 \cdot (12, p_{2} - 1)}{(12, (p_{1} - 1)(p_{2} - 1))}, p_{1} - p_{2}\right)^{-1},$$

$$\Delta_{2} = \frac{24}{(12, p_{2} - 1)}.$$

The expression for Δ_1 seems to be complicated but it actually depends only on $p_1 \mod 12$ and $p_2 \mod 12$. For instance, if $p_2 \equiv 1 \mod 12$, then Δ_1 is simplified to $\frac{24}{(12,p_1-1)}$. A complete system of representatives of $\mathscr{S}(p_1^{\alpha_1}p_2^{\alpha_2})$ is given by

$$r_{p_1}e_{p_1} + r_{p_2}e_{p_2} - (r_{p_1} + r_{p_2})e_1$$

with

$$0 \le r_{p_1} < \Delta_1, \quad 0 \le r_{p_2} < \Delta_2$$

As a consequence, for any $k \in \frac{1}{2}\mathbb{Z}$ there are exactly $\Delta_1 \Delta_2$ characters $\chi \colon \Gamma_0(\widetilde{p_1^{\alpha_1}p_2^{\alpha_2}}) \to \mathbb{C}^{\times}$ of eta-quotients of level $p_1^{\alpha_1}p_2^{\alpha_2}$, weight k and cover index 2 when $p_1, p_2 \geq 5$ are distinct primes and $\alpha_1, \alpha_2 \geq 1$.

7. Dimension formulas for weights 1/2, 1, 3/2 and small levels

In this section, we give values of all $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi)$ that can be calculated by Theorem 4.2 (throughout this section, t in this theorem always takes the value 0) for k = 1/2, 1 and 3/2, $N \in \mathbb{Z}_{\geq 1}$ and χ being the character of an eta-quotient of level N, weight k and cover index 2. There are only finitely many such formulas since the right-hand side of (21) tends to 2 when $N \to +\infty$ and the set of χ is finite for each N and k (cf. Corollary 6.9). The following lemma gives an upper bound of N that should be considered.

Lemma 7.1. Let N be a positive integer and let m, ε_2 , ε_3 and ε_{∞} be defined in (9), (10), (11) and (12) respectively. If $N \ge 1024$, we have

$$\varepsilon_2 < \frac{m}{96}, \quad \varepsilon_3 < \frac{m}{128}, \quad \varepsilon_\infty \le \frac{m}{32}.$$

Consequently, in Theorem 4.2, if $k \leq 3/2$, t = 0 and $N \geq 1024$, then (21) does not hold.

For ε_{∞} , a stronger estimate⁸ is $\varepsilon_{\infty} \leq m \cdot N^{-1/2}$ with $N \in \mathbb{Z}_{\geq 1}$ which we will prove below.

⁸This is the best estimate if we want it to hold for all N since it becomes equality when N is a square.

Proof. Suppose $N = \prod_{1 \le j \le s} p_j^{\alpha_j}$ where $p_1 < p_2 < \cdots < p_s$ are primes, $s \in \mathbb{Z}_{\ge 0}$ and $\alpha_j \in \mathbb{Z}_{\ge 1}$. Note that for $p = p_j$, $\alpha = \alpha_j$ we have

$$\frac{p^{[\alpha/2]} + p^{[\alpha/2-1/2]}}{p^{\alpha} + p^{\alpha-1}} = \begin{cases} \frac{2}{p^{\alpha/2}(p^{1/2} + p^{-1/2})} & \text{if } 2 \nmid \alpha, \\ \frac{1}{p^{\alpha/2}} & \text{if } 2 \mid \alpha. \end{cases}$$

Thereby, according to [29, Corollary 6.3.24(b)]

(41)
$$\frac{\varepsilon_{\infty}}{m} = \prod_{1 \le j \le s} \frac{p_j^{[\alpha_j/2]} + p_j^{[\alpha_j/2-1/2]}}{p_j^{\alpha_j} + p_j^{\alpha_j-1}} \le \prod_{1 \le j \le s} \frac{1}{p_j^{\alpha_j/2}} = N^{-1/2}$$

Now we begin to prove the inequalities for ε_2 and ε_3 , so suppose $N \ge 1024$. If s = 1, 2, 3, then $\varepsilon_2/m \le 2^3/1025 < 1/96$, $\varepsilon_3/m \le 2^3/1025 < 1/128$. If $s \ge 4$, we consider two subcases: $\varepsilon_2 = 0$ and $\varepsilon_2 \ne 0$. If $\varepsilon_2 = 0$, obviously $\varepsilon_2/m < 1/96$. If $\varepsilon_2 \ne 0$, then $\left(\frac{-4}{p_j}\right) \ne -1$ and hence $p_j = 2$ or $p_j \equiv 1 \mod 4$ for $1 \le j \le s$. If $p_1 \ne 2$, then $p_1^{\alpha_1} \ge 5$, $p_2^{\alpha_2} \ge 13$ and $p_j^{\alpha_j} \ge 17$ for $j \ge 3$. Therefore

$$\frac{\varepsilon_2}{m} \le \frac{2}{5+1} \cdot \frac{2}{13+1} \cdot \left(\frac{2}{17+1}\right)^{s-2} < \frac{1}{96}$$

If $p_1 = 2$, then $p_2^{\alpha_2} \ge 5$, $p_3^{\alpha_3} \ge 13$ and $p_j^{\alpha_j} \ge 17$ for $j \ge 4$. Therefore

$$\frac{\varepsilon_2}{m} \le \frac{1}{2+1} \cdot \frac{2}{5+1} \cdot \frac{2}{13+1} \cdot \left(\frac{2}{17+1}\right)^{s-3} < \frac{1}{96}.$$

This proves the inequality for ε_2 . To prove the inequality for ε_3 when $s \ge 4$, we consider the subcases $\varepsilon_3 = 0$ and $\varepsilon_3 \neq 0$ separately and proceed as in the proof for ε_2 .

Finally, with the notation of Theorem 4.2 suppose that $k \leq 3/2$, t = 0 and $N \geq 1024$. Assume by contradiction that (21) holds, then

$$\frac{3}{2} \ge k > 2 - \frac{6}{m}\varepsilon_2 - \frac{8}{m}\varepsilon_3 - \frac{12}{m}\varepsilon_\infty > 2 - \frac{6}{96} - \frac{8}{128} - \frac{12}{32}$$

which is absurd. Hence (21) does not hold.

The algorithm for presenting all $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi)$ described above is as follows:

- (a) Let k range over $\{1/2, 1, 3/2\}$ and let N range over the positive integers ≤ 1023 . We do not miss anything according to Lemma 7.1.
- (b) Given k, N, we arrange \mathscr{B}_N in any order and calculate the sequence $\Delta_1, \ldots, \Delta_t$ (cf. Definition 6.7).
- (c) Let $\mathbf{r} = (r_n)_{n|N}$ range over the vectors described in Corollary 6.9. According to this corollary, the corresponding $\chi_{\mathbf{r}}$ ranges over exactly all characters of eta-quotients of level N, weight k and cover index 2.
- (d) Given k, N and \mathbf{r} , we calculate $\mathbf{x} = (x_c)_{c|N}$ via (19).

(e) We determine whether (21) holds. If it holds, then we calculate $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi_{\mathbf{r}})$ by (22) where t = 0. If it does not hold, $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi_{\mathbf{r}})$ can not be calculated by Theorem 4.2. Nevertheless we can obtain an upper bound and a lower bound of $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi_{\mathbf{r}})$ (cf. Remark 4.3).

Remark 7.2. For a fixed k, one can shrink the range of N by picking out those $N \leq 1023$ such that $k > 2 - \frac{6}{m}\varepsilon_2 - \frac{8}{m}\varepsilon_3 - \frac{12}{m}\varepsilon_{\infty}$. Hence, for k = 1/2, N need only to range over $\{1 \leq N \leq 21\} \cup \{24, 25, 27, 32, 36, 49, 50\}$; for k = 1, N need only to range over $\{1 \leq N \leq 22\} \cup \{24 \leq N \leq 32\} \cup \{34, 36, 37, 39, 40, 45, 48, 49, 50, 54, 64, 72, 75, 81, 98, 100, 121, 169\}$; for k = 3/2, N need only to range over a subset of $\{1 \leq N \leq 529\}$.

All the above steps can be easily implemented in any computer algebra system. We program them using SageMath. See Appendix A for the usage of the code.

7.1. The weight 1/2. The dimensions obtained by the above algorithm are summarized in Table 1.

TABLE 1. For each N, a means the total number of characters (cf. Corollary 6.9) and v means the number of characters χ such that $\dim_{\mathbb{C}} M_{1/2}(\Gamma_0(N), \chi)$ can be computed using Theorem 4.2, d_j means the number of spaces of dimension j for j = 0, 1, 2.

N	a	v	d_0	d_1	d_2	N	a	v	d_0	d_1	d_2
1	1	1	0	1	0	2	24	24	20	4	0
3	12	12	10	2	0	4	192	146	136	10	0
5	6	6	4	2	0	6	288	160	148	12	0
7	4	4	2	2	0	8	384	214	198	16	0
9	36	18	15	3	0	10	48	32	24	8	0
12	1152	318	286	32	0	13	2	2	0	2	0
16	384	127	108	18	1	18	288	56	42	12	2
20	384	6	0	6	0	24	2304	24	0	24	0
25	6	3	1	1	1	27	36	2	0	2	0
32	384	10	0	8	2	36	1152	12	0	6	6
49	4	1	0	0	1	50	48	4	0	0	4

Example 7.3. Let us consider the case N = 50. We have $\mathscr{B}_{50} = \{2, 5\}$ and $(\Delta_1, \Delta_2) = (8, 6)$ as one can verify directly. Thus, there are totally 48 characters of eta-quotients of level 50, weight 1/2 and cover index 2. They are exactly the characters on $\Gamma_0(50)^2$ of the following eta-quotients:

$$\eta(\tau)^{1-r_2-r_5}\eta(2\tau)^{r_2}\eta(5\tau)^{r_5}, \quad 0 \le r_2 < 8, \ 0 \le r_5 < 6$$

Among the 48 characters, for precisely the four characters χ given by

(42)
$$(r_2, r_5) = (0, 0), (1, 0), (2, 0), (7, 4)$$

one can compute $\dim_{\mathbb{C}} M_{1/2}(\Gamma_0(50), \chi)$ using Theorem 4.2 (that is, (21) holds with t = 0) and the four dimensions all equal 2. For the other 44 characters χ our method only gives that $0 \leq \dim_{\mathbb{C}} M_{1/2}(\Gamma_0(50), \chi) \leq 4$ (cf. Remark 4.3). Let $\chi_1, \chi_2, \chi_3, \chi_4$ denote respectively the characters given by (42). Then according to (14) and Lemma 6.1 we have

$$M_{1/2}(\Gamma_0(50), \chi_1) = \mathbb{C}\eta(\tau) \oplus \mathbb{C}\eta(25\tau),$$

$$M_{1/2}(\Gamma_0(50), \chi_2) = \mathbb{C}\eta(2\tau) \oplus \mathbb{C}\eta(50\tau),$$

$$M_{1/2}(\Gamma_0(50), \chi_3) = \mathbb{C}\eta(\tau)^{-1}\eta(2\tau)^2 \oplus \mathbb{C}\eta(25\tau)^{-1}\eta(50\tau)^2.$$

Finding explicit generators of $M_{1/2}(\Gamma_0(50), \chi_4)$ seems to be a bit harder since $\eta(\tau)^{-10}\eta(2\tau)^7\eta(5\tau)^4$ is only a weakly holomorphic modular form. However, we have $7e_2 + 4e_5 - 11e_1 \sim -e_2 + e_1$ module Z(50) which can be verified by (39). It follows that

$$M_{1/2}(\Gamma_0(50),\chi_4) = \mathbb{C}\eta(\tau)^2\eta(2\tau)^{-1} \oplus \mathbb{C}\eta(25\tau)^2\eta(50\tau)^{-1}.$$

Finally note that $M_{1/2}(\Gamma_0(50), \chi_j) = S_{1/2}(\Gamma_0(50), \chi_j)$ for j = 1, 2, while the Eisenstein subspace of $M_{1/2}(\Gamma_0(50), \chi_j)$ is nontrivial for j = 3, 4.

Example 7.4. We consider another example N = 20 in which there are totally 384 characters. They have been described in Example 6.13 with p = 5, $\alpha = 1$. We have $\mathscr{B}_{20} = \{2, 4, 5\}$ and $(\Delta_1, \Delta_2, \Delta_3) = (8, 8, 6)$. Therefore the characters are exactly those on $\Gamma_0(20)^2$ of the following eta-quotients:

$$\eta(\tau)^{1-r_2-r_4-r_5}\eta(2\tau)^{r_2}\eta(4\tau)^{r_4}\eta(5\tau)^{r_5}, \quad 0 \le r_2 < 8, \ 0 \le r_4 < 8, \ 0 \le r_5 < 6.$$

Among the 384 characters, for precisely the six characters χ given by

(43)
$$(r_2, r_4, r_5) = (1, 6, 1), (3, 0, 5), (3, 2, 5), (5, 6, 0), (7, 0, 4), (7, 2, 4)$$

one can compute $\dim_{\mathbb{C}} M_{1/2}(\Gamma_0(20), \chi)$ using Theorem 4.2 and the six dimensions all equal 1. For the other 378 characters χ our method only gives that $0 \leq \dim_{\mathbb{C}} M_{1/2}(\Gamma_0(20), \chi) \leq 2$ (cf. Remark 4.3). Let $\chi_1, \chi_2, \chi_3, \chi_4, \chi_5, \chi_6$ denote respectively the characters given by (43). One can verify using (14) and Lemma 6.1 that

$$\begin{split} M_{1/2}(\Gamma_0(20),\chi_1) &= \mathbb{C}\eta(5\tau)^{-2}\eta(10\tau)^5\eta(20\tau)^{-2}, \qquad M_{1/2}(\Gamma_0(20),\chi_2) = \mathbb{C}\eta(5\tau)^2\eta(10\tau)^{-1}, \\ M_{1/2}(\Gamma_0(20),\chi_3) &= \mathbb{C}\eta(10\tau)^{-1}\eta(20\tau)^2, \qquad M_{1/2}(\Gamma_0(20),\chi_4) = \mathbb{C}\eta(\tau)^{-2}\eta(2\tau)^5\eta(4\tau)^{-2}, \\ M_{1/2}(\Gamma_0(20),\chi_5) &= \mathbb{C}\eta(\tau)^2\eta(2\tau)^{-1}, \qquad M_{1/2}(\Gamma_0(20),\chi_6) = \mathbb{C}\eta(2\tau)^{-1}\eta(4\tau)^2. \end{split}$$

Remark 7.5. There are totally 188 nonzero spaces $M_{1/2}(\Gamma_0(N), \chi)$ whose dimensions can be computed by Theorem 4.2 according to Table 1. Note that we restrict ourselves to the characters χ of eta-quotients of cover index 2, so the dimension formulas summarized in Table 1 are not direct consequences of the well known theorem of Serre and Stark [14] that gives an explicit basis of $M_{1/2}(\Gamma_0(N), \chi_1\chi)$ where $4 \mid N, \chi_1$ is the character of Euler theta function $\theta(\tau) =$
$\eta(\tau)^{-2}\eta(2\tau)^5\eta(4\tau)^{-2}$ and χ is the character induced by any even Dirichlet character $(\mathbb{Z}/N\mathbb{Z})^{\times} \to \mathbb{C}^{\times}$. For instance, when N = 12, there are two even Dirichlet characters and hence the theorem of Serre and Stark gives explicit bases of two spaces $M_{1/2}(\Gamma_0(12), \chi)$. As a comparison, there are totally 1152 characters of eta-quotients of weight 1/2, level 12 and cover index 2 among which the dimensions of 318 spaces can be computed by Theorem 4.2—286 spaces are zero-dimensional and 32 spaces are one-dimensional.

Remark 7.6. Zagier observed that there are exactly 14 primitive (which means the greatest common divisor of all the *n* with $r_n \neq 0$ is equal to 1) eta-quotients of integral exponents that are modular forms of weight 1/2. This and the similar conjectures on higher weights were proved by Mersmann in his Master's thesis. See [21] for a simpler proof. It seems that all the generators of the 188 nonzero spaces $M_{1/2}(\Gamma_0(N), \chi)$ whose dimensions can be computed by Theorem 4.2 (cf. Table 1) take the form $f(m\tau)$ where f is one of the 14 primitive eta-quotients and m is a suitable positive integer. This is verified by the SageMath program for N = 4, 20, 50. See Appendix A for how to find the code doing this.

7.2. The weight 1. The dimensions obtained by the algorithm described above are summarized in Table 2. See Appendix A for the SageMath code that generates this kind of tables. Note that the characters are originally defined on $\Gamma_0(N)^2$ since we are considering eta-quotients of cover index 2 but they actually descend to characters on the matrix groups $\Gamma_0(N)$ since the weight is 1.

Table 2: For each N, a means the total number of characters (cf. Corollary 6.9) and v means the number of characters χ such that $\dim_{\mathbb{C}} M_1(\Gamma_0(N), \chi)$ can be computed using Theorem 4.2, d_j means the number of spaces of dimension j for $0 \leq j \leq 8$.

N	a	v	d_0	d_1	d_2	d_3	d_4	d_5	d_6	d_7	d_8
1	1	1	0	1	0	0	0	0	0	0	0
2	24	24	17	$\overline{7}$	0	0	0	0	0	0	0
3	12	12	7	5	0	0	0	0	0	0	0
4	192	173	142	31	0	0	0	0	0	0	0
5	6	6	3	3	0	0	0	0	0	0	0
6	288	250	189	60	1	0	0	0	0	0	0
7	4	4	1	3	0	0	0	0	0	0	0
8	384	331	247	83	1	0	0	0	0	0	0
9	36	29	17	11	1	0	0	0	0	0	0
10	48	46	20	22	4	0	0	0	0	0	0
11	12	1	0	1	0	0	0	0	0	0	0
12	1152	947	617	305	24	1	0	0	0	0	0
13	2	2	0	1	1	0	0	0	0	0	0

N	a	v	d_0	d_1	d_2	d_3	d_4	d_5	d_6	d_7	d_8
14	96	21	0	20	1	0	0	0	0	0	0
15	72	18	0	16	2	0	0	0	0	0	0
16	384	293	166	108	18	1	0	0	0	0	0
17	6	4	0	4	0	0	0	0	0	0	0
18	288	217	113	75	25	3	1	0	0	0	0
19	4	3	0	3	0	0	0	0	0	0	0
20	384	96	0	72	24	0	0	0	0	0	0
21	24	14	0	8	6	0	0	0	0	0	0
22	96	1	0	0	1	0	0	0	0	0	0
24	2304	658	0	480	176	0	2	0	0	0	0
25	6	4	1	1	1	1	0	0	0	0	0
26	48	4	0	0	4	0	0	0	0	0	0
27	36	15	0	9	5	1	0	0	0	0	0
28	384	11	0	0	10	1	0	0	0	0	0
30	576	2	0	0	0	0	2	0	0	0	0
32	384	146	0	100	36	8	2	0	0	0	0
36	1152	412	0	198	168	30	15	0	1	0	0
37	2	1	0	0	1	0	0	0	0	0	0
40	768	2	0	0	0	0	2	0	0	0	0
45	72	2	0	0	0	0	2	0	0	0	0
48	2304	58	0	0	0	16	40	0	2	0	0
49	4	2	0	0	1	0	1	0	0	0	0
50	48	26	0	0	19	0	0	$\overline{7}$	0	0	0
54	288	1	0	0	0	0	0	0	1	0	0
64	384	18	0	0	0	0	16	0	2	0	0
72	2304	2	0	0	0	0	0	0	0	0	2
75	72	2	0	0	0	0	0	0	2	0	0
81	36	1	0	0	0	0	0	0	1	0	0
98	96	1	0	0	0	0	0	0	0	0	1
121	12	1	0	0	0	0	0	0	1	0	0

We focus on the level 98 in the rest of this subsection. According to Table 2, there are totally 96 characters χ of eta-quotients of weight 1, level 98 and cover index 2. They correspond to 96 independent spaces $M_1(\Gamma_0(98), \chi)$ among which the dimension of exactly one space can be computed by Theorem 4.2, namely, the space where χ is the character of $\eta(\tau)^{-15}\eta(2\tau)^{16}\eta(7\tau)$. For this χ we have dim_C $M_1(\Gamma_0(98), \chi) = 8$. One can show that $\chi \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} -7 \\ d \end{pmatrix}$ by, for instance, a result of Gordon, Hughes and Newman (cf. [39, Theorem 1.64]) or by checking directly for $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ being in a set of generators of $\Gamma_0(98)$. Thus, an explicit basis can be given using Eisenstein series constructed by Weisinger [40]. See also [29, Theorem 8.5.23]. This shows there is no cusp form in $M_1(\Gamma_0(98), \chi)$, which can also be proved using the tools developed in [15]. One may ask: is the space $M_1(\Gamma_0(98), \chi)$ generated by holomorphic eta-quotients? The answer is no. According to (14), the functions

$$\begin{aligned} \eta(\tau)^{-1}\eta(2\tau)^2\eta(7\tau)^{-1}\eta(14\tau)^2 &= q + q^2 + q^4 + q^7 + q^8 + q^9 + 2q^{11} + q^{14} + \dots \\ \eta(\tau)^2\eta(2\tau)^{-1}\eta(7\tau)^2\eta(14\tau)^{-1} &= 1 - 2q + 2q^4 - 2q^7 + 4q^8 - 2q^9 - 4q^{11} + 6q^{16} + \dots \\ \eta(7\tau)^{-1}\eta(14\tau)^2\eta(49\tau)^{-1}\eta(98\tau)^2 &= q^7 + q^{14} + q^{28} + q^{49} + q^{56} + q^{63} + 2q^{77} + q^{98} + \dots \\ \eta(7\tau)^2\eta(14\tau)^{-1}\eta(49\tau)^2\eta(98\tau)^{-1} &= 1 - 2q^7 + 2q^{28} - 2q^{49} + 4q^{56} - 2q^{63} - 4q^{77} + 6q^{112} + \dots \end{aligned}$$

are holomorphic eta-quotients of level 98. Using Lemma 6.1 the characters of these four modular forms are equal to χ and hence they belong to $M_1(\Gamma_0(98), \chi)$. By their *q*-coefficients these four functions are linearly independent. However there are no other holomorphic eta-quotients of cover index 2 in $M_1(\Gamma_0(98), \chi)$ so this space can not be generated by a set consisting of all eta-quotients. This can be seen as follows. Let $f(\tau) = \prod_{n|98} \eta(n\tau)^{r_n}$ be in $M_1(\Gamma_0(98), \chi)$ with $r_n \in \mathbb{Z}$. Then $x_c \in \mathbb{Z}_{\geq 0}$ for any $c \mid 98$ where x_c is given by (19). According to Theorem 3.2 (applied to f), (14) and the complete set of representatives of $\Gamma_0(98) \setminus \mathbb{P}^1(\mathbb{Q})$ described in the proof of Proposition 3.4 we have $x_1 + x_2 + 6x_7 + 6x_{14} + x_{49} + x_{98} = 336$ of which there are only finitely many solutions $(x_c)_{c|98}$. For each solution, we obtain $(r_n)_{n|98}$ via (27) and then check whether $r_n \in \mathbb{Z}$ and the character corresponding to $(r_n)_{n|98}$ is the same as χ using Lemma 6.1. In this way we find that there are exactly four holomorphic eta-quotients of cover index 2 in $M_1(\Gamma_0(98), \chi)$ (totally 69 holomorphic eta-quotients, 57 of which are primitive).

7.3. The weight 3/2. There are totally 17862 spaces $M_{3/2}(\Gamma_0(N), \chi)$ whose dimensions can be calculated using Theorem 4.2. Among these spaces, the largest level is N = 400 and the largest dimension is 48. See Appendix A for the usage of SageMath code that gives these data. The following observation will be used later: the dimensions of these 17862 spaces are all greater than 1 when N > 36.

8. Application: one-dimensional spaces and Eisenstein series of rational weights

In [16, Section 7], the author gives infinitely many identities whose left-hand sides are Eisenstein series and right-hand sides are eta-quotients of rational weights for the levels 2 and 3. Using Theorem 4.2 (with t = 0), we find new identities of this kind for the level 4 in this section.

First let us rewrite Theorem 4.2 more explicitly for N = 4. We have m = 6, $\varepsilon_2 = \varepsilon_3 = 0$. Let r_1, r_2, r_4 be rational numbers and set $k = \frac{1}{2}(r_1 + r_2 + r_4)$. Let

(44)
$$\begin{pmatrix} x_1 \\ x_2 \\ x_4 \end{pmatrix} = \begin{pmatrix} 4 & 2 & 1 \\ 1 & 2 & 1 \\ 1 & 2 & 4 \end{pmatrix} \cdot \begin{pmatrix} r_1 \\ r_2 \\ r_4 \end{pmatrix}$$

Our aim is to find an Eisenstein series $E(\tau)$ of weight k equal to $f(\tau) = \eta(\tau)^{r_1} \eta(2\tau)^{r_2} \eta(4\tau)^{r_4}$. We only consider the Eisenstein series defined at the cusp i ∞ , so assume $x_4 = 0$ which means f does

not vanish at the cusp i ∞ . Since f must be a holomorphic eta-quotient, we assume $x_1, x_2 \ge 0$. Therefore,

 $x_1 = -6r_2 - 15r_4 \ge 0$, $x_2 = -3r_4 \ge 0$, hence $k = \frac{1}{2}(-r_2 - 3r_4) \ge 0$.

Now the first inequality in (21) becomes $k > -4 + 2\left(\left\{\frac{x_1}{24}\right\} + \left\{\frac{x_2}{24}\right\}\right)$, which holds since $k \ge 0$. Consequently, (22) with t = 0 is applicable. We summarize:

Proposition 8.1. Let r_2 and r_4 be rational numbers satisfying $-2r_2 - 5r_4 \ge 0$ and $-r_4 \ge 0$. Let $D \in \mathbb{Z}_{\ge 1}$ such that Dr_2 , $Dr_4 \in 2\mathbb{Z}$. Then we have

$$\dim_{\mathbb{C}} M_k(\Gamma_0(4), \chi) = \left[\frac{-2r_2 - 5r_4}{8}\right] + \left[\frac{-r_4}{8}\right] + 1,$$

where $\chi \colon \widetilde{\Gamma_0(4)^D} \to \mathbb{C}^{\times}$ is the character of $\eta(\tau)^{-2r_2-4r_4}\eta(2\tau)^{r_2}\eta(4\tau)^{r_4}$ and $k = \frac{1}{2}(-r_2-3r_4)$.

Proof. We have shown that (22) with t = 0 holds. A simplification gives the desired formula.

Corollary 8.2. Let the notation and assumptions be as in Proposition 8.1. Then $\dim_{\mathbb{C}} M_k(\Gamma_0(4), \chi) = 1$ if and only if

 $0 \le -2r_2 - 5r_4 < 8$ and $0 \le -r_4 < 8$.

Moreover, suppose $\dim_{\mathbb{C}} M_k(\Gamma_0(4), \chi) = 1$; then k > 2 if and only if $(-2r_2 - 5r_4) + (-r_4) > 8$.

Note that when $\dim_{\mathbb{C}} M_k(\Gamma_0(4), \chi) = 1$, then $M_k(\Gamma_0(4), \chi) = \mathbb{C}\eta(\tau)^{-2r_2-4r_4}\eta(2\tau)^{r_2}\eta(4\tau)^{r_4}$. Hence if $M_k(\Gamma_0(4), \chi)$ contains an Eisenstein series, then an identity that relates the eta-quotient and the Eisenstein series follows. In the rest we only consider the case k > 2 in which nonholomorphic Eisenstein series are avoided. Similar identities can also be derived in the cases k < 2, which we will do in the future work.

Recall the definition of Eisenstein series of rational weights greater than 2:

Definition 8.3 (Definition 3.1, [16]). Let D, N be positive integers, and let $k \in \frac{1}{D}\mathbb{Z}$ that is greater than 2. Let $\chi: \widetilde{\Gamma_0(N)^D} \to \mathbb{C}^{\times}$ be a finite index character such that $\chi(\widetilde{-I}) = \mathfrak{e}(-k/2)$. Let $s \in \mathbb{P}^1(\mathbb{Q})$ and $\gamma_s \in \mathrm{SL}_2(\mathbb{Z})$ such that $\gamma_s(\mathrm{i}\infty) = s$. If $\chi(\widetilde{\gamma_s}\widetilde{T}^{w_s}\widetilde{\gamma_s}^{-1}) = 1$, then we define the Eisenstein series $E_{\gamma_s,k}$ on the group $\widetilde{\Gamma_0(N)^D}$, of weight k, with character χ and at cusp γ_s as

$$E_{\gamma_s,k}(\tau) = \sum_{\gamma \in \widetilde{\gamma_s} \langle \widetilde{T}^{w_s} \rangle \widetilde{\gamma_s}^{-1} \setminus \widetilde{\Gamma_0(N)^D}} \chi(\gamma)^{-1} \cdot 1|_k \widetilde{\gamma_s}^{-1} \gamma.$$

In the above definition, w_s is the width (c.f. Section 3). If s = a/c where a, c are coprime integers, then $w_s = \frac{N}{(N,c^2)}$. The basic properties of these $E_{\gamma_s,k}$ are collected in [16, Section 3].

Now let N = 4 and let χ be as in Proposition 8.1. Since $\chi(\tilde{T}) = 1$ the Eisenstein series $E_{I,k}(\tau)$ is well-defined provided that k > 2, that is, $(-2r_2 - 5r_4) + (-r_4) > 8$. The main result of this section is the following:

Theorem 8.4. Let r_2 and r_4 be rational numbers satisfying

(45)
$$0 \le -2r_2 - 5r_4 < 8, \quad 0 \le -r_4 < 8, \quad (-2r_2 - 5r_4) + (-r_4) > 8.$$

Let D be a positive integer such that Dr_2 , $Dr_4 \in 2\mathbb{Z}$. Let $\chi \colon \widetilde{\Gamma_0(4)^D} \to \mathbb{C}^{\times}$ be the character of $\eta(\tau)^{-2r_2-4r_4}\eta(2\tau)^{r_2}\eta(4\tau)^{r_4}$. Set $k = \frac{1}{2}(-r_2-3r_4)$ and

$$E_{I,k}(\tau) = E_{I,k}(\tau; r_2, r_4) = \sum_{\gamma \in \langle \widetilde{T} \rangle \setminus \widetilde{\Gamma_0(4)^D}} \chi(\gamma)^{-1} \cdot 1|_k \gamma.$$

Then we have

(46)
$$\eta(\tau)^{-2r_2-4r_4}\eta(2\tau)^{r_2}\eta(4\tau)^{r_4} = \frac{1}{2D}E_{I,k}(\tau;r_2,r_4).$$

Proof. Corollary 8.2 shows that $\dim_{\mathbb{C}} M_k(\Gamma_0(4), \chi) = 1$. Let x_1, x_2 and x_4 be as in (44) with $r_1 = -2r_2 - 4r_4$. Then the assumptions imply that $x_1 \ge 0, x_2 \ge 0$ and $x_4 = 0$. Thus $\eta(\tau)^{-2r_2-4r_4}\eta(2\tau)^{r_2}\eta(4\tau)^{r_4}$ is holomorphic at all cusps and hence is in $M_k(\Gamma_0(4), \chi)$. By [16, Theorem 3.3] we have $E_{I,k} \in M_k(\Gamma_0(4), \chi)$. It follows that $\eta(\tau)^{-2r_2-4r_4}\eta(2\tau)^{r_2}\eta(4\tau)^{r_4} = c \cdot E_{I,k}$. Let $\Im \tau \to +\infty$; we find that $c = \frac{1}{2D}$ again by [16, Theorem 3.3].

In the rest of this section we will write out the Fourier expansion of $\frac{1}{2D}E_{I,k}(\tau)$ which will give us a more explicit form of the identity (46).

Lemma 8.5. For any rational numbers r_2, r_4 with $k = \frac{1}{2}(-r_2 - 3r_4) > 2$ we have

$$\frac{1}{2D}E_{I,k}(\tau;r_2,r_4) = 1 + \sum_{\substack{4|c>0,d\in\mathbb{Z}\\(c,d)=1}} \mathfrak{e}\left(-\frac{1}{24}P(c,d;r_2,r_4)\right) \cdot (c\tau+d)^{-k}$$

where

$$P(c, d; r_2, r_4) = r_2 \cdot \left(\Psi\begin{pmatrix}a & 2b\\c/2 & d\end{pmatrix} - 2\Psi\begin{pmatrix}a & b\\c & d\end{pmatrix}\right) + r_4 \cdot \left(\Psi\begin{pmatrix}a & 4b\\c/4 & d\end{pmatrix} - 4\Psi\begin{pmatrix}a & b\\c & d\end{pmatrix}\right)$$

(47)
$$= r_2 \cdot (12s(-d, c/2) - 24s(-d, c) + 3) + r_4 \cdot (12s(-d, c/4) - 48s(-d, c) + 9)$$

with a, b being any integers such that ad - bc = 1.

Proof. According to [16, Lemma 3.2],

$$E_{I,k}(\tau; r_2, r_4) = D \cdot \sum_{\substack{4 \mid c \in \mathbb{Z}, d \in \mathbb{Z} \\ (c,d)=1}} \chi \widetilde{\binom{a \ b}{c \ d}}^{-1} \cdot (c\tau + d)^{-k}$$

where a, b are any integers such that ad - bc = 1. By Remark 2.2, $\chi\left(-I\right) = \mathfrak{e}\left(-k/2\right)$. It follows that

$$\chi \widetilde{\left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right)}^{-1} \cdot (c\tau + d)^{-k} = \chi \widetilde{\left(\begin{smallmatrix} -a & -b \\ -c & -d \end{smallmatrix}\right)}^{-1} \cdot (-c\tau - d)^{-k}.$$

42

Thus

$$\frac{1}{2D}E_{I,k}(\tau; r_2, r_4) = 1 + \sum_{\substack{4|c>0, d \in \mathbb{Z} \\ (c,d)=1}} \chi(\widetilde{\left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right)^{-1}} \cdot (c\tau + d)^{-k}.$$

Substituting (6) and (5) in the above expression gives the desired formula.

Proposition 8.6. With the notation being as above, we have

$$\frac{1}{2D}E_{I,k}(\tau; r_2, r_4) = 1 + e^{-\pi i k/2} \frac{(2\pi)^k}{\Gamma(k)} \sum_{n \in \mathbb{Z}_{\ge 1}} n^{k-1} a(n) \cdot q^n,$$

where

$$a(n) = \sum_{c \in \mathbb{Z}_{\geq 1}} \frac{1}{(4c)^k} \sum_{\substack{0 \le d < 4c \\ (d,4c) = 1}} \mathfrak{e}\left(\frac{dn}{4c} - \frac{1}{24}P(4c,d;r_2,r_4)\right).$$

Proof. For $d \in \mathbb{Z}$, $c \in \mathbb{Z}_{\geq 1}$ set

$$f(d,c) = \begin{cases} \mathfrak{e}\left(-\frac{1}{24}P(c,d;r_2,r_4)\right) & \text{if } 4 \mid c \text{ and } (d,c) = 1; \\ 0 & \text{if } 4 \nmid c \text{ and } (d,c) = 1; \\ f\left(\frac{d}{(d,c)},\frac{c}{(d,c)}\right) & \text{if } (d,c) > 1. \end{cases}$$

Thus, $\frac{1}{2D}E_{I,k}(\tau; r_2, r_4) = 1 + \zeta(k)^{-1} \cdot \sum_{c \in \mathbb{Z}_{\geq 1}, d \in \mathbb{Z}} f(d, c)(c\tau + d)^{-k}$ since $\sum_{t \mid (c,d)} \mu(t) = 0$ if (c, d) > 1 and $\sum_t \frac{\mu(t)}{t^k} = \zeta(k)^{-1}$, where ζ is the Riemann zeta function and μ is the Möbius function. Note that f(d + c, c) = f(d, c). This fact, together with the Lipschitz summation formula

$$\sum_{n\in\mathbb{Z}}\frac{1}{(\tau+n)^s} = e^{-\pi i s/2}\frac{(2\pi)^s}{\Gamma(s)}\sum_{n\geq 1}n^{s-1}e^{2\pi i n\tau}, \quad \tau\in\mathfrak{H}, \, \Re(s)>1$$

implies that

$$\sum_{d\in\mathbb{Z}} f(d,c)(c\tau+d)^{-k} = c^{-k} \sum_{0\le d_1 < c} f(d_1,c) \sum_{d_0\in\mathbb{Z}} (\tau+\frac{d_1}{c}+d_0)^{-k}$$
$$= e^{-\pi i k/2} \frac{(2\pi)^k}{\Gamma(k)} c^{-k} \sum_{n\in\mathbb{Z}_{\ge 1}} n^{k-1} \sum_{0\le d < c} f(d,c) \mathfrak{e}\left(\frac{dn}{c}\right) q^n.$$

Therefore,

(48)
$$\frac{1}{2D}E_{I,k}(\tau; r_2, r_4) = 1 + e^{-\pi i k/2} \frac{(2\pi)^k}{\Gamma(k)\zeta(k)} \sum_{n \in \mathbb{Z}_{\ge 1}} n^{k-1} \sum_{c \in \mathbb{Z}_{\ge 1}} \frac{1}{c^k} \sum_{0 \le d < c} f(d, c) \mathfrak{e}\left(\frac{dn}{c}\right) q^n.$$

By the definition of f(d, c) we find that

$$\begin{split} \sum_{c \in \mathbb{Z}_{\geq 1}} \frac{1}{c^k} \sum_{0 \leq d < c} f(d,c) \mathfrak{e}\left(\frac{dn}{c}\right) &= \sum_{t \in \mathbb{Z}_{\geq 1}} \sum_{\substack{c \in \mathbb{Z}_{\geq 1}, 0 \leq d < c \\ (d,c) = t}} \frac{1}{(tc)^k} f(td,tc) \mathfrak{e}\left(\frac{dn}{c}\right) \\ &= \zeta(k) \cdot \sum_{\substack{c \in \mathbb{Z}_{\geq 1}, 0 \leq d < 4c \\ (d,4c) = 1}} \frac{1}{(4c)^k} f(d,4c) \mathfrak{e}\left(\frac{dn}{4c}\right) \\ &= \zeta(k) \cdot \sum_{\substack{c \in \mathbb{Z}_{\geq 1}, 0 \leq d < 4c \\ (d,4c) = 1}} \frac{1}{(4c)^k} \mathfrak{e}\left(\frac{dn}{4c} - \frac{1}{24} P(4c,d;r_2,r_4)\right). \end{split}$$

Inserting this into (48) gives the desired formula.

Remark 8.7. The reader may compare the above proposition with [16, Theorem 6.2].

We can now give an equivalent statement of Theorem 8.4 in a form that gives Fourier coefficients of certain infinite q-products:

Corollary 8.8. Let r_2 and r_4 be rational numbers satisfying (45). Let D be a positive integer such that Dr_2 , $Dr_4 \in 2\mathbb{Z}$. Let $\sum_{n \in \mathbb{Z}_{\geq 0}} c(n)q^n$ be the holomorphic branch of the D-th root of the infinite product

$$\prod_{n \in \mathbb{Z}_{\geq 1}} (1 - q^n)^{-2Dr_2 - 4Dr_4} \prod_{n \in \mathbb{Z}_{\geq 1}} (1 - q^{2n})^{Dr_2} \prod_{n \in \mathbb{Z}_{\geq 1}} (1 - q^{4n})^{Dr_4}$$

with c(0) = 1. Then for $n \ge 1$ we have

$$c(n) = e^{-\pi i k/2} \frac{(2\pi)^k}{\Gamma(k)} n^{k-1} \sum_{c \in \mathbb{Z}_{\ge 1}} \frac{1}{(4c)^k} \sum_{\substack{0 \le d < 4c \\ (d, 4c) = 1}} \mathfrak{e}\left(\frac{dn}{4c} - \frac{1}{24} P(4c, d; r_2, r_4)\right).$$

(See (47) for the definition of $P(4c, d; r_2, r_4)$.)

Proof. There are exactly D holomorphic branches and the one with the leading term 1 is the etaquotient $\eta(\tau)^{-2r_2-4r_4}\eta(2\tau)^{r_2}\eta(4\tau)^{r_4}$. Now the assertion follows from Theorem 8.4 and Proposition 8.6.

Example 8.9. Let us consider the special case $r_2 = 0$, $r_4 = -\frac{3}{2}$ for which (45) holds. Note that $k = \frac{1}{2}(-r_2 - 3r_4) = \frac{9}{4}$. Set D = 4. Then Corollary 8.8 shows

$$\frac{\prod_{n\in\mathbb{Z}_{\geq 1}}(1-q^n)^6}{\sqrt{\prod_{n\in\mathbb{Z}_{\geq 1}}(1-q^{4n})^3}} = 1 + \mathrm{e}^{-9\pi\mathrm{i}/8}\frac{(2\pi)^{\frac{9}{4}}}{\Gamma(\frac{9}{4})} \sum_{n\in\mathbb{Z}_{\geq 1}} n^{\frac{5}{4}} \left(\sum_{\substack{c\in\mathbb{Z}_{\geq 1}}}\frac{1}{(4c)^{\frac{9}{4}}}\sum_{\substack{0\leq d<4c\\(d,4c)=1}} \mathfrak{e}\left(\frac{dn}{4c} - \frac{1}{24}P(4c,d;0,-\frac{3}{2})\right)\right) \cdot q^n$$

where the square root is the principal branch $\sqrt{\prod_{n \in \mathbb{Z}_{\geq 1}} (1 - q^{4n})^3} = 1 + o(1)$ and

$$P(4c,d;0,-\frac{3}{2}) = -18s(-d,c) + 72s(-d,4c) - \frac{27}{2}.$$

(See (4) for the definition of s(-d, c).) For numerical check (of this and the next examples) using SageMath programs, see Appendix A.

Example 8.10. Corollary 8.8 contains as well some identities involving ordinary eta-quotients, that is, eta-quotients with integral exponents. For instance, set $r_2 = 7$, $r_4 = -4$ and hence $k = \frac{1}{2}(-r_2 - 3r_4) = \frac{5}{2}$ and D = 2. For this setting, (45) holds. The identity reads

$$\prod_{n \in \mathbb{Z}_{\geq 1}} (1 - q^n)^2 (1 - q^{2n})^7 (1 - q^{4n})^{-4}$$

= $1 + e^{-5\pi i/4} \frac{(2\pi)^{\frac{5}{2}}}{\Gamma(\frac{5}{2})} \sum_{n \in \mathbb{Z}_{\geq 1}} n^{\frac{3}{2}} \sum_{c \in \mathbb{Z}_{\geq 1}} \frac{1}{(4c)^{\frac{5}{2}}} \sum_{\substack{0 \le d < 4c \\ (d, 4c) = 1}} \mathfrak{e}\left(\frac{dn}{4c} - \frac{1}{24}P(4c, d; 7, -4)\right) \cdot q^n$

where

$$P(4c, d; 7, -4) = -48s(-d, c) + 84s(-d, 2c) + 24s(-d, 4c) - 15.$$

We will return to this function, i.e., $\eta(\tau)^2 \eta(2\tau)^7 \eta(4\tau)^{-4}$ in the next section (c.f. Section 9.4.1) and show that it is, among others, a Hecke eigenform in a generalized sense. (In [20, Section 4], the authors obtained some Hecke eigenforms in the generalized sense. However, for N = 4, only Hecke eigenforms of weight 1/2, 1, 3/2 are obtained. The current example gives a Hecke eigenform of weight 5/2 which can not be achieved using methods in [20].)

9. Application: An extension of Martin's list of multiplicative eta-quotients

In 1996 Martin [17] obtained the complete list of integral-weight holomorphic eta-quotients that are eigenforms for all Hecke operators. This generalized a previous result of Dummit, Kisilevsky and McKay [41]. The term "eta-quotient" in [17] refers to one whose multiplier system is induced by a Dirichlet character (necessarily real) and the Hecke operators are in the sense of [18].

In this section, we provide an algorithm for finding out all holomorphic eta-quotients $f(\tau) = \prod_{n\geq 1} \eta(n\tau)^{r_n}$ of integral exponents r_n (integral or half-integral weights) and of arbitrary multiplier systems such that either of the following conditions holds:

- (a) $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi)$ can be computed using Theorem 4.2 (with t = 0) and is equal to 1,
- (b) $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi)$ can be computed using Theorem 4.2 (with t = 0) and is greater than 1, f is a cusp form, k > 2 and there exist $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) - 1$ linearly independent Eisenstein series in $M_k(\Gamma_0(N), \chi)$ (c.f. Definition 8.3),

where $k = \frac{1}{2} \sum_{n} r_n$ is the weight, N is the least common multiple of all n with $r_n \neq 0$ (when $r_n = 0$ for all n, let N = 1) and χ is the character $\Gamma_0(N) \to \mathbb{C}^{\times}$ of f. (Note that throughout this section $\widetilde{\Gamma_0(N)} = \widetilde{\Gamma_0(N)^2}$ as in Section 6.)

If f satisfies either of the above conditions, we say it is *admissible*. If f satisfies the condition (a) then we say it is *admissible of type I*; if it satisfies (b) then we say it is *admissible of type II*. Moreover, if f satisfies either of the conditions

- (a') $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) = 1,$
- (b') $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) \ge 2$ but $\dim_{\mathbb{C}} S_k(\Gamma_0(N), \chi) = 1$ and f is a cusp form,

we say f is weakly admissible. Thus an admissible eta-quotient must be a weakly admissible etaquotient. Similarly, we can define the concept of weakly admissible eta-quotients of type I and type II. For each weakly admissible f, we will show that it is a Hecke eigenform in the sense of Wohlfahrt [19]; c.f. Theorem 9.19. Therefore, our list of admissible eta-quotients can be considered as an extension of Martin's list.

9.1. The algorithm for finding admissible eta-quotients. Throughout this subsection, let $f(\tau) = \prod_{n\geq 1} \eta(n\tau)^{r_n}$ be a holomorphic eta-quotient where r_n are integers and are equal to 0 for all but finitely many n. Let $k = \frac{1}{2} \sum_n r_n$ be the weight and N, the level⁹, be the least common multiple of $\{n \in \mathbb{Z}_{\geq 1} : r_n \neq 0\}$ (with the convention N = 1 if $r_n = 0$ for all n). Let $\chi: \widetilde{\Gamma_0(N)} \to \mathbb{C}^{\times}$ be the character of f (c.f. (6) with D = 2). Since f is holomorphic at all cusps we have $x_c \geq 0$ for all $c \mid N$ (x_c are defined in (19)). By Remark 4.3 we have

$$\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) \ge \frac{k-1}{12}m + \frac{1}{4}\varepsilon_2 + \frac{1}{3}\varepsilon_3 + \sum_{c|N} \phi(c, N/c) \cdot \left(\frac{1}{2} - \left\{\frac{x_c}{24}\right\}\right)$$
$$> \frac{k-1}{12}m - \frac{1}{2}\sum_{c|N} \phi(c, N/c)$$

where $m, \varepsilon_2, \varepsilon_3$ are given by (9), (10), (11) respectively.

Lemma 9.1. If $\left(\frac{(k-1)N}{12} - \frac{3\sqrt{N}}{2}\right) \cdot \prod_{p|N} \left(1 + \frac{1}{p}\right) \ge 1$, then f is not admissible. As a consequence, there are only finitely many admissible eta-quotients.

Proof. Assume by contradiction that f is admissible. If the condition (a) holds, then

(49)
$$1 = \dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) > \frac{k-1}{12}m - \frac{1}{2}\sum_{c|N} \phi(c, N/c).$$

By (41) we have $\varepsilon_{\infty} = \sum_{c|N} \phi(c, N/c) \leq m \cdot N^{-1/2}$. This, together with (49) and (9), implies that $\left(\frac{(k-1)N}{12} - \frac{\sqrt{N}}{2}\right) \cdot \prod_{p|N} \left(1 + \frac{1}{p}\right) < 1$ which contradicts the assumption. On the other hand, if the

⁹The reader should notice that this usage of "eta quotient of level N" is different from that in the second half of Section 2. In the usage here any eta-quotient has a uniquely determined level while in the usage before, an eta-quotient, say $\eta(\tau)\eta(2\tau)$, may be regarded as of level 2, or of level 4, etc.

condition (b) holds. Since k > 2, there are totally¹⁰ n_0 Eisenstein series in $M_k(\Gamma_0(N), \chi)$, where

(50)
$$n_0 = \# \left\{ s \in \Gamma_0(N) \setminus \mathbb{P}^1(\mathbb{Q}) \colon \chi(\widetilde{\gamma_s} \widetilde{T}^{w_s} \widetilde{\gamma_s}^{-1}) = 1 \right\}$$

(w_s is the width; c.f. Section 3, γ_s is any matrix in $\operatorname{SL}_2(\mathbb{Z})$ such that $\gamma_s(i\infty) = s$) and these n_0 functions are \mathbb{C} -linearly independent. See [16, Theorem 3.3]. By condition (b) we have $n_0 \geq \dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) - 1$. Since in the current case f is a cusp form, these n_0 Eisenstein series and f together constitute a basis of $M_k(\Gamma_0(N), \chi)$ and $n_0 = \dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) - 1$. Hence

(51)
$$n_0 + 1 = \dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) > \frac{k-1}{12}m - \frac{1}{2}\sum_{c|N} \phi(c, N/c).$$

This inequality, together with (41), (9) and the fact $n_0 \leq \varepsilon_{\infty} = \sum_{c|N} \phi(c, N/c)$, implies that $\left(\frac{(k-1)N}{12} - \frac{3\sqrt{N}}{2}\right) \cdot \prod_{p|N} \left(1 + \frac{1}{p}\right) < 1$ which contradicts the assumption. Therefore, we have shown that both of conditions (a) and (b) lead to contradictions, so f is not admissible.

Now we begin to prove that there are only finitely many admissible eta-quotients. Since $f \in M_k(\Gamma_0(N), \chi)$ is a holomorphic eta-quotient of integral exponents, $2k \in \mathbb{Z}_{\geq 0}$. For k = 0, we must have f = 1. For k = 1/2 and 1, we have shown in Remark 7.2 that there are only finitely many N such that $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi)$ can be calculated using Theorem 4.2. For each pair (k, N), the holomorphic eta-quotients of level N and weight k constitute a finite set since the cardinality of this set does not exceed the cardinality of the solution set of $\sum_{c|N} \phi(c, N/c)x_c = 2mk$, $x_c \in \mathbb{Z}_{\geq 0}$ (c.f. Proposition 5.1 and Remark 5.2) which is finite. This proves that there are only finitely many admissible eta-quotients of weights 0, 1/2 and 1. For each weight $k \geq 3/2$, there exists an N_0 such that when $N > N_0$ we have $\left(\frac{(k-1)N}{12} - \frac{3\sqrt{N}}{2}\right) \cdot \prod_{p|N} \left(1 + \frac{1}{p}\right) \geq 1$. Thus, the level of an admissible eta-quotient of weight k does not exceed N_0 and hence there are finitely many of them. Finally, if $k \geq 19$ and $N \geq 2$, we have

$$\left(\frac{(k-1)N}{12} - \frac{3\sqrt{N}}{2}\right) \cdot \prod_{p|N} \left(1 + \frac{1}{p}\right) \ge \frac{3}{2}(N - \sqrt{N}) \cdot \prod_{p|N} \left(1 + \frac{1}{p}\right) \ge 1,$$

and if $k \ge 31$, N = 1, then the left-hand side still ≥ 1 . It follows that if the weight of f is greater than or equal to 31, then f is not admissible. Therefore, there are only finitely many admissible eta-quotients.

Corollary 9.2. If N > 400, then f is not admissible. Moreover, if N > 36, then f is not admissible of type I.

Proof. Suppose N > 400; we have $k \neq 0$. If k = 1/2, 1 or 3/2, then by Tables 1, 2 and Section 7.3, dim_C $M_k(\Gamma_0(N), \chi)$ can not be calculated using Theorem 4.2 with t = 0 (that is, the first

¹⁰These n_0 functions, in the rational weight (k > 2) situation, constitute a basis of the Eisenstein space which, as in the integral weight case, is defined as the orthogonal complement of the space of cusp forms under the Petersson inner product. In addition, one can prove that $n_0 = |R|$ where R is (18) with f_0 arbitrarily chosen.

inequality in (21) does not hold) for any χ . Thus, f is not admissible¹¹. Otherwise, if $k \ge 2$, then obviously $\left(\frac{(k-1)N}{12} - \frac{3\sqrt{N}}{2}\right) \cdot \prod_{p|N} \left(1 + \frac{1}{p}\right) \ge 1$. It follows from this and Lemma 9.1 that f is not admissible.

Now we begin to prove the latter assertion; suppose N > 36. Certainly $k \neq 0$. According to Table 1, Table 2 and Subsection 7.3, if k = 1/2, 1 or 3/2, then $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) \geq 2$ provided it can be calculated using Theorem 4.2 (with t = 0). Hence f is not admissible of type I. If $k \geq 2$, then for $36 < N \leq 400$ but $N \neq 49$ we have

$$\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) > \frac{k-1}{12}m - \frac{1}{2}\sum_{c|N} \phi(c, N/c) > 1.$$

See Appendix A for the SageMath code verifying this. If N = 49, then $\varepsilon_3 = 2$ (c.f. (11)), and hence

$$\dim_{\mathbb{C}} M_k(\Gamma_0(49), \chi) \ge \frac{k-1}{12}m + \frac{1}{4}\varepsilon_2 + \frac{1}{3}\varepsilon_3 + \sum_{c|N} \phi(c, N/c) \cdot \left(\frac{1}{2} - \left\{\frac{x_c}{24}\right\}\right)$$
$$> \frac{2-1}{12} \cdot 56 + \frac{2}{3} - \frac{1}{2}\sum_{c|49} \phi(c, 49/c) > 1.$$

Therefore f is not admissible of type I for $k \ge 2$, $36 < N \le 400$.

According to the above corollary and Lemma 9.1, the following algorithm can be used to present all admissible nonconstant eta-quotients.

- (a) Let N range over $\{1, 2, 3, 4, \dots, 400\}$.
- (b) For a fixed N, set $K_N = \left\{ k \in \frac{1}{2}\mathbb{Z} : \frac{1}{2} \le k < 1 + \frac{12}{m} + \frac{18}{\sqrt{N}} \right\}$, where m is defined in (9).
- (c) Let N be fixed as above. For any $c \mid N$, let x_c range over nonnegative integers such that $\sum_{c \mid N} \phi(c, N/c) x_c \in 2m \cdot K_N$.
- (d) Let N and $(x_c)_{c|N}$ be fixed. Determine the sequence $(r_n)_{n|N}$ by (27) and set $k = \frac{1}{2} \sum_{n|N} r_n$, $g = \prod_{n|N} \eta(n\tau)^{r_n}$. Since $\sum_{c|N} \phi(c, N/c) x_c = m \cdot \sum_{n|N} r_n$ (c.f. Remark 5.2) we have $k \in K_N$.
- (e) If r_n are not all integers, or they are all integers but N is not the least common multiple of $\{r_n \neq 0 : n \mid N\}$, then we continue to the next value of $(x_c)_{c|N}$ in Step (d) since in these cases either g is not an eta-quotient of integral exponents or g has been considered in a previous step.
- (f) We check whether (21) with t = 0 holds. If it does not hold, g is not admissible so we go to Step (d) and consider the next value of $(x_c)_{c|N}$.
- (g) We check whether $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) = 1$ via (22). If this is the case, then we find an admissible g. We record it, go to Step (d) and consider the next value of $(x_c)_{c|N}$.
- (h) We check whether k > 2, g is a cusp form (equivalent to $x_c > 0$ for all $c \mid N$) and $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) = n_0 + 1$ where n_0 is given in (50). If this is the case, then g is

¹¹Although f is possibly weakly admissible.

admissible which we record; otherwise g is not admissible. We have completely determined whether this g is admissible or not so we go to Step (d) and consider the next value of $(x_c)_{c|N}$.

(i) Since for each N there are only finitely many values of $(x_c)_{c|N}$, this algorithm halts in finitely many steps.

Proposition 9.3. The above algorithm gives all admissible nonconstant eta-quotients with no repetition.

Proof. First, it is immediate that the functions g recorded in Steps (g) or (h) are admissible nonconstant eta-quotients. Second, suppose g_1 and g_2 are functions recorded in Steps (g) or (h) corresponding to different pairs $(N_1, (x_c)_{c|N_1})$ and $(N_2, (x'_c)_{c|N_2})$. If $N_1 \neq N_2$, then the exponents of g_1 are different from those of g_2 since their least common multiples are different. Hence $g_1 \neq g_2$. Otherwise $N_1 = N_2$ but their is some $c \mid N_1$ such that $x_c \neq x'_c$. Thus the sequences of exponents of g_1 and g_2 are different by Proposition 5.1, and hence $g_1 \neq g_2$ as well. Finally, let $f = \prod_{n \mid N_0} \eta(n\tau)^{r_n}$ be any admissible nonconstant eta-quotient whose level is N_0 and weight is k_0 . By Corollary 9.2 we have $1 \leq N_0 \leq 400$; by Lemma 9.1 we have $k_0 \in K_{N_0}$. Define $(y_c)_{c\mid N_0}$ by (19) (with $N = N_0$ and x_c replaced by y_c). It is immediate that $y_c \in \mathbb{Z}$. Moreover, $y_c \geq 0$ since f is holomorphic at all cusps. We have $\sum_{c\mid N_0} \phi(c, N_0/c)y_c = 2mk_0$ (c.f. Remark 5.2) and hence $\sum_{c\mid N_0} \phi(c, N_0/c)y_c \in 2m \cdot K_{N_0}$ where $m = N_0 \prod_{p\mid N_0} (1 + 1/p)^{-1}$. It follows that in Steps (a) and (c) the variable N can actually take the value N_0 and $(x_c)_{c\mid N}$ can take the value $(y_c)_{c\mid N_0}$. The corresponding g in Step (d) is equal to f according to Proposition 5.1. Therefore, f is recorded in some step of the algorithm which concludes the proof. □

Remark 9.4. If Step (h) is removed, then the remaining part of the algorithm gives all admissible nonconstant eta-quotients of type I. If this is the task, then we can shrink the set that N ranges over in Step (a) to $\{1, 2, 3, \ldots, 36\}$ by the latter assertion of Corollary 9.2. In the following subsections, we will only consider such eta-quotients because finding those of type II need more effort and the program takes rather long time¹² which we will do in a future paper. The simplest example of type II is $\eta(\tau)^{24}$ since it is a cusp form, dim_C $M_{12}(SL_2(\mathbb{Z})) = 2$ and there is an Eisenstein series in $M_{12}(SL_2(\mathbb{Z}))$. As a comparison, $\eta(\tau)^r$ is an admissible eta-quotient of type I for $r = 0, 1, 2, \ldots, 23$ since dim_C $M_{r/2}(SL_2(\mathbb{Z}), \chi_{\eta^r}) = 1$. See [20, Section 5].

9.2. The table of admissible eta-quotients of type I. For the SageMath program realizing the algorithm for finding admissible eta-quotients of type I in the last subsection, see Appendix A. Here (Table 3 below) we present all such eta-quotients for levels N = 1, p_1 , p_1^2 , p_1^3 and p_1p_2 where p_1 , p_2 are distinct primes.

 $^{^{12}}$ We have provided the SageMath code that can produce admissible eta-quotients of type II. See Appendix A for the usage.

Table 3: All admissible eta-quotients of type I of level $N = 1, p_1, p_1^2, p_1^3$ or p_1p_2 where p_1, p_2 are distinct primes. An entry $n_1^{r_1}n_2^{r_2}\cdots n_t^{r_t}$ represents the eta-quotient $\eta(n_1\tau)^{r_1}\eta(n_2\tau)^{r_2}\dots\eta(n_t\tau)^{r_t}$ and k is the weight.

N = 1								
1^r where $r =$	$0, 1, 2, \ldots, 23$							
N = 2, k = 1	1/2							
$1^{-1}2^2$	$1^{0}2^{1}$	$1^2 2^{-1}$						
N = 2, k = 1	_							
$1^{-2}2^4$	$1^{-1}2^3$	$1^{0}2^{2}$	$1^{1}2^{1}$	$1^{3}2^{-1}$	$1^4 2^{-2}$			
N = 2, k = 3	3/2							
$1^{-3}2^{6}$	$1^{-2}2^{5}$	$1^{-1}2^4$	$1^{0}2^{3}$	$1^{1}2^{2}$	$1^{2}2^{1}$			
$1^{4}2^{-1}$	$1^{5}2^{-2}$	$1^{6}2^{-3}$						
N = 2, k = 2								
$1^{-4}2^{8}$	$1^{-3}2^{7}$	$1^{-2}2^{6}$	$1^{-1}2^5$	$1^{0}2^{4}$	$1^{1}2^{3}$			
$1^{2}2^{2}$	$1^{3}2^{1}$	$1^{5}2^{-1}$	$1^{6}2^{-2}$	$1^7 2^{-3}$	$1^{8}2^{-4}$			
N = 2, k = 5/2								
$1^{-5}2^{10}$	$1^{-4}2^{9}$	$1^{-3}2^{8}$	$1^{-2}2^{7}$	$1^{-1}2^{6}$	$1^{0}2^{5}$			
$1^{1}2^{4}$	$1^{2}2^{3}$	$1^{3}2^{2}$	$1^{4}2^{1}$	$1^{6}2^{-1}$	$1^7 2^{-2}$			
$1^{8}2^{-3}$	$1^{9}2^{-4}$	$1^{10}2^{-5}$						
N = 2, k = 3	}							
$1^{-6}2^{12}$	$1^{-5}2^{11}$	$1^{-4}2^{10}$	$1^{-3}2^{9}$	$1^{-2}2^{8}$	$1^{-1}2^{7}$			
$1^{0}2^{6}$	$1^{1}2^{5}$	$1^{2}2^{4}$	$1^{3}2^{3}$	$1^{4}2^{2}$	$1^{5}2^{1}$			
$1^7 2^{-1}$	$1^{8}2^{-2}$	$1^{9}2^{-3}$	$1^{10}2^{-4}$	$1^{11}2^{-5}$	$1^{12}2^{-6}$			
N = 2, k = 7	7/2							
$1^{-7}2^{14}$	$1^{-6}2^{13}$	$1^{-5}2^{12}$	$1^{-4}2^{11}$	$1^{-3}2^{10}$	$1^{-2}2^{9}$			
$1^{-1}2^{8}$	$1^{0}2^{7}$	$1^{1}2^{6}$	$1^{2}2^{5}$	$1^{3}2^{4}$	$1^{4}2^{3}$			
$1^{5}2^{2}$	$1^{6}2^{1}$	$1^{8}2^{-1}$	$1^{9}2^{-2}$	$1^{10}2^{-3}$	$1^{11}2^{-4}$			
$1^{12}2^{-5}$	$1^{13}2^{-6}$	$1^{14}2^{-7}$						
N = 2, k = 4	Ł							
$1^{-7}2^{15}$	$1^{-6}2^{14}$	$1^{-5}2^{13}$	$1^{-4}2^{12}$	$1^{-3}2^{11}$	$1^{-2}2^{10}$			
$1^{-1}2^9$	$1^{0}2^{8}$	$1^{1}2^{7}$	$1^{2}2^{6}$	$1^{3}2^{5}$	$1^{4}2^{4}$			
$1^{5}2^{3}$	$1^{6}2^{2}$	$1^{7}2^{1}$	$1^{9}2^{-1}$	$1^{10}2^{-2}$	$1^{11}2^{-3}$			
$1^{12}2^{-4}$	$1^{13}2^{-5}$	$1^{14}2^{-6}$	$1^{15}2^{-7}$					
N = 2, k = 9	0/2							
$1^{-5}2^{14}$	$1^{-4}2^{13}$	$1^{-3}2^{12}$	$1^{-2}2^{11}$	$1^{-1}2^{10}$	$1^{0}2^{9}$			
$1^{1}2^{8}$	$1^{2}2^{7}$	$1^{3}2^{6}$	$1^{4}2^{5}$	$1^{5}2^{4}$	$1^{6}2^{3}$			
$1^{7}2^{2}$	$1^{8}2^{1}$	$1^{10}2^{-1}$	$1^{11}2^{-2}$	$1^{12}2^{-3}$	$1^{13}2^{-4}$			
				То	be continued			

Table 3. (c	continued)				
$1^{14}2^{-5}$					
$N = 2, k = 1^{-3}2^{13}$ $1^{3}2^{7}$ $1^{9}2^{1}$	$ \begin{array}{c} = 5 \\ 1^{-2} 2^{12} \\ 1^{4} 2^{6} \\ 1^{11} 2^{-1} \end{array} $	$ 1^{-1}2^{11} \\ 1^{5}2^{5} \\ 1^{12}2^{-2} $	$1^{0}2^{10}$ $1^{6}2^{4}$ $1^{13}2^{-3}$		$\frac{1^2 2^8}{1^8 2^2}$
$N = 2, k = 1^{-1}2^{12}$ $1^{5}2^{6}$ $1^{12}2^{-1}$	$= \frac{11/2}{1^{0}2^{11}}$ $1^{6}2^{5}$	$1^{1}2^{10}$ $1^{7}2^{4}$	$1^{2}2^{9}$ $1^{8}2^{3}$	$\frac{1^3 2^8}{1^9 2^2}$	$1^{4}2^{7}$ $1^{10}2^{1}$
$N = 2, k = 1^{12} + 1^{12} + 1^{7} + 2^{5}$	= 6 $1^{2}2^{10}$ $1^{8}2^{4}$	$1^{3}2^{9}$ $1^{9}2^{3}$	$1^{4}2^{8}$ $1^{10}2^{2}$	$1^{5}2^{7}$ $1^{11}2^{1}$	$1^{6}2^{6}$
$N = 2, k = 1^{3}2^{10}$ $1^{9}2^{4}$	$= \frac{13/2}{1^{4}2^{9}} \\ 1^{10}2^{3}$	$1^{5}2^{8}$	$1^{6}2^{7}$	$1^{7}2^{6}$	$1^{8}2^{5}$
$N = 2, k = 1^5 2^9$	$= 7 1^{6}2^{8}$	$1^{7}2^{7}$	$1^{8}2^{6}$	$1^{9}2^{5}$	
$N = 2, k = 1^7 2^8$	= 15/2 $1^{8}2^{7}$				
$N = 3, k = 1^0 3^1$	= 1/2				
$N = 3, k = 1^{-1}3^3$	= 1 $1^0 3^2$	$1^{1}3^{1}$	$1^{3}3^{-1}$		
$N = 3, k = 1^{-1}3^4$	= 3/2 $1^0 3^3$	$1^{1}3^{2}$	$1^{2}3^{1}$	$1^4 3^{-1}$	
$N = 3, k = 1^{-2}3^{6}$ $1^{5}3^{-1}$	$= \frac{2}{1^{-1}3^5} \\ 1^{6}3^{-2}$	$1^{0}3^{4}$	$1^{1}3^{3}$	$1^{2}3^{2}$	$1^{3}3^{1}$
$N = 3, k = 1^{-2}3^{7}$ $1^{4}3^{1}$	5/2 $1^{-1}3^{6}$ $1^{6}3^{-1}$	$ 1^{0}3^{5} \\ 1^{7}3^{-2} $	$1^{1}3^{4}$	$1^{2}3^{3}$	$1^{3}3^{2}$
$N = 3, k = 1^{-2}3^{8}$ $1^{4}3^{2}$	= 3 $1^{-1}3^{7}$ $1^{5}3^{1}$	$ 1^{0}3^{6} \\ 1^{7}3^{-1} $	$ 1^{1}3^{5} \\ 1^{8}3^{-2} $	$1^{2}3^{4}$	$1^{3}3^{3}$
$N = 3, k = 1^{-1}3^8$	= 7/2 $1^0 3^7$	$1^{1}3^{6}$	$1^{2}3^{5}$	$1^{3}3^{4}$	$1^{4}3^{3}$
					To be contir

Table 3. (cor	ntinued)				
$1^{5}3^{2}$	$1^{6}3^{1}$	$1^{8}3^{-1}$			
N = 3, k = 4 $1^{1}3^{7}$ $1^{7}3^{1}$	$1^{2}3^{6}$	$1^{3}3^{5}$	$1^{4}3^{4}$	$1^{5}3^{3}$	$1^{6}3^{2}$
N = 3, k = 9 $1^2 3^7$	$\frac{0}{2}$ 1 ³ 3 ⁶	$1^{4}3^{5}$	$1^{5}3^{4}$	$1^{6}3^{3}$	$1^{7}3^{2}$
N = 3, k = 5 $1^4 3^6$	$1^{5}3^{5}$	$1^{6}3^{4}$			
N = 3, k = 1 $1^5 3^6$	$\frac{1/2}{1^{6}3^{5}}$				
$N = 4, k = 1$ $1^{-2}2^{5}4^{-2}$	1/2 $1^{-1}2^{3}4^{-1}$	$1^{0}2^{-1}4^{2}$	$1^{0}2^{0}4^{1}$	$1^{0}2^{2}4^{-1}$	$1^{1}2^{-1}4^{1}$
N = 4, k = 1 $1^{-4}2^{10}4^{-4}$ $1^{-1}2^{1}4^{2}$ $1^{0}2^{0}4^{2}$ $1^{1}2^{0}4^{1}$	$\begin{array}{c} 1^{-3}2^{7}4^{-2} \\ 1^{-1}2^{2}4^{1} \\ 1^{0}2^{1}4^{1} \\ 1^{1}2^{2}4^{-1} \end{array}$	$ 1^{-3}2^{8}4^{-3} \\ 1^{-1}2^{4}4^{-1} \\ 1^{0}2^{3}4^{-1} \\ 1^{2}2^{-2}4^{2} $	$ \begin{array}{r} 1^{-2}2^{5}4^{-1} \\ 1^{-1}2^{5}4^{-2} \\ 1^{0}2^{4}4^{-2} \\ 1^{2}2^{-1}4^{1} \end{array} $	$ \begin{array}{r} 1^{-2}2^{6}4^{-2} \\ 1^{0}2^{-2}4^{4} \\ 1^{1}2^{-2}4^{3} \\ 1^{2}2^{1}4^{-1} \end{array} $	$ \begin{array}{r} 1^{-2}2^{7}4^{-3} \\ 1^{0}2^{-1}4^{3} \\ 1^{1}2^{-1}4^{2} \\ 1^{3}2^{-2}4^{1} \end{array} $
$N = 4, k = 3$ $1^{-6}2^{15}4^{-6}$ $1^{-4}2^{12}4^{-5}$ $1^{-2}2^{4}4^{1}$ $1^{-1}2^{1}4^{3}$ $1^{0}2^{-3}4^{6}$ $1^{0}2^{4}4^{-1}$ $1^{1}2^{0}4^{2}$ $1^{2}2^{-1}4^{2}$ $1^{3}2^{-1}4^{1}$	$\frac{3}{2}$ $\frac{1^{-5}2^{12}4^{-4}}{1^{-3}2^{7}4^{-1}}$ $\frac{1^{-2}2^{6}4^{-1}}{1^{-1}2^{2}4^{2}}$ $\frac{1^{0}2^{-2}4^{5}}{1^{0}2^{5}4^{-2}}$ $\frac{1^{1}2^{1}4^{1}}{1^{2}2^{0}4^{1}}$ $\frac{1^{3}2^{1}4^{-1}}{1^{3}2^{1}4^{-1}}$	$1^{-5}2^{13}4^{-5}$ $1^{-3}2^{8}4^{-2}$ $1^{-2}2^{7}4^{-2}$ $1^{-1}2^{3}4^{1}$ $1^{0}2^{-1}4^{4}$ $1^{0}2^{6}4^{-3}$ $1^{1}2^{3}4^{-1}$ $1^{2}2^{2}4^{-1}$ $1^{4}2^{-3}4^{2}$	$1^{-4}2^{9}4^{-2}$ $1^{-3}2^{9}4^{-3}$ $1^{-2}2^{8}4^{-3}$ $1^{-1}2^{5}4^{-1}$ $1^{0}2^{0}4^{3}$ $1^{1}2^{-3}4^{5}$ $1^{1}2^{4}4^{-2}$ $1^{2}2^{3}4^{-2}$ $1^{4}2^{-2}4^{1}$	$1^{-4}2^{10}4^{-3}$ $1^{-3}2^{10}4^{-4}$ $1^{-2}2^{9}4^{-4}$ $1^{-1}2^{6}4^{-2}$ $1^{0}2^{1}4^{2}$ $1^{1}2^{-2}4^{4}$ $1^{2}2^{-3}4^{4}$ $1^{3}2^{-3}4^{3}$ $1^{4}2^{0}4^{-1}$	$1^{-4}2^{11}4^{-4}$ $1^{-2}2^{3}4^{2}$ $1^{-1}2^{0}4^{4}$ $1^{-1}2^{7}4^{-3}$ $1^{0}2^{2}4^{1}$ $1^{1}2^{-1}4^{3}$ $1^{2}2^{-2}4^{3}$ $1^{3}2^{-2}4^{2}$ $1^{5}2^{-3}4^{1}$
$N = 4, k = 2$ $1^{-7}2^{17}4^{-6}$ $1^{-5}2^{11}4^{-2}$ $1^{-4}2^{10}4^{-2}$ $1^{-3}2^{6}4^{1}$ $1^{-2}2^{2}4^{4}$ $1^{-2}2^{9}4^{-3}$ $1^{-1}2^{2}4^{3}$ $1^{-1}2^{9}4^{-4}$ $1^{0}2^{2}4^{2}$ $1^{1}2^{-4}4^{7}$ $1^{1}2^{2}4^{1}$	$\begin{array}{c} 1^{-7}2^{18}4^{-7}\\ 1^{-5}2^{12}4^{-3}\\ 1^{-4}2^{11}4^{-3}\\ 1^{-3}2^{8}4^{-1}\\ 1^{-2}2^{3}4^{3}\\ 1^{-2}2^{10}4^{-4}\\ 1^{-1}2^{3}4^{2}\\ 1^{0}2^{-3}4^{7}\\ 1^{0}2^{3}4^{1}\\ 1^{1}2^{-3}4^{6}\\ 1^{1}2^{4}4^{-1} \end{array}$	$1^{-6}2^{14}4^{-4} \\ 1^{-5}2^{13}4^{-4} \\ 1^{-4}2^{12}4^{-4} \\ 1^{-3}2^{9}4^{-2} \\ 1^{-2}2^{4}4^{2} \\ 1^{-2}2^{11}4^{-5} \\ 1^{-1}2^{4}4^{1} \\ 1^{0}2^{-2}4^{6} \\ 1^{0}2^{5}4^{-1} \\ 1^{1}2^{-2}4^{5} \\ 1^{1}2^{5}4^{-2} \\ 1^{2}54^{-2$	$1^{-6}2^{15}4^{-5} \\ 1^{-5}2^{14}4^{-5} \\ 1^{-4}2^{13}4^{-5} \\ 1^{-3}2^{10}4^{-3} \\ 1^{-2}2^{5}4^{1} \\ 1^{-1}2^{-1}4^{6} \\ 1^{-1}2^{6}4^{-1} \\ 1^{0}2^{-1}4^{5} \\ 1^{0}2^{6}4^{-2} \\ 1^{1}2^{-1}4^{4} \\ 1^{1}2^{6}4^{-3} \\ 1^{-1}2^{6}4^{-3} \\ 1^{-1}2^{-1}4^{-1} \\ 1^{-1}2^{-1} \\ 1^$	$\begin{array}{c} 1^{-6}2^{16}4^{-6} \\ 1^{-5}2^{15}4^{-6} \\ 1^{-4}2^{14}4^{-6} \\ 1^{-3}2^{11}4^{-4} \\ 1^{-2}2^{7}4^{-1} \\ 1^{-1}2^{0}4^{5} \\ 1^{-1}2^{7}4^{-2} \\ 1^{0}2^{0}4^{4} \\ 1^{0}2^{7}4^{-3} \\ 1^{1}2^{0}4^{3} \\ 1^{2}2^{-4}4^{6} \end{array}$	$1^{-6}2^{17}4^{-7}$ $1^{-4}2^{9}4^{-1}$ $1^{-3}2^{5}4^{2}$ $1^{-3}2^{12}4^{-5}$ $1^{-2}2^{8}4^{-2}$ $1^{-1}2^{1}4^{4}$ $1^{-1}2^{8}4^{-3}$ $1^{0}2^{1}4^{3}$ $1^{0}2^{8}4^{-4}$ $1^{1}2^{1}4^{2}$ $1^{2}2^{-3}4^{5}$ be continued

51

()					
$1^2 2^{-2} 4^4$	$1^2 2^{-1} 4^3$	$1^2 2^0 4^2$	$1^2 2^1 4^1$	$1^2 2^3 4^{-1}$	$1^2 2^4 4^{-2}$
$1^2 2^5 4^{-3}$	$1^{3}2^{-4}4^{5}$	$1^{3}2^{-3}4^{4}$	$1^{3}2^{-2}4^{3}$	$1^3 2^{-1} 4^2$	$1^{3}2^{0}4^{1}$
$1^{3}2^{2}4^{-1}$	$1^{3}2^{3}4^{-2}$	$1^4 2^{-4} 4^4$	$1^4 2^{-3} 4^3$	$1^4 2^{-2} 4^2$	$1^4 2^{-1} 4^1$
$1^4 2^1 4^{-1}$	$1^4 2^2 4^{-2}$	$1^{5}2^{-4}4^{3}$	$1^5 2^{-3} 4^2$	$1^5 2^{-2} 4^1$	$1^{5}2^{0}4^{-1}$
$1^{6}2^{-4}4^{2}$	$1^{6}2^{-3}4^{1}$	$1^{6}2^{-1}4^{-1}$	$1^7 2^{-4} 4^1$		
N = 4, k =	5/2				
$1^{-7}2^{16}4^{-4}$	$1^{-7}2^{17}4^{-5}$	$1^{-7}2^{18}4^{-6}$	$1^{-6}2^{13}4^{-2}$	$1^{-6}2^{14}4^{-3}$	$1^{-6}2^{15}4^{-4}$
$1^{-6}2^{16}4^{-5}$	$1^{-6}2^{17}4^{-6}$	$1^{-6}2^{18}4^{-7}$	$1^{-5}2^{11}4^{-1}$	$1^{-5}2^{12}4^{-2}$	$1^{-5}2^{13}4^{-3}$
$1^{-5}2^{14}4^{-4}$	$1^{-5}2^{15}4^{-5}$	$1^{-5}2^{16}4^{-6}$	$1^{-5}2^{17}4^{-7}$	$1^{-4}2^{7}4^{2}$	$1^{-4}2^{8}4^{1}$
$1^{-4}2^{10}4^{-1}$	$1^{-4}2^{11}4^{-2}$	$1^{-4}2^{12}4^{-3}$	$1^{-4}2^{13}4^{-4}$	$1^{-4}2^{14}4^{-5}$	$1^{-4}2^{15}4^{-6}$
$1^{-4}2^{16}4^{-7}$	$1^{-3}2^{4}4^{4}$	$1^{-3}2^{5}4^{3}$	$1^{-3}2^{6}4^{2}$	$1^{-3}2^{7}4^{1}$	$1^{-3}2^{9}4^{-1}$
$1^{-3}2^{10}4^{-2}$	$1^{-3}2^{11}4^{-3}$	$1^{-3}2^{12}4^{-4}$	$1^{-3}2^{13}4^{-5}$	$1^{-3}2^{14}4^{-6}$	$1^{-2}2^{2}4^{5}$
$1^{-2}2^{3}4^{4}$	$1^{-2}2^{4}4^{3}$	$1^{-2}2^{5}4^{2}$	$1^{-2}2^{6}4^{1}$	$1^{-2}2^{8}4^{-1}$	$1^{-2}2^{9}4^{-2}$
$1^{-2}2^{10}4^{-3}$	$1^{-2}2^{11}4^{-4}$	$1^{-2}2^{12}4^{-5}$	$1^{-2}2^{13}4^{-6}$	$1^{-1}2^{0}4^{6}$	$1^{-1}2^{1}4^{5}$
$1^{-1}2^{2}4^{4}$	$1^{-1}2^{3}4^{3}$	$1^{-1}2^{4}4^{2}$	$1^{-1}2^{5}4^{1}$	$1^{-1}2^{7}4^{-1}$	$1^{-1}2^{8}4^{-2}$
$1^{-1}2^{9}4^{-3}$	$1^{-1}2^{10}4^{-4}$	$1^{-1}2^{11}4^{-5}$	$1^{0}2^{-1}4^{6}$	$1^{0}2^{0}4^{5}$	$1^{0}2^{1}4^{4}$
$1^{0}2^{2}4^{3}$	$1^{0}2^{3}4^{2}$	$1^{0}2^{4}4^{1}$	$1^{0}2^{6}4^{-1}$	$1^{0}2^{7}4^{-2}$	$1^{0}2^{8}4^{-3}$
$1^{0}2^{9}4^{-4}$	$1^{0}2^{10}4^{-5}$	$1^{1}2^{-3}4^{7}$	$1^{1}2^{-2}4^{6}$	$1^{1}2^{-1}4^{5}$	$1^{1}2^{0}4^{4}$
$1^{1}2^{1}4^{3}$	$1^{1}2^{2}4^{2}$	$1^{1}2^{3}4^{1}$	$1^{1}2^{5}4^{-1}$	$1^{1}2^{6}4^{-2}$	$1^{1}2^{7}4^{-3}$
$1^{1}2^{8}4^{-4}$	$1^{2}2^{-4}4^{7}$	$1^{2}2^{-3}4^{6}$	$1^{2}2^{-2}4^{5}$	$1^{2}2^{-1}4^{4}$	$1^{2}2^{0}4^{3}$
$1^{2}2^{1}4^{2}$	$1^{2}2^{2}4^{1}$	$1^{2}2^{4}4^{-1}$	$1^{2}2^{5}4^{-2}$	$1^{2}2^{6}4^{-3}$	$1^{2}2^{7}4^{-4}$
$1^{3}2^{-5}4^{7}$	$1^{3}2^{-4}4^{6}$	$1^{3}2^{-3}4^{5}$	$1^{3}2^{-2}4^{4}$	$1^{3}2^{-1}4^{3}$	$1^{3}2^{0}4^{2}$
$1^{3}2^{1}4^{1}$	$1^{3}2^{3}4^{-1}$	$1^{3}2^{4}4^{-2}$	$1^{3}2^{5}4^{-3}$	$1^{4}2^{-5}4^{6}$	$1^{4}2^{-4}4^{5}$
$1^{4}2^{-3}4^{4}$	$1^{4}2^{-2}4^{3}$	$1^{4}2^{-1}4^{2}$	$1^{4}2^{0}4^{1}$	$1^{4}2^{2}4^{-1}$	$1^{4}2^{3}4^{-2}$
$1^{4}2^{4}4^{-3}$	$1^{5}2^{-5}4^{5}$	$1^{5}2^{-4}4^{4}$	$1^{5}2^{-3}4^{3}$	$1^{5}2^{-2}4^{2}$	$1^{5}2^{-1}4^{1}$
$1^{5}2^{1}4^{-1}$	$1^{5}2^{2}4^{-2}$	$1^{6}2^{-5}4^{4}$	$1^{6}2^{-4}4^{3}$	$1^{6}2^{-3}4^{2}$	$1^{6}2^{-2}4^{1}$
$1^{6}2^{0}4^{-1}$	$1^7 2^{-5} 4^3$	$1^7 2^{-4} 4^2$	$1^7 2^{-3} 4^1$		
N = 4, k =	3				
$1^{-7}2^{15}4^{-2}$	$1^{-7}2^{16}4^{-3}$	$1^{-7}2^{17}4^{-4}$	$1^{-6}2^{13}4^{-1}$	$1^{-6}2^{14}4^{-2}$	$1^{-6}2^{15}4^{-3}$
$1^{-6}2^{16}4^{-4}$	$1^{-6}2^{17}4^{-5}$	$1^{-5}2^{9}4^{2}$	$1^{-5}2^{10}4^{1}$	$1^{-5}2^{12}4^{-1}$	$1^{-5}2^{13}4^{-2}$
$1^{-5}2^{14}4^{-3}$	$1^{-5}2^{15}4^{-4}$	$1^{-5}2^{16}4^{-5}$	$1^{-5}2^{17}4^{-6}$	$1^{-4}2^{7}4^{3}$	$1^{-4}2^{8}4^{2}$
$1^{-4}2^{9}4^{1}$	$1^{-4}2^{11}4^{-1}$	$1^{-4}2^{12}4^{-2}$	$1^{-4}2^{13}4^{-3}$	$1^{-4}2^{14}4^{-4}$	$1^{-4}2^{15}4^{-5}$
$1^{-4}2^{16}4^{-6}$	$1^{-4}2^{17}4^{-7}$	$1^{-3}2^{5}4^{4}$	$1^{-3}2^{6}4^{3}$	$1^{-3}2^{7}4^{2}$	$1^{-3}2^{8}4^{1}$
$1^{-3}2^{10}4^{-1}$	$1^{-3}2^{11}4^{-2}$	$1^{-3}2^{12}4^{-3}$	$1^{-3}2^{13}4^{-4}$	$1^{-3}2^{14}4^{-5}$	$1^{-3}2^{15}4^{-6}$
$1^{-3}2^{16}4^{-7}$	$1^{-2}2^{4}4^{4}$	$1^{-2}2^{5}4^{3}$	$1^{-2}2^{6}4^{2}$	$1^{-2}2^{7}4^{1}$	$1^{-2}2^{9}4^{-1}$
$1^{-2}2^{10}4^{-2}$	$1^{-2}2^{11}4^{-3}$	$1^{-2}2^{12}4^{-4}$	$1^{-2}2^{13}4^{-5}$	$1^{-2}2^{14}4^{-6}$	$1^{-2}2^{15}4^{-7}$
$1^{-1}2^{2}4^{5}$	$1^{-1}2^{3}4^{4}$	$1^{-1}2^{4}4^{3}$	$1^{-1}2^{5}4^{2}$	$1^{-1}2^{6}4^{1}$	$1^{-1}2^{8}4^{-1}$
$1^{-1}2^{9}4^{-2}$	$1^{-1}2^{10}4^{-3}$	$1^{-1}2^{11}4^{-4}$	$1^{-1}2^{12}4^{-5}$	$1^{-1}2^{13}4^{-6}$	$1^{0}2^{1}4^{5}$
$1^{0}2^{2}4^{4}$	$1^{0}2^{3}4^{3}$	$1^{0}2^{4}4^{2}$	$1^{0}2^{5}4^{1}$	$1^{0}2^{7}4^{-1}$	$1^{0}2^{8}4^{-2}$
$1^{0}2^{9}4^{-3}$	$1^{0}2^{10}4^{-4}$	$1^{0}2^{11}4^{-5}$	$1^{0}2^{12}4^{-6}$	$1^{1}2^{-1}4^{6}$	$1^{1}2^{0}4^{5}$
$1^{1}2^{1}4^{4}$	$1^{1}2^{2}4^{3}$	$1^{1}2^{3}4^{2}$	$1^{1}2^{4}4^{1}$	$1^{1}2^{6}4^{-1}$	$1^{1}2^{7}4^{-2}$
				Т	o be continued

Table 3. (continued)

	/				
$1^{1}2^{8}4^{-3}$	$1^{1}2^{9}4^{-4}$	$1^{1}2^{10}4^{-5}$	$1^{2}2^{-2}4^{6}$	$1^{2}2^{-1}4^{5}$	$1^{2}2^{0}4^{4}$
$1^{2}2^{1}4^{3}$	$1^2 2^2 4^2$	$1^{2}2^{3}4^{1}$	$1^{2}2^{5}4^{-1}$	$1^{2}2^{6}4^{-2}$	$1^{2}2^{7}4^{-3}$
$1^{2}2^{8}4^{-4}$	$1^{2}2^{9}4^{-5}$	$1^{3}2^{-4}4^{7}$	$1^{3}2^{-3}4^{6}$	$1^{3}2^{-2}4^{5}$	$1^{3}2^{-1}4^{4}$
$1^{3}2^{0}4^{3}$	$1^{3}2^{1}4^{2}$	$1^{3}2^{2}4^{1}$	$1^{3}2^{4}4^{-1}$	$1^{3}2^{5}4^{-2}$	$1^{3}2^{6}4^{-3}$
$1^{3}2^{7}4^{-4}$	$1^{4}2^{-5}4^{7}$	$1^{4}2^{-4}4^{6}$	$1^{4}2^{-3}4^{5}$	$1^{4}2^{-2}4^{4}$	$1^{4}2^{-1}4^{3}$
$1^{4}2^{0}4^{2}$	$1^{4}2^{1}4^{1}$	$1^{4}2^{3}4^{-1}$	$1^{4}2^{4}4^{-2}$	$1^{4}2^{5}4^{-3}$	$1^{5}2^{-6}4^{7}$
$1^{5}2^{-5}4^{6}$	$1^{5}2^{-4}4^{5}$	$152^{-3}4^{4}$	$1^{5}2^{-2}4^{3}$	$152^{-1}4^{2}$	$1^{5}2^{0}4^{1}$
$1^{5}2^{2}4^{-1}$	$1^{6}2^{-6}4^{6}$	$1^{6}2^{-5}4^{5}$	$1^{6}2^{-4}4^{4}$	$1^{6}2^{-3}4^{3}$	$1^{6}2^{-2}4^{2}$
$1^{6}2^{-1}4^{1}$	$1^{7}2^{-6}4^{5}$	$1^{7}2^{-5}4^{4}$	$1^{7}2^{-4}4^{3}$		
N = 4, k =	7/2				
$1^{-7}2^{15}4^{-1}$	$1^{-7}2^{16}4^{-2}$	$1^{-6}2^{12}4^{1}$	$1^{-6}2^{14}4^{-1}$	$1^{-6}2^{15}4^{-2}$	$1^{-6}2^{16}4^{-3}$
$1^{-5}2^{10}4^2$	$1^{-5}2^{11}4^{1}$	$1^{-5}2^{13}4^{-1}$	$1^{-5}2^{14}4^{-2}$	$1^{-5}2^{15}4^{-3}$	$1^{-5}2^{16}4^{-4}$
$1^{-4}2^{9}4^{2}$	$1^{-4}2^{10}4^{1}$	$1^{-4}2^{12}4^{-1}$	$1^{-4}2^{13}4^{-2}$	$1^{-4}2^{14}4^{-3}$	$1^{-4}2^{15}4^{-4}$
$1^{-4}2^{16}4^{-5}$	$1^{-3}2^{7}4^{\overline{3}}$	$1^{-3}2^{8}4^{\overline{2}}$	$1^{-3}2^{9}4^{\overline{1}}$	$1^{-3}2^{11}4^{-1}$	$1^{-3}2^{12}\bar{4}^{-2}$
$1^{-3}2^{13}4^{-3}$	$1^{-3}2^{14}4^{-4}$	$1^{-3}2^{15}4^{-5}$	$1^{-3}2^{16}4^{-6}$	$1^{-2}2^{6}4^{\overline{3}}$	$1^{-2}2^{7}4^{2}$
$1^{-2}2^{8}4^{1}$	$1^{-2}2^{10}4^{-1}$	$1^{-2}2^{11}4^{-2}$	$1^{-2}2^{12}4^{-3}$	$1^{-2}2^{13}4^{-4}$	$1^{-2}2^{14}4^{-5}$
$1^{-2}2^{15}4^{-6}$	$1^{-2}2^{16}4^{-7}$	$1^{-1}2^{4}4^{4}$	$1^{-1}2^{5}4^{3}$	$1^{-1}2^{6}4^{2}$	$1^{-1}2^{7}4^{1}$
$1^{-1}2^{9}4^{-1}$	$1^{-1}2^{10}4^{-2}$	$1^{-1}2^{11}4^{-3}$	$1^{-1}2^{12}4^{-4}$	$1^{-1}2^{13}4^{-5}$	$1^{-1}2^{14}4^{-6}$
$1^{-1}2^{15}4^{-7}$	$1^{0}2^{3}4^{4}$	$1^{0}2^{4}4^{3}$	$1^{0}2^{5}4^{2}$	$1^{0}2^{6}4^{1}$	$1^{0}2^{8}4^{-1}$
$1^{0}2^{9}4^{-2}$	$1^{0}2^{10}4^{-3}$	$1^{0}2^{11}4^{-4}$	$1^{0}2^{12}4^{-5}$	$1^{0}2^{13}4^{-6}$	$1^{0}2^{14}4^{-7}$
$1^{1}2^{1}4^{5}$	$1^{1}2^{2}4^{4}$	$1^{1}2^{3}4^{3}$	$1^{1}2^{4}4^{2}$	$1^{1}2^{5}4^{1}$	$1^{1}2^{7}4^{-1}$
$1^{1}2^{8}4^{-2}$	$1^{1}2^{9}4^{-3}$	$1^{1}2^{10}4^{-4}$	$1^{1}2^{11}4^{-5}$	$1^{1}2^{12}4^{-6}$	$1^2 2^0 4^5$
$1^{2}2^{1}4^{4}$	$1^2 2^2 4^3$	$1^2 2^3 4^2$	$1^{2}2^{4}4^{1}$	$1^{2}2^{6}4^{-1}$	$1^{2}2^{7}4^{-2}$
$1^{2}2^{8}4^{-3}$	$1^{2}2^{9}4^{-4}$	$1^{2}2^{10}4^{-5}$	$1^{3}2^{-2}4^{6}$	$1^{3}2^{-1}4^{5}$	$1^{3}2^{0}4^{4}$
$1^{3}2^{1}4^{3}$	$1^{3}2^{2}4^{2}$	$1^{3}2^{3}4^{1}$	$1^{3}2^{5}4^{-1}$	$1^{3}2^{6}4^{-2}$	$1^{3}2^{7}4^{-3}$
$1^{4}2^{-3}4^{6}$	$1^{4}2^{-2}4^{5}$	$1^4 2^{-1} 4^4$	$1^4 2^0 4^3$	$1^4 2^1 4^2$	$1^4 2^2 4^1$
$1^{4}2^{4}4^{-1}$	$1^{5}2^{-5}4^{7}$	$1^{5}2^{-4}4^{6}$	$1^{5}2^{-3}4^{5}$	$1^{5}2^{-2}4^{4}$	$1^5 2^{-1} 4^3$
$1^5 2^0 4^2$	$1^5 2^1 4^1$	$1^{6}2^{-6}4^{7}$	$1^{6}2^{-5}4^{6}$	$1^{6}2^{-4}4^{5}$	$1^{6}2^{-3}4^{4}$
$1^{6}2^{-2}4^{3}$	$1^7 2^{-7} 4^7$	$1^7 2^{-6} 4^6$	$1^7 2^{-5} 4^5$		
N = 4, k =	4				
$1^{-6}2^{15}4^{-1}$	$1^{-5}2^{12}4^{1}$	$1^{-5}2^{14}4^{-1}$	$1^{-5}2^{15}4^{-2}$	$1^{-4}2^{11}4^{1}$	$1^{-4}2^{13}4^{-1}$
$1^{-4}2^{14}4^{-2}$	$1^{-4}2^{15}4^{-3}$	$1^{-3}2^{9}4^{2}$	$1^{-3}2^{10}4^{1}$	$1^{-3}2^{12}4^{-1}$	$1^{-3}2^{13}4^{-2}$
$1^{-3}2^{14}4^{-3}$	$1^{-3}2^{15}4^{-4}$	$1^{-2}2^{8}4^{2}$	$1^{-2}2^{9}4^{1}$	$1^{-2}2^{11}4^{-1}$	$1^{-2}2^{12}4^{-2}$
$1^{-2}2^{13}4^{-3}$	$1^{-2}2^{14}4^{-4}$	$1^{-2}2^{15}4^{-5}$	$1^{-1}2^{6}4^{3}$	$1^{-1}2^{7}4^{2}$	$1^{-1}2^{8}4^{1}$
$1^{-1}2^{10}4^{-1}$	$1^{-1}2^{11}4^{-2}$	$1^{-1}2^{12}4^{-3}$	$1^{-1}2^{13}4^{-4}$	$1^{-1}2^{14}4^{-5}$	$1^{-1}2^{15}4^{-6}$
$1^{0}2^{5}4^{3}$	$1^{0}2^{6}4^{2}$	$1^{0}2^{7}4^{1}$	$1^{0}2^{9}4^{-1}$	$1^{0}2^{10}4^{-2}$	$1^{0}2^{11}4^{-3}$
$1^{0}2^{12}4^{-4}$	$1^{0}2^{13}4^{-5}$	$1^{0}2^{14}4^{-6}$	$1^{0}2^{15}4^{-7}$	$1^{1}2^{3}4^{4}$	$1^{1}2^{4}4^{3}$
$1^{1}2^{5}4^{2}$	$1^{1}2^{6}4^{1}$	$1^{1}2^{8}4^{-1}$	$1^{1}2^{9}4^{-2}$	$1^{1}2^{10}4^{-3}$	$1^{1}2^{11}4^{-4}$
$1^{1}2^{12}4^{-5}$	$1^2 2^2 4^4$	$1^2 2^3 4^3$	$1^2 2^4 4^2$	$1^2 2^5 4^1$	$1^2 2^7 4^{-1}$
$1^{2}2^{8}4^{-2}$	$1^2 2^9 4^{-3}$	$1^{3}2^{0}4^{5}$	$1^{3}2^{1}4^{4}$	$1^{3}2^{2}4^{3}$	$1^{3}2^{3}4^{2}$
$1^{3}2^{4}4^{1}$	$1^{3}2^{6}4^{-1}$	$1^4 2^{-1} 4^5$	$1^4 2^0 4^4$	$1^4 2^1 4^3$	$1^4 2^2 4^2$
				Т	o be continued

Table 3. (continued)

Table 3. (cor	ntinued)				
$1^4 2^3 4^1$	$1^{5}2^{-3}4^{6}$	$1^5 2^{-2} 4^5$	$1^5 2^{-1} 4^4$	$1^5 2^0 4^3$	$1^{6}2^{-4}4^{6}$
$1^{6}2^{-3}4^{5}$	$1^7 2^{-6} 4^7$				
N = 4, k = 9)/2				
$1^{-4}2^{14}4^{-1}$	$1^{-3}2^{11}4^{1}$	$1^{-3}2^{13}4^{-1}$	$1^{-3}2^{14}4^{-2}$	$1^{-2}2^{10}4^{1}$	$1^{-2}2^{12}4^{-1}$
$1^{-2}2^{13}4^{-2}$	$1^{-2}2^{14}4^{-3}$	$1^{-1}2^{8}4^{2}$	$1^{-1}2^{9}4^{1}$	$1^{-1}2^{11}4^{-1}$	$1^{-1}2^{12}4^{-2}$
$1^{-1}2^{13}4^{-3}$	$1^{-1}2^{14}4^{-4}$	$1^{0}2^{7}4^{2}$	$1^{0}2^{8}4^{1}$	$1^{0}2^{10}4^{-1}$	$1^{0}2^{11}4^{-2}$
$1^{0}2^{12}4^{-3}$	$1^{0}2^{13}4^{-4}$	$1^{0}2^{14}4^{-5}$	$1^{1}2^{5}4^{3}$	$1^{1}2^{6}4^{2}$	$1^{1}2^{7}4^{1}$
$1^{1}2^{9}4^{-1}$	$1^{1}2^{10}4^{-2}$	$1^{1}2^{11}4^{-3}$	$1^{2}2^{4}4^{3}$	$1^2 2^5 4^2$	$1^2 2^6 4^1$
$1^{2}2^{8}4^{-1}$	$1^{3}2^{2}4^{4}$	$1^{3}2^{3}4^{3}$	$1^{3}2^{4}4^{2}$	$1^{3}2^{5}4^{1}$	$1^4 2^1 4^4$
$1^4 2^2 4^3$	$1^5 2^{-1} 4^5$				
N = 4, k = 5	,)				
$1^{-2}2^{13}4^{-1}$	$1^{-1}2^{10}4^{1}$	$1^{-1}2^{12}4^{-1}$	$1^{-1}2^{13}4^{-2}$	$1^{0}2^{9}4^{1}$	$1^{0}2^{11}4^{-1}$
$1^{0}2^{12}4^{-2}$	$1^{0}2^{13}4^{-3}$	$1^{1}2^{7}4^{2}$	$1^{1}2^{8}4^{1}$	$1^{1}2^{10}4^{-1}$	$1^{2}2^{6}4^{2}$
$1^{2}2^{7}4^{1}$	$1^{3}2^{4}4^{3}$				
N = 4 $k = 1$	1/2				
$1^{0}2^{12}4^{-1}$	$1^{1}2^{9}4^{1}$				
N = 5, k = 1	/2				
$1^{0}5^{1}$					
N = 5, k = 1	_				
$1^{0}5^{2}$	$1^{1}5^{1}$				
$N = 5 \ k = 3$	8/2				
$1^{0} = 0, n = 0$ $1^{0} 5^{3}$	$1^{1}5^{2}$	$1^{2}5^{1}$			
	2.0	2.0			
N = 5, k = 2	11-3	12-2	13-1		
1.9.	1.2	1-9-	1.2.		
N = 5, k = 5	5/2		4 4		
$1^{1}5^{4}$	$1^{2}5^{3}$	$1^{3}5^{2}$	$1^{4}5^{1}$		
N = 5, k = 3	3				
$1^{2}5^{4}$	$1^{3}5^{3}$	$1^{4}5^{2}$			
$N = 5. \ k = 7$	7/2				
$1^{3}5^{4}$	$1^{4}5^{3}$				
N - 6 k - 1	/2				
$1^{-1}2^{1}3^{2}6^{-1}$	$1^{-1}2^{2}3^{1}6^{-1}$	$1^{0}2^{0}3^{-1}6^{2}$	$1^{0}2^{0}3^{0}6^{1}$	$1^{0}2^{0}3^{2}6^{-1}$	$1^{1}2^{-1}3^{-1}6^{2}$
$1^{2}2^{-1}3^{-1}6^{1}$	1 200	1200	1200	1200	1200
N C 1 1					
N = 0, k = 1 1-20204 <i>c</i> -2	1-203026-1	1-20303c-2	1 - 204 2 1 c - 1	1 - 20492c - 2	1-1919161
1 - 2 - 3 - 0 - 1 - 1 - 1 - 1 - 2 - 2 - 0	$1 -2^{-} 3^{-} 0^{-1}$	$1 -2^{\circ} 3^{\circ} 0^{-2}$ 1 -1 9 1 9 4 c -2	1 - 2 - 3 - 0 - 1 - 1 - 1 - 2 - 2 - 1 - 2 - 2 - 1 - 2 - 2	1 - 2 - 3 - 0 - 1 - 1 - 1 - 2 - 2 - 0 - 2 - 1 - 1 - 1 - 2 - 2 - 0 - 2 - 1 - 1 - 1 - 1 - 2 - 2 - 2 - 2 - 2	1 -2 - 3 - 0 - 1 - 1 - 1 - 2 - 2 - 1 - 0 - 0 - 0 - 0 - 0 - 0 - 0 - 0 - 0
$1 2 3^{-}0^{-}$ $1^{-1}9^{2}9^{2}\kappa^{-1}$	$1 - 2 - 3^{\circ} 0^{-1}$ 1 - 1 - 1 - 2 - 2 - 3 - 2	$1 2 3 0 -1 \\ 1 - 1 9 3 2 1 \\ \kappa - 1 -1 0 -1 -1$	$1 2 3 0^{-1}$ $109 - 120 \beta^{3}$	$1 2 3^{\circ} 0^{-1} \\ 10 90 9 9 9 9 9 9 9 9 $	エーム ひ ⁻ 0° 1090マー1 ₆ 3
1 2 3 0	1 2 3 0	1 200	12 30	1 4 J U T-	1 2 0 U
				10	be continued

To be continued

$1^{0}2^{0}3^{0}6^{2}$	$1^{0}2^{0}3^{1}6^{1}$	$1^{0}2^{0}3^{3}6^{-1}$	$1^{0}2^{0}3^{4}6^{-2}$	$1^{0}2^{1}3^{-1}6^{2}$	$1^{0}2^{1}3^{0}6^{1}$
$1^{0}2^{1}3^{1}6^{0}$	$1^{0}2^{1}3^{2}6^{-1}$	$1^{0}2^{2}3^{1}6^{-1}$	$1^{0}2^{3}3^{0}6^{-1}$	$1^{1}2^{-1}3^{-2}6^{4}$	$1^{1}2^{-1}3^{-1}6^{3}$
$1^{1}2^{-1}3^{0}6^{2}$	$1^{1}2^{-1}3^{1}6^{1}$	$1^{1}2^{0}3^{-1}6^{2}$	$1^{1}2^{0}3^{0}6^{1}$	$1^{1}2^{0}3^{2}6^{-1}$	$1^{1}2^{1}3^{-1}6^{1}$
$1^{1}2^{1}3^{1}6^{-1}$	$1^2 2^{-2} 3^{-2} 6^4$	$1^2 2^{-1} 3^{-2} 6^3$	$1^2 2^{-1} 3^{-1} 6^2$	$1^2 2^{-1} 3^0 6^1$	$1^2 2^{-1} 3^1 6^0$
$1^2 2^{-1} 3^2 6^{-1}$	$1^{2}2^{0}3^{-1}6^{1}$	$1^{3}2^{-2}3^{-2}6^{3}$	$1^{3}2^{-2}3^{-1}6^{2}$	$1^{3}2^{-1}3^{-1}6^{1}$	$1^{4}2^{-2}3^{-2}6^{2}$
$1^{4}2^{-2}3^{-1}6^{1}$					
N = 6, k =	3/2				
$1^{-3}2^{4}3^{4}6^{-2}$	$1^{-3}2^{4}3^{5}6^{-3}$	$1^{-3}2^{5}3^{2}6^{-1}$	$1^{-3}2^{5}3^{3}6^{-2}$	$1^{-3}2^{5}3^{4}6^{-3}$	$1^{-2}2^{2}3^{3}6^{0}$
$1^{-2}2^{2}3^{4}6^{-1}$	$1^{-2}2^{2}3^{5}6^{-2}$	$1^{-2}2^{3}3^{1}6^{1}$	$1^{-2}2^{3}3^{2}6^{0}$	$1^{-2}2^{3}3^{3}6^{-1}$	$1^{-2}2^{3}3^{4}6^{-2}$
$1^{-2}2^{3}3^{5}6^{-3}$	$1^{-2}2^{4}3^{-1}6^{2}$	$1^{-2}2^{4}3^{0}6^{1}$	$1^{-2}2^{4}3^{1}6^{0}$	$1^{-2}2^{4}3^{2}6^{-1}$	$1^{-2}2^{4}3^{3}6^{-2}$
$1^{-2}2^{4}3^{4}6^{-3}$	$1^{-2}2^{5}3^{1}6^{-1}$	$1^{-2}2^{5}3^{2}6^{-2}$	$1^{-1}2^{0}3^{2}6^{2}$	$1^{-1}2^{0}3^{3}6^{1}$	$1^{-1}2^{1}3^{0}6^{3}$
$1^{-1}2^{1}3^{1}6^{2}$	$1^{-1}2^{1}3^{2}6^{1}$	$1^{-1}2^{1}3^{3}6^{0}$	$1^{-1}2^{1}3^{4}6^{-1}$	$1^{-1}2^{1}3^{5}6^{-2}$	$1^{-1}2^{2}3^{-2}6^{4}$
$1^{-1}2^{2}3^{-1}6^{3}$	$1^{-1}2^{2}3^{0}6^{2}$	$1^{-1}2^{2}3^{1}6^{1}$	$1^{-1}2^{2}3^{2}6^{0}$	$1^{-1}2^{2}3^{3}6^{-1}$	$1^{-1}2^{2}3^{4}6^{-2}$
$1^{-1}2^{2}3^{5}6^{-3}$	$1^{-1}2^{3}3^{-1}6^{2}$	$1^{-1}2^{3}3^{0}6^{1}$	$1^{-1}2^{3}3^{1}6^{0}$	$1^{-1}2^{3}3^{2}6^{-1}$	$1^{-1}2^{3}3^{3}6^{-2}$
$1^{-1}2^{4}3^{1}6^{-1}$	$1^{-1}2^{4}3^{2}6^{-2}$	$1^{0}2^{-1}3^{0}6^{4}$	$1^{0}2^{-1}3^{1}6^{3}$	$1^{0}2^{-1}3^{2}6^{2}$	$1^{0}2^{0}3^{-1}6^{4}$
$1^{0}2^{0}3^{0}6^{3}$	$1^{0}2^{0}3^{1}6^{2}$	$1^{0}2^{0}3^{2}6^{1}$	$1^{0}2^{0}3^{4}6^{-1}$	$1^{0}2^{1}3^{-2}6^{4}$	$1^{0}2^{1}3^{-1}6^{3}$
$1^{0}2^{1}3^{0}6^{2}$	$1^{0}2^{1}3^{1}6^{1}$	$1^{0}2^{1}3^{2}6^{0}$	$1^{0}2^{1}3^{3}6^{-1}$	$1^{0}2^{1}3^{4}6^{-2}$	$1^{0}2^{2}3^{-1}6^{2}$
$1^{0}2^{2}3^{0}6^{1}$	$1^{0}2^{2}3^{1}6^{0}$	$1^{0}2^{2}3^{2}6^{-1}$	$1^{0}2^{2}3^{3}6^{-2}$	$1^{0}2^{3}3^{-1}6^{1}$	$1^{0}2^{3}3^{1}6^{-1}$
$1^{0}2^{3}3^{2}6^{-2}$	$1^{0}2^{4}3^{0}6^{-1}$	$1^{1}2^{-1}3^{-2}6^{5}$	$1^{1}2^{-1}3^{-1}6^{4}$	$1^{1}2^{-1}3^{0}6^{3}$	$1^{1}2^{-1}3^{1}6^{2}$
$1^{1}2^{-1}3^{2}6^{1}$	$1^{1}2^{-1}3^{3}6^{0}$	$1^{1}2^{0}3^{-2}6^{4}$	$1^{1}2^{0}3^{-1}6^{3}$	$1^{1}2^{0}3^{0}6^{2}$	$1^{1}2^{0}3^{1}6^{1}$
$1^{1}2^{0}3^{3}6^{-1}$	$1^{1}2^{0}3^{4}6^{-2}$	$1^{1}2^{1}3^{-2}6^{3}$	$1^{1}2^{1}3^{-1}6^{2}$	$1^{1}2^{1}3^{0}6^{1}$	$1^{1}2^{1}3^{1}6^{0}$
$1^{1}2^{1}3^{2}6^{-1}$	$1^{1}2^{1}3^{3}6^{-2}$	$1^{1}2^{2}3^{-1}6^{1}$	$1^{1}2^{2}3^{1}6^{-1}$	$1^{1}2^{3}3^{0}6^{-1}$	$1^2 2^{-2} 3^{-2} 6^5$
$1^2 2^{-2} 3^{-1} 6^4$	$1^2 2^{-2} 3^0 6^3$	$1^2 2^{-1} 3^{-3} 6^5$	$1^2 2^{-1} 3^{-2} 6^4$	$1^2 2^{-1} 3^{-1} 6^3$	$1^2 2^{-1} 3^0 6^2$
$1^2 2^{-1} 3^1 6^1$	$1^2 2^{-1} 3^2 6^0$	$1^2 2^{-1} 3^3 6^{-1}$	$1^2 2^{-1} 3^4 6^{-2}$	$1^{2}2^{0}3^{-2}6^{3}$	$1^{2}2^{0}3^{-1}6^{2}$
$1^2 2^0 3^0 6^1$	$1^{2}2^{0}3^{2}6^{-1}$	$1^{2}2^{1}3^{-1}6^{1}$	$1^{2}2^{1}3^{1}6^{-1}$	$1^2 2^2 3^{-1} 6^0$	$1^{2}2^{2}3^{0}6^{-1}$
$1^{3}2^{-2}3^{-3}6^{5}$	$1^{3}2^{-2}3^{-2}6^{4}$	$1^{3}2^{-2}3^{-1}6^{3}$	$1^{3}2^{-2}3^{0}6^{2}$	$1^{3}2^{-2}3^{1}6^{1}$	$1^{3}2^{-1}3^{-2}6^{3}$
$1^{3}2^{-1}3^{-1}6^{2}$	$1^{3}2^{-1}3^{0}6^{1}$	$1^{3}2^{-1}3^{1}6^{0}$	$1^{3}2^{-1}3^{2}6^{-1}$	$1^{3}2^{0}3^{-2}6^{2}$	$1^{3}2^{0}3^{-1}6^{1}$
$1^{3}2^{0}3^{1}6^{-1}$	$1^{3}2^{1}3^{-1}6^{0}$	$1^{4}2^{-3}3^{-3}6^{5}$	$1^{4}2^{-3}3^{-2}6^{4}$	$1^{4}2^{-2}3^{-3}6^{4}$	$1^{4}2^{-2}3^{-2}6^{3}$
$1^{4}2^{-2}3^{-1}6^{2}$	$1^{4}2^{-2}3^{0}6^{1}$	$1^{4}2^{-2}3^{1}6^{0}$	$1^{4}2^{-2}3^{2}6^{-1}$	$1^{4}2^{-1}3^{-2}6^{2}$	$1^{4}2^{-1}3^{-1}6^{1}$
$1^5 2^{-3} 3^{-3} 6^4$	$1^{5}2^{-3}3^{-2}6^{3}$	$1^5 2^{-3} 3^{-1} 6^2$	$1^{5}2^{-2}3^{-2}6^{2}$	$1^{5}2^{-2}3^{-1}6^{1}$	
N = 6, k =	2				
$1^{-4}2^{6}3^{5}6^{-3}$	$1^{-4}2^{6}3^{6}6^{-4}$	$1^{-3}2^{4}3^{3}6^{0}$	$1^{-3}2^{4}3^{4}6^{-1}$	$1^{-3}2^{4}3^{5}6^{-2}$	$1^{-3}2^{5}3^{2}6^{0}$
$1^{-3}2^{5}3^{3}6^{-1}$	$1^{-3}2^{5}3^{4}6^{-2}$	$1^{-3}2^{5}3^{5}6^{-3}$	$1^{-3}2^{5}3^{6}6^{-4}$	$1^{-2}2^{2}3^{2}6^{2}$	$1^{-2}2^{2}3^{3}6^{1}$
$1^{-2}2^{2}3^{4}6^{0}$	$1^{-2}2^{3}3^{0}6^{3}$	$1^{-2}2^{3}3^{1}6^{2}$	$1^{-2}2^{3}3^{2}6^{1}$	$1^{-2}2^{3}3^{3}6^{0}$	$1^{-2}2^{3}3^{4}6^{-1}$
$1^{-2}2^{3}3^{5}6^{-2}$	$1^{-2}2^{4}3^{-1}6^{3}$	$1^{-2}2^{4}3^{0}6^{2}$	$1^{-2}2^{4}3^{1}6^{1}$	$1^{-2}2^{4}3^{2}6^{0}$	$1^{-2}2^{4}3^{3}6^{-1}$
$1^{-2}2^{4}3^{4}6^{-2}$	$1^{-2}2^{4}3^{5}6^{-3}$	$1^{-2}2^{5}3^{3}6^{-2}$	$1^{-2}2^{5}3^{4}6^{-3}$	$1^{-1}2^{0}3^{2}6^{3}$	$1^{-1}2^{0}3^{3}6^{2}$
$1^{-1}2^{1}3^{1}6^{3}$	$1^{-1}2^{1}3^{2}6^{2}$	$1^{-1}2^{1}3^{3}6^{1}$	$1^{-1}2^{1}3^{4}6^{0}$	$1^{-1}2^{2}3^{0}6^{3}$	$1^{-1}2^{2}3^{1}6^{2}$
$1^{-1}2^{2}3^{2}6^{1}$	$1^{-1}2^{2}3^{3}6^{0}$	$1^{-1}2^{2}3^{4}6^{-1}$	$1^{-1}2^{3}3^{-2}6^{4}$	$1^{-1}2^{3}3^{-1}6^{3}$	$1^{-1}2^{3}3^{0}6^{2}$
$1^{-1}2^{3}3^{1}6^{1}$	$1^{-1}2^{3}3^{2}6^{0}$	$1^{-1}2^{3}3^{3}6^{-1}$	$1^{-1}2^{3}3^{4}6^{-2}$	$1^{-1}2^{3}3^{5}6^{-3}$	$1^{-1}2^{4}3^{0}6^{1}$
$1^{-1}2^{4}3^{1}6^{0}$	$1^{-1}2^{4}3^{2}6^{-1}$	$1^{-1}2^{4}3^{3}6^{-2}$	$1^{-1}2^{4}3^{4}6^{-3}$	$1^{0}2^{-1}3^{2}6^{3}$	$1^{0}2^{-1}3^{3}6^{2}$
$1^{0}2^{0}3^{1}6^{3}$	$1^{0}2^{0}3^{2}6^{2}$	$1^{0}2^{0}3^{3}6^{1}$	$1^{0}2^{1}3^{-1}6^{4}$	$1^{0}2^{1}3^{0}6^{3}$	$1^{0}2^{1}3^{1}6^{2}$

Table 3. (continued)

To be continued

	nemaca)				
$1^{0}2^{1}3^{2}6^{1}$	$1^{0}2^{1}3^{3}6^{0}$	$1^{0}2^{1}3^{4}6^{-1}$	$1^{0}2^{2}3^{-2}6^{4}$	$1^{0}2^{2}3^{-1}6^{3}$	$1^{0}2^{2}3^{0}6^{2}$
$1^{0}2^{2}3^{1}6^{1}$	$1^{0}2^{2}3^{2}6^{0}$	$1^{0}2^{2}3^{3}6^{-1}$	$1^{0}2^{2}3^{4}6^{-2}$	$1^{0}2^{2}3^{5}6^{-3}$	$1^{0}2^{3}3^{-2}6^{3}$
$1^{0}2^{3}3^{-1}6^{2}$	$1^{0}2^{3}3^{0}6^{1}$	$1^{0}2^{3}3^{1}6^{0}$	$1^{0}2^{3}3^{2}6^{-1}$	$1^{0}2^{3}3^{3}6^{-2}$	$1^{0}2^{3}3^{4}6^{-3}$
$1^{0}2^{4}3^{1}6^{-1}$	$1^{0}2^{4}3^{2}6^{-2}$	$1^{1}2^{-1}3^{0}6^{4}$	$1^{1}2^{-1}3^{1}6^{3}$	$1^{1}2^{-1}3^{2}6^{2}$	$1^{1}2^{-1}3^{3}6^{1}$
$1^{1}2^{0}3^{-1}6^{4}$	$1^1 2^0 3^0 6^3$	$1^1 2^0 3^1 6^2$	$1^{1}2^{0}3^{2}6^{1}$	$1^{1}2^{0}3^{4}6^{-1}$	$1^{1}2^{1}3^{-2}6^{4}$
$1^{1}2^{1}3^{-1}6^{3}$	$1^{1}2^{1}3^{0}6^{2}$	$1^{1}2^{1}3^{1}6^{1}$	$1^{1}2^{1}3^{2}6^{0}$	$1^{1}2^{1}3^{3}6^{-1}$	$1^{1}2^{1}3^{4}6^{-2}$
$1^{1}2^{2}3^{-2}6^{3}$	$1^{1}2^{2}3^{-1}6^{2}$	$1^{1}2^{2}3^{0}6^{1}$	$1^1 2^2 3^1 6^0$	$1^{1}2^{2}3^{2}6^{-1}$	$1^{1}2^{2}3^{3}6^{-2}$
$1^{1}2^{3}3^{-1}6^{1}$	$1^{1}2^{3}3^{1}6^{-1}$	$1^{1}2^{3}3^{2}6^{-2}$	$1^2 2^{-2} 3^0 6^4$	$1^2 2^{-2} 3^1 6^3$	$1^2 2^{-2} 3^2 6^2$
$1^2 2^{-1} 3^{-1} 6^4$	$1^2 2^{-1} 3^0 6^3$	$1^2 2^{-1} 3^1 6^2$	$1^2 2^{-1} 3^2 6^1$	$1^2 2^{-1} 3^3 6^0$	$1^2 2^0 3^{-3} 6^5$
$1^2 2^0 3^{-2} 6^4$	$1^{2}2^{0}3^{-1}6^{3}$	$1^2 2^0 3^0 6^2$	$1^2 2^0 3^1 6^1$	$1^2 2^0 3^3 6^{-1}$	$1^{2}2^{0}3^{4}6^{-2}$
$1^{2}2^{1}3^{-2}6^{3}$	$1^{2}2^{1}3^{-1}6^{2}$	$1^2 2^1 3^0 6^1$	$1^2 2^1 3^1 6^0$	$1^{2}2^{1}3^{2}6^{-1}$	$1^{2}2^{1}3^{3}6^{-2}$
$1^2 2^2 3^{-2} 6^2$	$1^{2}2^{2}3^{-1}6^{1}$	$1^2 2^2 3^1 6^{-1}$	$1^2 2^2 3^2 6^{-2}$	$1^2 2^3 3^{-1} 6^0$	$1^{2}2^{3}3^{0}6^{-1}$
$1^{3}2^{-2}3^{-2}6^{5}$	$1^{3}2^{-2}3^{-1}6^{4}$	$1^{3}2^{-2}3^{0}6^{3}$	$1^{3}2^{-2}3^{1}6^{2}$	$1^{3}2^{-2}3^{2}6^{1}$	$1^{3}2^{-2}3^{3}6^{0}$
$1^{3}2^{-1}3^{-3}6^{5}$	$1^{3}2^{-1}3^{-2}6^{4}$	$1^{3}2^{-1}3^{-1}6^{3}$	$1^{3}2^{-1}3^{0}6^{2}$	$1^{3}2^{-1}3^{1}6^{1}$	$1^{3}2^{-1}3^{2}6^{0}$
$1^{3}2^{-1}3^{3}6^{-1}$	$1^{3}2^{-1}3^{4}6^{-2}$	$1^{3}2^{0}3^{-3}6^{4}$	$1^{3}2^{0}3^{-2}6^{3}$	$1^{3}2^{0}3^{-1}6^{2}$	$1^3 2^0 3^0 6^1$
$1^{3}2^{0}3^{2}6^{-1}$	$1^{3}2^{0}3^{3}6^{-2}$	$1^{3}2^{1}3^{-2}6^{2}$	$1^{3}2^{1}3^{-1}6^{1}$	$1^{3}2^{1}3^{1}6^{-1}$	$1^{3}2^{2}3^{-1}6^{0}$
$1^{3}2^{2}3^{0}6^{-1}$	$1^4 2^{-3} 3^{-2} 6^5$	$1^{4}2^{-3}3^{-1}6^{4}$	$1^{4}2^{-3}3^{0}6^{3}$	$1^{4}2^{-2}3^{-3}6^{5}$	$1^4 2^{-2} 3^{-2} 6^4$
$1^4 2^{-2} 3^{-1} 6^3$	$1^4 2^{-2} 3^0 6^2$	$1^{4}2^{-2}3^{1}6^{1}$	$1^{4}2^{-2}3^{2}6^{0}$	$1^{4}2^{-2}3^{3}6^{-1}$	$1^4 2^{-1} 3^{-3} 6^4$
$1^4 2^{-1} 3^{-2} 6^3$	$1^4 2^{-1} 3^{-1} 6^2$	$1^{4}2^{-1}3^{0}6^{1}$	$1^4 2^{-1} 3^1 6^0$	$1^4 2^0 3^{-2} 6^2$	$1^4 2^0 3^{-1} 6^1$
$1^5 2^{-3} 3^{-4} 6^6$	$1^5 2^{-3} 3^{-3} 6^5$	$1^{5}2^{-3}3^{-2}6^{4}$	$1^{5}2^{-3}3^{-1}6^{3}$	$1^{5}2^{-3}3^{0}6^{2}$	$1^{5}2^{-2}3^{-3}6^{4}$
$1^{5}2^{-2}3^{-2}6^{3}$	$1^{6}2^{-4}3^{-4}6^{6}$	$1^{6}2^{-4}3^{-3}6^{5}$			
N = 6, k =	5/2				
$1^{-3}2^{4}3^{2}6^{2}$	$1^{-3}2^{4}3^{3}6^{1}$	$1^{-3}2^{4}3^{4}6^{0}$	$1^{-3}2^{5}3^{4}6^{-1}$	$1^{-3}2^{5}3^{5}6^{-2}$	$1^{-2}2^{2}3^{3}6^{2}$
1-2030202					
1 236	$1^{-2}2^{3}3^{3}6^{1}$	$1^{-2}2^{3}3^{4}6^{0}$	$1^{-2}2^{4}3^{1}6^{2}$	$1^{-2}2^{4}3^{2}6^{1}$	$1^{-2}2^{4}3^{3}6^{0}$
$1^{-2}2^{3}3^{4}6^{-1}$ $1^{-2}2^{4}3^{4}6^{-1}$	$\frac{1^{-2}2^33^36^1}{1^{-2}2^53^56^{-3}}$	$\frac{1^{-2}2^3 3^4 6^0}{1^{-1}2^1 3^3 6^2}$	$\frac{1^{-2}2^4 3^1 6^2}{1^{-1}2^2 3^2 6^2}$	$\frac{1^{-2}2^4 3^2 6^1}{1^{-1}2^2 3^3 6^1}$	$1^{-2}2^43^36^0$ $1^{-1}2^33^06^3$
$ \begin{array}{c} 1 & 2^{3}3^{2}6^{2} \\ 1^{-2}2^{4}3^{4}6^{-1} \\ 1^{-1}2^{3}3^{1}6^{2} \end{array} $	$ 1^{-2}2^{3}3^{3}6^{1} \\ 1^{-2}2^{5}3^{5}6^{-3} \\ 1^{-1}2^{3}3^{2}6^{1} $	$ \begin{array}{r}1^{-2}2^{3}3^{4}6^{0}\\1^{-1}2^{1}3^{3}6^{2}\\1^{-1}2^{3}3^{3}6^{0}\end{array} $	$ \begin{array}{r} 1^{-2} 2^4 3^1 6^2 \\ 1^{-1} 2^2 3^2 6^2 \\ 1^{-1} 2^3 3^4 6^{-1} \end{array} $	$ 1^{-2}2^{4}3^{2}6^{1} \\ 1^{-1}2^{2}3^{3}6^{1} \\ 1^{-1}2^{4}3^{2}6^{0} $	$ 1^{-2}2^{4}3^{3}6^{0} \\ 1^{-1}2^{3}3^{0}6^{3} \\ 1^{-1}2^{4}3^{3}6^{-1} $
$ \begin{array}{c} 1 & 2^{2} 3^{2} 6^{2} \\ 1^{-2} 2^{4} 3^{4} 6^{-1} \\ 1^{-1} 2^{3} 3^{1} 6^{2} \\ 1^{-1} 2^{4} 3^{4} 6^{-2} \end{array} $	$ \begin{array}{r} 1^{-2}2^{3}3^{3}6^{1} \\ 1^{-2}2^{5}3^{5}6^{-3} \\ 1^{-1}2^{3}3^{2}6^{1} \\ 1^{-1}2^{4}3^{5}6^{-3} \end{array} $	$\begin{array}{c} 1^{-2}2^{3}3^{4}6^{0} \\ 1^{-1}2^{1}3^{3}6^{2} \\ 1^{-1}2^{3}3^{3}6^{0} \\ 1^{0}2^{1}3^{1}6^{3} \end{array}$	$\frac{1^{-2}2^{4}3^{1}6^{2}}{1^{-1}2^{2}3^{2}6^{2}}\\\frac{1^{-1}2^{3}3^{4}6^{-1}}{1^{0}2^{1}3^{2}6^{2}}$	$\begin{array}{c} 1^{-2}2^{4}3^{2}6^{1} \\ 1^{-1}2^{2}3^{3}6^{1} \\ 1^{-1}2^{4}3^{2}6^{0} \\ 1^{0}2^{1}3^{3}6^{1} \end{array}$	$ \begin{array}{r}1^{-2}2^{4}3^{3}6^{0}\\1^{-1}2^{3}3^{0}6^{3}\\1^{-1}2^{4}3^{3}6^{-1}\\1^{0}2^{2}3^{0}6^{3}\end{array} $
$1^{-2}2^{3}3^{6}6^{2}$ $1^{-2}2^{4}3^{4}6^{-1}$ $1^{-1}2^{3}3^{1}6^{2}$ $1^{-1}2^{4}3^{4}6^{-2}$ $1^{0}2^{2}3^{1}6^{2}$	$1^{-2}2^{3}3^{3}6^{1}$ $1^{-2}2^{5}3^{5}6^{-3}$ $1^{-1}2^{3}3^{2}6^{1}$ $1^{-1}2^{4}3^{5}6^{-3}$ $1^{0}2^{2}3^{2}6^{1}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{0}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2} \end{array}$
$ \begin{array}{c} 1 & 2^{2} 3^{2} 6^{2} \\ 1^{-2} 2^{4} 3^{4} 6^{-1} \\ 1^{-1} 2^{3} 3^{1} 6^{2} \\ 1^{-1} 2^{4} 3^{4} 6^{-2} \\ 1^{0} 2^{2} 3^{1} 6^{2} \\ 1^{0} 2^{3} 3^{1} 6^{1} \end{array} $	$1^{-2}2^{3}3^{3}6^{1}$ $1^{-2}2^{5}3^{5}6^{-3}$ $1^{-1}2^{3}3^{2}6^{1}$ $1^{-1}2^{4}3^{5}6^{-3}$ $1^{0}2^{2}3^{2}6^{1}$ $1^{0}2^{3}3^{2}6^{0}$	$\begin{array}{c} 1^{-2}2^33^46^0\\ 1^{-1}2^13^36^2\\ 1^{-1}2^33^36^0\\ 1^02^13^16^3\\ 1^02^23^36^0\\ 1^02^33^36^{-1}\end{array}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3} \end{array}$
$1^{-2}2^{3}3^{6}6^{2}$ $1^{-2}2^{4}3^{4}6^{-1}$ $1^{-1}2^{3}3^{1}6^{2}$ $1^{-1}2^{4}3^{4}6^{-2}$ $1^{0}2^{2}3^{1}6^{2}$ $1^{0}2^{3}3^{1}6^{1}$ $1^{1}2^{-1}3^{2}6^{3}$	$1^{-2}2^{3}3^{3}6^{1}$ $1^{-2}2^{5}3^{5}6^{-3}$ $1^{-1}2^{3}3^{2}6^{1}$ $1^{-1}2^{4}3^{5}6^{-3}$ $1^{0}2^{2}3^{2}6^{1}$ $1^{0}2^{3}3^{2}6^{0}$ $1^{1}2^{0}3^{1}6^{3}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{0}$ $1^{0}2^{3}3^{3}6^{-1}$ $1^{1}2^{0}3^{2}6^{2}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2} \end{array}$
$1^{-2}2^{+3}4^{-1}$ $1^{-1}2^{3}3^{1}6^{2}$ $1^{-1}2^{4}3^{4}6^{-2}$ $1^{0}2^{2}3^{1}6^{2}$ $1^{0}2^{3}3^{1}6^{1}$ $1^{1}2^{-1}3^{2}6^{3}$ $1^{1}2^{1}3^{2}6^{1}$	$1^{-2}2^{3}3^{3}6^{1}$ $1^{-2}2^{5}3^{5}6^{-3}$ $1^{-1}2^{3}3^{2}6^{1}$ $1^{-1}2^{4}3^{5}6^{-3}$ $1^{0}2^{2}3^{2}6^{1}$ $1^{0}2^{3}3^{2}6^{0}$ $1^{1}2^{0}3^{1}6^{3}$ $1^{1}2^{1}3^{3}6^{0}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{0}$ $1^{0}2^{3}3^{3}6^{-1}$ $1^{1}2^{0}3^{2}6^{2}$ $1^{1}2^{2}3^{-2}6^{4}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1} \end{array}$
$1^{-2}2^{+3}4^{-1}$ $1^{-1}2^{3}3^{1}6^{2}$ $1^{-1}2^{4}3^{4}6^{-2}$ $1^{0}2^{2}3^{1}6^{2}$ $1^{0}2^{3}3^{1}6^{1}$ $1^{1}2^{-1}3^{2}6^{3}$ $1^{1}2^{1}3^{2}6^{1}$ $1^{1}2^{2}3^{2}6^{0}$	$\begin{array}{c} 1^{-2}2^33^36^1\\ 1^{-2}2^53^56^{-3}\\ 1^{-1}2^33^26^1\\ 1^{-1}2^43^56^{-3}\\ 1^{0}2^23^26^1\\ 1^{0}2^33^26^0\\ 1^{1}2^03^{1}6^3\\ 1^{1}2^{1}3^{3}6^0\\ 1^{1}2^23^{3}6^{-1} \end{array}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{0}$ $1^{0}2^{3}3^{3}6^{-1}$ $1^{1}2^{0}3^{2}6^{2}$ $1^{1}2^{2}3^{-2}6^{4}$ $1^{1}2^{2}3^{4}6^{-2}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{1}2^{3}3^{0}6^{1}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{1}2^{3}3^{1}6^{0}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1} \end{array}$
$1^{-2}2^{-3}2^{-6}2^{-1}$ $1^{-2}2^{4}3^{4}6^{-1}$ $1^{-1}2^{3}3^{1}6^{2}$ $1^{-1}2^{4}3^{4}6^{-2}$ $1^{0}2^{2}3^{1}6^{2}$ $1^{0}2^{3}3^{1}6^{1}$ $1^{1}2^{-1}3^{2}6^{3}$ $1^{1}2^{1}3^{2}6^{1}$ $1^{1}2^{2}3^{2}6^{0}$ $1^{1}2^{3}3^{3}6^{-2}$	$1^{-2}2^{3}3^{3}6^{1}$ $1^{-2}2^{5}3^{5}6^{-3}$ $1^{-1}2^{3}3^{2}6^{1}$ $1^{-1}2^{4}3^{5}6^{-3}$ $1^{0}2^{2}3^{2}6^{1}$ $1^{0}2^{3}3^{2}6^{0}$ $1^{1}2^{0}3^{1}6^{3}$ $1^{1}2^{1}3^{3}6^{0}$ $1^{1}2^{2}3^{3}6^{-1}$ $1^{1}2^{3}3^{4}6^{-3}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{0}$ $1^{0}2^{3}3^{3}6^{-1}$ $1^{1}2^{0}3^{2}6^{2}$ $1^{1}2^{2}3^{-2}6^{4}$ $1^{1}2^{2}3^{4}6^{-2}$ $1^{2}2^{-2}3^{2}6^{3}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{1}2^{3}3^{0}6^{1}$ $1^{2}2^{-1}3^{1}6^{3}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{1}2^{3}3^{1}6^{0}$ $1^{2}2^{-1}3^{2}6^{2}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1}\\ 1^{2}2^{0}3^{-1}6^{4} \end{array}$
$1 22^{3}2^{6}2$ $1^{-2}2^{4}3^{4}6^{-1}$ $1^{-1}2^{3}3^{1}6^{2}$ $1^{-1}2^{4}3^{4}6^{-2}$ $1^{0}2^{2}3^{1}6^{2}$ $1^{0}2^{3}3^{1}6^{1}$ $1^{1}2^{-1}3^{2}6^{3}$ $1^{1}2^{1}3^{2}6^{1}$ $1^{1}2^{3}3^{3}6^{-2}$ $1^{2}2^{0}3^{0}6^{3}$	$1^{-2}2^{3}3^{3}6^{1}$ $1^{-2}2^{5}3^{5}6^{-3}$ $1^{-1}2^{3}3^{2}6^{1}$ $1^{-1}2^{4}3^{5}6^{-3}$ $1^{0}2^{2}3^{2}6^{1}$ $1^{0}2^{3}3^{2}6^{0}$ $1^{1}2^{0}3^{1}6^{3}$ $1^{1}2^{1}3^{3}6^{0}$ $1^{1}2^{2}3^{3}6^{-1}$ $1^{1}2^{3}3^{4}6^{-3}$ $1^{2}2^{0}3^{1}6^{2}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{0}$ $1^{0}2^{3}3^{3}6^{-1}$ $1^{1}2^{0}3^{2}6^{2}$ $1^{1}2^{2}3^{-2}6^{4}$ $1^{1}2^{2}3^{4}6^{-2}$ $1^{2}2^{-2}3^{2}6^{3}$ $1^{2}2^{0}3^{2}6^{1}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{1}2^{3}3^{0}6^{1}$ $1^{2}2^{-1}3^{1}6^{3}$ $1^{2}2^{1}3^{-2}6^{4}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{1}2^{3}3^{1}6^{0}$ $1^{2}2^{-1}3^{2}6^{2}$ $1^{2}2^{1}3^{-1}6^{3}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1}\\ 1^{2}2^{0}3^{-1}6^{4}\\ 1^{2}2^{1}3^{0}6^{2} \end{array}$
$1^{-2}2^{-3}2^{-6}2^{-1}$ $1^{-1}2^{3}3^{1}6^{2}$ $1^{-1}2^{4}3^{4}6^{-2}$ $1^{0}2^{2}3^{1}6^{2}$ $1^{0}2^{3}3^{1}6^{1}$ $1^{1}2^{-1}3^{2}6^{3}$ $1^{1}2^{1}3^{2}6^{1}$ $1^{1}2^{2}3^{2}6^{0}$ $1^{1}2^{3}3^{3}6^{-2}$ $1^{2}2^{0}3^{0}6^{3}$ $1^{2}2^{1}3^{1}6^{1}$	$\begin{array}{c} 1^{-2}2^33^36^1\\ 1^{-2}2^53^56^{-3}\\ 1^{-1}2^33^26^1\\ 1^{-1}2^43^56^{-3}\\ 1^{0}2^23^26^1\\ 1^{0}2^33^26^0\\ 1^{1}2^03^{1}6^3\\ 1^{1}2^{1}3^{3}6^0\\ 1^{1}2^23^36^{-1}\\ 1^{1}2^33^46^{-3}\\ 1^{2}2^{0}3^{1}6^2\\ 1^{2}2^{1}3^{2}6^0\end{array}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{-1}$ $1^{0}2^{3}3^{3}6^{-1}$ $1^{1}2^{0}3^{2}6^{2}$ $1^{1}2^{2}3^{-2}6^{4}$ $1^{1}2^{2}3^{4}6^{-2}$ $1^{2}2^{-2}3^{2}6^{3}$ $1^{2}2^{0}3^{2}6^{1}$ $1^{2}2^{1}3^{3}6^{-1}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{1}2^{3}3^{0}6^{1}$ $1^{2}2^{-1}3^{1}6^{3}$ $1^{2}2^{1}3^{-2}6^{4}$ $1^{2}2^{1}3^{4}6^{-2}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{1}2^{3}3^{1}6^{0}$ $1^{2}2^{-1}3^{2}6^{2}$ $1^{2}2^{1}3^{-1}6^{3}$ $1^{2}2^{2}3^{-3}6^{4}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1}\\ 1^{2}2^{0}3^{-1}6^{4}\\ 1^{2}2^{1}3^{0}6^{2}\\ 1^{2}2^{2}3^{-2}6^{3} \end{array}$
$1^{2}2^{3}3^{6}6^{2}$ $1^{-2}2^{4}3^{4}6^{-1}$ $1^{-1}2^{3}3^{1}6^{2}$ $1^{-1}2^{4}3^{4}6^{-2}$ $1^{0}2^{3}3^{1}6^{1}$ $1^{1}2^{-1}3^{2}6^{3}$ $1^{1}2^{1}3^{2}6^{1}$ $1^{1}2^{2}3^{2}6^{0}$ $1^{1}2^{3}3^{3}6^{-2}$ $1^{2}2^{0}3^{0}6^{3}$ $1^{2}2^{1}3^{1}6^{1}$ $1^{2}2^{2}3^{-1}6^{2}$	$1^{-2}2^{3}3^{3}6^{1}$ $1^{-2}2^{5}3^{5}6^{-3}$ $1^{-1}2^{3}3^{2}6^{1}$ $1^{-1}2^{4}3^{5}6^{-3}$ $1^{0}2^{2}3^{2}6^{1}$ $1^{0}2^{3}3^{2}6^{0}$ $1^{1}2^{0}3^{1}6^{3}$ $1^{1}2^{1}3^{3}6^{0}$ $1^{1}2^{2}3^{3}6^{-1}$ $1^{1}2^{3}3^{4}6^{-3}$ $1^{2}2^{0}3^{1}6^{2}$ $1^{2}2^{1}3^{2}6^{0}$ $1^{2}2^{2}3^{0}6^{1}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{-1}$ $1^{0}2^{3}3^{3}6^{-1}$ $1^{1}2^{0}3^{2}6^{2}$ $1^{1}2^{2}3^{-2}6^{4}$ $1^{1}2^{2}3^{-4}6^{-2}$ $1^{2}2^{-2}3^{2}6^{3}$ $1^{2}2^{0}3^{2}6^{1}$ $1^{2}2^{1}3^{3}6^{-1}$ $1^{2}2^{2}3^{1}6^{0}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{2}2^{-1}3^{1}6^{3}$ $1^{2}2^{1}3^{-2}6^{4}$ $1^{2}2^{1}3^{4}6^{-2}$ $1^{2}2^{2}3^{2}6^{-1}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{1}2^{3}3^{1}6^{0}$ $1^{2}2^{-1}3^{2}6^{2}$ $1^{2}2^{1}3^{-1}6^{3}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{3}6^{-2}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1}\\ 1^{2}2^{0}3^{-1}6^{4}\\ 1^{2}2^{1}3^{0}6^{2}\\ 1^{2}2^{2}3^{-2}6^{3}\\ 1^{2}2^{2}3^{4}6^{-3}\end{array}$
$1^{2}2^{3}3^{6}6^{2}$ $1^{-2}2^{4}3^{4}6^{-1}$ $1^{-1}2^{3}3^{1}6^{2}$ $1^{-1}2^{4}3^{4}6^{-2}$ $1^{0}2^{3}3^{1}6^{1}$ $1^{1}2^{-1}3^{2}6^{3}$ $1^{1}2^{1}3^{2}6^{1}$ $1^{1}2^{2}3^{2}6^{0}$ $1^{1}2^{3}3^{3}6^{-2}$ $1^{2}2^{0}3^{0}6^{3}$ $1^{2}2^{1}3^{1}6^{1}$ $1^{2}2^{2}3^{-1}6^{2}$ $1^{2}2^{3}3^{1}6^{-1}$	$\begin{array}{c} 1^{-2}2^33^36^1\\ 1^{-2}2^53^56^{-3}\\ 1^{-1}2^33^26^1\\ 1^{-1}2^43^56^{-3}\\ 1^{0}2^23^26^1\\ 1^{0}2^33^26^0\\ 1^{1}2^{0}3^{1}6^3\\ 1^{1}2^{1}3^{3}6^0\\ 1^{1}2^23^36^{-1}\\ 1^{1}2^33^46^{-3}\\ 1^{2}2^{0}3^{1}6^2\\ 1^{2}2^{1}3^{2}6^0\\ 1^{2}2^{2}3^{0}6^1\\ 1^{2}2^{3}3^{2}6^{-2} \end{array}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{-1}$ $1^{0}2^{3}3^{3}6^{-1}$ $1^{1}2^{0}3^{2}6^{2}$ $1^{1}2^{2}3^{-2}6^{4}$ $1^{2}2^{-2}3^{2}6^{3}$ $1^{2}2^{0}3^{2}6^{1}$ $1^{2}2^{1}3^{3}6^{-1}$ $1^{2}2^{2}3^{1}6^{0}$ $1^{3}2^{-2}3^{0}6^{4}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{1}2^{3}3^{0}6^{1}$ $1^{2}2^{-1}3^{1}6^{3}$ $1^{2}2^{1}3^{-2}6^{4}$ $1^{2}2^{1}3^{4}6^{-2}$ $1^{2}2^{2}3^{2}6^{-1}$ $1^{3}2^{-2}3^{1}6^{3}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{1}2^{3}3^{1}6^{0}$ $1^{2}2^{-1}3^{2}6^{2}$ $1^{2}2^{1}3^{-1}6^{3}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{-3}3^{2}6^{2}$ $1^{3}2^{-2}3^{2}6^{2}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1}\\ 1^{2}2^{0}3^{-1}6^{4}\\ 1^{2}2^{1}3^{0}6^{2}\\ 1^{2}2^{2}3^{-2}6^{3}\\ 1^{2}2^{2}3^{4}6^{-3}\\ 1^{3}2^{-1}3^{-1}6^{4}\end{array}$
$1 22^{3}2^{6}2$ $1^{-2}2^{4}3^{4}6^{-1}$ $1^{-1}2^{3}3^{1}6^{2}$ $1^{-1}2^{4}3^{4}6^{-2}$ $1^{0}2^{2}3^{1}6^{2}$ $1^{0}2^{3}3^{1}6^{1}$ $1^{1}2^{-1}3^{2}6^{3}$ $1^{1}2^{1}3^{2}6^{1}$ $1^{1}2^{3}3^{3}6^{-2}$ $1^{2}2^{0}3^{0}6^{3}$ $1^{2}2^{1}3^{1}6^{1}$ $1^{2}2^{2}3^{-1}6^{2}$ $1^{2}2^{3}3^{1}6^{-1}$ $1^{3}2^{-1}3^{0}6^{3}$	$\begin{array}{c} 1^{-2}2^33^36^1\\ 1^{-2}2^53^56^{-3}\\ 1^{-1}2^33^26^1\\ 1^{-1}2^43^56^{-3}\\ 1^{0}2^23^26^1\\ 1^{0}2^33^26^0\\ 1^{1}2^03^{1}6^3\\ 1^{1}2^{1}3^{3}6^0\\ 1^{1}2^23^36^{-1}\\ 1^{1}2^33^46^{-3}\\ 1^{2}2^{0}3^{1}6^2\\ 1^{2}2^{1}3^{2}6^0\\ 1^{2}2^23^{0}6^1\\ 1^{2}2^{3}3^{2}6^{-2}\\ 1^{3}2^{-1}3^{1}6^2\end{array}$	$\begin{array}{c} 1^{-2}2^{3}3^{4}6^{0}\\ 1^{-1}2^{1}3^{3}6^{2}\\ 1^{-1}2^{3}3^{3}6^{0}\\ 1^{0}2^{1}3^{1}6^{3}\\ 1^{0}2^{2}3^{3}6^{0}\\ 1^{0}2^{3}3^{3}6^{-1}\\ 1^{1}2^{0}3^{2}6^{2}\\ 1^{1}2^{2}3^{-2}6^{4}\\ 1^{1}2^{2}3^{-2}6^{4}\\ 1^{2}2^{-2}3^{2}6^{3}\\ 1^{2}2^{0}3^{2}6^{1}\\ 1^{2}2^{1}3^{3}6^{-1}\\ 1^{2}2^{2}3^{1}6^{0}\\ 1^{3}2^{-2}3^{0}6^{4}\\ 1^{3}2^{-1}3^{2}6^{1} \end{array}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{1}2^{2}-13^{1}6^{3}$ $1^{2}2^{-1}3^{1}6^{3}$ $1^{2}2^{1}3^{-2}6^{4}$ $1^{2}2^{1}3^{4}6^{-2}$ $1^{2}2^{2}3^{2}6^{-1}$ $1^{3}2^{-2}3^{1}6^{3}$ $1^{3}2^{-1}3^{3}6^{0}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{2}3^{-1}6^{3}$ $1^{2}2^{-1}3^{2}6^{2}$ $1^{2}2^{1}3^{-1}6^{3}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-2}6^{2}$ $1^{3}2^{-2}3^{2}6^{2}$ $1^{3}2^{0}3^{-2}6^{4}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1}\\ 1^{2}2^{0}3^{-1}6^{4}\\ 1^{2}2^{1}3^{0}6^{2}\\ 1^{2}2^{2}3^{-2}6^{3}\\ 1^{2}2^{2}3^{4}6^{-3}\\ 1^{3}2^{-1}3^{-1}6^{4}\\ 1^{3}2^{0}3^{-1}6^{3}\\ \end{array}$
$1 \ 2^{2} 3^{2} 6^{2} \\ 1^{-2} 2^{4} 3^{4} 6^{-1} \\ 1^{-1} 2^{3} 3^{1} 6^{2} \\ 1^{-1} 2^{4} 3^{4} 6^{-2} \\ 1^{0} 2^{2} 3^{1} 6^{2} \\ 1^{0} 2^{3} 3^{1} 6^{1} \\ 1^{1} 2^{-1} 3^{2} 6^{3} \\ 1^{1} 2^{1} 3^{2} 6^{1} \\ 1^{1} 2^{2} 3^{2} 6^{0} \\ 1^{1} 2^{3} 3^{3} 6^{-2} \\ 1^{2} 2^{0} 3^{0} 6^{3} \\ 1^{2} 2^{1} 3^{1} 6^{1} \\ 1^{2} 2^{2} 3^{-1} 6^{2} \\ 1^{2} 2^{3} 3^{1} 6^{-1} \\ 1^{3} 2^{-1} 3^{0} 6^{3} \\ 1^{3} 2^{0} 3^{0} 6^{2} \\ 1^{-1} 2^{3} 3^{-1} 6^{2} \\ 1^{3} 2^{-1} 3^{0} 6^{3} \\ 1^{3} 2^{0} 3^{0} 6^{2} \\ 1^{-1} 2^{3} 3^{-1} 6^{2} \\ 1^{-1} 2^{-1} 3^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{$	$\begin{array}{c} 1^{-2}2^33^36^1\\ 1^{-2}2^53^56^{-3}\\ 1^{-1}2^33^26^1\\ 1^{-1}2^43^56^{-3}\\ 1^{0}2^23^26^1\\ 1^{0}2^33^26^0\\ 1^{1}2^03^{1}6^3\\ 1^{1}2^{1}3^{3}6^0\\ 1^{1}2^23^{3}6^{-1}\\ 1^{1}2^33^{4}6^{-3}\\ 1^{2}2^{0}3^{1}6^2\\ 1^{2}2^{1}3^{2}6^0\\ 1^{2}2^{2}3^{0}6^1\\ 1^{2}2^{3}3^{2}6^{-2}\\ 1^{3}2^{-1}3^{1}6^2\\ 1^{3}2^{0}3^{1}6^1\end{array}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{-1}$ $1^{1}2^{0}3^{2}6^{2}$ $1^{1}2^{2}3^{-2}6^{4}$ $1^{1}2^{2}3^{-2}6^{4}$ $1^{2}2^{-2}3^{2}6^{3}$ $1^{2}2^{0}3^{2}6^{1}$ $1^{2}2^{2}3^{1}6^{0}$ $1^{3}2^{-2}3^{0}6^{4}$ $1^{3}2^{-1}3^{2}6^{1}$ $1^{3}2^{0}3^{3}6^{-1}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{2}2^{-1}3^{1}6^{3}$ $1^{2}2^{1}3^{-2}6^{4}$ $1^{2}2^{2}3^{2}6^{-1}$ $1^{3}2^{-2}3^{1}6^{3}$ $1^{3}2^{-1}3^{3}6^{0}$ $1^{3}2^{1}3^{-3}6^{4}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{2}2^{-1}3^{2}6^{2}$ $1^{2}2^{1}3^{-1}6^{3}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-2}6^{2}$ $1^{3}2^{0}3^{-2}6^{4}$ $1^{3}2^{1}3^{-2}6^{3}$	$\begin{array}{c} 1^{-2}2^43^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1}\\ 1^{2}2^{0}3^{-1}6^{4}\\ 1^{2}2^{1}3^{0}6^{2}\\ 1^{2}2^{2}3^{-2}6^{3}\\ 1^{3}2^{-1}3^{-1}6^{4}\\ 1^{3}2^{0}3^{-1}6^{3}\\ 1^{3}2^{1}3^{-1}6^{2}\end{array}$
$1 \ 2^{2}3^{2}6^{2} \\ 1^{-2}2^{4}3^{4}6^{-1} \\ 1^{-1}2^{3}3^{1}6^{2} \\ 1^{-1}2^{4}3^{4}6^{-2} \\ 1^{0}2^{3}3^{1}6^{1} \\ 1^{1}2^{-1}3^{2}6^{3} \\ 1^{1}2^{1}3^{2}6^{1} \\ 1^{1}2^{2}3^{2}6^{0} \\ 1^{1}2^{3}3^{3}6^{-2} \\ 1^{2}2^{0}3^{0}6^{3} \\ 1^{2}2^{1}3^{1}6^{1} \\ 1^{2}2^{2}3^{-1}6^{2} \\ 1^{2}2^{3}3^{1}6^{-1} \\ 1^{3}2^{-1}3^{0}6^{3} \\ 1^{3}2^{1}3^{0}6^{1} \\ 1^{3}2^{1}3^{0} \\ 1^{3}2^{1}3^{0} \\ 1^{3}2^{1}3^{0} \\ 1^{3}2^{1}3^{0} \\ 1^{3}2^{1}3^{0} \\ 1^{3}2^{1}3^{0} \\ 1^{3}2^{1}3^{1}6^{1} \\ 1^{3}2^{1}3^{1}6^{$	$\begin{array}{c} 1^{-2}2^33^36^1\\ 1^{-2}2^53^56^{-3}\\ 1^{-1}2^33^26^1\\ 1^{-1}2^43^56^{-3}\\ 1^{0}2^23^26^1\\ 1^{0}2^33^26^0\\ 1^{1}2^{0}3^{1}6^3\\ 1^{1}2^{1}3^{3}6^0\\ 1^{1}2^23^36^{-1}\\ 1^{1}2^33^46^{-3}\\ 1^{2}2^{0}3^{1}6^2\\ 1^{2}2^{1}3^{2}6^0\\ 1^{2}2^{3}3^{2}6^{-2}\\ 1^{3}2^{-1}3^{1}6^2\\ 1^{3}2^{0}3^{1}6^1\\ 1^{3}2^{1}3^{1}6^0\\ \end{array}$	$\begin{array}{c} 1^{-2}2^{3}3^{4}6^{0}\\ 1^{-1}2^{1}3^{3}6^{2}\\ 1^{-1}2^{3}3^{3}6^{0}\\ 1^{0}2^{1}3^{1}6^{3}\\ 1^{0}2^{2}3^{3}6^{-1}\\ 1^{0}2^{3}3^{3}6^{-1}\\ 1^{1}2^{0}3^{2}6^{2}\\ 1^{1}2^{2}3^{-2}6^{4}\\ 1^{1}2^{2}3^{4}6^{-2}\\ 1^{2}2^{-2}3^{2}6^{3}\\ 1^{2}2^{0}3^{2}6^{1}\\ 1^{2}2^{1}3^{3}6^{-1}\\ 1^{2}2^{2}3^{1}6^{0}\\ 1^{3}2^{-1}3^{2}6^{1}\\ 1^{3}2^{0}3^{3}6^{-1}\\ 1^{3}2^{0}3^{-1}\\ 1^{3}2^{2}3^{-2}6^{2} \end{array}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{2}2^{-1}3^{1}6^{3}$ $1^{2}2^{1}3^{-2}6^{4}$ $1^{2}2^{1}3^{4}6^{-2}$ $1^{2}2^{2}3^{2}6^{-1}$ $1^{3}2^{-2}3^{1}6^{3}$ $1^{3}2^{-1}3^{3}6^{0}$ $1^{3}2^{1}3^{-3}6^{4}$ $1^{3}2^{2}3^{-1}6^{1}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{1}2^{3}3^{1}6^{0}$ $1^{2}2^{-1}3^{2}6^{2}$ $1^{2}2^{1}3^{-1}6^{3}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-2}6^{3}$ $1^{3}2^{1}3^{-2}6^{3}$ $1^{4}2^{-3}3^{0}6^{4}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1}\\ 1^{2}2^{0}3^{-1}6^{4}\\ 1^{2}2^{1}3^{0}6^{2}\\ 1^{2}2^{2}3^{-2}6^{3}\\ 1^{2}2^{2}3^{4}6^{-3}\\ 1^{3}2^{-1}3^{-1}6^{4}\\ 1^{3}2^{0}3^{-1}6^{3}\\ 1^{3}2^{1}3^{-1}6^{2}\\ 1^{4}2^{-3}3^{1}6^{3}\\ \end{array}$
$1 \ 2^{2} 3^{2} 6^{2} \\ 1^{-2} 2^{4} 3^{4} 6^{-1} \\ 1^{-1} 2^{3} 3^{1} 6^{2} \\ 1^{-1} 2^{4} 3^{4} 6^{-2} \\ 1^{0} 2^{2} 3^{1} 6^{2} \\ 1^{0} 2^{3} 3^{1} 6^{1} \\ 1^{1} 2^{-1} 3^{2} 6^{3} \\ 1^{1} 2^{1} 3^{2} 6^{1} \\ 1^{1} 2^{2} 3^{2} 6^{0} \\ 1^{1} 2^{3} 3^{3} 6^{-2} \\ 1^{2} 2^{0} 3^{0} 6^{3} \\ 1^{2} 2^{1} 3^{1} 6^{1} \\ 1^{2} 2^{2} 3^{-1} 6^{2} \\ 1^{2} 2^{3} 3^{1} 6^{-1} \\ 1^{3} 2^{-1} 3^{0} 6^{3} \\ 1^{3} 2^{0} 3^{0} 6^{2} \\ 1^{3} 2^{1} 3^{0} 6^{1} \\ 1^{4} 2^{-3} 3^{2} 6^{2} \\ 1^{-1} 2^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} 3^{-1} 3^{-1} \\ 1^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-1} \\ 1^{-1} 3^{-$	$\begin{array}{c} 1^{-2}2^33^36^1\\ 1^{-2}2^53^56^{-3}\\ 1^{-1}2^33^26^1\\ 1^{-1}2^43^56^{-3}\\ 1^{0}2^23^26^1\\ 1^{0}2^33^26^0\\ 1^{1}2^03^{1}6^3\\ 1^{1}2^{1}3^{3}6^0\\ 1^{1}2^23^36^{-1}\\ 1^{1}2^33^46^{-3}\\ 1^{2}2^{0}3^{1}6^2\\ 1^{2}2^{1}3^{2}6^0\\ 1^{2}2^{2}3^{0}6^1\\ 1^{2}2^{3}3^{2}6^{-2}\\ 1^{3}2^{-1}3^{1}6^2\\ 1^{3}2^{0}3^{1}6^1\\ 1^{3}2^{1}3^{1}6^0\\ 1^{4}2^{-2}3^{-1}6^4\end{array}$	$\begin{array}{c} 1^{-2}2^33^{4}6^{0}\\ 1^{-1}2^{1}3^{3}6^{2}\\ 1^{-1}2^{3}3^{3}6^{0}\\ 1^{0}2^{1}3^{1}6^{3}\\ 1^{0}2^{2}3^{3}6^{-1}\\ 1^{0}2^{3}3^{3}6^{-1}\\ 1^{1}2^{0}3^{2}6^{2}\\ 1^{1}2^{2}3^{-2}6^{4}\\ 1^{1}2^{2}3^{-2}6^{4}\\ 1^{2}2^{-2}3^{2}6^{3}\\ 1^{2}2^{0}3^{2}6^{1}\\ 1^{2}2^{1}3^{3}6^{-1}\\ 1^{2}2^{2}3^{1}6^{0}\\ 1^{3}2^{-1}3^{2}6^{1}\\ 1^{3}2^{0}3^{3}6^{-1}\\ 1^{3}2^{0}3^{-1}\\ 1^{3}2^{3}2^{-2}6^{2}\\ 1^{4}2^{-2}3^{0}6^{3}\\ \end{array}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{2}2^{-1}3^{1}6^{3}$ $1^{2}2^{1}3^{-2}6^{4}$ $1^{2}2^{1}3^{4}6^{-2}$ $1^{2}2^{2}3^{2}6^{-1}$ $1^{3}2^{-2}3^{1}6^{3}$ $1^{3}2^{-1}3^{3}6^{0}$ $1^{3}2^{1}3^{-3}6^{4}$ $1^{3}2^{2}3^{-1}6^{1}$ $1^{4}2^{-2}3^{1}6^{2}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{2}2^{-1}3^{2}6^{2}$ $1^{2}2^{1}3^{-1}6^{3}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-2}6^{2}$ $1^{3}2^{0}3^{-2}6^{4}$ $1^{3}2^{1}3^{-2}6^{3}$ $1^{4}2^{-3}3^{0}6^{4}$ $1^{4}2^{-2}3^{2}6^{1}$	$\begin{array}{c} 1^{-2}2^{4}3^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1}\\ 1^{2}2^{0}3^{-1}6^{4}\\ 1^{2}2^{1}3^{0}6^{2}\\ 1^{2}2^{2}3^{-2}6^{3}\\ 1^{2}2^{2}3^{-4}6^{-3}\\ 1^{3}2^{-1}3^{-1}6^{4}\\ 1^{3}2^{0}3^{-1}6^{3}\\ 1^{3}2^{1}3^{-1}6^{2}\\ 1^{4}2^{-3}3^{1}6^{3}\\ 1^{4}2^{-1}3^{-3}6^{5} \end{array}$
$\begin{array}{c} 1 & 2^{2} 3^{2} 6^{2} \\ 1^{-2} 2^{4} 3^{4} 6^{-1} \\ 1^{-1} 2^{3} 3^{1} 6^{2} \\ 1^{-1} 2^{4} 3^{4} 6^{-2} \\ 1^{0} 2^{2} 3^{1} 6^{2} \\ 1^{0} 2^{3} 3^{1} 6^{1} \\ 1^{1} 2^{-1} 3^{2} 6^{3} \\ 1^{1} 2^{1} 3^{2} 6^{1} \\ 1^{1} 2^{2} 3^{2} 6^{0} \\ 1^{1} 2^{3} 3^{3} 6^{-2} \\ 1^{2} 2^{0} 3^{0} 6^{3} \\ 1^{2} 2^{1} 3^{1} 6^{1} \\ 1^{2} 2^{2} 3^{-1} 6^{2} \\ 1^{2} 2^{3} 3^{1} 6^{-1} \\ 1^{3} 2^{-1} 3^{0} 6^{3} \\ 1^{3} 2^{0} 3^{0} 6^{2} \\ 1^{3} 2^{1} 3^{0} 6^{1} \\ 1^{4} 2^{-3} 3^{2} 6^{2} \\ 1^{4} 2^{-1} 3^{-2} 6^{4} \end{array}$	$\begin{array}{c} 1^{-2}2^33^36^1\\ 1^{-2}2^53^56^{-3}\\ 1^{-1}2^33^26^1\\ 1^{-1}2^43^56^{-3}\\ 1^{0}2^23^26^1\\ 1^{0}2^33^26^0\\ 1^{1}2^03^{1}6^3\\ 1^{1}2^{1}3^{3}6^0\\ 1^{1}2^23^{3}6^{-1}\\ 1^{1}2^33^{4}6^{-3}\\ 1^{2}2^{0}3^{1}6^2\\ 1^{2}2^{1}3^{2}6^0\\ 1^{2}2^{3}3^{2}6^{-2}\\ 1^{3}2^{-1}3^{1}6^2\\ 1^{3}2^{0}3^{1}6^1\\ 1^{3}2^{1}3^{1}6^0\\ 1^{4}2^{-2}3^{-1}6^4\\ 1^{4}2^{-1}3^{-1}6^3\end{array}$	$1^{-2}2^{3}3^{4}6^{0}$ $1^{-1}2^{1}3^{3}6^{2}$ $1^{-1}2^{3}3^{3}6^{0}$ $1^{0}2^{1}3^{1}6^{3}$ $1^{0}2^{2}3^{3}6^{-1}$ $1^{1}2^{0}3^{2}6^{2}$ $1^{1}2^{2}3^{-2}6^{4}$ $1^{1}2^{2}3^{-2}6^{4}$ $1^{2}2^{-2}3^{2}6^{3}$ $1^{2}2^{0}3^{2}6^{1}$ $1^{2}2^{2}3^{1}6^{0}$ $1^{3}2^{-2}3^{0}6^{4}$ $1^{3}2^{-1}3^{2}6^{1}$ $1^{3}2^{0}3^{3}6^{-1}$ $1^{3}2^{2}3^{-2}6^{2}$ $1^{4}2^{-2}3^{0}6^{3}$ $1^{4}2^{-1}3^{0}6^{2}$	$1^{-2}2^{4}3^{1}6^{2}$ $1^{-1}2^{2}3^{2}6^{2}$ $1^{-1}2^{3}3^{4}6^{-1}$ $1^{0}2^{1}3^{2}6^{2}$ $1^{0}2^{2}3^{4}6^{-1}$ $1^{0}2^{3}3^{4}6^{-2}$ $1^{1}2^{0}3^{3}6^{1}$ $1^{1}2^{2}3^{-1}6^{3}$ $1^{2}2^{-1}3^{1}6^{3}$ $1^{2}2^{-1}3^{-2}6^{4}$ $1^{2}2^{1}3^{4}6^{-2}$ $1^{2}2^{2}3^{2}6^{-1}$ $1^{3}2^{-2}3^{1}6^{3}$ $1^{3}2^{-1}3^{3}6^{0}$ $1^{3}2^{1}3^{-3}6^{4}$ $1^{3}2^{2}3^{-1}6^{1}$ $1^{4}2^{-2}3^{1}6^{2}$ $1^{4}2^{0}3^{-3}6^{4}$	$1^{-2}2^{4}3^{2}6^{1}$ $1^{-1}2^{2}3^{3}6^{1}$ $1^{-1}2^{4}3^{2}6^{0}$ $1^{0}2^{1}3^{3}6^{1}$ $1^{0}2^{3}3^{-1}6^{3}$ $1^{0}2^{4}3^{3}6^{-2}$ $1^{1}2^{1}3^{0}6^{3}$ $1^{1}2^{2}3^{0}6^{2}$ $1^{1}2^{3}3^{1}6^{0}$ $1^{2}2^{-1}3^{2}6^{2}$ $1^{2}2^{1}3^{-1}6^{3}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-3}6^{4}$ $1^{2}2^{2}3^{-2}6^{2}$ $1^{3}2^{0}3^{-2}6^{4}$ $1^{3}2^{1}3^{-2}6^{3}$ $1^{4}2^{-3}3^{0}6^{4}$ $1^{4}2^{-2}3^{2}6^{1}$ $1^{4}2^{0}3^{-2}6^{3}$	$\begin{array}{c} 1^{-2}2^43^{3}6^{0}\\ 1^{-1}2^{3}3^{0}6^{3}\\ 1^{-1}2^{4}3^{3}6^{-1}\\ 1^{0}2^{2}3^{0}6^{3}\\ 1^{0}2^{3}3^{0}6^{2}\\ 1^{0}2^{4}3^{4}6^{-3}\\ 1^{1}2^{1}3^{1}6^{2}\\ 1^{1}2^{2}3^{1}6^{1}\\ 1^{1}2^{3}3^{2}6^{-1}\\ 1^{2}2^{0}3^{-1}6^{4}\\ 1^{2}2^{1}3^{0}6^{2}\\ 1^{2}2^{2}3^{-2}6^{3}\\ 1^{3}2^{-1}3^{-1}6^{4}\\ 1^{3}2^{0}3^{-1}6^{3}\\ 1^{3}2^{1}3^{-1}6^{2}\\ 1^{4}2^{-3}3^{1}6^{3}\\ 1^{4}2^{-1}3^{-3}6^{5}\\ 1^{5}2^{-3}3^{-2}6^{5} \end{array}$

Table 3. (continued)

$N = 6, k = 3$ $1^{-2}2^{4}3^{3}6^{1} 1^{-1}$ $1^{0}2^{3}3^{1}6^{2} 1^{0}2$ $1^{1}2^{2}3^{2}6^{1} 1^{1}2$ $1^{2}2^{0}3^{2}6^{2} 1^{2}2$ $1^{2}2^{2}3^{0}6^{2} 1^{2}2$ $1^{3}2^{0}3^{0}6^{3} 1^{3}2$ $1^{4}2^{-2}3^{1}6^{3} 1^{4}2$	$12^{3}3^{2}6^{2}$ $23^{3}6^{1}$ $23^{3}6^{0}$ $23^{3}6^{3}$ $23^{1}6^{1}$ $20^{3}16^{2}$ $2^{-1}3^{-1}6^{4}$	$1^{-1}2^{3}3^{3}6^{1}$ $1^{0}2^{3}3^{3}6^{0}$ $1^{1}2^{3}3^{2}6^{0}$ $1^{2}2^{1}3^{1}6^{2}$ $1^{2}2^{2}3^{2}6^{0}$ $1^{3}2^{0}3^{2}6^{1}$	$1^{-1}2^{4}3^{4}6^{-1}$ $1^{1}2^{1}3^{2}6^{2}$ $1^{1}2^{3}3^{3}6^{-1}$ $1^{2}2^{1}3^{2}6^{1}$ $1^{2}2^{2}3^{3}6^{-1}$ $1^{3}2^{1}3^{-2}6^{4}$	$1^{0}2^{2}3^{2}6^{2}$ $1^{1}2^{2}3^{0}6^{3}$ $1^{1}2^{3}3^{4}6^{-2}$ $1^{2}2^{1}3^{3}6^{0}$ $1^{3}2^{-1}3^{1}6^{3}$ $1^{3}2^{1}3^{-1}6^{3}$	$1^{0}2^{2}3^{3}6^{1}$ $1^{1}2^{2}3^{1}6^{2}$ $1^{2}2^{0}3^{1}6^{3}$ $1^{2}2^{2}3^{-1}6^{3}$ $1^{3}2^{-1}3^{2}6^{2}$ $1^{3}2^{1}3^{0}6^{2}$
$N = 6, \ k = 7/2 \\ 1^1 2^2 3^2 6^2 \qquad 1^2 2$	$2^{1}3^{2}6^{2}$	$1^2 2^2 3^1 6^2$	$1^2 2^2 3^2 6^1$		
$N = 7, \ k = 1/2$ $1^0 7^1$					
$N = 7, \ k = 1$ 1 ⁰ 7 ² 1 ¹ 7	71				
$N = 7, \ k = 3/2 \\ 1^0 7^3 \qquad 1^1 7$	72	$1^{2}7^{1}$			
$N = 7, \ k = 2 \\ 1^1 7^3 \qquad 1^2 7$	72	$1^{3}7^{1}$			
$N = 7, \ k = 5/2 \\ 1^2 7^3 \qquad 1^3 7$	72				
$N = 8, \ k = 1/2 \\ 1^0 2^{-2} 4^5 8^{-2} 1^0 2$	$2^{-1}4^{3}8^{-1}$	$1^{0}2^{0}4^{-1}8^{2}$	$1^{0}2^{0}4^{0}8^{1}$	$1^{0}2^{0}4^{2}8^{-1}$	$1^{0}2^{1}4^{-1}8^{1}$
$\begin{array}{c} N=8,k=1\\ 1^{-2}2^{3}4^{3}8^{-2} & 1^{-2}\\ 1^{-1}2^{0}4^{5}8^{-2} & 1^{-1}\\ 1^{-1}2^{3}4^{-2}8^{2} & 1^{-1}\\ 1^{0}2^{-2}4^{5}8^{-1} & 1^{0}2\\ 1^{0}2^{-1}4^{5}8^{-2} & 1^{0}2\\ 1^{0}2^{1}4^{-2}8^{3} & 1^{0}2\\ 1^{0}2^{2}4^{1}8^{-1} & 1^{0}2\\ 1^{1}2^{-1}4^{1}8^{1} & 1^{1}2\\ 1^{2}2^{-3}4^{5}8^{-2} & 1^{2}2 \end{array}$	$\begin{array}{c} 22^{4}4^{1}8^{-1} \\ 12^{1}4^{3}8^{-1} \\ 12^{3}4^{-1}8^{1} \\ 2^{-2}4^{6}8^{-2} \\ 04^{-1}8^{3} \\ 14^{-1}8^{2} \\ 2^{3}4^{-2}8^{1} \\ 2^{-1}4^{3}8^{-1} \\ 2^{-2}4^{3}8^{-1} \end{array}$	$\begin{array}{c} 1^{-2}2^{5}4^{-3}8^{2} \\ 1^{-1}2^{1}4^{4}8^{-2} \\ 1^{-1}2^{3}4^{1}8^{-1} \\ 1^{0}2^{-2}4^{7}8^{-3} \\ 1^{0}2^{0}4^{0}8^{2} \\ 1^{0}2^{1}4^{0}8^{1} \\ 1^{1}2^{-3}4^{6}8^{-2} \\ 1^{1}2^{0}4^{-1}8^{2} \\ 1^{2}2^{-1}4^{-1}8^{2} \end{array}$	$\begin{array}{c} 1^{-2}2^{5}4^{-2}8^{1} \\ 1^{-1}2^{2}4^{-1}8^{2} \\ 1^{-1}2^{4}4^{-2}8^{1} \\ 1^{0}2^{-1}4^{1}8^{2} \\ 1^{0}2^{0}4^{1}8^{1} \\ 1^{0}2^{1}4^{2}8^{-1} \\ 1^{1}2^{-2}4^{4}8^{-1} \\ 1^{1}2^{0}4^{0}8^{1} \\ 1^{2}2^{-1}4^{0}8^{1} \end{array}$	$\begin{array}{c} 1^{-2}2^{5}4^{0}8^{-1}\\ 1^{-1}2^{2}4^{0}8^{1}\\ 1^{0}2^{-3}4^{7}8^{-2}\\ 1^{0}2^{-1}4^{2}8^{1}\\ 1^{0}2^{0}4^{3}8^{-1}\\ 1^{0}2^{2}4^{-2}8^{2}\\ 1^{1}2^{-2}4^{5}8^{-2}\\ 1^{1}2^{0}4^{2}8^{-1}\\ 1^{2}2^{-1}4^{2}8^{-1}\\ \end{array}$	$\begin{array}{c} 1^{-2}2^{6}4^{-3}8^{1}\\ 1^{-1}2^{2}4^{2}8^{-1}\\ 1^{0}2^{-3}4^{8}8^{-3}\\ 1^{0}2^{-1}4^{4}8^{-1}\\ 1^{0}2^{0}4^{4}8^{-2}\\ 1^{0}2^{2}4^{-1}8^{1}\\ 1^{1}2^{-1}4^{0}8^{2}\\ 1^{1}2^{1}4^{-1}8^{1}\\ 1^{2}2^{0}4^{-1}8^{1} \end{array}$
$N = 8, k = 3/2$ $1^{-3}2^{5}4^{3}8^{-2} \qquad 1^{-3}$ $1^{-3}2^{8}4^{-4}8^{2} \qquad 1^{-3}$ $1^{-2}2^{3}4^{3}8^{-1} \qquad 1^{-2}$ $1^{-2}2^{4}4^{3}8^{-2} \qquad 1^{-2}$	$\frac{32^{6}4^{1}8^{-1}}{32^{8}4^{-3}8^{1}}$ $\frac{32^{8}4^{-3}8^{1}}{2^{2}3^{4}4^{8-2}}$ $\frac{32^{2}4^{-3}8^{3}}{2^{2}2^{5}4^{-3}8^{3}}$	$\frac{1^{-3}2^{6}4^{2}8^{-2}}{1^{-3}2^{8}4^{-1}8^{-1}}$ $\frac{1^{-2}2^{3}4^{5}8^{-3}}{1^{-2}2^{5}4^{-2}8^{2}}$	$ \frac{1^{-3}2^{7}4^{-3}8^{2}}{1^{-3}2^{9}4^{-4}8^{1}} \\ \frac{1^{-2}2^{4}4^{-1}8^{2}}{1^{-2}2^{5}4^{-1}8^{1}} $	$ \frac{1^{-3}2^{7}4^{-2}8^{1}}{1^{-2}2^{2}4^{5}8^{-2}} \\ \frac{1^{-2}2^{4}4^{0}8^{1}}{1^{-2}2^{5}4^{1}8^{-1}} \\ $ To	$\frac{1^{-3}2^{7}4^{0}8^{-1}}{1^{-2}2^{2}4^{6}8^{-3}}$ $\frac{1^{-2}2^{4}4^{2}8^{-1}}{1^{-2}2^{5}4^{2}8^{-2}}$ be continued

Table 3. (continued)

 $1^5 2^{-3} 3^{-1} 6^4$

 $1^5 2^{-2} 3^{-3} 6^5$

To be continued

Table 5. (COI	tillueu)				
$1^{-2}2^{6}4^{-4}8^{3}$	$1^{-2}2^{6}4^{-3}8^{2}$	$1^{-2}2^{6}4^{-2}8^{1}$	$1^{-2}2^{6}4^{0}8^{-1}$	$1^{-2}2^{7}4^{-4}8^{2}$	$1^{-2}2^{7}4^{-3}8^{1}$
$1^{-2}2^{7}4^{-1}8^{-1}$	$1^{-2}2^{8}4^{-4}8^{1}$	$1^{-1}2^{-1}4^{7}8^{-2}$	$1^{-1}2^{-1}4^{8}8^{-3}$	$1^{-1}2^{0}4^{5}8^{-1}$	$1^{-1}2^{0}4^{6}8^{-2}$
$1^{-1}2^{0}4^{7}8^{-3}$	$1^{-1}2^{1}4^{1}8^{2}$	$1^{-1}2^{1}4^{2}8^{1}$	$1^{-1}2^{1}4^{4}8^{-1}$	$1^{-1}2^{1}4^{5}8^{-2}$	$1^{-1}2^{1}4^{6}8^{-3}$
$1^{-1}2^24^{-1}8^3$	$1^{-1}2^{2}4^{0}8^{2}$	$1^{-1}2^{2}4^{1}8^{1}$	$1^{-1}2^{2}4^{3}8^{-1}$	$1^{-1}2^{2}4^{4}8^{-2}$	$1^{-1}2^{3}4^{-2}8^{3}$
$1^{-1}2^{3}4^{-1}8^{2}$	$1^{-1}2^{3}4^{0}8^{1}$	$1^{-1}2^{3}4^{2}8^{-1}$	$1^{-1}2^{3}4^{3}8^{-2}$	$1^{-1}2^{4}4^{-3}8^{3}$	$1^{-1}2^{4}4^{-2}8^{2}$
$1^{-1}2^{4}4^{-1}8^{1}$	$1^{-1}2^{4}4^{1}8^{-1}$	$1^{-1}2^{5}4^{-3}8^{2}$	$1^{-1}2^{5}4^{-2}8^{1}$	$1^{-1}2^{5}4^{0}8^{-1}$	$1^{-1}2^{6}4^{-3}8^{1}$
$1^{0}2^{-3}4^{7}8^{-1}$	$1^{0}2^{-3}4^{8}8^{-2}$	$1^{0}2^{-2}4^{4}8^{1}$	$1^{0}2^{-2}4^{6}8^{-1}$	$1^{0}2^{-2}4^{7}8^{-2}$	$1^{0}2^{-2}4^{8}8^{-3}$
$1^{0}2^{-1}4^{2}8^{2}$	$1^{0}2^{-1}4^{3}8^{1}$	$1^{0}2^{-1}4^{5}8^{-1}$	$1^{0}2^{-1}4^{6}8^{-2}$	$1^{0}2^{-1}4^{7}8^{-3}$	$1^{0}2^{0}4^{1}8^{2}$
$1^{0}2^{0}4^{2}8^{1}$	$1^{0}2^{0}4^{4}8^{-1}$	$1^{0}2^{0}4^{5}8^{-2}$	$1^{0}2^{0}4^{6}8^{-3}$	$1^{0}2^{1}4^{-1}8^{3}$	$1^{0}2^{1}4^{0}8^{2}$
$1^{0}2^{1}4^{1}8^{1}$	$1^{0}2^{1}4^{3}8^{-1}$	$1^{0}2^{1}4^{4}8^{-2}$	$1^{0}2^{2}4^{-2}8^{3}$	$1^{0}2^{2}4^{-1}8^{2}$	$1^{0}2^{2}4^{0}8^{1}$
$1^{0}2^{2}4^{2}8^{-1}$	$1^{0}2^{2}4^{3}8^{-2}$	$1^{0}2^{3}4^{-3}8^{3}$	$1^{0}2^{3}4^{-2}8^{2}$	$1^{0}2^{3}4^{-1}8^{1}$	$1^{0}2^{3}4^{1}8^{-1}$
$1^{0}2^{4}4^{-3}8^{2}$	$1^{0}2^{4}4^{-2}8^{1}$	$1^{0}2^{4}4^{0}8^{-1}$	$1^{0}2^{5}4^{-3}8^{1}$	$1^{1}2^{-4}4^{8}8^{-2}$	$1^{1}2^{-4}4^{9}8^{-3}$
$1^{1}2^{-3}4^{6}8^{-1}$	$1^{1}2^{-3}4^{7}8^{-2}$	$1^{1}2^{-3}4^{8}8^{-3}$	$1^{1}2^{-2}4^{2}8^{2}$	$1^{1}2^{-2}4^{3}8^{1}$	$1^{1}2^{-2}4^{5}8^{-1}$
$1^{1}2^{-2}4^{6}8^{-2}$	$1^{1}2^{-2}4^{7}8^{-3}$	$1^{1}2^{-1}4^{0}8^{3}$	$1^{1}2^{-1}4^{1}8^{2}$	$1^{1}2^{-1}4^{2}8^{1}$	$1^{1}2^{-1}4^{4}8^{-1}$
$1^{1}2^{-1}4^{5}8^{-2}$	$1^{1}2^{0}4^{-1}8^{3}$	$1^1 2^0 4^0 8^2$	$1^{1}2^{0}4^{1}8^{1}$	$1^{1}2^{0}4^{3}8^{-1}$	$1^{1}2^{0}4^{4}8^{-2}$
$1^{1}2^{1}4^{-2}8^{3}$	$1^{1}2^{1}4^{-1}8^{2}$	$1^{1}2^{1}4^{0}8^{1}$	$1^{1}2^{1}4^{2}8^{-1}$	$1^{1}2^{2}4^{-2}8^{2}$	$1^{1}2^{2}4^{-1}8^{1}$
$1^{1}2^{2}4^{1}8^{-1}$	$1^{1}2^{3}4^{-2}8^{1}$	$1^{2}2^{-4}4^{7}8^{-2}$	$1^2 2^{-4} 4^8 8^{-3}$	$1^2 2^{-3} 4^5 8^{-1}$	$1^2 2^{-3} 4^6 8^{-2}$
$1^2 2^{-3} 4^7 8^{-3}$	$1^2 2^{-2} 4^1 8^2$	$1^2 2^{-2} 4^2 8^1$	$1^2 2^{-2} 4^4 8^{-1}$	$1^2 2^{-2} 4^5 8^{-2}$	$1^2 2^{-1} 4^{-1} 8^3$
$1^2 2^{-1} 4^0 8^2$	$1^2 2^{-1} 4^1 8^1$	$1^2 2^{-1} 4^3 8^{-1}$	$1^2 2^{-1} 4^4 8^{-2}$	$1^{2}2^{0}4^{-2}8^{3}$	$1^2 2^0 4^{-1} 8^2$
$1^2 2^0 4^0 8^1$	$1^{2}2^{0}4^{2}8^{-1}$	$1^{2}2^{1}4^{-2}8^{2}$	$1^{2}2^{1}4^{-1}8^{1}$	$1^{2}2^{1}4^{1}8^{-1}$	$1^{2}2^{2}4^{-2}8^{1}$
$1^{3}2^{-4}4^{6}8^{-2}$	$1^{3}2^{-3}4^{4}8^{-1}$	$1^{3}2^{-3}4^{5}8^{-2}$	$1^{3}2^{-2}4^{0}8^{2}$	$1^{3}2^{-2}4^{1}8^{1}$	$1^{3}2^{-2}4^{3}8^{-1}$
$1^{3}2^{-1}4^{-1}8^{2}$	$1^3 2^{-1} 4^0 8^1$	$1^{3}2^{-1}4^{2}8^{-1}$	$1^{3}2^{0}4^{-1}8^{1}$		
$N = 8 \ k = 2$)				
$1^{-3}2^{4}4^{5}8^{-2}$	$1^{-3}2^{4}4^{6}8^{-3}$	$1^{-3}2^{5}4^{3}8^{-1}$	$1^{-3}2^{5}4^{4}8^{-2}$	$1^{-3}2^{5}4^{5}8^{-3}$	$1^{-3}2^{6}4^{-1}8^{2}$
$1^{-3}2^{6}4^{0}8^{1}$	$1^{-3}2^{6}4^{2}8^{-1}$	$1^{-3}2^{6}4^{3}8^{-2}$	$1^{-3}2^{6}4^{4}8^{-3}$	$1^{-3}2^{7}4^{-3}8^{3}$	$1^{-3}2^{7}4^{-2}8^{2}$
$1^{-3}2^{7}4^{-1}8^{1}$	$1^{-3}2^{7}4^{1}8^{-1}$	$1^{-3}2^{8}4^{-4}8^{3}$	$1^{-3}2^{8}4^{-3}8^{2}$	$1^{-3}2^{8}4^{-2}8^{1}$	$1^{-3}2^{9}4^{-5}8^{3}$
$1^{-2}2^{2}4^{5}8^{-1}$	$1^{-2}2^{2}4^{6}8^{-2}$	$1^{-2}2^{3}4^{2}8^{1}$	$1^{-2}2^{3}4^{4}8^{-1}$	$1^{-2}2^{3}4^{5}8^{-2}$	$1^{-2}2^{3}4^{6}8^{-3}$
$1^{-2}2^{4}4^{0}8^{2}$	$1^{-2}2^{4}4^{1}8^{1}$	$1^{-2}2^{4}4^{3}8^{-1}$	$1^{-2}2^{4}4^{4}8^{-2}$	$1^{-2}2^{4}4^{5}8^{-3}$	$1^{-2}2^{5}4^{-1}8^{2}$
$1^{-2}2^{5}4^{0}8^{1}$	$1^{-2}2^{5}4^{2}8^{-1}$	$1^{-2}2^{5}4^{3}8^{-2}$	$1^{-2}2^{5}4^{4}8^{-3}$	$1^{-2}2^{6}4^{-3}8^{3}$	$1^{-2}2^{6}4^{-2}8^{2}$
$1^{-2}2^{6}4^{-1}8^{1}$	$1^{-2}2^{6}4^{1}8^{-1}$	$1^{-2}2^{6}4^{2}8^{-2}$	$1^{-2}2^{7}4^{-4}8^{3}$	$1^{-2}2^{7}4^{-3}8^{2}$	$1^{-2}2^{7}4^{-2}8^{1}$
$1^{-2}2^{7}4^{0}8^{-1}$	$1^{-2}2^{8}4^{-5}8^{3}$	$1^{-2}2^{8}4^{-4}8^{2}$	$1^{-2}2^{8}4^{-3}8^{1}$	$1^{-1}2^{-1}4^{7}8^{-1}$	$1^{-1}2^{0}4^{4}8^{1}$
$1^{-1}2^{0}4^{6}8^{-1}$	$1^{-1}2^{0}4^{7}8^{-2}$	$1^{-1}2^{1}4^{3}8^{1}$	$1^{-1}2^{1}4^{5}8^{-1}$	$1^{-1}2^{1}4^{6}8^{-2}$	$1^{-1}2^{1}4^{7}8^{-3}$
$1^{-1}2^{2}4^{1}8^{2}$	$1^{-1}2^{2}4^{2}8^{1}$	$1^{-1}2^{2}4^{4}8^{-1}$	$1^{-1}2^{2}4^{5}8^{-2}$	$1^{-1}2^{2}4^{6}8^{-3}$	$1^{-1}2^{3}4^{0}8^{2}$
$1^{-1}2^{3}4^{1}8^{1}$	$1^{-1}2^{3}4^{3}8^{-1}$	$1^{-1}2^{3}4^{4}8^{-2}$	$1^{-1}2^{3}4^{5}8^{-3}$	$1^{-1}2^{4}4^{-2}8^{3}$	$1^{-1}2^{4}4^{-1}8^{2}$
$1^{-1}2^{4}4^{0}8^{1}$	$1^{-1}2^{4}4^{2}8^{-1}$	$1^{-1}2^{4}4^{3}8^{-2}$	$1^{-1}2^{5}4^{-3}8^{3}$	$1^{-1}2^{5}4^{-2}8^{2}$	$1^{-1}2^{5}4^{-1}8^{1}$
$1^{-1}2^{5}4^{1}8^{-1}$	$1^{-1}2^{5}4^{2}8^{-2}$	$1^{-1}2^{6}4^{-4}8^{3}$	$1^{-1}2^{6}4^{-3}8^{2}$	$1^{-1}2^{6}4^{-2}8^{1}$	$1^{-1}2^{6}4^{0}8^{-1}$
$1^{-1}2^{7}4^{-4}8^{2}$	$1^{-1}2^{7}4^{-3}8^{1}$	$1^{-1}2^{7}4^{-1}8^{-1}$	$1^{-1}2^{8}4^{-4}8^{1}$	$1^{0}2^{-2}4^{7}8^{-1}$	$1^{0}2^{-1}4^{4}8^{1}$
$1^{0}2^{-1}4^{6}8^{-1}$	$1^{0}2^{-1}4^{7}8^{-2}$	$1^{0}2^{0}4^{3}8^{1}$	$1^{0}2^{0}4^{5}8^{-1}$	$1^{0}2^{0}4^{6}8^{-2}$	$1^{0}2^{0}4^{7}8^{-3}$
$1^{0}2^{1}4^{1}8^{2}$	$1^{0}2^{1}4^{2}8^{1}$	$1^{0}2^{1}4^{4}8^{-1}$	$1^{0}2^{1}4^{5}8^{-2}$	$1^{0}2^{1}4^{6}8^{-3}$	$1^{0}2^{2}4^{0}8^{2}$
$1^{0}2^{2}4^{1}8^{1}$	$1^{0}2^{2}4^{3}8^{-1}$	$1^{0}2^{2}4^{4}8^{-2}$	$1^{0}2^{2}4^{5}8^{-3}$	$1^{0}2^{3}4^{-2}8^{3}$	$1^{0}2^{3}4^{-1}8^{2}$
$1^{0}2^{3}4^{0}8^{1}$	$1^{0}2^{3}4^{2}8^{-1}$	$1^{0}2^{3}4^{3}8^{-2}$	$1^{0}2^{4}4^{-3}8^{3}$	$1^{0}2^{4}4^{-2}8^{2}$	$1^{0}2^{4}4^{-1}8^{1}$
$1^{0}2^{4}4^{1}8^{-1}$	$1^{0}2^{4}4^{2}8^{-2}$	$1^{0}2^{5}4^{-4}8^{3}$	$1^{0}2^{5}4^{-3}8^{2}$	$1^{0}2^{5}4^{-2}8^{1}$	$1^{0}2^{5}4^{0}8^{-1}$

Table 3. (continued)

$1^{0}2^{6}4^{-4}8^{2}$	$1^{0}2^{6}4^{-3}8^{1}$	$1^{0}2^{6}4^{-1}8^{-1}$	$1^{0}2^{7}4^{-4}8^{1}$	$1^{1}2^{-4}4^{8}8^{-1}$	$1^{1}2^{-3}4^{5}8^{1}$
$1^{1}2^{-3}4^{7}8^{-1}$	$1^{1}2^{-3}4^{8}8^{-2}$	$1^{1}2^{-2}4^{4}8^{1}$	$1^{1}2^{-2}4^{6}8^{-1}$	$1^{1}2^{-2}4^{7}8^{-2}$	$1^{1}2^{-2}4^{8}8^{-3}$
$1^{1}2^{-1}4^{2}8^{2}$	$1^{1}2^{-1}4^{3}8^{1}$	$1^{1}2^{-1}4^{5}8^{-1}$	$1^{1}2^{-1}4^{6}8^{-2}$	$1^{1}2^{-1}4^{7}8^{-3}$	$1^{1}2^{0}4^{1}8^{2}$
$1^{1}2^{0}4^{2}8^{1}$	$1^{1}2^{0}4^{4}8^{-1}$	$1^{1}2^{0}4^{5}8^{-2}$	$1^{1}2^{0}4^{6}8^{-3}$	$1^{1}2^{1}4^{-1}8^{3}$	$1^{1}2^{1}4^{0}8^{2}$
$1^{1}2^{1}4^{1}8^{1}$	$1^{1}2^{1}4^{3}8^{-1}$	$1^{1}2^{1}4^{4}8^{-2}$	$1^{1}2^{2}4^{-2}8^{3}$	$1^{1}2^{2}4^{-1}8^{2}$	$1^{1}2^{2}4^{0}8^{1}$
$1^{1}2^{2}4^{2}8^{-1}$	$1^{1}2^{2}4^{3}8^{-2}$	$1^{1}2^{3}4^{-3}8^{3}$	$1^{1}2^{3}4^{-2}8^{2}$	$1^{1}2^{3}4^{-1}8^{1}$	$1^{1}2^{3}4^{1}8^{-1}$
$1^{1}2^{4}4^{-3}8^{2}$	$1^{1}2^{4}4^{-2}8^{1}$	$1^{1}2^{4}4^{0}8^{-1}$	$1^{1}2^{5}4^{-3}8^{1}$	$1^{2}2^{-4}4^{7}8^{-1}$	$1^2 2^{-4} 4^8 8^{-2}$
$1^2 2^{-3} 4^4 8^1$	$1^{2}2^{-3}4^{6}8^{-1}$	$1^{2}2^{-3}4^{7}8^{-2}$	$1^{2}2^{-3}4^{8}8^{-3}$	$1^2 2^{-2} 4^2 8^2$	$1^2 2^{-2} 4^3 8^1$
$1^2 2^{-2} 4^5 8^{-1}$	$1^{2}2^{-2}4^{6}8^{-2}$	$1^{2}2^{-2}4^{7}8^{-3}$	$1^2 2^{-1} 4^1 8^2$	$1^2 2^{-1} 4^2 8^1$	$1^{2}2^{-1}4^{4}8^{-1}$
$1^2 2^{-1} 4^5 8^{-2}$	$1^{2}2^{-1}4^{6}8^{-3}$	$1^{2}2^{0}4^{-1}8^{3}$	$1^2 2^0 4^0 8^2$	$1^{2}2^{0}4^{1}8^{1}$	$1^{2}2^{0}4^{3}8^{-1}$
$1^{2}2^{0}4^{4}8^{-2}$	$1^{2}2^{1}4^{-2}8^{3}$	$1^{2}2^{1}4^{-1}8^{2}$	$1^{2}2^{1}4^{0}8^{1}$	$1^{2}2^{1}4^{2}8^{-1}$	$1^2 2^2 4^{-3} 8^3$
$1^2 2^2 4^{-2} 8^2$	$1^{2}2^{2}4^{-1}8^{1}$	$1^{3}2^{-5}4^{8}8^{-2}$	$1^{3}2^{-5}4^{9}8^{-3}$	$1^{3}2^{-4}4^{6}8^{-1}$	$1^{3}2^{-4}4^{7}8^{-2}$
$1^{3}2^{-4}4^{8}8^{-3}$	$1^{3}2^{-3}4^{2}8^{2}$	$1^{3}2^{-3}4^{3}8^{1}$	$1^{3}2^{-3}4^{5}8^{-1}$	$1^{3}2^{-3}4^{6}8^{-2}$	$1^{3}2^{-3}4^{7}8^{-3}$
$1^{3}2^{-2}4^{0}8^{3}$	$1^{3}2^{-2}4^{1}8^{2}$	$1^{3}2^{-2}4^{2}8^{1}$	$1^{3}2^{-2}4^{4}8^{-1}$	$1^{3}2^{-1}4^{-1}8^{3}$	$1^{3}2^{-1}4^{0}8^{2}$
$1^{3}2^{-1}4^{1}8^{1}$	$1^{3}2^{0}4^{-2}8^{3}$	_	_	-	-
N = 8, k = 5/2					
$1^{-3}2^{4}4^{5}8^{-1}$	$1^{-3}2^{5}4^{2}8^{1}$	$1^{-3}2^{5}4^{4}8^{-1}$	$1^{-3}2^{5}4^{5}8^{-2}$	$1^{-3}2^{6}4^{1}8^{1}$	$1^{-3}2^{6}4^{3}8^{-1}$
$1^{-3}2^{7}4^{-1}8^{2}$	$1^{-3}2^{7}4^{0}8^{1}$	$1^{-2}2^{3}4^{5}8^{-1}$	$1^{-2}2^{4}4^{2}8^{1}$	$1^{-2}2^{4}4^{4}8^{-1}$	$1^{-2}2^{4}4^{5}8^{-2}$
$1^{-2}2^{5}4^{1}8^{1}$	$1^{-2}2^{5}4^{3}8^{-1}$	$1^{-2}2^{5}4^{4}8^{-2}$	$1^{-2}2^{5}4^{5}8^{-3}$	$1^{-2}2^{6}4^{-1}8^{2}$	$1^{-2}2^{6}4^{0}8^{1}$
$1^{-2}2^{6}4^{2}8^{-1}$	$1^{-2}2^{7}4^{-2}8^{2}$	$1^{-2}2^{7}4^{-1}8^{1}$	$1^{-2}2^{8}4^{-4}8^{3}$	$1^{-1}2^{1}4^{6}8^{-1}$	$1^{-1}2^{2}4^{3}8^{1}$
$1^{-1}2^{2}4^{5}8^{-1}$	$1^{-1}2^{2}4^{6}8^{-2}$	$1^{-1}2^{3}4^{2}8^{1}$	$1^{-1}2^{3}4^{4}8^{-1}$	$1^{-1}2^{3}4^{5}8^{-2}$	$1^{-1}2^{3}4^{6}8^{-3}$
$1^{-1}2^{4}4^{0}8^{2}$	$1^{-1}2^{4}4^{1}8^{1}$	$1^{-1}2^{4}4^{3}8^{-1}$	$1^{-1}2^{4}4^{4}8^{-2}$	$1^{-1}2^{4}4^{5}8^{-3}$	$1^{-1}2^{5}4^{-1}8^{2}$
$1^{-1}2^{5}4^{0}8^{1}$	$1^{-1}2^{5}4^{2}8^{-1}$	$1^{-1}2^{5}4^{3}8^{-2}$	$1^{-1}2^{5}4^{4}8^{-3}$	$1^{-1}2^{6}4^{-3}8^{3}$	$1^{-1}2^{6}4^{-2}8^{2}$
$1^{-1}2^{6}4^{-1}8^{1}$	$1^{-1}2^{6}4^{1}8^{-1}$	$1^{-1}2^{7}4^{-4}8^{3}$	$1^{-1}2^{7}4^{-3}8^{2}$	$1^{-1}2^{7}4^{-2}8^{1}$	$1^{-1}2^{8}4^{-5}8^{3}$
$1^{0}2^{0}4^{6}8^{-1}$	$1^{0}2^{1}4^{3}8^{1}$	$1^{0}2^{1}4^{5}8^{-1}$	$1^{0}2^{1}4^{6}8^{-2}$	$1^{0}2^{2}4^{2}8^{1}$	$1^{0}2^{2}4^{4}8^{-1}$
$1^{0}2^{2}4^{5}8^{-2}$	$1^{0}2^{2}4^{6}8^{-3}$	$1^{0}2^{3}4^{0}8^{2}$	$1^{0}2^{3}4^{1}8^{1}$	$1^{0}2^{3}4^{3}8^{-1}$	$1^{0}2^{3}4^{4}8^{-2}$
$1^{0}2^{3}4^{5}8^{-3}$	$1^{0}2^{4}4^{-1}8^{2}$	$1^{0}2^{4}4^{0}8^{1}$	$1^{0}2^{4}4^{2}8^{-1}$	$1^{0}2^{4}4^{3}8^{-2}$	$1^{0}2^{4}4^{4}8^{-3}$
$1^{0}2^{5}4^{-3}8^{3}$	$1^{0}2^{5}4^{-2}8^{2}$	$1^{0}2^{5}4^{-1}8^{1}$	$1^{0}2^{5}4^{1}8^{-1}$	$1^{0}2^{5}4^{2}8^{-2}$	$1^{0}2^{6}4^{-4}8^{3}$
$1^{0}2^{6}4^{-3}8^{2}$	$1^{0}2^{6}4^{-2}8^{1}$	$1^{0}2^{6}4^{0}8^{-1}$	$1^{0}2^{7}4^{-5}8^{3}$	$1^{0}2^{7}4^{-4}8^{2}$	$1^{0}2^{7}4^{-3}8^{1}$
$1^{1}2^{-2}4^{7}8^{-1}$	$1^{1}2^{-1}4^{4}8^{1}$	$1^{1}2^{-1}4^{6}8^{-1}$	$1^{1}2^{-1}4^{7}8^{-2}$	$1^{1}2^{0}4^{3}8^{1}$	$1^{1}2^{0}4^{5}8^{-1}$
$1^{1}2^{0}4^{6}8^{-2}$	$1^{1}2^{0}4^{7}8^{-3}$	$1^{1}2^{1}4^{1}8^{2}$	$1^{1}2^{1}4^{2}8^{1}$	$1^{1}2^{1}4^{4}8^{-1}$	$1^{1}2^{1}4^{5}8^{-2}$
$1^{1}2^{1}4^{6}8^{-3}$	$1^{1}2^{2}4^{0}8^{2}$	$1^{1}2^{2}4^{1}8^{1}$	$1^{1}2^{2}4^{3}8^{-1}$	$1^{1}2^{2}4^{4}8^{-2}$	$1^{1}2^{2}4^{5}8^{-3}$
$1^{1}2^{3}4^{-2}8^{3}$	$1^{1}2^{3}4^{-1}8^{2}$	$1^{1}2^{3}4^{0}8^{1}$	$1^{1}2^{3}4^{2}8^{-1}$	$1^{1}2^{4}4^{-3}8^{3}$	$1^{1}2^{4}4^{-2}8^{2}$
$1^{1}2^{4}4^{-1}8^{1}$	$1^{1}2^{5}4^{-4}8^{3}$	$1^2 2^{-3} 4^7 8^{-1}$	$1^2 2^{-2} 4^4 8^1$	$1^2 2^{-2} 4^6 8^{-1}$	$1^{2}2^{-2}4^{7}8^{-2}$
$1^2 2^{-1} 4^3 8^1$	$1^2 2^{-1} 4^5 8^{-1}$	$1^2 2^{-1} 4^6 8^{-2}$	$1^2 2^{-1} 4^7 8^{-3}$	$1^2 2^0 4^1 8^2$	$1^2 2^0 4^2 8^1$
$1^{2}2^{0}4^{4}8^{-1}$	$1^2 2^1 4^0 8^2$	$1^2 2^1 4^1 8^1$	$1^{2}2^{2}4^{-2}8^{3}$	$1^{3}2^{-5}4^{8}8^{-1}$	$1^{3}2^{-4}4^{5}8^{1}$
$1^{3}2^{-4}4^{7}8^{-1}$	$1^{3}2^{-4}4^{8}8^{-2}$	$1^{3}2^{-3}4^{4}8^{1}$	$1^{3}2^{-3}4^{6}8^{-1}$	$1^{3}2^{-2}4^{2}8^{2}$	$1^{3}2^{-2}4^{3}8^{1}$
N = 8, k = 3					
$1^{-2}2^{5}4^{4}8^{-1}$	$1^{-2}2^{6}4^{1}8^{1}$	$1^{-1}2^{3}4^{5}8^{-1}$	$1^{-1}2^{4}4^{2}8^{1}$	$1^{-1}2^{4}4^{4}8^{-1}$	$1^{-1}2^{4}4^{5}8^{-2}$
$1^{-1}2^{5}4^{1}8^{1}$	$1^{-1}2^{5}4^{3}8^{-1}$	$1^{-1}2^{6}4^{-1}8^{2}$	$1^{-1}2^{6}4^{0}8^{1}$	$1^{0}2^{2}4^{5}8^{-1}$	$1^{0}2^{3}4^{2}8^{1}$
$1^{0}2^{3}4^{4}8^{-1}$	$1^{0}2^{3}4^{5}8^{-2}$	$1^{0}2^{4}4^{1}8^{1}$	$1^{0}2^{4}4^{3}8^{-1}$	$1^{0}2^{4}4^{4}8^{-2}$	$1^{0}2^{4}4^{5}8^{-3}$
				То	be continued

Table 3. (continued)

10010 0. (001.	(initiaca)				
$ \begin{array}{r} 1^{0}2^{5}4^{-1}8^{2} \\ 1^{1}2^{0}4^{6}8^{-1} \\ 1^{1}2^{3}4^{0}8^{2} \end{array} $	$ \begin{array}{r} 1^{0}2^{5}4^{0}8^{1} \\ 1^{1}2^{1}4^{3}8^{1} \\ 1^{1}2^{3}4^{1}8^{1} \\ \end{array} $	$ \frac{1^{0}2^{5}4^{2}8^{-1}}{1^{1}2^{1}4^{5}8^{-1}} \\ 1^{2}2^{-1}4^{6}8^{-1} $	$ \begin{array}{r} 1^{0}2^{6}4^{-2}8^{2} \\ 1^{1}2^{1}4^{6}8^{-2} \\ 1^{2}2^{0}4^{3}8^{1} \end{array} $	$ 1^{0}2^{6}4^{-1}8^{1} \\ 1^{1}2^{2}4^{2}8^{1} $	$\frac{1^{0}2^{7}4^{-4}8^{3}}{1^{1}2^{2}4^{4}8^{-1}}$
$N = 8, \ k = 7$ $1^{0}2^{4}4^{4}8^{-1}$	$\frac{1}{2}/2$ $1^{0}2^{5}4^{1}8^{1}$				
$N = 9, \ k = 1$ $1^{0}3^{0}9^{1}$	/2				
N = 9, k = 1 1 ⁻¹ 3 ⁴ 9 ⁻¹ 1 ² 3 ⁻¹ 9 ¹	$1^{0}3^{0}9^{2}$	$1^{0}3^{1}9^{1}$	$1^{0}3^{3}9^{-1}$	$1^{1}3^{-1}9^{2}$	$1^{1}3^{0}9^{1}$
N = 9, k = 3 1 ⁻² 3 ⁷ 9 ⁻² 1 ¹ 3 ⁰ 9 ²	$ \begin{array}{c} /2 \\ 1^{-1} 3^{3} 9^{1} \\ 1^{1} 3^{1} 9^{1} \end{array} $	$1^{-1}3^{5}9^{-1}$ $1^{1}3^{3}9^{-1}$	$ 1^{0}3^{1}9^{2} \\ 1^{2}3^{-1}9^{2} $	$\frac{1^0 3^2 9^1}{1^2 3^0 9^1}$	$1^{0}3^{4}9^{-1}$
N = 9, k = 2 $1^{-2}3^{7}9^{-1}$ $1^{0}3^{5}9^{-1}$ $1^{2}3^{1}9^{1}$	$1^{-2}3^{8}9^{-2}$ $1^{0}3^{6}9^{-2}$	$1^{-1}3^{4}9^{1}$ $1^{1}3^{1}9^{2}$	$1^{-1}3^{6}9^{-1}$ $1^{1}3^{2}9^{1}$	$\frac{1^{-1}3^{7}9^{-2}}{1^{1}3^{4}9^{-1}}$	$1^{0}3^{3}9^{1}$ $1^{2}3^{0}9^{2}$
N = 9, k = 5 $1^{-2}3^{8}9^{-1}$ $1^{0}3^{6}9^{-1}$		$\frac{1^{-1}3^59^1}{1^13^39^1}$	$1^{-1}3^{7}9^{-1} \\ 1^{1}3^{5}9^{-1}$	$\frac{1^{-1}3^89^{-2}}{1^23^{1}9^2}$	$1^{0}3^{4}9^{1}$
$N = 9, k = 3$ $1^{-1}3^{8}9^{-1}$	$1^{0}3^{7}9^{-1}$	$1^{0}3^{8}9^{-2}$	$1^{1}3^{4}9^{1}$		
N = 9, k = 7/2 1 ⁰ 3 ⁸ 9 ⁻¹					
$N = 10, k = 1^{0}2^{0}5^{-1}10^{2}$	$\frac{1/2}{1^0 2^0 5^0 10^1}$	$1^{0}2^{0}5^{2}10^{-1}$			
$\overline{N = 10, k} = 1^{-1}2^{2}5^{-1}10^{2}$ $1^{0}2^{1}5^{-1}10^{2}$ $1^{1}2^{0}5^{2}10^{-1}$	$1 \\ 1^{-1}2^{2}5^{0}10^{1} \\ 1^{0}2^{1}5^{0}10^{1} \\ 1^{2}2^{-1}5^{-1}10^{2} $	$\frac{1^{-1}2^{2}5^{1}10^{0}}{1^{0}2^{1}5^{1}10^{0}}$ $\frac{1^{2}2^{-1}5^{0}10^{1}}{1^{2}2^{-1}5^{0}10^{1}}$	$\frac{1^{-1}2^{2}5^{2}10^{-1}}{1^{0}2^{1}5^{2}10^{-1}}$ $\frac{1^{2}2^{-1}5^{1}10^{0}}{1^{0}}$	$\frac{1^{0}2^{0}5^{0}10^{2}}{1^{1}2^{0}5^{-1}10^{2}}\\1^{2}2^{-1}5^{2}10^{-1}$	$ \frac{1^{0}2^{0}5^{1}10^{1}}{1^{1}2^{0}5^{0}10^{1}} $
$N = 10, k = 1^{-1}2^{2}5^{0}10^{2}$ $1^{0}2^{2}5^{-1}10^{2}$ $1^{1}2^{1}5^{-1}10^{2}$ $1^{2}2^{-1}5^{2}10^{0}$	$3/2 \\ 1^{-1}2^{2}5^{1}10^{1} \\ 1^{0}2^{2}5^{0}10^{1} \\ 1^{1}2^{1}5^{0}10^{1} \\ 1^{2}2^{0}5^{-1}10^{2} $	$ \begin{array}{r} 1^{-1}2^{2}5^{2}10^{0} \\ 1^{0}2^{2}5^{1}10^{0} \\ 1^{1}2^{1}5^{1}10^{0} \\ 1^{2}2^{0}5^{0}10^{1} \end{array} $	$1^{0}2^{1}5^{0}10^{2}$ $1^{0}2^{2}5^{2}10^{-1}$ $1^{1}2^{1}5^{2}10^{-1}$ $1^{2}2^{0}5^{2}10^{-1}$	$ \begin{array}{c} 1^{0}2^{1}5^{1}10^{1} \\ 1^{1}2^{0}5^{0}10^{2} \\ 1^{2}2^{-1}5^{0}10^{2} \end{array} $	$ \begin{array}{r} 1^{0}2^{1}5^{2}10^{0} \\ 1^{1}2^{0}5^{1}10^{1} \\ 1^{2}2^{-1}5^{1}10^{1} \end{array} $
$N = 10, k = 1^0 2^2 5^1 10^1$	$2 1^{0}2^{2}5^{2}10^{0}$	$1^{1}2^{1}5^{0}10^{2}$	$1^{1}2^{1}5^{1}10^{1}$	$1^{1}2^{1}5^{2}10^{0}$	$1^2 2^0 5^0 10^2$
				To	be continued

Table 3. (continued)

$1^2 2^0 5^1 10^1$					
$N = 11, k = 1^{0}11^{3}$	3/2				
$N = 11, k = 1^{0}11^{4}$	$2 \\ 1^1 11^3$	$1^{3}11^{1}$			
$N = 11, k = 1^{11} = 1^{11}$	5/2 1 ⁴ 11 ¹				
$N = 13, k = 1^{0}13^{1}$	1/2				
N = 13, k = 1 1 ¹ 13 ¹					
$N = 14, k = 1^{0}2^{0}7^{-2}14^{4}$	$1 \\ 1^{0} 2^{0} 7^{-1} 14^{3}$	$1^{0}2^{0}7^{0}14^{2}$	$1^{0}2^{0}7^{3}14^{-1}$	$1^{0}2^{0}7^{4}14^{-2}$	
$\begin{split} N &= 14, k = \\ 1^{-2} 2^4 7^0 14^1 \\ 1^{-1} 2^3 7^0 14^1 \\ 1^0 2^1 7^0 14^2 \\ 1^0 2^2 7^2 14^{-1} \\ 1^2 2^{-1} 7^{-2} 14^4 \\ 1^3 2^{-1} 7^0 14^1 \end{split}$	$\begin{array}{c} 3/2\\ 1^{-2}2^{4}7^{1}14^{0}\\ 1^{-1}2^{3}7^{1}14^{0}\\ 1^{0}2^{1}7^{2}14^{0}\\ 1^{1}2^{0}7^{-2}14^{4}\\ 1^{2}2^{-1}7^{-1}14^{3}\\ 1^{3}2^{-1}7^{1}14^{0} \end{array}$	$\begin{array}{c} 1^{-2}2^{4}7^{2}14^{-1}\\ 1^{-1}2^{3}7^{2}14^{-1}\\ 1^{0}2^{1}7^{3}14^{-1}\\ 1^{1}2^{0}7^{-1}14^{3}\\ 1^{2}2^{-1}7^{0}14^{2}\\ 1^{4}2^{-2}7^{-1}14^{2} \end{array}$	$1^{-1}2^{2}7^{2}14^{0}$ $1^{0}2^{0}7^{0}14^{3}$ $1^{0}2^{1}7^{4}14^{-2}$ $1^{1}2^{0}7^{0}14^{2}$ $1^{2}2^{0}7^{-1}14^{2}$ $1^{4}2^{-2}7^{0}14^{1}$	$1^{-1}2^{2}7^{3}14^{-1}$ $1^{0}2^{1}7^{-2}14^{4}$ $1^{0}2^{2}7^{0}14^{1}$ $1^{1}2^{0}7^{3}14^{-1}$ $1^{2}2^{0}7^{0}14^{1}$ $1^{4}2^{-2}7^{1}14^{0}$	$1^{-1}2^{2}7^{4}14^{-2}$ $1^{0}2^{1}7^{-1}14^{3}$ $1^{0}2^{2}7^{1}14^{0}$ $1^{1}2^{0}7^{4}14^{-2}$ $1^{3}2^{-1}7^{-1}14^{2}$
$N = 14, k = 1^{0}2^{1}7^{0}14^{3}$	$2 \\ 1^0 2^1 7^3 14^0$	$1^{0}2^{3}7^{0}14^{1}$	$1^{0}2^{3}7^{1}14^{0}$	$1^{1}2^{0}7^{0}14^{3}$	$1^{3}2^{0}7^{0}14^{1}$
$N = 15, k = 1^{0}3^{0}5^{-1}15^{3}$	$1 \\ 1^0 3^0 5^0 15^2$	$1^{0}3^{0}5^{3}15^{-1}$			
$N = 15, k = 1^{-1}3^{3}5^{0}15^{1} \\ 1^{0}3^{1}5^{3}15^{-1} \\ 1^{2}3^{0}5^{0}15^{1}$	$\begin{array}{c} 3/2 \\ 1^{-1}3^35^{1}15^{0} \\ 1^{0}3^25^{0}15^{1} \\ 1^{3}3^{-1}5^{0}15^{1} \end{array}$	${\begin{array}{*{20}c} 1^{0}3^{0}5^{0}15^{3}\\ 1^{0}3^{2}5^{1}15^{0}\\ 1^{3}3^{-1}5^{1}15^{0}\end{array}}$	$ 1^{0}3^{1}5^{-1}15^{3} \\ 1^{1}3^{0}5^{-1}15^{3} $	$ 1^{0}3^{1}5^{0}15^{2} \\ 1^{1}3^{0}5^{0}15^{2} $	$ 1^{0}3^{1}5^{2}15^{0} \\ 1^{1}3^{0}5^{3}15^{-1} $
$N = 15, k = 1^{0}3^{1}5^{3}15^{0}$	$2 \\ 1^0 3^3 5^1 15^0$	$1^1 3^0 5^0 15^3$	$1^3 3^0 5^0 15^1$		
$N = 17, \ k = 1$ $1^0 17^2$					
$N = 17, \ k = 3/2$ $1^{1}17^{2} \qquad 1^{2}17^{1}$					
N = 19, k = 1 $1^0 19^2$					

Table 3. (continued)

Table 3. (continued)		
$N = 19, \ k = 3/2 \\ 1^1 19^2 \qquad 1^2 19^1$		
$N = 21, \ k = 1$ $1^{0}3^{0}7^{0}21^{2}$		
N = 21, k = 3/2 1 ⁰ 3 ¹ 7 ² 21 ⁰ 1 ⁰ 3 ² 7 ¹ 21 ⁰	$1^{1}3^{0}7^{0}21^{2}$	$1^2 3^0 7^0 2 1^1$
$N = 27, \ k = 1/2$ $1^0 3^0 9^0 27^1$		
$N = 27, \ k = 1$ 1 ⁰ 3 ⁰ 9 ¹ 27 ¹ 1 ⁰ 3 ¹ 9 ⁰ 27 ¹	$1^{0}3^{2}9^{-1}27^{1}$	
$N = 27, \ k = 3/2 \\ 1^0 3^0 9^2 27^1 \qquad 1^0 3^1 9^1 27^1$	$1^{0}3^{2}9^{0}27^{1}$	
$N = 27, \ k = 2$ $1^0 3^2 9^1 27^1$		

Remark 9.5. Each entry in Table 3 is different from others. For instance, although the eta-quotient $\eta(\tau)^3$, when considered as $\eta(\tau)^3 \eta(2\tau)^0$, generates the one-dimensional space $M_{3/2}(\Gamma_0(2), \chi)$ where $\chi: \widetilde{\Gamma_0(3)} \to \mathbb{C}^{\times}$ is the character of $\eta(\tau)^3 \eta(2\tau)^0$, we list $\eta(\tau)^3$ only in the group N = 1, not N = 2, k = 3/2 to avoid repetition. As a comparison, Tables 2 and 3 of [20], whose entries constitute a subset of Table 3 above, contain repeated eta-quotients. For example, $\eta(\tau)^{-1}\eta(2\tau)^2 = \eta(\tau)^{-1}\eta(2\tau)^2\eta(4\tau)^0$ is listed both in the groups corresponding to N = 2, k = 1/2 and N = 4, k = 1/2 there.

Remark 9.6. One should notice, e.g., that the formal product $1^{0}3^{0}5^{0}15^{2}$ which represents $\eta(15\tau)^{2}$ is in Table 3 but $1^{0}3^{0}5^{0}15^{1}$ is not. This is because $\dim_{\mathbb{C}} M_{1/2}(\Gamma_{0}(15), \chi)$ can not be computed using Theorem 4.2 (with t = 0) for any χ but $\dim_{\mathbb{C}} M_{1}(\Gamma_{0}(15), \chi)$ can with χ being the character of $\eta(15\tau)^{2}$ as one may directly see by comparing both sides of the first inequality of (21). See also Table 1 and Table 2. Nevertheless, we can prove that $\eta(15\tau)$ is weakly admissible, as the following proposition shows.

Proposition 9.7. Let f be an eta-quotient (of integral exponents). If there is a positive integer n such that f^n is weakly admissible, then f is weakly admissible. As a consequence, The m-th root, where $m \in \mathbb{Z}_{\geq 1}$, of an eta-quotient $\prod \eta(n\tau)^{r_n}$ in Table 3 is weakly admissible provided that $m \mid r_n$ for all n.

Proof. Let N, k be the level and weight of f respectively. Let $\chi: \widetilde{\Gamma_0(N)} \to \mathbb{C}^{\times}$ be the character of f. Then χ^n is the character¹³ of f^n . First suppose f^n is weakly admissible of type I, which means f^n is holomorphic at all cusps and $\dim_{\mathbb{C}} M_{nk}(\Gamma_0(N), \chi^n) = 1$. Then obviously f is holomorphic at all cusps so $f \in M_k(\Gamma_0(N), \chi)$. To prove $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi) = 1$, let $0 \neq g \in M_k(\Gamma_0(N), \chi)$ be arbitrary. Then $g^n \in M_{nk}(\Gamma_0(N), \chi^n)$ and hence $g^n = c \cdot f^n$ for some $c \in \mathbb{C}^{\times}$. It follows that $g = c_1 \cdot f$ where c_1 is an n-th root of c since \mathfrak{H} is connected and f is holomorphic on \mathfrak{H} . This proves that f is weakly admissible of type I. The case of type II is proved in a similar manner with spaces such as $M_k(\Gamma_0(N), \chi)$ replaced by $S_k(\Gamma_0(N), \chi)$.

Remark 9.8. It should be noted that if f^n is weakly admissible of type II, then we can only assert that f is weakly admissible; we can say nothing about whether f is of type I or II. For instance, consider $f(\tau) = \eta(\tau), n = 24$.

9.3. Generalized double coset operators. We recall the theory of generalized double coset operators developed in [20, Section 3]. This is a generalization of Wohlfahrt's extension of Hecke operators; c.f. [19]. Let l, N be positive integers. Let χ be the character (6) where D = 2 and r_n are integers. Let $(r'_n)_{n|N}$ be another sequence of integers such that $\sum_n r_n = \sum_n r'_n$ and let χ' be the character (6) with r_n replaced by r'_n (again D = 2). We define a formal expression (c.f. [20, eq. (28) and (30)])

(52)
$$T_l f(\tau) = l^{-\frac{k}{2}} \cdot \sum_{\substack{a|l\\(N,a)=1}} a^k \sum_{\substack{0 \le b < d\\(a,b,d)=1}} \chi\left(\underbrace{\widetilde{(N,b+ax \ z)}}_{-Nd \ y}\right)^{-1} \chi'\left(\underbrace{\widetilde{(N,b+ax \ z)}}_{N \ x}\right)^{-1} f\left(\frac{a\tau + b}{d}\right)$$

where $k \in \frac{1}{2}\mathbb{Z}$, $d = \frac{l}{a}$ and x, y, z are any integers (depending on a, b and d) such that (Nd, -Nb + ax) = 1 and (-Nb + ax)y + Ndz = 1. If the dependence on k, N, χ and χ' is important, then we also denote T_l by $T_{l;k,N,\chi,\chi'}$ or $T_{l;\chi,\chi'}$.

Theorem 9.9. For f being a meromorphic modular form of weight $k \in \frac{1}{2}\mathbb{Z}$ for the group $\Gamma_0(N)$ with character χ , $T_l f$ is well-defined (i.e., independent of the choices of x, y, z) and is a meromorphic modular form of the same weight for the same group with character χ' provided that the following three conditions hold:

(53)
$$l\sum_{n|N}\frac{N}{n}r_n \equiv \sum_{n|N}\frac{N}{n}r'_n \pmod{24},$$

(54)
$$\sum_{n|N} nr_n \equiv l \sum_{n|N} nr'_n \pmod{24},$$

(55)
$$l^{2|k'|} \cdot \prod_{2 \nmid r_n - r'_n} n \text{ is a perfect square,}$$

¹³Here χ^n means the character on $\widetilde{\Gamma_0(N)}$ that sends (γ, ε) to $\chi(\gamma, \varepsilon)^n$ where $\gamma \in \Gamma_0(N)$ and $\varepsilon \in \{\pm 1\}$.

where $k' = \frac{1}{2} \sum_{n} r_{n}$. Moreover, if the above conditions hold, then T_{l} maps $M_{k}(\Gamma_{0}(N), \chi)$ into $M_{k}(\Gamma_{0}(N), \chi')$ and $S_{k}(\Gamma_{0}(N), \chi)$ into $S_{k}(\Gamma_{0}(N), \chi')$ respectively.

Proof. This is a combination of the last assertion of [20, Theorem 3.9], [20, Proposition 3.2(4),(5)] and [20, eq. (28), (30)].

Theorem 9.10. Let $f_1 = \prod_n \eta(n\tau)^{r_n}$ and $f_2 = \prod_n \eta(n\tau)^{r'_n}$ be weakly admissible eta-quotients of the same type (e.g., functions listed in Table 3) of the same weight k and the same level N. Let χ and χ' be the characters of f_1 and f_2 on $\widetilde{\Gamma_0(N)}$ respectively. Then for any $l \in \mathbb{Z}_{\geq 1}$ satisfying (53), (54) and (55) (with k' = k), there exists a $c_l \in \mathbb{C}$ such that

$$T_{l;\chi,\chi'}f_1 = c_l \cdot f_2$$

Proof. An immediate consequence of Theorem 9.9.

Remark 9.11. This theorem is still valid in the following two cases:

- f_1 is weakly admissible of type II and f_2 is weakly admissible of type I,
- f_1 is a cusp form that is weakly admissible of type I and f_2 is weakly admissible of type II.

Remark 9.12. In the case N = 1 (hence $r_1 = r'_1$, $k = \frac{1}{2}r_1$ and $\chi = \chi'$) the identity reads $T_l\eta^{r_1} = c_l \cdot \eta^{r_1}$ where $r_1 = 0, 1, 2, \ldots 24$. This has been investigated in detail in [20, Section 5] where c_l and (52) are given in explicit forms.

We are mainly interested in the case $f_1 = f_2$ in Theorem 9.10.

Corollary 9.13. Let $f(\tau) = \prod_n \eta(n\tau)^{r_n}$ be a weakly admissible eta-quotient (e.g., a function listed in Table 3) of weight $k = \frac{1}{2} \sum_n r_n$ and level N. We define a submonoid of the multiplicative monoid of positive integers as follows:

$$L_f = \begin{cases} \{l \in \mathbb{Z}_{\geq 1} \colon l \equiv 1 \mod m_f\} & \text{if } k \in \mathbb{Z}, \\ \{l \in \mathbb{Z}_{\geq 1} \colon l \equiv 1 \mod m_f \text{ and } l \text{ is a perfect square}\} & \text{if } k \in \frac{1}{2} + \mathbb{Z}, \end{cases}$$

where

$$m_f = \frac{24}{(24, \sum_n (N/n)r_n, \sum_n nr_n)}$$
where $T_f = c_f \cdot f_f$ for some $c_f \in \mathbb{C}$

Then for any $l \in L_f$ we have $T_l f = c_l \cdot f$ for some $c_l \in \mathbb{C}$.

Proof. Elementary number theory shows that $l \in L_f$ if and only if (53), (54) and (55) with $r'_n = r_n$ and k' = k hold. Therefore the assertion follows from Theorem 9.10 with $f_1 = f_2 = f$.

Remark 9.14. The monoid L_f and the integer m_f actually depend only on $\mathbf{r} = (r_n)_{n|N}$, so we sometimes write $L_{\mathbf{r}}$ and $m_{\mathbf{r}}$ instead.

In another words, any weakly admissible eta-quotient f is an eigenform for the operators T_l indexed by the infinite monoid L_f . When $k \in \mathbb{Z}$ and the character of f is induced by some Dirichlet character, then by a result of Gordon, Hughes and Newman (cf. [39, Theorem 1.64]) we have $m_f = 1$ and hence $L_f = \mathbb{Z}_{\geq 1}$. Moreover, in this case, T_l is proportional to the usual Hecke operator of index l. Consequently, the functions in Table 3 are all Hecke eigenforms for appropriate infinite operator monoids in a generalized sense.

The identities $T_l f = c_l \cdot f$ become interesting only when we can express (52) in a more explicit form, at least one where x, y, z do not appear. The following theorem gives such a formula which is key to the current topic. The reader may compare this with a previous result of Wohlfahrt [19, eq. (7|3) and (10|3)].

Theorem 9.15. Let $N \in \mathbb{Z}_{>1}$. For each $n \mid N$, let r_n be an integer and set

(56)
$$\mathbf{r} = (r_n)_{n|N}, \quad k' = \frac{1}{2} \sum_{n|N} r_n, \quad x_N = \sum_{n|N} nr_n, \quad \Pi = \prod_{n|N} (Nn^{-1})^{|r_n|}$$

Let $\chi_{\mathbf{r}} \colon \widetilde{\Gamma_0(N)} \to \mathbb{C}^{\times}$ be the character of $\prod_{n|N} \eta(n\tau)^{r_n}$, i.e. (6). Let f be a meromorphic modular form on $\Gamma_0(N)$ of weight $k \in k' + 2\mathbb{Z}$ with character $\chi_{\mathbf{r}}$ and let $l \in L_{\mathbf{r}}$. Then

(a) we have the expansions (on $\Im \tau > Y_0$)

$$f(\tau) = \sum_{n \in \frac{x_N}{24} + \mathbb{Z}} c_f(n) q^n, \quad T_l f(\tau) = \sum_{n \in \frac{x_N}{24} + \mathbb{Z}} c_{T_l f}(n) q^n,$$

where Y_0 is some nonnegative number, $c_f(n)$, $c_{T_lf}(n) \in \mathbb{C}$ are uniquely determined and T_l is the operator defined in (52) with $\chi = \chi' = \chi_{\mathbf{r}}$;

(b) for $n \in \frac{x_N}{24} + \mathbb{Z}$ we have

(57)
$$c_{T_l f}(n) = l^{-\frac{k}{2}} \sum_{\substack{a|l, d=l/a \\ (a,N)=1}} \left(\frac{a}{\Pi}\right) a^k \cdot c_f\left(\frac{ln}{a^2}\right) \sum_{\substack{0 \le b < d \\ (a,b,d)=1}} \left(\frac{-Nb}{(a,d)}\right)^{2k} \mathfrak{e}\left(bd\left(\frac{n}{l} - \frac{x_N}{24}\right)\right) \psi_{l,\mathbf{r}}(a,b)$$

where

$$\psi_{l,\mathbf{r}}(a,b) = \begin{cases} \mathfrak{e}\left(-\frac{k+\delta}{4}(d-1) + \frac{(k+\delta)(l-1)(N-1)}{4}\right) & \text{ if } 2 \nmid l, \\\\ \mathfrak{e}\left(-\frac{k+\delta}{4}(a-1) - \frac{N(k+\delta)(1+\delta_1)}{4}b\right) & \text{ if } 2 \mid l, 2 \mid N, \\\\ 1 & \text{ if } 2 \mid l, 2 \nmid N, \end{cases}$$

with $\delta = 0$ ($\delta = 1$ respectively) if Π takes the form $2^{\alpha} \cdot (4m+1)$ ($2^{\alpha} \cdot (4m+3)$ respectively), $\alpha, m \in \mathbb{Z}_{\geq 0}, \delta_1 = 1$ if $2 \mid l, 4 \mid N, k \in \frac{1}{2} + \mathbb{Z}, v_2(\Pi) \equiv 1 \mod 2$ and $\delta_1 = 0$ otherwise.

A few words of caution are in order. Note that $\left(\frac{a}{\Pi}\right)$ and $\left(\frac{-Nb}{(a,d)}\right)$ refer to Kronecker-Jacobi symbols; they are not fractions. Also note that in (57) we have set $c_f\left(\frac{ln}{a^2}\right) = 0$ if $\frac{ln}{a^2} \notin \frac{x_N}{24} + \mathbb{Z}$. In addition, k

can be replaced by k' in the definition of $\psi_{l,\mathbf{r}}(a,b)$ and if f is the eta-quotient $\prod_{n|N} \eta(n\tau)^{r_n}$ itself, which is the case we will use below, then k = k'. Finally, $Y_0 = 0$ if f is holomorphic on \mathfrak{H} .

Remark 9.16. The formula (57) is equivalent to

(58)
$$T_l f(\tau) = l^{-\frac{k}{2}} \sum_{\substack{a|l, d=l/a \\ (a,N)=1}} \left(\frac{a}{\Pi}\right) a^k \sum_{\substack{0 \le b < d \\ (a,b,d)=1}} \left(\frac{-Nb}{(a,d)}\right)^{2k} \mathfrak{e}\left(-\frac{x_N}{24}bd\right) \psi_{l,\mathbf{r}}(a,b) \cdot f\left(\frac{a\tau+b}{d}\right)$$

which can be considered as an explicit form of (52) with $\chi = \chi'$.

Proof of Theorem 9.15 and Remark 9.16. We begin with the proof of (58). Set

$$R := \chi \left(\underbrace{\stackrel{-Nb+ax}{-Nd} z}_{-Nd} \right) \chi' \left(\underbrace{\stackrel{yy}{ay} \stackrel{by-dz}{by-dz}}_{w} \right) = \chi_{\mathbf{r}} \left(\underbrace{\stackrel{(-Nb+ax)ay+Nz}{-Nady+Ny} \stackrel{(-Nb+ax)(by-dz)+xz}_{-Nd(by-dz)+xy}}_{-Nd(by-dz)+xy} \right)$$

in (52). (Theorem 9.9 and the assumption $l \in L_{\mathbf{r}}$ ensure that T_l is well-defined.) Since ad = l and

$$(59) \qquad \qquad (-Nb+ax)y+Ndz = 1$$

we have

$$R = \chi_{\mathbf{r}} \left(\underbrace{\stackrel{a-N(l-1)z \ b-(l-1)xz}{-N(l-1)y \ d-(l-1)xy}}_{-N(l-1)y \ d-(l-1)xy} \right).$$

To derive (58) from (52), we need only to prove

(60)
$$R^{-1} = \left(\frac{a}{\Pi}\right) \left(\frac{-Nb}{(a,d)}\right)^{2k} \mathfrak{e}\left(-\frac{x_N}{24}bd\right) \psi_{l,\mathbf{r}}(a,b)$$

for which we need the Petersson's formula of χ_{η} :

(61)
$$\chi_{\eta}\left(\left(\begin{smallmatrix}a&b\\c&d\end{smallmatrix}\right),\varepsilon\right) = \begin{cases} \varepsilon \cdot \left(\frac{d}{|c|}\right) \mathfrak{e}\left(\frac{1}{24}\left((a-2d)c-bd(c^2-1)+(3d-3)c\right)\right) & \text{if } 2 \nmid c, \\ \varepsilon \cdot \left(\frac{c}{d}\right) \mathfrak{e}\left(\frac{1}{24}\left((a-2d)c-bd(c^2-1)+3d-3\right)\right) & \text{if } 2 \mid c. \end{cases}$$

For a proof of (61) see [42]. Since $r_n \in \mathbb{Z}$ we have

(62)
$$R = \prod_{n|N} \chi_{\eta}^{r_n} \left(\begin{smallmatrix} a - N(l-1)z & n(b-(l-1)xz) \\ -\frac{N}{n}(l-1)y & d-(l-1)xy \end{smallmatrix} \right).$$

The strategy¹⁴ is to insert (61) into (62) and then eliminate x, y, z using properties of Kronecker-Jacobi symbols. Inserting (61) into (62) gives

$$(63) \quad R = \prod_{\substack{n|N\\2\nmid\frac{N}{n}(l-1)y}} \left(\frac{d-(l-1)xy}{Nn^{-1}(l-1)|y|}\right)^{r_n} \cdot \prod_{\substack{n|N\\2\mid\frac{N}{n}(l-1)y}} \left(\frac{-Nn^{-1}(l-1)y}{d-(l-1)xy}\right)^{r_n} \cdot \mathfrak{e}\left(\frac{x_N}{24}bd\right)$$
$$\cdot \prod_{\substack{n|N\\2\mid\frac{N}{n}(l-1)y}} \mathfrak{e}\left(\frac{r_n}{8}(d-1-(l-1)xy)(-Nn^{-1})(l-1)y\right) \cdot \prod_{\substack{n|N\\2\mid\frac{N}{n}(l-1)y}} \mathfrak{e}\left(\frac{r_n}{8}(d-1-(l-1)xy)\right)$$

¹⁴The N = 1 case has been successfully worked out in the proof of [20, Lemma 5.2]. The strategy here is essentially the one there.

where we have used the fact

(64)
$$24 \mid (l-1) \sum_{n \mid N} nr_n, \quad 24 \mid (l-1) \sum_{n \mid N} Nn^{-1}r_n$$

which are consequences of the assumption $l \in L_{\mathbf{r}}$. Now the proof splits into three cases.

The case $2 \nmid l$. We must have $2 \mid \frac{N}{n}(l-1)y$ for any $n \mid N$. Therefore

(65)
$$R = \mathfrak{e}\left(\frac{x_N}{24}bd\right) \prod_{n|N} \left(\frac{-Nn^{-1}(l-1)y}{d-(l-1)xy}\right)^{r_n} \prod_{n|N} \mathfrak{e}\left(\frac{r_n}{8}(d-1-(l-1)xy)\right).$$

Note that if $k' \in \frac{1}{2} + \mathbb{Z}$ then *l* is an odd square and hence

$$\left(\frac{-Nn^{-1}(l-1)y}{d-(l-1)xy}\right) = \left(\frac{Nn^{-1}}{d-(l-1)xy}\right) \cdot \left(\frac{-(l-1)y}{d-(l-1)xy}\right) = \left(\frac{Nn^{-1}}{d-(l-1)xy}\right) \cdot \left(\frac{y}{d}\right).$$

Inserting this into (65) we find that

(66)
$$R = \mathfrak{e}\left(\frac{x_N}{24}bd\right) \cdot \left(\frac{\Pi}{d - (l - 1)xy}\right) \left(\frac{y}{d}\right)^{2k'} \mathfrak{e}\left(\frac{k'}{4}(d - 1 - (l - 1)xy)\right).$$

If $k' \in \mathbb{Z}$, the above expression is still valid. By (59) if $k' \in \frac{1}{2} + \mathbb{Z}$ we have

(67)
$$\left(\frac{y}{d}\right) = \left(\frac{-Nb + ax}{d}\right) = \left(\frac{-Nb}{(a,d)}\right)$$

For the last equality, see [20, p. 567, line -3]. If $k' \in \mathbb{Z}$, then trivially $\left(\frac{y}{d}\right)^{2k'} = \left(\frac{-Nb}{(a,d)}\right)^{2k'}$. For the calculation of $\left(\frac{\Pi}{d-(l-1)xy}\right)$, we set

(68)
$$\Pi' = \prod_{p \mid \Pi, 2 \nmid v_p(\Pi)} p.$$

There are four cases to consider. For the first case, $\Pi' \equiv 1 \mod 4$, we have

(69)
$$\left(\frac{\Pi}{d-(l-1)xy}\right) = \left(\frac{\Pi'}{d-(l-1)xy}\right) = \left(\frac{d-(l-1)xy}{\Pi'}\right) = \left(\frac{a}{\Pi'}\right) = \left(\frac{a}{\Pi'}\right).$$

The third equality above follows from

$$\left(\frac{d-(l-1)xy}{p}\right) = \left(\frac{d-lxy+xy}{p}\right) = \left(\frac{xy}{p}\right) = \left(\frac{a}{p}\right), \quad p \mid \Pi'$$

since by (59) we have $axy \equiv 1 \mod p$ and $lxy \equiv d \mod p$. Moreover, we have

(70)
$$\mathfrak{e}\left(-\frac{k'(l-1)xy}{4}\right) = \mathfrak{e}\left(-\frac{k'(l-1)(N-1)}{4}\right)$$

(When $k' \in \mathbb{Z}$, $l \equiv 3 \mod 4$ and $2 \nmid N$, (70) does not necessarily hold, but we can always choose x, y, z such that $2 \mid y$ for which (70) holds. We always make such a choice.) Inserting (67), (69) and (70) into (66) and replacing k' by k we obtain (60) as required. For the other three cases $\Pi' \equiv 3 \mod 4$, $\Pi' \equiv 2 \mod 8$ and $\Pi' \equiv 6 \mod 8$, we proceed similarly and obtain (60) likewise. However, if $2 \mid \Pi'$ there is an extra difficulty: a factor $\left(\frac{2}{ad-(l-1)axy}\right) = \left(\frac{2}{1+(l-1)N(dz-by)}\right)$ appears and

we should prove it equals 1. This is immediate if $l \equiv 1 \mod 4$ since $2 \mid \Pi'$ implies $2 \mid N$. Now suppose $l \equiv 3 \mod 4$. Assume by contradiction that $4 \nmid N$; we have $N \equiv 2 \mod 4$. By the first relation of (64) we have $4 \mid \sum_{n} nr_n$ from which $2 \mid \sum_{2 \nmid n} r_n$. Since $2 \mid \Pi'$ we have

$$v_2(\Pi) = \sum_{n|N} v_2(N/n) \cdot |r_n| = \sum_{2 \nmid n} |r_n| \equiv 1 \mod 2$$

which contradicts $2 \mid \sum_{2 \nmid n} r_n$. Therefore, $4 \mid N$ and hence $\left(\frac{2}{1+(l-1)N(dz-by)}\right) = 1$.

The case $2 \mid l, 2 \nmid N$. By (59) if $2 \mid d$ then $2 \nmid y$ and if $2 \nmid d$ then we can choose x, y, z such that $2 \nmid y$. (If (x, y, z) is a solution to (59) with $2 \mid y$, then (x, y + Nd, z - (-Nb + ax)) is another solution with $2 \nmid y + Nd$.) We always make such a choice. Thus (63) becomes

$$R = \mathfrak{e}\left(\frac{x_N}{24}bd\right) \prod_{n|N} \left(\frac{d - (l-1)xy}{Nn^{-1}(l-1)|y|}\right)^{r_n} \prod_{n|N} \mathfrak{e}\left(\frac{r_n}{8}(d-1 - (l-1)xy)(-Nn^{-1})(l-1)y\right).$$

Since $l \equiv 1 \mod m_{\mathbf{r}}$ we have $2 \nmid m_{\mathbf{r}}$. By the definition of $m_{\mathbf{r}}$ (c.f. Remark 9.14) we have

(71)
$$8 \mid \sum_{n} nr_{n}, \quad 8 \mid \sum_{n} Nn^{-1}r_{n}$$

It follows that $k' = \frac{1}{2} \sum_{n} r_n \in \mathbb{Z}$ and

(72)
$$R = \mathfrak{e}\left(\frac{x_N}{24}bd\right) \prod_{n|N} \left(\frac{d - (l-1)xy}{Nn^{-1}(l-1)|y|}\right)^{r_n}$$

Note that

$$\left(\frac{d - (l - 1)xy}{Nn^{-1}(l - 1)|y|}\right) = \left(\frac{d - (l - 1)xy}{Nn^{-1}}\right) \cdot \left(\frac{d - (l - 1)xy}{(l - 1)|y|}\right) = \left(\frac{d - (l - 1)xy}{Nn^{-1}}\right) \cdot \left(\frac{d}{(l - 1)|y|}\right)$$

where the last equality follows from the fact (l-1)|y| is odd and positive. Inserting this and $2k' \equiv 0 \mod 2$ into (72) we obtain

$$R = \mathfrak{e}\left(\frac{x_N}{24}bd\right) \cdot \left(\frac{d - (l - 1)xy}{\Pi}\right) = \mathfrak{e}\left(\frac{x_N}{24}bd\right) \cdot \left(\frac{a}{\Pi}\right)$$

where the last equality follows just as in the deduction of (69). We thus have arrived at (60).

The case $2 \mid l, 2 \mid N$. As in the last case, we have $8 \mid \sum_{n} nr_{n}$ and $8 \mid \sum_{n} Nn^{-1}r_{n}$. Moreover, y must be odd by (59). For any $n \mid N$ with $2 \nmid \frac{N}{n}(l-1)y$, that is, with $2 \nmid \frac{N}{n}$, we have

$$\left(\frac{d-(l-1)xy}{Nn^{-1}(l-1)|y|}\right) = \left(\frac{Nn^{-1}(l-1)|y|}{d-(l-1)xy}\right) \cdot \mathfrak{e}\left(-\frac{1}{8}(d-1-(l-1)xy) \cdot (Nn^{-1}(l-1)|y|-1)\right).$$

Choosing a solution (x, y, z) to (59) with y < 0 and inserting the above identity into (63), we find that

$$R = \mathfrak{e}\left(\frac{x_N}{24}bd\right) \cdot \prod_{n|N} \left(\frac{-Nn^{-1}(l-1)y}{d-(l-1)xy}\right)^{r_n} \cdot \prod_{\substack{n|N\\2|\frac{N}{n}}} \mathfrak{e}\left(-\frac{r_n}{8}(d-1-(l-1)xy)(-Nn^{-1}(l-1)xy)(-Nn^{-1}(l-1)y\right) + \prod_{\substack{n|N\\2|\frac{N}{n}}} \mathfrak{e}\left(\frac{r_n}{8}(d-1-(l-1)xy)(-Nn^{-1})(l-1)y\right) \cdot \prod_{\substack{n|N\\2|\frac{N}{n}}} \mathfrak{e}\left(\frac{r_n}{8}(d-1-(l-1)xy)\right)$$

We deal with the factor $\prod_{n|N} \left(\frac{-Nn^{-1}(l-1)y}{d-(l-1)xy}\right)^{r_n}$ as in previous cases and thus simplify the expression for R as

(73)
$$R = \mathfrak{e}\left(\frac{x_N}{24}bd\right) \cdot \left(\frac{\Pi}{d - (l-1)xy}\right) \cdot \left(\frac{d}{-(l-1)y}\right)^{2k'} \cdot \mathfrak{e}\left(\frac{k'}{4}(d - 1 - (l-1)xy) \cdot (-(l-1)y)\right)$$

There are six subcases: (c.f. (68) for Π')

(a) 2 | l, 4 | N, k' ∈ Z and 2 ∤ Π'.
(b) 2 | l, 4 | N, k' ∈ Z and 2 | Π'.
(c) 2 | l, 4 | N, k' ∈ ¹/₂ + Z and 2 ∤ Π'.
(d) 2 | l, 4 | N, k' ∈ ¹/₂ + Z and 2 | Π'. (Only in this subcase we have δ₁ = 1.)
(e) 2 | l, N ≡ 2 mod 4 and 2 ∤ Π'. (We must have k' ∈ Z in this subcase.)
(f) 2 | l, N ≡ 2 mod 4 and 2 | Π'. (We must have k' ∈ Z in this subcase.)

We will only give the proof of the more difficult subcases (b) and (d) and omit the other four since the strategies are the same. For (b), we first prove $8 \mid N$. Assume by contradiction that $8 \nmid N$; then $2^2 \parallel N$. It follows that

$$v_2(\Pi) \equiv \sum_{\substack{n|N\\v_2(n)=1}} v_2(N/n) \cdot |r_n| = \sum_{\substack{n|N\\v_2(n)=1}} |r_n| \equiv 1 \mod 2.$$

On the other hand, (71) still holds in this case, and hence

$$\sum_{\substack{n|N\\v_2(n)=0}} r_n \equiv \sum_{\substack{n|N\\v_2(n)=2}} r_n \equiv 0 \mod 2.$$

Therefore $\sum_{n|N} r_n \equiv 1 \mod 2$, that is, $k' \in \frac{1}{2} + \mathbb{Z}$, which contradicts the assumption in (b). Hence $8 \mid N$. Note that both $\Pi'/2$ and d - (l-1)xy are odd and positive, from which we deduce that

$$\begin{pmatrix} \overline{\Pi} \\ \overline{d - (l-1)xy} \end{pmatrix} = \begin{pmatrix} \overline{\Pi'} \\ \overline{d - (l-1)xy} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{2}{d - (l-1)xy} \end{pmatrix} \begin{pmatrix} \frac{d - (l-1)xy}{\Pi'/2} \end{pmatrix} \mathfrak{e} \begin{pmatrix} \frac{1}{4}(d - 1 - (l-1)xy) \cdot \frac{\Pi'/2 - 1}{2} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{2}{ad - (l-1)axy} \end{pmatrix} \begin{pmatrix} \frac{a}{\Pi'} \end{pmatrix} \mathfrak{e} \begin{pmatrix} \frac{1}{4}(d - 1 - (l-1)xy) \cdot \frac{\Pi'/2 - 1}{2} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{2}{1 + (l-1)N(dz - by)} \end{pmatrix} \begin{pmatrix} \frac{a}{\Pi} \end{pmatrix} \mathfrak{e} \begin{pmatrix} \frac{1}{4}(d - 1 - (l-1)xy) \cdot \frac{\Pi'/2 - 1}{2} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{a}{\Pi} \end{pmatrix} \mathfrak{e} \begin{pmatrix} \frac{1}{4}(a - 1) \cdot \frac{\Pi'/2 - 1}{2} \end{pmatrix}.$$

$$(74)$$

The last equality holds since by (59) and 8 | N we have $axy \equiv 1 \mod 4$ and hence

$$(a-1) - (d-1 - (l-1)xy) \equiv a - d + (l-1)a = (a^2 - 1)d \equiv 0 \mod 4.$$

An argument similar to the above deduction shows that

$$\mathfrak{e}\left(\frac{k'}{4}(d-1-(l-1)xy)\cdot(-(l-1)y)\right) = \mathfrak{e}\left(\frac{k'}{4}(a-1)(l-1)\right)$$

Inserting this and (74) into (73) and noting that a is odd, we obtain (60) in the subcase (b). For the subcase (d), l is a square. Since (N, a) = 1 and ad = l we have $v_2(d) = v_2(l) \equiv 0 \mod 2$ and hence $4 \mid d$. Thus if we let $d' = d/2^{v_2(d)}$ then

$$\left(\frac{d}{-(l-1)y}\right) = \left(\frac{d}{y}\right) = \left(\frac{d'}{y}\right) = \mathfrak{e}\left(\frac{y-1}{4} \cdot \frac{d'-1}{2}\right) \cdot \left(\frac{y}{d}\right).$$

Combining this and

$$\left(\frac{-Nb+ax}{d}\right)\left(\frac{y}{d}\right) = \left(\frac{(-Nb+ax)y}{d}\right) = \left(\frac{1-Ndz}{d}\right) = 1$$

we find that

(75)
$$\left(\frac{d}{-(l-1)y}\right) = \mathfrak{e}\left(\frac{y-1}{4} \cdot \frac{d'-1}{2}\right) \cdot \left(\frac{-Nb+ax}{d}\right) = \mathfrak{e}\left(\frac{y-1}{4} \cdot \frac{d'-1}{2}\right) \cdot \left(\frac{-Nb}{(a,d)}\right).$$

The calculation of $\left(\frac{\Pi}{d-(l-1)xy}\right)$ is similar to that in above cases (e.g. (69) or (74)) and the result is

(76)
$$\left(\frac{\Pi}{d-(l-1)xy}\right) = \left(\frac{2}{1+Nb}\right) \left(\frac{a}{\Pi}\right) \mathfrak{e} \left(\frac{1}{4}(l-1)(a-1) \cdot \frac{\Pi'/2 - 1}{2}\right).$$

For the factor $\mathfrak{e}\left(\frac{k'}{4}(d-1-(l-1)xy)\cdot(-(l-1)y)\right)$ in (73), note that $l-d=(a-1)d\equiv 0 \mod 8$, $2 \nmid y$ and hence

$$\mathfrak{e}\left(\frac{k'}{4}(d-1-(l-1)xy)\cdot(-(l-1)y)\right) = \mathfrak{e}\left(\frac{k'}{4}(l-1)^2(1-xy)(-y)\right)$$
$$= \mathfrak{e}\left(\frac{k'}{4}(1-xy)(-y)\right) = \mathfrak{e}\left(\frac{k'}{4}(x-y)\right).$$

Inserting this equality, (75) and (76) into (73) we obtain

(77)
$$R = \left(\frac{a}{\Pi}\right) \left(\frac{-Nb}{(a,d)}\right) \mathfrak{e}\left(\frac{x_N}{24}bd\right) \left(\frac{2}{1+Nb}\right) \\ \cdot \mathfrak{e}\left(\frac{1}{4}(l-1)(a-1) \cdot \frac{\Pi'/2 - 1}{2}\right) \mathfrak{e}\left(\frac{y-1}{4} \cdot \frac{d'-1}{2} + \frac{k'}{4}(x-y)\right).$$

Now we simplify the factor $\mathfrak{e}\left(\frac{y-1}{4} \cdot \frac{d'-1}{2} + \frac{k'}{4}(x-y)\right)$ in above. By (59) and the facts $2 \mid d, 4 \mid N$ we find that $(-Nb + ax)y \equiv 1 \mod 8$. This, together with $2 \nmid y$, implies $x \equiv ay + Nab \mod 8$. It follows that

(78)
$$\mathbf{\mathfrak{e}}\left(\frac{y-1}{4}\cdot\frac{d'-1}{2}+\frac{k'}{4}(x-y)\right) = \mathbf{\mathfrak{e}}\left(\frac{-2k'}{4}(y-1)\cdot\frac{d'-1}{2}+\frac{k'}{4}(ay-y+Nab)\right)$$
$$= \mathbf{\mathfrak{e}}\left(\frac{k'}{4}Nab\right)\mathbf{\mathfrak{e}}\left(\frac{k'}{4}y(a-d')\right)\mathbf{\mathfrak{e}}\left(\frac{k'}{4}(d'-1)\right).$$

Since $ad' = l/2^{v_2(d)} = l/2^{v_2(l)}$ is a square, $2 \nmid a$ and $2 \nmid d'$ we have

(79)
$$a - d' \equiv 0 \mod 8$$

In addition, since $k' \in \frac{1}{2} + \mathbb{Z}$, $2 \nmid a$ and $4 \mid N$,

(80)
$$\mathbf{\mathfrak{e}}\left(\frac{k'}{4}Nab\right) = \mathbf{\mathfrak{e}}\left(\frac{1}{8}Nab\right) = \mathbf{\mathfrak{e}}\left(\frac{1}{8}Nb\right)$$

Inserting (79) and (80) into (78), then combining (77) and noting that $\mathfrak{e}\left(\frac{1}{8}Nb\right)\left(\frac{2}{1+Nb}\right) = 1$, we arrive at (60) as required. This concludes the proof of (58).

Finally, we begin to prove Theorem 9.15. For Theorem 9.15(a), elementary Fourier analysis shows that

$$f(\tau) = \sum_{n \in \frac{1}{24}\mathbb{Z}} c_f(n)q^n, \quad T_l f(\tau) = \sum_{n \in \frac{1}{24}\mathbb{Z}} c_{T_l f}(n)q^n,$$

which converge on $\Im \tau > Y_0$ for some $Y_0 \ge 0$ since $f(\tau + 24) = f(\tau)$ and $T_l f(\tau + 24) = T_l f(\tau)$. By modularity $f|_{k}(\underbrace{1 \ 1}_{0 \ 1}) = \chi_{\mathbf{r}}(\underbrace{1 \ 1}_{0 \ 1}) \cdot f = \mathfrak{e}(\underbrace{x_N}{24}) \cdot f$ and $T_l f|_{k}(\underbrace{1 \ 1}_{0 \ 1}) = \chi_{\mathbf{r}}(\underbrace{1 \ 1}_{0 \ 1}) \cdot T_l f = \mathfrak{e}(\underbrace{x_N}{24}) \cdot T_l f$. In terms of Fourier coefficients,

$$c_f(n)\mathbf{\mathfrak{e}}(n) = c_f(n)\mathbf{\mathfrak{e}}\left(\frac{x_N}{24}\right), \quad c_{T_lf}(n)\mathbf{\mathfrak{e}}(n) = c_{T_lf}(n)\mathbf{\mathfrak{e}}\left(\frac{x_N}{24}\right).$$

It follows that $c_f(n) = 0$ unless $n \in \frac{x_N}{24} + \mathbb{Z}$ and so is $c_{T_lf}(n)$. Theorem 9.15(b), that is, (57), follows immediately from (58) whose details are omitted.

In the important case $x_N \equiv 0 \mod 24$ we have $c_f(n) = 0$ and $c_{T_lf}(n) = 0$ unless $n \in \mathbb{Z}$. Therefore, (57) becomes

$$(81) \qquad c_{T_lf}(n) = l^{-\frac{k}{2}} \sum_{\substack{a|l, a^2|ln\\(a,N)=1, d=l/a}} \left(\frac{a}{\Pi}\right) a^k \cdot c_f\left(\frac{ln}{a^2}\right) \sum_{\substack{0 \le b < d\\(a,b,d)=1}} \left(\frac{-Nb}{(a,d)}\right)^{2k} \mathfrak{e}\left(\frac{bdn}{l}\right) \psi_{l,\mathbf{r}}(a,b)$$

for $n \in \mathbb{Z}$. From this some elegant formulas for special cases follow:

Corollary 9.17. We keep the notations and assumptions of Theorem 9.15 and assume that $x_N \equiv 0 \mod 24$. Then

$$c_{T_l f}(0) = c_f(0) \cdot l^{-\frac{k}{2}} \sum_{\substack{a|l, d=l/a \\ (a,N)=1}} \left(\frac{a}{\Pi}\right) a^k \sum_{\substack{0 \le b < d \\ (a,b,d)=1}} \left(\frac{-Nb}{(a,d)}\right)^{2k} \psi_{l,\mathbf{r}}(a,b).$$

Moreover, if for any prime p not dividing N we have $p^2 \nmid l$ (e.g. l is square-free or $rad(l) \mid rad(N)$), then

(82)
$$c_{T_l f}(1) = c_f(l) \cdot l^{-\frac{k}{2}} \sum_{0 \le b < l} \psi_{l, \mathbf{r}}(1, b)$$

Finally, if $l = p^{\alpha}$ where p is a prime not dividing N and $\alpha \in \mathbb{Z}_{\geq 1}$, then

$$c_{T_{p^{\alpha}f}}(1) = p^{-\frac{\alpha k}{2}} \sum_{0 \le \beta \le [\alpha/2]} \left(\frac{p}{\Pi}\right)^{\beta} p^{k\beta} \cdot c_f(p^{\alpha-2\beta}) \sum_{\substack{0 \le b < p^{\alpha-\beta} \\ (p^{\beta},b)=1}} \left(\frac{-Nb}{p^{\beta}}\right)^{2k} \mathfrak{e}\left(\frac{b}{p^{\beta}}\right) \psi_{l,\mathbf{r}}(p^{\beta},b) = 0$$

Remark 9.18. The sum over b in the above formulas, (81) or (57) is either a partial sum of a geometric sequence or a Gauss sum according to whether k is integral or half-integral. It is not hard to derive an explicit formula for this sum but we exclude this task here. For the N = 1 case the reader may consult [20, eq. (58)] and [20, Lemma 5.6] for such formulas.

9.4. Identities involving Fourier coefficients of weakly admissible eta-quotients. The main theorem of Section 9 is the following:

Theorem 9.19. Let $f(\tau) = \prod_{n|N} \eta(n\tau)^{r_n} = \sum_n c_f(n)q^n$ be a weakly admissible eta-quotient (e.g., a function listed in Table 3) where N is the level. We keep the notations in (56) with k' = k. Then for any $l \in L_f$ (c.f. Corollary 9.13) there exists a unique $c_l \in \mathbb{C}$ such that

$$(83) \qquad l^{-\frac{k}{2}} \sum_{\substack{a|l,d=l/a\\(a,N)=1}} \left(\frac{a}{\Pi}\right) a^k \cdot c_f\left(\frac{ln}{a^2}\right) \sum_{\substack{0 \le b < d\\(a,b,d)=1}} \left(\frac{-Nb}{(a,d)}\right)^{2k} \mathfrak{e}\left(bd\left(\frac{n}{l} - \frac{x_N}{24}\right)\right) \psi_{l,\mathbf{r}}(a,b) = c_l \cdot c_f(n)$$

holds for any $n \in \frac{x_N}{24} + \mathbb{Z}$. Moreover, the number c_l is equal to the left-hand side of the above identity with n replaced by $\frac{x_N}{24}$.
Proof. By Corollary 9.13 $T_l f = c_l \cdot f$ and hence $c_{T_l f}(n) = c_l \cdot c_f(n)$ for any $n \in \frac{x_N}{24} + \mathbb{Z}$. Inserting (57) into this identity gives the desired formula. Letting $n = \frac{x_N}{24}$ and noting that $c_f\left(\frac{x_N}{24}\right) = 1$ we obtain the formula for c_l .

The identities (83) have been verified numerically for $l \leq 121$, $n - \frac{x_N}{24} \leq 40$ and f being any function listed in Table 3 by a SageMath program. See Appendix A for the code. The rest of this section is devoted to useful consequences of Theorem 9.19 and concrete examples.

9.4.1. Closed formulas for Fourier coefficients of eta-quotients. Let $f(\tau) = \prod_{n|N} \eta(n\tau)^{r_n}$ be a weakly admissible eta-quotient (e.g., a function listed in Table 3) where N is the level. We keep the notations in (56) with k' = k. Here let us consider the situation $x_N = 0$ where we have

$$f(\tau) = \prod_{n|N} \prod_{m \in \mathbb{Z}_{\geq 1}} (1 - q^{nm})^{r_n} = \sum_{n \in \mathbb{Z}_{\geq 0}} c_f(n) q^n, \quad \tau \in \mathfrak{H}.$$

We have given a series expression of $c_f(n)$ involving Dedekind sums for N = 4, $r_1 = 2$, $r_2 = 7$, $r_4 = -4$ in Example 8.10. This example is of course also listed in Table 3. In fact, there exist very simple closed formulas for $c_f(n)$ only involving Kronecker-Jacobi symbols at least for n square-free—thanks to Theorem 9.19.

Theorem 9.20. Let the notations and assumptions be as above. Let $l \in L_f$ (c.f. Corollary 9.13) satisfy

(84)
$$p \nmid N \implies p^2 \nmid l$$
, for any prime p.

(e.g. l is square-free or rad(l) | rad(N).) We require that at least one of l and N is odd. Then

$$c_{f}(l) = \begin{cases} -r_{1} \cdot \sum_{a \mid l, (a,N)=1} \left(\frac{a}{\Pi}\right) a^{k-1} & \text{if } 2 \mid l, 2 \nmid N, \\ -r_{1} \cdot \sum_{a \mid l, (a,N)=1} \left(\frac{a}{\Pi}\right) a^{k-1} \mathfrak{e}\left(\frac{(k+\delta)(l-d)}{4}\right) & \text{if } 2 \nmid l, \end{cases}$$

where d = l/a, $\Pi = \prod_{n|N} (Nn^{-1})^{|r_n|}$ and $\delta = 0$ ($\delta = 1$ respectively) if Π takes the form $2^{\alpha} \cdot (4m+1)$ ($2^{\alpha} \cdot (4m+3)$ respectively). In particular, if rad(l) | rad(N), then $c_f(l) = -r_1$.

Proof. A combination of (82) and (83) (with n = 1) gives $c_f(l) \cdot \sum_{0 \le b < l} \psi_{l,\mathbf{r}}(1,b) = l^{\frac{k}{2}} c_l \cdot c_f(1)$. By the last assertion of Theorem 9.19 we have

$$c_{l} = l^{-\frac{k}{2}} \sum_{a|l, (a,N)=1} \left(\frac{a}{\Pi}\right) a^{k} \cdot c_{f}(0) \sum_{0 \le b < d} \psi_{l,\mathbf{r}}(a,b).$$

Since $c_f(0) = 1$ and $c_f(1) = -r_1$ we find that

(85)
$$c_f(l) \cdot \sum_{0 \le b < l} \psi_{l,\mathbf{r}}(1,b) = -r_1 \cdot \sum_{a|l, (a,N)=1} \left(\frac{a}{\Pi}\right) a^k \cdot \sum_{0 \le b < d} \psi_{l,\mathbf{r}}(a,b).$$

The assumption at least one of l and N is odd ensures that $\sum_{0 \le b < l} \psi_{l,\mathbf{r}}(1,b) \ne 0$. Therefore the desired formulas follow by expanding $\sum_{0 \le b < l} \psi_{l,\mathbf{r}}(1,b)$ and $\sum_{0 \le b < d} \psi_{l,\mathbf{r}}(a,b)$ in (85).

Remark 9.21. The formulas for $c_f(l)$ in the above theorem seem like coefficients of Eisenstein series. This is actually the case since $x_N = 0$ implies f is not a cusp form. Since f is weakly admissible, then it must be of type I, that is, $\dim_{\mathbb{C}} M_k(\Gamma_0(N), \chi_{\mathbf{r}}) = 1$. Therefore f is in the Eisenstein space.

Remark 9.22. In the case of half-integral weights, Theorem 9.20 can be used only for those l with rad(l) | rad(N). However, there is no half-integral weight eta-quotient f in Table 3 such that $c_f(l)$ (l > 1) can be computed using Theorem 9.20. Nevertheless, it is still worthwhile to include the case of half-integral weights in Theorem 9.20 since there may exist weakly admissible eta-quotients not listed in Table 3 whose coefficients $c_f(l)$ can be computed using this theorem.

Example 9.23. Let us consider an example of level 12 which is not listed in Table 3. The reader may use SageMath programs described in Appendix A to produce this eta-quotient and check the assertions given here. Set

$$f(\tau) = \eta(\tau)^1 \eta(2\tau)^{-1} \eta(3\tau)^{-1} \eta(4\tau)^1 \eta(6\tau)^4 \eta(12\tau)^{-2} = \sum_{n \in \mathbb{Z}_{\geq 0}} c_f(n) q^n.$$

Let $(x_c)_{c|12}$ be as in Proposition 5.1. Hence $\operatorname{div}_{a/c} f = \frac{x_c}{24}$ for $c \mid 12$ and a coprime to c; c.f. (14). We have

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_6 \\ x_{12} \end{pmatrix} = \begin{pmatrix} 12 & 6 & 4 & 3 & 2 & 1 \\ 3 & 6 & 1 & 3 & 2 & 1 \\ 4 & 2 & 12 & 1 & 6 & 3 \\ 3 & 6 & 1 & 12 & 2 & 4 \\ 1 & 2 & 3 & 1 & 6 & 3 \\ 1 & 2 & 3 & 4 & 6 & 12 \end{pmatrix} \begin{pmatrix} r_1 \\ r_2 \\ r_3 \\ r_4 \\ r_6 \\ r_{12} \end{pmatrix} = \begin{pmatrix} 11 \\ 5 \\ 9 \\ 8 \\ 15 \\ 0 \end{pmatrix},$$

from which we see that f is a holomorphic eta-quotient. Moreover, f is admissible of type I. Thus Theorem 9.20 can be applied to f. Note that m_f in Corollary 9.13 is equal to 24 and hence

$$L_f = \{ l \in \mathbb{Z}_{\geq 1} \colon l \equiv 1 \bmod 24 \}.$$

Therefore (a, 12) = 1 for any $a \mid l$ where $l \in L_f$. Moreover,

$$k = 1$$
, $\Pi = 2^9 \cdot 3^3 = \text{square} \cdot 6$, $\delta = 1$.

It follows from Theorem 9.20 that

$$c_f(l) = -\sum_{a|l} \left(\frac{a}{6}\right) = -\prod_{p|l} \left(1 + \left(\frac{p}{6}\right)\right)$$

where $1 \leq l \equiv 1 \mod 24$ is square-free. Consequently, for such l, $-c_f(l)$ is either zero or a power of 2. Moreover, for any $\alpha \in \mathbb{Z}_{\geq 1}$ there exist infinitely many l such that $c_f(l) = -2^{\alpha}$.

9.4.2. Multiplicativity of weakly admissible eta-quotients. The Fourier coefficients $c_f(n)$ of an etaquotient f in Martin's list [17, Table I] satisfy $c_f(nm)c_f(1) = c_f(n)c_f(m)$ whenever n, m are coprime positive integers. This is the reason they are called multiplicative eta-quotients. The proper generalization to weakly admissible eta-quotients is stated in the following theorem.

Theorem 9.24. Let us keep the notations and assumptions of Theorem 9.19. Then the map $L_f \to \mathbb{C}$ that sends l to c_l is a multiplicative function, that is, we have $c_{l_1l_2} = c_{l_1} \cdot c_{l_2}$ for any $l_1, l_2 \in L_f$ with¹⁵ $(l_1, l_2) = 1$.

This theorem is an immediate consequence of the following general fact which in the special case N = 1 is due to Wohlfahrt [19].

Proposition 9.25. Let $N \in \mathbb{Z}_{\geq 1}$ and χ_1, χ_2, χ_3 be complex linear characters on the double cover $\widetilde{\Gamma_0(N)}$. Let $l_1, l_2 \in \mathbb{Z}_{\geq 1}$ such that χ_1 and χ_2 are $(1 \ 0 \ 0 \ l_1)$ -compatible, χ_2 and χ_3 are $(1 \ 0 \ l_2)$ -compatible. Then χ_1 and χ_3 are $(1 \ 0 \ l_1 \ l_2)$ -compatible. Moreover, if $(l_1, l_2) = 1$ then

(86)
$$T_{l_2;\chi_2,\chi_3} \circ T_{l_1;\chi_1,\chi_2} f = T_{l_1 l_2;\chi_1,\chi_3} f$$

for any meromorphic modular form f on $\Gamma_0(N)$ of weight $k \in \frac{1}{2}\mathbb{Z}$ and with character χ_1 .

For the concept of compatibility of multiplier systems, see [20, Lemma 3.1] for details. We say χ and χ' are $\widetilde{\binom{10}{0l}}$ -compatible if for any $\gamma \in \widetilde{\binom{10}{0l}}^{-1}\Gamma_0(N)\widetilde{\binom{10}{0l}} \cap \Gamma_0(N)$ we have $\chi\left(\widetilde{\binom{10}{0l}}\gamma\widetilde{\binom{10}{0l}}^{-1}\right) = \chi'(\gamma)$.

Proof of Proposition 9.25 and Theorem 9.24. First let us prove χ_1 and χ_3 are $(1 \ 0 \ l_1 l_2)$ -compatible. Let $g \in (1 \ 0 \ l_1 l_2)^{-1} \widetilde{\Gamma_0(N)} (1 \ 0 \ l_1 l_2) \cap \widetilde{\Gamma_0(N)}$ be arbitrary. We have

$$\chi_1\left(\left(\widetilde{\left(\begin{smallmatrix}1&0\\0&l_1l_2\end{smallmatrix}\right)}g\left(\widetilde{\left(\begin{smallmatrix}1&0\\0&l_1l_2\end{smallmatrix}\right)}^{-1}\right)\right) = \chi_1\left(\left(\widetilde{\left(\begin{smallmatrix}1&0\\0&l_1\end{smallmatrix}\right)}\left(\widetilde{\left(\begin{smallmatrix}1&0\\0&l_2\end{smallmatrix}\right)}g\left(\widetilde{\left(\begin{smallmatrix}1&0\\0&l_2\end{smallmatrix}\right)}^{-1}\right)\left(\widetilde{\left(\begin{smallmatrix}1&0\\0&l_2\end{smallmatrix}\right)}^{-1}\right)\right)$$
$$= \chi_2\left(\left(\widetilde{\left(\begin{smallmatrix}1&0\\0&l_2\end{smallmatrix}\right)}g\left(\widetilde{\left(\begin{smallmatrix}1&0\\0&l_2\end{smallmatrix}\right)}^{-1}\right)\right)$$
$$= \chi_3(g).$$

In the second equality we have used the assumption χ_1 and χ_2 are $\overbrace{\begin{pmatrix} 1 & 0 \\ 0 & l_1 \end{pmatrix}}^{0}$ -compatible and in the third equality the assumption χ_2 and χ_3 are $\overbrace{\begin{pmatrix} 1 & 0 \\ 0 & l_2 \end{pmatrix}}^{0}$ -compatible. This proves the $\overbrace{\begin{pmatrix} 1 & 0 \\ 0 & l_1 l_2 \end{pmatrix}}^{0}$ -compatibility of χ_1 and χ_3 .

¹⁵The gcd (l_1, l_2) is the usual greatest common divisor in the monoid of positive integers, not in the monoid L_f . These two concepts are different. For instance in the monoid $L = \{l \in \mathbb{Z}_{\geq 1} : l \equiv 1 \mod 24\}, (5 \cdot 29, 5 \cdot 53) = 5$ in the usual sense, but the unique common divisor of $5 \cdot 29$ and $5 \cdot 53$ in L is 1.

Assume that $(l_1, l_2) = 1$. To prove (86) we need an equivalent definition of $T_l f$ via double coset actions. For the proof of this equivalence see [20, eq. (28) and (30)]. Write

$$\widetilde{\Gamma_0(N)} \underbrace{\left(\begin{smallmatrix} 1 & 0 \\ 0 & l_i \end{smallmatrix}\right)}_{\gamma \in A_i} \widetilde{\Gamma_0(N)} = \bigsqcup_{\gamma \in A_i} \widetilde{\Gamma_0(N)} \widetilde{\gamma}$$

for i = 1, 2. Then

$$T_{l_1;\chi_1,\chi_2}f = \sum_{\gamma \in A_1} (\chi_1|_{l_1}\chi_2)^{-1}(\widetilde{\gamma}) \cdot f|_k \widetilde{\gamma},$$

where $\chi_1|_{l_1}\chi_2(h) = \chi_1(h_1)\chi_2(h_2)$ if we write $h = h_1(\overbrace{0\ l_1}^{n})h_2$ with $h_1, h_2 \in \widetilde{\Gamma_0(N)}$. The compatibility ensures that $\chi_1|_{l_1}\chi_2(h)$ is well-defined. Therefore

(87)
$$T_{l_2;\chi_2,\chi_3} \circ T_{l_1;\chi_1,\chi_2} f = \sum_{\gamma_1 \in A_1, \gamma_2 \in A_2} (\chi_2|_{l_2}\chi_3)^{-1} (\widetilde{\gamma_2}) (\chi_1|_{l_1}\chi_2)^{-1} (\widetilde{\gamma_1}) \cdot f|_k \widetilde{\gamma_1} \widetilde{\gamma_2}.$$

By [20, eq. (25)] with m = 1 we can choose the following A_i :

(88)
$$A_i = \left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} : a, d > 0, \ 0 \le b < d, \ ad = l_i, \ (N, a) = 1, \ (a, b, d) = 1 \right\}.$$

It follows from the assumption $(l_1, l_2) = 1$ and elementary number theory that $\tilde{\gamma}_1 \tilde{\gamma}_2 \in \widetilde{\Gamma_0(N)} \left(\begin{smallmatrix} 1 & 0 \\ 0 & l_1 l_2 \end{smallmatrix} \right) \widetilde{\Gamma_0(N)}$ if $\gamma_1 \in A_1$ and $\gamma_2 \in A_2$. Therefore the map

$$\sigma \colon A_1 \times A_2 \to \widetilde{\Gamma_0(N)} \backslash \widetilde{\Gamma_0(N)} \left(\underbrace{\begin{smallmatrix} 1 & 0 \\ 0 & l_1 l_2 \end{smallmatrix}}_{(\gamma_1, \gamma_2)} \right) \widetilde{\Gamma_0(N)} \cdot \widetilde{\gamma_1} \widetilde{\gamma_2}$$

is well-defined. By the theory of classical Hecke algebras, or more precisely, by Theorem 2.8.1(2) and Eq. (2.7.2) of [35], σ is a surjection and each inverse image $\sigma^{-1}(x)$ has the same cardinality. Since $\sigma^{-1}\left(\widetilde{\Gamma_0(N)}\left(\overset{1}{0}\overset{0}{l_1l_2}\right)\right) = \left\{\left(\begin{pmatrix}1 & 0\\ 0 & l_1\end{pmatrix}, \begin{pmatrix}1 & 0\\ 0 & l_2\end{pmatrix}\right)\right\}$ is a singleton, σ is a bijection which means the range of summation in the right-hand side of (87) is essentially a set of representatives of the coset space $\widetilde{\Gamma_0(N)}\setminus\widetilde{\Gamma_0(N)}\left(\overset{1}{0}\overset{0}{l_1l_2}\right)\widetilde{\Gamma_0(N)}$. Now we can define a map, which is key to the proof,

$$\xi \colon \widetilde{\Gamma_0(N)} \left(\underbrace{\begin{smallmatrix} 1 & 0 \\ 0 & l_1 l_2 \end{smallmatrix}}_{1 & l_1 l_2} \right) \widetilde{\Gamma_0(N)} \to \mathbb{C}$$
$$x \mapsto \chi_1 |_{l_1} \chi_2(x \widetilde{\gamma}_2^{-1}) \cdot \chi_2 |_{l_2} \chi_3(\widetilde{\gamma}_2)$$

where (γ_1, γ_2) is the uniquely determined pair in $A_1 \times A_2$ such that $\Gamma_0(N)x = \Gamma_0(N)\widetilde{\gamma_1}\widetilde{\gamma_2}$. It is not hard to see that

(89)
$$\xi(gxh) = \chi_1(g)\xi(x)\chi_3(h), \quad g,h \in \widetilde{\Gamma_0(N)}, x \in \widetilde{\Gamma_0(N)}(\underbrace{1 \ 0 \ 0 \ l_1 l_2}_{0 \ l_1 l_2})\widetilde{\Gamma_0(N)}, \quad \xi(\underbrace{1 \ 0 \ 0 \ l_1 l_2}_{0 \ l_1 l_2}) = 1$$

and that

(90)
$$\xi(\widetilde{\gamma}_1\widetilde{\gamma}_2) = \chi_1|_{l_1}\chi_2(\widetilde{\gamma}_1)\chi_2|_{l_2}\chi_3(\widetilde{\gamma}_2).$$

Since the map $x \mapsto \chi_1|_{l_1 l_2} \chi_3(x)$ also satisfies (89), we have $\xi = \chi_1|_{l_1 l_2} \chi_3$. Inserting this and (90) into (87) gives

$$T_{l_2;\chi_2,\chi_3} \circ T_{l_1;\chi_1,\chi_2} f = \sum_{\gamma_1 \in A_1, \gamma_2 \in A_2} \xi^{-1}(\widetilde{\gamma_1}\widetilde{\gamma_2}) \cdot f|_k \widetilde{\gamma_1}\widetilde{\gamma_2} = \sum_{\gamma_1 \in A_1, \gamma_2 \in A_2} (\chi_1|_{l_1l_2}\chi_3)^{-1}(\widetilde{\gamma_1}\widetilde{\gamma_2}) \cdot f|_k \widetilde{\gamma_1}\widetilde{\gamma_2}$$

Since σ is a bijection, we obtain (86) from the above equalities.

Finally we prove Theorem 9.24. Let f be as in this theorem. Then $T_{l_1}f = c_{l_1} \cdot f$, $T_{l_2}f = c_{l_2} \cdot f$ and $T_{l_1l_2}f = c_{l_1l_2} \cdot f$ by Corollary 9.13. Since $(l_1, l_2) = 1$ we have

$$c_{l_1 l_2} \cdot f = T_{l_1 l_2} f = T_{l_2}(T_{l_1} f) = T_{l_2}(c_{l_1} \cdot f) = c_{l_1} c_{l_2} \cdot f$$

where we have used Proposition 9.25 in the second equality. Since f is not identically zero we find $c_{l_1l_2} = c_{l_1} \cdot c_{l_2}$ as required.

Before looking at examples, let us simplify some notations. Set $x'_N = \frac{x_N}{24}$ and

$$S_{l,\mathbf{r}}(a) = \sum_{\substack{0 \le b < d \\ (a,b,d)=1}} \left(\frac{-Nb}{(a,d)}\right)^{2k} \mathfrak{e}\left(-\frac{(l-1)x'_N b}{a}\right) \psi_{l,\mathbf{r}}(a,b).$$

Then according to Theorem 9.19 we have

$$c_l = l^{-\frac{k}{2}} \sum_{a|l, (a,N)=1} \left(\frac{a}{\Pi}\right) a^k \cdot c_f \left(\frac{lx'_N}{a^2}\right) S_{l,\mathbf{r}}(a), \quad l \in L_f.$$

We recall that here and below f must be a weakly admissible eta-quotient.

Example 9.26. If $x'_N = 0$, that is, $x_N = \sum_{n|N} nr_n = 0$, then $c_f(0) = 1$ and hence Theorem 9.24 gives

$$\sum_{a|l_1,(a,N)=1} \left(\frac{a}{\overline{\Pi}}\right) a^k \cdot S_{l_1,\mathbf{r}}(a) \cdot \sum_{a|l_2,(a,N)=1} \left(\frac{a}{\overline{\Pi}}\right) a^k \cdot S_{l_2,\mathbf{r}}(a) = \sum_{a|l_1l_2,(a,N)=1} \left(\frac{a}{\overline{\Pi}}\right) a^k \cdot S_{l_1l_2,\mathbf{r}}(a)$$

for $l_1, l_2 \in L_f$ with $(l_1, l_2) = 1$.

Example 9.27. If $x'_N \in \mathbb{Z}_{\geq 0}$ and $(l, x'_N) = 1$, then $c_f\left(\frac{lx'_N}{a^2}\right) = 0$ unless $\frac{lx'_N}{a^2} \in \mathbb{Z}$ by Theorem 9.15(a). Therefore we have

$$\sum_{a^{2}|l_{1},(a,N)=1} \left(\frac{a}{\Pi}\right) a^{k} \cdot c_{f} \left(\frac{l_{1}x'_{N}}{a^{2}}\right) S_{l_{1},\mathbf{r}}(a) \cdot \sum_{a^{2}|l_{2},(a,N)=1} \left(\frac{a}{\Pi}\right) a^{k} \cdot c_{f} \left(\frac{l_{2}x'_{N}}{a^{2}}\right) S_{l_{2},\mathbf{r}}(a)$$
$$= \sum_{a^{2}|l_{1}l_{2},(a,N)=1} \left(\frac{a}{\Pi}\right) a^{k} \cdot c_{f} \left(\frac{l_{1}l_{2}x'_{N}}{a^{2}}\right) S_{l_{1}l_{2},\mathbf{r}}(a)$$

if $l_1, l_2 \in L_f$ with $(l_1, l_2) = 1$, $(l_1, x'_N) = 1$ and $(l_2, x'_N) = 1$. If in addition (84) holds for $l = l_1, l_2$, then the above identity becomes

(91)
$$c_f(l_1x'_N)S_{l_1,\mathbf{r}}(1) \cdot c_f(l_2x'_N)S_{l_2,\mathbf{r}}(1) = c_f(l_1l_2x'_N)S_{l_1l_2,\mathbf{r}}(1).$$

From this we obtain a property of $c_f(n)$ very closed to the multiplicativity property. (We exclude the half-integral case below for simplicity.)

Proposition 9.28. Let $f(\tau) = \prod_{n|N} \eta(n\tau)^{r_n} = \sum_n c_f(n)q^n$ be a weakly admissible eta-quotient (e.g., a function listed in Table 3 or Example 9.23) where N is the level. We keep the notations in (56) with k' = k. Assume that $x'_N = \frac{x_N}{24} \in \mathbb{Z}_{\geq 0}$ and $k \in \mathbb{Z}$. Let $l_1, l_2 \in L_f$ satisfy $(l_1, l_2) = 1$, $(l_1, x'_N) = 1$ and $(l_2, x'_N) = 1$. We require (84) holds for $l = l_1$ and l_2 . Moreover, if $N \equiv 0 \mod 2$ and $N(k + \delta) \equiv 2 \mod 4$, then we require $l_1 \equiv l_2 \equiv 1 \mod 2$. (See Theorem 9.15(b) for the definition of δ .) Then

$$c_f(l_1x'_N) \cdot c_f(l_2x'_N) = \varepsilon \cdot c_f(l_1l_2x'_N),$$

where $\varepsilon = u_{l_1,\mathbf{r}} \cdot u_{l_2,\mathbf{r}} \cdot u_{l_1l_2,\mathbf{r}}$ with

$$u_{l,\mathbf{r}} = \begin{cases} 1, & \text{if } 2 \mid l, \quad 2 \nmid N \text{ or } 4 \mid N(k+\delta) \\ 1, & \text{if } 2 \nmid l, \quad 4 \mid N(k+\delta)(l-1) \\ -1, & \text{if } 2 \nmid l, \quad 4 \nmid N(k+\delta)(l-1). \end{cases}$$

Proof. According to (91) we need to prove

(92)
$$S_{l_1 l_2, \mathbf{r}}(1) = u_{l_1, \mathbf{r}} \cdot u_{l_2, \mathbf{r}} \cdot u_{l_1 l_2, \mathbf{r}} \cdot S_{l_1, \mathbf{r}}(1) \cdot S_{l_2, \mathbf{r}}(1).$$

By the definition of $\psi_{l,\mathbf{r}}$ in Theorem 9.15, we have, since $k \in \mathbb{Z}$,

$$S_{l,\mathbf{r}}(1) = \begin{cases} l, & \text{if } 2 \mid l, \quad 2 \nmid N \text{ or } 4 \mid N(k+\delta) \\ 0, & \text{if } 2 \mid l, \quad 2 \mid N \text{ and } N(k+\delta) \equiv 2 \mod 4 \\ l, & \text{if } 2 \nmid l, \quad 4 \mid N(k+\delta)(l-1) \\ -l, & \text{if } 2 \nmid l, \quad 4 \nmid N(k+\delta)(l-1). \end{cases}$$

From this (92) follows. Substituting (92) in (91) we obtain

$$c_f(l_1x'_N) \cdot c_f(l_2x'_N)S_{l_1,\mathbf{r}}(1)S_{l_2,\mathbf{r}}(1) = \varepsilon \cdot c_f(l_1l_2x'_N)S_{l_1,\mathbf{r}}(1)S_{l_2,\mathbf{r}}(1)$$

By the assumption we have $S_{l_1,\mathbf{r}}(1)$ and $S_{l_2,\mathbf{r}}(1)$ are nonzero and hence the required relation follows.

If $x'_N = 0$, then the above proposition holds trivially. We believe there are many weakly admissible eta-quotients with $x'_N \in \mathbb{Z}_{\geq 1}$ to which we can apply the above proposition. The most well-known example is the discriminant function $\Delta(\tau) = \eta(\tau)^{24}$, which in our terminology is an admissible eta-quotient of type II. The only example in Table 3 is the one presented below.

Example 9.29. Set $f = \eta(3\tau)^2 \eta(9\tau)^{-1} \eta(27\tau) = \sum_{n \in \mathbb{Z}_{\geq 1}} c_f(n) q^n$. This is an admissible etaquotient of type I. We have

$$N = 27, \quad k = 1, \quad x_N = 24, \quad \Pi = 3^5, \quad \delta = 1.$$

Since (c.f. (19))

$$\begin{pmatrix} x_1 \\ x_3 \\ x_9 \\ x_{27} \end{pmatrix} = \begin{pmatrix} 27 & 9 & 3 & 1 \\ 3 & 9 & 3 & 1 \\ 1 & 3 & 9 & 3 \\ 1 & 3 & 9 & 27 \end{pmatrix} \begin{pmatrix} r_1 \\ r_3 \\ r_9 \\ r_{27} \end{pmatrix} = \begin{pmatrix} 16 \\ 16 \\ 0 \\ 24 \end{pmatrix},$$

we have $m_f = 3$ (c.f. Corollary 9.13) and hence $L_f = \{3n + 1 : n \in \mathbb{Z}_{\geq 0}\}$. It follows from Proposition 9.28 that $c_f(l_1)c_f(l_2) = c_f(l_1l_2)$ where $l_1, l_2 \equiv 1 \mod 3$ are square-free positive integers with $(l_1, l_2) = 1$.

Example 9.30. Let us give an example of admissible eta-quotient of type II. Set

$$f = \eta(\tau)\eta(2\tau)\eta(3\tau)\eta(6\tau)^3 = \sum_{n \in \mathbb{Z}_{\geq 1}} c_f(n)q^n.$$

We have

$$N = 6, \quad k = 3, \quad x_N = 24, \quad \Pi = 2^3 \cdot 3^2, \quad \delta = 0$$

Since

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_6 \end{pmatrix} = \begin{pmatrix} 6 & 3 & 2 & 1 \\ 3 & 6 & 1 & 2 \\ 2 & 1 & 6 & 3 \\ 1 & 2 & 3 & 6 \end{pmatrix} \begin{pmatrix} r_1 \\ r_2 \\ r_3 \\ r_6 \end{pmatrix} = \begin{pmatrix} 14 \\ 16 \\ 18 \\ 24 \end{pmatrix},$$

we have $m_f = 12$ and hence $L_f = \{12n + 1 : n \in \mathbb{Z}_{\geq 0}\}$. By Theorem 4.2, $\dim_{\mathbb{C}} M_3(\Gamma_0(6), \chi) = 2$ where $\chi \colon \Gamma_0(6) \to \mathbb{C}^{\times}$ is the character of f. (Of course this descends to a character of the matrix group $\Gamma_0(6)$ since $k \in \mathbb{Z}$.) Recall that the number of Eisenstein series in $M_3(\Gamma_0(6), \chi)$ is denoted by n_0 in (50). Since $\chi(\widetilde{\gamma_{a/c}}\widetilde{T}^{w_{a/c}}\widetilde{\gamma_{a/c}}^{-1}) = 1$ if and only if $x_c \equiv 0 \mod 24$ where $c \mid 6$ and (a, c) = 1, among the four cusps in $\Gamma_0(6) \setminus \mathbb{P}^1(\mathbb{Q})$ there is only one Eisenstein series which is defined at the cusp i ∞ . Therefore $n_0 = 1$. Since f is a cusp form, we find that it is admissible of type II. It follows from Proposition 9.28 (applied to this f) that $c_f(l_1)c_f(l_2) = c_f(l_1l_2)$ whenever $l_1, l_2 \equiv 1 \mod 12$ are square-free positive integers with $(l_1, l_2) = 1$. It is worthwhile to notice that for other pair (l_1, l_2) , the multiplicativity does not necessarily hold. For instance, $c_f(2) = -1$, $c_f(3) = -2$ but $c_f(6) = 4$.

9.4.3. An example of half-integral weight. The examples presented above are all of integral weights. To conclude this section we supplement an example of half-integral weight where we show how the Gauss sum \sum_{b} in (81) can be simplified. Let

(93)
$$f(\tau) = \eta(\tau)^{-7} \eta(2\tau)^{17} \eta(4\tau)^{-3} = \sum_{n \in \frac{5}{8} + \mathbb{Z}_{\ge 0}} c_f(n) q^n.$$

The statistics of f are

$$N = 4$$
, $k = \frac{7}{2}$, $x_1 = 3$, $x_2 = 24$, $x_4 = 15$, $\Pi = 2^{31}$, $\delta = 0$, $m_f = 8$.

Thus $L_f = \{l^2 : l \equiv 1 \mod 2\}$. By Theorem 4.2 we have $\dim_{\mathbb{C}} M_{7/2}(\Gamma_0(4), \chi) = 2$ where $\chi : \widetilde{\Gamma_0(4)} \to \mathbb{C}^{\times}$ is the character of f and by Definition 8.3 the Eisenstein series can only be given at the cusp 1/2. Thus, f is admissible of type II and Theorems 9.19, 9.24 can be applied.

Now we calculate c_{l^2} for $l^2 \in L_f$ (c.f. Theorem 9.19). We have, by definition,

(94)
$$c_{l^{2}} = l^{-\frac{7}{2}} \sum_{a|l^{2}} \left(\frac{a}{2}\right) a^{\frac{7}{2}} c_{f} \left(\frac{5l^{2}}{8a^{2}}\right) \mathfrak{e} \left(-\frac{7l^{2}}{8a} + \frac{7}{8}\right) \sum_{\substack{0 \le b < l^{2}/a \\ (a,b,l^{2}/a) = 1}} \left(\frac{-b}{(a,l^{2}/a)}\right) \mathfrak{e} \left(-\frac{5(l^{2}-1)b}{8a}\right).$$

Note that $c_f\left(\frac{5l^2}{8a^2}\right) = 0$ unless $\frac{5l^2}{8a^2} \in \frac{5}{8} + \mathbb{Z}$, that is, unless $a \mid l$ since a, l are odd. It follows that for such a we have $(a, l^2/a) = a$,

$$t := \frac{5(l^2 - 1) \cdot (a, l^2/a)}{8a} = \frac{5(l^2 - 1)}{8}$$

is an integer (depending on a and l) and

(95)
$$\sum_{\substack{0 \le b < l^2/a \\ (a,b,l^2/a)=1}} \left(\frac{-b}{(a,l^2/a)}\right) \mathfrak{e}\left(-\frac{5(l^2-1)b}{8a}\right) = \sum_{0 \le b < l^2/a} \left(\frac{-b}{a}\right) \mathfrak{e}\left(\frac{-tb}{a}\right) \mathfrak{e}\left(\frac{-tb}{a}\right) \mathfrak{e}\left(\frac{-b}{a}\right) \mathfrak{e}\left(\frac{-b}{a}\right)$$

is, up to a simple factor, a Gauss sum associated with the character $\left(\frac{\cdot}{(a,l^2/a)}\right)$. To express the values of these Gauss sums, we let, for any odd positive integer m,

$$\operatorname{rad}_{E}(m) = \prod_{\substack{p \mid m \\ v_{p}(m) \equiv 0 \mod 2}} p \qquad \operatorname{rad}_{O}(m) = \prod_{\substack{p \mid m \\ v_{p}(m) \equiv 1 \mod 2}} p$$
$$\operatorname{rad}(m) = \operatorname{rad}_{E}(m) \operatorname{rad}_{O}(m) \qquad \operatorname{rad}'(m) = \operatorname{rad}_{E}(m)^{2} \operatorname{rad}_{O}(m)$$
$$\operatorname{irad}(m) = m/\operatorname{rad}(m) \qquad \operatorname{irad}'(m) = m/\operatorname{rad}'(m)$$

where p denotes a prime. By [20, Lemma 5.6], if $irad(a) \nmid t$ then $\sum_{0 \leq b < a} \left(\frac{b}{a}\right) \mathfrak{e}\left(\frac{tb}{a}\right) = 0$; if $irad(a) \mid t$ then

(96)
$$\sum_{0 \le b < a} \left(\frac{b}{a}\right) \mathfrak{e}\left(\frac{tb}{a}\right) = \varepsilon_a \frac{a}{\sqrt{\mathrm{rad}'(a)}} \left(\frac{t/\mathrm{irad}'(a)}{\mathrm{rad}_O(a)}\right) \prod_{p \mid \mathrm{rad}_E(a)} \left(p - 1 - p\left(\frac{t/\mathrm{irad}(a)}{p}\right)^2\right),$$

where $\left(\frac{t/\operatorname{irad}'(a)}{\operatorname{rad}_O(a)}\right)$, $\left(\frac{t/\operatorname{irad}(a)}{p}\right)$ are Jacobi symbols and $\varepsilon_a = 1$ ($\varepsilon_a = i$ respectively) if $a \equiv 1 \mod 4$ ($a \equiv 3 \mod 4$ respectively). Inserting (96) and (95) into (94) and noting that $\mathfrak{e}\left(-\frac{7l^2}{8a}+\frac{7}{8}\right) = \mathfrak{e}\left(\frac{a-1}{8}\right)$, $\left(\frac{a}{2}\right)\varepsilon_a\mathfrak{e}\left(\frac{a-1}{8}\right) = 1$ we obtain the following formula.

Proposition 9.31. For $f(\tau) = \eta(\tau)^{-7} \eta(2\tau)^{17} \eta(4\tau)^{-3} = \sum_{n \in \frac{5}{8} + \mathbb{Z}_{\geq 0}} c_f(n) q^n$ and $1 \leq l \equiv 1 \mod 2$ we have

$$c_{l^{2}} = l^{-\frac{3}{2}} \sum_{\substack{a|l\\ \text{irad}(a)|t}} \frac{a^{\frac{5}{2}}}{\sqrt{\text{rad}'(a)}} \prod_{p|\text{rad}_{E}(a)} \left(p - 1 - p\left(\frac{t/\operatorname{irad}(a)}{p}\right)^{2}\right) \left(\frac{t/\operatorname{irad}'(a)}{\operatorname{rad}_{O}(a)}\right) c_{f}\left(\frac{5l^{2}}{8a^{2}}\right)$$

where $t = \frac{5(l^2-1)}{8}$. In particular, if l is square-free, then

$$c_{l^2} = l^{-\frac{3}{2}} \sum_{a|l} a^2 \cdot \left(\frac{t}{a}\right) c_f\left(\frac{5l^2}{8a^2}\right).$$

Remark 9.32. Since $\frac{a^{5/2}}{\sqrt{\operatorname{rad}'(a)}} \in \mathbb{Z}$ and $c_f(n) \in \mathbb{Z}$, we have $l^{3/2} \cdot c_{l^2} \in \mathbb{Z}$.

According to Theorem 9.24 (applied to (93)) we have $c_{l_1^2}c_{l_2^2} = c_{l_1^2l_2^2}$ whenever l_1, l_2 are coprime odd integers. In the special case l_1 and l_2 are square-free, this identity reads

$$(97) \quad \sum_{a|l_1} a^2 \cdot \left(\frac{5(l_1^2 - 1)/8}{a}\right) c_f\left(\frac{5l_1^2}{8a^2}\right) \cdot \sum_{a|l_2} a^2 \cdot \left(\frac{5(l_2^2 - 1)/8}{a}\right) c_f\left(\frac{5l_2^2}{8a^2}\right) \\ = \sum_{a|l_1l_2} a^2 \cdot \left(\frac{5(l_1^2l_2^2 - 1)/8}{a}\right) c_f\left(\frac{5l_1^2l_2^2}{8a^2}\right).$$

From this we find the genuine multiplicativity property of (93) as stated below. The reader may compare this to the integral-weight counterpart—Proposition 9.28.

Proposition 9.33. Let $f(\tau) = \eta(\tau)^{-7} \eta(2\tau)^{17} \eta(4\tau)^{-3} = \sum_{n \in \frac{5}{8} + \mathbb{Z}_{\geq 0}} c_f(n) q^n$. Let l_1, l_2 be coprime odd square-free integers. Then

$$c_f\left(\frac{5l_1^2}{8}\right)c_f\left(\frac{5l_2^2}{8}\right) = c_f\left(\frac{5l_1^2l_2^2}{8}\right).$$

Proof. First suppose l_1 and l_2 are primes. By (97) we have

$$(98) \quad \left(c_f\left(\frac{5l_1^2}{8}\right) + l_1^2 \cdot \left(\frac{5(l_1^2 - 1)/8}{l_1}\right)\right) \cdot \left(c_f\left(\frac{5l_2^2}{8}\right) + l_2^2 \cdot \left(\frac{5(l_2^2 - 1)/8}{l_2}\right)\right) \\ = c_f\left(\frac{5l_1^2l_2^2}{8}\right) + l_2^2 \cdot \left(\frac{5(l_1^2l_2^2 - 1)/8}{l_2}\right) c_f\left(\frac{5l_1^2}{8}\right) + l_1^2 \cdot \left(\frac{5(l_1^2l_2^2 - 1)/8}{l_1}\right) c_f\left(\frac{5l_2^2}{8}\right) + l_1^2 l_2^2 \cdot \left(\frac{5(l_1^2l_2^2 - 1)/8}{l_1l_2}\right).$$

It is not hard to prove, by the periodicity and multiplicativity of Jacobi symbols, that

$$\begin{pmatrix} \frac{5(l_1^2 l_2^2 - 1)/8}{l_1} \end{pmatrix} = \begin{pmatrix} \frac{5(l_1^2 - 1)/8}{l_1} \end{pmatrix}, \begin{pmatrix} \frac{5(l_1^2 l_2^2 - 1)/8}{l_2} \end{pmatrix} = \begin{pmatrix} \frac{5(l_2^2 - 1)/8}{l_2} \end{pmatrix}, \begin{pmatrix} \frac{5(l_1^2 l_2^2 - 1)/8}{l_1 l_2} \end{pmatrix} = \begin{pmatrix} \frac{5(l_1^2 - 1)/8}{l_1} \end{pmatrix} \cdot \begin{pmatrix} \frac{5(l_2^2 - 1)/8}{l_2} \end{pmatrix}.$$

Inserting these into (98) gives the desired identity in the case l_1 and l_2 are primes.

Second, suppose l_1 is a prime and l_2 is arbitrary. Again by (97) we have

$$(99) \quad \left(c_f\left(\frac{5l_1^2}{8}\right) + l_1^2 \cdot \left(\frac{5(l_1^2 - 1)/8}{l_1}\right)\right) \cdot \sum_{a|l_2} a^2 \cdot \left(\frac{5(l_2^2 - 1)/8}{a}\right) c_f\left(\frac{5l_2^2}{8a^2}\right) \\ = \sum_{a|l_2} a^2 \cdot \left(\frac{5(l_1^2l_2^2 - 1)/8}{a}\right) c_f\left(\frac{5l_1^2l_2^2}{8a^2}\right) + \sum_{a|l_2} l_1^2 a^2 \cdot \left(\frac{5(l_1^2l_2^2 - 1)/8}{l_1a}\right) c_f\left(\frac{5l_2^2}{8a^2}\right).$$

Since for $a \mid l_2$,

$$\left(\frac{5(l_1^2 - 1)/8}{l_1}\right) \cdot \left(\frac{5(l_2^2 - 1)/8}{a}\right) = \left(\frac{5(l_1^2 l_2^2 - 1)/8}{l_1 a}\right)$$

we obtain from (99) that

(100)
$$c_{f}\left(\frac{5l_{1}^{2}}{8}\right) \cdot \sum_{a|l_{2}} a^{2} \cdot \left(\frac{5(l_{2}^{2}-1)/8}{a}\right) c_{f}\left(\frac{5l_{2}^{2}}{8a^{2}}\right) = \sum_{a|l_{2}} a^{2} \cdot \left(\frac{5(l_{1}^{2}l_{2}^{2}-1)/8}{a}\right) c_{f}\left(\frac{5l_{1}^{2}l_{2}^{2}}{8a^{2}}\right)$$
$$= \sum_{a|l_{2}} a^{2} \cdot \left(\frac{5(l_{2}^{2}-1)/8}{a}\right) c_{f}\left(\frac{5l_{1}^{2}l_{2}^{2}}{8a^{2}}\right).$$

The desired identities for l_1 prime and l_2 arbitrary now follows from (100), the already proven case and the induction on the number of prime divisors of l_2 .

Finally, if l_1 and l_2 are arbitrary, then, by the already proven cases, both sides of the desired identity are equal to $\prod_{p|l_1l_2} c_f\left(\frac{5p^2}{8}\right)$ and hence the identity holds which concludes the proof. \Box

We have discussed consequences of Theorem 9.24 applied to (93). The above exploration is also valid for any admissible eta-quotient of half-integral weight. (Of course the integral weight case is simpler to deal with.) In addition, for (93) or any admissible eta-quotient of half-integral weight, we can as well simplify the left-hand side of (83) as we have just done for c_{l^2} . This will give more interesting identities about $c_f(n)$. We will not include more details about this due to the length of the paper.

10. Miscellaneous observations and open questions

More general multiplier systems. The dimension formulas stated in Theorem 4.2 concern multiplier systems induced by eta-quotients of fractional exponents. As a comparison, most of the formulas that have appeared in the literature concern multiplier systems induced by Dirichlet characters (in the case of integral weights) or Dirichlet characters times the multiplier system of a power of $\eta(\tau)^{-2}\eta(2\tau)^5\eta(4\tau)^{-2}$ (in the case of half-integral weights). It is meaningful to deduce an explicit formula for dim_C $M_k(\Gamma_0(N), \chi_1\chi_2)$ where $k \in \mathbb{Q}, \chi_1$ is induced by an eta-quotient of fractional exponents and χ_2 by a Dirichlet character.

Do generalized double coset operators exist for rational weights? Theorem 9.9 let us know for what numbers l does the expression $T_l f$ make sense. The multiplier systems involved are

required to be induced by eta-quotients of *integral* exponents. The question is: if the multiplier systems are induced by eta-quotients of fractional exponents, is there any nontrivial T_l (i.e. l > 1)? All identities presented in Section 9.4 are based on Theorem 9.9. Thus if we can find any T_l with l > 1 in the case of fractional exponents, then, taking into account of Theorem 4.2, we would obtain identities involving $c_f(n)$ for infinitely many eta-quotients f of *fractional* exponents. However, no such operator has been found up to now.

L-functions and Euler products. According to Theorem 9.24, it is natural to associate an L-function $L(f,s) = \sum_{l \in L_f} \frac{c_l}{l^s}$ with a weakly admissible eta-quotient f. This association is different from the usual one $\sum_n \frac{c_f(n)}{n^s}$. The identities presented in Section 9.4 can be rephrased as properties of this L(f,s). For instance, it has an Euler product. It is interesting to investigate L(f,s), e.g., the functional equations, the corresponding Weil's theorem, etc.

Listing all weakly admissible eta-quotients. Open question: to find out all weakly admissible eta-quotients (if there exist only finitely many). Another open question: for each weakly admissible eta-quotient, to deduce identities like the one in Proposition 9.33 from (83).

Appendix A. Usage of SageMath code

The SageMath code can be obtained from the repository [43]. It is an ipynb file which should be opened in a Sage Jupyter notebook. The code in the first two cells should be run after the file is opened. The first cell, which actually is the one in [44], contains Python/Sage functions dealing with eta-quotients among which we need those concerning the characters and Fourier coefficients of eta-quotients of fractional exponents. The second cell then contains Python/Sage functions on computing dimensions, on order-character relations, on checking identities and on generating tables in this paper.

References

- H. Petersson, Zur analytischen Theorie der Grenzkreisgruppen, Math. Ann. 115 (1) (1938) 175-204. doi:10.1007/BF01448938. URL https://doi.org/10.1007/BF01448938
- [2] G. Shimura, Introduction to the arithmetic theory of automorphic functions, Vol. No. 1 of Kanô Memorial Lectures, Iwanami Shoten Publishers, Tokyo; Princeton University Press, Princeton, NJ, 1971, publications of the Mathematical Society of Japan, No. 11.
- [3] D. Zagier, Traces des opérateurs de Hecke, in: Séminaire Delange-Pisot-Poitou, 17e année: 1975/76, Théorie des nombres: Fasc. 2, Secrétariat Math., Paris, 1977, pp. Exp. No. 23, 12.
- [4] D. Zagier, Correction to: "The Eichler-Selberg trace formula on SL₂(Z)" (introduction to modular forms, Appendix, pp. 44–54, Springer, Berlin, 1976) by S. Lang, in: Modular functions of one variable, VI (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976), Vol. Vol. 627 of Lecture Notes in Math., Springer, Berlin-New York, 1977, pp. 171–173.
- [5] H. Cohen, Trace des opérateurs de Hecke sur $\Gamma_0(N)$, in: Séminaire de Théorie des Nombres (1976–1977), Centre National de Recherche Scientifique, Laboratoire de Théorie des Nombres, Talence, 1977, pp. Exp. No. 4, 9.

[6] H. Hijikata, Explicit formula of the traces of Hecke operators for $\Gamma_0(N)$, J. Math. Soc. Japan 26 (1974) 56–82. doi:10.2969/jmsj/02610056.

URL https://doi.org/10.2969/jmsj/02610056

- [7] H. Cohen, J. Oesterlé, Dimensions des espaces de formes modulaires, in: Modular functions of one variable, VI (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976), Vol. Vol. 627 of Lecture Notes in Math., Springer, Berlin-New York, 1977, pp. 69–78.
- [8] G. Martin, Dimensions of the spaces of cusp forms and newforms on Γ₀(N) and Γ₁(N), J. Number Theory 112 (2) (2005) 298-331. doi:10.1016/j.jnt.2004.10.009.
 URL https://doi.org/10.1016/j.jnt.2004.10.009
- [9] J. Quer, Dimensions of spaces of modular forms for Γ_H(N), Acta Arith. 145 (4) (2010) 373-395.
 doi:10.4064/aa145-4-4.
 URL https://doi.org/10.4064/aa145-4-4
- [10] K. Martin, Refined dimensions of cusp forms, and equidistribution and bias of signs, J. Number Theory 188 (2018) 1-17. doi:10.1016/j.jnt.2018.01.015.
 URL https://doi.org/10.1016/j.jnt.2018.01.015
- [11] Y. Zhang, Y. Zhou, Dimension formulas for modular form spaces with character for Fricke groups, Acta Arith. 206 (4) (2022) 291-311. doi:10.4064/aa211126-17-11. URL https://doi.org/10.4064/aa211126-17-11
- T. Ibukiyama, Modular forms of rational weights and modular varieties, Abh. Math. Sem. Univ. Hamburg 70 (2000) 315-339. doi:10.1007/BF02940923.
 URL https://doi.org/10.1007/BF02940923
- [13] T. Ibukiyama, Graded rings of modular forms of rational weights, Res. Number Theory 6 (1) (2020) Paper No. 8, 13. doi:10.1007/s40993-019-0183-9.
 URL https://doi.org/10.1007/s40993-019-0183-9
- [14] J.-P. Serre, H. M. Stark, Modular forms of weight 1/2, in: Modular functions of one variable, VI (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976), Vol. Vol. 627 of Lecture Notes in Math., Springer, Berlin-New York, 1977, pp. 27–67.
- [15] P. Deligne, J.-P. Serre, Formes modulaires de poids 1, Ann. Sci. École Norm. Sup. (4) 7 (1974) 507-530. URL http://www.numdam.org/item?id=ASENS_1974_4_7_4_507_0
- [16] X.-J. Zhu, Holomorphic Eisenstein series of rational weights and special values of gamma function, Acta Arith. 210 (2023) 279-305. doi:10.4064/aa221110-1-4. URL https://doi.org/10.4064/aa221110-1-4
- [17] Y. Martin, Multiplicative η-quotients, Trans. Amer. Math. Soc. 348 (12) (1996) 4825-4856.
 doi:10.1090/S0002-9947-96-01743-6.
 URL https://doi.org/10.1090/S0002-9947-96-01743-6
- [18] A. O. L. Atkin, J. Lehner, Hecke operators on $\Gamma_0(m)$, Math. Ann. 185 (1970) 134–160. doi:10.1007/BF01359701.
 - URL https://doi.org/10.1007/BF01359701
- K. Wohlfahrt, Über Operatoren Heckescher Art bei Modulformen reeller Dimension, Math. Nachr. 16 (1957) 233-256. doi:10.1002/mana.19570160307. URL https://doi.org/10.1002/mana.19570160307
- [20] H.-G. Zhou, X.-J. Zhu, Double coset operators and eta-quotients, J. Number Theory 249 (2023) 537-601. doi:10.1016/j.jnt.2023.02.017. URL https://doi.org/10.1016/j.jnt.2023.02.017
- [21] S. Bhattacharya, Holomorphic eta quotients of weight 1/2, Adv. Math. 320 (2017) 1185-1200. doi:10.1016/j.aim.2017.09.017. URL https://doi.org/10.1016/j.aim.2017.09.017
- [22] SageMath, The Sage mathematics software system (version 9.2), The Sage Developers (2021). URL https://www.sagemath.org
- [23] H. Cohen, Number theory. Vol. I. Tools and Diophantine equations, Vol. 239 of Graduate Texts in Mathematics, Springer, New York, 2007.

84

- [24] M. Knopp, G. Mason, Revisions to "Parabolic generalized modular forms and their characters" [mr2553511], Int. J. Number Theory 8 (4) (2012) 859-864. doi:10.1142/S1793042112500509. URL https://doi.org/10.1142/S1793042112500509
- [25] M. Knopp, G. Mason, Generalized modular forms, J. Number Theory 99 (1) (2003) 1–28. doi:10.1016/S0022-314X(02)00065-3.
- URL https://doi.org/10.1016/S0022-314X(02)00065-3 [26] F. Freiter, P. M. Hill, Modular forms on SU(2, 1) with weight ¹ Pos. Nur
- [26] E. Freitag, R. M. Hill, Modular forms on SU(2,1) with weight ¹/₃, Res. Number Theory 9 (2) (2023) Paper No. 26, 20. doi:10.1007/s40993-022-00361-6.
 URL https://doi.org/10.1007/s40993-022-00361-6
- [27] Y. Lee, S. Lim, Zagier duality for real weights, Acta Arith. 207 (3) (2023) 235-249.
 doi:10.4064/aa220614-13-1.
 URL https://doi.org/10.4064/aa220614-13-1
- [28] T. Kubota, Topological covering of SL(2) over a local field, J. Math. Soc. Japan 19 (1967) 114-121. doi:10.2969/jmsj/01910114. URL https://doi.org/10.2969/jmsj/01910114
- [29] H. Cohen, F. Strömberg, Modular forms, Vol. 179 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2017, a classical approach. doi:10.1090/gsm/179.
- URL https://doi.org/10.1090/gsm/179
 [30] T. M. Apostol, Modular functions and Dirichlet series in number theory, 2nd Edition, Vol. 41 of Graduate Texts in Mathematics, Springer-Verlag, New York, 1990.
- URL https://doi.org/10.1007/978-1-4612-0999-7 [31] S. Bhattacharya, Special factors of holomorphic eta quotients, Adv. Math. 392 (2021) Paper No. 108019, 24.
- doi:10.1016/j.aim.2021.108019. URL https://doi.org/10.1016/j.aim.2021.108019
- [32] G. Shimura, On modular forms of half integral weight, Ann. of Math. (2) 97 (1973) 440–481. doi:10.2307/1970831.
 - URL https://doi.org/10.2307/1970831
- [33] E. Freitag, Complex analysis. 2, Universitext, Springer, Heidelberg, 2011, riemann surfaces, several complex variables, abelian functions, higher modular functions. doi:10.1007/978-3-642-20554-5. URL https://doi.org/10.1007/978-3-642-20554-5
- [34] J. M. Lee, Introduction to topological manifolds, 2nd Edition, Vol. 202 of Graduate Texts in Mathematics, Springer, New York, 2011. doi:10.1007/978-1-4419-7940-7. URL https://doi.org/10.1007/978-1-4419-7940-7
- [35] T. Miyake, Modular forms, english Edition, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2006, translated from the 1976 Japanese original by Yoshitaka Maeda.
- [36] F. Diamond, J. Shurman, A first course in modular forms, Vol. 228 of Graduate Texts in Mathematics, Springer-Verlag, New York, 2005.
- [37] W. Kohnen, G. Mason, On generalized modular forms and their applications, Nagoya Math. J. 192 (2008) 119– 136. doi:10.1017/S0027763000026003.
 - URL https://doi.org/10.1017/S0027763000026003
- [38] M. Newman, Construction and application of a class of modular functions. II, Proc. London Math. Soc. (3) 9 (1959) 373-387. doi:10.1112/plms/s3-9.3.373.
 URL https://doi.org/10.1112/plms/s3-9.3.373
- [39] K. Ono, The web of modularity: arithmetic of the coefficients of modular forms and q-series, Vol. 102 of CBMS Regional Conference Series in Mathematics, Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2004.
- [40] J. Weisinger, SOME RESULTS ON CLASSICAL EISENSTEIN SERIES AND MODULAR FORMS OVER FUNCTION I ProQuest LLC, Ann Arbor, MI, 1977, thesis (Ph.D.)-Harvard University. URL http://gateway.proquest.com/openurl?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:dissertat
- [41] D. Dummit, H. Kisilevsky, J. McKay, Multiplicative products of η-functions, in: Finite groups—coming of age (Montreal, Que., 1982), Vol. 45 of Contemp. Math., Amer. Math. Soc., Providence, RI, 1985, pp. 89–98.

doi:10.1090/conm/045/822235. URL https://doi.org/10.1090/conm/045/822235

- [42] M. I. Knopp, Modular functions in analytic number theory, Markham Publishing Co., Chicago, Ill., 1970.
- [43] X.-J. Zhu, SageMath code for the paper Dimension formulas for modular form spaces of rational weights, the classification (2024).

URL https://github.com/Xiao-Jie-Zhu/Dimension-formulas-for-modular-form-spaces-of-rational-weights [44] X.-J. Zhu, Code for Double coset operators and eta-quotients (2023).

URL https://github.com/Xiao-Jie-Zhu/double_coset_operators_and_eta_quotients

School of Mathematical Sciences, Key Laboratory of MEA (Ministry of Education) & Shanghai Key Laboratory of PMMP, East China Normal University, 500 Dongchuan Road, 200241, Shanghai, P. R. China

Email address: zhuxiaojiemath@outlook.com *URL*: https://orcid.org/0000-0002-6733-0755