

Quantized electrical, thermal, and spin transports of non-Hermitian clean and dirty two-dimensional topological insulators and superconductors

Sanjib Kumar Das¹ and Bitan Roy¹

¹*Department of Physics, Lehigh University, Bethlehem, Pennsylvania, 18015, USA*

(Dated: August 2, 2024)

From lattice-regularized models, devoid of any non-Hermitian (NH) skin effects, here we compute the electrical (σ_{xy}), thermal (κ_{xy}), and spin (σ_{xy}^{sp}) Hall, and the electrical (G_{xx}) and thermal (G_{xx}^{th}) longitudinal conductivities for appropriate NH planar topological insulators and superconductors related to all five non-trivial Altland-Zirbauer symmetry classes in the Hermitian limits. These models feature real eigenvalues over an extended NH parameter regime, only where the associated topological invariants remain quantized. In this regime, the NH quantum anomalous and spin Hall insulators show quantized σ_{xy} and G_{xx} , respectively, the NH $p + ip$ ($p \pm ip$) pairing shows half-quantized κ_{xy} (G_{xx}^{th}), while the NH $d + id$ pairing shows quantized κ_{xy} and σ_{xy}^{sp} in the clean and weak disorder (due to random pointlike charge impurities) regimes. We compute these quantities in experimentally realizable suitable six-terminal setups using the Kwant software package. But, in the strong disorder regime, all these topological responses vanish and with the increasing non-Hermiticity in the system this generic phenomenon occurs at weaker disorder.

Introduction. Topological crystals displaying the hallmark bulk-boundary correspondence in terms of robust gapless modes living on their boundaries [1–21] typically encounter the notorious non-Hermitian (NH) skin effect in open systems interacting with a bath or the environment [22–45]. It corresponds to an accumulation of all the right and left eigenvectors at opposite ends of the system with open boundary conditions. Even though topological invariants can be defined in NH systems their signatures on the the boundary modes get masked by the NH skin effect. Theoretically, this obstacle can be bypassed through the bi-orthogonal bulk-boundary correspondence, devoid of any NH skin effect [32]. Nevertheless, any direct manifestation of the topological invariants on experimentally measurable quantities even in a NH toy topological model remains illusive [46], leaving aside their actual measurements in open quantum crystals. In this quest, two-dimensional NH topological models stand as cornerstones, since in the Hermitian or closed systems their topological invariants can be computed and measured in six-terminal Hall-bar setups [47–56].

Here, we consider skin effect free NH generalization of square lattice models for all five non-trivial Hermitian Altland-Zirnbauer symmetry classes [57] that over an extended NH parameter regime feature real eigenvalue spectrum and non-vanishing topological invariants [58]. Using numerical Kwant software package [59–62], we compute their electrical (σ_{xy}), thermal (κ_{xy}), and spin (σ_{xy}^{sp}) Hall, and electrical (G_{xx}) and thermal (G_{xx}^{th}) longitudinal conductivities in a six-terminal setup (Fig. 1), to arrive at the following conclusions. The NH quantum anomalous Hall insulator (QAHI) [quantum spin Hall insulator (QSHI)] belonging to class A [class AII] in the Hermitian limit, shows quantized σ_{xy} [G_{xx}] in units of e^2/h (Fig. 2), where e (h) is the electrical charge (Planck’s constant). By contrast, the NH $p + ip$ ($p \pm ip$) paired state, belonging to class D (class DIII) in Hermitian




FIG. 1. Six-terminal setup with a rectangular scattering region (black sites) of length L and width D connected to six semi-infinite leads (red sites), for the calculation of electrical (σ_{xy}), thermal (κ_{xy}), and spin (σ_{xy}^{sp}) Hall, and electrical (G_{xx}) and thermal (G_{xx}^{th}) longitudinal conductivities. For the computation of σ_{xy} and G_{xx} , an electrical current (I_{el}) flows due to a voltage bias ($\Delta x \equiv \Delta V$) between the horizontal leads, and we extract voltages $x_i \equiv V_i$ on the vertical leads with $i = 2, 3, 5, 6$. Due to a temperature gradient ($\Delta x \equiv \Delta T$), a thermal current (I_{th}) flows between the horizontal leads and temperature develops at the vertical leads with $x_i \equiv T_i$, when we compute κ_{xy} and G_{xx}^{th} . While calculating σ_{xy}^{sp} , a spin current (I_{sp}) flows between the horizontal leads, subject to a magnetic field bias with $\Delta x \equiv \Delta H$, yielding magnetization at the vertical leads with $x_i \equiv m_i$. The scattering region is maintained at a fixed voltage V (for σ_{xy} and G_{xx}) or temperature T (for κ_{xy} and G_{xx}^{th}) or magnetic field H (for σ_{xy}^{sp}). Throughout, we set $L = 180$ and $D = 90$ (Figs. 2–5).

systems, supports half-quantized κ_{xy} (G_{xx}^{th}) in units of $\kappa_0 = \pi^2 k_B^2 T / (3h)$ as the temperature $T \rightarrow 0$, where k_B is the Boltzmann constant (Fig. 3). Finally, the NH $d + id$ paired state, falling in class C in the Hermitian limit, accommodates quantized κ_{xy} (in units of κ_0) and σ_{xy}^{sp} (in units of $\sigma_0^{sp} = h / (8\pi)$). See Fig. 4. These conclu-




FIG. 2. Electrical Hall conductivity (σ_{xy}) of a NH QAH I for (a) a fixed strength of non-Hermiticity (t_2) as a function of m_0 and (b) two fixed values of m_0 with the varying strength of t_2 , showing sharp quantization $\sigma_{xy} = C$ in units of e^2/h (within numerical accuracy), where C is the first Chern number. Electrical longitudinal conductivity (G_{xx}) of a NH QSHI for (c) a fixed t_2 as a function of m_0 and (d) two fixed values of m_0 with varying t_2 , showing $G_{xx} = C_{sp}$ (in units of e^2/h), where C_{sp} is the spin Chern number.

sions hold in the entire topological regime in a clean and weakly disordered (due to pointlike random charge impurities) systems. In the strong disorder regime, all these responses vanish, which with increasing non-Hermiticity in the system occur at weaker disorder (Fig. 5).

Model. For all the gapped topological phases belonging to any one of the five non-trivial Altland-Zirnbauer symmetry classes in two dimensions, the model Hamiltonian on a square lattice take the following universal form

$$H = \alpha \sum_{\mathbf{k}} \Psi_{\mathbf{k}}^\dagger \left(\sum_{j=1}^3 d_j(\mathbf{k}) \Gamma_j \right) \Psi_{\mathbf{k}} \equiv \alpha \sum_{\mathbf{k}} \Psi_{\mathbf{k}}^\dagger \mathcal{H}(\mathbf{k}) \Psi_{\mathbf{k}}. \quad (1)$$

The internal structure of the spinor $\Psi_{\mathbf{k}}$ with momentum $\mathbf{k} = (k_x, k_y)$ and representation of the Hermitian Γ matrices depend on the symmetry class, while always satisfying the anticommuting Clifford algebra $\{\Gamma_j, \Gamma_k\} = 2\delta_{jk}$. The third component of the $\mathbf{d}(\mathbf{k})$ -vector takes the form $d_3(\mathbf{k}) = m_0 - t_0 [\cos(k_x a) + \cos(k_y a)]$, and unless otherwise stated $d_1(\mathbf{k}) = t_1 \sin(k_x a)$ and $d_2(\mathbf{k}) = t_1 \sin(k_y a)$, where a is the lattice spacing. Within the topological regime ($|m_0/t_0| < 2$), for topological insulators (superconductors) $d_3(\mathbf{k})$ features a band inversion (Fermi surface) near the $\Gamma = (0, 0)$ and $M = (1, 1)\pi/a$ points of the Brillouin zone (BZ) when $0 < m_0/t_0 < 2$ and $-2 < m_0/t_0 < 0$, respectively, with $\alpha = 1$ ($\alpha = 1/2$ due to the Nambu doubling). Throughout, we set $t_1 = t_0 = 1$.

Note that $d_3(\mathbf{k})$ and Γ_3 do not break any crystal symmetry, and $\mathcal{H}_\perp(\mathbf{k}) = d_1(\mathbf{k})\Gamma_1 + d_2(\mathbf{k})\Gamma_2$ is also invariant under all the crystal symmetries. So, $\mathcal{H}(\mathbf{k})$ trans-




FIG. 3. Thermal Hall conductivity (κ_{xy}) of a NH $p + ip$ paired state for (a) a fixed non-Hermiticity (t_2) as a function of m_0 and (b) two fixed values of m_0 with varying t_2 , showing sharp half-quantization $\kappa_{xy} = C/2$ (in units of κ_0). Thermal longitudinal conductivity (G_{xx}^{th}) of a NH $p \pm ip$ superconductor for (c) a fixed t_2 with varying m_0 and (d) two fixed values of m_0 with varying t_2 , showing $G_{xx}^{th} = C_{sp}/2$ (in units of κ_0).

forms under the trivial A_{1g} representation. Then the anti-Hermitian operator $\mathcal{H}_{\text{AH}}(\mathbf{k}) = \Gamma_3 \mathcal{H}_\perp(\mathbf{k})/t_1$ does not break any crystallographic symmetry either. Finally, we introduce the desired NH topological operator

$$\mathcal{H}_{\text{NH}}(\mathbf{k}) = \mathcal{H}(\mathbf{k}) + t_2 \mathcal{H}_{\text{AH}}(\mathbf{k}) \quad (2)$$

that is symmetric under all crystal symmetries, such as reflections, rotations, and inversion. Hence, it is devoid of any NH skin effect for all parameter values [58]. Here, t_2 determines the strength of non-Hermiticity in the system. Eigenvalues of $\mathcal{H}_{\text{NH}}(\mathbf{k})$ are $\pm E_{\text{NH}}(\mathbf{k})$, where

$$E_{\text{NH}}(\mathbf{k}) = [(t_1^2 - t_2^2)\{d_1^2(\mathbf{k}) + d_2^2(\mathbf{k})\} + d_3^2(\mathbf{k})]^{1/2}. \quad (3)$$

They are purely real when $t_2 < t_1$. Then $\mathcal{H}_{\text{NH}}(\mathbf{k})$ fosters non-trivial topological invariants, when $|m_0/t_0| < 2$.

NH QAH I. For $\Gamma_j = \tau_j$ and $\Psi_{\mathbf{k}}^\dagger = (c_+, c_-)(\mathbf{k})$, where $c_\tau(\mathbf{k})$ is the fermionic annihilation operator on orbital with parity eigenvalue $\tau = \pm$ and momentum \mathbf{k} , we find the NH incarnation of the Qi-Wu-Zhang model for QAH I [63]. The Pauli matrices $\{\tau_\mu\}$ act on the orbital indices. Then the NH operator takes the form $\mathcal{H}_{\text{NH}}(\mathbf{k}) = \sum_{j=1}^3 d_j^{\text{NH}}(\mathbf{k}) \tau_j$, with $d_1^{\text{NH}}(\mathbf{k}) = t_1 \sin(k_x a) - it_2 \sin(k_y a)$, $d_2^{\text{NH}}(\mathbf{k}) = t_1 \sin(k_y a) + it_2 \sin(k_x a)$, and $d_3^{\text{NH}}(\mathbf{k}) = d_3(\mathbf{k})$. The first Chern number of $\mathcal{H}_{\text{NH}}(\mathbf{k})$ reads [58, 64]

$$C = \int_{\text{BZ}} \frac{d^2 \mathbf{k}}{4\pi} \left[\partial_{k_x} \hat{\mathbf{d}}^{\text{NH}}(\mathbf{k}) \times \partial_{k_y} \hat{\mathbf{d}}^{\text{NH}}(\mathbf{k}) \right] \cdot \hat{\mathbf{d}}^{\text{NH}}(\mathbf{k}), \quad (4)$$

where $\hat{\mathbf{d}}^{\text{NH}}(\mathbf{k}) = \mathbf{d}^{\text{NH}}(\mathbf{k})/\sqrt{[\mathbf{d}^{\text{NH}}(\mathbf{k})]^2}$. The momentum integral is performed over the first BZ. For any $|t_2| < 1$,




FIG. 4. Thermal Hall conductivity (κ_{xy}) of a NH $d + id$ pairing for (a) a fixed t_2 as a function of m_0 and (b) two fixed values of m_0 with varying t_2 , showing $\kappa_{xy} = C$ (in units of κ_0), where $C = 2$. Its spin Hall conductivity (σ_{xy}^{sp}) for (c) a fixed t_2 as a function of m_0 and (d) two fixed values of m_0 with varying t_2 shows $\sigma_{xy}^{sp} = C$ (in units of σ_0^{sp}).

we find $C = -1$ ($+1$) for $0 < m_0/t_0 < 2$ ($-2 < m_0/t_0 < 0$). Next, we discuss the ramification of non-trivial C on the quantized σ_{xy} , computed in a six-terminal setup (Fig. 1), detailed in the Supplemental Material [65].

A voltage gradient is applied between two horizontal leads, yielding a longitudinal electrical current (I_{el}) between them, with no current flowing between the transverse leads, serving as the voltage probes. The current-voltage relation is then given by $\mathbf{I}_{el} = \mathbf{G}_{el}\mathbf{V}$, with $\mathbf{V}^\top = (-\Delta V/2, V_2, V_3, \Delta V/2, V_5, V_6)$ and $\mathbf{I}_{el}^\top = (I_{el}, 0, 0, -I_{el}, 0, 0)$. Upon computing the conductance matrix \mathbf{G}_{el} using Kwant [59], containing only the transmission blocks of the scattering matrix, we extract different voltages from the current-voltage relation. Subsequently, we compute the transverse electrical resistance $R_{xy}^{el} = (V_2 + V_3 - V_5 - V_6)/(2I_{el})$, and find $\sigma_{xy} = 1/R_{xy}^{el} = C$ (in units of e^2/h). See Fig. 2(a) and (b).

NH QSHI. A NH generalization of the Bernevig-Hughes-Zhang model for QSHI is realized with $\mathbf{\Gamma} = (\Gamma_{01}, \Gamma_{32}, \Gamma_{03})$, where $\Gamma_{\mu\nu} = \sigma_\mu \tau_\nu$, and the Pauli matrices $\{\sigma_\mu\}$ operate on the spin index [5]. The four-component spinor reads as $\Psi_{\mathbf{k}}^\top = (c_\uparrow^\dagger, c_-^\dagger, c_\downarrow^\dagger, c_-^\dagger)(\mathbf{k})$, where $c_\tau^\sigma(\mathbf{k})$ is the fermionic annihilation operator with parity $\tau = \pm$, spin projection $\sigma = \uparrow, \downarrow$ in the z direction, and momentum \mathbf{k} . Following the same steps, highlighted for the NH QAHI model, we compute the Chern number for individual spin components, given by C_\uparrow and C_\downarrow . Within the topological regime, we find that the total Chern number of the model is $C_{total} = C_\uparrow + C_\downarrow = 0$. Nevertheless, we can define an invariant, the spin Chern number $C_{sp} = |C_\uparrow - C_\downarrow|$ which is non-trivial in the entire topological regime ($|m_0/t_0| < 2$ and $|t_2| < 1$), given by $C_{sp} = 2$.

The non-trivial C_{sp} leaves its footprint on the quantized G_{xx} for which we employ the same six-terminal arrangement, previously discussed for the NH QAHI. In this case, $\sigma_{xy} = 0$ as $C_{total} = 0$ therein. But, from the same current-voltage relationship, we now compute the longitudinal electrical resistance $R_{xx} = (V_3 - V_2)/I_{el} = (V_5 - V_6)/I_{el}$, in turn yielding $G_{xx} = R_{xx}^{-1} = C_{sp}$ (in units of e^2/h), as shown in Fig. 2(c) and (d).

NH $p + ip$ pairing. For a NH $p + ip$ paired state among spinless or spin-polarized fermions $\Psi_{\mathbf{k}}^\top = (c_{\mathbf{k}}, c_{-\mathbf{k}}^\dagger)$, where $c_{\mathbf{k}}$ ($c_{\mathbf{k}}^\dagger$) is the fermionic annihilation (creation) operator with momentum \mathbf{k} . The Pauli matrices $\{\tau_\mu\}$ operate on the Nambu or particle-hole index [21]. The effective single-particle NH Bogoliubov de-Gennes operator $\mathcal{H}_{NH}(\mathbf{k})$ has the topological invariant C [Eq. (4)], which manifests via a half-quantized thermal Hall conductivity (κ_{xy}). Now, a thermal current (I_{th}) flows between two horizontal leads, held at fixed but different temperatures (Fig. 1). Four vertical or transverse leads serve as the temperature probe, with no thermal current flowing between them. The thermal current-temperature relation is a matrix equation $\mathbf{I}_{th} = \mathbf{A}_{th}\mathbf{T}$, where $\mathbf{I}_{th}^\top = (I_{th}, 0, 0, -I_{th}, 0, 0)$ and $\mathbf{T}^\top = (-\Delta T/2, T_2, T_3, \Delta T/2, T_5, T_6)$. The elements of \mathbf{A}_{th} are

$$A_{th,ij} = \int_0^\infty \frac{E^2}{T} \left(-\frac{\partial f(E, T)}{\partial E} \right) \left[\delta_{ij} \mu_j - \text{Tr}(\mathbf{t}_{ij}^\dagger \mathbf{t}_{ij}) \right] dE, \quad (5)$$

where $f(E, T) = 1/(1 + \exp[E/(k_B T)])$ is the Fermi-Dirac distribution function, μ_j denotes the number of propagating modes in the j th lead, \mathbf{t}_{ij} is the transmission part of the scattering matrix between the leads i and j , and the trace (Tr) is taken over the conducting channels. From \mathbf{A}_{th} , we extract temperatures at various leads. The transverse thermal resistance is then given by $R_{xy}^{th} = (T_2 + T_3 - T_5 - T_6)/(2I_{th})$ [60–62, 66], from which we compute $\kappa_{xy} = (R_{xy}^{th})^{-1}$ at $T = 0.01$, showing $\kappa_{xy} = \alpha C$ in units of κ_0 [Fig. 3(a) and (b)], where $\alpha = 1/2$ accounts for the Nambu doubling [Eq. (1)]. Thus our proposed model NH $p + ip$ paired state features a half-quantized κ_{xy} in the entire topological regime.

NH $p \pm ip$ pairing. This paired state occurs among spin-1/2 fermions, with $p + ip$ and $p - ip$ pairing symmetries for opposite spin projections. The associated four-component spinor is $\Psi_{\mathbf{k}} = (\psi_{\mathbf{k}}, \sigma_2 \psi_{-\mathbf{k}}^*)$ with $\psi_{\mathbf{k}} = (c_\uparrow, c_\downarrow)(\mathbf{k})$, where $c_\sigma(\mathbf{k})$ is the fermionic annihilation operator with spin projection $\sigma = \uparrow, \downarrow$ and momentum \mathbf{k} . The involved Γ matrices are $\Gamma_1 = \sigma_0 \tau_1$, $\Gamma_2 = \sigma_3 \tau_2$, and $\Gamma_3 = \sigma_0 \tau_3$. Similar to the situation in NH QSHI, $C_{total} = 0$ but $C_{sp} = 2$ for the NH $p \pm ip$ paired. We numerically confirm that now $\kappa_{xy} = 0$. Nevertheless, from the longitudinal thermal resistance $R_{xx}^{th} = (T_3 - T_2)/I_{th} = (T_6 - T_5)/I_{th}$, we find that $G_{xx}^{th} = (R_{xx}^{th})^{-1} = \alpha C_{sp} \kappa_0$ with $\alpha = 1/2$, as shown in Fig. 3(c) and (d).

NH $d + id$ pairing. In the same four-component Nambu-doubled spinor basis, a spin-singlet NH $d + id$ paired state




FIG. 5. Disorder averaged (a) electrical Hall conductivity $\langle \sigma_{xy} \rangle$ in QAH, (b) electrical longitudinal conductivity $\langle G_{xx} \rangle$ in QSH, (c) thermal Hall conductivity $\langle \kappa_{xy} \rangle$ in $p+ip$ paired state, (d) thermal longitudinal conductivity $\langle G_{xx}^{th} \rangle$ in $p \pm ip$ superconductor, and (e) $\langle \kappa_{xy} \rangle$ and (f) spin Hall conductivity $\langle \sigma_{xy}^{sp} \rangle$ in $d+id$ paired state in Hermitian ($t_2 = 0$) and NH ($t_2 = 0.1, 0.5$) systems. The error bars correspond to the saturated (after averaging over a large number of disorder realizations) standard deviations [65].

is realized with $d_1(\mathbf{k}) = t_1[\cos(k_x a) - \cos(k_y a)]$, $d_2(\mathbf{k}) = t_1 \sin(k_x a) \sin(k_y a)$, and $\Gamma_j = \sigma_0 \tau_j$ for $j = 1, 2, 3$ [21, 67]. Within the topological regime ($|m_0/t_0| < 2$ and $|t_2| < 1$), the Chern number of the corresponding NH operator for each spin projection is $C = 2$, and thus its total Chern number is $2C = 4$. This NH paired state accommodates $\kappa_{xy} = C$ (in units of κ_0), as shown in Fig. 4(a) and (b). The spin-charge separation allows this NH paired state to harbor quantized spin Hall conductivity (σ_{xy}^{sp}), for which a spin current passes through the scattering region due to a difference in the magnetic field bias between the two horizontal leads. Then magnetization (m_j) develops in the vertical leads, which we compute from the spin current-magnetization matrix relation $\mathbf{I}_{sp} = \mathbf{G}_{sp} \mathbf{M}$, where $\mathbf{I}_{sp}^T = (I_{sp}, 0, 0, -I_{sp}, 0, 0)$ and $\mathbf{M}^T = (-\Delta H/2, m_2, m_3, \Delta H/2, m_5, m_6)$, and \mathbf{G}_{sp} is the spin conductance matrix, which we extract using Kwant [62]. From its solutions, we compute the transverse spin Hall resistance $R_{xy}^{sp} = (m_2 + m_3 - m_5 - m_6)/(2I_{sp})$, subsequently yielding $\sigma_{xy}^{sp} = (R_{xy}^{sp})^{-1} = C$ (in units of σ_0^{sp}), as shown in Fig. 4(c) and (d).

Disorder. Finally, we unfold the effects of disorder on all the topological responses, discussed so far in clean NH systems. We consider only pointlike charge impurities,

the dominant source of elastic scattering in any real material. In the above NH systems, we introduce the terms $V(\mathbf{r})\tau_0$, $V(\mathbf{r})\sigma_0\tau_0$, $V(\mathbf{r})\tau_3$, $V(\mathbf{r})\sigma_0\tau_3$, and $V(\mathbf{r})\sigma_0\tau_3$, respectively, where $V(\mathbf{r})$ is uniformly and randomly distributed within the range $[-W/2, W/2]$ at each site of the scattering region and W denotes the strength of disorder. The results are shown in Fig. 5. All the topological transport quantities (disorder averaged) retain robust (half-)quantized values in the weak disorder regime, while they all vanish in the strong disorder regime. Furthermore, with the introduction of non-Hermiticity in the system as $t_1 \rightarrow \sqrt{t_1^2 - t_2^2}$ [Eq. (3)], fermions become weakly dispersive and the latter event occurs at weaker disorder, which can be seen by comparing the results for $t_2 = 0$ and 0.5. However, even $L = 2D = 180$ systems [Fig. 1] turn out to be insufficient to compare the results for different finite t_2 values. Numerical simulations in larger systems become too time expensive for us to pursue at this stage. Nevertheless, most of the results from Fig. 5 strongly suggest that with increasing t_2 , the disappearance of the topological responses occurs at weaker disorder.

Discussions. Here we identify a family of NH operators, devoid of NH skin effects, for planar topological insulators and superconductors that manifests the topological invariant through (half-)quantized electrical, thermal,

and spin transport quantities, closely mimicking the ones previously found in Hermitian systems [61, 62]. Even though the requisite six-terminal Hall bar arrangement is well-developed by now, controlled synthesis of NH quantum crystals remains far from reality. Nevertheless, optical lattices of neutral atoms constitute a promising testbed for our theoretical predictions, where a plethora of topological band engineering protocols has been proposed [68] and realized [69]. Simplicity of our construction, in which the NH operators result from nearest-neighbor hopping modulations, makes it achievable on such highly tunable platforms that also harbor superfluids (charge-neutral superconductors). In such systems, the Hall conductivity can be obtained from the “heating effect” [70], for example, which can also be employed to measure quantized longitudinal transports.

Acknowledgments. S.K.D. was supported by the Startup Grant of B.R. from Lehigh University. B.R. was supported by NSF CAREER Grant No. DMR-2238679. Portions of this research were conducted on Lehigh University’s Research Computing infrastructure partially supported by NSF Award No. 2019035. We thank Vladimir Juričić for critical reading of the manuscript.

-
- [1] M. Z. Hasan and C. L. Kane, Colloquium: Topological insulators, *Rev. Mod. Phys.* **82**, 3045 (2010).
- [2] X.-L. Qi and S.-C. Zhang, Topological insulators and superconductors, *Rev. Mod. Phys.* **83**, 1057 (2011).
- [3] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, Classification of topological quantum matter with symmetries, *Rev. Mod. Phys.* **88**, 035005 (2016).
- [4] C. L. Kane and E. J. Mele, Z_2 Topological Order and the Quantum Spin Hall Effect, *Phys. Rev. Lett.* **95**, 146802 (2005).
- [5] B. A. Bernevig, T. L. Hughes, and S.-C. Zhang, Quantum Spin Hall Effect and Topological Phase Transition in HgTe Quantum Wells, *Science* **314**, 1757 (2006).
- [6] L. Fu, C. L. Kane, and E. J. Mele, Topological Insulators in Three Dimensions, *Phys. Rev. Lett.* **98**, 106803 (2007).
- [7] L. Fu and C. L. Kane, Topological insulators with inversion symmetry, *Phys. Rev. B* **76**, 045302 (2007).
- [8] J. E. Moore and L. Balents, Topological invariants of time-reversal-invariant band structures, *Phys. Rev. B* **75**, 121306 (2007).
- [9] R. Roy, Topological phases and the quantum spin Hall effect in three dimensions, *Phys. Rev. B* **79**, 195322 (2009).
- [10] S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. W. Ludwig, Topological insulators and superconductors: tenfold way and dimensional hierarchy, *New J. Phys.* **12**, 065010 (2010).
- [11] A. Kitaev, Periodic table for topological insulators and superconductors, *AIP Conf. Proc.* **1134**, 22 (2009).
- [12] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, Classification of topological insulators and superconductors in three spatial dimensions, *Phys. Rev. B* **78**, 195125 (2008).
- [13] L. Fu, Topological Crystalline Insulators, *Phys. Rev. Lett.* **106**, 106802 (2011).
- [14] R.-J. Slager, A. Mesaros, V. Juričić, and J. Zaanen, The space group classification of topological band-insulators, *Nat. Phys.* **9**, 98 (2012).
- [15] K. Shiozaki and M. Sato, Topology of crystalline insulators and superconductors, *Phys. Rev. B* **90**, 165114 (2014).
- [16] B. Bradlyn, L. Elcoro, J. Cano, M. G. Vergniory, Z. Wang, C. Felser, M. I. Aroyo, and B. A. Bernevig, Topological quantum chemistry, *Nature (London)* **547**, 298 (2017).
- [17] G. E. Volovik, *The Universe in a Helium Droplet* (Oxford University Press, Oxford, UK, 2009).
- [18] Y. Ando and L. Fu, Topological Crystalline Insulators and Topological Superconductors: From Concepts to Materials, *Annu. Rev. Condens. Matter Phys.* **6**, 361 (2015).
- [19] M. Sato and Y. Ando, Topological superconductors: a review, *Rep. Prog. Phys.* **80**, 076501 (2017).
- [20] T. Senthil, J. B. Marston, and M. P. A. Fisher, Spin quantum Hall effect in unconventional superconductors, *Phys. Rev. B* **60**, 4245 (1999).
- [21] N. Read and D. Green, Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect, *Phys. Rev. B* **61**, 10267 (2000).
- [22] L. E. F. F. Torres, Perspective on topological states of non-Hermitian lattices, *J. Phys. Mater.* **3**, 014002 (2019).
- [23] A. Ghatak and T. Das, New topological invariants in non-Hermitian systems, *J. Phys.: Condens. Matter* **31**, 263001 (2019).
- [24] E. J. Bergholtz, J. C. Budich, and F. K. Kunst, Exceptional topology of non-Hermitian systems, *Rev. Mod. Phys.* **93**, 015005 (2021).
- [25] K. Esaki, M. Sato, K. Hasebe, and M. Kohmoto, Edge states and topological phases in non-Hermitian systems, *Phys. Rev. B* **84**, 205128 (2011).
- [26] S.-D. Liang and G.-Y. Huang, Topological invariance and global Berry phase in non-Hermitian systems, *Phys. Rev. A* **87**, 012118 (2013).
- [27] S. Yao and Z. Wang, Edge States and Topological Invariants of Non-Hermitian Systems, *Phys. Rev. Lett.* **121**, 086803 (2018).
- [28] S. Yao, F. Song, and Z. Wang, Non-Hermitian Chern Bands, *Phys. Rev. Lett.* **121**, 136802 (2018).
- [29] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, Topological Phases of Non-Hermitian Systems, *Phys. Rev. X* **8**, 031079 (2018).
- [30] K. Kawabata, K. Shiozaki, and M. Ueda, Anomalous helical edge states in a non-Hermitian Chern insulator, *Phys. Rev. B* **98**, 165148 (2018).
- [31] H. Shen, B. Zhen, and L. Fu, Topological Band Theory for Non-Hermitian Hamiltonians, *Phys. Rev. Lett.* **120**, 146402 (2018).
- [32] F. K. Kunst, E. Edvardsson, J. C. Budich, and E. J. Bergholtz, Biorthogonal Bulk-Boundary Correspondence in Non-Hermitian Systems, *Phys. Rev. Lett.* **121**, 026808 (2018).
- [33] K. Kawabata, S. Higashikawa, Z. Gong, Y. Ashida, and M. Ueda, Topological unification of time-reversal and particle-hole symmetries in non-Hermitian physics, *Nat. Commun.* **10**, 297 (2019).
- [34] H. Zhou and J. Y. Lee, Periodic table for topological bands with non-Hermitian symmetries, *Phys. Rev. B* **99**, 235112 (2019).

- [35] K. Yokomizo and S. Murakami, Non-Bloch Band Theory of Non-Hermitian Systems, *Phys. Rev. Lett.* **123**, 066404 (2019).
- [36] K. Kawabata, K. Shiozaki, M. Ueda, and M. Sato, Symmetry and topology in non-hermitian physics, *Phys. Rev. X* **9**, 041015 (2019).
- [37] C. H. Lee and R. Thomale, Anatomy of skin modes and topology in non-Hermitian systems, *Phys. Rev. B* **99**, 201103 (2019).
- [38] N. Okuma, K. Kawabata, K. Shiozaki, and M. Sato, Topological Origin of Non-Hermitian Skin Effects, *Phys. Rev. Lett.* **124**, 086801 (2020).
- [39] K. Zhang, Z. Yang, and C. Fang, Correspondence between Winding Numbers and Skin Modes in Non-Hermitian Systems, *Phys. Rev. Lett.* **125**, 126402 (2020).
- [40] D. S. Borgnia, A. J. Kruchkov, and R.-J. Slager, Non-Hermitian Boundary Modes and Topology, *Phys. Rev. Lett.* **124**, 056802 (2020).
- [41] H. C. Wu, L. Jin, and Z. Song, Nontrivial topological phase with a zero Chern number, *Phys. Rev. B* **102**, 035145 (2020).
- [42] X.-Q. Sun, P. Zhu, and T. L. Hughes, Geometric Response and Disclination-Induced Skin Effects in Non-Hermitian Systems, *Phys. Rev. Lett.* **127**, 066401 (2021).
- [43] W. Zhu, W. X. Teo, L. Li, and J. Gong, Delocalization of topological edge states, *Phys. Rev. B* **103**, 195414 (2021).
- [44] A. Panigrahi, R. Moessner, and B. Roy, Non-Hermitian dislocation modes: Stability and melting across exceptional points, *Phys. Rev. B* **106**, L041302 (2022).
- [45] S. Manna and B. Roy, Inner skin effects on non-Hermitian topological fractals, *Commun. Phys.* **6**, 10 (2023).
- [46] R. Chaturvedi, V. Könye, E. M. Hankiewicz, J. van den Brink, and I. Cosma Fulga, Non-Hermitian topology of transport in Chern insulators, [arXiv:2406.14303](https://arxiv.org/abs/2406.14303).
- [47] K. v. Klitzing, G. Dorda, and M. Pepper, New Method for High-Accuracy Determination of the Fine-Structure Constant Based on Quantized Hall Resistance, *Phys. Rev. Lett.* **45**, 494 (1980).
- [48] M. Könnig, S. Wiedmann, C. Brüne, A. Roth, H. Buhmann, L. W. Molenkamp, X.-L. Qi, and S.-C. Zhang, Quantum Spin Hall Insulator State in HgTe Quantum Wells, *Science* **318**, 766 (2007).
- [49] I. Knez, R.-R. Du, and G. Sullivan, Evidence for Helical Edge Modes in Inverted InAs/GaSb Quantum Wells, *Phys. Rev. Lett.* **107**, 136603 (2011).
- [50] R. Yu, W. Zhang, H.-J. Zhang, S.-C. Zhang, X. Dai, and Z. Fang, Quantized Anomalous Hall Effect in Magnetic Topological Insulators, *Science* **329**, 61 (2010).
- [51] C.-Z. Chang, J. Zhang, X. Feng, J. Shen, Z. Zhang, M. Guo, K. Li, Y. Ou, P. Wei, L.-L. Wang, Z.-Q. Ji, Y. Feng, S. Ji, X. Chen, J. Jia, X. Dai, Z. Fang, S.-C. Zhang, K. He, Y. Wang, L. Lu, X.-C. Ma, and Q.-K. Xue, Experimental Observation of the Quantum Anomalous Hall Effect in a Magnetic Topological Insulator, *Science* **340**, 167 (2013).
- [52] C.-Z. Chang, W. Zhao, D. Y. Kim, H. Zhang, B. A. Assaf, D. Heiman, S.-C. Zhang, C. Liu, M. H. W. Chan, and J. S. Moodera, High-precision realization of robust quantum anomalous Hall state in a hard ferromagnetic topological insulator, *Nature Mater* **14**, 473 (2015).
- [53] M. Banerjee, M. Heiblum, V. Umansky, D. E. Feldman, Y. Oreg, and A. Stern, Observation of half-integer thermal Hall conductance, *Nature (London)* **559**, 205 (2018).
- [54] Y. Kasahara, T. Ohnishi, Y. Mizukami, O. Tanaka, S. Ma, K. Sugii, N. Kurita, H. Tanaka, J. Nasu, Y. Motome, T. Shibauchi, and Y. Matsuda, Majorana quantization and half-integer thermal quantum Hall effect in a Kitaev spin liquid, *Nature (London)* **559**, 227 (2018).
- [55] S. K. Srivastav, M. R. Sahu, K. Watanabe, T. Taniguchi, S. Banerjee, and A. Das, Universal quantized thermal conductance in graphene, *Sci. Adv.* **5**, eaaw5798 (2019).
- [56] G. Le Breton, R. Delagrangé, Y. Hong, M. Garg, K. Watanabe, T. Taniguchi, R. Ribeiro-Palau, P. Roulleau, P. Roche, and F. D. Parmentier, Heat Equilibration of Integer and Fractional Quantum Hall Edge Modes in Graphene, *Phys. Rev. Lett.* **129**, 116803 (2022).
- [57] A. Altland and M. R. Zirnbauer, Nonstandard symmetry classes in mesoscopic normal-superconducting hybrid structures, *Phys. Rev. B* **55**, 1142 (1997).
- [58] D. J. Salib, S. K. Das, and B. Roy, Model non-Hermitian topological operators without skin effect, [arXiv:2309.12310](https://arxiv.org/abs/2309.12310).
- [59] C. W. Groth, M. Wimmer, A. R. Akhmerov, and X. Waintal, Kwant: a software package for quantum transport, *New Journal of Physics* **16**, 063065 (2014).
- [60] I. C. Fulga, Y. Oreg, A. D. Mirlin, A. Stern, and D. F. Mross, Temperature Enhancement of Thermal Hall Conductance Quantization, *Phys. Rev. Lett.* **125**, 236802 (2020).
- [61] S. K. Das, S. Manna, and B. Roy, Topologically distinct atomic insulators, *Phys. Rev. B* **108**, L041301 (2023).
- [62] S. K. Das and B. Roy, Quantized thermal and spin transports of dirty planar topological superconductors, *Phys. Rev. B* **109**, 195403 (2024).
- [63] X.-L. Qi, Y.-S. Wu, and S.-C. Zhang, Topological quantization of the spin Hall effect in two-dimensional paramagnetic semiconductors, *Phys. Rev. B* **74**, 085308 (2006).
- [64] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, Quantized Hall Conductance in a Two-Dimensional Periodic Potential, *Phys. Rev. Lett.* **49**, 405 (1982).
- [65] See Supplemental Material at XXX-XXXX for the details of transport calculations and saturation of transport responses with the number of disorder realizations.
- [66] W. Long, H. Zhang, and Q.-f. Sun, Quantum thermal Hall effect in graphene, *Phys. Rev. B* **84**, 075416 (2011).
- [67] R. B. Laughlin, Magnetic induction of $d_{x^2-y^2} + id_{xy}$ order in high- T_c superconductors, *Phys. Rev. Lett.* **80**, 5188 (1998).
- [68] D.-W. Zhang, Y.-Q. Zhu, Y. X. Zhao, H. Yan, and S.-L. Zhu, Topological quantum matter with cold atoms, *Adv. Phys.* **67**, 253 (2018).
- [69] N. R. Cooper, J. Dalibard, and I. B. Spielman, Topological bands for ultracold atoms, *Rev. Mod. Phys.* **91**, 015005 (2019).
- [70] L. Asteria, D. T. Tran, T. Ozawa, M. Tarnowski, B. S. Rem, N. Fläschner, K. Sengstock, N. Goldman, and C. Weitenberg, Measuring quantized circular dichroism in ultracold topological matter, *Nat. Phys.* **15**, 449 (2019).