

# JOINT TAIL OF RANDOMLY WEIGHTED SUMS UNDER GENERALIZED QUASI ASYMPTOTIC INDEPENDENCE

DIMITRIOS G. KONSTANTINIDES, CHARALAMPOS D. PASSALIDIS

**ABSTRACT.** In this paper we revisited the classical problem of max-sum equivalence of randomly (weighted) sums in two dimensions. In opposite to the most papers in literature, we consider that there exist some interdependence between the primary random variables, which is achieved by a combination of a new dependence structure with some two-dimensional heavy-tailed classes of distributions. Further we introduce a new approach in two-dimensional regular varying distributions, and we study some closure properties of this, and of other two-dimensional classes. Our results contain the finite-time ruin probability in a two-dimensional discrete time risk model.

*Keywords:* Two-dimensional heavy-tailed distributions, joint max-sum equivalence, closure properties, Breiman's Theorem, ruin probability, interdependence.

*Mathematics Subject Classification:* Primary 62P05 ; Secondary 60G70.

## 1. INTRODUCTION

Let  $X_1, \dots, X_n$ ,  $n \in \mathbb{N}$ , be real valued random variables, with distributions  $F_1, \dots, F_n$  respectively, and we consider their tails  $\bar{F}_i(x) = 1 - F_i(x) = \mathbf{P}[X_i > x]$ , with  $\bar{F}_i(x) > 0$  for any  $x \in \mathbb{R}$  and any  $i = 1, \dots, n$ . A well-known question for heavy-tailed distributions is the validity of the asymptotic relation

$$\mathbf{P}[S_n > x] \sim \sum_{i=1}^n \bar{F}_i(x), \quad (1.1)$$

as  $x \rightarrow \infty$ , where  $S_n := \sum_{i=1}^n X_i$ , or furthermore the validity of

$$\mathbf{P}[S_n > x] \sim \mathbf{P}\left[\bigvee_{i=1}^n S_i > x\right] \sim \mathbf{P}\left[\bigvee_{i=1}^n X_i > x\right] \sim \sum_{i=1}^n \bar{F}_i(x), \quad (1.2)$$

as  $x \rightarrow \infty$ , where

$$\bigvee_{i=1}^n S_i := \max_{1 \leq k \leq n} \sum_{i=1}^k X_i, \quad \bigvee_{i=1}^n X_i := \max_{1 \leq i \leq n} X_i, \quad (1.3)$$

For the study of the relations (1.1) or (1.2) two factors play crucial role. The first factor is the distribution class to which belong the  $F_1, \dots, F_n$  and the second one is the dependence structure among these random variables. For some papers on this topic see [31], [59], [68].

This question is extended in the case of randomly weighted sums, where now we have some non-negative, non-degenerated to zero random variables  $\Theta_1, \dots, \Theta_n$ , whence the relation

---

*Date:* August 5, 2024.

(1.2) becomes

$$\mathbf{P}[S_n^\Theta > x] \sim \mathbf{P}\left[\bigvee_{i=1}^n S_i^\Theta > x\right] \sim \mathbf{P}\left[\bigvee_{i=1}^n \Theta_i X_i > x\right] \sim \sum_{i=1}^n \mathbf{P}[\Theta_i X_i > x], \quad (1.4)$$

as  $x \rightarrow \infty$ , where

$$S_n^\Theta := \sum_{i=1}^n \Theta_i X_i, \quad \bigvee_{i=1}^n S_i^\Theta := \max_{1 \leq k \leq n} \sum_{i=1}^k \Theta_i X_i, \quad \bigvee_{i=1}^n \Theta_i X_i := \max_{1 \leq i \leq n} \Theta_i X_i, \quad (1.5)$$

It is easily to observe that if the random weights  $\Theta_1, \dots, \Theta_n$  are degenerated to unity, then we return back to relation (1.2).

In most literature the random weights  $\Theta_1, \dots, \Theta_n$  are arbitrarily dependent and either among the  $X_1, \dots, X_n$  is assumed some dependence structure but they are independent of the random weights, or is assumed some dependence structure between  $(\Theta_i, X_i)$ , for the same  $i = 1, \dots, n$ , but the  $X_1, \dots, X_n$  are independent. For some papers about the asymptotic behavior of the randomly weighted sums see in [60], [61] and [66].

Recently, there were several attempts to extend this problem in two dimensional set-up. Namely, was arisen the question about the validity of

$$\mathbf{P}[S_n > x, T_m > y] \sim \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[X_i > x, Y_j > y], \quad (1.6)$$

as  $(x, y) \rightarrow (\infty, \infty)$ , or as  $x \wedge y \rightarrow \infty$ , where  $T_m = \sum_{j=1}^m Y_j$ , (with  $m \in \mathbb{N}$ ) and the question about the weighted sums becomes

$$\mathbf{P}[S_n^\Theta > x, T_m^\Delta > y] \sim \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[\Theta_i X_i > x, \Delta_j Y_j > y], \quad (1.7)$$

as  $(x, y) \rightarrow (\infty, \infty)$ , or as  $x \wedge y \rightarrow \infty$ , where  $T_m^\Delta = \sum_{j=1}^m \Delta_j Y_j$ , or if are true the asymptotic equivalencies for the maxima of the sums in two dimensions and the jointly maxima, with the

$$\bigvee_{j=1}^m T_j, \quad \bigvee_{j=1}^m Y_j, \quad \bigvee_{j=1}^m T_j^\Delta, \quad \bigvee_{j=1}^m \Delta_j Y_j,$$

defined in a similar way as in relations (1.3) and (1.5). The reason, for which we are interested in the joint behavior of the randomly weighted sums is the dependence, that can appear among these sums. Obviously, if  $S_n^\Theta, T_m^\Theta$  are independent, then there is no reason to study (1.7), because the problem can be reduced in (1.4) for each sum.

Relation (1.7) was studied in several papers where was examined one of the following cases:

1) The main random variables  $X_1, \dots, X_n$  and  $Y_1, \dots, Y_m$  are independent sequences of random variables and the two sequences are also independent. The random weights  $\Theta_1, \dots, \Theta_n, \Delta_1, \dots, \Delta_m$  are arbitrarily dependent, non-negative and non-degenerated to zero random variables, and independent of the main variables  $X_1, \dots, X_n$  and  $Y_1, \dots, Y_m$ . Therefore, the dependence between the two sums  $S_n^\Theta, T_m^\Delta$  comes from the random weights.

2) The random weights are still as in case 1), but the main random variables have dependence structure. Namely, the  $X_1, \dots, X_n$  have some dependence structure and the

$Y_1, \dots, Y_m$  have some dependence structure, but the two sequences of  $\{X_i\}$  and  $\{Y_j\}$  are mutually independent, see for example in [65].

3) The random weights are still as in case 1), but the main random variables  $\{X_i\}$  and  $\{Y_j\}$  represent each a sequence of independent random variables, but the pair  $(X_i, Y_i)$  has some dependence structure for any  $i = 1, \dots, n \wedge m$ .

In case 3) there exist another distinction between the papers, in these where the dependence among pairs  $(X_i, Y_i)$  is 'weak', that means it is close to asymptotic independence, see for example [46], and in those with arbitrary dependence among the pairs  $(X_i, Y_i)$ , see for example [9], however in most of the cases there exist assumption of identically distributed random pairs, as for example in [56] and [57].

In [36] was introduced the generalized tail asymptotic independence (GTAI), as an attempt to merge the 2) and 3) dependence structures, however as result was found that GTAI covers the case when each pair  $(X_i, Y_i)$  follows a special form of weak dependence. Further in [38], assuming the sums restricted by the case when  $m = n$ , was established the asymptotic

$$\begin{aligned} \mathbf{P}[S_n > x, T_n > y] &\sim \mathbf{P}\left[\bigvee_{i=1}^n S_i > x, \bigvee_{j=1}^n T_j > y\right] \sim \mathbf{P}\left[\bigvee_{i=1}^n X_i > x, \bigvee_{j=1}^n Y_j > y\right] \\ &\sim \sum_{i=1}^n \sum_{j=1}^n \mathbf{P}[X_i > x, Y_j > y], \end{aligned} \quad (1.8)$$

as  $x \wedge y \rightarrow \infty$ , under GTAI dependence structure (and TAI dependence structure on each sequence) and under some two-dimensional distribution class. Furthermore, under some conditions on the random weights, that are independent of main random variables, the asymptotic relation (1.8) is still true.

In this work we have two aims.

1) to establish partially relation (1.8) for some greater dependence structure, which was possible reducing the set-up from class  $(\mathcal{D} \cap \mathcal{L})^{(2)}$  to class  $\mathcal{C}^{(2)}$ . Further in our results it is not necessary  $n = m$ .

2) to generalize the conditions for the random weights, staying always in frame of weighted form of (1.8). This way, we generalize both [38, Th. 6.1] and Theorem 3.1 in this paper.

We plan this paper as follows. In section 2 we give some preliminary results for the distributions with heavy tails, with one or two dimensions and we introduce the class of regular variation in two dimensions. After depicting some dependence structures with some known results, a new dependence structure is introduced. Next, in section 3 we present the main result for the jointly asymptotic behavior of the random sums, together with some preliminary lemmas. In section 4 we study some closure properties of our two-dimensional classes, and we give some new results both in one and two-dimensional cases. Finally in section 5, similarly we provide the results for the randomly weighted sums as in case of dependence as also in case of GTAI structure, and we discuss the applications in ruin probabilities over finite time in a two-dimensional risk model with discrete time with stochastic discount factors.

## 2. PRELIMINARIES

For two real numbers  $x, y$  we denote  $x^+ := \max\{x, 0\}$ ,  $x \wedge y := \min\{x, y\}$  and  $x \vee y := \max\{x, y\}$ . With  $\mathbf{1}_A$  we symbolized the indicator function of some event  $A$ . The joint distribution of two random variables  $X, Y$  is given by  $\mathbf{F}(x, y) = \mathbf{P}[X \leq x, Y \leq y]$ . As

tail of the joint distribution we take the joint excess, namely  $\bar{\mathbf{F}}(x, y) = \bar{\mathbf{F}}_1(x, y) := \mathbf{P}[X > x, Y > y]$ , where for sake of easiness, for any vector  $\mathbf{b} = (b_1, b_2)$  we write  $\bar{F}_{\mathbf{b}}(x, y) := \mathbf{P}[X > b_1 x, Y > b_2 y]$ .

In what follows we use the notations: for two uni-variate positive functions  $f_1$  and  $g_1$ , with  $f_1(x) \sim g_1(x)$ , as  $x \rightarrow \infty$ , we mean that

$$\lim_{x \rightarrow \infty} \frac{f_1(x)}{g_1(x)} = 1,$$

with  $f_1(x) \lesssim g_1(x)$ , or  $g_1(x) \gtrsim f_1(x)$ , as  $x \rightarrow \infty$ , we mean that

$$\limsup_{x \rightarrow \infty} \frac{f_1(x)}{g_1(x)} \leq 1,$$

with  $f_1(x) = o[g_1(x)]$ , or  $f_1(x) = g_1(x) o(1)$ , as  $x \rightarrow \infty$ , we mean that

$$\lim_{x \rightarrow \infty} \frac{f_1(x)}{g_1(x)} = 0,$$

while with  $f_1(x) = O[g_1(x)]$ , or  $f_1(x) = g_1(x) O(1)$ , as  $x \rightarrow \infty$ , we mean that

$$\limsup_{x \rightarrow \infty} \frac{f_1(x)}{g_1(x)} < \infty,$$

and with  $f_1(x) \asymp g_1(x)$ , as  $x \rightarrow \infty$ , we mean that

$$f_1(x) = O[g_1(x)] \quad g_1(x) = O[f_1(x)],$$

as  $x \rightarrow \infty$ .

If there exist two positive two-variate functions  $f_2$  and  $g_2$ , the corresponding limit relations come as extension of the one-variate with the restriction that the limits of two-variate functions hold for  $x \wedge y \rightarrow \infty$ . For example, with  $f_2(x, y) \sim g_2(x, y)$ , as  $x \wedge y \rightarrow \infty$ , we mean that

$$\lim_{x \wedge y \rightarrow \infty} \frac{f_2(x, y)}{g_2(x, y)} = 1,$$

and  $f_2(x, y) = o[g_2(x, y)]$ , as  $x \wedge y \rightarrow \infty$ , if

$$\lim_{x \wedge y \rightarrow \infty} \frac{f_2(x, y)}{g_2(x, y)} = 0.$$

**2.1. Heavy-tailed Distributions.** The heavy-tailed distributions recently attract more interest of applied and theoretical probability community. For example in Risk Theory, Financial Mathematics, Risk Management, Branching Processes, Lévy Processes present the domains of applications with heavy tails, see for example [1], [2], [3], [53], [23], [39]. We give some of the most important classes of heavy-tailed distributions, with some of these classes to be used later, together with their properties.

We say that a distribution  $F$  is heavy tailed, symbolically  $F \in \mathcal{H}$  if for any  $\varepsilon > 0$  holds

$$\int_{-\infty}^{\infty} e^{\varepsilon y} F(dy) = \infty.$$

We say that a distribution  $F$  is long tailed, symbolically  $F \in \mathcal{L}$  if for any (or equivalently for some)  $a > 0$ , holds

$$\lim_{x \rightarrow \infty} \frac{\overline{F}(x-a)}{\overline{F}(x)} = 1.$$

It is well-known that  $F \in \mathcal{L}$  if and only if there exists some function  $a : [0, \infty) \rightarrow [0, \infty)$  such that  $a(x) \rightarrow \infty$ ,  $a(x) = o(x)$  and  $\overline{F}[x \pm a(x)] \sim \overline{F}(x)$ , as  $x \rightarrow \infty$ . Function  $a(x)$  is called insensitivity function for distribution  $F$ , see in [32] or in [24].

Let remind that for two random variables  $X_1$  and  $X_2$  with distributions  $F_1$  and  $F_2$  respectively, the distribution of the sum is defined as  $F_{X_1+X_2}(x) := \mathbf{P}[X_1 + X_2 \leq x]$  and its tail  $\overline{F}_{X_1+X_2}(x) := \mathbf{P}[X_1 + X_2 > x]$ . If the random variables are independent then we denote  $F_1 * F_2$  instead of  $F_{X_1+X_2}$ , which is called convolution of  $F_1$  and  $F_2$ . A distribution  $F$  with support  $\mathbb{R}_+$  is called subexponential, symbolically  $F \in \mathcal{S}$ , if

$$\lim_{x \rightarrow \infty} \frac{\overline{F^{*n}}(x)}{\overline{F}(x)} = n,$$

for any (or equivalently, for some) integer  $n \geq 2$ . With  $F^{*n}$  we depict the  $n$ -th convolution power of  $F$ . Both classes  $\mathcal{L}$  and  $\mathcal{S}$  were defined in [13].

If a distribution  $F$  has support  $\mathbb{R}$ , then we say that it is subexponential on the real axis, symbolically  $F \in \mathcal{S}$ , if  $F_+(x) \in \mathcal{S}$ , where  $F_+(x) := F(x) \mathbf{1}_{\{x>0\}}$ , see [51]. Therefore, if  $X_1, \dots, X_n$  are independent identically distributed real random variables with common distribution  $F \in \mathcal{S}$ , we find relation (1.1) and by elementary inequalities we get also (1.2). This is a main reason to make class  $\mathcal{S}$  important, see for applications in risk theory in [29], [44], among others.

Class  $\mathcal{D}$  was introduced in [21] as extension of the regular variation. We say that a distribution  $F$  belongs to the class of dominatedly varying distributions, symbolically  $F \in \mathcal{D}$ , if for any or for some  $b \in (0, 1)$  holds

$$\limsup_{x \rightarrow \infty} \frac{\overline{F}(bx)}{\overline{F}(x)} < \infty.$$

It is well-known that  $\mathcal{D} \not\subseteq \mathcal{S}$ ,  $\mathcal{S} \not\subseteq \mathcal{D}$  and  $\mathcal{D} \cap \mathcal{S} = \mathcal{D} \cap \mathcal{L} \subsetneq \mathcal{H}$ , see [27].

Another distribution class with heavy tails is class  $\mathcal{C}$ , of consistently varying distributions. We say that  $F$  is consistently varying and we write  $F \in \mathcal{C}$  if

$$\lim_{z \uparrow 1} \limsup_{x \rightarrow \infty} \frac{\overline{F}(zx)}{\overline{F}(x)} = 1.$$

Further, we say that a distribution  $F$  belongs to the class of regularly varying distributions, with index  $\alpha > 0$ , symbolically  $F \in \mathcal{R}_{-\alpha}$ , if holds

$$\lim_{x \rightarrow \infty} \frac{\overline{F}(tx)}{\overline{F}(x)} = t^{-\alpha},$$

for any  $t > 0$ . It is well-known that  $\mathcal{R} := \bigcup_{\alpha>0} \mathcal{R}_{-\alpha} \subsetneq \mathcal{C} \subsetneq \mathcal{D} \cap \mathcal{L} \subsetneq \mathcal{S} \subsetneq \mathcal{L} \subsetneq \mathcal{H}$ , see for example [32], [41].

Next for any distribution with upper unbounded support, we bring up the upper and lower Matuszewska indexes  $\alpha_F, \beta_F$  respectively, introduced in [48], which have important role on

the characterization of heavy-tailed and related distributions

$$J_F^+ := \inf \left\{ -\frac{\ln \bar{F}_*(v)}{\ln v} : v > 1 \right\} = -\lim_{y \rightarrow \infty} \frac{\ln \bar{F}_*(y)}{\ln y},$$

$$J_F^- := \sup \left\{ -\frac{\ln \bar{F}^*(v)}{\ln v} : v > 1 \right\} = -\lim_{y \rightarrow \infty} \frac{\ln \bar{F}^*(y)}{\ln y}.$$

where

$$\bar{F}_*(v) := \liminf_{x \rightarrow \infty} \frac{\bar{F}(vx)}{\bar{F}(x)}, \quad \bar{F}^*(v) := \limsup_{x \rightarrow \infty} \frac{\bar{F}(vx)}{\bar{F}(x)},$$

for any  $v > 1$ . The following inequalities hold  $0 \leq J_F^- \leq J_F^+$ . We have the equivalences:  $F \in \mathcal{D}$  if and only if  $J_F^+ < \infty$  and  $F \in \mathcal{R}_{-\alpha}$  for some  $\alpha > 0$  if and only if  $J_F^+ = J_F^- = \alpha$ .

**2.2. Two-dimensional Distribution Classes with Heavy Tails.** Several attempts appeared recently for the modelling of extreme events in many dimensions. The reason was the dependence structures among these events, that play crucial role on final outcome. Although the multivariate regular variation (*MRV*) is a well established multivariate extension of the regular variation to many dimensions, this is NOT the case for the rest distribution classes. The most popular attempts for multi-variate distribution classes are referred to subexponential distributions and to long tailed distributions. In literature we find at least four different definitions to multi-variate subexponential distributions, see [14], [38], [50], [54]. However, none has dominated at moment, although in [54] there exists an application to continuous time multivariate risk model.

In the present work we follow the two-dimensional definitions from the [38], and after reminding the definitions of  $\mathcal{L}^{(2)}$ ,  $\mathcal{S}^{(2)}$ ,  $\mathcal{D}^{(2)}$ ,  $\mathcal{C}^{(2)}$ , we introduce the two-dimensional regular variation with different way, than in case of *MRV* or of non-standard *MRV*.

The characterization of the bivariate classes is based on the joint excess of random variables, as also on the marginal distributions. This makes the extension of one-dimensional classes more direct and in the same time permits arbitrary dependence structure among the components. Further, with the notation  $\mathbf{a} = (a_1, a_2) > (0, 0)$ , we have in mind that  $\mathbf{a} \in [0, \infty)^2 \setminus \{\mathbf{0}\}$ , except it is referred differently.

Let consider initially the two-dimensional class of long tailed distributions. We say that a random pair  $(X, Y)$  with marginal distributions  $F, G$ , belongs to the class of bivariate long tailed, symbolically  $\mathbf{F} := (F, G) \in \mathcal{L}^{(2)}$ , if

- (1)  $F \in \mathcal{L}, G \in \mathcal{L}$ ,
- (2)

$$\lim_{x \wedge y \rightarrow \infty} \frac{\bar{\mathbf{F}}_1(x - a_1, y - a_2)}{\bar{\mathbf{F}}_1(x, y)} = \lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X > x - a_1, Y > y - a_2]}{\mathbf{P}[X > x, Y > y]} = 1,$$

for any, or equivalently, for all  $\mathbf{a} = (a_1, a_2) > (0, 0)$ .

As in one-dimensional case there exists insensitivity function, in two-dimensional case also exists  $\mathbf{a}$ -jointly insensitivity function. Concretely, for a pair of two non-decreasing functions  $a_F(x), a_G(y) > 0$ , for any  $x, y > 0$ , the two-dimensional distribution  $\mathbf{F} = (F, G)$ , with infinite right endpoints for both marginal distributions, is called  $(a_F, a_G)$ -joint insensitivity

distribution, if the following is true

$$\begin{aligned} & \sup_{|a_1| \leq a_F(x), |a_2| \leq a_G(y)} |\mathbf{P}[X > x - a_1, Y > y - a_2] - \mathbf{P}[X > x, Y > y]| \\ & = o(\mathbf{P}[X > x, Y > y]), \end{aligned} \quad (2.1)$$

as  $x \wedge y \rightarrow \infty$ . From [38] we find that if  $(F, G) \in \mathcal{L}^{(2)}$ , then there exists some bivariate function  $(a_F(x), a_G(y))$ , such that relation (2.1) holds. Furthermore, this means that

$$\lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X > x \pm a_F(x), Y > y \pm a_G(y)]}{\mathbf{P}[X > x, Y > y]} = 1,$$

but this function is NOT making  $a$ -insensitive the marginal distributions.

We say that the random pair  $(X, Y)$ , with marginal distributions  $F$  and  $G$  respectively, belongs to the class of bivariate subexponential distributions, symbolically  $(F, G) \in \mathcal{S}^{(2)}$ , if

- (1)  $F \in \mathcal{S}$  and  $G \in \mathcal{S}$ .
- (2)  $(F, G) \in \mathcal{L}^{(2)}$ .
- (3) It holds

$$\lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_1 + X_2 > x, Y_1 + Y_2 > y]}{\mathbf{P}[X > x, Y > y]} = 2^2, \quad (2.2)$$

where  $X_1$  and  $X_2$  are independent and identically distributed and  $Y_1$  and  $Y_2$  are independent and identically distributed, with distributions  $F$  and  $G$  respectively.

**Remark 2.1.** In case of  $d$ -variate distribution relation (2.2) becomes

$$\lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_{1,1} + X_{1,2} > x_1, \dots, X_{d,1} + X_{d,2} > x_d]}{\mathbf{P}[X_{1,1} > x_1, \dots, X_{d,1} > x_d]} = 2^d. \quad (2.3)$$

Another bivariate distribution class is the  $\mathcal{D}^{(2)}$  of the bivariate dominatedly varying distributions. For a random pair  $(X, Y)$  with marginal distributions  $F, G$ , we write  $(F, G) \in \mathcal{D}^{(2)}$ , if

- (1)  $F \in \mathcal{D}, G \in \mathcal{D}$ ,
- (2)

$$\limsup_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{F}}_{\mathbf{b}}(x, y)}{\overline{\mathbf{F}}_{\mathbf{1}}(x, y)} = \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X > b_1 x, Y > b_2 y]}{\mathbf{P}[X > x, Y > y]} < \infty,$$

for any, or equivalently for some,  $\mathbf{b} = (b_1, b_2) \in (0, 1)^2$  and  $\mathbf{1} = (1, 1)$ .

Let us denote  $(\mathcal{D} \cap \mathcal{L})^{(2)} := \mathcal{D}^{(2)} \cap \mathcal{L}^{(2)}$ .

Next, we say that the random pair  $(X, Y)$  with marginal distributions  $F, G$ , belongs to the class of bivariate consistently varying distributions, symbolically  $(F, G) \in \mathcal{C}^{(2)}$ , if

- (1)  $F \in \mathcal{C}, G \in \mathcal{C}$ ,
- (2)

$$\lim_{\mathbf{z} \uparrow \mathbf{1}} \limsup_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{F}}_{\mathbf{z}}(x, y)}{\overline{\mathbf{F}}_{\mathbf{1}}(x, y)} = \lim_{\mathbf{z} \uparrow \mathbf{1}} \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X > z_1 x, Y > z_2 y]}{\mathbf{P}[X > x, Y > y]} = 1, \quad (2.4)$$

with  $\mathbf{z} = (z_1, z_2)$  and  $\mathbf{1} = (1, 1)$ .

We can see that  $\mathcal{C}^{(2)} \subsetneq (\mathcal{D} \cap \mathcal{L})^{(2)}$ , see [38, Th. 2.1].

Now we have a new distribution class, which represents a new kind of two-variate regular variation. Except the definition we comment about the advantages and disadvantages in relation with the classical multivariate regular variation, introduced in [28].

**Definition 2.1.** *We say that the random pair  $(X, Y)$  with marginal distributions  $F, G$  respectively, follows two-variate regular variation, symbolically  $\mathbf{F} \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , with  $0 < \alpha_1, \alpha_2 < \infty$ , if*

- (1)  $F \in \mathcal{R}_{-\alpha_1}, G \in \mathcal{R}_{-\alpha_2}$ ,
- (2)

$$\lim_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{F}}_{\mathbf{t}}(x, y)}{\overline{\mathbf{F}}_{\mathbf{1}}(x, y)} = \lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X > t_1 x, Y > t_2 y]}{\mathbf{P}[X > x, Y > y]} = t_1^{-\alpha_1} t_2^{-\alpha_2}, \quad (2.5)$$

for any  $\mathbf{t} = (t_1, t_2)$ , with  $t_1, t_2 > 0$  and  $\mathbf{1} = (1, 1)$ .

In this definition, for the case  $\alpha_1 = \alpha_2$ , we say that we have typical, two-variate, regular variation, while in opposite case  $\alpha_1 \neq \alpha_2$ , we have a non-typical, two-variate, regular variation.

**Remark 2.2.** *Let restrict ourselves to two-dimensional case, say the BRV, for the depiction of the multi-variate regular variation. If the normalizing functions are*

$$U_X(x) = \left(\frac{1}{F}\right)^{\leftarrow}(x), \quad U_Y(x) = \left(\frac{1}{G}\right)^{\leftarrow}(x),$$

for  $x > 0$ , then the two-dimensional regular variation is defined as

$$\lim_{x \rightarrow \infty} x \mathbf{P} \left[ \left( \frac{X}{U_X(x)}, \frac{Y}{U_Y(x)} \right) \in \mathbb{B} \right] = \nu(\mathbb{B}),$$

for any Borel set  $\mathbb{B} \subsetneq [0, \infty]^2 \setminus \{\mathbf{0}\}$ , and the limit measure  $\nu$  is non-degenerated. Furthermore, it follows that the distributions  $F$  and  $G$  are regularly varying and the measure  $\nu$  is homogeneous in the sense of  $\nu(\mathbb{B}^\lambda) = \lambda^{-1} \nu(\mathbb{B})$ , where  $\mathbb{B}^\lambda = \{(\lambda^{1/\alpha_1} \kappa, \lambda^{1/\alpha_2} \mu) : (\kappa, \mu) \in \mathbb{B}\}$ , where  $\alpha_1, \alpha_2$  are the regular variation indexes of  $F, G$  respectively. If  $\alpha_1 = \alpha_2$ , then we have typical BRV.

The class MRV attracted the interest of several branches of applied probability and statistics, see for example [4], [7], [52].

Recently, risk theory gained popularity, as for example in [9], [62] in non-typical BRV, and for example in [12], [34], [67], in typical MRV.

Although, class BRV is well-established in standard case, there are several difficulties, coming from the non-standard case, that appear because of the variety of the normalizing functions. Usually, the joint behavior of randomly weighted sums from relations (1.6) and (1.7) is very difficult to be approached. The class  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$  tries to surpass this obstacle, but with loss of some dependence cases, as the asymptotic dependence, see Proposition 2.1 later in this section.

Let us present some examples for the building of the class  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ . At first, it is obvious that if  $F \in \mathcal{R}_{-\alpha_1}$  and  $G \in \mathcal{R}_{-\alpha_2}$ , with  $a_1, a_2 > 0$ , with their distributions stemming from independent random variables, then it holds  $\mathbf{F} \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ . An interesting case of dependence presents the strongly asymptotic independence, symbolically SAI, see [45], [46].



Without loss of generality, for sake of simplicity with less conditions, we consider two non-negative random variables  $X, Y$  with distributions  $F, G$  respectively. We say that the  $X, Y$  satisfy the SAI condition, if there exists a constant  $C \geq 0$  such that

$$\mathbf{P}[X > x, Y > y] = [C + o(1)] \overline{F}(x) \overline{G}(y), \quad (2.6)$$

as  $(x, y) \rightarrow (\infty, \infty)$ .

Under the previous conditions, if  $F \in \mathcal{R}_{-\alpha_1}$  and  $G \in \mathcal{R}_{-\alpha_2}$ , then for any  $t_1, t_2 > 0$  it holds

$$\lim_{x \wedge y \rightarrow \infty} \frac{\overline{F}_t(x, y)}{\overline{F}_1(x, y)} = \frac{C \overline{F}(t_1 x) \overline{G}(t_2 y)}{C \overline{F}(x) \overline{G}(y)},$$

from where, if  $C > 0$ , we obtain  $\mathbf{F} \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , but if  $C = 0$  we get indefiniteness, and therefore is not necessarily valid that the joint distribution of  $(X, Y)$  belongs to  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ . The counterexample of SAI with  $C = 0$ , shows that the marginal distribution alone can not imply the membership in  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ . Even more, this counterexample still holds for the classes  $\mathcal{L}^{(2)}, \mathcal{D}^{(2)}, \mathcal{C}^{(2)}$ .

Next, we present some relations with respect to the characterization of closure property in a two-dimensional class, as also of joint max-sum equivalence, as can be found in [38].

- (1) *Closure property with respect to sum.* Let  $X_1, X_2, Y_1, Y_2$  random variables with distributions  $F_1, F_2, G_1, G_2$  respectively. If the following conditions hold  $F_1, F_2, G_1, G_2 \in \mathcal{B}$  and for any  $k, l \in \{1, 2\}$  holds  $(F_k, G_l) \in \mathcal{B}^{(2)}$ , and  $(F_{X_1+X_2}, G_{Y_1+Y_2}) \in \mathcal{B}^{(2)}$ , then we say that  $\mathcal{B}^{(2)}$  is closed with respect to the sum, where  $\mathcal{B}^{(2)}$  represents some two-dimensional version of some class  $\mathcal{B}$ .
- (2) *Closure property with respect to convolution product.* If  $X, Y$  random variables with distributions  $F, G$  and  $(F, G) \in \mathcal{B}^{(2)}$  and  $(\Theta, \Delta)$  random pair, then we say that the two-dimensional class  $\mathcal{B}^{(2)}$  is closed with respect to convolution product if  $(\Theta X, \Delta Y) \in \mathcal{B}^{(2)}$ .
- (3) *Joint max-sum equivalence.* Let  $X_1, X_2, Y_1, Y_2$  be random variables, then we say that they are joint max-sum equivalent if it holds  $\mathbf{P}[X_1 + X_2 > x, Y_1 + Y_2 > x] \sim \sum_{k=1}^2 \sum_{l=1}^2 \mathbf{P}[X_k > x, Y_l > y]$ , as  $x \wedge y \rightarrow \infty$ .
- (4) *Closure properties with respect to finite mixture.* Let  $X_1, X_2, Y_1, Y_2$  random variables, with distributions  $F_1, F_2, G_1, G_2$  respectively. If  $F_1, F_2, G_1, G_2 \in \mathcal{B}$ , and from the  $\mathbf{F} = (F_1, G_1) \in \mathcal{B}^{(2)}$  and  $\mathbf{G} = (F_2, G_2) \in \mathcal{B}^{(2)}$  follows the inclusion

$$p \overline{\mathbf{F}} + (1 - p) \overline{\mathbf{G}} \in \mathcal{B}^{(2)}, \quad (2.7)$$

for any  $p \in (0, 1)$ , then we say that class  $\mathcal{B}^{(2)}$  is closed with respect to finite mixture.

**Remark 2.3.** Relation (2.7) means that the joint distribution has the properties of  $\mathcal{B}^{(2)}$ , but also that

$$p \overline{F}_1 + (1 - p) \overline{F}_2 \in \mathcal{B}, \quad p \overline{G}_1 + (1 - p) \overline{G}_2 \in \mathcal{B}, \quad (2.8)$$

for any  $p \in (0, 1)$ . The closure property with respect to finite mixture comes here compatibly with the closure with respect to sum. The reason is that in case of finite mixture in univariate case in some classes, these two properties are closely related, see for example [33] or [40], for the classes of convolution equivalence and strong subexponentiality respectively.

**Remark 2.4.** In one dimensional set up the closure properties are well studied with respect to convolution, convolution product, finite mixture or minima and maxima. For some papers

on this topic see [15], [16], [40], [58]. Furthermore a detailed account of closure properties for heavy tailed distributions we refer to [41]. In many dimensions there are only a few attempts, mostly with respect to convolution product of MRV, see [5] and [25]. An introduction of random vectors with heavy tails in relation with closure properties can be found in [37] and [17].

**2.3. Dependence Modeling.** Now, we introduce the dependence structures, which are useful in our presentation. We suggest two dependencies, that apply on only single sequence of random variables and next we provide another two structures, which apply on double sequences of random variables.

For a sequence of real, random variables  $X_1, \dots, X_n$  with distributions  $F_1, \dots, F_n$  respectively, we say that they are pairwise quasi-asymptotically independent, symbolically  $pQAI$ , if for any pair  $i, j = 1, \dots, n$ , with  $i \neq j$  holds the limit

$$\lim_{x \rightarrow \infty} \frac{\mathbf{P}[|X_i| > x, X_j > x]}{\overline{F}_i(x) + \overline{F}_j(x)} = 0, \quad (2.9)$$

while, for the same sequence of random variables, we say that they are Tail Asymptotic Independent, symbolically  $TAI$ , and in some works named as strong quasi-asymptotically independent, if for any pair  $i, j = 1, \dots, n$ , with  $i \neq j$  holds the limit

$$\lim_{x_i \wedge x_j \rightarrow \infty} \mathbf{P}[|X_i| > x_i \mid X_j > x_j] = 0. \quad (2.10)$$

From relations (2.9) and (2.10) we see easily the inclusion  $TAI \subset pQAI$ . The dependencies  $pQAI$  and  $TAI$  were introduced in [10] and [26] respectively, where the asymptotic formula (1.1) was proved for these two dependencies in the distribution classes  $\mathcal{C}$  and  $\mathcal{D} \cap \mathcal{L}$  respectively. These dependence structures are included in the concept of asymptotic independence, see for example [49], and used for randomly weighted sums, or for generalized moments of randomly weighted sums, see [19], [43], [63].

The following proposition shows that class  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$  is restricted in the quasi-asymptotic independence structure, for the random pair  $(X, Y)$ .

**Proposition 2.1.** *If  $(X, Y) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , then  $X$  and  $Y$  are quasi-asymptotically independent.*

**Proof.** We use the method of contradiction. Let assume that  $X$  and  $Y$  are not quasi-asymptotically independent. The, at least one of the following two relations does not hold

$$\lim_{x \rightarrow \infty} \frac{\mathbf{P}[X > x, |Y| > x]}{\overline{F}(x) + \overline{G}(x)} = 0, \quad \lim_{x \rightarrow \infty} \frac{\mathbf{P}[|X| > x, Y > x]}{\overline{F}(x) + \overline{G}(x)} = 0. \quad (2.11)$$

Without loss of generality, let  $\alpha_1 \leq \alpha_2$  and let assume that the first relation in (2.11) does not hold. Note also that

$$\lim_{x \rightarrow \infty} \frac{\mathbf{P}[X > x, |Y| > x]}{\overline{F}(x) + \overline{G}(x)} \leq 1.$$

Whence, we find

$$\limsup_{x \rightarrow \infty} \frac{\mathbf{P}[X > x, |Y| > x]}{\overline{F}(x) + \overline{G}(x)} =: c \in (0, 1].$$

Thus, there exists a sequence  $\{x_n, n \in \mathbb{N}\}$ , such that the convergence  $\lim_{n \rightarrow \infty} x_n = \infty$  and

$$\lim_{n \rightarrow \infty} \frac{\mathbf{P}[X > x_n, |Y| > x_n]}{\overline{F}(x_n) + \overline{G}(x_n)} = c,$$

hold. We choose now  $t$  small enough, such that  $1/c < t^{-\alpha_1}$ , which implies  $t \in (0, 1)$ . Then, we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\mathbf{P}[X > t x_n, Y > t x_n]}{\mathbf{P}[X > x_n, Y > x_n]} &\leq \lim_{n \rightarrow \infty} \frac{\mathbf{P}[X > t x_n, |Y| > t x_n]}{\mathbf{P}[X > x_n, Y > x_n]} \\ &= \lim_{n \rightarrow \infty} \frac{\mathbf{P}[X > t x_n, |Y| > t x_n]}{\overline{F}(t x_n) + \overline{G}(t x_n)} \frac{\overline{F}(t x_n) + \overline{G}(t x_n)}{\overline{F}(x_n) + \overline{G}(x_n)} \frac{\overline{F}(x_n) + \overline{G}(x_n)}{\mathbf{P}[X > x_n, Y > x_n]} \\ &\leq \lim_{n \rightarrow \infty} \max \left\{ \frac{\overline{F}(t x_n)}{\overline{F}(x_n)}, \frac{\overline{G}(t x_n)}{\overline{G}(x_n)} \right\} \frac{\overline{F}(x_n) + \overline{G}(x_n)}{\mathbf{P}[X > x_n, Y > x_n]} = \frac{t^{-\alpha_2}}{c} < t^{-\alpha_1} t^{-\alpha_2}. \end{aligned}$$

This implies that

$$\lim_{n \rightarrow \infty} \frac{\mathbf{P}[X > t x, Y > t x]}{\mathbf{P}[X > x, Y > x]} \neq t^{-\alpha_1} t^{-\alpha_2},$$

which contradicts with  $(X, Y) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ .  $\square$

**Remark 2.5.** Another observation to class  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$  is the following. Let us write

$$f(x, y) := \frac{\mathbf{P}[X > x, Y > y]}{\mathbf{P}[X > x] \mathbf{P}[Y > y]},$$

then, if  $(X, Y) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , with  $\alpha_1, \alpha_2 > 0$ , the function  $f(x, y)$  has some kind of 'bivariate slow variation', in the sense that for any  $t_1, t_2 > 0$  it holds

$$\lim_{x \wedge y \rightarrow \infty} \frac{f(t_1 x, t_2 y)}{f(x, y)} = \lim_{x \wedge y \rightarrow \infty} \frac{\frac{\mathbf{P}[X > t_1 x, Y > t_2 y]}{\mathbf{P}[X > t_1 x] \mathbf{P}[Y > t_2 y]}}{\frac{\mathbf{P}[X > x, Y > y]}{\mathbf{P}[X > x] \mathbf{P}[Y > y]}} = 1,$$

hence we find

$$\frac{\mathbf{P}[X > t_1 x, Y > t_2 y]}{\mathbf{P}[X > x] \mathbf{P}[Y > y]} = \frac{f(t_1 x, t_2 y) \overline{F}(t_1 x) \overline{G}(t_2 y)}{f(x, y) \overline{F}(x) \overline{G}(y)} \rightarrow t^{-\alpha_1} t^{-\alpha_2},$$

as  $x \wedge y \rightarrow \infty$ . By this is implied a generalization of the SAI dependence structure. Indeed, we get  $\mathbf{P}[X > x, Y > y] = f(x, y) \overline{F}(x) \overline{G}(y)$ , to become SAI, when the function  $f(x, y)$  takes a constant value, and the last relation is not necessarily an asymptotic one.

In the dependence structures with two sequences of random variables, we want to model simultaneously the dependence among the terms of each sequences and the dependence between the two sequences. Let two sequences of real, random variables  $X_1, \dots, X_n$  and  $Y_1, \dots, Y_m$ . We say that they are Generalized Tail Asymptotic Independent, symbolically GTAI, if

$$\lim_{x_i \wedge x_k \wedge y_j \rightarrow \infty} \mathbf{P}[|X_i| > x_i \mid X_k > x_k, Y_j > y_j] = 0,$$

for any  $i, k = 1, \dots, n, j = 1, \dots, m$ , with  $i \neq k$ , and if

$$\lim_{x_i \wedge y_k \wedge y_j \rightarrow \infty} \mathbf{P}[|Y_j| > y_j \mid Y_k > y_k, X_i > x_i] = 0,$$

for any  $j, k = 1, \dots, m, i = 1, \dots, n$ , with  $i \neq k$ .

This kind of dependence structure indicates that the probability of happening three extreme events is negligible in comparison with the probability of happening two extreme events, one in each sequence. Furthermore, we see that if the two sequences  $X_1, \dots, X_n$  and  $Y_1, \dots, Y_m$  are independent, then each one of them has *TAI* dependence structure. The *GTAI* dependence structure is found in [36].

Next, we introduce a new dependence structure between the two sequences, that generalizes the *GTAI*, but remains in the same spirit.

**Definition 2.2.** *Let two sequences of real, random variables  $X_1, \dots, X_n$  and  $Y_1, \dots, Y_m$ . We say that they are Generalized Quasi Asymptotically Independent, symbolically *GQAI*, if*

$$\lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[|X_i| > x, X_k > x, Y_j > y]}{\mathbf{P}[X_i > x, Y_j > y] + \mathbf{P}[X_k > x, Y_j > y]} = 0,$$

for any  $i, k = 1, \dots, n, j = 1, \dots, m$  with  $i \neq k$ , and if

$$\lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[|Y_j| > y, Y_k > y, X_i > x]}{\mathbf{P}[X_i > x, Y_j > y] + \mathbf{P}[X_i > x, Y_k > y]} = 0,$$

for any  $j, k = 1, \dots, m, i = 1, \dots, n$  with  $j \neq k$ .

**Remark 2.6.** *We can observe that  $GTAI \subsetneq GQAI$ , and if the two sequences are independent, then for any  $i, k = 1, \dots, n, j = 1, \dots, m$  with  $k \neq i$  we obtain*

$$\begin{aligned} 0 &= \lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[|X_i| > x, X_k > x, Y_j > y]}{\mathbf{P}[X_i > x, Y_j > y] + \mathbf{P}[X_k > x, Y_j > y]} \\ &= \lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[|X_i| > x, X_k > k] \mathbf{P}[Y_j > y]}{(\mathbf{P}[X_i > x] + \mathbf{P}[X_k > x]) \mathbf{P}[Y_j > y]} \\ &= \lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[|X_i| > x, X_k > k]}{\overline{F}_i(x) + \overline{F}_k(x)} = 0, \end{aligned}$$

hence, the  $X_1, \dots, X_n$  have the *pQAI* dependence structure and similarly the  $Y_1, \dots, Y_m$  have also the *pQAI* dependence structure.

Further, we see that the dependencies *GTAI* and *GQAI* contain the complete independence case, namely when the two sequences are independent and both sequences have independent terms.

The main target in this paper is the estimation of the asymptotic expressions (1.6) and (1.7), in first case of *GQAI* dependence in distribution class  $\mathcal{C}^{(2)}$ , and in second case of *GTAI* dependence in distribution class  $(\mathcal{D} \cap \mathcal{L})^{(2)}$ , for not necessarily common multitude of terms, namely for  $n \neq m$ .

### 3. JOINT TAIL BEHAVIOR OF RANDOM SUMS

In this section we show that the insensitivity property of the joint distribution tail of the random sums, with respect to *GQAI* dependence in the frame of distribution class  $\mathcal{C}^{(2)}$ .

**Theorem 3.1.** *Let two sequences of real, random variables  $X_1, \dots, X_n$  with distributions  $F_1, \dots, F_n \in \mathcal{C}$  and  $Y_1, \dots, Y_m$  with distributions  $G_1, \dots, G_m \in \mathcal{C}$  respectively. If both sequences  $X_1, \dots, X_n, Y_1, \dots, Y_m \in GQAI$ , and  $(X_i, Y_j) \in \mathcal{C}^{(2)}$ , for any  $i = 1, \dots, n$  and  $j = 1, \dots, m$ , then*

$$\mathbf{P}[S_n > x, T_m > y] \sim \mathbf{P}\left[\bigvee_{i=1}^n S_i > x, \bigvee_{j=1}^m T_j > y\right] \sim \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[X_i > x, Y_j > y] =: S(x, y), \quad (3.1)$$

as  $x \wedge y \rightarrow \infty$ .

**Proof.** We start with the upper inequality

$$\mathbf{P}[S_n > x, T_m > y] \leq [1 + o(1)]S(x, y), \quad (3.2)$$

as  $x \wedge y \rightarrow \infty$ . Let choose arbitrarily a constant  $\varepsilon \in (0, 1)$ , then we obtain

$$\begin{aligned} \mathbf{P}[S_n > x, T_m > y] &\leq \mathbf{P}\left[\bigcup_{i=1}^n \{X_i > (1 - \varepsilon)x\}, T_m > y\right] \\ &+ \mathbf{P}\left[S_n > x, T_m > y, \bigcap_{i=1}^n \{X_i \leq (1 - \varepsilon)x\}\right] =: I_1(x, y) + I_2(x, y). \end{aligned} \quad (3.3)$$

For the first term we find

$$\begin{aligned} I_1(x, y) &\leq \mathbf{P}\left[\bigcup_{i=1}^n \{X_i > (1 - \varepsilon)x\}, \bigcup_{j=1}^m \{Y_j > (1 - \varepsilon)y\}\right] \\ &+ \mathbf{P}\left[\bigcup_{i=1}^n \{X_i > (1 - \varepsilon)x\}, T_m > y, \bigcap_{j=1}^m \{Y_j \leq (1 - \varepsilon)y\}\right] =: I_{11}(x, y) + I_{12}(x, y). \end{aligned} \quad (3.4)$$

Next, we estimate the  $I_{11}(x, y)$

$$I_{11}(x, y) \leq \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[X_i > (1 - \varepsilon)x, Y_j > (1 - \varepsilon)y]. \quad (3.5)$$

For the term  $I_{12}(x, y)$  we obtain

$$\begin{aligned} I_{12}(x, y) &= \mathbf{P}\left[\bigcup_{i=1}^n \{X_i > (1 - \varepsilon)x\}, T_m > y, \bigcup_{l=1}^m \left\{Y_l > \frac{y}{m}\right\}, \bigcap_{j=1}^m \{Y_j \leq (1 - \varepsilon)y\}\right] \\ &\leq \sum_{i=1}^n \sum_{l=1}^m \mathbf{P}\left[X_i > (1 - \varepsilon)x, Y_l > \frac{y}{m}, T_m - Y_l > \varepsilon y\right] \leq \sum_{i=1}^n \sum_{l \neq j=1}^m \sum \end{aligned} \quad (3.6)$$

$$\begin{aligned} &\mathbf{P}\left[X_i > (1 - \varepsilon)x, Y_l > \frac{y}{m} \wedge \frac{\varepsilon y}{m-1}, Y_j > \frac{y}{m} \wedge \frac{\varepsilon y}{m-1}\right] = o\left(\sum_{i=1}^n \sum_{l \neq j=1}^m \sum \mathbf{P}\left[X_i > \right. \right. \\ &\left. \left. (1 - \varepsilon)x, Y_j > \frac{y}{m} \wedge \frac{\varepsilon y}{m-1}\right] + \mathbf{P}\left[X_i > (1 - \varepsilon)x, Y_l > \frac{y}{m} \wedge \frac{\varepsilon y}{m-1}\right]\right) = o[S(x, y)], \end{aligned}$$

as  $x \wedge y \rightarrow \infty$ , where in the pre-last step we take into account that  $X_1, \dots, X_n, Y_1, \dots, Y_m \in GQAI$  and in the last step used that

$$(X_i, Y_j) \in \mathcal{C}^{(2)} \subsetneq \mathcal{D}^{(2)}, \quad (3.7)$$

. Therefore from (3.5) and (3.6) taking into account also relation (3.4), we finally obtain the following upper asymptotic bound of the  $I_1(x, y)$

$$I_1(x, y) \leq [1 + o(1)] \sum_{i=1}^n \sum_{j=1}^m \mathbf{P} [X_i > x, Y_j > y], \quad (3.8)$$

as  $x \wedge y \rightarrow \infty$ .

Now we consider the estimation of term  $I_2(x, y)$

$$\begin{aligned} I_2(x, y) &\leq \mathbf{P} \left[ S_n > x, \bigcap_{i=1}^n \{X_i \leq (1 - \varepsilon)x\}, \bigcup_{j=1}^m \{Y_j > (1 - \varepsilon)y\} \right] \\ &\quad + \mathbf{P} \left[ S_n > x, T_m > y, \bigcap_{i=1}^n \{X_i \leq (1 - \varepsilon)x\}, \bigcap_{j=1}^m \{Y_j \leq (1 - \varepsilon)y\} \right] \\ &=: I_{21}(x, y) + I_{22}(x, y). \end{aligned} \quad (3.9)$$

But for the first term  $I_{21}(x, y)$  holds

$$\begin{aligned} I_{21}(x, y) &= \mathbf{P} \left[ S_n > x, \bigcap_{i=1}^n \{X_i \leq (1 - \varepsilon)x\}, \bigcup_{j=1}^m \{Y_j > (1 - \varepsilon)y\}, \bigcup_{k=1}^n \left\{ X_k > \frac{x}{n} \right\} \right] \\ &\leq \sum_{i=1}^n \sum_{j=1}^m \mathbf{P} \left[ X_k > \frac{x}{n}, S_n - X_k > \varepsilon x, Y_j > (1 - \varepsilon)y \right] \\ &\leq \sum_{i \neq k=1}^n \sum_{j=1}^m \mathbf{P} \left[ X_i > \frac{x}{n} \wedge \frac{\varepsilon x}{n-1}, X_k > \frac{x}{n} \wedge \frac{\varepsilon x}{n-1}, Y_j > (1 - \varepsilon)y \right] \\ &= o \left[ \sum_{i \neq k=1}^n \sum_{j=1}^m \left( \mathbf{P} \left[ X_i > \frac{x}{n} \wedge \frac{\varepsilon x}{n-1}, Y_j > (1 - \varepsilon)y \right] + \mathbf{P} \left[ X_k > \frac{x}{n} \wedge \frac{\varepsilon x}{n-1}, Y_j > (1 - \varepsilon)y \right] \right) \right], \end{aligned}$$

as  $x \wedge y \rightarrow \infty$ . Hence,

$$I_{21}(x, y) = o(1) \sum_{i=1}^n \sum_{j=1}^m \mathbf{P} [X_i > x, Y_j > y], \quad (3.10)$$

as  $x \wedge y \rightarrow \infty$ , where we used again (3.7) and that  $X_1, \dots, X_n, Y_1, \dots, Y_m \in GQAI$ .

For the  $I_{22}(x, y)$  we find

$$\begin{aligned}
I_{22}(x, y) &= \mathbf{P} \left[ S_n > x, T_m > y, \bigcap_{i=1}^n \{X_i \leq (1 - \varepsilon)x\}, \right. \\
&\quad \left. \bigcap_{j=1}^m \{Y_j > (1 - \varepsilon)y\}, \bigcup_{k=1}^n \left\{X_k > \frac{x}{n}\right\}, \bigcup_{l=1}^m \left\{Y_l > \frac{y}{m}\right\} \right] \\
&\leq \sum_{k=1}^n \sum_{l=1}^m \mathbf{P} \left[ X_k > \frac{x}{n}, S_n - X_k > \varepsilon x, Y_l > \frac{y}{m}, T_m - Y_l > \varepsilon y \right] \leq \sum_{i \neq k=1}^n \sum \\
&\quad \sum_{j \neq l=1}^m \mathbf{P} \left[ X_i > \frac{x}{n} \wedge \frac{\varepsilon x}{n-1}, X_k > \frac{x}{n} \wedge \frac{\varepsilon x}{n-1}, Y_j > \frac{y}{m} \wedge \frac{\varepsilon y}{m-1}, Y_l > \frac{y}{m} \wedge \frac{\varepsilon y}{m-1} \right] \\
&\leq \sum_{i=1}^n \sum_{j \neq l=1}^m \sum \mathbf{P} \left[ X_i > \frac{x}{n} \wedge \frac{\varepsilon x}{n-1}, Y_j > \frac{y}{m} \wedge \frac{\varepsilon y}{m-1}, Y_l > \frac{y}{m} \wedge \frac{\varepsilon y}{m-1} \right] \\
&= o(1) \sum_{i=1}^n \sum_{j \neq l=1}^m \sum \left( \mathbf{P} \left[ X_i > \frac{x}{n} \wedge \frac{\varepsilon x}{n-1}, Y_j > \frac{y}{m} \wedge \frac{\varepsilon y}{m-1} \right] \right. \\
&\quad \left. + \mathbf{P} \left[ X_i > \frac{x}{n} \wedge \frac{\varepsilon x}{n-1}, Y_l > \frac{y}{m} \wedge \frac{\varepsilon y}{m-1} \right] \right),
\end{aligned}$$

as  $x \wedge y \rightarrow \infty$ , whence from inclusion  $\mathcal{C}^{(2)} \subsetneq \mathcal{D}^{(2)}$  we obtain

$$I_{22}(x, y) = o(1) \sum_{i=1}^n \sum_{j=1}^m \mathbf{P} [X_i > x, Y_j > y], \quad (3.11)$$

as  $x \wedge y \rightarrow \infty$ . Therefore from relations (3.10) and (3.11) in combination with (3.9) we conclude

$$I_2(x, y) = o(1) \sum_{i=1}^n \sum_{j=1}^m \mathbf{P} [X_i > x, Y_j > y], \quad (3.12)$$

as  $x \wedge y \rightarrow \infty$ , and by (3.8) and (3.12) in combination with (3.3) we get (3.2).

Now we estimate the lower bound in the form

$$\mathbf{P}[S_n > x, T_m > y] \geq [1 - o(1)] \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[X_i > x, Y_j > y], \quad (3.13)$$

as  $x \wedge y \rightarrow \infty$ . Let remind that  $\mathcal{C}^{(2)} \subsetneq (\mathcal{D} \cap \mathcal{L})^{(2)}$ , that means for any sequences  $X_1, \dots, X_n$  and  $Y_1, \dots, Y_m$ , there exists some joint insensitivity function  $\mathbf{a}$ , such that

$$\mathbf{a} = (a_F, a_G) := \left( \bigwedge_{i=1}^n a_{F_i}, \bigwedge_{j=1}^m a_{G_j} \right), \quad (3.14)$$

therefore, this function represents an insensitivity function, for any distribution pair  $(F_i, G_j)$ , for any  $i = 1, \dots, n$  and  $j = 1, \dots, m$ .

For such a function  $\mathbf{a}$  we obtain

$$\mathbf{P}[S_n > x, T_m > y] \geq \mathbf{P} \left[ S_n > x, T_m > y, \bigvee_{i=1}^n X_i > x + a_F(x), \bigvee_{j=1}^m Y_j > y + a_G(y) \right], \quad (3.15)$$

Applying Bonferroni inequality twice in (3.15), we find

$$\begin{aligned} \mathbf{P}[S_n > x, T_m > y] &\geq \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[S_n > x, T_m > y, X_i > x + a_F(x), Y_j > y + a_G(y)] \\ &- \sum_{i < k=1}^n \sum_{j=1}^m \mathbf{P}[X_i > x + a_F(x), X_k > x + a_F(x), Y_j > y + a_G(y)] \\ &- \sum_{i=1}^n \sum_{j < l=1}^m \mathbf{P}[X_i > x + a_F(x), Y_j > y + a_G(y), Y_l > y + a_G(y)] =: \sum_{k=1}^3 P_k(x, y). \end{aligned} \quad (3.16)$$

By the assumption that  $X_1, \dots, X_n$  and  $Y_1, \dots, Y_m$  are *GQAI* and the insensitivity functions properties we conclude that

$$P_k(x, y) = o \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[X_i > x, Y_j > y] \right), \quad (3.17)$$

as  $x \wedge y \rightarrow \infty$  for  $k = 1, 2$ . For  $P_1(x, y)$  we write

$$\begin{aligned} P_1(x, y) &\geq \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[X_i > x + a_F(x), Y_j > y + a_G(y)] \\ &- \sum_{i \neq k=1}^n \sum_{j=1}^m \mathbf{P} \left[ X_i > x + a_F(x), X_k < -\frac{a_F(x)}{n}, Y_j > y + a_G(y) \right] \\ &- \sum_{i=1}^n \sum_{j \neq l=1}^m \mathbf{P} \left[ X_i > x + a_F(x), Y_j > y + a_G(y), Y_l < -\frac{a_G(y)}{m} \right] \\ &=: P_{11}(x, y) - P_{12}(x, y) - P_{13}(x, y). \end{aligned} \quad (3.18)$$

Now we estimate the  $P_{12}(x, y)$

$$\begin{aligned} P_{12}(x, y) &\leq \sum_{i \neq k=1}^n \sum_{j=1}^m \mathbf{P} \left[ X_i > \frac{a_F(x)}{n}, Y_j > y, |X_k| > \frac{a_G(y)}{n} \right] \\ &= o(1) \sum_{i \neq k=1}^n \sum_{j=1}^m \left( \mathbf{P} \left[ X_i > \frac{a_F(x)}{n}, Y_j > y \right] + \mathbf{P} \left[ X_k > \frac{a_F(x)}{n}, Y_j > y \right] \right) \\ &= o(1) \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[X_i > x, Y_j > y], \end{aligned}$$



as  $x \wedge y \rightarrow \infty$ , where we use the GQAI property and in last step we take into account the inclusion  $\mathcal{C}^{(2)} \subsetneq \mathcal{D}^{(2)}$ . By symmetry we find similarly

$$P_{13}(x, y) = o(1) \sum_{i=1}^n \sum_{j=1}^m \mathbf{P} [X_i > x, Y_j > y], \quad (3.19)$$

as  $x \wedge y \rightarrow \infty$ . Hence, from relations (3.18) - (3.19) and the fact that  $\mathcal{C}^{(2)} \subsetneq (\mathcal{D} \cap \mathcal{L})^{(2)}$  in combination with the properties of function  $\mathbf{a}$  we obtain

$$P_1(x, y) \geq [1 - o(1)] \sum_{i=1}^n \sum_{j=1}^m \mathbf{P} [X_i > x, Y_j > y], \quad (3.20)$$

as  $x \wedge y \rightarrow \infty$ . So, by relation (3.16) in combination with relations (3.17) and (3.20) we conclude that (3.13) is true. Therefore

$$\mathbf{P}[S_n > x, T_m > y] \sim \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[X_i > x, Y_j > y], \quad (3.21)$$

as  $x \wedge y \rightarrow \infty$ . Now it remains to use (3.21) and the inequality

$$\mathbf{P}[S_n > x, T_m > y] \leq \mathbf{P} \left[ \bigvee_{i=1}^n S_i > x, \bigvee_{j=1}^m T_j > y \right] \leq \mathbf{P} [S_n^+ > x, T_m^+ > y],$$

where  $S_n^+ = \sum_{i=1}^n X_i^+$ ,  $T_m^+ = \sum_{j=1}^m Y_j^+$ , to establish (3.1).  $\square$

**Remark 3.1.** Comparing Theorem 3.1 with relation (1.8) we find the following differences. In relation (1.8) the dependence structure is restricted as  $GTAI \subset GQAI$  (and in our Theorem we do not need the assumption of TAI in each sequence), while the distribution class is wider since  $\mathcal{C}^{(2)} \subsetneq (\mathcal{D} \cap \mathcal{L})^{(2)}$ . Furthermore relation (1.8) contains also the asymptotic behavior of the maxima, which is missing from Theorem 3.1. Another difference is that in Theorem 3.1 is not required the same multitude of summands, although in relation (1.8) is not permitted, which can be helpful in risk theory, since it allows two different counting processes.

The two-dimensional risk models become more and more popular, because of the need of insurance companies to operate several parallel business lines, see for example in [29], [47], [70].

Next, we have a direct corollary for a two-dimensional risk model, with discrete time. The surplus processes are of the form  $U_1(k, x) := x - \sum_{i=1}^k X_i$ ,  $U_2(k, y) := y - \sum_{j=1}^k Y_j$ , for any  $k = 1, \dots, n$ , where  $x, y$  are the initial surpluses in each of the two business lines, while  $X_i$  and  $Y_j$  are the claims of each business line during the  $i$ -th period. In contrast to one-dimensional case, there exist several ways to define the ruin time, or the ruin probability, see for example [11]. Let choose the following definition of the ruin probability

$$\begin{aligned} \tau_1(x) &:= \inf\{k = 1, \dots, n : U_1(k, x) < 0 \mid U_1(0) = x\}, \\ \tau_2(y) &:= \inf\{k = 1, \dots, n : U_2(k, y) < 0 \mid U_2(0) = y\}, \end{aligned}$$

and  $\tau_{and} := \max\{\tau_1(x), \tau_2(y)\}$ , that indicates the first moment, when both surpluses fall below zero, but not necessarily simultaneously. Hence we obtain

$$\psi_{and}(x, y, n) := \mathbf{P}[\tau_{and} \leq n] = \mathbf{P} \left[ \bigvee_{i=1}^n S_i > x, \bigvee_{j=1}^n T_j > y \right].$$

Next we obtain an asymptotic expression of the ruin probability in finite time for  $n = m$ .

**Corollary 3.1.** *Under the conditions of Theorem 3.1, with  $n = m$ , then*

$$\psi_{and}(x, y, n) \sim \sum_{i=1}^n \sum_{j=1}^n \mathbf{P}[X_i > x, Y_j > y],$$

as  $x \wedge y \rightarrow \infty$ .

#### 4. TWO-DIMENSIONAL CLOSURE PROPERTIES

Now we proceed to the closure properties. Next lemma studies the closure properties of sum with respect to classes of regularly and dominatedly varying distributions under the dependence structure  $QAI$ . For the regular variation class the corresponding closure property under independence can be found in [22, p. 278], while for arbitrarily dependent random variables, under more strict conditions can be found in [55, Lem. 4.4.2] and [42, Lem. 3.1]. For the class of consistently varying distributions the closure property in the independence case was given by [8] for non-negative variables and by [30] for real valued variables. Further, under more strict conditions in [69] we find closure property of convolution under arbitrary dependence in class  $\mathcal{C}$ .

**Lemma 4.1.** (1) *Let  $X_1, X_2$  real random variables, with distributions  $F_1, F_2 \in \mathcal{C}$  respectively, under  $QAI$  dependence. Then*

$$\overline{F}_{X_1+X_2}(x) \sim \overline{F}_1(x) + \overline{F}_2(x), \quad (4.1)$$

as  $x \rightarrow \infty$ , and  $F_{X_1+X_2} \in \mathcal{C}$ .

(2) *Under the conditions of part (1), with the only difference that  $F_1, F_2 \in \mathcal{R}_{-\alpha}$ , for some  $\alpha > 0$ , then  $F_{X_1+X_2} \in \mathcal{R}_{-\alpha}$ .*

**Proof.**

(1) In the case of  $QAI$  relation (4.1) follows from [10, Th. 3.1], for  $n = 2$ . Hence, for the closure property of  $\mathcal{C}$  with respect to convolution we obtain for all the distributions

$$\lim_{z \uparrow 1} \limsup_{x \rightarrow \infty} \frac{\overline{F}_{X_1+X_2}(zx)}{\overline{F}_{X_1+X_2}(x)} \geq 1, \quad (4.2)$$

from the elementary inequalities

$$\min \left\{ \frac{a}{c}, \frac{b}{d} \right\} \leq \frac{a+b}{c+d} \leq \max \left\{ \frac{a}{c}, \frac{b}{d} \right\}, \quad (4.3)$$

for any constants  $a, b, c, d > 0$  and from (4.1) we find

$$\begin{aligned} \lim_{z \uparrow 1} \limsup_{x \rightarrow \infty} \frac{\overline{F}_{X_1+X_2}(zx)}{\overline{F}_{X_1+X_2}(x)} &= \lim_{z \uparrow 1} \limsup_{x \rightarrow \infty} \frac{\overline{F}_1(zx) + \overline{F}_2(zx)}{\overline{F}_1(x) + \overline{F}_2(x)} \\ &\leq \lim_{z \uparrow 1} \limsup_{x \rightarrow \infty} \max \left\{ \frac{\overline{F}_1(zx)}{\overline{F}_1(x)}, \frac{\overline{F}_2(zx)}{\overline{F}_2(x)} \right\} = \max_{i \in \{1, 2\}} \left\{ \lim_{z \uparrow 1} \limsup_{x \rightarrow \infty} \frac{\overline{F}_i(zx)}{\overline{F}_i(x)} \right\} = 1, \end{aligned} \quad (4.4)$$

where in the last step we use the assumption  $F_1, F_2 \in \mathcal{C}$ . Whence by (4.2) and (4.4) we find  $F_{X_1+X_2} \in \mathcal{C}$ .

- (2) From the relation  $\mathcal{R}_{-\alpha}$  and by part (1) we obtain (4.1), hence together with (4.3) we conclude

$$\lim_{x \rightarrow \infty} \frac{\overline{F}_{X_1+X_2}(tx)}{\overline{F}_{X_1+X_2}(x)} = \lim_{x \rightarrow \infty} \frac{\overline{F}_1(tx) + \overline{F}_2(tx)}{\overline{F}_1(x) + \overline{F}_2(x)} \leq \max_{i \in \{1,2\}} \left\{ \lim_{x \rightarrow \infty} \frac{\overline{F}_i(tx)}{\overline{F}_i(x)} \right\} = t^{-\alpha}, \quad (4.5)$$

for any  $t > 0$ , since  $F_i \in \mathcal{R}_{-\alpha}$ . Furthermore

$$\lim_{x \rightarrow \infty} \frac{\overline{F}_{X_1+X_2}(tx)}{\overline{F}_{X_1+X_2}(x)} \geq \min_{i \in \{1,2\}} \left\{ \lim_{x \rightarrow \infty} \frac{\overline{F}_i(tx)}{\overline{F}_i(x)} \right\} = t^{-\alpha}, \quad (4.6)$$

for any  $t > 0$ , so by (4.5) and (4.6) we find  $F_{X_1+X_2} \in \mathcal{R}_{-\alpha}$ .  $\square$

In the next result we find closure property of the distribution classes  $\mathcal{C}^{(2)}$  and  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , with respect to convolution under  $GQAI$ , with additional restriction that the particular summands are  $pQAI$ . This way we have generalization of [38, Cor. 4.1] in case of  $GQAI$  instead of  $GTAI$  (and  $QAI$  instead of  $TAI$  for the summands) and with real random variables.

**Theorem 4.1.** (1) *Let  $X_1, X_2, Y_1, Y_2$  be four real random variables, with the distributions  $F_1, F_2, G_1, G_2 \in \mathcal{C}$  respectively, under  $GQAI$  dependence. If  $X_1, X_2$  are  $QAI$  and  $Y_1, Y_2$  are also  $QAI$  with  $(F_i, G_j) \in \mathcal{C}^{(2)}$  for any  $i, j = 1, 2$ , then we find  $(F_{X_1+X_2}, G_{Y_1+Y_2}) \in \mathcal{C}^{(2)}$ .*

- (2) *Under the conditions of part (1), with the only difference that  $F_1, F_2 \in \mathcal{R}_{-\alpha_1}$  and  $G_1, G_2 \in \mathcal{R}_{-\alpha_2}$ , for some  $\alpha_1, \alpha_2 > 0$  where  $(F_i, G_j) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , for any  $i, j = 1, 2$ , then  $(F_{X_1+X_2}, G_{Y_1+Y_2}) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ .*

**Proof.**

- (1) From Theorem 3.1 and  $\mathcal{L}^{(2)}$ , for  $n = m = 2$  we get

$$\mathbf{P}[X_1 + X_2 > x, Y_1 + Y_2 > y] \sim \sum_{i=1}^2 \sum_{j=1}^2 \mathbf{P}[X_i > x, Y_j > y], \quad (4.7)$$

as  $x \wedge y \rightarrow \infty$ . For any two-dimensional distributions is true the lower bound

$$\lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_1 + X_2 > z_1 x, Y_1 + Y_2 > z_2 y]}{\mathbf{P}[X_1 + X_2 > x, Y_1 + Y_2 > y]} \geq 1, \quad (4.8)$$

while by (4.7), applying the upper inequality in (4.3) thrice, we obtain

$$\begin{aligned} & \lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_1 + X_2 > z_1 x, Y_1 + Y_2 > z_2 y]}{\mathbf{P}[X_1 + X_2 > x, Y_1 + Y_2 > y]} \\ &= \lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{\sum_{i=1}^2 \sum_{j=1}^2 \mathbf{P}[X_i > z_1 x, Y_j > z_2 y]}{\sum_{i=1}^2 \sum_{j=1}^2 \mathbf{P}[X_i > x, Y_j > y]} \\ &\leq \max_{i, j \in \{1,2\}} \left\{ \lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_i > z_1 x, Y_j > z_2 y]}{\mathbf{P}[X_i > x, Y_j > y]} \right\} = 1, \end{aligned} \quad (4.9)$$

where in the last step we used that  $(F_i, G_j) \in \mathcal{C}^{(2)}$ , for any  $i, j = 1, 2$ . Thence, by (4.8) and (4.9) we find (2.4). Furthermore, by Lemma 4.1 we obtain  $F_{X_1+X_2} \in \mathcal{C}$  and  $G_{Y_1+Y_2} \in \mathcal{C}$ , which together with (2.4) gives  $(F_{X_1+X_2}, G_{Y_1+Y_2}) \in \mathcal{C}^{(2)}$ .

- (2) From  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)} \subsetneq \mathcal{C}^{(2)}$ , using Theorem 3.1, with  $n = m = 2$ , we reach to (4.7), and from this, through the application of upper inequality in (4.3) thrice, we finally get

$$\begin{aligned} \lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_1 + X_2 > t_1 x, Y_1 + Y_2 > t_2 y]}{\mathbf{P}[X_1 + X_2 > x, Y_1 + Y_2 > y]} &= \lim_{x \wedge y \rightarrow \infty} \frac{\sum_{i=1}^2 \sum_{j=1}^2 \mathbf{P}[X_i > t_1 x, Y_j > t_2 y]}{\sum_{i=1}^2 \sum_{j=1}^2 \mathbf{P}[X_i > x, Y_j > y]} \\ &\leq \max_{i, j \in \{1, 2\}} \left\{ \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_i > t_1 x, Y_j > t_2 y]}{\mathbf{P}[X_i > x, Y_j > y]} \right\} = t_1^{-\alpha_1} t_2^{-\alpha_2}, \end{aligned} \quad (4.10)$$

for any  $t_1, t_2 > 0$ , where in last step was used that

$$(F_i, G_j) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)},$$

for any  $i, j = 1, 2$ , and with similar way, by application of lower inequality in (4.3) thrice, we have

$$\begin{aligned} \lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_1 + X_2 > t_1 x, Y_1 + Y_2 > t_2 y]}{\mathbf{P}[X_1 + X_2 > x, Y_1 + Y_2 > y]} &\geq \min_{i, j \in \{1, 2\}} \left\{ \lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_i > t_1 x, Y_j > t_2 y]}{\mathbf{P}[X_i > x, Y_j > y]} \right\} \\ &= t_1^{-\alpha_1} t_2^{-\alpha_2}, \end{aligned} \quad (4.11)$$

thus by (4.10) and (4.11) we have (2.5). Next, by Lemma 4.1 we obtain the inclusions  $F_{X_1+X_2} \in \mathcal{R}_{-\alpha_1}$  and  $F_{Y_1+Y_2} \in \mathcal{R}_{-\alpha_2}$ , that together with (2.5) finally gives  $(F_{X_1+X_2}, G_{Y_1+Y_2}) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ .  $\square$

Now we study the closure properties of distribution classes with respect to product convolution in two dimensions. This work can help the extension of Theorem 3.1 to the direction of random weighted sums.

To show the closure property with respect to convolution product in distributions classes  $\mathcal{D}^{(2)}, \mathcal{C}^{(2)}, \mathcal{L}^{(2)}$  we employ the following assumption.

**Assumption 4.1.** *Let  $b : [0, \infty) \rightarrow (0, \infty)$  be a function, such that  $b(x) \rightarrow \infty$ ,  $b(x) = o(x)$ , as  $x \rightarrow \infty$  and*

$$\mathbf{P}[\Theta > b(x)] = o(\mathbf{P}[\Theta X > x, \Delta Y > y]), \quad (4.12)$$

*as  $x \rightarrow \infty$  and the  $y$  is free, and  $c : [0, \infty) \rightarrow (0, \infty)$  be a function, such that  $c(y) \rightarrow \infty$ ,  $c(y) = o(y)$ , as  $y \rightarrow \infty$  and*

$$\mathbf{P}[\Delta > c(y)] = o(\mathbf{P}[\Theta X > x, \Delta Y > y]), \quad (4.13)$$

*as  $y \rightarrow \infty$  and the  $x$  is free.*

**Remark 4.1.** *It is easy to see that if the  $\Theta$  and  $\Delta$  have distributions with bounded upper endpoint, then relations (4.12) and (4.13) are true directly. We observe that by relation (4.12) follows*

$$\mathbf{P}[\Theta > b(x)] = o(\mathbf{P}[\Theta X > x]), \quad (4.14)$$

*as  $x \rightarrow \infty$ , and by relation (4.13) follows*

$$\mathbf{P}[\Delta > c(y)] = o(\mathbf{P}[\Delta Y > y]), \quad (4.15)$$

*as  $y \rightarrow \infty$ . Therefore, from (4.14) and (4.15), in combination with the definition of functions  $b(x)$  and  $c(y)$ , by [58, Lem. 3.2] follows that if  $\Theta$  and  $\Delta$  have distributions with infinite upper*

endpoint, then

$$\mathbf{P}[\Theta > u x] = o(\mathbf{P}[\Theta X > x]), \quad \mathbf{P}[\Delta > u y] = o(\mathbf{P}[\Delta Y > y]), \quad (4.16)$$

as  $x \rightarrow \infty$  and as  $y \rightarrow \infty$  respectively, for any  $u > 0$ . In case  $\Theta$  and  $\Delta$  have distributions with finite upper endpoint relation (4.16) is true.

**Theorem 4.2.** *Let  $(X, Y)$  be a random pair with the distributions  $F, G$  respectively,  $(\Theta, \Delta)$ , be a random pair independent of  $(X, Y)$ , with non-negative, non-degenerated to zero marginal distributions, and hold Assumption 4.1. The following are true*

- (1) *If  $(X, Y) \in \mathcal{D}^{(2)}$ , then  $(\Theta X, \Delta Y) \in \mathcal{D}^{(2)}$ .*
- (2) *If  $(X, Y) \in \mathcal{L}^{(2)}$ , then  $(\Theta X, \Delta Y) \in \mathcal{L}^{(2)}$ .*
- (3) *If  $(X, Y) \in \mathcal{C}^{(2)}$ , then  $(\Theta X, \Delta Y) \in \mathcal{C}^{(2)}$ .*
- (4) *If  $(X, Y) \in (\mathcal{D} \cap \mathcal{L})^{(2)}$ , then  $(\Theta X, \Delta Y) \in (\mathcal{D} \cap \mathcal{L})^{(2)}$ .*

**Proof.**

- (1) Let  $\mathbf{b} = (b_1, b_2) \in (0, 1)^2$ , then

$$\begin{aligned} & \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[\Theta X > b_1 x, \Delta Y > b_2 y]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \quad (4.17) \\ &= \limsup_{x \wedge y \rightarrow \infty} \frac{\int_0^\infty \int_0^\infty \mathbf{P} \left[ X > \frac{b_1 x}{t}, Y > \frac{b_2 y}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \\ &= \limsup_{x \wedge y \rightarrow \infty} \frac{\left( \int_0^{b(x)} + \int_{b(x)}^\infty \right) \left( \int_0^{c(y)} + \int_{c(y)}^\infty \right) \mathbf{P} \left[ X > \frac{b_1 x}{t}, Y > \frac{b_2 y}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \\ &=: \limsup_{x \wedge y \rightarrow \infty} \frac{I_{11}(x, y) + I_{12}(x, y) + I_{21}(x, y) + I_{22}(x, y)}{\mathbf{P}[\Theta X > x, \Delta Y > y]}. \end{aligned}$$

But from Assumption 4.1 we obtain

$$\begin{aligned} I_{22}(x, y) &= \int_{b(x)}^\infty \int_{c(y)}^\infty \mathbf{P} \left[ X > \frac{b_1 x}{t}, Y > \frac{b_2 y}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds] \quad (4.18) \\ &\leq \mathbf{P}[\Theta > b(x), \Delta > c(y)] \leq \mathbf{P}[\Theta > b(x)] = o(\mathbf{P}[\Theta X > x, \Delta Y > y]), \end{aligned}$$

as  $x \rightarrow \infty$  and the  $y$  is free. For  $I_{12}(x, y)$  we find

$$I_{12}(x, y) \leq \mathbf{P}[\Theta \leq b(x), \Delta > c(y)] \leq \mathbf{P}[\Delta > c(y)] = o(\mathbf{P}[\Theta X > x, \Delta Y > y]), \quad (4.19)$$

as  $y \rightarrow \infty$  and the  $x$  is free. Through symmetry we obtain similarly

$$I_{21}(x, y) \leq \mathbf{P}[\Theta > b(x), \Delta \leq c(y)] \leq \mathbf{P}[\Theta > b(x)] = o(\mathbf{P}[\Theta X > x, \Delta Y > y]), \quad (4.20)$$

as  $x \rightarrow \infty$  and the  $y$  is free. Hence, by (4.17) together with (4.18) - (4.20) follows

$$\begin{aligned} & \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[\Theta X > b_1 x, \Delta Y > b_2 y]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \tag{4.21} \\ & \leq \limsup_{x \wedge y \rightarrow \infty} \frac{\int_0^{b(x)} \int_0^{c(y)} \mathbf{P} \left[ X > \frac{b_1 x}{t}, Y > \frac{b_2 y}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds]}{\int_0^{b(x)} \int_0^{c(y)} \mathbf{P} \left[ X > \frac{x}{t}, Y > \frac{y}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds]} \leq \limsup_{x \wedge y \rightarrow \infty} \\ & \sup_{t \in (0, b(x)), s \in (0, c(y))} \frac{\mathbf{P} \left[ X > \frac{b_1 x}{t}, Y > \frac{b_2 y}{s} \right]}{\mathbf{P} \left[ X > \frac{x}{t}, Y > \frac{y}{s} \right]} \leq \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X > b_1 x, Y > b_2 y]}{\mathbf{P}[X > x, Y > y]} < \infty, \end{aligned}$$

where in the last step used that  $(X, Y) \in \mathcal{D}^{(2)}$ . Next, since  $F \in \mathcal{D}$  and  $G \in \mathcal{D}$ , through [15, Th. 3.3(ii)] we conclude that  $\Theta X \in \mathcal{D}$  and  $\Delta Y \in \mathcal{D}$ , which in combination with (4.21) provided  $(\Theta X, \Delta Y) \in \mathcal{D}^{(2)}$ .

- (2) In case  $(X, Y) \in \mathcal{L}^{(2)}$ , for  $a_1, a_2 > 0$  is well known that for any two-dimensional distribution

$$\limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[\Theta X > x - a_1, \Delta Y > y - a_2]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \geq 1.$$

From the opposite side we find

$$\begin{aligned} & \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[\Theta X > x - a_1, \Delta Y > y - a_2]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} = \tag{4.22} \\ & \limsup_{x \wedge y \rightarrow \infty} \frac{\left( \int_0^{b(x)} + \int_{b(x)}^\infty \right) \left( \int_0^{c(y)} + \int_{c(y)}^\infty \right) \mathbf{P} \left[ X > \frac{x - a_1}{t}, Y > \frac{y - a_2}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \\ & = \limsup_{x \wedge y \rightarrow \infty} \frac{K_{11}(x, y) + K_{12}(x, y) + K_{21}(x, y) + K_{22}(x, y)}{\mathbf{P}[\Theta X > x, \Delta Y > y]}. \end{aligned}$$

Here also, similarly to (4.18) - (4.20), we find that the  $K_{12}(x, y)$ ,  $K_{21}(x, y)$ ,  $K_{22}(x, y)$  are of  $o(\mathbf{P}[\Theta X > x, \Delta Y > y])$  order of magnitude, as  $x \rightarrow \infty$  or  $y \rightarrow \infty$ , respectively. Thus, together with (4.22) we conclude

$$\begin{aligned} & \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[\Theta X > x - a_1, \Delta Y > y - a_2]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \leq \limsup_{x \wedge y \rightarrow \infty} \tag{4.23} \\ & \frac{\int_0^{b(x)} \int_0^{c(y)} \mathbf{P} \left[ X > \frac{x - a_1}{t}, Y > \frac{y - a_2}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds]}{\int_0^{b(x)} \int_0^{c(y)} \mathbf{P} \left[ X > \frac{x}{t}, Y > \frac{y}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds]} \leq \limsup_{x \wedge y \rightarrow \infty} \sup_{t \in (0, b(x)), s \in (0, c(y))} \\ & \frac{\mathbf{P} \left[ X > \frac{x - a_1}{t}, Y > \frac{y - a_2}{s} \right]}{\mathbf{P} \left[ X > \frac{x}{t}, Y > \frac{y}{s} \right]} = \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X > x - a_1, Y > y - a_2]}{\mathbf{P}[X > x, Y > y]} = 1, \end{aligned}$$

where in the last step we take into account that  $(X, Y) \in \mathcal{L}^{(2)}$ . By Assumption 4.1 and Remark 4.1, through [41, Cor. 5.1] we obtain  $\Theta X \in \mathcal{L}$ ,  $\Delta Y \in \mathcal{L}$ , which together with (4.23) provides  $(\Theta X, \Delta Y) \in \mathcal{L}^{(2)}$ .

(3) For any two-dimensional distribution holds the inequality

$$\lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[\Theta X > z_1 x, \Delta Y > z_2 y]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \geq 1, \quad (4.24)$$

from the opposite side we obtain

$$\begin{aligned} & \lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[\Theta X > z_1 x, \Delta Y > z_2 y]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \\ &= \lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{\left( \int_0^{b(x)} + \int_{b(x)}^\infty \right) \left( \int_0^{c(y)} + \int_{c(y)}^\infty \right) \mathbf{P} \left[ X > \frac{z_1 x}{t}, Y > \frac{z_2 y}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \\ &= \lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{J_{11}(x, y) + J_{12}(x, y) + J_{21}(x, y) + J_{22}(x, y)}{\mathbf{P}[\Theta X > x, \Delta Y > y]}. \end{aligned} \quad (4.25)$$

Similarly to (4.18) - (4.20), we find that the  $J_{12}(x, y)$ ,  $J_{21}(x, y)$ ,  $J_{22}(x, y)$  are of  $o(\mathbf{P}[\Theta X > x, \Delta Y > y])$  order of magnitude, as  $x \rightarrow \infty$  or  $y \rightarrow \infty$ , respectively. Therefore, together with relation (4.25) we obtain

$$\begin{aligned} & \lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[\Theta X > z_1 x, \Delta Y > z_2 y]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} \\ & \leq \lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{\int_0^{b(x)} \int_0^{c(y)} \mathbf{P} \left[ X > \frac{z_1 x}{t}, Y > \frac{z_2 y}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds]}{\int_0^{b(x)} \int_0^{c(y)} \mathbf{P} \left[ X > \frac{x}{t}, Y > \frac{y}{s} \right] \mathbf{P}[\Theta \in dt, \Delta \in ds]} \leq \lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \\ & \sup_{t \in (0, b(x)), s \in (0, c(y))} \frac{\mathbf{P} \left[ X > \frac{z_1 x}{t}, Y > \frac{z_2 y}{s} \right]}{\mathbf{P} \left[ X > \frac{x}{t}, Y > \frac{y}{s} \right]} = \lim_{z \uparrow 1} \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X > z_1 x, Y > z_2 y]}{\mathbf{P}[X > x, Y > y]} = 1, \end{aligned} \quad (4.26)$$

where in last step was used the condition  $(X, Y) \in \mathcal{C}^{(2)}$ . Now taking into consideration Assumption 4.1 and  $X \in \mathcal{C}$ ,  $Y \in \mathcal{C}$ , through Remark 4.1, because of (4.24), via [15, Th. 3.4(ii)], or [41, Prop. 5.3(ii)] we obtain  $\Theta X \in \mathcal{C}$ ,  $\Delta Y \in \mathcal{C}$ , which together with (4.26) provides  $(\Theta X, \Delta Y) \in \mathcal{C}^{(2)}$ .

(4) This case follows from parts (1) and (2).  $\square$

In part (3) of Theorem 4.2 we find a generalization of [38, Lem. 6.2], since now we have real-valued random variables  $X$  and  $Y$ . In next lemma we find an extension of Breiman's Theorem in case of distribution class  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ . For the class of regular variation in one dimension case, see [6], [18], [35], while for the class of multivariate regular variation, see [5], [25].

**Lemma 4.2.** *Let  $(X, Y)$  be a random pair with the distributions  $F, G$  respectively, such that  $(F, G) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ . Let  $(\Theta, \Delta)$  be a random pair independent of  $(X, Y)$  with non-negative, non-degenerated to zero distributions, such that  $\mathbf{E}[\Theta^{\alpha_1 + \varepsilon} \Delta^{\alpha_2 + \varepsilon}] < \infty$ , for some  $\varepsilon > 0$ . Under*

the Assumption 4.1 we have

$$\mathbf{P}[\Theta X > x, \Delta Y > y] \sim \mathbf{E}[\Theta^{\alpha_1} \Delta^{\alpha_2}] \mathbf{P}[X > x, Y > y], \quad (4.27)$$

as  $x \wedge y \rightarrow \infty$ . Furthermore  $(\Theta X, \Delta Y) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ .

**Proof.** Since for some  $\varepsilon > 0$  holds  $\mathbf{E}[\Theta^{\alpha_1+\varepsilon} \Delta^{\alpha_2+\varepsilon}] < \infty$ , from dominated convergence theorem and by  $(F, G) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$  we have

$$\begin{aligned} \mathbf{P}[\Theta X > x, \Delta Y > y] &= \int_0^\infty \int_0^\infty \mathbf{P}\left[X > \frac{x}{t}, Y > \frac{y}{s}\right] \mathbf{P}[\Theta \in dt, \Delta \in ds] \\ &= \left(\int_0^{b(x)} + \int_{b(x)}^\infty\right) \left(\int_0^{c(y)} + \int_{c(y)}^\infty\right) \mathbf{P}\left[X > \frac{x}{t}, Y > \frac{y}{s}\right] \mathbf{P}[\Theta \in dt, \Delta \in ds] \\ &=: \Lambda_{11}(x, y) + \Lambda_{1,2}(x, y) + \Lambda_{2,1}(x, y) + \Lambda_{2,2}(x, y). \end{aligned}$$

By Assumption 4.1 and similarly to (4.18) - (4.20), we find that the  $\Lambda_{12}(x, y)$ ,  $\Lambda_{21}(x, y)$ ,  $\Lambda_{22}(x, y)$  are of  $o(\mathbf{P}[\Theta X > x, \Delta Y > y])$  order of magnitude, as  $x \wedge y \rightarrow \infty$ . Hence, we by the  $\mathcal{R}^{(2)}$  property, and by definitions of functions  $b, c$ , we have

$$\begin{aligned} \mathbf{P}[\Theta X > x, \Delta Y > y] &\sim \int_0^{b(x)} \int_0^{c(y)} \mathbf{P}\left[X > \frac{x}{t}, Y > \frac{y}{s}\right] \mathbf{P}[\Theta \in dt, \Delta \in ds] \\ &\sim \int_0^{b(x)} \int_0^{c(y)} t^{-\alpha_1} s^{-\alpha_2} \mathbf{P}[X > x, Y > y] \mathbf{P}[\Theta \in dt, \Delta \in ds] \\ &= E[\Theta^{\alpha_1} \Delta^{\alpha_2} \mathbf{1}_{\{\Theta < b(x), \Delta < c(y)\}}] \mathbf{P}[X > x, Y > y], \end{aligned}$$

by the definition of functions  $b, c$ , namely tends to infinity, the indicator function tends to unity, as a result we have the relation (4.27). Further, for any  $t_1, t_2 > 0$  by (4.27) we have

$$\lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[\Theta X > t_1 x, \Delta Y > t_2 y]}{\mathbf{P}[\Theta X > x, \Delta Y > y]} = \lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{E}[\Theta^{\alpha_1} \Delta^{\alpha_2}] \mathbf{P}[X > t_1 x, Y > t_2 y]}{\mathbf{E}[\Theta^{\alpha_1} \Delta^{\alpha_2}] \mathbf{P}[X > x, Y > y]} = t_1^{-\alpha_1} t_2^{-\alpha_2},$$

where in the last step we used the fact that  $(X, Y) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ . By the assumption  $\mathbf{E}[\Theta^{\alpha_1+\varepsilon} \Delta^{\alpha_2+\varepsilon}] < \infty$ , for some  $\varepsilon > 0$ , we have that

$$\mathbf{E}[\Theta^{\alpha_1+\varepsilon}] < \infty, \quad \mathbf{E}[\Delta^{\alpha_2+\varepsilon}] < \infty,$$

for some  $\varepsilon > 0$ , and by Breiman's Theorem (see for example Proposition 5.2 of [41]), we have that  $\Theta X \in \mathcal{R}_{-\alpha_1}$  and  $\Delta Y \in \mathcal{R}_{-\alpha_2}$ , which in combination with the last equation provides  $(\Theta X, \Delta Y) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ .  $\square$

**Remark 4.2.** The previous result, except closure property, provides property (4.27) as well, which can be helpful to direct asymptotic expression of the joint asymptotic behavior of the randomly weighted sums, in the frame of class  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ . We observe that the random weights  $\Theta, \Delta$  are arbitrarily dependent, that can play crucial in actuarial applications, since they represent discount factors.

The following proposition establishes the closure property with respect to finite mixture in classes  $\mathcal{D}^{(2)}, \mathcal{L}^{(2)}, \mathcal{C}^{(2)}, \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ .



**Proposition 4.1.** *Let  $X_1, X_2, Y_1, Y_2$  be real random variables, with distributions  $F_1, F_2, G_1, G_2$  respectively and  $\mathbf{F} = (F_1, G_1)$  and  $\mathbf{G} = (F_2, G_2)$ .*

- (1) *Assume that  $\mathbf{F} = (F_1, G_1) \in \mathcal{D}^{(2)}$ .*
  - (i) *If  $\mathbf{G} \in \mathcal{D}^{(2)}$ , then  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{D}^{(2)}$ , for any  $p \in (0, 1)$ .*
  - (ii) *If  $\overline{\mathbf{G}}_1(x, y) = o[\overline{\mathbf{F}}_1(x, y)]$  and holds some of the following*
    - (a)  $\overline{F}_2(x) = o[\overline{F}_1(x)]$ ,  $\overline{G}_2(x) = o[\overline{G}_1(x)]$ , as  $x \rightarrow \infty$ ,
    - (b)  $\overline{F}_2(x) = o[\overline{F}_1(x)]$ , as  $x \rightarrow \infty$ ,  $G_2 \in \mathcal{D}$ ,
    - (c)  $F_2 \in \mathcal{D}$ ,  $\overline{G}_2(x) = o[\overline{G}_1(x)]$ , as  $x \rightarrow \infty$ ,
    - (d)  $G_2 \in \mathcal{D}$ ,  $G_2 \in \mathcal{D}$ ,*then  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{D}^{(2)}$ , for any  $p \in (0, 1)$ .*
- (2) *Assume that  $\mathbf{F} = (F_1, G_1) \in \mathcal{L}^{(2)}$ .*
  - (i) *If  $\mathbf{G} \in \mathcal{L}^{(2)}$ , then  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{L}^{(2)}$ , for any  $p \in (0, 1)$ .*
  - (ii) *If  $\overline{\mathbf{G}}_1(x, y) = o[\overline{\mathbf{F}}_1(x, y)]$  and holds some of the following*
    - (a)  $\overline{F}_2(x) = o[\overline{F}_1(x)]$ ,  $\overline{G}_2(x) = o[\overline{G}_1(x)]$ , as  $x \rightarrow \infty$ ,
    - (b)  $\overline{F}_2(x) = o[\overline{F}_1(x)]$ , as  $x \rightarrow \infty$ ,  $G_2 \in \mathcal{L}$ ,
    - (c)  $F_2 \in \mathcal{L}$ ,  $\overline{G}_2(x) = o[\overline{G}_1(x)]$ , as  $x \rightarrow \infty$ ,
    - (d)  $G_2 \in \mathcal{L}$ ,  $G_2 \in \mathcal{L}$ ,*then  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{L}^{(2)}$ , for any  $p \in (0, 1)$ .*
- (3) *Assume that  $\mathbf{F} = (F_1, G_1) \in (\mathcal{D} \cap \mathcal{L})^{(2)}$ .*
  - (i) *If  $\mathbf{G} \in (\mathcal{D} \cap \mathcal{L})^{(2)}$ , then  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in (\mathcal{D} \cap \mathcal{L})^{(2)}$ , for any  $p \in (0, 1)$ .*
  - (ii) *If  $\overline{\mathbf{G}}_1(x, y) = o[\overline{\mathbf{F}}_1(x, y)]$  and holds some of the following*
    - (a)  $\overline{F}_2(x) = o[\overline{F}_1(x)]$ ,  $\overline{G}_2(x) = o[\overline{G}_1(x)]$ , as  $x \rightarrow \infty$ ,
    - (b)  $\overline{F}_2(x) = o[\overline{F}_1(x)]$ , as  $x \rightarrow \infty$ ,  $G_2 \in (\mathcal{D} \cap \mathcal{L})$ ,
    - (c)  $F_2 \in (\mathcal{D} \cap \mathcal{L})$ ,  $\overline{G}_2(x) = o[\overline{G}_1(x)]$ , as  $x \rightarrow \infty$ ,
    - (d)  $G_2 \in (\mathcal{D} \cap \mathcal{L})$ ,  $G_2 \in (\mathcal{D} \cap \mathcal{L})$ ,*then  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in (\mathcal{D} \cap \mathcal{L})^{(2)}$ , for any  $p \in (0, 1)$ .*
- (4) *Assume that  $\mathbf{F} = (F_1, G_1) \in \mathcal{C}^{(2)}$ .*
  - (i) *If  $\mathbf{G} \in \mathcal{C}^{(2)}$ , then  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{C}^{(2)}$ , for any  $p \in (0, 1)$ .*
  - (ii) *If  $\overline{\mathbf{G}}_1(x, y) = o[\overline{\mathbf{F}}_1(x, y)]$  and holds some of the following*
    - (a)  $\overline{F}_2(x) = o[\overline{F}_1(x)]$ ,  $\overline{G}_2(x) = o[\overline{G}_1(x)]$ , as  $x \rightarrow \infty$ ,
    - (b)  $\overline{F}_2(x) = o[\overline{F}_1(x)]$ , as  $x \rightarrow \infty$ ,  $G_2 \in \mathcal{C}$ ,
    - (c)  $F_2 \in \mathcal{C}$ ,  $\overline{G}_2(x) = o[\overline{G}_1(x)]$ , as  $x \rightarrow \infty$ ,
    - (d)  $G_2 \in \mathcal{C}$ ,  $G_2 \in \mathcal{C}$ ,*then  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{C}^{(2)}$ , for any  $p \in (0, 1)$ .*
- (5) *Assume that  $\mathbf{F} = (F_1, G_1) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , for some  $\alpha_1 > 0$   $\alpha_2 > 0$ .*
  - (i) *If  $\mathbf{G} \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , then  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , for any  $p \in (0, 1)$ .*
  - (ii) *If  $\overline{\mathbf{G}}_1(x, y) = o[\overline{\mathbf{F}}_1(x, y)]$  and holds some of the following*
    - (a)  $\overline{F}_2(x) = o[\overline{F}_1(x)]$ ,  $\overline{G}_2(x) = o[\overline{G}_1(x)]$ , as  $x \rightarrow \infty$ ,
    - (b)  $\overline{F}_2(x) = o[\overline{F}_1(x)]$ , as  $x \rightarrow \infty$ ,  $G_2 \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ ,
    - (c)  $F_2 \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ ,  $\overline{G}_2(x) = o[\overline{G}_1(x)]$ , as  $x \rightarrow \infty$ ,
    - (d)  $G_2 \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ ,  $G_2 \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ ,*then  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , for any  $p \in (0, 1)$ .*

**Proof.**

(1) (i) Let  $\mathbf{b} = (b_1, b_2) \in (0, 1)^2$ , then

$$\begin{aligned} & \limsup_{x \wedge y \rightarrow \infty} \frac{p \overline{\mathbf{F}}_{\mathbf{b}}(x, y) + (1-p) \overline{\mathbf{G}}_{\mathbf{b}}(x, y)}{p \overline{\mathbf{F}}_1(x, y) + (1-p) \overline{\mathbf{G}}_1(x, y)} \\ &= \limsup_{x \wedge y \rightarrow \infty} \frac{p \mathbf{P}[X_1 > b_1 x, Y_1 > b_2 y] + (1-p) \mathbf{P}[X_2 > b_1 x, Y_2 > b_2 y]}{p \mathbf{P}[X_1 > x, Y_1 > y] + (1-p) \mathbf{P}[X_2 > x, Y_2 > y]} \quad (4.28) \\ &\leq \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_1 > b_1 x, Y_1 > b_2 y]}{\mathbf{P}[X_1 > x, Y_1 > y]} \bigvee \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_2 > b_1 x, Y_2 > b_2 y]}{\mathbf{P}[X_2 > x, Y_2 > y]} < \infty, \end{aligned}$$

for any  $p \in (0, 1)$ , where in the last step we used that  $\mathbf{F}, \mathbf{G} \in \mathcal{D}^{(2)}$ .

Next, since  $\mathbf{F}, \mathbf{G} \in \mathcal{D}^{(2)}$ , it follows that  $F_1, F_2, G_1, G_2 \in \mathcal{D}$ , whence by [41, Prop. 3.7(iv)] we obtain  $p \overline{F}_1 + (1-p) \overline{F}_2 \in \mathcal{D}$  and  $p \overline{G}_1 + (1-p) \overline{G}_2 \in \mathcal{D}$  for any  $p \in (0, 1)$ . This last in combination with relation (4.28), provides  $p \overline{\mathbf{F}} + (1-p) \overline{\mathbf{G}} \in \mathcal{D}^{(2)}$ , for any  $p \in (0, 1)$ .

(ii) From  $\overline{\mathbf{G}}_1(x, y) = o[\overline{\mathbf{F}}_1(x, y)]$  we obtain

$$\limsup_{x \wedge y \rightarrow \infty} \frac{p \overline{\mathbf{F}}_{\mathbf{b}}(x, y) + (1-p) \overline{\mathbf{G}}_{\mathbf{b}}(x, y)}{p \overline{\mathbf{F}}_1(x, y) + (1-p) \overline{\mathbf{G}}_1(x, y)} = \limsup_{x \wedge y \rightarrow \infty} \frac{p \overline{\mathbf{F}}_{\mathbf{b}}(x, y)}{p \overline{\mathbf{F}}_1(x, y)} < \infty. \quad (4.29)$$

Since  $\mathbf{F} \in \mathcal{D}^{(2)}$ , we find  $F_1, G_1 \in \mathcal{D}$  and by [41, Prop. 3.7(iv)] we find that each of the (a) - (d) implies  $p \overline{F}_1 + (1-p) \overline{F}_2 \in \mathcal{D}$  and  $p \overline{G}_1 + (1-p) \overline{G}_2 \in \mathcal{D}$  for any  $p \in (0, 1)$ . This, in combination with relation (4.29), imply that  $p \overline{\mathbf{F}} + (1-p) \overline{\mathbf{G}} \in \mathcal{D}^{(2)}$ , for any  $p \in (0, 1)$ .

(2) (i) Initially, for any  $\mathbf{a} = (a_1, a_2) > (0, 0)$ , then

$$\frac{p \mathbf{P}[X_1 > x - a_1, Y_1 > y - a_2] + (1-p) \mathbf{P}[X_2 > x - a_1, Y_2 > y - a_2]}{p \mathbf{P}[X_1 > x, Y_1 > y] + (1-p) \mathbf{P}[X_2 > x, Y_2 > y]} \geq 1.$$

for any  $x, y \in \mathbb{R}$ . If  $\mathbf{G} \in \mathcal{L}^{(2)}$  then

$$\begin{aligned} & \lim_{x \wedge y \rightarrow \infty} \frac{p \overline{\mathbf{F}}_1(x - a_1, y - a_2) + (1-p) \overline{\mathbf{G}}_1(x - a_1, y - a_2)}{p \overline{\mathbf{F}}_1(x, y) + (1-p) \overline{\mathbf{G}}_1(x, y)} \\ &= \lim_{x \wedge y \rightarrow \infty} \frac{p \mathbf{P}[X_1 > x - a_1, Y_1 > y - a_2] + (1-p) \mathbf{P}[X_2 > x - a_1, Y_2 > y - a_2]}{p \mathbf{P}[X_1 > x, Y_1 > y] + (1-p) \mathbf{P}[X_2 > x, Y_2 > y]} \quad (4.30) \\ &\leq \lim_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_1 > x - a_1, Y_1 > y - a_2]}{\mathbf{P}[X_1 > x, Y_1 > y]} \bigvee \limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[X_2 > x - a_1, Y_2 > y - a_2]}{\mathbf{P}[X_2 > x, Y_2 > y]} = 1, \end{aligned}$$

where in the last step we use that  $\mathbf{F}, \mathbf{G} \in \mathcal{L}^{(2)}$ , hence from relations (4.29) and (4.30) we obtain the joint property of  $\mathcal{L}^{(2)}$  and furthermore, because the marginal distributions belong to  $\mathcal{L}$ , by [41, Prop. 3.9(iii)], we have that the mixtures in (4.29) belong to class  $\mathcal{L}$ , thence we conclude  $p \overline{\mathbf{F}} + (1-p) \overline{\mathbf{G}} \in \mathcal{L}^{(2)}$ .

(ii) Since  $\overline{\mathbf{G}}_1(x, y) = o[\overline{\mathbf{F}}_1(x, y)]$  holds, we find

$$\limsup_{x \wedge y \rightarrow \infty} \frac{p \overline{\mathbf{F}}_1(x - a_1, y - a_2) + (1-p) \overline{\mathbf{G}}_1(x - a_1, y - a_2)}{p \overline{\mathbf{F}}_1(x, y) + (1-p) \overline{\mathbf{G}}_1(x, y)} = 1. \quad (4.31)$$

Additionally, we observe that by  $\mathbf{F} \in \mathcal{L}^{(2)}$ , follows that  $F_1, G_1 \in \mathcal{L}$  and then each of (a) - (d) implies the closure property with respect to uni-variate finite mixture

for class  $\mathcal{L}$  in relations (2.8), from [41, Prop. 3.9(iii)]. Thus, in combination with relation (4.31), we obtain  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{L}^{(2)}$ , for any  $p \in (0, 1)$ .

- (3) (i) It follows directly from (1)(i) and (2)(i).  
(ii) It follows directly from (1)(ii) and (2)(ii).  
(4) (i) Initially, we observe that

$$\lim_{z \uparrow 1} \lim_{x \wedge y \rightarrow \infty} \frac{p\overline{\mathbf{F}}_z(x, y) + (1-p)\overline{\mathbf{G}}_z(x, y)}{p\overline{\mathbf{F}}_1(x, y) + (1-p)\overline{\mathbf{G}}_1(x, y)} \geq 1, \quad (4.32)$$

for any  $p \in (0, 1)$ , for any distribution. From the other side, since  $\mathbf{F}, \mathbf{G} \in \mathcal{C}^{(2)}$  we can write

$$\begin{aligned} & \lim_{z \uparrow 1} \lim_{x \wedge y \rightarrow \infty} \frac{p\overline{\mathbf{F}}_z(x, y) + (1-p)\overline{\mathbf{G}}_z(x, y)}{p\overline{\mathbf{F}}_1(x, y) + (1-p)\overline{\mathbf{G}}_1(x, y)} \\ & \leq \lim_{z \uparrow 1} \lim_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{F}}_z(x, y)}{\overline{\mathbf{F}}_1(x, y)} \bigvee \lim_{z \uparrow 1} \lim_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{G}}_z(x, y)}{\overline{\mathbf{G}}_1(x, y)} = 1. \end{aligned} \quad (4.33)$$

From relations (4.32) and (4.33) and taking into account that  $F_1, F_2, G_1, G_2 \in \mathcal{C}$ , we find that relations (2.8) are satisfied for class  $\mathcal{C}$ , whence by [41, Prop. 3.5(iii)] is implied that  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{C}^{(2)}$  for any  $p \in (0, 1)$ .

(ii) From relation  $\overline{\mathbf{G}}_1(x, y) = o[\overline{\mathbf{F}}_1(x, y)]$ , having in mind that  $\mathcal{C}^{(2)} \subsetneq \mathcal{D}^{(2)}$ , we obtain

$$\lim_{z \uparrow 1} \lim_{x \wedge y \rightarrow \infty} \frac{p\overline{\mathbf{F}}_z(x, y) + (1-p)\overline{\mathbf{G}}_z(x, y)}{p\overline{\mathbf{F}}_1(x, y) + (1-p)\overline{\mathbf{G}}_1(x, y)} = \lim_{z \uparrow 1} \lim_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{F}}_z(x, y)}{\overline{\mathbf{F}}_1(x, y)} = 1, \quad (4.34)$$

for any  $p \in (0, 1)$ . Further, applying [41, Prop. 3.5(iii)], we find that, under each of (a) - (d), the finite mixtures in (2.8) are satisfied for class  $\mathcal{C}$ , which in combination with relation (4.34) implies that  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{C}^{(2)}$  for any  $p \in (0, 1)$ .

- (5) (i) For any  $\mathbf{t} = (t_1, t_2) \in (0, 1)^2$ , the inequality

$$\lim_{x \wedge y \rightarrow \infty} \frac{p\overline{\mathbf{F}}_{\mathbf{t}}(x, y) + (1-p)\overline{\mathbf{G}}_{\mathbf{t}}(x, y)}{p\overline{\mathbf{F}}_1(x, y) + (1-p)\overline{\mathbf{G}}_1(x, y)} \leq \lim_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{F}}_{\mathbf{t}}(x, y)}{\overline{\mathbf{F}}_1(x, y)} \bigvee \lim_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{G}}_{\mathbf{t}}(x, y)}{\overline{\mathbf{G}}_1(x, y)} = t_1^{-\alpha_1} t_2^{-\alpha_2}, \quad (4.35)$$

is true, for any  $p \in (0, 1)$ , where in the last step we used that  $\mathbf{F}, \mathbf{G} \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ . From the other side we find similarly

$$\lim_{x \wedge y \rightarrow \infty} \frac{p\overline{\mathbf{F}}_{\mathbf{t}}(x, y) + (1-p)\overline{\mathbf{G}}_{\mathbf{t}}(x, y)}{p\overline{\mathbf{F}}_1(x, y) + (1-p)\overline{\mathbf{G}}_1(x, y)} \geq \lim_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{F}}_{\mathbf{t}}(x, y)}{\overline{\mathbf{F}}_1(x, y)} \bigwedge \lim_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{G}}_{\mathbf{t}}(x, y)}{\overline{\mathbf{G}}_1(x, y)} = t_1^{-\alpha_1} t_2^{-\alpha_2}. \quad (4.36)$$

Further, since  $\mathbf{F}, \mathbf{G} \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ , we find that  $F_1, F_2 \in \mathcal{R}_{-\alpha_1}$  and  $G_1, G_2 \in \mathcal{R}_{-\alpha_2}$  and by [41, Prop. 3.3(iii)], we obtain that the finite mixtures  $p\overline{F}_1 + (1-p)\overline{F}_2 \in \mathcal{R}_{-\alpha_1}$  and  $p\overline{G}_1 + (1-p)\overline{G}_2 \in \mathcal{R}_{-\alpha_2}$ , for any  $p \in (0, 1)$ . Therefore, in combination with relations (4.35) and (4.36) we conclude  $p\overline{\mathbf{F}} + (1-p)\overline{\mathbf{G}} \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$  for any  $p \in (0, 1)$ .

(ii) Because of  $\overline{\mathbf{G}}_1(x, y) = o[\overline{\mathbf{F}}_1(x, y)]$  and taking into account that  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)} \subsetneq \mathcal{D}^{(2)}$ , we obtain for any  $\mathbf{t} = (t_1, t_2) \in (0, \infty)^2$ , the relation

$$\lim_{x \wedge y \rightarrow \infty} \frac{p \overline{\mathbf{F}}_{\mathbf{t}}(x, y) + (1-p) \overline{\mathbf{G}}_{\mathbf{t}}(x, y)}{p \overline{\mathbf{F}}_1(x, y) + (1-p) \overline{\mathbf{G}}_1(x, y)} = \lim_{x \wedge y \rightarrow \infty} \frac{\overline{\mathbf{F}}_{\mathbf{t}}(x, y)}{\overline{\mathbf{F}}_1(x, y)} = t_1^{-\alpha_1} t_2^{-\alpha_2}, \quad (4.37)$$

and from [41, Prop. 3.3(iii)], we find that for each of (a) - (d), the finite mixtures in (2.8) are satisfied, which in combination with relation (4.37) implies that  $p \overline{\mathbf{F}}_1 + (1-p) \overline{\mathbf{F}}_2 \in \mathcal{R}_{-\alpha_1}$  and  $p \overline{\mathbf{G}}_1 + (1-p) \overline{\mathbf{G}}_2 \in \mathcal{R}_{-\alpha_2}$  for any  $p \in (0, 1)$ .  $\square$

## 5. RANDOMLY WEIGHTED SUMS

In this section we study the joint behavior of randomly weighted sums. There are several work around the formula (1.7), see for example [46], [56], [65]. This joint behavior can be applied on several areas of actuarial science and financial mathematics. Indeed, in risk theory the main components  $X_i$  and  $Y_j$  represent gains or losses in  $i$ -th and  $j$ -th period, while the random weights  $\Theta_i$  and  $\Delta_j$  represent discount factors, that are allowed to be degenerated to some non-zero number, see for example [64]. In credit risk applications, the random variables  $X_i$  and  $Y_j$  represent the rate of default of the  $i$ -th and  $j$ -th debtor and the random weights  $\Theta_i$  and  $\Delta_j$  represent Bernoulli trials with values zero and unity, where the unity reflect the case of default.

Before the main result, we need a lemma, which examines the closure property for the dependencies  $GTAI$  and  $GQAI$  with respect to the convolution product under the distribution class  $\mathcal{D}^{(2)}$  for the main variables, see some results for  $TAI$  and  $QAI$  in [43, Th. 2.2].

**Lemma 5.1.** *Let  $X_1, \dots, X_n, Y_1, \dots, Y_m$  be real random variables with corresponding distributions  $F_1, \dots, F_n, G_1, \dots, G_m$ , from class  $\mathcal{D}$ , which are  $GTAI$ , (or  $GQAI$ ), and for any  $i = 1, \dots, n, j = 1, \dots, m$  hold  $(X_i, Y_j) \in \mathcal{D}^{(2)}$ . Let  $\Theta_1, \dots, \Theta_n, \Delta_1, \dots, \Delta_m$  be non-negative, non-degenerated to zero random variables with  $(\Theta_i, \Delta_j)$  satisfying Assumption 4.1, (or  $(\Theta_i X_i, \Delta_j Y_j)$  satisfying Assumption 4.1 respectively). If additionally we assume  $\Theta_1, \dots, \Theta_n, \Delta_1, \dots, \Delta_m$  are independent of  $X_1, \dots, X_n, Y_1, \dots, Y_m$ , then the products  $\Theta_1 X_1, \dots, \Theta_n X_n, \Delta_1 Y_1, \dots, \Delta_m Y_m$  are  $GTAI$ , (or  $GQAI$  respectively).*

**Proof.** Let begin with the  $GTAI$  case. We denote the maximum  $\widehat{\Theta} := \Theta_i \vee \Theta_k$ , for  $i \neq k \in \{1, \dots, n\}$  and by Assumption 4.1, and  $\widehat{b}(x_i \wedge x_k) := \max\{b_i(x_i \wedge x_k), b_k(x_i \wedge x_k)\}$ , where the functions  $b_i$  and  $b_k$  are defined according to Assumption 4.1 for the random weights  $\Theta_i$  and  $\Theta_k$ , respectively, we find that  $\widehat{b}(x_i \wedge x_k) \rightarrow \infty$  and  $\widehat{b}(x_i \wedge x_k) = o(x_i \wedge x_k)$ , as  $x_i \wedge x_k \rightarrow \infty$  and further for some  $j = 1, \dots, m$

$$\mathbf{P}[\widehat{\Theta} > \widehat{b}(x_i \wedge x_k)] = o(\mathbf{P}[\Theta_k X_k > x, \Delta_j Y_j > y]), \quad (5.1)$$

as  $x_i \wedge x_k \rightarrow \infty$ . Indeed, if  $\widehat{\Theta} = \Theta_k$ , then by Assumption 4.1 relation (5.1) is obvious. If  $\widehat{\Theta} = \Theta_i$ , we multiply and divide with  $\mathbf{P}[\Theta_i X_i > x_i, \Delta_j Y_j > y_j]$  to find that

$$\frac{\mathbf{P}[\Theta_i > \widehat{b}(x_i \wedge x_k)]}{\mathbf{P}[\Theta_i X_i > x_i, \Delta_j Y_j > y_j]},$$

tends to zero, while the ratio

$$\frac{\mathbf{P}[\Theta_i X_i > x_i, \Delta_j Y_j > y_j]}{\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]},$$

is bounded by

$$\frac{1}{\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]},$$

which is finite (because  $X, Y$  have distributions with infinity upper bound).

Hence, for a function  $c_j$ , defined in Assumption 4.1 for the random variable  $Y_j$ , for  $x_i, x_k, y_j > 0$  we obtain

$$\begin{aligned} \mathbf{P}[|\Theta_i X_i| > x_i, \Theta_k X_k > x_k, \Delta_j Y_j > y_j] &\leq \mathbf{P}\left[\widehat{\Theta}|X_i| > x_i, \widehat{\Theta} X_k > x_k, \Delta_j Y_j > y_j\right] \\ &= \int_0^\infty \int_0^\infty \mathbf{P}\left[|X_i| > \frac{x_i}{t}, X_k > \frac{x_k}{t}, Y_j > \frac{y_j}{s}\right] \mathbf{P}\left[\widehat{\Theta} \in dt, \Delta_j \in ds\right] \\ &= \left(\int_0^{\widehat{b}(x_i \wedge x_k)} + \int_{\widehat{b}(x_i \wedge x_k)}^\infty\right) \left(\int_0^{c_j(y_j)} + \int_{c_j(y_j)}^\infty\right) \mathbf{P}\left[|X_i| > \frac{x_i}{t}, X_k > \frac{x_k}{t}, Y_j > \frac{y_j}{s}\right] \\ &\quad \times \mathbf{P}\left[\widehat{\Theta} \in dt, \Delta_j \in ds\right] \\ &=: L_{11}(x_i, x_k, y_j) + L_{12}(x_i, x_k, y_j) + L_{21}(x_i, x_k, y_j) + L_{22}(x_i, x_k, y_j), \end{aligned} \tag{5.2}$$

Thence we find

$$\begin{aligned} L_{22}(x_i, x_k, y_j) &= \int_{\widehat{b}(x_i \wedge x_k)}^\infty \int_{c_j(y_j)}^\infty \mathbf{P}\left[|X_i| > \frac{x_i}{t}, X_k > \frac{x_k}{t}, Y_j > \frac{y_j}{s}\right] \mathbf{P}\left[\widehat{\Theta} \in dt, \Delta_j \in ds\right] \\ &\leq \mathbf{P}\left[\widehat{\Theta} > \widehat{b}(x_i \wedge x_k), \Delta_j > c_j(y_j)\right] \leq \mathbf{P}\left[\Delta_j > c_j(y_j)\right] \\ &= o(\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]), \end{aligned} \tag{5.3}$$

as  $x_i \wedge x_k \wedge y_j \rightarrow \infty$ . Next we calculate

$$\begin{aligned} L_{21}(x_i, x_k, y_j) &= \int_{\widehat{b}(x_i \wedge x_k)}^\infty \int_0^{c_j(y_j)} \mathbf{P}\left[|X_i| > \frac{x_i}{t}, X_k > \frac{x_k}{t}, Y_j > \frac{y_j}{s}\right] \mathbf{P}\left[\widehat{\Theta} \in dt, \Delta_j \in ds\right] \\ &\leq \mathbf{P}\left[\widehat{\Theta} > \widehat{b}(x_i \wedge x_k), \Delta_j \leq c_j(y_j)\right] \leq \mathbf{P}\left[\widehat{\Theta} > \widehat{b}(x_i \wedge x_k)\right] \\ &= o(\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]), \end{aligned}$$

as  $x_i \wedge x_k \wedge y_j \rightarrow \infty$ . Similarly we obtain

$$\begin{aligned} L_{12}(x_i, x_k, y_j) &\leq \mathbf{P}\left[\widehat{\Theta} > \widehat{b}(x_i \wedge x_k), \Delta_j \leq c_j(y_j)\right] \leq \mathbf{P}\left[\Delta_j > c_j(y_j)\right] \\ &= o(\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]), \end{aligned}$$

as  $x_i \wedge x_k \wedge y_j \rightarrow \infty$ . For the first term from the *GTAI* property of  $X_1, \dots, X_n, Y_1, \dots, Y_m$  and the definitions of  $\widehat{b}$  and  $c_j$  we have that

$$\begin{aligned} L_{11}(x_i, x_k, y_j) &\leq \mathbf{P}\left[|X_i| > \frac{x_i}{\widehat{b}(x_i \wedge x_k)}, X_k > \frac{x_k}{\widehat{b}(x_i \wedge x_k)}, Y_j > \frac{y_j}{c_j(y_j)}\right] \\ &= o(\mathbf{P}[X_k > x_k, Y_j > y_j]), \end{aligned} \tag{5.4}$$

as  $x_i \wedge x_k \wedge y_j \rightarrow \infty$ , where in the last step we use that  $(X_k, Y_j) \in \mathcal{D}^{(2)}$ , for any  $i = 1, \dots, n$ ,  $j = 1, \dots, m$ . Therefore by (5.2), together with (5.3) (5.4) we conclude that

$$\begin{aligned} \mathbf{P}[|\Theta_i X_i| > x_i, \Theta_k X_k > x_k, \Delta_j Y_j > y_j] &= o(\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]) \\ &+ o(\mathbf{P}[X_k > x_k, Y_j > y_j]) = o(\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]), \end{aligned} \quad (5.5)$$

as  $x_i \wedge x_k \wedge y_j \rightarrow \infty$ , where in last step we used that form assumptions follows

$$\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j] \asymp \mathbf{P}[X_k > x_k, Y_j > y_j], \quad (5.6)$$

as  $x_k \wedge y_j \rightarrow \infty$ . Indeed, since

$$\begin{aligned} &\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j] \quad (5.7) \\ &= \left( \int_0^{b_k(x_k)} + \int_{b_k(x_k)}^\infty \right) \left( \int_0^{c_j(y_j)} + \int_{c_j(y_j)}^\infty \right) \mathbf{P} \left[ X_k > \frac{x_k}{t}, Y_j > \frac{y_j}{s} \right] \mathbf{P}[\Theta_k \in dt, \Delta_j \in ds] \\ &=: M_{11}(x_k, y_j) + M_{12}(x_k, y_j) + M_{21}(x_k, y_j) + M_{22}(x_k, y_j), \end{aligned}$$

From definitions of functions  $b_k$  and  $c_j$ , see Assumption 4.1 and by  $(X_k, Y_j) \in \mathcal{D}^{(2)}$  we find

$$M_{11}(x_k, y_j) \asymp \mathbf{P}[X_k > x_k, Y_j > y_j], \quad (5.8)$$

as  $x_k \wedge y_j \rightarrow \infty$ . Further we observe that

$$\begin{aligned} M_{22}(x_k, y_j) &\leq \mathbf{P}[\Theta_k > b_k(x_k), \Delta_j > c_j(y_j)] \leq \mathbf{P}[\Delta_j > c_j(y_j)] \\ &= o(\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]), \end{aligned}$$

as  $y_j \rightarrow \infty$ , while  $x_k$  is free, where in the last step we employed Assumption 4.1. Next, we see that

$$\begin{aligned} M_{21}(x_k, y_j) &\leq \mathbf{P}[\Theta_k > b_k(x_k), \Delta_j \leq c_j(y_j)] \leq \mathbf{P}[\Theta_k > b_k(x_k)] \\ &= o(\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]), \end{aligned}$$

as  $x_k \rightarrow \infty$ , while  $y_j$  remains free. Similarly by symmetry we obtain

$$M_{12}(x_k, y_j) = o(\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]), \quad (5.9)$$

as  $y_j \rightarrow \infty$ , while  $x_k$  is free. Hence, from relations (5.8) - (5.9), together with (5.7), we get

$$\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j] \leq \mathbf{P} \left[ X_k > \frac{x_k}{b_k(x_k)}, Y_j > \frac{y_j}{c_j(y_j)} \right] \asymp \mathbf{P}[X_k > x_k, Y_j > y_j],$$

as  $x_k \wedge y_j \rightarrow \infty$ . As a result we become

$$\limsup_{x \wedge y \rightarrow \infty} \frac{\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]}{\mathbf{P}[X_k > x_k, Y_j > y_j]} < \infty. \quad (5.10)$$

From the other side, for some arbitrarily chosen  $\varepsilon \in (0, 1)$ , we obtain

$$\begin{aligned}
& \mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j] \tag{5.11} \\
& \geq \left( \int_\varepsilon^1 + \int_1^\infty \right) \left( \int_\varepsilon^1 + \int_1^\infty \right) \mathbf{P} \left[ X_k > \frac{x_k}{\theta_k}, Y_j > \frac{y_j}{\delta_j} \right] \mathbf{P}[\Theta_k \in d\theta_k, \Delta_j \in d\delta_j] \\
& \geq d_{1,1} \mathbf{P}[X_k > x_k, Y_j > y_j] \mathbf{P}[\Theta_k \in (\varepsilon, 1], \Delta_j \in (\varepsilon, 1]] \\
& + \int_1^\infty \int_\varepsilon^1 \mathbf{P} \left[ X_k > \frac{x_k}{\theta_k}, Y_j > y_j \right] \mathbf{P}[\Theta_k \in d\theta_k, \Delta_j \in d\delta_j] \\
& + \int_\varepsilon^1 \int_1^\infty \mathbf{P} \left[ X_k > x_k, Y_j > \frac{y_j}{\delta_j} \right] \mathbf{P}[\Theta_k \in d\theta_k, \Delta_j \in d\delta_j] \\
& + d_{2,2} \mathbf{P}[X_k > x_k, Y_j > y_j] \mathbf{P}[\Theta_k \in (1, \infty), \Delta_j \in (1, \infty)] \\
& \geq d_{1,1} \mathbf{P}[X_k > x_k, Y_j > y_j] \mathbf{P}[\Theta_k \in (\varepsilon, 1], \Delta_j \in (\varepsilon, 1]] \\
& + d_{1,2} \mathbf{P}[X_k > x_k, Y_j > y_j] \mathbf{P}[\Theta_k \in (\varepsilon, 1], \Delta_j \in (1, \infty)] \\
& + d_{2,1} \mathbf{P}[X_k > x_k, Y_j > y_j] \mathbf{P}[\Theta_k \in (1, \infty), \Delta_j \in (\varepsilon, 1]] \\
& + d_{2,2} \mathbf{P}[X_k > x_k, Y_j > y_j] \mathbf{P}[\Theta_k \in (1, \infty), \Delta_j \in (1, \infty)] \\
& \geq (d_{1,1} + d_{1,2} + d_{2,1} + d_{2,2}) \mathbf{P}[X_k > x_k, Y_j > y_j] (\mathbf{P}[\Theta_k \in (\varepsilon, 1], \Delta_j \in (\varepsilon, 1]] \\
& + \mathbf{P}[\Theta_k \in (\varepsilon, 1], \Delta_j \in (1, \infty)] + \mathbf{P}[\Theta_k \in (1, \infty), \Delta_j \in (\varepsilon, 1]] \\
& + \mathbf{P}[\Theta_k \in (1, \infty), \Delta_j \in (1, \infty)]) \rightarrow (d_{1,1} + d_{1,2} + d_{2,1} + d_{2,2}) \mathbf{P}[X_k > x_k, Y_j > y_j],
\end{aligned}$$

as  $\varepsilon \downarrow 0$ , where the  $d_{2,2} > 0$  follows by class  $\mathcal{D}^{(2)}$  property and the inclusion  $(\theta_k, \delta_j) \in (1, \infty) \times (1, \infty)$ . The inequalities  $d_{1,1}, d_{1,2}, d_{2,1} > 0$  follow from the intervals where the pairs  $(\theta_k, \delta_j)$  belong. Hence by (5.11) we obtain

$$\limsup_{x_k \wedge y_j \rightarrow \infty} \frac{\mathbf{P}[X_k > x_k, Y_j > y_j]}{\mathbf{P}[\Theta_k X_k > x_k, \Delta_j Y_j > y_j]} < \infty, \tag{5.12}$$

whence from relations (5.10) and (5.12) we find (5.6).

With similar handling, because of symmetry, we obtain

$$\mathbf{P}[|\Delta_j Y_j| > y_j, \Delta_k Y_k > y_k, \Theta_i X_i > x_i] = o(\mathbf{P}[\Theta_i X_i > x_i, \Delta_k Y_k > y_k]), \tag{5.13}$$

as  $x_i \wedge y_k \wedge y_j \rightarrow \infty$ . So from relations (5.5) and (5.13), we find that the  $\Theta_1 X_1, \dots, \Theta_n X_n$ , and  $\Delta_1 Y_1, \dots, \Delta_m Y_m$  are *GTAI*.

For the second case, with *GQAI* dependence, we follow the same route, with the only difference that the convergences are with  $x \wedge y \rightarrow \infty$ , and in relations (5.4) and (5.5) the last term takes the form  $o(\mathbf{P}[\Theta_i X_i > x, \Delta_j Y_j > y] + \mathbf{P}[\Theta_k X_k > x, \Delta_j Y_j > y])$ .  $\square$

Now we are ready to present a weighted form of Theorem 3.1, and the same time to generalize [38, Th. 6.1], since the random weights  $\Theta_i, \Delta_j$  are no more strictly positive and bounded from above and the main components  $X_i, Y_j$  are real random variables.

**Theorem 5.1.** (1) *Let  $X_1, \dots, X_n, Y_1, \dots, Y_m$  be GQAI, real random variables with corresponding distributions  $F_1, \dots, F_n, G_1, \dots, G_m$ , from class  $\mathcal{C}$ , and for any  $i =$*

$1, \dots, n, j = 1, \dots, m$  hold  $(X_i, Y_j) \in \mathcal{C}^{(2)}$ . Let  $\Theta_1, \dots, \Theta_n, \Delta_1, \dots, \Delta_m$  be non-negative, non-degenerated to zero random variables, independent of the main variables  $X_1, \dots, X_n, Y_1, \dots, Y_m$ , with  $(\Theta_i, \Delta_j)$  satisfying Assumption 4.1. Then

$$\mathbf{P}[S_n^\Theta > x, T_m^\Delta > y] \sim \mathbf{P}\left[\bigvee_{i=1}^n S_i^\Theta > x, \bigvee_{j=1}^m T_j^\Delta > y\right] \sim \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[\Theta_i X_i > x, \Delta_j Y_j > y], \quad (5.14)$$

as  $x \wedge y \rightarrow \infty$ .

- (2) If hold the assumptions of part (1), with the only differences that  $n = m$ , the  $X_1, \dots, X_n, Y_1, \dots, Y_m$  are GTAI, real, random variables with corresponding distributions  $F_1, \dots, F_n, G_1, \dots, G_n$ , from class  $\mathcal{D} \cap \mathcal{L}$ , and for any  $i, j = 1, \dots, n$ , hold  $(X_i, Y_j) \in (\mathcal{D} \cap \mathcal{L})^{(2)}$ . If further  $X_1, \dots, X_n$  are TAI and  $Y_1, \dots, Y_n$  are TAI and  $E[\Theta_i^p] < \infty$ , for some  $p > \bigvee_{i=1}^n J_{F_i}^+$ , and  $E[\Delta_j^q] < \infty$ , for some  $q > \bigvee_{j=1}^n J_{G_j}^+$  then

$$\begin{aligned} \mathbf{P}[S_n^\Theta > x, T_m^\Delta > y] &\sim \mathbf{P}\left[\bigvee_{i=1}^n S_i^\Theta > x, \bigvee_{j=1}^m T_j^\Delta > y\right] \sim \mathbf{P}\left[\bigvee_{i=1}^n \Theta_i X_i > x, \bigvee_{j=1}^n \Delta_j Y_j > y\right] \\ &\sim \sum_{i=1}^n \sum_{j=1}^n \mathbf{P}[\Theta_i X_i > x, \Delta_j Y_j > y], \end{aligned} \quad (5.15)$$

as  $x \wedge y \rightarrow \infty$ .

### Proof.

- (1) Rewriting Theorem 3.1, we find that  $Z_1, \dots, Z_n, W_1, \dots, W_m$  are GQAI with distributions from class  $\mathcal{C}$  and for any  $i = 1, \dots, n, j = 1, \dots, m$  hold  $(Z_i, W_j) \in \mathcal{C}^{(2)}$ , whence

$$\begin{aligned} \mathbf{P}\left[\sum_{i=1}^n Z_i > x, \sum_{j=1}^m W_j > y\right] &\sim \mathbf{P}\left[\bigvee_{k=1}^n \sum_{i=1}^k Z_i > x, \bigvee_{l=1}^m \sum_{j=1}^l W_j > y\right] \\ &\sim \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[Z_i > x, W_j > y], \end{aligned} \quad (5.16)$$

as  $x \wedge y \rightarrow \infty$ . From the assumptions of part (1), applying Theorem 4.2(2), since for any  $i = 1, \dots, n, j = 1, \dots, m$  holds  $(X_i, Y_j) \in \mathcal{C}^{(2)}$ , it follows that  $(\Theta_i X_i, \Delta_j Y_j) \in \mathcal{C}^{(2)}$ . Further, applying Lemma 5.1 we find out that the products  $\Theta_1 X_1, \dots, \Theta_n X_n, \Delta_1 Y_1, \dots, \Delta_m Y_m$  are GQAI, as  $\mathcal{C}^{(2)} \subsetneq \mathcal{D}^{(2)}$ . Hence, putting

$$Z_i := \Theta_i X_i, \quad W_j := \Delta_j Y_j, \quad (5.17)$$

for any  $i = 1, \dots, n$  and any  $j = 1, \dots, m$ , we find (5.14) through (5.16).

- (2) Repeating relation (1.8) through the  $Z_1, \dots, Z_n, W_1, \dots, W_n$  that are GTAI (and each one of these sequences are TAI) with distributions from class  $\mathcal{D} \cap \mathcal{L}$ , and with



any  $i, j = 1, \dots, n$ , finding that  $(Z_i, W_j) \in (\mathcal{D} \cap \mathcal{L})^{(2)}$  we obtain

$$\begin{aligned} \mathbf{P} \left[ \sum_{i=1}^n Z_i > x, \sum_{j=1}^n W_j > y \right] &\sim \mathbf{P} \left[ \bigvee_{k=1}^n \sum_{i=1}^k Z_i > x, \bigvee_{l=1}^n \sum_{j=1}^l W_j > y \right] \\ &\sim \mathbf{P} \left[ \bigvee_{i=1}^n Z_i > x, \bigvee_{j=1}^n W_j > y \right] \sim \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[Z_i > x, W_j > y], \end{aligned} \quad (5.18)$$

as  $x \wedge y \rightarrow \infty$ , see also [37, Th. 4.2]. Therefore, since  $(X_i, Y_j) \in (\mathcal{D} \cap \mathcal{L})^{(2)}$ , from Theorem 4.2(4), because of Assumption 4.1, we conclude  $(\Theta_i X_i, \Delta_j Y_j) \in (\mathcal{D} \cap \mathcal{L})^{(2)}$ . Further by Lemma 5.1 we obtain that the  $\Theta_1 X_1, \dots, \Theta_n X_n, \Delta_1 Y_1, \dots, \Delta_n Y_n$  are GTAI, and by the moment condition and the class  $\mathcal{D}$  (for the primary random variables) we have that  $\Theta_1 X_1, \dots, \Theta_n X_n$  are TAI and  $\Delta_1 Y_1, \dots, \Delta_n Y_n$  are TAI, by Theorem 2.2 of [43]. Thus, using (5.17) for any  $i, j = 1, \dots, n$ , we find (5.15) through (5.18).  $\square$

**Remark 5.1.** Comparing the two parts of Theorem 5.1 we realize that as the distribution class  $\mathcal{C}^{(2)}$  increases to  $(\mathcal{D} \cap \mathcal{L})^{(2)}$ , the dependence decreases from GQAI to GTAI and additionally we need  $n = m$ . However, it is remarkable, that in GTAI case we find the asymptotic behavior of the maximums, which is NOT possible in the part (1). Furthermore in the second part we obtain

$$\mathbf{P} \left[ \bigvee_{i=1}^n \Theta_i X_i > x, \bigvee_{j=1}^m \Delta_j Y_j > y \right] \sim \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}[\Theta_i X_i > x, \Delta_j Y_j > y],$$

as  $x \wedge y \rightarrow \infty$ , whose proof follows the same arguments of the part (2), (and without necessary TAI for each sequence!) namely we establish closure property of  $(\mathcal{D} \cap \mathcal{L})^{(2)}$  with respect to GTAI dependence of the products, in combination with [38, Th. 4.1].

The next result, with an extra condition on the moments of the random weights, provides a direct formula for calculation in the distribution class  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$ .

**Corollary 5.1.** (1) Let hold the assumptions of Theorem 5.1(1), with the only difference  $F_1, \dots, F_n \in \mathcal{R}_{-\alpha_1}$  and  $G_1, \dots, G_m \in \mathcal{R}_{-\alpha_2}$  with  $(X_i, Y_j) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$  for any  $i = 1, \dots, n, j = 1, \dots, m$ . Additionally we assume that  $\mathbf{E}[\Theta_i^{\alpha_1 + \varepsilon} \Delta_j^{\alpha_2 + \varepsilon}] < \infty$ , for any  $i = 1, \dots, n$  and  $j = 1, \dots, m$ , for some  $\varepsilon > 0$ . Then

$$\begin{aligned} \mathbf{P}[S_n^\Theta > x, T_m^\Delta > y] &\sim \mathbf{P} \left[ \bigvee_{i=1}^n S_i^\Theta > x, \bigvee_{j=1}^m T_j^\Delta > y \right] \\ &\sim \sum_{i=1}^n \sum_{j=1}^m \mathbf{E}[\Theta_i^{\alpha_1} \Delta_j^{\alpha_2}] \mathbf{P}[X_i > x, Y_j > y]. \end{aligned}$$

as  $x \wedge y \rightarrow \infty$ .

(2) Let hold the assumptions of Theorem 5.1(2), with the only difference  $F_1, \dots, F_n \in \mathcal{R}_{-\alpha_1}$  and  $G_1, \dots, G_m \in \mathcal{R}_{-\alpha_2}$  with  $(X_i, Y_j) \in \mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)}$  for any  $i, j = 1, \dots, n$ . Furthermore we assume that  $\mathbf{E}[\Theta_i^{\alpha_1 + \varepsilon} \Delta_j^{\alpha_2 + \varepsilon}] < \infty$ , for some  $\varepsilon > 0$ , for any  $i, j =$

1, \dots, n. Then

$$\begin{aligned} \mathbf{P}[S_n^\Theta > x, T_n^\Delta > y] &\sim \mathbf{P}\left[\bigvee_{i=1}^n S_i^\Theta > x, \bigvee_{j=1}^n T_j^\Delta > y\right] \sim \mathbf{P}\left[\bigvee_{i=1}^n \Theta_i X_i > x, \bigvee_{j=1}^n \Delta_j Y_j > y\right] \\ &\sim \sum_{i=1}^n \sum_{j=1}^n \mathbf{E}[\Theta_i^{\alpha_1} \Delta_j^{\alpha_2}] \mathbf{P}[X_i > x, Y_j > y], \end{aligned}$$

as  $x \wedge y \rightarrow \infty$ .

**Proof.** The arguments follow directly from application of Lemma 4.2 on Theorem 5.1, since  $\mathcal{R}_{(-\alpha_1, -\alpha_2)}^{(2)} \subsetneq \mathcal{C}^{(2)} \subsetneq (\mathcal{D} \cap \mathcal{L})^{(2)}$ .  $\square$

Now it remains to apply these results to discrete risk model with two-dimensions with random weights. The surplus processes take the form

$$U_1^\Theta(k, x) := x - \sum_{i=1}^k \Theta_i X_i, \quad U_2^\Delta(k, y) := y - \sum_{j=1}^k \Delta_j Y_j,$$

for some  $k = 1, \dots, n$ , with  $x, y$  the initial surpluses in each business lines, the  $X_i, Y_j$  are the claims in the  $i$ -th and  $j$ -th period respectively, and the  $\Theta_i$  and  $\Delta_j$  play the role of discount factor. With these notations we provide the formula for the ruin probability in two-dimensional discrete risk model, over finite time horizon

$$\psi_{and}(x, y, n) := \mathbf{P}[\tau_{and} \leq n] = \mathbf{P}\left[\bigvee_{i=1}^n S_i^\Theta > x, \bigvee_{j=1}^n T_j^\Delta > y\right].$$

**Corollary 5.2.** (1) Let hold the assumptions of Theorem 5.1[(1) or (2)]. Then

$$\psi_{and}(x, y, n) \sim \sum_{i=1}^n \sum_{j=1}^n \mathbf{P}[X_i > x, Y_j > y].$$

as  $x \wedge y \rightarrow \infty$ .

(2) Let hold the assumptions of Corollary 5.1[(1) or (2)]. Then

$$\psi_{and}(x, y, n) \sim \sum_{i=1}^n \sum_{j=1}^n \mathbf{E}[\Theta_i^{\alpha_1} \Delta_j^{\alpha_2}] \mathbf{P}[\Theta_i X_i > x, \Delta_j Y_j > y],$$

as  $x \wedge y \rightarrow \infty$ .

**Proof.** We can just repeat the arguments from Theorem 5.1, Corollary 5.1 and relation (5.19).  $\square$

**Acknowledgments.** We feel the pleasant duty to express deep gratitude to prof. Jinzhu Li for his useful comments that improved significantly the text.

## REFERENCES

- [1] ASMUSSEN, S. (2003) *Applied Probability and Queues* Springer, New York, 2nd ed.
- [2] ASMUSSEN, S., KLÜPPELBERG, C. (1996) Large deviations results for subexponential tails, with applications to insurance risk. *Stoch. Process. Appl.*, **64**, 103–125.
- [3] ATHREYA, K.B., NEY, P.E. (1972) *Branching Processes* Springer, New York.
- [4] BASRAK, B., DAVIS, R.A., MIKOSCH, T. (2002) A characterization of multivariate regular variation. *Ann. Appl. Probab.* **12**, 908–920.

- [5] BASRAK, B., DAVIS, R.A., MIKOSCH, T. (2002) Regular variation of GARCH processes. *Stoch. Process. Appl.* **99**, no. 1, 95–115.
- [6] BREIMAN L. (1965) On some limit theorems similar to arc-sin law. *Theory Probab. Appl.*, **10**, 323–331.
- [7] BURACZEWSKI, D., DAMEK, E., MIKOSCH, T. (2016) *Stochastic Models with Power-Law Tails* Springer, New York.
- [8] CAI, J., TANG, Q. (2004) On max-sum equivalence and convolution closure of heavy-tailed distributions and their applications. *J. Appl. Probab.* **41**, 117–130.
- [9] CHEN, Y., YANG, Y. (2019) Bivariate regular variation among randomly weighted sums in general insurance. *Eur. Actuar. J.* **9**, 301–322.
- [10] CHEN, Y., YUEN, K.C. (2009) Sums of pairwise quasi-asymptotic independent random variables with consistent variation. *Stochastic Models*, **25**, 76–89.
- [11] CHENG, D., YU, C. (2019) Uniform asymptotics for the ruin probabilities in a bidimensional renewal risk model with strongly subexponential claims. *Stochastics* **91**, Vol 1. 643–656.
- [12] CHENG, M., KONSTANTINIDES, D.G., WANG, D (2024) Multivariate regular varying insurance and financial risks in  $d$ -dimensional risk model. *J. Appl. Probab.*, **61**, no. 4.
- [13] CHISTYAKOV, V.P. (1964) A theorem on sums of independent positive random variables and its applications to branching random processes. *Theory Probab. Appl.*, **9**, 640–648.
- [14] CLINE, D.B.H., RESNICK, S. (1992) Multivariate subexponential distributions. *Stoch. Process. Appl.*, **42**, no.1, 49–72.
- [15] CLINE, D.B.H., SAMORODNITSKY, G. (1994) Subexponentiality of the product of independent random variables. *Stoch. Process. Appl.*, **49**, 75–98.
- [16] CUI, Z., WANG, Y. (2020) On the long tail property of product convolution. *Lith. Math. J.*, **60**, no. 2, 315–329.
- [17] DAS, B., FASEN-HARTMANN, V. (2023) Aggregating heavy-tailed random vectors: from finite sums to Levy processes. *arXiv:2301.10423v1*.
- [18] DENISOV, D., ZWART, B. (2007) On a theorem of Breiman and a class of random difference equations. *J. Appl. Probab.* **44**, 1031–1046.
- [19] DIRMA, M., NAKLIUDA, N., ŠIAULYS, J. (2023) Generalized moments of sums with heavy-tailed random summands. *Lith. Math. J.*, **63**, no. 3, 254–271.
- [20] EMBRECHTS, P., KLÜPELBERG, C. AND MIKOSCH, T. (1997) *Modelling Extremal Events for Insurance and Finance*. Springer, New York.
- [21] FELLER, W. (1969) One-sided analogues of Karamata’s regular variation. *L’enseignement Mathématique*, **15**, 107–121.
- [22] Feller, W. (1971) *An Introduction to Probability Theory and Its Applications* Vol. II, second edition, Wiley, New York.
- [23] FOSS, S., KORSHUNOV, D., PALMOWSKI, Z. (2024) Maxima over random time intervals for heavy-tailed compound renewal and Lévy processes. *Stoch. Process. Appl.*, **176**, 104422.
- [24] FOSS, S., KORSHUNOV, D., ZACHARY, S. (2013) *An Introduction to Heavy-Tailed and Subexponential Distributions*. Springer, New York, 2nd ed.
- [25] FOUGERES, A., MERCADIER, C. (2012) Risk measures and multivariate extensions of Breiman’s theorem. *J. Appl. Probab.*, **49**, no. 2, 364–384.
- [26] GELUK, J., TANG, Q. (2009) Asymptotic tail probabilities of sums of dependent subexponential random variables. *J. Theor. Probab.*, **22**, 871–882.
- [27] GOLDIE, C.M. (1978) Subexponential distributions and dominated variation tails *J. Appl. Probab.*, **15**, 440–442.
- [28] HAAN, L. DE, RESNICK, S. (1981) On the observation closet to the origin. *Stoch. Process. Appl.*, **11**, no. 3, 301–308.
- [29] JIANG, T., WANG, Y., CHEN, Y., XU, H. (2015) Uniform asymptotic estimate for finite-time ruin probabilities of a time-dependent bidimensional renewal model. *Insur. Math. Econom.*, **64**, 45–53.
- [30] KIZINEVIČ, E., SPRINDYS, J., ŠIAULYS, J. (2016) Randomly stopped sums with consistently varying distributions. *Modern Stochastics: Theory and Applications*, **3**, 165–179.
- [31] KO, B.W., TANG Q.H. (2008) Sums of dependent nonnegative random variables with subexponential tails. *J. Appl. Probab.*, **45**, 85–94.
- [32] KONSTANTINIDES, D.G. (2018) *Risk Theory. A Heavy Tail Approach*. World Scientific, New Jersey.

- [33] KONSTANTINIDES, D.G., LEIPUS, R., ŠIAULYS, J. (2023) On the non-closure property for strong subexponential distributions. *Nonlinear Analysis: Modelling and Control* **28**, no. 1, 97–115.
- [34] KONSTANTINIDES, D.G., LI, J. (2016) Asymptotic ruin probabilities for a multidimensional renewal risk model with multivariate regularly varying claims. *Insur. Math. and Econom.*, **69**, 38–44.
- [35] KONSTANTINIDES, D.G., MIKOSCH, T. (2005) Large Deviations and Ruin Probabilities for Solutions to Stochastic Recurrence Equations with Heavy-tailed Innovations. *Ann. Probab.*, **33**, 1.992–2.035.
- [36] KONSTANTINIDES, D.G., PASSALIDIS, C.D. (2024) Background risk model in presence of heavy tails under dependence. *Preprint, arXiv:2405.03014*.
- [37] KONSTANTINIDES, D.G., PASSALIDIS, C.D. (2024) Closure properties and heavy tails: random vectors in the presence of dependence *Preprint, arXiv:2402.09041*.
- [38] KONSTANTINIDES, D.G., PASSALIDIS, C.D. (2024) A new approach in two-dimensional heavy-tailed distributions. *Preprint, arXiv:2402.09040*.
- [39] KORSHUNOV, D. (2018) On subexponential tails for the maxima of negatively driven compound renewal and Lévy processes. *Stoch. Process. Appl.*, **128**, 1.316–1.332.
- [40] LEIPUS, R., ŠIAULYS, J. (2020) On a closure property of convolution equivalent class of distributions. *J. Math. Anal. Appl.*, **490**, no. 124226.
- [41] LEIPUS, R., ŠIAULYS, J., KONSTANTINIDES, D.G. (2023) *Closure Properties for Heavy-Tailed and Related Distributions: An Overview*. Springer Nature, Cham.
- [42] LEIPUS, R., SURGAILIS, D. (2007) On long-range dependence in regenerative processes based on a general ON/OFF scheme. *J. Appl. Probab.*, **44**, 379–392.
- [43] LI, J. (2013) On pairwise quasi-asymptotically independent random variables and their applications. *Stat. Prob. Lett.*, **83**, 2081–2087.
- [44] LI, J. (2017) A note on the finite-time ruin probabilities of a renewal risk model with Brownian perturbation. *Stat. Prob. Lett.*, **127**, 49–55.
- [45] LI, J. (2018) A revisit to asymptotic ruin probabilities of a two-dimensional renewal risk model. *Stat. Probab. Lett.*, **140**, 23–32.
- [46] LI, J. (2018b) On the joint tail behavior of randomly weighted sums of heavy-tailed random variables. *J. Multivar. Anal.*, **164**, 40–53.
- [47] LI, J., YANG, H. (2015) Asymptotic ruin probabilities for a bidimensional renewal risk model with constant interest rate and dependent claims. *J. Math. Anal. Appl.*, **426**, no. 1, 247–266.
- [48] MATUSZEWSKA, W. (1964) On generalization of regularly increasing functions. *Studia Mathematica*, **24**, 271–279.
- [49] MAULIK, K., RESNICK, S.I. (2004) Characterizations and examples of hidden regular variation. *Extremes*, **7**, 31–67.
- [50] OMEY, E. (2006) Subexponential distribution functions in  $R^d$ . *J. Math. Sci.*, **138**, no. 1, 5434–5449.
- [51] PAKES, A.G. (2004) Convolution equivalence and infinite divisibility. *J. Appl. Probab.*, **41**, 407–424.
- [52] RESNICK, S. (2007) *Heavy-Tail Phenomena. Probabilistic and Statistical Modeling*. Springer, New York.
- [53] ROLSKI, T., SCHMIDLI, H., SCHMIDT, V., TEUGELS, J.L. (1999) *Stochastic Processes for Insurance and Finance* Wiley, Chichester.
- [54] SAMORODNITSKY, G., SUN, J. (2016) Multivariate subexponential distributions and their applications. *Extremes*, **19**, no. 2, 171–196.
- [55] SAMORODNITSKY, G., TAQQU, M. (1994) *Stable Non-Gaussian Random Processes: Stochastic Models with Infinite Variance* Chapman and Hall, New York.
- [56] SHEN, X., DU, K. (2023) Uniform approximation for the tail behavior of bidimensional randomly weighted sums. *Methodol. Comput. Appl. Probab.*, **25**, no. 26.
- [57] SHEN, X., GE, M., FU, K.A. (2020) Approximation of the tail probabilities for bidimensional randomly weighted sums with dependent components. *Probab. Eng. Inf. Sci.*, **34**, no. 1, 112–130.
- [58] TANG, Q. (2006) The subexponentiality of products revisited. *Extremes*, **9**, 231–241.
- [59] TANG, Q. (2008) Insensitivity to negative dependence of asymptotic tail probabilities of sums and maxima of sums. *Stoch. Anal. Appl.*, **26**, 435–450.
- [60] TANG, Q., TSITSIAHVILI, G. (2003) Randomly weighted sums of subexponential random variables with application to ruin theory. *Extremes*, **6**, 171–188.
- [61] TANG, Q., YUAN, Z. (2014) Randomly weighted sums of subexponential random variables with application to capital allocation. *Extremes*, **17**, 467–493.

- [62] TANG, Q., YANG, Y. (2019) Interplay of insurance and financial risks in a stochastic environment. *Scand. Actuar. J.*, no. 5, 432–451.
- [63] WANG, K. (2011) Randomly weighted sums of dependent subexponential random variables. *Lith. Math. J.*, **51**, no. 4, 573–586.
- [64] WANG, H., SU, Q., YANG, Y. (2024) Asymptotics for ruin probabilities in a bidimensional discrete-time risk model with dependence and consistently varying net losses. *Stochastics*, **96**, no. 1, 667–695.
- [65] YANG, Y., CHEN, S., YUEN, C. (2024) Asymptotics for the joint tail probability of bidimensional randomly weighted sums with applications to insurance. *Sci. China Math.*, **67**, 163–186.
- [66] YANG, Y., LEIPUS, R., ŠIAULYS, J. (2012) Tail probability of randomly weighted sums of subexponential random variables under a dependence structure. *Stat. Probab. Lett.*, **82**, 1727–1736.
- [67] YANG, Y., SU, Q. (2023) Asymptotic behavior of ruin probabilities in a multidimensional risk model with investment and multivariate regularly varying claims. *J. Math. Anal. Appl.*, **525**, 127319.
- [68] YANG, Y., WANG, K., LEIPUS, R., ŠIAULYS, J. (2011) Tail behavior of sums and maxima of sums of dependent subexponential random variables. *Acta Appl. Math.*, **114**, 219–231.
- [69] YANG, Y., YUEN, K.C., LIU, J.-F. (2018) Asymptotics for ruin probabilities in Lévy-driven risk models with heavy-tailed claims. *J. Ind. Manag. Optim.*, **14**, 231–247.
- [70] YUAN, M., LU, D. (2023) Asymptotics for a time-dependent by-claim model with dependent subexponential claims. *Insur. Math. Econom.*, **112**, 120–141.

DEPT. OF STATISTICS AND ACTUARIAL-FINANCIAL MATHEMATICS, UNIVERSITY OF THE AEGEAN, KARLOVASSI, GR-83 200 SAMOS, GREECE

*Email address:* `konstant@aegean.gr`, `sasm23002@sas.aegean.gr`.