arXiv:2408.01097v1 [math.AP] 2 Aug 2024

One dimensional energy cascades in a fractional quasilinear NLS

Alberto Maspero^{*}, Federico Murgante[†]

Abstract

We consider the problem of transfer of energy to high frequencies in a quasilinear Schrödinger equation with sublinear dispersion, on the one dimensional torus. We exhibit initial data undergoing finite but arbitrary large Sobolev norm explosion: their initial norm is arbitrary small in Sobolev spaces of high regularity, but at a later time becomes arbitrary large. We develop a novel mechanism producing instability, which is based on extracting, via paradifferential normal forms, an effective equation driving the dynamics whose leading term is a non-trivial transport operator with non-constant coefficients. We prove that such operator is responsible for energy cascades via a positive commutator estimate inspired by Mourre's commutator theory.

Contents

1	Introduction 1.1 Scheme of the proof	2 4
2	Functional setting 2.1 Paradifferential calculus 2.2 Admissible transformations	6 9 14
3	Analysis of weak resonances	18
4	Paradifferential normal form 4.1 Block diagonalization	24 26 29 30 32
5	The effective equation	33
6	Instability via paradifferential Mourre theory6.1The Mourre operator	39 39 42 44
A	High frequency paradifferential calculus	46
В	Flows and conjugations	49

*International School for Advanced Studies (SISSA), Via Bonomea 265, 34136, Trieste, Italy. *Emails:* amaspero@sissa.it

[†]Università degli studi di Milano, Department of Math. Federigo Enriques, Via Saldini 50, 20133 Milano, Italy *Email:* federico.murgante@unimi.it

1 Introduction

A fundamental question in physics and mathematical analysis is to study how energy is transferred and redistributed from macro to micro scales in deterministic systems, in a process usually called energy cascade, which is central to understand the insurgence of turbulent dynamics. In nonlinear Hamiltonian PDEs, this phenomenon is characterized by the transfer of energy from low to high Fourier modes, initiated and sustained by nonlinear resonant and quasi-resonant interactions between different wave components. As first pointed out by Bourgain [18], an effective way to capture energy transfers is by constructing solutions whose Sobolev norms grow over time. On the one dimensional torus $\mathbb{T} := \mathbb{R}/2\pi\mathbb{Z}$, which is the setup of our paper, the Sobolev norms $H^s := H^s(\mathbb{T}, \mathbb{C})$, $s \in \mathbb{R}$, are defined by

$$||u(t)||_s^2 := \sum_{k \in \mathbb{Z}} \langle k \rangle^{2s} |u_k(t)|^2 , \quad \langle k \rangle := \max(1, |k|)$$

where $u_k(t) := \frac{1}{2\pi} \int_{\mathbb{T}} u(x) e^{-ikx} dx$ are the Fourier coefficients of u(t,x). Whenever s > 0, the H^s norms give greater weight to higher frequencies than to lower ones. Since the L^2 norm of a solution is typically preserved by mass conservation, a growth in the H^s norm over time indicates that energy is cascading towards increasingly higher frequencies. Thus, the challenge becomes constructing solutions of dispersive PDEs that exhibit growth in Sobolev norms. Remark that, in a dispersive wave model, the linearized waves merely oscillate over time and consequently any growth in Sobolev norms is purely a nonlinear mechanism, which makes the analysis particularly challenging.

As we will explain below, there are currently only two rigorous mechanisms known to produce energy cascades in nonlinear Hamiltonian PDEs, each with limited applications. The first one, pioneered by Colliander-Keel-Staffilani-Takaoka-Tao [21], exploits the dynamics of the so-called "toy model" and works for semilinear NLS on \mathbb{T}^d , $d \geq 2$, and some related models [21, 42, 41, 43, 40, 39, 36, 38]. The second one, discovered by Gérard-Grellier [33], leverages the peculiar integrable structure of the Szegő equation.

The purpose of this paper is to introduce a novel paradigm to construct energy cascades. Specifically, we consider a fractional *quasilinear* NLS (Nonlinear Schrödinger) equation, and employ it as a theoretical test-bed to explore our new mechanism. Precisely, we consider

$$\partial_t u = -\mathbf{i}|D|^\alpha u + |u|^2 u_x, \quad x \in \mathbb{T} \ , \quad \alpha \in (0,1) \ , \tag{1.1}$$

with $|D|^{\alpha}$ the Fourier multiplier defined by $|D|^{\alpha}e^{ikx} = |k|^{\alpha}e^{ikx}$, $k \in \mathbb{Z}$. Note that the equation is gauge invariant, so the L^2 -norm is preserved. Note also two important features of equation (1.1), which actually motivate its choice: a nonlinear transport term and a sublinear dispersion relation. These properties are characteristic of equations found in fluid dynamics, such as pure gravity water waves, the surface quasi-geostrophic, the β -plane equations. Moreover, equation (1.1) resembles the Majda–McLaughlin–Tabak model [55, 70] used to check the validity of weak turbulence for random waves. Our main result is the following:

Theorem 1.1. There exists $s_0 > \frac{3}{2}$ such that given any $s > 3s_0$, $0 < \delta \le 1$ and $K \ge 1$, there exists a solution $u(t) \in H^s(\mathbb{T}, \mathbb{C})$ of (1.1) and a time T > 0 such that

$$||u(0)||_s \le \delta \quad and \quad ||u(T)||_s \ge K .$$

Moreover $\sup_{0 \le t \le T} \|u(t)\|_{s_0} \le 2\delta.$

Theorem 1.1 guarantees the existence of a solution of (1.1) with smooth and arbitrary small initial data undergoing finite but arbitrary large Sobolev norm explosion. Such solution has constant L^2 -norm and stays small in the "low" H^{s_0} -norm. We do not know the fate of such solution after time T, and given that global existence for (1.1) is not established, we cannot exclude the possibility of finite-time blow-up. This phenomenon occurs in similar models such as the fractional KdV equation, where solutions with large initial data can develop shocks [22, 48, 46, 47, 68, 62, 49], resulting in the H^1 norm explodes while the L^{∞} stays bounded. However, these shock solutions appear distinct from those described in our Theorem1.1, for which we ensure that low Sobolev norms stay small.

On the other end, not every initial data gives rise to turbulent solutions of (1.1): consider for example the plane waves $ae^{i(kx-\omega t)}$ with $\omega = |k|^{\alpha} - a^2k$, which can be made of arbitrary small size. We also expect that KAM methods, like those developed in [6, 12, 28], would enable the construction of globally defined, small-amplitude, time quasi-periodic solutions, demonstrating the coexistence of stable and unstable dynamics.

As mentioned earlier, the primary novelty of this paper is to introduce a new mechanism producing energy cascades. At the core of this new mechanism is the nature of the quasi-resonant effective system that drives the unstable dynamics of (1.1). Such effective system, that we extract from (1.1) via quasi-linear normal forms, is the linear PDE with dominant *non-constant coefficient tranport operator*

$$\partial_t \zeta = -\mathbf{i} |D|^{\alpha} \zeta + \mathbf{i} \mathrm{Op}^{BW}((\mathbf{J}_1 + \mathfrak{v}(x))\xi) \zeta .$$
(1.2)

Here J_1 is a real number and v(x) a real valued function, both depending nonlinearly on the initial data u(0) (see (5.25) and (5.26)), and $Op^{BW}(\cdot)$ is a Bony-Weyl paradifferential operator (see (2.21)) of order one, coming from the nonlinearity of (1.1). The effective system (1.2) is very different from the one of the NLS of [21, 42, 41, 43, 40, 39, 36, 38], which consists in a large nonlinear ODE system –the so called toy-model– carrying orbits connecting very distant modes.

To show that (1.2) has solutions with growing Sobolev norms, we develop a positive commutator method, inspired by Mourre's commutator theory [61]. Precisely, we construct a paradifferential operator A, see (6.5), such that the commutator

$$i[\mathsf{A}, \operatorname{Op}^{BW}((\mathsf{J}_1 + \mathfrak{v}(x))\xi)]$$

is strictly positive on large frequencies up to a small remainder. This is possible provided the function $J_1 + v(x)$ does not have sign, a condition that we are able to impose by tuning the initial datum. We remark that positive commutator estimates has been exploited in [57, 58] to prove growth of Sobolev norms in linear time-dependent Schrödinger equations on \mathbb{R} , where the full strength of Mourre's theory [61] and local-energy decay estimates [65] are used; see also [19, 20] for zero-order pseudodifferential operators.

A fundamental contribution of this paper is to successfully extended positive commutator methods to a nonlinear setup. A further benefit of this approach is that it allow us to prove that the solution of Theorem (1.1) grows at an exponentially fast rate. This is due to the quasilinear nature of equation (1.1): for semilinear NLS, polynomial upper bounds in time are known (see e.g. [15, 67, 66, 63]), which become subpolynomial in time in the linear time-dependent case (see e.g. [17, 23, 59, 2, 5, 4]).

A second point of interest is that the paradifferential operator $\operatorname{Op}^{BW}((J_1 + \mathfrak{v}(x))\xi)\zeta$ is not Birkhoff-resonant, but only quasi-resonant. In principle it might be eliminated by a (formal) Birkhoff normal form procedure, but the required transformation is unbounded and not well defined in H^s , due to the quasi-linear nature of the problem. Consequently, in equation (1.1) energy cascades are due to quasi-resonant interactions rather than exact resonances; this is reminiscent, in wave turbulence, to the fact that are quasi-resonances (rather than resonances) to play a fundamental role in the rigorous derivation of the wave kinetic equation [25].

A final point of interest, which has been a main motivation for the choice of (1.1), is that important equations coming from fluids have a nonlinear transport term and a sublinear dispersion relation, such as the pure-gravity water waves equations which, in an appropriate complex variable u, read (see e.g. [11])

$$\partial_t u = \operatorname{Op}^{BW}(\mathrm{i}(1+\zeta(u;x))\omega(\xi) + V(u;x)\mathrm{i}\xi) u + R(u), \quad \omega(\xi) \sim |\xi|^{\frac{1}{2}}$$

with V(u; x) a real function and R(u) a smoothing remainder. This form is also the one that we obtain from (1.1) after paralinearizing (of course with different symbols and smoothing remainders) and in case $\alpha = \frac{1}{2}$, see Lemma 4.3.

Related literature: Whereas for linear time dependent equations several results are known [16, 24, 56, 1, 52, 54, 53, 3, 27, 45], for nonlinear systems the results are scarce and limited to essentially

two models: the semilinear Schrödinger equation (NLS) and the Szegő equation. Regarding the first, after the seminal works by Kuksin [50, 51], the breakthrough result by Colliander-Keel-Staffilani-Takaoka-Tao [21] for the NLS on \mathbb{T}^d , $d \geq 2$, identified the first mechanism of growth, based on the toy-model construction. Such mechanism was further exploited by Guardia-Kaloshin [41], Haus-Procesi [43], Guardia-Haus-Procesi [40], Guardia-Giuliani [36] and Giuliani [38]. All these results construct solutions starting with norm arbitrally small and becoming arbitrarily large at a later time. We also mention Hani [42] and Guardia-Haus-Hani-Maspero-Procesi [39] that construct solutions undergoing Sobolev norm inflation and starting arbitrary close to periodic or quasi-periodic orbits. Solutions with unbounded paths have been constructed by Hani-Pausader-Tzvetkov-Visciglia [44] for the NLS on $\mathbb{R} \times \mathbb{T}^2$, combining dispersive effects and the resonant toy-model construction.

The second known mechanism ensuring growth of Sobolev norms was pioneered by Gérard-Grellier [33] for the Szegő equation, exploiting its peculiar integrable structure [32]. We also mention Biasi-Evnin [7] for a truncated Szegő systems, Gérard-Lenzmann [35] for the integrable Calogero-Moser derivative NLS, and Gérard-Grellier [34] for a long time instability result for the cubic half-wave equation (exploiting a resonant approximation with the Szegő equation).

Finally we mention Guardia-Giuliani [37] for chains of infinite pendula, the recent numerical result by Gallone-Marian-Ponno-Ruffo [31] for the FPUT chain and Elgindi- Shikh Khalil [26] for a completely different norm inflation mechanism in L^{∞} .

1.1 Scheme of the proof

First note that, via energy methods, the local Cauchy theory ensures that initial data in H^s , $s > \frac{3}{2}$, give rise to local in time solutions. Additionally, if the initial size is $\delta \ll 1$, the solution remains of comparable size for times of order δ^{-2} , giving a lower bound to the time of growth. To see energy transfers, we need to observe the dynamics for longer time scales and this is a crucial difficulty. We shall now describe in more details the main ideas and methods of the proof.

Step 1: paradifferential normal form. The first step in our analysis is to transform equation (1.1) via a suitable quasi-linear normal form. We employ the method pioneered by Berti and Delort [9], further developed and extended in [30, 11, 29, 10, 13, 8, 60]. While previous applications of the Berti-Delort method aimed primarily at constructing a modified energy to establish *upper bounds* on the Sobolev norms of solutions, our approach leverages the method to extract an effective equation that have unstable solutions. Indeed, the Berti-Delort method provides also a normal form equation that governs the dynamics over long times.

In Section 4, we perform two paradifferential transformations to conjugate the original equation (1.1) to the normal form system (4.22), whose cubic component has the form

$$\partial_t w = -\mathbf{i}|D|^{\alpha} w + \mathrm{Op}^{BW} \left(\mathbf{i} \langle \underline{\mathbf{V}} \rangle (u(t); x) \xi + \mathbf{i} a_2^{(\alpha)}(u(t); x, \xi) \right) w + R_2(u(t)) w + h.o.t.$$
(1.3)

where $\langle \underline{\mathbf{V}} \rangle(u(t); x) := 2 \operatorname{Re} \left(\sum_{n \in \mathbb{N}} u_n(t) \overline{u_{-n}(t)} e^{i2nx} \right)$, $a_2^{(\alpha)}$ a symbol of order α and quadratic in u(t), and $R_2(u(t))$ a smoothing operator again quadratic in u. This normal form is significantly different from the one of [9] and of [30, 11, 10, 13, 60], where the symbol of the paradifferential operator has *constant* coefficients (at least at low homogeneity).

On the contrary, in (1.3), $\langle \underline{\mathbf{V}} \rangle(u;x)$ has non-constant coefficients, and additionally it depends on time through u(t). This is the term who will give rise to the paradifferential operator in (1.2). To do so, we need to remove (or at least simplify) such time dependence. The first natural attempt, i.e. replace in $\langle \underline{\mathbf{V}} \rangle(u(t);x)$ the function u(t) with its linear evolution $e^{-i|D|^{\alpha}t}u(0)$, fails because it produces an error that we cannot bound on the long time scales needed to see growth. Therefore, we need to study the nonlinear dynamics of at least two modes $u_n(t)$, $u_{-n}(t)$. So we fix the modes in $\Lambda := \{-1, 1\}$ and study the nonlinear dynamics of $u_1(t)$, $u_{-1}(t)$.

Step 2: the Λ -normal form. We decompose the solution as follows:

$$u(t) = u^{\top}(t) + u^{\perp}(t) \quad \text{where} \\ u^{\top}(t) := u_1(t)e^{ix} + u_{-1}(t)e^{-ix} \quad u^{\perp}(t) := \sum_{k \neq \pm 1} u_k(t)e^{ikx}.$$

This decomposition separates the *tangential modes* $u^{\top}(t)$ from the *normal modes* $u^{\perp}(t)$. To decouple the dynamics of these modes, we use a weak-normal form. The paradifferential operator in equation (1.3) vanishes when restricted to Λ (see (5.9)). Therefore, the dynamics of $u^{\top}(t)$ is governed by the smoothing operator $R_2(u)w$.

We further decouple the dynamics of the tangential and normal modes in $R_2(u)w$ by removing from this term two types of monomials $u_{j_1}^{\sigma_1} u_{j_2}^{\sigma_2} u_{j_3}^{\sigma_3} e^{ikx}$:

- (i) Monomials with $(j_1, j_2, j_3) \in \Lambda$ and $k \in \Lambda^c$:
 - This ensures that the set Λ remains invariant under the cubic part dynamics of (1.3).
 - It requires first-order Melnikov conditions:

$$|j_1|^{\alpha} - |j_2|^{\alpha} + |j_3|^{\alpha} - |k|^{\alpha} \neq 0$$
, $j_1 - j_2 + j_3 - k = 0$,

that actually we verify whenever one and only one among (j_1, j_2, j_3, k) is in Λ .

(ii) Monomials with exactly two indexes among (j_1, j_2, j_3) in Λ and the remaining one and k in Λ^c :

• This is needed so that the leading term in equation (1.3) is given by the skewadjoint paradifferential term $\operatorname{Op}^{BW}(i\langle \underline{\underline{V}}\rangle(u_1\overline{u_{-1}};x)\xi)w$ (whose monomials have exactly 2 indexes inside Λ and 2 outside).

• It requires second-order Melnikov conditions:

$$|j_1|^{\alpha} - |j_2|^{\alpha} + |j_3|^{\alpha} - |k|^{\alpha} \neq 0$$
, $j_1 - j_2 + j_3 - k = 0$,

when two indexes among (j_1, j_2, j_3, k) are in Λ and the other two in Λ^c , provided $j_1 \neq j_2$ or $j_1 \neq k$.

As a result, only integrable monomials of the form $|u_{j_1}|^2 u_{j_3} e^{ij_3x}$, with either $j_1, j_3 \in \Lambda$ or $j_1 \in \Lambda$, $j_3 \in \Lambda^c$ or viceversa are left in the smoothing operator $R_2(u)w$. Finally, in Proposition 4.10, we identify the remaining resonant integrable monomials via an a-posteriori identification argument à la Berti-Feola-Pusateri [11] (see also [10]), obtaining the explicit form 4.9.

Step 3: The effective equation. In Section 5 we extract the effective equation. Starting from the quasi-linear normal form in Section 4, the variables $z^{\top}(t)$ and $z^{\perp}(t)$ solve system (5.3)–(5.4), which has roughly the form

$$\begin{cases} \partial_t z^{\top} = -\mathbf{i} |D|^{\alpha} z^{\top} + Y_3^{(\Lambda)}(z^{\top}(t)) + \mathcal{O}(||z^{\perp}||_{s_0}^3, ||z||_{s_0}^5) \\ \partial_t z^{\perp} = \mathbf{i} |D|^{\alpha} z^{\perp} + \mathrm{Op}^{BW} \left(\mathbf{i} \langle \underline{\mathbf{V}} \rangle(z^{\top}(t); x) \xi + \mathbf{i} a_2^{(\alpha)}(z(t); x, \xi) \right) z^{\perp} + \mathcal{O}(||z^{\top}||_{s_0} ||z^{\perp}||_{s_0} ||z^{\perp}||_{s_0}) \end{cases}$$
(1.4)

where $Y_3^{(\Lambda)}(z^{\top})$ is the explicit *integrable* vector field (5.5), and the symbol of the transport operator in the equation for z^{\perp} is evaluated only on the tangential modes $z^{\top}(t)$.

To further understand the dynamics of system (1.4) and to extract from it the effective equation (1.2), we introduce a small parameter $\epsilon \ll 1$ and we consider special solutions of system (1.4), that we call *long-time controlled* (see Definition 5.2). They are characterized by two properties:

(i) Their initial data are small in L^2 , with most mass on the modes $z_1(0), z_{-1}(0)$:

$$||z^{\top}(0,\cdot)||_{L^2} \le \epsilon, \quad ||z^{\perp}(0,\cdot)||_{L^2} \le \epsilon^3;$$

(ii) Their high H^s -norms have large a-priori bounds:

$$||z(t)||_s \le \epsilon^{-\theta}$$
 with $0 < \theta \ll 1$.

The large a-priori bound above is not restrictive for our problem; if it fails, it means the solution has already grown.

We then prove that any long-time controlled solution, for times $|t| \leq \epsilon^{-2} \log(\epsilon^{-1})$, has:

• The modes $z_1(t)$ and $z_{-1}(t)$ evolving very close to the rotations:

$$z_{\pm 1}(t) = e^{-it(1\pm|z_{\pm 1}(0)|^2)} z_{\pm 1}(0) + \mathcal{O}(\epsilon^{3-\theta})$$

• The low norm of $z^{\perp}(t)$ staying very small, i.e. $||z^{\perp}(t)||_{s_0} \leq \epsilon^2$. One key idea to obtain this is to estimate $z^{\perp}(t)$ in L^2 , exploiting the cancellation coming from the skewadjointness of the paradifferential operator, then deducing a bound for $||z^{\perp}(t)||_{s_0}$ by interpolation with the large a-priori bound for $||z(t)||_s$.

Finally, we approximate the evolution of $z^{\top}(t)$ with the rotations $e^{-it(1\pm|z_{\pm 1}(0)|^2)}z_{\pm 1}(0)$ in the symbol $\langle \underline{V} \rangle(z^{\top}(t); x)$ obtaining a negligible remainder, and, after a space translation, we arrive to an effective system of the form (1.2), see Proposition 5.4.

Step 4: Growth of Sobolev norms. After this analysis, we have essentially reduced the problem to construct solutions of the effective equation (1.2) with growing Sobolev norms. Inspired by Mourre's commutator theory, we construct a paradifferential operator A, of order 2s and supported on high-frequencies, see (6.5), fulfilling the positive commutator estimate (Lemma 6.2)

$$\mathbf{i}[\mathsf{A}, \operatorname{Op}^{^{BW}}((\mathsf{J}_1 + \mathfrak{v}(x))\xi)] \ge \mathbf{I}_1 \operatorname{Op}^{^{BW}}\left(|\xi|^{2s}\eta_{\mathsf{R}}^2(\xi)\right) + h.o.t.$$
(1.5)

Here I_1 is a strictly positive real number depending on the initial data, see (6.9), and η_R a cut-off function on high frequencies. To obtain such positive commutator estimate, the main ingredient is to find a symbol $\mathfrak{a}(x,\xi)$ which is an escape-function for the dynamics of $(J_1 + \mathfrak{v}(x))\xi$, namely such that the Poisson bracket $\{\mathfrak{a}(x,\xi), (J_1 + \mathfrak{v}(x))\xi\}$ is strictly positive. This is possible provided the function $J_1 + \mathfrak{v}(x)$ does not have sign; since

$$\mathbf{J}_1 + \mathbf{v}(x) = \frac{|z_1(0)|^2 + |z_{-1}(0)|^2}{2} + 2\operatorname{Re}(z_1(0)\overline{z_{-1}(0)}e^{\mathbf{i}2x}),$$

it is enough to select the values of the initial modes $z_{\pm 1}(0)$ so that $\frac{|z_1(0)|^2 + |z_{-1}(0)|^2}{2} < 2|z_1(0)| |z_{-1}(0)|$. The same condition yields the strict positivity of the number I_1 in (1.5). An important point is that the operator A is chosen to be supported on very large $|\xi| \ge R \ge e^{-\frac{3+\theta}{1-\alpha}}$. This is required so that the dispersive term $-i|D|^{\alpha}$ and all the other lower order operators becomes perturbative with respect to the leading transport. To conclude, we define the functional $\mathcal{A}(t) := \langle Az^{\perp}, z^{\perp} \rangle$ and show that (1.5) leads to a lower bound for the dynamics of $\frac{d}{dt}\mathcal{A}(t)$, forcing $\mathcal{A}(t)$ to grow exponentially fast provided $\mathcal{A}(0)$ is not too small, a condition that can be imposed by well-preparing the initial data. Being $\mathcal{A}(t) \lesssim ||z^{\perp}(t)||_s^2$, growth of Sobolev norms follows.

Acknowledgments. We thank R. Grande for useful discussions. A. Maspero is supported by the European Union ERC CONSOLIDATOR GRANT 2023 GUnDHam, Project Number: 101124921 and by PRIN 2022 (2022HSSYPN) "TESEO - Turbulent Effects vs Stability in Equations from Oceanography". F. Murgante is supported by the ERC STARTING GRANT 2021 HamDyWWa, Project Number: 101039762. Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

2 Functional setting

In this section we introduce the paradifferential operators and smoothing remainders, following [9, 13]. We also introduce a new class of transformations, that we call *admissible transformations*, see Definition 2.10. They are maps $U \mapsto \mathbf{F}(U)$ whose main property is to be of regularity C^1 with respect to the internal variable. Consequently, the nonlinear map $U \mapsto \mathbf{F}(U)U$ results invertible. We shall prove that all the transformation generated along the normal form reduction of Section 4 are admissible.

Function spaces. Along the paper we deal with real parameters $s \ge s_0 \gg \rho$.

2 FUNCTIONAL SETTING

For $s \in \mathbb{R}$ we shall denote with $H^s(\mathbb{T}; \mathbb{C}^2)$ the space of couples of complex valued Sobolev functions in $H^s(\mathbb{T}, \mathbb{C})$ and with

$$H^s_{\mathbb{R}}(\mathbb{T};\mathbb{C}^2) := \left\{ U = \begin{pmatrix} u^+ \\ u^- \end{pmatrix} \in H^s(\mathbb{T};\mathbb{C}^2) \colon u^- = \overline{u^+} \right\} \,.$$

Given r > 0 we set $B_s(r)$ the ball or radius r in $H^s(\mathbb{T}, \mathbb{C}^2)$ and $B_{s,\mathbb{R}}(r)$ the ball or radius r in $H^s_{\mathbb{R}}(\mathbb{T}, \mathbb{C}^2)$. Given an interval $I \subset \mathbb{R}$ symmetric with respect to t = 0 and a Banach space X, we use the standard notation C(I, X) to denote the space of continuous functions with values in X. Given r > 0 we set $B_s(I; r)$ the ball of radius r in $C(I, H^s(\mathbb{T}, \mathbb{C}^2))$ and by $B_{s,\mathbb{R}}(I; r)$ the ball of radius r in $C(I, H^s_{\mathbb{R}}(\mathbb{T}, \mathbb{C}^2))$. We denote $L^2(\mathbb{T}, \mathbb{C}) := H^0(\mathbb{T}, \mathbb{C})$ and we define

$$\langle u, v \rangle_{L^2} := \frac{1}{2\pi} \int_{\mathbb{T}} u(x) \overline{v(x)} \,\mathrm{d}x \,. \tag{2.1}$$

Given $N \in \mathbb{N}_0$, we denote by $W^{N,\infty}(\mathbb{T})$ the space of continuous functions $u : \mathbb{T} \to \mathbb{C}$, 2π -periodic, whose derivatives up to order N are in L^{∞} , equipped with the norm

$$\|u\|_{W^{N,\infty}}:=\sum_{\ell=0}^N \|\partial_x^\ell u\|_{L^\infty}$$

For N = 0 the norm $\| \|_{W^{N,\infty}} = \| \|_{L^{\infty}}$.

We denote by $\tau_{\varsigma}, \varsigma \in \mathbb{R}$, and by $g_{\theta}, \theta \in \mathbb{T}$, the translation operator respectively the phase rotation given by

$$[\tau_{\varsigma}u](x) := u(x+\varsigma) , \qquad [\mathsf{g}_{\theta}\left(\frac{u}{u}\right)](x) := \begin{pmatrix} e^{\mathrm{i}\theta}u(x)\\ e^{-\mathrm{i}\theta}\overline{u}(x) \end{pmatrix} . \tag{2.2}$$

Symmetries of operators and vector fields. Given a linear operator A(U) acting on $L^2(\mathbb{T};\mathbb{C})$ we associate the linear operator defined by the relation

$$\overline{A}(U)[v] := \overline{A(U)[v]}, \quad \forall v : \mathbb{T} \to \mathbb{C}.$$

An operator A is *real* if $A = \overline{A}$. We say that a matrix of operators acting on $L^2(\mathbb{T}; \mathbb{C}^2)$ is *real-to-real*, if it has the form

$$R(U) = \begin{pmatrix} R_1(U) & R_2(U) \\ \overline{R_2(U)} & \overline{R_1(U)} \end{pmatrix}, \quad \forall U \in L^2_{\mathbb{R}}(\mathbb{T}, \mathbb{C}^2) .$$

$$(2.3)$$

A real-to-real matrix of operators R(U) acts in the subspace $L^2_{\mathbb{R}}(\mathbb{T}, \mathbb{C}^2)$. If R(U) and R'(U) are real-to-real operators then also $R(U) \circ R'(U)$ is real-to-real.

A matrix R(U) as in (2.3) is translation resp. gauge invariant if

$$\tau_{\varsigma} \circ R(U) = R(\tau_{\varsigma}U) \circ \tau_{\varsigma} , \ \forall \varsigma \in \mathbb{R} \quad \text{resp.} \quad \mathbf{g}_{\theta} \circ R(U) = R(\mathbf{g}_{\theta}U) \circ \mathbf{g}_{\theta} , \ \forall \theta \in \mathbb{T} .$$
(2.4)

Similarly we will say that a vector field

$$X(U) := \begin{pmatrix} X(U)^+ \\ X(U)^- \end{pmatrix} \text{ is real-to-real if } \overline{X(U)^+} = X(U)^-, \quad \forall U \in L^2_{\mathbb{R}}(\mathbb{T}, \mathbb{C}^2),$$
(2.5)

and translation resp. gauge invariant if

$$\tau_{\varsigma} \circ X = X \circ \tau_{\varsigma} , \quad \forall \varsigma \in \mathbb{R} , \qquad \mathbf{g}_{\theta} \circ X = X \circ \mathbf{g}_{\theta}, \quad \forall \theta \in \mathbb{T} .$$
 (2.6)

If R(U) in (2.3) is translation resp. gauge invariant, then the vector field X(U) := R(U)U is translation resp. gauge invariant as well.

Fourier expansion. Given a 2π -periodic function u(x) in $L^2(\mathbb{T}, \mathbb{C})$, we expand it in Fourier series as

$$u(x) = \sum_{j \in \mathbb{Z}} u_j e^{ijx}, \quad u_j := \frac{1}{2\pi} \int_{\mathbb{T}} u(x) e^{-ijx} \, \mathrm{d}x \,.$$
(2.7)

2 FUNCTIONAL SETTING

We shall expand a function $U \in L^2(\mathbb{T}; \mathbb{C}^2)$ as

$$U = \begin{pmatrix} u^+ \\ u^- \end{pmatrix} = \sum_{\sigma \in \pm} \sum_{j \in \mathbb{Z}} q^{\sigma} u_j^{\sigma} e^{i\sigma jx}, \quad u_j^{\sigma} := \frac{1}{2\pi} \int_{\mathbb{T}} u^{\sigma}(x) e^{-i\sigma jx} dx$$

where $q^+ := \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $q^- := \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

For
$$\vec{j} = (j_1, \dots, j_p) \in \mathbb{Z}^p$$
, $p \ge 1$, and $\vec{\sigma} = (\sigma_1, \dots, \sigma_p) \in \{\pm\}^p$ we denote $|\vec{j}| := \max(|j_1|, \dots, |j_p|)$ and

$$u_{\vec{j}}^{\vec{\sigma}} := u_{j_1}^{\sigma_1} \cdots u_{j_p}^{\sigma_p}, \qquad \vec{\sigma} \cdot \vec{j} := \sigma_1 j_1 + \cdots + \sigma_p j_p, \quad \vec{\sigma} \cdot \vec{1} := \sigma_1 + \cdots + \sigma_p$$

We also denote by \mathcal{P}_p the set of indexes

$$\mathcal{P}_p := \left\{ (\vec{j}, \vec{\sigma}) \in \mathbb{Z}^p \times \{\pm\}^p : \quad \vec{j} \cdot \vec{\sigma} = 0 \ , \quad \vec{\sigma} \cdot \vec{1} = 0 \right\} \ .$$

$$(2.8)$$

Fourier representation of homogeneous operators and vector fields. In the sequel we shall encounter matrices of linear operators, gauge and translational invariant, of the form

$$M(U) = \begin{pmatrix} M_{+}^{+}(U) & M_{+}^{-}(U) \\ M_{-}^{+}(U) & M_{-}^{-}(U) \end{pmatrix},$$
(2.9)

depending on U in a homogeneous way. We shall call them p-homogeneous if they are polynomials in U of order p. We write them in Fourier as

$$M(U)V = \begin{pmatrix} (M(U)V)^+ \\ (M(U)V)^- \end{pmatrix}, \quad (M(U)V)^{\sigma} = \sum_{\substack{\sigma k = \vec{\sigma}_p, \vec{j}_p + \sigma'j \\ \sigma = \vec{\sigma}_p, \vec{1} + \sigma'}} M_{\vec{j}_p, j, k}^{\vec{\sigma}_p, \sigma', \sigma} u_{\vec{j}_p}^{\vec{\sigma}_p} v_j^{\sigma'} e^{\mathbf{i}\sigma kx} ,$$

where the coefficients $M_{j_p,j,k}^{\vec{\sigma}_p,\sigma',\sigma} \in \mathbb{C}$ fulfill the following symmetric property: for any permutation π of $\{1,\ldots,p\}$, it results

$$M_{j_{\pi(1)},\dots,j_{\pi(p)},j,k}^{\sigma_{\pi(1)},\dots,\sigma_{\pi(p)},\sigma',\sigma} = M_{j_{1},\dots,j_{p},j,k}^{\sigma_{1},\dots,\sigma_{p},\sigma',\sigma}.$$
(2.10)

The operator M(U) is real-to-real, according to definition (2.3), if and only if its coefficients fulfill

$$\overline{M_{\vec{j}_{p},j,k}^{\vec{\sigma}_{p},\sigma',\sigma}} = M_{\vec{j}_{p},j,k}^{-\vec{\sigma}_{p},-\sigma',-\sigma} .$$
(2.11)

A (p + 1)-homogeneous vector field, which is gauge and translation invariant (see (2.6)), can be expressed in Fourier as: for any $\sigma = \pm$,

$$X(U)^{\sigma} = \sum_{k \in \mathbb{Z}} X(U)_{k}^{\sigma} e^{i\sigma kx}, \qquad X(U)_{k}^{\sigma} = \sum_{\substack{k\sigma = \vec{\sigma}_{p+1} \cdot \vec{j}_{p+1}, \\ \sigma = \vec{\sigma}_{p+1} \cdot \vec{1}}} X_{\vec{j}_{p+1},k}^{\vec{\sigma}_{p+1},\sigma} u_{\vec{j}_{p+1}}^{\vec{\sigma}_{p+1}}, \qquad (2.12)$$

the last sum being in $(\vec{j}_{p+1}, \vec{\sigma}_{p+1})$, and with coefficients $X_{\vec{j}_{p+1},k}^{\vec{\sigma}_{p+1},\sigma} \in \mathbb{C}$ satisfying the symmetry condition: for any permutation π of $\{1, \ldots, p+1\}$,

$$X_{j_{\pi(1)},\dots,j_{\pi(p+1)},k}^{\sigma_{\pi(1)},\dots,\sigma_{\pi(p+1)},\sigma} = X_{j_{1},\dots,j_{p+1},k}^{\sigma_{1},\dots,\sigma_{p+1},\sigma}.$$

The constraint of the indexes in (2.12) can also be written as $(\vec{j}_{p+1}, k, \vec{\sigma}_{p+1}, -\sigma) \in \mathcal{P}_{p+2}$ (recall (2.8)), and we shall often use this notation.

If X(U) is real-to-real, see (2.5), then

$$\overline{X(U)_k^+} = X(U)_k^-$$
 i.e. $\overline{X_{\vec{j}_{p+1},k}^{\vec{\sigma}_{p+1},+}} = X_{\vec{j}_{p+1},k}^{-\vec{\sigma}_{p+1},-}$.

2.1 Paradifferential calculus

In this section we introduce paradifferential and smoothing operators, following [9, 13].

Symbols. We define the class of symbols which we will use along the paper. They correspond to the autonomous symbols of Definition 3.3 in [9], where the dependence on time enters only through the function U = U(t). In view of this, we do not need to keep track on the regularity indexes in time and we fix K = K' = 0 with respect to Definition 3.3 of [9].

Definition 2.1 (Symbols). Let $m \in \mathbb{R}$, $N \in \mathbb{N}_0$, $p \in \mathbb{N}$, $s_0, r > 0$.

1. Hölder symbols. We denote by $\Gamma_{W^{N,\infty}}^m$ the space of functions $a : \mathbb{T} \times \mathbb{R} \to \mathbb{C}$, $a(x,\xi)$, which are C^{∞} with respect to ξ and such that, for any $\beta \in \mathbb{N}_0$, there exists a constant $C_{\beta} > 0$ such that

$$\left\|\partial_{\xi}^{\beta} a(\cdot,\xi)\right\|_{W^{N,\infty}} \le C_{\beta} \left<\xi\right>^{m-|\beta|}, \quad \forall \xi \in \mathbb{R}$$

We endow $\Gamma_{W^{N,\infty}}^m$ with the family of norms defined, for any $n \in \mathbb{N}_0$, by

$$|a|_{m,W^{N,\infty},n} := \max_{\beta \in \{0,\dots,n\}} \sup_{\xi \in \mathbb{R}} \left\| \langle \xi \rangle^{-m+|\beta|} \, \partial_{\xi}^{\beta} a(\cdot,\xi) \right\|_{W^{N,\infty}} \,. \tag{2.13}$$

2. p-Homogeneous symbols. We denote by $\widetilde{\Gamma}_p^m$ the space of p-linear maps from $(C^{\infty}(\mathbb{T};\mathbb{C}^2))^p$ to the space of C^{∞} functions from $\mathbb{T} \times \mathbb{R}$ to \mathbb{C} , $(x,\xi) \mapsto a_p(U;x,\xi)$ of the form

$$a_p(U; x, \xi) = \sum_{\substack{\vec{j} \in \mathbb{Z}^p \\ \vec{\sigma} \in \{\pm\}^p}} a_{\vec{j}}^{\vec{\sigma}}(\xi) u_{\vec{j}}^{\vec{\sigma}} e^{i(\vec{\sigma} \cdot \vec{j})x}, \qquad (2.14)$$

where $a_{\vec{i}}^{\vec{\sigma}}(\xi)$ are complex valued Fourier multipliers satisfying, for some $\mu \geq 0$,

$$|\partial_{\xi}^{\beta} a_{\vec{j}}^{\vec{\sigma}}(\xi)| \le C_{\beta} |\vec{j}|^{\mu} \langle \xi \rangle^{m-\beta}, \quad \forall \vec{j} \in \mathbb{Z}^{p}, \, \vec{\sigma} \in \{\pm\}^{p}, \, \beta \in \mathbb{N}_{0}.$$

$$(2.15)$$

We denote by $\widetilde{\Gamma}_0^m$ the space of constant coefficients symbols $\xi \mapsto a(\xi)$ which satisfy (2.15) with $\mu = 0$.

3. Non-homogeneous symbols. We denote by $\Gamma_{\geq p}^{m}[r]$ the space of functions $(U; x, \xi) \mapsto a(U; x, \xi)$, defined for $U \in B_{s_0}(r)$ for some s_0 large enough, with complex values, such that for any $\mathbf{s} \geq s_0$, there are C > 0, $r' := r'(\mathbf{s}) \in (0, r)$ and for any $U \in B_{s_0}(r') \cap H^{\mathbf{s}}(\mathbb{T}; \mathbb{C}^2)$, any $\beta \in \mathbb{N}_0$ and $N \leq \mathbf{s} - s_0$, one has the estimate

$$\left\|\partial_{\xi}^{\beta}a\left(U;\cdot,\xi\right)\right\|_{W^{N,\infty}} \le C\langle\xi\rangle^{m-\beta} \|U\|_{s_{0}}^{p-1}\|U\|_{s}.$$
(2.16)

In addition we require also the translation invariance property

$$a\left(\tau_{\varsigma}U; x, \xi\right) = a\left(U; x+\varsigma, \xi\right), \quad \forall \varsigma \in \mathbb{R},$$

$$(2.17)$$

where τ_{ς} is the translation operator in (2.2).

4. Symbols. We denote by $\Sigma\Gamma_0^m[r]$ the class of symbols of the form

$$a(U; x, \xi) = a_0(\xi) + a_2(U; x, \xi) + a_{\geq 4}(U; x, \xi)$$
(2.18)

where $a_0(\xi) \in \widetilde{\Gamma}_0^m$ is a Fourier multiplier, $a_2(U) \in \widetilde{\Gamma}_2^m$ and $a_{\geq 4}(U) \in \Gamma_{\geq 4}^m[r]$. We denote by $\Sigma \Gamma_2^m[r]$ the class of symbols of the form (2.18) with $a_0(\xi) = 0$. Finally sometimes we shall write $\Sigma \Gamma_4^m[r] \equiv \Gamma_{\geq 4}^m[r]$.

We say that a symbol $a(U; x, \xi)$ is real if it is real valued for any $U \in B_{s_0,\mathbb{R}}(I; r)$.

We also denote by $\widetilde{\mathcal{F}}_p$ (respectively $\mathcal{F}_{\geq p}[r]$) the subspace of $\widetilde{\Gamma}_p^0$ (respectively $\Gamma_{\geq p}^0[r]$) made of those symbols which are independent of ξ , and by $\widetilde{\mathcal{F}}_p^{\mathbb{R}}$ (respectively $\mathcal{F}_{\geq p}^{\mathbb{R}}[r]$) to denote functions in $\widetilde{\mathcal{F}}_p$ (respectively $\mathcal{F}_{\geq p}^{\mathbb{R}}[r]$) which are real valued.

FUNCTIONAL SETTING 2

If a is a symbol in Γ^m_{W^{N,∞}} then ∂_xa ∈ Γ^m<sub>W<sup>N-1,∞</sub> and ∂_ξa ∈ Γ^{m-1}<sub>W<sup>N,∞</sub>. If b is a symbol in Γ^{m'}_{W^{N,∞}} then ab ∈ Γ^{m+m'}_{W^{N,∞}}. If a ∈ Γ^m_{≥p}[r] and b ∈ Γ^{m'}_{≥q}[r], then ab ∈ Γ^{m+m'}_{≥p+q}[r].
p-homogeneous symbols in Γ^m_p and non-homogeneous symbols in Γ^m_{≥p}[r] are actually functions with values in Γ^m_{W^{N,∞}} for some N ∈ N, whose seminorms (2.13) are bounded by
</sub></sup></sub></sup>

$$|a_p|_{m,W^{N,\infty},n} \le C_n \, \|U\|_1^{p-1} \|U\|_{N+\mu+1} \,, \quad |a|_{m,W^{N,\infty},n} \le C_n \, \|U\|_{s_0}^{p-1} \|U\|_{\mathfrak{s}} \,, \quad N \le \mathfrak{s} - s_0$$

• A p-homogeneous symbol $a_p(U, x, \xi)$ is a non-homogeneous symbol, since (2.14)–(2.15) imply

$$\left\|\partial_{\xi}^{\beta}a_{p}\left(U;\cdot,\xi\right)\right\|_{W^{N,\infty}} \leq C\langle\xi\rangle^{m-\beta}\|U\|_{1}^{p-1}\|U\|_{N+\mu+1} , \qquad (2.19)$$

and (2.14) implies the translation invariance property (2.17).

Paradifferential quantization. Given $p \in \mathbb{N}_0$ we consider functions $\chi_p \in C^{\infty}(\mathbb{R}^p \times \mathbb{R}; \mathbb{R})$ and $\chi \in C^{\infty}(\mathbb{R} \times \mathbb{R}; \mathbb{R})$, even with respect to each of their arguments, satisfying, for $0 < \delta_0 \leq \frac{1}{10}$,

$$\sup \chi_p \subset \{(\xi',\xi) \in \mathbb{R}^p \times \mathbb{R}; |\xi'| \le \delta_0 \langle \xi \rangle \}, \qquad \chi_p(\xi',\xi) \equiv 1 \text{ for } |\xi'| \le \delta_0 \langle \xi \rangle/2, \\ \operatorname{supp} \chi \subset \{(\xi',\xi) \in \mathbb{R} \times \mathbb{R}; |\xi'| \le \delta_0 \langle \xi \rangle \}, \qquad \chi(\xi',\xi) \equiv 1 \text{ for } |\xi'| \le \delta_0 \langle \xi \rangle/2.$$

For p = 0 we set $\chi_0 \equiv 1$. We assume moreover that

$$|\partial_{\xi}^{\ell}\partial_{\xi'}^{\beta}\chi_{p}(\xi',\xi)| \leq C_{\ell,\beta}\langle\xi\rangle^{-\ell-|\beta|}, \ \forall \ell \in \mathbb{N}_{0}, \ \beta \in \mathbb{N}_{0}^{p}, \ |\partial_{\xi}^{\ell}\partial_{\xi'}^{\beta}\chi(\xi',\xi)| \leq C_{\ell,\beta}\langle\xi\rangle^{-\ell-\beta}, \ \forall \ell, \ \beta \in \mathbb{N}_{0}.$$

If $a(x,\xi)$ is a smooth symbol we define its Weyl quantization as the operator acting on a 2π -periodic function u(x) (written as in (2.7)) as

$$Op^{W}(a)u = \sum_{k \in \mathbb{Z}} \left(\sum_{j \in \mathbb{Z}} \hat{a}\left(k - j, \frac{k + j}{2}\right) u_{j} \right) e^{ikx}$$

where $\hat{a}(k,\xi)$ is the k^{th} -Fourier coefficient of the 2π -periodic function $x \mapsto a(x,\xi)$.

Definition 2.2. (Bony-Weyl quantization) If $a(U; x, \xi)$ is a symbol in $\widetilde{\Gamma}_p^m$, respectively in $\Gamma_{W^{N,\infty}}^m$ or $\Gamma^m_{\geq p}[r]$, we set

$$a_{\chi_{p}}(U;x,\xi) := \sum_{\substack{\vec{j} \in \mathbb{Z}^{p} \\ \vec{\sigma} \in \{\pm\}^{p}}} \chi_{p}(\vec{j},\xi) a_{\vec{j}}^{\vec{\sigma}}(\xi) u_{\vec{j}}^{\vec{\sigma}} e^{i(\vec{\sigma}\cdot\vec{j})x}, \quad a_{\chi}(U;x,\xi) := \sum_{j \in \mathbb{Z}} \chi(j,\xi) \hat{a}(U;j,\xi) e^{ijx}$$
(2.20)

where in the last equality $\hat{a}(U; j, \xi)$ stands for j^{th} Fourier coefficient of $a(U; x, \xi)$ with respect to the x variable, and we define the Bony-Weyl quantization of $a(U; \cdot)$ as

$$Op^{BW}(a(U; \cdot))v = Op^{W}(a_{\chi_{p}}(U; \cdot))v = \sum_{\substack{(\vec{j}, j, k) \in \mathbb{Z}^{p+2} \\ \vec{\sigma} \in \{\pm\}^{p} \\ \vec{\sigma} \cdot \vec{\tau} + j = k}} \chi_{p}\left(\vec{j}, \frac{j+k}{2}\right) a_{\vec{j}}^{\vec{\sigma}}\left(\frac{j+k}{2}\right) u_{\vec{j}}^{\vec{\sigma}}v_{j}e^{ikx}, \quad (2.21)$$

$$Op^{BW}(a(U; \cdot))v = Op^{W}(a_{\chi}(U; \cdot))v = \sum_{(j,k)\in\mathbb{Z}^{2}} \chi\left(k - j, \frac{j + k}{2}\right) \hat{a}\left(U; k - j, \frac{k + j}{2}\right) v_{j}e^{ikx}.$$
 (2.22)

Note that if $\chi\left(k-j,\frac{k+j}{2}\right) \neq 0$ then $|k-j| \leq \delta_0 \langle \frac{j+k}{2} \rangle$ and therefore, for $\delta_0 \in (0,1)$,

$$\frac{1-\delta_0}{1+\delta_0}|k| \leq |j| \leq \frac{1+\delta_0}{1-\delta_0}|k|\,,\quad \forall j,k\in\mathbb{Z}\,.$$

This relation shows that the action of a paradifferential operator does not spread much the Fourier support of functions.

• If a is a homogeneous symbol, the two definitions of quantization in (2.21) and (2.22) differ by a smoothing operator according to Definition 2.5 below.

2 FUNCTIONAL SETTING

• Definition 2.2 is independent of the cut-off functions χ_p , χ , up to smoothing operators that we define below (see Definition 2.5), see the remark at page 50 of [9].

• Given a paradifferential operator $A = \operatorname{Op}^{BW}(a(x,\xi))$ it results

$$\overline{A} = \operatorname{Op}^{BW}\left(\overline{a(x, -\xi)}\right), \quad A^{\top} = \operatorname{Op}^{BW}(a(x, -\xi)), \quad A^* = \operatorname{Op}^{BW}\left(\overline{a(x, \xi)}\right),$$

where A^{\top} and A^* denote respectively the transposed and adjoint operator with respect to the complex, respectively real, scalar product of $L^2(\mathbb{T}, \mathbb{C})$ in (2.1). It results $A^* = \overline{A}^{\top}$. • A paradifferential operator $A = \operatorname{Op}^{BW}(a(x, \xi))$ is real (i.e. $A = \overline{A}$) if

$$\overline{a(x,\xi)} = a^{\vee}(x,\xi) \quad \text{where} \quad a^{\vee}(x,\xi) := a(x,-\xi) \,. \tag{2.23}$$

• A matrix of paradifferential operators $\operatorname{Op}^{BW}(A(x,\xi))$ is real-to-real, i.e. (2.3) holds, if and only if the matrix of symbols $A(x,\xi)$ has the form

$$A(x,\xi) = \begin{pmatrix} \underline{a(x,\xi)} & \underline{b(x,\xi)} \\ \overline{b^{\vee}(x,\xi)} & \overline{a^{\vee}(x,\xi)} \end{pmatrix} = \begin{pmatrix} a(x,\xi) & \underline{0} \\ 0 & \overline{a^{\vee}(x,\xi)} \end{pmatrix} + \begin{pmatrix} \underline{0} & b(x,\xi) \\ \overline{b^{\vee}(x,\xi)} & 0 \end{pmatrix} .$$
(2.24)

• A real-to-real matrix of U-dependent paradifferential operators $\operatorname{Op}^{BW}(A(U; x, \xi))$ is gauge invariant, i.e. (2.4) holds, if and only if the symbols in (2.24) fulfill, with g_{θ} in (2.2),

$$a(U;x,\xi) = a(\mathsf{g}_{\theta}U;x,\xi) , \quad e^{\mathrm{i}2\theta} b(U;x,\xi) = b(\mathsf{g}_{\theta}U;x,\xi) , \quad \forall \theta \in \mathbb{T} , \qquad (2.25)$$

If, in addition, $a, b \in \widetilde{\Gamma}_p^m$, then $\operatorname{Op}^{BW}(a)$ in (2.21) have indexes restricted to $\vec{\sigma} \cdot 1 = 0$, whereas $\operatorname{Op}^{BW}(b)$ to $\vec{\sigma} \cdot 1 = 2$.

We will use also the notations

$$\operatorname{Op}_{\mathsf{vec}}^{BW}(a(x,\xi)) := \operatorname{Op}^{BW}\left(\begin{bmatrix} a(x,\xi) & 0\\ 0 & a^{\vee}(x,\xi) \end{bmatrix} \right) , \ \operatorname{Op}_{\mathsf{out}}^{BW}(b(x,\xi)) := \operatorname{Op}^{BW}\left(\begin{bmatrix} 0 & b(x,\xi)\\ \overline{b^{\vee}(x,\xi)} & 0 \end{bmatrix} \right)$$
(2.26)

Along the paper we shall use the following results concerning the action of a paradifferential operator in Sobolev spaces. We refer to [13, Theorem A.7] for the proof of (i) and to [9, Proposition 3.8] for the proof of (ii), (iii).

Theorem 2.3. (Continuity of Bony-Weyl operators) Let $m \in \mathbb{R}$, $p \in \mathbb{N}$, r > 0. Then:

(i) Let $a \in \Gamma_{L^{\infty}}^{m}$. Then $\operatorname{Op}^{BW}(a)$ extends to a bounded operator $H^{s} \to H^{s-m}$ for any $s \in \mathbb{R}$ satisfying the estimate, for any $u \in H^{s}$,

$$\|\operatorname{Op}^{BW}(a) \, u\|_{s-m} \lesssim \|a\|_{m,L^{\infty},4} \, \|u\|_{s} \, . \tag{2.27}$$

(ii) Let $a \in \widetilde{\Gamma}_p^m$. There is $s_0 > 0$ such that for any $s \in \mathbb{R}$, there is a constant C > 0, depending only on s and on (2.15) with $\ell = \beta = 0$, such that for any $U_1, \ldots, U_p \in H^{s_0}(\mathbb{T}, \mathbb{C}^2)$ and $v \in H^s(\mathbb{T}, \mathbb{C})$, one has

$$\|\operatorname{Op}^{^{BW}}(a(U_1,\ldots,U_p;\cdot))v\|_{s-m} \le C \prod_{j=1}^p \|U_j\|_{s_0} \|v\|_s, \qquad (2.28)$$

for $p \ge 1$, while for p = 0 the (2.28) holds by replacing the right hand side with $C ||v||_s$.

(iii) Let $a \in \Gamma_{\geq p}^{m}[r]$. There is $s_0 > 0$ such that for any $s \in \mathbb{R}$ there is a constant C > 0 such that for any $U \in B_{s_0}(r)$ one has

$$\|\operatorname{Op}^{BW}(a(U; \cdot))\|_{\mathcal{L}(H^{s}, H^{s-m})} \le C \|U\|_{s_{0}}^{p}.$$
(2.29)

Classes of *m*-operators and smoothing operators. We introduce *m*-operators and smoothing operators. This is a small adaptation of [9, 13] where we consider only autonomous maps, where again the time dependence is only through U(t). In particular we put K, K' = 0 with respect to the notation in [9, 13]. Given integers $(n_1, \ldots, n_{p+1}) \in \mathbb{N}^{p+1}$, we denote by $\max_2(n_1, \ldots, n_{p+1})$ the second largest among n_1, \ldots, n_{p+1} .

Definition 2.4 (Classes of *m*-operators). Let $m \in \mathbb{R}$, $p \in \mathbb{N}_0$ and r > 0.

1. p-homogeneous m-operators. We denote by $\widetilde{\mathcal{M}}_p^m$ the class of (p+1)-linear operators from $(C^{\infty}(\mathbb{T};\mathbb{C}^2))^p \times C^{\infty}(\mathbb{T};\mathbb{C})$ to $C^{\infty}(\mathbb{T};\mathbb{C})$ of the form $(U_1,\ldots,U_p,v) \to M_p(U_1,\ldots,U_p)v$, symmetric in (U_1,\ldots,U_p) , with Fourier expansion

$$M_{p}(U)v := M_{p}(U, \dots, U)v = \sum_{\substack{\vec{\sigma}_{p} \in \{\pm\}^{p} \\ k-j = \vec{\sigma}_{p} \cdot \vec{j}_{p}}} M_{\vec{j}_{p}, j, k}^{\vec{\sigma}_{p}} u_{\vec{j}_{p}}^{\vec{\sigma}_{p}} v_{j} e^{ikx}$$
(2.30)

that satisfy the following. There are $\mu \geq 0$, C > 0 such that for any $(j_p, j, k) \in \mathbb{Z}^{p+2}$, $\vec{\sigma}_p \in \{\pm\}^p$, one has

$$|M_{\vec{j}_p,j,k}^{\vec{\sigma}_p}| \le C \max_2\{\langle j_1 \rangle, \dots, \langle j_p \rangle, \langle j \rangle\}^{\mu} \max\{\langle j_1 \rangle, \dots, \langle j_p \rangle, \langle j \rangle\}^{m}.$$
(2.31)

2. Non-homogeneous m-operators. We denote by $\mathcal{M}_{\geq p}^m[r]$ the class of operators $(U, v) \mapsto M(U)v$ defined on $B_{s_0}(r) \times H^{s_0}(\mathbb{T}, \mathbb{C})$ for some $s_0 > 0$, which are linear in the variable v and such that the following holds true. For any $s \geq s_0$ there are C > 0 and $r' = r'(s) \in]0, r[$ such that for any $U \in B_{s_0}(r') \cap H^s(\mathbb{T}, \mathbb{C}^2)$, any $v \in H^s(\mathbb{T}, \mathbb{C})$, we have that

$$||M(U)v||_{s-m} \le C\left(||v||_s ||U||_{s_0}^p + ||v||_{s_0} ||U||_{s_0}^{p-1} ||U||_s\right).$$
(2.32)

In addition we require the translation invariance property

$$M(\tau_{\varsigma}U)[\tau_{\varsigma}v] = \tau_{\varsigma}(M(U)v), \quad \forall \varsigma \in \mathbb{R}.$$
(2.33)

where τ_{ς} is the translation operator in (2.2).

3. m-Operators. We denote by $\Sigma \mathcal{M}_0^m[r]$ the space of operators $(U,c) \to M(U)v$ of the form

$$M(U) = M_0 + M_2(U) + M_{\ge 4}(U)$$
(2.34)

where $M_p(U)$ in $\widetilde{\mathcal{M}}_p^m$, $p \in \{0, 2\}$, and $M_{\geq 4}(U)$ in $\mathcal{M}_{\geq 4}^m[r]$. We denote by $\Sigma \mathcal{M}_2^m[r]$ the operators of the form (2.34) with $M_0 = 0$. Finally sometimes we shall write $\Sigma \mathcal{M}_4^m[r] \equiv \mathcal{M}_{\geq 4}^m[r]$.

• A *p*-homogeneous *m*-operator M_p is a non-homogeneous *m*-operator. Indeed (2.31) implies the quantitative estimate: for $s_0 \ge \mu + 1 > 0$, for any $s \ge s_0$, any $U \in H^s(\mathbb{T}; \mathbb{C}^2)$, any $v \in H^s(\mathbb{T}; \mathbb{C})$

$$\|M_p(U)v\|_{s-m} \lesssim_s \|U\|_{s_0}^p \|V\|_s + \|U\|_{s_0}^{p-1} \|U\|_s \|V\|_{s_0}$$
(2.35)

which is (2.32) (see Lemma 2.8 and 2.9 in [13] for a proof). Moreover (2.33) follows from the Fourier restriction $k - j = \vec{\sigma}_p \cdot \vec{j}_p$ in (2.30).

• (Paradifferential operators as *m*-operators) If $a(U; x, \xi)$ is a symbol in $\Sigma\Gamma_0^m[r]$ then the paradifferential operator $\operatorname{Op}^{\mathbb{B}W}(a(U; x, \xi))$ is an *m*-operator $\Sigma\mathcal{M}_0^m[r]$. This is a consequence of Theorem 2.3–(*ii*)&(*iii*).

• We will meet vector fields of the form X(U) = M(U)U where M(U) is a matrix of *p*-homogeneous *m*-operators as in (2.9). In this case the relation between the Fourier coefficients of the vector field in (2.12) and those of the *m*-operator in (2.30) is given by

$$X_{j_1,\dots,j_p,j_{p+1},k}^{\sigma_1,\dots,\sigma_p,\sigma_{p+1},\sigma} = \frac{1}{p+1} \left(M_{j_1,\dots,j_p,j_{p+1},k}^{\sigma_1,\dots,\sigma_p,\sigma_{p+1},\sigma} + M_{j_{p+1},\dots,j_p,j_1,k}^{\sigma_{p+1},\dots,\sigma_p,\sigma_1,\sigma} + \dots + M_{j_1,\dots,j_{p+1},j_p,k}^{\sigma_1,\dots,\sigma_{p+1},\sigma_p,\sigma} \right) , \qquad (2.36)$$

namely they are obtained symmetrizing with respect to the second last index (j, σ') the coefficients $M_{j_p,j,k}^{\vec{\sigma}_p,\sigma',\sigma}$ of M(U).

If $m \leq 0$ the *m*-operators are referred to as smoothing operators.

Definition 2.5. (Smoothing operators) Let $\rho \geq 0$, $p \in \mathbb{N}_0$ and $q \in \{0,2\}$. We define the *p*-smoothing operators

$$\widetilde{\mathcal{R}}_p^{-\varrho} := \widetilde{\mathcal{M}}_p^{-\varrho}, \quad \mathcal{R}_{\geq p}^{-\varrho}[r] := \mathcal{M}_{\geq p}^{-\varrho}[r] , \quad \Sigma \mathcal{R}_q^{-\varrho}[r] := \Sigma \mathcal{M}_q^{-\varrho}[r] .$$

• In view of (2.31) a homogeneous *m*-operator in $\widetilde{\mathcal{M}}_p^m$ with the property that, on its support, $\max_{2}\{\langle j_{1}\rangle,\ldots,\langle j_{p}\rangle,\langle j\rangle\}\sim\max\{\langle j_{1}\rangle,\ldots,\langle j_{p}\rangle,\langle j\rangle\}$ is actually a smoothing operator in $\widetilde{\mathcal{R}}_{p}^{-\varrho}$ for any $\rho \geq 0$ satisfying (2.31) with $\mu \rightsquigarrow \mu + m + \rho$ and $m \rightsquigarrow -\rho$.

• The Definition 2.5 of smoothing operators is modeled to gather remainders which satisfy either the property $\max_2(n_1,\ldots,n_{p+1}) \sim \max(n_1,\ldots,n_{p+1})$ or arise as remainders of compositions of paradifferential operators, see Proposition 2.7 below, and thus have a fixed order ρ of regularization. **Composition theorems.** Let $D_x := \frac{1}{i} \partial_x$. The following is Definition 3.11 in [9].

Definition 2.6. (Asymptotic expansion of composition symbol) Let $\varrho \geq 0, m, m' \in \mathbb{R}$, r > 0. Consider symbols $a \in \Sigma\Gamma_p^m[r]$ and $b \in \Sigma\Gamma_{p'}^{m'}[r]$, $p, p' \in \{0, 2\}$. For U in $B_{\mathfrak{s}}(I; r)$ we define, for $\rho < \mathbf{s} - s_0$, the symbol

$$(a\#_{\varrho}b)(U;x,\xi) := \sum_{k=0}^{\varrho} \frac{1}{2^{k}} \sum_{\ell+\beta=k} \frac{(-1)^{\beta}}{\ell!\beta!} (\partial_{\xi}^{\ell} D_{x}^{\beta} a) \cdot (\partial_{\xi}^{\beta} D_{x}^{\ell} b)(U;x,\xi) .$$
(2.37)

- The symbol a#_ρb belongs to ΣΓ^{m+m'}_{p+p'}[r].
 We have that a#_ρb = ab + ¹/_{2i}{a, b} up to a symbol in ΣΓ^{m+m'-2}_{p+p'}[r], where

$$\{a,b\} := \partial_{\xi} a \partial_x b - \partial_x a \partial_{\xi} b \in \Sigma \Gamma^{m+m'-1}_{p+p'}[r]$$
(2.38)

denotes the Poisson bracket. Moreover if $a \in \Gamma_{W^{N,\infty}}^m$ and $b \in \Gamma_{W^{N,\infty}}^{m'}$ then $\{a,b\} \in \Gamma_{W^{N-1,\infty}}^{m+m'-1}$ with estimate

$$|\{a,b\}|_{m+m'-1,W^{N-1,\infty},n} \lesssim |a|_{m,W^{N,\infty},n+1} |b|_{m',W^{N,\infty},n+1}.$$
(2.39)

- Due to (2.18), the symbol $a \#_{\rho} b$ does not contain symbols of odd homogeneity.
- $\overline{a^{\vee}} \#_{\varrho} \overline{b^{\vee}} = \overline{a \#_{\varrho} b^{\vee}}$ where a^{\vee} is defined in (2.23).

The following proposition is proved in [13, Theorem A.8] and [9, Proposition 3.12].

Proposition 2.7. (Composition of Bony-Weyl operators) Let $m, m' \in \mathbb{R}$, $p, p' \in \{0, 2\}$, $\rho \ge 0$ and r > 0.

(i) Let $a \in \Gamma^m_{W^{\varrho,\infty}}$, $b \in \Gamma^{m'}_{W^{\varrho,\infty}}$. Then

$$\operatorname{Op}^{BW}(a) \operatorname{Op}^{BW}(b) = \operatorname{Op}^{BW}(a \#_{\varrho} b) + R(a, b)$$

where the linear operator $R(a,b): H^s \to H^{s-(m+m')+\varrho}, \forall s \in \mathbb{R}$, satisfies, for some $N = N(\varrho) > 0$,

$$\|R(a,b)u\|_{s-(m+m')+\varrho} \lesssim \left(|a|_{m,W^{\varrho,\infty},N} |b|_{m',L^{\infty},N} + |a|_{m,L^{\infty},N} |b|_{m',W^{\varrho,\infty},N}\right) \|u\|_{s} .$$
(2.40)

One can take N(2) = 7. (ii) Let $a \in \Sigma \Gamma_p^m[r]$, $b \in \Sigma \Gamma_{p'}^{m'}[r]$. Then

$$\operatorname{Op}^{BW}(a(U;x,\xi)) \circ \operatorname{Op}^{BW}(b(U;x,\xi)) = \operatorname{Op}^{BW}((a \#_{\varrho} b)(U;x,\xi)) + R(U)$$

where R(U) are smoothing operators in $\Sigma \mathcal{R}_{p+p'}^{-\varrho+m+m'}[r]$.

• Let $a(U) \in \Sigma \Gamma_p^m[r]$ and $b(U) \in \Sigma \Gamma_{p'}^{m'}[r]$, with the notation in (2.26), one has

$$\begin{bmatrix} \operatorname{Op}_{\mathsf{out}}^{BW}(b), \operatorname{Op}_{\mathsf{vec}}^{BW}(a) \end{bmatrix} = \operatorname{Op}_{\mathsf{out}}^{BW}\left(b \#_{\varrho} \overline{a^{\vee}} - a \#_{\varrho} b\right) + R(U) \\ \begin{bmatrix} \operatorname{Op}_{\mathsf{out}}^{BW}(a), \operatorname{Op}_{\mathsf{out}}^{BW}(b) \end{bmatrix} = \operatorname{Op}_{\mathsf{vec}}^{BW}\left(a \#_{\varrho} \overline{b^{\vee}} - b \#_{\varrho} \overline{a^{\vee}}\right) + R(U) \\ \begin{bmatrix} \operatorname{Op}_{\mathsf{vec}}^{BW}(a), \operatorname{Op}_{\mathsf{vec}}^{BW}(b) \end{bmatrix} = \operatorname{Op}_{\mathsf{vec}}^{BW}\left(a \#_{\varrho} b - b \#_{\varrho} a\right) + R(U)$$
(2.41)

where R(U) are real-to-real matrices of smoothing operators in $\Sigma \mathcal{R}_{p+p'}^{-\varrho+m'+m}[r]$. We conclude this section with the paralinearization of the product (see [9, Lemma 7.2]). Lemma 2.8. (Bony paraproduct decomposition) Let f, g, h be functions in $H^{\sigma}(\mathbb{T}; \mathbb{C})$ with $\sigma > \frac{1}{2}$. Then

$$fgh = \operatorname{Op}^{BW}(fg)h + \operatorname{Op}^{BW}(fh)g + \operatorname{Op}^{BW}(gh)f + R_1(f,g)h + R_2(f,h)g + R_3(g,h)f$$

where for j = 1, 2, 3, R_j is a homogeneous smoothing operator in $\widetilde{\mathcal{R}}_1^{-\varrho}$ for any $\varrho \geq 0$.

Composition of *m***-operators.** The following lemma, which is a consequence of Proposition 2.15 (items (ii) and (iv)) in [13], shall be used below.

Lemma 2.9. Let $m, m', m_0 \in \mathbb{R}$, $\varrho \geq 0$, r > 0, $p \in \{0, 2\}$. Let M(U) be a real-to-real matrix of *m*-operators in $\Sigma \mathcal{M}_2^m[r]$, $\mathbf{F}(U)$ be a real-to-real matrix of 0-operators $\mathcal{M}_{\geq 0}^0[r]$ and $\mathbf{p}(\xi)$ be a matrix of Fourier multipliers in $\widetilde{\Gamma}_0^{m_0}$. Then:

1. If c(U) is a 2-homogeneous symbol in $\widetilde{\Gamma}_2^{m'}$ and $c_{>4}(U)$ is a non-homogeneous symbol in $\Gamma_{>4}^m[r]$,

$$b_2(U; x, \xi) := c(-\mathrm{ip}(D)U; x, \xi), \quad and \quad \begin{cases} b_{\geq 4}(U; x, \xi) := c(M(U)U, U; x, \xi) \\ b'_{\geq 4}(U; x, \xi) := c_{\geq 4}(\mathbf{F}(U)U; x, \xi) \end{cases}$$

are symbols respectively in $\widetilde{\Gamma}_{2}^{m'}$ and $\Gamma_{\geq 4}^{m'}[r']$ for some r' > 0;

2. If Q(U) is a 2-homogeneous smoothing operator in $\widetilde{\mathcal{R}}_2^{-\varrho}$,

$$\widetilde{R}_{2}(U) := Q(-\mathrm{ip}(D)U, U) \in \widetilde{\mathcal{R}}_{2}^{-\varrho + \max\{0, m_{0}\}} \quad and \quad R_{\geq 4}(U) := Q(M(U)U, U) \in \mathcal{R}_{\geq 4}^{-\varrho + \max\{0, m\}}[r];$$

- $\begin{aligned} & 3. \ If \ R(U) \in \Sigma \mathcal{R}_2^{-\varrho}[r] \ and \ \mathsf{a}(U; x, \xi) \in \Sigma \Gamma_2^m[r], \ \varrho \geq m, \ then \\ & R(U) \circ \operatorname{Op}^{^{BW}}(\mathsf{a}(U; x, \xi)) \in \mathcal{R}_{\geq 4}^{-\varrho + m}[r], \qquad \operatorname{Op}^{^{BW}}(\mathsf{a}(U; x, \xi)) \circ R(U) \in \mathcal{R}_{\geq 4}^{-\varrho + m}[r]. \end{aligned}$
- 4. If M is in $\Sigma \mathcal{M}_p^m[r]$ and M' is in $\Sigma \mathcal{M}_{p'}^{m'}[r]$ then the composition $M \circ M'$ is in $\Sigma \mathcal{M}_{p+p'}^{m+\max(m',0)}[r]$.
- 5. If M(U) is in $\mathcal{M}_{\geq 4}^m[r]$, then $M(\mathbf{F}(U)U)$ is in $\mathcal{M}_{\geq 4}^m[r']$ for some r' > 0.

2.2 Admissible transformations

In this section we introduce a class of U-dependent transformations that are C^1 with respect to U.

Definition 2.10 (Admissible transformations). Let r > 0 and $m \ge 0$. We say that a real-to-real matrix $\mathbf{F}(U)$ of non-homogeneous 0-operators in $\mathcal{M}^{0}_{\ge 0}[r]$ is an m-admissible transformation if the following holds:

- (i) Linear invertibility: $\mathbf{F}(U)$ is linearly invertible and its inverse $\mathbf{F}(U)^{-1}$ is a real-to-real matrix of non-homogeneous 0-operators in $\mathcal{M}^0_{\geq 0}[r]$;
- (ii) Expansion: $\mathbf{F}(U)$ Id is a matrix of m-operators in $\Sigma \mathcal{M}_2^m[r]$ expanding as

$$\mathbf{F}(U) = \mathrm{Id} + \mathbf{F}_2(U) + \mathbf{F}_{\geq 4}(U), \quad \mathbf{F}_2(U) \in \widetilde{\mathcal{M}}_2^m, \quad \mathbf{F}_{\geq 4}(U) \in \mathcal{M}_{\geq 4}^m[r].$$
(2.42)

(iii) **Derivative:** there is $s_0 \ge 0$ such that for any $Z \in H^{s_0+m}_{\mathbb{R}}(\mathbb{T};\mathbb{C}^2)$ one has

$$U \mapsto \mathbf{F}(U)Z \in C^1\left(B_{s_0,\mathbb{R}}(r); H^{s_0}_{\mathbb{R}}(\mathbb{T}; \mathbb{C}^2)\right)$$

Moreover, for any $s \geq s_0 + m$, there is $C := C_s > 0$ such that for any $U \in B_{s_0,\mathbb{R}}(r) \cap H^s_{\mathbb{R}}(\mathbb{T};\mathbb{C}^2)$, $Z, \hat{U} \in H^s_{\mathbb{R}}(\mathbb{T};\mathbb{C}^2)$ one has

$$\| \left(\mathrm{d}_{U} \mathbf{F}(U)[\hat{U}] - \mathrm{d}_{U} \mathbf{F}_{2}(U)[\hat{U}] \right) Z \|_{s-m} = \| \mathrm{d}_{U} \mathbf{F}_{\geq 4}(U)[\hat{U}] Z \|_{s-m}$$

$$\leq C \left(\| U \|_{s_{0}}^{3} \| \hat{U} \|_{s_{0}} \| Z \|_{s} + \| U \|_{s_{0}}^{3} \| \hat{U} \|_{s} \| Z \|_{s_{0}} + \| U \|_{s} \| U \|_{s_{0}}^{2} \| \hat{U} \|_{s_{0}} \| Z \|_{s_{0}} \right).$$

$$(2.43)$$

Remark 2.11. (1) Property (i) is equivalent to say that there exists $s_0 \ge 0$ such that for any $s \ge s_0$ there is a constant $C := C_s > 0$ and $r = r_s > 0$ such that for any $U \in B_{s_0,\mathbb{R}}(r)$ and $V \in H^s(\mathbb{T};\mathbb{C}^2)$ one has

$$\|\mathbf{F}(U)V\|_{s} + \|\mathbf{F}^{-1}(U)V\|_{s} \le C\|V\|_{s} .$$
(2.44)

(2) Thanks to the bound in (2.44), $\mathbf{F}(U)$ conjugates any matrix $B_{\geq 4}(U)$ of 0-operators in $\mathcal{M}^{0}_{\geq 4}[r]$ into another matrix of 0-operators in $\mathcal{M}^{0}_{\geq 4}[r]$, namely $\mathbf{F}(U)B_{\geq 4}(U)\mathbf{F}(U)^{-1}$ is a matrix of 0-operators in $\mathcal{M}^{0}_{\geq 4}[r]$.

(3) Property (ii) implies that

$$\| \left[\mathbf{F}(U) - \mathrm{Id} \right] V \|_{s-m} + \| \left[\mathbf{F}^{-1}(U) - \mathrm{Id} \right] V \|_{s-m} \le C \| U \|_{s_0}^2 \| V \|_s$$
(2.45)

and that

$$\|\mathbf{d}_U \mathbf{F}_2(U)[\hat{U}]V\|_{s-m} \lesssim_s \|U\|_{s_0} \|\hat{U}\|_{s_0} \|V\|_s + \|U\|_{s_0} \|\hat{U}\|_s \|V\|_{s_0} + \|U\|_s \|\hat{U}\|_{s_0} \|V\|_{s_0} .$$
(2.46)

(4) The expansion (2.42) for $\mathbf{F}(U)$ implies the corresponding expansion for $\mathbf{F}(U)^{-1}$:

$$\mathbf{F}(U)^{-1} = \mathrm{Id} - \mathbf{F}_2(U) + \check{\mathbf{F}}_{\geq 4}(U),$$

where $\check{\mathbf{F}}_{\geq 4}(U) := -\mathbf{F}(U)^{-1}\mathbf{F}_{\geq 4}(U) + \mathbf{F}(U)^{-1}[\mathbf{F}(U) - \mathrm{Id}]\mathbf{F}_{2}(U)$ is a real-to-real matrix of 2*m*-operators in $\mathcal{M}_{\geq 4}^{2m}[r]$.

We now prove that admissible transformations are closed by composition.

Lemma 2.12. If $\mathbf{F}^{(1)}(U)$ is m_1 -admissible and $\mathbf{F}^{(2)}(U)$ is m_2 -admissible, then the composition $\mathbf{F}^{(1)}(U)\mathbf{F}^{(2)}(U)$ is a $m_1 + m_2$ -admissible transformation.

Proof. We set $m := m_1 + m_2$. (i) and (ii) follow from the composition properties of *m*-operators, see Lemma 2.9-4. Moreover we have the expansion

$$\mathbf{F}^{(1)}(U)\mathbf{F}^{(2)}(U) = \mathrm{Id} + \mathbf{F}_{2}^{(1)}(U) + \mathbf{F}_{2}^{(2)}(U) + \mathbf{F}_{\geq 4}^{(1,2)}(U)$$
(2.47)

where $\mathbf{F}_{\geq 4}^{(1,2)}(U) = \mathbf{F}_{\geq 4}^{(1)}(U) + \mathbf{F}_{\geq 4}^{(2)}(U) + \left(\mathbf{F}_{2}^{(1)}(U) + \mathbf{F}_{\geq 4}^{(1)}(U)\right) \left(\mathbf{F}_{2}^{(2)}(U) + \mathbf{F}_{\geq 4}^{(2)}(U)\right) \in \mathcal{M}_{\geq 4}^{m}[r]$. We prove item (*iii*). Note that, in view of the expansion (2.47), it is sufficient to prove that $\mathbf{F}_{\geq 4}^{(1,2)}(U)Z \in C^{1}\left(B_{s_{0},\mathbb{R}}(r); H_{\mathbb{R}}^{s_{0}}(\mathbb{T}; \mathbb{C}^{2})\right)$ for any $Z \in H_{\mathbb{R}}^{s_{0}+m}(\mathbb{T}; \mathbb{C}^{2})$ with estimate (2.43). First we compute the differential

$$d_{U}\mathbf{F}_{\geq 4}^{(1,2)}(U)[\hat{U}]Z = d_{U}\mathbf{F}_{\geq 4}^{(1)}(U)[\hat{U}]Z + d_{U}\mathbf{F}_{\geq 4}^{(2)}(U)[\hat{U}]Z + \left(d_{U}\mathbf{F}_{2}^{(1)}(U)[\hat{U}] + d_{U}\mathbf{F}_{\geq 4}^{(1)}(U)[\hat{U}]\right) \left(\mathbf{F}_{2}^{(2)}(U) + \mathbf{F}_{\geq 4}^{(2)}(U)\right) Z + \left(\mathbf{F}_{2}^{(1)}(U) + \mathbf{F}_{\geq 4}^{(1)}(U)\right) \left(d_{U}\mathbf{F}_{2}^{(2)}(U)[\hat{U}] + d_{U}\mathbf{F}_{\geq 4}^{(2)}(U)[\hat{U}]\right) Z.$$

Estimate (2.43) for $d_U \mathbf{F}_{\geq 4}^{(1,2)}(U)[\hat{U}]Z$ follows from the corresponding estimates for $d_U \mathbf{F}_{\geq 4}^{(1)}(U)[\hat{U}]Z$, $d_U \mathbf{F}_{\geq 4}^{(2)}(U)[\hat{U}]Z$ in (2.46) and (2.32)–(2.35) for $\mathbf{F}_2^{(1)}(U)$, $\mathbf{F}_2^{(2)}(U)$, $\mathbf{F}_{\geq 4}^{(1)}(U)$ and $\mathbf{F}_{\geq 4}^{(2)}(U)$.

Next we prove a local invertibility property of the nonlinear map $U \mapsto \mathbf{F}(U)U$ when $\mathbf{F}(U)$ is an admissible transformation.

Lemma 2.13. Let $\mathbf{F}(U)$ be a m-admissible transformation, and consider the nonlinear map $\mathcal{F}(U) := \mathbf{F}(U)U$. The following holds true:

(i) There exists $s'_0 \geq 0$ such that for any $s \geq s'_0$, the map \mathcal{F}^{-1} is locally invertible: namely there is r' > 0 and $\mathcal{F}^{-1} : B_{s'_0,\mathbb{R}}(r') \cap H^s_{\mathbb{R}}(\mathbb{T};\mathbb{C}^2) \to H^s_{\mathbb{R}}(\mathbb{T};\mathbb{C}^2)$ such that

$$\mathcal{F} \circ \mathcal{F}^{-1}(V) = V, \quad \mathcal{F}^{-1} \circ \mathcal{F}(U) = U, \quad \forall U, V \in B_{s'_{0},\mathbb{R}}(r') .$$

(ii) One has $\mathcal{F}^{-1}(V) = \mathbf{G}(V)V$ with $\mathbf{G}(V)$ a matrix of non-homogeneous 0-operators in $\mathcal{M}^{0}_{\geq 0}[r']$ such that $\mathbf{G}(V) - \mathrm{Id} \in \Sigma \mathcal{M}^{2m}_{2}[r']$ for some r' > 0 and expands as

$$\mathbf{G}(V) = \mathrm{Id} - \mathbf{F}_2(V) + \mathbf{G}_{\geq 4}(V) , \quad \mathbf{G}_{\geq 4}(V) \in \mathcal{M}_{\geq 4}^{2m}[r'] .$$
 (2.48)

Proof. Fix $s_0, r > 0$ the parameters given by Definition 2.10 associated to $\mathbf{F}(U)$.

(i) We define $U = \mathcal{F}^{-1}(V)$ as the unique solution of the equation $V = \mathcal{F}(U) = \mathbf{F}(U)U$, which thanks to the linear invertibility of $\mathbf{F}(U)$, it is equivalent to

$$\mathcal{G}(U;V) := \mathbf{F}(U)^{-1}V - U = 0$$

We apply the implicit function theorem. Clearly $\mathcal{G}(0;0) = 0$. By the property (*ii*) of admissible transformations, $\mathcal{G} \in C^1\left(B_{s_0,\mathbb{R}}(r) \times H^{s_0+m}_{\mathbb{R}}(\mathbb{T};\mathbb{C}^2); H^{s_0}_{\mathbb{R}}(\mathbb{T};\mathbb{C}^2)\right)$. Moreover $d_U\mathcal{G}(0;0) = -\mathrm{Id}$: $H^{s_0}_{\mathbb{R}} \to H^{s_0}_{\mathbb{R}}$. Then, in conclusion, there is $r_1 > 0$ such that for any $V \in B_{s_0+m,\mathbb{R}}(r_1)$ there is a unique $U := \mathcal{F}^{-1}(V) \in B_{s_0,\mathbb{R}}(r)$ such that

$$0 \equiv \mathcal{G}(\mathcal{F}^{-1}(V); V) = \mathbf{F}(\mathcal{F}^{-1}(V))^{-1}V - \mathcal{F}^{-1}(V) , \quad \forall V \in B_{s_0+m,\mathbb{R}}(r_1)$$
(2.49)

which implies that $\mathcal{F} \circ \mathcal{F}^{-1}(V) = V$. In addition, from equation (2.49) we get, for any $V \in B_{s_0+m,\mathbb{R}}(r_1) \cap H^s_{\mathbb{R}}(\mathbb{T},\mathbb{C}^2)$, $\mathcal{F}^{-1}(V)$ belongs to $H^s_{\mathbb{R}}(\mathbb{T},\mathbb{C}^2)$ for any $s \geq s_0 + m$ and

$$\|\mathcal{F}^{-1}(V)\|_s = \|\mathbf{F}^{-1}(\mathcal{F}(V))^{-1}V\|_s \le C_s \|V\|_s$$
.

Moreover, by (2.45), for some C > 1, we have also

$$\|\mathcal{F}(U)\|_{s_0+m} = \|\mathbf{F}(U)U\|_{s_0+m} \le C \|U\|_{s_0+m} \le r_1$$

for any $U \in B_{s_0+m,\mathbb{R}}(r_1/C)$. The thesis of item (i) follows by choosing $r' := r_1/C$.

(ii) It follows from (2.49)

$$\mathcal{F}^{-1}(V) = \mathbf{G}(V)V$$
, $\mathbf{G}(V) := \mathbf{F}(\mathcal{F}^{-1}(V))^{-1} \in \mathcal{M}^{0}_{\geq 0}[r'].$ (2.50)

Since by definition $r' = r_1/C \leq r_1$, by the implicit function theorem, $\mathcal{F}^{-1}(V) \in B_{s_0,\mathbb{R}}(r)$ for any $V \in B_{s_0+m,\mathbb{R}}(r')$. Then, since $\mathbf{F}(U)^{-1}$ is a real-to-real matrix of non-homogeneous 0-operators in $\mathcal{M}^0_{\geq 0}[r]$, it follows that $\mathbf{G}(V)$ is a real-to-real matrix of non-homogeneous 0-operators in $\mathcal{M}^0_{\geq 0}[r']$ (with $s_0 \rightsquigarrow s_0 + m$).

Next we show that $\mathbf{G}(V)$ expands as in (2.48). Put $\tilde{\mathcal{F}}^{-1}(V) := V - \mathbf{F}_2(V)V$. Then, using the expansion $\mathcal{F}(U) = U + \mathbf{F}_2(U)U + \mathbf{F}_{\geq 4}(U)U$ and Lemma 2.9, we get

$$(\widetilde{\mathcal{F}}^{-1} \circ \mathcal{F})(U) = U + \mathbf{F}'_{\geq 4}(U)U$$
, with $\mathbf{F}'_{\geq 4}(U) \in \mathcal{M}^{2m}_{\geq 4}[r].$

Substituting $U = \mathcal{F}^{-1}(V)$ and using (2.50) and Lemma 2.9, we obtain

$$\mathcal{F}^{-1}(V) = V - \mathbf{F}_2(V)V + \mathbf{G}_{\geq 4}(V)V, \quad \mathbf{G}_{\geq 4}(V) := -\mathbf{F}'_{\geq 4}(\mathcal{F}^{-1}(V))\mathbf{G}(V) \in \mathcal{M}^{2m}_{\geq 4}[r']$$

This proves the expansion in (2.48).

An immediate consequence of the above lemma is that the inverse \mathcal{F}^{-1} of an admissible transformation \mathcal{F} fulfills the estimate scritta male, controllo norma bassa

$$\|\mathcal{F}^{-1}(V)\|_s \le C_s \|V\|_s, \quad \text{for any } V \in B_{s'_0,\mathbb{R}}(r') \cap H^s(\mathbb{T};\mathbb{C}^2) .$$

$$(2.51)$$

We now show that the linear flows generated by two types of paradifferential operators are admissible transformations. Consider the flows

$$\begin{cases} \partial_{\tau} \Phi^{\tau}(U) = G(\tau, U) \Phi^{\tau}(U) \\ \Phi^{0}(U) = \mathrm{Id} \end{cases} \quad \text{where} \quad G(\tau, U) = \begin{cases} \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\frac{\beta(U;x)}{1 + \tau \beta_{x}(U;x)} \mathrm{i}\xi \right), & \beta \in \widetilde{\mathcal{F}}_{2}^{\mathbb{R}} \text{ or} \\ \operatorname{Op}_{\mathsf{out}}^{BW}(g(U;x,\xi)), & g \in \widetilde{\Gamma}_{2}^{0}. \end{cases}$$
(2.52)

FUNCTIONAL SETTING 2

Remark 2.14. The map $\Phi^{\tau}(U)$ is gauge invariant if the generator $G(U;\tau)$ is gauge invariant. Indeed $\Phi^{\tau}(\mathbf{g}_{\theta}U)\mathbf{g}_{\theta}$ and $\mathbf{g}_{\theta}\Phi(U)$ solve the same equation, thus coinciding.

The following lemma ensures that the flow map $\Phi^{\tau}(U)$ generated by $G(\tau, U)$ is an admissible transformation for any $\tau \in [0, 1]$.

Lemma 2.15. Let $\Phi^{\tau}(U)$ be the flow map in (2.52). Then:

- (i) if $G(\tau, U) = \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\frac{\beta(U;x)}{1 + \tau \beta_x(U;x)} \mathbf{i} \xi \right)$ then $\Phi^{\tau}(U)$ is a 2-admissible transformation; (ii) if $G(\tau, U) = \operatorname{Op}_{\mathsf{out}}^{BW}(g(U; x, \xi))$ then $\Phi^{\tau}(U)$ is a 0-admissible transformation.

Proof. Along the proof we put m = 2 if $G(\tau, U)$ is as in (i), and m = 0 in case (ii).

It is classical that $\Phi^{\tau}(U)$ is a matrix of 0-operators in $\mathcal{M}^0_{\geq 0}[r]$ as well as its linear inverse, see e.g. Lemma 3.16 of [14]. We prove now the expansion (2.42); first expand

$$G(\tau, U) = G_2(U) + G_{\geq 4}(\tau, U) = \begin{cases} \operatorname{Op}_{\mathtt{vec}}^{BW}(\beta(U; x) \mathrm{i}\xi) + G_{\geq 4}(\tau, U), \\ \operatorname{Op}_{\mathtt{out}}^{BW}(g(U; x, \xi)) \end{cases} , \qquad (2.53)$$

so the expansion of $\Phi^{\tau}(U)$ reads

$$\Phi^{\tau}(U) = \mathrm{Id} + \tau G_2(U) + \mathbf{F}_{\geq 4}(\tau, U), \quad \mathbf{F}_{\geq 4}(\tau, U) \in \mathcal{M}^m_{\geq 4}[r]$$

We prove now (*iii*). The differential $d_U \Phi^{\varsigma}(U)[\widehat{U}]$ fulfills the variational equation

$$\begin{cases} \partial_{\varsigma} \mathrm{d}_U \Phi^{\varsigma}(U)[\widehat{U}] = G(\varsigma, U) \mathrm{d}_U \Phi^{\varsigma}(U)[\widehat{U}] + \mathrm{d}_U G(\varsigma, U)[\widehat{U}] \Phi^{\varsigma}(U) \\ \mathrm{d}_U \Phi^0(U)[\widehat{U}] = 0 \end{cases}$$

whose solution is given by the Duhamel formula

$$d_{U}\Phi^{\varsigma}(U)[\widehat{U}] = \Phi^{\varsigma}(U) \int_{0}^{\varsigma} \Phi^{\tau}(U)^{-1} d_{U}G(\tau, U)[\widehat{U}] \Phi^{\tau}(U) d\tau$$

$$\stackrel{(2.53),(2.52)}{=} \varsigma d_{U}G_{2}(U)[\widehat{U}] + \varsigma \int_{0}^{\varsigma} G(\theta, U)\Phi^{\theta}(U)d_{U}G_{2}(U)[\widehat{U}] d\theta + \Phi^{\varsigma}(U) \int_{0}^{\varsigma} d_{U}G_{\geq 4}(\tau, U)[\widehat{U}] d\tau$$

$$+ \Phi^{\varsigma}(U) \int_{0}^{\varsigma} \int_{0}^{\tau} \Phi^{\theta}(U)^{-1} \left[d_{U}G(\tau, U)[\widehat{U}], G(\theta, U) \right] \Phi^{\theta}(U) d\theta d\tau \qquad (2.54)$$

where in the second equality we also used the expansion

$$\Phi^{\theta}(U)^{-1} d_U G(\tau, U)[\widehat{U}] \Phi^{\theta}(U)|_{\theta=\tau} = d_U G(\tau, U)[\widehat{U}] + \int_0^{\tau} \Phi^{\theta}(U)^{-1} \left[d_U G(\tau, U)[\widehat{U}], G(\theta, U) \right] \Phi^{\theta}(U) d\theta.$$

We claim that, for both choices of $G(\tau, U)$ in (2.52),

$$\sup_{\tau \in [0,1]} \| \mathbf{d}_U G(\tau, U)[\widehat{U}] W \|_{s-\frac{m}{2}} \lesssim \| U \|_{s_0} \| \widehat{U} \|_{s_0} \| W \|_{s},$$

$$\sup_{\tau \in [0,1]} \| \mathbf{d}_U G_{\geq 4}(\tau, U)[\widehat{U}] W \|_{s-\frac{m}{2}} \lesssim \| U \|_{s_0}^3 \| \widehat{U} \|_{s_0} \| W \|_{s}, \quad \forall s \in \mathbb{R}.$$

$$(2.55)$$

Inserting these estimates in (2.54) and using (2.32) for $\Phi^{\varsigma}(U)$ and (2.29) for $G(\tau, U)$, one checks that the term $\left(\mathrm{d}_U \Phi^{\tau}(U)[\hat{U}] - \tau \mathrm{d}_U G_2(U)[\hat{U}] \right) W$ fulfills (2.43). This also shows that, for any $W \in$ $H^{s_0+m}_{\mathbb{R}}(\mathbb{T},\mathbb{C}^2)$, the map $U \mapsto \Phi^{\tau}(U)W$ is of class $C^0(B_{s_0,\mathbb{R}}(r);H^{s_0}_{\mathbb{R}}(\mathbb{T};\mathbb{C}^2))$, and so, using (2.54), it is also of class $C^1(B_{s_0,\mathbb{R}}(r); H^{s_0}_{\mathbb{R}}(\mathbb{T}; \mathbb{C}^2)).$

We now prove (2.55)–(2.56). Consider first $G(\tau, U) = \operatorname{Op}_{out}^{BW}(g(U; x, \xi))$, for which (2.56) is trivial $(G_{\geq 4}(\tau, U) \equiv 0)$. Since $g(U; \cdot)$ is homogeneous of degree 2,

$$d_U G(\tau, U)[\widehat{U}] = \operatorname{Op}_{\mathsf{out}}^{BW} \left(2g(\widehat{U}, U; x, \xi) \right) = d_U G_2(U)[\widehat{U}]$$

and (2.55) follows from Theorem 2.3. Next we analyze the case $G(\tau, U) = \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\frac{\beta(U;x)}{1 + \tau \beta_{\tau}(U;x)} \mathrm{i}\xi \right)$. Its differential is given by

$$d_U G(\tau, U)[\widehat{U}] = 2 \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\beta(\widehat{U}, U; x) \mathrm{i}\xi \right) - \tau \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\frac{\beta(\widehat{U}, U; x)\beta_x(U; x)}{1 + \tau\beta_x(U; x)} - \tau \frac{\beta(U; x)\beta_x(\widehat{U}, U; x)}{(1 + \tau\beta_x(U; x))^2} \mathrm{i}\xi \right)$$
$$= d_U G_2(U)[\widehat{U}] + d_U G_{\geq 4}(\tau, U)[\widehat{U}]$$

3 ANALYSIS OF WEAK RESONANCES

Now notice that $\beta(\widehat{U}, U; x) \in \widetilde{\mathcal{F}}_2^{\mathbb{R}}$ and $\frac{\beta(\widehat{U}, U; x)\beta_x(U; x)}{1 + \tau \beta_x(U; x)} + \frac{\beta(U; x)\beta_x(\widehat{U}, U; x)}{(1 + \tau \beta_x(U; x))^2} \in L^{\infty}(\mathbb{T}; \mathbb{R})$ with bound

$$\sup_{\tau \in [0,1]} \|\frac{\beta(\widehat{U}, U; x)\beta_x(U; x)}{1 + \tau\beta_x(U; x)} + \frac{\beta(U; x)\beta_x(\widehat{U}, U; x)}{(1 + \tau\beta_x(U; x))^2}\|_{L^{\infty}} \lesssim \|\widehat{U}\|_{s_0} \|U\|_{s_0}^3$$

Then Theorem 2.3 gives (2.55) and (2.56).

Next we consider the flow map generated by a matrix of smoothing operators:

$$\begin{cases} \partial_{\tau} \Phi^{\tau}(U) = R(U) \Phi^{\tau}(U) \\ \Phi^{0}(U) = \mathrm{Id} \end{cases} \quad \text{where} \quad R(U) \in \widetilde{\mathcal{R}}_{2}^{-\varrho} . \tag{2.57}$$

Lemma 2.16. Let $\varrho \geq 0$. The flow map $\Phi^{\tau}(U)$ in (2.57) is a 0-admissible transformation.

Proof. The proof follows on the same lines of the previous one. The algebraic expansion (2.54) holds with $G(\tau, U) \rightsquigarrow R(U)$ and, since $d_U R(U)[\hat{U}] = 2R(U, \hat{U})$, we replace (2.55) and (2.56) with estimate (2.35) with $m = -\rho$ to get the bound (2.43).

3 Analysis of weak resonances

Equation (1.1) is Hamiltonian, with Hamiltonian function given by

$$\mathscr{H}(u) := \int_{\mathbb{T}} (|D|^{\alpha} u) \overline{u} + \frac{\mathrm{i}}{4} \int_{\mathbb{T}} |u|^2 \left(\overline{u} \, u_x - u \overline{u}_x \right) \mathrm{d}x \; .$$

Due to the gauge and translation invariance of equation (1.1), any sufficiently regular solution u(t) of (1.1) conserves the total mass and momentum, namely

$$\mathcal{M}(u(t)) := \frac{1}{2\pi} \|u(t)\|_{L^2}^2 \equiv \frac{1}{2\pi} \int_{\mathbb{T}} |u(t,x)|^2 dx = \sum_{k \in \mathbb{Z}} |u_k(t)|^2 = \mathcal{M}(u(0)),$$

$$\mathcal{P}(u(t)) := \frac{1}{2\pi} \int_{\mathbb{T}} i(\partial_x u(t,x)) \overline{u(t,x)} \, dx = -\sum_{k \in \mathbb{Z}} k |u_k(t)|^2 = \mathcal{P}(u(0)) .$$

(3.1)

In view of this we introduce the new variable

$$v(t,x) := e^{it\mathscr{P}(u(t))}u(t,x - \mathscr{M}(u(t))t) .$$

Clearly v(t, x) and u(t, x) have same Sobolev norms, same magnitude, mass and the momentum, i.e.

$$\|v(t,\cdot)\|_s = \|u(t,\cdot)\|_s , \quad \forall s \in \mathbb{R}$$

and

$$|v(t,x)| = |u(t,x - \mathscr{M}(u(t))t)|, \quad \mathscr{M}(v(t)) = \mathscr{M}(u(t)), \quad \mathscr{P}(v(t)) = \mathscr{P}(u(t)) ,$$

and one readily checks that $v(t, \cdot)$ fulfills the re-normalized equation

$$\partial_t v = -\mathbf{i}|D|^{\alpha}v + |v|^2 v_x - \mathscr{M}(v)v_x + \mathbf{i}\mathscr{P}(v)v .$$
(3.2)

This is the equation that we shall consider from now on, and we will relabel $v \rightarrow u$. Also (3.2) is a Hamiltonian PDE with Hamiltonian function

$$\widetilde{\mathscr{H}}(v) := \mathscr{H}(v) - \mathscr{M}(v)\mathscr{P}(v)$$
.

Remark 3.1. The reason we renormalize equation (1.1) is that the vector field of (3.2) does not contain integrable resonant monomials of the form $|u_k|^2 u_\ell e^{i\ell x}$ with $k \neq \ell$. Although not strictly necessary, it simplifies the analysis of the resonant part of (3.2) in Lemma 3.4.

$$\mathsf{R} := \{ (\vec{j}, \vec{\sigma}) \in \mathcal{P}_4 : \quad \sigma_1 |j_1|^{\alpha} + \sigma_2 |j_2|^{\alpha} + \sigma_3 |j_3|^{\alpha} + \sigma_4 |j_4|^{\alpha} = 0 \} \quad .$$
(3.3)

When $\alpha \in (0, 1)$ is irrational, one can expect the set R to contain only integrable resonances, namely indexes of the form $((k, k, \ell, \ell), (+, -, +, -))$ with $k, \ell \in \mathbb{Z}$ and their permutations. For α rational, instead, nonintegrable resonances do exist in general: for example, when $\alpha = \frac{1}{2}$, one has the non-integrable Zakharov-Dyachenko resonances [69]. We do not care if such non-integrable resonances exist or not, since, as we discussed in the introduction, our energy cascades will be due to *quasi-resonances*, rather than exact resonances. What we really are interested in, is to study the resonances between frequencies in a fixed set Λ and those in its complementary set, with at most two frequencies in Λ^c .

We shall now study resonant sets with indexes constrained to belong to certain subsets.

Definition 3.2. Given a set $\Lambda \subseteq \mathbb{Z}$ and $n \in \{0, \ldots, 4\}$, we denote by $\mathcal{P}_{\Lambda}^{(n)}$ the elements of \mathcal{P}_4 (see (2.8)) having exactly n indexes outside the set Λ :

$$\mathcal{P}_{\Lambda}^{(n)} := \{ (j_1, j_2, j_3, j_4, \vec{\sigma}) \in \mathcal{P}_4 : \text{ exactly } n \text{ indexes among } j_1, j_2, j_3, j_4 \text{ are outside } \Lambda \} .$$
(3.4)

We denote by $\mathsf{R}^{(n)}_{\Lambda}$ the subset of $\mathcal{P}^{(n)}_{\Lambda}$ made of 4-waves resonances: with **R** in (3.3),

$$\mathsf{R}^{(n)}_{\Lambda} := \{ (j_1, j_2, j_3, j_4, \vec{\sigma}) \in \mathsf{R} : exactly \ n \ indexes \ among \ j_1, j_2, j_3, j_4 \ are \ outside \ \Lambda \} \ .$$
(3.5)

We shall now study in detail the sets $\mathsf{R}^{(n)}_{\Lambda}$, n = 0, 1, 2, when Λ is given by

$$\Lambda := \{-1, +1\} . \tag{3.6}$$

Lemma 3.3. Let Λ in (3.6) and $\mathcal{P}_{\Lambda}^{(n)}$, $\mathsf{R}_{\Lambda}^{(n)}$ defined in (3.4) and (3.5).

(i) The set $\mathcal{P}^{(0)}_{\Lambda} \equiv \mathsf{R}^{(0)}_{\Lambda}$ and it contains only integrable resonances:

$$\mathsf{R}^{(0)}_{\Lambda} = \{ (\pi(\mathsf{k}, \mathsf{k}, \ell, \ell), \ \pi(+, -, +, -)), : \mathsf{k}, \ell \in \Lambda \ , \ \pi \in \mathcal{S}_4 \}$$
(3.7)

and S_4 is the symmetric group of permutations of four symbols.

(ii) The set $\mathsf{R}^{(1)}_{\Lambda} = \emptyset$. Moreover $\mathcal{P}^{(1)}_{\Lambda}$ has finite cardinality and there exists c > 0 such that

$$(\vec{j},\vec{\sigma}) \in \mathcal{P}_{\Lambda}^{(1)} \quad \Rightarrow \quad |\sigma_1|j_1|^{\alpha} + \sigma_2|j_2|^{\alpha} + \sigma_3|j_3|^{\alpha} + \sigma_4|j_4|^{\alpha}| \ge c .$$

$$(3.8)$$

(iii) The set

$$\mathsf{R}^{(2)}_{\Lambda} = \{ (\pi(\mathtt{k}, \mathtt{k}, \ell, \ell), \ \pi(+, -, +, -)) \colon \ \mathtt{k} \in \Lambda, \ \ell \in \Lambda^c, \ \pi \in \mathcal{S}_4 \} .$$
(3.9)

Moreover there exists c > 0 such that

$$(\vec{j},\vec{\sigma}) \in \mathcal{P}_{\Lambda}^{(2)} \setminus \mathsf{R}_{\Lambda}^{(2)} \quad \Rightarrow \quad |\sigma_1|j_1|^{\alpha} + \sigma_2|j_2|^{\alpha} + \sigma_3|j_3|^{\alpha} + \sigma_4|j_4|^{\alpha}| \ge \frac{c}{\max_{a=1,\dots,4} (|j_a|)^{1-\alpha}} .$$
(3.10)

Proof. The gauge condition $\sum_{a=1}^{4} \sigma_a = 0$ implies that exactly two σ_a 's are +, the other are -. So, up to permutation, we can always assume that $\sigma_1 = \sigma_3 = 1$ and $\sigma_2 = \sigma_4 = -1$.

(*i*) In this case all indexes $j_1, j_2, j_3, j_4 \in \Lambda$, so automatically $|j_1|^{\alpha} - |j_2|^{\alpha} + |j_3|^{\alpha} - |j_4|^{\alpha} = 0$, so $\mathcal{P}^{(0)}_{\Lambda} = \mathsf{R}^{(0)}_{\Lambda}$. Next the momentum condition $j_1 - j_2 + j_3 - j_4 = 0$ gives that either $j_1 = j_2 = \mathtt{k}, j_3 = j_4 = \ell$, yielding $((\mathtt{k}, \mathtt{k}, \ell, \ell), (+, -, +, -))$, or $j_1 = j_4 = \mathtt{k}, j_2 = j_3 = \ell$, yielding $((\mathtt{k}, \ell, \ell, \mathtt{k}), (+, -, +, -))$, which is a permutation of the previous one.

3 ANALYSIS OF WEAK RESONANCES

(ii) We can always assume that $j_1, j_2, j_3 \in \Lambda$ and $j_4 \in \Lambda^c$. Then the resonant condition $|j_1|^{\alpha}$ – $|j_2|^{\alpha} + |j_3|^{\alpha} - |j_4|^{\alpha}$ reduces to $|j_3|^{\alpha} - |j_4|^{\alpha}$, for which we have the lower bound

$$||j_3|^{\alpha} - |j_4|^{\alpha}| \ge \begin{cases} 2^{\alpha} - 1, & \text{if } |j_4| \ge 2, \\ 1, & \text{if } j_4 = 0 \end{cases}$$

This proves both $\mathsf{R}^{(1)}_{\Lambda} = \emptyset$ and (3.8). (*iii*) We have two different cases.

<u>Case I</u>: W.l.o.g. assume $j_1, j_3 \in \Lambda$, $j_2, j_4 \in \Lambda^c$. The momentum condition reads $j_1 + j_3 = j_2 + j_4$. We examine further subcases.

• If $j_2 = j_4 = 0$, then $||j_1|^{\alpha} - |j_2|^{\alpha} + |j_3|^{\alpha} - |j_4|^{\alpha}| = 2$.

• If $j_2 = 0$ and $j_4 \neq 0$, from the momentum condition we get $|j_4| \leq 2$, so actually $j_4 = \pm 2$. Then $||j_1|^{\alpha} - |j_2|^{\alpha} + |j_3|^{\alpha} - |j_4|^{\alpha}| = 2 - 2^{\alpha} > 0.$

• If $j_2, j_4 \neq 0$, then $|j_2|, |j_4| \ge 2$. Then $||j_1|^{\alpha} - |j_2|^{\alpha} + |j_3|^{\alpha} - |j_4|^{\alpha}| \ge 2(2^{\alpha} - 1) > 0$.

Hence in Case I there are no resonances and the lower bound (3.10) holds.

<u>Case II</u>: W.l.o.g. assume that $j_1, j_2 \in \Lambda$, $j_3, j_4 \in \Lambda^c$. The momentum condition reads $j_1 - j_2 =$ $j_4 - j_3$. Again we examine further subcases.

• If $j_1 = j_2 = k \in \Lambda$, then, by the momentum, $j_3 = j_4 = \ell \in \Lambda^c$ and they form an element of $\mathsf{R}^{(2)}_{\Lambda}$. All other cases in (3.9) are obtained by permutations.

• If $j_1 \neq j_2$, then $j_4 = j_3 \pm 2$. Consider the "+" case, the other being analogous. The term $||j_1|^{\alpha} - |j_2|^{\alpha} + |j_3|^{\alpha} - |j_4|^{\alpha}|$ reduces to

$$||j_{3}+2|^{\alpha}-|j_{3}|^{\alpha}| \geq \begin{cases} 2^{\alpha} & \text{if } j_{3}=0 \text{ or } j_{3}=-2\\ \frac{4^{\alpha}-2^{\alpha}}{\max\left(|j_{3}|, |j_{3}+2|\right)^{1-\alpha}} & \text{if } |j_{3}| \geq 3 \end{cases}$$

proving (3.10).

Projection of cubic vector fields. We introduce now projections of cubic vector fields on the sets $\mathcal{P}^{(n)}_{\Lambda}$ and $\mathsf{R}^{(n)}_{\Lambda}$. Recall that any real-to-real cubic vector field X(U), translation and gauge invariant, expand in Fourier as (see (2.12))

$$X(U)^{\sigma} = \sum_{(\vec{j},k,\vec{\sigma},-\sigma)\in\mathcal{P}_4} X^{\sigma_1,\sigma_2,\sigma_3,\sigma}_{j_1,j_2,j_3,k} u^{\sigma_1}_{j_1} u^{\sigma_2}_{j_2} u^{\sigma_3}_{j_3} e^{\mathrm{i}\sigma kx} , \quad X^{\sigma_{\pi(1)},\dots,\sigma_{\pi(3)},\sigma}_{j_{\pi(1)},\dots,j_{\pi(3)},k} = X^{\sigma_1,\dots,\sigma_3,\sigma}_{j_1,\dots,j_3,k}$$
(3.11)

for any permutation π of $\{1,2,3\}$. Given a subset $A \subseteq \mathcal{P}_4$, we denote by $\prod_A X$ the vector field obtained restricting the indexes to belong to A, namely

$$(\Pi_A X)(U)^{\sigma} := \sum_{(\vec{j},k,\vec{\sigma},-\sigma)\in A} X^{\sigma_1,\sigma_2,\sigma_3,\sigma}_{j_1,j_2,j_3,k} \ u^{\sigma_1}_{j_1} \ u^{\sigma_2}_{j_2} \ u^{\sigma_3}_{j_3} \ e^{\mathrm{i}\sigma kx} \ .$$
(3.12)

We now compute the projections of the cubic vector field in (3.2), that we denote by

$$X_3(U)^+ := |u|^2 u_x - \mathscr{M}(u)u_x + \mathrm{i}\mathscr{P}(u)u , \qquad (3.13)$$

on the sets $\mathsf{R}^{(n)}_{\Lambda}$ defined in (3.5) for n = 0, 1, 2.

Lemma 3.4. The cubic, translation and gauge invariant vector field $X_3(U)^+$ in (3.13) fulfills:

- (i) Structure: There exists a 2-homogeneous 1-operator $M^+_{\text{NLS}}(U) \in \widetilde{\mathcal{M}}_2^1$ such that $X_3(U)^+ =$ $M^+_{\rm NLS}(U)u;$
- (ii) **Resonances:** The projections of the vector field $X_3(U)^+$ on the sets $\mathsf{R}^{(n)}_{\Lambda}$, n = 0, 1, 2, defined in (3.5) are given by

$$(\Pi_{\mathsf{R}^{(0)}_{\Lambda}}X_{3})(U)^{+} = -\mathrm{i}|u_{1}|^{2}u_{1} e^{\mathrm{i}x} + \mathrm{i}|u_{-1}|^{2}u_{-1} e^{-\mathrm{i}x} ,$$

$$(\Pi_{\mathsf{R}^{(1)}_{\Lambda}}X_{3})(U)^{+} = 0 , \quad (\Pi_{\mathsf{R}^{(2)}_{\Lambda}}X_{3})(U)^{+} = 0 .$$
(3.14)

3 ANALYSIS OF WEAK RESONANCES

Proof. (i) Define $M^+_{\text{NLS}}(U)$ to be the operator

$$M^+_{\mathsf{NLS}}(U)v := \left(|u|^2 - \mathscr{M}(u)\right) \,\partial_x v + \mathrm{i}\mathscr{P}(u)v \,\,, \tag{3.15}$$

so that $M^+_{\mathsf{NLS}}(U)u = X_3(U)^+$. To prove that $M^+_{\mathsf{NLS}}(U) \in \widetilde{\mathcal{M}}_2^1$ we write it in Fourier as

$$M_{\text{NLS}}^{+}(U)v = \sum_{\substack{\sigma_{1}j_{1}+\sigma_{2}j_{2}+j=k\\\sigma_{1}+\sigma_{2}=0}} M_{j_{1},j_{2},j,k}^{\sigma_{1},\sigma_{2}} u_{j_{1}}^{\sigma_{1}} u_{j_{2}}^{\sigma_{2}} v_{j} e^{ikx},$$
$$M_{j_{1},j_{2},j,k}^{\sigma_{1},\sigma_{2}} := \begin{cases} \frac{i}{2}j & \text{if } j_{1} \neq j_{2}, \ j \neq k, \ \sigma_{1} \neq \sigma_{2} \\ -\frac{i}{2}j_{1} & \text{if } j_{1} = j_{2}, \ j = k \ , \ \sigma_{1} \neq \sigma_{2} \\ 0 & \text{otherwise} \end{cases}$$

The coefficients $M_{j_1,j_2,j,k}^{\sigma_1,\sigma_2}$ are symmetric in the first two indexes and fulfill (2.31) with m = 1 and $\mu = 0$.

(*ii*) As we shall compute the projectors using the definition (3.12), we need first to write $X_3(U)^+$ in the form (3.11). So expand $X_3(U)^+$ in (3.13) in Fourier, getting

$$X_3(U)^+ = \sum_{\substack{j_1 - j_2 + j_3 = k \\ j_1 \neq j_2}} i j_3 u_{j_1} \overline{u}_{j_2} u_{j_3} e^{ikx} - \sum_{\substack{j_1 = j_2, j_3 = k \\ j_1 = j_2, j_3 = k}} i j_2 |u_{j_2}|^2 u_{j_3} e^{ikx} .$$

$$X_{3}(U)^{+} = \sum_{\substack{j_{1}-j_{2}+j_{3}=k\\j_{1}\neq j_{2}}} ij_{3}u_{j_{1}}\overline{u}_{j_{2}}u_{j_{3}}e^{ikx} - \sum_{j_{1}=j_{2}, j_{3}=k} ij_{2}|u_{j_{2}}|^{2}u_{j_{3}}e^{ikx} = \sum_{(\vec{j},k,\vec{\sigma},-)\in\mathcal{P}_{4}} N_{\vec{j},k}^{\vec{\sigma},+}u_{\vec{j}}^{\vec{\sigma}}e^{ikx},$$

where

$$N_{j_1,j_2,j_3,k}^{\sigma_1,\sigma_2,\sigma_3,+} := \mathbf{i}(j_3\delta_{j_1\neq j_2} - j_2\delta_{j_1=j_2}\delta_{j_3=k})\delta_{(\sigma_1,\sigma_2,\sigma_3)=(+,-,+)} \ .$$

The coefficients of expansion (3.11) are obtained by symmetrization

$$X_{j_1,j_2,j_3,k}^{\sigma_1,\sigma_2,\sigma_3,+} = \frac{1}{6} \sum_{\pi \in \mathcal{S}_3} N_{j_{\pi(1)},j_{\pi(2)},j_{\pi(3)},k}^{\sigma_{\pi(1)},\sigma_{\pi(2)},\sigma_{\pi(3)},+}$$

yielding

$$X_{j_1,j_2,j_3,k}^{+,-,+,+} = \frac{i}{6} \left(j_3 \delta_{j_1 \neq j_2} + j_1 \delta_{j_3 \neq j_2} - j_2 (\delta_{j_1 = j_2} + \delta_{j_3 = j_2}) \right)$$
(3.16)

Projection on $\mathsf{R}^{(0)}_{\Lambda}$: We use the definition of projections in (3.12). In view of the characterization of $\mathsf{R}^{(0)}_{\Lambda}$ given in (3.7), we must consider only those monomials with indexes of the form $((k, k, \ell, \ell), (+, -, +, -))$ with $k, \ell \in \{\pm 1\}$ and their permutations. Once the last couple $(\ell, -)$ is fixed, than either $k = \ell$, giving the index $((\ell, \ell, \ell, \ell), (+, -, +, -))$ and its 3 permutations, or $k = -\ell$, giving $((-\ell, -\ell, \ell, \ell, \ell), (+, -, +, -))$ and its 6 permutations. Therefore we obtain

$$(\Pi_{\mathsf{R}^{(0)}_{\Lambda}}X_{3})(U)^{+} = \left(3X^{+,-,+,+}_{1,1,1,1}|u_{1}|^{2}u_{1} + 6X^{+,-,+,+}_{-1,-1,1,1}|u_{-1}|^{2}u_{1}\right)e^{\mathrm{i}x} \\ + \left(6X^{+,-,+,+}_{1,1,-1,-1}|u_{1}|^{2}u_{-1} + 3X^{+,-,+,+}_{-1,-1,-1,-1}|u_{-1}|^{2}u_{-1}\right)e^{-\mathrm{i}x} \\ \stackrel{(3.16)}{=} -\mathrm{i}|u_{1}|^{2}u_{1}e^{\mathrm{i}x} + \mathrm{i}|u_{-1}|^{2}u_{-1}e^{-\mathrm{i}x}$$

proving the first of (3.14).

Projection on $\mathsf{R}^{(1)}_{\Lambda}$: It is zero since $\mathsf{R}^{(1)}_{\Lambda} = \emptyset$ by Lemma 3.3 (*ii*).

<u>Projection on $\mathsf{R}^{(2)}_{\Lambda}$ </u>: In view of the characterization of $\mathsf{R}^{(2)}_{\Lambda}$ in (3.9), the monomials surviving the projection have indexes of the form $((k, k, \ell, \ell), (+, -, +, -))$ (and their permutations) with only

one among k and ℓ in Λ . Once the last index $(\ell, -)$ is fixed in either Λ or Λ^c , and k is fixed in the complementary set, there are 6 possible permutations. Hence we get

$$(\Pi_{\mathsf{R}^{(2)}_{\Lambda}}X_{3})(U)^{+} = \sum_{k \in \Lambda^{c}} 6X^{+,-,+,+}_{k,k,1,1} |u_{k}|^{2} u_{1} e^{\mathrm{i}x} + \sum_{k \in \Lambda^{c}} 6X^{+,-,+,+}_{k,k,k,-1,-1} |u_{k}|^{2} u_{-1} e^{-\mathrm{i}x} + \sum_{\ell \in \Lambda^{c}} \sum_{k=\pm 1} 6X^{+,-,+,+}_{k,k,\ell,\ell} |u_{k}|^{2} u_{\ell} e^{\mathrm{i}\ell x} \stackrel{(3.16)}{=} 0$$

proving the last of (3.14).

For later use, we prove a lemma about the projections on $\mathsf{R}^{(n)}_{\Lambda}$, n = 0, 1, 2, of cubic paradifferential vector fields. Precisely we have

Lemma 3.5. Let $a(Z; x, \xi)$ be a 2-homogeneous symbol in $\widetilde{\Gamma}_2^m$, $m \in \mathbb{R}$, with zero average and fulfilling $a(\mathbf{g}_{\theta}Z; \cdot) = a(Z; \cdot)$ for any $\theta \in \mathbb{T}$, where \mathbf{g}_{θ} in (2.2). Then

$$\Pi_{\mathsf{R}^{(n)}_{\Lambda}}\left[\operatorname{Op}_{\mathtt{vec}}^{\scriptscriptstyle BW}(a(Z;x,\xi))\,Z\right] = 0 \ , \quad n = 0, 1, 2 \ .$$

Proof. Recalling (2.26), $\left(\operatorname{Op}_{\mathsf{vec}}^{BW}(a(Z; x, \xi)) Z\right)^+ = \operatorname{Op}^{BW}(a(Z; x, \xi)) z$. Using definition (2.21) specialized to quadratic symbols fulfilling $a(\mathsf{g}_{\theta} Z; \cdot) = a(Z; \cdot), \forall \theta \in \mathbb{T}$, and the comments right below (2.25), we get

$$Op^{BW}(a(Z;x,\xi)) z = \sum_{j_1-j_2+j=k} \chi_2\left(j_1, j_2, \frac{j+k}{2}\right) a_{j_1,j_2}^{+,-}\left(\frac{j+k}{2}\right) z_{j_1}\overline{z}_{j_2} z_j e^{ikx} .$$

The point is that, when projecting on $\Pi_{\mathsf{R}^{(n)}_{\mathsf{A}}}$, n = 0, 1, 2, either the cut-off $\chi_2(\cdot, \cdot)$ or the coefficient $a_{j_1,j_2}^{+,-}$ vanish. Recall that $\chi_2(\xi',\xi) \equiv 0$ whenever $|\xi'| > \frac{\langle \xi \rangle}{10}$. <u>Case n = 0</u>: In this case $j_1, j_2, j, k \in \Lambda$, and $\chi_2\left(2, \frac{j+k}{2}\right) = 0$ for any choice of $j, k \in \{\pm 1\}$. <u>Case n = 1</u>: By Lemma 3.3 $\mathsf{R}^{(1)}_{\Lambda} = \emptyset$ and there is nothing to prove.

<u>Case n = 2</u>: By Lemma 3.3 the indexes j_1, j_2, j, k are pairwise equal. Assume first that $j_1 = j_2$, then $a_{j_1,j_1}^{+,-} = 0$ since $a(Z; \cdot)$ has zero-average in x. The case $j_1 = j \in \Lambda$ and $j_2 = k \in \Lambda^c$ violates the momentum conservation, as well as $j_1 = j \in \Lambda^c$,

 $j_2 = k \in \Lambda.$

In case $j_1 = k \in \Lambda$ and $j_2 = j \in \Lambda^c$, the cut-off vanishes since

$$\chi_2\left(\pm 1, j, \frac{j+k}{2}\right) \equiv 0 \quad \forall k \in \Lambda, \ j \in \Lambda^c$$

Analogously the case $j_1 = k \in \Lambda^c$, $j_2 = j \in \Lambda$ is ruled out, concluding the proof.

Identification argument. We prove an abstract identification argument in the spirit of [11, 10]. In section 4 we shall conjugate equation (3.2) with an admissible transformation. Without doing explicit computations, we shall a posteriori identify the explicit form of the resonant parts of the conjugated vector field thanks to the following proposition.

Proposition 3.6 (Identification of the resonant normal form). Let $\mathbf{F}(U)$ be a 2-admissible transformation (see Definition 2.10). There exist $r, s_0 > 0$ such that, provided $U(t) \in B_{s_0,\mathbb{R}}(I;r)$ is a solution of the system

$$\partial_t U = -\mathbf{i} \mathbf{\Omega}(D) U + X_3(U), \qquad \mathbf{\Omega}(D) := \begin{pmatrix} |D|^\alpha & 0\\ 0 & -|D|^\alpha \end{pmatrix}$$
(3.17)

where

$$X_3(U) = M_2(U)U , \quad M_2(U) \ a \ matrix \ of \ operators \ in \ \mathcal{M}_2^1 , \qquad (3.18)$$

3 ANALYSIS OF WEAK RESONANCES

then the variable $Z := \mathcal{F}(U) = \mathbf{F}(U)U$ solves

$$\partial_t Z = -\mathrm{i}\mathbf{\Omega}(D)Z + X_3(Z) + M_{\geq 4}(Z)Z . \qquad (3.19)$$

Here $\widetilde{M}_{\geq 4}(Z)$ is a matrix of non-homogeneous 7-operators in $\mathcal{M}_{\geq 4}^{7}[r]$, whereas $\widetilde{X}_{3}(Z)$ is a cubic vector field fulfilling

$$\Pi_{\mathsf{A}} X_3 = \Pi_{\mathsf{A}} X_3, \quad for \ any \ \mathsf{A} \subseteq \mathsf{R} \tag{3.20}$$

where R is the 4-waves resonant set in (3.3).

Proof. Defining $X(U) := -i\Omega(D)U + X_3(U)$, the variable Z solves the equation

$$\partial_t Z = \mathcal{F}^* X(Z) := \mathrm{d}_U \mathcal{F}(U) \left[X(U) \right]_{|U = \mathcal{F}^{-1}(Z)},$$

where to invert the nonlinear map \mathcal{F} we used Lemma 2.13.

Next we provide a Taylor expansion of the push-forward vector field \mathcal{F}^*X . Using the expansion (2.42) for $\mathcal{F}(U) = \mathbf{F}(U)U$, we get

$$d_{U}\mathcal{F}(U)\left[X(U)\right]$$

$$= -i\mathbf{\Omega}(D)U + X_{3}(U) + \mathbf{F}_{2}(U)\left[-i\mathbf{\Omega}(D)U\right] + d_{U}\mathbf{F}_{2}(U)\left[-i\mathbf{\Omega}(D)U\right]U + M_{\geq 4}(U)U$$

$$(3.21)$$

where, using the structure (3.18) of $X_3(U)$

$$M_{\geq 4}(U)W := -\mathbf{F}_{\geq 4}(U)\mathbf{i}\mathbf{\Omega}(D)W + \mathbf{F}_{\geq 4}(U)M_2(U)W + \mathbf{d}_U\mathbf{F}_{\geq 4}(U)[X(U)]W + \mathbf{F}_2(U)M_2(U)W + \mathbf{d}_U\mathbf{F}_2(U)[X_3(U)]W.$$
(3.22)

We prove in Lemma 3.7 below that $M_{\geq 4}(U)$ is a matrix of non-homogeneous operators in $\mathcal{M}^3_{\geq 4}[r]$. Next we compute (3.21) at

$$U = \mathcal{F}^{-1}(Z) \stackrel{(2.48)}{=} \mathbf{G}(Z)Z, \qquad \mathbf{G}(Z) - \mathrm{Id} = \underbrace{-\mathbf{F}_2(Z) + \mathbf{G}_{\geq 4}(Z)}_{=:\mathbf{G}_{\geq 2}(Z)} \in \Sigma \mathcal{M}_2^4[r], \qquad (3.23)$$

obtaining

$$\mathcal{F}^*X(Z) = -\mathrm{i}\Omega(D)Z + \widetilde{X}_3(Z) + \widetilde{M}_{\geq 4}(Z)Z$$

where

$$\widetilde{X}_{3}(Z) := X_{3}(Z) + \llbracket \mathbf{F}_{2}(Z)Z, -\mathbf{i}\mathbf{\Omega}(D)Z \rrbracket$$

$$\llbracket \mathbf{F}_{2}(Z)Z, -\mathbf{i}\mathbf{\Omega}(D)Z \rrbracket := \mathbf{i}\mathbf{\Omega}(D)\mathbf{F}_{2}(Z)Z + \mathbf{F}_{2}(Z)[-\mathbf{i}\mathbf{\Omega}(D)Z] + \mathbf{d}_{Z}\mathbf{F}_{2}(Z)[-\mathbf{i}\mathbf{\Omega}(D)Z]Z$$

$$(3.24)$$

and

$$\widetilde{M}_{\geq 4}(Z)W = -i\mathbf{\Omega}(D)\mathbf{G}_{\geq 4}(Z)W + [M_2(\mathcal{F}^{-1}(Z))\mathbf{G}(Z) - M_2(Z)]W - [\mathbf{F}_2(\mathcal{F}^{-1}(Z))i\mathbf{\Omega}(D)\mathbf{G}(Z) - \mathbf{F}_2(Z)i\mathbf{\Omega}(D)]W - [\mathbf{d}_U\mathbf{F}_2(\mathcal{F}^{-1}(Z))[i\mathbf{\Omega}(D)\mathcal{F}^{-1}(Z)]\mathbf{G}(Z) - \mathbf{d}_Z\mathbf{F}_2(Z)[i\mathbf{\Omega}(D)Z]]W + M_{\geq 4}(\mathcal{F}^{-1}(Z))\mathbf{G}(Z)W$$
(3.25)

We prove in Lemma 3.7 below that $\widetilde{M}_{\geq 4}(Z)$ belongs to $\mathcal{M}_{\geq 4}^{7}[r]$. This concludes the proof of (3.19). To prove (3.20) we note that

$$\llbracket \mathbf{F}_2(Z)Z, -\mathrm{i}\mathbf{\Omega}(D)Z \rrbracket^{\sigma} = \sum_{(\vec{j},k,\vec{\sigma},-\sigma)\in\mathcal{P}_4} -\mathrm{i}\left(\sigma_1|j_1|^{\alpha} + \sigma_2|j_2|^{\alpha} + \sigma_3|j_3|^{\alpha} - \sigma|k|^{\alpha}\right) \mathbf{F}_{\vec{j},k}^{\vec{\sigma},\sigma} z_{\vec{j}}^{\vec{\sigma}} e^{\mathrm{i}\sigma kx} ;$$

it then follows that, for any set $\mathsf{A}\subseteq\mathsf{R},$ one has

$$\Pi_{\mathsf{A}}\left[\!\left[\mathbf{F}_{2}(Z)Z\,,\,-\mathrm{i}\mathbf{\Omega}(D)Z\right]\!\right]\equiv0$$

which, together with (3.24), implies (3.20).

Lemma 3.7. There is r > 0 such that $M_{\geq 4}(U)$ defined in (3.22) is a matrix of 3-operators in $\mathcal{M}^3_{\geq 4}[r]$ and $\widetilde{M}_{\geq 4}(Z)$ defined in (3.25) is a matrix of 7-operators in $\mathcal{M}^7_{\geq 4}[r]$.

Proof. We need to show that each term in (3.22) and (3.25) fulfills (2.32) with p = 4, some $s_0 \ge 0$ and m equal 3 or 7. This is proved exploiting that each term is a composition of either m-operators or differentials of admissible transformations and therefore satisfying (2.43). As an example, we explicitly show how to bound the most difficult terms in (3.22) and (3.25). Recall that, by definition of admissible transformations, $\mathbf{F}(U)$ – Id is a matrix of 2-operators in $\Sigma \mathcal{M}_2^2[r]$ for some r > 0.

We start from $d_U \mathbf{F}_{\geq 4}(U)[X(U)]W$ in (3.22). Using (2.43) (with $s \rightsquigarrow s-1$ and m=2) and that $||X(U)||_{s-1} \leq ||U||_s$, we get

$$\begin{aligned} \| \mathbf{d}_U \mathbf{F}_{\geq 4}(U)[X(U)]W\|_{s-3} \lesssim & \|U\|_{s_0}^3 \|X(U)\|_{s_0} \|W\|_{s-1} + \|U\|_{s_0}^3 \|X(U)\|_{s-1} \|W\|_{s_0} \\ &+ \|U\|_{s_0}^2 \|U\|_{s-1} \|X(U)\|_{s_0} \|W\|_{s_0} \\ \lesssim & \|U\|_{s_0+1}^4 \|W\|_s + \|U\|_{s_0+1}^3 \|U\|_s \|W\|_{s_0+1} \end{aligned}$$

proving (2.32) with $s_0 \rightsquigarrow s_0 + 1$.

Now we consider the term in the third line of (3.25). Using the trilinearity of $(V, V', W) \mapsto d_U \mathbf{F}_2(V)[V']W$ and (3.23) we decompose it as

$$\begin{aligned} \left[\mathrm{d}_{U} \mathbf{F}_{2}(\mathcal{F}^{-1}(Z)) [\mathrm{i} \mathbf{\Omega}(D) \mathcal{F}^{-1}(Z)] \mathbf{G}(Z) - \mathrm{d}_{Z} \mathbf{F}_{2}(Z) [\mathrm{i} \mathbf{\Omega}(D) Z] \right] W \\ &= \mathrm{d}_{U} \mathbf{F}_{2}\left(\mathbf{G}_{\geq 2}(Z) Z \right) [\mathrm{i} \mathbf{\Omega}(D) \mathcal{F}^{-1}(Z)] \mathbf{G}(Z) W + \mathrm{d}_{U} \mathbf{F}_{2}(Z) [\mathrm{i} \mathbf{\Omega}(D) \mathcal{F}^{-1}(Z)] \mathbf{G}_{\geq 2}(Z) W \\ &+ \mathrm{d}_{U} \mathbf{F}_{2}(Z) [\mathrm{i} \mathbf{\Omega}(D) \mathbf{G}_{\geq 2}(Z)] W \end{aligned}$$
(3.26)

We bound each term in (3.26) separately. We shall repeatedly use that $\|\Omega(D)U\|_{s-\alpha} \leq \|U\|_s$. First, using (2.46) and then (2.35), (2.32), (2.51) and (3.23), we get

$$\begin{aligned} \| \mathbf{d}_{U} \mathbf{F}_{2} \left(\mathbf{G}_{\geq 2}(Z) Z \right) [\mathbf{i} \mathbf{\Omega}(D) \mathcal{F}^{-1}(Z)] \mathbf{G}(Z) W \|_{s-7} \\ \lesssim \| \mathbf{G}_{\geq 2}(Z) Z \|_{s_{0}} \| \mathbf{\Omega}(D) \mathcal{F}^{-1}(Z) \|_{s_{0}} \| \mathbf{G}(Z) W \|_{s-5} + \| \mathbf{G}_{\geq 2}(Z) Z \|_{s_{0}} \| \mathbf{\Omega}(D) \mathcal{F}^{-1}(Z) \|_{s-5} \| \mathbf{G}(Z) W \|_{s_{0}} \\ + \| \mathbf{G}_{\geq 2}(Z) Z \|_{s-5} \| \mathbf{\Omega}(D) \mathcal{F}^{-1}(Z) \|_{s_{0}} \| \mathbf{G}(Z) W \|_{s_{0}} \\ \lesssim \| Z \|_{s_{0}+4}^{4} \| W \|_{s} + \| Z \|_{s_{0}+4}^{3} \| Z \|_{s} \| W \|_{s_{0}+4}. \end{aligned}$$

$$(3.27)$$

Similarly one obtains

$$\|\mathbf{d}_U \mathbf{F}_2(Z) [\mathbf{i} \mathbf{\Omega}(D) \mathcal{F}^{-1}(Z)] \mathbf{G}_{\geq 2}(Z) W\|_{s-7} \lesssim \|Z\|_{s_0+4}^4 \|W\|_s + \|Z\|_{s_0+4}^3 \|Z\|_s \|W\|_{s_0+4}.$$
(3.28)

Finally, using (2.46) and then (2.35), (2.32) and (3.23), we get

$$\begin{aligned} |\mathbf{d}_{U}\mathbf{F}_{2}(Z)\left[\mathrm{i}\mathbf{\Omega}(D)\mathbf{G}_{\geq 2}(Z)Z\right]W\|_{s-7} \\ \lesssim \|Z\|_{s_{0}}\|\mathbf{\Omega}(D)\mathbf{G}_{\geq 2}(Z)Z\|_{s_{0}}\|W\|_{s-5} + \|Z\|_{s_{0}}\|\mathbf{\Omega}(D)\mathbf{G}_{\geq 2}(Z)Z\|_{s-5}\|W\|_{s_{0}} \\ &+ \|Z\|_{s-5}\|\mathbf{\Omega}(D)\mathbf{G}_{\geq 2}(Z)Z\|_{s_{0}}\|W\|_{s_{0}} \\ \lesssim \|Z\|_{s_{0}+5}^{4}\|W\|_{s} + \|Z\|_{s_{0}+5}^{3}\|Z\|_{s}\|W\|_{s_{0}+5}. \end{aligned}$$
(3.29)

Estimates (3.27), (3.28) and (3.29) prove that the operator in (3.26) is a non-homogeneous 7-operator in $\mathcal{M}_{\geq 4}^7[r]$.

4 Paradifferential normal form

The goal of this section is to use paradifferential transformations and Birkhoff normal forms, in the spirit of [9], to put the quasilinear equation (3.2) into a suitable normal form. However, the normal form that we shall obtain is rather different from the one of [9] and of [11, 10, 13, 60]; indeed, in these papers, the paradifferential part has symbols with constant coefficients (at least at low homogeneity), and the smoothing vector field is in Birkhoff normal form, namely supported only on resonant monomials. On the contrary, our normal form has to two important and different features,

4 PARADIFFERENTIAL NORMAL FORM

see Theorem 4.3: (i) the cubic part of the paradifferential vector field has a dominant transport term with *variable coefficients* and supported only on resonant sites, see (4.7), and (ii) the cubic smoothing vector field is in a suitable weak normal form, that we call Λ -normal form and we now introduce.

Definition 4.1 (A-normal form). Let $\Lambda = \{1, -1\}$ as in (3.6). A cubic, translation and gauge invariant vector field X(U) is said to be in

• weak- Λ normal form if all its monomials with at most two indexes outside Λ are resonant, i.e.

$$\Pi_{\mathcal{P}^{(n)}_\Lambda X} = \Pi_{\mathsf{R}^{(n)}_\Lambda X} \ , \quad n=0,1,2 \ ;$$

 strong-Λ normal form if in addition there are no resonant monomials with one or two indexes outside Λ, i.e.

$$\Pi_{\mathcal{P}^{(0)}_{\Lambda}} X = \Pi_{\mathsf{R}^{(0)}_{\Lambda}} X , \quad \Pi_{\mathcal{P}^{(1)}_{\Lambda}} X = \Pi_{\mathcal{P}^{(2)}_{\Lambda}} X = 0 ,$$

the sets $\mathcal{P}^{(n)}_{\Lambda}$, $\mathsf{R}^{(n)}_{\Lambda}$ being defined in (3.4) and (3.5).

Note that a cubic vector field in *strong-* Λ *normal form* is composed by monomials $u_{j_1}^{\sigma_1} u_{j_2}^{\sigma_2} u_{j_3}^{\sigma_3} e^{i\sigma kx}$ whose indexes $((j_1, j_2, j_3, k), (\sigma_1, \sigma_2, \sigma_3, -\sigma))$ are

- either in Λ and resonant, i.e. $((j_1, j_2, j_3, k), (\sigma_1, \sigma_2, \sigma_3, -\sigma)) \in \mathsf{R}^{(0)}_{\Lambda};$
- or at least three indexes are outside Λ , i.e. $((j_1, j_2, j_3, k), (\sigma_1, \sigma_2, \sigma_3, -\sigma)) \in \mathcal{P}_{\Lambda}^{(3)} \cup \mathcal{P}_{\Lambda}^{(4)}$.

To start the normal form procedure, it is convenient to write (3.2) as the system in the variable $U := \left(\frac{u}{u}\right)$ given by

$$\partial_t U = -\mathrm{i}\mathbf{\Omega}(D)U + X_3(U), \quad X_3(U) = \begin{pmatrix} |u|^2 u_x - \mathscr{M}(u)u_x + \mathrm{i}\mathscr{P}(u)u\\ |u|^2 \overline{u_x} - \mathscr{M}(u)\overline{u_x} - \mathrm{i}\mathscr{P}(u)\overline{u} \end{pmatrix}$$
(4.1)

where $\Omega(D)$ is defined in (3.17) and, with M_{NLS}^+ the 1-operator in $\widetilde{\mathcal{M}}_2^1$ in (3.15),

$$X_{3}(U) = M_{\text{NLS}}(U)U , \qquad M_{\text{NLS}}(U) := \begin{pmatrix} M_{\text{NLS}}^{+}(U) & 0\\ 0 & M_{\text{NLS}}^{+}(U) \end{pmatrix} .$$
(4.2)

The first step is to paralinearize such system.

Lemma 4.2 (Paralinearization). Fix $\rho \geq 0$ and $s_0 \geq \rho + 2$. If $u(t) \in H^{s_0}(\mathbb{T}, \mathbb{C})$ solves equation (3.2), then $U(t) = \begin{pmatrix} u(t) \\ \overline{u}(t) \end{pmatrix}$ solves the system in paradifferential form (recall the notation in (2.26))

$$\partial_t U = -\mathrm{i}\mathbf{\Omega}(D)U + \mathrm{Op}_{\mathsf{vec}}^{BW}(\mathrm{i}\,\underline{\mathsf{V}}(U;x)\xi + \mathrm{i}\,\underline{\mathsf{d}}(U;x))\,U + \mathrm{Op}_{\mathsf{out}}^{BW}(b(U;x))\,U + R_2(U)U \tag{4.3}$$

where:

- $\Omega(D)$ is the matrix of Fourier multipliers in (3.17);
- $V(U;x), \underline{d}(U;x), \in \widetilde{\mathcal{F}}_2^{\mathbb{R}}$ and $b(U;x) \in \widetilde{\mathcal{F}}_2$ are the zero-average, 2-homogeneous functions

$$\underline{\underline{V}}(U;x) := |u|^2 - \mathscr{M}(u) = \sum_{k_1 \neq k_2} u_{k_1} \overline{u}_{k_2} e^{i(k_1 - k_2)x},$$
(4.4)
$$\underline{d}(U;x) := \operatorname{Im}(u_x \overline{u}) - \mathscr{P}(u) = \operatorname{Im}\sum_{k_1 \neq k_2} i k_1 u_{k_1} \overline{u}_{k_2} e^{i(k_1 - k_2)x} ,$$

$$b(U;x) := u u_x = \sum_{k_1, k_2 \in \mathbb{Z}} i \frac{k_1 + k_2}{2} u_{k_1} u_{k_2} e^{i(k_1 + k_2)x},$$

where $\mathcal{M}(u)$, $\mathcal{P}(u)$ are the mass and momentum defined in (3.1);

• $R_2(U)$ is a real-to-real, gauge invariant matrix of smoothing operators in $\tilde{\mathcal{R}}_2^{-\varrho}$.

Proof. The nonlinearity $|u|^2 u_x$ is parallely a standard way using Lemma 2.8 and Proposition 2.7, getting a smoothing remainder R(U) whose coefficients fulfill (2.31) with $\mu \rightsquigarrow \rho+1$ and $m \rightsquigarrow -\rho$. Note also that, in view of the Bony quantization (2.20), (2.21) for homogeneous symbols

$$\mathscr{M}(u)u_x = \operatorname{Op}^{BW}(\mathscr{M}(u)\mathrm{i}\xi) u + R(U)u , \quad \mathscr{P}(u)u = \operatorname{Op}^{BW}(\mathscr{P}(u)) u + R(U)u$$

for some smoothing remainders in $\widetilde{\mathcal{R}}_2^{-\varrho}$. Finally, remark that equation (4.1) is real-to-real and gauge invariant. Since also the paradifferential operators in (4.3) are real-to-real and gauge invariant (see (2.24) and (2.25)), by difference so is the matrix of smoothing operators $R_2(U)$.

The main result of the section is the following normal form theorem.

Theorem 4.3. There exist $s_0, r > 0$ and a 2-admissible transformation $\mathbf{F}(U) \in \mathcal{M}^0_{\geq 0}[r]$ (see Definition 2.10) such that if $U(t) \in B_{s_0,\mathbb{R}}(I;r)$ solves (4.3) then the variable

$$Z := \mathcal{F}(U) := \mathbf{F}(U)U \quad solves \tag{4.5}$$

$$\partial_t Z = -\mathrm{i} \mathbf{\Omega}(D) Z + \mathrm{Op}_{\mathsf{vec}}^{BW} \left(\mathrm{i} \langle \underline{\Psi} \rangle(Z; x) \xi + \mathrm{i} a_2^{(\alpha)}(Z; x, \xi) \right) Z + R_2^{(\Lambda)}(Z) Z + \mathrm{Op}_{\mathsf{vec}}^{BW} \left(\mathrm{i} \widetilde{V}_{\geq 4}(Z; x) \xi + \mathrm{i} \widetilde{a}_{\geq 4}^{(\alpha)}(Z; x, \xi) \right) Z + \widetilde{B}_{\geq 4}(Z) Z$$

$$(4.6)$$

where:

- $\Omega(D)$ is the matrix of Fourier multipliers in (3.17);
- $\langle \underline{\mathbf{V}} \rangle(Z; x)$ is the zero-average, real valued function in $\widetilde{\mathcal{F}}_2^{\mathbb{R}}$ defined by

$$\langle \underline{\mathbf{V}} \rangle(Z;x) := 2 \operatorname{Re} \left(\sum_{n \in \mathbb{N}} z_n \, \overline{z_{-n}} \, e^{\mathrm{i} 2nx} \right) ;$$

$$(4.7)$$

- $a_2^{(\alpha)}(Z; x, \xi)$ is a zero average, gauge-invariant, real symbol in $\widetilde{\Gamma}_2^{\alpha}$;
- $\widetilde{V}_{\geq 4}(Z; x)$ is a real function in $\mathcal{F}_{\geq 4}^{\mathbb{R}}[r]$ and $\widetilde{a}_{\geq 4}^{(\alpha)}(Z; x, \xi)$ a real non-homogeneous symbol in $\Gamma_{\geq 4}^{\alpha}[r]$;

• $R_2^{(\Lambda)}(Z)$ is a real-to-real and gauge invariant matrix of smoothing operators in $\widetilde{\mathcal{R}}_2^{-4}$ such that the cubic vector field

$$X^{(\Lambda)}(Z) := R_2^{(\Lambda)}(Z)Z \tag{4.8}$$

is in strong- Λ normal form (see Definition 4.1). Precisely, with the notation in (3.12),

$$(\Pi_{\mathcal{P}^{(0)}_{\Lambda}}X^{(\Lambda)})(Z) = \begin{pmatrix} -i|z_{1}|^{2}z_{1} e^{ix} + i|z_{-1}|^{2}z_{-1} e^{-ix} \\ i|z_{1}|^{2}\overline{z_{1}} e^{-ix} - i|z_{-1}|^{2}\overline{z_{-1}} e^{ix} \end{pmatrix} , \qquad (4.9)$$
$$\Pi_{\mathcal{P}^{(1)}_{\Lambda}}X^{(\Lambda)} = \Pi_{\mathcal{P}^{(2)}_{\Lambda}}X^{(\Lambda)} = 0 .$$

• Finally $\widetilde{B}_{\geq 4}(Z)$ is a real-to-real matrix of 0-operators in $\mathcal{M}_{\geq 4}^0[r]$.

The rest of the section is devoted to the proof of Theorem 4.3.

4.1 Block diagonalization

The goal of this section is to remove the out-diagonal term $\operatorname{Op}_{out}^{BW}(b(U;x))$ from equation (4.3) up to quadratic smoothing operators and quartic bounded operators. Precisely we prove:

Proposition 4.4 (Block-diagonalization). Let $\varrho \geq 1 - \alpha$. There exist $s_0, r > 0$ and a 0-admissible transformation $\Psi(U) \in \mathcal{M}^0_{\geq 0}[r]$ (see Definition 2.10) such that if $U(t) \in B_{s_0,\mathbb{R}}(I;r)$ solves (4.3), then the variable

$$W := \Psi(U)U \quad solves \tag{4.10}$$

$$\partial_t W = -\mathrm{i}\mathbf{\Omega}(D)W + \mathrm{Op}_{\mathsf{vec}}^{BW}(\mathrm{i}\underline{V}(U;x)\xi + \mathrm{i}\underline{\mathbf{d}}(U;x))W + R_2(U)W + B_{\geq 4}(U)W$$
(4.11)

where:

- $\Omega(D)$ is the matrix of Fourier multipliers defined in (3.17);
- $\underline{V}(U;x)$ and $\underline{d}(U;x)$ are the zero average functions defined in (4.4);
- $R_2(U)$ is a real-to-real, gauge invariant matrix of homogeneous smoothing remainders in $\widetilde{\mathcal{R}}_2^{-\varrho}$;
- $B_{\geq 4}(U)$ is a real-to-real matrix of non-homogeneous bounded operators in $\mathcal{M}^{0}_{\geq 4}[r]$.

Proof. We define the map $\Psi(U)$ as the time-1 flow $\Psi(U) := \Psi^{\tau}(U)_{|\tau=1}$ of the paradifferential equation

$$\begin{cases} \partial_{\tau} \Psi^{\tau}(U) = G(U) \Psi^{\tau}(U) \\ \Psi^{0}(U) = \mathrm{Id}, \end{cases} \quad \text{where} \quad G(U) := \mathrm{Op}_{\mathsf{out}}^{^{BW}}(g_{2}(U; x, \xi)) \end{cases}$$

and with the 2-homogeneous symbol g_2 of the form

$$g_2(U; x, \xi) = \sum_{j_1, j_2 \in \mathbb{Z}} g_{j_1, j_2}(\xi) u_{j_1} u_{j_2} e^{\mathbf{i}(j_1 + j_2)x} \in \widetilde{\Gamma}_2^{-\alpha}$$
(4.12)

to be determined. By Lemma 2.15, $\Psi(U)$ is a 0-admissible transformation. Moreover, G is gauge invariant (see the bullet of formula (2.25)), so is Ψ^{τ} (Remark 2.14). The variable $W = \Psi(U)U$ solves

$$\partial_t W = \Psi(U) \operatorname{Op}_{\mathsf{vec}}^{BW}(-\mathrm{i}|\xi|^{\alpha} + \mathrm{i}\underline{\mathsf{V}}(U;x)\xi + \mathrm{i}\underline{\mathsf{d}}(U;x)) \Psi(U)^{-1}W$$
(4.13)

$$+\Psi(U) \left| \operatorname{Op}_{\mathsf{out}}^{BW}(b(U;x)) + R_2(U) \right| \Psi(U)^{-1}W$$
(4.14)

$$+ \left(\partial_t \Psi(U)\right) \Psi(U)^{-1} W . \tag{4.15}$$

We first expand (4.13). The Lie expansion formula (see e.g. Lemma A.1 of [11]) says that for any operator A(U), setting $\operatorname{Ad}_B[A] := [B, A]$, one has

$$\Psi(U)A(U)\Psi(U)^{-1} = A(U) + [G(U), A(U)] + \int_0^1 (1-\tau)\Psi^{\tau}(U)\operatorname{Ad}^2_{G(U)}[A(U)] (\Psi^{\tau}(U))^{-1} d\tau (4.16)$$

Applying this formula with $A = Op_{vec}^{BW}(-i|\xi|^{\alpha} + i\underline{V}\xi + i\underline{d})$, using formulas (2.41) we get

$$(4.13) = \operatorname{Op}_{\mathsf{vec}}^{BW}(-\mathrm{i}|\xi|^{\alpha} + \mathrm{i}\underline{\mathsf{V}}\xi + \mathrm{i}\underline{\mathsf{d}})W + \operatorname{Op}_{\mathsf{out}}^{BW}(\mathrm{i}(g_{2}\#_{\varrho}|\xi|^{\alpha} + |\xi|^{\alpha}\#_{\varrho}g_{2}))W + R_{2}'(U)W + B_{\geq 4}(U)W$$

where R'_2 is a matrix of smoothing remainders in $\widetilde{\mathcal{R}}_2^{-\varrho}$ (coming from the first of (2.41)), and the operator $B_{\geq 4}$ is given by

$$B_{\geq 4}(U) := \operatorname{Op}_{\mathsf{out}}^{BW} \left(\mathrm{i}(g_2 \#_{\varrho} \underline{\mathsf{V}} \xi - \underline{\mathsf{V}} \xi \#_{\varrho} g_2) - \mathrm{i}(g_2 \#_{\varrho} \underline{\mathsf{d}} + \underline{\mathsf{d}} \#_{\varrho} g_2) \right) + R'(U) + \int_0^1 (1 - \tau) \Psi^{\tau}(U) \operatorname{Ad}_{G(U)}^2 \left[\operatorname{Op}_{\mathsf{vec}}^{BW}(-\mathrm{i}|\xi|^{\alpha} + \mathrm{i}\underline{\mathsf{V}} \xi + \mathrm{i}\underline{\mathsf{d}}) \right] \left(\Psi^{\tau}(U) \right)^{-1} \mathrm{d}\tau ,$$

$$(4.17)$$

where R' is a matrix of smoothing operators in $\mathcal{R}_{\geq 4}^{-\varrho+(1-\alpha)}[r]$. We claim that $B_{\geq 4}$ is a nonhomogeneous bounded operator in $\mathcal{M}_{\geq 4}^0[r]$. Indeed, since $g_2 \in \widetilde{\Gamma}_2^{-\alpha}$, $\underline{\underline{V}}$ and $\underline{\underline{d}}$ belong to $\widetilde{\mathcal{F}}_2^{\mathbb{R}}$, and $-\varrho+1-\alpha \leq 0$, we get that both the first line of (4.17) and $\operatorname{Ad}_{G(U)}^2[\operatorname{Op}_{\operatorname{vec}}^{BW}(-i|\xi|^{\alpha}+i\underline{\underline{V}}\xi+i\underline{\underline{d}})]$ are matrices of 0-operators in $\widetilde{\mathcal{M}}_4^0$ and so in $\mathcal{M}_{\geq 4}^0[r]$ (use the symbolic calculus of Proposition 2.7 and the bullets after Definition 2.4). Finally, being Ψ^{τ} an admissible transformation, also the second line of (4.17) is a matrix of non-homogeneous 0-operators in $\mathcal{M}_{\geq 4}^0[r]$ (see Remark 2.11–(2)).

Consider now (4.14). Expanding as in (4.16) one see that the $\overline{2}$ -homogeneous component remains the unchanged, getting

$$(4.14) = \operatorname{Op}_{\mathsf{out}}^{BW}(b(U;x)) W + R_2(U)W + B_{\geq 4}(U)W$$

where $B_{\geq 4}(U)$ is another matrix of non-homogeneous 0-operators in $\mathcal{M}^0_{\geq 4}[r]$.

Finally we consider line (4.15). This time we use the Lie expansion (Lemma A.1 of [11])

$$(\partial_t \Psi(U)) \Psi(U)^{-1} = \partial_t G(U) + \int_0^1 (1-\tau) \Psi^{\tau}(U) \operatorname{Ad}_{G(U)} [\partial_t G(U)] (\Psi^{\tau}(U))^{-1} \mathrm{d}\tau .$$

Then, using that $g_2(U) \equiv g_2(U, U)$ is a symmetric function of U, we get that $\partial_t G(U) = \operatorname{Op}_{\mathsf{out}}^{BW}(\partial_t g_2(U; x, \xi)) = 2\operatorname{Op}_{\mathsf{out}}^{BW}(g_2(\partial_t U, U; x, \xi))$. Since U solves equation (4.1), we get

$$(\partial_t \Psi(U)) \Psi^{-1}(U) = \operatorname{Op}_{\mathsf{out}}^{BW}(2g_2(-\mathrm{i}\Omega(D)U, U; x, \xi))) + B_{\geq 4}(U)$$

4 PARADIFFERENTIAL NORMAL FORM

where, using also (4.2),

$$\begin{split} B_{\geq 4}(U) &:= \mathrm{Op}_{\mathsf{out}}^{^{BW}}(2g_2(M_{\mathtt{NLS}}(U)U, U; x, \xi)) \\ &+ \int_0^1 (1-\tau) \Psi^{\tau}(U) \operatorname{Ad}_{G(U)} \left[2\mathrm{Op}_{\mathtt{out}}^{^{BW}}(g_2(-\mathrm{i}\Omega(D)U + M_{\mathtt{NLS}}(U)U, U; x, \xi)) \right] (\Psi^{\tau}(U))^{-1} \mathrm{d}\tau \end{split}$$

By Lemma 2.9, the fact that Ψ^{τ} is an admissible transformation, and the bullets after Definition 2.4, we deduce that $B_{\geq 4}$ is a matrix of $(-\alpha)$ -operators in $\mathcal{M}_{\geq 4}^{-\alpha}[r]$.

In conclusion, we get that

$$\partial_{t}W = \operatorname{Op}_{\mathsf{vec}}^{BW}(-\mathbf{i}|\xi|^{\alpha} + \mathbf{i}\underline{\mathbb{V}}(U)\xi + \mathbf{i}\underline{\mathbb{d}}(U))W + \operatorname{Op}_{\mathsf{out}}^{BW}(\mathbf{i}[(g_{2}(U)\#_{\varrho}|\xi|^{\alpha} + |\xi|^{\alpha}\#_{\varrho}g_{2}(U) - 2g_{2}(\mathbf{\Omega}(D)U,U))] + b(U))W + (R_{2}(U) + R'_{2}(U))W + B_{\geq 4}(U)W$$

$$(4.18)$$

where $B_{\geq 4}(U)$ is a matrix of 0-operators in $\mathcal{M}^{0}_{\geq 4}[r]$. Then the thesis follows from the following lemma.

Lemma 4.5 (The out-diagonal homological equation). Let $\rho > 0$. There exists a symbol $g_2(U; x, \xi) \in \tilde{\Gamma}_2^{-\alpha}$ of the form (4.12) such that

$$r_2(U; \cdot) := i [(g_2(U) \#_{\varrho} |\xi|^{\alpha} + |\xi|^{\alpha} \#_{\varrho} g_2(U) - 2g_2(\mathbf{\Omega}(D)U, U))] + b(U) \in \widetilde{\Gamma}_2^{-\varrho}$$
(4.19)

and $r_2(U; \cdot)$ fulfills the second of (2.25).

Proof. Thanks to symbolic calculus formula (2.37) (see also (2.38)), we have that for any $g \in \widetilde{\Gamma}_2^m$, $m \in \mathbb{R}$,

$$\begin{cases} \mathbf{r}[g](U) := g(U) \#_{\varrho} |\xi|^{\alpha} + |\xi|^{\alpha} \#_{\varrho} g(U) - 2g(U) |\xi|^{\alpha} \in \widetilde{\Gamma}_{2}^{m+\alpha-2} \\ \mathbf{f}[g](U) := 2g(\mathbf{\Omega}(D)U, U) \in \widetilde{\Gamma}_{2}^{m} \end{cases}$$

Moreover if g fulfills the second of (2.25), so do r[g] and f[g]. Then the homological equation in (4.19) reads

$$r_2(U) = 2ig_2(U)|\xi|^{\alpha} + ir[g_2](U) - if[g_2](U) + b(U) \in \widetilde{\Gamma}_2^{-\varrho}$$

which we solve iteratively exploiting that $g \mapsto \mathbf{r}[g]$ and $g \mapsto \mathbf{f}[g]$ are linear. Namely we put $g_2 := g^{(1)} + g^{(2)} + \cdots + g^{(p)}$ with

$$\begin{split} g^{(1)}(U;x,\xi) &:= -\frac{b(U;x)}{2i|\xi|^{\alpha}} \in \widetilde{\Gamma}_{2}^{-\alpha}, \\ g^{(2)}(U;x,\xi) &:= -\frac{\mathrm{ir}[g^{(1)}](U;x,\xi) - \mathrm{if}[g^{(1)}](U;x,\xi)}{2i|\xi|^{\alpha}} \in \widetilde{\Gamma}_{2}^{-2\alpha} \\ &\vdots \\ g^{(p)}(U;x,\xi) &:= -\frac{\mathrm{ir}[g^{(p-1)}](U;x,\xi) - \mathrm{if}[g^{(p-1)}](U;x,\xi)}{2i|\xi|^{\alpha}} \in \widetilde{\Gamma}_{2}^{-p\alpha} \end{split}$$

With this choice we have $r_2(U) = i\mathbf{r}[g^{(p)}](U) - i\mathbf{f}[g^{(p)}](U) \in \widetilde{\Gamma}_2^{-p\alpha}$ which implies the thesis choosing $p > \varrho/\alpha$. Moreover, since b fulfills the second of (2.25) (recall (4.4)), so does $g^{(1)}$, and by construction each $g^{(\ell)}, \ell \geq 2$ and the symbol $r_2(U)$. In particular g_2 has the claimed form in (4.12).

Applying Lemma 4.5, equation (4.18) becomes

$$\partial_t W = \operatorname{Op}_{\mathsf{vec}}^{BW}(-\mathbf{i}|\xi|^{\alpha} + \mathbf{i}\underline{\mathbf{U}}(U)\xi + \mathbf{i}\underline{\mathbf{d}}(U))W + (R_2(U) + R_2'(U) + R_2''(U))W + B_{\geq 4}(U)W$$
(4.20)

where $R_2''(U) = \operatorname{Op}_{\mathsf{out}}^{BW}(r_2(U; \cdot)) \in \widetilde{\mathcal{R}}_2^{-\varrho}$ is the paradifferential operator of order $-\varrho$ coming from the symbol in (4.19). This proves the identity (4.11), renaming $R_2 + R_2' + R_2'' \rightsquigarrow R_2$.

Finally we prove that the matrices of smoothing operators are gauge invariant. Indeed each operator on the right of (4.13)–(4.15) is gauge invariant (recall Lemma 4.2), as well as the 2-homogeneous matrix of paradifferential operators in (4.20). Then, by difference, the 2-homogeneous smoothing operators $R_2 + R'_2 + R''_2$ are gauge invariant as well.

4.2 Reduction of the highest order

In this section we perform a transformation that reduces the symbol of the highest order paradifferential operator $\operatorname{Op}_{\mathsf{vec}}^{BW}(\underline{\mathbb{V}}(U;x)\mathrm{i}\xi)$ to its resonant normal form.

Proposition 4.6 (Paracomposition). Let $\rho \geq 1$. There are $s_0, r > 0$ and a 2-admissible transformation $\Phi(U) \in \mathcal{M}^0_{\geq 0}[r]$ (see Definition 2.10) such that if $U(t) \in B_{s_0,\mathbb{R}}(I;r)$ solves (4.3), then the variable

$$W_1 := \Phi(U)W \stackrel{(4.10)}{=} \Phi(U)\Psi(U)U \quad solves \tag{4.21}$$

$$\partial_t W_1 = -\operatorname{i} \mathbf{\Omega}(D) W + \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\operatorname{i} \langle \underline{\mathbf{V}} \rangle (U; x) \xi + \operatorname{i} V_{\geq 4}(U; x) \xi + \operatorname{i} a_2^{(\alpha)}(U; x, \xi) + \operatorname{i} a_{\geq 4}^{(\alpha)}(U; x, \xi) \right) W_1$$

$$+ R_2(U) W_1 + B_{\geq 4}(U) W_1$$

$$(4.22)$$

where:

• $\Omega(D)$ is the matrix of Fourier multipliers defined in (3.17);

• $\langle \underline{\mathbf{V}} \rangle (U; x)$ is the resonant part of the function $\underline{\mathbf{V}}(U; x)$ in (4.4), namely the zero-average, real valued function in (4.7);

• $V_{\geq 4}(U;x)$ is a real function in $\mathcal{F}_{\geq 4}^{\mathbb{R}}[r]$;

• $a_2^{(\alpha)}(U; x, \xi)$ is a zero average, gauge invariant (fulfills the first of (2.25)), real symbol in $\widetilde{\Gamma}_2^{\alpha}$ and $a_{\geq 4}^{(\alpha)}(U; x, \xi)$ a real non-homogeneous symbol in $\Gamma_{\geq 4}^{\alpha}[r]$;

• $R_2(U)$ is a real-to-real, gauge invariant matrix of homogeneous smoothing operators in $\widetilde{\mathcal{R}}_2^{-\varrho}$;

• $B_{\geq 4}(U)$ is a real-to-real matrix of 0-operators in $\mathcal{M}^{0}_{\geq 4}[r]$.

Proof. We define the transformation $\Phi(U)$ as the time-1 flow of the paradifferential equation

$$\begin{cases} \partial_{\tau} \Phi^{\tau}(U) = G(U) \Phi^{\tau}(U) \\ \Phi^{0}(U) = \mathrm{Id}, \end{cases} \quad \text{where} \quad G(U) := \mathrm{Op}_{\mathsf{vec}}^{BW} \left(\frac{\beta_{2}(U;x)}{1 + \tau(\beta_{2})_{x}(U;x)} \mathrm{i}\xi \right) \tag{4.23}$$

and β_2 is the real valued, 2-homogeneous function

$$\beta_2(U;x) := \sum_{|j_1| \neq |j_2|} \frac{1}{\mathbf{i}(|j_1|^\alpha - |j_2|^\alpha)} u_{j_1} \overline{u_{j_2}} e^{\mathbf{i}(j_1 - j_2)x}$$
(4.24)

whose coefficients fulfill (2.15) with $\mu = 1 - \alpha$. By Lemma 2.15, Φ is a 2-admissible transformation. Moreover, since β_2 fulfills the first of (2.25), G as well as Φ^{τ} , $\tau \in [0, 1]$, are gauge invariant (see the bullet of formula (2.25) and Remark 2.14).

Recalling (4.11), the variable $W_1 := \Phi(U)W$ solves

$$\partial_t W_1 = \Phi(U) \operatorname{Op}_{\mathsf{vec}}^{BW}(-\mathrm{i}|\xi|^{\alpha} + \mathrm{i}\underline{\mathsf{V}}(U)\xi + \mathrm{i}\underline{\mathsf{d}}(U)) \Phi(U)^{-1} W_1$$
(4.25)

$$+ \left(\partial_t \Phi(U)\right) \Phi(U)^{-1} W_1 \tag{4.26}$$

$$+ \Phi(U) \left[R_2(U) + B_{\geq 4}(U) \right] \Phi(U)^{-1} W_1 .$$
(4.27)

We now compute each term, starting from (4.25). By Proposition B.1–2 (with $\rho \rightarrow \rho + \alpha$) we get

$$\Phi(U) \operatorname{Op}_{\mathtt{vec}}^{BW}(-\mathrm{i}|\xi|^{\alpha}) \Phi(U)^{-1} = \operatorname{Op}_{\mathtt{vec}}^{BW}\left(-\mathrm{i}|\xi|^{\alpha} + \mathrm{i}a_{2}^{(\alpha)} + \mathrm{i}a_{\geq 4}^{(\alpha)}\right) + B_{\geq 4}(U) + R_{2}'(U)$$

where $a_2^{(\alpha)}$ is a real, zero average, gauge invariant symbol in $\widetilde{\Gamma}_2^{\alpha}$, $a_{\geq 4}^{(\alpha)}$ is a real symbol in $\Gamma_{\geq 4}^{\alpha}[r]$, $B_{\geq 4} = \operatorname{Op}_{\mathsf{vec}}^{BW}\left(\operatorname{ia}_{\geq 4}^{(\alpha-2)}\right) + R_{\geq 4}$ (see (B.2)) is a real-to-real matrix of 0-operators in $\mathcal{M}_{\geq 4}^0[r]$ and finally $R'_2(U)$ is a real-to-real, gauge invariant matrix of smoothing operators in $\widetilde{\mathcal{R}}_2^{-\varrho}$.

Then, by Proposition B.1–1, we get

$$\Phi(U)\operatorname{Op}_{\mathtt{vec}}^{BW}(\underline{i}\underline{V}\xi + \underline{i}\underline{d}) \Phi(U)^{-1} = \operatorname{Op}_{\mathtt{vec}}^{BW}(\underline{i}\underline{V}\xi + \underline{i}V'_{\geq 4}\xi + \underline{i}\underline{d}) + B_{\geq 4}(U)$$

with $V'_{\geq 4} \in \mathcal{F}^{\mathbb{R}}_{\geq 4}[r]$ and, thanks to $\varrho \geq 1$, $B_{\geq 4}$ a real-to-real matrix of 0-operators in $\mathcal{M}^{0}_{\geq 4}[r]$.

4 PARADIFFERENTIAL NORMAL FORM

Next we consider the term in (4.26). We apply Proposition B.1–4 and get

$$(\partial_t \Phi(U)) \Phi(U)^{-1} = \operatorname{Op}_{\mathsf{vec}}^{BW} \left(2\mathrm{i}\beta_2 (-\mathrm{i}\mathbf{\Omega}(D)U, U)\xi + \mathrm{i}V_{\geq 4}''(U)\xi \right) + B_{\geq 4}(U)$$

where $V_{\geq 4}'' \in \mathcal{F}_{\geq 4}^{\mathbb{R}}[r]$ and, using again $\varrho \geq 1$, $B_{\geq 4}$ a real-to-real matrix of 0-operators in $\mathcal{M}_{\geq 4}^0[r]$. Finally we consider line (4.27). By Proposition B.1–3 and Remark 2.11– (2)

$$(4.27) = R_2(U) + B_{>4}(U)$$

with $R_2(U)$ the same real-to-real, gauge invariant matrix of smoothing operators in $\mathcal{R}_2^{-\varrho}$ of Proposition 4.4 and with $B_{\geq 4}$ a real-to-real matrix of 0-operators in $\mathcal{M}_{\geq 4}^0[r]$.

Altogether we have the expansion

$$\begin{aligned} \partial_t W_1 = & \operatorname{Op}_{\mathsf{vec}}^{BW} \left(-\mathrm{i} |\xi|^{\alpha} + \mathrm{i} \underline{\mathbb{V}} \xi + 2\mathrm{i} \beta_2 (-\mathrm{i} \mathbf{\Omega}(D) U, U) \xi + \mathrm{i} a_2^{(\alpha)} \right) W_1 + (R_2(U) + R_2'(U)) W_1 \\ &+ \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\mathrm{i} V_{\geq 4} \xi + \mathrm{i} a_{\geq 4}^{(\alpha)} \right) W_1 + B_{\geq 4}(U) W_1 \;. \end{aligned}$$

One verifies that β_2 in (4.24) solves the homological equation

$$2\beta_2(-\mathrm{i}\mathbf{\Omega}(D)U, U; x) + \underline{\mathrm{V}}(U; x) = \langle \underline{\mathrm{V}} \rangle(U; x) ,$$

using the expressions of $\underline{\underline{V}}$ in (4.4), $\Omega(D)$ in (3.17), and $\langle \underline{\underline{V}} \rangle$ in (4.7). This proves the expansion in (4.22), renaming $R_2 + R'_2 \rightsquigarrow R_2$; note that we proved that it is gauge invariant being sum of gauge invariant operators.

4.3 The weak Λ -normal form

In this section we perform a Poincaré normal form, with the goal of putting the smoothing operator $R_2(U)W_1$ in (4.22) into weak- Λ normal form (see Definition (4.1)).

Proposition 4.7 (Weak- Λ normal form). Let $\varrho \geq 2 - \alpha$. There are $s_0, r > 0$ and a 0-admissible transformation $\Upsilon(U) \in \mathcal{M}^0_{\geq 0}[r]$ (see Definition 2.10) such that if $U(t) \in B_{s_0,\mathbb{R}}(I;r)$ solves (4.3), then the variable

$$Z := \Upsilon(U)W_1 \stackrel{(4.21),(4.10)}{=} \Upsilon(U)\Phi(U)\Psi(U)U \quad solves \tag{4.28}$$

$$\partial_t Z = -\operatorname{i} \mathbf{\Omega}(D) Z + \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\operatorname{i} \langle \underline{\mathbf{V}} \rangle(U; x) \xi + \operatorname{i} V_{\geq 4}(U; x) \xi + \operatorname{i} a_2^{(\alpha)}(U; x, \xi) + \operatorname{i} a_{\geq 4}^{(\alpha)}(U; x, \xi) \right) Z + R_2^{(\Lambda)}(U) Z + B_{\geq 4}(U) Z$$

$$(4.29)$$

where $\langle \underline{\underline{V}} \rangle$, $V_{\geq 4}$, $a_2^{(\alpha)}$ and $a_{\geq 4}^{(\alpha)}$ are the same symbols of Proposition 4.6, whereas

• $R_2^{(\Lambda)}(U)$ is a real-to-real, gauge invariant matrix of smoothing operators in $\widetilde{\mathcal{R}}_2^{-\varrho}$ such that the cubic vector field $X^{(\Lambda)}(Z) := R_2^{(\Lambda)}(Z)Z$ is in weak- Λ normal form, namely it fulfills

$$\Pi_{\mathcal{P}^{(n)}_{\Lambda}} X^{(\Lambda)} = \Pi_{\mathsf{R}^{(n)}_{\Lambda}} X^{(\Lambda)} , \quad n = 0, 1, 2 .$$
(4.30)

• $B_{\geq 4}(U)$ is a real-to-real matrix of 0-operators in $\mathcal{M}_{\geq 4}^0[r]$.

Proof. We look for a transformation $\Upsilon(U)$ as the time-1 flow of the equation

$$\partial_{\tau}\Upsilon^{\tau}(U) = \mathsf{Q}_2(U)\Upsilon^{\tau}(U), \quad \Upsilon^0(U) = \mathrm{Id}$$

where Q_2 is a matrix of smoothing operators in $\widetilde{\mathcal{R}}_2^{-\varrho+1-\alpha}$ to be determined. By Lemma 2.16, the map Υ^{τ} is a 0-admissible transformation. Recalling (4.22), the variable $Z := \Upsilon(U)W_1$ fulfills

$$\partial_t Z = \Upsilon(U) \left(-\mathrm{i}\Omega(D)\right) \Upsilon(U)^{-1} Z + \Upsilon(U) \mathrm{Op}_{\mathsf{vec}}^{BW} \left(\mathrm{i}\mathfrak{m}^{(1)}\right) \Upsilon(U)^{-1} Z + \Upsilon(U) \left(R_2(U) + B_{\geq 4}(U)\right) \Upsilon(U)^{-1} Z + \left(\partial_t \Upsilon(U)\right) \Upsilon(U)^{-1} Z$$

4 PARADIFFERENTIAL NORMAL FORM

where we set $\mathbf{m}^{(1)} := \langle \underline{\mathbf{V}} \rangle \xi + a_2^{(\alpha)} + V_{\geq 4} \xi + \widetilde{a}_{\geq 4}^{(\alpha)} \in \Sigma \Gamma_2^1[r]$. By Proposition B.2 (with $\rho \rightsquigarrow \rho - (1 - \alpha)$) we get

$$\partial_t Z = -\operatorname{i} \mathbf{\Omega}(D) Z + \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\operatorname{im}^{(1)} \right) Z + 2 \mathbb{Q}_2 \left(-\operatorname{i} \mathbf{\Omega}(D) U, U \right) Z + [\mathbb{Q}_2(U), -\operatorname{i} \mathbf{\Omega}(D)] Z + R_2(U) Z + B_{\geq 4}(U) Z + R_{\geq 4}(U) Z$$

$$(4.31)$$

where $B_{\geq 4}(U)$ is a real-to-real matrix of 0-operators in $\mathcal{M}_{\geq 4}^0[r]$ and $R_{\geq 4}(U)$ is a real-to-real matrix of smoothing operators in $\mathcal{R}_{\geq 4}^{-\varrho+2-\alpha}[r]$ which we shall regard as a 0-operator in $\mathcal{M}_{\geq 4}^0[r]$ since $\varrho \geq 2-\alpha$. To determine $\mathbb{Q}_2(U)$, expand the vector field $R_2(U)Z$ in (4.22) in Fourier components as

$$(R_2(U)Z)_k^{\sigma} = \sum_{\mathcal{P}_4} R_{j_1,j_2,j,k}^{\sigma_1,\sigma_2,\sigma',\sigma} u_{j_1}^{\sigma_1} u_{j_2}^{\sigma_2} z_j^{\sigma'}$$

where with the sum over \mathcal{P}_4 we mean that the indexes $(j_1, j_2, j, k, \sigma_1, \sigma_2, \sigma', -\sigma)$ belong to \mathcal{P}_4 . Below we use the same notation. Note that this writing is possible since $R_2(U)$ is gauge invariant.

Then we define

$$(R_{2}^{(\Lambda)}(U)Z)_{k}^{\sigma} := \sum_{\mathcal{P}_{4}} \Lambda_{j_{1},j_{2},j,k}^{\sigma_{1},\sigma_{2},\sigma',\sigma} u_{j_{1}}^{\sigma_{1}} u_{j_{2}}^{\sigma_{2}} z_{j}^{\sigma'}, \quad \Lambda_{j_{1},j_{2},j,k}^{\sigma_{1},\sigma_{2},\sigma',\sigma} := R_{j_{1},j_{2},j,k}^{\sigma_{1},\sigma_{2},\sigma',\sigma} \delta\Big((j_{1},j_{2},j,k,\sigma_{1},\sigma_{2},\sigma',-\sigma) \in \mathcal{C}\Big),$$

where $\mathcal{C} := \bigcup_{n=0}^{2} \mathsf{R}^{(n)}_{\Lambda} \cup \bigcup_{n=3}^{4} \mathcal{P}^{(n)}_{\Lambda}$. We choose $\mathsf{Q}_{2}(U)$ so that

$$2\mathbf{Q}_{2}\left(-\mathrm{i}\mathbf{\Omega}(D)U,U\right) + \left[\mathbf{Q}_{2}(U),-\mathrm{i}\mathbf{\Omega}(D)\right] + R_{2}(U) = R_{2}^{(\Lambda)}(U).$$
(4.32)

We claim that one can put, denoting $\vec{j} = (j_1, j_2), \ \vec{\sigma} = (\sigma_1, \sigma_2),$

$$(\mathbf{Q}_{2}(U)Z)_{k}^{\sigma} := \sum_{\mathcal{P}_{4}} \mathbf{Q}_{\vec{j},j,k}^{\vec{\sigma},\sigma',\sigma} u_{j_{1}}^{\sigma_{1}} u_{j_{2}}^{\sigma_{2}} z_{j}^{\sigma'}$$
(4.33)

where

$$\mathbb{Q}_{\vec{j},j,k}^{\vec{\sigma},\sigma',\sigma} := \begin{cases}
\frac{R_{\vec{j},j,k}^{\vec{\sigma},\sigma',\sigma}}{i(\sigma_1|j_1|^{\alpha} + \sigma_2|j_2|^{\alpha} + \sigma|j|^{\alpha} - \sigma|k|^{\alpha})}, & (\vec{j},j,k,\vec{\sigma},\sigma',-\sigma) \in \bigcup_{n=1}^{2} \left(\mathcal{P}_{\Lambda}^{(n)} \setminus \mathsf{R}_{\Lambda}^{(n)}\right) \\
0, & (\vec{j},j,k,\vec{\sigma},\sigma',-\sigma) \in \mathcal{C}
\end{cases} (4.34)$$

Lemma 4.8. $\mathbb{Q}_2(U)$ in (4.33)–(4.34) is a matrix of smoothing operators in $\widetilde{\mathcal{R}}_2^{-\varrho+1-\alpha}$ fulfilling (4.32).

Proof. As $R_2(U)$ is a smoothing operator in $\widetilde{\mathcal{R}}_2^{-\varrho}$, its coefficients fulfill the estimate: for some $\mu \geq 0$, C > 0,

$$\left| R_{\vec{j},j,k}^{\vec{\sigma},\sigma',\sigma} \right| \le C \frac{\max_2\{\langle j_1 \rangle, \langle j_2 \rangle, \langle j \rangle\}^{\mu}}{\max\{\langle j_1 \rangle, \langle j_2 \rangle, \langle j \rangle\}^{\varrho}}, \quad \forall (\vec{j},j,k,\vec{\sigma},\sigma',-\sigma) \in \mathcal{P}_4 ,$$

$$(4.35)$$

and satisfy the symmetric and reality properties (2.10) and (2.11).

Consider now the coefficients $Q_{\vec{j},j,k}^{\vec{\sigma},\sigma',\sigma}$ in (4.34). Clearly they satisfy the symmetric and reality properties (2.10) and (2.11). We now bound them. By (4.35), Lemma 3.3 and the momentum relation $\sigma k = \sigma_1 j_1 + \sigma_2 j_2 + \sigma' j$,

$$\left| \mathbf{Q}_{\vec{j},j,k}^{\vec{\sigma},\sigma',\sigma} \right| \le C \frac{\max_2\{\langle j_1 \rangle, \langle j_2 \rangle, \langle j_3 \rangle\}^{\mu}}{\max\{\langle j_1 \rangle, \langle j_2 \rangle, \langle j_3 \rangle\}^{\varrho - (1-\alpha)}} \qquad \forall (\vec{j},j,k,\vec{\sigma},\sigma',-\sigma) \in \mathcal{P}_{\Lambda}^{(1)} \cup (\mathcal{P}_{\Lambda}^{(2)} \setminus \mathsf{R}_{\Lambda}^{(2)}) ,$$

(recall that $\mathsf{R}^{(1)}_{\Lambda} = \emptyset$). This shows that $\mathsf{Q}_2(U)$ is a matrix of smoothing operators in $\widetilde{\mathcal{R}}_2^{-\varrho+1-\alpha}$.

It is clear that $\mathbb{Q}_2(U)$ fulfills (4.32), also noting that $\Pi_{\mathcal{P}^{(0)}_{\lambda}}(R_2(Z)Z) = \Pi_{\mathsf{R}^{(0)}_{\lambda}}(R_2(Z)Z)$ in view of Lemma 3.3 (i).

With such $Q_2(U)$, system (4.31) reduces to (4.29).

We prove now that the vector field $X^{(\Lambda)}(Z) = R_2^{(\Lambda)}(Z)Z$ is in weak- Λ normal form, i.e. it fulfills (4.30). Indeed the coefficients of the vector field $X^{(\Lambda)}$ are obtained as in (2.36) and, being the set $\mathcal C$ symmetric with respect to the first three indexes, they have the form

$$X_{j_{1},j_{2},j_{3},k}^{\sigma_{1},\sigma_{2},\sigma_{3},\sigma} = \frac{1}{3} \Big(R_{j_{1},j_{2},j_{3},k}^{\sigma_{1},\sigma_{2},\sigma_{3},\sigma} + R_{j_{3},j_{2},j_{1},k}^{\sigma_{3},\sigma_{2},\sigma_{1},\sigma} + R_{j_{1},j_{3},j_{2},k}^{\sigma_{1},\sigma_{3},\sigma_{2},\sigma} \Big) \delta\Big((j_{1},j_{2},j_{3},k,\sigma_{1},\sigma_{2},\sigma_{3},-\sigma) \in \mathcal{C} \Big) .$$

Proposition 4.7 is proved.

4.4 Identification and proof of Theorem 4.3

With the aid of paradifferential normal form, we have conjugated the original system (4.1) to the new system (4.29). The next steps are: (i) to write (4.29) as a system in the single variable Z(t), and (ii) to compute explicitly $\Pi_{\mathcal{P}_{i}^{(n)}}X^{(\Lambda)}$ in (4.30) for n = 0, 1, 2, deducing (4.9).

To achieve (i), recall that the map in (4.28) has the form

$$Z = \mathcal{F}(U) = \mathbf{F}(U)U, \quad \mathbf{F}(U) := \Upsilon(U)\Phi(U)\Psi(U)$$
(4.36)

with $\mathbf{F}(U)$ a 2-admissible transformation, being composition of admissible transformations (recall Propositions 4.4, 4.6, 4.7 and Lemma 2.12). Moreover Lemma 2.13 ensures that \mathcal{F} is locally invertible in a small ball $B_{s'_0}(r')$ for some $s'_0, r' > 0$, with inverse map \mathcal{F}^{-1} having the structure

$$U = \mathcal{F}^{-1}(Z) = \mathbf{G}(Z)Z, \quad \text{with } \mathbf{G}(Z) = \mathrm{Id} + \mathbf{G}_{\geq 2}(Z), \quad \mathbf{G}_{\geq 2}(Z) \in \Sigma \mathcal{M}_2^4[r'], \quad (4.37)$$

for some r' > 0. We then substitute U in the internal variables of the operators in (4.29). Consider first the 2-homogeneous operators. We have, using Lemma 2.9–1,

$$\langle \underline{\mathbf{V}} \rangle (\mathcal{F}^{-1}(Z); x) \xi - \langle \underline{\mathbf{V}} \rangle (Z; x) \xi \in \Gamma^{1}_{\geq 4}[r'], \quad a_{2}^{(\alpha)} (\mathcal{F}^{-1}(Z); x, \xi) - a_{2}^{(\alpha)}(Z; x, \xi) \in \Gamma^{\alpha}_{\geq 4}[r']$$

and, using Lemma 2.9–2, $R_2^{(\Lambda)}(\mathcal{F}^{-1}(Z)) - R_2^{(\Lambda)}(Z) \in \mathcal{R}_{\geq 4}^{-\varrho+4}[r']$. Then we substitute $U = \mathcal{F}^{-1}(Z)$ in the non-homogeneous operators $\operatorname{Op}_{\mathsf{vec}}^{BW}\left(\mathrm{i}V_{\geq 4}(U;x)\xi + \mathrm{i}a_{\geq 4}^{(\alpha)}(U;x,\xi)\right)$ and $B_{\geq 4}(U)$, applying Lemma 2.9–1& 5. In conclusion, setting $\varrho := 4$, we obtain the following:

Proposition 4.9. There are $s_0, r > 0$ such that if $U(t) \in B_{s_0,\mathbb{R}}(I; r)$ solves (4.3), then the variable Z(t) in (4.36) solves the system

$$\partial_t Z = -i\mathbf{\Omega}(D)Z + \operatorname{Op}_{\mathsf{vec}}^{BW} \left(i\langle \underline{\mathbf{V}} \rangle (Z; x) \xi + ia_2^{(\alpha)}(Z; x, \xi) \right) + X^{(\Lambda)}(Z) + \operatorname{Op}_{\mathsf{vec}}^{BW} \left(i\widetilde{V}_{\geq 4}(Z; x) \xi + i\widetilde{a}_{\geq 4}^{(\alpha)}(Z; x, \xi) \right) Z + \widetilde{B}_{\geq 4}(Z) Z$$

$$(4.38)$$

where $\langle \underline{\mathbf{V}} \rangle$ and $a_2^{(\alpha)}$ are the quadratic symbols in Proposition 4.6, $X^{(\Lambda)}(Z)$ is the cubic vector field in weak- Λ normal form of Proposition 4.7 is , whereas

- $\widetilde{V}_{\geq 4}(Z; x)$ is a real function in $\mathcal{F}_{\geq 4}^{\mathbb{R}}[r];$
- $\widetilde{a}_{>4}^{(\alpha)}(Z; x, \xi)$ is a real non-homogeneous symbol in $\Gamma_{\geq 4}^{\alpha}[r];$
- $\widetilde{B}_{\geq 4}(Z)$ is a real-to-real matrix of 0-operators in $\mathcal{M}^{0}_{\geq 4}[r]$.

The next step (*ii*) is to compute explicitly $\prod_{\mathcal{P}_{\lambda}^{(n)}} X^{(\Lambda)}$, n = 0, 1, 2:

Proposition 4.10. The vector field $X^{(\Lambda)}(Z)$ of Proposition 4.7 is actually in strong- Λ normal form (Definition 4.1) and fulfills (4.9).

Proof. We combine the abstract identification argument of Proposition 3.6 with the characterization of the resonant monomials of the original vector field X_3 in Lemma 3.4.

Precisely, we apply the identification result of Proposition 3.6 to the starting NLS equation (4.1) (which has the required structure in (3.18) in view of (4.2)) and with the admissible transformation $\mathbf{F}(U)$ in (4.36), getting that Z fulfills an equation of the form (3.19). Identifying the cubic vector field of (3.19) with the one of (4.38) we get the identity

$$\operatorname{Op}_{\mathtt{vec}}^{BW}\left(\mathrm{i}\langle \underline{\mathtt{V}}\rangle(Z;x)\xi + \mathrm{i}a_2^{(\alpha)}(Z;x,\xi)\right) + X^{(\Lambda)}(Z) = \widetilde{X}_3(Z) \ .$$

In addition, in view of (3.20), we have

$$\Pi_{\mathsf{R}^{(n)}_{\Lambda}}\left(\operatorname{Op}_{\mathtt{vec}}^{BW}\left(\mathrm{i}\langle\underline{\mathtt{V}}\rangle(Z;x)\xi + \mathrm{i}a_{2}^{(\alpha)}(Z;x,\xi)\right)Z + X^{(\Lambda)}\right) = \Pi_{\mathsf{R}^{(n)}_{\Lambda}}X_{3} , \quad n = 0, 1, 2 .$$

$$(4.39)$$

Now we apply Lemma 3.5 to the cubic vector field $\operatorname{Op}_{\mathsf{vec}}^{BW}\left(\mathrm{i}\langle \underline{\underline{V}}\rangle \xi + \mathrm{i}a_2^{(\alpha)}\right)Z$; this can be done since the symbols $\langle \underline{\mathbf{V}} \rangle(Z;x)\xi$ and $a_2^{(\alpha)}(Z;x,\xi)$ have both zero-average (Proposition 4.6) and are gauge invariant (i.e. fulfills the first of (2.25)). We conclude that

$$\Pi_{\mathsf{R}^{(n)}_{\Lambda}}\left[\operatorname{Op}_{\mathtt{vec}}^{BW}\left(\mathrm{i}\langle\underline{\mathtt{V}}\rangle(Z;x)\xi + \mathrm{i}a_{2}^{(\alpha)}(Z;x,\xi)\right)Z\right] = 0 , \quad n = 0, 1, 2 , \qquad (4.40)$$

from which we get immediately

$$\Pi_{\mathcal{P}^{(n)}_{\Lambda}} X^{(\Lambda)} \stackrel{(4.30)}{=} \Pi_{\mathsf{R}^{(n)}_{\Lambda}} X^{(\Lambda)} \stackrel{(4.39),(4.40)}{=} \Pi_{\mathsf{R}^{(n)}_{\Lambda}} X_3 , \quad n = 0, 1, 2 .$$

This last vector field is computed in Lemma 3.4, proving (4.9).

Proof of Theorem 4.3. It follows from Proposition 4.9 and 4.10.

$\mathbf{5}$ The effective equation

The goal of this section is to study the long-time dynamics of solutions of equation (4.6) fulfilling certain upper-bounds, that we call *long-time controlled*, see Definition 5.2. In view of the reality of system (4.6), we regard it as a scalar equation in z(t). We study separately the dynamics of the modes supported on Λ , namely $z_{\pm 1}(t)$, and those supported on Λ^c . Specifically we decompose

$$z(t) = z^{\top}(t) + z^{\perp}(t) , \quad z^{\top}(t) := z_1(t) e^{ix} + z_{-1}(t) e^{-ix} , \quad z^{\perp}(t) := \sum_{|j| \neq 1} z_j(t) e^{ijx} .$$
(5.1)

• **Parameters:** From now on we fix $\mathfrak{s}_0, \mathfrak{r} > 0$ as follows: $\mathfrak{s}_0 := \max\{s_0, s'_0\}$ and $\mathfrak{r} := \min\{r, r'\}$ where $s_0, r > 0$ are given in Theorem 4.3 whereas $s'_0, r' > 0$ are the parameters required to invert the map \mathcal{F} in (4.5), see (4.37). We also fix

$$s > 3\mathfrak{s}_0, \quad \theta \in (0, \theta_*), \quad \theta_* := \min\left(\frac{s - 3\mathfrak{s}_0}{2s - \mathfrak{s}_0}, \frac{1}{5}\right)$$
 (5.2)

The first step is the following one:

Lemma 5.1. If $Z(t) = \left(\frac{z(t)}{\overline{z}(t)}\right) \in B_{\mathfrak{s}_0,\mathbb{R}}(I;\mathfrak{r})$ solves (4.6), then the variables $\left(z^{\top}(t), z^{\perp}(t)\right)$ defined in (5.1) fulfill the system

$$\partial_t z^{\top} = -i|D|^{\alpha} z^{\top} + Y_3^{(\Lambda)}(z^{\top}) + Y_3^{\top}(z) + Y_{\geq 5}^{\top}(z)$$
(5.3)

$$\partial_t z^{\perp} = -i|D|^{\alpha} z^{\perp} + \operatorname{Op}^{BW}(im(z; x, \xi)) z^{\perp} + Y_3^{\perp}(z) + Y_{\geq 5}^{\perp}(z)$$
(5.4)

where

• $Y_3^{(\Lambda)}(z)$ is the integrable vector field

$$Y_3^{(\Lambda)}(z) := Y_3^{(\Lambda)}(z^{\top}) = -i|z_1|^2 z_1 e^{ix} + i|z_{-1}|^2 z_{-1} e^{-ix} ; \qquad (5.5)$$

• $Y_3^{\top}(z)$ and $Y_3^{\perp}(z)$ are cubic smoothing vector fields fulfilling: for any $\mathbf{s} \geq \mathfrak{s}_0$

$$\|Y_{3}^{\top}(z)\|_{\mathfrak{s}} \lesssim \|z^{\perp}\|_{\mathfrak{s}_{0}}^{3} , \qquad \|Y_{3}^{\perp}(z)\|_{\mathfrak{s}+4} \lesssim \left(\|z^{\top}\|_{\mathfrak{s}_{0}} + \|z^{\perp}\|_{\mathfrak{s}_{0}}\right) \|z^{\perp}\|_{\mathfrak{s}_{0}} \|z^{\perp}\|_{\mathfrak{s}} ; \qquad (5.6)$$

• $m(z; x, \xi)$ is the symbol in $\Sigma\Gamma^1_{\geq 2}[\mathfrak{r}]$ given by

$$\mathbf{m}(z;x,\xi) := \langle \underline{\mathbf{V}} \rangle(Z;x)\xi + a_2^{(\alpha)}(Z;x,\xi) + \widetilde{V}_{\geq 4}(Z;x)\xi + \widetilde{a}_{\geq 4}^{(\alpha)}(Z;x,\xi) , \qquad (5.7)$$

with $\langle \underline{\mathbf{V}} \rangle(Z; x)$ defined in (4.7).

• $Y_{\geq 5}^{\top}(z)$ and $Y_{\geq 5}^{\perp}(z)$ are non-homogeneous vector fields fulfilling the estimate: for any $\mathbf{s} \geq \mathfrak{s}_0$ there are $\overline{C} > 0$, $\mathbf{r} := \mathbf{r}(\mathbf{s}) \in (0, \mathfrak{r})$ and for any $z \in B_{\mathfrak{s}_0}(\mathbf{r}) \cap H^{\mathbf{s}}(\mathbb{T}, \mathbb{C})$,

$$\|Y_{\geq 5}^{\top}(z)\|_{\mathbf{s}} + \|Y_{\geq 5}^{\perp}(z)\|_{\mathbf{s}} \le C \|z\|_{\mathfrak{s}_{0}}^{4} \|z\|_{\mathbf{s}} .$$
(5.8)

Proof. We introduce the projectors

$$\Pi^{\top} z := \sum_{j=\pm 1} z_j \, e^{ijx} \, , \quad \Pi^{\perp} z := \sum_{j\neq \pm 1} z_j \, e^{ijx}$$

and compute the projections of the first component of each term in system (4.6). Since $(-i\Omega(D))^+ = -i|D|^{\alpha}$ is a Fourier multiplier, it commutes with the projectors. So consider the paradifferential vector field $(\operatorname{Op}_{\mathsf{vec}}^{BW}(\operatorname{im})Z)^+ = \operatorname{Op}^{BW}(\operatorname{im})z$. We decompose

$$\mathrm{Op}^{BW}(\mathrm{i}\,\mathrm{m}) = \Pi^{\top}\mathrm{Op}^{BW}(\mathrm{i}\,\mathrm{m})\,\Pi^{\top} + \Pi^{\top}\mathrm{Op}^{BW}(\mathrm{i}\,\mathrm{m})\,\Pi^{\perp} + \Pi^{\perp}\mathrm{Op}^{BW}(\mathrm{i}\,\mathrm{m})\,\Pi^{\top} + \Pi^{\perp}\mathrm{Op}^{BW}(\mathrm{i}\,\mathrm{m})\,\Pi^{\perp} \,.$$

Writing $\mathfrak{m}_2(z; x, \xi) := \langle \underline{\mathbf{U}} \rangle(Z; x)\xi + a_2^{(\alpha)}(Z; x, \xi), \ \mathfrak{m}_{\geq 4}(z; x, \xi) := \widetilde{V}_{\geq 4}(Z; x)\xi + \widetilde{a}_{\geq 4}^{(\alpha)}(Z; x, \xi)$, we claim that

$$\Pi^{\top} \operatorname{Op}^{BW}(\operatorname{i} \mathfrak{m}) \Pi^{\top} = \Pi^{\top} \operatorname{Op}^{BW}(\operatorname{i} \mathfrak{m}_{\geq 4}) \Pi^{\top} , \qquad (5.9)$$

$$\Pi^{\top} \operatorname{Op}^{BW}(\operatorname{im}) \Pi^{\perp} = \Pi^{\perp} \operatorname{Op}^{BW}(\operatorname{im}) \Pi^{\top} = 0 , \qquad (5.10)$$

$$\Pi^{\perp} \operatorname{Op}^{BW}(\operatorname{i} \mathfrak{m}) \Pi^{\perp} = \operatorname{Op}^{BW}(\operatorname{i} \mathfrak{m}) \Pi^{\perp} .$$
(5.11)

Proof of (5.9). We shall exploit that the symbol $m_2(z; x, \xi)$ has zero average in x (see Theorem 4.3). Using the definition (2.21) for 2-homogeneous paradifferential operators applied to the quadratic, gauge invariant, zero-average symbol $m_2(z; \cdot)$ we get

$$\Pi^{\top} \operatorname{Op}^{BW}(\operatorname{im}_{2}(z; x, \xi)) \Pi^{\top} z = \sum_{\substack{j_{1} - j_{2} + j = k \\ j_{1} \neq j_{2}, \ j, k \in \Lambda}} \chi_{2}\left(j_{1}, j_{2}, \frac{j+k}{2}\right) \operatorname{im}_{j_{1}, j_{2}}^{+, -}\left(\frac{j+k}{2}\right) z_{j_{1}} \overline{z}_{j_{2}} z_{j} e^{\operatorname{i}kx}$$

We show that the cut-off is always vanishing. Indeed, recalling that $\chi_2(\xi',\xi) \equiv 0$ when $|\xi'| \equiv \max(|\xi'_1|, |\xi'_2|) \geq \langle \xi \rangle / 10$, and using $\max(|j_1|, |j_2|) \geq 1$ (as j_1, j_2 cannot be both 0), $j = k - j_1 + j_2$ and $k \in \Lambda = \{\pm 1\}$, one has

$$\frac{1}{10} \langle \frac{j_1 - j_2 \pm 2}{2} \rangle = \frac{1}{10} \left(1 + \frac{|j_1 - j_2 \pm 2|}{2} \right) \le \frac{4 + 2\max(|j_1|, |j_2|)}{20} \le \frac{3\max(|j_1|, |j_2|)}{10} \le \max(|j_1|, |j_2|),$$
(5.12)

proving that $\chi_2\left(j_1, j_2, \frac{j+k}{2}\right) \equiv 0$. Consequently $\Pi^{\top} \operatorname{Op}^{BW}(\operatorname{im}_2) \Pi^{\top} = 0$ and (5.9) follows.

Proof of (5.10). Again we write explicitly the action of $\Pi^{\top} \operatorname{Op}^{BW}(\operatorname{im}) \Pi^{\perp}$, using the quantization (2.21) for the 2-homogeneous symbol $\mathfrak{m}_2(z;\cdot)$ and (2.22) for the non-homogeneous symbol $\mathfrak{m}_{\geq 4}(z;\cdot)$, getting

$$\Pi^{\top} \operatorname{Op}^{BW}(\operatorname{im}(z;\cdot)) \Pi^{\perp} z = \sum_{\substack{j_{1}-j_{2}+j=k\\j_{1}\neq j_{2}, \ j\in\Lambda^{c}, k\in\Lambda}} \chi_{2} \left(j_{1}, j_{2}, \frac{j+k}{2}\right) \operatorname{im}_{j_{1}, j_{2}}^{+, -} \left(\frac{j+k}{2}\right) z_{j_{1}} \overline{z}_{j_{2}} z_{j} e^{\operatorname{i}kx} + \sum_{j\in\Lambda^{c}, k\in\Lambda} \chi\left(k-j, \frac{j+k}{2}\right) \operatorname{im}_{\geq 4}\left(z; k-j, \frac{k+j}{2}\right) z_{j} e^{\operatorname{i}kx} .$$
(5.13)

Arguing as in (5.12), the first line of (5.13) vanishes. To deal with the second line, recall that also $\chi(\xi',\xi) \equiv 0$ when $|\xi'| \ge \langle \xi \rangle / 10$, so when $k \in \Lambda$ and $j \in \Lambda^c$ (so $|j-k| \ge 1$)

$$\frac{1}{10} \langle \frac{j+k}{2} \rangle = \frac{1}{10} \left(1 + \frac{|j\pm 1|}{2} \right) \le \frac{3+|j|}{20} \le \frac{4+|j-k|}{20} \le \frac{|j-k|}{4} \le |j-k| ,$$

proving that $\chi\left(k-j,\frac{j+k}{2}\right) \equiv 0$. In conclusion, also the second line of (5.13) vanishes, proving the first of (5.10). The second identity is analogous exchanging the roles of j and k. Proof of (5.11). It follows writing $\Pi^{\perp} = \mathrm{Id} - \Pi^{\top}$ and using the first of (5.10).

This concludes the analysis of the projection of the paradifferential vector field $Op^{BW}(im) z$.

We pass to the cubic vector field $X^{(\Lambda)}(Z)$ in (4.8). We set

$$Y_3^{(\Lambda)}(z) := (\Pi_{\mathcal{P}_\Lambda^{(0)}} X^{(\Lambda)})(Z)^+$$

which has the claimed form (5.5) in view of (4.9). Then we put

$$Y_3^{\top}(z) := \Pi^{\top} \left(X^{(\Lambda)}(Z)^+ - (\Pi_{\mathcal{P}_{\Lambda}^{(0)}} X^{(\Lambda)})(Z)^+ \right) , \quad Y_3^{\perp}(z) := \Pi^{\perp} X^{(\Lambda)}(Z)^+.$$

To prove estimates (5.6) we exploit that $X^{(\Lambda)}(Z)$ is in strong- Λ normal form, see (4.9). Estimate of $Y_3^{\top}(z)$. By definition

$$Y_{3}^{\top}(z) = \sum_{k \in \Lambda} \sum_{(\vec{j}, k, \vec{\sigma}, -) \in \mathcal{P} \setminus \mathcal{P}_{\Lambda}^{(0)}} X_{\vec{j}, k}^{\vec{\sigma}, +} z_{\vec{j}}^{\vec{\sigma}} e^{ikx} , \quad \vec{j} = (j_1, j_2, j_3), \quad \vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$$

By (4.9), $\Pi_{\mathcal{P}^{(1)}_{\Lambda}}X^{(\Lambda)} = \Pi_{\mathcal{P}^{(2)}_{\Lambda}}X^{(\Lambda)} = 0$, so, since $k \in \Lambda$, the only possibly remaining monomials are those with $(\vec{j}, k, \vec{\sigma}, -) \in \mathcal{P}_{\Lambda}^{(3)}$ and in addition $\vec{j} \in (\Lambda^c)^3$. Then, recalling (4.8), $Y_3^{\top}(z) =$ $\Pi^{\top} \left(R^{(\Lambda)}(Z^{\perp}) Z^{\perp} \right)^{+}, Z^{\perp} := \left(\frac{z^{\perp}}{z^{\perp}} \right), \text{ and the first estimate (5.6) follows from } \|Y_{3}^{\top}(z)\|_{\mathfrak{s}} \lesssim \|Y_{3}^{\top}(z)\|_{L^{2}}$ and estimate (2.35).

Estimate of $Y_3^{\perp}(z)$. Again by (4.9), we expand $Y_3^{\perp}(z)$ as

$$Y_3^{\perp}(z) = \sum_{k \in \Lambda^c} \sum_{(\vec{j}, k, \vec{\sigma}, -) \in \mathcal{P}_{\Lambda}^{(3)} \cup \mathcal{P}_{\Lambda}^{(4)}} X_{\vec{j}, k}^{\vec{\sigma}, +} z_{\vec{j}}^{\vec{\sigma}} e^{ikx}$$

Then either (i) two indexes among (j_1, j_2, j_3) belong to Λ^c and one to Λ , or (ii) all three indexes belong to Λ^c . Consequently $Y_3^{\perp}(z) = \Pi^{\perp} \left(R^{(\Lambda)}(Z^{\perp}) Z^{\perp} + R^{(\Lambda)}(Z^{\perp}) Z^{\top} + 2R^{(\Lambda)}(Z^{\perp}, Z^{\top}) Z^{\perp} \right)^+$, $Z^{\top} := \left(\frac{z}{z}^{\top}\right)$. The second estimate (5.6) follows again from estimate (2.35) (with $m \rightsquigarrow -4$), using also the trivial bound $\|z^{\top}\|_{\mathfrak{s}} \leq C_{\mathfrak{s},\mathfrak{s}_0}\|z^{\top}\|_{\mathfrak{s}_0}$. This concludes the analysis of the projection of $X^{(\Lambda)}(Z).$

Finally we consider the projections of the vector field $\widetilde{B}_{>4}(Z)Z$ in (4.6). We put

$$Y_{\geq 5}^{\top}(z) := \Pi^{\top} \big(\widetilde{B}_{\geq 4}(Z) Z \big)^{+} + \Pi^{\top} \operatorname{Op}^{BW}(\operatorname{im}_{\geq 4}) \Pi^{\top} z , \quad Y_{\geq 5}^{\perp}(z) := \Pi^{\perp} \big(\widetilde{B}_{\geq 4}(Z) Z \big)^{+} .$$

Estimate of $Y_{\geq 5}^{\perp}(z)$. It follows since $\widetilde{B}_{\geq 4}(Z)$ is a matrix of non-homogeneous 0-operators in $\mathcal{M}_{\geq 4}^0[r]$, see (2.32). Estimate of $Y_{\geq 5}^{\top}(z)$. As the previous one, using also (2.29) and $\|\Pi^{\top} z\|_{s} \lesssim \|z\|_{s-1}$.

The next step is to extract an effective system driving the dynamics of particular solutions of (5.3)-(5.4) which we call *long-time controlled*, see Definition 5.2 below. These solutions have two main features: (i) the initial data is supported mostly on Λ and (ii) they have a large a-priori bound on the high norm $\|\cdot\|_s$ for long times. These features allow us to propagate smallness of both tangential and normal modes in the low norm $\|\cdot\|_{\mathfrak{s}_0}$ for long times, and moreover to ensure that the normal modes keep having a size much smaller than the tangential ones, i.e. $\|z^{\perp}(t)\|_{\mathfrak{s}_0} \ll \|z^{\top}(t)\|_{L^2}$, see (5.17), (5.18). This is possible because of the normal form procedure of the previous section, and in particular because

- (i) the leading term in the dynamics of the low modes $z^{\top}(t)$ in (5.3) is the cubic integrable vector field $Y_3^{(\Lambda)}(z^{\top})$ (the non-explicit cubic term $Y_3^{\top}(z) = \mathcal{O}((z^{\perp})^3)$, hence its size is much smaller);
- (ii) in equation (5.4) for $z^{\perp}(t)$, the term $\operatorname{Op}^{BW}(\operatorname{im}(z; x, \xi)) z^{\perp}$ is skew-adjoint, hence it vanishes in a L^2 -energy estimate; consequently the dominant term becomes $Y_3^{\perp}(z)$ which, in view of (5.6), fulfills the quadratic estimate $||Y_3^{\perp}(z)||_{\mathfrak{s}_0} \lesssim ||z^{\top}||_{\mathfrak{s}_0} ||z^{\perp}||_{\mathfrak{s}_0}^2$ and therefore has a very small size. To obtain such estimate is the reason why we put $X^{(\Lambda)}(Z)$ in (4.8) in strong- Λ normal form, namely it does not contain monomials of the form $z_{j_1}^{\sigma_1} z_{j_2}^{\sigma_2} z_{j_3}^{\sigma_3} e^{ijx}$ supported in $\mathcal{P}^{(2)}_{\Lambda}$. Otherwise, $Y_3^{\perp}(z)$ would have had monomials with exactly two frequencies among (j_1, j_2, j_3) in Λ and one in Λ^c , and the estimate in (5.6) would have had an additional term $\|z^{\top}\|_{\mathfrak{s}_0}^2 \|z^{\perp}\|_{\mathfrak{s}}$, which is too large for the bootstrap lemma 5.3 below.

We now introduce precisely the notion of long-time controlled solutions.

Definition 5.2 (Long-time controlled solutions). Let s, θ as in (5.2). Let also $T_* > 0$ and $\epsilon \in (0, \mathfrak{r})$. We say that a solution $z(t) \in H^s(\mathbb{T}, \mathbb{C})$ of system (5.3)–(5.4) is long-time controlled with parameters $(s, \theta, T_*, \epsilon)$ if

(A1) at time 0 fulfills

$$||z^{\top}(0,\cdot)||_{L^2} \le \epsilon$$
, $||z^{\perp}(0,\cdot)||_{L^2} \le \epsilon^3$; (5.14)

(A2) it exists over the time interval $[0, T_{\star}]$ where it fulfills the large a-priori bound

$$\sup_{0 \le t \le T_{\star}} \|z(t)\|_s \le \epsilon^{-\theta} .$$
(5.15)

One crucial property of any long-time controlled solution is that its low norm $\|\cdot\|_{\mathfrak{s}_0}$ is automatically small for all $0 \leq t \leq T_{\star}$, as we shall now prove.

Lemma 5.3 (Bootstrap lemma). Let s, θ as in (5.2). Fix also $T_0 > 0$. There exists $\epsilon_{\star} = \epsilon_{\star}(\theta, T_0) > 0$ such that for any $\epsilon \in (0, \epsilon_{\star})$ the following holds true.

Let z(t) be a solution of (5.3)–(5.4) which is long-time controlled with parameters $(s, \theta, T_{\star}, \epsilon)$ (according to Definition 5.2) and with

$$T_{\star} \le \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right) \quad . \tag{5.16}$$

Then z(t) fulfills the improved L^2 -bound

$$||z^{\top}(t)||_{L^2} \le 2\epsilon$$
, $||z^{\perp}(t)||_{L^2} \le \epsilon^{3-\frac{3}{2}\theta}$, $\forall 0 \le t \le T_{\star}$ (5.17)

and the improved low-norm bound

$$||z(t)||_{\mathfrak{s}_0} \le 3\epsilon$$
, $||z^{\perp}(t)||_{\mathfrak{s}_0} \le \epsilon^2$, $\forall 0 \le t \le T_{\star}$. (5.18)

Proof. The proof is by a bootstrap argument. We assume the bound

$$||z^{\top}(t)||_{L^2} \le 10\epsilon, \quad ||z^{\perp}(t)||_{L^2} \le \epsilon^{3-2\theta}, \quad \forall 0 \le t \le T_{\star}$$
 (5.19)

and show that, provided $\epsilon \in (0, \epsilon_*)$ with ϵ_* sufficiently small, the better bound (5.17) holds.

First we bound $||z^{\perp}(t)||_{\mathfrak{s}_0}$. This is done interpolating the bound on $||z^{\perp}(t)||_{L^2}$ that we have by the bootstrap assumption (5.19) and the large bound that we have on $||z^{\perp}(t)||_s$ in (5.15), being z(t)long-time controlled by assumption. We obtain

$$\|z^{\perp}(t)\|_{\mathfrak{s}_{0}} \le \|z^{\perp}(t)\|_{L^{2}}^{1-\frac{\mathfrak{s}_{0}}{s}} \|z^{\perp}(t)\|_{s}^{\frac{\mathfrak{s}_{0}}{s}} \le \epsilon^{3-\theta(2-\frac{\mathfrak{s}_{0}}{s})-3\frac{\mathfrak{s}_{0}}{s}} \le \epsilon^{2}$$
(5.20)

which is possible for s, θ as in (5.2). Using again the first of (5.19) we also get

$$||z(t)||_{\mathfrak{s}_0} \le 11\epsilon , \qquad \forall 0 \le t \le T_\star . \tag{5.21}$$

Next we consider $||z^{\top}(t)||_{L^2}$ and prove the improved estimate (5.17). Recall that the function $z^{\top}(t)$ fulfills equation (5.3); since $Y_3^{(\Lambda)}(z)$ is integrable, we get that for all times $0 \le t \le T_{\star}$

$$\frac{\mathrm{d}}{\mathrm{d}t} \|z^{\top}(t)\|_{L^{2}}^{2} = 2 \underbrace{\operatorname{Re}\langle -\mathrm{i}|D|^{\alpha}z^{\top} + Y_{3}^{(\Lambda)}(z), z^{\top}\rangle}_{=0} + 2\operatorname{Re}\langle Y_{3}^{\top}(z) + Y_{\geq 5}^{\top}(z), z^{\top}\rangle \\ \stackrel{(5.6),(5.8)}{\leq} C(\|z^{\perp}(t)\|_{\mathfrak{s}_{0}}^{3} + \|z(t)\|_{\mathfrak{s}_{0}}^{5}) \|z^{\top}(t)\|_{L^{2}} \stackrel{(5.20),(5.21),(5.19)}{\leq} C\epsilon^{6}.$$

Then, since z(t) is long-time controlled, its initial datum $z^{\top}(0)$ is bounded by (5.14); hence for all times $0 \le t \le T_{\star} \le \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right)$,

$$\|z^{\top}(t)\|_{L^{2}}^{2} \leq \|z^{\top}(0)\|_{L^{2}}^{2} + |t|C\epsilon^{6} \leq \epsilon^{2} + CT_{0}\epsilon^{4}\log(\epsilon^{-1}) \leq 4\epsilon^{2}$$
(5.22)

provided $0 < \epsilon \leq \epsilon_{\star}$ and ϵ_{\star} is sufficiently small. This proves the first estimate in (5.17).

Next we bound $||z^{\perp}(t)||_{L^2}$. We exploit that the paradifferential operator in equation (5.4) is skew-adjoint, so we get, for all times $0 \le t \le T_{\star} \le \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right)$,

$$\frac{\mathrm{d}}{\mathrm{d}t} \|z^{\perp}(t)\|_{L^{2}}^{2} = 2 \underbrace{\operatorname{Re}\left(\left(-\mathrm{i}|D|^{\alpha} + \operatorname{Op}^{BW}(\mathrm{im}(z;\cdot))\right)z^{\perp}, z^{\perp}\right)}_{=0} + 2\operatorname{Re}\left(Y_{3}^{\perp}(z) + Y_{\geq 5}^{\perp}(z), z^{\perp}\right)}_{=0} \\ \underbrace{\sum_{i=0}^{(5.6), (5.8)} C\left(\|z(t)\|_{\mathfrak{s}_{0}} \|z^{\perp}(t)\|_{\mathfrak{s}_{0}}^{2} + \|z(t)\|_{\mathfrak{s}_{0}}^{5}\right)}_{\leq} \|z^{\perp}(t)\|_{0} \\ \underbrace{\sum_{i=0}^{(5.21), (5.20), (5.19)} C\epsilon^{8-2\theta}}_{\leq} .$$

Again, being z(t) long-time controlled, its initial datum $z^{\perp}(0)$ fulfills (5.14); hence for all times $0 \le t \le T_{\star} \le \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right)$ we bound

$$\|z^{\perp}(t)\|_{L^{2}}^{2} \leq \|z^{\perp}(0)\|_{L^{2}}^{2} + |t|C\epsilon^{8-2\theta} \leq \epsilon^{6} + CT_{0}\epsilon^{6-2\theta}\log(\epsilon^{-1}) \leq \epsilon^{2(3-\frac{3}{2}\theta)} , \qquad (5.23)$$

which is true shrinking ϵ_{\star} . Estimates (5.22) and (5.23) prove (5.17). Then, again by interpolation, we obtain the second of (5.18), which, together with (5.17), gives also the first of (5.18).

A second important property of any long-time controlled solution is that it fulfills an *effective* equation with a very precise structure: up to higher order corrections, for long times, the modes $z_{\pm 1}(t)$ rotate with constant speed, whereas $z^{\perp}(t)$ fulfills a linear Schrödinger equation whose Hamiltonian $-i|D|^{\alpha} + iOp^{BW}(v(x - J_1t)\xi)$ does not have constant coefficients. We shall show, in the next section, that this Hamiltonian is actually responsible for the growth of Sobolev norms of the solution. Precisely we prove the following result:

Proposition 5.4. Let s, θ as in (5.2). Fix also $T_0 > 0$. There exists $\epsilon_{\star} = \epsilon_{\star}(s, \theta, T_0) > 0$ such that for any $\epsilon \in (0, \epsilon_{\star})$ the following holds true. Let z(t) be a solution of (5.3)–(5.4) which is long-time controlled with parameters $(s, \theta, T_{\star}, \epsilon)$ (see Definition 5.2) and with T_{\star} fulfilling (5.16). Then $z(t) = (z_1(t), z_{-1}(t), z^{\perp}(t))$ fulfills the system

$$\begin{cases} \partial_t z_1 = -i(1+|z_1(0)|^2)z_1 + d_1(t) \\ \partial_t z_{-1} = -i(1-|z_{-1}(0)|^2)z_{-1} + d_{-1}(t) \\ \partial_t z^{\perp} = -i|D|^{\alpha} z^{\perp} + iOp^{BW}(\mathfrak{v}(x-J_1t)\xi + \mathfrak{V}(t;x)\xi + \mathfrak{b}(t;x,\xi)) z^{\perp} + Y(t) \end{cases}$$
(5.24)

where

• J_1 is the real number

$$J_1 := \frac{|z_1(0)|^2 + |z_{-1}(0)|^2}{2} , \qquad (5.25)$$

• the real valued function $\mathfrak{v}(x)$ is given by

$$\mathfrak{v}(x) := 2\operatorname{Re}\left(z_1(0)\,\overline{z_{-1}(0)}\,e^{\mathrm{i}2x}\right) \tag{5.26}$$

whereas the real valued, time dependent function V(t;x) fulfills the estimate

$$\|\mathbf{V}(t;\cdot)\|_{W^{2,\infty}} \le C\epsilon^{4-\theta} , \quad \forall 0 \le t \le T_{\star} ;$$
(5.27)

• the real valued symbol $\mathbf{b}(t; x, \xi) \in \Gamma_{W^{2,\infty}}^{\alpha}$ fulfills the estimate (recall (2.13)): for every $n \in \mathbb{N}_0$, there is $C_n > 0$ such that

$$\left| \mathbf{b}(t; \cdot) \right|_{\alpha, W^{2, \infty}, n} \le C_n \epsilon^2 , \quad \forall 0 \le t \le T_\star ; \tag{5.28}$$

• the functions $d_{\pm 1}(t)$ fulfill the estimates

$$|\mathbf{d}_{\pm 1}(t)| \le \epsilon^{5-\theta} , \quad \forall 0 \le t \le T_{\star} ;$$
(5.29)

• the vector field $Y(t) \equiv Y(t, x)$ fulfills the estimate

$$||Y(t; \cdot)||_s \le C\epsilon^{3-\theta} , \quad \forall 0 \le t \le T_\star .$$
(5.30)

Proof. We shall use that z(t), being long-time controlled with parameters $(s, \theta, T_{\star}, \epsilon)$ and with T_{\star} fulfilling (5.16), satisfies the bounds (5.17), (5.18).

Equations for $z_{\pm 1}(t)$. Write equation (5.3) in components, using the explicit expression of $Y_3^{(\Lambda)}$ in (5.5), to get the coupled system

$$\begin{cases} \partial_t z_1 = -iz_1 - i|z_1|^2 z_1 + \langle Y_3^{\top}(z) + Y_{\geq 5}^{\top}(z), e^{ix} \rangle \\ \partial_t z_{-1} = -iz_{-1} + i|z_{-1}|^2 z_{-1} + \langle Y_3^{\top}(z) + Y_{\geq 5}^{\top}(z), e^{-ix} \rangle \end{cases}$$
(5.31)

Consider the equation for z_1 . We write it as

$$\partial_t z_1 = -i(1+|z_1(0)|^2)z_1 + d_1(t), d_1(t) := -i\left(|z_1(t)|^2 - |z_1(0)|^2\right)z_1(t) + \langle Y_3^{\top}(z) + Y_{\geq 5}^{\top}(z), e^{ix}\rangle$$
(5.32)

giving the first equation in (5.24). We prove now that $d_1(t)$ fulfills the bound claimed in (5.29). First, using the first of (5.31) and assumption (5.16), we get for all times $0 \le t \le T_{\star} \le \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right)$

$$\frac{\mathrm{d}}{\mathrm{d}t}|z_1(t)|^2 = 2\mathrm{Re}\left(\langle Y_3^\top(z) + Y_{\geq 5}^\top(z), e^{ix}\rangle \overline{z}_1\right)$$

$$\stackrel{(5.6),(5.8)}{\leq} C\left(\|z^\perp(t)\|_{\mathfrak{s}_0}^3 + \|z(t)\|_{\mathfrak{s}_0}^5\right)\|z^\top(t)\|_0 \stackrel{(5.18),(5.21),(5.17)}{\leq} C\epsilon^6 ,$$

which implies, on the same time scale,

$$\left| \left(|z_1(t)|^2 - |z_1(0)|^2 \right) \right| \le C |t| \epsilon^6 \le C T_0 \, \epsilon^4 \log(\epsilon^{-1}) \, . \tag{5.33}$$

Hence we get that $d_1(t)$ in (5.32) is bounded for $0 \le t \le T_\star \le \frac{T_0}{\epsilon^2} \log(\epsilon^{-1})$ by

$$|\mathbf{d}_{1}(t)| \leq \left| \left(|z_{1}(t)|^{2} - |z_{1}(0)|^{2} \right) z_{1}(t) \right| + \left| \left\langle Y_{3}^{\top}(z) + Y_{\geq 5}^{\top}(z), e^{\mathrm{i}x} \right\rangle \right| \stackrel{(5.33), (5.17)}{\leq} CT_{0} \, \epsilon^{5} \log(\epsilon^{-1}) + C\epsilon^{5} \, , \tag{5.34}$$

proving (5.29) provided ϵ_{\star} is sufficiently small. An analogous argument proves that $z_{-1}(t)$ fulfills the second of (5.24).

A consequence, which we shall use in a moment, is that

$$z_{\pm 1}(t) = \mathbf{z}_{\pm 1}(t) + r_{\pm 1}(t) , \quad \text{where } \mathbf{z}_{\pm 1}(t) := e^{-it(1\pm|z_{\pm 1}(0)|^2)} z_{\pm 1}(0)$$
(5.35)

whereas

$$r_{\pm 1}(t) := \int_0^t e^{-\mathbf{i}(t-\tau)(1\pm|z_{\pm 1}(0)|^2)} \, \mathbf{d}_{\pm 1}(\tau) \, \mathrm{d}\tau$$

fulfill, by (5.34), (5.16) and eventually shrinking again ϵ_{\star} , the bounds

$$|r_{\pm 1}(t)| \le \epsilon^{3-\theta}, \qquad \forall 0 \le t \le T_{\star} . \tag{5.36}$$

Equation for $z^{\perp}(t)$. We start from equation (5.4) and we substitute the explicit expression of $z_{\pm}1(t)$ in (5.35). Consider first the symbol $\mathfrak{m}(z; x, \xi)$ in (5.7). We shall extract from its component $\langle \underline{V} \rangle(Z; x)$, defined in (4.7), the main contribution which is the one supported on $z_{\pm 1}(t)$. Precisely

$$\begin{split} \langle \underline{\mathbf{V}} \rangle (Z(t); x) &= 2 \operatorname{Re} \left(z_1(t) \, \overline{z_{-1}(t)} \, e^{\mathrm{i} 2x} \right) + 2 \operatorname{Re} \left(\sum_{n \ge 2} z_n(t) \, \overline{z_{-n}(t)} \, e^{\mathrm{i} 2nx} \right) \\ & \stackrel{(5.35)}{=} \underbrace{2 \operatorname{Re} \left(z_1(0) \, \overline{z_{-1}(0)} \, e^{\mathrm{i} 2x - 2 J_1 t} \right)}_{= \mathfrak{v}(x - J_1 t) \text{ by } (5.26), (5.25)} + \underbrace{2 \operatorname{Re} \left(\left(z_1(t) \, \overline{r_{-1}(t)} + r_1(t) \overline{z_{-1}(t)} + r_1(t) \overline{r_{-1}(t)} \right) e^{\mathrm{i} 2x} \right)}_{=: \mathbb{V}_1(t; x)} \\ & + \underbrace{2 \operatorname{Re} \left(\sum_{n \ge 2} z_n(t) \, \overline{z_{-n}(t)} \, e^{\mathrm{i} 2nx} \right)}_{=: \mathbb{V}_2(t; x)} . \end{split}$$

The functions $V_1(t;x)$ and $V_2(t;x)$ fulfill, by (5.14), (5.36) and (5.18), the bounds

$$\|\mathbf{V}_1(t;\cdot)\|_{W^{2,\infty}} \le C\epsilon^{4-\theta} , \quad \|\mathbf{V}_2(t;\cdot)\|_{W^{2,\infty}} \le C\epsilon^4 , \quad \forall 0 \le t \le T_\star .$$

$$(5.37)$$

Then we write $m(z; \cdot)$ in (5.7) as

$$\mathbf{m}(z(t); x, \xi) = \mathbf{v}(x - \mathbf{J}_1 t)\xi + \underbrace{(\mathbf{V}_1(t; x) + \mathbf{V}_2(t; x) + \widetilde{V}_{\geq 4}(z(t); x))}_{=:\mathbf{V}(t; x)} \xi + \underbrace{a_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{a}_{\geq 4}^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \widetilde{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{b}(t; x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi) + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{M}_2^{(\alpha)}(x(t); x, \xi)}_{=:\mathbf{M}_2^{(\alpha)}(x(t); x, \xi)} \xi + \underbrace{\mathbf{M}_2^{(\alpha)}(z(t); x, \xi)}_{=:\mathbf{M}_2^{(\alpha)}(x(t); x, \xi)}_{=:\mathbf{M}_2^$$

We bound V(t; x) using estimates (5.37) for V_1 and V_2 , and that

$$\|\widetilde{V}_{\geq 4}(z(t); \cdot)\|_{W^{2,\infty}} \stackrel{(2.16)}{\leq} C \|z(t)\|_{\mathfrak{s}_0}^4 \stackrel{(5.18)}{\leq} C\epsilon^4 , \quad \forall 0 \leq t \leq T_{\star} ,$$

getting the claimed bound (5.27).

The bound (5.28) for $b(t; x, \xi)$ follows from (2.19), (2.16) and (5.18).

Finally we put

$$Y(t,z) := Y_3^{\perp}(z(t)) + Y_{\geq 5}^{\perp}(z(t))$$

which fulfills the estimates (5.30) by (5.6), (5.8) and using (5.18) and (5.15).

6 Instability via paradifferential Mourre theory

The goal of this section is to give sufficient conditions on the initial datum z(0) ensuring that, if the corresponding solution z(t) is long-time controlled, than its high H^s -norm undergoes Sobolev norm explosion, becoming larger than $e^{-\theta}$. We will achive this via a positive commutator estimate.

We will focus on the third equation in (5.24); actually it is more convenient to work with the translated variable

$$\zeta(t,x) := z^{\perp}(t,x+J_1t) , \quad J_1 \text{ in } (5.25) .$$
(6.1)

Clearly one has

 $\|\zeta(t,\cdot)\|_s = \|z^{\perp}(t,\cdot)\|_s , \quad \forall t, \quad \forall s \in \mathbb{R} ,$

so it is equivalent to prove growth of Sobolev norms for $\zeta(t)$ and $z^{\perp}(t)$. The equation fulfilled by $\zeta(t)$ is easily derived from the third of (5.24) as

$$\partial_t \zeta = -i|D|^{\alpha} \zeta + iOp^{BW}((\mathbf{J}_1 + \boldsymbol{\mathfrak{v}}(x))\xi)\zeta + iOp^{BW}(\widetilde{\mathbf{V}}(t;x)\xi + \widetilde{\mathbf{b}}(t;x,\xi))\zeta + \widetilde{Y}(t)$$
(6.2)

where we defined the real valued function $\widetilde{V}(t;x)$, the real valued symbol $\widetilde{b}(t;x,\xi)$ and the vector field $\widetilde{Y}(t;x)$ as

$$\widetilde{\mathtt{V}}(t;x):=\mathtt{V}(t,x+\mathtt{J}_1t)\;,\quad \widetilde{\mathtt{b}}(t;x,\xi):=\mathtt{b}(t;x+\mathtt{J}_1t,\xi)\;,\quad \widetilde{Y}(t;x):=Y(t;x+\mathtt{J}_1t)\;.$$

It follows, by (5.27), (5.28) and (5.30), the estimates

$$\|\widetilde{\mathsf{V}}(t;\cdot)\|_{W^{2,\infty}} \le C\epsilon^{4-\theta} , \quad |\widetilde{\mathsf{b}}(t;\cdot)|_{\alpha,W^{2,\infty},n} \le C_n\epsilon^2 , \quad \|\widetilde{Y}(t;\cdot)\|_s \le C\epsilon^{3-\theta} , \quad \forall \ 0 \le t \le T_\star .$$
(6.3)

6.1 The Mourre operator

The leading term in equation (6.2) is the non-constant coefficient transport operator

Op^{BW}(
$$(J_1 + \mathfrak{v}(x))\xi$$
), J_1 in (5.25), $\mathfrak{v}(x)$ in (5.26). (6.4)

The crucial point is that, provided $z_1(0)$ and $z_{-1}(0)$ fulfill

$$J_1 \equiv \frac{|z_1(0)|^2 + |z_{-1}(0)|^2}{2} < 2|z_1(0)| |z_{-1}(0)| ,$$

corresponding to the function $J_1 + \mathfrak{v}(x)$ having a zero, the operator $\operatorname{Op}^{BW}((J_1 + \mathfrak{v}(x))\xi)$ admits a Mourre-conjugate operator, namely there exists an operator A such that the commutator $i[A, \operatorname{Op}^{BW}((J_1 + \mathfrak{v}(x))\xi)]$ is positive. Actually this also shows that the operator in (6.4) has a non-trivial absolutely continuous spectrum, although we shall not exploit directly this property.

Precisely, take s as in (5.2) and $\mathbb{R} \gg 1$ (to be fixed later) and define the *self-adjoint* operator

$$A := A_{s,\mathbf{R}} := \operatorname{Op}^{BW}(\mathfrak{a}(x,\xi)), \quad \mathfrak{a}(x,\xi) := \mathbf{a}(x) \, |\xi|^{2s} \, \eta_{\mathbf{R}}^{2}(\xi)$$

where $\mathbf{a}(x) := -\operatorname{Im}\left(z_{1}(0) \, \overline{z_{-1}(0)} \, e^{\mathrm{i}2x}\right)$ (6.5)

and $\eta_{\mathbf{R}}(\xi)$ the smooth step function

$$\eta_{\mathbf{R}}(\xi) := \eta\left(\frac{\xi}{\mathbf{R}}\right), \quad \eta(y) := \begin{cases} 0 & \text{if } y \le 1\\ \frac{e^{-\frac{1}{y-1}}}{e^{-\frac{1}{y-1}} + e^{-\frac{1}{2-y}}} & \text{if } y \in (1,2) \\ 1 & \text{if } y \ge 2 \end{cases}$$
(6.6)

Note that $\mathfrak{a}(x,\xi)$ is a symbol in $\Gamma^{2s}_{W^{2,\infty}}$, and for any $n \in \mathbb{N}_0$, there is $C_n > 0$ such that

$$|\mathfrak{a}|_{2s,W^{2,\infty},n} \le C_{s,n} |z_1(0)| |z_{-1}(0)| , \qquad |\mathfrak{a}|_{2s+1,W^{2,\infty},n} \le C_{s,n} \frac{|z_1(0)| |z_{-1}(0)|}{\mathfrak{R}} , \qquad (6.7)$$

as it follows from its definition and from Lemma A.1 with $a \to a(x)|\xi|^{2s}\eta_{\mathbb{R}}(\xi)$, $m \to 2s$, $N \to 2$ and $\nu \to 1$. Moreover we will ensure that $|z_1(0)z_{-1}(0)| > 0$, so that A is non trivial, see Remark 6.4.

The choice of the function $\mathfrak{a}(x,\xi)$ in (6.5) is motivated by the fact that it is an escape function for the symbol $(J_1 + \mathfrak{v}(x))\xi$ of the operator in (6.4); precisely one has the following result:

Lemma 6.1. Fix $s, \mathbb{R} > 1$. Let $\mathfrak{a}(x, \xi)$ as in (6.5) and J_1 , $\mathfrak{v}(x)$ as in (5.25), (5.26). Then

$$\{\mathfrak{a}(x,\xi), \, (\mathsf{J}_1 + \mathfrak{v}(x))\xi\} = \mathsf{I}_1 \, |\xi|^{2s} \, \eta^2_\mathsf{R}(\xi) + a(x,\xi) \tag{6.8}$$

where I_1 is the real number

$$I_{1} := 2|z_{1}(0)||z_{-1}(0)|\left(2|z_{1}(0)||z_{-1}(0)| - \frac{|z_{1}(0)|^{2} + |z_{-1}(0)|^{2}}{2}\right)$$
(6.9)

whereas $a(x,\xi)$ is a smooth, non-negative symbol having the structure

$$a(x,\xi) = a_1(x)\psi_1(\xi)^2 + a_2(x)\psi_2(\xi)^2 .$$
(6.10)

Here $a_j(x)$, j = 1, 2, are smooth, real valued, non-negative functions fulfilling

$$||a_j(x)||_{W^{3,\infty}} \le C\left(|z_1(0)|^4 + |z_{-1}(0)|^4\right) , \qquad (6.11)$$

and $\psi_j(\xi)$, j = 1, 2, are smooth, real valued symbols in $\widetilde{\Gamma}_0^s$ with support in $[\mathbb{R}, +\infty)$.

Proof. We compute, using (2.38), (6.5), (5.25), (5.26) and denoting $(\eta')_{\mathbb{R}}(\xi) := \eta'(\xi/\mathbb{R})$,

$$\{\mathfrak{a}(x,\xi), (\mathbf{J}_{1}+\mathfrak{v}(x))\xi\} = (2s\,\mathfrak{a}\,\mathfrak{v}_{x}-\mathfrak{v}\,\mathfrak{a}_{x}-\mathbf{J}_{1}\mathfrak{a}_{x})\,|\xi|^{2s}\,\eta_{\mathsf{R}}^{2} + \frac{2}{\mathsf{R}}\mathfrak{a}\mathfrak{v}_{x}\,|\xi|^{2s}\xi\,\eta_{\mathsf{R}}\,(\eta')_{\mathsf{R}} \\ = (\mathfrak{a}\,\mathfrak{v}_{x}-\mathfrak{v}\,\mathfrak{a}_{x}-\mathbf{J}_{1}\mathfrak{a}_{x})\,|\xi|^{2s}\,\eta_{\mathsf{R}}^{2} + (2s-1)\mathfrak{a}\mathfrak{v}_{x}|\xi|^{2s}\eta_{\mathsf{R}}^{2} + 2\mathfrak{a}\mathfrak{v}_{x}\,|\xi|^{2s}\eta_{\mathsf{R}}\,\frac{\xi}{\mathsf{R}}\,(\eta')_{\mathsf{R}} .$$
(6.12)

Now, using the explicit definition of $\mathbf{a}(x)$ in (6.5), of $\mathbf{v}(x)$ in (5.26) and of \mathbf{J}_1 in (5.25) and that $\mathbf{a}_x(x) = -2\operatorname{Re}\left(z_1(0)\overline{z_{-1}(0)}e^{i2x}\right), \ \mathbf{v}_x(x) = -4\operatorname{Im}\left(z_1(0)\overline{z_{-1}(0)}e^{i2x}\right), \ \text{we get the lower bound}$

$$\begin{aligned} \mathbf{a}\mathfrak{v}_{x} - \mathfrak{v}\mathbf{a}_{x} - \mathbf{J}_{1}\mathbf{a}_{x} &= 4\operatorname{Im}\left(z_{1}(0)\,\overline{z_{-1}(0)}\,e^{\mathrm{i}2x}\right)^{2} + 4\operatorname{Re}\left(z_{1}(0)\,\overline{z_{-1}(0)}\,e^{\mathrm{i}2x}\right)^{2} - \mathbf{a}_{x}\mathbf{J}_{1} \\ &\geq 4\left|z_{1}(0)\right|^{2}\left|z_{-1}(0)\right|^{2} - 2\mathbf{J}_{1}\left|z_{1}(0)\right|\left|z_{-1}(0)\right| \\ &\geq 2\left|z_{1}(0)\right|\left|z_{-1}(0)\right|\left(2\left|z_{1}(0)\right|\left|z_{-1}(0)\right| - \mathbf{J}_{1}\right) \equiv \mathbf{I}_{1}, \end{aligned}$$
(6.13)

where to pass from the first to the second line we also used that

$$|\mathbf{a}_x| \le 2|z_1(0)| |z_{-1}(0)|$$
.

Hence, adding and subtracting $I_1|\xi|^{2s}\eta_{\rm R}^2(\xi)$ in (6.12), we get the claimed formula (6.8) with

$$a(x,\xi) := \underbrace{(\mathbf{a}\mathbf{v}_{x} - \mathbf{v}\mathbf{a}_{x} - \mathbf{J}_{1}\mathbf{a}_{x} - \mathbf{I}_{1} + (2s-1)\mathbf{a}\mathbf{v}_{x})}_{=:a_{1}(x)} \underbrace{|\xi|^{2s}\eta_{\mathbf{R}}^{2}}_{=:\psi_{1}(\xi)^{2}} + \underbrace{2\mathbf{a}\mathbf{v}_{x}}_{=:a_{2}(x)} \underbrace{|\xi|^{2s}\eta_{\mathbf{R}}\frac{\xi}{\mathbf{R}}(\eta')_{\mathbf{R}}}_{=:\psi_{2}(\xi)^{2}} + \underbrace{2\mathbf{a}\mathbf{v}_{x}}_{=:\psi_{2}(\xi)^{2}} + \underbrace{2\mathbf{a}\mathbf{v}_{x}}_{=:\psi_{2}(\xi)^{2}}$$

Note that both $a_1(x)$ and $a_2(x)$ are non-negative functions in view of (6.13) and the fact that $\mathbf{av}_x = 4 \operatorname{Im} \left(z_1(0) \overline{z_{-1}(0)} e^{i2x} \right)^2 \ge 0$. They clearly are smooth, and estimate (6.11) follows from the definitions of $\mathbf{a}(x), \mathbf{v}(x)$ in (6.5), (5.26), of J_1 in (5.25) and I_1 in (6.9).

We claim that the functions $\psi_1(\xi) = |\xi|^s \eta_{\mathbb{R}}$ and $\psi_2(\xi) = |\xi|^s \sqrt{\eta_{\mathbb{R}}} \frac{\xi}{\mathbb{R}}(\eta')_{\mathbb{R}}$ are smooth symbols in $\widetilde{\Gamma}_0^s$ supported in $[\mathbb{R}, \infty)$. We prove the claim only for ψ_2 since the one for ψ_1 is trivial. First notice that ψ_2 is well defined since, by (6.6), one has $\xi(\eta')_{\mathbb{R}} \ge 0$. Define

$$f(y) := \sqrt{\eta(y) y \eta'(y)}$$
, $\operatorname{supp}(f) \subset [1, 2]$.

Then $\psi_2(\xi) = |\xi|^s f(\xi/\mathbb{R})$ and is supported in $[\mathbb{R}, 2\mathbb{R}]$. So we are left to prove that f(y) is a smooth function. It is easy to see that $\sqrt{y\eta(y)}$ is smooth on its support. The function

$$\sqrt{\eta'(y)} = \begin{cases} 0, & y \le 1\\ \frac{\sqrt{2y^2 - 6y + 5}}{e^{-\frac{1}{2-y}} + e^{-\frac{1}{y-1}}} \cdot \frac{e^{-\frac{1}{2(y-1)}}}{y-1} \cdot \frac{e^{-\frac{1}{2(2-y)}}}{2-y}, & y \in (1,2)\\ 0, & y \ge 2 \end{cases}$$

is smooth by direct inspection.

Thanks to Lemma 6.1, we now prove that the commutator between A in (6.5) and $\operatorname{Op}^{BW}((J_1 + \mathfrak{v}(x))\xi)$ is a non-negative operator up to a small remainder. In the following, given two operators A, B, we write $A \ge B$ with the meaning $\langle Au, u \rangle \ge \langle Bu, u \rangle$ for any $u \in \bigcap_s H^s$. Precisely we have:

Lemma 6.2. Fix $s, \mathbb{R} > 1$. Let $A \equiv A_{s,\mathbb{R}}$ be defined in (6.5). Then:

(i) Positive commutator: Let J_1 in (5.25) and v(x) in (5.26). One has

$$i[\mathsf{A}, \operatorname{Op}^{BW}((\mathsf{J}_1 + \mathfrak{v}(x))\xi)] \ge \mathsf{I}_1 \operatorname{Op}^{BW}(|\xi|^{2s}\eta_{\mathsf{R}}^2(\xi)) + \mathsf{R}$$
(6.14)

with I_1 in (6.9) and the operator $R: H^s \to H^{-s}$ with estimate

$$\|\mathsf{R}u\|_{-s} \le C_s \frac{|z_1(0)|^4 + |z_{-1}(0)|^4}{\mathsf{R}} \|u\|_s .$$
(6.15)

(ii) Upper bound: One has

$$\mathsf{A} \le 2|z_1(0)| |z_{-1}(0)| \operatorname{Op}^{^{BW}} \left(|\xi|^{2s} \eta_{\mathsf{R}}^2(\xi) \right) + \mathsf{R}$$
(6.16)

with $R: H^s \to H^{-s}$ satisfying the estimate

$$\|\mathsf{R}u\|_{-s} \le C_s \frac{|z_1(0)|^2 + |z_{-1}(0)|^2}{\mathsf{R}^2} \|u\|_s .$$
(6.17)

Proof. (i) First note that $(J_1 + \mathfrak{v}(x))\xi$ is a symbol in $\Gamma^1_{W^{2,\infty}}$ with seminorm

$$|(\mathbf{J}_1 + \mathbf{v}(x))\xi|_{1,W^{2,\infty},7} \le C\left(|z_1(0)|^2 + |z_{-1}(0)|^2\right) .$$
(6.18)

We now compute the commutator between A and $\operatorname{Op}^{BW}((J_1 + \mathfrak{v}(x))\xi)$. We use the composition Theorem 2.7 (i) regarding $\mathfrak{a}(x,\xi)$ as a symbol in $\Gamma^{2s+1}_{W^{2,\infty}}$ (so putting $m \rightsquigarrow 2s+1, m' \rightsquigarrow 1, \rho \rightsquigarrow 2$); we get

$$\mathbf{i}[\mathsf{A}, \operatorname{Op}^{BW}((\mathsf{J}_1 + \mathfrak{v}(x))\xi)] = \operatorname{Op}^{BW}(\{\mathfrak{a}(x,\xi), (\mathsf{J}_1 + \mathfrak{v}(x))\xi\}) + \check{R}$$
(6.19)

where the operator $\breve{R} \colon H^s \to H^{-s}$ satisfies

$$\|\breve{R}u\|_{-s} \lesssim |\mathfrak{a}|_{2s+1,W^{2,\infty},7} |(\mathsf{J}_1 + \mathfrak{v}(x))\xi|_{1,W^{2,\infty},7} ||u||_s \overset{(6.7),(6.18)}{\lesssim} \frac{|z_1(0)|^4 + |z_{-1}(0)|^4}{\mathsf{R}} ||u||_s .$$

Back to formula (6.19), the Poisson bracket $\{\mathfrak{a}(x,\xi), (J_1 + \mathfrak{v}(x))\xi\}$ was already computed in (6.8), hence

$$\operatorname{Op}^{BW}(\{\mathfrak{a}(x,\xi),(\mathsf{J}_1+\mathfrak{v}(x))\xi\}) = \mathbb{I}_1 \operatorname{Op}^{BW}(|\xi|^{2\mathfrak{s}}\eta_{\mathsf{R}}^2) + \operatorname{Op}^{BW}(a(x,\xi))$$
(6.20)

with $a(x,\xi)$ a smooth, non-negative symbol having the structure (6.10). Thanks to these properties we bound the operator $\operatorname{Op}^{BW}(a)$ from below using the strong Garding inequality A.2, getting

$$\langle \operatorname{Op}^{^{BW}}(a) u, u \rangle \ge -C \frac{\|a_1\|_{W^{3,\infty}} + \|a_2\|_{W^{3,\infty}}}{\mathbb{R}^2} \|u\|_s^2 \stackrel{(6.11)}{\ge} -C \frac{|z_1(0)|^4 + |z_{-1}(0)|^4}{\mathbb{R}^2} \langle \langle D \rangle^{2s} u, u \rangle .$$
(6.21)

We conclude by (6.19), (6.20), (6.21) that

$$i[\mathsf{A}, \operatorname{Op}^{BW}((\mathsf{J}_1 + \mathfrak{v}(x))\xi)] \ge \mathsf{I}_1 \operatorname{Op}^{BW}(|\xi|^{2\mathfrak{s}}\eta_{\mathsf{R}}^2) + \mathsf{R} , \quad \mathsf{R} := \breve{R} - C \frac{|z_1(0)|^4 + |z_{-1}(0)|^4}{\mathsf{R}^2} \langle D \rangle^{2s}$$

where the operator $\mathsf{R}: H^s \to H^{-s}$ fulfills the estimate (6.15).

(*ii*) Define the positive symbol $\tilde{a}(x,\xi) := (2|z_1(0)| |z_{-1}(0)| - \mathbf{a}(x)) |\xi|^{2s} \eta_{\mathbf{R}}^2(\xi)$ and apply again Garding's inequality A.2.

6.2 Growth of Sobolev norms

We now give sufficient conditions on the initial data of a long-time controlled solution z(t) ensuring growth of Sobolev norms.

Definition 6.3 (Well-prepared data). Fix s, θ as in (5.2). Fix also $\nu_0 \in (0, \frac{1}{2}), \epsilon > 0$.

We say that an initial datum $z(0) \in H^s(\mathbb{T}, \mathbb{C})$ is well prepared with parameters $(s, \theta, \nu_0, \epsilon)$ if

(B1) On the modes on Λ

$$2|z_1(0)||z_{-1}(0)| - \frac{|z_1(0)|^2 + |z_{-1}(0)|^2}{2} \ge \nu_0 \epsilon^2 ; \qquad (6.22)$$

(B2) On the modes on Λ^c

$$\langle \mathsf{A}_{s,\mathsf{R}} z^{\perp}(0), z^{\perp}(0) \rangle > \epsilon^{3-3\theta} , \quad with \; \mathsf{R} := \epsilon^{-(3+\theta)/(1-\alpha)}$$
(6.23)

and $A_{s,R}$ in (6.5).

Remark 6.4. Condition (6.22) ensures that $|z_1(0)z_{-1}(0)| > 0$, hence both $\mathfrak{v}(x)$ in (5.26) and the symbol $\mathfrak{a}(x,\xi)$ in (6.5) are non-trivial.

The next result proves that a solution z(t) which is long-time controlled for times $T_0 \epsilon^{-2} \log (\epsilon^{-1})$ with T_0 sufficiently large and whose initial datum is well-prepared, undergoes growth of Sobolev norms. Precisely:

Proposition 6.5. Fix s, θ as in (5.2). Fix also $\nu_0 \in (0, \frac{1}{2})$. There exists $\epsilon_1 = \epsilon_1(s, \theta, \nu_0) > 0$ such that for any $\epsilon \in (0, \epsilon_1)$, the following holds true. Let $z(t) \in H^s(\mathbb{T}, \mathbb{C})$ be a solution of system (5.3)–(5.4) such that

6 INSTABILITY VIA PARADIFFERENTIAL MOURRE THEORY

(i) it is long-time controlled with parameters $(s, \theta, T_{\star}, \epsilon)$ (see Definition 5.2), with

$$T_{\star} = \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right) , \quad T_0 := \frac{1}{\nu_0} ; \qquad (6.24)$$

(ii) its initial datum $z(0) \in H^s(\mathbb{T}, \mathbb{C})$ is well-prepared with parameters $(s, \theta, \nu_0, \epsilon)$ (see Definition 6.3).

Then the solution z(t) undergoes growth of Sobolev norms, i.e.

$$\sup_{|t| \le T_{\star}} \|z(t)\|_s \ge \frac{1}{\epsilon^{\theta}} .$$
(6.25)

The first step to prove such result is to define the \mathcal{A} -functional

$$\mathcal{A}(t) := \langle \mathsf{A}_{s,\mathsf{R}}\,\zeta(t),\zeta(t)\rangle, \quad \mathsf{A}_{s,\mathsf{R}} \text{ in } (6.5) , \quad \zeta(t) \text{ in } (6.1)$$
(6.26)

and exploit Lemma 6.2 to give a *lower bound* on the time derivative $\frac{d}{dt}\mathcal{A}(t)$. Precisely we have:

Lemma 6.6. Under the same assumptions of Propositon 6.5, there are a constant C > 0 and $\epsilon_1 = \epsilon_1(s, \theta, \alpha, \nu_0) > 0$ such that if $\epsilon \in (0, \epsilon_1)$ the \mathcal{A} -functional in (6.26), with \mathbb{R} in (6.23) fulfills: then

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{A}(t) \ge \epsilon^2 \nu_0 \left(\mathcal{A}(t) - C\epsilon^{3-2\theta}\right) , \qquad \forall 0 \le t \le \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right) . \tag{6.27}$$

Proof. First note that if z(t) is a long-time controlled solution with parameters $(s, \theta, T_{\star}, \epsilon)$ and has initial datum well prepared with parameters $(s, \theta, \nu_0, \nu_1, \epsilon)$ then the translated solution $\zeta(t)$ defined in (6.1) is long-time controlled and has initial data well-prepared with the same parameters.

From now on we shall simply denote $A \equiv A_{s,R}$. Since $\zeta(t)$ fulfills (6.2), we compute

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{A}(t) = \langle \mathrm{i}[\mathsf{A}, \mathrm{Op}^{BW}((\mathsf{J}_1 + \mathfrak{v}(x))\xi)]\zeta, \zeta \rangle$$
(6.28)

$$+ \langle \mathbf{i}[\mathsf{A}, \operatorname{Op}^{BW}(\widetilde{\mathsf{V}}(t; x)\xi)]\zeta, \zeta \rangle \tag{6.29}$$

+
$$\langle i[\mathsf{A}, \operatorname{Op}^{BW}\left(-|\xi|^{\alpha} + \widetilde{\mathsf{b}}(t; x, \xi)\right)]\zeta, \zeta\rangle$$
 (6.30)

$$+2\operatorname{Re}\left\langle \mathsf{A}\widetilde{Y}(t),\zeta\right\rangle \tag{6.31}$$

We shall use that, for well-prepared data, the number I_1 in (6.9) fulfills (see (6.22))

$$I_1 \ge 2|z_1(0)| |z_{-1}(0)|\nu_0 \epsilon^2, \tag{6.32}$$

whereas for long-time controlled solutions (see (5.14)), one has

$$|z_1(0)|^2 + |z_{-1}(0)|^2 \le \epsilon^2.$$
(6.33)

We first estimate the term (6.28) from below using Lemma 6.2. Precisely we get

$$\langle \mathbf{i}[\mathbf{A}, \mathrm{Op}^{BW}((\mathbf{J}_{1} + \boldsymbol{\mathfrak{v}}(x))\xi)]\zeta, \zeta \rangle \overset{(6.14), (6.15)}{\geq} \mathbf{I}_{1} \langle \mathrm{Op}^{BW}\left(|\xi|^{2s}\eta_{\mathsf{R}}^{2}(\xi)\right)\zeta, \zeta \rangle - C_{s}\frac{\epsilon^{4}}{\mathsf{R}}\|\zeta\|_{s}^{2}$$

$$\overset{(6.32)}{\geq} 2|z_{1}(0)| |z_{-1}(0)|\nu_{0} \epsilon^{2} \langle \mathrm{Op}^{BW}\left(|\xi|^{2s}\eta_{\mathsf{R}}^{2}(\xi)\right)\zeta, \zeta \rangle - C_{s}\frac{\epsilon^{4}}{\mathsf{R}}\|\zeta\|_{s}^{2}$$

$$\overset{(6.16), (6.17)}{\geq} \nu_{0} \epsilon^{2} \mathcal{A}(t) - C_{s}\frac{\epsilon^{4}}{\mathsf{R}}\|\zeta\|_{s}^{2} .$$

$$(6.34)$$

Next we estimate (6.29) from above. We first use estimate (A.2) (with $\nu = 0, m' = 1, m = 2s$),

$$|(6.29)| \le |\mathfrak{a}|_{2s,W^{2,\infty},7} |\widetilde{\mathfrak{V}}(t,\cdot)|_{1,W^{2,\infty},7} \|\zeta\|_s^2 \stackrel{(6.7),(6.3),(6.33)}{\le} C_s \epsilon^{6-\theta} \|\zeta\|_s^2 .$$
(6.35)

Next we estimate (6.30) from above. We use again estimate (A.2) (this time with $\nu = 1 - \alpha$, $m' = \alpha$, m = 2s, thinking $\mathfrak{a}(x,\xi)$ as a symbol in $\Gamma_{W^{2,\infty}}^{2s+1-\alpha}$ supported on high frequencies) to bound

$$|(6.30)| \leq \frac{1}{\mathsf{R}^{1-\alpha}} \,|\mathfrak{a}|_{2s,W^{2,\infty},7} \,\,||\xi|^{\alpha} + \widetilde{\mathsf{b}}(t,\cdot)|_{\alpha,W^{2,\infty},7} \,\|\zeta\|_{s}^{2} \stackrel{(6.7),(6.3)}{\leq} C_{s} \frac{\epsilon^{2}}{\mathsf{R}^{1-\alpha}} \,\|\zeta\|_{s}^{2} \,\,. \tag{6.36}$$

Finally we estimate (6.31) from above. We use estimate (2.27) to bound

$$|(6.31)| \le \|\mathsf{A}\widetilde{Y}(t)\|_{-s} \|\zeta\|_{s} \le C_{s} |\mathfrak{a}|_{2s,L^{\infty},7} \|\widetilde{Y}(t)\|_{s} \|\zeta\|_{s} \stackrel{(6.7),(6.3)}{\le} C_{s} \epsilon^{5-\theta} \|\zeta\|_{s} .$$
(6.37)

Then (6.27) follows from (6.34), (6.35), (6.36) and (6.37), choosing **R** as in (6.23), and using that $\zeta(t)$, being long-time controlled, fulfills $\|\zeta(t)\|_s \leq \epsilon^{-\theta}$ and provided ϵ is sufficiently small.

We are finally able to prove Proposition 6.5.

Proof of Proposition 6.5. Let $z(t) \in H^s(\mathbb{T}, \mathbb{C})$ be a solution of system (5.3)–(5.4) whose initial datum $z(0) \in H^s(\mathbb{T}; \mathbb{C})$ is well-prepared with parameters $(s, \theta, \nu_0, \epsilon)$ and which is long-time controlled with parameters $(s, \theta, T_\star, \epsilon)$, T_\star in (6.24). By Lemma 6.6, provided $\epsilon > 0$ is sufficiently small, the functional $\mathcal{A}(t)$ in (6.26) fulfills the inequality (6.27). Integrating in time, we get

$$\mathcal{A}(t) \ge e^{\nu_0 \epsilon^2 t} \left(\mathcal{A}(0) - C \epsilon^{3-2\theta} \right) + C \epsilon^{3-3\theta} , \qquad 0 \le t \le \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right)$$

A sufficient condition for $\mathcal{A}(t)$ to grow in time is that $\mathcal{A}(0) > C\epsilon^{3-2\theta}$; this condition is fulfilled for well-prepared initial data provided ϵ is sufficiently small; indeed by (6.23)

$$\mathcal{A}(0) = \langle \mathsf{A}\zeta(0), \zeta(0) \rangle = \langle \mathsf{A}z^{\perp}(0), z^{\perp}(0) \rangle > \epsilon^{3-3\theta} > 2C\epsilon^{3-2\theta} .$$

Then for some $C_s > 1$ we get that

$$\frac{1}{2}\epsilon^{3-3\theta} e^{\nu_0 \epsilon^2 t} \leq \mathcal{A}(t) \leq C_s \epsilon^2 \|z(t)\|_s^2 .$$

Hence, when $t = \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right)$, eventually shrinking ϵ , one gets

$$||z(t)||_{s}^{2} \geq \frac{1}{2C_{s}} \epsilon^{1-3\theta} e^{\nu_{0}T_{0}\log(\epsilon^{-1})} \stackrel{(6.24)}{\geq} \frac{1}{\epsilon^{2\theta}}$$

yielding (6.25).

6.3 Conclusion and proof of Theorem 1.1

Fix s, θ as in (5.2). We give now an example of a well-prepared initial data.

Lemma 6.7. Let $\rho_1, \rho_{-1} > 0$ in the non-empty region limited by

$$\rho_1^2 + \rho_{-1}^2 \le 1 , \qquad \nu_0 := 2\rho_1\rho_{-1} - \frac{\rho_1^2 + \rho_{-1}^2}{2} > 0 .$$
(6.38)

There exists $\epsilon_0 > 0$ and, for any $\epsilon \in (0, \epsilon_0)$, an interval $I(\epsilon)$ such that the initial datum

$$z(0) := \epsilon \rho_1 e^{\mathbf{i}x} + \epsilon \rho_{-1} e^{-\mathbf{i}x} + \rho e^{\mathbf{i}3\mathbb{N}x} + \mathbf{i}\rho e^{\mathbf{i}(3\mathbb{N}+2)x} , \quad \mathbb{N} := \lceil \mathbb{R} \rceil$$
(6.39)

with $\mathbf{R} = \epsilon^{-(3+\theta)/(1-\alpha)}$ and $\rho \in I(\epsilon)$, fulfills:

- well-prepared: z(0) in (6.39) is a well-prepared initial datum with parameters $(s, \theta, \nu_0, \epsilon)$ (according to Definition 6.3);
- L^2 -smallness: the bounds in (5.14) holds true;

• H^s -smallness: z(0) fulfills the high norm bound

$$\|z(0)\|_s \le \epsilon^\theta . \tag{6.40}$$

Proof. We first prove that each of the three claimed properties gives a restriction on the choice of ρ . Then we prove that such conditions are compatible.

Well-prepared: Condition (B1) follows immediately from (6.38). We now check condition (B2). Using the definition of paradifferential operator in (2.21), the form of A in (6.5) and of z(0) in (6.39), we get

$$\begin{split} \langle \mathsf{A}\Pi^{\perp} z(0), \Pi^{\perp} z(0) \rangle &= \sum_{k} \epsilon^{2} \rho_{1} \rho_{-1} \, \left| k + 1 \right|^{2s} \, \eta_{\mathsf{R}}^{2} (k+1) \, \chi_{2} (1, -1, k+1) \, \mathrm{Im} \left(\overline{z}_{k}^{\perp}(0) \, z_{k+2}^{\perp}(0) \right) \\ &= \epsilon^{2} \rho_{1} \rho_{-1} \, \left| \mathsf{3N} + 1 \right|^{2s} \, \underbrace{\eta_{\mathsf{R}}^{2} (\mathsf{3N} + 1)}_{=1} \underbrace{\chi_{2} (1, -1, \mathsf{3N} + 1)}_{=1} \, \rho^{2} = \epsilon^{2} \rho_{1} \rho_{-1} \, \left| \mathsf{3N} + 1 \right|^{2s} \, \rho^{2} \, . \end{split}$$

Then (6.23) is fulfilled provided $\rho_1 \rho_{-1} 3^{2s} \mathbb{R}^{2s} \rho^2 \ge \epsilon^{1-3\theta}$, which using (6.23) gives

$$\rho \ge \frac{\epsilon^{\frac{1}{2} - \frac{3}{2}\theta + s\frac{3+\theta}{1-\alpha}}}{3^s \sqrt{\rho_1 \rho_{-1}}}.$$
(6.41)

This proves that z(0) is well prepared.

<u> L^2 -smallness</u>: The first condition in (5.14) is satisfied thanks to the first assumption in (6.38) and the second condition in (5.14) is satisfied provided that

$$\rho \le \frac{\epsilon^3}{\sqrt{2}}.\tag{6.42}$$

<u> H^s -smallness</u>: The condition (6.40) is satisfied provided that

$$(\rho_1^2 + \rho_{-1}^2)\epsilon^2 \le \frac{\epsilon^{2\theta}}{2}$$
 and $\rho^2(3N+1)^{2s} + \rho^2(3N+3)^{2s} \le \frac{\epsilon^{2\theta}}{2}.$

The first condition follows automatically from (6.38) and taking ϵ sufficiently small, while the second one, using $N \leq R + 1$ and (6.23), is fulfilled for example for

$$\rho \le \frac{\epsilon^{\theta + s \frac{3+\theta}{1-\alpha}}}{6^{s}2}.\tag{6.43}$$

Note also that, since $s \ge 3\mathfrak{s}_0 \ge 1$, for ϵ small enough the second condition (6.42) is less restrictive than the third one (6.43). Note that, provided ϵ is small enough and using $\theta < \frac{1}{5}$, conditions (6.41) and (6.43) are compatible. Then, taking

$$\rho \in I(\epsilon) := \left(\frac{1}{3^s \sqrt{\rho_1 \rho_{-1}}} \epsilon^{\frac{1}{2} - \frac{3}{2}\theta + s\frac{3+\theta}{1-\alpha}}, \frac{\epsilon^{\theta + s\frac{3+\theta}{1-\alpha}}}{6^s 2}\right),$$

the datum z(0) satisfies all the claimed conditions.

We now show that any solution of system (4.6) with a well prepared initial datum as in Lemma 6.7 undergoes Sobolev norm explosion. Precisely we have:

Lemma 6.8. Fix s, θ as in (5.2). There exists $\epsilon_2 > 0$ such that, provided $\epsilon \in (0, \epsilon_2)$ the following holds true. Let $z(0) \in H^s(\mathbb{T}, \mathbb{C})$ as in Lemma 6.7 and so well-prepared with parameters $(s, \theta, \nu_0, \epsilon)$, for some $\nu_0 \in (0, \frac{1}{2})$. Consider the solution z(t) of system (5.3)–(5.4) with initial datum z(0). Denote by

$$0 < T_1 := T_1(\epsilon; z(0)) := \inf \left\{ t \ge 0 : \| z(t) \|_s \ge \epsilon^{-\theta} \right\} .$$
(6.44)

Then T_1 is finite and bounded by $T_1 \leq \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right)$, $T_0 = \nu_0^{-1}$. Moreover one has

$$\sup_{0 \le t \le T_1} \|z(t)\|_{\mathfrak{s}_0} \le 3\epsilon \ , \quad \|z(0)\|_s \le \epsilon^{\theta} \ , \quad \|z(T_1)\|_s \ge \epsilon^{-\theta} \ . \tag{6.45}$$

Proof. Define $\epsilon_2 := \min(\epsilon_{\star}, \epsilon_0, \epsilon_1, \mathfrak{r})$ with ϵ_{\star} of Lemma 5.3, ϵ_0 of Lemma 6.7 and ϵ_1 of Proposition 6.5. First note that the solution z(t) is long-time controlled with parameters $(s, \theta, T_1, \epsilon)$ (see Definition 5.2); indeed condition (A1) holds true in view of the L^2 -smallness of Lemma 6.7, whereas condition (A2) holds true with $T_{\star} \rightsquigarrow T_1$ by the minimality of T_1 .

We now show that T_1 is finite and bounded by $\frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right)$. Assume by contradiction that $T_1 > T_0 \epsilon^{-2} \log(\epsilon^{-1})$. Then, by the very definition of T_1 ,

$$\sup_{0 \le t \le T_0 \epsilon^{-2} \log(\epsilon^{-1})} \|z(t)\|_s \le \epsilon^{-\theta} ,$$

namely the solution z(t) is long-time controlled also with parameters $(s, \theta, \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right), \epsilon)$. Then, since by Lemma 6.7 the initial data z(0) is well prepared, Proposition 6.5 applies and therefore

$$\sup_{0 \le t \le T_0 \epsilon^{-2} \log(\epsilon^{-1})} \|z(t)\|_s \ge \epsilon^{-\theta}$$

contradicting the minimality of T_1 . This proves that $T_1 \leq \frac{T_0}{\epsilon^2} \log\left(\frac{1}{\epsilon}\right)$.

To control the low norm $||z(t)||_{\mathfrak{s}_0}$, we apply the bootstrap lemma 5.3 with the parameter $T_{\star} = T_1$ that we have just proved satisfy the required condition (5.16). The last two inequalities of (6.45) follow by (6.40) and (6.44).

We conclude with:

Proof of Theorem 1.1. Recall that the variables u(t) and z(t) are related by the admissible transformation $Z(t) = \mathcal{F}(U(t)) \equiv \mathbf{F}(U(t))U(t)$ in (4.36). By Lemma 2.13, the map $Z = \mathcal{F}(U)$ is locally invertible provided $||Z||_{\mathfrak{s}_0} \leq r'$ is sufficiently small, and has the form $\mathcal{F}^{-1}(Z) = \mathbf{G}(Z)Z$ for some $\mathbf{G}(Z)$ fulfilling the bound in (2.44).

So consider $Z(0) = \left(\frac{z(0)}{\overline{z}(0)}\right)$ with z(0) as in Lemma 6.7 and therefore fulfilling $||Z(0)||_{\mathfrak{s}_0} \leq \epsilon^{\theta} \leq \mathfrak{r}$. We define

$$U(0) := \mathcal{F}^{-1}(Z(0)) = \mathbf{G}(Z(0))Z(0)$$
.

We take U(0) as the initial data for equation (1.1); by (2.44), its Sobolev norm

$$||U(0)||_s \le C_s ||Z(0)||_s \stackrel{(6.45)}{\le} C_s \epsilon^{\theta}$$
.

Consider now the solution U(t) of (1.1) with initial data U(0). By Theorem 4.3, $Z(t) = \mathcal{F}(U(t))$ is the solution of equation (4.6) with initial datum Z(0) of Lemma 6.7; consequently, in view of Lemma 5.1 and Lemma 6.8, z(t) has a small $H^{\mathfrak{s}_0}$ -norm for all times $0 \leq t \leq T_1$, but large H^s -norm at time T_1 . We deduce that $U(t) = \mathcal{F}^{-1}(Z(t))$ fulfills the bound

$$||U(t)||_{\mathfrak{s}_0} \le C_{\mathfrak{s}_0} ||Z(t)||_{\mathfrak{s}_0} \le C_{\mathfrak{s}_0} \epsilon < \mathfrak{r}, \quad \forall 0 \le t \le T_1 .$$

At time T_1 , we bound from below the H^s -norm of $U(T_1)$ using the identity $Z(T_1) = \mathcal{F}(U(T_1))$, the fact that $||U(T_1)||_{\mathfrak{s}_0} \leq \mathfrak{r}$ and estimate (2.44), to get

$$||U(T_1)||_s \ge C_s^{-1} ||Z(T_1)||_s \stackrel{(6.45)}{\ge} C_s^{-1} \epsilon^{-\theta}$$
.

Given arbitrary $\delta \in (0,1)$ and $K \ge 1$, shrink ϵ to conclude the proof of Theorem 1.1.

A High frequency paradifferential calculus

In this section we consider paradifferential operators with symbols supported only on high frequencies and prove a commutator estimate and a Garding inequality keeping track of the size of the support of the symbols. **Lemma A.1.** Let $N \in \mathbb{N}_0$, $m \in \mathbb{R}$ and $\mathbb{R} \geq 1$. If $a \in \Gamma^m_{W^{N,\infty}}$, then

$$a_{\mathbf{R}}(x,\xi) := a(x,\xi) \eta_{\mathbf{R}}(\xi), \quad \eta_{\mathbf{R}} \text{ in } (6.6)$$

is a symbol in $\Gamma_{W^{N,\infty}}^{m+\nu}$ for any $\nu \geq 0$ with quantitative bound

$$|a_{\mathbf{R}}|_{m+\nu,W^{N,\infty},n} \le C_n \, \mathbf{R}^{-\nu} \, |a|_{m,W^{N,\infty},n} \quad \text{for any } n \in \mathbb{N}_0.$$
(A.1)

In addition, if $N \geq 2$ and $b \in \Gamma_{W^{2,\infty}}^{m'}$, $m' \in \mathbb{R}$, one has the commutator estimate

$$\|[\operatorname{Op}^{BW}(a_{\mathbb{R}}), \operatorname{Op}^{BW}(b)]u\|_{s-m-m'-\nu+1} \le C\mathbb{R}^{-\nu} |a|_{m,W^{2,\infty},7} |b|_{m',W^{2,\infty},7} ||u||_{s} .$$
(A.2)

Proof. For any $\alpha, \beta \in \mathbb{N}_0$, $\alpha \leq N$, $\beta \leq n$, we have

$$\begin{split} \left| (\partial_x^{\alpha} \partial_{\xi}^{\beta} a_{\mathbf{R}}(x,\xi) \right| \lesssim \sum_{\beta_1 + \beta_2 = \beta} \left| \partial_x^{\alpha} \partial_{\xi}^{\beta_1} a(x,\xi) \right| \left| \partial_{\xi}^{\beta_2} \eta_{\mathbf{R}}(\xi) \right| \\ \lesssim \sum_{\beta_1 + \beta_2 = \beta} \left| a \right|_{m,W^{N,\infty},n} \langle \xi \rangle^{m-\beta_1} \frac{1}{\mathbf{R}^{\beta_2}} \left| \eta^{(\beta_2)} \left(\frac{\xi}{\mathbf{R}}\right) \right| \\ \lesssim \left| a \right|_{m,W^{N,\infty},n} \sum_{\beta_1 + \beta_2 = \beta} \langle \xi \rangle^{m-\beta_1 - \beta_2 + \nu} \sup_{\xi} \left| \langle \xi \rangle^{-\nu} \langle \frac{\xi}{\mathbf{R}} \rangle^{\beta_2} \eta^{(\beta_2)} \left(\frac{\xi}{\mathbf{R}}\right) \right| \\ \lesssim \frac{1}{\mathbf{R}^{\nu}} \left| a \right|_{m,W^{N,\infty},n} \langle \xi \rangle^{m-\beta+\nu} , \end{split}$$

where in the last step we used that the function $\langle \frac{\xi}{R} \rangle^{\beta_2} \eta^{(\beta_2)}(\frac{\xi}{R})$ is uniformly bounded on \mathbb{R} and has support on $\xi \geq R$.

We prove now (A.2). By Theorem 2.7 with $\rho = 2$ we have

$$[\operatorname{Op}^{BW}(a_{\mathbb{R}}), \operatorname{Op}^{BW}(b)] = \operatorname{Op}^{BW}(\{a_{\mathbb{R}}, b\}) + R^{-2}(a_{\mathbb{R}}, b).$$

We now bound both terms in the above equation regarding $a_{\mathbb{R}}$ as a symbol in $\Gamma_{W^{N,\infty}}^{m+\nu}$ and $\{a_{\mathbb{R}}, b\}$ as a symbol in $\Gamma_{W^{N-1,\infty}}^{m+m'+\nu-1}$. By (2.27) and (2.39), we get

$$\begin{split} \| \operatorname{Op}^{BW}(\{a_{\mathbb{R}}, b\}) \, u \|_{s-m-m'-\nu+1} &\lesssim |\{a_{\mathbb{R}}, b\}|_{m+m'+\nu-1, L^{\infty}, 4} \| u \|_{s} \\ &\lesssim |a_{\mathbb{R}}|_{m+\nu, W^{1,\infty}, 5} |b|_{m', W^{1,\infty}, 5} \| u \|_{s} \\ &\stackrel{(A.1)}{\lesssim} \mathbb{R}^{-\nu} \, |a|_{m, W^{1,\infty}, 5} \, |b|_{m', W^{1,\infty}, 5} \| u \|_{s} \,. \end{split}$$
(A.3)

Next we estimate the norm of $R^{-2}(a_{\mathbb{R}}, b)$ using (2.40):

$$\|R^{-2}(a_{\mathbb{R}},b)u\|_{s-m-m'-\nu+2} \lesssim |a_{\mathbb{R}}|_{m+\nu,W^{2,\infty},7} \|b\|_{m',W^{2,\infty},7} \|u\|_{s}$$

$$\stackrel{(A.1)}{\lesssim} \mathbb{R}^{-\nu} |a|_{m,W^{2,\infty},7} |b|_{m',W^{2,\infty},7} \|u\|_{s}$$
(A.4)

In conclusion (A.2) follows from (A.3), (A.4).

In the following we shall use a well-known cancellation which is a direct consequence of Proposition 2.7: if $a \in \Gamma_{W^{2,\infty}}^{m}$, $b \in \Gamma_{W^{2,\infty}}^{m'}$, with $m, m' \in \mathbb{R}$, then

$$\operatorname{Op}^{BW}(b) \circ \operatorname{Op}^{BW}(a) \circ \operatorname{Op}^{BW}(b) = \operatorname{Op}^{BW}(ab^2) + R^{-2}(a,b),$$
(A.5)

where $R^{-2}(a,b)$ is a bounded operator $H^s \to H^{s-(m+2m')+2}$, $\forall s \in \mathbb{R}$, satisfying, for any $u \in H^s$,

$$\|R^{-2}(a,b)\|_{s-(m+m')+2} \lesssim |a|_{m,W^{2,\infty},8} \|b\|_{m',W^{2,\infty},8}^2 \|u\|_s.$$
(A.6)

In the next lemma we prove a simplified version of the strong Garding inequality adapted to our setting.

Lemma A.2 (Strong Garding's inequality). Let $\mathbb{R} \geq 1$, $a(x) \in W^{3,\infty}$ and $a(x) \geq 0$. Let $\psi(\xi) \in \widetilde{\Gamma}_0^m$, m > 0, a real valued Fourier multiplier with supp $\psi \subseteq [\mathbb{R}, +\infty)$. Then there is C > 0 such that

$$\langle \operatorname{Op}^{BW}(a(x)\psi^{2}(\xi)) u, u \rangle \geq -C \frac{\|a\|_{W^{3,\infty}}}{\mathbb{R}^{2}} \|u\|_{m}^{2} .$$
 (A.7)

Proof. Arguing as in Lemma A.1 one shows that, for any $n \in \mathbb{N}_0$,

$$|\psi|_{m+1,L^{\infty},n} \le C_n \frac{1}{\mathbf{R}} |\psi|_{m,L^{\infty},n} .$$
(A.8)

We apply now the composition formula (A.5) regarding $\psi(\xi)$ as a symbol in $\widetilde{\Gamma}_0^{m+1}$:

$$\operatorname{Op}^{BW}(\psi) \circ \operatorname{Op}^{BW}(a) \circ \operatorname{Op}^{BW}(\psi) = \operatorname{Op}^{BW}(a\psi^2) + \mathsf{R}_1$$
(A.9)

with $\mathsf{R}_1 \colon H^m \to H^{-m}$ fulfilling, by (A.6),

$$\|\mathsf{R}_{1}u\|_{-m} \lesssim \|a\|_{W^{2,\infty}} \|\psi\|_{m+1,L^{\infty},8}^{2} \|u\|_{m} \lesssim \frac{(A.8)}{\mathsf{R}^{2}} \frac{1}{\mathsf{R}^{2}} \|a\|_{W^{2,\infty}} \|u\|_{m} .$$
(A.10)

Then observe that $\operatorname{Op}^{BW}(\psi) = \operatorname{Op}^{W}(\psi) = \psi(D)$ and $\operatorname{Op}^{BW}(a) = \operatorname{Op}^{W}(a) + \operatorname{Op}^{W}(a_{\chi} - a)$, where a_{χ} is the cut-offed symbol defined in (2.20), so

$$\operatorname{Op}^{BW}(\psi) \circ \operatorname{Op}^{BW}(a) \circ \operatorname{Op}^{BW}(\psi) = \psi(D) \circ \operatorname{Op}^{W}(a) \circ \psi(D) + \mathsf{R}_{2}$$
(A.11)

where $\mathsf{R}_2 := \psi(D) \circ \operatorname{Op}^W(a_{\chi} - a) \circ \psi(D)$. Now we prove that R_2 is bounded $H^m \to H^{-m}$. First note that, by the definitions (2.20) and (2.22), for any $v \in H^{-1}$,

$$\begin{split} \|\mathrm{Op}^{W}(a_{\chi}-a)\,v\|_{1}^{2} &\lesssim \sum_{j} \langle j \rangle^{2} \left| \sum_{k} \widehat{a}_{j-k} \left(1-\chi(k-j,\frac{j+k}{2}) \right) v_{k} \right|^{2} \\ &\lesssim \sum_{j} \left| \sum_{k} \langle j-k \rangle^{2} \left| \widehat{a}_{j-k} \right| \left(1-\chi(k-j,\frac{j+k}{2}) \right) \frac{1}{\langle k \rangle} |v_{k}| \right|^{2} \\ &\lesssim \sum_{j} \left| \sum_{k} \langle j-k \rangle^{2} \left| \widehat{a}_{j-k} \right| \frac{1}{\langle k \rangle} |v_{k}| \right|^{2} \\ &\lesssim \|a\|_{3}^{2} \|v\|_{-1}^{2} \lesssim \|a\|_{W^{3,\infty}}^{2} \|v\|_{-1}^{2} \end{split}$$

where to pass from the first to the second line we used that, on the support of $1 - \chi(k - j, \frac{j+k}{2})$, one has

$$\langle k \rangle, \langle j \rangle \lesssim \langle j - k \rangle + \langle j + k \rangle \lesssim \langle j - k \rangle$$
,

and to pass from the third to the last line we used Young's inequality for convolution of sequences. Thus we get, for any $u \in H^m$,

$$\begin{aligned} \|\mathsf{R}_{2}u\|_{-m} &\lesssim |\psi|_{m+1,L^{\infty},0} \, \|\mathsf{Op}^{W}(a_{\chi}-a) \circ \psi(D)u\|_{1} \\ &\lesssim |\psi|_{m+1,L^{\infty},0} \, \|a\|_{W^{3,\infty}} \, \|\psi(D)u\|_{-1} \\ &\lesssim |\psi|_{m+1,L^{\infty},0}^{2} \, \|a\|_{W^{3,\infty}} \, \|u\|_{m} \, \overset{(\mathrm{A}.8)}{\lesssim} \, \frac{1}{\mathsf{R}^{2}} \|a\|_{W^{3,\infty}} \|u\|_{m} \, . \end{aligned}$$
(A.12)

In conclusion, combining (A.9) and (A.11) and since $Op^W(a) = a \ge 0$ and $\psi(D)$ is self-adjoint, we have that

$$0 \le \langle \psi(D) \circ a \circ \psi(D)u, u \rangle = \langle \operatorname{Op}^{BW}(a\psi^2)u, u \rangle + \langle (\mathsf{R}_1 - \mathsf{R}_2)u, u \rangle$$

and (A.7) follows by (A.10) and (A.12).

B Flows and conjugations

In this section we collect some results about the conjugation of paradifferential operators and smoothing remainders under flows, following [9, 11, 13, 60].

Conjugation by a flow generated by a real symbol of order one. Given a function $\beta \in \widetilde{\mathcal{F}}_2^{\mathbb{R}}$ gauge invariant, i.e. $\beta(\mathbf{g}_{\theta}U; \cdot) = \beta(U; \cdot)$ for any $\theta \in \mathbb{T}$, consider the flow $\Phi^{\tau}(u), \tau \in [-1, 1]$ defined by (4.23). It is standard (see e.g. Lemma 3.22 in [9]) that, for any $U \in B_{s_0,\mathbb{R}}(r)$ with $s_0 > 0$ sufficiently large and r > 0 sufficiently small, the operator $\Phi^{\tau}(U) \in \mathcal{L}(H^s(\mathbb{T}, \mathbb{C}^2))$ for any $s \in \mathbb{R}$ with the quantitative estimate: there is a constant C(s) > 0 such that for any $W \in H^s(\mathbb{T}, \mathbb{C}^2)$, $\|\Phi^{\tau}(U)W\|_s + \|\Phi^{\tau}(U)^{-1}W\|_s \leq C(s)\|W\|_s$. Following [9], we define the path of diffeomorphism of \mathbb{T} via

 $\Psi(U,\tau;x) := x + \tau\beta(U;x) \quad \text{with inverse} \quad \Psi^{-1}(U,\tau;y) := y + \check{\beta}(U,\tau;y), \quad \ \check{\beta} \in \mathcal{F}^{\mathbb{R}}_{\geq 2}[r]$

and set $\Psi(U; x) := \Psi(U, 1; x)$.

Proposition B.1 (Conjugations for a transport flow). Let $m \in \mathbb{R}$, $\rho > 0$, and let $\Phi(U)$ be the flow generated by (4.23).

1. Space conjugation of a para-differential operator: Let $a \in \Sigma\Gamma_2^m[r]$ be a real symbol and $a^{(m)}(U; x, \xi) := a(U; y, \xi \ \partial_y \Psi^{-1}(U; y))|_{y=\Psi(U; x)} \in \Sigma\Gamma_2^m[r]$. Then

$$\begin{split} \Phi(U) \circ \operatorname{Op}_{\mathsf{vec}}^{BW}(a(U;x,\xi)) \circ \Phi(U)^{-1} &= \operatorname{Op}_{\mathsf{vec}}^{BW}\left(a^{(m)}(U;x,\xi) + a^{(m-2)}_{\geq 4}(U;x,\xi)\right) + R_{\geq 4}(U) \\ &= \operatorname{Op}_{\mathsf{vec}}^{BW}\left(a(U;x,\xi) + a^{(m)}_{\geq 4}(U;x,\xi)\right) + R_{\geq 4}(U), \end{split}$$
(B.1)

where $a_{\geq 4}^{(m-2)}(U; x, \xi)$ and $a_{\geq 2}^{(m)}(U; x, \xi)$ are non-homogeneous real symbols in $\Gamma_{\geq 4}^{m-2}[r]$ respectively $\Gamma_{\geq 4}^m[r]$, whereas $R_{\geq 4}(U)$ is a real-to-real matrix of smoothing operators in $\mathcal{R}_{\geq 4}^{-\varrho+m}[r]$. In addition if $a(U; x, \xi) = V(U; x)\xi$ for some $V \in \widetilde{\mathcal{F}}_2^{\mathbb{R}}[r]$ then in (B.1) $a_{\geq 4}^{(m-2)} \equiv 0$ and $a_{\geq 4}^{(m)}(U; x, \xi) = V_{\geq 4}'(U; x)\xi$ for a suitable function $V_{\geq 4}' \in \mathcal{F}_{\geq 4}^{\mathbb{R}}[r]$.

2. Space conjugation of a Fourier multiplier Let $\omega(\xi) \in \widetilde{\Gamma}_0^{\alpha}$ be a real Fourier multiplier. Then

$$\Phi(U) \circ \operatorname{Op}_{\mathsf{vec}}^{BW}(\mathrm{i}\omega) \circ \Phi(U)^{-1} = \operatorname{Op}_{\mathsf{vec}}^{BW} \Big(\mathrm{i} \big(\omega + a_2^{(\alpha)}(U; x, \xi) + a_{\geq 4}^{(\alpha)}(U; x, \xi) + a_{\geq 4}^{(\alpha-2)}(U; x, \xi) \big) \Big) + R_2(U) + R_{\geq 4}(U),$$
(B.2)

where

- $a_2^{(\alpha)}(U; x, \xi)$ is a real, zero-average, gauge invariant symbol in $\widetilde{\Gamma}_2^{\alpha}$;
- $a_{\geq 4}^{(\alpha)}(U; x, \xi)$ is a real non-homogeneous symbol in $\Gamma_{\geq 4}^{\alpha}[r]$ and $a_{\geq 4}^{(\alpha-2)}(U; x, \xi)$ is a non-homogeneous symbol in $\Gamma_{>4}^{\alpha-2}[r]$;

• $R_2(U)$ is a real-to-real, gauge invariant matrix of smoothing operators in $\widetilde{\mathcal{R}}_2^{-\varrho+m}$, and $R_{\geq 4}(U)$ is a real-to-real matrix of non-homogeneous smoothing operators in $\mathcal{R}_{\geq 4}^{-\varrho+m}$.

3. Space conjugation of a smoothing remainder: If $R_2(U)$ is a real-to-real matrix of smoothing operators in $\tilde{\mathcal{R}}_2^{-\varrho}[r]$ then

$$\Phi(U) \circ R_2(U) \circ \Phi(U)^{-1} = R_2(U) + R_{\ge 4}(U),$$

where $R_{\geq 4}(U)$ is a real-to-real matrix of smoothing operators in $\mathcal{R}_{\geq 4}^{-\varrho+1}[r]$.

4. Conjugation of ∂_t : If U is a solution of (4.1) then

$$(\partial_t \Phi(U)) \Phi(U)^{-1} = \mathrm{i} \operatorname{Op}_{\mathsf{vec}}^{BW}(2\beta(-\mathrm{i}\Omega(D)U, U; x) \xi + \mathrm{i} \mathsf{V}_{\geq 4}(U; x)\xi) + R_{\geq 4}(U),$$

where $\Omega(D)$ is the matrix of real Fourier multipliers in (3.17), $V_{\geq 4}(U; x)$ is a real function in $\mathcal{F}_{\geq 4}^{\mathbb{R}}[r]$ and $R_{\geq 4}(U)$ is a real-to-real matrix of smoothing operators in $\mathcal{R}_{\geq 4}^{-\varrho}[r]$.

Proof. During the proof we shall denote $b := \frac{\beta}{1 + \tau \beta_x}$.

1. Follows by Lemmas A4 and A5 in [11].

2. We first define the operator $P^{\tau}(U) := \Phi^{\tau}(U) \circ \operatorname{Op}_{\mathsf{vec}}^{BW}(\mathrm{i}\omega) \circ (\Phi^{\tau}(U))^{-1}$. Note that $P^{\tau}(U)$ is gauge invariant being composition of gauge invariant operators. By Theorem 3.27 in [9] (actually adapting that result when the function β is 2-homogeneous rather than 1-homogeneous), we have for any $\tau \in [0, 1]$

$$P^{\tau}(U) = \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\mathrm{i}\omega_{\Phi}^{(m)} + \mathrm{i}\omega^{(m-2)} \right) + R(U,\tau)$$

=
$$\operatorname{Op}_{\mathsf{vec}}^{BW} \left(\mathrm{i}\omega + \mathrm{i}\omega_{2}^{(m)} + \mathrm{i}a_{\geq 4}^{(m)} + \mathrm{i}\omega_{2}^{(m-2)} + \mathrm{i}a_{\geq 4}^{(m-2)} \right) + R_{2}(U,\tau) + R_{\geq 4}(U,\tau)$$
(B.3)

where $\omega_{\Phi}^{(m)} = \omega + \omega_2^{(m)} + a_{\geq 4}^{(m)}$ is a real symbol in $\Sigma \Gamma_0^m[r]$, $\omega^{(m-2)} = \omega_2^{(m-2)} + a_{\geq 4}^{(m-2)}$ is a symbol in $\Sigma \Gamma_2^{m-2}[r]$ and $R = R_2 + R_{\geq 4} \in \Sigma \mathcal{R}_2^{-\varrho+m}[r]$.

To identify the quadratic component of $P^1(U)$ we use the Taylor expansion $P^1(U) = P^0(U) + \partial_\tau P^\tau(U)|_{\tau=0} + \int_0^1 (1-\tau) \partial_\tau^2 P^\tau(U) d\tau$ and exploit that $P^\tau(U)$ fulfills the Heisenberg equation $\partial_\tau P^\tau(U) = [G(U,\tau), P^\tau(U)], \quad P^0(U) = \operatorname{Op}_{\mathsf{vec}}^{BW}(i\omega)$. Using that $G(U,0) = \operatorname{Op}_{\mathsf{vec}}^{BW}(ib(U)\xi)$ and the paradifferential structure of $P^\tau(U)$ in (B.3), we obtain

$$P^{1}(U) = \operatorname{Op}_{\mathsf{vec}}^{BW}(\mathrm{i}\omega) + [\operatorname{Op}_{\mathsf{vec}}^{BW}(\mathrm{i}b(U)\xi), \operatorname{Op}_{\mathsf{vec}}^{BW}(\mathrm{i}\omega)] + M_{\geq 4}(U)$$

with $M_{\geq 4}(U)$ a *m*-operator in $\mathcal{M}^m_{\geq 4}[r]$. Now we use the composition Theorem 2.7 (with $\rho \rightsquigarrow \rho + 1$) and formula (2.37) to expand the commutator as

$$P^{1}(U) = \operatorname{Op}_{\mathsf{vec}}^{BW} \left(\mathrm{i}\omega + \mathrm{i}a_{2}^{(m)} \right) + R_{2}(U) + M_{\geq 4}(U)$$
(B.4)

with $a_2^{(m)}(U; x, \xi)$ the real, zero-average symbol

$$a_{2}^{(m)}(U;x,\xi) := \sum_{k=1}^{\varrho+1} \frac{(-1)^{k} - 1}{2^{k} k!} (D_{x}^{k} \beta) \left(\partial_{\xi}^{k} \omega\right) \mathrm{i}\xi \in \widetilde{\Gamma}_{2}^{m} , \qquad (B.5)$$

and $R_2(U) \in \widetilde{\mathcal{R}}_2^{-\varrho+m}$. Identifying the quadratic components of $P^1(U)$ in (B.3) and (B.4) we get that

$$\operatorname{Op}^{BW}\left(\omega_{2}^{(m)}+\omega_{2}^{(m-2)}\right) = \operatorname{Op}^{BW}\left(a_{2}^{(m)}\right) + \tilde{R}_{2}(U)$$

and therefore we get the thesis. Since $\beta(U)$ is gauge invariant (fulfills the first of (2.25)), so is $a_2^{(m)}$ in (B.5). Finally, since $P^1(U)$ is gauge invariant, also $R_2(U)$ in (B.4) is gauge invariant by difference. 3. It follows as in [9], Remark at pag 89 (see also [60, Proposition A.2] for details).

4. Differentiating (4.23) with respect to time, we get

$$(\partial_t \Phi^1(U)) \Phi^1(U)^{-1} = \int_0^1 \Phi^1(U) \, [\Phi^\tau(U)]^{-1} \operatorname{Op}_{\mathsf{vec}}^{^{BW}}(\partial_t b(U,\tau;x)\mathrm{i}\xi) \, \Phi^\tau(U) [\Phi^1(U)]^{-1} \, \mathrm{d}\tau.$$

We claim that that

$$\partial_t b(U,\tau;x) = \beta(-\mathrm{i}\mathbf{\Omega}(D)U;x) + \mathsf{V}_{\geq 4}(U,\tau;x), \quad \mathsf{V}_{\geq 4} \in \mathcal{F}_{\geq 4}^{\mathbb{R}}[r] .$$
(B.6)

Differentiating $b(U(t), \tau; x)$ with respect to t and using that, by equation (4.1), $\partial_t \beta(U) = 2\beta(\partial_t U, U) = 2\beta(X(U), U)$ with $X(U) = -i\Omega(D)U + X_3(U)$ we get

$$\begin{split} \partial_t b(U,\tau;x) &= 2\beta(-\mathrm{i}\mathbf{\Omega}(D)U,U;x) \\ &+ \underbrace{2\beta(M_{\mathrm{NLS}}(U)U,U;x) - 2\tau \left[\frac{\beta(U;x)\beta_x(X(U),U;x)}{(1+\tau\beta_x(U;x))^2} + \frac{\beta_x(U;x)\beta(X(U),U;x)}{(1+\tau\beta_x(U;x))}\right]}_{=:\mathsf{V}_{\geq 4}(U,\tau;x)} \,. \end{split}$$

Then (B.6) follows using Lemma 2.9–1 for each internal composition, getting that $V_{\geq 4}(U, \tau; x)$ is a function in $\mathcal{F}_{\geq 4}^{\mathbb{R}}[r]$.

50

Conjugation by flows generated by linear smoothing operators. In this section we study the conjugation rules for a flow $\Upsilon(U) := \Upsilon^{\tau}(U)|_{\tau=1}$ generated by

$$\partial_{\tau} \Upsilon(U) = Q(U) \circ \Upsilon^{\tau}(U), \quad \Phi^{0}_{Q}(U) = \mathrm{Id} \,, \tag{B.7}$$

with Q(U) a matrix of smoothing operators in $\widetilde{\mathcal{R}}_2^{\varrho}$. It is standard (see e.g. [60, Lemma A.4]) for any $U \in H^{s_0}(\mathbb{T}, \mathbb{C}^2)$ with $s_0 > 0$ sufficiently large, the problem (B.7) admits a unique solution $\Upsilon^{\tau}(U)$ fulfilling: for any $s \geq s_0$ there is r > 0 such that for any $U \in B_{s_0,\mathbb{R}}(r)$, and $V \in H^s(\mathbb{T}; \mathbb{C}^2)$

$$\|\Upsilon^{\tau}(U)V\|_{s} + \|\Upsilon^{-\tau}(U)V\|_{s} \lesssim_{s} \|V\|_{s} + \|V\|_{s_{0}}\|U\|_{s}\|U\|_{s_{0}},$$

uniformly in $\tau \in [-1,1]$. We denote the inverse of $\Phi_Q(u)$ as $\Upsilon(U)^{-1} = \Upsilon^{\tau}(U)|_{\tau=-1}$.

The following result is a small variation of [60, Proposition A.5] and we omit the proof.

Proposition B.2 (Conjugation by flows generated by smoothing operators). Let $m \in \mathbb{R}$, $\varrho, \varrho', r > 0$. Let Q(U) be a matrix of smoothing operators in $\tilde{\mathcal{R}}_2^{-\varrho}$ and $\Upsilon(U)$ be the flow generated by Q(U) as in (B.7). Then the following holds:

i) Space conjugation: If $a \in \Sigma \Gamma_2^m[r]$, then

$$\Upsilon(U) \circ \operatorname{Op}_{\mathsf{vec}}^{BW}(a(U; x, \xi)) \circ \Upsilon(U)^{-1} - \operatorname{Op}_{\mathsf{vec}}^{BW}(a(U; x, \xi)) \in \mathcal{R}_{\geq 4}^{-\varrho + \max\{m, 0\}}[r],$$

$$\Upsilon(U) \circ (-\mathrm{i}\Omega(D)) \circ \Upsilon(U)^{-1} - (-\mathrm{i}\Omega(D) + [Q(U), -\mathrm{i}\Omega(D)]) \in \mathcal{R}_{>4}^{-\varrho + \alpha}[r].$$

These matrices of operators are real-to-real provided Q(U) is.

ii) Conjugation of smoothing operators: If R(U) is a real-to-real matrix of smoothing operators in $\Sigma \mathcal{R}_2^{-\varrho'}[r]$, then

$$\Upsilon(U) \circ R(U) \circ \Upsilon(U)^{-1} - R(U) \in \mathcal{R}_{>4}^{-\min\{\varrho, \varrho'\}}[r]$$

and it is real-to-real.

iii) Conjugation of ∂_t : If U is a solution of (4.1) then

$$(\partial_t \Upsilon(U)) \circ \Upsilon(U)^{-1} - 2Q(-\mathrm{i}\Omega(D)U, U) \in \mathcal{R}_{\geq 4}^{-\varrho+1}[r]$$

and it is real-to-real.

References

- Bambusi D., Grébert B., Maspero A. Robert D. Reducibility of the quantum harmonic oscillator in d-dimensions with polynomial time-dependent perturbation. Anal. PDE, 11(3):775–799, 2018.
- [2] D. Bambusi, B. Grébert, A. Maspero, and D. Robert. Growth of Sobolev norms for abstract linear Schrödinger equations. J. Eur. Math. Soc. (JEMS), 2020. doi: 10.4171/JEMS/1017
- [3] Bambusi D., Grébert B., Maspero A. Robert D., Villegas-Blas C. Longtime dynamics for the Landau Hamiltonian with a time dependent magnetic field. ArXiv 2024
- [4] Bambusi D., Langella B. Growth of Sobolev norms in quasi integrable quantum systems. ArXiv 2022.
- [5] Bambusi D., Langella B., Montalto R. Growth of Sobolev norms for unbounded perturbations of the Schrödinger equation on flat tori. J. Diff. Eq., 318: 344–358, 2022.
- [6] Baldi P., Berti M., Haus E., Montalto R., Time quasi-periodic gravity water waves in finite depth. Invent. Math. 214(2): 739-911, 2018.
- Biasi A., Evnin O. Turbulent cascades in a truncation of the cubic Szegő equation and related systems. Analysis & PDE, 15(1), 217–243, 2022.
- [8] Berti M., Cuccagna S., Gancedo F., Scrobogna S., Paralinearization and extended lifespan for solutions of the α -SQG sharp front equation. arXiv 2310.15963, 2023.
- Berti M., Delort J.-M., Almost Global Solutions of Capillary-gravity Water Waves Equations on the Circle. UMI Lecture Notes 2018, ISBN 978-3-319-99486-4.

- [10] Berti M., Feola R., Franzoi L., Quadratic life span of periodic gravity-capillary water waves. Water Waves 3(1): 85-115, 2021.
- [11] Berti M., Feola R., Pusateri F., Birkhoff Normal Form and Long Time Existence for Periodic Gravity Water Waves. Comm. Pure Applied Math., 76(7), 1416–1494, 2023.
- [12] Berti M., Franzoi L., Maspero A., Pure gravity traveling quasi-periodic water waves with constant vorticity. Comm. Pure Applied Math., 77(2), 990-1064, 2024.
- [13] Berti M., Maspero A., Murgante F., Local well posedness of the Euler-Korteweg equations on T^d. J. Dyn. Diff. Equat., 33: 1475-1513, 2021.
- Berti M., Maspero A., Murgante F., Hamiltonian Birkhoff normal form for gravity-capillary water waves with constant vorticity: almost global existence. preprint: arXiv:2212.12255,
- [15] Bourgain, J., On the growth in time of higher Sobolev norms of smooth solutions of Hamiltonian PDE.. Int. Math. Res. Notices 6:277–304, 1996.
- Bourgain J. Growth of Sobolev norms in linear Schrödinger equations with quasi-periodic potential. Comm. Math. Phys., 204(1):207-247, 1999.
- [17] J. Bourgain. On growth of Sobolev norms in linear Schrödinger equations with smooth time dependent potential. J. Anal. Math., 77:315–348, 1999.
- [18] Bourgain J. Problems in Hamiltonian PDE's. Geom. Funct. Anal., Special Volume, Part I:32–56, 2000.
- [19] Colin de Verdière Y. Spectral theory of pseudodifferential operators of degree 0 and an application to forced linear waves. Anal. PDE, 13(5):1521–1537, 2020.
- [20] Colin de Verdière Y., Saint-Raymond L. Attractors for two-dimensional waves with homogeneous Hamiltonians of degree 0. Comm. Pure Appl. Math., 73(2):421–462, 2020.
- [21] Colliander J., Keel M., Staffilani G., Takaoka H., Tao T. Transfer of energy to high frequencies in the cubic defocusing nonlinear Schrödinger equation. Invent. Math., 181(1):39–113, 2010.
- [22] Constantin A. and Escher J. Wave breaking for nonlinear nonlocal shallow water equations. Acta Math. 181 229–43, 1998.
- [23] Delort J.M. Growth of Sobolev norms of solutions of linear Schrödinger equations on some compact manifolds. Int. Math. Res. Not. IMRN, (12):2305–2328, 2010.
- [24] Delort J.M. Growth of Sobolev norms for solutions of time dependent Schrödinger operators with harmonic oscillator potential. Comm. Partial Differential Equations, 39(1):1–33, 2014.
- [25] Deng, Y., Hani, Z. Full derivation of the wave kinetic equation. Invent. math. 233, 543–724, 2023.
- [26] Elgindi T., Shikh Khalil K. Strong Ill-Posedness in L^{∞} for the Riesz Transform Problem arxiv.org/abs/2207.04556
- [27] Faou E., Raphael P. On weakly turbulent solutions to the perturbed linear harmonic oscillator. American Journal of Mathematics, 145(5): 1465–1507, 2023.
- [28] Feola R., Giuliani F. Quasi-periodic traveling waves on an infinitely deep fluid under gravity. Memoires American Mathematical Society, Vol. 295, 2024.
- [29] Feola R., Grebert B., Iandoli F. Long time solutions for quasi-linear Hamiltonian perturbations of Schrödinger and Klein-Gordon equations on tori Analysis & PDE, 16(5):1133–1203, 2023.
- [30] Feola R., Iandoli F. Long time existence for fully nonlinear NLS with small Cauchy data on the circle. Annali SNS, 22(1), 2021.
- [31] Gallone M., Marian M., Ponno A., Ruffo S. Burgers turbulence in the Fermi-Pasta-Ulam-Tsingou chain. Phys. Rev. Lett. 129, 114101, 2022.
- [32] Gérard P., Grellier S. The cubic Szegő equation. Ann Sci Éc Norm Supér 43(4), 761–810, 2010
- [33] Gérard P., Grellier S. The cubic Szegő equation and Hankel operators. Astérisque, (389):vi+112, 2017.
- [34] Gérard P., Grellier S. Effective integrable dynamics for a certain nonlinear wave equation. Anal PDE, 5, 1139–1155, 2012.
- [35] Gérard P., Lenzmann E. The Calogero-Moser Derivative Nonlinear Schrödinger Equation CPAM, 2024. doi:10.1002/cpa.22203
- [36] Giuliani F., Guardia M. Sobolev norms explosion for the cubic NLS on irrational tori. Nonlinear Analysis, 220, 112865, 2022.
- [37] Giuliani F., Guardia M. Arnold diffusion in Hamiltonian systems on infinite lattices. CPAM, 77(8): 3333–3426, 2024.
- [38] Giuliani F. Sobolev instability in the cubic NLS equation with convolution potentials on irrational tori. arXiv:2308.13468, 2023.
- [39] Guardia M., Haus E., Hani Z., Maspero A., Procesi M. Strong nonlinear instability and growth of Sobolev norms near quasiperiodic finite-gap tori for the 2D cubic NLS equation. J. Eur. Math. Soc. (JEMS), 25(4) 1497–1551, 2022.

- [40] Guardia M., Haus E., Procesi M. Growth of Sobolev norms for the analytic NLS on T². Adv. Math., 301:615–692, 2016.
- [41] Guardia M., Kaloshin V. Growth of Sobolev norms in the cubic defocusing nonlinear Schrödinger equation. J. Eur. Math. Soc. (JEMS), 17(1):71–149, 2015.
- [42] Hani, Z. Long-time instability and unbounded Sobolev orbits for some periodic nonlinear Schrödinger equations. Arch. Ration. Mech. Anal., 211(3):929–964, 2014.
- [43] Haus E., Procesi M. Growth of Sobolev norms for the quintic NLS on \mathbb{T}^2 . Anal. PDE, 8(4):883–922, 2015.
- [44] Hani Z., Pausader B., Tzvetkov N., and Visciglia N. Modified scattering for the cubic Schrödinger equation on product spaces and applications. Forum Math. Pi, 3:e4, 63, 2015.
- [45] Haus E., Maspero A. Growth of Sobolev norms in time dependent semiclassical anharmonic oscillators. J. Funct. Anal., 278(2), 108316, 2020.
- [46] Hur, V. Wave breaking in the Whitham equation. Adv. Math. 317, 410–437, 2017.
- [47] Hur V., Tao L. Wave breaking for the Whitham equation with fractional dispersion. Nonlinearity 27(12), 2937–2949, 2014.
- [48] Klein C., Saut J.-C. A numerical approach to blow-up issues for dispersive perturbations of Burgers' equation. Phys. D 295(296), 46–65, 2015.
- [49] Klein C., Saut J.-C., Wang, Y. On the modified fractional Korteweg-de Vries and related equations. Nonlinearity, 35(3), 1170, 2022.
- [50] Kuksin, S. Growth and oscillations of solutions of nonlinear Schrödinger equation. Comm. Math. Phys. 178, 265–280, 1996.
- [51] Kuksin, S. On turbulence in nonlinear Schrödinger equations. Geom. Funct. Anal. 7,783–822, 1997.
- [52] Liang Z., Zhao Z. and Zhou Q. 1-d quantum harmonic oscillator with time quasi-periodic quadratic perturbation: reducibility and growth of Sobolev norms. J. Math. Pures Appl. 146(1): 158–182 (2021).
- [53] Liang Z., Luo J., Zhao Z. Symplectic Normal Form and Growth of Sobolev Norm. arXiv:2312.16492, 2023.
- [54] Luo J., Liang Z. and Zhao Z. Growth of Sobolev Norms in 1-d Quantum Harmonic Oscillator with Polynomial Time Quasi-periodic Perturbation. Commun. Math. Phys. 392, 1–23, 2022.
- [55] Majda, A.J., McLaughlin, D.W. and Tabak, E.G. A one-dimensional model for dispersive wave turbulence. J Nonlinear Sci 7, 9–44, 1997.
- [56] Maspero A. Lower bounds on the growth of Sobolev norms in some linear time dependent Schrödinger equations. Math. Res. Lett., 26(4):1197–1215, 2019.
- [57] Maspero A. Growth of Sobolev norms in linear Schrödinger equations as a dispersive phenomenon. Advances in math, 411(A) 2022.
- [58] Maspero A. Generic transporters for the linear time dependent quantum Harmonic oscillator on ℝ. IMRN, rnac174 2022.
- [59] A. Maspero and D. Robert. On time dependent Schrödinger equations: Global well-posedness and growth of Sobolev norms. J. Fun. Anal., 273(2):721 – 781, 2017.
- [60] Montalto M., Murgante F., Scrobogna S. Quadratic lifespan for the sublinear α -SQG sharp front problem. arXiv:2402.06364, 2024.
- [61] Mourre E. Absence of singular continuous spectrum for certain selfadjoint operators. Comm. Math. Phys., 78(3):391–408, 1980/81.
- [62] Oh, SJ., Pasqualotto, F. Gradient Blow-Up for Dispersive and Dissipative Perturbations of the Burgers Equation. Arch Rational Mech Anal 248, 54, 2024.
- [63] Planchon, F., Tzvetkov, N., Visciglia, N. On the growth of Sobolev norms for NLS on 2- and 3-dimensional manifolds. Anal. PDE 10, 1123–1147, 2017.
- [64] Saut J.-C., Wang Y. Global dynamics of small solutions to the modified fractional Korteweg-de Vries and fractional cubic nonlinear Schrödinger equations. Commun. Part. Differ. Equ. 46(10), 1851–1891, 2021.
- [65] Sigal I., Soffer A. Local decay and velocity bounds for quantum propagation. preprint (Princeton), 1988. http://www.math.toronto.edu/sigal/publications/SigSofVelBnd.pdf
- [66] Sohinger, V. Bounds on the growth of high Sobolev norms of solutions to nonlinear Schrödinger equations on S¹. Differential Integral Equations, 24(7-8):653–718, 2011.
- [67] Staffilani G.. On the growth of high Sobolev norms of solutions for KdV and Schrödinger equations. Duke Math. J., 86:109–142, 1997.
- [68] Yang R. Shock formation for the Burgers-Hilbert equation. SIAM J. Math. Anal. 53(5), 2021.
- [69] Zakharov V.E. and Dyachenko A.I. Is free-surface hydrodynamics an integrable system? Physics Letters A 190, 144-148, 1994.
- [70] Zakharov V., Diasc F, Pushkarev A. One-dimensional wave turbulence. Physics Reports 398, 1 65, 2004.