

Model Independent Tests of the Hadronic Vacuum Polarization Contribution to the Muon $g-2$

Luca Di Luzio,¹ Alexander Keshavarzi,² Antonio Masiero,^{3, 1} and Paride Paradisi^{3, 1}

¹*Istituto Nazionale di Fisica Nucleare, Sezione di Padova, Via F. Marzolo 8, 35131 Padova, Italy*

²*Department of Physics and Astronomy, The University of Manchester, Manchester M13 9PL, U.K.*

³*Dipartimento di Fisica e Astronomia ‘G. Galilei’,*

Università di Padova, Via F. Marzolo 8, 35131 Padova, Italy

The hadronic vacuum polarization (HVP) contributions to the muon $g-2$ are the crucial quantity to resolve whether new physics is present or not in the comparison between the Standard Model (SM) prediction and experimental measurements at Fermilab. They are commonly and historically determined via dispersion relations using a vast catalogue of experimentally measured, low-energy $e^+e^- \rightarrow$ hadrons cross section data as input. These dispersive estimates result in a SM prediction that exhibits a muon $g-2$ discrepancy of more than 5σ when compared to experiment. However, recent lattice QCD evaluations of the HVP and a new hadronic cross section measurement from the CMD-3 experiment favor a no-new-physics scenario and, therefore, exhibit a common tension with the previous $e^+e^- \rightarrow$ hadrons data. This study explores the current and future implications of these two scenarios on other observables that are also sensitive to the HVP contributions in the hope that they may provide independent tests of the current tensions observed in the muon $g-2$.

Introduction.— The anomalous magnetic moment of the muon, $a_\mu \equiv (g_\mu - 2)/2$, stands amongst the best probes of the Standard Model (SM) of particle physics and its possible extensions. Recently, the Muon $g-2$ Experiment at Fermilab announced a new experimental (exp) measurement of a_μ [1, 2] which, when combined with the previous and consistent results from the same experiment [3–6] (and from the earlier Muon $g-2$ Experiment at Brookhaven [7–9]), results in a new world average of $a_\mu^{\text{exp}} = 116592059(22) \times 10^{-11}$ with an unprecedented precision of 190 parts-per-billion (ppb).

The most recent, community-approved SM prediction of a_μ from the Muon $g-2$ Theory Initiative [10–30] relies on the hadronic vacuum polarization (HVP) contribution, a_μ^{HVP} , being entirely evaluated using the data-driven, dispersive approach [15–25, 31–45]. This method uses measured $e^+e^- \rightarrow$ hadrons cross section data (σ_{had}) as input and the leading-order (LO) HVP contribution was found by the Theory Initiative to be [10, 15–20]

$$(a_\mu^{\text{HVP}})_{e^+e^-}^{\text{TI}} = (693.1 \pm 4.0) \times 10^{-10}. \quad (1)$$

This, when combined with the updated values for the other SM contributions, resulted in a SM prediction of $a_\mu^{\text{SM}} = 116591810(43) \times 10^{-11}$ [10–30]. Comparing a_μ^{SM} with the current experimental world average leads to

$$\Delta a_\mu \equiv a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (24.9 \pm 4.8) \times 10^{-10}, \quad (2)$$

yielding a discrepancy of 5.1σ and implying the presence of physics beyond the SM (BSM).

a_μ^{HVP} can now be calculated from first-principles lattice QCD methods [10] (see also [46–57]). In particular, the BMW lattice QCD collaboration (BMWc) first computed a_μ^{HVP} with sub per-cent precision [56] and found $(a_\mu^{\text{HVP}})_{\text{Lattice}}^{\text{BMW}} = (707.5 \pm 5.5) \times 10^{-10}$, which is evidently larger than the Theory Initiative value [10]. They have since updated their result to [57]

$$(a_\mu^{\text{HVP}})_{\text{Lattice}}^{\text{BMW}} = (714.1 \pm 3.3) \times 10^{-10}, \quad (3)$$

which reduces Δa_μ to 0.9σ (fully consistent with a_μ^{exp} at the current level of precision) at the expense of generating a 4.0σ tension with Eq. (1). These results have been partially confirmed by other lattice QCD studies of the so-called window quantities [47, 58–62], which further increases the tension between lattice QCD evaluations and $(a_\mu^{\text{HVP}})_{e^+e^-}^{\text{TI}}$ in the intermediate window [63] where the lattice QCD computations are most proficient.

Increasing the hadronic cross section to accommodate Δa_μ in $(a_\mu^{\text{HVP}})_{e^+e^-}^{\text{TI}}$ requires shifts below ~ 1 GeV to avoid inconsistency with the electroweak precision fits [64–68]. Indeed, a large portion of the differences observed between $(a_\mu^{\text{HVP}})_{e^+e^-}^{\text{TI}}$ and lattice QCD evaluations dominantly originate from the light-quark-connected contributions (which in-turn are dominated by $\pi^+\pi^-$) [69, 70]. The potential for light new physics affecting the interpretation of e^+e^- data was also investigated in [71–75].

Recently, the CMD-3 experiment released a new measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section [76, 77]. It covers the energy range from 0.32 to 1.2 GeV, therefore capturing the dominant ρ -resonance region, and is $\sim 2\sigma$ – 4σ higher than all previous $e^+e^- \rightarrow \pi^+\pi^-$ cross section measurements in the same region [78–89]. Although $(a_\mu^{\pi\pi})_{e^+e^-}^{\text{CMD3}}$ covers a limited energy range, it goes in the same direction as the BMWc result of Eq. (3), thereby also suggesting consistency between a_μ^{exp} and a_μ^{SM} .

Several updates and improvements on all fronts are expected soon which may help shed light on the current puzzling picture. New results of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section are expected from the BaBar [90], Belle II [91], BESIII [92], CMD-3 [93], KLOE [94] and SND [95] experiments in the next 2–4 years, with theoretical improvements also expected for the higher-order QED corrections which are paramount for these measurements (see e.g. [96]). New full HVP determinations from other lattice QCD collaborations are also expected soon, as well as new dispersive determinations of the HVP incorporating new $e^+e^- \rightarrow$ hadrons cross section data. In ad-




FIG. 1. Representative Feynman diagrams for observables that are sensitive to the leading HVP contribution: leptonic $g-2$ (diagram (a)), running of α and $\sin^2 \theta_W$ (diagram (b)), and Muonium HFS (diagrams (c) and (d)).

dition, a direct experimental measurement of a_μ^{HVP} , has been proposed and is in preparation by the MUonE experiment [97–101].

This letter adds to this catalogue of probes other observables that are sensitive to HVP effects and, therefore, be used as additional indirect tools to scrutinize the differences in a_μ^{HVP} . The new data from CMD-3 are used to perform numerical analyses and provide analytical expressions relating the HVP contributions of other observables to a_μ^{HVP} . The current and future implications of the CMD-3 data (and, by data-based proxy, the increase of $(a_\mu^{\text{HVP}})^{\text{BMW}}$ compared to dispersive estimates) are assessed and quantified for the electron, muon and tau $g-2$, the running of the QED coupling constant (α), the low-energy weak mixing angle ($\sin^2 \theta_W(0)$) and the Muonium hyperfine splitting (HFS). The conclusions drawn focus upon how new, precise experimental measurements of these observables could be sensitive to the differences observed due to the CMD-3 data in a_μ^{HVP} and, therefore, provide additional robust tests of the current tensions.

Impact of the CMD-3 data.— The CMD-3 measurement of the $\pi^+\pi^-$ final state [76] is larger than all previous $\pi^+\pi^-$ measurements by 2σ - 4σ , resulting in it being comparatively larger than the compilation by the KNT collaboration in 2019 (referred to as KNT19) [20]. Given it is not expected that the differences between these measurements will be reconciled, and that no new compilation of $\pi^+\pi^-$ measurements including the CMD-3 data has yet been performed, two scenarios can be compared and analysed: (1) $\sigma_{\text{had}} = \text{KNT19}$. The previous data are correct and the new CMD-3 data are not used. (2) $\sigma_{\text{had}} = \text{KNT19}/\text{CMD-3}$. The CMD-3 data are correct therefore are substituted into and replace the KNT19 data only in their available energy range.

The electron, muon and tau $g-2$, the running of the QED coupling constant, the low-energy weak mixing angle and the Muonium HFS are all sensitive to HVP effects (see e.g. Fig. 1). To understand the impact of the CMD-3 data on the HVP contributions to these observable, $O_{e^+e^-}^{\text{HVP}}$, the difference between the scenario (1) and scenario (2) is calculated as

$$\delta O^{\text{CMD3}} \equiv (O_{e^+e^-}^{\text{HVP}})^{\text{CMD3}} - (O_{e^+e^-}^{\text{HVP}})^{\text{KNT19}}. \quad (4)$$

All results are given in Table I. Other than the region of the replaced $\pi^+\pi^-$ data, the two scenarios are correlated

for all common KNT19 data by a factor given by ρ_{12} . The statistical significance of each δO^{CMD3} in Table I and depicted in Fig. 2 includes this correlation coefficient. The results for each $O_{e^+e^-}^{\text{HVP}}$ are discussed in the following, starting with the muon $g-2$ as the original source of the tensions that motivates this study.

Muon $g-2$.— The LO HVP contribution to the muon $g-2$ is calculated via

$$(a_\mu^{\text{HVP}})_{e^+e^-} = \frac{1}{4\pi^3} \int_{m_{\pi^0}^2}^\infty ds K_\mu(s) \sigma_{\text{had}}(s). \quad (5)$$

The positive-definite kernel function $K_\mu(s)$ is given by

$$K_\mu(s) = \int_0^1 dx \frac{x^2(1-x)}{x^2 + (1-x)s/m_\mu^2}, \quad (6)$$

with $K_\mu(s) \approx m_\mu^2/3s$ in the high-energy limit $\sqrt{s} \gg m_\mu$.

The KNT19 evaluation (scenario (1)) was one of several inputs to the estimate found in Eq. (1) and found the LO HVP contribution to be [20]

$$(a_\mu^{\text{HVP}})^{\text{KNT19}}_{e^+e^-} = (692.8 \pm 2.4) \times 10^{-10}. \quad (7)$$

Using scenario (2) (substituting in the CMD-3 data) [76] increases this estimate to

$$(a_\mu^{\text{HVP}})^{\text{CMD3}}_{e^+e^-} = (714.5 \pm 3.4) \times 10^{-10}, \quad (8)$$

which is in better agreement with the BMWc value of Eq. (3). This results in a shift in a_μ^{HVP} of

$$\begin{aligned} \delta a_\mu^{\text{CMD3}} &\equiv (a_\mu^{\text{HVP}})^{\text{CMD3}}_{e^+e^-} - (a_\mu^{\text{HVP}})^{\text{KNT19}}_{e^+e^-} \\ &= (21.7 \pm 3.6) \times 10^{-10}, \end{aligned} \quad (9)$$

exhibiting a tension of 6.1σ . In the following, the phenomenological implications of the above shift on $O_{e^+e^-}^{\text{HVP}}$ (see Table I) are analyzed, followed by a discussion regarding the required experimental and theoretical improvements needed to probe the disagreement in Eq. (9).

Electron $g-2$.— The anomalous magnetic moment of the electron, a_e , is commonly used to determine the value of the fine-structure constant, α (and vice versa). Recent, more-refined atomic-physics experiments using Cesium

$O_{e^+e^-}^{\text{HVP}}$	(1) KNT19	(2) KNT19/CMD3	Correlation, ρ_{12}	Difference, δO^{CMD3}	Significance (σ)
$a_\mu^{\text{HVP}} \times 10^{10}$	692.8 ± 2.4	714.5 ± 3.4	0.280	21.7 ± 3.6	6.1
$a_e^{\text{HVP}} \times 10^{14}$	186.1 ± 0.7	192.0 ± 0.0	0.257	6.0 ± 1.0	6.2
$a_\tau^{\text{HVP}} \times 10^8$	332.8 ± 1.4	340.2 ± 2.1	0.546	7.4 ± 1.8	4.2
$\Delta\alpha_{\text{had}}^{(5)}(M_Z^2) \times 10^4$	276.1 ± 1.1	277.5 ± 1.2	0.908	1.4 ± 0.5	2.8
$\sin^2\theta_W(0) \times 10^4$	2386.0 ± 1.4	2386.4 ± 1.5	0.996	0.4 ± 0.1	2.9
$\nu_{\text{HFS}}^{\text{HVP}} (\text{Hz})$	540.5 ± 1.9	557.0 ± 2.7	0.297	16.5 ± 2.8	5.9

TABLE I. HVP contributions to the studied observables [first column] using scenario (1) KNT19 data [second column], or scenario (2) KNT19/CMD3 data [third column]. The fourth column gives the degree of correlation between scenario (1) and scenario (2) from the common KNT19 data (and other factors). The fifth column lists the difference or shift induced by the CMD-3 data as $\delta O^{\text{CMD3}} \equiv (O_{e^+e^-}^{\text{HVP}})^{\text{CMD3}} - (O_{e^+e^-}^{\text{HVP}})^{\text{KNT19}}$, where the error accounts for the correlation ρ_{12} . The sixth column then quantifies the statistical significance of this shift in standard deviations.




FIG. 2. The tension observed in standard deviations (σ) between dispersive evaluations of HVP contributions to observables sensitive to such effects when the $e^+e^- \rightarrow \text{hadrons}$ data used as input as those given in scenario (1) (KNT19) [20] or scenario (2) KNT19/CMD3 [76]. The black points correspond to scenario (1) and the red points show the difference in σ due to scenario (2). The results leading to these tensions are given in Table I.

(Cs) and Rubidium (Rb) interferometry have led to the following results for direct measurements of α :

$$\alpha(\text{Cs}) = 1/137.035999046(27) \quad [102], \quad (10)$$

$$\alpha(\text{Rb}) = 1/137.035999206(11) \quad [103], \quad (11)$$

in disagreement with each other by 5.5σ . Yet, a_e can be extracted and used as another precision test of the SM and its possible extensions [104, 105]. Comparing the resulting SM predictions, a_e^{SM} , with the latest experimental measurement of $a_e^{\text{exp}} = (115\,965\,218\,059 \pm 13) \times 10^{-14}$ [106], yields

$$(\Delta a_e)_{\text{Cs}} = (-102.0 \pm 26.4) \times 10^{-14} \Rightarrow -3.9\sigma, \quad (12)$$

$$(\Delta a_e)_{\text{Rb}} = (33.8 \pm 16.1) \times 10^{-14} \Rightarrow 2.1\sigma, \quad (13)$$

where $\Delta a_e \equiv a_e^{\text{exp}} - a_e^{\text{SM}}$.

The impact of the CMD-3 data on the HVP contribution to a_e^{SM} can first be estimated by reconsidering the high-energy approximation of Eq. (6) and relating a_μ^{HVP} and a_e^{HVP} by a quadratic-scaling between the lepton masses. In doing so, and by employing the value of $\delta a_\mu^{\text{CMD3}}$ quoted in Eq. (9), results in

$$\delta a_e^{\text{CMD3}} \approx \delta a_\mu^{\text{CMD3}} \left(\frac{m_e}{m_\mu} \right)^2 \approx (5.1 \pm 0.8) \times 10^{-14}. \quad (14)$$

This is in good agreement with the numerical result in Table I, which shows a 6.2σ shift from the CMD-3 data

compared to the previous data. It follows that the electron $g-2$ will only be sensitive to the increase observed on a_μ^{HVP} if the current discrepancy in the measurements of α is resolved and a precision at the level of $\mathcal{O}(10^{-14})$ is achieved for Δa_e .

Consider the sources of error in the determinations of $(\Delta a_e)_{\text{Cs}}$ and $(\Delta a_e)_{\text{Rb}}$: It follows that should the ex-

Δa_e error source	Value	% of Δa_e error
Five-loop QED, $\delta a_e^{\text{QED}5}$	6×10^{-14}	5% (Cs)/13% (Rb)
Hadronic, δa_e^{HAD}	1×10^{-14}	< 1%
$\alpha(\text{Cs}), \delta a_e^{\alpha(\text{Cs})}$	22×10^{-14}	70%
$\alpha(\text{Rb}), \delta a_e^{\alpha(\text{Rb})}$	9×10^{-14}	28%
Experiment, δa_e^{exp}	13×10^{-14}	24% (Cs)/59% (Rb)

perimental resolutions on α and a_e^{exp} improve by roughly one order of magnitude, the total uncertainties of Δa_e should reach the $\mathcal{O}(10^{-14})$ level and the study of electron $g-2$ will become sensitive to the increase observed in a_μ^{HVP} due to CMD-3 (and BMWc). Fortunately, an improvement on δa_e^{exp} by a factor of ~ 5 is expected in the next few years.¹ On the same timescale, new measurements of $\alpha(\text{Rb})$ and $\alpha(\text{Cs})$ by the Paris and Berkeley collaborations, respectively, will significantly reduce the systematic effects that were dominant in their previous measurements.² Such improvements should be sufficient to probe the shift induced by CMD-3 (and BMWc) through the electron $g-2$.

Additionally, $\delta a_e^{\text{QED}5}$ reflects the current 5σ discrepancy between the two independent calculations of the gauge-invariant set of 5-loop QED diagrams with no lepton loops by Kinoshita et al. [107] and Volkov [108, 109]. As a non-negligible source of uncertainty on Δa_e , and with expected improvements in the experimental inputs, resolving this difference may soon become crucial in this context.³

Tau g-2.— The shift in a_τ^{SM} due to the CMD-3 data is estimated by assuming dominant effects at the ρ -peak:

$$\begin{aligned} \delta a_\tau^{\text{CMD}3} &\approx 0.63 (m_\rho^2/m_\mu^2) \delta a_\mu^{\text{CMD}3} \\ &\approx (7.2 \pm 1.4) \times 10^{-8}, \end{aligned} \quad (15)$$

which is in very good agreement with the numerical estimate in Table I. Compared to the electron and muon $g-2$, the larger mass of the tau lepton increases the weight of the hadronic contributions to higher energies. As such, the influence of the $\pi^+\pi^-$ and ρ -resonance contribution is reduced in $(a_\tau^{\text{HVP}})_{e^+e^-}$, resulting in a much larger degree of correlation between the two scenarios for the tau ($\sim 55\%$) compared to the muon ($\sim 28\%$).

¹ G. Gabrielse, private communication.

² S. Guellati-Khélifa and H. Mueller, private communications.

³ S. Laporta, work in progress.

This means the significance of $\delta a_\tau^{\text{CMD}3}$ is 4.2σ compared to more than 6σ for $(a_e^{\text{HVP}})_{e^+e^-}$ and $(a_\mu^{\text{HVP}})_{e^+e^-}$.

As $\delta a_\tau^{\text{CMD}3}$ is roughly one order of magnitude below the expected future experimental sensitivity of $\mathcal{O}(10^{-6})$ at Belle-II [110], an experiment-theory comparison is unlikely to shed light on the current discrepancies in the SM predictions of the muon $g-2$ without significant improvement in measurements of a_τ^{exp} .

The running of α .— Hadronic effects to the running QED coupling at the Z -boson mass, $\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)$, are a principal component of the electroweak precision fit. The shift in $\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)$ due to the CMD-3 data is estimated as

$$\begin{aligned} \delta\Delta\alpha_{\text{had}}^{(5)}(M_Z^2) &\approx \frac{3\pi}{\alpha} \frac{m_\rho^2}{m_\mu^2} \delta a_\mu^{\text{CMD}3} \\ &\approx (1.5 \pm 0.3) \times 10^{-4}, \end{aligned} \quad (16)$$

in full agreement with Table I. The relevant kernel function in the dispersive evaluation of $\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)$ exhibits an s -dependent weighting that is much higher in energy than any of the dispersion relations used to evaluate $(a_l^{\text{HVP}})_{e^+e^-}$. As a result, the CMD-3 increase has less of an impact than in any of the $g-2$ cases, exhibiting a 2.8σ deviation. The magnitude of the shift is comparable with the current uncertainty of $\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)$, making it difficult for $\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)$ to probe $\delta a_\mu^{\text{CMD}3}$. However, looking ahead, a future e^+e^- collider, e.g. FCC-ee [111, 112], is expected to reach an unprecedented precision on $\alpha(M_Z^2)$ of $\mathcal{O}(10^{-5})$, which would provide sensitivity to the shift in Eq. (16).

The running of $\sin^2\theta_W$.— The low-energy weak mixing angle, defined as $\sin^2\theta_W(0)$ at zero momentum transfer ($q^2 = 0$), has been measured at parity-violating electron scattering and atomic physics asymmetry experiments [113, 114]. Following the prescription in [66], $\sin^2\theta_W(0)$ can be connected with the high-energy LEP measurement of the same quantity, $\sin^2\theta_W(M_Z)$, by including quantum effects from the $\gamma - Z$ mixing [115].

The shift from the CMD-3 data is estimated via

$$\begin{aligned} \delta \sin^2\theta_W(0) &\approx k' \sin^2\theta_W(M_Z) \frac{3\pi}{\alpha} \frac{m_\rho^2}{m_\mu^2} \delta a_\mu^{\text{CMD}3} \\ &\approx (0.4 \pm 0.1) \times 10^{-4}, \end{aligned} \quad (17)$$

where $\sin^2\theta_W(M_Z) = 0.23119(14)$ [113, 114] and $k' = 1.14$ [115]. This is again in full agreement with the numerical estimate in Table I. The comparison exhibits a similar significance (2.9σ) as for $\delta\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)$.

Unfortunately, current experimental errors prevent $\sin^2\theta_W(0)$ to probe the HVP contributions, which would require a precision on both $\sin^2\theta_W(0)$ and $\sin^2\theta_W(M_Z)$ at the $\sim 10^{-5}$ level. While future e^+e^- high-energy colliders aim at a resolution on $\sin^2\theta_W(M_Z)$ much better than $\mathcal{O}(10^{-5})$ [112], experimental progress on $\sin^2\theta_W(0)$ planned at both MESA (P2) in Mainz [116] and JLab (Møller) [117] is projected to reach only $\sim \mathcal{O}(10^{-4})$.

The Muonium HFS.— An alternative method to extract a_μ^{HVP} is via Muonium, a bound state of an antimuon and an electron ($M \equiv \mu^+ e^-$). The Muonium HFS of the 1S ground state is described by the expression [118–121]

$$\frac{\nu_{\text{HFS}}}{\nu_F} = 1 + a_\mu + \Delta_{\text{HFS}}^{\text{QED}} + \Delta_{\text{HFS}}^{\text{weak}} + \Delta_{\text{HFS}}^{\text{HVP}}, \quad (18)$$

where ν_F is given by

$$\nu_F = \frac{16}{3} \frac{m_e}{m_\mu} \frac{R_\infty h c \alpha^2}{(1 + m_e/m_\mu)^3}, \quad (19)$$

and R_∞ is the Rydberg constant. The quantities $\Delta_{\text{HFS}}^{\text{QED}}$ and $\Delta_{\text{HFS}}^{\text{weak}}$ in Eq. (18) refer to corrections stemming from QED and weak contributions, respectively. $\Delta_{\text{HFS}}^{\text{HVP}}$ accounts for HVP effects [120] which are given by [120]

$$\Delta_{\text{HFS}}^{\text{HVP}} = \frac{1}{2\pi^3} \int_{m_\pi^2}^\infty ds K_{\text{Mu}}(s) \sigma_{\text{had}}(s), \quad (20)$$

where, for $s \gg m_\mu^2$, the kernel function $K_{\text{Mu}}(s)$ reads

$$K_{\text{Mu}}(s) \approx \frac{m_\mu^2}{s} \left(\frac{9}{2} \log \frac{s}{m_\mu^2} + \frac{15}{4} \right) \frac{m_e}{m_\mu}. \quad (21)$$

Again, as the dominant effects to σ_{had} arise at the ρ -peak, $s \approx m_\rho^2 \gg m_\mu^2$. Using Eqs. (5) and (20) results in

$$\begin{aligned} \Delta_{\text{HFS}}^{\text{HVP}} &\approx 6 \frac{m_\rho^2}{m_\mu^2} K_{\text{Mu}}(m_\rho^2) (a_\mu^{\text{HVP}})_{e^+ e^-} \\ &\approx 0.63 (a_\mu^{\text{HVP}})_{e^+ e^-}. \end{aligned} \quad (22)$$

Therefore, using also Eq. (18), the entire HVP contribution to ν_{HFS} can be expressed as

$$\nu_{\text{HFS}}^{\text{HVP}} = (a_\mu^{\text{HVP}} + \Delta_{\text{HFS}}^{\text{HVP}}) \nu_F \approx 1.63 \nu_F a_\mu^{\text{HVP}}, \quad (23)$$

where the two terms are depicted in the third and fourth diagrams of Fig. 1. Therefore, the impact of the CMD-3 data on $\nu_{\text{HFS}}^{\text{HVP}}$ is estimated to be

$$\delta\nu_{\text{HFS}}^{\text{CMD3}} \approx (15.8 \pm 2.6) \text{ Hz}, \quad (24)$$

in excellent agreement with the numerical evaluation in Table I and exhibits a 5.9σ discrepancy in the comparison. This makes the Muonium HFS among the most sensitive probes of $\delta a_\mu^{\text{CMD3}}$.⁴ To be sensitive to this shift would require a measurement at a precision of $\mathcal{O}(1)$ Hz. Fortunately, the MuSEUM experiment at J-PARC aims to reduce the uncertainty of the current measurement $\nu_{\text{HFS}}^{\text{exp}} = (4463302776 \pm 51)$ Hz [123, 124] by one order of magnitude [125], well below the shift of Eq. (24).

⁴ In Ref. [122], it has been also proposed to determine a_μ^{HVP} from the measurement of ν_{HFS} . However, $e^+ e^- \rightarrow$ hadrons data have been used to estimate $\Delta_{\text{HFS}}^{\text{HVP}}$, making the procedure inconsistent.

Other sources of uncertainty are also set to improve. The uncertainty on ν_F is fully dominated by the m_e/m_μ ratio (the uncertainties of R_∞ and α are negligible) [118]. Here, m_e/m_μ is obtained from the measurement of the 1S–2S transition frequency in Muonium given by

$$\nu_{1\text{S}-2\text{S}} = \frac{3}{4} \frac{R_\infty hc}{(1 + m_e/m_\mu)} [1 + \delta_{1\text{S}-2\text{S}}], \quad (25)$$

where the muon mass enters as a recoil contribution. As the hadronic corrections to $\delta_{1\text{S}-2\text{S}}$ [126] do not spoil the extraction of m_e/m_μ at the required precision, the dominant uncertainty on m_e/m_μ arises from the experimental error in $\nu_{1\text{S}-2\text{S}}$ [127]. This induces an error on ν_F of 4×10^3 Hz. Experimental progress is ongoing at the MuMASS experiment to improve this precision by three orders of magnitude [128]. Additionally, the dominant theory uncertainty in ν_{HFS} arises from unknown three-loop QED contributions [119] to $\delta_{\text{HFS}}^{\text{QED}} (\delta_{1\text{S}-2\text{S}})$ and amounts to about 70 Hz (7 Hz) [119]. A complete three-loop QED calculation would bring the related uncertainty to a negligible level [129].

Conclusions and outlook.— Recent lattice QCD calculations and new measurements by the CMD-3 collaboration show a significant tension with the low-energy $e^+ e^- \rightarrow$ hadrons data in the determination of the leading HVP contribution to the muon $g-2$. Several model-independent tests of the HVP contribution are possible through the electron and tau $g-2$, the running of α and of the weak mixing angle, and the Muonium HFS. Both numerical analyses based on state-of-the-art $e^+ e^- \rightarrow$ hadrons data and approximate analytical expressions relating the HVP contributions of the considered observables have revealed the electron $g-2$ and the Muonium HFS to be the most promising prospects (see Fig. 2), offering unique opportunities to shed light on the current $g-2$ puzzle.

To achieve this ambitious goal, both experimental and theoretical improvements are needed. For the $g-2$ of the electron, the required precision level of $\mathcal{O}(10^{-14})$ is the expected future resolution for a_e^{exp} and α from atomic physics. In the case of the Muonium HFS, the MuSEUM experiment is expected to improve the HFS ground-state measurement by one order of magnitude, while the MuMASS experiment is expected to determine the m_e/m_μ ratio with a precision three orders of magnitude better than the present situation. Finally, a better theoretical control of QED effects (five loop contributions in the case of the electron $g-2$ and three loops in the case of Muonium HFS) is necessary. Although challenging, the above program seems to be feasible and it should be pursued with high priority to resolve significant tensions in one of the most stringent probes of the SM.

Acknowledgments.— We thank G. Colangelo, P. Crivelli, R. Frezzotti, G. Gabrielse, S. Guellati-Khélifa, M. Incagli, K. Kirch, S. Laporta, L. Lellouch, V. Lubicz, H. Mueller, M. Neubert, M. Passera, F. Piccinini, M. Pospelov, N. Tantalo and G. Venanzoni for private

communications and useful discussions. Special thanks are extended to D. Nomura and T. Teubner for their collaboration with AK in producing the KNT19 data. The work of AK is supported by The Royal Society (URF\R1\231503). The work of LDL and PP is supported by the European Union – Next Generation EU and by the Italian Ministry of University and Research (MUR) via the PRIN 2022 project n. 2022K4B58X – Ax-

ionOrigins. This work received funding by the INFN Iniziative Specifiche APINE and TASP and from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreements n. 860881 – HIDDeN, n. 101086085 – ASYMMETRY. This work was also partially supported by the Italian MUR Departments of Excellence grant 2023-2027 “Quantum Frontiers”.

-
- [1] D. P. Aguillard et al. (Muon g-2), *Phys. Rev. Lett.* **131**, 161802 (2023), arXiv:2308.06230 [hep-ex].
 - [2] D. P. Aguillard et al. (Muon g-2), (2024), arXiv:2402.15410 [hep-ex].
 - [3] B. Abi et al. (Muon g-2), *Phys. Rev. Lett.* **126**, 141801 (2021), arXiv:2104.03281 [hep-ex].
 - [4] T. Albahri et al. (Muon g-2), *Phys. Rev. Accel. Beams* **24**, 044002 (2021), arXiv:2104.03240 [physics.acc-ph].
 - [5] T. Albahri et al. (Muon g-2), *Phys. Rev. A* **103**, 042208 (2021), arXiv:2104.03201 [hep-ex].
 - [6] T. Albahri et al. (Muon g-2), *Phys. Rev. D* **103**, 072002 (2021), arXiv:2104.03247 [hep-ex].
 - [7] G. W. Bennett et al. (Muon g-2), *Phys. Rev. D* **73**, 072003 (2006), arXiv:hep-ex/0602035.
 - [8] G. W. Bennett et al. (Muon g-2), *Phys. Rev. Lett.* **89**, 101804 (2002), [Erratum: Phys.Rev.Lett. 89, 129903 (2002)], arXiv:hep-ex/0208001.
 - [9] G. W. Bennett et al. (Muon g-2), *Phys. Rev. Lett.* **92**, 161802 (2004), arXiv:hep-ex/0401008.
 - [10] T. Aoyama et al., *Phys. Rept.* **887**, 1 (2020), arXiv:2006.04822 [hep-ph].
 - [11] T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, *Phys. Rev. Lett.* **109**, 111808 (2012), arXiv:1205.5370 [hep-ph].
 - [12] T. Aoyama, T. Kinoshita, and M. Nio, *Atoms* **7**, 28 (2019).
 - [13] A. Czarnecki, W. J. Marciano, and A. Vainshtein, *Phys. Rev.* **D67**, 073006 (2003), [Erratum: Phys. Rev. **D73**, 119901 (2006)], arXiv:hep-ph/0212229 [hep-ph].
 - [14] C. Gnendiger, D. Stöckinger, and H. Stöckinger-Kim, *Phys. Rev.* **D88**, 053005 (2013), arXiv:1306.5546 [hep-ph].
 - [15] M. Davier, A. Hoecker, B. Malaescu, and Z. Zhang, *Eur. Phys. J.* **C77**, 827 (2017), arXiv:1706.09436 [hep-ph].
 - [16] A. Keshavarzi, D. Nomura, and T. Teubner, *Phys. Rev.* **D97**, 114025 (2018), arXiv:1802.02995 [hep-ph].
 - [17] G. Colangelo, M. Hoferichter, and P. Stoffer, *JHEP* **02**, 006 (2019), arXiv:1810.00007 [hep-ph].
 - [18] M. Hoferichter, B.-L. Hoid, and B. Kubis, *JHEP* **08**, 137 (2019), arXiv:1907.01556 [hep-ph].
 - [19] M. Davier, A. Hoecker, B. Malaescu, and Z. Zhang, *Eur. Phys. J.* **C80**, 241 (2020), [Erratum: Eur. Phys. J. **C80**, 410 (2020)], arXiv:1908.00921 [hep-ph].
 - [20] A. Keshavarzi, D. Nomura, and T. Teubner, *Phys. Rev.* **D101**, 014029 (2020), arXiv:1911.00367 [hep-ph].
 - [21] A. Kurz, T. Liu, P. Marquard, and M. Steinhauser, *Phys. Lett.* **B734**, 144 (2014), arXiv:1403.6400 [hep-ph].
 - [22] K. Melnikov and A. Vainshtein, *Phys. Rev. D* **70**, 113006 (2004), arXiv:hep-ph/0312226 [hep-ph].
 - [23] P. Masjuan and P. Sánchez-Puertas, *Phys. Rev.* **D95**, 054026 (2017), arXiv:1701.05829 [hep-ph].
 - [24] G. Colangelo, M. Hoferichter, M. Procura, and P. Stoffer, *JHEP* **04**, 161 (2017), arXiv:1702.07347 [hep-ph].
 - [25] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold, and S. P. Schneider, *JHEP* **10**, 141 (2018), arXiv:1808.04823 [hep-ph].
 - [26] A. Gérardin, H. B. Meyer, and A. Nyffeler, *Phys. Rev.* **D100**, 034520 (2019), arXiv:1903.09471 [hep-lat].
 - [27] J. Bijnens, N. Hermansson-Truedsson, and A. Rodríguez-Sánchez, *Phys. Lett.* **B798**, 134994 (2019), arXiv:1908.03331 [hep-ph].
 - [28] G. Colangelo, F. Hagelstein, M. Hoferichter, L. Laub, and P. Stoffer, *JHEP* **03**, 101 (2020), arXiv:1910.13432 [hep-ph].
 - [29] T. Blum, N. Christ, M. Hayakawa, T. Izubuchi, L. Jin, C. Jung, and C. Lehner, *Phys. Rev. Lett.* **124**, 132002 (2020), arXiv:1911.08123 [hep-lat].
 - [30] G. Colangelo, M. Hoferichter, A. Nyffeler, M. Passera, and P. Stoffer, *Phys. Lett.* **B735**, 90 (2014), arXiv:1403.7512 [hep-ph].
 - [31] S. J. Brodsky and E. De Rafael, *Phys. Rev.* **168**, 1620 (1968).
 - [32] B. E. Lautrup and E. De Rafael, *Phys. Rev.* **174**, 1835 (1968).
 - [33] B. Krause, *Phys. Lett. B* **390**, 392 (1997), arXiv:hep-ph/9607259.
 - [34] F. Jegerlehner, Vol. 274 (Springer, Cham, 2017).
 - [35] F. Jegerlehner, *EPJ Web Conf.* **118**, 01016 (2016), arXiv:1511.04473 [hep-ph].
 - [36] F. Jegerlehner, *EPJ Web Conf.* **166**, 00022 (2018), arXiv:1705.00263 [hep-ph].
 - [37] F. Jegerlehner, *EPJ Web Conf.* **218**, 01003 (2019), arXiv:1711.06089 [hep-ph].
 - [38] F. Jegerlehner, *EPJ Web Conf.* **199**, 01010 (2019), arXiv:1809.07413 [hep-ph].
 - [39] S. Eidelman and F. Jegerlehner, *Z. Phys. C* **67**, 585 (1995), arXiv:hep-ph/9502298.
 - [40] M. Benayoun, P. David, L. DelBuono, O. Leitner, and H. B. O’Connell, *Eur. Phys. J. C* **55**, 199 (2008), arXiv:0711.4482 [hep-ph].
 - [41] M. Benayoun, P. David, L. DelBuono, and F. Jegerlehner, *Eur. Phys. J. C* **72**, 1848 (2012), arXiv:1106.1315 [hep-ph].
 - [42] M. Benayoun, P. David, L. DelBuono, and F. Jegerlehner, *Eur. Phys. J. C* **73**, 2453 (2013), arXiv:1210.7184 [hep-ph].
 - [43] M. Benayoun, P. David, L. DelBuono, and F. Jegerlehner, *Eur. Phys. J. C* **75**, 613 (2015), arXiv:1507.02943 [hep-ph].
 - [44] M. Benayoun, L. Delbuono, and F. Jegerlehner, *Eur. Phys. J. C* **80**, 81 (2020), [Erratum: Eur.Phys.J.C 80,

- 244 (2020)], arXiv:1903.11034 [hep-ph].
- [45] M. Davier, A. Hoecker, B. Malaescu, and Z. Zhang, *Eur. Phys. J. C* **71**, 1515 (2011), [Erratum: Eur.Phys.J.C 72, 1874 (2012)], arXiv:1010.4180 [hep-ph].
- [46] S. Borsanyi et al. (Budapest-Marseille-Wuppertal), *Phys. Rev. Lett.* **121**, 022002 (2018), arXiv:1711.04980 [hep-lat].
- [47] T. Blum, P. A. Boyle, V. Gülpers, T. Izubuchi, L. Jin, C. Jung, A. Jüttner, C. Lehner, A. Portelli, and J. T. Tsang (RBC, UKQCD), *Phys. Rev. Lett.* **121**, 022003 (2018), arXiv:1801.07224 [hep-lat].
- [48] D. Giusti, V. Lubicz, G. Martinelli, F. Sanfilippo, and S. Simula, *Phys. Rev. D* **99**, 114502 (2019), arXiv:1901.10462 [hep-lat].
- [49] C. T. H. Davies et al. (Fermilab Lattice, LATTICE-HPQCD, MILC), *Phys. Rev. D* **101**, 034512 (2020), arXiv:1902.04223 [hep-lat].
- [50] A. Gérardin, M. Cè, G. von Hippel, B. Hörrz, H. B. Meyer, D. Mohler, K. Ottnad, J. Wilhelm, and H. Wittig, *Phys. Rev. D* **100**, 014510 (2019), arXiv:1904.03120 [hep-lat].
- [51] B. Chakraborty et al. (Fermilab Lattice, LATTICE-HPQCD, MILC), *Phys. Rev. Lett.* **120**, 152001 (2018), arXiv:1710.11212 [hep-lat].
- [52] T. Blum, P. A. Boyle, V. Gülpers, T. Izubuchi, L. Jin, C. Jung, A. Jüttner, C. Lehner, A. Portelli, and J. T. Tsang (RBC, UKQCD), *Phys. Rev. Lett.* **121**, 022003 (2018), arXiv:1801.07224 [hep-lat].
- [53] E. Shintani and Y. Kuramashi, *Phys. Rev. D* **100**, 034517 (2019), arXiv:1902.00885 [hep-lat].
- [54] C. Aubin, T. Blum, C. Tu, M. Golterman, C. Jung, and S. Peris, *Phys. Rev. D* **101**, 014503 (2020), arXiv:1905.09307 [hep-lat].
- [55] D. Giusti and S. Simula, PoS **LATTICE2019**, 104 (2019), arXiv:1910.03874 [hep-lat].
- [56] S. Borsanyi et al., *Nature* **593**, 51 (2021), arXiv:2002.12347 [hep-lat].
- [57] A. Boccaletti et al., (2024), arXiv:2407.10913 [hep-lat].
- [58] C. Lehner and A. S. Meyer, *Phys. Rev. D* **101**, 074515 (2020), arXiv:2003.04177 [hep-lat].
- [59] C. Alexandrou et al. (Extended Twisted Mass), *Phys. Rev. D* **107**, 074506 (2023), arXiv:2206.15084 [hep-lat].
- [60] T. Blum et al. (RBC, UKQCD), *Phys. Rev. D* **108**, 054507 (2023), arXiv:2301.08696 [hep-lat].
- [61] S. Kuberski, M. Cè, G. von Hippel, H. B. Meyer, K. Ottnad, A. Risch, and H. Wittig, *JHEP* **03**, 172 (2024), arXiv:2401.11895 [hep-lat].
- [62] C. T. H. Davies et al. (Fermilab Lattice, MILC, HPQCD), *Phys. Rev. D* **106**, 074509 (2022), arXiv:2207.04765 [hep-lat].
- [63] G. Colangelo, A. X. El-Khadra, M. Hoferichter, A. Keshavarzi, C. Lehner, P. Stoffer, and T. Teubner, *Phys. Lett. B* **833**, 137313 (2022), arXiv:2205.12963 [hep-ph].
- [64] M. Passera, W. J. Marciano, and A. Sirlin, *Phys. Rev. D* **78**, 013009 (2008), arXiv:0804.1142 [hep-ph].
- [65] A. Crivellin, M. Hoferichter, C. A. Manzari, and M. Montull, *Phys. Rev. Lett.* **125**, 091801 (2020), arXiv:2003.04886 [hep-ph].
- [66] A. Keshavarzi, W. J. Marciano, M. Passera, and A. Sirlin, *Phys. Rev. D* **102**, 033002 (2020), arXiv:2006.12666 [hep-ph].
- [67] B. Malaescu and M. Schott, *Eur. Phys. J. C* **81**, 46 (2021), arXiv:2008.08107 [hep-ph].
- [68] G. Colangelo, M. Hoferichter, and P. Stoffer, *Phys. Lett. B* **814**, 136073 (2021), arXiv:2010.07943 [hep-ph].
- [69] G. Benton, D. Boito, M. Golterman, A. Keshavarzi, K. Maltman, and S. Peris, *Phys. Rev. Lett.* **131**, 251803 (2023), arXiv:2306.16808 [hep-ph].
- [70] G. Benton, D. Boito, M. Golterman, A. Keshavarzi, K. Maltman, and S. Peris, *Phys. Rev. D* **109**, 036010 (2024), arXiv:2311.09523 [hep-ph].
- [71] L. Di Luzio, A. Masiero, P. Paradisi, and M. Passera, *Phys. Lett. B* **829**, 137037 (2022), arXiv:2112.08312 [hep-ph].
- [72] L. Darmé, G. Grilli di Cortona, and E. Nardi, *JHEP* **06**, 122 (2022), arXiv:2112.09139 [hep-ph].
- [73] A. Crivellin and M. Hoferichter, *Phys. Rev. D* **108**, 013005 (2023), arXiv:2211.12516 [hep-ph].
- [74] L. Darmé, G. Grilli di Cortona, and E. Nardi, *Phys. Rev. D* **108**, 095056 (2023), arXiv:2212.03877 [hep-ph].
- [75] N. M. Coyle and C. E. M. Wagner, *JHEP* **12**, 071 (2023), arXiv:2305.02354 [hep-ph].
- [76] F. V. Ignatov et al. (CMD-3), *Phys. Rev. D* **109**, 112002 (2024), arXiv:2302.08834 [hep-ex].
- [77] F. V. Ignatov et al. (CMD-3), *Phys. Rev. Lett.* **132**, 231903 (2024), arXiv:2309.12910 [hep-ex].
- [78] M. N. Achasov et al., *J. Exp. Theor. Phys.* **103**, 380 (2006), arXiv:hep-ex/0605013.
- [79] V. M. Aul'chenko et al., *JETP Lett.* **84**, 413 (2006), arXiv:hep-ex/0610016.
- [80] R. R. Akhmetshin et al. (CMD-2), *Phys. Lett. B* **648**, 28 (2007), arXiv:hep-ex/0610021.
- [81] F. Ambrosino et al. (KLOE), *Phys. Lett. B* **670**, 285 (2009), arXiv:0809.3950 [hep-ex].
- [82] F. Ambrosino et al. (KLOE), *Phys. Lett. B* **700**, 102 (2011), arXiv:1006.5313 [hep-ex].
- [83] D. Babusci et al. (KLOE), *Phys. Lett. B* **720**, 336 (2013), arXiv:1212.4524 [hep-ex].
- [84] A. Anastasi et al. (KLOE-2), *JHEP* **03**, 173 (2018), arXiv:1711.03085 [hep-ex].
- [85] B. Aubert et al. (BaBar), *Phys. Rev. Lett.* **103**, 231801 (2009), arXiv:0908.3589 [hep-ex].
- [86] J. P. Lees et al. (BaBar), *Phys. Rev. D* **86**, 032013 (2012), arXiv:1205.2228 [hep-ex].
- [87] M. Ablikim et al. (BESIII), *Phys. Lett. B* **753**, 629 (2016), [Erratum: Phys.Lett.B 812, 135982 (2021)], arXiv:1507.08188 [hep-ex].
- [88] T. Xiao, S. Dobbs, A. Tomaradze, K. K. Seth, and G. Bonvicini, *Phys. Rev. D* **97**, 032012 (2018), arXiv:1712.04530 [hep-ex].
- [89] M. N. Achasov et al. (SND), *JHEP* **01**, 113 (2021), arXiv:2004.00263 [hep-ex].
- [90] Zhiqing ZHANG, “News from BABAR,” Muon $g-2$ Theory Initiative Spring 2024 Meeting, URL: https://indico.cern.ch/event/1400808/contributions/5902094/attachments/2842140/4968393/Zhang_HVP240422.pdf (2024).
- [91] Hisaki Hayashii, “Status and plans regarding $g-2$ at belle II,” Muon $g-2$ Theory Initiative Spring 2024 Meeting, URL: https://indico.cern.ch/event/1400808/contributions/5902135/attachments/2841580/4968265/g-2-theory_mni-workshop_240422_v2.pdf (2024).
- [92] Riccardo Aliberti, “Status and Plans for Experimental Inputs to HVP at BESIII,” Muon $g-2$ Theory Initiative Spring 2024 Meeting, URL: <https://indico.cern.ch/event/1400808/>

- [contributions/5902095/attachments/2841881/4967869/Muon_g-2_spring_meeting_2024.pdf](https://indico.cern.ch/event/1400808/contributions/5902095/attachments/2841881/4967869/Muon_g-2_spring_meeting_2024.pdf) (2024).
- [93] Ivan Logashenko, “CMD2/3 report (on $\pi^+\pi^-$),” Muon $g-2$ Theory Initiative Spring 2024 Meeting, URL: https://indico.cern.ch/event/1400808/contributions/5902096/attachments/2842193/4968485/Logashenko_CMD_TI_2024.pdf (2024).
- [94] Giuseppe Mandaglio, “Hadron physics results at KLOE-2,” Sixth Plenary Workshop of the Muon $g-2$ Theory Initiative (2023), URL: https://indico.cern.ch/event/1400808/contributions/5902134/attachments/2841711/4967471/talk_IL.pdf (2023).
- [95] Andrey Kupich, “Preliminary results of the $e^+e^- \rightarrow \pi^+\pi^-$ analysis with SND at VEPP-2000,” Muon $g-2$ Theory Initiative Spring 2024 Meeting, URL: https://indico.cern.ch/event/1400808/contributions/5902134/attachments/2841711/4967471/talk_IL.pdf (2024).
- [96] J. P. Lees et al. (BaBar), *Phys. Rev. D* **108**, L111103 (2023), arXiv:2308.05233 [hep-ex].
- [97] C. M. Carloni Calame, M. Passera, L. Trentadue, and G. Venanzoni, *Phys. Lett. B* **746**, 325 (2015), arXiv:1504.02228 [hep-ph].
- [98] G. Abbiendi et al., *Eur. Phys. J. C* **77**, 139 (2017), arXiv:1609.08987 [hep-ex].
- [99] G. Abbiendi, Letter of Intent: the MUonE project, Tech. Rep. (CERN, Geneva, 2019).
- [100] P. Banerjee et al., *Eur. Phys. J. C* **80**, 591 (2020), arXiv:2004.13663 [hep-ph].
- [101] A. Masiero, P. Paradisi, and M. Passera, *Phys. Rev. D* **102**, 075013 (2020), arXiv:2002.05418 [hep-ph].
- [102] R. H. Parker, C. Yu, W. Zhong, B. Estey, and H. Müller, *Science* **360**, 191 (2018), arXiv:1812.04130 [physics.atom-ph].
- [103] L. Morel, Z. Yao, P. Cladé, and S. Guellati-Khélifa, *Nature* **588**, 61 (2020).
- [104] G. F. Giudice, P. Paradisi, and M. Passera, *JHEP* **11**, 113 (2012), arXiv:1208.6583 [hep-ph].
- [105] A. Crivellin, M. Hoferichter, and P. Schmidt-Wellenburg, *Phys. Rev. D* **98**, 113002 (2018), arXiv:1807.11484 [hep-ph].
- [106] X. Fan, T. G. Myers, B. A. D. Sukra, and G. Gabrielse, *Phys. Rev. Lett.* **130**, 071801 (2023), arXiv:2209.13084 [physics.atom-ph].
- [107] T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, *Phys. Rev. D* **91**, 033006 (2015), [Erratum: *Phys. Rev. D* 96, 019901 (2017)], arXiv:1412.8284 [hep-ph].
- [108] S. Volkov, *Phys. Rev. D* **100**, 096004 (2019), arXiv:1909.08015 [hep-ph].
- [109] S. Volkov, (2024), arXiv:2404.00649 [hep-ph].
- [110] J. Bernabeu, G. A. Gonzalez-Sprinberg, J. Papavassiliou, and J. Vidal, *Nucl. Phys. B* **790**, 160 (2008), arXiv:0707.2496 [hep-ph].
- [111] P. Janot, *JHEP* **02**, 053 (2016), [Erratum: *JHEP* 11, 164 (2017)], arXiv:1512.05544 [hep-ph].
- [112] A. Blondel and P. Janot, *Eur. Phys. J. Plus* **137**, 92 (2022), arXiv:2106.13885 [hep-ex].
- [113] M. Tanabashi et al. (Particle Data Group), *Phys. Rev. D* **98**, 030001 (2018).
- [114] P. Gambino and A. Sirlin, *Phys. Rev. D* **49**, 1160 (1994), arXiv:hep-ph/9309326.
- [115] J. Erler and R. Ferro-Hernández, *JHEP* **03**, 196 (2018), arXiv:1712.09146 [hep-ph].
- [116] D. Becker et al., *Eur. Phys. J. A* **54**, 208 (2018), arXiv:1802.04759 [nucl-ex].
- [117] J. Benesch et al. (MOLLER), (2014), arXiv:1411.4088 [nucl-ex].
- [118] E. Tiesinga, P. J. Mohr, D. B. Newell, and B. N. Taylor, *Rev. Mod. Phys.* **93**, 025010 (2021).
- [119] M. I. Eides, *Phys. Lett. B* **795**, 113 (2019), arXiv:1812.10881 [hep-ph].
- [120] A. Czarnecki, S. I. Eidelman, and S. G. Karshenboim, *Phys. Rev. D* **65**, 053004 (2002), arXiv:hep-ph/0107327.
- [121] S. G. Karshenboim and V. A. Shelyuto, *Phys. Lett. B* **517**, 32 (2001), arXiv:hep-ph/0107328.
- [122] C. Delaunay, B. Ohayon, and Y. Soreq, *Phys. Rev. Lett.* **127**, 251801 (2021), arXiv:2106.11998 [hep-ph].
- [123] F. G. Mariam et al., *Phys. Rev. Lett.* **49**, 993 (1982).
- [124] W. Liu et al., *Phys. Rev. Lett.* **82**, 711 (1999).
- [125] P. Strasser et al., *Hyperfine Interact.* **237**, 124 (2016).
- [126] J. L. Friar, J. Martorell, and D. W. L. Sprung, *Phys. Rev. A* **59**, 4061 (1999), arXiv:nucl-th/9812053.
- [127] V. Meyer et al., *Phys. Rev. Lett.* **84**, 1136 (2000), arXiv:hep-ex/9907013.
- [128] P. Crivelli, *Hyperfine Interact.* **239**, 49 (2018), arXiv:1811.00310 [physics.atom-ph].
- [129] M. I. Eides and V. A. Shelyuto, *Int. J. Mod. Phys. A* **31**, 1645034 (2016).