On the Burde–de Rham theorem for finitely presented pro-p groups

Yasushi MIZUSAWA, Ryoto TANGE, and Yuji TERASHIMA

Abstract. We consider the Burde–de Rham theorem for finitely presented pro-p groups under the assumption that the total degrees of all relators are 0. We also give some concrete examples including higher-dimensional cases under Iwasawa theoretic conditions, and consider some cohomological interpretations.

0. Introduction

In this paper, we study linear representations of finitely presented pro-p groups from the knot theoretic viewpoint. In knot theory, Burde and de Rham independently discovered the constructions of linear representations of knot groups using the zero of Alexander polynomials ([Bur67; deR67; Kit13]). Generalizations to higher-dimensional linear representations are studied ([SW11]), and are still actively studied from cohomological viewpoints ([HPS01; HP15]).

From the viewpoint of arithmetic topology [Mor24], we introduce arithmetic twisted Alexander polynomials and show the arithmetic Burde–de Rham theorem which describes a relation between zeros of the arithmetic twisted Alexander polynomials and extensions of representations for pro-p Galois groups of number fields.

The analogy between the Alexander polynomial and the algebraic p-adic L-function was pointed out by Mazur ([Maz64]), and based on this analogy, various studies have developed. One of the differences between knot groups and finitely presented pro-p Galois groups of number fields is the deficiency of group presentations, namely the difference in the number of generators and relators of group presentations. The knot groups can always transform into having the deficiency one (called the Wirtinger presentation), whereas the pro-p Galois groups cannot. In this paper, we consider the case of group presentations for any deficiency.

Another concern is the number theoretic counterpart of the Alexander polynomial for higher-dimensional cases. Based on the ideas of twisted Alexander invariant in knot theory ([Lin01; Wad94]), we consider the counterpart using Fitting ideals. We also give some cohomological viewpoints, which we expect the relations with infinitesimal deformations of representations ([Maz89]). We believe that this study will enrich the field of number theory (e.g., Iwasawa theory, zeros of p-adic L-functions, etc.), and strengthen the bridge between number theory and knot theory (e.g., multiple residue symbols, Massey products, etc. ([AC22; Mor04; Sha07a; Sha07b])).

This paper is organized as follows. In Section 1, we recall the complete differential modules and introduce some properties that we use later. In Section 2, we prove the Burde–de Rham theorem for finitely presented pro-p groups including higher-dimensional cases, and consider the existence of non-standard extensions of homomorphisms under the assumption that the total degrees of all relators are 0. In Section 3, we give some explicit examples of extensions of homomorphisms for Galois groups under Iwasawa theoretic settings. Lastly, in Section 4, we recall the group cohomology of local systems and study some cohomological interpretations of extensions of homomorphisms, and consider generalizations of the relation with zeros of algebraic p-adic L-functions. Section 4 is seen as a sequel of the works by Morishita ([Mor06], [Mor24, Chapter 14]), where the knot theoretic part is mainly due to Hironaka ([Hir97]) and Lê ([Le94]).

Notation. For a pro-p group G, we denote by $[h,g] = h^{-1}g^{-1}hg$ the commutator of g, $h \in G$. We denote by $|\cdot|_p$ the p-adic absolute value normalized by $|p|_p = p^{-1}$. For a closed subgroup H of G, [H,G] denotes the minimal closed subgroup containing $\{[h,g] \mid h \in H, g \in G\}$. For a, b in a commutative ring R, $a \doteq b$ means a = bu for some unit $u \in R^{\times}$.

1. Complete differential modules

Let G and H be profinite groups, and let $\psi \colon G \to H$ be a continuous homomorphism. Let p be a prime number. For simplicity, we also denote by the same ψ the algebra homomorphism $\mathbb{Z}_p[[G]] \to \mathbb{Z}_p[[H]]$ of the complete group algebras over the p-adic integer \mathbb{Z}_p induced by ψ .

The complete ψ -differential module A_{ψ} is defined as the quotient module of the left free $\mathbb{Z}_p[[H]]$ -module $\bigoplus_{g \in G} \mathbb{Z}_p[[H]] dg$ on the symbols dg $(g \in G)$ by the left $\mathbb{Z}_p[[H]]$ -submodule generated by elements of the form $d(xy) - dx - \psi(x)dy$ for any $x, y \in G$:

$$A_{\psi} := (\bigoplus_{g \in G} \mathbb{Z}_p[[H]] dg) / \langle d(xy) - dx - \psi(x) dy \ (x, y \in G) \rangle_{\mathbb{Z}_p[[H]]}.$$

The map $d: G \to A_{\psi}$ defined by $g \mapsto dg$ is a ψ -differential, namely, we have

$$d(xy) = dx + \psi(x)dy$$

for any $x, y \in G$, and the following universal property holds:

For any left $\mathbb{Z}_p[[H]]$ -module M, and for any ψ -differential $\partial \colon G \to M$, there exists a unique $\mathbb{Z}_p[[H]]$ -homomorphism $f \colon A_{\psi} \to M$ such that $f \circ d = \partial$.

Next, suppose that G is a finitely presented pro-p group, and choose a presentation

$$(1.1) G = \langle g_1, \dots, g_n \mid r_1 = \dots = r_m = 1 \rangle^{\operatorname{pro}-p}.$$

Let F_n be the free pro-p group on g_1, \ldots, g_n , and let $\pi \colon F_n \to G$ be the natural homomorphism. We also denote by the same g_i for the image of g_i in G, and the same π for the algebra homomorphism $\mathbb{Z}_p[[F_n]] \to \mathbb{Z}_p[[G]]$ of complete group algebras induced by π . Then we can describe the presentation matrix of the complete ψ -differential module A_{ψ} using the pro-p Fox free differential calculus $\frac{\partial}{\partial g_i} \colon \mathbb{Z}_p[[F_n]] \to \mathbb{Z}_p[[F_n]]$ of complete group algebras ([Iha86, Section 2], [Mor02, Section 2], [Mor24, Section 10.3], [NSW20, Section V.6]), as follows:

Proposition ([Mor24, Corollary 10.3.6]). The complete ψ -differential module A_{ψ} has a free resolution over $\mathbb{Z}_p[[H]]$:

$$\mathbb{Z}_p[[H]]^m \xrightarrow{Q_{\psi \circ \pi}} \mathbb{Z}_p[[H]]^n \longrightarrow A_{\psi} \to 0$$

whose presentation matrix $Q_{\psi \circ \pi} \in \mathcal{M}(m, n; \mathbb{Z}_p[[H]])$ is given by

$$Q_{\psi \circ \pi} := \left((\psi \circ \pi) \left(\frac{\partial r_j}{\partial g_i} \right) \right) \in \mathcal{M}(m, n; \mathbb{Z}_p[[H]]).$$

For later use, we remark on some basic rules for Fox derivatives ([Iha86, Section 2.1]):

1.
$$\frac{\partial g_j}{\partial g_i} = \begin{cases} 1 & (i=j) \\ 0 & (i \neq j) \end{cases};$$

2.
$$\frac{\partial(\eta+\xi)}{\partial g_i} = \frac{\partial\eta}{\partial g_i} + \frac{\partial\xi}{\partial g_i}, \quad (\eta, \xi \in \mathbb{Z}_p[[F_n]]);$$

3.
$$\frac{\partial(\eta\xi)}{\partial g_i} = \frac{\partial\eta}{\partial g_i} \cdot \epsilon_{\mathbb{Z}_p[[F_n]]}(\xi) + \eta \cdot \frac{\partial\xi}{\partial g_i} \quad (\eta, \ \xi \in \mathbb{Z}_p[[F_n]]),$$
where $\epsilon_{\mathbb{Z}_p[[F_n]]} \colon \mathbb{Z}_p[[F_n]] \to \mathbb{Z}_p$ is the augmentation map;

4.
$$\frac{\partial(w^{-1})}{\partial g_i} = -w^{-1} \cdot \frac{\partial w}{\partial g_i} \quad (w \in F_n).$$

Now, for a positive integer $\ell \in \mathbb{Z}_{\geq 1}$, and for a given homomorphism $\varphi \colon F_n \to \mathrm{GL}(\ell; \overline{\mathbb{Q}_p})$, consider a tensor product homomorphism $\widetilde{\varphi} := \varphi \otimes (\psi \circ \pi) \colon F_n \to \mathrm{GL}(\ell; \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[H]])$ defined by

$$(1.2) \widetilde{\varphi}(x) := (\varphi \otimes (\psi \circ \pi))(x) := \varphi(x) \cdot (\psi \circ \pi)(x) = \operatorname{GL}(\ell; \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[H]])$$

for any $x \in F_n$. Consider the $m\ell \times n\ell$ matrix

$$Q_{\widetilde{\varphi}} = \left(\widetilde{\varphi}\left(\frac{\partial r_j}{\partial g_i}\right)\right) \in \mathcal{M}(m\ell, n\ell; \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[H]]),$$

as in Proposition. When $\mathbb{Z}_p[[H]]$ is a Noetherian UFD, for a non-negative integer d with $d < n\ell$, we define the d-th Fitting ideal $E_d(A_{\widetilde{\varphi}}) \in \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[H]]$ of $A_{\widetilde{\varphi}}$ as the ideal generated by the $(n\ell-d)$ -minors of $Q_{\widetilde{\varphi}}$. If $n\ell-d > m\ell$, we let $E_d(A_{\widetilde{\varphi}}) := 0$. These ideals depend only on $A_{\widetilde{\varphi}}$ and are independent of the choice of a presentation ([Nor76, Section 3.1, Theorem 1]). Since the GCD of the generators of $E_d(A_{\widetilde{\varphi}})$ is well-defined up to multiplication by a unit of $\overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[H]]$, we denote the GCD of generators of $E_d(A_{\widetilde{\varphi}})$ by $\Delta_d(A_{\widetilde{\varphi}}) \in \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[H]]$. If $\mathrm{Im} \varphi \subset \mathrm{GL}(\ell, \mathbb{Z}_p)$, then $E_d(A_{\widetilde{\varphi}})$ can be defined as an ideal of $\mathbb{Z}_p[[H]]$, and so $\Delta_d(A_{\widetilde{\varphi}})$ is the GCD in $\mathbb{Z}_p[[H]]$.

2. Burde-de Rham Theorem for finitely presented pro-p groups

Denote $\Gamma := \mathbb{Z}_p$, and suppose that we can take a surjective homomorphism $\alpha \colon G \to \Gamma$ such that $\alpha(g_i) = \gamma$ for all i, where γ is a topological generator of Γ .

For an integer $k \in \mathbb{Z}_{\geq 2}$, and for a given homomorphism $\varphi_{k-2} \colon F_n \to \operatorname{GL}(k-1; \overline{\mathbb{Q}_p})$, consider a tensor product homomorphism $\widetilde{\varphi_{k-2}} := \varphi_{k-2} \otimes (\alpha \circ \pi) \colon F_n \to \operatorname{GL}(k-1; \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]])$ as (1.2). For a given $a_{k-1} \in \overline{\mathbb{Q}_p}^{\times}$ such that $|a_{k-1} - 1|_p < 1$, let $\overline{\beta_{k-1}} \colon F_n \to \overline{\mathbb{Q}_p}^{\oplus (k-1)}$ be a crossed homomorphism with respect to $\widetilde{\varphi_{k-2}}|_{\gamma=a_{k-1}} \colon F_n \to \operatorname{GL}(k-1; \overline{\mathbb{Q}_p})$, namely satisfying the relation

$$\overline{\beta_{k-1}}(xy) = \overline{\beta_{k-1}}(x) + \widetilde{\varphi_{k-2}}(x)|_{\gamma = a_{k-1}} \overline{\beta_{k-1}}(y) = \overline{\beta_{k-1}}(x) + a_{k-1}^{(\alpha_{\gamma} \circ \pi)(x)} \cdot \varphi_{k-2}(x) \cdot \overline{\beta_{k-1}}(y)$$

for any $x, y \in F_n$, where $\alpha_{\gamma} \colon G \to \mathbb{Z}_p$ maps $x \in G$ to the power of γ of $\alpha(x)$. Note that by the definition of the complete differential module, the crossed homomorphism $\overline{\beta_{k-1}} \colon F_n \to \overline{\mathbb{Q}_p}^{\oplus (k-1)}$ can be regarded as a $\overline{\mathbb{Q}_p}$ -module homomorphism from $A_{\widetilde{\varphi_{k-2}}}$ to $\overline{\mathbb{Q}_p}$. Namely, we have an isomorphism between the set of a crossed homomorphism with respect to $\widehat{\varphi_{k-2}}|_{\gamma=a_{k-1}}$ and the set $\operatorname{Hom}_{\overline{\mathbb{Q}_p}}(A_{\widetilde{\varphi_{k-2}}}, \overline{\mathbb{Q}_p}^{\oplus (k-1)})$ as vector spaces over $\overline{\mathbb{Q}_p}$. Using the crossed homomorphism $\overline{\beta_{k-1}} \colon F_n \to \overline{\mathbb{Q}_p}^{\oplus (k-1)}$, we define a homomorphism $\varphi_{k-1} \colon F_n \to \operatorname{GL}(k; \overline{\mathbb{Q}_p})$ as

$$\varphi_{k-1}(g_1) = \begin{pmatrix} \widetilde{\varphi_{k-2}}(g_1)|_{\gamma = a_{k-1}} & \overline{\beta_{k-1}}(g_1) \\ 0 & 1 \end{pmatrix}, \cdots, \varphi_{k-1}(g_n) = \begin{pmatrix} \widetilde{\varphi_{k-2}}(g_n)|_{\gamma = a_{k-1}} & \overline{\beta_{k-1}}(g_n) \\ 0 & 1 \end{pmatrix}.$$

We focus on when $\varphi_{k-1} \colon F_n \to \operatorname{GL}(k; \overline{\mathbb{Q}_p})$ can be extended to G as a representation $\rho_{k-1} \colon G \to \operatorname{GL}(k; \overline{\mathbb{Q}_p})$, where $\rho_{k-1} \circ \pi = \varphi_{k-1}$. We say such a representation $\rho_{k-1} \colon G \to \operatorname{GL}(k; \overline{\mathbb{Q}_p})$ an extension of a homomorphism $\varphi_{k-1} \colon F_n \to \operatorname{GL}(k; \overline{\mathbb{Q}_p})$.

We have the following main results, which is an analogue of the Burde-de Rham theorem including higher-dimensional cases ([Kit13; HPS01; HP15; SW11]):

Theorem 1. Let G be a finitely presented pro-p group with the total degrees of all relators being 0. Let $\varphi_{k-2} \colon F_n \to \mathrm{GL}(k-1; \overline{\mathbb{Q}_p})$ be a homomorphism that can be extended to G. Then the homomorphism defined by

$$\varphi_{k-1} \colon F_n \ni g_i \mapsto \begin{pmatrix} \widetilde{\varphi_{k-2}}(g_i)|_{\gamma = a_{k-1}} & \overline{\beta_{k-1}}(g_i) \\ \mathbf{0} & 1 \end{pmatrix} \in \mathrm{GL}(k; \overline{\mathbb{Q}_p})$$

for all i, can be extended to G as a representation $\rho_{k-1}: G \to \operatorname{GL}(k; \overline{\mathbb{Q}_p})$ if and only if $Q_{\widetilde{\varphi_{k-2}}}|_{\gamma=a_{k-1}} \cdot \mathbf{b} = 0$, where $\mathbf{b} = {}^t(\overline{\beta_{k-1}}(g_1), \dots, \overline{\beta_{k-1}}(g_n)) \in \overline{\mathbb{Q}_p}^{\oplus n(k-1)}$.

Moreover, we have $\dim_{\overline{\mathbb{Q}_p}^{\oplus (k-1)}} \operatorname{Hom}_{\overline{\mathbb{Q}_p}}(A_{\widetilde{\varphi_{k-2}}}, \overline{\mathbb{Q}_p}^{\oplus (k-1)}) \geq k$, namely, we have at least k

Moreover, we have $\dim_{\overline{\mathbb{Q}_p}^{\oplus(k-1)}} \operatorname{Hom}_{\overline{\mathbb{Q}_p}}(A_{\widetilde{\varphi_{k-2}}}, \overline{\mathbb{Q}_p}^{\oplus(k-1)}) \geq k$, namely, we have at least k extensions $\rho_{k-1} \colon G \to \operatorname{GL}(k; \overline{\mathbb{Q}_p})$ of a homomorphism $\varphi_{k-1} \colon F_n \to \operatorname{GL}(k; \overline{\mathbb{Q}_p})$ if and only if $a_{k-1} \in \overline{\mathbb{Q}_p}^{\times}$ is a zero of $\Delta_{k-1}(A_{\widetilde{\varphi_{k-2}}}) \in \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]]$.

Proof. The proof is based on [HPS01, Section 4]. We use the following Lemma:

Lemma. For $w = w(g_1, \ldots, g_n) \in F_n$, we have

$$\varphi_{k-1}(w) = \begin{pmatrix} \widetilde{\varphi_{k-2}}(w)|_{\gamma = a_{k-1}} & \sum_{i=1}^{n} \left(\widetilde{\varphi_{k-2}} \left(\frac{\partial w}{\partial g_i} \right) \right) \Big|_{\gamma = a_{k-1}} \cdot \overline{\beta_{k-1}}(g_i) \end{pmatrix} \in GL(k; \overline{\mathbb{Q}_p}).$$

Proof. Due to [Iha86, Section 2], since the classical Fox derivative $\frac{\partial}{\partial g_i}$: $\mathbb{Z}[F_n] \to \mathbb{Z}[F_n]$ of group algebras is continuous with respect to the topology of $\mathbb{Z}[F_n]$, which is induced from the topology of $\mathbb{Z}_p[[F_n]]$ via the canonical injection $\mathbb{Z}[F_n] \to \mathbb{Z}_p[[F_n]]$, it is enough to prove the classical case.

We denote by the same π for the algebra homomorphism $\mathbb{Z}[F_n] \to \mathbb{Z}[G]$ of group algebras induced by $\pi \colon F_n \to G$, and the same α for the algebra homomorphism $\mathbb{Z}[G] \to \mathbb{Z}[\Gamma]$ of group algebras induced by $\alpha \colon G \to \Gamma$. For simplicity, denote $\overline{\eta} := (\alpha \circ \pi)(\eta) \in \mathbb{Z}[[\Gamma]]$, and $\partial_i \eta := \frac{\partial \eta}{\partial g_i} \in \mathbb{Z}[[F_n]]$ for $\eta \in \mathbb{Z}[[F_n]]$. We prove by induction on the dimension k and the length l of the word w.

For k=2 and for l=1, since

$$\partial_i(g_h) = \begin{cases} 1 & (i=h) \\ 0 & (i \neq h) \end{cases}, \ \partial_i(g_h^{-1}) = \begin{cases} -g_h^{-1} & (i=h) \\ 0 & (i \neq h) \end{cases},$$

we have

$$\varphi_1(g_h) = \begin{pmatrix} a & \overline{\beta_1}(g_h) \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \overline{g_h}|_{\gamma=a} & \sum_{i=1}^n (\overline{\partial_i g_h})|_{\gamma=a} \cdot \overline{\beta_1}(g_i) \\ 0 & 1 \end{pmatrix},$$

$$\varphi_1(g_h^{-1}) = \begin{pmatrix} a^{-1} & -a^{-1} \cdot \overline{\beta_1}(g_h) \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \overline{g_h^{-1}}|_{\gamma=a} & \sum_{i=1}^n (\overline{\partial_i g_h^{-1}})|_{\gamma=a} \cdot \overline{\beta_1}(g_i) \\ 0 & 1 \end{pmatrix},$$

and so the claim holds.

Next, assume that the claim holds for k and l-1. Namely, for any $w_{l-1} \in F_n$ with the length l-1, assume we have

$$\varphi_{k-1}(w_{l-1}) = \begin{pmatrix} \widetilde{\varphi_{k-2}}(w_{l-1})|_{\gamma = a_{k-1}} & \sum_{i=1}^{n} \widetilde{\varphi_{k-2}} \left(\partial_{i} w_{l-1} \right)|_{\gamma = a_{k-1}} \cdot \overline{\beta_{k-1}}(g_{i}) \\ \mathbf{0} & 1 \end{pmatrix} \in GL(k; \overline{\mathbb{Q}_{p}}).$$

Then for the case of k and l, since

$$\partial_i(w_{l-1}g_h) = \begin{cases} \partial_i w_{l-1} + w_{l-1} & (i=h) \\ \partial_i w_{l-1} & (i \neq h) \end{cases}, \ \partial_i(w_{l-1}g_h^{-1}) = \begin{cases} \partial_i w_{l-1} - w_{l-1}g_h^{-1} & (i=h) \\ \partial_i w_{l-1} & (i \neq h) \end{cases},$$

we have

$$\varphi_{k-1}(w_{l-1}g_h) = \varphi_{k-1}(w_{l-1}) \cdot \varphi_{k-1}(g_h)$$

$$= \begin{pmatrix} \widetilde{\varphi_{k-2}}(w_{l-1}g_h)|_{\gamma=a_{k-1}} & \sum_{i=1}^n \widetilde{\varphi_{k-2}}(\partial_i w_{l-1})|_{\gamma=a_{k-1}} \cdot \overline{\beta_{k-1}}(g_i) - \widetilde{\varphi_{k-2}}(w_{l-1})|_{\gamma=a_{k-1}} \cdot \overline{\beta_{k-1}}(g_h) \\ \mathbf{0} & 1 \end{pmatrix}$$

$$= \begin{pmatrix} \widetilde{\varphi_{k-2}}(w_{l-1}g_h)|_{\gamma=a_{k-1}} & \sum_{i=1}^n \widetilde{\varphi_{k-2}} \left(\partial_i (w_{l-1}g_h) \right)|_{\gamma=a_{k-1}} \cdot \overline{\beta_{k-1}}(g_i) \\ \mathbf{0} & 1 \end{pmatrix} \in GL(k; \overline{\mathbb{Q}_p}),$$

$$\varphi_{k-1}(w_{l-1}g_h^{-1}) = \varphi_{k-1}(w_{l-1}) \cdot \varphi_{k-1}(g_h)^{-1}$$

$$= \begin{pmatrix} \widetilde{\varphi_{k-2}}(w_{l-1}g_h^{-1})|_{\gamma=a_{k-1}} & \sum_{i=1}^n \widetilde{\varphi_{k-2}}(\partial_i w_{l-1})|_{\gamma=a_{k-1}} \cdot \overline{\beta_{k-1}}(g_i) - \widetilde{\varphi_{k-2}}(w_{l-1})|_{\gamma=a_{k-1}} \cdot a^{-1} \cdot \overline{\beta_{k-1}}(g_h) \\ \mathbf{0} & 1 \end{pmatrix}$$

$$= \begin{pmatrix} \widetilde{\varphi_{k-2}}(w_{l-1}g_h^{-1})|_{\gamma=a_{k-1}} & \sum_{i=1}^n \widetilde{\varphi_{k-2}}\left(\partial_i (w_{l-1}g_h^{-1})\right)|_{\gamma=a_{k-1}} \cdot \overline{\beta_{k-1}}(g_i) \\ \mathbf{0} & 1 \end{pmatrix} \in \mathrm{GL}(k; \overline{\mathbb{Q}_p}),$$

and so the claim holds.

Lastly, for the case of k+1 and l-1, the claim holds using a calculation similar to the one above.

By the assumption that the total degree $deg(r_j)$ of a relator r_j of G is 0 for all j, we have

$$\widetilde{\varphi_{k-2}}(r_j)|_{\gamma=a_{k-1}} = a_{k-1}^{\deg(r_j)} \cdot \varphi_{k-2}(r_j) = \varphi_{k-2}(r_j).$$

Hence, by Lemma, the homomorphism $\varphi_{k-1} \colon F_n \to \operatorname{GL}(k; \overline{\mathbb{Q}_p})$ can be extended to G as a representation $\rho_{k-1} \colon G \to \operatorname{GL}(k; \overline{\mathbb{Q}_p})$ if and only if

$$\sum_{i=1}^{n} \left(\widetilde{\varphi_{k-2}} \left(\frac{\partial r_j}{\partial g_i} \right) \right) \Big|_{\gamma = a_{k-1}} \cdot \overline{\beta_{k-1}}(g_i) = 0$$

for all $j=1,\ldots,m$. This means that the following linear system has a non-zero solution $\mathbf{b}_{k-1} \in \overline{\mathbb{Q}_p}^{\oplus n(k-1)}$:

$$(2.1) Q_{\widetilde{\varphi_{k-2}}}|_{\gamma=a_{k-1}} \cdot \mathbf{b}_{k-1} = 0.$$

In order to have at least k extensions of ρ_{k-1} , we need to have at least k solutions of (2.1), which are linearly independent. This condition is equivalent to

$$\operatorname{rank}(Q_{\widetilde{\varphi_{k-2}}}|_{\gamma = a_{k-1}}) \le n(k-1) - k < n(k-1) - (k-1),$$

which also means that $\Delta_{k-1}(A_{\widetilde{\varphi_{k-2}}}) = 0$, and $a_{k-1} \in \overline{\mathbb{Q}_p}^{\times}$ is a zero of $\Delta_{k-1}(A_{\widetilde{\varphi_{k-2}}}) \in \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]]$.

Remark 1. In the spirit of arithmetic topology, we call $\Delta_{k-1}(A_{\widetilde{\varphi_{k-2}}}) \in \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]]$ the arithmetic twisted Alexander polynomial associated with ρ_{k-2} which may be regarded as a knottheoretic analogue of the twisted Alexander invariant associated with a representation of a knot group ([Lin01; Wad94]) or the *L*-function associated with the universal representation of a knot group ([KMTT18; KMTT22; MTTU17]). These knot invariants would be interesting to compare in our future study.

It may be interesting to note that the knot theoretic analogue of Theorem 1 for higher-dimensional cases are proven by Heusener and Porti [HP15, Theorem 1.4]. Their result gives a partially positive answer to the rigidity questions of Mazur ([Maz00, p.452 Proposition, p.454 Question 3]). Their idea of using the non-vanishing of the cup products is reminiscent of Sharifi's works ([Sha07a; Sha07b]), and would also be of interest to us in our future study.

3. Examples

Let us consider some examples under Iwasawa theoretic conditions. Let K_{∞}/K a \mathbb{Z}_p -extension, and Σ the set of primes of K ramifying in K_{∞}/K . Let S be a finite set consisting of places of K, and let $(K_{\infty})_S$ (resp. K_S) be the maximal pro-p-extension of K_{∞} (resp. K) unramified outside S. Put $G_S(K) = \operatorname{Gal}(K_S/K)$ and $\widetilde{G}_S(K) = \operatorname{Gal}((K_{\infty})_S/K)$. It is known ([NSW20, Theorem 10.7.12], [Sha63] (resp. [EM22; BLM13; Miz19; Sal08])) that the Galois group $G_S(K)$ (resp. $\widetilde{G}_S(K)$) is finitely presented as a pro-p group. Moreover, if $\Sigma \subset S$, then we have $(K_{\infty})_S = K_S$ and $\widetilde{G}_S(K) = G_S(K)$. Hence, we focus on the Galois $\widetilde{G}_S(K)$ having the same presentation as (1.1):

$$\widetilde{G}_S(K) = \langle g_1, \dots, g_n \mid r_1 = \dots = r_m = 1 \rangle^{\operatorname{pro}-p}.$$

The total degrees $\deg(r_j)$ of all relators r_j of $\widetilde{G}_S(K)$ are always 0 since the ramified pro-p extension $(K_\infty)_S/K$ contains the \mathbb{Z}_p -extension K_∞/K . Since S contains the set of primes of K over p, we can take the profinite group H in Section 1 to be the Galois group $\operatorname{Gal}(K_\infty/K) \simeq \mathbb{Z}_p = \Gamma$. Moreover, the complete ψ -differential module A_ψ becomes a module over $\Lambda := \mathbb{Z}_p[[\Gamma]] \simeq \mathbb{Z}_p[[T]]$ by the pro-p Magnus isomorphism ([Mor24, Lemma 9.3.1]). Since Γ is an abelian group, and Λ is a Noetherian UFD, the GCD $\Delta_d(A_\psi)$ of the d-th Fitting ideal $(d \geq 0)$ of A_ψ is defined, and we can regard $\Delta_d(A_\psi)$ as an element in $\overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]]$.

Denote $N := \operatorname{Ker}(\psi : \widetilde{G}_S(K) \to \Gamma)$, so that we have a short exact sequence of pro-p groups

$$1 \longrightarrow N \longrightarrow \widetilde{G}_S(K) \stackrel{\psi}{\longrightarrow} \Gamma \longrightarrow 1.$$

Then we have the exact sequence of left Λ -modules

$$(3.1) 0 \longrightarrow N^{\mathrm{ab}} \xrightarrow{\theta_1} A_{\psi} \xrightarrow{\theta_2} \mathbb{Z}_p[[\Gamma]] \xrightarrow{\epsilon_{\mathbb{Z}_p[[H]]}} \mathbb{Z}_p \longrightarrow 0,$$

where $N^{ab} = H_1(N, \mathbb{Z}_p)$ is the maximal pro-p quotient of the abelianization N/[N, N] of N, θ_1 is the homomorphism induced by $n \mapsto dn$ for any $n \in N$, and θ_2 is the homomorphism induced by $dg \mapsto \psi(g) - 1$ for any $g \in \widetilde{G}_S(K)$ ([Mor24, Section 10.4]). We call the left Λ -module N^{ab} the Iwasawa module. By (3.1), and since the kernel $\operatorname{Ker}(\epsilon_{\Lambda} \colon \Lambda \to \mathbb{Z}_p)$ of the augmentation map $\epsilon_{\Lambda} \colon \Lambda \to \mathbb{Z}_p$ is Λ -isomorphic to Λ , we have a Λ -isomorphism $A_{\psi} \simeq N^{ab} \oplus \Lambda$. Therefore, we have $\Delta_d(N^{ab}) = \Delta_{d+1}(A_{\psi})$ for any $d \geq 0$.

Here are some examples of extensions of homomorphism for finitely presented pro-p Galois groups $G_S(K)$ over \mathbb{Z}_p -extensions. For simplicity, we consider the case where φ_0 is the constant map to 1, so that we have $\widetilde{\varphi_0} = \alpha \circ \pi$. We used SageMath [S24] for calculations of group presentations and Fox derivatives, and Mathematica [W24] for calculations of linear algebra.

Example 1. Let p be an odd prime number, and $S = \{q_1, q_2\}$ a set of prime numbers $q_1 \equiv q_2 \equiv 1 \pmod{p}$, $q_1 \not\equiv 1 \pmod{p^2}$, $q_2 \not\equiv 1 \pmod{p^2}$, and $G_{\emptyset}((\mathbb{Q}_{\{q_1\}}\mathbb{Q}_{\{q_2\}})_{\infty})$ is trivial. Then due to [MO13, Theorem 1], the Galois group $G_S(\mathbb{Q}_{\infty}) = \operatorname{Gal}((\mathbb{Q}_{\infty})_S/\mathbb{Q}_{\infty})$ has a presentation

$$G_S(\mathbb{Q}_{\infty}) = \langle x, y \mid x^{p^2} = 1, y^{-1}xy = x^{1+p}\rangle^{\text{pro}-p}.$$

Then the presentation of the Galois group $\widetilde{G}_S(\mathbb{Q}) = \operatorname{Gal}((\mathbb{Q}_{\infty})_S/\mathbb{Q})$ can be described as follows:

$$\widetilde{G}_{S}(\mathbb{Q}) = \left\langle g_{1}, g_{2}, g_{3} \middle| \begin{array}{l} (g_{2}g_{1}^{-1})^{p^{2}} = 1, (g_{3}g_{1}^{-1})^{-1}g_{2}g_{1}^{-1}g_{3}g_{1}^{-1} = (g_{2}g_{1}^{-1})^{1+p}, \\ g_{1}g_{2} = g_{2}g_{1}, g_{1}(g_{3}g_{1}^{-1})g_{1}^{-1} = (g_{3}g_{1}^{-1})^{1+p}(g_{2}g_{1}^{-1})^{p} \right\rangle^{\operatorname{pro}-p},$$

where $\alpha(g_i) = \gamma$ for all i. Then the matrix $Q_{\widetilde{\varphi_0}}$ is given by

$$Q_{\widetilde{\varphi_0}} = \begin{pmatrix} -p^2 & p^2 & 0\\ p & -p & 0\\ \gamma - 1 & -(\gamma - 1) & 0\\ -\{\gamma - (1+2p)\} & -p & \gamma - (1+p) \end{pmatrix},$$

and the GCD $\Delta_1(A_{\widetilde{\varphi_0}})$ of all 2-minors of Q_{α} is given by

$$\Delta_1(A_{\widetilde{\varphi_0}}) \doteq \gamma - (1+p) \in \mathbb{Z}_p[[\Gamma]].$$

Let $a_1 = 1 + p \in \overline{\mathbb{Q}_p}^{\times}$ be a zero of $\Delta_1(A_{\widetilde{\varphi_0}})$. Then

$$Q_{\widetilde{\varphi_0}}|_{\gamma=1+p} = \begin{pmatrix} -p^2 & p^2 & 0\\ p & -p & 0\\ p & -p & 0\\ p & -p & 0 \end{pmatrix},$$

and hence $\mathbf{b} = {}^t(b_1, b_2, b_3)$ is a solution of $Q_{\widetilde{\varphi_0}}|_{\gamma=1+p} \cdot \mathbf{b} = \mathbf{0}$ if and only if $b_1 = b_2$. Therefore, the homomorphism $\varphi_1 \colon F_3 \to \mathrm{GL}(2; \overline{\mathbb{Q}_p})$ defined by

$$\varphi_1(g_1) = \varphi_1(g_2) = \begin{pmatrix} 1+p & b_1 \\ 0 & 1 \end{pmatrix}, \ \varphi_1(g_3) = \begin{pmatrix} 1+p & b_3 \\ 0 & 1 \end{pmatrix}$$

can be extended to a representation $\rho_1 : G_S(\mathbb{Q}) \to GL(2; \overline{\mathbb{Q}_p})$, and ρ_1 is non-abelian if and only if $b_1 \neq b_3$.

Example 2. Let $K_{\infty}/\mathbb{Q}(\sqrt{-q_1q_2})$ be the cyclotomic \mathbb{Z}_2 -extension of $\mathbb{Q}(\sqrt{-q_1q_2})$ with prime numbers $q_1 \equiv 3 \pmod{8}$ and $q_2 \equiv 7 \pmod{16}$. By [Miz10], the Galois group $N := \operatorname{Gal}((K_{\infty})_{\emptyset}/K_{\infty})$ is generated by x, y, z with three relators

$$r_1 = x^2[x, y], \quad r_2 = x^{-2}[y, z], \quad r_3 = [x, z],$$

and $\widetilde{G}_{\emptyset}(\mathbb{Q}(\sqrt{-q_1q_2})) = \operatorname{Gal}((\mathbb{Q}(\sqrt{-q_1q_2}))_{\infty})_{\emptyset}/\mathbb{Q}(\sqrt{-q_1q_2})) = N \rtimes \Gamma$ is generated by x, y, z, γ with six relators r_1, r_2, r_3 ,

$$r_4 = \gamma x \gamma^{-1} x^{-1}, \quad r_5 = x^{C_1} y^{-C_0} z^{1-C_1} \gamma z^{-1} \gamma^{-1}, \quad r_6 = \gamma y^{-1} \gamma^{-1} y z,$$

with some $C_0, C_1 \in 2\mathbb{Z}_2$. Put $g_1 = \gamma$, $g_2 = \gamma x$, $g_3 = \gamma y$. Since $z = (\gamma y^{-1} \gamma^{-1} y)^{-1}$, the Galois group $\widetilde{G}_{\emptyset}(\mathbb{Q}(\sqrt{-q_1q_2}))$ is generated by g_1, g_2, g_3 with five relators r_1, r_2, r_3, r_4, r_5 . Then we have

$$\begin{split} r_1 &= g_1^{-1} g_2 g_3^{-1} g_2 g_1^{-1} g_3, \\ r_2 &= (g_2^{-1} g_1)^2 g_3^{-1} g_1^2 g_3^{-1} g_1^{-1} g_3^2 g_1^{-1}, \\ r_3 &= g_2^{-1} g_1^2 g_3^{-1} g_1^{-1} g_3 g_1^{-1} g_2 g_3^{-1} g_1 g_3 g_1^{-1}, \\ r_4 &= g_2 g_1^{-1} g_2^{-1} g_1, \\ r_5 &= (g_1^{-1} g_2)^{C_1} (g_1^{-1} g_3)^{-C_0} (g_3^{-1} g_1 g_3 g_1^{-1})^{1-C_1} g_1^2 g_3^{-1} g_1^{-1} g_3 g_1^{-1}, \end{split}$$

and

$$Q_{\widetilde{\varphi_0}} = \begin{pmatrix} -2\gamma^{-1} & 2\gamma^{-1} & 0 \\ 2\gamma^{-1} & -2\gamma^{-1} & 0 \\ 0 & 0 & 0 \\ -1+\gamma^{-1} & 1-\gamma^{-1} & 0 \\ \gamma + C_1 - 2 + (C_0 - 2C_1 + 1)\gamma^{-1} & C_1\gamma^{-1} & -\gamma - C_1 + 2 + (-C_0 + C_1 - 1)\gamma^{-1} \end{pmatrix}.$$

The non-zero 2-minors of $Q_{\widetilde{\varphi_0}}$ are essentially

$$2\gamma^{-1}(-\gamma - C_1 + 2 + (-C_0 + C_1 - 1)\gamma^{-1})$$

and

$$(1-\gamma^{-1})(-\gamma-C_1+2+(-C_0+C_1-1)\gamma^{-1}).$$

Their GCD $\Delta_1(A_{\widetilde{\varphi_0}})$ is given by

$$\Delta_1(A_{\widetilde{\varphi_0}}) = \gamma^2 + (C_1 - 2)\gamma + (C_0 - C_1 + 1) \in \mathbb{Z}_p[[\Gamma]].$$

Let $a_1 = a \in \overline{\mathbb{Q}_p}^{\times}$ be a zero of $\Delta_1(A_{\widetilde{\varphi_0}})$. Then

$$Q_{\widetilde{\varphi_0}}|_{\gamma=a} = \begin{pmatrix} -2a^{-1} & 2a^{-1} & 0\\ 2a^{-1} & -2a^{-1} & 0\\ 0 & 0 & 0\\ -1+a^{-1} & 1-a^{-1} & 0\\ -C_1a^{-1} & C_1a^{-1} & 0 \end{pmatrix},$$

and hence $\mathbf{b} = {}^t(b_1, b_2, b_3)$ is a solution of $Q_{\widetilde{\varphi_0}}|_{\gamma=a} \cdot \mathbf{b} = \mathbf{0}$ if and only if $b_1 = b_2$. Therefore, the homomorphism $\varphi_1 \colon F_3 \to \mathrm{GL}(2; \overline{\mathbb{Q}_p})$ defined by

$$\varphi_1(g_1) = \varphi_1(g_2) = \begin{pmatrix} a & b_1 \\ 0 & 1 \end{pmatrix}, \ \varphi_1(g_3) = \begin{pmatrix} a & b_3 \\ 0 & 1 \end{pmatrix},$$

can be extended to a representation $\rho_1 : \widetilde{G}_{\emptyset}(\mathbb{Q}(\sqrt{-q_1q_2})) \to \mathrm{GL}(2; \overline{\mathbb{Q}_p})$, and ρ_1 is non-abelian if and only if $a \neq 1$ and $b_1 \neq b_3$.

Example 3. Suppose $p \neq 2$, and let K be a real quadratic field satisfying the assumption of [Kom89, Theorem]. Then the Galois group $G_{\{p\}}(K)$ is generated by x, y with a single relator

$$r = (y^{d}xy^{-d})(y^{d-1}xy^{-(d-1)})^{C_{d-1}} \cdots (yxy^{-1})^{C_{1}}xh$$

with some $h \in [N, N]$, where $N := \operatorname{Ker}(\alpha \colon G_{\{p\}}(K) \to \Gamma \ ; \ x \mapsto 1, \ y \mapsto \gamma)$, and with some $C_i \in \mathbb{Z}_p$, where $C_i \equiv (-1)^{d-i} \binom{d}{i} \pmod{p}$. Put $g_1 = y, \ g_2 = yx$. Then $x = g_1^{-1}g_2$, and

$$(\alpha \circ \pi) \left(\frac{\partial (y^{j}xy^{-j})}{\partial g_{i}} \right) = (\alpha \circ \pi) \left(\frac{\partial y}{\partial g_{i}} \right) + \gamma(\alpha \circ \pi) \left(\frac{\partial (y^{j-1}xy^{-j})}{\partial g_{i}} \right)$$

$$= (\alpha \circ \pi) \left(\frac{\partial y}{\partial g_{i}} \right) + \gamma \left((\alpha \circ \pi) \left(\frac{\partial (y^{j-1}xy^{-(j-1)})}{\partial g_{i}} \right) + (\alpha \circ \pi) \left(\frac{\partial y^{-1}}{\partial g_{i}} \right) \right)$$

$$= (\alpha \circ \pi) \left(\frac{\partial y}{\partial g_{i}} \right) + \gamma \left((\alpha \circ \pi) \left(\frac{\partial (y^{j-1}xy^{-(j-1)})}{\partial g_{i}} \right) - \gamma^{-1}(\alpha \circ \pi) \left(\frac{\partial y}{\partial g_{i}} \right) \right)$$

$$= \gamma(\alpha \circ \pi) \left(\frac{\partial (y^{j-1}xy^{-(j-1)})}{\partial g_{i}} \right)$$

$$= \begin{cases} -\gamma^{-1+j} & (i=1) \\ \gamma^{-1+j} & (i=2) \end{cases}.$$

Since $(\alpha \circ \pi) \left(\frac{\partial h}{\partial g_i} \right) = 0$,

$$(\alpha \circ \pi) \left(\frac{\partial r}{\partial g_i} \right) = \sum_{j=0}^d C_j \cdot (\alpha \circ \pi) \left(\frac{\partial (y^j x y^{-j})}{\partial g_i} \right) = (-1)^{i+1} \gamma^{-1} \sum_{j=0}^d C_j \gamma^j$$

where $C_d = 1$. The GCD $\Delta_1(A_{\widetilde{\varphi_0}})$ of 1-minors of $Q_{\widetilde{\varphi_0}}$ is given by

$$\Delta_1(A_{\widetilde{\varphi_0}}) = \sum_{j=0}^d C_j \gamma^j \in \mathbb{Z}_p[[\Gamma]].$$

Let $a_1 = a \in \overline{\mathbb{Q}_p}^{\times}$ be a zero of $\Delta_1(A_{\widetilde{\varphi_0}})$. Then

$$Q_{\widetilde{\varphi_0}}|_{\gamma=a} = \begin{pmatrix} 0 & 0 \end{pmatrix}$$

and hence the homomorphism $\varphi_1 \colon F_2 \to \mathrm{GL}(2; \overline{\mathbb{Q}_p})$ defined by

$$\varphi_1(g_1) = \begin{pmatrix} a & b_1 \\ 0 & 1 \end{pmatrix}, \ \varphi_1(g_2) = \begin{pmatrix} a & b_2 \\ 0 & 1 \end{pmatrix}$$

for any b_1 and b_2 can be extended to a representation $\rho_1 \colon G_{\{p\}}(K) \to \mathrm{GL}(2; \overline{\mathbb{Q}_p})$, and ρ_1 is non-abelian if and only if $a \neq 1$ and $b_1 \neq b_2$.

Next, we consider extensions of homomorphisms to $GL(3; \overline{\mathbb{Q}_p})$ using Example 1.

Example 4. For the same Galois group $\widetilde{G}_S(\mathbb{Q})$ and the representation $\rho_1 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(2; \mathbb{Q}_p)$ in Example 1, assume $b_1 = b_3 = b$. Namely, we consider extensions of the homomorphism $\varphi_2 \colon F_3 \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$. Consider the homomorphism $\widetilde{\varphi_1} \colon F_3 \to \mathrm{GL}(2; \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]])$ defined by

$$\widetilde{\varphi_1}(g_1) = \widetilde{\varphi_1}(g_2) = \widetilde{\varphi_1}(g_3) = \begin{pmatrix} (1+p) \cdot \gamma & b \cdot \gamma \\ 0 & \gamma \end{pmatrix}.$$

Then the matrix $Q_{\widetilde{\varphi_1}}$ is given by

$$Q_{\widetilde{\varphi_1}} = \begin{pmatrix} -p^2 & 0 & p^2 & 0 & 0 & 0 \\ 0 & -p^2 & 0 & p^2 & 0 & 0 \\ p & 0 & -p & 0 & 0 & 0 \\ 0 & p & 0 & -p & 0 & 0 \\ 0 & p & 0 & -p & 0 & 0 \\ (1+p)\gamma - 1 & b\gamma & -(1+p)\gamma + 1 & -b\gamma & 0 & 0 \\ 0 & \gamma - 1 & 0 & -\gamma + 1 & 0 & 0 \\ -(1+p)\gamma + (1+2p) & -b\gamma & -p & 0 & (1+p)(\gamma - 1) & b\gamma \\ 0 & -\gamma + (1+2p) & 0 & -p & 0 & \gamma - (1+p) \end{pmatrix},$$

and the GCD $\Delta_2(A_{\widetilde{\varphi_1}})$ of all (6-2)-minors of $Q_{\widetilde{\varphi_1}}$ is given by

$$\Delta_2(A_{\widetilde{\varphi_1}}) \doteq (\gamma - 1)\{\gamma - (1+p)\} \in \mathbb{Z}_p[[\Gamma]].$$

(i) Let $a_2 = 1$. Then

$$Q_{\widetilde{\varphi_1}}|_{\gamma=1} = \begin{pmatrix} -p^2 & 0 & p^2 & 0 & 0 & 0 \\ 0 & -p^2 & 0 & p^2 & 0 & 0 \\ p & 0 & -p & 0 & 0 & 0 \\ 0 & p & 0 & -p & 0 & 0 \\ p & b & -p & -b & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ p & -b & -p & 0 & 0 & b \\ 0 & 2p & 0 & -p & 0 & -p \end{pmatrix},$$

and hence $\mathbf{b} = {}^{t}(b_{11}, b_{12}, b_{21}, b_{22}, b_{31}, b_{32})$ is a solution of $Q_{\widetilde{\varphi_{1}}}|_{\gamma=1} \cdot \mathbf{b} = \mathbf{0}$ if and only if $b_{11} = b_{21}$, and $b_{12} = b_{22} = b_{32}$, namely,

$$\mathbf{b} = b_{11} \cdot \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + b_{12} \cdot \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \\ 0 \\ 1 \end{pmatrix} + b_{31} \cdot \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}.$$

Therefore, the homomorphism $\varphi_2 \colon F_3 \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$ defined by

$$\varphi_2(g_1) = \varphi_2(g_2) = \begin{pmatrix} 1+p & b & b_{11} \\ 0 & 1 & b_{12} \\ 0 & 0 & 1 \end{pmatrix}, \ \varphi_2(g_3) = \begin{pmatrix} 1+p & b & b_{31} \\ 0 & 1 & b_{12} \\ 0 & 0 & 1 \end{pmatrix},$$

can be extended to a non-abelian representation $\rho_2 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$, and ρ_2 is non-abelian if and only if $b_{11} \neq b_{31}$.

(ii) Let $a_2 = 1 + p$. Then

$$Q_{\widetilde{\varphi_1}}|_{\gamma=1+p} = \begin{pmatrix} -p^2 & 0 & p^2 & 0 & 0 & 0\\ 0 & -p^2 & 0 & p^2 & 0 & 0\\ p & 0 & -p & 0 & 0 & 0\\ 0 & p & 0 & -p & 0 & 0\\ p(2+p) & (1+p)b & -p(2+p) & -(1+p)b & 0 & 0\\ 0 & p & 0 & -p & 0 & 0\\ -p^2 & -(1+p)b & -p & 0 & p(1+p) & (1+p)b\\ 0 & p & 0 & -p & 0 & 0 \end{pmatrix},$$

and hence $\mathbf{b} = {}^t(b_{11}, b_{12}, b_{21}, b_{22}, b_{31}, b_{32})$ is a solution of $Q_{\widetilde{\varphi_1}}|_{\gamma=1+p} \cdot \mathbf{b} = \mathbf{0}$ for $b_{11} = b_{21}$, $b_{12} = b_{22}$, and $b_{31} = b_{11} + \frac{1}{p} \cdot b \cdot b_{12} - \frac{1}{p} \cdot b \cdot b_{32}$, namely,

$$\mathbf{b} = b_{11} \cdot \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + b_{12} \cdot \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \\ \frac{1}{p} \cdot b \\ 0 \end{pmatrix} + b_{32} \cdot \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ -\frac{1}{p} \cdot b \\ 1 \end{pmatrix}.$$

Therefore, the homomorphism $\varphi_2 \colon F_3 \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$ defined by

$$\varphi_2(g_1) = \varphi_2(g_2) = \begin{pmatrix} (1+p)^2 & (1+p) \cdot b & b_{11} \\ 0 & 1+p & b_{12} \\ 0 & 0 & 1 \end{pmatrix}, \ \varphi_2(g_3) = \begin{pmatrix} (1+p)^2 & (1+p) \cdot b & b_{31} \\ 0 & 1+p & b_{32} \\ 0 & 0 & 1 \end{pmatrix},$$

can be extended to a representation $\rho_2 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$, and ρ_2 is non-abelian if and only if $b_{11} \neq b_{31}$ or $b_{12} \neq b_{32}$.

Example 5. For the same Galois group $\widetilde{G}_S(\mathbb{Q})$ and the representation $\rho_1 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(2; \overline{\mathbb{Q}_p})$ in Example 1, consider the homomorphism $\widetilde{\varphi_1} \colon F_3 \to \mathrm{GL}(2; \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]])$ defined by

$$\widetilde{\varphi_1}(g_1) = \widetilde{\varphi_1}(g_2) = \begin{pmatrix} (1+p) \cdot \gamma & b_1 \cdot \gamma \\ 0 & \gamma \end{pmatrix}, \ \widetilde{\varphi_1}(g_3) = \begin{pmatrix} (1+p) \cdot \gamma & b_3 \cdot \gamma \\ 0 & \gamma \end{pmatrix}.$$

Then the matrix $Q_{\widetilde{\varphi_1}}$ is given by

$$Q_{\widetilde{\varphi_1}} = \begin{pmatrix} -p^2 & 0 & p^2 & 0 & 0 & 0 \\ 0 & -p^2 & 0 & p^2 & 0 & 0 \\ p & -(b_1-b_3) & -p & b_1-b_3 & 0 & 0 \\ 0 & p & 0 & -p & 0 & 0 \\ (1+p)\gamma-1 & b_1\gamma & -(1+p)\gamma+1 & -b_1\gamma & 0 & 0 \\ 0 & \gamma-1 & 0 & -\gamma+1 & 0 & 0 \\ -(1+p)\gamma+(1+2p) & \{p(b_1-b_3)-b_3\}\gamma-\frac{1}{2}p(9+p)(b_1-b_3) & -p & 4p(b_1-b_3) & (1+p)(\gamma-1) & b_1\gamma+\frac{1}{2}p(1+p)(b_1-b_3) \\ 0 & -\gamma+(1+2p) & 0 & -p & 0 & \gamma-(1+p) \end{pmatrix},$$

and the GCD $\Delta_2(A_{\widetilde{\varphi_1}})$ of all (6-2)-minors of $Q_{\widetilde{\varphi_1}}$ is given by

$$\Delta_2(A_{\widetilde{o}_1}) = \gamma - (1+p) \in \mathbb{Z}_p[[\Gamma]].$$

Let $a_2 = 1 + p$, which is also a zero of $\Delta_1(A_{\widetilde{\varphi_0}}) = \gamma - (1 + p) \in \mathbb{Z}_p[[\Gamma]]$ (cf. Example 1). Then

$$Q_{\widetilde{\varphi_1}}|_{\gamma=1+p} = \begin{pmatrix} -p^2 & 0 & p^2 & 0 & 0 & 0 \\ 0 & -p^2 & 0 & p^2 & 0 & 0 \\ p & -(b_1-b_3) & -p & b_1-b_3 & 0 & 0 \\ 0 & p & 0 & -p & 0 & 0 \\ p(2+p) & (1+p)b_1 & -p(2+p) & -(1+p)b_1 & 0 & 0 \\ 0 & p & 0 & -p & 0 & 0 \\ -p^2 & -(1+p)b_3 + \frac{1}{2}p(-7+p)(b_1-b_3) & -p & 4p(b_1-b_3) & p(1+p) & \frac{1}{2}(1+p)\{2b_1+p(b_1-b_3)\} \\ 0 & p & 0 & -p & 0 & 0 \end{pmatrix},$$

and hence $\mathbf{b} = {}^t(b_{11}, b_{12}, b_{21}, b_{22}, b_{31}, b_{32})$ is a solution of $Q_{\widetilde{\varphi_1}}|_{\gamma=1+p} \cdot \mathbf{b} = \mathbf{0}$ if and only if $b_{11} = b_{21}$, $b_{12} = b_{22}$, and $b_{31} = b_{11} + \{\frac{1}{p} \cdot b_3 - \frac{1}{2}(b_1 - b_3)\} \cdot b_{12} + \{-\frac{1}{p} \cdot b_1 - \frac{1}{2}(b_1 - b_3)\} \cdot b_{32}$, namely,

$$\mathbf{b} = b_{11} \cdot \begin{pmatrix} 1\\0\\1\\0\\1\\0 \end{pmatrix} + b_{12} \cdot \begin{pmatrix} 0\\1\\0\\1\\\frac{1}{p} \cdot b_3 - \frac{1}{2}(b_1 - b_3)\\0 \end{pmatrix} + b_{32} \cdot \begin{pmatrix} 0\\0\\0\\-\frac{1}{p} \cdot b_1 - \frac{1}{2}(b_1 - b_3)\\1 \end{pmatrix}.$$

Therefore, the homomorphism $\varphi_2 \colon F_3 \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$ defined by

$$\varphi_2(g_1) = \varphi_2(g_2) = \begin{pmatrix} (1+p)^2 & (1+p) \cdot b_1 & b_{11} \\ 0 & 1+p & b_{12} \\ 0 & 0 & 1 \end{pmatrix}, \ \varphi_2(g_3) = \begin{pmatrix} (1+p)^2 & (1+p) \cdot b_3 & b_{31} \\ 0 & 1+p & b_{32} \\ 0 & 0 & 1 \end{pmatrix},$$

can be extended to a representation $\rho_2 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$, and ρ_2 is non-abelian if and only if $b_1 \neq b_3$, $b_{11} \neq b_{31}$, or $b_{12} \neq b_{32}$.

4. Cohomological interpretations

In this Section, we focus on cohomological interpretations of our observations so far. Generalizations to higher-dimensional cases are due to [HP15, Section 4].

For a given $a_{k-1} \in \overline{\mathbb{Q}_p}^{\times}$ such that $|a_{k-1} - 1|_p < 1$, note that we have the relation

$$\overline{\beta_{k-1}}(xy) = \overline{\beta_{k-1}}(x) + \widetilde{\varphi_{k-2}}(x)|_{\gamma = a_{k-1}} \overline{\beta_{k-1}}(y) = \overline{\beta_{k-1}}(x) + a_{k-1}^{(\alpha_{\gamma} \circ \pi)(x)} \cdot \varphi_{k-2}(x) \cdot \overline{\beta_{k-1}}(y)$$

for any $x, y \in F_n$, where $\overline{\beta_{k-1}} \colon F_n \to \overline{\mathbb{Q}_p}^{\oplus (k-1)}$ is the crossed homomorphism, $\widetilde{\varphi_{k-2}} := \varphi_{k-2} \otimes (\alpha \circ \pi) \colon F_n \to \mathrm{GL}(k-1; \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]])$ is the tensor product homomorphism, and $\alpha_\gamma \colon G \to \mathbb{Z}_p$

maps $g \in G$ to the power of γ of $\alpha(g)$. In particular, when $\rho_{k-2} \colon G \to \mathrm{GL}(k-1; \overline{\mathbb{Q}_p})$ is a representation such that $\rho_{k-2} \circ \pi = \widehat{\varphi_{k-2}}|_{\gamma = a_{k-1}}$,

$$\beta_{k-1}(xy) = \beta_{k-1}(x) + \rho_{k-2}(x)\beta_{k-1}(y),$$

for any $x, y \in G$, where $\beta_{k-1} : G \to \overline{\mathbb{Q}_p}^{\oplus (k-1)}$ is a crossed homomorphism such that $\beta_{k-1} \circ \pi = \overline{\beta_{k-1}}$. We call $\beta_{k-1} : G \to \overline{\mathbb{Q}_p}^{\oplus (k-1)}$ a 1-cocycle with coefficients $(\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}$.

On the other hand, if there exists $v \in \overline{\mathbb{Q}_p}^{\oplus (k-1)}$ such that

$$\overline{\beta_{k-1}}(x) = \widetilde{\varphi_{k-2}}(x)|_{\gamma = a_{k-1}}v - v = (\widetilde{\varphi_{k-2}}(x)|_{\gamma = a_{k-1}} - I_{k-1})v = (a_{k-1}^{(\alpha_{\gamma} \circ \pi)(x)} \cdot \varphi_{k-2}(x) - I_{k-1}) \cdot v$$

for any $x \in F_n$, where $I_{k-1} \in \mathrm{GL}(k-1; \overline{\mathbb{Q}_p})$ is the $(k-1) \times (k-1)$ identity matrix, namely,

(4.1)
$$\beta_{k-1}(x) = \rho_{k-2}(x)v - v = (\rho_{k-2}(x) - I_{k-1})v,$$

for any $x \in G$, we call $\beta_{k-1} \colon G \to \overline{\mathbb{Q}_p}^{\oplus (k-1)}$ a 1-coboundary with coefficients $(\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}$. We respectively denote the set of continuous 1-cocyles and continuous 1-coboundaries by $Z^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}})$ and $B^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}})$. We can easily see that $B^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}) \subset Z^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}})$, and we define the first continuous cohomology group $H^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}})$ with coefficients $(\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}$ by

$$H^1(G,(\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}) := Z^1(G,(\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}})/B^1(G,(\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}).$$

Denote the first cohomology class of β_{k-1} in $H^1(G,(\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}})$ by $[\beta_{k-1}]$.

From now on, let us consider the relation between the zeros of the arithmetic twisted Alexander polynomial $\Delta_{k-1}(A_{\widetilde{\varphi_{k-2}}}) \in \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]]$ and the non-vanishment of $H^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}})$. This may be seen as a generalization of [Mor24, Corollary 14.4.5].

Theorem 2. If $a_{k-1} \in \overline{\mathbb{Q}_p}^{\times}$ is a zero of $\Delta_{k-1}(A_{\widetilde{\varphi_{k-2}}}) \in \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]]$, then $H^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}})$ does not vanish.

Proof. Since we have an isomorphism $Z^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}) \simeq \operatorname{Hom}_{\overline{\mathbb{Q}_p}}(A_{\widetilde{\varphi_{k-2}}}, \overline{\mathbb{Q}_p}^{\oplus (k-1)})$ as vector spaces over $\overline{\mathbb{Q}_p}$, and $\dim_{\overline{\mathbb{Q}_p}}(A_{\widetilde{\varphi_{k-1}}}, \overline{\mathbb{Q}_p}) \geq k$ by Theorem 1, we have

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus (k-1)}} Z^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}) \ge k.$$

On the other hand, since we have a surjective linear map

$$(4.2) F: \overline{\mathbb{Q}_p}^{\oplus (k-1)} \to B^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}),$$

defined by $F(v) = \beta_{k-1,v}$, where $\beta_{k-1,v}(x) = \rho_{k-2}(x)v - v$ for any $x \in G$, we have

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus (k-1)}} B^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}) \le k-1.$$

Hence, we have $H^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}) \neq 0$.

For the converse, we have the following:

Theorem 3. Assume the matrices $\{\rho_{k-2}(x)\}_{x\in G}\subset \mathrm{GL}(k-1,\overline{\mathbb{Q}_p})$ do not have a simultaneous eigenvector with simultaneous eigenvalue 1. If $H^1(G,(\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}})$ does not vanish, then $a_{k-1}\in\overline{\mathbb{Q}_p}^{\times}$ is a zero of $\Delta_{k-1}(A_{\widetilde{\varphi_{k-2}}})\in\overline{\mathbb{Q}_p}\otimes_{\mathbb{Z}_p}\mathbb{Z}_p[[\Gamma]]$.

Proof. Since $H^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}) \neq 0$, we have

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus (k-1)}} B^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}) < \dim_{\overline{\mathbb{Q}_p}^{\oplus (k-1)}} Z^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}).$$

Due to the assumption, since we do not have a vector $v \in \overline{\mathbb{Q}_p}^{\oplus (k-1)}$ such that $\rho_{k-2}(x)v = v$ for any $x \in G$, we have $\operatorname{Ker} F = 0$, where $F : \overline{\mathbb{Q}_p}^{\oplus (k-1)} \to B^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}})$ is the surjective linear map (4.2). Hence, we have

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus (k-1)}} B^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}) = k-1.$$

This means that

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus (k-1)}} \operatorname{Hom}_{\overline{\mathbb{Q}_p}}(A_{\widetilde{\varphi_{k-2}}}, \overline{\mathbb{Q}_p}^{\oplus (k-1)}) = \dim_{\overline{\mathbb{Q}_p}^{\oplus (k-1)}} Z^1(G, (\overline{\mathbb{Q}_p}^{\oplus (k-1)})_{\rho_{k-2}}) \ge k$$

and by Theorem 1,
$$a_{k-1} \in \overline{\mathbb{Q}_p}^{\times}$$
 is a zero of $\Delta_{k-1}(A_{\widetilde{\varphi_{k-2}}}) \in \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]]$.

Here are some examples. Note that we have an example which does not satisfy the assumption of Theorem 3 but the claim holds (Example 7), and the claim does not hold (Example 10).

Example 6. Consider the representation $\rho_1 : \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(2; \overline{\mathbb{Q}_p})$ constructed in Example 1, which is defined by

$$\rho_1(g_1) = \rho_1(g_2) = \begin{pmatrix} 1+p & b_1 \\ 0 & 1 \end{pmatrix}, \ \rho_1(g_3) = \begin{pmatrix} 1+p & b_3 \\ 0 & 1 \end{pmatrix}.$$

Let $\beta_{1,(b_1,b_3)}: \widetilde{G}_S(\mathbb{Q}) \to \overline{\mathbb{Q}_p}$ be a 1-cocycle such that $\beta_{1,(b_1,b_3)}(g_1) = \beta_{1,(b_1,b_3)}(g_2) = b_1$ and $\beta_{1,(b_1,b_3)}(g_3) = b_3.$

If $b_1 = b_3 = b$, namely if ρ is abelian, then $\beta_{1,(b,b)}$ is a 1-coboundary since we can take $v \in \overline{\mathbb{Q}_p}$

as $\frac{b}{(1+p)-1} = \frac{1}{p} \cdot b$ which satisfies (4.1). If $b_1 \neq b_3$, namely if ρ is non-abelian, then $\beta_{1,(b_1,b_3)}$ is not a 1-coboundary since we cannot take $v \in \overline{\mathbb{Q}_p}$ which satisfies (4.1).

Since the cocycles are unique up to adding a coboundary and up to multiplying by a non-zero scalar, we have two different cocycles $\beta_{1,(1,1)}$ and $\beta_{1,(1,0)}$, where the cohomology class $[\beta_{1,(1,1)}]$ is trivial, whereas $[\beta_{1,(1,0)}]$ is non-trivial. Therefore, we have

$$\dim_{\overline{\mathbb{Q}_p}} Z^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p})_{\rho_0}) = 2,$$

$$\dim_{\overline{\mathbb{Q}_p}} B^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p})_{\rho_0}) = 1,$$

$$\dim_{\overline{\mathbb{Q}_p}} H^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p})_{\rho_0}) = 1.$$

Example 7. Consider the representation $\rho_2 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$ constructed in Example 4 (i), which is defined by

$$\rho_2(g_1) = \rho_2(g_2) = \begin{pmatrix} 1+p & b & b_{11} \\ 0 & 1 & b_{12} \\ 0 & 0 & 1 \end{pmatrix}, \ \rho_2(g_3) = \begin{pmatrix} 1+p & b & b_{31} \\ 0 & 1 & b_{12} \\ 0 & 0 & 1 \end{pmatrix}.$$

Let $\beta_{2,(b_{11},b_{12},b_{31},b_{32})} : \widetilde{G}_S(\mathbb{Q}) \to \overline{\mathbb{Q}_p}^{\oplus 2}$ be a 1-cocycle with coefficient φ_2 such that $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}(g_1) = \beta_{2,(b_{11},b_{12},b_{31},b_{32})}(g_2) = {}^t(b_{11},b_{12})$ and $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}(g_3) = {}^t(b_{31},b_{32})$.

If $b_{11} = b_{31}$ and $b_{12} = b_{32} = 0$, then $\beta_{2,(b_{11},0,b_{11},0)}$ is a 1-coboundary since we can take $v \in \overline{\mathbb{Q}_p}^{\oplus 2}$ as $\frac{1}{p} \begin{pmatrix} b_{11} \\ 0 \end{pmatrix}$ which satisfies (4.1).

For the rest of the cases, $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}$ is not a 1-coboundary since we cannot take $v \in \overline{\mathbb{Q}_p}^{\oplus 2}$ which satisfies (4.1).

Hence, we have three different cocycles $\beta_{2,(1,0,1,0)}$, $\beta_{2,(0,1,0,1)}$, and $\beta_{2,(0,0,1,0)}$, where the cohomology class $[\beta_{2,(1,0,1,0)}]$ is trivial, whereas $[\beta_{2,(0,1,0,1)}]$ and $[\beta_{2,(0,0,1,0)}]$ are non-trivial. Therefore, we have

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} Z^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1}) = 3,$$

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} B^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1}) = 1,$$

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} H^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1}) = 2.$$

Example 8. Consider the representation $\rho_2 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$ constructed in Example 4 (ii), which is defined by

$$\rho_2(g_1) = \rho_2(g_2) = \begin{pmatrix} (1+p)^2 & (1+p) \cdot b & b_{11} \\ 0 & 1+p & b_{12} \\ 0 & 0 & 1 \end{pmatrix}, \ \rho_2(g_3) = \begin{pmatrix} (1+p)^2 & (1+p) \cdot b & b_{31} \\ 0 & 1+p & b_{32} \\ 0 & 0 & 1 \end{pmatrix}.$$

Let $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}: \widetilde{G}_{S}(\mathbb{Q}) \to \overline{\mathbb{Q}_{p}}^{\oplus 2}$ be a 1-cocycle with coefficient φ_{2} such that $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}(g_{1}) = \beta_{2,(b_{11},b_{12},b_{31},b_{32})}(g_{2}) = {}^{t}(b_{11},b_{12})$ and $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}(g_{3}) = {}^{t}(b_{31},b_{32})$.

If $b_{11} = b_{31}$ and $b_{12} = b_{32} = 0$, then $\beta_{2,(b_{11},0,b_{11},0)}$ is a 1-coboundary since we can take $v \in \overline{\mathbb{Q}_p}^{\oplus 2}$ as $\frac{1}{p(2+p)} \binom{b_{11}}{0}$ which satisfies (4.1).

If $b_{11} = b_{31} = 0$ and $b_{12} = b_{32}$, then $\beta_{2,(0,b_{12},0,b_{12})}$ is a 1-coboundary since we can take $v \in \overline{\mathbb{Q}_p}^{\oplus 2}$ as $\frac{1}{p^2(2+p)} \begin{pmatrix} -(1+p) \cdot b \cdot b_{12} \\ p(2+p) \cdot b_{12} \end{pmatrix}$ which satisfies (4.1).

For the rest of the cases, $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}$ is not a 1-coboundary since we cannot take $v \in \overline{\mathbb{Q}_p}^{\oplus 2}$ which satisfies (4.1).

Hence, we have three different cocycles $\beta_{2,(1,0,1,0)}$, $\beta_{2,(0,1,0,1)}$, and $\beta_{2,(0,0,-\frac{1}{p}\cdot b,1)}$, where the cohomology classes $[\beta_{2,(1,0,1,0)}]$ and $[\beta_{2,(0,1,0,1)}]$ are trivial, whereas $[\beta_{2,(0,0,-\frac{1}{p}\cdot b,1)}]$ is non-trivial.

Therefore, we have

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} Z^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})) = 3,$$

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} B^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})) = 2,$$

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} H^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})) = 1.$$

Example 9. Consider the representation $\rho_2 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$ constructed in Example 5, which is defined by

$$\rho_2(g_1) = \rho_2(g_2) = \begin{pmatrix} (1+p)^2 & (1+p) \cdot b_1 & b_{11} \\ 0 & 1+p & b_{12} \\ 0 & 0 & 1 \end{pmatrix}, \ \rho_2(g_3) = \begin{pmatrix} (1+p)^2 & (1+p) \cdot b_3 & b_{31} \\ 0 & 1+p & b_{32} \\ 0 & 0 & 1 \end{pmatrix}.$$

Let $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}$: $\widetilde{G}_{S}(\mathbb{Q}) \to \overline{\mathbb{Q}_{p}}^{\oplus 2}$ be a 1-cocycle with coefficient φ_{2} such that $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}(g_{1}) = \beta_{2,(b_{11},b_{12},b_{31},b_{32})}(g_{2}) = {}^{t}(b_{11},b_{12})$ and $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}(g_{3}) = {}^{t}(b_{31},b_{32})$.

If $b_{11} = b_{31}$ and $b_{12} = b_{32} = 0$, then $\beta_{2,(b_{11},0,b_{11},0)}$ is a 1-coboundary since we can take $v \in \overline{\mathbb{Q}_p}^{\oplus 2}$ as $\frac{1}{p(2+p)} \binom{b_{11}}{0}$ which satisfies (4.1).

If $b_{11} = 0$, $b_{12} = b_{32}$, and $b_{31} = -\frac{1+p}{p} \cdot (b_1 - b_3) \cdot b_{12}$, then $\beta_{2,(0,b_{12},-\frac{1+p}{p}\cdot(b_1-b_3)\cdot b_{12},b_{12})}$ is a 1-coboundary since we can take $v \in \overline{\mathbb{Q}_p}^{\oplus 2}$ as $\frac{1}{p^2(2+p)} \begin{pmatrix} -(1+p) \cdot b_1 \cdot b_{12} \\ p(2+p) \cdot b_{12} \end{pmatrix}$ which satisfies (4.1).

For the rest of the cases, $\beta_{2,(b_{11},b_{12},b_{31},b_{32})}$ is not a 1-coboundary since we cannot take $v \in \overline{\mathbb{Q}_p}^{\oplus 2}$ which satisfies (4.1).

Hence, we have three different cocycles $\beta_{2,(1,0,1,0)}$, $\beta_{2,(0,1,-\frac{1+p}{p}\cdot(b_1-b_3),1)}$, and $\beta_{2,(0,0,-\frac{1}{p}\cdot b_3-\frac{1}{2}(b_1-b_3),1)}$, where the cohomology classes $[\beta_{2,(1,0,1,0)}]$ and $[\beta_{2,(0,1,-\frac{1+p}{p}\cdot(b_1-b_3),1)}]$ are trivial, whereas $[\beta_{2,(0,0,-\frac{1}{p}\cdot b_1-\frac{1}{2}(b_1-b_3),1)}]$ is non-trivial. Therefore, we have

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} Z^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})) = 3,$$

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} B^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})) = 2,$$

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} H^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})) = 1.$$

Remark 2. Since

$$\frac{1}{4} \cdot \begin{pmatrix} b^2 \\ 2b \\ b^2 \\ 2b \end{pmatrix} = \frac{b^2}{4} \cdot \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + \frac{b}{2} \cdot \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \end{pmatrix} + 0 \cdot \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}
= \frac{b^2}{4} \cdot \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + \frac{b}{2} \cdot \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \end{pmatrix} + 0 \cdot \begin{pmatrix} 0 \\ 0 \\ -\frac{1}{p} \cdot b \\ 1 \end{pmatrix},
\frac{1}{4} \cdot \begin{pmatrix} b_1^2 \\ 2b_1 \\ b_3^2 \\ 2b_3 \end{pmatrix} = \frac{b_1^2}{4} \cdot \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + \frac{b_1}{2} \cdot \begin{pmatrix} 0 \\ -\frac{1+p}{p} \cdot (b_1 - b_3) \\ 1 \end{pmatrix} + \frac{b_1 - b_3}{2} \cdot \begin{pmatrix} 0 \\ -\frac{1}{p} \cdot b_1 - \frac{1}{2}(b_1 - b_3) \\ 1 \end{pmatrix},$$

the cocycle $\beta_{\operatorname{Sym}^2(\rho_1)}$ obtained from the second symmetric power representation $\operatorname{Sym}^2(\rho_1) \colon \widetilde{G}_S(\mathbb{Q}) \to \operatorname{GL}(3; \overline{\mathbb{Q}_p})$ of the representation ρ_1 for the case of Example 8 is contained in the trivial cohomology class in $H^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})$, namely

$$[\beta_{\operatorname{Sym}^2(\rho_1)}] = 0 \in H^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1}),$$

and the representation ρ_1 for the case of Example 7 and Example 9 is contained in the non-trivial cohomology class in $H^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})$, namely

$$[\beta_{\operatorname{Sym}^2(\rho_1)}] \neq 0 \in H^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1}).$$

Example 10. For the same Galois group $\widetilde{G}_S(\mathbb{Q})$ and the representation $\rho_1 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(2; \overline{\mathbb{Q}_p})$ in Example 1, assume $b_1 = b_3 = b$. Namely, we consider extensions of the homomorphism $\varphi_2 \colon F_3 \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$. Consider the homomorphism $\widetilde{\varphi_1} \colon F_3 \to \mathrm{GL}(2; \overline{\mathbb{Q}_p} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]])$ defined by

$$\widetilde{\varphi_1}(g_1) = \widetilde{\varphi_1}(g_2) = \widetilde{\varphi_1}(g_3) = \begin{pmatrix} (1+p) \cdot \gamma & b \cdot \gamma \\ 0 & \gamma \end{pmatrix}.$$

Then the matrix $Q_{\widetilde{\varphi_1}}$ is given by

$$Q_{\widetilde{\varphi_1}} = \begin{pmatrix} -p^2 & 0 & p^2 & 0 & 0 & 0 \\ 0 & -p^2 & 0 & p^2 & 0 & 0 \\ p & 0 & -p & 0 & 0 & 0 \\ 0 & p & 0 & -p & 0 & 0 \\ 0 & p & 0 & -p & 0 & 0 \\ (1+p)\gamma-1 & b\gamma & -(1+p)\gamma+1 & -b\gamma & 0 & 0 \\ 0 & \gamma-1 & 0 & -\gamma+1 & 0 & 0 \\ -(1+p)\gamma+(1+2p) & -b\gamma & -p & 0 & (1+p)(\gamma-1) & b\gamma \\ 0 & -\gamma+(1+2p) & 0 & -p & 0 & \gamma-(1+p) \end{pmatrix},$$

and the GCD $\Delta_2(A_{\widetilde{\omega_1}})$ of all (6-2)-minors of $Q_{\widetilde{\omega_1}}$ is given by

$$\Delta_2(A_{\widetilde{\varphi_1}}) \doteq (\gamma - 1)\{\gamma - (1+p)\} \in \mathbb{Z}_p[[\Gamma]].$$

Let $a_2 \neq 0$, 1, 1+p. Then $\mathbf{b} = {}^t(b_{11}, b_{12}, b_{21}, b_{22}, b_{31}, b_{32})$ is a solution of $Q_{\widetilde{\varphi_1}}|_{\gamma=a_2} \cdot \mathbf{b} = \mathbf{0}$ if and only if $b_{11} = b_{21} = b_{31}$, and $b_{12} = b_{22} = b_{32}$, namely,

$$\mathbf{b} = b_{11} \cdot \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + b_{12} \cdot \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \\ 0 \\ 1 \end{pmatrix}.$$

Therefore, the homomorphism $\varphi_2 \colon F_3 \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$ defined by

$$\varphi_2(g_1) = \varphi_2(g_2) = \varphi_2(g_3) = \begin{pmatrix} a_2 \cdot (1+p) & a_2 \cdot b & b_{11} \\ 0 & a_2 & b_{12} \\ 0 & 0 & 1 \end{pmatrix},$$

can be extended to a representation $\rho_2 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(3; \overline{\mathbb{Q}_p}).$

Let $a_2 = \frac{1}{1+p}$, which is not a zero of $\Delta_2(A_{\widetilde{\varphi_1}})$, but the inverse is a zero of $\Delta_2(A_{\widetilde{\varphi_1}})$. Let b = 0. Consider the representation $\rho_2 \colon \widetilde{G}_S(\mathbb{Q}) \to \mathrm{GL}(3; \overline{\mathbb{Q}_p})$, which is defined by

$$\rho_2(g_1) = \rho_2(g_2) = \rho_2(g_3) = \begin{pmatrix} 1 & 0 & b_{11} \\ 0 & \frac{1}{1+p} & b_{12} \\ 0 & 0 & 1 \end{pmatrix}.$$

Let $\beta_{2,(b_{11},b_{12})} \colon \widetilde{G}_S(\mathbb{Q}) \to \overline{\mathbb{Q}_p}^{\oplus 2}$ be a 1-cocycle with coefficient φ_2 such that $\beta_{2,(b_{11},b_{12})}(g_1) = \beta_{2,(b_{11},b_{12})}(g_2) = \beta_{2,(b_{11},b_{12})}(g_3) = {}^t(b_{11},b_{12}).$

If $b_{11} \neq 0$ and $b_{12} = 0$, then $\beta_{2,(b_{11},0)}$ is not a 1-coboundary since we cannot take $v = {}^{t}(v_{1}, v_{2}) \in \overline{\mathbb{Q}_{p}}^{\oplus 2}$ which satisfies (4.1), namely

$$\begin{pmatrix} (1-1) \cdot v_1 + 0 \cdot v_2 \\ \left(\frac{1}{1+p} - 1\right) \cdot v_2 \end{pmatrix} = \begin{pmatrix} b_{11} \\ 0 \end{pmatrix}.$$

If $b_{11} = 0$ and $b_{12} \neq 0$, then $\beta_{2,(0,b_{12})}$ is a 1-coboundary since we can take $v = {}^t(v_1, v_2) \in \overline{\mathbb{Q}_p}^{\oplus 2}$ as $v = \begin{pmatrix} 1 \\ -\frac{1+p}{p} \cdot b_{12} \end{pmatrix}$ which satisfies (4.1), namely

$$\begin{pmatrix} (1-1) \cdot v_1 + 0 \cdot v_2 \\ \left(\frac{1}{1+p} - 1\right) \cdot v_2 \end{pmatrix} = \begin{pmatrix} 0 \\ b_{12} \end{pmatrix}.$$

Therefore, we have

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} Z^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})) = 2,$$

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} B^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})) = 1,$$

$$\dim_{\overline{\mathbb{Q}_p}^{\oplus 2}} H^1(\widetilde{G}_S(\mathbb{Q}), (\overline{\mathbb{Q}_p}^{\oplus 2})_{\rho_1})) = 1.$$

Acknowledgements. The authors would like to thank Takahiro Kitayama, Masanori Morishita and Yoshikazu Yamaguchi for helpful communications. This work is partially supported by JSPS KAKENHI Grant Numbers JP22K03268, JP22H01117, and JP21K03240.

References

- [AC22] Eric Ahlqvist and Magnus Carlson. "Massey products in the étale cohomology of number fields". arXiv:2207.06353 (2022) (1).
- [BLM13] Julien Blondeau, Philippe Lebacque, and Christian Maire. "On the cohomological dimension of some pro-p-extensions above the cyclotomic \mathbb{Z}_p -extension of a number field". *Mosc. Math. J.* 13 (2013), pp. 601–619 (6).
- [Bur67] Gerhard Burde. "Darstellungen von Knotengruppen". Math. Ann. 173 (1967), pp. 24–33 (1).
- [deR67] Georges de Rham. "Introduction aux polynômes d'un nœud". Enseign. Math. (2) 13 (1967), pp. 187–194 (1).
- [EM22] Abdelaziz El Habibi and Yasushi Mizusawa. "On pro-p-extensions of number fields with restricted ramification over intermediate \mathbb{Z}_p -extensions". J. Number Theory 231 (2022), pp. 214–238 (6).
- [HP15] Michael Heusener and Joan Porti. "Representations of knot groups into $SL_n(\mathbb{C})$ and twisted Alexander polynomials". *Pac. J. Math.* 277.2 (2015), pp. 313–354 (1, 3, 6, 11).
- [HPS01] Michael Heusener, Joan Porti, and Eva Suárez Peiró. "Deformations of reducible representations of 3-manifold groups into $SL_2(\mathbb{C})$ ". J. Reine Angew. Math. 530 (2001), pp. 191–227 (1, 3, 4).
- [Hir97] Eriko Hironaka. "Alexander stratifications of character varieties". Ann. Inst. Fourier (Grenoble). Vol. 47. 2. 1997, pp. 555–583 (1).
- [Iha86] Yasutaka Ihara. "On Galois representations arising from towers of coverings of $\mathbf{P}^1 \setminus \{0, 1, \infty\}$ ". *Invent. Math.* 86.3 (1986), pp. 427–459 (2, 4).
- [Kit13] Teruaki Kitano. "Linear Representations of a Knot Group over a Finite Ring and Alexander Polynomial as an Obstruction". *RIMS Kôkyûroku* 1836 (2013), pp. 65–74 (1, 3).
- [KMTT18] Takahiro Kitayama, Masanori Morishita, Ryoto Tange, and Yuji Terashima. "On certain *L*-functions for deformations of knot group representations". *Trans. Amer. Math. Soc.* 370.5 (2018), pp. 3171–3195 (6).
- [KMTT22] Takahiro Kitayama, Masanori Morishita, Ryoto Tange, and Yuji Terashima. "On adjoint homological Selmer modules for SL₂-representations of knot groups". *Int. Math. Res. Not. IMRN* (2022) (6).
- [Kom89] Keiichi Komatsu. "On the maximal p-extensions of real quadratic fields unramified outside p". J. Algebra 123.1 (1989), pp. 240–247 (8).

- [Le94] Tkhang Le. "Varieties of representations and their cohomology-jump subvarieties for knot groups". Russian Acad. Sci. Sb. Math. 78.1 (1994), pp. 187–209 (1).
- [Lin01] Xiao Song Lin. "Representations of knot groups and twisted Alexander polynomials". Acta Math. Sin. (Engl. Ser.) 17 (2001), pp. 361–380 (1, 6).
- [Maz64] Barry Mazur. "Remarks on the Alexander Polynomial" (1964). URL: https://sites.harvard.edu/barry-mazur/older-material/(1).
- [Maz89] Barry Mazur. "Deforming galois representations". Galois Groups over \mathbb{Q} . Springer. 1989, pp. 385–437 (1).
- [Maz00] Barry Mazur. The theme of p-adic variation. Mathematics: frontiers and perspectives, Amer. Math. Soc., Providence, RI, 2000, pp. 433–459 (6).
- [Miz10] Yasushi Mizusawa. "On the maximal unramified pro-2-extension over the cyclotomic \mathbb{Z}_2 -extension of an imaginary quadratic field". J. Théor. Nombres Bordeaux 22.1 (2010), pp. 115–138 (7).
- [Miz19] Yasushi Mizusawa. "On pro-p link groups of number fields". Trans. Amer. Math. Soc. 372.10 (2019), pp. 7225–7254 (6).
- [MO13] Yasushi Mizusawa and Manabu Ozaki. "On tame pro-p Galois groups over basic \mathbb{Z}_p -extensions". $Math.\ Z.\ 273.3\ (2013),\ pp.\ 1161-1173\ (7).$
- [Mor02] Masanori Morishita. "On certain analogies between knots and primes". *J. Reine Angew. Math.* 550 (2002), pp. 141–167 (2).
- [Mor04] Masanori Morishita. "Milnor invariants and Massey products for prime numbers". Compos. Math. 140.1 (2004), pp. 69–83 (1).
- [Mor06] Masanori Morishita. "On the Alexander stratification in the deformation space of Galois characters". Kyushu J. Math. 60.2 (2006), pp. 405–414 (1).
- [Mor24] Masanori Morishita. Knots and Primes: An Introduction to Arithmetic Topology. Universitext, Springer, London, second ed., 2024 (1, 2, 6, 12).
- [MTTU17] Masanori Morishita, Yu Takakura, Yuji Terashima, and Jun Ueki. "On the universal deformations for SL₂-representations of knot groups". *Tohoku Math. J.* 69.1 (2017), pp. 67–84 (6).
- [NSW20] Jürgen Neukirch, Alexander Schmidt, and Kay Wingberg. Cohomology of number fields. Vol. 323. Springer Science & Business Media, second ed., 2020 (2, 6).
- [Nor76] Douglas G. Northcott. Finite free resolutions. 71. Cambridge Univ. Press, 1976 (3).
- [S24] The Sage Developers. SageMath, the Sage Mathematics Software System (Version 10.7). 2024. URL: https://www.sagemath.org (6).
- [Sal08] Landry Salle. "Sur les pro-p-extensions à ramification restreinte au-dessus de la \mathbb{Z}_p -extension cyclotomique d'un corps de nombres". J. Théor. Nombres Bordeaux 20.2 (2008), pp. 485–523 (6).
- [Sha63] Igor R. Shafarevich. "Extensions with prescribed ramification points". *Publ. Math. IHÉS* 18 (1963), pp. 71–92 (6).
- [Sha07a] Romyar T. Sharifi. "Iwasawa theory and the Eisenstein ideal". $Duke\ Math.\ J.$ 137.1 (2007), pp. 63–101 (1, 6).
- [Sha07b] Romyar T. Sharifi. "Massey products and ideal class groups". J. Reine Angew. Math. 603 (2007), pp. 1–33 (1, 6).

- [SW11] Daniel S. Silver and Susan G. Williams. "On a Theorem of Burde and de Rham".

 J. Knot Theory Ramifications 20.05 (2011), pp. 713–720 (1, 3).
- [Wad94] Masaaki Wada. "Twisted Alexander polynomial for finitely presentable groups". Topology 33.2 (1994), pp. 241–256 (1, 6).
- [W24] Wolfram Research, Inc. *Mathematica*, (Version 14.0). Champaign, IL. 2024. URL: https://www.wolfram.com/mathematica (6).

Yasushi Mizusawa mizusawa@rikkyo.ac.jp

Department of Mathematics, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima-ku, 171-8501 Tokyo, Japan

Ryoto Tange rtange.math@gmail.com

Department of Mathematics, School of Education, Waseda University, 1-104, Totsuka-cho, Shinjuku-ku, Tokyo, 169-8050, Japan

Yuji Terashima yujiterashima@tohoku.ac.jp

Graduate School of Science, Tohoku University, 6-3, Aoba, Aramaki-aza, Aoba-ku, Sendai, 980-8578, Japan