

Type II orientifold flux vacua in 3D

Álvaro Arboleya, Adolfo Guarino and Matteo Morittu

Departamento de Física, Universidad de Oviedo,
Avda. Federico García Lorca 18, 33007 Oviedo, Spain.

Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA)
Calle de la Independencia 13, 33004 Oviedo, Spain.

alvaroarbo@gmail.com , adolfo.guarino@uniovi.es ,
morittumatteo@uniovi.es

Abstract

We initiate a systematic study of type II orientifold flux vacua in three dimensions including gauge and metric fluxes, O-planes and D-branes. We derive simple flux models (we dub them RSTU-models) that admit a description in terms of gauged supergravities with half-maximal $\mathcal{N} = 8$ supersymmetry in three dimensions. As a landscape appetizer, we present various multi-parametric families of supersymmetric and non-supersymmetric AdS_3 and Mkw_3 vacua. Remarkably, negative masses turn out to be always absent in the spectrum of scalar fluctuations, thus making all the vacua perturbatively stable within half-maximal supergravity. We provide examples of non-supersymmetric type IIB AdS_3 flux vacua which feature parametrically-controlled scale separation and come along with integer-valued conformal dimensions of the would-be dual CFT_2 operators. We also comment on the implications of our results in light of the Swampland Program.

Contents

1	Motivation	1
2	A crash course on half-maximal 3D supergravity	3
2.1	Gaugings, embedding tensor and quadratic constraints	4
2.2	Fermion masses and scalar potential	5
3	Type II orientifolds	6
3.1	Warming up: M-theory fluxes/embedding tensor dictionary	8
3.2	Type IIA with O2-planes	10
3.3	Type IIB with O3-planes	13
3.4	Type IIA with O4-planes	16
3.5	Type IIB with O5-planes	19
3.6	Type IIA with O6-planes	22
3.7	Type IIB with O7-planes	25
3.8	Type IIA with O8-planes	28
3.9	Type IIB with O9-planes: type I/Heterotic	31
4	Flux models and vacua	34
4.1	Invariant sectors	34
4.1.1	The \mathbb{Z}_2^2 invariant sector: the eight-chiral model	34
4.1.2	The $\text{SO}(3)$ invariant sector: the RSTU-model	35
4.2	Flux vacua in the eight-chiral model: type IIA with O2-planes	36
4.2.1	Fluxes and tadpoles	36
4.2.2	Mkw ₃ flux vacua	37
4.2.3	Adding O6-planes to type IIA with O2-planes	37
4.3	Flux vacua in the RSTU-model: a type IIB with O5-planes appetizer	39
4.3.1	Fluxes and tadpoles	40
4.3.2	Exploring the landscape of Mkw ₃ and AdS ₃ flux vacua	41
4.3.3	On the conformal dimension of dual operators and scale separation	46
5	Summary and future directions	49
A	Spinorial representations and Γ-matrices of $\text{SO}(8)_{\mathbb{R}}$	50
B	Gauge algebra for the type IIB with O5 vacua in Table 27	51

1 Motivation

The study of flux compactifications has been and continues being one of the most active research areas in string phenomenology. While most of the effort has been devoted to derive semi-realistic models of particle physics and cosmology in four dimensions (4D), the advent of the so-called Swampland Program [1] has broadened the interest to other dimensions. As stated in the abstract of [1]:

“The Swampland Program aims to distinguish effective theories which can be completed into quantum gravity in the ultraviolet from those which cannot.”

Since it is not an exclusive program in 4D, we will rather investigate simpler three-dimensional effective theories where some of the conjectures that have been stated in the literature might be easier to test. This motivates us to work within the context of supersymmetric gravity, *i.e.* supergravity, in three dimensions (3D). Supergravity theories in three dimensions are special in that the dreibein and the gravitino field(s) describe non-propagating degrees of freedom. All the massless propagating degrees of freedom are either spin-0 scalars or spin- $\frac{1}{2}$ fermions. Still, it is convenient to formulate three-dimensional supergravities in terms of a dreibein and the gravitino fields in order to present the theories in a manifestly covariant form [2].

Flux compactifications down to three dimensions have been considered in the literature, nonetheless, in much less detail than their four-dimensional counterparts (see [3] for a comprehensive review). Paralleling previous constructions in 4D, examples of $\mathcal{N} = 1$ (two real supercharges) flux models in 3D have been put forward in the context of type IIA [4] and type IIB [5] orientifold reductions on seven-dimensional internal spaces with G_2 -holonomy and co-calibrated G_2 -structure, respectively. The structure of three-dimensional Minkowski (Mkw₃) and anti-de Sitter (AdS₃) flux vacua was also investigated in [4, 5] (see also [6, 7]), where both supersymmetric and non-supersymmetric vacua were shown to exist with interesting phenomenological properties like full close string moduli stabilisation at scale-separated AdS₃ vacua. The small amount of supersymmetry of the $\mathcal{N} = 1$ type II orientifold models in [4, 5] allows for intricate systems of intersecting Op -planes and Dp -branes in the compactification scheme. While this enriches the structure of extrema of the scalar potential in the effective 3D supergravity arising upon compactification, it also makes an analytic study of such extrema unfeasible due to the complexity of the extremisation conditions.

A systematic and analytic study of AdS₃ vacua within the context of 3D supergravity with a larger $\mathcal{N} = 8$ supersymmetry (sixteen real supercharges or, equivalently, half-maximal) has been carried out in [8]. However, and despite the large amount of supersymmetry, a complete classification of extrema of the scalar potential of $\mathcal{N} = 8$ supergravity remains out of computational reach. Still, employing the method put forward in [9] which exploits the homogeneity of the scalar geometry in half-maximal supergravity, ref. [8] succeeded in finding all the AdS₃ vacua preserving $\mathcal{N} = 8$ supersymmetry. These are AdS₃ vacua preserving all the supersymmetries of the supergravity theories where they live. However, the string-theoretic embedding (if any) of these 3D supergravities remains generically unknown.

In this work we will combine the two approaches discussed above. We will first identify which $\mathcal{N} = 8$ (half-maximal) supergravities in three dimensions arise from type II orientifold reductions with background gauge and metric fluxes, Op -planes and Dp -branes. Demanding half-maximal supersymmetry will translate into having a *single type* of coincident Op/Dp sources, *e.g.*, type IIA with O2/D2 or type IIB with O5/D5. The formulation of the type II flux models within the embedding tensor formalism of [10, 11] (see also [8]) will allow us to carry out a systematic study of vacua using the method of [9], to compute the complete mass spectrum within half-maximal supergravity at the various vacua, and also to test some conjectures within the Swampland Program. As an application, we will provide examples of scale-separated type IIB AdS₃ flux vacua which are non-supersymmetric and perturbatively stable within half-maximal supergravity, and that come along with integer-valued conformal dimensions of the would-be dual CFT₂ operators.

The paper is organised as follows. In Section 2 we describe half-maximal gauged supergravity theories in three dimensions. We present their field content; we introduce the embedding tensor formalism and so the gauging procedure; we explicitly display the bosonic part of the complete Lagrangian and conveniently express its scalar potential in terms of the fermionic mass matrices. In Section 3 we establish a precise correspondence between single Op -plane orientifold reductions of M-theory/type II strings in the presence of background fluxes and 3D half-maximal supergravities. In particular, relying on group-theoretical arguments we build a dictionary identifying M/string-theoretic and 3D gauged supergravity quantities, *e.g.* fluxes

and embedding tensor components. Equipped with such a well-posed correspondence, in Section 4 we prepare the ground for a systematic exploration of the landscape of half-maximal gauged supergravities in three dimensions. More specifically, we discuss two interesting invariant sectors (or truncations) of the full $\mathcal{N} = 8$ 3D theory: a $\mathbb{Z}_2 \times \mathbb{Z}_2$ invariant sector and an $\text{SO}(3)$ invariant sector. After connecting the former with the type IIA with O2/O6-plane models of [4, 6, 7], we study the $\text{SO}(3)$ invariant sector (we dub it RSTU-model) for the type IIB with an O5-plane duality frame. We investigate its vacuum structure and comment on the properties of the (surprisingly) rich structure of flux vacua that we find. In Section 5 we draw our conclusions and discuss future research lines in light of the Swampland Program. A couple of technical appendices are also included.

2 A crash course on half-maximal 3D supergravity

The study of all possible half-maximal ($\mathcal{N} = 8$) supergravities in three dimensions (3D) was initiated in [10, 11] and more recently completed in [8]. Starting from the ungauged theory of [2], which features an $\text{SO}(8, n)$ global symmetry, the most general $\mathcal{N} = 8$ supergravity in 3D was constructed by applying the so-called gauging procedure. Upon this procedure, a subgroup $G \subset \text{SO}(8, n)$ is promoted from global to local. The theory then becomes a gauged supergravity, and a scalar potential and scalar-dependent mass terms for the various fermionic fields are generated by the gauging. The field content of the half-maximal gauged supergravities consists, first of all, of the $\mathcal{N} = 8$ supergravity multiplet which is composed of the dreibein $e_\mu{}^a$ and eight gravitini $\psi_\mu{}^{\mathcal{A}}$ with $\mathcal{A} = 1, \dots, 8$. The latter transform in the spinorial representation of the R-symmetry group $\text{SO}(8)_R$. In addition, the theory can also include n matter multiplets. Each matter multiplet is composed of eight real scalars $\phi^{\mathcal{I}}$ (spin-0) and eight two-component Majorana fermions $\chi^{\mathcal{A}}$ (spin- $\frac{1}{2}$) with $\mathcal{I} = 1, \dots, 8$ and $\mathcal{A} = 1, \dots, 8$ being associated with the vectorial and conjugate-spinorial representations of $\text{SO}(8)_R$. In this work we will focus on the case $n = 8$. This is the largest value for which the duality group of three-dimensional half-maximal supergravity can be embedded into the one of maximal ($\mathcal{N} = 16$) supergravity, *i.e.* $\text{SO}(8, 8) \subset \text{E}_{8(8)}$. In this case the scalars in the theory serve as coordinates in the coset space

$$\mathcal{M}_{\text{scal}} = \frac{\text{SO}(8, 8)}{\text{SO}(8) \times \text{SO}(8)}, \quad (2.1)$$

which can be viewed as a maximal subspace of the scalar geometry $\text{E}_{8(8)}/\text{SO}(16)$ of the maximal theory [2]: the scalar geometry in (2.1) comprises 64 scalars which are half of the 128 scalars in the $\text{E}_{8(8)}/\text{SO}(16)$ scalar geometry of the maximal theory.

The complete Lagrangian of the $\mathcal{N} = 8$ gauged supergravities in 3D can be found in [8]. (We closely follow the notation and conventions therein). Its bosonic part, which will be the relevant one for this work, is¹

$$\mathcal{L}_{\text{bos}} = -\frac{1}{4} e R - \frac{1}{32} e D_\mu M_{MN} D^\mu M^{MN} - e V + \mathcal{L}_{\text{CS}}. \quad (2.2)$$

The first term in (2.2) is the standard Einstein–Hilbert term for the dreibein field. The second term is a kinetic term for the scalar fields in the theory. Using the standard coset construction, the scalar-dependent matrix M_{MN} entering (2.2) is obtained from a coset representative

$$\mathcal{V}_M{}^P(\phi) \in \frac{\text{SO}(8, 8)}{\text{SO}(8) \times \text{SO}(8)} \quad (2.3)$$

¹The term $-\frac{1}{32} e D_\mu M_{MN} D^\mu M^{MN}$ in (2.2) precisely corresponds to the contribution $+\frac{1}{4} e P_\mu{}^{I r} P^{\mu I r}$ in eq. (2.12) of [8].

as $M_{MN} = \mathcal{V}_M^P \mathcal{V}_N^Q \delta_{PQ}$, where M and \underline{M} denote fundamental (vectorial) indices of $\text{SO}(8,8)$ and $\text{SO}(8) \times \text{SO}(8)$, respectively.² The third term in (2.2) is a scalar potential whose explicit expression will be given in a moment. Finally, the last term in (2.2) denotes a topological term for the vector fields $A_\mu^{MN} = A_\mu^{[MN]}$ in the theory which formally transform in the **120** (adjoint) irreducible representation (irrep) of $\text{SO}(8,8)$. Note that in three dimensions vectors are dual to scalars and therefore do not carry an independent dynamics (see [8] for more details).

2.1 Gaugings, embedding tensor and quadratic constraints

The gaugings of half-maximal supergravity are encoded in the so-called embedding tensor

$$\Theta_{MN|PQ} = \theta_{MNPQ} + 2(\eta_{M[P}\theta_{Q]N} - \eta_{N[P}\theta_{Q]M}) + 2\eta_{M[P}\eta_{Q]N}\theta, \quad (2.4)$$

which consists of three irreducible representations (irreps) of the duality group $\text{SO}(8,8)$: $\theta_{MNPQ} = \theta_{[MNPQ]} \in \mathbf{1820}$, $\theta_{MN} = \theta_{(MN)} \in \mathbf{135}$ (with $\theta_M^M = 0$) and $\theta \in \mathbf{1}$. The η_{MN} matrix in (2.4) is the non-degenerate $\text{SO}(8,8)$ invariant matrix that is used to raise/lower vectorial indices. In this work we will alternate between two different basis for $\text{SO}(8,8)$: the light-cone (LC) basis and the Lorentzian (Ltz) basis. The η_{MN} matrix is

$$\eta_{MN}\Big|_{\text{LC}} = \begin{pmatrix} 0 & \mathbb{I} \\ \mathbb{I} & 0 \end{pmatrix} \quad \text{and} \quad \eta_{MN}\Big|_{\text{Ltz}} = \begin{pmatrix} -\mathbb{I} & 0 \\ 0 & \mathbb{I} \end{pmatrix}, \quad (2.5)$$

in the two different basis. While the LC basis will be relevant to establish the dictionary between the embedding tensor and the various type II flux parameters, the Lorentzian basis will be more adequate to compute the fermionic mass terms and, from them, the scalar potential in (2.2). Both basis are related by the $\text{SO}(16)$ rotation

$$\eta_{MN}\Big|_{\text{Ltz (LC)}} = R_M^P R_N^Q \eta_{PQ}\Big|_{\text{LC (Ltz)}} \quad \text{with} \quad R_M^N = \frac{1}{\sqrt{2}} \begin{pmatrix} -\mathbb{I} & \mathbb{I} \\ \mathbb{I} & \mathbb{I} \end{pmatrix}. \quad (2.6)$$

The subgroup $G \subset \text{SO}(8,8)$ of the duality group that is gauged (and then becomes a local symmetry of the theory) upon the gauging procedure has generators of the form

$$X_{MN} = \Theta_{MN|PQ} L^{PQ}, \quad (2.7)$$

with

$$(L^{PQ})_R^T = 2\delta_R^{[P} \eta^{Q]T} \quad \text{and} \quad (L^{PQ})_{RS}{}^{TU} = 8\delta_{[R}^{[P} \eta^{Q][T} \delta_S^{U]}, \quad (2.8)$$

being the generators of $\text{SO}(8,8)$ in the vectorial and adjoint representations, respectively. Using (2.8), the gauge group generators (2.7) are

$$X_{MNR}{}^T \equiv \Theta_{MN|PQ} (L^{PQ})_R^T \quad \text{and} \quad X_{MNR}{}^{ST} \equiv \Theta_{MN|PQ} (L^{PQ})_{RS}{}^{TU}, \quad (2.9)$$

in the fundamental (vectorial) and adjoint representations, respectively. The fundamental representation enters the covariant derivatives of the scalars in (2.2), namely,

$$D_\mu M_{MN} = \partial_\mu M_{MN} - 2g A_\mu^{PQ} X_{PQ(M}{}^R M_{N)R}, \quad (2.10)$$

where g denotes the gauge coupling in the supergravity. The adjoint representation determines the gauge brackets

$$[X_{MN}, X_{PQ}] = -\frac{1}{2} X_{MNPQ}{}^{RS} X_{RS}, \quad (2.11)$$

²In our conventions, the coset representative $\mathcal{V}_M^N(\phi)$ transforms under global $\text{SO}(8,8)$ transformations from the left and under local $\text{SO}(8) \times \text{SO}(8)$ transformations from the right (underlined index).

and closure of the gauge brackets (2.11) imposes a set of quadratic constraints (QC's) of the form

$$X_{RSM}{}^K \Theta_{KN|PQ} + X_{RSN}{}^K \Theta_{MK|PQ} + X_{RSP}{}^K \Theta_{MN|KQ} + X_{RSQ}{}^K \Theta_{MN|PK} = 0 . \quad (2.12)$$

Each embedding tensor configuration solving the QC's in (2.12) gives rise to a consistent half-maximal gauged supergravity in three dimensions. Lastly, as discussed in Section 6 of [8], the set of QC's in (2.12) must be supplemented with two additional ones, namely

$$\begin{aligned} 48 \theta \theta_{MN} + \theta_M{}^{PQR} \theta_{NPQR} - \frac{1}{16} \eta_{MN} \theta^{PQRS} \theta_{PQRS} &= 0 , \\ \theta_{M_1 M_2 M_3 M_4} \theta_{M_5 M_6 M_7 M_8} [\Gamma^{M_1 \dots M_8}]_{\dot{A} \dot{B}} &= 0 , \end{aligned} \quad (2.13)$$

where $[\Gamma^{M_1 \dots M_8}]_{\dot{A} \dot{B}}$ denotes the (self-dual) eight-fold antisymmetric product of Γ -matrices of $\text{SO}(8, 8)$, for a half-maximal supergravity to be embeddable into the maximal theory. We will come back to these additional constraints when analysing gaugings of half-maximal supergravity arising from M-theory/type II/type I flux compactifications.

2.2 Fermion masses and scalar potential

The gauging procedure generates scalar-dependent mass terms for the various fermionic fields in the theory [8]. These take the form

$$\mathcal{L}_{\text{mass}} = \frac{1}{2} e A_1^{AB} \bar{\psi}_\mu{}^{\mathcal{A}} \gamma^{\mu\nu} \psi_\nu{}^{\mathcal{B}} + i e A_2^{A\dot{B}s} \bar{\chi}^{\dot{B}s} \gamma^\mu \psi_\mu{}^{\mathcal{A}} + \frac{1}{2} e A_3^{A\dot{r}\dot{B}s} \bar{\chi}^{\dot{A}r} \chi^{\dot{B}s} , \quad (2.14)$$

where the index $r = 1, \dots, n$ runs over the $n = 8$ matter multiplets and the indices $\mathcal{A}, \dot{\mathcal{A}}$ refer, together with \mathcal{I} , to the (trianality-related) spinorial, conjugate-spinorial and vectorial representations of $\text{SO}(8)_R$. To obtain the various fermionic mass terms in (2.14) we will proceed as follows. Using the Lorentzian basis in which η_{MN} takes the block-diagonal form in (2.5) and the $\text{SO}(8, 8)$ vectorial index splits as $M = (\mathcal{I}, r)$, we will start by building the coset representative as

$$\mathcal{V}_M{}^N(\phi) = e^{\phi_{\mathcal{I}r}} (L^{\mathcal{I}r})_M{}^N , \quad (2.15)$$

where $\phi_{\mathcal{I}r}$ are the 64 physical scalars associated with the non-compact generators $L^{\mathcal{I}r}$. Note that, amongst the L^{MN} generators in (2.8), those of type $L^{\mathcal{I}\mathcal{J}}$ and L^{rs} span the compact $\text{SO}(8) \times \text{SO}(8) \subset \text{SO}(8, 8)$ in the denominator of (2.3) and, therefore, have no physical scalars associated. Equipped with the coset representative (2.15) in the Lorentzian basis, we will dress up the embedding tensor components in (2.4) to obtain the so-called T -tensor components³

$$\begin{aligned} T_{\underline{MNPQ}} &= (\mathcal{V}^{-1})_{\underline{M}}{}^R (\mathcal{V}^{-1})_{\underline{N}}{}^S (\mathcal{V}^{-1})_{\underline{P}}{}^T (\mathcal{V}^{-1})_{\underline{Q}}{}^U \theta_{RSTU} , \\ T_{\underline{MN}} &= (\mathcal{V}^{-1})_{\underline{M}}{}^R (\mathcal{V}^{-1})_{\underline{N}}{}^S \theta_{RS} , \\ T &= \theta . \end{aligned} \quad (2.16)$$

Finally, the fermionic mass terms in (2.14) can be extracted from (2.16) upon taking suitable projections into irreps of the R-symmetry group using $\text{SO}(8)_R$ invariant $\gamma^{(p)}$ -forms (see Appendix A). The result is

$$\begin{aligned} A_1^{AB} &= -\frac{1}{48} [\gamma^{\mathcal{I}\mathcal{J}\mathcal{K}\mathcal{L}}]_{AB} T_{\underline{\mathcal{I}\mathcal{J}\mathcal{K}\mathcal{L}}} - \frac{1}{4} \delta^{AB} (T_{\underline{\mathcal{I}\mathcal{I}}} - 4T) , \\ A_2^{A\dot{B}s} &= -\frac{1}{12} [\gamma^{\mathcal{I}\mathcal{J}\mathcal{K}}]_{A\dot{B}} T_{\underline{\mathcal{I}\mathcal{J}\mathcal{K}s}} - \frac{1}{2} [\gamma^{\mathcal{I}}]_{A\dot{B}} T_{\underline{\mathcal{I}s}} , \\ A_3^{A\dot{r}\dot{B}s} &= \frac{1}{48} \delta^{rs} [\gamma^{\mathcal{I}\mathcal{J}\mathcal{K}\mathcal{L}}]_{\dot{A}\dot{B}} T_{\underline{\mathcal{I}\mathcal{J}\mathcal{K}\mathcal{L}}} + \frac{1}{2} [\gamma^{\mathcal{I}\mathcal{J}}]_{\dot{A}\dot{B}} T_{\underline{\mathcal{I}\mathcal{J}rs}} - 2 \delta^{A\dot{B}} \left[\delta^{rs} \left(T - \frac{1}{8} T_{\underline{qq}} \right) + T_{\underline{rs}} \right] . \end{aligned} \quad (2.17)$$

³Importantly, the non-singlet embedding tensor components θ_{MNPQ} and θ_{MN} must be given in the Lorentzian basis too for the contractions in (2.16) to be well-defined.

In terms of the above fermionic mass terms, the scalar potential of the gauged supergravity can be expressed as

$$V(\phi) = -\frac{1}{4}A_1^{AB}A_1^{AB} + \frac{1}{8}A_2^{A\dot{B}s}A_2^{A\dot{B}s} . \quad (2.18)$$

Selecting a particular embedding tensor configuration $\Theta_{MN|PQ}$ in (2.4) that solves the QC's in (2.12) amounts to select a particular gauged supergravity, namely, a particular theory. Then, the extremisation of the corresponding scalar potential (2.18) yields the structure of maximally symmetric vacua, *i.e.* Anti de Sitter (AdS₃), Minkowski (Mkw₃) or de Sitter (dS₃), of such a theory. Once a vacuum has been found, the spectrum of fermions around that vacuum is obtained directly from (2.17). The spectrum of vectors and scalars can be computed from the mass matrices⁴ in eqs. (2.22) – (2.23) of [8].

3 Type II orientifolds

Type II orientifolds have been studied in [12] within the context of supersymmetric compactifications. The orientifold action is a composition of both worldsheet and target space transformations. In particular, a worldsheet orientation-reversal parity transformation Ω_P , a worldsheet fermion number projector for left-moving fermions $(-1)^{F_L}$ (which is not always needed [12]) and an internal target space involution σ . The internal involution σ may leave certain submanifolds of the internal space invariant. The product of one of such invariant submanifolds with the external spacetime is referred to as an orientifold Op -plane, where p denotes the total number of spatial dimensions filled by the orientifold plane.

Op -planes have negative tension, are electrically (magnetically) charged under a $C_{(p+1)}$ ($C_{(7-p)}$) gauge potential and do not have an associated dynamics. Being charged objects, they enter charge conservation conditions known as tadpole cancellation conditions (we refer the reader to [12] for more details) of the form

$$DF_{(8-p)} + \beta_{8-p} H_{(1)} \wedge F_{(8-p)} - H_{(3)} \wedge F_{(6-p)} = J_{Op/Dp} , \quad (3.1)$$

where β_{8-p} is a constant, $D \equiv d + \omega$ is a nilpotent ($D^2 = 0$) twisted exterior derivative on the internal space and $J_{Op/Dp}$ denotes the net current of charged Op -plane/ Dp -brane sources in the smeared limit. Similarly to (3.1), and in the absence of NS5-branes, there is a condition for the NS-NS three-form flux of the form

$$DH_{(3)} + \alpha H_{(1)} \wedge H_{(3)} = 0 , \quad (3.2)$$

with α being a constant. The parameters β_p in (3.1) are not independent. In particular, one has that $\beta_p = \beta - \frac{4-p}{2}\alpha$ in the type IIA context and $\beta_p = \beta - \frac{5-p}{2}\alpha$ in the type IIB one for arbitrary parameters α and β . Then, as we will show, the matching between the tadpole conditions (3.1)-(3.2) and the quadratic constraints (2.12) of the gauged supergravity requires an (α, β) -dependent embedding of the dilaton flux $H_{(1)}$ into the embedding tensor. The attentive reader might have noticed the unconventional terms $H_{(1)} \wedge F_{(8-p)}$ and $H_{(1)} \wedge H_{(3)}$ in (3.1) and (3.2), respectively. Starting from a standard formulation of type II supergravity in the string frame, these terms can be generated by performing non-standard field redefinitions $F_{(p)} \rightarrow e^{-\beta_p \Phi} F_{(p)}$ and $H_{(3)} \rightarrow e^{-\alpha \Phi} H_{(3)}$ involving different powers of the type II dilaton e^Φ .

⁴These mass matrices were obtained assuming that the vacuum solution is placed at the origin of moduli space, namely, at $\mathcal{V}_M^{\mathcal{N}}(0) = \mathbb{I}$. Note that this does not imply a lack of generality in the context of 3D gauged supergravity since the coset space (2.3) is a homogeneous space and, therefore, a vacuum at $\mathcal{V}_M^{\mathcal{N}}(\phi) \neq \mathbb{I}$ can be brought to the origin of moduli space by acting with a $SO(8, 8)$ transformation. Also notice that the original embedding tensor $\Theta_{MN|PQ}$ yielding the vacuum outside the origin must be transformed accordingly for the T -tensor (2.16) and, therefore, the scalar potential (2.18) and the mass spectra, to remain invariant.

The need for this unconventional frame stems from the fact that, when considering type IIA models, the metric fluxes ω and the dilaton flux $H_{(1)}$ must be treated on equal footing as they both come from metric fluxes in 11D. The parameters α and β will determine the embedding of $H_{(1)}$ inside the embedding tensor of the three-dimensional gauged supergravities.⁵ If $H_{(1)} = 0$ one can set $\alpha = \beta = 0$ and recover the standard string frame.

The tadpole cancellation conditions in (3.1) and (3.2) involve metric fluxes ω specifying the twist of the geometry, *i.e.* a twisted \mathbb{T}^7 torus, as well as gauge fluxes $H_{(3)}$ and $F_{(p)}$, and a dilaton flux $H_{(1)}$. However, the coexistence of metric ω and one-form $H_{(1)}$ or $F_{(1)}$ fluxes in a compactification scheme poses some issues. On the one hand, metric fluxes ω arise from Scherk–Schwarz (SS) reductions on twisted tori [13]. These fluxes have a geometric origin and can be thought of as operators that map an internal p -form into a $(p + 1)$ -form, *e.g.* $(DF)_{mnpq} = \partial_{[m} F_{npq]} + \omega_{[mn}{}^r F_{r|pq]}$, and specify the twist of fields (gauge potentials) with legs along the internal space. Since the dilaton Φ is a scalar (no legs along the internal \mathbb{T}^7), a dilaton flux $H_{(1)}$ cannot be generated by the ordinary SS procedure. The same holds for a flux $F_{(1)}$. On the other hand, $H_{(1)}$ and $F_{(1)}$ background fluxes are perfectly valid ingredients in a standard flux compactification on a \mathbb{T}^7 torus. Therefore, we are left with two possible well-defined scenarios:

i) Ordinary Scherk–Schwarz reduction. In this scenario one has

$$\omega_{mn}{}^p \neq 0 \quad \text{and} \quad H_{(1)} = F_{(1)} = 0 . \quad (3.3)$$

Since the metric fluxes ω are identified with the structure constants of the Lie group specifying the twist, they must obey standard Jacobi identities of the form

$$\omega_{[mn}{}^r \omega_p]r{}^q = 0 , \quad (3.4)$$

together with an additional “unimodularity” or trace condition [13]

$$\omega_{mr}{}^r = 0 . \quad (3.5)$$

In ordinary SS reductions, the unimodularity condition (3.5) ensures consistency of the truncation of the higher-dimensional action. For non-unimodular gaugings there is still a consistent truncation of the equations of motion but, in general, not of the action [14, 15].⁶

ii) Standard flux compactification. In this scenario one has

$$\omega_{mn}{}^p = 0 \quad \text{and} \quad H_{(1)}, F_{(1)} \neq 0 . \quad (3.6)$$

In what follows we will consider all the fluxes at the same time when discussing the fluxes/embedding tensor dictionary, quadratic constraints on fluxes, etc., but always bearing in mind that one of the two scenarios above must hold for a given three-dimensional gauged supergravity to describe a truncation of a type II supergravity action. One could of course adopt a bottom-up approach and activate *any* embedding tensor component associated with a (generically non-unimodular) metric flux ω as well as gauge fluxes $H_{(1)}$, $H_{(3)}$ and $F_{(p)}$

⁵It will also occur that the matching between the tadpole conditions (3.1) and (3.2) and the quadratic constraints of the gauged supergravity (2.12) will allow for additional parameters, we denote them γ and δ , in the embedding of the dilaton flux $H_{(1)}$ inside the embedding tensor. This ambiguity would ultimately be fixed by performing a full-fledged dimensional reduction of type II supergravity in presence of $H_{(1)}$.

⁶An example of a non-unimodular gauging still describing a consistent truncation at the level of the action can be found in [16]. However, this counterexample to the general statement includes an additional duality twist (see *e.g.* [17]) that introduces global pathologies in the higher-dimensional background.

	$E_{8(8)} / \text{Maximal}$	$SO(8, 8)$	Half-Maximal	$SL(8) \times \mathbb{R}_1$
	–	16	–	$\mathbf{8}'_{-1} \oplus \mathbf{8}_{+1}$
Scalars	248	120	L^{MN}	$\mathbf{28}'_{-2} \oplus (\mathbf{63} + \mathbf{1})_0 \oplus \mathbf{28}_{+2}$
		128'	–	$\mathbf{8}_{-3} \oplus \mathbf{56}_{-1} \oplus \mathbf{56}'_{+1} \oplus \boxed{\mathbf{8}'_{+3}}$
Embedding Tensor / Fluxes	1	1	θ	$\mathbf{1}_0$
	3875	135	θ_{MN}	$\mathbf{36}'_{-2} \oplus \mathbf{63}_0 \oplus \mathbf{36}_{+2}$
		1820	θ_{MNPQ}	$\mathbf{70}_{-4} \oplus \mathbf{28}'_{-2} \oplus \mathbf{420}'_{-2}$ $\mathbf{720}_0 \oplus \mathbf{63}_0 \oplus \mathbf{1}_0$ $\mathbf{70}_{+4} \oplus \mathbf{28}_{+2} \oplus \mathbf{420}_{+2}$
		1920'	–	$\mathbf{8}'_{-5} \oplus \mathbf{216}_{-3} \oplus \mathbf{8}_{-3}$ $\mathbf{56}_{-1} \oplus \mathbf{504}_{-1} \oplus \mathbf{168}_{-1}$ $\mathbf{56}'_{+1} \oplus \mathbf{504}'_{+1} \oplus \mathbf{168}'_{+1}$ $\mathbf{8}_{+5} \oplus \mathbf{216}'_{+3} \oplus \mathbf{8}'_{+3}$

Table 1: M-theory branching rules for the embedding chain $E_{8(8)} \supset SO(8, 8) \supset SL(8) \times \mathbb{R}_1$. The subscripts in the third column indicate \mathbb{R}_1 -charges. We have highlighted the scalars $e_A{}^B$, $C_{(3)}$ and $C_{(6)}$ in (3.8) as well as their associated fluxes $\omega_{AB}{}^C$, $G_{(4)}$ and $G_{(7)}$ in (3.9). Also the physical internal derivatives ∂_A have been put in a box for their quick identification. Note that the spinorial representations $\mathbf{128}'$ and $\mathbf{1920}'$ of $SO(8, 8)$ give rise to $SL(8) \times \mathbb{R}_1$ irreps with an odd \mathbb{R}_1 -charge. These irreps are projected out when truncating maximal to half-maximal supergravity and are marked with “–” as they do not give rise to scalars or embedding tensor deformations in half-maximal supergravity.

provided the quadratic constraints (2.11) hold. However, the uplift of the three-dimensional gauged supergravity described by such an embedding tensor would generically deviate from an ordinary SS reduction or a standard flux compactification. These more exotic scenarios are beyond the scope of this work.

3.1 Warming up: M-theory fluxes/embedding tensor dictionary

In the absence of sources, the dimensional reduction of 11D supergravity on \mathbb{T}^8 yields the ungauged 3D supergravity with $E_{8(8)}$ global U-duality symmetry [2]. This $E_{8(8)}$ symmetry group becomes manifest when 11D supergravity is reformulated in the form of an exceptional field theory living in a generalised $(3 + 248)$ -dimensional spacetime with an extended set of internal coordinates in the $\mathbf{248}$ (adjoint) irrep of $E_{8(8)}$ [18]. The eight physical internal coordinates of 11D supergravity are then part of the $\mathbf{248} \in E_{8(8)}$. On the other hand, standard Kaluza–Klein (KK) reduction of 11D supergravity on \mathbb{T}^8 comes along with a global $GL(8) = SL(8) \times \mathbb{R}_1$ symmetry descending from the ordinary internal diffeomorphisms. This

implies that the eight physical internal coordinates of 11D supergravity y^A , with $A = 1, \dots, 8$, must transform as an eight-dimensional irrep of $\text{SL}(8)$. A quick inspection of Table 1 then confirms that, up to a conventional choice, one has that

$$y^A \in \mathbf{8}_{-3} \quad \text{and} \quad \partial_A \in \mathbf{8}'_{+3} . \quad (3.7)$$

After having group-theoretically identified the physical derivatives ∂_A , we proceed to identify the internal components of the elfbein $e_A{}^B$ as well as the three-form $C_{(3)} \equiv C_{ABC} = C_{[ABC]}$ and six-form $C_{(6)} \equiv C_{A_1 \dots A_6} = C_{[A_1 \dots A_6]}$ gauge potentials of 11D supergravity. An inspection of Table 1 reveals that

$$e_A{}^B \in (\mathbf{63} + \mathbf{1})_0 \quad , \quad C_{(3)} \in \mathbf{56}'_{+1} \quad \text{and} \quad C_{(6)} \in \mathbf{28}_{+2} , \quad (3.8)$$

where 28 compact scalars must be subtracted from $e_A{}^B$ upon gauge-fixing of the internal $\text{SO}(8)$ local symmetry, namely, $e_A{}^B \in \text{GL}(8)/\text{SO}(8)$. Lastly, metric and gauge fluxes are constructed by applying exterior derivatives on the scalars in (3.8). This yields

$$\omega_{AB}{}^C \in (\mathbf{216}' + \mathbf{8}')_{+3} \quad , \quad G_{(4)} \in \mathbf{70}_{+4} \quad \text{and} \quad G_{(7)} \in \mathbf{8}_{+5} . \quad (3.9)$$

Importantly, bosonic (spinorial) irreps of $\text{SO}(8, 8)$ give rise to $\text{SL}(8) \times \mathbb{R}_1$ irreps with an even (odd) \mathbb{R}_1 -charge and are kept (projected out) when truncating maximal to half-maximal supergravity (see Table 1). Therefore, since the physical internal derivatives in (3.7) are spinorial in this sense (\mathbb{R}_1 -charge +3), only the spinorial scalars $C_{(3)}$ (\mathbb{R}_1 -charge +1) in (3.8) will produce a bosonic flux $G_{(4)}$ (\mathbb{R}_1 -charge +4) in (3.9) suitable to enter the embedding tensor of half-maximal supergravity. Summarising, the half-maximal supergravities obtained from M-theory will contain physical scalars and gauge fluxes of the form

$$\begin{aligned} \text{Scalars} & : e_A{}^B \in \frac{\text{GL}(8)}{\text{SO}(8)} \quad , \quad C_{(6)} , \\ \text{Fluxes} & : G_{(4)} . \end{aligned} \quad (3.10)$$

This simple spinorial/bosonic grading will help us later when discussing more complicated flux models arising from type II orientifold reductions.

Before moving to discuss specific algebraic properties of the M-theory gaugings, let us perform a precise counting of bosonic and spinorial scalars in the M-theory context. As we have already seen, bosonic scalars arise from $e_A{}^B \in \text{GL}(8)/\text{SO}(8)$ and $C_{(6)}$ adding up to 64 scalars. However, there are only 56 spinorial scalars arising from $C_{(3)}$. The difference is explained by the 8 additional spinorial scalars dual to the vectors $e_\mu{}^A$. All together, there are 64 bosonic and 64 spinorial scalars as required by the $\text{E}_{8(8)}/\text{SO}(16)$ scalar geometry of the maximal theory.

Since half-maximal supergravity does not contain spinorial irreps of $\text{SO}(8, 8)$, only the gauge flux $G_{(4)}$ and the 64 bosonic scalars $e_A{}^B \in \text{GL}(8)/\text{SO}(8)$ and $C_{(6)}$ (all of them with an even \mathbb{R}_1 -charge) are present in the theory. The resulting gauging in 3D has a simple embedding tensor (2.4) specified by the non-zero components

$$\theta^{ABCD} = \frac{1}{4!} \varepsilon^{ABCDEFGH} G_{EFGH} , \quad (3.11)$$

where we have made the $\text{SO}(8, 8)$ index splitting $T_M = (T_A, T^A)$ in light-cone coordinates. This is nothing but the branching $\mathbf{16} \rightarrow \mathbf{8}'_{-1} \oplus \mathbf{8}_{+1}$ under $\text{SO}(8, 8) \supset \text{SL}(8) \times \mathbb{R}_1$ in Table 1. The embedding tensor (3.11) satisfies the QC's in (2.12) and specifies an Abelian gauge group of dimension 28 for the non-zero generators $X^{AB} = \theta^{ABCD} L_{CD}$ in the light-cone basis.⁷

⁷In the light-cone basis there are two maximal Abelian subgroups spanned by L^{AB} and L_{AB} , respectively.

	Half-Maximal	$SL(8) \times \mathbb{R}_1$	$SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$
	–	$\mathbf{8}'_{-1} \oplus \mathbf{8}_{+1}$	$[\mathbf{7}'_{(-1,-1)} \oplus \mathbf{1}_{(+7,-1)}] \oplus [\mathbf{7}_{(+1,+1)} \oplus \mathbf{1}_{(-7,+1)}]$
Scalars	L^{MN}	$\mathbf{28}'_{-2} \oplus (\mathbf{63} + \mathbf{1})_0 \oplus \mathbf{28}_{+2}$	$[\mathbf{21}'_{(-2,-2)} \oplus \mathbf{7}'_{(+6,-2)}] \oplus [\mathbf{7}'_{(-8,0)} \oplus (\mathbf{48} + \mathbf{1})_{(0,0)} \oplus \mathbf{7}_{(+8,0)}]$ $\mathbf{1}_{(0,0)} \oplus [\mathbf{7}_{(-6,+2)} \oplus \mathbf{21}_{(+2,+2)}]$
	–	$\mathbf{8}_{-3} \oplus \mathbf{56}_{-1} \oplus \mathbf{56}'_{+1} \oplus \boxed{\mathbf{8}'_{+3}}$	$[\mathbf{7}_{(+1,-3)} \oplus \mathbf{1}_{(-7,-3)}] \oplus [\mathbf{35}_{(+3,-1)} \oplus \mathbf{21}_{(-5,-1)}]$ $[\mathbf{35}'_{(-3,+1)} \oplus \mathbf{21}'_{(+5,+1)}] \oplus [\boxed{\mathbf{7}'_{(-1,+3)}} \oplus \mathbf{1}_{(+7,+3)}]$
Embedding Tensor / Fluxes	θ	$\mathbf{1}_0$	$\mathbf{1}_{(0,0)}$
	θ_{MN}	$\mathbf{36}'_{-2} \oplus \mathbf{63}_0 \oplus \mathbf{36}_{+2}$	$[\mathbf{28}'_{(-2,-2)} \oplus \mathbf{7}'_{(+6,-2)} \oplus \mathbf{1}_{(+14,-2)}]$ $[\mathbf{7}'_{(-8,0)} \oplus \mathbf{48}_{(0,0)} \oplus \mathbf{1}_{(0,0)} \oplus \mathbf{7}_{(+8,0)}]$ $[\mathbf{28}_{(+2,+2)} \oplus \mathbf{7}_{(-6,+2)} \oplus \mathbf{1}_{(-14,+2)}]$
	θ_{MNPQ}	$\mathbf{70}_{-4} \oplus \mathbf{28}'_{-2} \oplus \mathbf{420}'_{-2}$	$[\mathbf{35}_{(-4,-4)} \oplus \mathbf{35}'_{(+4,-4)}] \oplus [\mathbf{21}'_{(-2,-2)} \oplus \mathbf{7}'_{(+6,-2)}]$ $[\mathbf{35}'_{(-10,-2)} \oplus \mathbf{21}'_{(-2,-2)} \oplus \mathbf{224}'_{(-2,-2)} \oplus \mathbf{140}'_{(+6,-2)}]$
		$\mathbf{720}_0 \oplus \mathbf{63}_0 \oplus \mathbf{1}_0$	$[\mathbf{140}'_{(-8,0)} \oplus \mathbf{392}_{(0,0)} \oplus \mathbf{48}_{(0,0)} \oplus \mathbf{140}_{(+8,0)}]$ $[\mathbf{7}'_{(-8,0)} \oplus \mathbf{48}_{(0,0)} \oplus \mathbf{1}_{(0,0)} \oplus \mathbf{7}_{(+8,0)}] \oplus \mathbf{1}_{(0,0)}$
		$\mathbf{70}_{+4} \oplus \mathbf{28}_{+2} \oplus \mathbf{420}_{+2}$	$[\mathbf{35}_{(-4,+4)} \oplus \mathbf{35}'_{(+4,+4)}] \oplus [\mathbf{7}_{(-6,+2)} \oplus \mathbf{21}_{(+2,+2)}]$ $[\mathbf{35}_{(+10,+2)} \oplus \mathbf{21}_{(+2,+2)} \oplus \mathbf{224}_{(+2,+2)} \oplus \mathbf{140}_{(-6,+2)}]$
	–	$\mathbf{8}'_{-5} \oplus \mathbf{8}_{-3} \oplus \mathbf{216}_{-3}$	$[\mathbf{7}'_{(-1,-5)} \oplus \mathbf{1}_{(+7,-5)}] \oplus [\mathbf{1}_{(-7,-3)} \oplus \mathbf{7}_{(+1,-3)}]$ $[\mathbf{48}_{(-7,-3)} \oplus \mathbf{140}_{(+1,-3)} \oplus \mathbf{7}_{(+1,-3)} \oplus \mathbf{21}_{(+9,-3)}]$
		$\mathbf{56}_{-1} \oplus \mathbf{504}_{-1} \oplus \mathbf{168}_{-1}$	$[\mathbf{21}_{(-5,-1)} \oplus \mathbf{35}_{(+3,-1)}]$ $[\mathbf{224}_{(-5,-1)} \oplus \mathbf{210}_{(+3,-1)} \oplus \mathbf{35}_{(+3,-1)} \oplus \mathbf{35}'_{(+11,-1)}]$ $[\mathbf{7}_{(-13,-1)} \oplus \mathbf{21}_{(-5,-1)} \oplus \mathbf{28}_{(-5,-1)} \oplus \mathbf{112}_{(+3,-1)}]$
		$\mathbf{56}'_{+1} \oplus \mathbf{504}'_{+1} \oplus \mathbf{168}'_{+1}$	$[\mathbf{21}'_{(+5,+1)} \oplus \mathbf{35}'_{(-3,+1)}]$ $[\mathbf{224}'_{(+5,+1)} \oplus \mathbf{210}'_{(-3,+1)} \oplus \mathbf{35}'_{(-3,+1)} \oplus \mathbf{35}_{(-11,+1)}]$ $[\mathbf{7}'_{(+13,+1)} \oplus \mathbf{21}'_{(+5,+1)} \oplus \mathbf{28}'_{(+5,+1)} \oplus \mathbf{112}'_{(-3,+1)}]$
		$\mathbf{8}_{+5} \oplus \mathbf{8}'_{+3} \oplus \mathbf{216}'_{+3}$	$[\mathbf{7}_{(+1,+5)} \oplus \mathbf{1}_{(-7,+5)}] \oplus [\mathbf{7}'_{(-1,+3)} \oplus \mathbf{1}_{(+7,+3)}]$ $[\mathbf{48}_{(+7,+3)} \oplus \mathbf{140}'_{(-1,+3)} \oplus \mathbf{7}'_{(-1,+3)} \oplus \mathbf{21}'_{(-9,+3)}]$

Table 2: Type IIA branching rules for the embedding $SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1 \subset SL(8) \times \mathbb{R}_1$. The subscripts in the third column indicate $(\mathbb{R}_2, \mathbb{R}_1)$ -charges. We have highlighted the scalars e_m^n , Φ , $B_{(2)}$, $B_{(6)}$ and $C_{(p)}$ in (3.13), as well as their associated fluxes in (3.14). The Romans mass parameter $F_{(0)} \in \mathbf{1}_{(-14,+2)}$ has also been highlighted and the physical internal derivatives ∂_m have been put in a box for their quick identification. This table should be understood as a continuation of Table 1.

3.2 Type IIA with O2-planes

The M-theory reduction on $\mathbb{T}^8 = \mathbb{T}^7 \times \mathbb{S}^1$ can be reinterpreted as a (massless) type IIA reduction on \mathbb{T}^7 with the coordinate along the \mathbb{S}^1 being the M-theory coordinate. The $GL(8)$ diffeomorphisms on \mathbb{T}^8 are then reduced to the $GL(7)$ diffeomorphisms on \mathbb{T}^7 and coordinates

split as $y^A = (y^m, y^8)$ with $m = 1, \dots, 7$. The relevant branching rules for the M-theory \Leftrightarrow type IIA with O2 correspondence are summarised in Table 2.

The seven physical coordinates and their associated derivatives follow from the branching rules in Table 2. They are identified with

$$y^m \in \mathbf{7}_{(+1,-3)} \quad \text{and} \quad \partial_m \in \mathbf{7}'_{(-1,+3)}. \quad (3.12)$$

On the other hand, the type IIA dilaton Φ , the internal components of the zehnbein $e_m{}^n$ and the various type IIA gauge potentials $B_{(2)}$, $B_{(6)}$ and $C_{(1)}$, $C_{(3)}$, $C_{(5)}$ are identified as

$$\begin{aligned} e_m{}^n &\in (\mathbf{48} + \mathbf{1})_{(0,0)} \quad , \quad C_{(1)} \in \mathbf{7}'_{(-8,0)} \quad , \quad \Phi \in \mathbf{1}_{(0,0)} \quad , \\ C_{(3)} &\in \mathbf{35}'_{(-3,+1)} \quad , \quad B_{(2)} \in \mathbf{21}'_{(+5,+1)} \quad , \quad B_{(6)} \in \mathbf{7}_{(-6,+2)} \quad , \quad C_{(5)} \in \mathbf{21}_{(+2,+2)} \quad , \end{aligned} \quad (3.13)$$

where 21 compact scalars must be subtracted from $e_m{}^n$ upon gauge-fixing of the internal $\text{SO}(7)$ local symmetry, namely, $e_m{}^n \in \text{GL}(7)/\text{SO}(7)$. While the bosonic scalars in (3.13) correctly add up to 64, the number of the spinorial ones turns out to be 56. The 7 + 1 missing spinorial scalars are dual to the three-dimensional vectors $e_\mu{}^n$ and C_μ .⁸ The metric and gauge fluxes obtained by applying exterior derivatives on the scalars in (3.13) are

$$\begin{aligned} \omega_{mn}{}^p &\in (\mathbf{140}' + \mathbf{7}')_{(-1,+3)} \quad , \quad F_{(2)} \in \mathbf{21}'_{(-9,+3)} \quad , \quad H_{(1)} \in \mathbf{7}'_{(-1,+3)} \quad , \\ F_{(4)} &\in \mathbf{35}_{(-4,+4)} \quad , \quad H_{(3)} \in \mathbf{35}'_{(+4,+4)} \quad , \quad H_{(7)} \in \mathbf{1}_{(-7,+5)} \quad , \quad F_{(6)} \in \mathbf{7}_{(+1,+5)}. \end{aligned} \quad (3.14)$$

Finally, the massive version of type IIA supergravity admits an additional deformation parameter known as the Romans mass $F_{(0)}$ [19]. This parameter does not follow from a gauge potential and it is group-theoretically identified with

$$F_{(0)} \in \mathbf{1}_{(-14,+2)}. \quad (3.15)$$

Again, since half-maximal supergravity does not contain spinorial irreps of $\text{SO}(8,8)$, we must keep scalars and fluxes with an even \mathbb{R}_1 -charge. The result is that only the 64 physical scalars and gauge fluxes

$$\begin{aligned} \text{Scalars} &: e_m{}^n \in \frac{\text{GL}(7)}{\text{SO}(7)} \quad , \quad C_{(1)} \quad , \quad \Phi \quad , \quad B_{(6)} \quad , \quad C_{(5)} \quad , \\ \text{Fluxes} &: H_{(3)} \quad , \quad F_{(4)} \quad , \quad F_{(0)} \quad , \end{aligned} \quad (3.16)$$

are present in the theory. These fluxes induce a gauging in 3D specified by the embedding tensor components

$$\theta^{mnpq} = \frac{1}{3!} \epsilon^{mnpqrst} H_{rst} \quad , \quad \theta^{mnp8} = \frac{1}{4!} \epsilon^{mnpqrst} F_{qrst} \quad , \quad \theta^{88} = F_{(0)} \quad , \quad (3.17)$$

provided the QC's in (2.12) hold. An explicit computation of (2.12) for the embedding tensor (3.17) yields

$$F_{(0)} H_{(3)} = 0 \quad , \quad (3.18)$$

so $F_{(0)}$ and $H_{(3)}$ can not be simultaneously turned on in a compactification preserving half-maximal supersymmetry in 3D. The embedding tensor components in (3.17) result in the gauge brackets

$$\begin{aligned} [X^{mn}, X^{8p}] &= \frac{2}{3!} \epsilon^{mnpqrst} H_{qrs} X^8{}_t \quad , \\ [X^{8m}, X^{8n}] &= 2 \left(\frac{1}{4!} \epsilon^{mnpqrst} F_{pqrs} X^8{}_t - F_{(0)} X^{mn} \right) \quad , \end{aligned} \quad (3.19)$$

⁸Since we are not including $C_{(7)}$ in the democratic formulation of type IIA supergravity, the vector C_μ that would be captured by a purely internal $C_{(7)}$ must be taken into account explicitly.

Fields	e_n^p	$C_{(1)}$	Φ	$C_{(3)}$	$B_{(2)}$	$B_{(6)}$	$C_{(5)}$
Ω_P	+	-	+	+	-	+	-
σ_{O2}	+	-	+	-	+	+	-
$\mathcal{O}_{\mathbb{Z}_2}$	+	+	+	-	-	+	+
Fluxes	ω_{mn}^p	$F_{(2)}$	$H_{(1)}$	$F_{(4)}$	$H_{(3)}$	$H_{(7)}$	$F_{(6)}$
$\mathcal{O}_{\mathbb{Z}_2}$	-	-	-	+	+	-	-

Table 3: Grading of type IIA fields and fluxes under the O2-plane orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P \sigma_{O2}$. The Romans mass parameter $F_{(0)}$ is not included in the table since it does not originate from a gauge potential.

which generically involve the $21 + 7 + 7$ generators $\{X^{mn}, X^{8p}, X^8_t\}$. Solving (3.18) by setting $F_{(0)} = 0$ trivialises $X^8_t = 0$ and the remaining algebra spanned by $\{X^{mn}, X^{8p}\}$ reduces to the 28-dimensional Abelian gauging of M-theory. On the contrary, solving (3.18) by setting $H_{(3)} = 0$ does not trivialise any generator and yields a 2-step nilpotent algebra of dimension 35 with a 28-dimensional center.

Orientifold interpretation and O2-planes

The half-maximal supergravity models that we have just derived can be realised more string-theoretically in terms of massive type IIA orientifold reductions including O2-planes/D2-branes. Three-dimensional Lorentz invariance requires these sources to be located at

$$\begin{array}{c|c|c|c|c|c|c|c} x^0 & x^1 & x^2 & y^1 & y^2 & y^3 & y^4 & y^5 & y^6 & y^7 \\ \hline \times & \times & \times & & & & & & & \end{array} \quad (3.20)$$

thus filling the 3D external spacetime completely. From a group-theoretical viewpoint, the orientifold action $\mathcal{O}_{\mathbb{Z}_2}$ in the string theory side is precisely the \mathbb{Z}_2 symmetry that truncates from maximal to half-maximal supergravity. More specifically, this \mathbb{Z}_2 projects out those fields and fluxes sitting in spinorial representations of $\text{SO}(8, 8)$ when branching $\text{E}_{8(8)} \supset \text{SO}(8, 8)$ (see Table 1).

The location of the O2-plane in (3.20) is compatible with the $\text{SL}(7)$ covariance of the massive type IIA models. The internal target space involution σ_{O2} reflects all the coordinates (and derivatives) on \mathbb{T}^7 transverse to the O2-plane, namely,

$$\begin{aligned} \sigma_{O2} : \quad y^m &\rightarrow -y^m \\ \partial_m &\rightarrow -\partial_m \end{aligned} \quad \text{with} \quad m = 1, \dots, 7. \quad (3.21)$$

The full orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P \sigma_{O2}$ acts on the various type IIA fields and fluxes as summarised in Table 3. Observe that the set of $\mathcal{O}_{\mathbb{Z}_2}$ -even fields and fluxes in Table 3 precisely matches the ones in (3.16).

The string-theoretic interpretation of the half-maximal supergravity models also allows for a better understanding of the QC's in (3.18). The orientifold $\mathcal{O}_{\mathbb{Z}_2}$ we are considering here allows for O2-planes (and D2-branes). As a result, the flux combination $H_{(3)} \wedge F_{(4)}$ entering (3.1) can be used to cancel a tadpole for $C_{(5)}$ and, therefore, its value is totally unrestricted

	Half-Maximal	$SL(6) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$
	–	$[(\mathbf{6}'_{(-1,-1,-1)} \oplus \mathbf{1}_{(+6,-1,-1)}) \oplus \mathbf{1}_{(0,+7,-1)}] \oplus [(\mathbf{6}_{(+1,+1,+1)} \oplus \mathbf{1}_{(-6,+1,+1)}) \oplus \mathbf{1}_{(0,-7,+1)}]$
Scalars	L^{MN}	$[(\mathbf{15}'_{(-2,-2,-2)} \oplus \mathbf{6}'_{(+5,-2,-2)}) \oplus (\mathbf{6}'_{(-1,+6,-2)} \oplus \mathbf{1}_{(+6,+6,-2)})]$ $\mathbf{1}_{(0,0,0)} \oplus [(\mathbf{6}'_{(-7,0,0)} \oplus (\mathbf{35} + \mathbf{1})_{(0,0,0)} \oplus \mathbf{6}_{(+7,0,0)}) \oplus \mathbf{1}_{(0,0,0)}]$ $(\mathbf{6}'_{(-1,-8,0)} \oplus \mathbf{1}_{(+6,-8,0)}) \oplus (\mathbf{6}_{(+1,+8,0)} \oplus \mathbf{1}_{(-6,+8,0)})]$ $[(\mathbf{6}_{(+1,-6,+2)} \oplus \boxed{\mathbf{1}_{(-6,-6,+2)}}) \oplus (\mathbf{15}_{(+2,+2,+2)} \oplus \mathbf{6}_{(-5,+2,+2)})]$
	–	$[(\mathbf{6}_{(+1,+1,-3)} \oplus \mathbf{1}_{(-6,+1,-3)}) \oplus \mathbf{1}_{(0,-7,-3)}]$ $[(\mathbf{20}_{(+3,+3,-1)} \oplus \mathbf{15}_{(-4,+3,-1)}) \oplus (\mathbf{15}_{(+2,-5,-1)} \oplus \mathbf{6}_{(-5,-5,-1)}]$ $[(\mathbf{15}'_{(-2,+5,+1)} \oplus \mathbf{6}'_{(+5,+5,+1)}) \oplus (\mathbf{20}_{(-3,-3,+1)} \oplus \mathbf{15}'_{(+4,-3,+1)})]$ $[(\boxed{\mathbf{6}'_{(-1,-1,+3)}} \oplus \mathbf{1}_{(+6,-1,+3)}) \oplus \mathbf{1}_{(0,+7,+3)}]$

Table 4: Type IIB branching rules for the embedding $SL(6) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1 \subset SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$. The subscripts in the third column indicate $(\mathbb{R}_3, \mathbb{R}_2, \mathbb{R}_1)$ -charges. We have highlighted the scalars e_m^n , Φ , $B_{(2)}$, $B_{(6)}$ and $C_{(p)}$ listed in (3.25)-(3.26). The physical internal derivatives $\tilde{\partial}_7 \equiv \partial^7$ and ∂_m have been put in a box for their quick identification too. This table should be understood as a continuation of Table 2.

from the 3D supergravity perspective. In other words, O2-planes/D2-branes are compatible with half-maximal supersymmetry in 3D. However, once O2-planes/D2-branes are present, no additional sources can be added to the compactification scheme without causing a further breaking of supersymmetry. In light of this, the QC in (3.18) is nothing but the absence of (a net charge of) O6-planes/D6-branes. Note also that the cancellation of a potential tadpole for $C_{(3)}$ due to O4-planes/D4-branes would require $H_{(3)} \wedge F_{(2)} \neq 0$, but this is simply not possible since the orientifold action $\mathcal{O}_{\mathbb{Z}_2}$ we are considering projects out $F_{(2)}$ (see Table 3). Finally, let us observe that less than half-maximally supersymmetric models including simultaneously O2-planes/D2-branes and O6-planes/D6-branes (thus violating the condition (3.18)) have been investigated in [4] and [6, 7].

3.3 Type IIB with O3-planes

Type IIB reductions on \mathbb{T}^7 down to half-maximal supergravity in 3D can be studied similarly to the M-theory and massive type IIA cases. However, type IIB reductions come along with an additional complication: there is no orientifold action $\mathcal{O}_{\mathbb{Z}_2}$ treating the seven internal coordinates on \mathbb{T}^7 on equal footing.

In this section we will consider the orientifold action induced by an O3-plane filling the external spacetime and a single one-cycle inside

$$\mathbb{T}^7 = \mathbb{T}_1^2 \times \mathbb{T}_2^2 \times \mathbb{T}_3^2 \times \mathbb{S}^1 . \quad (3.22)$$

Taking as a starting point the O2-plane of the type IIA models in the previous section and applying a T-duality along one coordinate – we denote that coordinate as y^7 which specifies the \mathbb{S}^1 in (3.22) – one is left with an O3-plane in the type IIB theory with a dual type IIB coordinate $\tilde{y}^7 \equiv y_7$ (notice the change of position of the index). The presence of the O3-plane breaks $SL(7)$ covariance down to a subgroup $SL(6) \subset SL(7)$. The relevant branching rules for the type IIA with O2 \Leftrightarrow type IIB with O3 correspondence are summarised in Table 4. The

Fluxes	Flux components	Embedding tensor
ω	$\omega_{7m}{}^n \in (\mathbf{35} + \mathbf{1})_{(-6,-6,+2)}$	$\Theta^{7n8}{}_m$
	$\omega_{mn}{}^7 \in \mathbf{15}'_{(+4,+4,+4)}$	Θ^{pqrs}
$H_{(1)}$	$H_7 \in \mathbf{1}_{(-6,-6,+2)}$	$\frac{1}{3\gamma-\alpha+\beta}\Theta^{78}, \frac{1}{\gamma}\Theta^{7m8}{}_m$
$F_{(1)}$	$F_7 \in \mathbf{1}_{(0,-14,+2)}$	Θ^{88}
$H_{(3)}$	$H_{mnp} \in \mathbf{20}_{(-3,+4,+4)}$	$-\Theta^{qrs7}$
$F_{(3)}$	$F_{mnp} \in \mathbf{20}_{(+3,-4,+4)}$	Θ^{qrs8}
$F_{(5)}$	$F_{mnpq7} \in \mathbf{15}_{(-4,-4,+4)}$	Θ^{rs78}

Table 5: Type IIB fluxes in the O3-plane duality frame of half-maximal supergravity and their identification with embedding tensor components.

original type IIA internal derivatives $\partial_m \in \mathbf{7}'_{(-1,+3)} \in \text{SL}(7) \times \mathbb{R}_2 \times \mathbb{R}_1$ in (3.7) branch under $\text{SL}(6) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$ as

$$\begin{aligned} \mathbf{7}'_{(-1,+3)} &\rightarrow \mathbf{6}'_{(-1,-1,+3)} \oplus \mathbf{1}_{(+6,-1,+3)} \quad , \\ \partial_m &\rightarrow \partial_m \oplus \partial_7 \quad . \end{aligned} \quad (3.23)$$

Performing a T-duality along the type IIA coordinate y^7 replaces $\partial_7 \rightarrow \tilde{\partial}_7$ and leaves ∂_m unaffected. As a result, the physical derivatives in the new type IIB duality frame differ from the original type IIA ones and are now identified as

$$\tilde{\partial}_7 \equiv \partial^7 \in \mathbf{1}_{(-6,-6,+2)} \quad \text{and} \quad \partial_m \in \mathbf{6}'_{(-1,-1,+3)} \quad . \quad (3.24)$$

Unlike for the type IIA models in the previous section, now the physical derivatives in the type IIB models have a mixed nature: $\tilde{\partial}_7$ is bosonic (\mathbb{R}_1 charge +2) and ∂_m are spinorial (\mathbb{R}_1 charge +3).

Let us move to group-theoretically identify the internal components of the various type IIB fields. A detailed analysis of bosonic scalars yields

$$\begin{aligned} e_m{}^n &\in (\mathbf{35} + \mathbf{1})_{(0,0,0)} \quad , \quad C_{mnpqr7} \in \mathbf{6}_{(-5,+2,+2)} \quad , \quad C_{(0)} \in \mathbf{1}_{(+6,-8,0)} \quad , \\ e_7{}^7 &\in \mathbf{1}_{(0,0,0)} \quad , \quad C_{mnpq} \in \mathbf{15}_{(+2,+2,+2)} \quad , \quad B_{m7} \in \mathbf{6}'_{(-7,0,0)} \quad , \\ \Phi &\in \mathbf{1}_{(0,0,0)} \quad , \quad C_{m7} \in \mathbf{6}'_{(-1,-8,0)} \quad , \quad B_{mnpqr7} \in \mathbf{6}_{(+1,-6,+2)} \quad , \end{aligned} \quad (3.25)$$

where 15 compact scalars must be subtracted from $e_m{}^n$ upon gauge-fixing of the internal $\text{SO}(6)$ local symmetry, namely, $e_m{}^n \in \text{GL}(6)/\text{SO}(6)$. Note also that there are $64 - 1 = 63$ bosonic scalars in (3.25), the missing one being dual to the vector field $e_\mu{}^7$. The spinorial

Fields	Ω_P	$(-1)^{F_L}$	σ_{O3}	$\mathcal{O}_{\mathbb{Z}_2}$
e_m^n, e_7^7	+	+	+	+
e_m^7, e_7^m			-	-
Φ	+	+	+	+
B_{mn}	-	+	+	-
B_{m7}			-	+
B_{mnpqrs}	-	+	+	-
B_{mnpqr7}			-	+

Fields	Ω_P	$(-1)^{F_L}$	σ_{O3}	$\mathcal{O}_{\mathbb{Z}_2}$
C_{mnpq}	-	-	+	+
C_{mnp7}			-	-
$C_{(0)}$	-	-	+	+
C_{mn}	+	-	+	-
C_{m7}			-	+
C_{mnpqrs}	+	-	+	-
C_{mnpqr7}			-	+

Table 6: Grading of the various type IIB fields under the O3-plane orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P (-1)^{F_L} \sigma_{O3}$. The $\mathcal{O}_{\mathbb{Z}_2}$ -even fields match the ones in (3.25), whereas the $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields match the ones in (3.26).

scalars are given by

$$\begin{aligned}
e_m^7 &\in \mathbf{6}'_{(+5,+5,+1)} \quad , \quad e_7^m \in \mathbf{6}_{(-5,-5,-1)} \quad , \quad C_{mnp7} \in \mathbf{20}_{(-3,-3,+1)} \quad , \\
B_{mn} &\in \mathbf{15}'_{(-2,+5,+1)} \quad , \quad B_{mnpqrs} \in \mathbf{1}_{(+6,-1,+3)} \quad , \quad C_{mn} \in \mathbf{15}'_{(+4,-3,+1)} \quad , \\
C_{mnpqrs} &\in \mathbf{1}_{(0,+7,+3)} \quad ,
\end{aligned} \tag{3.26}$$

adding up to $64 - 6 = 58$. The six missing spinorial scalars are dual to the vector fields e_μ^m . Finally, acting with the partial derivatives in (3.24) on the scalar fields (3.25)-(3.26) produces the set of bosonic fluxes listed in Table 5 together with additional spinorial ones which are projected out of the half-maximal theory.

Orientifold interpretation and O3-planes

The type IIB flux models that we have just described can be string-theoretically interpreted as type IIB orientifold reductions including O3-planes (and D3-branes) placed as

$$\begin{array}{c|c|c|c|c|c|c}
x^0 & x^1 & x^2 & y^1 & y^2 & y^3 & y^4 & y^5 & y^6 & \tilde{y}^7 \\
\hline
\times & \times & \times & & & & & & & \times
\end{array} \tag{3.27}$$

The location of the O3-plane in (3.27) is consistent with the $SL(6)$ covariance of the type IIB models. The internal target space involution σ_{O3} now reflects the six coordinates (and derivatives) on the $\mathbb{T}^6 = \mathbb{T}_1^2 \times \mathbb{T}_2^2 \times \mathbb{T}_3^2$ in (3.22) transverse to the O3-plane, namely,

$$\begin{aligned}
\sigma_{O3} : \quad \tilde{y}^7 &\rightarrow \tilde{y}^7 \quad , \quad y^m \rightarrow -y^m \\
\tilde{\partial}_7 &\rightarrow \tilde{\partial}_7 \quad , \quad \partial_m \rightarrow -\partial_m
\end{aligned} \quad \text{with} \quad m = 1, \dots, 6 \quad , \tag{3.28}$$

and the full orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P (-1)^{F_L} \sigma_{O3}$ acts on the various type IIB fields and fluxes as summarised in Table 6. Note that the $\mathcal{O}_{\mathbb{Z}_2}$ -even and $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields in Table 6 consistently match the ones in (3.25) and (3.26), respectively. The same matching holds at the level of the fluxes.

Half-Maximal	$SL(5) \times SL(2) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$
-	$[(\mathbf{5}', \mathbf{1})_{(-2,-1,-1)} \oplus (\mathbf{1}, \mathbf{2})_{(+5,-1,-1)} \oplus (\mathbf{1}, \mathbf{1})_{(0,+7,-1)}] \oplus [(\mathbf{5}, \mathbf{1})_{(+2,+1,+1)} \oplus (\mathbf{1}, \mathbf{2})_{(-5,+1,+1)} \oplus (\mathbf{1}, \mathbf{1})_{(0,-7,+1)}]$
Scalars	$[(\mathbf{10}', \mathbf{1})_{(-4,-2,-2)} \oplus (\mathbf{5}', \mathbf{2})_{(+3,-2,-2)} \oplus (\mathbf{1}, \mathbf{1})_{(+10,-2,-2)} \oplus ((\mathbf{5}', \mathbf{1})_{(-2,+6,-2)} \oplus (\mathbf{1}, \mathbf{2})_{(+5,+6,-2)})]$ $(\mathbf{1}, \mathbf{1})_{(0,0,0)} \oplus [((\mathbf{5}', \mathbf{2})_{(-7,0,0)} \oplus (\mathbf{24} + \mathbf{1}, \mathbf{1})_{(0,0,0)} \oplus (\mathbf{1}, \mathbf{3} + \mathbf{1})_{(0,0,0)} \oplus (\mathbf{5}, \mathbf{2})_{(+7,0,0)})]$ $((\mathbf{5}', \mathbf{1})_{(-2,-8,0)} \oplus (\mathbf{1}, \mathbf{2})_{(+5,-8,0)}) \oplus ((\mathbf{5}, \mathbf{1})_{(+2,+8,0)} \oplus (\mathbf{1}, \mathbf{2})_{(-5,+8,0)})]$ $[(\mathbf{5}, \mathbf{1})_{(+2,-6,+2)} \oplus \boxed{(\mathbf{1}, \mathbf{2})_{(-5,-6,+2)}} \oplus ((\mathbf{10}, \mathbf{1})_{(+4,+2,+2)} \oplus (\mathbf{5}, \mathbf{2})_{(-3,+2,+2)} \oplus (\mathbf{1}, \mathbf{1})_{(-10,+2,+2)})]$
	$[(\mathbf{5}, \mathbf{1})_{(+2,+1,-3)} \oplus (\mathbf{1}, \mathbf{2})_{(-5,+1,-3)} \oplus (\mathbf{1}, \mathbf{1})_{(0,-7,-3)}]$ $[(\mathbf{5}, \mathbf{1})_{(-8,+3,-1)} \oplus (\mathbf{10}, \mathbf{2})_{(-1,+3,-1)} \oplus (\mathbf{10}', \mathbf{1})_{(+6,+3,-1)}]$ $((\mathbf{5}, \mathbf{2})_{(-3,-5,-1)} \oplus (\mathbf{10}, \mathbf{1})_{(+4,-5,-1)} \oplus (\mathbf{1}, \mathbf{1})_{(-10,-5,-1)})]$ $[(\mathbf{5}', \mathbf{2})_{(+3,+5,+1)} \oplus (\mathbf{10}', \mathbf{1})_{(-4,+5,+1)} \oplus (\mathbf{1}, \mathbf{1})_{(+10,+5,+1)}]$ $((\mathbf{5}', \mathbf{1})_{(+8,-3,+1)} \oplus (\mathbf{10}', \mathbf{2})_{(+1,-3,+1)} \oplus (\mathbf{10}, \mathbf{1})_{(-6,-3,+1)})]$ $[(\mathbf{5}', \mathbf{1})_{(-2,-1,+3)} \oplus \boxed{(\mathbf{1}, \mathbf{2})_{(+5,-1,+3)}} \oplus (\mathbf{1}, \mathbf{1})_{(0,+7,+3)}]$

Table 7: Type IIA branching rules for the embedding $SL(5) \times SL(2) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1 \subset SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$. The subscripts in the third column indicate $(\mathbb{R}_3, \mathbb{R}_2, \mathbb{R}_1)$ -charges. We have highlighted the scalars e_m^n , Φ , $B_{(2)}$, $B_{(6)}$ and $C_{(p)}$ listed in (3.32)-(3.33). The physical internal derivatives $\tilde{\partial}_i \equiv \partial^i$ and $\partial_{\tilde{a}}$ have been put in a box for their quick identification too. This table should be understood as a continuation of Table 2.

The string-theoretic realisation of the half-maximal supergravity models again allows us to better understand the QC's restricting the type IIB fluxes. These include the nilpotency condition ($D^2 = 0$) of the $D = d + \omega$ twisted exterior derivative on the internal space, as well as sourceless Bianchi identities of the form $DH_{(1)} = 0$, $DF_{(1)} = 0$, $DH_{(3)} + \alpha H_{(1)} \wedge H_{(3)} = 0$ and $DF_{(3)} + \beta_3 H_{(1)} \wedge F_{(3)} - H_{(3)} \wedge F_{(1)} = 0$. These follow from the total absence of 7-branes (the first two), NS5-branes (the third one) and 5-branes (the fourth one) in the compactification scheme. More relevant are the Bianchi identities involving O3/D3-sources in (3.1), namely, $DF_{(5)} - H_{(3)} \wedge F_{(3)} = J_{O3/D3}$. Since the orientifold action we are considering here is generated by O3-planes that extend along the internal direction \tilde{y}^7 , we don't expect any QC in the supergravity restricting the number of such sources. Indeed, an explicit computation of the QC's in (2.12) using the embedding tensor/flux correspondence in Table 5 shows that

$$DF_{(5)} - H_{(3)} \wedge F_{(3)} \Big|_{dy^{m_1} \wedge \dots \wedge dy^{m_6}} = \text{unrestricted} , \quad (3.29)$$

while any other component must vanish. In other words, O3/D3-sources threading the submanifold whose Poincaré dual is $dy^{m_1} \wedge \dots \wedge dy^{m_6}$ are compatible with the half-maximal supersymmetry of the type IIB flux models.

3.4 Type IIA with O4-planes

In this section we will consider the orientifold action induced by an O4-plane filling the external spacetime and a single two-cycle inside $\mathbb{T}^7 = \mathbb{T}_1^2 \times \mathbb{T}_2^2 \times \mathbb{T}_3^2 \times \mathbb{S}^1$ in (3.22). We will proceed as before and, starting from the type IIA models with an O2-plane, we will perform two T-dualities along the coordinates in an internal two-torus. Without loss of generality we select the internal two-torus $\mathbb{T}_1^2 \subset \mathbb{T}^7$ in (3.22). The presence of the O4-plane necessarily breaks the $SL(7)$ covariance down to a subgroup $SL(5) \times SL(2) \subset SL(7)$. The relevant branching rules for the type IIA with O2 \Leftrightarrow type IIA with O4 correspondence are summarised in Table 7. The

Fluxes	Flux components	Embedding tensor
ω	$\omega_{ij}{}^j \in (\mathbf{1}, \mathbf{2})_{(-5, -6, +2)}$	$\Theta^{ij8}{}_j$
	$\omega_{i\hat{a}}{}^{\hat{b}} \in (\mathbf{24} + \mathbf{1}, \mathbf{2})_{(-5, -6, +2)}$	$-\Theta^{i\hat{b}8}{}_{\hat{a}}$
	$\omega_{\hat{a}\hat{b}}{}^i \in (\mathbf{10}', \mathbf{2})_{(+1, +4, +4)}$	$\Theta^{\hat{c}\hat{d}e}{}_j$
$H_{(1)}$	$H_i \in (\mathbf{1}, \mathbf{2})_{(-5, -6, +2)}$	$\frac{1}{2\gamma+\delta-\beta}\Theta^{i8}, \frac{1}{\delta}\Theta^{ij8}{}_j, \frac{1}{\gamma}\Theta^{i\hat{a}8}{}_{\hat{a}}$
$H_{(3)}$	$H_{ij\hat{a}} \in (\mathbf{5}', \mathbf{1})_{(-12, -6, +2)}$	$\Theta^{ij8}{}_{\hat{a}}$
	$H_{\hat{a}\hat{b}\hat{c}} \in (\mathbf{10}, \mathbf{1})_{(-6, +4, +4)}$	$\Theta^{\hat{d}\hat{e}i}{}_j$
$F_{(2)}$	$F_{ij} \in (\mathbf{1}, \mathbf{1})_{(0, -14, +2)}$	Θ^{88}
	$F_{\hat{a}\hat{b}} \in (\mathbf{10}', \mathbf{1})_{(+6, -4, +4)}$	$-\Theta^{\hat{c}\hat{d}e}{}_8$
$F_{(4)}$	$F_{i\hat{a}\hat{b}\hat{c}} \in (\mathbf{10}, \mathbf{2})_{(-1, -4, +4)}$	$\Theta^{i\hat{d}e}{}_8$
$F_{(6)}$	$F_{ij\hat{a}\hat{b}\hat{c}\hat{d}} \in (\mathbf{5}, \mathbf{1})_{(-8, -4, +4)}$	$\Theta^{ij\hat{e}8}$

Table 8: Type IIA fluxes in the O4-plane duality frame of half-maximal supergravity and their identification with embedding tensor components. Note that the Romans mass is not present in this duality frame as it corresponds to the spinorial representation $F_{(0)} \in (\mathbf{1}, \mathbf{1})_{(+10, -9, +3)}$.

original type IIA internal derivatives $\partial_m \in \mathbf{7}'_{(-1, +3)} \in \text{SL}(7) \times \mathbb{R}_2 \times \mathbb{R}_1$ in (3.7) now branch under $\text{SL}(5) \times \text{SL}(2) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$ as

$$\begin{aligned} \mathbf{7}'_{(-1, +3)} &\rightarrow (\mathbf{5}', \mathbf{1})_{(-2, -1, +3)} \oplus (\mathbf{1}, \mathbf{2})_{(+5, -1, +3)} \quad , \\ \partial_m &\rightarrow \partial_{\hat{a}} \oplus \partial_i \quad , \end{aligned} \quad (3.30)$$

and the physical derivatives in this new type IIA duality frame are identified as

$$\partial_{\hat{a}} \in (\mathbf{5}', \mathbf{1})_{(-2, -1, +3)} \quad \text{and} \quad \tilde{\partial}_i \equiv \partial^i \in (\mathbf{1}, \mathbf{2})_{(-5, -6, +2)} \quad . \quad (3.31)$$

Note again the mixed nature of the physical derivatives in these type IIA models: $\tilde{\partial}_i$ are bosonic (\mathbb{R}_1 charge +2) and $\partial_{\hat{a}}$ are spinorial (\mathbb{R}_1 charge +3).

We continue with the group-theoretical identification of the internal components of the various type IIA fields. The bosonic scalars are identified as

$$\begin{aligned} e_i{}^j &\in (\mathbf{1}, \mathbf{3} + \mathbf{1})_{(0, 0, 0)} \quad , \quad B_{i\hat{a}} \in (\mathbf{5}', \mathbf{2})_{(-7, 0, 0)} \quad , \quad C_{ij\hat{a}} \in (\mathbf{5}', \mathbf{1})_{(-2, -8, 0)} \\ e_{\hat{a}}{}^{\hat{b}} &\in (\mathbf{24} + \mathbf{1}, \mathbf{1})_{(0, 0, 0)} \quad , \quad B_{ij\hat{a}\hat{b}\hat{c}\hat{d}} \in (\mathbf{5}, \mathbf{1})_{(+2, -6, +2)} \quad , \quad C_{\hat{a}\hat{b}\hat{c}} \in (\mathbf{10}, \mathbf{1})_{(+4, +2, +2)} \\ \Phi &\in (\mathbf{1}, \mathbf{1})_{(0, 0, 0)} \quad , \quad C_i \in (\mathbf{1}, \mathbf{2})_{(+5, -8, 0)} \quad , \quad C_{i\hat{a}\hat{b}\hat{c}\hat{d}} \in (\mathbf{5}, \mathbf{2})_{(-3, +2, +2)} \end{aligned} \quad (3.32)$$

where 10 + 1 compact scalars must be subtracted from $e_{\hat{a}}{}^{\hat{b}}$ and $e_i{}^j$ upon gauge-fixing of the internal $\text{SO}(5) \times \text{SO}(2)$ local symmetry, namely, $e_{\hat{a}}{}^{\hat{b}} \in \text{GL}(5)/\text{SO}(5)$ and $e_i{}^j \in \text{GL}(2)/\text{SO}(2)$.

Fields	Ω_P	$(-1)^{FL}$	σ_{O4}	$\mathcal{O}_{\mathbb{Z}_2}$
$e_i^j, e_{\hat{a}}^{\hat{b}}$	+	+	+	+
$e_i^{\hat{a}}, e_{\hat{a}}^i$			-	-
Φ	+	+	+	+
$B_{ij}, B_{\hat{a}\hat{b}}$	-	+	+	-
$B_{i\hat{a}}$			-	+
$B_{i\hat{a}\hat{b}\hat{c}\hat{d}\hat{e}}$	+	+	-	-
$B_{ij\hat{a}\hat{b}\hat{c}\hat{d}}$			+	+

Fields	Ω_P	$(-1)^{FL}$	σ_{O4}	$\mathcal{O}_{\mathbb{Z}_2}$
$C_{\hat{a}}$	-	-	-	-
C_i			+	+
$C_{ij\hat{a}}, C_{\hat{a}\hat{b}\hat{c}}$	+	-	-	+
$C_{i\hat{a}\hat{b}}$			+	-
$C_{i\hat{a}\hat{b}\hat{c}\hat{d}}$	-	-	+	+
$C_{ij\hat{a}\hat{b}\hat{c}}, C_{\hat{a}\hat{b}\hat{c}\hat{d}\hat{e}}$			-	-

Table 9: Grading of the various type IIA fields under the O4-plane orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P (-1)^{FL} \sigma_{O4}$. The $\mathcal{O}_{\mathbb{Z}_2}$ -even fields match the ones in (3.32), whereas the $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields match the ones in (3.33).

On the other hand, there are $64 - 2 - 1 = 61$ bosonic scalars in (3.32) with the $2 + 1$ missing scalars being dual to the vectors e_{μ}^i and C_{μ} . The spinorial scalars are identified as

$$\begin{aligned}
e_i^{\hat{a}} &\in (\mathbf{5}, \mathbf{2})_{(-3, -5, -1)} & B_{\hat{a}\hat{b}} &\in (\mathbf{10}', \mathbf{1})_{(-4, +5, +1)} & C_{i\hat{a}\hat{b}} &\in (\mathbf{10}', \mathbf{2})_{(+1, -3, +1)} \\
e_{\hat{a}}^i &\in (\mathbf{5}', \mathbf{2})_{(+3, +5, +1)} & B_{i\hat{a}\hat{b}\hat{c}\hat{d}\hat{e}} &\in (\mathbf{1}, \mathbf{2})_{(+5, -1, +3)} & C_{ij\hat{a}\hat{b}\hat{c}} &\in (\mathbf{10}, \mathbf{1})_{(-6, -3, +1)} \\
B_{ij} &\in (\mathbf{1}, \mathbf{1})_{(-10, -5, -1)} & C_{\hat{a}} &\in (\mathbf{5}', \mathbf{1})_{(+8, -3, +1)} & C_{\hat{a}\hat{b}\hat{c}\hat{d}\hat{e}} &\in (\mathbf{1}, \mathbf{1})_{(0, +7, +3)}
\end{aligned} \tag{3.33}$$

They add up to $64 - 5 = 59$ with the 5 missing scalars being dual to the vector fields $e_{\mu}^{\hat{a}}$. Lastly, an explicit computation gives rise to the bosonic fluxes in Table 8 together with additional spinorial fluxes which are projected out of the half-maximal theory.

Orientifold interpretation and O4-planes

The above class of type IIA flux models can be string-theoretically interpreted as type IIA orientifold reductions including O4-planes (and D4-branes) placed as

$$\begin{array}{c|c|c|c|c|c|c|c|c}
x^0 & x^1 & x^2 & \tilde{y}^1 & \tilde{y}^2 & y^3 & y^4 & y^5 & y^6 & y^7 \\
\hline
\times & \times & \times & \times & \times & & & & &
\end{array} \tag{3.34}$$

The location of the O4-plane in (3.34) is this time compatible with the $SL(5) \times SL(2)$ covariance of the type IIA models. The internal target space involution σ_{O4} reflects the five coordinates (and derivatives) on the $\mathbb{T}_2^2 \times \mathbb{T}_3^3 \times \mathbb{S}^1 \subset \mathbb{T}^7$ in (3.22) transverse to the O4-plane. This is

$$\begin{aligned}
\sigma_{O4}: \quad \tilde{y}^i &\rightarrow \tilde{y}^i & , & \quad y^{\hat{a}} &\rightarrow -y^{\hat{a}} \\
\tilde{\partial}_i &\rightarrow \tilde{\partial}_i & , & \quad \partial_{\hat{a}} &\rightarrow -\partial_{\hat{a}}
\end{aligned} \quad \text{with} \quad i = 1, 2 \quad , \quad \hat{a} = 3, \dots, 7. \tag{3.35}$$

At the level of the type IIA fields, the full orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P (-1)^{FL} \sigma_{O4}$ acts as displayed in Table 9. As a check of consistency, the $\mathcal{O}_{\mathbb{Z}_2}$ -even and $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields in Table 9

	Half-Maximal	$SL(4) \times SL(3) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$
	–	$[(\mathbf{4}', \mathbf{1})_{(-3,-1,-1)} \oplus (\mathbf{1}, \mathbf{3}')_{(4,-1,-1)} \oplus \mathbf{1}_{(0,+7,-1)}] \oplus [(\mathbf{4}, \mathbf{1})_{(3,+1,+1)} \oplus (\mathbf{1}, \mathbf{3})_{(-4,+1,+1)} \oplus \mathbf{1}_{(0,-7,+1)}]$
Scalars	L^{MN}	$[(\mathbf{6}, \mathbf{1})_{(-6,-2,-2)} \oplus (\mathbf{4}', \mathbf{3}')_{(1,-2,-2)} \oplus (\mathbf{1}, \mathbf{3})_{(8,-2,-2)} \oplus ((\mathbf{4}', \mathbf{1})_{(-3,+6,-2)} \oplus (\mathbf{1}, \mathbf{3}')_{(4,+6,-2)})]$ $(\mathbf{1}, \mathbf{1})_{(0,0,0)} \oplus [((\mathbf{4}', \mathbf{3})_{(-7,0,0)} \oplus (\mathbf{15} + \mathbf{1}, \mathbf{1})_{(0,0,0)} \oplus (\mathbf{1}, \mathbf{8} + \mathbf{1})_{(0,0,0)} \oplus (\mathbf{4}, \mathbf{3}')_{(7,0,0)})]$ $((\mathbf{4}', \mathbf{1})_{(-3,-8,0)} \oplus (\mathbf{1}, \mathbf{3}')_{(4,-8,0)}) \oplus ((\mathbf{4}, \mathbf{1})_{(3,+8,0)} \oplus (\mathbf{1}, \mathbf{3})_{(-4,+8,0)})]$ $[(\mathbf{4}, \mathbf{1})_{(3,-6,+2)} \oplus \boxed{(\mathbf{1}, \mathbf{3})_{(-4,-6,+2)}} \oplus ((\mathbf{6}, \mathbf{1})_{(6,+2,+2)} \oplus (\mathbf{4}, \mathbf{3})_{(-1,+2,+2)} \oplus (\mathbf{1}, \mathbf{3}')_{(-8,+2,+2)})]$
	–	$[(\mathbf{4}, \mathbf{1})_{(3,+1,-3)} \oplus (\mathbf{1}, \mathbf{3})_{(-4,+1,-3)} \oplus (\mathbf{1}, \mathbf{1})_{(0,-7,-3)}]$ $[(\mathbf{4}, \mathbf{3}')_{(-5,+3,-1)} \oplus (\mathbf{6}, \mathbf{3})_{(2,+3,-1)} \oplus (\mathbf{4}', \mathbf{1})_{(9,+3,-1)} \oplus (\mathbf{1}, \mathbf{1})_{(-12,+3,-1)}]$ $((\mathbf{4}, \mathbf{3})_{(-1,-5,-1)} \oplus (\mathbf{6}, \mathbf{1})_{(6,-5,-1)} \oplus (\mathbf{1}, \mathbf{3}')_{(-8,-5,-1)})]$ $((\mathbf{4}', \mathbf{3}')_{(1,+5,+1)} \oplus (\mathbf{6}, \mathbf{1})_{(-6,+5,+1)} \oplus (\mathbf{1}, \mathbf{3})_{(8,+5,+1)})]$ $[(\mathbf{4}', \mathbf{3})_{(5,-3,+1)} \oplus (\mathbf{6}, \mathbf{3}')_{(-2,-3,+1)} \oplus (\mathbf{4}, \mathbf{1})_{(-9,-3,+1)} \oplus (\mathbf{1}, \mathbf{1})_{(12,-3,+1)}]$ $[(\mathbf{4}', \mathbf{1})_{(-3,-1,+3)} \oplus \boxed{(\mathbf{1}, \mathbf{3}')_{(4,-1,+3)}} \oplus (\mathbf{1}, \mathbf{1})_{(0,+7,+3)}]$

Table 10: Type IIB branching rules for the embedding $SL(4) \times SL(3) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1 \subset SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$. The subscripts in the third column indicate $(\mathbb{R}_3, \mathbb{R}_2, \mathbb{R}_1)$ -charges. We have highlighted the scalars e_m^n , Φ , $B_{(2)}$, $B_{(6)}$ and $C_{(p)}$ listed in (3.39)-(3.40). The physical internal derivatives $\tilde{\partial}_i \equiv \partial^i$ and $\partial_{\hat{a}}$ have been put in a box for their quick identification too. This table should be understood as a continuation of Table 2.

are in one-to-one correspondence with the scalars in (3.32) and (3.33), respectively. The same matching holds also at the level of the fluxes.

The set of QC's that descends from (2.12) upon plugging the embedding tensor/flux dictionary in Table 8 is interpreted as follows. There is the nilpotency condition ($D^2 = 0$) of the $D = d + \omega$ twisted exterior derivative on the internal space, as well as sourceless Bianchi identities of the form $DH_{(1)} = 0$ and $DH_{(3)} + \alpha H_{(1)} \wedge H_{(3)} = 0$ that follow from the total absence of NS7-branes (the first one) and NS5-branes (the second one) in the compactification scheme. In addition, there are Bianchi identities of the form $DF_{(2)} + \beta_2 H_{(1)} \wedge F_{(2)} = 0$ and $DF_{(6)} = 0$ reflecting the absence of D6-branes and D2-branes, respectively. Lastly, there is a non-trivial Bianchi identity (3.1) of the form $DF_{(4)} + \beta_4 H_{(1)} \wedge F_{(4)} - H_{(3)} \wedge F_{(2)} = J_{O4/D4}$ involving O4/D4-sources. Since the orientifold action is generated by O4-planes that extend along the internal two-torus \mathbb{T}_1^2 in (3.22), the QC's in (2.12) yield

$$DF_{(4)} - H_{(3)} \wedge F_{(2)} \Big|_{dy^{\hat{a}_1} \wedge \dots \wedge dy^{\hat{a}_5}} = \text{unrestricted} , \quad (3.36)$$

while any other component vanishes. Equivalently, O4/D4-sources threading the submanifold whose Poincaré dual is $dy^{\hat{a}_1} \wedge \dots \wedge dy^{\hat{a}_5}$ are compatible with the half-maximal supersymmetry of the type IIA flux models under inspection.

3.5 Type IIB with O5-planes

Let us continue with the orientifold action induced by an O5-plane filling the external space-time and a three-cycle inside \mathbb{T}^7 . Taking again as a starting point the O2-plane of the type IIA models but now applying three T-dualities along three type IIA coordinates y^i , one is left with an O5-plane in the type IIB theory with dual type IIB coordinates $\tilde{y}^i \equiv y_i$ (notice the position change of the index i). The presence of the O5-plane breaks the $SL(7)$ covariance down to a

Fluxes	Flux components	Embedding tensor
ω	$\omega_{ij}{}^k \in (\mathbf{1}, \mathbf{6} + \mathbf{3})_{(-4, -6, +2)}$	$\Theta^{ij8}{}_k$
	$\omega_{\hat{a}\hat{b}}{}^i \in (\mathbf{6}, \mathbf{3}')_{(-2, +4, +4)}$	$\Theta^{\hat{c}\hat{d}jk}$
	$\omega_{\hat{a}i}{}^{\hat{b}} \in (\mathbf{15} + \mathbf{1}, \mathbf{3})_{(-4, -6, +2)}$	$\Theta^{i\hat{b}8}{}_{\hat{a}}$
$H_{(1)}$	$H_i \in (\mathbf{1}, \mathbf{3})_{(-4, -6, +2)}$	$\frac{1}{3\gamma+2\alpha-\beta}\Theta^{i8}, \frac{1}{\gamma+\alpha}\Theta^{ij8}{}_j, \frac{1}{\gamma}\Theta^{i\hat{a}8}{}_{\hat{a}}$
$F_{(1)}$	$F_{\hat{a}} \in (\mathbf{4}', \mathbf{1})_{(+9, -4, +4)}$	$-\Theta^{\hat{b}\hat{c}\hat{d}8}$
$H_{(3)}$	$H_{\hat{a}\hat{b}\hat{c}} \in (\mathbf{4}, \mathbf{1})_{(-9, +4, +4)}$	$-\Theta^{\hat{d}ijk}$
	$H_{ij\hat{c}} \in (\mathbf{4}', \mathbf{3}')_{(-11, -6, +2)}$	$\Theta^{ij8}{}_{\hat{c}}$
$F_{(3)}$	$F_{ijk} \in (\mathbf{1}, \mathbf{1})_{(0, -14, +2)}$	$-\Theta^{88}$
	$F_{\hat{a}\hat{b}k} \in (\mathbf{6}, \mathbf{3})_{(+2, -4, +4)}$	$\Theta^{\hat{c}\hat{d}k8}$
$F_{(5)}$	$F_{\hat{a}\hat{b}\hat{c}ij} \in (\mathbf{4}, \mathbf{3}')_{(-5, -4, +4)}$	$\Theta^{\hat{d}ij8}$
$F_{(7)}$	$F_{\hat{a}\hat{b}\hat{c}\hat{d}ijk} \in (\mathbf{1}, \mathbf{1})_{(-12, -4, +4)}$	$-\Theta^{ijk8}$

Table 11: Type IIB fluxes in the O5-plane duality frame of half-maximal supergravity and their identification with embedding tensor components.

subgroup $\text{SL}(4) \times \text{SL}(3) \subset \text{SL}(7)$. The relevant branching rules for the type IIA with O2 \Leftrightarrow type IIB with O5 correspondence are summarised in Table 10. The original type IIA internal derivatives $\partial_m \in \mathbf{7}'_{(-1, +3)} \in \text{SL}(7) \times \mathbb{R}_2 \times \mathbb{R}_1$ in (3.7) branch under $\text{SL}(4) \times \text{SL}(3) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$ as

$$\begin{aligned} \mathbf{7}'_{(-1, +3)} &\rightarrow (\mathbf{1}, \mathbf{3}')_{(+4, -1, +3)} \oplus (\mathbf{4}', \mathbf{1})_{(-3, -1, +3)} \quad , \\ \partial_m &\rightarrow \partial_i \oplus \partial_{\hat{a}} \quad , \end{aligned} \quad (3.37)$$

where, without loss of generality, we take $i = 2, 4, 6$ and $\hat{a} = 1, 3, 5, 7$. Performing three T-dualities along the type IIA coordinates y^i replaces $\partial_i \rightarrow \tilde{\partial}_i$ and leaves $\partial_{\hat{a}}$ unaffected. The physical derivatives in the new type IIB duality frame differ from the original type IIA ones and are identified as

$$\tilde{\partial}_i \equiv \partial^i \in (\mathbf{1}, \mathbf{3})_{(-4, -6, +2)} \quad \text{and} \quad \partial_{\hat{a}} \in (\mathbf{4}', \mathbf{1})_{(-3, -1, +3)} \quad . \quad (3.38)$$

Note the maximally-mixed nature of the physical derivatives in these type IIB models: $\tilde{\partial}_i$ are bosonic (\mathbb{R}_1 charge +2) whereas $\partial_{\hat{a}}$ are spinorial (\mathbb{R}_1 charge +3). This will translate into a more diverse set of type IIB fluxes, as compared to the M-theory and previous type IIA cases, and also a richer structure of flux vacua.

The next step is to group-theoretically identify the internal components of the various

Fields	Ω_P	σ_{O5}	$\mathcal{O}_{\mathbb{Z}_2}$
$e_i^j, e_{\hat{a}}^{\hat{b}}$	+	+	+
$e_i^{\hat{a}}, e_{\hat{a}}^i$		-	-
Φ	+	+	+
$B_{ij}, B_{\hat{a}\hat{b}}$	-	+	-
$B_{i\hat{a}}$		-	+
$B_{ij\hat{a}\hat{b}\hat{c}\hat{d}}$	-	+	-
$B_{ijk\hat{a}\hat{b}\hat{c}}$		-	+

Fields	Ω_P	σ_{O5}	$\mathcal{O}_{\mathbb{Z}_2}$
$C_{\hat{a}\hat{b}\hat{c}\hat{d}}, C_{ij\hat{a}\hat{b}}$	-	+	-
$C_{i\hat{a}\hat{b}\hat{c}}, C_{ijk\hat{a}}$		-	+
$C_{(0)}$	-	+	-
$C_{ij}, C_{\hat{a}\hat{b}}$	+	+	+
$C_{i\hat{a}}$		-	-
$C_{ij\hat{a}\hat{b}\hat{c}\hat{d}}$	+	+	+
$C_{ijk\hat{a}\hat{b}\hat{c}}$		-	-

Table 12: Grading of the type IIB fields under the O5-plane orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P \sigma_{O5}$. $\mathcal{O}_{\mathbb{Z}_2}$ -even fields match the ones in (3.39), whereas $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields match the ones in (3.40).

type IIB fields. A careful analysis shows that they are given by

$$\begin{aligned}
e_i^j &\in (\mathbf{1}, \mathbf{8} + \mathbf{1})_{(0,0,0)} & , & & C_{i\hat{a}\hat{b}\hat{c}} &\in (\mathbf{4}, \mathbf{3})_{(-1,+2,+2)} & , \\
e_{\hat{a}}^{\hat{b}} &\in (\mathbf{15} + \mathbf{1}, \mathbf{1})_{(0,0,0)} & , & & C_{ijk\hat{a}} &\in (\mathbf{4}', \mathbf{1})_{(-3,-8,0)} & , \\
\Phi &\in (\mathbf{1}, \mathbf{1})_{(0,0,0)} & , & & C_{ij} &\in (\mathbf{1}, \mathbf{3}')_{(+4,-8,0)} & , \\
B_{i\hat{a}} &\in (\mathbf{4}', \mathbf{3})_{(-7,0,0)} & , & & C_{\hat{a}\hat{b}} &\in (\mathbf{6}, \mathbf{1})_{(+6,+2,+2)} & , \\
B_{ijk\hat{a}\hat{b}\hat{c}} &\in (\mathbf{4}, \mathbf{1})_{(+3,-6,+2)} & , & & C_{ij\hat{a}\hat{b}\hat{c}\hat{d}} &\in (\mathbf{1}, \mathbf{3}')_{(-8,+2,+2)} & ,
\end{aligned} \tag{3.39}$$

where $3 + 6$ compact scalars must be subtracted from e_i^j and $e_{\hat{a}}^{\hat{b}}$ upon gauge-fixing of the internal $\text{SO}(3) \times \text{SO}(4)$ local symmetry, namely, $e_i^j \in \text{GL}(3)/\text{SO}(3)$ and $e_{\hat{a}}^{\hat{b}} \in \text{GL}(4)/\text{SO}(4)$. The spinorial scalars are identified as

$$\begin{aligned}
e_i^{\hat{a}} &\in (\mathbf{4}, \mathbf{3})_{(-1,-5,-1)} & , & & C_{\hat{a}\hat{b}\hat{c}\hat{d}} &\in (\mathbf{1}, \mathbf{1})_{(0,+7,+3)} & , \\
e_{\hat{a}}^i &\in (\mathbf{4}', \mathbf{3}')_{(+1,+5,+1)} & , & & C_{ij\hat{a}\hat{b}} &\in (\mathbf{6}, \mathbf{3}')_{(-2,-3,+1)} & , \\
B_{ij} &\in (\mathbf{1}, \mathbf{3}')_{(-8,-5,-1)} & , & & C_{(0)} &\in (\mathbf{1}, \mathbf{1})_{(+12,-3,+1)} & , \\
B_{\hat{a}\hat{b}} &\in (\mathbf{6}, \mathbf{1})_{(-6,+5,+1)} & , & & C_{i\hat{a}} &\in (\mathbf{4}', \mathbf{3})_{(+5,-3,+1)} & , \\
B_{ij\hat{a}\hat{b}\hat{c}\hat{d}} &\in (\mathbf{1}, \mathbf{3}')_{(+4,-1,+3)} & , & & C_{ijk\hat{a}\hat{b}\hat{c}} &\in (\mathbf{4}, \mathbf{1})_{(-9,-3,+1)} & .
\end{aligned} \tag{3.40}$$

In order to add up to 64, the $64 - 3 = 61$ bosonic scalars (even \mathbb{R}_1 -charges) in (3.39) must be completed with 3 additional scalars dual to the vectors e_{μ}^i . Similarly, the $64 - 4 = 60$ spinorial scalars (odd \mathbb{R}_1 -charges) in (3.40) must be completed with 4 additional scalars dual to the vectors $e_{\mu}^{\hat{a}}$. Since there are both bosonic and spinorial physical derivatives in (3.38), the set of type IIB fluxes that appear when acting upon the scalars (3.39)-(3.40) is very diverse. An explicit computation yields the bosonic fluxes in Table 11 together with additional spinorial fluxes which are projected out of the half-maximal theory. The fluxes in Table 11 therefore specify the embedding tensor (2.4) and gauge brackets (2.11) of the type IIB half-maximal supergravity models under consideration.

Orientifold interpretation and O5-planes

These type IIB flux models can be string-theoretically interpreted as type IIB orientifold reductions including O5-planes (and D5-branes) located as

$$\frac{x^0 \quad x^1 \quad x^2 \mid y^1 \quad \tilde{y}^2 \mid y^3 \quad \tilde{y}^4 \mid y^5 \quad \tilde{y}^6 \mid y^7}{\times \quad \times \quad \times \mid \quad \times \mid \quad \times \mid \quad \times \mid y^7} \quad (3.41)$$

The O5-plane is filling the external spacetime together with the three internal directions \tilde{y}^i with $i = 2, 4, 6$. The internal target space involution σ_{O5} then reflects the four transverse coordinates (and their derivatives) on \mathbb{T}^7 , namely,

$$\sigma_{\text{O5}} : \quad \begin{aligned} \tilde{y}^i &\rightarrow \tilde{y}^i & , & \quad y^{\hat{a}} \rightarrow -y^{\hat{a}} \\ \tilde{\partial}_i &\rightarrow \tilde{\partial}_i & , & \quad \partial_{\hat{a}} \rightarrow -\partial_{\hat{a}} \end{aligned} \quad \text{with} \quad i = 2, 4, 6 \quad , \quad \hat{a} = 1, 3, 5, 7 . \quad (3.42)$$

The corresponding orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P \sigma_{\text{O5}}$ acts on the type IIB fields as displayed in Table 12. As expected, the $\mathcal{O}_{\mathbb{Z}_2}$ -even and $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields in Table 12 precisely match the ones in (3.39) and (3.40), respectively. The same matching holds at the level of the fluxes.

The string-theoretic realisation of the half-maximal supergravity models again allows us to better understand the QC's restricting the type IIB fluxes. These include the nilpotency condition ($D^2 = 0$) of the $D = d + \omega$ twisted exterior derivative on the internal space, as well as sourceless Bianchi identities of the form $DH_{(1)} = 0$, $DF_{(1)} + \beta_1 H_{(1)} \wedge F_{(1)} = 0$, $DH_{(3)} + \alpha H_{(1)} \wedge H_{(3)} = 0$ and $DF_{(5)} + \beta_5 H_{(1)} \wedge F_{(5)} - H_{(3)} \wedge F_{(3)} = 0$. These follow from the total absence of 7-branes (the first two), NS5-branes (the third one) and D3-branes (the fourth one) in the compactification scheme. More relevant are the Bianchi identities involving O5/D5-sources in (3.1), namely, $DF_{(3)} + \beta_3 H_{(1)} \wedge F_{(3)} - H_{(3)} \wedge F_{(1)} = J_{\text{O5/D5}}$. Since the orientifold action we are considering here is generated by O5-planes that extend along the three internal directions \tilde{y}^i , we don't expect any QC in the supergravity restricting the number of such sources. Indeed, an explicit computation of the QC's in (2.12) using the embedding tensor/flux correspondence in Table 11 shows that

$$DF_{(3)} - H_{(3)} \wedge F_{(1)} \Big|_{dy^{\hat{a}} \wedge dy^{\hat{b}} \wedge dy^{\hat{c}} \wedge dy^{\hat{d}}} = \text{unrestricted} , \quad (3.43)$$

while any other component must vanish. In other words, O5/D5-sources threading the sub-manifold whose Poincaré dual is $dy^{\hat{a}} \wedge dy^{\hat{b}} \wedge dy^{\hat{c}} \wedge dy^{\hat{d}}$ are compatible with the half-maximal supersymmetry of the type IIB flux models under consideration.

3.6 Type IIA with O6-planes

The next orientifold action to be considered is the one induced by an O6-plane filling the external spacetime and a four-cycle inside \mathbb{T}^7 . This case can be obtained from the O2-plane setup by performing four T-dualities along four coordinates $y^{\hat{a}}$. Without loss of generality, we will take such four coordinates to span the four-cycle $\mathbb{T}_2^2 \times \mathbb{T}_3^2 \subset \mathbb{T}^7$ in (3.22). The O6-plane then breaks the $\text{SL}(7)$ covariance down to a subgroup $\text{SL}(3) \times \text{SL}(4) \subset \text{SL}(7)$. The relevant branching rules for the type IIA with O2 \Leftrightarrow type IIA with O6 correspondence are summarised in Table 13. The original type IIA internal derivatives $\partial_m \in \mathbf{7}'_{(-1,+3)} \in \text{SL}(7) \times \mathbb{R}_2 \times \mathbb{R}_1$ in (3.7) branch under $\text{SL}(4) \times \text{SL}(3) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$ as

$$\begin{aligned} \mathbf{7}'_{(-1,+3)} &\rightarrow (\mathbf{1}, \mathbf{3}')_{(+4,-1,+3)} \oplus (\mathbf{4}', \mathbf{1})_{(-3,-1,+3)} , \\ \partial_m &\rightarrow \partial_i \oplus \partial_{\hat{a}} . \end{aligned} \quad (3.44)$$

	Half-Maximal	$SL(4) \times SL(3) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$
	–	$[(\mathbf{4}', \mathbf{1})_{(-3,-1,-1)} \oplus (\mathbf{1}, \mathbf{3}')_{(+4,-1,-1)} \oplus \mathbf{1}_{(0,+7,-1)}] \oplus [(\mathbf{4}, \mathbf{1})_{(+3,+1,+1)} \oplus (\mathbf{1}, \mathbf{3})_{(-4,+1,+1)} \oplus \mathbf{1}_{(0,-7,+1)}]$
Scalars	L^{MN}	$[(\mathbf{6}, \mathbf{1})_{(-6,-2,-2)} \oplus (\mathbf{4}', \mathbf{3}')_{(+1,-2,-2)} \oplus (\mathbf{1}, \mathbf{3})_{(+8,-2,-2)} \oplus ((\mathbf{4}', \mathbf{1})_{(-3,+6,-2)} \oplus (\mathbf{1}, \mathbf{3}')_{(+4,+6,-2)})]$ $(\mathbf{1}, \mathbf{1})_{(0,0,0)} \oplus [((\mathbf{4}', \mathbf{3})_{(-7,0,0)} \oplus (\mathbf{15} + \mathbf{1}, \mathbf{1})_{(0,0,0)} \oplus (\mathbf{1}, \mathbf{8} + \mathbf{1})_{(0,0,0)} \oplus (\mathbf{4}, \mathbf{3}')_{(+7,0,0)})]$ $((\mathbf{4}', \mathbf{1})_{(-3,-8,0)} \oplus (\mathbf{1}, \mathbf{3}')_{(+4,-8,0)}) \oplus ((\mathbf{4}, \mathbf{1})_{(+3,+8,0)} \oplus (\mathbf{1}, \mathbf{3})_{(-4,+8,0)})]$ $[[\boxed{(\mathbf{4}, \mathbf{1})_{(+3,-6,+2)}} \oplus (\mathbf{1}, \mathbf{3})_{(-4,-6,+2)}] \oplus ((\mathbf{6}, \mathbf{1})_{(+6,+2,+2)} \oplus (\mathbf{4}, \mathbf{3})_{(-1,+2,+2)} \oplus (\mathbf{1}, \mathbf{3}')_{(-8,+2,+2)})]$
	–	$[(\mathbf{4}, \mathbf{1})_{(+3,+1,-3)} \oplus (\mathbf{1}, \mathbf{3})_{(-4,+1,-3)} \oplus (\mathbf{1}, \mathbf{1})_{(0,-7,-3)}]$ $[(\mathbf{4}, \mathbf{3}')_{(-5,+3,-1)} \oplus (\mathbf{6}, \mathbf{3})_{(+2,+3,-1)} \oplus (\mathbf{4}', \mathbf{1})_{(+9,+3,-1)} \oplus (\mathbf{1}, \mathbf{1})_{(-12,+3,-1)}]$ $((\mathbf{4}, \mathbf{3})_{(-1,-5,-1)} \oplus (\mathbf{6}, \mathbf{1})_{(+6,-5,-1)} \oplus (\mathbf{1}, \mathbf{3}')_{(-8,-5,-1)})]$ $((\mathbf{4}', \mathbf{3}')_{(+1,+5,+1)} \oplus (\mathbf{6}, \mathbf{1})_{(-6,+5,+1)} \oplus (\mathbf{1}, \mathbf{3})_{(+8,+5,+1)})]$ $[(\mathbf{4}', \mathbf{3})_{(+5,-3,+1)} \oplus (\mathbf{6}, \mathbf{3}')_{(-2,-3,+1)} \oplus (\mathbf{4}, \mathbf{1})_{(-9,-3,+1)} \oplus (\mathbf{1}, \mathbf{1})_{(+12,-3,+1)}]$ $[(\mathbf{4}', \mathbf{1})_{(-3,-1,+3)} \oplus \boxed{(\mathbf{1}, \mathbf{3}')_{(+4,-1,+3)}}] \oplus (\mathbf{1}, \mathbf{1})_{(0,+7,+3)}]$

Table 13: Type IIA branching rules for the embedding $SL(4) \times SL(3) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1 \subset SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$. The subscripts in the third column indicate $(\mathbb{R}_3, \mathbb{R}_2, \mathbb{R}_1)$ -charges. We have highlighted the scalars e_m^n , Φ , $B_{(2)}$, $B_{(6)}$ and $C_{(p)}$ listed in (3.46)-(3.47). The physical internal derivatives $\tilde{\partial}_{\hat{a}} \equiv \partial^{\hat{a}}$ and ∂_i have been put in a box for their quick identification too. This table should be understood as a continuation of Table 2.

Then, the physical derivatives in the type IIA with O6-planes duality frame are identified as

$$\partial_i \in (\mathbf{1}, \mathbf{3}')_{(+4,-1,+3)} \quad \text{and} \quad \tilde{\partial}_{\hat{a}} \equiv \partial^{\hat{a}} \in (\mathbf{4}, \mathbf{1})_{(+3,-6,+2)}. \quad (3.45)$$

Note again the maximally-mixed nature of the physical derivatives in this type IIA duality frame: $\tilde{\partial}_{\hat{a}}$ are bosonic (\mathbb{R}_1 charge +2) and ∂_i are spinorial (\mathbb{R}_1 charge +3).

The group-theoretical identification of the internal components of the various type IIA fields results as follows. The bosonic scalars are identified as

$$\begin{aligned}
e_i^j &\in (\mathbf{1}, \mathbf{8} + \mathbf{1})_{(0,0,0)} & , & & C_{i\hat{a}\hat{b}\hat{c}\hat{d}} &\in (\mathbf{1}, \mathbf{3}')_{(+4,-8,0)} & , \\
e_{\hat{a}}^{\hat{b}} &\in (\mathbf{15} + \mathbf{1}, \mathbf{1})_{(0,0,0)} & , & & C_{ijk\hat{a}\hat{b}} &\in (\mathbf{6}, \mathbf{1})_{(+6,+2,+2)} & , \\
\Phi &\in (\mathbf{1}, \mathbf{1})_{(0,0,0)} & , & & C_{\hat{a}\hat{b}\hat{c}} &\in (\mathbf{4}', \mathbf{1})_{(-3,-8,0)} & , \\
B_{i\hat{a}} &\in (\mathbf{4}, \mathbf{3}')_{(+7,0,0)} & , & & C_{ij\hat{a}} &\in (\mathbf{4}, \mathbf{3})_{(-1,+2,+2)} & , \\
B_{ij\hat{a}\hat{b}\hat{c}\hat{d}} &\in (\mathbf{1}, \mathbf{3})_{(+8,-2,-2)} & , & & C_i &\in (\mathbf{1}, \mathbf{3}')_{(-8,+2,+2)} & ,
\end{aligned} \quad (3.46)$$

where 3 + 6 compact scalars must be subtracted from e_i^j and $e_{\hat{a}}^{\hat{b}}$ upon gauge-fixing of the internal $SO(3) \times SO(4)$ local symmetry, namely, $e_i^j \in GL(3)/SO(3)$ and $e_{\hat{a}}^{\hat{b}} \in GL(4)/SO(4)$. There are $64 - 4 = 60$ bosonic scalars in (3.46), the four missing scalars being dual to the vectors $e_{\mu}^{\hat{a}}$. In addition, the spinorial scalars are given by

$$\begin{aligned}
e_i^{\hat{a}} &\in (\mathbf{4}', \mathbf{3}')_{(+1,+5,+1)} & , & & B_{\hat{a}\hat{b}} &\in (\mathbf{6}, \mathbf{1})_{(+6,-5,-1)} & , & & C_{ijk} &\in (\mathbf{1}, \mathbf{1})_{(0,+7,+3)} & , \\
e_{\hat{a}}^i &\in (\mathbf{4}, \mathbf{3})_{(-1,-5,-1)} & , & & B_{ijk\hat{a}\hat{b}\hat{c}} &\in (\mathbf{4}', \mathbf{1})_{(+9,+3,-1)} & , & & C_{i\hat{a}\hat{b}} &\in (\mathbf{6}, \mathbf{3}')_{(-2,-3,+1)} & , \\
B_{ij} &\in (\mathbf{1}, \mathbf{3})_{(+8,+5,+1)} & , & & C_{ij\hat{a}\hat{b}\hat{c}} &\in (\mathbf{4}', \mathbf{3})_{(+5,-3,+1)} & , & & C_{\hat{a}} &\in (\mathbf{4}, \mathbf{1})_{(-9,-3,+1)} & ,
\end{aligned} \quad (3.47)$$

Fluxes	Flux components	Embedding tensor
ω	$\omega_{\hat{a}\hat{b}}^{\hat{c}} \in (\mathbf{20} + \mathbf{4}, \mathbf{1})_{(+3, -6, +2)}$	$-\Theta^{\hat{a}\hat{b}\hat{8}}_{\hat{c}}$
	$\omega_{\hat{a}i}^{\hat{j}} \in (\mathbf{4}, \mathbf{8} + \mathbf{1})_{(+3, -6, +2)}$	$\Theta^{\hat{a}\hat{j}\hat{8}}_i$
	$\omega_{ij}^{\hat{a}} \in (\mathbf{4}', \mathbf{3})_{(+5, +4, +4)}$	$\Theta^{k\hat{b}\hat{c}\hat{d}}$
$H_{(1)}$	$H_{\hat{a}} \in (\mathbf{4}, \mathbf{1})_{(+3, -6, +2)}$	$\frac{1}{3\gamma+\alpha+\beta}\Theta^{\hat{a}\hat{8}}, \frac{1}{\gamma}\Theta^{\hat{a}\hat{b}\hat{8}}_{\hat{b}}, \frac{1}{\gamma+\alpha}\Theta^{\hat{a}i\hat{8}}_i$
$F_{(0)}$	$F_{(0)} \in (\mathbf{1}, \mathbf{1})_{(-12, -4, +4)}$	$-\Theta^{ijk\hat{8}}$
$H_{(3)}$	$H_{i\hat{a}\hat{b}} \in (\mathbf{6}, \mathbf{3}')_{(+10, -6, +2)}$	$\Theta^{\hat{a}\hat{b}\hat{8}}_i$
	$H_{ijk} \in (\mathbf{1}, \mathbf{1})_{(+12, +4, +4)}$	$-\Theta^{\hat{a}\hat{b}\hat{c}\hat{d}}$
$F_{(2)}$	$F_{i\hat{a}} \in (\mathbf{4}, \mathbf{3}')_{(-5, -4, +4)}$	$\Theta^{jk\hat{a}\hat{8}}$
$F_{(4)}$	$F_{\hat{a}\hat{b}\hat{c}\hat{d}} \in (\mathbf{1}, \mathbf{1})_{(0, -14, +2)}$	$\Theta^{\hat{8}\hat{8}}$
	$F_{ij\hat{a}\hat{b}} \in (\mathbf{6}, \mathbf{3})_{(+2, -4, +4)}$	$\Theta^{k\hat{a}\hat{b}\hat{8}}$
$F_{(6)}$	$F_{ijk\hat{a}\hat{b}\hat{c}} \in (\mathbf{4}', \mathbf{1})_{(+9, -4, +4)}$	$-\Theta^{\hat{a}\hat{b}\hat{c}\hat{8}}$

Table 14: Type IIA fluxes in the O6-plane duality frame of half-maximal supergravity and their identification with embedding tensor components.

adding up to $64 - 3 - 1 = 60$. The $3 + 1$ missing scalars are dual to the vectors e_μ^i and C_μ . Lastly, an explicit computation yields the bosonic fluxes in Table 14 together with additional spinorial fluxes which are projected out of the half-maximal theory.

Orientifold interpretation and O6-planes

These type IIA flux models can be string-theoretically interpreted as type IIA orientifold reductions including O6-planes (and D6-branes) placed as

$$\begin{array}{c|c|c|c|c|c|c|c|c|c} x^0 & x^1 & x^2 & \tilde{y}^1 & y^2 & \tilde{y}^3 & y^4 & \tilde{y}^5 & y^6 & \tilde{y}^7 \\ \hline \times & \times & \times & \times & & \times & & \times & & \times \end{array} \quad (3.48)$$

The O6-plane is filling the external spacetime together with the four internal directions $\tilde{y}^{\hat{a}}$ with $\hat{a} = 1, 3, 5, 7$. The internal target space involution σ_{O6} then reflects the three transverse coordinates (and their derivatives) on \mathbb{T}^7 , namely,

$$\begin{aligned} \sigma_{\text{O6}} : \quad & \tilde{y}^{\hat{a}} \rightarrow \tilde{y}^{\hat{a}} \quad , \quad y^i \rightarrow -y^i \\ & \tilde{\partial}_{\hat{a}} \rightarrow \tilde{\partial}_{\hat{a}} \quad , \quad \partial_i \rightarrow -\partial_i \end{aligned} \quad \text{with} \quad \hat{a} = 1, 3, 5, 7 \quad , \quad i = 2, 4, 6 \quad . \quad (3.49)$$

The corresponding orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P \sigma_{\text{O6}}$ acts on the type IIB fields as summarised in Table 15. The $\mathcal{O}_{\mathbb{Z}_2}$ -even and $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields in Table 15 match the ones in (3.46) and (3.47), respectively. The same matching holds at the level of the fluxes.

Fields	Ω_P	σ_{O6}	$\mathcal{O}_{\mathbb{Z}_2}$
$e_i^j, e_{\hat{a}}^{\hat{b}}$	+	+	+
$e_i^{\hat{a}}, e_{\hat{a}}^i$		-	-
Φ	+	+	+
$B_{ij}, B_{\hat{a}\hat{b}}$	-	+	-
$B_{i\hat{a}}$		-	+
$B_{ijk\hat{a}\hat{b}\hat{c}}$	+	-	-
$B_{ij\hat{a}\hat{b}\hat{c}\hat{d}}$		+	+

Fields	Ω_P	σ_{O6}	$\mathcal{O}_{\mathbb{Z}_2}$
$C_{\hat{a}}$	-	+	-
C_i		-	+
$C_{ij\hat{a}}, C_{\hat{a}\hat{b}\hat{c}}$	+	+	+
$C_{ijk}, C_{i\hat{a}\hat{b}}$		-	-
$C_{ij\hat{a}\hat{b}\hat{c}}$	-	+	-
$C_{i\hat{a}\hat{b}\hat{c}\hat{d}}, C_{ijk\hat{a}\hat{b}}$		-	+

Table 15: Grading of the various type IIA fields under the O6-plane orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P \sigma_{O6}$. The $\mathcal{O}_{\mathbb{Z}_2}$ -even fields match the ones in (3.46), whereas the $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields match the ones in (3.47).

Employing the embedding tensor/flux dictionary in Table 14, the computation of the QC's in (2.12) produces the following constraints on the type IIA fluxes. There is the by now standard nilpotency condition ($D^2 = 0$) of the $D = d + \omega$ twisted exterior derivative on the internal space, as well as sourceless Bianchi identities of the form $DH_{(1)} = 0$ and $DH_{(3)} + \alpha H_{(1)} \wedge H_{(3)} = 0$ forbidding the presence of NS7-branes and NS5-branes in the compactification scheme, respectively. There are also Bianchi identities of the form $DF_{(4)} + \beta_4 H_{(1)} \wedge F_{(4)} - H_{(3)} \wedge F_{(2)} = 0$ and $DF_{(6)} + \beta_6 H_{(1)} \wedge F_{(6)} - H_{(3)} \wedge F_{(4)} = 0$ reflecting the absence of D4-branes and D2-branes, respectively. Finally there is a non-trivial Bianchi identity (3.1) of the form $DF_{(2)} + \beta_2 H_{(1)} \wedge F_{(2)} - H_{(3)} \wedge F_{(0)} = J_{O6/D6}$ involving O6/D6-sources. Recalling that the orientifold action is generated by O6-planes extending along the $y^{\hat{a}}$ coordinates, the QC's in (2.12) yield

$$DF_{(2)} - H_{(3)} \wedge F_{(0)} \Big|_{dy^i \wedge dy^j \wedge dy^k} = \text{unrestricted} , \quad (3.50)$$

while any other component vanishes. Therefore, O6/D6-sources threading the submanifold whose Poincaré dual is $dy^i \wedge dy^j \wedge dy^k$ are compatible with the half-maximal supersymmetry of the type IIA flux models.

3.7 Type IIB with O7-planes

The next duality frame we are considering is type IIB with an O7-plane filling the external spacetime and an internal five-cycle. We will choose it to be $\mathbb{T}_2^2 \times \mathbb{T}_3^2 \times \mathbb{S}^1 \subset \mathbb{T}^7$ in (3.22) without loss of generality. This type IIB with an O7-plane duality frame can be generated from the O2-plane case by applying this time five T-dualities along the aforementioned five-cycle. The presence of the O7-plane breaks $SL(7)$ covariance down to a subgroup $SL(2) \times SL(5) \subset SL(7)$. The relevant branching rules for the type IIA with O2 \Leftrightarrow type IIB with O7 correspondence are summarised in Table 16. The original type IIA internal derivatives $\partial_m \in \mathbf{7}'_{(-1,+3)} \in SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$ in (3.7) branch under $SL(5) \times SL(2) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$ as

$$\begin{aligned} \mathbf{7}'_{(-1,+3)} &\rightarrow (\mathbf{5}', \mathbf{1})_{(-2,-1,+3)} \oplus (\mathbf{1}, \mathbf{2})_{(+5,-1,+3)} , \\ \partial_m &\rightarrow \partial_{\hat{a}} \oplus \partial_i , \end{aligned} \quad (3.51)$$

Half-Maximal	$SL(5) \times SL(2) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$
-	$[(\mathbf{5}', \mathbf{1})_{(-2,-1,-1)} \oplus (\mathbf{1}, \mathbf{2})_{(+5,-1,-1)} \oplus (\mathbf{1}, \mathbf{1})_{(0,+7,-1)}] \oplus [(\mathbf{5}, \mathbf{1})_{(+2,+1,+1)} \oplus (\mathbf{1}, \mathbf{2})_{(-5,+1,+1)} \oplus (\mathbf{1}, \mathbf{1})_{(0,-7,+1)}]$
Scalars	L^{MN} $[(\mathbf{10}', \mathbf{1})_{(-4,-2,-2)} \oplus (\mathbf{5}', \mathbf{2})_{(+3,-2,-2)} \oplus (\mathbf{1}, \mathbf{1})_{(+10,-2,-2)} \oplus ((\mathbf{5}', \mathbf{1})_{(-2,+6,-2)} \oplus (\mathbf{1}, \mathbf{2})_{(+5,+6,-2)})]$ $(\mathbf{1}, \mathbf{1})_{(0,0,0)} \oplus [((\mathbf{5}', \mathbf{2})_{(-7,0,0)} \oplus (\mathbf{24}, \mathbf{1})_{(0,0,0)} \oplus (\mathbf{1}, \mathbf{1})_{(0,0,0)} \oplus (\mathbf{1}, \mathbf{3})_{(0,0,0)})]$ $(\mathbf{5}, \mathbf{2})_{(+7,0,0)} \oplus (\mathbf{1}, \mathbf{1})_{(0,0,0)} \oplus ((\mathbf{5}', \mathbf{1})_{(-2,-8,0)} \oplus (\mathbf{1}, \mathbf{2})_{(+5,-8,0)}) \oplus ((\mathbf{5}, \mathbf{1})_{(+2,+8,0)} \oplus (\mathbf{1}, \mathbf{2})_{(-5,+8,0)})]$ $[(\mathbf{5}, \mathbf{1})_{(+2,-6,+2)} \oplus (\mathbf{1}, \mathbf{2})_{(-5,-6,+2)}] \oplus [(\mathbf{10}, \mathbf{1})_{(+4,+2,+2)} \oplus (\mathbf{5}, \mathbf{2})_{(-3,+2,+2)} \oplus (\mathbf{1}, \mathbf{1})_{(-10,+2,+2)}]$
	-

Table 16: Type IIB branching rules for the embedding $SL(5) \times SL(2) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1 \subset SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$. The subscripts in the third column indicate $(\mathbb{R}_3, \mathbb{R}_2, \mathbb{R}_1)$ -charges. We have highlighted the scalars e_m^n , Φ , $B_{(2)}$, $B_{(6)}$ and $C_{(p)}$ listed in (3.53)-(3.54). The physical internal derivatives $\tilde{\partial}_{\hat{a}} \equiv \partial^{\hat{a}}$ and ∂_i have been put in a box for their quick identification too. This table should be understood as a continuation of Table 2.

where, without loss of generality, we take $i = 1, 2$ and $\hat{a} = 3, 4, 5, 6, 7$. The physical derivatives in this type IIB duality frame are given by

$$\tilde{\partial}_{\hat{a}} \equiv \partial^{\hat{a}} \in (\mathbf{5}, \mathbf{1})_{(+2,-6,+2)} \quad \text{and} \quad \partial_i \in (\mathbf{1}, \mathbf{2})_{(+5,-1,+3)} . \quad (3.52)$$

Note that the derivatives $\tilde{\partial}_{\hat{a}}$ are bosonic (with \mathbb{R}_1 charge +2) whereas ∂_i are spinorial (with \mathbb{R}_1 charge +3).

The group-theoretical identification of the internal components of the various type IIB fields yields the following bosonic scalars

$$\begin{aligned}
e_i^j &\in (\mathbf{1}, \mathbf{3} + \mathbf{1})_{(0,0,0)} & , & & C_{i\hat{a}\hat{b}\hat{c}\hat{d}\hat{e}} &\in (\mathbf{1}, \mathbf{2})_{(+5,-8,0)} & , \\
e_{\hat{a}}^{\hat{b}} &\in (\mathbf{24} + \mathbf{1}, \mathbf{1})_{(0,0,0)} & , & & C_{\hat{a}\hat{b}\hat{c}\hat{d}} &\in (\mathbf{5}', \mathbf{1})_{(-2,-8,0)} & , \\
\Phi &\in (\mathbf{1}, \mathbf{1})_{(0,0,0)} & , & & C_{ij\hat{a}\hat{b}} &\in (\mathbf{10}, \mathbf{1})_{(+4,+2,+2)} & , \\
B_{i\hat{a}} &\in (\mathbf{5}, \mathbf{2})_{(+7,0,0)} & , & & C_{i\hat{a}} &\in (\mathbf{5}, \mathbf{2})_{(-3,+2,+2)} & , \\
B_{i\hat{a}\hat{b}\hat{c}\hat{d}\hat{e}} &\in (\mathbf{1}, \mathbf{2})_{(-5,-6,+2)} & , & & C_{(0)} &\in (\mathbf{1}, \mathbf{1})_{(-10,+2,+2)} & ,
\end{aligned} \quad (3.53)$$

where $10 + 1$ compact scalars must be subtracted from $e_{\hat{a}}^{\hat{b}}$ and e_i^j upon gauge-fixing of the internal $SO(5) \times SO(2)$ local symmetry, namely, $e_{\hat{a}}^{\hat{b}} \in GL(5)/SO(5)$ and $e_i^j \in GL(2)/SO(2)$. The bosonic scalars add up to $64 - 5 = 59$ with the 5 missing scalars being dual to the five vectors $e_{\mu}^{\hat{a}}$. The fermionic scalars are given by

$$\begin{aligned}
e_i^{\hat{a}} &\in (\mathbf{5}', \mathbf{2})_{(+3,+5,+1)} & , & & B_{ij} &\in (\mathbf{1}, \mathbf{1})_{(+10,+5,+1)} & , & & C_{ij\hat{a}\hat{b}\hat{c}\hat{d}} &\in (\mathbf{5}', \mathbf{1})_{(+8,-3,+1)} \\
e_{\hat{a}}^i &\in (\mathbf{5}, \mathbf{2})_{(-3,-5,-1)} & , & & B_{ij\hat{a}\hat{b}\hat{c}\hat{d}} &\in (\mathbf{5}', \mathbf{1})_{(-2,-1,+3)} & , & & C_{ij} &\in (\mathbf{1}, \mathbf{1})_{(0,+7,+3)} \\
B_{\hat{a}\hat{b}} &\in (\mathbf{10}, \mathbf{1})_{(+4,-5,-1)} & , & & C_{\hat{a}\hat{b}} &\in (\mathbf{10}, \mathbf{1})_{(-6,-3,+1)} & , & & C_{i\hat{a}\hat{b}\hat{c}} &\in (\mathbf{10}', \mathbf{2})_{(+1,-3,+1)} \\
&&&&&&&&&& (3.54)
\end{aligned}$$

Fluxes	Flux components	Embedding tensor
ω	$\omega_{\hat{a}\hat{b}}^{\hat{c}} \in (\mathbf{45} + \mathbf{5}, \mathbf{1})_{(+2, -6, +2)}$	$\Theta^{\hat{a}\hat{b}8}_{\hat{c}}$
	$\omega_{\hat{a}\hat{i}}^{\hat{j}} \in (\mathbf{5}, \mathbf{3} + \mathbf{1})_{(+2, -6, +2)}$	$-\Theta^{\hat{a}\hat{j}8}_{\hat{i}}$
	$\omega_{\hat{i}\hat{j}}^{\hat{a}} \in (\mathbf{5}', \mathbf{1})_{(+8, +4, +4)}$	$\Theta^{\hat{b}\hat{c}\hat{d}\hat{e}}$
$H_{(1)}$	$H_{\hat{a}} \in (\mathbf{5}, \mathbf{1})_{(+2, -6, +2)}$	$\frac{1}{3\gamma-\beta}\Theta^{\hat{a}8}, \frac{1}{\gamma-\alpha}\Theta^{\hat{a}\hat{i}8}_{\hat{i}}, \frac{1}{\gamma}\Theta^{\hat{a}\hat{b}8}_{\hat{b}}$
$H_{(3)}$	$H_{\hat{i}\hat{a}\hat{b}} \in (\mathbf{10}, \mathbf{2})_{(+9, -6, +2)}$	$\Theta^{\hat{a}\hat{b}8}_{\hat{i}}$
$F_{(1)}$	$F_{\hat{a}} \in (\mathbf{5}, \mathbf{1})_{(-8, -4, +4)}$	$\Theta^{\hat{a}\hat{i}\hat{j}8}$
$F_{(3)}$	$F_{\hat{i}\hat{a}\hat{b}} \in (\mathbf{10}, \mathbf{2})_{(-1, -4, +4)}$	$\Theta^{\hat{j}\hat{a}\hat{b}8}$
$F_{(5)}$	$F_{\hat{a}\hat{b}\hat{c}\hat{d}\hat{e}} \in (\mathbf{1}, \mathbf{1})_{(0, -14, +2)}$	Θ^{88}
	$F_{\hat{i}\hat{j}\hat{a}\hat{b}\hat{c}} \in (\mathbf{10}', \mathbf{1})_{(+6, -4, +4)}$	$-\Theta^{\hat{a}\hat{b}\hat{c}8}$

Table 17: Type IIB fluxes in the O7-plane duality frame of half-maximal supergravity and their identification with embedding tensor components.

adding up to $64 - 2 = 62$. The 2 missing spinorial scalars are dual to the vectors $e_{\mu}^{\hat{i}}$. Lastly, an explicit computation yields the bosonic fluxes in Table 17 together with additional spinorial fluxes which are projected out of the half-maximal theory.

Orientifold interpretation and O7-planes

This class of type IIB flux models has a string-theoretic description in terms of type IIB orientifold reductions including O7-planes (and D7-branes) placed at

$$\begin{array}{c|c|c|c|c|c|c|c} x^0 & x^1 & x^2 & y^1 & y^2 & \tilde{y}^3 & \tilde{y}^4 & \tilde{y}^5 & \tilde{y}^6 & \tilde{y}^7 \\ \times & \times & \times & & & \times & \times & \times & \times & \times \end{array} \quad (3.55)$$

The O7-plane fills the external spacetime together with the five internal directions $\tilde{y}^{\hat{a}}$ with $\hat{a} = 3, \dots, 7$. The internal target space involution σ_{O7} reflects the two transverse coordinates (and their derivatives) on \mathbb{T}^7 , namely,

$$\sigma_{O7}: \quad \begin{array}{l} \tilde{y}^{\hat{a}} \rightarrow \tilde{y}^{\hat{a}} \quad , \quad y^{\hat{i}} \rightarrow -y^{\hat{i}} \\ \tilde{\partial}_{\hat{a}} \rightarrow \tilde{\partial}_{\hat{a}} \quad , \quad \partial_{\hat{i}} \rightarrow -\partial_{\hat{i}} \end{array} \quad \text{with} \quad \hat{a} = 3, \dots, 7 \quad , \quad \hat{i} = 1, 2. \quad (3.56)$$

The orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P (-1)^{F_L} \sigma_{O7}$ acts on the type IIB fields as summarised in Table 18. Note again the perfect matching between the $\mathcal{O}_{\mathbb{Z}_2}$ -even and $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields in Table 18 and those in (3.53) and (3.54), respectively. The same matching holds at the level of the fluxes.

The QC's in (2.12) reduce to the following string-theoretic conditions on the type IIB fluxes. First of all, there is the nilpotency condition ($D^2 = 0$) of the $D = d + \omega$ twisted exterior derivative on the internal space. There are also the sourceless Bianchi identities

Fields	Ω_P	$(-1)^{F_L}$	σ_{O7}	$\mathcal{O}_{\mathbb{Z}_2}$
$e_i^j, e_{\hat{a}}^{\hat{b}}$	+	+	+	+
$e_i^{\hat{a}}, e_{\hat{a}}^i$			-	-
Φ	+	+	+	+
$B_{ij}, B_{\hat{a}\hat{b}}$	-	+	+	-
$B_{i\hat{a}}$			-	+
$B_{i\hat{a}\hat{b}\hat{c}\hat{d}\hat{e}}$	-	+	-	+
$B_{ij\hat{a}\hat{b}\hat{c}\hat{d}}$			+	-

Fields	Ω_P	$(-1)^{F_L}$	σ_{O7}	$\mathcal{O}_{\mathbb{Z}_2}$
$C_{\hat{a}\hat{b}\hat{c}\hat{d}}, C_{ij\hat{a}\hat{b}}$	-	-	+	+
$C_{i\hat{a}\hat{b}\hat{c}}$			-	-
$C_{(0)}$	-	-	+	+
$C_{ij}, C_{\hat{a}\hat{b}}$	+	-	+	-
$C_{i\hat{a}}$			-	+
$C_{i\hat{a}\hat{b}\hat{c}\hat{d}\hat{e}}$	+	-	-	+
$C_{ij\hat{a}\hat{b}\hat{c}\hat{d}}$			+	-

Table 18: Grading of the various type IIB fields under the O7-plane orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P (-1)^{F_L} \sigma_{O7}$. The $\mathcal{O}_{\mathbb{Z}_2}$ -even fields match the ones in (3.53), whereas the $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields match the ones in (3.54).

$DH_{(1)} = 0$ and $DH_{(3)} + \alpha H_{(1)} \wedge H_{(3)} = 0$ that follow from the absence of NS7-branes and NS5-branes in the compactification scheme, respectively. Moreover, sourceless Bianchi identities $DF_{(5)} + \beta_5 H_{(1)} \wedge F_{(5)} - H_{(3)} \wedge F_{(3)} = 0$ and $DF_{(3)} + \beta_3 H_{(1)} \wedge F_{(3)} - H_{(3)} \wedge F_{(1)} = 0$ appear associated with the absence of D3-branes and D5-branes. Lastly, one finds Bianchi identities involving O7/D7-sources in (3.1), namely, $DF_{(1)} + \beta_1 H_{(1)} \wedge F_{(1)} = J_{O7/D7}$. Since the orientifold action is generated by O7-planes that extend along the five internal directions $\tilde{y}^{\hat{a}}$, there are no QC's restricting the number of such sources, namely,

$$DF_{(1)} \Big|_{dy^i \wedge dy^j} = \text{unrestricted} , \quad (3.57)$$

while any other component must vanish. Therefore, O7/D7-sources threading the submanifold whose Poincaré dual is $dy^i \wedge dy^j$ are compatible with the half-maximal supersymmetry of the type IIB flux models.

3.8 Type IIA with O8-planes

In this section we will consider the orientifold action induced by an O8-plane filling the external spacetime and a single six-cycle inside \mathbb{T}^7 . Without loss of generality we will select $\mathbb{T}_1^2 \times \mathbb{T}_2^2 \times \mathbb{T}_3^2 \subset \mathbb{T}^7$ in (3.22). This O8-plane case is generated by performing six T-dualities on the O2-plane setup along such a particular six-cycle. The presence of the O8-plane breaks $SL(7)$ covariance down to a subgroup $SL(6) \subset SL(7)$. The relevant branching rules for the type IIA with O2 \Leftrightarrow type IIA with O8 correspondence are summarised in Table 19. The original type IIA internal derivatives $\partial_m \in \mathbf{7}'_{(-1,+3)} \in SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$ in (3.7) branch under $SL(6) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$ as

$$\begin{aligned} \mathbf{7}'_{(-1,+3)} &\rightarrow \mathbf{6}'_{(-1,-1,+3)} \oplus \mathbf{1}_{(+6,-1,+3)} , \\ \partial_m &\rightarrow \partial_m \oplus \partial_7 , \end{aligned} \quad (3.58)$$

with $m = 1, \dots, 6$. As a result, the physical derivatives in this type IIA duality frame are identified as

$$\tilde{\partial}_m \equiv \partial^m \in \mathbf{6}_{(+1,-6,+2)} \quad \text{and} \quad \partial_7 \in \mathbf{1}_{(+6,-1,+3)} . \quad (3.59)$$

	Half-Maximal	$SL(6) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1$
	–	$[(\mathbf{6}'_{(-1,-1,-1)} \oplus \mathbf{1}_{(+6,-1,-1)}) \oplus \mathbf{1}_{(0,+7,-1)}] \oplus [(\mathbf{6}_{(+1,+1,+1)} \oplus \mathbf{1}_{(-6,+1,+1)}) \oplus \mathbf{1}_{(0,-7,+1)}]$
Scalars	L^{MN}	$[(\mathbf{15}'_{(-2,-2,-2)} \oplus \mathbf{6}'_{(+5,-2,-2)}) \oplus (\mathbf{6}'_{(-1,+6,-2)} \oplus \mathbf{1}_{(+6,+6,-2)})]$ $\mathbf{1}_{(0,0,0)} \oplus [(\mathbf{6}'_{(-7,0,0)} \oplus \mathbf{35}_{(0,0,0)} \oplus \mathbf{1}_{(0,0,0)} \oplus \mathbf{6}_{(+7,0,0)}) \oplus \mathbf{1}_{(0,0,0)}]$ $(\mathbf{6}'_{(-1,-8,0)} \oplus \mathbf{1}_{(+6,-8,0)}) \oplus (\mathbf{6}_{(+1,+8,0)} \oplus \mathbf{1}_{(-6,+8,0)})]$ $[(\mathbf{6}_{(+1,-6,+2)} \oplus \mathbf{1}_{(-6,-6,+2)}) \oplus (\mathbf{15}_{(+2,+2,+2)} \oplus \mathbf{6}_{(-5,+2,+2)})]$
	–	$[(\mathbf{6}_{(+1,+1,-3)} \oplus \mathbf{1}_{(-6,+1,-3)}) \oplus \mathbf{1}_{(0,-7,-3)}]$ $[(\mathbf{20}_{(+3,+3,-1)} \oplus \mathbf{15}_{(-4,+3,-1)}) \oplus (\mathbf{15}_{(+2,-5,-1)} \oplus \mathbf{6}_{(-5,-5,-1)})]$ $[(\mathbf{15}'_{(-2,+5,+1)} \oplus \mathbf{6}'_{(+5,+5,+1)}) \oplus (\mathbf{20}_{(-3,-3,+1)} \oplus \mathbf{15}'_{(+4,-3,+1)})]$ $[(\mathbf{6}'_{(-1,-1,+3)} \oplus \mathbf{1}_{(+6,-1,+3)}) \oplus \mathbf{1}_{(0,+7,+3)}]$

Table 19: Type IIA branching rules for the embedding $SL(6) \times \mathbb{R}_3 \times \mathbb{R}_2 \times \mathbb{R}_1 \subset SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$. The subscripts in the third column indicate $(\mathbb{R}_3, \mathbb{R}_2, \mathbb{R}_1)$ -charges. We have highlighted the scalars e_m^n , Φ , $B_{(2)}$, $B_{(6)}$ and $C_{(p)}$ listed in (3.60)-(3.61). The physical internal derivatives $\tilde{\partial}_m \equiv \partial^m$ and ∂_7 have been put in a box for their quick identification too. This table should be understood as a continuation of Table 2.

As in previous cases, the physical derivatives in this type IIA frame are of mixed nature: $\tilde{\partial}_m$ are bosonic (\mathbb{R}_1 charge +2) and ∂_7 is spinorial (\mathbb{R}_1 charge +3).

The group-theoretical identification of the internal components of the various type IIA fields gives

$$\begin{aligned}
e_m^n &\in (\mathbf{35} + \mathbf{1})_{(0,0,0)} & , & & B_{mnpqr} &\in \mathbf{1}_{(-6,-6,+2)} & , \\
e_7^7 &\in \mathbf{1}_{(0,0,0)} & , & & C_{mnpqr} &\in \mathbf{6}'_{(-1,-8,0)} & , \\
\Phi &\in \mathbf{1}_{(0,0,0)} & , & & C_{mn7} &\in \mathbf{15}_{(+2,+2,+2)} & , \\
B_{m7} &\in \mathbf{6}_{(+7,0,0)} & , & & C_m &\in \mathbf{6}_{(-5,+2,+2)} & ,
\end{aligned} \tag{3.60}$$

where 15 compact scalars must be subtracted from e_m^n upon gauge-fixing of the internal $SO(6)$ local symmetry, namely, $e_m^n \in GL(6)/SO(6)$. The number of bosonic scalars in (3.60) is $64 - 6 - 1 = 57$, the $6 + 1$ missing scalars being dual to the vectors e_μ^m and C_μ . The spinorial scalars are given by

$$\begin{aligned}
e_m^7 &\in \mathbf{6}_{(-5,-5,-1)} & , & & B_{mnpqr7} &\in \mathbf{6}'_{(-1,-1,+3)} & , & & C_7 &\in \mathbf{1}_{(0,+7,+3)} & , \\
e_7^m &\in \mathbf{6}'_{(+5,+5,+1)} & , & & C_{mnpq7} &\in \mathbf{15}'_{(+4,-3,+1)} & , & & & & \\
B_{mn} &\in \mathbf{15}_{(+2,-5,-1)} & , & & C_{mnp} &\in \mathbf{20}_{(-3,-3,+1)} & . & & & &
\end{aligned} \tag{3.61}$$

They add up to $64 - 1 = 63$ with the missing spinorial scalar being dual to the vector e_μ^7 . Finally, an explicit computation yields the bosonic fluxes displayed in Table 20 together with additional spinorial fluxes which are projected out of the half-maximal theory.

Fluxes	Flux components	Embedding tensor
ω	$\omega_{mn}{}^p \in (\mathbf{84} + \mathbf{6})_{(+1,-6,+2)}$	$\Theta^{mn8}{}_p$
	$\omega_{m7}{}^7 \in \mathbf{6}_{(+1,-6,+2)}$	$\Theta^{m78}{}_7$
$H_{(1)}$	$H_m \in \mathbf{6}_{(+1,-6,+2)}$	$\frac{1}{\gamma}\Theta^{m8}, \frac{3}{\gamma+\beta}\Theta^{mn8}{}_n, \frac{1}{\delta}\Theta^{m78}{}_7$
$H_{(3)}$	$H_{mn7} \in \mathbf{15}_{(+8,-6,+2)}$	$-\Theta^{mn8}{}_7$
$F_{(2)}$	$F_{mn} \in \mathbf{15}_{(-4,-4,+4)}$	Θ^{mn78}
$F_{(4)}$	$F_{mnp7} \in \mathbf{20}_{(+3,-4,+4)}$	Θ^{mnp8}
$F_{(6)}$	$F_{mnpqrs} \in \mathbf{1}_{(0,-14,+2)}$	Θ^{88}

Table 20: Type IIA fluxes in the O8-plane duality frame of half-maximal supergravity and their identification with embedding tensor components. Note that the Romans mass is not present in this duality frame as it corresponds to the spinorial representation $F_{(0)} \in \mathbf{1}_{(-6,+1,+5)}$.

Orientifold interpretation and O8-planes

The type IIA flux models here can be string-theoretically described as type IIA orientifold reductions including O8-planes (and D8-branes) located at

$$\begin{array}{c|c|c|c|c|c|c|c|c|c} x^0 & x^1 & x^2 & \tilde{y}^1 & \tilde{y}^2 & \tilde{y}^3 & \tilde{y}^4 & \tilde{y}^5 & \tilde{y}^6 & y^7 \\ \hline \times & \times & \times & \times & \times & \times & \times & \times & \times & \times \end{array} \quad (3.62)$$

The O8-plane fills the external spacetime and the six internal directions \tilde{y}^m . The internal target space involution σ_{O8} reflects the unique transverse coordinate (and its derivative) on \mathbb{T}^7 , namely,

$$\begin{aligned} \sigma_{O8} : \quad \tilde{y}^m &\rightarrow \tilde{y}^m \quad , \quad y^7 \rightarrow -y^7 \\ \tilde{\partial}_m &\rightarrow \tilde{\partial}_m \quad , \quad \partial_7 \rightarrow -\partial_7 \end{aligned} \quad \text{with} \quad m = 1, \dots, 6. \quad (3.63)$$

The corresponding orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P (-1)^{F_L} \sigma_{O8}$ acts on the type IIA fields as summarised in Table 21. The $\mathcal{O}_{\mathbb{Z}_2}$ -even and $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields in Table 21 match the ones in (3.60) and (3.61), respectively. The same matching holds at the level of the fluxes.

Upon use of the embedding tensor/flux dictionary in Table 20, the computation of the QC's in (2.12) gives rise to the following constraints on the type IIA fluxes. There is the nilpotency condition ($D^2 = 0$) of the $D = d + \omega$ twisted exterior derivative on the internal space. In addition, there are sourceless Bianchi identities of the form $DH_{(1)} = 0$, $DH_{(3)} + \alpha H_{(1)} \wedge H_{(3)} = 0$, $DF_{(2)} + \beta_2 H_{(1)} \wedge F_{(2)} = 0$ and $DF_{(4)} + \beta_4 H_{(1)} \wedge F_{(4)} - H_{(3)} \wedge F_{(2)} = 0$ reflecting the absence of NS7-branes, NS5-branes, D6-branes and D4-branes in the compactification scheme, respectively. Note also that the constraint $DF_{(6)} = 0$, reflecting the absence of D2-branes, trivialises with the permitted fluxes in Table 20. One is then left with no non-trivial constraints on the gauge fluxes. In other words, no flux-induced tadpole can be generated for the O8/D8 sources. Therefore, these type IIA flux models satisfy the extra constraints in (2.13) and are embeddable into maximal three-dimensional supergravity.

Fields	Ω_P	$(-1)^{F_L}$	σ_{O8}	$\mathcal{O}_{\mathbb{Z}_2}$
e_7^7, e_m^n	+	+	+	+
e_7^m, e_m^7			-	-
Φ	+	+	+	+
B_{mn}	-	+	+	-
B_{m7}			-	+
B_{mnpqr7}	+	+	-	-
B_{mnpqrs}			+	+

Fields	Ω_P	$(-1)^{F_L}$	σ_{O8}	$\mathcal{O}_{\mathbb{Z}_2}$
C_m	-	-	+	+
C_7			-	-
C_{mnp}	+	-	+	-
C_{mn7}			-	+
C_{mnpq7}	-	-	-	-
C_{mnpqr}			+	+

Table 21: Grading of the various type IIA fields under the O8-plane orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P (-1)^{F_L} \sigma_{O8}$. The $\mathcal{O}_{\mathbb{Z}_2}$ -even fields match the ones in (3.60), whereas the $\mathcal{O}_{\mathbb{Z}_2}$ -odd fields match the ones in (3.61).

3.9 Type IIB with O9-planes: type I/Heterotic

Let us now consider the duality frame of the type I theory, *i.e.* type IIB with O9-planes. This duality frame is obtained upon further T-dualisation of the $y^{\hat{a}}$ coordinates in (3.42). As a result of this procedure, the physical coordinates and their associated derivatives turn out to be

$$\tilde{y}^m \in \mathbf{7}'_{(+6,-2)} \quad \text{and} \quad \tilde{\partial}_m \in \mathbf{7}_{(-6,+2)} , \quad (3.64)$$

so that they all become bosonic as far as their $SO(8,8)$ origin is concerned (see Table 22). A group-theoretical identification of the internal components of the type IIB fields yields

$$\begin{aligned} e_m^n &\in (\mathbf{48} + \mathbf{1})_{(0,0)} \quad , \quad C_{(2)} \in \mathbf{21}_{(+2,+2)} \quad , \\ \Phi &\in \mathbf{1}_{(0,0)} \quad , \quad C_{(6)} \in \mathbf{7}'_{(-8,0)} \quad , \end{aligned} \quad (3.65)$$

together with

$$\begin{aligned} B_{(2)} &\in \mathbf{21}_{(-5,-1)} \quad , \quad C_{(0)} \in \mathbf{1}_{(+7,+3)} \quad , \\ B_{(6)} &\in \mathbf{7}'_{(-1,+3)} \quad , \quad C_{(4)} \in \mathbf{35}'_{(-3,+1)} \quad , \end{aligned} \quad (3.66)$$

where, as before, 21 compact scalars must be subtracted from e_m^n upon gauge-fixing of the internal $SO(7)$ local symmetry, namely, $e_m^n \in GL(7)/SO(7)$. The $64 - 7 = 57$ bosonic scalars in (3.65) must be completed with 7 additional scalars dual to the vectors e_μ^m in order to match the 64 scalars of half-maximal supergravity. The spinorial scalars in (3.66) correctly add up to 64. Acting on the scalars (3.65) and (3.66) with the physical derivatives in (3.64) produces fluxes of the form

$$\begin{aligned} \omega_{mn}{}^p &\in (\mathbf{140} + \mathbf{7})_{(-6,+2)} \quad , \quad F_{(3)} \in \mathbf{35}_{(-4,+4)} \quad , \\ H_{(1)} &\in \mathbf{7}_{(-6,+2)} \quad , \quad F_{(7)} \in \mathbf{1}_{(-14,+2)} \quad , \end{aligned} \quad (3.67)$$

together with

$$\begin{aligned} H_{(3)} &\in \mathbf{35}_{(-11,+1)} \quad , \quad F_{(1)} \in \mathbf{7}_{(+1,+5)} \quad , \\ H_{(7)} &\in \mathbf{1}_{(-7,+5)} \quad , \quad F_{(5)} \in \mathbf{21}'_{(-9,+3)} \quad . \end{aligned} \quad (3.68)$$

	Half-Maximal	$SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$
	-	$[\mathbf{7}'_{(-1,-1)} \oplus \mathbf{1}_{(+7,-1)}] \oplus [\mathbf{7}_{(+1,+1)} \oplus \mathbf{1}_{(-7,+1)}]$
Scalars	L^{MN}	$[\mathbf{21}'_{(-2,-2)} \oplus \mathbf{7}'_{(+6,-2)}] \oplus [\mathbf{7}'_{(-8,0)} \oplus (\mathbf{48} + \mathbf{1})_{(0,0)} \oplus \mathbf{7}_{(+8,0)}]$ $\mathbf{1}_{(0,0)} \oplus \boxed{\mathbf{7}_{(-6,+2)}} \oplus \mathbf{21}_{(+2,+2)}$
	-	$[\mathbf{7}_{(+1,-3)} \oplus \mathbf{1}_{(-7,-3)}] \oplus [\mathbf{35}_{(+3,-1)} \oplus \mathbf{21}_{(-5,-1)}]$ $[\mathbf{35}'_{(-3,+1)} \oplus \mathbf{21}'_{(+5,+1)}] \oplus [\mathbf{7}'_{(-1,+3)} \oplus \mathbf{1}_{(+7,+3)}]$
Embedding Tensor / Fluxes	θ	$\mathbf{1}_{(0,0)}$
	θ_{MN}	$[\mathbf{28}'_{(-2,-2)} \oplus \mathbf{7}'_{(+6,-2)} \oplus \mathbf{1}_{(+14,-2)}]$ $[\mathbf{7}'_{(-8,0)} \oplus \mathbf{48}_{(0,0)} \oplus \mathbf{1}_{(0,0)} \oplus \mathbf{7}_{(+8,0)}]$ $[\mathbf{28}_{(+2,+2)} \oplus \mathbf{7}_{(-6,+2)} \oplus \mathbf{1}_{(-14,+2)}]$
	θ_{MNPQ}	$[\mathbf{35}_{(-4,-4)} \oplus \mathbf{35}'_{(+4,-4)}] \oplus [\mathbf{21}'_{(-2,-2)} \oplus \mathbf{7}'_{(+6,-2)}]$ $[\mathbf{35}'_{(-10,-2)} \oplus \mathbf{21}'_{(-2,-2)} \oplus \mathbf{224}'_{(-2,-2)} \oplus \mathbf{140}'_{(+6,-2)}]$
		$[\mathbf{140}'_{(-8,0)} \oplus \mathbf{392}_{(0,0)} \oplus \mathbf{48}_{(0,0)} \oplus \mathbf{140}_{(+8,0)}]$ $[\mathbf{7}'_{(-8,0)} \oplus \mathbf{48}_{(0,0)} \oplus \mathbf{1}_{(0,0)} \oplus \mathbf{7}_{(+8,0)}] \oplus \mathbf{1}_{(0,0)}$
		$[\mathbf{35}_{(-4,+4)} \oplus \mathbf{35}'_{(+4,+4)}] \oplus [\mathbf{7}_{(-6,+2)} \oplus \mathbf{21}_{(+2,+2)}]$ $[\mathbf{35}_{(+10,+2)} \oplus \mathbf{21}_{(+2,+2)} \oplus \mathbf{224}_{(+2,+2)} \oplus \mathbf{140}_{(-6,+2)}]$
		$[\mathbf{7}'_{(-1,-5)} \oplus \mathbf{1}_{(+7,-5)}] \oplus [\mathbf{1}_{(-7,-3)} \oplus \mathbf{7}_{(+1,-3)}]$ $[\mathbf{48}_{(-7,-3)} \oplus \mathbf{140}_{(+1,-3)} \oplus \mathbf{7}_{(+1,-3)} \oplus \mathbf{21}_{(+9,-3)}]$
	$[\mathbf{21}_{(-5,-1)} \oplus \mathbf{35}_{(+3,-1)}]$ $[\mathbf{224}_{(-5,-1)} \oplus \mathbf{210}_{(+3,-1)} \oplus \mathbf{35}_{(+3,-1)} \oplus \mathbf{35}'_{(+11,-1)}]$ $[\mathbf{7}_{(-13,-1)} \oplus \mathbf{21}_{(-5,-1)} \oplus \mathbf{28}_{(-5,-1)} \oplus \mathbf{112}_{(+3,-1)}]$	
$[\mathbf{21}'_{(+5,+1)} \oplus \mathbf{35}'_{(-3,+1)}]$ $[\mathbf{224}'_{(+5,+1)} \oplus \mathbf{210}'_{(-3,+1)} \oplus \mathbf{35}'_{(-3,+1)} \oplus \mathbf{35}_{(-11,+1)}]$ $[\mathbf{7}'_{(+13,+1)} \oplus \mathbf{21}'_{(+5,+1)} \oplus \mathbf{28}'_{(+5,+1)} \oplus \mathbf{112}'_{(-3,+1)}]$		
$[\mathbf{7}_{(+1,+5)} \oplus \mathbf{1}_{(-7,+5)}] \oplus [\mathbf{7}'_{(-1,+3)} \oplus \mathbf{1}_{(+7,+3)}]$ $[\mathbf{48}_{(+7,+3)} \oplus \mathbf{140}'_{(-1,+3)} \oplus \mathbf{7}'_{(-1,+3)} \oplus \mathbf{21}'_{(-9,+3)}]$		

Table 22: Type I branching rules for the embedding $SO(8,8) \supset SL(7) \times \mathbb{R}_2 \times \mathbb{R}_1$. The subscripts in the third column indicate $(\mathbb{R}_2, \mathbb{R}_1)$ -charges. We have highlighted the scalars e_m^n , Φ , $B_{(2)}$, $B_{(6)}$ and $C_{(p)}$ listed in (3.65)-(3.66) as well as their associated fluxes in (3.67)-(3.68). The physical internal derivatives $\tilde{\partial}_m$ have been put in a box for their quick identification.

Truncating away the scalars in (3.66) and the fluxes in (3.68), which descend from spinorial representations of $SO(8,8)$, the field content of the type I models is given by

$$\begin{aligned}
\text{Scalars} & : e_m^n \in \frac{GL(7)}{SO(7)} \quad , \quad \Phi \quad , \quad C_{(2)} \quad , \quad C_{(6)} \quad , \\
\text{Fluxes} & : \omega_{mn}{}^p \quad , \quad H_{(1)} \quad , \quad F_{(3)} \quad , \quad F_{(7)} \quad .
\end{aligned} \tag{3.69}$$

Note that the content of fields and fluxes in (3.69) becomes the universal NS-NS sector

Fields	e_n^p	Φ	$B_{(2)}$	$B_{(6)}$	$C_{(0)}$	$C_{(2)}$	$C_{(4)}$	$C_{(6)}$
Ω_P	+	+	-	-	-	+	-	+
σ_{O9}	+	+	+	+	+	+	+	+
$\mathcal{O}_{\mathbb{Z}_2}$	+	+	-	-	-	+	-	+
Fluxes	ω_{mn}^p	$H_{(1)}$	$H_{(3)}$	$H_{(7)}$	$F_{(1)}$	$F_{(3)}$	$F_{(5)}$	$F_{(7)}$
$\mathcal{O}_{\mathbb{Z}_2}$	+	+	-	-	-	+	-	+

Table 23: Grading of the various type IIB fields and fluxes under the O9-plane orientifold action $\mathcal{O}_{\mathbb{Z}_2} = \Omega_P \sigma_{O9}$.

of the Heterotic string upon a S-duality: $\Phi \rightarrow -\Phi$, $C_{(2)} \rightarrow B_{(2)}$ and $C_{(6)} \rightarrow B_{(6)}$, with the corresponding fluxes also being exchanged. The fluxes in (3.69) induce a gauging with embedding tensor components

$$\begin{aligned}
\theta^{mn8}{}_p &= \omega_{mn}^p + \frac{\gamma - \alpha + \beta}{3} \delta_{[m}^p H_{n]} & , & & \theta^{mnp8} &= F_{mnp} & , & & (3.70) \\
\theta^{m8} &= \gamma H_m & , & & \theta^{88} &= \frac{1}{7!} \varepsilon^{mnpqrst} F_{mnpqrst} & . & &
\end{aligned}$$

An explicit computation of the QC's in (2.12) yields

$$\omega_{[mn}{}^r \omega_{p]r}{}^s = 0 \quad , \quad \omega_{[mn}{}^r F_{pq]r} + (\beta - \alpha) H_{[m} F_{npq]} = 0 \quad , \quad \omega_{mn}{}^r H_r = 0 . \quad (3.71)$$

Orientifold interpretation and O9-planes

The O9-plane is somehow special. Being placed as

$$\begin{array}{c|c|c|c|c|c|c|c|c|c}
x^0 & x^1 & x^2 & \tilde{y}^1 & \tilde{y}^2 & \tilde{y}^3 & \tilde{y}^4 & \tilde{y}^5 & \tilde{y}^6 & \tilde{y}^7 \\
\hline
\times & \times & \times & \times & \times & \times & \times & \times & \times & \times
\end{array} \quad (3.72)$$

it does fill the entire ten-dimensional spacetime. As a consequence, the internal target space involution σ_{O9} does not reflect any of the internal coordinates (and derivatives) on \mathbb{T}^7 , namely,

$$\begin{aligned}
\sigma_{O9} : \quad \tilde{y}^m &\rightarrow \tilde{y}^m & \text{with} & & m &= 1, \dots, 7 . & (3.73) \\
\tilde{\partial}_m &\rightarrow \tilde{\partial}_m & & & & &
\end{aligned}$$

In the string-theoretic language of orientifold reductions, the QC's in (3.71) are reinterpreted as follows. The first condition is again the nilpotency condition ($D^2 = 0$) of the twisted exterior derivative $D = d + \omega$ on the internal space. The second and third conditions are interpreted as the sourceless Bianchi identities $DF_{(3)} + \beta_3 H_{(1)} \wedge F_{(3)} = 0$ and $DH_{(1)} = 0$ forbidding D5-branes and NS7-branes, respectively. In the S-dual Heterotic picture, these map into the absence of NS5-branes and (again) NS7-branes. Finally, although in principle permitted by half-maximal supersymmetry, no flux-induced tadpole can be defined for the O9/D9 sources. Therefore, these type I/Heterotic flux models are embeddable into maximal three-dimensional supergravity.

4 Flux models and vacua

In the previous section we have established a precise correspondence between single Op -plane orientifold reductions of type II strings in the presence of background fluxes and half-maximal gauged supergravities in three dimensions. Equipped with this correspondence we can start charting the landscape of half-maximal flux compactifications in three dimensions.

4.1 Invariant sectors

Half-maximal supergravity contains 64 scalars. In the light-cone basis of $\text{SO}(8, 8)$ they parameterise the scalar geometry (2.1) in terms of a coset representative of the form

$$\mathcal{V} = \begin{pmatrix} \mathbb{I} & 0 \\ \mathbf{b} & \mathbb{I} \end{pmatrix} \begin{pmatrix} \mathbf{e} & 0 \\ 0 & \mathbf{e}^{-T} \end{pmatrix} \in \frac{\text{SO}(8, 8)}{\text{SO}(8) \times \text{SO}(8)}, \quad (4.1)$$

where $\mathbf{e} \in \text{GL}(8)/\text{SO}(8)$ and $\mathbf{b} = -\mathbf{b}^T$ are 8×8 matrices containing 36 and 28 scalars, respectively. The scalar-dependent and symmetric matrix M_{MN} entering (2.2) takes the form

$$M = \mathcal{V} \mathcal{V}^T = \begin{pmatrix} \mathbf{g} & -\mathbf{g} \mathbf{b} \\ \mathbf{b} \mathbf{g} & \mathbf{g}^{-1} - \mathbf{b} \mathbf{g} \mathbf{b} \end{pmatrix} \in \text{SO}(8, 8), \quad (4.2)$$

with $\mathbf{g} = \mathbf{e} \mathbf{e}^T$. In addition, there is a (worrying) large number of both metric and gauge flux components which complicates the analysis of the vacuum structure of the models. In order to derive more tractable models, still able to capture relevant physics, we will require invariance of the fields and fluxes under additional \mathbb{Z}_2^2 and $\text{SO}(3)$ symmetries.

4.1.1 The \mathbb{Z}_2^2 invariant sector: the eight-chiral model

The first symmetry we will mod out the theory by is a \mathbb{Z}_2^2 discrete symmetry. It consists of two independent elements acting on the vector index $A = 1, \dots, 8$ of $\text{SL}(8) \subset \text{SO}(8, 8)$ as

$$\begin{aligned} \mathbb{Z}_2^{(\alpha)} : (y^1, y^2, y^3, y^4, y^5, y^6, y^7, y^8) &\rightarrow (-y^1, -y^2, -y^3, -y^4, y^5, y^6, y^7, y^8) \\ \mathbb{Z}_2^{(\beta)} : (y^1, y^2, y^3, y^4, y^5, y^6, y^7, y^8) &\rightarrow (-y^1, -y^2, y^3, y^4, -y^5, -y^6, y^7, y^8) \end{aligned} \quad (4.3)$$

together with the additional elements \mathbb{I} and $\mathbb{Z}_2^{(\alpha)} \mathbb{Z}_2^{(\beta)}$. This \mathbb{Z}_2^2 discrete symmetry leaves invariant $8 + 8$ non-compact scalars within the coset space (2.3). In the Lorentzian basis, these are associated to the generators $L^{\mathcal{I}r}$ in (2.15) with $(\mathcal{I}, r) = (1, 1), (2, 2), \dots, (8, 8)$ as well as $(\mathcal{I}, r) = (1, 2), (2, 1), (3, 4), (4, 3), (5, 6), (6, 5), (7, 8), (8, 7)$. The scalar geometry describing the \mathbb{Z}_2^2 invariant sector is identified with

$$\mathcal{M}_{\text{scal}} = \left[\frac{\text{SL}(2)}{\text{SO}(2)} \right]^8 \subset \frac{\text{SO}(8, 8)}{\text{SO}(8) \times \text{SO}(8)}, \quad (4.4)$$

and comprises eight copies of the Poincaré disk $\text{SL}(2)/\text{SO}(2)$ parameterised by eight complex scalars

$$T_I = B_I + iA_I \quad \text{and} \quad U_I = u_I + i\mu_I \quad \text{with} \quad I = 1, \dots, 4. \quad (4.5)$$

Using the upper-half plane parameterisation of the Poincaré disk, *i.e.* $A_I > 0, \mu_I > 0$, the coset representative in (4.1) adopts the block-diagonal structure

$$\mathbf{e} = \begin{pmatrix} \mathbf{e}_1 & 0 & 0 & 0 \\ 0 & \mathbf{e}_2 & 0 & 0 \\ 0 & 0 & \mathbf{e}_3 & 0 \\ 0 & 0 & 0 & \mathbf{e}_4 \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} \mathbf{b}_1 & 0 & 0 & 0 \\ 0 & \mathbf{b}_2 & 0 & 0 \\ 0 & 0 & \mathbf{b}_3 & 0 \\ 0 & 0 & 0 & \mathbf{b}_4 \end{pmatrix}, \quad (4.6)$$

with

$$\mathbf{e}_I = (A_I \mu_I)^{-\frac{1}{2}} \begin{pmatrix} 1 & 0 \\ u_I & \mu_I \end{pmatrix} \quad , \quad \mathbf{b}_I = \begin{pmatrix} 0 & B_I \\ -B_I & 0 \end{pmatrix} . \quad (4.7)$$

In the M-theory duality frame of Section 3.1, the maximal subgroup $\mathrm{GL}(8) \subset \mathrm{SO}(8, 8)$ is realised purely geometrically. The parameterisation in (4.6)-(4.7) then gives rise to a matrix \mathbf{g} in (4.2) that is interpreted as the metric on the internal \mathbb{T}^8 . It is

$$ds_8^2 = \sum_{I=1}^4 \frac{A_I^{-1}}{\mathrm{Im}U_I} \left[(\eta^{2I-1})^2 + |U_I|^2 (\eta^{2I})^2 + 2 (\mathrm{Re}U_I) \eta^{2I-1} \eta^{2I} \right] , \quad (4.8)$$

where $\eta^A = \bigoplus_{I=1}^4 (\eta^{2I-1}, \eta^{2I})$ denotes a basis of one-forms in the internal $\mathbb{T}^8 = \bigotimes_{I=1}^4 \mathbb{T}_I^2$. Looking at the metric (4.8), the scalars A_I play the role of Kähler moduli parameterising the size of the four $\mathbb{T}_I^2 \subset \mathbb{T}^8$, whereas U_I are complex structure moduli parameterising the shape. Note that $\sqrt{|\mathbf{g}|} = (A_1 A_2 A_3 A_4)^{-1}$ in agreement with the previous identification of moduli fields in the M-theory duality frame. A similar identification holds in the type IIA with O2-planes duality frame of Section 3.2 that appears after dimensional reduction, except that the radius of the 8th circle on \mathbb{T}^8 is no longer interpreted geometrically but becomes the type IIA dilaton Φ . Further moving to other Op -plane duality frames involves the action of T-dualities which invert the radius of the internal circles upon which they act. As a result of this inversion, a single T-duality along a circle on $\mathbb{T}_I^2 \subset \mathbb{T}^8$ exchanges the role of the corresponding Kähler A_I and complex structure $\mathrm{Im}U_I$ moduli.

Constructing the scalar-dependent matrix in (4.2) from (4.6)-(4.7), and truncating away the vectors in the theory, the Lagrangian (2.2) reduces to an Einstein-scalar model of the form

$$\mathcal{L} = -\frac{1}{4} e R + \frac{1}{8} e \sum_{I=1}^4 \frac{(\partial B_I)^2 + (\partial A_I)^2}{A_I^2} + \frac{1}{8} e \sum_{I=1}^4 \frac{(\partial u_I)^2 + (\partial \mu_I)^2}{\mu_I^2} - e V , \quad (4.9)$$

where the scalar potential V depends on the M-theory/string fluxes compatible with the specific duality frame under consideration. From now on, we will refer to the \mathbb{Z}_2^2 invariant sector of half-maximal supergravity as the eight-chiral model. This model contains the eight dilatons (A_I, μ_I) associated with the Cartan subalgebra of $\mathrm{SO}(8, 8)$ and their partners (B_I, u_I) . From (4.6)-(4.7) one sees that $(A_I, \mu_I; u_I)$ belong to $\mathfrak{e} \in \mathrm{GL}(8)/\mathrm{SO}(8)$ whereas the axions B_I belong to the completion into $\mathrm{SO}(8, 8)/(\mathrm{SO}(8) \times \mathrm{SO}(8))$. The eight-chiral model presented here provides a natural extension of the models investigated in [4, 5] which only included the eight dilatons (A_I, μ_I) (see Section 4.2).

4.1.2 The $\mathrm{SO}(3)$ invariant sector: the RSTU-model

The second symmetry we will mod out the theory by is a $\mathrm{SO}(3)$ symmetry embedded in the $\mathrm{SO}(8, 8)$ duality group of half-maximal supergravity as

$$\mathrm{SO}(8, 8) \supset \mathrm{SO}(2, 2) \times \mathrm{SO}(6, 6) \supset \mathrm{SO}(2, 2) \times \mathrm{SO}(2, 2) \times \mathrm{SO}(3) . \quad (4.10)$$

It then follows from (4.10) that the commutant of $\mathrm{SO}(3)$ within $\mathrm{SO}(8, 8)$ is $\mathrm{SO}(2, 2) \times \mathrm{SO}(2, 2)$. Using the fact that $\mathrm{SO}(2, 2) \sim \mathrm{SL}(2) \times \mathrm{SL}(2)$, the scalar geometry describing the $\mathrm{SO}(3)$ invariant sector of half-maximal supergravity is given by

$$\mathcal{M}_{\mathrm{scal}} = \left[\frac{\mathrm{SL}(2)}{\mathrm{SO}(2)} \right]^4 \subset \frac{\mathrm{SO}(8, 8)}{\mathrm{SO}(8) \times \mathrm{SO}(8)} , \quad (4.11)$$

and consists of four copies of the Poincaré disk parameterised by four complex scalars which we will denote as R, S, T and U . It is worth mentioning here that the $\mathrm{SO}(3)$ invariant sector is

contained within the \mathbb{Z}_2^2 invariant sector previously discussed. More concretely, it is obtained upon the identifications

$$T_4 \equiv R \quad , \quad U_4 \equiv S \quad , \quad T_1 = T_2 = T_3 \equiv T \quad , \quad U_1 = U_2 = U_3 \equiv U \quad . \quad (4.12)$$

This sector of the theory – we will refer to it as the RSTU-model – is the three-dimensional analogue of the extensively studied STU-model in four dimensions [20, 21, 22, 23, 24, 25, 26, 27, 9]. The Einstein-scalar Lagrangian describing this sector is given by (4.9) subject to the identifications in (4.12).

4.2 Flux vacua in the eight-chiral model: type IIA with O2-planes

The eight-chiral model has previously been considered in [4, 6] within the context of type IIA orientifold reductions with O2-planes, discussed in Section 3.2. However, refs. [4, 6] set the axions to zero, *i.e.* $u_I = B_I = 0$ in (4.7) so that $\mathbf{b} = 0$ in (4.6). This followed from imposing an extra \mathbb{Z}_2^* symmetry on the models acting as

$$\mathbb{Z}_2^* : (y^1, y^2, y^3, y^4, y^5, y^6, y^7, y^8) \rightarrow (-y^1, y^2, -y^3, y^4, -y^5, y^6, -y^7, y^8) \quad . \quad (4.13)$$

In what follows it will prove convenient to redefine the eight dilatons (A_I, μ_I) entering the e_I matrices in (4.7) so that \mathbf{e} in (4.6) becomes

$$\begin{aligned} \mathbf{e} &= \text{diag} \left(\frac{1}{\sqrt{A_1 \mu_1}}, \sqrt{\frac{\mu_1}{A_1}}, \frac{1}{\sqrt{A_2 \mu_2}}, \sqrt{\frac{\mu_2}{A_2}}, \frac{1}{\sqrt{A_3 \mu_3}}, \sqrt{\frac{\mu_3}{A_3}}, \frac{1}{\sqrt{A_4 \mu_4}}, \sqrt{\frac{\mu_4}{A_4}} \right) \\ &\equiv \frac{1}{\sqrt{\sigma}} \text{diag} \left(\sqrt{\rho_1}, \sqrt{\rho_2}, \sqrt{\rho_3}, \sqrt{\rho_4}, \sqrt{\rho_5}, \sqrt{\rho_6}, \sqrt{\rho_7}, \frac{1}{\sqrt{\rho_1 \rho_2 \rho_3 \rho_4 \rho_5 \rho_6 \rho_7}} \right) , \end{aligned} \quad (4.14)$$

in terms of eight new dilatons (σ, ρ_m) with $m = 1, \dots, 7$. This redefinition will make some features of the type IIA flux vacua more manifest.

4.2.1 Fluxes and tadpoles

The $\mathbb{Z}_2^* \times \mathbb{Z}_2^2$ invariance of the supergravity models we have discussed is string-theoretically compatible with demanding G_2 holonomy on the internal space of the corresponding type IIA reductions (see [4, 6] for an explicit realisation). As a consequence, the $\mathbb{Z}_2^* \times \mathbb{Z}_2^2$ invariant gauge fluxes $H_{(3)}$ and $F_{(4)}$ have an expansion

$$H_{(3)} = \sum_{m=1}^7 h_m \varphi_{(3)}^m \quad , \quad F_{(4)} = \sum_{m=1}^7 f_m \varphi_{(4)}^m \quad (4.15)$$

in terms of flux parameters (h_m, f_m) and basis elements $\varphi_{(3)}^m$ and $\varphi_{(4)}^m$. These 7 + 7 basis elements specify the G_2 invariant three-form and four-form as

$$\begin{aligned} \varphi_{(3)} &= dy^{127} + dy^{347} + dy^{567} + dy^{235} + dy^{145} + dy^{136} + dy^{246} \quad , \\ \varphi_{(4)} &= dy^{3456} + dy^{1256} + dy^{1234} + dy^{1467} + dy^{2367} + dy^{2457} + dy^{1357} \quad , \end{aligned} \quad (4.16)$$

where we have introduced the notation $dy^{127} = dy^1 \wedge dy^2 \wedge dy^7$, etc.⁹ Together with the gauge fluxes in (4.15), the type IIA with O2-planes duality frame also allowed for the Romans mass

⁹Note that $\varphi_{(4)} = \star_{7d} \varphi_{(3)}$.

parameter $F_{(0)}$. Importantly, the set of QC's of half-maximal supergravity gave in this case the tadpole cancellation condition in (3.18), namely,

$$F_{(0)} H_{(3)} = 0, \quad (4.17)$$

related to the absence of O6/D6 sources in the compactification scheme. This condition is solved by setting

$$F_{(0)} = 0 \quad \text{or} \quad h_m = 0. \quad (4.18)$$

In other words, $F_{(0)}$ and $H_{(3)}$ cannot be simultaneously activated in a type IIA reduction with O2-planes while preserving half-maximal supersymmetry.

4.2.2 Mkw₃ flux vacua

An explicit computation of the scalar potential using the gauge fluxes in (4.15), the Romans mass $F_{(0)}$ and the e matrix in (4.14) yields

$$g^{-2} V_{\mathcal{N}=8} = \frac{\sigma^4}{32} \left[\sum_{m=1}^7 \left(\frac{h_m}{\sqrt{\rho_{(4)}^m}} - f_m \sqrt{\rho_{(4)}^m} \right)^2 \right] + \frac{\sigma^2}{32} F_{(0)}^2 \prod_{m=1}^7 \rho_m^2, \quad (4.19)$$

where we have introduced the short-hand notation $\rho_{(4)}^m$ ($m = 1, \dots, 7$) for the seven quartic terms $\rho_{(4)}^1 \equiv \rho^3 \rho^4 \rho^5 \rho^6$, $\rho_{(4)}^2 \equiv \rho^1 \rho^2 \rho^5 \rho^6$, \dots , $\rho_{(4)}^7 \equiv \rho^1 \rho^3 \rho^5 \rho^7$ in one-to-one correspondence with the basis elements for $\varphi_{(4)}$ in (4.16). As a result, the scalar potential in (4.19) is non-negative definite.¹⁰

Solving the constraint (4.17) by choosing $F_{(0)} = 0$ in (4.18) kills the second term in (4.19) and leaves an overall σ^4 dependence in the scalar potential. In this case, the extremisation of V with respect to σ forces the bracket in (4.19) to vanish. This gives rise to a Minkowski vacuum ($V_0 \equiv \langle V \rangle = 0$) at

$$\rho_{(4)}^m = \frac{h_m}{f_m}, \quad (4.21)$$

with the overall modulus σ being the (unstabilised) massless ‘‘no-scale’’ direction of [4]. Note also that, having $F_{(0)} = 0$, these Minkowski vacua can be embedded in the M-theory setup of Section 3.1. On the contrary, solving the constraint (4.17) by choosing $H_{(3)} = 0$ in (4.18) retains both the σ^4 and σ^2 terms in the scalar potential (4.19). In this case, since both terms are positive definite, the extremisation of V requires them to vanish separately. However, this cannot be achieved unless all the fluxes are zero (trivialising V) or the dilatons vanish (singular behaviour). Therefore, there are no vacua whenever $H_{(3)} = 0$.

4.2.3 Adding O6-planes to type IIA with O2-planes

Let us consider the possibility of adding O6-planes to the type IIA with O2 duality frame. The first type of O6-planes we will consider is the O6-plane discussed in Section 3.6. The location of this O6-plane was displayed in (3.48) and its associated orientifold action given in

¹⁰Up to a term accounting for O6/D6 sources (to be discussed in a moment), the scalar potential (4.19) maps into the one in [4, 6] via a simple redefinition of the eight dilatons:

$$(\rho_1, \rho_2, \rho_3, \rho_4, \rho_5, \rho_6, \rho_7) = e^{-\frac{1}{8}y + \frac{27}{16\sqrt{7}}x} \left(\frac{1}{\tilde{s}_1 \tilde{s}_4 \tilde{s}_6}, \tilde{s}_2 \tilde{s}_3 \tilde{s}_4 \tilde{s}_6, \frac{1}{\tilde{s}_2 \tilde{s}_4 \tilde{s}_5}, \tilde{s}_1 \tilde{s}_3 \tilde{s}_4 \tilde{s}_5, \frac{1}{\tilde{s}_3 \tilde{s}_5 \tilde{s}_6}, \tilde{s}_1 \tilde{s}_2 \tilde{s}_5 \tilde{s}_6, \frac{1}{\tilde{s}_1 \tilde{s}_2 \tilde{s}_3} \right), \quad (4.20)$$

together with $\sigma^8 = e^{3y + \frac{5\sqrt{7}}{2}x}$.

	External s-t			\mathbb{T}_1^2		\mathbb{T}_2^2		\mathbb{T}_3^2		\mathbb{S}^1
	x^0	x^1	x^2	y^1	y^2	y^3	y^4	y^5	y^6	y^7
O6 _* -plane	×	×	×	×		×		×		×
O6 _α -plane	×	×	×	×	×	×	×			
O6 _β -plane	×	×	×	×	×			×	×	
O6 _{αβ} -plane	×	×	×			×	×	×	×	
O6 _{*α} -plane	×	×	×		×		×	×		×
O6 _{*β} -plane	×	×	×		×	×			×	×
O6 _{*αβ} -plane	×	×	×	×			×		×	×
O2-planes	×	×	×							

Table 24: Multiple O2/O6-planes configuration. The combined action of the three independent O6_α, O6_β and O6_{*} orientifold actions, together with the O2-plane orientifold action, breaks half-maximal $\mathcal{N} = 8$ supergravity into minimal $\mathcal{N} = 1$ supergravity.

(3.49). In order to distinguish this O6-plane from other types (to be introduced in a moment) we will denote it as O6_{*} and its orientifold action as σ_{O6_*} . Note that σ_{O6_*} descends from \mathbb{Z}_2^* in (4.13). Together with the O6_{*}-plane, we will also introduce other types of O6-planes placed differently in the internal \mathbb{T}^7 . Following the construction in [4], let us consider two more types of O6-planes – we will denote them as O6_α and O6_β – whose orientifold actions $\sigma_{\text{O6}_\alpha}$ and σ_{O6_β} are inherited from $\mathbb{Z}_2^{(\alpha)}$ and $\mathbb{Z}_2^{(\beta)}$ in (4.3). Finally, we will also consider four more types of O6-planes that appear upon composition of O6_{*}, O6_α and O6_β. The resulting O2/O6-plane system is summarised in Table 24.

As it was shown in Section 3.2, the O2-plane in the type IIA with O2 duality frame was responsible for halving maximal into half-maximal supergravity, *i.e.* $\mathcal{N} = 16 \xrightarrow{\text{O2}} \mathcal{N} = 8$. Adding now the independent O6_{*}, O6_α and O6_β further triggers three additional halvings, namely

$$\mathcal{N} = 16 \xrightarrow{\text{O2}} \mathcal{N} = 8 \xrightarrow{\text{O6}_*} \mathcal{N} = 4 \xrightarrow{\text{O6}_\alpha} \mathcal{N} = 2 \xrightarrow{\text{O6}_\beta} \mathcal{N} = 1. \quad (4.22)$$

This breaking of supersymmetry from half-maximal to minimal due to the presence of O6-planes translates into the violation of the QC's of half-maximal supergravity in (4.17). From (3.1) one finds

$$-F_{(0)} H_{(3)} = J_{\text{O6/D6}} \neq 0, \quad (4.23)$$

so that both the Romans mass $F_{(0)}$ and the gauge flux $H_{(3)}$ can be activated simultaneously. Having O6-planes in the compactification scheme produces a new contribution to the scalar potential of the form

$$g^{-2} V_{\text{O6}} = \frac{\sigma^3}{16} F_{(0)} \sum_{m=1}^7 \left(h_m \rho_{(3)}^m \sqrt{\rho_{(4)}^m} \right), \quad (4.24)$$

where we have again introduced a short-hand notation $\rho_{(3)}^m$ ($m = 1, \dots, 7$) for the seven cubic terms $\rho_{(3)}^1 \equiv \rho^1 \rho^2 \rho^7$, $\rho_{(3)}^2 \equiv \rho^3 \rho^4 \rho^7$, \dots , $\rho_{(3)}^7 \equiv \rho^2 \rho^4 \rho^6$ in one-to-one correspondence with the basis elements of $\varphi_{(3)}$ in (4.16). The contribution in (4.24) was shown to descend from the

	External s-t			\mathbb{T}_1^2		\mathbb{T}_2^2		\mathbb{T}_3^2		\mathbb{S}^1	Internal Diff
	x^0	x^1	x^2	y^1	y^2	y^3	y^4	y^5	y^6	y^7	
O2-plane	×	×	×								SL(7)
O3-plane	×	×	×							×	SL(6)
O5-plane	×	×	×		×		×		×		SL(3) × SL(4)
O6-plane	×	×	×	×		×		×		×	SL(3) × SL(4)
O8-plane	×	×	×	×	×	×	×	×	×		SL(6)
O9-plane	×	×	×	×	×	×	×	×	×	×	SL(7)

Table 25: Single Op -plane duality frames compatible with the $\text{SO}(3)$ invariance of the RSTU-models. In all the cases the orientifold action $\mathcal{O}_{\mathbb{Z}_2}$ amounts to a discrete \mathbb{Z}_2 symmetry that halves maximal into half-maximal supergravity.

ten-dimensional DBI action of the O6/D6 sources in [4]. Being proportional to $F_{(0)} H_{(3)}$, the contribution V_{O6} is not part of the scalar potential of half-maximal supergravity due to the QC's in (4.17).

The scalar potential of the minimal supergravity models, *i.e.* $V_{\mathcal{N}=1} = V_{\mathcal{N}=8} + V_{\text{O6}}$,¹¹ has been shown to accommodate AdS_3 vacua ($V_0 \equiv \langle V \rangle < 0$) with interesting phenomenological properties, like scale separation between the AdS_3 scale $L^2 = -2/V_0$ and the scale of the internal space [4, 6] or between different internal cycles (anisotropy) [7].

Our next goal will be to show that the simpler half-maximal supergravity models arising from a single stack of Op -planes already suffice to accommodate both Mkw_3 and AdS_3 flux vacua with interesting phenomenology. We will illustrate this in the context of type IIB reductions with O5-planes.

4.3 Flux vacua in the RSTU-model: a type IIB with O5-planes appetizer

Let us now investigate half-maximal RSTU-models with a single class of Op -planes. The group-theoretical embedding (4.10) of the $\text{SO}(3)$ symmetry of the RSTU-models implies the set of branching rules summarised in Table 26. In particular, using the light-cone basis for $\text{SO}(8, 8)$, the vector index M of the **16** splits as

$$T_M = (T_A; T^A) \rightarrow (T_i, T_a, T_7, T_8; T^i, T^a, T^7, T^8), \quad (4.25)$$

with $a = 1, 3, 5$ and $i = 2, 4, 6$ transforming as two copies of the vector representation of $\text{SO}(3)$. As a result, there exist $\text{SO}(3)$ invariant tensors of the form

$$\delta_{ab}, \delta_{ij}, \delta_{ai}, \delta_{ia}, \delta_{77}, \delta_{78}, \delta_{88}, \epsilon_{abc}, \epsilon_{abk}, \epsilon_{ajk}, \epsilon_{ijk}, \quad (4.26)$$

which can be used to construct the set of $\text{SO}(3)$ invariant fluxes in a given Op -plane duality frame (see Table 25). As an example, we will focus the rest of the discussion on the type IIB with O5-plane duality frame presented in Section 3.5.

¹¹This potential matches the corresponding expression in [4, 6] upon the field redefinition (4.20).

SO(8, 8)	SL(2) × SL(2) × SO(6, 6)	SL(2) × SL(2) × SL(2) × SL(2) × SO(3)
16	(1, 1; 12) ⊕ (2, 2; 1)	(1, 1; 2, 2; 3) ⊕ (2, 2; 1, 1; 1)
1	(1, 1; 1)	(1, 1; 1, 1; 1)
135	(1, 1; 1) ⊕ (3, 3; 1) (1, 1; 77) ⊕ (2, 2; 12)	(1, 1; 1, 1; 1) ⊕ (3, 3; 1, 1; 1) [(1, 1; 3, 3; 1) ⊕ (1, 1; 1, 3; 3) ⊕ (1, 1; 3, 1; 3) (1, 1; 1, 1; 5) ⊕ (1, 1; 3, 3; 5)] ⊕ (2, 2; 2, 2; 3)
1820	(1, 1; 1) (1, 3; 66) ⊕ (3, 1; 66) (1, 1; 495) ⊕ (2, 2; 220) (2, 2; 12)	(1, 1; 1, 1; 1) [(1, 3; 1, 3; 1) ⊕ (1, 3; 3, 1; 1) ⊕ ...] [(3, 1; 1, 3; 1) ⊕ (3, 1; 3, 1; 1) ⊕ ...] [2(1, 1; 1, 1; 1) ⊕ (1, 1; 3, 3; 1) (1, 1; 1, 5; 1) ⊕ (1, 1; 5, 1; 1) ⊕ ...] [(2, 2; 2, 2; 1) ⊕ (2, 2; 4, 4; 1) ⊕ ...] (2, 2; 2, 2; 3)

Table 26: Branching rules for $\text{SO}(8, 8) \supset \text{SL}(2)^2 \times \text{SO}(6, 6) \supset \text{SL}(2)^4 \times \text{SO}(3)$. The ellipsis in the bottom-right box denote additional non-singlets under $\text{SO}(3)$.

4.3.1 Fluxes and tadpoles

The set of type IIB fluxes compatible with an O5-plane was presented in Table 11. In there, the $\text{SL}(4) \times \text{SL}(3)$ covariance of the O5-plane duality frame forced a splitting of the **16** of $\text{SO}(8, 8)$ of the form

$$T_M = (T_A; T^A) \rightarrow (T_i, T_{\hat{a}}, T_8; T^i, T^{\hat{a}}, T^8), \quad (4.27)$$

with $\hat{a} = 1, 3, 5, 7$ and $i = 2, 4, 6$. This splitting followed from the O5-plane orientifold action in (3.42). A quick comparison with the $\text{SO}(3)$ splitting in (4.25) then yields $\hat{a} = (a, 7)$. Using the $\text{SO}(3)$ invariant tensors in (4.26) one can construct the following set of $\text{SO}(3)$ invariant metric fluxes

$$\begin{aligned} \omega_{ab}{}^k &= \omega_1 \epsilon_{ab}{}^k & , & & \omega_{7a}{}^i &= \omega_2 \delta_a^i & , & & \omega_{7i}{}^a &= \omega_3 \delta_i^a & , \\ \omega_{ia}{}^7 &= \omega_4 \delta_{ia} & , & & \omega_{ia}{}^b &= \omega_5 \epsilon_{ia}{}^b & , & & \omega_{ij}{}^k &= \omega_6 \epsilon_{ij}{}^k & , \end{aligned} \quad (4.28)$$

as well as gauge fluxes

$$\begin{aligned} H_{abc} &= h_{31} \epsilon_{abc} & , & & H_{aij} &= h_{32} \epsilon_{aij} & , & & F_{ijk} &= f_{31} \epsilon_{ijk} & , & & F_{ia7} &= f_{32} \delta_{ia} & , & & F_{ibc} &= f_{33} \epsilon_{ibc} & , \\ & & & & F_{abij7} &= f_5 \delta_{ai} \delta_{bj} & , & & F_{abcijk7} &= f_7 \epsilon_{abc} \epsilon_{ijk} & , \end{aligned} \quad (4.29)$$

in terms of $6 + 7 = 13$ arbitrary flux parameters.¹² These 13 fluxes are a subset of the 158 embedding tensor components of half-maximal supergravity which are $\text{SO}(3)$ singlets (see

¹²A proper antisymmetrisation of indices is understood in (4.28) and (4.29).

Table 26). The rest are known as “non-geometric” fluxes (see [28]) and their study goes beyond the scope of this work.

The type IIB flux parameters in (4.28) and (4.29) are restricted by the QC’s of half-maximal supergravity in (2.12). An explicit computation yields

$$\begin{aligned} \omega_3 \omega_4 + \omega_5 (\omega_5 + \omega_6) &= 0 \quad , \quad \omega_4 (2\omega_5 + \omega_6) = 0 \\ \omega_1 \omega_3 - \omega_2 (\omega_5 + \omega_6) &= 0 \quad , \quad \omega_3 (2\omega_5 + \omega_6) = 0 \quad , \quad \omega_1 \omega_4 = 0 \quad , \\ \omega_2 \omega_4 - \omega_1 (\omega_5 + \omega_6) &= 0 \quad , \quad \omega_1 \omega_3 - 2\omega_2 \omega_5 = 0 \end{aligned} \quad (4.30)$$

which originate from the nilpotency condition ($D^2 = 0$) of the $D = d + \omega$ twisted exterior derivative on the internal space. In addition, one also finds

$$\omega_3 h_{32} = 0 \quad , \quad 2\omega_2 h_{32} - \omega_3 h_{31} = 0 \quad , \quad (4.31)$$

coming from $DH_{(3)} = 0$ and reflecting the absence of NS5-branes, as well as

$$\omega_2 f_{31} + (2\omega_5 + \omega_6) f_{32} + 2\omega_3 f_{33} = 0 \quad , \quad \omega_1 f_{31} - (2\omega_5 + \omega_6) f_{33} - 2\omega_4 f_{32} = 0 \quad , \quad (4.32)$$

originating from $DF_{(3)} = 0$ and reflecting the absence of O5/D5 sources different from the ones in (3.41). More concretely, the unrestricted O5/D5 sources entering (3.43) yield

$$DF_{(3)} \Big|_{dy^{\hat{a}} \wedge dy^{\hat{b}} \wedge dy^{\hat{c}} \wedge dy^{\hat{d}}} = \omega_1 f_{32} - \omega_2 f_{33} = J_{\text{O5/D5}} \quad . \quad (4.33)$$

Finally, there is a last condition of the form

$$3\omega_4 f_5 - h_{31} f_{31} + 3h_{32} f_{33} = 0 \quad , \quad (4.34)$$

which originates from $DF_{(5)} - H_{(3)} \wedge F_{(3)} = 0$ and reflects the absence of O3/D3 sources. The flux parameters must satisfy the algebraic conditions (4.30)-(4.32) and (4.34) for them to specify a well-defined gauging of half-maximal supergravity.

4.3.2 Exploring the landscape of Mkw₃ and AdS₃ flux vacua

Let us now investigate the vacuum structure of the RSTU-model constructed in the type IIB with O5 duality frame. The scalar potential of the half-maximal models is very lengthy (and not very enlightening) so we will not present it here explicitly. Still, it possesses some interesting features that we will exploit in order to explore its vacuum structure.

- The first feature is generic and follows from the general form of the scalar potential (2.18) in half-maximal supergravity: it is quadratic in the embedding tensor (flux parameters) and involves high powers of the scalar fields.
- The second property applies to the type IIB with O5 duality frame we are discussing here. The scalar potential of the RSTU-model turns out to be invariant under non-compact SL(2) transformations (scalings and shifts) on the complex scalars, of the form

$$R' = \lambda_R R + c_R \quad , \quad S' = \lambda_S S \quad , \quad T' = \lambda_T T + c_T \quad , \quad U' = \lambda_U U \quad , \quad (4.35)$$

with the λ ’s and c ’s being real parameters, accompanied by the following transformation of the metric fluxes

$$\begin{aligned} \omega'_1 &= \left(\frac{\lambda_R \lambda_S \lambda_T^3}{\lambda_U} \right)^{\frac{1}{2}} \omega_1 \quad , \quad \omega'_2 = \lambda_T^2 \omega_2 \quad , \quad \omega'_3 = \frac{\lambda_T}{\lambda_S} \omega_3 \quad , \\ \omega'_4 &= \frac{\lambda_R}{\lambda_U} \omega_4 \quad , \quad \omega'_5 = \left(\frac{\lambda_R \lambda_T}{\lambda_S \lambda_U} \right)^{\frac{1}{2}} \omega_5 \quad , \quad \omega'_6 = \left(\frac{\lambda_R \lambda_T}{\lambda_S \lambda_U} \right)^{\frac{1}{2}} \omega_6 \quad , \end{aligned} \quad (4.36)$$

ID	Type	SUSY	ω						$H_{(3)}$		$F_{(3)}$			$F_{(5)}$	$F_{(7)}$
			ω_1	ω_2	ω_3	ω_4	ω_5	ω_6	h_{31}	h_{32}	f_{31}	f_{32}	f_{33}	f_5	f_7
vac 1 †	Mkw ₃	$\mathcal{N} = 0, 4$	κ	ξ	0	0	0	0	0	0	0	κ	$-\xi$	0	0
vac 2 †		$\mathcal{N} = 0$	0	κ	0	0	0	0	0	0	0	0	$-\kappa$	0	0
vac 3 *		$\mathcal{N} = 0$	0	κ	κ	0	0	0	0	0	0	0	0	0	0
vac 4 *†	AdS ₃	$\mathcal{N} = 4$	0	0	0	0	0	κ	0	0	$\pm\kappa$	0	0	0	$-\kappa$
vac 5 *†		$\mathcal{N} = 0$	0	0	0	0	0	κ	0	0	$\pm\kappa$	0	0	0	κ
vac 6 *	AdS ₃	$\mathcal{N} = 3$	κ	0	0	0	$-\kappa$	κ	0	0	$\mp\kappa$	0	$\pm\kappa$	0	-2κ
vac 7 *		$\mathcal{N} = 1$	κ	0	0	0	$-\kappa$	κ	0	0	$\mp\kappa$	0	$\pm\kappa$	0	2κ
vac 8 *†	AdS ₃	$\mathcal{N} = 1$	0	0	0	0	$-\kappa$	κ	0	0	$\pm\kappa$	0	0	0	κ
vac 9 *†		$\mathcal{N} = 0$	0	0	0	0	$-\kappa$	κ	0	0	$\pm\kappa$	0	0	0	$-\kappa$
vac 10	AdS ₃	$\mathcal{N} = 0$	0	2κ	κ	0	0	0	0	0	κ	$\pm\kappa$	$-\kappa$	0	$\pm\kappa$
vac 11	AdS ₃	$\mathcal{N} = 0$	0	2κ	κ	0	0	0	0	0	κ	$\pm\kappa$	$-\kappa$	0	$\mp\kappa$
vac 12 *†	AdS ₃	$\mathcal{N} = 4$	0	0	$\pm\kappa$	$\pm\kappa$	κ	-2κ	0	0	$\pm 2\kappa$	0	0	0	2κ
vac 13 *†		$\mathcal{N} = 1$	0	0	$\pm\kappa$	$\pm\kappa$	κ	-2κ	0	0	$\pm 2\kappa$	0	0	0	-2κ
vac 14 *†		$\mathcal{N} = 0$	0	0	$\pm\kappa$	$\pm\kappa$	κ	-2κ	0	0	$\mp 2\kappa$	0	0	0	2κ
vac 15 *†		$\mathcal{N} = 0$	0	0	$\pm\kappa$	$\pm\kappa$	κ	-2κ	0	0	$\mp 2\kappa$	0	0	0	-2κ

Table 27: Fluxes in the type IIB with O5-planes duality frame producing a vacuum at the origin of moduli space, *i.e.* $\langle R \rangle = \langle S \rangle = \langle T \rangle = \langle U \rangle = i$. The **vac 1** is generically non-supersymmetric but becomes $\mathcal{N} = 4$ supersymmetric when $\kappa = \pm\xi$. The vacuum solutions with an asterisk * satisfy the extra self-dual constraint (4.41) and admit an embedding into maximal supergravity. Those with a dagger † satisfy the anti-self-dual condition (4.42). If both conditions are satisfied simultaneously then (4.43) holds.

and of the gauge fluxes

$$\begin{aligned}
h'_{31} &= \left(\frac{\lambda_R \lambda_S \lambda_T^3}{\lambda_U^3} \right)^{\frac{1}{2}} h_{31} \quad , \quad h'_{32} = \left(\frac{\lambda_R \lambda_T}{\lambda_S \lambda_U^3} \right)^{\frac{1}{2}} h_{32} \quad , \\
f'_{31} &= \frac{\lambda_R}{\lambda_S} f_{31} \quad , \quad f'_{32} = \left(\frac{\lambda_R \lambda_U \lambda_T^3}{\lambda_S} \right)^{\frac{1}{2}} f_{32} \quad , \quad f'_{33} = \lambda_R \lambda_T f_{33} \quad , \\
f'_5 &= \left(\frac{\lambda_R \lambda_T}{\lambda_S \lambda_U} \right)^{\frac{1}{2}} [\lambda_T f_5 + c_T (2\omega_5 + \omega_6)] \quad , \quad f'_7 = \left(\frac{\lambda_R \lambda_T}{\lambda_S \lambda_U^3} \right)^{\frac{1}{2}} [\lambda_T f_7 + 3 c_T h_{32}] \quad .
\end{aligned} \tag{4.37}$$

- The third feature, which actually can be inferred from (4.37), is that the scalar potential does not depend on the scalar $\text{Re}R = B_4$. This explains why the transformation of the gauge fluxes in (4.37) does not depend on the constant shift c_R in (4.35).

Charting the landscape of flux vacua: the strategy

The invariance of the scalar potential under the simultaneous transformations (4.35) and (4.36)-(4.37) allows us to set vacuum expectation values (VEV's) of the form

$$\langle R \rangle = i \quad , \quad \langle S \rangle = \langle u_4 \rangle + i \quad , \quad \langle T \rangle = i \quad , \quad \langle U \rangle = \langle u \rangle + i \quad , \quad (4.38)$$

when searching for the extrema of V without loss of generality. In other words, a given vacuum of the scalar potential can be brought to the form in (4.38) by applying a duality transformation at the expense of changing the flux configuration accordingly. Note that real shifts of S and U cannot be reabsorbed by a flux redefinition. In our scanning of critical points of the scalar potential we will restrict to vacua with

$$\langle u_4 \rangle = \langle u \rangle = 0 \quad . \quad (4.39)$$

This amounts to search for vacua at the origin of the moduli space, which can afterwards be moved to a different location in field space by performing the duality transformations in (4.35) at the expense of changing fluxes as in (4.36)-(4.37).

The main advantage of searching for critical points at the origin of moduli space is that the extremisation conditions

$$\partial V \Big|_{\langle R \rangle = \langle S \rangle = \langle T \rangle = \langle U \rangle = i} = 0 \quad , \quad (4.40)$$

become a set of algebraic equations which are quadratic on the flux parameters. In addition, there are the constraints (4.30)-(4.34) required by half-maximal supergravity which are also quadratic on the flux parameters. Therefore, putting together (4.40) and (4.30)-(4.34), one is left with a complicated algebraic set of quadratic conditions on the flux parameters. This system can be solved in full generality with the help of the algebra software SINGULAR [29]. The outcome is the set of 15 vacua summarised in Table 27. The gauge algebra induced by the fluxes that are activated in Table 27 is presented in the Appendix B.

Supersymmetry and stability of flux vacua

Table 27 contains the complete set of Mkw₃ ($V_0 \equiv \langle V \rangle = 0$) and AdS₃ ($V_0 \equiv \langle V \rangle < 0$) flux vacua located at the origin of moduli space, *i.e.*, $\langle R \rangle = \langle S \rangle = \langle T \rangle = \langle U \rangle = i$. For each vacuum, the complete set of gravitino masses – from which we extract the number \mathcal{N} of preserved supersymmetries – is presented in Table 28. The spectrum of scalar masses – from which we assess the perturbative stability of the solutions – is summarised in Table 29.

There are three Mkw₃ vacua which are generically non-supersymmetric. However, one of them – labelled **vac 1** in Table 27 – features an enhancement of supersymmetry to $\mathcal{N} = 4$ when the flux parameters are adjusted to $\kappa = \mp \xi$ so that four gravitini become massless (see Table 28). The remaining Mkw₃ vacua, labelled **vac 2** and **vac 3**, have the same gravitino (see Table 28) and scalar (see Table 29) mass spectra but, as we will see in a moment, **vac 3** is embeddable into maximal supergravity while **vac 2** is not. On the other hand, amongst the twelve AdS₃ vacua we find examples preserving $\mathcal{N} = 0, 1, 3, 4$ supersymmetry. Curiously, there are four groups of AdS₃ vacua in which each member has a different amount of preserved supersymmetry, and therefore different gravitino masses, but they all have the same spectrum of scalar fluctuations. These four groups are **vac 4,5**, **vac 6,7**, **vac 8,9** and **vac 12,13,14,15** in Tables 27, 28 and 29.

In order to assess the perturbative (in)stability of the non-supersymmetric vacua, we have computed the scalar mass spectrum at each flux vacuum including *all* the scalars in half-maximal supergravity (see Table 29). Surprisingly (at least to us), we find that only

ID	Gravitino spectrum
vac 1	$g^{-2}m_{3/2}^2 = \left[\frac{(\kappa \pm \xi)^2}{16} \right]_{(4)}$
vac 2	$g^{-2}m_{3/2}^2 = \left(\frac{9\kappa^2}{16} \right)_{(2)}, \left(\frac{\kappa^2}{16} \right)_{(6)}$
vac 3	
vac 4	$m_{3/2}^2 L^2 = 1_{(4)}, 9_{(4)}$
vac 5	$m_{3/2}^2 L^2 = 9_{(4)}, 25_{(4)}$
vac 6	$m_{3/2}^2 L^2 = 1_{(3)}, 9_{(4)}, 25_{(1)}$
vac 7	$m_{3/2}^2 L^2 = 1_{(1)}, 9_{(4)}, 25_{(3)}$
vac 8	$m_{3/2}^2 L^2 = 1_{(1)}, 9_{(1)}, 25_{(3)}, 49_{(3)}$
vac 9	$m_{3/2}^2 L^2 = 9_{(4)}, 25_{(4)}$
vac 10	$m_{3/2}^2 L^2 = 9_{(3)}, 49_{(3)}, 81_{(1)}, 169_{(1)}$
vac 11	$m_{3/2}^2 L^2 = 25_{(6)}, 49_{(1)}, 225_{(1)}$
vac 12	$m_{3/2}^2 L^2 = 1_{(4)}, 9_{(1)}, 25_{(3)}$
vac 13	$m_{3/2}^2 L^2 = 1_{(1)}, 9_{(4)}, 49_{(3)}$
vac 14	$m_{3/2}^2 L^2 = 9_{(7)}, 25_{(1)}$
vac 15	$m_{3/2}^2 L^2 = 9_{(1)}, 25_{(7)}$

Table 28: Gravitino masses at the type IIB with O5-planes vacua of Table 27. The subscript in $n_{(s)}$ denotes the multiplicity of the mass n in the spectrum. For the AdS₃ vacua we have normalised the gravitino spectrum using the AdS₃ radius $L^2 = -2/V_0$. Note that all the normalised gravitino masses are integer-valued for the AdS₃ vacua. A “massless” gravitino has $m_{3/2}^2 = 0$ in Mkw₃ and $m_{3/2}^2 L^2 = 1$ in AdS₃. The number \mathcal{N} of such gravitini correspond to the number of preserved supersymmetries at the corresponding vacuum in Table 27.

non-negative masses appear in the scalar spectra ensuring the perturbative stability of all the non-supersymmetric vacua within half-maximal supergravity. For those non-supersymmetric vacua that admit an embedding into maximal supergravity (marked with * in Table 27), unstable modes (tachyons) violating the Breitenlohner–Freedman (BF) bound for stability in AdS₃ [30] could still appear in the maximal theory which has a larger E₈₍₈₎/SO(16) scalar geometry.

Embedding into maximal supergravity

The embedding tensor configurations associated with vacua marked with an asterisk * in Table 27 satisfy the extra QC’s in (2.13) and specify a consistent gauging of maximal supergravity. More concretely, the first condition in (2.13) trivialises in our type IIB with O5 duality frame. The second condition in (2.13) is non-trivial and, by virtue of the self-dual (SD) nature of $[\Gamma^{M_1 \dots M_8}]_{\dot{A}\dot{B}}$, amounts to compute the self-dual part of the SO(8, 8) eight-form

ID	Scalar spectrum
vac 1	$g^{-2} m^2 = 0_{(30)}, \left(\frac{\kappa^2}{16}\right)_{(9)}, \left(\frac{\kappa^2}{4}\right)_{(9)}, \left(\frac{\xi^2}{4}\right)_{(9)}, \left(\frac{9\kappa^2}{16}\right)_{(1)}, \left[\frac{(\kappa-2\xi)^2}{16}\right]_{(3)}, \left[\frac{(\kappa+2\xi)^2}{16}\right]_{(3)}$
vac 2	$g^{-2} m^2 = \left(\frac{\kappa^2}{4}\right)_{(15)}, 0_{(49)}$
vac 3	
vac 4	$m^2 L^2 = 8_{(19)}, 0_{(45)}$ $\Delta = 4_{(19)}, 2_{(45)}$
vac 5	
vac 6	$m^2 L^2 = 8_{(10)}, 4_{(18)}, 0_{(36)}$ $\Delta = 4_{(10)}, (1 + \sqrt{5})_{(18)}, 2_{(36)}$
vac 7	
vac 8	$m^2 L^2 = 24_{(10)}, 8_{(25)}, 0_{(29)}$ $\Delta = 6_{(10)}, 4_{(25)}, 2_{(29)}$
vac 9	
vac 10	$m^2 L^2 = 80_{(3)}, 48_{(9)}, 24_{(4)}, 8_{(7)}, 0_{(41)}$ $\Delta = 10_{(3)}, 8_{(9)}, 6_{(4)}, 4_{(7)}, 2_{(41)}$
vac 11	
vac 12	$m^2 L^2 = 15_{(8)}, 8_{(19)}, 3_{(8)}, 0_{(29)}$ $\Delta = 5_{(8)}, 4_{(19)}, 3_{(8)}, 2_{(29)}$
vac 13	
vac 14	
vac 15	

Table 29: Scalar masses at the type IIB with O5-planes vacua of Table 27. The subscript in $n_{(s)}$ denotes the multiplicity of the mass n in the spectrum. For the AdS₃ vacua we have normalised the scalar spectrum using the AdS₃ radius $L^2 = -2/V_0$. In addition, for each AdS₃ vacuum, the conformal dimension Δ of the would-be CFT₂ dual operators are also indicated. These correspond to the larger root of $m^2 L^2 = \Delta(\Delta - 2)$. Note that all but **vac 6** and **vac 7** AdS₃ solutions come along with integer-valued Δ 's.

$\theta_{[M_1 M_2 M_3 M_4]} \theta_{[M_5 M_6 M_7 M_8]}$. The result is just a single condition

$$\theta_{[M_1 M_2 M_3 M_4]} \theta_{[M_5 M_6 M_7 M_8]} \Big|_{\text{SD}} = 0 \implies \omega_1 f_{32} - \omega_2 f_{33} = 0. \quad (4.41)$$

A direct comparison with (4.33) then shows that, as expected, an embedding into maximal supergravity is possible whenever the net charge of unrestricted O5/D5 sources of the type in (3.41) vanishes, namely, $J_{\text{O5/D5}} = 0$. These were precisely the sources causing the breaking of supersymmetry from maximal to half-maximal.

Although not necessary to have an embedding into maximal supergravity, it is also interesting to evaluate whether the anti-self-dual (ASD) part of the SO(8, 8) eight-form $\theta_{[M_1 M_2 M_3 M_4]} \theta_{[M_5 M_6 M_7 M_8]}$ vanishes. This implies two additional conditions which are purely geometrical since they only involve metric fluxes

$$\theta_{[M_1 M_2 M_3 M_4]} \theta_{[M_5 M_6 M_7 M_8]} \Big|_{\text{ASD}} = 0 \implies \begin{aligned} \omega_2 \omega_3 &= 0 \\ \omega_2 \omega_4 - 2 \omega_1 \omega_5 &= 0 \end{aligned}. \quad (4.42)$$

The embedding tensors associated with the flux vacua marked with a dagger † in Table 27 satisfy (4.42). Finally, if both (4.41) and (4.42) hold, the corresponding embedding tensor satisfies

$$\theta_{[M_1 M_2 M_3 M_4} \theta_{M_5 M_6 M_7 M_8]} = 0 . \quad (4.43)$$

As explained in [31], a generalised Scherk–Schwarz ansatz with twist matrices that obey the so-called *section constraint* of the $O(8,8)$ enhanced double field theory (DFT) can only reproduce gaugings whose embedding tensor satisfies (4.43). It would be interesting to understand the role of the geometric conditions (4.42) in the $O(8,8)$ -DFT context. This would also open up the possibility to undertake a full Kaluza–Klein spectral analysis of the corresponding AdS_3 vacua along the lines of [32, 33, 34], which we though leave to future work.

4.3.3 On the conformal dimension of dual operators and scale separation

In the context of type IIA supergravity, the supersymmetric and scale-separated AdS_4 vacua of DGKT [26] come along with a peculiar feature: the conformal dimensions of the would-be operators dual to the closed string sector are integer-valued [35] (see also [36, 37, 38]). More concretely, the authors of [35] showed that, independently of the details of the compactification, the conformal dimensions of the low-lying scalar primaries of the CFT_3 dual to supersymmetric AdS_4 DGKT flux vacua on general Calabi–Yau three-folds (CY₃'s) are always integer-valued. When considering instead non-supersymmetric configurations (see *e.g.* [38]), integer-valued conformal dimensions are not guaranteed but, in light of the AdS swampland conjecture [39], one might argue that such vacua are unstable and so not suitable for applying the AdS/CFT correspondence. More general type IIA flux models including not only gauge but also metric fluxes have been investigated in [40]. In this case, the conformal dimensions of the would-be CFT_3 operators dual to the light scalars defining the corresponding AdS_4 vacua cease to be (only) integer-valued. Differently from the type IIA DGKT-like vacua, such models with metric fluxes do not admit scale separation and one might then wonder whether integer-valued conformal dimensions are actually related to scale separation in four dimensions. Similar questions on supersymmetric AdS_4 flux vacua with scale separation and integer-valued conformal dimensions of would-be dual operators have been investigated in the mirror-symmetric type IIB context of [41]. When considering one-loop corrections to the complex-structure moduli space of the mirror-dual of DGKT, it was shown in [41] that the conformal dimensions of the would-be CFT_3 dual operators get also corrected and no longer take integer values. Only in the strict large complex structure regime where the aforementioned corrections are ignored, the integer-valued conformal dimensions of [35, 36, 37] are recovered. In addition, ref. [41] also considered supersymmetric AdS_4 vacua in type IIB models including non-geometric fluxes [28]. For these more exotic supersymmetric vacua, the conformal dimensions of the would-be dual operators were (numerically) determined to be non-integer-valued.

The above story about simultaneously having scale-separated AdS vacua and integer-valued conformal dimensions, as in DGKT, seems to change when moving to the context of $\text{AdS}_3/\text{CFT}_2$. In this context, various examples of $\mathcal{N} = 1$ supersymmetric AdS_3 flux vacua have been constructed upon compactification of type IIA [4] and type IIB [5] supergravity on (co-calibrated) G_2 orientifolds. For the parametrically scale-separated type IIA flux vacua of [4], it was noticed in [36] that the light single-trace operators of the would-be dual CFT_2 have non-integer (actually irrational) conformal dimensions. On the other hand, the type IIB flux vacua of [5] allow for integer-valued (and rational) conformal dimensions but do not exhibit a parametrically-controlled hierarchy between the size of the internal space and AdS_3 . Therefore, and differently from what happens in four dimensions, the phenomenon of having integer-valued conformal dimensions seems to be in tension with having scale-separated AdS flux vacua in three dimensions. In the rest of this section we will argue that the type IIB

		$\langle R \rangle$		$\langle S \rangle$		$\langle T \rangle$		$\langle U \rangle$		
ID	$V_0 \equiv \langle V \rangle$	B_4	A_4	u_4	μ_4	B	A	u	μ	
vac 1	0		$-\frac{f_{33}}{\omega_2} A$	0	$\frac{\omega_1}{f_{32}} \mu$			0		
vac 2	0		$-\frac{f_{33}}{\omega_2} A$	0				0		
vac 3	0			0	$\frac{\omega_2}{\omega_3} A^{-1}$			0		
vac 4	$-\frac{g^2}{32} \frac{\omega_6^6}{f_{31}^2 f_7^2}$		$-\frac{f_{31}^2 f_7}{\omega_6^3} \mu_4$	0		$\frac{f_5}{\omega_6}$	$-\frac{f_7}{\omega_6} \mu$	0		
vac 5			$\frac{f_{31}^2 f_7}{\omega_6^3} \mu_4$				$\frac{f_7}{\omega_6} \mu$			
vac 6	$-\frac{g^2}{2} \frac{\omega_6^6}{f_{31}^2 f_7^2}$		$\frac{\omega_1 f_{31}^2}{\omega_6^3} \mu^{-1}$	0		$-\frac{f_5}{\omega_6}$	$-\frac{f_7}{2\omega_6} \mu$	0		
vac 7							$\frac{2\omega_1}{f_7} \mu^{-1}$			$\frac{f_7}{2\omega_6} \mu$
vac 8	$-\frac{g^2}{32} \frac{\omega_6^6}{f_{31}^2 f_7^2}$		$\frac{f_{31}^2 f_7}{\omega_6^3} \mu_4$	0		$-\frac{f_5}{\omega_6}$	$\frac{f_7}{\omega_6} \mu$	0		
vac 9			$-\frac{f_{31}^2 f_7}{\omega_6^3} \mu_4$				$-\frac{f_7}{\omega_6} \mu$			
vac 10	$-\frac{g^2}{32} \frac{\omega_3^6 f_{33}^6}{f_{31}^2 f_{32}^2 f_7^2}$		$-\frac{f_{31} (f_{32}^3 f_7)^{\frac{1}{2}}}{\omega_3^2 f_{33}}$	0		$\frac{\omega_3 f_{33}^2}{f_{31} (f_{32}^3 f_7)^{\frac{1}{2}}}$	$-\frac{(f_{32}^3 f_7)^{\frac{1}{2}}}{\omega_3 f_{33}}$	0	$\left(\frac{f_{32}}{f_7}\right)^{\frac{1}{2}}$	
vac 11			$-\frac{f_{31} (-f_{32}^3 f_7)^{\frac{1}{2}}}{\omega_3^2 f_{33}}$				$\frac{\omega_3 f_{33}^2}{f_{31} (-f_{32}^3 f_7)^{\frac{1}{2}}}$		$-\frac{(-f_{32}^3 f_7)^{\frac{1}{2}}}{\omega_3 f_{33}}$	$\left(-\frac{f_{32}}{f_7}\right)^{\frac{1}{2}}$
vac 12	$-2g^2 \frac{\omega_5^6}{f_{31}^2 f_7^2}$		$\frac{f_{31} f_7}{4\omega_3 \omega_5} \mu$	0			$\frac{f_7}{2\omega_5} \mu$	0		
vac 13			$-\frac{f_{31} f_7}{4\omega_3 \omega_5} \mu$				$\frac{2\omega_5^2}{\omega_3 f_{31}} \mu$			$-\frac{f_7}{2\omega_5} \mu$
vac 14			$-\frac{f_{31} f_7}{4\omega_3 \omega_5} \mu$				$-\frac{2\omega_5^2}{\omega_3 f_{31}} \mu$			$\frac{f_7}{2\omega_5} \mu$
vac 15			$\frac{f_{31} f_7}{4\omega_3 \omega_5} \mu$				$-\frac{2\omega_5^2}{\omega_3 f_{31}} \mu$			$-\frac{f_7}{2\omega_5} \mu$

Table 30: Type IIB fields as a function of the fluxes after undoing the “go to the origin” process. The empty boxes in the table indicate that the corresponding field is not fixed by the extremisation of the scalar potential V . For a given vacuum solution, the fluxes appearing in the table must be considered as independent. Those which do not appear are either zero or, otherwise, can be expressed in terms of the independent ones. In particular: $\omega_5 = -\omega_6$ and $f_{33} = -\omega_1 f_{31} / \omega_6$ at **vac 6,7**; $\omega_5 = -\omega_6$ at **vac 8,9**; $\omega_2 = -2\omega_3 f_{33} / f_{31}$ at **vac 10,11**; $\omega_4 = \omega_5^2 / \omega_3$ and $\omega_6 = -2\omega_5$ at **vac 12,13,14,15**. These additional relations are necessary for the QC’s (4.30)-(4.32) and (4.34) of half-maximal supergravity to hold.

vacua labelled as **vac 10** and **vac 11** in Table 27 – and which are non-supersymmetric – come along with integer-valued conformal dimensions of would-be dual operators and, at the same time, allow for parametrically-controlled scale separation.

AdS₃ vacua: on integer-valued conformal dimensions

The complete scalar mass spectra of the type IIB with O5 flux vacua of Table 27 are reported in Table 29. Quite surprisingly, and with the exception of **vac 6,7**, the rest of AdS₃ vacua

therein exhibit all integer-valued conformal dimensions for the would-be low-lying CFT_2 dual operators. This observation places itself within the on-going discussion on the relation between (non-)supersymmetric AdS flux vacua and their CFT potential description, and the attempts of understanding what makes the aforementioned conformal dimensions to be integer-valued. By looking at Table 27, one observes that the groups **vac 4,5** and **vac 12,13,14,15** possess a member with $\mathcal{N} = 4$ supersymmetry. This large amount of supersymmetry might help to understand the integer nature of the Δ 's at those groups of AdS_3 vacua. However, the groups **vac 8,9** and **vac 10,11** have members with smaller $\mathcal{N} = 1$ or $\mathcal{N} = 0$ supersymmetry, respectively, making the appearance of integer-valued Δ 's more puzzling in these cases. It would be interesting to understand the ultimate reason behind these spectral structures.

AdS₃ vacua: on scale separation and (an)isotropy

Let us now look at some phenomenological aspects of the set of type IIB vacua with a single O5-plane listed in Table 27. In particular, we will deal with the phenomenon of scale separation between the AdS_3 external space and the internal space, as well as between different internal cycles (anisotropy). To this end we will resort to the more standard picture of having the VEV's of the scalar fields as a function of the flux parameters. This amounts to undo the duality transformation (4.35)-(4.37) that we implemented to go to the origin of moduli space when extremising the potential. The result is summarised in Table 30.

The four dilatons in the RSTU-model allow for two characteristic length scales in the compactification. Accordingly to the factorisation in (3.22), these correspond to the size of the internal $\mathbb{T}^6 = \mathbb{T}_1^2 \times \mathbb{T}_2^2 \times \mathbb{T}_3^2 \subset \mathbb{T}^7$ and the one of the additional circle $\mathbb{S}^1 \subset \mathbb{T}^7$. Upon the exchange of Kähler and complex structure moduli in (4.14), which follows from the three T-dualities taking an O2-plane into the O5-plane of interest, such characteristic length scales are respectively identified as

$$L_6^2 = \frac{1}{\mu} \quad \text{and} \quad L_1^2 = \frac{1}{A_4 \mu_4} . \quad (4.44)$$

Importantly, we observe that **vac 10** and **vac 11** in Table 30 stabilise all the dilatons in the RSTU-model so that the two scales in (4.46) become well-defined and fully determined by the flux parameters.¹³ In particular, imposing

$$\pm f_{32} f_7 > 0 \quad , \quad \omega_3 f_{31} > 0 \quad , \quad \omega_3 f_{33} < 0 \quad \text{and} \quad f_{31} f_{33} < 0 \quad (4.45)$$

for the solutions to be physically acceptable, one finds

$$L^2 = \frac{64 (f_{32}^3 f_7)^2}{g^2 (\omega_3 f_{33})^6} f_{31}^2 \quad , \quad L_6^2 = \frac{1}{\mu} = \left(\pm \frac{f_7}{f_{32}} \right)^{\frac{1}{2}} \quad , \quad L_1^2 = \frac{1}{A_4 \mu_4} = -\frac{\omega_3}{f_{33}} \quad , \quad (4.46)$$

with the upper (lower) choice of sign for the **vac 10** (**vac 11**) solution and where, as before, $L^2 = -2/V_0$ denotes the AdS_3 radius.¹⁴ In addition, the string coupling is given by

$$g_s^2 \propto \frac{\mu_4}{A_4} = \frac{(-\omega_3 f_{33})^3}{(f_{32}^3 f_7)} \frac{4}{f_{31}^2} \quad , \quad (4.47)$$

and can be set to a small value while also guaranteeing control over the (string-frame) high-curvature corrections. The three scales in (4.46) can be tuned separately: after (ω_3, f_{33}) set L_1 and (f_7, f_{32}) set L_6 , the remaining flux f_{31} scales L and g_s inversely to each other.

¹³The axions B and B_4 are potentially stabilised by other, *e.g.* quantum, effects.

¹⁴Here all flux parameters are meant to be quantised, *i.e.*, $f_{(p)} = (2\pi)^{p-1} N_{(p)}$ with $N_{(p)} \in \mathbb{Z}$.

Then we can achieve any kind of parametrically-controlled scale separation between AdS_3 , \mathbb{T}^6 and \mathbb{S}^1 while keeping g_s small. For example, we can tune fluxes so that $L \gg L_6 \sim L_1$ yielding a scale-separated vacuum with an isotropic \mathbb{T}^7 . Another interesting case is to tune fluxes such that $L \sim L_1 \gg L_6$ so that the internal torus becomes anisotropic and the effective description of the vacuum is of the type $\text{AdS}_3 \times \mathbb{S}^1$ investigated in [7]. Lastly, for **vac 10** and **vac 11**, the tadpole cancellation condition (4.33) becomes

$$J_{\text{O5/D5}} = 2 (\omega_3 f_{33}) \left(\frac{f_{33}}{f_{31}} \right) > 0 , \quad (4.48)$$

requiring the presence of D5-branes in the compactification scheme.

5 Summary and future directions

In this work we have identified which $\mathcal{N} = 8$ (half-maximal) supergravities in three dimensions arise from type II orientifold reductions including background gauge and metric fluxes, Op -planes and Dp -branes. Demanding half-maximal supersymmetry only allowed for a single type of coincident Op -planes/ Dp -branes, *e.g.* type IIA with O2/D2 or type IIB with O5/D5 . Relying on group-theoretical arguments, we presented the dictionary between M/string theory fluxes and the embedding tensor of the three-dimensional half-maximal supergravity that results from the compactification. This was done for all the possible type IIA and type IIB duality frames (see Section 3). Formulating the various type II flux models within the embedding tensor formalism [10, 11] allowed us to initiate a systematic study of their vacuum structure. This also gave us the possibility to compute the complete scalar mass spectrum within half-maximal supergravity and to establish connections with the Swampland Program, for example, with the AdS conjecture [39].

We then focused on the particular type IIB with O5 duality frame and considered its $\text{SO}(3)$ invariant sector, deriving what we dubbed the RSTU-model. Making a combined use of the method put forward in [9] and of algebraic geometry techniques, we investigated the landscape of the RSTU-model. We found a surprisingly rich structure of vacua consisting of Mkw_3 and AdS_3 solutions with and without supersymmetry (see Table 27). In all the vacua, the masses of all the scalar fields in the coset geometry (2.1) of half-maximal supergravity turned out to be non-negative definite, establishing the perturbative stability of all the vacua (irrespective of supersymmetry) within half-maximal supergravity. Some of these vacua – in particular the ones marked with $*$ in Table 27 – can be embedded into maximal supergravity since the corresponding embedding tensors satisfy the self-duality condition (4.41). In other words, they are vacua of a maximal $\mathcal{N} = 16$ supergravity in three dimensions. In the absence of supersymmetry, it would be important to compute the masses of all the scalar fields in the $\text{E}_{8(8)}/\text{SO}(16)$ scalar geometry of the maximal theory in order to assess the stability of such solutions. In any case, for non-supersymmetric vacua like **vac 10** and **vac 11** in Table 27 which cannot be embedded into maximal supergravity due to the need of O5/D5 sources to cancel the flux-induced tadpole (4.48), we have established their perturbative stability within half-maximal supergravity. It would be interesting to investigate mechanisms for such vacua to decay in light of the AdS conjecture [39]. Besides being perturbatively stable within half-maximal supergravity, **vac 10** and **vac 11** display two interesting properties. Firstly, they provide examples of AdS vacua with parametrically-controlled scale separation between AdS_3 and the internal cycles \mathbb{T}^6 and \mathbb{S}^1 within the regime of validity of the supergravity approximation and in a classical type IIB context. Secondly, and despite being non-supersymmetric AdS_3 vacua, all the would-be dual CFT_2 operators turned out to have integer-valued conformal dimensions. These results motivate a systematic exploration of the landscape of vacua of

the various half-maximal flux models presented in Section 3. It would also be interesting to have a (real) superpotential formulation thereof along the lines of [42] and [4, 5].

There are some directions we would like to investigate in the future. For example, it would be interesting to explore lower-supersymmetric scenarios arising from the simultaneous presence of Op -planes (and Dp -branes) of various types, *e.g.*, type IIA with three types of $O4$ -planes breaking supersymmetry down to $\mathcal{N} = 2$, or type IIA with $O2/O6$ -planes to capture the $\mathcal{N} = 1$ models of [4, 6, 7]. Having a superpotential formulation of the simple flux models presented in this work would provide a good starting point to construct less supersymmetric models. These less supersymmetric models would have new terms in the superpotential enriching the dynamics of the moduli fields and, perhaps, opening the possibility to realise the phenomenon of fixed-point annihilation [43, 44] whose top-down realisation is still lacking.¹⁵ In addition to this, and for the flux models satisfying the algebraic condition (4.43), it would be interesting to reformulate them in an $O(8, 8)$ -DFT context. This would open up the possibility to carry out a systematic study of KK modes at the corresponding AdS_3 flux vacua along the lines of [45, 46, 47]. Tracking the masses of the KK modes while moving the AdS_3 vacua in field space by tuning the flux parameters would allow us to perform quantitative tests of the distance conjecture of [48]. Another interesting question has to do with the smearing of the sources in the AdS_3 flux vacua we have presented. We are assuming that smeared Op -planes are a consistent ingredient of flux compactifications and that there exists a full-fledged solution in string theory that is described by such an approximation at low energies. Building upon the results in, *e.g.*, [49, 50, 51, 52], it would be interesting to take into account the backreaction of such Op -planes (order by order in g_s) and see whether it can always be considered as a small perturbation around the smeared solution. Due to the non-linear character of the backreaction near the Op -planes, a conclusive answer can not be given yet. We hope to come back to these and other related issues in the near future.

Acknowledgements

We are grateful to Gabriel Larios and Fotis Farakos for interesting conversations. This work is supported by the Spanish national grant MCIU-22-PID2021-123021NB-I00.

A Spinorial representations and Γ -matrices of $SO(8)_R$

In this appendix we collect our conventions for the set of Γ -matrices of $SO(8)_R$ (we adopt conventions in Appendix A of [53]). In the Majorana representation they are of the form

$$[\Gamma^{\mathcal{I}}]_{\mu}{}^{\nu} = \begin{pmatrix} 0 & [\gamma^{\mathcal{I}}]_{\mathcal{A}\dot{\mathcal{B}}} \\ [\bar{\gamma}^{\mathcal{I}}]_{\dot{\mathcal{A}}\mathcal{B}} & 0 \end{pmatrix}, \quad (\text{A.1})$$

where $\mathcal{I} = 1, \dots, 8$ is the vector index of $SO(8)_R$ and $\mathcal{A}(\dot{\mathcal{A}}) = 1, \dots, 8$ is the left-handed (right-handed) Majorana-Weyl index. Majorana indices are lowered/raised using the invariant matrix $\mathcal{C} \equiv \mathcal{C}_{\mu\nu}$ and its inverse $\mathcal{C}^{-1} \equiv \mathcal{C}^{\mu\nu}$, with

$$\mathcal{C}_{\mu\nu} = \begin{pmatrix} \mathbb{I}_{\mathcal{A}\mathcal{B}} & 0 \\ 0 & \mathbb{I}_{\dot{\mathcal{A}}\dot{\mathcal{B}}} \end{pmatrix}. \quad (\text{A.2})$$

¹⁵The phenomenon of fixed-point annihilation is holographically described by the merging (and subsequent disappearance) of two AdS vacua in field space as a consequence of the tuning of a parameter in the superpotential. This phenomenon does not take place in the RSTU-model of Section 4.3, as it can be deduced by looking at Table 30.

This implies that left-handed and right-handed Majorana-Weyl indices are trivially raised and lowered. Lastly, using the above set of $\text{SO}(8)_R$ invariant matrices, $\Gamma^{(p)}$ -forms can be introduced as

$$[\Gamma^{\mathcal{I}_1 \dots \mathcal{I}_p}] = \Gamma^{[\mathcal{I}_1} \dots \Gamma^{\mathcal{I}_p]} \mathcal{C} . \quad (\text{A.3})$$

All the results in the main text have been derived using the following set of $[\gamma^{\mathcal{I}}]_{\mathcal{A}\dot{\mathcal{B}}}$ matrices

$$\begin{aligned} \gamma^1 &= i\sigma_2 \otimes i\sigma_2 \otimes i\sigma_2 & , & & \gamma^2 &= \mathbb{I}_2 \otimes \sigma_3 \otimes i\sigma_2 & , \\ \gamma^3 &= \mathbb{I}_2 \otimes \sigma_1 \otimes i\sigma_2 & , & & \gamma^4 &= \sigma_1 \otimes i\sigma_2 \otimes \mathbb{I}_2 & , \\ \gamma^5 &= \sigma_3 \otimes i\sigma_2 \otimes \mathbb{I}_2 & , & & \gamma^6 &= i\sigma_2 \otimes \mathbb{I}_2 \otimes \sigma_1 & , \\ \gamma^7 &= i\sigma_2 \otimes \mathbb{I}_2 \otimes \sigma_3 & , & & \gamma^8 &= \mathbb{I}_2 \otimes \mathbb{I}_2 \otimes \mathbb{I}_2 & , \end{aligned} \quad (\text{A.4})$$

in terms of the Pauli matrices $\sigma_{1,2,3}$. It then follows that $\bar{\gamma}^{\mathcal{I}} = (\gamma^{\mathcal{I}})^T$ and the following Clifford algebra holds

$$[\gamma^{\mathcal{I}}]_{\mathcal{A}\dot{\mathcal{A}}} [\bar{\gamma}^{\mathcal{J}}]^{\dot{\mathcal{A}}\mathcal{B}} + [\gamma^{\mathcal{J}}]_{\mathcal{A}\dot{\mathcal{A}}} [\bar{\gamma}^{\mathcal{I}}]^{\dot{\mathcal{A}}\mathcal{B}} = 2\delta^{\mathcal{I}\mathcal{J}} \delta_{\mathcal{A}}^{\mathcal{B}} . \quad (\text{A.5})$$

Vector indices of $\text{SO}(8)_R$ are then trivially raised/lowered too. Finally, the set of $\text{SO}(8)_R$ invariant $\gamma^{(p)}$ -forms entering (2.17) are straightforwardly extracted from (A.3). They are of the form

$$\begin{aligned} p=4 : & \quad [\gamma^{\mathcal{I}\mathcal{J}\mathcal{K}\mathcal{L}}]_{\mathcal{A}\mathcal{B}} = \gamma^{[\mathcal{I}\bar{\gamma}^{\mathcal{J}}\gamma^{\mathcal{K}}\bar{\gamma}^{\mathcal{L}}]} & , & & [\gamma^{\mathcal{I}\mathcal{J}\mathcal{K}\mathcal{L}}]_{\dot{\mathcal{A}}\dot{\mathcal{B}}} &= \bar{\gamma}^{[\mathcal{I}\gamma^{\mathcal{J}}\bar{\gamma}^{\mathcal{K}}\gamma^{\mathcal{L}}]} & , \\ p=3 : & \quad [\gamma^{\mathcal{I}\mathcal{J}\mathcal{K}}]_{\mathcal{A}\dot{\mathcal{B}}} = \gamma^{[\mathcal{I}\bar{\gamma}^{\mathcal{J}}\gamma^{\mathcal{K}}]} & , & & & & \\ p=2 : & \quad [\gamma^{\mathcal{I}\mathcal{J}}]_{\dot{\mathcal{A}}\dot{\mathcal{B}}} = \bar{\gamma}^{[\mathcal{I}\gamma^{\mathcal{J}}]} & . & & & & \end{aligned} \quad (\text{A.6})$$

B Gauge algebra for the type IIB with O5 vacua in Table 27

The embedding tensors giving rise to the type IIB with O5 flux vacua in Table 27 generically activate 73 (out of the 120) generators in the gauge algebra (2.11). These are generators of the form

$$X^{AB} \quad , \quad X_m{}^i \quad , \quad X_{\hat{a}}{}^{\hat{b}} \quad \text{and} \quad X^8{}_A . \quad (\text{B.1})$$

The gauge algebra they span is described by the following non-trivial commutation relations induced by the fluxes

$$\begin{aligned} [X^{ij}, X^{k\hat{c}}] &= -2\tilde{\omega}_{\hat{c}\hat{d}}{}^{ij} X^k{}_{\hat{d}} & , & & & & \\ [X^{ij}, X^{k8}] &= -2\omega_{ij}{}^h X^{kh} - 2\tilde{F}_7\epsilon^{ijk} X^8{}_8 - 2\tilde{F}_7\epsilon^{ijh} X^k{}_h & , & & & & \\ [X^{ij}, X^{\hat{c}\hat{d}}] &= 2\tilde{\omega}_{\hat{c}\hat{e}}{}^{ij} X^{\hat{d}}{}_{\hat{e}} - (c \leftrightarrow d) & , & & & & \\ [X^{ij}, X^{\hat{c}8}] &= -2\omega_{ij}{}^k X^{\hat{c}k} + 2\tilde{\omega}_{\hat{a}\hat{c}}{}^{ih} X^k{}_h & , & & & & \\ [X^{i\hat{a}}, X^{k8}] &= 2\omega_{\hat{b}i}{}^{\hat{a}} X^k{}_{\hat{b}} - 2\tilde{\omega}_{\hat{a}\hat{b}}{}^{ik} X^8{}_{\hat{b}} + 2\tilde{F}_{(3)i\hat{a}\hat{b}} X^k{}_{\hat{b}} & , & & & & \\ [X^{i\hat{a}}, X^{\hat{c}8}] &= 2\omega_{\hat{b}i}{}^{\hat{a}} X^{\hat{c}\hat{b}} + 2\tilde{F}_{(3)i\hat{a}\hat{c}} X^8{}_8 + 2\tilde{\omega}_{\hat{a}\hat{c}}{}^i X^8{}_j + 2\tilde{F}_{(3)i\hat{a}\hat{d}} X^{\hat{c}}{}_{\hat{d}} & , & & & & \\ [X^{i8}, X^{k8}] &= -2\omega_{ik}{}^h X^{8h} - 2\tilde{F}_0 X^{ik} - 2\tilde{F}_7\epsilon^{ikh} X^8{}_h & , & & & & \\ [X^{i8}, X^{\hat{c}\hat{d}}] &= 2\omega_{\hat{b}i}{}^{\hat{c}} X^{\hat{d}\hat{b}} + 2\tilde{F}_{(3)i\hat{c}\hat{b}} X^{\hat{d}}{}_{\hat{b}} - (c \leftrightarrow d) & , & & & & \\ [X^{i8}, X^{\hat{c}8}] &= 2\omega_{\hat{b}i}{}^{\hat{c}} X^{8\hat{b}} + 2\tilde{F}_0 X^{\hat{c}i} + 2\tilde{F}_{(3)i\hat{c}\hat{b}} X^8{}_{\hat{b}} & , & & & & \\ [X^{\hat{a}8}, X^{\hat{c}8}] &= -2\tilde{F}_0 X^{\hat{a}\hat{c}} + 2\tilde{F}_{(3)i\hat{a}\hat{c}} X^8{}_i & , & & & & \end{aligned} \quad (\text{B.2})$$

and

$$\begin{aligned}
[X_i^j, X^{k8}] &= -2\omega_{jh}^i X^k{}_h - 2\omega_{jk}^i X^8{}_8 \quad , \\
[X_i^8, X^{kh}] &= -2\omega_{kj}^i X^h{}_j - (k \leftrightarrow h) \quad , \\
[X_i^8, X^{k8}] &= -2\omega_{kh}^i X^8{}_h + 2\tilde{F}_0 X^k{}_i + 2\delta_i^k \tilde{F}_0 X^8{}_8 \quad , \\
[X_{\hat{a}}^i, X^{k8}] &= 2\omega_{\hat{a}i}^{\hat{b}} X^k{}_{\hat{b}} \quad , \\
[X_{\hat{a}}^i, X^{\hat{c}8}] &= 2\omega_{\hat{a}i}^{\hat{d}} X^{\hat{c}}{}_{\hat{d}} + 2\omega_{\hat{a}i}^{\hat{c}} X^8{}_8 \quad , \\
[X_{\hat{a}}^{\hat{b}}, X^{k8}] &= -2\omega_{\hat{a}j}^{\hat{b}} X^k{}_j - 2\omega_{\hat{a}k}^{\hat{b}} X^8{}_8 \quad , \\
[X_{\hat{a}}^8, X^{kh}] &= 2\omega_{\hat{a}k}^{\hat{b}} X^h{}_{\hat{b}} - (k \leftrightarrow h) \quad , \\
[X_{\hat{a}}^8, X^{k\hat{c}}] &= 2\omega_{\hat{a}k}^{\hat{d}} X^{\hat{c}}{}_{\hat{d}} + 2\omega_{\hat{a}h}^{\hat{c}} X^k{}_h \quad , \\
[X_{\hat{a}}^8, X^{k8}] &= 2\omega_{\hat{a}k}^{\hat{b}} X^8{}_{\hat{b}} + 2\tilde{F}_0 X^k{}_{\hat{a}} \quad , \\
[X_{\hat{a}}^8, X^{\hat{c}8}] &= -2\omega_{\hat{a}i}^{\hat{c}} X^8{}_i + 2\tilde{F}_0 X^{\hat{c}}{}_{\hat{a}} + 2\delta_{\hat{a}}^{\hat{c}} \tilde{F}_0 X^8{}_8 \quad ,
\end{aligned} \tag{B.3}$$

where we have introduced the flux-dependent quantities

$$\begin{aligned}
\tilde{\omega}_{\hat{a}\hat{b}}{}^{ij} &\equiv \frac{1}{2!} \epsilon^{\hat{a}\hat{b}\hat{c}\hat{d}} \epsilon^{ijk} \omega_{\hat{c}\hat{d}}{}^k \quad , & \tilde{F}_7 &\equiv \frac{1}{3!4!} \epsilon^{\hat{a}\hat{b}\hat{c}\hat{d}} \epsilon^{ijk} F_{\hat{a}\hat{b}\hat{c}\hat{d}ijk} \quad , \\
\tilde{F}_{(3)i\hat{a}\hat{b}} &\equiv \frac{1}{2!} \epsilon^{\hat{a}\hat{b}\hat{c}\hat{d}} F_{i\hat{c}\hat{d}} \quad , & \tilde{F}_0 &\equiv \frac{1}{3!} \epsilon^{ijk} F_{ijk} \quad .
\end{aligned} \tag{B.4}$$

References

- [1] E. Palti, *The Swampland: Introduction and Review*, *Fortsch. Phys.* **67** (2019) 1900037, [[1903.06239](#)].
- [2] N. Marcus and J. H. Schwarz, *Three-Dimensional Supergravity Theories*, *Nucl. Phys. B* **228** (1983) 145.
- [3] M. Grana, *Flux compactifications in string theory: A Comprehensive review*, *Phys.Rept.* **423** (2006) 91–158, [[hep-th/0509003](#)].
- [4] F. Farakos, G. Tringas and T. Van Riet, *No-scale and scale-separated flux vacua from IIA on G2 orientifolds*, *Eur. Phys. J. C* **80** (2020) 659, [[2005.05246](#)].
- [5] M. Emelin, F. Farakos and G. Tringas, *Three-dimensional flux vacua from IIB on co-calibrated G2 orientifolds*, *Eur. Phys. J. C* **81** (2021) 456, [[2103.03282](#)].
- [6] F. Farakos, M. Morittu and G. Tringas, *On/off scale separation*, *JHEP* **10** (2023) 067, [[2304.14372](#)].
- [7] F. Farakos and M. Morittu, *Scale-separated AdS₃ × S¹ vacua from IIA orientifolds*, *Eur. Phys. J. C* **84** (2024) 98, [[2311.08991](#)].
- [8] N. S. Deger, C. Eloy and H. Samtleben, *N = (8, 0) AdS vacua of three-dimensional supergravity*, *JHEP* **10** (2019) 145, [[1907.12764](#)].
- [9] G. Dibitetto, A. Guarino and D. Roest, *Charting the landscape of N=4 flux compactifications*, *JHEP* **1103** (2011) 137, [[1102.0239](#)].
- [10] H. Nicolai and H. Samtleben, *N=8 matter coupled AdS(3) supergravities*, *Phys. Lett. B* **514** (2001) 165–172, [[hep-th/0106153](#)].
- [11] B. de Wit, I. Herger and H. Samtleben, *Gauged locally supersymmetric D = 3 nonlinear sigma models*, *Nucl. Phys. B* **671** (2003) 175–216, [[hep-th/0307006](#)].

- [12] P. Koerber and D. Tsimpis, *Supersymmetric sources, integrability and generalized-structure compactifications*, *JHEP* **08** (2007) 082, [[0706.1244](#)].
- [13] J. Scherk and J. H. Schwarz, *How to Get Masses from Extra Dimensions*, *Nucl.Phys.* **B153** (1979) 61–88.
- [14] E. Bergshoeff, U. Gran, R. Linares, M. Nielsen, T. Ortin and D. Roest, *The Bianchi classification of maximal $D = 8$ gauged supergravities*, *Class. Quant. Grav.* **20** (2003) 3997–4014, [[hep-th/0306179](#)].
- [15] C. M. Hull and R. A. Reid-Edwards, *Flux compactifications of string theory on twisted tori*, *Fortsch. Phys.* **57** (2009) 862–894, [[hep-th/0503114](#)].
- [16] J.-P. Derendinger, P. M. Petropoulos and N. Prezas, *Axionic symmetry gaugings in $N=4$ supergravities and their higher-dimensional origin*, *Nucl. Phys. B* **785** (2007) 115–134, [[0705.0008](#)].
- [17] A. Dabholkar and C. Hull, *Duality twists, orbifolds, and fluxes*, *JHEP* **09** (2003) 054, [[hep-th/0210209](#)].
- [18] O. Hohm and H. Samtleben, *Exceptional field theory. III. $E_{8(8)}$* , *Phys. Rev. D* **90** (2014) 066002, [[1406.3348](#)].
- [19] L. J. Romans, *Massive $N=2a$ Supergravity in Ten-Dimensions*, *Phys. Lett. B* **169** (1986) 374.
- [20] S. Kachru, M. B. Schulz and S. Trivedi, *Moduli stabilization from fluxes in a simple IIB orientifold*, *JHEP* **10** (2003) 007, [[hep-th/0201028](#)].
- [21] J.-P. Derendinger, C. Kounnas, P. M. Petropoulos and F. Zwirner, *Superpotentials in IIA compactifications with general fluxes*, *Nucl.Phys.* **B715** (2005) 211–233, [[hep-th/0411276](#)].
- [22] O. DeWolfe, A. Giriyavets, S. Kachru and W. Taylor, *Enumerating flux vacua with enhanced symmetries*, *JHEP* **02** (2005) 037, [[hep-th/0411061](#)].
- [23] P. G. Camara, A. Font and L. E. Ibanez, *Fluxes, moduli fixing and MSSM-like vacua in a simple IIA orientifold*, *JHEP* **09** (2005) 013, [[hep-th/0506066](#)].
- [24] G. Villadoro and F. Zwirner, *$N=1$ effective potential from dual type-IIA $D6/O6$ orientifolds with general fluxes*, *JHEP* **0506** (2005) 047, [[hep-th/0503169](#)].
- [25] J. P. Derendinger, C. Kounnas, P. M. Petropoulos and F. Zwirner, *Fluxes and gaugings: $N=1$ effective superpotentials*, *Fortsch. Phys.* **53** (2005) 926–935, [[hep-th/0503229](#)].
- [26] O. DeWolfe, A. Giriyavets, S. Kachru and W. Taylor, *Type IIA moduli stabilization*, *JHEP* **07** (2005) 066, [[hep-th/0505160](#)].
- [27] G. Aldazabal, P. G. Camara, A. Font and L. Ibanez, *More dual fluxes and moduli fixing*, *JHEP* **0605** (2006) 070, [[hep-th/0602089](#)].
- [28] J. Shelton, W. Taylor and B. Wecht, *Nongeometric flux compactifications*, *JHEP* **0510** (2005) 085, [[hep-th/0508133](#)].
- [29] W. Decker, G.-M. Greuel, G. Pfister and H. Schönemann, “SINGULAR 4-3-0 — A computer algebra system for polynomial computations.” <http://www.singular.uni-kl.de>, 2022.
- [30] P. Breitenlohner and D. Z. Freedman, *Positive Energy in anti-De Sitter Backgrounds and Gauged Extended Supergravity*, *Phys. Lett. B* **115** (1982) 197–201.
- [31] O. Hohm, E. T. Musaev and H. Samtleben, *$O(d+1, d+1)$ enhanced double field theory*, *JHEP* **10** (2017) 086, [[1707.06693](#)].
- [32] E. Malek and H. Samtleben, *Kaluza-Klein Spectrometry for Supergravity*, *Phys. Rev. Lett.* **124** (2020) 101601, [[1911.12640](#)].

- [33] E. Malek and H. Samtleben, *Kaluza-Klein Spectrometry from Exceptional Field Theory*, *Phys. Rev. D* **102** (2020) 106016, [[2009.03347](#)].
- [34] G. Larios Plaza, *Spectroscopy and truncations from holography and exceptional dualities*. PhD thesis, U. Autonoma, Madrid (main), 2022.
- [35] J. P. Conlon, S. Ning and F. Revello, *Exploring the holographic Swampland*, *JHEP* **04** (2022) 117, [[2110.06245](#)].
- [36] F. Apers, M. Montero, T. Van Riet and T. Wrase, *Comments on classical AdS flux vacua with scale separation*, *JHEP* **05** (2022) 167, [[2202.00682](#)].
- [37] F. Apers, J. P. Conlon, S. Ning and F. Revello, *Integer conformal dimensions for type IIA flux vacua*, *Phys. Rev. D* **105** (2022) 106029, [[2202.09330](#)].
- [38] J. Quirant, *Noninteger conformal dimensions for type IIA flux vacua*, *Phys. Rev. D* **106** (2022) 066017, [[2204.00014](#)].
- [39] H. Ooguri and C. Vafa, *Non-supersymmetric AdS and the Swampland*, *Adv. Theor. Math. Phys.* **21** (2017) 1787–1801, [[1610.01533](#)].
- [40] A. Font, A. Herráez and L. E. Ibáñez, *On scale separation in type II AdS flux vacua*, *JHEP* **03** (2020) 013, [[1912.03317](#)].
- [41] E. Plauschinn, *Mass spectrum of type IIB flux compactifications — comments on AdS vacua and conformal dimensions*, *JHEP* **02** (2023) 257, [[2210.04528](#)].
- [42] X. de la Ossa, M. Larfors, M. Magill and E. E. Svanes, *Superpotential of three dimensional $\mathcal{N} = 1$ heterotic supergravity*, *JHEP* **01** (2020) 195, [[1904.01027](#)].
- [43] D. B. Kaplan, J.-W. Lee, D. T. Son and M. A. Stephanov, *Conformality Lost*, *Phys. Rev. D* **80** (2009) 125005, [[0905.4752](#)].
- [44] A. F. Faedo, C. Hoyos, D. Mateos and J. G. Subils, *Holographic Complex Conformal Field Theories*, *Phys. Rev. Lett.* **124** (2020) 161601, [[1909.04008](#)].
- [45] C. Eloy, *Kaluza-Klein spectrometry for AdS₃ vacua*, *SciPost Phys.* **10** (2021) 131, [[2011.11658](#)].
- [46] C. Eloy and G. Larios, *Nonsupersymmetric stable marginal deformations in AdS₃/CFT₂*, *Phys. Rev. D* **108** (2023) L121901, [[2309.03261](#)].
- [47] C. Eloy and G. Larios, *Charting the Conformal Manifold of Holographic CFT₂'s*, [2405.17542](#).
- [48] H. Ooguri and C. Vafa, *On the Geometry of the String Landscape and the Swampland*, *Nucl. Phys. B* **766** (2007) 21–33, [[hep-th/0605264](#)].
- [49] D. Junghans, *O-Plane Backreaction and Scale Separation in Type IIA Flux Vacua*, *Fortsch. Phys.* **68** (2020) 2000040, [[2003.06274](#)].
- [50] D. Junghans, *A note on O6 intersections in AdS flux vacua*, *JHEP* **02** (2024) 126, [[2310.17695](#)].
- [51] M. Emelin, F. Farakos and G. Tringas, *O6-plane backreaction on scale-separated Type IIA AdS₃ vacua*, *JHEP* **07** (2022) 133, [[2202.13431](#)].
- [52] M. Emelin, *Consistency conditions for O-plane unsmearing from second-order perturbation theory*, [2407.12717](#).
- [53] A. Guarino, *On new maximal supergravity and its BPS domain-walls*, *JHEP* **02** (2014) 026, [[1311.0785](#)].