GRADIENT FLOW IN PARAMETER SPACE IS EQUIVALENT TO LINEAR INTERPOLATION IN OUTPUT SPACE

THOMAS CHEN AND PATRICIA MUÑOZ EWALD

ABSTRACT. We prove that the usual gradient flow in parameter space that underlies many training algorithms for neural networks in deep learning can be continuously deformed into an adapted gradient flow which yields (constrained) Euclidean gradient flow in output space. Moreover, if the Jacobian of the outputs with respect to the parameters is full rank (for fixed training data), then the time variable can be reparametrized so that the resulting flow is simply linear interpolation, and a global minimum can be achieved.

1. INTRODUCTION

At the core of most algorithms currently used for training neural networks is gradient descent, whose theoretical counterpart in the continuum limit is gradient flow. This flow is defined in a space of parameters \mathbb{R}_{θ}^{K} with respect to a cost function defined in $\mathbb{R}_{\underline{x}}^{QN}$. The cost is non-convex as a function of the parameters, and standard gradient flow in \mathbb{R}_{θ}^{K} might not converge to a global minimum. Even changing perspective to $\mathbb{R}_{\underline{x}}^{QN}$, where the cost can be chosen to be convex, the resulting associated flow is complicated.

In [\[Che23\]](#page-10-0), it was discussed in detail how the choice of the Riemannian structure in \mathbb{R}_{θ}^{K} influences the training dynamics. In this context, a modified gradient flow that induces the Euclidean gradient flow in output space \mathbb{R}_{x}^{QN} was introduced and contrasted with the standard Euclidean gradient flow in parameter space; it was shown that both flows exhibit the same critical sets. In the work at hand, we extend these results by proving that those two flows are in fact homotopy equivalent to one another (Theorem [2.3\)](#page-3-0). Moreover, we prove that if the Jacobian matrix exhibits no rank loss, then the Euclidean flow in output space is reparametrization equivalent to linear interpolation with respect to a suitable time variable (Proposition [2.4\)](#page-5-0). When there is rank loss, an expression for the deviation from linear interpolation is given (Proposition [2.5\)](#page-6-0).

In section [3,](#page-6-1) we provide some applications: First, we note that instead of modifying the metric in \mathbb{R}_{θ}^K , one could prescribe a convenient path in $\mathbb{R}_{\underline{x}}^{QN}$ and then find an associated path in parameter space. Next, we show that neural collapse occurs in the output layer for the trivialized dynamics studied in [2.2](#page-5-1) (Corollary [3.2\)](#page-8-0). Finally, we rephrase Theorem [2.3](#page-3-0) in terms of the neural tangent kernel.

We briefly comment on related work in section [4.](#page-9-0) A short appendix on generalized inverses for rank deficient matrices is provided as well.

2. Main results

Consider a family of functions

$$
f_{\underline{\theta}} : \mathbb{R}^M \to \mathbb{R}^Q \tag{2.1}
$$

Date: August 2, 2024.

parametrized by a vector of parameters $\underline{\theta} \in \mathbb{R}^K$. Suppose we wish to optimize $f_{\underline{\theta}}$ with respect to a given cost function $C: \mathbb{R}^K \to \mathbb{R}$. A natural approach is to pick a starting point $\underline{\theta}_0 \in \mathbb{R}^K$ and construct a path

$$
\underline{\theta}(s) : [0, \infty) \to \mathbb{R}^K \tag{2.2}
$$

via the gradient flow

$$
\partial_s \underline{\theta}(s) = -\nabla_{\underline{\theta}} \mathcal{C},
$$

$$
\underline{\theta}(0) = \underline{\theta}_0.
$$
 (2.3)

We will refer to (2.3) as the *standard gradient flow*.

When f_{θ} is a neural network, the optimization problem has the following structure: Consider a set $\mathcal{X}_0 \subset \mathbb{R}^M$ (the training data), such that $|\mathcal{X}_0| = N$. We can form a data matrix $X_0 \in \mathbb{R}^{M \times N}$ by making the vectors in \mathcal{X}_0 into the columns of X_0 . Then we can consider a function

$$
f(\underline{\theta}, X_0) : \mathbb{R}^K \times \mathbb{R}^{M \times N} \to \mathbb{R}^{Q \times N},
$$
\n(2.4)

defined by its action on each column of X_0 : for $i = 1, \dots, N$,

$$
f(\underline{\theta}, X_0)_i = f_{\underline{\theta}}((X_0)_i) \in \mathbb{R}^Q.
$$
\n
$$
(2.5)
$$

In the context of supervised learning, each data point $x \in \mathcal{X}_0$ is associated to a desired output $y(x) \in \mathbb{R}^Q$, and so from a data matrix X_0 we can form the corresponding matrix of labels, $Y \in \mathbb{R}^{Q \times N}$,

$$
Y_i = y((X_0)_i). \t\t(2.6)
$$

A common choice for cost function is the squared loss,

$$
\mathcal{C}(\underline{\theta}, X_0, Y) = \frac{1}{N} \| f(\underline{\theta}, X_0) - Y \|_{\mathcal{L}^2}^2,
$$
\n(2.7)

where $\lVert \cdot \rVert_{\mathcal{L}^2}$ is the Frobenius or Hilbert-Schmidt norm.

For fixed training data $X_0 = [x_1 \cdots x_j \cdots x_N]$, we can also consider the function

$$
\underline{x}: \mathbb{R}^K \to \mathbb{R}^{QN} \tag{2.8}
$$

from parameter space to output space defined by

$$
\underline{x}(\underline{\theta}) := (f_{\underline{\theta}}(x_1)^T, \cdots, f_{\underline{\theta}}(x_N)^T)^T.
$$
\n(2.9)

We define the correspondent vector of labels $y \in \mathbb{R}^{QN}$,

$$
\underline{y} := (y(x_1)^T, \cdots, y(x_N)^T)^T,\tag{2.10}
$$

In this case, the cost (2.7) becomes

$$
\mathcal{C}(\underline{x}(\underline{\theta})) = \frac{1}{N} |\underline{x}(\underline{\theta}) - \underline{y}|^2.
$$
\n(2.11)

The Jacobian $D[\underline{\theta}] \in \mathbb{R}^{QN \times K}$ is

$$
(D[\underline{\theta}])_{jk} := \frac{\partial \underline{x}_j[\underline{\theta}]}{\partial \theta_k},\tag{2.12}
$$

and we assume from now on that the partial derivatives in [\(2.12\)](#page-1-2) exist and are Lipschitz continuous for all $\underline{\theta} \in \mathbb{R}^K$. We will often abbreviate $D = D[\underline{\theta}]$.

For $\underline{\theta}(s) \in \mathbb{R}^K$, we can consider the associated path

$$
\underline{x}(s) : [0, \infty) \to \mathbb{R}^{QN} \tag{2.13}
$$

given by

$$
\underline{x}(s) := \underline{x}(\theta(s))\tag{2.14}
$$

from [\(2.9\)](#page-1-3). The associated Jacobian is now also time dependent, and we will write $D(s) := D[\underline{\theta}(s)]$, or simply D for convenience. The chain rule gives

$$
\partial_s \underline{x}(s) = D(s) \partial_s \underline{\theta}(s) \tag{2.15}
$$

and

$$
\nabla_{\underline{\theta}} \mathcal{C} = D^T \nabla_{\underline{x}} \mathcal{C}.
$$
\n(2.16)

Remark 2.1. It is clear that

$$
\{\underline{\theta} : \nabla_{\underline{x}(\underline{\theta})} C = 0\} \subset \{\underline{\theta} : \nabla_{\underline{\theta}} C = 0\},\tag{2.17}
$$

however, the two sets are not necessarily identical. It is known that if rank $D[\underline{\theta}] = QN = \min\{K, QN\}$, then $\nabla_{\underline{x}}\mathcal{C}=0$ if, and only if, $\nabla_{\underline{\theta}}\mathcal{C}=0$. Moreover, if C is the squared loss, then $\nabla_{\underline{x}}\mathcal{C}=0$ corresponds to a global minimum with zero loss. See e.g. [\[Che23,](#page-10-0) [CE23,](#page-10-1) [KMTM24\]](#page-11-0) for more details.

2.1. Adapted gradient flow. By a simple computation using [\(2.15\)](#page-2-0) and [\(2.16\)](#page-2-1), if $\mathcal{Q}(s)$ satisfies the standard gradient flow [\(2.3\)](#page-1-0), then

$$
\partial_s \underline{x} = -DD^T \nabla_{\underline{x}} \mathcal{C}.
$$
\n(2.18)

We will refer to (2.18) as *standard gradient flow in output space*. Suppose instead we wished to modify $\theta(s)$ so that the associated path $x(s)$ satisfies the Euclidean gradient flow. We recall here results in previous work by one of the authors, in the overparametrized case:

Lemma 2.2 ([\[Che23\]](#page-10-0)). Let $K \ge QN$, and let $\underline{x}(s) = \underline{x}(\underline{\theta}(s))$ be defined as in [\(2.9\)](#page-1-3). When D is full rank, setting

$$
\partial_s \underline{\theta}(s) = -(D^T D)^+ \nabla_{\underline{\theta}} \mathcal{C}
$$
\n(2.19)

yields

$$
\partial_s \underline{x}(s) = -\nabla_{\underline{x}} \mathcal{C}.\tag{2.20}
$$

If we allow for $rank(D) \leq QN$, then letting

$$
\partial_s \underline{\theta}(s) = -D^T \psi,\tag{2.21}
$$

for ψ satisfying

$$
DD^T \psi = \mathcal{P}_{\text{ran}(DD^T)} \nabla_x \mathcal{C},\tag{2.22}
$$

results in

$$
\partial_s \underline{x}(s) = -\mathcal{P}_{\text{ran}(DD^T)} \nabla_{\underline{x}} \mathcal{C}.\tag{2.23}
$$

We will now condense these two separate cases into the same. First, note that

 ψ :

$$
= D(D)^{+} (DD^{T})^{+} (D^{T})^{+} \nabla_{\underline{\theta}} \mathcal{C}
$$
\n(2.24)

satisfies [\(2.22\)](#page-2-3), as the following computation shows:

$$
DD^{T}\tilde{\psi} = D \underbrace{D^{T}D(D)^{+}}_{\stackrel{(\mathbf{A},\mathbf{I}^{(2)})}{=}} D^{T} D(D)^{+}} (DD^{T})^{+} (D^{T})^{+} \nabla_{\underline{\theta}} C
$$

\n
$$
= DD^{T} (DD^{T})^{+} (D)^{+} (D^{T})^{+} D^{T} \nabla_{\underline{x}} C
$$

\n
$$
= DD^{T} (D^{T})^{+} \underbrace{(D)^{+} (D^{T})^{+} D^{T}}_{\stackrel{(\mathbf{A},\mathbf{I}^{(2)})}{=}} (D)^{+}
$$

\n
$$
= DD^{T} (DD^{T})^{+} \nabla_{\underline{x}} C
$$

\n
$$
= \mathcal{P}_{\text{ran}(DD^{T})} \nabla_{\underline{x}} C.
$$

\n(2.25)

Moreover, we can rewrite

$$
-(D^T D)^+ \nabla_{\underline{\theta}} C = -(D^T D)^+ (D^T D)(D^T D)^+ \nabla_{\underline{\theta}} C \n= -(D^T D)(D^T D)^+ (D^T D)^+ \nabla_{\underline{\theta}} C \n= -D^T [D(D)^+ (D^T)^+ (D)^+ (D^T)^+] \nabla_{\underline{\theta}} C \n= -D^T [D(D)^+ (DD^T)^+ (D^T)^+] \nabla_{\underline{\theta}} C, \n= -D^T \tilde{\psi}.
$$
\n(2.26)

Therefore in general, for $K \geq QN$ and $\text{rank}(D) \leq QN$,

$$
\partial_s \underline{\theta} = -(D^T D)^+ \nabla_{\underline{\theta}} \mathcal{C}
$$
\n(2.27)

implies

$$
\partial_s \underline{x} = -\mathcal{P}_{\text{ran}(DD^T)} \nabla_{\underline{x}} \mathcal{C}.
$$
\n(2.28)

We will refer to [\(2.27\)](#page-3-1) as *adapted gradient flow* in \mathbb{R}_{θ}^{K} and [\(2.28\)](#page-3-2) as *constrained Euclidean gradient flow* in $\mathbb{R}_{\underline{x}}^{QN}$.

It was further shown in [\[Che23,](#page-10-0) Theorem 4.2] that the standard gradient flow [\(2.3\)](#page-1-0) and the adapted gradient flow [\(2.27\)](#page-3-1) have the same equilibrium points. We extend this to a family of interpolating gradient fields.

Theorem 2.3. Let $C : \mathbb{R}^K \to \mathbb{R}$ depend on $\underline{\theta} \in \mathbb{R}^K$ only through $\underline{x}(\underline{\theta})$, for $\underline{x} : \mathbb{R}^K \to \mathbb{R}^{QN}$ defined as in [\(2.9\)](#page-1-3), and let $D[\underline{\theta}] \in \mathbb{R}^{QN \times K}$ be the Jacobian defined in [\(2.12\)](#page-1-2). There is a one-parameter family of vector fields in \mathbb{R}^K interpolating between

$$
-\nabla_{\underline{\theta}}\mathcal{C} \tag{2.29}
$$

and

$$
-(D^T D)^+ \nabla_{\underline{\theta}} \mathcal{C};\tag{2.30}
$$

this interpolation preserves singularities.

Moreover, this induces an interpolation between the vector fields for the standard gradient flow in output space (2.18) and the constrained Euclidean gradient flow (2.28) which preserves the equilibrium points of these flows.

Proof. Define the family of vector fields

$$
V_{\underline{\theta},\alpha} := -A_{\underline{\theta},\alpha} \nabla_{\underline{\theta}} \mathcal{C}, \quad \alpha \in [0,1], \tag{2.31}
$$

for

$$
A_{\underline{\theta},\alpha} := (\alpha(D^T D)^+ + (1 - \alpha)\mathbf{1}_{K \times K}). \tag{2.32}
$$

Then

$$
V_{\theta,0} = -\nabla_{\theta} \mathcal{C}
$$
\n(2.33)

and

$$
V_{\underline{\theta},1} = -(D^T D)^+ \nabla_{\underline{\theta}} \mathcal{C}.\tag{2.34}
$$

For $\alpha \in [0,1)$, note that $A_{\underline{\theta},\alpha} > 0$ for all $\underline{\theta}$. For $\alpha = 1$, writing $\mathcal{P}_{\text{ran}(D^{T})} = (D)^{+}D$, we have

$$
\nabla_{\underline{\theta}} \mathcal{C} = (D)^{+} D \nabla_{\underline{\theta}} \mathcal{C}, \qquad (2.35)
$$

since [\(2.16\)](#page-2-1) shows $\nabla_{\theta} \mathcal{C} \in \text{ran}(D^T)$, and also

$$
\ker(D^T D)^{+} = \ker(D^T D)^{+} = \ker(D^T D) = \ker((D)^{+} D),
$$
\n(2.36)

where we used [\(A.11\)](#page-10-3) and Lemma [A.2.](#page-10-4) Thus, $V_{\theta,\alpha} = 0$ if, and only if, $\nabla_{\theta} C = 0$, and the vector fields $V_{\theta,\alpha}$ are singular at the same points $\theta \in \mathbb{R}^K$, for all $\alpha \in [0, 1]$.

Suppose now that

$$
\partial_s \underline{\theta}(s) = V_{\underline{\theta}(s),\alpha},\tag{2.37}
$$

so that for $\underline{x}(s) = \underline{x}(\underline{\theta}(s)),$

$$
\partial_s \underline{x}(s) = D[\underline{\theta}(s)] V_{\underline{\theta}(s),\alpha}.
$$
\n(2.38)

We wish to write $\partial_s \underline{x}$ in terms of $\nabla_{\underline{x}} C$, so we compute

$$
D V_{\underline{\theta},\alpha} = -D \left(\alpha (D^T D)^+ + (1 - \alpha) \mathbf{1}_{K \times K} \right) \nabla_{\underline{\theta}} C
$$

= -\left(\alpha D (D^T D)^+ + (1 - \alpha) D \right) D^T \nabla_{\underline{x}} C
= -\left(\alpha D (D^T D)^+ D^T + (1 - \alpha) D D^T \right) \nabla_{\underline{x}} C
(*)

$$
\stackrel{(*)}{=} -(\alpha \mathcal{P}_{\text{ran}(DD^T)} + (1 - \alpha)DD^T) \nabla_{\underline{x}} \mathcal{C}
$$

= -(\alpha \mathbf{1}_{QN \times QN} + (1 - \alpha)DD^T) \mathcal{P}_{\text{ran}(DD^T)} \nabla_{\underline{x}} \mathcal{C},

where $(*)$ follows from $(A.7)$, $(A.8)$ and $(A.12)$. Therefore,

$$
\partial_s \underline{x}(s) = V_{\underline{x}(s),\alpha} \tag{2.40}
$$

for vector fields in \mathbb{R}^{QN} defined along the path $\underline{x}(s)$,

$$
V_{\underline{x}(s),\alpha} := -\left(\alpha \mathbf{1}_{QN \times QN} + (1-\alpha)D(s)D(s)^{T}\right) \mathcal{P}_{\text{ran}(D(s)D(s)^{T})} \nabla_{\underline{x}} \mathcal{C}.
$$
\n(2.41)

Observe that

$$
V_{\underline{\theta}(s),0} = -\nabla_{\underline{\theta}} \mathcal{C}
$$
\n(2.42)

and

$$
V_{\underline{x}(s),1} = -\mathcal{P}_{\text{ran}(DD^T)} \nabla_{\underline{x}} \mathcal{C},\tag{2.43}
$$

so that in fact $V_{\underline{x}(s),\alpha}$, $\alpha \in [0,1]$, interpolates between the standard dynamics given by (2.3) at $\alpha = 0$, and constrained Euclidean gradient flow [\(2.28\)](#page-3-2) for $\underline{x}(s)$ at $\alpha = 1$. Moreover, for $\alpha \in (0,1]$, $(\alpha \mathbf{1}_{QN \times QN} +$ $(1-\alpha)DD^{T}$) is positive-definite, and for $\alpha=0$, DD^{T} is injective in the range of $\mathcal{P}_{\text{ran}(DD^{T})}$, so

$$
(\alpha \mathbf{1} + (1 - \alpha) D D^T) \mathcal{P}_{\text{ran}(DD^T)} \nabla_{\underline{x}} \mathcal{C} = 0 \text{ if, and only if, } \mathcal{P}_{\text{ran}(DD^T)} \nabla_{\underline{x}} \mathcal{C} = 0. \tag{2.44}
$$

Thus, the equilibrium points of [\(2.40\)](#page-4-0) are the same for all $\alpha \in [0,1]$.

We add the following remarks highlighting the geometric interpretation of the above results. If DD^T has full rank QN , then $V_{\underline{x}(s),\alpha}$ is the gradient vector field with respect to the Riemannian metric on $T \mathbb{R}^{QN}$ with tensor

$$
\left(\alpha \mathbf{1}_{QN \times QN} + (1 - \alpha)DD^T\right)^{-1} \,. \tag{2.45}
$$

This means that $V_{\underline{x}(s),\alpha}$ can equivalently be considered as the gradient fields obtained from the family of Riemannian structures interpolating between the Euclidean structure on $T\mathbb{R}^{QN}$, and the metric structure induced by the Euclidean structure on parameter space.

If DD^T is rank-deficient, rank $(DD^T) < QN$, we let $\mathcal{V} \subset T\mathbb{R}^{QN}$ denote the vector subbundle whose fibers are given by the range of DD^T . In this situation, $V_{\underline{x}(s),\alpha}$ is the gradient with respect to the bundle metric h_{α} on V with tensor

$$
[h_{\alpha}] = (\alpha \mathbf{1}_{QN \times QN} + (1 - \alpha)DD^T)^{-1} \Big|_{\text{ran}(DD^T[\underline{\theta}])}.
$$
\n(2.46)

The triple $(\mathbb{R}^{QN}, V, h_{\alpha})$ defines a sub-Riemannian manifold with a family of bundle metrics for $\alpha \in [0, 1]$; see [\[Che23\]](#page-10-0) for a related discussion and further details.

2.2. Reparametrization. Assuming rank $D = QN \leq K$, the adapted gradient flow in output space resulting from [\(2.27\)](#page-3-1) is simply Euclidean gradient flow,

$$
\partial_s \underline{x}(s) = -\nabla_{\underline{x}} \mathcal{C}(\underline{x}(s)).\tag{2.47}
$$

For

$$
\mathcal{C}(\underline{x}(\underline{\theta})) = \frac{1}{2N} \left| \underline{x}(\underline{\theta}) - \underline{y} \right|_{\mathbb{R}^{QN}}^2
$$
\n(2.48)

we have

$$
\partial_s \underline{x} = \frac{1}{N} (\underline{x}(s) - \underline{y}),\tag{2.49}
$$

which is solved by

$$
\underline{x}(s) = \underline{y} + e^{-\frac{s}{N}}(\underline{x}(0) - \underline{y}),\tag{2.50}
$$

and we see that $\underline{x}(s) \to y$ as $s \to \infty$.

Suppose instead that we wanted to impose a linear interpolation

$$
\hat{\underline{x}}(t) = \underline{y} + (1 - t)(\underline{x}_0 - \underline{y}).\tag{2.51}
$$

Then letting

$$
t := 1 - e^{-\frac{s}{N}},\tag{2.52}
$$

$$
s = -N \ln(1 - t),
$$
\n(2.53)

and

$$
\underline{\tilde{x}}(t) := \underline{x}(-N\ln(1-t)),\tag{2.54}
$$

for $\underline{x}(s)$ a solution to [\(2.47\)](#page-5-2), we have

$$
\partial_t \tilde{\underline{x}}(t) = \partial_t \underline{x}(-N \ln(1-t))
$$

= $-\nabla_{\underline{x}} \mathcal{C}[\tilde{\underline{x}}(t)] \frac{N}{1-t}$
= $-\frac{1}{1-t} (\tilde{\underline{x}}(t) - \underline{y}).$ (2.55)

Note that $\hat{x}(t)$ solves [\(2.55\)](#page-5-3), as

$$
\partial_t \hat{\underline{x}}(t) = -(\hat{\underline{x}}_0 - \underline{y}) \n= -\frac{1}{1-t}(\hat{\underline{x}}(t) - \underline{y}).
$$
\n(2.56)

Thus, we have proved the following:

Proposition 2.4. If $\underline{x}(s)$ is a solution to Euclidean gradient flow [\(2.47\)](#page-5-2), then

$$
\underline{\tilde{x}}(t) := \underline{x}(-N\ln(1-t)) = \underline{y} + (1-t)(\underline{x}_0 - \underline{y}),\tag{2.57}
$$

and $\underline{\tilde{x}}(t) \to \underline{y}$ as $t \to 1$. In particular, this holds for $\underline{x}(s) = \underline{x}(\underline{\theta}(s))$ defined as in [\(2.9\)](#page-1-3) when $\underline{\theta}(s)$ is a solution to the adapted gradient flow [\(2.27\)](#page-3-1) and rank $D[\underline{\theta}(s)] = QN \leq K$.

On the other hand, if $rank(D) < QN$, the reparametrized Euclidean gradient flow in output space provides a concrete criterion whereby the effect of rank loss can be measured.

Proposition 2.5. In general, the constrained Euclidean gradient flow in output space with time reparametrization as in [\(2.54\)](#page-5-4) satisfies

$$
\partial_t \underline{\tilde{x}}(t) = -\frac{1}{1-t} \mathcal{P}_t(\underline{\tilde{x}}(t) - \underline{y}), \quad \underline{\tilde{x}}(0) = \underline{x}_0, \quad t \in [0, 1), \tag{2.58}
$$

where $\underline{\tilde{x}}(t) = \underline{x}[\theta(s(t))]$ and $\mathcal{P}_t := \mathcal{P}_{\text{ran}(DD^T)}[\theta(s(t))]$. When $\text{rank}(D) < QN$, the deviation from linear interpolation is given by

$$
\underline{\tilde{x}}(t) - \left((1-t)\underline{x}_0 - t\underline{y} \right) = \int_0^t dt' \, \mathcal{U}_{t,t'} \, \frac{1-t}{1-t'} \, \mathcal{P}_{t'}^\perp \left(\underline{x}_0 - \underline{y} \right),\tag{2.59}
$$

where the linear propagator $\mathcal{U}_{t,t'}$ is determined by

$$
\partial_t \mathcal{U}_{t,t'} = \frac{1}{1-t} \mathcal{P}_t \mathcal{U}_{t,t'}, \quad \mathcal{U}_{t',t'} = \mathbf{1}_{QN \times QN}, \quad t,t' \in [0,1).
$$
 (2.60)

Proof. The equation (2.58) is straightforwardly obtained in a similar way as (2.55) .

To prove [\(2.59\)](#page-6-3), we write

$$
\underline{\tilde{x}}(t) = (1 - t)\underline{x}_0 + t\underline{y} + (1 - t)R(t)
$$
\n(2.61)

so that

$$
\partial_t \underline{\tilde{x}}(t) = -(\underline{x}_0 - \underline{y}) - R(t) + (1 - t)\partial_t R(t). \qquad (2.62)
$$

On the other hand,

$$
\partial_t \underline{\tilde{x}}(t) = -\frac{1}{1-t} \mathcal{P}_t(\underline{\tilde{x}}(t) - \underline{y})
$$
\n
$$
= -\frac{1}{1-t} (\tilde{x}(t) - \underline{y}) + \frac{1}{1-t} \mathcal{P}_t^{\perp}(\tilde{x}(t) - \underline{y})
$$
\n
$$
= -\frac{1}{1-t} ((1-t)\underline{x}_0 + t\underline{y} - \underline{y} + (1-t)R(t)) + \frac{1}{1-t} \mathcal{P}_t^{\perp}((1-t)\underline{x}_0 + t\underline{y} - \underline{y} + (1-t)R(t)))
$$
\n
$$
= -(\underline{x}_0 - \underline{y}) - R(t) + \mathcal{P}_t^{\perp}(\underline{x}_0 - \underline{y} + R(t)),
$$
\n(2.63)

where we used (2.61) to pass to the third line. Comparing the last line with (2.62) , we find

$$
\partial_t R(t) = \frac{1}{1-t} \mathcal{P}_t^{\perp} R(t) + \frac{1}{1-t} \mathcal{P}_t^{\perp} (\underline{x}_0 - \underline{y}). \tag{2.64}
$$

Solving this matrix valued linear ODE for $R(t)$ using the Duhamel (or variation of constants) formula, and substituting the resulting expression in (2.61) yields (2.59) , as claimed. □

3. Applications

3.1. Prescribed paths in output space. In the previous section, we discussed how changes in $\partial_s \theta(s)$ influence $\partial_s \underline{x}(s)$. It could be interesting to take a different point of view and find $\underline{\theta}(s)$ from a given path $x(s)$. In general, from (2.15) ,

$$
\mathcal{P}_{\text{ran}(D^T)} \partial_s \underline{\theta} = (D)^+ \partial_s \underline{x},\tag{3.1}
$$

and so assuming $\partial_s \underline{\theta} = \mathcal{P}_{\text{ran}(D^T)} \partial_s \underline{\theta},^1$ $\partial_s \underline{\theta} = \mathcal{P}_{\text{ran}(D^T)} \partial_s \underline{\theta},^1$ we can write

$$
\partial_s \underline{\theta}(s) = (D)^+ \partial_s \underline{x}(s). \tag{3.2}
$$

¹Note that this is satisfied by the standard gradient flow $\partial_s \underline{\theta} = -\nabla_{\underline{\theta}} C = -D^T \nabla_{\underline{x}} C$.

Thus, we could prescribe a path

$$
\underline{x}(s), \quad s \in [0, T],
$$

\n
$$
\partial_s \underline{x}(s) \in \text{ran}(D(s)),
$$

\n
$$
\underline{x}(T) = y,
$$
\n(3.3)

and then search for a corresponding $\underline{\theta}_*$ which realizes

$$
\underline{x}(\underline{\theta}_*) = \underline{x}(T) \tag{3.4}
$$

by solving

$$
\partial_s \underline{\theta}(s) = (D(s))^+ \partial_s \underline{x}(s),
$$

\n
$$
\underline{\theta}(0) = \underline{\theta}_0,
$$

\n
$$
\underline{x}(s) \text{ satisfies (3.3) with } \underline{x}(0) = \underline{x}(\underline{\theta}_0),
$$

\n(3.5)

for some initialization $\underline{\theta}_0 \in \mathbb{R}^K$. Observe that

$$
ran(D) = (ker(DT))\perp = (ker(D)+)\perp,
$$
\n(3.6)

and so all equilibrium points of [\(3.5\)](#page-7-1) satisfy $\partial_s \underline{x} = 0$.

As an example, motivated by the results in section [2.2,](#page-5-1) one could take $x(s)$ to be the linear interpolation (2.51) and check if, for a given $\underline{\theta}_0$,

$$
(\underline{x}_0 - \underline{y}) \in \text{ran } D(s),\tag{3.7}
$$

for all $s \in [0, T]$.

Remark 3.1. Whether or not a path $\underline{x}(s)$ satisfies the constraint in [\(3.3\)](#page-7-0) (or [\(3.7\)](#page-7-2) in the example above) could depend on the choice of $\underline{\theta}_0$. In general, properties of the Jacobian matrix $D[\underline{\theta}(s)]$ that determine the dynamics of $\underline{x}(s)$, such as rank and the subspace ran (DD^T) , might depend on the initialization even within a class $\{\underline{\theta}_0 : \underline{x}(\underline{\theta}_0) = \underline{x}_0\}.$

3.2. Final layer collapse. We turn to the case of a classification task. Assume without any loss of generality that the components of \tilde{x} are ordered by way of

$$
\underline{\tilde{x}} = (\dots, \dots, \tilde{x}_{j,1}^T, \dots, \tilde{x}_{j,i_j}^T, \dots, \tilde{x}_{j,N_j}^T, \dots,)^T \in \mathbb{R}^{QN}
$$
\n(3.8)

where each $\tilde{x}_{j,i_j} \in \mathbb{R}^Q$ belongs to the class labeled by $y_j \in \mathbb{R}^Q$, for $j = 1, \ldots, Q$, $i_j = 1, \ldots, N_j$, and $\sum_{j=1}^{Q} N_j = N$. Then, defining the class averages

$$
\overline{\tilde{x}_j}(t) := \frac{1}{N_j} \sum_{i_j=1}^{N_j} \tilde{x}_{j,i_j}(t)
$$
\n(3.9)

and the deviations

$$
\Delta \tilde{x}_{j,i_j}(t) := \tilde{x}_{j,i_j}(t) - \overline{\tilde{x}_j}(t),\tag{3.10}
$$

the \mathcal{L}^2 cost [\(2.7\)](#page-1-1) decomposes into

$$
\mathcal{C}[\underline{\tilde{x}}(t)] = \frac{1}{N} \sum_{j=1}^{Q} \sum_{i_j}^{N_j} |\tilde{x}_{j,i_j}(t) - y_j|^2
$$

=
$$
\frac{1}{N} \sum_{j=1}^{Q} \sum_{i_j}^{N_j} (|\tilde{x}_{j,i_j}(t) - \overline{\tilde{x}_j}(t)|^2 + |\overline{\tilde{x}_j}(t) - y_j|^2)
$$

=
$$
\frac{1}{N} \sum_{j=1}^{Q} \sum_{i_j}^{N_j} |\Delta \tilde{x}_{j,i_j}(t)|^2 + \sum_{j=1}^{Q} |\overline{\tilde{x}_j}(t) - y_j|^2.
$$
 (3.11)

This implies that zero loss training is achievable if and only if the class averages in the output layer are matched to the reference outputs, $\overline{\tilde{x}_j}(t) \to y_j$ for all $j = 1, \ldots, Q$, and all deviations vanish, $\Delta \tilde{x}_{j,i_j}(t) \to 0$ for all $j = 1, \ldots, Q, i_j = 1, \ldots, N_j$. The latter implies that the image of training data in the output layer contracts to one point per class.

Thus, Proposition [2.4](#page-5-0) allows us to arrive at the following conclusion.

Corollary 3.2. If $rank(D) = QN$, then zero loss minimization is achieved, whereby the cluster averages converge to the reference outputs,

$$
\lim_{t \to 1^{-}} \overline{\tilde{x}_j}(t) = y_j,
$$
\n(3.12)

and the deviations converge to zero,

$$
\lim_{t \to 1^{-}} \Delta \tilde{x}_{j,i_j}(t) = 0, \tag{3.13}
$$

for all $j = 1, ..., Q$ and $i_j = 1, ..., N_j$.

The decomposition of the cost function [\(3.11\)](#page-8-1) is related to the phenomenon known as neural collapse ([\[PHD20,](#page-11-1) [HPD22\]](#page-11-2)), which takes place on the penultimate layer of a neural network. For work on the relationship between neural collapse and final layer collapse, see e.g. [\[EW22\]](#page-10-7).

3.3. Tangent kernel. A related point of view is to study the neural tangent kernel (NTK) introduced by [\[JGH18\]](#page-11-3). It can be written as

$$
\Theta(x, y, \underline{\theta}) = Df_{\underline{\theta}}(x)(Df_{\underline{\theta}}(y))^T \in \mathbb{R}^{Q \times Q}, \tag{3.14}
$$

for a function $f_{\theta} : \mathbb{R}^M \to \mathbb{R}^Q$ and $Df_{\theta}(x)$ the Jacobian of f_{θ} at $x \in \mathbb{R}^M$. Then for a set of training data $\mathcal{X}_0 = \{x_1, \dots, x_N\}$ and $D[\underline{\theta}]$ defined as in section [2](#page-0-0) we can write $DD^T \in \mathbb{R}^{QN \times QN}$ as a block matrix with blocks of size $Q \times Q$,

$$
(D[\underline{\theta}]D[\underline{\theta}]^T)i, j = \Theta(x_i, x_j, \underline{\theta}), \quad i, j = 1, \cdots, N.
$$
\n(3.15)

The kernel $\Theta(x, y, \underline{\theta})$ is positive-definite with respect to the chosen training data at $\underline{\theta}$ if, ^{[2](#page-8-2)} for all $(u_i)_{i=1}^N \in$ $\bigoplus^N{\mathbb R}^Q,$

$$
\sum_{i=1}^{N} |u_i|^2 > 0 \text{ implies } \sum_{i,j=1}^{N} u_i^T \Theta(x_i, x_j, \underline{\theta}) u_j > 0.
$$
 (3.16)

From [\(3.15\)](#page-8-3) we see that his holds if, and only if, DD^T is a positive-definite matrix.

 2 This is stronger than the definition given in [\[JGH18\]](#page-11-3), which gives a similar statement for all functions in a given function class F.

Thus, Theorem [2.3](#page-3-0) suggests that, by changing the metric in parameter space, the NTK can be modified into a simpler kernel K which preserves the equilibrium points of the flow. When Θ is positive-definite, K satisfies

$$
K(x_i, x_j) = \delta_{ij} \mathbf{1}_{Q \times Q},\tag{3.17}
$$

so that it is constant throughout training at least when evaluated on \mathcal{X}_0 . This trivializes the training dynamics in output space, and holds for any neural network, or indeed any family of functions parametrized by $\underline{\theta}$.

4. Some related work

Linear paths. Some works, starting with [\[GVS15\]](#page-11-4), interpolate the initial parameters \mathcal{Q}_0 and final parameters $\underline{\theta}_f$ obtained after training the network with usual methods such as stochastic gradient descent,

$$
\underline{\theta}_{\alpha} = (1 - \alpha)\underline{\theta}_{0} + \alpha \underline{\theta}_{f}, \ \alpha \in [0, 1], \tag{4.1}
$$

and study the behavior of the loss function on these linear slices of parameter space. Recent works, such as $[LBZ+21]$ $[LBZ+21]$, provide empirical and theoretical investigations of the monotonic linear interpolation (MLI) property, whereby the loss decreases monotonically along the path θ_{α} . This is a different approach from the one taken in this paper, where the linear interpolation happens in output space.

In [\[GKRBZ24\]](#page-10-8), the authors find linear trajectories of probability distributions when studying generative models and modifying the cost function.

The Jacobian [\(2.12\)](#page-1-2). As already discussed in subsection [3.3,](#page-8-4) the neural tangent kernel introduced by [\[JGH18\]](#page-11-3) is closely related to the Jacobian studied in this paper. It inspired works on linear approximations of gradient flow/descent and the so called kernel or lazy regime, see e.g. $[COB19, WGL+20]$ $[COB19, WGL+20]$ $[COB19, WGL+20]$ $[COB19, WGL+20]$.

Certain works study the rank of D more explicitly: In [GLMW22] [GLMW22] [GLMW22] , the authors study the functional dimension of ReLU neural networks, defined in terms of the rank of the Jacobian. In [\[KMTM24\]](#page-11-0), the authors show that for mildly overparametrized ReLU neural networks with scalar output $(Q = 1)$ most activation patterns correspond to regions in parameter space where the Jacobian is full rank.

Acknowledgments: T.C. gratefully acknowledges support by the NSF through the grant DMS-2009800, and the RTG Grant DMS-1840314 - Analysis of PDE. P.M.E. was supported by NSF grant DMS-2009800 through T.C., and UT Austin's GS Summer fellowship.

Appendix A. Pseudoinverse

We recall here the definition and properties of the pseudoinverse of a (possibly rank-deficient) matrix. For proofs and more detail, we refer the reader to [\[CM09\]](#page-10-10), chapter 1.

Definition A.1. The pseudoinverse of $A \in \mathbb{R}^{m \times n}$ is the unique matrix $(A)^+ \in \mathbb{R}^{n \times m}$ such that

$$
A(A)^{+}A = A, \tag{A.1}
$$

$$
(A)^{+}A(A)^{+} = (A)^{+},\tag{A.2}
$$

$$
(A(A)^{+})^{T} = A(A)^{+},\tag{A.3}
$$

$$
((A)^{+}A)^{T} = (A)^{+}A.
$$
\n(A.4)

When A is full rank, there are simple expressions for the pseudoinverse: When rank(A) = $m =$ $\min\{m, n\}$, A is surjective, so AA^T is invertible and we have $(A)^+ = A^T (AA^T)^{-1}$. On the other hand, when $\text{rank}(A) = n = \min\{m, n\}, A$ is injective and $A^T A$ is invertible and we have $(A)^+ = (A^T A)^{-1} A^T$. In general, there exist many algorithms to construct the pseudoinverse, for instance using singular value decomposition; see e.g. [\[CM09\]](#page-10-10), chapter 12.

The matrices A and $(A)^+$ as above satisfy a few basic properties:

$$
((A)^{+})^{+} = A, \tag{A.5}
$$

$$
(AT)+ = ((A)+)T,
$$
\n(A.6)

$$
(A^T A)^+ = (A)^+ (A^T)^+.
$$
\n(A.7)

Moreover, the pseudoinverse gives a convenient way to write projectors: Consider a subspace $V \subset \mathbb{R}^d$. We will denote by \mathcal{P}_V the orthogonal projector onto V, which satisfies $\mathcal{P}_V = \mathcal{P}_V^2 = \mathcal{P}_V^T$. Then

$$
A(A)^{+} = \mathcal{P}_{\text{ran}(A)},\tag{A.8}
$$

$$
(A)^{+}A = \mathcal{P}_{\text{ran}(A^{T})}. \tag{A.9}
$$

When considering the subspaces associated to A , it is useful to know the relations

$$
ran(A)^{+} = ran A^{T}, \qquad (A.10)
$$

$$
\ker(A)^{+} = \ker A^{T}.
$$
\n(A.11)

The following is a useful identity:

$$
A = AA^T(A^T)^+ = (A^T)^+A^T A. \tag{A.12}
$$

Similar identities can be found by substituting A^T and $(A)⁺$ in place of A.

Finally, note that it is not true in general that $(A)^+A = A(A)^+$; a sufficient condition is $A^T = A$. We make use of this fact, along with the following lemma, in the main text.

Lemma A.2. ker(
$$
A^T A
$$
) = ker((A)⁺ A).

Proof. We expand

$$
A^T A = A^T A (A)^+ A, \tag{A.13}
$$

so

$$
\ker((A)^{+}A) \subset \ker(A^{T}A). \tag{A.14}
$$

Similarly,

$$
(A)^{+}A \stackrel{(\mathbf{A},12)}{=} (A)^{+}(A^{T})^{+}A^{T}A \tag{A.15}
$$

and so

$$
\ker(A^T A) \subset \ker((A)^+ A). \tag{A.16}
$$

□

REFERENCES

- [CE23] Thomas Chen and Patricia Muñoz Ewald. Non-approximability of constructive global \mathcal{L}^2 minimizers by gradient descent in deep learning, 2023. [arXiv:2311.07065.](https://arxiv.org/abs/2311.07065)
- [Che23] Thomas Chen. Global \mathcal{L}^2 minimization at uniform exponential rate via geometrically adapted gradient descent in deep learning, 2023. [arXiv:2311.15487.](https://arxiv.org/abs/2311.15487)

[CM09] Stephen L Campbell and Carl D Meyer. Generalized inverses of linear transformations. SIAM, 2009.

- [COB19] Lenaic Chizat, Edouard Oyallon, and Francis Bach. On lazy training in differentiable programming. Advances in neural information processing systems, 32, 2019.
- [EW22] Weinan E and Stephan Wojtowytsch. On the emergence of simplex symmetry in the final and penultimate layers of neural network classifiers. In Joan Bruna, Jan Hesthaven, and Lenka Zdeborova, editors, Proceedings of the 2nd Mathematical and Scientific Machine Learning Conference, volume 145 of Proceedings of Machine Learning Research, pages 270–290. PMLR, 16–19 Aug 2022.
- [GKRBZ24] Hyemin Gu, Markos A. Katsoulakis, Luc Rey-Bellet, and Benjamin J. Zhang. Combining wasserstein-1 and wasserstein-2 proximals: robust manifold learning via well-posed generative flows, 2024. [arXiv:2407.11901.](https://arxiv.org/abs/2407.11901)
- [GLMW22] J Elisenda Grigsby, Kathryn Lindsey, Robert Meyerhoff, and Chenxi Wu. Functional dimension of feedforward relu neural networks, 2022. [arXiv:2209.04036.](https://arxiv.org/abs/2209.04036)
- [GVS15] Ian J. Goodfellow, Oriol Vinyals, and Andrew M. Saxe. Qualitatively characterizing neural network optimization problems, 2015. [arXiv:1412.6544.](https://arxiv.org/abs/1412.6544)
- [HPD22] X.Y. Han, Vardan Papyan, and David L. Donoho. Neural collapse under MSE loss: Proximity to and dynamics on the central path. In International Conference on Learning Representations, 2022.
- [JGH18] Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. Advances in neural information processing systems, 31, 2018.
- [KMTM24] Kedar Karhadkar, Michael Murray, Hanna Tseran, and Guido Montúfar. Mildly overparameterized relu networks have a favorable loss landscape, 2024. [arXiv:2305.19510.](https://arxiv.org/abs/2305.19510)
- [LBZ+21] James R Lucas, Juhan Bae, Michael R Zhang, Stanislav Fort, Richard Zemel, and Roger B Grosse. On monotonic linear interpolation of neural network parameters. In Marina Meila and Tong Zhang, editors, Proceedings of the 38th International Conference on Machine Learning, volume 139 of Proceedings of Machine Learning Research, pages 7168–7179. PMLR, 18–24 Jul 2021.
- [PHD20] Vardan Papyan, XY Han, and David L Donoho. Prevalence of neural collapse during the terminal phase of deep learning training. Proceedings of the National Academy of Sciences, 117(40):24652–24663, 2020.
- [WGL+20] Blake Woodworth, Suriya Gunasekar, Jason D. Lee, Edward Moroshko, Pedro Savarese, Itay Golan, Daniel Soudry, and Nathan Srebro. Kernel and rich regimes in overparametrized models. In Jacob Abernethy and Shivani Agarwal, editors, Proceedings of Thirty Third Conference on Learning Theory, volume 125 of Proceedings of Machine Learning Research, pages 3635–3673. PMLR, 09–12 Jul 2020.

(T. Chen) Department of Mathematics, University of Texas at Austin, Austin TX 78712, USA Email address: tc@math.utexas.edu

(P. Muñoz Ewald) Department of Mathematics, University of Texas at Austin, Austin TX 78712, USA Email address: ewald@utexas.edu