

Symmetric black-to-white hole solutions with a cosmological constant

Zhong-Wen Feng^{1,*} Qing-Quan Jiang^{1,†} Yi Ling^{2,3,1,‡}
Xiao-Ning Wu^{4,5,§} and Zhangping Yu^{2,3,¶}

1 School of Physics and Astronomy, China West Normal University, Nanchong 637009, China

2 Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

3 School of Physics, University of Chinese Academy of Sciences, Beijing 100049, China

*4 Institute of Mathematics, Academy of Mathematics and System
Science and Hua Loo-Keng Key Laboratory of Mathematics,
Chinese Academy of Sciences, Beijing 100190, China*

*5 School of Mathematical Sciences, University of
Chinese Academy of Sciences, Beijing 100049, China*

For a system with a Hamiltonian constraint, we demonstrate that its dynamics preserves invariant with respect to the choice of the lapse function, regardless of whether the Hamiltonian incorporates quantum corrections. Applying this observation to the interior of black-to-white holes, we analyze its dynamics with different choices of the lapse function. The results explicitly show that the leading-order expansion of both metrics in Refs. [1, 2] and Refs. [3, 4] exhibits identical behavior near the transition surface, therefore in this sense the black-to-white hole model proposed in Refs. [1, 2] may be interpreted as a coarse-grained version of the solution within the framework of loop quantum gravity. The black-to-white hole solutions with exact symmetry between the black hole and white hole regions are constructed by appropriately fixing the quantum parameters in the effective theory of loop quantum gravity. This approach circumvents the issue of amplification of mass, which could arise from a mass difference between the black hole and white hole. Finally, the black-to-white hole solutions with a cosmological constant are constructed. The numerical solutions for the interior of black-to-white hole with a cosmological constant are obtained, and their symmetric behavior is also discussed.

* Email: zwfengphy@cwnu.edu.cn

† Email: qqjiangphys@yeah.net

‡ Email: lingy@ihep.ac.cn

§ Email: wuxn@amss.ac.cn

¶ Email: yuzp@ihep.ac.cn

I. INTRODUCTION

The successful detection of gravitational waves emanating from black holes [5] and the observation of the shadow of black holes [6, 7] have announced the coming of a new age for black hole physics. As a theory that predicts the existence of black holes, the general relativity (GR) plays a pivotal role in scrutinizing their properties including the structure of spacetime, causality as well as the thermodynamic properties. Nonetheless, as one delves deeper into the evolution of a massive star into a black hole, a quandary emerges—the gravitational collapse inexorably culminates and a singularity unavoidably appears at the center of a black hole, wherein all matter converges to a region with infinite curvature [8, 9], which signals the ultraviolet incompleteness of classical GR as a theory for the description of the spacetime of a black hole. The emergence of a singularity also implies that existing physical theories fail under such extreme conditions, leading to a significant gap in our understanding on the physical processes inside the horizon of a black hole. It is widely believed that as the curvature of spacetime approaches the Planckian scale, the quantum effects of gravity would become significant enough to change the geometry of spacetime. Therefore, quantum gravity (QG) would offer a promising way for us to come out from the enigma of singularity of spacetime [10–14]. Before a complete theory of quantum gravity is established, people have attempted to construct black holes without singularity (which are called as non-singular black holes or regular black holes) at both the phenomenological level and the semi-classical level. At the phenomenological level, one may construct the metric for regular black holes by introducing the exotic matter fields into the standard Einstein field equations, and the presence of such exotic matter fields might ultimately ascribe to the effects of QG. In history, the famous Bardeen black hole [15] and Hayward black hole [16] are prominent examples in this route. Recently, some novel solutions for regular black holes have also been constructed in this manner [17–35]. In particular, a black-to-white hole solution characterized by the sub-Planckian curvature is constructed in Refs. [1, 2]. The key observation leading to a sub-Planckian curvature is that the parameter l , which could be understood as the regulator reflecting the scale that the quantum corrections of gravity start to have an impact, may not have the same magnitude as the fundamental scale-Planck length, but depending on the mass of the black hole as well. As a result, one finds that the maximal value of the Kretschmann scalar curvature is not only finite, but also bounded from above by the Planck

energy density. Inspired by this work, some novel regular black holes with sub-Planckian curvature have been investigated in Refs. [36–48]. Nevertheless, we intend to point out that all the metrics of regular black holes proposed at the phenomenological level assume the existence of unknown exotic matter fields. Usually, one just derives the form of their energy stress tensor by requiring that Einstein equations should be satisfied, while their Lagrangian in terms of such matter fields is unknown. Essentially, one is urged to provide the theoretical foundation for such regular black holes by solving the equations of motion with matter fields as fundamental variables. It is worthwhile to point out that recently some typical regular black holes have been derived in the framework of gravity coupled to nonlinear electrodynamics [49–60], and other modified gravity theories, for instance, see Refs. [61–70]. On the contrary, the semi-classical approach is devoted to construct the effective equations of motion at first, which may receive significant quantum corrections of gravity due to quantum geometry and thus exhibit distinct behavior from that in classical theory of GR, and then try to derive the black-to-white metric as the solution to these modified equations. Usually, such metrics may look more complicated, but their theoretical foundation is more robust. Recently, such effective equations of motion for Schwarzschild black hole have been constructed in the framework of loop quantum gravity (LQG) [3, 4]. Previously, the relevant works on applying LQG techniques to the singularity problem of black hole can be found in literatures, for instance in Refs. [71–86]. In Refs. [3, 4], one significant improvement is incorporating new quantum parameters into the semi-classical Hamiltonian constraint. These parameters are set to Dirac observables, which means that along each dynamic trajectory they are constants, thereby enabling the simultaneous depiction of the influence of LQG on Schwarzschild black hole. Particularly, it is interesting enough to notice that the quantum parameters exhibit the same mass-dependent behavior as that in Refs. [1, 2]. In this new effective theory, singularity is eliminated and the black hole may undergo a transition to a white hole, presenting a novel avenue for addressing the singularity problem of black holes, and some subsequent works can be found in Refs. [87–100].

The motivations of the present work are following. Firstly, we intend to reveal some connections between two black-to-white solutions in Refs. [1, 2] and Refs. [3, 4]. On one hand, the metric form of the black-to-white solution in Refs. [1, 2] are quite simple and elegant, but proposed at the phenomenological level. In addition, its metric form only applies to the interior of the horizon, and if one tries to extend it to the region outside horizon, one would

find that the energy conditions would be violated everywhere, as pointed out in Ref. [48]. Here our effort may provide a theoretical foundation for this solution and embed it into the effective theory of black holes in the context of LQG. On the other hand, the metric form of the black-to-white solution in Refs. [3, 4] is obtained by explicitly solving the effective equations of motion which are derived in the context of LQG, thus with a robust theoretical foundation. Nevertheless, we notice that in comparison with Refs. [1, 2] a different lapse function is considered in Refs. [3, 4], which is very helpful for one to write down the analytical expression of solutions. However, in this coordinate system, the location of the transition surface does not occur at the original singularity with $T_{cl} = -\infty$, but at some place with finite T_q . Although it is understood that the time coordinate T_{cl} in classical theory is different from that T_q in semi-classical theory, this scenario perhaps is not quite intuitive as that in Ref. [1], where the transition surface from black hole to white hole occurs exactly at the location of the original singularity, namely at $T_{cl} \approx T_q$. Or in another word, the singularity is replaced by a transition from a black hole to a white hole. The scenario depicted in Ref. [1] is manifestly similar to that in loop quantum cosmology (LQC), where the cosmological singularity is replaced by a bounce from an attracting universe to an expanding universe [101–106]. Furthermore, the values of the quantum parameters considered in Refs. [3, 4] transition surface at finite T always result in spacetime asymmetry between the black hole region and the white hole region, which would lead to the problem of amplification of mass [4, 81]. Although the amplification of mass in Schwarzschild case is tiny and thus can be ignored, as pointed out in Ref. [4], whether the amplification of mass is always small enough in general cases is not justified yet. Inspired by the above works, the following questions can be raised: Is it possible to provide a more intuitive picture for the black hole to white hole transition where the original singularity is replaced by the transition surface? Could one construct a black-to-white hole solution with spacetime symmetry between black hole and white hole region such that one need not worry about the amplification of mass even in the general cases? All of the above questions will be answered in the affirmative in this work.

Subsequently, we also intend to apply this universal framework to other types of black holes in a spacetime with a cosmological constant. In light of the above motivations, in this paper we will firstly show that the dynamics of the geometry inside the horizon is always independent of the choice of the lapse function no matter if the quantum correction is taken

into account. This observation links the black-to-white solution in Refs. [1, 2] to the one in Refs. [3, 4]. The leading order expansion of the former solution near the transition surface is identical to that of the latter solution with the same lapse function. Then, we focus on the symmetry between the black hole and the white hole. It is found that the effective metric may exhibit a symmetric behavior when the quantum parameters are appropriately fixed such that the trouble of mass amplification can be avoided. Finally, we extend the above discussion to construct black-to-white hole solutions with a cosmological constant. The numerical solutions for the interior of black-to-white hole with a cosmological constant are obtained and the symmetric behavior of the metric between the black hole region and the white hole region is also discussed.

The paper is organized as follows. In section II, we show that the dynamics of a system with Hamiltonian constraint is equivalent under the different choice of the lapse function, no matter if this system is classical or semi-classical. Then, we apply this to discuss the dynamics of the interior of Schwarzschild black hole and the corresponding black-to-white hole with quantum corrections. We will focus on the symmetry between the black hole region and white hole region with different lapse functions. In section III, we extend this discussion to construct black-to-white hole solutions with a cosmological constant, showing the effect of the cosmological constant on the geometry and analyzing the spacetime symmetry. In particular, we propose an iteration method to obtain the values of quantum parameters in the presence of a cosmological constant to guarantee that the area enclosed by two plaquettes on the transition surface remains to be the area gap. The paper ends with conclusions and discussions in section IV.

To simplify the notation, in this work we adopt the Planck units $G = k_B = c = \hbar = 1$.

II. THE DYNAMICS OF THE INTERIOR REGION OF SCHWARZSCHILD BLACK HOLE

A. The classical dynamics in canonical formalism

In this section we will demonstrate that for a system with Hamiltonian constraint, one is free to choose the lapse function to investigate its dynamics since the system is invariant under the re-parameterization of the time coordinate.

It is well known that for a system with Hamiltonian constraint, one is free to choose

the lapse function which corresponds to the specification of the time coordinate. To see this, consider two different lapse functions N_1 and N_2 such that the resultant Hamiltonians are $H(N_1) = N_1 h$ and $H(N_2) = N_2 h$, respectively, with the Hamiltonian constraint $h \approx 0$. The corresponding time coordinates are denoted as T_1 and T_2 , respectively. The reparameterization invariance requires that the spacetime geometry is described by the same metric, namely $ds^2 = -N_1^2 dT_1^2 + \dots = -N_2^2 dT_2^2 + \dots$.

Obviously, the above relation holds if and only if two time coordinates are transformed as follows

$$\frac{dT_1}{dT_2} = \frac{N_2}{N_1}. \quad (1)$$

With the above condition, it is straightforward to show that the canonical equations of motion with lapse function N_1 are equivalent to those with lapse function N_2 . Specifically, taking the canonical equation for variable b (which will appear as the variable for Schwarzschild black hole in the next) as an example, one has

$$\frac{db}{dT_1} = \frac{\partial(N_1 h)}{\partial p_b} = \left(N_1 \frac{\partial h}{\partial p_b} + h \frac{\partial N_1}{\partial p_b} \right) = N_1 \frac{\partial h}{\partial p_b}. \quad (2)$$

On the other hand, applying the relation in Eq. (1), one can directly derive the canonical equation with T_2 as the time coordinate as follows

$$\frac{db}{dT_2} = \frac{dT_1}{dT_2} \frac{db}{dT_1} = \frac{N_2}{N_1} \frac{db}{dT_1} = \frac{N_2}{N_1} N_1 \frac{\partial h}{\partial p_b} = N_2 \frac{\partial h}{\partial p_b} = \frac{\partial(N_2 h)}{\partial p_b}. \quad (3)$$

Therefore, if the solution to Eq. (2) is $b(T_1)$, then $b(T_2) := b(T_1(T_2))$ constitutes the solution to Eq. (3), where the time transformation $T_1(T_2)$ is given by Eq. (1). The equations of motion, as well as the solutions for other canonical variables, can be derived analogously. Thus, it is concluded that for a system with Hamiltonian constraint, one is free to choose the lapse function N , which corresponds to choose a specific time coordinate for the system, but at root all the dynamics with different lapse functions are equivalent. In particular, we stress that the above argument is applicable to all the systems with Hamiltonian constraint, including the effective theory of black holes after the quantum correction is taken into account, where just a different form of Hamiltonian is introduced.

Next, we briefly review the canonical formalism of gravity vacuum in terms of the Ashtekar variables for a spacetime with spherical symmetry closely following the scheme presented in Refs. [3, 4]. In a homogeneous but non-isotropic Kantowski-Sachs spacetime, the connection

as the configuration variable and the spatial triad as the conjugate variable are reduced to the following form [107]

$$A_a^i \tau_i dx^a = \bar{c} \tau_3 dx + \bar{b} r_o \tau_2 d\theta - \bar{b} r_o \tau_1 \sin \theta d\phi + \tau_3 \cos \theta d\phi, \quad (4)$$

$$E_i^a \tau^i \partial_a = \bar{p}_c r_o^2 \tau_3 \sin \theta \partial_x + \bar{p}_b r_o \tau_2 \sin \theta \partial_\theta - \bar{p}_b r_o \tau_1 \partial_\phi, \quad (5)$$

where the triad variables are defined as $\bar{c} = c/L_o$, $\bar{b} = b/r_o$, $\bar{p}_c = p_c/r_o^2$, and $\bar{p}_b = p_b/L_o r_o$. L_o is the length of the fiducial cell and r_o is the radius in the fiducial metric, while τ_i are SU(2) generators related to Pauli spin matrices σ_i . As a result, the phase space is described by two pairs of conjugate variables (c, p_c) and (b, p_b) . The corresponding Poisson brackets are given by $\{c, p_c\} = 2\gamma$ and $\{b, p_b\} = \gamma$ with the Barbero-Immirzi parameter γ [3, 4].

The interior of the spacetime inside the horizon is homogeneous such that the metric is only time-dependent, which takes the form as

$$ds^2 = -N^2 dT^2 + \frac{p_b^2}{|p_c| L_o^2} dx^2 + |p_c| d\Omega^2, \quad (6)$$

where $d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$ represents the line element of the spherical surface, and T is the time coordinate associated with the lapse function N . Plugging the form of the metric into the action, one can derive the Hamiltonian formalism with a constraint. According to Refs. [74, 75], the classical Hamiltonian constraint is

$$\mathcal{H} = -\frac{N}{2\gamma^2} \frac{\text{sgn}(p_c)}{\sqrt{|p_c|}} [2bc p_c + (b^2 + \gamma^2) p_b]. \quad (7)$$

In order to derive the canonical equations, one specifies the lapse function as [74, 75]

$$N(T) = \frac{\gamma \text{sgn}(p_c) |p_c(T)|^{\frac{1}{2}}}{b}, \quad (8)$$

and denotes the corresponding time variable as T . By substituting Eq. (8) into Eq. (7), the Hamiltonian reads as [81]

$$\mathcal{H}(N(T)) = -\frac{1}{2\gamma} \left[2c p_c + \left(b + \frac{\gamma^2}{b} \right) p_b \right], \quad (9)$$

which leads to the equations of motion as follows

$$\dot{b} = \{b, \mathcal{H}\} = \gamma \frac{\partial \mathcal{H}}{\partial p_b} = -\frac{1}{2b} (b^2 + \gamma^2), \quad (10a)$$

$$\dot{c} = \{c, \mathcal{H}\} = 2\gamma \frac{\partial \mathcal{H}}{\partial p_c} = -2c, \quad (10b)$$

$$\dot{p}_b = \{p_b, \mathcal{H}\} = -\gamma \frac{\partial \mathcal{H}}{\partial b} = \frac{p_b}{2b^2} (b^2 - \gamma^2), \quad (10c)$$

$$\dot{p}_c = \{p_c, \mathcal{H}\} = -2\gamma \frac{\partial \mathcal{H}}{\partial c} = 2p_c, \quad (10d)$$

where the ‘‘dot’’ denotes time derivative with respect to T . By solving the dynamical equations above with the Hamiltonian constraint $H(N) \approx 0$ and properly fixing the integration constants, one obtains the solutions as

$$p_c(T) = 4m^2 e^{2T}, c(T) = \frac{\gamma L_o}{4m} e^{-2T}, p_b(T) = -2mL_o e^T (e^{-T} - 1)^{\frac{1}{2}}, b(T) = \gamma (e^{-T} - 1)^{\frac{1}{2}}, \quad (11)$$

which give rise to the metric for the interior of Schwarzschild black hole as follows

$$ds^2 = -\frac{4m^2 e^{2T}}{e^{-T} - 1} dT^2 + (e^{-T} - 1) dx^2 + (4m^2 e^{2T})^2 d\Omega^2. \quad (12)$$

In this coordinate system, the event horizon of Schwarzschild black hole is located at $T = 0$ and the singularity occurs at $T = -\infty$, which indicates the covering region of metric (12) is $(-\infty, 0)$.

Next we intend to link the metric form adopted in Refs. [1, 2] to the above metric by explicit coordinate transformation. Based on the above discussion, we choose a new lapse function as follows

$$N'(\tau) = \frac{2\gamma \operatorname{sgn}(p_c) |p_c|^{\frac{1}{4}}}{b}, \quad (13)$$

and the corresponding time coordinate is denoted as τ . By substituting Eq. (13) into Eq. (7), the corresponding Hamiltonian reads as

$$\mathcal{H}(N') = -\frac{1}{\gamma} \left[2p_c^{3/4} c + \frac{p_b}{p_c^{1/4}} \left(b + \frac{\gamma^2}{b} \right) \right]. \quad (14)$$

The equations of motion are

$$\dot{b}(\tau) = -\frac{1}{p_c^{1/4} b} (b^2 + \gamma^2), \quad (15a)$$

$$\dot{c}(\tau) = -4p_c^{-1/4} c \quad (15b)$$

$$\dot{p}_b(\tau) = \frac{p_b}{p_c^{1/4} b^2} (b^2 - \gamma^2), \quad (15c)$$

$$\dot{p}_c(\tau) = 4p_c^{3/4}, \quad (15d)$$

where the ‘‘dot’’ denotes time derivative with respect to τ and $H \approx 0$ is used to simplify the equations. The time transformation is given by

$$\frac{d\tau}{dT} = \frac{N}{N'} = \frac{p_c^{1/4}}{2} = \sqrt{\frac{m}{2}} e^{\frac{T}{2}}, \quad (16)$$

and one gets $\tau = \sqrt{2me^{\frac{T}{2}}}$ and $T = 2 \ln(\tau/\sqrt{2m})$. The integration constant is determined by requiring the singularity to be located at $\tau = 0$. The solutions to Eq. (15) are then

$$b(\tau) = \frac{\gamma\sqrt{2m-\tau^2}}{\tau}, c(\tau) = -\frac{L_o m \gamma}{\tau^4}, p_b(\tau) = L_o \tau \sqrt{2m-\tau^2}, p_c(\tau) = \tau^4. \quad (17)$$

Notably, $p_c c$ is a Dirac observable, so that one can define $p_c c = L_o m \gamma$. Now, plugging Eq. (17) into Eq. (6), one finds the metric is expressed as

$$ds^2 = -\frac{4\tau^4}{2m-\tau^2} d\tau^2 + \frac{2m-\tau^2}{\tau^2} dx^2 + \tau^4 d\Omega^2, \quad (18)$$

which is nothing but the metric form for the interior of Schwarzschild black hole presented in Refs. [1, 2]. Now the event horizon is located at $\tau = \sqrt{2m}$ and the singularity occurs at $\tau = 0$. The original region of the interior of Schwarzschild black hole is $0 < \tau < \sqrt{2m}$. It is observed in Refs. [1, 2] that if one introduces some quantum parameter l into the metric to eliminate the singularity, then one may extend the time coordinate of metric (18) to $-\sqrt{2m} < \tau < \sqrt{2m}$, leading to a white hole region

$$ds^2 = -\frac{4(\tau^2+l)^2}{2m-\tau^2} d\tau^2 + \frac{2m-\tau^2}{\tau^2+l} dx^2 + (\tau^2+l)^2 d\Omega^2, \quad (19)$$

where $l \sim m^{1/3}$ is a constant depending on the mass of black hole. It is noticed that this black-to-white hole exhibits an exact symmetric behavior between the black hole region and the white hole region since the metric is invariant under the transformation $\tau \rightarrow -\tau$. Conversely, it is not manifest to extend spacetime to a white hole region simply by introducing some quantum parameter into the metric in Eq. (12) and then by time extension. The above discussion indicates that the choice of the lapse function does not alter the geometry of spacetime (the similar conclusion is also presented in Ref. [108]), but might constrain the possible extension of spacetime coordinate after some quantum corrections are taken into account. In next subsection, we firstly review the effective black-to-white hole solution for the interior based on the Hamiltonian (7) and the lapse function (8), and then investigate its dynamics by changing the lapse function to Eq. (13).

B. Effective dynamics of the interior Schwarzschild black-to-white hole

The holonomy correction plays a core role in LQG and LQC. The key point is that the local variables which are defined at specific point of spacetime should be replaced by the non-local variables which are defined over the small region of spacetime, such as the holonomy of the connection. This replacement may change the evolution picture of the universe as well as the black hole dramatically. Following Refs. [3, 4], we first consider lapse function (8) with time coordinate T . By implementing the replacement $b \rightarrow \sin(\delta_b b)/\delta_b$ and $c \rightarrow \sin(\delta_c c)/\delta_c$ [74, 76], where the quantum parameters δ_b and δ_c reflect the QG effect with holonomy corrections. The effective lapse function is expressed as

$$N_{\text{eff}}(T) = \frac{\gamma \operatorname{sgn}(p_c) \delta_b |p_c(T)|^{\frac{1}{2}}}{\sin(\delta_b b)}, \quad (20)$$

where we denote the corresponding time variable by T for simplicity, but it should be kept in mind that now the time coordinate is different from that in classical theory. The effective Hamiltonian is then formulated as

$$\mathcal{H}_{\text{eff}}(N_{\text{eff}}) = -\frac{1}{2\gamma} \left[2 \frac{\sin(\delta_c c)}{\delta_c} p_c + p_b \left(\frac{\sin(\delta_b b)}{\delta_b} + \frac{\gamma^2 \delta_b}{\sin(\delta_b b)} \right) \right]. \quad (21)$$

It is evident that for $\delta_b \rightarrow 0$ and $\delta_c \rightarrow 0$, the classical lapse function and the Hamiltonian are restored. However, for non-vanishing δ_b and δ_c , both the lapse function and the Hamiltonian are distinct from the classical cases (i.e., Eq. (7) and Eq. (8)), thus a different dynamical behavior is expected. As a matter of fact, the equations of motion for connection and triad components are derived as follows:

$$\dot{b} = -\frac{1}{2} \left(\frac{\sin(\delta_b b)}{\delta_b} + \frac{\gamma^2 \delta_b}{\sin(\delta_b b)} \right), \quad (22a)$$

$$\dot{c} = -2 \frac{\sin(\delta_c c)}{\delta_c} \quad (22b)$$

$$\dot{p}_b = \frac{p_b}{2} \cos(\delta_b b) \left(1 - \frac{\gamma^2 \delta_b^2}{\sin^2(\delta_b b)} \right), \quad (22c)$$

$$\dot{p}_c = 2p_c \cos(\delta_c c). \quad (22d)$$

The solutions are

$$\cos(\delta_b b(T)) = b_0 \tanh \left[\frac{1}{2} \left(b_0 T + 2 \tanh^{-1} \left(\frac{1}{b_0} \right) \right) \right], \quad (23a)$$

$$\tan \left(\frac{\delta_c c(T)}{2} \right) = \mp \frac{\gamma L_0 \delta_c}{8m} e^{-2T}, \quad (23b)$$

$$p_b(T) = -2m\gamma L_o \frac{\delta_b \sin(\delta_b b(T))}{\sin^2(\delta_b b(T)) + \gamma^2 \delta_b^2}, \quad (23c)$$

$$p_c(T) = 4m^2 \left(e^{2T} + \frac{\gamma^2 L_o^2 \delta_c^2}{64m^2} e^{-2T} \right), \quad (23d)$$

where $b_0 = \sqrt{1 + \gamma^2 \delta_b^2}$. From Eq. (23d), it is found that p_c takes a nonzero minimum value at special time $T_{\mathcal{T}}$

$$l^2 := p_c|_{\min} = p_c(T_{\mathcal{T}}) = L_o \gamma m \delta_c, \quad (24)$$

$$T_{\mathcal{T}} = \frac{1}{2} \ln \left(\frac{l^2}{8m^2} \right). \quad (25)$$

The space-like hypersurface at $T = T_{\mathcal{T}}$ is denoted by \mathcal{T} . The analysis in Refs. [3, 4] demonstrates that this hypersurface is a transition surface that separates the trapped region and anti-trapped region. In the classical limit $\delta_b, \delta_c \rightarrow 0$, one has $T_{\mathcal{T}} \rightarrow -\infty$, corresponding to the singularity of the classical black hole. In this sense, the transition surface \mathcal{T} supplants the classical singularity in the quantum corrected geometry. The interior region of the classical black hole, which corresponds to $-\infty < T < 0$, now maps to the region $T_{\mathcal{T}} < T < T_{\text{BH}}$, with $T_{\text{BH}} = 0$ being the location of the black hole horizon. In the absence of the singularity, now it can be extended to a white hole region, corresponding to $T_{\text{WH}} < T < T_{\mathcal{T}}$, where $T_{\text{WH}} = -(4/b_0) \tanh^{-1}(1/b_0)$ denotes the location of the white hole horizon.

Next, let us consider the dynamics with the lapse function in Eq. (13). Once the quantum correction is taken into account, the effective lapse function and effective Hamiltonian become

$$N'_{\text{eff}}(\tau) = \frac{2\gamma \operatorname{sgn}(p_c) \delta_b |p_c(\tau)|^{\frac{1}{4}}}{\sin(\delta_b b)}, \quad (26)$$

$$\mathcal{H}_{\text{eff}}(N'_{\text{eff}}) = -\frac{1}{\gamma} \left[2p_c^{3/4} \frac{\sin(\delta_c c)}{\delta_c} + \frac{p_b}{p_c^{1/4}} \left(\frac{\sin(\delta_b b)}{\delta_b} + \frac{\gamma^2 \delta_b}{\sin(\delta_b b)} \right) \right]. \quad (27)$$

Here the corresponding time variable is still denoted by τ . The new equations of motion now read

$$\dot{b}(\tau) = -\frac{1}{p_c^{1/4}} \left(\frac{\sin(\delta_b b)}{\delta_b} + \frac{\gamma^2 \delta_b}{\sin(\delta_b b)} \right), \quad (28a)$$

$$\dot{c}(\tau) = -4p_c^{-1/4} \frac{\sin(\delta_c c)}{\delta_c}, \quad (28b)$$

$$\dot{p}_b(\tau) = p_c^{-1/4} p_b \cos(\delta_b b) \left(1 - \frac{\gamma^2 \delta_b^2}{\sin^2(\delta_b b)} \right), \quad (28c)$$

$$\dot{p}_c(\tau) = 4p_c^{3/4} \cos(\delta_c c). \quad (28d)$$

Two time coordinates T and τ transform as follows

$$\frac{d\tau}{dT} = \frac{N_{\text{eff}}}{N'_{\text{eff}}} = \frac{p_c^{1/4}}{2} = \sqrt{\frac{m}{2}} \left(e^{2T} + \frac{\gamma^2 L_o^2 \delta_c^2}{64m^2} e^{-2T} \right)^{1/4}, \quad (29)$$

$$\tau(T) = \frac{\Gamma\left(\frac{7}{8}\right)}{\Gamma\left(\frac{3}{8}\right)} \sqrt{\pi l} - \frac{\left(64m^2 e^{2T} + \frac{l^4}{m^2} e^{-2T}\right)^{1/4}}{2 \left(1 + \frac{64m^4}{l^4} e^{4T}\right)^{1/4}} {}_2F_1\left(-\frac{1}{4}, -\frac{1}{8}; \frac{7}{8}; \frac{64m^4}{l^4} e^{4T}\right), \quad (30)$$

where l is given by Eq. (24) and ${}_2F_1$ is the hypergeometric function. Integration constant is fixed by requiring the transition surface to be located at $\tau_{\mathcal{T}} = 0$. Then, the solutions to Eq. (28) are given by Eq. (23) composed with time transformation $T(\tau)$, where $T(\tau)$ is the inverse function of Eq. (30). The horizons of black hole and white hole are now located at $\tau_{\text{BH}} = \tau(T_{\text{BH}})$ and $\tau_{\text{WH}} = \tau(T_{\text{WH}})$, respectively.

Following Refs. [3, 4], the quantum parameter δ_b is considered to be the length of each link constituting the plaquette within the $\theta - \phi$ 2-sphere, while δ_c is the fractional length of the links in the x -direction within the plaquette in the $\theta - x$ and $\phi - x$ planes in a fiducial cell. By requiring that the area enclosed by these two plaquettes on the transition surface \mathcal{T} is equal to the area gap Δ :

$$2\pi\delta_c\delta_b|p_b|_{\mathcal{T}} = \Delta, \quad 4\pi\delta_b^2 p_c|_{\mathcal{T}} = \Delta, \quad (31)$$

the quantum parameters δ_b and δ_c can be constrained in the large m limit as follows:

$$\delta_b = \left(\frac{\sqrt{\Delta}}{\sqrt{2\pi\gamma^2 m}} \right)^{1/3}, \quad L_o\delta_c = \frac{1}{2} \left(\frac{\gamma\Delta^2}{4\pi^2 m} \right)^{1/3}. \quad (32)$$

It is noteworthy that both quantum parameters are proportional to $m^{-1/3}$, whereas then l goes as $m^{1/3}$. Now we consider the dynamics of the interior with respect to the time coordinate τ . The time evolution of p_c , p_b , c and b in the interior of black-to-white hole are illustrated in Fig. 1.

In Fig. 1, the red dashed curves represent the dynamics of the metric components and their conjugate variables of classical Schwarzschild black hole. It should be noted that all these curves terminate at $\tau = 0$ due to the existence of the singularity, and the rest can be obtained by the time evolution from $-\tau$ to 0 and from τ to 0. The black solid curves describe the evolution of the metric components for a quantum corrected Schwarzschild




FIG. 1. Time (τ) evolution of p_c , p_b , c and b in the Schwarzschild interior for $m = 10^6$, $L_o = \Delta = \gamma = 1$, $\delta_b = \left(\sqrt{\Delta}/\sqrt{2\pi\gamma^2 m}\right)^{1/3}$ and $L_o\delta_c = (\gamma\Delta^2/4\pi^2 m)^{1/3}/2$. The black solid curve represents the effective Schwarzschild black-to-white hole case, while the red dashed curve denotes the classical Schwarzschild black hole case.

black-to-white hole, which evolves in time from τ_{BH} to τ_{WH} . It is evident that these curves are continuous and the metric components are not vanishing at $\tau = 0$, indicating that the singularity is erased by QG effect, and this opens a window for matter passing from the black hole region to the white hole region.

C. The symmetry of the geometry between black hole and white hole

From the perspective of symmetry, Fig. 1 shows that $p_c(\tau)$ and $p_b(\tau)$ are approximately even functions, whereas $c(\tau)$ and $b(\tau)$ are approximately odd functions. Therefore, there exists an approximate symmetry between the black hole region and the white hole region. However, this symmetry is not exact. In fact, when the quantum parameters δ_b and δ_c are set as Eq. (32), one can derive the ratio of the black hole mass to the white hole mass as

$$\frac{m_{\text{WH}}}{m_{\text{BH}}} = 1 + \mathcal{O}\left[\left(\frac{\ell_{\text{Pl}}}{m}\right)^{\frac{2}{3}} \ln\left(\frac{m}{\ell_{\text{Pl}}}\right)\right], \quad (33)$$

which implies that the masses of the black hole and white hole are equal only in the limit of large m .

We intend to propose a different scheme to fix quantum parameters to guarantee that the metric of the black-to-white hole exhibits an exact symmetric behavior between the black hole region and the white hole region. As a result, the mass of the black hole is exactly equal to that of the white hole. Utilizing Eq. (23), it is observed that

$$p_c(T_{\text{BH}}) - p_c(T_{\text{WH}}) = \frac{1}{16m^2} (e^{-2T_{\text{WH}}} - 1) (64m^4 e^{2T_{\text{WH}}} - l^4). \quad (34)$$

Then, by setting $p_c(T_{\text{BH}}) = p_c(T_{\text{WH}})$, one obtains that $8m^2 e^{T_{\text{WH}}} = l^2$, or

$$\exp\left[-\frac{4}{b_0} \tanh^{-1}\left(\frac{1}{b_0}\right)\right] = \frac{L_o \gamma \delta_c}{8m}, \quad (35)$$

where $b_0 = \sqrt{1 + \gamma^2 \delta_b^2}$. Note that this condition can also be written as $\cos(\delta_b b(T_{\mathcal{T}})) = 0$ or $T_{\text{WH}} = 2T_{\mathcal{T}}$. Therefore, if we set δ_b as before, but change the setup of $L_o \delta_c$ as above, then the quantum parameters become

$$\delta_b = \left(\frac{\sqrt{\Delta}}{\sqrt{2\pi\gamma^2 m}}\right)^{1/3}, \quad L_o \delta_c = \frac{8m}{\gamma} \exp\left[-\frac{4}{b_0} \tanh^{-1}\left(\frac{1}{b_0}\right)\right]. \quad (36)$$

Now, one finds that the radius of black hole horizon $r_{\text{BH}} = \sqrt{p_c(T_{\text{BH}})}$ is equal to the radius of white hole horizon $r_{\text{WH}} = \sqrt{p_c(T_{\text{WH}})}$. Obviously, the leading order of $L_o \delta_c$ is consistent with Eq. (32), namely $L_o \delta_c = \frac{1}{2}(\gamma \Delta^2 / 4\pi^2 m)^{1/3} + \mathcal{O}(m^{-4/3})$. One can treat the set in Eq. (36) as the modification of Eq. (32). Furthermore, by virtue of the solutions presented in Eq. (23), one can directly verify that $p_b(T)$ and $p_c(T)$ are even functions with respect to $T = T_{\mathcal{T}}$, while $b(T)$ and $c(T)$ are odd functions with respect to $T = T_{\mathcal{T}}$. In fact, we have

$$p_c(T_{\mathcal{T}} + T) = 4m^2 (e^{T_{\text{WH}}+2T} + e^{2T_{\text{WH}}} e^{-T_{\text{WH}}-2T}) = p_c(T_{\mathcal{T}} - T), \quad (37)$$

$$\begin{aligned} c(T_{\mathcal{T}} + T) + c(T_{\mathcal{T}} - T) &= \frac{2}{\delta_c} [\tan^{-1}(\mp e^{T_{\text{WH}}} e^{-T_{\text{WH}}-2T}) + \tan^{-1}(\mp e^{T_{\text{WH}}} e^{-T_{\text{WH}}+2T})] \\ &= \frac{2}{\delta_c} \left(n + \frac{\pi}{2}\right). \end{aligned} \quad (38)$$

The symmetry of $b(T)$ and $p_b(T)$ can be similarly verified. Additionally, one can confirm that the time transformation $\tau(T)$ is an odd function with respect to $T = T_{\mathcal{T}}$. Consequently, $p_b(\tau)$ and $p_c(\tau)$ are even functions with respect to $\tau = 0$, while $b(\tau)$ and $c(\tau)$ are odd functions with respect to $\tau = 0$. Thus, the metric exhibits the symmetry under the

transformation $\tau \rightarrow -\tau$ as well. We will prove this symmetry again with a different manner in section III C, where a cosmological constant is involved.

Thanks to the symmetry of the metric, all curvature scalars are even functions and attain their extremal values at $\tau = 0$. Recall that \mathcal{T} corresponds to the singularity of classical black hole, one expects that $\tau = 0$ should be a point with the maximal value. One can justify this through numerical analysis, as demonstrated in Fig. 2, where the Kretschmann scalar has a maximal value at \mathcal{T} .




FIG. 2. Time (τ) evolution of Kretschmann scalar in Schwarzschild interior for $m = 10^6$, $L_o = \Delta = \gamma = 1$, $\delta_b = \left(\sqrt{\Delta}/\sqrt{2\pi\gamma^2 m}\right)^{1/3}$ and $L_o\delta_c = \frac{8m}{\gamma} \exp\{-[4\tanh^{-1}(1/b_0)]/b_0\}$. The black solid curve represents the effective Schwarzschild black-to-white hole case, while the red dashed curve denotes the classical Schwarzschild black hole case.

Next, we intend to embed the metric proposed in Refs. [1, 2] into the framework of LQG by showing that it can be viewed as the approximation solution of the above equations. Let us examine the behavior of the metric near the transition surface. The components of metric can be expressed as

$$-N_{\text{eff}}^2(\tau) = -4l(b_0^2 - 1) - 4(b_0^2 - 1)(4 + b_0^4)\tau^2 + \mathcal{O}(\tau^4), \quad (39)$$

$$\frac{p_b(\tau)^2}{|p_c(\tau)|L_o^2} = \frac{4(b_0^2 - 1)m^2}{b_0^4 l^2} - \frac{4(b_0^2 - 1)(b_0^4 - 2b_0^2 + 8)m^2}{b_0^4 l^3}\tau^2 + \mathcal{O}(\tau^4), \quad (40)$$

$$p_c(\tau) = l^2 + 8l\tau^2 + \mathcal{O}(\tau^6), \quad (41)$$

where b_0 and l satisfy condition (35). Note that $l \sim m^{1/3}$, $b_0 \sim 1$ and $(b_0^2 - 1) = \gamma^2 \delta_b^2 \sim m^{-2/3}$,

the effective metric can be written as

$$\begin{aligned} ds^2 &= -N_{\text{eff}}^2(\tau) d\tau^2 + \frac{p_b(\tau)^2}{|p_c(\tau)| L_o^2} dx^2 + |p_c(\tau)| d\Omega^2 \\ &\sim -\left(m^{-\frac{1}{3}} + m^{-\frac{2}{3}}\tau^2\right) d\tau^2 + \left(m^{\frac{2}{3}} - m^{\frac{1}{3}}\tau^2\right) dx^2 + \left(m^{\frac{2}{3}} + m^{\frac{1}{3}}\tau^2\right) d\Omega^2 + \mathcal{O}(\tau^4). \end{aligned} \quad (42)$$

On the other hand, the metric (19) which is given in Refs. [1, 2] can also be written as

$$\begin{aligned} ds^2 &= -\frac{4(\tau^2 + l)^2}{2m - \tau^2} d\tau^2 + \frac{2m - \tau^2}{\tau^2 + l} dx^2 + (\tau^2 + l)^2 d\Omega^2 \\ &= \left(-\frac{2l^2}{m} - \frac{l^2 + 4lm}{m^2}\tau^2\right) d\tau^2 + \left(\frac{2m}{l} - \frac{l + 2m}{l^2}\tau^2\right) dx^2 + (l^2 + 2l\tau^2) d\Omega^2 + \mathcal{O}(\tau^4) \\ &\sim -\left(m^{-\frac{1}{3}} + m^{-\frac{2}{3}}\tau^2\right) d\tau^2 + \left(m^{\frac{2}{3}} - m^{\frac{1}{3}}\tau^2\right) dx^2 + \left(m^{\frac{2}{3}} + m^{\frac{1}{3}}\tau^2\right) d\Omega^2 + \mathcal{O}(\tau^4). \end{aligned} \quad (43)$$

Therefore, the metric (19) can be regarded as a coarse-grained version of the metric (42), which is derived by solving the effective dynamical equations.

Using the approximate metric (42), it is easy to show that the leading terms of the maximal value of the curvature scalars are mass independent:

$$R^2|_{\mathcal{T}} = 4 \left(1 + m^{-\frac{2}{3}}\right)^2, \quad (44)$$

$$R_{ab}R^{ab}|_{\mathcal{T}} = 2 \left(2 + 2m^{-\frac{2}{3}} + m^{-\frac{4}{3}}\right), \quad (45)$$

$$K^2|_{\mathcal{T}} = 4 \left(3 + m^{-\frac{3}{4}}\right), \quad (46)$$

$$C_{abcd}C^{abcd}|_{\mathcal{T}} = \frac{4}{3} \left(2 - m^{-\frac{2}{3}}\right)^2. \quad (47)$$

This is a significant characteristic of quantum corrected black holes, suggesting that the curvature scalars are bounded at the Planck scale [109]. The quantum parameter l is not at the Planck scale, but goes as $m^{1/3}$, defining a ‘‘Planck star’’ [110–113].

III. THE DYNAMICS OF SCHWARZSCHILD-(ANTI)DE SITTER SOLUTION

In this section we extend the above analysis on the interior of Schwarzschild black-to-white hole to Schwarzschild-(anti)de Sitter ((A)dS) spacetime with a cosmological constant, thereby obtaining the effective internal solution of Schwarzschild-(A)dS black-to-white holes in the framework of LQG with new quantum parameters. Previously, the dynamics of Schwarzschild-(A)dS black holes with old quantum parameters has been investigated in Refs. [78, 80, 114].

A. Classical dynamics of the interior Schwarzschild-(A)dS solution

The Einstein-Hilbert action with a cosmological constant is given by

$$S = \frac{1}{16\pi} \int d^4x \sqrt{-g} (R - 2\Lambda), \quad (48)$$

where Λ is the cosmological constant. With the same algebra, one can derive the Hamiltonian constraint in terms of Ashtekar variables as [80]:

$$\mathcal{H}^\Lambda = -\frac{N}{2\gamma^2} \frac{\text{sgn}(p_c)}{\sqrt{|p_c|}} [2bc p_c + (b^2 + \gamma^2) p_b - \Lambda p_b p_c]. \quad (49)$$

In parallel with the Schwarzschild case, we firstly consider the lapse function Eq. (8), which was studied in subsection II A. According to Eq. (48) and Eq. (49), the equations of motion now read as

$$\dot{b} = -\frac{1}{2b} (b^2 + 1 - \Lambda p_c), \quad (50a)$$

$$\dot{c} = -2c + \Lambda \frac{p_b}{b}, \quad (50b)$$

$$\dot{p}_b = \frac{p_b}{2b^2} (b^2 - 1 + \Lambda p_c), \quad (50c)$$

$$\dot{p}_c = 2p_c, \quad (50d)$$

where the “dot” denotes time derivative with respect to T and without loss of generality the Immirzi parameter γ has been set to unity here. It is straightforward to obtain the solutions to Eq. (50) as follows

$$b = \sqrt{\left(1 - \frac{\Lambda}{3} r_h^2\right) e^{-T} - 1 + \frac{\Lambda}{3} r_h^2 e^{2T}}, \quad (51a)$$

$$c = -\frac{\Lambda}{6} L_o r_h (2e^T + e^{-2T}) + \frac{L_o}{2r_h} e^{-2T}, \quad (51b)$$

$$p_b = -L_o r_h e^T \sqrt{\left(1 - \frac{\Lambda}{3} r_h^2\right) e^{-T} - 1 + \frac{\Lambda}{3} r_h^2 e^{2T}}, \quad (51c)$$

$$p_c = r_h^2 e^{2T}. \quad (51d)$$

Two integration constants are fixed by requiring that the black hole horizon lies at $T = 0$ and by always satisfying the Hamiltonian constraint. If one introduces the time coordinate $\tau := r_h e^T$ and $m := \frac{r_h}{2} \left(1 - \frac{\Lambda}{3} r_h^2\right)$, then the metric in the interior Schwarzschild-(A)dS black hole takes the following familiar form

$$ds^2 = -\left(\frac{2m}{\tau} - 1 + \frac{\Lambda}{3} \tau^2\right)^{-1} d\tau^2 + \left(\frac{2m}{\tau} - 1 + \frac{\Lambda}{3} \tau^2\right) dx^2 + \tau^2 d\Omega^2. \quad (52)$$

The black hole horizon lies at $T = 0$ and the singularity occurs at $T = -\infty$, so the black hole interior corresponds to $-\infty < T < 0$. Obviously these solutions reduce to the metric of Schwarzschild black hole in Eq. (11) if $\Lambda = 0$.

Next, to compare the results in semi-classical theory, we choose a new lapse function as follows

$$\hat{N}(\tau_\Lambda) = \frac{\gamma \operatorname{sgn}(p_c)}{b}. \quad (53)$$

The corresponding canonical equations of motion are then

$$\dot{b} = -\frac{1}{2b\sqrt{p_c}}(b^2 + 1 - \Lambda p_c), \quad (54a)$$

$$\dot{c} = \frac{1}{\sqrt{p_c}}\left(-2c + \Lambda \frac{p_b}{b}\right), \quad (54b)$$

$$\dot{p}_b = \frac{p_b}{2b^2\sqrt{p_c}}(b^2 - 1 + \Lambda p_c), \quad (54c)$$

$$\dot{p}_c = 2\sqrt{p_c}, \quad (54d)$$

where the ‘‘dot’’ denotes time derivative with respect to τ_Λ and $H \approx 0$ is used to simplify the equations. Between these two coordinate systems, the time coordinate transforms as

$$\frac{d\tau_\Lambda}{dT} = \frac{N}{\hat{N}} = \sqrt{p_c} = r_h e^T, \quad (55)$$

$$\tau_\Lambda = r_h e^T \quad \text{and} \quad T = \ln(\tau_\Lambda/r_h). \quad (56)$$

The solutions are given by

$$b = \sqrt{\frac{2m}{\tau_\Lambda} - 1 + \frac{\Lambda}{3}\tau_\Lambda^2}, \quad (57a)$$

$$c = L_o \left(\frac{m}{\tau_\Lambda^2} - \frac{\Lambda}{3}\tau_\Lambda \right), \quad (57b)$$

$$p_b = -L_o \tau_\Lambda \sqrt{\frac{2m}{\tau_\Lambda} - 1 + \frac{\Lambda}{3}\tau_\Lambda^2}, \quad (57c)$$

$$p_c = \tau_\Lambda^2. \quad (57d)$$

Substituting Eq. (57) into expression (6), one obtains the line element (52) again. Now, the singularity occurs at $\tau_\Lambda = 0$ and the black hole interior corresponds to $0 < \tau_\Lambda < r_h$.

B. Effective dynamics of the interior Schwarzschild-(A)dS black-to-white hole

The effective theory for the evolution of the interior of the Schwarzschild-(A)dS black hole can be introduced similarly as the Schwarzschild case. In this subsection, the effective

dynamics will be considered based on the Hamiltonian with the modification of the lapse function (53). Classically, the interior of the black hole corresponds to $0 < \tau_\Lambda < r_h$. By replacement $b \rightarrow \sin(\delta_b b)/\delta_b$, $c \rightarrow \sin(\delta_c c)/\delta_c$, the resulting effective Hamiltonian is given by

$$\mathcal{H}_{\text{eff}}^\Lambda(\hat{N}) = -\frac{\delta_b}{2G\gamma \sin(\delta_b b)\sqrt{p_c}} \left[2\frac{\sin(\delta_b b)}{\delta_b} \frac{\sin(\delta_c c)}{\delta_c} p_c + \left(\frac{\sin^2(\delta_b b)}{\delta_b^2} + \gamma^2 \right) p_b - \Lambda p_b p_c \right]. \quad (58)$$

This Hamiltonian leads to equations of motion:

$$\dot{b} = -\frac{\delta_b}{2\sqrt{p_c} \sin(\delta_b b)} \left(\frac{\sin^2(\delta_b b)}{\delta_b^2} + \gamma^2 - \Lambda p_c \right), \quad (59a)$$

$$\dot{c} = \frac{1}{\sqrt{p_c}} \left(-2\frac{\sin(\delta_c c)}{\delta_c} + \Lambda p_b \frac{\delta_b}{\sin(\delta_b b)} \right), \quad (59b)$$

$$\dot{p}_b = \frac{p_b \cos(\delta_b b) \delta_b^2}{2\sqrt{p_c} \sin^2(\delta_b b)} \left(\frac{\sin^2(\delta_b b)}{\delta_b^2} + \Lambda p_c - \gamma^2 \right), \quad (59c)$$

$$\dot{p}_c = 2\sqrt{p_c} \cos(\delta_c c), \quad (59d)$$

where the Hamiltonian constraint $\mathcal{H}_{\text{eff}}^\Lambda \approx 0$ is used to simplify the equations of motion. Note that these equations reduce to Eq. (54) in the classical limit $\delta_b, \delta_c \rightarrow 0$.

Due to the difficulty of solving these equations analytically, the numerical analysis of these equations is performed. Firstly, we remark that in the presence of the cosmological constant the setup of δ_b and δ_c may be relevant to the value of Λ in general case, thus we intend to provide a strategy to fix δ_b and δ_c for a given cosmological constant numerically. The key point is to require that the area enclosed by two plaquettes on the transition surface remains to be the area gap Δ , which can be implemented by an iterative method. We elaborate this strategy as following. According to Eq. (31), one gets

$$\delta_b = \sqrt{\frac{\Delta}{4\pi p_c(\delta_b, \delta_c)|_{\mathcal{T}}}}, \quad \delta_c = \frac{\Delta}{2\pi \delta_b |p_b(\delta_b, \delta_c)|_{\mathcal{T}}}. \quad (60)$$

We require that the values of δ_b and δ_c satisfy the above equations in the presence of the cosmological constant. However, it is noticed that in general case the value of p_b and p_c on the transition surface is unknown since we have no analytic solutions for them as in Schwarzschild case. Thus, numerically we first set initial values $\delta_{b,0}, \delta_{c,0}$ for δ_b and δ_c as those without cosmological constant, and then plug them into the equations of motion with the cosmological constant and numerically find the solutions for p_b and p_c on the transition surface. Subsequently, we substitute p_b and p_c into Eq. (60) and obtain the values of $\delta_{b,1}$,

$\delta_{c,1}$ as the new values of δ_b and δ_c for next iteration. Concisely, we may apply the following iterative equations

$$\delta_{b,n} = \sqrt{\frac{\Delta}{4\pi p_c(\delta_{b,n-1}, \delta_{c,n-1})|\mathcal{T}}}, \quad \delta_{c,n} = \frac{\Delta}{2\pi\delta_{b,n-1}|p_b(\delta_{b,n-1}, \delta_{c,n-1})|\mathcal{T}}, \quad (61)$$

and we stop the iteration until the area enclosed by two plaquettes on the transition surface \mathcal{T} is very close to the area gap Δ (numerically we require $\sigma := (|\Delta_n - \Delta|)/\Delta < 10^{-3}$), where $p_c(\delta_{b,n-1}, \delta_{c,n-1})$ and $p_b(\delta_{b,n-1}, \delta_{c,n-1})$ are understood as the solutions to the equations of motion with $\delta_{b,n-1}$ and $\delta_{c,n-1}$.

Next, we apply this strategy to the Schwarzschild-AdS case with a negative cosmological constant. It is worth emphasizing that in numerical simulation and in the Planck units, $\Lambda = -1$ leads to very large vacuum energy density which lies at the Planck energy density level. Therefore, we may restrict our analysis within the range $-1 < \Lambda < 0$. In addition, in the effective region of QG we are mainly concerned with the large mass of black holes. Without loss of generality we still fix the mass as $m = 10^6$. In Fig. 3, we show the variation of δ_b and δ_c with Λ . It is observed that δ_b becomes smaller with the increase of $|\Lambda|$ but δ_c becomes larger. Specially, when $|\Lambda| > 0.3$, both δ_b and δ_c can approximately be fitted by linear functions of Λ .




FIG. 3. Relationships between δ_b , δ_c and Λ in the Schwarzschild-AdS case. δ_b and δ_c can approximately be fitted by linear functions of Λ when $|\Lambda| > 0.3$. The linear fit function of δ_b is $\delta_b \approx 0.00635 - 0.00117|\Lambda|$ while the linear fit function of δ_c is $\delta_c \approx 0.00209 + 0.000847|\Lambda|$.

Now, we illustrate the specific solutions for the case with large mass and small cosmological constant. Let us set $m = 10^6$ and $\Lambda = -10^{-3}$. By iteration method we find the quantum parameters should be set as $\delta_b = 7.410 \times 10^{-3}$ and $\delta_c = 1.476 \times 10^{-3}$ to guarantee




FIG. 4. Time (τ_Λ) evolution of p_c , p_b , c and b in the quantum corrected Schwarzschild-AdS interior for $m = 10^6$, $\Lambda = -10^{-3}$, $L_o = \Delta = \gamma = 1$, $\delta_b = 7.410 \times 10^{-3}$ and $\delta_c = 1.476 \times 10^{-3}$. The black solid curve represents the effective Schwarzschild-AdS black-to-white hole case, while the red dashed curve denotes the classical Schwarzschild-AdS black hole case. The horizon of classical Schwarzschild-AdS black hole lies at $r_h \approx 1817.12$.

that the area enclosed by two plaquettes on the transition surface \mathcal{T} is the area gap Δ . In comparison with the values in the Schwarzschild case which are $\delta_b = 7.362 \times 10^{-3}$ and $\delta_c = 1.468 \times 10^{-3}$, we notice both quantities just have a small shift. Correspondingly, the evolution of p_c , p_b , c , and b are obtained by solving the equations of motion, and the results are illustrated in Fig. 4. It is noticed that the evolution of all the variables can pass through $\tau_\Lambda = 0$, signaling that the original singularity at $\tau_\Lambda = 0$ disappears. The variable p_c in Fig. 4(a) takes the minimal value $p_c|_{\min} \approx 1450$ at $\tau_\Lambda = 0$, which is slightly lower than the value in Schwarzschild case $(p_c|_{\min})_{\text{Sch}} = \gamma L_o m \delta_c \approx 1468$, but still of the same order of magnitude. Therefore, in this case the presence of the negative cosmological constant does not significantly alter the minimal value of p_c , and thus the radius of transition surface $l = \sqrt{p_c|_{\min}}$ also goes as $m^{1/3}$. On the contrary, we point out that the radius of classical




FIG. 5. Time (τ_Λ) evolution of p_c , p_b , c and b in the quantum corrected Schwarzschild-AdS interior for $m = 10^6$, $\Lambda = -1$, $L_o = \Delta = \gamma = 1$, $\delta_b = 5.216 \times 10^{-3}$ and $\delta_c = 2.913 \times 10^{-3}$. The black solid curve represents the effective Schwarzschild-AdS black-to-white hole case, while the red dashed curve denotes the classical Schwarzschild-AdS black hole case. The horizon of classical Schwarzschild-AdS black hole lies at $r_h \approx 181.7$.

Schwarzschild-AdS black hole $r_h = (-6m/\Lambda)^{1/3} + \mathcal{O}(m^{-1/3}) \approx 1817.12$ also goes as $m^{1/3}$ and thus is dramatically less than the radius of classical Schwarzschild black hole with the same mass since $(r_h)_{\text{Sch}} = 2m = 2 \times 10^6$. It is noticed that l and r_h are of the same order of the magnitude of mass, implying that the transition surface is closer to the horizon in large m limit in comparison with that in the Schwarzschild case, and the effective dynamics is significantly contrast to the classical dynamics in the interior of Schwarzschild-AdS black hole. In addition, according to Fig. 4(b), it can be found that the classical case (red dashed curve) is strictly symmetric with respect to $\tau = 0$, whereas the quantization case (black solid curve) is not, which suggests that the effective theory with a cosmological constant after quantization breaks the symmetry of spacetime.

Then, we consider the specific solutions with large mass and large cosmological constant




FIG. 6. Time (τ_Λ) evolution of p_c , p_b , c and b in the quantum corrected Schwarzschild-dS interior for $m = 10^6$, $\Lambda = (1/9) \times 10^{-12}$, $L_o = \Delta = \gamma = 1$, $\delta_b = 7.362 \times 10^{-3}$ and $\delta_c = 1.468 \times 10^{-3}$. The black solid curve represents the effective Schwarzschild-dS black-to-white hole case, while the red dashed curve denotes the classical Schwarzschild-dS black hole case. The horizon of classical Schwarzschild-dS black hole lies at $r_h = 3 \times 10^6$.

with $\Lambda = -1$ and $m = 10^6$. In this case we find $\delta_b = 5.216 \times 10^{-3}$ and $\delta_c = 2.913 \times 10^{-3}$ by iteration method, and the evolution of spacetime is illustrated in Fig. 5. It is worthy to notice that the minimal value of p_c is larger than the value in $\Lambda = -10^{-3}$ case, which means the transition surface is closer to the event horizon in extreme case. In fact, it is found that $l = \sqrt{p_c|_{\min}} \approx 54.0$ while $r_{\text{BH}} = \sqrt{p_c|_{\text{BH}}} \approx 179.8$, so that we have $l \approx 0.3r_{\text{BH}}$. Furthermore, the radius of the white hole horizon $r_{\text{WH}} = \sqrt{p_c|_{\text{WH}}} \approx 286.0$ is significantly larger than r_{BH} . We recall that, in the Schwarzschild case with the same mass $m = 10^6$, the black hole mass agrees with the white hole mass, namely $m_{\text{BH}} = m_{\text{WH}} + \mathcal{O}(1)$. This relation does not hold when $\Lambda = -1$, and the negative cosmological constant disrupts the approximate symmetry between black hole and white hole. This can also be seen from Fig. 5(a) and Fig. 5(b): these two figures exhibit manifest asymmetry between the black hole region and the white hole region. Of course, we remark that in general this discrepancy depends on the ratio of

the mass of black hole to the square root of the cosmological constant, namely $m/\sqrt{|\Lambda|}$. Numerically, we find when $m/\sqrt{|\Lambda|} > 10^8$, then the effect of the cosmological constant is limited and the dynamics is similar to the Schwarzschild case, while if $m/\sqrt{|\Lambda|} < 10^6$, then the effect of the cosmological constant becomes notable, and may leads to manifest asymmetry between the black hole region and the white hole region.

Next, we turn to consider the Schwarzschild-dS black holes with a positive cosmological constant, which is slightly different from the Schwarzschild-AdS case. The classical Schwarzschild-dS spacetime has two horizons when $0 < 9\Lambda m^2 < 1$, one degenerate horizon when $9\Lambda m^2 = 1$, and no horizon when $9\Lambda m^2 > 1$. For simplicity, we consider $9\Lambda m^2 = 1$, and similar conclusions can be obtained for the case where $0 < 9\Lambda m^2 < 1$. When $9\Lambda m^2 = 1$, the event horizon lies at $r_h = 3m = 1/\sqrt{\Lambda}$. This is the so called Nariai black hole [115]. This black hole exhibits distinct evolutionary behavior with different values of mass. One extremal case is that the mass of the black hole is very large, for instance, setting the mass of the black hole to be 10^6 , then $\Lambda = (1/9) \times 10^{-12}$. The evolution of such black-to-white hole is illustrated in Fig. 6. Similarly, by requiring that quantum parameters δ_b and δ_c satisfy relation (31) and employing the iterative method, we find that $\delta_b = 7.362 \times 10^{-3}$ and $\delta_c = 1.468 \times 10^{-3}$, which are almost the same values as those given by Eq. (32) for $m = 10^6$. This arises from the fact that the cosmological constant is extremely small such that its effects may be neglected. Therefore, in large mass limit the effective dynamics of Schwarzschild-dS black-to-white hole is very similar to the effective dynamics of Schwarzschild black-to-white hole.

C. The symmetry of the geometry between black hole and white hole

In the previous subsection, it has been demonstrated that the presence of a cosmological constant may lead to an asymmetry between the black hole region and the white hole region and its impact depends on the ratio $m/\sqrt{|\Lambda|}$. In section II C, it is shown that in Schwarzschild case, if δ_b and δ_c satisfy the condition $\cos(\delta_b b) = 0$ at transition surface \mathcal{T} , the metric is exactly symmetric between the black hole region and the white hole region. In this subsection, we intend to prove that this conclusion holds true even in the presence of a cosmological constant. In fact, one can show that $p_b(\tau_\Lambda)$ and $p_c(\tau_\Lambda)$ are even functions, whereas $b(\tau_\Lambda)$ and $c(\tau_\Lambda)$ are odd functions if $\cos(\delta_b b(0)) = 0$ is satisfied. Thus, the metric, which depends on p_b and p_c , is also an even function, and remains invariant under the transformation $\tau \rightarrow -\tau$.

The proof can be demonstrated as follows. In order to show that b and c are odd functions, while p_b and p_c are even functions, we need to prove the following equations:

$$b(\tau_\Lambda) + b(-\tau_\Lambda) = 2b(0), \quad (62a)$$

$$c(\tau_\Lambda) + c(-\tau_\Lambda) = 2c(0), \quad (62b)$$

$$p_b(\tau_\Lambda) = p_b(-\tau_\Lambda), \quad (62c)$$

$$p_c(\tau_\Lambda) = p_c(-\tau_\Lambda). \quad (62d)$$

Then, by defining $\tilde{b}(\tau_\Lambda) = 2b(0) - b(-\tau_\Lambda)$, $\tilde{c}(\tau) = 2c(0) - c(-\tau_\Lambda)$, $\tilde{p}_b(\tau_\Lambda) = p_b(-\tau_\Lambda)$ and $\tilde{p}_c(\tau_\Lambda) = p_c(-\tau_\Lambda)$, we need to prove that $\tilde{b}(\tau_\Lambda) = b(\tau_\Lambda)$, $\tilde{c}(\tau_\Lambda) = c(\tau_\Lambda)$, $\tilde{p}_b(\tau_\Lambda) = p_b(\tau_\Lambda)$ and $\tilde{p}_c(\tau_\Lambda) = p_c(\tau_\Lambda)$. This can be shown by demonstrating that they are the same set of solutions to Eq. (59). Actually, based on the condition that $\cos(\delta_b b) = 0$ at \mathcal{T} , one can derive the following corollary

$$\cos(\delta_b b(0)) = 0 \quad \text{and} \quad \delta_b b(0) = n_b \pi + \frac{\pi}{2}. \quad (63)$$

From Eq. (59d), since p_c takes the minimal value at \mathcal{T} , one obtains

$$\cos(\delta_c c(0)) = 0 \quad \text{and} \quad \delta_c c(0) = n_c \pi + \frac{\pi}{2}. \quad (64)$$

Thus, substituting $\tilde{b}(\tau_\Lambda)$, $\tilde{c}(\tau_\Lambda)$, $\tilde{p}_b(\tau_\Lambda)$ and $\tilde{p}_c(\tau_\Lambda)$ into the right hand side of Eq. (59a), one gets

$$\begin{aligned} & - \frac{\delta_b}{2\sqrt{\tilde{p}_c(\tau_\Lambda)} \sin(\delta_b \tilde{b}(\tau_\Lambda))} \left[\frac{\sin^2(\delta_b \tilde{b}(\tau_\Lambda))}{\delta_b^2} + \gamma^2 - \Lambda \tilde{p}_c(\tau_\Lambda) \right] \\ &= - \frac{\delta_b}{2\sqrt{p_c(-\tau_\Lambda)} \sin[\pi - \delta_b b(-\tau_\Lambda)]} \left\{ \frac{\sin^2[\pi - \delta_b b(-\tau_\Lambda)]}{\delta_b^2} + \gamma^2 - \Lambda p_c(-\tau_\Lambda) \right\} \\ &= - \frac{\delta_b}{2\sqrt{p_c(-\tau_\Lambda)} \sin[\delta_b b(-\tau_\Lambda)]} \left\{ \frac{\sin^2[\delta_b b(-\tau_\Lambda)]}{\delta_b^2} + \gamma^2 - \Lambda p_c(-\tau_\Lambda) \right\} \\ &= \left. \frac{db}{d\tau_\Lambda} \right|_{-\tau_\Lambda} = \left. \frac{d\tilde{b}}{d\tau_\Lambda} \right|_{\tau_\Lambda}, \end{aligned} \quad (65)$$

which means that $\tilde{b}(\tau_\Lambda)$ also satisfies equation Eq. (59a). The other three equations can be verified similarly. We have

$$\frac{1}{\sqrt{\tilde{p}_c(\tau_\Lambda)}} \left[-2 \frac{\sin(\delta_c \tilde{c}(\tau_\Lambda))}{\delta_c} + \Lambda \tilde{p}_b(\tau_\Lambda) \frac{\delta_b}{\sin(\delta_b \tilde{b}(\tau_\Lambda))} \right] = \left. \frac{d\tilde{c}}{d\tau_\Lambda} \right|_{\tau_\Lambda}, \quad (66a)$$

$$\frac{\tilde{p}_b(\tau_\Lambda) \cos(\delta_b \tilde{b}(\tau_\Lambda)) \delta_b^2}{2\sqrt{\tilde{p}_c(\tau_\Lambda)} \sin^2(\delta_b \tilde{b}(\tau_\Lambda))} \left[\frac{\sin^2(\delta_b \tilde{b}(\tau_\Lambda))}{\delta_b^2} + \Lambda \tilde{p}_c(\tau_\Lambda) - \gamma^2 \right] = \left. \frac{d\tilde{p}_b}{d\tau_\Lambda} \right|_{\tau_\Lambda}, \quad (66b)$$

$$2\sqrt{\tilde{p}_c(\tau_\Lambda)} \cos(\delta_c \tilde{c}(\tau_\Lambda)) = \left. \frac{d\tilde{p}_c}{d\tau_\Lambda} \right|_{\tau_\Lambda}. \quad (66c)$$

The above expressions indicate that $\tilde{b}(\tau_\Lambda)$, $\tilde{c}(\tau_\Lambda)$, $\tilde{p}_b(\tau_\Lambda)$ and $\tilde{p}_c(\tau_\Lambda)$ are also solutions to Eq. (59). Furthermore, these two sets of solutions share the same initial conditions $\tilde{b}(0) = b(0)$, $\tilde{c}(0) = c(0)$, $\tilde{p}_b(0) = p_b(0)$, $\tilde{p}_c(0) = p_c(0)$, which implies that they must be identical to each other. Therefore, one gets $\tilde{b}(\tau_\Lambda) = b(\tau_\Lambda)$, $\tilde{c}(\tau_\Lambda) = c(\tau_\Lambda)$, $\tilde{p}_b(\tau_\Lambda) = p_b(\tau_\Lambda)$, $\tilde{p}_c(\tau_\Lambda) = p_c(\tau_\Lambda)$.

So far, we have shown that $p_b(\tau_\Lambda)$ and $p_c(\tau_\Lambda)$ are even functions, whereas $b(\tau_\Lambda)$ and $c(\tau_\Lambda)$ are odd functions if $\cos(\delta_b b(0)) = 0$ is satisfied. Although we prove the above property under a specific lapse function (53), it is worthwhile to emphasize that the proof holds true under any lapse function. As a matter of fact, the conditions $\cos(\delta_b b) = 0$ at \mathcal{T} do not depend on the choice of lapse function. Using the same method, one can also prove the conclusion mentioned in section II C that $p_b(T)$ and $p_c(T)$, which are the solutions to Eq. (23), are even functions with respect to $T = T_{\mathcal{T}}$, whereas $b(T)$ and $c(T)$ are odd functions with respect to $T = T_{\mathcal{T}}$ if $\cos(\delta_b b(T_{\mathcal{T}})) = 0$.

Next we discuss the value of quantum parameters which may give rise to symmetric solutions. Once δ_b is fixed, the corresponding value of δ_c leading to symmetric solutions can be obtained by numerically solving Eq. (59) with condition $\cos(\delta_b b(0)) = 0$, which is denoted

TABLE I. Values of δ_b and δ_c for different Schwarzschild and Schwarzschild-(A)dS black-to-white holes. We set $m = 10^6$, and δ_b , δ_{c1} are obtained through iteration while δ_{c2} is obtained by fixing δ_b and then numerically solving Eq. (59) with condition $\cos(\delta_b b(0)) = 0$.

The cosmological constant	δ_b	δ_{c1}	δ_{c2}
$\Lambda = 0$	7.362×10^{-3}	1.468×10^{-3}	1.469×10^{-3}
$\Lambda = -10^{-3}$	7.410×10^{-3}	1.476×10^{-3}	1.623×10^{-3}
$\Lambda = -1$	5.216×10^{-3}	2.913×10^{-3}	5.532×10^{-4}
$\Lambda = (1/9) \times 10^{-12}$	7.362×10^{-3}	1.468×10^{-3}	1.469×10^{-3}

as δ_{c2} . The various values of δ_b and δ_{c2} leading to symmetric solutions for different black-to-white holes are listed in Tab. I, where for comparison the value of δ_{c1} which is obtained through iteration is also given. The time evolution of p_c , p_b , c and b with quantum parameters δ_b , δ_{c2} is illustrated in Figs. 7-9. In those figures, the black solid curve represents the effective Schwarzschild-(A)dS black-to-white hole case, while the red dashed curve denotes the classical Schwarzschild-(A)dS black hole case. It is manifestly observed that all the functions exhibit the symmetric behavior as they perform in the analytic analysis.

About the symmetric solutions we present the following remarks. Firstly, as the cosmological constant is relatively small comparing with the mass of the black hole (numerically, the ratio $m/\sqrt{|\Lambda|} > 10^8$), then we find δ_{c1} and δ_{c2} are quite close. It indicates that the symmetric solutions also satisfy the requirement that the area enclosed by two plaquettes on the transition surface \mathcal{T} is the area gap Δ very well. On the other hand, as the cosmological constant is relatively large comparing with the mass of the black hole (numerically, the ratio $m/\sqrt{|\Lambda|} < 10^6$), then we find δ_{c1} and δ_{c2} are not close (See the data for $\Lambda = -1$ in Tab. I). In this case we find that the area enclosed by two plaquettes on the transition surface \mathcal{T} has




FIG. 7. Time (τ_Λ) evolution of p_c , p_b , c and b in the quantum corrected Schwarzschild-AdS interior for $m = 10^6$, $\Lambda = -10^{-3}$, $L_o = \Delta = \gamma = 1$, $\delta_b \approx 7.410 \times 10^{-3}$ and $\delta_{c2} \approx 1.623 \times 10^{-3}$.




FIG. 8. Time (τ_Λ) evolution of p_c , p_b , c and b in the quantum corrected Schwarzschild-AdS interior for $m = 10^6$, $\Lambda = -1$, $L_o = \Delta = \gamma = 1$, $\delta_b \approx 5.216 \times 10^{-3}$ and $\delta_{c2} \approx 5.532 \times 10^{-4}$.




FIG. 9. Time (τ_Λ) evolution of p_c , p_b , c and b in the quantum corrected Schwarzschild-dS interior for $m = 10^6$, $\Lambda = (1/9) \times 10^{-12}$, $L_o = \Delta = \gamma = 1$, $\delta_b \approx 7.362 \times 10^{-3}$ and $\delta_{c2} \approx 1.468 \times 10^{-3}$.




FIG. 10. The Kretschmann scalar in the quantum corrected Schwarzschild-(A)dS interior for (a) $m = 10^6$, $\Lambda = -10^{-3}$, $\delta_b \approx 7.410 \times 10^{-3}$ and $\delta_{c2} \approx 1.623 \times 10^{-3}$. (b) $m = 10^6$, $\Lambda = -1$, $\delta_b \approx 5.216 \times 10^{-3}$ and $\delta_{c2} \approx 5.532 \times 10^{-4}$. (c) $m = 10^6$, $\Lambda = (1/9) \times 10^{-12}$, $\delta_b \approx 7.362 \times 10^{-3}$ and $\delta_{c2} \approx 1.469 \times 10^{-3}$. The black solid curve represents the effective Schwarzschild-(A)dS black-to-white hole case, while the red dashed curve denotes the classical Schwarzschild-(A)dS black hole case.

a discrepancy with the area gap Δ (For $\Lambda = -1$, $\Delta_{\text{symmetry}} \approx 0.17\Delta$). Secondly, we have adopted the method of finding δ_{c2} by fixing δ_b to look for symmetric solutions. Alternatively, one may adjust both parameters δ_b and δ_c to be subject to the symmetric condition. Then for a large cosmological constant, the discrepancy between the area of plaquette on the transition surface and the area gap Δ would become smaller.

Finally, we discuss the Kretschmann scalars of the quantum corrected Schwarzschild-(A)dS black-to-white holes, which are plotted for various quantum parameters in Fig. 10. It can be seen that the Kretschmann scalars are even function due to the symmetry of metric. They are finite in the entire interior region, and reach a single maximum at \mathcal{T} , indicating that the classical singularity is removed and matter is allowed to travel from the black hole to the white hole.

IV. CONCLUSIONS AND DISCUSSIONS

In this paper, we have investigated the dynamics of the black-to-white holes in the interior of the event horizon in the framework of LQG, where an effective theory with holonomy corrections is proposed at the semi-classical level. This work has made progress on the following topics. Firstly, we have explicitly shown that for any system with Hamiltonian constraint, the different choices of the lapse function lead to the same dynamics even at the semi-classical level where the canonical equations of motions are modified due to the quantum correction. Of course this result is not surprising since we know that the Hamiltonian constraint should lead to a re-parameterization invariance for the time coordinate. Secondly, inspired by this observation, we have looked into the dynamics of Schwarzschild black-to-white hole with different choices of the lapse function, and explicitly demonstrated that the black-to-white solution with the metric in Eq. (19), which is proposed based on the phenomenological consideration, can be viewed as the coarse-grained version of the black-to-white hole solution with the metric in Eq. (42), since both metrics exhibit the same behavior at the leading order expansion near the transition surface. In this sense, we have provided the theoretical foundation for the metric in Eq. (19) by embedding it into the framework of LQG and treating it as the approximation or the coarse-grained version of the solution to the equations of motion in the semi-classical theory. Thirdly, inspired by the exact symmetry appearing in the metric described by Eq. (19), we have constructed the black-to-white hole solutions with exact symmetry between the black hole region and the white hole region by properly fixing the quantum parameters in the effective theory of LQG, and thus the geometry of black hole and white hole is symmetric, needless to worry about the amplification of mass which might appear due to the mass difference between the black hole and the white hole. Remarkably, we have also demonstrated that with appropriate choice of lapse function (or, the time coordinate), the transition surface from the black hole to the white hole is exactly located at $\tau = 0$, which previously locates the presence of the singularity, as demonstrated in Fig. 1(b). In this sense, the picture of replacing the singularity with a bounce from the black hole to the white hole is more intuitive and vivid in comparison with the previous work where different lapse functions are employed. As a result, in this coordinate system the dynamics of the interior black-to-white hole is plotted with a picture analogous to that in loop quantum cosmology, where a cosmological singularity is replaced

by the big bounce at the same time moment. The analogy between the dynamics of the black hole interior and that of the universe has been revealed more evidently.

Subsequently, the effective theory has been applied to investigate the interior dynamics of a black hole in the presence of a cosmological constant. We have proposed an iteration method to fix the value of quantum parameters to guarantee that the area enclosed by two plaquettes on the transition surface remains to be the area gap. By numerically solving the equations of motion, we demonstrate that quantum gravity effects can eliminate the singularity in spacetime and extend time-like coordinates from the black hole region to the white hole region, thus providing the possibility for particles to reach other universes. More importantly, it is found that the geometry as well as the physical property of the black hole interior is different from that of the Schwarzschild interior. For Schwarzschild-AdS black holes, the radius of the event horizon of black hole is not the same as that of the white hole because the presence of a negative cosmological constant breaks this symmetry. More importantly, the location of both the transition surface and the event horizon goes as $m^{1/3}$ for larger m , indicating that spacetime bounces near event horizon. On the other hand, for Schwarzschild-dS black-to-white holes, the time evolution of p_c , p_b , c and b is similar to those of quantities in Schwarzschild black-to-white case (see Fig. 6), suggesting that both of them have similar dynamical properties. Finally, in order to fix the asymmetric problem of the Schwarzschild (A)dS metric, a scheme has been proposed to modify the quantum parameters δ_c and δ_b . It is shown when the quantum parameters satisfy the relation $\cos(\delta_b b(\tau_T)) = 0$, the symmetry between the black hole region and the white hole region may be restored. In addition, as the ratio of $m/\sqrt{\Lambda} > 10^8$, the symmetric solutions are subject to the condition that the area enclosed by two plaquettes on the transition surface remains to be the area gap.

In the end, we remark that we have only focused on the interior dynamics of the black hole in this paper. The effective spacetime structure outside the event horizon and how they match to the interior region should be studied in detail, and we leave these for further investigation.

ACKNOWLEDGMENTS

We are very grateful to Guoping Li, Kai Li, Pan Li, Yongge Ma, Jun Nian, Wenbin Pan, Yu Tian and Hongbao Zhang for helpful discussions. Specially we thank Xiangdong Zhang

and Cong Zhang for their correspondence with detailed comments and suggestions, which help us to improve the manuscript greatly. This work is supported in part by the Natural Science Foundation of China (Grant Nos. 12035016, 12275275, 12105231 and 12275350). It is also supported by Beijing Natural Science Foundation (Grant No. 1222031) and the Innovative Projects of Science and Technology (E2545BU210) at IHEP, and by Natural Science Foundation of Sichuan Province (Grant Nos. 2024NSFSC0456 and 2023NSFSC1348), the Sichuan Youth Science and Technology Innovation Research Team (Grant No. 21CXTD0038).

-
- [1] E. Bianchi, M. Christodoulou, F. D’Ambrosio, H. M. Haggard, and C. Rovelli, *Class. Quant. Grav.* **35**, 225003 (2018), [arXiv:1802.04264](#).
 - [2] F. D’Ambrosio and C. Rovelli, *Class. Quant. Grav.* **35**, 215010 (2018), [arXiv:1803.05015](#).
 - [3] A. Ashtekar, J. Olmedo, and P. Singh, *Phys. Rev. Lett.* **121**, 241301 (2018), [arXiv:1806.00648](#).
 - [4] A. Ashtekar, J. Olmedo, and P. Singh, *Phys. Rev. D* **98**, 126003 (2018), [arXiv:1806.02406](#).
 - [5] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. Lett.* **116**, 061102 (2016), [arXiv:1602.03837](#).
 - [6] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. D* **100**, 104036 (2019), [arXiv:1903.04467](#).
 - [7] D. Psaltis *et al.* (Event Horizon Telescope), *Phys. Rev. Lett.* **125**, 141104 (2020), [arXiv:2010.01055](#).
 - [8] R. Penrose, *Phys. Rev. Lett.* **14**, 57 (1965).
 - [9] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 1973).
 - [10] E. Witten, *Phys. Rev. D* **44**, 314 (1991).
 - [11] C. Rovelli, in *15th International Conference on General Relativity and Gravitation (GR15)* (1997) pp. 281–331, [arXiv:gr-qc/9803024](#).
 - [12] M. Bojowald, R. Goswami, R. Maartens, and P. Singh, *Phys. Rev. Lett.* **95**, 091302 (2005), [arXiv:gr-qc/0503041](#).
 - [13] A. Saini and D. Stojkovic, *Phys. Rev. D* **89**, 044003 (2014), [arXiv:1401.6182](#).

- [14] A. Ashtekar and E. Bianchi, *Rept. Prog. Phys.* **84**, 042001 (2021), [arXiv:2104.04394](#).
- [15] J. Bardeen, in *Proceedings of the 5th International Conference on Gravitation and the Theory of Relativity* (Tbilisi, 1968).
- [16] S. A. Hayward, *Phys. Rev. Lett.* **96**, 031103 (2006), [arXiv:gr-qc/0506126](#).
- [17] L. Xiang, Y. Ling, and Y. G. Shen, *Int. J. Mod. Phys. D* **22**, 1342016 (2013), [arXiv:1305.3851](#).
- [18] L. Balart and E. C. Vagenas, *Phys. Lett. B* **730**, 14 (2014), [arXiv:1401.2136](#).
- [19] Z.-Y. Fan and X. Wang, *Phys. Rev. D* **94**, 124027 (2016), [arXiv:1610.02636](#).
- [20] X. Li, Y. Ling, Y.-G. Shen, C.-Z. Liu, H.-S. He, and L.-F. Xu, *Annals Phys.* **396**, 334 (2018), [arXiv:1611.09016](#).
- [21] S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **96**, 104008 (2017), [arXiv:1708.05226](#).
- [22] Y. He and M.-S. Ma, *Phys. Lett. B* **774**, 229 (2017), [arXiv:1709.09473](#).
- [23] M. E. Rodrigues and M. V. de Sousa Silva, *JCAP* **06**, 025 (2018), [arXiv:1802.05095](#).
- [24] M. V. de Sousa Silva and M. E. Rodrigues, *Eur. Phys. J. C* **78**, 638 (2018), [arXiv:1808.05861](#).
- [25] A. Simpson and M. Visser, *JCAP* **02**, 042 (2019), [arXiv:1812.07114](#).
- [26] C. Zhang and X. Zhang, *Phys. Rev. D* **101**, 086002 (2020), [arXiv:1912.07278](#).
- [27] G. Melgarejo, E. Contreras, and P. Bargueño, *Phys. Dark Univ.* **30**, 100709 (2020).
- [28] N. Burzillà, B. L. Giacchini, T. d. P. Netto, and L. Modesto, *Eur. Phys. J. C* **81**, 462 (2021), [arXiv:2012.11829](#).
- [29] A. Bonanno, A.-P. Khosravi, and F. Saueressig, *Phys. Rev. D* **103**, 124027 (2021), [arXiv:2010.04226](#).
- [30] M. Estrada and F. Tello-Ortiz, *EPL* **135**, 20001 (2021), [arXiv:2012.05068](#).
- [31] R. Torres, (2022), [arXiv:2208.12713](#).
- [32] C. Lan, H. Yang, Y. Guo, and Y.-G. Miao, *Int. J. Theor. Phys.* **62**, 202 (2023), [arXiv:2303.11696](#).
- [33] J. T. S. S. Junior, F. S. N. Lobo, and M. E. Rodrigues, *Class. Quant. Grav.* **41**, 055012 (2024), [arXiv:2310.19508](#).
- [34] J. Mazza and S. Liberati, *JHEP* **03**, 199 (2023), [arXiv:2301.04697](#).
- [35] O. Luongo and H. Quevedo, *Class. Quant. Grav.* **41**, 125011 (2024), [arXiv:2305.11185](#).
- [36] C. Rovelli and P. Martin-Dussaud, *Class. Quant. Grav.* **35**, 147002 (2018), [arXiv:1803.06330](#).

- [37] S. Brahma and D.-h. Yeom, *Class. Quant. Grav.* **35**, 205007 (2018), [arXiv:1804.02821](#).
- [38] R. Carballo-Rubio, F. Di Filippo, S. Liberati, and M. Visser, *Phys. Rev. D* **98**, 124009 (2018), [arXiv:1809.08238](#).
- [39] N. Bodendorfer, F. M. Mele, and J. Münch, *Class. Quant. Grav.* **38**, 095002 (2021), [arXiv:1912.00774](#).
- [40] E. Alesci, S. Bahrami, and D. Pranzetti, *Phys. Lett. B* **797**, 134908 (2019), [arXiv:1904.12412](#).
- [41] J. Ben Achour, S. Brahma, S. Mukohyama, and J. P. Uzan, *JCAP* **09**, 020 (2020), [arXiv:2004.12977](#).
- [42] Y. Ling and M.-H. Wu, *Class. Quant. Grav.* **40**, 075009 (2023), [arXiv:2109.05974](#).
- [43] Y. Ling and M.-H. Wu, *Chin. Phys. C* **46**, 025102 (2022), [arXiv:2109.12938](#).
- [44] A. Rignon-Bret and C. Rovelli, *Phys. Rev. D* **105**, 086003 (2022), [arXiv:2108.12823](#).
- [45] Y. Ling and M.-H. Wu, *Symmetry* **14**, 2415 (2022), [arXiv:2205.08919](#).
- [46] W. Zeng, Y. Ling, and Q.-Q. Jiang, *Chin. Phys. C* **47**, 085103 (2023), [arXiv:2207.07529](#).
- [47] W. Zeng, Y. Ling, Q.-Q. Jiang, and G.-P. Li, *Phys. Rev. D* **108**, 104072 (2023), [arXiv:2308.00976](#).
- [48] Z. Feng, Y. Ling, X. Wu, and Q. Jiang, *Sci. China Phys. Mech. Astron.* **67**, 270412 (2024), [arXiv:2308.15689](#).
- [49] E. Ayon-Beato and A. Garcia, *Phys. Rev. Lett.* **80**, 5056 (1998), [arXiv:gr-qc/9911046](#).
- [50] M. Dehghani, *Phys. Rev. D* **99**, 104036 (2019).
- [51] E. Ayon-Beato and A. Garcia, *Phys. Lett. B* **464**, 25 (1999), [arXiv:hep-th/9911174](#).
- [52] K. A. Bronnikov, *Phys. Rev. D* **63**, 044005 (2001), [arXiv:gr-qc/0006014](#).
- [53] I. Dymnikova, *Class. Quant. Grav.* **21**, 4417 (2004), [arXiv:gr-qc/0407072](#).
- [54] L. Balart and E. C. Vagenas, *Phys. Rev. D* **90**, 124045 (2014), [arXiv:1408.0306](#).
- [55] A. Allahyari, M. Khodadi, S. Vagnozzi, and D. F. Mota, *JCAP* **02**, 003 (2020), [arXiv:1912.08231](#).
- [56] M. Guerrero and D. Rubiera-Garcia, *Phys. Rev. D* **102**, 024005 (2020), [arXiv:2005.08828](#).
- [57] S. I. Kruglov, *Annals Phys.* **428**, 168449 (2021), [arXiv:2104.08099](#).
- [58] D. Kokoška and M. Ortaggio, *Phys. Rev. D* **104**, 124051 (2021), [arXiv:2109.08886](#).
- [59] A. Ali and K. Saifullah, *Eur. Phys. J. C* **82**, 131 (2022).

- [60] S. Capozziello and G. G. L. Nashed, *Class. Quant. Grav.* **40**, 205023 (2023), [arXiv:2309.08894](#).
- [61] J. W. Moffat, *Eur. Phys. J. C* **75**, 175 (2015), [arXiv:1412.5424](#).
- [62] L. Modesto and L. Rachwal, (2016), [arXiv:1605.04173](#).
- [63] A. Simpson, P. Martin-Moruno, and M. Visser, *Class. Quant. Grav.* **36**, 145007 (2019), [arXiv:1902.04232](#).
- [64] R. Carballo-Rubio, F. Di Filippo, S. Liberati, and M. Visser, *Phys. Rev. D* **101**, 084047 (2020), [arXiv:1911.11200](#).
- [65] R. Carballo-Rubio, F. Di Filippo, S. Liberati, and M. Visser, *JHEP* **02**, 122 (2022), [arXiv:2111.03113](#).
- [66] A. Bakopoulos, C. Charmousis, and P. Kanti, *JCAP* **05**, 022 (2022), [arXiv:2111.09857](#).
- [67] D.-f. Zeng, *Nucl. Phys. B* **977**, 115722 (2022), [arXiv:2112.12531](#).
- [68] Z.-C. Li and H. Lu, *Eur. Phys. J. C* **83**, 755 (2023), [arXiv:2303.16924](#).
- [69] J. Nian, (2023), [arXiv:2312.14287](#).
- [70] W.-C. Lin, D.-h. Yeom, and D. Stojkovic, (2023), [arXiv:2310.15508](#).
- [71] H. Nicolai, K. Peeters, and M. Zamaklar, *Class. Quant. Grav.* **22**, R193 (2005), [arXiv:hep-th/0501114](#).
- [72] M. Bojowald and R. Swiderski, *Class. Quant. Grav.* **23**, 2129 (2006), [arXiv:gr-qc/0511108](#).
- [73] D. Cartin and G. Khanna, *Phys. Rev. D* **73**, 104009 (2006), [arXiv:gr-qc/0602025](#).
- [74] A. Ashtekar and M. Bojowald, *Class. Quant. Grav.* **23**, 391 (2006), [arXiv:gr-qc/0509075](#).
- [75] L. Modesto, *Class. Quant. Grav.* **23**, 5587 (2006), [arXiv:gr-qc/0509078](#).
- [76] C. G. Boehmer and K. Vandersloot, *Phys. Rev. D* **76**, 104030 (2007), [arXiv:0709.2129](#).
- [77] D.-W. Chiou, *Phys. Rev. D* **78**, 064040 (2008), [arXiv:0807.0665](#).
- [78] J. Brannlund, S. Kloster, and A. DeBenedictis, *Phys. Rev. D* **79**, 084023 (2009), [arXiv:0901.0010](#).
- [79] M. Reuter and F. Saueressig, *New J. Phys.* **14**, 055022 (2012), [arXiv:1202.2274](#).
- [80] N. Dadhich, A. Joe, and P. Singh, *Class. Quant. Grav.* **32**, 185006 (2015), [arXiv:1505.05727](#).
- [81] A. Corichi and P. Singh, *Class. Quant. Grav.* **33**, 055006 (2016), [arXiv:1506.08015](#).
- [82] J. Olmedo, S. Saini, and P. Singh, *Class. Quant. Grav.* **34**, 225011 (2017), [arXiv:1707.07333](#).
- [83] J. Cortez, W. Cuervo, H. A. Morales-Técotl, and J. C. Ruelas, *Phys. Rev. D* **95**, 064041 (2017), [arXiv:1704.03362](#).

- [84] A. Perez, *Rept. Prog. Phys.* **80**, 126901 (2017), [arXiv:1703.09149](#).
- [85] A. Yonika, G. Khanna, and P. Singh, *Class. Quant. Grav.* **35**, 045007 (2018), [arXiv:1709.06331](#).
- [86] A. Joe and P. Singh, *Class. Quant. Grav.* **32**, 015009 (2015), [arXiv:1407.2428](#).
- [87] P. Martin-Dussaud and C. Rovelli, *Class. Quant. Grav.* **36**, 245002 (2019), [arXiv:1905.07251](#).
- [88] J. G. Kelly, R. Santacruz, and E. Wilson-Ewing, *Phys. Rev. D* **102**, 106024 (2020), [arXiv:2006.09302](#).
- [89] C. Zhang, Y. Ma, S. Song, and X. Zhang, *Phys. Rev. D* **102**, 041502 (2020), [arXiv:2006.08313](#).
- [90] C. Zhang, Y. Ma, S. Song, and X. Zhang, *Phys. Rev. D* **105**, 024069 (2022), [arXiv:2107.10579](#).
- [91] X. Zhang, *Universe* **9**, 313 (2023), [arXiv:2308.10184](#).
- [92] M. Geiller, E. R. Livine, and F. Sartini, *SciPost Phys.* **10**, 022 (2021), [arXiv:2010.07059](#).
- [93] R. G. Daghigh, M. D. Green, and G. Kunstatter, *Phys. Rev. D* **103**, 084031 (2021), [arXiv:2012.13359](#).
- [94] A. Ashtekar, *Universe* **6**, 21 (2020), [arXiv:2001.08833](#).
- [95] D. K. Hong, W.-C. Lin, and D.-h. Yeom, *Phys. Rev. D* **106**, 104011 (2022), [arXiv:2207.03183](#).
- [96] B. Elizaga Navascués, A. García-Quismondo, and G. A. Mena Marugán, *Phys. Rev. D* **106**, 063516 (2022), [arXiv:2207.04677](#).
- [97] J. Lewandowski, Y. Ma, J. Yang, and C. Zhang, *Phys. Rev. Lett.* **130**, 101501 (2023), [arXiv:2210.02253](#).
- [98] M. Han, C. Rovelli, and F. Soltani, *Phys. Rev. D* **107**, 064011 (2023), [arXiv:2302.03872](#).
- [99] C. Zhang, Y. Ma, and J. Yang, *Phys. Rev. D* **108**, 104004 (2023), [arXiv:2302.02800](#).
- [100] G. Fu, D. Zhang, P. Liu, X.-M. Kuang, and J.-P. Wu, *Phys. Rev. D* **109**, 026010 (2024), [arXiv:2301.08421](#).
- [101] M. Bojowald, *Phys. Rev. Lett.* **86**, 5227 (2001), [arXiv:gr-qc/0102069](#).
- [102] A. Ashtekar, M. Bojowald, and J. Lewandowski, *Adv. Theor. Math. Phys.* **7**, 233 (2003), [arXiv:gr-qc/0304074](#).
- [103] A. Ashtekar and J. Lewandowski, *Class. Quant. Grav.* **21**, R53 (2004), [arXiv:gr-qc/0404018](#).

- [104] T. Thiemann, [Lect. Notes Phys. **721**, 185 \(2007\)](#), [arXiv:hep-th/0608210](#).
- [105] T. Thiemann, (2001), [arXiv:gr-qc/0110034](#).
- [106] T. Thiemann, [Lect. Notes Phys. **631**, 41 \(2003\)](#), [arXiv:gr-qc/0210094](#).
- [107] A. Ashtekar, [Phys. Rev. D **36**, 1587 \(1987\)](#).
- [108] G. Ongole, P. Singh, and A. Wang, [Phys. Rev. D **109**, 026015 \(2024\)](#), [arXiv:2311.10166](#).
- [109] The discrepancy between the numerical results based on Eq. (42) and the above analytical results comes from the fact that many factors have been discarded in Eq. (42). Their contribution to the curvature may not small, but mass independent indeed.
- [110] C. Rovelli and F. Vidotto, [Int. J. Mod. Phys. D **23**, 1442026 \(2014\)](#), [arXiv:1401.6562](#).
- [111] A. Barrau and C. Rovelli, [Phys. Lett. B **739**, 405 \(2014\)](#), [arXiv:1404.5821](#).
- [112] T. De Lorenzo, C. Pacilio, C. Rovelli, and S. Speziale, [Gen. Rel. Grav. **47**, 41 \(2015\)](#), [arXiv:1412.6015](#).
- [113] C. Rovelli, [Nature Astron. **1**, 0065 \(2017\)](#), [1708.01789](#).
- [114] J. Lin and X. Zhang, [Phys. Rev. D **110**, 026002 \(2024\)](#), [arXiv:2402.13638](#).
- [115] H. Nariai, [Gen. Rel. Grav. **31**, 963 \(1951\)](#).