

3D HARD SPHERE BOLTZMANN EQUATION: EXPLICIT STRUCTURE AND THE TRANSITION PROCESS FROM POLYNOMIAL TAIL TO GAUSSIAN TAIL

YU-CHU LIN, HAITAO WANG, AND KUNG-CHIEN WU

ABSTRACT. We study the Boltzmann equation with hard sphere in a near-equilibrium setting. The initial data is compactly supported in the space variable and has a polynomial tail in the microscopic velocity. We show that the solution can be decomposed into a particle-like part (polynomial tail) and a fluid-like part (Gaussian tail). The particle-like part decays exponentially in both space and time, while the fluid-like part corresponds to the behavior of the compressible Navier-Stokes equation, which dominates the long time behavior and exhibits rich wave motion. The nonlinear wave interactions in the fluid-like part are precisely characterized and therefore we are able to distinguish the linear and nonlinear wave of the solution. It is notable that although the solution has polynomial tail in the velocity initially, the transition process from the polynomial to the Gaussian tail can be quantitatively revealed due to the collision with the background global Maxwellian.

CONTENTS

| | | |
|----|--|----|
| 1. | Introduction | 1 |
| 2. | Preliminaries | 8 |
| 3. | Global wave structure | 13 |
| 4. | Exponential decay outside the wave cone | 18 |
| 5. | The transition from polynomial tail to Gaussian tail | 21 |
| 6. | Some convolution estimates | 26 |
| | References | 59 |

1. INTRODUCTION

1.1. The model. The Boltzmann equation is a fundamental model in the collisional kinetic theory, which describes the evolution of a phase space distribution function of moderately dilute gas. Precisely, the Boltzmann equation reads

$$(1) \quad \begin{cases} \partial_t F + \xi \cdot \nabla_x F = Q(F, F), \\ F(0, x, \xi) = F_0(x, \xi), \end{cases} \quad (t, x, \xi) \in \mathbb{R}^+ \times \mathbb{R}^3 \times \mathbb{R}^3,$$

where $F(t, x, \xi)$ is the distribution function for particles at time $t > 0$, position $x = (x_1, x_2, x_3) \in \mathbb{R}^3$ and microscopic velocity $\xi = (\xi_1, \xi_2, \xi_3) \in \mathbb{R}^3$, and initial data $F_0(x, \xi) \geq 0$ is given. Here the Boltzmann collision operator $Q(\cdot, \cdot)$ is given by

$$Q(G, F) = \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |q \cdot n| [G(\xi'_*) F(\xi') - G(\xi_*) F(\xi)] dnd\xi_*$$

where $q = \xi - \xi_*$ is the relative velocity and the post-collisional velocities satisfy

$$\xi' = \xi - [(\xi - \xi_*) \cdot n] n, \quad \xi'_* = \xi_* + [(\xi - \xi_*) \cdot n] n.$$

Date: August 6, 2024.

Key words and phrases. Boltzmann equation, Maxwellian, polynomial tail.

2020 Mathematics Subject Classification: 35Q20; 82C40.

It is well known that the global Maxwellians are the steady solutions to the Boltzmann equation. In the perturbation regime near the Maxwellian, we look for the solution in the form of

$$(2) \quad F = \mathcal{M} + f, \quad F_0(x, \xi) = \mathcal{M} + \varepsilon f_0,$$

where $\varepsilon > 0$ sufficiently small, with a perturbation function f to \mathcal{M} . Here the global Maxwellian \mathcal{M} is normalized as

$$\mathcal{M} = \frac{1}{(2\pi)^{3/2}} \exp\left(-\frac{|\xi|^2}{2}\right).$$

Substituting (2) into (1), the perturbation function f satisfies the equation

$$(3) \quad \begin{cases} \partial_t f + \xi \cdot \nabla_x f = \mathcal{L}f + Q(f, f), \\ f|_{t=0} = F_0(x, \xi) - \mathcal{M} = \varepsilon f_0, \quad (x, \xi) \in \mathbb{R}^3 \times \mathbb{R}^3, \end{cases}$$

where

$$\mathcal{L}f = Q(\mathcal{M}, f) + Q(f, \mathcal{M}).$$

In fact, \mathcal{L} can be split into

$$\mathcal{L}f = -\nu + \mathcal{K},$$

with

$$\begin{aligned} \nu(\xi) &= \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |q \cdot \omega| \mathcal{M}(\xi_*) d\omega d\xi_* \sim (1 + |\xi|), \\ \mathcal{K}f &= \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |q \cdot n| [\mathcal{M}(\xi'_*) f(\xi') + f(\xi'_*) \mathcal{M}(\xi') - f(\xi_*) \mathcal{M}(\xi)] dnd\xi_*. \end{aligned}$$

Note that this kind of perturbation (2) allows the initial data to have a polynomial tail in the microscopic velocity, which is different from the standard perturbation, $F = \mathcal{M} + \sqrt{\mathcal{M}}f$, where initial data is assumed to have a Gaussian tail.

It is known that there are extensive studies on the standard perturbation, including the global existence, time-asymptotic behavior, and even the pointwise structure, see [7, 16, 27, 28] and the references therein. It would be very interesting to see that can we still obtain the precise space-time structure of the solution for initial data with a polynomial tail?

Moreover, since the perturbation setting describes the collisions between a small amount of released particles and the ambient particles that have reached thermal equilibrium, the physical intuition suggests that the distribution of the released particles will also approach thermal equilibrium over a long period, namely, it will become close to a Gaussian in terms of the microscopic velocity. It is a challenge problem that is it possible to give a quantitative description of the transition process?

The main goal of this paper is to answer the above two questions. Specifically, we will construct a pointwise estimate of solution to (3) with respect to all variables, space, time, and velocity. The estimate not only exhibits the wave motion in space-time, but also reveals how the solution transitions from a polynomial tail to a Gaussian tail.

1.2. Review of previous works. Concerning the polynomial tail perturbation for collisional kinetic equation, there has been substantial progress recently. For the torus case, it was initiated by Gualdani-Mischler-Mouhot [17] for the Boltzmann equation with hard sphere. It was then generalized by [8] for Landau equation, by [1, 18] for the Boltzmann equation without angular cutoff, and by [5] for the Boltzmann equation with soft potentials. For the whole space case, we refer to [9, 6] for the non-cutoff Boltzmann equation and [13] for the cutoff Boltzmann equation. These works mainly focused on global existence and large time decay of the solution, whereas our study aim at providing a more quantitative description of the structure of the solution.

Next, we review some space-time pointwise results closely related to the current study. It was initiated by Liu [22] for 1D viscous conservation laws, then developed to multi-dimensional compressible Navier-Stokes equation [10, 12, 23, 24]. There are two key ingredients. The first is the construction of Green's function for the linearized system, where rich wave phenomena, such as dissipative Huygens wave, diffusion wave, Riesz wave are identified. The other one is the careful estimate of nonlinear wave couplings between the above basic wave patterns. As is known,

the long time behavior of the Boltzmann equation is governed by macroscopic fluid. There are some parallel results for Boltzmann equation with standard Gaussian tail. The result for 1D hard sphere Boltzmann equation was constructed by Liu-Yu [25]. As the nonlinear interaction is strong in 1D, the authors need to extract the so called “kinetic Burger equations” to close the nonlinear problem. Later, [26, 27] obtained the explicit structure of Green’s function for the linearized Boltzmann equation with hard sphere in 3D. Recently, [20] constructed the explicit structure of the relativistic Boltzmann equation for “hard ball”, an exponentially sharp ansatz similar to structure in [10] was justified. In these works, the nonlinear interactions have been estimated to the extent necessary to close the nonlinear ansatz.

The transition from polynomial tail to Gaussian tail is related to the decay estimates for large velocities in the Boltzmann equation. For space homogeneous case, there are extensive studies on L_v^1 moments or pointwise decay, both for polynomial and exponential weight, see [2, 3, 11, 15] and references therein. For space inhomogeneous case, the results are relatively fewer. Generation of polynomial moments in L_v^1 or pointwise sense was established under suitable moment bound conditions for hard potential with or without cutoff, see [4, 17, 19]. Different from the conditional results, our result provides a dynamic process for Gaussian tail generation in the perturbation regime.

1.3. Notations. Before stating our main results, we introduce some notations used in this paper. We denote $w_\beta(\xi) = \langle \xi \rangle^\beta = (1 + |\xi|^2)^{\beta/2}$, $\beta \in \mathbb{R}$. For the microscopic variable ξ , we denote

$$|g|_{L_\xi^p} = \left(\int_{\mathbb{R}^3} |g|^p d\xi \right)^{1/p} \text{ if } 1 \leq p < \infty, \quad |g|_{L_\xi^\infty} = \sup_{\xi \in \mathbb{R}^3} |g(\xi)|,$$

and the weighted norms can be defined by

$$|g|_{L_{\xi,\beta}^p} = \left(\int_{\mathbb{R}^3} |\langle \xi \rangle^\beta |g|^p d\xi \right)^{1/p} \text{ if } 1 \leq p < \infty, \quad |g|_{L_{\xi,\beta}^\infty} = \sup_{\xi \in \mathbb{R}^3} |\langle \xi \rangle^\beta g(\xi)|.$$

The L_ξ^2 inner product in \mathbb{R}^3 will be denoted by $\langle \cdot, \cdot \rangle_\xi$, i.e.,

$$\langle f, g \rangle_\xi = \int f(\xi) \overline{g(\xi)} d\xi.$$

For the Boltzmann equation with hard sphere, the natural norm in ξ is $|\cdot|_{L_\xi^2}$, which is defined as

$$|g|_{L_\xi^2} = \left| \langle \xi \rangle^{\frac{1}{2}} g \right|_{L_\xi^2}.$$

For the space variable x , we have similar notations, namely,

$$|g|_{L_x^p} = \left(\int_{\mathbb{R}^3} |g|^p dx \right)^{1/p} \text{ if } 1 \leq p < \infty, \quad |g|_{L_x^\infty} = \sup_{x \in \mathbb{R}^3} |g(x)|.$$

Finally, with \mathcal{X} and \mathcal{Y} being two normed spaces, we define

$$\|g\|_{\mathcal{X}\mathcal{Y}} = \|g|_{\mathcal{Y}}\|_{\mathcal{X}},$$

and for simplicity, we denote

$$\|g\|_{L^2} = \|g\|_{L_\xi^2 L_x^2} = \left(\int_{\mathbb{R}^3} |g|_{L_x^2}^2 d\xi \right)^{1/2}.$$

For any two functions $f(x, t)$ and $g(x, t)$, we define the space-time convolution as

$$f(x, t) *_{x,t} g(x, t) = \int_0^t \int_{\mathbb{R}^3} f(x-y, t-\tau) g(y, \tau) dy d\tau.$$

For any two real numbers a and b , we define $a \wedge b = \min\{a, b\}$.

For simplicity of notations, hereafter, we abbreviate $\leq C$ to \lesssim , where C is a constant depending only on fixed numbers.

1.4. Main theorem and significant points of our results. In order to achieve our goal, we introduce the decomposition $f = f_1 + \sqrt{\mathcal{M}}f_2$. Here f_1 corresponds to the part with only polynomial tail, while $\sqrt{\mathcal{M}}f_2$ has Gaussian tail. Heuristically, the closer a distribution function to a Gaussian, the closer the behavior resembles macroscopic hydrodynamics. Therefore, one may expect f_1 behaves like particle, and f_2 behaves like fluid.

The following coupled system designed for f_1 and f_2 is to realize one's intuition:

$$(4) \quad \begin{cases} \partial_t f_1 + \xi \cdot \nabla_x f_1 = -\nu(\xi) f_1 + \mathcal{K}_s f_1 + Q(f_1, f_1) + Q(f_1, \sqrt{\mathcal{M}}f_2) + Q(\sqrt{\mathcal{M}}f_2, f_1), \\ \partial_t f_2 + \xi \cdot \nabla_x f_2 = Lf_2 + \mathcal{K}_b f_1 + \Gamma(f_2, f_2), \end{cases}$$

with initial data

$$f_1(0, x, \xi) = f_{1,0}(x, \xi) = \varepsilon f_0(x, \xi), \quad f_2(0, x, \xi) = f_{2,0}(x, \xi) = 0,$$

where

$$Lg = \frac{1}{\sqrt{\mathcal{M}}} \left[Q(\mathcal{M}, \sqrt{\mathcal{M}}g) + Q(\sqrt{\mathcal{M}}g, \mathcal{M}) \right]$$

is the so called linearized Boltzmann collision operator,

$$\Gamma(f_2, f_2) = \frac{1}{\sqrt{\mathcal{M}}} Q(\sqrt{\mathcal{M}}f_2, \sqrt{\mathcal{M}}f_2)$$

is the nonlinear Boltzmann collision operator, and $\mathcal{K} = \mathcal{K}_s + \mathcal{M}^{1/2}\mathcal{K}_b$ with

$$\mathcal{K}_s := \chi_{\{|\xi| \geq R\}} \mathcal{K}, \quad \mathcal{K}_b := \mathcal{M}^{-1/2} \chi_{\{|\xi| < R\}} \mathcal{K}$$

for a constant $R > 0$ large enough, $\chi_{\{\cdot\}}$ being the indicator function.

It is noted that similar decomposition was also employed in [6] for Boltzmann equation without angular cutoff. Based on this decomposition, we have the main theorem as follows:

Theorem 1. *Let $\beta > 4$ be sufficiently large. Assume that $f_0 \in L_{\xi, \beta}^\infty L_x^\infty$ is compactly supported in the variable x for all ξ*

$$f_0(x, \xi) \equiv 0 \text{ for } |x| \geq 1, \xi \in \mathbb{R}^3.$$

Then for any fixed $\delta > 0$, there exists $\varepsilon > 0$ small enough such that the solution f of (3) exists for $t > 0$ and it can be decomposed as $f = f_1 + \sqrt{\mathcal{M}}f_2$, where f_1 and f_2 satisfy (4) with the following estimates:

For f_1 , we have

$$|\mathbf{1}_{\{\langle \xi \rangle \leq t\}} f_1| \lesssim \varepsilon \langle \xi \rangle^{-\beta} e^{-\bar{c}_0 [(\xi)^2 + (t+|x|)]}$$

and

$$|\mathbf{1}_{\{\langle \xi \rangle > t\}} f_1| \lesssim \varepsilon \langle \xi \rangle^{-\beta} e^{-\bar{c}_0 [(\xi)t + (t+|x|)]}$$

for some positive constant \bar{c}_0 .

For f_2 , we have

$$|f_2|_{L_{\xi, \beta}^\infty} \lesssim \varepsilon \left[\mathbf{1}_{\{|x| \leq ct\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D(1+t)}} \right] \\ + \varepsilon^2 \mathbf{1}_{\{|x| \leq (c+\delta)t\}} \left[(1+t)^{-2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} + (1+t)^{-\frac{5}{2}} \left(1 + \frac{(|x| - ct)^2}{1+t} \right)^{-1} \right]$$

for some positive constants $\hat{D}, \hat{D}_0, \hat{c}_\delta$. Here $\mathbf{c} = \sqrt{5/3}$ is the sound speed associated with global Maxwellian \mathcal{M} .

Several remarks on the main theorem are in order:

- The main result is the combinations of Theorems 9, 10, and 15. In Theorem 9, we obtain the global wave structure, which is accurate in the time-like region but only shows polynomial decay for f_2 in the space-like region. In Theorem 10, the estimate in space-like region is further improved to be exponentially sharp. Theorem 15 describes the dynamic process of transition from polynomial tail to Gaussian tail for f_1 .
- The result shows that the polynomial tail part f_1 decays exponentially in both space and time, while Gaussian tail part $\sqrt{\mathcal{M}}f_2$ exhibits rich wave phenomena and dominates the solution at large time. This is consistent with our intuition: the polynomial and Gaussian tail parts are associated with particle-like and fluid-like behaviors, respectively.
- In the estimate of f_2 , we see that the terms consist of ε and ε^2 orders. The ε order terms represent linear waves, such as Huygens, diffusion, and Riesz waves, as given by the Green's function. The ε^2 order terms arise from nonlinear interactions between these basic waves. They consist of polynomial versions of Huygens and diffusion waves, primarily concentrate inside the acoustic wave cone. Compared to the linear waves, the nonlinear waves not only have a ε^2 order magnitude, but also decay faster by $(1+t)^{-1/2}$ than their linear counterparts. However, previous works [10, 12, 20, 23, 24] only showed that they have the same decay rates. Thus, our results provide a more accurate description for the nonlinear effect, based on sharper estimates of nonlinear wave couplings.
- The polynomial tail of the solution is fully captured by f_1 part. The pointwise estimate of f_1 in velocity variable thus shows how the polynomial tail transitions to a Gaussian tail. Specifically, f_1 only exhibits a polynomial tail initially, as time evolves it immediately acquires an exponential tail. As time continues to evolve, the particles with velocity $\langle \xi \rangle < t$ will become Gaussian tail, while the non-Gaussian part is of the order e^{-t^2} and any moments generated by the non-Gaussian part are bounded by e^{-t^2} . Therefore, as time tends to infinity, the distribution function will eventually transit to a Gaussian tail; but however, the transition cannot be completed in any finite time.
- The assumption that the initial data has compact support in space is unessential. It is not hard to generalize to the case where the initial data decays polynomially in space, but in this case, the space-like behavior should be modified accordingly. We do not pursue this, as our focus is on the quantitative description of wave motion and the transition to a Gaussian tail.

1.5. Ideas and strategies. We now outline the ideas and strategies for the proof of the main theorem.

1.5.1. Global wave structure. Let us begin with Theorem 9, the global wave structure of the solution. Consider the coupled system (4). Let \mathbb{S}^t and \mathbb{G}^t be the solution operators to the damped transport equation and (standard) linearized Boltzmann equation respectively, namely, $g(t) \equiv \mathbb{S}^t g_0$ solves

$$\partial_t g + \xi \cdot \nabla_x g + \nu(\xi)g = 0, \quad g(0, x, \xi) = g_0,$$

and $g(t) \equiv \mathbb{G}^t g_0$ solves

$$\partial_t g + \xi \cdot \nabla_x g - Lg = 0, \quad g(0, x, \xi) = g_0.$$

Then by Duhamel principle, solutions f_1 and f_2 to (4) satisfy the following coupled integral system (5)

$$\begin{cases} f_1(t) = \varepsilon \mathbb{S}^t f_0 + \int_0^t \mathbb{S}^{t-\tau} \mathcal{K}_s f_1(\tau) d\tau + \int_0^t \mathbb{S}^{t-\tau} \left[Q(f_1, f_1) + Q(f_1, \sqrt{\mathcal{M}}f_2) + Q(\sqrt{\mathcal{M}}f_2, f_1) \right](\tau) d\tau, \\ f_2(t) = \int_0^t \mathbb{G}^{t-\tau} \mathcal{K}_b f_1(\tau) d\tau + \int_0^t \mathbb{G}^{t-\tau} \Gamma(f_2, f_2)(\tau) d\tau. \end{cases}$$

The essential step for constructing global wave structure is to find out the accurate ansatz for the solution.

We neglect the nonlinear effects for the time being. In the equation for f_1 , if R is chosen sufficiently large, the integral operator \mathcal{K}_s can be regarded as a perturbation of the damped transport operator. This results in an exponential decay of f_1 in both space and time. We

then substitute the estimate of f_1 into the integral equation for f_2 , requiring consideration of $\int_0^t \mathbb{G}^{t-\tau} \mathcal{K}_b f_1(\tau) d\tau$. The explicit structure of Green's function \mathbb{G} is constructed in [26, 27] (we stated it in Lemma 8), showing rich wave structures, including a dissipative Huygens wave (acoustic wave described by a moving heat kernel), a diffusion wave (thermal wave described by a stationary heat kernel), a Riesz wave (related to vorticity of macroscopic fluid, described by a polynomial analogue of diffusion wave confined in the wave cone), and a space-time exponential decay term. The estimate of f_2 is given by the convolution of the Green's function and the source term $\mathcal{K}_b f_1$, inheriting a structure similar to that of the Green's function (see Lemmas 19-21). This indicates that f_1 can be viewed as particle-like wave, while f_2 can be viewed as fluid-like wave.

Next, we incorporate nonlinear effects. In designing the coupled system, we intentionally placed all nonlinear terms involving f_1 in the first equation of (4), as it describes the particle-like behavior. The term $\Gamma(f_2, f_2)$ is included in the second equation, as it is associated with the fluid-like behavior. The key term is $\int_0^t \mathbb{G}^{t-\tau} \Gamma(f_2, f_2)(\tau) d\tau$, which accounts for the nonlinear wave interactions. A fundamental property is that the nonlinear operator Γ is purely microscopic, and when acted upon by the Green's operator, it gains an extra $\frac{1}{2}$ -order time decay. Substituting the linear estimates leads to the convolution estimates:

$$\frac{(\text{Diffusion+Huygens+Riesz+Exponential decay})}{\sqrt{1+t}} \underset{(x,t)}{*} (\text{Diffusion+Huygens+Riesz+Exponential decay})^2.$$

The main effort is to provide sharp estimates of these convolutions. Without delving into the details, we provide a heuristic explanation of the interaction process here. We illustrate this with the convolution of two Huygens waves.

$$\int_0^t \int_{\mathbb{R}^3} (1+t-s)^{-5/2} e^{-\frac{(|x-y|-(t-s))^2}{D_0(t-s)}} (1+s)^{-4} e^{-\frac{(|y|-s)^2}{C_1(s+1)}} dy ds,$$

where $c = 1$ for simplicity. The following figure is to explain the interaction process:




FIGURE 1. Interaction between two Huygens waves

Here, the source $(1+s)^{-4} e^{-\frac{(|y|-s)^2}{C_1(s+1)}}$ is plotted as a forward cone with thickness \sqrt{s} , and the propagator $(1+t-s)^{-5/2} e^{-\frac{(|x-y|-(t-s))^2}{D_0(t-s)}}$ is plotted as a backward cone with thickness $\sqrt{t-s}$. The space is represented in 2D in the figure. The interaction essentially occurs in the following space-time region:

$$\{(y, s) \mid ||x-y| - (t-s)| \leq O(1)\sqrt{t-s} \text{ and } ||y|-s| \leq O(1)\sqrt{s}\}.$$

Inside this region, the exponential term in Huygens wave is not effective, and the decay is mainly due to time decay factor. The key point is the sharp estimate of the volume for this space-time

region. In Section 6.2, we first provide some heuristic calculations. In fact, our rigorous estimates are greatly motivated by the heuristics. We identify the strong interaction region (which appears in the regions D_4 and D_5 of Section 6) and perform very careful estimates there. The results match those obtained by the heuristic argument.

Through these sharp convolution estimates, we propose an appropriate ansatz, which is exponential decay for f_1 and polynomially sharp for f_2 as the main focus here is the region inside wave cone, where only polynomial decay can be expected. Justifying the ansatz involves even more complicated convolution estimates, but the underlying idea remains similar. Additionally, the damped transport operator \mathbb{S}^t is used to compensate the loss of velocity decay from the nonlinear operator Γ in the justification for f_1 (see Lemmas 4 and 5). Our result distinguishes between the linear and nonlinear parts of the solution, significantly improving upon previous results in [10, 12, 20, 23, 24], where the nonlinear couplings and the linear part have the same decay rate.

1.5.2. Exponential decay outside the acoustic wave cone. In Theorem 9, we obtain space-time exponential decay for f_1 , but for f_2 , we only achieve a polynomial decay estimate. This is because we designed a polynomial-type ansatz to facilitate the control of the nonlinear part of f_2 . This ansatz is accurate for the structure inside the acoustic wave cone. However, since the initial data has compact support in space and f_2 corresponds to a fluid structure that propagates at a finite speed, we expect the solution to decay exponentially outside the wave cone.

To observe the behavior of f_2 outside the cone, we multiply a suitable weight function on f_2 and prove an L^∞ bound of the weighted solution through regularization and energy estimates. This approach for obtaining the space asymptotic behavior of the Boltzmann equation was developed in our previous work [21], and here it is adapted to handle f_2 .

It is worth mentioning that in our previous work, we proved exponential decay outside a wave cone $|x| < Mt$ for a sufficiently large M . Here, through careful calculation of the micro projection, we show that f_2 indeed decays space-time exponentially outside the wave cone $|x| < (\mathbf{c} + \delta)t$ for any positive δ (see Theorem 10), where \mathbf{c} is the sound speed. This result is more consistent with physical reality.

1.5.3. The transition from polynomial tail to Gaussian tail. In the decomposition $f = f_1 + \sqrt{\mathcal{M}}f_2$, the latter already exhibits Gaussian tail. Therefore, studying the transition process is equivalent to examining the generation of the Gaussian tail for f_1 . The mechanism for generating velocity decay comes from $e^{-\nu(\xi)t}$ in the damped transport operator \mathbb{S}^t :

$$\mathbb{S}^t g_0 = e^{-\nu(\xi)t} g_0(x - \xi t, \xi).$$

At first glance, it seems that f_1 can gain arbitrary velocity decay as time evolves. However, the upper bound for velocity decay is limited by the coupling between f_1 and $\sqrt{\mathcal{M}}f_2$, specifically by the term

$$\int_0^t \mathbb{S}^{t-\tau} \left[Q(f_1, \sqrt{\mathcal{M}}f_2) + Q(\sqrt{\mathcal{M}}f_2, f_1) \right](\tau) d\tau$$

in the second equation of (5). The velocity weight will ultimately be slowed down by $\sqrt{\mathcal{M}}f_2$ as it decay at most as a Gaussian. We design a suitable weighted function

$$e^{\kappa(\xi) \min\{\langle \xi \rangle \wedge t\}}$$

to capture this feature. By carefully analyzing the commutator between this weight function and the \mathcal{K}_s , Q operators, we complete the description of dynamic transition process (see Theorem 15).

It is interesting to note that the mechanism for gaining velocity weight and the limitation of the maximal generation both stem from collisions with the global Maxwellian. This is entirely consistent with our physical intuition.

1.6. Organization of the paper. The rest of this paper is organized as follows: In Section 2, we first prepare some basic properties of the integral operator \mathcal{K} , and present some estimates for the damped transport equation and the linearized Boltzmann equation. In Section 3, we construct the global wave structures of the solution, fully utilizing sharp nonlinear wave interactions. In Section 4, we apply the weighted energy estimate to prove that the solution indeed decays exponentially

in space-time outside the wave cone. In Section 5, we provide a quantitative description of how solution approaches a Gaussian tail in terms of the microscopic velocity. Finally, we present the proof of all kinds of wave interactions in Section 6.

2. PRELIMINARIES

To begin with, we study some essential properties of the collision frequency $\nu(\xi)$, the operator \mathcal{K} and the collision operator Q . It is well known that there exist two positive constants ν_0 and ν_1 such that

$$\nu_0 \langle \xi \rangle \leq \nu(\xi) = \int_{\mathbb{R}^3} \int_{S^2} |q \cdot n| \mathcal{M}(\xi_*) dnd\xi_* \leq \nu_1 \langle \xi \rangle$$

for all $\xi \in \mathbb{R}^3$. In [5, Lemma 2.1 and Lemma 6.2], the following estimates of the collision kernel have been proved.

Lemma 2 ([5]). *For any $\beta > 4$, there exists a constant $C_\beta > 0$ depending only on β such that*

$$\int_{\mathbb{R}^3} \int_{S^2} |\xi - \xi_*| \frac{w_\beta(\xi)}{w_\beta(\xi'_*)} e^{-\frac{1}{4}|\xi'^2} dnd\xi_* \leq \frac{C}{\beta} \nu(\xi) + C_\beta \frac{\nu(\xi)}{(1 + |\xi|)^2}$$

and

$$\int_{\mathbb{R}^3} \int_{S^2} |\xi - \xi_*| \frac{w_\beta(\xi)}{w_\beta(\xi') w_\beta(\xi'_*)} dnd\xi_* \leq \frac{C}{\beta} \nu(\xi) + C_\beta \frac{\nu(\xi)}{(1 + |\xi|)^2}.$$

where $C > 0$ is a universal constant independent of β . Here $w_\beta(\xi) = \langle \xi \rangle^\beta$.

Using this lemma, one can get the estimates for the operators \mathcal{K} and Q .

Lemma 3 ([5]). *For any $\beta > 4$, there exists a constant $C_\beta > 0$ depending only on β such that*

$$\langle \xi \rangle^\beta |\mathcal{K}f| \leq \left(\frac{C}{\beta} \nu(\xi) + C_\beta \frac{\nu(\xi)}{(1 + |\xi|)^2} \right) |f|_{L_{\xi, \beta}^\infty}$$

and

$$\left| \langle \xi \rangle^\beta Q(f, g) \right| \leq C_\beta \nu(\xi) |f|_{L_{\xi, \beta}^\infty} |g|_{L_{\xi, \beta}^\infty}.$$

Moreover, for any $R > 0$,

$$\chi_{\{|\xi| \geq R\}} \langle \xi \rangle^\beta |\mathcal{K}f| \leq \left(\frac{C}{\beta} + \frac{C_\beta}{R^2} \right) \nu(\xi) |f|_{L_{\xi, \beta}^\infty}.$$

In order to study the first equation of (4), we introduce the damped transport operator \mathbb{S}^t , that is, $\mathbb{S}^t g_0$ is the solution of the equation:

$$\partial_t g + \xi \cdot \nabla_x g + \nu(\xi) g = 0, \quad g(0, x, \xi) = g_0.$$

Lemma 4. *Let $\beta > 4$ and $M > 0$. Assume that $U(t, x, \xi)$ and $h(t, x, \xi)$ satisfy*

$$\left| \nu(\xi)^{-1} U(t, x, \xi) \right|_{L_{\xi, \beta}^\infty} \leq A e^{-\frac{t+|x|}{c}}$$

and

$$|h(t, x, \xi)|_{L_{\xi, \beta}^\infty} \leq B e^{-\frac{t+|x|}{c}}$$

for some constants $A, c > 0$ with $\frac{\nu_0}{2} > \frac{1}{c}$. If g satisfies the integral equation

$$g(t, x, \xi) = \int_0^t \mathbb{S}^{t-\tau} (\mathcal{K}_s h + U)(\tau, x, \xi) d\tau,$$

then

$$|g(t, x, \xi)|_{L_{\xi, \beta}^\infty} \leq 2(A + \eta B) e^{-\frac{t+|x|}{c}},$$

where

$$\eta = \eta(\beta, R) = \left(\frac{C}{\beta} + \frac{C_\beta}{R^2} \right).$$

Proof. Let $y = x - (t - \tau)\xi$, we have

$$\begin{aligned}
\left| \langle \xi \rangle^\beta g(t, x, \xi) \right| &= \left| \int_0^t e^{-\nu(\xi)(t-\tau)} \langle \xi \rangle^\beta [\mathcal{K}_s h(\tau, x - \xi(t-\tau), \xi) + U(\tau, x - \xi(t-\tau), \xi)] d\tau \right| \\
&\leq \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) (\nu(\xi))^{-1} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} \langle \xi \rangle^\beta [|\mathcal{K}_s h(\tau, y, \xi)|] d\tau \\
&\quad + \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) (\nu(\xi))^{-1} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} \langle \xi \rangle^\beta |U(\tau, y, \xi)| d\tau \\
&\leq \eta \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) \cdot \sup_{y \in \mathbb{R}^3} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} |h(\tau, y, \xi)|_{L_{\xi, \beta}^\infty} d\tau \\
&\quad + \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) \cdot \sup_{y \in \mathbb{R}^3} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} |\nu^{-1}(\xi) U(\tau, y, \xi)|_{L_{\xi, \beta}^\infty} d\tau \\
&\leq 2(A + \eta B) e^{-\frac{t+|x|}{c}}.
\end{aligned}$$

The proof is completed. \square

Lemma 5. Let $\alpha > 0$ and $\rho > 0$. Assume that $V(t, x, \xi)$ satisfies

$$\left| \nu(\xi)^{-1} V(t, x, \xi) \right|_{L_{\xi, \beta}^\infty} \leq (1+t)^{-\alpha} \left(1 + \frac{|x|^2}{1+t} \right)^{-\rho}.$$

If g satisfies the integral equation

$$g(t, x, \xi) = \int_0^t \mathbb{S}^{t-\tau} V(\tau, x, \xi) d\tau,$$

then

$$|g(t, x, \xi)|_{L_{\xi, \beta}^\infty} \leq C_{\alpha, \rho} (1+t)^{-\alpha} \left(1 + \frac{|x|^2}{1+t} \right)^{-\rho}$$

for some constant $C_{\alpha, \rho} > 0$.

Proof. The proof is similar to Lemma 4. It suffices to verify that

$$\begin{aligned}
&\int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) \sup_{y \in \mathbb{R}^3} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} (1+\tau)^{-\alpha} \left(1 + \frac{|y|^2}{1+\tau} \right)^{-\rho} d\tau \\
&\leq C_{\alpha, \rho} (1+t)^{-\alpha} \left(1 + \frac{|x|^2}{1+t} \right)^{-\rho}.
\end{aligned}$$

Let $0 \leq \tau \leq t$. If $|y| > |x|/2$, then

$$\left(1 + \frac{|y|^2}{1+\tau} \right)^{-\rho} \leq \left(1 + \frac{|x|^2}{4(1+t)} \right)^{-\rho} \leq 4^\rho \left(1 + \frac{|x|^2}{1+t} \right)^{-\rho},$$

and thus

$$e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} (1+\tau)^{-\alpha} \left(1 + \frac{|y|^2}{1+\tau} \right)^{-\rho} \leq 4^\rho (1+\tau)^{-\alpha} \left(1 + \frac{|x|^2}{1+t} \right)^{-\rho}.$$

If $|y| \leq |x|/2$, then

$$e^{-\frac{\nu_0}{2}|x-y|} \leq e^{-\frac{\nu_0|x|}{4}} \leq C_\rho (1+|x|^2)^{-\rho} \leq C_\rho \left(1 + \frac{|x|^2}{1+t} \right)^{-\rho},$$

and so

$$e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} (1+\tau)^{-\alpha} \left(1 + \frac{|y|^2}{1+\tau} \right)^{-\rho} \leq C_\rho (1+\tau)^{-\alpha} \left(1 + \frac{|x|^2}{1+t} \right)^{-\rho}.$$

Therefore,

$$\begin{aligned}
& \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) \sup_{y \in \mathbb{R}^3} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} (1+\tau)^{-\alpha} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-\rho} d\tau \\
& \leq (C_\rho + 4^\rho) \left(1 + \frac{|x|^2}{1+t}\right)^{-\rho} \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) (1+\tau)^{-\alpha} d\tau \\
& \leq C_\alpha (C_\rho + 4^\rho) (1+t)^{-\alpha} \left(1 + \frac{|x|^2}{1+t}\right)^{-\rho}.
\end{aligned}$$

□

Lemma 6. Let $\alpha > 0$ and $\rho > 0$. Assume that $W(t, x, \xi)$ satisfies

$$\left| \nu(\xi)^{-1} W(t, x, \xi) \right|_{L_{\xi, \beta}^\infty} \leq (1+t)^{-\alpha} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho}.$$

If g satisfies the integral equation

$$g(t, x, \xi) = \int_0^t \mathbb{S}^{t-\tau} W(\tau, x, \xi) d\tau,$$

then

$$|g(t, x, \xi)|_{L_{\xi, \beta}^\infty} \leq C_{\alpha, \rho} (1+t)^{-\alpha} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho}$$

for some constant $C_{\alpha, \rho} > 0$.

Proof. The proof is similar to Lemma 4 and it suffices to verify that

$$\begin{aligned}
& \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) \sup_{y \in \mathbb{R}^3} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} (1+\tau)^{-\alpha} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-\rho} d\tau \\
& \leq C_{\alpha, \rho} (1+t)^{-\alpha} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho}
\end{aligned}$$

We consider two cases $|x| \geq \mathbf{c}t$ and $|x| \leq \mathbf{c}t$.

Case 1: $|x| \geq \mathbf{c}t$. We split \mathbb{R}^3 into two parts

$$\left\{ y \in \mathbb{R}^3 : ||y| - \mathbf{c}\tau| > \frac{|x| - \mathbf{c}t}{2} \right\} \text{ and } \left\{ y \in \mathbb{R}^3 : ||y| - \mathbf{c}\tau| \leq \frac{|x| - \mathbf{c}t}{2} \right\}.$$

If $||y| - \mathbf{c}\tau| \leq \frac{|x| - \mathbf{c}t}{2}$, then

$$\begin{aligned}
|x-y| & \geq ||x| - |y|| = ||x| - \mathbf{c}\tau + \mathbf{c}\tau - |y|| \geq ||x| - \mathbf{c}\tau| - ||y| - \mathbf{c}\tau| \\
& \geq |x| - \mathbf{c}t - \frac{|x| - \mathbf{c}t}{2} = \frac{|x| - \mathbf{c}t}{2},
\end{aligned}$$

and thus

$$e^{-\frac{\nu_0}{2}|x-y|} \leq e^{-\frac{\nu_0||x| - \mathbf{c}t|}{4}} \leq C_\rho \left(1 + (|x| - \mathbf{c}t)^2\right)^{-\rho} \leq C_\rho \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-\rho}.$$

If $||y| - \mathbf{c}\tau| > \frac{|x| - \mathbf{c}t}{2}$, then

$$\left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-\rho} \leq \left(1 + \frac{(|x| - \mathbf{c}t)^2}{4(1+t)}\right)^{-\rho} \leq 4^\rho \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-\rho}.$$

Therefore,

$$\begin{aligned}
& \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) \sup_{y \in \mathbb{R}^3} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} (1+\tau)^{-\alpha} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{(1+\tau)}\right)^{-\rho} d\tau \\
& \leq (C_\rho + 4^\rho) \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-\rho} \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) (1+\tau)^{-\alpha} d\tau \\
& \leq C_\alpha (C_\rho + 4^\rho) (1+t)^{-\alpha} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-\rho}.
\end{aligned}$$

Case 2: $|x| \leq \mathbf{c}t$. If $0 \leq \tau \leq \frac{t}{2} + \frac{|x|}{2\mathbf{c}}$, then

$$t - \tau \geq t - \left(\frac{t}{2} + \frac{|x|}{2\mathbf{c}}\right) = \frac{\mathbf{c}t - |x|}{2\mathbf{c}} \geq 0,$$

and thus

$$e^{-\nu_0(t-\tau)} \leq e^{-\frac{\nu_0}{2\mathbf{c}}(\mathbf{c}t - |x|)} \leq C_\rho \left(1 + (\mathbf{c}t - |x|)^2\right)^{-\rho} \leq C_\rho \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho}.$$

It implies that

$$\begin{aligned}
& \int_0^{\frac{t}{2} + \frac{|x|}{2\mathbf{c}}} e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) \sup_{y \in \mathbb{R}^3} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} (1+\tau)^{-\alpha} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-\rho} d\tau \\
& \leq C_\rho \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho} \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) (1+\tau)^{-\alpha} d\tau \\
& \leq C_\alpha C_\rho (1+\tau)^{-\alpha} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho}.
\end{aligned}$$

If $\frac{t}{2} + \frac{|x|}{2\mathbf{c}} \leq \tau \leq t$, $|y| \leq \frac{\mathbf{c}\tau + |x|}{2}$, then

$$\mathbf{c}\tau - |y| \geq \frac{\mathbf{c}\tau - |x|}{2} \geq \frac{\mathbf{c}t - |x|}{4} \geq 0,$$

and so

$$\left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-\rho} \leq \left(1 + \frac{(\mathbf{c}t - |x|)^2}{16(1+t)}\right)^{-\rho} \leq 16^\rho \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho}.$$

If $\frac{t}{2} + \frac{|x|}{2\mathbf{c}} \leq \tau \leq t$, $|y| > \frac{\mathbf{c}\tau + |x|}{2}$, then

$$|x - y| \geq |y| - |x| \geq \frac{\mathbf{c}\tau - |x|}{2} \geq \frac{\mathbf{c}t - |x|}{4},$$

and thus

$$e^{-\frac{\nu_0}{2}|x-y|} \leq e^{-\frac{\nu_0}{8}(\mathbf{c}t - |x|)} \leq C_\rho \left(1 + (\mathbf{c}t - |x|)^2\right)^{-\rho} \leq C_\rho \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho}.$$

Consequently,

$$\begin{aligned}
& \int_{\frac{t}{2} + \frac{|x|}{2\mathbf{c}}}^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) \sup_{y \in \mathbb{R}^3} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} (1+\tau)^{-\alpha} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{(1+\tau)}\right)^{-\rho} d\tau \\
& \leq (16^\rho + C_\rho) \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho} \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) (1+\tau)^{-\alpha} d\tau \\
& \leq C_\alpha (16^\rho + C_\rho) (1+t)^{-\alpha} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho}.
\end{aligned}$$

Combining all the discussion, there exists a constant $C_{\alpha,\rho} > 0$ such that

$$\begin{aligned} & \int_0^t e^{-\frac{\nu(\xi)}{2}(t-\tau)} \nu(\xi) \sup_{y \in \mathbb{R}^3} e^{-\frac{\nu_0}{2}(t-\tau+|x-y|)} (1+\tau)^{-\alpha} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-\rho} d\tau \\ & \leq C_{\alpha,\rho} (1+t)^{-\alpha} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-\rho}, \end{aligned}$$

as desired. \square

For the second equation of decomposition (4), we list some basic properties of the linearized Boltzmann collision operator L and nonlinear operator Γ as below.

It is well known that the null space of L

$$\ker(L) = \text{span} \{\chi_0, \chi_1, \chi_2, \chi_3, \chi_4\},$$

is a five-dimensional vector space, where

$$\chi_0 = \mathcal{M}^{1/2}, \quad \chi_i = \xi_i \mathcal{M}^{1/2}, \quad \chi_4 = \frac{1}{\sqrt{6}} (|\xi|^2 - 3) \mathcal{M}^{1/2}, \quad i = 1, 2, 3.$$

Let P_0 be the orthogonal projection with respect to the L_ξ^2 inner product onto $\ker(L)$, and $P_1 \equiv \text{Id} - P_0$. That is, for any $g \in L_\xi^2$,

$$P_0 g = \sum_{i=0}^4 \langle \chi_i, g \rangle_\xi \chi_i, \quad P_1 g = g - P_0 g.$$

The solution of the wave propagation is connected to the operator $P_0(\xi \cdot \omega)P_0$ for $\omega \in \mathbb{S}^2$

$$(6) \quad \left\{ \begin{array}{l} P_0 \xi \cdot \omega E_j = \lambda_j E_j, \\ \lambda_0 = \mathbf{c}, \quad \lambda_1 = -\mathbf{c}, \quad \lambda_2 = \lambda_3 = \lambda_4 = 0, \quad \mathbf{c} = \sqrt{\frac{5}{3}}, \\ E_0 = \sqrt{\frac{3}{10}} \chi_0 + \sqrt{\frac{1}{2}} \omega \cdot \bar{\chi} + \sqrt{\frac{1}{5}} \chi_4, \\ E_1 = \sqrt{\frac{3}{10}} \chi_0 - \sqrt{\frac{1}{2}} \omega \cdot \bar{\chi} + \sqrt{\frac{1}{5}} \chi_4, \\ E_2 = -\sqrt{\frac{2}{5}} \chi_0 + \sqrt{\frac{3}{5}} \chi_4, \\ E_3 = \omega_1 \cdot \bar{\chi}, \\ E_4 = \omega_2 \cdot \bar{\chi}. \end{array} \right.$$

where $\bar{\chi} = (\chi_1, \chi_2, \chi_3)$, and $\{\omega_1, \omega_2, \omega\}$ is an orthonormal basis of \mathbb{R}^3 .

Lemma 7 ([26]). *The collision operator L consists of a multiplicative operator $\nu(\xi)$ and an integral operator K , namely,*

$$Lf = -\nu(\xi)f + Kf,$$

where $\nu(\xi) \sim (1 + |\xi|)$ and

$$|Kg|_{L_{\xi, \eta+1}^\infty} \leq C |g|_{L_{\xi, \eta}^\infty}, \quad \eta \in \mathbb{R},$$

for some universal constant $C > 0$.

The nonlinear operator Γ has the following estimate

$$|\nu^{-1}(\xi)\Gamma(g, h)|_{L_{\xi, \eta}^\infty} \leq C |g|_{L_{\xi, \eta}^\infty} |h|_{L_{\xi, \eta}^\infty}, \quad \eta \in \mathbb{R},$$

for some universal constant $C > 0$.

It is vital to get the space-time structure of the solution of the linearized Boltzmann equation

$$\partial_t h + \xi \cdot \nabla_x h = Lh,$$

and we denote \mathbb{G}^t as its solution operator.

Lemma 8 ([26, 27]). *The solution $h = \mathbb{G}^t h_0$ of the equation*

$$\partial_t h + \xi \cdot \nabla_x h = Lh, \quad h(0, x, \xi) = h_0(x, \xi).$$

satisfies

$$|\mathbb{G}^t h_0|_{L_{\xi, \beta}^\infty} \leq \mathcal{C}_1 \left[\begin{array}{l} (1+t)^{-2} e^{-\frac{(|x|-\mathbf{c}t)^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \\ + \mathbf{1}_{\{|x| \leq \mathbf{c}t\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \end{array} \right] *_x |h_0|_{L_{\xi, \beta}^\infty}$$

for any $\beta > 3/2$ and some constants $\mathcal{C}_1, c_0, D_0 > 0$, where $h_0 \in L_{\xi, \beta}^\infty(L_x^\infty \cap L_x^1)$. Moreover, if $\mathbb{P}_1 h_0 = 0$, then

$$|\mathbb{G}^t h_0|_{L_{\xi, \beta}^\infty} \leq \mathcal{C}_1 \left[\begin{array}{l} (1+t)^{-5/2} e^{-\frac{(|x|-\mathbf{c}t)^2}{D_0(1+t)}} + (1+t)^{-2} e^{-\frac{|x|^2}{D_0(1+t)}} \\ + \mathbf{1}_{\{|x| \leq \mathbf{c}t\}} (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \end{array} \right] *_x |h_0|_{L_{\xi, \beta}^\infty}.$$

3. GLOBAL WAVE STRUCTURE

In this section we will prove the global wave structure of the equation (3) which is described in the following theorem.

Theorem 9. *Let $\beta > 4$ be sufficiently large. Assume that $f_0 \in L_{\xi, \beta}^\infty L_x^\infty$ is compactly supported in the variable x for all ξ*

$$f_0(x, \xi) \equiv 0 \text{ for } |x| \geq 1, \quad \xi \in \mathbb{R}^3.$$

Then there exists $\varepsilon > 0$ small enough such that the solution f of (3) exists for $t > 0$ and it can be written as $f = f_1 + \sqrt{\mathcal{M}}f_2$, where f_1 and f_2 satisfy (4) with

$$|f_1|_{L_{\xi, \beta}^\infty} \leq 2\mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} e^{-\frac{\nu_0}{2}(t+|x|)},$$

and

$$\begin{aligned} & |f_2|_{L_{\xi, \beta}^\infty} \\ & \leq \mathfrak{B} \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \left[\begin{array}{l} (1+t)^{-2} e^{-\frac{(|x|-\mathbf{c}t)^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \\ + \mathbf{1}_{\{|x| \leq \mathbf{c}t\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \end{array} \right] \\ & + 2\mathfrak{C} \left(\mathfrak{B} \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \right)^2 \left[(1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-\frac{5}{2}} \left(1 + \frac{(|x|-\mathbf{c}t)^2}{1+t}\right)^{-1} \right] \end{aligned}$$

for some positive constants $\mathfrak{B}, \mathcal{C}_1, \mathfrak{C}, \hat{D}_0, \hat{c}_0$.

Proof. To clarify the space-time structures of the solutions f_1 and f_2 to (4), we design an iteration $\{f^n\}$ with $f^n = f_1^n + \sqrt{\mathcal{M}}f_2^n$, as follows:

$$(7) \quad \begin{cases} \partial_t f_1^{n+1} + \xi \cdot \nabla_x f_1^{n+1} + \nu(\xi) f_1^{n+1} = \mathcal{K}_s f_1^n + U(f_1^n, f_2^n) \\ \partial_t f_2^{n+1} + \xi \cdot \nabla_x f_2^{n+1} = L f_2^{n+1} + \mathcal{K}_b f_1^{n+1} + \Gamma(f_2^n, f_2^n) \end{cases}$$

with

$$f_1^{n+1}(0, x, \xi) = \varepsilon f_0(x, \xi), \quad f_2^{n+1}(0, x, \xi) = 0, \quad f_1^0(t, x, \xi) = f_2^0(t, x, \xi) = 0,$$

where

$$U(f_1^n, f_2^n) = Q(f_1^n, f_1^n) + Q\left(f_1^n, \sqrt{\mathcal{M}}f_2^n\right) + Q\left(\sqrt{\mathcal{M}}f_2^n, f_1^n\right).$$

We shall study the space-time structures of f_1^n and f_2^n . Now we rewrite (7) as integral forms:

$$\begin{cases} f_1^{n+1} = \varepsilon \mathbb{S}^t f_0 + \int_0^t \mathbb{S}^{t-\tau} [\mathcal{K}_s f_1^n(\tau) + U(f_1^n, f_2^n)(\tau)] d\tau =: f_{1,1}^{n+1} + f_{1,2}^{n+1}, \\ f_2^{n+1} = \int_0^t \mathbb{G}^{t-\tau} \mathcal{K}_b(f_1^{n+1})(\tau) d\tau + \int_0^t \mathbb{G}^{t-\tau} \Gamma(f_2^n, f_2^n)(\tau) d\tau =: f_{2,1}^{n+1} + f_{2,2}^{n+1}. \end{cases}$$

First, one can see that $f_1^1 = f_{1,1}^1$, $f_2^1 = f_{2,1}^1$ and $f_{1,2}^1 = f_{2,2}^1 = 0$. Since f_0 is compactly supported in the unit ball with respect to the variable x uniformly for all ξ , we have

$$|f_1^1|_{L_{\xi,\beta}^\infty} \leq \mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} e^{-\frac{\nu_0}{2}(t+|x|)}$$

for some constant \mathcal{C}_1 . In view of Lemmas 3, 8 and Lemmas 19 – 21, we get

$$\begin{aligned} |f_2^1|_{L_{\xi,\beta}^\infty} &= \left| \int_0^t \mathbb{G}^{t-\tau} \mathcal{K}_b(f_1^1)(\tau) d\tau \right|_{L_{\xi,\beta}^\infty} \\ &\leq 2\mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \left[(2\pi)^{\frac{3}{4}} e^{\frac{R^2}{4}} \left(\frac{C\nu_1}{\beta} (1+R) + C_\beta \right) \right] \\ &\quad \cdot \mathcal{C}_1 \left[(1+t)^{-2} e^{-\frac{(|x|-ct)^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \right. \\ &\quad \left. + \mathbf{1}_{\{|x|\leq ct\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \right]^{*_{x,t}} e^{-\frac{\nu_0}{2}(t+|x|)} \\ &\leq 2\mathcal{C}_2 \mathcal{C}_1^2 \left[(2\pi)^{\frac{3}{4}} e^{\frac{R^2}{4}} \left(\frac{C\nu_1}{\beta} (1+R) + C_\beta \right) \right] \\ &\quad \cdot \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \left[(1+t)^{-2} e^{-\frac{(|x|-ct)^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \right. \\ &\quad \left. + \mathbf{1}_{\{|x|\leq ct\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \right] \\ &\leq \mathfrak{B} \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \left[(1+t)^{-2} e^{-\frac{(|x|-ct)^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \right. \\ &\quad \left. + \mathbf{1}_{\{|x|\leq ct\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \right]. \end{aligned}$$

In fact, by definition of $f_{1,1}^n$, we have $f_{1,1}^n = f_{1,1}^1$ for all n and so

$$|f_{1,1}^n|_{L_{\xi,\beta}^\infty} \leq \mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} e^{-\frac{\nu_0}{2}(t+|x|)}.$$

Now we choose $\beta > 4$ and $R > 0$ sufficiently large such that the constant $\eta(\beta, R)$ defined in Lemma 4 satisfies

$$\eta(\beta, R) = \frac{C}{\beta} + \frac{C_\beta}{R^2} < \frac{1}{8}.$$

Fixing such β and R , we shall prove that if $\varepsilon > 0$ is sufficiently small, we have

$$|f_{1,2}^n|_{L_{\xi,\beta}^\infty} \leq \mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} e^{-\frac{\nu_0}{2}(t+|x|)},$$

$$|f_{2,1}^n|_{L_{\xi,\beta}^\infty} \leq \mathfrak{B} \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \left[(1+t)^{-2} e^{-\frac{(|x|-ct)^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \right. \\ \left. + \mathbf{1}_{\{|x|\leq ct\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \right],$$

$$|f_{2,2}^n|_{L_{\xi,\beta}^\infty} \leq 2\mathfrak{C} \left(\mathfrak{B} \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \left[(1+t)^{-2} \left(1 + \frac{|x|^2}{1+t} \right)^{-\frac{3}{2}} + (1+t)^{-\frac{5}{2}} \left(1 + \frac{(|x|-ct)^2}{1+t} \right)^{-1} \right],$$

for all $n \in \mathbb{N}$ by induction on n , here the large constant $\mathfrak{C} > 0$ will be determined later.

By induction hypothesis and Lemma 3, we have

$$\begin{aligned} |\nu^{-1} U(f_1^n, f_2^n)|_{L_{\xi,\beta}^\infty} &\leq C \left(|f_1^n|_{L_{\xi,\beta}^\infty}^2 + 2|f_1^n|_{L_{\xi,\beta}^\infty} |f_2^n|_{L_{\xi,\beta}^\infty} \right) \\ &\leq C \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \left(4\mathcal{C}_1 + \mathfrak{B} \left(1 + 2\mathfrak{C} \mathfrak{B} \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right) \right) \end{aligned}$$

$$\cdot \mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} e^{-\frac{\nu_0}{2}(t+|x|)}.$$

By Lemma 4,

$$\begin{aligned} |f_{1,2}^{n+1}|_{L_{\xi, \beta}^\infty} &= \left| \int_0^t \langle \xi \rangle^\beta \mathbb{S}^{t-\tau} [\mathcal{K}_s f_1^n(\tau) + U(f_1^n, f_2^n)(\tau)] d\tau \right| \\ &\leq \left[4\eta(\beta, R) + 2C\varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \left(4\mathcal{C}_1 + \mathfrak{B} \left(1 + 2\mathfrak{C}\mathfrak{B}\varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \right) \right) \right] \\ &\quad \cdot \mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} e^{-\frac{\nu_0}{2}(t+|x|)} \\ &\leq \mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} e^{-\frac{\nu_0}{2}(t+|x|)}, \end{aligned}$$

after we choose $\varepsilon > 0$ sufficiently small such that

$$2C\varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \left(4\mathcal{C}_1 + 2\mathfrak{C}\mathfrak{B} \left(1 + \mathfrak{B}\varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \right) \right) < 1/2.$$

Therefore,

$$|f_{1,2}^{n+1}|_{L_{\xi, \beta}^\infty} \leq \mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} e^{-\frac{\nu_0}{2}(t+|x|)},$$

and so

$$|f_1^{n+1}|_{L_{\xi, \beta}^\infty} \leq 2\mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} e^{-\frac{\nu_0}{2}(t+|x|)}.$$

Likewise $f_{2,1}^1$, we have

$$\begin{aligned} |f_{2,1}^{n+1}|_{L_{\xi, \beta}^\infty} &= \left| \int_0^t \mathbb{G}^{t-\tau} \mathcal{K}_b (f_1^{n+1})(\tau) d\tau \right|_{L_{\xi, \beta}^\infty} \\ &\leq 2\mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \left[(2\pi)^{\frac{3}{4}} e^{\frac{R^2}{4}} \left(\frac{C\nu_1}{\beta} (1+R) + C_\beta \right) \right] \\ &\quad \cdot \mathcal{C}_1 \left[\begin{array}{l} (1+t)^{-2} e^{-\frac{(|x|-\mathbf{c}t)^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \\ + \mathbf{1}_{\{|x| \leq \mathbf{c}t\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \end{array} \right] *_{x,t} e^{-\frac{\nu_0}{2}(t+|x|)} \\ &\leq 2\mathcal{C}_2 \mathcal{C}_1^2 \left[(2\pi)^{\frac{3}{4}} e^{\frac{R^2}{4}} \left(\frac{C\nu_1}{\beta} (1+R) + C_\beta \right) \right] \\ &\quad \cdot \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \left[\begin{array}{l} (1+t)^{-2} e^{-\frac{(|x|-\mathbf{c}t)^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \\ + \mathbf{1}_{\{|x| \leq \mathbf{c}t\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \end{array} \right] \\ &\leq \mathfrak{B}\varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \left[\begin{array}{l} (1+t)^{-2} e^{-\frac{(|x|-\mathbf{c}t)^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \\ + \mathbf{1}_{\{|x| \leq \mathbf{c}t\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \end{array} \right]. \end{aligned}$$

Also, there exists a constant $\mathcal{C}_3 > 1$ depending on \widehat{D}_0 and \widehat{c}_0 such that

$$\begin{aligned} &|f_{2,1}^{n+1}|_{L_{\xi, \beta}^\infty} \\ (8) \quad &\leq 2\mathcal{C}_2 \mathcal{C}_1^2 \left[(2\pi)^{\frac{3}{4}} e^{\frac{R^2}{4}} \left(\frac{C\nu_1}{\beta} (1+R) + C_\beta \right) \right] \\ &\quad \cdot \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \left[\begin{array}{l} (1+t)^{-2} e^{-\frac{(|x|-\mathbf{c}t)^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \\ + \mathbf{1}_{\{|x| \leq \mathbf{c}t\}} (1+t)^{-\frac{3}{2}} \left(1 + \frac{|x|^2}{1+t} \right)^{-\frac{3}{2}} + e^{-\frac{t+|x|}{\widehat{c}_0}} \end{array} \right] \\ &\leq 2\mathcal{C}_3 \mathcal{C}_2 \mathcal{C}_1^2 \left[(2\pi)^{\frac{3}{4}} e^{\frac{R^2}{4}} \left(\frac{C\nu_1}{\beta} (1+R) + C_\beta \right) \right] \end{aligned}$$

$$\begin{aligned} & \cdot \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \left[(1+t)^{-\frac{3}{2}} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1} \right] \\ & \leq \mathfrak{B}\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \left[(1+t)^{-\frac{3}{2}} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1} \right]. \end{aligned}$$

Since $f_{2,2}^{n+1}$ satisfies the equation

$$\partial_t f_{2,2}^{n+1} + \xi \cdot \nabla_x f_{2,2}^{n+1} = L f_{2,2}^{n+1} + \Gamma(f_2^n, f_2^n) \text{ with } f_{2,2}^{n+1}(0, x, \xi) = 0,$$

we further decompose $f_{2,2}^{n+1}$ into two parts as $f_{2,2}^{n+1} = h_1^{n+1} + h_2^{n+1}$ where h_1^{n+1}, h_2^{n+1} satisfy the equations

$$\begin{cases} \partial_t h_1^{n+1} + \xi \cdot \nabla_x h_1^{n+1} + \nu(\xi) h_1^{n+1} = \Gamma(f_2^n, f_2^n), \\ \partial_t h_2^{n+1} + \xi \cdot \nabla_x h_2^{n+1} + \nu(\xi) h_2^{n+1} = K f_{2,2}^{n+1}, \end{cases}$$

with $h_1^{n+1}(0, x, \xi) = h_2^{n+1}(0, x, \xi) = 0$. By induction hypothesis and (8),

$$\begin{aligned} |f_2^n|_{L_{\xi,\beta}^\infty} &= |f_{2,1}^n + f_{2,2}^n|_{L_{\xi,\beta}^\infty} \\ &\leq \left[\mathfrak{B}\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} + 2\mathfrak{C} \left(\mathfrak{B}\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \right] \\ &\quad \cdot \left[(1+t)^{-2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1} + (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} \right] \\ &\leq 2\mathfrak{B}\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \left[(1+t)^{-2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1} + (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} \right], \end{aligned}$$

so that

$$\begin{aligned} |\nu^{-1}\Gamma(f_2^n, f_2^n)|_{L_{\xi,\beta}^\infty} &\leq C |f_2^n|_{L_{\xi,\beta}^\infty}^2 \\ (9) \quad &\leq 8C \left(\mathfrak{B}\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \\ &\quad \cdot \left[(1+t)^{-4} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-2} + (1+t)^{-3} \left(1 + \frac{|x|^2}{1+t}\right)^{-3} \right]. \end{aligned}$$

In view of Lemmas 5 and 6, we have

$$\begin{aligned} & \left| \langle \xi \rangle^\beta h_1^{n+1} \right| \\ (10) \quad &= \left| \langle \xi \rangle^\beta \int_0^t \mathbb{S}^{t-\tau} \Gamma(f_2^n, f_2^n)(\tau) d\tau \right| \\ &\leq 8C^2 \left(\mathfrak{B}\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \\ &\quad \cdot \left[(1+t)^{-4} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{(1+t)}\right)^{-2} + (1+t)^{-3} \left(1 + \frac{|x|^2}{(1+t)}\right)^{-3} \right] \\ &\leq \mathfrak{C} \left(\mathfrak{B}\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \left[(1+t)^{-4} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{(1+t)}\right)^{-2} + (1+t)^{-3} \left(1 + \frac{|x|^2}{(1+t)}\right)^{-3} \right]. \end{aligned}$$

As for h_2^{n+1} , we have

$$\left| \langle \xi \rangle^\beta h_2^{n+1} \right| = \left| \langle \xi \rangle^\beta \int_0^t \mathbb{S}^{t-\tau} K(f_{2,2}^{n+1})(\tau, x, \xi) d\tau \right|.$$

It follows from Lemma 7 that

$$\left| \nu(\xi)^{-1} \langle \xi \rangle^\beta K f_{2,2}^{n+1} \right|_{L_\xi^\infty} \leq \frac{C}{\nu_0} |f_{2,2}^{n+1}|_{L_{\xi,\beta-1}^\infty}.$$

In view of Lemma 8 and (9), together with convolution estimates in Section 6.2, to be more specific, Lemmas 22–27, we obtain

$$\begin{aligned} |f_{2,2}^{n+1}|_{L_{\xi,\beta-1}^\infty} &= \left| \int_0^t \langle \xi \rangle^{\beta-1} \mathbb{G}^{t-\tau} \Gamma(f_2^n, f_2^n)(\tau) d\tau \right| \\ &\leq \mathcal{C}_1 \nu_1 \left[(1+t)^{-\frac{5}{2}} e^{-\frac{(|x|-\mathbf{ct})^2}{D_0(1+t)}} + (1+t)^{-2} e^{-\frac{|x|^2}{D_0(1+t)}} \right. \\ &\quad \left. + \mathbf{1}_{\{|x| \leq \mathbf{ct}\}} (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \right] *_{x,t} |\nu^{-1} \Gamma(f_2^n, f_2^n)|_{L_{\xi,\beta}^\infty} \\ &\leq \mathcal{C}_1 \nu_1 8C \left(\mathfrak{B}_\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \left[(1+t)^{-\frac{5}{2}} e^{-\frac{(|x|-\mathbf{ct})^2}{D_0(1+t)}} + (1+t)^{-2} e^{-\frac{|x|^2}{D_0(1+t)}} \right. \\ &\quad \left. + \mathbf{1}_{\{|x| \leq \mathbf{ct}\}} (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \right] \\ &\quad *_{x,t} \left[(1+t)^{-4} \left(1 + \frac{(|x|-\mathbf{ct})^2}{1+t}\right)^{-2} + (1+t)^{-3} \left(1 + \frac{|x|^2}{1+t}\right)^{-3} \right] \\ &\leq 8C^2 \mathcal{C}_1 \nu_1 \left(\mathfrak{B}_\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \\ &\quad \cdot \left[(1+t)^{-\frac{5}{2}} \left(1 + \frac{(|x|-\mathbf{ct})^2}{1+t}\right)^{-1} + (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} \right]. \end{aligned}$$

Therefore, using Lemmas 5 and 6,

$$\begin{aligned} &|h_2^{n+1}|_{L_{\xi,\beta}^\infty} \\ (11) \quad &\leq C \cdot \frac{C}{\nu_0} 8C^2 \mathcal{C}_1 \nu_1 \left(\mathfrak{B}_\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \\ &\quad \cdot \left[(1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-\frac{5}{2}} \left(1 + \frac{(|x|-\mathbf{ct})^2}{1+t}\right)^{-1} \right] \\ &\leq \mathfrak{C} \left(\mathfrak{B}_\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \left[(1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-\frac{5}{2}} \left(1 + \frac{(|x|-\mathbf{ct})^2}{1+t}\right)^{-1} \right] \end{aligned}$$

after we choose $\mathfrak{C} > 0$ sufficiently large. Combining (10) and (11) gives

$$|f_{2,2}^{n+1}|_{L_{\xi,\beta}^\infty} \leq 2\mathfrak{C} \left(\mathfrak{B}_\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \left[(1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-\frac{5}{2}} \left(1 + \frac{(|x|-\mathbf{ct})^2}{1+t}\right)^{-1} \right].$$

Consequently, we get the desired estimates for $f_{1,2}^n, f_{2,1}^n, f_{2,2}^n$ by induction on n .

Finally, it is straightforward to prove that $\{(f_1^n, f_2^n)\}$ is a Cauchy sequence in $L_{\xi,\beta}^\infty (L_x^2 \cap L_x^\infty)$. As a consequence, (f_1^n, f_2^n) converges to (f_1, f_2) in $L_{\xi,\beta}^\infty (L_x^2 \cap L_x^\infty)$ and (f_1, f_2) solves the problem (4) with

$$|f_1|_{L_{\xi,\beta}^\infty} \leq 2\mathcal{C}_1 \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} e^{-\frac{\nu_0}{2}(t+|x|)},$$

and

$$\begin{aligned} &|f_2|_{L_{\xi,\beta}^\infty} \\ &\leq \mathfrak{B}_\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \left[(1+t)^{-2} e^{-\frac{(|x|-\mathbf{ct})^2}{D_0(1+t)}} + (1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} \right. \\ &\quad \left. + \mathbf{1}_{\{|x| \leq \mathbf{ct}\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} \right] \end{aligned}$$

$$+ 2\mathfrak{C} \left(\mathfrak{B}\varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right)^2 \left[(1+t)^{-2} \left(1 + \frac{|x|^2}{1+t} \right)^{-\frac{3}{2}} + (1+t)^{-\frac{5}{2}} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t} \right)^{-1} \right].$$

The proof of Theorem 9 is completed. \square

4. EXPONENTIAL DECAY OUTSIDE THE WAVE CONE

In Theorem 9, the estimate for f_1 is exponentially sharp, while the estimate of f_2 is only polynomially sharp. The reason is that, to facilitate the closure of nonlinearity, we focused on the structure inside sound wave cone, so we chose a polynomial ansatz. However, since our initial data is compactly supported in space and f_2 represents the fluid part with an essentially finite propagation speed. Therefore, we expect a faster decay in the space-like region.

In this section, we will improve the behavior of f_2 outside the sound wave cone. Indeed, we can prove that f_2 has exponential decay both in space and time there. The result is stated as follows.

Theorem 10. *Under the same assumption of Theorem 9, for any $0 < \delta \ll 1$, if $\varepsilon > 0$ is sufficiently small, there exists a large positive constant D depending on δ , such that for $|x| > (\mathbf{c} + \delta)t$,*

$$|f_2|_{L_{\xi,\beta}^\infty} \leq C_\delta \varepsilon e^{-\frac{\langle x \rangle + t}{D}},$$

where the constant $C_\delta > 0$ is independent of time and $C_\delta \rightarrow \infty$ as $\delta \rightarrow 0$.

To attain this end, we consider the weighted nonlinear equation corresponding to (4). That is, let $u = f_w := wf = u_1 + \sqrt{\mathcal{M}}u_2$, $u_i = wf_i$ ($i = 1, 2$), where the weight function is given by

$$w(t, x) = \exp\left(\frac{\langle x \rangle - Mt}{\ell}\right),$$

$\mathbf{c} < M < \mathbf{c} + 1$ and sufficiently large $\ell > 0$. Note that by Theorem 9, f_1 decays in space and time exponentially, so that u_1 can be controlled if ℓ is large enough, that is, there exists $c_0 > 0$ such that

$$\|u_1\|_{L_{\xi,\beta}^\infty L_x^\infty}, \|u_1\|_{L_{\xi,\beta}^\infty L_x^2} \lesssim \varepsilon e^{-c_0 t}.$$

In view of (4), u_2 satisfies the equation

$$\begin{cases} \partial_t u_2 + \xi \cdot \nabla_x u_2 - w^{-1} (\partial_t w + \xi \cdot \nabla_x w) u_2 = Lu_2 + \mathcal{K}_b u_1 + \Gamma(f_2, u_2) \\ u_2(0, x, \xi) = 0. \end{cases}$$

After choosing $\ell > 0$ large such that $M\ell^{-1}$ is small, we have

$$\tilde{\nu}(t, x, \xi) = \nu(\xi) + w^{-1} (\partial_t w + \xi \cdot \nabla_x w) \geq \frac{3}{4} \nu(\xi).$$

Under this situation, we are ready to estimate u_2 . Let $T > 0$ be a finite number. Denote

$$C_{u_2, T}^\infty = \varepsilon^{-1} \sup_{0 \leq t \leq T} \|u_2\|_{L_{\xi,\beta}^\infty L_x^\infty}, \quad C_{u_2, T}^2 = \varepsilon^{-1} \sup_{0 \leq t \leq T} \|u_2\|_{L_{\xi,\beta}^\infty L_x^2}.$$

Also denote

$$C_{f_2}^\infty = \varepsilon^{-1} \sup_{t \geq 0} (1+t)^{3/2} \|f_2\|_{L_{\xi,\beta}^\infty L_x^\infty}, \quad C_{f_2}^2 = \varepsilon^{-1} \sup_{t \geq 0} (1+t)^{3/4} \|f_2\|_{L_{\xi,\beta}^\infty L_x^2},$$

which are finite due to Theorem 9.

To estimate u_2 , we design a Picard-type iteration: the zeroth order approximation $u_2^{(0)}$ is defined as

$$\partial_t u_2^{(0)} + \xi \cdot \nabla_x u_2^{(0)} + \tilde{\nu}(\xi) u_2^{(0)} = \mathcal{K}_b u_1 + \Gamma(f_2, u_2), \quad u_2^{(0)}(0, x, \xi) = 0,$$

and define the j th order approximation $u_2^{(j)}$, $j \geq 1$, inductively as

$$\partial_t u_2^{(j)} + \xi \cdot \nabla_x u_2^{(j)} + \tilde{\nu}(\xi) u_2^{(j)} = \mathcal{K} u_2^{(j-1)}, \quad u_2^{(j-1)}(0, x, \xi) = 0.$$

Thus, the wave part and the remainder part are defined respectively as follows:

$$W_w^{(m)} = \sum_{j=0}^m u_2^{(j)}, \quad \mathcal{R}_w^{(m)} = u_2 - W_w^{(m)},$$

$\mathcal{R}_w^{(m)}$ solving the equation

$$\partial_t \mathcal{R}_w^{(m)} + \xi \cdot \nabla_x \mathcal{R}_w^{(m)} + \tilde{\nu}(\xi) \mathcal{R}_w^{(m)} = K \mathcal{R}_w^{(m)} + K u_2^{(m)}, \quad \mathcal{R}_w^{(m)}(0, x, \xi) = 0.$$

In fact, $m = 6$ is enough to get the H_x^2 regularization estimate for the remainder part. Following a similar argument as [21], we have

Lemma 11. *Let $\beta > 4$ be large enough. Then for $0 \leq t \leq T$,*

$$\left\| u_2^{(j)} \right\|_{L_{\xi, \beta}^\infty L_x^\infty} \lesssim \varepsilon e^{-c_0 t} + \varepsilon^2 C_{u_2, T}^\infty (1+t)^{-\frac{3}{2}}, \quad \left\| u_2^{(j)} \right\|_{L_{\xi, \beta}^\infty L_x^2} \lesssim \varepsilon e^{-c_0 t} + \varepsilon^2 C_{u_2, T}^2 (1+t)^{-\frac{3}{2}},$$

for all integers $j \geq 0$.

Lemma 12. *For $0 \leq t \leq T$,*

$$\left\| u_2^{(6)} \right\|_{L_\xi^2 H_x^2} \lesssim \varepsilon e^{-c_0 t} + \varepsilon^2 C_{u_2, T}^2 (1+t)^{-\frac{3}{2}}.$$

Proposition 13. *(the H_x^2 regularization estimate for $\mathcal{R}_w^{(6)}$) Let $0 < \delta \ll 1$. Then for $M = \mathbf{c} + 25\delta$ and large $\ell > 0$, the corresponding weighted remainder $\mathcal{R}_w^{(6)}$ satisfies*

$$\left\| \mathcal{R}_w^{(6)} \right\|_{L_\xi^2 H_x^2} \leq C_\delta (\varepsilon + \varepsilon^2 C_{u_2, T}^2)$$

for some positive constant C_δ , $C_\delta \rightarrow \infty$ as $\delta \rightarrow 0$.

Proof. Note that

$$w^{-1} (\partial_t w + \xi \cdot \nabla_x w) = -\frac{M}{\ell} + \frac{\xi \cdot x}{\ell \langle x \rangle},$$

$$\left| \partial_{x_i} [w^{-1} (\partial_t w + \xi \cdot \nabla_x w)] \right| \leq \frac{2|\xi|}{\ell \langle x \rangle}, \quad \left| \partial_{x_i x_j}^2 [w^{-1} (\partial_t w + \xi \cdot \nabla_x w)] \right| \leq \frac{6|\xi|}{\ell \langle x \rangle^2}$$

for $1 \leq i, j \leq 3$. Let $0 < \delta \ll 1$. Consider the quantity

$$B(t) := \left\| \mathcal{R}_w^{(6)} \right\|_{L_\xi^2 L_x^2}^2 + \frac{\delta^2}{4\mathfrak{D}^4} \sum_{i=1}^3 \left\| \partial_{x_i} \mathcal{R}_w^{(6)} \right\|_{L_\xi^2 L_x^2}^2 + \left(\frac{\delta^2}{4\mathfrak{D}^4} \right)^2 \sum_{i,j=1}^3 \left\| \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right\|_{L_\xi^2 L_x^2}^2,$$

where the constant $\mathfrak{D} = \left(\int_{\mathbb{R}^3} \sum_{j=0}^4 |\xi| \chi_j^2 d\xi \right)^{1/2} > 1$. The direct computation gives

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \left\| \mathcal{R}_w^{(6)} \right\|_{L_\xi^2 L_x^2}^2 &= \int_{\mathbb{R}^3} \left\langle \left(-\frac{M}{\ell} + \frac{\xi \cdot x}{\ell \langle x \rangle} \right) \left(P_0 \mathcal{R}_w^{(6)} + P_1 \mathcal{R}_w^{(6)} \right), \left(P_0 \mathcal{R}_w^{(6)} + P_1 \mathcal{R}_w^{(6)} \right) \right\rangle_\xi dx \\ &\quad + \int_{\mathbb{R}^n} \left\langle L \mathcal{R}_w^{(6)}, \mathcal{R}_w^{(6)} \right\rangle_\xi + \int_{\mathbb{R}^3} \left\langle K u_2^{(6)}, \mathcal{R}_w^{(6)} \right\rangle_\xi dx, \\ \frac{1}{2} \frac{d}{dt} \left\| \partial_{x_i} \mathcal{R}_w^{(6)} \right\|_{L_\xi^2 L_x^2}^2 &= \int_{\mathbb{R}^3} \left\langle \left(-\frac{M}{\ell} + \frac{\xi \cdot x}{\ell \langle x \rangle} \right) \left(P_0 \partial_{x_i} \mathcal{R}_w^{(6)} + P_1 \partial_{x_i} \mathcal{R}_w^{(6)} \right), \left(P_0 \partial_{x_i} \mathcal{R}_w^{(6)} + P_1 \partial_{x_i} \mathcal{R}_w^{(6)} \right) \right\rangle_\xi dx \\ &\quad + \int_{\mathbb{R}^3} \left\langle L \left(\partial_{x_i} \mathcal{R}_w^{(6)} \right), \partial_{x_i} \mathcal{R}_w^{(6)} \right\rangle_\xi dx \\ &\quad + \int_{\mathbb{R}^3} \left\langle \mathcal{R}_w^{(6)} \partial_{x_i} [w^{-1} (\partial_t w + \xi \cdot \nabla_x w)], \partial_{x_i} \mathcal{R}_w^{(6)} \right\rangle_\xi dx + \int_{\mathbb{R}^3} \left\langle K \partial_{x_i} u_2^{(6)}, \partial_{x_i} \mathcal{R}_w^{(6)} \right\rangle_\xi dx, \\ \frac{1}{2} \frac{d}{dt} \left\| \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right\|_{L_\xi^2 L_x^2}^2 &= \int_{\mathbb{R}^3} \left\langle \left(-\frac{M}{\ell} + \frac{\xi \cdot x}{\ell \langle x \rangle} \right) \left(P_0 \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} + P_1 \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right), \left(P_0 \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} + P_1 \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right) \right\rangle_\xi dx \\ &\quad + \int_{\mathbb{R}^3} \left\langle L \left(\partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right), \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right\rangle_\xi + \int_{\mathbb{R}^3} \left\langle \partial_{x_i} \mathcal{R}_w^{(6)} \cdot \partial_{x_j} [w^{-1} (\partial_t w + \xi \cdot \nabla_x w)], \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right\rangle_\xi dx \end{aligned}$$

$$\begin{aligned}
& + \int_{\mathbb{R}^3} \left\langle \partial_{x_j} \mathcal{R}_w^{(6)} \cdot \partial_{x_i} [w^{-1} (\partial_t w + \xi \cdot \nabla_x w)], \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right\rangle_{\xi} dx \\
& + \int_{\mathbb{R}^3} \left\langle \mathcal{R}_w^{(6)} \partial_{x_i x_j}^2 [w^{-1} (\partial_t w + \xi \cdot \nabla_x w)], \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right\rangle_{\xi} dx \\
& + \int_{\mathbb{R}^3} \left\langle K \left(\partial_{x_i x_j}^2 u_2^{(6)} \right), \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right\rangle_{\xi} dx.
\end{aligned}$$

Using the fact that in (6)

$$P_0 \left(\frac{x \cdot \xi}{|x|} \right) P_0 g = \lambda_1 \langle g, E_1 \rangle_{\xi} E_1 + \lambda_2 \langle g, E_2 \rangle_{\xi} E_2, \quad x \neq 0,$$

where $\lambda_1 = -\lambda_2 = \mathbf{c}$, we have

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} B(t) \\
= & -\frac{1}{\ell} \left(M - \mathbf{c} - \delta - 3\delta - \frac{81\delta^3}{4\mathfrak{D}^4} \right) \left\| P_0 \mathcal{R}_w^{(6)} \right\|_{L_{\xi}^2 L_x^2}^2 - \left(\nu_0 - \frac{1}{\ell} - \frac{\mathfrak{D}^2}{4\ell\delta} - \frac{3\delta}{\ell\mathfrak{D}^2} - \frac{81\delta^3}{4\ell\mathfrak{D}^6} \right) \left\| P_1 \mathcal{R}_w^{(6)} \right\|_{L_{\sigma}^2 L_x^2}^2 \\
& + \frac{\delta^2}{4\mathfrak{D}^4} \sum_{i=1}^3 \left\{ -\frac{1}{\ell} (M - \mathbf{c} - \delta - 6\delta) \left\| P_0 \partial_{x_i} \mathcal{R}_w^{(6)} \right\|_{L_{\xi}^2 L_x^2}^2 - \left(\nu_0 - \frac{1}{\ell} - \frac{\mathfrak{D}^2}{4\ell\delta} - \frac{6\delta^3}{4\ell\mathfrak{D}^6} \right) \left\| P_1 \partial_{x_i} \mathcal{R}_w^{(6)} \right\|_{L_{\sigma}^2 L_x^2}^2 \right\} \\
& + \left(\frac{\delta^2}{4\mathfrak{D}^4} \right)^2 \sum_{i,j=1}^3 \left\{ -\frac{1}{\ell} (M - \mathbf{c} - 4\delta) \left\| P_0 \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right\|_{L_{\xi}^2 L_x^2}^2 - \left(\nu_0 - \frac{1}{\ell} - \frac{\mathfrak{D}^2}{4\ell\delta} - \frac{3\delta}{\ell\mathfrak{D}^2} \right) \left\| P_1 \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right\|_{L_{\sigma}^2 L_x^2}^2 \right\} \\
& + C \left\| u_2^{(6)} \right\|_{L_{\xi}^2 L_x^2} \left\| \mathcal{R}_w^{(6)} \right\|_{L_{\xi}^2 L_x^2} + \frac{\delta^2}{4\mathfrak{D}^4} \sum_{i=1}^3 \left[C \left\| \partial_{x_i} u_2^{(6)} \right\|_{L_{\xi}^2 L_x^2} \left\| \partial_{x_i} \mathcal{R}_w^{(6)} \right\|_{L_{\xi}^2 L_x^2} \right] \\
& + \left(\frac{\delta^2}{4\mathfrak{D}^4} \right)^2 \sum_{i,j=1}^3 \left[C \left\| \partial_{x_i x_j}^2 u_2^{(6)} \right\|_{L_{\xi}^2 L_x^2} \left\| \partial_{x_i x_j}^2 \mathcal{R}_w^{(6)} \right\|_{L_{\xi}^2 L_x^2} \right] \\
\leq & 3C \left[\left\| u_2^{(6)} \right\|_{L_{\xi}^2 L_x^2}^2 + \frac{\delta^2}{4\mathfrak{D}^4} \sum_{i=1}^3 \left\| \partial_{x_i} u_2^{(6)} \right\|_{L_{\xi}^2 L_x^2}^2 + \left(\frac{\delta^2}{4\mathfrak{D}^4} \right)^2 \sum_{i,j=1}^3 \left\| \partial_{x_i x_j}^2 u_2^{(6)} \right\|_{L_{\xi}^2 L_x^2}^2 \right]^{1/2} \sqrt{B(t)} \\
\leq & 3C \left\| u_2^{(6)} \right\|_{L_{\xi}^2 H_x^2} \sqrt{B(t)}
\end{aligned}$$

if we choose $M = \mathbf{c} + 25\delta$, and then choose $\ell > 0$ sufficiently large such that $\nu_0 - \frac{166}{\ell} - \frac{\mathfrak{D}^2}{4\ell\delta} > 0$. Under this choice, we have

$$\sqrt{B(t)} \leq \int_0^t 3C \left\| u_2^{(6)} \right\|_{L_{\xi}^2 H_x^2} d\tau \lesssim \varepsilon + \varepsilon^2 C_{u_2, T}^2$$

by Lemma 12, so that

$$\left\| \mathcal{R}_w^{(6)} \right\|_{L_{\xi}^2 H_x^2} \leq C_{\delta} (\varepsilon + \varepsilon^2 C_{u_2, T}^2),$$

for some positive constant C_{δ} , $C_{\delta} \rightarrow \infty$ as $\delta \rightarrow 0$. The proof of this proposition is complete. \square

Therefore, the Sobolev inequality implies that

$$\left\| \mathcal{R}_w^{(6)} \right\|_{L_{\xi}^2 L_x^{\infty}} \lesssim C_{\delta} (\varepsilon + \varepsilon^2 C_{u_2, T}^2).$$

Combining this with Lemma 11, we have

$$\begin{aligned}
\|u_2\|_{L_{\xi}^2 L_x^{\infty}} & \leq \|W^{(6)}\|_{L_{\xi}^2 L_x^{\infty}} + \left\| \mathcal{R}_w^{(6)} \right\|_{L_{\xi}^2 L_x^{\infty}} \\
& \lesssim \|W^{(6)}\|_{L_{\xi, \beta}^{\infty} L_x^{\infty}} + \left\| \mathcal{R}_w^{(6)} \right\|_{L_{\xi}^2 L_x^{\infty}} \\
& \lesssim \varepsilon e^{-c_0 t} + \varepsilon^2 C_{u_2, T}^{\infty} (1+t)^{-\frac{3}{2}} + C_{\delta} (\varepsilon + \varepsilon^2 C_{u_2, T}^2).
\end{aligned}$$

According to the wave-remainder decomposition, $u_2 = W^{(7)} + \mathcal{R}_w^{(7)}$ and

$$\mathcal{R}_w^{(7)} = \int_0^t \exp\left(-\int_\tau^t \tilde{\nu}(r, x - (t-r)\xi, \xi) dr\right) \left(K\mathcal{R}_w^{(6)}\right)(\tau) d\tau.$$

Since

$$\left\|\mathcal{R}_w^{(7)}\right\|_{L_\xi^\infty L_x^\infty} \lesssim \int_0^t e^{-\frac{3}{4}\nu_0(t-\tau)} \left\|\mathcal{R}_w^{(6)}\right\|_{L_\xi^2 L_x^\infty}(\tau) d\tau,$$

we get

$$\|u_2\|_{L_\xi^\infty L_x^\infty} \lesssim \varepsilon e^{-c_0 t} + \varepsilon^2 C_{u_2, T}^\infty (1+t)^{-\frac{3}{2}} + C_\delta (\varepsilon + \varepsilon^2 C_{u_2, T}^2).$$

By finite steps of bootstrap argument, we obtain the desired $L_{\xi, \beta}^\infty L_x^\infty$ estimate for u_2 . Similarly, by Lemma 11, Proposition 13, and the bootstrap argument, we obtain $L_{\xi, \beta}^\infty L_x^2$ estimate for u_2 as well. We summarize the estimates for u_2 as below.

Proposition 14. *Let $\beta > 4$ be large enough and let $0 < \delta \ll 1$. Then for $M = \mathbf{c} + 25\delta$ and large $\ell > 0$, the corresponding u_2 satisfies*

$$\begin{aligned} C_{u_2, T}^2 &\leq C_1 [1 + \varepsilon C_{u_2, T}^2 + C_\delta (1 + \varepsilon C_{u_2, T}^2)] \\ C_{u_2, T}^\infty &\leq C_2 [1 + \varepsilon C_{u_2, T}^\infty + C_\delta (1 + \varepsilon C_{u_2, T}^2)] \end{aligned}$$

for some positive constants C_1 and C_2 dependent on $\|f_0\|_{L_{\xi, \beta}^\infty(L_x^\infty \cap L_x^2)}$ but independent of T and δ .

Now, we are ready to prove Theorem 10. For any fixed $0 < \delta \ll 1$, we take $M = \mathbf{c} + 25\delta$, and consider the weight function

$$w(x, t) = \exp\left(\frac{\langle x \rangle - Mt}{\ell}\right),$$

with $\ell > 0$ being chosen large. In view of Proposition 14, choosing $\varepsilon > 0$ sufficiently small gives

$$|u_2|_{L_{\xi, \beta}^\infty} \leq CC_\delta \varepsilon,$$

for some positive constant C dependent on $\|f_0\|_{L_{\xi, \beta}^\infty(L_x^\infty \cap L_x^2)}$.

Note that $|x| > (M + \delta)t = (\mathbf{c} + 26\delta)t$, we have

$$\begin{aligned} \langle x \rangle - Mt &= \frac{\frac{\delta}{2M} \langle x \rangle}{2 + \frac{\delta}{2M}} + \frac{2 \langle x \rangle}{2 + \frac{\delta}{2M}} - Mt \\ &> \frac{\frac{\delta}{2M} \langle x \rangle}{2 + \frac{\delta}{2M}} + \frac{\frac{\delta}{2} t}{2 + \frac{\delta}{2M}} \geq \frac{\delta (\langle x \rangle + t)}{4M + \delta}. \end{aligned}$$

Then for $|x| > (\mathbf{c} + 26\delta)t$,

$$e^{\frac{\delta(\langle x \rangle + t)}{(4M + \delta)\ell}} |f_2|_{L_{\xi, \beta}^\infty} \leq |u_2|_{L_{\xi, \beta}^\infty} \leq CC_\delta \varepsilon$$

which implies that

$$|f_2|_{L_{\xi, \beta}^\infty} \leq CC_\delta \varepsilon e^{-\frac{\delta(\langle x \rangle + t)}{(4M + \delta)\ell}}.$$

The proof of Theorem 10 is complete.

5. THE TRANSITION FROM POLYNOMIAL TAIL TO GAUSSIAN TAIL

In this section, we study the behavior of f_1 in the microscopic variable as t increases. We provide a quantitative description of how the velocity variable transitions from a polynomial tail to a Gaussian tail. The result is as follows:

Theorem 15. *Let $\beta > 4$, $R > 0$ be sufficiently large and $0 < \kappa < \min\{1/4, \nu_0/2\}$. Assume that f_0 satisfies the same condition as in Theorem 9. If $\varepsilon > 0$ is sufficiently small, then there exists a constant $\bar{C}_\beta > 0$ only depending on β such that*

$$|f_1(t, x, \xi)| \leq \bar{C}_\beta \varepsilon \langle \xi \rangle^{-\beta} e^{-\kappa \rho(\xi, t)} \|f_0(x, \xi)\|_{L_{\xi, \beta}^\infty L_x^\infty}$$

for all $t \geq 0$, $x \in \mathbb{R}^3$, $\xi \in \mathbb{R}^3$, where $\rho(\xi, t) = \langle \xi \rangle (\langle \xi \rangle \wedge t)$. Consequently, for any fixed $t > 0$, then

$$|\mathbf{1}_{\{\langle \xi \rangle \leq t\}} f_1| \lesssim \varepsilon \langle \xi \rangle^{-\beta} e^{-\kappa \langle \xi \rangle^2}$$

and

$$|\mathbf{1}_{\{\langle \xi \rangle > t\}} f_1| \lesssim \varepsilon \langle \xi \rangle^{-\beta} e^{-\kappa t^2}.$$

Firstly, we need some estimate for the weight function $e^{\kappa \rho(\xi, t)}$. For simplicity, we define

$$(12) \quad \rho(\xi, t) = \bar{\rho}(|\xi|, t), \quad \xi \in \mathbb{R}^3, \quad t \geq 0,$$

with $\bar{\rho}(z, t) = (1 + z^2)^{1/2} \left((1 + z^2)^{1/2} \wedge t \right)$, $t \geq 0$. Now, we give an inequality regarding the function $\bar{\rho}$.

Lemma 16. *Let $\bar{\rho}(z, t)$ be a function defined by*

$$\bar{\rho}(z, t) = (1 + z^2)^{1/2} \left((1 + z^2)^{1/2} \wedge t \right)$$

for $z \in \mathbb{R}$ and $t \geq 0$. Then

$$\bar{\rho}(a, t) + \bar{\rho}(b, t) \geq \bar{\rho}\left(\sqrt{a^2 + b^2}, t\right)$$

for all $a, b \geq 0$.

Proof. If either $a = 0$ or $b = 0$, it is trivial. By symmetry, we may assume that $b > a > 0$ and so $\sqrt{a^2 + b^2} > b > a > 0$. In the following we discuss the inequality in four cases.

Case 1: $t \geq \sqrt{a^2 + b^2} > b > a > 0$. Then

$$\bar{\rho}(a, t) + \bar{\rho}(b, t) = 2 + a^2 + b^2 = 1 + \bar{\rho}\left(\sqrt{a^2 + b^2}, t\right) \geq \bar{\rho}\left(\sqrt{a^2 + b^2}, t\right).$$

Case 2: $\sqrt{a^2 + b^2} > b > a > t$. Then

$$\begin{aligned} \bar{\rho}\left(\sqrt{a^2 + b^2}, t\right) &= (1 + a^2 + b^2)^{1/2} t \leq (1 + a^2 + 1 + b^2)^{1/2} t \\ &\leq (1 + a^2)^{1/2} t + (1 + b^2)^{1/2} t = \bar{\rho}(a, t) + \bar{\rho}(b, t). \end{aligned}$$

Case 3: $\sqrt{a^2 + b^2} > t > b > a$. Then

$$\bar{\rho}(a, t) + \bar{\rho}(b, t) = 2 + a^2 + b^2 \geq (1 + a^2 + b^2)^{1/2} t = \bar{\rho}\left(\sqrt{a^2 + b^2}, t\right).$$

Case 4: $\sqrt{a^2 + b^2} > b > t > a$. Then

$$\begin{aligned} &[\bar{\rho}(a, t) + \bar{\rho}(b, t)]^2 - \left[\bar{\rho}\left(\sqrt{a^2 + b^2}, t\right)\right]^2 \\ &= \left[1 + a^2 + (1 + b^2)^{1/2} t\right]^2 - (1 + a^2 + b^2) t^2 \\ &= (1 + a^2)^2 + 2(1 + a^2)(1 + b^2)^{1/2} t - a^2 t^2 \geq 2(1 + a^2) t^2 - a^2 t^2 \geq 0. \end{aligned}$$

As $a = b > 0$, it is a consequence of Case 1-Case 3. Gathering all the cases, the proof is complete. \square

According to this lemma, we can prove the following weighted estimate regarding Q and \mathcal{K} .

Lemma 17. *Let $\beta > 4$, $\kappa > 0$ and let $\rho(\xi, t)$ be defined by (12). Then there exists a constant $C_\beta > 0$ depending only on β such that*

$$(13) \quad \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |(\xi - \xi_*) \cdot n| \frac{\langle \xi \rangle^\beta}{\langle \xi_* \rangle^\beta \langle \xi' \rangle^\beta} \frac{e^{\kappa \rho(\xi, t)}}{e^{\kappa \rho(\xi_*, t)} e^{\kappa \rho(\xi', t)}} dnd\xi_* \leq \left(\frac{C}{\beta} + \frac{C_\beta}{\langle \xi \rangle^2} \right) \nu(\xi).$$

Moreover, if $0 < \kappa < \frac{1}{4}$, we have

$$(14) \quad \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |(\xi - \xi_*) \cdot n| \frac{\langle \xi \rangle^\beta}{\langle \xi_* \rangle^\beta} e^{-\frac{1}{4}|\xi'|^2} \frac{e^{\kappa \rho(\xi, t)}}{e^{\kappa \rho(\xi_*, t)} e^{\kappa \rho(\xi', t)}} dnd\xi_* \leq \left(\frac{C}{\beta} + \frac{C_\beta}{\langle \xi \rangle^2} \right) \nu(\xi).$$

Proof. Let $\eta = \xi' - \xi$ and $\omega = \xi_* - \xi'$. Then $\eta \perp \omega$, $\xi' = \xi + \eta$, $\xi_* = \xi + \eta + \omega$, $\xi'_* = \xi + \omega$. By change of variables,

$$\begin{aligned} & \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |(\xi - \xi_*) \cdot n| \frac{\langle \xi \rangle^\beta}{\langle \xi'_* \rangle^\beta \langle \xi' \rangle^\beta} \frac{e^{\kappa\rho(\xi,t)}}{e^{\kappa\rho(\xi'_*,t)} e^{\kappa\rho(\xi',t)}} dnd\xi_* \\ & \leq \int_{\mathbb{R}^3} \int_{\{\omega : \eta \perp \omega\}} \frac{1}{|\eta|} \frac{\langle \xi \rangle^\beta}{\langle \xi + \omega \rangle^\beta \langle \xi + \eta \rangle^\beta} \frac{e^{\kappa\rho(\xi,t)}}{e^{\kappa\rho(\xi+\omega,t)} e^{\kappa\rho(\xi+\eta,t)}} d\omega d\eta =: \text{I}. \end{aligned}$$

We split ξ into two parts

$$\xi = \xi_{\parallel} + \xi_{\perp}, \quad \xi_{\perp} = \frac{(\xi \cdot \eta) \eta}{|\eta|^2}, \quad \xi_{\parallel} \parallel \omega, \quad \xi_{\perp} \perp \omega, \quad |\xi|^2 = |\xi_{\parallel}|^2 + |\xi_{\perp}|^2.$$

Then

$$\begin{aligned} \text{I} &= \int_{\mathbb{R}^3} \frac{\langle \xi \rangle^\beta e^{\kappa[\rho(\xi,t) - \rho(\xi+\eta,t)]}}{\langle \xi + \eta \rangle^\beta |\eta|} \int_{\{\omega : \eta \perp \omega\}} \frac{e^{-\kappa(1+|\xi_{\parallel}+\omega|^2+|\xi_{\perp}|^2)^{1/2} \left((1+|\xi_{\parallel}+\omega|^2+|\xi_{\perp}|^2)^{1/2} \wedge t \right)}}{\left(1 + |\xi_{\parallel} + \omega|^2 + |\xi_{\perp}|^2\right)^{\beta/2}} d\omega d\eta \\ &= \int_{\mathbb{R}^3} \frac{\langle \xi \rangle^\beta e^{\kappa[\rho(\xi,t) - \rho(\xi+\eta,t)]}}{\langle \xi + \eta \rangle^\beta |\eta|} \int_{\mathbb{R}^2} \frac{e^{-\kappa(1+|\varpi|^2+|\xi_{\perp}|^2)^{1/2} \left[(1+|\varpi|^2+|\xi_{\perp}|^2)^{1/2} \wedge t \right]}}{\left(1 + |\varpi|^2 + |\xi_{\perp}|^2\right)^{\beta/2}} d\varpi d\eta. \end{aligned}$$

Making a change of variable by $\varpi = \sqrt{1 + |\xi_{\perp}|^2} z$ gives

$$\begin{aligned} & \int_{\mathbb{R}^2} \frac{e^{-\kappa(1+|\varpi|^2+|\xi_{\perp}|^2)^{1/2} \left[(1+|\varpi|^2+|\xi_{\perp}|^2)^{1/2} \wedge t \right]}}{\left(1 + |\varpi|^2 + |\xi_{\perp}|^2\right)^{\beta/2}} d\varpi \\ &= \left(1 + |\xi_{\perp}|^2\right)^{-\frac{\beta-2}{2}} \int_{\mathbb{R}^2} \frac{e^{-\kappa(1+|\xi_{\perp}|^2)^{1/2} (1+|z|^2)^{1/2} \left[(1+|\xi_{\perp}|^2)^{1/2} (1+|z|^2)^{1/2} \wedge t \right]}}{\left(1 + |z|^2\right)^{\beta/2}} dz \\ &= \frac{2\pi}{\langle \xi_{\perp} \rangle^{\beta-2}} \int_1^\infty \frac{e^{-\kappa \langle \xi_{\perp} \rangle \zeta \left[\langle \xi_{\perp} \rangle \zeta \wedge t \right]}}{\zeta^{\beta-1}} d\zeta. \end{aligned}$$

If $\langle \xi_{\perp} \rangle \geq t$, then

$$\begin{aligned} \frac{2\pi}{\langle \xi_{\perp} \rangle^{\beta-2}} \int_1^\infty \frac{e^{-\kappa \langle \xi_{\perp} \rangle \zeta \left[\langle \xi_{\perp} \rangle \zeta \wedge t \right]}}{\zeta^{\beta-1}} d\zeta &= \frac{2\pi}{\langle \xi_{\perp} \rangle^{\beta-2}} \int_1^\infty \frac{e^{-\kappa \langle \xi_{\perp} \rangle t \zeta}}{\zeta^{\beta-1}} d\zeta \\ &\leq \frac{2\pi}{\beta-2} \frac{e^{-\kappa \langle \xi_{\perp} \rangle t}}{\langle \xi_{\perp} \rangle^{\beta-2}} = \frac{2\pi}{\beta-2} \frac{e^{-\kappa\rho(\xi_{\perp},t)}}{\langle \xi_{\perp} \rangle^{\beta-2}}. \end{aligned}$$

If $\langle \xi_{\perp} \rangle < t$, then $\langle \xi_{\perp} \rangle \zeta < t$ for $\zeta < t / \langle \xi_{\perp} \rangle$ and thus

$$\begin{aligned} \frac{2\pi}{\langle \xi_{\perp} \rangle^{\beta-2}} \int_1^\infty \frac{e^{-\kappa \langle \xi_{\perp} \rangle \zeta \left[\langle \xi_{\perp} \rangle \zeta \wedge t \right]}}{\zeta^{\beta-1}} d\zeta &= \frac{2\pi}{\langle \xi_{\perp} \rangle^{\beta-2}} \left[\int_1^{\frac{t}{\langle \xi_{\perp} \rangle}} \frac{e^{-\kappa \langle \xi_{\perp} \rangle^2 \zeta^2}}{\zeta^{\beta-1}} d\zeta + \int_{\frac{t}{\langle \xi_{\perp} \rangle}}^\infty \frac{e^{-\kappa \langle \xi_{\perp} \rangle \zeta t}}{\zeta^{\beta-1}} d\zeta \right] \\ &\leq \frac{2\pi}{\beta-2} \frac{1}{\langle \xi_{\perp} \rangle^{\beta-2}} \left(e^{-\kappa \langle \xi_{\perp} \rangle^2} + e^{-\kappa t^2} \right) \leq \frac{4\pi}{\beta-2} \frac{e^{-\kappa\rho(\xi_{\perp},t)}}{\langle \xi_{\perp} \rangle^{\beta-2}}. \end{aligned}$$

Therefore,

$$\text{I} \leq \frac{4\pi}{\beta-2} \int_{\mathbb{R}^3} \frac{\langle \xi \rangle^\beta}{\langle \xi + \eta \rangle^\beta \langle \xi_{\perp} \rangle^{\beta-2}} \frac{e^{\kappa[\rho(\xi,t) - \rho(\xi+\eta,t) - \rho(\xi_{\perp},t)]}}{|\eta|} d\eta.$$

Observe that $|\xi_{\perp}|^2 + |\xi + \eta|^2 = |\xi_{\perp}|^2 + |\xi|^2 + 2\xi \cdot \eta + |\eta|^2 = |\xi_{\perp}|^2 + |\xi|^2 + 2\xi_{\perp} \cdot \eta + |\eta|^2 = |\xi|^2 + |\xi_{\perp} + \eta|^2$. Hence,

$$\rho(\xi, t) - \rho(\xi + \eta, t) - \rho(\xi_{\perp}, t) = \bar{\rho}(|\xi|, t) - \bar{\rho}(|\xi + \eta|, t) - \bar{\rho}(|\xi_{\perp}|, t)$$

$$\leq \bar{\rho} \left(\sqrt{|\xi_{\perp}|^2 + |\xi + \eta|^2}, t \right) - \bar{\rho}(|\xi + \eta|, t) - \bar{\rho}(|\xi_{\perp}|, t) \leq 0,$$

the last inequality being valid due to Lemma 16. It follows

$$\text{I} \leq \frac{4\pi}{\beta - 2} \int_{\mathbb{R}^3} \frac{1}{|\eta|} \frac{\langle \xi \rangle^{\beta}}{(\xi + \eta)^{\beta} \langle \xi_{\perp} \rangle^{\beta-2}} d\eta = \frac{4\pi}{\beta - 2} \int_{\mathbb{R}^3} \frac{1}{|\xi' - \xi|} \frac{\langle \xi \rangle^{\beta}}{\langle \xi' \rangle^{\beta} \langle \xi_{\perp} \rangle^{\beta-2}} d\xi'.$$

According to the estimate in the proof of [5, Lemma 2.11], there exists a constant $C_{\beta} > 0$ depending only on β such that

$$\text{I} \leq \frac{4\pi}{\beta - 2} \int_{\mathbb{R}^3} \frac{1}{|\xi' - \xi|} \frac{\langle \xi \rangle^{\beta}}{\langle \xi' \rangle^{\beta} \langle \xi_{\perp} \rangle^{\beta-2}} d\xi' \leq \left(\frac{C}{\beta} + \frac{C_{\beta}}{\langle \xi \rangle^2} \right) \nu(\xi),$$

where $C > 0$ is a universal constant. This completes the proof of (13).

For the estimate of (14), applying the same argument as (13), one gets

$$\begin{aligned} & \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |(\xi - \xi_*) \cdot n| \frac{\langle \xi \rangle^{\beta}}{\langle \xi_* \rangle^{\beta}} e^{-\frac{1}{4}|\xi|^2} \frac{e^{\kappa\rho(\xi,t)}}{e^{\kappa\rho(\xi_*,t)} e^{\kappa\rho(\xi',t)}} dnd\xi_* \\ & \leq \frac{4\pi}{\beta - 2} \int_{\mathbb{R}^3} \frac{1}{|\xi' - \xi|} \frac{\langle \xi \rangle^{\beta}}{e^{\frac{1}{4}|\xi|^2} \langle \xi_{\perp} \rangle^{\beta-2}} d\xi'. \end{aligned}$$

Then one can modify the argument in [5, Lemma 2.11] (in fact, it is easier) to conclude our result. \square

Corollary 18. *Let $\beta > 4$, $\kappa > 0$ and let $\rho(\xi, t)$ be defined by (12). Then there exists a constant $C''_{\beta} > 0$ depending only on β such that*

$$e^{\kappa\rho(\xi,t)} \langle \xi \rangle^{\beta} |Q(g, h)| \leq C''_{\beta} \nu(\xi) \left| e^{\kappa\rho(\xi,t)} g \right|_{L_{\xi, \beta}^{\infty}} \left| e^{\kappa\rho(\xi,t)} h \right|_{L_{\xi, \beta}^{\infty}}.$$

Moreover, for any $R > 0$, and $0 < \kappa < \frac{1}{4}$, we have

$$\chi_{\{|\xi| \geq R\}} e^{\kappa\rho(\xi,t)} \langle \xi \rangle^{\beta} |\mathcal{K}f| \leq \left(\frac{C}{\beta} + \frac{C_{\beta}}{R^2} \right) \nu(\xi) \left| e^{\kappa\rho(\xi,t)} f \right|_{L_{\xi, \beta}^{\infty}}.$$

Proof. We only prove the estimate of Q since the estimate of \mathcal{K} is similar. By definition of Q ,

$$\begin{aligned} e^{\kappa\rho(\xi,t)} \langle \xi \rangle^{\beta} |Q(g, h)| & \leq \left| e^{\kappa\rho(\xi,t)} g \right|_{L_{\xi, \beta}^{\infty}} \left| e^{\kappa\rho(\xi,t)} h \right|_{L_{\xi, \beta}^{\infty}} \\ & \cdot \left[\int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |(\xi - \xi_*) \cdot n| \frac{\langle \xi \rangle^{\beta}}{\langle \xi_* \rangle^{\beta} \langle \xi' \rangle^{\beta}} \frac{e^{\kappa\rho(\xi,t)}}{e^{\kappa\rho(\xi_*,t)} e^{\kappa\rho(\xi',t)}} dnd\xi_* \right. \\ & \left. + \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |(\xi - \xi_*) \cdot n| \frac{1}{\langle \xi_* \rangle^{\beta} e^{\kappa\rho(\xi_*, \tau)}} dnd\xi_* \right]. \end{aligned}$$

In view of Lemma 17,

$$\int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |(\xi - \xi_*) \cdot n| \frac{\langle \xi \rangle^{\beta}}{\langle \xi_* \rangle^{\beta} \langle \xi' \rangle^{\beta}} \frac{e^{\kappa\rho(\xi,t)}}{e^{\kappa\rho(\xi_*,t)} e^{\kappa\rho(\xi',t)}} dnd\xi_* \leq C'_{\beta} \nu(\xi)$$

for some $C'_{\beta} > 0$. By straightforward computation,

$$\begin{aligned} & \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |(\xi - \xi_*) \cdot n| \frac{1}{\langle \xi_* \rangle^{\beta} e^{\kappa\rho(\xi_*, \tau)}} dnd\xi_* \\ & \leq 4\pi \left[\int_{|\xi_*| \leq |\xi|} |\xi - \xi_*| \frac{1}{\langle \xi_* \rangle^{\beta}} d\xi_* + \int_{|\xi_*| > |\xi|} |\xi - \xi_*| \frac{1}{\langle \xi_* \rangle^{\beta}} d\xi_* \right] \\ & \leq (4\pi)^2 \left(\frac{2|\xi|}{\beta - 3} + \frac{2}{\beta - 4} \frac{1}{\langle \xi \rangle^{\beta-4}} \right). \end{aligned}$$

Hence,

$$e^{\kappa\rho(\xi,t)} \langle \xi \rangle^\beta |Q(g, h)| \leq C''_\beta \nu(\xi) \left| e^{\kappa\rho(\xi,t)} g \right|_{L_{\xi,\beta}^\infty} \left| e^{\kappa\rho(\xi,t)} h \right|_{L_{\xi,\beta}^\infty}$$

for some constant $C''_\beta > 0$ only depending upon β , as desired. \square

Proof of Theorem 15. In virtue of (4), f_1 can be expressed as

$$f_1 = \mathbb{S}^t f_0 + \int_0^t \mathbb{S}^{t-\tau} \left[K_s f_1 + Q(f_1, f_1) + Q(f_1, \sqrt{\mathcal{M}} f_2) + Q(\sqrt{\mathcal{M}} f_2, f_1) \right] (\tau) d\tau.$$

Let $T > 0$ be any number and denote

$$u(t, x, \xi) = \langle \xi \rangle^\beta e^{\kappa\rho(\xi,t)} f_1(t, x, \xi).$$

Multiplying $\langle \xi \rangle^\beta e^{\kappa\rho(\xi,t)}$ on both sides of the integral equation gives

$$\begin{aligned} \langle \xi \rangle^\beta e^{\kappa\rho(\xi,t)} f_1 &= \varepsilon \mathbb{S}^t \left(\langle \xi \rangle^\beta e^{\kappa\rho(\xi,t)} f_0 \right) + \int_0^t \langle \xi \rangle^\beta e^{\kappa\rho(\xi,t)} \mathbb{S}^{t-\tau} Q(f_1, f_1) (\tau) d\tau \\ &\quad + \int_0^t \langle \xi \rangle^\beta e^{\kappa\rho(\xi,t)} \mathbb{S}^{t-\tau} \left[K_s f_1 + Q(f_1, \sqrt{\mathcal{M}} f_2) + Q(\sqrt{\mathcal{M}} f_2, f_1) \right] (\tau) d\tau \\ &=: \text{I} + \text{II} + \text{III}. \end{aligned}$$

At first, it is easy to see

$$(15) \quad |\text{I}| \leq \varepsilon \left\| \langle \xi \rangle^\beta e^{-\nu_0 \langle \xi \rangle t + \kappa \langle \xi \rangle t} |f_0| \right\|_{L_\xi^\infty L_x^\infty} \leq \varepsilon \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty},$$

since $\kappa\rho(\xi, t) \leq \kappa \langle \xi \rangle t$.

As for II and III, we can find

$$e^{-\nu(\xi)(t-\tau)} e^{\kappa\rho(\xi,t)} \leq e^{-(\nu_0 - \kappa)(\xi)(t-\tau)} e^{\kappa\rho(\xi,\tau)}$$

for all ξ and $0 \leq \tau \leq t$. To see this, for $\langle \xi \rangle > t$, we have $\rho(\xi, t) = \langle \xi \rangle t$ and thus

$$e^{-\nu(\xi)(t-\tau)} e^{\kappa\rho(\xi,t)} = e^{-\nu(\xi)(t-\tau)} e^{\kappa \langle \xi \rangle t} \leq e^{-(\nu_0 - \kappa)(\xi)(t-\tau)} e^{\kappa \langle \xi \rangle \tau} = e^{-(\nu_0 - \kappa)(\xi)(t-\tau)} e^{\kappa\rho(\xi,\tau)};$$

for $\langle \xi \rangle \leq t$, we have $\rho(\xi, t) = \langle \xi \rangle^2$, so that

$$\begin{aligned} e^{-\nu(\xi)(t-\tau)} e^{\kappa\rho(\xi,t)} &\leq e^{-\nu_0 \langle \xi \rangle (t-\tau)} e^{\kappa \langle \xi \rangle^2} = e^{-(\nu_0 - \kappa)(\xi)(t-\tau)} e^{\kappa \langle \xi \rangle ((\xi) - t) + \kappa \langle \xi \rangle \tau} \\ &\leq e^{-(\nu_0 - \kappa)(\xi)(t-\tau)} e^{\kappa\rho(\xi,\tau)}. \end{aligned}$$

Therefore,

$$|\text{III}| \leq \int_0^t e^{-(\nu_0 - \kappa)(\xi)(t-\tau)} e^{\kappa\rho(\xi,\tau)} \langle \xi \rangle^\beta |Q(f_1, f_1) (\tau)| d\tau,$$

$$|\text{III}| \leq \int_0^\infty e^{-(\nu_0 - \kappa)(\xi)(t-\tau)} \langle \xi \rangle^\beta e^{\kappa\rho(\xi,\tau)} \left| \left[K_s f_1 + Q(f_1, \sqrt{\mathcal{M}} f_2) + Q(\sqrt{\mathcal{M}} f_2, f_1) \right] (\tau) \right| d\tau.$$

By Corollary 18, we have

$$(16) \quad |\text{III}| \leq C''_\beta \sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty}^2 \int_0^t e^{-(\nu_0 - \kappa)(\xi)(t-\tau)} \nu(\xi) d\tau \leq \frac{C''_\beta \nu_1}{\nu_0 - \kappa} \left(\sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty} \right)^2.$$

Regarding III, in view of Theorem 9 and Corollary 18, together the fact that

$$e^{\kappa\rho(\xi,\tau)} \sqrt{\mathcal{M}} = e^{\kappa \langle \xi \rangle ((\xi) \wedge \tau)} e^{-\frac{|\xi|^2}{4}} \leq e^{\kappa \langle \xi \rangle^2 - \frac{|\xi|^2}{4}} \leq e^{1/4},$$

we have

$$\begin{aligned} &\int_0^\infty e^{-(\nu_0 - \kappa)(\xi)(t-\tau)} \langle \xi \rangle^\beta e^{\kappa\rho(\xi,\tau)} \left| Q(f_1, \sqrt{\mathcal{M}} f_2) + Q(\sqrt{\mathcal{M}} f_2, f_1) \right| d\tau \\ &\leq 2C''_\beta \sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty} \cdot \sup_{0 \leq t \leq T} \left\| e^{\kappa\rho(\xi,\tau)} \sqrt{\mathcal{M}} f_2 \right\|_{L_{\xi,\beta}^\infty L_x^\infty} \int_0^t e^{-(\nu_0 - \kappa)(\xi)(t-\tau)} \nu(\xi) d\tau \\ &\leq \left(\frac{\tilde{C}_\beta \varepsilon}{\nu_0 - \kappa} \|f_0\|_{L_{\xi,\beta}^\infty L_x^\infty} \right) \sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty}. \end{aligned}$$

By Corollary 18,

$$\begin{aligned}
& \int_0^\infty e^{-(\nu_0 - \kappa)\langle \xi \rangle (t - \tau)} \langle \xi \rangle^\beta e^{\kappa \rho(\xi, \tau)} |K_s f_1| d\tau \\
& \leq \left(\frac{C}{\beta} + \frac{C_\beta}{R^2} \right) \sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty} \int_0^\infty e^{-(\nu_0 - \kappa)\langle \xi \rangle (t - \tau)} \nu(\xi) d\tau \\
& \leq \frac{\nu_1}{\nu_0 - \kappa} \left(\frac{C}{\beta} + \frac{C_\beta}{R^2} \right) \sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty} \leq \frac{1}{4} \sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty},
\end{aligned}$$

after choosing $\beta > 4$ and $R > 0$ sufficiently large. Hence,

$$(17) \quad \text{III} \leq \left(\frac{1}{4} + \frac{\tilde{C}_\beta \varepsilon}{\nu_0 - \kappa} \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \right) \sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty}.$$

Combining (15), (16) and (17), we have

$$\begin{aligned}
\sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty} & \leq \varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} + \left(\frac{1}{4} + \frac{\tilde{C}_\beta \varepsilon}{\nu_0 - \kappa} \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} \right) \sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty} \\
& \quad + \frac{C''_\beta \nu_1}{\nu_0 - \kappa} \left(\sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty} \right)^2.
\end{aligned}$$

We may assume that $\frac{C''_\beta \nu_1}{\nu_0 - \kappa} > 1$. Choosing $\varepsilon > 0$ sufficiently small such that

$$\frac{\tilde{C}_\beta \varepsilon}{\nu_0 - \kappa} \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} < \frac{1}{4} \quad \text{and} \quad \left(\frac{4C''_\beta \nu_1}{\nu_0 - \kappa} \right)^2 \varepsilon \|f_0(x, \xi)\|_{L_{\xi, \beta}^\infty L_x^\infty} < 1,$$

we obtain

$$\sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty} \leq 2\varepsilon \|f_0\|_{L_{\xi, \beta}^\infty L_x^\infty} + \frac{2C''_\beta \nu_1}{\nu_0 - \kappa} \left(\sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty} \right)^2.$$

Since $\|u(0, x, \xi)\|_{L_\xi^\infty L_x^\infty} = \|f_1(0, x, \xi)\|_{L_{\xi, \beta}^\infty L_x^\infty} = \varepsilon \|f_0(x, \xi)\|_{L_{\xi, \beta}^\infty L_x^\infty}$,

$$\sup_{0 \leq t \leq T} \|u\|_{L_\xi^\infty L_x^\infty} \leq \left(\frac{4C''_\beta \nu_1}{\nu_0 - \kappa} \right) \varepsilon \|f_0(x, \xi)\|_{L_{\xi, \beta}^\infty L_x^\infty},$$

for any finite $T > 0$. Consequently,

$$\langle \xi \rangle^\beta e^{\kappa \rho(\xi, t)} |f_1(t, x, \xi)| = |u(t, x, \xi)| \leq \bar{C}_\beta \varepsilon \|f_0(x, \xi)\|_{L_{\xi, \beta}^\infty L_x^\infty}$$

for some constant $\bar{C}_\beta > 0$ depending only on β , i.e.,

$$|f_1(t, x, \xi)| \leq \bar{C}_\beta \varepsilon \langle \xi \rangle^{-\beta} e^{-\kappa \rho(\xi, t)} \|f_0(x, \xi)\|_{L_{\xi, \beta}^\infty L_x^\infty}$$

for all $t \geq 0$, $x \in \mathbb{R}^3$, $\xi \in \mathbb{R}^3$. □

6. SOME CONVOLUTION ESTIMATES

In this section, we will compute the interactions between different wave patterns, which are essential for determining the precise space-time structure of the solution. Although these estimates appear complicated, there is a clear physical picture behind them (see Section 6.2 for some illustrations). The proofs in fact aim to translate this heuristic picture into refined convolution estimates.

To facilitate the estimates, we decompose space-time domain into the following 5 regions:

$$\begin{aligned}
D_1 &= \{|x| \leq \sqrt{1+t}\}, \\
D_2 &= \{ct - \sqrt{1+t} \leq |x| \leq ct + \sqrt{1+t}\}, \\
D_3 &= \{|x| \geq ct + \sqrt{1+t}\}, \\
D_4 &= \left\{ \sqrt{1+t} \leq |x| \leq \frac{1}{2}ct \right\},
\end{aligned}$$

$$D_5 = \left\{ \frac{1}{2}ct \leq |x| \leq ct - \sqrt{1+t} \right\}.$$

6.1. Linear interaction.

Lemma 19 (Diffusion wave convolved with exponential decay).

$$(1+t)^{-3/2} e^{-\frac{|x|^2}{D_0(1+t)}} *_{x,t} e^{-\frac{t+|x|}{c_0}} \lesssim (1+t)^{-3/2} e^{-\frac{|x|^2}{D(1+t)}} + e^{-\frac{t+|x|}{\hat{c}}},$$

for some constants \hat{c} and $\hat{D} > 0$.

Lemma 20 (Huygens wave convolved with exponential decay).

$$\begin{aligned} A &= (1+t)^{-2} e^{-\frac{(|x|-ct)^2}{D_0(1+t)}} *_{x,t} e^{-\frac{t+|x|}{c_0}} \\ &\lesssim (1+t)^{-2} e^{-\frac{(|x|-ct)^2}{D(1+t)}} + e^{-\frac{t+|x|}{\hat{c}}}, \end{aligned}$$

for some constants \hat{c} and $\hat{D} > 0$.

Lemma 21 (Riesz wave convolved with exponential decay).

$$\begin{aligned} B &= \mathbf{1}_{\{|x| \leq ct\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} *_{x,t} e^{-\frac{t+|x|}{c_0}} \\ &\lesssim \mathbf{1}_{\{|x| < ct\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} + (1+t)^{-2} e^{-\frac{(|x|-ct)^2}{D_0(1+t)}}, \end{aligned}$$

for some constants $\hat{c}_0, \hat{D}_0 > 0$.

The proof of Lemma 19 is easy and hence we omit it.

Proof of Lemma 20 (Huygens wave convolved with exponential decay). We rewrite

$$A = \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{-2} e^{-\frac{(|y|-c\tau)^2}{D_0(1+\tau)}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau.$$

We discuss the integral A in each domain D_i ($1 \leq i \leq 5$) for which (x, t) belongs to.

Case 1: $(x, t) \in D_1$. Direct computation gives

$$\begin{aligned} A &\lesssim \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+\tau)^{-2} e^{-\frac{t}{2c_0} - \frac{|x-y|}{c_0}} dy d\tau + \int_{\frac{t}{2}}^t \int_{|y| \leq \frac{c\tau}{2}} (1+\tau)^{-2} e^{-\frac{(|y|-c\tau)^2}{D_0(1+\tau)}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau \\ &\quad + \int_{\frac{t}{2}}^t \int_{|y| > \frac{c\tau}{2}} (1+\tau)^{-2} e^{-\frac{(|y|-c\tau)^2}{D_0(1+\tau)}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau \\ &\lesssim e^{-\frac{t}{2c_0}} + e^{-\frac{c^2 t}{32D_0}} \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+\tau)^{-2} e^{-\frac{|x-y|}{c_0}} dy d\tau \\ &\quad + \int_{\frac{t}{2}}^t \int_{|y| > \frac{c\tau}{2}} (1+\tau)^{-2} e^{-\frac{(|y|-c\tau)^2}{D_0(1+\tau)}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau \\ &\lesssim e^{-\frac{t}{2c_0}} + e^{-\frac{ct}{32D_0}} + \int_{\frac{t}{2}}^t \int_{|y| > \frac{c\tau}{2}} (1+\tau)^{-2} e^{-\frac{(|y|-c\tau)^2}{D_0(1+\tau)}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau. \end{aligned}$$

Note that if $\frac{t}{2} \leq \tau \leq t$, $|y| > \frac{c\tau}{2}$, then

$$|x-y| \geq |y| - |x| \geq \frac{ct}{4} - \sqrt{1+t} \geq \frac{ct}{8} + \frac{ct}{8} - \sqrt{1+t} \geq \frac{ct}{8}$$

for $t \geq 40$, so that

$$\int_{\frac{t}{2}}^t \int_{|y| > \frac{c\tau}{2}} (1+\tau)^{-2} e^{-\frac{(|y|-c\tau)^2}{D_0(1+\tau)}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau$$

$$\begin{aligned}
&\lesssim e^{-\frac{ct}{8c_0}} \int_{\frac{t}{2}}^t \int_{|y| > \frac{ct}{2}} (1+\tau)^{-2} e^{-\frac{(|y|-c\tau)^2}{D_0(1+\tau)}} dy d\tau \\
&\lesssim e^{-\frac{ct}{8c_0}} \int_{\frac{t}{2}}^t (1+\tau)^{-2+\frac{5}{2}} d\tau \lesssim e^{-\frac{ct}{16c_0}},
\end{aligned}$$

for $t \geq 40$. Consequently,

$$A \lesssim e^{-\frac{t}{2c_0}} + e^{-\frac{ct}{32D_0}} + e^{-\frac{ct}{16c_0}} \lesssim e^{-\frac{t+|x|}{c}}$$

for all $t \geq 0$, where $\frac{1}{c} = \frac{1}{2} \min\{\frac{1}{2c_0}, \frac{c}{32D_0}, \frac{c}{16c_0}\}$.

Case 2: $(x, t) \in D_2$.

$$\begin{aligned}
A &\lesssim \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+\tau)^{-2} e^{-\frac{t}{2c_0}} e^{-\frac{|x-y|}{c_0}} dy d\tau + \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+t)^{-2} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau \\
&\lesssim e^{-\frac{t}{2c_0}} + (1+t)^{-2} \lesssim (1+t)^{-2} \lesssim (1+t)^{-2} e^{-\frac{(|x|-ct)^2}{D(1+t)}}.
\end{aligned}$$

Case 3: $(x, t) \in D_3$. We split the integral A into four parts

$$\begin{aligned}
A &= \int_0^{\frac{t}{2}} \left(\int_{|y| \leq \frac{|x|+c\tau}{2}} + \int_{|y| > \frac{|x|+c\tau}{2}} \right) (\dots) dy d\tau + \int_{\frac{t}{2}}^t \left(\int_{|y| \leq \frac{|x|+c\tau}{2}} + \int_{|y| > \frac{|x|+c\tau}{2}} \right) (\dots) dy d\tau \\
&=: A_{11} + A_{12} + A_{21} + A_{22}.
\end{aligned}$$

Note that if $|y| \leq \frac{|x|+c\tau}{2}$, then

$$|x-y| \geq |x|-|y| \geq \frac{|x|-c\tau}{2} = \frac{|x|-ct}{2} + \frac{c(t-\tau)}{2}.$$

if $|y| > \frac{|x|+c\tau}{2}$, then

$$|y|-c\tau > \frac{|x|-c\tau}{2} = \frac{|x|-ct}{2} + \frac{c(t-\tau)}{2}.$$

It immediately follows that

$$\begin{aligned}
A_{11} &\lesssim \int_0^{\frac{t}{2}} \int_{|y| \leq \frac{|x|+c\tau}{2}} (1+\tau)^{-2} e^{-\frac{|x-y|}{2c_0}} e^{-\frac{t}{2c_0}} e^{-\frac{|x-y|}{2c_0}} dy d\tau \\
&\lesssim \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+\tau)^{-2} e^{-\frac{|x-y|}{2c_0}} e^{-\frac{t}{2c_0} - \frac{1}{2c_0} \left(\frac{|x|}{2} - \frac{ct}{4}\right)} dy d\tau \lesssim e^{-\frac{t}{2c_0} - \frac{1}{2c_0} \left(\frac{|x|}{2} - \frac{ct}{4}\right)} \lesssim e^{-\frac{t}{2c_0} - \frac{|x|}{8c_0}}, \\
A_{12} &\lesssim \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+\tau)^{-2} e^{-\frac{(|x|-ct)^2}{4D_0(1+t)}} e^{-\frac{t}{2c_0}} e^{-\frac{|x-y|}{c_0}} \lesssim e^{-\frac{(|x|-ct)^2}{4D_0(1+t)}} e^{-\frac{t}{2c_0}}, \\
A_{22} &\lesssim \int_{\frac{t}{2}}^t \int_{|y| \geq \frac{|x|+c\tau}{2}} (1+t)^{-2} e^{-\frac{(|x|-ct)^2}{4D_0(1+t)}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau \lesssim (1+t)^{-2} e^{-\frac{(|x|-ct)^2}{4D_0(1+t)}}.
\end{aligned}$$

As for A_{21} ,

$$\begin{aligned}
A_{21} &\lesssim \int_{\frac{t}{2}}^t \int_{|y| \leq \frac{|x|+c\tau}{2}} (1+\tau)^{-2} e^{-\frac{(|y|-c\tau)^2}{D_0(1+\tau)}} e^{-\frac{|x|-ct}{2c_0}} dy d\tau \\
&\lesssim e^{-\frac{|x|-ct}{2c_0}} \int_{\frac{t}{2}}^t (1+\tau)^{-2+\frac{5}{2}} d\tau \lesssim (1+t)^{\frac{3}{2}} e^{-\frac{|x|-ct}{2c_0}} \lesssim (1+t)^{\frac{3}{2}} e^{-\frac{\sqrt{1+t}}{4c_0}} e^{-\frac{|x|-ct}{4c_0}} \\
&\lesssim (1+t)^{-2} e^{-\frac{|x|-ct}{4c_0}},
\end{aligned}$$

since $|x|-ct \geq \sqrt{1+t}$. Now we discuss two cases: (i) $|x|-ct \leq \frac{1}{2}(1+t)$, and (ii) $|x|-ct \geq \frac{1}{2}(1+t)$. For case (i),

$$|x|-ct \geq \frac{2(|x|-ct)^2}{1+t},$$

which follows that

$$e^{-\frac{|x|-\mathbf{c}t}{4c_0}} \lesssim e^{-\frac{(|x|-\mathbf{c}t)^2}{2c_0(1+t)}}.$$

For case (ii), it is easy to see

$$(1+t) \leq 2(|x| - \mathbf{c}t)$$

and thus

$$|x| = ||x| - \mathbf{c}t + \mathbf{c}t| \leq (|x| - \mathbf{c}t) + \mathbf{c}t \leq (1 + 2\mathbf{c})(|x| - \mathbf{c}t) < 4(|x| - \mathbf{c}t),$$

so that

$$e^{-\frac{|x|-\mathbf{c}t}{4c_0}} = e^{-\frac{|x|-\mathbf{c}t}{8c_0}} e^{-\frac{|x|-\mathbf{c}t}{8c_0}} \lesssim e^{-\frac{(1+t)}{16c_0}} e^{-\frac{|x|}{32c_0}} \lesssim e^{-\frac{t}{16c_0}} e^{-\frac{|x|}{32c_0}}.$$

Hence,

$$A_{21} \lesssim (1+t)^{-2} e^{-\frac{(|x|-\mathbf{c}t)^2}{2c_0(1+t)}} + e^{-\frac{t}{16c_0}} e^{-\frac{|x|}{32c_0}}.$$

Combining all above estimates, we have

$$A \lesssim (1+t)^{-2} e^{-\frac{(|x|-\mathbf{c}t)^2}{D(1+t)}} + e^{-\frac{t+|x|}{\hat{c}}}$$

for some constants \hat{c} and $\hat{D} > 0$.

Case 4: $(x, t) \in D_4$. We split the integral into two parts

$$A = \left(\int_0^{\frac{2}{3}t} + \int_{\frac{2}{3}t}^t \right) \int_{\mathbb{R}^3} (\dots) dy d\tau =: A_1 + A_2.$$

For A_1 ,

$$A_1 \lesssim \int_0^{\frac{2}{3}t} \int_{\mathbb{R}^3} (1+\tau)^{-2} e^{-\frac{|x-y|}{c_0}} e^{-\frac{t}{3c_0}} dy d\tau \lesssim e^{-\frac{t}{3c_0}} \lesssim e^{-\frac{t}{6c_0}} e^{-\frac{|x|}{3c_0c}}$$

since $\sqrt{1+t} \leq |x| \leq \mathbf{c}t/2$.

For A_2 , we decompose \mathbb{R}^3 into two parts

$$A_2 = \int_{\frac{2}{3}t}^t \left(\int_{|y| \leq \frac{|x|+\mathbf{c}\tau}{2}} + \int_{|y| > \frac{|x|+\mathbf{c}\tau}{2}} \right) (\dots) dy d\tau =: A_{21} + A_{22}.$$

If $\frac{2}{3}t \leq \tau \leq t$, $|y| \leq \frac{|x|+\mathbf{c}\tau}{2}$, then

$$\mathbf{c}\tau - |y| \geq \frac{\mathbf{c}\tau - |x|}{2} \geq \frac{\mathbf{c}t}{3} - \frac{\mathbf{c}t}{4} = \frac{\mathbf{c}t}{12} \geq \frac{\mathbf{c}t - |x|}{12}.$$

If $\frac{2}{3}t \leq \tau \leq t$, $|y| > \frac{|x|+\mathbf{c}\tau}{2}$, then

$$|x-y| \geq |y| - |x| \geq \frac{\mathbf{c}\tau - |x|}{2} \geq \frac{\mathbf{c}t}{12} \geq \frac{\mathbf{c}t - |x|}{12}.$$

Hence,

$$\begin{aligned} A_{21} &\lesssim \int_{\frac{2}{3}t}^t \int_{|y| \leq \frac{|x|+\mathbf{c}\tau}{2}} (1+\tau)^{-2} e^{-\frac{(|y|-\mathbf{c}\tau)^2}{D_0(1+\tau)}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau \\ &\lesssim (1+t)^{-2} e^{-\frac{(\mathbf{c}t-|x|)^2}{144D_0(1+t)}}, \end{aligned}$$

and

$$\begin{aligned} A_{22} &\lesssim \int_{\frac{2}{3}t}^t \int_{|y| > \frac{|x|+\mathbf{c}\tau}{2}} (1+t)^{-2} e^{-\frac{t-\tau}{c_0}} e^{-\frac{|x-y|}{2c_0}} e^{-\frac{\mathbf{c}t-|x|}{24c_0}} dy d\tau \\ &\lesssim (1+t)^{-2} e^{-\frac{\mathbf{c}t-|x|}{24c_0}}. \end{aligned}$$

Since $t \geq 1$ and $\mathbf{c}t - |x| > \frac{\mathbf{c}t}{2} \geq \frac{\mathbf{c}}{4}(1+t)$ for $(x, t) \in D_4$,

$$\mathbf{c}(1+t) \leq 4(\mathbf{c}t - |x|),$$

and thus

$$|x| = ||x| - \mathbf{c}t + \mathbf{c}t| \leq \mathbf{c}t - |x| + \mathbf{c}t \leq 5(\mathbf{c}t - |x|),$$

which implies that

$$e^{-\frac{ct-|x|}{24c_0}} = e^{-\frac{ct-|x|}{48c_0}} e^{-\frac{ct-|x|}{48c_0}} \lesssim e^{-\frac{c(1+t)}{192c_0}} e^{-\frac{|x|}{240c_0}} \lesssim e^{-\frac{ct}{192c_0}} e^{-\frac{|x|}{240c_0}}.$$

Therefore,

$$A_{22} \lesssim e^{-\frac{t}{96c_0}} e^{-\frac{|x|}{192c_0}}.$$

Combining this with A_1 and A_{11} , we get the desired estimate

$$A \lesssim e^{-\frac{t}{6c_0}} e^{-\frac{|x|}{3c_0c}} + (1+t)^{-2} e^{-\frac{(ct-|x|)^2}{144D_0(1+t)}} + e^{-\frac{t}{96c_0}} e^{-\frac{|x|}{192c_0}}.$$

Case 5: $(x, t) \in D_5$. We split the integral A into three parts

$$A = \left(\int_0^{\frac{t}{2}} + \int_{\frac{t}{2}}^{\frac{t}{2} + \frac{|x|}{2c}} + \int_{\frac{t}{2} + \frac{|x|}{2c}}^t \right) \int_{\mathbb{R}^3} (\dots) dyd\tau =: A_1 + A_2 + A_3.$$

It immediately follows that

$$A_1 \lesssim \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+\tau)^{-2} e^{-\frac{|x-y|}{c_0}} e^{-\frac{t}{2c_0}} dyd\tau \lesssim e^{-\frac{t}{2c_0}} \lesssim e^{-\frac{t}{4c_0} - \frac{|x|}{4c_0c}},$$

and

$$\begin{aligned} A_2 &\lesssim \int_{\frac{t}{2}}^{\frac{t}{2} + \frac{|x|}{2c}} \int_{\mathbb{R}^3} (1+\tau)^{-2} e^{-\frac{(|y|-c\tau)^2}{D_0(1+\tau)}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dyd\tau \\ &\lesssim (1+t)^{-2} \int_{\frac{t}{2}}^{\frac{t}{2} + \frac{|x|}{2c}} \int_{\mathbb{R}^3} e^{-\frac{|x-y|}{c_0}} e^{-\frac{t-\tau}{2c_0}} e^{-\frac{1}{2c_0}(\frac{t}{2} - \frac{|x|}{2c})} dyd\tau \\ &\lesssim (1+t)^{-2} e^{-\frac{ct-|x|}{4c_0c}}. \end{aligned}$$

For A_3 , we decompose \mathbb{R}^3 into two parts

$$A_3 = \int_{\frac{t}{2} + \frac{|x|}{2c}}^t \left(\int_{|y| \leq \frac{|x|+c\tau}{2}} + \int_{|y| > \frac{|x|+c\tau}{2}} \right) (\dots) dyd\tau = A_{31} + A_{32}.$$

If $\frac{t}{2} + \frac{|x|}{2c} \leq \tau \leq t$, $|y| \leq \frac{|x|+c\tau}{2}$, then

$$c\tau - |y| \geq \frac{c\tau - |x|}{2} \geq \frac{ct - |x|}{4}.$$

If $\frac{t}{2} + \frac{|x|}{2c} \leq \tau \leq t$, $|y| > \frac{|x|+c\tau}{2}$, then

$$|x-y| \geq |y| - |x| > \frac{c\tau - |x|}{2} \geq \frac{ct - |x|}{4}.$$

Hence,

$$\begin{aligned} A_3 &= A_{31} + A_{32} \\ &\lesssim \int_{\frac{t}{2} + \frac{|x|}{2c}}^t \int_{|y| \leq \frac{|x|+c\tau}{2}} (1+t)^{-2} e^{-\frac{(ct-|x|)^2}{16D_0(1+t)}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dyd\tau \\ &\quad + \int_{\frac{t}{2} + \frac{|x|}{2c}}^t \int_{|y| > \frac{|x|+c\tau}{2}} (1+t)^{-2} e^{-\frac{|x-y|}{2c_0}} e^{-\frac{t-\tau}{c_0}} e^{-\frac{ct-|x|}{8c_0}} dyd\tau \\ &\lesssim (1+t)^{-2} e^{-\frac{(ct-|x|)^2}{16D_0(1+t)}} + (1+t)^{-2} e^{-\frac{ct-|x|}{8c_0}}. \end{aligned}$$

Since $\sqrt{1+t} \leq ct - |x| \leq \frac{ct}{2} \leq \frac{c}{2}(1+t)$, we have

$$ct - |x| \geq \frac{2(ct - |x|)^2}{c(1+t)},$$

and thus

$$e^{-\frac{ct-|x|}{4c_0c}} \lesssim e^{-\frac{(ct-|x|)^2}{2c_0c^2(1+t)}}, \quad e^{-\frac{ct-|x|}{8c_0}} \lesssim e^{-\frac{(ct-|x|)^2}{4c_0c(1+t)}}.$$

Consequently,

$$A_2 \lesssim (1+t)^{-2} e^{-\frac{(ct-|x|)^2}{2c_0c^2(1+t)}}, \quad A_3 \lesssim (1+t)^{-2} e^{-\frac{(ct-|x|)^2}{16D_0(1+t)}} + (1+t)^{-2} e^{-\frac{(ct-|x|)^2}{4c_0c(1+t)}}.$$

Combining this with A_1 , we get the desired estimate. \square

Proof of Lemma 21 (*Riesz wave convolved with exponential decay*). We compute the integral for two cases: $|x| > ct$ and $|x| \leq ct$.

Case 1: $|x| > ct$. Let $0 \leq \tau \leq t$. If $|x| > ct$ and $|y| < c\tau$, then

$$|x-y| \geq |x| - |y| \geq |x| - ct + ct - |y| \geq |x| - ct \geq 0.$$

Hence,

$$\begin{aligned} & \int_0^t \int_{\mathbb{R}^3} e^{-\frac{(t-\tau)+|x-y|}{c_0}} 1_{\{|y| < c\tau\}} (1+\tau)^{-3/2} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3/2} dyd\tau \\ & \lesssim e^{-\frac{|x|-ct}{2c_0}} \int_0^t \int_{\mathbb{R}^3} e^{-\frac{t-\tau}{c_0}} e^{-\frac{|x-y|}{2c_0}} (1+\tau)^{-3/2} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3/2} dyd\tau \\ & = e^{-\frac{|x|-ct}{2c_0}} \left(\int_0^{\frac{t}{2}} + \int_{\frac{t}{2}}^t \right) \int_{\mathbb{R}^3} (\dots) dyd\tau =: B_1 + B_2. \end{aligned}$$

It immediately follows that

$$B_1 \leq e^{-\frac{t}{2c_0}} e^{-\frac{|x|-ct}{2c_0}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} e^{-\frac{|x-y|}{2c_0}} (1+\tau)^{-3/2} dyd\tau \lesssim e^{-\frac{t}{2c_0}} e^{-\frac{|x|-ct}{2c_0}}.$$

As for B_2 ,

$$\begin{aligned} B_2 & = e^{-\frac{|x|-ct}{2c_0}} \int_{\frac{t}{2}}^t \left(\int_{|y| \leq \frac{|x|}{2}} + \int_{|y| > \frac{|x|}{2}} \right) e^{-\frac{t-\tau}{c_0}} e^{-\frac{|x-y|}{2c_0}} (1+\tau)^{-3/2} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3/2} dyd\tau \\ & \leq e^{-\frac{|x|-ct}{2c_0}} e^{-\frac{|x|}{8c_0}} \int_{\frac{t}{2}}^t \int_{|y| \leq \frac{|x|}{2}} e^{-\frac{t-\tau}{c_0}} e^{-\frac{|x-y|}{4c_0}} dyd\tau \\ & \quad + (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} e^{-\frac{|x|-ct}{2c_0}} \int_{\frac{t}{2}}^t \int_{|y| > \frac{|x|}{2}} e^{-\frac{t-\tau}{c_0}} e^{-\frac{|x-y|}{2c_0}} dyd\tau \\ & \lesssim e^{-\frac{|x|-ct}{2c_0}} e^{-\frac{|x|}{8c_0}} + (1+t)^{-3} e^{-\frac{|x|-ct}{2c_0}} \lesssim (1+t)^{-3} e^{-\frac{|x|-ct}{2c_0}}. \end{aligned}$$

since $|x| > ct$.

Now, we consider the cases: $0 \leq |x| - ct \leq 2c(1+t)$ and $|x| - ct \geq 2c(1+t)$. For $0 \leq |x| - ct \leq 2c(1+t)$, we have

$$|x| - ct \geq \frac{(|x| - ct)^2}{2c(1+t)},$$

so that

$$e^{-\frac{|x|-ct}{2c_0}} \leq e^{-\frac{(|x|-ct)^2}{4c_0c(1+t)}}.$$

For $|x| - ct \geq 2c(1+t)$, it follows that

$$|x| - ct = \frac{|x|}{2} + \frac{|x|}{2} - ct \geq \frac{|x|}{2} + \frac{ct}{2} \geq \frac{|x|}{2} + \frac{t}{2},$$

and thus

$$e^{-\frac{|x|-ct}{2c_0}} \leq e^{-\frac{|x|+t}{4c_0}}.$$

Consequently,

$$\int_0^t \int_{\mathbb{R}^3} e^{-\frac{(t-\tau)+|x-y|}{c_0}} 1_{\{|y| < c\tau\}} (1+\tau)^{-3/2} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3/2} dyd\tau$$

$$\lesssim (1+t)^{-3} e^{-\frac{(|x|-ct)^2}{4c_0(1+t)}} + e^{-\frac{|x|+t}{4c_0}}.$$

Case 2: $|x| < ct$. Direct computation gives

$$\begin{aligned} B &\lesssim e^{-\frac{t}{2c_0}} \int_0^{t/2} \int_{\mathbb{R}^3} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau \\ &\quad + (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} \int_{t/2}^t \int_{|y| > \frac{|x|}{2}} e^{-\frac{(t-\tau)+|x-y|}{c_0}} dy d\tau \\ &\quad + (1+t)^{-3/2} e^{-\frac{|x|}{4c_0}} \int_{t/2}^t \int_{|y| \leq \frac{|x|}{2}} e^{-\frac{t-\tau}{c_0}} e^{-\frac{|x-y|}{2c_0}} dy d\tau \\ &\lesssim e^{-\frac{t}{2c_0}} + (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} \\ &\lesssim e^{-\frac{t+|x|}{4c_0}} + \mathbf{1}_{\{|x| < ct\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2}. \end{aligned}$$

To sum up,

$$B \lesssim \mathbf{1}_{\{|x| < ct\}} (1+t)^{-3/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3/2} + e^{-\frac{t+|x|}{c_0}} + (1+t)^{-2} e^{-\frac{(|x|-ct)^2}{D_0(1+t)}}$$

for some constants $\widehat{D}_0, \widehat{c}_0 > 0$. □

6.2. Nonlinear wave interaction.

Lemma 22 (Diffusion wave convolved with Diffusion wave).

$$\begin{aligned} &(1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} *_{x,t} (1+t)^{-3} \left(1 + \frac{|x|^2}{1+t}\right)^{-3} \\ &\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}. \end{aligned}$$

Lemma 23 (Space-time exponential decay convolved with Huygens wave).

$$\begin{aligned} &e^{-\frac{t+|x|}{c_0}} *_{x,t} (1+t)^{-4} \left(1 + \frac{(|x|-ct)^2}{1+t}\right)^{-2} \\ &\lesssim (1+t)^{-4} \left(1 + \frac{(|x|-ct)^2}{1+t}\right)^{-2}. \end{aligned}$$

Lemma 24 (Riesz wave convolved with Huygens wave).

$$\begin{aligned} I &= \mathbf{1}_{\{|x| \leq ct\}} (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} *_{x,t} (1+t)^{-4} \left(1 + \frac{(|x|-ct)^2}{1+t}\right)^{-2} \\ &\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(|x|-ct)^2}{1+t}\right)^{-1}. \end{aligned}$$

Lemma 25 (Huygens wave convolved with Huygens wave).

$$J = (1+t)^{-5/2} e^{-\frac{(|x|-ct)^2}{D_0(1+t)}} *_{x,t} (1+t)^{-4} \left(1 + \frac{(|x|-ct)^2}{1+t}\right)^{-2}$$

$$\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(|x| - ct)^2}{1+t}\right)^{-1}.$$

Lemma 26 (Huygens wave convolved with diffusion wave).

$$\begin{aligned} K &= (1+t)^{-5/2} e^{-\frac{(|x| - ct)^2}{D_0(1+t)}} *_{x,t} (1+t)^{-3} \left(1 + \frac{|x|^2}{1+t}\right)^{-3} \\ &\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(|x| - ct)^2}{1+t}\right)^{-1}. \end{aligned}$$

Lemma 27 (Diffusion wave convoluted with Huygens wave).

$$\begin{aligned} L &= (1+t)^{-2} e^{-\frac{|x|^2}{D_0(1+t)}} *_{x,t} (1+t)^{-4} \left(1 + \frac{(|x| - ct)^2}{1+t}\right)^{-2} \\ &\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(|x| - ct)^2}{1+t}\right)^{-1}. \end{aligned}$$

We omit the proof of Lemma 22 and Lemma 23. Before we proceed to the detailed proof of Lemmas 24-27, let us present some heuristic calculations to help understand the mechanism of nonlinear wave interactions. We use two examples for illustration: the convolution of two Huygens waves (Lemma 25) and convolution of diffusion wave with a Huygens wave (Lemma 27).

Heuristic estimate for Lemma 25. Set $\mathbf{c} = 1$ for simplicity. Then

$$(18) \quad J = \int_0^t \int_{\mathbb{R}^3} (1+t-s)^{-5/2} e^{-\frac{(|x-y| - (t-s))^2}{D_0(t-s)}} (1+s)^{-4} \left(1 + \frac{(|y| - s)^2}{1+s}\right)^{-2} dy ds.$$

We can interpret $(1+t-s)^{-5/2} e^{-\frac{(|x-y| - (t-s))^2}{D_0(t-s)}}$ as a receiver located at (x, t) that can only receive signals along the wave cone concentrated on $|x - y| = (t - s)$ with thickness $\sqrt{t - s}$. Similarly, $(1+s)^{-4} \left(1 + \frac{(|y| - s)^2}{1+s}\right)^{-2}$ can be viewed as a sender located at $(0, 0)$ that sends signals along the wave cone concentrated on $|y| = s$ with thickness \sqrt{s} .




FIGURE 2. Interaction between two Huygens waves

During the interaction process, the interaction becomes strong in the following space-time region (See Figure 2(a))

$$(19) \quad E = \{(y, s) \mid ||x - y| - (t - s)| \leq O(1)\sqrt{t - s} \text{ and } ||y| - s| \leq O(1)\sqrt{s}\}.$$

Inside this region, the space decay terms in the convolution are of order $O(1)$, and the decay is mainly from time factor. The key point is the sharp estimate of the volume for the strong interaction region.

If $|x| > t$, as s increases from 0 to t , the receiving and sending wave cones are almost disjoint, so the interaction is very weak.

Now we focus the case where $O(1)\sqrt{t} < |x| < t - O(1)\sqrt{t}$. The interaction process starts at $s = (t - |x|)/2$ and ends at $s = (t + |x|)/2$ (See Figure 2(b)). To satisfy the condition in (19), one has

$$\begin{cases} ||x - y| - (t - s)| \leq O(1)\sqrt{t - s}, \\ ||y| - s| \leq O(1)\sqrt{s}. \end{cases}$$

Let $r = |y|$, θ be the angle between x and y , and set $O(1) = 1$. The above constraints are equivalent to

$$(20) \quad \begin{cases} |\sqrt{|x|^2 + r^2 - 2r|x|\cos\theta} - (t - s)| \leq \sqrt{t - s}, \\ |r - s| \leq \sqrt{s}. \end{cases}$$

The first equation of (20) implies

$$\frac{|x|^2 + r^2 - (t - s)^2 - (t - s)}{2r|x|} - \frac{(t - s)^{3/2}}{r|x|} \leq \cos\theta \leq \frac{|x|^2 + r^2 - (t - s)^2 - (t - s)}{2r|x|} + \frac{(t - s)^{3/2}}{r|x|}$$

This leads to

$$\Delta \cos\theta \approx \frac{(t - s)^{3/2}}{r|x|}.$$

For s large, $r \approx s$, $\Delta r \approx \sqrt{s}$. Then inside the strong interaction region, (18) is approximated by

$$\begin{aligned} J &\approx \int_{\frac{t-|x|}{2}}^{\frac{t+|x|}{2}} (1+t-s)^{-5/2} (1+s)^{-4} \underbrace{s^2 \sqrt{s}}_{r^2 dr} \underbrace{\frac{(t-s)^{3/2}}{s|x|}}_{\sin\theta d\theta} ds \\ &\lesssim \int_{\frac{t-|x|}{2}}^{\frac{t+|x|}{2}} (1+t-s)^{-1} (1+s)^{-5/2} ds \frac{1}{|x|} \\ &\lesssim \left(\int_{\frac{t-|x|}{2}}^{\frac{t}{2}} (1+t)^{-1} (1+s)^{-5/2} ds + \int_{\frac{t}{2}}^{\frac{t+|x|}{2}} (1+t-s)^{-1} (1+t)^{-5/2} ds \right) \frac{1}{|x|} \\ &\lesssim \left[(1+t)^{-1} (1+t-|x|)^{-3/2} + (1+t)^{-5/2} \ln \frac{t}{t-|x|} \right] \frac{1}{|x|}. \end{aligned}$$

When $\sqrt{t} \leq |x| \leq t/2$, it is bounded by

$$(1+t)^{-5/2} |x|^{-1} \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t} \right)^{-3/2}.$$

When $t/2 \leq |x| \leq t - \sqrt{t}$, it is bounded by

$$t^{-2} (1+t-|x|)^{-3/2} + (1+t)^{-5/2} \frac{t^2}{(t-|x|)^2} (1+t)^{-1} \lesssim t^{-5/2} \left(1 + \frac{(t-|x|)^2}{1+t} \right)^{-1}.$$

This is the desired estimate in Lemma 25. \square

Heuristic estimate for Lemma 26. Set $\mathbf{c} = 1$, and one has

$$K = \int_0^t \int_{\mathbb{R}^3} (1+t-s)^{-5/2} e^{-\frac{(|x-y|-(t-s))^2}{D_0(1+t-s)}} (1+s)^{-3} \left(1 + \frac{|y|^2}{1+s} \right)^{-3} dy ds.$$

Similar as heuristics argument of Lemma 25, we view $(1+t-s)^{-5/2}e^{-\frac{(|x-y|-(t-s))^2}{D_0(1+t-s)}}$ as a receiver and $(1+s)^{-3}\left(1+\frac{|y|^2}{1+s}\right)^{-3}$ as a sender, to have the following figure representing the interaction:




FIGURE 3. Interaction between Huygens and diffusion wave

For $O(1)\sqrt{t} \leq |x| \leq t - O(1)\sqrt{t}$, the interaction occurs at $s \approx t - |x|$ and its duration is of the order \sqrt{s} . So the volume of the interaction region in space-time is approximated by

$$\underbrace{s^{3/2}}_{\text{vol of space}} \cdot \underbrace{\sqrt{s}}_{\text{vol of time}}$$

The contribution of the integrand during the interaction is approximated by

$$(1+t-s)^{-5/2}(1+s)^{-3} \Big|_{s \approx t-|x|}.$$

So we conclude

$$\begin{aligned} K &\approx (1+t-s)^{-5/2}(1+s)^{-3} s^2 \Big|_{s \approx t-|x|} \approx (1+|x|)^{-5/2}(1+t-|x|)^{-1} \\ &\lesssim \begin{cases} (1+t)^{-1}(1+|x|)^{-5/2} \lesssim (1+t)^{-2} \left(1+\frac{|x|^2}{1+t}\right)^{-3/2}, & \sqrt{t} \leq |x| \leq t/2, \\ (1+t)^{-5/2}(1+t-|x|)^{-1} \lesssim (1+t)^{-5/2} \left(1+\frac{(t-|x|)^2}{1+t}\right)^{-1}, & t/2 \leq |x| \leq t-\sqrt{t}. \end{cases} \end{aligned}$$

This yields the desired estimate in Lemma 26. \square

Other convolutions can be estimated heuristically in a similar manner. It turns out that the convolutions in Lemmas 25 and 26 are the dominated ones among all the nonlinear wave couplings.

We now begin the rigorous proofs, transforming the previous heuristic calculations into refined (and complex) convolution estimates!

Proof of Lemma 24. (Riesz wave convolved with Huygens wave).

Case 1: $(x, t) \in D_1$. Direct computation gives

$$I = \left(\int_0^{\frac{t}{2}} + \int_{\frac{t}{2}}^t \right) \int_{\mathbb{R}^3} (\dots) dy d\tau$$

$$\begin{aligned}
&\lesssim (1+t)^{-2} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+\tau)^{-4} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\quad + (1+t)^{-4} \int_{\frac{t}{2}}^t \int_{|x-y| \leq \mathbf{c}(t-\tau)} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-\frac{3}{2}} dy d\tau \\
&\lesssim (1+t)^{-2} + (1+t)^{-4} (1+t)^{3/4} \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.
\end{aligned}$$

Case 2: $(x, t) \in D_2$. We split the integral I into two parts

$$I = \left(\int_0^{\frac{t}{4}} + \int_{\frac{t}{4}}^t \right) \int_{\mathbb{R}^3} (\dots) dy d\tau =: I_1 + I_2.$$

For I_2 , one can see that

$$\begin{aligned}
I_2 &\lesssim (1+t)^{-4} \int_{\frac{t}{4}}^t \int_{\mathbb{R}^3} \mathbf{1}_{\{|x-y| \leq \mathbf{c}(t-\tau)\}} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-\frac{3}{2}} dy d\tau \\
&\lesssim (1+t)^{-4} \int_{\frac{t}{4}}^t (1+t-\tau)^{-1/4} d\tau \lesssim (1+t)^{-4} (1+t)^{3/4} \\
&\lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}.
\end{aligned}$$

For I_1 , we further decompose \mathbb{R}^3 into two parts

$$I_1 = \left(\int_0^{\frac{t}{4}} \int_{|y| \leq \frac{2}{3}|x|} + \int_0^{\frac{t}{4}} \int_{|y| > \frac{2}{3}|x|} \right) (\dots) dy d\tau = I_{11} + I_{12}.$$

If $|y| \leq \frac{2}{3}|x|$, then we have

$$|x-y| \geq |x| - |y| \geq \frac{|x|}{3},$$

and thus

$$\begin{aligned}
I_{11} &\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} \int_0^{\frac{t}{4}} \int_{\mathbb{R}^3} (1+\tau)^{-4} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} \int_0^{\frac{t}{4}} (1+\tau)^{-4} (1+\tau)^{\frac{5}{2}} d\tau \\
&\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.
\end{aligned}$$

If $|y| > \frac{2}{3}|x|$ and $0 \leq \tau \leq \frac{t}{4}$,

$$|y| - \mathbf{c}\tau \geq \frac{2}{3}|x| - \mathbf{c}\tau \geq \frac{|x|}{3} + \frac{1}{3}(\mathbf{c}t - \sqrt{1+t}) - \frac{\mathbf{c}t}{4} \geq \frac{|x|}{3} + \frac{1}{24}\mathbf{c}t$$

for $t \geq 40$. We mention here that $(x, t) \in D_2$ with $t \geq \frac{2+2\sqrt{1+\mathbf{c}^2}}{\mathbf{c}^2}$ ($\doteq 3.15$) implies that $|x| \geq \sqrt{1+t}$. Hence for $t \geq 40$,

$$\begin{aligned}
I_{12} &\lesssim (1+|x|^2)^{-\frac{3}{2}} \int_0^{\frac{t}{4}} \int_{|y| > \frac{2}{3}|x|} \mathbf{1}_{\{|x-y| \leq \mathbf{c}(t-\tau)\}} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-\frac{3}{2}} \\
&\quad \cdot (1+\tau)^{-\frac{5}{2}} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-\frac{1}{2}} dy d\tau
\end{aligned}$$

$$\begin{aligned}
&\lesssim (1+|x|^2)^{-\frac{3}{2}} \int_0^{\frac{t}{4}} \int_{\{|y|>\frac{2}{3}|x|\} \cap \{|x-y|\leq c(t-\tau)\}} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-\frac{3}{2}} \\
&\quad \cdot (1+\tau)^{-\frac{5}{2}} \left(1 + \frac{|x-y|^2}{1+t-\tau}\right)^{-\frac{1}{2}} dyd\tau \\
&\lesssim (1+|x|^2)^{-\frac{3}{2}} \int_0^{\frac{t}{4}} (1+\tau)^{-\frac{5}{2}} \int_{\mathbb{R}^3} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-2} dyd\tau \\
&\lesssim (1+t)^{-1/2} (1+|x|^2)^{-\frac{3}{2}} \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}},
\end{aligned}$$

and for $0 \leq t \leq 40$,

$$\begin{aligned}
I_{12} &\lesssim (1+t)^{-\frac{3}{2}} \int_0^{\frac{t}{4}} \int_{\mathbb{R}^3} (1+\tau)^{-4} \left(1 + \frac{(|y| - c\tau)^2}{1+\tau}\right)^{-2} dyd\tau \\
&\lesssim C \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - ct)^2}{1+t}\right)^{-1}.
\end{aligned}$$

Thereupon,

$$I_1 \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(|x| - ct)^2}{1+t}\right)^{-1},$$

and so

$$I \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(|x| - ct)^2}{1+t}\right)^{-1}.$$

Case 3: $(x, t) \in D_3$. We split the integral I into two parts

$$I = \left(\int_0^{\frac{t}{2}} + \int_{\frac{t}{2}}^t \right) \int_{\mathbb{R}^3} (\dots) dyd\tau =: I_1 + I_2.$$

For I_1 , we decompose \mathbb{R}^3 into two parts as follows:

$$I_1 = \int_0^{\frac{t}{2}} \left(\int_{|y| \leq \frac{2}{3}|x|} + \int_{|y| > \frac{2}{3}|x|} \right) (\dots) dyd\tau =: I_{11} + I_{12}.$$

If $|y| \leq \frac{2}{3}|x|$, then

$$|x-y| \geq \frac{|x|}{3},$$

and thus

$$\begin{aligned}
I_{11} &\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+\tau)^{-4} \left(1 + \frac{(|y| - c\tau)^2}{1+\tau}\right)^{-2} dyd\tau \\
&\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.
\end{aligned}$$

If $|y| > \frac{2}{3}|x|$ and $0 \leq \tau \leq \frac{t}{2}$, we have

$$|y| - c\tau \geq \frac{2}{3}|x| - \frac{ct}{2} = \frac{1}{6}|x| + \frac{1}{2}(|x| - ct) \geq \frac{1}{6}ct,$$

so that

$$\begin{aligned}
I_{12} &\lesssim (1+|x|^2)^{-\frac{3}{2}} \int_0^{\frac{t}{2}} \int_{|y|>\frac{2}{3}|x|} \mathbf{1}_{\{|x-y|\leq c(t-\tau)\}} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-\frac{3}{2}} \\
&\quad \cdot (1+\tau)^{-\frac{5}{2}} \left(1 + \frac{(|y|-c\tau)^2}{1+\tau}\right)^{-\frac{1}{2}} dyd\tau \\
&\lesssim (1+|x|^2)^{-\frac{3}{2}} \int_0^{\frac{t}{2}} \int_{\{|y|\geq\frac{2}{3}|x|\} \cap \{|x-y|\leq c(t-\tau)\}} (1+t-\tau)^{-2} (1+\tau)^{-\frac{5}{2}} \cdot \left(1 + \frac{|x-y|^2}{(1+t-\tau)}\right)^{-2} dyd\tau \\
&\lesssim (1+|x|^2)^{-\frac{3}{2}} \int_0^{\frac{t}{2}} (1+\tau)^{-\frac{5}{2}} \cdot \int_{\mathbb{R}^3} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{(1+t-\tau)}\right)^{-2} dyd\tau \\
&\lesssim (1+t)^{-1/2} (1+|x|^2)^{-\frac{3}{2}} \lesssim (1+t)^{-1/2} \left(\frac{1+t}{2} + \frac{|x|^2}{2}\right)^{-\frac{3}{2}} \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}
\end{aligned}$$

since $|x| \geq \sqrt{1+t}$. Therefore,

$$I_1 \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.$$

For I_2 , we decompose \mathbb{R}^3 into two parts

$$I_2 = \int_{\frac{t}{2}}^t \left(\int_{|y|-c\tau \leq \frac{|x|-c\tau}{2}} + \int_{|y|-c\tau > \frac{|x|-c\tau}{2}} \right) (\dots) dyd\tau = I_{21} + I_{22}.$$

If $|y| - c\tau \leq \frac{|x|-c\tau}{2}$, we have

$$|x-y| \geq |x| - \left(\frac{|x|-c\tau}{2}\right) - c\tau \geq \frac{|x|-c\tau}{2},$$

and thus

$$\begin{aligned}
I_{21} &\lesssim (1+(|x|-c\tau)^2)^{-\frac{3}{2}} \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-1/2} (1+\tau)^{-4} \left(1 + \frac{(|y|-c\tau)^2}{1+\tau}\right)^{-2} dyd\tau \\
&\lesssim (1+t)^{-\frac{3}{2}} \left(1 + \frac{(|x|-c\tau)^2}{1+t}\right)^{-\frac{3}{2}} \int_{\frac{t}{2}}^t (1+t-\tau)^{-1/2} (1+\tau)^{-4} (1+\tau)^{\frac{5}{2}} d\tau \\
&\lesssim (1+t)^{-5/2} \left(1 + \frac{(|x|-c\tau)^2}{1+t}\right)^{-\frac{3}{2}}.
\end{aligned}$$

If $|y| - c\tau > \frac{|x|-c\tau}{2}$, then

$$\begin{aligned}
I_{22} &\lesssim (1+t)^{-4} \left(1 + \frac{(|x|-c\tau)^2}{1+t}\right)^{-2} \int_{\frac{t}{2}}^t \int_{\{|x-y|\leq c(t-\tau)\}} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-\frac{3}{2}} dyd\tau \\
&\lesssim (1+t)^{-4} \left(1 + \frac{(|x|-c\tau)^2}{1+t}\right)^{-2} \int_{\frac{t}{2}}^t (1+t-\tau)^{-1/2} \ln(1+t-\tau) d\tau \\
&\lesssim (1+t)^{-3} \left(1 + \frac{(|x|-c\tau)^2}{1+t}\right)^{-2}.
\end{aligned}$$

Therefore,

$$I_2 \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t} \right)^{-\frac{3}{2}}.$$

Combining all the estimates, we get desired estimate

$$I \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t} \right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t} \right)^{-\frac{3}{2}}.$$

Case 4: $(x, t) \in D_4$. In this region $\sqrt{1+t} \leq |x| \leq \frac{\mathbf{c}t}{2}$. We split the integral I into three parts:

$$\begin{aligned} I &= \int_0^t \int_{\mathbb{R}^3} (1+t-\tau)^{-4} \left(1 + \frac{(|x-y| - \mathbf{c}(t-\tau))^2}{1+t-\tau} \right)^{-2} (1+\tau)^{-2} \left(1 + \frac{|y|^2}{D_0(1+\tau)} \right)^{-\frac{3}{2}} \mathbf{1}_{\{|y| \leq \mathbf{c}\tau\}} dy d\tau \\ &= \left(\int_0^{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}} + \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{t - \frac{|x|}{2\mathbf{c}}} + \int_{t - \frac{|x|}{2\mathbf{c}}}^t \right) \int_{\mathbb{R}^3} (\dots) dy d\tau =: I_1 + I_2 + I_3. \end{aligned}$$

For I_1 , we decompose \mathbb{R}^3 into two parts

$$I_1 = \int_0^{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}} \left(\int_{|y| \leq \frac{\mathbf{c}(t-\tau) - |x|}{2}} + \int_{|y| > \frac{\mathbf{c}(t-\tau) - |x|}{2}} \right) =: I_{11} + I_{12}.$$

If $0 \leq \tau \leq \frac{t}{2} - \frac{|x|}{2\mathbf{c}}$ and $|y| \leq \frac{\mathbf{c}(t-\tau) - |x|}{2}$, then

$$\mathbf{c}(t-\tau) - |x-y| \geq \mathbf{c}(t-\tau) - |x| - |y| \geq \frac{\mathbf{c}t - |x|}{4},$$

so that

$$\begin{aligned} I_{11} &\lesssim (1+t+|x|)^{-4} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t} \right)^{-2} \int_0^{\frac{t}{2}} \int_{|y| \leq \mathbf{c}\tau} (1+\tau)^{-2} \left(1 + \frac{|y|^2}{D_0(1+\tau)} \right)^{-\frac{3}{2}} dy d\tau \\ &\lesssim (1+t)^{-4} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t} \right)^{-2} (1+t)^{3/4} \lesssim (1+t)^{-13/4} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t} \right)^{-2}. \end{aligned}$$

If $0 \leq \tau \leq \frac{t}{2} - \frac{|x|}{2\mathbf{c}}$ and $|y| > \frac{\mathbf{c}(t-\tau) - |x|}{2}$, then

$$|y| > \frac{\mathbf{c}(t-\tau) - |x|}{2} \geq \frac{\mathbf{c}t - |x|}{4},$$

and thus

$$\begin{aligned} I_{12} &\lesssim \left(1 + \frac{t}{2} + \frac{|x|}{2\mathbf{c}} \right)^{-4} \left(1 + (\mathbf{c}t - |x|)^2 \right)^{-\frac{3}{2}} \int_0^{\frac{t}{2} - \frac{|x|}{\mathbf{c}}} \int_{\mathbb{R}^3} (1+\tau)^{-1/2} \left(1 + \frac{(|x-y| - \mathbf{c}(t-\tau))^2}{1+t-\tau} \right)^{-2} dy d\tau \\ &\lesssim (1+t)^{-4} \left(1 + (\mathbf{c}t - |x|)^2 \right)^{-\frac{3}{2}} \int_0^{\frac{t}{2}} (1+\tau)^{-1/2} (1+t-\tau)^{\frac{5}{2}} d\tau \\ &\lesssim (1+t)^{-4} \left(\frac{(\mathbf{c}t - |x|)^2}{2} + \frac{1+t}{2} \right)^{-\frac{3}{2}} (1+t)^{\frac{7}{2}} \lesssim (1+t)^{-5/2} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t} \right)^{-1} \end{aligned}$$

since $\mathbf{c}t - |x| \geq \frac{\mathbf{c}t}{2}$. Therefore,

$$I_1 \lesssim (1+t)^{-5/2} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t} \right)^{-1}.$$

For I_2 , we use the spherical coordinates to obtain

$$\begin{aligned}
I_2 &\lesssim \int_{\frac{t}{2}-\frac{|x|}{2c}}^{t-\frac{|x|}{2c}} \int_0^\infty \int_0^\pi (1+t-\tau)^{-4} \left(1 + \frac{\left(\sqrt{|x|^2 + r^2 - 2r|x|\cos\theta} - \mathbf{c}(t-\tau) \right)^2}{1+t-\tau} \right)^{-2} \\
&\quad \cdot (1+\tau)^{-2} \left(1 + \frac{|r|^2}{(1+\tau)} \right)^{-\frac{3}{2}} \mathbf{1}_{\{r \leq c\tau\}} r^2 \sin\theta d\theta dr d\tau \\
&\lesssim \int_{\frac{t}{2}-\frac{|x|}{2c}}^{t-\frac{|x|}{2c}} \int_0^\infty \int_{||x|-r|}^{|x|+r} (1+t-\tau)^{-4} \left(1 + \frac{(z - \mathbf{c}(t-\tau))^2}{1+t-\tau} \right)^{-2} z (1+\tau)^{-2} \\
&\quad \cdot \left(1 + \frac{r^2}{(1+\tau)} \right)^{-\frac{3}{2}} \frac{r}{|x|} \mathbf{1}_{\{r \leq c\tau\}} dz dr d\tau \\
&\lesssim |x|^{-1} \int_{\frac{t}{2}-\frac{|x|}{2c}}^{t-\frac{|x|}{2c}} \int_0^\infty (1+t-\tau)^{-4+\frac{3}{2}} (1+\tau)^{-2} \cdot r \left(1 + \frac{r^2}{(1+\tau)} \right)^{-\frac{3}{2}} \mathbf{1}_{\{r \leq c\tau\}} dr d\tau \\
&\lesssim |x|^{-1} \int_{\frac{t}{2}-\frac{|x|}{2c}}^{t-\frac{|x|}{2c}} (1+t-\tau)^{-\frac{5}{2}} (1+\tau)^{-1} d\tau \\
&\lesssim |x|^{-1} \left(1 + \frac{|x|}{2c} \right)^{-\frac{3}{2}} \left(1 + \frac{t}{2} - \frac{|x|}{2c} \right)^{-1} \\
&\lesssim (1+t)^{-1/2} (1+|x|)^{-3} \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t} \right)^{-\frac{3}{2}}.
\end{aligned}$$

by setting $z = \sqrt{|x|^2 + r^2 - 2r|x|\cos\theta}$ and $\sin\theta d\theta = \frac{z}{r|x|} dz$.

For I_3 , we decompose \mathbb{R}^3 into two parts

$$I_3 = \int_{t-\frac{|x|}{2c}}^t \left(\int_{|y| \leq \frac{|x| - \mathbf{c}(t-\tau)}{2}} + \int_{|y| > \frac{|x| - \mathbf{c}(t-\tau)}{2}} \right) =: I_{31} + I_{32}.$$

If $t - \frac{|x|}{2c} \leq \tau \leq t$, $|y| \leq \frac{|x| - \mathbf{c}(t-\tau)}{2}$, then

$$|x-y| - \mathbf{c}(t-\tau) \geq |x| - |y| - \mathbf{c}(t-\tau) \geq \frac{|x| - \mathbf{c}(t-\tau)}{2} \geq \frac{|x|}{4}.$$

If $t - \frac{|x|}{2c} \leq \tau \leq t$, $|y| > \frac{|x| - \mathbf{c}(t-\tau)}{2}$, then

$$|y| > \frac{|x| - \mathbf{c}(t-\tau)}{2} \geq \frac{|x|}{4}.$$

Hence,

$$\begin{aligned}
I_{31} &\lesssim \left(1 + |x|^2 \right)^{-\frac{3}{2}} \int_{t-\frac{|x|}{2c}}^t \int_{|y| \leq \frac{|x| - \mathbf{c}(t-\tau)}{2}} (1+t-\tau)^{-\frac{5}{2}} (1+\tau)^{-2} \left(1 + \frac{|y|^2}{D_0(1+\tau)} \right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-1/2} \left(1 + |x|^2 \right)^{-\frac{3}{2}} \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{(1+t)} \right)^{-\frac{3}{2}},
\end{aligned}$$

since $|x| \geq \sqrt{1+t}$, and

$$I_{32} \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{(1+t)} \right)^{-\frac{3}{2}} \int_{t-\frac{|x|}{2c}}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-4} \left(1 + \frac{(|x-y| - \mathbf{c}(t-\tau))^2}{1+t-\tau} \right)^{-2} dy d\tau$$

$$\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{(1+t)}\right)^{-\frac{3}{2}}.$$

Therefore,

$$I_3 \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{(1+t)}\right)^{-\frac{3}{2}}.$$

Combining all the estimates yields

$$I \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{(1+t)}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(ct - |x|)^2}{1+t}\right)^{-1}.$$

Case 5: $(x, t) \in D_5$. In this region $ct/2 \leq |x| \leq ct - \sqrt{1+t}$. Now we split the integral I into three parts

$$\begin{aligned} I &= \int_0^t \int_{\mathbb{R}^3} (1+t-\tau)^{-4} \left(1 + \frac{(|x-y| - \mathbf{c}(t-\tau))^2}{1+t-\tau}\right)^{-2} (1+\tau)^{-2} \left(1 + \frac{|y|^2}{D_0(1+\tau)}\right)^{-\frac{3}{2}} \mathbf{1}_{\{|y| \leq \mathbf{c}\tau\}} dy d\tau \\ &= \left(\int_0^{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}} + \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{\mathbf{c}}))} + \int_{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{\mathbf{c}}))}^t \right) \int_{\mathbb{R}^3} (\dots) dy d\tau =: I_1 + I_2 + I_3. \end{aligned}$$

For I_1 , the estimate is the same as the I_1 of Case 4, so

$$I_1 \lesssim (1+t)^{-5/2} \left(1 + \frac{(ct - |x|)^2}{1+t}\right)^{-1}.$$

For I_2 , we use the spherical coordinates to obtain

$$\begin{aligned} I_2 &= \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{\mathbf{c}}))} \int_0^\infty \int_0^\pi (1+t-\tau)^{-4} \left(1 + \frac{\left(\sqrt{|x|^2 + r^2 - 2r|x|\cos\theta} - \mathbf{c}(t-\tau)\right)^2}{1+t-\tau}\right)^{-2} \\ &\quad \cdot (1+\tau)^{-2} \left(1 + \frac{r^2}{D_0(1+\tau)}\right)^{-\frac{3}{2}} \mathbf{1}_{\{r \leq \mathbf{c}\tau\}} r^2 \sin\theta d\theta dr d\tau \\ &= \frac{1}{|x|} \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{\mathbf{c}}))} \int_0^\infty \int_{||x|-r|}^{|x|+r} (1+t-\tau)^{-4} \left(1 + \frac{(z - \mathbf{c}(t-\tau))^2}{1+t-\tau}\right)^{-2} z (1+\tau)^{-2} \\ &\quad \cdot r \left(1 + \frac{r^2}{D_0(1+\tau)}\right)^{-\frac{3}{2}} \mathbf{1}_{\{r \leq \mathbf{c}\tau\}} dz dr d\tau \\ &\lesssim |x|^{-1} \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{\mathbf{c}}))} (1+t-\tau)^{-\frac{5}{2}} (1+\tau)^{-1} d\tau \\ &\lesssim |x|^{-1} \left(1 + \frac{3|x|}{4\mathbf{c}} - \frac{t}{4}\right)^{-\frac{3}{2}} \left(1 + \frac{t}{2} - \frac{|x|}{2\mathbf{c}}\right)^{-1} \\ &\lesssim |x|^{-1} (1+t)^{-1/2} \left(1 + \frac{t}{4} - \frac{|x|}{4\mathbf{c}} + \frac{|x|}{\mathbf{c}} - \frac{t}{2}\right)^{-1} \left(1 + \frac{t}{2} - \frac{|x|}{2\mathbf{c}}\right)^{-1} \\ &\lesssim (1+t)^{-5/2} \left(1 + \frac{(ct - |x|)^2}{1+t}\right)^{-1} \end{aligned}$$

by setting $z = \sqrt{|x|^2 + r^2 - 2r|x|\cos\theta}$ and $\sin\theta d\theta = \frac{z}{r|x|} dz$.

For I_3 , we decompose \mathbb{R}^3 into two parts

$$I_3 = \int_{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{c}))}^t \left(\int_{|y| \leq \frac{|x| - c(t-\tau)}{2}} + \int_{|y| > \frac{|x| - c(t-\tau)}{2}} \right) (\dots) dy d\tau =: I_{31} + I_{32}.$$

If $\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{c})) \leq \tau \leq t$, $|y| \leq \frac{|x| - c(t-\tau)}{2}$, then

$$|x - y| - c(t - \tau) \geq \frac{|x| - c(t - \tau)}{2} \geq \frac{ct + |x|}{8} = \frac{ct - |x|}{8} + \frac{|x|}{4}.$$

If $\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{c})) \leq \tau \leq t$, $|y| > \frac{|x| - c(t-\tau)}{2}$, then

$$|y| > \frac{|x| - c(t - \tau)}{2} \geq \frac{ct - |x|}{8} + \frac{|x|}{4}.$$

Hence,

$$\begin{aligned} I_{31} &\lesssim \left(1 + (ct - |x|)^2\right)^{-2} \int_{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{c}))}^t \int_{|y| \leq c\tau} (1 + t - \tau)^{-2} (1 + \tau)^{-2} \left(1 + \frac{|y|^2}{D_0(1 + \tau)}\right)^{-\frac{3}{2}} dy d\tau \\ &\lesssim \left(1 + (ct - |x|)^2\right)^{-2} \int_{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{c}))}^t (1 + t - \tau)^{-2} (1 + \tau)^{-1/4} d\tau \\ &\lesssim (1 + t)^{-1/4} \left(1 + (ct - |x|)^2\right)^{-2} \lesssim (1 + t)^{-9/4} \left(1 + \frac{(ct - |x|)^2}{1 + t}\right)^{-2}. \end{aligned}$$

On the other hand,

$$\begin{aligned} I_{31} &\lesssim (1 + |x|^2)^{-2} \int_{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{c}))}^t \int_{|y| \leq c\tau} (1 + t - \tau)^{-2} (1 + \tau)^{-2} \left(1 + \frac{|y|^2}{D_0(1 + \tau)}\right)^{-\frac{3}{2}} dy d\tau \\ &\lesssim (1 + t)^{-17/4}. \end{aligned}$$

By interpolation, one has

$$I_{31} \lesssim (1 + t)^{-13/4} \left(1 + \frac{(ct - |x|)^2}{1 + t}\right)^{-1}.$$

For I_{32} ,

$$\begin{aligned} I_{32} &\lesssim \int_{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{c}))}^t (1 + |x| - c(t - \tau))^{-3} (1 + \tau)^{-1/2} \\ &\quad \cdot \int_{|y| > \frac{|x| - c(t-\tau)}{2}} (1 + t - \tau)^{-4} \left(1 + \frac{(|x - y| - c(t - \tau))^2}{1 + t - \tau}\right)^{-2} dy d\tau \\ &\lesssim \int_{\frac{1}{2}(t + \frac{3}{2}(t - \frac{|x|}{c}))}^t (1 + |x| - c(t - \tau))^{-3} (1 + t - \tau)^{-4 + \frac{5}{2}} (1 + \tau)^{-\frac{1}{2}} d\tau \\ &\lesssim (1 + t)^{-7/2} \lesssim (1 + t)^{-1/2} (1 + |x|)^{-3} \lesssim (1 + t)^{-2} \left(1 + \frac{|x|^2}{(1 + t)}\right)^{-\frac{3}{2}}, \end{aligned}$$

so that

$$I_3 \lesssim (1 + t)^{-2} \left(1 + \frac{|x|^2}{(1 + t)}\right)^{-\frac{3}{2}} + (1 + t)^{-5/2} \left(1 + \frac{(ct - |x|)^2}{1 + t}\right)^{-1}.$$

To sum up,

$$I \leq (1 + t)^{-2} \left(1 + \frac{|x|^2}{(1 + t)}\right)^{-\frac{3}{2}} + (1 + t)^{-5/2} \left(1 + \frac{(ct - |x|)^2}{1 + t}\right)^{-1}.$$

□

Proof of Lemma 25. (Huygens wave convolved with Huygens wave).

Case 1: $(x, t) \in D_1 \cup D_2$. Direct computation gives

$$\begin{aligned}
J &\lesssim \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+t)^{-5/2} (1+\tau)^{-4} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\quad + \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y| - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+t)^{-4} dy d\tau \\
&\lesssim (1+t)^{-5/2} \int_0^{\frac{t}{2}} (1+\tau)^{-4} (1+\tau)^{\frac{5}{2}} d\tau + (1+t)^{-4} \int_{\frac{t}{2}}^t (1+t-\tau)^{-5/2} (1+t-\tau)^{\frac{5}{2}} d\tau \\
&\lesssim (1+t)^{-5/2}.
\end{aligned}$$

Case 2: $(x, t) \in D_3$. We split the integral J into four parts

$$\begin{aligned}
J &= \int_0^{\frac{t}{2}} \left(\int_{|y| - \mathbf{c}\tau \leq \frac{|x| - \mathbf{c}t}{2}} + \int_{|y| - \mathbf{c}\tau > \frac{|x| - \mathbf{c}t}{2}} \right) (\dots) dy d\tau + \int_{\frac{t}{2}}^t \left(\int_{|y| - \mathbf{c}\tau \leq \frac{|x| - \mathbf{c}t}{2}} + \int_{|y| - \mathbf{c}\tau > \frac{|x| - \mathbf{c}t}{2}} \right) (\dots) dy d\tau \\
&= J_{11} + J_{12} + J_{21} + J_{22}.
\end{aligned}$$

Note that if $|y| - \mathbf{c}\tau \leq \frac{|x| - \mathbf{c}t}{2}$, then

$$|x - y| - \mathbf{c}(t - \tau) \geq |x| - |y| - \mathbf{c}(t - \tau) \geq \frac{|x| - \mathbf{c}t}{2}.$$

Hence,

$$\begin{aligned}
J_{11} &\lesssim \int_0^{\frac{t}{2}} \int_{|y| - \mathbf{c}\tau \leq \frac{|x| - \mathbf{c}t}{2}} (1+t)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2}{D_0(1+t)}} (1+\tau)^{-4} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\lesssim \int_0^{\frac{t}{2}} (1+t)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2}{D_0(1+t)}} (1+\tau)^{-4} (1+\tau)^{\frac{5}{2}} d\tau \lesssim (1+t)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2}{D_0(1+t)}}.
\end{aligned}$$

For J_{12} , we have

$$\begin{aligned}
J_{12} &\lesssim \int_0^{\frac{t}{2}} \int_{|y| - \mathbf{c}\tau > \frac{|x| - \mathbf{c}t}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y| - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+\tau)^{-4} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\lesssim \int_0^{\frac{t}{2}} (1+t-\tau)^{-5/2} (1+t-\tau)^{\frac{5}{2}} (1+\tau)^{-2} \left(1 + (|x| - \mathbf{c}t)^2\right)^{-2} d\tau \\
&\lesssim (1 + |x| - \mathbf{c}t)^{-4},
\end{aligned}$$

and

$$\begin{aligned}
J_{12} &\lesssim \int_0^{\frac{t}{2}} \int_{|y| - \mathbf{c}\tau > \frac{|x| - \mathbf{c}t}{2}} (1+t)^{-5/2} (1+\tau)^{-2} (1 + |y| - \mathbf{c}\tau)^{-4} dy d\tau \\
&\lesssim \int_0^{\frac{t}{2}} (1+t)^{-5/2} (1+\tau)^{-2} \int_{\mathbf{c}\tau + \frac{|x| - \mathbf{c}t}{2}}^{\infty} (1+r - \mathbf{c}\tau)^{-4} r^2 dr d\tau \\
&\lesssim \int_0^{\frac{t}{2}} (1+t)^{-5/2} (1+\tau)^{-2} \int_{\frac{|x| - \mathbf{c}t}{2}}^{\infty} (1+r)^{-4} (r + \mathbf{c}\tau)^2 dr d\tau \\
&\lesssim \int_0^{\frac{t}{2}} (1+t)^{-5/2} (1+\tau)^{-2} \left[r^{-1} + \mathbf{c}\tau \cdot r^{-2} + (\mathbf{c}\tau)^2 r^{-3} \right]_{r=\frac{|x| - \mathbf{c}t}{2}} d\tau \\
&\lesssim (1+t)^{-5/2} \left[(1 + |x| - \mathbf{c}t)^{-1} + \ln(2+t) (1 + |x| - \mathbf{c}t)^{-2} + t (1 + |x| - \mathbf{c}t)^{-3} \right]
\end{aligned}$$

$$\begin{aligned} &\lesssim (1+t)^{-5/2} (1+|x|-\mathbf{c}t)^{-1} \left[1 + \ln(2+t) (1+\sqrt{1+t})^{-1} + t(1+\sqrt{1+t})^{-2} \right] \\ &\lesssim (1+t)^{-5/2} (1+|x|-\mathbf{c}t)^{-1}, \end{aligned}$$

which implies that

$$\begin{aligned} J_{12} &= (J_{12})^{\frac{1}{3}} (J_{12})^{\frac{2}{3}} \lesssim \left[(1+|x|-\mathbf{c}t)^{-4} \right]^{\frac{1}{3}} \left[(1+t)^{-5/2} (1+|x|-\mathbf{c}t)^{-1} \right]^{\frac{2}{3}} \\ &\lesssim (1+t)^{-\frac{5}{3}} (1+|x|-\mathbf{c}t)^{-2} \\ &\lesssim (1+t)^{-\frac{8}{3}} \left(1 + \frac{(|x|-\mathbf{c}t)^2}{1+t} \right)^{-1}. \end{aligned}$$

For J_{21} and J_{22} , it follows that

$$\begin{aligned} J_{21} &\lesssim \int_{\frac{t}{2}}^t \int_{|y|-\mathbf{c}\tau \leq \frac{|x|-\mathbf{c}t}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x|-\mathbf{c}t)^2}{2D_0(1+t)}} e^{-\frac{(|x-y|-\mathbf{c}(t-\tau))^2}{2D_0(1+t-\tau)}} (1+t)^{-4} \left(1 + \frac{(|y|-\mathbf{c}\tau)^2}{1+\tau} \right)^{-2} dyd\tau \\ &\lesssim (1+t)^{-4} e^{-\frac{(|x|-\mathbf{c}t)^2}{2D_0(1+t)}} \int_{\frac{t}{2}}^t (1+t-\tau)^{-5/2} (1+t-\tau)^{\frac{5}{2}} d\tau \\ &\lesssim (1+t)^{-3} e^{-\frac{(|x|-\mathbf{c}t)^2}{2D_0(1+t)}}, \end{aligned}$$

$$\begin{aligned} J_{22} &\lesssim \int_{\frac{t}{2}}^t \int_{|y|-\mathbf{c}\tau > \frac{|x|-\mathbf{c}t}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y|-\mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+t)^{-4} \left(1 + \frac{(|y|-\mathbf{c}\tau)^2}{1+\tau} \right)^{-2} dyd\tau \\ &\lesssim \int_{\frac{t}{2}}^t (1+t-\tau)^{-5/2} (1+t-\tau)^{\frac{5}{2}} (1+t)^{-4} \left(1 + \frac{(|x|-\mathbf{c}t)^2}{1+t} \right)^{-2} d\tau \\ &\lesssim (1+t)^{-3} \left(1 + \frac{(|x|-\mathbf{c}t)^2}{1+t} \right)^{-2}. \end{aligned}$$

Gathering all the estimates, we can conclude that

$$J \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x|-\mathbf{c}t)^2}{1+t} \right)^{-1}.$$

Case 3: $(x, t) \in D_4$. We split the integral into four parts

$$\begin{aligned} J &= \left(\int_0^{\frac{t}{4}-\frac{|x|}{4\mathbf{c}}} + \int_{\frac{t}{4}-\frac{|x|}{4\mathbf{c}}}^{\frac{t}{2}} + \int_{\frac{t}{2}}^{\frac{t}{2}+\frac{1}{4}(t+\frac{|x|}{\mathbf{c}})} + \int_{\frac{t}{2}+\frac{1}{4}(t+\frac{|x|}{\mathbf{c}})}^t \right) \int_{\mathbb{R}^3} (\dots) dyd\tau \\ &=: J_1 + J_2 + J_3 + J_4. \end{aligned}$$

For J_1 , we decompose \mathbb{R}^3 into two parts

$$J_1 = \int_0^{\frac{t}{4}-\frac{|x|}{4\mathbf{c}}} \left(\int_{|y| \leq \frac{\mathbf{c}t-|x|}{2}} + \int_{|y| > \frac{\mathbf{c}t-|x|}{2}} \right) \int_{\mathbb{R}^3} (\dots) dyd\tau =: J_{11} + J_{12}.$$

If $0 \leq \tau \leq \frac{t}{4} - \frac{|x|}{4\mathbf{c}}$, $|y| \leq \frac{\mathbf{c}t-|x|}{2}$, then

$$\mathbf{c}(t-\tau) - |x-y| \geq \mathbf{c}(t-\tau) - |x| - |y| \geq \frac{\mathbf{c}t-|x|}{4} \geq \frac{\mathbf{c}t}{8}.$$

Hence,

$$\begin{aligned} J_{11} &\lesssim \int_0^{\frac{t}{4}-\frac{|x|}{4\mathbf{c}}} \int_{|y| \leq \frac{\mathbf{c}t-|x|}{2}} (1+t-s)^{-5/2} e^{-\frac{(|x|-\mathbf{c}t)^2}{2D_0(1+t)}} (1+\tau)^{-4} \left(1 + \frac{(|y|-\mathbf{c}\tau)^2}{1+\tau} \right)^{-2} dyd\tau \\ &\lesssim (1+t)^{-5/2} e^{-\frac{(|x|-\mathbf{c}t)^2}{2D_0(1+t)}} \int_0^{\frac{t}{4}-\frac{|x|}{4\mathbf{c}}} (1+\tau)^{-4} (1+\tau)^{\frac{5}{2}} d\tau \lesssim (1+t)^{-5/2} e^{-\frac{(|x|-\mathbf{c}t)^2}{2D_0(1+t)}}, \end{aligned}$$

and

$$\begin{aligned}
J_{12} &\lesssim \int_0^{\frac{t}{4} - \frac{|x|}{4c}} \int_{|y| > \frac{ct - |x|}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y| - c(t-\tau))^2}{D_0(1+t-\tau)}} (1+\tau)^{-2} \left(1 + (c\tau - |x|)^2\right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-4} \int_0^{\frac{t}{4} - \frac{|x|}{4c}} (1+t-\tau)^{-5/2} (1+t-\tau)^{\frac{5}{2}} (1+\tau)^{-2} d\tau \\
&\lesssim (1+t)^{-4} \lesssim (1+t)^{-1} (1+|x|)^{-3} \lesssim (1+t)^{-5/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}},
\end{aligned}$$

since $\sqrt{1+t} \leq |x| \leq \frac{ct}{2}$, so that

$$J_1 \lesssim (1+t)^{-5/2} e^{-\frac{(|x| - ct)^2}{2D_0(1+t)}} + (1+t)^{-5/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.$$

For J_2 and J_3 ,

$$\begin{aligned}
J_2 &\lesssim \int_{\frac{t}{4} - \frac{|x|}{4c}}^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+t)^{-5/2} e^{-\frac{(|x-y| - c(t-\tau))^2}{D_0(1+t-\tau)}} (1+\tau)^{-4} \left(1 + \frac{(|y| - c\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-5/2} \left(1 + \frac{t}{4} - \frac{|x|}{4c}\right)^{-4} \int_0^t \int_{\mathbb{R}^3} e^{-\frac{(|x-y| - c(t-\tau))^2}{D_0(1+t-\tau)}} \left(1 + \frac{(|y| - c\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-5/2} (1+t)^{-4} (1+t)^3 \lesssim (1+t)^{-7/2} \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}},
\end{aligned}$$

$$\begin{aligned}
J_3 &\lesssim \int_{\frac{t}{2}}^{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{c})} \int_{\mathbb{R}^3} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y| - c(t-\tau))^2}{D_0(1+t-\tau)}} (1+t)^{-4} \left(1 + \frac{(|y| - c\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\lesssim \left(1 + \frac{1}{4} \left(t - \frac{|x|}{c}\right)\right)^{-5/2} (1+t)^{-4} \int_0^t \int_{\mathbb{R}^3} e^{-\frac{(|x-y| - c(t-\tau))^2}{D_0(1+t-\tau)}} \left(1 + \frac{(|y| - c\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-13/2} (1+t)^3 \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}},
\end{aligned}$$

due to Lemma 5.4 of [24].

For J_4 , we decompose \mathbb{R}^3 into two parts

$$J_4 = \int_{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{c})}^t \left(\int_{|y| \leq \frac{|x| + ct}{2}} + \int_{|y| > \frac{|x| + ct}{2}} \right) (\dots) dy d\tau =: J_{41} + J_{42}.$$

If $\frac{t}{2} + \frac{1}{4} \left(t + \frac{|x|}{c}\right) \leq \tau \leq t$, $|y| \leq \frac{|x| + ct}{2}$, then

$$c\tau - |y| \geq \frac{ct}{2} + \frac{1}{4}(ct + |x|) - \frac{|x| + ct}{2} \geq \frac{ct - |x|}{4} \geq \frac{ct}{8}.$$

If $\frac{t}{2} + \frac{1}{4} \left(t + \frac{|x|}{c}\right) \leq \tau \leq t$, $|y| > \frac{|x| + ct}{2}$, then

$$\begin{aligned}
|x-y| - c(t-\tau) &\geq |y| - |x| - c(t-\tau) \geq \frac{|x| + ct}{2} - |x| - ct + \frac{ct}{2} + \frac{1}{4}(ct + |x|) \\
&\geq \frac{ct - |x|}{4} \geq \frac{ct}{8}.
\end{aligned}$$

Hence,

$$J_{41} \lesssim \int_{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{c})}^t \int_{|y| \leq \frac{|x| + ct}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y| - c(t-\tau))^2}{D_0(1+t-\tau)}} (1+t)^{-4} \left(1 + \frac{(ct - |x|)^2}{1+t}\right)^{-2} dy d\tau$$

$$\begin{aligned}
&\lesssim (1+t)^{-4} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-2} (1+t) \\
&\lesssim (1+t)^{-5} \lesssim (1+t)^{-2} (1+|x|)^{-3} \lesssim (1+t)^{-7/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}, \\
J_{42} &\lesssim \int_{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{\mathbf{c}})}^t \int_{|y| > \frac{|x| + \mathbf{c}t}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(\mathbf{c}t - |x|)^2}{2D_0(1+t)}} e^{-\frac{(|x-y| - \mathbf{c}(t-\tau))^2}{2D_0(1+t-\tau)}} (1+t)^{-4} dy d\tau \\
&\lesssim (1+t)^{-4} e^{-\frac{(\mathbf{c}t - |x|)^2}{2D_0(1+t)}} \int_{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{\mathbf{c}})}^t (1+t-\tau)^{-\frac{5}{2} + \frac{5}{2}} d\tau \\
&\lesssim (1+t)^{-3} e^{-\frac{t}{\mathbf{c}}} \lesssim (1+t)^{-7/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.
\end{aligned}$$

for some $C > 0$, so that

$$J_4 \lesssim (1+t)^{-7/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.$$

As a result,

$$J \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2}{2D_0(1+t)}}$$

for $(x, t) \in D_4$.

Case 4: $(x, t) \in D_5$. We split the integral J into five parts

$$J = \left(\int_0^{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}} + \int_{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}}^{\frac{t}{2}} + \int_{\frac{t}{2}}^{\frac{|x|}{2\mathbf{c}} + \frac{1}{4}(t + \frac{|x|}{\mathbf{c}})} + \int_{\frac{|x|}{2\mathbf{c}} + \frac{1}{4}(t + \frac{|x|}{\mathbf{c}})}^{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{\mathbf{c}})} + \int_{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{\mathbf{c}})}^t \right) \int_{\mathbb{R}^3} (\dots) dy d\tau =: \sum_{i=1}^5 J_i.$$

For J_1 , we decompose \mathbb{R}^3 into two parts

$$J_1 = \int_0^{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}} \left(\int_{|y| \leq \frac{\mathbf{c}t - |x|}{2}} + \int_{|y| > \frac{\mathbf{c}t - |x|}{2}} \right) (\dots) dy d\tau =: J_{11} + J_{12}.$$

If $0 \leq \tau \leq \frac{t}{4} - \frac{|x|}{4\mathbf{c}}$, $|y| \leq \frac{\mathbf{c}t - |x|}{2}$, then

$$\mathbf{c}(t-\tau) - |x-y| \geq \mathbf{c}(t-\tau) - (|x| + |y|) \geq \frac{\mathbf{c}t - |x|}{4}.$$

If $0 \leq \tau \leq \frac{t}{4} - \frac{|x|}{4\mathbf{c}}$, $|y| > \frac{\mathbf{c}t - |x|}{2}$, then

$$|y| - \mathbf{c}\tau \geq \frac{\mathbf{c}t - |x|}{4}.$$

Hence,

$$\begin{aligned}
J_{11} &\lesssim \int_0^{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}} \int_{|y| \leq \frac{\mathbf{c}t - |x|}{2}} (1+t)^{-5/2} e^{-\frac{(\mathbf{c}t - |x|)^2}{D_0(1+t)}} (1+\tau)^{-4} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-5/2} e^{-\frac{(\mathbf{c}t - |x|)^2}{D_0(1+t)}} \int_0^{\frac{\mathbf{c}t - |x|}{4}} (1+\tau)^{-4 + \frac{5}{2}} d\tau \lesssim (1+t)^{-5/2} e^{-\frac{(\mathbf{c}t - |x|)^2}{D_0(1+t)}}.
\end{aligned}$$

For J_{12} , we have

$$\begin{aligned}
J_{12} &\lesssim \int_0^{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}} \int_{|y| > \frac{\mathbf{c}t - |x|}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y| - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+\tau)^{-2} \left(1 + (\mathbf{c}t - |x|)^2\right)^{-2} dy d\tau \\
&\lesssim \left(1 + (\mathbf{c}t - |x|)^2\right)^{-2} \int_0^{\frac{\mathbf{c}t - |x|}{4}} (1+\tau)^{-2} d\tau
\end{aligned}$$

$$\lesssim (1 + \mathbf{c}t - |x|)^{-4},$$

and

$$\begin{aligned} J_{12} &\lesssim \int_0^{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}} (1+t)^{-5/2} (1+\tau)^{-2} \int_{\frac{\mathbf{c}t - |x|}{2}}^{\infty} (1+r - \mathbf{c}\tau)^{-4} r^2 dr d\tau \\ &\lesssim \int_0^{\frac{\mathbf{c}t - |x|}{4}} (1+t)^{-5/2} (1+\tau)^{-2} (1 + \mathbf{c}t - |x|)^{-1} d\tau \\ &\lesssim (1+t)^{-5/2} (1 + \mathbf{c}t - |x|)^{-1}, \end{aligned}$$

which implies that

$$\begin{aligned} J_{12} &\lesssim (J_{12})^{\frac{1}{3}} (J_{12})^{\frac{2}{3}} \lesssim \left[(1 + \mathbf{c}t - |x|)^{-4} \right]^{\frac{1}{3}} \left[(1+t)^{-5/2} (1 + \mathbf{c}t - |x|)^{-1} \right]^{\frac{2}{3}} \\ &\lesssim (1+t)^{-\frac{5}{3}} (1 + \mathbf{c}t - |x|)^{-2} \lesssim (1+t)^{-\frac{8}{3}} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t} \right)^{-1}. \end{aligned}$$

Therefore, we get

$$J_1 \lesssim (1+t)^{-5/2} e^{-\frac{(\mathbf{c}t - |x|)^2}{D_0(1+t)}} + (1+t)^{-\frac{8}{3}} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t} \right)^{-1}.$$

For J_2 , we use the spherical coordinates to obtain

$$\begin{aligned} J_2 &\lesssim \int_{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}}^{\frac{t}{2}} \int_0^{\infty} \int_0^{\pi} (1+t-\tau)^{-5/2} e^{-\frac{(\sqrt{|x|^2 + r^2 - 2r|x|\cos\theta} - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+\tau)^{-4} \left(1 + \frac{(r - \mathbf{c}\tau)^2}{1+\tau} \right)^{-2} r^2 \sin\theta d\theta dr d\tau \\ &\lesssim \int_{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}}^{\frac{t}{2}} \int_0^{\infty} \int_{||x|-r|}^{|x|+r} (1+t-\tau)^{-5/2} e^{-\frac{(z - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+\tau)^{-4} \left(1 + \frac{(r - \mathbf{c}\tau)^2}{1+\tau} \right)^{-2} rz \frac{1}{|x|} dz dr d\tau \\ &\lesssim \int_{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}}^{\frac{t}{2}} \int_0^{\infty} \int_0^{\infty} (1+t-\tau)^{-5/2} e^{-\frac{(z - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+\tau)^{-4} \left(1 + \frac{(r - \mathbf{c}\tau)^2}{1+\tau} \right)^{-2} rz \frac{1}{|x|} dz dr d\tau \\ &\lesssim \int_{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}}^{\frac{t}{2}} \int_0^{\infty} (1+t-\tau)^{-\frac{5}{2} + \frac{3}{2}} (1+\tau)^{-4} \left(1 + \frac{(r - \mathbf{c}\tau)^2}{1+\tau} \right)^{-2} \frac{r}{|x|} dr d\tau \\ &\lesssim (1+t)^{-\frac{5}{2} + \frac{3}{2}} |x|^{-1} \int_{\frac{t}{4} - \frac{|x|}{4\mathbf{c}}}^{\frac{t}{2}} (1+\tau)^{-4 + \frac{3}{2}} d\tau \\ &\lesssim (1+t)^{-2} (1 + \mathbf{c}t - |x|)^{-\frac{3}{2}} \lesssim (1+t)^{-2} (1 + \mathbf{c}t - |x|)^{\frac{1}{2}} (1 + \mathbf{c}t - |x|)^{-2} \\ &\lesssim (1+t)^{-5/2} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t} \right)^{-1}, \end{aligned}$$

by setting $z = \sqrt{|x|^2 + r^2 - 2r|x|\cos\theta}$ and $\sin\theta d\theta = \frac{z}{r|x|} dz$.

For J_3 , we use the spherical coordinates again to obtain

$$\begin{aligned} J_3 &\lesssim \int_{\frac{t}{2}}^{\frac{3|x|}{4\mathbf{c}} + \frac{t}{4}} \int_0^{\infty} \int_0^{\pi} (1+t-\tau)^{-5/2} e^{-\frac{(\sqrt{|x|^2 + r^2 - 2r|x|\cos\theta} - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+t)^{-4} \left(1 + \frac{(r - \mathbf{c}\tau)^2}{1+\tau} \right)^{-2} r^2 \sin\theta d\theta dr d\tau \\ &\lesssim \int_{\frac{t}{2}}^{\frac{3|x|}{4\mathbf{c}} + \frac{t}{4}} \int_0^{\infty} \int_{||x|-r|}^{|x|+r} (1+t-\tau)^{-5/2} e^{-\frac{(z - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+t)^{-4} \left(1 + \frac{(r - \mathbf{c}\tau)^2}{1+\tau} \right)^{-2} rz \frac{1}{|x|} dz dr d\tau \\ &\lesssim (1+t)^{-4} |x|^{-1} \int_{\frac{t}{2}}^{\frac{3|x|}{4\mathbf{c}} + \frac{t}{4}} \int_0^{\infty} (1+t-\tau)^{-\frac{5}{2} + \frac{3}{2}} \left(1 + \frac{(r - \mathbf{c}\tau)^2}{1+\tau} \right)^{-2} r dr d\tau \end{aligned}$$

$$\begin{aligned}
&\lesssim (1+t)^{-7/2} \int_{\frac{t}{2}}^{\frac{3|x|}{4c} + \frac{t}{4}} (1+t-\tau)^{-1} d\tau \\
&\lesssim (1+t)^{-7/2} \left[\ln \left(1 + \frac{t}{2} \right) - \ln \left(1 + \frac{ct - |x|}{4c} \right) \right] \\
&\lesssim (1+t)^{-5/2} \left(1 + \frac{(ct - |x|)^2}{1+t} \right)^{-1}.
\end{aligned}$$

For J_4 , similar to J_3 , we have

$$\begin{aligned}
J_4 &\lesssim (1+t)^{-7/2} \int_{\frac{3|x|}{4c} + \frac{t}{4}}^{\frac{|x|}{4c} + \frac{3t}{4}} (1+t-\tau)^{-1} d\tau \\
&\lesssim (1+t)^{-7/2} \left[\left(1 + \frac{3(ct - |x|)}{4c} \right)^{-1} \left(1 + \frac{(ct - |x|)}{2c} \right) \right] \\
&\lesssim (1+t)^{-5/2} \left(1 + \frac{(ct - |x|)^2}{1+t} \right)^{-1}.
\end{aligned}$$

For J_5 , we decompose \mathbb{R}^3 into two parts

$$J_5 = \int_{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{c})}^t \left(\int_{|y| \leq \frac{|x|+ct}{2}} + \int_{|y| > \frac{|x|+ct}{2}} \right) (\dots) dy d\tau =: J_{51} + J_{52}.$$

If $\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{c}) \leq \tau \leq t$, $|y| \leq \frac{|x|+ct}{2}$, then

$$c\tau - |y| \geq \frac{ct}{2} + \frac{1}{4}(ct + |x|) - \frac{|x| + ct}{2} \geq \frac{ct - |x|}{4}.$$

If $\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{c}) \leq \tau \leq t$, $|y| > \frac{|x|+ct}{2}$, then

$$|x - y| - c(t - \tau) \geq |y| - |x| - c(t - \tau) \geq \frac{|x| + ct}{2} - |x| - ct + \frac{ct}{2} + \frac{1}{4}(ct + |x|) \geq \frac{ct - |x|}{4}.$$

Hence,

$$\begin{aligned}
J_{51} &\lesssim \int_{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{c})}^t \int_{|y| \leq \frac{|x|+ct}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y|-c(t-\tau))^2}{D_0(1+t-\tau)}} (1+t)^{-4} \left(1 + \frac{(ct - |x|)^2}{1+t} \right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-4} \left(1 + \frac{(ct - |x|)^2}{1+t} \right)^{-2} (ct - |x|) \\
&\lesssim (1+t)^{-3} \left(1 + \frac{(ct - |x|)^2}{1+t} \right)^{-2},
\end{aligned}$$

$$\begin{aligned}
J_{52} &\lesssim \int_{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{c})}^t \int_{|y| > \frac{|x|+ct}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(ct-|x|)^2}{2D_0(1+ct-|x|)}} e^{-\frac{(|x-y|-c(t-\tau))^2}{2D_0(1+t-\tau)}} (1+t)^{-4} dy d\tau \\
&\lesssim (1+t)^{-4} e^{-\frac{(ct-|x|)}{2D_0}} \int_{\frac{t}{2} + \frac{1}{4}(t + \frac{|x|}{c})}^t (1+t-\tau)^{-\frac{5}{2} + \frac{5}{2}} d\tau \\
&\lesssim (1+t)^{-3} e^{-\frac{(ct-|x|)}{2D_0}}.
\end{aligned}$$

Gathering all the estimates, we have

$$J \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t} \right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(ct - |x|)^2}{1+t} \right)^{-1}.$$

□

Proof of Lemma 26. (Huygens wave convolved with diffusion wave).

Case 1: $(x, t) \in D_1$. Direct computation gives

$$\begin{aligned}
K &\lesssim (1+t)^{-5/2} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+\tau)^{-3} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3} dy d\tau \\
&\quad + (1+t)^{-3} \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-5/2} \left(1 + \frac{(|x-y| - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}\right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-5/2} \int_0^{\frac{t}{2}} (1+\tau)^{-3} (1+\tau)^{\frac{3}{2}} dy d\tau + (1+t)^{-3} \int_{\frac{t}{2}}^t (1+t-\tau)^{-5/2} (1+t-\tau)^{\frac{5}{2}} d\tau \\
&\lesssim (1+t)^{-5/2} + (1+t)^{-3} (1+t) \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.
\end{aligned}$$

Case 2: $(x, t) \in D_2$. We split the integral K into two parts

$$K = \left(\int_0^{\frac{3}{4}t} + \int_{\frac{3}{4}t}^t \right) \int_{\mathbb{R}^3} (\dots) dy d\tau =: K_1 + K_2.$$

For K_1 , it is easy to see

$$\begin{aligned}
K_1 &\lesssim \int_0^{\frac{3}{4}t} \int_{\mathbb{R}^3} (1+t)^{-5/2} (1+\tau)^{-3} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3} dy d\tau \\
&\lesssim \int_0^{\frac{3}{4}t} (1+t)^{-5/2} (1+\tau)^{-3} (1+\tau)^{\frac{3}{2}} d\tau \lesssim (1+t)^{-5/2} \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}.
\end{aligned}$$

For K_2 , we decompose \mathbb{R}^3 into two parts

$$K_2 = \int_{\frac{3}{4}t}^t \left(\int_{|y| \leq \frac{|x|}{2}} + \int_{|y| > \frac{|x|}{2}} \right) (\dots) dy d\tau =: K_{21} + K_{22}.$$

It readily follows that

$$\begin{aligned}
K_{22} &\lesssim (1+t)^{-3} \left(1 + \frac{|x|^2}{1+t}\right)^{-3} \int_{\frac{3}{4}t}^t \int_{|y| > \frac{|x|}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y| - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} dy d\tau \\
&\lesssim (1+t)^{-3} \left(1 + \frac{|x|^2}{1+t}\right)^{-3} \int_{\frac{3}{4}t}^t (1+t-\tau)^{-5/2} (1+t-\tau)^{\frac{5}{2}} d\tau \\
&\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3}.
\end{aligned}$$

If $\frac{3}{4}t \leq \tau \leq t$, $|y| \leq \frac{|x|}{2}$, then

$$\begin{aligned}
|x-y| - \mathbf{c}(t-\tau) &\geq |x| - |y| - \mathbf{c}(t-\tau) \geq \frac{|x|}{2} - \frac{\mathbf{c}t}{4} \geq \frac{1}{2}(\mathbf{c}t - \sqrt{1+t}) - \frac{\mathbf{c}t}{4} \\
&\geq \frac{\mathbf{c}t}{8} + \left(\frac{\mathbf{c}t}{8} - \frac{\sqrt{1+t}}{2}\right) \geq \frac{\mathbf{c}t}{8}
\end{aligned}$$

for $t \geq 11$. Hence, there exists a universal constant $C > 0$ such that

$$K_{21} \lesssim (1+t)^{-3} e^{-\frac{t}{C}} \int_{\frac{3}{4}t}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-5/2} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3} dy d\tau$$

$$\begin{aligned}
&\lesssim (1+t)^{-3} e^{-\frac{t}{C}} \int_{\frac{3}{4}t}^t (1+t-\tau)^{-5/2} (1+\tau)^{\frac{3}{2}} d\tau \\
&\lesssim e^{-\frac{t}{C}} \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}
\end{aligned}$$

for $t \geq 11$, and

$$\begin{aligned}
K_{21} &\lesssim (1+t)^{-3} \int_{\frac{3}{4}t}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-5/2} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3} dy d\tau \\
&\lesssim C \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}
\end{aligned}$$

for $0 \leq t \leq 11$. Therefore we can conclude that

$$K \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3} + (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}.$$

Case 3: $(x, t) \in D_3$. We split the integral K into four parts

$$\begin{aligned}
K &= \int_0^{\frac{t}{2}} \left(\int_{|y| \leq \frac{|x| - \mathbf{c}(t-\tau)}{2}} + \int_{|y| > \frac{|x| - \mathbf{c}(t-\tau)}{2}} \right) (\cdots) dy d\tau + \int_{\frac{t}{2}}^t \left(\int_{|y| \leq \frac{|x| - \mathbf{c}(t-\tau)}{2}} + \int_{|y| > \frac{|x| - \mathbf{c}(t-\tau)}{2}} \right) (\cdots) dy d\tau \\
&= K_{11} + K_{12} + K_{21} + K_{22}.
\end{aligned}$$

If $|y| \leq \frac{|x| - \mathbf{c}(t-\tau)}{2}$, then

$$|x - y| - \mathbf{c}(t - \tau) \geq |x| - |y| - \mathbf{c}(t - \tau) \geq \frac{|x| - \mathbf{c}(t - \tau)}{2} = \frac{|x| - \mathbf{c}t + \mathbf{c}\tau}{2}.$$

Hence,

$$\begin{aligned}
K_{11} &\lesssim \int_0^{\frac{t}{2}} \int_{|y| \leq \frac{|x| - \mathbf{c}(t-\tau)}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2 + (\mathbf{c}\tau)^2}{4D_0(1+t-\tau)}} (1+\tau)^{-3} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3} dy d\tau \\
&\lesssim (1+t)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2}{4D_0(1+t)}} \int_0^{\frac{t}{2}} (1+\tau)^{-3} (1+\tau)^{\frac{3}{2}} dy d\tau \lesssim (1+t)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2}{4D_0(1+t)}},
\end{aligned}$$

$$\begin{aligned}
K_{12} &\lesssim (1+t)^{-5/2} \int_0^{\frac{t}{2}} \int_{|y| > \frac{|x| - \mathbf{c}(t-\tau)}{2}} (1+\tau + |y|^2)^{-3} dy d\tau \\
&\lesssim (1+t)^{-5/2} \int_0^{\frac{t}{2}} \int_{\frac{|x| - \mathbf{c}(t-\tau)}{2}}^{\infty} (1+r)^{-6} r^2 dr d\tau \lesssim (1+t)^{-5/2} \int_0^{\frac{t}{2}} (1+|x| - \mathbf{c}t + \mathbf{c}\tau)^{-3} d\tau \\
&\lesssim (1+t)^{-5/2} (1+|x| - \mathbf{c}t)^{-2} \lesssim (1+t)^{-7/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1},
\end{aligned}$$

$$\begin{aligned}
K_{21} &\lesssim \int_{\frac{t}{2}}^t \int_{|y| \leq \frac{|x| - \mathbf{c}(t-\tau)}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2 + (\mathbf{c}\tau)^2}{4D_0(1+t-\tau)}} (1+\tau)^{-3} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3} dy d\tau \\
&\lesssim e^{-\frac{(|x| - \mathbf{c}t)^2}{4D_0(1+t)}} e^{-\frac{t}{C}} \int_{\frac{t}{2}}^t (1+t-\tau)^{-5/2} (1+\tau)^{-\frac{3}{2}} d\tau \lesssim e^{-\frac{t}{C}} e^{-\frac{(|x| - \mathbf{c}t)^2}{4D_0(1+t)}}
\end{aligned}$$

for some $C > 0$, and

$$K_{22} \lesssim (1+t)^{-3} \left(1 + \frac{(|x| - \mathbf{c}t)^2 + \mathbf{c}^2 t^2}{1+t}\right)^{-3} \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y| - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} dy d\tau$$

$$\begin{aligned}
&\lesssim (1+t)^{-3} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t} + t\right)^{-3} \int_{\frac{t}{2}}^t (1+t-\tau)^{-5/2} (1+t-\tau)^{\frac{5}{2}} d\tau \\
&\lesssim (1+t)^{-4} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}.
\end{aligned}$$

It implies that

$$K \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}.$$

Case 4: $(x, t) \in D_4$. First note that $\sqrt{1+t} \leq |x| \leq \mathbf{c}t/2$ in this region. Now we split the integral K into three parts

$$K = \left(\int_0^{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}} + \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{t - \frac{|x|}{2\mathbf{c}}} + \int_{t - \frac{|x|}{2\mathbf{c}}}^t \right) \int_{\mathbb{R}^3} (\dots) dyd\tau =: K_1 + K_2 + K_3.$$

For K_1 , we decompose \mathbb{R}^3 into two parts

$$K_1 = \int_0^{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}} \left(\int_{|y| \leq \frac{\mathbf{c}(t-\tau) - |x|}{2}} + \int_{|y| > \frac{\mathbf{c}(t-\tau) - |x|}{2}} \right) (\dots) dyd\tau =: K_{11} + K_{12}.$$

If $0 \leq \tau \leq \frac{t}{2} - \frac{|x|}{2\mathbf{c}}$, $|y| \leq \frac{\mathbf{c}(t-\tau) - |x|}{2}$, then

$$\mathbf{c}(t-\tau) - |x| - |y| \geq \frac{\mathbf{c}(t-\tau) - |x|}{2} \geq \frac{\mathbf{c}t - |x|}{4}.$$

If $0 \leq \tau \leq \frac{t}{2} - \frac{|x|}{2\mathbf{c}}$, $|y| > \frac{\mathbf{c}(t-\tau) - |x|}{2}$, then

$$|y| > \frac{\mathbf{c}t - |x|}{4}.$$

Hence, we have

$$\begin{aligned}
K_{11} &\lesssim \left(1 + \frac{t}{2} + \frac{|x|}{2\mathbf{c}}\right)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2}{16D_0(1+t)}} \int_0^{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}} \int_{|y| \leq \frac{\mathbf{c}(t-\tau) - |x|}{2}} (1+\tau)^{-3} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-3} dyd\tau \\
&\lesssim (1+t)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2}{16D_0(1+t)}} \int_0^{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}} (1+\tau)^{-3+\frac{3}{2}} d\tau \lesssim (1+t)^{-5/2} e^{-\frac{(|x| - \mathbf{c}t)^2}{4D_0(1+t)}},
\end{aligned}$$

$$\begin{aligned}
K_{12} &\lesssim (1+t)^{-5/2} \int_0^{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}} \int_{|y| \geq \frac{\mathbf{c}(t-\tau) - |x|}{2}} (1+\tau)^{-3} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3} dyd\tau \\
&\lesssim (1+t)^{-5/2} \int_0^{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}} \int_{\frac{\mathbf{c}(t-\tau) - |x|}{2}}^{\infty} r^{-6} r^2 dr d\tau \lesssim (1+t)^{-5/2} (1 + \mathbf{c}t - |x|)^{-2} \\
&\lesssim (1+t)^{-7/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1},
\end{aligned}$$

so that

$$K_1 \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}.$$

For K_2 , we use the spherical coordinates to obtain

$$\begin{aligned}
K_2 &= \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^t \int_0^{\infty} \int_0^{\pi} (1+t-\tau)^{-5/2} e^{-\frac{(\sqrt{|x|^2 + r^2 - 2r|x|\cos\theta} - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+\tau)^{-3} \left(1 + \frac{r^2}{1+\tau}\right)^{-3} r^2 \sin\theta d\theta dr d\tau \\
&= \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^t \int_0^{\infty} \int_{||x|-r|}^{|x|+r} (1+t-\tau)^{-5/2} e^{-\frac{(z - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+\tau)^{-3} \left(1 + \frac{r^2}{1+\tau}\right)^{-3} r^2 \frac{dz}{r|x|} dr d\tau
\end{aligned}$$

$$\begin{aligned}
&\lesssim \int_{\frac{t}{2} - \frac{|x|}{2c}}^{t - \frac{|x|}{2c}} \int_0^\infty \int_{||x|-r|}^{|x|+r} (1+|x|)^{-5/2} e^{-\frac{c\tau - (ct-z)^2}{D_0(1+t)}} (1+t)^{-3} \left(1 + \frac{r^2}{1+t}\right)^{-3} r \frac{zdz}{|x|} dr d\tau \\
&\lesssim \frac{1}{|x|} \int_0^\infty \int_{||x|-r|}^{|x|+r} (1+|x|)^{-5/2} \sqrt{1+t} (1+t)^{-3} \left(1 + \frac{r^2}{1+t}\right)^{-3} r z dz dr \\
&\lesssim (1+|x|)^{-7/2} (1+t)^{-\frac{5}{2}} \int_0^\infty \int_{||x|-r|}^{|x|+r} \left(1 + \frac{r^2}{1+t}\right)^{-3} r z dz dr \\
&\lesssim (1+|x|)^{-7/2} (1+t)^{-\frac{5}{2}} \left[\int_0^{|x|} \int_{|x|-r}^{|x|+r} z r \left(1 + \frac{r^2}{1+t}\right)^{-3} dz dr + \int_{|x|}^\infty \int_{r-|x|}^{r+|x|} z r \left(1 + \frac{r^2}{1+t}\right)^{-3} dz dr \right] \\
&\lesssim (1+|x|)^{-7/2} (1+t)^{-\frac{5}{2}} \left[\int_0^{|x|} r^2 |x| \left(1 + \frac{r^2}{1+t}\right)^{-3} dr + \int_{|x|}^\infty r^2 |x| \left(1 + \frac{r^2}{1+t}\right)^{-3} dr \right] \\
&\lesssim (1+|x|)^{-7/2} (1+t)^{-\frac{5}{2}} \left[|x| (1+t)^{\frac{3}{2}} + |x| (1+t)^{\frac{3}{2}} \left(1 + \frac{|x|}{\sqrt{1+t}}\right)^{-3} \right] \\
&\lesssim (1+|x|)^{-5/2} (1+t)^{-1} \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.
\end{aligned}$$

by setting $z = \sqrt{|x|^2 + r^2 - 2r|x|\cos\theta}$ and $\sin\theta d\theta = \frac{z}{r|x|} dz$.

For K_3 , we split \mathbb{R}^3 into two parts

$$K_3 = \int_{t - \frac{|x|}{2c}}^t \left(\int_{|y| \leq \frac{|x| - c(t-\tau)}{2}} + \int_{|y| > \frac{|x| - c(t-\tau)}{2}} \right) (\dots) dy d\tau =: K_{31} + K_{32}.$$

If $t - \frac{|x|}{2c} < \tau \leq t$, $|y| \leq \frac{|x| - c(t-\tau)}{2}$, then

$$|x-y| - c(t-\tau) \geq |x| - |y| - c(t-\tau) \geq \frac{|x| - c(t-\tau)}{2} \geq \frac{|x|}{4}.$$

If $t - \frac{|x|}{2c} < \tau \leq t$, $|y| > \frac{|x| - c(t-\tau)}{2}$, then

$$|y| > \frac{|x| - c(t-\tau)}{2} \geq \frac{|x|}{4}.$$

Hence,

$$\begin{aligned}
K_{31} &\lesssim (1+|x|^2)^{-\frac{5}{2}} \int_{t - \frac{|x|}{2c}}^t \int_{\mathbb{R}^3} (1+\tau)^{-3} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3} dy d\tau \\
&\lesssim (1+|x|^2)^{-\frac{5}{2}} \int_{t - \frac{|x|}{2c}}^t (1+\tau)^{-\frac{3}{2}} d\tau \\
&\lesssim (1+t)^{-3} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}, \\
K_{32} &\lesssim \left(1+t - \frac{|x|}{2c}\right)^{-3} \left(1 + \frac{|x|^2}{1+t}\right)^{-3} \int_{t - \frac{|x|}{2c}}^t \int_{|y| > \frac{|x| - c(t-\tau)}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y| - c(t-\tau))^2}{D_0(1+t-\tau)}} dy d\tau \\
&\lesssim \left(1+t - \frac{|x|}{2c}\right)^{-3} \left(1 + \frac{|x|^2}{1+t}\right)^{-3} \int_{t - \frac{|x|}{2c}}^t (1+t-\tau)^{-5/2} (1+t-\tau)^{\frac{5}{2}} d\tau \\
&\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-3},
\end{aligned}$$

so that

$$K_3 \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.$$

To sum up,

$$K \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}.$$

Case 5: $(x, t) \in D_5$. We split the integral K into three parts

$$K = \left(\int_0^{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}} + \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{\frac{3}{2}(t - \frac{|x|}{\mathbf{c}})} + \int_{\frac{3}{2}(t - \frac{|x|}{\mathbf{c}})}^t \right) \int_{\mathbb{R}^3} (\dots) dyd\tau =: K_1 + K_2 + K_3.$$

For K_1 , the estimate is the same as the term K_1 of Case 4 and so

$$K_1 \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}.$$

For K_2 , we use the spherical coordinates to obtain

$$\begin{aligned} K_2 &\lesssim \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{\frac{3}{2}(t - \frac{|x|}{\mathbf{c}})} \int_{\mathbb{R}^3} (1+t)^{-5/2} e^{-\frac{(|x-y| - \mathbf{c}(t-\tau))^2}{D_0(1+t-\tau)}} (1+\tau)^{-3} \left(1 + \frac{|y|^2}{1+\tau}\right)^{-3} dyd\tau \\ &\lesssim \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{\frac{3}{2}(t - \frac{|x|}{\mathbf{c}})} \int_0^\infty \int_0^\pi (1+t)^{-5/2} e^{-\frac{(\sqrt{|x|^2+r^2-2r|x|\cos\theta} - \mathbf{c}(t-\tau))^2}{D_0(1+t)}} (1+\tau)^{-3} \left(1 + \frac{r^2}{1+\tau}\right)^{-3} r^2 \sin\theta d\theta dr d\tau \\ &\lesssim \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{\frac{3}{2}(t - \frac{|x|}{\mathbf{c}})} \int_0^\infty \int_{||x|-r|}^{|x|+r} (1+t)^{-5/2} e^{-\frac{(z - \mathbf{c}(t-\tau))^2}{D_0(1+t)}} z (1 + \mathbf{c}t - |x|)^{-3} \left(1 + \frac{r^2}{1+\tau}\right)^{-3} \frac{r}{|x|} dz dr d\tau \\ &\lesssim (1+t)^{-7/2} (1 + \mathbf{c}t - |x|)^{-3} \int_{\frac{t}{2} - \frac{|x|}{2\mathbf{c}}}^{\frac{3}{2}(t - \frac{|x|}{\mathbf{c}})} \int_0^\infty \int_{||x|-r|}^{|x|+r} e^{-\frac{(c\tau - (ct-z))^2}{D_0(1+t)}} z \left(1 + \frac{r^2}{1+\tau}\right)^{-3} rdz dr d\tau \\ &\lesssim (1+t)^{-7/2} (1 + \mathbf{c}t - |x|)^{-3} (1+t)^{\frac{1}{2}} \\ &\quad \cdot \left[\int_0^{|x|} \int_{|x|-r}^{|x|+r} z \left(1 + \frac{r^2}{1 + \mathbf{c}t - |x|}\right)^{-3} rdz dr + \int_{|x|}^\infty \int_{r-|x|}^{r+|x|} z \left(1 + \frac{r^2}{1 + \mathbf{c}t - |x|}\right)^{-3} rdz dr \right] \\ &\lesssim (1+t)^{-3} (1 + \mathbf{c}t - |x|)^{-3} \left[\int_0^{|x|} r^2 |x| \left(1 + \frac{r^2}{1 + \mathbf{c}t - |x|}\right)^{-3} dr + \int_{|x|}^\infty r^2 |x| \left(1 + \frac{r^2}{1 + \mathbf{c}t - |x|}\right)^{-3} dr \right] \\ &\lesssim (1+t)^{-3} (1 + \mathbf{c}t - |x|)^{-3} \left[|x| (1 + \mathbf{c}t - |x|)^{\frac{3}{2}} + |x| (1 + \mathbf{c}t - |x|)^3 (1 + |x|)^{-3} \right] \\ &\lesssim (1+t)^{-2} (1 + \mathbf{c}t - |x|)^{-\frac{3}{2}} \lesssim (1+t)^{-2} (1 + \mathbf{c}t - |x|)^{\frac{1}{2}} (1 + \mathbf{c}t - |x|)^{-2} \\ &\lesssim (1+t)^{-5/2} \left(1 + \frac{(\mathbf{c}t - |x|)^2}{1+t}\right)^{-1}. \end{aligned}$$

For K_3 , we decompose \mathbb{R}^3 into two parts

$$K_3 = \int_{\frac{3}{2}(t - \frac{|x|}{\mathbf{c}})}^t \left(\int_{|y| \leq \frac{|x| - \mathbf{c}(t-\tau)}{2}} + \int_{|y| > \frac{|x| - \mathbf{c}(t-\tau)}{2}} \right) (\dots) dyd\tau = K_{31} + K_{32}.$$

If $\frac{3}{2}(t - \frac{|x|}{\mathbf{c}}) \leq \tau \leq t$, $|y| \leq \frac{|x| - \mathbf{c}(t-\tau)}{2}$, then

$$|x-y| - \mathbf{c}(t-\tau) \geq |x| - |y| - \mathbf{c}(t-\tau) \geq \frac{|x| - \mathbf{c}(t-\tau)}{2} \geq \frac{\mathbf{c}t - |x|}{4}.$$

If $\frac{3}{2}\left(t - \frac{|x|}{c}\right) \leq \tau \leq t$, $|y| > \frac{|x| - c(t-\tau)}{2}$, then

$$|y| > \frac{|x| - c(t-\tau)}{2} \geq \frac{ct - |x|}{4}.$$

Hence, we use the spherical coordinates to obtain

$$\begin{aligned} K_{31} &\lesssim \int_{\frac{3}{2}\left(t - \frac{|x|}{c}\right)}^t \int_0^{\frac{|x| - c(t-\tau)}{2}} \int_0^\pi (1+t-\tau)^{-5/2} e^{-\frac{(ct-|x|)^2}{32D_0(1+t-\tau)}} e^{-\frac{(\sqrt{|x|^2+r^2-2r|x|\cos\theta}-c(t-\tau))^2}{2D_0(1+t-\tau)}} \\ &\quad \cdot (1+\tau)^{-3} \left(1 + \frac{r^2}{1+\tau}\right)^{-3} r^2 \sin\theta d\theta dr d\tau \\ &\lesssim e^{-\frac{(ct-|x|)^2}{32D_0(1+t)}} \int_{\frac{3}{2}\left(t - \frac{|x|}{c}\right)}^t \int_0^{\frac{|x| - c(t-\tau)}{2}} \int_{|x|-r}^{|x|+r} (1+t-\tau)^{-5/2} e^{-\frac{(z-c(t-\tau))^2}{2D_0(1+t-\tau)}} z (1+\tau)^{-3} \\ &\quad \cdot \left(1 + \frac{r^2}{1+\tau}\right)^{-3} \frac{r}{|x|} dz dr d\tau \\ &\lesssim \frac{1}{|x|} e^{-\frac{(ct-|x|)^2}{32D_0(1+t)}} \int_{\frac{3}{2}\left(t - \frac{|x|}{c}\right)}^t (1+t-\tau)^{-5/2} (1+\tau)^{-3} (1+t-\tau)^{\frac{3}{2}} (1+\tau) d\tau \\ &\lesssim \frac{1}{|x|} e^{-\frac{(ct-|x|)^2}{32D_0(1+t)}} \left(\int_{\frac{3}{2}\left(t - \frac{|x|}{c}\right)}^{\frac{1}{2}\left(t + \frac{3}{2}\left(t - \frac{|x|}{c}\right)\right)} + \int_{\frac{1}{2}\left(t + \frac{3}{2}\left(t - \frac{|x|}{c}\right)\right)}^t \right) (1+t-\tau)^{-1} (1+\tau)^{-2} d\tau \\ &\lesssim \frac{e^{-\frac{(ct-|x|)^2}{32D_0(1+t)}}}{|x|} \left[\left(1 + \frac{3|x|}{4c} - \frac{t}{4}\right)^{-1} \left(1 + \frac{3}{2}\left(t - \frac{|x|}{c}\right)\right)^{-1} + (1+t)^{-2} \ln\left(1 + \frac{3|x|}{4c} - \frac{t}{4}\right) \right] \\ &\lesssim \frac{e^{-\frac{(ct-|x|)^2}{32D_0(1+t)}}}{1+t} (1+t)^{-3/2} \lesssim (1+t)^{-5/2} \left(1 + \frac{(ct-x)^2}{1+t}\right)^{-1}, \end{aligned}$$

by setting $z = \sqrt{|x|^2 + r^2 - 2r|x|\cos\theta}$ and $\sin\theta d\theta = \frac{z}{r|x|} dz$. As for K_{32} , direct computation gives

$$\begin{aligned} K_{32} &\lesssim \int_{\frac{3}{2}\left(t - \frac{|x|}{c}\right)}^t \int_{|y| > \frac{|x| - c(t-\tau)}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y|-c(t-\tau))^2}{D_0(1+t-\tau)}} (1+|y|^2)^{-3} dy d\tau \\ &\lesssim \left(\int_{\frac{3}{2}\left(t - \frac{|x|}{c}\right)}^{\frac{1}{2}\left(t + \frac{3}{2}\left(t - \frac{|x|}{c}\right)\right)} + \int_{\frac{1}{2}\left(t + \frac{3}{2}\left(t - \frac{|x|}{c}\right)\right)}^t \right) \int_{|y| > \frac{|x| - c(t-\tau)}{2}} (\dots) dy d\tau \\ &= K_{321} + K_{322}, \end{aligned}$$

we then have

$$\begin{aligned} K_{321} &\lesssim \int_{\frac{3}{2}\left(t - \frac{|x|}{c}\right)}^{\frac{1}{2}\left(t + \frac{3}{2}\left(t - \frac{|x|}{c}\right)\right)} \int_{|y| > \frac{|x| - c(t-\tau)}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y|-c(t-\tau))^2}{D_0(1+t-\tau)}} (1+|y|^2)^{-3} dy d\tau \\ &\lesssim \int_{\frac{3}{2}\left(t - \frac{|x|}{c}\right)}^{\frac{1}{2}\left(t + \frac{3}{2}\left(t - \frac{|x|}{c}\right)\right)} (1+t-\tau)^{-5/2} (1+|x| - c(t-\tau))^{-3} d\tau \\ &\lesssim (1+t)^{-5/2} (1+ct - |x|)^{-2} \\ &\lesssim (1+t)^{-7/2} \left(1 + \frac{(ct - |x|)^2}{1+t}\right)^{-1}. \end{aligned}$$

and

$$K_{322} \lesssim \int_{\frac{1}{2}\left(t + \frac{3}{2}\left(t - \frac{|x|}{c}\right)\right)}^t \int_{|y| > \frac{|x| - c(t-\tau)}{2}} (1+t-\tau)^{-5/2} e^{-\frac{(|x-y|-c(t-\tau))^2}{D_0(1+t-\tau)}} (1+|y|^2)^{-3} dy d\tau$$

$$\begin{aligned}
&\lesssim \int_{\frac{1}{2}(t+\frac{3}{2}(t-\frac{|x|}{c}))}^t (1+|x|-\mathbf{c}(t-\tau))^{-6} d\tau \\
&\lesssim (1+|x|+\mathbf{c}t)^{-5} \\
&\lesssim (1+t)^{-\frac{7}{2}} \left(1+\frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.
\end{aligned}$$

It implies that

$$K_3 \lesssim (1+t)^{-5/2} \left(1+\frac{(\mathbf{c}t-|x|)^2}{1+t}\right)^{-1} + (1+t)^{-2} \left(1+\frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.$$

As a consequence,

$$K \lesssim (1+t)^{-5/2} \left(1+\frac{(\mathbf{c}t-|x|)^2}{1+t}\right)^{-1} + (1+t)^{-2} \left(1+\frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.$$

□

Proof of Lemma 27. (Diffusion wave convolution with Huygens wave).

Case 1: $(x, t) \in D_1$. We split the integral into two parts to obtain

$$\begin{aligned}
L &= \left(\int_0^{\frac{t}{2}} + \int_{\frac{t}{2}}^t \right) \int_{\mathbb{R}^3} (\dots) dy d\tau \\
&\lesssim (1+t)^{-2} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+\tau)^{-4} \left(1+\frac{(|y|-\mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\
&\quad + (1+t)^{-4} \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-2} \left(1+\frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-2} \int_0^{\frac{t}{2}} (1+\tau)^{-4} (1+\tau)^{\frac{5}{2}} d\tau + (1+t)^{-4} \int_{t/2}^t (1+t-\tau)^{-\frac{1}{2}} d\tau \\
&\lesssim (1+t)^{-2} \lesssim (1+t)^{-2} \left(1+\frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.
\end{aligned}$$

Case 2: $(x, t) \in D_2$. We split the integral into two parts:

$$L = \left(\int_0^{\frac{t}{2}} + \int_{\frac{t}{2}}^t \right) \int_{\mathbb{R}^3} (\dots) dy d\tau =: L_1 + L_2.$$

First one can see

$$\begin{aligned}
L_2 &\lesssim (1+t)^{-4} \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-2} \left(1+\frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-2} dy d\tau \\
&\lesssim (1+t)^{-7/2} \left(1+\frac{(|x|-\mathbf{c}t)^2}{1+t}\right)^{-1}.
\end{aligned}$$

As for L_1 , we further decompose \mathbb{R}^3 into two parts:

$$L_1 = \int_0^{\frac{t}{2}} \left(\int_{|y| \leq \frac{2}{3}|x|} + \int_{|y| > \frac{2}{3}|x|} \right) (\dots) dy d\tau =: L_{11} + L_{12}.$$

If $|y| \leq \frac{2}{3}|x|$, then we have

$$|x-y| \geq |x| - |y| \geq \frac{|x|}{3},$$

and thus

$$\begin{aligned} L_{11} &\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-2} \int_0^{\frac{t}{2}} \int_{|y| \leq \frac{2}{3}|x|} (1+\tau)^{-4} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dyd\tau \\ &\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-2} \int_0^{\frac{t}{2}} (1+\tau)^{-4} (1+\tau)^{\frac{5}{2}} d\tau \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-2}. \end{aligned}$$

If $|y| \geq \frac{2}{3}|x|$ and $0 \leq \tau \leq \frac{t}{2}$, then

$$|y| - \mathbf{c}\tau \geq \frac{2}{3}|x| - \mathbf{c}\tau \geq \frac{2}{3}(\mathbf{c}t - \sqrt{1+t}) - \frac{\mathbf{c}t}{2} \geq \frac{\mathbf{c}t}{12} + \frac{\mathbf{c}t}{12} - \frac{2}{3}\sqrt{1+t} \geq \frac{\mathbf{c}t}{12}$$

for $t \geq 40$. Hence,

$$\begin{aligned} &\int_0^{\frac{t}{2}} \int_{|y| \geq \frac{2}{3}|x|} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-2} (1+\tau)^{-4} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dyd\tau \\ &\lesssim (1+t)^{-4} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-2} (1+\tau)^{-2} dyd\tau \\ &\lesssim (1+t)^{-9/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1} \end{aligned}$$

whenever $t \geq 40$, and

$$\begin{aligned} &\int_0^{\frac{t}{2}} \int_{|y| \geq \frac{2}{3}|x|} (1+t-\tau)^{-2} \left(1 + \frac{|x-y|^2}{D_0(1+t-\tau)}\right)^{-2} (1+\tau)^{-4} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dyd\tau \\ &\lesssim C \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1} \end{aligned}$$

whenever $0 \leq t \leq 40$. Therefore, we get

$$L_{12} \lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}.$$

Combining all the estimates, we have

$$L \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-2} + (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-1}.$$

Case 3: $(x, t) \in D_3$. We split the integral into two parts

$$L = \left(\int_0^{\frac{t}{2}} + \int_{\frac{t}{2}}^t \right) \int_{\mathbb{R}^3} (\dots) dyd\tau =: L_1 + L_2.$$

For L_1 , we further decompose the space domain into two parts:

$$L_1 = \int_0^{\frac{t}{2}} \left(\int_{|y| \leq \frac{2}{3}|x|} + \int_{|y| > \frac{2}{3}|x|} \right) (\dots) dyd\tau =: L_{11} + L_{12}.$$

If $|y| \leq \frac{2}{3}|x|$, then

$$|x-y| \geq \frac{|x|}{3},$$

and thus

$$L_{11} \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+\tau)^{-4} \left(1 + \frac{(|y| - \mathbf{c}\tau)^2}{1+\tau}\right)^{-2} dyd\tau$$

$$\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}}.$$

If $|y| \geq \frac{2}{3}|x|$ and $0 \leq \tau \leq \frac{t}{2}$, we have

$$|y| - \mathbf{c}\tau \geq \frac{2}{3}|x| - \frac{\mathbf{c}t}{2} = \frac{1}{6}|x| + \frac{1}{2}(|x| - \mathbf{c}t),$$

so that

$$\begin{aligned} L_{12} &\lesssim \int_0^{\frac{t}{2}} \int_{|y| \geq \frac{2}{3}|x|} (1+t-\tau)^{-2} e^{-\frac{|x-y|^2}{D_0(1+t-\tau)}} (1+\tau)^{-2} \left(1+\tau + (|x| - \mathbf{c}t)^2\right)^{-2} dyd\tau \\ &\lesssim \left(1 + (|x| - \mathbf{c}t)^2\right)^{-2} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+t-\tau)^{-2} e^{-\frac{|x-y|^2}{D_0(1+t-\tau)}} (1+\tau)^{-2} dyd\tau \\ &\lesssim \left(1 + (|x| - \mathbf{c}t)^2\right)^{-2} (1+t)^{-1/2} \lesssim \left(\frac{1+t}{2} + \frac{(|x| - \mathbf{c}t)^2}{2}\right)^{-2} (1+t)^{-1/2} \\ &\lesssim (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-2} \end{aligned}$$

since $|x| - \mathbf{c}t \geq \sqrt{1+t}$ for $(x, t) \in D_3$. Therefore, we obtain

$$L_1 \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-2}.$$

Next for L_2 , we decompose \mathbb{R}^3 into two parts

$$L_2 = \int_{\frac{t}{2}}^t \left(\int_{|y| - \mathbf{c}\tau \leq \frac{|x| - \mathbf{c}t}{2}} + \int_{|y| - \mathbf{c}\tau > \frac{|x| - \mathbf{c}t}{2}} \right) (\dots) dyd\tau =: L_{21} + L_{22}.$$

If $|y| - \mathbf{c}\tau \leq \frac{|x| - \mathbf{c}t}{2}$, we have

$$|x - y| \geq |x| - \left(\frac{|x| - \mathbf{c}t}{2}\right) - \mathbf{c}\tau \geq \frac{|x| - \mathbf{c}t}{2},$$

so

$$\begin{aligned} L_{21} &\lesssim \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-2} \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-2} e^{-\frac{|x-y|^2}{2D_0(1+t-\tau)}} (1+\tau)^{-4} dyd\tau \\ &\lesssim (1+t)^{-7/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-2}, \end{aligned}$$

and

$$\begin{aligned} L_{22} &\lesssim \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-2} \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-2} e^{-\frac{|x-y|^2}{D_0(1+t-\tau)}} (1+\tau)^{-4} dyd\tau \\ &\lesssim (1+t)^{-7/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-2}. \end{aligned}$$

Therefore,

$$L \lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + (1+t)^{-5/2} \left(1 + \frac{(|x| - \mathbf{c}t)^2}{1+t}\right)^{-2}.$$

Case 4: $(x, t) \in D_4 \cup D_5$. Observe that

$$\frac{\sqrt{1+t}}{2\mathbf{c}} \leq \frac{|x|}{2\mathbf{c}} \leq \frac{t}{2}.$$

We split the integral into three parts:

$$L = \left(\int_0^{\frac{|x|}{2c}} + \int_{\frac{|x|}{2c}}^{\frac{t}{2}} + \int_{\frac{t}{2}}^t \right) \int_{\mathbb{R}^3} (\dots) dy d\tau =: L_1 + L_2 + L_3.$$

For L_1 , we decompose \mathbb{R}^3 into two parts $\{|y| \leq \frac{3}{4}|x|\}$ and $\{|y| > \frac{3}{4}|x|\}$. If $|y| \geq \frac{3}{4}|x|$ and $0 \leq \tau \leq \frac{|x|}{2c}$, then

$$|y| - c\tau \geq \frac{1}{4}|x|.$$

Thus,

$$\begin{aligned} L_1 &\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} \int_0^{\frac{t}{2}} \int_{|y| \leq \frac{3|x|}{4}} (1+\tau)^{-4} \left(1 + \frac{(|y| - c\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\ &\quad + \left(1 + |x|^2\right)^{-2} \int_0^{\frac{t}{2}} \int_{|y| > \frac{3|x|}{4}} (1+t-\tau)^{-2} e^{-\frac{|x-y|^2}{D_0(1+t-\tau)}} (1+\tau)^{-2} dy d\tau \\ &\lesssim (1+t)^{-2} \left(1 + \frac{|x|^2}{1+t}\right)^{-\frac{3}{2}} + \left(\frac{|x|^2}{2} + \frac{1+t}{2}\right)^{-2} (1+t)^{-1/2} \\ &\lesssim (1+t)^{-2} \left(1 + \frac{|x|}{1+t}\right)^{-\frac{3}{2}}. \end{aligned}$$

As for L_2 and L_3 , it immediately follows that

$$\begin{aligned} L_2 &\lesssim \int_{\frac{|x|}{2c}}^{\frac{t}{2}} \int_{\mathbb{R}^3} (1+t-\tau)^{-2} e^{-\frac{|x-y|^2}{D_0(1+t-\tau)}} (1+\tau)^{-4} \left(1 + \frac{(|y| - c\tau)^2}{1+\tau}\right)^{-2} dy d\tau \\ &\lesssim (1+t)^{-2} \left(1 + \frac{|x|}{2c}\right)^{-2} \int_{\frac{|x|}{2c}}^{\frac{t}{2}} (1+\tau)^{-2} \int_{\mathbb{R}^3} e^{-\frac{|x-y|^2}{D_0(1+t-\tau)}} dy d\tau \\ &\lesssim (1+t)^{-2} \left(1 + \frac{|x|}{2c}\right)^{-3} (1+t)^{\frac{3}{2}} \\ &\lesssim (1+t)^{-5/2} \left(1 + \frac{|x|^2}{1+t}\right)^{-2}, \end{aligned}$$

and

$$\begin{aligned} L_3 &\lesssim (1+t)^{-4} \int_{\frac{t}{2}}^t \int_{\mathbb{R}^3} (1+t-\tau)^{-2} e^{-\frac{|x-y|^2}{D_0(1+t-\tau)}} dy d\tau \\ &\lesssim (1+t)^{-7/2} \lesssim \left(1 + |x|^2\right)^{-\frac{3}{2}} (1+t)^{-1/2} \lesssim (1+t)^{-2} \left(1 + \frac{|x|}{1+t}\right)^{-\frac{3}{2}}, \end{aligned}$$

since $\sqrt{1+t} \leq |x| \leq ct$. Therefore, we have

$$L \lesssim (1+t)^{-2} \left(1 + \frac{|x|}{1+t}\right)^{-\frac{3}{2}}.$$

□

Acknowledgments:

This work is partially supported by the National Key R&D Program of China under grant 2022YFA1007300. Y.-C. Lin is supported by the National Science and Technology Council under the grant NSTC 112-2115-M-006-006-MY2. H.-T. Wang is supported by NSFC under Grant No. 12371220 and 12031013, the Strategic Priority Research Program of Chinese Academy of Sciences under Grant No. XDA25010403. K.-C. Wu is supported by the National Science and Technology Council under the grant NSTC 112-2628-M-006-006 -MY4 and National Center for Theoretical Sciences.

REFERENCES

- [1] R. Alonso, Y. Morimoto, W. Sun and T. Yang, Non-cutoff Boltzmann equation with polynomial decay perturbations, *Rev. Mat. Iberoam.*, 37 (2021), 189-292.
- [2] R. Alonso, J. A. Cañizo, I. Gamba, and C. Mouhot, A new approach to the creation and propagation of exponential moments in the Boltzmann equation, *Comm. Partial Differential Equations*, 38 (2013), no. 1, 155-169.
- [3] A. V. Bobylev, Moment inequalities for the Boltzmann equation and applications to spatially homogeneous problems, *J. Statist. Phys.*, 88 (1997), no. 5-6, 1183-1214.
- [4] S. Cameron and S. Snelson, Velocity decay estimates for Boltzmann equation with hard potentials, *Nonlinearity*, 33(6), (2020), 2941.
- [5] C. Cao, Cutoff Boltzmann equation with polynomial perturbation near Maxwellian, *J. Funct. Anal.*, 282 (2022), 179-184.
- [6] C. Cao, R.-J. Duan and Z.-G. Li, Time velocity decay of solution to the Non-cutoff Boltzmann equation in the whole space, *Commun. Math. Anal. Appl.*, 3 (2024), 61-120.
- [7] C. Cercignani, R. Illner and M. Pulvirenti, *The Mathematical Theory of Dilute Gases*, Springer (1994).
- [8] K. Carrapatoso, I. Tristani and K.-C. Wu, Cauchy Problem and Exponential Stability for the Inhomogeneous Landau Equation, *Archive for Rational Mechanics and Analysis*, 221 (2016), 363-418.
- [9] K. Carrapatoso and P. Gervais, Noncutoff Boltzmann equation with soft potentials in the whole space, *Pure and Applied Analysis*, 6 (2024), 253-303.
- [10] S.-J. Deng and S.-H. Yu, Green's function and pointwise convergence for compressible Navier-Stokes equations, *Quart. Appl. Math.*, 75 (2017), No. 3, 433-503.
- [11] L. Desvillettes, Some applications of the method of moments for the homogeneous Boltzmann and Kac equations, *Arch. Rational Mech. Anal.*, 123 (1993), no. 4, 387-404.
- [12] L.-L. Du and Z.-G. Wu, Solving the non-isentropic Navier-Stokes equations in odd space dimensions: the Green function method, *J. Math. Phys.* 58 (2017), No. 10, 101506, 38.
- [13] R.-J. Duan, Z.-G. Li and S.-Q. Liu, Global mild solutions with polynomial tail for the Boltzmann equation in the whole space, arXiv:2212.04676v2.
- [14] R. Ellis and M. Pinsky, The first and second fluid approximations to the linearized Boltzmann equation, *J. Math. Pure. App.*, 54 (1975), 125-156.
- [15] I. M. Gamba, V. Panferov, and C. Villani, Upper Maxwellian bounds for the spatially homogeneous Boltzmann equation, *Arch. Rational Mech. Anal.* 194 (2009), no. 1, 253-282.
- [16] R. Glassey: *The Cauchy Problem in Kinetic Theory*. SIAM, Philadelphia (1996)
- [17] M.P. Gualdani, S. Mischler and C. Mouhot, Factorization of non-symmetric operators and exponential H-theorem, *Mémoire de la Société Mathématique de France*, 153 (2017).
- [18] F. Hérau, D. Tonon and I. Tristani, Regularization estimates and Cauchy theory for inhomogeneous Boltzmann equation for hard potentials without cut-off, *Comm. Math. Phys.*, 377 (2020), 697-771.
- [19] C. Imbert, C. Mouhot, and L. Silvestre, Decay estimates for large velocities in the Boltzmann equation without cutoff, *Journal de l'École polytechnique-Mathématiques* 7 (2020): 143-183.
- [20] Y.-C. Li and M.-Y. Zhong, Green's Function and Pointwise Space-time Behaviors of the Three-Dimensional Relativistic Boltzmann Equation, arXiv:2405.06280.
- [21] Y.-C. Lin, M.-J. Lyu, H.-T. Wang, K.-C. Wu, Space-time behavior of the Boltzmann equation with soft potentials, *J. Differential Equations*, 322 (2022), 180-236.
- [22] T.-P. Liu, *Interactions of nonlinear hyperbolic waves*, Nonlinear analysis (Taipei, 1989), 171-183, World Sci. Publ., Teaneck, NJ, 1991.
- [23] T.-P. Liu and S.-N. Noh, Wave propagation for the compressible Navier-Stokes equations, *J. Hyperbolic Differ. Equ.*, 12 (2015), 385-445.
- [24] T.-P. Liu and W.-K. Wang, The Pointwise Estimates of Diffusion Wave for the Navier-Stokes Systems in Odd Multi-Dimensions, *Comm. Math. Phys.*, 196 (1998), 145-173.
- [25] T.-P. Liu, S.-H. Yu, The Green's function and large time behavior of solutions for one-dimensional Boltzmann equation, *Commun. Pure App. Math.*, 57 (2004), 1543-1608.
- [26] T.-P. Liu and S.-H. Yu, The Green's function of Boltzmann equation, 3D waves, *Bull. Inst. Math. Acad. Sin. (N.S.)*, 1 (2006), 1-78.
- [27] T.-P. Liu and S.-H. Yu, Solving Boltzmann equation, part I: Green's function, *Bull. Inst. Math. Acad. Sin.*, 6 (2011), 115-243.
- [28] S. Ukai and T. Yang, *Mathematical theory of Boltzmann equation*, Lecture Notes Series 8 (2006).

YU-CHU LIN, DEPARTMENT OF MATHEMATICS, NATIONAL CHENG KUNG UNIVERSITY, TAINAN, TAIWAN
Email address: `yuchu@mail.ncku.edu.tw`

HAITAO WANG, SCHOOL OF MATHEMATICAL SCIENCES, INSTITUTE OF NATURAL SCIENCES, MOE-LSC, IMA-SHANGHAI, SHANGHAI JIAO TONG UNIVERSITY, SHANGHAI, CHINA
Email address: `haitallica@sjtu.edu.cn`

KUNG-CHIEN WU, DEPARTMENT OF MATHEMATICS, NATIONAL CHENG KUNG UNIVERSITY, TAINAN, TAIWAN AND NATIONAL CENTER FOR THEORETICAL SCIENCES, NATIONAL TAIWAN UNIVERSITY, TAIPEI, TAIWAN
Email address: `kungchienwu@gmail.com`