- 1---+:-

ASYMPTOTICS OF INTEGRAL POINTS, EQUIVARIANT COMPACTIFICATIONS AND EQUIDISTRIBUTIONS FOR HOMOGENEOUS SPACES

RUNLIN ZHANG

ABSTRACT. Let **U** be a homogeneous variety over \mathbb{Q} of a linear algebraic group. Let \mathcal{U} be an integral model and assume the existence of infinitely many integral points. Then one would like to give an asymptotic count with the help of some height function. In many cases, with the help of measure rigidity of unipotent flows, we reduce this problem to one on equivariant birational geometry. For instance, we show that if **G** and **H** are both connected, semisimple, simply connected and without compact factors, then \mathbf{G}/\mathbf{H} is strongly Hardy-Littlewood with respect to certain height function. We also show that when **H** is "large" in **G** and both **G** and **H** are connected, reductive and without non-trivial \mathbb{Q} -characters, then for every equivariant height, the asymptotic of integral points is the same as the volume asymptotic up to a constant. Three concrete examples with explicit heights are also provided to illustrate our approach.

CONTENTS

1. Introduction	1
Notations	7
2. Arithmetic and Geometry of Equivariant Compactifications	7
3. Equidistribution and Orbital Counting	14
4. Equidistribution and Nondivergence	18
5. Good Height Functions	24
6. Equivariant Compactifications and Focusing	28
7. Example I, Representation of a binary quadratic form by a quaternary form	46
8. Example II, $(n, 1)$ -splitting of \mathbb{Z}^{n+1}	49
9. Example III, the space of triangles	51
Appendix A. Details on the Examples I, II, III	53
Appendix B. Divergence of Translates of Homogeneous Closed Subsets	68
Appendix C. Indices of Various Conditions	78
Ackonwledgement	79
References	79

1. INTRODUCTION

Given a set of polynomials with integral coefficients, it is natural to ask whether solutions exist and when they do, whether the number of solutions is finite or infinite. And when there are infinitely many solutions, can one count them with the help of some height function? These problems are sometimes referred to as Diophantine problems and, depending on the situations, could be extremely difficult.

In this paper we consider the special case when the equations have rich symmetries: their complex solutions allow a transitive action of some linear algebraic group. There are many works on when an integral solution exist in this context (see [BR95, CTX09, EV08] for a small sample of researches). Assuming that integral solutions do exist, the question remained is to count them and we are mainly interested in the case when there are infinitely many solutions. It turns out that the set of integral solutions has a nice structure: it is preserved by the action of an arithmetic subgroup and decomposes into finitely many orbits. Thus, the task of counting integral solutions reduces to counting the orbits of

RUNLIN ZHANG

an arithmetic group. It was observed by [DRS93] that such a question is related to equidistribution of homogeneous measures on a special type of homogeneous spaces: arithmetic quotients of real points of a linear algebraic groups. Ergodic theoretic methods then kick in and solve the problem in many cases [EM93, EMS96]. It is also realized that the height function one use to count the solutions matters.

The purpose of the present paper is to continue the approach of [EMS96, Zha21] via unipotent flows and reduces the counting problem in many cases to geometric questions about the height function. In the remaining part of this introduction, we will present some theorems as well as problems one encounters in this process.

As far as counting integral points on homogeneous varieties is concerned, there are many existing works. Besides those already mentioned, some relevant works are [Sha00, HT03, Mau07, GOS09, BO12, GN12, CLT12, MG14, OS14, KK18, SZ19, Zha19], touching on different aspects of the question.

1.1. Hardy-Littlewood varieties. The notion of strongly Hardy-Littlewood varieties are introduced in [BR95] (compare [Bir62]) for affine varieties. Such varieties are defined by certain nice local-to-global properties. To simplify some discussions, we specialize to the case when **U** is isomorphic to \mathbf{G}/\mathbf{H} for some semisimple linear algebraic group **G** together with a semisimple Q-subgroup **H**. In particular, we assume that $\mathbf{U}(\mathbb{Q}) \neq \emptyset$.

1.1.1. Integral points. Let **U** be an affine homogeneous variety over \mathbb{Q} . Let $\mathbf{U}(\mathbb{A}_{\mathbb{Q}})$ be the topological space of adelic points. Given an open compact subset $B_f \subset \mathbf{U}(\mathbb{A}_{\mathbb{Q},f})$ of finite adeles and a connected component B_{∞} of $\mathbf{U}(\mathbb{R})$, one define the set of \mathscr{B} -integral points $\mathbf{U}_{\mathscr{B}}(\mathbb{Z})$ to be the intersection of $\mathscr{B} := B_{\infty} \times B_f$ with the diagonal embedding to $\mathbf{U}(\mathbb{Q})$ in the set of adeles.

1.1.2. Tamagawa measures. Let ω be a non-vanishing top-degree differential form on **U** over \mathbb{Q} . It is actually **G**-invariant and induces measures ω_p on each $\mathbf{U}(\mathbb{Q}_p)$ ($\mathbb{Q}_{\infty} := \mathbb{R}$) and, in fact, the products over finite p's induce a measure ω_f on $\mathbf{U}(\mathbb{A}_{\mathbb{Q},f})$. By the product formula and the triviality of global regular functions on **U**, the measure $\omega_f \otimes \omega_{\infty}$ is independent of the choice of ω .

1.1.3. The definition. Let $l: \mathbf{U}(\mathbb{R}) \to \mathbb{R}_{>0}$ be a proper function and

$$B_R := \left\{ x \in \mathbf{U}(\mathbb{R}), \ l(x) \le R \right\}.$$

We say that **U** is a strongly Hardy-Littlewood variety (with respect to l) if for any choice of \mathscr{B} as above,

$$\lim_{R \to \infty} \frac{\# \mathbf{U}_{\mathscr{B}}(\mathbb{Z}) \cap B_R}{\omega_f(B_f) \cdot \omega_\infty(B_R)} = 1$$

In [BR95], l is induced from an embedding of **U** into some affine space \mathbf{A}^n .

1.1.4. Theorems. Now, assume further that **G** and **H** are both connected and simply connected. Also assume that the real points of every non-trivial \mathbb{Q} -factor of **G** is non-compact. Note that we have assumed $\mathbf{U}(\mathbb{Q}) \neq \emptyset$. The following is [BR95, Theorem 0.3]:

Theorem 1.1. Assume that for any congruence subgroup $\Gamma \subset \mathbf{G}(\mathbb{Q})$ and any rational point $x \in \mathbf{U}(\mathbb{Q})$, one has

$$\lim_{R \to \infty} \frac{\#\Gamma.x \cap B_R}{\omega_{\infty}(B_R) \cap \mathbf{G}(\mathbb{R}).x} = \frac{\operatorname{Vol}(\mathbf{H}_x(\mathbb{R})/\mathbf{H}_x(\mathbb{R}) \cap \Gamma)}{\operatorname{Vol}(\mathbf{G}(\mathbb{R})/\Gamma)}$$
(1)

where \mathbf{H}_x denotes the stabilizer of x in **G** and volumes are induced from compatible choices of Haar measures on $\mathbf{G}(\mathbb{R}), \mathbf{H}_x(\mathbb{R})$ and $\mathbf{G}(\mathbb{R})/\mathbf{H}_x(\mathbb{R})$. Then **U** is strongly Hardy–Littlewood with respect to l.

We remark that if $\Gamma' \subset \Gamma$ is a finite-index subgroup, then the truth of Equa.(1) for Γ' implies Equa.(1) for Γ . Equa.(1) has been verified in many cases. For instance, when **H** is maximal in a semisimple **G** without compact Q-factors, all "algebraic height functions" l would work. What is proved in this paper (See Section 6.8) is that

Theorem 1.2. Assumptions same as above. There exists l such that Equa.(1) holds. In particular, U is always strongly Hardy-Littlewood for some l.

Already when $\mathbf{G} = \mathbf{SL}_n$ and $\mathbf{H} = \mathbf{SL}_{n_1} \times ... \times \mathbf{SL}_{n_k}$ with $\sum n_i \leq n$, this seems unknown before. Generalizations can be obtained by combining Theorem 6.43 with [WX16]. 1.2. Manin conjecture for a pair. In the previous section, there is no emphasis on which height function l one should use.

In the context of integral points in projective varieties, roughly speaking, Manin conjecture makes prediction on the asymptotics of integral points with respect to heights associated to the anti-canonical bundle. In the case of a pair, one should replace the anti-canonical line bundle by the log anti-canonical line bundle.

1.2.1. Log anti-canonical line bundles. Let **X** be a **G**-equivariant compactification of $\mathbf{U} \cong \mathbf{G}/\mathbf{H}$ and $\mathbf{D} := \mathbf{X} \setminus \mathbf{U}$ be the boundary. We assume that **X** is smooth and **D** is a simple normal crossing divisor. Let $\mathcal{K}_{\mathbf{X}}$ be the canonical line bundle. Then $(\mathcal{K}_{\mathbf{X}} \otimes \mathcal{O}_{\mathbf{X}}(\mathbf{D}))^{\vee}$ is called the *log anti-canonical* line bundle.

1.2.2. *Heights.* Without loss of generality, we only consider the height at infinity here (see Remark 2.4). Let $\mathbf{D} = \sum_{\alpha \in \mathscr{A}} \mathbf{D}_{\alpha}$ be the decomposition of \mathbf{D} into absolutely irreducible components. Let $\mathbf{L} := \sum \lambda_{\alpha} \mathbf{D}_{\alpha}$ with $\lambda_{\alpha} > 0$ for all $\alpha \in \mathscr{A}$. View the line bundle $\mathcal{O}_{\mathbf{X}}(\mathbf{L})$ over \mathbb{R} in the category of smooth manifolds and equip it with a smooth metric $\|\cdot\|$. Let $\mathbf{1}_{\mathbf{L}}$ be the canonical section of $\mathcal{O}_{\mathbf{X}}(\mathbf{L})$ and define for $x \in \mathbf{U}(\mathbb{R})$,

$$\operatorname{Ht}(x) := \|\mathbf{1}_{\mathbf{L}}(x)\|^{-1}.$$

Points of height bounded by a fixed number is a compact subset of $\mathbf{U}(\mathbb{R})$.

1.2.3. Volume. Let $\mathbf{D}' := \sum d_{\alpha} \mathbf{D}_{\alpha}$ be a divisor supported on \mathbf{D} . We do not require \mathbf{D}' to be effective or $d_{\alpha} \geq 0$, so $\mathbf{1}_{\mathbf{D}'}$ is only a rational section that is regular on \mathbf{U} . Let $\mathcal{K}_{\mathbf{X}}(\mathbf{D}') := \mathcal{K}_{\mathbf{X}} \otimes \mathcal{O}_{\mathbf{X}}(\mathbf{D}')$. Assume that $\mathcal{K}_{\mathbf{X}}(\mathbf{D}')$ (again, viewed in the category of smooth manifolds) is equipped with a smooth metric $\|\cdot\|$, then a measure, denoted as Vol, on $\mathbf{U}(\mathbb{R})$ is locally defined by

$$\operatorname{Vol} = \frac{|\omega|}{\|\omega \otimes \mathbf{1}_{\mathbf{D}'}\|}, \text{ for any non-vanishing local volume form } \omega.$$

1.2.4. Haar measure. In the homogeneous setting, a natural choice of measure is the $\mathbf{G}(\mathbb{R})$ -invariant Haar measure. For this we assume the existence of a non-zero **G**-invariant top-degree differential form on **U**. Let ω_0 be such a form and let $\mathbf{m}_{\mathbf{G}/\mathbf{H}}$ be the measure on $\mathbf{U}(\mathbb{R})$ induced form it, then $\mathbf{m}_{\mathbf{G}/\mathbf{H}}$ is $\mathbf{G}(\mathbb{R})$ invariant. Let $\mathbf{D}' := -\operatorname{div}(\omega_0)$, then $\mathbf{1}_{\mathbf{D}'}$ is identified with ω_0^{\vee} , the unique rational section of $\mathcal{O}_{\mathbf{X}}(\mathbf{D}') \cong \mathcal{K}_{\mathbf{X}}^{\vee}$ (the dual of $\mathcal{K}_{\mathbf{X}}$) such that the natural pairing between ω_0 and ω_0^{\vee} is the constant function 1. Now define a smooth metric on $\mathcal{K}_{\mathbf{X}}(\mathbf{D}')$ by imposing $\|(\omega_0 \otimes \omega_0^{\vee})(x)\| = 1$ for every x. Then the Vol from the last subsection coincides with $\mathbf{m}_{\mathbf{G}/\mathbf{H}}$ here.

1.2.5. The prediction. Fix an integral model \mathcal{X} of \mathbf{X} and \mathcal{D} of \mathbf{D} . Then $\mathcal{U} := \mathcal{X} \setminus \mathcal{D}$ is an integral model of \mathbf{U} . Or more generally, one can take $\mathcal{U}(\mathbb{Z}) := \mathbf{U}_{\mathscr{B}}(\mathbb{Z})$ as above.

Let $(D_{\alpha'})_{\alpha' \in \mathscr{A}'}$ be consisting of connected components of $\mathbf{D}_{\alpha}(\mathbb{R})$ as α varies in \mathscr{A} . The analytic Clemens complex $\mathscr{C}_{\mathbb{R}}^{\mathrm{an}}$, elements of which consist of connected components of non-empty intersection of $D_{\alpha'}$'s, encodes the intersection patterns among boundary divisors.

Depending on the integral model, let $\mathscr{C}_{\mathbb{R},\mathbb{Z}}^{an}$ be the modified analytic Clemens complex consisting of faces $F \in \mathscr{C}_{\mathbb{R}}^{an}$ with

$$\left(\bigcap_{\alpha\in F} D_{\alpha}\right) \bigcap \left(\bigcup_{x\in\mathcal{U}(\mathbb{Z})} \overline{\mathbf{G}(\mathbb{R})^{\circ}.x}\right) \neq \emptyset.$$

That is to say, we only take closures of connected components of $\mathbf{U}(\mathbb{R})$ where there exist at least one integral point. Let $E_{\mathbb{Q}}(\mathbf{U})$ be the abelian group of invertible functions on \mathbf{U} modulo constant functions \mathbb{Q}^{\times} and

 $b := \max\left\{ \operatorname{rk}\operatorname{Pic}_{\mathbb{Q}}(\mathbf{U}) - \operatorname{rk} E_{\mathbb{Q}}(\mathbf{U}) + \#F \mid F \text{ is a maximal face of } \mathscr{C}_{\mathbb{R},\mathbb{Z}}^{\mathrm{an}} \right\}.$

When **G** is semisimple and simply connected and $H^1(\mathbb{Q}, \mathbf{H})$ is trivial, the modified analytic Clemens complex coincide with the usual one. But in general, they could be different (see [Wil22]).

Here is a natural question inspired by Manin conjecture.

Question 1.3. Let Ht be a height function associated with a log anti-canonical divisor. Assume $\mathcal{U}(\mathbb{Z}) \neq \emptyset$. Does there exist a positive number c > 0 such that

$$\# \{ x \in \mathcal{U}(\mathbb{Z}) \mid \operatorname{Ht}(x) \le R \} \sim cR(\log R)^{b-1}$$

as $R \to +\infty$?

Here is what we can do on the positive side (combine Theorem 6.27, Lemma 5.2 and Theorem 2.13).

Theorem 1.4. Assume the following:

- G and H are connected and reductive, and both have no non-trivial Q-characters;
- the identity component of the centralizer of **H** in **G** is contained in **H**;
- the projection of **H** to the compact Q-factor of **G** is surjective.

Then the log anti-canonical divisor is big and Question 1.3 admits a positive answer.

Note that under the assumptions made in the theorem, $\operatorname{rk}\operatorname{Pic}_{\mathbb{Q}}(\mathbf{U}) = \operatorname{rk} E_{\mathbb{Q}}(\mathbf{U}) = 0$.

1.3. Weighted counting. In this subsection, we assume that **G** is semisimple without compact \mathbb{Q} -simple factors and **H** is reductive and connected. However, as opposed to previous discussions, we are mainly interested in the case when $\mathbf{H}(\mathbb{R})\Gamma/\Gamma$ has infinite volume. In this case, the number of integral points of bounded height is no longer expected to be asymptotic to the volume. To remedy the situation, we consider weighted versions of the counting problem.

We also assume $\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ}$ is contained in \mathbf{H} throughout this subsection. First we discuss an analogue of Theorem 1.2. Let ω_{∞} be an invariant measure on $\mathbf{U}(\mathbb{R})$ induced from some invariant gauge form.

Theorem 1.5. There exists a height function Ht as constructed in Section 1.2 such that for every $x \in U(\mathbb{Q})$ and every arithmetic lattice Γ of \mathbf{G} , one has

$$\sum_{\in \Gamma.x \cap B_{R,x}} \mathbf{w}_y \sim \omega_\infty(B_{R,x})$$

where $B_{R,x} := \{ y \in \mathbf{G}(\mathbb{R})^{\circ} . x, \operatorname{Ht}(y) \leq R \}.$

See Theorem 6.26. For a more general height, our result would be much weaker. Let (\mathbf{X}, \mathbf{D}) be a smooth **G**-pair over \mathbb{Q} such that **U** is **G**-equivariantly isomorphic to $\mathbf{X} \setminus \mathbf{D}$. Let Ht be associated with some divisor whose support is equal to **D**. Let $\mathbf{G} := \mathbf{G}(\mathbb{R})^{\circ}$, $\mathbf{H}_x := \mathbf{H}_x(\mathbb{R}) \cap \mathbf{G}$ and assume that Γ is contained in **G**.

Let

 $b' := \max\left\{ \#F \mid F \text{ is a maximal face of } \mathscr{C}_{\mathbb{R},\mathbb{Z}}^{\mathrm{an}} \right\}.$

The following will be proved in Theorem 6.27.

Theorem 1.6. Let ψ be a non-negative compactly supported function whose support is large enough. For $x \in \mathbf{U}(\mathbb{Q})$, define

$$\mathbf{w}_x := \sum_{h \in \mathbf{H}_x / \mathbf{H}_x(\mathbb{R})^{\circ}} \langle \psi, h_* \mathbf{m}_{[\mathbf{H}_x(\mathbb{R})^{\circ}]} \rangle^{-1}.$$

Then there exists c > 0 such that

$$\sum_{x \in \mathcal{U}(\mathbb{Z}), \operatorname{Ht}(x) \leq R} \mathbf{w}_x \sim c \cdot R(\log R)^{b'-1}$$

1.4. **Examples.** We provide three examples of different flavours. The first two examples are spherical varieties but the third one is not.

1.4.1. Example I. Let (Q_1, \mathbb{Q}^2) and (Q_2, \mathbb{Q}^4) be two quadratic forms represented by symmetric matrices M_{Q_i} for i = 1, 2. Let $\mathbf{Mat}_{4,2}(\mathbb{Z})$ be the set of 4-by-2 matrices with integral coefficients and

$$\mathcal{U}(\mathbb{Z}) := \left\{ M \in \mathbf{Mat}_{4,2}(\mathbb{Z}) \mid M^{\mathrm{tr}} M_{Q_2} M = M_{Q_1} \right\}.$$

satisfying:

- Both (Q_1, \mathbb{R}^2) and (Q_2, \mathbb{R}^4) are split. And (Q_1, \mathbb{Q}^2) is Q-anisotropic;
- $\mathcal{U}(\mathbb{Z})$ is non-empty.

Hence we can find (Q'_1, \mathbb{Q}^2) such that (Q_2, \mathbb{Q}^4) is isomorphic to $(Q_1, \mathbb{Q}^2) \oplus (Q'_1, \mathbb{Q}^2)$. We further assume that

• the orthogonal group of (Q_1, \mathbb{Q}^2) is not isomorphic to the orthogonal group of (Q'_1, \mathbb{Q}^2) as linear algebraic groups over \mathbb{Q} .

Let $\|\cdot\|$ denote the Euclidean norm of a matrix.

Theorem 1.7. As R tends to infinity, there exists some constant c > 0 such that

$$\# \{ M \in \mathcal{U}(\mathbb{Z}), \|M\| \le R \} \sim cR^2 \log R.$$

See Theorem 7.1 for more details.

1.4.2. Example II. Let $\Lambda(n, 1)$ denote the set of splittings of \mathbb{Z}^{n+1} as a direct sum of two subgroups, one of which has rank n. Each element of $\Lambda(n, 1)$ can be associated with (v, M) where $v \in \mathbb{Z}^{n+1}$ is an integral vector and M is an element in $\wedge^n \mathbb{Z}^{n+1}$. Moreover, v, M are unique up to sign. Below $\|\cdot\|$ denotes the natural Euclidean norm.

Theorem 1.8. For two positive integers λ_1, λ_2 , we have

$$\#\{[(v,M)] \in \Lambda(n,1), \|v\|^{\lambda_1} \|M\|^{\lambda_2} \le R\} \sim \begin{cases} c_{\lambda_1,\lambda_2} \cdot R^{\frac{n+1}{\min\{\lambda_1,\lambda_2\}}} \log(R), & \lambda_1 = \lambda_2 \\ c_{\lambda_1,\lambda_2} \cdot R^{\frac{n+1}{\min\{\lambda_1,\lambda_2\}}}, & \lambda_1 \neq \lambda_2 \end{cases}$$

 $m \perp 1$

for some $c_{\lambda_1,\lambda_2} > 0$.

See Theorem 8.1 below for details.

1.4.3. Example III. The third example concerns with the space of triangles. Let $\mathcal{M}_3(\mathbb{Z})$ denote the set of three ordered linearly independent lines in \mathbb{Q}^3 . For an element $(\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3) \in \mathcal{M}_3(\mathbb{Z})$, we take \mathbf{v}_i to be a non-zero vector on \mathbf{l}_i . Here are two ways to measure the "height" of such an element:

$$\begin{aligned} &\operatorname{Ht}_{1}(\mathbf{l}_{1}, \mathbf{l}_{2}, \mathbf{l}_{3}) := \frac{\prod_{i} \|\mathbf{v}_{i}\|^{2}}{\|\mathbf{v}_{1} \wedge \mathbf{v}_{2}\| \|\mathbf{v}_{1} \wedge \mathbf{v}_{3}\| \|\mathbf{v}_{2} \wedge \mathbf{v}_{3}\|} \\ &\operatorname{Ht}_{2}(\mathbf{l}_{1}, \mathbf{l}_{2}, \mathbf{l}_{3}) := \frac{\|\mathbf{v}_{1} \wedge \mathbf{v}_{2}\|^{2} \|\mathbf{v}_{1} \wedge \mathbf{v}_{3}\|^{2} \|\mathbf{v}_{2} \wedge \mathbf{v}_{3}\|^{2}}{(\prod_{i} \|\mathbf{v}_{i}\|) \cdot \|\mathbf{v}_{1} \wedge \mathbf{v}_{2} \wedge \mathbf{v}_{3}\|^{3}} \end{aligned}$$

Theorem 1.9. Let $\kappa_1, \kappa_2 > 0$ and $\operatorname{Ht}(\mathbf{x}) := \operatorname{Ht}_1(\mathbf{x})^{\kappa_1} \operatorname{Ht}_2(\mathbf{x})^{\kappa_2}$ for $\mathbf{x} \in \mathbf{M}_3(\mathbb{R})$. Then for some constant $c_{\kappa_1,\kappa_2} > 0$,

$$\sum_{\{\mathbf{x}\in\mathcal{M}_3(\mathbb{Z})\mid \mathrm{Ht}(x)\leq R\}} \mathbf{w}_x \sim \begin{cases} c_{\kappa_1,\kappa_2} \cdot R^{\frac{8}{3}\kappa_1^{-1}} \cdot \log(R) & \kappa_1 = \kappa_2 \\ c_{\kappa_1,\kappa_2} \cdot R^{\frac{8}{3}\max\{\kappa_1^{-1},\kappa_2^{-1}\}} & \kappa_1 \neq \kappa_2. \end{cases}$$

See Theorem 9.1 below for more details.

1.5. Outline of the proof and future directions.

1.5.1. Outline of the proof. Let me explain what goes into the proof.

First, it is known that if \mathbf{U} admits transitive action of a linear algebraic group, then its set of integral points admits an action of some arithmetic subgroup and furthermore, there are only finitely many orbits (see Theorem 2.3).

So now one should count points on a fixed orbit, say $\Gamma.x$. It is discovered in [DRS93] that a folding/unfolding argument transforms this orbital counting problem into an equidistribution problem. We elaborate on this in Section 3.

This equidistribution problem is about limiting behaviour of certain average of translates of a fixed homogeneous measure, which looks like:

$$\frac{1}{\operatorname{Vol}(B_{R,x})}\int_{[g]\in B_{R,x}}g_*\mathrm{m}_{\mathbf{H}_x(\mathbb{R})\Gamma/\Gamma}\operatorname{Vol}([g]).$$

Here \mathbf{H}_x is the stabilizer of x in \mathbf{G} , Vol refers to certain natural measure on $\mathbf{G}(\mathbb{R})/\mathbf{H}_x(\mathbb{R})$ and $B_{R,x}$ are those co-sets $g\mathbf{H}_x(\mathbb{R})$ satisfying $\mathrm{Ht}(g.x) \leq R$.

The study of the limiting behaviors of the integrand in the above expression is the subject of [EMS96] and [Zha21]. Perhaps the most crucial input behind these works is the classification of unipotent-invariant ergodic probability measures due to Ratner [Ra91].

Theorems from [EMS96, Zha21] are most useful when **H** (and hence \mathbf{H}_x) is large in **G**. When this is not the case, one may do the centralizer trick by considering, for instance, $\mathbf{G} \times \mathbf{Z}/\mathbf{H} \cdot \Delta(\mathbf{Z})$ where **Z** is contained in the centralizer of **H** in **G**. This trick works better when **H** is connected and reductive.

RUNLIN ZHANG

Anyway, assume **H** is large in **G** in some sense, then one can try to analyze the possible limiting measures of $(g_*m_{\mathbf{H}_x(\mathbb{R})\Gamma/\Gamma})$ as g varies. Preferably, the limiting behaviours are controlled by finitely many vectors in finitely many linear representations. Or as a weaker goal, explain the failure of tending to the full G-invariant Haar measure by finitely many conditions. Then one can try interpreting these conditions in the language of algebraic varieties and their closed subvarieties, and finally, by resolution of a pair, into language of smooth projective varieties and their simple normal crossing divisors. The reader might jump ahead to Section 5 for more rigorous discussions.

Once this is done, the equidistribution theorem of [CLT12] becomes very useful. When applying [CLT12], it is crucial to have some knowledge about the log anti-canonical divisor. Under some (necessary) conditions, we show that its support is the full boundary (see Theorem 2.13 and 2.14).

1.5.2. What remains to be done? We list some questions that are related to the discussion of the current paper. Of course, one natural task is to verify the log Manin conjecture recently proposed by [San23]. Here we are a bit more specific.

Let **U** be a homogeneous variety over \mathbb{Q} isomorphic to \mathbf{G}/\mathbf{H} . Assume $\mathbf{U}(\mathbb{Q}) \neq \emptyset$. For $x \in \mathbf{U}(\mathbb{Q})$, we let \mathbf{H}_x be the stabilizer of x in \mathbf{G} . Let $\mathbf{G} := \mathbf{G}(\mathbb{R})^\circ$ and $\mathbf{H}_x := \mathbf{H}_x(\mathbb{R}) \cap \mathbf{G}$. Let Γ be an arithmetic subgroup contained in \mathbf{G} . We let $\mathbf{m}_{[\mathbf{H}_x]}$, $\mathbf{m}_{[\mathbf{G}]}$ and $\mathbf{m}_{\mathbf{G}/\mathbf{H}_x}$ to denote invariant measures (so we need to assume that they exists) on $\mathbf{H}_x\Gamma/\Gamma$, \mathbf{G}/Γ and \mathbf{G}/\mathbf{H}_x respectively. We require these measures to be compatible. Namely, if $\mathbf{m}_{\mathbf{H}_x}$ and $\mathbf{m}_{\mathbf{G}}$ are the corresponding invariant measures on \mathbf{H}_x and \mathbf{G} respectively that induce $\mathbf{m}_{[\mathbf{H}_x]}$ and $\mathbf{m}_{[\mathbf{G}]}$, then for every compactly supported function f on \mathbf{G} , one has

$$\int \left(\int f(gh) \operatorname{m}_{\operatorname{H}_{x}}(h)\right) \operatorname{m}_{\operatorname{G/H}_{x}}([g]) = \int f(x) \operatorname{m}_{\operatorname{G}}(x)$$

1.5.3. Finite-volume case. Assume that **G** and **H** are both connected. Assume all the non-trivial \mathbb{Q} -factors of **G** are non-compact over \mathbb{R} . The case when **H** is not connected is less understood. Assume further that **G** and **H** has no non-trivial \mathbb{Q} -characters. Equivalently, $\mathbf{G}(\mathbb{R})/\Gamma$ and $\mathbf{H}(\mathbb{R})\Gamma/\Gamma$ have finite volume.

We expect that there exists a class of good heights (or, good boundary divisors) such that

$$\# \{ y \in \Gamma.x \mid \operatorname{Ht}(y) \le R \} \sim \frac{|\mathbf{m}_{[\mathbf{H}_x]}|}{|\mathbf{m}_{[\mathbf{G}]}|} \cdot \mathbf{m}_{\mathbf{G}/\mathbf{H}_x} \left(\{ y \in \mathbf{G}.x \mid \operatorname{Ht}(y) \le R \} \right)$$
(2)

for every $x \in \mathbf{U}(\mathbb{Q})$ and every arithmetic lattice Γ (of course, it is sufficient to verify this for some finite-index subgroup of Γ).

Question 1.10. Assume that **H** is large in **G** and all the invariant measures exist. Let (\mathbf{X}, \mathbf{D}) be a smooth G-pair over \mathbb{Q} with $\mathbf{X} \setminus \mathbf{D}$ equivariantly isomorphic to **U**. Is Equa.(2) true for the log anti-canonical height on (\mathbf{X}, \mathbf{D}) ?

Being large does not have a precise definition. For instance, one may say that \mathbf{H} is large in \mathbf{G} if one of the following is true:

- 1. **G** is semisimple and \mathbf{G}/\mathbf{H} is a spherical variety;
- 2. **G** and **H** are reductive and the identity component of the centralizer of **H** in **G** is contained in **H**.

1.5.4. Infinite-volume case. Assume that **G** is connected, semisimple and all \mathbb{Q} -simple factors are noncompact. Assume **H** is connected, reductive and has non-trivial \mathbb{Q} -characters. In this case, we no longer expect the asymptotic of integers is the same as the volume asymptotic on G/H. However, we do have a replacement of the volume asymptotic.

Recall $\mathscr{P}_{\mathbf{H}}^{\max}$ is the collection of maximal parabolic Q-subgroups containing **H**. For each $\mathbf{P} \in \mathscr{P}_{\mathbf{H}}^{\max}$, we fix an integral vector¹ $\mathbf{v}_{\mathbf{P}} \in \wedge^{\dim \mathbf{P}} \mathfrak{g}$ that lifts the Lie algebra of **P**. Let Ht be a height function on $\mathbf{U}(\mathbb{R})$ and let $x \in \mathbf{U}(\mathbb{Q})$. Let \mathbf{H}'_x be the largest connected Q-subgroup of \mathbf{H}_x where all the Q-characters of **H** vanish. For $\eta, R > 0$, define the following region

$$B_{\eta,R,x} := \left\{ [g] \in \mathcal{G}/\mathcal{H}'_x \mid \operatorname{Ht}(g.x) \le R, \ \|\operatorname{Ad}(g)\mathbf{v}_{\mathbf{P}}\| \ge \eta, \ \forall \ \mathbf{P} \in \mathscr{P}_{\mathbf{H}_x}^{\max} \right\}.$$

We expect that in good situations, the asymptotic of $m_{G/H'_{-}}(B_{\eta,R,x})$ as $R \to +\infty$ is independent of η .

¹Implicitly, an integral structure $\mathfrak{g}_{\mathbb{Z}}$ of \mathfrak{g} is fixed.

$$\# \{ y \in \Gamma.x \mid \operatorname{Ht}(y) \le R \} \sim \frac{1}{\left| \operatorname{m}_{[G]} \right|} \cdot \operatorname{m}_{G/\operatorname{H}'_{x}}(B_{\eta,R,x}) \quad as \ R \to +\infty$$

hold for every $x \in \mathbf{U}(\mathbb{Q})$ and every arithmetic subgroup Γ ?

Question 1.12. Given $\eta > 0$, what is the asymptotic of $m_{G/H'_{\pi}}(B_{\eta,R,x})$?

In the case of Example III (Section 1.4.3) discussed above, it seems that $m_{G/H'_x}(B_{\eta,R,x})$ can be expressed in terms of the norms of sections of certain line bundles.

NOTATIONS

For an algebraic group **G** over a field k, let the gothic letter \mathfrak{g} denote its Lie algebra over k. When the notation looks cumbersome, we also use Lie(**G**). Let $\mathfrak{X}^*(\mathbf{G})$ (resp. $\mathfrak{X}^*_k(\mathbf{G})$) be the group of characters (resp. k-characters) of **G**. Similarly \mathfrak{X}^k_* is the group of cocharacters over k. Following [BS73], we let

$$^{\circ}\mathbf{G}:=\bigcap_{\alpha\in\mathfrak{X}_{k}^{*}(\mathbf{G})}\ker\alpha^{2}.$$

Let $\mathbf{S}_{\mathbf{G}}$ be the quotient \mathbb{Q} -split torus $\mathbf{G}/{}^{\circ}\mathbf{G}$ and p^{spl} be the natural quotient morphism $\mathbf{G} \to \mathbf{S}_{\mathbf{G}}$.

Assume G to be a connected linear algebraic group and let $\mathbf{R}_u(G)$ be its unipotent radical, then G is equal to

$$\mathbf{G}^{\mathrm{red}} \ltimes \mathbf{R}_{\mathbf{u}}(\mathbf{G}) = (\mathbf{G}^{\mathrm{ss}} \cdot \mathbf{Z}(\mathbf{G})) \ltimes \mathbf{R}_{\mathbf{u}}(\mathbf{G}) = \left(\mathbf{G}^{\mathrm{cpt}} \cdot \mathbf{G}^{\mathrm{nc}} \cdot \mathbf{Z}(\mathbf{G})^{\mathrm{an}} \cdot \mathbf{Z}(\mathbf{G})^{\mathrm{spl}}\right) \ltimes \mathbf{R}_{\mathbf{u}}(\mathbf{G})$$

where $\mathbf{G}^{\mathrm{red}}$ is a connected reductive \mathbb{Q} -subgroup (called a Levi subgroup) lifting $\mathbf{G}/\mathbf{R}_{\mathbf{u}}(\mathbf{G})$, $\mathbf{G}^{\mathrm{ss}} = [\mathbf{G}^{\mathrm{red}}, \mathbf{G}^{\mathrm{red}}]$ is a connected semisimple \mathbb{Q} -subgroup, $\mathbf{Z}(\mathbf{G})$ is the identity component of the center of $\mathbf{G}^{\mathrm{red}}$ (called the central torus), $\mathbf{G}^{\mathrm{cpt}}$ is the product of \mathbb{Q} -simple factors of \mathbf{G}^{ss} that are \mathbb{R} -anisotropic, \mathbf{G}^{nc} is the product of \mathbb{Q} -simple factors of \mathbf{G}^{ss} that are \mathbb{R} -anisotropic. A \mathbb{Q} -torus \mathbf{T} can be written uniquely as an almost direct product of $\mathbf{T}^{\mathrm{an}} \cdot \mathbf{T}^{\mathrm{spl}}$, a \mathbb{Q} -anisotropic subtorus and a \mathbb{Q} -split subtorus. This explains the notation $\mathbf{Z}(\mathbf{G})^{\mathrm{an}}$ and $\mathbf{Z}(\mathbf{G})^{\mathrm{spl}}$ above.

Let $\mathbb{A}_{\mathbb{Q}}$ denote the topological adelic ring of the field of rational numbers \mathbb{Q} . $\operatorname{Gal}_{\mathbb{Q}}$ denotes the Galois group of the extension $\overline{\mathbb{Q}}/\mathbb{Q}$.

We say that (\mathbf{X}, \mathbf{D}) is a **G**-pair over \mathbb{Q} if **X** is a projective variety over \mathbb{Q} , **D** is a closed subvariety over \mathbb{Q} and **X** is equipped with the action of **G** such that **D** is preserved under this action. A pair (\mathbf{X}, \mathbf{D}) is said to be a smooth pair over \mathbb{Q} if **X** is a smooth projective variety over \mathbb{Q} and **D** is a simple/strict normal crossing divisor over \mathbb{Q} .

For locally finite measures (ν_n) and ν on a locally compact second countable Hausdorff space X, following [SZ19], we say that $[\nu_n]$ converges to $[\nu]$ and write $\lim_{n\to\infty} [\nu_n] = [\nu]$ if there exists $(a_n) \subset \mathbb{R}_{>0}$ such that $\lim_{n\to\infty} a_n\nu_n = \nu$ under the weak-* topology.

2. ARITHMETIC AND GEOMETRY OF EQUIVARIANT COMPACTIFICATIONS

For this section we further assume the following notations:

- X is a smooth projective variety over Q of dimension d;
- D ⊂ X is a reduced simple normal crossing divisor over Q and U is the complement of D in X;
- (D_α)_{α∈𝒜} are the irreducible components of D over the algebraic closure Q. This set and its index set 𝒜 are equipped with an action of Gal_Q;
- **G** is a linear algebraic group over \mathbb{Q} and **H** is a \mathbb{Q} -subgroup of **G**.

2.1. Integral points, adelic heights and orbits of arithmetic groups.

RUNLIN ZHANG

2.1.1. Integral points. Let Val be the set of valuations on \mathbb{Q} up to equivalence, identified with the set of prime numbers and the Archimedean place $\{\infty\}$. By convention $\mathbb{Q}_{\infty} := \mathbb{R}$. Let Val_f denote only the set of finite primes. For a finite set $S \subset \operatorname{Val}_f$, let \mathbb{Z}_S denote the set of rational numbers that are integral outside S.

Consider the space of adelic points given by certain restricted product

$$\mathbf{U}(\mathbb{A}_{\mathbb{Q}}) := \prod_{p \in \mathrm{Val}}' \mathbf{U}(\mathbb{Q}_p) imes \mathbf{U}(\mathbb{R})$$

equipped with adelic topology (see [Con12] for details). Let \mathscr{K}_f be a non-empty open compact subset in $\prod_{p \in \operatorname{Val}_f} \mathbf{U}(\mathbb{Q}_p)$. More concretely, there exists a finite set $S \subset \operatorname{Val}_f$ and a smooth model $(\mathscr{X}, \mathcal{D})$ of (\mathbf{X}, \mathbf{D}) over Spec \mathbb{Z}_S such that $\mathscr{K}_f = K_S \times \prod_{p \notin S} K_p$ for some open compact subset $K_S \subset \prod_{p \in S} \mathbf{U}(\mathbb{Q}_p)$ and

$$K_p = (\mathcal{X} \setminus \mathcal{D})(\mathbb{Z}_p), \quad \forall p \notin S.$$
 (3)

For $\mathscr{K} = \mathscr{K}_f \times K_\infty$ where K_∞ is a union of some connected components of $\mathbf{U}(\mathbb{R})$, define

$$\mathbf{U}_{\mathscr{K}}(\mathbb{Z}) := \mathbf{X}(\mathbb{Q}) \bigcap \mathscr{K}_f \times K_{\infty}$$

This adelic point of view contains the classical case of taking the integral points of some integral model as a special case.

2.1.2. Adelic heights. As a reference, see [CLT10].

Let (L, s) be a line bundle over **X** together with a global section $s \in \Gamma(\mathbf{X}, L)$. An *adelic metric* $(\|\cdot\|_p)_{p \in Val}$ is the following:

- 1. for each $p \in \text{Val}$, the $\|\cdot\|_p$ is a smooth metric on the analytic line bundle $\mathcal{L}(\mathbb{Q}_p)$ over the analytic manifold $\mathbf{X}(\mathbb{Q}_p)$;
- 2. there exist a finite set $S \subset \operatorname{Val}_f$ and a smooth model $\mathcal{L} \to \mathcal{X}$ over $\operatorname{Spec} \mathbb{Z}_S$ such that $\|\cdot\|_p$ is defined by

$$\|s(x)\|_{p} \le 1 \iff s(x) \in \tilde{x}^{*}\mathcal{L}$$

$$\tag{4}$$

where $x \in \mathbf{X}(\mathbb{Q}_p)$ and $\tilde{x} \in \mathcal{X}(\mathbb{Z}_p)$ is a lift of x.

By saying that a metric $\|\cdot\|_p$ is *smooth*, we mean that for every nonzero local section s, the map $x \mapsto \|s(x)\|_p$ is a smooth function on the locus where s does not vanish. For simplicity, L together with this adelic metric is called an *adelic line bundle*.

If $L = \mathcal{O}_{\mathbf{X}}(\mathbf{L})$ for some effective divisor \mathbf{L} and $\mathbf{s}_{\mathbf{L}}$ is the canonical section, then for some finite set S,

$$\|\mathbf{s}_{\mathbf{L}}(x)\|_{p} = 1, \quad \forall p \notin S, \ x \in K_{p}.$$
(5)

Indeed, since $\mathbf{s}_{\mathbf{L}}$ is a local generator of L over \mathbb{Q} , it is also a local generator of \mathcal{L} over \mathbf{Z}_S for some finite S. Take S larger such that Equa.(4) also holds. Then Equa.(5) follows from Equa.(4) above.

Assume that the support of L is contained in D. We define an *adelic height function* on $U(\mathbb{A}_{\mathbb{O}})$ by

$$\operatorname{Ht}\left((x_p)\right) := \left(\prod_{p \in \operatorname{Val}} \left\|\mathbf{s}_{\mathbf{L}}(x_p)\right\|_p\right)^{-1}$$

Now we choose a finite $S \subset \operatorname{Val}_f$ large enough such that Equa.(3, 4, 5) hold. Moreover, the integral models \mathcal{X} appearing in Equa.(3) and (4) coincide over \mathbb{Z}_S . Then

$$\operatorname{Ht}\left((x_p)\right) = \left(\left\|\mathbf{s}_{\mathbf{L}}(x)\right\|_{\infty} \cdot \prod_{p \in S} \left\|\mathbf{s}_{\mathbf{L}}(x_p)\right\|_p\right)^{-1}, \ \forall x \in \mathscr{K}.$$

A great advantage of adelic height is that, by the product formula, this height is independent of the choice of sections.

2.1.3. Orbits of arithmetic subgroups. Keep the finite set $S \subset \operatorname{Val}_f$ from the last subsection. Assume that **X** is a **G**-equivariant compactification of \mathbf{G}/\mathbf{H} and **D** is the complement of \mathbf{G}/\mathbf{H} . Let \mathscr{K} and an adelic line bundle $(L, (\|\cdot\|_p)_{p \in \operatorname{Val}})$ be given as above. By enlarging S, we assume further that we have a smooth model \mathcal{G} of **G** over \mathbb{Z}_S and that the action $G \times \mathbf{U} \to \mathbf{U}$ over \mathbb{Q} extends to $\mathcal{G} \times \mathcal{U} \to \mathcal{U}$ over \mathbb{Z}_S . Note the following:

Lemma 2.1. For $p \in \operatorname{Val}_f$, the subgroup

$$\operatorname{Stab}_{\mathbf{G}}(\left\|\cdot\right\|_{p}) := \left\{ g \in \mathbf{G}(\mathbb{Q}_{p}) \mid \left\|\mathbf{s}_{\mathbf{L}}\right\|_{g.x} = \left\|\mathbf{s}_{\mathbf{L}}\right\|_{x}, \ \forall x \in K_{p} \right\}$$

contains an open, compact subgroup of $\mathbf{G}(\mathbb{Q}_p)$.

Proof. Indeed, the function $x \mapsto ||s_{\mathbf{L}}(x)||_p$ is smooth and hence locally constant on $\mathcal{U}(\mathbb{Q}_p)$ and in particular on K_p . By compactness and the totally disconnectedness of p-adic topology, we can cover K_p by disjoint open sets such that $||s_{\mathbf{L}}(x)||_p$ is constant on each open set. By choosing a small enough open compact subgroup $K \subset \mathbf{G}(\mathbb{Q}_p)$, we can guarantee that each open subset is K-stable. Such a K is thus contained in $\mathrm{Stab}_{\mathbf{G}}(||\cdot||_p)$, proving the claim.

Let

$$\Gamma_p := \begin{cases} \text{an open, compact subgroup of } \operatorname{Stab}_{\mathbf{G}}(\|\cdot\|_p), & p \in S \\ \mathcal{G}(\mathbb{Z}_p), & p \notin S, \end{cases}$$

and

$$\Gamma := \mathbf{G}(\mathbb{Q}) \cap \prod_{p \in \operatorname{Val}_f} \Gamma_p.$$

Lemma 2.2. The Γ defined above is an arithmetic subgroup of $\mathbf{G}(\mathbb{Q})$ that preserves $\mathbf{U}_{\mathscr{K}}(\mathbb{Z})$. Moreover,

$$\operatorname{Ht}_p(x) = \operatorname{Ht}_p(\gamma . x_p), \ \forall p \in \operatorname{Val}_f, \ x = (x_p) \in \mathbf{U}_{\mathscr{K}}(\mathbb{Z}), \ \gamma \in \mathbf{I}$$

where $\operatorname{Ht}_{p}(x) := \|\mathbf{s}_{\mathbf{L}}(x_{p})\|_{p}^{-1}$.

The theorem below follows from [GMB13, Proposition 7.13, 7.14]. Special cases can be proved using reduction theory (see [Bor19, Remark 6.4]).

Theorem 2.3. There are only finitely many Γ -orbits on $\mathbf{U}_{\mathscr{K}}(\mathbb{Z})$.

We sketch a proof following [WX16] assuming H is connected for the sake of completeness.

Sketch of proof assuming \mathbf{H} to be connected. The conclusion is going to follow from a spreading-out argument, finiteness of Shafarevich–Tate groups and finiteness of class numbers of linear algebraic groups. We define

$$x \sim_{\text{loc}} y \iff \Gamma_p \cdot x = \Gamma_p \cdot y, \ \forall p \in \text{Val}_f$$

for two elements $x = (x_p)$ and $y = (y_p)$ in $\mathbf{U}_{\mathscr{K}}(\mathbb{Z})$. Let $[x]_{\mathrm{loc}}$ denote the equivalence class containing x. As a first step, we note that $[x]_{\mathrm{loc}}$ is finite (compare [BR95, Lemma 1.6.4]). Indeed, for $p \notin S$, there always exists $\gamma_p \in \Gamma_p$ such that $y_p = \gamma_p x_p$ by Hensel's lemma and Lang's theorem on triviality of the Galois cohomology group $H^1(F, \mathbf{H})$ for a finite field F and connected \mathbf{H} . So it remains to show that for each $p \in S$, $K_p \cap \mathbf{U}(\mathbb{Q})$ is finite modulo the equivalence relation defined by $x \sim y$ iff $\Gamma_p . x = \Gamma_p . y$, which follows from the finiteness of K_p/Γ_p .

We define an equivalence relation $\sim_{\mathbb{Q}}$ on $\mathbf{U}_{\mathscr{K}}(\mathbb{Z})$ by

$$x \sim_{\mathbb{Q}} y \iff \mathbf{G}(\mathbb{Q}).x = \mathbf{G}(\mathbb{Q}).y$$

Then we show that for each x, the number of equivalence classes $[x]_K / \sim_{\mathbb{Q}}$ is finite. Indeed, by inspecting the following commutative diagram

$$1 \longrightarrow \mathbf{H}(\mathbb{Q}) \longrightarrow \mathbf{G}(\mathbb{Q}) \longrightarrow \mathbf{U}(\mathbb{Q}) \longrightarrow H^{1}(\mathbb{Q}, \mathbf{H})$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$1 \longrightarrow \mathbf{H}(\mathbb{A}_{\mathbb{Q}}) \longrightarrow \mathbf{G}(\mathbb{A}_{\mathbb{Q}}) \longrightarrow \mathbf{U}(\mathbb{A}_{\mathbb{Q}}) \longrightarrow \prod_{p \in \operatorname{Val}_{f}} H^{1}(\mathbb{Q}_{p}, \mathbf{H}),$$

we have an injection

$$[x]_{\mathrm{loc}} / \sim_{\mathbb{Q}} \hookrightarrow \ker \left(H^1(\mathbb{Q}, \mathbf{H}) \to \prod_p H^1(\mathbb{Q}_p, \mathbf{H}) \right).$$

But the latter is finite (See [BS64, Theorem 7.1]).

For $y \in [x]_{\text{loc}}$, let $[y]_{\text{loc},\mathbb{Q}} := [x]_{\text{loc}} \cap [y]_{\mathbb{Q}}$ denote the equivalence class containing y. As the final step, we show that this is finite for every $x \in \mathbf{U}_{\mathscr{K}}(\mathbb{Z})$ and $y \in [x]_{\text{loc}}$. For each $z \in [y]_{\text{loc},\mathbb{Q}}$, we find $q_z \in \mathbf{G}(\mathbb{Q})$ and $\gamma_z \in \prod \Gamma_p$ such that $z = q_z y = \gamma_z y$. Then the map

$$[y]_{\mathrm{loc},\mathbb{Q}} \to \mathbf{H}_y(\mathbb{Q}) \backslash \mathbf{H}_y(\mathbb{A}_{\mathbb{Q}}) / \mathbf{H}_y(\mathbb{A}_{\mathbb{Q}}) \cap \prod \Gamma_p$$
$$z \mapsto q_z^{-1} \gamma_z$$

is well-defined and injective. As the right hand side is finite (see [PRR23, Theorem 5.8] or [Bor63, Theorem 5.1]), we are done.

Remark 2.4. In light of Lemma 2.2 and Theorem 2.3, it makes no difference to count with respect to the adelic height or the height at infinity.

2.2. Invariant gauge forms. Assume that (\mathbf{X}, \mathbf{D}) is a smooth **G**-pair over \mathbb{Q} with $\mathbf{U} = \mathbf{X} \setminus \mathbf{D}$ isomorphic to \mathbf{G}/\mathbf{H} **G**-equivariantly. Let $o \in \mathbf{U}(\mathbb{Q})$ denote the identity coset. Let $\mathcal{T}_{\mathbf{X}}$ be the tangent bundle of \mathbf{X} , $\mathcal{T}_{\mathbf{X}}^*$ be its dual and $\mathcal{K}_{\mathbf{X}} := \det(\mathcal{T}_{\mathbf{X}}^*)$ be the canonical line bundle. Let $\langle \cdot, \cdot \rangle$ denote the natural pairing between $\mathcal{T}_{\mathbf{X}}$ and $\mathcal{T}_{\mathbf{X}}^*$ and their wedge products.

An element $g \in \mathbf{G}(\mathbb{C})$ acts on $\mathbf{X}(\mathbb{C})$ and hence on tangent vectors by $(g, v) \mapsto g_* v$ and differential forms by $(g, \omega) \mapsto g^* \omega$. From the definition, one has that for every analytically open subset $\mathcal{O} \subset \mathbf{X}(\mathbb{C})$, $\omega \in \Gamma(\mathcal{O}, \mathcal{T}^*_{\mathbf{X}})$ and $\partial \in \Gamma(\mathcal{O}, \mathcal{T}_{\mathbf{X}})$,

$$\langle \omega, \partial \rangle_{g.x} = \langle g^* \omega, (g^{-1})_* \partial \rangle_x, \ \forall x \in g^{-1} \mathcal{O}, \ g \in \mathbf{G}(\mathbb{C}).$$

Identify the fibre $\mathcal{T}_o(\mathbf{X})$ (resp. $\mathcal{T}_o^*(\mathbf{X})$) of the tangent (resp. cotangent) bundle at o with $\mathfrak{g}/\mathfrak{h}$ (resp. $(\mathfrak{g}/\mathfrak{h})^*$). The natural pairing on $((\mathfrak{g}/\mathfrak{h})^*, \mathfrak{g}/\mathfrak{h})$ is then identified with that on $(\mathcal{T}_o^*(\mathbf{X}), \mathcal{T}_o(\mathbf{X}))$. Let $\Delta_{\mathbf{G}}$ (resp. $\Delta_{\mathbf{H}}$) denote the determinant character of the adjoint action of \mathbf{G} (resp. \mathbf{H}) on \mathfrak{g} (resp. \mathfrak{h}), that is, $\Delta_{\mathbf{G}}(g) = \det(\mathrm{Ad}(g), \mathfrak{g})$ and $\Delta_{\mathbf{H}}(h) = \det(\mathrm{Ad}(h), \mathfrak{h})$.

Since **H** stabilizes o, its actions on $\mathcal{T}_{\mathbf{X}}$ and $\mathcal{T}_{\mathbf{X}}^*$ induce actions on $\mathcal{T}_o(\mathbf{X})$ and $\mathcal{T}_o^*(\mathbf{X})$. These actions are identified with the adjoint action of **H** on $\mathfrak{g}/\mathfrak{h}$ and $(\mathfrak{g}/\mathfrak{h})^*$ respectively. Therefore,

$$\det(h_*, \mathcal{T}_o(\mathbf{X})) = \det(\mathrm{Ad}(h), \mathfrak{g}/\mathfrak{h}) = \frac{\det(\mathrm{Ad}(h), \mathfrak{g})}{\det(\mathrm{Ad}(h), \mathfrak{h})} = \frac{\Delta_{\mathbf{G}}(h)}{\Delta_{\mathbf{H}}(h)}.$$

and

$$\det((h^{-1})^*, \mathcal{T}_o^*(\mathbf{X})) = \det(h_*, \mathcal{T}_o(\mathbf{X}))^{-1} = \frac{\Delta_{\mathbf{H}}(h)}{\Delta_{\mathbf{G}}(h)}$$

Define $(\Delta_{\mathbf{G}/\mathbf{H}})^{-1} := \Delta_{\mathbf{G}}^{-1} \Delta_{\mathbf{H}}$, an element of $\mathfrak{X}^*_{\mathbb{Q}}(\mathbf{H})$.

For a Q-character α of **H**, let \mathcal{L}_{α} be the GIT quotient of $\mathbf{G} \times \operatorname{Spec} \mathbb{Q}[x]$ by the right action of **H** defined by

$$(g,\lambda) \cdot h := (gh, \alpha(h)^{-1}\lambda)$$

It is a **G**-linearized line bundle over **G**/**H**. On the other hand, given a **G**-linearized line bundle \mathcal{L} over **G**/**H**, the (left) action of **H** on the fibre of *o* gives a \mathbb{Q} -character $\alpha_{\mathcal{L}}$ of **H**. One can check the following:

Lemma 2.5. Let \mathcal{L} be a **G**-linearized line bundle, then $\mathcal{L} \cong \mathcal{L}_{\alpha_{\mathcal{L}}}$. In particular, $\mathcal{K}_{\mathbf{G}/\mathbf{H}} \cong \mathcal{L}_{\Delta_{\mathbf{G}/\mathbf{H}}^{-1}}$.

Lemma 2.6. Let χ be a \mathbb{Q} -character of \mathbf{H} ,

$$\Gamma(\mathbf{G}/\mathbf{H}, \mathcal{L}_{\chi}) \cong \left\{ f \in \mathbb{Q}[\mathbf{G}] \mid f(gh) = \chi^{-1}(h)f(g), \ \forall (g, h) \in \mathbf{G} \times \mathbf{H} \right\}$$

By [KKV89, 2.2, 2.3], the non-vanishing global sections of $\mathcal{K}_{\mathbf{G}/\mathbf{H}}$ consist of **G**-eigenvectors. Combined with the lemma above, we have **Lemma 2.7.** The non-vanishing global sections of $\mathcal{K}_{\mathbf{G}/\mathbf{H}}$ consist of

$$\Gamma(\mathbf{G}/\mathbf{H},\mathcal{K}_{\mathbf{G}/\mathbf{H}})^{\times} \cong \left\{ f \in \mathbb{Q}[\mathbf{G}] \mid f = \lambda \chi, \ \exists \lambda \in \mathbb{Q}^{\times}, \ \chi \in \mathfrak{X}_{\mathbb{Q}}^{*}(\mathbf{G}) \ with \ \chi|_{\mathbf{H}} = \Delta_{\mathbf{G}/\mathbf{H}} \right\}.$$

Hence we may label, up to a scalar, non-vanishing global sections of $\mathcal{K}_{\mathbf{G}/\mathbf{H}}$ as $\omega_{\mathbf{G}/\mathbf{H}}^{\chi}$ with χ ranging over extensions of $\Delta_{\mathbf{G}/\mathbf{H}}$ to \mathbf{G} . In particular, $\mathcal{K}_{\mathbf{G}/\mathbf{H}}$ is trivial iff $\Delta_{\mathbf{H}}$ extends to a \mathbb{Q} -character on \mathbf{G} .

If both $\Delta_{\mathbf{G}/\mathbf{H}}$ and χ are trivial characters, we abbreviate $\omega_{\mathbf{G}/\mathbf{H}} := \omega_{\mathbf{G}/\mathbf{H}}^{\chi}$ and call it an *invariant* gauge form. One may wish to compare the discussions above with those in [Rag72, Chapter 1] or [Wei82, Chapter 2].

Lemma 2.8. If $\mathcal{K}_{\mathbf{G}/\mathbf{H}}$ is trivial, then \mathbf{H} is observable in \mathbf{G} .

Proof. By Lemma 2.7, $\Delta_{\mathbf{H}}^{-1}$ extends to a character β on \mathbf{G} and take some non-zero vector \mathbf{v}_{β} in the one dimensional representation where \mathbf{G} acts by β . Let $\mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\beta}$ be the stabilizer of $\mathbf{v}_{\mathfrak{h}} \otimes \mathbf{v}_{\beta}$, where $\mathbf{v}_{\mathfrak{h}} \in \wedge^{\dim \mathbf{H}} \mathfrak{g}$ is a lift of \mathfrak{h} .

Then $\mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\beta}$ is observable in \mathbf{G} by definition. On the other hand, \mathbf{H} is contained in $\mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\beta}$ as a normal subgroup and is hence observable in $\mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\beta}$. By the transitivity of observability, \mathbf{H} is observable in \mathbf{G} .

Remark 2.9. The converse is not true. For instance, if $G := SL_3$ and H is generated by elements of the form

$$\left[\begin{array}{ccc}t&\\&t\\&&t^{-2}\end{array}\right]\cdot\left[\begin{array}{ccc}1&\\&1&s\\&&1\end{array}\right].$$

Then **H** is observable in **G** yet $\Delta_{\mathbf{H}}$ is not extendable to **G**.

2.3. Boundary components of anti-canonical divisors. Assumptions as in the last subsection and assume that $\mathcal{K}_{\mathbf{G}/\mathbf{H}}$ is a trivial line bundle. Here we are interested in the anti-canonical divisor $-\operatorname{div}(\omega_{\mathbf{G}/\mathbf{H}}^{\chi})$ on \mathbf{X} where $\chi \in \mathfrak{X}_{\mathbb{Q}}^{*}(\mathbf{G})$ is a fixed extension of $\Delta_{\mathbf{G}/\mathbf{H}}$.

Note that \mathfrak{g} naturally maps to $\Gamma(\mathbf{X}, \mathcal{T}_{\mathbf{X}})$, inducing

$$\wedge^{d} \mathfrak{g} \to \Gamma(\mathbf{X}, \wedge^{d} \mathcal{T}_{\mathbf{X}}) \cong \Gamma(\mathbf{X}, (\mathcal{K}_{\mathbf{X}})^{*}).$$

For $\mathbf{v} = v_1 \wedge ... \wedge v_d \in \wedge^d \mathfrak{g}$, let $\partial_{\mathbf{v}}$ denote its image. Then for each $\mathbf{v} \in \wedge^d \mathfrak{g}$,

$$\varphi_{\mathbf{v}}(x) := \langle \omega_{\mathbf{G}/\mathbf{H}}^{\chi}, \partial_{\mathbf{v}} \rangle_x$$

defines a rational function on **X** that is regular on **U**. Thus, to understand the boundary behaviour of $\omega_{\mathbf{G}/\mathbf{H}}^{\chi}$, it suffices to understand that of $\varphi_{\mathbf{v}}$ and $\partial_{\mathbf{v}}$.

Take a nonzero element $\mathbf{l}_{\mathfrak{h}} \in \wedge^{d}(\mathfrak{g}/\mathfrak{h})^{*} \hookrightarrow \wedge^{d}\mathfrak{g}^{*}$. So $\left(\omega_{\mathbf{G}/\mathbf{H}}^{\chi}\right)_{o} = c_{\chi}\mathbf{l}_{\mathfrak{h}}$ for some $c_{\chi} \neq 0$.

Lemma 2.10. For $g \in \mathbf{G}(\mathbb{C})$, $\varphi_{\mathbf{v}}(g.o) = c_{\chi}\chi(g) \cdot \langle \mathbf{l}_{\mathfrak{h}}, \mathrm{Ad}(g)^{-1}\mathbf{v} \rangle$.

Proof. This follows from a direct computation:

$$\begin{split} \langle \omega_{\mathbf{G}/\mathbf{H}}^{\chi}, \partial_{\mathbf{v}} \rangle_{g.o} &= \chi(g) \cdot \langle (g^{-1})^* \omega_{\mathbf{G}/\mathbf{H}}^{\chi}, \partial_{\mathbf{v}} \rangle_{g.o} \\ &= \chi(g) \cdot \langle \omega_{\mathbf{G}/\mathbf{H}}^{\chi}, (g^{-1})_* \partial_{\mathbf{v}} \rangle_o \\ &= \chi(g) \cdot \langle \omega_{\mathbf{G}/\mathbf{H}}^{\chi}, \partial_{\mathrm{Ad}(g)^{-1}\mathbf{v}} \rangle_o \\ &= c_{\chi} \chi(g) \cdot \langle \mathbf{l}_{\mathfrak{h}}, \mathrm{Ad}(g)^{-1} \mathbf{v} \rangle. \end{split}$$

2.3.1. Dual. To simplify the conclusion of Lemma 2.10, we use a dual operation.

Fix $\omega_0 \in \wedge^{\dim \mathbf{G}} \mathfrak{g}^*$ and $\omega_0^* \in \wedge^{\dim \mathbf{G}} \mathfrak{g}$ with $\langle \omega_0^*, \omega_0 \rangle = 1$. For two positive integers $l + l' = \dim \mathbf{G}$, we have a natural linear isomorphism:

$$\wedge^{l}\mathfrak{g} \to \wedge^{l'}\mathfrak{g}^{*} \cong (\wedge^{l'}\mathfrak{g})^{*}$$
$$\mathbf{v} \mapsto \mathbf{v}^{\vee}$$

defined by

$$\mathbf{v}\wedge\mathbf{w}=\langle\mathbf{v}^ee,\mathbf{w}
angle\omega_0^*,\ orall\,\mathbf{w}\in\wedge^{l'}\mathfrak{g}$$

Note that $\langle \mathbf{v}^{\vee}, \mathbf{w} \rangle = \langle \mathbf{v}, \mathbf{w}^{\vee} \rangle$. Furthermore,

$$\langle (\operatorname{Ad}(g)\mathbf{v})^{\vee}, \mathbf{w} \rangle \omega_0^* = \operatorname{Ad}(g)\mathbf{v} \wedge \mathbf{w} = \operatorname{Ad}(g)(\mathbf{v} \wedge \operatorname{Ad}(g)^{-1}\mathbf{w}) = \operatorname{Ad}(g) \left(\langle \mathbf{v}^{\vee}, \operatorname{Ad}(g^{-1})\mathbf{w} \rangle \omega_0^* \right) = \Delta_{\mathbf{G}}(g) \langle \mathbf{v}^{\vee}, \operatorname{Ad}(g^{-1})\mathbf{w} \rangle \omega_0^*.$$

Thus, $(\operatorname{Ad}(g)\mathbf{v})^{\vee} = \Delta_{\mathbf{G}}(g) \operatorname{Ad}^*(g)\mathbf{v}^{\vee}$. So we see that $(\cdot)^{\vee}$ intertwines Ad and $\operatorname{Ad}^* \otimes \Delta_{\mathbf{G}}$, or equivalently, $\operatorname{Ad} \otimes \Delta_{\mathbf{G}}^{-1}$ and Ad^* .

Now take a $\mathbb{Q}\text{-vector }\mathbf{v}_\mathfrak{h}$ lifting \mathfrak{h} such that $\mathbf{v}_\mathfrak{h}^\vee=l_\mathfrak{h}.$ Then we have

$$\begin{split} \langle \mathbf{l}_{\mathfrak{h}}, \mathrm{Ad}(g)^{-1} \mathbf{v} \rangle &= \langle \mathrm{Ad}^*(g) \mathbf{v}_{\mathfrak{h}}^{\vee}, \mathbf{v} \rangle = \Delta_{\mathbf{G}}(g)^{-1} \langle (\mathrm{Ad}(g) \mathbf{v}_{\mathfrak{h}})^{\vee}, \mathbf{v} \rangle \\ &= \Delta_{\mathbf{G}}(g)^{-1} \langle \mathrm{Ad}(g) \mathbf{v}_{\mathfrak{h}}, \mathbf{v}^{\vee} \rangle. \end{split}$$

Combined with Lemma 2.10 above,

Lemma 2.11. For $g \in \mathbf{G}(\mathbb{C})$, $\varphi_{\mathbf{v}}(g.o) = c_{\chi}\chi(g)\Delta_{\mathbf{G}}(g)^{-1}\langle \operatorname{Ad}(g)\mathbf{v}_{\mathfrak{h}}, \mathbf{v}^{\vee} \rangle$.

2.3.2. The conclusion. For $\mathbf{v} \in \wedge^d \mathfrak{g}$ and $\alpha \in \mathscr{A}$, we define integers $d_{\alpha}, d_{\alpha}^{\mathbf{v}}$ and $d_{\alpha}^{\varphi, \mathbf{v}}$ by

$$-\operatorname{div}(\omega_{\mathbf{G}/\mathbf{H}}^{\chi}) := \sum_{\alpha \in \mathscr{A}} d_{\alpha} \mathbf{D}_{\alpha}, \ \operatorname{div}(\partial_{\mathbf{v}}) := \sum_{\alpha \in \mathscr{A}} d_{\alpha}^{\mathbf{v}} \mathbf{D}_{\alpha}, \ -\operatorname{div}(\varphi_{\mathbf{v}}) := \sum_{\alpha \in \mathscr{A}} d_{\alpha}^{\varphi,\mathbf{v}} \mathbf{D}_{\alpha}.$$

By the definition of $\varphi_{\mathbf{v}}$, we have

$$-\operatorname{div}(\omega_{\mathbf{G}/\mathbf{H}}^{\chi}) = \operatorname{div}\partial_{\mathbf{v}} - \operatorname{div}(\varphi_{\mathbf{v}}), \text{ or } d_{\alpha} = d_{\alpha}^{\mathbf{v}} + d_{\alpha}^{\varphi,\mathbf{v}}, \,\forall \, \alpha \in \mathscr{A}.$$
(6)

Each \mathbf{D}_{α} has dimension smaller than d, one sees that $d_{\alpha}^{\mathbf{v}} \geq 1$ for every \mathbf{v} and α . As a corollary of Lemma 2.11 we have

Theorem 2.12. Let $\alpha \in \mathscr{A}$, $x \in U(\mathbb{C})$ and \mathbf{H}_x be the stabilizer of x in \mathbf{G} . Let (g_n) be a sequence in $\mathbf{G}(\mathbb{C})$ such that

- (1) $\lim_{n\to\infty} g_n \cdot x \in \mathbf{D}_{\alpha}(\mathbb{C})^{\circ}$ and $|\chi(g_n)\Delta_{\mathbf{G}}(g_n)^{-1}| = 1;$
- (2) $(\operatorname{Ad}(g_n).\mathbf{v}_{\mathfrak{h}_x})$ is bounded away from **0**.

Then $d_{\alpha} \geq 1$. If additionally

(3) $(\operatorname{Ad}(g_n).\mathbf{v}_{\mathfrak{h}_x})$ diverges,

then $d_{\alpha} \geq 2$.

Here $\mathbf{D}_{\alpha}(\mathbb{C})^{\circ}$ denotes the points in $\mathbf{D}_{\alpha}(\mathbb{C})$ that do not lie in any other $\mathbf{D}_{\beta}(\mathbb{C})$.

Proof. Without loss of generality assume x = o and $\mathbf{H}_x = \mathbf{H}$. Note that after passing to a subsequence,

 $(\operatorname{Ad}(g_n).\mathbf{v}_{\mathfrak{h}})$ bounded away from $\mathbf{0} \implies l(\operatorname{Ad}(g_n).\mathbf{v}_{\mathfrak{h}}) > c, \exists l \in (\wedge^d \mathfrak{g})^*, c > 0;$

$$(\operatorname{Ad}(g_n).\mathbf{v}_{\mathfrak{h}})$$
 diverges $\implies l(\operatorname{Ad}(g_n).\mathbf{v}_{\mathfrak{h}})$ diverges, $\exists l \in (\wedge^d \mathfrak{g})^*$.

By Lemma 2.11 and our assumption, the first line implies the existence of $\mathbf{v} \in \wedge^d \mathfrak{g}$ such that $d_{\alpha}^{\varphi, \mathbf{v}} \ge 0$, implying $d_{\alpha} \ge 1 + d_{\alpha}^{\varphi, \mathbf{v}} \ge 1$. Similarly, the second line implies $d_{\alpha} \ge 2$.

There are two special cases that we would like to mention.

Theorem 2.13. Assume that **H** is reductive and $\chi \cdot \Delta_{\mathbf{G}}^{-1}$ is trivial. Then $d_{\alpha} \geq 1$ for every $\alpha \in \mathscr{A}$. If additionally, **G** is reductive and $\mathbf{Z}_{\mathbf{G}}(\mathbf{H}^{\circ})^{\circ} \subset \mathbf{H}$, then $d_{\alpha} \geq 2$ for every $\alpha \in \mathscr{A}$.

Note that when **H** is reductive, $\delta_{\mathbf{H}}$ is trivial and hence $\mathcal{K}_{\mathbf{G}/\mathbf{H}}$ is trivial. And if χ corresponds to the trivial extension of $\delta_{\mathbf{H}}$, then $\chi \cdot \Delta_{\mathbf{G}}^{-1}$ is trivial.

Proof. If **G** is also reductive, then $\mathbf{G}.\mathbf{v}_{\mathfrak{h}}$ is closed. Indeed, for $\mathbf{H}' := \mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\mathbf{1}}$, the stabilizer of $\mathbf{v}_{\mathfrak{h}}$, the identity component of its centralizer in **G** is in \mathbf{H}' , thus the closedness follows from the result of [Kem78]. For every $\alpha \in \mathscr{A}$ and $x \in \mathbf{D}_{\alpha}(\mathbb{C})^{\circ}$, find $(g_n) \subset \mathbf{G}(\mathbb{C})$ with $\lim g_{n}.o = x$. Then Theorem 2.12 applies and completes the proof.

Now assume moreover that $\mathbf{Z}_{\mathbf{G}}(\mathbf{H}^{\circ})^{\circ} \subset \mathbf{H}$. Let (g_n) and x be the same as above. Thus (g_n) is unbounded modulo \mathbf{H} , and under our assumption, this in turn implies that (g_n) is also unbounded modulo the stabilizer of $\mathbf{v}_{\mathfrak{h}}$. Note that identity component of the normalizer of a connected reductive group (taken to be \mathbf{H}° here) is the same as the identity connected component of the group generated by its centralizer and itself. So $(\mathrm{Ad}(g_n).\mathbf{v}_{\mathfrak{h}})$ diverges and we conclude by Theorem 2.12.

When **G** may not be reductive, we still claim that $\mathbf{0} \notin \overline{\mathbf{G}(\mathbb{C}).\mathbf{v}_{\mathfrak{h}}}$, which is sufficient by discussion above. Find a Levi decomposition over \mathbb{Q} : $\mathfrak{g} = \mathfrak{l} \ltimes \mathfrak{u}$ such that $\mathfrak{h} \subset \mathfrak{l}$. Here \mathfrak{u} denotes the Lie algebra of the unipotent radical $\mathbf{R}_{\mathbf{u}}(\mathbf{G})$ of **G** and \mathfrak{l} is the Lie algebra of a maximal reductive subgroup **L** of **G**.

Decompose $\wedge^{\dim \mathbf{H}} \mathfrak{g} = \wedge^{\dim \mathbf{H}} \mathfrak{l} \oplus W$ where

$$W := \left\{ v \in \wedge^{\dim \mathbf{H}} \mathfrak{g} \mid v \wedge \mathbf{u} = \mathbf{0}, \ \forall \, \mathbf{u} \in \wedge^{\dim \mathfrak{u}} \mathfrak{u} \right\}$$

For an element $g \in \mathbf{G}(\mathbb{C})$, written as $g = u \cdot l$ with $u \in \mathbf{R}_{\mathbf{u}}(\mathbb{C})$ and $l \in \mathbf{L}(\mathbb{C})$, we have

$$\operatorname{Ad}(g)\mathbf{v}_{\mathfrak{h}} = \operatorname{Ad}(ul)\mathbf{v}_{\mathfrak{h}} = \operatorname{Ad}(l)\mathbf{v}_{\mathfrak{h}} + w$$

for some $w \in W$. Hence $\|\operatorname{Ad}(ul)\mathbf{v}_{\mathfrak{h}}\|$ is bounded from below by some positive multiple of $\|\operatorname{Ad}(l)\mathbf{v}_{\mathfrak{h}}\|$. But $\operatorname{Ad}(\mathbf{L}(\mathbb{C}))\mathbf{v}_{\mathfrak{h}}$ is closed and can never approach 0. So our claim follows.

Therefore, if **G** and **H** are both semisimple and the centralizer of **H** in **G** is finite, the log anticanonical bundle of **X** is big by [HTT15, Proposition 5.1]. The reductivity assumption on **H** can not be dropped (see Example 2.15 below). Also, in the second part, the assumption $\mathbf{Z}_{\mathbf{G}}(\mathbf{H}^{\circ})^{\circ} \subset \mathbf{H}$ is necessary. Nevertheless, we have

Theorem 2.14. Let $\Gamma_{\mathbb{C}} \subset \mathbf{G}(\mathbb{Q}(i))$ be an arithmetic subgroup of $\operatorname{Res}_{\mathbb{Q}/\mathbb{Q}(i)}(\mathbf{G})$ and $\alpha \in \mathscr{A}$. Assume that for some $x \in \mathbf{U}(\mathbb{Q}(i))$ and sequence (γ_n) in $\Gamma_{\mathbb{C}}$ one has $\lim_{n\to\infty} \gamma_n \cdot x \in \mathbf{D}_{\alpha}(\mathbb{C})^\circ$, then $d_{\alpha} \geq 1$. Assume additional that (γ_n) can be chosen to be unbounded modulo $\mathbf{N}_{\mathbf{G}}(\mathbf{H})^1(\mathbb{C}) \cap \Gamma_{\mathbb{C}}$, then $d_{\alpha} \geq 2$.

Here $\operatorname{Res}_{\mathbb{Q}/\mathbb{Q}(i)}$ denotes the restriction of scalar of a linear algebraic group (see [Spr98, 11.4,12.4]), and $\mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\mathbf{1}}$ is the stabilizer of $\mathbf{v}_{\mathfrak{h}}$ under the adjoint action.

Related results are obtained in [HT99, Theorem 2.7] for bi-equivariant compactifications of the additive groups \mathbf{G}_a^n .

Proof. Take $x \in \mathbf{U}(\mathbb{Q}(i))$ and let \mathbf{H}_x be the stabilizer of x in \mathbf{G} , which is a subgroup defined over $\mathbb{Q}(i)$. Let $\mathbf{v}_{\mathfrak{h}_x} \in (\wedge^{\dim \mathbf{H}}\mathfrak{g}) \otimes_{\mathbb{Q}} \mathbb{Q}(i)$ be a $\mathbb{Q}(i)$ -vector lifting \mathfrak{h}_x . Define $\varphi_{\mathbf{v}}^x$ by modifying the definition of $\varphi_{\mathbf{v}}$ with o replaced by x and \mathbf{H} replaced by \mathbf{H}_x . Let $\Gamma_{\mathbb{C}}$ be an arithmetic subgroup of $\operatorname{Res}_{\mathbb{Q}(i)/\mathbb{Q}}(\mathbf{G})$ obtained by the restriction of scalars. We view it as a subgroup of $\mathbf{G}(\mathbb{C})$. Note that $\chi \Delta_{\mathbf{G}}^{-1}$ is defined over $\mathbb{Q}(i)$ and thus $(\chi \Delta_{\mathbf{G}}^{-1})(\Gamma_{\mathbb{C}})$ is discrete in \mathbb{C} , implying that $(\chi \Delta_{\mathbf{G}}^{-1})(\Gamma_{\mathbb{C}})$ lies in the unit circle of \mathbb{C} . We have $|\chi(\gamma)\Delta_{\mathbf{G}}(\gamma)^{-1}| = 1$ for $\gamma \in \Gamma_{\mathbb{C}}$. As $\operatorname{Ad}(\Gamma_{\mathbb{C}})\mathbf{v}_{\mathfrak{h}_x}$ is discrete in $\wedge^{\dim \mathbf{H}}\mathfrak{g} \otimes \mathbb{C}$, by similar arguments as those used in the previous theorem, we complete the proof.

Example 2.15. Let $\mathbf{G} := \mathbf{SL}_2$ and \mathbf{H} be generated by upper triangular unipotent matrices. Thus, \mathbf{G}/\mathbf{H} has an invariant gauge form and can be identified with $\mathbf{A}_{\mathbb{Q}}^2 \setminus \{0,0\}$. Note that \mathbf{G}/\mathbf{H} naturally embeds in $\mathbf{P}_{\mathbb{Q}}^2$ as an open and dense subvariety. The action of \mathbf{G} on \mathbf{G}/\mathbf{H} extends to $\mathbf{P}_{\mathbb{Q}}^2$.

Let **X** be the blow-up of $\mathbf{P}_{\mathbb{Q}}^2$ at $\{[0:0:1]\}$, then $\mathbf{X} = \mathbf{U} \sqcup \mathbf{P} \sqcup \mathbf{E}$ where **P** is the proper transform of a hyperplane section and **E** is the exceptional divisor. By explicit computation, $-\omega_{\mathbf{G}/\mathbf{H}} = -\mathbf{E} + 2\mathbf{P}$. Therefore, not all d_{α} 's are positive here. Nevertheless, one can verify that $-\omega_{\mathbf{G}/\mathbf{H}}$ is still an effective and big divisor (See [FZ13]).

2.4. Analytic Clemens complex. Keep the assumptions in the last section. Also assume that $\max\{d_{\alpha}\} > 0$. As a reference, see [CLT10, Section 3.1]. Since we are assuming **D** is a simple normal crossing divisor rather than just normal crossing, the discussions are simplified. For instance, if $\sigma(\mathbf{D}_{\alpha}) \neq \mathbf{D}_{\alpha}$ for $\sigma_{\neq id} \in \operatorname{Gal}(\mathbb{C}/\mathbb{R})$, then $\mathbf{D}_{\alpha}(\mathbb{C}) \cap \mathbf{X}(\mathbb{R}) = \emptyset$.

Let the analytic irreducible components of $\mathbf{D}(\mathbb{R})$ be indexed as (D_{α}) for $\alpha \in \mathscr{A}_{\mathbb{R}}^{\mathrm{an}}$. There is a natural map from $\mathscr{A}_{\mathbb{R}}^{\mathrm{an}}$ to \mathscr{A} . For a subset $I \subset \mathscr{A}_{\mathbb{R}}^{\mathrm{an}}$, let $D_I := \bigcap_{\alpha \in I} D_{\alpha}$. Since $\mathbf{G} := \mathbf{G}(\mathbb{R})^{\circ}$ is connected, each connected component of D_I is G-invariant.

We define a partially ordered set by

$$\mathscr{C}_{\mathbb{R}}^{\mathrm{an}} := \{ (I, Z) \mid I \subset \mathscr{A}_{\mathbb{R}}^{\mathrm{an}}, Z \text{ is a connected component of } D_I \}, (I, Z) \prec (I', Z') \iff I \subset I', Z' \subset Z.$$

$$(7)$$

For $x \in \mathbf{U}(\mathbb{R})$, let

$$\mathscr{A}_{\mathbb{R},x}^{\mathrm{an}} := \left\{ \alpha \in \mathscr{A}_{\mathbb{R}}^{\mathrm{an}} \mid D_{\alpha} \cap \overline{\mathrm{G.}x} \neq \emptyset \right\}$$

And $\mathscr{C}_{\mathbb{R},x}^{\mathrm{an}}$ is defined similarly as $\mathscr{C}_{\mathbb{R}}^{\mathrm{an}}$. Note that G.x is exactly the connected component of $\mathbf{U}(\mathbb{R})$ containing x.

Now we take $\mathbf{L} := \sum_{\alpha \in \mathscr{A}} \lambda_{\alpha} \mathbf{D}_{\alpha}$ to be another divisor defined over \mathbb{R} such that $\lambda_{\alpha} \geq 0$ for all $\alpha \in \mathscr{A}$ and

$$d_{\alpha} > 0 \implies \lambda_{\alpha} > 0.$$

Thus points of bounded heights (associated with **L**) have finite volume. For every $x \in \mathbf{U}(\mathbb{R})$ such that $\lambda_{\alpha} > 0$ for some $\alpha \in \mathscr{A}_{\mathbb{R},x}^{\mathrm{an}}$, let

$$a_x := \max_{\alpha \in \mathscr{A}_{\mathbb{R},x}^{\mathrm{an}}, \, \lambda_{\alpha} \neq 0} \left\{ \frac{d_{\alpha} - 1}{\lambda_{\alpha}} \right\}$$

and

$$\mathscr{C}_{\mathbb{R},x}^{\mathrm{an}}(\mathbf{L}) := \left\{ (I,Z) \in \mathscr{C}_{\mathbb{R},x}^{\mathrm{an}} \mid \frac{d_{\alpha}-1}{\lambda_{\alpha}} = a_x, \, \forall \, \alpha \in I \right\}.$$

Also let $b_x := \dim \mathscr{C}^{\mathrm{an}}_{\mathbb{R},x}(\mathbf{L}).$

Let Vol denote the measure on $\mathbf{U}(\mathbb{R})$ induced from $\omega_{\mathbf{G}/\mathbf{H}}^{\chi}$. Let $\mathrm{Ht}_{\mathbf{L}}$ be the height at ∞ associated with certain smooth metric on \mathbf{L} together with its canonical section. For R > 0, let $B_{R,x} := \{y \in \mathrm{G.}x, \, \mathrm{Ht}(y) < R\}$. The following is proved in [CLT10, Theorem 4.7, Corollary 4.8]:

Theorem 2.16. Take $x \in \mathbf{U}(\mathbb{R})$ such that $\lambda_{\alpha} > 0$ for some $\alpha \in \mathscr{A}_{\mathbb{R},x}^{\mathrm{an}}$. Assume $a_x \ge 0$ and $b_x \ge 1$. As R tends to infinity, there exists $c_x > 0$ such that

$$\operatorname{Vol}(B_{R,x}) \sim \begin{cases} c_x R^{a_x} \log(R)^{b_x - 1} & a_x \neq 0 \\ c_x \log(R)^{b_x} & a_x = 0. \end{cases}$$

The family of probability measures on $\mathbf{X}(\mathbb{R})$

$$\mu_{R,x} := \frac{\mathbf{1}_{B_{R,x}} \cdot \operatorname{Vol}}{\operatorname{Vol}(B_{R,x})}$$

has a limit ν under the weak-* topology. Furthermore, ν is a sum of smooth measures on Z as (I, Z) varies over faces of $\mathscr{C}_{\mathbb{R},x}^{\mathrm{an}}(\mathbf{L})$ of dimension b_x .

3. Equidistribution and Orbital Counting

In this section, we explain the relation between equidistribution and orbital counting. Compared to [DRS93, EM93], the possibility of focusing is also considered. On the other hand, we put more restrictions on the height functions, which do not apply to those coming from Riemannian metrics on symmetric spaces.

3.1. Orbital counting follows from equidistribution. Let G be a locally compact second countable topological group and H be a closed subgroup. Let Γ be a discrete subgroup of G. Let m_G (resp. m_H) be a left G-invariant locally finite measure on G (resp. on H). Let $o \in G/H$ denote the identity coset.

Assumption 3.1. We assume that

- (1) G/Γ has a G-invariant locally finite measure $m_{[G]}$;
- (2) $H\Gamma/\Gamma$ is closed in G/Γ and supports an H-invariant locally finite measure $m_{[H]}$;
- (3) there exists a G-invariant locally finite measure $m_{G/H}$ on G/H and that the triple $(m_G, m_H, m_{G/H})$ is compatible, that is,

$$\int f(x) \operatorname{m}_{G}(x) = \int \int f(gh) \operatorname{m}_{H}(h) \operatorname{m}_{G/H}([g])$$
(8)

for every compactly supported continuous function f on G.

Item (1) always holds in the arithmetic setting. By [Rag72, Lemma 1.4], item (3) above is equivalent to the coincidence of the modular functions Δ_G and Δ_H on H. For instance, if G is a semisimple Lie group, then Δ_G is trivial and such a $m_{G/H}$ exists iff Δ_H is trivial.

Let $l: G/H \to \mathbb{R}_{>0}$ be a continuous function and for a positive number R, define

$$B_R := \{ x \in G/H \mid l(x) \le R \}.$$

If we wish to emphasize the role of l, an upper index will be added B_R^l . Assume that $m_{G/H}(B_R)$ is finite for each R.

A family of $(B_R)_{R\geq 1}$ is said to be *well-rounded* (see [EM93, Proposition 1.3]) iff for every $0 < \varepsilon < 1$, there exists an open neighborhood $\mathcal{O}_{\varepsilon}$ of id $\in G$ such that

$$(1-\varepsilon)\mathbf{m}_{G/H}\left(\bigcup_{g\in\mathcal{O}_{\varepsilon}}gB_R\right) \le \mathbf{m}_{G/H}(B_R) \le (1+\varepsilon)\mathbf{m}_{G/H}\left(\bigcap_{g\in\mathcal{O}_{\varepsilon}}gB_R\right).$$
(9)

Remark 3.2. This is different from the notion of (two-sided) well-roundedness as introduced in [GN12] which does not seem to hold in many examples considered in this paper.

Here we assume that one can find $\mathcal{O}_{\varepsilon}$ such that additionally the following is true for all R > 1:

$$B_{(1-\varepsilon)R} \subset \bigcap_{g \in \mathcal{O}_{\varepsilon}} gB_R \subset \bigcup_{g \in \mathcal{O}_{\varepsilon}} gB_R \subset B_{(1+\varepsilon)R}.$$
(10)

This property allows us to prove counting results in the presence of focusing and it is satisfied by all the heights considered in this paper.

A family $(C_R)_{R\geq 1}$ of positive real numbers is said to have a *polynomial asymptotic* if there exist $c_1 > 0, c_2, c_3 \geq 0$ such that $(c_1 R^{c_2} (\ln R)^{c_3}) / C_R \to 1$ as $R \to +\infty$.

For the application when $m_{[H]}$ is an infinite measure, we consider a weighted variant. Assume that $\mathbf{w} : G/H \to \mathbb{R}_{>0}$ satisfies the following: For every $\varepsilon > 0$, there exists an open neighborhood $\mathcal{O}_{\varepsilon}$ of $id \in G$ such that

$$(1-\varepsilon)\mathbf{w}_{gx} \le \mathbf{w}_x \le (1+\varepsilon)\mathbf{w}_{gx}, \quad \forall x \in G/H, \ \forall g \in \mathcal{O}_{\varepsilon}.$$
(11)

The weighted orbital counting problem is to study the asymptotic of the function $\Phi_R^{\mathbf{w}} : G/\Gamma \to \mathbb{R}$ defined by

$$\Phi_R^{\mathbf{w}}([g]) := \frac{1}{C_R} \sum_{x \in g \Gamma. o \cap B_R} \mathbf{w}_x$$

as R tends to $+\infty$.

Theorem 3.3. Assume that Assumption 3.1 is met, Equa.(9,10,11) hold and (C_R) has polynomial asymptotic. Assume further that there exists a nonzero locally finite measure μ_{∞} on G/Γ such that in the weak-* topology,

$$\lim_{R \to \infty} \frac{1}{C_R} \int_{[g] \in B_R} \left(\mathbf{w}_{g.o} g_* \mathbf{m}_{[H]} \right) \, \mathbf{m}_{G/H}([g]) = \mu_{\infty}.$$
(12)

Then there exists a positive continuous function f_{∞} such that $\mu_{\infty} = f_{\infty} \cdot m_{[G]}$ and $\lim_{R \to \infty} \Phi_{R}^{\mathbf{w}}([g]) = f_{\infty}([g])$ for every $[g] \in G/\Gamma$. Actually, if Equa.(12) holds only against compactly supported continuous functions supported in some fixed open neighborhood of the identity coset, then we still have that $\mu_{\infty} = f_{\infty} \cdot m_{[G]}$ when restricted to this neighborhood with f_{∞} continuous and $\lim_{R \to \infty} \Phi_{R}^{\mathbf{w}}([g]) = f_{\infty}([g])$ in this neighborhood.

3.2. **Proof of the theorem.** The proof follows similar lines as [EM93, Section 5] or [DRS93, Section 2] with some modifications.

3.2.1. Step 1, limit exists and is continuous. Fix $[g_0] \in G/\Gamma$, we first show that the limit $\lim_{R\to\infty} \Phi_R^{\mathbf{w}}([g_0])$ exists and is continuous.

For $0 < \varepsilon < 1$, choose $\mathcal{O}_{\varepsilon}$ satisfying Equa.(9, 10, 11). By shrinking to a smaller one we assume that $\mathcal{O}_{\varepsilon} = \mathcal{O}_{\varepsilon}^{-1}$.

Then we choose $V'_{\varepsilon} \subset V_{\varepsilon} \subset \mathcal{O}_{\varepsilon}$ to be two families of open neighborhoods of identity such that the closure of V'_{ε} is contained in V_{ε} and

$$\lim_{\varepsilon \to 0} \frac{\mu_{\infty}(V_{\varepsilon}'[g_0])}{\mu_{\infty}(V_{\varepsilon}[g_0])} = \lim_{\varepsilon \to 0} \frac{\mathrm{m}_{[G]}(V_{\varepsilon}'[g_0])}{\mathrm{m}_{[G]}(V_{\varepsilon}[g_0])} = 1.$$

Then we choose a continuous function f_{ε} with $1_{V'_{\varepsilon}[g_0]} \leq f_{\varepsilon} \leq 1_{V_{\varepsilon}[g_0]}$. By an unfolding and folding argument as in [DRS93, EM93], we have

$$\begin{split} \langle \Phi_R^{\mathbf{w}}, f_{\varepsilon} \rangle_{\mathbf{m}_{[G]}} &:= \int_{x \in G/\Gamma} \Phi_R^{\mathbf{w}}(x) f_{\varepsilon}(x) \operatorname{m}_{[G]}(x) \\ &= \frac{1}{C_R} \int_{[g] \in B_R} \mathbf{w}_{g.o} \left(\int_{x \in G/\Gamma} f_{\varepsilon}(gx) \operatorname{m}_{[H]}(x) \right) \operatorname{m}_{G/H}([g]) \\ &= \int_{x \in G/\Gamma} f_{\varepsilon}(x) \mu_R(x) \end{split}$$

where

$$\mu_R := \frac{1}{C_R} \int_{[g] \in B_R} \left(\mathbf{w}_{g.o} g_* \mathbf{m}_{[H]} \right) \mathbf{m}_{G/H}([g]).$$

Therefore,

$$\limsup_{R \to \infty} \langle \Phi_R^{\mathbf{w}}, \mathbf{1}_{V_{\varepsilon}'[g_0]} \rangle_{\mathbf{m}_{[G]}} \leq \lim_{R \to \infty} \langle \Phi_R^{\mathbf{w}}, f_{\varepsilon} \rangle_{\mathbf{m}_{[G]}}
= \int f_{\varepsilon}(x) \mu_{\infty}(x) \leq \liminf_{R \to \infty} \langle \Phi_R^{\mathbf{w}}, \mathbf{1}_{V_{\varepsilon}[g_0]} \rangle_{\mathbf{m}_{[G]}}.$$
(13)

Also note that

$$\mu_{\infty}(V_{\varepsilon}'[g_0]) \le \int f_{\varepsilon}(x)\mu_{\infty}(x) \le \mu_{\infty}(V_{\varepsilon}[g_0]).$$
(14)

Assume ε to be small enough so that the natural map $\mathcal{O}_{\varepsilon} \to \mathcal{O}_{\varepsilon}[g_0]$ is a homeomorphism. Then $\lambda_0 \mathbf{m}_G|_{\mathcal{O}_{\varepsilon}}$ is identified with $\mathbf{m}_{[G]}|_{\mathcal{O}_{\varepsilon}[g_0]}$ under this homeomorphism for some $\lambda_0 > 0$.

Now we can start to estimate:

(by

$$\begin{split} \langle \Phi_R^{\mathbf{w}}, \mathbf{1}_{V_{\varepsilon}'[g_0]} \rangle_{\mathbf{m}_{[G]}} &= \frac{1}{C_R} \int_{[g] \in V_{\varepsilon}'[g_0]} \left(\sum_{x \in g\Gamma.o \cap B_R} \mathbf{w}_x \right) \, \mathbf{m}_{[G]}([g]) \\ &= \frac{1}{C_R} \int_{g \in V_{\varepsilon}'} \left(\sum_{x \in g_0 \Gamma.o \cap g^{-1}B_R} \mathbf{w}_{gx} \right) \, \lambda_0 \mathbf{m}_G(g) \\ (\text{by Equa.}(11)) &\geq \frac{\mathbf{m}_{[G]}(V_{\varepsilon}'[g_0])}{(1+\varepsilon)C_R} \left(\sum_{x \in g_0 \Gamma.o \cap (\bigcap_{g \in V_{\varepsilon}'}g^{-1}B_R)} \mathbf{w}_x \right) \\ (\text{by Equa.}(10)) &\geq \frac{\mathbf{m}_{[G]}(V_{\varepsilon}'[g_0])}{(1+\varepsilon)C_R} \left(\sum_{x \in g_0 \Gamma.o \cap B_{(1-\varepsilon)R}} \mathbf{w}_x \right) \\ \text{definition of } \Phi_R^{\mathbf{w}}) &= \frac{\mathbf{m}_{[G]}(V_{\varepsilon}'[g_0])}{1+\varepsilon} \frac{C_{(1-\varepsilon)R}}{C_R} \Phi_{(1-\varepsilon)R}^{\mathbf{w}}([g_0]). \end{split}$$

By taking the $\limsup_{R\to\infty}$ and making use of Equa.(13, 14), we get

$$\begin{split} & \operatorname{m}_{[G]}(V_{\varepsilon}'[g_0]) \limsup_{R \to \infty} \frac{C_{(1-\varepsilon)R}}{C_R} \Phi_{(1-\varepsilon)R}^{\mathbf{w}}([g_0]) \\ & \leq (1+\varepsilon) \limsup_{R \to \infty} \langle \Phi_R^{\mathbf{w}}, \mathbf{1}_{V_{\varepsilon}'[g_0]} \rangle_{\operatorname{m}_{[G]}} \leq (1+\varepsilon) \mu_{\infty}(V_{\varepsilon}[g_0]). \end{split}$$

Similarly we have

$$m_{[G]}(V_{\varepsilon}[g_0]) \liminf_{R \to \infty} \frac{C_{(1+\varepsilon)R}}{C_R} \Phi_{(1+\varepsilon)R}^{\mathbf{w}}([g_0]) \ge (1-\varepsilon)\mu_{\infty}(V_{\varepsilon}'[g_0]).$$

Combining these two while replacing $(1 - \varepsilon)R$ by R in the first inequality and $(1 + \varepsilon)R$ by R in the second inequality, we get

$$(1-\varepsilon)\liminf_{R\to\infty}\frac{C_R}{C_{R(1+\varepsilon)^{-1}}}\frac{\mu_{\infty}(V_{\varepsilon}'[g_0])}{\mathrm{m}_{[G]}(V_{\varepsilon}[g_0])}\leq\liminf_{R\to\infty}\Phi_R^{\mathbf{w}}([g_0])$$
$$\leq\limsup_{R\to\infty}\Phi_R^{\mathbf{w}}([g_0])\leq(1+\varepsilon)\limsup_{R\to\infty}\frac{C_{R(1-\varepsilon)^{-1}}}{C_R}\frac{\mu_{\infty}(V_{\varepsilon}[g_0])}{\mathrm{m}_{[G]}(V_{\varepsilon}'[g_0])}$$

By our assumptions, the $\limsup_{\varepsilon \to 0}$ of the left end coincide with the $\liminf_{\varepsilon \to 0}$ of the right end. Hence we must have

$$\lim_{R \to \infty} \Phi_R^{\mathbf{w}}([g_0]) = \lim_{\varepsilon \to 0} \frac{\mu_{\infty}(V_{\varepsilon}[g_0])}{\mathrm{m}_{[G]}(V_{\varepsilon}[g_0])}.$$

Call this limit $f([g_0])$. It is continuous because for $u \in \mathcal{O}_{\varepsilon}$,

$$f(u[g_0]) = \lim_{R \to \infty} \frac{1}{C_R} \sum_{x \in ug_0 \Gamma.o \cap B_R} \mathbf{w}_x = \lim_{R \to \infty} \frac{1}{C_R} \sum_{x \in g_0 \Gamma.o \cap u^{-1}B_R} \mathbf{w}_{ux}$$
$$\geq \lim_{R \to \infty} \frac{1}{C_R} \sum_{x \in g_0 \Gamma.o \cap B_{(1-\varepsilon)R}} \mathbf{w}_x (1+\varepsilon)^{-1} \geq (1+\varepsilon)^{-1} \liminf_{R \to \infty} \frac{C_{(1-\varepsilon)R}}{C_R} f([g_0]),$$

which converges to $f([g_0])$ as $\varepsilon \to 0$. A similar argument shows that the limit is no larger than $f([g_0])$. Thus $\lim_{u\to \mathrm{id}} f(u[g_0]) = f([g_0])$, proving the continuity.

By similar arguments, it is not hard to see that for any two g_1, g_0 in G, there exists a positive constant c_{g_1,g_0} such that $f([g_1g_0]) \ge c_{g_1,g_0}f([g_0])$. Therefore f is either constantly equal to 0 (which will be excluded below) or strictly positive.

3.2.2. Step 2, absolute continuity. Now we switch our attention to μ_{∞} and show that it is absolutely continuous with respect to $m_{[G]}$. That is to say, we need to show $\mu_{\infty}(E) = 0$ whenever $m_{[G]}(E) = 0$.

Recall that m_G is the Haar measure on G whose restriction to a fundamental set induces some constant multiple, which we may just assume to be 1 for simplicity, of $m_{[G]}$. By a Fubini-type argument, we first observe that if $m_{[G]}(E) = 0$, then for any probability measure λ on G/Γ and any bounded nonempty open set U in G, we have

$$\left(\int_{u\in U} u_*\lambda \operatorname{m}_G(u)\right)(E) = 0.$$

Now take a bounded measurable set E with $m_{[G]}(E) = 0$, there exists a family of shrinking bounded measurable sets $(E_i)_{i \in \mathbb{Z}^+}$ such that $E = \cap E_i$ and $\mu_{\infty}(\partial E_i) = 0$. For $u \in \mathcal{O}_{\varepsilon}$,

$$u_*\mu_{\infty}(E_i) = \lim_{R \to \infty} \frac{1}{C_R} \int_{[g] \in u \cdot B_R} g_* \mathrm{m}_{\mathrm{H}}(E_i) \mathbf{w}_{u^{-1}[g]} \mathrm{m}_{G/H}([g])$$

$$\leq (1-\varepsilon)^{-1} \lim_{R \to \infty} \frac{1}{C_R} \int_{[g] \in B_{(1+\varepsilon)R}} g_* \mathrm{m}_{\mathrm{H}}(E_i) \mathbf{w}_{[g]} \mathrm{m}_{G/H}([g])$$

$$\leq (1-\varepsilon)^{-1} \left(\limsup_{R \to \infty} \frac{C_{(1+\varepsilon)R}}{C_R}\right) \mu_{\infty}(E_i).$$

A lower bound for $u_*\mu_{\infty}(E_i)$ can be similarly obtained. Thus we have positive number ε' , which converges to 0 as ε does so, such that for every $u \in \mathcal{O}_{\varepsilon}$,

$$(1 - \varepsilon')\mu_{\infty}(E_i) \le u_*\mu_{\infty}(E_i) \le (1 + \varepsilon')\mu_{\infty}(E_i)$$

Hence

$$\left|\mu_{\infty}(E_{i}) - \frac{1}{\mathrm{m}_{G}(\mathcal{O}_{\varepsilon})} \int_{\mathcal{O}_{\varepsilon}} u_{*}\mu_{\infty}(E_{i}) \mathrm{m}_{G}(u)\right| \leq \varepsilon' \mu_{\infty}(E_{i}).$$

Letting i tend to infinity, we get

$$\left|\mu_{\infty}(E) - \frac{1}{\mathrm{m}_{G}(\mathcal{O}_{\varepsilon})} \int_{u \in \mathcal{O}_{\varepsilon}} u_{*}\mu_{\infty}(E)\mathrm{m}_{G}(u)\right| \leq \varepsilon' \mu_{\infty}(E).$$

However, by our observation, the measure of E with respect to "Haar average" of any probability measure is 0. So we are left with

$$\mu_{\infty}(E) \le \varepsilon' \mu_{\infty}(E_i)$$

for ε' arbitrarily small, which forces $\mu_{\infty}(E) = 0$. This ends the proof of absolute continuity.

3.2.3. Step 3, completing the proof. Write $\mu_{\infty} = \psi \cdot m_{[G]}$ for some non-negative measurable function ψ . By what has been shown in Step 1, for any $[g_0] \in G/\Gamma$,

$$f([g_0]) = \lim_{\varepsilon \to 0} \frac{\mu_{\infty}(V_{\varepsilon}[g_0])}{\mathrm{m}_{[G]}(V_{\varepsilon}[g_0])} = \lim_{\varepsilon \to 0} \frac{1}{\mathrm{m}_G(V_{\varepsilon})} \int_{u \in V_{\varepsilon}} \psi(u[g_0]) \,\mathrm{m}_G(u).$$

Therefore $f = \psi$ almost surely. As $\mu_{\infty} = \psi \cdot m_{[G]}$ is nonzero, we see that f can not be the 0 function. As mentioned towards the end of step 1, this implies that f is strictly positive.

4. Equidistribution and Nondivergence

In this section we collect some results on equidistribution and nondivergence of homogeneous measures on a special type of homogeneous spaces: arithmetic quotients of real points of linear algebraic groups over \mathbb{Q} . The structure of linear algebraic groups over \mathbb{Q} and reduction theory for arithmetic subgroups are well-understood (see [BHC62, Spr98, Bor19]). For instance, arithmetic quotients have finite volume iff the algebraic group has no non-torsion \mathbb{Q} -characters. The study of dynamics of subgroup action on such homogeneous spaces comes later. The full classification of unipotent-invariant ergodic measures is obtained by Ratner [Ra91] with shorter proof given by [MT94]. The analysis of ergodic components of a unipotent-invariant measure is also possible by the linearization method developed in [DM91]. In addition to these works, the results presented below (mainly taken from [EMS96, Zha21, Zha22]) also rely on the theory of (C, α) -functions of [KM98] (see [EMS97] for a slightly different approach).

4.1. Notations. Throughout this section we adopt the following notations:

- For a linear algebraic group **G** over \mathbb{R} , write $G := \mathbf{G}(\mathbb{R})^{\circ}$. When it is understood that there is some ambient algebraic group **G** over \mathbb{R} , let $L := \mathbf{L}(\mathbb{R}) \cap \mathbf{G}$ for an \mathbb{R} -subgroup **L**. If **H** is another \mathbb{R} -subgroup contained in **L**, let $L^{\mathrm{H}} := L^{\circ} \cdot \mathrm{H}$.
- Let $\Gamma \subset G \cap \mathbf{G}(\mathbb{Q})$ be an arithmetic lattice;
- For an observable Q-subgroup L of G, let $m_{[L]}$ denote a locally finite L-invariant measure supported on $L\Gamma/\Gamma$. For other $L^{\circ} \subset L \subset L$, we similarly define $m_{[L]}$, supported on $L\Gamma/\Gamma$.
- For an algebraic subgroup L of G, let $Z_G(L)$ be the centralizer of L in G.

We fix an integral structure (that is, a lattice) $\mathfrak{g}_{\mathbb{Z}}$ on \mathfrak{g} and hence on its exterior powers. Whenever an arithmetic lattice is fixed, we further require that $\mathfrak{g}_{\mathbb{Z}}$ is preserved by $\operatorname{Ad}(\Gamma)$. Also, Euclidean metrics on these spaces are fixed. For every parabolic \mathbb{Q} -subgroup \mathbf{P} , take $\mathbf{v}_{\mathbf{P}}$ to be the unique (up to ± 1) primitive integral vector in $\wedge^{\dim \mathbf{P}} \mathfrak{g}_{\mathbb{Z}}$ that lifts its Lie algebra. Also define

 $\mathscr{P}_{\mathbf{H}} := \left\{ \mathrm{proper \ parabolic} \ \mathbb{Q} \mathrm{-subgroups} \ \mathrm{of} \ \mathbf{G} \ \mathrm{containing} \ \mathbf{H} \right\}.$

 $\mathscr{P}_{\mathbf{H}}^{\max} := \{ \mathrm{maximal \ proper \ parabolic} \ \mathbb{Q} \mathrm{-subgroups \ of} \ \mathbf{G} \ \mathrm{containing} \ \mathbf{H} \} \, .$

Recall that a subgroup \mathbf{L} of a linear algebraic group \mathbf{G} is said to be *observable* iff there exists a finite-dimensional representation of \mathbf{G} and a vector whose stabilizer subgroup is equal to \mathbf{L} . If \mathbf{L} is assumed to be a Q-subgroup, then $L\Gamma$ is closed iff \mathbf{L} is observable in \mathbf{G} ([Wei98, Corollary 7]). Reductive groups and unipotent groups are automatically observable in any ambient group. If one is only interested in the case when $m_{[H]}$ is finite, which implies the observability, then one can ignore the word observable or replace it by having finite invariant measure on the quotient by an arithmetic subgroup.

Related to observable subgroups is the notion of *epimorphic subgroups*. A subgroup \mathbf{L} of \mathbf{G} is said to be epimorphic iff the smallest observable subgroup containing \mathbf{L} is \mathbf{G} . Alternatively, for every v in a finite-dimensional representation of \mathbf{G} , v fixed by \mathbf{L} implies that v is fixed by \mathbf{G} .

4.2. Subgroup convergence and equidistribution.

Definition 4.1. Let \mathbf{L} be a connected linear algebraic group over \mathbb{Q} and (\mathbf{M}_n) be a sequence of connected observable \mathbb{Q} -subgroups of \mathbf{L} , we say that (\mathbf{M}_n) converges to \mathbf{L} iff there is no proper connected observable subgroup of \mathbf{L} that contains \mathbf{M}_n for infinitely many n's.

Note that (\mathbf{M}_n) always converges after passing to a subsequence. It is not true that if (\mathbf{M}_n) converges to **L**, then the associated homogeneous measures also converge. Below is a special case when this does happen.

Theorem 4.2. Let **H** and **L** be two observable connected Q-subgroups of a connected linear algebraic group **G** over Q. Let (γ_n) be a sequence in Γ such that $(\gamma_n \mathbf{H} \gamma_n^{-1})$ converges to **L**. Then there exists a sequence of positive numbers (a_n) such that $\lim_{n\to\infty} \frac{1}{a_n} (\gamma_n)_* \mathbf{m}_{[\mathbf{H}^\circ]} = \mathbf{m}_{[\mathbf{L}^\circ]}$ under the weak-* topology.

This is proved in [Zha21, Theorem 1.3] building on previous work of [EMS96] (see also [OS14, SZ19, Zha19]).

The asymptotic of (a_n) is unique. Let ψ be a compactly supported non-negative function on G/Γ with $\langle \psi, m_{[L^\circ]} \rangle \neq 0$, the conclusion of last theorem unwraps to the following:

$$\lim_{n \to \infty} \frac{\langle f, (\gamma_n)_* \mathbf{m}_{[\mathrm{H}^\circ]} \rangle}{\langle \psi, (\gamma_n)_* \mathbf{m}_{[\mathrm{H}^\circ]} \rangle} = \frac{\langle f, \mathbf{m}_{[\mathrm{L}^\circ]} \rangle}{\langle \psi, \mathbf{m}_{[\mathrm{L}^\circ]} \rangle}.$$

4.2.1. Definition of p^{sta} . For a connected linear algebraic group \mathbf{G} , let $\mathbf{G}^{\text{ari}} := \left(\overline{\overline{\Gamma}}\right)^{\circ}$, the identity component of the Zariski closure of Γ . Let p^{sta} be the natural quotient map $\mathbf{G} \to \mathbf{G}/\mathbf{G}^{\text{ari}}$. Actually \mathbf{G}^{ari} is independent of the choice of the arithmetic subgroup Γ .

More explicitly, recall

$$\mathbf{G} = (\mathbf{G}^{\mathrm{ss}} \cdot \mathbf{Z}(\mathbf{G})) \ltimes \mathbf{R}_{\mathbf{u}}(\mathbf{G}) = ((\mathbf{G}^{\mathrm{cpt}} \cdot \mathbf{G}^{\mathrm{nc}}) \cdot \mathbf{Z}(\mathbf{G})) \ltimes \mathbf{R}_{\mathbf{u}}(\mathbf{G}).$$

Also, $\mathbf{Z}(\mathbf{G})$ is an almost direct product $\mathbf{Z}(\mathbf{G})^{\mathrm{cpt}} \cdot \mathbf{Z}(\mathbf{G})^{\mathrm{spl}} \cdot \mathbf{Z}(\mathbf{G})^{\mathrm{ari}}$ where $\mathbf{Z}(\mathbf{G})^{\mathrm{spl}}$ is a \mathbb{Q} -split torus, $\mathbf{Z}(\mathbf{G})^{\mathrm{cpt}}$ is an \mathbb{R} -anisotropic \mathbb{Q} -torus and $\mathbf{Z}(\mathbf{G})^{\mathrm{ari}}$ is the identity component of the Zariski closure of some/any arithmetic lattice of $\mathbf{Z}(\mathbf{G})$. In terms of the character groups, these tori can be described as follows:

Let $\Lambda_0 \leq \mathfrak{X}^*(\mathbf{Z}(\mathbf{G}))$ be the primitive subgroup fixed by $\operatorname{Gal}_{\mathbb{Q}}$. It admits a unique $\operatorname{Gal}_{\mathbb{Q}}$ -stable complementary primitive subgroup Λ_1 (in the sense that $\Lambda_0 \cap \Lambda_1 = \{0\}$ and $\Lambda_0 + \Lambda_1$ is of finite-index in $\mathfrak{X}^*(\mathbf{Z}(\mathbf{G}))$). Then $\mathbf{Z}(\mathbf{G})^{\operatorname{spl}}$ is the common kernel of characters in Λ_1 . Similarly, let Λ_2 be the primitive subgroup of $\mathfrak{X}^*(\mathbf{Z}(\mathbf{G}))$ fixed by the smaller subgroup $\operatorname{Gal}_{\overline{\mathbb{Q}}\cap\mathbb{R}}$ and Λ_3 be the $\operatorname{Gal}_{\overline{\mathbb{Q}}\cap\mathbb{R}}$ -stable complement of Λ_2 in $\mathfrak{X}^*(\mathbf{Z}(\mathbf{G}))$. As $\operatorname{Gal}_{\overline{\mathbb{Q}}\cap\mathbb{R}}$ is normal in $\operatorname{Gal}_{\mathbb{Q}}$, Λ_2 and Λ_3 are $\operatorname{Gal}_{\mathbb{Q}}$ -stable. Then $\mathbf{Z}(\mathbf{G})^{\operatorname{cpt}}$ corresponds to Λ_2 and $\mathbf{Z}^{\operatorname{ari}}$ corresponds to the primitive subgroup containing $\Lambda_0 + \Lambda_3$.

With these notations, \mathbf{G}^{ari} is the normal subgroup $(\mathbf{G}^{\text{nc}} \cdot \mathbf{Z}(\mathbf{G})^{\text{ari}}) \ltimes \mathbf{R}_{\mathbf{u}}(\mathbf{G})$ by Borel density lemma.

4.2.2. Criteria on convergence to the full group.

Lemma 4.3. Let **G** be a connected linear algebraic group over \mathbb{Q} , **N** be a connected normal \mathbb{Q} -subgroup of **G** and **H** be an observable \mathbb{Q} -subgroup contained in **N**. Let Γ be an arithmetic lattice of **G**. The followings are equivalent:

- (1) There exists $(g_n) \subset G$ such that $[(g_n)_* m_{[H^\circ]}] \rightarrow [m_{[N^\circ]}];$
- (2) There exists $(\gamma_n) \subset \Gamma$ such that $[(\gamma_n)_* \mathbf{m}_{[\mathrm{H}^\circ]}] \to [\mathbf{m}_{[\mathrm{N}^\circ]}];$
- (3) $p^{\text{sta}}(\mathbf{H}) = p^{\text{sta}}(\mathbf{N})$ and any proper \mathbb{Q} -subgroup that is normalized by $\mathbf{N} \cdot \mathbf{G}^{\text{ari}}$ and contains \mathbf{H} must also contain \mathbf{N} .

Proof. Part (1) and (2) are easily seen to be equivalent. That (2) implies (3) is quite direct. Let us explain why (3) implies (2).

By [Zha21, Lemma 4.14], if (2) were not true, then we can find an observable Q-subgroup L that is properly contained in N, contains H and is normalized by some arithmetic subgroup Γ' . Hence L is normalized by \mathbf{G}^{ari} . But $p^{\text{sta}}(\mathbf{H}) = p^{\text{sta}}(\mathbf{N})$, thus L is normalized by \mathbf{NG}^{ari} , which can not be true by (3).

In the special case of $\mathbf{N} = \mathbf{G}$ we get

Lemma 4.4. Let **G** be a connected linear algebraic group over \mathbb{Q} and **H** be an observable \mathbb{Q} -subgroup. Let Γ be an arithmetic lattice. The followings are equivalent:

- (1) There exists $(g_n) \subset \mathbf{G}$ such that $[(g_n)_*\mathbf{m}_{[\mathbf{H}^\circ]}] \to [\mathbf{m}_{[\mathbf{G}]}];$
- (2) There exists $(\gamma_n) \subset \Gamma$ such that $[(\gamma_n)_* \mathbf{m}_{[\mathrm{H}^\circ]}] \to [\mathbf{m}_{[\mathrm{G}]}];$
- (3) p^{sta} restricted to **H** is surjective and **H** is not contained in any proper normal \mathbb{Q} -subgroup of **G**.

4.3. Non-divergence criteria.

4.3.1. Non-divergence of homogeneous closed subsets. Recall that a Q-group L is said to be Q-anisotropic iff the set of Q-cocharacters of L is finite, which is equivalent to $\mathbf{L}(\mathbb{R})/\Gamma$ being compact for some/any arithmetic subgroup Γ of L. The following is from [Zha21, Theorem 1.7].

Theorem 4.5. Assume that **H** and **G** are reductive linear algebraic groups over \mathbb{Q} and Γ is an arithmetic subgroup of **G** contained in G. If $\mathbf{Z}_{\mathbf{G}}(\mathbf{H}^{\circ})/\mathbf{Z}(\mathbf{H})$ is \mathbb{Q} -anisotropic, then there exists a bounded subset $\mathcal{B}_{\Gamma} \subset \mathbf{G}$ such that

$$\mathbf{G} = \mathcal{B}_{\Gamma} \cdot \Gamma \cdot \mathbf{H}^{\circ}.$$

Moreover, if \mathbf{L} is an observable \mathbb{Q} -subgroup containing \mathbf{H} , then \mathbf{L} is reductive.

Here is a more general version.

Theorem 4.6. Let \mathbf{G} be a connected linear algebraic group over \mathbb{Q} , \mathbf{H} be a connected observable \mathbb{Q} subgroup and Γ be an arithmetic subgroup of \mathbf{G} contained in \mathbf{G} . Fix a maximal reductive subgroup $\mathbf{G}^{\mathrm{red}}$ of \mathbf{G} and write $\mathbf{G} = \mathbf{G}^{\mathrm{red}} \ltimes \mathbf{R}_{\mathbf{u}}(\mathbf{G})$. Also fix a Cartan involution and hence a maximal compact
subgroup \mathbf{K} of $\mathbf{G}^{\mathrm{red}}(\mathbb{R})$. Given a sequence $(g_n) \subset \mathbf{G}$, after passing to a subsequence, there exist a
sequence $(h_n) \subset \mathbf{H}^\circ$, $(\gamma_n) \subset \Gamma$ and a parabolic \mathbb{Q} -subgroup \mathbf{P} such that the following holds. Write $g_n h_n \gamma_n^{-1} = k_n a_n p_n$ using horospherical coordinates of (\mathbf{P}, \mathbf{K}) . Then

- (1) (p_n) is bounded;
- (2) $\alpha(a_n) \to 0$ for every $\alpha \in \Delta^{\text{red}}(\mathbf{A}_{\mathbf{P},\mathrm{K}},\mathbf{P})$;
- (3) if (a_n) is unbounded, there exist a \mathbb{Q} -representation \mathbf{V} of \mathbf{G} factoring through $\mathbf{G}/\mathbf{R}_{\mathbf{u}}(\mathbf{G})$ and $\mathbf{v} \in \mathbf{V}(\mathbb{Q})$ such that the line spanned by \mathbf{v} is preserved by \mathbf{P} , \mathbf{v} is fixed by $\gamma_n \mathbf{H} \gamma_n^{-1}$ for all n and $\lim_{n\to\infty} a_n \cdot \mathbf{v} = \mathbf{0}$;
- (4) $(\gamma_n \mathbf{H} \gamma_n^{-1})$ strongly converges to some observable subgroup of **G**.

The theorem is restated and proved in the appendix (see Theorem B.1).

Corollary 4.7. Let $\mathbf{G}, \mathbf{H}, \Gamma$ be the same as in the last theorem. Let (g_n) be a sequence in \mathbf{G} such that $(g_n \mathrm{H}^\circ \Gamma/\Gamma)$ diverges topologically, then there exists a \mathbb{Q} -representation $\rho : \mathbf{G} \to \mathbf{SL}(\mathbf{V})$, an integral structure $\mathbf{V}(\mathbb{Z})$ on \mathbf{V} and a sequence of non-zero vectors $(v_n) \subset \mathbf{V}(\mathbb{Z})$ that are $\rho(\mathbf{H})$ -fixed such that $\rho(g_n).v_n \to \mathbf{0}$.

4.3.2. Non-divergence of a bounded piece. Let **G** be a connected linear algebraic group over \mathbb{Q} and **H** be a \mathbb{Q} -subgroup. For a non-empty open bounded subset \mathcal{O}_{H} of $H^{\circ}\Gamma/\Gamma$, let $m_{\mathcal{O}_{H}}$ be the restriction of $m_{[H]}$ to \mathcal{O}_{H} and let $m_{\mathcal{O}_{H}}^{1}$ be the unique probability measure proportional to $m_{\mathcal{O}_{H}}$.

Theorem 4.8. Fix a non-empty open bounded subset \mathcal{O}_{H} of $H^{\circ}\Gamma/\Gamma$. Then

(1) for every bounded set $\mathcal{B} \subset G/\Gamma$, there exists $\eta_{\mathcal{B}} > 0$ such that for every $g \in {}^{\circ}G$,

 $g\mathcal{O}_{\mathrm{H}} \cap \mathcal{B} \neq \emptyset \implies \|\mathrm{Ad}(g)\mathbf{v}_{\mathbf{P}}\| > \eta_{\mathcal{B}}, \ \forall \mathbf{P} \in \mathscr{P}_{\mathbf{H}^{\circ}}^{\mathrm{max}}.$

(2) for every $\eta > 0$, there exists a bounded set $\mathcal{B}_{\eta} \subset G/\Gamma$ such that for every $g \in {}^{\circ}G$,

 $\|\mathrm{Ad}(g)\mathbf{v}_{\mathbf{P}}\| > \eta, \ \forall \, \mathbf{P} \in \mathscr{P}_{\mathbf{H}^{\circ}}^{\mathrm{max}} \implies g\mathcal{O}_{\mathrm{H}} \cap \mathcal{B}_{\eta} \neq \emptyset.$

For every $\varepsilon \in (0,1)$, by enlarging \mathcal{B}_{η} if possible, one can arrange that for every $g \in {}^{\circ}G$,

$$\|\operatorname{Ad}(g)\mathbf{v}_{\mathbf{P}}\| > \eta, \ \forall \mathbf{P} \in \mathscr{P}_{\mathbf{H}^{\circ}}^{\max} \implies (g_* \mathbf{m}_{\mathcal{O}_{\mathbf{H}}}^1)(\mathcal{B}_{\eta}) > 1 - \varepsilon.$$

Part (1) follows directly. It remains to prove part (2). Due to the (C, α) -good property, it suffices to prove the first claim of part (2). The special case when **G** is semisimple is proved in [Zha22, Theorem 1.3]. In order to reduce the general case to this, we need a little preparation.

21

4.3.3. Parabolic subgroups and radicals. Let r_1 denote the dimension of $\mathbf{R}(\mathbf{G})$, the radical of \mathbf{G} . Let $p^{ss} : \mathbf{G} \to \mathbf{G}/\mathbf{R}(\mathbf{G})$ be the natural quotient morphism. For a non-negative integer l, let $\wedge^l dp^{ss}$ be the induced morphism between exterior powers of Lie algebras $\wedge^l \mathfrak{g} \to \wedge^l \mathfrak{g}/\mathfrak{r}(\mathfrak{g})$. Fix an integral vector $\mathbf{v}_{\mathfrak{r}} \in \wedge^r \mathfrak{g}$ that lifts the Lie algebra $\mathfrak{r}(\mathfrak{g})$ of $\mathbf{R}(\mathbf{G})$.

For every $\omega \in \wedge^l \mathfrak{g}/\mathfrak{r}(\mathfrak{g})$, choose some $\widetilde{\omega} \in \wedge^l \mathfrak{g}$ with $\wedge^l dp^{ss}(\widetilde{\omega}) = \omega$. Define a map φ from $\wedge^l \mathfrak{g}/\mathfrak{r}(\mathfrak{g})$ to $\wedge^{l+r}\mathfrak{g}$ by

$$\omega \mapsto \varphi(\omega) := \widetilde{\omega} \wedge \mathbf{v}_{\mathfrak{r}}$$

One can check that $\varphi(\omega)$ is independent of the choice of $\tilde{\omega}$ and φ is a linear injection. Thus, there exists C > 1 such that

$$C^{-1} \left\| \varphi(\omega) \right\| \le \left\| \omega \right\| \le C \left\| \varphi(\omega) \right\|, \quad \forall \, \omega \in \wedge^l \mathfrak{g}/\mathfrak{r}(\mathfrak{g}).$$

Moreover, φ is °G-equivariant since °G fixes the vector $\mathbf{v}_{\mathfrak{r}}.$

In addition to this, we need another simple fact:

Lemma 4.9. Every parabolic subgroup of **G** contains $\mathbf{R}(\mathbf{G})$. Consequently, the map $\mathbf{P} \mapsto \mathbf{P}/\mathbf{R}(\mathbf{G})$ defines a bijection between parabolic \mathbb{Q} -subgroups of **G** and those of $\mathbf{G}/\mathbf{R}(\mathbf{G})$. It also induces a bijection between $\mathscr{P}_{\mathbf{H}^{\circ}}^{\max}$ and $\mathscr{P}_{p^{\mathrm{ss}}(\mathbf{H}^{\circ})}^{\max}$.

By [Spr98, Theorem 6.2.7], a parabolic subgroup contains some Borel subgroup and hence $\mathbf{R}(\mathbf{G})$. The rest of the claim follows from this.

Therefore, we find some $C_1 > 1$ such that for every $\mathbf{P} \in \mathscr{P}_{p^{\mathrm{ss}}(\mathbf{H}^\circ)}^{\mathrm{max}}$ and every $g \in {}^{\circ}\mathbf{G}$,

$$C_1^{-1} \|\operatorname{Ad}(g)\mathbf{v}_{\mathbf{Q}}\| \le \|\operatorname{Ad}(p^{\operatorname{ss}}(g))\mathbf{v}_{\mathbf{P}}\| \le C_1 \|\operatorname{Ad}(g)\mathbf{v}_{\mathbf{Q}}\|$$
(15)

where $\mathbf{Q} := (p^{\mathrm{ss}})^{-1}(\mathbf{P})$ is an element in $\mathscr{P}_{\mathbf{H}^{\circ}}^{\mathrm{max}}$. Note that $\mathbf{v}_{\mathbf{Q}} = c_{\mathbf{P}}\varphi(\mathbf{v}_{\mathbf{P}})$ for some positive constant $c_{\mathbf{P}}$ bounded away from 0 and $+\infty$ as \mathbf{P} varies.

4.3.4. Proof of Theorem 4.8. For simplicity, write $\mathbf{G}' := \mathbf{G}/\mathbf{R}(\mathbf{G})$ and $\mathbf{G}' := \mathbf{G}'(\mathbb{R})^{\circ}$.

The morphism p^{ss} induces $\overline{p}^{ss} : G/\Gamma \to G'/p^{ss}(\Gamma)$. By [Zha22, Theorem 1.3], there exists a bounded subset $\mathcal{B}'_{\eta} \subset G'/p^{ss}(\Gamma)$ such that for every $g \in G'$,

$$\|\operatorname{Ad}(g)\mathbf{v}_{\mathbf{P}}\| > \eta, \ \forall \, \mathbf{P} \in \mathscr{P}_{p^{\operatorname{ss}}(\mathbf{H}^{\circ})}^{\max} \implies g\bar{p}^{\operatorname{ss}}(\mathcal{O}_{\mathrm{H}}) \cap \mathcal{B}'_{\eta} \neq \emptyset.$$
(16)

On the other hand, take a bounded open subset $\mathcal{B}^{\text{spl}} \subset \mathbf{S}_{\mathbf{G}}(\mathbb{R})^{\circ}$ containing $\overline{p}^{\text{spl}}(\mathcal{O}_{\mathrm{H}})$. Let $\overline{p}_{\mathcal{B}}^{\text{ss}}$ be the restriction of \overline{p}^{ss} to the closure of $(\overline{p}^{\text{spl}})^{-1}(\mathcal{B}^{\text{spl}})$. Then $\overline{p}_{\mathcal{B}}^{\text{ss}}$ is a proper map. Define a bounded subset of G/Γ by

$$\mathcal{B}_{\eta} := (\overline{p}^{\mathrm{ss}})^{-1} \left(\mathcal{B}'_{C_{1}^{-1}\eta} \right) \bigcap (\overline{p}^{\mathrm{spl}})^{-1} \left(\mathcal{B}^{\mathrm{spl}} \right).$$

Now fix some $g \in {}^{\circ}G$ such that $\|\operatorname{Ad}(g)\mathbf{v}_{\mathbf{P}}\| > \eta$ for every $\mathbf{P} \in \mathscr{P}_{\mathbf{H}^{\circ}}^{\max}$. Since $\overline{p}^{\operatorname{spl}}(g\mathcal{O}_{\mathrm{H}}) = \overline{p}^{\operatorname{spl}}(\mathcal{O}_{\mathrm{H}})$, the containment $g\mathcal{O}_{\mathrm{H}} \subset (\overline{p}^{\operatorname{spl}})^{-1}(\mathcal{B}^{\operatorname{spl}})$ is assured. For the other part, by Equa.(15),

 $\|\mathrm{Ad}(p^{\mathrm{ss}}(g))\mathbf{v}_{\mathbf{P}}\| > C_1^{-1}\eta, \quad \forall \, \mathbf{P} \in \mathscr{P}_{p^{\mathrm{ss}}(\mathbf{H}^\circ)}^{\max}.$

Hence we are done by Equa.(16).

4.4. Refined versions. In this subsection, **G** is a connected linear algebraic group over \mathbb{Q} and **H** is a \mathbb{Q} -subgroup. Let $\mathbf{H}^1 := {}^{\circ}\mathbf{G} \cap \mathbf{H}$.

4.4.1. Explicit form of (a_n) . Assume the natural injective morphism $\mathbf{H}/\mathbf{H}^1 \to \mathbf{S}_{\mathbf{G}}$ is an isomorphism. We choose a lift $\widetilde{\mathbf{S}}^{\mathbf{H}}$, a \mathbb{Q} -split torus, of $\mathbf{S}_{\mathbf{G}}$ in \mathbf{H} . Then the natural product map

$$\widetilde{\mathbf{S}}^{\mathbf{H}}(\mathbb{R})^{\circ} \times \mathbf{H}^{\mathbf{1}}(\mathbb{R})^{\circ} \to \mathbf{H}(\mathbb{R})^{\circ}$$

is a homeomorphism. From this we deduce that the natural map

$$\widetilde{\mathbf{S}}^{\mathbf{H}}(\mathbb{R})^{\circ} \times \mathbf{H}^{\mathbf{1}}(\mathbb{R})^{\circ} / \mathbf{H}^{\mathbf{1}}(\mathbb{R})^{\circ} \cap \Gamma \to \mathbf{H}(\mathbb{R})^{\circ} / \mathbf{H}(\mathbb{R})^{\circ} \cap \Gamma$$
(17)

is a homeomorphism.

Based on this, for $g \in {}^{\circ}\mathbf{G}$ (which, one can verify, is connected) and $\eta > 0$, define

$$\Omega^{\mathbf{1}}_{\mathbf{H},g,\eta} := \left\{ [h] \in \mathbf{H}^{\mathbf{1}}(\mathbb{R})^{\circ} / \mathbf{H}^{\mathbf{1}}(\mathbb{R})^{\circ} \cap \Gamma \mid \|\mathrm{Ad}(gh)\mathbf{v}_{\mathbf{P}}\| > \eta, \ \forall \, \mathbf{P} \in \mathscr{P}_{\mathbf{H}^{\circ}}^{\max} \right\}.$$
(18)

Theorem 4.10. Assume additionally that **H** is observable. Let (g_n) be a sequence in °G such that $\lim_{n\to\infty} [(g_n)_* \mathfrak{m}_{[\mathbf{H}^\circ]}] = [\mathfrak{m}_{[\mathbf{G}]}]$. Then for every $\eta > 0$,

$$\lim_{n \to \infty} \frac{1}{\mathbf{m}_{[(\mathbf{H}^1)^\circ]} \left(\Omega^{\mathbf{1}}_{\mathbf{H},g_n,\eta}\right)} (g_n)_* \mathbf{m}_{[\mathbf{H}^\circ]} = \mathbf{m}_{[\mathbf{G}]}.$$
 (19)

In Equa.(19), the Haar measures are chosen as follows: Choose some Haar measure m_S on $\widetilde{\mathbf{S}}^{\mathbf{H}}(\mathbb{R})^{\circ}$ and $m_{[(\mathbf{H}^1)^{\circ}]}$ on $\mathbf{H}^1(\mathbb{R})^{\circ}/\mathbf{H}^1(\mathbb{R})^{\circ} \cap \Gamma$. By Equa.(17), define $m_{[\mathbf{H}^{\circ}]} := m_S \otimes m_{[(\mathbf{H}^1)^{\circ}]}$. Likewise, one has

$$\widetilde{\mathbf{S}}^{\mathbf{H}}(\mathbb{R})^{\circ} \times {}^{\circ}\mathrm{G}/\Gamma \cong \mathrm{G}/\Gamma.$$

And define $m_{[G]} := m_S \otimes m_{[\circ G]}^1$. The truth of Equa.(19) is independent of the choice of m_S and $m_{[(H^1)^\circ]}$.

4.4.2. *Two lemmas.* The proof of Theorem 4.10 is essentially contained in [Zha21] by using Theorem 4.8. We outline a proof here for the sake of completeness. Theorem 4.10 can be directly deduced from the following two lemmas:

Lemma 4.11. Assumption as in Theorem 4.10. There exists (η_n) diverging to $+\infty$ such that for any $\eta > 0$,

$$\lim_{n \to \infty} \frac{\mathrm{m}_{[(\mathrm{H}^{1})^{\circ}]} \left(\Omega^{1}_{\mathbf{H}, g_{n}, \eta_{n}} \right)}{\mathrm{m}_{[(\mathrm{H}^{1})^{\circ}]} \left(\Omega^{1}_{\mathbf{H}, g_{n}, \eta} \right)} = 1.$$

Lemma 4.12. Assumption as in Theorem 4.10. Let (η_n) be a sequence of real numbers diverging to $+\infty$. Then for every $(h_n) \subset \Omega^1_{\mathbf{H},q_n,\eta_n}$ and non-empty open bounded subset $\mathcal{O}_{\mathbf{H}} \subset \mathbf{H}^\circ \Gamma / \Gamma$,

$$\lim_{n \to \infty} (g_n h_n)_* \mathbf{m}_{\mathcal{O}_{\mathrm{H}}}^{\mathbf{1}} = \mathbf{m}_{\mathcal{O}_{\mathrm{H}}}^{\mathbf{1}}$$

where

$$\mathcal{O}_{\mathrm{H}}^{\mathrm{spl}} := (\overline{p}^{\mathrm{spl}})^{-1} \left(\overline{p}^{\mathrm{spl}}(\mathcal{O}_{\mathrm{H}}) \right).$$

In the above, $m_{\mathcal{O}_H}$ (resp. $m_{\mathcal{O}_H^{spl}}$) denotes the restriction of $m_{[H^\circ]}$ (resp. $m_{[G]}$) to \mathcal{O}_H (resp. \mathcal{O}_H^{spl}). The upper index $(\cdot)^1$ is used to denote the unique probability measure proportional to (\cdot) .

4.4.3. Proof of Lemma 4.12. By Theorem 4.8, $g_n h_n \omega_n = \delta_n \gamma_n$ for some bounded sequence $(\omega_n) \subset \mathcal{H}$ lifting elements in $\mathcal{O}_{\mathcal{H}}$, some bounded sequence (δ_n) in G and $(\gamma_n) \subset \Gamma$. Passing to a subsequence, may assume (ω_n) converges to ω_{∞} . Thus it is sufficient to establish the claim with g_n replaced by γ_n and $\mathcal{O}_{\mathcal{H}}$ replaced by $\omega_{\infty}^{-1}\mathcal{O}_{\mathcal{H}} =: \mathcal{O}'_{\mathcal{H}}$.

By the work of [EMS96] (see [Zha21, Section 4] for details), it is sufficient to show that $(\gamma_n \mathbf{H}^{\circ} \gamma_n^{-1})$ is not contained in any proper \mathbb{Q} -subgroup of \mathbf{G} for infinitely many *n*'s. So assume otherwise and we seek for a contradiction. Let \mathbf{L} be such a \mathbb{Q} -subgroup. If \mathbf{L} is observable, then this would violate the assumption that $\lim [(g_n)_* \mathbf{m}_{[\mathbf{H}^{\circ}]}]] = [\mathbf{m}_{[\mathbf{G}]}]$. So let us assume that \mathbf{L} is epimorphic in \mathbf{G} . By Lemma 4.13 below, we assume that $\mathbf{L} = \mathbf{P}$ is a maximal proper parabolic \mathbb{Q} -subgroup.

Find an infinite subsequence (n_k) such that $\gamma_{n_k} \mathbf{H}^{\circ} \gamma_{n_k}^{-1} \subset \mathbf{P}$. Let $\mathbf{P}_k := \gamma_{n_k}^{-1} \mathbf{P} \gamma_{n_k}$, then by assumption

$$\|\operatorname{Ad}(g_{n_k}h_{n_k})\mathbf{v}_{\mathbf{P}_k}\| \to +\infty.$$

On the other hand, there exists C > 1 such that

$$C^{-1} \left\| \operatorname{Ad}(\gamma_{n_k}) \mathbf{v}_{\mathbf{P}_k} \right\| \le \left\| \operatorname{Ad}(g_{n_k} h_{n_k}) \mathbf{v}_{\mathbf{P}_k} \right\| \le C \left\| \operatorname{Ad}(\gamma_{n_k}) \mathbf{v}_{\mathbf{P}_k} \right\|.$$

Therefore, $\|\operatorname{Ad}(\gamma_{n_k})\mathbf{v}_{\mathbf{P}_k}\| \to +\infty$. On the other hand, $\mathbf{P} = \gamma_{n_k}\mathbf{P}_k\gamma_{n_k}^{-1}$, thus for some C > 1,

$$C^{-1} \|\mathbf{v}_{\mathbf{P}}\| \le \|\operatorname{Ad}(\gamma_{n_k})\mathbf{v}_{\mathbf{P}_k}\| \le C \|\mathbf{v}_{\mathbf{P}}\|,$$

which are bounded and unbounded at the same time, leading to a contradiction.

Lemma 4.13. Let \mathbf{A} be a linear algebraic group over \mathbb{Q} . Every proper epimorphic \mathbb{Q} -subgroup of \mathbf{A} is contained in a proper parabolic \mathbb{Q} -subgroup.

Proof. Let \mathbf{C} be epimorphic in \mathbf{A} , we need to show that \mathbf{C} is contained in some proper parabolic \mathbb{Q} -subgroup.

Let U be the non-trivial unipotent radical of C and let $\mathbf{v}_{\mathbf{U}} \in \wedge^{\dim \mathbf{U}} \mathfrak{a}$ be a vector lifting the Lie algebra of U.

When **A** is reductive, $\mathbf{v}_{\mathbf{U}}$ is an unstable **A**-vector. Thus, the stabilizer of the line spanned by $\mathbf{v}_{\mathbf{U}}$ is contained in a parabolic Q-subgroup **P** of **A** (see [Kem78, Corollary 5.1]). In particular, **C** is contained in **P**.

In general, consider the morphism $p^{\text{red}} : \mathbf{A} \to \mathbf{A}/\mathbf{R}_{\mathbf{u}}(\mathbf{A})$. Then $p^{\text{red}}(\mathbf{C})$ is epimorphic in $\mathbf{A}/\mathbf{R}_{\mathbf{u}}(\mathbf{A})$ and hence is either contained in a proper maximal parabolic \mathbb{Q} -subgroup of $\mathbf{A}/\mathbf{R}_{\mathbf{u}}(\mathbf{A})$ or equal the full $\mathbf{A}/\mathbf{R}_{\mathbf{u}}(\mathbf{A})$. In the former case, we are done by taking the preimage. In the latter case, we see that \mathbf{U} is contained in $\mathbf{R}_{\mathbf{u}}(\mathbf{A})$. Hence \mathbf{A}/\mathbf{C} is affine by [Gro97, Theorem 7.1]. Thus \mathbf{C} is observable, a contradiction.

4.4.5. Proof of Lemma 4.11. Thanks to the properties of convex polytopes (See [SZ19, Section 4]), it suffices to show that there exists (η_n) diverging to $+\infty$ such that $\Omega^1_{\mathbf{H},g_n,\eta_n} \neq \emptyset$.

By assumption, there exist $(h_n) \subset \mathrm{H}^\circ$, $(b_n) \subset \mathrm{G}$ bounded and $(\gamma_n) \subset \Gamma$ such that

 $g_n h_n = b_n \gamma_n$ for all n.

Since $p^{\operatorname{spl}}(h_n) = p^{\operatorname{spl}}(g_n^{-1}b_n\gamma_n) = p^{\operatorname{spl}}(b_n)$ remains bounded, we can find $(\omega_n) \subset \operatorname{H}^\circ$ bounded and $(h'_n) \subset (\operatorname{H}^1)^\circ$ such that $h_n = h'_n\omega_n$. Therefore, it suffices to find diverging (η_n) such that $\Omega^1_{\mathbf{H},\gamma_n,\eta_n} \neq \emptyset$.

For simplicity, let $\mathrm{H}^{10} := \mathrm{H}^{1}(\mathbb{R})^{\circ}$, $\mathrm{S}_{\mathrm{H}^{10}} := \mathrm{H}^{10}/\mathrm{H}^{10} \cap {}^{\circ}(\mathrm{H}^{1})(\mathbb{R})$ and $p : \mathrm{H}^{10} \to \mathrm{S}_{\mathrm{H}^{10}}$ be the natural quotient map. One sees that p factors through $\overline{p} : \mathrm{H}^{10}/\mathrm{H}^{10} \cap \Gamma \to \mathrm{S}_{\mathrm{H}^{10}}$. Then for any $g \in {}^{\circ}\mathrm{G}$, $\Omega^{1}_{\mathrm{H},g,\eta}$ descends along \overline{p} , i.e, $\Omega^{1}_{\mathrm{H},g,\eta} = \overline{p}^{-1} \left(\exp\left(\mathcal{P}^{1}_{\mathrm{H},g,\eta}\right) \right)$ where

$$\mathcal{P}_{\mathbf{H},g,\eta}^{\mathbf{1}} := \left\{ x \in \operatorname{Lie}(\mathbf{S}_{\mathbf{H}^{10}}) \mid \left\| \operatorname{Ad}(g\tilde{t})\mathbf{v}_{\mathbf{P}} \right\| > \eta, \ \forall \, \mathbf{P} \in \mathscr{P}_{\mathbf{H}^{\circ}}^{\max}, \ \tilde{t} \in p^{-1}(\exp(x)) \right\}.$$

For $\mathbf{P} \in \mathscr{P}_{\mathbf{H}^{\circ}}^{\max}$, let $l_{\mathbf{P}}$ be the induced linear functional on $\operatorname{Lie}(S_{\mathbf{H}^{10}})$ such that

 $\operatorname{Ad}(\widetilde{t})\mathbf{v}_{\mathbf{P}} = \exp(l_{\mathbf{P}}(x)) \cdot \mathbf{v}_{\mathbf{P}}, \ \forall \, \widetilde{t} \in p^{-1}(\exp(x)).$

Let Φ denote the collection of linear functionals l on $\text{Lie}(\mathbf{S}_{\mathbf{H}^{10}})$ such that $l = l_{\mathbf{P}}$ for some $\mathbf{P} \in \mathscr{P}_{\mathbf{H}^{\circ}}^{\max}$. Then

$$\mathcal{P}_{\mathbf{H},g,\eta}^{\mathbf{1}} = \left\{ x \in \operatorname{Lie}(\mathcal{S}_{\mathcal{H}^{\mathbf{10}}}) \mid l(x) > \log(\eta) - \log \inf_{\mathbf{P} \in \mathscr{P}_{\mathbf{H}^{o}}^{\max}, \, l_{\mathbf{P}} = l} \left\| \operatorname{Ad}(g) \mathbf{v}_{\mathbf{P}} \right\|, \, \forall \, l \in \Phi \right\}.$$

After passing to a subsequence, we assume that for each $l \in \Phi$,

 $\inf_{\mathbf{P}\in\mathscr{P}_{\mathbf{H}^{o}}^{\max},\, l_{\mathbf{P}}=l}\|\mathrm{Ad}(\gamma_{n})\mathbf{v}_{\mathbf{P}}\| \quad \text{either remains constant or diverges to infinity.}$

Let Φ_{∞} consist of those $l \in \Phi$ with (inf $||\operatorname{Ad}(\gamma_n)\mathbf{v}_{\mathbf{P}}||$) diverging to infinity and let $\widetilde{\Phi}$ be its complement. Let Φ_0 be consisting of $l_0 \in \widetilde{\Phi}$ such that there exists $(a_l)_{l \in \widetilde{\Phi}} \subset \mathbb{Z}_{\geq 0}$ such that $\sum a_l l = 0$ and $a_{l_0} \neq 0$. Let Φ_1 be the complement of Φ_0 in $\widetilde{\Phi}$. By [Zha21, Lemma 3.4], if $\Phi_0 = \emptyset$, then $\mathcal{P}^1_{\mathbf{H},\gamma_n,\eta_n}$ is non-empty for some (η_n) diverging to $+\infty$ and hence the proof is complete.

Note that by assumption, $\gamma_n \mathbf{H}^{\circ} \gamma_n^{-1}$ can not be contained in any proper observable Q-subgroup of **G** for infinitely many *n*'s. Let us assume that Φ_0 is non-empty and find a proper observable Q-subgroup containing $\gamma_n \mathbf{H}^{\circ} \gamma_n^{-1}$ for infinitely many *n*, which is a contradiction.

So take $(l_1, ..., l_k) \subset \widetilde{\Phi}$ and $(a_1, ..., a_k) \subset \mathbb{Z}^+$ with $\sum a_i l_i = 0$. As $(\inf \|\operatorname{Ad}(\gamma_n) \mathbf{v_P}\|)$ is bounded and $(\operatorname{Ad}(\gamma_n) \mathbf{v_P})$ is discrete, after passing to a subsequence, for each i, we find $(\mathbf{P}_n^i) \subset \mathscr{P}_{\mathbf{H}}^{\max}$ such that

$$\operatorname{Ad}(\gamma_n)\mathbf{v}_{\mathbf{P}_n^i} = \operatorname{Ad}(\gamma_1)\mathbf{v}_{\mathbf{P}_1^i} \quad \forall i = 1, ..., k, \ n \in \mathbb{Z}^+$$

Let $v_0 := \operatorname{Ad}(\gamma_1) \bigotimes_{i=1}^k \mathbf{v}_{\mathbf{P}_1^i}^{\otimes a_i}$, then v_0 lies in a **G**-representation space and is a $\gamma_n \mathbf{H}^{\circ} \gamma_n^{-1}$ -weight vector. Since $\sum a_i l_i = 0$, v_0 is fixed by $\gamma_n \mathbf{H}^{10} \gamma_n^{-1}$ for all n. Replacing v_0 by $v_0^{\otimes m}$ for some positive integer m, we assume that $\gamma_n \mathbf{H}^1 \gamma_n^{-1}$ fixes v_0 for all n.

RUNLIN ZHANG

Let $\beta_n \in \mathfrak{X}^*(\mathbf{H}^\circ)$ be such that $\gamma_n h \gamma_n^{-1} v_0 = \beta_n(h) v_0$ for every $h \in \mathbf{H}^\circ$. By rigidity of diagonalizable groups (see [Spr98, Proposition 3.2.8]), there are only finitely many possibilities for β_n . By passing to a subsequence, assume $\beta_n = \beta_1$ for all n. As β_1 trivializes on \mathbf{H}^1 , it factors through some $\overline{\beta} \in \mathfrak{X}^*(\mathbf{S}_{\mathbf{G}})$. Let W_β be a one-dimensional representation of \mathbf{G} , factoring through $\mathbf{S}_{\mathbf{G}}$, defined by

$$g \cdot w_{\beta} := \overline{\beta}^{-1} \left(p^{\operatorname{spl}}(g) \right) w_{\beta}, \quad \forall w_{\beta} \in W_{\beta}.$$

Fix some non-zero $w_{\beta} \in W_{\beta}$. Then $\gamma_n \mathbf{H}^{\circ} \gamma_n^{-1}$ is contained in the stabilizer subgroup of $v_0 \otimes w_{\beta}$ for all n. And we are done.

5. Good Height Functions

In this section, we reduce the counting problem to various properties of equivariant compactifications. Those actually satisfying these properties will be constructed in the next section under suitable assumptions.

Notations from Section 4.1 are inherited except that \mathbf{H}^1 is replaced by its identity component here. Also, for a smooth **G**-pair (\mathbf{X}, \mathbf{D}) over \mathbb{Q} such that $\mathbf{X} \setminus \mathbf{D}$ is **G**-equivariantly isomorphic to **U**, notations/definitions from Section 2.4 are kept.

5.1. Notations and assumptions. We start with some notations and assumptions.

- **G** is a connected linear algebraic group over \mathbb{Q} and let $\mathbf{G}^1 := (^{\circ}\mathbf{G})^{\circ}$;
- U is a variety over \mathbb{Q} that is homogeneous under some action of G and assume that $\mathbf{U}(\mathbb{Q}) \neq \emptyset$;
- Assume the canonical line bundle $\mathcal{K}_{\mathbf{U}}$ on \mathbf{U} is trivial and $\omega_{\mathbf{U}}$ is an invariant gauge form on \mathbf{U} ;
- for every $x \in \mathbf{U}(\mathbb{Q})$, let \mathbf{H}_x (resp. \mathbf{H}_x^1) be the stabilizer of x in \mathbf{G} (resp. \mathbf{G}^1). We assume that \mathbf{H}_x 's are connected.

Recall $G := \mathbf{G}(\mathbb{R})^{\circ}$ and for every \mathbb{R} -subgroup \mathbf{H} of \mathbf{G} , $\mathbf{H} := \mathbf{H}(\mathbb{R}) \cap \mathbf{G}$. Note that $\mathbf{G}^{1} = \mathbf{G} \cap \mathbf{G}^{1}(\mathbb{R})$ is connected but in general, \mathbf{H} could have different connected components. For $x \in \mathbf{U}(\mathbb{Q})$, choose $\mathbf{m}_{\mathbf{G}/\mathbf{H}_{x}}$, an invariant measure on $\mathbf{G}/\mathbf{H}_{x}$ that is identified with the measure induced from $\omega_{\mathbf{U},\infty}$ under the orbit map. We choose invariant measures $\mathbf{m}_{\mathbf{G}}$ on \mathbf{G} and $\mathbf{m}_{\mathbf{H}_{x}}$ on \mathbf{H}_{x} such that the triple $(\mathbf{m}_{\mathbf{G}}, \mathbf{m}_{\mathbf{H}_{x}}, \mathbf{m}_{\mathbf{G}/\mathbf{H}_{x}})$ is compatible. We similarly ask $(\mathbf{m}_{\mathbf{G}^{1}}, \mathbf{m}_{\mathbf{H}_{x}^{1}}, \mathbf{m}_{\mathbf{G}^{1}/\mathbf{H}_{x}^{1}})$ to be compatible. When $\mathbf{G}/\mathbf{H}_{x}$ is isomorphic to $\mathbf{G}^{1}/\mathbf{H}_{x}^{1}$, we further ask $\mathbf{m}_{\mathbf{G}^{1}/\mathbf{H}_{x}^{1}}$ to be identified with $\omega_{\mathbf{U},\infty}$. We also require that $\mathbf{m}_{[\mathbf{H}_{x}]}$ is compatible with $\mathbf{m}_{\mathbf{H}_{x}}$, namely the triple $\mathbf{m}_{[\mathbf{H}_{x}]}$, $\mathbf{m}_{\mathbf{H}_{x}}$ and the counting measure on $\Gamma \cap \mathbf{H}_{x}$ are compatible.

5.2. Definition of good and ok heights. Let $B_R^{\text{Ht}} := \{x \in \mathbf{U}(\mathbb{R}), \text{Ht}(x) \leq R\}$ for a function $\text{Ht} : \mathbf{U}(\mathbb{R}) \to \mathbb{R}_{\geq 0}$. And if $x \in \mathbf{U}(\mathbb{R})$, let $B_{R,x}^{\text{Ht}} := \{y \in \text{G.}x, \text{Ht}(y) \leq R\}$. The upper index Ht may be dropped if the function is clear from the context. Note that any arithmetic lattice Γ of \mathbf{G} has a finite-index subgroup contained in \mathbf{G} and hence in ${}^{\circ}\mathbf{G} = \mathbf{G}^1$.

5.2.1. Good and ok heights. If $\operatorname{Pic}(\mathbf{U})$ is torsion, we say that a function $\operatorname{Ht} : \mathbf{U}(\mathbb{R}) \to \mathbb{R}_{\geq 0}$ is **good** if for every $x \in \mathbf{U}(\mathbb{Q})$ and every arithmetic subgroup Γ of **G** contained in G^1 , one has that as R tends to infinity,

$$\#\Gamma . x \cap B_R^{\mathrm{Ht}} \sim \frac{|\mathbf{m}_{[\mathrm{H}_x^1]}|}{|\mathbf{m}_{[\mathrm{G}^1]}|} \mathbf{m}_{\mathrm{G}^1/\mathrm{H}_x^1} \left(B_{R,x}^{\mathrm{Ht}} \right).$$

The function Ht is said to be **ok** if the above asymptotic holds up to some constant c > 0 possibly dependent on x and Γ .

For $\eta > 0, g \in \mathbf{G}$ and $x \in \mathbf{U}(\mathbb{Q})$, let

$$\widetilde{B}_{R,x,\eta}^{\mathrm{Ht}} := \left\{ [g] \in \mathrm{G}^{1}/\mathrm{H}_{x} \cap \Gamma \ \Big| \ \mathrm{Ht}(g.x) \leq R, \ \|\mathrm{Ad}(g)\mathbf{v}_{\mathbf{P}}\| > \eta, \ \forall \ \mathbf{P} \in \mathscr{P}_{\mathbf{H}_{x}^{\circ}}^{\mathrm{max}} \right\}$$

where $\mathscr{P}_{\mathbf{H}_{\tau}^{\circ}}^{\max}$ denotes the set of maximal proper parabolic \mathbb{Q} -subgroups containing \mathbf{H}_{x}° .

Without assuming $\operatorname{Pic}(\mathbf{U})$ to be torsion, we say that a function $\operatorname{Ht} : \mathbf{U}(\mathbb{R}) \to \mathbb{R}_{\geq 0}$ is **good** if for every $x \in \mathbf{U}(\mathbb{Q}), \eta > 0$ and every arithmetic subgroup Γ of **G** contained in \mathbf{G}^1 , one has

$$\#\Gamma.x \cap B_R^{\mathrm{Ht}} \sim \frac{1}{|\mathbf{m}_{[\mathbf{G}^1]}|} \mathbf{m}_{[\mathbf{G}^1]_x} \left(\widetilde{B}_{R,x,\eta}^{\mathrm{Ht}} \right)$$

where $m_{[G^1]_x}$ denotes the G¹-invariant measure on $G^1/H_x \cap \Gamma$ compatible with m_{G^1} . The function Ht is said to be **ok** if the above asymptotic holds up to some constant c > 0 possibly dependent on x and

 Γ . One can check that this is compatible with the above when $\operatorname{Pic}(\mathbf{U})$ is torsion (that is, when $m_{[H_x^1]}$ is a finite measure).

5.2.2. Weighted versions. Unfortunately, in this paper we are not able to exhibit any good or ok height when $\text{Pic}(\mathbf{U})$ is not torsion. Nevertheless, here is a weighted version. For $\eta > 0$ and $y \in \mathbf{U}(\mathbb{Q})$, let

$$\Omega_{y,\eta} := \left\{ [h] \in \mathbf{H}_y^1/\mathbf{H}_y^1 \cap \Gamma \mid \|\mathrm{Ad}(h)\mathbf{v}_{\mathbf{P}}\| \ge \eta, \ \forall \, \mathbf{P} \in \mathscr{P}_{\mathbf{H}_y^\circ}^{\max} \right\}.$$

We say that $\text{Ht} : \mathbf{U}(\mathbb{R}) \to \mathbb{R}_{\geq 0}$ is **good with weights** if for every $x \in \mathbf{U}(\mathbb{Q})$, every arithmetic subgroup Γ contained in G^1 , there exists $\eta > 0$ such that for every c > 0 as $R \to +\infty$,

$$\sum_{y \in \Gamma.x \cap B_R^{\mathrm{Ht}}} \mathbf{w}_y \sim \frac{1}{\left|\mathbf{m}_{[\mathrm{G}^1]}\right|} \mathbf{m}_{\mathrm{G}^1/\mathrm{H}_x^1} \left(\mathrm{G}^1.x \cap B_R^{\mathrm{Ht}}\right) \tag{20}$$

where $\mathbf{w}_{y}^{-1} := \max \left\{ m_{[\mathrm{H}_{y}^{1}]} \left(\Omega_{y,\eta} \right), c \right\}$. Here we need to assume that $m_{[\mathrm{H}_{y}^{1}]}$ is the same as $m_{[\mathrm{H}_{x}^{1}]}$ when conjugating by γ .

The naive definition of "ok with weights" does not seem natural for us. Here is a slightly different one. We say that Ht is **ok with weights** if for every $x \in \mathbf{U}(\mathbb{Q})$ and every arithmetic subgroup Γ contained in G^1 , there exists a compactly supported non-negative function $\psi : \mathrm{G}^1/\Gamma \to \mathbb{R}$ such that

- (1) $\langle \psi, (\gamma h)_* \mathbf{m}_{[(\mathbf{H}^1_{\pi})^{\circ}]} \rangle \neq 0$ for all $\gamma \in \Gamma$ and $h \in \mathbf{H}^1_x$;
- (2) for some constant c > 0,

$$\sum_{y \in \Gamma.x \cap B_R^{\mathrm{Ht}}} \mathbf{w}_y^{\psi} \sim c \cdot \frac{1}{\left| \mathbf{m}_{[\mathrm{G}^1]} \right|} \mathbf{m}_{\mathrm{G}^1/\mathrm{H}_x^1} \left(\mathrm{G}^1.x \cap B_R^{\mathrm{Ht}} \right)$$

where for $y = \gamma . x$, we set $\mathbf{w}_y^{\psi} := \sum_{h \in \mathrm{H}^1_x/(\mathrm{H}^1_x)^{\circ}} \langle \psi, (\gamma h)_* \mathrm{m}_{[\mathrm{H}^{\circ}_x]} \rangle^{-1}$.

We usually take $\mathbf{G}^1 = \mathbf{G}$ in the discussion of heights that are ok with weights.

5.3. Conditions implying good heights. Here are some additional conditions on the pair (\mathbf{X}, \mathbf{D}) that might give good heights. We always assume that Ht is as constructed in Section 2.1.2. So implicitly there is some $\mathbf{L} := \sum \lambda_{\alpha} \mathbf{D}_{\alpha}$, a divisor supported on \mathbf{D} satisfying:

• For every $\alpha \in \mathscr{A}_{\mathbb{R},\mathbb{O}}^{\mathrm{an}}$, $\lambda_{\alpha} \geq 0$ and if $d_{\alpha} - 1 \geq 0$, then $\lambda_{\alpha} > 0$.

Let $\mathbf{B} := \sum_{\alpha \in \mathscr{B}} \mathbf{D}_{\alpha}$ be a closed \mathbb{Q} -subvariety of \mathbf{D} that is a union of irreducible divisors indexed by some subset \mathscr{B} of \mathscr{A} . We let $\mathscr{B}^{\mathrm{an}}_{\mathbb{R}}$ be the subset of $\mathscr{A}^{\mathrm{an}}_{\mathbb{R}}$ corresponding to sets contained in $\mathbf{B}(\mathbb{R})$. $\mathscr{B}^{\mathrm{an}}_{\mathbb{R},x}$ and $\mathscr{B}^{\mathrm{an}}_{\mathbb{R},\mathbb{O}}$ are defined in a similar way.

5.3.1. Conditions for good heights. We say that $(\mathbf{X}, \mathbf{D}, \mathbf{B})$ satisfies condition (B1) if

(B1) For every $x \in \mathbf{U}(\mathbb{Q})$ and $(g_n) \subset \mathbf{G}$ such that $\lim g_n \cdot x \in \mathbf{B}(\mathbb{R})$, one has

$$\lim_{n \to \infty} \left[(g_n)_* \mathbf{m}_{[\mathbf{H}_x^\circ]} \right] = \left[\mathbf{m}_{[\mathbf{G}]} \right].$$

For $x \in \mathbf{U}(\mathbb{Q}), R > 0$ and $\mathrm{Ht} : \mathbf{U}(\mathbb{R}) \to \mathbb{R}_{\geq 0}$, let

$$\nu_{R,x} := \frac{1}{\omega_{\mathbf{U},\infty} \left(B_{R,x}^{\mathrm{Ht}} \right)} \cdot \omega_{\mathbf{U},\infty} |_{B_{R,x}^{\mathrm{Ht}}}$$

be a probability measure on $\mathbf{X}(\mathbb{R})$.

We say that $(\mathbf{X}, \mathbf{D}, \mathbf{B}, \mathrm{Ht})$ satisfies condition (BH1) if

(BH1) For every $x \in \mathbf{U}(\mathbb{Q}), \nu_{\infty,x} := \lim_{R \to \infty} \nu_{R,x}$ exists and $\operatorname{supp}(\nu_{\infty,x}) \subset \mathbf{B}(\mathbb{R})$.

Lemma 5.1. Assume that (B1) and (BH1) hold. If $Pic(\mathbf{U})$ is torsion, then Ht is good. In general, Ht is good with weights.

5.3.2. Proof of Lemma 5.1. For $x \in \mathbf{U}(\mathbb{Q})$, by assumption, there exists some sequence $(g_n) \subset {}^{\circ}\mathbf{G}$ such that $\lim [(g_n)_* \mathbf{m}_{[\mathbf{H}_x^{\circ}]}] = [\mathbf{m}_{[\mathbf{G}]}]$, hence the natural injective morphism $\mathbf{H}_x/\mathbf{H}_x^1 \to \mathbf{S}_{\mathbf{G}}$ is an isomorphism by Lemma 4.4. So Theorem 4.10 is applicable. For simplicity, we write $\widetilde{\mathbf{S}}_x$ for $\widetilde{\mathbf{S}}^{\mathbf{H}_x}$ and $\Omega_{x,g,\eta}^1$ for $\Omega^1_{\mathbf{H}_x,g,\eta}$. Thus for every $\eta > 0$,

$$\lim_{n \to \infty} \frac{(g_n)_* \mathbf{m}_{[\mathrm{H}_x^{1}]}}{\mathbf{m}_{[(\mathrm{H}_x^{1})^\circ]} \left(\Omega_{x,g_n,\eta}^{\mathbf{1}}\right)} = \mathbf{m}_{[\mathrm{G}]}$$

In the following, for $g \in G$, we let $s_x \in H_x$ be such that $p^{spl}(g) = p^{spl}(s_x)$ and let $g_x := gs_x^{-1}$. In other words, we choose $g_x \in {}^{\circ}G$ such that $g_x \in gH_x$. Note that the coset space $g_x H_x^1$ is uniquely determined and $\Omega^1_{x,g_x,\eta}$ is independent of the choice of the coset representatives.

For a subset $\mathcal{B} \subset G/\Gamma$, let

$$\mathcal{B}_x := \left(p^{\mathrm{spl}} |_{\widetilde{\mathbf{S}}_x(\mathbb{R})^{\circ}} \right)^{-1} \left(p^{\mathrm{spl}}(\mathcal{B}) \right)$$

Assume that \mathcal{B} is non-empty, open, bounded and contains the identity coset, then there exists $\eta_{\mathcal{B}} > 0$ such that for every $g \in G$,

$$\{[h] \in \mathrm{H}_x^{\circ} \Gamma / \Gamma \mid g_x[h] \in \mathcal{B}\} \subset \mathcal{B}_x \times \Omega_{x, g_x, \eta_{\mathcal{B}}}$$

under the natural homeomorphism as in Equa.(17). One can check that this condition is independent of the choice of g_x . In particular, $g_* \mathbf{m}_{[\mathbf{H}_x^{\circ}]}|_{\mathcal{B}} = 0$ and actually $g_* \mathbf{m}_{[\mathbf{H}_x]}|_{\mathcal{B}} = 0$ if $\Omega_{x,g_x,\eta_{\mathcal{B}}} = \emptyset$. For any c > 0, let

$$\mathbf{w}_{g.x} := \left(\max\{ \operatorname{m}_{[(\operatorname{H}_{x}^{1})^{\circ}]} \left(\Omega_{x,g_{x},\eta} \right), c \} \right)^{-1}$$

Note that when $g = \gamma \in \Gamma$ and $y = \gamma . x$, $\mathbf{w}_{\gamma . x}$ coincides with the \mathbf{w}_y as defined following Equa.(20) since $\gamma_* \mathbf{m}_{[\mathbf{H}_x]}(\mathcal{B}) \neq 0$. In virtue of Theorem 3.3², it only remains to prove that

$$\frac{1}{\mathrm{m}_{\mathrm{G/H}_x}(B_{R,x}^{\mathrm{Ht}})} \int_{[g] \in B_{R,x}^{\mathrm{Ht}}} \mathbf{w}_{g.x} \cdot \left(g_* \mathrm{m}_{[\mathrm{H}_x]}\right) \, \mathrm{m}_{\mathrm{G/H}_x} \text{ converges to } C_0 \mathrm{m}_{[\mathrm{G}]}$$

against all test functions $\psi \in C_c(\mathcal{B})$. Here $C_0 := |\mathrm{H}_x/\mathrm{H}_x^{\circ}(\mathrm{H}_x \cap \Gamma)|$.

Now fix $\psi \in C_c(\mathcal{B})$ and $\varepsilon > 0$. By assumption and Theorem 4.10, Lemma 4.11, for R sufficiently large, we find Good \sqcup Bad = $B_{R,x}^{\text{Ht}}$ such that

1.

$$\frac{\mathrm{m}_{\mathrm{G/H}_x}(\mathrm{Bad})}{\mathrm{m}_{\mathrm{G/H}_x}(B_{R,x}^{\mathrm{Ht}})} \le \varepsilon \left(\left\| \psi \right\|_{\mathrm{sup}} \cdot C_0 \cdot \max\{\mathrm{m}_{\widetilde{\mathbf{S}}_x(\mathbb{R})^\circ}(\mathcal{B}_x), \mathrm{m}_{[\mathrm{G}]}(\mathcal{B})\} \right)^{-1}$$

2.

$$\left| \langle \psi, \frac{(g)_* \mathbf{m}_{[\mathbf{H}_x]}}{\mathbf{m}_{[(\mathbf{H}_x^1)^\circ]} \left(\Omega_{x,g_x,\eta}^1 \right)} \rangle - \langle \psi, C_0 \cdot \mathbf{m}_{[\mathbf{G}]} \rangle \right| \le \varepsilon, \; \forall \, g.x \in \text{Good}$$

3. $m_{[(H_x^1)^\circ]}(\Omega_{x,g_x,\eta}^1) > c$ for every $g.x \in \text{Good}$. Note that for every $g \in \mathcal{G}$,

$$\begin{aligned} \mathbf{w}_{g.x} \left| \langle \psi, g_* \mathbf{m}_{[\mathbf{H}_x]} \rangle \right| &\leq C_0 \cdot \mathbf{w}_{g.x} \mathbf{m}_{[(\mathbf{H}_x^1)^\circ]}(\Omega_{x,g_x,\eta_{\mathcal{B}}}) \cdot \|\psi\|_{\sup} \mathbf{m}_{\widetilde{\mathbf{S}}_x(\mathbb{R})^\circ}(\mathcal{B}_x) \\ &\leq C_0 \cdot \|\psi\|_{\sup} \mathbf{m}_{\widetilde{\mathbf{S}}_x(\mathbb{R})^\circ}(\mathcal{B}_x). \end{aligned}$$

Therefore, item 1 above implies that

$$\begin{aligned} & \left| \frac{1}{\mathrm{m}_{\mathrm{G/H}_{x}}(B_{R,x}^{\mathrm{Ht}})} \int_{[g]\in\mathrm{Bad}} \mathbf{w}_{g.x} \langle \psi, g_{*}\mathrm{m}_{[\mathrm{H}_{x}]} \rangle \,\mathrm{m}_{\mathrm{G/H}_{x}} - C_{0} \langle \psi, \mathrm{m}_{[\mathrm{G}]} \rangle \right| \\ & \leq \frac{\mathrm{m}_{\mathrm{G/H}_{x}}(\mathrm{Bad})}{\mathrm{m}_{\mathrm{G/H}_{x}}(B_{R,x}^{\mathrm{Ht}})} \left(C_{0} \cdot \|\psi\|_{\mathrm{sup}} \,\mathrm{m}_{\widetilde{\mathbf{S}}_{x}(\mathbb{R})^{\circ}}(\mathcal{B}_{x}) + C_{0} \cdot \|\psi\|_{\mathrm{sup}} \,\mathrm{m}_{[\mathrm{G}]}(\mathcal{B}) \right) \\ & \leq 2\varepsilon. \end{aligned}$$

 $^{^{2}}$ The remaining hypothesis either follows from Theorem 2.16 or can be verified directly.

On the other hand, $\mathbf{w}_{g.x}^{-1} = m_{[(\mathbf{H}_x^1)^\circ]} \left(\Omega_{x,g_x,\eta}^1\right)$ when $g.x \in \text{Good}$. Thus,

$$\left| \frac{1}{\mathbf{m}_{\mathrm{G/H}_{x}}(B_{R,x}^{\mathrm{Ht}})} \int_{[g]\in\mathrm{Good}} \mathbf{w}_{g.x} \langle \psi, g_{*}\mathbf{m}_{[\mathrm{H}_{x}]} \rangle \, \mathbf{m}_{\mathrm{G/H}_{x}} - C_{0} \langle \psi, \mathbf{m}_{[\mathrm{G}]} \rangle \right|$$

$$\leq \frac{\mathbf{m}_{\mathrm{G/H}_{x}}(\mathrm{Good})}{\mathbf{m}_{\mathrm{G/H}_{x}}(B_{R,x}^{\mathrm{Ht}})} \varepsilon \leq \varepsilon.$$

The proof is thus complete.

5.3.3. When the log anti-canonical height is good.

(B2) For every $x \in \mathbf{U}(\mathbb{Q})$ and $y \in \overline{\mathbf{G}.x}$, one has $\overline{\mathbf{G}.y} \cap \mathbf{B}(\mathbb{R}) \neq \emptyset$.

(K1) For every $\alpha \in \mathscr{A}_{\mathbb{R},\mathbb{O}}^{\mathrm{an}}, d_{\alpha} - 1 > 0.$

Lemma 5.2. If condition (B2) and (K1) hold, then condition (BH1) holds for the log anti-canonical height $Ht = Ht_{-(\mathbf{K}_{\mathbf{X}}+\mathbf{D})}$.

Proof. Fix $x \in \mathbf{U}(\mathbb{Q})$ and take $\mathbf{L} = -(\mathbf{K}_{\mathbf{X}} + \mathbf{D})$. By condition (K1), $\lambda_{\alpha} = d_{\alpha} - 1 > 0$ and hence $\frac{d_{\alpha} - 1}{\lambda_{\alpha}} = 1$ for every $\alpha \in \mathscr{A}_{\mathbb{R},x}^{\mathrm{an}}$. Thus, $\mathscr{C}_{\mathbb{R},x}^{\mathrm{an}} = \mathscr{C}_{\mathbb{R},x}^{\mathrm{an}}(\mathbf{L})$. By Theorem 2.16, $\nu_{\infty,x}$ exists and is supported on D_F as F varies over maximal faces of $\mathscr{C}_{\mathbb{R},x}^{\mathrm{an}}$. It remains to argue that $D_F \subset \mathbf{B}(\mathbb{R})$ for every such F.

Indeed, since D_F is G-invariant and closed, $D_F \cap D_\beta \neq \emptyset$ for some $\beta \in \mathscr{B}_{\mathbb{R},x}^{\mathrm{an}}$ by condition (B2). By maximality of F, we must have $\beta \in F$ and therefore, $D_F \subset D_\beta \subset \mathbf{B}(\mathbb{R})$.

5.3.4. When good heights exists.

(B4) For every $x \in \mathbf{U}(\mathbb{Q}), d_{\alpha} - 1 > 0$ for some $\alpha \in \mathscr{B}_{\mathbb{R},x}^{\mathrm{an}}$. In particular, $\mathscr{B}_{\mathbb{R},x}^{\mathrm{an}}$ is non-empty.

Lemma 5.3. If condition (B4) holds, then there exists an effective divisor \mathbf{L} supported on \mathbf{D} such that (BH1) holds for $\mathrm{Ht} := \mathrm{Ht}_{\mathbf{L}}$.

Proof. For every $x \in \mathbf{U}(\mathbb{Q})$, choose $\beta_x \in \mathscr{B}_{\mathbb{R},x}$ with $d_{\beta_x} - 1 > 0$. Choose $(\lambda_\alpha)_{\alpha \in \mathscr{A}}$ such that

$$\min_{x \in \mathbf{U}(\mathbb{Q})} \left\{ \frac{d_{\beta_x} - 1}{\lambda_{\beta_x}} \right\} > \max_{\alpha \in \mathscr{A} \setminus \{\beta_x\}} \left\{ \frac{d_\alpha - 1}{\lambda_\alpha} \right\}.$$

For instance, one can pick

$$\lambda_{\alpha} := \begin{cases} d_{\alpha} - 1 & \text{if } \alpha \in \{\beta_x\} \\ 2(d_{\alpha} - 1) & \text{if } d_{\alpha} - 1 > 0 \text{ and } \alpha \notin \{\beta_x\} \\ 1 & \text{otherwise} \end{cases}$$

Then by Theorem 2.16, for every $x \in \mathbf{U}(\mathbb{Q})$, $\operatorname{supp}(\nu_{\infty,x}) \subset \mathbf{B}(\mathbb{R})$.

5.3.5. Ok heights. Assume that \mathbf{H}_x° has no non-trivial \mathbb{Q} -characters. For $x \in \mathbf{U}(\mathbb{Q})$ and $\Gamma \subset \mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, let $\Psi_x : \mathbf{G}.x \to \operatorname{Prob}(\mathbf{G}/\Gamma)$ be defined by $\Psi_x(g.x) := g_* \mathbf{m}_{[\mathbf{H}_x]}^1$. Let $\mu_{R,x} := \operatorname{Bary}((\Psi_x)_*(\nu_{R,x})) \in \operatorname{Prob}(\mathbf{G}/\Gamma)$. Here $\operatorname{Bary} : \operatorname{Prob}(\operatorname{Prob}(\mathbf{G}/\Gamma)) \to \operatorname{Prob}(\mathbf{G}/\Gamma)$ is defined by $\nu \mapsto \int_{\phi \in \operatorname{Prob}(\mathbf{G}/\Gamma)} \phi \nu(\phi)$.

(H1) For every $x \in \mathbf{U}(\mathbb{Q})$ and arithmetic subgroup $\Gamma \subset \mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, \mathbf{H}_x° has no non-trivial \mathbb{Q} -characters and $\lim_{R\to\infty} \mu_{R,x}$ exists in $\operatorname{Prob}(\mathbf{G}/\Gamma)$.

The following is a direct corollary of Theorem 3.3 and 2.16.

Lemma 5.4. If (H1) holds, then Ht is ok.

- (D1) For every $x \in \mathbf{U}(\mathbb{Q})$ and arithmetic subgroup $\Gamma \subset \mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, \mathbf{H}_x° has no non-trivial \mathbb{Q} -characters and Ψ_x extends continuously to $\overline{\mathbf{G}.x} \to \operatorname{Prob}(\mathbf{G}/\Gamma) \cup \{0\}$ where the closure is taken in $\mathbf{X}^{\operatorname{cor}}$, the manifold with corners associated with (\mathbf{X}, \mathbf{D}) (see Section 6.4.2).
- (S1) For every $x \in \mathbf{U}(\mathbb{Q})$ and arithmetic subgroup Γ in $\mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, there exists a bounded subset B of \mathbf{G}/Γ such that $g\mathbf{H}_x^\circ\Gamma/\Gamma$ intersects with B for every $g \in \mathbf{G}$.

Lemma 5.5. If (D1) and (S1) hold, then (H1) holds for $\text{Ht} = \text{Ht}_{\mathbf{L}}$ for every \mathbf{L} from Section 2.4, namely, $\lambda_{\alpha} \geq 0$ and $d_{\alpha} > 0 \implies \lambda_{\alpha} > 0$ where $\mathbf{L} = \sum \lambda_{\alpha} \mathbf{D}_{\alpha}$.

Proof. Indeed, Ψ_x extends continuously to $\overline{G.x} \to \operatorname{Prob}(G/\Gamma)$. By Theorem 6.13, $\lim \nu_{R,x}$ exists in $\operatorname{Prob}(X^{\operatorname{cor}})$. Applying $\operatorname{Bary} \circ (\Psi_x)_*$, we get that $\lim \mu_{R,x}$ exists in $\operatorname{Prob}(G/\Gamma)$.

With more care, (S1) can be weakened as

(DS1) There exists a closed subset D of $\mathbf{B}(\mathbb{R})$ such that for every (g_n) and every $x \in \mathbf{U}(\mathbb{Q})$ with lim $g_n.x$ not in D, there exists a bounded subset B of G/Γ such that $g_n \mathrm{H}_x^{\circ} \Gamma/\Gamma$ intersects with B for every n. Moreover, for every $F \subset \mathscr{C}_{\mathbb{R},\mathbb{Q}}^{\mathrm{an}}$, if Leb_F denotes a smooth measure on D_F , then Leb_F(B) = 0.

With a similar proof, noting that $\mu_{\infty,x}\left((\Psi_x)^{-1}(\{\mathbf{0}\})\right) = 0$, one has:

Lemma 5.6. If (D1) and (DS1) hold, then (H1) holds for $Ht = Ht_L$ for every L from Section 2.4.

5.3.6. Ok with weights. Assume that **G** has no non-trivial \mathbb{Q} -characters. Take $x \in \mathbf{U}(\mathbb{Q})$ and an arithmetic subgroup Γ . Now we allow \mathbf{H}_x° to have non-trivial \mathbb{Q} -characters. Take a non-negative compactly supported function ψ on G/Γ such that $\langle \psi, g_* \mathbf{m}_{[\mathbf{H}_x^{\circ}]} \rangle \neq 0$ for all $g \in G$. Such a ψ exists if (S1) holds. Now we define

$$\begin{split} \Psi_x^{\psi} &: \mathbf{G}.x \to \operatorname{Prob}^{\psi}(\mathbf{G}/\Gamma) \\ g.x &\mapsto \alpha_g^{\psi}\left(\mathbf{m}_{[\mathbf{H}_x]}^{\psi}\right). \end{split}$$

See Section 6.4.5 for the definition of $\alpha_g^{\psi}\left(\mathbf{m}_{[\mathbf{H}_x]}^{\psi}\right)$ and $\operatorname{Prob}^{\psi}(\mathbf{G}/\Gamma)$. Let

 $\mu_{R,x}^{\psi} := \operatorname{Bary}\left((\Psi_x^{\psi})_*(\nu_{R,x})\right) \in \operatorname{Prob}^{\psi}(\mathcal{G}/\Gamma).$

(H2) For every $x \in \mathbf{U}(\mathbb{Q})$ and arithmetic subgroup $\Gamma \subset \mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, there exists a non-negative compactly supported continuous function ψ on \mathbf{G}/Γ such that $\langle \psi, g_*\mathbf{m}_{[\mathbf{H}_x^o]} \rangle \neq 0$ for all $g \in \mathbf{G}$ and for every such ψ , $\lim_{R\to\infty} \mu_{R,x}^{\psi}$ exists in $\operatorname{Prob}^{\psi}(\mathbf{G}/\Gamma)$.

Lemma 5.7. Assume that **G** has no non-trivial \mathbb{Q} -characters. If (H2) holds, then Ht is ok with weights.

The proof is analogous to that in Section 5.3.5, except that there are weights and Theorem 3.3 should be applied to $H = H_x^{\circ}$ rather than $H = H_x$.

(D2) For every $x \in \mathbf{U}(\mathbb{Q})$ and arithmetic subgroup $\Gamma \subset \mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, there exists ψ satisfying the paragraph above and Ψ_x^{ψ} extends continuously to $\overline{\mathbf{G}.x} \to \operatorname{Prob}^{\psi}(\mathbf{G}/\Gamma)$ where the closure is taken in X^{cor}, the manifold with corners associated with (\mathbf{X}, \mathbf{D}) .

Lemma 5.8. Assume that **G** has no non-trivial \mathbb{Q} -characters. If (D2) and (S1) hold, then (H2) holds for every Ht = Ht_L.

The proof is analogous to that of Lemma 5.5 with $\operatorname{Prob}^{\psi}(G/\Gamma)$ in the place of $\operatorname{Prob}(G/\Gamma)$.

6. Equivariant Compactifications and Focusing

Notations from Section 4.1 and 5.1 are inherited. Furthermore, **G** and **H** are assumed to be connected linear algebraic groups over \mathbb{Q} and **H** is assumed to be observable in **G**.

We will construct compactifications (and heights) of \mathbf{G}/\mathbf{H} satisfying various properties from Section 5. Two guiding examples are

- (1) **G** and **H** are connected reductive and $\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ} \subset \mathbf{H}$;
- (2) \mathbf{G} is \mathbb{Q} -split semisimple and \mathbf{H} contains a maximal unipotent subgroup of \mathbf{G} .

6.1. Finiteness of intermediate subgroups. Let

 $INT(\mathbf{H}, \mathbf{G}) := \{ \mathbf{H} \subset \mathbf{L} \subsetneq \mathbf{G} \mid \mathbf{L} \text{ is connected} \}$

be the set of intermediate connected closed subgroups between \mathbf{H} and \mathbf{G} . Let $\mathrm{INT}_{\mathbb{Q}}(\mathbf{H}, \mathbf{G})$ (resp. INT^{obs}(\mathbf{H}, \mathbf{G})) collect elements in $\mathrm{INT}(\mathbf{H}, \mathbf{G})$ that are defined over \mathbb{Q} (resp. that are observable). Let $\mathrm{INT}_{\Gamma}(\mathbf{H}, \mathbf{G})$ denote the even smaller subset consisting of $\mathbf{L} \in \mathrm{INT}_{\mathbb{Q}}^{\mathrm{obs}}(\mathbf{H}, \mathbf{G})$ such that $(\gamma_n \mathbf{H} \gamma_n^{-1})$ converges to \mathbf{L} (see Definition 4.1) for some sequence $(\gamma_n) \subset \Gamma$. We say that (\mathbf{G}, \mathbf{H}) satisfies condition (F1) if (F1) $INT_{\Gamma}(\mathbf{H}, \mathbf{G})$ is finite.

Lemma 6.1. Assume G is reductive and that one of the following is true:

- (1) **H** is reductive and $\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ}$ is contained in **H**;
- (2) **H** contains a maximal unipotent subgroup.

Then (F1) holds.

From the proof, $INT(\mathbf{H}, \mathbf{G})$ is finite in case (1), which is no longer true in case (2).

6.1.1. Proof of Lemma 6.1, part 1. Under our assumptions, the central torus $\mathbf{Z}(\mathbf{G})$ of \mathbf{G} is contained in \mathbf{H} , so it suffices to show this for $\mathbf{H}/\mathbf{Z}(\mathbf{G}) \subset \mathbf{G}/\mathbf{Z}(\mathbf{G})$. When $\mathbf{H}/\mathbf{Z}(\mathbf{G})$ is semisimple, this already follows from [EMV09, Lemma 3.1].

In general, every intermediate observable subgroup is reductive by Theorem 4.5. Then, up to $\mathbf{G}(\overline{\mathbb{Q}})$ -conjugacy, there are only finitely many $\mathbf{L} \in \mathrm{INT}^{\mathrm{obs}}(\mathbf{H}, \mathbf{G})$ (see for instance, [EMV09, Lemma A.1] and the beginning part of [ZZ21, Section 4]). Fix finitely many representatives $\{\mathbf{L}_1, ..., \mathbf{L}_k\} \subset \mathrm{INT}^{\mathrm{obs}}(\mathbf{H}, \mathbf{G})$. By Lemma 6.6, for each i = 1, ..., k, find $(f_i^i)_{j=1,...,l_i} \subset \mathbf{Z}(\mathbf{H}, \mathbf{L}_i)(\overline{\mathbb{Q}})$ such that

$$\mathbf{Z}(\mathbf{H},\mathbf{L}_i)(\overline{\mathbb{Q}}) = \bigcup_{j=1}^{l_k} \mathbf{L}_i(\overline{\mathbb{Q}}) f_j^i \mathbf{H}(\overline{\mathbb{Q}})$$

where we used the fact that \mathbf{H} is of finite-index in $N_{\mathbf{G}}(\mathbf{H})$ under our assumptions. Now we claim that

$$\text{INT}^{\text{obs}}(\mathbf{H}, \mathbf{G}) = \left\{ (f_j^i)^{-1} \mathbf{L}_i f_j^i \mid i = 1, ..., k; \ j = 1, ..., l_i \right\}$$

Indeed, for $\mathbf{L} \in \text{INT}^{\text{obs}}(\mathbf{H}, \mathbf{G})$, we find $g \in \mathbf{G}(\overline{\mathbb{Q}})$ and $i \in \{1, ..., k\}$ such that $g\mathbf{L}g^{-1} = \mathbf{L}_i$. So $g \in \mathbf{Z}(\mathbf{H}, \mathbf{L}_i)(\overline{\mathbb{Q}})$. Find $l \in \mathbf{L}_i(\overline{\mathbb{Q}})$, $h \in \mathbf{H}(\overline{\mathbb{Q}})$ and $j \in \{1, ..., l_i\}$ such that $g = lf_i^i h$. Then,

$$\mathbf{L} = g^{-1} \mathbf{L}_i g = (f_j^i h)^{-1} \mathbf{L}_i f_j^i h = (f_j^i)^{-1} \mathbf{L}_i f_j^i.$$

So we are done.

6.1.2. Proof of Lemma 6.1, part 2. Assume that **H** contains a maximal unipotent subgroup and $\mathbf{L} \in \operatorname{INT}_{\Gamma}(\mathbf{H}, \mathbf{G})$. By Section 6.3.1, $\mathbf{L} = \mathbf{M}_{\mathbf{L}} \cdot \mathbf{T}_{\mathbf{L}}$ with $\mathbf{M}_{\mathbf{L}}$ normalized by the torus $\mathbf{T}_{\mathbf{L}}$ and $\mathfrak{l} = \mathfrak{m}_{\mathfrak{l}} \oplus \mathfrak{t}_{\mathfrak{l}}$. When **L** is defined over \mathbb{Q} , both $\mathbf{M}_{\mathbf{L}}$ and $\mathbf{T}_{\mathbf{L}}$ can be arranged to be over \mathbb{Q} . Let $\pi : \mathbf{L} \to \mathbf{L}/\mathbf{M}_{\mathbf{L}}$ be the natural quotient morphism.

Let $(\gamma_n) \subset \Gamma$ be such that $(\gamma_n \mathbf{H} \gamma_n^{-1})$ converges to **L**. Then $(\pi(\gamma_n \mathbf{H} \gamma_n^{-1})) = (\pi(\mathbf{H}))$ also converges to $\mathbf{L}/\mathbf{M}_{\mathbf{L}}$, which is impossible unless $\pi(\mathbf{H}) = \mathbf{L}/\mathbf{M}_{\mathbf{L}}$. Hence $\mathbf{L} = \mathbf{M}_{\mathbf{L}} \cdot \mathbf{H}$. This implies that **L** is determined by $\mathbf{M}_{\mathbf{L}}$. So there are only finitely many possibilities.

6.2. Compactifications using intermediate subgroups.

6.2.1. Condition (F2). To detect nondivergence we introduce

(F2) there are only finitely many parabolic \mathbb{Q} -subgroups containing \mathbf{H} , i.e., $\mathscr{P}_{\mathbf{H}}$ is finite.

For simplicity we abbreviate INT := INT_Γ(**H**, **G**) and INTP := INT_Γ(**H**, **G**) $\cup \mathscr{P}_{\mathbf{H}}$ in this subsection. Clearly INTP is finite if INT_Q^{obs}(**H**, **G**) is, but the converse does not hold. For instance **G** = **SL**_n $(n \geq 3)$ and **H** is a maximal unipotent Q-subgroup. Another case is when **G** = **SL**_n ×**D** $(n \geq 3)$ with **D** embedded in **SL**_n as a Q-anisotropic maximal Q-torus and **H** is the diagonal embedding of **D** in **G**. For a parabolic Q-subgroup **P**, we let $\mathscr{V}(\mathbf{P})$ consist of all triples $(\rho, \mathbf{V}_{\rho}, \mathbf{v}_{\rho})$ where

- 1. \mathbf{V}_{ρ} is a finite dimensional vector space over \mathbb{Q} ;
- 2. $\rho: \mathbf{G} \to \mathbf{G}/\mathbf{R}_{\mathbf{u}}(\mathbf{G}) \to \mathbf{GL}(\mathbf{V}_{\rho})$ is a \mathbb{Q} -representation of \mathbf{G} factoring through $\mathbf{G}/\mathbf{R}_{\mathbf{u}}(\mathbf{G})$;
- 3. $\mathbf{v}_{\rho} \in \mathbf{V}_{\rho}(\mathbb{Q})$ is not fixed by $\rho(\mathbf{P})$ but the line spanned by \mathbf{v}_{ρ} is preserved by $\rho(\mathbf{P})$.

By abuse of notation, we simply write an element of $\mathscr{V}(\mathbf{P})$ as ρ or \mathbf{v}_{ρ} when no confusion might arise. And we let $\alpha_{\mathbf{v}_{\rho}}$ denote the Q-character of \mathbf{P} attached to \mathbf{v}_{ρ} . Let $\pi'_{\mathbf{P}} : \mathbf{P} \to \mathbf{S}'_{\mathbf{P}}$ be a Q-split quotient torus of \mathbf{P} such that every Q-character of \mathbf{P} factors through $\pi'_{\mathbf{P}}$. By rigidity of diagonalizable groups,

$$\left\{\pi'_{\mathbf{P}}(g\mathbf{H}g^{-1}) \mid g \in \mathbf{Z}(\mathbf{H}, \mathbf{P})\right\} = \{\mathbf{S}_1, ..., \mathbf{S}_k\}$$

is finite. For each \mathbf{S}_i , let

$$\mathscr{V}(\mathbf{P})^{\mathbf{S}_i,\mathbf{H}} := \{ \rho \in \mathscr{V}(\mathbf{P}) \mid \mathbf{v}_{\rho} \text{ is fixed by } \mathbf{S}_i \text{ and } \mathbf{H} \}$$

We find a finite subset $\mathscr{V}_i \subset \mathscr{V}(\mathbf{P})^{\mathbf{S}_i,\mathbf{H}}$ such that $\{\alpha_{\mathbf{v}_{\rho}} \mid \rho \in \mathscr{V}(\mathbf{P})^{\mathbf{S}_i,\mathbf{H}}\}$ is contained in the cone spanned by $\{\alpha_{\mathbf{v}_{\rho}} \mid \rho \in \mathscr{V}_i\}$. Let $\mathscr{V}(\mathbf{H},\mathbf{P}) := \bigcup \mathscr{V}_i$ and $\mathscr{V}(\mathbf{H}) := \bigcup_{\mathbf{P} \in \mathscr{P}_{\mathbf{H}}} \mathscr{V}(\mathbf{H},\mathbf{P})$.

6.2.2. Definition of the compactification. For a finite dimensional vector space \mathbf{V} over \mathbb{Q} , let $\mathbf{P}(\mathbf{V})$ be the associated projective variety over \mathbb{Q} . For $\mathbf{L} \in \text{INT}$, let $\mathbf{v}_{\mathfrak{l}}$ be a vector in $\mathbf{V}_{\mathfrak{l}} := \wedge^{\dim \mathbf{L}} \mathfrak{g}$ that lifts \mathfrak{l} . In order that $\mathbf{v}_{\mathfrak{l}}$ is fixed by \mathbf{H} we need to assume that the modular character of \mathbf{L} is trivial on \mathbf{H} . If \mathbf{H} is assumed to have no non-trivial \mathbb{Q} -characters, then this automatically holds. To get an embedding of \mathbf{G}/\mathbf{H} , we fix a rational \mathbf{G} -representation $\mathbf{V}_{\mathbf{L}}$ and $\mathbf{v}_{\mathbf{H}} \in \mathbf{V}_{\mathbf{H}}(\mathbb{Q})$ such that the stabilizer of $\mathbf{v}_{\mathbf{H}}$ in \mathbf{G} is exactly \mathbf{H} .

Definition 6.2. Assume both \mathbf{G}, \mathbf{H} are connected and (F1), (F2) hold. Moreover, assume that the modular character of \mathbf{L} is trivial on \mathbf{H} for every $\mathbf{L} \in \text{INT}$. Let

$$\begin{split} \iota_{\mathbf{H}}^{\mathrm{INTP}} &: \mathbf{G}/\mathbf{H} \to \mathbf{V}_{\mathbf{H}} \oplus \bigoplus_{\mathbf{L} \in \mathrm{INT}} \mathbf{V}_{\mathfrak{l}} \oplus \bigoplus_{\rho \in \mathscr{V}(\mathbf{H})} \mathbf{V}_{\rho} \to \mathbf{P}(\mathbf{V}_{\mathbf{H}} \oplus \mathbb{Q}) \oplus \bigoplus_{\mathbf{L} \in \mathrm{INT} \,\sqcup \, \mathscr{V}(\mathbf{H})} \mathbf{P}(\mathbf{V}_{\mathfrak{l}} \oplus \mathbb{Q}) \\ & q\mathbf{H} \mapsto q.\mathbf{v}_{\mathbf{H}} \oplus (\oplus q.\mathbf{v}_{\mathfrak{l}}) \oplus (\oplus q.\mathbf{v}_{\rho}) \end{split}$$

Let $\mathbf{X}_{\mathbf{H}}^{\mathrm{INTP}}$ be the Zariski closure of $\iota_{\mathbf{H}}^{\mathrm{INTP}}(\mathbf{G}/\mathbf{H})$ and $\mathbf{X}_{\mathrm{H}}^{\mathrm{INTP}}$ be the analytic closure of G.o in $\mathbf{X}_{\mathbf{H}}^{\mathrm{INTP}}(\mathbb{R})$ where o denotes the image of the identity coset.

By extending every representation ρ here to $(\rho \oplus 1, \mathbf{V} \oplus \mathbb{Q})$, we see that $\iota_{\mathbf{H}}^{\text{INTP}}$ is **G**-equivariant.

6.2.3. Nondivergence. Every element \mathbf{v} in $X_{\mathrm{H}}^{\mathrm{INTP}}$ can be written as $\oplus[\mathbf{v}^{\mathbf{L}}: t^{\mathbf{L}}]$ with $t^{\mathbf{L}} \in \{0, 1\}$. Let (g_n) be a sequence in G and $\mathbf{v} = \lim_{n \to \infty} g_n.o.$ If $\mathbf{v}^{\rho} = 0$ for some $\rho \in \mathscr{V}(\mathbf{H})$, then $(g_n \mathrm{H} \Gamma / \Gamma)$ diverges topologically in G/Γ by Mahler's criterion. The converse is also true by Theorem 4.6:

Lemma 6.3. Assume (F1) and (F2) hold. If (g_n) is a sequence in G such that $\mathbf{v} = \lim_{n \to \infty} g_n$ o exists in $X_{\mathrm{H}}^{\mathrm{INTP}}$ and $\mathbf{v}^{\rho} \neq \mathbf{0}$ for every $\rho \in \mathscr{V}(\mathbf{H})$, then $g_n \mathrm{H}^{\circ} \Gamma / \Gamma$ intersects with some bounded set in G/ Γ for all n.

Proof. By Theorem 4.6, if the conclusion were not true, then there exist $(\gamma_n) \subset \Gamma$, $(h_n) \subset \mathrm{H}^\circ$, a parabolic Q-subgroup **P** containing all $\gamma_n \mathbf{H} \gamma_n^{-1}$ and a rational **P**-eigenvector **v** that is fixed by $\gamma_n \mathbf{H} \gamma_n^{-1}$ and lives in a **G**-representation that factors through $\mathbf{G}/\mathbf{R}_{\mathbf{u}}(\mathbf{G})$ such that $g_n h_n \gamma_n^{-1} \cdot \mathbf{v} \to \mathbf{0}$. So $g_n \gamma_n^{-1} \cdot \mathbf{v} \to \mathbf{0}$. Since $\gamma_n^{-1} \mathbf{P} \gamma_n \in \mathscr{P}_{\mathbf{H}}$ for all n and $\mathscr{P}_{\mathbf{H}}$ is assumed to be finite, $\gamma_n^{-1} \mathbf{P} \gamma_n = \gamma_1^{-1} \mathbf{P} \gamma_1$ for all n after passing to a subsequence.

Let $\mathbf{P}' := \gamma_1^{-1} \mathbf{P} \gamma_1 \in \mathscr{P}_{\mathbf{H}}$ and $\mathbf{v}' := \gamma_1^{-1} \cdot \mathbf{v}$. Then \mathbf{v}' is a \mathbf{P}' -eigenvector fixed by $\gamma_1^{-1} \gamma_n \mathbf{H} \gamma_n^{-1} \gamma_1$ for all n. Passing to a subsequence, we find i_0 such that $\pi'_{\mathbf{P}'}(\gamma_1^{-1} \gamma_n \mathbf{H} \gamma_n^{-1} \gamma_1) = \mathbf{S}_{i_0}$ for all n. So $d\alpha_{\mathbf{v}'}$ is a $\mathbb{R}_{\geq 0}$ -linear combination of $\{d\alpha_{\mathbf{v}_{\rho}}, \rho \in \mathscr{V}_{i_0}\}$. In particular, passing to a further subsequence, we can find a fixed $\rho_0 \in \mathscr{V}_{i_0} \subset \mathscr{V}(\mathbf{H}, \mathbf{P}')$ such that (note that $\gamma_n^{-1} \gamma_1 \in \mathbf{P}' \cap \Gamma$)

$$g_n.\mathbf{v}' = \pm g_n \gamma_n^{-1} \gamma_1.\mathbf{v}' \to \mathbf{0} \implies g_n.\mathbf{v}_{\rho_0} \to \mathbf{0}$$

In particular, $\mathbf{v}^{\rho_0} = \mathbf{0}$ and this is a contradiction.

Definition 6.4. Assume condition (F1) and (F2) hold and that the modular character of **L** is trivial on **H** for every $\mathbf{L} \in INT$. Let $\mathbf{B}_{\mathbf{H}}^{INTP}$ be the **G**-invariant closed \mathbb{Q} -subvariety of $\mathbf{X}_{\mathbf{H}}^{INTP}$ consisting of elements of the form $\bigoplus_{\mathbf{L} \in \{\mathbf{H}\} \sqcup INT \sqcup \mathscr{V}(\mathbf{H})} [\mathbf{v}^{\mathbf{L}} : t^{\mathbf{L}}]$ with $t^{\mathbf{L}} = 0$ for every $\mathbf{L} \in INT$.

Lemma 6.5. Assume that **G** is reductive, or more generally, the unipotent radical of **G** commutes with **G**. Also assume that Condition (F1), (F2) hold and that the modular character of **L** is trivial on **H** for every $\mathbf{L} \in \text{INT}$. Let (g_n) be a sequence in **G** such that $(g_n.o)$ converges to $\mathbf{v} \in X_{\text{H}}^{\text{INTP}}$, then

$$\lim_{n \to \infty} \left[(g_n)_* \mathbf{m}_{[\mathrm{H}]} \right] = \left[\mathbf{m}_{[\mathrm{G}]} \right] \iff \mathbf{v} \in \mathrm{B}_{\mathrm{H}}^{\mathrm{INTP}}$$

In particular, assuming the equivalent conditions in Lemma 4.4, $\mathbf{B}_{\mathbf{H}}^{\mathrm{INTP}}$ is non-empty.

Proof. The non-trivial direction is " \Leftarrow ". So we assume $g_n \cdot \mathbf{v}_{\mathfrak{l}} \to \infty$ for all $\mathbf{L} \in INT$.

By Theorem 4.6, we can find $(h_n) \subset \mathrm{H}^\circ$, $(\gamma_n) \subset \Gamma$, a parabolic Q-subgroup **P** and an observable Q-subgroup **O** such that $\gamma_n \mathbf{H} \gamma_n^{-1}$ is contained in **P** and strongly converges to **O**. Replacing γ_n by $\gamma_1^{-1} \gamma_n$ if necessary, we assume $\mathbf{H} \subset \mathbf{O}$ and hence $\mathbf{O} \in \mathrm{INT}$. Thus, $\gamma_n^{-1} \mathbf{O} \gamma_n \in \mathrm{INT}$ for all n. But INT is finite, by passing to a subsequence, we assume that $\gamma_n^{-1} \mathbf{O} \gamma_n = \gamma_1^{-1} \mathbf{O} \gamma_1 =: \mathbf{O}'$ for all n. So $\gamma_n^{-1} \gamma_1$

normalizes \mathbf{O}' , implying $\gamma_n^{-1}\gamma_1 \cdot \mathbf{v}_{\mathfrak{o}'} = \pm \mathbf{v}_{\mathfrak{o}'}$. Thus $g_n \cdot \mathbf{v}_{\mathfrak{o}'} = \pm g_n h_n \gamma_n^{-1} \gamma_1 \cdot \mathbf{v}_{\mathfrak{o}'}$. Let $g_n h_n \gamma_n^{-1} \gamma_1 = k_n a_n p_n$ using the horospherical coordinate attached to \mathbf{P}' (and some choice of maximal reductive subgroup of \mathbf{G} and Cartan involution), then Theorem 4.6, together with our assumption on \mathbf{G} , asserts that $\alpha(a_n)$ is bounded for all character appearing in the Lie algebra of \mathbf{P}' and in particular, of \mathbf{O}' . Since (p_n) is bounded, we have that $g_n h_n \gamma_n^{-1} \gamma_1 \cdot \mathbf{v}_{\mathfrak{o}'} = k_n a_n p_n \cdot \mathbf{v}_{\mathfrak{o}'}$ is bounded. This is a contradiction against our assumption.

6.3. Conjugacy of subgroups. Consider the following closed subvariety of G:

$$\mathbf{Z}(\mathbf{H}, \mathbf{L}) := \left\{ g \in \mathbf{G} \mid g\mathbf{H}g^{-1} \subset \mathbf{L} \right\}$$

for two Q-subgroups \mathbf{H}, \mathbf{L} of \mathbf{G} . From the definition, it admits an action of \mathbf{L} by multiplying from the left and an action of $\mathbf{N}_{\mathbf{G}}(\mathbf{H})$ from the right. One naturally wonders whether the action has finitely many orbits. We formulate two conditions related to this. We say that (\mathbf{G}, \mathbf{H}) satisfies (C1) or (N1) if

- (C1) $\mathbf{Z}(\mathbf{H}, \mathbf{L})$ decomposes into finitely many orbits under the action of $\mathbf{L} \times \mathbf{N}_{\mathbf{G}}(\mathbf{H})$ for every $\mathbf{L} \in$ INT (\mathbf{H}, \mathbf{G}) ;
- (N1) $\Gamma \cap \mathbf{H}$ is a finite index subgroup of $\Gamma \cap \mathbf{N}_{\mathbf{G}}(\mathbf{H})$ for an(y) arithmetic subgroup Γ .

Lemma 6.6. Assume that one of the following is true:

- (1) **G** and **H** are connected reductive and $\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ} \subset \mathbf{H}$;
- (2) **H** contains a maximal unipotent subgroup of **G**, which is a \mathbb{Q} -split semisimple group.
- Then (C1) and (N1) hold.

Item 1 is true by [EMS96, Lemma 5.2] (cf. [Ric67, Theorem 8.1]). Item 2 will be proved in the following subsections. Note that (2) might fail if **G** is not Q-split, for instance, when $\mathbf{G} = \operatorname{Res}_{\mathbb{Q}(\sqrt{2})/\mathbb{Q}}(\mathbf{SL}_2)$ (but the discussion in the last subsection still applies). For now, we note the following consequence of these conditions plus Theorem 2.3:

Proposition 6.7. Assume condition (C1) holds. Let Γ be an arithmetic subgroup of \mathbf{G} , then there exists a finite-index subgroup Γ' with the following property: The action of $\mathbf{L} \cap \Gamma' \times \mathbf{N}_{\mathbf{G}}(\mathbf{H}) \cap \Gamma'$ on $\mathbf{Z}(\mathbf{H}, \mathbf{L}) \cap \Gamma'$ has finitely many orbits for every \mathbf{L} . In particular, if (N1) also holds, then the action of $\mathbf{L} \cap \Gamma' \times \mathbf{H} \cap \Gamma'$ has finitely many orbits.

Indeed, Γ' can be taken to be the intersection of Γ with any congruence subgroup.

Remark 6.8. The statement [EMS96, Lemma 5.2] seems to suggest that (C1) holds whenever **G** is reductive. This is wrong. One can find unipotent subgroups **H** and **L** of $\mathbf{G} := \mathbf{SL}_6$ such that (C1) fails (even after replacing the action of **L** by that of $\mathbf{N}_{\mathbf{G}}(\mathbf{L})$).

6.3.1. Structure of **L** containing \mathbf{U}_{\max} . Fix a Borel subgroup **B** containing a maximal unipotent subgroup \mathbf{U}_{\max} and a maximal torus **T** in **B**. Let Φ be the set of (non-zero) weights. Thus $\mathfrak{g} = (\bigoplus_{\alpha \in \Phi} \mathfrak{g}_{\alpha}) \oplus \mathfrak{t}$. Let **L** be a connected closed subgroup containing \mathbf{U}_{\max} , then one can show that its Lie algebra is

$$\mathfrak{l} = (\oplus_{\alpha \in \Phi}, \mathfrak{g}_{\alpha}) \oplus (\mathfrak{l} \cap \mathfrak{t}), \quad \exists \Phi_{\mathfrak{l}} \subset \Phi.$$

$$\tag{21}$$

Let $\mathfrak{m}_{\mathfrak{l}}$ be the Lie subalgebra generated by $\bigoplus_{\alpha \in \Phi_{\mathfrak{l}}} \mathfrak{g}_{\alpha}$ and $\mathbf{M}_{\mathbf{L}}$ the corresponding connected algebraic subgroup. Then there exists a torus $\mathbf{T}_{\mathbf{L}}$ in \mathbf{T} such that $\mathbf{L} = \mathbf{M}_{\mathbf{L}} \cdot \mathbf{T}_{\mathbf{L}}$ and $\mathfrak{l} = \mathfrak{m}_{\mathfrak{l}} \oplus \mathfrak{t}_{\mathfrak{l}}$. So we get an injection

$$INT(\mathbf{U}, \mathbf{G}) \hookrightarrow 2^{\Phi} \times \{ subtori \text{ of } \mathbf{T} \}$$
$$\mathbf{L} \mapsto (\Phi_{\mathfrak{l}}, \mathbf{T}_{\mathbf{L}}).$$

6.3.2. Proof of Lemma 6.6: verify (N1). Now we explain that $\mathbf{N}_{\mathbf{G}}(\mathbf{H}) \cap \Gamma$ is virtually contained in \mathbf{H} under the assumption that \mathbf{G} is \mathbb{Q} -split and \mathbf{H} contains a maximal unipotent subgroup \mathbf{U}_{\max} . It suffices to show that $\mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\circ}/\mathbf{H}^{\circ}$ is a \mathbb{Q} -split torus (without assuming \mathbf{G} to be \mathbb{Q} -split, $\mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\circ}/\mathbf{H}^{\circ}$ is only proved to be a \mathbb{Q} -torus). As \mathbf{G} is \mathbb{Q} -split, we assume that \mathbf{U}_{\max} and \mathbf{T} are defined over \mathbb{Q} .

By Equa.(21), **H** is normalized by **T**. Hence $N_{\mathbf{G}}(\mathbf{H})$ contains a Borel subgroup, implying that it is a parabolic subgroup **P**. In particular, it is connected. By assumption **H** contains $\mathbf{R}_{\mathbf{u}}(\mathbf{P})$ and so $\mathbf{H}/\mathbf{R}_{\mathbf{u}}(\mathbf{P})$ is a normal subgroup of $\mathbf{P}/\mathbf{R}_{\mathbf{u}}(\mathbf{P})$. But $\mathbf{H}/\mathbf{R}_{\mathbf{u}}(\mathbf{P})$ contains a maximal unipotent subgroup of $\mathbf{P}/\mathbf{R}_{\mathbf{u}}(\mathbf{P})$, so $\mathbf{H}/\mathbf{R}_{\mathbf{u}}(\mathbf{P})$ contains every semisimple factor of $\mathbf{P}/\mathbf{R}_{\mathbf{u}}(\mathbf{P})$. Hence \mathbf{P}/\mathbf{H} is a quotient of the central torus of $\mathbf{P}/\mathbf{R}_{\mathbf{u}}(\mathbf{P})$, which is \mathbb{Q} -split since **G** is \mathbb{Q} -split. 6.3.3. Proof of Lemma 6.6: verify (C1). Condition (C1) is a geometric one and holds without assuming **G** to be \mathbb{Q} -split.

Take $g \in \mathbf{Z}(\mathbf{H}, \mathbf{L})(\overline{\mathbb{Q}})$, so $g\mathbf{U}_{\max}g^{-1} \subset g\mathbf{H}g^{-1} \subset \mathbf{L}$. As both $g\mathbf{U}_{\max}g^{-1}$ and \mathbf{U}_{\max} are maximal unipotent subgroups of \mathbf{L} , there exists $l_q \in \mathbf{L}(\overline{\mathbb{Q}})$ such that

$$l_g g \mathbf{U}_{\max}(l_g g)^{-1} = \mathbf{U}_{\max} \subset \mathbf{H}_g := l_g g \mathbf{H}(l_g g)^{-1} \subset \mathbf{L}.$$

Thus $l_g g$ belongs to the normalizer of \mathbf{U}_{\max} , which is exactly \mathbf{B} . By the structure of subgroups containing \mathbf{U}_{\max} , \mathbf{H} is normalized by \mathbf{T} and hence \mathbf{B} . So we conclude that $l_g g \in \mathbf{N}_{\mathbf{G}}(\mathbf{H})$ and $\mathbf{Z}(\mathbf{H}, \mathbf{L})(\overline{\mathbb{Q}}) = \mathbf{L}(\overline{\mathbb{Q}}) \cdot \mathbf{N}_{\mathbf{G}}(\mathbf{H})(\overline{\mathbb{Q}})$.

6.3.4. Another compactification by intermediate groups. For each $\mathbf{L} \in \text{INT}$, we fix a rational **G**-representation $\mathbf{V}_{\mathbf{L}}$ and $\mathbf{v}_{\mathbf{L}} \in \mathbf{V}_{\mathbf{L}}(\mathbb{Q})$ such that the stabilizer of $\mathbf{v}_{\mathbf{L}}$ in **G** is exactly **L**. Here we do not need to assume that the modular character of **L** is trivial on **H**.

Definition 6.9. Assume both G, H are connected and (F1), (F2) hold. Let

$$\ell_{\mathbf{H}}^{\mathrm{IMTP}} : \mathbf{G}/\mathbf{H} \to \bigoplus_{\mathbf{L}\in\mathrm{INT}} \mathbf{V}_{\mathbf{L}} \oplus \bigoplus_{\rho\in\mathscr{V}(\mathbf{H})} \mathbf{V}_{\rho} \to \bigoplus_{\mathbf{L}\in\mathrm{INT}\,\sqcup\,\mathscr{V}(\mathbf{H})} \mathbf{P}(\mathbf{V}_{\mathbf{L}}\oplus\mathbb{Q})$$
$$g\mathbf{H} \mapsto (\oplus g.\mathbf{v}_{\mathbf{L}}) \oplus (\oplus g.\mathbf{v}_{\rho})$$

Let $\mathbf{X}_{\mathbf{H}}^{\mathrm{IMTP}}$ be the Zariski closure of $\iota_{\mathbf{H}}^{\mathrm{IMTP}}(\mathbf{G}/\mathbf{H})$ and $\mathbf{X}_{\mathbf{H}}^{\mathrm{IMTP}}$ be the analytic closure of G.o in $\mathbf{X}_{\mathbf{H}}^{\mathrm{IMTP}}(\mathbb{R})$ where o denotes the image of the identity coset. Define $\mathbf{B}_{\mathbf{H}}^{\mathrm{IMTP}}$ in the same way as Definition 6.4.

Statement analogous to Lemma 6.5 also holds assuming additionally (C1) and (N1).

6.3.5. Focusing and intermediate subgroups. For a sequence (g_n) in G, assuming $g_n H^{\circ} \Gamma / \Gamma$ intersects with some bounded set in G/ Γ for all n, then $g_n h_n = \delta_n \gamma_n$ for some $(h_n) \subset H^{\circ}, (\gamma_n) \subset \Gamma$ and bounded $(\delta_n) \subset G$. So the limiting behaviour of $g_n H^{\circ} \Gamma / \Gamma$ is essentially the same as $\gamma_n H^{\circ} \Gamma / \Gamma$. A sequence (γ_n) in Γ is said to be **clean** if for every $\mathbf{L} \in INT$, one of the following holds

- 1. $(\gamma_n \cdot \mathbf{v_L})$ is bounded, or equivalently, (γ_n) is bounded modulo **L**;
- 2. $(\gamma_n \cdot \mathbf{v_L})$ diverges, or equivalently, (γ_n) diverges modulo **L**.

For a clean sequence (γ_n) , let

 $\mathscr{L}((\gamma_n)) := \{ \mathbf{L} \in \text{INT} \mid (\gamma_n \cdot \mathbf{v}_{\mathbf{L}}) \text{ is bounded } \}.$

Lemma 6.10. Let (γ_n) be a clean sequence in Γ . Further assume condition (C1) and (N1) hold. Then for every $\mathbf{L}_1, \mathbf{L}_2 \in \mathscr{L}((\gamma_n))$, there exists $\mathbf{L}' \in \mathscr{L}((\gamma_n))$ that is contained in $\mathbf{L}_1 \cap \mathbf{L}_2$. In particular, $\mathscr{L}((\gamma_n))$ has a unique minimum element, denoted as $\mathbf{L}((\gamma_n))$.

Proof. Since $(\gamma_n \mathbf{v}_{\mathbf{L}_1})$ and $(\gamma_n \mathbf{v}_{\mathbf{L}_2})$ are bounded, we must have that $(\gamma_n (\mathbf{v}_{\mathbf{L}_1} \oplus \mathbf{v}_{\mathbf{L}_2}))$ is bounded, that is, (γ_n) is bounded modulo $\mathbf{L}_1 \cap \mathbf{L}_2$. Without loss of generality we assume (γ_n) is contained in $\mathbf{L}_3 := (\mathbf{L}_1 \cap \mathbf{L}_2)^\circ$. But we do not know whether \mathbf{L}_3 belongs to INT. If not, then by Lemma 4.4, at least one of the following is true

- 1. **H** is contained in a proper normal \mathbb{Q} -subgroup of \mathbf{L}_3 ;
- 2. $\mathbf{HL}_{3}^{\mathrm{ari}}$ is a proper subgroup of \mathbf{L}_{3} .

Let \mathbf{L}_4 be the identity component of the proper normal Q-subgroup in case 1 and be $\mathbf{HL}_3^{\operatorname{ari}}$ in case 2. Note that \mathbf{L}_4 is observable in **G**. In the second case, (γ_n) is clearly bounded modulo \mathbf{L}_4 by the definition of $\mathbf{L}_3^{\operatorname{ari}}$. For the first case, we invoke the condition (C1), (N1) and Proposition 6.7 to see that $\gamma_n = l_{\gamma_n}^4 f_{i_n} h_{\gamma_n}$ for some $l_{\gamma_n}^4 \in \mathbf{L}_4 \cap \Gamma$, $h_{\gamma_n} \subset \mathbf{H} \cap \Gamma$ and f_{i_n} belongs to certain finite subset of $\mathbf{Z}(\mathbf{H}, \mathbf{L}_4)$. Since (γ_n) belongs to \mathbf{L}_3 , (f_{i_n}) is contained in $\mathbf{Z}(\mathbf{H}, \mathbf{L}_4) \cap \mathbf{L}_3$. And since \mathbf{L}_4 is normal in \mathbf{L}_3 , we find that $\gamma_n = f_{i_n}(f_{i_n}^{-1}l_{\gamma_n}^4f_{i_n})h_{\gamma_n}$ is bounded modulo \mathbf{L}_4 .

Continuing this process one will end with the smallest element in $\mathscr{L}((\gamma_n))$.

6.4. Resolution of singularities and manifolds with corners. Assume $m_{[H]}$ is finite for the moment. It is not true that in general, the natural map³

$$\Psi_{\mathbf{H}} : \mathbf{G}.o \to \operatorname{Prob}(\mathbf{G}/\Gamma)$$
$$g.x \mapsto g_* \mathbf{m}_{[\mathbf{H}]}^{\mathbf{1}}$$

extends to a continuous map from X_{H}^{IMTP} to $Prob(G/\Gamma) \sqcup \{0\}$. This is because the limiting measure of $(\gamma_n)_* m_{[H]}^1$ is of the form $m_{[L^\circ H]}^1$ but the stabilizer of $m_{[L^\circ H]}$ may not contain L even though it contains a Zariski-dense subset.

The issue would disappear if one had assumed that $L^{\circ}H = L$ for all $\mathbf{L} \in INT_{\Gamma}(\mathbf{H}, \mathbf{G})$. This is indeed the case, for instance, when \mathbf{G} is semisimple and \mathbf{H} contains a maximal \mathbb{R} -split torus (see [Mat64] or [BT65, Theorem 14.4]). As we are not assuming this, one is forced to pass to the non-algebraic world. It will be shown (see Theorem 6.16 and 6.20 below) that under some assumptions $\Psi_{\mathbf{H}}$ extends to the closure in a manifold with corners.

6.4.1. *Resolution of singularities.* As a first step, one applies the resolution of singularity due to Hironaka [Hir64]. The desired statement can be found in [Wo05, Theorem 1.0.2] and the lifting of algebraic group actions is explained in [Kol07, Proposition 3.9.1].

Theorem 6.11. Let (\mathbf{X}, \mathbf{D}) be a **G**-pair such that $\mathbf{X} \setminus \mathbf{D}$ is homogeneous. There exists a smooth **G**-pair $(\widetilde{\mathbf{X}}, \widetilde{\mathbf{D}})$ over \mathbb{Q} and a **G**-equivariant morphism $\pi : \widetilde{\mathbf{X}} \to \mathbf{X}$ such that π is an isomorphism restricted to $\pi^{-1}(\mathbf{X} \setminus \mathbf{D}) = \widetilde{\mathbf{X}} \setminus \widetilde{\mathbf{D}}$.

Assume (F1), (F2) hold and $\mathbf{B}_{\mathbf{H}}^{\mathrm{IMTP}}$ is non-empty. Apply Theorem 6.11 to the pair $(\mathbf{X}_{\mathbf{H}}^{\mathrm{IMTP}}, \mathbf{B}_{\mathbf{H}}^{\mathrm{IMTP}})$. We get $\pi_1 : (\mathbf{X}_1, \mathbf{B}_1) \to (\mathbf{X}_{\mathbf{H}}^{\mathrm{IMTP}}, \mathbf{B}_{\mathbf{H}}^{\mathrm{IMTP}})$ where $(\mathbf{X}_1, \mathbf{B}_1)$ is a smooth pair over \mathbb{Q} . Let \mathbf{D}_1 be the preimage of $\mathbf{D}_{\mathbf{H}}^{\mathrm{IMTP}}$ under π_1 where $\mathbf{D}_{\mathbf{H}}^{\mathrm{IMTP}} := \mathbf{X}_{\mathbf{H}}^{\mathrm{IMTP}} \setminus \iota_{\mathbf{H}}^{\mathrm{IMTP}}(\mathbf{G}/\mathbf{H})$. Apply Theorem 6.11 again to the pair $(\mathbf{X}_1, \mathbf{D}_1)$ to get $\pi_2 : (\mathbf{X}_2, \mathbf{D}_2) \to (\mathbf{X}_1, \mathbf{D}_1)$ where $(\mathbf{X}_2, \mathbf{D}_2)$ is a smooth pair. If \mathbf{B}_2 denotes the preimage of the (Cartier) divisor \mathbf{B}_1 , then \mathbf{B}_2 is a union of certain irreducible components of \mathbf{D}_2 . Define $\widetilde{\mathbf{X}}_{\mathbf{H}}^{\mathrm{IMTP}} := \mathbf{X}_2, \widetilde{\mathbf{D}}_{\mathbf{H}}^{\mathrm{IMTP}} := \mathbf{D}_2$ and $\widetilde{\mathbf{B}}_{\mathbf{H}}^{\mathrm{IMTP}} := \mathbf{B}_2$. If one is only interested in whether $(g_n)_* \mathbf{m}_{[\mathrm{H}]}$ converges to the full Haar measure, then it is not necessary to pass to the non-algebraic world.

6.4.2. Manifolds with corners. Let (\mathbf{X}, \mathbf{D}) be a smooth **G**-pair over \mathbb{Q} with $\mathbf{X} \setminus \mathbf{D}$ being homogeneous under the action of **G**. We take $o \in (\mathbf{X} \setminus \mathbf{D})(\mathbb{Q})$. Let X be the analytic closure of G.o and D be its intersection with $\mathbf{D}(\mathbb{R})$. We define in this section the associated "manifold with corners", denoted as $(X^{cor}, \mathbf{D}^{cor})$, together with an equivariant continuous map π^{cor} onto (X, D).

For i = 1, 2, ..., d, let $\mathbf{e}_i(x_1, ..., x_d) := x_i$ be the standard coordinate functions on \mathbb{R}^d . Then X can be glued by local charts $(\mathcal{O}_i, \varphi_i)_{i \in \mathscr{I}}$, where (\mathcal{O}_i) is an open covering of X, such that for each $i \in \mathscr{I}$,

1. $\varphi_i : \mathcal{O}_i \to \varphi_i(\mathcal{O}_i)$ is a homeomorphism onto a connected open neighborhood of $\mathbf{0} \in \mathbb{R}^d$;

2. there exists $0 \leq r_i \leq d$ such that

$$\varphi_i(\mathbf{D} \cap \mathcal{O}_i) = \varphi(\mathcal{O}_i) \cap \left(\bigcup_{j=1,\dots,r_i} \{\mathbf{e}_j = 0\}\right).$$

By shrinking \mathcal{O}_i 's if necessary, we further assume the following: For every (i, j) such that $\mathcal{O}_{ij} := \mathcal{O}_i \cap \mathcal{O}_j \neq \emptyset$, there exist $I_{ij} \subset \{1, ..., r_i\}$, $I_{ji} \subset \{1, ..., r_j\}$ and a bijection $\tau_{ij} : I_{ij} \to I_{ji}$ such that

- 1. $\varphi_i(\mathcal{O}_{ij}) \cap \{\mathbf{e}_k = 0, \forall k \in I_{ij}\} \neq \emptyset, \varphi_j(\mathcal{O}_{ji}) \cap \{\mathbf{e}_k = 0, \forall k \in I_{ji}\} \neq \emptyset;$
- 2. for every $k \in \{1, ..., r_i\} \setminus I_{ij}, \varphi_i(\mathcal{O}_{ij}) \cap \{\mathbf{e}_k = 0\} = \emptyset$, every $k \in \{1, ..., r_j\} \setminus I_{ji}, \varphi_j(\mathcal{O}_{ji}) \cap \{\mathbf{e}_k = 0\} = \emptyset$;
- 3. for every $k \in I_{ji}$, there is a number $\epsilon_k^{ij} \in \{-1, 1\}$ such that the homeomorphism

$$\varphi_j \circ \varphi_i^{-1} : \varphi_i(\mathcal{O}_{ij}) \to \varphi_j(\mathcal{O}_{ij})$$

sends $\mathbf{e}_{\tau_{ij}^{-1}(k)} > 0$ to $\mathbf{e}_k \epsilon_k^{ij} > 0$.

³This was called Ψ_o in Section 5.3.5.

We define a new topological space X^{cor} by "cutting along the boundaries". For $\kappa = \kappa(\cdot) \in \{-1, 1\}^{r_i}$ and $i \in \mathscr{I}$, let

$$\mathcal{O}_i^{\kappa} := \left\{ x \in \mathcal{O}_i, \ \kappa(k) \mathbf{e}_k(x) \ge 0, \ \forall \ k = 1, ..., r_i \right\};$$
$$\mathcal{O}_i^{\kappa, \circ} := \left\{ x \in \mathcal{O}_i, \ \kappa(k) \mathbf{e}_k(x) > 0, \ \forall \ k = 1, ..., r_i \right\}.$$

To distinguish \mathcal{O}_i^{κ} (resp. \mathcal{O}_i) as an independent topological space from a subset of X, let $\iota_i^{\kappa} : \mathcal{O}_i^{\kappa} \to X$ (resp. $\iota_i : \mathcal{O}_i \to X$) be the natural inclusion map. Their inverses are well defined and continuous on the image.

We define an equivalence relation \sim on $\bigsqcup_{i \in \mathscr{I}, \kappa \in \{-1,1\}^{r_i}} \mathcal{O}_i^{\kappa}$ as follows: $x \sim y$ iff there exist $i, j \in \mathscr{I}$ and $\kappa_i \in \{-1, 1\}^{r_i}, \kappa_j \in \{-1, 1\}^{r_j}$ such that

$$x \in \mathcal{O}_i^{\kappa_i}, \ y \in \mathcal{O}_j^{\kappa_j}, \ \iota_i^{\kappa_i}(\mathcal{O}_i^{\kappa_i,\circ}) \cap \iota_j^{\kappa_j}(\mathcal{O}_j^{\kappa_j,\circ}) \neq \emptyset \text{ and } \iota_i^{\kappa_i}(x) = \iota_j^{\kappa_j}(y)$$

Let $X^{cor} := \bigsqcup_{i,\kappa} \mathcal{O}_i^{\kappa} / \sim$ be equipped with the quotient topology.

Note that \mathcal{O}_i^{κ} can be viewed as an open subset of X^{cor}. Also, the natural continuous map $\bigsqcup_{i,\kappa} \mathcal{O}_i^{\kappa} \to X$ factors through some map $\pi^{cor} : X^{cor} \to X$, which is continuous by the definition of quotient topology. Let D^{cor} be the preimage of D under π^{cor} . From the construction we see that

Lemma 6.12. The restriction of π^{cor} to $X^{cor} \setminus D^{cor}$, or to each \mathcal{O}_i^{κ} , is a homeomorphism onto its image.

As the structures are compatible with the group action, one can also check that the G-action on X lifts to a continuous G-action on X^{cor}.

Let us also point out that the proof of Theorem 2.16 in [CLT10] also yields similar equidistribution statements on the manifolds with corners:

Theorem 6.13. Under the weak-* topology, the family of probability measures on X^{cor}

$$\mu_{R,x} := \frac{\mathbf{1}_{B_{R,x}} \cdot \operatorname{Vol}}{\operatorname{Vol}(B_{R,x})}$$

has a limit ν in $\operatorname{Prob}(X^{\operatorname{cor}})$ as $R \to +\infty$. Moreover, the support of ν is equal to the union of $Z^{\operatorname{cor}} := (\pi^{\operatorname{cor}})^{-1}(Z)$ as (I, Z) varies over faces of $\mathscr{C}^{\operatorname{an}}_{\mathbb{R},x}(\mathbf{L})$ of dimension b_x .

6.4.3. Measure compactifications, finite volume cases. Whereas Theorem 4.2 gives a satisfactory description for the limiting behaviour of $(g_n)_* \mathbf{m}_{[\mathrm{H}^\circ]}$, what is really needed for the counting problem is the limiting behaviour of $(g_n)_* \mathbf{m}_{[\mathrm{H}]}$. For this assume **H** to be connected and apply the following corollary to the weak approximation property [PR94, Theorem 7.7]:

Lemma 6.14. Let A be a connected linear algebraic group. Under the analytic topology, $A(\mathbb{Q})$ is dense in $A(\mathbb{R})$.

By Lemma 6.14, we choose a finite subset $(c_i^{\mathbf{H}})_{i \in I} \subset \mathbf{H}(\mathbb{Q})$ such that

$$\mathbf{H} = \bigsqcup_{i \in I} c_i^{\mathbf{H}} \mathbf{H}^{\circ},$$

and let

$$\Gamma^{\mathrm{sma}} := \bigcap_{i \in I} c_i^{\mathbf{H}} \Gamma(c_i^{\mathbf{H}})^{-1}.$$

 Γ^{sma} is a finite index subgroup of Γ . We define two equivalence relations on I. We say that $i \sim_1 j$ iff the image of $c_i^{\mathbf{H}}$ in $\mathbf{H}/\mathbf{H}^{\circ}(\mathbf{H} \cap \Gamma)$ equals to that of $c_j^{\mathbf{H}}$. For $\mathbf{L} \in \text{INT}_{\Gamma}(\mathbf{H}, \mathbf{G})$, we say that $i \sim_{\mathbf{L}} j$ iff the image of $c_i^{\mathbf{H}}$ in $\mathbf{H}/\mathbf{H} \cap (\mathbf{L}^{\circ} \cdot (\mathbf{L} \cap \Gamma))$ equals to that of $c_j^{\mathbf{H}}$. One can check that

$$\mathbf{H}\Gamma = \bigsqcup_{i \in I/\sim_1} c_i^{\mathbf{H}} \mathbf{H}^{\circ} \Gamma, \quad \mathbf{L}^{\mathbf{H}}\Gamma = \bigsqcup_{i \in I/\sim_{\mathbf{L}}} c_i^{\mathbf{H}} \mathbf{L}^{\circ} \Gamma$$

where $L^{H} := L^{\circ}H$. Hence,

$$\begin{split} \mathbf{m}_{[\mathbf{H}]}^{\mathbf{1}} &= \left| I / \sim_{1} \right|^{-1} \sum_{[i] \in I / \sim_{\mathbf{1}}} (c_{i}^{\mathbf{H}})_{*} \mathbf{m}_{[\mathbf{H}^{\circ}]}^{\mathbf{1}}, \\ \mathbf{m}_{[\mathbf{L}^{\mathbf{H}}]}^{\mathbf{1}} &= \left| I / \sim_{\mathbf{L}} \right|^{-1} \sum_{[i] \in I / \sim_{\mathbf{L}}} (c_{i}^{\mathbf{H}})_{*} \mathbf{m}_{[\mathbf{L}^{\circ}]}^{\mathbf{1}}. \end{split}$$

Lemma 6.15. Assume **H** is connected and $m_{[H]}$ is finite. Let (γ_n) be a sequence in Γ^{sma} such that $(\gamma_n \mathbf{H} \gamma_n^{-1})$ converges to **L**, then $\lim_{n\to\infty} (\gamma_n)_* m_{[H]}^1 = m_{[L^H]}^1$.

To avoid redundancy, the reader is referred to Lemma 6.18 below for the proof in a similar case.

6.4.4. The measure compactification is dominated by manifolds with corners. For the sake of simplicity, write $X := \widetilde{X}_{H}^{IMTP,cor}$ and $D := \widetilde{D}_{H}^{IMTP,cor}$ in this subsection.

Theorem 6.16. Assume (F1), (F2), (C1) and (N1) hold and that **H** is connected with no non-trivial \mathbb{Q} -characters. Let (x_n) be a sequence in X\D converging to some $x_{\infty} \in X$. Then $(\Psi_{\mathbf{H}}(x_n))$ is convergent in $\operatorname{Prob}(G/\Gamma) \sqcup \{\mathbf{0}\}$. Consequently, $\Psi_{\mathbf{H}}$ extends to a continuous map $\overline{\Psi}_{\mathbf{H}} : X \to \operatorname{Prob}(G/\Gamma) \sqcup \{\mathbf{0}\}$.

Such an extension $\overline{\Psi}_{\mathbf{H}}$ is necessarily G-equivariant.

Proof. Let π denote the natural projection $X \to X_H^{IMTP}$. Write

$$\pi(\mathbf{x}_{\infty}) =: \mathbf{v}_{\infty} = \bigoplus_{\mathbf{L} \in \text{INT} \cup \mathscr{V}(\mathbf{H})} [\mathbf{v}_{\infty}^{\mathbf{L}} : t_{\infty}^{\mathbf{L}}].$$

If $\mathbf{v}_{\infty}^{\rho} = \mathbf{0}$ for some $\rho \in \mathscr{V}(\mathbf{H})$, then $(\Psi_{\mathbf{H}}(x_n))$ converges to **0**. Assume otherwise, by Lemma 6.3, we find $(\gamma_n) \subset \Gamma^{\text{sma}}$ and a bounded sequence $(\delta_n) \subset \mathbf{G}$ such that $x_n = \delta_n \gamma_n .o.$ Let $\mathbf{L}_{\infty} := \mathbf{L}((\gamma_n))$ by Lemma 6.10, noting that (γ_n) is a clean sequence since $(\delta_n \gamma_n [\mathbf{v}_{\mathbf{L}} : 1])$ converges. Let $\delta_{\infty} \in \mathbf{G}$ be such that $\lim \delta_n \gamma_n \mathbf{v}_{\mathbf{L}_{\infty}} = \delta_{\infty} \mathbf{v}_{\mathbf{L}_{\infty}}$. As we are going to apply Proposition 6.7 (to Γ^{sma} in the place of Γ), we may replace Γ^{sma} by a finite-index subgroup from the beginning and assume that the conclusion of Proposition 6.7 holds with $\Gamma' = \Gamma^{\text{sma}}$.

By assumption, $(\gamma_n \cdot \mathbf{v}_{\mathbf{L}_{\infty}})$ is bounded. Take a finite set $\mathcal{F} \subset \Gamma^{\text{sma}}$ such that for every *n*, there exists $\gamma \in \mathcal{F}$ such that $\gamma_n \cdot \mathbf{v}_{\mathbf{L}_{\infty}} = \gamma \cdot \mathbf{v}_{\mathbf{L}_{\infty}}$. By ignoring finitely many terms, we can find a subsequence (γ_{n_k}) and $\gamma \in \mathcal{F}$ such that $\gamma_{n_k} \mathbf{v}_{\mathbf{L}_{\infty}} = \gamma \cdot \mathbf{v}_{\mathbf{L}_{\infty}}$ for all *k*. We claim that $(\gamma^{-1}\gamma_{n_k}\mathbf{H}\gamma_{n_k}^{-1}\gamma)$ converges to \mathbf{L}_{∞} .

Write $l_k := \gamma^{-1} \gamma_{n_k} \in \mathbf{L}_{\infty} \cap \Gamma^{\text{sma}}$. If the claim were not true, by passing to a further subsequence, we may assume that $l_k \mathbf{H} l_k^{-1}$'s are all contained in some $\mathbf{L} \in \text{INT}_{\Gamma}(\mathbf{H}, \mathbf{G})$, a proper subgroup of \mathbf{L}_{∞} . By Proposition 6.7 and passing to a further subsequence, we assume $l_k = l'_k f h_k$ where $(l'_k) \subset \mathbf{L} \cap \Gamma^{\text{sma}}$, $(h_k) \subset \mathbf{H} \cap \Gamma^{\text{sma}}$ and $f \in \mathbf{Z}(\mathbf{H}, \mathbf{L}_{\infty}) \cap \Gamma^{\text{sma}}$. Note that actually f is contained in \mathbf{L}_{∞} . Write $\mathbf{L}_f := f^{-1}\mathbf{L}f$, then it is also an element of $\text{INT}_{\Gamma}(\mathbf{H}, \mathbf{G})$ contained in \mathbf{L}_{∞} . But (l_k) , and hence (γ_{n_k}) , is bounded modulo \mathbf{L}_f . This is a contradiction against the minimality of \mathbf{L}_{∞} and the claim is proved.

Fix $\gamma \in \mathcal{F}$ and (γ_{n_k}) with $\gamma_{n_k} \cdot \mathbf{v}_{\mathbf{L}_{\infty}} = \gamma \cdot \mathbf{v}_{\mathbf{L}_{\infty}}$. Then (δ_{n_k}) converges to $\delta_{\infty} \gamma^{-1}$ modulo $\gamma \mathbf{L}_{\infty} \gamma^{-1}$ or equivalently, $\gamma \mathbf{L}_{\infty} \gamma^{-1}$. Let $\mathcal{F}_{\gamma} \subset \gamma \mathbf{L}_{\infty} \gamma^{-1}$ be a finite subset such that every limit point of (δ_n) in $G/\gamma \mathbf{L}_{\infty}^{\mathrm{H}} \gamma^{-1}$ is equal to the image of δc for some $c \in \mathcal{F}_{\gamma}$. By Theorem 6.15, we have

$$\lim_{k \to \infty} \delta_{n_k} \gamma_{n_k} \mathbf{m}_{[\mathrm{H}]}^{\mathbf{1}} \in \left\{ (\delta_{\infty} c \gamma)_* \mathbf{m}_{[\mathrm{L}_{\infty}^{\mathrm{H}}]}^{\mathbf{1}} \mid c \in \mathcal{F}_{\gamma} \right\}.$$

Up to now, we have shown that whenever (x_n) converges to x_{∞} , all limit points of $\Psi_{\mathbf{H}}(x_n)$ are contained in the following finite set:

$$F := \left\{ (\delta_{\infty} c \gamma)_* \mathbf{m}^{\mathbf{1}}_{[\mathbf{L}_{\infty}^{\mathbf{H}}]} \mid \gamma \in \mathcal{F}, \ c \in \mathcal{F}_{\gamma} \right\}.$$

For each $\nu \in F$, there exists a neighborhood $\mathcal{N}_{\nu} \subset \operatorname{Prob}(G/\Gamma)$ such that $\mathcal{N}_{\nu} \cap \mathcal{N}_{\mu} = \emptyset$ for $\nu \neq \mu \in F$. By discussion above, there exists an open neighborhood $\mathcal{N}_{x_{\infty}}$ of x_{∞} in X^{cor} such that

$$\Psi_{\mathbf{H}}(\mathcal{N}_{x_{\infty}} \setminus \mathbf{D}) \subset \bigsqcup_{\nu \in F} \mathcal{N}_{\nu}.$$

By construction, there exists a smaller neighborhood $\mathcal{N}'_{x_{\infty}}$ of x_{∞} such that $\mathcal{N}'_{x_{\infty}} \setminus D$ is connected (this is the only place where the construction of manifolds with corners is needed!). Thus $\Psi_{\mathbf{H}}(\mathcal{N}'_{x_{\infty}} \setminus D) \subset \mathcal{N}_{\nu}$ for a unique $\nu \in F$. This shows that $\lim_{n\to\infty} \Psi_{\mathbf{H}}(x_n)$ exists and is equal to ν for all sequences (x_n) converging to x_{∞} . And the proof is now complete. 6.4.5. Measure compactifications, infinite volume cases. For this subsection, we assume that **G** and **H** are connected and reductive, and $\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ} \subset \mathbf{H}$. Here we are mainly interested in the case when $\mathbf{m}_{[\mathrm{H}]}$ is infinite.

By Theorem 4.5, for any arithmetic subgroup $\Gamma \subset G \cap \mathbf{G}(\mathbb{Q})$, there exists an open bounded set $\mathcal{B}_{\Gamma} \subset G$ such that $G = \mathcal{B}_{\Gamma} \cdot \Gamma \cdot H$. Therefore, for a positive function ψ whose support is large enough, one has $\langle \psi, \nu \rangle > 0$ for all ν of the form $g_*\mathfrak{m}_{[\mathbf{L}^\circ]}$ with $g \in G$ and $\mathbf{L} \in INT_{\Gamma}(\mathbf{H}, \mathbf{G})$. Fix such a ψ , define

$$\operatorname{Prob}^{\psi}(\mathcal{G}/\Gamma) := \left\{ \nu \in \operatorname{Meas}(\mathcal{G}/\Gamma) \mid \langle \psi, \nu \rangle = 1 \right\},$$
(22)

equipped with the weak-* topology where $\text{Meas}(G/\Gamma)$ denotes the set of locally finite Borel measures on G/Γ . We will be concerned with measures $\nu \in \text{Meas}(G/\Gamma)$ such that $\nu = \sum_{i=1}^{k} \nu_i$ where

- each supp ν_i is connected and supp $\nu_i \cap \text{supp } \nu_j = \emptyset$ for $i \neq j$;
- $\langle \psi, \nu_i \rangle \neq 0$ for all *i*.

For such a measure ν , let $\nu^{\psi} := k^{-1} \sum_{i=1}^{k} \nu_i / \langle \psi, \nu_i \rangle = k^{-1} \sum_{i=1}^{k} \nu_i^{\psi} \in \operatorname{Prob}^{\psi}(G/\Gamma).$

Remark 6.17. The definition is made so that Lemma 6.18 below holds. If $\mathbf{H}(\mathbb{R}) \to \mathbf{G}(\mathbb{R})/\mathbf{G}(\mathbb{R})^{\circ}$ is surjective, then we could have defined $\nu^{\psi} := \frac{\nu}{\langle \psi, \nu \rangle}$.

For $g \in G$, if both ν and $g_*\nu$ enjoy the above properties, we define $\alpha_g^{\psi}(\nu^{\psi}) := (g_*\nu)^{\psi} \in \operatorname{Prob}^{\psi}(G/\Gamma)$. Let

$$X_{\mathbf{H}}^{\mathrm{meas},\psi} := \overline{\alpha_{\mathrm{G}}^{\psi}\left(\mathrm{m}_{[\mathrm{H}]}^{\psi}\right)} \subset \mathrm{Prob}^{\psi}(\mathrm{G}/\Gamma).$$

Similar to Section 6.4.3, one has

$$\begin{split} \mathbf{m}^{\psi}_{[\mathbf{H}]} &= |I/\sim_1|^{-1} \sum_{[i] \in I/\sim_1} ((c^{\mathbf{H}}_i)_* \mathbf{m}_{[\mathbf{H}^\circ]})^{\psi}, \\ \mathbf{m}^{\psi}_{[\mathbf{L}^{\mathbf{H}}]} &= |I/\sim_{\mathbf{L}}|^{-1} \sum_{[i] \in I/\sim_{\mathbf{L}}} ((c^{\mathbf{H}}_i)_* \mathbf{m}_{[\mathbf{L}^\circ]})^{\psi}. \end{split}$$

Lemma 6.18. Let $(\gamma_n) \subset \Gamma^{\text{sma}}$ be a sequence such that $(\gamma_n \mathbf{H} \gamma_n^{-1})$ converges to $\mathbf{L} \in \text{INT}(\mathbf{H}, \mathbf{G})$, then $\lim_{n \to \infty} \alpha_{\gamma_n}^{\psi} \left(\mathbf{m}_{[\mathbf{H}]}^{\psi} \right) = \mathbf{m}_{[\mathbf{L}^{\mathbf{H}}]}^{\psi}$.

Proof. For every n, i, let $\gamma_n^i := (c_i^{\mathbf{H}})^{-1} \gamma_n c_i^{\mathbf{H}} \in \Gamma$. Then

$$(\gamma_n c_i^{\mathbf{H}})_* \mathbf{m}_{[\mathbf{H}^\circ]} = (c_i^{\mathbf{H}} \gamma_n^i)_* \mathbf{m}_{[\mathbf{H}^\circ]}.$$

Also, $\left(\gamma_n^i \mathbf{H}(\gamma_n^i)^{-1}\right)$ converges to $(c_i^{\mathbf{H}})^{-1} \mathbf{L} c_i^{\mathbf{H}}$, which is equal to \mathbf{L} as $\mathbf{H} \subset \mathbf{L}$. By Theorem 4.2 and 4.5, $\lim \alpha_{\gamma_n^i}^{\psi} \left(\mathbf{m}_{[\mathbf{H}^\circ]}^{\psi}\right) = \mathbf{m}_{[\mathbf{L}^\circ]}^{\psi}$ and so $\lim \alpha_{\gamma_n c_i^{\mathbf{H}}}^{\psi} \left(\mathbf{m}_{[\mathbf{H}^\circ]}^{\psi}\right) = \left((c_i^{\mathbf{H}})_* \mathbf{m}_{[\mathbf{L}^\circ]}\right)^{\psi}$. Consequently,

$$\lim_{n \to \infty} \alpha_{\gamma_n}^{\psi} \left(\mathbf{m}_{[\mathbf{H}]}^{\psi} \right) = |I/\sim_1|^{-1} \sum_{[i] \in I/\sim_1} \lim \left((\gamma_n c_i^{\mathbf{H}})_* \mathbf{m}_{[\mathbf{H}^\circ]} \right)^{\psi}$$
$$= |I/\sim_1|^{-1} \sum_{[i] \in I/\sim_1} \left((c_i^{\mathbf{H}})_* \mathbf{m}_{[\mathbf{L}^\circ]} \right)^{\psi}$$
$$= |I/\sim_{\mathbf{L}}|^{-1} \sum_{[i] \in I/\sim_{\mathbf{L}}} \left((c_i^{\mathbf{H}})_* \mathbf{m}_{[\mathbf{L}^\circ]} \right)^{\psi} = \mathbf{m}_{[\mathbf{L}^{\mathbf{H}}]}^{\psi}$$

Define an equivalence relation \sim on INT := INT_{Γ}(**H**, **G**) by

$$\mathbf{L}_1 \sim \mathbf{L}_2 \iff \mathbf{L}_1^{\mathrm{H}} \Gamma = g \mathbf{L}_2^{\mathrm{H}} \Gamma, \ \exists g \in \mathbf{G}$$

Lemma 6.19. We can decompose $X_{\mathbf{H}}^{\mathrm{meas},\psi} = \bigsqcup_{[\mathbf{L}] \in \mathrm{INT} / \sim} \alpha_{\mathrm{G}}^{\psi} \left(\mathbf{m}_{[\mathrm{L}^{\mathrm{H}}]}^{\psi} \right)$. In particular, the map $(g,\nu) \mapsto \alpha_{g}^{\psi}(\nu)$ defines a continuous action of G on $X_{\mathbf{H}}^{\mathrm{meas},\psi}$.

Proof. It suffices to show every $\nu \in X_{\mathbf{H}}^{\mathrm{meas},\psi}$ is equal to $\alpha_g^{\psi}\left(\mathbf{m}_{[\mathbf{L}^{\mathrm{H}}]}^{\psi}\right)$ for some $\mathbf{L} \in \mathrm{INT}_{\Gamma}(\mathbf{H},\mathbf{G})$ and $g \in \mathbf{G}$. Indeed, every sequence (g_n) in \mathbf{G} , after passing to a subsequence, can be written as $g_n = \delta_n \gamma_n h_n$ with (δ_n) converging to some $\delta_{\infty}, (\gamma_n) \subset \Gamma^{\mathrm{sma}}$ with $\gamma_n \mathbf{H} \gamma_n^{-1}$ converging to some $\mathbf{L} \in \mathrm{INT}$ and $(h_n) \subset \mathbf{H}$. By Lemma 6.18, $\lim \alpha_{g_n}^{\psi}\left(\mathbf{m}_{[\mathbf{H}]}^{\psi}\right) = \alpha_{\delta_{\infty}}^{\psi}\left(\mathbf{m}_{[\mathbf{L}^{\mathrm{H}}]}^{\psi}\right)$. So we are done.

Define $\Psi_{\mathbf{H}}^{\psi}(g.o) := \alpha_g^{\psi}\left(\mathbf{m}_{[\mathbf{H}]}^{\psi}\right)$. Let $\mathbf{X} := \widetilde{\mathbf{X}}_{\mathbf{H}}^{\mathrm{IMTP,cor}}$ and $\mathbf{D} := \widetilde{\mathbf{D}}_{\mathbf{H}}^{\mathrm{IMTP,cor}}$ as before. Similar to Theorem 6.16, we have:

Theorem 6.20. Assume that **G**, **H** are connected and reductive, $\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ} \subset \mathbf{H}$ and ψ is as above Equa.(22) where $\operatorname{Prob}^{\psi}(\mathbf{G}/\Gamma)$ is defined. Let (x_n) be a sequence in $X \setminus D$ converging to some $x_{\infty} \in X$. Then $\left(\Psi_{\mathbf{H}}^{\psi}(x_n)\right)$ is convergent in $\operatorname{Prob}^{\psi}(\mathbf{G}/\Gamma)$. Consequently, $\Psi_{\mathbf{H}}^{\psi}$ extends to a continuous map $\overline{\Psi}_{\mathbf{H}}^{\psi}$: $X \to \operatorname{Prob}^{\psi}(\mathbf{G}/\Gamma)$.

Note that conditions (F1), (F2), (C1) and (N1) hold under the assumptions made here.

6.5. Compactifications of U. Let \mathbf{U} be a variety over \mathbb{Q} equipped with a transitive \mathbf{G} -action and assume that there exists $o \in \mathbf{U}(\mathbb{Q})$ such that the stabilizer of o in \mathbf{G} is \mathbf{H} . Let \mathbf{H}_x denote the stabilizer of $x \in \mathbf{U}(\mathbb{Q})$. For different rational points, \mathbf{H}_x 's may not be isomorphic to each other. So there seems no canonical choice of \mathbf{H}_x to identify \mathbf{U} with \mathbf{G}/\mathbf{H}_x .

6.5.1. Conjugacy between intermediate subgroups. While \mathbf{H}_x and \mathbf{H}_y may not be isomorphic for different $x, y \in \mathbf{U}(\mathbb{Q})$, the set of intermediate groups are related.

Lemma 6.21. Let x, y be two rational points on $\mathbf{U}(\mathbb{Q})$. Find $f \in \mathbf{G}(\overline{\mathbb{Q}})$ such that fy = x. Assume \mathbf{L} is a \mathbb{Q} -subgroup normalized by \mathbf{H}_y . Then $f\mathbf{L}f^{-1}$ is defined over \mathbb{Q} . Therefore, $\mathbf{L} \to f\mathbf{L}f^{-1}$ induces bijections between $\mathrm{INT}_{\mathbb{Q}}(\mathbf{H}_y, \mathbf{G})$ and $\mathrm{INT}_{\mathbb{Q}}(\mathbf{H}_x, \mathbf{G})$; $\mathrm{INT}_{\mathbb{Q}}^{\mathrm{obs}}(\mathbf{H}_y, \mathbf{G})$ and $\mathrm{INT}_{\mathbb{Q}}^{\mathbb{Q}}(\mathbf{H}_x, \mathbf{G})$; $\mathcal{P}_{\mathbf{H}_y}$ and $\mathcal{P}_{\mathbf{H}_x}$.

Proof. For $\sigma \in \text{Gal}_{\mathbb{Q}}$, we have

$$fy = x = \sigma(x) = \sigma(fy) = \sigma(f)y \implies f^{-1}\sigma(f) \in \mathbf{H}_y.$$

Let $h_{\sigma} := f^{-1}\sigma(f)$, then $\sigma(f\mathbf{L}f^{-1}) = fh_{\sigma}\mathbf{L}h_{\sigma}^{-1}f^{-1} = f\mathbf{L}f^{-1}$ for all $\sigma \in \text{Gal}_{\mathbb{Q}}$. So $f\mathbf{L}f^{-1}$ is defined over \mathbb{Q} . The rest of the claim follows from this.

6.5.2. Convergent intermediate subgroups. However, it is not clear whether conjugation by f maps INT_Γ(\mathbf{H}_y, \mathbf{G}) to INT_Γ(\mathbf{H}_x, \mathbf{G}). Consequently, though Lemma 6.5 or its analogue for \mathbf{X}^{IMTP} is great for counting a single Γ-orbit, it needs to be modified for counting integral points, which may consist of different Γ-orbits. Define

$$\operatorname{INT}_{\mathbf{G}(\overline{\mathbb{Q}})}^{\operatorname{cvg}}(\mathbf{H},\mathbf{G}) = \left\{ \mathbf{L} \in \operatorname{INT}_{\mathbb{Q}}(\mathbf{H},\mathbf{G}) \mid \mathbf{L} = g\mathbf{L}'g^{-1}, \ \exists g \in \mathbf{G}(\overline{\mathbb{Q}}), \ \mathbf{L}' \in \operatorname{INT}_{\Gamma}(\mathbf{H},\mathbf{G}) \right\}.$$

Lemma 6.22. Assumptions same as in Lemma 6.21. Then $\mathbf{L} \to f\mathbf{L}f^{-1}$ induces bijections between $\operatorname{INT}_{\mathbf{G}(\overline{\mathbb{Q}})}^{\operatorname{cvg}}(\mathbf{H}_y, \mathbf{G})$ and $\operatorname{INT}_{\mathbf{G}(\overline{\mathbb{Q}})}^{\operatorname{cvg}}(\mathbf{H}_x, \mathbf{G})$.

Proof. This follows from Lemma 4.4 and Lemma 6.21 above.

We formulate a condition that guarantees the finiteness of this set:

(N2) $\mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\circ} \subset \mathbf{N}_{\mathbf{G}}(\mathbf{L})$ for every $\mathbf{L} \in \mathrm{INT}_{\mathbb{O}}^{\mathrm{obs}}(\mathbf{H},\mathbf{G})$.

Lemma 6.23. Assume that one of the following is true:

- (1) **G** and **H** are reductive and $\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ} \subset \mathbf{H}$;
- (2) **H** contains a maximal unipotent subgroup of \mathbf{G} .
- Then (N2) holds.

The example $\mathbf{SL}_n \times \mathbf{D}$ modulo the diagonal embedding of \mathbf{D} mentioned in Section 6.2.2 also satisfies this.

Proof. Part 1 is easy as $\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ} = \mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\circ}$. Part 2 follows from the structure of subgroups containing some maximal unipotent subgroup.

Lemma 6.24. Assume (C1), (F1) and (N2) hold, then $INT_{\mathbf{G}}^{cvg}(\mathbf{H}, \mathbf{G})$ is finite.

Proof. By (C1), for each $\mathbf{L} \in INT_{\Gamma}(\mathbf{H}, \mathbf{G})$, find finitely many elements $(f_i)_{i \in \mathscr{I}_{\mathbf{L}}}$ such that

$$\mathbf{Z}(\mathbf{H},\mathbf{L})(\overline{\mathbb{Q}}) = \bigsqcup_{i \in \mathscr{J}_{\mathbf{L}}} \mathbf{L}(\overline{\mathbb{Q}}) f_i \mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\circ}(\overline{\mathbb{Q}}).$$

For $g \in \mathbf{Z}(\mathbf{H}, \mathbf{L})(\overline{\mathbb{Q}})$ such that $g^{-1}\mathbf{L}g$ is defined over \mathbb{Q} , written as $l_g f_i n_g$ according to this decomposition,

$$g^{-1}\mathbf{L}g = n_g^{-1}(f_i^{-1}\mathbf{L}f_i)n_g = f_i^{-1}\mathbf{L}f_i$$

by (N2) applied to $n_g^{-1}f_i^{-1}\mathbf{L}f_in_g \in \mathrm{INT}^{\mathrm{obs}}_{\mathbb{Q}}(\mathbf{H}, \mathbf{G})$. Therefore,

$$\operatorname{INT}_{\mathbf{G}(\overline{\mathbb{Q}})}^{\operatorname{cvg}}(\mathbf{H},\mathbf{G}) \subset \left\{ f_i^{-1} \mathbf{L} f_i \mid \mathbf{L} \in \operatorname{INT}, \ i \in \mathscr{J}_{\mathbf{L}}, \ f_i^{-1} \mathbf{L} f_i \text{ is defined over } \mathbb{Q} \right\}$$

is finite.

6.5.3. Refined compactifications and condition (B1). We similarly define (abbreviate $\operatorname{INT}_{\mathbf{G}(\overline{\mathbb{Q}})}^{\operatorname{cvg}} := \operatorname{INT}_{\mathbf{G}(\overline{\mathbb{Q}})}^{\operatorname{cvg}}(\mathbf{H}, \mathbf{G})$)

$$\mathbf{X}_{\mathbf{H}}^{\mathrm{IMTP}'} \subset \prod_{\mathbf{L} \in \mathrm{INT}_{\mathbf{G}(\overline{\mathbb{Q}})}^{\mathrm{evg}} \sqcup \mathscr{V}(\mathbf{H})} \mathbf{P}(\mathbf{V}_{\mathbf{L}} \oplus \mathbb{Q})$$
(23)

and

$$\mathbf{X}_{\mathbf{H}}^{\mathrm{INTP}'} \subset \mathbf{P}(\mathbf{V}_{\mathbf{H}} \oplus \mathbb{Q}) \times \prod_{\mathbf{L} \in \mathrm{INT}_{\mathbf{G}(\mathbb{Q})}^{\mathrm{evg}}} \mathbf{P}(\mathbf{V}_{\mathfrak{l}} \oplus \mathbb{Q}) \times \prod_{\rho \in \mathscr{V}(\mathbf{H})} \mathbf{P}(\mathbf{V}_{\rho} \oplus \mathbb{Q})$$
(24)

which admit natural morphisms onto $\mathbf{X}_{\mathbf{H}}^{\mathrm{IMTP}}$ or $\mathbf{X}^{\mathrm{INTP}}$. In defining $\mathbf{B}_{\mathbf{H}}^{\mathrm{IMTP}'}$, we replace the index $\mathrm{INT}_{\Gamma}(\mathbf{H}, \mathbf{G})$ by the larger $\mathrm{INT}_{\mathbf{G}(\overline{\mathbb{Q}})}^{\mathrm{cvg}}(\mathbf{H}, \mathbf{G})$. For simplicity refer $\left(\widetilde{\mathbf{X}}_{\mathbf{H}}^{\mathrm{IMTP}'}, \widetilde{\mathbf{D}}_{\mathbf{H}}^{\mathrm{IMTP}'}, \widetilde{\mathbf{B}}_{\mathbf{H}}^{\mathrm{IMTP}'}\right)$ as the log smooth IMTP' compactification of U. Log smooth INTP' compactification is similarly defined. We have the following:

Lemma 6.25. Consider one of the following two situations:

- (1) Assume (F1), (F2), (C1), (N1) and (N2) hold. Let (**X**, **D**, **B**) be the log smooth IMTP' compactification;
- (2) Assume that the unipotent radical of **G** commutes with **G**. Assume (F1), (F2), (C1), (N2) hold and that the modular character of **L** is trivial on **H** for all $\mathbf{L} \in INT^{cvg}_{\mathbf{G}(\overline{\mathbb{Q}})}(\mathbf{H}, \mathbf{G})$. Let $(\mathbf{X}, \mathbf{D}, \mathbf{B})$ be the log smooth INTP' compactification.

Let $x \in \mathbf{U}(\mathbb{Q})$ and $(g_n) \subset \mathbf{G}$ be such that $(g_n \cdot x)$ converges to some $\mathbf{v} \in \mathbf{X}$. Then

$$\lim_{n \to \infty} \left[(g_n)_* \mathbf{m}_{[\mathbf{H}_x^\circ]} \right] = \left[\mathbf{m}_{[\mathbf{G}]} \right] \iff \mathbf{v} \in \mathbf{B}.$$

In particular, condition (B1) holds for $(\mathbf{X}, \mathbf{D}, \mathbf{B})$.

Proof. The proof is similar to that of Lemme 6.5 and is omitted.

6.6. Wrap-up. By combining efforts made so far, we have arrived at a few theorems.

Theorem 6.26. Consider one of the following two situations:

- (1) Assume (F1), (F2), (C1), (N1) and (N2) hold. Let (**X**, **D**, **B**) be the log smooth IMTP' compactification;
- (2) Assume that the unipotent radical of **G** commutes with **G**. Assume (F1), (F2), (C1), (N2) hold and that the modular character of **L** is trivial on **H** for all $\mathbf{L} \in INT^{cvg}_{\mathbf{G}(\overline{\mathbb{Q}})}(\mathbf{H}, \mathbf{G})$. Let $(\mathbf{X}, \mathbf{D}, \mathbf{B})$ be the log smooth INTP' compactification.

Also, assume equivalent conditions in Lemma 4.4 hold for $\mathbf{H}_x \subset \mathbf{G}$ for all $x \in \mathbf{U}(\mathbb{Q})$. Then condition (B4) holds for $(\mathbf{X}, \mathbf{D}, \mathbf{B})$. Therefore, there exists \mathbf{L} supported on \mathbf{D} such that $\operatorname{Ht}_{\mathbf{L}}$ is good if $\operatorname{Pic}(\mathbf{U})$ is torsion and is good with weights in general.

□ :=

Proof. By assumption, there exists $(\gamma_n) \subset \Gamma$ that is unbounded modulo the normalizer of \mathbf{H}_x . So Condition (B4) holds by applying Theorem 2.14. Lemma 5.3 shows the existence of height satisfying (BH1). (B1) holds by Lemma 6.25. Lemma 5.1 then concludes the proof.

Theorem 6.27. Assume **G** and **H** are connected and reductive and $\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ} \subset \mathbf{H}$. Also, assume that the projection of **H** to the compact \mathbb{Q} -factor of **G** is surjective. Then (B2) and (K1) hold for $(\mathbf{X}, \mathbf{D}, \mathbf{B})$, the log smooth IMTP' compactification. In particular, the log anti-canonical height is good if Pic(U) is torsion and is good with weights in general. Every other height is ok if $\mathbf{m}_{[\mathbf{H}]}$ is finite and is ok with weights otherwise.

Note that under these assumptions, (F1), (F2), (C1), (N1), (N2) and equivalent conditions in Lemma 4.4 hold for \mathbf{H}_x for all $x \in \mathbf{U}(\mathbb{Q})$.

Proof. Condition (K1) follows from Theorem 2.13. By Theorem 4.5, every limit measure μ is non-zero and hence homogeneous. By Lemma 4.4, there exists $(\gamma_n) \subset \Gamma$ such that $\left[(\gamma_n)_* \mathbf{m}_{[\mathbf{m}_{H_x^o}]}\right]$ converges to $\left[\mathbf{m}_{[G]}\right]$ for every $x \in \mathbf{U}(\mathbb{Q})$. Hence (B2) holds. Lemma 5.2 gives condition (BH1) and Lemma 6.25 gives (B1). So we are done by invoking Lemma 5.1.

If $m_{[H]}$ is finite, then **H** and **G** have no non-trivial Q-characters under our assumption. Also, thanks to Theorem 6.16 and 4.5, condition (S1) and (D1) hold, which implies (H1) by Lemma 5.5. Then one can conclude with Lemma 5.4.

In the infinite-volume case, apply Theorem 6.20 and 4.5 to get (S1) and (D2). Then we get (H2) by Lemma 5.8. Finally one concludes with Lemma 5.7.

6.7. Lift equidistributions. We would like to extend Theorem 6.26 to a more general setting. This is necessary to prove Theorem 1.2 in full generality.

6.7.1. Standing assumptions. Take **G** to be a connected linear algebraic group over \mathbb{Q} , **U** to be a homogeneous variety under **G** with a point $o \in \mathbf{U}(\mathbb{Q})$ and **H** to be the stabilizer of o in **G**. Assume

Assumption 6.28. $\mathbf{R}_{\mathbf{u}}(\mathbf{G})$ commutes with the reductive part of \mathbf{G} . Namely, $\mathbf{G} = \mathbf{G}^{\mathrm{red}} \times \mathbf{R}_{\mathbf{u}}(\mathbf{G})$.

Thus **G** can be written as $(\mathbf{G}^{\text{cpt}} \cdot \mathbf{G}^{\text{nc}} \cdot \mathbf{Z}(\mathbf{G})) \times \mathbf{R}_{\mathbf{u}}(\mathbf{G})$. Under these assumptions, here are a few natural quotient morphisms:

• $p^{\text{stau}}: \mathbf{G} \to \mathbf{G}/\mathbf{G}^{\text{nc}}; p_1: \mathbf{G} \to \mathbf{G}_1 := \mathbf{G}/\mathbf{Z}(\mathbf{G})^{\text{spl}}, p_{12}: \mathbf{G}_1 \to \mathbf{G}_2 := \mathbf{G}/(\mathbf{G}^{\text{cpt}}\cdot\mathbf{Z}(\mathbf{G})) \times \mathbf{R}_{\mathbf{u}}(\mathbf{G}),$ and $p_2: \mathbf{G} \to \mathbf{G}_2$ is the composition $p_{12} \circ p_1$.

Regarding \mathbf{H} , we assume that

Assumption 6.29. H is a connected observable \mathbb{Q} -subgroup of G, p^{stau} is surjective restricted to H, $p_1(\mathbf{H})$ is observable and $\mathscr{P}_{\mathbf{H}}^{\max}$ is finite.

Note that under this assumption, \mathbf{G}/\mathbf{H} is a homogeneous space under \mathbf{G}^{nc} . However, we get some advantage by considering the larger automorphism subgroup. Also, since every parabolic Q-subgroup of \mathbf{G} must contain $(\mathbf{G}^{\mathrm{cpt}} \cdot \mathbf{Z}(\mathbf{G})) \times \mathbf{R}_{\mathbf{u}}(\mathbf{G})$, thus $\mathscr{P}_{\mathbf{H}}^{\mathrm{max}}$ being finite is equivalent to $\mathscr{P}_{\mathbf{H}_1}^{\mathrm{max}}$ being finite, which is again equivalent to $\mathscr{P}_{\mathbf{H}_2}^{\mathrm{max}}$ being finite.

Remark 6.30. By Lemma 4.4, p^{stau} being surjective is a necessary assumption for the existence of $(\gamma_n) \subset \Gamma$ such that $[(\gamma_n)_* \mathbf{m}_{[\mathbf{H}^\circ]}]$ converges to $[\mathbf{m}_{[\mathbf{G}]}]$ since maximal proper subgroups of a unipotent group is normal. Of course, when \mathbf{G}^{red} acts non-trivially on $\mathbf{R}_{\mathbf{u}}(\mathbf{G})$, this is no longer necessary.

Recall that the stabilizer of a general point $x \in \mathbf{U}(\mathbb{Q})$ in **G** is denoted as \mathbf{H}_x . Its image in \mathbf{G}_i is denoted as $\mathbf{H}_{x,i}$ for i = 1, 2. We have assumed $\mathbf{H}_1 := p_1(\mathbf{H})$ to be observable, thus $\mathbf{H}_2 := p_2(\mathbf{H})$ is also observable (see Lemma 6.37) and $\mathbf{m}_{[\mathbf{H}_x^\circ]}$ makes sense.

Assumption 6.31. There exists (and we fix such) a G_2 -pair (X, B) over \mathbb{Q} with X being a G-equivariant compactification of G_2/H_2 such that

1. for every $x \in (\mathbf{G}_2/\mathbf{H}_2)(\mathbb{Q})$, the analytic closure of $\mathbf{G}_2.x$ intersects with $\mathbf{B}(\mathbb{R})$;

2. for every $x \in (\mathbf{G}_2/\mathbf{H}_2)(\mathbb{Q})$ and $(g_n) \subset \mathbf{G}_2$ such that every limit point of $(g_n.x)$ is contained in $\mathbf{B}(\mathbb{R})$, we have

$$\lim_{n \to \infty} \left[(g_n)_* \mathbf{m}_{[\mathbf{H}_{2,x}^\circ]} \right] = \left[\mathbf{m}_{[\mathbf{G}_2]} \right]$$

where $\mathbf{H}_{2,x}$ denotes the stabilizer of x in \mathbf{G}_2 .

The compactification of $\mathbf{G}_2/\mathbf{H}_2$ as in the assumption can be easily upgraded to one of $\mathbf{U} \cong \mathbf{G}/\mathbf{H}$ with the same conclusion. Namely, for every $x \in \mathbf{U}(\mathbb{Q})$ and every $(g_n) \subset \mathbf{G}$ with $\lim g_n . x \in \mathbf{B}(\mathbb{R})$, one has $\lim_{n\to\infty} \left[p_2(g_n)_* \mathbf{m}_{[\mathbf{H}_{x,2}^o]} \right] = \left[\mathbf{m}_{[\mathbf{G}_2]} \right]$ and such a sequence (g_n) does exist. Note that if ϕ is the quotient morphism $\mathbf{G}/\mathbf{H} \to \mathbf{G}_2/\mathbf{H}_2$ and $x \in \mathbf{G}/\mathbf{H}(\mathbb{Q})$, then $\mathbf{H}_{x,2} = \mathbf{H}_{2,\phi(x)}$.

• Let Γ be an arithmetic subgroup of **G** and $\Gamma_i := p_i(\Gamma)$.

Note that Γ_i is an arithmetic subgroup of \mathbf{G}_i for i = 1, 2 (see [PRR23, Theorem 4.7]).

6.7.2. Main theorem and outline of the proof.

Theorem 6.32. There exist a smooth **G**-pair (\mathbf{X}, \mathbf{D}) over \mathbb{Q} such that $\mathbf{U} := \mathbf{X} \setminus \mathbf{D}$ is equivariantly isomorphic to \mathbf{G}/\mathbf{H} over \mathbb{Q} , and $\mathbf{B} \subset \mathbf{D}$, a union of irreducible components of \mathbf{D} defined over \mathbb{Q} , such that (B1) and (B4) hold.

Proof of Theorem 6.32. We explain that Theorem 6.32 follows from Lemma 6.35 and 6.39 below. We fix some $x \in \mathbf{U}(\mathbb{Q})$.

Let $(\mathbf{X}_1, \mathbf{D}_1, \mathbf{B}_1)$ be the one obtained by Lemma 6.39. $(\mathbf{X}_3, \mathbf{B}_3)$ is constructed by firstly applying Theorem 6.11 to $(\mathbf{X}_1, \mathbf{B}_1)$ to get $(\mathbf{X}_2, \mathbf{D}_2, \mathbf{B}_2)$ $(\mathbf{D}_2$ is the inverse image of \mathbf{D}_1). Apply Theorem 6.11 again to $(\mathbf{X}_2, \mathbf{D}_2)$ to get $(\mathbf{X}_3, \mathbf{D}_3, \mathbf{B}_3)$ where \mathbf{B}_3 is the inverse image of \mathbf{B}_2 . Thus $(\mathbf{X}_3, \mathbf{D}_3)$ is a smooth **G**-pair over \mathbb{Q} and \mathbf{B}_3 is a union of irreducible components of \mathbf{D}_3 that is over \mathbb{Q} .

Now assume that $(g_n.x)$ converges to a point in $\mathbf{B}_3(\mathbb{R})$ for some $(g_n) \subset \mathbf{G}$. By Lemma 6.39, we find $(h_n) \subset \mathbf{H}_x^\circ$, a bounded sequence (b_n) in \mathbf{G} and $(\gamma_n) \subset \Gamma$ such that $g_n h_n \omega_n = b_n \gamma_n$ and

$$\|\operatorname{Ad}(p_1(\gamma_n))\mathbf{v}_{\mathbf{P}}\| \to \infty, \quad \forall \, \mathbf{P} \in \mathscr{P}_{\mathbf{H}_{x,1}}^{\max}.$$

By Lemma 6.39 again,

$$\lim_{n \to \infty} \left[p_1(\gamma_n)_* \mathbf{m}_{[\mathbf{H}_{x,1}^\circ]} \right] = \left[\mathbf{m}_{[\mathbf{G}_1]} \right].$$

Hence $(p_1(\gamma_n)\mathbf{H}_{x,1}p_1(\gamma_n)^{-1})$ converges to \mathbf{G}_1 . Passing to a subsequence if necessary, assume that $(p_1(\gamma_n)\mathbf{H}_{x,1}p_1(\gamma_n)^{-1})$ strongly converges (see the next subsection for the definition) to some \mathbf{L} . Then by Lemma 6.34, \mathbf{L} is observable in \mathbf{G}_1 and hence $\mathbf{L} = \mathbf{G}_1$. Invoking Lemma 6.35, we have that $(\gamma_n \mathbf{H}_x \gamma_n^{-1})$ strongly converges to \mathbf{G} , showing that

$$\lim_{n \to \infty} \left[(\gamma_n)_* \mathbf{m}_{[\mathbf{H}_x^\circ]} \right] = \lim_{n \to \infty} \left[(g_n)_* \mathbf{m}_{[\mathbf{H}_x^\circ]} \right] = \left[\mathbf{m}_{[\mathbf{G}]} \right].$$

This proves condition (B1).

By Lemma 6.39, we can find $(\gamma_n) \subset \Gamma$ with $\lim \gamma_n . x \in \mathbf{B}_3(\mathbb{R})$. Thus, some irreducible component B of $\mathbf{B}_3(\mathbb{R})$ is contained in $\overline{\mathbf{G}.x}$. So we have some sequence $(g'_n) \subset \mathbf{G}$ such that $g'_n . x$ converges to B° (the complement in B of other boundary components). By (B1), $\lim [(g'_n)_* \mathbf{m}_{[\mathbf{H}_x^\circ]}] = [\mathbf{m}_{[\mathbf{G}]}]$. In particular, $g'_n . x = b_n \gamma'_n . x$ for some bounded (b_n) in \mathbf{G} and $(\gamma'_n) \subset \Gamma$. Thus we are done by Theorem 2.14 since (γ'_n) can not be bounded modulo the normalizer of \mathbf{H} . This proves (B4).

6.7.3. Strongly convergence and observability. Let us introduce the notion of strongly convergence and a simple criterion of observability. Strongly convergence is indeed stronger than convergence from Definition 4.1.

Definition 6.33. Let (\mathbf{A}_n) be a sequence of connected \mathbb{Q} -subgroups of a connected linear algebraic group \mathbf{C} over \mathbb{Q} and \mathbf{E} be another connected \mathbb{Q} -subgroup of \mathbf{C} . We say that (\mathbf{A}_n) strongly converges to \mathbf{E} if

- 1. every \mathbf{A}_n is contained in \mathbf{E} ;
- 2. for every subsequence (\mathbf{A}_{n_k}) , \mathbf{E} is the smallest \mathbb{Q} -subgroup containing all \mathbf{A}_{n_k} 's.

Lemma 6.34. Assume that \mathbf{A} is a connected linear algebraic group over \mathbb{Q} with no nontrivial \mathbb{Q} -characters and \mathbf{C} is a connected observable \mathbb{Q} -subgroup. Let (γ_n) be a sequence in some fixed arithmetic subgroup of \mathbf{A} and \mathbf{E} be a connected \mathbb{Q} -subgroup of \mathbf{A} such that

- (1) $\inf_{\mathbf{P}\in\mathscr{P}_{\mathbf{C}}^{\max}} \|\operatorname{Ad}(\gamma_n)\mathbf{v}_{\mathbf{P}}\| \to \infty;$ (2) $(\gamma_n \mathbf{C} \gamma_n^{-1})$ strongly converges to \mathbf{E} .

Then \mathbf{E} is observable in \mathbf{A} .

Proof. If **E** is contained some maximal proper parabolic \mathbb{Q} subgroup **P**, then **C** is contained in $\mathbf{P}_n := \gamma_n^{-1} \mathbf{P} \gamma_n$ for all *n*. Thus $\|\operatorname{Ad}(\gamma_n) \mathbf{v}_{\mathbf{P}_n}\| \simeq \|\operatorname{Ad}(\gamma_n \gamma_n^{-1}) \mathbf{v}_{\mathbf{P}}\|$ is bounded, contradicting our first assumption.

So **E**, and hence $\mathbf{ER}_{\mathbf{u}}(\mathbf{A})$ are not contained in any parabolic \mathbb{Q} -subgroup. Thus $\mathbf{ER}_{\mathbf{u}}(\mathbf{A})/\mathbf{R}_{\mathbf{u}}(\mathbf{A}) \subset \mathbb{Q}$ $\mathbf{A}/\mathbf{R}_{\mathbf{u}}(\mathbf{A})$ is reductive and hence observable. Equivalently, $\mathbf{ER}_{\mathbf{u}}(\mathbf{A})$ is observable in \mathbf{A} . But \mathbf{E} is observable in $\mathbf{ER}_{\mathbf{u}}(\mathbf{A})$ since $\mathbf{ER}_{\mathbf{u}}(\mathbf{A})/\mathbf{E}$ is a homogeneous space under a unipotent group and hence affine. So we are done.

6.7.4. Lift strong convergence. Strongly convergence, unlike convergence, is liftable.

Lemma 6.35. Let (γ_n) be a sequence in Γ and (\mathbf{L}_n) be a sequence of connected observable \mathbb{Q} -subgroups of **G**. Assume that p^{stau} is surjective restricted to any \mathbf{L}_n . The followings are equivalent:

- (1) (\mathbf{L}_n) strongly converges to \mathbf{G} ;
- (2) $(p_1(\mathbf{L}_n))$ strongly converges to \mathbf{G}_1 ;
- (3) $(p_2(\mathbf{L}_n))$ strongly converges to \mathbf{G}_2 .

Proof. The nontrivial implication is $3. \implies 1$.

By passing to a subsequence if necessary, we assume that (\mathbf{L}_n) strongly converges to \mathbf{L} . Thus $\mathbf{G}_2 = p_2(\mathbf{L})$. Write $\mathbf{L} = (\mathbf{L}^{\mathrm{nc}} \cdot \mathbf{L}^{\mathrm{cpt}}) \cdot \mathbf{R}(\mathbf{L})$. Then the image of $\mathbf{R}(\mathbf{L})$ under p_2 is contained in the radical of $p_2(\mathbf{L})$, which is trivial. Similarly, \mathbf{L}^{cpt} is contained in the kernel of p_2 . Thus, the subgroup \mathbf{L}^{nc} of \mathbf{G}^{nc} must be the full \mathbf{G}^{nc} . In particular, \mathbf{L} contains the kernel of p^{stau} . But p^{stau} is surjective restricted to \mathbf{L}_n and hence to \mathbf{L} , implying that \mathbf{L} is equal to \mathbf{G} .

 \square

Remark 6.36. The analogous statement is wrong replacing "strongly converges" by "converges". For instance, let $\mathbf{G} := \mathbf{SL}_2 \times \mathbf{G}_m$, \mathbf{G}_m be embedded in \mathbf{SL}_2 as the diagonal torus, and \mathbf{H} be the diagonal embedding of \mathbf{G}_m in \mathbf{G} . So here $p_1 = p_2 : \mathbf{G} \to \mathbf{SL}_2 = \mathbf{G}_2$ are the natural projection. Let \mathbf{N} be the upper triangular unipotent subgroup of \mathbf{SL}_2 and let $\mathbf{N}' := \mathbf{N} \times \{\mathrm{id}\}$ in \mathbf{G} . Take $\gamma_n := \left(\begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}, \mathrm{id} \right)$. Then one can verify that $\mathbf{H} \cdot \mathbf{N}'$ is an observable subgroup of \mathbf{G} and $(\gamma_n \mathbf{H} \gamma_n^{-1})$ converges to $\mathbf{H} \cdot$ N'. However, $p_2(HN')$, being the upper triangular Borel subgroup, is not observable in SL_2 and $(p_2(\gamma_n \mathbf{H} \gamma_n^{-1}))$ actually converges to \mathbf{SL}_2 .

6.7.5. Lift equidistribution through compact fibrations.

Lemma 6.37. Let A be a linear algebraic group over \mathbb{Q} , B be a \mathbb{Q} -anisotropic normal subgroup of A and $\pi: \mathbf{A} \to \mathbf{A}/\mathbf{B} =: \overline{\mathbf{A}}$ be the natural quotient. Let \mathbf{C} be a \mathbb{Q} -subgroup of \mathbf{A} . If \mathbf{C} is observable in **A**, then $\pi(\mathbf{C})$ is observable in $\overline{\mathbf{A}}$.

Proof. Let Γ be an arithmetic subgroup of **A** and $\overline{\pi}$ be the induced quotient map $\mathbf{A}(\mathbb{R})/\Gamma \to \overline{\mathbf{A}}(\mathbb{R})/\overline{\Gamma}$ where $\overline{\Gamma} := \pi(\Gamma)$ is an arithmetic subgroup of $\overline{\mathbf{A}}$. By [Wei98, Corollary 7], it is sufficient to show that $\pi(\mathbf{C})(\mathbb{R})\overline{\Gamma}/\overline{\Gamma}$, or equivalently, $\pi(\mathbf{C}(\mathbb{R}))\overline{\Gamma}/\overline{\Gamma}$ is closed in $\overline{\mathbf{A}}(\mathbb{R})/\overline{\Gamma}$. This follows from the properness of $\overline{\pi}$.

The equivalence between the convergence when applying p_{12} does hold, which leads to the following:

Lemma 6.38. Let (g_n) be a sequence in G and $x \in \mathbf{U}(\mathbb{Q})$. If $\left[p_2(g_n)_* \mathbf{m}_{[\mathrm{H}_{x,2}^\circ]}\right]$ converges to $[\mathbf{m}_{[\mathrm{G}_2]}]$, then $\left[p_1(g_n)_* \mathbf{m}_{[\mathbf{H}_{x,1}^\circ]}\right]$ converges to $\left[\mathbf{m}_{[\mathbf{G}_1]}\right]$.

Proof. As p_{12} is proper, $p_1(g_n) H_{x_1}^{\circ} \Gamma_1 / \Gamma_1$ intersects with some bounded set for all n. We may assume $p_1(g_n) = \lambda_n \in \Gamma_1$. So $(p_{12}(\lambda_n \mathbf{H}_{x,1} \lambda_n^{-1}))$ converges to \mathbf{G}_2 . After passing to a subsequence assume that $(\lambda_n \mathbf{H}_{x,1} \lambda_n^{-1})$ converges to **L**, an observable Q-subgroup of **G**₁. By Lemma 6.37, $p_{12}(\mathbf{L})$ is observable and has to be equal to \mathbf{G}_2 . But $\mathbf{LG}_1^{nc} = \mathbf{G}_1^{nc}$ by assumption. Hence \mathbf{L} has to be \mathbf{G}_1 by arguments similar to those in the proof of Lemma 6.35. And we conclude by invoking Theorem 4.2.

6.7.6. The general case. For $x \in \mathbf{U}(\mathbb{Q})$, fix a non-empty open bounded subset $\Omega_x \subset \mathrm{H}^*_{\mathbf{x}} \Gamma / \Gamma$. Up to now, it only remains to prove the following:

Lemma 6.39. There exist a **G**-pair (\mathbf{X}, \mathbf{D}) over \mathbb{Q} and a closed **G**-invariant \mathbb{Q} -subvariety **B** of **D** with $\mathbf{X} \setminus \mathbf{D}$ being **G**-equivariantly isomorphic to **U** such that the following holds. For every $x \in \mathbf{U}(\mathbb{Q})$, the analytic closure of G.x intersects with $\mathbf{B}(\mathbb{R})$. Moreover, for every $x \in \mathbf{U}(\mathbb{Q})$ and $(g_n) \subset \mathbf{G}$ such that every limit point of $(g_n x)$ is contained in $\mathbf{B}(\mathbb{R})$, we have

- (1) $\lim_{n\to\infty} \left| (p_1(g_n))_* \mathbf{m}_{[\mathbf{H}_{x,1}^\circ]} \right| = \left[\mathbf{m}_{[\mathbf{G}_1]} \right];$
- (2) there exists a sequence (h_n) in H_x° such that (2.1) $(g_n h_n \Omega_x)$ intersects with some bounded subset of G/Γ for all n; (2.2) for every $\mathbf{P} \in \mathscr{P}_{\mathbf{H}_{x,1}}^{\max}$, $\|\operatorname{Ad}(p_1(g_nh_n))\mathbf{v}_{\mathbf{P}}\| \to \infty$.

6.7.7. Splitting of parabolic characters. Let $\mathbf{M}_{\mathbf{H}} := (\mathbf{H} \cap (\mathbf{G}^{ss}\mathbf{Z}(\mathbf{G})^{an}\mathbf{R}_{\mathbf{u}}(\mathbf{G})))^{\circ}$. Then $\mathbf{H}/\mathbf{M}_{\mathbf{H}}$ is a Q-split torus.

Let $\mathbf{M}_{\mathbf{H},1} := p_1(\mathbf{M}_{\mathbf{H}})$, then $\mathbf{H}/\mathbf{M}_{\mathbf{H}}$ naturally surjects onto $\mathbf{H}_1/\mathbf{M}_{\mathbf{H},1}$. Therefore, $\mathbf{H}_1/\mathbf{M}_{\mathbf{H},1}$ is also a Q-split torus. So there exists a Q-split subtorus $\widetilde{\mathbf{S}}_1^{\mathbf{H}}$ of \mathbf{H}_1 that maps to $\mathbf{H}_1/\mathbf{M}_{\mathbf{H},1}$ surjectively with finite kernel. Choose another \mathbb{Q} -split subtorus $\mathbf{S}_{\mathbf{H}}^1$ of $\mathbf{M}_{\mathbf{H},1}$ such that the restriction of $p_{\mathbf{H}_1}^{\mathrm{spl}}$ to $\widetilde{\mathbf{S}}_{\mathbf{H}_1} := \mathbf{S}_{\mathbf{H}}^1 \cdot \widetilde{\mathbf{S}}_1^{\mathbf{H}}$ is surjective onto $\mathbf{S}_{\mathbf{H}_1}$ with finite kernel. By definition

$$\operatorname{Lie}(\widetilde{\mathbf{S}}_{\mathbf{H}_1}) = \operatorname{Lie}(\mathbf{S}_{\mathbf{H}}^1) \oplus \operatorname{Lie}(\widetilde{\mathbf{S}}_1^{\mathbf{H}}).$$

Note that the image of $\mathbf{M}_{\mathbf{H},1}$ under $p_{\mathbf{H}_1}^{\mathrm{spl}}$ is the same as that of $\mathbf{S}_{\mathbf{H}}^1$: certainly it contains $p_{\mathbf{H}_1}^{\mathrm{spl}}(\mathbf{S}_{\mathbf{H}}^1)$, but it can not be strictly larger as it has to be disjoint from the image of $\widetilde{\mathbf{S}}_{1}^{\mathbf{n}}$.

For a Q-eigenvector **v** of **H**₁, let $l_{\mathbf{v}}$ (resp. $\alpha_{\mathbf{v}}$) be the associated linear functional (resp. character) on $\operatorname{Lie}(\mathbf{S}_{\mathbf{H}_1})$ (resp. **H**).

Let

$$\mathscr{P}_{\mathbf{H}_{1}}^{+} := \left\{ \otimes^{\mathbf{n}_{\mathbf{P}}} \mathbf{v}_{\mathbf{P}} \mid \mathbf{n}_{\bullet} \in \operatorname{Map}^{*}(\mathscr{P}_{\mathbf{H}_{1}}^{\max}, \mathbb{Z}_{\geq 0}) \right\}$$
(25)

where $\operatorname{Map}^*(\mathscr{P}_{\mathbf{H}_1}^{\max}, \mathbb{Z}_{\geq 0})$ denotes the collection of set-theoretic maps from $\mathscr{P}_{\mathbf{H}_1}^{\max}$ to $\mathbb{Z}_{\geq 0}$ excluding the

zero map. Note that $\mathscr{P}_{\mathbf{H}_{1}}^{\max}$ can be naturally viewed as a subset of $\mathscr{P}_{\mathbf{H}_{1}}^{+}$. For a subset \mathscr{P} of $\mathscr{P}_{\mathbf{H}_{1}}^{+}$, let the corresponding $\mathcal{P} := \{l_{\mathbf{v}}, \mathbf{v} \in \mathscr{P}\}$. A subset of a Q-vector space is said to be *non-degenerate* iff $\mathbb{Q}_{>0}$ -linear combinations of this subset equal to the whole space.

Lemma 6.40. Let A be a connected linear algebraic group over \mathbb{Q} with no non-trivial \mathbb{Q} -characters. Let **B** be an observable \mathbb{Q} -subgroup of **A**. Define $\mathscr{P}^+_{\mathbf{A}}$ in the same way as $\mathscr{P}^+_{\mathbf{H}_1}$ above. And \mathcal{P} corresponding to some $\mathscr{P} \subset \mathscr{P}^+_{\mathbf{B}}$ is viewed as a subset of $\operatorname{Lie}(\mathbf{S}_{\mathbf{B}})^{\vee}$ here. Then $\mathcal{P}_{\mathbf{B}}$, corresponding to $\mathscr{P}^{\max}_{\mathbf{B}}$, is nondegenerate.

Before presenting the proof, we firstly deduce that

Lemma 6.41. There exists a finite subset $\mathscr{P} \subset \mathscr{P}^+_{\mathbf{H}_1}$ such that \mathcal{P} is non-degenerate and for each $\mathbf{v} \in \mathscr{P}$, either Lie($\mathbf{S}_{\mathbf{H}}^1$) $\subset \ker(l_{\mathbf{v}})$ or Lie($\widetilde{\mathbf{S}}_1^{\mathbf{H}}$) $\subset \ker(l_{\mathbf{v}})$.

Proof. Indeed, it is easy to find a finite subset of $\operatorname{Lie}(\widetilde{\mathbf{S}}_{\mathbf{H}_1})^{\vee}$ satisfying the required properties. By Lemma 6.40, replacing this subset by suitable \mathbb{Z}^+ -multiples one gets a subset of $\mathcal{P}_{H_1}^+$ without losing any required properties.

Henceforth, we fix such a choice of $\mathscr{P}_o = \mathscr{P}_o^1 \sqcup \mathscr{P}_o^2$ where

$$\mathcal{P}_{o}^{1} := \left\{ \mathbf{v} \in \mathcal{P}_{o}, \operatorname{Lie}(\mathbf{S}_{\mathbf{H}}^{1}) \subset \ker(l_{\mathbf{v}}) \right\},$$

$$\mathcal{P}_{o}^{2} := \left\{ \mathbf{v} \in \mathcal{P}_{o}, \operatorname{Lie}(\widetilde{\mathbf{S}}_{1}^{\mathbf{H}}) \subset \ker(l_{\mathbf{v}}) \right\}.$$
(26)

Replacing elements $\mathbf{v} \in \mathscr{P}_o$ by $\mathbf{v}^{\otimes 2}$ if necessary, we assume that each $\alpha_{\mathbf{v}} = \beta^2$ for some \mathbb{Q} -character β of **H**. Note that for $\mathbf{v} \in \mathscr{P}_o^1$, the character $\alpha_{\mathbf{v}}$ vanishes on $\mathbf{M}_{\mathbf{H}}$ (since it factors through $p_{\mathbf{H}_1}^{\mathrm{spl}}(\mathbf{M}_{\mathbf{H},1})$, which is the same as $p_{\mathbf{H}_1}^{\mathrm{spl}}(\mathbf{S}_{\mathbf{H}}^1))$ and factors through $\mathbf{S}_{\mathbf{G}}$.

6.7.8. Proof of Lemma 6.40. For the purpose of the proof, we may consider $\mathscr{P}_{\mathbf{B}}$, including all the proper parabolic \mathbb{Q} -subgroups containing \mathbf{B} , not just the maximal ones. This is because the character associated to a parabolic \mathbb{Q} -subgroup can be expressed as \mathbb{Q}_{\geq} -linear combinations of the characters of maximal parabolic \mathbb{Q} -subgroups containing it.

By [Gro97, Theorem 7.3], we find a highest weight \mathbb{Q} -vector \mathbf{v} and a parabolic \mathbb{Q} -subgroup \mathbf{P}_1 preserving $\mathbb{C}.\mathbf{v}$ such that \mathbf{v} is fixed by \mathbf{B} , \mathbf{B} is contained in \mathbf{P}_1 and $\mathbf{R}_u(\mathbf{B})$ is contained in $\mathbf{R}_u(\mathbf{P}_1)$. Here χ is the \mathbb{Q} -character of \mathbf{P}_1 associated with \mathbf{v} .

Let \mathbf{B}^{red} be a maximal reductive \mathbb{Q} -subgroup of \mathbf{B} and $\mathbf{\tilde{S}}_{\mathbf{B}}$ be the \mathbb{Q} -split part of the central torus of \mathbf{B}^{red} . Find a maximal reductive \mathbb{Q} -subgroup $\mathbf{L}_{\mathbf{P}_1}$ of \mathbf{P}_1 containing \mathbf{B}^{red} . Then $\mathbf{B}^{\text{red}} \subset \mathbf{L}_{\mathbf{P}_1}^{\chi} := \mathbf{L}_{\mathbf{P}_1} \cap \ker \chi$. Let $\mathbf{S}_{\mathbf{P}_1}$ be the \mathbb{Q} -split part of the central torus of $\mathbf{L}_{\mathbf{P}_1}$ and $\mathbf{M}_{\mathbf{P}_1} := (^{\circ}\mathbf{L}_{\mathbf{P}_1})^{\circ}$. Then $\mathbf{L}_{\mathbf{P}_1}$ is an almost direct product $\mathbf{M}_{\mathbf{P}_1} : \mathbf{S}_{\mathbf{P}_1}$.

Let $\mathbf{S}_{\mathbf{P}_1}^{\chi} := \mathbf{S}_{\mathbf{P}_1} \cap \ker \chi$ and $\widetilde{\mathbf{S}}_{\mathbf{B}}' := \widetilde{\mathbf{S}}_{\mathbf{B}} \cdot \mathbf{S}_{\mathbf{P}_1}^{\chi}$, contained in the centralizer of \mathbf{B}^{red} . Then $\widetilde{\mathbf{S}}_{\mathbf{B}}'$ is an almost direct product of two subtori: $\mathbf{S}_1 := \mathbf{S}_{\mathbf{P}_1}^{\chi}$ and some $\mathbf{S}_2 \subset \mathbf{M}_{\mathbf{P}_1} \cap \widetilde{\mathbf{S}}_{\mathbf{B}}'$.

The proof below actually proves the conclusion with **B** replaced by $\mathbf{B}' := \widetilde{\mathbf{S}}'_{\mathbf{B}} \cdot \mathbf{B} \cdot \mathbf{R}_{\mathbf{u}}(\mathbf{P}_1)$, which is a stronger claim. For this purpose, we are interested in two types of parabolic Q-subgroups.

The first type \mathscr{P}_1 consists of parabolic Q-subgroups containing \mathbf{P}_1 . The other type \mathscr{P}_2 is constructed as follows. Take a Q-cocharacter $\mathbf{a}_t : \mathbf{G}_m \to \mathbf{S}_2$, let

$$\mathbf{Q}_{\mathbf{a}_t}^1 := \left\{ x \in \mathbf{L}_{\mathbf{P}_1} \ \Big| \ \lim_{t \to 0} \mathbf{a}_t x \mathbf{a}_t^{-1} \text{ exists } \right\}, \quad \mathbf{Q}_{\mathbf{a}_t} := \mathbf{Q}_{\mathbf{a}_t}^1 \ltimes \mathbf{R}_{\mathbf{u}}(\mathbf{P}_1)$$

Then $\mathbf{Q}_{\mathbf{a}_t}^1$ is a parabolic \mathbb{Q} -subgroup of $\mathbf{L}_{\mathbf{P}_1}$ and $\mathbf{Q}_{\mathbf{a}_t}$ is a parabolic \mathbb{Q} -subgroup of \mathbf{A} . Define \mathscr{P}_2 to be the collection of all $\mathbf{Q}_{\mathbf{a}_t}$'s constructed this way. One sees from the definition that every element from $\mathscr{P}_1 \cup \mathscr{P}_2$ contains \mathbf{B}' . Let \mathcal{P}'_1 and \mathcal{P}'_2 be the associated linear functionals (denoted as $l'_{\mathbf{P}}$ as \mathbf{P} varies in $\mathscr{P}'_1 \cup \mathscr{P}'_2$) on $\operatorname{Lie}(\widetilde{\mathbf{S}}'_{\mathbf{B}})^{\vee}$.

First we explain that \mathcal{P}'_1 is non-degenerate when restricted to $\text{Lie}(\mathbf{S}_1)$. Let $\mathbf{S}_{\max} \subset \mathbf{L}_{\mathbf{P}_1}$ be a maximal \mathbb{Q} -split torus of \mathbf{A} containing $\mathbf{\widetilde{S}}_{\mathbf{B}}$ and $\mathbf{S}_{\mathbf{P}_1}$. Also fix a \mathbb{Q} -minimal parabolic subgroup \mathbf{P}_{\min} containing \mathbf{S}_{\max} that is contained in \mathbf{P}_1 . So

$$\widetilde{\mathbf{S}}_{\mathbf{B}}' = \mathbf{S}_1 \cdot \mathbf{S}_2 \subset \widetilde{\mathbf{S}}_{\mathbf{B}} \cdot \mathbf{S}_{\mathbf{P}_1} \subset \mathbf{S}_{\max} \subset \mathbf{P}_{\min} \subset \mathbf{P}_1.$$

Let Δ be the set of simple roots. Then one can find $I \subset \Delta$ such that $\mathbf{S}_{\mathbf{P}_1} = (\bigcap_{\alpha \in I} \ker \alpha)^\circ$. Moreover, χ is a non-zero $\mathbb{Q}_{\geq 0}$ -linear combination of $l'_{\mathbf{P}}$ as \mathbf{P} varies over maximal parabolic \mathbb{Q} -subgroups containing \mathbf{P}_1 . If \mathcal{P}'_1 is not non-degenerate inside $\operatorname{Lie}(\mathbf{S}_1)^{\vee}$, then $l'_{\mathbf{P}}(v) > 0$ for some $v_{\neq 0} \in \operatorname{Lie}(\mathbf{S}_1)$ and for each maximal parabolic \mathbb{Q} -subgroup \mathbf{P} containing \mathbf{P}_1 , implying that $d\chi(\mathbf{v}) > 0$, a contradiction.

Next we show that \mathcal{P}'_2 is non-degenerate restricted to $\text{Lie}(\mathbf{S}_2)$. Take $\mathbf{Q} = \mathbf{Q}_{\mathbf{a}_t} \in \mathscr{P}_2$. Let $l''_{\mathbf{Q}}$ be the linear functional on $\text{Lie}(\mathbf{S}_2)$ associated with $\mathbf{Q}_{\mathbf{a}_t}^1$. Since the determinant character of the adjoint action of \mathbf{S}_2 on $\mathbf{R}_{\mathbf{u}}(\mathbf{P}_1)$ is trivial, $l''_{\mathbf{Q}}$ is equal to $l'_{\mathbf{Q}}$ restricted to $\text{Lie}(\mathbf{S}_2)$. Since $\mathbf{L}_{\mathbf{P}_1}$ is reductive, it admits an involution whose restriction to \mathbf{S}_2 brings each element to its inverse. Therefore, if $\{l''_{\mathbf{Q}}\}_{\mathbf{Q}\in\mathscr{P}_2}$ is not non-degenerate, then there exists a \mathbb{Q} -cocharacter $\mathbf{b}_t : \mathbf{G}_m \to \mathbf{S}_2$ such that $l''_{\mathbf{Q}}$ vanishes on the image of $d\mathbf{b}_t$ for all $\mathbf{Q}_{\mathbf{a}_t} = \mathbf{Q} \in \mathscr{P}_2$. In particular, this is true when $\mathbf{a}_t = \mathbf{b}_t$, which is a contradiction.

The general case follows since $\text{Lie}(\mathbf{S}_2)$ is contained in the common kernel of elements from \mathcal{P}'_1 .

6.7.9. Definition of the compactification. For $\mathbf{v} \in \mathscr{P}_o^1$, let $V_{\mathbf{v}}$ be the \mathbf{G}_1 -representation where \mathbf{v} lives. The character $\alpha_{\mathbf{v}} : \mathbf{H} \to \mathbf{G}_m$ defined by

$$p_1(h).\mathbf{v} = \alpha_{\mathbf{v}}(h)\mathbf{v}, \ \forall h \in \mathbf{H}$$

factors through some $\beta_{\mathbf{v}} : \mathbf{S}_{\mathbf{G}} \to \mathbf{G}_m$. Now we define a Q-linear action $\rho_{\mathbf{v}}$ of \mathbf{G} on $V_{\mathbf{v}}$ twisting the \mathbf{G}_1 -representation by $\beta_{\mathbf{v}}^{-1}$:

$$\rho_{\mathbf{v}}(g) \cdot w := \beta_{\mathbf{v}}(p^{\operatorname{spl}}(g))^{-1} \cdot p_1(g).w, \ \forall g \in \mathbf{G}, \ w \in V_{\mathbf{v}}.$$

Thus for every $h \in \mathbf{H}$,

$$\rho_{\mathbf{v}}(h)\mathbf{v} = \beta_{\mathbf{v}}(p^{\mathrm{spl}}(h))^{-1} \cdot \alpha_{\mathbf{v}}(h)\mathbf{v} = \mathbf{v}$$

implying \mathbf{v} is fixed by \mathbf{H} .

Let $(\mathbf{X}_1, \mathbf{B}_1)$ be as offered by Assumption 6.31 (by the remark after the assumption, we assume this is a **G**-pair). Recall that $\mathbf{G}/\mathbf{H} \cong \mathbf{U}$ with identity coset sent to $o \in \mathbf{U}(\mathbb{Q})$. For the rest of this subsection, let **X** be the Zariski closure of

$$\begin{aligned} \mathbf{U} &\hookrightarrow \mathbf{X}_1 \times \bigoplus_{\mathbf{v} \in \mathscr{P}_o^1} V_{\mathbf{v}} \subset \mathbf{X}_1 \times \prod_{\mathbf{v} \in \mathcal{P}_1^1} \mathbf{P}(V_{\mathbf{v}} \oplus \mathbb{Q}) \\ g.o &\mapsto (g.o, \oplus \rho_{\mathbf{v}}(g)\mathbf{v}), \end{aligned}$$

And let

$$\mathbf{B} := \left\{ \left(x, \prod [\mathbf{x}_{\mathbf{v}} : t_{\mathbf{v}}] \right) \mid x \in \mathbf{B}_1; \ \mathbf{t}_{\mathbf{v}} = 0, \ \forall \, \mathbf{v} \in \mathscr{P}_o^1 \right\}.$$

Let \mathbf{D} be the complement of \mathbf{U} in \mathbf{X} .

6.7.10. Proof of Lemma 6.39, I, equidistribution. Take $x \in \mathbf{U}(\mathbb{Q})$ and $(g_n) \subset \mathbf{G}$ with $\lim g_n . x \in \mathbf{B}(\mathbb{R})$. Find $g_x \in \mathbf{G}(\overline{\mathbb{Q}})$ such that $x = g_x.o.$ So $(g_ng_x.o)$ converges to some point in $\mathbf{B}(\mathbb{R})$. By our assumption, we may and do require that g_n 's and g_x are contained in $\mathbf{G}^{ss}\mathbf{Z}(\mathbf{G})^{an} \times \mathbf{R}_{\mathbf{u}}(\mathbf{G})$. Item 1 of Lemma 6.39 holds for free by Lemma 6.38. In particular, replacing g_n by $g_nh_n^1$ for some $(h_n^1) \subset \mathbf{H}_x^\circ$, there are $(b_{n,1})$ bounded in $\mathbf{G}_1, (\gamma_{n,1}) \subset \Gamma_1$ such that $p_1(g_n) = b_{n,1}\gamma_{n,1}$ and $\lim_{n\to\infty} \left[(\gamma_{n,1})_* \mathbf{m}_{[\mathbf{H}_{x,1}^\circ]} \right] = [\mathbf{m}_{[\mathbf{G}_1]}].$

Now turn to item 2. Namely, we need to find $(h_n) \subset \mathbf{H}_x(\mathbb{R})^\circ$ such that

- 1. $(g_n h_n \Omega_x)$ is nondivergent in G/ Γ ;
- 2. for every $\mathbf{P} \in \mathscr{P}_{\mathbf{H}_{n,1}}^{\max}$, $\|\operatorname{Ad}(p_1(g_nh_n))\mathbf{v}_{\mathbf{P}}\| \to \infty$.

Thanks to Lemma 6.21, for every Q-subgroup **F** of **G**₁ that is normalized by **H**₁, $p_1(g_x)\mathbf{F}p_1(g_x)^{-1}$ is defined over Q. In particular, conjugating by $p_1(g_x)$ induces bijections between $\mathscr{P}_{\mathbf{H}_1}^{\max}$ and $\mathscr{P}_{\mathbf{H}_x,1}^{\max}$. The proof of Lemma 6.21 also shows that conjugating by $p_1(g_x)$ induces bijections between $\mathfrak{X}^*_{\mathbb{Q}}(\mathbf{H}_1)$ and $\mathfrak{X}^*_{\mathbb{Q}}(\mathbf{H}_{x,1})$.

For i = 1, 2, let $\overline{\mathcal{P}}_o^i$ be the characters of \mathbf{H}_1 associated with \mathscr{P}_o^i and $\overline{\mathcal{P}}_x^i$ (resp. \mathscr{P}_x^i) be the subset of $\mathfrak{X}^*_{\mathbb{Q}}(\mathbf{H}_{x,1})$ (resp. $\mathscr{P}^+_{\mathbf{H}_{x,1}}$) induced from conjugating by $p_1(g_x)$. By assumption we have

$$\|p_1(g_n g_x) \cdot \mathbf{v}\| \to +\infty, \quad \forall \mathbf{v} \in \mathscr{P}_o^1.$$
$$\|p_1(g_n) \cdot \mathbf{v}\| \to +\infty, \quad \forall \mathbf{v} \in \mathscr{P}_x^1.$$
(27)

Then

We choose a Q-split subtorus $\widetilde{\mathbf{S}}_{\mathbf{H}_{x,1}} \subset \mathbf{H}_{x,1}$ such that $p_{\mathbf{H}_{x,1}}^{\mathrm{spl}}$ restricted to $\widetilde{\mathbf{S}}_{\mathbf{H}_{x,1}}$ is surjective onto $\mathbf{S}_{\mathbf{H}_{x,1}}$ with finite kernel. For i = 1, 2, let \mathcal{P}_x^i collect the linear functionals on $\mathrm{Lie}(\mathbf{S}_{\mathbf{H}_{x,1}})$ induced from $\overline{\mathcal{P}}_x^i$. Let $\mathbf{S}_{\mathbf{H}_x}^1$ be the identity component of the common kernel of $\overline{\mathcal{P}}_x^1$ in $\widetilde{\mathbf{S}}_{\mathbf{H}_{x,1}}$ and $\widetilde{\mathbf{S}}_1^{\mathbf{H}_x}$ be that of $\overline{\mathcal{P}}_x^2$. Then $\widetilde{\mathbf{S}}_{\mathbf{H}_{x,1}} = \mathbf{S}_{\mathbf{H}_x}^1 \cdot \widetilde{\mathbf{S}}_1^{\mathbf{H}_x}$ is an almost direct product of two Q-subtori. We need the following lemma:

Lemma 6.42. Let \mathbf{A} be a linear algebraic group over \mathbb{Q} and (γ_n) be contained in a fixed arithmetic subgroup of \mathbf{A} . Let \mathbf{C} be a \mathbb{Q} -subgroup and assume that $\mathscr{P}_{\mathbf{C}}^{\max}$ is finite. Let $\widetilde{\mathbf{S}}_{\mathbf{C}}$ be a \mathbb{Q} -split subtorus of \mathbf{C} that surjects onto $\mathbf{S}_{\mathbf{C}}$ under $p_{\mathbf{C}}^{\mathrm{spl}}$ with a finite kernel. After passing to a subsequence, one of the following holds:

(1) there exists $(z_n) \subset \operatorname{Lie}(\widetilde{\mathbf{S}}_{\mathbf{C}}(\mathbb{R}))$ such that

$$\|\operatorname{Ad}(\gamma_n \exp(z_n))\mathbf{v}_{\mathbf{P}}\| \to \infty, \ \forall \, \mathbf{P} \in \mathscr{P}_{\mathbf{C}}^{\max};$$

(2) there exists $\mathbf{v} \in \mathscr{P}^+_{\mathbf{C}}$ fixed by \mathbf{C} such that $(\|\gamma_n \cdot \mathbf{v}\|)$ is bounded.

Proof. The proof is essentially contained in Section 4.4.5. Indeed, by [Zha21, Lemma 3.4], if the first alternative were not true, then there are positive integers (a_i) and $(\mathbf{P}_i) \subset \mathscr{P}_{\mathbf{C}}^{\max}$ such that $\mathbf{v} := \otimes \mathbf{v}_{\mathbf{P}_i}^{\otimes a_i}$ is fixed by \mathbf{C} and $(\gamma_n.\mathbf{v})$ is bounded.

By our assumption, Lemma 6.42 is applicable to $\mathbf{A} = \mathbf{G}_1$, $\mathbf{C} = \mathbf{H}_{x,1}$ and $\gamma_n = \gamma_{n,1}$. As the second alternative violates the equidistribution towards $\mathbf{m}_{[G_1]}$, there exist $(s_n) \subset \operatorname{Lie}(\mathbf{S}^1_{\mathbf{H}_x}(\mathbb{R}))$ and $(t_n) \subset \operatorname{Lie}(\widetilde{\mathbf{S}}^{\mathbf{H}_x}_1(\mathbb{R}))$ such that

$$\|\operatorname{Ad}(p_1(g_n)\exp(s_n+t_n))\mathbf{v}_{\mathbf{P}}\| \to \infty, \ \forall \, \mathbf{P} \in \mathscr{P}_{\mathbf{H}_{x,1}}^{\max}.$$

Therefore,

$$||p_1(g_n)\exp(s_n+t_n).\mathbf{v}|| \to \infty, \ \forall \, \mathbf{v} \in \mathscr{P}^+_{\mathbf{H}_{x,1}}.$$

Now observe that one can replace (t_n) by zero. Indeed, for every $\mathbf{v} \in \mathscr{P}^2_x$,

$$\|p_1(g_n)\exp(s_n)\cdot\mathbf{v}\| = \|p_1(g_n)\exp(s_n+t_n)\cdot\mathbf{v}\| \to \infty.$$

On the other hand, by Equa.(27),

$$||p_1(g_n)\exp(s_n).\mathbf{v}|| = ||p_1(g_n).\mathbf{v}|| \to \infty$$

for every $\mathbf{v} \in \mathscr{P}^1_x$. Find $(h_n) \subset \mathbf{M}_{\mathbf{H}_x}(\mathbb{R})^\circ$ $(\mathbf{M}_{\mathbf{H}_x} := (\mathbf{H}_x \cap {}^\circ \mathbf{G})^\circ)$ such that $\exp(s_n) = p_1(h_n)$, then

$$\|\operatorname{Ad}(p_1(g_nh_n))\mathbf{v}_{\mathbf{P}}\| \to \infty, \ \forall \, \mathbf{P} \in \mathscr{P}_{\mathbf{H}_{x,1}}^{\max}.$$
(28)

To complete the proof, it only remains to verify that $(g_n h_n \Omega_x)$ is nondivergent in G/Γ . Since $p^{\text{spl}}(g_n h_n \Omega_x) = p^{\text{spl}}(\Omega_x)$ is bounded in $\mathbf{S}_{\mathbf{G}}(\mathbb{R})$, it suffices to show that $(p_1(g_n h_n)\Omega_{1,x})$ is nondivergent for every nonempty open bounded subset $\Omega_{1,x}$ of $\mathrm{H}^{\circ}_{x,1}\Gamma_1/\Gamma_1$, which is true by Equa.(28) and Theorem 4.8.

6.7.11. *Proof of Lemma 6.39, II, non-empty intersection.* Here we prove the remaining assertion of Lemma 6.39, namely,

Claim. For $x \in U(\mathbb{Q})$, the closure of G.x under the analytic topology intersects with $\mathbf{B}(\mathbb{R})$ non-trivially.

Proof. By Assumption 6.31, let (g_n) be a sequence in G such that every limit point of $(g_n.x)$ is contained in $\mathbf{B}_1(\mathbb{R})$. We will build a new sequence $(g'_n) \subset \mathbf{G}$ such that $(g'_n.x)$ converges to some point in $\mathbf{B}(\mathbb{R})$.

By Lemma 6.38,

$$\lim_{n \to \infty} \left[p_1(g_n)_* \mathbf{m}_{[\mathbf{H}_{x,1}^\circ]} \right] = \left[\mathbf{m}_{[\mathbf{G}_1]} \right].$$
⁽²⁹⁾

There exist a bounded sequence $(\delta_n) \subset G_1$, a sequence $(\gamma_n) \subset \Gamma_1$ and $(h_n) \subset H_{x,1}^{\circ}$ such that $p_1(g_n) = \delta_n \gamma_n h_n$.

Apply Lemma 6.42 with $\mathbf{C} = \mathbf{H}_{x,1}$, $\mathbf{A} = \mathbf{G}_1$. The second alternative can not hold because of Equa.(29). Therefore, there exists a sequence (z_n) of $\operatorname{Lie}(\widetilde{\mathbf{S}}_{\mathbf{H}_{x,1}}(\mathbb{R}))$ such that

$$\|\operatorname{Ad}(p_1(g_n)\exp(z_n))\mathbf{v}_{\mathbf{P}}\| \to \infty, \ \forall \, \mathbf{P} \in \mathscr{P}_{\mathbf{H}_{x,1}}^{\max}.$$
(30)

Recall $g_x \in {}^{\circ}\mathbf{G}(\overline{\mathbb{Q}})$ and $x = g_x.o.$ Also take $g'_n \in {}^{\circ}\mathbf{G}$ such that $p_1(g'_n) = p_1(g_n) \exp(z_n)$. We shall show that the sequence $(g'_n.x) = (g'_ng_x.o)$ converges to some point in $\mathbf{B}(\mathbb{R})$. For this, we only need to check that

$$\|\rho_{\mathbf{v}}(g'_n g_x)\mathbf{v}\| = \|p_1(g'_n g_x)\mathbf{v}\| \to \infty, \quad \forall \, \mathbf{v} \in \mathscr{P}^1_o.$$

As vectors in $\mathscr{P}_o^1 \subset \mathscr{P}_{\mathbf{H}_1}^+$ are obtained by tensoring powers of $\mathbf{v}_{\mathbf{P}}$'s for some $\mathbf{P} \in \mathscr{P}_{\mathbf{H}_1}^{\max}$, the above holds if

$$\|\operatorname{Ad}(p_1(g'_ng_x))\mathbf{v}_{\mathbf{P}}\| \to \infty, \quad \forall \, \mathbf{P} \in \mathscr{P}_{\mathbf{H}_1}^{\max}.$$

By Lemma 6.21, for every $\mathbf{P} \in \mathscr{P}_{\mathbf{H}_1}^{\max}$, $\mathbf{P}_x := p_1(g_x)\mathbf{P}p_1(g_x)^{-1}$ belongs to $\mathscr{P}_{\mathbf{H}_{x,1}}^{\max}$. So there is a non-zero complex number $c_{\mathbf{P}}$ such that $\operatorname{Ad}(p_1(g_x))\mathbf{v}_{\mathbf{P}} = c_{\mathbf{P}}\mathbf{v}_{\mathbf{P}_x}$. So

$$\|\operatorname{Ad}(p_1(g'_ng_x))\mathbf{v}_{\mathbf{P}}\| = |c_{\mathbf{P}}| \cdot \|\operatorname{Ad}(p_1(g_n)\exp(z_n))\mathbf{v}_{\mathbf{P}_x}\|,$$

which diverges to $+\infty$ by Equa.(30).

6.8. Proof of Theorem 1.2. Let us prove something more general:

Theorem 6.43. Let \mathbf{G} be a connected semisimple linear algebraic group over \mathbb{Q} without compact factors and \mathbf{H} be a connected reductive \mathbb{Q} -subgroup without non-trivial \mathbb{Q} -characters. Then there exists a smooth \mathbf{G} -pair (\mathbf{X}, \mathbf{D}) with $\mathbf{U} := \mathbf{X} \setminus \mathbf{D}$ equivariantly isomorphic to \mathbf{G}/\mathbf{H} and an effective divisor \mathbf{L} supported on \mathbf{D} such that for any smooth metric on $\mathcal{O}_{\mathbf{X}}(\mathbf{L})$, the associated height function $\mathrm{Ht} := \mathrm{Ht}_{\mathbf{L}}$ satisfies:

$$\lim_{R \to \infty} \frac{\#\Gamma . x \cap B_R^{\mathrm{Ht}}}{\omega_{\infty}(B_R^{\mathrm{Ht}} \cap \mathrm{G.}x)} = \frac{|\mathbf{m}_{[\mathrm{H}_x]}|}{|\mathbf{m}_{[\mathrm{G}]}|}, \quad \forall x \in \mathbf{U}(\mathbb{Q}), \ \forall \ arithmetic \ subgroup \ \Gamma.$$
(31)

Remark 6.44. This result, combined with [WX16, Theorem 4.3], gives generalizations of Theorem 1.2.

Proof. Write $\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ}$ as an almost direct product:

$$\mathbf{Z}_{\mathbf{G}}(\mathbf{H})^{\circ} = \mathbf{Z}(\mathbf{H})^{\circ} \cdot \mathbf{M}_{\mathbf{Z}} \cdot \mathbf{T}_{\mathbf{Z}} = \mathbf{Z}(\mathbf{H})^{\circ} \cdot \mathbf{M}_{\mathbf{Z}} \cdot \mathbf{T}_{\mathbf{Z}}^{\mathrm{an}} \cdot \mathbf{T}_{\mathbf{Z}}^{\mathrm{spl}}$$

where $\mathbf{M}_{\mathbf{Z}}$ is semisimple, $\mathbf{T}_{\mathbf{Z}}^{\mathrm{an}}$ is a Q-anisotropic torus and $\mathbf{T}_{\mathbf{Z}}^{\mathrm{spl}}$ is a Q-split torus. Let $\widetilde{\mathbf{G}} := \mathbf{G} \times \mathbf{M}_{\mathbf{Z}} \times \mathbf{T}_{\mathbf{Z}}^{\mathrm{an}} \times \mathbf{T}_{\mathbf{Z}}^{\mathrm{spl}}$. It acts on $\mathbf{U} \cong \mathbf{G}/\mathbf{H}$ by $(a, b_1, b_2, b_3).g\mathbf{H} := agb^{-1}\mathbf{H}$ with $b := b_1b_2b_3$. Write $\mathbf{F} := \mathbf{M}_{\mathbf{Z}} \times \mathbf{T}_{\mathbf{Z}}^{\mathrm{an}} \times \mathbf{T}_{\mathbf{Z}}^{\mathrm{spl}}$ for simplicity. Let $\iota_o : \mathbf{F} \to \mathbf{G}$ be the natural product map. And let $\Delta_{\iota_o} : \mathbf{F} \to \widetilde{\mathbf{G}}$ be defined by $f \mapsto (\iota_o(f), f)$. For a general $x \in \mathbf{U}(\mathbb{Q})$, fix $g_x \in \mathbf{G}(\overline{\mathbb{Q}})$ with $x = g_x.o$. Let $\iota_x(f) := g_x \iota_o(f) g_x^{-1}$ and Δ_{ι_x} be defined similarly. Similar to the proof of Lemma 6.21, one shows that ι_x is actually defined over \mathbb{Q} . Then, the stabilizer of x in $\widetilde{\mathbf{G}}$ is $\mathbf{H}_x \times \Delta_{\iota_x}(\mathbf{F})$. Let $\widetilde{\Gamma} := \Gamma \times \mathbf{M}_{\mathbf{Z}} \cap \Gamma \times \mathbf{T}_{\mathbf{Z}}^{\mathrm{an}} \cap \Gamma$.

We now apply discussions in previous sections to the pair $(\widetilde{\mathbf{G}}, \widetilde{\mathbf{H}})$ in the place of (\mathbf{G}, \mathbf{H}) . Suppose that $(\widetilde{\mathbf{G}}, \widetilde{\mathbf{H}})$ satisfies the assumption listed in Section 6.7.1. Theorem 6.32 is applicable. Note that $\operatorname{Pic}(\mathbf{U})$ is torsion under our assumption. Combined with Lemma 5.1 and 5.3, this shows that for every x and Γ ,

$$\lim_{R \to \infty} \frac{\# \widetilde{\Gamma}.x \cap B_R^{\mathrm{Ht}}}{\omega_{\infty}(B_R^{\mathrm{Ht}} \cap \widetilde{\mathrm{G}}.x)} = \frac{\left| \mathbf{m}_{[\widetilde{\mathrm{H}}_x^1]} \right|}{\left| \mathbf{m}_{[\widetilde{\mathrm{G}}^1]} \right|}, \quad \forall x \in \mathbf{U}(\mathbb{Q}), \; \forall \text{ arithmetic subgroup } \Gamma.$$

Note that:

•
$$\widetilde{\Gamma}.x = \Gamma.x$$
, $\widetilde{G}.x = G.x$;
• $\widetilde{\mathbf{G}}^{1} = \mathbf{G} \times \mathbf{M}_{\mathbf{Z}} \times \mathbf{T}_{\mathbf{Z}}^{\mathrm{an}}$ and $\widetilde{\mathbf{H}}_{x}^{1} = \mathbf{H}_{x} \times \Delta_{\iota_{x}}(\mathbf{M}_{\mathbf{Z}} \times \mathbf{T}_{\mathbf{Z}}^{\mathrm{an}})$ and hence
 $\left| \mathbf{m}_{[\widetilde{\mathbf{H}}_{x}^{1}]} \right| = \left| \mathbf{m}_{[\mathbf{H}_{x}]} \right| \cdot \left| \mathbf{m}_{[\mathbf{M}_{\mathbf{Z}}]} \right| \cdot \left| \mathbf{m}_{[\mathbf{T}_{\mathbf{Z}}^{\mathrm{an}}]} \right|$ and $\left| \mathbf{m}_{[\widetilde{\mathbf{G}}^{1}]} \right| = \left| \mathbf{m}_{[\mathbf{G}]} \right| \cdot \left| \mathbf{m}_{[\mathbf{M}_{\mathbf{Z}}]} \right| \cdot \left| \mathbf{m}_{[\mathbf{T}_{\mathbf{Z}}^{\mathrm{an}}]} \right|$.

Thus Equa.(31) follows.

Now it only remains to verify the assumptions. As $\widetilde{\mathbf{G}}$ is reductive, Assumption 6.28 holds. As for Assumption 6.29,

- (1) $\widetilde{\mathbf{H}}$ being reductive implies the observability of $\widetilde{\mathbf{H}}$ and $p_1(\widetilde{\mathbf{H}})$;
- (2) $\mathbf{G} = \mathbf{G}^{\mathrm{nc}}$ by assumption, so p^{stau} , the quotient of $\widetilde{\mathbf{G}}$ by $\widetilde{\mathbf{G}}^{\mathrm{nc}}$, factors through that by \mathbf{G} . Thus $p^{\mathrm{stau}}|_{\widetilde{\mathbf{H}}}$ is surjective by the definition of $\widetilde{\mathbf{H}}$;
- (3) It is direct to verify that the identity component of the centralizer of $p_2(\tilde{\mathbf{H}})$ is contained in itself and hence $\mathscr{P}_{p_2(\tilde{\mathbf{H}})}$ is finite, which is equivalent to $\mathscr{P}_{\tilde{\mathbf{H}}}^{\max}$ being finite.

Finally, let us take care of Assumption 6.31. By Theorem 6.26 and Lemma 6.25, this reduces to the fact that the identity component of the centralizer of $p_2(\widetilde{\mathbf{H}})$ in $p_2(\widetilde{\mathbf{G}})$ is contained in $p_2(\widetilde{\mathbf{H}})$.

7. Example I, Representation of a binary quadratic form by a quaternary form

Let (Q_1, \mathbb{Q}^2) be a \mathbb{Q} -anisotropic quadratic from of rank 2 and (Q_2, \mathbb{Q}^4) be a rank 4 quadratic form. Let \mathbf{U}_{Q_1,Q_2} be the \mathbb{Q} -variety defined by

$$\mathbf{U}_{Q_1,Q_2}(\mathbb{Q}) := \left\{ \phi \in \operatorname{Hom}_{\mathbb{Q}}(\mathbb{Q}^2,\mathbb{Q}^4), \ \phi^*Q_2 = Q_1 \right\}.$$

We assume that

• (Q_1, \mathbb{R}^2) has signature (1, 1) and (Q_2, \mathbb{R}^4) has signature (2, 2);

• $\mathbf{U}_{Q_1,Q_2}(\mathbb{Q}) \neq \emptyset.$

Thus we can find (Q'_1, \mathbb{Q}^2) such that $(Q_2, \mathbb{Q}^4) \cong (Q_1, \mathbb{Q}^2) \oplus (Q'_1, \mathbb{Q}^2)$. We further assume that

• $\mathbf{SO}_{Q'_1}$ is not isomorphic to \mathbf{SO}_{Q_1} as an algebraic group over $\mathbb Q$.

For i = 1, 2, let $M_{Q_i} \in \operatorname{Mat}_2(\mathbb{Q})$ or $\operatorname{Mat}_4(\mathbb{Q})$ be symmetric matrices representing Q_i : $Q_i(v, w) = v^{\operatorname{tr}} M_{Q_i} w$ if $v, w \in \mathbb{Q}^2$ or \mathbb{Q}^4 are written as column vectors. Then U_{Q_1,Q_2} is naturally embedded into $\operatorname{Mat}_{4,2}$ as

$$\left\{ M \in \mathbf{Mat}_{4,2}(\mathbb{Q}) \mid M^{\mathrm{tr}} M_{Q_2} M = M_{Q_1} \right\}.$$

Let \mathcal{U}_{Q_1,Q_2} be the integral model by taking closure of \mathbf{U}_{Q_1,Q_2} in $\mathcal{M}\mathbf{at}_{4,2}$ which is naturally an affine space over \mathbb{Z} . Let ω be an invariant gauge form on \mathbf{U}_{Q_1,Q_2} over \mathbb{Q} . For each prime p, let $|\omega|_p$ be the Haar measure on $\mathbf{U}_{Q_1,Q_2}(\mathbb{Q}_p)$ induced from ω . For a matrix M with real coefficients, let ||M|| be the Euclidean norm of M.

Theorem 7.1. Write $\mathcal{U} := \mathcal{U}_{Q_1,Q_2}$ for simplicity. Then as R tends to infinity,

$$\# \{ M \in \mathcal{U}(\mathbb{Z}), \ \|M\| \le R \} \sim \prod_{p \in \operatorname{Val}_f} \omega_p \left(\mathcal{U}(\mathbb{Z}_p) \right) \cdot \omega_\infty \left(\{ M \in \mathcal{U}(\mathbb{R}), \ \|M\| \le R \} \right).$$

Namely, \mathbf{U}_{Q_1,Q_2} with this embedding is strongly Hardy-Littlewood. Also, there exists some constant c > 0 such that

$$# \{ M \in \mathcal{U}(\mathbb{Z}), \|M\| \le R \} \sim cR^2 \log R.$$

By [WX16, Theorem 4.3, Lemma 4.1] (one should use replace \mathbf{SO}_{Q_2} by its simply connected cover when applying this result and verify that the stabilizer group remains connected. Details are omitted), for the first part, it suffices to verify certain orbital counting statement which then is implied by equidistribution towards the full Haar measure generically in certain sense. At this point, let us note that \mathbf{U}_{Q_1,Q_2} is homogeneous under the action of $\mathbf{G} := \mathbf{SO}_{Q_2} \times \mathbf{SO}_{Q_1}$. The stabilizer of $\phi_0 \in \mathbf{U}_{Q_1,Q_2}(\mathbb{Q})$ is of the form $\mathbf{H}_{\phi_0} = \mathbf{H}_1 \times \mathbf{H}_2$ where \mathbf{H}_1 is the subgroup of \mathbf{SO}_{Q_2} preserving the Q_2 -orthogonal complement of the image of ϕ_0 and \mathbf{H}_2 is suitable diagonal embedding of \mathbf{SO}_{Q_1} . Under the assumptions imposed above, the only connected intermediate \mathbb{Q} -group is the maximal torus \mathbf{T}_{ϕ_0} containing \mathbf{H}_{ϕ_0} . By equidistribution theorems, the Haar measure supported on $\mathbf{H}_{\phi_0}(\mathbb{R})\Gamma/\Gamma$, where Γ is an(y) arithmetic lattice of \mathbf{G} , when pushed by (g_n) , equidistributes towards the full Haar measure unless (g_n) is bounded modulo \mathbf{T}_{ϕ_0} which one can show is generically not true. Our main purpose here is to explain how this is done using the approach taken in the current paper.

Let $\mathbf{X}^{\mathbf{Mat}}$ denote the Zariski closure of \mathbf{U}_{Q_1,Q_2} inside $\mathbf{P}(\mathbf{Mat}_{4,2} \oplus \mathbb{Q}) \cong \mathbf{P}^8_{\mathbb{Q}}$. It is covered by another "incidence compactification" which we now describe.

For a matrix M, let $\langle M \rangle$ be the linear subspace spanned by the column vectors of M. Then \mathbf{X}^{INC} is the Zariski closure of

$$\{([M:1], \langle M \rangle), M \in \mathbf{U}_{Q_1, Q_2}\}$$
 in $\mathbf{X}^{\mathbf{Mat}} \times \mathbf{Gr}(2, 4)$.

Let \mathbf{D}^{INC} be the complement of \mathbf{U}_{Q_1,Q_2} in \mathbf{X}^{INC} . The pair $(\mathbf{X}^{\text{INC}}, \mathbf{D}^{\text{INC}})$ is still not smooth. Luckily, it is equipped with a natural stratification, components of which one can try blowing up.

For simplicity, we often work with complex-valued points and a typical point in \mathbf{X}^{INC} is written as $([M : \lambda], \langle N \rangle)$ where N is some 4-by-2 matrix.

At the level of complex points, one sees that the functions $\operatorname{rk}(\langle N \rangle, Q_2)$, $\operatorname{rk}(M)$ and $\operatorname{rk}(\langle M^{\operatorname{tr}} \rangle, Q_1)$ are $\mathbf{G}(\mathbb{C})$ -invariant. Note that $f \leq c$, for f being one of these functions and c an integer, is indeed (locally) defined by zeros of certain polynomials and defines a closed **G**-invariant subvariety over \mathbb{Q} .

Definition A. For a tuple of integers (a, b, c),

• let $\mathbf{S}_{a,b,c}$ be the subvariety of \mathbf{X}^{INC} whose complex-valued points consist of $([M:\lambda], \langle N \rangle)$ such that

 $\operatorname{rk}(\langle N \rangle, Q_2) = a, \operatorname{rk}(M) = b, \operatorname{rk}(\langle M^{\operatorname{Tr}} \rangle, Q_1) = c;$

• let $\mathbf{D}_{a,b,c}$ be the Zariski closure of $\mathbf{S}_{a,b,c}$, whose complex-valued points consist of $([M:\lambda], \langle N \rangle)$ such that

 $\operatorname{rk}(\langle N\rangle,Q_2) \leq a, \ \operatorname{rk}(M) \leq b, \ \operatorname{rk}(\langle M^{\operatorname{Tr}}\rangle,Q_1) \leq c.$

RUNLIN ZHANG

When c = 0, $\mathbf{S}_{a,b,0}$, if non-empty, splits into two disjoint irreducible components, denoted as $\mathbf{S}_{a,b,0^+}$ and $\mathbf{S}_{a,b,0^-}$. Similarly, $\mathbf{D}_{a,b,0}$ is also a disjoint union of two irreducible components $\mathbf{D}_{a,b,0^+}$ and $\mathbf{D}_{a,b,0^-}$. As they are disjoint and we will be working inside affine open subvarieties where only one of them appears, we will sometimes not distinguish them.

The values of these functions are subject to certain constraints. Some easy-to-observe ones are


$$\begin{split} \mathrm{rk}(\langle N \rangle, Q_2^0) &\in \{0, 1, 2\}; \ \mathrm{rk}(M) \in \{1, 2\}; \ \mathrm{rk}(\langle M^{\mathrm{Tr}} \rangle, Q_1^0) \in \{0, 1, 2\} \\ \mathrm{rk}(\langle M^{\mathrm{Tr}} \rangle, Q_1^0) &\leq \mathrm{rk}(M). \end{split}$$

Working harder, one finds that

Lemma 7.2. The space \mathbf{X}^{INC} decomposes into non-empty locally closed subvarieties:

$$\mathbf{X}^{\text{INC}} = \mathbf{S}_{2,2,2} \bigsqcup \mathbf{S}_{2,1,0} \bigsqcup \mathbf{S}_{1,1,1} \bigsqcup \mathbf{S}_{1,1,0} \bigsqcup \mathbf{S}_{0,2,2} \bigsqcup \mathbf{S}_{0,1,1} \bigsqcup \mathbf{S}_{0,1,0}.$$

Moreover, \mathbf{X}^{INC} has the following stratified structure:



where $A \rightarrow B$ means "A is contained in the closure of B".

Now, a smooth pair can be obtained from the following three-step blowup processes:

where \mathbf{E}_i denotes the exceptional divisor of π_i and \mathbf{D}^+ the birational transform of a closed subvariety, which is well defined as long as $\mathbf{D} \setminus \mathbf{C}$ ($\mathbf{C} :=$ center of the blowup) is dense in \mathbf{D} .

Let \mathbf{D}^i be the complement of \mathbf{U}_{Q_1,Q_2} in \mathbf{X}^i . Let \mathbf{E}_2^1 (resp. \mathbf{E}_2^2) be the inverse image of \mathbf{D}_{011}^+ (resp. \mathbf{D}_{110}^+) under π_2 . Let \mathbf{E}_3^0 (resp. $\mathbf{E}_3^1, \mathbf{E}_3^2$) be the total transform of $\mathbf{E}_2^1 \cap \mathbf{D}_{022}^{++}$ (resp. $\mathbf{D}_{111}^{++}, \mathbf{E}_2^2 \cap \mathbf{D}_{210}^{++}$). Let (\mathcal{L}_0, s_0) be the line bundle and the global section over $\mathbf{X}^{\mathbf{Mat}}$ that underlies the definition of $\|\cdot\|$

and let $(\widetilde{\mathcal{L}}_0, \widetilde{s}_0)$ be their pull-backs over \mathbf{X}^3 .

Lemma 7.3. $(\mathbf{X}^3, \mathbf{D}^3)$ is a smooth pair. Moreover, the irreducible components of \mathbf{D}^3 are

$$(\mathbf{E}_1^{++}, \mathbf{D}_{210}^{++}, \mathbf{E}_3^2, (\mathbf{E}_2^2)^+, \mathbf{E}_3^1, (\mathbf{E}_2^1)^+, \mathbf{E}_3^0, \mathbf{D}_{022}^+))$$

which in this order are labeled as $(\mathbf{D}_1^3, ..., \mathbf{D}_8^3)$. Also \mathbf{D}_1^3 intersects with the rest of the divisors transversally, and for the other i's, \mathbf{D}_i^3 , besides \mathbf{D}_1^3 , only intersects with \mathbf{D}_{i+1}^3 . Moreover,

$$-\operatorname{div}(\omega_{\mathbf{U}}) = 2\mathbf{D}_{1}^{3} + 1\mathbf{D}_{2}^{3} + 3\mathbf{D}_{3}^{3} + 5\mathbf{D}_{4}^{3} + 3\mathbf{D}_{5}^{3} + 7\mathbf{D}_{6}^{3} + 5\mathbf{D}_{7}^{3} + 3\mathbf{D}_{8}^{3} =: \sum d_{\alpha}\mathbf{D}_{\alpha}^{3}$$

and

$$\operatorname{div}(\widetilde{s}_0) = 1\mathbf{D}_1^3 + 1\mathbf{D}_2^3 + 2\mathbf{D}_3^3 + 3\mathbf{D}_4^3 + 1\mathbf{D}_5^3 + 3\mathbf{D}_6^3 + 2\mathbf{D}_7^3 + 1\mathbf{D}_8^3 =: \sum \lambda_{\alpha} \mathbf{D}_{\alpha}^3.$$

Consequently,

$$\left(\frac{d_{\alpha}-1}{\lambda_{\alpha}}\right)_{\alpha=1}^{8} = (1,0,1,4/3,2,2,2,2)$$

The proof of Lemma 7.3, which can be found in the appendix, relies on explicit construction of blowups and explicit calculation of the divisor of the invariant gauge form.

Note that $G := \mathbf{SO}_{Q_2}(\mathbb{R})$ is connected in analytic topology. Also, \mathbf{H}_x denotes the stabilizer subgroup of \mathbf{G} of some point $x \in \mathbf{U}(\mathbb{Q})$. Thanks to our \mathbb{R} -split assumption,

Lemma 7.4. For every $\alpha \in \{1, ..., 8\}$, $\mathbf{D}^3_{\alpha}(\mathbb{R}) \neq \emptyset$ and $\mathbf{U}(\mathbb{R})$ consists of a single G-orbit. Consequently, for every $\alpha = 1, ..., 8$ and $x \in \mathbf{U}(\mathbb{Q})$, the analytic closure of G.x intersects with $\mathbf{D}_{\alpha}(\mathbb{R})$.

Proof. That $\mathbf{D}^3_{\alpha}(\mathbb{R}) \neq \emptyset$ follows from the construction (see appendix for details). It is easy to see that the analytic closure of $\mathbf{U}(\mathbb{R})$ intersects with every $\mathbf{D}^3_{\alpha}(\mathbb{R})$. However, G.x is equal to $\mathbf{U}(\mathbb{R})$ thanks to the following "exact" sequence

$$H_x \to G \to U(\mathbb{R}) \to H^1(Gal(\mathbb{C}/\mathbb{R}), \mathbf{H}_x)$$

and the triviality of $H^1(\text{Gal}(\mathbb{C}/\mathbb{R}), \mathbf{H}_x)$.

Let Γ be some arithmetic lattice.

Lemma 7.5. For every $x \in \mathbf{U}(\mathbb{Q})$ and every sequence (g_n) of G, the following two are equivalent:

- 1. every limit point of $(g_n.x)$ is contained in $\bigcup_{\alpha\neq 2} \mathbf{D}^3_{\alpha}(\mathbb{R})$;
- 2. under the weak-* topology, $\lim_{n\to\infty} (g_n)_* \mathbf{m}_{[\mathbf{H}_x]}^1 = \mathbf{m}_{[\mathbf{G}]}^1$.

Proof. By our assumption, the only intermediate connected closed \mathbb{Q} -subgroup between \mathbf{H}_x and \mathbf{G} is the centralizer of \mathbf{H}_x , a maximal torus in \mathbf{G} . And a sequence (g_n) is bounded modulo this maximal torus iff the limit of $(g_n.x)$ is contained in $\mathbf{D}_2^3(\mathbb{R})^\circ \cup \mathbf{U}(\mathbb{R})$. The conclusion thus follows from Theorem 4.2.

Proof of Theorem 7.1. For the first part, by [WX16, Theorem 4.3] and the fact that $\operatorname{Pic}(\mathbf{H})$ and hence $\operatorname{Br}(\mathbf{U})/\operatorname{Br}(\mathbb{Q})$ is trivial, it is sufficient to show that for every $x \in \mathbf{U}(\mathbb{Q})$, and every arithmetic lattice Γ of \mathbf{G} ,

$$\frac{\# \{ M \in \Gamma.x \mid \|M\| \le R \}}{\omega_{\infty} \{ M \in \mathbf{G}.x \mid \|M\| \le R \}} \sim \frac{|\mathbf{m}_{[\mathbf{H}_x]}|}{|\mathbf{m}_{[\mathbf{G}]}|}$$

where ω_{∞} is identified with the Haar measure m_{G/H_x} which is assumed to be compatible with the Haar measures m_G and m_{H_x} .

By Lemma 7.3 and Theorem 2.16, the limits

$$\nu := \lim_{R \to \infty} \frac{\mathrm{m}_{\mathrm{G/H}_x} \cdot \mathbf{1}_{B_{R,x}}}{\mathrm{m}_{\mathrm{G/H}_x}(B_{R,x})}$$

exists and is supported on $\bigcup_{i=5}^{7} \mathbf{D}_{i}^{3}(\mathbb{R}) \cap \mathbf{D}_{i+1}^{3}(\mathbb{R})$. Here $B_{R,x} := \{M \in G.x, \|M\| \leq R\}$. It only remains to invoke Lemma 5.1.

The second part follows from Theorem 2.16 and the calculation of $\left(\frac{\lambda_{\alpha}-1}{d_{\alpha}}\right)$ as in Lemma 7.3.

8. EXAMPLE II,
$$(n, 1)$$
-Splitting of \mathbb{Z}^{n+1}

Let

$$\Lambda(n,1) := \left\{ (v,M) \mid v \in \mathbb{Z}^n, \, M \in \operatorname{Prim}^n(\mathbb{Z}^{n+1}), \, \mathbb{Z}^{n+1} = \mathbb{Z}.v \oplus M \right\}$$

where $\operatorname{Prim}^{n}(\mathbb{Z}^{n+1})$ denotes the collection of the rank-*n* subgroups *M* of \mathbb{Z}^{n+1} such that \mathbb{Z}^{n+1}/M is torsion-free.

Theorem 8.1. For two positive integers λ_1, λ_2 , we have

$$#\{(v,M) \in \Lambda(n,1), \|v\|^{\lambda_1} \|M\|^{\lambda_2} \le R\} \sim \begin{cases} c_{\lambda_1,\lambda_2} \cdot R^{\frac{n+1}{\lambda_1}} \log(R), & \lambda_1 = \lambda_2\\ c_{\lambda_1,\lambda_2} \cdot R^{\frac{n+1}{\min\{\lambda_1,\lambda_2\}}}, & \lambda_1 \neq \lambda_2 \end{cases}$$

for some $c_{\lambda_1,\lambda_2} > 0$.

Here $\|\cdot\|$ denotes the covolume in the \mathbb{R} -subspace spanned by the subgroup with respect to the standard Euclidean norm.

Remark 8.2. As one sees in the proof, in the case $\lambda_1 \neq \lambda_2$, focusing happens.

Let $\mathbf{G} := \mathbf{SL}_{n+1}$, which naturally acts on \mathbb{Q}^{n+1} and its dual $(\mathbb{Q}^{n+1})^{\vee}$. Let $(\mathbf{e}_1, ..., \mathbf{e}_{n+1})$ be the standard basis of \mathbb{Q}^{n+1} and $(\mathbf{e}_1^{\vee}, ..., \mathbf{e}_{n+1}^{\vee})$ be its dual basis so that $(\mathbb{Q}^{n+1})^{\vee}$ is identified with \mathbb{Q}^{n+1} under this basis. Note that \mathbb{Q}^{n+1} naturally embeds in $\mathbf{P}_{\mathbb{Q}}^{n+1}$ by $\mathbf{x} \mapsto [\mathbf{x}:1]$. Let $y_0 := ([\mathbf{e}_1:1], [\mathbf{e}_1^{\vee}:1])$ and **H** be its stabilizer in **G**.

Let **X** be the Zariski closure of $\mathbf{U} := \mathbf{G}.y_0$ in $\mathbf{P}_{\mathbb{Q}}^{n+1} \times \mathbf{P}_{\mathbb{Q}}^{n+1}$ and **D** be the complement of **U** in **X**. Write $([\mathbf{x}:s], [\boldsymbol{\alpha}:t])$ for a point in **X**. Define the following subvarieties:

- 1. \mathbf{D}_1 is defined by s = 0 and \mathbf{D}_2 by t = 0;
- 2. \mathbf{S}_1 (resp. \mathbf{S}_2) is defined by s = 0 but $t \neq 0$ (resp. t = 0 but $s \neq 0$);
- 3. \mathbf{S}_1^* (resp. \mathbf{S}_1^0) is the subvariety of \mathbf{S}_1 such that $\boldsymbol{\alpha} \neq \mathbf{0}$ (resp. $\boldsymbol{\alpha} = \mathbf{0}$). \mathbf{S}_2^* and \mathbf{S}_2^0 are similarly defined.

Lemma 8.3. The pair (\mathbf{X}, \mathbf{D}) is smooth. Moreover, \mathbf{D}_1 and \mathbf{D}_2 are the irreducible components of \mathbf{D} whose intersection is non-empty and irreducible.

As for real points, we have

Lemma 8.4. With respect to the analytic topology, $\mathbf{U}(\mathbb{R})$ is connected and dense in $\mathbf{X}(\mathbb{R})$.

For convenience we fix some representatives in each G-orbit as follows

- $y_0 := ([\mathbf{e}_1 : 1], [\mathbf{e}_1^{\vee} : 1])$ as above is a representative of $\mathbf{U}(\mathbb{R})$;
- $y_1^* := ([\mathbf{e}_2:0], [\mathbf{e}_1^{\vee}:1]) \in \mathbf{S}_1^*(\mathbb{R}) \text{ and } y_2^* := ([\mathbf{e}_1:1], [\mathbf{e}_2^{\vee}:0]) \in \mathbf{S}_2^*(\mathbb{R});$ $y_1^0 := ([\mathbf{e}_2:0], [\mathbf{0}:1]) \in \mathbf{S}_1^0(\mathbb{R}) \text{ and } y_2^0 := ([\mathbf{0}:1], [\mathbf{e}_2^{\vee}:0]) \in \mathbf{S}_2^0(\mathbb{R});$
- $y_{12} := ([\mathbf{e}_1 : 0], [\mathbf{e}_2^{\vee} : 0]) \in \mathbf{D}_1(\mathbb{R}) \cap \mathbf{D}_2(\mathbb{R}).$

Lemma 8.5. Under the G-action, we have

$$\begin{aligned} \mathbf{X}(\mathbb{R}) &= \mathbf{U}(\mathbb{R}) \bigsqcup \mathbf{S}_1^*(\mathbb{R}) \bigsqcup \mathbf{S}_1^0(\mathbb{R}) \bigsqcup \mathbf{S}_2^*(\mathbb{R}) \bigsqcup \mathbf{S}_2^0(\mathbb{R}) \bigsqcup (\mathbf{D}_1(\mathbb{R}) \cap \mathbf{D}_2(\mathbb{R})) \\ &= \mathbf{G}.y_0 \bigsqcup \mathbf{G}.y_1^* \bigsqcup \mathbf{G}.y_1^0 \bigsqcup \mathbf{G}.y_2^* \bigsqcup \mathbf{G}.y_2^0 \bigsqcup \mathbf{G}.y_{12}. \end{aligned}$$

Let $\Gamma := \mathbf{SL}_{n+1}(\mathbb{Z})$. Let \mathbf{M}_1 be the stabilizer of \mathbf{e}_1 in \mathbf{G} and \mathbf{M}_2 be the stabilizer of \mathbf{e}_1^{\vee} . We define a map Φ from $\mathbf{X}(\mathbb{R})$ to $\operatorname{Prob}(G/\Gamma) \cup \{\mathbf{0}\}$ (here **0** denotes the zero measure) by

$$\Phi(x) = \begin{cases} g_* \mathbf{m}_{[\mathrm{H}]}^{\mathbf{1}} & x = g.y_0 \in \mathbf{U}(\mathbb{R}) \\ g_* \mathbf{m}_{[\mathrm{M}_2]}^{\mathbf{1}} & x = g.y_1^* \in \mathbf{S}_1^*(\mathbb{R}) \\ g_* \mathbf{m}_{[\mathrm{M}_1]}^{\mathbf{1}} & x = g.y_2^* \in \mathbf{S}_2^*(\mathbb{R}) \\ \mathbf{0} & x \in \mathbf{S}_1^0(\mathbb{R}) \cup \mathbf{S}_2^0(\mathbb{R}) \\ \mathbf{m}_{[\mathrm{G}]}^{\mathbf{1}} & x \in \mathbf{D}_1(\mathbb{R}) \cap \mathbf{D}_2(\mathbb{R}). \end{cases}$$

Lemma 8.6. The map Φ is well-defined (independent of the choice of g) and continuous.

Let $\omega_{\mathbf{U}}$ be the **G**-invariant gauge form on **U**.

Lemma 8.7. The anti-canonical divisor is $-\operatorname{div}(\omega_{\mathbf{U}}) = (n+2)\mathbf{D}_1 + (n+2)\mathbf{D}_2$.

See Section A.2.2 for the proof.

Finally, note the metric line bundle underlying the height function.

Lemma 8.8. There exist smooth metrics $\|\cdot\|$ on $\mathcal{O}_{\mathbf{X}}(\mathbf{D}_i)$ for i = 1, 2 such that for $(v, M) \in \Lambda(n, 1)$,

/2

$$\|\mathbf{1}_{\mathbf{D}_{1}}\|_{(v,M)} = \left(\sum v_{i}^{2}\right)^{-1/2}, \quad \|\mathbf{1}_{\mathbf{D}_{2}}\|_{(v,M)} = \left(\sum \alpha_{i}^{2}\right)^{-1}$$

where $v = \sum v_i \mathbf{e}_i$ and $M = \sum \alpha_i \mathbf{e}_i^{\vee}$.

See Section A.2.4 for details on this construction.

Proof of Theorem 8.1. From Lemma 8.8, there exists a smooth metric $\|\cdot\|$ on $\mathcal{O}_{\mathbf{X}}(\lambda_1\mathbf{D}_1 + \lambda_2\mathbf{D}_2)$ such that

$$\left\|\mathbf{1}_{\mathbf{D}_{1}}^{\otimes\lambda_{1}}\otimes\mathbf{1}_{\mathbf{D}_{2}}^{\otimes\lambda_{2}}\right\|_{(v,M)}=\left\|v\right\|^{-\lambda_{1}}\left\|M\right\|^{-\lambda_{2}}$$

By Lemma 8.6 and 5.6, there exists some c > 0 such that

$$\#\{(v,M) \in \Lambda(n,1), \|v\|^{\lambda_1} \|M\|^{\lambda_2} \le R\} \sim c \cdot m_{G/H}(B_{R,y_0}).$$

Thus the conclusion follows from Theorem 2.16 and Lemma 8.7.

9. Example III, the space of triangles

In this section we consider the space of three linearly independent lines on \mathbb{Q}^3 . In fact, there exists a \mathbb{Q} -variety \mathbf{M}_3 such that

$$\mathbf{M}_3(\mathbb{Q}) = \left\{ (\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3) \in (\mathbf{P}^2(\mathbb{Q}))^3 \mid (\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3) \text{ are linearly independent} \right\}.$$

One can show that \mathbf{M}_3 is dense in $\mathbf{P}^2_{\mathbb{Q}} \times \mathbf{P}^2_{\mathbb{Q}} \times \mathbf{P}^2_{\mathbb{Q}}$. Let \mathbf{D}^0 be its complement. $\mathbf{P}^2_{\mathbb{Q}}$ has a natural integral model over \mathbb{Z} , denoted as $\mathbf{P}^2_{\mathbb{Z}}$, where $\mathbf{P}^2_{\mathbb{Q}}$ is viewed as the generic fibre. Let \mathcal{D}^0 be the closure of \mathbf{D}^0 in $\mathbf{P}^2_{\mathbb{Z}} \times \mathbf{P}^2_{\mathbb{Z}} \times \mathbf{P}^2_{\mathbb{Z}}$. Let \mathcal{M}_3 be the complement of \mathcal{D}^0 in $\mathbf{P}^2_{\mathbb{Z}} \times \mathbf{P}^2_{\mathbb{Z}} \times \mathbf{P}^2_{\mathbb{Z}}$. Then one can verify that, viewed as a subset of $\mathbf{M}_3(\mathbb{Q})$ as above,

$$\mathcal{M}_{3}(\mathbb{Z}) = \left\{ (\mathbf{l}_{1}, \mathbf{l}_{2}, \mathbf{l}_{3}) \in \mathbf{M}_{3}(\mathbb{Q}) \mid \exists \mathbf{v}_{i} \in \mathbb{Z}^{3} \text{ with } \mathbf{l}_{i} = \mathbb{Q}.\mathbf{v}_{i}, \text{ det}(\mathbf{v}_{1}, \mathbf{v}_{2}, \mathbf{v}_{3}) = \pm 1 \right\} \\ = \left\{ (\mathbf{l}_{1}, \mathbf{l}_{2}, \mathbf{l}_{3}) \in \mathbf{M}_{3}(\mathbb{Q}) \mid \mathbb{Z}^{3} = (\mathbf{l}_{1} \cap \mathbb{Z}^{3}) \oplus (\mathbf{l}_{2} \cap \mathbb{Z}^{3}) \oplus (\mathbf{l}_{3} \cap \mathbb{Z}^{3}) \right\}.$$

To measure the complexity of an element in $\mathcal{M}_3(\mathbb{Z})$, we introduce the following functions:

For a discrete \mathbb{Z} -submodule Λ of \mathbb{R}^3 , let

$$\|\Lambda\| := \operatorname{Vol}(\Lambda \otimes_{\mathbb{Z}} \mathbb{R}/\Lambda)$$

where $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ is naturally embedded in \mathbb{R}^3 as a linear subspace and Vol is induced from the standard Euclidean metric on \mathbb{R}^3 . By convention, $\|\{\mathbf{0}\}\| := 1$. Let $(\Lambda_1, \Lambda_2, \Lambda_3)$ be three rank one free \mathbb{Z} -submodules of \mathbb{R}^3 that form an \mathbb{R} -linear basis of \mathbb{R}^3 . For $I \subset \{1, 2, 3\}$, let $\Lambda_I := \bigoplus_{i \in I} \Lambda_i$. By default, $\Lambda_{\emptyset} := \{\mathbf{0}\}$. For $I, J \subset \{1, 2, 3\}$,

$$d_{IJ}((\Lambda_i)_{i=1}^3) := \frac{\|\Lambda_I\| \cdot \|\Lambda_J\|}{\|\Lambda_{I \cap J}\| \|\Lambda_{I \cup J}\|}.$$

Finally, for $\mathbf{x} = (\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3) \in \mathbf{M}_3(\mathbb{R})$, take some non-zero $\mathbf{v}_i \in \mathbf{l}_i$, and let $\Lambda_i := \mathbb{Z} \cdot \mathbf{v}_i$. For l = 1, 2, define

$$\operatorname{Ht}_{l}(\mathbf{x}) := \prod_{I,J \subset \{1,2,3\}, |I|=|J|=l} d_{IJ}((\Lambda_{i})_{i=1}^{3}).$$

One can check that this definition is independent of the choice of Λ_i 's.

Now we define weights for points in $\mathcal{M}_3(\mathbb{Z})$. Take some $\mathbf{x} = (\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3) \in \mathcal{M}_3(\mathbb{Z})$ and let $\Lambda_i := \mathbf{l}_i \cap \mathbb{Z}^3$. Consider, for $\eta > 0$,

$$\Omega_{\mathbf{x},\eta} := \left\{ \mathbf{t} = (t_1, t_2, t_3) \in \mathbb{R}^3 \ \middle| \ t_1 + t_2 + t_3 = 0, \ \sum_{i \in I} t_i \ge -\ln \|\Lambda_I\| + \ln \eta, \ \forall I \subset \{1, 2, 3\} \right\}$$

where summation over an empty set is set to be 0. The weight function is just

$$\mathbf{w}_{\mathbf{x}} = \min\{\operatorname{Vol}(\Omega_{\mathbf{x},\eta})^{-1}, 1\}.$$

Theorem 9.1. Let $\kappa_1, \kappa_2 > 0$ and $\operatorname{Ht}(\mathbf{x}) := \operatorname{Ht}_1(\mathbf{x})^{\kappa_1} \operatorname{Ht}_2(\mathbf{x})^{\kappa_2}$ for $\mathbf{x} \in \mathbf{M}_3(\mathbb{R})$. Then for $\eta > 0$ small enough and some constant $c_{\kappa_1,\kappa_2} > 0$,

$$\sum_{\{\mathbf{x}\in\mathcal{M}_3(\mathbb{Z})\mid \operatorname{Ht}(x)\leq R\}} \mathbf{w}_x \sim \begin{cases} c_{\kappa_1,\kappa_2} \cdot R^{\frac{8}{3}\max\{\kappa_1^{-1},\kappa_2^{-1}\}} & \kappa_1\neq\kappa_2\\ c_{\kappa_1,\kappa_2} \cdot R^{\frac{8}{3}\kappa_1^{-1}} \cdot \log(R) & \kappa_1=\kappa_2. \end{cases}$$

Remark 9.2. Without weights, we expect an additional $\log(R)^2$ factor in the asymptotic. We hope to discuss this in a future work. This counting problem has been studied in [SZ19] with respect to a different height function.

It will be clear that the counting problem is naturally related to the embedding of M_3 into the following incidence variety:

$$\mathbf{X}^{\text{INC}} = \left\{ (\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3, \mathbf{l}_{12}, \mathbf{l}_{13}, \mathbf{l}_{23}) \in (\mathbf{P}^2)^3 \times (\mathbf{Gr}_{2,3})^3 \mid \mathbf{l}_I \subset \mathbf{l}_J, \ \forall I \subset J \right\}.$$

The morphism $\mathbf{M}_3 \rightarrow \mathbf{X}^{\text{INC}}$ is given by mapping $(\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3)$ to $(\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3, \mathbf{l}_{12}, \mathbf{l}_{13}, \mathbf{l}_{23})$ where \mathbf{l}_{ij} is the unique plane spanned by l_i and l_j . One can check that the morphism is an open embedding with dense image. As points in \mathbf{X}^{INC} are tuple of points and lines in \mathbf{P}^2 satisfying incidence conditions, they describe "triangles" in \mathbf{P}^2 . The space is thus sometimes referred to as "space of triangles". It is also related to certain counting problem studied by Schubert, see [Sem54, RS84, Rob88] for further discussions.

The space \mathbf{X}^{INC} is naturally stratified. Let

$$\begin{split} \mathbf{D}_{123}^1 &:= \left\{ \mathbf{l}_1 = \mathbf{l}_2 = \mathbf{l}_3 \right\}, \ \mathbf{D}_{123}^2 &:= \left\{ \mathbf{l}_{13} = \mathbf{l}_{13} = \mathbf{l}_{23} \right\}, \\ \mathbf{D}_{12,3} &:= \left\{ \mathbf{l}_1 = \mathbf{l}_2, \, \mathbf{l}_{13} = \mathbf{l}_{23} \right\}, \ \mathbf{D}_{13,2} &:= \left\{ \mathbf{l}_1 = \mathbf{l}_3, \, \mathbf{l}_{12} = \mathbf{l}_{23} \right\}, \\ \mathbf{D}_{23,1} &:= \left\{ \mathbf{l}_2 = \mathbf{l}_3, \, \mathbf{l}_{12} = \mathbf{l}_{13} \right\}. \end{split}$$

Also let the corresponding $\mathbf{S}^{\star}_{\bullet}$ denote points contained in $\mathbf{D}^{\star}_{\bullet}$ but not the other \mathbf{D} 's. Let $\mathbf{X}^1 := \mathbf{Bl}_{\mathbf{D}^1_{123}}(\mathbf{X}^{\mathrm{INC}})$ and \mathbf{D}^1 be the complement of \mathbf{M}_3 in \mathbf{X}^1 .

Lemma 9.3. The pair $(\mathbf{X}^1, \mathbf{D}^1)$ is smooth. Irreducible components of \mathbf{D}^1 are

$$(\mathbf{E},\mathbf{D}^+_{12,3},\mathbf{D}^+_{13,2},\mathbf{D}^+_{23,1},(\mathbf{D}^2_{123})^+)$$

where **E** denotes the exceptional divisor, same as the inverse image of \mathbf{D}_{123}^1 . Moreover, the intersection pattern is given as follows: $\mathbf{D}_{ij,k}^+$'s do not intersect with each other and for each i, j, k, the triple $\left(\mathbf{D}_{ij,k}^+, \mathbf{E}, (\mathbf{D}_{123}^2)^+\right)$ intersects transversally.

The proof is in Section A.3.6. We henceforth label $(\mathbf{E}, \mathbf{D}_{12,3}^+, \mathbf{D}_{13,2}^+, \mathbf{D}_{23,1}^+, (\mathbf{D}_{123}^2)^+)$ as $(\mathbf{D}_1, ..., \mathbf{D}_5)$. To continue, note that $\mathbf{G} := \mathbf{SL}_3$ acts on \mathbf{M}_3 and also \mathbf{X}^{INC} by linear transformations, making \mathbf{M}_3 into a homogeneous space. If $o \in \mathbf{M}_3(\mathbb{Z})$ is the base point $(\mathbb{Z}\mathbf{e}_1, \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_3)$, then the stabilizer **H** of oin **G** is equal to the full diagonal torus in SL_3 .

Let $\omega_{\mathbf{M}_3}$ be the invariant gauge form, then

Lemma 9.4. In \mathbf{X}^1 , the anti-canonical divisor is $-\operatorname{div}(\omega_{\mathbf{M}_3}) = 9\mathbf{D}_1 + 6\mathbf{D}_2 + 6\mathbf{D}_3 + 6\mathbf{D}_4 + 9\mathbf{D}_5$.

See Section A.3.9 for the proof.

Let ψ be a non-negative function on G/Γ (here $G := \mathbf{SL}_3(\mathbb{R})$ and $\Gamma := \mathbf{SL}_3(\mathbb{Z})$) whose support is large enough. Now we construct a continuous map from $\mathbf{X}^{\text{INC}}(\mathbb{R})$ to $\text{Prob}^{\psi}(G/\Gamma)$. Let

- $\mathbf{x}_1 := (\mathbb{Z}\mathbf{e}_1, \mathbb{Z}\mathbf{e}_1, \mathbb{Z}\mathbf{e}_1, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_3, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}(\mathbf{e}_2 + \mathbf{e}_3)) \text{ in } \mathbf{S}^1_{123}(\mathbb{R});$
- $\mathbf{x}_2 := (\mathbb{Z}\mathbf{e}_1, \mathbb{Z}\mathbf{e}_1, \mathbb{Z}\mathbf{e}_3, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_3, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_3)$ in $\mathbf{S}_{12,3}(\mathbb{R})$;
- $\mathbf{x}_3 := (\mathbb{Z}\mathbf{e}_1, \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_1, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_3, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_2) \text{ in } \mathbf{S}_{13,2}(\mathbb{R});$
- $\mathbf{x}_4 := (\mathbb{Z}\mathbf{e}_1, \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_2 \oplus \mathbb{Z}\mathbf{e}_3) \text{ in } \mathbf{S}_{23,1}(\mathbb{R});$
- $\mathbf{x}_5 := (\mathbb{Z}\mathbf{e}_1, \mathbb{Z}\mathbf{e}_2, \mathbb{Z}(\mathbf{e}_1 + \mathbf{e}_2), \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_2, \mathbb{Z}\mathbf{e}_1 \oplus \mathbb{Z}\mathbf{e}_2) \text{ in } \mathbf{S}^2_{123}(\mathbb{R}).$

Lemma 9.5. We have

$$\mathbf{X}^{\mathrm{INC}}(\mathbb{R}) = \mathrm{G.}o \bigsqcup \mathrm{G.}x_2 \bigsqcup \mathrm{G.}x_3 \bigsqcup \mathrm{G.}x_4 \bigsqcup (\mathbf{D}_{123}^1(\mathbb{R}) \cup \mathbf{D}_{123}^2(\mathbb{R})).$$

For a partition $\{i, j\} \sqcup \{k\}$ of $\{1, 2, 3\}$, let \mathbf{H}_{ij} be the simultaneous stabilizer in **G** of the line spanned by \mathbf{e}_k and the plane spanned by $\mathbf{e}_i, \mathbf{e}_j$. Define $\Phi_{\mathbf{H}} : \mathbf{X}^{\text{INC}}(\mathbb{R}) \to \text{Prob}^{\psi}(\mathbf{G}/\Gamma)$ by

$$\Phi_{\mathbf{H}}(\mathbf{x}) := \begin{cases} \alpha_{g}^{\psi}(\mathbf{m}_{[\mathrm{H}]}) & \text{if } \mathbf{x} = g.o \in \mathbf{M}_{3}(\mathbb{R}) \\ \alpha_{g}^{\psi}(\mathbf{m}_{[\mathrm{H}_{12}]}) & \text{if } \mathbf{x} = g.x_{2} \in \mathbf{S}_{12,3}(\mathbb{R}) \\ \alpha_{g}^{\psi}(\mathbf{m}_{[\mathrm{H}_{13}]}) & \text{if } \mathbf{x} = g.x_{3} \in \mathbf{S}_{13,2}(\mathbb{R}) \\ \alpha_{g}^{\psi}(\mathbf{m}_{[\mathrm{H}_{23}]}) & \text{if } \mathbf{x} = g.x_{4} \in \mathbf{S}_{23,1}(\mathbb{R}) \\ \mathbf{m}_{[\mathrm{G}]}^{\psi} & \text{if } \mathbf{x} \in \mathbf{D}_{123}^{1}(\mathbb{R}) \cup \mathbf{D}_{123}^{2}(\mathbb{R}) \end{cases}$$

Lemma 9.6. The map $\Phi_{\mathbf{H}}$ is well-defined and continuous.

This is a corollary to discussions in Section A.3.10. For $\{i, j\} \subset \{1, 2, 3\}$, let

$$\mathbf{X}_{ij}^{\mathrm{INC}} := \left\{ (\mathbf{l}_i, \mathbf{l}_j, \mathbf{l}_{ij}) \in \mathbf{P}^2 \times \mathbf{P}^2 \times \mathbf{Gr}_{2,3} \, \big| \, \mathbf{l}_i, \mathbf{l}_j \subset \mathbf{l}_{ij} \right\}$$

together with a natural morphism $\pi_{ij} : \mathbf{X}^{\text{INC}} \to \mathbf{X}_{ij}^{\text{INC}}$. Let \mathbf{M}_{ij} denote the open subvariety where \mathbf{l}_i and \mathbf{l}_j are linearly independent and \mathbf{D}_{ij} be its complement.

Similarly, for $\{\{i, j\}, \{i, k\}\} \subset \{\{1, 2\}, \{1, 3\}, \{2, 3\}\}$, we define $\mathbf{X}_{ij,ik}^{\text{INC}}$, a subvariety of $\mathbf{P}^2 \times \mathbf{Gr}_{2,3} \times \mathbf{Gr}_{2,3}$, and $\pi_{ij,ik} : \mathbf{X}^{\text{INC}} \to \mathbf{X}_{ij,ik}^{\text{INC}}$. Let $\mathbf{M}_{ij,ik}$ denote the open subvariety where $\mathbf{l}_{ij} \neq \mathbf{l}_{ik}$ and $\mathbf{D}_{ij,ik}$ its complement.

Lemma 9.7. The pair $(\mathbf{X}_{ij}^{\text{INC}}, \mathbf{D}_{ij})$ is smooth. Moreover, there exists a smooth metric on $\mathcal{O}_{\mathbf{X}_{ij}^{\text{INC}}}(\mathbf{D}_{ij})$, as a real analytic line bundle, such that if \mathbf{v}_i and \mathbf{v}_j are non-zero vectors on \mathbf{l}_i and \mathbf{l}_j respectively, then for $\mathbf{x} = (\mathbf{l}_i, \mathbf{l}_j, \mathbf{l}_{ij})$ in $\mathbf{M}_{ij}(\mathbb{R})$,

$$\left\|\mathbf{1}_{\mathbf{D}_{ij}}\right\|_{\mathbf{x}} = \frac{\left\|\mathbf{v}_{i} \wedge \mathbf{v}_{j}\right\|}{\left\|\mathbf{v}_{i}\right\| \left\|\mathbf{v}_{j}\right\|}.$$

Similarly, the pair $(\mathbf{X}_{ij,ik}^{\text{INC}}, \mathbf{D}_{ij,ik})$ is smooth. And $\mathcal{O}_{\mathbf{X}_{ij,ik}^{\text{INC}}}(\mathbf{D}_{ij,ik})$ can be equipped with a smooth metric such that

$$\left\|\mathbf{1}_{\mathbf{D}_{ij,ik}}\right\|_{\mathbf{x}} = \frac{\left\|\mathbf{v}_{i}\right\| \left\|\mathbf{v}_{i} \wedge \mathbf{v}_{j} \wedge \mathbf{v}_{k}\right\|}{\left\|\mathbf{v}_{i} \wedge \mathbf{v}_{j}\right\| \left\|\mathbf{v}_{i} \wedge \mathbf{v}_{k}\right\|}$$

where \mathbf{l}_{ij} is spanned by $\mathbf{v}_i, \mathbf{v}_j$ and \mathbf{l}_{ik} is spanned by $\mathbf{v}_i, \mathbf{v}_k$.

Let \mathbf{s}_{ij} be the pull-back of the section $\mathbf{1}_{\mathbf{D}_{ij}}$ to a section of the pull-back of $\mathcal{O}_{\mathbf{X}_{ij}^{\text{INC}}}(\mathbf{D}_{ij})$ to a line bundle over \mathbf{X}^1 . Similarly define $\mathbf{s}_{ij,ik}$.

Lemma 9.8. For $\{i, j\} \subset \{1, 2, 3\}$, div $(\mathbf{s}_{ij}) = \mathbf{E} + \mathbf{D}_{ij,k}^+$ where k is such that $\{i, j, k\} = \{1, 2, 3\}$. And div $(\mathbf{s}_{ij,ik}) = (\mathbf{D}_{123}^2)^+ + \mathbf{D}_{ik,i}^+$.

Proof of Theorem 9.1. By Lemma 5.1, it suffices to check condition (B1) and (BH1). (B1) follows from Lemma 9.6. For any (κ_1, κ_2) ,

$$\max\left\{\frac{9-1}{\kappa_1}, \frac{9-1}{\kappa_2}\right\} > \frac{6-1}{\kappa_1 + \kappa_2},$$

hence the limit

$$\nu := \lim_{R \to \infty} \frac{\mathbf{m}_{\mathrm{G/H}} \cdot \mathbf{1}_{B_{R,o}}}{\mathbf{m}_{\mathrm{G/H}}(B_{R,o})}$$

exists and is supported on $\mathbf{E}(\mathbb{R}) \cup (\mathbf{D}_{123}^2)^+(\mathbb{R})$ by Lemma 9.4 and Theorem 2.16. Consequently, (BH1) holds.

Remark 9.9. Let $\mathbf{L} := \sum_{i=1}^{5} \lambda_i \mathbf{D}_i$ be a divisor on \mathbf{X}^1 with $\lambda_i > 0$. If we count with respect to a height function associated with \mathbf{L} , then the focusing phenomenon happens exactly when $\min\{\lambda_1, \lambda_2\} > \frac{4}{3}\min\{\lambda_3, \lambda_4, \lambda_5\}$.

Appendix A. Details on the Examples I, II, III

A.1. Example I. Here we fill in the various missing details from Section 7.

A.1.1. Change of coordinates.

Definition A. • Define quadratic forms Q_1^0 and Q_2^0 by $Q_1^0(x_1, x_2) = 2x_1x_2$ and $Q_2^0(x_1, x_2, x_3, x_4) = 2x_1x_4 + 2x_2x_3$;

 Identify vectors with column vectors and linear maps with matrices. Let M_{Q_i⁰} (i = 1, 2) be the matrix representations of Q_i⁰ (i = 1, 2), we have

$$M_{Q_1^0} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad M_{Q_2^0} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix};$$

• Let x_0 be the point $([M_0 : \lambda_0], \langle N_0 \rangle)$ with $\lambda_0 = 1$,

$$M_0 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad N_0 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}.$$

Note that (Q_1, \mathbb{R}^2) is isomorphic to (Q_1^0, \mathbb{R}^2) and (Q_2, \mathbb{R}^4) is isomorphic to (Q_2^0, \mathbb{R}^4) . Therefore, over \mathbb{R} , we may and do replace Q_2 , Q_1 by Q_2^0 and Q_1^0 in the definition of \mathbf{X}^{INC} .

A.1.2. Local coordinates: ideals of the closure. Without loss of generality, we assume that $([M : \lambda], \langle N \rangle)$ lies in an open affine subvariety **O** where, since $\langle M \rangle$ is contained in $\langle N \rangle$, they take the form

$$M = \begin{bmatrix} 1 & \beta_1 \\ \alpha_2 & \beta_2 \\ x_1 + \alpha_2 y_1 & \beta_1 x_1 + \beta_2 y_1 \\ x_2 + \alpha_2 y_2 & \beta_1 x_2 + \beta_2 y_2 \end{bmatrix}, N = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ x_1 & y_1 \\ x_2 & y_2 \end{bmatrix}.$$
 (32)

Hence we regard \mathbf{O} as a closed subvariety of the affine space

Spec $\mathbb{R}[\alpha_2, \beta_1, \beta_2, \lambda, x_1, x_2, y_1, y_2].$

We need to find the (prime) ideals that correspond to \mathbf{X}^{INC} and various strata. By restricting to the open subvariety $\lambda \neq 0, \beta_2 - \beta_1 \alpha_2 \neq 0$, we have the equations (write $M = [\boldsymbol{\alpha}, \boldsymbol{\beta}]$)

$$\begin{cases} Q_{2}^{0}(\boldsymbol{\alpha},\boldsymbol{\alpha}) = 0\\ Q_{2}^{0}(\boldsymbol{\alpha},\boldsymbol{\beta}) = \lambda^{2} \iff \begin{cases} Q_{2}^{0}(\boldsymbol{\alpha},\boldsymbol{\alpha}) = 0\\ Q_{2}^{0}(\boldsymbol{\alpha},\boldsymbol{\beta}) = \lambda^{2}\\ Q_{2}^{0}(\boldsymbol{\beta},\boldsymbol{\beta}) = 0 \end{cases} & \begin{cases} Q_{2}^{0}(\boldsymbol{\alpha},\boldsymbol{\alpha}) = 0\\ Q_{2}^{0}(\boldsymbol{\alpha},\boldsymbol{\beta}-\beta_{1}\boldsymbol{\alpha}) = \lambda^{2}\\ Q_{2}^{0}(\boldsymbol{\beta}-\beta_{1}\boldsymbol{\alpha}) + 2\beta_{1}\lambda^{2} = 0 \end{cases} \\ \Leftrightarrow & \begin{cases} (x_{2}+\alpha_{2}y_{2}) + \alpha_{2}(x_{1}+\alpha_{2}y_{1}) = 0\\ (\beta_{2}-\beta_{1}\alpha_{2})(y_{2}+x_{1}+2\alpha_{2}y_{1}) - \lambda^{2}\\ (\beta_{2}-\beta_{1}\alpha_{2})[(\beta_{2}-\beta_{1}\alpha_{2})y_{1}+\beta_{1}(y_{2}+x_{1}+2\alpha_{2}y_{1})] = 0 \end{cases} \\ \Leftrightarrow & \begin{cases} (x_{2}+\alpha_{2}y_{2}) + \alpha_{2}(x_{1}+\alpha_{2}y_{1}) = 0\\ (\beta_{2}-\beta_{1}\alpha_{2})(y_{2}+x_{1}+2\alpha_{2}y_{1}) - \lambda^{2} = 0 \end{cases} & (1)\\ (\beta_{2}-\beta_{1}\alpha_{2})(y_{2}+x_{1}+2\alpha_{2}y_{1}) - \lambda^{2} = 0 \end{cases} \\ (2) \end{cases} \end{cases}$$

The last " \iff " is because $\beta_2 - \beta_1 \alpha_2$ is invertible in this open subvariety. If $\mathcal{I}_{\mathbf{X}^{\text{INC}}}$ stands for the prime ideal corresponding to **O**, then we have seen that

$$\begin{cases} (x_2 + \alpha_2 y_2) + \alpha_2 (x_1 + \alpha_2 y_1), \\ (\beta_2 - \beta_1 \alpha_2) (y_2 + x_1 + 2\alpha_2 y_1) - \lambda^2, \\ (\beta_2 - \beta_1 \alpha_2) y_1 + \beta_1 (y_2 + x_1 + 2\alpha_2 y_1) \end{cases} \subset \mathcal{I}_{\mathbf{X}^{\text{INC}}}$$

It will be shown that $\mathcal{I}_{\mathbf{X}^{\text{INC}}}$ is actually generated by these polynomials. This follows once we know that the ideal generated by the left hand side, denoted as \mathcal{I}_0 , is prime. To see this, we first do a few change of variables.

By equations above, we eliminate x_2 and replace β_2 and y_2 by

$$\beta_2' := \beta_2 - \beta_1 \alpha_2, \ y_2' := y_2 + x_1 + 2\alpha_2 y_1.$$
(33)

Then **O** is the closed subvariety of Spec $\mathbb{R}[\lambda, \beta_1, \beta'_2, y_1, y'_2, \alpha_2, x_1]$ corresponding to the smallest prime ideal containing

$$\mathcal{I}_0 = \langle \beta_2' y_2' - \lambda^2, \beta_2' y_1 + \beta_1 y_2' \rangle. \tag{34}$$

Lemma A.1. In $R := \mathbb{C}[x, y, a, b, w]$, the ideal I generated by $xy - w^2$ and xa + yb is prime. Hence \mathcal{I}_0 is a prime ideal and $\mathcal{I}_0 = \mathcal{I}_{\mathbf{X}^{\text{INC}}}$.

Proof. Once we know \mathcal{I}_0 is prime, the closed subscheme X_0 cut out by \mathcal{I}_0 is irreducible. In particular, the closure of $X_0 \cap \{\lambda \neq 0\}$ is X_0 . But \mathbf{X}^{INC} , by definition, is the closure of $\mathbf{X}^{\text{INC}} \cap \{\lambda \neq 0\} = X_0 \cap \{\lambda \neq 0\}$, so $X_0 = \mathbf{X}^{\text{INC}}$ and $\mathcal{I}_0 = \mathcal{I}_{\mathbf{X}^{\text{INC}}}$.

$$x \cdot \phi_0 = \phi_1(xy - w^2) + \phi_2(xa + yb). \tag{35}$$

Quotienting by $\langle x \rangle$, we have

$$\overline{\phi}_1 w^2 = \overline{\phi}_2 y b.$$

Since $\mathbb{C}[y, w, a, b]$ is a unique factorization domain, we have

$$\phi_1 = x\phi_3 + yb\phi_4, \ \phi_2 = x\phi_5 + w^2\phi_4$$

for some $\phi_3, \phi_4, \phi_5 \in \mathbb{R}$. Inserting back to Equa.(35), we have

$$\begin{aligned} x \cdot \phi_0 &= x \cdot [\phi_3(xy - w^2) + \phi_5(xa + yb) + \phi_4(y^2b + w^2a)] \\ \implies \phi_0 &= \phi_3(xy - w^2) + \phi_5(xa + yb) + \phi_4(y^2b + w^2a) \\ &= \phi_3(xy - w^2) + \phi_5(xa + yb) + \phi_4(-a(xy - w^2) + y(xa + yb)) \in I \end{aligned}$$

So x is not a zero divisor modulo I. By inverting x, we see that $\langle I \rangle = \langle y - x^{-1}w^2, a + x^{-1}yb \rangle$ in $\mathbb{R}[x, x^{-1}, y, w, a, b]$. In particular, it is a prime ideal. So if $\phi \cdot \psi \in I$, then there exists n such that $x^n \phi \in I$ or $x^n \psi \in I$. But x is not a zero divisor modulo I, thus, $\phi \in I$ or $\psi \in I$. This shows that I is prime.

A.1.3. Local coordinates: ideals of the stratum. Now we give the explicit local equations of $\mathbf{D}_{a,b,c}$. Let $I_{a,b,c}$ be the ideal corresponding to $\mathbf{D}_{a,b,c}$ in the local affine subscheme given by Spec $\mathbb{R}[\lambda, \beta_1, \beta'_2, y_1, y'_2, \alpha_2, x_1]$, same as last subsection.

Lemma A.2.

$$\begin{split} I_{2,1,0^+} &= \langle \lambda, \beta_1, \beta_2' \rangle, \ I_{1,1,1} = \langle \lambda, \beta_2', y_2' \rangle, \ I_{0,2,2} = \langle \lambda, y_1, y_2' \rangle, \ I_{1,1,0^+} = \langle \lambda, \beta_1, \beta_2', y_2' \rangle, \\ I_{0,1,1} &= \langle \lambda, y_1, y_2', \beta_2' \rangle, \ I_{0,1,0^+} = \langle \lambda, \beta_1, \beta_2', y_1, y_2' \rangle. \end{split}$$

Proof. Via column operations, transform M into

$$M' = \begin{bmatrix} 1 & 0\\ \alpha_2 & \beta'_2\\ x_1 + \alpha_2 y_1 & \beta'_2 y_1\\ x_2 + \alpha_2 y_2 & \beta'_2 y_2 \end{bmatrix}.$$

Thus $\operatorname{rk}(M) \leq 1$ iff all two minors of M' vanishes, which happens exactly when $\beta'_2 = 0$.

To detect $rk(\langle N \rangle, Q_2^0)$, write $N = [\mathbf{x}, \mathbf{y}]$ (as in Equa.(32)) and

$$B_N := \begin{bmatrix} Q_2^0(\mathbf{x}, \mathbf{x}) & Q_2^0(\mathbf{x}, \mathbf{y}) \\ Q_2^0(\mathbf{x}, \mathbf{y}) & Q_2^0(\mathbf{y}, \mathbf{y}) \end{bmatrix} = \begin{bmatrix} 2x_1 & y_2 + x_1 \\ y_2 + x_1 & 2y_1 \end{bmatrix}.$$

Thus the ideal corresponding to $\operatorname{rk}(\langle N \rangle, Q_2^0) \leq 1$ is the radical of $\langle x_2y_1 - (y_2 + x_1)^2, \lambda \rangle$, which, after eliminating x_2 , is equal to $\langle (y_2 + x_1 + 2\alpha_2y_1)^2, \lambda \rangle$. So its radical is $\langle y'_2 = y_2 + x_1 + 2\alpha_2y_1, \lambda \rangle$. And the ideal corresponding to $\operatorname{rk}(\langle N \rangle, Q_2^0) = 0$ is then $\langle x_2, y_1, y_2 + x_1, \lambda \rangle$, which, after eliminating x_2 , is equal to $\langle y_1, y'_2, \lambda \rangle$.

To see $\operatorname{rk}([M^{\operatorname{Tr}}], Q_1^0)$, first we perform row operation on M to get

$$M'' = \begin{bmatrix} 1 & \beta_1 \\ 0 & \beta'_2 \\ 0 & \beta'_2 y_1 \\ 0 & \beta'_2 y_2 \end{bmatrix} = \begin{bmatrix} \mathbf{v}_1 \\ \mathbf{v}_2 \\ \mathbf{v}_3 \\ \mathbf{v}_4 \end{bmatrix}.$$

Then we compute

$$B_{M''} = \begin{bmatrix} Q_1^0(\mathbf{v}_1, \mathbf{v}_1) & Q_1^0(\mathbf{v}_1, \mathbf{v}_2) & Q_1^0(\mathbf{v}_1, \mathbf{v}_3) & Q_1^0(\mathbf{v}_1, \mathbf{v}_4) \\ Q_1^0(\mathbf{v}_1, \mathbf{v}_2) & Q_1^0(\mathbf{v}_2, \mathbf{v}_2) & Q_1^0(\mathbf{v}_2, \mathbf{v}_3) & Q_1^0(\mathbf{v}_2, \mathbf{v}_4) \\ Q_1^0(\mathbf{v}_1, \mathbf{v}_3) & Q_1^0(\mathbf{v}_2, \mathbf{v}_3) & Q_1^0(\mathbf{v}_3, \mathbf{v}_3) & Q_1^0(\mathbf{v}_3, \mathbf{v}_4) \\ Q_1^0(\mathbf{v}_1, \mathbf{v}_4) & Q_1^0(\mathbf{v}_2, \mathbf{v}_4) & Q_1^0(\mathbf{v}_2, \mathbf{v}_3) & Q_1^0(\mathbf{v}_4, \mathbf{v}_4) \end{bmatrix} \\ = \begin{bmatrix} 2\beta_1 & \beta_2' & y_1\beta_2' & y_2\beta_2' \\ \beta_2' & 0 & 0 & 0 \\ y_1\beta_2' & 0 & 0 & 0 \\ y_2\beta_2' & 0 & 0 & 0 \end{bmatrix}.$$

Thus the ideal for $\operatorname{rk}(\langle M^{\operatorname{Tr}}\rangle, Q_1^0) \leq 1$ is $\langle \beta_2', \lambda \rangle$ and the ideal for $\operatorname{rk}(\langle M^{\operatorname{Tr}}\rangle, Q_1^0) = 0$ is $\langle \beta_1, \beta_2', \lambda \rangle$. Now it is not hard to check the claim using these computations.

A.1.4. Explicit constructions of a resolution, step I. By Lemma A.1, \mathbf{X}^{INC} restricted to this affine open subvariety corresponds to

$$\langle \beta_2' y_2' - \lambda^2, \beta_2' y_1 + \beta_1 y_2' \rangle \subset \mathbb{R}[\beta_1, \beta_2', \lambda, y_1, y_2'] \times \mathbb{R}[\alpha_2, x_1].$$

As the variables α_2, x_1 play no role in the calculation, we will omit them for simplicity. So we work inside a 5-dimensional affine space.

First we blow up \mathbf{D}_{010^+} . Then $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{X}^{\mathrm{INC}}$ can be viewed as a subvariety of $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{A}^5 \subset \mathbf{A}^5 \times \mathbf{P}^4$. If $(x_1 : x_2 : ... : x_5)$ (resp. $(x_1, x_2, ..., x_5)$) is a typical point of \mathbf{P}^4 (resp. \mathbf{A}^5), we let \mathbf{U}_i (resp. \mathbf{A}^5_i) be the open subvariety of $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{A}^5$ (resp. \mathbf{A}^5) with $x_i \neq 0$. Therefore, $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{X}^{\mathrm{INC}} \cap \mathbf{U}_i$ is viewed as a subvariety of $\mathbf{A}^1 \times \mathbf{A}^4$. Recall that we are going to ignore α_2, x_1 .

A.1.5. U₁. On U₁ := { $\widetilde{\lambda} \neq 0$ }, we let $\widetilde{\beta}_1, \widetilde{\beta}_2, \widetilde{y}_1, \widetilde{y}_2$ be the coordinates of \mathbf{A}^4 . So U₁ is identified with Spec $\mathbb{R}[\lambda] \times \operatorname{Spec} \mathbb{R}[\widetilde{\beta}_1, \widetilde{\beta}_2, \widetilde{y}_1, \widetilde{y}_2]$ and we have

- 1. $\beta_1 = \lambda \widetilde{\beta}_1, \ \beta'_2 = \lambda \widetilde{\beta}_2, \ y_1 = \lambda \widetilde{y}_1, \ y'_2 = \lambda \widetilde{y}_1;$ 2. The ideal of $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{X}^{\mathrm{INC}}$ on \mathbf{U}_1 is $\langle \widetilde{\beta}_2 \widetilde{y}_2 1, \widetilde{\beta}_2^2 \widetilde{y}_1 + \widetilde{\beta}_1 \rangle;$
- 3. The proper transforms of \mathbf{D}_{210} , \mathbf{D}_{111} and \mathbf{D}_{022} are contained in the complement;
- 4. The exceptional divisor \mathbf{E}_1 is given by $\langle \lambda \rangle$.

According to 2, we can eliminate $\tilde{\beta}_1 = -\tilde{\beta}_2^2 \tilde{y}_1$ and view $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{X}^{\text{INC}}$ on \mathbf{U}_1 as a closed subvariety of Spec $\mathbb{R}[\lambda, \widetilde{\beta}_2, \widetilde{y}_1, \widetilde{y}_2]$ defined by the ideal $\langle \widetilde{\beta}_2 \widetilde{y}_2 - 1 \rangle$. With this new coordinate,

- 3'. The proper transforms of $\mathbf{D}_{210}, \mathbf{D}_{111}, \mathbf{D}_{022}$ are contained in the complement;
- 4'. The exceptional divisor \mathbf{E}_1 is given by $\langle \lambda \rangle$.

So it is clear that we have arrived at a smooth pair on U_1 .

A.1.6. U₂. Similarly U₂ = { $\widetilde{\beta}_1 \neq 0$ } is identified with Spec $\mathbb{R}[\beta_1] \times \text{Spec } \mathbb{R}[\widetilde{\lambda}, \widetilde{\beta}_2, \widetilde{y}_1, \widetilde{y}_2]$. And

- 1. $\lambda = \beta_1 \widetilde{\lambda}, \beta'_2 = \beta_1 \widetilde{\beta}_2, y_1 = \beta_1 \widetilde{y}_1, y'_2 = \beta_1 \widetilde{y}_2;$ 2. The ideal of $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{X}^{\text{INC}}$ on \mathbf{U}_2 is $\langle \widetilde{\beta}_2 \widetilde{y}_2 \widetilde{\lambda}^2, \widetilde{\beta}_2 \widetilde{y}_1 + \widetilde{y}_2 \rangle;$ 3. The proper transform of \mathbf{D}_{210} is contained in the complement of $\mathbf{U}_2;$
- 4. The proper transform \mathbf{D}_{111}^+ is $\langle \widetilde{\lambda}, \widetilde{\beta}_2, \widetilde{y}_2 \rangle$, \mathbf{D}_{022}^+ is $\langle \widetilde{\lambda}, \widetilde{y}_1, \widetilde{y}_2 \rangle$ and \mathbf{D}_{011}^+ is $\langle \widetilde{\lambda}, \widetilde{y}_1, \widetilde{y}_2, \widetilde{\beta}_2 \rangle$; 5. The exceptional divisor \mathbf{E}_1 is given by $\langle \beta_1 \rangle$.

Eliminate $\widetilde{y}_2 = -\widetilde{\beta}_2 \widetilde{y}_1$ by 2. So view $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{X}^{\mathrm{INC}}$ on \mathbf{U}_2 as the closed subvariety of Spec $\mathbb{R}[\beta_1, \widetilde{\lambda}, \widetilde{\beta}_2, \widetilde{y}_1]$ defined by $\langle \tilde{\beta}_2^2 \tilde{y}_1 + \tilde{\lambda}^2 \rangle$. The proper transforms of divisors (on U₂) become

$$\mathbf{D}_{210}^{+}:\emptyset;\ \mathbf{D}_{111}^{+}:\langle\widetilde{\lambda},\widetilde{\beta}_{2}\rangle;\ \mathbf{D}_{022}^{+}:\langle\widetilde{\lambda},\widetilde{y}_{1}\rangle;\ \mathbf{D}_{011}^{+}:\langle\widetilde{\lambda},\widetilde{\beta}_{2},\widetilde{y}_{1}\rangle;\ \mathbf{E}_{1}:\langle\beta_{1}\rangle$$

A.1.7. U₃. U₃ is identified with Spec $\mathbb{R}[\beta'_2] \times \operatorname{Spec} \mathbb{R}[\widetilde{\lambda}, \widetilde{\beta}_1, \widetilde{y}_1, \widetilde{y}_2]$. And

- 1. $\lambda = \beta_1 \widetilde{\lambda}, \beta'_2 = \beta_1 \widetilde{\beta}_2, y_1 = \beta_1 \widetilde{y}_1, y'_2 = \beta_1 \widetilde{y}_2;$ 2. The ideal of $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{X}^{\mathrm{INC}}$ is $\langle \widetilde{y}_2 \widetilde{\lambda}^2, \widetilde{y}_1 + \widetilde{\beta}_1 \widetilde{y}_2 \rangle;$ 3. The proper transforms of $\mathbf{D}_{210}, \mathbf{D}_{111}$ are contained in the complement;
- 4. The proper transform \mathbf{D}_{022}^+ is $\langle \lambda \rangle$;

5. The exceptional divisor \mathbf{E}_1 is given by $\langle \beta_2' \rangle$.

Eliminating $\tilde{y}_2 = \tilde{\lambda}^2$, $\tilde{y}_1 = -\tilde{\beta}_1 \tilde{y}_2$ by 2, we see that $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{X}^{\mathrm{INC}} \cap \mathbf{U}_3$ is isomorphic to the affine space Spec $\mathbb{R}[\beta'_2, \tilde{\lambda}, \tilde{\beta}_1]$, with

- 3'. The proper transforms of $\mathbf{D}_{210}, \mathbf{D}_{111}$ are contained in the complement;
- 4'. The proper transform \mathbf{D}_{022}^+ is $\langle \lambda \rangle$;
- 5'. The exceptional divisor \mathbf{E}_1 is given by $\langle \beta'_2 \rangle$.

So we have arrived at a smooth pair.

A.1.8. U₄. U₄ is identified with Spec $\mathbb{R}[y_1] \times \operatorname{Spec} \mathbb{R}[\widetilde{\lambda}, \widetilde{\beta}_1, \widetilde{\beta}_2, \widetilde{y}_2]$. It turns out that $\widetilde{\beta}_2 = -\widetilde{\beta}_1 \widetilde{y}_2$ can be eliminated and we get


- 1. $\mathbf{Bl}_{\mathbf{D}_{010^+}} \mathbf{X}^{\mathrm{INC}} \cap \mathbf{U}_4$ is a closed subvariety of $\operatorname{Spec} \mathbb{R}[y_1, \widetilde{\lambda}, \widetilde{\beta}_1, \widetilde{y}_2]$ defined by $\langle \widetilde{y}_2^2 \widetilde{\beta}_1 + \widetilde{\lambda}^2 \rangle$;
- 2. The proper transform of \mathbf{D}_{022} is contained in the complement;
- 3. The proper transform \mathbf{D}_{210}^+ is $\langle \widetilde{\lambda}, \widetilde{\beta}_1 \rangle$, \mathbf{D}_{111}^+ is $\langle \widetilde{\lambda}, \widetilde{y}_2 \rangle$ and \mathbf{D}_{110}^+ is $\langle \widetilde{\lambda}, \widetilde{\beta}_1, \widetilde{y}_2 \rangle$;
- 4. The exceptional divisor \mathbf{E}_1 is given by $\langle y_1 \rangle$.

A.1.9. U₅. U₅ is identified with Spec $\mathbb{R}[y'_2] \times \text{Spec } \mathbb{R}[\widetilde{\lambda}, \widetilde{\beta}_1, \widetilde{\beta}_2, \widetilde{y}_1]$. It turns out that both $\widetilde{\beta}_1 = \widetilde{\lambda}^2, \widetilde{\beta}_2 = -\widetilde{\beta}_2 \widetilde{y}_1$ can be eliminated and

- 1. $\mathbf{Bl}_{\mathbf{D}_{010^{\pm}}} \mathbf{X}^{\text{INC}}$ on U₅ is identified with the affine space Spec $\mathbb{R}[y'_2, \widetilde{\lambda}, \widetilde{y}_1]$;
- 2. The proper transforms of \mathbf{D}_{022} and \mathbf{D}_{111} are contained in the complement of U₅;
- 3. The proper transform \mathbf{D}_{210}^+ is $\langle \lambda \rangle$.
- 4. The exceptional divisor \mathbf{E}_1 is $\langle y'_2 \rangle$.

So we also get a smooth pair here.

A.1.10. Summary. On U₁, U₃, U₅, our $\mathbf{Bl}_{\mathbf{D}_{0,1,0^+}} \mathbf{X}^{\text{INC}}$ is already a smooth pair. Below is the intersection patterns of the proper transforms of various boundary components:



where the arrow means "is contained in", or more precisely, " is the intersection of ". The exceptional divisor \mathbf{E}_1 , which does not appear, intersects all of them transversally.

A.1.11. Explicit constructions of a resolution, step II. Call $\mathbf{X}^1 := \mathbf{Bl}_{\mathbf{D}_{0,1,0^+}} \mathbf{X}^{\text{INC}}$. As the second step, we blow up $\mathbf{D}_{011}^+ \bigsqcup \mathbf{D}_{110}^+$ on \mathbf{U}_2 and \mathbf{U}_4 . Let \mathbf{E}_2^1 (resp. \mathbf{E}_2^2) the exceptional divisor corresponding to \mathbf{D}_{011}^+ (resp. \mathbf{D}_{110}^+).

A.1.12. U₂₁. Over U₂ we would have U₂₁ \cup U₂₂ \cup U₂₃, with **D**⁺₀₁₁ being blown up.

 U_{21} is identified with $\operatorname{Spec} \mathbb{R}[\beta_1, \widetilde{\lambda}] \times \operatorname{Spec} \mathbb{R}[\widetilde{\beta}_2, \widetilde{\widetilde{y}}_1]$. We have

- 1. The ideal of $\mathbf{Bl}_{\mathbf{D}_{011}^+} \mathbf{X}^1 \cap \mathbf{U}_{21}$ is $\langle \tilde{\beta}_2^2 \lambda \tilde{\widetilde{y}}_1 + 1 \rangle$;
- 2. The proper transforms of \mathbf{D}_{210}^+ , \mathbf{D}_{022}^+ and \mathbf{D}_{111}^+ are contained in the complement;
- 3. The proper transform \mathbf{E}_1^+ is $\langle \beta_1 \rangle$.
- 4. The exceptional divisor \mathbf{E}_2^1 is in the complement.

A.1.13. U₂₂. U₂₂ is identified with Spec $\mathbb{R}[\beta_1, \widetilde{\beta}_2] \times \operatorname{Spec} \mathbb{R}[\widetilde{\lambda}, \widetilde{\widetilde{y}}_1]$. We have

- 1. The ideal of $\mathbf{Bl}_{\mathbf{D}_{011}^+} \mathbf{X}^1 \cap \mathbf{U}_{22}$ is $\langle \widetilde{\beta}_2 \widetilde{\widetilde{y}}_1 + \widetilde{\widetilde{\lambda}}^2 \rangle$;
- 2. The proper transforms of \mathbf{D}_{210}^+ and \mathbf{D}_{111}^+ are contained in the complement of U_{21} ;
- 3. The proper transform \mathbf{D}_{022}^{++} (of \mathbf{D}_{022}^{+}) is $\langle \lambda, \tilde{y}_1 \rangle$.
- 4. The proper transform \mathbf{E}_{1}^{+} is $\langle \beta_{1} \rangle$ (strictly speaking, \mathbf{E}_{1}^{+} should be the intersection of the zero of $\langle \beta_{1} \rangle$ with $\mathbf{Bl}_{\mathbf{D}_{011}^{+}} \mathbf{X}^{1}$. Similar abuse of notation also appears below). The exceptional divisor \mathbf{E}_{2}^{1} is $\langle \widetilde{\beta}_{2}, \widetilde{\widetilde{\lambda}} \rangle$.

- A.1.14. U₂₃. U₂₃ is identified with Spec $\mathbb{R}[\beta_1, \widetilde{y}_1] \times \text{Spec} \mathbb{R}[\widetilde{\widetilde{\lambda}}, \widetilde{\widetilde{\beta}}_2]$ and

 - The ideal of Bl_{D⁺₀₁₁} X¹ ∩ U₂₃ is ⟨β̃₂²ŷ₁ + β̃²⟩;
 The proper transforms of D⁺₂₁₀ and D⁺₀₂₂ are contained in the complement of U₂₁;
 - 3. The proper transform \mathbf{D}_{111}^{++} (of \mathbf{D}_{111}^{+}) is $\langle \widetilde{\lambda}, \widetilde{\beta}_2 \rangle$;
 - 4. The proper transform \mathbf{E}_1^+ is $\langle \beta_1 \rangle$. The exceptional divisor \mathbf{E}_2^1 is $\langle \tilde{y}_1, \tilde{\lambda} \rangle$.
- A.1.15. U₄₁. Over U₄ we would have U₄₁ \cup U₄₂ \cup U₄₃, with D⁺₁₁₀ being blown up.

 U_{41} is identified with $\operatorname{Spec} \mathbb{R}[y_1, \widetilde{\lambda}] \times \operatorname{Spec} \mathbb{R}[\widetilde{\beta}_1, \widetilde{\widetilde{y}}_2]$ and

- 1. The ideal of $\mathbf{Bl}_{\mathbf{D}_{110}^+} \mathbf{X}^1 \cap \mathbf{U}_{41}$ is $\langle \widetilde{\lambda} \widetilde{\beta}_1 \widetilde{\widetilde{y}}_2^2 + 1 \rangle$;
- 2. The proper transforms of \mathbf{D}_{210}^+ , \mathbf{D}_{022}^+ and \mathbf{D}_{111}^+ are contained in the complement;
- 3. The proper transform \mathbf{E}_1^+ is $\langle y_1 \rangle$;
- 4. The exceptional divisor \mathbf{E}_2^2 is in the complement.
- A.1.16. U₄₂. U₄₂ is identified with Spec $\mathbb{R}[y_1, \widetilde{\beta}_1] \times \text{Spec } \mathbb{R}[\widetilde{\lambda}, \widetilde{\widetilde{y}}_2]$ and

 - The ideal of **Bl**_{D⁺₁₁₀} **X**¹ ∩ U₄₂ is ⟨*y*²₂β₁ + *k*² ⟩;
 The proper transforms of **D**⁺₂₁₀ and **D**⁺₀₂₂ are contained in the complement of U₄₂;
 - 3. The proper transform \mathbf{D}_{111}^{++} (of \mathbf{D}_{111}^{+}) is $\langle \widetilde{\lambda}, \widetilde{\widetilde{y}}_2 \rangle$;
 - 4. The proper transform \mathbf{E}_1^+ is $\langle y_1 \rangle$. The exceptional divisor \mathbf{E}_2^2 is $\langle \widetilde{\beta}_1, \widetilde{\lambda} \rangle$.
- A.1.17. U₄₃. U₄₃ is identified with Spec $\mathbb{R}[y_1, \tilde{y}_2] \times \text{Spec } \mathbb{R}[\tilde{\lambda}, \tilde{\beta}_1]$ and
 - 1. The ideal of $\mathbf{Bl}_{\mathbf{D}_{110}^+} \mathbf{X}^1 \cap \mathbf{U}_{43}$ is $\langle \widetilde{\widetilde{\beta}}_1 \widetilde{y}_2 + \widetilde{\widetilde{\lambda}}^2 \rangle$;
 - 2. The proper transforms of \mathbf{D}_{111}^+ and \mathbf{D}_{022}^+ are contained in the complement;
 - 3. The proper transform \mathbf{D}_{210}^{++} of \mathbf{D}_{210}^{+} is $\langle \widetilde{\lambda}, \widetilde{\beta}_1 \rangle$;
 - 4. The proper transform \mathbf{E}_1^+ is $\langle y_1 \rangle$. The exceptional divisor \mathbf{E}_2^2 is $\langle \tilde{y}_2, \tilde{\lambda} \rangle$.

A.1.18. Summary. On U₂₁ and U₄₁, we already have a good pair. It remains to deal with U₂₂, U₂₃, U₄₂ and U_{43} . The intersection pattern after this blowup is as follows:

$$\mathbf{D}_{210}^{++}\longleftrightarrow \mathbf{E}_{2}^{2}\longleftrightarrow \mathbf{D}_{111}^{++}\longleftrightarrow \mathbf{E}_{2}^{1}\longleftrightarrow \mathbf{D}_{022}^{++}$$

where \leftrightarrow means "intersects with". As before, \mathbf{E}_1^+ intersects every divisor here transversally.

A.1.19. Explicit constructions of a resolution, step III. Call $\mathbf{X}^2 := \mathbf{Bl}_{\mathbf{D}_{011}^+ \sqcup \mathbf{D}_{110}^+} \mathbf{X}^1$. As the final step, we blow up $\mathbf{C} := (\mathbf{E}_2^1 \cap \mathbf{D}_{022}^{++}) \bigsqcup \mathbf{D}_{111}^{++} \bigsqcup (\mathbf{D}_{210}^{++} \cap \mathbf{E}_2^2)$. Let \mathbf{E}_3^0 (resp. $\mathbf{E}_3^1, \mathbf{E}_3^2$) be the total transform of $\mathbf{E}_{2}^{1} \cap \mathbf{D}_{022}^{++}$ (resp. $\mathbf{D}_{111}^{++}, \mathbf{D}_{210}^{++} \cap \mathbf{E}_{2}^{2}$).

A.1.20. U₂₂₁. Over U₂₂ we are going to have $U_{221} \cup U_{222} \cup U_{223}$.

- U_{221} is identified with $\operatorname{Spec} \mathbb{R}[\beta_1, \widetilde{\widetilde{\lambda}}] \times \operatorname{Spec} \mathbb{R}[\widetilde{\beta}_2, \widetilde{\widetilde{\beta}}_1]$ and

 - The ideal of **Bl_C X²** ∩ U₂₂₁ is ([≈]_{β2} [∞]₁ + 1);
 The proper transforms of **D**⁺⁺₂₁₀, **D**⁺⁺₁₁₁, **D**⁺⁺₀₂₂ and **E**¹₂ are in the complement of U₂₂₁;
 - 3. The proper transform \mathbf{E}_1^{++} is $\langle \beta_1 \rangle$. The exceptional divisor \mathbf{E}_3^0 is $\langle \tilde{\lambda} \rangle$.

A.1.21. U₂₂₂. U₂₂₂ is identified with Spec $\mathbb{R}[\beta_1, \widetilde{\beta}_2] \times \operatorname{Spec} \mathbb{R}[\widetilde{\lambda}, \widetilde{\widetilde{g}}_1]$. It turns out that one can eliminate $\widetilde{\widetilde{y}}_1 = -\widetilde{\widetilde{\lambda}}^2$. After this is done, we have:

- Bl_C X² ∩ U₂₂₂ is identified with the affine space Spec ℝ[β₁, β̃₂, λ̃];
 The proper transforms of D⁺⁺₂₁₀, D⁺⁺₁₁₁ and E¹₂ are in the complement of U₂₂₂. The proper transform \mathbf{D}_{022}^{+++} is $\langle \widetilde{\widetilde{\lambda}} \rangle$;

3. The proper transform \mathbf{E}_1^{++} is $\langle \beta_1 \rangle$. The exceptional divisor \mathbf{E}_3^0 is $\langle \tilde{\beta}_2 \rangle$.

A.1.22. U₂₂₃. U₂₂₃ is identified as Spec $\mathbb{R}[\beta_1, \widetilde{\widetilde{y}}_1] \times \operatorname{Spec} \mathbb{R}[\widetilde{\widetilde{\lambda}}, \widetilde{\widetilde{\beta}}_2]$ and we can eliminate $\widetilde{\widetilde{\beta}}_2 = -\widetilde{\widetilde{\lambda}}^2$. Then we have

- 1. **Bl**_C **X**² \cap U₂₂₃ is identified with the affine space Spec $\mathbb{R}[\beta_1, \tilde{\tilde{y}}_1, \tilde{\tilde{\lambda}}];$ 2. The proper transforms of **D**⁺⁺₂₁₀, **D**⁺⁺₁₁₁ and **D**⁺⁺₀₂₂ are in the complement of U₂₂₃;
- 3. The proper transform \mathbf{E}_1^{++} is $\langle \beta_1 \rangle$, $(\mathbf{E}_2^1)^+$ is $\langle \widetilde{\lambda} \rangle$ and the exceptional divisor \mathbf{E}_3^0 is $\langle \widetilde{\widetilde{y}}_1 \rangle$.

A.1.23. U₂₃₁. Over U₂₃ we are going to have $U_{231} \cup U_{232}$.

 U_{231} is identified as $\operatorname{Spec} \mathbb{R}[\beta_1, \widetilde{y}_1, \widetilde{\lambda}] \times \operatorname{Spec} \mathbb{R}[\widetilde{\beta}_2]$ and

- The ideal of Bl_C X² ∩ U₂₃₁ is (^{≈²}β₂ŷ₁ + 1);
 The proper transforms of D⁺⁺₂₁₀, D⁺⁺₁₁₁, D⁺⁺₀₂₂ and E¹₂ are in the complement of U₂₂₁;
- 3. The proper transform \mathbf{E}_1^{++} is $\langle \beta_1 \rangle$. The exceptional divisor \mathbf{E}_3^1 is $\langle \tilde{\lambda} \rangle$

A.1.24. U₂₃₂. U₂₃₂ is identified as Spec $\mathbb{R}[\beta_1, \widetilde{y}_1, \widetilde{\beta}_2] \times \text{Spec } \mathbb{R}[\widetilde{\widetilde{\lambda}}]$ and one eliminates $\widetilde{y}_1 = -\widetilde{\widetilde{\lambda}}^2$. After that we below: that, we have:

- Bl_C X² ∩ U₂₃₂ is identified with the affine space Spec ℝ[β₁, β₂, λ);
 The proper transforms of D⁺⁺₂₁₀, D⁺⁺₁₁₁ and D⁺⁺₀₂₂ are in the complement of U₂₃₂;
- 3. The proper transform \mathbf{E}_1^{++} is $\langle \beta_1 \rangle$, $(\mathbf{E}_2^1)^+$ is $\langle \tilde{\lambda} \rangle$ and the exceptional divisor \mathbf{E}_3^1 is $\langle \tilde{\beta}_2 \rangle$.

A.1.25. U₄₂₁. Over U₄₂ we have U₄₂₁ \cup U₄₂₂ with the ideal $\langle \widetilde{\lambda}, \widetilde{\widetilde{y}}_2 \rangle$ of **D**⁺⁺₁₁₁ being blown up. U_{421} is identified as $\operatorname{Spec} \mathbb{R}[y_1, \widetilde{\beta}_1, \widetilde{\lambda}] \times \operatorname{Spec} \mathbb{R}[\widetilde{\widetilde{y}}_2]$ and we have

- The ideal of Bl_C X² is ⟨^{≈2}<sub>y₂β₁ + 1⟩;
 The proper transforms of D⁺⁺₂₁₀, D⁺⁺₁₁₁, D⁺⁺₀₂₂ and E²₂ are in the complement of U₄₂₁;
 </sub>
- 3. The proper transform \mathbf{E}_1^{++} is $\langle y_1 \rangle$. The exceptional divisor \mathbf{E}_3^1 is $\langle \widetilde{\lambda} \rangle$.

A.1.26. U₄₂₂. U₄₂₂ is identified as Spec $\mathbb{R}[y_1, \widetilde{\beta}_1, \widetilde{\widetilde{y}}_2] \times \text{Spec } \mathbb{R}[\widetilde{\widetilde{\lambda}}]$ and we eliminate $\widetilde{\beta}_1 = -\widetilde{\widetilde{\lambda}}^2$. Then,

- 1. **Bl**_C **X**² \cap U₄₂₂ is identified with the affine space Spec $\mathbb{R}[y_1, \tilde{\tilde{y}}_2, \tilde{\tilde{\lambda}}];$ 2. The proper transforms of **D**⁺⁺₂₁₀, **D**⁺⁺₁₁₁ and **D**⁺⁺₀₂₂ are in the complement of U₄₂₂;
- 3. The proper transform \mathbf{E}_1^{++} is $\langle y_1 \rangle$ and $(\mathbf{E}_2^2)^+$ is $\langle \widetilde{\widetilde{\lambda}} \rangle$. The exceptional divisor \mathbf{E}_3^1 is $\langle \widetilde{\widetilde{y}}_2 \rangle$.

A.1.27. U₄₃₁. Over U₄₃, we are going to have U₄₃₁ \cup U₄₃₂ \cup U₄₃₃, with the ideal $\langle \tilde{\lambda}, \tilde{\beta}_1, \tilde{y}_2 \rangle$ of $\mathbf{D}_{210}^{++} \cap \mathbf{E}_2^2$ being blown up.

 U_{431} is identified as $\operatorname{Spec} \mathbb{R}[y_1, \widetilde{\widetilde{\lambda}}] \times \operatorname{Spec} \mathbb{R}[\widetilde{\widetilde{\beta}}_1, \widetilde{\widetilde{y}}_2]$ and we have

- The ideal of Bl_C X² is (*ỹ*₂*β̃*₁ + 1);
 The proper transforms of D⁺⁺₂₁₀, D⁺⁺₁₁₁, D⁺⁺₀₂₂ and (E²₂)⁺ are in the complement of U₂₂₁;
- 3. The proper transform \mathbf{E}_1^{++} is $\langle y_1 \rangle$. The exceptional divisor \mathbf{E}_3^2 is $\langle \tilde{\lambda} \rangle$.

A.1.28. U₄₃₂. U₄₃₂ is identified as Spec $\mathbb{R}[\tilde{y}_1, \tilde{\beta}_1] \times \text{Spec } \mathbb{R}[\tilde{\tilde{\lambda}}, \tilde{\tilde{y}}_2]$. One can eliminate $\tilde{\tilde{y}}_2 = -\tilde{\tilde{\lambda}}^2$ here after which we have

- 1. **Bl**_C **X**² is identified with the affine space Spec $\mathbb{R}[\tilde{y}_1, \widetilde{\tilde{\beta}}_1, \widetilde{\tilde{\lambda}}];$ 2. The proper transforms of **D**⁺⁺₂₁₀, **D**⁺⁺₁₁₁ and **D**⁺⁺₀₂₂ are in the complement of U₄₃₂;
- 3. The proper transform \mathbf{E}_1^{++} is $\langle y_1 \rangle$, $(\mathbf{E}_2^2)^+$ is $\langle \widetilde{\widetilde{\lambda}} \rangle$. The exceptional divisor \mathbf{E}_3^0 is $\langle \widetilde{\widetilde{\beta}}_1 \rangle$.

A.1.29. U₄₃₃. U₄₃₃ is identified with Spec $\mathbb{R}[\widetilde{y}_1, \widetilde{y}_2] \times \text{Spec } \mathbb{R}[\widetilde{\widetilde{\lambda}}, \widetilde{\widetilde{\beta}}_1]$ and we eliminate $\widetilde{\widetilde{\beta}}_1 = -\widetilde{\widetilde{\lambda}}^2$ here. After that,

- 1. **Bl**_C **X**² is identified with the affine space Spec $\mathbb{R}[\tilde{y}_1, \tilde{y}_2, \tilde{\tilde{\lambda}}];$ 2. The proper transforms of **D**⁺⁺₁₁₁, **D**⁺⁺₀₂₂ and (**E**¹₂)⁺ are in the complement of U₄₃₃. The proper
- transform \mathbf{D}_{210}^{+++} is $\langle \widetilde{\widetilde{\lambda}} \rangle$; 3. The proper transform \mathbf{E}_1^{++} is $\langle y_1 \rangle$. The exceptional divisor \mathbf{E}_3^2 is $\langle \widetilde{y}_2 \rangle$.

Finally, we have obtained a smooth pair everywhere. Let $\mathbf{X}^3 := \mathbf{Bl}_{\mathbf{C}} \mathbf{X}^2$. Since each center of the blowup is G-invariant, all the blowup morphisms above are G-equivariant. Here is an illustration of the intersection pattern. As above, \mathbf{E}_1^{++} intersects everything transversally.

$$\mathbf{D}_{210}^{+++} \longleftrightarrow \mathbf{E}_3^2 \longleftrightarrow (\mathbf{E}_2^2)^+ \longleftrightarrow \mathbf{E}_3^1 \longleftrightarrow (\mathbf{E}_2^1)^+ \longleftrightarrow \mathbf{E}_3^0 \longleftrightarrow \mathbf{D}_{022}^{+++}$$

A.1.30. Explicit invariant gauge form. First we present the invariant gauge form on $\mathbf{SL}_2(\mathbb{R}) \times \mathbf{SL}_2(\mathbb{R})$. In coordinates, elements in $\mathbf{SL}_2(\mathbb{R}) \times \mathbf{SL}_2(\mathbb{R})$ can be written as

$$\left(\begin{bmatrix} u_1 & u_2 \\ u_3 & u_4 \end{bmatrix}, \begin{bmatrix} w_1 & w_2 \\ w_3 & w_4 \end{bmatrix} \right), \text{ satisfying } u_1 u_4 - u_2 u_3 = w_1 w_4 - w_2 w_3 = 1.$$
(36)

Then the invariant form is given by

$$\omega_0 := \frac{\mathrm{d}u_1 \wedge \mathrm{d}u_2 \wedge \mathrm{d}u_3}{u_1} \wedge \frac{\mathrm{d}w_1 \wedge \mathrm{d}w_2 \wedge \mathrm{d}w_3}{w_1}$$

Let us put one more restriction:

$$f = u_1 w_4 + u_4 w_1 - u_2 w_3 - u_3 w_2 = 0, (37)$$

which is invariant under the $\mathbf{SL}_2(\mathbb{R}) \times \mathbf{SL}_2(\mathbb{R})$ action: $(g,h) \cdot (A,B) := (gAh^{-1}, gBh^{-1})$. One can verify that

$$\omega_1 := \frac{\mathrm{d}u_1 \wedge \mathrm{d}u_2 \wedge \mathrm{d}u_3 \wedge \mathrm{d}w_1 \wedge \mathrm{d}w_2}{(u_1 w_2 - u_2 w_1)u_1}$$

satisfies

$$\omega_1 \wedge \mathrm{d}f = \omega_0$$
 restricted to $f = 0$

And hence ω_1 is an invariant volume form on $\{f = 0\}$.

Let $\mathbf{u}' := (u'_1, u'_2, u'_3, u'_4) := (u_1, -u_2, u_3, u_4)$ and $\mathbf{w}' := (w'_1, w'_2, w'_3, w'_4) := (w_1, -w_2, w_3, w_4)$, then Equa.(36, 37) become

$$Q_2^0(\mathbf{u}', \mathbf{u}') = Q_2^0(\mathbf{w}', \mathbf{w}') = 2, \ Q_2^0(\mathbf{u}', \mathbf{w}') = 0.$$
(38)

Let us further set

$$\boldsymbol{\gamma} := rac{\mathbf{u}' + i\mathbf{w}'}{2}, \ \boldsymbol{\theta} := rac{\mathbf{u}' - i\mathbf{w}'}{2}$$

Then Equa.(38) becomes

$$Q_2^0(oldsymbol{\gamma},oldsymbol{\gamma})=Q_2^0(oldsymbol{ heta},oldsymbol{ heta})=0,\;Q_2^0(oldsymbol{\gamma},oldsymbol{ heta})=1$$

which is exactly the model of homogeneous space discussed above in Section A.1.1. Using these new variables,

$$\omega_1 = \frac{\mathrm{d}\gamma_1 \wedge \mathrm{d}\theta_1 \wedge \mathrm{d}\gamma_2 \wedge \mathrm{d}\theta_2 \wedge (\mathrm{d}\gamma_3 + \mathrm{d}\theta_3)}{(\gamma_1 + \theta_1)(\gamma_1 \theta_2 - \gamma_2 \theta_1)}$$

Recall the coordinates as introduced in Equa.(33) and

$$\begin{bmatrix} \gamma_1 & \theta_1 \\ \gamma_2 & \theta_2 \\ \gamma_3 & \theta_3 \\ \gamma_4 & \theta_4 \end{bmatrix} : 1] = \begin{bmatrix} 1 & \beta_1 \\ \alpha_2 & \beta_2 \\ x_1 + \alpha_2 y_1 & \beta_1 x_1 + \beta_2 y_1 \\ x_2 + \alpha_2 y_2 & \beta_1 x_1 + \beta_2 y_2 \end{bmatrix} : \lambda].$$

We have

$$\begin{cases} \gamma_1 = \frac{1}{\lambda} \implies d\gamma_1 = -\lambda^{-1} d\lambda \\ \theta_1 = \frac{\beta_1}{\lambda} \implies d\theta_1 = \pm \lambda^{-1} d\beta_1 \mod d\lambda \\ \gamma_2 = \frac{\alpha_2}{\lambda} \implies d\gamma_2 = \lambda^{-1} d\alpha_2 \mod d\lambda \\ \theta_2 = \frac{\beta_2}{\lambda} \implies d\theta_2 = \lambda^{-1} d\beta'_2 \mod d\lambda, d\beta_1, d\alpha_2 \end{cases}$$

and finally,

$$\gamma_3 + \theta_3 = \lambda^{-1} (x_1 + \alpha_2 y_1 + \beta_1 x_1 + \beta_2 y_1)$$

$$\implies d\gamma_3 + d\theta_3 = \pm \frac{1 + \beta_1}{\lambda} dx_1 \mod d\lambda, d\beta_1, d\alpha_2, d\beta'_2$$

$$\gamma_1 + \theta_1 = \lambda^{-1} (1 + \beta_1), \quad \gamma_1 \theta_2 - \gamma_2 \theta_1 = \lambda^{-2} \beta'_2.$$

In sum, we get the invariant gauge form on \mathbf{X}^{INC} :

$$\omega_1 = \pm \frac{1}{\lambda^3 \beta_2'} \mathrm{d}\lambda \wedge \mathrm{d}\beta_1 \wedge \mathrm{d}\beta_2' \wedge \mathrm{d}x_1 \wedge \mathrm{d}\alpha_2.$$

Similar to the above, we can ignore the variables x_1 and α_2 and focus on

$$\omega_2 := \frac{1}{\lambda^3 \beta_2'} \mathrm{d}\lambda \wedge \mathrm{d}\beta_1 \wedge \mathrm{d}\beta_2'. \tag{39}$$

We will compute poles of ω_2 , or equivalently, ω_1 , along the boundaries in \mathbf{X}^3 .

A.1.31. The order of pole along \mathbf{E}_1^{++} is 2. This can be checked on U_1 where

$$\beta_1 = \lambda \widetilde{\beta}_1, \ \beta'_2 = \lambda \widetilde{\beta}'_2, \ \widetilde{\beta}_1 = -\widetilde{\beta}_2^2 \widetilde{y}_1.$$

In particular,

$$\mathrm{d}\widetilde{\beta}_1 = -\widetilde{\beta}_2^2 \mathrm{d}\widetilde{y}_1 \mod \mathrm{d}\widetilde{\beta}_2.$$

Consequently,

$$\omega_2 = \frac{\widetilde{\beta}_2}{\lambda^2} \mathrm{d}\lambda \wedge \mathrm{d}\widetilde{y}_1 \wedge \mathrm{d}\widetilde{\beta}_2. \tag{40}$$

Since the local equation of \mathbf{E}_1 is given by $\lambda = 0$, we get $\text{Pole}(\omega_2, \mathbf{E}_1) = 2$. But the blowup morphisms are trivial over U_1 hence $\text{Pole}(\omega_2, \mathbf{E}_1^{++}) = 2$.

A.1.32. The order of pole along \mathbf{D}_{022}^{+++} is 3. This can be found on U₃ where $\lambda = \tilde{\lambda}\beta'_2$, $\beta_1 = \tilde{\beta}_1\beta'_2$. Thus,

$$\omega_2 = \frac{1}{\widetilde{\lambda}^3 \beta_2'^4} d(\widetilde{\lambda}\beta_2') \wedge d(\widetilde{\beta}_1\beta_2') \wedge d\beta_2' = \frac{1}{\widetilde{\lambda}^3 (\beta_2')^2} d\widetilde{\lambda} \wedge d\widetilde{\beta}_1 \wedge d\beta_2'.$$

Since the local equation of \mathbf{D}_{022}^+ is given by $\tilde{\lambda} = 0$, we have $\operatorname{Pole}(\omega_2, \mathbf{D}_{022}^+) = 3$. But the blowup morphism is trivial over U₃, so $\operatorname{Pole}(\omega_2, \mathbf{D}_{022}^{+++}) = 3$.

A.1.33. The order of pole along \mathbf{D}_{210}^{+++} is 1. This can be checked on U₅ where one has the coordinates $y'_2, \tilde{y}_1, \tilde{\lambda}$ and $\tilde{\beta}_2 = \tilde{\lambda}^2, \tilde{\beta}_1 = -\tilde{\beta}_2 \tilde{y}_1 = -\tilde{\lambda}^2 \tilde{y}_1$ are eliminated. Also, $\lambda = \tilde{\lambda} y'_2, \beta_1 = \tilde{\beta}_1 y'_2$ and $\beta_2 = \tilde{\beta}_2 y'_2$. The local equation of \mathbf{D}_{210}^+ is $\tilde{\lambda} = 0$. Then one can compute

$$\omega_{2} = \frac{1}{y_{2}^{\prime 4} \tilde{\lambda}^{3} \tilde{\beta}_{2}} \mathbf{d}(\tilde{\lambda} y_{2}^{\prime}) \wedge \mathbf{d}(\tilde{\beta}_{1} y_{2}^{\prime}) \wedge \mathbf{d}(\tilde{\beta}_{2} y_{2}^{\prime})$$
$$= \pm \frac{1}{(y_{2}^{\prime})^{2} \tilde{\lambda}} \mathbf{d} y_{2}^{\prime} \wedge \mathbf{d} \tilde{y}_{1} \wedge \mathbf{d} \tilde{\lambda}.$$

Now one sees directly that $\operatorname{Pole}(\omega_2, \mathbf{D}_{210}^{+++}) = \operatorname{Pole}(\omega_2, \mathbf{D}_{210}^{+}) = 1.$

A.1.34. The order of pole along \mathbf{E}_3^0 is 5, along $(\mathbf{E}_2^1)^+$ is 7. These can be checked on U₂₂₂. First we have that on U₂, the coordinates are given by $\beta_1, \tilde{\lambda}, \tilde{\beta}_2, \tilde{y}_1$ and $\tilde{y}_2 = -\tilde{\beta}_2 \tilde{y}_1$ is eliminated. The local equation of \mathbf{X}^1 is $\tilde{\beta}_2^2 \tilde{y}_1 + \tilde{\lambda}^2 = 0$. And one has the relations:

$$\lambda = \widetilde{\lambda}\beta_1, \ \beta_2 = \widetilde{\beta}_2\beta_1, \ y_i = \widetilde{y}_i\beta_1, \ i = 1, 2$$

From this one can compute

$$\omega_2 = \frac{1}{\widetilde{\lambda}^3 \beta_1^2 \widetilde{\beta}_2} \mathrm{d} \widetilde{\lambda} \wedge \mathrm{d} \beta_1 \wedge \mathrm{d} \widetilde{\beta}_2$$

On U₂₂ one has coordinates $\beta_1, \widetilde{\beta}_2, \widetilde{\widetilde{\lambda}}, \widetilde{\widetilde{y}}_1$ with $\widetilde{\lambda} = \widetilde{\beta}_2 \widetilde{\widetilde{\lambda}}, \ \widetilde{y}_1 = \widetilde{\beta}_2 \widetilde{\widetilde{y}}_1$. Local equation of \mathbf{X}^2 is $\widetilde{\beta}_2 \widetilde{\widetilde{y}}_1 + \widetilde{\widetilde{\lambda}}^2 = 0$. Then

$$\omega_2 = \frac{1}{(\widetilde{\widetilde{\lambda}})^3 \beta_1^2 \widetilde{\beta}_2^3} \mathrm{d} \widetilde{\widetilde{\lambda}} \wedge \mathrm{d} \beta_1 \wedge \mathrm{d} \widetilde{\beta}_2.$$

Finally over U₂₂₂, we have the coordinates $\beta_1, \tilde{\tilde{y}}_1, \tilde{\tilde{\tilde{\lambda}}}$ and $\tilde{\tilde{\beta}}_2 = -\tilde{\tilde{\tilde{\lambda}}}^2$ is eliminated. And $\mathbf{X}^3 \cap U_{222}$ is the full affine space here. One has $\tilde{\tilde{\lambda}} = \tilde{\tilde{y}}_1 \tilde{\tilde{\tilde{\lambda}}}, \tilde{\tilde{\beta}}_2 = \tilde{\tilde{y}}_1 \tilde{\tilde{\beta}}_2$. The local equation of \mathbf{E}_3^0 is $\tilde{\tilde{y}}_1 = 0$ and that of $(\mathbf{E}_2^1)^+$ is $\widetilde{\lambda} = 0$. Now compute

$$\begin{split} \omega_2 &= \frac{1}{(\widetilde{\widetilde{\lambda}})^3 \beta_1^2 (\widetilde{\widetilde{\beta}}_2)^3 (\widetilde{\widetilde{y}}_1)^6} \mathbf{d}(\widetilde{\widetilde{y}}_1 \widetilde{\widetilde{\widetilde{\lambda}}}) \wedge \mathbf{d}\beta_1 \wedge \mathbf{d}(\widetilde{\widetilde{y}}_1 \widetilde{\widetilde{\widetilde{\lambda}}}^2) \\ &= \pm \frac{1}{(\widetilde{\widetilde{\lambda}})^7 \beta_1^2 (\widetilde{\widetilde{y}}_1)^5} \mathbf{d}\widetilde{\widetilde{y}}_1 \wedge \mathbf{d}\beta_1 \wedge \mathbf{d}\widetilde{\widetilde{\widetilde{\lambda}}}. \end{split}$$

Hence $Pole(\omega_2, \mathbf{E}_3^0) = 5$ and $Pole(\omega_2, (\mathbf{E}_2^1)^+) = 7$.

A.1.35. The order of pole along \mathbf{E}_3^1 is 3, along $(\mathbf{E}_2^2)^+$ is 5. We shall check this on U₄₂₂.

To cut the story short, on U₄₂₂, we have coordinates $y_1, \tilde{\tilde{y}}_2, \tilde{\tilde{\lambda}}$ and \mathbf{X}^3 is the full affine space. The local equation of \mathbf{E}_3^1 is $\tilde{\tilde{y}}_2 = 0$ and that of $(\mathbf{E}_2^2)^+$ is $\tilde{\tilde{\lambda}} = 0$. The relations with the old coordinates are

$$\begin{split} \lambda &= y_1 \widetilde{\lambda} = y_1 \widetilde{\beta}_1 \widetilde{\widetilde{\lambda}} = -y_1 (\widetilde{\widetilde{\lambda}})^3 \widetilde{\widetilde{y}}_2, \\ \beta_1 &= -y_1 (\widetilde{\widetilde{\lambda}})^2, \ \beta_2' = -y_1 (\widetilde{\widetilde{\lambda}})^4 \widetilde{\widetilde{y}}_2 \end{split}$$

Now we can compute

$$\begin{split} \omega_{2} &= \pm \frac{\mathrm{d}(y_{1}(\widetilde{\widetilde{\lambda}})^{3}\widetilde{\widetilde{y}_{2}}) \wedge \mathrm{d}(y_{1}(\widetilde{\widetilde{\lambda}})^{4}\widetilde{\widetilde{y}_{2}}) \wedge \mathrm{d}(y_{1}(\widetilde{\widetilde{\lambda}})^{2})}{y_{1}^{3}(\widetilde{\widetilde{\lambda}})^{9}(\widetilde{\widetilde{y}_{2}})^{3}y_{1}(\widetilde{\widetilde{\lambda}})^{4}\widetilde{\widetilde{y}_{2}}} \\ &= \pm \frac{1}{y_{1}^{4}(\widetilde{\widetilde{\lambda}})^{13}(\widetilde{\widetilde{y}_{2}})^{4}} \mathrm{d}(y_{1}(\widetilde{\widetilde{\lambda}})^{3}\widetilde{\widetilde{y}_{2}}) \wedge \left(y_{1}(\widetilde{\widetilde{\lambda}})^{3}\widetilde{\widetilde{y}_{2}}\right) \mathrm{d}\widetilde{\widetilde{\lambda}} \wedge \mathrm{d}(y_{1}(\widetilde{\widetilde{\widetilde{\lambda}}})^{2}) \\ &= \pm \frac{1}{y_{1}^{2}(\widetilde{\widetilde{\lambda}})^{5}(\widetilde{\widetilde{y}_{2}})^{3}} \mathrm{d}\widetilde{\widetilde{y}_{2}} \wedge \mathrm{d}\widetilde{\widetilde{\widetilde{\lambda}}} \wedge \mathrm{d}y_{1}. \end{split}$$

Hence $\operatorname{Pole}(\omega_2, \mathbf{E}_3^1) = 3$ and $\operatorname{Pole}(\omega_2, (\mathbf{E}_2^2)^+) = 5$.

A.1.36. The order of pole along \mathbf{E}_3^2 is 3. This will be checked on U_{432} . We have here coordinates $y_1, \tilde{\widetilde{\beta}}_1, \tilde{\widetilde{\lambda}}$ and there are no further constraints on \mathbf{X}^3 . Moreover, the local equation of \mathbf{E}_3^2 is $\tilde{\widetilde{\beta}}_1 = 0$. The relations with the old coordinates are given by

$$\begin{split} \lambda &= y_1 \widetilde{\lambda} = y_1 \widetilde{y}_2 \widetilde{\widetilde{\lambda}} = y_1 \widetilde{\widetilde{\beta}}_1^2 \widetilde{\widetilde{y}}_2^{\widetilde{\widetilde{\lambda}}} \widetilde{\widetilde{\lambda}} = -y_1 (\widetilde{\widetilde{\beta}}_1)^2 (\widetilde{\widetilde{\lambda}})^3, \\ \beta_1 &= y_1 \widetilde{\beta}_1 = y_1 \widetilde{y}_2 \widetilde{\widetilde{\beta}}_1 = -y_1 (\widetilde{\widetilde{\beta}}_1)^2 (\widetilde{\widetilde{\widetilde{\lambda}}})^2, \\ \beta_2' &= y_1 \widetilde{\beta}_2 = -y_1 \widetilde{\beta}_1 \widetilde{y}_2 = -y_1 \widetilde{\widetilde{\beta}}_1 \widetilde{y}_2^2 = -y_1 (\widetilde{\widetilde{\beta}}_1)^3 (\widetilde{\widetilde{y}}_2)^2 = -y_1 (\widetilde{\widetilde{\beta}}_1)^3 (\widetilde{\widetilde{\widetilde{\lambda}}})^4. \end{split}$$

Therefore,

$$\begin{split} \omega_{2} &= \pm \frac{\mathrm{d}(y_{1}(\widetilde{\widetilde{\beta}}_{1})^{2}(\widetilde{\widetilde{\widetilde{\lambda}}})^{3}) \wedge \mathrm{d}(y_{1}(\widetilde{\widetilde{\beta}}_{1})^{2}(\widetilde{\widetilde{\widetilde{\lambda}}})^{2}) \wedge \mathrm{d}(y_{1}(\widetilde{\widetilde{\beta}}_{1})^{3}(\widetilde{\widetilde{\widetilde{\lambda}}})^{4})}{\left(y_{1}^{3}(\widetilde{\widetilde{\beta}}_{1})^{6}(\widetilde{\widetilde{\widetilde{\lambda}}})^{9}\right) \cdot \left(y_{1}(\widetilde{\widetilde{\beta}}_{1})^{3}(\widetilde{\widetilde{\widetilde{\lambda}}})^{4}\right)} \\ &= \pm \frac{1}{y_{1}^{4}(\widetilde{\widetilde{\beta}}_{1})^{9}(\widetilde{\widetilde{\widetilde{\lambda}}})^{13}} \left(y_{1}(\widetilde{\widetilde{\beta}}_{1})^{2}(\widetilde{\widetilde{\widetilde{\lambda}}})^{2}\right) \mathrm{d}^{\widetilde{\widetilde{\lambda}}} \wedge \mathrm{d}(y_{1}(\widetilde{\widetilde{\beta}}_{1})^{2}(\widetilde{\widetilde{\widetilde{\lambda}}})^{2}) \wedge \mathrm{d}(y_{1}(\widetilde{\widetilde{\beta}}_{1})^{3}(\widetilde{\widetilde{\widetilde{\lambda}}})^{4}) \\ &= \pm \frac{1}{y_{1}^{3}(\widetilde{\widetilde{\beta}}_{1})^{7}(\widetilde{\widetilde{\widetilde{\lambda}}})^{11}} \mathrm{d}^{\widetilde{\widetilde{\lambda}}} \wedge (\widetilde{\widetilde{\widetilde{\lambda}}})^{2} \mathrm{d}(y_{1}(\widetilde{\widetilde{\beta}}_{1})^{2}) \wedge (\widetilde{\widetilde{\lambda}})^{4} \mathrm{d}(y_{1}(\widetilde{\widetilde{\beta}}_{1})^{3}) \\ &= \pm \frac{1}{y_{1}^{2}(\widetilde{\widetilde{\beta}}_{1})^{3}(\widetilde{\widetilde{\widetilde{\lambda}}})^{5}} \mathrm{d}^{\widetilde{\widetilde{\lambda}}} \wedge \mathrm{d}y_{1} \wedge \mathrm{d}^{\widetilde{\widetilde{\beta}}}_{1}, \end{split}$$

from which one sees that $Pole(\omega_2, \mathbf{E}_3^2) = 3$.

A.2. Example II. Here we provide details for the example presented in Section 8. Recall that **X** is the Zariski closure of $\mathbf{U} := \mathbf{G}.y_0$ with $y_0 := ([\mathbf{e}_1 : 1], [\mathbf{e}_1^{\vee} : 1])$ in $\mathbf{P}_{\mathbb{Q}}^{n+1} \times \mathbf{P}_{\mathbb{Q}}^{n+1}$ and **D** is the complement of **U** in **X**. If a general (closed) point in **X** is denoted as $[\mathbf{v} : s], [\boldsymbol{\alpha} : t]$, then \mathbf{D}_1 is defined by s = 0 and \mathbf{D}_2 by t = 0.

A.2.1. Local equations of stratum. We define the following affine open subvarieties of X:

$$\begin{aligned} \mathbf{O}_1 : & \{ ([1:v_2:\ldots:v_{n+1}:s], [\alpha_1:1:\alpha_3:\ldots:\alpha_{n+1}:t]) \} \\ \mathbf{O}_2 : & \{ ([v_1:v_2:\ldots:v_{n+1}:1], [1:\alpha_2:\ldots:\alpha_{n+1}:t]) \} ; \\ \mathbf{O}_3 : & \{ ([1:v_2:\ldots:v_{n+1}:s], [\alpha_1:\alpha_2:\ldots:\alpha_{n+1}:1]) \} . \end{aligned}$$

Using these local coordinates, one finds that⁴

Lemma A.3. As a closed subvariety of Spec $\mathbb{Q}[v_2, ..., v_{n+1}, \alpha_1, \alpha_3, ..., \alpha_{n+1}, s, t]$, the ideal of \mathbf{O}_1 is generated by

$$f_1 := \alpha_1 + v_2 + \sum_{i=3}^{n+1} v_i \alpha_i - st.$$

The ideal of $\mathbf{D}_1 \cap \mathbf{O}_1$ (resp. $\mathbf{D}_2 \cap \mathbf{O}_1$) is generated by f_1 , s (resp. f_1 , t). As a closed subvariety of Spec $\mathbb{Q}[v_1, ..., v_{n+1}, \alpha_2, ..., \alpha_{n+1}, t]$, the ideal of \mathbf{O}_2 is generated by

$$f_2 := v_1 + \sum_{i=2}^{n+1} v_i \alpha_i - t.$$

The ideal of $\mathbf{D}_2 \cap \mathbf{O}_2$ is generated by f_2 , t and $\mathbf{D}_1 \cap \mathbf{O}_2$ is empty. As a closed subvariety of Spec $\mathbb{Q}[v_2, ..., v_{n+1}, \alpha_1, ..., \alpha_{n+1}, s]$, the ideal of \mathbf{O}_3 is generated by

$$f_3 := \alpha_1 + \sum_{i=2}^{n+1} v_i \alpha_i - s.$$

⁴Compared to the last subsection, it is direct to verify the ideals generated by f_i is prime.

The ideal of $\mathbf{D}_1 \cap \mathbf{O}_3$ is generated by f_2 , s and $\mathbf{D}_2 \cap \mathbf{O}_3$ is empty.

A.2.2. The divisor of the invariant gauge form. It suffices to work with O_1 . In the ambient affine space, the differential form

$$\omega_1 := \operatorname{d}(\frac{1}{s}) \wedge \operatorname{d}(\frac{v_2}{s}) \wedge \dots \wedge \operatorname{d}(\frac{v_{n+1}}{s}) \wedge \operatorname{d}(\frac{\alpha_1}{t}) \wedge \operatorname{d}(\frac{1}{t}) \wedge \operatorname{d}(\frac{\alpha_3}{t}) \wedge \dots \wedge \operatorname{d}(\frac{\alpha_{n+1}}{t})$$
$$= \pm s^{-(n+2)} t^{-(n+2)} \operatorname{d} s \wedge \operatorname{d} v_2 \wedge \dots \wedge \operatorname{d} v_{n+1} \wedge \operatorname{d} t \wedge \operatorname{d} \alpha_1 \wedge \operatorname{d} \alpha_3 \wedge \dots \wedge \operatorname{d} \alpha_{n+1}$$

is **G**-invariant. Then, the unique (up to a scalar) solution $\omega_{\mathbf{U}}$ on $f_1 = 0$ to

$$\omega_1 = \omega_{\mathbf{U}} \wedge \mathrm{d}f_1$$

is the invariant gauge form. Since

$$\mathrm{d}f_1 = \mathrm{d}v_2 + \left(\mathrm{d}\alpha_1 + \sum_{i=3}^{n+1} (\alpha_i \mathrm{d}v_i + v_i \mathrm{d}\alpha_i) - t\mathrm{d}s - s\mathrm{d}t\right),\,$$

we find that up to a scalar,

$$\omega_{\mathbf{U}} = s^{-(n+2)} t^{-(n+2)} \mathrm{d}s \wedge \mathrm{d}v_3 \wedge \ldots \wedge \mathrm{d}v_{n+1} \wedge \mathrm{d}t \wedge \mathrm{d}\alpha_1 \wedge \mathrm{d}\alpha_3 \wedge \ldots \wedge \mathrm{d}\alpha_{n+1}.$$

And it is direct to check that

$$\operatorname{Pole}(\omega_{\mathbf{U}}, \mathbf{D}_1) = \operatorname{Pole}(\omega_{\mathbf{U}}, \mathbf{D}_2) = n + 2.$$

A.2.3. Measure compactification. The lemma below follows from the equidistribution theorem of [EMS96] and the nondivergence theorem of [DM91] (stated more precisely in [DGU21, Theorem 4.6]). Note that there are exactly two rational parabolic subgroups containing **H**. And for a sequence (g_n) in G, to test whether adjoint orbits based at the two vectors representing the Lie algebras of these two parabolic subgroups go to 0, is the same as to test whether $(g_n \mathbf{e}_1)$ and $(g_n \mathbf{e}_1^{\vee})$ go to 0.

Lemma A.4. Let (g_n) be a sequence in G. By passing to a subsequence, we are in exactly one of the following situations:

1. $(g_n \cdot \mathbf{e}_1)$ or $(g_n \mathbf{e}_1^{\vee})$ tends to $\mathbf{0}$, in which case,

$$\lim_{n \to \infty} (g_n)_* \mathbf{m}^{\mathbf{1}}_{[\mathrm{H}]} = \mathbf{0};$$

2. (g_n) converges to $\delta \in G$ modulo H, in which case,

$$\lim_{n \to \infty} (g_n)_* \mathbf{m}_{[\mathrm{H}]}^1 = \delta_* \mathbf{m}_{[\mathrm{H}]}^1;$$

3. (g_n) is unbounded modulo H, but converges to $\delta \in G$ modulo M_1 , in which case,

$$\lim_{n \to \infty} (g_n)_* \mathbf{m}_{[\mathrm{H}]}^1 = \delta_* \mathbf{m}_{[\mathrm{M}_1]}^1$$

4. (g_n) is unbounded modulo H, but converges to $\delta \in G$ modulo M₂, in which case,

$$\lim_{n \to \infty} (g_n)_* \mathbf{m}_{[\mathrm{H}]}^1 = \delta_* \mathbf{m}_{[\mathrm{M}_2]}^1;$$

5. $(g_n.\mathbf{e}_1)$ and $(g_n.\mathbf{e}_1^{\vee})$ both tend to infinity, in which case,

$$\lim_{n \to \infty} (g_n)_* \mathbf{m}^{\mathbf{1}}_{[\mathrm{H}]} = \mathbf{m}^{\mathbf{1}}_{[\mathrm{G}]}$$

To deduce Lemma 8.6 from Lemma A.4, one observes that (g_n) is bounded modulo M_1 iff $(g_n.e_1)$ is bounded away from **0** and infinity and similarly; it is bounded modulo M_2 iff $(g_n.e_1^{\vee})$ is bounded away from **0** and infinity.

A.2.4. The metric line bundles. Let ρ be a non-negative smooth function on \mathbb{R}^{n+1} such that

$$\rho(v) = \begin{cases} 0 & \text{if } \|v\| \ge 1\\ 1 & \text{if } \|v\| \le 0.5 \end{cases}$$

For a positive integer κ , we define a smooth metric $\|\cdot\|_{2\kappa}$ on $\mathcal{O}_{\mathbf{X}}(\mathbf{D}_1)$ by

$$\|\mathbf{1}_{\mathbf{D}_{1}}([\mathbf{v}:s],[\boldsymbol{\alpha}:t])\|_{2\kappa} := \left(\frac{s^{2\kappa}}{\sum v_{i}^{2\kappa} + s^{2\kappa}\rho(\frac{v_{1}}{s},...,\frac{v_{n+1}}{s})}\right)^{\frac{1}{2\kappa}}.$$

Similarly, define a smooth metric $\|\cdot\|_{2\kappa}$ on $\mathcal{O}_{\mathbf{X}}(\mathbf{D}_2)$ by

$$\|\mathbf{1}_{\mathbf{D}_2}([\mathbf{v}:s],[\boldsymbol{\alpha}:t])\|_{2\kappa} \coloneqq \left(\frac{t^{2\kappa}}{\sum \alpha_i^{2\kappa} + t^{2\kappa}\rho(\frac{\alpha_1}{t},...,\frac{\alpha_{n+1}}{t})}\right)^{\frac{1}{2\kappa}}$$

One can check they are indeed smooth metrics using local coordinates. Let us check the one on $\mathcal{O}_{\mathbf{X}}(\mathbf{D}_1)$. On $\mathbf{O}_1 \cup \mathbf{O}_3$, we have

$$\begin{split} & \left\| \frac{1}{s} \mathbf{1}_{\mathbf{D}_{1}} ([1:v_{2}:\ldots:v_{n+1}:s],[\boldsymbol{\alpha}:t]) \right\|_{2\kappa} \\ & = \left(\frac{1}{(1+v_{2}^{2\kappa}+\ldots+v_{n+1}^{2\kappa})+s^{2\kappa}\rho(\frac{1}{s},\frac{v_{2}}{s},\ldots,\frac{v_{n+1}}{s})} \right)^{\frac{1}{2\kappa}} \end{split}$$

Note that ρ vanishes when $|s| \leq 1$. On \mathbf{O}_2 , we have

$$\|\mathbf{1}_{\mathbf{D}_{1}}([v_{1}:...:v_{n+1}:1],[\boldsymbol{\alpha}:t])\|_{2\kappa} = \left(\frac{1}{\sum v_{i}^{2\kappa} + \rho(v_{1},...,v_{n+1})}\right)^{\frac{1}{2\kappa}}$$

Since $\mathcal{O}_{\mathbf{X}}(\mathbf{D}_1)$ is generated by $\mathbf{1}_{\mathbf{D}_1}$ on \mathbf{O}_2 and generated by $\frac{1}{s}\mathbf{1}_{\mathbf{D}_1}$ on $\mathbf{O}_1 \cup \mathbf{O}_3$, the above computation shows the smoothness of the metric on $\mathcal{O}_{\mathbf{X}}(\mathbf{D}_1)$. Note that ρ vanishes when evaluated on integral points of \mathbf{M}_3 .

A.3. Example III. Here we provide details for the example presented in Section 9. Recall that we are concerned with certain homogeneous variety $\mathbf{M}_3 \cong \mathbf{G}/\mathbf{H}$ with $\mathbf{G} = \mathbf{SL}_3$ and \mathbf{H} being the full diagonal torus, which is compactified by

$$\mathbf{X}^{\text{INC}} = \left\{ (\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3, \mathbf{l}_{12}, \mathbf{l}_{13}, \mathbf{l}_{23}) \in (\mathbf{P}^2)^3 \times (\mathbf{Gr}_{2,3})^3 \mid \mathbf{l}_I \subset \mathbf{l}_J, \ \forall I \subset J \right\}.$$

It is equipped with **G**-invariant closed subvarieties:

$$\begin{split} \mathbf{D}_{123}^1 &:= \left\{ \mathbf{l}_1 = \mathbf{l}_2 = \mathbf{l}_3 \right\}, \ \mathbf{D}_{123}^2 &:= \left\{ \mathbf{l}_{13} = \mathbf{l}_{13} = \mathbf{l}_{23} \right\}, \\ \mathbf{D}_{12,3} &:= \left\{ \mathbf{l}_1 = \mathbf{l}_2, \, \mathbf{l}_{13} = \mathbf{l}_{23} \right\}, \ \mathbf{D}_{13,2} &:= \left\{ \mathbf{l}_1 = \mathbf{l}_3, \, \mathbf{l}_{12} = \mathbf{l}_{23} \right\}, \\ \mathbf{D}_{23,1} &:= \left\{ \mathbf{l}_2 = \mathbf{l}_3, \, \mathbf{l}_{12} = \mathbf{l}_{13} \right\}. \end{split}$$

A.3.1. An open affine subvariety. Using Plucker coordinates, we identify $\mathbf{Gr}_{2,3}$ with \mathbf{P}^2 . So \mathbf{X}^{INC} is a closed subvariety of $(\mathbf{P}^2)^6$. For our purpose, we may restrict to the open subvariety \mathbf{O} where all the first coordinates are non-zero. That is, points of the form

 $\{([1:a_2:a_3], [1:b_2:b_3], [1:c_2:c_3], [1:\alpha_{13}:\alpha_{23}], [1:\beta_{13}:\beta_{23}], [1:\gamma_{13}:\gamma_{23}])\}$

that are contained in \mathbf{X}^{INC} . Thus, \mathbf{O} is a closed subvariety of

 $\operatorname{Spec} \mathbb{R}[a_2, a_3, b_2, b_3, c_2, c_3, \alpha_{13}, \alpha_{23}, \beta_{13}, \beta_{23}, \gamma_{13}, \gamma_{23}]$

defined by the prime ideal generated by (these functions come from the incidence relations)

$$\begin{split} f_1 &:= \alpha_{23} - a_2 \alpha_{13} + a_3, \ f_2 &:= \beta_{23} - a_2 \beta_{13} + a_3, \ f_3 &:= \alpha_{23} - b_2 \alpha_{13} + b_3, \\ f_4 &:= \gamma_{23} - b_2 \gamma_{13} + b_3, \ f_5 &:= \beta_{23} - c_2 \beta_{13} + c_3, \ f_6 &:= \gamma_{23} - c_2 \gamma_{13} + c_3. \end{split}$$

That they indeed generate a prime ideal will be clear later.

A.3.2. The invariant gauge form. We compute the invariant gauge form on \mathbf{M}_3 under these coordinates. Set

$$\eta_{1} := \frac{1}{(\gamma_{23} - a_{2}\gamma_{13} + a_{3})^{3}} da_{2} \wedge da_{3} \wedge d\gamma_{13} \wedge d\gamma_{23};$$

$$\eta_{2} := \frac{1}{(\beta_{23} - b_{2}\beta_{13} + b_{3})^{3}} db_{2} \wedge db_{3} \wedge d\beta_{13} \wedge d\beta_{23};$$

$$\eta_{3} := \frac{1}{(\alpha_{23} - c_{2}\alpha_{13} + c_{3})^{3}} dc_{2} \wedge dc_{3} \wedge d\alpha_{13} \wedge d\alpha_{23}.$$

Lemma A.5. For i = 1, 2, 3, each η_i is **G**-invariant.

Proof. We only present the proof for η_1 . The other two cases are similar.

We write \boldsymbol{a} (resp. $\boldsymbol{\gamma}$) as a shorthand for (a_2, a_3) (resp. $(\gamma_{13}, \gamma_{23})$). Also set $d\boldsymbol{a} := da_2 \wedge da_3$ and $d\boldsymbol{\gamma} := d\gamma_{13} \wedge d\gamma_{23}$. Let $\phi_1(\boldsymbol{a}, \boldsymbol{\gamma}) := (\gamma_{23} - a_2\gamma_{13} + a_3)^3$ and

$$g_1 := \begin{bmatrix} 1 & x & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, g_2 := \begin{bmatrix} 1 & 0 & x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, g_3 := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & x \\ 0 & 0 & 1 \end{bmatrix}, g_4 := \begin{bmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ y & z & 1 \end{bmatrix}.$$

It suffices to verify that η_1 is invariant under g_1, g_2, g_3, g_4 for all x, y, z. We only verify the g_2 -invariance and the other cases are similar.

For functions $f = a_2, a_3, \boldsymbol{a}, \gamma_{13}, \gamma_{23}$ or $\boldsymbol{\gamma}$, let $f^g := g_2^*(f)$. Since

$$\begin{bmatrix} 1 & 0 & x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 1 + xa_3 \\ a_2 \\ a_3 \end{bmatrix},$$

we have

$$a_2^g = \frac{a_2}{1 + xa_3}, \ a_3^g = \frac{a_3}{1 + xa_3}, \ \text{and} \ \mathrm{d} \boldsymbol{a}^g = (1 + xa_3)^{-3} \mathrm{d} \boldsymbol{a}.$$

Under the basis $(\mathbf{e}_1 \wedge \mathbf{e}_2, \mathbf{e}_1 \wedge \mathbf{e}_3, \mathbf{e}_2 \wedge \mathbf{e}_3)$, we have

$$\left(\wedge^{2} \begin{bmatrix} 1 & 0 & x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}\right) \begin{bmatrix} 1 \\ \gamma_{13} \\ \gamma_{23} \end{bmatrix} = \begin{bmatrix} 1 & 0 & -x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ \gamma_{13} \\ \gamma_{23} \end{bmatrix} = \begin{bmatrix} 1 - x\gamma_{23} \\ \gamma_{13} \\ \gamma_{23} \end{bmatrix}.$$

Thus,

$$\gamma_{13}^g = \frac{\gamma_{13}}{1 - x\gamma_{23}}, \ \gamma_{23}^g = \frac{\gamma_{23}}{1 - x\gamma_{23}}, \ \mathrm{d}\gamma^g = (1 - x\gamma_{23})^{-3}\mathrm{d}\gamma.$$

On the other hand,

$$\phi_1(\boldsymbol{a}^g, \boldsymbol{\gamma}^g) = (1 - x\gamma_{23})^{-1}\gamma_{23} - (1 + xa_3)^{-1}a_2 \cdot (1 - x\gamma_{23})^{-1}\gamma_{13} + (1 + xa_3)^{-1}a_3$$
$$= (1 - x\gamma_{23})^{-1}(1 + xa_3)^{-1}\phi_1(\boldsymbol{a}, \boldsymbol{\gamma}).$$

It follows that

$$\phi_1^{-3}(\boldsymbol{a}^g,\boldsymbol{\gamma}^g)\mathrm{d}\boldsymbol{a}^g\wedge\mathrm{d}\boldsymbol{\gamma}^g=\phi_1^{-3}(\boldsymbol{a},\boldsymbol{\gamma})\mathrm{d}\boldsymbol{a}\wedge\mathrm{d}\boldsymbol{\gamma}.$$

That is to say, η_1 is g_2 -invariant.

It can be checked that f_i 's intersect transversally in an open set. Thus, the solution ω to the equation

$$\eta_1 \wedge \eta_2 \wedge \eta_3 = \omega \wedge \mathrm{d}f_1 \wedge \dots \wedge \mathrm{d}f_6 \tag{41}$$

is unique when restricted to the common zero set of f_i 's. As η_i 's and df_i 's are **G**-invariant. It follows that such an ω would also be **G**-invariant.

Lemma A.6. Let

$$\omega_{\mathbf{M}_3} := \frac{\mathrm{d}a_2 \wedge \mathrm{d}a_3 \wedge \mathrm{d}b_2 \wedge \mathrm{d}c_2 \wedge \mathrm{d}\alpha_{13} \wedge \mathrm{d}\gamma_{13}}{(a_2 - b_2)^6 (\alpha_{13} - \gamma_{13})^3 (\beta_{13} - \alpha_{13})^6 (a_2 - c_2)^4}.$$

Then $\omega := \omega_{\mathbf{M}_3}$ solves Equa.(41) above.

Proof. Note that by $f_1 = \ldots = f_6 = 0$, we have

$$\begin{split} \gamma_{23} - a_2\gamma_{13} + a_3 &= b_2\gamma_{13} - b_3 - a_2\gamma_{13} + a_3 = b_2\gamma_{13} + \alpha_{23} - b_2\alpha_{13} - a_2\gamma_{13} + a_3 \\ &= (b_2 - a_2)\gamma_{13} + (a_2 - b_2)\alpha_{13} = (a_2 - b_2)(\alpha_{13} - \gamma_{13}); \\ \beta_{23} - b_2\beta_{13} + b_3 &= a_2\beta_{13} - a_3 - b_2\beta_{13} + b_2\alpha_{13} - \alpha_{23} \\ &= (a_2 - b_2)\beta_{13} - a_3 + b_2\alpha_{13} - a_2\alpha_{13} + a_3 = (a_2 - b_2)(\beta_{13} - \alpha_{13}); \\ \alpha_{23} - c_2\alpha_{13} + c_3 &= a_2\alpha_{13} - a_3 - c_2\alpha_{13} + c_2\beta_{13} - \beta_{23} \\ &= (a_2 - c_2)\alpha_{13} + c_2\beta_{13} - a_2\beta_{13} = (a_2 - c_2)(\alpha_{13} - \beta_{13}). \end{split}$$

And modulo (the kernel of) $\omega_{\mathbf{M}_3} \wedge \cdot$, we have

$$df_1 \wedge df_2 \wedge df_3 \wedge df_4 \wedge df_5 \wedge df_6$$

$$\equiv d\alpha_{23} \wedge (d\beta_{23} - a_2 d\beta_{13}) \wedge db_3 \wedge d\gamma_{23} \wedge (-c_2 d\beta_{13} + d\beta_{23}) \wedge dc_3$$

$$\equiv (a_2 - c_2) d\alpha_{23} \wedge d\beta_{23} \wedge d\beta_{13} \wedge db_3 \wedge d\gamma_{23} \wedge dc_3.$$

Putting these computations together yields the result.

A.3.3. New coordinates. We perform a few change of variables and eliminate a few redundant variables to make the description of \mathbf{O} clearer. The explicit formula of the group action in the new coordinates will be quite complicated but we are not concerned about it.

Using $f_1, f_2, f_3, f_4, f_5 = 0$, we can eliminate

$$\begin{cases} \alpha_{23} = a_2 \alpha_{13} - a_3, \\ \beta_{23} = a_2 \beta_{13} - a_3, \\ b_3 = b_2 \alpha_{13} - a_2 \alpha_{13} + a_3, \\ \gamma_{23} = b_2 \gamma_{13} - b_2 \alpha_{13} + a_2 \alpha_{13} - a_3, \\ c_3 = c_2 \beta_{13} - a_2 \beta_{13} + a_3 \end{cases}$$

And $f_6 = 0$ is equivalent to

$$(b_2\gamma_{13} - b_2\alpha_{13} + a_2\alpha_{13} - a_3) - c_2\gamma_{13} + (c_2\beta_{13} - a_2\beta_{13} + a_3) = 0$$

$$\iff b_2(\gamma_{13} - \alpha_{13}) + a_2(\alpha_{13} - \beta_{13}) + c_2(\beta_{13} - \gamma_{13}) = 0$$

$$\iff -(a_2 - c_2)(\beta_{13} - \alpha_{13}) + (b_2 - c_2)(\gamma_{13} - \alpha_{13}) = 0.$$

Now we replace $\beta_{13}, \gamma_{13}, a'_2$ and b'_2 by

$$\beta'_{13} := \beta_{13} - \alpha_{13}, \ \gamma'_{13} := \gamma_{13} - \alpha_{13}, \ a'_2 := a_2 - c_2, \ b'_2 := b_2 - c_2.$$

Hence the open subvariety $\mathbf{O} \subset \mathbf{X}^{\text{INC}}$ is identified with the zero set of the prime ideal

$$\langle -a_2'\beta_{13}' + \gamma_{13}'b_2'\rangle \text{ in } \operatorname{Spec} \mathbb{Q}[a_3, c_2, \alpha_{13}] \times \operatorname{Spec} \mathbb{Q}[a_2', b_2', \beta_{13}', \gamma_{13}'].$$

Thus **O** is singular exactly when $a'_2 = \beta'_{13} = \gamma'_{13} = b'_2 = 0$. The first three coordinates a_3, c_2, α_{13} have no effect on the discussions below and are often omitted for simplicity.

A.3.4. Invariant gauge form in the new coordinates. Under the new coordinates, we have

$$\omega_{\mathbf{M}_3} = \pm (\mathrm{d}a_3 \wedge \mathrm{d}c_2 \wedge \mathrm{d}\alpha_{13}) \wedge \frac{\mathrm{d}a'_2 \wedge \mathrm{d}b'_2 \wedge \mathrm{d}\gamma'_{13}}{(a'_2 - b'_2)^6 (\gamma'_{13})^3 (\beta'_{13})^6 (a'_2)^4}.$$

A.3.5. Boundary stratum in the new coordinates. We list the ideal for \mathbf{O} and various boundary stratum intersecting \mathbf{O} :

The ideal of
$$\begin{cases} \mathbf{O} \text{ is generated by } & -a'_{2}\beta'_{13} + \gamma'_{13}b'_{2} \\ \mathbf{D}_{123}^{1} \text{ is generated by } & a'_{2}, b'_{2} \\ \mathbf{D}_{12,3} \text{ is generated by } & a'_{2} - b'_{2}, \beta'_{13} - \gamma'_{13} \\ \mathbf{D}_{13,2} \text{ is generated by } & a'_{2}, \gamma'_{13} \\ \mathbf{D}_{23,1} \text{ is generated by } & b'_{2}, \beta'_{13} \\ \mathbf{D}_{2123}^{2} \text{ is generated by } & \beta'_{13}, \gamma'_{13} \end{cases}$$

A.3.6. Blowup. We blow up the ideal $\langle a'_2, b'_2 \rangle$ of \mathbf{D}^1_{123} and verify in local coordinates that the obtained pair $(\mathbf{X}^1, \mathbf{D}^1)$ is, as has been pointed out in [RS84], smooth. Also we calculate the invariant gauge form in this smooth pair.

Naturally $\mathbf{Bl}_{\mathbf{D}_{123}^1}(\mathbf{O})$ is the union of two affine open subvarieties $\mathbf{U}_1 \cup \mathbf{U}_2$. For simplicity, coordinates a_3, c_2, α_{13} are omitted below.

A.3.7. Blowup, U₁. Here we have coordinates $\beta'_{13}, \gamma'_{13}, a'_2, \tilde{b}_2$ with $b'_2 = a'_2 \tilde{b}_2$. And the equation defining $\mathbf{Bl}_{\mathbf{D}_{123}^1}(\mathbf{O})$ is $\beta'_{13} = \gamma'_{13}\widetilde{b}_2$. Thus we can eliminate β'_{13} and treat \mathbf{U}_1 as $\operatorname{Spec} \mathbb{Q}[\gamma'_{13}, a'_2, \widetilde{b}_2]$. One can compute that (let \mathbf{E} denote the exceptional divisor)

- 1. $\omega_{\mathbf{M}_3} = \pm (a_2')^{-9} (1 \tilde{b}_2)^{-6} (\gamma_{13}')^{-9} (\tilde{b}_2)^{-6} \cdot \mathrm{d}a_2' \wedge \mathrm{d}\tilde{b}_2 \wedge \mathrm{d}\gamma_{13}';$ 2. $\mathbf{D}_{12,3}^+ = \{1 - \tilde{b}_2 = 0\}, \mathbf{D}_{13,2}^+$ is in the complement, $\mathbf{D}_{23,1}^+ = \{\tilde{b}_2 = 0\};$ 3. $\mathbf{E} = \{a_2' = 0\}, (\mathbf{D}_{123}^2)^+ = \{\gamma_{13}' = 0\}.$

A.3.8. Blowup, U₂. Here we have coordinates $\beta'_{13}, \gamma'_{13}, b'_2, \tilde{a}_2$ with $a'_2 = b'_2 \tilde{a}_2$. And the equation defining $\mathbf{Bl}_{\mathbf{D}_{123}^1}(\mathbf{O})$ in \mathbf{U}_1 is $\gamma'_{13} = \beta'_{13}\tilde{a}_2$. Thus we can eliminate γ'_{13} and treat \mathbf{U}_2 as $\operatorname{Spec} \mathbb{Q}[\beta'_{13}, b'_2, \tilde{a}_2]$. One can compute that (let \mathbf{E} denote the exceptional divisor)

- 1. $\omega_{\mathbf{M}_3} = \pm (b'_2)^{-9} (1 \tilde{a}_2)^{-6} (\beta'_{13})^{-9} (\tilde{a}_2)^{-6} \cdot db'_2 \wedge d\tilde{a}_2 \wedge d\beta'_{13};$ 2. $\mathbf{D}^+_{12,3} = \{1 \tilde{a}_2 = 0\}, \mathbf{D}^+_{13,2} = \{\tilde{a}_2 = 0\}, \mathbf{D}^+_{23,1}$ is in the complement; 3. $\mathbf{E} = \{b'_2 = 0\}, (\mathbf{D}^2_{123})^+ = \{\beta'_{13} = 0\}.$

A.3.9. The divisor of the invariant gauge form. Labeling the boundary divisors on \mathbf{X}^2 as

$$\mathbf{D}_1 := \mathbf{E}, \ \mathbf{D}_2 := \mathbf{D}_{12,3}^+, \ \mathbf{D}_3 := \mathbf{D}_{13,2}^+, \ \mathbf{D}_4 := \mathbf{D}_{23,1}^+, \ \mathbf{D}_5 := (\mathbf{D}_{123}^2)^+$$

From the description above, one gets

$$-\operatorname{div}(\omega_{\mathbf{M}_3}) = 9\mathbf{D}_1 + 6\mathbf{D}_2 + 6\mathbf{D}_3 + 6\mathbf{D}_4 + 9\mathbf{D}_5$$

A.3.10. Mapping to the measure compactification space. Let $\Gamma \leq \mathbf{G}(\mathbb{Q})$ be an arithmetic subgroup. We have the following, implied by [SZ19, Theorem 2.4].

Theorem A.7. Let (g_n) be a sequence in G.

1. If (g_n) is unbounded modulo H_{ij} for every $\{i, j\} \subset \{1, 2, 3\}$, then

$$\lim_{n \to \infty} \left[(g_n)_* \mathbf{m}_{[\mathrm{H}]} \right] = \left[\mathbf{m}_{[\mathbf{G}]} \right];$$

2. If (g_n) is unbounded modulo H but convergent to $\delta \in G$ modulo H_{ij} for some (necessarily unique) $\{i, j\}$, then

$$\lim_{n \to \infty} \left[(g_n)_* \mathbf{m}_{[\mathbf{H}]} \right] = \left[\delta_* \mathbf{m}_{[\mathbf{H}_{ij}]} \right]$$

As a corollary, we have

Corollary A.8. Let X^{meas} be the closure of $\{\alpha_{G}^{\psi}(m_{[H]}^{\psi})\}$ in $\text{Prob}^{\psi}(G/\Gamma)$. Then

$$X^{\text{meas}} = \alpha_{\text{G}}^{\psi}(\mathbf{m}_{[\text{H}]}^{\psi}) \bigsqcup \left(\alpha_{\text{G}}^{\psi}(\mathbf{m}_{[\text{H}_{12}]}^{\psi}) \cup \alpha_{\text{G}}^{\psi}(\mathbf{m}_{[\text{H}_{13}]}^{\psi}) \cup \alpha_{\text{G}}^{\psi}(\mathbf{m}_{[\text{H}_{23}]}^{\psi}) \right) \bigsqcup \left\{ \mathbf{m}_{[\text{G}]}^{\psi} \right\}.$$

Appendix B. Divergence of Translates of Homogeneous Closed Subsets

Let **G** be a connected linear algebraic group over \mathbb{Q} , **H** be a connected observable \mathbb{Q} -subgroup, and Γ be an arithmetic subgroup of **G**. Fix a maximal reductive connected Q-subgroup **G**^{red} of **G** and hence $\mathbf{G} = \mathbf{G}^{\mathrm{red}} \ltimes \mathbf{R}_{\mathbf{u}}(\mathbf{G})$. Also fix a Cartan involution and hence a maximal compact subgroup K of $\mathbf{G}^{\mathrm{red}}(\mathbb{R})$. Depending on this choice, each parabolic \mathbb{Q} -subgroup \mathbf{P} is associated with a \mathbb{Q} -split subtorus $\mathbf{A}_{\mathbf{P},K}$ that is isomorphic to $\mathbf{S}_{\mathbf{P}}$ under $p_{\mathbf{P}}^{\text{spl}}$. Unlike the main body of the paper, in this appendix, the Roman letter L is used to denote $\mathbf{L}(\mathbb{R})^{\circ}$ for an algebraic group over \mathbb{R} .

Let $\Phi^{\mathrm{red}}(\mathbf{A}_{\mathbf{P},\mathrm{K}},\mathbf{P})$ be the non-trivial characters of $\mathbf{A}_{\mathbf{P},\mathrm{K}}$ appearing in $\mathfrak{r}_{\mathbf{u}}(\mathfrak{p})/\mathfrak{r}_{\mathbf{u}}(\mathfrak{g}) \subset \mathfrak{g}/\mathfrak{r}_{\mathbf{u}}(\mathfrak{g})$. Let $\Delta^{\mathrm{red}}(\mathbf{A}_{\mathbf{P},K},\mathbf{P})$ be the subset of cardinality dim $\mathbf{A}_{\mathbf{P},K}$ whose $\mathbb{Z}_{>0}$ -span equals to $\Phi^{\mathrm{red}}(\mathbf{A}_{\mathbf{P},K},\mathbf{P})$. When **G** is already reductive, the superscript "red" is dropped.

Theorem B.1. Given a sequence $(g_n) \subset G$, after passing to a subsequence, there exist a sequence $(h_n) \subset H$, $(\gamma_n) \subset \Gamma$ and a parabolic Q-subgroup **P** such that the following holds. Write $g_n h_n \gamma_n^{-1} = k_n a_n p_n$ using horospherical coordinates of (**P**, K). Then

- (1) (p_n) is bounded;
- (2) $\alpha(a_n) \to 0$ for every $\alpha \in \Delta^{\operatorname{red}}(\mathbf{A}_{\mathbf{P},\mathrm{K}},\mathbf{P});$
- (3) if (a_n) is unbounded, there exist a \mathbb{Q} -representation \mathbf{V} of \mathbf{G} factoring through $\mathbf{G}/\mathbf{R}_{\mathbf{u}}(\mathbf{G})$ and $\mathbf{v} \in \mathbf{V}(\mathbb{Q})$ such that the line spanned by \mathbf{v} is preserved by \mathbf{P} , \mathbf{v} is fixed by $\gamma_n \mathbf{H} \gamma_n^{-1}$ for all n and $\lim_{n\to\infty} a_n \cdot \mathbf{v} = \mathbf{0}$;
- (4) $(\gamma_n \mathbf{H} \gamma_n^{-1})$ strongly converges to some observable subgroup of **G**.

Compared to [Zha22, Theorem 5.2], 3 and 4 are new. Note that whereas one quotients Γ from the left in [Zha22], here we choose to quotient from the right. Also, when one has $\alpha(a_n) \to \infty$ in [Zha22], we have $\alpha(a_n) \to 0$ here. We shall follow the logic of [Zha22] by first reducing to the case of $\mathbf{G} = \mathbf{SL}_N$. And this special case will be handled with the help of "canonical polygons". The proof is independent from the work of [DGU23] and can actually be used to give an alternative treatment of their main results. The connection between canonical polygons and nondivergence property of unipotent flows has been noted in [dS23].

For simplicity, by saying the (\mathbf{P}, \mathbf{K}) coordinate of some element, we mean its **horospherical co-ordinate** associated to (\mathbf{P}, \mathbf{K}) (see [Zha22, Lemma 2.1], [BJ06, Chapter 9] or [BS73, Proposition 1.5]).

B.1. Reduction to the reductive case. We explain how the general case follows from the case when **G** is reductive. Indeed, by projecting everything to $p^{\text{red}} : \mathbf{G} \to \mathbf{G}/\mathbf{R}_{\mathbf{u}}(\mathbf{G})$, after passing to a subsequence, we find $(h_n) \subset \mathbf{H}, (\lambda_n) \subset \Gamma$ and a parabolic Q-subgroup **P** such that if $g_n h_n \lambda_n^{-1} = k_n a_n q_n$ is the (\mathbf{P}, \mathbf{K}) coordinate of $g_n h_n \lambda_n^{-1}$, then

- 1. $q_n = b_n u_n$ for some (b_n) bounded in P and $(u_n) \subset \mathbf{R}_{\mathbf{u}}(\mathbf{G})(\mathbb{R})$.
- 2. $\alpha(a_n) \to 0$ for every $\alpha \in \Delta^{\operatorname{red}}(\mathbf{A}_{\mathbf{P},\mathrm{K}},\mathbf{P});$
- 3. there exist a Q-representation V of G factoring through p^{red} and $\mathbf{v} \in \mathbf{V}(\mathbb{Q})$ such that the line spanned by v is preserved by P, v is fixed by $\gamma_n \mathbf{H} \gamma_n^{-1}$ for all n and $\lim_{n\to\infty} a_n \cdot \mathbf{v} = \mathbf{0}$;
- 4. $(p^{\text{red}}(\lambda_n \mathbf{H} \lambda_n^{-1}))$ strongly converges to some observable subgroup of $\mathbf{G}/\mathbf{R}_{\mathbf{u}}(\mathbf{G})$.

Write $u_n = b'_n \lambda'_n$ for some bounded sequence $(b'_n) \subset \mathbf{R}_{\mathbf{u}}(\mathbf{G})(\mathbb{R})$ and $(\lambda'_n) \subset \mathbf{R}_{\mathbf{u}}(\mathbf{G})(\mathbb{R}) \cap \Gamma$. Let $\gamma_n := \lambda'_n \lambda_n$. It only remains to verify item 4. in the theorem. Since (λ'_n) are contained in the kernel of p^{red} , we still have $(p^{\text{red}}(\gamma_n \mathbf{H}\gamma_n^{-1}))$ strongly converges to some observable subgroup \mathbf{F} of $\mathbf{G}/\mathbf{R}_{\mathbf{u}}(\mathbf{G})$. After passing to a subsequence assume $(\gamma_n \mathbf{H}\gamma_n^{-1})$ strongly converges to \mathbf{L} . Then $p^{\text{red}}(\mathbf{L}) = \mathbf{F}$. Note that $(p^{\text{red}})^{-1}(\mathbf{F})$ is observable in \mathbf{G} by assumption. On the other hand, $(p^{\text{red}})^{-1}(\mathbf{F}) = \mathbf{L} \cdot \mathbf{R}_{\mathbf{u}}(\mathbf{G})$ and hence $(p^{\text{red}})^{-1}(\mathbf{F})/\mathbf{L} \cong \mathbf{R}_{\mathbf{u}}(\mathbf{G})/\mathbf{U}$ for some \mathbb{Q} -subgroup \mathbf{U} of $\mathbf{R}_{\mathbf{u}}(\mathbf{G})$. Therefore, \mathbf{L} is observable in $(p^{\text{red}})^{-1}(\mathbf{F})$ and hence in \mathbf{G} . This completes the proof.

B.2. Reduction to the SL_N case. Let **G** be a connected reductive linear algebraic group over \mathbb{Q} . Without loss of generality, we assume that **G** is a \mathbb{Q} -subgroup of some \mathbf{SL}_N that is invariant under taking transpose. We let $K := \mathbf{G}(\mathbb{R}) \cap \mathbf{SO}_N(\mathbb{R})$ be the maximal compact subgroup of $\mathbf{G}(\mathbb{R})$ that is transpose invariant.

The concept of Siegel sets can be generalized to reductive groups as follows. Write $\mathbf{G} = \mathbf{G}^{ss} \cdot \mathbf{Z}(\mathbf{G})^{an} \cdot \mathbf{Z}(\mathbf{G})^{spl}$. Then every Siegel set \mathfrak{S} of \mathbf{G} is of the form $\mathfrak{S}' \cdot F \cdot \mathbf{Z}(\mathbf{G})^{spl}(\mathbb{R})^{\circ}$ where $\mathfrak{S} \subset \mathbf{G}^{ss}(\mathbb{R})$ is a Siegel set for \mathbf{G}^{ss} and $F \subset \mathbf{Z}(\mathbf{G})^{an}(\mathbb{R})$ is a compact subset that is left invariant under the maximal compact subtorus. For every \mathbb{Q} -minimal parabolic subgroup \mathbf{P} of \mathbf{G} , there exist finitely many $c_1, ..., c_l \in \mathbf{G}(\mathbb{Q})$ and a Siegel set \mathfrak{S} associated with (\mathbf{P}, \mathbf{K}) such that

$$\mathbf{G}(\mathbb{R}) = \bigcup_{i=1}^{l} \mathfrak{S} \cdot c_i \cdot \Gamma.$$

The main argument of [Zha22, Section 5] carries through. We sketch the proof below. Without loss of generality, assume $\Gamma = \mathbf{SL}_N(\mathbb{Z}) \cap \mathbf{G}(\mathbb{R})$.

By the \mathbf{SL}_N case, after passing to a subsequence, there are $(h_n) \subset \mathrm{H}, (\gamma_n) \subset \mathbf{SL}_N(\mathbb{Z})$ and a parabolic \mathbb{Q} -subgroup $\mathbf{Q} \subset \mathbf{SL}_N$ such that if the $(\mathbf{Q}, \mathrm{K}')$ coordinate (for simplicity we write $\mathrm{K}' := \mathbf{SO}_N(\mathbb{R})$) of $g_n h_n \gamma_n^{-1}$ is $k_n a_n q_n$, then

- 1. (q_n) is bounded;
- 2. $\alpha(a_n) \to 0$ for every $\alpha \in \Delta(\mathbf{A}_{\mathbf{Q},\mathrm{K}'},\mathbf{Q});$
- 3. $\gamma_n \mathbf{H} \gamma_n^{-1} \subset \mathbf{Q}$ for all n. And there exist a \mathbf{SL}_N -representation and a \mathbb{Q} -vector \mathbf{v} fixed by $\gamma_n \mathbf{H} \gamma_n^{-1}$ and the line spanned by \mathbf{v} is preserved by \mathbf{Q} with a non-trivial character;
- 4. $(\gamma_n \mathbf{H} \boldsymbol{\gamma}_n^{-1})$ strongly converges to an observable subgroup \mathbf{L} of \mathbf{SL}_N .

Note that $a_n \cdot \mathbf{v} \to \mathbf{0}$ follows automatically.

B.2.1. Step 1. Assume $\mathfrak{sl}_N(\mathbb{Z}) \cap \operatorname{Lie}(\mathbf{R}_{\mathbf{u}}(\mathbf{Q})) = \bigoplus \mathbb{Z} w_i^{\mathbf{Q}}$ and let $\mathbf{w} := \bigoplus w_i^{\mathbf{Q}}$. Then $\lim_{n \to \infty} g_n h_n \gamma_n^{-1} \cdot \mathbf{w} = \mathbf{0}$.

By geometric invariant theory, there exist $n_0 \in \mathbb{Z}^+$ and $\mathbf{a}_t \in \mathfrak{X}^{\mathbb{Q}}_*(\mathbf{G})$ such that

$$\lim_{t \to \infty} \mathbf{a}_t \gamma_{n_0}^{-1} \cdot \mathbf{w} = 0$$

Let (similar notations are also used for cocharacters other than \mathbf{a}_t)

$$\mathbf{Q}_{\mathbf{a}_{t}} := \left\{ x \in \mathbf{SL}_{N} \mid \lim_{t \to \infty} \mathbf{a}_{t} x \mathbf{a}_{t}^{-1} \text{ exists} \right\};$$
$$\mathbf{P}_{\mathbf{a}_{t}} := \left\{ x \in \mathbf{G} \mid \lim_{t \to \infty} \mathbf{a}_{t} x \mathbf{a}_{t}^{-1} \text{ exists} \right\}.$$

Then $\mathbf{R}_{\mathbf{u}}(\mathbf{Q}_{\mathbf{a}_{t}})$ contains $\mathbf{R}_{\mathbf{u}}(\gamma_{n_{0}}^{-1}\mathbf{Q}\gamma_{n_{0}})$, or equivalently, $\mathbf{Q}_{\mathbf{a}_{t}} \subset \gamma_{n_{0}}^{-1}\mathbf{Q}\gamma_{n_{0}}$. Replacing \mathbf{Q} by $\gamma_{n_{0}}^{-1}\mathbf{Q}\gamma_{n_{0}}$ and γ_{n} by $\gamma_{n_{0}}^{-1}\gamma_{n}$, we assume without loss of generality that $\gamma_{n_{0}}$ is the identity element.

B.2.2. Step 2. Let $\mathbf{L}_{\mathbf{P}_{\mathbf{a}_{t}},K}$ (resp. $\mathbf{L}_{\mathbf{Q}_{\mathbf{a}_{t}},K'}$) be the maximal reductive subgroup of $\mathbf{P}_{\mathbf{a}_{t}}$ (resp. $\mathbf{Q}_{\mathbf{a}_{t}}$) that is invariant under taking transpose. Let $\mathbf{A}_{\mathbf{P}_{\mathbf{a}_{t}},K}$ be the central torus of $\mathbf{L}_{\mathbf{P}_{\mathbf{a}_{t}},K}$, which is conjugate to a maximal \mathbb{Q} -split torus \mathbf{S}_{\max} in $\mathbf{P}_{\mathbf{a}_{t}}$. We have $\mathbf{L}_{\mathbf{P}_{\mathbf{a}_{t}},K} \subset \mathbf{L}_{\mathbf{Q}_{\mathbf{a}_{t}},K'}$ and $\mathbf{R}_{\mathbf{u}}(\mathbf{P}_{\mathbf{a}_{t}}) \subset \mathbf{R}_{\mathbf{u}}(\mathbf{Q}_{\mathbf{a}_{t}})$. There exists $\mathbf{b}_{t} \in \mathfrak{X}^{\mathbb{Q}}_{*}(\mathbf{S}_{\max})$ such that

- 1. the centralizer of $\{\mathbf{b}_t\}$ in \mathbf{SL}_N is equal to the centralizer of \mathbf{S}_{\max} in \mathbf{SL}_N . In particular $\mathbf{P}_{\mathbf{b}_t}$ is a \mathbb{Q} -minimal parabolic subgroup of \mathbf{G} ;
- 2. $\mathbf{Q}_{\mathbf{b}_t} \subset \mathbf{Q}_{\mathbf{a}_t}$. In particular, $\mathbf{P}_{\mathbf{b}_t} \subset \mathbf{P}_{\mathbf{a}_t}$.

Thus $\mathbf{A}_{\mathbf{P}_{\mathbf{b}_{t}},\mathbf{K}}$ is contained in $\mathbf{A}_{\mathbf{Q}_{\mathbf{b}_{t}},\mathbf{K}'}$. Consequently, the $(\mathbf{P}_{\mathbf{b}_{t}},\mathbf{K})$ coordinate of $g \in \mathbf{G}(\mathbb{R})$ is the same as its $(\mathbf{Q}_{\mathbf{b}_{t}},\mathbf{K}')$ coordinate.

B.2.3. Step 3. Because $\mathbf{P}_{\mathbf{b}_t}$ is \mathbb{Q} -minimal, by passing to a subsequence, there exist $c_1 \in \mathbf{G}(\mathbb{Q}), (\lambda_n) \subset \Gamma$ and a Siegel set $\mathfrak{S}(\mathbf{P}_{\mathbf{b}_t})$ associated to $(\mathbf{P}_{\mathbf{b}_t}, \mathbf{K})$ such that

$$g_n h_n \in \mathfrak{S}(\mathbf{P}_{\mathbf{b}_t}) c_1 \lambda_n, \quad \forall n \in \mathbb{Z}^+$$

In particular, if $g_n h_n \lambda_n^{-1} c_1^{-1} = k'_n a'_n p'_n$ under $(\mathbf{P}_{\mathbf{b}_t}, \mathbf{K})$ coordinate, then (p'_n) is bounded and for some $t_0 > 0$, for every n and $\alpha \in \Delta(\mathbf{A}_{\mathbf{P}_{\mathbf{b}_t}, \mathbf{K}}, \mathbf{P}_{\mathbf{b}_t})$, one has $\alpha(a'_n) < t_0$.

Let $\Phi(\mathbf{A}_{\mathbf{Q}_{\mathbf{b}_{t}},\mathbf{K}'},\mathfrak{sl}_{N})$ be the collection of non-trivial weights of $\mathbf{A}_{\mathbf{Q}_{\mathbf{b}_{t}},\mathbf{K}'}$ appearing in the adjoint action on \mathfrak{sl}_{N} . Passing to a subsequence, assume that for every $\beta \in \Phi(\mathbf{A}_{\mathbf{Q}_{\mathbf{b}_{t}},\mathbf{K}'},\mathfrak{sl}_{N})$, the sequence $(\beta(a'_{n}))$ is bounded (away from 0 and $+\infty$), or converges to 0, or diverges to $+\infty$. Define a parabolic \mathbb{Q} -subgroup by

$$\mathbf{Q}_{(a'_n)} := \left\{ x \in \mathbf{SL}_N \mid \left(a'_n x a'_n^{-1} \right) \text{ is bounded} \right\}.$$

One notes that $\mathbf{P}_{\mathbf{b}_t}$ is contained in $\mathbf{Q}_{(a'_n)}$.

Since (a'_n) is a sequence in $A_{\mathbf{P}_{\mathbf{b}_t},\mathbf{K}}$, which is conjugate to S_{\max} in $\mathbf{P}_{\mathbf{b}_t}$, we can choose $\mathbf{b}'_t \in \mathfrak{X}^{\mathbb{Q}}_*(\mathbf{S}_{\max})$ such that

$$\mathbf{Q}_{\mathbf{b}'_t} \subset \mathbf{Q}_{(a'_n)}, \ \mathbf{P}_{\mathbf{b}'_t} = \mathbf{P}_{\mathbf{b}_t}, \ \mathbf{Z}_{\mathbf{SL}_N}(\{\mathbf{b}'_t\}) = \mathbf{Z}_{\mathbf{SL}_N}(\mathbf{S}_{\max}).$$

Then the $(\mathbf{P}_{\mathbf{b}_t} = \mathbf{P}_{\mathbf{b}'_t}, \mathbf{K})$ coordinate of $g_n h_n \lambda_n^{-1} c_1^{-1}$ is the same as its $(\mathbf{Q}_{\mathbf{b}'_t}, \mathbf{K}')$ coordinate. From the definition of $\mathbf{Q}_{(a'_n)}$, one deduces that for any weight $\alpha \in \Phi(\mathbf{A}_{\mathbf{P}_{\mathbf{b}'_t}, \mathbf{K}}, \mathbf{Q}_{(a'_n)})$ and in particular $\alpha \in \Phi(\mathbf{A}_{\mathbf{P}_{\mathbf{b}'_t}, \mathbf{K}}, \mathbf{Q}_{\mathbf{b}'_t})$, we have $\alpha(a'_n) < t_1$ for some $t_1 > 0$ and for all n. We conclude from here that $g_n h_n \lambda_n^{-1} c_1^{-1}$ is contained in a Siegel set $\mathfrak{S}(\mathbf{Q}_{\mathbf{b}'_t})$ associated to $(\mathbf{Q}_{\mathbf{b}'_t}, \mathbf{K}')$. B.2.4. Step 4. Though $\mathbf{P}_{\mathbf{b}_t} = \mathbf{P}_{\mathbf{b}'_t}$, it is not clear whether $\mathbf{Q}_{\mathbf{b}_t}$ and $\mathbf{Q}_{\mathbf{b}'_t}$ contain a common Borel subgroup. Find $w_1 \in \mathbf{SL}_N(\mathbb{Q})$ such that $\mathbf{Q}' := w_1 \mathbf{Q}_{\mathbf{b}'_t} w_1^{-1} \cap \mathbf{Q}_{\mathbf{b}_t}$ is a parabolic subgroup. Thus $g_n h_n \lambda_n^{-1} c_1^{-1} w_1^{-1}$ is contained in some Siegel set of $(w_1 \mathbf{Q}_{\mathbf{b}'_t} w_1^{-1}, \mathbf{K}')$ and hence some Siegel set $\mathfrak{S}(\mathbf{Q}')$ associated to $(\mathbf{Q}', \mathbf{K}')$. On the other hand, $(g_n h_n \gamma_n^{-1})$ is known to be contained in some Siegel set attached to $(\mathbf{Q}, \mathbf{K}')$ and hence in some Siegel set $\mathfrak{S}(\mathbf{Q}')'$ attached to $(\mathbf{Q}', \mathbf{K}')$. Replacing by a larger Siegel set if necessary, assume $\mathfrak{S}(\mathbf{Q}') = \mathfrak{S}(\mathbf{Q}')'$. Hence

$$g_n h_n \gamma_n^{-1} \in \mathfrak{S}(\mathbf{Q}') \cap \mathfrak{S}(\mathbf{Q}') w_1 c_1 \lambda_n \gamma_n^{-1}$$

In particular, this intersection is non-empty. But for every $q \in \mathbf{SL}_N(\mathbb{Q})$, the set $\{\gamma \in \mathbf{SL}_N(\mathbb{Z}), \mathfrak{S}(\mathbf{Q}') \cap \mathfrak{S}(\mathbf{Q}')q\gamma\}$ is finite. Therefore, after passing to a subsequence, there exists $\gamma'_1 \in \mathbf{SL}_N(\mathbb{Z})$ such that $\lambda_n \gamma_n^{-1} = \gamma'_1$ for all n.

B.2.5. Step 5. Let $\mathbf{Q}'' := \gamma'_1 \mathbf{Q} \gamma'_1^{-1}$, which contains $\lambda_n \mathbf{H} \lambda_n^{-1}$ for all *n*. As in Step 1 and 2, find some maximal \mathbb{Q} -split torus \mathbf{S}'_{\max} of \mathbf{G} and $\mathbf{b}''_t \in \mathfrak{X}^{\mathbb{Q}}_*(\mathbf{S}'_{\max})$ such that

1. $\mathbf{Z}_{\mathbf{SL}_N}(\{\mathbf{b}_t''\}) = \mathbf{Z}_{\mathbf{SL}_N}(\mathbf{S}'_{\max})$. In particular, $\mathbf{P}_{\mathbf{b}_t''}$ is a \mathbb{Q} -minimal parabolic subgroup; 2. $\mathbf{Q}_{\mathbf{b}_t''} \subset \mathbf{Q}''$.

By assumption, $g_n h_n \lambda_n^{-1} = g_n h_n \gamma_n^{-1} \gamma_1'^{-1}$ belongs to some Siegel set associated to $(\mathbf{Q}'', \mathbf{K}')$ and hence to $(\mathbf{Q}_{\mathbf{b}''_{t}}, \mathbf{K}')$. Let

$$g_n h_n \lambda_n^{-1} = k_n'' a_n'' p_n''$$

be the $(\mathbf{P}_{\mathbf{b}''_{t}}, \mathbf{K})$ coordinate, which is the same as the $(\mathbf{Q}_{\mathbf{b}''_{t}}, \mathbf{K}')$ coordinate. We find that

- 1. (p''_n) is bounded;
- 2. there exists $t_2 > 0$ such that for every n and $\alpha \in \Delta(\mathbf{A}_{\mathbf{P}_{\mathbf{b}''},\mathbf{K}},\mathbf{P}_{\mathbf{b}''})$, we have $\alpha(a''_n) < t_2$;

3.
$$\mathbf{Q}'' = \mathbf{Q}_{(q_n h_n \lambda_n^{-1})} = \mathbf{Q}_{(a_n'')}.$$

Choose $\mathbf{c}_t \in \mathfrak{X}^{\mathbb{Q}}_*(\mathbf{S}'_{\max})$ such that $\mathbf{Q}_{(a''_n)} = \mathbf{Q}_{\mathbf{c}_t}$. After passing to a subsequence, assume that for every $\alpha \in \Delta(\mathbf{A}_{\mathbf{P}_{\mathbf{b}'_t},\mathbf{K}},\mathbf{P}_{\mathbf{b}'_t})$, either $(\alpha(a''_n))$ converges to 0 or is bounded away from 0. Let I'' be those α such that $(\alpha(a''_n))$ is bounded away from 0. Then $(\mathbf{P}_{\mathbf{b}'_t})_{I''} = \mathbf{P}_{\mathbf{c}_t} = \mathbf{Q}_{\mathbf{c}_t} \cap \mathbf{G} = \mathbf{Q}'' \cap \mathbf{G}$. In particular, $\lambda_n \mathbf{H} \lambda_n^{-1}$ is contained in $\mathbf{P}_{\mathbf{c}_t}$ for all n. Also, if $g_n h_n \lambda_n^{-1} = k''_n a'''_n p'''_n$ is the $(\mathbf{P}_{\mathbf{c}_t}, \mathbf{K})$ coordinate, then (p''_n) is bounded and $\alpha(a'''_n) \to 0$ for all $\alpha \in \Delta(\mathbf{A}_{\mathbf{P}_{\mathbf{c}_t}}, \mathbf{K}, \mathbf{P}_{\mathbf{c}_t})$.

Finally, by assumption, there exist certain \mathbf{SL}_N -representation and a \mathbb{Q} vector \mathbf{v}'' fixed by $\lambda_n \mathbf{H} \lambda_n^{-1}$ and the line spanned by \mathbf{v}'' is preserved by \mathbf{Q}'' with a non-trivial character, which must be nontrivial on \mathbf{c}_t since $\mathbf{Q}'' = \mathbf{Q}_{\mathbf{c}_t}$. In particular, \mathbf{v}'' is preserved by $\mathbf{P}_{\mathbf{c}_t}$ with a non-trivial character. So we are done.

B.3. The case of SL_N . Fix an observable \mathbb{Q} -subgroup \mathbf{H} of \mathbf{SL}_N and a sequence $(g_n) \subset \mathbf{SL}_N(\mathbb{R})$. We prove Theorem B.1 in this case using the idea of canonical polygons (see [Cas04, Gra84]). For simplicity we abbreviate $\mathbf{A}_{\mathbf{P},\mathbf{K}'}$ as $\mathbf{A}_{\mathbf{P}}$ below and continue call $\mathbf{K}' := \mathbf{SO}_N(\mathbb{R})$.

B.3.1. Lattice flags. A subgroup Δ of \mathbb{Z}^N is said to be primitive iff $(\Delta \otimes \mathbb{Q}) \cap \mathbb{Z}^N = \Delta$. Let $\operatorname{Prim}(\mathbb{Z}^N)$ be the collection of all primitive subgroups of \mathbb{Z}^N . For a \mathbb{Q} -character α of \mathbf{H} , let $\operatorname{Prim}_{\alpha}^{\mathbf{H}}(\mathbb{Z}^N)$ denote those $\Delta = \mathbb{Z}\mathbf{v}_1 \oplus \ldots \oplus \mathbb{Z}\mathbf{v}_k$ with

$$h.\mathbf{v}_1 \wedge ... \wedge \mathbf{v}_k = \alpha(h)\mathbf{v}_1 \wedge ... \wedge \mathbf{v}_k, \quad \forall h \in \mathbf{H}.$$

Let $\operatorname{Prim}^{\mathbf{H}}(\mathbb{Z}^N)$ be the union of these $\operatorname{Prim}^{\mathbf{H}}_{\alpha}(\mathbb{Z}^N)$'s.

A totally ordered subset of $\operatorname{Prim}(\mathbb{Z}^N)$ (without loss of generality, assumed to contain $\{0\}$ and \mathbb{Z}^N) is referred to as a **lattice flag**, written as

$$\mathcal{F} = \left\{ \{\mathbf{0}\} = \Delta_0 \subsetneq \Delta_1 \subsetneq ... \subsetneq \Delta_l = \mathbb{Z}^N \right\}$$

The number l-1 is called the rank of \mathcal{F} . For i = 1, ..., l, we define

$$\overline{\Delta_i} := \left(\Delta_i / \Delta_{i-1}\right)' := \left(\frac{\|\Delta_i\|}{\|\Delta_{i-1}\|}\right)^{-\frac{1}{\operatorname{rank}(\Delta_i) - \operatorname{rank}(\Delta_{i-1})}} \cdot \Delta_i / \Delta_{i-1},\tag{42}$$

which has covolume one in the quotient Euclidean space $\Delta_i \otimes \mathbb{R}/\Delta_{i-1} \otimes \mathbb{R}$. We also define a parabolic \mathbb{Q} -subgroup by

$$\mathbf{P}_{\mathcal{F}} := \{ g \in \mathbf{SL}_N(\mathbb{C}) \mid g \text{ preserves } \Delta_i \otimes \mathbb{C}, \ \forall i \}.$$

Given a subset \mathcal{I} of $\{0, 1, ..., N\}$ containing $\{0, N\}$, ordered as $\mathcal{I} = \{0 = i_0 < i_1 < ... < i_l = N\}$, let $\mathcal{F}_{\mathcal{I}}$ be the **standard flag** associated to \mathcal{I} defined by

$$\mathcal{F}_{\mathcal{I}} := \left\{ \{\mathbf{0}\} \subsetneq \oplus_{i=1}^{i_1} \mathbb{Z} \mathbf{e}_i \subsetneq ... \subsetneq \oplus_{i=1}^{i_{l-1}} \mathbb{Z} \mathbf{e}_i \subsetneq \mathbb{Z}^N \right\}.$$

For simplicity we write $\mathbf{P}_{\mathcal{I}} := \mathbf{P}_{\mathcal{F}_{\mathcal{I}}}$.

B.3.2. Quotient lattices. Fix some $\mathcal{I} = \{0 = i_0 < i_1 < ... < i_l = N\}$ for this subsection. Define, for k = 1, ..., l,

$$\Lambda_k^{\text{Std}} := \bigoplus_{i=1}^{\iota_k} \mathbb{Z}.\mathbf{e}_i, \quad V_k := \Lambda_k^{\text{Std}} \otimes_{\mathbb{Z}} \mathbb{R}, \quad \overline{V}_k := V_k/V_{k-1}, \quad \overline{V}_k(\mathbb{Z}) := \Lambda_k^{\text{Std}}/\Lambda_{k-1}^{\text{Std}}$$

and $\pi_k : \mathcal{P}_{\mathcal{I}} := \mathbf{P}_{\mathcal{I}}(\mathbb{R})^\circ \to \mathbf{SL}(\overline{V}_k)$ by

$$p \mapsto c_p \cdot (p|_{V_k} \pmod{V_{k-1}})$$

where c_p is the unique positive real number such that the right hand side has determinant one. π_k descends to a continuous map

$$\overline{\pi}_k: \mathrm{P}_{\mathcal{I}}/\mathrm{P}_{\mathcal{I}} \cap \mathbf{SL}_N(\mathbb{Z}) \to \mathbf{SL}(\overline{V}_k)/\mathbf{SL}(\overline{V}_k(\mathbb{Z})).$$

that is equivariant with respect to π_k . The natural map

$$p \mapsto \left\{ \{ \mathbf{0} \} = p \Lambda_0^{\text{Std}} \subsetneq p \Lambda_1^{\text{Std}} \subsetneq \dots \subsetneq p \Lambda_l^{\text{Std}} = \mathbb{Z}^N \right\}$$

induces a bijection between

$$P_{\mathcal{I}}/P_{\mathcal{I}} \cap \mathbf{SL}_N(\mathbb{Z}) \cong LF_{\mathcal{I}} := \{(\Delta_{\star}) \text{ is a lattice flag } \mid (\Delta_i)_{\mathbb{R}} = V_i, \forall i = 1, ..., l\}.$$

If we additionally identity

$$\mathbf{SL}(\overline{V}_k) / \mathbf{SL}(\overline{V}_k(\mathbb{Z})) \cong \left\{ \text{ unimodular lattices in } \overline{V}_k \right\}$$

by $g \mapsto g \overline{V}_k(\mathbb{Z})$, then $\overline{\pi}_k$ is nothing but

$$(\Delta_{\star}) \mapsto \overline{\Delta}_k$$

(

defined in the last subsection (see Equa.(42)).

Let $\overline{p}^{\text{spl}} : P_{\mathcal{I}}/P_{\mathcal{I}} \cap \mathbf{SL}_N(\mathbb{Z}) \to S_{\mathbf{P}_{\mathcal{I}}}$. It is direct to check the following

Lemma B.2. For each \mathcal{I} , the map $P_{\mathcal{I}}/P_{\mathcal{I}} \cap \mathbf{SL}_N(\mathbb{Z}) \to S_{\mathbf{P}_{\mathcal{I}}} \times \prod_{k=1}^l \mathbf{SL}(V_k) / \mathbf{SL}(V_k(\mathbb{Z}))$ defined by $\Phi_{\mathcal{I}} := \overline{p}^{\text{spl}} \times \prod_k \overline{\pi}_k$ is a proper continuous map.

For k = 1, ..., l, define $j_k := i_k - i_{k-1}$ and $f_k, \varphi_k, \alpha_k : \mathbf{A}_{\mathbf{P}_{\mathcal{I}}} \to \mathbb{R}^+$ by

$$f_k(a) := \left\| a \Lambda_k^{\text{Std}} \right\|, \quad \varphi_k(a) := \left(\frac{\left\| a \Lambda_k^{\text{Std}} \right\|}{\left\| a \Lambda_{k-1}^{\text{Std}} \right\|} \right)^{\frac{1}{j_k}}, \quad \alpha_k(a) := \varphi_k(a) / \varphi_{k+1}(a).$$

One can check that

$$\Delta(\mathbf{A}_{\mathbf{P}_{\mathcal{I}}}, \mathbf{P}_{\mathcal{I}}) = \{\alpha_k \mid k = 1, ..., l - 1\}.$$
(43)

B.3.3. Canonical flags. In this subsection we fix an element $g \in \mathbf{SL}_N(\mathbb{R})$. Define

$$\operatorname{Plot}^{\mathbf{H}}(g) := \left\{ (\operatorname{rank}(\Delta), \log(\|g\Delta\|)) \mid \Delta \in \operatorname{Prim}^{\mathbf{H}}(\mathbb{Z}^N) \right\}$$
$$\operatorname{Poly}^{\mathbf{H}}(g) := \operatorname{closure of the convex hull of } \operatorname{Plot}^{\mathbf{H}}(g).$$

 $\operatorname{Poly}^{\mathbf{H}}(q)$ is a polygon of finitely many sides. Let

$$\mathcal{E}^{\mathbf{H}}(g) := \left\{ \Delta \in \operatorname{Prim}^{\mathbf{H}}(\mathbb{Z}^N) \mid (\operatorname{rank} \Delta, \log(\|g\Delta\|)) \text{ is an extreme point of } \operatorname{Poly}^{\mathbf{H}}(g) \right\}.$$

Lemma B.3. For every $g \in \mathbf{SL}_N(\mathbb{R})$, $\mathcal{E}^{\mathbf{H}}(g)$ is a lattice flag containing $\{\mathbf{0}\}$ and \mathbb{Z}^N .

A proof can be found in [Cas04, Section 4]. Let $r_q := \operatorname{rank}(\mathcal{E}^{\mathbf{H}}(q))$ and write

$$\mathcal{E}^{\mathbf{H}}(g) = \left\{ \{\mathbf{0}\} = \Delta_0(g) \subsetneq \Delta_1(g) \subsetneq \dots \subsetneq \Delta_{r_g+1}(g) = \mathbb{Z}^N \right\},\$$

called the **canonical flag**. Define

$$\begin{aligned} \text{type}(g) &:= \left\{ 0 = \text{rank}(\Delta_0(g)) < \text{rank}(\Delta_1(g)) < \dots < \text{rank}(\Delta_{r_g+1}(g)) = N \right\};\\ \text{slope}_k(g) &:= \frac{\log \|(g\Delta_k(g)\|) - \log(\|g\Delta_{k-1}(g)\|)}{\text{rank}(\Delta_k(g)) - \text{rank}(\Delta_{k-1}(g))}, \ k = 1, \dots, r_g + 1;\\ d_k(g) &:= \text{slope}_{k+1}(g) - \text{slope}_k(g), \ k = 1, \dots, r_g. \end{aligned}$$

B.3.4. Variation of canonical flags. Fixing some $g \in \mathbf{SL}_N(\mathbb{R})$, we would like to search for a better one among $\mathcal{E}^{\mathbf{H}}(gh)$ as h varies in H by (locally) optimizing certain quantities. This part is an extra key step required to treat general connected observable subgroups instead of just those without non-trivial O-characters.

Find a \mathbb{Q} -split subtorus $\mathbf{S}^{\mathbf{H}}$ such that \mathbf{H} is an almost direct product of $(^{\circ}\mathbf{H})^{\circ}$ and $\mathbf{S}^{\mathbf{H}}$. Thus the Lie algebra $\mathfrak{h} = {}^{\circ}\mathfrak{h} \oplus \mathfrak{s}^{\mathfrak{h}}$. Write $\mathfrak{s}^{\mathfrak{h}}_{\mathbb{R}}$ for the \mathbb{R} -span of $\mathfrak{s}^{\mathfrak{h}}$. For a \mathbb{Q} -character α of \mathbf{H} , we regard $d\alpha$ as a linear functional on $\mathfrak{s}^{\mathfrak{h}}_{\mathbb{R}}$ since it vanishes on $\mathfrak{h}_{\mathbb{R}}$. Let $\Phi^{\mathrm{Std}}_{\mathbf{H}}$ collect all $d\alpha$ such that $\mathrm{Prim}^{\mathbf{H}}_{\alpha}(\mathbb{Z}^N)$ is non-zero.

For $\lambda \in \mathbb{R}$, let CVX_{λ} be the collection of all convex closed polygons \mathcal{P} such that

1. $[0, N] \times \mathbb{R}_{>0} \subset \mathcal{P} \subset \{(x, y), y \ge \lambda, 0 \le x \le N\};$

2. $(x, y) \in \text{Extre}(\mathcal{P}) \implies x \in \{0, 1, ..., N\} \text{ and } \{(0, 0), (N, 0)\} \subset \text{Extre}(\mathcal{P}).$

Equip \cup CVX_{λ} with the Chabauty topology. Note that Poly^H(g) belongs to CVX_{λ} for some λ . Given some $\mathcal{P} \in \mathrm{CVX}_{\lambda}$ for some $\lambda \in \mathbb{R}$, we let

- 1. $y_i(\mathcal{P}) := \inf \{ y \mid (i, y) \in \mathcal{P} \}$ for $i \in \{0, ..., N\}$. So $y_0 = y_N = 0$;
- 2. $s_i(\mathcal{P}) := y_i(\mathcal{P}) y_{i-1}(\mathcal{P}) \text{ for } i \in \{1, ..., N\};$ 3. $d_i(\mathcal{P}) = s_{i+1}(\mathcal{P}) s_i(\mathcal{P}) = y_{i+1}(\mathcal{P}) + y_{i-1}(\mathcal{P}) 2y_i(\mathcal{P}) \text{ for } i \in \{1, ..., N-1\}.$

Thus $d_i(g) = d_{\operatorname{rank}(\Delta_i(g))}(\operatorname{Poly}^{\mathbf{H}}(g))$. Also let $\tau_{\mathcal{P}}$ be a permutation on $\{1, ..., N-1\}$ such that

1. $\{d_i(\mathcal{P}) \mid i = 1, ..., N - 1\} = \{d_{\tau_{\mathcal{P}}(1)}(\mathcal{P}) \ge d_{\tau_{\mathcal{P}}(2)}(\mathcal{P}) \ge \ge d_{\tau_{\mathcal{P}}(N-1)}(\mathcal{P})\};$ 2. if $d_{\tau_{\mathcal{P}}(i)}(\mathcal{P}) = d_{\tau_{\mathcal{P}}(i+1)}(\mathcal{P})$ then $\tau_{\mathcal{P}}(i) < \tau_{\mathcal{P}}(i+1)$.

Define a partial order \prec on $\cup \text{CVX}_{\lambda}$ by $\mathcal{P} \prec \mathcal{P}'$ iff $\mathcal{P} = \mathcal{P}'$ or $y_{\tau_{\mathcal{P}}(m)}(\mathcal{P}) < y_{\tau_{\mathcal{P}'}(m)}(\mathcal{P}')$ with m :=min $\{i \mid y_{\tau_{\mathcal{P}}(i)}(\mathcal{P}) \neq y_{\tau_{\mathcal{P}'}(i)}(\mathcal{P}')\}$. The following explains why it is possible to find a local maxima.

Lemma B.4. Let $g \in \mathbf{SL}_N(\mathbb{R}), \lambda \in \mathbb{R}$ and $\mathcal{P} \in \mathrm{CVX}_{\lambda}$. If there exists a sequence $(h_n) \subset \mathrm{H}$ such that

$$\mathcal{P} = \lim_{n \to \infty} \operatorname{Poly}^{\mathbf{H}}(gh_n),$$

then there exists $h \in H$ such that $\mathcal{P} = \operatorname{Poly}^{\mathbf{H}}(gh)$.

Proof. Let $V_{\mathfrak{s}} := \bigcap_{\mathrm{d}\alpha \in \Phi_{\mathfrak{s}}^{\mathrm{Std}}} \ker(\mathrm{d}\alpha)$ and choose some complementary \mathbb{R} -linear subspace $W_{\mathfrak{s}}$ such that $\mathfrak{s}^{\mathfrak{h}}_{\mathbb{R}} = V_{\mathfrak{s}} \oplus W_{\mathfrak{s}}$. Without loss of generality, assume the existence of $(w_n) \subset W_{\mathfrak{s}}$ such that

$$\mathcal{P} = \lim_{n \to \infty} \operatorname{Poly}^{\mathbf{H}}(g \exp(w_n)).$$

So we can find $\lambda' \in \mathbb{R}$ such that $\operatorname{Poly}^{\mathbf{H}}(g \exp(w_n)) \in \operatorname{CVX}_{\lambda'}$ for all *n*. In particular, there exists $C_1 \in \mathbb{R}$ such that $d\alpha(w_n) > C_1$ for all n and $d\alpha \in \Phi_{\mathbf{H}}^{\text{Std}}$.

Passing to a subsequence, assume that for every $d\alpha \in \Phi_{\mathbf{H}}^{\mathrm{Std}}$, either $d\alpha(w_n) \to +\infty$ or remains bounded. Let $\Phi_{\text{bdd}} \subset \Phi_{\mathbf{H}}^{\text{Std}}$ correspond to the bounded ones and Φ_{∞} be its complement. Let $W' := \bigcap_{\mathrm{d}\alpha \in \Phi_{\text{bdd}}} \ker d\alpha \cap W_{\mathfrak{s}}$ and take another subspace W'' such that $W_{\mathfrak{s}} = W' \oplus W''$. Write $w_n = w'_n + w''_n$. Then (w''_n) is contained in some bounded set $B \subset W''$ and $\mathrm{d}\alpha(w'_n) \to +\infty$ for every $\mathrm{d}\alpha \in \Phi_{\infty}$.

Find $u_0 \in W'$ such that for every $v \in B$ one has

$$\inf_{d\alpha \in \Phi_{\infty}} \inf_{\Delta \in \operatorname{Prim}_{\alpha}^{\mathbf{H}}(\mathbb{Z}^{N})} \log\left(\|g \exp(u_{0} + v)\Delta\| \right) > 0.$$
(44)

Define $w_n^{\text{new}} := u_0 + w_n''$. Since $\|g \exp(w_n^{\text{new}})\Delta\| = \|g \exp(w_n)\Delta\|$ for every $\Delta \in \text{Prim}_{\alpha}^{\mathbf{H}}(\mathbb{Z}^N)$ and $d\alpha \in \Phi_{bdd}$, we have (by Equa.(44))

$$\operatorname{Poly}^{\mathbf{H}}(g \exp(w_n)) = \operatorname{Poly}^{\mathbf{H}}(g \exp(w_n^{\operatorname{new}}))$$

Being bounded, we can select a subsequence such that (w_n^{new}) converges to some w_{∞} . Then $\mathcal{P} = \text{Poly}^{\mathbf{H}}(g \exp(w_{\infty}))$.

Thanks to Lemma B.4, for every $g \in \mathbf{SL}_N(\mathbb{R})$, there exists (and we fix such an) $h_g \in \mathbf{H}$ such that a local maxima with respect to \prec is taken at $\operatorname{Poly}^{\mathbf{H}}(gh_g)$ among $\{\operatorname{Poly}^{\mathbf{H}}(gh), h \in \mathbf{H}\}$.

Lemma B.5. Take some $g \in \mathbf{SL}_N(\mathbb{R})$. For $C \in \mathbb{R}$, let $\mathscr{I}_C := \{i = 1, ..., r_g, d_i(gh_g) > C\}$. For $\Delta \in \mathrm{Prim}^{\mathbf{H}}(\mathbb{Z}^N)$, we let α_Δ be the unique $\alpha \in \mathfrak{X}^*_{\mathbb{Q}}(\mathbf{H})$ such that $\Delta \in \mathrm{Prim}^{\mathbf{H}}_{\alpha}(\mathbb{Z}^N)$ and regard its differential $d\alpha_\Delta$ as a linear functional on $\mathfrak{s}^{\mathfrak{h}}_{\mathbb{R}}$. For every $C \in \mathbb{R}$ such that \mathscr{I}_C is non-empty, there exists a set of positive integers $(a_i)_{i \in \mathscr{I}_C}$ such that $\sum_{i \in \mathscr{I}_C} a_i d\alpha_{\Delta_i(gh_g)} = 0$.

Proof. If the conclusion were wrong for some $C \in \mathbb{R}$, then **0** would not belong to the relative interior of the convex hull of $\{d\alpha_{\Delta_i(gh_g)}, i \in \mathscr{I}_C\}$, which is non-empty. Therefore, there exists $w_0 \in \mathfrak{s}^{\mathfrak{h}}_{\mathbb{R}}$ such that $d\alpha_{\Delta_i(gh_g)}(w_0) \geq 0$ for all $i \in \mathscr{I}_C$ and $d\alpha_{\Delta_i(gh_g)}(w_0) > 0$ for some $i \in \mathscr{I}_C$. For $\varepsilon > 0$ small enough, the polygon $\operatorname{Poly}^{\mathbf{H}}(gh_g) \prec \operatorname{Poly}^{\mathbf{H}}(gh_g \exp(\varepsilon w_0))$ strictly. This is a contradiction against the choice of h_g .

B.3.5. Weights. By passing to a subsequence, assume without loss of generality that for all $n \in \mathbb{Z}^+$, $r_{g_n h_{g_n}} = r$ for some r, type $(g_n h_{g_n}) =$ type $(g_1 h_{g_1})$ and $\alpha_{\Delta_k(g_n h_{g_n})}$ remains constant for each k. So there exist $(\lambda_n) \subset \mathbf{SL}_N(\mathbb{Z})$ and $\mathcal{I} = \{0 = i_0 < i_1 < ... < i_{r+1} = N\} =$ type $(g_n h_{g_n})$ with $\mathcal{E}^{\mathbf{H}}(g_n h_{g_n}) = \lambda_n . \mathcal{F}_{\mathcal{I}}$ for every $n \in \mathbb{Z}^+$. Hence

$$\mathcal{F}_{\mathcal{I}} = \lambda_n^{-1} \mathcal{E}^{\mathbf{H}}(g_n h_{g_n}) = \mathcal{E}^{\lambda_n^{-1} \mathbf{H} \lambda_n}(g_n h_{g_n} \lambda_n).$$

Also note that $\lambda_n^{-1} \mathbf{H} \lambda_n \subset \mathbf{P}_{\mathcal{I}}$ for all n.

Now that \mathcal{I} is fixed, it is safe to adopt the notation from Section B.3.2 with l = r + 1. Using the adjoint action, define $\operatorname{Prim}^{\mathbf{H}}(\mathfrak{sl}_N(\mathbb{Z}))$ to be the primitive subgroups of $\mathfrak{sl}_N(\mathbb{Z})$ whose \mathbb{C} -spans are Ad(**H**)-invariant. For $\alpha \in \mathfrak{X}^*_{\mathbb{Q}}(\mathbf{H})$, define $\operatorname{Prim}^{\mathbf{H}}_{\alpha}(\mathfrak{sl}_N(\mathbb{Z}))$ similarly to $\operatorname{Prim}^{\mathbf{H}}_{\alpha}(\mathbb{Z}^N)$ and let $\mathcal{V}^{\operatorname{Ad}}_{\alpha,n}(\mathbb{Z}) := \lambda_n^{-1}\operatorname{Prim}^{\mathbf{H}}_{\alpha}(\mathfrak{sl}_N(\mathbb{Z}))$. To make notations uniform, define

$$\begin{aligned} \mathcal{V}_{\alpha,n}^{\mathrm{Std}}(\mathbb{Z}) &:= \lambda_n^{-1} \operatorname{Prim}_{\alpha}^{\mathbf{H}}(\mathbb{Z}^N), \quad \mathcal{V}_{\alpha,n}(\mathbb{Z}) := \mathcal{V}_{\alpha,n}^{\mathrm{Ad}}(\mathbb{Z}) \bigsqcup \mathcal{V}_{\alpha,n}^{\mathrm{Std}}(\mathbb{Z}), \quad \mathcal{V}_{\alpha}(\mathbb{Z}) &:= \lambda_n \cdot \mathcal{V}_{\alpha,n}(\mathbb{Z}); \\ \Phi_{\mathbf{H}}^{\mathrm{Ad}} &:= \left\{ \mathrm{d}\alpha : \mathfrak{s}_{\mathbb{R}}^{\mathfrak{h}} \to \mathbb{R} \mid \mathcal{V}_{\alpha,n}^{\mathrm{Ad}}(\mathbb{Z}) \neq \{\{\mathbf{0}\}\} \right\}, \quad \Phi_{\mathbf{H}} := \Phi_{\mathbf{H}}^{\mathrm{Std}} \bigcup \Phi_{\mathbf{H}}^{\mathrm{Ad}}. \end{aligned}$$

B.3.6. Nondivergence. Using $(\mathbf{P}_{\mathcal{I}}, \mathbf{K}')$ coordinates, write $g_n h_{g_n} \lambda_n = k_n a_n p_n$.

Lemma B.6. For every n and $\alpha \in \Delta(A_{\mathbf{P}_{\mathcal{I}}}, \mathbf{P}_{\mathcal{I}})$, one has $\alpha(a_n) \leq 1$. Moreover, for every non-empty bounded open subset \mathcal{O}_{H} , there exists a compact subset \mathscr{C} of $P_{\mathcal{I}}/P_{\mathcal{I}} \cap \mathbf{SL}_N(\mathbb{Z})$ such that $p_n \lambda_n^{-1}[\mathcal{O}_{\mathrm{H}}]\lambda_n \cap \mathscr{C} \neq \emptyset$ for all n.

Here $[\mathcal{O}_{\mathrm{H}}]$ denotes the image of \mathcal{O}_{H} in $\mathrm{P}_{\mathcal{E}^{\mathbf{H}}(g_{n}h_{g_{n}})}/\mathrm{P}_{\mathcal{E}^{\mathbf{H}}(g_{n}h_{g_{n}})} \cap \mathbf{SL}_{N}(\mathbb{Z})$ and $\lambda_{n}^{-1}[\mathcal{O}_{\mathrm{H}}]\lambda_{n}$ denotes the further image under the map from $\mathrm{P}_{\mathcal{E}^{\mathbf{H}}(g_{n}h_{g_{n}})}/\mathrm{P}_{\mathcal{E}^{\mathbf{H}}(g_{n}h_{g_{n}})} \cap \mathbf{SL}_{N}(\mathbb{Z})$ to $\mathrm{P}_{\mathcal{I}}/\mathrm{P}_{\mathcal{I}} \cap \mathbf{SL}_{N}(\mathbb{Z})$ induced by $x \mapsto \lambda_{n}^{-1}x\lambda_{n}$.

Proof of the first part of Lemma B.6. For $\alpha \in \Delta(A_{\mathbf{P}_{\mathcal{I}}}, \mathbf{P}_{\mathcal{I}})$, find k = 1, ..., r + 1 such that $\alpha(a_n) = \alpha_k(a_n)$ by Equa.(43). Then

$$\begin{aligned} \alpha(a_n) &= \varphi_k(a_n)\varphi_{k+1}(a_n)^{-1} = \left(\frac{\|a_n\Lambda_{k+1}^{\text{Std}}\|}{\|a_n\Lambda_k^{\text{Std}}\|}\right)^{-j_{k+1}^{-1}} \cdot \left(\frac{\|a_n\Lambda_k^{\text{Std}}\|}{\|a_n\Lambda_{k-1}^{\text{Std}}\|}\right)^{j_k^{-1}} \\ &= \left(\frac{\|g_nh_{g_n}\lambda_n\Lambda_{k+1}^{\text{Std}}\|}{\|g_nh_{g_n}\lambda_n\Lambda_k^{\text{Std}}\|}\right)^{-j_{k+1}^{-1}} \cdot \left(\frac{\|g_nh_{g_n}\lambda_n\Lambda_{k-1}^{\text{Std}}\|}{\|g_nh_{g_n}\lambda_n\Lambda_{k-1}^{\text{Std}}\|}\right)^{j_k^{-1}} \\ &= \exp\left(-\frac{\ln\|g_nh_{g_n}\Delta_{k+1}(g_nh_{g_n})\| - \ln\|g_nh_{g_n}\Delta_k(g_nh_{g_n})\|}{j_{k+1}} + \frac{\ln\|g_nh_{g_n}\Delta_k(g_nh_{g_n})\| - \ln\|g_nh_{g_n}\Delta_{k-1}(g_nh_{g_n})\|}{j_k}\right) \\ &= \exp(-d_k(g_nh_{g_n})) \le 1. \end{aligned}$$

The last inequality follows from the convexity of the polygons.

To prove the second part, recall the following nondivergence criterion proved in [Zha22, Theorem 4.1] based on Kleinbock-Margulis [KM98] and some geometry of numbers.

Theorem B.7. Fix some non-empty bounded open subset \mathcal{O}_H of H and a smooth probability measure $\mathfrak{m}_{\mathcal{O}}$ on $[\mathcal{O}_H]$. Also fix $k \in \{1, ..., r\}$ and some $\varepsilon \in (0, 1)$. If there exists $\delta > 0$ such that for every $n \in \mathbb{Z}^+$, one has

$$\|\pi_k(p_n)\Delta\| \ge \delta, \quad \forall \Delta \in \operatorname{Prim}^{\pi_k(\lambda_n^{-1}\mathbf{H}\lambda_n)}(\overline{V}_k(\mathbb{Z})),$$

then there exists a compact subset $\mathscr{C}_k \subset \mathbf{SL}(\overline{V}_k)/\mathbf{SL}(\overline{V}_k(\mathbb{Z}))$ such that

$$\mathbf{m}_{\mathcal{O}}\left\{x \in [\mathcal{O}_{\mathrm{H}}] \mid \overline{\pi}_{k}(p_{n}\lambda_{n}x\lambda_{n}^{-1}) \notin \mathscr{C}_{k}\right\} < \varepsilon.$$

Proof of the second part of Lemma B.6. By rigidity of diagonalizable groups (see [Spr98, 3.2.8]),

$$\left\{\overline{p}^{\mathrm{spl}}(p_n\lambda_n^{-1}[\mathcal{O}_{\mathrm{H}}]\lambda_n) = \overline{p}^{\mathrm{spl}}(\lambda_n^{-1}[\mathcal{O}_{\mathrm{H}}]\lambda_n) \mid n \in \mathbb{Z}^+\right\}$$

consists of finitely many bounded subsets of $S_{\mathbf{P}_{\mathcal{I}}}$ and thus remains bounded. In light of Lemma B.2 and Theorem B.7, it suffices to show that

$$\|\pi_k(p_n)\Delta\| \ge \delta, \quad \forall \Delta \in \operatorname{Prim}^{\pi_k(\lambda_n^{-1}\mathbf{H}\lambda_n)}(\overline{V}_k(\mathbb{Z})),$$

for some $\delta > 0$ independent of k, n. Fix such a Δ , let $\widetilde{\Delta} \in \operatorname{Prim}^{\lambda_n^{-1}\mathbf{H}\lambda_n}(\mathbb{Z}^N)$ be the unique element such that

$$\Lambda_{k-1}^{\mathrm{Std}} \subset \widetilde{\Delta} \subset \Lambda_k^{\mathrm{Std}}, \quad \Delta = \widetilde{\Delta} / \Lambda_{k-1}^{\mathrm{Std}}.$$

By assumption we have

$$\left(\operatorname{rank}(\widetilde{\Delta}), \log \left\| g_n h_{g_n} \lambda_n \widetilde{\Delta} \right\| \right) \in \operatorname{Plot}^{\lambda_n^{-1} \mathbf{H} \lambda_n}(g_n h_{g_n} \lambda_n) = \operatorname{Plot}^{\mathbf{H}}(g_n h_{g_n})$$

Therefore,

$$\begin{aligned} \frac{\log \|\pi_k(p_n)\Delta\|}{\operatorname{rank}(\widetilde{\Delta}) - i_{k-1}} &= \frac{\log \left\| p_n \widetilde{\Delta} \right\| - \log \|p_n \Lambda_{k-1}^{\operatorname{Std}} \|}{\operatorname{rank}(\widetilde{\Delta}) - i_{k-1}} \\ &= \frac{\log \left\| k_n a_n p_n \widetilde{\Delta} \right\| - \log \|k_n a_n p_n \Lambda_{k-1}^{\operatorname{Std}} \|}{\operatorname{rank}(\widetilde{\Delta}) - i_{k-1}} - \log(\varphi_k(a_n)) \\ &= \frac{\log \left\| k_n a_n p_n \widetilde{\Delta} \right\| - \log \|g_n h_{g_n} \Delta_{k-1}(g_n h_{g_n}) \|}{\operatorname{rank}(\widetilde{\Delta}) - i_{k-1}} - \log(\varphi_k(a_n)) \\ &\geq \frac{\log \|g_n h_{g_n} \Delta_k(g_n h_{g_n}) \| - \log \|g_n h_{g_n} \Delta_{k-1}(g_n h_{g_n}) \|}{j_k} - \log(\varphi_k(a_n)) \\ &= \frac{\log \|p_n \Lambda_k^{\operatorname{Std}} \| - \log \|p_n \Lambda_{k-1}^{\operatorname{Std}} \|}{j_k} = 0. \end{aligned}$$

That is to say, $\|\pi_k(p_n)\Delta\| \ge 1$ and we are done.

B.3.7. Deep in the polytopes. Let $\mathcal{V}^*_{\alpha,n}(\mathbb{Z}) := \mathcal{V}_{\alpha,n}(\mathbb{Z}) \setminus \{\{\mathbf{0}\}\}$ and

$$M_{\alpha,n} := \inf_{\Delta \in \mathcal{V}_{\alpha,n}^*(\mathbb{Z})} \log \|p_n \Delta\|, \quad \mathrm{d}\alpha \in \Phi_{\mathbf{H}}.$$

By Lemma B.6, $M_{\alpha,n} > C_2$ for some $C_2 \in \mathbb{R}$ and for all $d\alpha \in \Phi_{\mathbf{H}}$ and n. Passing to a subsequence, assume that for each $d\alpha \in \Phi_{\mathbf{H}}$, either $(M_{\alpha,n})$ remains bounded or diverges to $+\infty$. Let $\Phi(\text{bdd})$ correspond to those bounded ones and Φ_{∞} collect the rest. Let $M_n(\infty) := \inf_{d\alpha \in \Phi_{\infty}} M_{\alpha,n}$, which diverges to $+\infty$. Passing to a further subsequence, assume $M_n(\infty)$ is positive for each n. Let

$$\Phi_0 := \{ d\alpha \in \Phi(bdd) \mid -d\alpha \in Cone(\Phi(bdd)) \}, \ \Phi_1 := \Phi(bdd) \setminus \Phi_0$$

where $\operatorname{Cone}(-)$ refers to the $\mathbb{R}_{\geq 0}$ -linear span of a subset. Note that $\operatorname{Cone}(\Phi_0)$ is a \mathbb{R} -linear subspace and the convex hull of Φ_1 is compact and disjoint from $\operatorname{Cone}(\Phi_0)$. So there exists $w_0 \in \mathfrak{s}^{\mathfrak{h}}_{\mathbb{R}}$ such that

$$\mathrm{d}\alpha(w_0) \begin{cases} = 0 & \forall \,\mathrm{d}\alpha \in \Phi_0 \\ > 0 & \forall \,\mathrm{d}\alpha \in \Phi_1 \end{cases}.$$

Also, fix $C_3 > 0$ such that

$$|\mathrm{d}\alpha(w_0)| < C_3, \quad \forall \,\mathrm{d}\alpha \in \Phi_\mathbf{H}.$$

Define a sequence of positive numbers (κ_n) by

$$\kappa_n := \frac{M_n(\infty)}{2C_3}.$$

Using $(\mathbf{P}_{\mathcal{I}}, \mathbf{K}')$ coordinates, write

$$g_n h'_{g_n} \lambda_n = k'_n a'_n p'_n$$
, where $h'_{g_n} := h_{g_n} \exp(\kappa_n w_0)$.

 Then^5

$$k_n = k'_n, \quad a_n p_n \cdot (\lambda_n^{-1} \exp(\kappa_n w_0) \lambda_n) = a'_n p'_n.$$

Moreover, by Lemma B.5, the character associated to $\Delta_k(g_n h_{g_n}) = \lambda_n \Lambda_k^{\text{Std}}$ belongs to Φ_0 for each k = 1, ..., r. We have

$$\lambda_n^{-1} \exp(\kappa_n w_0) \lambda_n \Lambda_k^{\text{Std}} = \Lambda_k^{\text{Std}}.$$
(45)

Hence $\lambda_n^{-1} \exp(\kappa_n w_0) \lambda_n \in {}^{\circ} \mathbf{P}_{\mathcal{I}}$ and

$$a'_n = a_n, \quad p'_n = p_n \cdot (\lambda_n^{-1} \exp(\kappa_n w_0) \lambda_n).$$

For $d\alpha \in \Phi_{\mathbf{H}}$ and $\Delta \in \mathcal{V}_{\alpha,n}(\mathbb{Z})$, we have

$$\log \|p'_n \Delta\| = \kappa_n \mathrm{d}\alpha(w_0) + \log \|p_n \Delta\|.$$
(46)

We have already seen by Equa.(45) that:

$$\log \left\| g_n h'_{g_n} \lambda_n \Lambda_k^{\text{Std}} \right\| = \log \left\| g_n h_{g_n} \lambda_n \Lambda_k^{\text{Std}} \right\|$$

Next we claim that

Claim. Let $\Delta \in \lambda_n^{-1} \operatorname{Prim}^{\mathbf{H}}(\mathbb{Z}^N)$. The point $(\operatorname{rank}(\Delta), \log \|g_n h'_{g_n} \lambda_n \Delta\|)$ lies above the piecewise-linear line connecting $(\operatorname{rank}(\Lambda_k^{\operatorname{Std}}), \log \|g_n h'_{g_n} \lambda_n \Lambda_k^{\operatorname{Std}}\|)$ as k goes from 0 to r + 1.

Proof of the claim. First we assume that Δ is compatible with $\mathcal{F}_{\mathcal{I}}$, namely $\Lambda_{k-1}^{\text{Std}} \subset \Delta \subset \Lambda_k^{\text{Std}}$ for some k, and prove

$$\frac{\log \left\|g_n h'_{g_n} \lambda_n \Delta\right\| - \log \left\|g_n h'_{g_n} \lambda_n \Lambda_{k-1}^{\mathrm{Std}}\right\|}{\mathrm{rank}(\Delta) - i_{k-1}} \ge \frac{\log \left\|g_n h'_{g_n} \lambda_n \Lambda_k^{\mathrm{Std}}\right\| - \log \left\|g_n h'_{g_n} \lambda_n \Lambda_{k-1}^{\mathrm{Std}}\right\|}{i_k - i_{k-1}}.$$
 (47)

Since Δ is assumed to be compatible with $\mathcal{F}_{\mathcal{I}}$, this is equivalent to

$$\frac{\log \|p_n'\Delta\| - \log \|p_n'\Lambda_{k-1}^{\operatorname{Std}}\|}{\operatorname{rank}(\Delta) - i_{k-1}} \ge \frac{\log \|p_n'\Lambda_k^{\operatorname{Std}}\| - \log \|p_n'\Lambda_{k-1}^{\operatorname{Std}}\|}{i_k - i_{k-1}},$$

which, thanks to Equa.(46), is again equivalent to

$$\kappa_n \mathrm{d}\alpha(w_0) + \log \|p_n \Delta\| \ge 0. \tag{48}$$

For $d\alpha \in \Phi(bdd)$,

 $\kappa_n \mathrm{d}\alpha(w_0) + \log \|p_n \Delta\| \ge \log \|p_n \Delta\| \ge 0.$

For $d\alpha \in \Phi_{\infty}$,

$$\kappa_n \mathrm{d}\alpha(w_0) + \log \|p_n \Delta\| \ge \log \|p_n \Delta\| - C_3 \kappa_n \ge 0.$$

The truth of Equa.(48) and hence of Equa.(47) is verified.

⁵Strictly speaking $k_n = k'_n$ is only true modulo $K' \cap M_{\mathbf{P},K'}$. As this does not affect the proof, we choose to ignore it.

Now we drop the compatibility assumption on Δ . Fix some m < l such that Δ is contained in Λ_l^{Std} but not $\Lambda_{l-1}^{\text{Std}}$; Δ contains Λ_m^{Std} but not $\Lambda_{m+1}^{\text{Std}}$. Let $\Delta_j := \Delta \cap \Lambda_{m+j}$ as j ranges over $\{0, 1, \dots, l-m\}$ and $i'_j := \operatorname{rank}(\Delta_j)$. So $\Delta_0 = \Lambda_m$ and $\Delta_{l-m} = \Delta$. We are going to explain that for $j = 0, 1, \dots, l-m-1$,

$$\frac{\log \|g_n h'_{g_n} \lambda_n \Delta_{j+1}\| - \log \|g_n h'_{g_n} \lambda_n \Delta_j\|}{i'_{j+1} - i'_j} \ge \frac{\log \|g_n h'_{g_n} \lambda_n \Lambda_{m+j+1}^{\text{Std}}\| - \log \|g_n h'_{g_n} \lambda_n \Lambda_{m+j}^{\text{Std}}\|}{i_{m+j+1} - i_{m+j}}, \quad (49)$$

which is sufficient to conclude the proof since $i'_{j+1} - i'_j \leq i_{m+j+1} - i_{m+j}$. As the argument has nothing to do with the explicit form of $g_n h'_{g_n} \lambda_n$, we abbreviate $\|g_n h'_{g_n} \lambda_n . \Lambda\| =: \|\Lambda\|_g$ for every discrete subgroup Λ . For two primitive subgroups Λ_1, Λ_2 of \mathbb{Z}^N , let

$$\Lambda_1 + \Lambda_2 := (\Lambda_1 + \Lambda_2) \otimes \mathbb{Q} \cap \mathbb{Z}^N$$

which is again primitive. We have that (see [Cas04, Corollary 4.2])

$$\frac{\left\|\Delta_{j+1}\right\|_{g}}{\left\|\Delta_{j}\right\|_{g}} \geq \frac{\left\|\Delta+'\Lambda_{m+j}^{\mathrm{Std}}\right\|_{g}}{\left\|\Lambda_{m+j}^{\mathrm{Std}}\right\|_{g}}$$
$$\implies \frac{\log\left\|\Delta_{j+1}\right\|_{g} - \log\left\|\Delta_{j}\right\|_{g}}{i'_{j+1} - i'_{j}} \geq \frac{\log\left\|\Delta+'\Lambda_{m+j}^{\mathrm{Std}}\right\|_{g} - \log\left\|\Lambda_{m+j}^{\mathrm{Std}}\right\|_{g}}{\operatorname{rank}(\Delta+'\Lambda_{m+j}^{\mathrm{Std}}) - i_{m+j}}$$

Equa.(49) then follows by applying Equa.(47) to $\Delta + \Lambda^{\text{Std}}_{m+j}$, which is indeed compatible with $\mathcal{F}_{\mathcal{I}}$.

By the claim and Equa.(45),

$$\operatorname{Poly}^{\lambda_n^{-1}\mathbf{H}\lambda_n}(g_nh_{g_n}\lambda_n) = \operatorname{Poly}^{\lambda_n^{-1}\mathbf{H}\lambda_n}(g_nh'_{g_n}\lambda_n)$$
$$\mathcal{E}^{\lambda_n^{-1}\mathbf{H}\lambda_n}(g_nh_{g_n}\lambda_n) = \mathcal{E}^{\lambda_n^{-1}\mathbf{H}\lambda_n}(g_nh'_{g_n}\lambda_n) = \left\{\{0\} \subsetneq \Lambda_1^{\operatorname{Std}} \subsetneq ... \subsetneq \Lambda_{r+1}^{\operatorname{Std}} = \mathbb{Z}^N\right\}.$$

Also for $d\alpha \in \Phi_1$ and $\Delta \in \mathcal{V}_{\alpha,n}(\mathbb{Z})$,

$$\log \|p'_n \Delta\| \ge \kappa_n \mathrm{d}\alpha(w_0) + C_2$$

diverges to $+\infty$. Similarly, $\log \|p'_n \Delta\|$ with $\Delta \in \mathcal{V}_{\alpha,n}(\mathbb{Z})$ has a common lower bound for other $d\alpha \in \Phi_{\mathbf{H}}$. Repeating the proof of the second part of Lemma B.6, we have that

 $(p'_n \lambda_n^{-1}[\mathcal{O}_{\mathrm{H}}]\lambda_n)$ is nondivergent in $\mathrm{P}_{\mathcal{I}}/\mathrm{P}_{\mathcal{I}} \cap \mathbf{SL}_N(\mathbb{Z})$.

As a consequence, we find some $(\lambda'_n) \subset P_{\mathcal{I}} \cap \mathbf{SL}_N(\mathbb{Z})$, bounded sequence (b_n) in $P_{\mathcal{I}}$ and another bounded sequence (ω_n) in \mathcal{O}_H such that

$$p_n'\lambda_n^{-1}\omega_n\lambda_n = b_n\lambda_n'.$$

Letting $h''_{g_n} := h'_{g_n} \omega_n$ and $\gamma_n := \lambda'_n \lambda_n^{-1}$, we have

$$g_n h''_{g_n} \gamma_n^{-1} = k_n a_n b_n, \quad \gamma_n \mathbf{H} \gamma_n^{-1} \subset \mathbf{P}_{\mathcal{I}} \text{ fixes } \mathbf{v}^{\mathcal{I}}.$$

Rewrite $k_n a_n b_n = k_n a''_n b''_n$ using $(\mathbf{P}_{\mathcal{I}}, \mathbf{K}')$ coordinates, then (a''_n) (resp. (b''_n)) is bounded away from (a_n) (resp. (b_n)). In particular, (b''_n) is bounded and there exists $C_4 > 0$ such that $\alpha(a''_n) < C_4$ for all $\alpha \in \Delta(\mathbf{A}_{\mathbf{P}_{\mathcal{I}}}, \mathbf{P}_{\mathcal{I}})$. For $d\alpha \in \Phi_{\mathbf{H}}$, let $M'_{\alpha,n} := \inf \{ \log \|p'_n \Delta\| \mid \Delta \in \mathcal{V}^*_{\alpha,n}(\mathbb{Z}) \}$. By what we know about $(M_{\alpha,n})$ and Equa.(46), $(M'_{\alpha,n})$ diverges to $+\infty$ for $\alpha \in \Phi_{\infty} \cup \Phi_1$ and $(M'_{\alpha,n} = M_{\alpha,n})$ remains bounded for $\alpha \in \Phi_0$. In particular, if $(M'_{\alpha,n})$ is bounded, then $\sum_{i=0}^m a_i d\alpha_i = 0$ for some $(a_i)_{i=0}^m \subset \mathbb{R}^+$ and $(d\alpha_i)_{i=0}^m \subset \Phi_{\mathbf{H}}$ with $\alpha_0 = \alpha$ and $(M'_{\alpha_i,n})$ bounded for each *i*. As α_i 's are all Q-characters, we may take (a_i) 's to be positive integers. Let

$$M_{\alpha,n}'' := \inf \left\{ \log \|\gamma_n \Delta\| \mid \Delta \in \mathcal{V}_{\alpha}^*(\mathbb{Z}) \right\}.$$

Since

 $\log \|\gamma_n \Delta\| = \log \left\| b_n^{-1} p'_n \lambda_n^{-1} \omega_n \Delta \right\| \text{ is bounded from } \log \left\| p'_n \lambda_n^{-1} \Delta \right\|,$

the divergence/boundedness of $(M'_{\alpha,n})$ is the same as that of $(M''_{\alpha,n})$.

Lemma B.8. Assume that $(\gamma_n \mathbf{H} \gamma_n^{-1})$ has a subsequence converging to \mathbf{L} . Then \mathbf{L} is observable in \mathbf{SL}_N .

RUNLIN ZHANG

Proof. It follows from the proof of [Zha21, Lemma 4.10]. Let us briefly recall how. Denote by **O** the observable hull of **L** in \mathbf{SL}_N . It suffices to show that **L** is normalized by **O**. Assume $\Delta_{\mathbf{L}} \in \operatorname{Prim}(\mathfrak{sl}_N(\mathbb{Z}))$ spans the Lie algebra of **L**. By abuse of notation, also view $\Delta_{\mathbf{L}}$ as a vector in the appropriate wedged vector space. Then we only need to show that the line spanned by $\Delta_{\mathbf{L}}$ is preserved by **O**. Passing to a subsequence, assume for some fixed $d\alpha \in \Phi_{\mathbf{H}}, \gamma_n^{-1}\Delta_{\mathbf{L}} \in \mathcal{V}_{\alpha}(\mathbb{Z})$ for all $n \in \mathbb{Z}^+$. But it is a tautology that

$$\left(\log \left\|\gamma_n \cdot (\gamma_n^{-1} \Delta_{\mathbf{L}})\right\|\right)$$
 is bounded.

So we can find $(q_i) \subset \mathbb{Z}^+$, $(d\alpha_i) \in \Phi_{\mathbf{H}}$ and $(\Delta_i) \subset \gamma_n \mathcal{V}_{\alpha_i}(\mathbb{Z})$ with $\Delta_0 = \Delta_{\mathbf{L}}$ and $\sum q_i d\alpha_i = 0$. Therefore,

$$\bigotimes \Delta_i^{\otimes q_i} \text{ is fixed by } \gamma_n \mathbf{H} \gamma_n^{-1} \text{ for every } n$$
$$\implies \bigotimes \Delta_i^{\otimes q_i} \text{ is fixed by } \mathbf{O}$$
$$\implies \text{ the line spanned by } \Delta_0 = \Delta_{\mathbf{L}} \text{ is fixed by } \mathbf{O}.$$

So we are done.

Finally, passing to a subsequence, assume that for each $\alpha \in \Delta(\mathbf{A}_{\mathbf{P}_{\mathcal{I}}}, \mathbf{P}_{\mathcal{I}})$, either $(\alpha(a''_n))$ remains bounded away from 0 and ∞ or converges to 0. Let J collect the bounded ones. For simplicity let $\mathbf{P}_J := (\mathbf{P}_{\mathcal{I}})_J$. If we write

$$g_n h_{g_n}^{\prime\prime} \gamma_n^{-1} = k_n^J a_n^J b_n^J$$

using $(\mathbf{P}_J, \mathbf{K}')$ coordinates, then (b_n^J) is bounded and $\alpha(a_n^J) \to 0$ for all $\alpha \in \Delta(\mathbf{A}_{\mathbf{P}_J}, \mathbf{P}_J)$.

To conclude proof, we note that by Equa.(43),

$$J \cong \{k = 1, ..., l - 1 \mid \alpha_k(a_n) \not\to 0\} = \{k = 1, ..., l - 1 \mid -\log \alpha_k(a_n) \not\to +\infty\}.$$

But

$$\begin{aligned} &-\log \alpha_k(a_n) = \log \varphi_{k+1}(a_n) - \log \varphi_k(a_n) \\ &= \frac{\log \left\| a_n \Lambda_{k+1}^{\text{Std}} \right\| - \log \left\| a_n \Lambda_k^{\text{Std}} \right\|}{i_{k+1} - i_k} - \frac{\log \left\| a_n \Lambda_k^{\text{Std}} \right\| - \log \left\| a_n \Lambda_{k-1}^{\text{Std}} \right\|}{i_k - i_{k-1}} \\ &= \frac{\log \left\| g_n h_{g_n} \lambda_n \Lambda_{k+1}^{\text{Std}} \right\| - \log \left\| g_n h_{g_n} \lambda_n \Lambda_k^{\text{Std}} \right\|}{i_{k+1} - i_k} - \frac{\log \left\| g_n h_{g_n} \lambda_n \Lambda_k^{\text{Std}} \right\| - \log \left\| g_n h_{g_n} \lambda_n \Lambda_{k-1}^{\text{Std}} \right\|}{i_k - i_{k-1}} \\ &= d_k(g_n h_{g_n}). \end{aligned}$$

Choose $C_5 \in \mathbb{R}$ such that

 $d_k(g_n h_{q_n}) \leq C_5, \quad \forall n, \forall k \in J.$

Passing to a subsequence we may assume that $d_k(g_nh_{g_n}) > C_5$ for all n and $k \notin J$. By Lemma B.5 applied to $C = C_5$, we find $(a_k)_{k \in \{1,...,l-1\}\setminus J} \subset \mathbb{Z}^+$ such that $\sum_{k \notin J} a_k d\alpha_{\Delta_k(g_nh_{g_n})} = 0$ (recall we had assumed that $\alpha_{\Delta_k(g_nh_{g_n})}$ is independent of n for each k). Therefore, $\sum_{k \notin J} a_k df_k = 0$. By abuse of notation, treat Λ_k^{Std} as a vector for every k. If $\mathbf{v} := \otimes_{k \notin J} (\Lambda_k^{\text{Std}})^{\otimes a_k}$, then \mathbf{v} is fixed by $\gamma_n \mathbf{H} \gamma_n^{-1}$ for all n. Note that Λ_k^{Std} for $k \notin J$ and in particular \mathbf{v} are nontrivial eigenvectors of \mathbf{P}_J . The proof of Theorem B.1 in the \mathbf{SL}_N case is thus complete with $\mathbf{P} = \mathbf{P}_J$.

APPENDIX C. INDICES OF VARIOUS CONDITIONS

(B1) For every $x \in \mathbf{U}(\mathbb{Q})$ and $(g_n) \subset \mathbf{G}$ such that $\lim g_n \cdot x \in \mathbf{B}(\mathbb{R})$, one has

$$\lim_{n \to \infty} \left[(g_n)_* \mathbf{m}_{[\mathbf{H}_x^\circ]} \right] = \left[\mathbf{m}_{[\mathbf{G}]} \right]$$

- (BH1) For every $x \in \mathbf{U}(\mathbb{Q}), \nu_{\infty,x} := \lim_{R \to \infty} \nu_{R,x}$ exists and $\operatorname{supp}(\nu_{\infty,x}) \subset \mathbf{B}(\mathbb{R})$.
- (B2) For every $x \in \mathbf{U}(\mathbb{Q})$ and $y \in \overline{\mathbf{G}.x}$, one has $\overline{\mathbf{G}.y} \cap \mathbf{B}(\mathbb{R}) \neq \emptyset$.
- (K1) For every $\alpha \in \mathscr{A}_{\mathbb{R},\mathbb{Q}}^{\mathrm{an}}, d_{\alpha} 1 > 0.$
- (B4) For every $x \in \mathbf{U}(\mathbb{Q}), d_{\alpha} 1 > 0$ for some $\alpha \in \mathscr{B}_{\mathbb{R},x}^{\mathrm{an}}$. In particular, $\mathscr{B}_{\mathbb{R},x}^{\mathrm{an}}$ is non-empty.
- (H1) For every $x \in \mathbf{U}(\mathbb{Q})$ and arithmetic subgroup $\Gamma \subset \mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, \mathbf{H}_x° has no non-trivial \mathbb{Q} -characters and $\lim_{R\to\infty} \mu_{R,x}$ exists in $\operatorname{Prob}(\mathbf{G}/\Gamma)$.

- (D1) For every $x \in \mathbf{U}(\mathbb{Q})$ and arithmetic subgroup $\Gamma \subset \mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, \mathbf{H}_x° has no non-trivial \mathbb{Q} -characters and Ψ_x extends continuously to $\overline{\mathbf{G}.x} \to \operatorname{Prob}(\mathbf{G}/\Gamma) \cup \{\mathbf{0}\}$ where the closure is taken in $\mathbf{X}^{\operatorname{cor}}$, the manifold with corners associated with (\mathbf{X}, \mathbf{D}) (see Section 6.4.2).
- (S1) For every $x \in \mathbf{U}(\mathbb{Q})$ and arithmetic subgroup Γ in $\mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, there exists a bounded subset B of \mathbf{G}/Γ such that $g\mathbf{H}_x^\circ\Gamma/\Gamma$ intersects with B for every $g \in \mathbf{G}$.
- (DS1) There exists a closed subset D of $\mathbf{B}(\mathbb{R})$ such that for every (g_n) and every $x \in \mathbf{U}(\mathbb{Q})$ with lim $g_n.x$ not in D, there exists a bounded subset B of G/Γ such that $g_n \mathrm{H}^{\circ}_x \Gamma/\Gamma$ intersects with B for every n. Moreover, for every $F \subset \mathscr{C}^{\mathrm{an}}_{\mathbb{R},\mathbb{Q}}$, if Leb_F denotes a smooth measure on D_F , then Leb_F(B) = 0.
- (H2) For every $x \in \mathbf{U}(\mathbb{Q})$ and arithmetic subgroup $\Gamma \subset \mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, there exists a non-negative compactly supported continuous function ψ on \mathbf{G}/Γ such that $\langle \psi, g_*\mathbf{m}_{[\mathbf{H}_x^o]} \rangle \neq 0$ for all $g \in \mathbf{G}$ and for every such ψ , $\lim_{R\to\infty} \mu_{R,x}^{\psi}$ exists in $\operatorname{Prob}^{\psi}(\mathbf{G}/\Gamma)$.
- (D2) For every $x \in \mathbf{U}(\mathbb{Q})$ and arithmetic subgroup $\Gamma \subset \mathbf{G}(\mathbb{Q}) \cap \mathbf{G}$, there exists ψ satisfying the paragraph above and Ψ_x^{ψ} extends continuously to $\overline{\mathbf{G}.x} \to \operatorname{Prob}^{\psi}(\mathbf{G}/\Gamma)$ where the closure is taken in X^{cor}, the manifold with corners associated with (\mathbf{X}, \mathbf{D}) .
- (C1) $\mathbf{Z}(\mathbf{H}, \mathbf{L})$ decomposes into finitely many orbits under the action of $\mathbf{L} \times \mathbf{N}_{\mathbf{G}}(\mathbf{H})$ for every $\mathbf{L} \in$ INT (\mathbf{H}, \mathbf{G}) ;
- (N1) $\Gamma \cap \mathbf{H}$ is a finite index subgroup of $\Gamma \cap \mathbf{N}_{\mathbf{G}}(\mathbf{H})$ for an(y) arithmetic subgroup Γ .
- (F1) INT_{Γ}(**H**, **G**) is finite.
- (F2) there are only finitely many parabolic Q-subgroups containing **H**, i.e., $\mathscr{P}_{\mathbf{H}}$ is finite.
- (N2) $\mathbf{N}_{\mathbf{G}}(\mathbf{H})^{\circ} \subset \mathbf{N}_{\mathbf{G}}(\mathbf{L})$ for every $\mathbf{L} \in INT(\mathbf{H}, \mathbf{G})$.

Ackonwledgement

We are grateful to useful discussions with Jinpeng An, Yue Gao, Zhizhong Huang, Osama Khalil, Fei Xu and Pengyu Yang. We also thank Osama Khalil, Hee Oh, Fei Xu for comments and questions on an earlier version of this paper. The author is supported by NSFC, grant #12201013.

References

- [BHC62] Armand Borel and Harish-Chandra, Arithmetic subgroups of algebraic groups, Ann. of Math. (2) 75 (1962), 485–535. MR 0147566 18
- [Bir62] B. J. Birch, Forms in many variables, Proc. Roy. Soc. London Ser. A 265 (1961/62), 245–263. MR 150129 2
- [BJ06] Armand Borel and Lizhen Ji, Compactifications of symmetric and locally symmetric spaces, Mathematics: Theory & Applications, Birkhäuser Boston, Inc., Boston, MA, 2006. MR 2189882 69
- [BO12] Yves Benoist and Hee Oh, Effective equidistribution of S-integral points on symmetric varieties, Ann. Inst. Fourier (Grenoble) 62 (2012), no. 5, 1889–1942. MR 3025156 2
- [Bor63] Armand Borel, Some finiteness properties of adele groups over number fields, Inst. Hautes Études Sci. Publ. Math. (1963), no. 16, 5–30. MR 202718 10
- [Bor19] _____, Introduction to arithmetic groups, University Lectures Series, vol. 73, American Mathematical Soc., 2019. 9, 18
- [BR95] Mikhail Borovoi and Zeév Rudnick, Hardy-Littlewood varieties and semisimple groups, Invent. Math. 119 (1995), no. 1, 37–66. MR 1309971 1, 2, 9
- [BS64] A. Borel and J.-P. Serre, Théorèmes de finitude en cohomologie galoisienne, Comment. Math. Helv. 39 (1964), 111–164. MR 181643 10
- [BS73] _____, Corners and arithmetic groups, Comment. Math. Helv. 48 (1973), 436–491, Avec un appendice: Arrondissement des variétés à coins, par A. Douady et L. Hérault. MR 0387495 7, 69
- [BT65] Armand Borel and Jacques Tits, Groupes réductifs, Inst. Hautes Études Sci. Publ. Math. (1965), no. 27, 55–150. MR 207712 33
- [Cas04] Bill Casselman, Stability of lattices and the partition of arithmetic quotients, Asian J. Math. 8 (2004), no. 4, 607–637. MR 2127941 71, 73, 77
- [CLT10] Antoine Chambert-Loir and Yuri Tschinkel, Igusa integrals and volume asymptotics in analytic and adelic geometry, Confluentes Math. 2 (2010), no. 3, 351–429. MR 2740045 8, 13, 14, 34
- [CLT12] _____, Integral points of bounded height on partial equivariant compactifications of vector groups, Duke Math. J. 161 (2012), no. 15, 2799–2836. MR 2999313 2, 6
- [Con12] Brian Conrad, Weil and Grothendieck approaches to adelic points, Enseign. Math. (2) 58 (2012), no. 1-2, 61–97. MR 2985010 8

RUNLIN ZHANG

- [CTX09] Jean-Louis Colliot-Thélène and Fei Xu, Brauer-Manin obstruction for integral points of homogeneous spaces and representation by integral quadratic forms, Compos. Math. 145 (2009), no. 2, 309–363, With an appendix by Dasheng Wei and Xu. MR 2501421 1
- [DGU21] Christopher Daw, Alexander Gorodnik, and Emmanuel Ullmo, Convergence of measures on compactifications of locally symmetric spaces, Math. Z. 297 (2021), no. 3-4, 1293–1328. MR 4229603 64
- [DGU23] _____, The space of homogeneous probability measures on $\overline{\Gamma \setminus X}_{\max}^S$ is compact, Math. Ann. **386** (2023), no. 1-2, 987–1016, With an appendix by Jialun Li. MR 4585166 69
- [DM91] S. G. Dani and G. A. Margulis, Asymptotic behaviour of trajectories of unipotent flows on homogeneous spaces, Proc. Indian Acad. Sci. Math. Sci. 101 (1991), no. 1, 1–17. MR 1101994 18, 64
- [DRS93] W. Duke, Z. Rudnick, and P. Sarnak, Density of integer points on affine homogeneous varieties, Duke Math. J. 71 (1993), no. 1, 143–179. MR 1230289 2, 5, 14, 15, 16
- [dS23] Nicolas de Saxcé, Non-divergence in the space of lattices, Groups Geom. Dyn. 17 (2023), no. 3, 993–1003. MR 4622400 69
- [EM93] Alex Eskin and Curt McMullen, Mixing, counting, and equidistribution in Lie groups, Duke Math. J. 71 (1993), no. 1, 181–209. MR 1230290 2, 14, 15, 16
- [EMS96] Alex Eskin, Shahar Mozes, and Nimish Shah, Unipotent flows and counting lattice points on homogeneous varieties, Ann. of Math. (2) 143 (1996), no. 2, 253–299. MR 1381987 2, 5, 18, 19, 22, 31, 64
- [EMS97] A. Eskin, S. Mozes, and N. Shah, Non-divergence of translates of certain algebraic measures, Geom. Funct. Anal. 7 (1997), no. 1, 48–80. MR 1437473 18
- [EMV09] M. Einsiedler, G. Margulis, and A. Venkatesh, Effective equidistribution for closed orbits of semisimple groups on homogeneous spaces, Invent. Math. 177 (2009), no. 1, 137–212. MR 2507639 29
- [EV08] Jordan S. Ellenberg and Akshay Venkatesh, Local-global principles for representations of quadratic forms, Invent. Math. 171 (2008), no. 2, 257–279. MR 2367020 1
- [FZ13] Baohua Fu and De-Qi Zhang, A characterization of compact complex tori via automorphism groups, Math. Ann. 357 (2013), no. 3, 961–968. MR 3118620 13
- [GMB13] Philippe Gille and Laurent Moret-Bailly, Actions algébriques de groupes arithmétiques, Torsors, étale homotopy and applications to rational points, London Math. Soc. Lecture Note Ser., vol. 405, Cambridge Univ. Press, Cambridge, 2013, pp. 231–249. MR 3077171 9
- [GN12] Alexander Gorodnik and Amos Nevo, Counting lattice points, J. Reine Angew. Math. 663 (2012), 127–176. MR 2889708 2, 15
- [GOS09] Alexander Gorodnik, Hee Oh, and Nimish Shah, Integral points on symmetric varieties and Satake compactifications, Amer. J. Math. 131 (2009), no. 1, 1–57. MR 2488484 2
- [Gra84] Daniel R. Grayson, Reduction theory using semistability, Comment. Math. Helv. 59 (1984), no. 4, 600–634. MR 780079 71
- [Gro97] Frank D. Grosshans, Algebraic homogeneous spaces and invariant theory, Lecture Notes in Mathematics, vol. 1673, Springer-Verlag, Berlin, 1997. MR 1489234 23, 43
- [Hir64] Heisuke Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II, Ann. of Math. (2) 79 (1964), 109–203; 79 (1964), 205–326. MR 199184 33
- [HT99] Brendan Hassett and Yuri Tschinkel, Geometry of equivariant compactifications of Gⁿ_a, Internat. Math. Res. Notices (1999), no. 22, 1211–1230. MR 1731473 13
- [HT03] _____, Integral points and effective cones of moduli spaces of stable maps, Duke Math. J. 120 (2003), no. 3, 577–599. MR 2030096 2
- [HTT15] Brendan Hassett, Sho Tanimoto, and Yuri Tschinkel, Balanced line bundles and equivariant compactifications of homogeneous spaces, Int. Math. Res. Not. IMRN (2015), no. 15, 6375–6410. MR 3384482 13
- [Kem78] George R. Kempf, Instability in invariant theory, Ann. of Math. (2) 108 (1978), no. 2, 299–316. MR 506989 12, 23
- [KK18] Dubi Kelmer and Alex Kontorovich, Effective equidistribution of shears and applications, Math. Ann. 370 (2018), no. 1-2, 381–421. MR 3747491 2
- [KKV89] Friedrich Knop, Hanspeter Kraft, and Thierry Vust, The Picard group of a G-variety, Algebraische Transformationsgruppen und Invariantentheorie, DMV Sem., vol. 13, Birkhäuser, Basel, 1989, pp. 77–87. MR 1044586 10
- [KM98] D. Y. Kleinbock and G. A. Margulis, Flows on homogeneous spaces and Diophantine approximation on manifolds, Ann. of Math. (2) 148 (1998), no. 1, 339–360. MR 1652916 18, 75
- [Kol07] János Kollár, Lectures on resolution of singularities, Annals of Mathematics Studies, vol. 166, Princeton University Press, Princeton, NJ, 2007. MR 2289519 33
- [Mat64] Hideya Matsumoto, Quelques remarques sur les groupes de Lie algébriques réels, J. Math. Soc. Japan 16 (1964), 419–446. MR 183816 33
- [Mau07] François Maucourant, Homogeneous asymptotic limits of Haar measures of semisimple linear groups and their lattices, Duke Math. J. 136 (2007), no. 2, 357–399. MR 2286635 2
- [MG14] Amir Mohammadi and Alireza Salehi Golsefidy, Translate of horospheres and counting problems, Amer. J. Math. 136 (2014), no. 5, 1301–1346. MR 3263899 2
- [MT94] G. A. Margulis and G. M. Tomanov, Invariant measures for actions of unipotent groups over local fields on homogeneous spaces, Invent. Math. 116 (1994), no. 1-3, 347–392. MR 1253197 18
- [OS14] Hee Oh and Nimish A. Shah, Limits of translates of divergent geodesics and integral points on one-sheeted hyperboloids, Israel J. Math. 199 (2014), no. 2, 915–931. MR 3219562 2, 19

- [PR94] Vladimir Platonov and Andrei Rapinchuk, Algebraic groups and number theory, Pure and Applied Mathematics, vol. 139, Academic Press, Inc., Boston, MA, 1994, Translated from the 1991 Russian original by Rachel Rowen. MR 1278263 34
- [PRR23] Vladimir Platonov, Andrei Rapinchuk, and Igor Rapinchuk, Algebraic groups and number theory. Vol. I, second ed., Cambridge Studies in Advanced Mathematics, vol. 205, Cambridge University Press, Cambridge, [2023] ©2023. MR 4615820 10, 40
- [Rag72] M. S. Raghunathan, Discrete subgroups of Lie groups, Springer-Verlag, New York-Heidelberg, 1972, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 68. MR 0507234 11, 15
- [Ra91] Marina Ratner, On Raghunathan's measure conjecture, Ann. of Math. (2) 134 (1991), no. 3, 545–607. MR 1135878 5, 18
- [Ric67] R. W. Richardson, Jr., Conjugacy classes in Lie algebras and algebraic groups, Ann. of Math. (2) 86 (1967), 1–15. MR 217079 31
- [Rob88] Joel Roberts, Old and new results about the triangle varieties, Algebraic geometry (Sundance, UT, 1986), Lecture Notes in Math., vol. 1311, Springer, Berlin, 1988, pp. 197–219. MR 951647 52
- [RS84] Joel Roberts and Robert Speiser, Enumerative geometry of triangles. I, Comm. Algebra 12 (1984), no. 9-10, 1213–1255. MR 738546 52, 68
- [San23] Tim Santens, Manin's conjecture for integral points on toric varieties, arXiv e-prints (2023), arXiv:2312.13914. 6
- [Sem54] J. G. Semple, The triangle as a geometric variable, Mathematika 1 (1954), 80-88. MR 66677 52
- [Sha00] Nimish A. Shah, Counting integral matrices with a given characteristic polynomial, Sankhyā Ser. A 62 (2000), no. 3, 386–412, Ergodic theory and harmonic analysis (Mumbai, 1999). MR 1803465 2
- [Spr98] T. A. Springer, Linear algebraic groups, second ed., Progress in Mathematics, vol. 9, Birkhäuser Boston, Inc., Boston, MA, 1998. MR 1642713 13, 18, 21, 24, 75
- [SZ19] Uri Shapira and Cheng Zheng, Limiting distributions of translates of divergent diagonal orbits, Compositio Mathematica 155 (2019), no. 9, 1747–1793. 2, 7, 19, 23, 51, 68
- [Wei82] André Weil, Adeles and algebraic groups, Progress in Mathematics, vol. 23, Birkhäuser, Boston, MA, 1982, With appendices by M. Demazure and Takashi Ono. MR 670072 11
- [Wei98] Barak Weiss, Finite-dimensional representations and subgroup actions on homogeneous spaces, Israel J. Math. 106 (1998), 189–207. MR 1656889 18, 41
- [Wil22] Florian Wilsch, Integral points of bounded height on a certain toric variety, arXiv e-prints (2022), arXiv:2202.10909. 3
- [Wo05] Jarosł aw Wł odarczyk, Simple Hironaka resolution in characteristic zero, J. Amer. Math. Soc. 18 (2005), no. 4, 779–822. MR 2163383 33
- [WX16] Dasheng Wei and Fei Xu, Counting integral points in certain homogeneous spaces, J. Algebra 448 (2016), 350–398. MR 3438314 2, 9, 46, 47, 49
- [Zha19] Runlin Zhang, Limiting distribution of translates of the orbit of a maximal \mathbb{Q} -torus from identity on $\mathrm{SL}_N(\mathbb{R})/\mathrm{SL}_N(\mathbb{Z})$, Math. Ann. **375** (2019), no. 3-4, 1231–1281. MR 4023376 2, 19
- [Zha21] _____, Translates of homogeneous measures associated with observable subgroups on some homogeneous spaces, Compos. Math. 157 (2021), no. 12, 2657–2698. MR 4356197 2, 5, 18, 19, 20, 22, 23, 44, 78
- [Zha22] _____, Equidistribution of translates of a homogeneous measure on the Borel-Serre compactification, Discrete Contin. Dyn. Syst. 42 (2022), no. 4, 2053–2071. MR 4385787 18, 20, 21, 69, 75
- [ZZ21] Han Zhang and Runlin Zhang, Nondivergence on homogeneous spaces and rigid totally geodesics, arXiv e-prints (2021), arXiv:2111.02002. 29

College of Mathematics and Statistics, Center of Mathematics, Chongqing University, 401331, Chongqing, China.

 $Email \ address: \verb"zhangrunlin@outlook.com"$