Liquid drop with capillarity and rotating traveling waves

Pietro Baldi, Vesa Julin, Domenico Angelo La Manna

Abstract. We consider the free boundary problem for a 3-dimensional, incompressible, irrotational liquid drop of nearly spherical shape with capillarity. We study the problem from scratch, extending some classical results from the flat case (capillary water waves) to the spherical geometry: the reduction to a problem on the boundary, its Hamiltonian structure, the analyticity and tame estimates for the Dirichlet-Neumann operator in Sobolev class, and a linearization formula for it, both with the method of the good unknown of Alinhac and by a differential geometry approach. Then we prove the bifurcation of traveling waves, which are nontrivial (i.e., nonspherical) fixed profiles rotating with constant angular velocity.

Contents

1	Introduction and main results	1
2	Notations and parametrization of the geometric objects2.1Notations for differential operators on surfaces2.2Parametrization of the geometric objects	5 5 6
3	The water wave equations on the unit sphere3.1Reduction to an equivalent problem on the unit sphere3.2Hamiltonian structure	10 10 12
4	Shape derivative of the Dirichlet-Neumann operator4.1Proof by the method of the good unknown of Alinhac4.2Proof via geometric argument4.3Another proof of the Hamiltonian structure	16 17 23 27
5	Analyticity and tame estimates for the Dirichlet-Neumann operator5.1Analytic dependence in low norm5.2Sobolev spaces on domains and boundaries5.3Estimates in higher norms	28 28 30 36
6	Traveling waves6.1The linearized operator at zero6.2Symmetries and bifurcation from a simple eigenvalue6.3Arithmetics of simple eigenvalues	49 51 55 57
7	Appendix	59

1 Introduction and main results

We consider the free boundary problem for a liquid drop with capillarity, a problem already considered by Lord Rayleigh [37]. We do not consider gravity forces; in fact, a liquid drop with capillarity falling in the vacuum under gravity is described by the same system, as gravity can be removed from the equations by considering a reference frame that falls together with the drop.

We assume that the liquid drop occupies the time-dependent, open bounded region $\Omega_t \subset \mathbb{R}^3$ with smooth boundary $\partial \Omega_t$ for some time interval $t \in (0, T)$, and that the velocity vector field uand the pressure p are defined in Ω_t . Since the only effecting force is the surface tension, then Ω_t , u and p satisfy

$$\partial_t u + u \cdot \nabla u + \nabla p = 0 \quad \text{in } \Omega_t, \tag{1.1}$$

$$\operatorname{div} u = 0 \quad \text{in } \Omega_t, \tag{1.2}$$

$$p = \sigma_0 H_{\Omega_t} \quad \text{on } \partial \Omega_t, \tag{1.3}$$

$$V_t = \langle u, \nu_{\Omega_t} \rangle \quad \text{on } \partial \Omega_t, \tag{1.4}$$

where V_t is the normal velocity of the boundary, ν_{Ω_t} is the unit outer normal of the boundary, H_{Ω_t} is the mean curvature of the boundary $\partial\Omega_t$, σ_0 is the capillarity coefficient, and $\langle \cdot, \cdot \rangle$ denotes the scalar product of vectors in \mathbb{R}^3 . Equations (1.1), (1.2) are the Euler equations of incompressible fluid mechanics, (1.3) gives the pressure at the boundary in terms of capillarity, and (1.4) is the assumption that the movement of the boundary $\partial\Omega_t$ in its normal direction is due to the movement of the liquid particles on $\partial\Omega_t$, that is, the velocity of $\partial\Omega_t$ and the vector field u must have the same normal component at the boundary $\partial\Omega_t$. We call (1.1) the dynamics equation, (1.2) the incompressibility condition, (1.3) the condition for the pressure at the boundary, and (1.4) the kinematic condition. All four together form the free boundary problem for the motion of a drop of incompressible fluid with capillarity. The unknowns are the domain Ω_t , the velocity vector field u, and the pressure p.

An important property of system (1.1)-(1.4) is the conservation of the total energy

$$E(t) = \frac{1}{2} \int_{\Omega_t} |u|^2 dx + \sigma_0 \operatorname{Area}(\partial \Omega_t), \qquad (1.5)$$

which means that E(t) = E(0) for all $t \in (0, T)$; this follows from a straightforward calculation. The total fluid mass, i.e., the volume of Ω_t , and the velocity of the fluid barycenter are also conserved quantities.

If the velocity u is zero, then by (1.1) and (1.3) the mean curvature H_{Ω_t} is constant, and therefore by the Alexandrov theorem [4] the drop is the ball. Moreover, if the velocity is small (in C^1 -sense) and the droplet is uniformly C^2 -regular, then by [15] the drop is nearly spherical, i.e., a small perturbation of the ball. Hence, if we study solutions with small velocity, we may reduce to nearly spherical geometry. In particular, the drop is star-shaped.

In this paper we study the case when the vorticity is zero and Ω_t is star-shaped with respect to the origin, i.e., the boundary $\partial \Omega_t$ is the graph of a radial function over the sphere. We thus always assume that the domain Ω_t , or Ω when the time plays no role, is of the form

$$\partial\Omega_t = \{(1+h(t,x))x : x \in \mathbb{S}^2\} \quad \text{or} \quad \partial\Omega = \{(1+h(x))x : x \in \mathbb{S}^2\},\tag{1.6}$$

where $h(t, \cdot) : \mathbb{S}^2 \to (-1, \infty)$, or $h : \mathbb{S}^2 \to (-1, \infty)$ when time plays no role, is the elevation function. In particular, $\partial\Omega$ is diffeomorphic to the sphere with diffeomorphism $\gamma : \mathbb{S}^2 \to \partial\Omega$,

$$\gamma(x) = (1+h(x))x.$$
 (1.7)

In the case of moving boundary we denote $\gamma_t : \mathbb{S}^2 \to \partial \Omega_t$,

$$\gamma_t(x) = (1 + h(t, x))x.$$
 (1.8)

Since the vorticity is zero and Ω_t is diffeomorphic to the ball, there exists a velocity potential $\Phi: \Omega_t \to \mathbb{R}$ such that

$$u = \nabla \Phi \quad \text{in } \Omega_t. \tag{1.9}$$

Like in the case of water wave equations, we may use these assumptions to reduce system (1.1)-(1.4) into a system of two equations written in terms of the elevation function h in (1.6) and the function $\psi(t, \cdot) : \mathbb{S}^2 \to \mathbb{R}$ defined as the pullback by γ_t of the velocity potential at the boundary,

$$\psi(t,x) := \Phi(t,\gamma_t(x)). \tag{1.10}$$

In order to write these two equations we need to introduce further notation. Assuming sufficient regularity to exchange partial derivatives, one has $\partial_t u = \nabla \partial_t \Phi$ and $u \cdot \nabla u = \nabla (\frac{1}{2} |\nabla \Phi|^2)$. Hence the dynamics equation (1.1) becomes

$$\partial_t \Phi + \frac{1}{2} |\nabla \Phi|^2 + p = 0 \quad \text{in } \Omega_t, \tag{1.11}$$

which is an equation for equivalence classes with respect to the equivalence relation $f \sim g$ iff $\nabla f = \nabla g$. The incompressibility condition (1.2) becomes the Laplace equation

$$\Delta \Phi = 0 \quad \text{in } \Omega_t. \tag{1.12}$$

We define the Dirichlet-Neumann operator $G(h)\psi$, where h is the elevation function as in (1.6) and $\psi : \mathbb{S}^2 \to \mathbb{R}$ is a generic Dirichlet datum, as

$$G(h)\psi(x) := \langle (\nabla\Phi)(\gamma(x)), \nu_{\Omega}(\gamma(x)) \rangle$$
(1.13)

at all points $\gamma(x) \in \partial\Omega$, i.e., for all $x \in \mathbb{S}^2$, where $\Phi : \Omega \to \mathbb{R}$ is the solution of the boundary value problem

$$\Delta \Phi = 0 \text{ in } \Omega \quad \text{and} \quad \Phi(\gamma(x)) = \psi(x) \text{ for all } \gamma(x) \in \partial\Omega, \text{ i.e., for all } x \in \mathbb{S}^2.$$
(1.14)

Finally, we use the notation

$$H(h)(x) := H_{\Omega}(\gamma(x)) \quad \text{for } x \in \mathbb{S}^2, \tag{1.15}$$

where $\gamma : \mathbb{S}^2 \to \partial \Omega$ is defined in (1.7). We derive the formula of the mean curvature H(h) in terms of the elevation function h in Lemma 2.2 as the precise explicit formula is difficult to find in literature.

Under the assumptions of zero vorticity and star-shapedness of Ω_t , the free boundary problem (1.1)-(1.4) can be formulated as the system of two equations on \mathbb{S}^2

$$\partial_t h = \frac{\sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2}}{1+h} G(h)\psi, \tag{1.16}$$

$$\partial_t \psi - \frac{1}{2} \Big(G(h)\psi + \frac{\langle \nabla_{\mathbb{S}^2}\psi, \nabla_{\mathbb{S}^2}h \rangle}{(1+h)\sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2}h|^2}} \Big)^2 + \frac{|\nabla_{\mathbb{S}^2}\psi|^2}{2(1+h)^2} + \sigma_0 H(h) = 0, \tag{1.17}$$

where $\nabla_{\mathbb{S}^2}$ denotes the tangential gradient on \mathbb{S}^2 and, moreover, system (1.16), (1.17) is a Hamiltonian system. In the flat case $x \in \mathbb{R}^2$ or $x \in \mathbb{T}^2$, this is an old and well-known observation [45], [20]. In the spherical case $x \in \mathbb{S}^2$ the equivalence of system (1.1)-(1.4) and system (1.16), (1.17) is proved in [11, 38] but we give the argument in order to be self-contained, see Proposition 3.1. The Hamiltonian structure of (1.16), (1.17) is proved in [11] but we also prove it with a slightly more general argument, see Proposition 3.2 and Lemma 3.4.

Our aim is to study the motion of a drop when Ω_t is nearly spherical, i.e., when the elevation function in (1.6) is small. To this aim, we first study the linearization of equations (1.16), (1.17). The first main result of this paper is Theorem 4.1, which gives an explicit formula of the shape derivative of the Dirichlet-Neumann operator $G(h)\psi$ defined in (1.13) and (1.14), i.e., given a function $\eta: \mathbb{S}^2 \to \mathbb{R}$, we compute a formula for

$$G'(h)[\eta]\psi = \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} G(h+\varepsilon\eta)\psi.$$
(1.18)

The calculations for (1.18) are rather heavy, and we give two different methods to derive the desired formula. First, we consider the method of the good unknown of Alinhac from the theory of water waves [3], [30] and adapt it to the nearly spherical geometry. Interestingly, the method does not generalize from water waves to the drop trivially, because, on one side, the natural homogeneous extension of the diffeomorphism γ in (1.7) generates a singularity at the origin, which needs to be handled with care, and, on the other hand, the method shows a sort of rigidity, as it seems

it does not work with other extensions; see Remark 4.8 for a more detailed explanation of this (surprising!) difficulty. Then we also give a completely different argument for (1.18), which relies only on differential geometry. This geometric method is more direct and has the advantage that it can be adopted to more general setting as the method itself does not rely on the spherical symmetry. Then again, the fact that the reference manifold is the sphere and not the plane, as in the flat case, makes the calculations technically more challenging as the derivatives do not commute. On the other hand, the advantage of using the good unknown of Alinhac is that it does not require any knowledge in differential geometry and therefore we choose to give both arguments.

The second main result of the paper is Theorem 5.19, where we prove that the Dirichlet-Neumann operator in (1.13) depends analytically on the elevation function and prove tame estimates for it in Sobolev class. With respect to the flat case, the spherical geometry causes two technical difficulties. First, we need to derive functional inequalities, such as the product estimate, for high order Sobolev norms. A natural way to define high order Sobolev space on \mathbb{S}^2 is via the spherical harmonics, but spherical harmonics do not have the same nice properties as the complex exponentials on \mathbb{R}^n . For example, the product of two spherical harmonic functions is not a spherical harmonic. We overcome this issue by considering Sobolev spaces $H^s(\mathbb{S}^2)$ defined by partition of unity, rectification and extension following [33, 41]. We also consider non-isotropic versions of the spaces $H^s(\mathbb{R}^3_+)$ that are able to catch higher and fractional regularity in the tangential directions. These results in Section 5 are somewhat related to the idea of the "convolution by horizontal layers" in [17]. Second, we need to localize the argument near the boundary, since the homogeneous extension of the diffeomorphism γ in (1.7) has a singularity at the origin.

As an application of the analyticity proved in Section 5 we show in Theorem 6.14 the existence of traveling waves, i.e., non-trivial solutions of system (1.16), (1.17) which have a fixed profile that is rotating around the x_3 -axis with constant angular velocity. To the best of our knowledge this is the first existence result for three-dimensional capillary traveling waves on S^2 . An important consequence of Theorem 6.14 is that it provides the first example of global-in-time solutions for system (1.16), (1.17), for which, for general initial data, only local existence [11, 17, 39], continuation criteria and a priori estimates [29, 40] have been proved.

We prove Theorem 6.14 by applying the classical bifurcation theorem due to Crandall-Rabinowitz [21] for simple eigenvalues of the linearized operator. The set-up of the problem differs from the flat case and also the linearized operator is different due to the curvature of the sphere. This leads us to analyze solutions of certain Diophantine equation specific of the spherical geometry, see (6.48). By a careful arithmetical argument using prime number factorization, we are able to find infinitely many choices of angular velocity producing a one-dimensional kernel of the linearized operator.

Related literature. The Hamiltonian structure of the water wave equations has been proved to hold also with constant vorticity in [43]. With the method of the good unknown of Alinhac of [3], a formula for the shape derivative of the Dirichlet-Neumann operator is proved in [30] in the flat case. See also the recent work [28] in conical domains. A corresponding paralinearization formula is in [3] for the flat case, and in [39] for S^2 . The analyticity of the Dirichlet-Neumann operator as a function of the elevation function is proved in the flat case in dimension 2 and 3 by many authors, see, for example, the classical works [14, 16], the works [18, 19, 30], and the recent papers [10, 24]. See also the related paper [1]. Tame estimates for the Dirichlet-Neumann operator are proved in [2, 30] for the flat case; for related estimates on S^2 see [11]. The literature about traveling waves in the flat case is huge, for both pure gravity and gravity-capillary case, for both periodic profiles and solitary waves, both finite and infinite depth, with and without viscosity. For a comprehensive review of the existing literature we refer to the recent survey [26]; here we just mention the pioneering works [32, 36], the papers [18, 19, 25, 42, 44], and the recent works [9, 31]. We also mention [35], where rotating travelling waves are obtained for a 2-dimensional drop.

Acknowledgements. We thank Alberto Maspero for some interesting discussions. This work is supported by Italian GNAMPA, by Italian PRIN 2022E9CF89 *Geometric Evolution Problems* and Shape Optimization, by Italian PRIN 2020XB3EFL Hamiltonian and dispersive PDEs and the Academy of Finland grant 314227.

2 Notations and parametrization of the geometric objects

Throughout the paper, LHS means "left-hand side", RHS means "right-hand side", $\mathbb{N} = \{1, 2, \ldots\}$, and $\mathbb{N}_0 = \{0, 1, \ldots\}$.

2.1 Notations for differential operators on surfaces

We denote the fluid domain by $\Omega_t \subset \mathbb{R}^3$, or simply by Ω when time does not play any role, and assume it is star-shaped with respect to the origin. We may thus parametrize the boundary by the elevation (or height) function $h : \mathbb{S}^2 \to \mathbb{R}$ as in (1.6), and denote the associated diffeomorphism by $\gamma : \mathbb{S}^2 \to \partial \Omega$ as in (1.7). We always assume without mentioning that the elevation function satisfies h(x) > -1 for all $x \in \mathbb{S}^2$ in order to (1.6) make sense. If we assume that the Lipschitz norm of the elevation function is small, i.e., $\|h\|_{W^{1,\infty}(\mathbb{S}^2)} \leq c$ for some c < 1, then we call the domain Ω nearly spherical.

We assume that Ω is C^2 -regular, which for us means that the elevation function h is of class $C^2(\mathbb{S}^2)$, and we denote its outer unit normal by ν_{Ω} , the mean curvature by H_{Ω} and the second fundamental form by B_{Ω} . We use orientation for which H_{Ω} is nonnegative for convex sets. We denote the tangent plane at $x \in \partial \Omega$ by $T_x(\partial \Omega)$, which we may identify with the plane in \mathbb{R}^3

$$T_x(\partial\Omega) = \{ y \in \mathbb{R}^3 : \langle y, \nu_\Omega(x) \rangle = 0 \}.$$

We define the projection on $T_x(\partial\Omega)$ as

$$\Pi_{T_x(\partial\Omega)} = I - \nu_{\Omega}(x) \otimes \nu_{\Omega}(x), \tag{2.1}$$

where I is the identity matrix and \otimes denotes the tensor product (given $a, b \in \mathbb{R}^3$, $a \otimes b$ is the 3×3 matrix with entry $a_i b_j$ in the row i and column j, i.e., $a \otimes b$ is the matrix product ab^T of the column matrix a with the row matrix b^T). Note that in (2.1) we identify the projection map with the matrix representing it. We also remark that (2.1) is symmetric. We may split a given vector $a \in \mathbb{R}^3$ into normal and tangential components with respect to a fixed tangent plane $T_x(\partial\Omega)$ for $x \in \partial\Omega$ as

$$a_{\nu} = a_{\nu_{\Omega}(x)} = \langle a, \nu_{\Omega}(x) \rangle \,\nu_{\Omega}(x) \quad \text{and} \quad a_{\partial\Omega} = a - a_{\nu} = \Pi_{T_x(\partial\Omega)} a. \tag{2.2}$$

Then $|a|^2 = |a_{\nu}|^2 + |a_{\partial\Omega}|^2$. For a vector field $F : \mathbb{R}^3 \to \mathbb{R}^3$, F_{ν} and $F_{\partial\Omega}$ define normal and tangential vector fields on $\partial\Omega$. For a scalar function $f : \mathbb{R}^3 \to \mathbb{R}$, we define the normal and tangential gradient fields as

$$\nabla_{\nu} f(x) = \langle \nabla f(x), \nu_{\Omega}(x) \rangle \,\nu_{\Omega}(x) \quad \text{and} \quad \nabla_{\partial\Omega} f(x) = \Pi_{T_x(\partial\Omega)} \nabla f(x), \tag{2.3}$$

and we denote

$$D_{\partial\Omega}f(x) = [\nabla_{\partial\Omega}f(x)]^T = [\nabla f(x)]^T \Pi_{T_x(\partial\Omega)}$$
(2.4)

the transpose of the vector $\nabla_{\partial\Omega} f(x)$; thus, $\nabla_{\partial\Omega} f(x)$ is a vector of \mathbb{R}^3 , i.e., a column, while $D_{\partial\Omega} f(x)$ is a 1×3 matrix, i.e., a row. Since we assume that the domain Ω is at least C^2 -regular, we may extend any regular vector field $F : \partial\Omega \to \mathbb{R}^3$ to \mathbb{R}^3 and define the tangential differential at $x \in \partial\Omega$ as

$$D_{\partial\Omega}F(x) := DF(x) - DF(x)\nu_{\Omega}(x) \otimes \nu_{\Omega}(x) = DF(x)\Pi_{T_x(\partial\Omega)}, \qquad (2.5)$$

where DF(x) denotes the (Euclidian) differential (i.e., the Jacobian matrix) of the extension. Thus, $D_{\partial\Omega}F(x)$ is a 3 × 3 matrix, whose k-th row is $D_{\partial\Omega}F_k(x)$, the transpose of the tangential gradient of the k-th component F_k of the vector field F. It is easy to see that definition (2.5) is independent of the chosen extension. We remark that this definition, unlike the covariant derivative, does not generalize to tensors, but, since we only deal with first order derivatives of vector fields, definition (2.5) is enough for us. For a scalar function $f: \partial\Omega \to \mathbb{R}$, using any extension of it, the tangential gradient is defined in (2.3), and we define the tangential Hessian as

$$D^2_{\partial\Omega}f := D_{\partial\Omega}(\nabla_{\partial\Omega}f). \tag{2.6}$$

We also define the Laplace-Beltrami on $\partial \Omega$ as the trace of the tangential Hessian

$$\Delta_{\partial\Omega} f(x) = \operatorname{Tr} D^2_{\partial\Omega} f(x).$$
(2.7)

For a vector field $F : \partial \Omega \to \mathbb{R}^3$ we define the tangential divergence as the trace of the tangential differential

$$\operatorname{div}_{\partial\Omega} F = \operatorname{Tr}(D_{\partial\Omega}F) = \operatorname{div} F - \langle (DF)\nu_{\Omega}, \nu_{\Omega} \rangle, \qquad (2.8)$$

and note that it holds

$$\Delta_{\partial\Omega} f = \operatorname{div}_{\partial\Omega} (\nabla_{\partial\Omega} f) \tag{2.9}$$

and

$$H_{\Omega} = \operatorname{div}_{\partial\Omega} \nu_{\Omega}. \tag{2.10}$$

If f is defined in a neighborhood of $\partial \Omega$, it holds

$$\Delta_{\partial\Omega} f = \Delta f - \langle D^2 f \,\nu_{\Omega}, \nu_{\Omega} \rangle - H_{\Omega} \langle \nabla f, \nu_{\Omega} \rangle.$$
(2.11)

Finally, the divergence theorem for hypersurfaces states that

$$\int_{\partial\Omega} \operatorname{div}_{\partial\Omega} F \, d\mathcal{H}^2 = \int_{\partial\Omega} H_\Omega \langle F, \nu_\Omega \rangle \, d\mathcal{H}^2, \qquad (2.12)$$

where $d\mathcal{H}^2$ is the 2-dimensional Hausdorff measure (see, e.g., [34, 6]).

When Ω is the unit ball B_1 , its boundary is the unit sphere \mathbb{S}^2 , and we write $\nabla_{\mathbb{S}^2}$, div $_{\mathbb{S}^2}$, etc. instead of $\nabla_{\partial B_1}$, div $_{\partial B_1}$, etc. For the unit sphere, it is natural to extend a given scalar function $f: \mathbb{S}^2 \to \mathbb{R}$ to $\mathbb{R}^3 \setminus \{0\}$ in a homogeneous way as

$$\mathcal{E}_0 f(x) := f(\sigma(x)), \quad \mathcal{E}_1 f(x) := |x| f(\sigma(x)), \quad \text{where} \quad \sigma(x) := \frac{x}{|x|}, \tag{2.13}$$

for all $x \in \mathbb{R}^3 \setminus \{0\}$, and similarly for vector fields $F : \mathbb{S}^2 \to \mathbb{R}^3$. Note that the gradient and Hessian of 0-homogeneous extensions satisfy

$$\langle \nabla(\mathcal{E}_0 f)(x), x \rangle = 0, \quad D^2(\mathcal{E}_0 f)(x)x + \nabla(\mathcal{E}_0 f)(x) = 0$$
(2.14)

for all $x \in \mathbb{R}^3 \setminus \{0\}$ (differentiate the identity $\mathcal{E}_0 f(\lambda x) = \mathcal{E}_0 f(x)$ with respect to λ , and then with respect to x_k). For the unit sphere, the link between the tangential differential operators defined above and the corresponding classical differential operators for 0-homogeneous extensions become particularly simple: one has

$$\nabla_{\mathbb{S}^2} f(x) = \nabla(\mathcal{E}_0 f)(x), \qquad D_{\mathbb{S}^2} F(x) = D(\mathcal{E}_0 F)(x), \qquad D_{\mathbb{S}^2}^2 f(x) = D^2(\mathcal{E}_0 f)(x) + \nabla(\mathcal{E}_0 f)(x) \otimes x,$$

$$\Delta_{\mathbb{S}^2} f(x) = \Delta(\mathcal{E}_0 f)(x), \quad \operatorname{div}_{\mathbb{S}^2} F(x) = \operatorname{div}(\mathcal{E}_0 f)(x) \qquad (2.15)$$

for all $x \in \mathbb{S}^2$. From (2.14) and (2.15) it follows, in particular, that

$$\langle \nabla_{\mathbb{S}^2} f(x), x \rangle = 0, \tag{2.16}$$

$$\langle D_{\mathbb{S}^2}^2 f(x)v, x \rangle = -\langle \nabla_{\mathbb{S}^2} f(x), v \rangle$$
(2.17)

for all $x \in \mathbb{S}^2$, all $v \in T_x(\mathbb{S}^2)$.

2.2 Parametrization of the geometric objects

Now we write the normal unit vector and the mean curvature in terms of the elevation function. Let us first consider the diffeomorphism $\gamma : \mathbb{S}^2 \to \partial\Omega$ in (1.7), and take its 1-homogeneous extension

$$\mathcal{E}_1 \gamma(x) = x \left(1 + \mathcal{E}_0 h(x) \right), \tag{2.18}$$

defined on $\mathbb{R}^3 \setminus \{0\}$, where $\mathcal{E}_0, \mathcal{E}_1$ are defined in (2.13). Its Jacobian matrix is

$$D(\mathcal{E}_1\gamma) = (1 + \mathcal{E}_0h)I + x \otimes \nabla \mathcal{E}_0h.$$
(2.19)

The advantage of the extension in (2.18) is that its Jacobian matrix $D(\mathcal{E}_1\gamma)$ is invertible, and its inverse can be immediately calculated by observing that, by (2.16), $M = x \otimes \nabla(\mathcal{E}_0 h)$ is a nilpotent matrix satisfying $M^2 = 0$. Thus

$$[D(\mathcal{E}_1\gamma)]^{-1} = \frac{I}{1+\mathcal{E}_0h} - \frac{x \otimes \nabla \mathcal{E}_0h}{(1+\mathcal{E}_0h)^2}.$$
(2.20)

We also calculate the determinant

$$\det D(\mathcal{E}_1\gamma) = (1 + \mathcal{E}_0h)^3 \tag{2.21}$$

and the transpose of the inverse matrix

$$[D(\mathcal{E}_1\gamma)]^{-T} = \frac{I}{1+\mathcal{E}_0h} - \frac{(\nabla \mathcal{E}_0h) \otimes x}{(1+\mathcal{E}_0h)^2}.$$
(2.22)

Note that on \mathbb{S}^2 one has

$$D(\mathcal{E}_1\gamma) = (1+h)I + x \otimes \nabla_{\mathbb{S}^2}h \quad \text{on } \mathbb{S}^2$$
(2.23)

and

$$[D(\mathcal{E}_1\gamma)]^{-T} = \frac{I}{1+h} - \frac{(\nabla_{\mathbb{S}^2}h) \otimes x}{(1+h)^2} \quad \text{on } \mathbb{S}^2.$$
(2.24)

The tangent plane $T_{\gamma(x)}(\gamma(\mathbb{S}^2))$ to the surface $\gamma(\mathbb{S}^2)$ at the point $\gamma(x) \in \gamma(\mathbb{S}^2)$ is

$$T_{\gamma(x)}(\gamma(\mathbb{S}^2)) = \{ D_{\mathbb{S}^2}\gamma(x)v : v \in T_x(\mathbb{S}^2) \}, \quad \text{for all } x \in \mathbb{S}^2.$$

$$(2.25)$$

By (2.5), one has $D_{\mathbb{S}^2}\gamma(x)v = D\tilde{\gamma}(x)(I - x \otimes x)v = D\tilde{\gamma}(x)v$ for all $v \in T_x(\mathbb{S}^2)$, for any extension $\tilde{\gamma}$ of γ ; in particular, this holds for $\tilde{\gamma} = \mathcal{E}_1\gamma$. For all $v \in T_x(\mathbb{S}^2)$ one has

$$0 = \langle x, v \rangle = \langle x, [D\tilde{\gamma}(x)]^{-1} D\tilde{\gamma}(x) v \rangle = \langle [D\tilde{\gamma}(x)]^{-T} x, D\tilde{\gamma}(x) v \rangle, \quad \tilde{\gamma} = \mathcal{E}_1 \gamma_{+}$$

where $[D\tilde{\gamma}(x)]^{-T}$ is in (2.24). Hence, for $x \in \mathbb{S}^2$, the vector

$$N(\gamma)(x) := [D(\mathcal{E}_1\gamma)(x)]^{-T}x$$
(2.26)

satisfies $\langle N(\gamma)(x), w \rangle = 0$ for all $w \in T_{\gamma(x)}(\gamma(\mathbb{S}^2))$, i.e., it is orthogonal to the tangent plane (2.25). It is also easy to see that it points outside the domain Ω (note that, for h = 0 one has $N(\gamma)(x) = x$). Therefore the outward unit normal to the surface $\gamma(\mathbb{S}^2)$ at $\gamma(x)$ is, by (2.24) and (2.26),

$$\nu_{\Omega}(\gamma(x)) = \frac{N(\gamma)(x)}{|N(\gamma)(x)|} = \frac{(1+h(x))x - \nabla_{\mathbb{S}^2}h(x)}{\sqrt{(1+h(x))^2 + |\nabla_{\mathbb{S}^2}h(x)|^2}}$$
(2.27)

for $x \in \mathbb{S}^2$. For future purpose it is convenient to introduce the notation

$$J = J(h) = \sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2}.$$
(2.28)

In order to deal with the forthcoming computation, in the next lemma we give some general differentiation rules.

Lemma 2.1. Assume that the domain Ω and the elevation function $h : \mathbb{S}^2 \to \mathbb{R}$ are as in (1.6), and let γ be as in (1.7). For any scalar function $f : \partial \Omega \to \mathbb{R}$, let \tilde{f} be its pullback by γ , namely $\tilde{f}(x) = f(\gamma(x))$. Then

$$(\nabla_{\partial\Omega}f)(\gamma(x)) = \frac{\nabla_{\mathbb{S}^2}\tilde{f}(x)}{1+h} + \frac{\langle \nabla_{\mathbb{S}^2}\tilde{f}, \nabla_{\mathbb{S}^2}h\rangle}{(1+h)J}\nu_{\Omega}(\gamma(x))$$

$$= \frac{\nabla_{\mathbb{S}^2}\tilde{f}(x)}{1+h} - \frac{\langle \nabla_{\mathbb{S}^2}\tilde{f}, \nabla_{\mathbb{S}^2}h\rangle}{(1+h)J^2}\nabla_{\mathbb{S}^2}h + \frac{\langle \nabla_{\mathbb{S}^2}\tilde{f}, \nabla_{\mathbb{S}^2}h\rangle}{J^2}x,$$
(2.29)

where J is defined in (2.28). For a vector field $F : \partial \Omega \to \mathbb{R}^3$ we denote similarly $\tilde{F}(x) = F(\gamma(x))$. Then

$$(D_{\partial\Omega}F)(\gamma(x)) = \frac{D_{\mathbb{S}^2}\tilde{F}(x)}{1+h} - \frac{(D_{\mathbb{S}^2}\tilde{F})\nabla_{\mathbb{S}^2}h}{(1+h)J^2} \otimes \nabla_{\mathbb{S}^2}h + \frac{(D_{\mathbb{S}^2}\tilde{F})\nabla_{\mathbb{S}^2}h}{J^2} \otimes x,$$
(2.30)

$$(\operatorname{div}_{\partial\Omega}F)(\gamma(x)) = \frac{\operatorname{div}_{\mathbb{S}^2}\tilde{F}(x)}{1+h} - \frac{\langle (D_{\mathbb{S}^2}\tilde{F})\nabla_{\mathbb{S}^2}h, \nabla_{\mathbb{S}^2}h\rangle}{(1+h)J^2} + \frac{\langle (D_{\mathbb{S}^2}\tilde{F})\nabla_{\mathbb{S}^2}h, x\rangle}{J^2}.$$
(2.31)

Proof. We only need to prove (2.29), because (2.30) follows by applying (2.29) to the components of F, and (2.31) follows from (2.30) by applying the trace, since $\operatorname{div}_{\partial\Omega} F = \operatorname{Tr}(D_{\partial\Omega}F)$.

To prove (2.29), we extend $f : \partial \Omega \to \mathbb{R}$ by taking its 0-homogeneous extension, which we denote f_0 , namely we define $f_0(x) = f(\gamma(x/|x|))$ for all $x \in \mathbb{R}^3 \setminus \{0\}$. We also define $h_0 : \mathbb{R}^3 \setminus \{0\} \to \mathbb{R}$, $h_0(x) = (\mathcal{E}_0 h)(x) = h(x/|x|)$. Note that $\langle \nabla f_0, x \rangle = \langle \nabla h_0, x \rangle = 0$ on $\mathbb{R}^3 \setminus \{0\}$ and $\nabla h_0 = \nabla_{\mathbb{S}^2} h$ on \mathbb{S}^2 . We also extend γ as $\mathcal{E}_1 \gamma$ defined in (2.18).

For $\lambda > 0$ it holds $(f_0 \circ \mathcal{E}_1 \gamma)(\lambda x) = (f_0 \circ \mathcal{E}_1 \gamma)(x)$. Therefore for $x \in \mathbb{S}^2$ it holds

$$\nabla (f_0 \circ \mathcal{E}_1 \gamma)(x) = \nabla_{\mathbb{S}^2} (f_0 \circ \mathcal{E}_1 \gamma)(x) = \nabla_{\mathbb{S}^2} \tilde{f}(x).$$

On the other hand, one has

$$\nabla (f_0 \circ \mathcal{E}_1 \gamma)(x) = [D(\mathcal{E}_1 \gamma)(x)]^T (\nabla f_0)(\mathcal{E}_1 \gamma(x)).$$

Then, by (2.24) and the orthogonality $\langle \nabla_{\mathbb{S}^2} \tilde{f}, x \rangle = 0$, we have, for all $x \in \mathbb{S}^2$,

$$(\nabla f_0)(\gamma(x)) = [D(\mathcal{E}_1\gamma)(x)]^{-T} \nabla_{\mathbb{S}^2} \tilde{f}(x) = \frac{\nabla_{\mathbb{S}^2} \tilde{f}(x)}{1+h(x)}.$$

Then (2.29) follows from the identity

$$(\nabla_{\partial\Omega}f)(\gamma(x)) = (\nabla f_0)(\gamma(x)) - \langle (\nabla f_0)(\gamma(x)), \nu_{\Omega}(\gamma(x)) \rangle \nu_{\Omega}(\gamma(x))$$

(see definitions (2.1), (2.3)) and formula (2.27).

Note that in equation (1.17), as well as in (1.3), we have the mean curvature, which has an explicit formula in terms of the elevation function h. Such a formula follows from a classical calculation, which is, however, hard to find in literature in its complete version; see the recent paper [23]. For the sake of completeness, we give the calculations in the next lemma.

Lemma 2.2. Assume that the domain Ω and the elevation function $h : \mathbb{S}^2 \to \mathbb{R}$ are as in (1.6). Then the parametrization of the mean curvature is

$$H(h)(x) = H_{\Omega}(\gamma(x)) = -\frac{\Delta_{\mathbb{S}^2}h}{(1+h)J} + \frac{2}{J} + \frac{\langle (D_{\mathbb{S}^2}^2h)\nabla_{\mathbb{S}^2}h, \nabla_{\mathbb{S}^2}h \rangle}{(1+h)J^3} + \frac{|\nabla_{\mathbb{S}^2}h|^2}{J^3},$$
(2.32)

where J is defined in (2.28).

Proof. Let us recall formula (2.27) for the outer normal $\nu_{\Omega}(\gamma(x))$. We define the vector field $n: \partial\Omega \to \mathbb{R}^3$ such that, for $x \in \mathbb{S}^2$,

$$n(\gamma(x)) = \tilde{n}(x) = (1 + h(x))x - \nabla_{\mathbb{S}^2} h(x).$$
(2.33)

Then $\nu_{\Omega}(y) = \varphi(y)n(y)$ for $y \in \partial\Omega$, where $\varphi = 1/|n|$, and

$$H_{\Omega} = \operatorname{div}_{\partial\Omega} \nu_{\Omega} = \varphi \operatorname{div}_{\partial\Omega} n + \langle \nabla_{\partial\Omega} \varphi, n \rangle.$$

Since n(y) is in the direction of $\nu_{\Omega}(y)$, the last term is zero. Also, $|n(\gamma(x))| = J$. It remains to calculate $(\operatorname{div}_{\partial\Omega}n)(\gamma(x))$. To this aim, we apply formula (2.31) to the vector fields n, \tilde{n} . We first calculate $D_{\mathbb{S}^2}\tilde{n}$. By (2.33), (2.5), (2.6), (2.16), we have

$$D_{\mathbb{S}^2}\tilde{n}(x) = -D_{\mathbb{S}^2}^2 h + (1+h)I_{\mathbb{S}^2} + x \otimes \nabla_{\mathbb{S}^2} h, \qquad (2.34)$$

where $I_{\mathbb{S}^2} = I - x \otimes x$ (i.e., $I_{\mathbb{S}^2}$ is the matrix $\Pi_{T_x(\mathbb{S}^2)}$ in (2.1)) By (2.34), (2.8), (2.7), (2.16), we obtain

$$\begin{aligned} \operatorname{div}_{\mathbb{S}^2} \tilde{n}(x) &= -\Delta_{\mathbb{S}^2}^2 h + 2(1+h), \\ \langle (D_{\mathbb{S}^2} \tilde{n}) \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} h \rangle &= -\langle (D_{\mathbb{S}^2}^2 h) \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} h \rangle + (1+h) |\nabla_{\mathbb{S}^2} h|^2. \end{aligned}$$

Moreover, by (2.34) and (2.17),

$$\langle (D_{\mathbb{S}^2}\tilde{n})\nabla_{\mathbb{S}^2}h, x \rangle = -\langle (D_{\mathbb{S}^2}^2h)\nabla_{\mathbb{S}^2}h, x \rangle + |\nabla_{\mathbb{S}^2}h|^2 = 2|\nabla_{\mathbb{S}^2}h|^2.$$

We use (2.31) and the above calculations to deduce that

$$(\operatorname{div}_{\partial\Omega}n)(\gamma(x)) = \frac{\operatorname{div}_{\mathbb{S}^2}\tilde{n}(x)}{1+h} - \frac{\langle (D_{\mathbb{S}^2}\tilde{n})\nabla_{\mathbb{S}^2}h, \nabla_{\mathbb{S}^2}h\rangle}{(1+h)J^2} + \frac{\langle (D_{\mathbb{S}^2}\tilde{n})\nabla_{\mathbb{S}^2}h, x\rangle}{J^2}$$
$$= -\frac{\Delta_{\mathbb{S}^2}^2h}{1+h} + 2 + \frac{\langle (D_{\mathbb{S}^2}^2h)\nabla_{\mathbb{S}^2}h, \nabla_{\mathbb{S}^2}h\rangle}{(1+h)J^2} + \frac{|\nabla_{\mathbb{S}^2}h|^2}{J^2}.$$

Since $|n(\gamma(x))| = J$, the lemma is proved.

To conclude this subsection we parametrize the Laplace-Beltrami operator on $\partial\Omega$. This is needed in subsection 4.2.

Lemma 2.3. Assume that the domain Ω and the elevation function $h : \mathbb{S}^2 \to \mathbb{R}$ are as in (1.6). Let $f : \partial \Omega \to \mathbb{R}$ and denote $\tilde{f}(x) = f(\gamma(x))$. The parametrization of the Laplace-Beltrami operator is

$$(\Delta_{\partial\Omega}f)(\gamma(x)) = \frac{\Delta_{\mathbb{S}^2}\tilde{f}}{(1+h)^2} - \frac{\langle (D_{\mathbb{S}^2}^2\tilde{f})\nabla_{\mathbb{S}^2}h, \nabla_{\mathbb{S}^2}h\rangle}{(1+h)^2J^2} - 2\frac{\langle \nabla_{\mathbb{S}^2}\tilde{f}, \nabla_{\mathbb{S}^2}h\rangle}{(1+h)J^2} + H(h)\frac{\langle \nabla_{\mathbb{S}^2}\tilde{f}, \nabla_{\mathbb{S}^2}h\rangle}{(1+h)J},$$

where J is defined in (2.28) and the mean curvature $H(h) = H_{\Omega}(\gamma(x))$ is calculated in Lemma 2.2. Proof. By (2.29) and (2.27), it holds

$$(\nabla_{\partial\Omega}f)(\gamma(x)) = \underbrace{\frac{\nabla_{\mathbb{S}^2}\tilde{f}(x)}{1+h}}_{=\tilde{F}_1(x)} + \underbrace{\frac{\langle\nabla_{\mathbb{S}^2}\tilde{f},\nabla_{\mathbb{S}^2}h\rangle}{(1+h)J}\nu_{\Omega}(\gamma(x))}_{=\tilde{F}_2(x)}.$$

Recalling (2.9), we have to calculate $\Delta_{\partial\Omega} f = \operatorname{div}_{\partial\Omega}(\nabla_{\partial\Omega} f) = \operatorname{div}_{\partial\Omega}(F_1 + F_2)$, where F_1, F_2 are defined by the equality $F_i(\gamma(x)) = \tilde{F}_i(x)$, i = 1, 2, for $x \in \mathbb{S}^2$. We immediately notice that, since F_2 is of the form $\varphi \nu_{\Omega}$ for a scalar function φ , then, by (2.8), (2.10), we have $\operatorname{div}_{\partial\Omega}(\varphi \nu_{\Omega}) = \varphi H_{\Omega}$, and therefore

$$(\operatorname{div}_{\partial\Omega}F_2)(\gamma(x)) = H(h) \frac{\langle \nabla_{\mathbb{S}^2} \tilde{f}, \nabla_{\mathbb{S}^2} h \rangle}{(1+h)J}, \qquad (2.35)$$

where H(h) is defined in (1.15). In order to parametrize div $\partial_{\Omega}F_1$, first, by (2.5), (2.6), we calculate

$$D_{\mathbb{S}^2}\tilde{F}_1 = \frac{D_{\mathbb{S}^2}^2\tilde{f}}{1+h} - \frac{\nabla_{\mathbb{S}^2}\tilde{f}\otimes\nabla_{\mathbb{S}^2}h}{(1+h)^2}.$$

By (2.8), (2.7), (2.17), this yields

$$\begin{aligned} \operatorname{div}_{\mathbb{S}^{2}}\tilde{F}_{1} &= \frac{\Delta_{\mathbb{S}^{2}}f}{1+h} - \frac{\langle \nabla_{\mathbb{S}^{2}}f, \nabla_{\mathbb{S}^{2}}h \rangle}{(1+h)^{2}}, \\ \langle (D_{\mathbb{S}^{2}}\tilde{F}_{1})\nabla_{\mathbb{S}^{2}}h, \nabla_{\mathbb{S}^{2}}h \rangle &= \frac{\langle (D_{\mathbb{S}^{2}}^{2}\tilde{f})\nabla_{\mathbb{S}^{2}}h, \nabla_{\mathbb{S}^{2}}h \rangle}{1+h} - \frac{|\nabla_{\mathbb{S}^{2}}h|^{2}}{(1+h)^{2}} \langle \nabla_{\mathbb{S}^{2}}\tilde{f}, \nabla_{\mathbb{S}^{2}}h \rangle, \\ \langle (D_{\mathbb{S}^{2}}\tilde{F}_{1})\nabla_{\mathbb{S}^{2}}h, x \rangle &= \frac{\langle (D_{\mathbb{S}^{2}}^{2}\tilde{f})\nabla_{\mathbb{S}^{2}}h, x \rangle}{1+h} = -\frac{\langle \nabla_{\mathbb{S}^{2}}\tilde{f}, \nabla_{\mathbb{S}^{2}}h \rangle}{1+h}. \end{aligned}$$

We have then, by (2.31) and (2.28), that

$$\begin{split} (\operatorname{div}_{\partial\Omega}F_{1})(\gamma(x)) \\ &= \frac{\Delta_{\mathbb{S}^{2}}\tilde{f}}{(1+h)^{2}} - \frac{\langle \nabla_{\mathbb{S}^{2}}\tilde{f}, \nabla_{\mathbb{S}^{2}}h \rangle}{(1+h)^{3}} - \frac{\langle (D_{\mathbb{S}^{2}}^{2}\tilde{f})\nabla_{\mathbb{S}^{2}}h, \nabla_{\mathbb{S}^{2}}h \rangle}{(1+h)^{2}J^{2}} + \frac{|\nabla_{\mathbb{S}^{2}}h|^{2}\langle \nabla_{\mathbb{S}^{2}}\tilde{f}, \nabla_{\mathbb{S}^{2}}h \rangle}{(1+h)^{3}J^{2}} - \frac{\langle \nabla_{\mathbb{S}^{2}}\tilde{f}, \nabla_{\mathbb{S}^{2}}h \rangle}{(1+h)J^{2}} \\ &= \frac{\Delta_{\mathbb{S}^{2}}\tilde{f}}{(1+h)^{2}} - \frac{\langle (D_{\mathbb{S}^{2}}^{2}\tilde{f})\nabla_{\mathbb{S}^{2}}h, \nabla_{\mathbb{S}^{2}}h \rangle}{(1+h)^{2}J^{2}} - \frac{2\langle \nabla_{\mathbb{S}^{2}}\tilde{f}, \nabla_{\mathbb{S}^{2}}h \rangle}{(1+h)J^{2}}. \end{split}$$

This and (2.35) yield the claim.

We notice that we may define the elliptic operator $\mathcal{L}_h : C^2(\mathbb{S}^2) \to C(\mathbb{S}^2)$,

$$\mathcal{L}_{h}[\varphi] = \Delta_{\mathbb{S}^{2}}\varphi - \frac{\langle (D_{\mathbb{S}^{2}}^{2}\varphi)\nabla_{\mathbb{S}^{2}}h, \nabla_{\mathbb{S}^{2}}h\rangle}{J^{2}}, \qquad (2.36)$$

and the Laplace-Beltrami operator in Lemma 2.3 can be written as

$$(\Delta_{\partial\Omega}f)(\gamma(x)) = \frac{\mathcal{L}_h[\tilde{f}]}{(1+h)^2} + \left(-\frac{\mathcal{L}_h[h]}{(1+h)^2 J^2} + \frac{|\nabla_{\mathbb{S}^2}h|^2}{(1+h)J^4}\right) \langle \nabla_{\mathbb{S}^2}\tilde{f}, \nabla_{\mathbb{S}^2}h \rangle.$$
(2.37)

3 The water wave equations on the unit sphere

In this section we prove that for the star-shaped liquid drop with zero vorticity the system of equations (1.1)-(1.4) is equivalent to (1.16), (1.17). We then show that the equations (1.16), (1.17) have Hamiltonian structure. These results are classical for the water wave equations in nearly flat case.

3.1 Reduction to an equivalent problem on the unit sphere

In this subsection we show that the free boundary problem for the capillary liquid drop can be formulated as system (1.16), (1.17). In the flat case where $x \in \mathbb{R}^2$ or $x \in \mathbb{T}^2$, this is an old and well-known observation; we show here the analogue observation for the spherical case $x \in \mathbb{S}^2$.

Proposition 3.1. Under the condition of zero vorticity curl u = 0 and star-shapedness of Ω_t , system (1.1)-(1.4) is equivalent to system (1.16), (1.17), where ψ is defined in(1.10) and the elevation function h is in (1.6).

Proof. First we show that (1.1)-(1.4) imply (1.16), (1.17). We begin by observing that the parametrization of equation (1.4) is given by (1.16): the boundary $\partial \Omega_t$ in (1.6) is described by γ_t in (1.8), and the normal velocity V_t appearing in the left hand side of (1.4) is, by definition, the normal component at $\gamma_t(x) \in \partial \Omega_t$ of the time derivative $\partial_t \gamma$, namely

$$V_t(\gamma_t(x)) = \langle \partial_t \gamma_t(x), \nu_{\Omega_t}(\gamma_t(x)) \rangle = \frac{(1+h)}{J} \partial_t h \quad \text{on } \mathbb{S}^2,$$
(3.1)

where we have used formula (2.27) for the unit normal, the orthogonality property (2.16) and the definition (2.28) of J. On the other hand, the term $\langle u, \nu_{\Omega} \rangle$ in the right hand side of (1.4) can be written by using assumption (1.9) and the definition (1.13) of the Dirichlet-Neumann operator, i.e.,

$$\langle u(t,\gamma_t(x)),\nu_{\Omega_t}(\gamma_t(x))\rangle = \langle (\nabla\Phi)(t,\gamma_t(x)),\nu_{\Omega_t}(\gamma_t(x))\rangle = G(h)\psi(x)$$
(3.2)

for all $x \in \mathbb{S}^2$. Thus (1.4) becomes (1.16).

We proceed to derive (1.17). By continuity, equation (1.11) also holds on the boundary $\partial \Omega_t$ and, under the pressure condition (1.3), we have

$$\partial_t \Phi + \frac{1}{2} |\nabla \Phi|^2 + \sigma_0 H_{\Omega_t} = 0 \quad \text{on } \partial \Omega_t.$$
(3.3)

We recall that Φ, ψ satisfy identity (1.10). By differentiating (1.10) with respect to time, we have

$$(\partial_t \Phi)(t, \gamma_t(x)) + \langle (\nabla \Phi)(t, \gamma_t(x)), \partial_t \gamma_t(x) \rangle = \partial_t \psi(t, x)$$

for all $x \in \mathbb{S}^2$. Therefore (3.3) becomes

$$\partial_t \psi(t,x) - \langle (\nabla \Phi)(t,\gamma_t(x)), \partial_t \gamma_t(x) \rangle + \frac{1}{2} |(\nabla \Phi)(t,\gamma_t(x))|^2 + \sigma_0 H(h)(x) = 0$$

for all $x \in \mathbb{S}^2$, where H(h)(x) is defined in (1.15). We then split the gradient $\nabla \Phi$ into its normal $\nabla_{\nu} \Phi$ and tangential $\nabla_{\partial \Omega} \Phi$ part defined in (2.3). Using the definition of the Dirichlet-Neumann operator (1.13), we have

$$(\nabla_{\nu}\Phi)(t,\gamma_t(x)) = G(h)\psi(x)\nu_{\Omega}(\gamma_t(x)).$$
(3.4)

Therefore we deduce that

$$\begin{aligned} \partial_t \psi(t,x) - G(h)\psi \langle \partial_t \gamma_t, \nu_{\Omega_t}(\gamma_t(x)) \rangle &- \langle \nabla_{\partial \Omega_t} \Phi, \partial_t \gamma_t \rangle \\ &+ \frac{1}{2} (G(h)(\psi))^2 + \frac{1}{2} |\nabla_{\partial \Omega_t} \Phi|^2 + \sigma_0 H(h)(x) = 0. \end{aligned}$$

Now $\langle \partial_t \gamma_t, \nu_{\Omega_t}(\gamma_t) \rangle = G(h)\psi$ by (3.1), (3.2) and (1.4). Hence

$$\partial_t \psi - \frac{1}{2} (G(h)\psi)^2 - \langle \nabla_{\partial\Omega_t} \Phi, \partial_t \gamma_t \rangle + \frac{1}{2} |\nabla_{\partial\Omega_t} \Phi|^2 + \sigma_0 H(h) = 0.$$
(3.5)

We write the tangential part $(\nabla_{\partial\Omega_t} \Phi)(t, \gamma_t(x))$ using (2.29) and (1.10), and obtain

$$(\nabla_{\partial\Omega}\Phi)(t,\gamma_t(x)) = \frac{\nabla_{\mathbb{S}^2}\psi}{1+h} - \frac{\langle\nabla_{\mathbb{S}^2}\psi,\nabla_{\mathbb{S}^2}h\rangle}{(1+h)J^2}\nabla_{\mathbb{S}^2}h + \frac{\langle\nabla_{\mathbb{S}^2}\psi,\nabla_{\mathbb{S}^2}h\rangle}{J^2}x,$$
(3.6)

with $x \in \mathbb{S}^2$. By (3.6), we calculate

$$|(\nabla_{\partial\Omega}\Phi)(t,\gamma_t(x))|^2 = \frac{|\nabla_{\mathbb{S}^2}\psi|^2}{(1+h)^2} - \frac{\langle\nabla_{\mathbb{S}^2}\psi,\nabla_{\mathbb{S}^2}h\rangle^2}{(1+h)^2J^2}.$$
(3.7)

Moreover, since $\partial_t \gamma_t = x \partial_t h$, we have

$$\langle (\nabla_{\partial\Omega}\Phi)(t,\gamma_t(x)), \partial_t\gamma_t(x)\rangle = \frac{\langle \nabla_{\mathbb{S}^2}\psi, \nabla_{\mathbb{S}^2}h\rangle}{J^2} \partial_t h = \frac{\langle \nabla_{\mathbb{S}^2}\psi, \nabla_{\mathbb{S}^2}h\rangle}{(1+h)J}G(h)\psi,$$
(3.8)

where in the last equality we have used (1.16). Combining (3.5) with (3.7) and (3.8) yields

$$\partial_t \psi - \frac{1}{2} (G(h)\psi)^2 - \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle}{(1+h)J} G(h)\psi - \frac{1}{2} \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle^2}{(1+h)^2 J^2} + \frac{1}{2} \frac{|\nabla_{\mathbb{S}^2} \psi|^2}{(1+h)^2} + \sigma_0 H(h) = 0.$$

Then (1.17) follows by noticing that the three terms with minus sign form a square.

Now we prove that (1.16), (1.17) imply (1.1)-(1.4) with curl u = 0. Suppose that two functions h and ψ , defined on $(0, T) \times \mathbb{S}^2$, satisfy the kinematic equation (1.16) and the dynamics equation (1.17). Define the set Ω_t as in (1.6) and $\Phi(t, \cdot)$ in Ω_t as the solution of the Laplace problem (1.14). Then $\Phi(t, \cdot)$ satisfies the incompressibility condition $\Delta \Phi = 0$ in Ω_t . By (1.16) and (1.13), equation (1.4) is also satisfied. From (1.17), using (1.16), we obtain (3.3). Now we define p on the closure of Ω_t as

$$p := -\partial_t \Phi - \frac{1}{2} |\nabla \Phi|^2 \quad \text{in } \overline{\Omega_t} = \Omega_t \cup \partial \Omega_t.$$
(3.9)

Then the dynamics equation (1.11) in the open domain Ω_t trivially holds. From (3.9) at the boundary $\partial \Omega_t$ and (3.3) (which is an identity for points of the boundary $\partial \Omega_t$) we deduce that $p = \sigma_0 H_{\Omega_t}$ on $\partial \Omega_t$, i.e., (1.3).

3.2 Hamiltonian structure

In this subsection we prove that equations (1.16) and (1.17) form a Hamiltonian system. Similarly as for Proposition 3.1, also the Hamiltonian structure of the water wave system is an old and well-known result in the flat case $x \in \mathbb{R}^2$ or $x \in \mathbb{T}^2$, which goes back to [45, 20]. In this subsection we prove the analogue result for the spherical case $x \in \mathbb{S}^2$.

We remark that, concerning the Hamiltonian structure, with the spherical geometry there is a difference with respect to the flat case: while on \mathbb{R}^2 or \mathbb{T}^2 the elevation h and the value ψ of the velocity potential at the boundary are Darboux coordinates of the system, on \mathbb{S}^2 this is not true. However, (h, ψ) fail to be Darboux coordinates only because of a "wrong" multiplicative factor $(1+h)^2$ (see Lemma 3.2 below), and it is not difficult to obtain Darboux coordinates with a simple change of coordinate (Lemma 3.4).

We start with writing the energy (1.5) in terms of h, ψ . For $\partial \Omega = \gamma(\mathbb{S}^2)$, where γ is in (1.7), the area formula gives

Area
$$(\partial \Omega) = \int_{\mathbb{S}^2} (1+h)\sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2}h|^2} \, d\sigma =: U(h),$$
 (3.10)

where the last identity defines U(h). Recall that, given $\psi : \mathbb{S}^2 \to \mathbb{R}$, we denote $\Phi : \Omega \to \mathbb{R}$ the solution of problem (1.14) We extend γ to $\mathbb{R}^3 \setminus \{0\}$ by $\mathcal{E}_1 \gamma$ defined in (2.18). If h is smooth, then $\mathcal{E}_1 \gamma$ is smooth in $\mathbb{R}^3 \setminus \{0\}$, and it can be extended to a Lipschitz continuous map in the whole \mathbb{R}^3 , mapping 0 to itself and B_1 onto Ω bijectively, which in this subsection, with a little abuse of notation, we also denote by $\mathcal{E}_1 \gamma$. Therefore the function

$$\Phi(x) = \Phi(\mathcal{E}_1 \gamma(x)) \quad \text{for } x \in B_1 \tag{3.11}$$

is Lipschitz continuous in B_1 , it satisfies

$$\nabla \tilde{\Phi}(x) = [D(\mathcal{E}_1 \gamma)(x)]^T (\nabla \Phi)(\mathcal{E}_1 \gamma(x)) \quad \text{in } B_1 \setminus \{0\},$$
(3.12)

and hence, by arguing as in [22] (section 6.3, page 320), it is a weak solution of the Dirichlet problem

$$\operatorname{div}(P\nabla\Phi) = 0 \quad \text{in } B_1, \quad \Phi = \psi \quad \text{on } \mathbb{S}^2, \tag{3.13}$$

where the matrix P(x) is

$$P(x) := \det(D(\mathcal{E}_1\gamma)(x))[D(\mathcal{E}_1\gamma)(x)]^{-1}[D(\mathcal{E}_1\gamma)(x)]^{-T}$$
(3.14)

in $\mathbb{R}^3 \setminus \{0\}$ (note that P(x) is not defined at x = 0, but this is irrelevant for weak solutions). We use (2.20), (2.21) and (2.22) to calculate

$$P(x) = (1 + \mathcal{E}_0 h)I - (\nabla \mathcal{E}_0 h) \otimes x - x \otimes (\nabla \mathcal{E}_0 h) + \frac{|\nabla \mathcal{E}_0 h|^2}{1 + \mathcal{E}_0 h} x \otimes x$$
(3.15)

for $x \in \mathbb{R}^3 \setminus \{0\}$. If *h* is Lipschitz continuous with a norm $\|h\|_{W^{1,\infty}(\mathbb{S}^2)} \leq C$, then it is straightforward to see that the matrix *P* is uniformly elliptic on $B_1 \setminus \{0\}$, i.e., $c_0|\xi|^2 \leq \langle P(x)\xi,\xi \rangle \leq C_0|\xi|^2$ for all $\xi \in \mathbb{R}^3$, all $0 < |x| \leq 1$, for some constants $0 < c_0 < C_0$ independent of x,ξ . Therefore the differential operator in (3.13) is uniformly elliptic, and the weak solution of problem (3.13) is unique, i.e., $\tilde{\Phi}$ is its unique weak solution.

By the change of variable $y = \mathcal{E}_1 \gamma(x)$, the divergence theorem and (3.13), we have

$$\int_{\Omega} |\nabla \Phi|^2 \, dx = \int_{B_1} \langle P \nabla \tilde{\Phi}, \nabla \tilde{\Phi} \rangle \, dx = \int_{\mathbb{S}^2} \psi \langle P \nabla \tilde{\Phi}, x \rangle \, d\sigma. \tag{3.16}$$

By the definition (1.13) of the Dirichlet-Neumann, the differentiation rule (3.12), the first identity in (2.27) for the unit normal ν_{Ω} , formula (2.26) for the normal vector N, and definition (3.14) of the matrix P, we get

$$G(h)\psi(x) = \langle [D(\mathcal{E}_{1}\gamma)(x)]^{-T}\nabla\tilde{\Phi}(x), \frac{[D(\mathcal{E}_{1}\gamma)(x)]^{-T}x}{|[D(\mathcal{E}_{1}\gamma)(x)]^{-T}x|} \rangle$$
$$= \frac{\langle P(x)\nabla\tilde{\Phi}(x), x \rangle}{|[D(\mathcal{E}_{1}\gamma)(x)]^{-T}x|\det[D(\mathcal{E}_{1}\gamma)(x)]}$$
(3.17)

for $x \in \mathbb{S}^2$. The matrix $[D(\mathcal{E}_1\gamma)(x)]^{-T}$ on \mathbb{S}^2 is given by (2.24), and therefore

$$[D(\mathcal{E}_1\gamma)]^{-T}x = \frac{x}{1+h} - \frac{\nabla_{\mathbb{S}^2}h}{(1+h)^2} \quad \text{and} \quad |[D(\mathcal{E}_1\gamma)(x)]^{-T}x| = \frac{J}{(1+h)^2}$$
(3.18)

on \mathbb{S}^2 , where J is in (2.28). Thus, by (3.17), (3.18) and (2.21), we deduce that

$$G(h)\psi = \frac{\langle P\nabla\bar{\Phi}, x\rangle}{(1+h)J}$$
(3.19)

on \mathbb{S}^2 , where J is in (2.28), P is in (3.14), and $\tilde{\Phi}$ is the solution of (3.13). By (3.16) and (3.19) we have that

$$\frac{1}{2} \int_{\Omega} |\nabla \Phi|^2 \, dx = \frac{1}{2} \int_{\mathbb{S}^2} \psi \left(G(h)\psi \right) (1+h) \sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2} \, d\sigma =: K(h,\psi), \tag{3.20}$$

where the last identity defines K. Hence the energy (1.5) written in terms of h, ψ is

$$\mathcal{H}(h,\psi) = K(h,\psi) + \sigma_0 U(h), \qquad (3.21)$$

with K in (3.20) and U in (3.10). With the same calculations above, given h, ψ_1, ψ_2 , with corresponding $\Phi_i, \tilde{\Phi}_i, i = 1, 2$, one has

$$\int_{\Omega} \langle \nabla \Phi_1, \nabla \Phi_2 \rangle \, dx = \int_{B_1} \langle P \nabla \tilde{\Phi}_1, \nabla \tilde{\Phi}_2 \rangle \, dx = \int_{\mathbb{S}^2} \psi_2 \langle P \nabla \tilde{\Phi}_1, x \rangle \, d\sigma = \int_{\mathbb{S}^2} \psi_2 G(h) \psi_1(1+h) J \, d\sigma,$$

and therefore G(h) satisfies

$$\int_{\mathbb{S}^2} \psi_1 G(h) \psi_2 \, d\mu_h = \int_{\mathbb{S}^2} \psi_2 G(h) \psi_1 \, d\mu_h, \quad d\mu_h = (1+h)\sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2} \, d\sigma, \tag{3.22}$$

for all $\psi_1, \psi_2 \in H^{\frac{1}{2}}(\mathbb{S}^2)$.

Proposition 3.2. System (1.16), (1.17) is system

$$\partial_t h = \frac{\partial_\psi \mathcal{H}(h,\psi)}{(1+h)^2}, \quad \partial_t \psi = -\frac{\partial_h \mathcal{H}(h,\psi)}{(1+h)^2}, \quad (3.23)$$

where $\partial_h \mathcal{H}, \partial_{\psi} \mathcal{H}$ are the gradients of \mathcal{H} with respect to the $L^2(\mathbb{S}^2)$ scalar product.

We provide two different proofs of Proposition 3.2. The first proof is here and uses Hadamard's formula; the second proof is in subsection 4.3, and uses the formula (4.1) of the shape derivative of the Dirichlet-Neumann operator.

First proof of Proposition 3.2. By linearity and (3.22), we have

$$\partial_{\psi} \mathcal{H}(h,\psi) = \partial_{\psi} K(h,\psi) = (1+h) J G(h) \psi, \qquad (3.24)$$

with J in (2.28). Hence (1.16) is the first equation in (3.23). To calculate $\partial_h \mathcal{H}$, we consider $\partial_h U$ and $\partial_h K$ separately. Concerning the potential energy U in (3.10), its derivative with respect to h in direction η is

$$\begin{split} \partial_h U(h)[\eta] &= \int_{\mathbb{S}^2} \eta \sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2} \, d\sigma + \int_{\mathbb{S}^2} (1+h) \frac{(1+h)\eta + \langle \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \eta \rangle}{\sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2}} \, d\sigma \\ &= \int_{\mathbb{S}^2} \eta \frac{2(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2}{\sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2}} \, d\sigma - \int_{\mathbb{S}^2} \eta \operatorname{div}_{\mathbb{S}^2} \left(\frac{(1+h)\nabla_{\mathbb{S}^2} h}{\sqrt{(1+h)^2 + |\nabla h|^2}} \right) d\sigma \\ &= \int_{\mathbb{S}^2} \eta \frac{2(1+h)^2}{\sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2}} \, d\sigma - \int_{\mathbb{S}^2} \eta (1+h) \operatorname{div}_{\mathbb{S}^2} \left(\frac{\nabla_{\mathbb{S}^2} h}{\sqrt{(1+h)^2 + |\nabla h|^2}} \right) d\sigma \\ &= \int_{\mathbb{S}^2} \eta (1+h) \operatorname{div}_{\mathbb{S}^2} \left(\frac{(1+h)x - \nabla_{\mathbb{S}^2} h}{\sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2}} \right) d\sigma, \end{split}$$

where we have used the divergence theorem (2.12) on \mathbb{S}^2 with $F = (1+h)J^{-1}h_1\nabla_{\mathbb{S}^2}h$ where J is defined in (2.28), (2.16), and div $_{\mathbb{S}^2}(x) = 2$. Hence

$$\partial_h U(h) = (1+h) \operatorname{div}_{\mathbb{S}^2} \left(\frac{(1+h)x - \nabla_{\mathbb{S}^2} h}{\sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2}} \right) = (1+h)^2 H(h)$$
(3.25)

on \mathbb{S}^2 , where H(h) is given by (2.32). To prove the second identity in (3.25), apply formula $\operatorname{div}_{\mathbb{S}^2}(\varphi F) = \varphi \operatorname{div}_{\mathbb{S}^2}F + \langle \nabla_{\mathbb{S}^2}\varphi, F \rangle$ to $\varphi = (1+h)J^{-1}$, F = x and to $\varphi = J^{-1}$, $F = \nabla_{\mathbb{S}^2}h$, where J is in (2.28), and use the fact that, for any extension of h, one has $\nabla(J^{-1}) = -J^{-2}\nabla J$ and $\nabla J = J^{-1}\{(1+h)\nabla h + (D^2h)\nabla h\}.$

To compute the derivative of $K(h, \psi)$ with respect to h in direction η , we consider $K(h + \varepsilon \eta, \psi)$, given by (3.20) with h replaced by $h + \varepsilon \eta$. To this aim, we define

$$\gamma_{\varepsilon}(x) = (1 + h(x) + \varepsilon \eta(x))x \tag{3.26}$$

for $x \in \mathbb{S}^2$, we extend $h, \eta, \gamma_{\varepsilon}$ as $\mathcal{E}_0 h, \mathcal{E}_0 \eta, \mathcal{E}_1 \gamma_{\varepsilon}$, with $\mathcal{E}_0, \mathcal{E}_1$ in (2.18), we denote $\Omega_{\varepsilon} = (\mathcal{E}_1 \gamma_{\varepsilon})(B_1)$, and we denote Φ_{ε} the solution of problem

$$\Delta \Phi_{\varepsilon} = 0 \quad \text{in } \Omega_{\varepsilon}, \qquad \Phi_{\varepsilon} = \psi \circ \gamma_{\varepsilon}^{-1} \quad \text{on } \partial \Omega_{\varepsilon}, \quad \text{i.e.,} \quad \Phi_{\varepsilon}(\gamma_{\varepsilon}(x)) = \psi(x) \quad \forall x \in \mathbb{S}^{2}.$$
(3.27)

For $\varepsilon = 0$, this is problem (1.14), and we write Φ, γ, Ω instead of $\Phi_0, \gamma_0, \Omega_0$. Denote $\dot{\Phi} = \partial_{\varepsilon}|_{\varepsilon=0} \Phi_{\varepsilon}$. Thus, by (3.20),

$$K(h + \varepsilon \eta, \psi) = \frac{1}{2} \int_{\Omega_{\varepsilon}} |\nabla \Phi_{\varepsilon}|^2 \, dx.$$
(3.28)

To differentiate (3.28) with respect to ε , we use the following formula (see, e.g., [27]).

Lemma 3.3 (Hadamard formula, or Reynolds transport theorem). Let $\Omega \subset \mathbb{R}^3$ be as in (1.6), and assume that $\beta_{\varepsilon} : \mathbb{R}^3 \to \mathbb{R}^3$ is a one-parameter family of diffeomorphisms, differentiable with respect to the parameter ε , such that $\beta_0(x) = x$, and let $\frac{d}{d\varepsilon}\Big|_{\varepsilon=0}\beta_{\varepsilon}(x) = X(x)$. Assume that a family of functions $u(\cdot, \varepsilon) : \Omega_{\varepsilon} \to \mathbb{R}$ is differentiable with respect to ε , and denote $\dot{u} = \partial_{\varepsilon}u(\cdot, 0)$. Then

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0}\int_{\Omega_{\varepsilon}}u(x,\varepsilon)\,dx=\int_{\partial\Omega}u\langle X,\nu_{\Omega}\rangle\,d\sigma+\int_{\Omega}\dot{u}\,dx.$$

To apply Lemma 3.3, we define the family of diffeomorphisms $\beta_{\varepsilon} : \mathbb{R}^3 \to \mathbb{R}^3$ by

$$\beta_{\varepsilon}(x) = \left(1 + \varepsilon \frac{\mathcal{E}_0 \eta(x)}{1 + \mathcal{E}_0 h(x)}\right) x$$

for $x \in \mathbb{R}^3 \setminus \{0\}$, and $\beta_{\varepsilon}(0) = 0$. Hence $\Omega_{\varepsilon} = \beta_{\varepsilon}(\Omega)$ and $(\mathcal{E}_1 \gamma) \circ \beta_{\varepsilon} = \beta_{\varepsilon} \circ (\mathcal{E}_1 \gamma) = \mathcal{E}_1 \gamma_{\varepsilon}$. For all $x \in \mathbb{R}^3 \setminus \{0\}$, it holds

$$X(x) = \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \beta_{\varepsilon}(x) = \frac{\mathcal{E}_0 \eta(x)}{1 + \mathcal{E}_0 h(x)} x,$$

and, at $y = \gamma(x) \in \partial \Omega$, with $x \in \mathbb{S}^2$, one has

$$X(\gamma(x)) = \frac{\mathcal{E}_0 \eta(\gamma(x))}{1 + \mathcal{E}_0 h(\gamma(x))} \gamma(x) = \mathcal{E}_0 \eta(x) x, \qquad (3.29)$$

because from y = (1 + h(x))x it follows that $\mathcal{E}_0 h(y) = \mathcal{E}_0 h(x) = h(x)$. By (3.28) and Lemma 3.3, we have

$$\partial_{h}K(h,\psi)[\eta] = \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \frac{1}{2} \int_{\Omega_{\varepsilon}} |\nabla\Phi_{\varepsilon}|^{2} dx$$

$$= \frac{1}{2} \int_{\partial\Omega} |\nabla\Phi|^{2} \langle X, \nu_{\Omega} \rangle \, d\sigma + \int_{\Omega} \langle \nabla\Phi, \nabla\dot{\Phi} \rangle \, dx$$

$$= \frac{1}{2} \int_{\partial\Omega} |\nabla\Phi|^{2} \langle X, \nu_{\Omega} \rangle \, d\sigma + \int_{\partial\Omega} \dot{\Phi} \, \langle \nabla\Phi, \nu_{\Omega} \rangle \, d\sigma, \qquad (3.30)$$

where we have integrated by parts and used that Φ is harmonic in Ω . By (3.29), (2.27) and the area formula, the change of variable $y = \gamma(x)$, $d\sigma(y) = (1+h)Jd\sigma(x)$ gives

$$\int_{\partial\Omega} |\nabla\Phi|^2 \langle X, \nu_{\Omega} \rangle \, d\sigma = \int_{\mathbb{S}^2} |\nabla\Phi(\gamma(x))|^2 (1+h(x))^2 \eta(x) \, d\sigma.$$

From the identity $\Phi_{\varepsilon}(\gamma_{\varepsilon}(x)) = \psi(x), x \in \mathbb{S}^2$, it follows that $\dot{\Phi}(\gamma(x)) + \langle (\nabla \Phi)(\gamma(x)), \eta(x)x \rangle = 0$ for $x \in \mathbb{S}^2$, namely, by (3.29),

$$\dot{\Phi}(y) = -\langle \nabla \Phi(y), X(y) \rangle$$

at $y = \gamma(x) \in \partial\Omega$, with $x \in \mathbb{S}^2$. Hence, by (1.13), (3.29) and the area formula, the last integral in (3.30) is

$$\int_{\partial\Omega} \dot{\Phi} \langle \nabla\Phi, \nu_{\Omega} \rangle \, d\sigma = -\int_{\partial\Omega} \langle \nabla\Phi, X \rangle \langle \nabla\Phi, \nu_{\Omega} \rangle \, d\sigma = -\int_{\mathbb{S}^2} \langle (\nabla\Phi)(\gamma(x)), x \rangle \eta(1+h) JG(h) \psi \, d\sigma.$$

Thus, by (3.30),

$$\partial_h K(h,\psi) = \frac{|(\nabla\Phi) \circ \gamma|^2 (1+h)^2}{2} - \langle (\nabla\Phi) \circ \gamma, x \rangle (1+h) J G(h) \psi.$$
(3.31)

By (3.4), (2.27), (3.6), we have

$$\langle (\nabla \Phi) \circ \gamma, x \rangle = \langle (\nabla_{\partial \Omega} \Phi) \circ \gamma, x \rangle + \langle (\nabla_{\nu} \Phi) \circ \gamma, x \rangle = \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle}{J^2} + \frac{1+h}{J} G(h) \psi$$

on S^2 , and, by (3.4) and (3.7),

$$|(\nabla\Phi)\circ\gamma|^2 = |(\nabla_{\partial\Omega}\Phi)\circ\gamma|^2 + |(\nabla_{\nu}\Phi)\circ\gamma|^2 = \left(G(h)\psi\right)^2 + \frac{|\nabla_{\mathbb{S}^2}\psi|^2}{(1+h)^2} - \frac{\langle\nabla_{\mathbb{S}^2}\psi,\nabla_{\mathbb{S}^2}h\rangle^2}{(1+h)^2J^2}$$

on \mathbb{S}^2 . Using also (3.25), it follows that (1.17) is the second equation in (3.23).

Now we show that, with a simple change of variable, the factor $(1+h)^{-2}$ can be removed from (3.23), so that we obtain a Hamiltonian system written in Darboux coordinates.

Lemma 3.4. Consider a change of variable of the form

$$h = f(\eta), \quad \psi = g(\eta)\varpi, \tag{3.32}$$

where f, g are real-valued functions of one real variable, f invertible, g never vanishing, with

$$(1+f(\eta))^2 f'(\eta)g(\eta) = 1.$$
(3.33)

Then system (3.23) is transformed into the Hamiltonian system

$$\begin{cases} \partial_t \eta = \partial_{\varpi} \mathcal{H}_1(\eta, \varpi), \\ \partial_t \varpi = -\partial_{\eta} \mathcal{H}_1(\eta, \varpi), \end{cases}$$
(3.34)

where

$$\mathcal{H}_1(\eta, \varpi) = \mathcal{H}(f(\eta), g(\eta)\varpi). \tag{3.35}$$

Proof. Differentiating (3.35) gives

$$\partial_{\eta}\mathcal{H}_{1}(\eta,\varpi) = \partial_{h}\mathcal{H}(h,\psi)f'(\eta) + \partial_{\psi}\mathcal{H}(h,\psi)g'(\eta)\varpi, \quad \partial_{\varpi}\mathcal{H}_{1}(\eta,\varpi) = \partial_{\psi}\mathcal{H}(h,\psi)g(\eta),$$

where $(h, \psi) = (f(\eta), g(\eta)\varpi)$, whence

$$\partial_h \mathcal{H}(h,\psi) = \frac{\partial_\eta \mathcal{H}_1(\eta,\varpi)}{f'(\eta)} - \frac{\partial_\varpi \mathcal{H}_1(\eta,\varpi)}{g(\eta)f'(\eta)}g'(\eta)\varpi, \quad \partial_\psi \mathcal{H}(h,\psi) = \frac{\partial_\varpi \mathcal{H}_1(\eta,\varpi)}{g(\eta)}.$$

Also,

$$\partial_t h = f'(\eta)\partial_t \eta, \quad \partial_t \psi = g'(\eta)\varpi\partial_t \eta + g(\eta)\partial_t \varpi.$$

Hence (3.23) becomes

$$\partial_t \eta = \frac{\partial_{\varpi} \mathcal{H}_1(\eta, \varpi)}{a(\eta)}, \quad \partial_t \varpi = -\frac{\partial_{\eta} \mathcal{H}_1(\eta, \varpi)}{a(\eta)}, \quad \text{where } a(\eta) = (1 + f(\eta))^2 f'(\eta) g(\eta),$$

and this is (3.34) if f, g satisfy (3.33).

Special cases of transformations (3.32) satisfying (3.33) are

(i) $f(\eta) = \eta$, $g(\eta) = (1+\eta)^{-2}$, which is the change of variable $\psi = \varpi/(1+h)^2$ with h unchanged;

(*ii*)
$$f(\eta) = (1+3\eta)^{\frac{1}{3}} - 1$$
, $g(\eta) = 1$, which is $(1+h)^3 = (1+3\eta)$, with ψ unchanged.

The transformation (i) offers the convenience of not having to change h in the Dirichlet-Neumann operator G(h) and in the mean curvature H(h). The transformation (ii) also has some advantage, because, in that case, the conservation of the total fluid mass becomes a zero average condition for the new elevation function η . This nice feature of (ii), however, only concerns the mass conservation, because the conservation of the barycenter velocity becomes a condition involving $(1 + 3\eta)^{\frac{4}{3}}$, which does not seem to be better than $(1 + h)^4$.

4 Shape derivative of the Dirichlet-Neumann operator

The Dirichlet-Neumann operator $G(h)\psi$ defined in (1.13) is linear in ψ and, as we prove below (Theorem 5.19), it depends analytically on h in suitable Sobolev spaces. Hence $G(h)\psi$ is differentiable with respect to h; in this section we prove the following formula for its derivative.

Theorem 4.1. There exists $\delta_0 > 0$ such that, for $h, \eta \in H^3(\mathbb{S}^2)$, $\psi \in H^{\frac{5}{2}}(\mathbb{S}^2)$, $||h||_{H^3(\mathbb{S}^2)} < \delta_0$, the Fréchet derivative of $G(h)\psi$ with respect to h in direction η is

$$G'(h)[\eta]\psi = b\eta + \langle B, \nabla_{\mathbb{S}^2}\eta \rangle - G(h)(W\eta), \tag{4.1}$$

where

$$W = \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle}{J^2} + \frac{(1+h)G(h)\psi}{J}, \quad B = \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle \nabla_{\mathbb{S}^2} h}{(1+h)J^3} - \frac{\nabla_{\mathbb{S}^2} \psi}{(1+h)J}, \quad (4.2)$$

$$b = \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle}{J^3} - \frac{2G(h)\psi}{1+h} - \frac{\operatorname{div}_{\mathbb{S}^2} \{(1+h)(\nabla_{\mathbb{S}^2} \psi - W\nabla_{\mathbb{S}^2} h)\}}{(1+h)^2 J},$$
(4.3)

and J is in (2.28).

We give two independent proofs of Theorem 4.1. The first proof, in subsection 4.1, follows the method of the "good unknown of Alinhac"; the second proof, in subsection 4.2, is based on a geometric argument.

4.1 Proof by the method of the good unknown of Alinhac

In this section we prove Theorem 4.1 by adapting the approach of Alazard, Métivier, and Lannes to the nearly spherical geometry. We follow, as long as possible, the proof in Lannes' book [30]; however, the adaptation of that technique to the sphere has some delicate aspects; we postpone to Remark 4.8 a technical explanation of that (surprising!) difficulty.

Let $h \in W^{1,\infty}(\mathbb{S}^2)$, with $||h||_{W^{1,\infty}(\mathbb{S}^2)} \leq 1/2$, and let $\psi \in H^{\frac{1}{2}}(\mathbb{S}^2)$. Then problem (1.14) has a unique weak solution $\Phi \in H^1(\Omega)$, problem (3.13) has a unique weak solution $\tilde{\Phi} \in H^1(B_1)$, and the two problems are equivalent, as $\Phi \in H^1(\Omega)$ is the weak solution of (1.14) if and only if $\tilde{\Phi} \in H^1(B_1)$ in (3.11) is the weak solution of (3.13). The Dirichlet-Neumann $G(h)\psi$ is defined in (1.13), and it satisfies (3.19), with P in (3.15) and J in (2.28). Inserting the formula (3.15) into (3.19), using (2.28), (2.16), and the identity $\langle \nabla \tilde{\Phi}, \nabla_{\mathbb{S}^2} h \rangle = \langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle$, we obtain

$$G(h)\psi = \frac{J\langle \nabla \dot{\Phi}, x \rangle}{(1+h)^2} - \frac{\langle \nabla_{\mathbb{S}^2}\psi, \nabla_{\mathbb{S}^2}h \rangle}{(1+h)J} \quad \text{on } \mathbb{S}^2.$$

$$(4.4)$$

To compute the derivative of $G(h)\psi$ with respect to h in direction η using formula (4.4), we have to study the derivative of $\tilde{\Phi}$ with respect to h in direction η . Let $\tilde{\Phi}_{\varepsilon}$ be the weak solution of problem

$$\tilde{\Phi}_{\varepsilon} \in H^1(B_1), \quad \operatorname{div}\left(P_{\varepsilon}\nabla\tilde{\Phi}_{\varepsilon}\right) = 0 \quad \text{in } B_1, \quad \tilde{\Phi}_{\varepsilon} = \psi \quad \text{on } \mathbb{S}^2,$$

$$(4.5)$$

where P_{ε} is the matrix we obtain by replacing h with $h + \varepsilon \eta$ in (3.15). Given $\psi \in H^{\frac{1}{2}}(\mathbb{S}^2)$, the map $h \mapsto \tilde{\Phi}$ is analytic from the ball (5.12) into $H^1(B_1)$, see Proposition 5.2 below. Hence problem (4.5) can be differentiated with respect to ε , and, at $\varepsilon = 0$, we obtain

$$f_1 \in H^1(B_1), \quad \operatorname{div}(P_1 \nabla f_0) + \operatorname{div}(P_0 \nabla f_1) = 0 \quad \operatorname{in} B_1, \quad f_1 = 0 \quad \operatorname{on} \mathbb{S}^2,$$
 (4.6)

where, to shorten the notation, we denote

$$P_0 := P, \quad P_1 := \partial_{\varepsilon} P_{\varepsilon}|_{\varepsilon=0} = P'(h)[\eta], \quad f_0 := \tilde{\Phi}, \quad f_1 := \partial_{\varepsilon} \tilde{\Phi}_{\varepsilon}|_{\varepsilon=0} = \tilde{\Phi}'(h)[\eta].$$
(4.7)

Differentiating (3.15) with respect to h in direction η we get

$$P_1 = (\mathcal{E}_0\eta)I - (\nabla \mathcal{E}_0\eta) \otimes x - x \otimes \nabla \mathcal{E}_0\eta + \left(\frac{2\langle \nabla \mathcal{E}_0h, \nabla \mathcal{E}_0\eta \rangle}{1 + \mathcal{E}_0h} - \frac{|\nabla \mathcal{E}_0h|^2 \mathcal{E}_0\eta}{(1 + \mathcal{E}_0h)^2}\right)x \otimes x$$
(4.8)

in $\mathbb{R}^3 \setminus \{0\}$. To adapt the method of the "good unknown of Alinhac" in [30] to the unit ball B_1 , we replace the vertical partial derivative ∂_z of the flat case with the radial derivative operator

$$D_x := \langle x, \nabla \rangle, \tag{4.9}$$

and we look for a scalar function α defined in $B_1^* := B_1 \setminus \{0\}$ such that

$$\operatorname{div}\left(P_1 \nabla f_0\right) = \operatorname{div}\left(P_0 \nabla(\alpha D_x f_0)\right) \quad \text{in } B_1^*.$$
(4.10)

Identity (4.10) implies that also the term div $(P_1 \nabla f_0)$ appearing in (4.6) can be expressed as the operator div $(P_0 \nabla \cdot)$ applied to a scalar function; as a consequence, the second item of (4.6) becomes an identity of the form div $(P_0 \nabla v) = 0$ where v is a scalar function, and this is the identity one has in the definition of the Dirichlet-Neumann operator. We prove that equation (4.10) has an explicit solution, given by the following lemma.

Lemma 4.2. Let $h, \psi, \eta, P_0, P_1, f_0, f_1$ be as above. Then the function

$$\alpha = -\frac{\mathcal{E}_0 \eta}{1 + \mathcal{E}_0 h} \quad in \ \mathbb{R}^3 \setminus \{0\}$$

$$\tag{4.11}$$

solves equation (4.10).

Before starting with the proof of Lemma 4.2, we recall the following basic commutator rules, whose proof is elementary.

Lemma 4.3. One has

(i)
$$\nabla(D_x\varphi) = D_x(\nabla\varphi) + \nabla\varphi,$$
 (ii) div $(D_xg) = D_x(\operatorname{div} g) + \operatorname{div} g_y$
(iii) div $(\varphi g) = \varphi \operatorname{div} g + \langle \nabla\varphi, g \rangle,$ (iv) div $(x\varphi) = 3\varphi + D_x\varphi$

for all scalar functions φ and all vector-valued functions g.

In the next lemma we study div $(P_0 \nabla(\alpha D_x f_0))$ for any scalar function α , using the fact that f_0 solves the elliptic equation in (3.13), i.e.,

div
$$(P_0 \nabla f_0) = 0$$
 in B_1^* , (4.12)

and the identity

$$D_x P_0 = 0 \quad \text{in } \mathbb{R}^3 \setminus \{0\},\tag{4.13}$$

where $D_x P_0$ is the matrix obtained by applying D_x to each entry of P_0 . Identity (4.13) holds because P_0 is homogeneous of degree 0: $\mathcal{E}_0 h$ has degree 0, its gradient $\nabla(\mathcal{E}_0 h)$ has degree -1, the identity map x has degree 1, and each term in (3.15) has degree 0.

Lemma 4.4. Let P_0, f_0 be as above, and let α be any scalar function defined in $\mathbb{R}^3 \setminus \{0\}$. Then

$$\operatorname{div}\left(P_0\nabla(\alpha D_x f_0)\right) = \operatorname{div}\left(P\nabla f_0\right) \quad in \ B_1^*,\tag{4.14}$$

where

$$\tilde{P} = P_0(\nabla \alpha) \otimes x + (x \otimes \nabla \alpha)P_0 - (\alpha + D_x \alpha)P_0 \quad in \ \mathbb{R}^3 \setminus \{0\}.$$
(4.15)

Proof. By the formula of the gradient of a product of two functions, one has

$$\operatorname{div}\left(P_0\nabla(\alpha D_x f_0)\right) = \operatorname{div}\left(P_0(\nabla \alpha) D_x f_0\right) + \operatorname{div}\left(P_0\alpha \nabla D_x f_0\right).$$
(4.16)

Applying Lemma 4.3(iii) to the last term of (4.16) gives

$$(4.16) = \operatorname{div} \left(P_0(\nabla \alpha) D_x f_0 \right) + \alpha \operatorname{div} \left(P_0 \nabla D_x f_0 \right) + \langle \nabla \alpha, P_0 \nabla D_x f_0 \rangle.$$

$$(4.17)$$

The second term in the RHS of (4.17) vanishes, because

$$\operatorname{div}\left(P_0\nabla D_x f_0\right) = 0. \tag{4.18}$$

Let us prove (4.18). Applying D_x to identity (4.12), and using 4.3(*ii*) and (4.12), one has

$$0 = D_x \operatorname{div} (P_0 \nabla f_0) = \operatorname{div} (D_x (P_0 \nabla f_0)) - \operatorname{div} (P_0 \nabla f_0) = \operatorname{div} (D_x (P_0 \nabla f_0)).$$
(4.19)

Using the formula of the gradient of a product, then using (4.13), and then Lemma 4.3(i), one has

$$D_x(P_0\nabla f_0) = (D_x P_0)\nabla f_0 + P_0 D_x \nabla f_0 = P_0 D_x \nabla f_0 = P_0 \nabla D_x f_0 - P_0 \nabla f_0.$$
(4.20)

By (4.19), the divergence of (4.20) is

$$0 = \operatorname{div}\left(D_x(P_0 \nabla f_0)\right) = \operatorname{div}\left(P_0 \nabla D_x f_0\right) - \operatorname{div}\left(P_0 \nabla f_0\right).$$
(4.21)

By (4.12), the last term in (4.21) is zero, and this proves (4.18).

Concerning the last term in (4.17), we observe that, by (4.20),

$$P_0 \nabla D_x f_0 = D_x (P_0 \nabla f_0) + P_0 \nabla f_0.$$
(4.22)

Hence, by (4.18) and (4.22), identity (4.17) becomes

$$(4.16) = \operatorname{div} \left(P_0(\nabla \alpha) D_x f_0 \right) + \langle \nabla \alpha, D_x(P_0 \nabla f_0) \rangle + \langle \nabla \alpha, P_0 \nabla f_0 \rangle.$$

$$(4.23)$$

To study the second term in the RHS of (4.23), we observe that

$$D_x\{\langle \nabla \alpha, P_0 \nabla f_0 \rangle\} = \langle D_x \nabla \alpha, P_0 \nabla f_0 \rangle + \langle \nabla \alpha, D_x (P_0 \nabla f_0) \rangle, \tag{4.24}$$

and therefore, using (4.24) to substitute the second term in the RHS of (4.23), we get

$$(4.16) = \operatorname{div}\left(P_0(\nabla\alpha)D_x f_0\right) + D_x\{\langle\nabla\alpha, P_0\nabla f_0\rangle\} - \langle D_x\nabla\alpha, P_0\nabla f_0\rangle + \langle\nabla\alpha, P_0\nabla f_0\rangle.$$
(4.25)

To study the third term in the RHS of (4.25), we note that, by Lemma 4.3(iii), (4.12) and Lemma 4.3(i), we have

$$\operatorname{div}\left((D_x \alpha) P_0 \nabla f_0\right) = (D_x \alpha) \operatorname{div}\left(P_0 \nabla f_0\right) + \langle \nabla D_x \alpha, P_0 \nabla f_0 \rangle$$
$$= \langle \nabla D_x \alpha, P_0 \nabla f_0 \rangle$$
$$= \langle D_x \nabla \alpha, P_0 \nabla f_0 \rangle + \langle \nabla \alpha, P_0 \nabla f_0 \rangle.$$
(4.26)

Using (4.26) to substitute the third term in the RHS of (4.25), we obtain

$$(4.16) = \operatorname{div}\left(P_0(\nabla\alpha)D_x f_0\right) + D_x\{\langle\nabla\alpha, P_0\nabla f_0\rangle\} - \operatorname{div}\left((D_x\alpha)P_0\nabla f_0\right) + 2\langle\nabla\alpha, P_0\nabla f_0\rangle.$$
(4.27)

To study the second term in the RHS of (4.27), we observe that identity Lemma 4.3(*iv*) applied to the scalar function $\langle \nabla \alpha, P_0 \nabla f_0 \rangle$ gives

$$\operatorname{div}\left(x\langle\nabla\alpha, P_0\nabla f_0\rangle\right) = 3\langle\nabla\alpha, P_0\nabla f_0\rangle + D_x\{\langle\nabla\alpha, P_0\nabla f_0\rangle\}.$$
(4.28)

Using (4.28) to substitute the second term in the RHS of (4.27), we get

$$(4.16) = \operatorname{div}\left(P_0(\nabla\alpha)D_x f_0\right) + \operatorname{div}\left(x\langle\nabla\alpha, P_0\nabla f_0\rangle\right) - \operatorname{div}\left((D_x\alpha)P_0\nabla f_0\right) - \langle\nabla\alpha, P_0\nabla f_0\rangle.$$
(4.29)

Concerning the last term in (4.29), we note that, by Lemma 4.3(iii) and (4.12), one has

$$\operatorname{div}\left(\alpha P_0 \nabla f_0\right) = \alpha \operatorname{div}\left(P_0 \nabla f_0\right) + \langle \nabla \alpha, P_0 \nabla f_0 \rangle = \langle \nabla \alpha, P_0 \nabla f_0 \rangle.$$
(4.30)

Thus all the terms in (4.29) are in divergence form. Moreover, the third term in the RHS of (4.29) is of the form div $(M\nabla f_0)$ for some 3×3 matrix M. The first term in the RHS of (4.29) is also of the same form because $(\nabla \alpha)D_x f_0 = (\nabla \alpha)\langle x, \nabla f_0 \rangle = (\nabla \alpha)x^T \nabla f_0 = ((\nabla \alpha) \otimes x) \nabla f_0$. Since $\langle \nabla \alpha, P_0 \nabla f_0 \rangle = (\nabla \alpha)^T P_0 \nabla f_0$, the second term in the RHS of (4.29) is of the same form too (x is a column, and $(\nabla \alpha)^T$ is a row). By (4.30), the last term in (4.29) is also of the form div $(M \nabla f_0)$. The sum of all the terms in (4.29) gives (4.14) with \tilde{P} given by (4.15).

By (3.15), the matrix \tilde{P} in (4.15) can be explicitly calculated in terms of h and α . We get

$$\tilde{P} = -(\alpha + D_x \alpha)(1 + \mathcal{E}_0 h)I + \alpha \left[(\nabla \mathcal{E}_0 h) \otimes x + x \otimes \nabla \mathcal{E}_0 h \right] + (1 + \mathcal{E}_0 h) \left[(\nabla \alpha) \otimes x + x \otimes \nabla \alpha \right] \\ + \left(\frac{|\nabla \mathcal{E}_0 h|^2}{1 + \mathcal{E}_0 h} (D_x \alpha - \alpha) - 2 \langle \nabla \mathcal{E}_0 h, \nabla \alpha \rangle \right) x \otimes x \quad \text{in } \mathbb{R}^3 \setminus \{0\}.$$

$$(4.31)$$

Lemma 4.4 and formula (4.31) hold for any scalar function α . When α is the function in (4.11), one has Lemma 4.2, which can now be proved.

Proof of Lemma 4.2. Let α be the function in (4.11). Then a straightforward calculation shows that the matrices \tilde{P} in (4.31) and P_1 in (4.8) coincide. Hence (4.10) follows from Lemma 4.4.

Remark 4.5. Formula (4.11) for α such that $\tilde{P} = P_1$ is suggested, for example, by a comparison of the coefficients of the identity matrix in (4.31) and in (4.8): the coefficients are equal if $-(\alpha + D_x \alpha)(1 + \mathcal{E}_0 h) = \mathcal{E}_0 \eta$ in $\mathbb{R}^3 \setminus \{0\}$. The given functions $\mathcal{E}_0 h$ and $\mathcal{E}_0 \eta$ are homogeneous of degree 0; this suggests to look for α in the same class. For α homogeneous of degree 0, one has $D_x \alpha = 0$, and formula (4.11) for α follows immediately.

Now we use Lemma 4.2 to calculate the shape derivative $G'(h)[\eta]\psi$. Using (4.10) to replace the term div $(P_1 \nabla f_0)$ with div $(P_0 \nabla (\alpha D_x f_0))$ in (4.6), one obtains

$$\operatorname{div}(P_0 \nabla w) = 0 \quad \text{in } B_1^*, \qquad w := f_1 + \alpha D_x f_0 \quad \text{in } B_1^*, \tag{4.32}$$

where α is the function in (4.11). Since the elliptic problems of the form (3.13) have uniqueness of solution in $H^1(B_1)$, we want w to belong to $H^1(B_1)$. We already know that $f_1 \in H^1(B_1)$, see (4.6). Concerning $\alpha D_x f_0$, we have the following result. **Lemma 4.6.** Let $h, \eta \in H^3(\mathbb{S}^2)$, $\psi \in H^{\frac{3}{2}}(\mathbb{S}^2)$, and let $||h||_{H^3(\mathbb{S}^2)} < \delta_0$, where δ_0 is the constant in Proposition 5.16 below. Then $\alpha D_x f_0 \in H^1(B_1)$.

Proof. The function $f_0 = \tilde{\Phi} \in H^1(B_1)$ in (4.7) is the weak solution of (3.13). Hence the function Φ satisfying (3.11) is in $H^1(\Omega)$, and it is the weak solution of (1.14). Thus Φ is harmonic in Ω , and therefore $\Phi \in C^{\infty}(\Omega)$. We consider the partition of unity (5.19), and we study $\alpha D_x f_0$ in the ball $\bar{B}_{\rho} = \{x \in \mathbb{R}^3 : |x| \leq \rho\}, \ \rho = 1 - (\delta/2)$, and in the annulus $\mathcal{A} = B_1 \setminus \bar{B}_{\rho}$, where δ is the constant in (5.19).

Let $K = (\mathcal{E}_1 \gamma)(B_\rho)$ (we put $\mathcal{E}_1 \gamma(0) = 0$). Then K is a compact subset of Ω , and $\Phi, \nabla \Phi, D^2 \Phi$ are bounded on K. From (4.11), (4.9), (3.12), (2.19), and $\langle \nabla(\mathcal{E}_0 h), x \rangle = 0$ it follows that

$$\alpha D_x f_0 = -(\mathcal{E}_0 \eta) \langle x, (\nabla \Phi) \circ (\mathcal{E}_1 \gamma) \rangle$$

in $B_1^* = B_1 \setminus \{0\}$. Its gradient is

$$\nabla(\alpha D_x f_0) = -\nabla(\mathcal{E}_0 \eta) \langle x, (\nabla \Phi) \circ (\mathcal{E}_1 \gamma) \rangle - (\mathcal{E}_0 \eta) \{ (\nabla \Phi) \circ (\mathcal{E}_1 \gamma) + [D(\mathcal{E}_1 \gamma)]^T [(D^2 \Phi) \circ (\mathcal{E}_1 \gamma)] x \}$$

in B_1^* , and it belongs to $L^{\infty}(B_{\rho}) \subset L^2(B_{\rho})$ because, by homogeneity, in B_1^* one has

$$|\nabla(\mathcal{E}_0\eta)(x)||x| \le \|\nabla_{\mathbb{S}^2}\eta\|_{L^{\infty}(\mathbb{S}^2)}, \quad |\mathcal{E}_0\eta| \le \|\eta\|_{L^{\infty}(\mathbb{S}^2)}, \quad |D(\mathcal{E}_1\gamma)| \le 1 + \|h\|_{W^{1,\infty}(\mathbb{S}^2)},$$

while

$$|(\nabla\Phi)\circ(\mathcal{E}_1\gamma)| \le \|\nabla\Phi\|_{L^{\infty}(K)}, \quad |(D^2\Phi)\circ(\mathcal{E}_1\gamma)| \le \|D^2\Phi\|_{L^{\infty}(K)}$$

in B_{ρ} . Note that $\|h\|_{W^{1,\infty}(\mathbb{S}^2)} \leq C \|h\|_{H^3(\mathbb{S}^2)}$, see (5.55), (5.56).

By Proposition 5.16, the solution $f_0 = \tilde{\Phi}$ of the elliptic problem (3.13) satisfies $||f_0||_{X_0^{2,0}} \leq C ||\psi||_{H^{\frac{3}{2}}(\mathbb{S}^2)} < \infty$, with $|| \cdot ||_{X_0^{2,0}}$ defined in (5.61); note that the radial cut-off ζ_0 appearing in (5.61) satisfies $\zeta_0 = 1$ in the annulus \mathcal{A} , see definition (5.57). This implies that $f_0 \in H^2(\mathcal{A})$ and therefore $\nabla(\alpha D_x f_0) \in L^2(\mathcal{A})$.

By (4.6), $f_1 \in H^1(B_1)$, and, by Lemma 4.6, $\alpha D_x f_0 \in H^1(B_1)$. Hence w in (4.32) is in $H^1(B_1)$. By (4.6), $f_1 = 0$ on \mathbb{S}^2 , and, by (4.11), $\alpha = -\eta/(1+h)$ on \mathbb{S}^2 , while, by (4.4), $D_x f_0 = \langle x, \nabla \tilde{\Phi} \rangle$ on \mathbb{S}^2 is

$$D_x f_0 = \frac{(1+h)^2 G(h)\psi}{J} + \frac{(1+h)\langle \nabla_{\mathbb{S}^2}\psi, \nabla_{\mathbb{S}^2}h\rangle}{J^2} = (1+h)W \text{ on } \mathbb{S}^2,$$
(4.33)

where W is defined in (4.2). Therefore w in (4.32) is $w = -\eta W$ on \mathbb{S}^2 . Thus w is the weak solution of the problem

$$w \in H^{1}(B_{1}), \quad \operatorname{div}(P_{0}\nabla w) = 0 \text{ in } B_{1}, \quad w = -\eta W \text{ on } \mathbb{S}^{2}.$$
 (4.34)

By (4.34), formula (3.19) with $(-\eta W)$ in the role of ψ and w in that of $\tilde{\Phi}$ gives

$$-G(h)(\eta W) = Z\langle P_0 \nabla w, x \rangle \quad \text{on } \mathbb{S}^2, \quad Z := Z(h) = \frac{1}{(1+h)J}, \tag{4.35}$$

with J in (2.28).

Lemma 4.7. Let $h, \eta \in H^3(\mathbb{S}^2)$, $\psi \in H^{\frac{5}{2}}(\mathbb{S}^2)$, and let $||h||_{H^3(\mathbb{S}^2)} < \delta_0$, where δ_0 is the constant in Proposition 5.16 below. Then

(i) $\eta W \in H^{\frac{3}{2}}(\mathbb{S}^2)$,

(*ii*) $w, f_1, D_x f_0 \in H^2(\mathcal{A})$ in some annulus $\mathcal{A} = \{x \in \mathbb{R}^3 : c < |x| < 1\}, c \in (0, 1),$

(iii) the trace at \mathbb{S}^2 of the gradients $(\nabla w)|_{\mathbb{S}^2}, (\nabla f_1)|_{\mathbb{S}^2} (\nabla D_x f_0)|_{\mathbb{S}^2} \in H^{\frac{1}{2}}(\mathbb{S}^2)$ is well-defined.

Proof. For $\|h\|_{H^3(\mathbb{S}^2)} < \delta_0$, $\psi \in H^{\frac{5}{2}}(\mathbb{S}^2)$ and $\eta \in H^{\frac{3}{2}}(\mathbb{S}^2)$, one has $J \in H^{\frac{3}{2}}(\mathbb{S}^2)$ by Lemma 5.18, $G(h)\psi \in H^{\frac{3}{2}}(\mathbb{S}^2)$ by Theorem 5.19, and $W, \eta W \in H^{\frac{3}{2}}(\mathbb{S}^2)$ by (4.2), (5.53), (5.55). As a consequence, by Proposition 5.16, the solution w of the elliptic problem (4.34) satisfies $\|w\|_{X_0^{2,0}} \leq C\|\eta W\|_{H^{\frac{3}{2}}(\mathbb{S}^2)} < \infty$, with $\|\cdot\|_{X_0^{2,0}}$ defined in (5.61). This implies that $w \in H^2(\mathcal{A})$ for some annulus

 $\mathcal{A}, \nabla w \in H^1(\mathcal{A})$, and the trace $(\nabla w)|_{\mathbb{S}^2} \in H^{\frac{1}{2}}(\mathbb{S}^2)$ is well-defined. By Proposition 5.16, taking also η in $H^3(\mathbb{S}^2)$ (so that $h + \varepsilon \eta$ is in the required ball for ε small enough), recalling the definition (4.7) of f_1 , we have that $\|f_1\|_{X_0^{2,0}}$ is finite, and therefore $f_1 \in H^2(\mathcal{A})$ and $(\nabla f_1)|_{\mathbb{S}^2} \in H^{\frac{1}{2}}(\mathbb{S}^2)$. Since $\alpha D_x f_0 = w - f_1$, by triangular inequality $\|\alpha D_x f_0\|_{X_0^{2,0}}$ is also finite. For $\eta = -(1+h)$, one has $\alpha = 1$, whence $D_x f_0 \in H^2(\mathcal{A})$ and $(\nabla D_x f_0)|_{\mathbb{S}^2} \in H^{\frac{1}{2}}(\mathbb{S}^2)$.

Lemma 4.7 implies that any identity in B_1^* involving $\nabla D_x f_0, \nabla w, \nabla f_1$ also holds on \mathbb{S}^2 by taking the trace of the involved functions.

To calculate the derivative of $G(h)\psi$ with respect to h in direction η , we differentiate identity (3.19). Recalling the definition of P_0, P_1, f_0, f_1, Z in (4.7), (4.35), one has

$$G'(h)[\eta]\psi = \underbrace{Z'(h)[\eta]\langle P_0\nabla f_0, x\rangle}_{E_1} + \underbrace{Z\langle P_1\nabla f_0, x\rangle}_{E_2} + \underbrace{Z\langle P_0\nabla f_1, x\rangle}_{E_3} \quad \text{on } \mathbb{S}^2.$$
(4.36)

Calculation of E_1 . By the definition (4.35) of Z, we have

$$Z'(h)[\eta] = -\frac{\{2(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2\}\eta + (1+h)\langle \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \eta \rangle}{(1+h)^2 J^3}$$

on S², and, by (3.19), $\langle P_0 \nabla f_0, x \rangle = (1+h) J G(h) \psi$. Hence E_1 in (4.36) is

$$E_1 = -\frac{\{2(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2\}G(h)\psi}{(1+h)J^2} \eta - \frac{G(h)\psi}{J^2} \langle \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \eta \rangle.$$
(4.37)

Calculation of E_2 . By formula (4.8), recalling notation (4.9), we calculate

$$\langle P_1 \nabla f_0, x \rangle = \eta D_x f_0 - \langle \nabla_{\mathbb{S}^2} \eta, \nabla f_0 \rangle + \left(\frac{2 \langle \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \eta \rangle}{1+h} - \frac{|\nabla_{\mathbb{S}^2} h|^2 \eta}{(1+h)^2} \right) D_x f_0 \tag{4.38}$$

on \mathbb{S}^2 . Now $D_x f_0$ on \mathbb{S}^2 is given by (4.33), and $\langle \nabla_{\mathbb{S}^2} \eta, \nabla f_0 \rangle = \langle \nabla_{\mathbb{S}^2} \eta, \nabla_{\mathbb{S}^2} \psi \rangle$ on \mathbb{S}^2 because $f_0 = \psi$ on \mathbb{S}^2 . Hence E_2 in (4.36) is

$$E_2 = \frac{W}{J} \left(1 - \frac{|\nabla_{\mathbb{S}^2} h|^2}{(1+h)^2} \right) \eta - \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} \eta \rangle}{(1+h)J} + \frac{2W \langle \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \eta \rangle}{(1+h)J}.$$
(4.39)

Calculation of E_3 . To calculate the term E_3 in (4.36), we use the definition (4.32) of w to write f_1 as the difference $f_1 = w - \alpha D_x f_0$ in B_1^* . Thus,

$$\langle P_0 \nabla f_1, x \rangle = \langle P_0 \nabla w, x \rangle - \langle P_0 \nabla (\alpha D_x f_0), x \rangle$$
(4.40)

in B_1^* , and therefore, by the discussion following Lemma 4.7, also on S². Hence, by (4.35) and (4.40), the term E_3 in (4.36) is

$$E_3 = -G(h)(\eta W) - Z\langle P_0 \nabla(\alpha D_x f_0), x \rangle$$
(4.41)

on $\mathbb{S}^2.$ Since α and D_xf_0 are scalar functions, one has

$$\langle P_0 \nabla(\alpha D_x f_0), x \rangle = (D_x f_0) \langle P_0 \nabla \alpha, x \rangle + \alpha \langle P_0 \nabla D_x f_0, x \rangle$$
(4.42)

in B_1^* , and therefore on \mathbb{S}^2 . Hence, by (4.41) and (4.42),

$$E_3 = -G(h)(\eta W) - \underbrace{Z(D_x f_0) \langle P_0 \nabla \alpha, x \rangle}_{E_4} - \underbrace{Z\alpha \langle P_0 \nabla D_x f_0, x \rangle}_{E_5}$$
(4.43)

on \mathbb{S}^2 . To study E_4 , by (3.15), (4.11), we calculate

$$\langle P_0 \nabla \alpha, x \rangle = \frac{\langle \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \eta \rangle}{1+h} - \frac{|\nabla_{\mathbb{S}^2} h|^2 \eta}{(1+h)^2}$$

on \mathbb{S}^2 , while $D_x f_0$ and Z on \mathbb{S}^2 are in (4.33), (4.35). Therefore E_4 in (4.43) is

$$E_4 = \frac{W\langle \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \eta \rangle}{(1+h)J} - \frac{|\nabla_{\mathbb{S}^2} h|^2 W\eta}{(1+h)^2 J}.$$
(4.44)

The term E_5 in (4.43) contains derivatives of f_0 of second order; we use the elliptic equation (4.12) to express them in terms of h, ψ . Denoting $g = P_0 \nabla f_0$, and g_j its *j*-th component, by (4.12) one has

$$0 = \operatorname{div} g(y) = \sum_{j} \langle \nabla g_j(y), e_j \rangle = \sum_{j} \langle \Pi_{T_x(\mathbb{S}^2)} [\nabla g_j(y)], e_j \rangle + \sum_{j} \langle \nabla g_j(y), x \rangle \langle x, e_j \rangle$$

for $y \in B_1^*$, $x \in \mathbb{S}^2$. Taking the trace at the sphere, i.e., $y \in \mathbb{S}^2$, and recalling (2.8), we find

$$0 = \operatorname{div} \left(P_0 \nabla f_0 \right) = \operatorname{div}_{\mathbb{S}^2} \left(P_0 \nabla f_0 \right) + \langle x, D_x (P_0 \nabla f_0) \rangle$$
(4.45)

on \mathbb{S}^2 . We recall that the tangential divergence div $_{\mathbb{S}^2}(P_0 \nabla f_0)$ depends only on the restriction of $P_0 \nabla f_0$ to \mathbb{S}^2 , which now we calculate. By (4.33), and because $f_0 = \psi$ on \mathbb{S}^2 , one has

$$\nabla f_0 = \nabla_{\mathbb{S}^2} f_0 + \langle \nabla f_0, x \rangle x = \nabla_{\mathbb{S}^2} f_0 + (D_x f_0) x = \nabla_{\mathbb{S}^2} \psi + W(1+h) x \tag{4.46}$$

on \mathbb{S}^2 . By (4.46), using formulas (3.15), (4.2), (2.28) of P_0, W, J , we obtain

$$(P_0 \nabla f_0)|_{\mathbb{S}^2} = (1+h)(\nabla_{\mathbb{S}^2} \psi - W \nabla_{\mathbb{S}^2} h) + (1+h)J(G(h)\psi) x$$
(4.47)

on \mathbb{S}^2 (note that the *x* component in (4.47) is also given by (3.19)). Now we consider the scalar product $\langle x, D_x(P_0 \nabla f_0) \rangle$ in B_1^* . By (4.13) and Lemma 4.3(*i*), one has

$$D_x(P_0 \nabla f_0) = (D_x P_0) \nabla f_0 + P_0 D_x \nabla f_0 = P_0 D_x \nabla f_0 = P_0 \nabla D_x f_0 - P_0 \nabla f_0$$

in B_1^* , and therefore also on \mathbb{S}^2 . Hence, taking the scalar product with x, and using (3.19) (or (4.47)) to write $\langle x, P_0 \nabla f_0 \rangle$, we get

$$\langle x, D_x(P_0 \nabla f_0) \rangle = \langle x, P_0 \nabla D_x f_0 \rangle - (1+h) JG(h) \psi$$
(4.48)

on \mathbb{S}^2 . By (4.48), (4.45), (4.47), we get

$$\langle x, P_0 \nabla D_x f_0 \rangle = (1+h) J G(h) \psi - \operatorname{div}_{\mathbb{S}^2} \{ (4.47) \}.$$
 (4.49)

Moreover, by (2.5), (2.8), we have div $_{\mathbb{S}^2}\{\varphi(x)x\} = 2\varphi(x)$ for any scalar function φ on \mathbb{S}^2 ; we apply it to $\varphi = (1+h)JG(h)\psi$. Therefore, using (4.49), (4.47) and formulas (4.35), (4.11) of Z, α , the term E_5 in (4.43) is

$$E_5 = \left(\frac{\operatorname{div}_{\mathbb{S}^2}\{(1+h)(\nabla_{\mathbb{S}^2}\psi - W\nabla_{\mathbb{S}^2}h)\}}{(1+h)^2J} + \frac{G(h)\psi}{1+h}\right)\eta.$$
(4.50)

By (4.36), (4.37), (4.39), (4.43), (4.44), (4.50), we obtain (4.1) with W in (4.2) and

$$b = -\frac{\{2(1+h)^2 + |\nabla_{\mathbb{S}^2}h|^2\}G(h)\psi}{(1+h)J^2} + \frac{W}{J} - \frac{\operatorname{div}_{\mathbb{S}^2}\{(1+h)(\nabla_{\mathbb{S}^2}\psi - W\nabla_{\mathbb{S}^2}h)\}}{(1+h)^2J} - \frac{G(h)\psi}{1+h},$$

$$B = \left(\frac{W}{(1+h)J} - \frac{G(h)\psi}{J^2}\right)\nabla_{\mathbb{S}^2}h - \frac{\nabla_{\mathbb{S}^2}\psi}{(1+h)J}.$$

Note that, in computing b, the terms $\frac{W}{J} \frac{|\nabla_{\mathbb{S}^2} h|^2}{(1+h)^2}$ in E_2 in (4.39) and in E_4 in (4.44) cancel out, and also note that, in computing the coefficient of $\nabla_{\mathbb{S}^2} h$ in B, the term $\frac{W}{(1+h)J}$ appears with coefficient 2 in E_2 and with coefficient 1 in E_4 . Finally, using the definition of W, J in (4.2), (2.28), we obtain the formulas for B, b in (4.2), (4.3). The proof of Theorem 4.1 is complete.

Remark 4.8. To transform the free boundary $\partial\Omega$ into the fixed sphere \mathbb{S}^2 , and the fluid domain Ω into the unit ball B_1 , we have used the diffeomorphism $\mathcal{E}_1\gamma$ in (2.18), which is a homogeneous function. Working with homogeneous functions is certainly easier than with general functions, because nice simplifications occur in the calculations of integrals and derivatives: see, for example, properties (2.15).

However, homogeneous functions also have some unpleasant drawbacks in our free boundary problem: the diffeomorphism $\mathcal{E}_1\gamma$ in (2.18) is homogeneous of degree 1, and, in general, it is merely Lipschitz around the origin, even if $h \in C^{\infty}(\mathbb{S}^2)$. Its partial derivatives $\partial_x^{\beta}(\mathcal{E}_1\gamma)$ of order $|\beta| \in \mathbb{N}$ are homogeneous of degree $1 - |\beta|$; in particular, partial derivatives of second order have a singularity of type $|x|^{-1}$ around the origin, so that they are not in $L^{\infty}(B_1)$, and partial derivatives of third order are like $|x|^{-2}$ around the origin, so that they are not even in $L^2(B_1)$. As a consequence, even if $h, \psi \in C^{\infty}(\mathbb{S}^2)$, the solution $\tilde{\Phi}$ of the transformed problem (3.13) cannot have, in general, a high Sobolev regularity $H^m(B_1)$, because of the (artificially created) singularity at the origin, while the corresponding solution Φ of the original problem (1.14) is harmonic in Ω , and, for Φ , the origin is a point like all the other internal points, with no special role.

In some sense, we could say that $\mathcal{E}_1 \gamma$ gives to the origin the role of "the bottom" of the flat case, where "the bottom" is no longer a surface, but it collapses to one single point.

A natural way to overcome this problem is to modify $\mathcal{E}_1 \gamma$ around the origin: for example, consider

$$\gamma_{reg}: B_1 \to \Omega, \quad \gamma_{reg}(x) := (1 + \chi(x)\mathcal{E}_0 h(x))x_g$$

where $\chi \in C^{\infty}(\mathbb{R}^3)$ is a cut-off function such that $\chi(x) = 1$ for $|x| \ge 1/2$, $\chi(x) = 0$ for $|x| \le 1/4$, and $0 \le \chi \le 1$. Thus, $\gamma_{reg} = \mathcal{E}_1 \gamma$ in the annulus 1/2 < |x| < 1, and $\gamma_{reg}(x) = x$ in the ball |x| < 1/4. The map γ_{reg} has no geometric singularity, and, for *h* regular enough, the composition with γ_{reg} preserves the Sobolev regularity, so that $\tilde{\Phi}$ has the same Sobolev regularity as Φ .

So why not using γ_{reg} instead of $\mathcal{E}_1\gamma$? Because surprisingly, unlike in the flat case, the method of the good unknown of Alinhac (at least, the one used in this section) does not work with γ_{reg} . The point at which the method fails is in the proof of Lemma 4.2: the candidate α can be determined by an argument similar to that in Remark 4.5, but then the corresponding matrices \tilde{P} and P_1 do not coincide: they coincide for the terms involving the derivatives of η , but there are terms in η that do not cancel.

For this reason, we have used $\mathcal{E}_1\gamma$, and found ways to overcome the geometric singularities at the origin. This is why we use only the Sobolev space $H^1(B_1)$, and sometimes $H^2(B_1)$, but the higher, and fractional, regularity is treated differently, see subsection 5.3.

4.2 Proof via geometric argument

In this section we give another, independent proof of formula (4.1), using an argument relying on geometry. Here we assume that Ω is star-shaped. The proof is divided in different subsections.

We consider $\gamma_{\varepsilon}, \Omega_{\varepsilon}, \Phi_{\varepsilon}$ as in (3.26), (3.27). Arguing as in [13, Proof of Proposition 8.1] we deduce that $\varepsilon \mapsto \Phi_{\varepsilon}$ is smooth. We then denote $\dot{\Phi} := \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \Phi_{\varepsilon}$ and note that it is harmonic in Ω . Indeed, this follows from the fact that by linearity $\Phi_{\varepsilon} - \Phi$ is harmonic away from the boundary of $\partial\Omega$ and thus

$$\dot{\Phi} = \lim_{\varepsilon \to 0} \frac{\Phi_{\varepsilon} - \Phi}{\varepsilon}$$

is harmonic in Ω . Similarly we denote $\dot{\nu} = \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \nu_{\Omega_{\varepsilon}}(\gamma_{\varepsilon}(x))$ for $x \in \mathbb{S}^2$. We also define, like in Lemma 3.3, the vector field $X : \partial\Omega \to \mathbb{R}^3$ associated with the change of the domain such that, for $x \in \mathbb{S}^2$,

$$X(\gamma(x)) = \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \gamma_{\varepsilon}(x) = \eta(x) x, \qquad (4.51)$$

which is (3.29). Using these notations we may write the Dirichlet-Neumann operator as

$$G(h + \varepsilon \eta)\psi = \langle (\nabla \Phi_{\varepsilon})(\gamma_{\varepsilon}(x)), \nu_{\Omega_{\varepsilon}} \rangle.$$

We may then write (1.18) by differentiating the above and have

$$G'(h)[\eta]\psi = \overbrace{\langle \nabla \dot{\Phi}(\gamma(x)), \nu_{\Omega} \rangle}^{=A_1} + \overbrace{\langle D^2 \Phi(\gamma(x)), X(\gamma(x)), \nu_{\Omega} \rangle}^{=A_2} + \overbrace{\langle \nabla \Phi(\gamma(x)), \dot{\nu}(\gamma(x)) \rangle}^{=A_3}.$$
(4.52)

The calculations for each term is rather cumbersome. We will treat them separately in different subsections.

4.2.1 Calculations of the term A_1

We begin by calculating the term $A_1 = \langle \nabla \dot{\Phi}(\gamma(x)), \nu_{\Omega} \rangle$ in (4.52) and show that it can be written as

$$A_1 = G(h)(-\eta W), (4.53)$$

where $W : \mathbb{S}^2 \to \mathbb{R}$ is in (4.2). We begin by recalling that $\dot{\Phi}$ is harmonic. Therefore, by the definition of the Dirichlet-Neumann operator in (1.13), in order to identify the term A_1 we need to show that $\dot{\Phi}(\gamma(x)) = -\eta W(x)$ for $x \in \mathbb{S}^2$, where W is given by (4.2). To this aim we recall that Φ_{ε} has the boundary values $\Phi_{\varepsilon}(\gamma_{\varepsilon}(x)) = \psi(x)$ for all ε and $x \in \mathbb{S}^2$. We differentiate this with respect to ε and obtain

$$0 = \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \Phi_{\varepsilon}(\gamma_{\varepsilon}(x)) = \dot{\Phi}(\gamma(x)) + \langle \nabla \Phi(\gamma(x)), x \rangle \eta(x).$$

Let us decompose the gradient of Φ into the normal and the tangential components as in (2.3) and by the above it holds

$$\dot{\Phi}(\gamma(x)) = -\left(\left\langle \nabla \Phi(\gamma(x)), \nu_{\Omega} \right\rangle \left\langle \nu_{\Omega}(\gamma(x)), x \right\rangle + \left\langle \nabla_{\partial \Omega} \Phi(\gamma(x)), x \right\rangle \right) \eta(x).$$
(4.54)

We may simplify this by using the definition (1.13), i.e., $\langle \nabla \Phi(\gamma(x)), \nu_{\Omega} \rangle = G(h)\psi$ and the formula of the normal (2.27) which implies $\langle \nu_{\Omega}(\gamma(x)), x \rangle = \frac{1+h}{J}$, where J is defined in (2.28). Therefore the first term on the RHS in (4.54) is

$$\langle \nabla \Phi(\gamma(x)), \nu_{\Omega} \rangle \langle \nu_{\Omega}(\gamma(x)), x \rangle = \frac{1+h}{J} G(h) \psi.$$

To deal with the last term in (4.54), we use (3.6) and get

$$\langle \nabla_{\partial\Omega} \Phi(\gamma(x)), x \rangle = \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle}{J^2}$$

The two above equalities and (4.54) imply the formulas (4.53) and (4.2).

4.2.2 Calculations of the term A_2

This term is the most cumbersome to calculate and we show that it has the form

$$A_{2} = -\eta \frac{\mathcal{L}_{h}(\psi)}{(1+h)J} + \frac{\eta}{J^{2}} \langle \nabla_{\mathbb{S}^{2}} \left(G(h)\psi \right), \nabla_{\mathbb{S}^{2}}h \rangle + \frac{\eta}{(1+h)J^{3}} \left(\Delta_{\mathbb{S}^{2}}h - 2\frac{\langle (D_{\mathbb{S}^{2}}^{2}h)\nabla_{\mathbb{S}^{2}}h, \nabla_{\mathbb{S}^{2}}h \rangle}{J^{2}} - \frac{2(1+h)|\nabla_{\mathbb{S}^{2}}h|^{2}}{J^{2}} - (1+h) \right) \langle \nabla_{\mathbb{S}^{2}}\psi, \nabla_{\mathbb{S}^{2}}h \rangle + \frac{\eta}{(1+h)J^{3}} \langle (D_{\mathbb{S}^{2}}^{2}h)\nabla_{\mathbb{S}^{2}}h, \nabla_{\mathbb{S}^{2}}\psi \rangle - \eta \frac{1+h}{J}H(h)G(h)\psi.$$

$$(4.55)$$

We begin by decomposing vector field X in (4.51) into the normal and tangential components as in (2.3). Recalling that $X(\gamma(x)) = \eta(x) x$ and the formula for the normal (2.27) we have

$$X_{\nu}(\gamma(x)) = \frac{\eta(x)}{J} \langle ((1+h)x - \nabla_{\mathbb{S}^2}hx), \nu_{\Omega}(\gamma(x)) \rangle = \eta(x) \left(\frac{1+h}{J}\right) \nu_{\Omega}(\gamma(x)),$$

$$X_{\partial\Omega}(\gamma(x)) = (X - X_{\nu})(\gamma(x)) = \eta(x)x - \eta(x) \left(\frac{1+h}{J}\right) \left(\frac{(1+h)x - \nabla_{\mathbb{S}^2}h}{J}\right) \qquad (4.56)$$

$$= \eta(x) \frac{|\nabla_{\mathbb{S}^2}h|^2}{J^2} x + \eta(x) \left(\frac{1+h}{J^2}\right) \nabla_{\mathbb{S}^2}h.$$

Then we write

$$A_2 = \langle D^2 \Phi(\gamma(x)) X, \nu_\Omega \rangle = \langle X, \nu_\Omega \rangle \langle D^2 \Phi(\gamma(x)) \nu_\Omega, \nu_\Omega \rangle + \langle D^2 \Phi(\gamma(x)) X_{\partial\Omega}, \nu_\Omega \rangle.$$
(4.57)

Let us first calculate the first term on the RHS of (4.57). First, we have by (4.56) that

$$\langle X, \nu_{\Omega} \rangle = X_{\nu} = \eta(x) \left(\frac{1+h}{J}\right)$$

We proceed by recalling that the function Φ is harmonic in Ω and have by the formula (2.11)

$$\langle D^2 \Phi(\gamma(x))\nu_{\Omega},\nu_{\Omega}\rangle = -\Delta_{\partial\Omega}\Phi - H_{\Omega}\langle \nabla\Phi,\nu_{\Omega}\rangle.$$

The mean curvature is calculated in Lemma 2.2, the Laplace-Beltrami is calculated in Lemma 2.3 (recall that $\Phi(\gamma(s)) = \psi(x)$) and by definition of the Dirichlet-Neumann operator (1.13) it holds $\langle \nabla \Phi, \nu_{\Omega} \rangle \langle \gamma(x) \rangle = G(h)\psi$. Therefore it holds by (2.37)

$$\langle X, \nu_{\Omega} \rangle \langle D^{2} \Phi(\gamma(x)) \nu_{\Omega}, \nu_{\Omega} \rangle = -\eta \frac{1+h}{J} \left(\Delta_{\partial\Omega} \Phi + H_{\Omega}(\gamma(x)) G(h) \psi \right)$$

$$= -\eta \frac{\mathcal{L}_{h}(\psi)}{(1+h)J} + \eta \left(\frac{\mathcal{L}_{h}(h)}{(1+h)J^{3}} - \frac{|\nabla_{\mathbb{S}^{2}} h|^{2}}{(1+h)J^{5}} \right) \langle \nabla_{\mathbb{S}^{2}} \psi, \nabla_{\mathbb{S}^{2}} h \rangle$$

$$- \eta(x) \frac{1+h}{J} H(h) \left(G(h)(\psi) \right),$$

$$(4.58)$$

where the operator \mathcal{L}_h is defined in (2.37).

Let us then calculate the last term in (4.57). We differentiate $\langle \nabla \Phi, \nu_{\Omega} \rangle$ in the direction of $X_{\partial\Omega}$, which of course is on the tangent plane, and have

$$\langle \nabla_{\partial\Omega} \langle \nabla \Phi, \nu_{\Omega} \rangle, X_{\partial\Omega} \rangle = \langle D^2 \Phi X_{\partial\Omega}, \nu_{\Omega} \rangle + \langle D_{\partial\Omega} \nu_{\Omega} X_{\partial\Omega}, \nabla \Phi \rangle$$

Therefore

$$\langle D^2 \Phi X_{\partial\Omega}, \nu_{\Omega} \rangle = \langle \nabla_{\partial\Omega} \langle \nabla \Phi, \nu_{\Omega} \rangle, X_{\partial\Omega} \rangle - \langle D_{\partial\Omega} \nu_{\Omega} X_{\partial\Omega}, \nabla \Phi \rangle.$$
(4.59)

Let us treat the first term on the RHS of (4.59). First we write

$$\langle \nabla_{\partial\Omega} \langle \nabla \Phi, \nu_{\Omega} \rangle, X_{\partial\Omega} \rangle = \langle \nabla_{\partial\Omega} \langle \nabla \Phi, \nu_{\Omega} \rangle, X \rangle.$$

We have by (2.29), by $\langle \nabla \Phi, \nu_{\Omega} \rangle(\gamma(x)) = (G(h)\psi)(x)$, and by $X(\gamma(x)) = \eta(x)x$ that

$$\langle \nabla_{\partial\Omega} \langle \nabla\Phi, \nu_{\Omega} \rangle, X_{\partial\Omega} \rangle = \langle \nabla_{\partial\Omega} \langle \nabla\Phi, \nu_{\Omega} \rangle, X \rangle = \frac{\eta(x)}{J^2} \langle \nabla_{\mathbb{S}^2} \big(G(h)\psi \big), \nabla_{\mathbb{S}^2} h \rangle.$$
(4.60)

We begin by treating the second term on the RHS of (4.59) by recalling that the differential of the normal is the second fundamental form $B_{\partial\Omega} = D_{\partial\Omega}\nu_{\Omega}$. In particular, $D_{\partial\Omega}\nu_{\Omega}$ is symmetric, because $(D_{\partial\Omega}\nu_{\Omega})\nu_{\Omega} = 0$, and $D\nu_{\Omega}$ is the Hessian of the signed distance, therefore it is a symmetric matrix. Hence for all vector fields F, Ψ one has

$$\langle (D_{\partial\Omega}\nu_{\Omega})F,\Psi\rangle = \langle (D_{\partial\Omega}\nu_{\Omega})F,\Psi_{\partial\Omega}\rangle$$

Moreover, by definition, $(D_{\partial\Omega}\nu_{\Omega})F = (D_{\partial\Omega}\nu_{\Omega})F_{\partial\Omega}$, so that $D_{\partial\Omega}\nu_{\Omega}$ is symmetric. Thus

$$\langle D_{\partial\Omega}\nu_{\Omega} X_{\partial\Omega}, \nabla\Phi \rangle = \langle D_{\partial\Omega}\nu_{\Omega}\nabla_{\partial\Omega}\Phi, X_{\partial\Omega} \rangle$$

We write the normal by using the vector field n defined in (2.33) as $\nu_{\Omega} = \frac{n}{|n|}$. Then it holds $D_{\partial\Omega}\nu_{\Omega} = \frac{1}{|n|}D_{\partial\Omega}n + n \otimes (\nabla_{\partial\Omega}1/|n|)$ and using $\langle n, X_{\partial\Omega} \rangle = 0$ we have

$$\langle D_{\partial\Omega}\nu_{\Omega}\nabla\Phi, X_{\partial\Omega}\rangle = \frac{1}{|n|} \langle D_{\partial\Omega}n\,\nabla_{\partial\Omega}\Phi, X_{\partial\Omega}\rangle.$$

We use (2.29) and $\Phi(\gamma(x)) = \psi(x)$ to deduce that

$$(D_{\partial\Omega}n)(\gamma(x))(\nabla_{\partial\Omega}\Phi)(\gamma(x)) = D_{\partial\Omega}n(\gamma(x))\left(\frac{\nabla_{\mathbb{S}^2}\psi(x)}{1+h} + \frac{\langle\nabla_{\mathbb{S}^2}\psi,\nabla_{\mathbb{S}^2}h\rangle}{(1+h)^2J}\nu_{\Omega}(\gamma(x))\right)$$
$$= \frac{1}{1+h}D_{\partial\Omega}n(\gamma(x))\nabla_{\mathbb{S}^2}\psi(x)$$

because $D_{\partial\Omega}n(y)\nu_{\Omega}(y) = 0$, and then (2.30) and have

$$D_{\partial\Omega}n(\gamma(x))\nabla_{\mathbb{S}^2}\psi = \frac{1}{1+h}D_{\mathbb{S}^2}\tilde{n}\nabla_{\mathbb{S}^2}\psi - \frac{\langle\nabla_{\mathbb{S}^2}\psi,\nabla_{\mathbb{S}^2}h\rangle}{(1+h)J}D_{\mathbb{S}^2}\tilde{n}\nabla_{\mathbb{S}^2}h$$

Recall that by (4.56) it holds $X_{\partial\Omega}(\gamma(x)) = \eta \frac{|\nabla_{\mathbb{S}^2} h|^2}{J^2} x + \eta \left(\frac{1+h}{J^2}\right) \nabla_{\mathbb{S}^2} h$. Then by $|n|(\gamma(x)) = J$ we may write

$$\begin{split} \langle D_{\partial\Omega}\nu_{\Omega}\nabla\Phi, X_{\partial\Omega}\rangle(\gamma(x)) &= \frac{1}{(1+h)} \langle D_{\partial\Omega}n(\gamma(x))\nabla_{\mathbb{S}^{2}}\psi, X_{\partial\Omega}(\gamma(x))\rangle \\ &= \eta(x)\frac{|\nabla_{\mathbb{S}^{2}}h|^{2}}{(1+h)^{2}J^{3}} \Big(\langle D_{\mathbb{S}^{2}}\tilde{n}\nabla_{\mathbb{S}^{2}}\psi, x\rangle - \frac{\langle \nabla_{\mathbb{S}^{2}}\psi, \nabla_{\mathbb{S}^{2}}h\rangle}{J^{2}} \langle D_{\mathbb{S}^{2}}\tilde{n}\nabla_{\mathbb{S}^{2}}h, x\rangle \Big) \\ &+ \frac{\eta}{(1+h)J^{3}} \Big(\langle D_{\mathbb{S}^{2}}\tilde{n}\nabla_{\mathbb{S}^{2}}\psi, \nabla_{\mathbb{S}^{2}}h\rangle - \frac{\langle \nabla_{\mathbb{S}^{2}}\psi, \nabla_{\mathbb{S}^{2}}h\rangle}{J^{2}} \langle D_{\mathbb{S}^{2}}\tilde{n}\nabla_{\mathbb{S}^{2}}h, \nabla_{\mathbb{S}^{2}}h\rangle \Big). \end{split}$$
(4.61)

By (2.17) it holds $(D_{\mathbb{S}^2}\tilde{n})^T x = 2\nabla_{\mathbb{S}^2}h$. Using this and recalling $J = \sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2}h|^2}$ we may write the first term on the RHS of (4.61) as

$$\eta \frac{|\nabla_{\mathbb{S}^2} h|^2}{(1+h)^2 J^3} \left(\langle D_{\mathbb{S}^2} \tilde{n} \nabla_{\mathbb{S}^2} \psi, x \rangle - \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle}{J^2} \langle D_{\mathbb{S}^2} \tilde{n} \nabla_{\mathbb{S}^2} h, x \rangle \right) = 2\eta \frac{|\nabla_{\mathbb{S}^2} h|^2}{J^5} \langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle.$$
(4.62)

We use (2.34) and have

$$\begin{split} \langle D_{\mathbb{S}^2} \tilde{n} \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle &= -\langle D_{\mathbb{S}^2}^2 h \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \psi \rangle + (1+h) \langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle, \\ \langle D_{\mathbb{S}^2} \tilde{n} \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} h \rangle &= -\langle D_{\mathbb{S}^2}^2 h \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} h \rangle + (1+h) |\nabla_{\mathbb{S}^2} h|^2. \end{split}$$

We may then write the last term in (4.61) as

$$\frac{\eta(x)}{(1+h)J^3} \left(\langle D_{\mathbb{S}^2} \tilde{n} \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle - \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle}{J^2} \langle D_{\mathbb{S}^2} \tilde{n} \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} h \rangle \right) \\
= -\frac{\eta(x)}{(1+h)J^3} \langle D_{\mathbb{S}^2}^2 h \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \psi \rangle + \frac{\eta(x)}{J^3} \langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle \\
+ \frac{\eta(x)}{(1+h)J^5} \langle D_{\mathbb{S}^2}^2 h \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} h \rangle \langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle - \eta(x) \frac{|\nabla_{\mathbb{S}^2} h|^2}{J^5} \langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle.$$
(4.63)

The formula (4.55) then follows by combining (4.57), (4.58), (4.59), (4.60), (4.61), (4.62) and (4.63).

4.2.3 Calculations of the term A_3

We show that the term $A_3 = \langle \nabla \Phi(\gamma(x)), \dot{\nu}(\gamma(x)) \rangle$ in (4.52), where $\dot{\nu} = \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \nu_{\Omega_{\varepsilon}}(\gamma_{\varepsilon}(x))$ can be written as

$$A_3 = \eta \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle}{J^3} - \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} \eta \rangle}{(1+h)J} + \frac{\langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle \langle \nabla_{\mathbb{S}^2} \eta, \nabla_{\mathbb{S}^2} h \rangle}{(1+h)J^3}.$$
(4.64)

To this aim we use the result in [13], equation (8.6), where a general formula for calculation of $\dot{\nu}$ is derived, and we have

$$\langle \nabla \Phi(\gamma(x)), \dot{\nu}(\gamma(x)) \rangle = -\langle D_{\partial\Omega} X(\gamma(x)) \nabla_{\partial\Omega} \Phi(\gamma(x)), \nu_{\partial\Omega}(\gamma(x)) \rangle$$

Recall that $X(\gamma(x)) = \eta(x) x$. We then have $D_{\mathbb{S}^2} \tilde{X}(x) = \eta I_{\mathbb{S}^2} + x \otimes \nabla_{\mathbb{S}^2} \eta$, where $I_{\mathbb{S}^2} = I - x \otimes x$, and therefore by applying the second equality in Lemma 2.1 we deduce

$$D_{\partial\Omega}X(\gamma(x)) = \frac{\eta}{1+h}I_{\mathbb{S}^2} + \frac{1}{1+h}x \otimes \nabla_{\mathbb{S}^2}\eta - \frac{\eta(x)}{(1+h)J^2}\nabla_{\mathbb{S}^2}h \otimes \nabla_{\mathbb{S}^2}h - \frac{\langle \nabla_{\mathbb{S}^2}\eta, \nabla_{\mathbb{S}^2}h\rangle}{(1+h)J^2}x \otimes \nabla_{\mathbb{S}^2}h + \frac{\eta(x)}{J^2}\nabla_{\mathbb{S}^2}h \otimes x + \frac{\langle \nabla_{\mathbb{S}^2}\eta, \nabla_{\mathbb{S}^2}h\rangle}{J^2}x \otimes x$$

Therefore it holds by the first equality in Lemma 2.1, by $\Phi(\gamma(x)) = \psi(x)$, and by the formula of the normal (2.27) that

$$\begin{split} -\langle D_{\partial\Omega}X(\gamma(x))\nabla_{\partial\Omega}\Phi(\gamma(x)),\nu_{\partial\Omega}(\gamma(x))\rangle &= -\frac{1}{1+h}\langle D_{\partial\Omega}X(\gamma(x))\nabla_{\mathbb{S}^{2}}\psi,\nu_{\partial\Omega}(\gamma(x))\rangle\rangle\\ &= -\frac{1}{J}\langle D_{\partial\Omega}X(\gamma(x))\nabla_{\mathbb{S}^{2}}\psi,x\rangle + \frac{1}{(1+h)J}\langle D_{\partial\Omega}X(\gamma(x))\nabla_{\mathbb{S}^{2}}\psi,\nabla_{\mathbb{S}^{2}}h\rangle\\ &= -\frac{\langle \nabla_{\mathbb{S}^{2}}\psi,\nabla_{\mathbb{S}^{2}}\eta\rangle}{(1+h)J} + \frac{\langle \nabla_{\mathbb{S}^{2}}\psi,\nabla_{\mathbb{S}^{2}}h\rangle\langle\nabla_{\mathbb{S}^{2}}\eta,\nabla_{\mathbb{S}^{2}}h\rangle}{(1+h)J^{3}}\\ &+ \frac{\eta}{(1+h)^{2}J}\langle \nabla_{\mathbb{S}^{2}}\psi,\nabla_{\mathbb{S}^{2}}h\rangle - \frac{\eta|\nabla_{\mathbb{S}^{2}}h|^{2}}{(1+h)^{2}J^{3}}\langle \nabla_{\mathbb{S}^{2}}\psi,\nabla_{\mathbb{S}^{2}}h\rangle. \end{split}$$

The formula (4.64) then follows from above and by recalling that $J = \sqrt{(1+h)^2 + |\nabla_{\mathbb{S}^2} h|^2}$ which then simplifies the last terms as

$$\frac{\eta}{(1+h)^2} \langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle - \frac{\eta |\nabla_{\mathbb{S}^2} h|^2}{(1+h)^2 J^3} \langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle = \frac{\eta}{J^3} \langle \nabla_{\mathbb{S}^2} \psi, \nabla_{\mathbb{S}^2} h \rangle.$$

4.2.4 Conclusion

By (4.52), (4.53), (4.55), (4.64), we obtain (4.1) with W, B in (4.2) and

$$b = -\frac{\mathcal{L}_{h}(\psi)}{(1+h)J} + \frac{1}{J^{2}} \langle \nabla_{\mathbb{S}^{2}} \left(G(h)(\psi) \right), \nabla_{\mathbb{S}^{2}} h \rangle + \frac{\langle \nabla_{\mathbb{S}^{2}} \psi, \nabla_{\mathbb{S}^{2}} h \rangle}{(1+h)J^{3}} \left(\Delta_{\mathbb{S}^{2}} h - 2 \frac{\langle D_{\mathbb{S}^{2}}^{2} h \nabla_{\mathbb{S}^{2}} h, \nabla_{\mathbb{S}^{2}} h \rangle}{J^{2}} - \frac{2(1+h)|\nabla_{\mathbb{S}^{2}} h|^{2}}{J^{2}} \right)$$
(4.65)
$$+ \frac{\langle (D_{\mathbb{S}^{2}}^{2} h) \nabla_{\mathbb{S}^{2}} h, \nabla_{\mathbb{S}^{2}} \psi \rangle}{(1+h)J^{3}} - \frac{1+h}{J} H(h) G(h) \psi,$$

where the operator \mathcal{L}_h is defined in (2.36), and the mean curvature H(h) is calculated in Lemma 2.2. Finally, a long, but straightforward, calculation shows that the function b in (4.65) coincides with b in (4.3). This completes the proof of Theorem 4.1.

4.3 Another proof of the Hamiltonian structure

In this short subsection we provide an alternative proof of Proposition 3.2, which makes use of formula (4.1) instead of relying on Hadamard's formula (Lemma 3.3) to compute $\partial_h K(h, \psi)$.

Second proof of Proposition 3.2. The only difference with respect to the first proof concerns the calculation of $\partial_h K$. Recalling the definition (3.20) of $K(h, \psi)$, we have

$$\partial_h K(h,\psi)[\eta] = \frac{1}{2} \int_{\mathbb{S}^2} \eta J(h) \psi G(h) \psi \, d\sigma + \frac{1}{2} \int_{\mathbb{S}^2} (1+h) J'(h)[\eta] \psi G(h) \psi \, d\sigma + \frac{1}{2} \int_{\mathbb{S}^2} (1+h) J(h) \psi G'(h)[\eta] \psi \, d\sigma,$$
(4.66)

where J(h) = J is in (2.28). The second term in the RHS of (4.66) is equal to

$$\frac{1}{2} \int_{\mathbb{S}^2} (1+h) \frac{(1+h)\eta + \langle \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \eta \rangle}{J} \psi G(h) \psi \, d\sigma$$
$$= \frac{1}{2} \int_{\mathbb{S}^2} \eta \frac{(1+h)^2}{J} \psi G(h) \psi \, d\sigma - \frac{1}{2} \int_{\mathbb{S}^2} \eta \operatorname{div}_{\mathbb{S}^2} \left(\frac{(1+h)\nabla_{\mathbb{S}^2} h}{J} \, \psi G(h) \psi \right) d\sigma,$$

where we have used the divergence theorem (2.12) on \mathbb{S}^2 . For the last term of (4.66), we use (4.1) to replace $G'(h)[\eta]\psi$ with $b\eta + \langle B, \nabla_{\mathbb{S}^2}\eta \rangle - G(h)(W\eta)$, and then we apply the divergence theorem (2.12) on \mathbb{S}^2 to the integral containing $\langle B, \nabla_{\mathbb{S}^2}\eta \rangle$, and (3.22) to that containing $G(h)(W\eta)$. Hence the last term of (4.66) is

$$\frac{1}{2} \int_{\mathbb{S}^2} \eta \left\{ (1+h)J(b\psi - WG(h)\psi) - \operatorname{div}_{\mathbb{S}^2} \left((1+h)J\psi B \right) \right\} d\sigma$$

Thus we have proved that

$$\partial_h K(h,\psi) = \frac{1}{2} \left(J + \frac{(1+h)^2}{J} \right) \psi G(h)\psi + \frac{1}{2} (1+h) J \left(b\psi - WG(h)\psi \right) - \frac{1}{2} \operatorname{div}_{\mathbb{S}^2} \left(\frac{1+h}{J} (\nabla_{\mathbb{S}^2} h) \psi G(h)\psi + (1+h) J \psi B \right).$$
(4.67)

By definition (4.2) of W and B, the term in the tangential divergence in (4.67) is

$$\frac{1+h}{J}(\nabla_{\mathbb{S}^2}h)\psi G(h)\psi + (1+h)J\psi B = -\psi(\nabla_{\mathbb{S}^2}\psi - W\nabla_{\mathbb{S}^2}h).$$

Inserting the last identity in (4.67), and using definitions (4.2), (4.3), after some cancellations it remains

$$\partial_h K(h,\psi) = \frac{|\nabla_{\mathbb{S}^2}\psi|^2}{2} - \frac{1}{2} \left((1+h)G(h)\psi + \frac{\langle\nabla_{\mathbb{S}^2}\psi,\nabla_{\mathbb{S}^2}h\rangle}{J} \right)^2,\tag{4.68}$$

and the proof is complete.

5 Analyticity and tame estimates for the Dirichlet-Neumann operator

In this section we show that the Dirichlet-Neumann operator depends analytically on the elevation function h, and we prove tame estimates for it. As a first step, we prove that the solution $\tilde{\Phi}$ of the elliptic problem (3.13) is analytic in h, by writing it as a power series $\tilde{\Phi} = u = \sum u_n$ of h and its derivatives, see (5.3), (5.4), and estimating each term u_n of the series. Such estimates are rather simple to obtain in low norm, see Lemma 5.1. On the contrary, estimates in higher norm are much heavier to obtain. To this aim, we first recall in subsection 5.2 some general notions about Sobolev spaces on domains of \mathbb{R}^n and on their boundaries, mainly following the approach of Triebel's book [41], and we list several useful properties of Sobolev norms. The proof of such properties is in the Appendix. Then, in subsection 5.3, we estimate the high Sobolev norm of each term u_n of the power series of $\tilde{\Phi}$, see Lemma 5.14, and we obtain tame estimates for the Dirichlet-Neumann operator, see Theorem 5.19.

5.1 Analytic dependence in low norm

The matrix P(x) appearing in (3.13) is defined in (3.14), it is written in (3.15) in terms of h, and, for $||h||_{L^{\infty}(\mathbb{S}^2)} < 1$, it is given by the series $P = \sum_{n=0}^{\infty} P_n$, where $P_0 = I$,

$$P_1 = (\mathcal{E}_0 h)I - (\nabla \mathcal{E}_0 h) \otimes x - x \otimes \nabla \mathcal{E}_0 h, \quad P_n = (-\mathcal{E}_0 h)^{n-2} |\nabla \mathcal{E}_0 h|^2 x \otimes x \quad \forall n \ge 2.$$
(5.1)

Since $\mathcal{E}_0 h$ is a 0-homogeneous function of $x \in \mathbb{R}^3 \setminus \{0\}$, the matrix $(\nabla \mathcal{E}_0 h) \otimes x$, its transpose and the matrix $|\nabla \mathcal{E}_0 h|^2 x \otimes x$, and therefore all matrices P_n , are 0-homogeneous functions of x. Hence

$$\|\mathcal{E}_0 h\|_{L^{\infty}(B_1)} = \|h\|_{L^{\infty}(\mathbb{S}^2)}, \quad \|(\nabla \mathcal{E}_0 h) \otimes x\|_{L^{\infty}(B_1)} = \|\nabla \mathcal{E}_0 h\|_{L^{\infty}(\mathbb{S}^2)} = \|\nabla_{\mathbb{S}^2} h\|_{L^{\infty}(\mathbb{S}^2)},$$

and

$$\begin{aligned} \|P_1\|_{L^{\infty}(B_1)} &\leq \|h\|_{L^{\infty}(\mathbb{S}^2)} + 2\|\nabla_{\mathbb{S}^2}h\|_{L^{\infty}(\mathbb{S}^2)}, \\ \|P_n\|_{L^{\infty}(B_1)} &\leq \|h\|_{L^{\infty}(\mathbb{S}^2)}^{n-2} \|\nabla_{\mathbb{S}^2}h\|_{L^{\infty}(\mathbb{S}^2)}^2 \quad \forall n \geq 2. \end{aligned}$$
(5.2)

Thus, for any fixed $\delta_0 \in (0,1)$, the series $\sum P_n$ is totally convergent in the norm of $L^{\infty}(B_1)$ uniformly for h in the ball $\|h\|_{L^{\infty}(\mathbb{S}^2)} \leq \delta_0$.

We want to expand the solution $\tilde{\Phi}$ of the elliptic problem (3.13) in powers of h. Thus, we consider the series $u = \sum_{n=0}^{\infty} u_n$ whose terms are recursively defined in the following way: u_0 is the unique solution of the Laplace problem

$$u_0 \in H^1(B_1), \quad \Delta u_0 = 0 \text{ in } B_1, \quad u_0 = \psi \text{ on } \mathbb{S}^2,$$
(5.3)

namely u_0 is the harmonic extension of ψ to the open unit ball, and, for $n \ge 1$, u_n is recursively defined as the unique solution of

$$-\Delta u_n = \operatorname{div} g_n \quad \text{in } B_1, \quad u_n \in H_0^1(B_1)$$
(5.4)

with datum

$$g_n := \sum_{k=0}^{n-1} P_{n-k} \nabla u_k, \quad n \ge 1.$$
(5.5)

In the next lemma we estimate $\|\nabla u_n\|_{L^2(B_1)}$ using the inequality

$$\|P_n\|_{L^{\infty}(B_1)} \le (2\tau)^n \quad \forall n \ge 1, \quad \tau := \|h\|_{L^{\infty}(\mathbb{S}^2)} + \|\nabla_{\mathbb{S}^2}h\|_{L^{\infty}(\mathbb{S}^2)} = \|h\|_{W^{1,\infty}(\mathbb{S}^2)}.$$
 (5.6)

Note that (5.6) is less accurate than (5.2), but using (5.2) instead of (5.6) in proving Lemma 5.1 gives no relevant improvement.

Lemma 5.1. The functions u_n defined in (5.3), (5.4), (5.5) satisfy

$$\|\nabla u_n\|_{L^2(B_1)} \le C \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} ((1+C)2\tau)^n \tag{5.7}$$

for all $n \ge 0$, where τ is defined in (5.6) and C is a universal positive constant.

Proof. By the standard theory of harmonic functions and elliptic PDEs, one has

$$\|\nabla u_0\|_{L^2(B_1)} \le C \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)}, \quad \|\nabla u_n\|_{L^2(B_1)} \le C \|g_n\|_{L^2(B_1)}, \quad n \ge 1,$$
(5.8)

where C is a universal positive constant. Thus (5.7) for n = 0 holds by the first inequality in (5.8). Now assume that, for a given $n \ge 1$, inequality (5.7) holds for u_0, \ldots, u_{n-1} . By (5.5), (5.6) and the induction assumption, one has

$$\|g_n\|_{L^2(B_1)} \le \sum_{k=0}^{n-1} \|P_{n-k}\|_{L^{\infty}(B_1)} \|\nabla u_k\|_{L^2(B_1)} \le C \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} (2\tau)^n \sum_{k=0}^{n-1} (1+C)^k.$$
(5.9)

Since

$$\sum_{k=0}^{n-1} (1+C)^k = \frac{(1+C)^n - 1}{(1+C) - 1} < \frac{(1+C)^n}{C},$$
(5.10)

using the second inequality in (5.8) we obtain (5.7).

From (5.7) it follows that, for any $\rho \in (0, 1)$, the series $\sum \nabla u_n$ converges totally in $L^2(B_1)$, uniformly in h in the ball $(1 + C)2\tau \leq \rho$. Moreover, u_n (as well as P_n and g_n) depends on h in a *n*-homogeneous way, namely $u_n(\lambda h) = \lambda^n u_n(h)$ for all $\lambda \in \mathbb{R}$, and it coincides with a *n*-linear, continuous, symmetric map

$$\mathring{u}_n: W^{1,\infty}(\mathbb{S}^2) \times \ldots \times W^{1,\infty}(\mathbb{S}^2) \to H^1_0(B_1), \quad (h_1,\ldots,h_n) \mapsto \mathring{u}_n[h_1,\ldots,h_n]$$
(5.11)

evaluated at (h, \ldots, h) , i.e., $u_n(h) = \mathring{u}_n[h, \ldots, h]$. Explicitly, one has $P_n(h) = \mathring{P}_n[h, \ldots, h]$, where $\mathring{P}_1 = P_1$ and, for $n \ge 2$, \mathring{P}_n is the *n*-linear, continuous, symmetric map

$$\overset{\circ}{P}_{n}[h_{1},\ldots,h_{n}] = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_{n}} (-1)^{n} \langle \nabla \mathcal{E}_{0} h_{\sigma(1)}, \nabla \mathcal{E}_{0} h_{\sigma(2)} \rangle (\mathcal{E}_{0} h_{\sigma(3)}) \cdots (\mathcal{E}_{0} h_{\sigma(n)}) x \otimes x_{n}$$

where \mathfrak{S}_n is the set of the permutations of $\{1, \ldots, n\}$; $g_n(h) = \mathring{g}_n[h, \ldots, h]$, where

$$\mathring{g}_{n}[h_{1},\ldots,h_{n}] = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_{n}} \sum_{k=0}^{n-1} \mathring{P}_{n-k}[h_{\sigma(k+1)},\ldots,h_{\sigma(n)}] \nabla \{\mathring{u}_{k}[h_{\sigma(1)},\ldots,h_{\sigma(k)}]\},\$$

and $\mathring{u}_n[h_1,\ldots,h_n]$ is the solution of problem (5.4) with $\mathring{g}_n[h_1,\ldots,h_n]$ instead of g_n . Adapting the proof of Lemma 5.1, one easily obtains

$$\|\nabla\{\hat{u}_n[h_1,\ldots,h_n]\}\|_{L^2(B_1)} \le C \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} \prod_{j=1}^n ((1+C)2\|h_j\|_{W^{1,\infty}(\mathbb{S}^2)}).$$

Moreover, since $u_n(h) = \mathring{u}_n[h, \ldots, h]$, the derivative of u_n with respect to h in direction η is $u'_n(h)[\eta] = n\mathring{u}_n[h, \ldots, h, \eta]$, its second derivative is $u''_n(h)[\eta_1, \eta_2] = n(n-1)\mathring{u}_n[h, \ldots, h, \eta_1, \eta_2]$, and so on. Thus, we have proved the following analyticity result.

Proposition 5.2. For every $\psi \in H^{\frac{1}{2}}(\mathbb{S}^2)$, the map $h \mapsto u := \sum_{n=0}^{\infty} u_n$ from the ball

$$\{h \in W^{1,\infty}(\mathbb{S}^2) : \|h\|_{L^{\infty}(\mathbb{S}^2)} + \|\nabla_{\mathbb{S}^2}h\|_{L^{\infty}(\mathbb{S}^2)} < \delta_0\}$$
(5.12)

into $H^1(B_1)$ is well-defined and analytic, with

$$\begin{aligned} \|u\|_{H^{1}(B_{1})} &\leq C \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^{2})}, \quad \|u'(h)[\eta]\|_{H^{1}(B_{1})} \leq C \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^{2})} \|\eta\|_{W^{1,\infty}(\mathbb{S}^{2})}, \\ \|u''(h)[\eta_{1},\eta_{2}]\|_{H^{1}(B_{1})} &\leq C \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^{2})} \|\eta_{1}\|_{W^{1,\infty}(\mathbb{S}^{2})} \|\eta_{2}\|_{W^{1,\infty}(\mathbb{S}^{2})}, \end{aligned}$$

where the radius $\delta_0 \in (0,1)$ and the constant C > 0 are universal constants.

5.2 Sobolev spaces on domains and boundaries

In this subsection we consider both standard and non-isotropic versions of Sobolev spaces of functions on bounded open subsets of \mathbb{R}^n and on their smooth boundary, and we prove some useful properties for them; these properties will be used in subsection 5.3 for the sphere \mathbb{S}^2 and for the unit ball B_1 with various cut-off functions. We follow the approach of Triebel's book [41], i.e., localization, rectification and extension. The results of this subsection are classical, but it is nontrivial to find a complete proof for all of them in literature. The proofs are collected in the Appendix. Concerning the notation, in this subsection (and only in this one) n denotes the dimension of the Euclidean space \mathbb{R}^n .

• On \mathbb{R}^n , for $s \in \mathbb{R}$, we consider the usual Sobolev spaces (Bessel potential spaces)

$$H^{s}(\mathbb{R}^{n}) := \{ u \in \mathcal{S}'(\mathbb{R}^{n}) : \|u\|_{H^{s}(\mathbb{R}^{n})} < \infty \}, \quad \|u\|_{H^{s}(\mathbb{R}^{n})}^{2} := \int_{\mathbb{R}^{n}} |\hat{u}(\xi)|^{2} (1 + |\xi|^{2})^{s} \, d\xi, \quad (5.13)$$

where $\mathcal{S}'(\mathbb{R}^n)$ is the space of tempered distributions on \mathbb{R}^n , and \hat{u} is the Fourier transform of u on \mathbb{R}^n . We also consider the following non-isotropic version of those spaces, where the last real

variable x_n plays a distinguished role. For $x \in \mathbb{R}^n$, denote $x' = (x_1, \ldots, x_{n-1})$, so that $x = (x', x_n)$. For $s, r \in \mathbb{R}$, we define

$$H^{s,r}(\mathbb{R}^{n}) := \{ u \in \mathcal{S}'(\mathbb{R}^{n}) : \|u\|_{H^{s,r}(\mathbb{R}^{n})} < \infty \}, \\ \|u\|_{H^{s,r}(\mathbb{R}^{n})}^{2} := \int_{\mathbb{R}^{n}} |\hat{u}(\xi)|^{2} (1 + |\xi'|^{2})^{r} (1 + |\xi|^{2})^{s} d\xi = \|\Lambda'_{r}u\|_{H^{s}(\mathbb{R}^{n})}^{2},$$
(5.14)

where Λ'_r is the Fourier multiplier of symbol $(1 + |\xi'|^2)^{\frac{r}{2}}$. In subsection 5.3 these spaces will be used for $s \in \{0, 1, 2\}$ and $r \ge 0$ real. They are used by Triebel, see [41], Definition 4.2.1, page 218; see also Lannes [30], Definition 2.11.

• On any open set $\Omega \subseteq \mathbb{R}^n$, following [41], Definition 1 in Section 3.2.2, we define

$$H^{s}(\Omega) := \{ u \in \mathcal{S}'(\Omega) : \| u \|_{H^{s}(\Omega)} < \infty \}, \\ |u\|_{H^{s}(\Omega)} := \inf \{ \| v \|_{H^{s}(\mathbb{R}^{n})} : v \in H^{s}(\mathbb{R}^{n}), \ v|_{\Omega} = u \}, \\ H^{s,r}(\Omega) := \{ u \in \mathcal{S}'(\Omega) : \| u \|_{H^{s,r}(\Omega)} < \infty \},$$
(5.15)

$$\|u\|_{H^{s,r}(\Omega)} := \inf \{ \|v\|_{H^{s,r}(\mathbb{R}^n)} : v \in H^{s,r}(\mathbb{R}^n), \ v|_{\Omega} = u \},$$
(5.16)

with the convention that the infimum of the empty set is ∞ . In other words, $H^s(\Omega)$ is the set of the restrictions to Ω of the elements of $H^s(\mathbb{R}^n)$, and similarly for $H^{s,r}(\Omega)$. We mainly deal with the case when Ω is the open half space

$$\mathbb{R}^{n}_{+} := \{ (x', x_{n}) \in \mathbb{R}^{n} : x_{n} > 0 \}.$$
(5.17)

• On smooth compact manifolds, we recall the construction and the definition in [41], Sections 3.2.1 and 3.2.2. Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with smooth boundary $\Gamma = \partial \Omega$. Fix N open balls K_1, \ldots, K_N that cover Γ , namely $\Gamma \subset \bigcup_{j=1}^N K_j$, such that $\Gamma \cap K_j$ is nonempty for all $j = 1, \ldots, N$. For every ball K_j , fix a function \mathbf{f}_j such that

- (i) $\mathbf{f}_i: \mathbb{R}^n \to \mathbb{R}^n$ is a C^{∞} diffeomorphism of \mathbb{R}^n onto itself,
- (*ii*) \mathbf{f}_j maps the ball K_j onto a bounded open subset $A_j = \mathbf{f}_j(K_j)$ of \mathbb{R}^n ,
- (*iii*) f_j maps the center p_j of the ball K_j into 0,
- (iv) \mathbf{f}_j maps $\Gamma \cap K_j$ onto the (n-1)-dimensional open subset $\mathbf{f}_j(\Gamma \cap K_j) = \{y \in A_j : y_n = 0\} = A_j \cap \mathbb{R}_0^n$ of the hyperplane $\mathbb{R}_0^n := \{y \in \mathbb{R}^n : y_n = 0\},$
- (v) \mathbf{f}_j maps $\Omega \cap K_j$ onto the simply connected open subset $\mathbf{f}_j(\Omega \cap K_j) = \{y \in A_j : y_n > 0\} = A_j \cap \mathbb{R}^n_+$ of the half space \mathbb{R}^n_+ ,
- (vi) the Jacobian matrix $D\mathbf{f}_j(x)$ is invertible at every point x in the closure \overline{K}_j of K_j ,
- (vii) the Jacobian matrix $Df_j(p_j)$ of f_j at the center p_j of K_j is the identity matrix.

Also fix a smooth open set K_0 such that $\overline{K}_0 \subset \Omega$ and $\overline{\Omega} \subset \bigcup_{j=0}^N K_j$, and fix a resolution of unity $\{\psi_j\}_{j=0}^N$, where $\psi_j : \mathbb{R}^n \to \mathbb{R}$ is a C^{∞} function with $\operatorname{supp}(\psi_j) \subset K_j$, and $\sum_{j=0}^N \psi_j(x) = 1$ for all $x \in \overline{\Omega}$. Thus the restriction of ψ_1, \ldots, ψ_N to Γ is a resolution of unity for Γ , and, in addition, $\sum_{j=1}^N \psi_j(x) = 1$ for all $x \in \overline{\Omega} \setminus K_0$. On Γ , for $s \in \mathbb{R}$, we define

$$H^{s}(\Gamma) := \{ u \in \mathcal{S}'(\Gamma) : (\psi_{j}u)(\mathsf{g}_{j}(\cdot,0)) \in H^{s}(\mathbb{R}^{n-1}) \; \forall j = 1, \dots, N \}, \\ \|u\|_{H^{s}(\Gamma)} := \sum_{j=1}^{N} \|(\psi_{j}u)(\mathsf{g}_{j}(\cdot,0))\|_{H^{s}(\mathbb{R}^{n-1})}, \quad \mathsf{g}_{j} := \mathsf{f}_{j}^{-1},$$
(5.18)

where $(\psi_j u)(\mathbf{g}_j(\cdot, 0))$ is defined on \mathbb{R}^{n-1} , extended to 0 in the set of all $y' \in \mathbb{R}^{n-1}$ such that $\mathbf{g}_j(y', 0) \notin K_j$, in which case one has $\psi_j(\mathbf{g}_j(y', 0)) = 0$.

In subsection 5.3 these spaces will be used for $\Gamma = \mathbb{S}^2$. For $\Gamma = \mathbb{S}^{n-1}$, the construction can be more explicit and adapted to the spherical geometry. Given any $\delta \in (0, 1/4)$, we choose K_0 as the ball $B_{1-\delta}(0)$ of center the origin and radius $1 - \delta$, and $K_j = B_{2\delta}(p_j)$, $j = 1, \ldots, N$, where N is sufficiently large and the points $p_j \in S^{n-1}$ are appropriately positioned to obtain the covering property above. In this way one has

$$\sum_{j=1}^{N} \psi_j(x) = 1 \quad \forall x \in \mathbb{R}^n, \ 1 - \delta \le |x| \le 1,$$
(5.19)

namely $(K_j)_{j=1}^N$ covers the sphere \mathbb{S}^{n-1} and the closed annulus $1 - \delta \leq |x| \leq 1$, and $(\psi_j)_{j=1}^N$ is a resolution of unity for \mathbb{S}^{n-1} and for the closed annulus.

The diffeomorphisms \mathbf{f}_j can be defined in several ways; here we choose a simple construction that is convenient for the $H^{s,r}(\mathbb{R}^n_+)$ norms, i.e., we essentially introduce local spherical coordinates y, where the last coordinate y_n gives the signed distance of the point x from \mathbb{S}^{n-1} , and the coordinates $(y_1, \ldots, y_{n-1}) = y'$ determine the unit vector x/|x|, using the stereographical projection of x from the origin to the hyperplane $y_n = -1$. More precisely, we define \mathbf{f}_j as the composition $\mathbf{f}_j(x) := \mathbf{f}(\mathbf{R}_j x)$ of a rotation \mathbf{R}_j of \mathbb{R}^n mapping the center p_j of the ball K_j into the "South Pole" $p_0 := (0, \ldots, 0, -1)$ with the function

$$\mathbf{f}(x) := \left(-\frac{x'}{x_n}, 1 - |x|\right),\tag{5.20}$$

where $x' = (x_1, \ldots, x_{n-1})$. The function **f** is well defined and smooth in the half space $\{x \in \mathbb{R}^n : x_n < 0\}$, bijective onto the half space $\{y \in \mathbb{R}^n : y_n < 1\}$, with inverse

$$\mathbf{g}(y) := (1 - y_n)(1 + |y'|^2)^{-\frac{1}{2}}(y', -1), \tag{5.21}$$

because, if $\mathbf{f}(x) = y$, then $|x| = 1 - y_n$, $x' = -x_n y'$, whence $(1 - y_n)^2 = |x|^2 = |x'|^2 + x_n^2 = x_n^2(1 + |y'|^2)$; for $x_n < 0$, one obtains $x = \mathbf{g}(y)$. We also denote

$$\mathbf{g}_j(y) := \mathbf{R}_j^{-1} \mathbf{g}(y). \tag{5.22}$$

For $\delta < 1/2$, the ball $B_{2\delta}(p_0)$ is contained in $\{x \in \mathbb{R}^n : x_n < 0\}$, therefore \mathbf{f}_j is well defined in the ball K_j , with image $A_j := \mathbf{f}_j(K_j) = \mathbf{f}(B_{2\delta}(p_0)) =: A_0$. One has $\mathbf{f}_j(p_j) = \mathbf{f}(p_0) = 0$. The Jacobian matrix satisfies $D\mathbf{f}(p_0) = I$ and $|D\mathbf{f}(x) - I| \leq C\delta$ for all $x \in B_{2\delta}(p_0)$, where I is the identity matrix and C is a constant independent of δ . Hence, given any $\varepsilon > 0$, for N sufficiently large and δ sufficiently small one has $D\mathbf{f}_j(p_j) = \mathbf{R}_j$ and $|D\mathbf{f}_j(x) - D\mathbf{f}_j(p_j)| \leq \varepsilon$ for all $x \in K_j$. Note that $D\mathbf{f}_j(p_j)$ is the rotation matrix \mathbf{R}_j , and not the identity matrix appearing in item (vii) above; however, this gives the same result for the application to differential operators. By construction, one has $\mathbf{f}_j(B_1 \cap K_j) = A_0 \cap \mathbb{R}^n_+$ and $\mathbf{f}_j(\mathbb{S}^{n-1} \cap K_j) = A_0 \cap \mathbb{R}^n_0$.

If two balls K_j, K_ℓ have nonempty intersection, for $x \in K_j \cap K_\ell$ one has $\mathbf{f}_j(x) \in \mathbf{f}_j(K_j \cap K_\ell) \subseteq \mathbf{f}_j(K_j) = A_0$ and $\mathbf{f}_\ell(x) \in \mathbf{f}_\ell(K_j \cap K_\ell) \subseteq A_0$. If, in addition, $x \in B_1 \cap K_j \cap K_\ell$, then $\mathbf{f}_j(x) \in \mathbb{R}^n_+ \cap \mathbf{f}_j(K_j \cap K_\ell) \subseteq \mathbb{R}^n_+ \cap A_0$ and $\mathbf{f}_\ell(x) \in \mathbb{R}^n_+ \cap \mathbf{f}_\ell(K_j \cap K_\ell) \subseteq \mathbb{R}^n_+ \cap A_0$. The transition map

$$\mathbf{f}_{\ell} \circ \mathbf{g}_j = \mathbf{f}_{\ell} \circ \mathbf{f}_j^{-1} : \mathbf{f}_j(K_j \cap K_{\ell}) \to \mathbf{f}_{\ell}(K_j \cap K_{\ell})$$

can be explicitly calculated using (5.20), (5.21) and the basic property $|\mathbb{R}_{\ell}\mathbb{R}_{j}^{-1}x| = |x|$ of rotation matrices. We find that

$$\mathbf{f}_{\ell} \circ \mathbf{g}_{j}(y) = (\mathbf{T}_{\ell j}(y'), y_{n}), \quad \mathbf{T}_{\ell j}(y') = -\frac{[\mathbf{R}_{\ell}\mathbf{R}_{j}^{-1}(y', -1)]'}{[\mathbf{R}_{\ell}\mathbf{R}_{j}^{-1}(y', -1)]_{n}}$$
(5.23)

for all $y = (y', y_n)$ in the domain $\mathbf{f}_j(K_j \cap K_\ell)$ of $\mathbf{f}_\ell \circ \mathbf{g}_j$. Formula (5.23) shows the remarkable property that the transition map $\mathbf{f}_\ell \circ \mathbf{g}_j$ does not move the last variable y_n , and it acts on y' as a function independent of y_n .

To estimate the $H^{s,r}(\mathbb{R}^n_+)$ norms of products and compositions of functions, it is convenient to extend \mathbf{f} , \mathbf{g} and $\mathbf{f}_{\ell} \circ \mathbf{g}_j$ outside their original domains. The first order Taylor expansion of $\mathbf{f}(x)$ around p_0 is $x - p_0$; therefore we define

$$\mathbf{f}(x) := x - p_0 + \varphi_\delta(x - p_0)(\mathbf{f}(x) - x + p_0) \quad \forall x \in \mathbb{R}^n,$$
(5.24)

where $\varphi_{\delta}(x-p_0) := \varphi(\frac{|x-p_0|}{2\delta})$ for some cut-off function $\varphi \in C^{\infty}(\mathbb{R})$ such that $\varphi(t) = 1$ for $t \leq 1$, $\varphi(t) = 0$ for $t \geq 2$, and $0 \leq \varphi \leq 1$. Taking δ sufficiently small, the function $\tilde{\mathbf{f}}$ is a diffeomorphism of \mathbb{R}^n because $|D\tilde{\mathbf{f}}(x) - I| \leq 1/2$ for all $x \in \mathbb{R}^n$. Also, $\tilde{\mathbf{f}} = \mathbf{f}$ in $B_{2\delta}(p_0)$. We define $\tilde{\mathbf{g}}$ as the inverse of $\tilde{\mathbf{f}}$ in (5.24). We also define $\tilde{\mathbf{f}}_j := \tilde{\mathbf{f}} \circ \mathbf{R}_j$ and $\tilde{\mathbf{g}}_j$ as its inverse function.

Concerning the transition map, of course $\hat{\mathbf{f}}_{\ell} \circ \tilde{\mathbf{g}}_j$ is an extension of (5.23); however, it is convenient to introduce another, more explicit extension, preserving the property of not moving y_n and acting on y' independently on y_n . The affine function $y' \mapsto \mathbb{R}_{\ell}\mathbb{R}_j^{-1}(y',-1) = \mathbb{R}_{\ell}\mathbb{R}_j^{-1}(0,-1) + \mathbb{R}_{\ell}\mathbb{R}_j^{-1}(y',0)$ is defined for all $y' \in \mathbb{R}^{n-1}$, and its value at y' = 0 is

$$\mathbf{R}_{\ell}\mathbf{R}_{j}^{-1}(0,-1) = \mathbf{R}_{\ell}\mathbf{R}_{j}^{-1}p_{0} = \mathbf{R}_{\ell}p_{j} = \mathbf{R}_{\ell}p_{\ell} + \mathbf{R}_{\ell}(p_{j}-p_{\ell}) = p_{0} + \mathbf{R}_{\ell}(p_{j}-p_{\ell}),$$
(5.25)

with $|\mathbf{R}_{\ell}(p_j - p_{\ell})| = |p_j - p_{\ell}| < 4\delta$ because p_j, p_{ℓ} are the centers of the two intersecting balls K_j, K_{ℓ} of radius 2δ . We write the rotation matrix $\mathbf{R}_{\ell}\mathbf{R}_j^{-1}$ as $\begin{pmatrix}\mathbf{A} & \mathbf{b} \\ \mathbf{c} & \mathbf{d} \end{pmatrix}$, where \mathbf{A} is an $(n-1) \times (n-1)$ matrix, \mathbf{b} is a column with n-1 components, \mathbf{c} is a row with n-1 components, and \mathbf{d} is a scalar. Thus

$$\begin{aligned} \mathbf{A}y' &= [\mathbf{R}_{\ell}\mathbf{R}_{j}^{-1}(y',0)]', \qquad \mathbf{b} = [\mathbf{R}_{\ell}\mathbf{R}_{j}^{-1}(0,1)]' = -[\mathbf{R}_{\ell}(p_{j}-p_{\ell})]', \\ \mathbf{c}y' &= [\mathbf{R}_{\ell}\mathbf{R}_{j}^{-1}(y',0)]_{n}, \qquad \mathbf{d} = [\mathbf{R}_{\ell}\mathbf{R}_{j}^{-1}(0,1)]_{n} = 1 - [\mathbf{R}_{\ell}(p_{j}-p_{\ell})]_{n} \end{aligned}$$
(5.26)

for all $y' \in \mathbb{R}^{n-1}$, where we have used (5.25) and the fact that $p'_0 = 0, (p_0)_n = -1$. Hence $T_{\ell j}(y')$ in (5.23) becomes

$$\mathsf{T}_{\ell j}(y') = \frac{-\mathsf{b} + \mathsf{A}y'}{\mathsf{d} - \mathsf{c}y'}.$$
(5.27)

Lemma 5.3. Let $\delta < 1/8$. Then the scalar d and the matrix Ad – bc are invertible, one has

$$|\mathbf{d} - 1| < 4\delta, \quad |\mathbf{c}| = |\mathbf{b}| < 4\delta, \quad |(\mathbf{A}\mathbf{d} - \mathbf{b}\mathbf{c})^{-1}y'| \le \lambda |y'| \quad \forall y' \in \mathbb{R}^{n-1}, \quad \lambda := \mathbf{d}^{-1}, \quad (5.28)$$

and, for all $|y'| \leq 1$, one has the converging Taylor expansion

$$\frac{-\mathbf{b} + \mathbf{A}y'}{\mathbf{d} - \mathbf{c}y'} = -\lambda\mathbf{b} + \lambda^2(\mathbf{A}\mathbf{d} - \mathbf{b}\mathbf{c})y'(1 + \mathbf{S}(y')), \quad \mathbf{S}(y') := \sum_{m=1}^{\infty} (\lambda\mathbf{c}y')^m.$$
(5.29)

Proof. By (5.26), the vector $(\mathbf{b}, \mathbf{d} - 1)$ is the opposite of $\mathbf{R}_{\ell}(p_j - p_{\ell})$, and its norm is $|(\mathbf{b}, \mathbf{d} - 1)| = |\mathbf{R}_{\ell}(p_j - p_{\ell})| = |p_j - p_{\ell}| < 4\delta$. This gives the bound for $|\mathbf{b}|, |\mathbf{d} - 1|$ in (5.28). Since $\mathbf{R}_{\ell}\mathbf{R}_j^{-1}$ is a rotation matrix, all its columns and rows have unitary norm. In particular, taking the *n*th row and *n*th column, one has $|\mathbf{b}|^2 + \mathbf{d}^2 = 1$ and $|\mathbf{c}|^2 + \mathbf{d}^2 = 1$, whence $|\mathbf{c}| = |\mathbf{b}|$.

To study the matrix Ad - bc, we observe that

$$\begin{pmatrix} \mathbf{A} & \mathbf{b} \\ \mathbf{c} & \mathbf{d} \end{pmatrix} \begin{pmatrix} \mathbf{d}I' & 0 \\ -\mathbf{c} & 1 \end{pmatrix} = \begin{pmatrix} \mathbf{A}\mathbf{d} - \mathbf{b}\mathbf{c} & \mathbf{b} \\ 0 & \mathbf{d} \end{pmatrix},$$
(5.30)

where I' is the $(n-1) \times (n-1)$ identity matrix. The first matrix in (5.30) is the rotation $\mathbb{R}_{\ell}\mathbb{R}_{j}^{-1}$, whose determinant is 1, while the determinant of the second matrix is d^{n-1} . Hence the determinant of the third matrix in (5.30) is d^{n-1} , which is nonzero, and this is also the product of d with the determinant of Ad – bc. This proves that the matrix Ad – bc is invertible.

Now let $z' \in \mathbb{R}^{n-1}$. We consider the product of (5.30) with the vector (z', 0), and we calculate its norm. The matrix in the RHS of (5.30) times the vector (z', 0) gives $((\mathsf{Ad} - \mathsf{bc})z', 0)$, whose norm is $|(\mathsf{Ad} - \mathsf{bc})z'|$. The matrix product in the LHS of (5.30) times the same vector (z', 0) gives $\mathsf{R}_{\ell}\mathsf{R}_{j}^{-1}(\mathsf{d}z', -\mathsf{c}z')$, whose norm is equal to the norm of $(\mathsf{d}z', -\mathsf{c}z')$, which is $\geq \mathsf{d}|z'|$. This proves that $|(\mathsf{Ad} - \mathsf{bc})z'| \geq \mathsf{d}|z'|$ for all $z' \in \mathbb{R}^{n-1}$. Since $\mathsf{Ad} - \mathsf{bc}$ is invertible, taking $z' = (\mathsf{Ad} - \mathsf{bc})^{-1}y'$ gives the third inequality of (5.28).

Formula (5.29) is obtained by writing $(\mathbf{d} - \mathbf{c}y')^{-1} = \lambda(1 - \lambda \mathbf{c}y')^{-1}$ as a power series, which converges for $\lambda |\mathbf{c}y'| < 1$. One has $\lambda \leq (1 - 4\delta)^{-1} < 2$ and $|\mathbf{c}| < 4\delta$, whence $\lambda |\mathbf{c}y'| < 8\delta < 1$ for all $|y'| \leq 1$.

We define

$$\tilde{\mathbf{T}}_{\ell j}(y') := -\lambda \mathbf{b} + \lambda^2 (\operatorname{Ad} - \operatorname{bc}) y' \left(1 + \varphi_{\delta}(y') S(y') \right), \quad \varphi_{\delta}(y') := \varphi \left(\frac{|y'|}{4\delta} \right), \tag{5.31}$$

where $\varphi \in C^{\infty}(\mathbb{R})$ is a smooth cut-off function such that $\varphi(t) = 1$ for $t \leq 1$, $\varphi(t) = 0$ for $t \geq 2$, and $0 \leq \varphi \leq 1$. Hence $\tilde{T}_{\ell j} = T_{\ell j}$ where $\varphi_{\delta} = 1$, i.e., in the ball $|y'| \leq 4\delta$. Since $\delta < 1/4$, the set $A_0 = \mathbf{f}(B_{2\delta}(p_0))$ is contained in the ball $|y| < 4\delta$, and

$$(\tilde{\mathsf{T}}_{\ell j}(y'), y_n) = (\mathsf{T}_{\ell j}(y'), y_n) = \mathbf{f}_{\ell}(\mathbf{g}_j(y)) \quad \forall y = (y', y_n) \in A_0.$$
(5.32)

Lemma 5.4. There exists a universal constant $\delta_0 \in (0, 1/8)$ such that, for $0 < \delta \leq \delta_0$, if $K_j \cap K_\ell \neq \emptyset$, then the map $\tilde{\mathsf{T}}_{\ell j}$ in (5.31) is a C^{∞} diffeomorphism of \mathbb{R}^{n-1} that coincides with $\mathsf{T}_{\ell j}$ in (5.23) in the ball $|y'| \leq 4\delta$, and coincides in $|y'| \geq 8\delta$ with the affine map $y' \mapsto -\lambda \mathsf{b} + \lambda^2 (\mathsf{Ad} - \mathsf{bc})y'$.

Proof. Let us prove that $\tilde{\mathsf{T}}_{\ell j}$ is a bijection of \mathbb{R}^{n-1} . Given $x' \in \mathbb{R}^{n-1}$, one has $\tilde{\mathsf{T}}_{\ell j}(y') = x'$ if and only if $y' = \Phi(y')$, where $\Phi(y') := \lambda^{-2} (\mathsf{Ad} - \mathsf{bc})^{-1} (x' + \lambda \mathsf{b}) - y' \varphi_{\delta}(y') \mathsf{S}(y')$. The Jacobian matrix $D\Phi(y')$ vanishes for $|y'| \geq 8\delta$, and, for $\delta < 1/16$, one has

$$|D\Phi(y')z'| \le C\delta|z'| \quad \forall y', z' \in \mathbb{R}^{n-1},$$

where the constant C depends only on $\|\varphi'\|_{L^{\infty}(\mathbb{R})}$; we can assume that, say, $\|\varphi'\|_{L^{\infty}(\mathbb{R})} \leq 10$, so that C is a universal constant. Hence, for $C\delta \leq 1/2$, Φ is a contraction mapping on \mathbb{R}^{n-1} , and it has a unique fixed point. This proves that $\tilde{T}_{\ell j}$ is a bijection of \mathbb{R}^{n-1} . The C^{∞} regularity of $\tilde{T}_{\ell j}$ and of its inverse map $\tilde{T}_{\ell j}^{-1}$ follows from a standard implicit function argument.

Now that Sobolev spaces on domains are defined, we recall and/or prove some useful properties.

Lemma 5.5. Let $r \ge 0$ be any real number, let s = 0 or s = 1, and let $u \in H^{s+1,r}(\mathbb{R}^n_+)$. Then

$$\begin{aligned} \|u\|_{H^{s+1,r}(\mathbb{R}^{n}_{+})} &\leq C \|\Delta u\|_{H^{s-1,r}(\mathbb{R}^{n}_{+})} + C^{r+1} \|\Delta u\|_{H^{s-1,0}(\mathbb{R}^{n}_{+})} + C \|u(\cdot,0)\|_{H^{s+\frac{1}{2}+r}(\mathbb{R}^{n-1})} \\ &+ C^{r+1} \|u(\cdot,0)\|_{H^{s+\frac{1}{2}}(\mathbb{R}^{n-1})} + C^{r+1} \|u\|_{H^{s,0}(\mathbb{R}^{n}_{+})} \end{aligned}$$
(5.33)

where C > 0 is a universal constant.

Proof. See the Appendix.

Remark 5.6. The constant C in (5.33) is universal because we estimate the $H^{s+1,r}(\mathbb{R}^n_+)$ norm of u only in the cases s = 0 and s = 1; in general, for s arbitrarily varying in \mathbb{N}_0 , the constant C would depend on s in a nontrivial way, related to the operator norm of the extension operators involved in the proof. Here, however, we are only interested in high (and possibly non integer)

Sobolev regularity with respect to the variable x', corresponding to tangential directions, and in one or two derivatives with respect to the variable x_n , corresponding to the normal direction.

Lemma 5.7. On \mathbb{R}^n and \mathbb{R}^n_+ one has the following properties.

(i) Density. For s, r ∈ ℝ, the Schwartz class S(ℝⁿ) is dense in H^{s,r}(ℝⁿ), and the set S(ℝⁿ₊) := {f|_{ℝⁿ₊} : f ∈ S(ℝⁿ)} of the restrictions to ℝⁿ₊ of the Schwartz functions of ℝⁿ is dense in H^{s,r}(ℝⁿ₊).
(ii) Embedding. Let s, r ∈ ℝ, with s > 1/2, s + r > n/2. Then H^{s,r}(ℝⁿ) ⊂ L[∞](ℝⁿ) and H^{s,r}(ℝⁿ₊) ⊂ L[∞](ℝⁿ₊), with

$$\|v\|_{L^{\infty}(\mathbb{R}^{n})} \leq C_{s,r} \|v\|_{H^{s,r}(\mathbb{R}^{n})}, \quad \|u\|_{L^{\infty}(\mathbb{R}^{n}_{+})} \leq C_{s,r} \|u\|_{H^{s,r}(\mathbb{R}^{n}_{+})},$$
(5.34)

for all $v \in H^{s,r}(\mathbb{R}^n)$, all $u \in H^{s,r}(\mathbb{R}^n_+)$, where the constant $C_{s,r}$ depends on s,r. (*iii*) Partial derivatives. The partial derivatives (in the sense of distributions) satisfy

$$\|\partial_{x_k} u\|_{H^{s,r}(\mathbb{R}^n_+)} \le \|u\|_{H^{s,r+1}(\mathbb{R}^n_+)} \le \|u\|_{H^{s+1,r}(\mathbb{R}^n_+)}, \quad k = 1, \dots, n-1,$$
(5.35)

$$\|\partial_{x_n} u\|_{H^{s,r}(\mathbb{R}^n_+)} \le \|u\|_{H^{s+1,r}(\mathbb{R}^n_+)},\tag{5.36}$$

for all $u \in H^{s+1,r}(\mathbb{R}^n_+)$, for all $s, r \in \mathbb{R}$.

(iv) Equivalent norms. For every real $r \ge 0$, one has

$$\|u\|_{H^{0,r}(\mathbb{R}^{n}_{+})}^{2} = \int_{0}^{\infty} \|u(\cdot, x_{n})\|_{H^{r}(\mathbb{R}^{n-1})}^{2} dx_{n} \quad \forall u \in H^{0,r}(\mathbb{R}^{n}_{+}),$$
(5.37)

$$\frac{1}{2} \|u\|_{H^{1,r}(\mathbb{R}^n_+)}^2 \le \|u\|_{H^{0,r+1}(\mathbb{R}^n_+)}^2 + \|\partial_{x_n}u\|_{H^{0,r}(\mathbb{R}^n_+)}^2 \le \|u\|_{H^{1,r}(\mathbb{R}^n_+)}^2 \quad \forall u \in H^{1,r}(\mathbb{R}^n_+).$$
(5.38)

In particular, for r = 0,

$$\|u\|_{H^{0,0}(\mathbb{R}^{n}_{+})} = \|u\|_{L^{2}(\mathbb{R}^{n}_{+})}, \quad \frac{1}{2}\|u\|_{H^{1,0}(\mathbb{R}^{n}_{+})}^{2} \leq \sum_{|\alpha| \leq 1} \|\partial_{x}^{\alpha}u\|_{L^{2}(\mathbb{R}^{n}_{+})}^{2} \leq \|u\|_{H^{1,0}(\mathbb{R}^{n}_{+})}^{2}.$$
(5.39)

(v) Trace. Let $s, r \in \mathbb{R}$, with s > 1/2. On \mathbb{R}^n , the trace operator

$$T: H^{s,r}(\mathbb{R}^n) \to H^{s+r-\frac{1}{2}}(\mathbb{R}^{n-1}), \quad u \mapsto Tu = u(\cdot, 0)$$
(5.40)

is a well defined, bounded, linear operator, with

$$\|u(\cdot,0)\|_{H^{s+r-\frac{1}{2}}(\mathbb{R}^{n-1})} \le C_s \|u\|_{H^{s,r}(\mathbb{R}^n)}$$
(5.41)

for all $u \in H^{s,r}(\mathbb{R}^n)$ (first defined on $\mathcal{S}(\mathbb{R}^n)$), then extended to $H^{s,r}(\mathbb{R}^n)$ by density), where the constant C_s depends on s and it is independent of r. On \mathbb{R}^n_+ , for $s = 1, r \in \mathbb{R}$, the trace operator

$$T: H^{1,r}(\mathbb{R}^n_+) \to H^{r+\frac{1}{2}}(\mathbb{R}^{n-1}), \quad u \mapsto Tu = u(\cdot, 0)$$
 (5.42)

is a well defined, bounded, linear operator, with

$$\|u(\cdot,0)\|_{H^{r+\frac{1}{2}}(\mathbb{R}^{n-1})} \le C\|u\|_{H^{1,r}(\mathbb{R}^{n}_{+})}$$
(5.43)

for all $u \in H^{1,r}(\mathbb{R}^n_+)$, where $C = \sqrt{2\pi}$.

(vi) Product estimate. Let $r_0, r \in \mathbb{R}$, with $r_0 + 1 > n/2$ and $r \ge 0$. One has

$$\|uv\|_{H^{1,r}(\mathbb{R}^{n}_{+})} \leq C_{r_{0}}\|u\|_{H^{1,r_{0}}(\mathbb{R}^{n}_{+})}\|v\|_{H^{1,r}(\mathbb{R}^{n}_{+})} + C_{r}\|u\|_{H^{1,r}(\mathbb{R}^{n}_{+})}\|v\|_{H^{1,r_{0}}(\mathbb{R}^{n}_{+})}$$
(5.44)

for all $u, v \in H^{1,r}(\mathbb{R}^n_+) \cap H^{1,r_0}(\mathbb{R}^n_+)$, where C_{r_0} depends on r_0 and it is independent of r, while C_r depends on r, r_0 , and it is an increasing function of r. For $0 \le r \le r_0$, one also has

$$\|uv\|_{H^{1,r}(\mathbb{R}^{n}_{+})} \leq C_{r_{0}}\|u\|_{H^{1,r_{0}}(\mathbb{R}^{n}_{+})}\|v\|_{H^{1,r}(\mathbb{R}^{n}_{+})}$$
(5.45)

for all $u \in H^{1,r_0}(\mathbb{R}^n_+)$, $v \in H^{1,r}(\mathbb{R}^n_+)$. Estimates (5.44) and (5.45) can be written together as

$$\|uv\|_{H^{1,r}(\mathbb{R}^{n}_{+})} \leq C_{r_{0}}\|u\|_{H^{1,r_{0}}(\mathbb{R}^{n}_{+})}\|v\|_{H^{1,r}(\mathbb{R}^{n}_{+})} + (\chi_{r>r_{0}})C_{r}\|u\|_{H^{1,r}(\mathbb{R}^{n}_{+})}\|v\|_{H^{1,r_{0}}(\mathbb{R}^{n}_{+})},$$
(5.46)

where $(\chi_{r>r_0}) = 1$ if $r > r_0$, $(\chi_{r>r_0}) = 0$ if $r \le r_0$, C_r is increasing in r, and C_{r_0} does not depend on r. The same estimates also hold with \mathbb{R}^n_+ replaced by \mathbb{R}^n .

(vii) Multiplication by bounded functions with bounded derivatives. One has

$$\|fu\|_{H^{0,r}(\mathbb{R}^{n}_{+})} \leq 2\|f\|_{L^{\infty}(\mathbb{R}^{n}_{+})}\|u\|_{H^{0,r}(\mathbb{R}^{n}_{+})} + C_{r}\|f\|_{W^{0,b,\infty}(\mathbb{R}^{n}_{+})}\|u\|_{H^{0,0}(\mathbb{R}^{n}_{+})}$$
(5.47)

for all $f \in W^{0,b,\infty}(\mathbb{R}^n)$, $u \in H^{0,r}(\mathbb{R}^n_+)$, $r \ge 0$ real, where b is the smallest integer such that $b \ge r$,

$$||f||_{W^{0,b,\infty}(\mathbb{R}^{n}_{+})} := \underset{x_{n} \in (0,\infty)}{\operatorname{ess \, sup}} ||f(\cdot, x_{n})||_{W^{b,\infty}(\mathbb{R}^{n-1})},$$
(5.48)

 $W^{0,b,\infty}(\mathbb{R}^n_+)$ is the space of functions whose norm (5.48) is finite, and C_r is increasing in r. Also,

$$\|fu\|_{H^{1,r}(\mathbb{R}^n_+)} \le C_0 \|f\|_{W^{1,\infty}(\mathbb{R}^n_+)} \|u\|_{H^{1,r}(\mathbb{R}^n_+)} + C_r \|f\|_{W^{1,b,\infty}(\mathbb{R}^n_+)} \|u\|_{H^{1,0}(\mathbb{R}^n_+)}$$
(5.49)

for all $u \in H^{1,r}(\mathbb{R}^n_+)$, all $f \in W^{0,b,\infty}(\mathbb{R}^n_+)$ with $\partial_{x_n} f \in W^{0,b,\infty}(\mathbb{R}^n_+)$, where

$$\|f\|_{W^{1,b,\infty}(\mathbb{R}^n_+)} := \|f\|_{W^{0,b+1,\infty}(\mathbb{R}^n_+)} + \|\partial_{x_n}f\|_{W^{0,b,\infty}(\mathbb{R}^n_+)}$$
(5.50)

and $W^{1,b,\infty}(\mathbb{R}^n_+)$ is the space of functions whose norm (5.50) is finite.

(viii) Composition with functions of x'. Let $s \in \{0,1\}$, $r \ge 0$ real, $r_0 > (n-1)/2$. Let $f : \mathbb{R}^{n-1} \to \mathbb{R}^{n-1}$, f(x') = a + Ax' + g(x') be a diffeomorphisms of \mathbb{R}^{n-1} , with $a \in \mathbb{R}^{n-1}$, $A \in \operatorname{Mat}_{(n-1)\times(n-1)}(\mathbb{R})$, $g \in H^{r+1+r_0+s}(\mathbb{R}^{n-1})$. One has

$$\|x \mapsto u(f(x'), x_n)\|_{H^{s,r}(\mathbb{R}^n_+)} \le C_{r,f}(\|u\|_{H^{s,r}(\mathbb{R}^n_+)} + \|g\|_{H^{1+r_0+r+s}(\mathbb{R}^{n-1})}\|u\|_{H^{s,0}(\mathbb{R}^n_+)})$$
(5.51)

for all $u \in H^{s,r}(\mathbb{R}^n_+)$, where $C_{r,f}$ depends on $r, |A|, ||g||_{H^{1+r_0}(\mathbb{R}^{n-1})}$, and it is increasing in r.

Proof. See the Appendix.

The next lemma deals with Sobolev spaces $H^{s}(\mathbb{S}^{d})$ on the sphere.

Lemma 5.8. The Sobolev spaces $H^{s}(\mathbb{S}^{d})$ defined in (5.18) with $\Gamma = \mathbb{S}^{d}$, d = n - 1, satisfy the following properties.

(i) Interpolation inequality. Let $s_0, s_1 \in \mathbb{R}$, $s = s_0(1 - \vartheta) + s_1\vartheta$, $0 < \vartheta < 1$. Then

$$\|u\|_{H^{s}(\mathbb{S}^{d})} \leq \|u\|_{H^{s_{0}}(\mathbb{S}^{d})}^{1-\vartheta} \|u\|_{H^{s_{1}}(\mathbb{S}^{d})}^{\vartheta}$$
(5.52)

for all $u \in H^{s_1}(\mathbb{S}^d)$.

(ii) Product estimate. Let $s, s_0 \in \mathbb{R}$, with $s \ge 0$ and $s_0 > d/2$. Then

$$\|uv\|_{H^{s}(\mathbb{S}^{d})} \leq C_{s_{0}} \|u\|_{H^{s_{0}}(\mathbb{S}^{d})} \|v\|_{H^{s}(\mathbb{S}^{d})} + (\chi_{s>s_{0}})C_{s} \|u\|_{H^{s}(\mathbb{S}^{d})} \|v\|_{H^{s_{0}}(\mathbb{S}^{d})},$$
(5.53)

where $(\chi_{s>s_0}) = 1$ if $s > s_0$, $(\chi_{s>s_0}) = 0$ if $s \le s_0$, the constant C_{s_0} depends on s_0 and it is independent of s, while C_s depends on s, s_0 and it is increasing in s.

(iii) Power estimate. Let s, s_0 be as in (ii). Then

$$\begin{aligned} \|u^{m}\|_{H^{s}(\mathbb{S}^{d})} &\leq (C_{s_{0}}\|u\|_{H^{s_{0}}(\mathbb{S}^{d})})^{m-1}\|u\|_{H^{s}(\mathbb{S}^{d})} \\ &+ (\chi_{s>s_{0}})(m-1)C_{s}(C_{s_{0}}\|u\|_{H^{s_{0}}(\mathbb{S}^{d})})^{m-2}\|u\|_{H^{s_{0}}(\mathbb{S}^{d})}\|u\|_{H^{s}(\mathbb{S}^{d})} \end{aligned}$$
(5.54)

for all integers $m \geq 2$, where C_{s_0}, C_s are the constants in (ii).

(iv) Tangential gradient. Let $s \in \mathbb{R}$, with $s \geq 0$. Then

$$\|\nabla_{\mathbb{S}^d} u\|_{H^s(\mathbb{S}^d)} \le C_0 \|u\|_{H^{s+1}(\mathbb{S}^d)} + C_s \|u\|_{H^1(\mathbb{S}^d)},\tag{5.55}$$

with C_0 independent of s and C_s increasing in s.

(v) Embedding. Let $s_0 \in \mathbb{R}$, with $s_0 > d/2$. Then

$$\|u\|_{L^{\infty}(\mathbb{S}^d)} \le C_{s_0} \|u\|_{H^{s_0}(\mathbb{S}^d)}.$$
(5.56)

Proof. See the Appendix.

5.3 Estimates in higher norms

In subsection 5.1, see Lemma 5.1 and Proposition 5.2, we have studied the low norm of the solution $\tilde{\Phi}$ of problem (3.13) expanded as power series $\tilde{\Phi} = u = \sum u_n$, with u_n defined in (5.3), (5.4). Now in this subsection we study the higher Sobolev tangential regularity of u and $\langle x, \nabla u \rangle$ on \mathbb{S}^2 . To this aim, we use partitions of unity, local rectifications, smooth cut-off functions, and Sobolev spaces $H^s(\mathbb{S}^2)$ and $H^{s,r}(\mathbb{R}^3_+)$ defined in subsection 5.2, studying the functions u_n only at points of B_1 sufficiently close to the boundary \mathbb{S}^2 .

Remark 5.9. In general, the function $\mathcal{E}_0 h$ (appearing in the definition (5.4) of u_n for $n \geq 1$) is not in $H^2(B_1)$ even if h is C^{∞} on \mathbb{S}^2 , because of the singularity of homogeneous functions at the origin: at any point x = ry with 0 < r < 1 and $y \in \mathbb{S}^2$, one has $\partial_x^{\alpha}(\mathcal{E}_0 h)(x) = r^{-2}\partial_x^{\alpha}(\mathcal{E}_0 h)(y)$ for all multi-indices $|\alpha| = 2$, and the $L^2(B_1)$ integrability of second order partial derivatives is lost. For this reason, to study higher Sobolev regularity we only consider points of B_1 sufficiently close to the boundary \mathbb{S}^2 . See also the related discussion in Remark 4.8.

Let $\delta, K_j, \psi_j, \mathbf{f}_j, \mathbf{f}, \mathbf{g}, \mathbf{R}_j, \mathbf{g}_j, p_j, A_0, p_0, \mathbb{R}^3_+, \mathbb{R}^3_0$ be as in subsection 5.2. We decompose the annulus $1-\delta < |x| < 1-(\delta/2)$ into a sequence of annuli $\rho_{k+1} < |x| < \rho_k$, where $\rho_0 = 1-(\delta/2)$, the sequence (ρ_k) is decreasing and it converges to $1-\delta$, and for each annulus we introduce a smooth, radial cut-off function ζ_k in the following way. Let $\varphi \in C^{\infty}(\mathbb{R},\mathbb{R})$ with $\varphi = 0$ on $(-\infty,0], \varphi = 1$ on $[1,\infty)$, and $0 \le \varphi \le 1$. We define

$$\rho_k := 1 - \frac{\delta}{2} \sum_{\ell=0}^k \frac{1}{2^\ell}, \quad \zeta_k(x) := \varphi \Big(\frac{|x| - \rho_{k+1}}{\rho_k - \rho_{k+1}} \Big) \quad \forall x \in \mathbb{R}^3, \quad k \in \mathbb{N}_0.$$
(5.57)

Thus $\zeta_k(x) = 0$ for $|x| \le \rho_{k+1}$, $\zeta_k(x) = 1$ for $|x| \ge \rho_k$, and

$$|\zeta_k(x)| \le 1, \quad |\partial_x^{\alpha} \zeta_k(x)| \le C_{\alpha} 2^{k|\alpha|} \quad \forall x \in \mathbb{R}^3, \quad k \in \mathbb{N}_0, \quad \alpha \in \mathbb{N}_0^3, \tag{5.58}$$

where the constant C_{α} depends on the length $|\alpha|$ of the multi-index α (and also on φ and δ , which we consider as fixed). We note that $\zeta_{k+1} = 1$ where $\zeta_k \neq 0$, i.e.,

$$\zeta_{k+1} = 1 \text{ on } \operatorname{supp}(\zeta_k), \quad \forall k \in \mathbb{N}_0.$$
(5.59)

We also define

$$\zeta^*(x) := \varphi\left(\frac{|x| - 1 + 2\delta}{\delta}\right) \quad \forall x \in \mathbb{R}^3,$$
(5.60)

and we note that $\zeta^*(x) = 0$ for $|x| \leq 1 - 2\delta$, $\zeta^*(x) = 1$ for $|x| \geq 1 - \delta$, and, in particular, $\zeta^* = 1$ on the support of ζ_k , for all $k \in \mathbb{N}_0$.

For $s \in \{0, 1, 2\}, r \in \mathbb{R}, k \in \mathbb{N}_0$, for any function f, we denote

$$\|f\|_{X_k^{s,r}} := \sum_{j=1}^N \|(f\zeta_k \psi_j) \circ \mathsf{g}_j\|_{H^{s,r}(\mathbb{R}^3_+)}.$$
(5.61)

Before studying the norm (5.61) of u_n for general r, we consider the case of low norm r = 0, which is directly related to Lemma 5.1. For notation convenience, we define g_n also for n = 0 by setting $g_0 := 0$.

Lemma 5.10. The functions u_n, g_n defined in (5.3), (5.4), (5.5) satisfy

$$\|u_n\|_{X_{\iota}^{1,0}} \le C_0 2^k \mu z^n, \quad \|g_n\|_{X_{\iota}^{0,0}} \le \mu z^n \tag{5.62}$$

for all $n, k \in \mathbb{N}_0$, where

$$\mu := \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)}, \quad z := (1+C)2\tau, \tag{5.63}$$

 C_0 does not depend on n, k, h, ψ , and C, τ are given by Lemma 5.1.

Proof. For $j \in \{1, ..., N\}$, the function $\psi_j \circ \mathbf{g}_j$ vanishes in $\mathbb{R}^3_+ \setminus A_0$, and, by (5.58), the function $a_{kj} := (\zeta_k \psi_j) \circ \mathbf{g}_j$ satisfies

$$\|a_{kj}\|_{L^{\infty}(\mathbb{R}^{3}_{+})} \leq 1, \quad \|\partial_{y}^{\alpha}a_{kj}\|_{L^{\infty}(\mathbb{R}^{3}_{+})} \leq C_{\alpha}2^{k|\alpha|}$$
(5.64)

for all $\alpha \in \mathbb{N}_0^3$. Hence, by (5.39),

$$\begin{split} \|(u_n\zeta_k\psi_j)\circ \mathbf{g}_j\|_{H^{1,0}(\mathbb{R}^3_+)} &\leq 2\sum_{|\alpha|\leq 1} \|\partial_y^{\alpha}\{(u_n\zeta_k\psi_j)\circ \mathbf{g}_j\}\|_{L^2(\mathbb{R}^3_+)} \\ &= 2\sum_{|\alpha|\leq 1} \|\partial_y^{\alpha}\{(u_n\zeta_k\psi_j)\circ \mathbf{g}_j\}\|_{L^2(A_0\cap\mathbb{R}^3_+)} \\ &\leq 2\sum_{|\alpha|\leq 1} \|\partial_y^{\alpha}(u_n\circ \mathbf{g}_j)\|_{L^2(A_0\cap\mathbb{R}^3_+)} + C_12^k\|u_n\circ \mathbf{g}_j\|_{L^2(A_0\cap\mathbb{R}^3_+)} \\ &\leq C_0\sum_{|\beta|\leq 1} \|\partial_x^{\beta}u_n\|_{L^2(K_j\cap B_1)} + C_02^k\|u_n\|_{L^2(K_j\cap B_1)}, \end{split}$$

for some constants C_1, C_0 independent of n, k, h, ψ ; in the last inequality we have used the fact that \mathbf{g}_j is a diffeomorphism of $A_0 \cap \mathbb{R}^3_+$ onto $K_j \cap B_1$. Taking the sum over $j = 1, \ldots, N$ and using (5.7), we obtain the first inequality in (5.62). The second inequality in (5.62) is proved similarly, using (5.39), (5.9), (5.10); there is no factor 2^k because the derivatives of ζ_k are not involved. \Box

Lemma 5.11. For all $k \in \mathbb{N}_0$, $n \in \mathbb{N}$, the functions u_n, g_n defined in (5.4), (5.5) satisfy

$$\|u_n\|_{X_k^{2,0}} \le C_0 2^{3k} \mu z^n + C_0 \|g_n\|_{X_k^{1,0}},$$
(5.65)

where μ, z are defined in (5.63). For $0 \le b - 1 < r \le b$, with b integer and r real, one has

$$\|u_n\|_{X_k^{2,r}} \le C_0 \|g_n\|_{X_k^{1,r}} + C_r \|g_n\|_{X_k^{1,0}} + C_{r,k} \mu z^n + C_{r,k} \Big(\sum_{\ell=1}^{b-1} \|g_n\|_{X_{k+\ell}^{1,r-\ell}} + \|g_n\|_{X_{k+b}^{1,0}} \Big), \quad (5.66)$$

where C_r depends on r, $C_{r,k}$ depends on r, k, both C_r and $C_{r,k}$ are increasing functions of r, and C_0 is independent of r, k.

Proof. Let $j \in \{1, ..., N\}$, and consider the diffeomorphism $\mathbf{f}_j : B_1 \cap K_j \to A_0 \cap \mathbb{R}^3_+$ and its inverse \mathbf{g}_j . Since $-\Delta u_n = \operatorname{div} g_n$, taking test functions in $C_c^{\infty}(B_1 \cap K_j)$, recalling the transformation rule for the gradient

$$(\nabla u_n) \circ \mathbf{g}_j(y) = [D\mathbf{g}_j(y)]^{-T} \nabla (u_n \circ \mathbf{g}_j)(y), \quad y \in A_0 \cap \mathbb{R}^3_+,$$
(5.67)

and following the same kind of calculations leading to (3.14), (3.13), one finds that

$$-\operatorname{div}\{\mathsf{P}\nabla(u_n\circ\mathsf{g}_j)\}=\operatorname{div}\{(\det D\mathsf{g}_j)(D\mathsf{g}_j)^{-1}(g_n\circ\mathsf{g}_j)\}\quad\text{in }A_0\cap\mathbb{R}^3_+,\tag{5.68}$$

where

$$\mathbf{P}(y) := \mathbf{p}(y)\mathbf{Q}(y), \quad \mathbf{p}(y) := \det(D\mathbf{g}_j(y)) = \det(D\mathbf{g}(y)), \tag{5.69}$$

$$\mathbf{Q}(y) := [D\mathbf{g}_j(y)]^{-1} [D\mathbf{g}_j(y)]^{-T} = [D\mathbf{g}(y)]^{-1} [D\mathbf{g}(y)]^{-T}$$
(5.70)

in $A_0 \cap \mathbb{R}^3_+$. The last identities in (5.69) and (5.70) hold because $g_j = \mathbb{R}_j^{-1} \circ g$ and \mathbb{R}_j is a rotation. By (5.21), in A_0 one has

$$\mathbf{Q}(y) = \begin{bmatrix} (1+y_1^2)\mathbf{q}(y) & y_1y_2\mathbf{q}(y) & 0\\ y_1y_2\mathbf{q}(y) & (1+y_2^2)\mathbf{q}(y) & 0\\ 0 & 0 & 1 \end{bmatrix}, \quad \mathbf{q}(y) \coloneqq \frac{1+y_1^2+y_2^2}{(1-y_3)^2}, \quad \mathbf{p}(y) = \frac{(1-y_3)^2}{(1+y_1^2+y_2^2)^{\frac{3}{2}}}.$$
 (5.71)

Since the (3,3) entry of the matrix Q is 1, and since derivatives with respect to y_3 play a separate role, it is convenient to work with Q instead of P. Dividing by p, (5.68) becomes

$$-\operatorname{div}\{\operatorname{Q}\nabla(u_n\circ\operatorname{\mathbf{g}}_j)\}=\operatorname{p}^{-1}\langle\nabla\operatorname{p},\operatorname{Q}\nabla(u_n\circ\operatorname{\mathbf{g}}_j)\rangle+\operatorname{p}^{-1}\operatorname{div}\{\operatorname{p}(D\operatorname{\mathbf{g}}_j)^{-1}(g_n\circ\operatorname{\mathbf{g}}_j)\}\quad\text{in }A_0\cap\mathbb{R}^3_+.$$
 (5.72)

Given any two functions v, a, one has

$$\operatorname{div}(\mathbf{Q}\nabla(va)) = a\operatorname{div}(\mathbf{Q}\nabla v) + 2\langle \mathbf{Q}\nabla a, \nabla v \rangle + v\operatorname{div}(\mathbf{Q}\nabla a).$$
(5.73)

At the point y = 0, one has Dg(0) = I (identity matrix), whence Q(0) = I, p(0) = 1. To take advantage of the fact that Q(y) is close to Q(0) = I for y close to 0, we decompose Q in the l.h.s. of (5.73) as the sum I + (Q - I), and (5.73) becomes

$$\Delta(va) = \operatorname{div}\{(I - \mathsf{Q})\nabla(va)\} + a\operatorname{div}(\mathsf{Q}\nabla v) + 2\langle\mathsf{Q}\nabla a, \nabla v\rangle + v\operatorname{div}(\mathsf{Q}\nabla a).$$
(5.74)

For $k \in \mathbb{N}_0$, applying (5.74) with $v = u_n \circ \mathbf{g}_j$, $a = (\zeta_k \psi_j) \circ \mathbf{g}_j$, and using (5.72) to substitute the second term in the r.h.s. of (5.74), we get

$$\Delta w_{nkj} = \operatorname{div}\{(I - \mathbf{Q})\nabla w_{nkj}\} - a_{kj}\mathbf{p}^{-1}\langle \nabla \mathbf{p}, \mathbf{Q}\nabla v_{nj}\rangle - a_{kj}\mathbf{p}^{-1}\operatorname{div}\{\mathbf{p}(D\mathbf{g}_j)^{-1}(g_n \circ \mathbf{g}_j)\} + 2\langle \mathbf{Q}\nabla a_{kj}, \nabla v_{nj}\rangle + v_{nj}\operatorname{div}(\mathbf{Q}\nabla a_{kj})$$
(5.75)

in $A_0 \cap \mathbb{R}^3_+$, where

$$v_{nj} := u_n \circ \mathbf{g}_j, \quad a_{kj} := (\zeta_k \psi_j) \circ \mathbf{g}_j, \quad w_{nkj} := a_{kj} v_{nj} = (u_n \zeta_k \psi_j) \circ \mathbf{g}_j.$$
(5.76)

Since ψ_j is compactly supported in K_j , the support of $\psi_j \circ \mathbf{g}_j$ is a compact set contained in A_0 . Since $\psi_j \circ \mathbf{g}_j$ appears in every term of (5.75), identity (5.75) holds in the whole half-space \mathbb{R}^3_+ (it becomes "0 = 0" in $\mathbb{R}^3_+ \setminus A_0$), where w_{nkj} is extended to \mathbb{R}^3_+ by setting $w_{nkj} = 0$ in $\mathbb{R}^3_+ \setminus A_0$, and \mathbb{Q} , \mathbf{p} in (5.70), (5.69) are defined in \mathbb{R}^3 if we replace \mathbf{g} with its extension $\tilde{g} = \tilde{\mathbf{f}}^{-1}$, see (5.24); with a little abuse of notation, we denote the extensions without changing letters. Note that, by (5.24), one has $\tilde{\mathbf{f}}(x) = x - p_0$ for all x outside the ball $B_{4\delta}(p_0)$, and therefore $\mathbf{p} = 1$, $\mathbb{Q} = I$ outside the set $\mathbf{f}(B_{4\delta}(p_0))$, which is a neighborhood of $A_0 = \mathbf{f}(B_{2\delta}(p_0))$.

Since $u_n \in H^1_0(B_1)$, since ζ_k, ψ_j are smooth, and ψ_j is compactly supported in K_j , one has

$$w_{nkj} \in H^1_0(\mathbb{R}^3_+).$$
 (5.77)

Then, by Lemma 5.5, for every real $r \ge 0$ one has

$$\|w_{nkj}\|_{H^{2,r}(\mathbb{R}^3_+)} \le C \|\Delta w_{nkj}\|_{H^{0,r}(\mathbb{R}^3_+)} + C^{r+1} \|\Delta w_{nkj}\|_{L^2(\mathbb{R}^3_+)} + C^{r+1} \|w_{nkj}\|_{H^1(\mathbb{R}^3_+)},$$
(5.78)

where C > 0 is the universal constant in (5.33), and Δw_{nkj} is given by (5.75). Now we estimate each term in the right-hand side of (5.75).

First term. By (5.71), the third column and the third row of the matrix $(I - \mathbf{Q}(y))$ vanish in $A_0 \cap \mathbb{R}^3_+$, and div $\{(I - \mathbf{Q}) \nabla w_{nkj}\}$ is the sum of terms of the form $\partial_{y_i}(c(y)\partial_{y_\ell}w_{nkj})$ with $i, \ell \in \{1, 2\}$, without derivatives of w_{nkj} with respect to y_3 . Hence, by (5.35) and (5.47), one has

$$\begin{aligned} \|\operatorname{div}\{(I-\mathsf{Q})\nabla w_{nkj}\}\|_{H^{0,r}(\mathbb{R}^{3}_{+})} &\leq C_{0}\|I-\mathsf{Q}\|_{L^{\infty}(\mathbb{R}^{3}_{+})}\|w_{nkj}\|_{H^{0,r+2}(\mathbb{R}^{3}_{+})} \\ &+ C_{r}\|I-\mathsf{Q}\|_{W^{0,b,\infty}(\mathbb{R}^{3}_{+})}\|w_{nkj}\|_{H^{0,1}(\mathbb{R}^{3}_{+})}, \end{aligned}$$

where C_0, C_r, b are given by the application of (5.47). As explained in Section 5.2, given any $\varepsilon > 0$, the radius 2δ of the balls K_1, \ldots, K_N can be chosen sufficiently small to have $C_0 ||I - \mathbb{Q}||_{L^{\infty}(\mathbb{R}^3_+)} \leq \varepsilon$; note that such a δ is independent on r. Hence

$$\|\operatorname{div}\{(I-\mathbf{Q})\nabla w_{nkj}\}\|_{H^{0,r}(\mathbb{R}^3_+)} \le \varepsilon \|w_{nkj}\|_{H^{0,r+2}(\mathbb{R}^3_+)} + C_r \|w_{nkj}\|_{H^{0,1}(\mathbb{R}^3_+)},$$
(5.79)

where C_r is another constant depending on r, increasing in r, which incorporates the factor $||I - \mathbb{Q}||_{W^{0,b,\infty}(\mathbb{R}^3_+)}$.

Second term. By (5.76), (5.59) and (5.19), the second term in the r.h.s. of (5.75) is

$$a_{kj}\mathbf{p}^{-1}\langle \nabla \mathbf{p}, \mathbf{Q}\nabla v_{nj}\rangle = \sum_{\ell \in \mathcal{A}_j} a_{kj}\mathbf{p}^{-1}\langle \nabla \mathbf{p}, \mathbf{Q}\nabla \{(u_n\zeta_{k+1}\psi_\ell) \circ \mathbf{g}_j\}\rangle,$$
(5.80)

where $\mathcal{A}_j := \{\ell \in \{1, \ldots, N\} : K_\ell \cap K_j \neq \emptyset\}$. Identity (5.80) holds because, by (5.59), $\zeta_{k+1} \circ \mathbf{g}_j = 1$ in the support of $\zeta_k \circ \mathbf{g}_j$, by (5.19), and because, if the balls K_ℓ, K_j are disjoint, then $\psi_\ell \circ \mathbf{g}_j = 0$ in the support of $\psi_j \circ \mathbf{g}_j$. Moreover, for every $\ell \in \mathcal{A}_j$, one has

$$(u_n\zeta_{k+1}\psi_\ell)\circ\mathbf{g}_j = (u_n\zeta_{k+1}\psi_\ell)\circ\mathbf{g}_\ell\circ\mathbf{f}_\ell\circ\mathbf{g}_j = w_{n,k+1,\ell}\circ(\mathbf{f}_\ell\circ\mathbf{g}_j),\tag{5.81}$$

where $w_{n,k+1,\ell}$ is defined as in (5.76), and the transition map $\mathbf{f}_{\ell} \circ \mathbf{g}_j$ in A_0 is given by formula (5.23). We consider the extension of the transition map to \mathbb{R}^3_+ given in (5.31), (5.32). By Lemma 5.4, the function $\tilde{\mathbf{T}}_{\ell j}$ is a diffeomorphism of \mathbb{R}^2 , and it is the sum of an affine map and a smooth function with compact support. Hence, by the composition estimate (5.51), we get

$$\|w_{n,k+1,\ell} \circ (\mathbf{f}_{\ell} \circ \mathbf{g}_j)\|_{H^{1,r}(\mathbb{R}^3_{+})} \le C_r \|w_{n,k+1,\ell}\|_{H^{1,r}(\mathbb{R}^3_{+})},$$
(5.82)

where the constant C_r is increasing in r. Concerning a_{kj} , one has estimate (5.64). Also, by (5.57), $\zeta_k(\mathbf{g}_j(y))$ depends only on $|\mathbf{g}_j(y)|$; moreover $\mathbf{g}_j(y) = \mathbf{R}_j^{-1}\mathbf{g}(y)$, and $|\mathbf{g}_j(y)| = |\mathbf{g}(y)| = 1-y_3$ by (5.21). Hence $\zeta_k \circ \mathbf{g}_j$ depends only on y_3 , and not on y_1, y_2 , so that $\partial_y^{\alpha}\{(\zeta_k\psi_j) \circ \mathbf{g}_j\} = (\zeta_k \circ \mathbf{g}_j)\partial_y^{\alpha}(\psi_j \circ \mathbf{g}_j)$ for all multi-indices $\alpha = (\alpha_1, \alpha_2, 0)$. As a consequence, by (5.76) and (5.48) one has

$$\|a_{kj}\|_{W^{0,b,\infty}(\mathbb{R}^3_+)} = \|(\zeta_k\psi_j) \circ \mathsf{g}_j\|_{W^{0,b,\infty}(\mathbb{R}^3_+)} \le \|\psi_j \circ \mathsf{g}_j\|_{W^{0,b,\infty}(\mathbb{R}^3_+)} \le C_r$$
(5.83)

for some constant C_r dependent on r, increasing in r (in fact, C_r depends on b). By (5.47), (5.81), (5.82), (5.64), (5.83), each term of the sum in the r.h.s. of (5.80) satisfies

$$\|a_{kj}\mathbf{p}^{-1}\langle\nabla\mathbf{p},\mathbf{Q}\nabla\{(u_n\zeta_{k+1}\psi_\ell)\circ\mathbf{g}_j\}\rangle\|_{H^{0,r}(\mathbb{R}^3_+)} \le C_r\|w_{n,k+1,\ell}\|_{H^{1,r}(\mathbb{R}^3_+)}.$$
(5.84)

The sum of (5.84) over $\ell \in \mathcal{A}_i$ gives an estimate of the second term in the r.h.s. of (5.75).

Third term. We write the third term in the r.h.s. of (5.75) as

$$a_{kj}\mathbf{p}^{-1}\operatorname{div}\{\mathbf{p}(D\mathbf{g}_{j})^{-1}(g_{n}\circ\mathbf{g}_{j})\} = \mathbf{p}^{-1}\operatorname{div}\{\mathbf{p}(D\mathbf{g}_{j})^{-1}\gamma_{nkj}\} - \langle (D\mathbf{g}_{j})^{-1}(g_{n}\circ\mathbf{g}_{j}), \nabla a_{kj}\rangle, \quad (5.85)$$

where

$$\gamma_{nkj} := a_{kj}(g_n \circ \mathbf{g}_j) = (g_n \zeta_k \psi_j) \circ \mathbf{g}_j.$$
(5.86)

By (5.47), (5.35), (5.36), (5.49), the first term in the r.h.s. of (5.85) satisfies

$$\|\mathbf{p}^{-1}\operatorname{div}\{\mathbf{p}(D\mathbf{g}_{j})^{-1}\gamma_{nkj}\}\|_{H^{0,r}(\mathbb{R}^{3}_{+})} \leq C_{0}\|\gamma_{nkj}\|_{H^{1,r}(\mathbb{R}^{3}_{+})} + C_{r}\|\gamma_{nkj}\|_{H^{1,0}(\mathbb{R}^{3}_{+})}.$$
(5.87)

Using (5.76), (5.59) and (5.19) as in (5.80), and using the identity $g_j = g_\ell \circ f_\ell \circ g_j$ as in (5.81), we write the last term of (5.85) as

$$\langle (D\mathbf{g}_j)^{-1}(g_n \circ \mathbf{g}_j), \nabla a_{kj} \rangle = \sum_{\ell \in \mathcal{A}_j} \langle (D\mathbf{g}_j)^{-1}((g_n \zeta_{k+1} \psi_\ell) \circ \mathbf{g}_j), \nabla a_{kj} \rangle$$
$$= \sum_{\ell \in \mathcal{A}_j} \langle (D\mathbf{g}_j)^{-1}(\gamma_{n,k+1,\ell} \circ (\mathbf{f}_\ell \circ \mathbf{g}_j)), \nabla a_{kj} \rangle, \tag{5.88}$$

where $\gamma_{n,k+1,\ell}$ is defined as in (5.86). Hence, using (5.47), (5.64), (5.83), and Lemma 5.4 and (5.51) for the composition with the transition map $\mathbf{f}_{\ell} \circ \mathbf{g}_j$, one proves that each term of the sum in (5.88) satisfies

$$\|\langle (D\mathbf{g}_j)^{-1}(\gamma_{n,k+1,\ell} \circ (\mathbf{f}_\ell \circ \mathbf{g}_j)), \nabla a_{kj} \rangle\|_{H^{0,r}(\mathbb{R}^3_+)} \le C_r 2^k \|\gamma_{n,k+1,\ell}\|_{H^{0,r}(\mathbb{R}^3_+)}.$$
 (5.89)

The constant C_r in (5.89) depends on r because we have estimated a composition; the factor 2^k is due to the presence of the first derivative $\partial_{y_3}(\zeta_{k+1} \circ \mathbf{g}_j)$. From (5.87) and (5.89) we obtain an estimate for (5.85).

Fourth and fifth term. Proceeding as in (5.80)-(5.84), one proves that the fourth and fifth term in (5.75) are the sum over $\ell \in \mathcal{A}_j$ of terms satisfying the same bound (5.84), but with $C_r 2^{2k}$ instead of C_r . The factor 2^{2k} is due to the presence of the second derivative of ζ_{k+1} with respect to y_3 .

Estimate of u. By (5.75), (5.79), (5.84), (5.87), (5.89), and (5.84) with factor 2^{2k} , one has

$$\begin{split} \|\Delta w_{nkj}\|_{H^{0,r}(\mathbb{R}^{3}_{+})} &\leq \varepsilon \|w_{nkj}\|_{H^{0,r+2}(\mathbb{R}^{3}_{+})} + C_{r}\|w_{nkj}\|_{H^{1,0}(\mathbb{R}^{3}_{+})} + C_{r}2^{2k}\sum_{\ell\in\mathcal{A}_{j}}\|w_{n,k+1,\ell}\|_{H^{1,r}(\mathbb{R}^{3}_{+})} \\ &+ C_{0}\|\gamma_{nkj}\|_{H^{1,r}(\mathbb{R}^{3}_{+})} + C_{r}\|\gamma_{nkj}\|_{H^{1,0}(\mathbb{R}^{3}_{+})} + C_{r}2^{k}\sum_{\ell\in\mathcal{A}_{j}}\|\gamma_{n,k+1,\ell}\|_{H^{0,r}(\mathbb{R}^{3}_{+})}. \end{split}$$
(5.90)

By (5.78), (5.90) and $(5.90)|_{r=0}$, we get

$$\|w_{nkj}\|_{H^{2,r}(\mathbb{R}^{3}_{+})} \leq \varepsilon C_{0} \|w_{nkj}\|_{H^{0,r+2}(\mathbb{R}^{3}_{+})} + \varepsilon C_{r} \|w_{nkj}\|_{H^{0,2}(\mathbb{R}^{3}_{+})} + C_{r} \|w_{nkj}\|_{H^{1,0}(\mathbb{R}^{3}_{+})}$$

$$+ C_{r} 2^{2k} \sum_{\ell \in \mathcal{A}_{j}} \|w_{n,k+1,\ell}\|_{H^{1,r}(\mathbb{R}^{3}_{+})} + C_{0} \|\gamma_{nkj}\|_{H^{1,r}(\mathbb{R}^{3}_{+})} + C_{r} \|\gamma_{nkj}\|_{H^{1,0}(\mathbb{R}^{3}_{+})}$$

$$+ C_{r} 2^{k} \sum_{\ell \in \mathcal{A}_{j}} \|\gamma_{n,k+1,\ell}\|_{H^{0,r}(\mathbb{R}^{3}_{+})}$$

$$(5.91)$$

for some C_0, C_r (possibly different than above). Recalling definitions (5.76), (5.86) of w_{nkj}, γ_{nkj} , taking the sum of (5.91) over $j = 1, \ldots, N$, and using notation (5.61), we obtain

$$\begin{aligned} \|u_n\|_{X_k^{2,r}} &\leq \varepsilon a_0 \|u_n\|_{X_k^{0,r+2}} + \varepsilon b_r \|u_n\|_{X_k^{0,2}} + C_r \|u_n\|_{X_k^{1,0}} + C_r 2^{2k} \|u_n\|_{X_{k+1}^{1,r}} \\ &+ C_0 \|g_n\|_{X_k^{1,r}} + C_r \|g_n\|_{X_k^{1,0}} + C_r 2^k \|g_n\|_{X_{k+1}^{0,r}}, \end{aligned}$$
(5.92)

where we have denoted by a_0 and b_r the constants appearing with the factor ε ; also, b_r, C_r are increasing functions of r. For r = 0, (5.92) becomes

$$\|u_n\|_{X_k^{2,0}} \le \varepsilon(a_0 + b_0) \|u_n\|_{X_k^{0,2}} + C_0 \|u_n\|_{X_k^{1,0}} + C_0 2^{2k} \|u_n\|_{X_{k+1}^{1,0}} + 2C_0 \|g_n\|_{X_k^{1,0}} + C_0 2^k \|g_n\|_{X_{k+1}^{0,0}}.$$
(5.93)

Since $||u_n||_{X_k^{0,2}} \leq ||u_n||_{X_k^{2,0}}$, we fix ε (and therefore the half-radius δ of the balls K_j and their cardinality N) such that $\varepsilon(a_0 + b_0) \leq 1/2$, so that the first term in the r.h.s. of (5.93) can be absorbed by the l.h.s. of (5.93). The second, the third and the fifth term in the r.h.s. of (5.93) are estimated by (5.62), and we get (5.65). Note that ε, δ, N do not depend on r, j, k.

Since $a_0 \varepsilon \leq 1/2$ and $\|\cdot\|_{H^{0,r+2}(\mathbb{R}^3_+)} \leq \|\cdot\|_{H^{2,r}(\mathbb{R}^3_+)}$, the first term in the r.h.s. of (5.92) can be absorbed by the l.h.s. of (5.92). Also, we use (5.65) to estimate the second term in the r.h.s. of (5.92), and we use (5.62) to estimate the third one. Thus,

$$\|u_n\|_{X_k^{2,r}} \le C_r 2^{3k} \mu z^n + C_r 2^{2k} \|u_n\|_{X_{k+1}^{1,r}} + C_r \|g_n\|_{X_k^{1,0}} + C_0 \|g_n\|_{X_k^{1,r}} + C_r 2^k \|g_n\|_{X_{k+1}^{0,r}}.$$
 (5.94)

For r = 0, (5.94), (5.62) give (5.65). For $0 < r \le 1$, we consider (5.94), we use the fact that

$$\|u_n\|_{X_{k+1}^{1,r}} \le \|u_n\|_{X_{k+1}^{1,1}} \le \|u_n\|_{X_{k+1}^{2,0}}, \quad \|g_n\|_{X_{k+1}^{0,r}} \le \|g_n\|_{X_{k+1}^{0,1}} \le \|g_n\|_{X_{k+1}^{1,0}}$$

and we apply (5.65) (with k+1 instead of k) to bound $||u||_{X^{2,0}_{k+1}}$; in this way we obtain

$$\|u_n\|_{X_k^{2,r}} \le C_{r,k}\mu z^n + C_{r,k}\|g_n\|_{X_{k+1}^{1,0}} + C_0\|g_n\|_{X_k^{1,r}} + C_r\|g_n\|_{X_k^{1,0}}, \quad 0 < r \le 1.$$
(5.95)

Now we prove (5.66) by induction on b. For b = 1, (5.66) is (5.95). Assume that (5.66) is proved for some integer $b \in \mathbb{N}$, and let $b < r \le b+1$. Apply (5.94), use the inequality $||u||_{X_{k+1}^{1,r}} \le ||u||_{X_{k+1}^{2,r-1}}$ and use the induction assumption (5.66) (with r - 1, k + 1 instead of r, k) to estimate $||u||_{X_{k+1}^{2,r-1}}$. This gives (5.66) with b + 1 instead of b (and constants $C_{r,k}$ possibly different than those of the induction assumption). Note that the constant C_0 in (5.66) is not changed by the induction step. Hence (5.66) holds for all integers $b \ge 1$.

In the next lemma we estimate the norm (5.61) of u_0 , which was not included in Lemma 5.11.

Lemma 5.12. For $k \in \mathbb{N}_0$, u_0 in (5.3) satisfies

$$\|u_0\|_{X_k^{2,0}} \le C_0 \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)} + C_0 2^{3k} \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} \le C_k \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)}.$$
(5.96)

For $0 \le b - 1 < r \le b$, with b integer and r real, one has

$$\|u_0\|_{X_k^{2,r}} \le C_0 \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^2)} + C_{r,k} \Big(\sum_{\ell=1}^{b-1} \|\psi\|_{H^{r+\frac{3}{2}-\ell}(\mathbb{S}^2)} + \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)} \Big)$$
(5.97)

$$\leq C_0' \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^2)} + C_{r,k}' \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)},\tag{5.98}$$

where $C'_0 = 1 + C_0$, C_0 is independent of r, k, while $C_{r,k}, C'_{r,k}$ depend on r, k, and they are increasing functions of r.

Proof. We closely follow the proof of Lemma 5.11. Since u_0 solves (5.3), the function w_{0kj} (defined as in (5.76)) satisfies (5.75) with $g_0 = 0$, that is, without the third term in the r.h.s., and

$$w_{0kj} \in H^1(\mathbb{R}^3_+), \quad w_{0kj} = (\psi\psi_j) \circ \mathbf{g}_j \quad \text{on } \mathbb{R}^2 \times \{0\}$$
(5.99)

instead of (5.77) (note that $\zeta_k \circ \mathbf{g}_j(\cdot, 0) = 1$ for all $k \in \mathbb{N}_0$). Applying Lemma 5.5 gives

$$|w_{0kj}||_{H^{2,r}(\mathbb{R}^{3}_{+})} \leq C \|\Delta w_{0kj}\|_{H^{0,r}(\mathbb{R}^{3}_{+})} + C^{r+1} \|\Delta w_{0kj}\|_{L^{2}(\mathbb{R}^{3}_{+})} + C \|w_{0kj}(\cdot,0)\|_{H^{r+\frac{3}{2}}(\mathbb{R}^{2})} + C^{r+1} \|w_{0kj}\|_{H^{1}(\mathbb{R}^{3}_{+})},$$

$$(5.100)$$

where C > 0 is the universal constant in (5.33), Δw_{0kj} is given by (5.75) (with $g_0 = 0$) and $w_{0kj}(\cdot, 0)$ is given by (5.99). Now Δw_{0kj} satisfies (5.90) without the terms with $\gamma_{nkj}, \gamma_{n,k+1,\ell}$, which are zero for n = 0. Thus, by (5.100), (5.90) and (5.90)|_{r=0}, we get

$$\begin{aligned} \|w_{0kj}\|_{H^{2,r}(\mathbb{R}^{3}_{+})} &\leq \varepsilon C_{0} \|w_{0kj}\|_{H^{0,r+2}(\mathbb{R}^{3}_{+})} + \varepsilon C_{r} \|w_{0kj}\|_{H^{0,2}(\mathbb{R}^{3}_{+})} + C_{r} \|w_{0kj}\|_{H^{1,0}(\mathbb{R}^{3}_{+})} \\ &+ C_{r} 2^{2k} \sum_{\ell \in \mathcal{A}_{j}} \|w_{0,k+1,\ell}\|_{H^{1,r}(\mathbb{R}^{3}_{+})} + C \|(\psi\psi_{j}) \circ \mathbf{g}_{j}\|_{H^{r+\frac{3}{2}}(\mathbb{R}^{2})} + C^{r+1} \|(\psi\psi_{j}) \circ \mathbf{g}_{j}\|_{H^{\frac{3}{2}}(\mathbb{R}^{2})} \tag{5.101}$$

instead of (5.91). Hence u_0 satisfies (5.92) with $g_0 = 0$ and with the additional terms $C_0 \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^2)}$ and $C_r \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)}$ in the r.h.s., and we obtain

$$\|u_0\|_{X_k^{2,r}} \le C_0 \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^2)} + C_{r,k} \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} + C_r \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)} + C_{r,k} \|u_0\|_{X_{k+1}^{1,r}}$$
(5.102)

instead of (5.94). Then (5.96), (5.97) are proved in the same way as (5.65), (5.66).

Inequality (5.98) is deduced from (5.97) by the interpolation inequality (5.52), using the fact that

$$|\psi|_{r} \le |\psi|_{r_{0}}^{1-\vartheta}|\psi|_{r_{1}}^{\vartheta} = \frac{1}{\lambda} (\lambda^{p}|\psi|_{r_{0}})^{1-\vartheta}|\psi|_{r_{1}}^{\vartheta} \le \frac{1}{\lambda} (\lambda^{p}|\psi|_{r_{0}} + |\psi|_{r_{1}}) = \lambda^{p-1}|\psi|_{r_{0}} + \frac{|\psi|_{r_{1}}}{\lambda}$$
(5.103)

for any $\lambda > 0$, with $r = (1 - \vartheta)r_0 + \vartheta r_1$, $0 < \vartheta < 1$, $p = 1/(1 - \vartheta)$, $|\psi|_r = ||\psi||_{H^r(\mathbb{S}^2)}$, and choosing λ sufficiently large, depending on r, k.

Lemma 5.13. The matrices P_1, P_2, \ldots in (5.1) and the functions u_0, u_1, \ldots in (5.3), (5.4) satisfy

$$\begin{aligned} \|P_m \nabla u_\sigma\|_{X_k^{1,r}} &\leq m \big(C_0 \|h\|_{H^{r_0+2}(\mathbb{S}^2)} \big)^m \big(\|u_\sigma\|_{X_k^{2,r}} + C_{r,k} \|u_\sigma\|_{X_{k+1}^{1,r}} \big) \\ &+ (\chi_{r>r_0}) m \big(C_0 \|h\|_{H^{r_0+2}(\mathbb{S}^2)} \big)^{m-1} \|h\|_{H^{r+2}(\mathbb{S}^2)} \big(C_r \|u_\sigma\|_{X_k^{2,r_0}} + C_{r,k} \|u_\sigma\|_{X_{k+1}^{1,r_0}} \big) \end{aligned}$$
(5.104)

for all integers $m \in \mathbb{N}$, $\sigma, k \in \mathbb{N}_0$, for all real $r \ge 0$, $r_0 > 1/2$, where $(\chi_{r>r_0}) = 1$ for $r > r_0$ and $(\chi_{r>r_0}) = 0$ for $r \le r_0$. The constant $C_{r,k}$ depends on r, k and not on m, σ ; C_r depends on r and not on k, m, σ ; $C_{r,k}, C_r$ are increasing in r; C_0 is independent of r, k, m, σ ; $C_r, C_{r,k}, C_0$ depend on r_0 (which we consider as fixed).

Proof. By (5.1), $P_m(x)$ is homogeneous of order 0 in x. Hence, recalling (5.21), (5.22), and denoting $y' = (y_1, y_2)$, one has

$$g_j(y', y_3) = (1 - y_3)g_j(y', 0), \quad P_m \circ g_j(y', y_3) = P_m \circ g_j(y', 0)$$

for all $y = (y', y_3) \in A_0$. Therefore, in \mathbb{R}^3_+ , the product $P_m \psi_j$ satisfies

$$(P_m\psi_j)\circ \mathbf{g}_j(y)=P_m\circ \mathbf{g}_j(y',0)\,\psi_j\circ \mathbf{g}_j(y)\quad \forall y\in \mathbb{R}^3_+,$$

because $\psi_j \circ \mathbf{g}_j = 0$ outside A_0 . By (5.19), $1 = \sum_{\ell=1}^N \psi_\ell(x)$ at any point $x = \mathbf{g}_j(y', 0)$. Also, $\zeta_k \circ \mathbf{g}_j(y) = (\zeta^* \zeta_k) \circ \mathbf{g}_j(y)$ for all $y \in \mathbb{R}^3_+$, because ζ^* in (5.60) is equal to 1 on the support of ζ_k , for all $k \in \mathbb{N}_0$. Hence

$$(P_m\zeta_k\psi_j)\circ\mathbf{g}_j(y) = \sum_{\ell\in\mathcal{A}_j} (P_m\psi_\ell)\circ\mathbf{g}_j(y',0)\,\zeta^*\circ\mathbf{g}_j(y)a_{kj}(y)$$
$$= \sum_{\ell\in\mathcal{A}_j} (P_m\psi_\ell)\circ\mathbf{g}_\ell(\tilde{\mathsf{T}}_{\ell j}(y'),0)\,\zeta^*\circ\mathbf{g}_\ell(\tilde{\mathsf{T}}_{\ell j}(y'),y_3)\,a_{kj}(y) \tag{5.105}$$

in \mathbb{R}^3_+ , where a_{kj} is defined in (5.76). We have used the identity $\mathbf{g}_j = \mathbf{g}_\ell \circ \mathbf{f}_\ell \circ \mathbf{g}_j$ in A_0 , where the transition map $\mathbf{f}_\ell \circ \mathbf{g}_j$ satisfies (5.23) in A_0 , and $\tilde{\mathsf{T}}_{\ell j}(y')$ is defined for all $y' \in \mathbb{R}^2$ by (5.31) and it satisfies (5.32). Again, identity (5.105) holds in \mathbb{R}^3_+ (and not only in $A_0 \cap \mathbb{R}^3_+$) because $\psi_j \circ \mathbf{g}_j = 0$ outside A_0 . Moreover,

$$\zeta^* \circ \mathsf{g}_{\ell}(\tilde{\mathsf{T}}_{\ell j}(y'), y_3) = \zeta^* \circ \mathsf{g}_{\ell}(0, y_3)$$

because ζ^* is a radial function and, therefore, $\zeta^* \circ \mathbf{g}_{\ell}$ is a function of y_3 only, independent of y'. By (5.67), using the definitions in (5.76), we obtain

$$(P_m\zeta_k\psi_j\nabla u_{\sigma})\circ \mathbf{g}_j(y) = \sum_{\ell\in\mathcal{A}_j} (P_m\psi_\ell)\circ \mathbf{g}_\ell(\tilde{\mathbf{T}}_{\ell j}(y'),0)\,\zeta^*\circ \mathbf{g}_\ell(0,y_3)\,a_{kj}(y)[D\mathbf{g}_j(y)]^{-T}\nabla v_{\sigma j}(y)$$
$$= \sum_{\ell\in\mathcal{A}_j} (P_m\psi_\ell)\circ \mathbf{g}_\ell(\tilde{\mathbf{T}}_{\ell j}(y'),0)\,\zeta^*\circ \mathbf{g}_\ell(0,y_3)[D\mathbf{g}_j(y)]^{-T}\big(\nabla w_{\sigma kj}(y)-v_{\sigma j}(y)\nabla a_{kj}(y)\big) \quad (5.106)$$

in \mathbb{R}^3_+ . Moreover, using (5.19) and (5.59), and recalling the notations in (5.76), $v_{\sigma j} \nabla a_{kj}$ in (5.106) can be replaced by $\sum_{i \in \mathcal{A}_j} w_{\sigma,k+1,i} \nabla a_{kj}$. By (5.38), (5.37), (5.51), the function $y \mapsto (P_m \psi_\ell) \circ g_\ell(\tilde{T}_{\ell j}(y'), 0) \zeta^* \circ g_\ell(0, y_3)$ (which is the product of a function of y' with a function of y_3) satisfies

$$\| y \mapsto (P_m \psi_{\ell}) \circ g_{\ell}(\tilde{\mathsf{T}}_{\ell j}(y'), 0) \, \zeta^* \circ g_{\ell}(0, y_3) \|_{H^{1,r}(\mathbb{R}^3_+)} \leq C_0 \| (P_m \psi_{\ell}) \circ g_{\ell}(\tilde{\mathsf{T}}_{\ell j}(\cdot), 0) \|_{H^{r+1}(\mathbb{R}^2)}$$

$$\leq C_r \| (P_m \psi_{\ell}) \circ g_{\ell}(\cdot, 0) \|_{H^{r+1}(\mathbb{R}^2)}.$$
(5.107)

Hence, by (5.106), (5.107), (5.46), (5.35), (5.36), recalling definitions (5.61), (5.18) of norms $\| \|_{X_k^{s,r}}$ and $\| \|_{H^r(\mathbb{S}^2)}$, and summing over j, ℓ, i , we obtain

$$\begin{aligned} \|P_m \nabla u_\sigma\|_{X_k^{1,r}} &\leq C_{r_0} \|P_m\|_{H^{r_0+1}(\mathbb{S}^2)} \|u_\sigma\|_{X_k^{2,r}} + (\chi_{r>r_0}) C_r \|P_m\|_{H^{r+1}(\mathbb{S}^2)} \|u_\sigma\|_{X_k^{2,r_0}} \\ &+ C_{r,k} \|P_m\|_{H^{r_0+1}(\mathbb{S}^2)} \|u_\sigma\|_{X_{k+1}^{1,r}} + (\chi_{r>r_0}) C_{r,k} \|P_m\|_{H^{r+1}(\mathbb{S}^2)} \|u_\sigma\|_{X_{k+1}^{1,r_0}}, \end{aligned}$$
(5.108)

where the first and second term in the r.h.s. of (5.108) comes from estimating the terms with $\nabla w_{\sigma kj}$ in (5.106), while the third and fourth term in the r.h.s. of (5.108) comes from estimating the terms with $v_{\sigma j} \nabla a_{kj}$ in (5.106), namely with $w_{\sigma,k+1,i} \nabla a_{kj}$. The real number r_0 in (5.108) comes from the application of (5.46), and it has to satisfy $r_0 + 1 > 3/2$. As a consequence, to proceed with estimating the r.h.s. of (5.108), we have to estimate $\|P_m\|_{H^{r+1}(\mathbb{S}^2)}$ for $r \ge r_0$, namely $\|P_m\|_{H^s(\mathbb{S}^2)}$ for $s \ge r_0 + 1$. By (5.1), (5.52), (5.53), (5.54), (5.55), P_m on \mathbb{S}^2 satisfies

$$\|P_1\|_{H^s(\mathbb{S}^2)} \le C_{s_0} \|h\|_{H^{s+1}(\mathbb{S}^2)} + C_s \|h\|_{H^{s_0+1}(\mathbb{S}^2)}, \|P_m\|_{H^s(\mathbb{S}^2)} \le C_s(m-1)(C_{s_0} \|h\|_{H^{s_0}(\mathbb{S}^2)})^{m-2} \|h\|_{H^{s_0+1}(\mathbb{S}^2)} \|h\|_{H^{s+1}(\mathbb{S}^2)} \quad \forall m = 2, 3, \dots,$$
 (5.109)

for all real $s \ge s_0$, where s_0 is given by (5.53) and it has to satisfy $s_0 > 1$. For m = 1 and m = 2, (5.109) follows easily from (5.1), (5.53), (5.55); for $m \ge 3$, observe that $P_m = (-h)^{m-2}P_2$ and use the estimate for P_2 already proved, together with (5.53), (5.54), (5.52). Estimate (5.109) implies the less accurate inequality

$$\|P_m\|_{H^s(\mathbb{S}^2)} \le C_s m(C_0 \|h\|_{H^{s_0+1}(\mathbb{S}^2)})^{m-1} \|h\|_{H^{s+1}(\mathbb{S}^2)} \quad \forall m = 1, 2, \dots,$$
(5.110)

for all real $s \ge s_0$, where C_0 does not depend on s, m, h. Taking $s_0 \le r_0 + 1$, inequality (5.104) follows from (5.108) and (5.110).

In (5.104) we have used only Sobolev norms $\|h\|_{H^r(\mathbb{S}^2)}$ of h, while in (5.6), (5.7), (5.62), (5.63), (5.65), (5.66) one finds $\|h\|_{L^{\infty}(\mathbb{S}^2)}, \|\nabla_{\mathbb{S}^2}h\|_{L^{\infty}(\mathbb{S}^2)}$ in τ, z defined in (5.6), (5.63). We note that, by (5.55), (5.56), one has

$$z \le C_0 \|h\|_{H^{s_0+1}(\mathbb{S}^2)} \le C_0 \|h\|_{H^{r_0+2}(\mathbb{S}^2)},\tag{5.111}$$

where $1 < s_0 \leq r_0 + 1$, for some constant C_0 depending on s_0 (which we consider as fixed).

Lemma 5.14. The functions u_n in (5.4) satisfy

$$\begin{aligned} \|u_n\|_{X_k^{2,r}} &\leq (Bc_0 z_0)^n \left(\|\psi\|_{H^{\frac{3}{2}+r}(\mathbb{S}^2)} + A_{r,k} \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)} \right) \\ &+ (\chi_{r>r_0}) A_{r,k} B^n (c_0 z_0)^{n-1} \|h\|_{H^{2+r}(\mathbb{S}^2)} \|\psi\|_{H^{r_0+\frac{3}{2}}(\mathbb{S}^2)}, \quad z_0 := \|h\|_{H^{r_0+2}(\mathbb{S}^2)}, \quad (5.112) \end{aligned}$$

for all $n \in \mathbb{N}$, $k \in \mathbb{N}_0$, all real $r \ge 0$, where $r_0 \in (\frac{1}{2}, 1]$. The constants B, c_0 are independent of r, n, k; $A_{r,k}$ depends on r, k, it is increasing in r, and it is independent of n; c_0 is the maximum between the constant C_0 in (5.104) and the constant C_0 in (5.111); $B, c_0, A_{r,k}$ depend on r_0 .

Proof. We start with the case r = 0. For n = 1, by (5.65), (5.63), (5.111), (5.5), one has

$$\|u_1\|_{X_k^{2,0}} \le C_k z_0 \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} + C_0 \|P_1 \nabla u_0\|_{X_k^{1,0}},$$

and, by (5.104), (5.96), (5.62),

$$\|P_1 \nabla u_0\|_{X_k^{1,0}} \le C_k c_0 z_0 \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)}.$$
(5.113)

Hence (5.112) holds for r = 0, n = 1 if $B \ge 1$ and $A_{0,k} \ge C_k$, for some C_k depending on k. Now assume that (5.112) with r = 0 holds for u_1, \ldots, u_{n-1} , for some $n \ge 2$. By (5.65), (5.63), (5.111), (5.5), one has

$$\|u_n\|_{X_k^{2,0}} \le C_k (c_0 z_0)^n \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} + C_0 \sum_{\sigma=0}^{n-1} \|P_{n-\sigma} \nabla u_\sigma\|_{X_k^{1,0}},$$

and, by (5.104),

$$\sum_{\sigma=0}^{n-1} \|P_{n-\sigma} \nabla u_{\sigma}\|_{X_{k}^{1,0}} \leq \sum_{\sigma=0}^{n-1} (n-\sigma) (c_{0}z_{0})^{n-\sigma} (\|u_{\sigma}\|_{X_{k}^{2,0}} + C_{k}\|u_{\sigma}\|_{X_{k+1}^{1,0}})$$

Using (5.96), the induction assumption (5.112) for u_1, \ldots, u_{n-1} , and (5.62), we obtain that

$$\|u_n\|_{X_k^{2,0}} \le (c_0 z_0)^n \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)} \Big(C_k n + C_k \sum_{\sigma=1}^{n-1} (n-\sigma) + C_0 (1+A_{0,k}) \sum_{\sigma=1}^{n-1} (n-\sigma) B^{\sigma} \Big).$$

Now $n + \sum_{\sigma=1}^{n-1} (n-\sigma) = \frac{1}{2}n(n+1) \le n^2$, and

$$\sum_{\sigma=1}^{n-1} (n-\sigma) B^{\sigma} = B^n \sum_{\sigma=1}^{n-1} \frac{n-\sigma}{B^{n-\sigma}} = B^n \sum_{m=1}^{n-1} \frac{m}{B^m} < B^n \sum_{m=1}^{\infty} \frac{m}{B^m} = \frac{B^{n-1}}{(1-B^{-1})^2} \le 4B^{n-1}$$
(5.114)

for $B \ge 2$. Hence (5.112) with r = 0 holds for u_n if $C_k n^2 + C_0 (1 + A_{0,k}) 4B^{n-1} \le (1 + A_{0,k})B^n$. Now $C_k n^2 \le \frac{1}{2}(1 + A_{0,k})B^n$ if $A_{0,k} \ge 2C_k$ and $B \ge 3$ (so that $n^2 \le B^n$ for any $n \in \mathbb{N}$), and $C_0(1 + A_{0,k})4B^{n-1} \le \frac{1}{2}(1 + A_{0,k})B^n$ if $B \ge 8C_0$. This proves that, for $A_{0,k}$ larger than some constant C_k (depending on k), and B larger than some constant C_0 (independent of k), (5.112) with r = 0 holds for all $n \in \mathbb{N}$.

Consider the case $0 < r \le 1$. For n = 1, by (5.66), (5.63), (5.111), (5.5), one has

$$\|u_1\|_{X_k^{2,r}} \le C_{r,k} c_0 z_0 \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} + C_0 \|P_1 \nabla u_0\|_{X_k^{1,r}} + C_r \|P_1 \nabla u_0\|_{X_k^{1,0}} + C_{r,k} \|P_1 \nabla u_0\|_{X_{k+1}^{1,0}}$$

By (5.104), (5.98), (5.96), (5.62), one has

$$\|P_1 \nabla u_0\|_{X_k^{1,r}} \le c_0 z_0 (C_0 \|\psi\|_{H^{\frac{3}{2}+r}(\mathbb{S}^2)} + C_{r,k} \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)}) + (\chi_{r>r_0}) C_{r,k} \|h\|_{H^{r+2}(\mathbb{S}^2)} \|\psi\|_{H^{r_0+\frac{3}{2}}(\mathbb{S}^2)},$$

where we have used the fact that $||u_0||_{X_{k+1}^{1,r}} \leq ||u_0||_{X_{k+1}^{2,r-1}} \leq ||u_0||_{X_{k+1}^{2,0}}$. The term $||P_1 \nabla u_0||_{X_k^{1,0}}$ has been estimated in (5.113). Hence

$$\|u_1\|_{X_k^{2,r}} \le c_0 z_0(C_0 \|\psi\|_{H^{\frac{3}{2}+r}(\mathbb{S}^2)} + C_{r,k} \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)}) + (\chi_{r>r_0})C_{r,k} \|h\|_{H^{r+2}(\mathbb{S}^2)} \|\psi\|_{H^{r_0+\frac{3}{2}}(\mathbb{S}^2)},$$

and (5.112) with $r \in (0,1]$ holds for u_1 if $B \ge C_0$ and $A_{r,k} \ge C_{r,k}$. Assume that (5.112) with $r \in (0,1]$ holds for u_1, \ldots, u_{n-1} , for some $n \ge 2$. By (5.66), (5.63), (5.111), (5.5), one has

$$\begin{aligned} \|u_n\|_{X_k^{2,r}} &\leq C_{r,k} (c_0 z_0)^n \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} \\ &+ \sum_{\sigma=0}^{n-1} \Big(C_0 \|P_{n-\sigma} \nabla u_\sigma\|_{X_k^{1,r}} + C_r \|P_{n-\sigma} \nabla u_\sigma\|_{X_k^{1,0}} + C_{r,k} \|P_{n-\sigma} \nabla u_\sigma\|_{X_{k+1}^{1,0}} \Big), \end{aligned}$$

and, by (5.104), this is

$$\leq C_{r,k}(c_0 z_0)^n \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} + \sum_{\sigma=0}^{n-1} (n-\sigma)(c_0 z_0)^{n-\sigma} \Big(C_0 \|u_\sigma\|_{X_k^{2,r}} + C_r \|u_\sigma\|_{X_k^{2,0}} + C_{r,k} \|u_\sigma\|_{X_{k+1}^{2,0}} + C_{r,k} \|u_\sigma\|_{X_{k+2}^{2,0}} \Big) + (\chi_{r>r_0}) \sum_{\sigma=0}^{n-1} (n-\sigma)(c_0 z_0)^{n-\sigma-1} \|h\|_{H^{r+2}(\mathbb{S}^2)} \Big(C_r \|u_\sigma\|_{X_k^{2,r_0}} + C_{r,k} \|u_\sigma\|_{X_{k+1}^{2,0}} \Big),$$

where we have used the elementary inequalities

$$\|u_{\sigma}\|_{X^{1,0}_{k+1}} \le \|u_{\sigma}\|_{X^{1,r}_{k+1}} \le \|u_{\sigma}\|_{X^{2,r-1}_{k+1}} \le \|u_{\sigma}\|_{X^{2,0}_{k+1}}, \quad \|u_{\sigma}\|_{X^{1,0}_{k+2}} \le \|u_{\sigma}\|_{X^{2,0}_{k+2}}$$

(one has $r-1 \leq 0$ and also $r_0 - 1 \leq 0$). Using bounds (5.96) and (5.98) for u_0 , the induction assumption (5.112) with $r \in (0, 1]$ for u_1, \ldots, u_{n-1} , the already proved estimate (5.112) with r = 0 for u_1, \ldots, u_{n-1} , the inequalities $n \leq B^{n-1}$ and (5.114), we obtain that

$$\begin{aligned} \|u_n\|_{X_k^{2,r}} &\leq C_0 B^{n-1} (c_0 z_0)^n \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^2)} \\ &+ \left\{ C_0 A_{r,k} + C_{r,k} (1 + A_{0,k} + A_{0,k+1} + A_{0,k+2}) \right\} B^{n-1} (c_0 z_0)^n \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)} \\ &+ (\chi_{r>r_0}) \left\{ C_0 A_{r,k} + C_r (1 + A_{r_0,k}) + C_{r,k} (1 + A_{0,k+1}) \right\} B^{n-1} (c_0 z_0)^{n-1} \|h\|_{H^{r+2}(\mathbb{S}^2)} \|\psi\|_{H^{r_0+\frac{3}{2}}(\mathbb{S}^2)}. \end{aligned}$$

One has $C_0B^{n-1} \leq B^n$ if $B \geq C_0$; one has $C_0A_{r,k}B^{n-1} \leq \frac{1}{2}A_{r,k}B^n$ if $B \geq 2C_0$; and one has $C_{r,k}(1 + A_{0,k} + A_{0,k+1} + A_{0,k+2})B^{n-1} \leq \frac{1}{2}A_{r,k}B^n$ if $B \geq 1$ and $A_{r,k} \geq 2C_{r,k}(1 + A_{0,k} + A_{0,k+1} + A_{0,k+2})$. This proves that (5.112) with $r \in (0, r_0]$ holds for all $n \in \mathbb{N}$ if $A_{r,k}$ is larger than some constant depending on r, k (involving the constants $A_{0,k}, A_{0,k+1}, A_{0,k+2}$ already fixed in the r = 0 case) and B is larger than some constant independent of r, k. Similarly, (5.112) with $r \in (r_0, 1]$ holds for all $n \in \mathbb{N}$ if $A_{r,k}$ is larger than some constant depending on r, k (involving the constants).

 $A_{0,k}, A_{0,k+1}, A_{0,k+2}$ already fixed in the r = 0 case and also the constant $A_{r_0,k}$ already fixed in the $r = r_0$ case), and B is larger than some constant independent of r, k.

Now we prove, by induction on b, that (5.112) holds for all real $r \in (b-1, b]$, all $n \in \mathbb{N}$, all $b \in \mathbb{N}$. For b = 1 we have just proved it. Let $b \in \mathbb{N}$, $b \ge 2$, and assume that (5.112) holds for all $r \in (\beta - 1, \beta]$, all $n \in \mathbb{N}$, all $\beta = 1, \ldots, b-1$; we have to prove that (5.112) holds for $r \in (b-1, b]$. Let $b - 1 < r \le b$. For n = 1, by (5.66), (5.63), (5.111), (5.5), one has

$$\|u_1\|_{X_k^{2,r}} \le C_{r,k} c_0 z_0 \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} + C_0 \|P_1 \nabla u_0\|_{X_k^{1,r}} + C_r \|P_1 \nabla u_0\|_{X_k^{1,0}} + C_{r,k} \sum_{\ell=1}^{b-1} \|P_1 \nabla u_0\|_{X_{k+\ell}^{1,r-\ell}} + C_{r,k} \|P_1 \nabla u_0\|_{X_{k+b}^{1,0}}.$$
(5.115)

By (5.104), (5.98), one has

$$\begin{split} \|P_{1}\nabla u_{0}\|_{X_{k}^{1,r}} &\leq c_{0}z_{0}(C_{0}\|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^{2})} + C_{r,k}\|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^{2})}) + (\chi_{r>r_{0}})C_{r,k}\|h\|_{H^{r+2}(\mathbb{S}^{2})}\|\psi\|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})},\\ \|P_{1}\nabla u_{0}\|_{X_{k+\ell}^{1,r-\ell}} &\leq c_{0}z_{0}(C_{0}\|\psi\|_{H^{r-\ell+\frac{3}{2}}(\mathbb{S}^{2})} + C_{r,k,\ell}\|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^{2})})\\ &+ (\chi_{r-\ell>r_{0}})C_{r,k,\ell}\|h\|_{H^{r-\ell+2}(\mathbb{S}^{2})}\|\psi\|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})}, \end{split}$$

where we have used the inequalities

$$\|u_0\|_{X^{1,r}_{k+1}} \le \|u_0\|_{X^{2,r-1}_{k+1}}, \quad C_{r,k}\|\psi\|_{H^{r-1+\frac{3}{2}}(\mathbb{S}^2)} \le \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^2)} + C'_{r,k}\|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)}$$

(see (5.103)), and similarly for the terms with $r - \ell$. The terms $||P_1 \nabla u_0||_{X_k^{1,0}}$ and $||P_1 \nabla u_0||_{X_{k+b}^{1,0}}$ are estimated in (5.113). Hence (5.115) gives

$$\|u_1\|_{X_k^{2,r}} \le c_0 z_0 (C_0 \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^2)} + C_{r,k} \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)}) + (\chi_{r>r_0}) C_{r,k} \|h\|_{H^{r+2}(\mathbb{S}^2)} \|\psi\|_{H^{r_0+\frac{3}{2}}(\mathbb{S}^2)},$$

and (5.112) with $r \in (b-1, b]$ holds for u_1 if $B \ge C_0$ and $A_{r,k} \ge C_{r,k}$. Now assume that (5.112) with $r \in (b-1, b]$ holds for u_1, \ldots, u_{n-1} , for some $n \ge 2$. By (5.66), (5.63), (5.111), (5.5), one has

$$\begin{aligned} \|u_n\|_{X_k^{2,r}} &\leq C_{r,k} (c_0 z_0)^n \|\psi\|_{H^{\frac{1}{2}}(\mathbb{S}^2)} + \sum_{\sigma=0}^{n-1} \left(C_0 \|P_{n-\sigma} \nabla u_\sigma\|_{X_k^{1,r}} + C_r \|P_{n-\sigma} \nabla u_\sigma\|_{X_k^{1,0}} \right) \\ &+ C_{r,k} \sum_{\ell=1}^{b-1} \|P_{n-\sigma} \nabla u_\sigma\|_{X_{k+\ell}^{1,r-\ell}} + C_{r,k} \|P_{n-\sigma} \nabla u_\sigma\|_{X_{k+b}^{1,0}} \right), \end{aligned}$$

and, using (5.104), (5.114), the induction estimate for $||u_{\sigma}||_{X_{k}^{2,r}}$, the estimates for u_{σ} already proved, and the interpolation inequality (5.103), this is

$$\leq C_0 B^{n-1} (c_0 z_0)^n \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^2)} + (C_0 A_{r,k} + C_{r,k,A}) B^{n-1} (c_0 z_0)^n \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^2)} + (\chi_{r>r_0}) (C_0 A_{r,k} + C_{r,k,A}) B^{n-1} (c_0 z_0)^{n-1} \|h\|_{H^{r+2}(\mathbb{S}^2)} \|\psi\|_{H^{r_0+\frac{3}{2}}(\mathbb{S}^2)},$$

where C_0 is independent of r, b, n, k, while $C_{r,k,A}$ is independent of n, and it depends on r, k and on the constants $A_{r_0,k}$, $A_{0,k+b}$, $A_{0,k+b+1}$, $A_{r-\ell,k+\ell}$, $A_{r-\ell,k+\ell+1}$, $\ell = 1, \ldots, b-1$, all of which have been already fixed. Hence (5.112) with $r \in (b-1, b]$ holds for u_n if $C_0 B^{n-1} \leq B^n$, $C_0 A_{r,k} B^{n-1} \leq \frac{1}{2} A_{r,k} B^n$, and $C_{r,k,A} B^{n-1} \leq \frac{1}{2} A_{r,k} B^n$. These conditions are satisfied for $B \geq 2C_0$ and $A_{r,k} \geq 2C_{r,k,A}$. This completes the proof of the induction step.

The assumption $r_0 \leq 1$ in Lemma 5.14 is not really necessary, but only convenient.

In the next lemma we observe that what we have proved for the function u_n can be easily adapted to the *n*-linear map \mathring{u}_n in (5.11).

Lemma 5.15. The n-linear map \mathring{u}_n in (5.11) satisfies

$$\begin{aligned} \| \mathring{u}_{n}[h_{1}, \dots, h_{n}] \|_{X_{k}^{2,r}} &\leq B^{n} c_{0}^{n} \Big(\prod_{j=1}^{n} \| h_{j} \|_{H^{r_{0}+2}(\mathbb{S}^{2})} \Big) \Big(\| \psi \|_{H^{\frac{3}{2}+r}(\mathbb{S}^{2})} + A_{r,k} \| \psi \|_{H^{\frac{3}{2}}(\mathbb{S}^{2})} \Big) \\ &+ (\chi_{r>r_{0}}) A_{r,k} B^{n} c_{0}^{n-1} \frac{1}{n} \Big(\sum_{j=1}^{n} \| h_{j} \|_{H^{r+2}(\mathbb{S}^{2})} \prod_{i \neq j} \| h_{i} \|_{H^{r_{0}+2}(\mathbb{S}^{2})} \Big) \| \psi \|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})} \tag{5.116}$$

for all h_1, \ldots, h_n , all $n \in \mathbb{N}$, $k \in \mathbb{N}_0$, all real $r \ge 0$, where $r_0 \in (\frac{1}{2}, 1]$. The constants $B, c_0, A_{r,k}, c_0$ are those of Lemma 5.14.

Proof. The proof is a simple adaptation of the previous lemmas.

Proposition 5.16. For any $r_0 \in (\frac{1}{2}, 1]$, there exists $\delta_0 > 0$ such that, for $||h||_{H^{r_0+2}(\mathbb{S}^2)} \leq \delta_0$, the series $u = \sum_{n=0}^{\infty} u_n$, with u_n in (5.3), (5.4), satisfies

$$\|u\|_{X_{k}^{2,r}} \leq C_{0} \|\psi\|_{H^{\frac{3}{2}+r}(\mathbb{S}^{2})} + C_{r,k} \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^{2})} + (\chi_{r>r_{0}})C_{r,k} \|h\|_{H^{r+2}(\mathbb{S}^{2})} \|\psi\|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})}$$
(5.117)

for all $k \in \mathbb{N}_0$, all real $r \ge 0$, where the constant $C_{r,k}$ depends on r, k and it is increasing in r, while C_0 and δ_0 are independent of r, k. For any $\psi \in H^{\frac{3}{2}+r}(\mathbb{S}^2)$, the map $h \mapsto u$ is analytic from the ball $\{h \in H^{r+2}(\mathbb{S}^2) \cap H^{r_0+2}(\mathbb{S}^2) : \|h\|_{H^{r_0+2}(\mathbb{S}^2)} < \delta_0\}$ to the Banach space $\{u \in H^1(B_1) : \|u\|_{X_k^{2,r}} < \infty\}$ endowed with the norm $\|u\|_{H^1(B_1)} + \|u\|_{X_k^{2,r}}$. The derivative of u with respect to h in direction η satisfies

$$\begin{aligned} \|u'(h)[\eta]\|_{X_{k}^{2,r}} &\leq C_{0} \|\eta\|_{H^{r_{0}+2}(\mathbb{S}^{2})} (\|\psi\|_{H^{\frac{3}{2}+r}(\mathbb{S}^{2})} + C_{r,k} \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^{2})}) \\ &+ (\chi_{r>r_{0}})C_{r,k} (\|\eta\|_{H^{r+2}(\mathbb{S}^{2})} + \|\eta\|_{H^{r_{0}+2}(\mathbb{S}^{2})} \|h\|_{H^{r+2}(\mathbb{S}^{2})}) \|\psi\|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})}, \quad (5.118)\end{aligned}$$

and the second derivative satisfies

$$\begin{aligned} \|u''(h)[\eta_{1},\eta_{2}]\|_{X_{k}^{2,r}} &\leq C_{0}\|\eta_{1}\|_{H^{r_{0}+2}(\mathbb{S}^{2})}\|\eta_{2}\|_{H^{r_{0}+2}(\mathbb{S}^{2})}(\|\psi\|_{H^{\frac{3}{2}+r}(\mathbb{S}^{2})} + C_{r,k}\|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^{2})}) \\ &+ (\chi_{r>r_{0}})C_{r,k}\{\|\eta_{1}\|_{H^{r+2}(\mathbb{S}^{2})}\|\eta_{2}\|_{H^{r_{0}+2}(\mathbb{S}^{2})} + \|\eta_{1}\|_{H^{r_{0}+2}(\mathbb{S}^{2})}\|\eta_{2}\|_{H^{r+2}(\mathbb{S}^{2})} \\ &+ \|\eta_{1}\|_{H^{r_{0}+2}(\mathbb{S}^{2})}\|\eta_{2}\|_{H^{r_{0}+2}(\mathbb{S}^{2})}\|h\|_{H^{r+2}(\mathbb{S}^{2})}\}\|\psi\|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})}. \end{aligned}$$
(5.119)

Proof. To prove (5.117), use (5.96), (5.98), (5.112), with $Bc_0 ||h||_{H^{r_0+2}(\mathbb{S}^2)} \leq 1/2$, and fix $\delta_0 := 1/(2Bc_0)$. To prove (5.118) and (5.119), recall that $u'_n(h)[\eta] = n \mathring{u}_n[h, \ldots, h, \eta]$ and $u''_n(h)[\eta_1, \eta_2] = n(n-1)\mathring{u}_n[h, \ldots, h, \eta_1, \eta_2]$ and use (5.116).

Lemma 5.17. For $||h||_{H^{r_0+2}(\mathbb{S}^2)} \leq \delta_0$, the function u in Proposition 5.16 satisfies

$$\begin{aligned} \|(\nabla u)\|_{\mathbb{S}^{2}}\|_{H^{r+\frac{1}{2}}(\mathbb{S}^{2})} &\leq C_{0}\|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^{2})} + C_{r}\|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^{2})} + (\chi_{r>r_{0}})C_{r}\|h\|_{H^{2+r}(\mathbb{S}^{2})}\|\psi\|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})} \\ &\leq C_{0}'\|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^{2})} + (\chi_{r>r_{0}})C_{r}(1+\|h\|_{H^{2+r}(\mathbb{S}^{2})})\|\psi\|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})} \tag{5.120} \end{aligned}$$

for all real $r \ge 0$, where C_r depends on r and it is increasing in r, while C_0, C'_0, δ_0 are independent of r; δ_0, r_0 are given by Proposition 5.16. Also,

$$\|\langle \nabla u, x \rangle\|_{\mathbb{S}^2}\|_{H^{r+\frac{1}{2}}(\mathbb{S}^2)} \le RHS \text{ of } (5.120).$$
(5.121)

Moreover,

$$\begin{aligned} \| (\nabla \{u'(h)[\eta]\}) \|_{\mathbb{S}^{2}} \|_{H^{r+\frac{1}{2}}(\mathbb{S}^{2})} &\leq C_{0} \|\eta\|_{H^{r_{0}+2}(\mathbb{S}^{2})} \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^{2})} \\ &+ (\chi_{r>r_{0}}) C_{r}(\|\eta\|_{H^{r+2}(\mathbb{S}^{2})} + \|h\|_{H^{r+2}(\mathbb{S}^{2})} \|\eta\|_{H^{r_{0}+2}(\mathbb{S}^{2})}) \|\psi\|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})}, \end{aligned}$$
(5.122)
$$\| (\nabla \{u''(h)[\eta_{1},\eta_{2}]\}) \|_{\mathbb{S}^{2}} \|_{H^{r+\frac{1}{2}}(\mathbb{S}^{2})} &\leq C_{0} \|\eta_{1}\|_{H^{r_{0}+2}(\mathbb{S}^{2})} \|\eta_{2}\|_{H^{r_{0}+2}(\mathbb{S}^{2})} \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^{2})} \\ &+ (\chi_{r>r_{0}}) C_{r}\{\|\eta_{1}\|_{H^{r+2}(\mathbb{S}^{2})} \|\eta_{2}\|_{H^{r_{0}+2}(\mathbb{S}^{2})} + \|\eta_{1}\|_{H^{r_{0}+2}(\mathbb{S}^{2})} \|\eta_{2}\|_{H^{r+2}(\mathbb{S}^{2})} \\ &+ \|h\|_{H^{r+2}(\mathbb{S}^{2})} \|\eta_{1}\|_{H^{r_{0}+2}(\mathbb{S}^{2})} \|\eta_{2}\|_{H^{r_{0}+2}(\mathbb{S}^{2})}\} \|\psi\|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})}, \end{aligned}$$
(5.123)

$$\|\langle \nabla\{u'(h)[\eta]\}, x\rangle\|_{\mathbb{S}^2}\|_{H^{r+\frac{1}{2}}(\mathbb{S}^2)} \le RHS \text{ of } (5.122), \tag{5.124}$$

$$\|\langle \nabla\{u''(h)[\eta_1,\eta_2]\}, x\rangle|_{\mathbb{S}^2}\|_{H^{r+\frac{1}{2}}(\mathbb{S}^2)} \le RHS \text{ of } (5.123).$$
(5.125)

Proof. By (5.57), one has $\nabla u = \nabla(u\zeta_0)$ on \mathbb{S}^2 , because $\zeta_0 = 1$ on $|x| \ge 1 - (\delta/2)$. By the definition (5.18) of the $H^r(\mathbb{S}^2)$ norm and the trace estimate (5.43), one has

$$\begin{split} \|(\nabla u)|_{\mathbb{S}^{2}}\|_{H^{r+\frac{1}{2}}(\mathbb{S}^{2})} &= \|(\nabla(u\zeta_{0}))|_{\mathbb{S}^{2}}\|_{H^{r+\frac{1}{2}}(\mathbb{S}^{2})} \\ &= \sum_{j=1}^{N} \|\{\psi_{j}\nabla(u\zeta_{0})\}\circ\mathsf{g}_{j}(\cdot,0)\|_{H^{r+\frac{1}{2}}(\mathbb{R}^{2})} \leq C\sum_{j=1}^{N} \|\{\psi_{j}\nabla(u\zeta_{0})\}\circ\mathsf{g}_{j}\|_{H^{1,r}(\mathbb{R}^{3}_{+})}. \end{split}$$

We proceed similarly as in the proof of the previous lemmas. One has $\psi_j \nabla(u\zeta_0) = \nabla(u\zeta_0\psi_j) - u\zeta_0\nabla\psi_j$. Concerning the term $\nabla(u\zeta_0\psi_j)$, one has $\{\nabla(u\zeta_0\psi_j)\} \circ \mathbf{g}_j = (D\mathbf{g}_j)^{-T}\nabla\{(u\zeta_0\psi_j) \circ \mathbf{g}_j\}$, and, by (5.49), (5.35), (5.36), (5.61),

$$\begin{split} &\sum_{j=1}^{N} \| (D\mathbf{g}_{j})^{-T} \nabla \{ (u\zeta_{0}\psi_{j}) \circ \mathbf{g}_{j} \} \|_{H^{1,r}(\mathbb{R}^{3}_{+})} \\ &\leq C_{0} \sum_{j=1}^{N} \| \nabla \{ (u\zeta_{0}\psi_{j}) \circ \mathbf{g}_{j} \} \|_{H^{1,r}(\mathbb{R}^{3}_{+})} + C_{r} \sum_{j=1}^{N} \| \nabla \{ (u\zeta_{0}\psi_{j}) \circ \mathbf{g}_{j} \} \|_{H^{1,0}(\mathbb{R}^{3}_{+})} \\ &\leq C_{0} \sum_{j=1}^{N} \| (u\zeta_{0}\psi_{j}) \circ \mathbf{g}_{j} \|_{H^{2,r}(\mathbb{R}^{3}_{+})} + C_{r} \sum_{j=1}^{N} \| (u\zeta_{0}\psi_{j}) \circ \mathbf{g}_{j} \|_{H^{2,0}(\mathbb{R}^{3}_{+})} = C_{0} \| u \|_{X^{2,r}_{0}} + C_{r} \| u \|_{X^{2,0}_{0}}. \end{split}$$

Concerning the term $u\zeta_0\nabla\psi_j$, by (5.19) one has $(u\zeta_0\nabla\psi_j)\circ \mathbf{g}_j = ((\nabla\psi_j)\circ \mathbf{g}_j)\sum_{\ell\in\mathcal{A}_j}(u\zeta_0\psi_\ell)\circ \mathbf{g}_j = ((\nabla\psi_j)\circ \mathbf{g}_j)\sum_{\ell\in\mathcal{A}_j}(u\zeta_0\psi_\ell)\circ \mathbf{g}_\ell \circ (\mathbf{f}_\ell\circ \mathbf{g}_j)$, where $\mathcal{A}_j = \{\ell: K_j\cap K_\ell\neq\emptyset\}$, and, by (5.49), (5.32), Lemma 5.4, (5.51), (5.61),

$$\begin{split} &\sum_{j=1}^{N} \sum_{\ell \in \mathcal{A}_{j}} \| ((\nabla \psi_{j}) \circ \mathbf{g}_{j}) (u\zeta_{0}\psi_{\ell}) \circ \mathbf{g}_{\ell} \circ (\mathbf{f}_{\ell} \circ \mathbf{g}_{j}) \|_{H^{1,r}(\mathbb{R}^{3}_{+})} \\ &\leq C_{r} \sum_{\ell=1}^{N} \sum_{j \in \mathcal{A}_{\ell}} \| (u\zeta_{0}\psi_{\ell}) \circ \mathbf{g}_{\ell} \circ (\mathbf{f}_{\ell} \circ \mathbf{g}_{j}) \|_{H^{1,r}(\mathbb{R}^{3}_{+})} \\ &\leq C_{r}' \sum_{\ell=1}^{N} \| (u\zeta_{0}\psi_{\ell}) \circ \mathbf{g}_{\ell} \|_{H^{1,r}(\mathbb{R}^{3}_{+})} = C_{r}' \| u \|_{X^{1,r}_{0}} \leq C_{r}' \| u \|_{X^{2,r-1}_{0}} \end{split}$$

Hence

$$\|(\nabla u)|_{\mathbb{S}^2}\|_{H^{r+\frac{1}{2}}(\mathbb{S}^2)} \le C_0 \|u\|_{X_0^{2,r}} + C_r(\|u\|_{X_0^{2,r-1}} + \|u\|_{X_0^{2,0}}).$$

Applying (5.117) with k = 0, and then using the interpolation inequality (5.52) like in (5.103) if r - 1 > 0, we obtain the first inequality in (5.120). The second inequality in (5.120) is obtained from the first one because

$$C_{r} \|\psi\|_{H^{\frac{3}{2}}(\mathbb{S}^{2})} \leq C_{r_{0}} \|\psi\|_{H^{r+\frac{3}{2}}(\mathbb{S}^{2})} + (\chi_{r>r_{0}})C_{r} \|\psi\|_{H^{r_{0}+\frac{3}{2}}(\mathbb{S}^{2})}$$
(5.126)

for all $r \ge 0$. To prove (5.126), consider the two cases $r \in [0, r_0]$ and $r > r_0$ separately, and use the fact that $C_r \le C_{r_0}$ for $r \in [0, r_0]$. To prove (5.121), use (5.120) and the product estimate (5.53). The proof of (5.122), (5.123), (5.124), (5.125) is similar, using (5.118), (5.119), (5.53).

Now that the radial derivative $\langle \nabla u, x \rangle|_{\mathbb{S}^2}$ of the solution $u = \tilde{\Phi}$ of problem (3.13) has been estimated, we can estimate the Dirichlet-Neumann operator using identity (4.4). To this purpose, we estimate the other terms appearing in (4.4).

and

Lemma 5.18. Given $s_0 > 1$, there exists $\delta_0 > 0$ such that, for $\|h\|_{H^{s_0+1}(\mathbb{S}^2)} \leq \delta_0$, one has

$$\begin{split} &\|(1+h)^{\alpha}\|_{H^{s}(\mathbb{S}^{2})} \leq C_{s}(1+\|h\|_{H^{s}(\mathbb{S}^{2})}), \quad \alpha \in \{-1,-2\}, \\ &\|\{(1+h)^{2}+|\nabla_{\mathbb{S}^{2}}h|^{2}\}^{\pm\frac{1}{2}}\|_{H^{s}(\mathbb{S}^{2})} \leq C_{s}(1+\|h\|_{H^{s+1}(\mathbb{S}^{2})}), \\ &\|\langle\nabla_{\mathbb{S}^{2}}h,\nabla_{\mathbb{S}^{2}}\psi\rangle\|_{H^{s}(\mathbb{S}^{2})} \leq C_{s_{0}}\|h\|_{H^{s_{0}+1}(\mathbb{S}^{2})}\|\psi\|_{H^{s+1}(\mathbb{S}^{2})} + (\chi_{s>s_{0}})C_{s}\|h\|_{H^{s+1}(\mathbb{S}^{2})}\|\psi\|_{H^{s_{0}+1}(\mathbb{S}^{2})} \\ \end{aligned}$$

for all real $s \ge 0$. The constant C_s depends on s, s_0 and it is increasing in s, while C_{s_0}, δ_0 depend on s_0 and are independent of s.

Proof. To estimate $(1+h)^{\alpha}$, we write it as a power series around h = 0 and apply estimate (5.54) to each term of the series. We estimate $(1+g)^p$, $p = \pm \frac{1}{2}$, in the same way, with $g = 2h + h^2 + |\nabla_{\mathbb{S}^2} h|^2$, then we estimate g by (5.53), (5.55). The estimate for $\langle \nabla_{\mathbb{S}^2} h, \nabla_{\mathbb{S}^2} \psi \rangle$ follows from (5.53), (5.55). \Box

Theorem 5.19 (Tame estimate and analyticity of the Dirichlet-Neumann operator). Let $1 < s_0 \leq 3/2$. There exists $\delta_0 > 0$ such that, for $\|h\|_{H^{s_0+\frac{3}{2}}(\mathbb{S}^2)} \leq \delta_0$, the Dirichlet-Neumann operator $G(h)\psi$ satisfies

$$\|G(h)\psi\|_{H^{s}(\mathbb{S}^{2})} \leq C_{0}\|\psi\|_{H^{s+1}(\mathbb{S}^{2})} + (\chi_{s>s_{0}})C_{s}(1+\|h\|_{H^{s+\frac{3}{2}}(\mathbb{S}^{2})})\|\psi\|_{H^{s_{0}+1}(\mathbb{S}^{2})}$$
(5.127)

for all real $s \ge 1/2$. The constant C_s depends on s, s_0 and it is increasing in s, while C_0, δ_0 depend on s_0 and they are independent of s. For any $\psi \in H^{s+1}(\mathbb{S}^2)$, the map $h \mapsto G(h)\psi$ is analytic from the ball $\{h \in H^{s_0+\frac{3}{2}}(\mathbb{S}^2) \cap H^{s+\frac{3}{2}}(\mathbb{S}^2) : \|h\|_{H^{s_0+\frac{3}{2}}(\mathbb{S}^2)} < \delta_0\}$ to $H^s(\mathbb{S}^2)$, and its first and second derivative satisfy

$$\|G'(h)[\eta]\psi\|_{H^{s}(\mathbb{S}^{2})} \leq C_{0}\|\eta\|_{H^{s_{0}+\frac{3}{2}}(\mathbb{S}^{2})}\|\psi\|_{H^{s+1}(\mathbb{S}^{2})} + (\chi_{s>s_{0}})C_{s}(\|\eta\|_{H^{s+\frac{3}{2}}(\mathbb{S}^{2})} + \|h\|_{H^{s+\frac{3}{2}}(\mathbb{S}^{2})}\|\eta\|_{H^{s_{0}+\frac{3}{2}}(\mathbb{S}^{2})})\|\psi\|_{H^{s_{0}+1}(\mathbb{S}^{2})}$$

$$(5.128)$$

and

$$\begin{aligned} \|G''(h)[\eta_{1},\eta_{2}]\psi\|_{H^{s}(\mathbb{S}^{2})} &\leq C_{0}\|\eta_{1}\|_{H^{s_{0}+\frac{3}{2}}(\mathbb{S}^{2})}\|\eta_{2}\|_{H^{s_{0}+\frac{3}{2}}(\mathbb{S}^{2})}\|\psi\|_{H^{s+1}(\mathbb{S}^{2})} \\ &+ (\chi_{s>s_{0}})C_{s}\{\|\eta_{1}\|_{H^{s+\frac{3}{2}}(\mathbb{S}^{2})}\|\eta_{2}\|_{H^{s_{0}+\frac{3}{2}}(\mathbb{S}^{2})} + \|\eta_{1}\|_{H^{s_{0}+\frac{3}{2}}(\mathbb{S}^{2})}\|\eta_{2}\|_{H^{s+\frac{3}{2}}(\mathbb{S}^{2})} \\ &+ \|h\|_{H^{s+\frac{3}{2}}(\mathbb{S}^{2})}\|\eta_{1}\|_{H^{s_{0}+\frac{3}{2}}(\mathbb{S}^{2})}\|\eta_{2}\|_{H^{s_{0}+\frac{3}{2}}(\mathbb{S}^{2})}\}\|\psi\|_{H^{s_{0}+1}(\mathbb{S}^{2})}. \end{aligned}$$
(5.129)

Proof. To prove (5.127), consider the Dirichlet-Neumann formula (4.4), use estimate (5.121) with $r = s - \frac{1}{2}$, $r_0 = s_0 - \frac{1}{2}$ for the radial derivative $\langle \nabla \tilde{\Phi}, x \rangle = \langle \nabla u, x \rangle$ on \mathbb{S}^2 , use Lemma 5.18 for the other terms appearing in (4.4), and apply the product estimate (5.53). The proof of (5.128) and (5.129) is similar, using estimates (5.124) and (5.125).

As for $r_0 \leq 1$ in Lemma 5.14, the assumption $s_0 \leq 3/2$ in Theorem 5.19 is not really necessary, but only convenient.

6 Traveling waves

In this section we construct traveling waves. Let h, ψ be functions of the form

$$h(t,x) = \eta(R(\omega t)x), \quad \psi(t,x) = \beta(R(\omega t)x), \quad t \in \mathbb{R}, \ x \in \mathbb{S}^2,$$
(6.1)

where $\eta, \beta : \mathbb{S}^2 \to \mathbb{R}$ are scalar functions defined on \mathbb{S}^2 , independent of time, $\omega \in \mathbb{R}$ is the angular velocity parameter, and $R(\vartheta)$ is the rotation matrix

$$R(\vartheta) = \begin{pmatrix} \cos\vartheta & -\sin\vartheta & 0\\ \sin\vartheta & \cos\vartheta & 0\\ 0 & 0 & 1 \end{pmatrix}.$$
 (6.2)

The transformation law for the time derivative $\partial_t h$, $\partial_t \psi$ is the following one.

Lemma 6.1. The matrix $R(\vartheta)$ in (6.2) satisfies

$$\partial_{\vartheta} R(\vartheta) R^{T}(\vartheta) = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} =: \mathcal{J}$$
(6.3)

for all $\vartheta \in \mathbb{R}$. The function h(t, x) defined in (6.1) satisfies

$$\partial_t h(t,x) = \omega \langle \mathcal{J}y, \nabla_{\mathbb{S}^2} \eta(y) \rangle \tag{6.4}$$

where $y = R(\omega t)x$, for all $x \in \mathbb{S}^2$, all $t, \omega \in \mathbb{R}$.

Proof. The proof is a straightforward calculation.

For the other terms of the equations, the time variable plays the role of a parameter. We have the following transformation laws.

Lemma 6.2. Let $M \in Mat_{3\times 3}(\mathbb{R})$ be an orthogonal matrix, i.e., $MM^T = M^TM = I$. Let $h(x) = \eta(Mx)$ and $\psi(x) = \beta(Mx)$ for all $x \in \mathbb{S}^2$, where $h, \psi, \eta, \beta : \mathbb{S}^2 \to \mathbb{R}$. Then

$$\mathcal{E}_0 h)(x) = (\mathcal{E}_0 \eta)(Mx) \quad \forall x \in \mathbb{R}^3 \setminus \{0\},$$
(6.5)

$$\nabla_{\mathbb{S}^2} h(x) = (c_0 \eta)(Mx) \quad \forall x \in \mathbb{R} \setminus \{0\},$$

$$\nabla_{\mathbb{S}^2} h(x) = M^T (\nabla_{\mathbb{S}^2} \eta)(Mx) \quad \forall x \in \mathbb{S}^2,$$

$$\nabla_{\mathbb{S}^2} h(x) = |(\nabla_{\mathbb{S}^2} \eta)(Mx)| \quad \forall x \in \mathbb{S}^2,$$

$$(6.6)$$

$$\nabla_{\mathbb{S}^2} h(x)| = |(\nabla_{\mathbb{S}^2} \eta)(Mx)| \quad \forall x \in \mathbb{S}^2,$$
(6.7)

$$\Delta_{\mathbb{S}^2} h(x) = (\Delta_{\mathbb{S}^2} \eta)(Mx) \quad \forall x \in \mathbb{S}^2,$$
(6.8)

$$\langle (D_{\mathbb{S}^2}^2 h)(x) \nabla_{\mathbb{S}^2} h(x), \nabla_{\mathbb{S}^2} h(x) \rangle = \langle (D_{\mathbb{S}^2}^2 \eta)(Mx) (\nabla_{\mathbb{S}^2} \eta)(Mx), (\nabla_{\mathbb{S}^2} \eta)(Mx) \rangle \quad \forall x \in \mathbb{S}^2, \tag{6.9}$$

$$H(h)(x) = H(\eta)(Mx) \quad \forall x \in \mathbb{S}^2,$$

$$\nabla_{\text{reg}} \psi(x) = /(\nabla_{\text{reg}} x)(Mx) \quad (\nabla_{\text{reg}} \beta)(Mx)) \quad \forall x \in \mathbb{S}^2$$
(6.10)

$$\langle \nabla_{\mathbb{S}^2} h(x), \nabla_{\mathbb{S}^2} \psi(x) \rangle = \langle (\nabla_{\mathbb{S}^2} \eta)(Mx), (\nabla_{\mathbb{S}^2} \beta)(Mx) \rangle \quad \forall x \in \mathbb{S}^2,$$
(6.11)

$$P_h(x) = M^T P_\eta(Mx) M \quad \forall x \in \mathbb{R}^3 \setminus \{0\},$$
(6.12)

$$(G(h)\psi)(x) = (G(\eta)\beta)(Mx) \quad \forall x \in \mathbb{S}^2,$$
(6.13)

where $\mathcal{E}_0 h$ is the 0-homogeneous extension of h, and $P_h(x) = (1 + h_0(x))I - \nabla h_0(x) \otimes x - x \otimes I$ $\nabla h_0(x) + (1+h_0(x))^{-1} |\nabla h_0(x)|^2 x \otimes x$, with $h_0 = \mathcal{E}_0 h$, is the matrix in (3.14) and (3.15).

Proof. All the properties can be easily proved using the identities $MM^T = M^TM = I$, which imply that the map $x \mapsto Mx$ is an isometry; to prove (6.13), use also the uniqueness of the solution of the elliptic problem (3.13).

By Lemma 6.1 and Lemma 6.2, system (1.16), (1.17) for the unknowns h, ψ satisfying ansatz (6.1) becomes the equation

$$\mathcal{F}(\omega, u) = 0 \tag{6.14}$$

for the unknown $u = (\eta, \beta)$ on \mathbb{S}^2 , where

$$\mathcal{F}(\omega, u) := (\mathcal{F}_1(\omega, u), \mathcal{F}_2(\omega, u)), \tag{6.15}$$

$$\mathcal{F}_1(\omega, u) := \omega \mathcal{M}\eta - \frac{\sqrt{(1+\eta)^2 + |\nabla_{\mathbb{S}^2}\eta|^2}}{1+\eta} G(\eta)\beta,$$
(6.16)

$$\mathcal{F}_2(\omega, u) := \omega \mathcal{M}\beta - \frac{1}{2} \Big(G(\eta)\beta + \frac{\langle \nabla_{\mathbb{S}^2}\beta, \nabla_{\mathbb{S}^2}\eta \rangle}{(1+\eta)\sqrt{(1+\eta)^2 + |\nabla_{\mathbb{S}^2}\eta|^2}} \Big)^2 + \frac{|\nabla_{\mathbb{S}^2}\beta|^2}{2(1+\eta)^2} + \sigma_0 H_\eta, \quad (6.17)$$

and \mathcal{M} is the linear operator

$$\mathcal{M}f(x) := \langle \mathcal{J}x, \nabla_{\mathbb{S}^2} f(x) \rangle, \quad x \in \mathbb{S}^2,$$
(6.18)

for any $f: \mathbb{S}^2 \to \mathbb{R}$. Note that, since $\mathcal{J}x$ belongs to the tangent space $T_x(\mathbb{S}^2)$, one also has

$$\mathcal{M}f(x) = \langle \mathcal{J}x, \nabla \tilde{f}(x) \rangle = (x_1 \partial_{x_2} - x_2 \partial_{x_1}) \tilde{f}(x) \quad \forall x \in \mathbb{S}^2,$$
(6.19)

for any extension \tilde{f} of f to an open neighborhood of \mathbb{S}^2 . We also observe that $\mathcal{F}(\omega, 0) = 0$ for all $\omega \in \mathbb{R}$.

Lemma 6.3. Let $s_0, s \in \mathbb{R}$, $s \ge s_0 > 1$, and let

$$U := \{ u = (\eta, \beta) : \eta \in H^{s + \frac{3}{2}}(\mathbb{S}^2, \mathbb{R}), \quad \beta \in H^{s + 1}(\mathbb{S}^2, \mathbb{R}), \quad \|\eta\|_{H^{s_0 + \frac{3}{2}}(\mathbb{S}^2)} < \delta_0 \},$$

where δ_0 is the constant in Theorem 5.19. Then $\mathcal{F}_1(\omega, u) \in H^s(\mathbb{S}^2, \mathbb{R})$, $\mathcal{F}_2(\omega, u) \in H^{s-\frac{1}{2}}(\mathbb{S}^2, \mathbb{R})$ for all $u \in U$, $\omega \in \mathbb{R}$, and the map

$$\mathcal{F}: \mathbb{R} \times U \to H^s(\mathbb{S}^2, \mathbb{R}) \times H^{s - \frac{1}{2}}(\mathbb{S}^2, \mathbb{R})$$

is analytic.

Proof. It is a straightforward consequence of the properties and estimates proved in Section 5. \Box

6.1 The linearized operator at zero

We calculate the linearized operator $L := \partial_u \mathcal{F}(\omega, 0)$ at u = 0, which is the linear operator

$$L: H^{s+\frac{3}{2}}(\mathbb{S}^2) \times H^{s+1}(\mathbb{S}^2) \to H^s(\mathbb{S}^2) \times H^{s-\frac{1}{2}}(\mathbb{S}^2),$$
$$L(\eta,\beta) = \begin{pmatrix} \omega \mathcal{M}\eta - G(0)\beta\\ \omega \mathcal{M}\beta - \sigma_0(2\eta + \Delta_{\mathbb{S}^2}\eta) \end{pmatrix} = \begin{pmatrix} \omega \mathcal{M} & -G(0)\\ -\sigma_0(2 + \Delta_{\mathbb{S}^2}) & \omega \mathcal{M} \end{pmatrix} \begin{pmatrix} \eta\\ \beta \end{pmatrix}.$$
(6.20)

The operators G(0) and $2 + \Delta_{\mathbb{S}^2}$ are diagonalized by the real spherical harmonics, with

$$G(0)\varphi_{\ell} = \ell\varphi_{\ell}, \quad -(2 + \Delta_{\mathbb{S}^2})\varphi_{\ell} = (\ell + 2)(\ell - 1)\varphi_{\ell} \quad \forall \varphi_{\ell} \in \mathcal{H}_{\ell}(\mathbb{S}^2, \mathbb{R}), \quad \ell \in \mathbb{N}_0,$$

where $\mathcal{H}_{\ell}(\mathbb{S}^2, \mathbb{R})$ is the space of the real spherical harmonics of degree ℓ ; as is well known, it is a vector space of dimension $(2\ell + 1)$ on \mathbb{R} . The operator \mathcal{M} can also be block-diagonalized by real spherical harmonics; in particular, the restriction of \mathcal{M} to $\mathcal{H}_{\ell}(\mathbb{S}^2, \mathbb{R})$ can be represented by a block-diagonal matrix with ℓ 2-blocks $\begin{pmatrix} 0 & -m \\ m & 0 \end{pmatrix}$, $m = 1, \ldots, \ell$, and one 1-block 0 (using complex spherical harmonics, \mathcal{M} becomes diagonal with complex eigenvalues $im, m = -\ell, \ldots, \ell$).

As is well known, an $L^2(\mathbb{S}^2, \mathbb{R})$ orthonormal basis of $\mathcal{H}_{\ell}(\mathbb{S}^2, \mathbb{R})$ is given by the classical real spherical harmonics

$$Y_{\ell,m}^{(\cos)}(\theta,\phi) = c_{\ell}^{(m)}(\sin\theta)^m P_{\ell}^{(m)}(\cos\theta)\cos(m\phi), \quad m = 0,\dots,\ell,$$

$$Y_{\ell,m}^{(\sin)}(\theta,\phi) = c_{\ell}^{(m)}(\sin\theta)^m P_{\ell}^{(m)}(\cos\theta)\sin(m\phi), \quad m = 1,\dots,\ell,$$
 (6.21)

commonly written as functions of the angles $\theta \in [0, \pi]$, $\phi \in [0, 2\pi]$ expressing any point $x \in \mathbb{S}^2$ in spherical coordinates $x_1 = \sin \theta \cos \phi$, $x_2 = \sin \theta \sin \phi$, $x_3 = \cos \theta$. Here $P_{\ell}^{(m)}(t)$ is the *m*-th derivative of the ordinary Legendre polynomial $P_{\ell}(t)$, which is a polynomial of degree ℓ with real coefficients, with parity $P_{\ell}(-t) = (-1)^{\ell} P_{\ell}(t)$, and $c_{\ell}^{(m)} \in \mathbb{R}$ is a normalizing coefficient; see, e.g., [7], Example 2.48 in Section 2.11. For $a_m, b_m \in \mathbb{R}$, the sum $a_m \cos(m\phi) + b_m \sin(m\phi)$ is the real part of the complex number $(a_m - ib_m)e^{im\phi}$. Hence any linear combination of (6.21) with real coefficients a_m, b_m can be written as

$$\sum_{m=0}^{\ell} a_m Y_{\ell,m}^{(\cos)}(\theta,\phi) + \sum_{m=1}^{\ell} b_m Y_{\ell,m}^{(\sin)}(\theta,\phi)$$

= $a_0 c_{\ell}^{(0)} P_{\ell}(\cos\theta) + \sum_{m=1}^{\ell} c_{\ell}^{(m)} P_{\ell}^{(m)}(\cos\theta) \operatorname{Re} \{(a_m - ib_m)(\sin\theta)^m e^{im\phi}\},$

which, in Cartesian coordinates, becomes

$$= a_0 c_{\ell}^{(0)} P_{\ell}(x_3) + \sum_{m=1}^{\ell} c_{\ell}^{(m)} P_{\ell}^{(m)}(x_3) \operatorname{Re} \{ (a_m - ib_m)(x_1 + ix_2)^m \}.$$

Hence the functions

$$\varphi_{\ell,0}(x) := c_{\ell}^{(0)} P_{\ell}(x_3), \qquad \varphi_{\ell,m}^{(\text{Re}\,)}(x) := c_{\ell}^{(m)} P_{\ell}^{(m)}(x_3) \text{Re}\left[(x_1 + ix_2)^m\right], \quad m = 1, \dots, \ell,$$
$$\varphi_{\ell,m}^{(\text{Im}\,)}(x) := c_{\ell}^{(m)} P_{\ell}^{(m)}(x_3) \text{Im}\left[(x_1 + ix_2)^m\right], \quad m = 1, \dots, \ell, \qquad (6.22)$$

with $x \in \mathbb{S}^2$, form an $L^2(\mathbb{S}^2, \mathbb{R})$ orthonormal basis of the real vector space $\mathcal{H}_\ell(\mathbb{S}^2, \mathbb{R})$. For notation convenience, we denote

$$\varphi_{\ell,m} := \varphi_{\ell,m}^{(\text{Re})}, \quad \varphi_{\ell,-m} := \varphi_{\ell,m}^{(\text{Im})}, \quad m = 1, \dots, \ell.$$
(6.23)

Thus, $\{\varphi_{\ell,m} : m = -\ell, \dots, \ell\}$ is an $L^2(\mathbb{S}^2, \mathbb{R})$ orthonormal basis of $\mathcal{H}_\ell(\mathbb{S}^2, \mathbb{R})$. This is the basis of Legendre real spherical harmonics in Cartesian coordinates.

Lemma 6.4. One has $\mathcal{M}\varphi_{\ell,m} = -m\varphi_{\ell,-m}$ for all $m = -\ell, \ldots, \ell$, all $\ell \in \mathbb{N}_0$.

Proof. To apply (6.19), we observe that the functions in (6.22) have a natural extension, which we write without changing the notation, to a neighborhood of the sphere; such extensions are simply obtained by extending the validity of the formulae in (6.22). In general, these extensions are neither harmonic nor homogeneous, but (6.19) holds without requiring those properties. One has

$$(x_1\partial_{x_2} - x_2\partial_{x_1})\{(x_1 + ix_2)^m\} = im(x_1 + ix_2)^m$$

for all $m \in \mathbb{N}$, all $(x_1, x_2) \in \mathbb{R}^2$, and

$$(x_1\partial_{x_2} - x_2\partial_{x_1})\operatorname{Re}\{(x_1 + ix_2)^m\} = -m\operatorname{Im}\{(x_1 + ix_2)^m\}, (x_1\partial_{x_2} - x_2\partial_{x_1})\operatorname{Im}\{(x_1 + ix_2)^m\} = m\operatorname{Re}\{(x_1 + ix_2)^m\}.$$

Therefore, by (6.19), we obtain

$$\mathcal{M}\varphi_{\ell,0}(x) = 0, \quad \mathcal{M}\varphi_{\ell,m}^{(\mathrm{Re})}(x) = -m\varphi_{\ell,m}^{(\mathrm{Im})}, \quad \mathcal{M}\varphi_{\ell,m}^{(\mathrm{Im})}(x) = m\varphi_{\ell,m}^{(\mathrm{Re})}, \quad m = 1, \dots, \ell.$$
(6.24)
lling the notation in (6.23), this completes the proof.

Recalling the notation in (6.23), this completes the proof.

Given $(f,g) \in H^s(\mathbb{S}^2) \times H^{s-\frac{1}{2}}(\mathbb{S}^2)$, we study the equation $L(\eta,\beta) = (f,g)$. We use the real spherical harmonics $(\varphi_{\ell,m})$ of Lemma 6.4 to decompose

$$\eta = \sum_{(\ell,m)\in\mathcal{T}} \hat{\eta}_{\ell,m} \varphi_{\ell,m}, \quad \mathcal{T} = \bigcup_{\ell\in\mathbb{N}_0} \mathcal{T}_\ell, \quad \mathcal{T}_\ell = \{(\ell,m) : m = -\ell, \dots, \ell\},$$
(6.25)

with $\hat{\eta}_{\ell,m} \in \mathbb{R}$, and similarly for β, f, g . Hence

$$\mathcal{M}\eta = \sum_{(\ell,m)\in\mathcal{T}} \hat{\eta}_{\ell,m}(-m)\varphi_{\ell,-m} = \sum_{(\ell,m)\in\mathcal{T}} m\hat{\eta}_{\ell,-m}\varphi_{\ell,m}.$$

One has $L(\eta, \beta) = (f, g)$ if and only if

$$-\omega m \hat{\eta}_{\ell,m} - \ell \hat{\beta}_{\ell,-m} = \hat{f}_{\ell,-m}, \quad \sigma_0(\ell+2)(\ell-1)\hat{\eta}_{\ell,m} + \omega m \hat{\beta}_{\ell,-m} = \hat{g}_{\ell,m} \quad \forall (\ell,m) \in \mathcal{T}, \quad (6.26)$$

that is,

$$L_{\ell,m}\begin{pmatrix} \hat{\eta}_{\ell,m}\\ \hat{\beta}_{\ell,-m} \end{pmatrix} = \begin{pmatrix} \hat{f}_{\ell,-m}\\ \hat{g}_{\ell,m} \end{pmatrix} \quad \forall (\ell,m) \in \mathcal{T}, \quad L_{\ell,m} := \begin{pmatrix} -\omega m & -\ell\\ \sigma_0(\ell+2)(\ell-1) & \omega m \end{pmatrix}.$$
(6.27)

One has

$$\det L_{\ell,m} = -\omega^2 m^2 + \sigma_0 (\ell + 2)(\ell - 1)\ell, \qquad (6.28)$$

and bifurcation can only occur at values of ω such that (6.28) vanishes at some (ℓ, m) . Thus, we assume that

$$\omega = \sqrt{\sigma_0} \frac{\sqrt{(\ell_0 + 2)(\ell_0 - 1)\ell_0}}{m_0} \tag{6.29}$$

for some fixed integers ℓ_0, m_0 , with $\ell_0 \geq 2$ and $1 \leq m_0 \leq \ell_0$. With ω in (6.29), a pair (ℓ, m) gives det $L_{\ell m} = 0$ if and only if

$$(\ell+2)(\ell-1)\ell = c_0 m^2, \quad c_0 := (\ell_0 + 2)(\ell_0 - 1)\ell_0 m_0^{-2}.$$
 (6.30)

Lemma 6.5. Let $\ell_0, m_0 \in \mathbb{N}$, with $\ell_0 \geq 2$, $1 \leq m_0 \leq \ell_0$. Let $S \subset \mathcal{T}$ be the set of the pairs $(\ell, m) \in \mathcal{T}$ satisfying (6.30). Then S has a finite number of elements, which are $(\ell, m) = (\ell_0, m_0)$, $(\ell_0, -m_0)$, (1, 0), (0, 0), and possibly finitely many other pairs, all of which with $\ell \leq c_0$. Moreover, assuming (6.29), there exists a constant C > 0, depending on ℓ_0, σ_0 , such that

$$|\det L_{\ell,m}| \ge C\ell^3 \quad \forall (\ell,m) \in \mathcal{T} \setminus S.$$
 (6.31)

Proof. For any $(\ell, m) \in \mathcal{T}$ one has $m^2 \leq \ell^2$. If $(\ell, m) \in S$, then $(\ell+2)(\ell-1)\ell = c_0m^2 \leq c_0\ell^2$. For $\ell \geq 2$ one has $\ell^2 \leq (\ell+2)(\ell-1)$, and we deduce that $\ell \leq c_0$ for all $(\ell, m) \in S$ with $\ell \geq 2$. The bound $\ell \leq c_0$ holds also for $\ell = 0, 1$ because $c_0 \geq (\ell_0 + 2)(\ell_0 - 1)\ell_0^{-1} \geq \ell_0 \geq 2$.

bound $\ell \leq c_0$ holds also for $\ell = 0, 1$ because $c_0 \geq (\ell_0 + 2)(\ell_0 - 1)\ell_0^{-1} \geq \ell_0 \geq 2$. For $(\ell, m) \in \mathcal{T} \setminus S$, one has det $L_{\ell,m} \neq 0$. Moreover, det $L_{\ell,m} \geq \det L_{\ell\ell}$ because $|m| \leq \ell$, and for $\ell \geq 2c_0$ it holds det $L_{\ell\ell} = \sigma_0(\ell+2)(\ell-1)\ell - \sigma_0c_0\ell^2 \geq \sigma_0\ell^3 - \sigma_0c_0\ell^2 \geq \frac{1}{2}\sigma_0\ell^3$. On the other hand, min{ $|\det L_{\ell m}|\ell^{-3}: (\ell, m) \in \mathcal{T} \setminus S, \ \ell < 2c_0$ } is also positive because it is the minimum of a finite set of positive numbers.

For $(\ell, m) \in \mathcal{T} \setminus S$, one has det $L_{\ell,m} \neq 0$, and system (6.27) with $(\hat{f}_{\ell,-m}, \hat{g}_{\ell,m}) = (0,0)$ has only the trivial solution $(\hat{\eta}_{\ell,m}, \hat{\beta}_{\ell,-m}) = (0,0)$. For $(\ell,m) \in S$, we distinguish $\ell = 0$ from $\ell > 0$. For $(\ell,m) = (0,0)$, system (6.27) with $(\hat{f}_{0,0}, \hat{g}_{0,0}) = (0,0)$ has solutions $(\hat{\eta}_{0,0}, \hat{\beta}_{0,0}) = (0,\lambda), \lambda \in \mathbb{R}$. For $(\ell,m) \in S$ with $\ell \geq 1$, (6.27) with $(\hat{f}_{\ell,-m}, \hat{g}_{\ell,m}) = (0,0)$ has solutions $(\hat{\eta}_{\ell,m}, \hat{\beta}_{\ell,-m}) = \lambda(\ell, -\omega m),$ $\lambda \in \mathbb{R}$. Hence the kernel of the linear operator L is the finite dimensional space

$$V := \ker L = \left\{ \begin{pmatrix} \eta \\ \beta \end{pmatrix} = \lambda_{0,0} \begin{pmatrix} 0 \\ \varphi_{0,0} \end{pmatrix} + \sum_{\substack{(\ell,m) \in S \\ \ell \ge 1}} \lambda_{\ell,m} \begin{pmatrix} \ell \varphi_{\ell,m} \\ -\omega m \varphi_{\ell,-m} \end{pmatrix} : \lambda_{\ell,m} \in \mathbb{R} \right\}.$$
(6.32)

Its orthogonal complement in $L^2(\mathbb{S}^2) \times L^2(\mathbb{S}^2)$ (we denote $L^2(\mathbb{S}^2) = L^2(\mathbb{S}^2, \mathbb{R})$) is the vector space

$$W := \left\{ \begin{pmatrix} \eta \\ \beta \end{pmatrix} = \lambda_{0,0} \begin{pmatrix} \varphi_{0,0} \\ 0 \end{pmatrix} + \sum_{\substack{(\ell,m) \in S \\ \ell \ge 1}} \lambda_{\ell,m} \begin{pmatrix} \omega m \varphi_{\ell,m} \\ \ell \varphi_{\ell,-m} \end{pmatrix} + \sum_{\substack{(\ell,m) \in \mathcal{T} \setminus S \\ \hat{\beta}_{\ell,m} \varphi_{\ell,m}}} \begin{pmatrix} \hat{\eta}_{\ell,m} \varphi_{\ell,m} \\ \hat{\beta}_{\ell,m} \varphi_{\ell,m} \end{pmatrix} \\ : \lambda_{\ell,m}, \hat{\eta}_{\ell,m}, \hat{\beta}_{\ell,m} \in \mathbb{R}, \ (\eta,\beta) \in L^2(\mathbb{S}^2) \times L^2(\mathbb{S}^2) \right\}.$$

Thus $L^2(\mathbb{S}^2) \times L^2(\mathbb{S}^2) = V \oplus W$, and V and W are orthogonal with respect to the scalar product of $L^2(\mathbb{S}^2) \times L^2(\mathbb{S}^2)$. Moreover

$$H^{s+\frac{3}{2}}(\mathbb{S}^2) \times H^{s+1}(\mathbb{S}^2) = V \oplus W^s, \quad W^s := W \cap (H^{s+\frac{3}{2}}(\mathbb{S}^2) \times H^{s+1}(\mathbb{S}^2)).$$

For $(\ell, m) \in \mathcal{T} \setminus S$, given any $\hat{f}_{\ell,-m}, \hat{g}_{\ell,m}$, there exists a unique solution of system (6.27), which is

$$\hat{\eta}_{\ell,m} = \frac{\omega m f_{\ell,-m} + \ell \hat{g}_{\ell,m}}{\det L_{\ell,m}}, \quad \hat{\beta}_{\ell,-m} = \frac{-\sigma_0(\ell+2)(\ell-1)f_{\ell,-m} - \omega m \hat{g}_{\ell,m}}{\det L_{\ell,m}}.$$
(6.33)

For $(\ell, m) \in S$, we distinguish the cases $\ell = 0$ and $\ell > 0$. For $(\ell, m) = (0, 0)$, system (6.27) has a solution only if $\hat{f}_{0,0} = 0$, and, in that case, the solutions are the pairs $(\hat{\eta}_{0,0}, \hat{\beta}_{0,0})$ with

$$\hat{\eta}_{0,0} = -(2\sigma_0)^{-1}\hat{g}_{0,0}, \quad \hat{\beta}_{0,0} \in \mathbb{R}.$$
(6.34)

For $(\ell, m) \in S$ with $\ell \geq 1$, system (6.27) has a solution only if $\omega m \hat{f}_{\ell,-m} + \ell \hat{g}_{\ell,m} = 0$, and, in that case, the solutions are the pairs $(\hat{\eta}_{\ell,m}, \hat{\beta}_{\ell,-m})$ with

$$\hat{\beta}_{\ell,m} = -\ell^{-1}(\omega m \hat{\eta}_{\ell,m} + \hat{f}_{\ell,-m}), \quad \hat{\eta}_{\ell,m} \in \mathbb{R}.$$
(6.35)

Hence the range of L is contained in the space

$$R := \left\{ \begin{pmatrix} f \\ g \end{pmatrix} = \hat{g}_{0,0} \begin{pmatrix} 0 \\ \varphi_{0,0} \end{pmatrix} + \sum_{\substack{(\ell,m) \in S \\ \ell \ge 1}} \hat{f}_{\ell,-m} \begin{pmatrix} \varphi_{\ell,-m} \\ -\omega m \ell^{-1} \varphi_{\ell,m} \end{pmatrix} + \sum_{\substack{(\ell,m) \in \mathcal{T} \setminus S \\ \hat{g}_{\ell,m} \varphi_{\ell,m}}} \begin{pmatrix} \hat{f}_{\ell,m} \varphi_{\ell,m} \\ \hat{g}_{\ell,m} \varphi_{\ell,m} \end{pmatrix} \\ : \hat{f}_{\ell,m}, \hat{g}_{\ell,m} \in \mathbb{R}, \ (f,g) \in L^2(\mathbb{S}^2) \times L^2(\mathbb{S}^2) \right\}.$$
(6.36)

The orthogonal complement of R with respect to the scalar product of $L^2(\mathbb{S}^2) \times L^2(\mathbb{S}^2)$ is the finite-dimensional space

$$Z := \left\{ \begin{pmatrix} f \\ g \end{pmatrix} = \lambda_{0,0} \begin{pmatrix} \varphi_{0,0} \\ 0 \end{pmatrix} + \sum_{\substack{(\ell,m) \in S \\ \ell \ge 1}} \lambda_{\ell,m} \begin{pmatrix} \omega m \varphi_{\ell,-m} \\ \ell \varphi_{\ell,m} \end{pmatrix} : \lambda_{\ell,m} \in \mathbb{R} \right\}.$$
 (6.37)

Thus, $L^2(\mathbb{S}^2) \times L^2(\mathbb{S}^2) = R \oplus Z$, and R and Z are orthogonal with respect to the scalar product of $L^2(\mathbb{S}^2) \times L^2(\mathbb{S}^2)$. Moreover

$$H^{s}(\mathbb{S}^{2}) \times H^{s-\frac{1}{2}}(\mathbb{S}^{2}) = R^{s} \oplus Z, \quad R^{s} := R \cap (H^{s}(\mathbb{S}^{2}) \times H^{s-\frac{1}{2}}(\mathbb{S}^{2})).$$

Let $L|_{W^s}: W^s \to R^s$ be the restriction of L to W^s .

Lemma 6.6. The linear map $L|_{W^s}: W^s \to R^s$ is bijective. Its inverse $(L|_{W^s})^{-1}: R^s \to W^s$ is bounded, with

$$\|(L|_{W^s})^{-1}(f,g)\|_{H^{s+\frac{3}{2}}(\mathbb{S}^2)\times H^{s+1}(\mathbb{S}^2)} \le C_s\|(f,g)\|_{H^s(\mathbb{S}^2)\times H^{s-\frac{1}{2}}(\mathbb{S}^2)}$$
(6.38)

for all $(f,g) \in \mathbb{R}^s$. The constant C_s depends on σ_0, ℓ_0, s .

To prove the lemma, we use spectral norms for the spaces $H^{s}(\mathbb{S}^{2})$. Given any $\eta \in H^{s}(\mathbb{S}^{2})$ written as the series of spherical harmonics (6.25), we define

$$\|\eta\|_{*,s} := \left(|\hat{\eta}_{0,0}|^2 + \sum_{(\ell,m)\in\mathcal{T}\setminus\mathcal{T}_0} |\hat{\eta}_{\ell,m}|^2 \ell^{2s}\right)^{\frac{1}{2}}.$$
(6.39)

By [33], Remark 7.6 in Section 7.3, (6.39) and (5.18) are equivalent norms on $H^{s}(\mathbb{S}^{2})$.

Proof of Lemma 6.6. The map $L|_{W^s}$ is injective on W^s by construction; we prove that it is surjective onto R^s . Let $(f,g) \in R^s$, with coefficients $(\hat{f}_{\ell,m}, \hat{g}_{\ell,m})$. For $(\ell,m) \in \mathcal{T} \setminus S$, the solution $(\hat{\eta}_{\ell,m}, \hat{\beta}_{\ell,m})$ of system (6.27) is uniquely determined by (6.33). For $(\ell,m) = (0,0)$, the infinitely many solutions $(\hat{\eta}_{0,0}, \hat{\beta}_{0,0})$ of system (6.27) are given by (6.34), and the condition $(\eta, \beta) \in W$ selects just one of these, which is $\hat{\beta}_{0,0} = 0$. For $(\ell,m) \in S$ with $\ell \geq 1$, the infinitely many solutions $(\hat{\eta}_{\ell,m}, \hat{\beta}_{\ell,-m})$ of (6.27) are given by (6.35), and the condition $(\eta, \beta) \in W$ selects just one of these, which is

$$\hat{\eta}_{\ell,m} = \frac{-\omega m \hat{f}_{\ell,-m}}{\ell^2 + \omega^2 m^2} = \frac{\ell \hat{g}_{\ell,m}}{\ell^2 + \omega^2 m^2}, \quad \hat{\beta}_{\ell,-m} = \frac{-\ell \hat{f}_{\ell,-m}}{\ell^2 + \omega^2 m^2}.$$
(6.40)

Hence the inversion problem $L(\eta, \beta) = (f, g), (\eta, \beta) \in W^s$ has a unique candidate solution (η, β) determined by its coefficients $(\hat{\eta}_{\ell,m}, \hat{\beta}_{\ell,m})$. We have to prove that this candidate is an element of W^s . For $(\ell, m) \in \mathcal{T} \setminus S$, formula (6.33) and bound (6.31) imply that

$$|\hat{\eta}_{\ell,m}| \le C\ell^{-2}(|\hat{f}_{\ell,-m}| + |\hat{g}_{\ell,m}|), \quad |\hat{\beta}_{\ell,-m}| \le C(\ell^{-1}|\hat{f}_{\ell,-m}| + \ell^{-2}|\hat{g}_{\ell,m}|), \tag{6.41}$$

for some constant C > 0 depending on σ_0, ℓ_0 . From (6.41) it follows that

$$\ell^{s+\frac{3}{2}}|\hat{\eta}_{\ell,m}| + \ell^{s+1}|\hat{\beta}_{\ell,-m}| \le 2C(\ell^s|\hat{f}_{\ell,-m}| + \ell^{s-\frac{1}{2}}|\hat{g}_{\ell,m}|)$$
(6.42)

for all $(\ell, m) \in \mathcal{T} \setminus S$. For $(\ell, m) = (0, 0)$ one has $|\hat{\eta}_{0,0}| = (2\sigma_0)^{-1} |\hat{g}_{0,0}|$, $\hat{\beta}_{0,0} = 0$. For $(\ell, m) \in S$ with $\ell \geq 1$, (6.35) implies that $|\hat{\eta}_{\ell,m}| \leq \ell^{-1} |\hat{g}_{\ell,m}|$ and $|\hat{\beta}_{\ell,m}| \leq \ell^{-1} |\hat{f}_{\ell,m}|$, but ℓ is in the bounded interval $1 \leq \ell \leq c_0$, see Lemma 6.5, and therefore $\ell^{s+\frac{1}{2}} \leq c_0 \ell^{s-\frac{1}{2}}$. This implies that inequality (6.42) also holds for $(\ell, m) \in S$ with $\ell \geq 1$, with a (possibly different) constant C depending on σ_0, ℓ_0 . As a consequence, one has

$$\|\eta\|_{*,s+\frac{3}{2}}^{2} + \|\beta\|_{*,s+1}^{2} \le C(\|f\|_{*,s}^{2} + \|g\|_{*,s-\frac{1}{2}}^{2}).$$

Hence $(\eta, \beta) \in W^s$, the inverse map $(L|_{W^s})^{-1} : \mathbb{R}^s \to W^s$ is well-defined and bounded, and, from the equivalence of the norms (6.39) and (5.18), the proof is complete.

Remark 6.7. The dependence of the constant C_s in (6.38) on s is (only) due to the fact that we have used equivalent norms.

6.2 Symmetries and bifurcation from a simple eigenvalue

The set S in Lemma 6.5 has at least the 4 elements (0,0), (1,0), $(\ell_0, \pm m_0)$, and consequently the kernel of L has dimension dim $V \ge 4$. In this subsection we use the symmetries of equation (6.14) to reduce the problem to the case of bifurcation from a simple eigenvalue.

First, we observe that the space of functions that are even in x_3 is an invariant set for the map $\mathcal{F}(\omega, \cdot)$.

Lemma 6.8. If (η, β) is even in x_3 , then also $\mathcal{F}(\omega, \eta, \beta)$ is an even function of x_3 .

Proof. Let M = diag(1, 1, -1) be the 3×3 matrix that maps $x = (x_1, x_2, x_3) \mapsto Mx = (x_1, x_2, -x_3)$ for all $x \in \mathbb{R}^3$. Then M is an orthogonal matrix, i.e., $MM^T = M^TM = I$, and Lemma 6.2 applies to M. Now a function η defined on \mathbb{S}^2 is even in x_3 if $\eta(Mx) = \eta(x)$ for all $x \in \mathbb{S}^2$; in the notation of Lemma 6.2, this means that $\eta(x) = \eta(Mx) = h(x)$, i.e., $\eta = h$. Hence, if η, β are even in x_3 , then all the properties of Lemma 6.2 hold with $h = \eta$ and $\psi = \beta$. In particular, $G(\eta)\beta$, $\langle \nabla_{\mathbb{S}^2}\eta, \nabla_{\mathbb{S}^2}\beta \rangle$, $|\nabla_{\mathbb{S}^2}\eta|^2$, H_{η} are all even in x_3 . By (6.18), $\mathcal{M}\eta$, $\mathcal{M}\beta$ are also even in x_3 because $M^T \mathcal{J}M = \mathcal{J}$. Recalling the definition (6.16), (6.17) of \mathcal{F} , the proof is complete.

Lemma 6.9. The spherical harmonics $\varphi_{\ell,m}$ of Lemma 6.4 satisfy

$$\varphi_{\ell,m}(x_1, x_2, -x_3) = (-1)^{\ell-m} \varphi_{\ell,m}(x) \quad \forall x = (x_1, x_2, x_3) \in \mathbb{S}^2, \quad \forall (\ell, m) \in \mathcal{T}.$$

Proof. The ordinary Legendre polynomial $P_{\ell}(t)$ of degree ℓ has parity $P_{\ell}(-t) = (-1)^{\ell} P_{\ell}(t)$. Hence its *m*-th derivative has parity $(-1)^{\ell-m}$, i.e.,

$$P_{\ell}^{(m)}(-t) = (-1)^{\ell-m} P_{\ell}^{(m)}(t), \quad m = 1, \dots, \ell.$$

Then, by (6.22), $\varphi_{\ell,0}$ has parity $(-1)^{\ell}$ as a function of x_3 , and $\varphi_{\ell,m}^{(\text{Re})}, \varphi_{\ell,m}^{(\text{Im})}, m = 1, \ldots, \ell$, have parity $(-1)^{\ell-m}$ in x_3 . Since $(-1)^{\ell+m} = (-1)^{\ell-m}$, by (6.23) we obtain the thesis.

If a function η is even in x_3 , then, by Lemma 6.9, only the spherical harmonics $\varphi_{\ell,m}$ that are even in x_3 appear in its expansion, i.e., $\hat{\eta}_{\ell,m} = 0$ for all $(\ell,m) \in \mathcal{T}$ such that $\ell - m$ is odd, and only coefficients $\hat{\eta}_{\ell,m}$ with $\ell - m$ even can be nonzero.

Now we consider the parity with respect to x_2 , and prove that $\mathcal{F}(\omega, \cdot)$ changes that parity, as it maps any (even, odd) pair into an (odd, even) one.

Lemma 6.10. If η is even in x_2 and β is odd in x_2 , then $\mathcal{F}_1(\omega, \eta, \beta)$ is odd in x_2 and $\mathcal{F}_2(\omega, \eta, \beta)$ is even in x_2 .

Proof. Consider the matrix M = diag(1, -1, 1) that maps $x = (x_1, x_2, x_3)$ into $Mx = (x_1, -x_2, x_3)$ for all $x \in \mathbb{R}^3$. Then M is an orthogonal matrix, i.e., $MM^T = M^TM = I$, and Lemma 6.2 applies to M. Let η, β be defined on \mathbb{S}^2 , and let $h(x) := \eta(Mx), \psi(x) := \beta(Mx)$ for all $x \in \mathbb{S}^2$. Assume that η is even in x_2 and that β is odd in x_2 . Then $h = \eta$ and $\psi = -\beta$. By Lemma 6.2, we deduce that $(1 + \eta), (1 + \eta)^2, |\nabla_{\mathbb{S}^2}\eta|^2, H_{\eta}, |\nabla_{\mathbb{S}^2}\beta|^2$ are even in x_2 , while $G(\eta)\beta$ and $\langle \nabla_{\mathbb{S}^2}\beta, \nabla_{\mathbb{S}^2}\eta \rangle$ are odd in x_2 . Also, $\mathcal{M}\eta$ is odd in x_2 and $\mathcal{M}\beta$ is even in x_2 , because $(\mathcal{M}\eta)(Mx) = \langle \mathcal{J}Mx, (\nabla_{\mathbb{S}^2}\eta)(Mx) \rangle = \langle M^T \mathcal{J}Mx, \nabla_{\mathbb{S}^2}h(x) \rangle$, similarly $(\mathcal{M}\beta)(Mx) = \langle M^T \mathcal{J}Mx, \nabla_{\mathbb{S}^2}\psi(x) \rangle$, and $M^T \mathcal{J}M = -\mathcal{J}$. By the definition (6.16), (6.17) of $\mathcal{F}_1, \mathcal{F}_2$, the proof is complete.

Lemma 6.11. The spherical harmonics $\varphi_{\ell,m}$ of Lemma 6.4 satisfy

$$arphi_{\ell,m}(x_1, -x_2, x_3) = arphi_{\ell,m}(x), \quad m = 0, \dots, \ell, \ arphi_{\ell,m}(x_1, -x_2, x_3) = -arphi_{\ell,m}(x), \quad m = -\ell, \dots, -1,$$

for all $x = (x_1, x_2, x_3) \in \mathbb{S}^2$, all $\ell \in \mathbb{N}_0$.

Proof. For m = 0, the spherical harmonic $\varphi_{\ell,0}$ in (6.22) does not depend on x_2 , therefore it is even in x_2 . For $m = 1, \ldots, \ell$, one has $\overline{w}^m = \overline{w^m}$ for all $w \in \mathbb{C}$, that is, complex conjugation and *m*th power commute in \mathbb{C} , whence

$$\operatorname{Re}\left\{(x_1 - ix_2)^m\right\} = \operatorname{Re}\left\{(x_1 + ix_2)^m\right\}, \quad \operatorname{Im}\left\{(x_1 - ix_2)^m\right\} = -\operatorname{Im}\left\{(x_1 + ix_2)^m\right\}$$

for all $(x_1, x_2) \in \mathbb{R}^2$, and, recalling (6.22),

$$\varphi_{\ell,m}^{(\text{Re})}(x_1, -x_2, x_3) = \varphi_{\ell,m}^{(\text{Re})}(x), \quad \varphi_{\ell,m}^{(\text{Im})}(x_1, -x_2, x_3) = -\varphi_{\ell,m}^{(\text{Im})}(x)$$

for all $x \in \mathbb{S}^2$. By (6.23) we obtain the thesis.

If η is even in x_2 and β is odd in x_2 , then, by Lemma 6.11, their coefficients satisfy $\hat{\eta}_{\ell,m} = 0$ for m < 0 and $\hat{\beta}_{\ell,m} = 0$ for $m \ge 0$, and only coefficients $\hat{\eta}_{\ell,m}$ with $m \ge 0$ and $\hat{\beta}_{\ell,m}$ with m < 0 can be nonzero.

We put together the properties of parity with respect to x_2 and x_3 , and define the two subspaces

$$X := \{ f \in L^2(\mathbb{S}^2, \mathbb{R}) : f = \text{even}(x_2), \text{ even}(x_3) \}, Y := \{ f \in L^2(\mathbb{S}^2, \mathbb{R}) : f = \text{odd}(x_2), \text{ even}(x_3) \}.$$
(6.43)

Hence any $\eta \in X$, $\beta \in Y$ have expansion

$$\eta = \sum_{(\ell,m)\in\mathcal{T}_X} \hat{\eta}_{\ell,m}\varphi_{\ell,m}, \quad \beta = \sum_{(\ell,m)\in\mathcal{T}_Y} \hat{\beta}_{\ell,m}\varphi_{\ell,m}, \tag{6.44}$$
$$\mathcal{T}_X := \{(\ell,m)\in\mathbb{N}_0^2 : 0 \le m \le \ell, \ \ell-m = \operatorname{even}\}, \\\mathcal{T}_Y := \{(\ell,m)\in\mathbb{N}_0\times\mathbb{Z} : -\ell \le m \le -1, \ \ell-m = \operatorname{even}\}.$$

By Lemmas 6.3, 6.8 and 6.10, the domain and codomain of the map \mathcal{F} can be restricted to the subspaces $X \times Y$ and $Y \times X$ respectively, namely

$$\mathcal{F}_{\rm res}: \mathbb{R} \times (U \cap (X \times Y)) \to (H^s(\mathbb{S}^2) \times H^{s-\frac{1}{2}}(\mathbb{S}^2)) \cap (Y \times X), \tag{6.45}$$

where the index "res" indicates this restriction. The linearized operator $L_{\text{res}} = \partial_u \mathcal{F}_{\text{res}}(\omega, 0)$ is L restricted to $X \times Y$. The kernel of L_{res} is $V_{\text{res}} := V \cap (X \times Y)$, its complement in $X \times Y$ is $W_{\text{res}} := W \cap (X \times Y)$, the range of L_{res} is contained in $R_{\text{res}} := R \cap (Y \times X)$, whose complement in $Y \times X$ is $Z_{\text{res}} := Z \cap (Y \times X)$. We calculate

$$V_{\rm res} = V \cap (X \times Y) = \left\{ \begin{pmatrix} \eta \\ \beta \end{pmatrix} = \sum_{(\ell,m) \in S_{\rm res}} \lambda_{\ell,m} \begin{pmatrix} \ell \varphi_{\ell,m} \\ -\omega m \varphi_{\ell,-m} \end{pmatrix} : \lambda_{\ell,m} \in \mathbb{R} \right\},\tag{6.46}$$

$$S_{\text{res}} := \{ (\ell, m) \in S : \ell \ge 1, \ m \ge 0, \ \ell - m = \text{even} \},$$
(6.47)

and we note that among the 4 elements $(0,0), (1,0), (\ell_0, \pm m_0)$ of S listed in Lemma 6.5, only (ℓ_0, m_0) belongs to S_{res} . Hence, if S contains only those 4 elements, then V_{res} is a 1-dimensional space, and ω in (6.29) is a simple eigenvalue of L_{res} . Now we check that Lemma 6.6 also holds on the restricted spaces $W_{\text{res}}^s := W^s \cap (X \times Y), R_{\text{res}}^s := R^s \cap (Y \times X).$

Lemma 6.12. The map $L_{\text{res}}|_{W^s_{\text{res}}}: W^s_{\text{res}} \to R^s_{\text{res}}$ is invertible, with bounded inverse satisfying estimate (6.38) for all $(f,g) \in W^s_{\text{res}}$.

Proof. Let $(f,g) \in \mathbb{R}^s_{\text{res}} = \mathbb{R}^s \cap (Y \times X)$. By Lemma 6.6, we already know that there exists a unique $(\eta, \beta) = (L|_{W^s})^{-1}(f,g) \in W^s$ such that $L(\eta, \beta) = (f,g)$, with inequality (6.38). We only have to prove that $(\eta, \beta) \in X \times Y$. The coefficients of (η, β) are determined by those of (f,g) by explicit formulas: they are given by (6.33) for $(\ell, m) \in \mathcal{T} \setminus S$, by (6.40) for $(\ell, m) \in S$ with $\ell \geq 1$, and by (6.34) with $\hat{\beta}_{0,0} = 0$ for $(\ell, m) = (0, 0)$. If $(\ell, m) \in \mathcal{T}$ and $\ell - m$ is an odd integer, then $\hat{f}_{\ell,-m} = \hat{g}_{\ell,m} = 0$ because both f and g are even in x_3 , and hence $\hat{\eta}_{\ell,m} = \hat{\beta}_{\ell,-m} = 0$ from the explicit formulas just mentioned. This implies that both η and β are even in x_3 . If $(\ell, m) \in \mathcal{T}$

with m < 0, then $\hat{g}_{\ell,m} = 0$ because g is even in x_2 and $\hat{f}_{\ell,-m} = 0$ because -m > 0 and f is odd in x_2 . Then, again from the explicit formulas, $\hat{\eta}_{\ell,m} = \hat{\beta}_{\ell,-m} = 0$ for m < 0. This implies that η is even in x_2 . Moreover, for m = 0, one has $\hat{f}_{\ell,0} = 0$ because f is odd in x_2 , and therefore, from the explicit formulas, $\hat{\beta}_{\ell,0} = 0$. Hence β is odd in x_2 .

To obtain the bifurcation from a simple eigenvalue, it only remains to check the following transversality property. Recall that ω is given by (6.29), and it is nonzero.

Lemma 6.13. Let $(\ell, m) \in S$, with $\ell \geq 1$ and $1 \leq |m| \leq \ell$. Let $\eta = \ell \varphi_{\ell,m}$ and $\beta = -\omega m \varphi_{\ell,-m}$. Then the pair $(\mathcal{M}\eta, \mathcal{M}\beta)$ does not belong to R.

Proof. By Lemma 6.4, one has $\mathcal{M}\eta = -\ell m \varphi_{\ell,-m}$ and $\mathcal{M}\beta = -\omega m^2 \varphi_{\ell,m}$. Hence, by (6.36), the pair $(\mathcal{M}\eta, \mathcal{M}\beta)$ belongs to R if and only if

$$\begin{pmatrix} -\ell m \varphi_{\ell,-m} \\ -\omega m^2 \varphi_{\ell,m} \end{pmatrix} = \lambda \begin{pmatrix} \varphi_{\ell,-m} \\ -\omega m \ell^{-1} \varphi_{\ell,m} \end{pmatrix} \quad \exists \lambda \in \mathbb{R}$$

and, since $\omega \neq 0$, this is possible only for m = 0.

By the classical results of bifurcation from a simple eigenvalue, we obtain the following result.

Theorem 6.14. Let $\sigma_0 > 0$. Let ℓ_0, m_0 be integers with $\ell_0 \ge 2, 1 \le m_0 \le \ell_0$ and $\ell_0 - m_0$ even. Assume that the Diophantine equation in the unknowns (ℓ, m)

$$m_0^2(\ell+2)(\ell-1)\ell = (\ell_0+2)(\ell_0-1)\ell_0 m^2$$
(6.48)

has only the solution $(\ell, m) = (\ell_0, m_0)$ in the finite set

$$T = \{(\ell, m) \in \mathbb{Z}^2 : 1 \le \ell \le c_0, \quad 0 \le m \le \ell, \quad \ell - m = even\},\$$

$$c_0 = (\ell_0 + 2)(\ell_0 - 1)\ell_0 m_0^{-2}.$$
 (6.49)

Then the value

$$\omega_* = \sqrt{\sigma_0} \frac{\sqrt{(\ell_0 + 2)(\ell_0 - 1)\ell_0}}{m_0} \tag{6.50}$$

for the angular velocity parameter ω is a bifurcation point. The set of nontrivial solutions of equation $\mathcal{F}(\omega, u) = 0$ near $(\omega_*, 0)$ in $\mathbb{R} \times (V_{\text{res}} \oplus W^s_{\text{res}})$ is a unique analytic curve with parametric representation on the 1-dimensional space V_{res} .

Proof. Equation (6.48) is (6.30), and (6.50) is (6.29). As observed in Lemma 6.5, equation (6.48) has no solution with $\ell > c_0$, and therefore the set S_{res} in (6.47) has only one element, the pair (ℓ_0, m_0) . Recalling (6.20), the mixed second derivative $\partial^2_{\omega u} \mathcal{F}(\omega, 0)$ is the operator $(\eta, \beta) \mapsto (\mathcal{M}\eta, \mathcal{M}\beta)$. Thus, by the analysis above, the thesis follows from a direct application of the classical theory of bifurcation from a simple eigenvalue; see, e.g., [5], Theorem 4.1 in Section 5.4, and [12]. The use of symmetries to obtain a simple eigenvalue is also contained, e.g., in [5], Sections 6.3 and 6.4.

6.3 Arithmetics of simple eigenvalues

Using the prime factor decomposition of integers, it is not difficult to see that there exist both pairs (ℓ_0, m_0) that satisfy the assumptions of Theorem 6.14 and pairs that do not satisfy them. By direct check, we have studied the following few cases of small integers.

Lemma 6.15. (i) For $(\ell_0, m_0) = (2, 2)$, (3, 3), (4, 2), (4, 4), (5, 5), (6, 4), (6, 6), (7, 5), (7, 7), the assumptions of Theorem 6.14 are satisfied, and hence the set S_{res} in (6.47) has only one element, the pair (ℓ_0, m_0) itself.

(ii) For $(\ell_0, m_0) = (3, 1)$, (5, 3), (5, 1), (16, 16), the assumptions of Theorem 6.14 are not satisfied, and the corresponding set S_{res} is, respectively,

$$\begin{split} S_{\rm res} &= \{(3,1),\,(10,6),\,(16,12)\}, \quad S_{\rm res} = \{(5,3),\,(8,6)\},\\ S_{\rm res} &= \{(5,1),\,(8,2),\,(126,120)\}, \quad S_{\rm res} = \{(16,16),\,(10,8)\}. \end{split}$$

Proof. Using the prime factor decomposition, the proof is a bit long but completely elementary. \Box

As the previous lemma shows, it seems hard to guess a simple criterion that determines whether a given pair (ℓ_0, m_0) satisfies the assumptions of Theorem 6.14. Nonetheless, again using prime numbers, we can prove that there are infinitely many pairs (ℓ_0, m_0) that satisfy those assumptions.

Proposition 6.16. (i) For every prime integers $2 < p_1 < p_2 < \cdots < p_n$, $n \ge 1$, given $\ell_0 = p_1 p_2 \cdots p_n$, the pair $(\ell_0, m_0) = (\ell_0, \ell_0)$ satisfies the assumptions of Theorem 6.14.

(ii) For every prime integer $\ell_0 \ge 11$, the pair $(\ell_0, m_0) = (\ell_0, \ell_0 - 2)$ satisfies the assumptions of Theorem 6.14.

(iii) For every prime integer p > 3, given $\ell_0 = 2p$, the pair $(\ell_0, m_0) = (\ell_0, \ell_0)$ satisfies the assumptions of Theorem 6.14.

In particular, there are infinitely many pairs (ℓ_0, m_0) satisfying the assumptions of Theorem 6.14.

Proof. (i) For $m_0 = \ell_0 > 0$, equation (6.48) becomes

$$\ell_0(\ell+2)(\ell-1)\ell = (\ell_0+2)(\ell_0-1)m^2, \tag{6.51}$$

and c_0 in (6.49) becomes $c_0 = (\ell_0^2 + \ell_0 - 2)\ell_0^{-1} < \ell_0 + 1$. Assume that $(\ell, m) \in T$ solves (6.51). Recall that ℓ_0 is the product $p_1 \cdots p_n$, and observe that, for any $i \in \{1, \ldots, n\}$, p_i divides neither $\ell_0 + 2$ nor $\ell_0 - 1$, since $p_1 > 2$. Hence, for all i, the prime p_i must divide m^2 , and therefore m. Thus $m = \kappa \ell_0$ for some integer $\kappa \ge 1$. Since $m \le \ell \le c_0 < \ell_0 + 1$, we immediately have $\kappa = 1$. Hence $\ell_0 = m \le \ell < \ell_0 + 1$, whence $\ell = \ell_0$.

(*ii*) For $m_0 = \ell_0 - 2 > 0$, equation (6.48) becomes

$$(\ell_0 - 2)^2 (\ell + 2)(\ell - 1)\ell = (\ell_0 + 2)(\ell_0 - 1)\ell_0 m^2, \tag{6.52}$$

and c_0 in (6.49) becomes $c_0 = \varphi(\ell_0)\ell_0$, where φ is the function $\varphi(x) = (x+2)(x-1)(x-2)^{-2} = 1+5(x-2)^{-1}+4(x-2)^{-2}$, decreasing in $x \in (2,\infty)$. For $\ell_0 \ge 11$, one has $c_0 = \varphi(\ell_0)\ell_0 \le \varphi(11)\ell_0$, and $\varphi(11) = 130/81 < 2$. Assume that $(\ell, m) \in T$ solves (6.52), and let $\ell_0 \ge 11$ be prime. Then, by (6.52), ℓ_0 must divide one of the factors $\ell - 1$, ℓ , $\ell + 2$. We consider the three cases.

Case one. Assume that ℓ_0 divide ℓ . Hence $\ell = b\ell_0$ for some integer $b \ge 1$. Since $(\ell, m) \in T$, one has $b\ell_0 = \ell \le c_0 \le \varphi(11)\ell_0 < 2\ell_0$, whence b = 1. Then $\ell = \ell_0$, which is the trivial solution.

Case two. Assume that ℓ_0 divides $\ell - 1$. Hence $\ell - 1 = b\ell_0$ for some integer $b \ge 1$. Since $(\ell, m) \in T$, one has $b\ell_0 < b\ell_0 + 1 = \ell \le c_0 \le \varphi(11)\ell_0 < 2\ell_0$, whence b = 1. Then $\ell = \ell_0 + 1$, and (6.52) becomes

$$(\ell_0 - 2)^2 (\ell_0 + 3)(\ell_0 + 1) = (\ell_0 + 2)(\ell_0 - 1)m^2.$$
(6.53)

Now $gcd(\ell_0+2, \ell_0+1) = 1$ and $gcd(\ell_0+2, \ell_0+3) = 1$ (consecutive integers). Also, $gcd(\ell_0+2, \ell_0-2) \in \{1, 2, 4\}$ (their difference is 4), but $\ell_0 \pm 2$, like ℓ_0 , is odd. Hence $gcd(\ell_0 + 2, \ell_0 - 2) = 1$. As a consequence, any divisor of $\ell_0 + 2$ divides the RHS of (6.53) and it does not divide the LHS of (6.53), a contradiction.

Case three. Assume that ℓ_0 divides $\ell + 2$. Hence $\ell + 2 = b\ell_0$ for some integer $b \ge 1$. Since $(\ell, m) \in T$, one has $b\ell_0 - 2 = \ell \le c_0 \le \varphi(11)\ell_0$, whence $b\ell_0 \le \varphi(11)\ell_0 + 2$ and $b \le \varphi(11) + (2/\ell_0) \le \varphi(11) + (2/11) < 2$. Therefore b = 1. Then $\ell = \ell_0 - 2$, and (6.52) becomes

$$(\ell_0 - 2)^3 (\ell_0 - 3) = (\ell_0 + 2)(\ell_0 - 1)m^2.$$
(6.54)

Now since ℓ_0 is odd we have by arguing as above $\gcd(\ell_0-2,\ell_0+2) = 1$. Moreover $\gcd(\ell_0-2,\ell_0-1) = 1$ (consecutive integers). Hence $(\ell_0-2)^3$ divides m^2 , namely $m^2 = b(\ell_0-2)^3$ for some integer $b \ge 1$. Thus $(\ell_0-2)^3 \le b(\ell_0-2)^3 = m^2 \le \ell^2 = (\ell_0-2)^2$, but this is impossible for $\ell_0 > 3$.

(*iii*) As observed above, for $m_0 = \ell_0 > 0$, equation (6.48) becomes (6.51), and c_0 in (6.49) becomes $c_0 = (\ell_0^2 + \ell_0 - 2)\ell_0^{-1} < \ell_0 + 1$. Let $m_0 = \ell_0 = 2p$, with p prime. Then (6.51) becomes

$$p(\ell+2)(\ell-1)\ell = (p+1)(2p-1)m^2.$$
(6.55)

Assume that $(\ell, m) \in T$ solves (6.55). Since p is prime, gcd(p, 2p - 1) = 1, and gcd(p, p + 1) = 1, we have that p divides m^2 and therefore m. Hence $m = \kappa p$ for some integer $\kappa \ge 1$, and, since $m \le \ell \le c_0 < \ell_0 + 1 = 2p + 1$, one has $\kappa \in \{1, 2\}$. If $\kappa = 2$, then $m = 2p = \ell_0$, and $\ell_0 = m \le \ell \le c_0 < \ell_0 + 1$, whence $\ell = \ell_0$. Therefore $(\ell, m) = (\ell_0, m_0)$, which is the trivial solution. It remains to study the case $\kappa = 1$, i.e., m = p. For m = p, (6.55) gives

$$(\ell + 2)(\ell - 1)\ell = (p+1)(2p-1)p.$$
 (6.56)

Since p is prime and p > 3, we have that p divides exactly one of the three factors on the LHS of (6.56). We consider the three cases.

Case one. Assume that p divides ℓ . Hence $\ell \in \{p, 2p\}$ because $\ell \leq c_0 < 2p + 1$. Identity (6.56) with $\ell = p$ gives $p^2 + 1 = 0$, a contradiction; (6.56) with $\ell = 2p$ gives 4 = 1, a contradiction.

Case two. Assume that p divides $\ell - 1$. Hence $\ell - 1$ is an integer multiple of p, and, since $\ell - 1 \leq c_0 - 1 < \ell_0 = 2p$, one has $\ell - 1 = p$. Plugging $\ell = p + 1$ into (6.56) gives p = 4, which is not a prime number, a contradiction.

Case three. Assume that p divides $\ell + 2$. Hence $\ell + 2 \in \{p, 2p\}$ because $\ell + 2 \leq c_0 + 2 < 2p + 3 < 3p$. For $\ell + 2 = p$, (6.56) gives (p+7)(p-1) = 0, a contradiction. For $\ell + 2 = 2p$, (6.56) gives $6p^2 - 21p + 13 = 0$, but this polynomial has no integer root, a contradiction.

7 Appendix

Proof of Lemma 5.5. This is essentially Theorem 4.2.2 of [41], page 221, in the case m = 1, $A = \Delta$, p = q = 2, $\sigma = 0$, $s \in \{0, 1\}$, adapted to the non-isotropic spaces $H^{s,r}(\mathbb{R}^n_+)$. Note that the spaces $H^{s,r}(\mathbb{R}^n_+)$ are already used by Triebel in Step 1 of the proof of Theorem 4.2.2. The proof of our lemma consists in slightly modifying the proof in [41], using $r \in [0, \infty)$ as the higher regularity parameter, and paying attention to the constants in the various inequalities along the proof. Also, the two functions u_0, u_1 in Step 2 in [41], page 222, corresponding to low/high frequency components of the function u, must be treated separately regarding their regularity in x'; in particular, the factor C^r in (5.33) comes from the estimate of the term u_0 .

Proof of Lemma 5.7. (i) Proof of density. Let $s, r \in \mathbb{R}$. Since $1 \le 1 + |\xi'|^2 \le 1 + |\xi|^2$, one has

$$(1+|\xi|^2)^{r_0} \le (1+|\xi'|^2)^r \le (1+|\xi|^2)^{r_1} \quad \forall \xi \in \mathbb{R}^n$$

where $r_0 := \min\{0, r\}$ and $r_1 := \max\{0, r\}$. Hence

$$\|u\|_{H^{s+r_0}(\mathbb{R}^n)} \le \|u\|_{H^{s,r}(\mathbb{R}^n)} \le \|u\|_{H^{s+r_1}(\mathbb{R}^n)}$$
(7.1)

for all $u \in H^{s+r_1}(\mathbb{R}^n)$. By Fourier truncation, it is immediate to prove that $H^{s+r_1}(\mathbb{R}^n)$ is dense in $H^{s,r}(\mathbb{R}^n)$: given $u \in H^{s,r}(\mathbb{R}^n)$, let $\hat{u}_N = \hat{u}$ in the ball $|\xi| < N$, and $\hat{u}_N = 0$ outside that ball. Let u_N be the Fourier anti-transform of \hat{u}_N . Then $u_N \in H^{\sigma}(\mathbb{R}^n)$ for all $\sigma \in \mathbb{R}$, and $||u - u_N||_{H^{s,r}(\mathbb{R}^n)}$ tends to zero as $N \to \infty$. Given $\varepsilon > 0$, fix N such that $||u - u_N||_{H^{s,r}(\mathbb{R}^n)} < \varepsilon/2$. As is known (see, e.g., [41], Theorem 2.3.3, page 48), $\mathcal{S}(\mathbb{R}^n)$ is dense in $H^{\sigma}(\mathbb{R}^n)$ for all $\sigma \in \mathbb{R}$. Hence there exists $\varphi \in \mathcal{S}(\mathbb{R}^n)$ such that $||u_N - \varphi||_{H^{s+r_1}(\mathbb{R}^n)} < \varepsilon/2$. By the second inequality in (7.1) and triangular inequality, one has $||u - \varphi||_{H^{s,r}(\mathbb{R}^n)} < \varepsilon$. This proves that $\mathcal{S}(\mathbb{R}^n)$ is dense in $H^{s,r}(\mathbb{R}^n)$.

On \mathbb{R}^n_+ , let $u \in H^{s,r}(\mathbb{R}^n_+)$, and let $v \in H^{s,r}(\mathbb{R}^n)$ with $v|_{\mathbb{R}^n_+} = u$. Given $\varepsilon > 0$, take $\varphi \in \mathcal{S}(\mathbb{R}^n)$ such that $\|v - \varphi\|_{H^{s,r}(\mathbb{R}^n)} < \varepsilon$, and let $f := \varphi|_{\mathbb{R}^n_+}$. Since $v - \varphi \in H^{s,r}(\mathbb{R}^n)$ and $(v - \varphi)|_{\mathbb{R}^n_+} = u - f$, one has

$$||u - f||_{H^{s,r}(\mathbb{R}^n_+)} = \inf\{||w||_{H^{s,r}(\mathbb{R}^n)} : w \in H^{s,r}(\mathbb{R}^n), \ w|_{\mathbb{R}^n_+} = u - f\} \le ||v - \varphi||_{H^{s,r}(\mathbb{R}^n)} < \varepsilon.$$

This proves that the set $\{f : f = \varphi|_{\mathbb{R}^n_+} \exists \varphi \in \mathcal{S}(\mathbb{R}^n)\}$ is dense in $H^{s,r}(\mathbb{R}^n_+)$.

(*ii*) Proof of (5.34). Let v be a function in the Schwartz class $\mathcal{S}(\mathbb{R}^n)$, and let \hat{v} be its Fourier transform on \mathbb{R}^n . By Fourier inversion formula, $\|v\|_{L^{\infty}(\mathbb{R}^n)} \leq \|\hat{v}\|_{L^1(\mathbb{R}^n)}$. Multiplying and dividing $\hat{v}(\xi)$ by $(1 + |\xi|^2)^{\frac{s}{2}}(1 + |\xi'|^2)^{\frac{r}{2}}$, and applying Hölder's inequality, one has

$$\|\hat{v}\|_{L^{1}(\mathbb{R}^{n})} \leq C_{s,r} \|v\|_{H^{s,r}(\mathbb{R}^{n})}, \quad C^{2}_{s,r} = \int_{\mathbb{R}^{n}} \frac{1}{(1+|\xi|^{2})^{s}(1+|\xi'|^{2})^{r}} d\xi$$

For every fixed $\xi' \in \mathbb{R}^{n-1}$, the change of variable $\xi_n = (1 + |\xi'|^2)^{\frac{1}{2}t}$ gives

$$\int_{\mathbb{R}} \frac{1}{(1+|\xi|^2)^s} d\xi_n = \int_{\mathbb{R}} \frac{1}{(1+|\xi'|^2+\xi_n^2)^s} d\xi_n = (1+|\xi'|^2)^{\frac{1}{2}-s} \int_{\mathbb{R}} \frac{1}{(1+t^2)^s} dt.$$
(7.2)

The last integral converges for 2s > 1 (in particular, for s = 1 its value is π). Hence the integral $C_{s,r}^2$ is finite if $2(r + s - \frac{1}{2}) > n - 1$, i.e., r + s > n/2. The assumption $v \in \mathcal{S}(\mathbb{R}^n)$ is removed by density. This proves (5.34) on \mathbb{R}^n . On \mathbb{R}^n_+ , let $u \in H^{s,r}(\mathbb{R}^n_+)$, and let $v \in H^{s,r}(\mathbb{R}^n)$ with $v|_{\mathbb{R}^n_+} = u$. Then

$$||u||_{L^{\infty}(\mathbb{R}^{n}_{+})} = ||v||_{\mathbb{R}^{n}_{+}}||_{L^{\infty}(\mathbb{R}^{n}_{+})} \le ||v||_{L^{\infty}(\mathbb{R}^{n})} \le C_{s,r}||v||_{H^{s,r}(\mathbb{R}^{n})}.$$

Taking the infimum over all such v gives the second embedding inequality (5.34).

(*iii*) Proof of (5.35), (5.36). Inequality (5.36) is trivial on \mathbb{R}^n , i.e., with \mathbb{R}^n_+ replaced by \mathbb{R}^n . On \mathbb{R}^n_+ , let $u \in H^{s+1,r}(\mathbb{R}^n_+)$. For any $v \in H^{s+1,r}(\mathbb{R}^n)$ with $v|_{\mathbb{R}^n_+} = u$, the distribution $\partial_{x_n} v$ belongs to $H^{s,r}(\mathbb{R}^n)$ and it coincides with the distribution $\partial_{x_n} u$ in \mathbb{R}^n_+ . Hence the set $\{\partial_{x_n} v : v \in H^{s+1,r}(\mathbb{R}^n), v|_{\mathbb{R}^n_+} = u\}$ is a subset of $\{w \in H^{s,r}(\mathbb{R}^n) : w|_{\mathbb{R}^n_+} = \partial_{x_n} u\}$, and, by definition (5.16),

$$\begin{split} \|\partial_{x_n} u\|_{H^{s,r}(\mathbb{R}^n_+)} &= \inf\{\|w\|_{H^{s,r}(\mathbb{R}^n)} : w \in H^{s,r}(\mathbb{R}^n), \, w|_{\mathbb{R}^n_+} = \partial_{x_n} u\} \\ &\leq \inf\{\|\partial_{x_n} v\|_{H^{s,r}(\mathbb{R}^n)} : v \in H^{s+1,r}(\mathbb{R}^n), \, v|_{\mathbb{R}^n_+} = u\} \\ &\leq \inf\{\|v\|_{H^{s+1,r}(\mathbb{R}^n)} : v \in H^{s+1,r}(\mathbb{R}^n), \, v|_{\mathbb{R}^n_+} = u\} = \|u\|_{H^{s+1,r}(\mathbb{R}^n_+)}. \end{split}$$

The proof of (5.35) is similar.

(iv) Proof of (5.37), (5.38), (5.39). Proof of (5.37). On \mathbb{R}^n , one has

$$\|v\|_{H^{0,r}(\mathbb{R}^n)}^2 = \int_{\mathbb{R}} \|v(\cdot, x_n)\|_{H^r(\mathbb{R}^{n-1})}^2 dx_n \quad \forall v \in H^{0,r}(\mathbb{R}^n)$$

by Fubini/Tonelli and Plancherel. On \mathbb{R}^n_+ , given $u \in H^{0,r}(\mathbb{R}^n_+)$, its trivial extension u_0 defined as $u_0 = u$ in \mathbb{R}^n_+ and $u_0 = 0$ in $\mathbb{R}^n \setminus \mathbb{R}^n_+$ belongs to $H^{0,r}(\mathbb{R}^n)$ and it satisfies

$$\|u_0\|_{H^{0,r}(\mathbb{R}^n)}^2 = \int_{\mathbb{R}} \|u_0(\cdot, x_n)\|_{H^r(\mathbb{R}^{n-1})}^2 dx_n = \int_0^\infty \|u(\cdot, x_n)\|_{H^r(\mathbb{R}^{n-1})}^2 dx_n.$$

Hence $||u_0||_{H^{0,r}(\mathbb{R}^n)} \leq ||v||_{H^{0,r}(\mathbb{R}^n)}$ for any other $v \in H^{0,r}(\mathbb{R}^n)$ such that $v|_{\mathbb{R}^n_+} = u$, and therefore u_0 realizes the infimum of definition (5.16). *Proof of* (5.38). On \mathbb{R}^n , one has

$$\|v\|_{H^{1,r}(\mathbb{R}^n)}^2 = \|v\|_{H^{0,r+1}(\mathbb{R}^n)}^2 + \|\partial_{x_n}v\|_{H^{0,r}(\mathbb{R}^n)}^2 \quad \forall v \in H^{1,r}(\mathbb{R}^n)$$
(7.3)

(use Fourier transform and split $1 + |\xi|^2$ as $(1 + |\xi'|^2) + \xi_n^2$). On \mathbb{R}^n_+ , using (5.37) and (7.3), for $u \in H^{1,r}(\mathbb{R}^n_+)$ and $v \in H^{1,r}(\mathbb{R}^n)$ with $v|_{\mathbb{R}^n_+} = u$, one has

$$\begin{aligned} \|u\|_{H^{0,r+1}(\mathbb{R}^{n}_{+})}^{2} + \|\partial_{x_{n}}u\|_{H^{0,r}(\mathbb{R}^{n}_{+})}^{2} \\ &= \int_{0}^{\infty} \|u(\cdot,x_{n})\|_{H^{r+1}(\mathbb{R}^{n-1})}^{2} dx_{n} + \int_{0}^{\infty} \|\partial_{x_{n}}u(\cdot,x_{n})\|_{H^{r}(\mathbb{R}^{n-1})}^{2} dx_{n} \\ &\leq \int_{\mathbb{R}} \|v(\cdot,x_{n})\|_{H^{r+1}(\mathbb{R}^{n-1})}^{2} dx_{n} + \int_{\mathbb{R}} \|\partial_{x_{n}}v(\cdot,x_{n})\|_{H^{r}(\mathbb{R}^{n-1})}^{2} dx_{n} = \|v\|_{H^{1,r}(\mathbb{R}^{n})}^{2}.\end{aligned}$$

Taking the infimum over all $v \in H^{1,r}(\mathbb{R}^n)$ with $v|_{\mathbb{R}^n_+} = u$ gives the second inequality in (5.38). To prove the first inequality in (5.38), let $u \in H^{1,r}(\mathbb{R}^n_+)$, and assume that u is the restriction to \mathbb{R}^n_+ of a continuous function on \mathbb{R}^n . Consider the extension u_R defined by reflection as

$$u_R = u \quad \text{if } x_n \ge 0, \qquad u_R(x', x_n) = u(x', -x_n) \quad \text{if } x_n < 0.$$
 (7.4)

Then $\partial_{x_n} u_R$, as a distribution on \mathbb{R}^n , is $\partial_{x_n} u_R = \partial_{x_n} u$ in \mathbb{R}^n_+ , and $(\partial_{x_n} u_R)(x', x_n) = -\partial_{x_n} u(x', x_n)$ for $x_n < 0$, because, integrating by parts with test functions on \mathbb{R}^n , the two boundary terms at $x_n = 0$ cancel out. Hence $u_R \in H^{1,r}(\mathbb{R}^n)$, and

$$\|u_R\|_{H^{1,r}(\mathbb{R}^n)}^2 = \int_{\mathbb{R}} \|u_R(\cdot, x_n)\|_{H^{r+1}(\mathbb{R}^{n-1})}^2 dx_n + \int_{\mathbb{R}} \|\partial_{x_n} u_R(\cdot, x_n)\|_{H^r(\mathbb{R}^{n-1})}^2 dx_n$$
$$= 2\int_0^\infty \|u(\cdot, x_n)\|_{H^{r+1}(\mathbb{R}^{n-1})}^2 dx_n + 2\int_0^\infty \|\partial_{x_n} u(\cdot, x_n)\|_{H^r(\mathbb{R}^{n-1})}^2 dx_n.$$
(7.5)

Also, $||u||_{H^{1,r}(\mathbb{R}^n_+)} \leq ||u_R||_{H^{1,r}(\mathbb{R}^n)}$ by definition (5.16), and, using (5.37), we obtain the first inequality in (5.38). The continuity assumption for u is removed by density. *Proof of* (5.39). It follows immediately from (5.37), (5.38).

(v) Proof of (5.43). Let v be a function in the Schwartz class $\mathcal{S}(\mathbb{R}^n)$, let \hat{v} be its Fourier transform on \mathbb{R}^n , and let g(x') := v(x', 0) be the restriction of v to the hyperplane $x_n = 0$. Let \hat{g} be the Fourier transform of g on \mathbb{R}^{n-1} . One has

$$\hat{g}(\xi') = \int_{\mathbb{R}} \hat{v}(\xi', \xi_n) \, d\xi_n \quad \forall \xi' \in \mathbb{R}^{n-1}$$
(7.6)

because, by Fourier inversion formula, at any point x = (x', 0) one has

$$g(x') = v(x', 0) = v(x) = \int_{\mathbb{R}^n} \hat{v}(\xi) e^{i2\pi\langle\xi, x\rangle} \, d\xi = \int_{\mathbb{R}^{n-1}} \left(\int_{\mathbb{R}} \hat{v}(\xi', \xi_n) \, d\xi_n \right) e^{i2\pi\langle\xi', x'\rangle} \, d\xi'.$$

By (7.6), multiplying and dividing $\hat{v}(\xi)$ by $(1 + |\xi|^2)^{\frac{s}{2}}$, applying Hölder's inequality and (7.2), for every $\xi' \in \mathbb{R}^{n-1}$ one has

$$\begin{aligned} |\hat{g}(\xi')|^2 &\leq \left(\int_{\mathbb{R}} \frac{1}{(1+|\xi|^2)^s} \, d\xi_n\right) \left(\int_{\mathbb{R}} |\hat{v}(\xi',\xi_n)|^2 (1+|\xi|^2)^s \, d\xi_n\right) \\ &= C_s^2 (1+|\xi'|^2)^{\frac{1}{2}-s} \int_{\mathbb{R}} |\hat{v}(\xi)|^2 (1+|\xi|^2)^s \, d\xi_n, \quad C_s^2 := \int_{\mathbb{R}} \frac{1}{(1+t^2)^s} \, dt \end{aligned}$$

Multiplying $|\hat{g}(\xi')|^2$ by $(1+|\xi'|^2)^{\sigma}$, with $\sigma = r+s-\frac{1}{2}$, and integrating in $d\xi'$ over \mathbb{R}^{n-1} , we obtain $\|g\|_{H^{\sigma}(\mathbb{R}^{n-1})} \leq C_s \|v\|_{H^{s,r}(\mathbb{R}^n)}$. Since $\mathcal{S}(\mathbb{R}^n)$ is dense in $H^{s,r}(\mathbb{R}^n)$, the last inequality allows one to define the trace operator on $H^{s,r}(\mathbb{R}^n)$, with bound (5.41).

On \mathbb{R}^n_+ , consider $s = 1, r \in \mathbb{R}$, let $u \in H^{1,r}(\mathbb{R}^n_+)$, and assume that u is the restriction to \mathbb{R}^n_+ of a continuous function. Consider the extension u_R of u defined by reflection in (7.4). Then $u_R \in H^{1,r}(\mathbb{R}^n)$ and, by (7.5), (5.38), one has $||u||_{H^{1,r}(\mathbb{R}^n_+)} \leq ||u_R||_{H^{1,r}(\mathbb{R}^n)} \leq \sqrt{2}||u||_{H^{1,r}(\mathbb{R}^n_+)}$. We define $Tu := Tu_R$, and we note that this definition gives a bounded linear operator of $H^{1,r}(\mathbb{R}^n_+)$ to $H^{r+\frac{1}{2}}(\mathbb{R}^{n-1})$, with bound (5.43), such that $Tu = u(\cdot, 0)$ if u is continuous.

(vi) Proof of (5.44), (5.45). Proof of (5.44). On \mathbb{R}^n , let $u, v \in H^{1,r}(\mathbb{R}^n)$, $r \ge r_0$. The Fourier transform of the product uv is the convolution $\int_{\mathbb{R}^n} \hat{u}(\xi - \eta)\hat{v}(\eta) d\eta$, and

$$\|uv\|_{H^{1,r}(\mathbb{R}^n)}^2 = \int_{\mathbb{R}^n} \left| \int_{\mathbb{R}^n} \hat{u}(\xi - \eta) \hat{v}(\eta) (1 + |\xi|^2)^{\frac{1}{2}} (1 + |\xi'|^2)^{\frac{r}{2}} \, d\eta \right|^2 d\xi.$$
(7.7)

Now

$$(1+|\xi|^2)^{\frac{1}{2}} \le (1+|\xi-\eta|^2)^{\frac{1}{2}} + (1+|\eta|^2)^{\frac{1}{2}},\tag{7.8}$$

$$(1+|\xi'|^2)^{\frac{r}{2}} \le 4(1+|\xi'-\eta'|^2)^{\frac{r}{2}} + C_r(1+|\eta'|^2)^{\frac{r}{2}} \quad \forall \xi, \eta \in \mathbb{R}^n.$$

$$(7.9)$$

To prove (7.8), consider its square, use triangular inequality $|\xi| \leq |\xi - \eta| + |\eta|$ and its square to bound $1 + |\xi|^2$, and use the fact that $|x| \leq (1 + |x|^2)^{\frac{1}{2}}$ for $x = \eta$ and $x = \xi - \eta$. Inequality (7.9) is also elementary, and it is proved in [8], inequality (B.10) and Lemma B.5, where the slightly stronger version with $|\eta'|^r$ instead of $(1 + |\eta'|^2)^{\frac{r}{2}}$ is proved, and where the constant C_r is written explicitly; C_r is a continuous increasing function of $r \in [0, \infty)$, with $C_r = 1$ for $r \in [0, 1]$ and $C_r \to \infty$ as $r \to \infty$. The product of the LHS of (7.8) and (7.9) is bounded by the product of the corresponding RHS, which is the sum of 4 products. Then the internal integral in (7.7) is bounded by the sum of 4 integrals, and, since $(a_1 + \ldots + a_4)^2 \leq 4(a_1^2 + \ldots + a_4^2)$ for any $a_i \in \mathbb{R}$, one has

$$(7.7) \le 4 \sum_{i,j=1,2} \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} |\hat{u}(\xi - \eta)| |\hat{v}(\eta)| \sigma_i \tau_j \, d\eta \right)^2 d\xi,$$

where σ_1, σ_2 are the two terms in the RHS of (7.8) and τ_1, τ_2 are those in the RHS of (7.9). In the case $\sigma_1 \tau_1$, multiplying and dividing by $\sigma_2 (1 + |\eta'|^2)^{\frac{\tau_0}{2}}$, and applying Hölder's inequality, we get

$$\int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} |\hat{u}(\xi - \eta)| |\hat{v}(\eta)| \sigma_1 \tau_1 \, d\eta \right)^2 d\xi \le 4^2 C_{r_0}^2 \|u\|_{H^{1,r}(\mathbb{R}^n)}^2 \|v\|_{H^{1,r_0}(\mathbb{R}^n)}^2,$$

where

$$C_{r_0}^2 := \int_{\mathbb{R}^n} \frac{1}{(1+|\eta|^2)(1+|\eta'|^2)^{r_0}} \, d\eta = \int_{\mathbb{R}^{n-1}} \frac{\pi}{(1+|\eta'|^2)^{r_0+\frac{1}{2}}} \, d\eta \tag{7.10}$$

is finite for $2(r_0 + \frac{1}{2}) > n - 1$. The second identity in (7.10) holds by (7.2). In the case $\sigma_1 \tau_2$, we multiply and divide by $\sigma_2(1+|\xi'-\eta'|^2)^{\frac{r_0}{2}}$ and, proceeding similarly, we find that the corresponding integral is bounded by the square of $C_{r_0}C_r ||u||_{H^{1,r_0}(\mathbb{R}^n)} ||v||_{H^{1,r}(\mathbb{R}^n)}$, where C_r is the constant in (7.9), and C_{r_0} is in (7.10). In fact, using (7.2) and Hölder's inequality, for every $\xi \in \mathbb{R}^n$ one has

$$\int_{\mathbb{R}^{n}} \frac{1}{(1+|\eta|^{2})(1+|\xi'-\eta'|^{2})^{r_{0}}} d\eta = \int_{\mathbb{R}^{n-1}} \frac{\pi}{(1+|\eta'|^{2})^{\frac{1}{2}}(1+|\xi'-\eta'|^{2})^{r_{0}}} d\eta' \\
\leq \pi \Big(\int_{\mathbb{R}^{n-1}} \frac{1}{(1+|\eta'|^{2})^{\frac{p}{2}}} d\eta' \Big)^{\frac{1}{p}} \Big(\int_{\mathbb{R}^{n-1}} \frac{1}{(1+|\xi'-\eta'|^{2})^{r_{0}q}} d\eta' \Big)^{\frac{1}{q}} = C_{r_{0}}^{2} \quad (7.11)$$

with $p = 2r_0 + 1$, $\frac{1}{p} + \frac{1}{q} = 1$. The cases $\sigma_2 \tau_1$, $\sigma_2 \tau_2$ are analogous. This proves (5.44) with \mathbb{R}^n instead of \mathbb{R}^n_+ . To obtain (5.44), let $u, v \in H^{1,r}(\mathbb{R}^n_+)$, and assume that they are the restriction to \mathbb{R}^n_+ of continuous functions. Consider the extensions u_R, v_R of u, v as in (7.4). Then the product $u_R v_R$ satisfies the estimate (5.44) on \mathbb{R}^n that we have just proved. Thus, by (7.5), (5.38), we deduce that the product uv satisfies (5.44). The continuity assumption is removed by density. *Proof of* (5.45). To prove (5.45), we proceed similarly, but without exceeding r in the powers of η' : for example, in the case $\sigma_1 \tau_1$, we multiply and divide by $\sigma_2(1 + |\eta'|^2)^{\frac{r}{2}}(1 + |\xi' - \eta'|^2)^{\frac{r_0 - r}{2}}$. As above, we use (7.2) and Hölder's inequality, and we choose Hölder's exponents p, q giving the power $r_0 + \frac{1}{2}$ like in (7.10), (7.11).

(vii) Proof of (5.47), (5.49). Proof of (5.47). On \mathbb{R}^n , one has

$$\|gv\|_{H^{r}(\mathbb{R}^{n})} \leq 2\|g\|_{L^{\infty}(\mathbb{R}^{n})} \|v\|_{H^{r}(\mathbb{R}^{n})} + C_{r}\|g\|_{W^{b,\infty}(\mathbb{R}^{n})} \|v\|_{L^{2}(\mathbb{R}^{n})}$$
(7.12)

for all $v \in H^r(\mathbb{R}^n)$, all $g \in W^{b,\infty}(\mathbb{R}^n)$, where $r \ge 0$ is real, b is the smallest integer such that $b \ge r$, and C_r is an increasing function of r. Estimate (7.12) is proved, e.g., in Lemma B.2 of [8]. Then (5.47) follows by using identity (5.37) and applying estimate (7.12) (with \mathbb{R}^{n-1} instead of \mathbb{R}^n) to the product $f(\cdot, x_n)u(\cdot, x_n)$ in the integral, and Hölder's inequality (or triangular inequality for the $L^2(0,\infty)$ norm). Proof of (5.49). Use the first inequality in (5.38), estimate separately each product fu, $(\partial_{x_n} f)u$ and $f(\partial_{x_n} u)$ by (5.47), then use the second inequality in (5.38).

(viii) Proof of (5.51). On \mathbb{R}^n , given a diffeomorphism f of \mathbb{R}^n of the form f(x) = a + Ax + g(x), one has

$$\|u \circ f\|_{H^{r}(\mathbb{R}^{n})} \leq C_{r,f}(\|u\|_{H^{r}(\mathbb{R}^{n})} + \|g\|_{H^{r+1+r_{0}}(\mathbb{R}^{n})}\|u\|_{L^{2}(\mathbb{R}^{n})})$$
(7.13)

for all $u \in H^r(\mathbb{R}^n)$, all real $r \geq 0$, where $C_{r,f}$ depends on $r, |A|, ||g||_{H^{1+r_0}(\mathbb{R}^n)}$ and $r_0 > n/2$. Estimate (7.13) is classical, and it can be proved in this way. For r = 0, use the change of variable f(x) = y in the integral giving the $L^2(\mathbb{R}^d)$ norm of $u \circ f$, and the $L^{\infty}(\mathbb{R}^n)$ norm of the Jacobian determinant det $D(f^{-1})$ of the inverse transformation, to get $||u \circ f||_{L^2(\mathbb{R}^n)} \leq C_0 ||u||_{L^2(\mathbb{R}^n)}$ for some C_0 depending on f. For r = 1, we note that

$$\nabla(u \circ f)(x) = [Df(x)]^T (\nabla u) \circ f(x) = A^T (\nabla u) \circ f(x) + [Dg(x)]^T (\nabla u) \circ f(x)$$
(7.14)

for all $x \in \mathbb{R}^n$, then we proceed like in the r = 0 case to estimate the $L^2(\mathbb{R}^d)$ norm of (7.14), and we get $||u \circ f||_{H^1(\mathbb{R}^n)} \leq C_1 ||u||_{H^1(\mathbb{R}^n)}$ for some $C_1 \geq C_0$ depending on f. Then, by the classical interpolation theorems for linear operators, $||u \circ f||_{H^r(\mathbb{R}^n)} \leq C_1 ||u||_{H^r(\mathbb{R}^n)}$ for all real $r \in [0, 1]$. Assume that (7.13) holds for some real $r \geq 0$. The norm $||u \circ f||_{H^{r+1}(\mathbb{R}^n)}$ is controlled by $||u \circ f||_{H^r(\mathbb{R}^n)} + ||\nabla(u \circ f)||_{H^r(\mathbb{R}^n)}$, and we use identity (7.14) and triangular inequality. To estimate the product $(Dg)^T(\nabla u) \circ f$, we use the product estimate

$$\|vw\|_{H^{r}(\mathbb{R}^{n})} \leq C_{r_{0}}\|v\|_{H^{r_{0}}(\mathbb{R}^{n})}\|w\|_{H^{r}(\mathbb{R}^{n})} + C_{r}\|v\|_{H^{r+r_{0}}(\mathbb{R}^{n})}\|w\|_{L^{2}(\mathbb{R}^{n})}$$
(7.15)

with $v = (Dg)^T$ and $w = (\nabla u) \circ f$. Estimate (7.15) is proved similarly as (5.44), (5.45), using (7.9) (in fact, the proof of (7.15) is simpler because no coordinate has a special role). By the interpolation inequality of $H^r(\mathbb{R}^n)$ spaces, the product $\|g\|_{H^{r+1+r_0}} \|u\|_{H^1}$ is bounded by $\|g\|_{H^{r+2+r_0}} \|u\|_{H^0}$ $+\|g\|_{H^{r_0+1}} \|u\|_{H^{r+1}}$, and (7.13) follows by induction. On \mathbb{R}^n_+ , estimate (5.51) is immediately deduced from (5.37), (5.38), applying (7.13) with \mathbb{R}^{n-1} instead of \mathbb{R}^n ; for s = 1, we also use the fact that $\partial_{x_n} \{u(f(x'), x_n)\} = (\partial_{x_n} u)(f(x'), x_n)$.

Proof of Lemma 5.8. Proof of (5.52). Recall the definition (5.18) of the norm, apply the standard interpolation inequality $||f||_{H^s(\mathbb{R}^d)} \leq ||f||_{H^{s_0}(\mathbb{R}^d)}^{1-\vartheta} ||f||_{H^{s_1}(\mathbb{R}^d)}^{\vartheta}$ of the spaces $H^s(\mathbb{R}^d)$ to each function $f = (u\psi_j) \circ g_j(\cdot, 0)$, then use Hölder's inequality

$$\sum_{j=1}^{N} a_j b_j \le \left(\sum_{j=1}^{N} a_j^p\right)^{\frac{1}{p}} \left(\sum_{j=1}^{N} b_j^q\right)^{\frac{1}{q}}$$

with $p = 1/(1 - \vartheta), q = 1/\vartheta, a_j = ||(u\psi_j) \circ \mathbf{g}_j||_{H^{s_0}(\mathbb{R}^d)}^{1-\vartheta}, b_j = ||(u\psi_j) \circ \mathbf{g}_j||_{H^{s_1}(\mathbb{R}^d)}^{\vartheta}.$ *Proof of* (5.53). By (5.19),

$$(uv\psi_j)\circ\mathsf{g}_j(\cdot,0)=\sum_{\ell\in\mathcal{A}_j}\tilde{u}_{\ell j}\,\tilde{v}_j,\quad \tilde{u}_{\ell j}:=(u\psi_\ell)\circ\mathsf{g}_j(\cdot,0),\quad \tilde{v}_j:=(v\psi_j)\circ\mathsf{g}_j(\cdot,0).$$

On \mathbb{R}^d , one has the well-known product estimate

$$\|\tilde{u}_{\ell j}\tilde{v}_{j}\|_{H^{s}(\mathbb{R}^{d})} \leq C_{s_{0}}\|\tilde{u}_{\ell j}\|_{H^{s_{0}}(\mathbb{R}^{d})}\|\tilde{v}_{j}\|_{H^{s}(\mathbb{R}^{d})} + (\chi_{s \geq s_{0}})C_{s}\|\tilde{u}_{\ell j}\|_{H^{s}(\mathbb{R}^{d})}\|\tilde{v}_{j}\|_{H^{s_{0}}(\mathbb{R}^{d})},$$
(7.16)

which can be proved similarly as (5.46) (in fact, the proof of (7.16) is slightly easier). Since $\mathbf{g}_j = \mathbf{g}_\ell \circ \mathbf{f}_\ell \circ \mathbf{g}_j$, one has

$$\tilde{u}_{\ell j} = (u\psi_{\ell}) \circ \mathbf{g}_{\ell} \circ \mathbf{f}_{\ell} \circ \mathbf{g}_{j}(\cdot, 0) = (u\psi_{\ell}) \circ \mathbf{g}_{\ell}(\tilde{\mathsf{T}}_{\ell j}(\cdot), 0) = \tilde{u}_{\ell} \circ \tilde{\mathsf{T}}_{\ell j},$$

where $\tilde{u}_{\ell} := (u\psi_{\ell}) \circ g_{\ell}(\cdot, 0)$ and $\tilde{T}_{\ell j}$ is defined in (5.31). By Lemma 5.4 and the composition estimate (5.51), one has

$$\|\tilde{u}_{\ell j}\|_{H^{s_0}(\mathbb{R}^d)} \le C_{s_0} \|\tilde{u}_{\ell}\|_{H^{s_0}(\mathbb{R}^d)}, \quad \|\tilde{u}_{\ell j}\|_{H^s(\mathbb{R}^d)} \le C_s \|\tilde{u}_{\ell}\|_{H^s(\mathbb{R}^d)}.$$
(7.17)

Using (7.17) into (7.16) and taking the sum over j, ℓ gives (5.53).

Proof of (5.54). For m = 2, (5.54) is (5.53) with v = u; (5.54) with m + 1 instead of m is deduced from (5.54) by applying (5.53) with $v = u^m$.

Proof of (5.55). Let $u \in H^{s+1}(\mathbb{S}^d)$. Denote $u_0 := \mathcal{E}_0 u$ its 0-homogeneous extension to $\mathbb{R}^n \setminus \{0\}$, n = d + 1, so that $\nabla_{\mathbb{S}^d} u = \nabla u_0$ on \mathbb{S}^d . By (5.18), we have to estimate the sum over $j = 1, \ldots, N$ of $\|((\nabla u_0)\psi_j) \circ \mathbf{g}_j(\cdot, 0)\|_{H^s(\mathbb{R}^d)}$. The function $((\nabla u_0)\psi_j) \circ \mathbf{g}_j(\cdot, 0)$ is the result of the composition $T(\mathbf{G}_j(\Psi_j(\nabla u_0)))$, where T is the trace operator $f \mapsto f(\cdot, 0)$, \mathbf{G}_j is the composition operator $f \mapsto f \circ \mathbf{g}_j$, and Ψ_j is the multiplication operator $f \mapsto f\psi_j$. The composition of these operators with the gradient operator satisfies the following identities. First, one has $\Psi_j(\nabla f) = \nabla(\Psi_j f) - (\nabla \psi_j) f$. Second, $\mathbf{G}_j(\nabla f) = (D\mathbf{g}_j)^{-T} \nabla(\mathbf{G}_j f)$. Third, $T(\nabla f) = (\nabla_{y'}(Tf), T(\partial_{y_n} f))$. Applying these identities, one has that $((\nabla u_0)\psi_j) \circ \mathbf{g}_j(\cdot, 0) = S_{1,j} + S_{2,j} + S_{3,j}$, where $S_{1,j}$ is the product of the matrix $T((D\mathbf{g}_j)^{-T})$ with the vector $(\nabla_{y'}\{T(\mathbf{G}_j(\Psi_j u_0))\}, 0), S_{2,j}$ is the product of the same matrix $T((D\mathbf{g}_j)^{-T})$ with the vector $(0, T(\partial_{y_n} \{ \mathbf{G}_j(\Psi_j u_0) \}))$, and $S_{3,j}$ is $-T(\mathbf{G}_j\{(\nabla \psi_j) u_0\})$. By (7.12),

$$\begin{split} \|S_{1,j}\|_{H^{s}(\mathbb{R}^{d})} &\leq C_{0} \|\nabla_{y'} \{ T(\mathsf{G}_{j}(\Psi_{j}u_{0})) \} \|_{H^{s}(\mathbb{R}^{d})} + C_{s} \|\nabla_{y'} \{ T(\mathsf{G}_{j}(\Psi_{j}u_{0})) \} \|_{L^{2}(\mathbb{R}^{d})} \\ &\leq C_{0} \|T(\mathsf{G}_{j}(\Psi_{j}u_{0}))\|_{H^{s+1}(\mathbb{R}^{d})} + C_{s} \|T(\mathsf{G}_{j}(\Psi_{j}u_{0}))\|_{H^{1}(\mathbb{R}^{d})}, \end{split}$$

and therefore, recalling the definition (5.18) of the norm on \mathbb{S}^d , the sum over $j = 1, \ldots, N$ of the terms $S_{1,j}$ is bounded by $C_0 ||u||_{H^{s+1}(\mathbb{S}^d)} + C_s ||u||_{H^1(\mathbb{S}^d)}$. Concerning $S_{2,j}$, we observe that $\partial_{y_n}(\mathsf{G}_j u_0) = 0$ because, by (5.21), (5.22), the vector $\partial_{y_n} \mathsf{g}_j(y)$ is parallel to $\mathsf{g}_j(y)$, and $\langle x, \nabla u_0(x) \rangle = 0$, as u_0 is a 0-homogeneous function. Hence $T(\partial_{y_n}\{\mathsf{G}_j(\Psi_j u_0)\}) = T(\partial_{y_n}(\mathsf{G}_j\psi_j))T(\mathsf{G}_j u_0)$. We proceed similarly as in the proof of (5.53): we insert $1 = \sum_{\ell \in \mathcal{A}_j} \psi_\ell$, and write $\mathsf{g}_j = \mathsf{g}_\ell \circ \mathsf{f}_\ell \circ \mathsf{g}_j$, so that $\mathsf{g}_j(\cdot, 0) = \mathsf{g}_\ell(\tilde{\mathsf{T}}_{\ell_j}(\cdot), 0)$, with $\tilde{\mathsf{T}}_{\ell_j}$ defined in (5.31). Then we use the product estimate (7.12), Lemma 5.4 and the composition estimate (5.51), and we obtain that the sum over $j = 1, \ldots, N$ of the terms $||S_{2,j}||_{H^s(\mathbb{R}^d)}$ is bounded by the sum over j, ℓ with $\ell \in \mathcal{A}_j$ of $C_s ||T(\mathsf{G}_\ell(\Psi_\ell u_0))||_{H^s(\mathbb{R}^d)}$. The terms $S_{3,j}$ are estimated similarly as $S_{2,j}$. Thus we obtain

$$\|\nabla_{\mathbb{S}^d} u\|_{H^s(\mathbb{S}^d)} \le C_0 \|u\|_{H^{s+1}(\mathbb{S}^d)} + C_s \|u\|_{H^1(\mathbb{S}^d)} + C_s \|u\|_{H^s(\mathbb{S}^d)}.$$

To absorb the last term, we use (5.52) with $s_0 = 0$, $s_1 = s + 1$, and proceed like in (5.103).

Proof of (5.56). On \mathbb{R}^d one has the embedding inequality $||f||_{L^{\infty}(\mathbb{R}^d)} \leq C_{s_0} ||f||_{H^{s_0}(\mathbb{R}^d)}$. On \mathbb{S}^d , by (5.19), one has $u = \sum_{j=1}^N u\psi_j$, and

$$\begin{split} \|u\|_{L^{\infty}(\mathbb{S}^{d})} &\leq \sum_{j=1}^{N} \|u\psi_{j}\|_{L^{\infty}(\mathbb{S}^{d})} = \sum_{j=1}^{N} \|(u\psi_{j}) \circ \mathbf{g}_{j}(\cdot, 0)\|_{L^{\infty}(\mathbb{R}^{d})} \\ &\leq C_{s_{0}} \sum_{j=1}^{N} \|(u\psi_{j}) \circ \mathbf{g}_{j}(\cdot, 0)\|_{H^{s_{0}}(\mathbb{R}^{d})} = C_{s_{0}} \|u\|_{H^{s_{0}}(\mathbb{S}^{d})}, \end{split}$$

because $K_j \cap \mathbb{S}^2 = \{ g_j(y', 0) : (y', 0) \in A_0 \}$ and $\psi_j = 0$ outside $K_j, \psi_j \circ g_j = 0$ outside A_0 . \Box

References

- [1] T. Alazard, N. Burq, C. Zuily, *Cauchy theory for the water waves system in an analytic framework*, to appear in Tokyo Journal of Mathematics.
- [2] T. Alazard, J.-M. Delort, Sobolev estimates for two dimensional gravity water waves, Asterisque, 374, viii + 241 (2015).
- [3] T. Alazard, G. Metivier, Paralinearization of the Dirichlet to Neumann operator, and regularity of the three dimensional water waves, Comm. Partial Differential Equations 34, no. 10-12, 1632-1704, (2009).
- [4] A.D. Alexandrov, A characteristic property of spheres, Ann. Mat. Pura Appl. (4) 58 (1962), 303-315.
- [5] A. Ambrosetti, G. Prodi, A Primer of Nonlinear Analysis, Cambridge University Press, Cambridge (1993).
- [6] L. Ambrosio, N. Fusco, D. Pallara, Functions of bounded variation and free discontinuity problems, Oxford Mathematical Monographs, (2000)
- [7] K. Atkinson, W. Han, Spherical Harmonics and Approximations on the Unit Sphere: An Introduction, Lecture Notes in Mathematics 2044, Springer-Verlag Berlin Heidelberg (2012).
- [8] P. Baldi, E. Haus, Size of data in implicit function problems and singular perturbations for nonlinear Schrödinger systems, Ann. Inst. H. Poincaré Anal. Non Linéaire 40, 1051-1091 (2023).
- M. Berti, L. Franzoi, A. Maspero, Traveling quasi-periodic water waves with constant vorticity, Arch. Ration. Mech. Anal. 240 (2021), 99-202.
- [10] M. Berti, A. Maspero, P. Ventura, On the analyticity of the Dirichlet-Neumann operator and Stokes waves, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. 33 no. 3, 611-650 (2022).

- [11] K. Beyer, M. Günther, On the Cauchy Problem for a Capillary Drop. Part I: Irrotational Motions, Math. Meth. Appl. Sci. 21, 1149-1183 (1998).
- [12] B. Buffoni, J. Toland. Analytic Theory of Global Bifurcation, Princeton University Press (2003).
- [13] F. Cagnetti, M.G. Mora, M. Morini, A second order minimality condition for the Mumford-Shah functional, Calc. Var. 33, 37-74, (2008).
- [14] A.P. Calderon, Cauchy integrals on Lipschitz curves and related operators, Proc. Nat. Acad. Sci. USA 75 (1977), 1324-1327.
- [15] G. Ciraolo, F. Maggi, On the shape of compact hypersurfaces with almost-constant mean curvature. Comm. Pure Appl. Math. 70 (2017), 665–716.
- [16] R. Coifman, Y. Meyer, Nonlinear harmonic analysis and analytic dependence, Proc. Symp. Pure Math., Vol. 43, 71-78, (1985).
- [17] D. Coutand, S. Shkoller, Well-posedness of the free-surface incompressible Euler equations with or without surface tension, J. Amer. Math. Soc. 20, 829–930 (2007).
- [18] W. Craig, D. Nicholls, Travelling two and three dimensional capillary gravity water waves, SIAM J. Math. Anal. 32, no. 2, 323-359, (2000).
- [19] W. Craig, U. Schanz, C. Sulem, The modulational regime of three-dimensional water waves and the Davey-Stewartson system, Annales de l'I.H.P. Analyse non lineaire 14, no.5, 615–667, (1997).
- [20] W. Craig, C. Sulem, Numerical simulation of gravity waves, J. Comput. Phys., 108(1): 73–83, 1993.
- [21] M. G. Crandall, P. H. Rabinowitz, Bifurcation from simple eigenvalues, J. Funct. Anal. 8 (1971), 321–340.
- [22] L.C. Evans, *Partial differential equations*, American Mathematical Society, (2010).
- [23] N. Fusco, D.A. La Manna, A remark on a conjecture on the symmetric Gaussian Problem, Proceedings of the Edinburgh Mathematical Society 67, (2023), 643-661.
- [24] M. Groves, D. Nilsson, S. Pasquali, E. Wahlén, Analytical Study of a generalised Dirichlet-Neumann operator and application to three-dimensional water waves on Beltrami flows, preprint on arXiv:2307.01702.
- [25] M.D. Groves, S.-M. Sun, Fully localised solitary-wve solutions of the three-dimensional gravitycapillary water-wave problem, Arch. Ration. Mech. Anal. 188 (2008), 1, 1-91.
- [26] S. Haziot, V. Hur, W. Strauss, J. Toland, E. Wahlén, S. Walsh, M. Wheeler, Traveling water waves – The ebb and flow of two centuries, Quart. Appl. Math. 80 (2022), 317-401.
- [27] A. Henrot, M. Pierre, Shape Variation and Optimization: A Geometrical Analysis. EMS Tracts in Mathematics, 28. EMS, 2018.
- [28] Y. Huang, A. Karakhanyan, The Dirichlet-Neumann Operator for Taylor's Cone, preprint available at https://arxiv.org/pdf/2402.07005.
- [29] V. Julin, D.A. La Manna, A Priori Estimates for the Motion of Charged Liquid Drop: A Dynamic Approach via Free Boundary Euler Equations, J. Math. Fluid Mech. 26: 48,(2024).
- [30] D. Lannes, The Water Waves Problem: Mathematical Analysis and Asymptotics, American Mathematical Soc., (2013).
- [31] G. Leoni, I. Tice, Traveling wave solutions to the free boundary incompressible Navier-Stokes equations, Comm. Pure Appl. Math. 76, 2474-2576 (2023).
- [32] T. Levi-Civita, Determinazione rigorosa della onde irrotazionali periodiche in acqua profonda, Rend. Accad. Lincei 33, pp. 141-150, (1924).
- [33] J.L. Lions, E. Magenes, Non-homogeneous boundary value problems and applications, Die Grundlehren d. math. Wissenschaften. Band 183, 1972.
- [34] F. Maggi, Sets of Finite Perimeter and Geometric Variational Problems, Cambridge University Press, 2012.
- [35] G. Moon, Y. Wu. Global Bifurcation of Steady Surface Capillary Waves on a 2D Droplet, preprint available at https://arxiv.org/abs/2407.16794, (2024).
- [36] A. Nekrasov, On steady waves, Izv. Ivanovo-Voznesenk. Politekhn. 3, 1921.
- [37] Lord Rayleigh, Proc. Royal Soc. London 29 (1879), 71, The theory of Sound II. MacMillan Co., London (1896).

- [38] C. Shao, Longtime Dynamics of Irrotational Spherical Water Drops: Initial Notes, preprint available at https://arxiv.org/pdf/2301.00115.
- [39] C. Shao. On the Cauchy Problem of Spherical Capillary Water Waves. preprint available at https://arxiv.org/pdf/2310.07113.
- [40] J. Shatah, C. Zeng, Geometry and a priori estimates for free boundary problems of the Euler's equation, Comm. Pure Appl. Math. 61 (5), 698-744, 2008.
- [41] H. Triebel, *Theory of Function Spaces*, Birkhäuser 2010, Springer Basel AG, reprint of the 1983 edition.
- [42] E. Wahlén, Steady periodic capillary waves with vorticity, Ark. Mat. 44 (2006), 367-387.
- [43] E. Wahlén, A Hamiltonian formulation of water waves with constant vorticity, Letters in Mathematical Physics, 79 no.3, 303-315(2007).
- [44] S. Walsh, Steady stratified periodic gravity waves with surface tension II: Global bifurcation, Discr. Cont. Dyn. Syst. 34 (2014), 3287-3315.
- [45] V.E. Zakharov, Stability of periodic waves of finite amplitude on the surface of a deep fluid, J.Appl. Mech. Tech. Phys. 9, 190-194 (1968).

Pietro Baldi

Dipartimento di Matematica e Applicazioni "R. Caccioppoli" University of Naples Federico II Via Cintia, Monte Sant'Angelo, 80126 Naples, Italy pietro.baldi@unina.it

Vesa Julin

Department of Mathematics and Statistics University of Jyväskylä P.O. Box 35, 40014 Jyväskylä, Finland vesa.julin@jyu.fi

Domenico Angelo La Manna

Dipartimento di Matematica e Applicazioni "R. Caccioppoli" University of Naples Federico II Via Cintia, Monte Sant'Angelo, 80126 Naples, Italy domenicoangelo.lamanna@unina.it