

Hybrid Surrogate Models: Circumventing Gibbs Phenomenon for Partial Differential Equations with Finite Shock-Type Discontinuities

Juan-Esteban Suarez Cardona^[0000-0002-1990-2413] and
Shashank Reddy^[0009-0006-7037-6312] and
Michael Hecht^[0000-0001-9214-8253]

Abstract We introduce the concept of *Hybrid Surrogate Models* (HSMs) – combining multivariate polynomials with Heavyside functions – as approximates of functions with finitely many jump discontinuities. We exploit the HSMs for formulating a variational optimization approach, solving non-regular partial differential equations (PDEs) with non-continuous shock-type solutions. The HSM technique simultaneously obtains a parametrization of the position and the height of the shocks as well as the solution of the PDE. We show that the HSM technique circumvents the notorious Gibbs phenomenon, which limits the accuracy that classic numerical methods reach. Numerical experiments, addressing linear and non-linearly propagating shocks, demonstrate the strong approximation power of the HSM technique.

Key words: Hybrid Surrogates, Non-Smooth PDEs, Gibbs Phenomenon, Variational Formulation

1 Introduction

Solving partial differential equations (PDEs) with discontinuous solutions is crucial in various fields, including droplet physics [17], shock formation in hyperbolic systems [14], and singularity formation in magneto-hydrodynamics [4]. The challenge lies in the inherent difficulty of numerically approximating non-continuous functions, as smooth approximations struggle to resolve discontinuities, due to the notorious Gibbs phenomenon [10]. Gibbs phenomenon manifests as spurious oscillatory behavior in various approximations, such as Fourier series [18], polynomials [3], and even neural networks [9]. Classic numerical methods, including finite differences [12], finite elements [6], finite volumes [7], and particle methods [13], incorporate

Juan-Esteban Suarez Cardona
Center for Advanced Systems Understanding (CASUS), Görlitz, Germany, e-mail: j.suarez-cardona@hzdr.de

Shashank Reddy
Institute of Information Technology, Bengalore, India e-mail: Shashank.Reddy@iiitb.ac.in

Michael Hecht
Center for Advanced Systems Understanding (CASUS), Görlitz, Germany
University of Wrocław, Mathematical Institute, Wrocław, Poland e-mail: m.hecht@hzdr.de

smooth approximations within their formulations, thereby inheriting the limitations associated with the Gibbs phenomenon. State-of-the-art methods, such as Discontinuous Galerkin methods (DGM) [5] with non-linear limiters or unfitted DGM [15], have been proven to be effective in controlling spurious oscillations associated with high-order finite element schemes. Recent advancements in machine learning demonstrate promising results by leveraging the flexibility of neural networks (NNs), approximating discontinuous PDE solutions [16].

Complementary to the aforementioned methods, here, we present the *Hybrid Surrogate Models* (HSMs) as a novel approach, circumventing the Gibbs phenomenon. The HSMs inherently incorporate discontinuities into the surrogate model, combining the Heaviside function with multivariate polynomials. Remarkably, this eliminates the need for limiters or adaptive discretizations for suppressing the Gibbs phenomenon.

This work is a follow-up of the *Sobolev PINNs* (SC-PINNs) [1] and the *Polynomial Surrogate Models* (PSMs) [2], exploiting the therein delivered technique of *Sobolev cubatures*.

1.1 Basic Definitions and Notation

We denote with $\mathcal{B} = \Omega \times (0, T)$ the product space of the d -dimensional open hypercube $\Omega = (-1, 1)^d \subseteq \mathbb{R}^d$, $d \in \mathbb{N}$ and a temporal domain $[0, T] \subseteq \mathbb{R}_{\geq 0}$, $T > 0$, and with $\overline{\mathcal{B}}$ its closure. We assume the discontinuities $S_i \subseteq \mathbb{R}^{d+1}$ to be d -dimensional regular (smooth) manifolds given as hypergraphs $S_i = \{(x, t) \in \overline{\mathcal{B}} : t = s_i(x)\}$, with $s_i : \Omega \rightarrow [0, T]$ to be at least of class $s \in C^1(\Omega, [0, T])$, such that $L_i(x, t) = s_i(x) - t$ is a regular level-set function, i.e., $\nabla_{x,t} L(x, t) \neq 0$, for all $(x, t) \in L_i^{-1}(0) = S_i$, $i = 1, \dots, l$.

Let $BV(\mathcal{B})$ be the space of *Bounded Variations* [11]. For the i -th parametrization s_i we associate the Heaviside function $H_{s_i} \in BV(\mathcal{B})$ as:

$$H_{s_i}(x, t) = \begin{cases} 1 & , t \leq s_i(x) \\ 0 & , t > s_i(x) \end{cases} . \quad (1)$$

We set $\mathbb{S} = \bigcup_{i=1}^l S_i$ and $\mathfrak{h}^T H_{\mathbb{S}} = \sum_{i=1}^l h_i H_{s_i}$, where $\mathfrak{h} = \{h_i\}_{i=1}^l \subseteq \mathbb{R}$ denotes the constant heights. We define the space of functions with l jump discontinuities at \mathbb{S} as

$$M_l(\mathcal{B}) = \{f \in BV(\mathcal{B}) : f = g + \mathfrak{h}^T H_{\mathbb{S}}, \text{ for some } g \in C^0(\mathcal{B})\} . \quad (2)$$

We denote with $\Pi_{n,d}$ the space of d -variate polynomials of maximum l_∞ -degree $n \in \mathbb{N}$, spanned by some polynomial basis $\{q_\alpha\}_{\alpha \in A_{n,d}}$. Hereby, $A_{n,d} = \{\alpha \in \mathbb{N}^d : \|\alpha\|_\infty \leq n\}$ denotes the associated multi-index set, which we assume to be lexicographically ordered. In particular, we consider three possible bases [3]:

- i) The canonical polynomials $x^\alpha = x^{\alpha_1} \cdots x^{\alpha_d}$, $\alpha \in A_{n,d}$.

- ii) The Chebyshev polynomials of the first kind $T_\alpha = T_{\alpha_1} \cdots T_{\alpha_d}$, $\alpha \in A_{n,d}$, with $T_k(x) = \arccos(k \cos(\pi x))$, $x \in [-1, 1]$, for $k \in \mathbb{N}$.
- iii) The Lagrange basis L_α with $L_\alpha(p_\beta) = \delta_{\alpha,\beta}$, defined with respect to a given set of interpolation nodes $p_\beta \in P_{n,d} \subseteq [-1, 1]^d$, both enumerated by $\alpha, \beta \in A_{n,d}$.

1.1.1 Sobolev Cubatures

We shortly recap the notion of *Sobolev Cubatures* from our prior work [1, 2]: Let $P_{n,d+1} = P_{n,d} \times P_T \subseteq \mathcal{B}$, where $P_T \subseteq [0, T]$ denotes the re-scaled 1D Legendre grid. We denote with $\{L_\alpha\}_{\alpha \in A_{n,d+1}} \in \Pi_{n,d+1}$ the Lagrange basis with respect to $P_{n,d+1}$ and with $\omega_\alpha = \|L_\alpha\|_{L^2(\mathcal{B})}^2$ the associated Gauss-Legendre weights. We introduce $\mathbb{W}_{n,d+1} = \text{diag}(\{\omega_\alpha\}_{\alpha \in A_{n,d+1}}) \in \mathbb{R}^{|A_{n,d+1}| \times |A_{n,d+1}|}$ as the weight matrix and $\mathbb{D}_x, \mathbb{D}_t \in \mathbb{R}^{|A_{n,d+1}| \times |A_{n,d+1}|}$ as the polynomial differential matrices with respect to the Lagrange basis. For $\beta \in \mathbb{N}^{d+1}$, with l_1 -norm $\|\beta\|_1 = k \in \mathbb{N}$ we introduce the β -th differential operator and the k -th weight matrix by

$$\mathbb{D}^\beta = \prod_{i=1}^{d+1} \mathbb{D}_{\beta_i}, \quad \mathbb{W}_{n,d+1}^k = \sum_{\|\beta\|_1 \leq k} (\mathbb{D}^\beta)^T \mathbb{W}_{n,d+1} (\mathbb{D}^\beta). \quad (3)$$

The *Sobolev cubatures*, approximating the H^k -norm $\|\cdot\|_{H^k(\mathcal{B})}$ result as:

$$\|u\|_{H^k(\mathcal{B})} \approx \mathbf{u}^T \mathbb{W}_{n,d+1}^k \mathbf{u}, \quad u \in H^k(\mathcal{B}), \quad (4)$$

where $\mathbf{u} = u(P_{n,d+1}) \in \mathbb{R}^{|A_{n,d+1}|}$ denotes the evaluation vector.

1.2 Contribution

We introduce the *Hybrid Surrogate Models* (PSMs) $\psi_\theta \in \Psi_\theta(\mathcal{B})$, parametrized by coefficients $\theta \in \Theta$, approximating functions with $l \in \mathbb{N}$ jump discontinuities. The HSMs are given by combining a smooth surrogate $Q_\xi \in \Pi_{n,d+1}$ with Heaviside functions [11] $H_s \in BV(\mathcal{B})$. The HSMs induce the construction of a "stitching" map $\Phi : M_l(\mathcal{B}) \rightarrow C^0(\mathcal{B})$ that subtracts the detected byproduct-parametrization of the discontinuities from the solution, resulting in a continuous output that can be approximated with smooth polynomial surrogates.

The HSM solutions are computed by finding the optimal coefficients $\theta \in \Theta$, minimizing an associated discretized loss $\mathcal{L}^n : \Theta \rightarrow \mathbb{R}_{\geq 0}$, formulated for detecting the discontinuities. Here, we focus on two challenges:

- **Reconstruction problem:** Given a function, $g \in M_l(\mathcal{B})$ we seek for a HSM approximate $g \approx \psi_\theta \in \Psi_\theta(\mathcal{B}) \subseteq M_l(\mathcal{B})$, minimizing an associated discretized reconstruction loss $\mathcal{L}_0^n : \Theta \rightarrow \mathbb{R}_{\geq 0}$.

- **1D transport equation problem:** Given an initial condition, $u_0 \in M_l(\Omega)$ we want to find an HSM approximate $\psi_\theta \in \Psi_\theta(\mathcal{B})$ minimizing a discretized PDE loss $\mathcal{L}_{\text{PDE}}^n : \Theta \rightarrow \mathbb{R}_{\geq 0}$ associated to the transport equation with Dirichlet boundary conditions $g \in C^0(\partial\Omega \times (0, T])$ and constant velocity $v \in \mathbb{R}$, given by:

$$\partial_t u + v \partial_x u = 0, \quad \text{in } \mathcal{B}. \quad (5)$$

The capability of the HSM technique is demonstrated in numerical experiments.

2 Hybrid Surrogate Models HSMs

We detail the notion of the *Hybrid Surrogate Models* and introduce the notion of distributional derivative for the Heaviside function: To start with we consider

$$Q_\xi(x, t) = \sum_{\alpha \in A_{n,d+1}} \xi_\alpha T_\alpha(x, t) \in \Pi_{n,d+1}, \quad s_i(x) = \sum_{\alpha \in A_{m,d}} (C_i)_\alpha x^\alpha \in \Pi_{m,d}$$

a polynomial Q_ξ in *Chebyshev* basis and the approximation s of the parametrizations $s_i^* \in C^1(\bar{\Omega}; [0, T])$ in canonical basis. We define the space of HSMs $\Psi_\theta(\mathcal{B}) \subseteq M_l(\mathcal{B})$ as:

$$\Psi_\theta(\mathcal{B}) = \{Q_\xi + \mathfrak{h}^T H_\mathbb{S} : Q_\xi \in \Pi_{n,d+1}\}, \quad (6)$$

where $\mathfrak{h}^T H_\mathbb{S} = \sum_{i=0}^l h_i H_{s_i} \in BV(\mathcal{B})$ with constant hights $\mathfrak{h} = \{h_i\}_{i=1}^l \subseteq \mathbb{R}$. We assume the elements $\psi_\theta \in \Psi_\theta(\mathcal{B})$ to be parametrized by $\theta = \{\xi, \beta, C\} \in \Theta \subseteq \mathbb{R}^p$, with $p = |A_{n,d}| + l + l \cdot |A_{m,d}|$.

2.1 Distributional Derivative

For leveraging the HSMs solving PDEs, we define the differential operators acting on the HSMs in the distributional sense. When restricting to the 1D setting $d = 1$ it suffices to provide the notion only with respect to time $t \in [0, T]$. The definition involves the spaces of test functions $\mathcal{D}(\mathcal{B})$ and the space of distributions $\mathcal{D}'(\mathcal{B})$ [11].

Lemma 1 *Let $d = 1$, $\Omega = (-1, 1)$ and $H_s \in BV(\mathcal{B})$ be the Heaviside function with jump discontinuity parametrized by $s \in C^1(\bar{\Omega}; [0, T])$, Eq. (1).*

(i) *Let $F \in \mathcal{D}'((0, T))$ be the distribution on time dependent functions, given by*

$$F[\phi(x, \cdot)] = \int_0^T H_s(x, t) \phi(x, t) dt, \quad \phi(x, \cdot) \in \mathcal{D}((0, T)), \quad x \in \Omega.$$

Then the distributional derivative $F'[\phi(x, \cdot)] = -F[\partial_t \phi(x, \cdot)]$ is equivalently given by

$$F'[\phi(x, \cdot)] = \phi(x, s(x)) = \int_0^T \phi(x, t) \delta_s(t) dt,$$

implying the distributional derivative $\partial_t H_s(x, t) = \delta_s(t)$ to be given by the Dirac function.

(ii) The distributional derivatives $\partial_x H_s(x, t), \partial_t H_s(x, t)$ satisfy the relation

$$\partial_x H_s(x, t) = \partial_t H_s(x, t) \partial_x s(x), \quad (7)$$

Proof. (i) is a standard result and can be found in [11]. To show (ii) we note that by our assumptions on the parametrization $s \in C^1(\bar{\Omega}; [0, T])$, section 1.1, the inverse function theorem [8] applies. Hence, $x = s^{-1}(t)$ for all $(x, t) \in S$. Consequently, when applying Fubini's theorem and (i), the following holds:

$$\int_{\Omega} \int_0^T \partial_x H_s(x, t) \phi(x, t) dt dx = - \int_0^T \phi(s^{-1}(t), t) dt = - \int_{\Omega} \phi(x, s(x)) \partial_x s(x) dx. \quad (8)$$

On the other side, by (i) the distributional derivative $\partial_t H_s(x, t)$ is given by

$$\int_{\Omega} \int_0^T \partial_t H_s(x, t) \phi(x, t) dt = - \int_{\Omega} \phi(x, s(x)) dx, \quad (9)$$

yielding the claim. \square

We provided all the ingredients for addressing the reconstruction problem.

3 Reconstruction Problem and Discretized Loss

We state a variational optimization problem, providing HSM approximates of non-continuous functions $g \in M_l(\mathcal{B})$. To do so, we introduce reconstruction loss functional $\mathcal{L}_0 : \Psi_{\theta}(\mathcal{B}) \rightarrow \mathbb{R}_{\geq 0}$, by

$$\inf_{\psi_{\theta} \in \Psi_{\theta}(\mathcal{B})} \mathcal{L}_0[\psi], \text{ with } \mathcal{L}_0[\psi_{\theta}] = \int_{\Omega} \int_0^T \left(\psi_{\theta}(x, t) - g(x, t) \right)^2 dt dx. \quad (10)$$

For the sake of simplicity, we assume that $g \in M_l(\mathcal{B})$ possesses only one single discontinuity, $l = 1$, parametrized by $s^* \in C^1(\bar{\Omega}; [0, T])$ with constant height $h^* \in \mathbb{R}$. Given $\psi_{\theta} \in \Psi(\mathcal{B})$ and its approximate $s \approx s^*$, $s \in \Pi_{m, d+1}$ we define the subdomains

$$U_-(s) = \{(x, t) \in \mathcal{B} : t \leq s(x)\}, \quad U_+(s) = \{(x, t) \in \mathcal{B} : t > s(x)\} \quad (11)$$

Applying the definition of $\Psi_\theta(\mathcal{B})$ from Eq. (6), the reconstruction loss \mathcal{L}_0 is equivalent to:

$$\mathcal{L}_0[\psi_\theta] = \int_{U_-} (Q_\xi + h - g)^2 dt dx + \int_{U_+} (Q_\xi - g)^2 dt dx. \quad (12)$$

We exploit the Sobolev cubatures for discretizing the loss.

Definition 1 (discretized reconstruction loss). *Let $\mathbf{u}_\pm(s) = \{(x_\alpha, t_\alpha) \in P_{n,d+1} : (x_\alpha, t_\alpha) \in U_\pm(s)\}$ be the left/right training sets, $A_{n,d+1}^\pm = \{\alpha \in A_{n,d+1} : (x_\alpha, t_\alpha) \in U_\pm(s)\}$, and $\mathbb{T}_{n,d+1}^\pm(s) = (T_i(\mathbf{u}_\pm(s)))_{i \in A_{n,d+1}}$ denote the left/right Vandermonde matrices with respect to the Chebyshev basis, and $\mathbb{W}_{n,d+1}^{k\pm} = (\mathbb{W}_{n,d+1}^k)_{i,j \in A_{n,d+1}^\pm}$ be the corresponding left/right Sobolev weight matrices. The discretized reconstruction loss $\mathcal{L}_0^n : \Theta \rightarrow \mathbb{R}_{\geq 0}$, Eq. (12) with Sobolev cubatures of order $k > 0$ is defined as*

$$\mathcal{L}_0^n[\theta] = (\mathbf{r}_{n,d+1}^-)^T \mathbb{W}_{n,d+1}^{k-} \mathbf{r}_{n,d+1}^- + (\mathbf{r}_{n,d+1}^+)^T \mathbb{W}_{n,d+1}^{k+} \mathbf{r}_{n,d+1}^+, \quad (13)$$

where the left and right residuals are given by $\mathbf{r}_{n,d+1}^- = \mathbb{T}_{n,d+1}^- \xi + h - g(P_{n,d+1}^-)$, $\mathbf{r}_{n,d+1}^+ = \mathbb{T}_{n,d+1}^+ \xi - g(P_{n,d+1}^+)$.

We further extend the concept by addressing the PDE problem below.

4 Transport Equation in 1D with Discontinuous Solution

We focus on exploiting the HSMs for solving the 1D transport equation, Eq. (5), with an initial condition $u_0 \in M_l(\Omega)$, possessing one single discontinuity at $x_0 \in \Omega$, constant velocity $v \in \mathbb{R}$, and Dirichlet boundary conditions $g \in C^0(\partial\Omega \times [0, T])$. The corresponding solution $u \in M_l(\mathcal{B})$ has a single jump discontinuity, $l = 1$, parametrized by $s^* \in C^1(\bar{\Omega}; [0, T])$, and a constant height $h^* \in \mathbb{R}$.

Stating the problem in a distributional sense is to seek for a solution $u \in M_l(\mathcal{B})$, minimizing the loss

$$\mathcal{L}_{\text{PDE}}[u] = \sup_{\phi \in D(\mathcal{B})} \left(\int_{\Omega} \int_0^T (\partial_t u(x, t) + v \partial_x u(x, t)) \phi(x, t) dt dx \right)^2. \quad (14)$$

We propose to derive an approximate surrogate solution $\psi_\theta \in \Psi_\theta(\mathcal{B})$ by minimizing the discrete loss $\mathcal{L}^n : \Theta \rightarrow \mathbb{R}_{\geq 0}$, given by

$$\mathcal{L}^n[\theta] = \mathcal{L}_{\text{PDE}}^n[\theta] + \mathcal{L}_\partial^n[\theta] + \mathcal{L}_0^n[\theta],$$

where $\mathcal{L}_\partial^n[\theta]$, $\mathcal{L}_0^n[\theta]$ are the Sobolev cubature discretizations of the boundary and reconstruction losses

$$\mathcal{L}_\partial[\psi_\theta] = \int_{\partial\Omega} \int_0^T (\psi_\theta - u_0)^2 dt dx, \quad \mathcal{L}_0[\psi_\theta] = \int_{\Omega} (\psi_\theta(x, 0) - g(x))^2 dx.$$

The discrete PDE loss, Eq. (14), results when restricting the space of test functions to the polynomial space $\Pi_{n,d+1}$:

$$\mathcal{L}_{\text{PDE}}^n[\theta] = \sum_{\alpha \in A_{n,d+1}} \left(\int_{\Omega} \int_0^T (\partial_t \psi_\theta + v \partial_x \psi_\theta) L_\alpha dt dx \right)^2. \quad (15)$$

The following Lemma, gives us an equivalent equation for the discrete PDE loss, by exploiting the structure of the HSM surrogates.

Lemma 2 *Let $\psi_\theta \in \Psi_\theta(\mathcal{B})$ parametrized by $\theta = \{\xi, h, C\}$, then the discretized PDE loss associated to Eq. (15) is equivalent to:*

$$\begin{aligned} \mathcal{L}_{\text{PDE}}^n[\theta] &= \sum_{\alpha \in A_{n,d+1}} \left(\int_{\Omega} \int_0^T (\partial_t Q_\xi + v \partial_x Q_\xi) L_\alpha(x, t) dt dx \right. \\ &\quad \left. - h \int_{\Omega} ((1 + v \partial_x s(x)) L_\alpha(x, s(x))) dx \right)^2 \end{aligned} \quad (16)$$

Proof. Due to the linearity of the PDE operator $\partial \cdot + v \partial_x \cdot$, the definition of the HSMs, and the smoothness of $Q_\xi \in \Pi_{n,d+1}$ we obtain the first term, $\partial_t Q_\xi + v \partial_x Q_\xi$, in Eq. (16) in the strong sense. The discontinuous part, Eq. (17) follows from Lemma (1). \square

Definition 2 (discretized PDE loss). *Given the ingredients of Eq. (14) we define the discretized PDE loss, Eq. (16), in terms of the Sobolev cubatures of order $k = 0$ as:*

$$\begin{aligned} \mathcal{L}_{\text{PDE}}^n[\theta] &= \sum_{\alpha \in A_{n,d+1}} \left(\sum_{\gamma \in A_{n,d+1}} (\mathbb{D}_t \mathbb{T}_{n,d+1} \xi + v \mathbb{D}_x \mathbb{T}_{n,d+1} \xi)_\gamma \omega_\gamma \right. \\ &\quad \left. - h \sum_{\gamma \in A_{n,d}} ((1 + v \partial_x s(P_{n,d})) L_\alpha(s(P_{n,d}))_\gamma) \omega_\gamma \right)^2, \end{aligned}$$

where $\mathbb{T}_{n,d+1} = (T_\alpha(P_{n,d+1}))_{\alpha \in A_{n,d}} \in \mathbb{R}^{|A_{n,d+1}| \times |A_{n,d+1}|}$ denotes the Vandermonde matrix with respect to the Chebyshev polynomials and the Legendre grid.

Definition 2 can be extended to Sobolev cubatures of arbitrary order $k \in \mathbb{Z}$ [2]. The setup $k = -1$ is used in Experiment 2, while Definition 1 with $k = 0$ is the setting of our implementation for Experiment 1.

5 Numerical Experiments

We designed experiments, addressing the reconstruction problem of a non-linearly propagating shock, and the transport equation in 1D with discontinuous initial conditions.

We compare the HSMs with the methods mentioned in the introduction: PSMs [2], SC-PINNs [1], and standard NNs with ReLU and Sine activation functions. All models were trained over the same number of points. For HSMs, SC-PINNs, and PSMs we used Legendre grids of degree $n_x, n_t = 72$ in the x and t direction, respectively, while for the standard NNs, the training points were randomly sampled from \mathcal{B} . The NN-based models have an architecture of 5 fully connected layers with 50 activation functions each. The errors are measured in the l^1 -norm for $N = 300^2$ equidistant points $\{x_i, t_i\}_{i=1}^N \subseteq \mathcal{B}$.




Fig. 1: Ground truth for the numerical experiments

EXPERIMENT 1. (*Reconstruction Problem of a Discontinuous Function*) We consider the setting with $d = 1$ of approximating a discontinuous function $g \in BV(\mathcal{B})$ with a single jump discontinuity, $l = 1$, constant height $h^* = 5$, and analytical smooth part given by $\Phi(g)(x, t) = \sin(5x - f(t))$ with $f(t) = 0.3t^2 - 0.3$.





Fig. 2: Approximation errors for the reconstruction problem

Table 2b reports the errors. HSM reaches at least two orders of magnitude better accuracy, while requiring one order of magnitude less computational time compared to the NN based models. When considering the solutions at $t = 0.33$ in Fig. 2a, we observe that PSM, NN_Sine and SC_PINN run into Gibbs phenomenon. The NN_ReLU avoids the Gibbs phenomenon but fails to capture the discontinuity accurately, unlike the HSM that fits the discontinuity optimally.

We continue our investigations:

EXPERIMENT 2. (Transport equation in 1D with discontinuous initial condition)
 We keep the design of Experiment 1 and consider the 1D transport equation, with an initial condition possessing a discontinuity at $s_0 = -0.3$, a smooth part given by $\Phi(u_0(x)) = \sin(5x)$ and a constant height $h^* = 5$. We assume Dirichlet boundary conditions given by the analytical solution $u(x, t) = u_0(x - vt)$ and velocity $v = 1$.



(a) Predictions at $t = 0$

| method | l_1 -error | runtime (s) |
|---------|-------------------------------|---------------------|
| PSM | $3.77 \cdot 10^{-2}$ | $t = \mathbf{0.89}$ |
| SC-PINN | $1.44 \cdot 10^{-3}$ | $t = 214.47$ |
| NN-ReLU | $6.07 \cdot 10^{-3}$ | $t = 342.45$ |
| NN-Sine | $1.75 \cdot 10^{-2}$ | $t = 404.78$ |
| HSM | $\mathbf{1.45 \cdot 10^{-4}}$ | $t = 29.00$ |

(b) Approximation errors in \mathcal{B}

Fig. 3: Approximation of the 1D transport equation

Table 3b reports the errors. As expected all the methods except for the NN_ReLU and the HSM run into Gibbs phenomenon. All the methods are outperformed by the HSM, both in accuracy and in runtime. Fig. 3a clearly shows the HSM, circumventing the Gibbs phenomenon optimally.

6 Conclusions

The *Hybrid Surrogate Models (HSMs)* showed remarkable efficiency and approximation power when dealing with discontinuous functions, avoiding the Gibbs phenomenon while providing an approximated parametrization of the discontinuity as a byproduct. Extending the present 1D implementations to multiple shocks and higher dimensions holds great promise for advancing the field of shock-type PDE solvers.

Acknowledgements This research was partially funded by the Center of Advanced Systems Understanding (CASUS), which is financed by Germany's Federal Ministry of Education and Research (BMBF) and by the Saxon Ministry for Science, Culture, and Tourism (SMWK) with tax funds on the basis of the budget approved by the Saxon State Parliament.

References

- [1] J.-E. Cardona and M. Hecht. Polynomial differentiation decreases the training time complexity of physics-informed neural networks and strengthens their approximation power. *Machine Learning: Science and Technology*, 4, 10 2023.
- [2] J.-E. S. Cardona, P.-A. Hofmann, and M. Hecht. Learning partial differential equations by spectral approximates of general Sobolev spaces, 2023.
- [3] N. Carothers. Approximation theory - a short course. 11 2000.
- [4] D. Chae and S. Weng. Singularity formation for the incompressible Hall-MHD equations without resistivity. *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*, 33(4):1009–1022, 2016.
- [5] B. Cockburn. Discontinuous Galerkin methods. *Journal of Applied Mathematics and Mechanics / Zeitschrift für Angewandte Mathematik und Mechanik*, 83(11):731–754, 2003.
- [6] A. Ern and J.-L. Guermond. *Theory and practice of finite elements*, volume 159. Springer, 2004.
- [7] R. Eymard, T. Gallouët, and R. Herbin. Finite volume methods. *Handbook of numerical analysis*, 7:713–1018, 2000.
- [8] R. S. Hamilton. The inverse function theorem of Nash and Moser. *Bulletin (New Series) of the American Mathematical Society*, 7(1):65 – 222, 1982.
- [9] J. Hopfield. Neural networks and physical systems with emergent collective computational abilities. *Proceedings of the National Academy of Sciences of the United States of America*, 79:2554–8, 05 1982.
- [10] A. J. Jerri. *The Gibbs Phenomenon in Fourier Analysis, Splines and Wavelet Approximations*. Mathematics and Its Applications. Springer New York, NY, 1 edition, 1998.
- [11] R. P. Kanwal. *Distributional Derivatives of Functions with Jump Discontinuities*, pages 99–138. Birkhäuser Boston, Boston, MA, 2004.
- [12] R. J. LeVeque. *Finite difference methods for ordinary and partial differential equations: steady-state and time-dependent problems*. SIAM, 2007.
- [13] S. Li and W. K. Liu. *Meshfree particle methods*. Springer Science & Business Media, 2007.
- [14] S. Nakane. *Formation of Shocks for Nonlinear Partial Differential Equations of First Order*, pages 199–204. De Gruyter, Berlin, Boston, 1994.
- [15] C.-W. Shu. A brief survey on discontinuous Galerkin methods in computational fluid dynamics. *Advances in Mechanics*, 43:541–554, 11 2013.
- [16] Y.-H. Tseng and M.-C. Lai. A discontinuity and cusp capturing PINN for Stokes interface problems with discontinuous viscosity and singular forces, 2023.
- [17] X. Wang, B. Xu, Z. Chen, D. Del Col, D. Li, L. Zhang, X. Mou, Q. Liu, Y. Yang, and Q. Cao. Review of droplet dynamics and dropwise condensation enhancement: Theory, experiments and applications. *Advances in Colloid and Interface Science*, 305:102684, 2022.
- [18] A. Zygmund. *Trigonometric Series*. Cambridge Mathematical Library. Cambridge University Press, 3 edition, 2003.