

KK-RIGIDITY OF SIMPLE NUCLEAR C*-ALGEBRAS

CHRISTOPHER SCHAFHAUSER

ABSTRACT. It is shown that if A and B are unital separable simple nuclear \mathcal{Z} -stable C*-algebras and there is a unital embedding $A \rightarrow B$ which is invertible on KK -theory and traces, then $A \cong B$. In particular, two unital separable simple nuclear \mathcal{Z} -stable C*-algebras which either have real rank zero or unique trace are isomorphic if and only if they are homotopy equivalent. It is further shown that two finite strongly self-absorbing C*-algebras are isomorphic if and only if they are KK -equivalent in a unit-preserving way.

CONTENTS

Introduction	1
1. Preliminaries	3
2. Classification of approximate embeddings	5
3. Proof of Theorem A	9
4. The strongly self-absorbing case	11
References	14

INTRODUCTION

A conjecture of Elliott from the early 1990s ([13]) states that unital separable simple nuclear C*-algebras are classified up to isomorphism by K -theoretic invariants, in direct analogy with the Connes–Haagerup classification of separably acting injective factors ([6, 21]). After several decades of work by the C*-algebra community at large, a positive answer was reached: unital separable simple nuclear \mathcal{Z} -stable C*-algebras satisfying the universal coefficient theorem (UCT) are classified by their Elliott invariant, consisting of a suitable combination of K -theory and traces ([24, 34, 19, 20, 14, 44, 4]—see also [17, 2]).

Without significantly enlarging the invariant, \mathcal{Z} -stability is necessary for such a classification result ([45]), but the role of the UCT is far more mysterious. There is no known example of a separable nuclear C*-algebra failing the UCT, and the question of whether one exists is perhaps the most important open problem in the theory of nuclear C*-algebras, dating back to [39]. The purpose of this paper is to examine the role of the UCT in the classification theorem.

In the infinite setting, the Kirchberg–Phillips theorem ([24, 34]) states that two unital separable simple nuclear purely infinite C*-algebras (i.e., unital Kirchberg algebras) A and B are isomorphic if and only if there is an invertible element in

Date: September 9, 2024.

2020 *Mathematics Subject Classification.* 46L35.

This work was partially supported by NSF grants DMS-2000129 and DMS-2400178.

$KK(A, B)$ which preserves the unit on K_0 . In the same spirit, in the finite setting, one would hope for a classification result without the UCT using some combination of KK -theory and traces as the invariant. The following result may be viewed as proof of concept.

Theorem A. *If A and B are unital separable simple nuclear \mathcal{Z} -stable C^* -algebras and there is a unital embedding $A \rightarrow B$ which induces an invertible element in KK -theory and an invertible map on traces, then $A \cong B$.*

The condition that an embedding $A \rightarrow B$ is invertible in KK -theory is, loosely speaking, saying that it is invertible up to a weak notion of homotopy. In particular, if there is a homotopy equivalence $A \rightarrow B$ which induces a bijection on traces, then $A \cong B$. If A and B are C^* -algebras for which projections separate traces, then any homotopy equivalence $A \rightarrow B$ will induce a bijection on traces, leading to the following homotopy rigidity result. Note that, in particular, this holds if A and B either have real rank zero or unique trace.

Corollary B. *If A and B are unital separable simple nuclear \mathcal{Z} -stable C^* -algebras for which projections separate traces, then $A \cong B$ if and only if A and B are homotopy equivalent.*

For strongly self-absorbing C^* -algebras, a stronger result holds. Recall that a unital separable infinite dimensional C^* -algebra A is *strongly self-absorbing* if $A \cong A \otimes A$ via an isomorphism approximately unitarily equivalent to the first factor embedding. Such C^* -algebras are always simple, nuclear, and \mathcal{Z} -stable, and the finite ones have unique trace. Infinite strongly self-absorbing C^* -algebras are Kirchberg algebras, and hence, as noted above, they are isomorphic if and only if they are KK -equivalent in a unit-preserving way. It turns out that this also holds in the finite case.

Theorem C. *If A and B are finite strongly self-absorbing C^* -algebras, then $A \cong B$ if and only if there is an invertible $\kappa \in KK(A, B)$ such that the induced map $\kappa_0: K_0(A) \rightarrow K_0(B)$ sends $[1_A]$ to $[1_B]$.*

Note that strong self-absorption is necessary in the previous theorem: the result does not extend to unital separable simple nuclear \mathcal{Z} -stable finite C^* -algebras, even if one assumes the algebras have a unique trace. Indeed, there is extra information encoded in the real character $K_0(A) \rightarrow \mathbb{R}$ induced by the unique trace on such a C^* -algebra A . This can be seen in the setting of unital simple AF algebras with unique trace. For instance, take $K_0(A) = \mathbb{Q} + \mathbb{Q}\sqrt{2} \subseteq \mathbb{R}$ with the order inherited from \mathbb{R} and unit $[1_A] = 1$, and take $K_0(B) = \mathbb{Q} \oplus \mathbb{Q}$ with the strict ordering from the first coordinate and unit $[1_B] = (1, 0)$. It is easy to see that A and B are KK -equivalent in a unit-preserving way but are not isomorphic.

In general, any non-UCT classification result in the finite case must account for some interaction between KK -theory and traces, the same way that the Elliott invariant includes the natural pairing $T \times K_0 \rightarrow \mathbb{R}$. The following is the most optimistic candidate for a compatibility condition. Note that this is true under the UCT since the K -theory map $\kappa_*: K_*(A) \rightarrow K_*(B)$ induced by κ and the trace map γ form an isomorphism between the Elliott invariants of A and B .¹

¹Conversely, if A and B satisfy the UCT and (κ_*, γ) is an isomorphism of the Elliott invariants, then κ_* lifts to an invertible $\kappa \in KK(A, B)$ by [39, Corollary 7.5], and the pair (κ, γ) satisfies the hypotheses of Conjecture D.

Conjecture D. *Suppose A and B are unital separable simple nuclear \mathcal{Z} -stable C^* -algebras. If there is an invertible $\kappa \in KK(A, B)$ and an affine homeomorphism $\gamma: T(B) \rightarrow T(A)$ with $\kappa_0([1_A]) = [1_B]$ and $\langle \gamma(\tau), x \rangle = \langle \tau, \kappa_0(x) \rangle$ for all $\tau \in T(B)$ and $x \in K_0(A)$, then $A \cong B$.*

This conjecture may be too naive. It may be that some stronger, and perhaps more mysterious, compatibility condition is needed between KK and traces. Any modification of the pairing map for which the conjecture is true would be most welcome. Of course, whatever the “correct” compatibility condition is should be automatic for a pair (κ, γ) induced by a $*$ -homomorphism. This is, in some sense, why no compatibility condition is needed in Theorem A, where it is assumed that one starts with a $*$ -homomorphism inducing κ and γ .

The growing interest in group actions on “classifiable” C^* -algebras makes such a result more pressing. Even under the somewhat restrictive condition where an equivariant UCT may be possible (i.e., for actions in a suitable bootstrap class), there are typically no satisfactory formulas for equivariant KK -theory. To my knowledge, the only known equivariant UCTs are for certain finite group actions ([28, 32]) and for actions of compact connected Lie groups with torsion free fundamental group ([38]), and even in these cases, the formulas obtained are rather complicated. For this reason, obtaining classification results in the finite setting which do not depend on the UCT seems crucial for further development of the equivariant classification theory.²

Acknowledgments. A significant portion of this work was completed during my stay at the Fields Institute in the fall of 2023 during the thematic program on “Operator Algebras and Applications,” and I thank the institute and the organizers of the program for facilitating such a stimulating research environment. In addition, this project substantially benefited from my participation in the AIM SQuaRE program “von Neumann algebraic techniques in the classification of C^* -algebras” during the period 2019–2023 with José Carrión, Kristin Courtney, Jamie Gabe, Aaron Tikuisis, and Stuart White; I thank all of them for countless conversations over the last several years and thank AIM for their support. I also thank Eloísa Grifo for a helpful discussion about triangulated categories, which played a role in an earlier attempt at the results in this paper.

1. PRELIMINARIES

Throughout this article, A and B will denote unital separable simple nuclear \mathcal{Z} -stable C^* -algebras. Although some of these conditions can be relaxed in some of the intermediate results, they are crucial for the results stated in the introduction. For the sake of clarity, I chose to keep the same hypotheses on A and B throughout instead of aiming for maximal generality.

A unital separable simple nuclear \mathcal{Z} -stable C^* -algebra A is either purely infinite or stably finite ([37, Theorem 4.1.10(ii)]), and in the latter case, the tracial state space of A , $T(A)$, is non-empty ([22, Corollary 5.12]). In the infinite case, all of the results in this paper are standard consequences of the Kirchberg–Phillips theorem, so the reader may prefer to assume the C^* -algebras are finite throughout the paper.

The paper will freely use basic results from KK -theory. A thorough treatment can be found in [1] or [23]. For a much more condensed treatment, highlighting the

²For equivariant classification results in the infinite setting, see [18].

aspects of KK -theory most used in the classification theory, see [2, Section 5.1]. At various points in the paper, it will be necessary to work with KK -groups $KK(D, E)$ when E is not σ -unital—however, the first variable D will always be separable. In this case, one can define

$$(1.1) \quad KK(D, E) = \varinjlim KK(D, E_0),$$

where the limit is taken over all separable subalgebras $E_0 \subseteq E$.³ See [2, Appendix B] for further discussion.

For a unital C^* -algebra D , let $T(D)$ denote the set of tracial states on D , viewed as a weak*-compact convex subset of D^* . If D and E are unital C^* -algebras and $\phi: D \rightarrow E$ is a unital *-homomorphism, write $T(\phi): T(E) \rightarrow T(D)$ for the continuous affine map $\tau \mapsto \tau\phi$. Also, write $[1_D]$ for the class of the unit of D in the group $K_0(D)$. For an element $\kappa \in KK(D, E)$, the induced map $K_0(D) \rightarrow K_0(E)$ will be denoted by κ_0 .

When A and B are separable, the group $KK(A, B)$ carries a natural (often non-Hausdorff) topology ([8]), and the group $KL(A, B)$ can be defined as the quotient of $KK(A, B)$ by the closure of $\{0\}$. The advantage of working with KL over KK is that two *-homomorphisms $\phi, \psi: A \rightarrow B$ which are approximately unitarily equivalent may define different classes in $KK(A, B)$, but they will coincide in $KL(A, B)$; in fact, this is the reason KL -groups were initially introduced by Rørdam in [35] (where they were only defined when A satisfies the UCT).

The usual constructions of the topology on $KK(A, B)$ depend on the separability of B . To define $KL(A, B)$ when B is non-separable, following [2, Definition 5.5], define

$$(1.2) \quad Z_{KK(A, B)} = \left\{ (\text{ev}_\infty^B)_*(\kappa) : \begin{array}{l} \kappa \in KK(A, C(\mathbb{N}^\dagger, B)) \text{ with} \\ (\text{ev}_n^B)_*(\kappa) = 0 \text{ for all } n \in \mathbb{N} \end{array} \right\} \subseteq KK(A, B),$$

where $\mathbb{N}^\dagger = \mathbb{N} \cup \{\infty\}$ is the one-point compactification of \mathbb{N} (with \mathbb{N} viewed as a discrete space) and, for $n \in \mathbb{N}^\dagger$, $\text{ev}_n^B: C(\mathbb{N}^\dagger, B) \rightarrow B$ denotes the evaluation map $f \mapsto f(n)$. Then define $KL(A, B) = KK(A, B)/Z_{KK(A, B)}$. When B is separable, $Z_{KK(A, B)}$ coincides with the closure of $\{0\}$ by [8, Theorem 3.5], and so the above definition of $KL(A, B)$ coincides with the usual one in the case when B is separable. As with KK -theory, any map $\kappa \in KL(A, B)$ will induce a map $\kappa_0: K_0(A) \rightarrow K_0(B)$; note that if $\tilde{\kappa} \in KK(A, B)$ is a lift of κ , then $\tilde{\kappa}_0 = \kappa_0$, so there is no ambiguity caused by overusing the notation in this way.

The main tool in the paper is the trace-kernel extension, which first arose somewhat implicitly in the work of Winter ([47, 49]) and Matui and Sato ([31]) as a tool for proving \mathcal{Z} -stability and has since been heavily used in both the classification and regularity theories. Suppose A is a unital separable simple nuclear \mathcal{Z} -stable finite C^* -algebra. As noted above, this implies $T(A)$ is non-empty. Let $\ell^\infty(A)$ denote the C^* -algebra of bounded sequences in A and let

$$(1.3) \quad A^\infty = \ell^\infty(A) / \{a \in \ell^\infty(A) : \lim_{n \rightarrow \infty} \sup_{\tau \in T(A)} \tau(a_n^* a_n) = 0\}$$

be the uniform tracial sequence algebra of A . Also, let

$$(1.4) \quad A_\infty = \ell^\infty(A) / \{a \in \ell^\infty(A) : \lim_{n \rightarrow \infty} \|a_n\| = 0\}$$

³When E is σ -unital, the limit definition in (1.1) agrees with the standard definition; see [2, Proposition B.6], for example.

denote the norm sequence algebra of A . Then the trace-kernel extension of A is the extension

$$(1.5) \quad 0 \longrightarrow J_A \xrightarrow{j_A} A_\infty \xrightarrow{q_A} A^\infty \longrightarrow 0,$$

where q_A is defined on representing sequences by $(a_n)_{n=1}^\infty \mapsto (a_n)_{n=1}^\infty$, $J_A = \ker(q_A)$ is the trace-kernel ideal of A , and j_A is the inclusion map.

For future use, recall the central surjectivity result of [27, Theorem 3.3], stating that q_A restricts to a surjective map $A_\infty \cap A' \rightarrow A^\infty \cap A'$. The more general result below follows from [27, Propositions 4.5(iii) and 4.6].⁴

Proposition 1.1. *With the notation above, if $S \subseteq A_\infty$ is a separable subset, then q_A restricts to a surjective map $A_\infty \cap S' \rightarrow A^\infty \cap q_A(S)'$.*

Given a function $r: \mathbb{N} \rightarrow \mathbb{N}$ with $r(n) \rightarrow \infty$ as $n \rightarrow \infty$, let $r^*: A_\infty \rightarrow A_\infty$ be the *-homomorphism defined on representing sequences by $(a_n)_{n=1}^\infty \mapsto (a_{r(n)})_{n=1}^\infty$. If D is a separable C*-algebra, $\psi_\infty: D \rightarrow A_\infty$ is a *-homomorphism, and r is a map as above, call the composition $r^*\psi_\infty$ a reparameterization of ψ_∞ . This will be used in the “intertwining via reparameterization” result from [16, Theorem 4.3], which is modeled on the asymptotic version in [34, Proposition 1.3.7], where it is attributed to Kirchberg. Briefly, if ψ_∞ is unitarily equivalent to all of its reparameterizations, then there is a *-homomorphism $\psi: D \rightarrow A$ such that ψ_∞ is unitarily equivalent to $\iota_A\psi$, where ι_A is the canonical inclusion of A into A_∞ as constant sequences.

2. CLASSIFICATION OF APPROXIMATE EMBEDDINGS

Let A and B be unital separable simple nuclear \mathcal{Z} -stable finite C*-algebras. The goal of this section is to describe all unital embeddings $A \rightarrow B_\infty$ in terms of traces and KK -theory. The results of this section are all essentially from [3] and [2].

The classification is broken into three parts, collected in Theorems 2.1, 2.2, and 2.5. Although not quoted directly in this paper, it should be noted that the latter two results depend crucially on the Elliott–Kucerovsky characterization of (nuclearly) absorbing extensions and representations in terms of pure largeness ([15]), along with a series of results in the regularity theory allowing one to access pure largeness from conditions such as \mathcal{Z} -stability ([29, 33]).

The first step is a classification of unital embeddings $A \rightarrow B^\infty$ due to Castillejos, Evington, Tikuisis, and White in [3], which is based on the notion of complemented partitions of unity (CPoU), developed in their collaboration with Winter in [4].

Theorem 2.1 ([3, Theorem A]). *Let A and B be unital separable simple nuclear \mathcal{Z} -stable finite C*-algebras.*

- (i) *If $\theta, \rho: A \rightarrow B^\infty$ are unital embeddings with $T(\theta) = T(\rho)$, then θ and ρ are unitarily equivalent.*
- (ii) *If $\gamma: T(B^\infty) \rightarrow T(A)$ is a continuous affine map, then there is a unital embedding $\theta: A \rightarrow B^\infty$ with $T(\theta) = \gamma$.*

The second step of the classification is to decide when a given unital embedding $A \rightarrow B^\infty$ lifts along q_B to a unital embedding $A \rightarrow B_\infty$. The idea goes back to [40], where the question is addressed when B is the universal UHF algebra \mathcal{Q} (and with ultrapowers in place of sequence algebras).

⁴These results are stated in [27] with ultrapowers in place of sequence algebras, but after replacing all ultralimits with sequential limits, the same proofs apply here.

Theorem 2.2 ([2, Theorem 7.10(i)]). *Let A and B be unital separable simple nuclear \mathcal{Z} -stable finite C^* -algebras and let $\theta: A \rightarrow B^\infty$ be a unital embedding. There is a unital embedding $\psi: A \rightarrow B_\infty$ such that $q_B\psi = \theta$ if and only if there is $\kappa \in KK(A, B_\infty)$ such that*

$$(2.1) \quad \kappa_0([1_A]) = [1_{B_\infty}] \in K_0(B_\infty) \quad \text{and} \quad (q_B)_*(\kappa) = [\theta] \in KK(A, B^\infty).$$

The final step is to describe all unital embeddings $A \rightarrow B_\infty$ with prescribed tracial data up to unitary equivalence, assuming at least one such embedding exists. The result is essentially given in [2], but it will be convenient to set this up slightly differently. For two unital embeddings $\phi, \psi: A \rightarrow B_\infty$, if ϕ and ψ are unitarily equivalent, then $T(\phi) = T(\psi)$. Conversely, when $T(\phi) = T(\psi)$, there is a secondary obstruction to unitary equivalence in KL -theory, as described below.

Suppose D and E are C^* -algebras with D separable, $I \subseteq E$ is an ideal, and $\phi, \psi: D \rightarrow E$ are $*$ -homomorphisms whose pointwise difference $\phi - \psi$ has range contained in I —the pair (ϕ, ψ) is often called a (D, I) -Cuntz pair. Then ϕ and ψ have a formal difference $[\phi, \psi] \in KK(D, I)$, and hence also define an element $\langle \phi, \psi \rangle \in KL(D, I)$. This can be used to define what is called the Cuntz–Thomsen picture of KK -theory in [2] (see [7, 43]), which essentially views KK -theory as the set of Cuntz pairs up to a suitable notion of homotopy.

Definition 2.3. Suppose A and B are unital separable simple nuclear \mathcal{Z} -stable finite C^* -algebras. If $\phi, \psi: A \rightarrow B_\infty$ are unital embeddings with $T(\phi) = T(\psi)$, define

$$(2.2) \quad \langle \phi, \psi \rangle = [\text{ad}(u)\phi, \psi] \in KL(A, J_B),$$

where $u \in B_\infty$ is a unitary with $q_B \text{ad}(u)\phi = q_B\psi$.

The following result implies that the secondary invariant $\langle \phi, \psi \rangle$ is Definition 2.3 is well-defined.

Proposition 2.4 (cf. [2, Remark 7.5]). *Suppose A and B are unital separable simple nuclear \mathcal{Z} -stable finite C^* -algebras and $\phi, \psi: A \rightarrow B_\infty$ are unital embeddings with $T(\phi) = T(\psi)$.*

- (i) *There is a unitary $u \in B_\infty$ such that $q_B \text{ad}(u)\phi = q_B\psi$.*
- (ii) *The element $[\text{ad}(u)\phi, \psi] \in KL(A, J_B)$ is independent of the choice of unitary u .⁵*

Proof. To see the existence of u , note that Theorem 2.1(i) implies there is a unitary $\bar{u} \in B^\infty$ with $\text{ad}(\bar{u})q_B\phi = q_B\psi$. By [3, Proposition 2.1], the unitary group of B^∞ is path-connected, and hence there is a unitary $u \in B_\infty$ such that $q_B(u) = \bar{u}$. Then $q_B \text{ad}(u)\phi = q_B\psi$.

Suppose now that $u, v \in B_\infty$ are unitaries with $q_B \text{ad}(u)\phi = q_B\psi = q_B \text{ad}(v)\phi$. Then $q_B(v^*u) \in B^\infty \cap q_B(\phi(A))'$. By another application of [3, Proposition 2.1], the unitary group of $B^\infty \cap q_B(\phi(A))'$ is path-connected. Also, by Proposition 1.1, q_B restricts to a surjective $*$ -homomorphism $B_\infty \cap \phi(A)' \rightarrow B^\infty \cap q_B(\phi(A))'$. Hence there is a unitary $w \in B_\infty \cap \phi(A)'$ with $q_B(w) = q_B(v^*u)$. Then $vwu^* \in J_B + \mathbb{C}1_{B_\infty}$ is a unitary, and so

$$(2.3) \quad [\text{ad}(u)\phi, \psi] = [\text{ad}(vwu^*)\text{ad}(u)\phi, \psi] = [\text{ad}(v)\phi, \psi],$$

⁵The same proof shows that $\langle \phi, \psi \rangle$ can be viewed as a well-defined element of $KL(A, J_B)$, but this will not be needed.

where the first equality follows from [2, Proposition 5.4(ii)], for example, and the second equality uses that $\text{ad}(w)\phi = \phi$ by the choice of w . \square

The final piece of the classification of unital embeddings $A \rightarrow B_\infty$ is given below. Note that with the well-definedness of the secondary invariant $\langle \phi, \psi \rangle$ established, the result is essentially the same as [2, Theorem 7.10].

Theorem 2.5 (cf. [2, Theorem 7.10]). *Suppose A and B are unital separable simple nuclear \mathcal{Z} -stable finite C*-algebras.*

- (i) *Let $\phi, \psi: A \rightarrow B_\infty$ be unital embeddings. Then ϕ and ψ are unitarily equivalent if and only if $T(\phi) = T(\psi)$ and $\langle \phi, \psi \rangle = 0$.*
- (ii) *If $\psi: A \rightarrow B_\infty$ is a unital embedding and $\kappa \in KL(A, J_B)$ such that $\kappa_0([1_A]) = 0$, then there is a unital embedding $\phi: A \rightarrow B_\infty$ such that $T(\phi) = T(\psi)$ and $\langle \phi, \psi \rangle = \kappa$.*

Proof. The forward direction of (i) is immediate using the well-definedness of $\langle \phi, \psi \rangle$ (Proposition 2.4). Indeed, if $u \in B_\infty$ is a unitary with $\text{ad}(u)\phi = \psi$, then

$$(2.4) \quad \langle \phi, \psi \rangle = [\text{ad}(u)\phi, \psi] = [\psi, \psi] = 0.$$

For the converse, suppose $\langle \phi, \psi \rangle = 0$ and choose a unitary $u \in B_\infty$ such that $q_B \text{ad}(u)\phi = q_B \psi$, so $\langle \phi, \psi \rangle = [\text{ad}(u)\phi, \psi]$. Then the maps $\text{ad}(u)\phi$ and ψ both lift $\theta = q_B \psi: A \rightarrow B^\infty$, and $\tau\theta \in T(A)$ is faithful for all $\tau \in T(B^\infty)$ by the simplicity of A . Since $[\text{ad}(u)\phi, \psi] = 0$ in $KL(A, J_B)$, [2, Theorem 7.10(iii)] implies $\text{ad}(u)\phi$ and ψ are unitarily equivalent. In particular, ϕ and ψ are unitarily equivalent.

The proof of (ii) will use the same deunitization trick used in the proof of [2, Theorem 7.10] (and also in [40, 41]). For any C*-algebra D , let $\iota_2: D \rightarrow M_2(D)$ denote the inclusion $\iota_2(d) = d \oplus 0_D$. Given another C*-algebra E and a *-homomorphism $\rho: D \rightarrow E$, the induced *-homomorphism $M_2(D) \rightarrow M_2(E)$ is also denoted by ρ .

Let $\tilde{\kappa} \in KK(A, J_B)$ be a lift of κ . Define $\theta = q_B \psi: A \rightarrow B^\infty$ and note that θ is a unital embedding since A is simple. Then the unitization of $\iota_2\theta$ is faithful and hence full as A is simple. Also, since $q_B \iota_2 \psi = \iota_2 \theta$, applying [2, Proposition 7.9] with $\iota_2 \psi$ and $\iota_2 \theta$ in place of ψ and θ shows that there is a *-homomorphism $\phi': A \rightarrow M_2(B_\infty)$ such that $q_B \phi' = \iota_2 \theta$ and $[\phi', \iota_2 \psi] = (\iota_2)_*(\tilde{\kappa})$. Then by [2, Proposition 5.4(iv)],

$$(2.5) \quad (j_B \iota_2)_*(\tilde{\kappa}) = [\phi'] - [\iota_2 \psi] \in KK(A, M_2(B_\infty)),$$

and since $\kappa_0([1_A]) = 0$, it follows that

$$(2.6) \quad [\phi'(1_A)] = [\iota_2(\psi(1_A))] = [\iota_2(1_{B_\infty})] \in K_0(M_2(B_\infty)).$$

The assumptions on B imply that B has stable rank one ([36, Theorem 6.7]), and hence so does B_∞ .⁶ So there is a unitary $u \in M_2(B_\infty)$ with $\phi'(1_A) = u \iota_2(1_{B_\infty}) u^*$ (cf. [1, Proposition 6.5.1]).

Note that $q_B(u) \in M_2(B^\infty)$ commutes with $\iota_2(1_{B^\infty})$ since

$$(2.7) \quad q_B(\phi'(1_A)) = \iota_2(\theta(1_A)) = \iota_2(1_{B^\infty}) = q_B(\iota_2(1_{B_\infty})).$$

Using that $M_2(B^\infty) \cap \{\iota_2(1_{B^\infty})\}'$ has path-connected unitary group by [3, Proposition 2.1] and that q_B restricts to a surjective map

$$(2.8) \quad M_2(B_\infty) \cap \{\iota_2(1_{B_\infty})\}' \rightarrow M_2(B^\infty) \cap \{\iota_2(1_{B^\infty})\}'$$

⁶By [30, Lemma 19.2.2], for every element $b \in B_\infty$, there is a unitary $u \in B_\infty$ with $b = u|b|$. Then b is the limit of the invertible elements $u(|b| + \epsilon 1_{B_\infty})$ as $\epsilon \rightarrow 0^+$.

by Proposition 1.1, there is a unitary $v \in B_\infty$ commuting with $\iota_2(1_{B_\infty})$ such that $q_B(v) = q_B(u)$. Then $w = uv^*$ is a unitary in $M_2(J_B) + \mathbb{C}1_{M_2(B_\infty)}$ satisfying $\phi'(1_A) = w\iota_2(1_{B_\infty})w^*$.

There is a unital embedding $\phi: A \rightarrow B_\infty$ such that $\text{ad}(w)\iota_2\phi = \phi'$. By construction, $q_B\iota_2\phi = q_B\iota_2\psi$, and hence $q_B\phi = q_B\psi$. As noted in [2, Proposition 6.3(i)], $T(q_B)$ is invertible, and it follows that $T(\phi) = T(\psi)$. Also,

$$(2.9) \quad [\iota_2\phi, \iota_2\psi] = [\phi', \iota_2\psi] = (\iota_2)_*(\tilde{\kappa}) \in KK(A, M_2(J_B)),$$

using [2, Proposition 5.4(ii)] for the first equality. Then since $(\iota_2)_*$ is an isomorphism by the stability of KK -theory, $[\phi, \psi] = \tilde{\kappa} \in KK(A, J_B)$, and in particular, $\langle \phi, \psi \rangle = \kappa \in KL(A, J_B)$. \square

The existence result in the previous theorem is optimal, as the following proposition shows.

Proposition 2.6. *Suppose A and B are unital separable simple nuclear \mathcal{Z} -stable finite C^* -algebras and $\phi, \psi: A \rightarrow B$ are unital embeddings with $T(\phi) = T(\psi)$. Then the map $\langle \phi, \psi \rangle_0: K_0(A) \rightarrow K_0(J_B)$ induced by $\langle \phi, \psi \rangle$ vanishes on $[1_A]$.*

Proof. After replacing ϕ with a unitary conjugate of ϕ , assume $q_B\phi = q_B\psi$, so that $\langle \phi, \psi \rangle = [\phi, \psi]$. As noted in [2, Proposition 5.4(iv)], $(j_B)_*([\phi, \psi]) = [\phi] - [\psi]$, and since ϕ and ψ are unital, this implies

$$(2.10) \quad K_0(j_B)(\langle \phi, \psi \rangle_0([1_A])) = K_0(\phi)([1_A]) - K_0(\psi)([1_A]) = 0.$$

Also, a CPoU argument implies $K_1(B^\infty) = 0$ (see [2, Proposition 6.7], which is obtained by applying [3, Proposition 2.1] to matrix algebras over B^∞), and hence the six-term exact sequence in K -theory induced by the trace-kernel extension implies $K_0(j_B)$ is injective. The result follows. \square

The following result collects some arithmetic properties of the secondary invariants $\langle \phi, \psi \rangle \in KL(A, J_B)$ for later use.

Proposition 2.7. *Let A_0, A , and B be unital separable simple nuclear \mathcal{Z} -stable finite C^* -algebras. Suppose $\alpha: A_0 \rightarrow A$ is a unital embedding and $\phi, \psi, \rho: A \rightarrow B$ are unital embeddings with $T(\phi) = T(\psi) = T(\rho)$. Then*

- (i) $\langle \phi, \psi \rangle + \langle \psi, \rho \rangle = \langle \phi, \rho \rangle$,
- (ii) $-\langle \phi, \psi \rangle = \langle \psi, \phi \rangle$, and
- (iii) $\alpha^*(\langle \phi, \psi \rangle) = \langle \phi\alpha, \psi\alpha \rangle$.

Proof. Note that (ii) is immediate from (i) and the fact that $\langle \psi, \psi \rangle = 0$, which in turn follows from the fact that $[\psi, \psi] = 0$ and the well-definedness of $\langle \psi, \psi \rangle$. Also, (iii) follows directly from the definition of $\langle \cdot, \cdot \rangle$, together with the functoriality of KL -theory in the first variable. It remains to show (i). Fix unitaries $u, v \in B_\infty$ such that $q_B\text{ad}(u)\phi = q_B\psi$ and $q_B\text{ad}(v)\psi = q_B\rho$. Since KL -theory is invariant under unitary equivalence,

$$(2.11) \quad [\text{ad}(u)\phi, \psi] = \text{ad}(v)_*([\text{ad}(u)\phi, \psi]) = [\text{ad}(vu)\phi, \text{ad}(v)\psi].$$

Also,

$$(2.12) \quad [\text{ad}(vu)\phi, \text{ad}(v)\psi] + [\text{ad}(v)\psi, \rho] = [\text{ad}(vu)\phi, \rho].$$

Combining these two equations proves (i). \square

3. PROOF OF THEOREM A

The basic strategy behind the proof of Theorem A (and essentially all abstract isomorphism results in C*-algebra theory) is Elliott's intertwining argument. To show that a given unital embedding $\phi: A \rightarrow B$ is approximately unitarily equivalent to an isomorphism, it is enough to construct a unital embedding $\psi: B \rightarrow A$ which is a two-sided inverse for ϕ up to approximate unitary equivalence. In fact, by exploiting the symmetry between A and B , it will be enough to construct a unital embedding $\psi: B \rightarrow A$ which is a left inverse for ϕ up to approximate unitary equivalence. Indeed, repeating the argument with ψ in place of ϕ will give a left inverse for ψ , which will necessarily be approximately unitarily equivalent to ϕ by elementary algebra.

Some care is needed in this argument, however. Once a left inverse ψ is constructed, one must know that $[\psi]$ and $T(\psi)$ are invertible in order to repeat the argument to construct a left inverse of ψ . Since T is invariant under approximate unitary equivalence, it will follow that $T(\phi)T(\psi) = T(\psi\phi) = \text{id}_{T(A)}$, and hence the invertibility of $T(\phi)$ implies that of $T(\psi)$. However, this argument does not work on KK since KK is not invariant under approximate unitary equivalence. The salient point is that the approximate unitary invariance of KL does imply $[\psi]$ is invertible in $KL(B, A)$. Using that the secondary invariant appearing in Theorem 2.5 takes values in KL instead of KK , this will be sufficient to construct a left inverse for ψ , making use of the fact that a right inverse ϕ is already known to exist.

The following lemma will be applied twice in the proof of Theorem 3.2: once as written (when constructing a left inverse ψ for ϕ) and once after reserving the roles of A and B (when constructing a left inverse ϕ' for ψ).

Lemma 3.1. *Suppose A and B are unital separable simple nuclear \mathcal{Z} -stable finite C*-algebras and $\phi: A \rightarrow B$ is a unital embedding such that $[\phi] \in KL(A, B)$ and $T(\phi): T(B) \rightarrow T(A)$ are invertible. If there exists a unital embedding $\psi'_\infty: B \rightarrow A_\infty$ such that $T(\psi'_\infty\phi) = T(\iota_A)$, then there exists a unital embedding $\psi: B \rightarrow A$ such that $\psi\phi$ is approximately unitarily equivalent to id_A .*

Proof. Since $[\phi]$ is invertible in $KL(A, B)$, the natural transformation

$$(3.1) \quad \phi^*: KL(B, \cdot) \rightarrow KL(A, \cdot)$$

is also invertible. By Proposition 2.6, $\langle \iota_A, \psi'_\infty\phi \rangle \in KL(A, J_A)$ vanishes on the unit in K_0 . Since ϕ is unital, this also holds for $(\phi^*)^{-1}(\langle \iota_A, \psi'_\infty\phi \rangle)$ in $KL(B, J_A)$. Apply Theorem 2.5(ii) to obtain a unital embedding $\psi_\infty: B \rightarrow A_\infty$ such that $T(\psi_\infty) = T(\psi'_\infty)$ and

$$(3.2) \quad \langle \psi_\infty, \psi'_\infty \rangle = (\phi^*)^{-1}(\langle \iota_A, \psi'_\infty\phi \rangle) \in KL(B, J_A).$$

Applying ϕ^* to both sides of (3.2) yields

$$(3.3) \quad \langle \psi_\infty\phi, \psi'_\infty\phi \rangle = \phi^*(\langle \psi_\infty, \psi'_\infty \rangle) = \langle \iota_A, \psi'_\infty\phi \rangle,$$

using Proposition 2.7(iii) for the first equality. Then Proposition 2.7(i) and (ii) imply $\langle \psi_\infty\phi, \iota_A \rangle = 0$, and hence $\psi_\infty\phi$ is unitarily equivalent to ι_A by Theorem 2.5(i).

Let $r: \mathbb{N} \rightarrow \mathbb{N}$ be a function with $r(n) \rightarrow \infty$ as $n \rightarrow \infty$. Then $r^*\psi_\infty\phi$ is unitarily equivalent to $r^*\iota_A = \iota_A$, which, in turn, is unitarily equivalent to $\psi_\infty\phi$. As $T(\phi)$ is invertible, this implies $T(r^*\psi_\infty) = T(\psi_\infty)$. Also,

$$(3.4) \quad \phi^*(\langle r^*\psi_\infty, \psi_\infty \rangle) = \langle r^*\psi_\infty\phi, \psi_\infty\phi \rangle = 0,$$

where the first equality follows from Proposition 2.7(iii) and the second equality follows from Theorem 2.5(i) using that $r^*\psi_\infty\phi$ and $\psi_\infty\phi$ are unitarily equivalent. Since ϕ^* is invertible, (3.4) implies $\langle r^*\psi_\infty, \psi_\infty \rangle = 0$, and then another application of Theorem 2.5(i) yields that $r^*\psi_\infty$ and ψ_∞ are unitarily equivalent. As r was arbitrary, [16, Theorem 4.3] implies that there is a unital embedding $\psi: B \rightarrow A$ such that $\iota_A\psi$ is unitarily equivalent to ψ_∞ .

Note that $\iota_A\psi\phi$ is unitarily equivalent to $\psi_\infty\phi$ and hence also to ι_A . It follows that $\psi\phi$ and id_A are approximately unitarily equivalent as morphisms $A \rightarrow A$. \square

The following result is a more precise formulation of Theorem A. Although the algebras in the statement are not assumed to be finite, the new part of the theorem is the finite case. The infinite case follows from a direct application of the Kirchberg–Phillips theorem.

Theorem 3.2. *If A and B are unital separable simple nuclear \mathcal{Z} -stable C^* -algebras and $\phi: A \rightarrow B$ is a unital embedding with $[\phi] \in KK(A, B)$ and $T(\phi): T(B) \rightarrow T(A)$ invertible, then there is an isomorphism $A \rightarrow B$ approximately unitarily equivalent to ϕ .*

Proof. Note that since $T(A) \cong T(B)$, either A and B are both finite or both infinite. In the infinite case, A and B are purely infinite, and the result follows from the Kirchberg–Phillips theorem. For the rest of the proof, assume A and B are finite.

The first step is to construct a unital embedding $\psi'_\infty: B \rightarrow A_\infty$ as in Lemma 3.1. Define

$$(3.5) \quad \gamma = T(\phi)^{-1}T(q_A\iota_A): T(A^\infty) \rightarrow T(B)$$

and note that γ is continuous and affine. By Theorem 2.1(ii), there is a unital embedding $\theta: B \rightarrow A^\infty$ such that $T(\theta) = \gamma$. Then $T(\theta\phi) = T(q_A\iota_A)$. By Theorem 2.1(i), $\theta\phi$ and $q_A\iota_A$ are unitarily equivalent.

As $[\phi]$ is invertible in $KK(A, B)$, so is the natural transformation

$$(3.6) \quad \phi^*: KK(B, \cdot) \rightarrow KK(A, \cdot).$$

Let $\kappa = (\phi^*)^{-1}([\iota_A]) \in KK(B, A_\infty)$. Then $\kappa_0([1_B]) = [1_{A_\infty}] \in K_0(A_\infty)$ since both ι_A and ϕ are unital. Also,

$$(3.7) \quad \phi^*((q_A)_*(\kappa)) = [q_A\iota_A] = [\theta\phi] = \phi^*([\theta]) \in KK(A, A^\infty).$$

Since ϕ^* is invertible, this implies $(q_A)_*(\kappa) = [\theta]$, and hence by Theorem 2.2, there is a unital embedding $\psi'_\infty: B \rightarrow A_\infty$ such that $q_A\psi'_\infty = \theta$. Since $q_A\iota_A$ is unitarily equivalent to $\theta\phi = q_A\psi'_\infty\phi$,

$$(3.8) \quad T(\iota_A)T(q_A) = T(q_A\iota_A) = T(q_A\psi'_\infty\phi) = T(\psi'_\infty\phi)T(q_A),$$

and since $T(q_A): T(A^\infty) \rightarrow T(A_\infty)$ is invertible (see [2, Proposition 6.3(i)], for example), $T(\iota_A) = T(\psi'_\infty\phi)$.

By Lemma 3.1, there is a unital embedding $\psi: B \rightarrow A$ such that $\psi\phi$ is approximately unitarily equivalent to id_A . In particular, as KL and T are invariant under approximate unitary equivalence,

$$(3.9) \quad [\psi] = [\phi]^{-1} \in KL(B, A) \quad \text{and} \quad T(\psi) = T(\phi)^{-1}: T(A) \rightarrow T(B)$$

are invertible. Let $\phi'_\infty = \iota_B\phi: B \rightarrow A_\infty$ and note that

$$(3.10) \quad T(\phi'_\infty\psi) = T(\iota_B\phi\psi) = T(\psi)T(\phi)T(\iota_B) = T(\iota_B).$$

Applying Lemma 3.1 with the roles of A and B reversed and with ψ and ϕ'_∞ in place of ϕ and ψ'_∞ , there is a unital embedding $\phi': A \rightarrow B$ such that $\phi'\psi$ is approximately unitarily equivalent to id_B .

Note that $\phi'\psi\phi$ is approximately unitarily equivalent to both $\text{id}_B\phi = \phi$ and $\phi'\text{id}_A = \phi'$, and in particular, ϕ and ϕ' are approximately unitarily equivalent. It follows that ϕ and ψ are inverses up to approximate unitary equivalence. Finally, Elliott's intertwining argument (see [37, Corollary 2.3.4], for example), implies that ϕ is approximately unitarily equivalent to an isomorphism $A \rightarrow B$. \square

The homotopy rigidity result in the introduction is now immediate.

Proof of Corollary B. Suppose that A and B are unital separable simple nuclear \mathcal{Z} -stable C*-algebras for which projections separate traces and let $\phi: A \rightarrow B$ be a homotopy equivalence with homotopy inverse $\psi: B \rightarrow A$. Then ϕ is an embedding since A is simple and $[\phi] \in KK(A, B)$ is invertible with inverse $[\psi] \in KK(B, A)$. By Theorem 3.2, it suffices to show that ϕ is unital and $T(\phi)$ is invertible.

Since $\phi\psi$ is homotopic to id_B , there is a continuous path of projections from $\phi(\psi(1_B))$ to 1_B , and hence $\phi(\psi(1_B)) = 1_B$. Therefore,

$$(3.11) \quad 1_B = \phi(\psi(1_B)) \leq \phi(1_A) \leq 1_B,$$

and hence ϕ is unital. Note that by symmetry, ψ is also unital.

It remains to show $T(\phi)$ is invertible. Fix $\tau \in T(A)$. If $p \in A$ is a projection, then since $\psi\phi$ is homotopic to id_A , $\psi(\phi(p))$ is homotopic to p through a path of projections in A . In particular, $\psi(\phi(p))$ and p are unitarily equivalent, and hence $\tau(\psi(\phi(p))) = \tau(p)$. Since this holds for all projections $p \in A$ and projections in A separate traces, $\tau\psi\phi = \tau$. Therefore, $T(\phi)T(\psi) = \text{id}_{T(A)}$. By symmetry, $T(\psi)T(\phi) = \text{id}_{T(B)}$. So $T(\phi)$ is invertible with inverse $T(\psi)$. \square

4. THE STRONGLY SELF-ABSORBING CASE

Strongly self-absorbing C*-algebras were introduced by Toms and Winter in [46] (cf. [25]). A unital separable infinite dimensional C*-algebra A is strongly self-absorbing if the first factor embedding $\text{id}_A \otimes 1_A: A \rightarrow A \otimes A$ is approximately unitarily equivalent to an isomorphism. Any such C*-algebra A has an approximately inner half flip in the sense that the embeddings

$$(4.1) \quad \text{id}_A \otimes 1_A, 1_A \otimes \text{id}_A: A \rightarrow A \otimes A$$

are approximately unitarily equivalent ([46, Proposition 1.5]), and then A is simple and nuclear by [26, Lemma 3.10] (cf. [12, Propositions 2.7 and 2.8]). Also, A is either purely infinite or has unique trace, as is noted in [46, Theorem 1.7]. Finally, note that A is \mathcal{Z} -stable by main result of [48].

Strongly self-absorbing C*-algebras are very rigid, which makes them a natural test case for general problems about simple nuclear C*-algebras, a viewpoint promoted by Winter (see [50, Section 4.7], for example). Part of what makes the classification of strongly self-absorbing C*-algebras more accessible than the general case is that for any two strongly self-absorbing C*-algebras A and B , there is at most one unital embedding $A \rightarrow B_\infty$ up to unitary equivalence. This reduces the classification problem to proving an approximate existence theorem, characterizing when there exists a unital embedding $A \rightarrow B_\infty$. This reduction will be used in the form of the following well-known result.

Proposition 4.1. *If A and B are strongly self-absorbing C^* -algebras, the following statements are equivalent.*

- (i) $A \otimes B \cong B$.
- (ii) *There is a unital embedding $A \rightarrow B$.*
- (iii) *There is a unital embedding $A \rightarrow B_\infty$.*

Proof. The downward implications are clear. For (iii) \Rightarrow (i), by the Choi–Effros lifting theorem ([5, Theorem 3.10]), any unital embedding $A \rightarrow B_\infty$ lifts to a sequence of unital completely positive maps $\phi_n: A \rightarrow B$, which necessarily satisfy

$$(4.2) \quad \lim_{n \rightarrow \infty} \|\phi_n(a_1 a_2) - \phi_n(a_1)\phi_n(a_2)\| = 0, \quad a_1, a_2 \in A.$$

Then B is A -stable by [9, Proposition 4.2]. \square

The main extra ingredient that allows for the classification of strongly self-absorbing C^* -algebras in Theorem C is the following result of Dadarlat and Winter, which is a special case of the UCT satisfied by self-absorbing C^* -algebras.

Theorem 4.2 ([9, Theorem 3.4]). *If A is a strongly self-absorbing C^* -algebra and E is a separably A -stable C^* -algebra in the sense of [41, Definition 1.4], then there is an isomorphism $KK(A, E) \rightarrow K_0(E): \kappa \mapsto \kappa_0([1_A])$.*

Proof. Since E is separably A -stable, the set of separable A -stable subalgebras of E is cofinal in the set of separable subalgebras of E . Therefore,

$$(4.3) \quad KK(A, E) = \varinjlim KK(A, E_0) \quad \text{and} \quad K_0(E) = \varinjlim K_0(E_0),$$

with the limits taking over all separable A -stable subalgebras $E_0 \subseteq E$. By the naturality of the map $\kappa \mapsto \kappa_0([1_A])$, it suffices to prove the result when E is separable and A -stable. This is the content of [9, Theorem 3.4]. \square

The previous result will be applied when E is the tracial sequence algebra of a finite strongly self-absorbing C^* -algebra. The key observation is that since all finite strongly self-absorbing C^* -algebras have the same tracial sequence algebra, if A and B are finite strongly self-absorbing C^* -algebras, then B^∞ is separably A -stable.

Proposition 4.3. *Suppose A is a finite strongly self-absorbing C^* -algebra. Then $A^\infty \cong R^\infty$, where R is the separably acting hyperfinite II_1 factor.*

Proof. This is a standard consequence of Connes’s theorem and Kaplansky’s density theorem. Let τ denote the unique trace on A and note that $\pi_\tau(A)''$ is a separably acting finite von Neumann algebra, which is injective as A is nuclear ([11, Theorem 6.4(iv) \Rightarrow (v) and Theorem 6.2(i) \Rightarrow (ii)]). Further, since τ is the unique trace on A , $\pi_\tau(A)''$ is a factor (see [10, Theorem 6.7.3], for example). Since A is simple, and infinite dimensional, $\pi_\tau(A)''$ is also infinite dimensional. Therefore, $\pi_\tau(A)''$ is a separably acting injective II_1 factor, and by Connes’s theorem ([6, Theorem 7.1]), it is isomorphic to R .

Identify $\pi_\tau(A)''$ with R and, abusing notation, let τ also denote the unique trace on R . Define $\pi_\tau^\infty: A^\infty \rightarrow R^\infty$ on representing sequences by $(a_n)_{n=1}^\infty \mapsto (\pi_\tau(a_n))_{n=1}^\infty$. Since π_τ is isometric with respect to the 2-norms, π_τ^∞ is faithful. For any $b \in R^\infty$ represented by a bounded sequence $(b_n)_{n=1}^\infty \subseteq R$, by Kaplansky’s density theorem,⁷ there is a sequence $(a_n)_{n=1}^\infty \subseteq A$ such that $\|\pi_\tau(a_n) - b_n\|_2 < 1/n$ and $\|a_n\| \leq \|b_n\|$.

⁷This is using the standard fact that the strong topology on the unit ball of R is induced by the 2-norm $\|b\|_2 = \tau(b^*b)^{1/2}$. See [42, Proposition III.5.3], for example.

In particular, $(a_n)_{n=1}^\infty \subseteq A$ is a bounded sequence and hence defines an element $a \in A^\infty$. By construction, $\pi_\tau^\infty(a) = b$, proving π_τ^∞ is surjective. \square

The following result provides a one-sided version of Theorem C, characterizing when one finite strongly self-absorbing C*-algebra unitaly embeds into another.

Theorem 4.4. *For finite strongly self-absorbing C*-algebras A and B , there is a unital embedding $A \rightarrow B$ if and only if there is $\kappa \in KK(A, B)$ with $\kappa_0([1_A]) = [1_B]$.*

Proof. Applying Proposition 4.3 to both A and B implies $A^\infty \cong B^\infty$. In particular, there is a unital embedding $\theta: A \rightarrow B^\infty$. Further, this isomorphism also implies B^∞ is a quotient of $\ell^\infty(A)$, so B^∞ is separably A -stable by [41, Proposition 1.12]. Note that

$$(4.4) \quad (q_{B \wr B})_*(\kappa)_0([1_A]) = K_0(q_{B \wr B})(\kappa_0([1_A])) = [1_{B^\infty}] = K_0(\theta)([1_A]).$$

Then Theorem 4.2 implies $(q_{B \wr B})_*(\kappa) = [\theta]$, and Theorem 2.2 implies there is a unital embedding $A \rightarrow B^\infty$. The result follows from Proposition 4.1. \square

Symmetrizing the previous theorem gives Theorem C.

Proof of Theorem C. Let A and B be finite strongly self-absorbing C*-algebras and let $\kappa \in KK(A, B)$ be invertible with $\kappa_0([1_A]) = [1_B]$. Applying Theorem 4.4 to both κ and κ^{-1} , there are unital embeddings $A \rightarrow B$ and $B \rightarrow A$. Then two applications of Proposition 4.1 yield $A \cong A \otimes B \cong B$. \square

As a consequence of Theorem C, both the classification problem and embedding problem for strongly self-absorbing C*-algebras can be reduced to the infinite case.

Corollary 4.5. *Suppose A and B are finite strongly self-absorbing C*-algebras.*

- (i) *There is a unital embedding $A \rightarrow B$ if and only if there is a unital embedding $A \rightarrow B \otimes \mathcal{O}_\infty$.*
- (ii) *$A \cong B$ if and only if $A \otimes \mathcal{O}_\infty \cong B \otimes \mathcal{O}_\infty$.*

Proof. In both parts, the forward direction is trivial. To see the reverse direction in (i), suppose $\phi: A \rightarrow B \otimes \mathcal{O}_\infty$ is a unital embedding. Recall that the unital embedding $\text{id}_B \otimes 1_{\mathcal{O}_\infty}: B \rightarrow B \otimes \mathcal{O}_\infty$ is a KK -equivalence. Then $\phi^*([\text{id}_B \otimes 1_{\mathcal{O}_\infty}]^{-1})$ is an element of $KK(A, B)$ preserving the unit on K_0 , and hence Theorem 4.4 implies there is a unital embedding $A \rightarrow B$. Then the reverse direction of (ii) follows by symmetrizing (i) and applying Proposition 4.1 to obtain $A \cong A \otimes B \cong B$. \square

The reduction in Corollary 4.5 has a connection to the quasidiagonality problem for finite strongly self-absorbing C*-algebras. Let \mathcal{Q} denote the universal UHF algebra. In Winter's survey article [50], several variations of the quasidiagonality problem are considered, including the following questions ([50, Sections 4.9 and 4.10]).

- QDQ_{finite s.s.a.}**: Are all finite strongly self-absorbing C*-algebras quasidiagonal?
- QDQ_{infinite s.s.a.}**: Is $\mathcal{Q} \otimes \mathcal{O}_\infty$ the second largest strongly self-absorbing C*-algebra?

Winter also asks in [50, Question 7.2] if there are any formal implications between these problems. The following result is the implication

$$(4.5) \quad \mathbf{QDQ}_{\text{infinite s.s.a.}} \implies \mathbf{QDQ}_{\text{finite s.s.a.}}$$

Corollary 4.6. *If every strongly self-absorbing C^* -algebra not isomorphic to \mathcal{O}_2 unitaly embeds into $\mathcal{Q} \otimes \mathcal{O}_\infty$, then every finite strongly self-absorbing C^* -algebra is quasidiagonal.*

Proof. If A is a finite strongly self-absorbing C^* -algebra, then $A \not\cong \mathcal{O}_2$. So by assumption, there is a unital embedding $A \rightarrow \mathcal{Q} \otimes \mathcal{O}_\infty$. By Corollary 4.5(i), there is then a unital embedding $A \rightarrow \mathcal{Q}$, and as \mathcal{Q} is quasidiagonal, so is A . \square

REFERENCES

- [1] Bruce Blackadar. *K-theory for operator algebras*, volume 5 of *Mathematical Sciences Research Institute Publications*. Cambridge University Press, Cambridge, second edition, 1998.
- [2] José R. Carrión, James Gabe, Christopher Schafhauser, Aaron Tikuisis, and Stuart White. Classifying $*$ -homomorphisms I: Unital simple nuclear C^* -algebras. arXiv:2307.06480.
- [3] Jorge Castillejos, Samuel Evington, Aaron Tikuisis, and Stuart White. Classifying maps into uniform tracial sequence algebras. *Münster J. Math.*, 14(2):265–281, 2021.
- [4] Jorge Castillejos, Samuel Evington, Aaron Tikuisis, Stuart White, and Wilhelm Winter. Nuclear dimension of simple C^* -algebras. *Invent. Math.*, 224(1):245–290, 2021.
- [5] Man Duen Choi and Edward G. Effros. The completely positive lifting problem for C^* -algebras. *Ann. of Math. (2)*, 104(3):585–609, 1976.
- [6] Alain Connes. Classification of injective factors. Cases II_1 , II_∞ , III_λ , $\lambda \neq 1$. *Ann. of Math. (2)*, 104(1):73–115, 1976.
- [7] Joachim Cuntz. A new look at KK -theory. *K-Theory*, 1(1):31–51, 1987.
- [8] Marius Dadarlat. On the topology of the Kasparov groups and its applications. *J. Funct. Anal.*, 228(2):394–418, 2005.
- [9] Marius Dadarlat and Wilhelm Winter. On the KK -theory of strongly self-absorbing C^* -algebras. *Math. Scand.*, 104(1):95–107, 2009.
- [10] Jacques Dixmier. *C^* -algebras*. North-Holland Mathematical Library, Vol. 15. North-Holland Publishing Co., Amsterdam-New York-Oxford, 1977. Translated from the French by Francis Jellett.
- [11] Edward G. Effros and E. Christopher Lance. Tensor products of operator algebras. *Adv. Math.*, 25(1):1–34, 1977.
- [12] Edward G. Effros and Jonathan Rosenberg. C^* -algebras with approximately inner flip. *Pacific J. Math.*, 77(2):417–443, 1978.
- [13] George A. Elliott. The classification problem for amenable C^* -algebras. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994)*, pages 922–932. Birkhäuser, Basel, 1995.
- [14] George A. Elliott, Guihua Gong, Huaxin Lin, and Zhuang Niu. On the classification of simple amenable C^* -algebras with finite decomposition rank, II. arXiv:1507.03437.
- [15] George A. Elliott and Dan Kucerovsky. An abstract Voiculescu–Brown–Douglas–Fillmore absorption theorem. *Pacific J. Math.*, 198(2):385–409, 2001.
- [16] James Gabe. A new proof of Kirchberg’s \mathcal{O}_2 -stable classification. *J. Reine Angew. Math.*, 761:247–289, 2020.
- [17] James Gabe. Classification of \mathcal{O}_∞ -stable C^* -algebras. *Mem. Amer. Math. Soc.*, 293(1461):v+115, 2024.
- [18] James Gabe and Gábor Szabó. The dynamical Kirchberg–Phillips theorem. *Acta Math.*, 232(1):1–77, 2024.
- [19] Guihua Gong, Huaxin Lin, and Zhuang Niu. A classification of finite simple amenable \mathcal{Z} -stable C^* -algebras, I: C^* -algebras with generalized tracial rank one. *C. R. Math. Acad. Sci. Soc. R. Can.*, 42(3):63–450, 2020.
- [20] Guihua Gong, Huaxin Lin, and Zhuang Niu. A classification of finite simple amenable \mathcal{Z} -stable C^* -algebras, II: C^* -algebras with rational generalized tracial rank one. *C. R. Math. Acad. Sci. Soc. R. Can.*, 42(4):451–539, 2020.
- [21] Uffe Haagerup. Connes’ bicentralizer problem and uniqueness of the injective factor of type III_1 . *Acta Math.*, 158(1-2):95–148, 1987.
- [22] Uffe Haagerup. Quasitraces on exact C^* -algebras are traces. *C. R. Math. Acad. Sci. Soc. R. Can.*, 36(2-3):67–92, 2014.

- [23] Kjeld Knudsen Jensen and Klaus Thomsen. *Elements of KK-theory*. Mathematics: Theory & Applications. Birkhäuser Boston, Inc., Boston, MA, 1991.
- [24] Eberhard Kirchberg. The classification of purely infinite C^* -algebras using Kasparov's theory. Manuscript available at <https://ivv5hpp.uni-muenster.de/u/echters/ekneu1.pdf>.
- [25] Eberhard Kirchberg. Central sequences in C^* -algebras and strongly purely infinite algebras. In *Operator Algebras: The Abel Symposium 2004*, volume 1 of *Abel Symp.*, pages 175–231. Springer, Berlin, 2006.
- [26] Eberhard Kirchberg and N. Christopher Phillips. Embedding of exact C^* -algebras in the Cuntz algebra \mathcal{O}_2 . *J. Reine Angew. Math.*, 525:17–53, 2000.
- [27] Eberhard Kirchberg and Mikael Rørdam. Central sequence C^* -algebras and tensorial absorption of the Jiang–Su algebra. *J. Reine Angew. Math.*, 695:175–214, 2014.
- [28] Manuel Köhler. Universal coefficient theorems in equivariant KK -theory. Ph.D. Thesis, Georg-August-Universität Göttingen, 2010. Manuscript available at <http://hdl.handle.net/11858/00-1735-0000-0006-B6A9-9>.
- [29] Dan Kucerovsky and P. W. Ng. The corona factorization property and approximate unitary equivalence. *Houston J. Math.*, 32(2):531–550, 2006.
- [30] Terry A. Loring. *Lifting solutions to perturbing problems in C^* -algebras*, volume 8 of *Fields Institute Monographs*. American Mathematical Society, Providence, RI, 1997.
- [31] Hiroki Matui and Yasuhiko Sato. Strict comparison and \mathcal{Z} -absorption of nuclear C^* -algebras. *Acta Math.*, 209(1):179–196, 2012.
- [32] Ralf Meyer and George Nadareishvili. A universal coefficient theorem for actions of finite groups on C^* -algebras. arXiv:2406.11787.
- [33] Eduard Ortega, Francesc Perera, and Mikael Rørdam. The corona factorization property, stability, and the Cuntz semigroup of a C^* -algebra. *Int. Math. Res. Not. IMRN*, (1):34–66, 2012.
- [34] N. Christopher Phillips. A classification theorem for nuclear purely infinite simple C^* -algebras. *Doc. Math.*, 5:49–114, 2000.
- [35] Mikael Rørdam. Classification of certain infinite simple C^* -algebras. *J. Funct. Anal.*, 131(2):415–458, 1995.
- [36] Mikael Rørdam. The stable and the real rank of \mathcal{Z} -absorbing C^* -algebras. *Internat. J. Math.*, 15(10):1065–1084, 2004.
- [37] Mikael Rørdam and Erling Størmer. *Classification of nuclear C^* -algebras. Entropy in operator algebras*, volume 126 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2002. Operator Algebras and Non-commutative Geometry, 7.
- [38] Jonathan Rosenberg and Claude Schochet. The Künneth theorem and the universal coefficient theorem for equivariant K -theory and KK -theory. *Mem. Amer. Math. Soc.*, 62(348):vi+95, 1986.
- [39] Jonathan Rosenberg and Claude Schochet. The Künneth theorem and the universal coefficient theorem for Kasparov's generalized K -functor. *Duke Math. J.*, 55(2):431–474, 1987.
- [40] Christopher Schafhauser. A new proof of the Tikuisis–White–Winter theorem. *J. Reine Angew. Math.*, 759:291–304, 2020.
- [41] Christopher Schafhauser. Subalgebras of simple AF-algebras. *Ann. of Math. (2)*, 192(2):309–352, 2020.
- [42] Masamichi Takesaki. *Theory of operator algebras. I*. Springer-Verlag, New York-Heidelberg, 1979.
- [43] Klaus Thomsen. Homotopy classes of $*$ -homomorphisms between stable C^* -algebras and their multiplier algebras. *Duke Math. J.*, 61(1):67–104, 1990.
- [44] Aaron Tikuisis, Stuart White, and Wilhelm Winter. Quasidiagonality of nuclear C^* -algebras. *Ann. of Math. (2)*, 185(1):229–284, 2017.
- [45] Andrew S. Toms. On the classification problem for nuclear C^* -algebras. *Ann. of Math. (2)*, 167(3):1029–1044, 2008.
- [46] Andrew S. Toms and Wilhelm Winter. Strongly self-absorbing C^* -algebras. *Trans. Amer. Math. Soc.*, 359(8):3999–4029, 2007.
- [47] Wilhelm Winter. Decomposition rank and \mathcal{Z} -stability. *Invent. Math.*, 179(2):229–301, 2010.
- [48] Wilhelm Winter. Strongly self-absorbing C^* -algebras are \mathcal{Z} -stable. *J. Noncommut. Geom.*, 5(2):253–264, 2011.
- [49] Wilhelm Winter. Nuclear dimension and \mathcal{Z} -stability of pure C^* -algebras. *Invent. Math.*, 187(2):259–342, 2012.

- [50] Wilhelm Winter. QDQ vs. UCT. In *Operator algebras and applications—the Abel Symposium 2015*, volume 12 of *Abel Symp.*, pages 327–348. Springer, [Cham], 2017.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEBRASKA—LINCOLN, LINCOLN, NE, USA
Email address: `cschafhauser2@unl.edu`