

TRACES OF HECKE OPERATORS VIA HYPERGEOMETRIC CHARACTER SUMS

JEROME W. HOFFMAN, WEN-CHING WINNIE LI, LING LONG, AND FANG-TING TU

ABSTRACT. In this paper we obtain explicit formulas for the traces of Hecke operators on spaces of cusp forms in certain instances related to arithmetic triangle groups. These expressions are in terms of hypergeometric character sums over finite fields, a theory developed largely by Greene [28], Katz [36], Beukers-Cohen-Mellit [14], and Fuselier-Long-Ramakrishna-Swisher-Tu [25]. Our approach, in contrast to the previous works, is uniform and more geometric, and it works equally well for forms on elliptic modular curves and Shimura curves. The same method can be applied to obtain eigenvalues of Hecke operators as well.

CONTENTS

1. Introduction and main results	2
1.1. Introduction	2
1.2. Main results	3
Acknowledgements	6
2. Automorphic sheaves	7
2.1. Complex automorphic sheaves	7
2.2. ℓ -adic automorphic sheaves	8
3. Groups	18
3.1. Hypergeometric functions over \mathbf{C}	18
3.2. A motivating example	19
3.3. The groups to be considered	19
3.4. Isomorphisms using rigidity	23
3.5. Comparison Theorem	24
3.6. The ℓ -adic sheaves isomorphism	26
4. Hypergeometric character sums	26
4.1. Hypergeometric functions over finite fields	26
4.2. Beukers-Cohen-Mellit's finite Hypergeometric functions	28
4.3. For groups in a) list	29
4.4. For groups in b) list	30
4.5. Some properties of $H_p(HD(\Gamma); t)$	33
4.6. Special fiber at 1	35
4.7. Some character sum identities for the $k = 1$ cases	37
4.8. Some character sum identities for the $k = 2$ cases	38
5. Contributions from the singular fibers	39
5.1. The CM structure at elliptic points	39
5.2. Contributions from elliptic points	44
5.3. Contributions from cusps	48

2020 *Mathematics Subject Classification*. 11F11, 11F80, 11L05, 33C20.

Key words and phrases. Frobenius traces, arithmetic triangle groups, hypergeometric character sums, automorphic forms, Hecke operators.

6. Verification for the lowest weight cases	48
7. Further consequences	49
7.1. Other subgroups	49
7.2. Eigenvalues of Hecke operators	52
Appendix: G -sheaves	53
References	56

1. INTRODUCTION AND MAIN RESULTS

1.1. Introduction. In this paper we obtain explicit formulas for the traces of Hecke operators on spaces of cusp forms in certain instances related to arithmetic triangle groups. These expressions are in terms of hypergeometric character sums over finite fields, a theory developed largely by Greene [28], Katz [36], Beukers-Cohen-Mellit [14], and Fuselier-Long-Ramakrishna-Swisher-Tu [25].

Earlier works in this direction include [3] by Ahlgren and Ono for $\Gamma_0(8)$, [2] by Ahlgren for $\Gamma_0(4)$, [24] by Frechette, Ono and Papanikolas for newforms of $\Gamma_0(8)$, [26] by Fuselier for $\Gamma_0(1)$ and [42, 43] by Lennon for $\Gamma_0(3)$ and $\Gamma_0(9)$. The method in these papers is to combine (i) counting formulas for elliptic curves over finite fields, due to Schoof [57], and (ii) the Selberg trace formula. That proof is a descendant of the paper of Ihara [33] wherein he computed the traces of Hecke operators for $\mathrm{SL}(2, \mathbf{Z})$. Ihara did not use character sums and his counting of elliptic curves was based on results of Deuring. The paper [54] by Ono and Saad did it for $\Gamma_0(2)$ and $\Gamma_0(4)$, following Zagier's work [69] for $\Gamma_0(1)$. Their method uses the Rankin-Cohen brackets of Zagier's mock modular form.

In contrast to the works above, our approach, inspired by [45] by the last three authors and Scholl [56], is more geometric. It is based on Eichler-Shimura theory [61] as interpreted by Deligne [19] for elliptic modular groups and by Kuga-Shimura [40], Shimura [60] and Ohta [50, 51, 52] for the quaternion case. Let Γ be an arithmetic triangle group. Shimura's theory of canonical models implies that there is a smooth and projective algebraic curve X_Γ defined over an number field attached to it. For the groups considered in this paper, the number field is \mathbf{Q} . The key point is that the trace of the Hecke operator T_p acting on $S_{k+2}(\Gamma)$, the space of cusp forms¹ of weight $k+2$, is equal to the trace of Frobenius Frob_p acting on the H^1 of this curve X_Γ with coefficients in a suitable ℓ -adic sheaf, denoted by $V^k(\Gamma)_\ell$; see §2 for the setup. The Grothendieck-Lefschetz theorem implies

$$(1) \quad -\mathrm{Tr}(T_p | S_{k+2}(\Gamma)) = -\mathrm{Tr}(\mathrm{Frob}_p | H^1(X_\Gamma, V^k(\Gamma)_\ell)) = \sum_{\lambda \in X_\Gamma(\mathbf{F}_p)} \mathrm{Tr}(\mathrm{Frob}_\lambda | (V^k(\Gamma)_\ell)_\lambda),$$

and the main results of this paper express the terms on the right-hand side in terms of hypergeometric character sums attached to a hypergeometric datum $HD(\Gamma)$.

The novelty of this approach is twofold:

1. We give a unified treatment for Γ elliptic modular and Γ arising from quaternion algebras, specifically the quaternion algebra over \mathbf{Q} of discriminant 6. The latter continues the study of 4-dimensional Galois representations admitting quaternionic multiplications as initiated in [8, 44]. We make use of a certain family of abelian varieties with quaternion structures proposed in [18] and we extend some of the results of [45]. Moreover, if Γ' is a congruence subgroup of a group Γ studied by our approach such that $X_{\Gamma'}$ is defined over \mathbf{Q} and the canonical covering map $X_{\Gamma'} \rightarrow X_\Gamma$ is explicit, then our method also expresses the traces

¹By a cusp form we mean a holomorphic modular form which vanishes at all cusps of Γ . In particular, if Γ is cocompact, then a cusp form is just a modular form.


of the Hecke operators on $S_{k+2}(\Gamma')$ in terms of hypergeometric character sums attached to $HD(\Gamma)$. In particular, we obtain the Hecke traces for the groups cited above this way.

- Suppose $S_{k+2}(\Gamma)$ is m -dimensional. Our method used to compute the right-hand side of (1) can be easily extended to obtain the traces of $(\text{Frob}_p)^r$, $1 \leq r \leq m$, which in turn give rise to the eigenvalues of T_p on $S_{k+2}(\Gamma)$. Modular forms for a congruence subgroup arising from a quaternion algebra over \mathbf{Q} do not admit the usual Fourier expansion. This has been a handicap to understanding the arithmetic of modular forms on Shimura curves, as showcased in Yang's paper [68], where examples of Hecke eigenvalues for small primes are computed. Our method opens up a door to explore this territory.

1.2. **Main results.** The arithmetic triangle groups

$$(e_1, e_2, e_3) = \langle g_1, g_2 \mid g_1^{e_1} = g_2^{e_2} = (g_1 g_2)^{e_3} = id \rangle$$

have been classified by Takeuchi, [63], [64], and each can be realized as a discrete subgroup of $\text{PSL}_2(\mathbf{R})$ acting on the complex upper half plane \mathfrak{H} . The groups we consider are of two types: (1) those with noncompact fundamental regions, and therefore cusps, are commensurable with $\text{SL}_2(\mathbf{Z})$, and (2) those with compact fundamental regions, and therefore no cusps, are commensurable with $O_{B_6}^1$, the group of norm-1 elements of a maximal order (which is unique up to conjugation) in the quaternion algebra B_6 over \mathbf{Q} of discriminant 6. We also consider related groups gotten by adding Atkin-Lehner involutions. Note that modular forms on these quaternion groups are more difficult to understand because of the lack of q -expansions. In those cases, the Jacquet-Langlands (JL) correspondence allows one to relate these modular forms to cusp forms on congruence subgroups of $\text{SL}_2(\mathbf{Z})$, and we have utilized the JL correspondence to verify the calculations done in this paper related to the quaternions. Diagrams of the triangle groups and their interrelationships appear in Figures 1 and 2. In all cases, the curves X_Γ are of genus 0 and defined over \mathbf{Q} .



On the right, the group $(2,4,6)$ arises from the indefinite quaternion algebra $B_6 = \left(\frac{-1,3}{\mathbf{Q}}\right)$ with a maximal order $O_{B_6} = \mathbf{Z} + \mathbf{Z}I + \mathbf{Z}J + \mathbf{Z}\frac{1+I+J+IJ}{2}$, where $I^2 = -1, J^2 = 3, IJ = -JI$. Its elliptic points of order 2, 4, 6 are the fixed points of the Atkin-Lehner involutions w_6, w_2, w_3 , induced from the elements $3I + IJ, 1 + I, (3 + 3I + J + IJ)/2$ of O_{B_6} , respectively. As a group, $(2,4,6)$ is generated by the reduced norm 1 elements of O_{B_6} together with w_2, w_3 and w_6 modulo the center.

For some triangle groups Γ on the list, we associate hypergeometric data $HD(\Gamma) = \{\alpha(\Gamma), \beta(\Gamma)\}$ defined over \mathbf{Q} , to which Beukers-Cohen-Mellit [14] introduced hypergeometric character sums $H_p(HD(\Gamma), \lambda)$ for $\lambda \in \mathbf{F}_p^\times$ (cf. §4.1). We shall express the trace of T_p on $S_{k+2}(\Gamma)$ in terms of these

hypergeometric character sums. Our main result is as follows. For integers $m \geq 1$, let $F_m(S, T)$ be the degree- m polynomial in S and T defined by the recursive relation

$$(2) \quad F_{m+1}(S, T) = (S - T)F_m(S, T) - T^2 F_{m-1}(S, T), \quad F_0(S, T) = 1, \quad F_1(S, T) = S.$$

Then with $S = u^2 + uv + v^2$ and $T = uv$, we have

$$F_m(u^2 + uv + v^2, uv) = \sum_{i=0}^{2m} u^i v^{2m-i}.$$

Theorem 1. For $\Gamma = (2, 4, 6), (2, \infty, \infty), (2, 3, \infty), (2, 4, \infty), (2, 6, \infty)$, the table below describes the hypergeometric datum $HD(\Gamma) = \{\alpha(\Gamma), \beta(\Gamma)\}$ and the choice of a generator $\lambda = \lambda(\Gamma)$ of the field of \mathbf{Q} -rational functions on X_Γ by its values at each elliptic point of given order and each cusp:

Γ	$(2, \infty, \infty)$	$(2, 3, \infty)$	$(2, 4, \infty)$	$(2, 6, \infty)$	$(2, 4, 6)$
λ	$(1, 0, \infty)$	$(1, \infty, 0)$	$(1, \infty, 0)$	$(1, \infty, 0)$	$(1, \infty, 0)$
$\alpha(\Gamma)$	$\{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\}$	$\{\frac{1}{2}, \frac{1}{6}, \frac{5}{6}\}$	$\{\frac{1}{2}, \frac{1}{4}, \frac{3}{4}\}$	$\{\frac{1}{2}, \frac{1}{3}, \frac{2}{3}\}$	$\{\frac{1}{2}, \frac{1}{4}, \frac{3}{4}\}$
$\beta(\Gamma)$	$\{1, 1, 1\}$	$\{1, 1, 1\}$	$\{1, 1, 1\}$	$\{1, 1, 1\}$	$\{1, \frac{5}{6}, \frac{7}{6}\}$

Given $k \geq 2$ an even integer and a fixed prime ℓ , the terms on the right-hand side of (1) for almost all primes $p \neq \ell$ where X_Γ has good reduction can be explicitly expressed as follows.

For λ not corresponding to an elliptic point or a cusp,

$$(3) \quad \text{Tr}(\text{Frob}_\lambda \mid (V^k(\Gamma)_\ell)_{\bar{\lambda}}) = F_{k/2}(a_\Gamma(\lambda, p), p),$$

where

$$(4) \quad a_\Gamma(\lambda, p) = \begin{cases} \left(\frac{1-1/\lambda}{p}\right) H_p(HD(\Gamma), 1/\lambda) & \text{if } \Gamma \neq (2, 4, 6); \\ \left(\frac{-3(1-1/\lambda)}{p}\right) p H_p(HD(\Gamma), 1/\lambda) & \text{if } \Gamma = (2, 4, 6), \end{cases}$$

with $\left(\frac{\cdot}{p}\right)$ denoting the Legendre symbol. The contribution of λ corresponding to a cusp is 1. Let $\lambda = \lambda(z) \in X_\Gamma(\mathbf{F}_p)$ correspond to an elliptic point z of X_Γ of order N_z , which is a CM-point with complex multiplication (CM) by an imaginary quadratic field $K_z = \mathbf{Q}(\sqrt{d_z})$ listed below:

Γ	$(2, \infty, \infty)$	$(2, 3, \infty)$	$(2, 4, \infty)$	$(2, 6, \infty)$	$(2, 4, 6)$
$\sqrt{d_z}$	$\sqrt{-4}, -, -$	$\sqrt{-4}, \sqrt{-3}, -$	$\sqrt{-8}, \sqrt{-4}, -$	$\sqrt{-3}, \sqrt{-3}, -$	$\sqrt{-24}, \sqrt{-4}, \sqrt{-3}$

TABLE 1. CM fields of the elliptic points

When p is unramified in K_z , the contribution from $\lambda(z)$ in (1) is

$$(5) \quad \begin{cases} (-p)^{k/2} & \text{if } p \text{ is inert in } K_z; \\ \sum_{-\frac{k}{2N_z} \leq i \leq \frac{k}{2N_z}} p^{k/2} (\alpha_{z,p}^2/p)^{iN_z} & \text{if } p \text{ splits in } K_z. \end{cases}$$

In the latter case, upon picking any prime ideal \wp of the ring of integers of K_z above p , $\alpha_{z,p}$ can be chosen as any generator of the principal ideal \wp when $N_z > 2$. When $N_z = 2$, $\alpha_{z,p}^2$ can be taken as any root of $T^2 - \left(\frac{-3}{p}\right)^u p^u H_p(HD(\Gamma), 1)T + p^2 = 0$, where $u = 1$ if $\Gamma = (2, 4, 6)$, and $u = 0$ else.

In fact, we may choose $\alpha_{z,p}$ to be the Jacobi sum $J_\omega\left(\frac{1}{3}, \frac{1}{3}\right)$ (resp. $J_\omega\left(\frac{1}{4}, \frac{1}{4}\right)$) if $N_z \in \{3, 6\}$ (resp. $N_z = 4$). Here ω is any generator of the group of characters of \mathbf{F}_p^\times , and a Jacobi sum $J_\omega(a, b)$, defined as $\sum_{x \in \mathbf{F}_p} \omega^{(p-1)a}(x) \omega^{(p-1)b}(1-x)$, is a fundamental building block for hypergeometric character sums. See §4.1-§4.2 for more detail.

Here is an example illustrating our main result. For $\Gamma = (2, 4, 6)$ the lowest k with nontrivial $S_{k+2}(2, 4, 6)$ is $k = 6$, in which case $S_8(2, 4, 6) = \langle h_4^2 \rangle$ is 1-dimensional, where h_4 generates $S_4(2, 2, 3, 3)$ (cf. [9, §3.1]). By Jacquet-Langlands correspondence [34, 31, 27, 68], h_4^2 corresponds to the normalized weight-8 level 6 cuspidal newform $f_{6.8.a.a}$ in the L-functions and Modular Forms Database (LMFDB) label notation, which we will adopt throughout this paper. For primes $p > 5$, denote by $a_p(h_4^2)$ the eigenvalue of T_p on h_4^2 , which is equal to the p th Fourier coefficient $a_p(f_{6.8.a.a})$ of $f_{6.8.a.a}$. The theorem above gives

$$\begin{aligned} -a_p(h_4^2) = -a_p(f_{6.8.a.a}) &= \sum_{\lambda \in \mathbf{F}_p, \lambda \neq 0, 1} (a_\Gamma(\lambda, p)^3 - 2pa_\Gamma(\lambda, p)^2 - p^2a_\Gamma(\lambda, p) + p^3) \\ &\quad + p((pH_p(HD(\Gamma); 1))^2 - p^2) + \left(\binom{-1}{p} + \binom{-3}{p} + \binom{-6}{p} \right) p^3. \end{aligned}$$

More generally, under the Jacquet-Langlands correspondence, the space $S_{k+2}(2, 4, 6)$ is sent to $S_{k+2}^{new}(\Gamma_0(6), -, -)$, the subspace of newforms on $\Gamma_0(6)$ which are eigenfunctions with eigenvalues -1 of the Atkin-Lehner operators ω_2 and ω_3 , hence we obtain a formula for the traces of Hecke operators on the latter space.

For an index-2 subgroup $\Gamma \in \{(2, 6, 6), (2, 2, 2, 3), (3, 4, 4)\}$ of $(2, 4, 6)$, using the pullback of the hypergeometric sheaf on $X_{(2,4,6)}$ along the explicit \mathbb{Q} -rational covering map $X_\Gamma \rightarrow X_{(2,4,6)}$, we obtain the Hecke trace formulae for $S_{k+2}(\Gamma)$, which in turn gives rise to those for $S_{k+2}(2, 2, 3, 3)$ by inclusion-exclusion. Similar to $S_{k+2}(2, 4, 6)$, the Jacquet-Langlands correspondence maps $S_{k+2}(\Gamma)$ to the subspace of $S_{k+2}^{new}(\Gamma_0(6))$ consisting of common eigenfunctions of ω_2 and ω_3 with certain prescribed eigenvalues. See §7.1 for details. In particular, as a corollary of Theorem 14, the trace of T_p on $S_4(2, 2, 2, 3)$ for any prime $p > 5$ yields the following character sum identity

$$(6) \quad pH_p \left(\left\{ \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \right\}, \left\{ 1, 1, \frac{1}{6}, \frac{5}{6} \right\}; 1 \right) = \left(\frac{-1}{p} \right) a_p(f_{6.4.a.a}) + \left(\frac{-2}{p} \right) p.$$

The recent paper [4] provides an explicit hypergeometric-modularity method to express certain types of hypergeometric character sums in terms of Hecke eigenvalues of modular forms.

Explicit Hecke trace formulae have many applications. For instance, knowing Hecke trace can lead to special values of hypergeometric functions. As an example, it follows from the reasoning in Yang [68] that the Hecke trace $a_7(h_4^2)$ above yields the identity

$${}_3F_2 \left[\begin{matrix} \frac{1}{2} & \frac{1}{4} & \frac{3}{4} \\ \frac{5}{6} & \frac{7}{6} \end{matrix}; \frac{2^{10} \cdot 3^3 \cdot 5^6 \cdot 7}{11^4 \cdot 23^4} \right] = \frac{11 \cdot 23}{140\sqrt{3}} \frac{2^{1/3}(4 + 2\sqrt{2})}{7^{7/6}} \frac{\Gamma(7/6)\Gamma(13/24)\Gamma(19/24)}{\Gamma(5/6)\Gamma(17/24)\Gamma(23/24)}.$$

More can be said when the triangle group Γ is a matrix group not containing $-I_2$, such as $\Gamma = (3, \infty, \infty) \cong \Gamma_1(3)$ and $\Gamma = (\infty, \infty, \infty) \cong \Gamma_1(4)$, see §3 for more details. In these cases we obtain explicit Hecke trace formulas for almost all p and all $k \geq 2$. To describe our result, recall the identity (see for example [23])

$$(7) \quad \sum_{i=0}^k u^i v^{k-i} = \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} (-1)^j \binom{k-j}{j} (uv)^j \cdot (u+v)^{k-2j}.$$

Theorem 2. For $\Gamma = (e_\infty, \infty, \infty)$ with $e_\infty \in \{3, \infty\}$, let $HD(3, \infty, \infty) = \{\{\frac{1}{3}, \frac{2}{3}\}, \{1, 1\}\}$ and $HD(\infty, \infty, \infty) = \{\{\frac{1}{2}, \frac{1}{2}\}, \{1, 1\}\}$. Use $\Gamma_1(3)$ and $\Gamma_1(4)$ as their models in $\mathrm{SL}_2(\mathbf{Z})$ with level $N_\Gamma = 3, 4$ respectively, and choose the generator $\lambda = \lambda(\Gamma)$ of $\mathbf{Q}(X_\Gamma)$ so that λ takes values $\infty, 1, 0$ at the vertices of X_Γ of order e_∞, ∞, ∞ , respectively. Then for all integers $k \geq 1$, the contributions of

the right hand side of (1) at $\lambda \in \mathbf{F}_p \setminus \{0, 1\}$ are given by

$$(8) \quad \sum_{j=0}^{k/2} (-1)^j \binom{k-j}{j} p^j \cdot H_p(HD(\Gamma); 1/\lambda)^{k-2j},$$

The contributions at $\lambda = 0, 1, \infty$ are distinguished in two cases:

(1) For $\Gamma_1(4)$, they are $1, \left(\frac{-1}{p}\right)^k, \frac{1+(-1)^k}{2}$, respectively.

(2) For $\Gamma_1(3)$, those from $\lambda = 0$ and 1 are 1 and $\left(\frac{-3}{p}\right)^k$, and from $\lambda = \infty$ it is

$$\begin{cases} 0 & \text{if } p \equiv -1 \pmod{3}, \quad k \equiv 1 \pmod{2}; \\ (-p)^{k/2} & \text{if } p \equiv -1 \pmod{3}, \quad k \equiv 0 \pmod{2}; \\ (-1)^k \sum_{\substack{0 \leq i \leq k \\ k \equiv 2i \pmod{3}}} p^i \cdot J_\omega\left(\frac{1}{3}, \frac{1}{3}\right)^{k-2i} & \text{if } p \equiv 1 \pmod{3}, \end{cases}$$

where $\widehat{\mathbf{F}}_p^\times = \langle \omega \rangle$ and $J_\omega(a, b) = \sum_{x \in \mathbf{F}_p} \omega^{(p-1)a}(x) \omega^{(p-1)b}(1-x)$.

In particular, when $p \equiv -1 \pmod{N_\Gamma}$ and k is odd, $\text{Tr}(T_p | S_{k+2}(\Gamma)) = 0$.

We end this section with an outline of the proofs of Theorems 1 and 2, and the roles of the remaining sections. The trace formulae, and automorphic sheaves $V^k(\Gamma)$ over both the complex and ℓ -adic fields are discussed in §2. In particular, Proposition 3 and Remark 3 reduce the computation at generic points to $k = 2$ for Theorem 1 and $k = 1$ for Theorem 2, respectively. For certain arithmetic triangle groups Γ , it follows from the Rigidity Theorem 5 and the Comparison Theorem 8 in §3 that the automorphic sheaves $V^2(\Gamma)_\ell$ or $V^1(\Gamma)_\ell$, up to twists by rank-1 sheaves, could be replaced by hypergeometric sheaves for which the Galois action on a stalk has Frobenius traces explicitly expressed by hypergeometric character sums. In §3.3 we identify the list of such Γ whose associated hypergeometric data are defined over \mathbb{Q} ; these are the groups in Theorems 1 and 2. We recall in §4 the hypergeometric sheaves and derive their basic properties, following the works of Katz [36] and Beukers-Cohen-Mellit [14]. Using the information obtained, we determine, for each Γ , the required twist of the corresponding ℓ -adic hypergeometric sheaf so that the resulting sheaf is isomorphic to the automorphic sheaf $V^i(\Gamma)_\ell$ for $i = 2$ or 1 , as appropriate. This gives the contributions from generic points. The contributions from singular points are the theme of §5. All elliptic points are CM points. We first determine the CM structure at each elliptic point, with results summarized in Table 8 in §5.1. Then in §5.2 we compute the contribution at p from each elliptic point, which mainly depends on the order of the point and the behavior of p in the CM field. The contribution from each cusp is computed in §5.3. For k even it is 1, as done in Scholl [56], while for k odd, it is ± 1 or 0, depending on the type of the reduction at p of the degenerate elliptic curve at the singular point. This completes the proofs of Theorems 1 and 2. In §6 we provide an independent verification of the Hecke trace formula for $S_{k+2}(\Gamma)$ with smallest $k \geq 1$ by proving the hypergeometric character sum identities directly. The two novelties mentioned in §1.1 are addressed in §7. The Appendix at the end clarifies the G -Sheaves and the definition of the pushforward sheaves appearing in this paper.

ACKNOWLEDGEMENTS

Li is partially supported by Simons Foundation grant #355798, Long is supported in part by the Simons Foundation grant #MP-TSM-00002492 and the LSU Michael F. and Roberta Nesbit McDonald Professorship, Tu is supported by the NSF grant DMS #2302531. While working on this project, Li, Long, and Tu were hosted by the Institute of Mathematics, Academia Sinica, Taiwan in

the summer of 2023, Tu was also hosted at Oregon State University during the fall of 2023, and Li visited the University of Hong Kong in summer 2024. They would like to thank these Institutions for their hospitality and support.

2. AUTOMORPHIC SHEAVES

2.1. Complex automorphic sheaves.

2.1.1. For a Fuchsian subgroup Γ of $\mathrm{SL}_2(\mathbf{R})$ of first kind, let $X_\Gamma := X_\Gamma^{an} = \Gamma \backslash \mathfrak{H}^*$, where $\mathfrak{H}^* = \mathfrak{H}$ or the union of \mathfrak{H} and the cusps of Γ according to whether Γ is cocompact or not. Let $p : \mathfrak{H}^* \rightarrow X_\Gamma$ be the projection. Note that \mathfrak{H}^* carries a canonical topology, and p is a continuous map. In this paper, two kinds of Γ are considered:

1. Γ is commensurable with $\mathrm{SL}_2(\mathbf{Z})$. Then the cusps of Γ is the set of rational numbers and the point ∞ .
2. Γ is commensurable to the set O_B^1 of norm 1 elements in a maximal order O_B of an indefinite quaternion algebra B over \mathbf{Q} . That is, we choose once and for all an embedding $B \subset M_2(\mathbf{R})$, which induces an embedding $\theta : O_B^1 \subset \mathrm{SL}_2(\mathbf{R})$. In this case, the set of cusps is empty, so $\mathfrak{H}^* = \mathfrak{H}$.

We also have the set of elliptic points of Γ . We have a finite set $S = S_c \cup S_e \subset X_\Gamma$ of points which are cusps or elliptic points, that is, their preimages in \mathfrak{H}^* are cusps or elliptic points of Γ , respectively. Let

$$X_\Gamma^\circ = X_\Gamma - (S_c \cup S_e), \quad Y_\Gamma = X_\Gamma - S_c = \Gamma \backslash \mathfrak{H},$$

the complement of the set of cusps and elliptic points, and the complement of the set of cusps, respectively. Note that $p : \mathfrak{H} - \{\text{elliptic points of } \Gamma\} \rightarrow X_\Gamma^\circ$ is a covering space in the sense of topology. In particular, if Γ has no elliptic points, then $p : \mathfrak{H} \rightarrow Y_\Gamma$ is the universal covering of Y_Γ . The action of Γ on \mathfrak{H}^* is via the quotient $\bar{\Gamma}$ in $\mathrm{PSL}_2(\mathbf{R})$. For the groups in this paper, $\bar{\Gamma} = \Gamma/\Gamma \cap \{\pm I\}$. Therefore, if Γ is torsion-free, then $p : \mathfrak{H} \rightarrow Y_\Gamma$ is the universal covering of Y_Γ , and the fundamental group at any base-point x

$$\pi_1(Y_\Gamma, x) \cong \Gamma,$$

an isomorphism unique up to inner automorphism. In the quaternion cases this is $\pi_1(X_\Gamma, x) \cong \Gamma$ for Γ without torsion. In general, there is an epimorphism

$$\pi_1(X_\Gamma^\circ, x) \rightarrow \bar{\Gamma} \cong \mathrm{Aut}((\mathfrak{H} - \{\text{elliptic points of } \Gamma\})/X_\Gamma^\circ).$$

We consider local systems of \mathbf{C} -vector spaces of finite dimension on X_Γ° and their extensions to constructible sheaves on X_Γ . By the usual dictionary, a local system on X_Γ° is equivalent to a finite-dimensional representation of $\pi_1(X_\Gamma^\circ, x)$. In this paper, all the local systems will be defined by representations of $\bar{\Gamma}$.

2.1.2. There are constructible sheaves of \mathbf{C} -vector spaces $V^k(\Gamma)_{\mathbf{C}}$ on X_Γ for integers $k \geq 1$ with the following properties:

1. Over the open subset X_Γ° these automorphic sheaves are local systems of rank $k + 1$. They are thus solution sheaves to a system of regular singular differential equations on X_Γ° . For Γ considered in this paper, these will be hypergeometric local systems when $k = 1$ or 2.
2. The sheaves $V^k(\Gamma)_{\mathbf{C}}$ on X_Γ are extensions of the sheaves on X_Γ° defined above:

$$V^k(\Gamma)_{\mathbf{C}} = \iota_*(V^k(\Gamma)_{\mathbf{C}} | X_\Gamma^\circ) \quad \text{for the inclusion map } \iota : X_\Gamma^\circ \rightarrow X_\Gamma.$$

3. Fundamental theorems of Eichler, Shimura and Kuga give:

$$H_{par}^1(\Gamma, \mathrm{Sym}^k(\mathbf{C}^2)) \cong H^1(X_\Gamma, V^k(\Gamma)_{\mathbf{C}}) \cong S_{k+2}(\Gamma) \oplus \overline{S_{k+2}(\Gamma)},$$

which is a Hodge decomposition of type $(k+1, 0), (0, k+1)$. The left-hand side is parabolic cohomology. Moreover

$$H^1(X_\Gamma, V^k(\Gamma)_\mathbf{C}) \cong \text{Im}(H_c^1(X_\Gamma^\circ, V^k(\Gamma)_\mathbf{C}) \rightarrow H^1(X_\Gamma^\circ, V^k(\Gamma)_\mathbf{C}))$$

the image of the compactly supported cohomology. The Hecke operators act on the spaces as geometric correspondences and these isomorphisms are equivariant for the Hecke actions.

When $\Gamma \subset \text{SL}_2(\mathbf{R})$ is an arbitrary Fuchsian subgroup of the first kind, these sheaves are discussed in detail in [11] by Bayer and Neukirch. The group Γ acts on $\mathfrak{H}^* \times \text{Sym}^k(\mathbf{C}^2)$, on the first factor by fractional linear transformations, on the second factor via $\varrho_k := \text{Sym}^k(\varrho_1)$, where ϱ_1 arises from the inclusion $\Gamma \subset \text{SL}_2(\mathbf{R})$ and the canonical action of $\text{SL}_2(\mathbf{R})$ on \mathbf{C}^2 . Then

$$(9) \quad V^k(\Gamma)_\mathbf{C} := p_*^\Gamma(\text{Sym}^k(\mathbf{C}^2)) = \Gamma \backslash \mathfrak{H}^* \times \text{Sym}^k(\mathbf{C}^2),$$

where the middle expression uses the Γ -invariant push-forward functor. See the Appendix on G -sheaves for the theory of the functor p_*^G . Note that if $-I \in \Gamma$ these sheaves are identically zero if k is odd. When k is even, the action factors through the projective group $\bar{\Gamma} = \Gamma/\{\pm I\}$, and this action is discontinuous. When Γ is torsion-free, Bayer and Neukirch establish the Eichler-Shimura isomorphisms with cusp forms for all k , even or odd. When Γ has torsion and k is even, they show that by going to a torsion-free normal subgroup Γ' of finite index and using the projection $q : X_{\Gamma'} \rightarrow X_\Gamma$,

$$V^k(\Gamma)_\mathbf{C} = q_*^{\Gamma/\Gamma'}(V^k(\Gamma')_\mathbf{C})$$

gives the Eichler-Shimura isomorphism. When $x \in X_\Gamma$ is not a cusp or elliptic point, the stalk $(V^k(\Gamma)_\mathbf{C})_x$ is isomorphic to $\text{Sym}^k(\mathbf{C}^2)$. For x a cusp or elliptic point,

$$(V^k(\Gamma)_\mathbf{C})_x = \left(\text{Sym}^k(\mathbf{C}^2)\right)^{\Gamma_\tau}$$

where $\tau \in \mathfrak{H}^*$ is any point with $p(\tau) = x$ and Γ_τ is the stabilizer of τ in Γ . The same holds for k odd and Γ not containing $-I$.

2.2. ℓ -adic automorphic sheaves.

2.2.1. For Γ as before, in this section we view the quotient X_Γ as the collection of complex points of a curve over a number field. Recall that $Y_\Gamma = \Gamma \backslash \mathfrak{H}$, which is X_Γ for Γ cocompact, and X_Γ with cusps removed for Γ non-cocompact. Let G be the algebraic group GL_2 or \mathbf{B}^\times according as Γ non-cocompact or cocompact arising from an indefinite quaternion algebra B defined over \mathbf{Q} . In both cases, $G(\mathbf{R}) = \text{GL}_2(\mathbf{R})$ which has two connected components $G(\mathbf{R})_\pm$ consisting of elements with positive and negative determinants, respectively. Moreover, $G(\mathbf{R})_+$ acts transitively on the upper half plane \mathfrak{H} via fractional linear transformations so that we may identify \mathfrak{H} with $G(\mathbf{R})_+/SO_2Z(\mathbf{R})$, where Z denotes the center of G . Thus $G(\mathbf{R})/SO_2Z(\mathbf{R})$ may be identified with \mathfrak{H}^\pm , the disjoint union of upper and lower half plane. Let $G(\mathbf{Q})_+ = G(\mathbf{Q}) \cap G(\mathbf{R})_+$. Write \mathbb{A} for the ring of adèles over \mathbf{Q} , and \mathbb{A}_f the subring of finite adèles. The group $G(\mathbf{Q})$ is diagonally embedded in $G(\mathbb{A})$. Denote by $G(\mathbf{Q})_{+,f}$ the diagonal imbedding of $G(\mathbf{Q})_+$ in $G(\mathbb{A}_f)$.

Let U be a compact-open subgroup of $G(\mathbb{A}_f)$. Shimura's theory of canonical models gives a curve Y_U defined over \mathbf{Q} whose complex points can be expressed as

$$Y_U := G(\mathbf{Q}) \backslash \mathfrak{H}^\pm \times G(\mathbb{A}_f)/U = G(\mathbf{Q})_+ \backslash \mathfrak{H} \times G(\mathbb{A}_f)/U.$$

It has finitely many connected components, parametrized by the double cosets

$$G(\mathbf{Q})_{+,f} \backslash G(\mathbb{A}_f)/U = \coprod_{i=1}^h G(\mathbf{Q})_{+,f} x_i U$$

with $h = |\widehat{\mathbf{Z}}^\times / \det U|$. Note that an element $\gamma \in G(\mathbf{Q})_{+,f}$ satisfies $\gamma x_i U = x_i U$ if and only if γ lies in $\Gamma_{i,U} := G(\mathbf{Q})_{+,f} \cap x_i U x_i^{-1}$, a congruence subgroup of $\mathrm{SL}_2(\mathbf{R})$. Thus we get

$$(10) \quad Y_U = \prod_{i=1}^h \Gamma_{i,U} \backslash \mathfrak{H} = \prod_{i=1}^h Y_{\Gamma_{i,U}},$$

showing that each connected component of Y_U is the curve of a congruence subgroup of $\mathrm{SL}_2(\mathbf{R})$. Conversely, all congruence subgroups Γ of $\mathrm{SL}_2(\mathbf{R})$ are known to arise as $\Gamma = G(\mathbf{Q})_{+,f} \cap U$ for some compact open subgroup U of $G(\mathbb{A}_f)$. We call U the *(adelic) group corresponding to Γ* .

Definition 1. *A compact open subgroup U of $G(\mathbb{A}_f)$ is called neat if all $\Gamma_{i,U}$ defined above are torsion-free.*

Note that, when $h > 1$, the curve Y_U while defined over \mathbf{Q} is not absolutely irreducible over \mathbf{Q} . Its connected components $Y_{\Gamma_{i,U}}$ are absolutely irreducible over an abelian extension of \mathbf{Q} given by Shimura's theory of canonical models.

Adelically, we may regard

$$Y_U = G(\mathbf{Q}) \backslash G(\mathbf{R}) G(\mathbb{A}_f) / \mathrm{SO}_2 \mathbf{Z}(\mathbf{R}) U = G(\mathbf{Q}) \backslash G(\mathbb{A}) / \mathrm{SO}_2 \mathbf{Z}(\mathbf{R}) U.$$

Denote by $S_k(U)$ the space of automorphic forms on $G(\mathbb{A})$, cuspidal if $G = \mathrm{GL}_2$, right invariant under $UZ(\mathbf{R})$, whose components at the place ∞ lie in the space of discrete series representation of $G(\mathbf{R})$ of weight k . Note that it decomposes into the direct sum $\bigoplus_{i=1}^h S_k(\Gamma_{i,U})$.

In the elliptic modular cases there are canonical compactifications X_U of Y_U by adding cusps of $\Gamma_{i,U}$. For the quaternion cases, we set $X_U = Y_U$.

Finally, there are constructible \mathbf{Q}_ℓ sheaves $V^k(U)_\ell$ on X_U to be elaborated in the next subsection. For both cases Eichler-Shimura theory takes the shape:

Theorem 3 (Deligne [19], Ohta [50]). *For integers $k \geq 0$, the action of the Galois group $G_{\mathbf{Q}}$ of \mathbf{Q} on $H^1(X_U \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V^k(U)_\ell)$ is unramified at the primes $p \neq \ell$ where X_U has good reduction. Moreover, we have*

$$\mathrm{Tr}(\mathrm{Frob}_p \mid H^1(X_U \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V^k(U)_\ell)) = \mathrm{Tr}(T_p \mid S_{k+2}(U)).$$

Combined with the Grothendieck-Lefschetz trace formula, we can express the trace of T_p on $S_{k+2}(U)$ at primes p where X_U has good reduction as a sum over the \mathbf{F}_p -points on the reduced curve:

$$(11) \quad -\mathrm{Tr}(T_p \mid S_{k+2}(U)) = \sum_{\lambda \in X_U(\mathbf{F}_p)} \mathrm{Tr}(\mathrm{Frob}_\lambda \mid (V^k(U)_\ell)_\lambda).$$

Write $V^k(\Gamma)_\ell$ for the restriction of the sheaf $V^k(U)_\ell$ on X_Γ . In case that X_Γ is defined over \mathbf{Q} , we have $X_U = X_\Gamma$ and $S_{k+2}(U) = S_{k+2}(\Gamma)$. Then the above theorem gives a geometric description of the trace of the Hecke operator T_p on $S_{k+2}(\Gamma)$.

Now we explain the notation $(V^k(U)_\ell)_\lambda$ on the right side of (11). Let X_U° be the complement in X_U of the cusps and elliptic points. Then our sheaves $V^k(U)_\ell$ are lisse on the open set X_U° , equivalent therefore to a representation of $\pi_1(X_U^\circ, \bar{\eta})$ where $\bar{\eta}$ is the generic geometric point $\bar{\eta} = \mathrm{Spec}(\overline{\mathbf{Q}(X_U)})$ of X_U° . Thus there is a canonical epimorphism

$$G_\eta := \mathrm{Gal}(\overline{\mathbf{Q}(X_U)} / \mathbf{Q}(X_U)) \rightarrow \pi_1(X_U^\circ, \bar{\eta})$$

so that we may regard all lisse sheaves on X_U° as modules for the Galois group G_η . Every algebraic point $t \in X_U$ gives a discrete valuation of the function field $\mathbf{Q}(X_U)$ with residue field $\kappa(t)$, so we may speak of the decomposition D_t and inertia I_t subgroups (well-defined up to conjugation) and their images in $\pi_1(X_U^\circ, \bar{\eta})$. As $t : \mathrm{Spec}(\kappa(t)) \rightarrow X_U$ gives a morphism of étale sites, we get

$t^*V^k(U)_\ell$, a sheaf for the étale site on $\text{Spec}(\overline{\kappa(t)})$. This is equivalent to a module over $G_{\kappa(t)} := \text{Gal}(\overline{\kappa(t)}/\kappa(t)) \cong D_t/I_t$ for an algebraic closure $\overline{\kappa(t)}$ of $\kappa(t)$. In other words, the stalk $(V^k(U)_\ell)_{\bar{t}}$ is the \mathbf{Q}_ℓ -vector space $\bar{t}^*V^k(U)_\ell$ with $G_{\kappa(t)}$ action. Here $\bar{t} : \text{Spec}(\overline{\kappa(t)}) \rightarrow \text{Spec}(\kappa(t)) \rightarrow X_U$.

In particular, if t is in the open set X_U° , then the inertia I_t acts trivially. Also, by choosing an embedding $\kappa(\bar{t}) \subset \kappa(\bar{\eta})$, we get a homomorphism

$$G_{\kappa(t)} = \pi_1(\text{Spec}(\kappa(t)), \bar{t}) \rightarrow \pi_1(X_U^\circ, \bar{\eta})$$

and any representation of the right-hand side gives a representation of the left hand side. This describes the stalk $(V^k(U)_\ell)_{\bar{t}}$ for $t \in X_U^\circ$.

Moreover if $\iota : X_U^\circ \hookrightarrow X_U$ is the inclusion, then our sheaves also have the property that

$$V^k(U)_\ell = \iota_*(V^k(U)_\ell | X_U^\circ).$$

This implies that for every $t \in X_U$, the stalk $(V^k(U)_\ell)_{\bar{t}}$ is the \mathbf{Q}_ℓ -space $(V^k(U)_\ell)_{\bar{\eta}}^{I_t}$ with $G_{\kappa(t)}$ action.

The structure of these stalks will be examined in more detail in §2.2.3.

The curves X_U are defined over \mathbf{Q} . In fact, they have canonical reductions modulo all but finitely many primes p . This follows because of the interpretation of these curves as moduli spaces. That is, we may regard $f : X_U \rightarrow S$ where $S = \text{Spec}(\mathbf{Z}[1/N])$ for an integer $N \geq 1$, where f is smooth and projective. Moreover, the sheaves $V^k(U)_\ell$ extend to these integral models, and for each $i \geq 0$, the sheaves $R^i f_* \mathcal{F}_\ell$ are lisse on S and compatible with base-change. This implies that for each prime $p \nmid N$ we have an isomorphism

$$H^1(X_U \otimes \bar{\mathbf{F}}_p, V^k(U)_\ell \otimes \bar{\mathbf{F}}_p) \cong H^1(X_U \otimes \bar{\mathbf{Q}}, V^k(U)_\ell \otimes \bar{\mathbf{Q}}),$$

respecting Galois actions. Namely, a Frobenius $\text{Frob}_p \in G_{\mathbf{Q}}$ acts on the left-hand side as the (inverse of the) p th power endomorphism. The Grothendieck-Lefschetz trace formula expresses the Frobenius trace as a sum of local traces in the fibers of the points $X_U(\mathbf{F}_p)$.

For $\lambda \in X_U(\mathbf{F}_p)$ in (10), the stalk $(V^k(U)_\ell)_{\bar{\lambda}}$ refers to the stalk of $V^k(U)_\ell$ at any algebraic point t of X_U which reduces to λ modulo p .

2.2.2. As in the previous subsection, let G be GL_2 or \mathbf{B}^\times for an indefinite quaternion algebra B defined over \mathbf{Q} . Let U be an open compact subgroup of $G(\mathbb{A}_f)$. In this subsection, we discuss the ℓ -adic sheaves $V^k(U)_\ell$ on X_U , following the construction by Ohta in [50]. Since it is crucial to this paper, we sketch the key idea below.

Denote by $\varrho_k : \text{GL}_2 \rightarrow \text{GL}_{k+1}$ the morphism of algebraic groups over \mathbf{Z} which is k th symmetric power of $\varrho_1 : \text{GL}_2 \rightarrow \text{GL}_2$, the identity map. So for any commutative ring R , $\varrho_k(R) : \text{GL}_2(R) \rightarrow \text{GL}_{k+1}(R)$ sends the matrix $A \in \text{GL}_2(R)$ to the R -linear transformation $R^{k+1} \rightarrow R^{k+1}$ given by the matrix which is the k th symmetric power of A .

When $G = \mathbf{B}^\times$, choose a real quadratic field F which splits B , namely $B \otimes_{\mathbf{Q}} F = \text{M}_2(F)$, and fix an embedding $\theta : B \rightarrow \text{M}_2(F)$. Since B is an inner form of $\text{M}_2(F)$, there is an element $a \in \text{GL}_2(F)$ so that the embedded image can be described as

$$\theta(B) = \{x \in \text{M}_2(F) : a x a^{-1} = x^\tau\},$$

where τ is the nontrivial automorphism of F over \mathbf{Q} . Hence every $x \in \theta(B^\times)$ satisfies

$$\varrho_k(a)\varrho_k(x)\varrho_k(a)^{-1} = \varrho_k(x)^\tau, \quad \text{for all } k \geq 1.$$

Let

$$Y_k = \{x \in \text{M}_{k+1}(F) : \varrho_k(a) x \varrho_k(a)^{-1} = x^\tau\} \subset \text{M}_{k+1}(F).$$

It is a central simple algebra over \mathbf{Q} with an involution. By construction

$$Y_k \otimes_{\mathbf{Q}} F \cong \text{M}_{k+1}(F).$$

The invertible elements of Y_k define an algebraic group over \mathbf{Q} , denoted Y_k^\times . Note that $Y_k^\times(\mathbf{Q}) = Y_k^\times$ and $Y_k^\times(F) = (Y_k \otimes_{\mathbf{Q}} F)^\times \cong \mathrm{GL}_{k+1}(F)$. Let $j : Y_k^\times \hookrightarrow \mathrm{M}_{k+1}(F)^\times = \mathrm{GL}_{k+1}(F)$ be the natural inclusion. There is a morphism of algebraic groups $\varrho'_k : \mathbf{B}^\times \rightarrow Y_k^\times$ with the property that

$$j \circ \varrho'_k(\mathbf{Q}) : \mathbf{B}^\times(\mathbf{Q}) = B^\times \rightarrow Y_k^\times(\mathbf{Q}) = Y_k^\times \hookrightarrow \mathrm{GL}_{k+1}(F)$$

is θ restricted to B^\times followed by $\varrho_k(F)$, and $\varrho'_k(F)$ is $\varrho_k(F)$. Thus if we identify B^\times with $\theta(B^\times) \subset \mathrm{GL}_2(F)$, then we may regard $\varrho'_k(\mathbf{Q})$ as the restriction of $\varrho'_k(F)$ to B^\times .

As such, there are two possibilities for Y_k :

(I) $Y_k = \mathrm{M}_{k+1}(\mathbf{Q})$,

(II) $Y_k = \mathrm{M}_{(k+1)/2}(D)$ for a (division) quaternion algebra D over \mathbf{Q} .

For $k \geq 2$ even, Y_k falls in case (I) and ϱ'_k is a $(k+1)$ -dimensional representation of the algebraic group \mathbf{B}^\times .

In the second case, we can use the regular representation $r : D^\times \rightarrow \mathrm{GL}_4$ and define a representation $\mathbf{B}^\times \rightarrow \mathrm{GL}_{2(k+1)}$ by composing ϱ'_k with the embedding

$$\mathrm{M}_{(k+1)/2}(D) \rightarrow \mathrm{M}_{(k+1)/2}(\mathrm{M}_4(\mathbf{Q})) = \mathrm{M}_{2(k+1)}(\mathbf{Q}).$$

Note that $Y_1 = \theta(B) = D$ falls in case (II). So ϱ'_1 is 4-dimensional.

Fix a positive integer k . Let g be an element in $G(\mathbf{Q})$ such that $U \subset g \prod_p G(\mathbf{Z}_p)g^{-1}$. Then $\Gamma = G(\mathbf{Q})_+ \cap U$ is contained in $gG(\mathbf{Z})g^{-1}$. In the representation space of ϱ_k (resp. ϱ'_k) if $G = \mathrm{GL}_2$ (resp. $G = \mathbf{B}^\times$), choose a lattice L_k stable under $\varrho_k(gRg^{-1} \cap G(\mathbf{Q}))$ (resp. $\varrho'_k(gRg^{-1} \cap G(\mathbf{Q}))$), where $R = \mathrm{M}_2(\mathbf{Z})$ for $G = \mathrm{GL}_2$ and R is a maximal order O_B for $G = \mathbf{B}^\times$.

Fix a prime ℓ . Define a tower of curves

$$\dots \rightarrow Y_{U_3} \rightarrow Y_{U_2} \rightarrow Y_{U_1} \rightarrow Y_{U_0} = Y_U$$

where

$$Y_n = U \cap g(\{x \in G(\mathbf{Z}_\ell) : x \equiv id \pmod{\ell^n}\} \prod_{p \neq \ell} G(\mathbf{Z}_p))g^{-1}.$$

Then each Y_{U_n} is a finite Galois covering of Y_U with Galois group

$$\mathrm{Aut}(Y_{U_n}/Y_U) = U/(Z(\mathbf{Q}) \cap U)U_n.$$

As a compact subgroup of the discrete group $Z(\mathbf{Q})$, $Z(\mathbf{Q}) \cap U$ is contained in $\{\pm id\}$ for U in general, and is trivial for U neat. It follows from our choice of g that the finite group $U/(Z(\mathbf{Q}) \cap U)U_n$ can be represented by elements in $gRg^{-1} \cap G(\mathbf{Q})$. The constant group-scheme $Y_{U_n} \times L_k/\ell^n L_k$ has an action of $U/(Z(\mathbf{Q}) \cap U)U_n$. On the first factor this is the Galois action resulting from the theory of canonical models; the action on the second factor is from ϱ_k or ϱ'_k as appropriate.

Assume U is neat as in Definition 1. The quotient

$$V_n(U, L_k) := (U/(Z(\mathbf{Q}) \cap U)U_n) \backslash (Y_{U_n} \times L_k/\ell^n L_k)$$

is a finite étale group-scheme over Y_U . These form a projective system with n . Taking projective limit yields a lisse \mathbf{Z}_ℓ -sheaf $V(U, L_k)$ on Y_U . Tensoring this with \mathbf{Q}_ℓ , we obtain a lisse \mathbf{Q}_ℓ -sheaf on Y_U independent of the choice of L_k . The sheaf $V^k(U)_\ell$ on X_U is the pushforward of this sheaf on Y_U via the inclusion map $\iota : Y_U \hookrightarrow X_U$. Recall that ι is surjective for $G = \mathbf{B}^\times$, and for $G = \mathrm{GL}_2$, the complement $X_U \setminus Y_U$ is the set of cusps. Restricting $V^k(U)_\ell$ to X_Γ gives the sheaf $V^k(\Gamma)_\ell$.

When the group U is neat, the curve Y_U admits a moduli interpretation, namely there is a family of abelian varieties $f : A_U \rightarrow Y_U$ defined over \mathbf{Q} such that each fiber is an elliptic curve for $G = \mathrm{GL}_2$, and a polarized abelian surface for $G = \mathbf{B}^\times$. The sheaves defined above can be expressed geometrically as well. As shown in [51, Theorem 3.3.3],

$$\varprojlim_n V_n(U, L_1) \otimes_{\mathbf{Z}_\ell} \mathbf{Q}_\ell \simeq R^1 f_* \mathbf{Q}_\ell$$

and hence we can make the identification

$$V^1(U)_\ell = \iota_* R^1 f_* \mathbf{Q}_\ell, \quad \iota : Y_U \hookrightarrow X_U.$$

Therefore $V^1(U)_\ell$ is a rank-2 or 4 sheaf according as $G = \mathrm{GL}_2$ or $G = \mathbf{B}^\times$. In the elliptic modular case, since $\varrho_k = \mathrm{Sym}^k(\varrho_1)$, we have

$$V^k(U)_\ell = \iota_* \mathrm{Sym}^k R^1 f_* \mathbf{Q}_\ell,$$

as given by Deligne [19]. Thus $V^k(U)_\ell$ has rank $k + 1$. For the quaternion case, as discussed before, $V^k(U)_\ell$ has rank $k + 1$ for k even, but the relation between $V^k(U)_\ell$ and $R^1 f_* \mathbf{Q}_\ell$ is less clear. Proposition (4.3.1) of [51] shows that $V^k(U)_\ell$ occurs as a component of the sheaf $R^w f_*^n \mathbf{Q}_\ell$ for suitable choices of w and n , whose Galois module structure can be expressed as

$$R^w f_*^n \mathbf{Q}_\ell = \bigwedge^w R^1 f_* \mathbf{Q}_\ell = \bigwedge^w (R^1 f_* \mathbf{Q}_\ell)^n,$$

in which the first equality follows from the structure of the cohomology of abelian varieties and the second from the Künneth formula. From this one sees that all the sheaves $R^w f_*^n \mathbf{Q}_\ell$ are calculated from tensor operations from the fundamental $V^1(U)_\ell$. We will see an expression for $V^{2k}(U)_\ell$ similar to the elliptic modular case in Proposition 3.

The sheaf $V^k(U)_\ell$ defined above is for U neat. For a general U , there is a neat compact open normal subgroup U' of U (such as U_n above for n large) for which $V^k(U')_\ell$ is defined. It follows from the definition of $V^k(U')_\ell$ that its stalk at any algebraic point of $Y_{U'}$ is $(\varprojlim_n L_k / \ell^n L_k) \otimes_{\mathbf{Z}_\ell} \mathbf{Q}_\ell$, on which $U / (\mathbf{Z}(\mathbf{Q}) \cap U) U'$ acts since by construction the lattice L_k is invariant under the action of ϱ_k or ϱ'_k on $gRg^{-1} \cap G(\mathbf{Q})$ and the quotient $U / (\mathbf{Z}(\mathbf{Q}) \cap U) U'$ can be represented by elements in $gRg^{-1} \cap G(\mathbf{Q})$. We say that $V^k(U')_\ell$ admits $U / (\mathbf{Z}(\mathbf{Q}) \cap U) U'$ -structure. This allows us to define $V^k(U)_\ell$ by pushforward from $V^k(U')_\ell$ as described below. Here k is even if $\mathbf{Z}(\mathbf{Q}) \cap U = \{\pm \mathrm{id}\}$ so that $\varrho_k(-\mathrm{id}) = (-\mathrm{id})^k = \mathrm{id}$.

As before, let X_U° be the open subset of X_U with cusps and elliptic points removed. Denote by $T_{U'}$ the preimage of X_U° under the covering map $p : X_{U'} \rightarrow X_U$. Note that $T_{U'} \subset X_{U'}^\circ$. Let $q : T_{U'} \rightarrow X_U^\circ$ be the restriction of p to $T_{U'}$. Then q is finite étale. When U is neat, we have $X_U^\circ = Y_U$ and $T_{U'} = Y_{U'} = X_{U'}^\circ$. The sheaf $V^k(U)_\ell$ restricted to Y_U defined before is the pushforward of $V^k(U')$ restricted to $Y_{U'}$ via the covering map $q : Y_{U'} \rightarrow Y_U$. Hence, as a generalization from neat to general U , we pushforward the restriction of $V^k(U')_\ell$ to $T_{U'}$ via q to get a lisse sheaf \mathcal{F}_ℓ on X_U° , and define the sheaf $V^k(U)_\ell$ on X_U as the pushforward of \mathcal{F}_ℓ via the inclusion $\iota : X_U^\circ \rightarrow X_U$. In other words, $V^k(U)_\ell = \iota_* q_*^{U / (\mathbf{Z}(\mathbf{Q}) \cap U) U'} (V^k(U')_\ell | T_{U'})$. Note that the sheaf \mathcal{F}_ℓ on X_U° is independent of the choice of U' , hence so is $V^k(U)_\ell$.

Lemma 1. *The sheaf $V^k(U)_\ell$ on X_U defined above agrees with the pushforward $p_*^{U / (\mathbf{Z}(\mathbf{Q}) \cap U) U'} V^k(U')_\ell$ via the covering map $p : X_{U'} \rightarrow X_U$. Here k is even if $\mathbf{Z}(\mathbf{Q}) \cap U = \{\pm \mathrm{id}\}$.*

Proof. By construction, this is true at all points of X_U° . Let t be a point of $X_U \setminus X_U^\circ$ and t' a point of $X_{U'}$ lying over it. Because U' is a normal subgroup of U with finite index, p is a Galois cover with Galois group $U / (\mathbf{Z}(\mathbf{Q}) \cap U) U'$, which acts transitively on each fiber of p . Denote by $\bar{\eta}$ and $\bar{\eta}'$ the generic geometric points of X_U and $X_{U'}$ respectively. Note that

$$U / (\mathbf{Z}(\mathbf{Q}) \cap U) U' \cong \mathrm{Aut}(X_{U'} / X_U).$$

Also, because we defined $V^k(U)_\ell$ via ι , and $V^k(U)_\ell$ is lisse on X_U° , we have

$$(V^k(U)_\ell)_{\bar{t}} = (V^k(U)_\ell)_{\bar{\eta}}^{I_{\bar{t}}}.$$

Similarly, we also have

$$(V^k(U')_\ell)_{\bar{t}'} = (V^k(U')_\ell)_{\bar{\eta}'}^{I_{t'}}.$$

Here I_t and $I_{t'}$ are the inertia subgroups of the decomposition groups D_t and $D_{t'}$ at t and t' , respectively. The stabilizer of t' in $U/(Z(\mathbf{Q}) \cap U)U'$ is $D_t/D_{t'}$ with the inertia subgroup $I(t'/t) = I_t/I_{t'}$. Since the stalk at t of $p_*^{U/(Z(\mathbf{Q}) \cap U)U'} V^k(U')_\ell$ is $((V^k(U')_\ell)_{\bar{t}'})^{I(t'/t)}$ the result now follows from

$$((V^k(U')_\ell)_{\bar{\eta}'}^{I_{t'}})^{I_t/I_{t'}} = ((V^k(U')_\ell)_{\bar{\eta}'}^{I_t})^{I_t/I_{t'}} = (V^k(U)_\ell)_{\bar{\eta}}^{I_t}.$$

□

Observe that the sheaf $V^k(U)_\ell$ defined above is compatible with taking the first cohomology.

Proposition 1. *With the above notation and assumption on the parity of k we have*

$$H^1(X_{U'} \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V^k(U')_\ell)^{U/(Z(\mathbf{Q}) \cap U)U'} = H^1(X_U \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, p_*^{U/(Z(\mathbf{Q}) \cap U)U'} V^k(U')_\ell) = H^1(X_U \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V^k(U)_\ell).$$

Proof. Because \mathbf{Q}_ℓ is flat over \mathbf{Z}_ℓ it follows from the definition of $V^k(U')_\ell$ that

$$H^1(X_{U'} \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V^k(U')_\ell)^{U/(Z(\mathbf{Q}) \cap U)U'} = H^1(X_{U'} \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V(U', L_k))^{U/(Z(\mathbf{Q}) \cap U)U'} \otimes_{\mathbf{Z}_\ell} \mathbf{Q}_\ell.$$

An elementary argument shows that

$$\begin{aligned} H^1(X_{U'} \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V(U', L_k))^{U/(Z(\mathbf{Q}) \cap U)U'} &= (\varinjlim_n H^1(X_{U'} \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V_n(U', L_k)))^{U/(Z(\mathbf{Q}) \cap U)U'} \\ &= \varinjlim_n H^1(X_{U'} \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V_n(U', L_k))^{U/(Z(\mathbf{Q}) \cap U)U'}, \end{aligned}$$

so the problem is reduced to prove that

$$H^1(X_{U'} \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V_n(U', L_k))^{U/(Z(\mathbf{Q}) \cap U)U'} \cong H^1(X_U \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, p_*^{U/(Z(\mathbf{Q}) \cap U)U'} V_n(U', L_k)).$$

For ℓ prime to the order of $U/(Z(\mathbf{Q}) \cap U)U'$, the higher direct images $R^i p_*^{U/(Z(\mathbf{Q}) \cap U)U'}(\mathcal{F}) = 0$ for $i > 0$ and any sheaf \mathcal{F} of \mathbf{Z}/ℓ^n -modules. A spectral sequence argument then gives the result (in the case of ordinary sheaf cohomology, see [29, ch. V, cor. of prop. 5.2.3, and th. 5.3.1]). □

In [50] Ohta proved Theorem 3 for U neat using the sheaf $V^k(U)_\ell$. For a general U , he chose a neat compact open normal subgroup U' of U for which Theorem 3 holds, then took the $U/(Z(\mathbf{Q}) \cap U)U'$ invariant part of both sides to show that the trace of the Hecke operator T_p on $S_{k+2}(U)$ is equal to the trace of Frob_p on $H^1(X_{U'} \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V^k(U')_\ell)^{U/(Z(\mathbf{Q}) \cap U)U'}$. Proposition 1 above implies that $H^1(X_{U'} \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V^k(U')_\ell)^{U/(Z(\mathbf{Q}) \cap U)U'}$ indeed is equal to $H^1(X_U \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, V^k(U)_\ell)$.

Remark 1. When the group Γ is a finite index subgroup of O_B^1 for a split or nonsplit indefinite quaternion algebra B over \mathbf{Q} , the group of Atkin-Lehner involutions is defined as the quotient $\Gamma^*/(\{\pm id\}\Gamma)$, where Γ^* is the normalizer of Γ in the group of elements in O_B with positive reduced norm. It is a finite elementary 2-group. A split B corresponds to $G = \text{GL}_2$. When Γ is $\Gamma_0(N)$ or $\Gamma_1(N)$, these are the familiar Atkin-Lehner operators W_Q in the literature (cf. [6, 7]), where Q is the highest power of a prime p dividing N . A nonsplit B corresponds to $G = \mathbf{B}^\times$. When $\Gamma = O_B^1$, the group $\Gamma^*/(\{\pm id\}\Gamma)$ is generated by r elements, which can be chosen to be any element in O_B with reduced norm p at the r places p where B ramifies (see §28.9 of [65]). More discussions are given in §5.1. We shall regard the Atkin-Lehner involutions as elements in $G(\mathbb{A}_f)$.

Let U be the open compact subgroup of $G(\mathbb{A}_f)$ corresponding to Γ and let U'' be the group generated by U and some Atkin-Lehner operators of Γ . Let U' be a neat normal open compact subgroup of U . Then the same argument as before shows that the sheaf $V^k(U')_\ell$ admits $U''/(Z(\mathbf{Q}) \cap U'')U'$ -structure so that the sheaf $V^k(U'')_\ell$ on $X_{U''}$ can be defined as the pushforward $q_*^{U''/(Z(\mathbf{Q}) \cap U'')U'} V^k(U')_\ell$ via the projection $q : X_{U'} \rightarrow X_{U''}$.

For U'' defined above, when the curve $X_{U''}$ is defined over \mathbf{Q} , Theorem 3 also holds for U'' .

2.2.3. The goal of this subsection is to investigate the Galois structure of the stalks of $V^k(U)_\ell$. We begin by proving a general statement for the quaternion case inspired by Theorem 3.1.2 of [8].

Proposition 2. *Let B be a quaternion algebra over \mathbf{Q} and K an extension of \mathbf{Q} such that $B_K = B \otimes_{\mathbf{Q}} K \cong M_2(K)$. Let V_K be a 4-dimensional K -vector space which is a principal B_K -module. Suppose a group H acts on V_K via the representation ρ_K which commutes with the B_K -action. Then, with the standard action of $M_2(K)$ on the 2-dimensional K -vector space K^2 , we have*

(1)

$$W_K := \text{Hom}_{M_2(K)}(K^2, V_K) = \text{Hom}_{B_K}(K^2, V_K)$$

is a 2-dimensional K -vector space endowed with the H -action via the representation σ_K induced from the H -action on V_K ;

(2) $\rho_K \cong \sigma_K \oplus \sigma_K$ as H -modules, and $\rho_K \cong \sigma_K \otimes \varrho_1(K)$ as $H \times \text{GL}_2(K)$ -modules.

Proof. The induced H -action on $W_K = \text{Hom}_{M_2(K)}(K^2, V_K)$ is given by the rule

$$(\sigma_K(h)f)(z) = \rho_K(h)(f(z)) \quad \text{for } h \in H, f \in W_K, \text{ and } z \in K^2.$$

We verify that $\sigma_K(h)f$ lies in W_K . Indeed, for $m \in M_2(K)$ and $z \in K^2$, we have

$$(\sigma_K(h)f)(mz) = \rho_K(h)(f(mz)) = \rho_K(h)(mf(z)) = m(\rho_K(h)(f(z))) = m(\sigma_K(h)f)(z)$$

due to the assumption that $\rho_K(h)$ preserves the B_K -action. This makes the canonical evaluation map

$$(*) \quad K^2 \otimes_K \text{Hom}_{M_2(K)}(K^2, V_K) \rightarrow V_K \quad \text{given by } (z, f) \mapsto f(z)$$

H -equivariant.

Next we show that $W_K = \text{Hom}_{M_2(K)}(K^2, V_K)$ is a 2-dimensional K -vector space. Since V_K is a principal $M_2(K)$ -module, denote it by $V_K = M_2(K)v$ for some nonzero $v \in V_K$ with the $M_2(K)$ -action given by left multiplication. Write the elements in K^2 as a column vector so that $M_2(K)$ acts on K^2 by the matrix product and $K^2 = M_2(K) \begin{pmatrix} 1 \\ 0 \end{pmatrix}$. Hence an element $f \in W_K$

is determined by $f \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix} \right) = \begin{pmatrix} a & b \\ c & d \end{pmatrix} v$ for some $a, b, c, d \in K$. As the stabilizer of $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ is the set $\left\{ \begin{pmatrix} 1 & x \\ 0 & y \end{pmatrix} : x, y \in K \right\}$, it follows that $\begin{pmatrix} 1 & x \\ 0 & y \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} v = f \left(\begin{pmatrix} 1 & x \\ 0 & y \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right) = f \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix} \right) =$

$\begin{pmatrix} a & b \\ c & d \end{pmatrix} v$ for all $x, y \in K$, which in turn implies $c = d = 0$. Therefore $f \mapsto \begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix}$ is a K -linear bijection between W_K and K^2 . We record this correspondence by writing $f_{(a,b)}$ for f . This proves that W_K is a 2-dimensional K -vector space on which H acts via σ_K , which is assertion (1). Notice

that $f_{(a,b)} \left(\begin{pmatrix} x & y \\ z & w \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right) = \begin{pmatrix} ax & by \\ az & bz \end{pmatrix} v$.

Observe that $K^2 \otimes_K W_K \cong W_K \oplus W_K$. Finally we prove the surjectivity of the evaluation map (*), from which the first assertion in (2) will follow. To see this, take any element $\begin{pmatrix} x & y \\ z & w \end{pmatrix} v \in V_K$ and note that

$$\begin{pmatrix} x & y \\ z & w \end{pmatrix} v = \begin{pmatrix} x & 0 \\ z & 0 \end{pmatrix} v + \begin{pmatrix} 0 & y \\ 0 & w \end{pmatrix} v = f_{(1,0)} \left(\begin{pmatrix} x & y \\ z & w \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right) + f_{(0,1)} \left(\begin{pmatrix} y & x \\ w & z \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right).$$

For the second assertion, observe that $\sigma_K \oplus \sigma_K = \sigma_K \otimes (1_K \oplus 1_K) = \sigma_K \otimes \varrho_1(K)$ as an $H \times \text{GL}_2(K)$ -module. \square

The main applications of the above proposition are to the following settings in which Γ arises from an indefinite quaternion algebra B defined over \mathbf{Q} :

- (a) $K = \mathbf{C}$ and Γ cocompact and torsion-free. Denote by A_z the fiber at z of the universal family of 2-dimensional abelian varieties over a modular curve X_Γ . Then we have a representation of $H = \pi_1(X_\Gamma, z)$ on A_z . It commutes with the action of the quaternions B_K on $V_K = H^1(A_z, \mathbf{C})$. It is known that V_K is a principal B_K -module. This gives

Corollary 1. *Assume that Γ is cocompact and torsion-free. Then*

1. $R^1 f_* \mathbf{C}$ is, in many ways, isomorphic to a direct sum $\sigma_{\Gamma, \mathbf{C}} \oplus \sigma_{\Gamma, \mathbf{C}}$ for a rank 2 local system $\sigma_{\Gamma, \mathbf{C}}$ on X_Γ° .
- 2.

$$V^k(\Gamma)_{\mathbf{C}} = \text{Sym}^k(\sigma_{\Gamma, \mathbf{C}}).$$

- (b) Same as above, except that we consider the universal abelian variety over X_U where U is the compact open subgroup of $G(\mathbb{A}_f)$ corresponding to Γ , z is a point of X_U , H is Grothendieck's $\pi_1(X_U, \bar{z})$, the field K is an extension of \mathbf{Q}_ℓ which splits B , and $V_K = H^1(A_{\bar{z}}, K)$. Then σ_K defines a local system of K -vector spaces of dimension 2 on X_U for the étale topology.
- (c) For the same K as in (b), this time we consider $V_K = H^1(A_{\bar{z}}, K)$ as an $H = \text{Gal}(\bar{\mathbf{Q}}/F)$ -module for a number field F over which the fiber $A_{\bar{z}}$ is defined, $F \otimes \mathbf{Q}_\ell$ contains K , and the actions of B_K and H on V_K commute. This is the representation ρ_K in Proposition 2. See more details below. An explicit description of the 2-dimensional representation σ_K occurring in $\rho_K = \sigma_K \oplus \sigma_K$ for the groups Γ in Theorem 1 is provided in Proposition 5.

Now we examine the structure of the stalks of $V^k(U)_\ell$ at algebraic points in X_U° . To ease our notation, we shall write $V^k(U)_{\bar{\lambda}, \ell}$ for the stalk $(V^k(U)_\ell)_{\bar{\lambda}}$ and denote by $\rho^k(U)_{\lambda, \ell}$ the Galois representation on this stalk.

We first study the case $G = \mathbf{B}^\times$ and U not containing $-\text{id}$. Let $\lambda \in X_U^\circ(L)$ for a number field L . Since $-\text{id}$ is not in U , $V^1(U)_\ell$ is defined as the pushforward of the sheaf $V^1(U')_\ell$ on $X_{U'}$ for a neat normal open compact subgroup U' of U via the covering map $X_{U'} \rightarrow X_U$. Thus the stalk of $V^1(U)_\ell$ at $\lambda \in X_U^\circ(L)$ is the same \mathbf{Q}_ℓ -space as the stalk of $V^1(U')_\ell$ at any point $\lambda' \in X_{U'}^\circ$ above λ . Here λ' is rational over a finite extension L' of L . As remarked before, Ohta proved that $V^1(U')_\ell = R^1 f_* \mathbf{Q}_\ell$ for a family of abelian surfaces $f : A_{U'} \rightarrow X_{U'}$. Moreover, the action of B on each fiber commutes with the Galois action. Hence B acts on $V^1(U)_{\bar{\lambda}, \ell}$ and commutes with the action of a finite index subgroup $G_{L'}$ of G_L .

Let E be a real quadratic extension of \mathbf{Q} which splits B . Fix a prime \mathfrak{l} of E above ℓ . Then Proposition 2 applied to $K = E_{\mathfrak{l}}$, the completion of E at \mathfrak{l} , $H = G_{L'}$, and $\rho_K = \rho^1(U)_{\lambda, \ell} \otimes_{\mathbf{Q}_\ell} E_{\mathfrak{l}}$ shows that, as a $G_{L'}$ -module, $\rho^1(U)_{\lambda, \ell} \otimes E_{\mathfrak{l}}$ decomposes as

$$\rho^1(U)_{\lambda, \ell} \otimes_{\mathbf{Q}_\ell} E_{\mathfrak{l}} = \sigma_{\lambda, \mathfrak{l}} \oplus \sigma_{\lambda, \mathfrak{l}} = \sigma_{\lambda, \mathfrak{l}} \otimes (1_{G_{L'}} \oplus 1_{G_{L'}}).$$

for a canonical representation $\sigma_{\lambda, \mathfrak{l}}$ of $G_{L'}$ on $E_{\mathfrak{l}}^2$.

We proceed to obtain a tensor product decomposition over G_F . When $\rho^1(U)_{\lambda, \ell} \otimes E_{\mathfrak{l}}$ is irreducible, using Clifford theory in the form of Theorem 2.3 of [44] we know that there are two 2-dimensional \mathfrak{l} -adic representations $\eta_{\lambda, \mathfrak{l}}$ and $\gamma_{\lambda, \mathfrak{l}}$ of G_F , where $\eta_{\lambda, \mathfrak{l}}$, up to a finite order character, is an extension of $\sigma_{\lambda, \mathfrak{l}}$ and $\gamma_{\lambda, \mathfrak{l}}$ has finite image, such that

$$(12) \quad \rho^1(U)_{\lambda, \ell} \otimes E_{\mathfrak{l}} \cong \eta_{\lambda, \mathfrak{l}} \otimes \gamma_{\lambda, \mathfrak{l}} \quad \text{as } G_F \text{ - modules.}$$

This also holds when $\rho^1(U)_{\lambda, \ell} \otimes E_{\mathfrak{l}}$ is reducible, for in this case, $\rho^1(U)_{\lambda, \ell} \otimes E_{\mathfrak{l}}$ is the sum of two degree-2 representations of G_F whose restrictions to $G_{L'}$ are equal to $\sigma_{\lambda, \mathfrak{l}}$. Thus they differ by a finite character ξ of G_F trivial on $G_{L'}$. We may take one component to be $\eta_{\lambda, \mathfrak{l}}$ and $\gamma_{\lambda, \mathfrak{l}} = 1 \oplus \xi$.

Remark 2. As an extension of $\sigma_{\lambda, \mathfrak{l}}$, the representation $\eta_{\lambda, \mathfrak{l}}$ in (12) is unique up to twisting by a finite order character $\mu_{\mathfrak{l}}$ of G_F , in which case $\gamma_{\lambda, \mathfrak{l}}$ is twisted by $\mu_{\mathfrak{l}}^{-1}$.

It follows from (12) that ²

$$(13) \quad \bigwedge^2 \rho^1(U)_{\lambda,\ell} \otimes E_{\mathfrak{l}} \cong (\mathrm{Sym}^2 \eta_{\lambda,\mathfrak{l}} \otimes \det \gamma_{\lambda,\mathfrak{l}}) \oplus (\det \eta_{\lambda,\mathfrak{l}} \otimes \mathrm{Sym}^2 \gamma_{\lambda,\mathfrak{l}})$$

as a sum of two degree-3 G_F -modules. In particular, $(\mathrm{Sym}^2 \eta_{\lambda,\mathfrak{l}} \otimes \det \gamma_{\lambda,\mathfrak{l}}) |_{G'_L}$ is a subrepresentation of $\sigma_{\lambda,\mathfrak{l}} \otimes \sigma_{\lambda,\mathfrak{l}}$. By Proposition 2 (2), $\rho^1(U)_{\lambda,\ell} \otimes E_{\mathfrak{l}}$ is also a $\mathrm{GL}_2(E_{\mathfrak{l}})$ -module. To see the $\mathrm{GL}_2(E_{\mathfrak{l}})$ -module structure of $\bigwedge^2 \rho^1(U)_{\lambda,\ell} \otimes E_{\mathfrak{l}}$, we regard (13) as $G_{L'}$ -modules. As such, on the right hand side, the first factor is $\mathrm{Sym}^2(\sigma_{\lambda,\mathfrak{l}} |_{G_{L'}}) \otimes 1_{E_{\mathfrak{l}}}$, which consists of three copies of a 1-dimensional $\mathrm{GL}_2(E_{\mathfrak{l}})$ -module, while the second factor reads $\det(\sigma_{\lambda,\mathfrak{l}} |_{G_{L'}}) \otimes \mathrm{Sym}^2(\varrho_1(E_{\mathfrak{l}}))$, which is a degree-3 irreducible $\mathrm{GL}_2(E_{\mathfrak{l}})$ -module.

On the other hand, by Proposition (4.3.1) and the proof of Proposition (4.4.2) of [51],

$$\bigwedge^2 \rho^1(U)_{\lambda,\ell} \otimes E_{\mathfrak{l}} = \bigwedge^2 \rho^1(U')_{\lambda',\ell} \otimes E_{\mathfrak{l}} \cong (R^2 f_* \mathbf{Q}_{\ell})_{\bar{\lambda}'} \otimes E_{\mathfrak{l}} \cong (V^2(U')_{\bar{\lambda}',\ell} \otimes E_{\mathfrak{l}}) \oplus E_{\mathfrak{l}}(-1)^3$$

as $G_{L'}$ -modules, where $E_{\mathfrak{l}}(-1)$ denotes the Tate twist of the constant sheaf $E_{\mathfrak{l}}$ by -1 . As $\mathrm{GL}_2(E_{\mathfrak{l}})$ -modules, on the right side, the first factor is of degree-3 irreducible generically, while the second factor is three copies of a 1-dimensional module. This shows that $V^2(U)_{\bar{\lambda},\ell} \otimes E_{\mathfrak{l}}$ and $\mathrm{Sym}^2 \eta_{\lambda,\mathfrak{l}} \otimes \det \gamma_{\lambda,\mathfrak{l}}$ agree as $G_{L'}$ -modules, and hence as G_F -modules, they agree up to a (finite order) character of G_F trivial on $G_{L'}$. Thus there is a finite order character $\chi_{\lambda,\mathfrak{l}}$ of G_F such that the representation $\rho^2_{\lambda,\ell} \otimes E_{\mathfrak{l}}$ of G_F on $V^2(U)_{\bar{\lambda},\ell} \otimes E_{\mathfrak{l}}$ is

$$(14) \quad \rho^2(U)_{\lambda,\ell} \otimes E_{\mathfrak{l}} = \chi_{\lambda,\mathfrak{l}} \otimes \mathrm{Sym}^2 \eta_{\lambda,\mathfrak{l}}.$$

We show that this is a general phenomenon.

Proposition 3. *Let U be a group generated by an open compact subgroup of $G(\mathbb{A}_f)$ and a subset of the Atkin-Lehner operators. Suppose $-\mathrm{id} \notin U$. Let λ be an algebraic point of X_U° .*

(1) *Suppose $G = B^{\times}$ for an indefinite quaternion algebra B defined over \mathbf{Q} . Let E be a real quadratic field that splits B . Fix a prime ℓ and a prime \mathfrak{l} of E above ℓ . Suppose the 4-dimensional ℓ -adic representation $\rho^1(U)_{\lambda,\ell}$ on the stalk $V^1(U)_{\bar{\lambda},\ell}$ is a G_F -module for some number field F . Then there is a 2-dimensional \mathfrak{l} -adic representation $\eta_{\lambda,\mathfrak{l}}$ and a finite order character $\chi_{\lambda,\mathfrak{l}}$ of G_F such that for all integers $k \geq 1$, the representation $\rho^{2k}(U)_{\lambda,\ell} \otimes_{\mathbf{Q}_{\ell}} E_{\mathfrak{l}}$ on the fiber $V^{2k}(U)_{\bar{\lambda},\ell} \otimes_{\mathbf{Q}_{\ell}} E_{\mathfrak{l}}$ is given by*

$$(15) \quad \rho^{2k}(U)_{\lambda,\ell} \otimes_{\mathbf{Q}_{\ell}} E_{\mathfrak{l}} = \chi_{\lambda,\mathfrak{l}}^k \otimes \mathrm{Sym}^{2k}(\eta_{\lambda,\mathfrak{l}})$$

as G_F -modules. Moreover, $\chi_{\lambda,\mathfrak{l}} \det(\eta_{\lambda,\mathfrak{l}}) = \epsilon_{\mathfrak{l}}$, where $\epsilon_{\mathfrak{l}}$ denotes the \mathfrak{l} -adic cyclotomic character.

(2) *Suppose $G = \mathrm{GL}_2$. Then for all integers $k \geq 1$,*

$$(16) \quad \rho^k(U)_{\lambda,\ell} = \mathrm{Sym}^k(\rho^1(U)_{\lambda,\ell}),$$

and $\det \rho^1(U)_{\lambda,\ell}$ is the cyclotomic character ϵ_{ℓ} .

In both cases, the determinant of the representation is the cyclotomic character raised to the degree of the representation.

Proof. Let ρ be a degree-2 characteristic zero representation of a group. We have, for $m \geq 4$,

$$(17) \quad \mathrm{Sym}^{m-2}(\rho) \otimes \mathrm{Sym}^2(\rho) \cong \mathrm{Sym}^m(\rho) \oplus (\det(\rho) \otimes \mathrm{Sym}^{m-2}(\rho)) \oplus (\det(\rho)^2 \otimes \mathrm{Sym}^{m-4}(\rho))$$

by checking the characters of the representations or the Clebsch Gordan coefficient formula.

²Assume V and W both 2-dimensional G -modules with a basis $\{v_1, v_2\}, \{w_1, w_2\}$ respectively. Then $V \otimes W$ has a basis $\{v_i \otimes w_j\}_{1 \leq i, j \leq 2}$ so that $\bigwedge^2(V \otimes W)$ has a basis $\{(v_{i_1} \otimes w_{j_1}) \wedge (v_{i_2} \otimes w_{j_2}) \mid i_1 \neq i_2 \text{ or } j_1 \neq j_2\}$. The vectors $(v_i \otimes w_1) \wedge (v_i \otimes w_2)$ with $i = 1, 2$ together with $(v_1 \otimes w_1) \wedge (v_2 \otimes w_2) + (v_2 \otimes w_1) \wedge (v_1 \otimes w_2)$ span a sub- G -module, which is isomorphic to $\mathrm{Sym}^2 V \otimes \det W$. Switching the roles of V and W will give the other one. It is independent of the choice of $\eta_{\lambda,\mathfrak{l}}$ in (12) by the remark above.

Case (1) $G = \mathbf{B}^\times$. Applying (17) to the representation $\rho = \varrho_1(E)$ of $\mathrm{GL}_2(E)$, we obtain the canonical decomposition

$$\varrho_{m-2}(E) \otimes \varrho_2(E) \cong \varrho_m(E) \oplus (\det \varrho_1(E) \otimes \varrho_{m-2}(E)) \oplus ((\det \varrho_1(E))^2 \otimes \varrho_{m-4}(E)).$$

As discussed in §2.2.2, the representations $\varrho'_m(\mathbf{Q})$ of B^\times for $m \geq 2$ even and $m = 1$ may be regarded as the restriction of $\varrho_m(E)$ to B^\times under the identification $B^\times = \theta(B^\times) \subset \mathrm{GL}_2(E)$. Thus the above identity holds with $\varrho'_j(E)$ replacing $\varrho_j(\mathbf{Q})$, which in turn gives rise to the following identity on sheaves

$$(18) \quad V^{2k-2}(U)_\ell \otimes V^2(U)_\ell \cong V^{2k}(U)_\ell \oplus (\mathbf{Q}_\ell(-1) \otimes V^{2k-2}(U)_\ell) \oplus (\mathbf{Q}_\ell(-2) \otimes V^{2k-4}(U)_\ell)$$

by the proof of Proposition (4.4.2) in [51] since the ℓ -adic rank-1 sheaf arising from $\det(\varrho'_1(\mathbf{Q}))$ is the Tate twist by -1 of the constant sheaf \mathbf{Q}_ℓ on which the Galois group acts by the cyclotomic character ϵ_ℓ . Also, $V^0(U)_\ell$ is the constant sheaf \mathbf{Q}_ℓ on X_U . The same holds when each sheaf is tensored with E_t .

We now prove (15) by induction on k . When $k = 1$, this is (14) proved above. Assume the statement holds for $2k - 2$, that is, as G_L -modules,

$$V^{2k-2}(U)_{\bar{\lambda}, \ell} \otimes E_t \cong \chi_{\lambda, t}^{k-1} \otimes \mathrm{Sym}^{2k-2}(\eta_{\lambda, t}).$$

The two surjective G_F -homomorphisms

$$(V^{2k-2}(U)_{\bar{\lambda}, \ell} \otimes E_t) \otimes (V^2(U)_{\bar{\lambda}, \ell} \otimes E_t) \rightarrow V^{2k}(U)_{\bar{\lambda}, \ell} \otimes E_t$$

from (18) tensored with E_t and

$$(\chi_{\lambda, t}^{k-1} \otimes \mathrm{Sym}^{2k-2}(\eta_{\lambda, t})) \otimes (\chi_{\lambda, t} \otimes \mathrm{Sym}^2(\eta_{\lambda, t})) \rightarrow \chi_{\lambda, t}^k \otimes \mathrm{Sym}^{2k}(\eta_{\lambda, t})$$

from (17) with $\rho = \eta_{\lambda, t}$ and $m = 2k$ both arise from the symmetrization map. As the left hand sides are isomorphic G_F -modules by induction hypothesis, so are the right hand sides. This proves (15).

Similarly, replacing the right hand sides of the above two surjective homomorphisms by the middle term from the right side of (18) and (17) respectively, we obtain two isomorphic G_F -modules $\det(\eta_{\lambda, t}) \otimes \chi_{\lambda, t}^k \otimes \mathrm{Sym}^{2k-2}(\eta_{\lambda, t})$ and $E_t(-1) \otimes V^{2k-2}(U)_{\bar{\lambda}, \ell} \otimes E_t$, which implies the identity $\chi_{\lambda, t} \det(\eta_{\lambda, t}) = \epsilon_t$ since the G_F action on $E_t(-1)$ is ϵ_t .

Case (2) $G = \mathrm{GL}_2$. This follows from Deligne [19]. \square

Remark 3. For the groups Γ considered in this paper, the curves X_Γ are defined over \mathbf{Q} , hence $X_\Gamma = X_U$ for the group U corresponding to Γ . Formulae (15) and (16) indicate that, when $-\mathrm{id} \notin U$, the Frobenius traces of $\rho^{2k}(U)_{\lambda, \ell}$ and $\rho^k(U)_{\lambda, \ell}$ can be computed from those of $\rho^2(U)_{\lambda, \ell}$ and $\rho^1(U)_{\lambda, \ell}$, using the relation given by (2) and (7), respectively. When U contains $-\mathrm{id}$, by definition, $\rho^{2k}(U)_{\lambda, \ell}$ is an extension of $\rho^{2k}(U')_{\lambda', \ell} = \chi_{\lambda', \ell}^k \otimes \mathrm{Sym}^{2k}(\eta_{\lambda', \ell})$ for some neat compact open normal subgroup U' of U , resulting from the sheaf $V^{2k}(U)_\ell$ being a push-forward of $V^{2k}(U')_\ell$. While $\rho^2(U)_{\lambda, \ell}$ may no longer be a symmetric square up to twist by a character, the recursive relation (18) holds over X_U° . In particular, this implies that if λ is in $X_U^\circ(\mathbf{Q})$, then at a good prime p , the trace of $\rho^{2k}(U)_{\lambda, \ell}$ at Frob_p is equal to $F_k(S, T)$ defined by (2) with S being the trace of $\rho^2(U)_{\lambda, \ell}(\mathrm{Frob}_p)$ and $T = p$, the same expression as the case U not containing $-\mathrm{id}$. This will be used in the proofs of the theorems listed in §1.1. Thus our work is reduced to determining the Frobenius traces of $\rho^2(U)_{\lambda, \ell}$ or $\rho^1(U)_{\lambda, \ell}$ as appropriate. A key contribution of this paper is to showcase certain groups U (or their corresponding Γ as discussed in §1.1) for which we can identify suitable hypergeometric sheaves with explicit Galois actions which are isomorphic to the automorphic sheaves in question.

We end this section by discussing the integrality of Frobenius traces of the Galois action on the stalk $V^k(U)_{\bar{\lambda}, \ell}$ at an algebraic point λ in X_U° . First assume the group $G = \mathbf{B}^\times$ for a definite

quaternion division algebra B defined over \mathbf{Q} . If A is a 2-dimensional abelian variety defined over a number field L with multiplication by B , then the action of $\text{Gal}(\bar{L}/L)$ on the first étale cohomology group factors as

$$\rho_\ell : \text{Gal}(\bar{L}/L) \rightarrow (B \otimes_{\mathbf{Q}} \mathbf{Q}_\ell)^\times = \mathbf{B}^\times(\mathbf{Q}_\ell) \subset \text{Aut}(H^1(A \otimes_L \bar{L}, \mathbf{Q}_\ell)) \sim \text{GL}_4(\mathbf{Q}_\ell).$$

This is because the commutant of B in $\text{End}(H^1(A \otimes_L \bar{L}, \mathbf{Q}_\ell)) \simeq \text{M}_4(\mathbf{Q}_\ell)$ is $B_\ell = B \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$. It follows from the construction of the ℓ -adic sheaf in §2.2.2 that, the stalks $V^k(U)_{\bar{\lambda}, \ell}$ in an L -rational point $\lambda \in X_U^\circ(L)$, as a Galois module, are in the shape

$$\varrho'_k(\mathbf{Q}_\ell) \circ \rho_\ell : \text{Gal}(\bar{L}/L) \rightarrow \mathbf{B}^\times(\mathbf{Q}_\ell) \rightarrow \text{GL}_m(\mathbf{Q}_\ell)$$

where $m = k + 1$ for k even and $m = 2(k + 1)$ for k odd, as explained in §2.2.2. From this, one can prove that the coefficients of the characteristic polynomials in unramified primes \mathfrak{p} of L

$$(19) \quad \det \left(1 - T \text{Frob}_{\mathfrak{p}} \mid V^k(U)_{\bar{\lambda}, \ell} \right) \in \mathbf{Z}[T]$$

have integer coefficients. We will not prove this here. The above reasoning assumes that U is neat. To do a general U , which may include some Atkin-Lehner operators as above, note that the Frobenius traces can be computed, as in the theory of Artin L -functions, by averaging the Frobenius traces in the fibers over λ in a covering $X_{U'} \rightarrow X_U$ for a neat U' .

A similar argument yields the same conclusion for $G = \text{GL}_2$. In this case A is an elliptic curve, $\text{End}(H^1(A \otimes_L \bar{L}, \mathbf{Q}_\ell)) \simeq \text{M}_2(\mathbf{Q}_\ell)$ and the Galois action on the stalk $V^k(U)_{\bar{\lambda}, \ell}$ has the shape $\text{Gal}(\bar{L}/L) \rightarrow \text{GL}_{k+1}(\mathbf{Q}_\ell)$.

3. GROUPS

3.1. Hypergeometric functions over \mathbf{C} . We first recall some relevant notation in hypergeometric functions. Let $\Gamma(x)$ denote the Gamma function. For $k \in \mathbf{Z}$ and $a \in \mathbf{C}$, define the Pochhammer symbol $(a)_k := \Gamma(a + k)/\Gamma(a) = a(a + 1) \cdots (a + k - 1)$. A hypergeometric datum is a pair of multi-sets $\alpha = \{a_1, \dots, a_n\}, \beta = \{b_1 = 1, b_2, \dots, b_n\}$ with $a_i, b_j \in \mathbf{Q}$ to which we associate the Hypergeometric function in variable $t \in \mathbf{C}$:

$$(20) \quad F(\alpha, \beta; t) = {}_nF_{n-1} \left[\begin{matrix} a_1 & a_2 & \cdots & a_n \\ b_2 & \cdots & b_n \end{matrix} ; t \right] := \sum_{k \geq 0} \frac{(a_1)_k \cdots (a_n)_k}{(b_1)_k \cdots (b_n)_k} t^k.$$

It satisfies the Fuchsian ordinary differential equation below with three regular singular points at $0, 1, \infty$:

$$[\theta(\theta + b_2 - 1) \cdots (\theta + b_n - 1) - t(\theta + a_1) \cdots (\theta + a_n)] F = 0, \quad \text{where } \theta := t \frac{d}{dt}.$$

The local solutions of this equation form a rank- n hypergeometric local system.

Theorem 4. *The local exponents of the above hypergeometric differential equation are*

$$(21) \quad \begin{aligned} &0, 1 - b_2, \dots, 1 - b_n && \text{at } t = 0, \\ &a_1, a_2, \dots, a_n && \text{at } t = \infty, \\ &0, 1, 2, \dots, n - 2, \gamma && \text{at } t = 1, \end{aligned}$$

respectively, where

$$(22) \quad \gamma = -1 + \sum_{j=1}^n b_j - \sum_{j=1}^n a_j.$$

See [13, §2] by Beukers and Heckman for more details. We say that a local monodromy matrix in $\mathrm{GL}_n(\mathbf{R})$ has exponents $(r_1, \dots, r_n) \in (\mathbf{Q}/\mathbf{Z})^n$ if its eigenvalues are $e^{2\pi i r_1}, \dots, e^{2\pi i r_n}$. In particular, the above theorem says the local monodromy at 1 with monodromy matrix M_1 is a pseudoreflection (cf. [13, 36]), namely the rank of $M_1 - I_n$ is 1, where I_n stands for the rank- n identity matrix.

Next we recall a couple useful transformation formulae. Combined with the Euler transformation formula (cf. [5, Theorem 2.2.5])

$$(23) \quad {}_2F_1 \left[\begin{matrix} a & b \\ & c \end{matrix}; t \right] = (1-t)^{c-a-b} {}_2F_1 \left[\begin{matrix} c-a & c-b \\ & c \end{matrix}; t \right],$$

a version of the Clausen formula ([5] or [45, Eqn. (39)]) says that for $a, b, t \in \mathbf{C}$,

$$(24) \quad (1-t)^{-\frac{1}{2}} {}_3F_2 \left[\begin{matrix} \frac{1}{2} & a-b+\frac{1}{2} & b-a+\frac{1}{2} \\ & a+b+\frac{1}{2} & \frac{3}{2}-a-b \end{matrix}; t \right] = {}_2F_1 \left[\begin{matrix} a & b \\ & a+b+\frac{1}{2} \end{matrix}; t \right] {}_2F_1 \left[\begin{matrix} 1-a & 1-b \\ & \frac{3}{2}-a-b \end{matrix}; t \right]$$

when both sides are convergent.

3.2. A motivating example. We demonstrate the main motivation of this section using the following classical example. Let $\Gamma = \mathrm{PSL}_2(\mathbf{Z}) \cong (2, 3, \infty)$ whose modular curve $X_\Gamma = X(1)$ is parameterized by the modular j -function. The open subset $X(1)^\circ := X(1) \setminus \{0, 1728, \infty\}$ admits the universal elliptic curve

$$(25) \quad \mathcal{E}_j : y^2 + xy = x^3 - \frac{36x+1}{j-1728}.$$

Through the fiber map $f : \mathcal{E}_j \mapsto j$, the local constant sheaf \mathbf{Q}_ℓ on \mathcal{E}_j gives rise to $\mathcal{F}_\ell = R^1 f_* \mathbf{Q}_\ell$ on $X(1)^\circ$. The Picard-Fuchs equation of this elliptic fibration, using an algorithm in [62, Theorem 1.5]

by Stienstra and Beukers, is hypergeometric with one solution given by $(1-t)^{-1/4} {}_2F_1 \left[\begin{matrix} \frac{1}{12} & \frac{5}{12} \\ & 1 \end{matrix}; t \right]$

with $t = 1728/j$. Note that the hypergeometric datum $\{\{\frac{1}{12}, \frac{5}{12}\}, \{1, 1\}\}$ is not defined over \mathbf{Q} . However

$$\begin{aligned} \left((1-t)^{-1/4} {}_2F_1 \left[\begin{matrix} \frac{1}{12} & \frac{5}{12} \\ & 1 \end{matrix}; t \right] \right)^2 &\stackrel{(23)}{=} {}_2F_1 \left[\begin{matrix} \frac{1}{12} & \frac{5}{12} \\ & 1 \end{matrix}; t \right] {}_2F_1 \left[\begin{matrix} \frac{11}{12} & \frac{7}{12} \\ & 1 \end{matrix}; t \right] \\ &\stackrel{(24)}{=} (1-t)^{-1/2} {}_3F_2 \left[\begin{matrix} \frac{1}{2} & \frac{1}{6} & \frac{5}{6} \\ & 1 & 1 \end{matrix}; t \right], \end{aligned}$$

in which the hypergeometric datum $\{\{\frac{1}{2}, \frac{1}{6}, \frac{5}{6}\}, \{1, 1, 1\}\}$ in the last expression is defined over \mathbf{Q} .

Here a datum $HD = \{\alpha = \{a_1, \dots, a_n\}, \beta = \{b_1 = 1, b_2, \dots, b_n\}\}$ with $a_i, b_j \in \mathbf{Q}^\times$ is said to be defined over \mathbf{Q} if the set of column vectors $\left\{ \begin{pmatrix} a_1 \\ b_1 \end{pmatrix}, \dots, \begin{pmatrix} a_n \\ b_n \end{pmatrix} \right\} \bmod \mathbf{Z}$ is invariant under multiplication by all $r \in (\mathbf{Z}/M\mathbf{Z})^\times$, where $M = M(HD)$, called the level of HD , is the least positive common denominators of $a_1, \dots, a_n, b_1, \dots, b_n$. See Definition 1 in [45, §2.2]. Also, in this paper we mainly consider primitive hypergeometric data HD , namely $a_i - b_j \notin \mathbf{Z}$ for all i, j , such that the corresponding local systems are irreducible.

3.3. The groups to be considered. We now restrict our discussion from congruence subgroups of $\mathrm{SL}_2(\mathbf{R})$ to a class of arithmetic triangle groups which have been classified by Takeuchi [63, 64]. Petersson [55] (see also [17] by Clark and Voight) provides an algorithm to express the generators as matrices over a totally real subfield of a cyclotomic field. Among all arithmetic triangle groups, we consider those whose corresponding rank-2 local systems V are hypergeometric with datum $\alpha_2 = \{a, b\}, \beta_2 = \{1, c\}$ such that either

- a) The datum $\{\alpha_2, \beta_2\}$ is defined over \mathbf{Q} ; or,

b) $\text{Sym}^2 V$ is also hypergeometric, which is very special due to the pseudoreflection requirement, and its corresponding datum is defined over \mathbf{Q} (as the example in §3.2).

The above criteria give rise to the following groups, classified according to the conditions they satisfy:

- a.I) (m, ∞, ∞) with $m = 3, \infty$,
- b.I) $(2, m, \infty)$ with $m = 3, 4, 6, \infty$,
- b.II) $(2, 4, 6), (2, 6, 6)$.

All groups are isomorphic to a subgroup of $\text{SL}_2(\mathbf{R}) \bmod \pm I_2$, while those in a.I) are the only ones isomorphic to a subgroup of $\text{SL}_2(\mathbf{R})$ (in fact $\text{SL}_2(\mathbf{Z})$) not containing $-I_2$.

We now describe how we arrive at the lists. Given a triangle group $\Gamma = (e_0, e_1, e_\infty)$, following Theorem 9 of [68] by Y. Yang we introduce the following hypergeometric parameters:

$$a = \frac{1}{2}\left(1 - \frac{1}{e_1} - \frac{1}{e_0} - \frac{1}{e_\infty}\right), \quad b = \frac{1}{2}\left(1 - \frac{1}{e_1} - \frac{1}{e_0} + \frac{1}{e_\infty}\right), \quad c = 1 - \frac{1}{e_0},$$

and

$$\tilde{a} = \frac{1}{2}\left(1 - \frac{1}{e_1} + \frac{1}{e_0} - \frac{1}{e_\infty}\right), \quad \tilde{b} = \frac{1}{2}\left(1 - \frac{1}{e_1} + \frac{1}{e_0} + \frac{1}{e_\infty}\right), \quad \tilde{c} = 1 + \frac{1}{e_0}.$$

Using these Yang wrote down an explicit basis for $S_k(\Gamma)$ in terms of ${}_2F_1\left[\begin{smallmatrix} a & b \\ c \end{smallmatrix}; t\right]$ and ${}_2F_1\left[\begin{smallmatrix} \tilde{a} & \tilde{b} \\ \tilde{c} \end{smallmatrix}; t\right]$

which satisfy ordinary differential equations that are projectively equivalent. When $e_0 = e_1 = \infty$, the corresponding arithmetic triangle group $(e_\infty, \infty, \infty)$ can be identified with an index-2 subgroup of a Hecke group up to conjugation in $\text{PSL}_2(\mathbf{R})$. We further check whether they have a realization in $\text{SL}_2(\mathbf{R})$ not containing $-I_2$. This process leads to the list a.I) above with $HD(\Gamma) = \{\alpha = \{\frac{1}{2}(1 - \frac{1}{e_\infty}), \frac{1}{2}(1 + \frac{1}{e_\infty})\}, \beta = \{1, 1\}\}$, in which the Hauptmodul is chosen such that the two cusps are located at $t = 0$ and $t = 1$.

If, in particular, $e_1 = 2$, then the Clausen formula recalled in (24) implies

$$(26) \quad {}_2F_1\left[\begin{smallmatrix} a & b \\ c \end{smallmatrix}; t\right] {}_2F_1\left[\begin{smallmatrix} \tilde{a} & \tilde{b} \\ \tilde{c} \end{smallmatrix}; t\right] = {}_3F_2\left[\begin{smallmatrix} \frac{1}{2} & \frac{1}{2} - \frac{1}{e_\infty} & \frac{1}{2} + \frac{1}{e_\infty} \\ 1 - \frac{1}{e_0} & 1 + \frac{1}{e_0} \end{smallmatrix}; t\right].$$

The hypergeometric datum corresponding to the right hand side of (26) is

$$(27) \quad HD := HD(\Gamma) := \left\{ \alpha = \left\{ \frac{1}{2}, \frac{1}{2} - \frac{1}{e_\infty}, \frac{1}{2} + \frac{1}{e_\infty} \right\}, \beta = \left\{ 1, 1 - \frac{1}{e_0}, 1 + \frac{1}{e_0} \right\} \right\}$$

in which α and β are self-dual. The datum is defined over \mathbf{Q} if $e_0, e_\infty \in \{3, 4, 6, \infty\}$. This leads to the groups listed in b.I) for Γ non-cocompact and b.II) for Γ cocompact.

As explained in §3.1, a hypergeometric sheaf is defined on \mathbf{P}^1 with singularities at $0, 1, \infty$. When we associate a hypergeometric sheaf to a triangle group Γ , the three singularities of the sheaf are set at the three vertices of the curve X_Γ (as a compact Riemann surface). For this to happen, the model for X_Γ should be a genus zero curve with the three singular points that are \mathbf{Q} -rational.

The models we choose for the non-cocompact groups are as follows

$$\begin{array}{cccc} (2, \infty, \infty) & (2, 3, \infty) & (2, 4, \infty) & (2, 6, \infty) \\ \Gamma_0(2)/\{\pm I\} & \text{PSL}_2(\mathbf{Z}) & \langle \Gamma_0(2), w_2 \rangle / \{\pm I\} & \langle \Gamma_0(3), w_3 \rangle / \{\pm I\} \end{array}$$

where w_2 and w_3 are Atkin-Lehner involutions, so that X_Γ are projective curves over \mathbf{Q} and all elliptic points and cusps are \mathbf{Q} -rational. Hence we may choose $\lambda(\Gamma)$ as described in the table in Theorem 1. For the cocompact group $(2, 4, 6)$, more explanation is needed. The quotient of the norm-1 group of $O_{B_6}^1$ by its center ± 1 gives the group $(2, 2, 3, 3)$ in the class II group 1b. It is normalized by the Atkin-Lehner involutions w_2, w_3 , and $w_6 = w_2 w_3$. The extended groups $\langle (2, 2, 3, 3), w_i \rangle$ for $i = 3, 2, 6$ respectively are the groups $(2, 6, 6), (3, 4, 4)$ and $(2, 2, 2, 3)$,

and $(2, 4, 6) = \langle (2, 2, 3, 3), w_2, w_3 \rangle$. The curve $X_{(2,2,3,3)}$ is a genus 0 curve defined over \mathbf{Q} with the canonical model given by $x^2 + 3y^2 + z^2 = 0$ (cf. [41]). As described in Baba-Granath [9], it is a 2-fold cover of the curves $X_{(2,6,6)}$, $X_{(3,4,4)}$, $X_{(2,2,2,3)}$, all projective lines over \mathbf{Q} , via the covering maps sending $[x : y : z] \in X_{(2,2,3,3)}$ to $[y : z]$, $[x : z]$, $[x : y]$, respectively. The curve $X_{(2,4,6)}$ is also a projective line over \mathbf{Q} ; it is covered by $X_{(2,6,6)}$, $X_{(3,4,4)}$, $X_{(2,2,2,3)}$ with respective covering maps

$$(28) \quad \pi_3 : [y : z] \mapsto [-3y^2 - z^2 : y^2], \quad \pi_2 : [x : z] \mapsto \left[x^2 : \frac{-x^2 - z^2}{3} \right], \quad \pi_6 : [x : y] \mapsto [x^2 : y^2].$$

These are the models we choose. As such, the three elliptic points on $X_{(2,4,6)}$ are $[0 : 1]$, $[1 : 0]$, $[-3 : 1]$, all \mathbf{Q} -rational; for the sake of convenience we may choose the Hauptmodul λ for the curve $X_{(2,4,6)}$ as given in Theorem 1. But $X_{(2,6,6)}$, $X_{(3,4,4)}$ and $X_{(2,2,2,3)}$ all have elliptic points rational only over a quadratic extension of \mathbf{Q} . For this reason, we did not include $(2, 6, 6)$ in Theorem 1. Since $X_{(2,6,6)}$ is a 2-fold cover of $X_{(2,4,6)}$ with an explicit covering map, we will use the pullback of the hypergeometric sheaf on $X_{(2,4,6)}$ to deal with $(2, 6, 6)$, see Theorem 14 in Section 7.1.

The rank-2 hypergeometric local systems $\mathcal{H}^1(\Gamma)_{\mathbf{C}} = \mathcal{H}(HD(\Gamma))_{\mathbf{C}}$ for groups in a.I) are listed in the Table 2 below.

$\Gamma = (e_0, e_1, e_\infty)$	Generators	Exponents	$HD(\Gamma)$	$\mathcal{H}^1(\Gamma)_{\mathbf{C}} = \mathcal{H}(HD(\Gamma))_{\mathbf{C}}$ a particular function
$(\infty, \infty, 3)$	$T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$	$(1, 1)$	$\alpha = \{\frac{1}{3}, \frac{2}{3}\}$	$\mathcal{H}(\{\frac{1}{3}, \frac{2}{3}\}, \{1, 1\})_{\mathbf{C}}$
$\cong \Gamma_1(3)$	$S := \begin{pmatrix} -2 & 3 \\ -3 & 4 \end{pmatrix}$ $(ST)^{-1}$	$(1, 1)$ $(\frac{1}{3}, \frac{2}{3})$	$\beta = \{1, 1\}$	${}_2F_1 \left[\begin{matrix} \frac{1}{3} & \frac{2}{3} \\ 1 \end{matrix} ; t \right]$
(∞, ∞, ∞)	$T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$	$(1, 1)$	$\alpha = \{\frac{1}{2}, \frac{1}{2}\}$	$\mathcal{H}(\{\frac{1}{2}, \frac{1}{2}\}, \{1, 1\})_{\mathbf{C}}$
$\cong \Gamma_1(4)$	$(ST)^{-1}$ $S := \begin{pmatrix} 1 & -1 \\ 4 & -3 \end{pmatrix}$	$(1, 1)$ $(\frac{1}{2}, \frac{1}{2})$	$\beta = \{1, 1\}$	${}_2F_1 \left[\begin{matrix} \frac{1}{2} & \frac{1}{2} \\ 1 \end{matrix} ; t \right]$

TABLE 2. Groups in a.I) list

Notice that the exponents of the generator S of $\Gamma = (\infty, \infty, \infty) \simeq \Gamma_1(4)$ are $(\frac{1}{2}, \frac{1}{2})$. This means that the point fixed by S is an *irregular* cusp, so the contribution of this cusp to $\text{Tr}(T_p | S_{k+2}(\Gamma))$ will depend on the parity of k as in the statement of Theorem 2.

For groups in b.I), we also write down the rank-2 local hypergeometric system $\mathcal{H}^1(\Gamma)_{\mathbf{C}}$. The groups Γ in b.II) arise from indefinite quaternion algebras defined over \mathbf{Q} . As explained in §2.1, $V^2(\Gamma)_{\mathbf{C}}$ is a 3-dimensional irreducible local system. For each case, we also identify a rank-2 hypergeometric local system $\mathcal{H}^1(\Gamma)_{\mathbf{C}}$ listed in the second to the last column of Table 3, in which branches for n th roots are chosen accordingly to make the local exponents on the third column as claimed. The symmetric square of $\mathcal{H}^1(\Gamma)_{\mathbf{C}}$, denoted by $\mathcal{H}^2(\Gamma)_{\mathbf{C}}$, is of the form $(1-t)^{-\frac{1}{2}}F(HD(\Gamma); t)$, where $F(HD(\Gamma); t)$, defined by (20), is as shown in the last column of Table 3. By construction, for each Γ in the b) list, the hypergeometric datum $HD(\Gamma)$ as given in (27) is defined over \mathbf{Q} . The data for these groups are listed in Table 3. We will omit $(2, 6, 6)$ for reasons explained earlier.

We take the representation of B_6 in $M_2(\mathbf{R})$ by matrices of size 2 over $\mathbf{Q}(\sqrt{3})$ and choose

$$(29) \quad s_4 = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 & 1 \\ -1 & -1 \end{pmatrix}, \quad s_6 = \frac{1}{2\sqrt{3}} \begin{pmatrix} 3 + \sqrt{3} & -3 + \sqrt{3} \\ 3 + \sqrt{3} & 3 - \sqrt{3} \end{pmatrix}, \quad s_2 = (s_4 s_6)^{-1}$$

(e_0, e_1, e_∞)	Generators	Expo.	M	$\mathcal{H}^1(\Gamma)_{\mathbf{C}}$ a particular function	$\mathcal{H}^2(\Gamma)_{\mathbf{C}} = \mathcal{L}(\Gamma)_{\mathbf{C}} \otimes \mathcal{H}(HD(\Gamma))_{\mathbf{C}}$ a particular function
$(\infty, 2, \infty)$	$T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ $S := \begin{pmatrix} -1 & 1 \\ -2 & 1 \end{pmatrix}$ $(ST)^{-1}$	$(1, 1)$ $(\frac{1}{4}, \frac{3}{4})$ $(\frac{1}{5}, \frac{1}{5})$	4	$\mathcal{H}(\{\frac{1}{4}\}, \{1\})_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{4}, \frac{1}{4}\}, \{1, 1\})_{\mathbf{C}}$ $(1-t)^{-\frac{1}{4}} {}_2F_1 \left[\begin{matrix} \frac{1}{4} & \frac{1}{4} \\ 1 \end{matrix}; t \right]$	$\mathcal{H}(\{\frac{1}{2}\}, \{1\})_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\}, \{1, 1, 1\})_{\mathbf{C}}$ $(1-t)^{-\frac{1}{2}} {}_3F_2 \left[\begin{matrix} \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ 1 & 1 \end{matrix}; t \right]$
$(\infty, 2, 3)$	$T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ $S := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ $(ST)^{-1}$	$(1, 1)$ $(\frac{1}{4}, \frac{3}{4})$ $(\frac{1}{3}, \frac{2}{3})$	12	$\mathcal{H}(\{\frac{1}{4}\}, \{1\})_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{12}, \frac{5}{12}\}, \{1, 1\})_{\mathbf{C}}$ $(1-t)^{-\frac{1}{4}} {}_2F_1 \left[\begin{matrix} \frac{1}{12} & \frac{5}{12} \\ 1 \end{matrix}; t \right]$	$\mathcal{H}(\{\frac{1}{2}\}, \{1\})_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{2}, \frac{1}{6}, \frac{5}{6}\}, \{1, 1, 1\})_{\mathbf{C}}$ $(1-t)^{-\frac{1}{2}} {}_3F_2 \left[\begin{matrix} \frac{1}{2} & \frac{1}{6} & \frac{5}{6} \\ 1 & 1 \end{matrix}; t \right]$
$(\infty, 2, 4)$	$T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ $S := \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ -2 & 0 \end{pmatrix}$ $(ST)^{-1}$	$(1, 1)$ $(\frac{1}{4}, \frac{3}{4})$ $(\frac{3}{8}, \frac{5}{8})$	8	$\mathcal{H}(\{\frac{1}{4}\}, \{1\})_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{8}, \frac{3}{8}\}, \{1, 1\})_{\mathbf{C}}$ $(1-t)^{-\frac{1}{4}} {}_2F_1 \left[\begin{matrix} \frac{1}{8} & \frac{3}{8} \\ 1 \end{matrix}; t \right]$	$\mathcal{H}(\{\frac{1}{2}\}, \{1\})_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{2}, \frac{1}{4}, \frac{3}{4}\}, \{1, 1, 1\})_{\mathbf{C}}$ $(1-t)^{-\frac{1}{2}} {}_3F_2 \left[\begin{matrix} \frac{1}{2} & \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix}; t \right]$
$(\infty, 2, 6)$	$T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ $S := \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & 1 \\ -3 & 0 \end{pmatrix}$ $(ST)^{-1}$	$(1, 1)$ $(\frac{1}{4}, \frac{3}{4})$ $(\frac{5}{12}, \frac{7}{12})$	6	$\mathcal{H}(\{\frac{1}{4}\}, \{1\})_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{6}, \frac{1}{3}\}, \{1, 1\})_{\mathbf{C}}$ $(1-t)^{-\frac{1}{4}} {}_2F_1 \left[\begin{matrix} \frac{1}{6} & \frac{1}{3} \\ 1 \end{matrix}; t \right]$	$\mathcal{H}(\{\frac{1}{2}\}, \{1\})_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{2}, \frac{1}{3}, \frac{2}{3}\}, \{1, 1, 1\})_{\mathbf{C}}$ $(1-t)^{-\frac{1}{2}} {}_3F_2 \left[\begin{matrix} \frac{1}{2} & \frac{1}{3} & \frac{2}{3} \\ 1 & 1 \end{matrix}; t \right]$
$(6, 2, 4)$	s_6 s_2 s_4	$(\frac{1}{12}, \frac{11}{12})$ $(\frac{1}{4}, \frac{3}{4})$ $(\frac{3}{8}, \frac{5}{8})$	24	$\mathcal{K}_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{4}\}, \{1\})_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{24}, \frac{7}{24}\}, \{1, \frac{5}{6}\})_{\mathbf{C}}$ $t^{-\frac{1}{12}} (1-t)^{-\frac{1}{4}} {}_2F_1 \left[\begin{matrix} \frac{1}{24} & \frac{7}{24} \\ \frac{5}{6} \end{matrix}; t \right]$	$\mathcal{H}(\{\frac{1}{2}\}, \{1\})_{\mathbf{C}} \otimes \mathcal{H}(\{\frac{1}{2}, \frac{1}{4}, \frac{3}{4}\}, \{1, \frac{5}{6}, \frac{7}{6}\})_{\mathbf{C}}$ $(1-t)^{-\frac{1}{2}} {}_3F_2 \left[\begin{matrix} \frac{1}{2} & \frac{1}{4} & \frac{3}{4} \\ \frac{5}{6} & \frac{7}{6} \end{matrix}; t \right]$

TABLE 3. Groups in b) list except (2,6,6)
where $(\mathcal{K}_{\mathbf{C}})_{\bar{t}} = t^{-\frac{1}{12}}$ and $(\mathcal{H}(\{a\}, \{1\})_{\mathbf{C}})_{\bar{t}} = (1-t)^{-a}$.

with $s_4^4 = s_6^6 = s_2^2 = -I_2$ as matrices. See [12, 21] for more detail. They generate a group whose quotient mod $\pm I_2$ is isomorphic to $(2, 4, 6)$.

3.4. Isomorphisms using rigidity. For arithmetic triangle groups, the rigidity theorems may be applied to compare the automorphic sheaves described in §2 and hypergeometric local systems arising from local solutions of hypergeometric differential equations. More details are discussed in [36], where ℓ -adic hypergeometric sheaves on the algebraic group \mathbb{G}_m are also introduced.

Theorem 5. (Katz [36, Rigidity Theorem 3.5.4]) *Let \mathcal{F} and \mathcal{G} be two irreducible local systems on $(\mathbb{G}_m - \{s_0\})^{an}$ of the same rank $n \geq 1$, where s_0 is a fixed nonzero value, and satisfying*

- 1) *The local monodromies at s_0 of both \mathcal{F} and \mathcal{G} are pseudoreflections;*
- 2) *At both 0 and ∞ , \mathcal{F} and \mathcal{G} have the same characteristic polynomials of local monodromies respectively.*

Then \mathcal{F} and \mathcal{G} are isomorphic.

For the groups in a) list, the generator T listed in Table 2 is a pseudoreflection. From Theorems 4 and 5, and the information in Table 2, we have the following conclusion.

Theorem 6. *The local system $V^1(\Gamma)_{\mathbf{C}}$ is isomorphic to $\mathcal{H}^1(\Gamma)_{\mathbf{C}} = \mathcal{H}(HD(\Gamma))_{\mathbf{C}}$ given in the last column of Table 2.*

We now illustrate how to use Katz's rigidity theorem to relate $V^2(\Gamma)_{\mathbf{C}}$ and hypergeometric local system directly for b) groups (which also works for $(2, 6, 6)$). For both local systems, $s_0 = 1$.

Theorem 7. *For each group Γ in Table 3, let $V^2(\Gamma)_{\mathbf{C}}$ be the rank-3 sheaf as in §2.1. Then $V^2(\Gamma)_{\mathbf{C}}$ is isomorphic to $\mathcal{L}(\Gamma)_{\mathbf{C}} \otimes \mathcal{H}(HD(\Gamma))_{\mathbf{C}} = \mathcal{H}^2(\Gamma)_{\mathbf{C}}$, where $\mathcal{H}(HD(\Gamma))_{\mathbf{C}}$ and $\mathcal{H}^2(\Gamma)_{\mathbf{C}}$ are the hypergeometric sheaves in Table 3. Moreover $V^2(\Gamma)_{\mathbf{C}}$ is isomorphic to the symmetric square of $\mathcal{H}^1(\Gamma)_{\mathbf{C}}$ listed in the second to the last column in Table 3.*

Proof. By Clausen formula, $\text{Sym}^2 \mathcal{H}^1(\Gamma)_{\mathbf{C}}$ is isomorphic to $\mathcal{H}^2(\Gamma)_{\mathbf{C}}$. It remains to prove the first statement.

The groups $\Gamma = (2, m, \infty)$ in b.I) with $m = 3, 4, 6$ are isomorphic to the projective groups $\text{PSL}_2(\mathbf{Z})$, $\Gamma_0^+(2)/\{\pm I_2\}$, and $\Gamma_0^+(3)/\{\pm I_2\}$ respectively. They are generated by the matrices

$$T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad S := \begin{pmatrix} 0 & 1/\sqrt{a} \\ -\sqrt{a} & 0 \end{pmatrix}, \quad (ST)^{-1}, \quad \text{where } a = \lfloor \frac{m}{2} \rfloor.$$

Thus the fixed points of $(ST)^{-1}$ are $\frac{-1+\sqrt{-3}}{2}$, $\frac{-1+\sqrt{-1}}{2}$, $\frac{-1+\sqrt{-1/3}}{2}$ respectively. Also

$$\text{Sym}^2 T = \begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad \text{Sym}^2 S = \begin{pmatrix} 0 & 0 & 1/a \\ 0 & -1 & 0 \\ a & 0 & 0 \end{pmatrix}, \quad \text{Sym}^2 (ST)^{-1} = \begin{pmatrix} a & 2 & 1/a \\ -a & -1 & 0 \\ a & 0 & 0 \end{pmatrix}.$$

We choose a Hauptmodul λ for Γ (such as $\lambda = 1728/j$ for $\text{PSL}_2(\mathbf{Z})$) whose values at the singular points are as described in Theorem 1. Precisely, it has a simple zero at infinity, takes value 1 at the fixed point of S , and has a simple pole at that of ST . The sheaf $V^2(\Gamma)_{\mathbf{C}}$ is defined in §2.1. While $V^1(\Gamma)_{\mathbf{C}}$ is not defined because the lift of Γ in $\text{SL}_2(\mathbf{R})$ contains $-I_2$, nonetheless the above three symmetric square matrices can be used to describe $V^2(\Gamma)_{\mathbf{C}}$. Indeed the Jordan normal forms of the local monodromy matrices of $V^2(\Gamma)_{\mathbf{C}} \otimes \mathcal{L}(\Gamma)_{\mathbf{C}}$ are

$$\begin{array}{ccc} 1 & 0 & \infty \\ \hline \begin{pmatrix} 1 & & \\ & 1 & \\ & & -1 \end{pmatrix} & \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} & \begin{pmatrix} -1 & & \\ & -\zeta_m & \\ & & -\zeta_m^{-1} \end{pmatrix} \end{array}$$

23

In particular, the monodromy at 1 is a pseudoreflection. The monodromy matrices at $1, 0, \infty$ have characteristic polynomials $(T-1)^2(T+1)$, $(T-1)^3$, $(T+1)(T+\zeta_m)(T+\zeta_m^{-1})$ respectively, which coincide with those of the hypergeometric local system $\mathcal{H}(HD(\Gamma))_{\mathbf{C}}$ with datum $HD(\Gamma)$ (see Theorem 4). We conclude from Theorem 5 that they are isomorphic.

The same argument applies to the fourth group $\Gamma = (2, \infty, \infty)$ in b.I), which is isomorphic to $\Gamma_0(2)/\{\pm I_2\}$. The only difference is the generator $S = \begin{pmatrix} -1 & 1 \\ -2 & 1 \end{pmatrix}$ so that $\text{Sym}^2 S$ has eigenvalues $-1, 1, -1$ and $\text{Sym}^2(ST)^{-1}$ has eigenvalues $1, 1, 1$.

The remaining case can be verified in the like manner. From the exponents of the generators for $(2,4,6)$ we know the exponents of their symmetric squares are $\{\frac{1}{4}, 1, \frac{3}{4}\}$, $\{\frac{7}{6}, 1, \frac{5}{6}\}$, and $\{\frac{1}{2}, \frac{1}{2}, 1\}$ respectively, which coincide with the local exponents of the order-3 differential equation satisfied by $(1-t)^{-\frac{1}{2}} {}_3F_2 \left[\begin{matrix} \frac{1}{2} & \frac{1}{4} & \frac{3}{4} \\ \frac{7}{6} & \frac{5}{6} \end{matrix}; t \right]$. \square

3.5. Comparison Theorem. The results of this section appear in [32]. We include them here for the convenience of the reader. Let F be a number field and $R = O_F[1/N]$ the localization of the ring of integers of F for an integer $N \geq 1$. Let $S = \text{Spec}(R)$. We let $\eta = \text{Spec}(F)$, the generic point of S , and $\bar{\eta} = \text{Spec}(\bar{F})$ for an algebraic closure of F . Let X/S be an irreducible separated scheme, smooth and of finite type over S , with geometrically connected fibers. We can choose a geometric generic point $\bar{\xi} : \text{Spec}(\bar{F}(X)) \rightarrow X$ which lies over $\bar{\eta}$, where $F(X)$ is the function field of X . We let X_η and $X_{\bar{\eta}}$ be the schemes over $\text{Spec} F$ and $\text{Spec} \bar{F}$ obtained from X by base-change.

We consider two geometrically irreducible lisse $\bar{\mathbf{Q}}_\ell$ -sheaves (for the étale topology) \mathcal{F}, \mathcal{G} on X . These are equivalent to two ℓ -adic representations

$$\rho_{\mathcal{F}} : \pi_1(X, \bar{\xi}) \longrightarrow \text{GL}(V), \quad \rho_{\mathcal{G}} : \pi_1(X, \bar{\xi}) \longrightarrow \text{GL}(W)$$

for finite-dimensional $\bar{\mathbf{Q}}_\ell$ -vector spaces V, W . Geometrically irreducible means: the restrictions

$$\rho_{\mathcal{F}}|_{X_{\bar{\eta}}}, \quad \rho_{\mathcal{G}}|_{X_{\bar{\eta}}}$$

of $\pi_1(X_{\bar{\eta}}, \bar{\xi})$ are irreducible. Note that we have a surjective homomorphism

$$\pi_1(X_\eta, \bar{\xi}) \rightarrow \pi_1(X, \bar{\xi})$$

and an exact sequence

$$0 \longrightarrow \pi_1(X_{\bar{\eta}}, \bar{\xi}) \longrightarrow \pi_1(X_\eta, \bar{\xi}) \longrightarrow \text{Gal}(\bar{F}/F) \longrightarrow 0.$$

Now let $\varphi : R \rightarrow \mathbf{C}$ be an embedding. We obtain a scheme $X_{\varphi, \mathbf{C}}$ over \mathbf{C} . This also defines an analytic space $X_\varphi^{\text{an}} := X_\varphi(\mathbf{C})$. Since φ will be fixed, we will drop it from the notation. There is a canonical map

$$\pi_1(X^{\text{an}}, u) \rightarrow \pi_1(X_{\mathbf{C}}, u)$$

(left-hand side: topological fundamental group; right-hand side, the étale fundamental group) which identifies the right-hand side with the profinite completion of left-hand side (Riemann's existence theorem). In particular, this map has dense image. Here we can take u to be the geometric $u : \text{Spec}(\mathbf{C}) \rightarrow X$

$$\text{Spec}(\mathbf{C}) \rightarrow \text{Spec}(\bar{F}(X)) = \bar{\xi} \rightarrow X$$

where the first arrow is induced by some embedding $\bar{\varphi} : \bar{F}(X) \rightarrow \mathbf{C}$ which extends φ . It is known that there are isomorphisms $\pi_1(X_{\mathbf{C}}, u) = \pi_1(X_{\bar{\eta}}, u) = \pi_1(X_{\bar{\eta}}, \bar{\xi})$ induced by $\bar{\varphi}$. The first holds because F has characteristic 0; the second is a change in base-point. Choose an isomorphism $\iota : \bar{\mathbf{Q}}_\ell \cong \mathbf{C}$. The theorem that follows will not depend on this artificial choice.

Composing all these, we get representations

$$\rho_{\mathcal{F}}^{\text{an}} : \pi_1(X^{\text{an}}, u) \rightarrow \pi_1(X_{\mathbf{C}}, u) = \pi_1(X_{\bar{\eta}}, \bar{\xi}) \xrightarrow{\rho_{\mathcal{F}}} \text{GL}(V) \cong \text{GL}(V_{\mathbf{C}})$$

where $V_{\mathbf{C}} = V \otimes_{\bar{\mathbf{Q}}_{\ell, \iota}} \mathbf{C}$. We get a similar story for $\rho_{\mathcal{G}}^{\text{an}}$. We let \mathbf{F} and \mathbf{G} be the \mathbf{C} -local systems on X^{an} that arise from these representations of the fundamental group. Also $\mathcal{D}(\mathbf{F})$ and $\mathcal{D}(\mathbf{G})$ the regular holonomic \mathcal{D} -modules (=connections with regular singular points) corresponding to these by the Riemann-Hilbert correspondence.

Theorem 8. *Under these assumptions (and \mathcal{F}, \mathcal{G} geometrically irreducible), if the local systems \mathbf{F} and \mathbf{G} on X^{an} are isomorphic (equivalently if the \mathcal{D} -modules $\mathcal{D}(\mathbf{F})$ and $\mathcal{D}(\mathbf{G})$ are isomorphic), then there is a continuous character $\chi : \text{Gal}(\bar{F}/F) \rightarrow \bar{\mathbf{Q}}_{\ell}^{\times}$, such that $\mathcal{G}_{\eta} \cong \mathcal{F}_{\eta} \otimes \chi$.*

Proof. There is a matrix $M : V_{\mathbf{C}} \rightarrow W_{\mathbf{C}}$ that intertwines the representation of $\pi_1(X^{\text{an}}, u)$ given by \mathbf{F} and \mathbf{G} . Then $\iota^{-1}(M) : V \rightarrow W$ is a matrix that intertwines the representations of $\pi_1(X^{\text{an}}, u)$ in $\text{GL}(V)$ and $\text{GL}(W)$. But $\pi_1(X^{\text{an}}, u)$ has dense image in $\pi_1(X_{\bar{\eta}}, \bar{\xi})$ and the representations given by \mathcal{G} and \mathcal{F} on V, W are continuous. Thus by continuity $\iota^{-1}(M)$ will intertwine those representations. Therefore the representations

$$\rho_{\mathcal{F}} \mid X_{\bar{\eta}}, \quad \rho_{\mathcal{G}} \mid X_{\bar{\eta}}$$

of $\pi_1(X_{\bar{\eta}}, \bar{\eta})$ are isomorphic. From the exact sequence above, and the fact that these representations are isomorphic we get, by the lemma below, a character $\chi : \text{Gal}(\bar{F}/F) \rightarrow \bar{\mathbf{Q}}_{\ell}^{\times}$ and an isomorphism

$$(\rho_{\mathcal{G}} \mid X_{\eta}) = (\rho_{\mathcal{F}} \mid X_{\eta}) \otimes \chi$$

as representations of $\pi_1(X_{\eta}, \bar{\xi})$. □

The following is well-known.

Lemma 2. *Given an exact sequence of groups*

$$0 \longrightarrow H \xrightarrow{a} G \xrightarrow{b} G/H \longrightarrow 0$$

and two finite-dimensional representations $\rho : G \rightarrow \text{GL}(V)$ and $\sigma : G \rightarrow \text{GL}(W)$ where V, W are vector spaces over an algebraically closed field k . Suppose that $\rho \mid H$ and $\sigma \mid H$ are irreducible and isomorphic. Then there is a character $\chi : G/H \rightarrow \text{GL}_1(k) = k^{\times}$, such that $\sigma \cong \rho \otimes \chi := \rho \otimes (\chi \circ b)$. If k is a topological field and ρ, σ are continuous representations, then χ is a continuous character.

Proof. Let $Z = \text{Hom}_k(V, W)$. This is a finite-dimensional k -space with a G -action:

$$(g \cdot f)(v) := \sigma(g)f(\rho(g)^{-1}v), \quad \text{where } v \in V, f \in Z, g \in G.$$

A fixed vector f , namely $g \cdot f = f, \forall g \in G$, is a G -equivariant map $f : V \rightarrow W$. By hypothesis there is an H -fixed vector corresponding to an H -equivariant isomorphism $f : V \xrightarrow{\sim} W$. Since these H -spaces are irreducible, any nonzero H -equivariant map $f : V \rightarrow W$ is an isomorphism, and any two such differ by a scalar multiple (Schur's lemma). Thus, $Z^H = k \cdot f$. The G -action on Z induces a G/H -action on Z^H . Since this is one-dimensional, this is a character χ which is defined by $g \cdot f = \chi(g)f$ for all $g \in G$. One checks that $f : V \rightarrow W$ is an isomorphism which is equivariant for the $\rho \otimes \chi$ action on V and σ on W . □

We can give a stronger version of Theorem 8 if we assume in addition that X/S has a section, and that X/S is the complement in Z/S of a divisor with normal crossings D/S , where Z/S is proper and smooth. We also assume that ℓ is invertible on X and S . Under those assumptions, then we have an exact sequence ([1, Ch. XIII, Prop. 4.3, and Examples 4.4])

$$0 \longrightarrow \pi_1^{\mathbb{L}}(X_{\bar{\eta}}, \bar{\xi}) \longrightarrow \pi_1'(X, \bar{\xi}) \longrightarrow \pi_1(S, \bar{\eta}) \longrightarrow 0 .$$

Here \mathbb{L} is a set of primes invertible on S , namely coprime to N . $\pi_1^{\mathbb{L}}(X_{\bar{\eta}}, \bar{\xi})$ is the pro- \mathbb{L} -quotient of $\pi_1(X_{\bar{\eta}}, \bar{\xi})$. If K is the kernel of the canonical homomorphism $\pi_1(X, \bar{\xi}) \rightarrow \pi_1(S, \bar{\eta})$ and $N \subset K$

is the smallest normal subgroup such that K/N is a pro- \mathbb{L} -group, then $N \subset \pi_1(X, \bar{\xi})$ is a normal subgroup, and we denote $\pi'_1(X, \bar{\xi}) = \pi_1(X, \bar{\xi})/N$. If we assume that the representations $\rho_{\mathcal{F}}$ and $\rho_{\mathcal{G}}$ factor through $\pi'_1(X, \bar{\xi})$ and are geometrically irreducible, we can conclude that there exists a continuous character $\chi : \pi_1(S, \bar{\eta}) \rightarrow \mathbf{Q}_\ell^\times$ such that $\mathcal{G} \cong \mathcal{F} \otimes \chi$ on X .

3.6. The ℓ -adic sheaves isomorphism. The main application of this theorem in our paper is to the situation where X is a (modular) curve of genus 0 defined over \mathbf{Q} and the two local systems are an automorphic sheaf and a hypergeometric sheaf respectively. The isomorphism of local systems on the analytic space X^{an} results from rigidity, as shown in §3.4. Because our local systems arise from geometry, they have special properties. For instance, they belong to a strictly compatible family of rational ℓ -adic representations.

To determine the twisting character χ in the statement of the Comparison Theorem 8 it suffices to restrict the sheaves to a \mathbf{Q} -rational point of the curve X° . Then \mathcal{F} and \mathcal{G} define representations of $G_{\mathbf{Q}} := \text{Gal}(\bar{\mathbf{Q}}/\mathbf{Q})$. The twisting character χ in the statement of the Comparison Theorem 8 is an algebraic Hecke character of the number field \mathbf{Q} . The reader is referred to Serre [58] for more detail.

In particular, we recall Corollary 2.1.6 of [8].

Lemma 3. *If χ is a 1-dimensional representation of $G_{\mathbf{Q}}$ taking values in the group of units of a finite extension of \mathbf{Q}_ℓ , which is unramified almost everywhere and whose restriction to the decomposition group at ℓ is crystalline, then $\chi = \epsilon_\ell^m \xi$, where ϵ_ℓ is the ℓ -adic cyclotomic character of $G_{\mathbf{Q}}$, $m \in \mathbf{Z}$ and ξ is a character of $G_{\mathbf{Q}}$ of finite order.*

As we have seen in §2, the automorphic sheaves $V^k(\Gamma)_{\mathbf{C}}$ have ℓ -adic counterparts $V^k(\Gamma)_\ell$. For the groups Γ we consider, we have introduced the hypergeometric sheaves $\mathcal{H}(HD(\Gamma))_{\mathbf{C}}$ attached to the hypergeometric data $HD(\Gamma)$ in Tables 2 and 3. To these $HD(\Gamma)$, Katz has associated the ℓ -adic sheaves $\mathcal{H}(HD(\Gamma))_\ell$, whose behavior at $0, 1, \infty$ agrees with that of $\mathcal{H}(HD(\Gamma))_{\mathbf{C}}$ under the change of generator $\lambda \mapsto 1/\lambda$ (see [36]). Theorems 6 and 7 tell us how the two complex sheaves of different origin are related. Using the Comparison Theorem 8, we know that the same relation holds for their ℓ -adic counterparts up to twist by a character χ_Γ . In the next section we will give more details of the ℓ -adic hypergeometric sheaves and determine the character χ_Γ .

4. HYPERGEOMETRIC CHARACTER SUMS

Given a hypergeometric datum $HD = \{\alpha, \beta\}$ of length n , in §3 we associated the hypergeometric function $F(HD; t)$ over \mathbf{C} and a rank- n sheaf $\mathcal{H}(HD)_{\mathbf{C}}$ on $\mathbb{P}_{\mathbf{C}}^1 \setminus \{0, 1, \infty\}$ along with the monodromy representation of the fundamental group $\pi_1(\mathbb{P}_{\mathbf{C}}^1 \setminus \{0, 1, \infty\}, *)$ with a base point. When the elements in α, β are in \mathbf{Q}^\times and HD is primitive, for each prime ℓ , Katz in [36, 37] has constructed an ℓ -adic sheaf $\mathcal{H}(HD)_\ell$ of rank- n on \mathbb{G}_m together with the action of the absolute Galois group $G(M) = \text{Gal}(\bar{\mathbf{Q}}/\mathbf{Q}(\zeta_M))$ of the cyclotomic field $\mathbf{Q}(\zeta_M)$. Here $M = M(HD) = \text{lcd}(\alpha \cup \beta)$, the least positive common denominator of elements in α and β , is the level of HD . After changing the generator λ by $1/\lambda$, this is the ℓ -adic counterpart of the complex hypergeometric sheaf $\mathcal{H}(HD)_{\mathbf{C}}$. The Galois action on each stalk of $\mathcal{H}(HD)_\ell$ is described by its Frobenius traces which are given by hypergeometric character sums expressed either in terms of the period functions $\mathbb{P}(HD)$ over finite fields introduced in [25] or the $H_q(HD)$ -functions introduced by Beukers-Cohen-Mellit in [14] based on McCarthy's work [48]. These will be recalled below.

4.1. Hypergeometric functions over finite fields. Let \mathbf{F}_q be a finite field of odd characteristic and use $\widehat{\mathbf{F}}_q^\times$ to denote the group of multiplicative characters of \mathbf{F}_q^\times . We use ε_q or simply ε to denote the trivial character, and ϕ_q or ϕ to denote the quadratic character. For any $A \in \widehat{\mathbf{F}}_q^\times$, use \bar{A} to

denote its inverse, and extend A to \mathbf{F}_q by setting $A(0) = 0$. For any characters $A_i, B_i \in \widehat{\mathbf{F}_q^\times}$, $i = 1, \dots, n$ with $B_1 = \varepsilon$ in $\widehat{\mathbf{F}_q^\times}$, the $n\mathbb{P}_{n-1}$ -function (cf. [25]) is defined as follows:

$$(30) \quad n\mathbb{P}_{n-1} \left[\begin{array}{cccc} A_1 & A_2 & \cdots & A_n \\ & B_2 & \cdots & B_n \end{array} ; \lambda; q \right] \\ := \prod_{i=2}^n (-A_i B_i(-1)) \cdot \left(\frac{1}{q-1} \sum_{\chi \in \widehat{\mathbf{F}_q^\times}} \binom{A_1 \chi}{\chi} \binom{A_2 \chi}{B_2 \chi} \cdots \binom{A_n \chi}{B_n \chi} \chi(\lambda) + \delta(\lambda) \prod_{i=2}^n \binom{A_i}{B_i} \right),$$

where

$$\binom{A}{B} := -B(-1)J(A, \overline{B}), \quad J(A, B) := \sum_{t \in \mathbf{F}_q} A(t)B(1-t), \quad \text{and} \quad \delta(\lambda) := \begin{cases} 1, & \text{if } \lambda = 0, \\ 0, & \text{otherwise.} \end{cases}$$

Let $\alpha = \{a_1, \dots, a_n\}$, and $\beta = \{1, b_2, \dots, b_n\}$ be multi-sets with entries in \mathbf{Q}^\times . Denote the positive least common denominator of all a_i, b_j by $M := \text{lcd}(\alpha \cup \beta)$. For a finite field \mathbf{F}_q containing a primitive M th root of 1 and any $\lambda \in \mathbf{F}_q$, let ω be a generator of $\widehat{\mathbf{F}_q^\times}$. Following [45], we write

$$(31) \quad \mathbb{P}(\alpha, \beta; \lambda; \mathbf{F}_q; \omega) := n\mathbb{P}_{n-1} \left[\begin{array}{cccc} \omega^{(q-1)a_1} & \omega^{(q-1)a_2} & \cdots & \omega^{(q-1)a_n} \\ & \omega^{(q-1)b_2} & \cdots & \omega^{(q-1)b_n} \end{array} ; \lambda; \mathbf{F}_q \right].$$

Write $G_K = \text{Gal}(\overline{\mathbf{Q}}/K)$ for the absolute Galois group of a number field K . When $K = \mathbf{Q}(\zeta_M)$ where ζ_M is a primitive M th root of unity, we abbreviate $G_{\mathbf{Q}(\zeta_M)}$ by $G(M)$. Note that the Frobenius conjugacy classes Frob_φ over the degree-1 prime ideals φ of $\mathbf{Q}(\zeta_M)$ are dense in $G(M)$, hence irreducible Galois representations of $G(M)$ are uniquely determined by their Frobenius traces on almost all degree-1 prime ideals φ , which are those above the primes $p \equiv 1 \pmod{M}$. Denote by χ_d the quadratic character of $G_{\mathbf{Q}}$ corresponding to the quadratic field $\mathbf{Q}(\sqrt{d})$.

The following frequently used fact concerning extensions of a Galois representation results from the Frobenius reciprocity law.

Theorem 9. *Let K be a finite Galois extension of \mathbf{Q} and σ be a finite dimensional irreducible ℓ -adic representation of G_K . If for every $g \in G_{\mathbf{Q}}$, the conjugate $\sigma^g \cong \sigma$ as G_K -modules, then σ can be extended to a representation of $G_{\mathbf{Q}}$, which is unique up to a linear character of $G_{\mathbf{Q}}$ trivial on G_K .*

Now we describe the stalks of an ℓ -adic hypergeometric sheaf.

Theorem 10 (Katz [36, 37]). *Let ℓ be a prime. Given a primitive hypergeometric datum $HD = \{\alpha, \beta\}$ consisting of $\alpha = \{a_1, \dots, a_n\}$, $\beta = \{1, b_2, \dots, b_n\}$ with $a_i, b_j \in \mathbf{Q}^\times$ and $M := M(HD) = \text{lcd}(\alpha \cup \beta)$, for each $\lambda \in \mathbf{Z}[\zeta_M, 1/M] \setminus \{0\}$ the following hold for the ℓ -adic representation $\rho_{HD, \lambda, \ell}$ of $G(M)$ on the stalk at λ of the hypergeometric sheaf $\mathcal{H}(HD)_\ell$:*

i). $\rho_{HD, \lambda, \ell}$ is unramified at the prime ideals φ of $\mathbf{Z}[\zeta_M, 1/(M\ell)]$ such that its trace at the conjugacy class Frob_φ of the geometric Frobenius at φ in $G(M)$ with residue field κ_φ and norm $N(\varphi) = |\kappa_\varphi|$ is

$$(32) \quad \text{Tr}_{\rho_{HD, \lambda, \ell}}(\text{Frob}_\varphi) = (-1)^{n-1} \omega_\varphi^{(N(\varphi)-1)a_1} (-1) \mathbb{P}(HD; 1/\lambda; \kappa_\varphi; \omega_\varphi),$$

where ω_φ is a generator of $\widehat{\kappa_\varphi^\times}$ satisfying $\omega_\varphi(\zeta_M \pmod{\varphi}) = \zeta_M^{-1}$.

ii). When $\lambda \neq 1$, the stalk $(\mathcal{H}(HD)_\ell)_\lambda$ has dimension n and all roots of the characteristic polynomial of $\rho_{HD, \lambda, \ell}(\text{Frob}_\varphi)$ are algebraic numbers and have the same absolute value $N(\varphi)^{(n-1)/2}$ under all archimedean embeddings.

iii). When $\lambda = 1$, the dimension of $(\mathcal{H}(HD)_\ell)_{\bar{\lambda}}$ equals $n - 1$.

Remark 4. Recall that for $a = r/M$ with $r \in \mathbf{Z}$, $\omega_\varphi^{(N(\varphi)-1)a}$ is the character $\iota_\varphi(a) = \left(\frac{\cdot}{\varphi}\right)_M^r$, the r th power of the M th norm residue symbol at a nonzero unramified prime ideal φ as in Definition 5.9 of [25]. In particular, when φ varies, the sign $\omega_\varphi^{(N(\varphi)-1)a}(-1) = \iota_\varphi(a)(-1)$ defines a character of $G(M)$, denoted by $\phi(M, a)$. As the character $\phi(M, a)$ is invariant under conjugation by $G_{\mathbf{Q}}/G(M)$, it extends to a character of $G_{\mathbf{Q}}$ and our convention here is use the extension with minimal conductor. It is nontrivial only when $\text{ord}_2 M = -\text{ord}_2 a = r \geq 1$. Especially, when $\text{ord}_2 M = -\text{ord}_2 a = 1$, $\phi(M, a)(\text{Frob}_p) = \left(\frac{-1}{p}\right)$ is given by the Legendre symbol at odd primes p .

4.2. Beukers-Cohen-Mellit's finite Hypergeometric functions. For $q \equiv 1 \pmod{M}$, let

$$H_q(\alpha, \beta; \lambda; \omega) := \frac{1}{1-q} \sum_{k=0}^{q-2} \omega^k ((-1)^n \lambda) \prod_{j=1}^n \frac{\mathfrak{g}(\omega^{(q-1)a_j+k})}{\mathfrak{g}(\omega^{(q-1)a_j})} \frac{\mathfrak{g}(\bar{\omega}^{(q-1)b_j+k})}{\mathfrak{g}(\bar{\omega}^{(q-1)b_j})},$$

where $\mathfrak{g}(\chi) := \sum_{x \in \mathbf{F}_q} \Psi(x)\chi(x)$ denotes the Gauss sum of χ with respect to a fixed choice of non-trivial additive character Ψ of \mathbf{F}_q . However the above sum is independent of the choice of Ψ since $|\alpha| = |\beta|$. The function H_q is equivalent to the normalized \mathbb{P} -function (denoted by \mathbb{F} -function in [25]) when the pair α and β is primitive, in which case they are related by

$$(33) \quad \mathbb{P}(\alpha, \beta; \lambda; \mathbf{F}_q; \omega) = H_q(\alpha, \beta; \lambda; \omega) \cdot \prod_{i=2}^n \omega^{(q-1)a_i} (-1) J(\omega^{(q-1)a_i}, \bar{\omega}^{(q-1)b_i}).$$

By Weil [67], as \mathbf{F}_q runs through the residue fields of almost all prime ideals of $\mathbf{Q}(\zeta_M)$, these Jacobi sums define a Grössencharacter of $G(M)$. The above relation indicates that instead of using the \mathbb{P} -functions to describe the trace function of the Galois representations in Theorem 10, one can also use the H -functions.

When $HD = \{\alpha, \beta\}$ and λ are both defined over \mathbf{Q} , one can rewrite $H_q(\alpha, \beta; \lambda; \omega)$ by using the reflection and multiplication formulas of Gauss sums [14, Theorem 1.3]:

$$\mathfrak{g}(A)\mathfrak{g}(\bar{A}) = A(-1)q, \quad \forall A \neq \varepsilon, \quad \prod_{\substack{\eta \in \widehat{\mathbf{F}_q^\times} \\ \eta^m = \varepsilon}} \frac{\mathfrak{g}(A\eta)}{\mathfrak{g}(\eta)} = -\mathfrak{g}(A^m)A(m^{-m}), \quad \text{when } m \mid q-1.$$

Firstly write $\prod_{j=1}^n \frac{X - e^{2\pi i a_j}}{X - e^{2\pi i b_j}} = \frac{\prod_{j=1}^r (X^{p_j} - 1)}{\prod_{k=1}^s (X^{q_k} - 1)}$ where $p_j, q_k \in \mathbf{Z}_{>0}$ and $p_j \neq q_k$ for all j, k . Then

$$(34) \quad H_q(\alpha, \beta; \lambda; \omega) = \frac{(-1)^{r+s}}{1-q} \sum_{m=0}^{q-2} q^{-s(0)+s(m)} \prod_{j=1}^r \mathfrak{g}(\omega^{mp_j}) \prod_{k=1}^s \mathfrak{g}(\omega^{-mq_k}) \omega(N^{-1}\lambda),$$

where $s(m)$ is the multiplicity of $X - e^{\frac{2\pi im}{q-1}}$ in $\gcd(\prod_{j=1}^r (X^{p_j} - 1), \prod_{k=1}^s (X^{q_k} - 1))$, and

$$N = (-1)^{\sum_{k=1}^s q_k} \cdot \frac{p_1^{p_1} \cdots p_r^{p_r}}{q_1^{q_1} \cdots q_s^{q_s}}.$$

In this case H_q is \mathbf{Q} -valued, and hence is independent of the choice of ω , which will be dropped from the left hand side of (34). Also (34) holds now for \mathbf{F}_q as long as q is coprime to $\text{lcd}(\{1/M, \lambda\})$.

The next theorem follows from Theorems 10 and 9.

Theorem 11 (Katz, Beukers-Cohen-Mellit). *Let $HD = \{\alpha = \{a_1, \dots, a_n\}, \beta = \{1, b_2, \dots, b_n\}\}$ be a primitive hypergeometric datum defined over \mathbf{Q} with level $M = M(HD)$. Let $\lambda \in \mathbf{Z}[1/M] \setminus \{0\}$. Assume that exactly m elements in β are in \mathbf{Z} . Then, for each prime ℓ , there exists an ℓ -adic representation $\rho_{HD,\lambda,\ell}^{BCM}$ of $G_{\mathbf{Q}}$, unramified almost everywhere, with the following properties:*

- i). $\rho_{HD,\lambda,\ell}^{BCM}|_{G(M)} \cong \rho_{HD,\lambda,\ell}$.
- ii). For any prime $p \nmid \ell M$ such that $\text{ord}_p \lambda = 0$,

$$(35) \quad \text{Tr } \rho_{HD,\lambda,\ell}^{BCM}(\text{Frob}_p) = \phi(M, \sum_{i=1}^n a_i)(\text{Frob}_p) H_p(HD; 1/\lambda) \cdot p^{(n-m)/2} \in \mathbf{Z},$$

where $\phi(M, a)$, as explained in Remark 4, is a character of $G_{\mathbf{Q}}$ of order at most 2.

- iii). When $\lambda = 1$, $\rho_{HD,\lambda,\ell}^{BCM}$ is $(n-1)$ -dimensional and it has a subrepresentation, denoted by $\rho_{HD,\lambda,\ell}^{BCM, \text{prim}}$, of dimension $2\lfloor \frac{n-1}{2} \rfloor$. For each unramified prime p , all roots of the characteristic polynomial of $\rho_{HD,\lambda,\ell}^{BCM, \text{prim}}(\text{Frob}_p)$ have absolute value $p^{(n-1)/2}$.

Under the assumptions of Theorem 11, the relation (33) at a prime \wp of $\mathbf{Z}[\zeta_M]$ above a prime $p \equiv 1 \pmod{M}$ simplifies as $\mathbb{P}(HD; \lambda; \kappa_{\wp}; \omega_{\wp}) = H_p(HD; \lambda; \omega_{\wp}) p^{\frac{n-m}{2}} \phi(M, \sum_{i=2}^n a_i) (-1)^{n-1}$. This fact is incorporated in the expression of (35).

Under the above assumption, we now clarify the relation between $H_p(\alpha, \beta; 1/\lambda)$ and $H_q(\alpha, \beta; 1/\lambda)$ where $q = p^r$. Let $\phi(M, \sum_{i=1}^n a_i)(\text{Frob}_p) p^{(n-m)/2} u_i, 1 \leq i \leq d$, be the roots of the characteristic polynomial of $\rho_{HD,\lambda,\ell}^{BCM}(\text{Frob}_p)$ so that $H_p(\alpha, \beta; 1/\lambda) = \sum_{i=1}^d u_i$. Here $d = n$ for $\lambda \neq 1$, and $d = n-1$ for $\lambda = 1$. Then $\text{Tr } \rho_{HD,\lambda,\ell}^{BCM}((\text{Frob}_p)^r) = \phi(M, \sum_{i=1}^n a_i)((\text{Frob}_p)^r) q^{(n-m)/2} = \sum_{i=1}^d u_i^r$. On the other hand, let K be a finite Galois extension of \mathbf{Q} unramified at p such that any prime ideal \wp of K above p has residue field \mathbf{F}_q . Then

$$\text{Tr } (\rho_{HD,\lambda,\ell}^{BCM})|_{G_K}(\text{Frob}_{\wp}) = \phi(M, \sum_{i=1}^n a_i)(\text{Frob}_{\wp}) H_q(\alpha, \beta; 1/\lambda) \cdot q^{(n-m)/2}.$$

Since $\text{Frob}_{\wp} = (\text{Frob}_p)^r$, we conclude that $H_{p^r}(\alpha, \beta; 1/\lambda) = \sum_{i=1}^d u_i^r$.

Remark 5. When HD and λ are both defined over \mathbf{Q} , $\rho_{HD,\lambda,\ell}^{BCM}$ is an extension of $\rho_{HD,\lambda,\ell}$ from $G(M)$ to $G_{\mathbf{Q}}$, which is unique up to a character of $G_{\mathbf{Q}}/G(M) \cong \text{Gal}(\mathbf{Q}(\zeta_M)/\mathbf{Q})$ when $\rho_{HD,\lambda,\ell}$ is an irreducible $G(M)$ -module. Thus on each $G(M)$ -coset in $G_{\mathbf{Q}}$, $\rho_{HD,\lambda,\ell}^{BCM}$ is determined by the action at one Frobenius element belonging to that coset. This property is used to extend identities involving the \mathbb{P} -functions to identities involving the H_p -functions.

4.3. For groups in a) list. We now relate the groups of the form $\Gamma = (\infty, \infty, e_{\infty})$ in Figure 1a and Table 2 to elliptic curves. In the table below t is the Hauptmodul of Γ the same as λ in Theorem 3, and $E_{\Gamma,t}$ is a family of elliptic curves over points in X_{Γ} with $t \neq 0, 1, \infty$.

e_{∞}	$E_{\Gamma,t}$	$HD(\Gamma)$	$j_{\Gamma}(t)$	$c_4(\mathcal{E}_{j_{\Gamma}(t)})/c_4(E_{\Gamma,t})$
∞	$y^2 + xy + \frac{t}{16}y = x^3 + \frac{t}{16}x^2$	$\{\{\frac{1}{2}, \frac{1}{2}\}, \{1, 1\}\}$	$\frac{16(t^2 - 16t + 16)^3}{t^4(1-t)}$	$\left(\frac{4(t^2 - 16t + 16)}{(t-2)(t^2 + 32t - 32)}\right)^2$
3	$y^2 + xy + \frac{t}{27}y = x^3$	$\{\{\frac{1}{3}, \frac{2}{3}\}, \{1, 1\}\}$	$\frac{27(8t - 9)^3}{t^3(t-1)}$	$\left(-\frac{3(8t-9)}{(8t^2-36t+27)}\right)^2$

TABLE 4. Elliptic curves for groups in a) list

The universal elliptic curve has been recalled in (25). Both $\mathcal{E}_{j_\Gamma(t)}$ and $E_{\Gamma,t}$ have the same j -invariant and their c_4 -constants quotients are squares. Thus $E_{\Gamma,t}$ is isomorphic to the pullback of the universal family \mathcal{E}_j via the covering map $j_\Gamma(t)$.

Based on [39] by Koike and [53] by Ono (who used a model isogenous to the one above over $\mathbf{Q}(t)$) for the case $e_\infty = \infty$ and [14] by Beukers-Cohen-Mellit for the case $e_\infty = 3$, for $t \in \mathbf{Q} \setminus \{0, 1\}$ and any prime $p > 3$ not dividing the level of $E_{\Gamma,t}$, we have

$$(36) \quad \#E_{\Gamma,t}(\mathbf{F}_p) = p + 1 - H_p(HD(\Gamma); t).$$

For any elliptic curve defined over \mathbf{Q} and prime ℓ , the determinant of the associated ℓ -adic representation of $G_{\mathbf{Q}}$ is the cyclotomic character, denoted by ϵ_ℓ . Putting together, one has

Proposition 4. *Let $\Gamma = (\infty, \infty, e_\infty)$ where $e_\infty \in \{3, \infty\}$. Given $t \in \mathbf{Q} \setminus \{0, 1\}$ and prime ℓ , at any prime $p \neq \ell$ where the 2-dimensional representation $\rho_{HD(\Gamma), 1/t, \ell}^{BCM}$ of $G_{\mathbf{Q}}$ is unramified, the characteristic polynomial of $\rho_{HD(\Gamma), 1/t, \ell}^{BCM}(\text{Frob}_p)$ is $T^2 - H_p(HD(\Gamma); t)T + p$.*

Proposition 4 implies, for each Γ in a) list, the isomorphism between the stalks $(\mathcal{H}(HD(\Gamma))_\ell)_{\frac{1}{t}}$ of the hypergeometric sheaf and $(V^1(\Gamma)_\ell)_{\bar{t}}$ of the automorphic sheaf at $t \in \mathbf{Q} \setminus \{0, 1\}$.

This Proposition also holds for $\Gamma = (\infty, \infty, 2)$ with $E_{\Gamma,t} : y^2 = x(x^2 + x + \frac{t}{4})$ and $HD(\Gamma)' = \{\{\frac{1}{4}, \frac{3}{4}\}, \{1, 1\}\}$ such that at almost all primes p , the characteristic polynomial of $\rho_{HD(\Gamma)', 1/t, \ell}^{BCM}(\text{Frob}_p)$ is $T^2 - H_p(HD(\Gamma)'; t)T + p$, as shown in [14].

Remark 6. In the elliptic modular cases, we can define $V^1(\Gamma)_\ell$ even when Γ contains $-I_2$, as $R^1 f_* \bar{\mathbf{Q}}_\ell$ for a suitable family of elliptic curves $f : E \rightarrow X_\Gamma^\circ$, not unique in general. As Scholl proves, for computing the Hecke traces for modular forms of even weight, any such suitable family of elliptic curves will do. By suitable we mean that the j -invariant of $E \rightarrow X_\Gamma$ is the given map $X_\Gamma \rightarrow X_\Gamma(1)$ and model is defined over $\mathbf{Q}(j)$. See [56, Prop. 1.3].

4.4. For groups in b) list. Throughout the remaining discussion of this section, we assume all hypergeometric data under consideration are primitive for simplicity.

We begin by recalling the character sum analogue of the Clausen formula (26) established by Evans-Greene [22], and rephrased in a way compatible with our current setting in [45]. To ease the notation, we use $J_\omega(a, b)$ to denote $J(\omega^{(q-1)a}, \omega^{(q-1)b})$ below.

Theorem 12 (Evans-Greene [22]). *Let $a, b, s \in \mathbf{Q}$ such that $2s \equiv a + b \pmod{\mathbf{Z}}$. Let q be a prime power which is congruent to 1 modulo $\text{lcd}\{a, b, s, \frac{1}{2} + b - s\}$ and ω a generator of $\widehat{\mathbf{F}}_q^\times$. For any $t \in \mathbf{F}_q \setminus \{0, 1\}$, we have*

$$(37) \quad H_q \left(\left\{ -\frac{1}{2} + b - s, s \right\}, \{1, b\}; t; \omega \right) H_q \left(\left\{ -\frac{1}{2} - b + s, 1 - s \right\}, \{1, 1 - b\}; t; \omega \right) \\ = \phi_q(1 - t) H_q \left(\left\{ \frac{1}{2}, a, 1 - a \right\}, \{1, b, 1 - b\}; t; \omega \right) + q^{\delta(b)},$$

where $\delta(b) = 1$ if $b \in \mathbf{Z}$ and 0 otherwise.

When $t = 1$, we have

$$(38) \quad H_q \left(\left\{ \frac{1}{2}, a, 1 - a \right\}, \{1, b, 1 - b\}; 1; \omega \right) \\ = \frac{1}{q^{1-\delta(b)}} \frac{J_\omega(a + b, b - a)}{J_\omega(\frac{1}{2}, -b)} \left(J_\omega(s - b, \frac{1}{2} - s)^2 + J_\omega(\frac{1}{2} + s - b, s)^2 \right),$$

if $\omega^{(q-1)(a+b)}$ is a square in $\widehat{\mathbf{F}}_q^\times$; otherwise

$$(39) \quad H_q \left(\left\{ \frac{1}{2}, a, 1-a \right\}, \{1, b, 1-b\}; 1; \omega \right) = 0.$$

By Prop. 8.7 of [25], the two H_q -functions on the left side of (37) are related by

$$(40) \quad H_q \left(\left\{ -\frac{1}{2} + b - s, s \right\}, \{1, b\}; t; \omega \right) \\ = \overline{\omega}^{(q-1)b}(t) \phi_q(1-t) \frac{J_\omega(-s, s-b)}{J_\omega(\frac{1}{2}+s, \frac{1}{2}+b-s)} H_q \left(\left\{ -\frac{1}{2} - b + s, 1-s \right\}, \{1, 1-b\}; t; \omega \right).$$

For each group Γ in Theorem 1, the hypergeometric datum $HD(\Gamma) = \{\alpha(\Gamma), \beta(\Gamma)\}$ is self-dual. Thus one can find a (non-unique) pair (a, b) such that $\alpha(\Gamma) = \{\frac{1}{2}, a, 1-a\}$, $\beta(\Gamma) = \{1, b, 1-b\}$. With $s = \frac{a+b-1}{2}$, the hypergeometric datum $HD_1(\Gamma) := \{\{-\frac{1}{2} + b - s, s\}, \{1, b\}\}$ on the left hand side of (37) agrees with the datum associated to the ${}_2F_1$ listed in the second to the last column of Table 3.

Definition 2. For $\lambda \in \mathbf{Q} \setminus \{0, 1\}$, denote by $\chi_{1-1/\lambda}$ the quadratic character $(\frac{1-1/\lambda}{\cdot})_2$ of $G_{\mathbf{Q}}$. It is the $G_{\mathbf{Q}}$ -representation on the stalk at λ of the ℓ -adic sheaf $\mathcal{H}(\{1/2\}, \{1\})_\ell$.

Proposition 5. Fix a prime ℓ . For each group Γ in Table 5 below with $N(\Gamma) = \text{lcm}(4, M(HD_1(\Gamma)))$ and $\lambda \in \mathbf{Q}(\zeta_{N(\Gamma)}) \setminus \{0, 1\}$, let $\sigma_{\Gamma, \lambda, \ell} := \rho_{HD_1(\Gamma), \lambda, \ell}|_{G(N(\Gamma))}$, whose Frobenius trace at each prime ideal \wp of $\mathbf{Z}[\zeta_{N(\Gamma)}, 1/(N(\Gamma)\ell\lambda)]$ above a prime $p \equiv 1 \pmod{N(\Gamma)}$ is given in the table below, in which $J_\wp(HD_1(\Gamma)) := -J_\omega(\frac{7}{24}, -\frac{5}{6})$. Let $\bar{\sigma}_{\Gamma, \lambda, \ell} := \rho_{HD'_1(\Gamma), \lambda, \ell}|_{G(N(\Gamma))}$. Then

- 1) $\sigma_{\Gamma, \lambda, \ell}$ and $\bar{\sigma}_{\Gamma, \lambda, \ell}$ differ by $\chi_{1-1/\lambda}|_{G(N(\Gamma))}$ times a finite order character $\xi_{\Gamma, \lambda, \ell}$;
- 2) $\sigma_{\Gamma, \lambda, \ell} \otimes \bar{\sigma}_{\Gamma, \lambda, \ell}$ contains a 3-dimensional subrepresentation $\chi_{1-1/\lambda}|_{G(N(\Gamma))} \xi_{\Gamma, \lambda, \ell} \otimes \text{Sym}^2 \sigma_{\Gamma, \lambda, \ell}$, which is isomorphic to $\chi_{1-1/\lambda} \otimes \rho_{HD(\Gamma), \lambda, \ell}|_{G(N(\Gamma))}$ with determinant ϵ_ℓ^3 , where ϵ_ℓ is the cyclotomic character on $G(N(\Gamma))$.

Γ	$N(\Gamma)$	$(a(\Gamma), b(\Gamma))$	$HD_1(\Gamma)$	$HD'_1(\Gamma)$	$\text{Tr} \sigma_{\Gamma, \lambda, \ell}(\text{Frob}_\wp)$
$(\infty, 2, \infty)$	4	$(\frac{1}{2}, 1)$	$\{\{\frac{1}{4}, \frac{1}{4}\}, \{1, 1\}\}$	$\{\{\frac{3}{4}, \frac{3}{4}\}, \{1, 1\}\}$	$H_p(HD_1(\Gamma); 1/\lambda; \omega_\wp)$
$(\infty, 2, 3)$	12	$(\frac{1}{6}, 1)$	$\{\{\frac{1}{12}, \frac{5}{12}\}, \{1, 1\}\}$	$\{\{\frac{11}{12}, \frac{7}{12}\}, \{1, 1\}\}$	$H_p(HD_1(\Gamma); 1/\lambda; \omega_\wp)$
$(\infty, 2, 4)$	8	$(\frac{1}{4}, 1)$	$\{\{\frac{1}{8}, \frac{3}{8}\}, \{1, 1\}\}$	$\{\{\frac{7}{8}, \frac{5}{8}\}, \{1, 1\}\}$	$H_p(HD_1(\Gamma); 1/\lambda; \omega_\wp)$
$(\infty, 2, 6)$	12	$(\frac{1}{3}, 1)$	$\{\{\frac{1}{6}, \frac{1}{3}\}, \{1, 1\}\}$	$\{\{\frac{5}{6}, \frac{2}{3}\}, \{1, 1\}\}$	$H_p(HD_1(\Gamma); 1/\lambda; \omega_\wp)$
$(6, 2, 4)$	24	$(\frac{1}{4}, \frac{5}{6})$	$\{\{\frac{1}{24}, \frac{7}{24}\}, \{1, \frac{5}{6}\}\}$	$\{\{\frac{23}{24}, \frac{17}{24}\}, \{1, \frac{7}{6}\}\}$	$J_\wp(HD_1(\Gamma)) \times H_p(HD_1(\Gamma); 1/\lambda; \omega_\wp)$

TABLE 5. σ representations for groups in Table 3

Proof. By Remark 4, the characters $\phi(M(HD_1(\Gamma)), \frac{1}{2})$ and $\phi(M(HD(\Gamma)), \frac{1}{2})$ are trivial on $G(N(\Gamma))$, so they will not show up in the discussion below. Since a prime $p \equiv 1 \pmod{N(\Gamma)}$ splits completely in $\mathbf{Q}(\zeta_{N(\Gamma)})$, the prime ideals \wp above p have norm equal to p . The trace of $\sigma_{\Gamma, \lambda, \ell}(\text{Frob}_\wp)$ follows from (32) and (33). Statement 1) follows from (40) while statement 2) follows from the finite field Clausen formula (Theorem 12). More precisely, when $b = 1$, the quotient of the Jacobi sums on the right hand side of (40) is 1. In the case of $(2, 4, 6)$, letting $a = 1/4$ and $b = 5/6$ in (40) gives

the following trace function for $\bar{\sigma}_{\Gamma,\lambda,\ell}$:

$$\begin{aligned}\mathrm{Tr}\bar{\sigma}_{\Gamma,\lambda,\ell}(\mathrm{Frob}_\varphi) &= -J_\omega\left(\frac{17}{24}, \frac{5}{6}\right) H_p\left(\left\{\frac{17}{24}, \frac{23}{24}\right\}, \left\{1, \frac{7}{6}\right\}; 1/\lambda; \omega_\varphi\right) \\ &= -\phi_p(1-1/\lambda)\bar{\omega}_\varphi(1/\lambda)^{\frac{p-1}{6}} \cdot C(\Gamma, \omega_\varphi) \cdot J_\omega\left(\frac{7}{24}, -\frac{5}{6}\right) H_p(HD_1(\Gamma); 1/\lambda; \omega_\varphi) \\ &= \phi_p(1-1/\lambda)\bar{\omega}_\varphi(1/\lambda)^{\frac{p-1}{6}} \cdot C(\Gamma, \omega_\varphi) \cdot \mathrm{Tr}\sigma_{\Gamma,\lambda,\ell}(\mathrm{Frob}_\varphi)\end{aligned}$$

where $C(\Gamma, \omega_\varphi) = \frac{J_\omega(\frac{17}{24}, \frac{5}{6}) J_\omega(\frac{7}{24}, \frac{13}{24})}{J_\omega(\frac{7}{24}, -\frac{5}{6}) J_\omega(\frac{5}{24}, -\frac{1}{24})}$. By converting $J(A, B) = \mathfrak{g}(A)\mathfrak{g}(B)/\mathfrak{g}(AB)$ when AB is nontrivial and using identities on Gauss sums, one verifies that $C(\Gamma, \omega_\varphi) = 1$.

Thus $\bar{\sigma}_{\Gamma,\lambda,\ell} \cong \xi_{\Gamma,\lambda,\ell} \otimes \chi_{1-1/\lambda}|_{G(N(\Gamma))} \otimes \sigma_{\Gamma,\lambda,\ell}$, where $\xi_{\Gamma,\lambda,\ell}$ is a finite order character of $G_{N(\Gamma)}$; this is 1). Consequently,

$$\sigma_{\Gamma,\lambda,\ell} \otimes \bar{\sigma}_{\Gamma,\lambda,\ell} = \chi_{1-\frac{1}{\lambda}}|_{G(N(\Gamma))} \xi_{\Gamma,\lambda,\ell} \otimes (\sigma_{\Gamma,\lambda,\ell} \otimes \sigma_{\Gamma,\lambda,\ell}) \cong \chi_{1-\frac{1}{\lambda}}|_{G(N(\Gamma))} \xi_{\Gamma,\lambda,\ell} \otimes (\mathrm{Sym}^2 \sigma_{\Gamma,\lambda,\ell} \oplus \mathrm{Alt}^2 \sigma_{\Gamma,\lambda,\ell}),$$

showing that $\sigma_{\Gamma,\lambda,\ell} \otimes \bar{\sigma}_{\Gamma,\lambda,\ell}$ contains the degree-3 subrepresentation $\chi_{1-1/\lambda}|_{G(N(\Gamma))} \xi_{\Gamma,\lambda,\ell} \otimes \mathrm{Sym}^2 \sigma_{\Gamma,\lambda,\ell}$. On the other hand, it follows from the Frobenius traces of $\rho_{HD(\Gamma),\lambda,\ell}$ shown in Table 5 that

$$\begin{aligned}\mathrm{Tr}\sigma_{\Gamma,\lambda,\ell} \otimes \bar{\sigma}_{\Gamma,\lambda,\ell}(\mathrm{Frob}_\varphi) &= p^{1-\delta(b)} H_p(HD_1(\Gamma); 1/\lambda; \omega_\varphi) H_p(HD'_1(\Gamma); 1/\lambda; \omega_\varphi) \\ &= p^{1-\delta(b)} \phi_p(1-1/\lambda) H_p(HD(\Gamma); 1/\lambda; \omega_\varphi) + p \\ &= \chi_{1-1/\lambda}(\mathrm{Frob}_\varphi) \mathrm{Tr}\rho_{HD(\Gamma),\lambda,\ell}(\mathrm{Frob}_\varphi) + \epsilon_\ell(\mathrm{Frob}_\varphi).\end{aligned}$$

In the above formula, the second equality follows from the Clausen formula (37), and the third equality results from Theorem 10 and the relation (33). This proves that $\chi_{1-1/\lambda} \otimes \rho_{HD(\Gamma),\lambda,\ell}|_{G(N(\Gamma))} \cong \chi_{1-1/\lambda}|_{G(N(\Gamma))} \xi_{\Gamma,\lambda,\ell} \otimes \mathrm{Sym}^2 \sigma_{\Gamma,\lambda,\ell}$ and $\epsilon_\ell \cong \chi_{1-1/\lambda}|_{G(N(\Gamma))} \xi_{\Gamma,\lambda,\ell} \otimes \mathrm{Alt}^2 \sigma_{\Gamma,\lambda,\ell}$, which in turn implies that

$$\det(\chi_{1-1/\lambda}|_{G(N(\Gamma))} \xi_{\Gamma,\lambda,\ell} \otimes \mathrm{Sym}^2 \sigma_{\Gamma,\lambda,\ell}) = \det(\chi_{1-1/\lambda} \otimes \rho_{HD(\Gamma),\lambda,\ell}|_{G(N(\Gamma))}) = \epsilon_\ell^3.$$

This proves 2). \square

Proposition 5 is the hypergeometric sheaf version of Proposition 3. For each Γ in Proposition 5, the hypergeometric datum $HD(\Gamma)$ is defined over \mathbf{Q} so that the representation $\rho_{HD(\Gamma),\lambda,\ell}$ for λ in \mathbf{Q}^\times can be extended to $G_{\mathbf{Q}}$; we choose $\rho_{HD(\Gamma),\lambda,\ell}^{BCM}$ as its extension. This defines the ℓ -adic hypergeometric sheaf $\mathcal{H}(HD(\Gamma))_\ell$ on X_Γ . Recall from definition 2 that the characters $\chi_{1-1/\lambda}$ come from the sheaf $\mathcal{H}(\{1/2\}, \{1\})_\ell$ on X_Γ . These representations have \mathbf{Q} -valued traces. Other extensions differ from this by a character of $G_{\mathbf{Q}}/G(N(\Gamma))$. By the Isomorphism Theorem 7 over \mathbf{C} and the Comparison Theorem 8, $\mathcal{H}(\{1/2\}, \{1\})_\ell \otimes \mathcal{H}(HD(\Gamma))_\ell$ (upon changing λ to $1/\lambda$) and $V^2(\Gamma)_\ell$ differ by a character χ_Γ of $G_{\mathbf{Q}}$. Both sheaves are punctually pure of the same weight, which can be seen from Propositions 3 and 5. Thus, by Lemma 3 χ_Γ has finite order. From definition, the traces of $\mathcal{H}(\{1/2\}, \{1\})_\ell \otimes \mathcal{H}(HD(\Gamma))_\ell$ at rational stalks have values in \mathbf{Q} . The traces of $V^2(\Gamma)_\ell$ at rational stalks, which originated in the local zeta functions of abelian varieties as in [40], also have values in \mathbf{Q} (See (19)). Thus the order of χ_Γ is at most 2. We summarize the discussion in the following Corollary.

Corollary 2. $\mathcal{H}(\{1/2\}, \{1\})_\ell \otimes \mathcal{H}(HD(\Gamma))_\ell$ (upon changing λ by $1/\lambda$) and $V^2(\Gamma)_\ell$ differ by a character χ_Γ of order at most 2.

Proposition 6. Let Γ be a group in Table 5 with datum $HD(\Gamma) = \{\alpha(\Gamma), \beta(\Gamma)\}$. Write φ_Γ for the character $\phi(\mathrm{lcd}(HD(\Gamma)), \sum_{a_i \in \alpha(\Gamma)} a_i)$ of $G_{\mathbf{Q}}$ in Theorem 11 ii). For any $\lambda \in \mathbb{Q} \setminus \{0, 1\}$, the determinant of $\varphi_\Gamma \otimes \chi_{1-1/\lambda} \otimes \rho_{HD(\Gamma),\lambda,\ell}^{BCM}$ is ϵ_ℓ^3 for the first 4 groups and is $\chi_{-3} \otimes \epsilon_\ell^3$ for (2, 4, 6).

Proof. Recall that X_Γ is a genus zero curve defined over \mathbf{Q} . Using the model we choose, we identify $X_\Gamma^\circ(\mathbf{C})$ with $\mathbb{P}_{\mathbf{C}}^1 \setminus \{0, 1, \infty\}$. The representation $\chi_{1-1/\lambda} \otimes \rho_{HD(\Gamma), \lambda, \ell}^{BCM}$ is the Galois action on the stalk at λ of the rank-3 sheaf $\mathcal{L}(\Gamma)_\ell \otimes \mathcal{H}(HD(\Gamma))_\ell$, so its determinant, as λ varies, constitutes a rank-1 ℓ -adic sheaf, denoted by $\mathcal{F}(\Gamma)_\ell$, on $\mathbb{P}_{\mathbf{Q}}^1 \setminus \{0, 1, \infty\}$. According to Proposition 5, when restricted to $\mathbb{P}_{\mathbf{Q}(\zeta_{N(\Gamma)})}^1 \setminus \{0, 1, \infty\}$, $\mathcal{F}(\Gamma)_\ell$ is the constant sheaf ϵ_ℓ^3 . Therefore $\mathcal{F}(\Gamma)_\ell$ will be a constant sheaf with value ϵ_ℓ^3 times a character of $G_{\mathbf{Q}}/G(N(\Gamma))$ by Theorem 8. As φ_Γ is also a constant degree-1 sheaf on $\mathbb{P}_{\mathbf{Q}}^1 \setminus \{0, 1, \infty\}$, trivial on $G(N(\Gamma))$ by Remark 4, this proves that the determinant of $\varphi_\Gamma \otimes \chi_{1-1/\lambda} \otimes \rho_{HD(\Gamma), \lambda, \ell}^{BCM}$ is ϵ_ℓ^3 up to a character ψ_λ trivial on $G(N(\Gamma))$ and independent of λ . By Theorem 11 ii) the trace of $\varphi_\Gamma \otimes \chi_{1-1/\lambda} \otimes \rho_{HD(\Gamma), \lambda, \ell}^{BCM}(\text{Frob}_p)$ is $\left(\frac{1-1/\lambda}{p}\right) H_p(HD(\Gamma), 1/\lambda) p^{(3-m)/2}$ for primes $p \nmid N(\Gamma)\ell$. Here m is the number of 1's in $\beta(\Gamma)$.

Let S be the group of characters of $G_{\mathbf{Q}}/G(N(\Gamma)) \cong \text{Gal}(\mathbf{Q}(\zeta_{N(\Gamma)})/\mathbf{Q})$, which is a product of finitely many copies of $\mathbf{Z}/2\mathbf{Z}$. Hence any character in S has order at most 2. We proceed to show that ψ_λ is χ_{-3} for $\Gamma = (2, 4, 6)$ and trivial otherwise. As it is independent of λ , it suffices to choose a convenient λ and compute the determinant of the representation at a few primes using the Magma hypergeometric motive package implemented by Watkins [66] to make the final conclusion.

For $(2, \infty, \infty)$, $N(\Gamma) = 4$, $S = \langle \chi_{\pm 1} \rangle$. It suffices to consider $\lambda = 3$ at $p = 7$; the determinant is 7^3 . Hence ψ_λ is trivial.

For $(2, 3, \infty)$ and for $(2, 6, \infty)$, $N(\Gamma) = 12$, $S = \langle \chi_{\pm 1}, \chi_{\pm 3} \rangle$. It suffices to consider $\lambda = 3$ at $p = 7, 11$; the determinants are 7^3 and 11^3 , respectively. Hence ψ_λ is trivial.

For $(2, 4, \infty)$, $N(\Gamma) = 8$, the final choice is among trivial, $S = \langle \chi_{\pm 1}, \chi_{\pm 2} \rangle$. It suffices to consider $\lambda = 3$ at $p = 5, 7, 11$; the determinants are $5^3, 7^3$ and 11^3 , respectively. Hence ψ_λ is trivial.

For $(2, 4, 6)$, $M(\Gamma) = 24$, $S = \langle \chi_{\pm 1}, \chi_{\pm 2}, \chi_{\pm 3} \rangle$. It suffices to consider $\lambda = 3$ at $p = 5, 7, 11, 13, 17, 19$, and the corresponding determinants are $-5^3, 7^3, -11^3, 13^3, -17^3, 19^3$ from which we conclude $\psi_\lambda = \chi_{-3}$. \square

A straightforward computation shows that the quadratic character φ_Γ in Proposition 6 is equal to χ_{-1} for $\Gamma = (2, \infty, \infty), (2, 3, \infty)$ and $(2, 6, \infty)$, and is trivial for $\Gamma = (2, 4, \infty)$ and $(2, 4, 6)$. It follows from Proposition 3 and the definition of $V^2(\Gamma)_\ell$ as the pushforward of $V^2(U')_\ell$ for a neat normal compact open finite-index subgroup U' of the group U corresponding to Γ that, at a \mathbf{Q} -rational point λ of X_Γ° , $\det \rho^2(\Gamma)_{\lambda, \ell} = \epsilon_\ell^3$. This also can be seen from the fact that corresponding modular forms have trivial central characters. The representation at the stalk at λ of $\mathcal{H}(\{1/2\}, \{1\})_\ell \otimes \mathcal{H}(HD(\Gamma))_\ell$ is $\chi_{1-1/\lambda} \otimes \rho_{HD(\Gamma), \lambda, \ell}^{BCM}$, which is isomorphic to $\chi_\Gamma \otimes \rho^2(\Gamma)_{\lambda, \ell}$ for a character χ_Γ of order ≤ 2 by Corollary 2. The above proposition then determines $\chi_\Gamma^3 = \chi_\Gamma$. We record the conclusion below.

Theorem 13. *Let Γ be a group in Table 5 with datum $HD(\Gamma) = \{\alpha(\Gamma), \beta(\Gamma)\}$ as in Theorem 1. Then, upon changing λ to $1/\lambda$, $\mathcal{H}(\{1/2\}, \{1\})_\ell \otimes \mathcal{H}(HD(\Gamma))_\ell = \chi_\Gamma \otimes V^2(\Gamma)_\ell$ with $\chi_\Gamma = \chi_{-1}$ for $\Gamma = (2, \infty, \infty), (2, 3, \infty)$ and $(2, 6, \infty)$, $\chi_\Gamma = \chi_{-3}$ for $\Gamma = (2, 4, 6)$, and χ_Γ trivial for $\Gamma = (2, 4, \infty)$.*

4.5. Some properties of $H_p(HD(\Gamma); t)$. We explore properties of $H_p(HD(\Gamma); t)$ for Γ in Table 5 according to their type. The case $\Gamma = (2, 4, 6)$ of type b.II) will be treated separately. For the four groups Γ of type b.I), $HD(\Gamma) = \{\{\frac{1}{2}, a(\Gamma), 1 - a(\Gamma)\}, \{1, 1, 1\}\}$ where $a(\Gamma)$ is listed in Table 5. The discussion for these groups will be in terms of $d = d(\Gamma) = 1/a(\Gamma)$, which takes values 2, 3, 4, 6, respectively. Define κ_d as follows as in Table 6.

Lemma 4. *Let Γ , $d = d(\Gamma)$ and κ_d be as in Table 6 and $HD_2(\Gamma) = HD_2(d) := \{\{\frac{1}{d}, \frac{d-1}{d}\}, \{1, 1\}\}$. For any prime $p > 5$, $x \in \mathbf{F}_p \setminus \{0, 1\}$, the following identities hold:*

$$(41) \quad H_p(HD_2(d); x) = \left(\frac{\kappa_d}{p}\right) H_p(HD_2(d); 1-x),$$

Γ	$(\infty, 2, \infty)$	$(\infty, 2, 6)$	$(\infty, 2, 4)$	$(\infty, 2, 3)$
$d = d(\Gamma)$	2	3	4	6
κ_d	-1	-3	-2	-1

TABLE 6. κ_d values

$$(42) \quad H_p(HD_2(d); 1/2) = 0 \quad \text{if} \quad \binom{\kappa_d}{p} = -1.$$

Remark 7. Since $HD_2(\Gamma)$ is defined over \mathbf{Q} , for $\lambda \in \mathbf{Q}^\times$, the representation $\rho_{HD_2(\Gamma), \lambda, \ell}$ of $G(d)$ can be extended to $G_{\mathbf{Q}}$. It follows from Theorem 11 and (42) that $\rho_{HD_2(\Gamma), 2, \ell}^{BCM}$ has CM by $\mathbf{Q}(\sqrt{\kappa_d})$.

Proof. By (9.13) of [18], these identities hold when $p \equiv 1 \pmod{d}$. For $d \in \{3, 4, 6\}$, by Remark 5 both sides agree up to a sign for $p \equiv -1 \pmod{d}$. To decide the sign, we use the following three-step strategy.

Firstly, we relate the H_p function to truncated hypergeometric series. Theorem 4 of [47] says for $d \in \{3, 4, 6\}$ and $x \in \mathbf{F}_p \setminus \{0, 1\}$,

$$H_p \left(\left\{ \frac{1}{d}, \frac{d-1}{d} \right\}, \{1, 1\}; x \right) \equiv {}_2F_1 \left[\begin{matrix} \frac{1}{d} & \frac{d-1}{d} \\ 1 \end{matrix}; x \right]_{p-1} \pmod{p}.$$

Next, we find a terminating series which is p -adically close to the truncated series. Let p_0 be the least positive integer such that $n = \frac{pp_0-1}{d} \in \mathbb{Z}$. Note that $n \equiv -\frac{1}{d} \pmod{p}$ and $(-1)^k \binom{n}{k} = \frac{(-n)_k}{k!} \equiv \frac{(\frac{1}{d})_k}{k!} \pmod{p}$, $\binom{n+k}{k} = \frac{(1+n)_k}{k!} \equiv \frac{(\frac{d-1}{d})_k}{k!} \pmod{p}$ for $0 \leq k < p$. Thus

$${}_2F_1 \left[\begin{matrix} \frac{1}{d} & \frac{d-1}{d} \\ 1 \end{matrix}; x \right]_{p-1} \equiv {}_2F_1 \left[\begin{matrix} -n & n+1 \\ 1 \end{matrix}; x \right] \equiv \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} (-x)^k \pmod{p}.$$

At last we use identities for the terminating series to reach the goal. Presently, recall that

$$\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} (-x)^k = (-1)^n \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} (x-1)^k$$

(see (2.3.11) of [5]). For $d = 3$, when $p = 6k + 1$ (resp. $6k - 1$) for $k \in \mathbf{Z}$, $n = \frac{p-1}{3} = 2k$ (resp. $n = \frac{2p-1}{3} = 4k - 1$). Thus $(-1)^n = \binom{-3}{p}$. The other cases can be verified similarly. \square

The next Lemma is a special version of the Clausen formula that works for almost all q .

Lemma 5. For Γ of type b.I), $d = d(\Gamma) \in \{2, 3, 4, 6\}$, p being any prime not dividing $2d$, and $t \in \mathbf{F}_p \setminus \{0, 1, 1/2\}$, we have

$$(43) \quad H_p(HD_2(\Gamma); t)^2 = H_p \left(\left\{ \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1\}; t \right)^2 = H_p(HD(\Gamma); 4t(1-t)) + p.$$

Proof. By our assumption, $1 - 4t(1 - t) = (1 - 2t)^2 \neq 0, 1$. For $p \equiv 1 \pmod{2d}$, we have

$$\begin{aligned} & H_p \left(\left\{ \frac{1}{2}, \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1, 1\}; 4t(1 - t) \right) + p \\ & \stackrel{\text{Clausen}}{=} H_p \left(\left\{ 1 - \frac{1}{2d}, \frac{1}{2} + \frac{1}{2d} \right\}, \{1, 1\}; 4t(1 - t) \right) H_p \left(\left\{ \frac{1}{2d}, \frac{1}{2} - \frac{1}{2d} \right\}, \{1, 1\}; 4t(1 - t) \right) \\ & = H_p \left(\left\{ \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1\}; t \right)^2. \end{aligned}$$

In the above, we use [25, (9.12)] to get the last equality. For p with $p \not\equiv 1 \pmod{2d}$, the conclusion can be drawn using the three-step strategy outlined in the proof of Lemma 4 that

$$H_p \left(\left\{ \frac{1}{2}, \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1, 1\}; t \right) \equiv {}_3F_2 \left[\begin{matrix} \frac{1}{2} & \frac{1}{d} & \frac{d-1}{d} \\ & 1 & 1 \end{matrix}; t \right]_{p-1} \pmod{p}.$$

and

$${}_2F_1 \left[\begin{matrix} \frac{1}{d} & \frac{d-1}{d} \\ & 1 \end{matrix}; t \right]_{p-1}^2 \equiv {}_3F_2 \left[\begin{matrix} \frac{1}{2} & \frac{1}{d} & \frac{d-1}{d} \\ & 1 & 1 \end{matrix}; 4t(1 - t) \right]_{p-1} \pmod{p^2}.$$

In the last step, Clausen formula over \mathbf{C} is used. \square

4.6. Special fiber at 1. In this subsection we give a description for $H_p(HD(\Gamma); 1)$. This information will be useful for computing the contributions at the special fibers in §5. Firstly, we give an evaluation formula for the length-2 datum at $t = 1/2$.

Lemma 6. *Let $a, s \in \mathbf{Q}^\times$ such that $2s \equiv a \pmod{\mathbf{Z}}$. For any prime p congruent to 1 modulo $\text{lcd}\{a, \frac{1}{2}\}$,*

$$H_p \left(\{a, 1 - a\}, \{1, 1\}; \frac{1}{2} \right) = -\phi_p(-1) \cdot \left(J_\omega(s, \frac{1}{2} - s) + J_\omega(-s, \frac{1}{2} + s) \right),$$

when $2|(p-1)a$; otherwise

$$H_p \left(\{a, 1 - a\}, \{1, 1\}; \frac{1}{2}; \omega \right) = 0.$$

Proof. When $\omega^{(p-1)a}$ is a not square in \widehat{F}_q^\times , the evaluation formula [28, (4.5)] gives

$$H_p \left(\{a, 1 - a\}, \{1, 1\}; \frac{1}{2}; \omega \right) = 0.$$

If $\omega^{(p-1)a}$ is a square in $\widehat{\mathbf{F}}_p^\times$, combining the evaluation formula [28, (4.5)] and the Jacobi sum identities [25, (2.16), Lemma 2.8], we obtain firstly by [28, (4.5)]

$$H_p \left(\{a, 1 - a\}, \{1, 1\}; \frac{1}{2}; \omega \right) = -\omega^{(p-1)a}(2) \cdot \left(J_\omega(s, -a) + J_\omega\left(\frac{1}{2} + s, -a\right) \right)$$

Further by [25, Lem. 2.8], we get

$$\begin{aligned} & H_p \left(\{a, 1 - a\}, \{1, 1\}; \frac{1}{2}; \omega \right) \\ & = -\omega^{(p-1)a}(2) \cdot \left(\omega^{(p-1)s}(-1)J_\omega(s, s) + \omega^{(p-1)(\frac{1}{2}+s)}(-1)J_\omega\left(\frac{1}{2} + s, \frac{1}{2} + s\right) \right) \\ & = -\phi_p(-1) \cdot \left(J_\omega(s, \frac{1}{2} - s) + J_\omega(-s, \frac{1}{2} + s) \right). \end{aligned}$$

\square

Corollary 3. For Γ of type b.I), $d = d(\Gamma)$ and κ_d as in Table 6, and any prime $p \nmid d$, we have

$$(44) \quad H_p(HD(\Gamma); 1) = H_p\left(\left\{\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d}\right\}, \{1, 1, 1\}; 1\right) = 0, \quad \text{if } \left(\frac{\kappa_d}{p}\right) = -1;$$

$$(45) \quad H_p\left(HD_2(\Gamma); \frac{1}{2}\right)^2 = H_p\left(\left\{\frac{1}{d}, \frac{d-1}{d}\right\}, \{1, 1\}; \frac{1}{2}\right)^2 = H_p(HD(\Gamma); 1) + 2p, \quad \text{if } \left(\frac{\kappa_d}{p}\right) = 1.$$

For $\Gamma = (2, 4, 6)$ $H_p(HD(\Gamma); 1) = H_p\left(\left\{\frac{1}{2}, \frac{1}{4}, \frac{3}{4}\right\}, \left\{1, \frac{1}{6}, \frac{5}{6}\right\}; 1\right) = 0$ for primes p with $\left(\frac{-6}{p}\right) = -1$.

Proof. For each of the five groups Γ considered here, $HD(\Gamma)$ is defined over \mathbf{Q} , so that the degree-2 representation $\rho_{HD(\Gamma), 1, \ell}$ can be extended to $\rho_{HD(\Gamma), 1, \ell}^{BCM}$ of $G_{\mathbf{Q}}$. It follows from Theorem 11, ii) that at an unramified odd prime p , we have

$$\text{Tr} \rho_{HD(\Gamma), 1, \ell}^{BCM}(\text{Frob}_p) = \phi(\text{lcd}(HD(\Gamma)), \frac{1}{2})(\text{Frob}_p) p^u H_p(HD(\Gamma); 1)$$

where $u = 0$ (resp. 1) for Γ of type b.I) (resp. b.II)), and $\text{Tr} \rho_{HD_2(\Gamma), 2, \ell}^{BCM}(\text{Frob}_p) = H_p(HD_2(\Gamma); 1/2)$ for Γ of type b.I). Here $HD_2(\Gamma)$ is defined above Table 6.

When Γ is of type b.I), by Clausen formula (38) with $2s \equiv a \pmod{\mathbf{Z}}$ and (39), for each $p \equiv 1 \pmod{2d}$ we have

$$H_p\left(\left\{\frac{1}{2}, a, 1-a\right\}, \{1, 1, 1\}; 1\right) = J_{\omega}\left(s, \frac{1}{2} - s\right)^2 + J_{\omega}\left(-s, \frac{1}{2} + s\right)^2.$$

This shows that the degree-2 representation $\rho_{HD(\Gamma), 1, \ell}^{BCM}$ restricted to $G(2d)$ decomposes into the sum of two distinct characters of $G(2d)$. Thus, by Clifford theory, the 2-dimensional $\rho_{HD(\Gamma), 1, \ell}^{BCM}$ is an irreducible representation induced from a one-dimensional representation ξ of $G(2d)$ corresponding to the first Jacobi sum extended to an index-2 subgroup $H(2d)$ of $G_{\mathbf{Q}}$ generated by $G(2d)$ and all $\{\tau \in G_{\mathbf{Q}}/G(2d) \mid \xi^{\tau} \cong \xi\}$. To decide H , we use Lemma 5 with $a = \frac{1}{d}$ and $s = \frac{1}{2d}$:

$$(46) \quad H_p\left(\left\{a, 1-a\right\}, \{1, 1\}; \frac{1}{2}\right)^2 = J_{\omega}\left(s, \frac{1}{2} - s\right)^2 + J_{\omega}\left(-s, \frac{1}{2} + s\right)^2 + 2p \\ = H_p(HD(\Gamma); 1) + 2p, \quad \forall p \equiv 1 \pmod{2d}.$$

This shows that the representation $\rho_{HD_2(\Gamma), 2, \ell}^{BCM} \otimes \rho_{HD_2(\Gamma), 2, \ell}^{BCM}$, when restricted to $G(2d)$, agrees with the restriction to $G(2d)$ of the sum of the degree-2 representation $\rho_{HD(\Gamma), 1, \ell}^{BCM}$ and two copies of the degree-1 representation ϵ_{ℓ} . Since all representations involved are of $G_{\mathbf{Q}}$ and $\rho_{HD(\Gamma), 1, \ell}^{BCM}$ is irreducible, we see that $\rho_{HD_2(\Gamma), 2, \ell}^{BCM} \otimes \rho_{HD_2(\Gamma), 2, \ell}^{BCM}$ can be decomposed in the same way, except that each irreducible component is now replaced by a suitable twist by a character of $G_{\mathbf{Q}}/G(2d)$. On the other hand, since $\rho_{HD_2(\Gamma), 2, \ell}^{BCM}$ has CM by $\mathbb{Q}(\sqrt{\kappa_d})$ by Remark 7, so do $\rho_{HD_2(\Gamma), 2, \ell}^{BCM} \otimes \rho_{HD_2(\Gamma), 2, \ell}^{BCM}$ and its only degree-2 irreducible component. Therefore $\rho_{HD(\Gamma), 1, \ell}^{BCM}$ admits the same CM. This proves (44).

The identity (45) can be drawn by checking a few primes representing other cosets of $G(2d)$ in $G_{\mathbf{Q}}$. For instance for $d = 2$, we only need to check one unramified prime in $p \equiv 3 \pmod{4}$ case such as $p = 7$.

For $\Gamma = (2, 4, 6)$, $\rho_{HD(\Gamma), 1, \ell}$ is a representation of $G(12)$. If $p \equiv 1 \pmod{24}$, (38) gives

$$pH_p\left(\left\{\frac{1}{2}, \frac{1}{4}, \frac{3}{4}\right\}, \left\{1, \frac{1}{6}, \frac{5}{6}\right\}; 1\right) = \frac{J_{\omega}\left(\frac{5}{12}, \frac{11}{12}\right)}{J_{\omega}\left(\frac{1}{2}, \frac{5}{6}\right)} \left(J_{\omega}\left(\frac{1}{24}, \frac{7}{24}\right)^2 + J_{\omega}\left(\frac{13}{24}, \frac{19}{24}\right)^2 \right).$$

By the same argument as above, the representation $\rho_{HD(\Gamma),1,\ell}^{BCM}$ of $G_{\mathbf{Q}}$ is invariant under the twist by a quadratic character of $G_{\mathbf{Q}}$. To identify this quadratic character, it suffices to compute the values of $H_p(\{\frac{1}{2}, \frac{1}{4}, \frac{3}{4}\}, \{1, \frac{1}{6}, \frac{5}{6}\}; 1)$ for $p = 5, 7, 11, 13, 17, 19, 23$ which reveals that for these cases the value is 0 if and only if $\left(\frac{-6}{p}\right) = -1$. \square

Remark 8. The degree-2 representations $\rho_{HD(\Gamma),1,\ell}^{BCM}$ of $G_{\mathbf{Q}}$ in Corollary 3 have CM. Hence they are modular. In Table 7 below we list the corresponding weight-3 cuspidal Hecke eigenforms, identified using a result of Serre (see [45, Theorem 10]). Similarly for $\rho_{HD_2(\Gamma),2,\ell}^{BCM}$ for Γ of type b.I) in Lemma 4.

Γ	$H_p(HD_2(\Gamma); \frac{1}{2})$	$\text{Tr} \rho_{HD(\Gamma),1,\ell}^{BCM}(\text{Frob}_p)$	$\det \rho_{HD(\Gamma),1,\ell}^{BCM}$
$(2, \infty, \infty)$	$a_p(f_{64.2.a.a})$	$\left(\frac{-1}{p}\right) H_p(\{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\}, \{1, 1, 1\}; 1) = a_p(\eta(4\tau)^6)$	$\chi_{-1}\epsilon_{\ell}^2$
$(2, 3, \infty)$	$a_p(f_{36.2.a.a})$	$\left(\frac{-1}{p}\right) H_p(\{\frac{1}{2}, \frac{1}{3}, \frac{2}{3}\}, \{1, 1, 1\}; 1) = a_p(f_{48.3.e.a})$	$\chi_{-3}\epsilon_{\ell}^2$
$(2, 4, \infty)$	$a_p(f_{256.2.a.a})$	$H_p(\{\frac{1}{2}, \frac{1}{4}, \frac{3}{4}\}, \{1, 1, 1\}; 1) = a_p(f_{8.3.d.a})$	$\chi_{-2}\epsilon_{\ell}^2$
$(2, 6, \infty)$	$a_p(f_{676.2.a.a})$	$\left(\frac{-1}{p}\right) H_p(\{\frac{1}{2}, \frac{1}{6}, \frac{5}{6}\}, \{1, 1, 1\}; 1) = a_p(f_{144.3.g.a})$	$\chi_{-1}\epsilon_{\ell}^2$
$(2, 4, 6)$	—	$p H_p(\{\frac{1}{2}, \frac{1}{4}, \frac{3}{4}\}, \{1, \frac{1}{6}, \frac{5}{6}\}; 1) = a_p(f_{24.3.h.a})$	$\chi_{-6}\epsilon_{\ell}^2$

TABLE 7. $H_p(HD(\Gamma); 1)$

The identities in the two remaining subsections will be used in §6 to identify the character χ_{Γ} alluded to by Theorem 8 as mentioned in §3.6.

4.7. Some character sum identities for the $k = 1$ cases.

Lemma 7. For any odd prime $p > 3$, we have

$$1 + \sum_{t \in \mathbf{F}_p^{\times}} H_p\left(\left\{\frac{1}{2}, \frac{1}{2}\right\}, \{1, 1\}; t\right) = 0; \quad H_p\left(\left\{\frac{1}{2}, \frac{1}{2}\right\}, \{1, 1\}; 1\right) = \left(\frac{-1}{p}\right).$$

$$1 + \sum_{t \in \mathbf{F}_p^{\times}} H_p\left(\left\{\frac{1}{3}, \frac{2}{3}\right\}, \{1, 1\}; t\right) = 0; \quad H_p\left(\left\{\frac{1}{3}, \frac{2}{3}\right\}, \{1, 1\}; 1\right) = \left(\frac{-3}{p}\right).$$

Proof. By definition,

$$H_p\left(\left\{\frac{1}{2}, \frac{1}{2}\right\}, \{1, 1\}; t\right) = \frac{1}{1-p} \sum_{\chi \in \widehat{\mathbf{F}_p^{\times}}} G(\chi) \chi(t), \quad \text{where } G(\chi) = \frac{\mathfrak{g}(\phi\chi)^2}{\mathfrak{g}(\phi)^2} \mathfrak{g}(\bar{\chi})^2.$$

It follows from the orthogonality result $\sum_{t \in \mathbf{F}_p^{\times}} \chi(t) = \begin{cases} p-1, & \text{if } \chi = \varepsilon, \\ 0, & \text{if } \chi \neq \varepsilon, \end{cases}$ that

$$\sum_{t \in \mathbf{F}_p^{\times}} H_p\left(\left\{\frac{1}{2}, \frac{1}{2}\right\}, \{1, 1\}; t\right) = \frac{1}{1-p} \sum_{\chi \in \widehat{\mathbf{F}_p^{\times}}} G(\chi) \sum_{t \in \mathbf{F}_p^{\times}} \chi(t) = -G(\varepsilon) = -1.$$

When $t = 1$, by [25, (Eqn (4.5))]

$$H_p\left(\left\{\frac{1}{2}, \frac{1}{2}\right\}, \{1, 1\}; 1\right) = -\left(\frac{-1}{p}\right) \mathbb{P}\left(\left\{\frac{1}{2}, \frac{1}{2}\right\}, \{1, 1\}; 1; \mathbf{F}_p\right) = \left(\frac{-1}{p}\right).$$

Similarly for any prime $p \geq 3$, $1 + \sum_{t \in \mathbf{F}_p^\times} H_p \left(\left\{ \frac{1}{3}, \frac{2}{3} \right\}, \{1, 1\}; t \right) = 0$ with

$$H_p \left(\left\{ \frac{1}{3}, \frac{2}{3} \right\}, \{1, 1\}; t \right) = \frac{1}{1-p} \sum_{\chi \in \widehat{\mathbf{F}_p^\times}} \frac{\mathfrak{g}(\chi^3)}{\mathfrak{g}(\chi)} \mathfrak{g}(\bar{\chi}) \chi(3^{-3}t).$$

When $p \equiv 1 \pmod{3}$,

$$H_p \left(\left\{ \frac{1}{3}, \frac{2}{3} \right\}, \{1, 1\}; 1 \right) = -\mathbb{P} \left(\left\{ \frac{1}{3}, \frac{2}{3} \right\}, \{1, 1\}; 1; \mathbf{F}_p \right) = 1.$$

For primes $p \equiv 5 \pmod{6}$, by Theorem 9 it suffices to check one prime in the class. For $p = 5$, $H_p \left(\left\{ \frac{1}{3}, \frac{2}{3} \right\}, \{1, 1\}; 1 \right) = -1$. Thus the one dimensional representation $\rho_{\left\{ \frac{1}{3}, \frac{2}{3} \right\}, \{1, 1\}, 1, \ell}$ is isomorphic to χ_{-3} . \square

4.8. Some character sum identities for the $k = 2$ cases.

Lemma 8. *For each prime $p \geq 5$, the following identities hold:*

$$\begin{aligned} \Gamma = (2, 4, 6) : \quad & \sum_{t \in \mathbf{F}_p^\times} \phi(-3(1-t)) H_p \left(\left\{ \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \right\}, \left\{ 1, \frac{1}{6}, \frac{5}{6} \right\}; t \right) + \left(\binom{-6}{p} + \binom{-1}{p} + \binom{-3}{p} \right) = 0. \\ \Gamma = (2, 4, \infty) : \quad & \sum_{t \in \mathbf{F}_p^\times} \phi(1-t) H_p \left(\left\{ \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \right\}, \{1, 1, 1\}; t \right) + 1 + \left(\binom{-2}{p} + \binom{-1}{p} \right) p = 0. \\ \Gamma = (2, 6, \infty) : \quad & \sum_{t \in \mathbf{F}_p^\times} \phi(1-t) H_p \left(\left\{ \frac{1}{2}, \frac{1}{3}, \frac{2}{3} \right\}, \{1, 1, 1\}; t \right) + 1 + 2 \binom{-3}{p} p = 0. \\ \Gamma = (2, \infty, \infty) : \quad & \sum_{t \in \mathbf{F}_p^\times} \phi(1-t) H_p \left(\left\{ \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \right\}, \{1, 1, 1\}; t \right) + 2 + \binom{-1}{p} p = 0. \\ \Gamma = (2, 3, \infty) : \quad & \sum_{t \in \mathbf{F}_p^\times} \phi(1-t) H_p \left(\left\{ \frac{1}{2}, \frac{1}{6}, \frac{5}{6} \right\}, \{1, 1, 1\}; t \right) + 1 + \left(\binom{-3}{p} + \binom{-1}{p} \right) p = 0. \end{aligned}$$

Proof. We give a proof for the (2, 4, 6)-case. The other cases can be proved similarly. For $\chi \in \widehat{\mathbf{F}_p^\times}$,

$$(47) \quad \sum_{s, t \in \mathbf{F}_p} \phi((1-t)s(s-1)) \chi(st) = \phi(-1) J(\phi, \chi) J(\phi\chi, \phi) = \begin{cases} 1 & \text{if } \chi^2 = \varepsilon \\ p & \text{otherwise.} \end{cases}$$

$$H_p \left(\left\{ \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \right\}, \left\{ 1, \frac{1}{6}, \frac{5}{6} \right\}; t \right) = \frac{1}{p-1} \sum_s \phi(-s(1-s)) \sum_\chi \frac{\mathfrak{g}(\chi^4) \mathfrak{g}(\bar{\chi}) \mathfrak{g}(\bar{\chi}^6)}{\mathfrak{g}(\chi^2) \mathfrak{g}(\bar{\chi}^2) \mathfrak{g}(\bar{\chi}^3)} \chi(27st/4),$$

where we apply that $\frac{\mathfrak{g}(\phi\chi)\mathfrak{g}(\bar{\chi})}{\mathfrak{g}(\phi)} = J(\phi\chi, \bar{\chi}) = \phi(-1) J(\phi\chi, \phi) = \sum_{s \in \mathbf{F}_p} \phi(-s(1-s)) \chi(s)$. Thus,

$$\begin{aligned} & \sum_{t \in \mathbf{F}_p^\times} \phi(1-t) H_p \left(\left\{ \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \right\}, \left\{ 1, \frac{1}{6}, \frac{5}{6} \right\}; t \right) \\ &= \frac{1}{p-1} \sum_{t \in \mathbf{F}_p} \phi(1-t) \sum_{s \in \mathbf{F}_p} \phi(s(s-1)) \sum_\chi \frac{\mathfrak{g}(\chi^4) \mathfrak{g}(\bar{\chi}) \mathfrak{g}(\bar{\chi}^6)}{\mathfrak{g}(\chi^2) \mathfrak{g}(\bar{\chi}^2) \mathfrak{g}(\bar{\chi}^3)} \chi(27st/4) \\ &= \frac{1}{p-1} (1-p) \left(1 + \binom{3}{p} + p H_p \left(\left\{ \frac{1}{4}, \frac{3}{4} \right\}, \left\{ \frac{1}{6}, \frac{5}{6} \right\}; 1 \right) \right). \end{aligned}$$

It remains to prove

$$p \cdot H_p \left(\left\{ \frac{1}{4}, \frac{3}{4} \right\}, \left\{ \frac{1}{6}, \frac{5}{6} \right\}; 1 \right) = \frac{p}{1-p} \sum_{\chi \in \widehat{\mathbf{F}_p^\times}} \frac{\mathfrak{g}(\chi^4)\mathfrak{g}(\bar{\chi})\mathfrak{g}(\bar{\chi}^6)}{\mathfrak{g}(\chi^2)\mathfrak{g}(\bar{\chi}^2)\mathfrak{g}(\bar{\chi}^3)} \chi_{(27/4)} = \left(\frac{2}{p} \right).$$

Let $HD := \{ \{ \frac{1}{4}, \frac{3}{4} \}, \{ \frac{1}{6}, \frac{5}{6} \} \}$, which has level 12. Since HD is defined over \mathbf{Q} , the 1-dimensional representation $\rho_{HD,1,\ell}$ of $G(12)$ extends to a character $\rho_{HD,1,\ell}^{BCM}$ of $G_{\mathbf{Q}}$ satisfying $\rho_{HD,1,\ell}^{BCM}(\text{Frob}_p) = p H_p(HD; 1)$ at primes $p \neq 2, 3$ by Theorem 11 ii). The desired identity can then be reinterpreted as the equality between two characters $\rho_{HD,1,\ell}^{BCM}$ and $\chi_2 = \left(\frac{2}{\cdot} \right)$. For $p \equiv 1 \pmod{12}$, this follows from the identity of Helversen-Pasotto, see (8.1) of [25]. For the remaining primes, it suffices to check the equality at primes $p = 5, 7, 11$, which indeed holds. Therefore,

$$\sum_{t \in \mathbf{F}_p^\times} \phi(1-t) H_p \left(\left\{ \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \right\}, \left\{ 1, \frac{1}{6}, \frac{5}{6} \right\}; t \right) = -1 - \left(\frac{3}{p} \right) - \left(\frac{2}{p} \right),$$

which is equivalent to the first identity stated in the Lemma. \square

5. CONTRIBUTIONS FROM THE SINGULAR FIBERS

5.1. The CM structure at elliptic points. Let Γ be a group in Theorem 1 or Theorem 2. In this section we compute the contributions of Frobenius traces on stalks at elliptic points and cusps of Γ whenever they exist. Recall from Proposition 3 that our sheaves, over the complement X_Γ° of the cusps and elliptic points, are in the form $f_*^G(\mathcal{L}^k \otimes \text{Sym}^{2k}(\sigma))$ for a finite Galois G -covering $f : Y \rightarrow X_\Gamma = X$, where σ (resp. \mathcal{L}) is an ℓ -adic rank 2 (resp. rank 1) local system on $Y^\circ = f^{-1}(X^\circ)$. The sheaf σ arises from a family of abelian varieties: in the elliptic modular cases, these are families of elliptic curves; in the quaternion cases we obtain σ from the H^1 of the universal family of abelian varieties with quaternion structure by utilizing Proposition 2. In fact, \mathcal{L} is a finite order character on each stalk, and this is absent if $Y = X_{\Gamma'}$ where Γ' is torsion-free, and we are not considering extension of the field of definition of σ .

In principle, all information about cusps or elliptic points can be obtained from Tate's algorithm or its generalizations by Liu, [46], and by Bouw and Wewers, [16]. By ramified base extension we obtain a family of abelian varieties which has stable reduction (see [15] for a general reference). In the elliptic modular cases, the Néron fiber at a cusp will be of multiplicative type. At the elliptic points, in both the elliptic modular and quaternion cases, we have potential good reduction (see [59, Section 2]), so that the stable fiber is an abelian variety.

Cusps only occur in the elliptic modular cases. By Tate's algorithm, we have multiplicative or potentially multiplicative reduction, and the trace of Frobenius on the stalk of $V^k(\Gamma)_\ell$ at a cusp is 1 for k even. For k odd and $-id \notin \Gamma$, the contribution is ± 1 or 0.

To facilitate our discussion at elliptic points, we recall the moduli interpretation of the curves X_Γ when Γ arises from an indefinite quaternion algebra B defined over \mathbf{Q} . These results are due to Shimura and are summarized in the first chapter of Jordan's thesis [35], on which our discussion below is based. We fix an identification $\varphi : B \otimes_{\mathbf{Q}} \mathbf{R} = M_2(\mathbf{R})$. The group B^+ of elements in B with positive reduced norm acts on \mathfrak{H} via $\varphi(B^+)$. Let X_B be the Shimura canonical model of the Riemann surface $O_B^1 \backslash \mathfrak{H}$. The points $X_B(\mathbf{C}) = O_B^1 \backslash \mathfrak{H}$ are in canonical bijection with the equivalence classes of (A, i, Θ) , where A is a 2-dimensional abelian variety, $i : O_B \rightarrow \text{End}(A)$ is an embedding and Θ is a principal polarization. Then $(A, i, \Theta) \mapsto (A, \Theta)$ defines a map from X_B to the moduli space of principally polarized abelian surfaces, whose explicit form is described by Hashimoto in [30].

Concretely, for each $z \in \mathfrak{H}$ let

$$L_z = \varphi(O_B) \begin{pmatrix} z \\ 1 \end{pmatrix} \subset \mathbf{C}^2 \quad \text{and} \quad A_z = \mathbf{C}^2 / L_z.$$

Then A_z is an abelian surface and the elements $b \in O_B$ define endomorphisms of A_z by sending $w \in \mathbf{C}^2$ to $\varphi(b)w$ which carries the lattice L_z to itself. This defines $i_z : B \rightarrow \text{End}(A_z)$.

Let $\text{End}(A, i, \Theta)$ be the endomorphisms of this structure, and $\text{End}^0(A, i, \Theta) = \text{End}(A, i, \Theta) \otimes \mathbf{Q}$. Then it follows that

$$\text{End}(A, i, \Theta) = \text{End}_{O_B}(A) = \{\gamma \in \text{End}(A) \mid \gamma i(b) = i(b)\gamma \text{ for all } b \in O_B\}$$

and similarly for $\text{End}^0(A, i, \Theta)$.

For any $z \in \mathfrak{H}$ we see that $\text{End}(A_z, i_z, \Theta_z)$ consists of $M \in M_2(\mathbf{C})$ that commute with the image of $\varphi(O_B)$. This forces M to be a scalar matrix $\alpha \mathbf{1}_2$ for some $\alpha \in \mathbf{C}$ such that $\alpha L_z \subset L_z$. Now if $z = z_x$ is a fixed point of an element x of B^+ , by multiplying by an element of \mathbf{Q}^\times we can assume that $x \in O_B^+$, the set of elements in O_B with positive reduced norm. Write

$$\varphi(x) = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad x \in O_B^+$$

and let $\mu(x) = cz + d$. Then multiplication by $\mu(x)$ on \mathbf{C}^2 maps $L_z \rightarrow L_z$ because of

$$\mu(x)\varphi(O_B) \begin{pmatrix} z \\ 1 \end{pmatrix} = \varphi(O_B)(cz+d) \begin{pmatrix} \frac{az+b}{cz+d} \\ 1 \end{pmatrix} = \varphi(O_B) \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} z \\ 1 \end{pmatrix} = \varphi(O_B)\varphi(x) \begin{pmatrix} z \\ 1 \end{pmatrix} = \varphi(O_B x) \begin{pmatrix} z \\ 1 \end{pmatrix}.$$

Since x is not fixed by any parabolic elements in B^+ (if they exist), the map $x \mapsto \mu(x)$ is injective. We have shown:

Proposition 7. *For all $z \in \mathfrak{H}$ there is an embedding*

$$\text{Stab}_{O_B^+}(z) \hookrightarrow \text{End}(A_z, i_z, \Theta_z)$$

by sending $x \in \text{Stab}_{O_B^+}(z)$ with $\varphi(x) = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ to $\mu(x) = cz + d$.

For a general point z we have $\text{End}(A_z, i_z, \Theta_z) = \mathbf{Z}$. We say z is a CM point if there is an embedding $q : K \rightarrow B$, where K is an imaginary quadratic field, such that z is the fixed point of $q(K) \cap O_B^+$. An argument similar to the above shows that we have an embedding $K \subset \text{End}^0(A_z, i_z, \Theta_z)$. Then $\text{End}(A_z, i_z, \Theta_z)$ is an order in K . In fact K is a subfield of B which splits B . By Hasse's criterion, a quadratic field K splits B if and only if every prime which ramifies in B fails to split in K .

Proposition 8. *If the QM abelian surface (A, i, Θ) has CM by the imaginary quadratic field K , then A is isogenous to $E \times E$ where E is an elliptic curve with CM by K .*

The space $V_{\mathbf{Q}} = H^1(A_z, \mathbf{Q})$ is a 4-dimensional \mathbf{Q} -vector space, but because of the quaternion structure this becomes a 2-dimensional module over every quadratic subfield $F \subset B$. Fix a real subfield F ; this defines an isomorphism $B \otimes_{\mathbf{Q}} \mathbf{R} = M_2(\mathbf{R})$.

Now suppose (A_z, i_z, Θ_z) has CM by K . Because the endomorphisms in $\text{End}^0(A_z, i_z, \Theta_z)$ commute with B , they commute with F and we have an embedding

$$K \subset \text{End}^0(A_z, i_z, \Theta_z) \rightarrow \text{End}_F(V_{\mathbf{Q}}) \sim M_2(F).$$

One shows that the subalgebra of $M_2(F)$ commuting with K is K itself. For z corresponding to a point in $X_B(\overline{\mathbf{Q}})$, A_z has a model defined over a number field, say, $k(z)$. If L is a number field such that the $\text{Gal}(\overline{L}/L)$ action on the ℓ -adic Tate-module $H^1(A_z \otimes_{k(z)} \overline{\mathbf{Q}}, \mathbf{Q}_\ell) = V_{\mathbf{Q}} \otimes_{\mathbf{Q}} \mathbf{Q}_\ell = V_\ell$ commutes with the induced action of $\text{End}(A_z)$, we get a representation

$$\text{Gal}(\overline{L}/L) \rightarrow (K \otimes \mathbf{Q}_\ell)^\times \subset \text{GL}_{F \otimes \mathbf{Q}_\ell}(V_{\mathbf{Q}} \otimes_{\mathbf{Q}} \mathbf{Q}_\ell) = \text{GL}_{F \otimes \mathbf{Q}_\ell}(V_\ell) \sim \text{GL}_2(F \otimes \mathbf{Q}_\ell).$$

Because this is an abelian representation, this ℓ -adic representation is given by a pair of characters conjugated by suitable liftings of $\text{Gal}(K/\mathbf{Q})$. In fact, these are the characters giving the Galois action on the Tate module of the elliptic curve E_z such that A_z is isogenous to $E_z \times E_z$.

We now describe the Atkin-Lehner involutions. Let Γ be a finite index subgroup of O_B^1 and Γ^* be the normalizer of Γ in O_B^+ . For each $b \in \Gamma^*$ we normalize it by dividing by the positive square root of its reduced norm $n(b)$ so that $\tilde{b} := \frac{1}{\sqrt{n(b)}}b$ is an element in $(B \otimes_{\mathbf{Q}} \mathbf{R})^\times$ with reduced norm

1. The collection of \tilde{b} forms a group

$$\Gamma^+ = \left\{ \tilde{b} = \frac{1}{\sqrt{n(b)}}b \mid b \in \Gamma^* \right\}$$

which contains $\{\pm id\}\Gamma$ as a normal subgroup of finite index. In fact, the quotient $\Gamma^+/\{\pm id\}\Gamma$ is an elementary 2-group. Each non-identity element \tilde{b} in $\Gamma^+/\{\pm id\}\Gamma$ is called an Atkin-Lehner involution. Sometimes, by abuse of language, we will also call b an Atkin-Lehner involution.³ The discussion of fixed points of Atkin-Lehner involutions can be found in Ogg's paper [49].

When B does not split and $\Gamma = O_B^1$, the group $\Gamma^+/\Gamma \simeq (\mathbf{Z}/2\mathbf{Z})^r$, where r is the number of distinct primes dividing the discriminant D_B of B . For each $d > 1$ dividing D_B , there is a $b_d \in \Gamma^*$ with reduced norm d , and the Atkin-Lehner involutions of O_B^1 are \tilde{b}_d , $d|D_B$ with $d > 1$. The action b_d also has a moduli interpretation.

In what follows we restrict our attention to the quaternion algebras B pertaining to the groups considered in this paper. There are two cases: $B = B_6$ and $B = M_2(\mathbf{Q})$. The groups appearing in Theorem 1 are contained in $\text{PSL}_2(\mathbf{R})$. For a subgroup Γ of $\text{PSL}_2(\mathbf{R})$, denote by $\pm\Gamma$ its preimage in $\text{SL}_2(\mathbf{R})$.

First consider the case $B = B_6 = \left(\frac{-1,3}{\mathbf{Q}}\right)$ with discriminant 6 and a maximal order $O_{B_6} = \mathbf{Z} + \mathbf{Z}I + \mathbf{Z}J + \mathbf{Z}\frac{1+I+J+IJ}{2}$, where $I^2 = -1$, $J^2 = 3$, $IJ = -JI$. Fix the embedding $\varphi : B_6 \otimes_{\mathbf{Q}} \mathbf{R} \rightarrow M_2(\mathbf{R})$ given by $\varphi(I) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $\varphi(J) = \begin{pmatrix} \sqrt{3} & 0 \\ 0 & -\sqrt{3} \end{pmatrix}$.

The group $\pm(2, 4, 6)$ is generated by s_2 , s_4 and s_6 given in (29). It comes from Γ^+ for $\Gamma = O_B^1 = \pm(2, 2, 3, 3)$. The points $z = i$, $\frac{1+i}{\sqrt{3}+1}$, $\frac{\sqrt{6}i}{3+\sqrt{3}}$ in the upper half-plane correspond to the \mathbf{Q} -rational elliptic points on $X_{(2,4,6)}$ of order 4, 6, 2, respectively. These are CM points stabilized by $b_2 = 1 + I$, $b_3 = (3 + 3I + J + IJ)/2$ and $b_6 = 3I + IJ$ with respective reduced norm 2, 3, 6; these stabilizers induce the Atkin-Lehner involutions w_2, w_3, w_6 on the curve $X_{(2,2,3,3)}$ explained before. The normalized stabilizers are the Atkin-Lehner involutions $\tilde{b}_2, \tilde{b}_3, \tilde{b}_6$ in Γ^+ with respective order twice that of the elliptic points they fixed. Via $\varphi(\tilde{b}_2) = s_2$, $\varphi(\tilde{b}_3) = s_4$, $\varphi(\tilde{b}_6) = s_6$, they are viewed as elements in $\pm(2, 4, 6)$, and as such they generate the stabilizers of the respective elliptic points in $\pm(2, 4, 6)$. We proceed to examine the structure of the three abelian surfaces $A_z = \mathbf{C}^2/\varphi(O_{B_6}) \begin{pmatrix} z \\ 1 \end{pmatrix}$

in more detail. Below for each positive integer n , let $\zeta_n = e^{2\pi i/n}$.

1. $z = i$. Consider the sublattice of finite index $L = (\mathbf{Z} + \mathbf{Z}I) + (\mathbf{Z} + \mathbf{Z}I)J \subset O_B$. It defines an abelian variety $B_z := \mathbf{C}^2/\varphi(L) \begin{pmatrix} z \\ 1 \end{pmatrix}$ which is isogenous to A_z . We claim that B_z is isomorphic to $E_z \times E_z$, where E_z is the elliptic curve $\mathbf{C}/\mathbf{Z}[i]$, which has complex

³The group $\Gamma^+/\{\pm id\}$ is isomorphic to the group $N_B(\Gamma) := \{b \in B^+ \mid b\Gamma = \Gamma b\}/\mathbf{Q}^\times$, which contains $P(\Gamma) := \Gamma/(\Gamma \cap \{\pm id\})$ as a normal subgroup. The nontrivial elements in the quotient $N_B(\Gamma)/P(\Gamma)$ are called Atkin-Lehner involutions traditionally. This agrees with our definition because $N_B(\Gamma)/P(\Gamma) \simeq \Gamma^+/\{\pm id\}\Gamma$.

multiplication by $\mathbf{Z}[i]$. To see this note the following

$$\begin{aligned}\varphi((\mathbf{Z} + \mathbf{Z}I)) \begin{pmatrix} i \\ 1 \end{pmatrix} &= (\mathbf{Z} + \mathbf{Z}i) \begin{pmatrix} i \\ 1 \end{pmatrix} \\ \varphi((\mathbf{Z} + \mathbf{Z}I)J) \begin{pmatrix} i \\ 1 \end{pmatrix} &= (\mathbf{Z} + \mathbf{Z}i)\varphi(J) \begin{pmatrix} i \\ 1 \end{pmatrix}.\end{aligned}$$

The vectors $\begin{pmatrix} i \\ 1 \end{pmatrix}, \varphi(J) \begin{pmatrix} i \\ 1 \end{pmatrix}$ in \mathbf{C}^2 are \mathbf{C} -linearly independent. Therefore B_z is isomorphic to $E_z \times E_z$ where $E_z = \mathbf{C}/\mathbf{Z}[i]$ is defined over \mathbf{Q} with CM by $\mathbf{Z}[i]$.

In this case $b_2 = 1 + I$ acts on A_z and B_z as multiplication by $\mu(b_2) = 1 + i = \sqrt{2}\zeta_8$. Consequently, for $\ell > 2$, it induces an automorphism on the ℓ -adic Tate module of A_z . As $H^1(A_z \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, \mathbf{Q}_\ell)$ is isomorphic to two copies of $H^1(E_z \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, \mathbf{Q}_\ell)$ on which multiplication by $1 + i$ acts as an automorphism, after extending the scalar field to $\mathbf{Q}_\ell(\sqrt{2})$, we see that the action of $\tilde{b}_2 = \frac{1}{\sqrt{2}}(1 + I)$ on $H^1(A_z \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, \mathbf{Q}_\ell(\sqrt{2}))$ is two copies of that on $H^1(E_z \otimes_{\mathbf{Q}} \bar{\mathbf{Q}}, \mathbf{Q}_\ell(\sqrt{2}))$, an automorphism of order 8.

2. $z = \frac{1+i}{\sqrt{3+1}}$. Consider the sublattice of finite index $L = (\mathbf{Z} + \mathbf{Z}x) + (\mathbf{Z} + \mathbf{Z}x)y \subset O_B$, where $x, y \in O_B$ are defined as follows: $x = I + J + IJ$ and $y = J + IJ$. One checks that

$$x^2 = -3, \quad y^2 = 6, \quad xy + yx = 0.$$

Note that $\mathbf{Z} + \mathbf{Z}x$ is isomorphic to $\mathbf{Z}[\sqrt{-3}]$ which is of index 2 in the ring of integers of $\mathbf{Q}[\zeta_3]$. At $z = \frac{1+i}{\sqrt{3+1}}$, one finds

$$\varphi(x) \begin{pmatrix} z \\ 1 \end{pmatrix} = \sqrt{-3} \begin{pmatrix} z \\ 1 \end{pmatrix}, \quad \text{so} \quad \varphi(\mathbf{Z} + \mathbf{Z}x) \begin{pmatrix} z \\ 1 \end{pmatrix} = \mathbf{Z}[\sqrt{-3}] \begin{pmatrix} z \\ 1 \end{pmatrix}.$$

Similarly one finds

$$\varphi((\mathbf{Z} + \mathbf{Z}x)y) \begin{pmatrix} z \\ 1 \end{pmatrix} = \mathbf{Z}[\sqrt{-3}]\varphi(y) \begin{pmatrix} z \\ 1 \end{pmatrix}.$$

The vectors $\begin{pmatrix} z \\ 1 \end{pmatrix}$ and $\varphi(y) \begin{pmatrix} z \\ 1 \end{pmatrix}$ form a \mathbf{C} -linear basis of \mathbf{C}^2 . Therefore the lattice $\varphi(L) \subset \mathbf{C}^2$ is isomorphic to $(\mathbf{Z} + \mathbf{Z}\sqrt{-3}) \oplus (\mathbf{Z} + \mathbf{Z}\sqrt{-3})$, making it clear that the abelian surface $B_z := \mathbf{C}^2/\varphi(L) \begin{pmatrix} z \\ 1 \end{pmatrix}$ is isomorphic to E_z^2 where $E_z = \mathbf{C}/\mathbf{Z}[\sqrt{-3}]$ is an elliptic curve over \mathbf{Q} with CM by $\mathbf{Z}[\sqrt{-3}]$.

In this case $b_3 = (3 + 3I + J + IJ)/2$ acts on A_z and B_z by multiplication by $\mu(b_3) = 1 + \frac{1+\sqrt{-3}}{2} = \sqrt{3}\zeta_{12}$. By the same argument as in the previous case, for $\ell \geq 5$, the action of \tilde{b}_3 on the Tate module of A_z over $\mathbf{Q}_\ell(\sqrt{3})$ is two copies of that on the Tate module of E_z over $\mathbf{Q}_\ell(\sqrt{3})$, which is an automorphism of order 12.

3. $z = \frac{\sqrt{6}i}{3+\sqrt{3}}$. Define a sublattice of finite index $L = (\mathbf{Z} + \mathbf{Z}b_6) + (\mathbf{Z} + \mathbf{Z}b_6)J \subset O_B$. Since $b_6^2 = -6$, $\mathbf{Z} + \mathbf{Z}b_6$ is isomorphic to the maximal order $\mathbf{Z}[\sqrt{-6}]$ in the field $\mathbf{Q}(\sqrt{-6})$. Note that $b_6J = -Jb_6$. At the point $z = \sqrt{-6}/(3 + \sqrt{3})$ one finds

$$\varphi(\mathbf{Z} + \mathbf{Z}b_6) \begin{pmatrix} z \\ 1 \end{pmatrix} = \mathbf{Z}[\sqrt{-6}]\varphi(b_6) \begin{pmatrix} z \\ 1 \end{pmatrix}.$$

Similarly one finds

$$\varphi((\mathbf{Z} + \mathbf{Z}b_6)J) \begin{pmatrix} z \\ 1 \end{pmatrix} = \mathbf{Z}[\sqrt{-6}]\varphi(Jb_6) \begin{pmatrix} z \\ 1 \end{pmatrix}.$$

The vectors $\varphi(b_6) \begin{pmatrix} z \\ 1 \end{pmatrix}$ and $\varphi(Jb_6) \begin{pmatrix} z \\ 1 \end{pmatrix}$ form a \mathbf{C} -linear basis of \mathbf{C}^2 relative to which the lattice L is isomorphic to $\mathbf{Z}[\sqrt{-6}] \oplus \mathbf{Z}[\sqrt{-6}]$. This gives the isomorphism $B_z = \mathbf{C}^2 / \varphi(L) \begin{pmatrix} z \\ 1 \end{pmatrix} \simeq E_z \times E_z$ where E_z is the elliptic curve $\mathbf{C}/\mathbf{Z}[\sqrt{-6}]$, which is defined over $\mathbf{Q}(\sqrt{2})$ with CM by $\mathbf{Z}[\sqrt{-6}]$.

In this case $b_6 = 3I + IJ$ acts on A_z and B_z by multiplication by $\mu(b_6) = \sqrt{-6} = \sqrt{6}\zeta_4$. Similar to the previous two cases, for $\ell \geq 7$, the action of \tilde{b}_6 on the Tate module of A_z over $\mathbf{Q}_\ell(\sqrt{6})$ is two copies of that on the Tate module of E_z over the same field, which is an automorphism of order 4.

For $B = M_2(\mathbf{Q})$, we choose the standard maximal order $O_B = M_2(\mathbf{Z})$ so that $O_B^1 = \mathrm{SL}_2(\mathbf{Z})$. Using the basis $e_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $e_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $e_3 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, and $e_4 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ of $M_2(\mathbf{Z})$, we see that, for each $z \in \mathfrak{H}$,

$$O_B \begin{pmatrix} z \\ 1 \end{pmatrix} = \mathbf{Z}e_1 \begin{pmatrix} z \\ 1 \end{pmatrix} + \mathbf{Z}e_2 \begin{pmatrix} z \\ 1 \end{pmatrix} + \mathbf{Z}e_3 \begin{pmatrix} z \\ 1 \end{pmatrix} + \mathbf{Z}e_4 \begin{pmatrix} z \\ 1 \end{pmatrix} = \mathbf{Z} \begin{pmatrix} z \\ 0 \end{pmatrix} + \mathbf{Z} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \mathbf{Z} \begin{pmatrix} 0 \\ z \end{pmatrix} + \mathbf{Z} \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

This shows that the abelian surface $A_z = \mathbf{C}^2 / O_B \begin{pmatrix} z \\ 1 \end{pmatrix}$ decomposes as E_z^2 with the elliptic curve $E_z = \mathbf{C}/(\mathbf{Z} \cdot z + \mathbf{Z} \cdot 1)$. Hence for elliptic modular groups we shall replace A_z by E_z and follow the classical moduli interpretation.

The preimages in $\mathrm{SL}_2(\mathbf{R})$ of the four elliptic modular groups in Theorem 1 are $\pm(2, \infty, \infty) = \Gamma_0(2)$, $\pm(2, 3, \infty) = \mathrm{SL}_2(\mathbf{Z})$, $\pm(2, 4, \infty) = \Gamma_0(2)^+$, and $\pm(2, 6, \infty) = \Gamma_0(3)^+$. The two elliptic modular groups in Theorem 2 are subgroups of $\mathrm{SL}_2(\mathbf{Z})$. For each of them it is straightforward to compute stabilizers at elliptic points z and determine the elliptic curves E_z explicitly. In the table below, for each $\Gamma \subset \mathrm{SL}_2(\mathbf{R})$ considered in this paper, we summarize the information on its elliptic points z , the group $\langle T_z \rangle$ of stabilizers in Γ of z and the order of T_z . At each elliptic point z , A_z is isogenous to $E_z \times E_z$ for an elliptic curve $E_z = \mathbf{C}/L_z$. For each group except $(2, 4, 6)$, the lattice $L_z = \mathbb{Z}z \oplus \mathbb{Z}$; for $(2, 4, 6)$, $L_z = \mathbb{Z}[\sqrt{-6}]$, $\mathbb{Z}[\sqrt{-1}]$, $\mathbb{Z}[\sqrt{-3}]$ at elliptic points of orders 2, 4, 6 respectively, as computed above. Let $K_z = L_z \otimes_{\mathbb{Z}} \mathbf{Q}$ be the CM field of E_z . We also record $\mu(z)$, the action of T_z , before normalization by a scalar, on E_z as complex multiplication, and κ_z , the field of definition of E_z .

Γ	$\Gamma_1(3)$	$\Gamma_0(2)$	$\mathrm{SL}_2(\mathbf{Z})$	$\Gamma_0(2)^+$	$\Gamma_0(3)^+$	$\pm(2, 4, 6)$
z	$\frac{3+\sqrt{-3}}{6}$	$\frac{1+i}{2}$	$i, \frac{-1+\sqrt{-3}}{2}$	$\frac{i}{\sqrt{2}}, \frac{1+i}{2}$	$\frac{i}{\sqrt{3}}, \frac{3+\sqrt{-3}}{6}$	$\frac{\sqrt{-6}}{3+\sqrt{3}}, i, \frac{1+i}{1+\sqrt{3}}$
T_z	$\begin{pmatrix} 1 & -1 \\ 3 & -2 \end{pmatrix}$	$\begin{pmatrix} 1 & -1 \\ 2 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 \\ -1 & 0 \\ 0 & -1 \\ 1 & 1 \end{pmatrix}$	$\frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ -2 & 0 \\ 1 & 1/2 \\ -1 & 0 \end{pmatrix}$	$\frac{1}{\sqrt{3}} \begin{pmatrix} 0 & 1 \\ -3 & 0 \\ 1 & 1/3 \\ -1 & 0 \end{pmatrix}$	s_2, s_4, s_6 <i>cf.</i> (29)
$\#\langle T_z \rangle$	3	4	4, 6	4, 8	4, 12	4, 8, 12
$\mu(z)$	ζ_3	i	i, ζ_6	$-\sqrt{-2}, -(1+i)$	$-\sqrt{-3}, -(1+\zeta_6)$	$\sqrt{-6}, 1+i, 1+\zeta_6$
κ_z	\mathbf{Q}	\mathbf{Q}	\mathbf{Q}, \mathbf{Q}	\mathbf{Q}, \mathbf{Q}	\mathbf{Q}, \mathbf{Q}	$\mathbf{Q}(\sqrt{2}), \mathbf{Q}, \mathbf{Q}$
$K_z \cdot \kappa_z$	$\mathbf{Q}(\sqrt{-3})$	$\mathbf{Q}(i)$	$\mathbf{Q}(i),$ $\mathbf{Q}(\sqrt{-3})$	$\mathbf{Q}(\sqrt{-2}),$ $\mathbf{Q}(i)$	$\mathbf{Q}(\sqrt{-3}),$ $\mathbf{Q}(\sqrt{-3})$	$\mathbf{Q}(\sqrt{2}, \sqrt{-6}),$ $\mathbf{Q}(i), \mathbf{Q}(\sqrt{-3})$

TABLE 8. CM points information

Observe that the CM field K_z has class number 1 except at the elliptic point z of order 2 for the group $(2, 4, 6)$, in which case it has class number 2.

The interested reader may consult [9] by Baba and Granath for explicit models of genus-two curves with QM at points not admitting CM by -3 or -4 , as well as the discussion of the fields of moduli. For example, at the elliptic point z of order 2 for $(2,4,6)$ admitting CM by $\mathbf{Q}(\sqrt{-24})$, they give a curve C with $\text{Jac}(C) \simeq E_{z,1} \times E_{z,2}$ defined over $\mathbf{Q}(\sqrt{2})$.

5.2. Contributions from elliptic points. Let Γ be one of the groups considered in Theorems 1 and 2 and let z be an elliptic point of Γ of order N_z . (The group (∞, ∞, ∞) in Theorem 2 is the only group containing no elliptic points.) Then z is a \mathbf{Q} -rational point on the curve X_Γ . Fix a prime ℓ . In this subsection we compute the traces of the Frobenius elements Frob_p for primes $p \geq 7$ on $V^k(\Gamma)_{\bar{z}, \ell}$, the stalk at z of the automorphic sheaf on X_Γ , for integers $k \geq 1$. Since the groups Γ in Theorem 1 are projective, only even k will be considered, so we shall not distinguish Γ and $\pm\Gamma$. In the discussion below, we assume Γ and z are as listed in Table 8 and will use notation from that table. Recall that the abelian surface A_z , defined over κ_z , is isogenous to $E_z \times E_z$ for an elliptic curve E_z defined over κ_z . Denote by $\sigma_{z, \ell}$ the Tate module on E_z over $\overline{\mathbf{Q}}_\ell$. It is a degree-2 representation of G_{κ_z} .

The elliptic curve E_z , defined over the real field κ_z , has CM by the imaginary quadratic field K_z . Thus there is an ℓ -adic character ξ_z of the Galois group $G_{K_z \kappa_z}$ such that $\sigma_{z, \ell}$ is the induced representation $\text{Ind}_{G_{K_z \kappa_z}}^{G_{\kappa_z}} \xi_z$. Let X be a vector spanning the line on which $G_{K_z \kappa_z}$ acts via ξ_z . The complex conjugation $c \in G_{\mathbf{Q}}$ lies in G_{κ_z} but not $G_{K_z \kappa_z}$. Let $Y = (\sigma_{z, \ell}(c))X$. Then X and Y form a basis of $\sigma_{z, \ell}$ since $G_{\kappa_z} = G_{K_z \kappa_z} \cup G_{K_z \kappa_z}c$ and $\sigma_{z, \ell}$ is 2-dimensional and irreducible. Therefore we can represent $\sigma_{z, \ell}(g)$ for $g \in G_{\kappa_z}$ by a 2×2 matrix in the basis X, Y as follows:

$$\sigma_{z, \ell}(g) = \begin{pmatrix} \xi_z(g) & \\ & \xi_z^c(g) \end{pmatrix} \quad \text{if } g \in G_{K_z \kappa_z},$$

and

$$\sigma_{z, \ell}(c) = \begin{pmatrix} & \xi_z(c^2) \\ 1 & \end{pmatrix} = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}$$

since $c^2 = id$. Here $\xi_z^c : g \mapsto \xi_z(c^{-1}gc)$ is the conjugate of ξ_z by c . In particular we get the decomposition $\sigma_{z, \ell}|_{G_{K_z \kappa_z}} = \xi_z \oplus \xi_z^c$. As $\sigma_{z, \ell}$ is a Tate module of E_z , we have $\det \sigma_{z, \ell} = \epsilon_\ell|_{G_{\kappa_z}}$. Therefore $\xi_z \cdot \xi_z^c = \epsilon_\ell|_{G_{K_z \kappa_z}}$.

Let \mathfrak{p} be a nonzero prime ideal of κ_z unramified in $K_z \kappa_z$ and let \mathfrak{P} be a prime ideal of $\kappa_z K_z$ above \mathfrak{p} . As a consequence of the above, we obtain the following action of $\text{Frob}_\mathfrak{p}$ with respect to the basis X, Y :

(5.2.1) If \mathfrak{p} splits in $K_z \kappa_z$, then

$$\sigma_{z, \ell}(\text{Frob}_\mathfrak{p})X = a_\mathfrak{p}X, \quad \sigma_{z, \ell}(\text{Frob}_\mathfrak{p})Y = d_\mathfrak{p}Y, \quad a_\mathfrak{p}d_\mathfrak{p} = N\mathfrak{p}.$$

In fact, $(a_\mathfrak{p}, d_\mathfrak{p}) = (\xi_z(\text{Frob}_\mathfrak{P}), \xi_z^c(\text{Frob}_\mathfrak{P}))$ or $(\xi_z^c(\text{Frob}_\mathfrak{P}), \xi_z(\text{Frob}_\mathfrak{P}))$. Note that $\xi_z^c(\text{Frob}_\mathfrak{P}) = \xi_z(\text{Frob}_\mathfrak{P}^c)$.

(5.2.2) If \mathfrak{p} is inert in $K_z \kappa_z$, then

$$\sigma_{z, \ell}(\text{Frob}_\mathfrak{p})X = c_\mathfrak{p}Y, \quad \sigma_{z, \ell}(\text{Frob}_\mathfrak{p})Y = b_\mathfrak{p}X, \quad c_\mathfrak{p}b_\mathfrak{p} = -N\mathfrak{p}.$$

As the trace and determinant of $\sigma_{z, \ell}(\text{Frob}_\mathfrak{p})$ are \mathbf{Z} -valued, upon fixing an isomorphism $\iota : \overline{\mathbf{Q}}_\ell \rightarrow \mathbf{C}$, we shall regard $\xi_z(\text{Frob}_\mathfrak{P})$ as an algebraic integer and $\xi_z^c(\text{Frob}_\mathfrak{P})$ its complex conjugation. When \mathfrak{p} is inert in $\kappa_z K_z$, we have $\xi_z(\text{Frob}_\mathfrak{P}) = \xi_z^c(\text{Frob}_\mathfrak{P}) = -N(\mathfrak{p})$. This identifies ξ_z with a Hecke Grossencharacter of $\kappa_z K_z$ which takes values in the ring of integers of the field $\kappa_z K_z$.

On the other hand, as explained in §5.1, the stabilizer T_z , before normalization by a scalar to make $\det T_z = 1$, acts on E_z via multiplication by $\mu(z) \in K_z$ given in Table 8. It follows from the CM theory that CM by $\mu(z)$ on E_z commutes with the action of the Galois group $G_{K_z \kappa_z}$ on E_z . This is carried over to the space $\langle X, Y \rangle$ of $\sigma_{z, \ell}$. Consequently, X and Y are eigenvectors of T_z . We know T_z has one eigenvalue $\mu(z)/|\mu(z)| =: \zeta_z$, which is a primitive M_z th root of unity. Here M_z is

the order of T_z , equal to N_z for $\Gamma = \Gamma_1(3) = (3, \infty, \infty)$, and $2N_z$ otherwise. Since $\det T_z = 1$, the other eigenvalue of T_z is ζ_z^{-1} . Without loss of generality, we may assume

$$T_z X = \zeta_z X \quad \text{and} \quad T_z Y = \zeta_z^{-1} Y.$$

As explained in §2.2.2, the sheaf $V^k(\Gamma)_{\bar{z}, \ell}$ is defined as the pushforward of a sheaf $V^k(\Gamma')_{\bar{z}, \ell}$, where Γ' is a torsion-free normal subgroup of Γ of finite index. We first consider the groups Γ in Theorem 1 for which $k \geq 2$ is even. It follows from Proposition 3 and the discussion in the previous subsection that $V^k(\Gamma')_{\bar{z}, \ell} = \text{Sym}^k(\eta_{z, \ell})$, where $\eta_{z, \ell}$ is the restriction of $\sigma_{z, \ell}$ to a suitable Galois subgroup of G_{κ_z} depending on the moduli interpretation of $X_{\Gamma'}$. As the stabilizers of z in Γ form the cyclic group $\langle T_z \rangle$ of finite order, the space of $V^k(\Gamma)_{\bar{z}, \ell}$ is $\text{Sym}^k(\sigma_{z, \ell})^{T_z}$. We shall describe the action of $G_{\mathbf{Q}}$ on this space by computing the trace of Frob_p for primes $p \geq 7$.

In view of the action of T_z described above, the space $\text{Sym}^k(\sigma_{z, \ell})^{T_z}$ has a basis

$$\begin{aligned} & \{X^a Y^b \mid a + b = k, a, b \geq 0 \text{ and } a \equiv b \pmod{M_z}\} \\ & = \{(XY)^{k/2} X^{rN_z} Y^{-rN_z} \mid r \in \mathbf{Z}, -\frac{k}{2N_z} \leq r \leq \frac{k}{2N_z}\}. \end{aligned}$$

In particular, when $k = 2$, the space $\text{Sym}^2(\sigma_{z, \ell})^{T_z}$ is 1-dimensional, spanned by XY .

Now we proceed to compute the trace of Frob_p on $\text{Sym}^k(\sigma_{z, \ell})^{T_z}$. We shall split the discussion into two cases.

Case (I). $\kappa_z = \mathbf{Q}$, which holds for all z except the order 2 elliptic point for $\Gamma = (2, 4, 6)$. It follows from (5.2.1) and (5.2.2) that $G_{\mathbf{Q}}$ acts on $\langle XY \rangle$ via the character $\chi_{d_z} \epsilon_\ell$, where d_z is the discriminant of K_z as given in Table 1. Furthermore, if p is inert in K_z , then, by (5.2.2), $\text{Sym}^k \sigma_{z, \ell}(\text{Frob}_p)$ swaps $(XY)^{k/2} X^{rN_z} Y^{-rN_z}$ and $(XY)^{k/2} Y^{rN_z} X^{-rN_z}$ so that $(XY)^{k/2}$ is the only invariant vector in the chosen basis. Consequently the trace of the Frob_p action on $\text{Sym}^k(\sigma_{z, \ell})^{T_z}$ is $(-p)^{k/2}$ for p inert in K_z . If p splits in K_z , then, by (5.2.1), each basis vector $(XY)^{k/2} X^{rN_z} Y^{-rN_z}$ is an eigenvector of $\text{Sym}^k \sigma_{z, \ell}(\text{Frob}_p)$ with eigenvalue

$$p^{k/2} (\xi_z(\text{Frob}_{\mathfrak{P}}) \xi_z^c(\text{Frob}_{\mathfrak{P}})^{-1})^{\pm rN_z} = p^{k/2} (\xi_z(\text{Frob}_{\mathfrak{P}})^2/p)^{\pm rN_z},$$

where \mathfrak{P} is a prime ideal of K_z dividing p and the sign in front of r is independent of r . The second expression above uses $\xi_z(\text{Frob}_{\mathfrak{P}}) \xi_z^c(\text{Frob}_{\mathfrak{P}}) = N(\mathfrak{P}) = p$. Summing over $-\frac{k}{2N_z} \leq r \leq \frac{k}{2N_z}$ gives the trace of Frob_p on $\text{Sym}^k(\sigma_{z, \ell})^{T_z}$ as

$$(48) \quad \sum_{-\frac{k}{2N_z} \leq r \leq \frac{k}{2N_z}} p^{k/2} (\xi_z(\text{Frob}_{\mathfrak{P}})^2/p)^{rN_z},$$

independent of the choice of \mathfrak{P} and the choice of ξ_z or ξ_z^c .

Note that the CM field K_z as class number 1. By the CM theory, when $(p) = \mathfrak{P}\mathfrak{P}^c$ splits in K_r , the value $\xi_z(\text{Frob}_{\mathfrak{P}})$ generates the ideal \mathfrak{P} . It differs from an arbitrary generator $\alpha_{z, p}$ of \mathfrak{P} by a unit of K_z , which has order dividing $2N_z$ for all z with $\kappa_z = \mathbf{Q}$ except for the elliptic point of order 2 for the group $(2, 6, \infty)$. So for the non-exceptional z we may replace $\xi_z(\text{Frob}_{\mathfrak{P}})$ in (48) by any generator $\alpha_{z, p}$ of a prime ideal of K_z above a splitting p , as described in Theorems 1 and 2. The exceptional z , namely the order 2 elliptic point of $(2, 6, \infty)$, has coordinate $\lambda(z) = 1$ as given in Theorem 1. It is the image of the unique point with coordinate $t = 1/2$ on the 2-fold cover $X_{(3, \infty, \infty)}$ of $X_{(2, 6, \infty)}$ under the projection $\lambda = -4(t - 1)t$. The elliptic curve at the point $t = 1/2$ on $X_{(3, \infty, \infty)}$ is $E_{(3, \infty, \infty), 1/2}$ defined by $x^3 = y^2 + xy + \frac{1/2}{27}y$ as given in Table 4. It is isogenous to E_z . So we have $\text{Tr} \sigma_{z, \ell}(\text{Frob}_p) = H_p(\{1/3, 2/3\}, \{1, 1\}; 1/2)$ by Proposition 4. Therefore at primes p splitting in $\mathbf{Q}(\sqrt{-3})$, $\alpha_{z, p} = \xi_z(\text{Frob}_{\mathfrak{P}})$ or $\xi_z^c(\text{Frob}_{\mathfrak{P}})$ is a root of

$$x^2 - H_p(\{1/3, 2/3\}, \{1, 1\}; 1/2)x + p = 0.$$

The hypergeometric datum $\{\{1/3, 2/3\}, \{1, 1\}\}$ is denoted by $HD_2(\Gamma)$ for $\Gamma = (2, 6, \infty)$ in §4.5. Corollary 3 in §4.6 states that $H_p(HD_2(\Gamma); 1/2)^2 = H_p(HD(\Gamma), 1) + 2p$ for p splitting in $\mathbf{Q}(\sqrt{-3})$. In other words, at primes p splitting in K_z , $\alpha_{z,p}^2 = \xi_z(\text{Frob}_{\mathfrak{P}})^2$ or $\xi_z^c(\text{Frob}_{\mathfrak{P}})^2$ is a root of

$$x^2 - H_p(HD(\Gamma); 1)x + p^2 = 0.$$

In fact, the above expression holds at the elliptic point z of order $N_z = 2$ for $\Gamma = (2, \infty, \infty), (2, 3, \infty), (2, 4, \infty)$, and $(2, 6, \infty)$ by the same corollary.

We have shown that the contributions of Frobenius traces at elliptic points z of Γ with $\kappa_z = \mathbf{Q}$ for even weight are as described in Theorems 1 and 2.

Case (II). $\kappa_z \neq \mathbf{Q}$, which occurs only at the order $N_z = 2$ elliptic point z for $(2, 4, 6)$. In this case $\kappa_z = \mathbf{Q}(\sqrt{2})$, $K_z = \mathbf{Q}(\sqrt{-6})$ and $\kappa_z K_z = \mathbf{Q}(\sqrt{2}, \sqrt{-6})$ as listed in Table 8. Let $\tau \in G_{\mathbf{Q}}$ be such that $\tau(\sqrt{2}) = -\sqrt{2}$ and $\tau(\sqrt{-6}) = \sqrt{-6}$. Note that $G_{\mathbf{Q}(\sqrt{-6})} = G_{\mathbf{Q}(\sqrt{2}, \sqrt{-6})} \cup G_{\mathbf{Q}(\sqrt{2}, \sqrt{-6})}\tau$ and $G_{\mathbf{Q}} = G_{\mathbf{Q}(\sqrt{-6})} \cup G_{\mathbf{Q}(\sqrt{-6})}c$. As τ gives rise to an isogeny from E_z to $\tau(E_z)$ over $\mathbf{Q}(\sqrt{2})$, the representation $\sigma_{z,\ell}$ is isomorphic to the conjugate $\sigma_{z,\ell}^\tau$. (The interested reader may refer to [10] by Baba and Granath for algebraic models for A_z , E_z and $\tau(E_z)$.) This implies that $\sigma_{z,\ell}$ can be extended to $G_{\mathbf{Q}}$. Also $\xi_z = \xi_z^\tau$ can be extended to $G_{\mathbf{Q}(\sqrt{-6})}$. There are two extensions of $\sigma_{z,\ell}$ to $G_{\mathbf{Q}}$; they differ by the quadratic character χ_2 , hence they induce the same action on the space of $V^k(\Gamma)_{\bar{z},\ell}$ for even $k \geq 2$.

Let σ'_z be an extension of $\sigma_{z,\ell}$ to $G_{\mathbf{Q}}$. By Clifford theory (cf. [44]), $\sigma'_z = \text{Ind}_{G_{\mathbf{Q}(\sqrt{-6})}}^{G_{\mathbf{Q}}} \psi_z$ for a character ψ_z of $G_{\mathbf{Q}(\sqrt{-6})}$ extending ξ_z . Since $\det \sigma'_z$ extends $\det \sigma_{z,\ell} = \epsilon_\ell|_{G_{\mathbf{Q}(\sqrt{2})}}$, we have $\det \sigma'_z = \epsilon_\ell$ or $\chi_2 \epsilon_\ell$. Its restriction to $G_{\mathbf{Q}(\sqrt{-6})}$ is $\psi_z \psi_z^c$.

To proceed, we need to determine the determinant of σ'_z , which is independent of the extension of σ_z we choose.

Lemma 9.

$$\det \sigma'_z = \epsilon_\ell \quad \text{and} \quad \psi_z \psi_z^c = \epsilon_\ell|_{G_{\mathbf{Q}(\sqrt{-6})}}.$$

Proof. Since χ_2 is nontrivial on $G_{\mathbf{Q}(\sqrt{-6})}$ and $\det \sigma'_z|_{G_{\mathbf{Q}(\sqrt{-6})}} = \psi_z \psi_z^c$, the two statements are equivalent. We shall prove $\det \sigma'_z = \epsilon_\ell$.

Let Γ' be a normal torsion-free subgroup of $\pm(2, 4, 6)$ of finite index so that $\text{Sym}^2(\Gamma)_\ell$ is the pushforward from $\text{Sym}^2(\Gamma')_\ell$. Then so are their respective determinant ρ and ρ' as rank-1 sheaves on the curves $X_{(2,4,6)}$ and $X_{\Gamma'}$ respectively. On the other hand, we know that $\rho' = \epsilon_\ell^3$ by Proposition 3, the same holds for the pushforward ρ at nonelliptic and hence all points on $X_{(2,4,6)}$ by the comparison theorem (Theorem 8). This shows that ρ restricted to the fiber at z , which is $\det \sigma'_z$, is ϵ_ℓ , as claimed. \square

A similar argument as before gives the following action of Frob_p for primes $p \geq 5$:

(5.2.3) If p splits in $K_z = \mathbf{Q}(\sqrt{-6})$, then

$$\sigma'_z(\text{Frob}_p)X = a_p X, \quad \sigma'_z(\text{Frob}_p)Y = d_p Y, \quad a_p d_p = p.$$

More precisely, let \mathfrak{P} be a prime ideal of K_z dividing p . Then $(a_p, d_p) = (\psi_z(\text{Frob}_{\mathfrak{P}}), \psi_z^c(\text{Frob}_{\mathfrak{P}}))$ or $(\psi_z^c(\text{Frob}_{\mathfrak{P}}), \psi_z(\text{Frob}_{\mathfrak{P}}))$. In both cases, we have $a_p d_p = \psi_z \psi_z^c(\text{Frob}_{\mathfrak{P}}) = \det \sigma'_z(\text{Frob}_p) = p$.

(5.2.4) If p is inert in K_z , then

$$\sigma'_z(\text{Frob}_p)X = c_p Y, \quad \sigma'_z(\text{Frob}_p)Y = b_p X, \quad c_p b_p = -p.$$

In all cases, we have

Corollary 4. *Let Γ be a group in Theorem 1 and z be an elliptic point of Γ . Then $G_{\mathbf{Q}}$ acts on the space $\text{Sym}^2(\sigma_{z,\ell})^{T_z} = \langle XY \rangle$ via the character $\chi_{d_z} \epsilon_\ell$, where d_z is the discriminant of the CM field K_z listed in Table 1.*

Similar computations as in Case (I) give similar results for Case (II). More precisely, at primes p inert in $K_z = \mathbf{Q}(\sqrt{-6})$, the trace of $\sigma'_z(\text{Frob}_p)$ on $\text{Sym}^k(\sigma_{z,\ell})^{T_z}$ is $(-p)^{k/2}$. When p splits in K_z , the trace of $\sigma'_z(\text{Frob}_p)$ on $\text{Sym}^k(\sigma_{z,\ell})^{T_z}$ is

$$(49) \quad \sum_{-\frac{k}{2N_z} \leq r \leq \frac{k}{2N_z}} p^{k/2} (\psi_z(\text{Frob}_{\mathfrak{P}})^2/p)^{rN_z},$$

independent of the choice of a degree-1 prime ideal \mathfrak{P} of K_z above p and the choice of ψ_z or ψ_z^c . In this case the field $K_z = \mathbf{Q}(\sqrt{-6})$ has class number 2 so that the square of every integral ideal of K_z is principal. Let \mathfrak{A} be a prime ideal of $\kappa_z K_z$ above \mathfrak{P} , and let \mathfrak{Q} be the prime ideal of κ_z divisible by \mathfrak{A} . If \mathfrak{P} splits in $\kappa_z K_z$, then $\psi_z(\text{Frob}_{\mathfrak{P}}) = \xi_z(\text{Frob}_{\mathfrak{A}}) = \xi_z^{\tau}(\text{Frob}_{\mathfrak{A}})$ so that $\psi_z(\text{Frob}_{\mathfrak{P}})^2 = \xi_z(\text{Frob}_{\mathfrak{A}})^2$. In this case, $\psi_z(\text{Frob}_{\mathfrak{P}})^2$, being in \mathfrak{A}^2 and invariant under τ , lies in \mathfrak{P}^2 . Furthermore, its norm is $\xi_z(\text{Frob}_{\mathfrak{A}})^2 \xi_z^c(\text{Frob}_{\mathfrak{A}})^2 = p^2$, which is the norm of \mathfrak{P}^2 . Therefore $\psi_z(\text{Frob}_{\mathfrak{P}})^2$ generates \mathfrak{P}^2 . Since the units of $\mathbf{Z}[\sqrt{-6}]$ are ± 1 and $N_z = 2$, we may replace $\psi_z(\text{Frob}_{\mathfrak{P}})^2$ by any generator $\alpha_{z,p}$ of \mathfrak{P}^2 . If \mathfrak{P} is inert in $\kappa_z K_z$, then p is inert in κ_z and $\mathfrak{Q} = (p)$ splits in $\kappa_z K_z$. So $\psi_z(\text{Frob}_{\mathfrak{P}})^2 = \xi_z(\text{Frob}_{\mathfrak{A}})$ has norm p^2 and it generates \mathfrak{P}^2 . By a similar argument, we may replace $\psi_z(\text{Frob}_{\mathfrak{P}})^2$ by any generator $\alpha_{z,p}$ of \mathfrak{P}^2 . On the other hand, as pointed out in Remark 8, $\rho_{HD(2,4,6),1,\ell}^{BCM}$ is the ℓ -adic representation of $G_{\mathbf{Q}}$ attached to the weight-3 modular form $f_{24,3,h,a}$, which has \mathbf{Z} -coefficients and CM by $\mathbf{Q}(\sqrt{-6}) = K_z$. Hence $\rho_{HD(2,4,6),1,\ell}^{BCM}$ is induced from an ℓ -adic character ν of G_{K_z} such that above primes p splitting in K_z , $\nu(\text{Frob}_{\mathfrak{P}})^2$ generates \mathfrak{P}^2 . Hence $\psi_z(\text{Frob}_{\mathfrak{P}})^2 = \pm \nu(\text{Frob}_{\mathfrak{P}})^2$. As $\text{Tr} \rho_{HD(2,4,6),1,\ell}^{BCM}(\text{Frob}_p) = p H_p(HD(2,4,6),1)$, we may choose $\alpha_{z,p}$ to be a root of

$$x^2 - \left(\frac{-3}{p}\right) p H_p(HD(2,4,6),1) x + p^2 = 0,$$

as asserted in Theorem 1.

Summarizing the above discussions, we have proved that the contributions from elliptic points of the groups in Theorem 1 for k even are as stated.

It remains to consider the unique elliptic point z of order $N_z = 3$ for $\Gamma = (3, \infty, \infty) = \Gamma_1(3)$ in Theorem 2. We have $\kappa_z = \mathbf{Q}$ and $K_z = \mathbf{Q}(\sqrt{-3})$. Since Γ does not contain $-I_2$, the sheaf $V^k(\Gamma)_{\ell}$ is defined as the pushforward of $V^k(\Gamma')_{\ell}$ and $V^k(\Gamma')_{\ell} = \text{Sym}^k(V^1(\Gamma')_{\ell})$. Therefore the space of $V^k(\Gamma)_{\bar{z},\ell}$ is $\text{Sym}^k(\sigma_{z,\ell})^{T_z}$, which has a basis (for $k \geq 1$)

$$\begin{aligned} & \{X^a Y^b \mid a+b=k, a,b \geq 0 \text{ and } a \equiv b \pmod{3}\} \\ & = \{X^r Y^{k-r} \mid r \in \mathbf{Z}, k \geq r \geq 0, \text{ and } 2r \equiv k \pmod{3}\} \end{aligned}$$

since T_z has order 3. Write $2r - k = 3m$. Then $m \equiv k \pmod{2}$, $-\frac{k}{3} \leq m \leq \frac{k}{3}$, $r = (k+3m)/2$ and $k-r = (k-3m)/2$. We rewrite the above basis as

$$\{X^{(k+3m)/2} Y^{(k-3m)/2} \mid m \equiv k \pmod{2}, -\frac{k}{3} \leq m \leq \frac{k}{3}\}.$$

At a prime p inert in K_z , $\sigma_z(\text{Frob}_p)$ swaps $X^{(k+3m)/2} Y^{(k-3m)/2}$ and $X^{(k-3m)/2} Y^{(k+3m)/2}$ up to scalar. This shows that the trace of $\sigma_z(\text{Frob}_p)$ on $\text{Sym}^k(\sigma_{z,\ell})^{T_z}$ is 0 for k odd and $(-p)^{k/2}$ for k even. For a prime p split in K_z , let \mathfrak{p} be a prime ideal of K_z above p . Because of the symmetry in m , we may assume $\sigma_z(\text{Frob}_{\mathfrak{p}})X = \xi_z(\text{Frob}_{\mathfrak{p}})X$ so that

$$\begin{aligned} \sigma_z(\text{Frob}_{\mathfrak{p}})X^{(k+3m)/2} Y^{(k-3m)/2} &= \xi_z(\text{Frob}_{\mathfrak{p}})^{(k+3m)/2} \xi_z^c(\text{Frob}_{\mathfrak{p}})^{(k-3m)/2} X^{(k+3m)/2} Y^{(k-3m)/2} \\ &= p^{(k-3m)/2} \xi_z(\text{Frob}_{\mathfrak{p}})^{3m} X^{(k+3m)/2} Y^{(k-3m)/2}. \end{aligned}$$

This shows that the trace of $\sigma_z(\text{Frob}_p)$ on $\text{Sym}^k(\sigma_{z,\ell})^{T_z}$ is

$$\sum_{m \equiv k \pmod{2}, -\frac{k}{3} \leq m \leq \frac{k}{3}} p^{(k-3m)/2} \xi_z(\text{Frob}_p)^{3m}.$$

By the CM theory, $\xi_z(\text{Frob}_p)$ is a generator of the ideal \mathfrak{p} , which is unique up to sixth roots of unity. When k is even, so is m and we can replace $\xi_z(\text{Frob}_p)$ in the above formula by any generator $\alpha_{z,p}$ of \mathfrak{p} . While for k odd, so is m , then $\alpha_{z,p}^{3m}$ agrees with $\xi_z(\text{Frob}_p)^{3m}$ up to sign. To determine $\xi_z(\text{Frob}_p)^3$, recall that every elliptic curve defined over \mathbf{Q} with a \mathbf{Q} -rational 3-torsion point and vanishing j -invariant is isomorphic to either $x^3 = y^2 + ay$ for some $a \in \mathbf{Q}$ or $x^3 = y^2 - (\frac{1}{2^4 \cdot 3 \cdot \sqrt{-3}})^2$ which is isomorphic to the curve above $\frac{1+\sqrt{-3}}{2}$ on $X_1(3)$. In either case, for each prime $p \equiv 1 \pmod{3}$ where the elliptic curve has a good reduction, $\xi_z(\text{Frob}_p)^3 = -J_\omega(1/3, 1/3)^3$.

Proposition 9. *The contributions of the Frobenius traces at the elliptic points for groups in Theorems 1 and 2 are as stated.*

5.3. Contributions from cusps. In this section we discuss the contribution from a cusp z of a group Γ in Theorems 1 and 2. Recall that $V^k(\Gamma)_\ell = \text{Sym}^k(V^1(\Gamma)_\ell)$. The inertia subgroup at z , denoted I_z , is cyclic generated by an element with Jordan form $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ or $\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}$. As explained in §2.2.1, the $G_{\mathbf{Q}}$ action on the stalk $V^1(\Gamma)_{\bar{z},\ell}$ arises from the action on the I_z -invariant space of a degree-2 representation ρ of the étale fundamental group of the curve X_Γ° at the generic point. Let $\{X, Y\}$ be a basis of the space of ρ . Then the space of $V^k(\Gamma)_{\bar{z},\ell}$ is 1-dimensional, spanned by Y^k , if 1 is an eigenvalue of I_z ; it is $\langle Y^k \rangle$ for k even and 0 for k odd if -1 is an eigenvalue of I_z .

The contribution at z is equal to 1 for k even, following the same argument and computation as in Scholl [56]. Hence we assume k odd and the group $\Gamma = \Gamma_1(3)$ or $\Gamma_1(4)$.

In Table 4 an explicit model for the “universal elliptic curve” $E_{\Gamma,t}$ with t running through the open curve X_Γ° for $\Gamma \in \{\Gamma_1(3), \Gamma_1(4)\}$ is given. The two cusps of $\Gamma_1(3)$ and the three cusps of $\Gamma_1(4)$ all have coordinates $t(z) \in \{0, 1, \infty\}$. The contribution from z is determined by the reduction type of the curve $E_{\Gamma,t(z)}$ at its unique singular point. More precisely, the trace of Frob_p on the 1-dimensional space $V^1(\Gamma)_{\bar{z},\ell}$ is 1, -1 , or 0 according as $E_{\Gamma,t(z)} \pmod{p}$ being split multiplicative reduction, nonsplit multiplicative reduction, or additive reduction. In the case of $\Gamma_1(3)$, when $t = 0$, the curve $E_{\Gamma_1(3),0}$, defined by $y^2 + xy = x^3$, has two distinct tangents over \mathbf{Q} at the singular point $(0,0)$. At $t = 1$, the singular point is $(-1/9, 1/27)$, which can be shifted to the origin $(0,0)$ in the equation $y^2 + xy = x^3 - x^2/3$. Hence, the curve has multiplicative reduction at primes $p \neq 3$, which splits if and only if -3 is a square mod p . This proves case (2) in Theorem 2 for $\Gamma = \Gamma_1(3)$.

We make similar computations for $E_{\Gamma_1(4),t}$. At $t = 0$, the curve admits a splitting multiplicative reduction at the singular point $(0,0)$ for all odd primes p . At $t = 1$, the singular point is $(-1/8, 1/32)$ where the curve has multiplicative reduction at odd primes, which splits if and only if -1 is a square mod p . At $t = \infty$, we set $t = a/b$ and make the change of variable $(x, y) \mapsto (x/b^3, y/b^2)$. By letting $b = 0$, the curve become $y^2 = x^3$, whose singularity $(0,0)$ is a cusp. Hence the contribution at this irregular cusp is zero. This proves case (1) of Theorem 2.

6. VERIFICATION FOR THE LOWEST WEIGHT CASES

Let $\Gamma = (e_1, e_2, e_3)$ be one of the groups in Theorem 1. We now verify the trace formula for the $k = 2$ case. Recall from Theorem 13 that $\chi_\Gamma \otimes V^2(\Gamma)_\ell = \varphi_\Gamma \otimes \mathcal{H}(\{1/2\}, \{1\})_\ell \otimes \mathcal{H}(HD(\Gamma))_\ell$ with $\chi_\Gamma = \chi_{-1}$ for $\Gamma = (2, \infty, \infty), (2, 3, \infty)$ and $(2, 6, \infty)$, $\chi_\Gamma = \chi_{-3}$ for $\Gamma = (2, 4, 6)$, and χ_Γ trivial for $\Gamma = (2, 4, \infty)$.

Firstly, we recall the dimension formula for $S_{2+k}(\Gamma)$, the space of holomorphic cusp forms for Γ of weight $2 + k$:

$$\dim S_{2+k}(\Gamma) = -1 - k + \sum_{j=1}^3 \left\lfloor \frac{k-2}{2} \left(1 - \frac{1}{e_j}\right) \right\rfloor, \quad \text{for } k \geq 2 \text{ even,}$$

(cf Theorem 4 of [68]). In particular, $S_4(\Gamma)$ is 0-dimensional for each Γ listed in the Theorem 1. Thus the left hand side of the Hecke trace formula (1) is 0 for each unramified prime p .

When $k = 2$, at each unramified prime p , the contributions from the elliptic points and cusps have been computed in §5. Namely, from each cusp the contribution is 1, and the contribution from an elliptic point z is $\chi_{d_z}(p)p$ as in Corollary 4. By Proposition 6 and Theorem 13 the contribution from a generic fiber is $\chi_\Gamma(p)\varphi_\Gamma(p)\phi_p(1 - 1/\lambda)p^u H_p(HD(\Gamma), 1/\lambda)$ where $u = 1$ when $\Gamma = (2, 4, 6)$ and $u = 0$ otherwise; also as in Proposition 6, $\varphi_\Gamma = \chi_{-1}$ for $(2, d, \infty)$ when $d = 2, 6$ or infinity and otherwise it is the trivial character. Putting together, when $k = 2$ the right hand side of (1) is of the form

$$\sum_{\lambda=2}^{p-1} \chi_\Gamma(p)\varphi_\Gamma(p)\phi_p(1 - 1/\lambda)p^u H_p(HD(\Gamma), 1/\lambda) + \text{the contributions from singular fibers.}$$

When $\Gamma = (2, 4, 6)$, by Table 1, at elliptic points of order 2, 4, 6, their contributions $\chi_{d_z}(p)p$ are $\chi_{-6}(p)p$, $\chi_{-1}(p)p$, $\chi_{-3}(p)p$ respectively. In this case $u = 1$. Thus for each unramified prime p

$$\sum_{\lambda=2}^{p-1} \chi_{-3}(p)\phi_p(1 - 1/\lambda)p H_p(HD(\Gamma), 1/\lambda) + \chi_{-6}(p)p + \chi_{-1}(p)p + \chi_{-3}(p)p = 0.$$

When $\Gamma = (2, 4, \infty)$, $u = 0$, and by Table 1, at elliptic points of order 2 and 4, their contributions are $\chi_{-2}(p)p$ and $\chi_{-1}(p)p$ respectively, and at the cusp ∞ , the contribution is 1. Putting together

$$\sum_{\lambda=2}^{p-1} \phi_p(1 - 1/\lambda)H_p(HD(\Gamma), 1/\lambda) + 1 + \chi_{-2}(p)p + \chi_{-1}(p)p = 0.$$

The same analysis applies to other cases.

Next we consider the $k = 1$ case for Theorem 2. Let $e_\infty \in \{\infty, 3\}$ and $\Gamma = (\infty, \infty, e_\infty)$. Note that $\dim S_3(\Gamma) = 0$ (see [20] for example). It follows from Proposition 4 that χ_Γ is trivial and

$$a_\Gamma(\lambda, p) = \begin{cases} H_p(\{\frac{1}{2}, \frac{1}{2}\}, \{1, 1\}; 1/\lambda) & \text{if } e_\infty = \infty \\ H_p(\{\frac{1}{3}, \frac{2}{3}\}, \{1, 1\}; 1/\lambda) & \text{if } e_\infty = 3. \end{cases}$$

When $e_\infty = \infty$, the group $\Gamma = \Gamma_1(4)$ has 3 cusps; as explained in §5.3, they contribute $1, \left(\frac{-1}{p}\right)^k$, and $\frac{1+(-1)^k}{2}$ respectively to the trace formula. Combined with Lemma 7, the trace formula gives value 0 for $k = 1$, consistent with $\dim S_3(\Gamma) = 0$.

The $e_\infty = 3$ case can be obtained similarly. In this case there are two regular cusps, whose contributions to the trace formula are 1 and $\left(\frac{-3}{p}\right)^k$ for each integer $k \geq 1$.

7. FURTHER CONSEQUENCES

7.1. Other subgroups. The key step in the proof of Theorems 1 and 2 is to explicitly relate the ℓ -adic sheaf $V^2(\Gamma)_\ell$ or $V^1(\Gamma)_\ell$ to the hypergeometric ℓ -adic sheaf attached to the datum $HD(\Gamma)$ as introduced by Katz [36] (See Theorem 7). This method applies to arithmetic triangle groups Γ satisfying the following two conditions: (i) X_Γ is a genus zero curve defined over \mathbf{Q} , (ii) Γ has three elliptic points and cusps altogether, and these points are all \mathbf{Q} -rational. None of the three index-2

subgroups $(2, 6, 6), (3, 4, 4), (2, 2, 2, 3)$ of $(2, 4, 6)$ listed in the diagram 1b satisfy these conditions. Since the automorphic sheaves on the subgroups are pullbacks of those on $(2, 4, 6)$, in the theorem below we demonstrate that, by using the pullback of the hypergeometric sheaf $\mathcal{H}(HD(2, 4, 6))_\ell$ along the explicit covering maps $\pi_i, i = 3, 2, 6$, over \mathbf{Q} given by (28) in §3.3, analogous Hecke trace formulae for these three subgroups can be obtained. This approach also applies to other groups in a similar situation, as long as explicit covering maps are known.

Theorem 14. *For the three index-2 subgroups $\Gamma = (2, 6, 6), (3, 4, 4), (2, 2, 2, 3)$ of $(2, 4, 6)$, choose the generator $t = t(\Gamma)$ of $\mathbf{Q}(X_\Gamma)$ with the following values at the elliptic points of given order:*

Γ	$(2, 6, 6)$	$(3, 4, 4)$	$(2, 2, 2, 3)$
t	$(0, 1/\sqrt{-3}, -1/\sqrt{-3})$	$(\infty, \sqrt{-1}, -\sqrt{-1})$	$(\sqrt{-3}, -\sqrt{-3}, \infty, 0)$
CM field discriminants	$(-4, -3, -3)$	$(-3, -4, -4)$	$(-24, -24, -4, -3)$
λ	$1 + 1/(3t^2)$	$1/(1 + t^2)$	$-t^2/3$

so that the generator $\lambda(2, 4, 6)$ of $\mathbf{Q}(X_{(2,4,6)})$ in Theorem 1 is a rational function of t given in the last row. Given any even integer $k \geq 2$ and a fixed prime ℓ , for almost all primes $p \neq \ell$ where X_Γ has good reduction, the explicit expression for the terms in the right-hand side of (1) as described in Theorem 1 holds with $a_\Gamma(t, p) = \text{Tr}(\text{Frob}_t | (V^k(\Gamma)_\ell)_{\bar{t}})$ in (3) at $t \in X_\Gamma(\mathbf{F}_p)$ not corresponding to an elliptic point given as follows:

For $\Gamma = (2, 6, 6)$,

$$a_\Gamma(t, p) = \begin{cases} \left(\frac{-3(1+3t^2)}{p}\right) pH_p(HD(2, 4, 6), 1/(1 + \frac{1}{3t^2})), & \text{if } t \neq \infty; \\ pH_p(HD(2, 4, 6), 1) + \left(\frac{-6}{p}\right) p, & \text{if } t = \infty \end{cases}$$

For $\Gamma = (3, 4, 4)$,

$$a_\Gamma(t, p) = \begin{cases} \left(\frac{-3(1+t^2)}{p}\right) pH_p(HD(2, 4, 6), 1 + t^2), & \text{if } t^2 \neq -1, 0; \\ \left(\frac{-3}{p}\right) pH_p(HD(2, 4, 6), 1) + \left(\frac{-6}{p}\right) p, & \text{if } t = 0; \end{cases}$$

For $\Gamma = (2, 2, 2, 3)$,

$$a_\Gamma(t, p) = \left(\frac{-3 - t^2}{p}\right) pH_p(HD(2, 4, 6), -3/t^2), \text{ if } t^2 \neq -3.$$

When $t \in \mathbf{F}_p$ corresponds to an elliptic point, the contribution from t is the same as (5).

Proof. The proof follows from noting the sheaf $V^k(\Gamma)_\ell$ on X_Γ is the pullback of the sheaf $V^k(2, 4, 6)_\ell$ on $X_{(2,4,6)}$ along the two-fold \mathbf{Q} -rational covering maps $X_\Gamma \rightarrow X_{(2,4,6)}$ arising from the canonical models of these Shimura curves as described in §3.3, as well as the known Frobenius traces given in Theorem 1. Details are omitted. \square

As an application, we have the following

Theorem 15. *For any given prime ℓ ,*

$$\rho_{\{\{\frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4}\}, \{1, 1, \frac{1}{6}, \frac{5}{6}\}\}; 1, \ell}^{BCM} \cong (\chi_{-1} \otimes \rho_{f_{6.4.a.a}, \ell}) \oplus \chi_{-2\ell}$$

as $G_{\mathbf{Q}}$ -modules, where $\rho_{f_{6.4.a.a}, \ell}$ is the ℓ -adic representation of $G_{\mathbf{Q}}$ associated with $f_{6.4.a.a}$. It is equivalent to (6) which says

$$pH_p\left(\left\{\frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4}\right\}, \left\{1, 1, \frac{1}{6}, \frac{5}{6}\right\}; 1\right) = \left(\frac{-1}{p}\right) a_p(f_{6.4.a.a}) + \left(\frac{-2}{p}\right) p.$$

Proof of (6). For any prime $p > 3$, by the relation between Gauss sum and Jacobi sum and $J(\phi\chi, \bar{\chi}) = \phi\chi(-1)J(\phi\chi, \phi)$,

$$\begin{aligned}
& pH_p \left(\left\{ \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \right\}, \left\{ 1, 1, \frac{1}{6}, \frac{5}{6} \right\}; 1 \right) \\
&= \frac{p}{1-p} \sum_{\chi} \frac{\mathfrak{g}(\phi\chi)\mathfrak{g}(\bar{\chi})}{\mathfrak{g}(\phi)} \cdot \frac{\mathfrak{g}(\phi\chi)\mathfrak{g}(\bar{\chi})}{\mathfrak{g}(\phi)} \frac{\mathfrak{g}(\chi^4)\mathfrak{g}(\bar{\chi})\mathfrak{g}(\bar{\chi}^6)}{\mathfrak{g}(\chi^2)\mathfrak{g}(\bar{\chi}^2)\mathfrak{g}(\bar{\chi}^3)} \chi(27/4) \\
&= \frac{p}{1-p} \sum_{\chi} \phi\chi(-1)J(\phi\chi, \phi) \cdot \frac{\mathfrak{g}(\phi\chi)\mathfrak{g}(\bar{\chi})}{\mathfrak{g}(\phi)} \frac{\mathfrak{g}(\chi^4)\mathfrak{g}(\bar{\chi})\mathfrak{g}(\bar{\chi}^6)}{\mathfrak{g}(\chi^2)\mathfrak{g}(\bar{\chi}^2)\mathfrak{g}(\bar{\chi}^3)} \chi(27/4) \\
&= - \sum_{t \in \mathbf{F}_p} \phi(-t(1-t)) pH_p \left(\left\{ \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \right\}, \left\{ 1, \frac{1}{6}, \frac{5}{6} \right\}; t \right) \\
&\stackrel{\text{Lem. 8}}{=} -\phi(3) \sum_{t \in \mathbf{F}_p} (\phi(-3t) + 1)\phi(1-t) pH_p \left(\left\{ \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \right\}, \left\{ 1, \frac{1}{6}, \frac{5}{6} \right\}; t \right) \\
&\quad - \left(\frac{-2}{p} \right) p - \left(\frac{-3}{p} \right) p - \left(\frac{-1}{p} \right) p \\
&\stackrel{-3t=s^2}{=} - \sum_{s \in \mathbf{F}_p} \phi(3(1+s^2/3)) pH_p \left(\left\{ \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \right\}, \left\{ 1, \frac{1}{6}, \frac{5}{6} \right\}; -s^2/3 \right) - \left(\frac{-2}{p} \right) p - \left(\frac{-3}{p} \right) p - \left(\frac{-1}{p} \right) p \\
&\stackrel{\text{Thm. 14}}{=} \phi(-1) \text{Tr}(T_p | S_4(2, 2, 2, 3)) + \left(\frac{-2}{p} \right) p.
\end{aligned}$$

□

Remark 9. If a different \mathbf{Q} -model on $X_{(2,6,6)}$ is chosen such that all three elliptic points are \mathbf{Q} -rational, then it has an associated datum $HD(2, 6, 6) = \{ \{ \frac{1}{2}, \frac{1}{3}, \frac{2}{3} \}, \{ 1, \frac{1}{6}, \frac{5}{6} \} \}$. The two \mathbf{Q} -models on $X_{(2,6,6)}$ are isomorphic over $\mathbf{Q}(\sqrt{-3})$ and not \mathbf{Q} . Consequently, the formula obtained using $H_p(HD(2, 6, 6); \lambda)$ agrees with that from Theorem 14 only for $p \equiv 1 \pmod{6}$. Similarly, $X_{(3,4,4)}$ has a different \mathbf{Q} -model to which one can associate a hypergeometric datum $HD(3, 4, 4) = \{ \{ \frac{1}{12}, \frac{1}{3} \}, \{ 1, \frac{3}{4} \} \}$, which is no longer defined over \mathbf{Q} . The expression using $H_p(HD(3, 4, 4); \lambda)$ agrees with that from Theorem 14 only for $p \equiv 1 \pmod{12}$. The curve $X_{(2,2,2,3)}$ does not have an associated hypergeometric datum because it has 4 elliptic points.

Remark 10. Theorem 14 combined with (1) gives an explicit description of the trace of the Hecke operator T_p on $S_{k+2}(\Gamma)$ for $\Gamma = (2, 6, 6), (3, 4, 4), (2, 2, 2, 3)$, which consists of forms in $S_{k+2}(2, 2, 3, 3)$ invariant under the involution w_i for $i = 3, 2, 6$ respectively. The intersection of any two out of these three spaces is the space of forms in $S_{k+2}(2, 2, 3, 3)$ invariant under all three Atkin-Lehner involutions w_2, w_3, w_6 , hence it is $S_{k+2}(2, 4, 6)$. Therefore combining Theorems 1 and 14, we also obtain an explicit expression of

$$\begin{aligned}
\text{Tr}(T_p | S_{k+2}(2, 2, 3, 3)) &= \text{Tr}(T_p | S_{k+2}(2, 6, 6)) + \text{Tr}(T_p | S_{k+2}(3, 4, 4)) \\
&\quad + \text{Tr}(T_p | S_{k+2}(2, 2, 2, 3)) - 2\text{Tr}(T_p | S_{k+2}(2, 4, 6))
\end{aligned}$$

in terms of hypergeometric sums attached to $HD(2, 4, 6)$. Our formulas are motivated in part by the results in [45] by the last three authors. For example it was shown there that for each prime $p > 5$,

$$-\text{Tr}(T_p | S_6(2, 2, 3, 3)) = p^2 H_p \left(\left\{ \frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \frac{2}{3} \right\}, \left\{ 1, \frac{1}{6}, \frac{5}{6}, 1, \frac{5}{6}, \frac{1}{6} \right\}; 1 \right) + pa_p(f_{18.4.a.a}) + \left(\frac{-1}{p} \right) p^2.$$

Under the Jacquet-Langlands correspondence [68], the space $S_{k+2}((2, 2, 3, 3))$ corresponds to the space of newforms in $\Gamma_0(6)$ of the same weight. Consequently, we obtain the following formulae for Hecke traces:

$$\begin{aligned}\mathrm{Tr}(T_p | S_{k+2}((2, 4, 6))) &= \mathrm{Tr}(T_p | S_{k+2}^{\mathrm{new}}(\Gamma_0(6), -, -)), \\ \mathrm{Tr}(T_p | S_{k+2}((2, 6, 6))) &= \mathrm{Tr}(T_p | S_{k+2}^{\mathrm{new}}(\Gamma_0(6), \pm, -)), \\ \mathrm{Tr}(T_p | S_{k+2}((3, 4, 4))) &= \mathrm{Tr}(T_p | S_{k+2}^{\mathrm{new}}(\Gamma_0(6), -, \pm)), \\ \mathrm{Tr}(T_p | S_{k+2}((2, 2, 2, 3))) &= \mathrm{Tr}(T_p | S_{k+2}^{\mathrm{new}}(\Gamma_0(6), \mathrm{sgn}, \mathrm{sgn})), \\ \mathrm{Tr}(T_p | S_{k+2}((2, 2, 3, 3))) &= \mathrm{Tr}(T_p | S_{k+2}^{\mathrm{new}}(\Gamma_0(6))).\end{aligned}$$

Here $S_{k+2}^{\mathrm{new}}(\Gamma_0(6), \mathrm{sgn}_1, \mathrm{sgn}_2)$ is the Atkin-Lehner subspace of the newforms on $\Gamma_0(6)$ with eigenvalues $\mathrm{sgn}_1 \cdot 1$ and $\mathrm{sgn}_2 \cdot 1$ for ω_2 and ω_3 , respectively.

7.2. Eigenvalues of Hecke operators. Let Γ be one of the groups discussed in this paper. Let p be a good prime for X_Γ . Suppose $a_r, 1 \leq r \leq m$, are the eigenvalues of T_p on $S_{k+2}(\Gamma)$. By Deligne [19] for Γ elliptic modular and Ohta [51] for Γ quaternionic, for each a_r there are two eigenvalues b_r and c_r of Frob_p on $H^1(X_\Gamma \otimes \bar{\mathbf{Q}}, V^k(\Gamma)_\ell)$ such that $a_r = b_r + c_r$. Furthermore, $b_r c_r = p^{k+1}$ since forms in $S_{k+2}(\Gamma)$ have trivial central character. Therefore to determine the m eigenvalues a_r of T_p , it suffices to compute the traces of $(\mathrm{Frob}_p)^r$ for $1 \leq r \leq m$ on the same cohomology space, which in turn gives rise to the values of $\sum_{1 \leq r \leq m} a_r^i$ for $i = 1, \dots, m$ from which we can solve for a_r . Since the action of Frob_p is unramified, the action of $(\mathrm{Frob}_p)^r$ agrees with that of an unramified $\mathrm{Frob}_{\mathfrak{P}_r}$ for a degree- r place \mathfrak{P}_r above p (of a number field). Consequently the trace of $(\mathrm{Frob}_p)^r$ for $1 \leq r \leq m$ can be computed from the local Frobenius traces at the points of reduction of X_Γ at \mathfrak{P}_r , that is,

$$\sum_{\lambda \in X_\Gamma(\mathbf{F}_{p^r})} \mathrm{Tr}(\mathrm{Frob}_\lambda | (V^k(\Gamma)_\ell)_{\bar{\lambda}}),$$

the same way as the case $r = 1$. In this way we express the eigenvalues of T_p on $S_{k+2}(\Gamma)$ in terms of hypergeometric character sums.

To illustrate our point, we exhibit two examples, the first one for the elliptic modular group $\Gamma_0(4)$ and the second for $(2,4,6)$ from the quaternion algebra B_6 .

Example 1. The space $S_8(\Gamma_0(4))$ has dimension two and all are old forms. Fix an odd prime p and let $a_{1,p}$ and $a_{2,p}$ be the eigenvalues of T_p on $S_8(\Gamma_0(4))$. By the trace formulae, we have the system of equations

$$\begin{aligned}-(a_{1,p} + a_{2,p}) &= \sum_{\lambda \in X_{\Gamma_0(4)}(\mathbf{F}_p)} \mathrm{Tr}(\mathrm{Frob}_\lambda | (V^6(\Gamma_0(4))_\ell)_{\bar{\lambda}}) \\ &= 3 + \sum_{\lambda=2}^{p-1} \sum_{j=0}^3 (-1)^j \binom{6-j}{j} p^j \cdot H_p \left(\left\{ \frac{1}{2}, \frac{1}{2} \right\}, \{1, 1\}; 1/\lambda \right)^{6-2j} \\ -(a_{1,p}^2 + a_{2,p}^2) &= -4p^7 + \sum_{\lambda \in X_{\Gamma_0(4)}(\mathbf{F}_{p^2})} \mathrm{Tr}(\mathrm{Frob}_\lambda | (V^6(\Gamma_0(4))_\ell)_{\bar{\lambda}}) \\ &= -4p^7 + 3 + \sum_{\lambda \in \mathbf{F}_{p^2} \setminus \{0,1\}} \sum_{j=0}^3 (-1)^j \binom{6-j}{j} p^j \cdot H_{p^2} \left(\left\{ \frac{1}{2}, \frac{1}{2} \right\}, \{1, 1\}; 1/\lambda \right)^{6-2j}.\end{aligned}$$

The computations give

p	5	7	11	13	17
$a_{1,p} + a_{2,p}$	-420	2032	2184	2764	29412
$a_{1,p}^2 + a_{2,p}^2$	88200	2064512	2384928	3819848	432532872

from which we get $a_{1,p} = a_{2,p} = \frac{1}{2}(a_{1,p} + a_{2,p})$, as it should be.

Example 2. The space $S_{24}(2, 4, 6)$ is 2-dimensional. Fix an odd prime $p > 3$ and let $a_{1,p}$ and $a_{2,p}$ be the eigenvalues of T_p on this space. By the trace formulae, we have the system of equations

$$\begin{aligned} -(a_{1,p} + a_{2,p}) &= \sum_{\lambda \in X_{(2,4,6)}(\mathbf{F}_p)} \text{Tr}(\text{Frob}_\lambda | (V^{22}(2, 4, 6)_\ell)_{\bar{\lambda}}) = CE(p) + \sum_{\lambda \in \mathbf{F}_p, \lambda \neq 0, -3} F_{11}(a_{(2,4,6)}(\lambda, p), p), \\ -(a_{1,p}^2 + a_{2,p}^2) &= -4p^{23} + CE(p^2) + \sum_{\lambda \in \mathbf{F}_{p^2}, \lambda \neq 0, -3} F_{11}(a_{(2,4,6)}(\lambda, p^2), p^2), \end{aligned}$$

where $CE(q)$ is the total contribution from the elliptic points in $X_{(2,4,6)}(\mathbf{F}_q)$. Its value for $q = p$ is as stated in Theorem 1, and for $q = p^2$ it is

$$CE(p^2) = \sum_{N=2,4,6} \sum_{-\frac{22}{2N} \leq i \leq \frac{22}{2N}} p^{22} (\alpha_{N,p^2}^2 / p^2)^{iN},$$

where $\alpha_{4,p^2} = J_\omega(\frac{1}{4}, \frac{1}{4})^2$, $\alpha_{6,p^2} = J_\omega(\frac{1}{3}, \frac{1}{3})^2$ for a generator ω of $\widehat{\mathbf{F}_{p^2}^\times}$, and α_{2,p^2}^2 is any root of $T^2 - p^2 H_{p^2}(HD(2, 4, 6), 1)T + p^4 = 0$. The computations give

p	5	7	11
$a_{1,p} + a_{2,p}$	25248156	5764462768	1017121470024
$a_{1,p}^2 + a_{2,p}^2$	70010194261011336	60171677733273590912	3068149691314205892000288

from which we get the eigenvalues $12624078 \pm 5184\beta$, $2882231384 \pm 129600\beta$, and $508560735012 \pm 31363200\beta$ with $\beta = \sqrt{1296640489}$ for $p = 5, 7, 11$, respectively. The two Hecke eigenforms in $S_{24}(2, 4, 6)$ correspond to the newforms in the Newform orbit 6.24.a.d in LMFDB.

APPENDIX: G -SHEAVES

Here we recall some of the theory of the functors p_*^G . If X is a space on which a group G acts and $p : X \rightarrow Y = G \backslash X$, then Grothendieck introduced functors

$$p_*^G : \{G\text{-sheaves on } X\} \rightarrow \{\text{sheaves on } Y\}$$

and studied their derived functors in section V of his Tohoku paper, [29]. He did this for ordinary topological spaces, but the definitions make sense in any generalized topology, such as the étale topology. By a G -sheaf on a G -space X we mean a sheaf F on X for which there are isomorphisms $\phi_g : g_* F \rightarrow F$, for all $g \in G$, which satisfy $\phi_{gh} = \phi_g \circ g_*(\phi_h)$ as maps $(gh)_* F = g_* h_* F \rightarrow g_* F \rightarrow F$. Then for any open U in Y , $p_*^G(F)(U) = F(p^{-1}U)^G$, a meaningful formula since $p^{-1}U$ is a G -invariant open subset of X . These functors are especially useful under a discontinuity hypothesis (condition (D) of Tohoku V).

Example: suppose that F is defined by a constant sheaf $X \times M$ for a G -module M . Then $p_*^G(F)$ is the sheaf defined by $G \backslash (X \times M)$ on $G \backslash X = Y$. For instance Bayer-Neukirch [11] let $G = \Gamma$ be a Fuchsian subgroup of $\text{SL}_2(\mathbf{R})$, $X = \mathfrak{H}^*$ the extended upper half-plane, and $M = \text{Sym}^k(\mathbf{C}^2)$ where Γ acts on \mathbf{C}^2 via its embedding in $\text{SL}_2(\mathbf{R})$. In this way they define sheaves $V^k(\Gamma) = p_*^\Gamma \text{Sym}^k(\mathbf{C}^2)$, at least when Γ is torsion-free. Note that if $-1 \in \Gamma$, these sheaves are identically 0 if k is odd. In

fact, it is really the quotient $\Gamma/\pm 1$ that acts. When k is even, $\text{Sym}^k(\mathbf{C}^2)$ is a $\Gamma/\pm 1$ -module, and condition (D) holds.

In general, if condition (D) holds, then the stalk $p_*^G(F)_y = F_x^{G_x}$ where $p(x) = y$, and $G_x \subset G$ is the isotropy subgroup of x , a finite group. Moreover, Grothendieck proved that under condition (D), if G acts fixed-point free, the functors p_*^G and p^{-1} induce an equivalence of categories

$$\{G - \text{sheaves on } X\} \cong \{\text{sheaves on } Y\}.$$

In particular $p^{-1}p_*^G(F) = F$.

In Tohoku, derived functors $R^i p_*^G$ and $R^i \Gamma_X^G$ are introduced. In particular he defines abelian groups $H^n(X; G, A)$ and sheaves of abelian groups $\mathbf{H}^n(G, A)$ on Y for any abelian G -sheaf A on X . Because of the two factorizations $\Gamma_X^G = \Gamma^G \Gamma_X = \Gamma_Y p_*^G$ there are two spectral sequences of composite functors converging to $H^n(X; G, A)$. The derived functors of Γ^G give group cohomology.

The definitions of these functors make sense in any topos. If E is a topos and G is a group in E , then there is a topos $B_{E,G}$ of G -objects in E . See SGA 4 IV, 2.3 and 2.4. In our application, we consider finite group G acting on a scheme X with quotient $p : X \rightarrow Y = G \backslash X$. The category of G -sheaves on $X_{\text{ét}}$ is a topos, by Giraud's criterion. From this one obtains functors p_*^G and factorizations $\Gamma_X^G = \Gamma^G \Gamma_X = \Gamma_Y p_*^G$ as above, so that the spectral sequences in Tohoku chapter V carry over to the étale topology.

Our application will be limited to a finite flat G -covering $p : X \rightarrow Y$ of nonsingular algebraic curves for a finite group G . We consider finite group G acting on the scheme X with quotient $Y = G \backslash X$ such that $p : X \rightarrow Y$ is faithfully flat and quasicompact. We assume that locally $X = \text{Spec}(A)$, for a Noetherian ring A , the action is given by an action G on a ring A , that $Y = \text{Spec}(B)$, $B = A^G$. This is faithfully flat and of finite type, which implies that A is a projective B -module of finite rank, hence locally free. We are going to work in the étale topology, and sheaf will generally mean constructible sheaf.

If in addition $p : X \rightarrow Y$ is étale, the functors p_*^G and $p^* = p^{-1}$ induce an equivalence of categories of constructible sheaves, and in particular $p^* p_*^G(F) = F$. Indeed, in this case, to give a G -sheaf F on X is the same as to give a descent datum on F . For more details, see [38]. Observe that, even though we still have descent for faithfully flat maps, to give a G -structure on F is not equivalent to give a descent datum.

First consider the case $X = \text{Spec}(L) \rightarrow Y = \text{Spec}(K)$ for a Galois extension of fields $K \subset L$. The étale site on $\text{Spec}(L)$ is equivalent to the category of continuous G_L -modules. Under this identity, the functors

$$p_*, p^* : \{\text{Sheaves on } X_{\text{ét}}\} \leftrightarrow \{\text{Sheaves on } Y_{\text{ét}}\}$$

become

$$\text{Ind, Res} : \{G_L - \text{modules}\} \leftrightarrow \{G_K - \text{modules}\}.$$

Note that the category of G_L -modules M with a $G = G(L/K)$ -action is equivalent to the category of G_K -modules, so to give a G_L -module M with a $G = G(L/K)$ -action is equivalent to give a G_K -module structure on M . If the sheaf F corresponds to the G_L -module M with a $G = G(L/K)$ -action, then $p_*^G(F)$ corresponds to the G_K -module

$$M = \text{Ind}_{G_K}^{G_L}(M)^{G(L/K)}.$$

Recall also that in étale topology one considers stalks in geometric points: spectra of algebraically closed fields. In this example, $M = F_{\bar{y}}$, where \bar{y} is the geometric point $\text{Spec}(\bar{L})$ of Y , so the above formula reads

$$p_*^G(F)_{\bar{y}} = F_{\bar{x}}.$$

A priori, $F_{\bar{x}}$ is only an G_L -module, and this isomorphism is as $G_L \subset G_K$ -modules. The existence of a G -structure on this sheaf means that in fact this G_L -module structure extends to a G_K -module structure.

We can consider more generally a G -covering $X \rightarrow Y = \text{Spec}(K)$, where $X = \text{Spec}(A)$ and A is an étale algebra over K , that is $A = \prod_{i=1}^g L_i$ where L_i are separable extensions of K . Then G acts transitively on the fields L_i . In this case, we have that

$$(p_*^G(F))_{\bar{y}} = F_{\bar{x}_i}.$$

Now let $x \in X$ be a closed point of the smooth curve X . The local ring $\mathcal{O}_{X,x}$ is a discrete valuation ring with residue field $\kappa(x)$. We consider the completion or henselization of this ring $\mathcal{O}_{X,x}^h$. More generally let $X = \text{Spec}(A)$, with a henselian dvr A . We let x, ξ be the closed and generic point of X , and $\bar{x}, \bar{\xi}$ be geometric points lying over them, corresponding to algebraic closures of $\kappa(x)$ and $\kappa(\xi)$ respectively.

The category of sheaves for the étale topology on X is equivalent to the category of triples $(M_{\bar{x}}, M_{\bar{\xi}}, \alpha)$, where $M_{\bar{x}}$ is a $G_{\kappa(x)}$ -module, $M_{\bar{\xi}}$ is a $G_{\kappa(\xi)}$ -module, and $\alpha : M_{\bar{x}} \rightarrow M_{\bar{\xi}}^{I_x}$ is a $G_{\kappa(x)}$ -morphism, where $I_x \subset G_{\kappa(\xi)}$ is the inertia subgroup, so $G_{\kappa(\xi)}/I_x \cong G_{\kappa(x)}$ ($M_{\bar{\xi}}^{I_x}$ is automatically a $G_{\kappa(\xi)}/I_x$ -module). Note that to say $\alpha : M_{\bar{x}} \rightarrow M_{\bar{\xi}}^{I_x}$ is equivalent to say that there is a morphism $\alpha' : M_{\bar{x}} \rightarrow M_{\bar{\xi}}$ which is compatible with the $G_{\kappa(\xi)}$ -module structures on source and target, since I_x acts trivially on the source.

Let $p : X = \text{Spec}(A) \rightarrow Y = \text{Spec}(B)$ where $B \subset A$ be a finite extension of henselian dvr's, with fraction fields $\kappa(\xi)$ and $\kappa(\eta)$ respectively. We assume that $\kappa(\xi)/\kappa(\eta)$ is a finite Galois extension with group $G = \text{Gal}(\kappa(\xi)/\kappa(\eta)) = G_{\kappa(\eta)}/G_{\kappa(\xi)}$. We also assume that A is the integral closure of B in $\kappa(\xi)$. It follows that $B = A^G$. Note that the inertia groups satisfy $I_x = I_y \cap G_{\kappa(\xi)}$.

We wish to calculate the functor p_*^G as a morphism $X \rightarrow Y$ in the étale topology, using the above description. A G -structure on a sheaf $F = (\alpha : M_{\bar{x}} \rightarrow M_{\bar{\xi}}^{I_x})$ on X is equivalent to giving a $G_{\kappa(\eta)}$ -structure on the diagram $\alpha' : M_{\bar{x}} \rightarrow M_{\bar{\xi}}$ of $G_{\kappa(\xi)}$ -modules. There are two stalks to compute: one at y and one at η . Since the map p is étale above η , the previous sections show that $p_*^G(F)_{\bar{\eta}} = M_{\bar{\xi}}$, now as a $G_{\kappa(\eta)}$ -module. For the stalk above y , a basic result in étale cohomology shows that

$$A_{\bar{y}} = H^0(\text{Spec}(\mathcal{O}_{Y,y}^{sh}), \tilde{A}), \quad \text{for any sheaf } A, \quad \tilde{A} = \text{Spec}(\mathcal{O}_{Y,y}^{sh}) \times_Y A,$$

where the superscript sh denotes strict henselization. We may therefore replace the rings A and B by their strict henselizations. Then $G_{\kappa(\xi)} = I_x$ and $G_{\kappa(\eta)} = I_y$. It is clear now that

$$p_*^G(F)_{\bar{y}} = M_{\bar{x}}^G = M_{\bar{x}}^{I_y/I_x},$$

for the strictly hensel case. The same formula is valid in the original (non strictly) hensel case, but now the finite group $\bar{G}_x := I_y/I_x$ gives the geometric ramification of the map and is a subquotient of G .

Summarizing: $p_*^G(F) = (\beta : N_{\bar{y}} \rightarrow N_{\bar{\eta}}^{I_y})$, where

$$N_{\bar{y}} = M_{\bar{x}}^{\bar{G}_x}, \quad N_{\bar{\eta}} = M_{\bar{\xi}}, \quad \beta = (\alpha')^{I_y}, \quad \bar{G}_x = I_y/I_x.$$

In particular $p_*^G(F)_{\bar{y}} = F_{\bar{x}}^{\bar{G}_x}$.

We can generalize this to a G -covering where A is now a product of the finite number of henselian discrete valuation rings A_i , corresponding to the discrete valuations of $\kappa(\xi)$ lying over the valuation ring $B \subset \kappa(\eta)$. Then $p_*^G(F)_{\bar{y}} = F_{\bar{x}_i}^{\bar{G}_{x_i}}$ where x_i is the closed point of A_i . Note that the A_i are permuted transitively by G .

Let $p : X \rightarrow Y$ be a finite flat G -covering of nonsingular algebraic curves defined over a field. The constructible sheaves F of interest to us will be such that on the complement $U \subset X$ of a finite number of points, $F|_U$ is lisse. This means that $F|_U$ is equivalent to a representation

$$\rho_F : \pi_1(U, \bar{\xi}) \rightarrow \text{Aut}(F_{\bar{\xi}}),$$

where $\bar{\xi}$ is an algebraic closure of the function field $\kappa(\xi)$ of the curve. Recall that there is a canonical epimorphism $G_{\kappa(\xi)} \rightarrow \pi_1(U, \bar{\xi})$, so $F_{\bar{\xi}}$ is a $G_{\kappa(\xi)}$ -module.

If $x \in X$ is a closed point, we obtain a morphism $f : S := \text{Spec}(\mathcal{O}_{X,x}^h) \rightarrow X$ and we may consider the constructible sheaf f^*F on S . This is described as in the previous paragraphs. If s, σ denote the special and generic point of S , then $F = (\alpha, M_{\bar{s}}, M_{\bar{\sigma}})$, where $M_{\bar{s}} = F_{\bar{x}}$, since $\kappa(s) = \kappa(x)$. To describe $M_{\bar{\sigma}}$ we must choose an embedding $\kappa(\sigma) \subset \kappa(\bar{\xi})$. Assume $\sigma \in U$, where F is lisse. Then we have an isomorphism $F_{\bar{\xi}} \cong F_{\bar{\sigma}} = M_{\bar{\sigma}}$ of $G_{\kappa(\sigma)}$ -modules. If F is a G -sheaf on X , then for the stalks we have $p_*^G(F)_{\bar{y}} = F_{\bar{x}}^{\bar{G}_x}$, where $x \in X$ is any point such that $p(x) = y$, and \bar{G}_x is the geometric inertia of the point x .

REFERENCES

- [1] *Revêtements étales et groupe fondamental (SGA 1)*, volume 3 of *Documents Mathématiques (Paris) [Mathematical Documents (Paris)]*. Société Mathématique de France, Paris, 2003. Séminaire de géométrie algébrique du Bois Marie 1960–61. [Algebraic Geometry Seminar of Bois Marie 1960–61], Directed by A. Grothendieck, With two papers by M. Raynaud, Updated and annotated reprint of the 1971 original [Lecture Notes in Math., 224, Springer, Berlin; MR0354651 (50 #7129)].
- [2] Scott Ahlgren. Gaussian hypergeometric series and combinatorial congruences. In *Symbolic computation, number theory, special functions, physics and combinatorics (Gainesville, FL, 1999)*, volume 4 of *Dev. Math.*, pages 1–12. Kluwer Acad. Publ., Dordrecht, 2001.
- [3] Scott Ahlgren and Ken Ono. A Gaussian hypergeometric series evaluation and Apéry number congruences. *J. Reine Angew. Math.*, 518:187–212, 2000.
- [4] Michael Allen, Brian Grove, Ling Long, and Fang-Ting Tu. The explicit hypergeometric-modularity method, arXiv:2404.00711 (2024).
- [5] George E. Andrews, Richard Askey, and Ranjan Roy. *Special functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1999.
- [6] A. O. L. Atkin and J. Lehner. Hecke operators on $\Gamma_0(m)$. *Math. Ann.*, 185:134–160, 1970.
- [7] A. O. L. Atkin and Wen Ch'ing Winnie Li. Twists of newforms and pseudo-eigenvalues of W -operators. *Invent. Math.*, 48(3):221–243, 1978.
- [8] A. O. L. Atkin, Wen-Ching Winnie Li, Tong Liu, and Ling Long. Galois representations with quaternion multiplication associated to noncongruence modular forms. *Trans. Amer. Math. Soc.*, 365(12):6217–6242, 2013.
- [9] Srinath Baba and Håkan Granath. Genus 2 curves with quaternionic multiplication. *Canad. J. Math.*, 60(4):734–757, 2008.
- [10] Srinath Baba and Håkan Granath. Quaternionic modular forms and exceptional sets of hypergeometric functions. *International Journal of Number Theory*, 11(02):631–643, 2015.
- [11] Pilar Báyer and Jürgen Neukirch. On automorphic forms and Hodge theory. *Math. Ann.*, 257(2):137–155, 1981.
- [12] Pilar Bayer and Artur Travesa. Uniformizing functions for certain Shimura curves, in the case $D = 6$. *Acta Arith.*, 126(4):315–339, 2007.
- [13] F. Beukers and G. Heckman. Monodromy for the hypergeometric function ${}_nF_{n-1}$. *Invent. Math.*, 95(2):325–354, 1989.
- [14] Frits Beukers, Henri Cohen, and Anton Mellit. Finite hypergeometric functions. *Pure Appl. Math. Q.*, 11(4):559–589, 2015.
- [15] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud. *Néron models*, volume 21 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1990.
- [16] Irene I. Bouw and Stefan Wewers. Computing L -functions and semistable reduction of superelliptic curves. *Glasg. Math. J.*, 59(1):77–108, 2017.
- [17] Pete L. Clark and John Voight. Algebraic curves uniformized by congruence subgroups of triangle groups. *Trans. Amer. Math. Soc.*, 371(1):33–82, 2019.
- [18] Alyson Deines, Jenny G. Fuselier, Ling Long, Holly Swisher, and Fang-Ting Tu. Generalized Legendre curves and quaternionic multiplication. *J. Number Theory*, 161:175–203, 2016.
- [19] Pierre Deligne. Formes modulaires et représentations l -adiques. In *Séminaire Bourbaki. Vol. 1968/69: Exposés 347–363*, volume 175 of *Lecture Notes in Math.*, pages Exp. No. 355, 139–172. Springer, Berlin, 1971.
- [20] Fred Diamond and Jerry Shurman. *A first course in modular forms*, volume 228 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005.
- [21] Noam D. Elkies. Shimura curve computations. In *Algorithmic number theory (Portland, OR, 1998)*, volume 1423 of *Lecture Notes in Comput. Sci.*, pages 1–47. Springer, Berlin, 1998.

- [22] Ron Evans and John Greene. Clausen’s theorem and hypergeometric functions over finite fields. *Finite Fields Appl.*, 15(1):97–109, 2009.
- [23] Sharon Frechette, Ken Ono, and Matthew Papanikolas. Combinatorics of traces of Hecke operators. *Proc. Natl. Acad. Sci. USA*, 101(49):17016–17020, 2004.
- [24] Sharon Frechette, Ken Ono, and Matthew Papanikolas. Gaussian hypergeometric functions and traces of Hecke operators. *Int. Math. Res. Not.*, (60):3233–3262, 2004.
- [25] Jenny Fuselier, Ling Long, Ravi Ramakrishna, Holly Swisher, and Fang-Ting Tu. Hypergeometric functions over finite fields. *Mem. Amer. Math. Soc.*, 280(1382), 2022.
- [26] Jenny G. Fuselier. Hypergeometric functions over \mathbb{F}_p and relations to elliptic curves and modular forms. *Proc. Amer. Math. Soc.*, 138(1):109–123, 2010.
- [27] Matthew Greenberg and John Voight. Computing systems of Hecke eigenvalues associated to Hilbert modular forms. *Math. Comp.*, 80(274):1071–1092, 2011.
- [28] John Greene. Hypergeometric functions over finite fields. *Trans. Amer. Math. Soc.*, 301(1):77–101, 1987.
- [29] Alexander Grothendieck. Sur quelques points d’algèbre homologique. *Tohoku Math. J. (2)*, 9:119–221, 1957.
- [30] Ki-ichiro Hashimoto. Explicit form of quaternion modular embeddings. *Osaka J. Math.*, 32(3):533–546, 1995.
- [31] Haruzo Hida. On abelian varieties with complex multiplication as factors of the Jacobians of Shimura curves. *Amer. J. Math.*, 103(4):727–776, 1981.
- [32] Jerome W. Hoffman and Fang-Ting Tu. Transformations of Hypergeometric Motives. *arXiv e-prints: 2003.05031*, 2020.
- [33] Yasutaka Ihara. Hecke polynomials as congruence ζ functions in elliptic modular case. *Ann. of Math. (2)*, 85:267–295, 1967.
- [34] Hervé Jacquet and Robert P. Langlands. *Automorphic forms on $GL(2)$* . Lecture Notes in Mathematics, Vol. 114. Springer-Verlag, Berlin-New York, 1970.
- [35] Bruce W. Jordan. On the Diophantine arithmetic of Shimura curves, thesis, Harvard University, 1981.
- [36] Nicholas M. Katz. *Exponential Sums and Differential Equations*, volume 124 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1990.
- [37] Nicholas M. Katz. Another look at the Dwork family. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II*, volume 270 of *Progr. Math.*, pages 89–126. Birkhäuser Boston, Boston, MA, 2009.
- [38] Max-Albert Knus and Manuel Ojanguren. *Théorie de la descente et algèbres d’Azumaya*. Lecture Notes in Mathematics, Vol. 389. Springer-Verlag, Berlin-New York, 1974.
- [39] Masao Koike. Hypergeometric series and elliptic curves over finite fields. *Sūrikaiseikikenkyūsho Kōkyūroku*, (843):27–35, 1993. Research into automorphic forms and L functions (Japanese) (Kyoto, 1992).
- [40] Michio Kuga and Goro Shimura. On the zeta function of a fibre variety whose fibres are abelian varieties. *Ann. of Math. (2)*, 82:478–539, 1965.
- [41] Akira Kurihara. On some examples of equations defining Shimura curves and the Mumford uniformization. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 25(3):277–300, 1979.
- [42] Catherine Lennon. Gaussian hypergeometric evaluations of traces of Frobenius for elliptic curves. *Proc. Amer. Math. Soc.*, 139(6):1931–1938, 2011.
- [43] Catherine Lennon. Trace formulas for Hecke operators, Gaussian hypergeometric functions, and the modularity of a threefold. *J. Number Theory*, 131(12):2320–2351, 2011.
- [44] Wen-Ching Winnie Li, Tong Liu, and Ling Long. Potentially GL_2 -type Galois representations associated to noncongruence modular forms. *Trans. Amer. Math. Soc.*, 371(8):5341–5377, 2019.
- [45] Wen-Ching Winnie Li, Ling Long, and Fang-Ting Tu. A Whipple ${}_7F_6$ formula revisited. *Matematica*, 1(2):480–530, 2022.
- [46] Qing Liu. Modèles minimaux des courbes de genre deux. *J. Reine Angew. Math.*, 453:137–164, 1994.
- [47] Ling Long. Some numeric hypergeometric supercongruences. In *Vertex operator algebras, number theory and related topics*, volume 753 of *Contemp. Math.*, pages 139–156. Amer. Math. Soc., Providence, RI, 2020.
- [48] Dermot McCarthy. Transformations of well-poised hypergeometric functions over finite fields. *Finite Fields Appl.*, 18(6):1133–1147, 2012.
- [49] A. P. Ogg. Real points on Shimura curves. In *Arithmetic and geometry, Vol. I*, volume 35 of *Progr. Math.*, pages 277–307. Birkhäuser Boston, Boston, MA, 1983.
- [50] Masami Ohta. On l -adic representations attached to automorphic forms. *Japan. J. Math. (N.S.)*, 8(1):1–47, 1982.
- [51] Masami Ohta. On the zeta function of an abelian scheme over the Shimura curve. *Japan. J. Math. (N.S.)*, 9(1):1–25, 1983.
- [52] Masami Ohta. On the zeta function of an abelian scheme over the Shimura curve. II. In *Galois groups and their representations (Nagoya, 1981)*, volume 2 of *Adv. Stud. Pure Math.*, pages 37–54. North-Holland, Amsterdam, 1983.
- [53] Ken Ono. Values of Gaussian hypergeometric series. *Trans. Amer. Math. Soc.*, 350(3):1205–1223, 1998.

- [54] Ken Ono and Hasan Saad. Some Eichler-Selberg Trace Formulas. *Hardy-Ramanujan Journal*, Volume 45 - 2022, February 2023.
- [55] Hans Petersson. Über die eindeutige Bestimmung und die Erweiterungsfähigkeit von gewissen Grenzkreisgruppen. *Abh. Math. Sem. Univ. Hamburg*, 12(1):180–199, 1937.
- [56] Anthony J. Scholl. The l -adic representations attached to a certain noncongruence subgroup. *J. Reine Angew. Math.*, 392:1–15, 1988.
- [57] René Schoof. Counting points on elliptic curves over finite fields. volume 7, pages 219–254. 1995. Les Dix-huitièmes Journées Arithmétiques (Bordeaux, 1993).
- [58] Jean-Pierre Serre. *Abelian l -adic representations and elliptic curves*, volume 7 of *Research Notes in Mathematics*. A K Peters, Ltd., Wellesley, MA, 1998. With the collaboration of Willem Kuyk and John Labute, Revised reprint of the 1968 original.
- [59] Jean-Pierre Serre and John Tate. Good reduction of abelian varieties. *Ann. of Math. (2)*, 88:492–517, 1968.
- [60] Goro Shimura. An ℓ -adic method in the theory of automorphic forms. *Collected Papers*, II:237–272, 1968.
- [61] Goro Shimura. *Introduction to the arithmetic theory of automorphic functions*, volume 11 of *Publications of the Mathematical Society of Japan*. Princeton University Press, Princeton, NJ, 1994. Reprint of the 1971 original, Kanô Memorial Lectures, 1.
- [62] Jan Stienstra and Frits Beukers. On the Picard-Fuchs equation and the formal Brauer group of certain elliptic $K3$ -surfaces. *Math. Ann.*, 271(2):269–304, 1985.
- [63] Kisao Takeuchi. Arithmetic triangle groups. *J. Math. Soc. Japan*, 29(1):91–106, 1977.
- [64] Kisao Takeuchi. Commensurability classes of arithmetic triangle groups. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 24(1):201–212, 1977.
- [65] John Voight. *Quaternion algebras*, volume 288 of *Graduate Texts in Mathematics*. Springer, Cham, [2021] ©2021.
- [66] Mark Watkins. Hypergeometric motives over \mathbb{Q} and their L -functions. <http://magma.maths.usyd.edu.au/~watkins/papers/known.pdf>, 2017. preprint.
- [67] André Weil. Jacobi sums as “Größencharaktere”. *Trans. Amer. Math. Soc.*, 73:487–495, 1952.
- [68] Yifan Yang. Schwarzian differential equations and Hecke eigenforms on Shimura curves. *Compos. Math.*, 149(1):1–31, 2013.
- [69] Don Zagier. *Modular forms of one variable*. Instituto de Matemática Pura e Aplicada, 2000.

Department of Mathematics, Louisiana State University, Baton Rouge, LA 70803, USA
hoffman@math.lsu.edu

Department of Mathematics, Pennsylvania State University, University Park, PA 16802, USA
wli@math.psu.edu, <http://www.math.psu.edu/wli/>

Department of Mathematics, Louisiana State University, Baton Rouge, LA 70803, USA
llong@lsu.edu, <https://www.math.lsu.edu/~llong/>

Department of Mathematics, Louisiana State University, Baton Rouge, LA 70803, USA
ftu@lsu.edu, <https://sites.google.com/view/ft-tu/>