Existence and uniqueness of quadratic and linear mean-variance equilibria in general semimartingale markets

Christoph Czichowsky^{*}

Martin Herdegen[†]

 gen^{\dagger} David Martins [‡]

August 5, 2024

Abstract

We revisit the classical topic of quadratic and linear mean-variance equilibria with both financial and real assets. The novelty of our results is that they are the first allowing for equilibrium prices driven by general semimartingales and hold in discrete as well as continuous time. For agents with quadratic utility functions, we provide necessary and sufficient conditions for the existence and uniqueness of equilibria. We complement our analysis by providing explicit examples showing non-uniqueness or non-existence of equilibria. We then study the more difficult case of linear mean-variance preferences. We first show that under mild assumptions, a linear mean-variance equilibrium corresponds to a quadratic equilibrium (for different preference parameters). We then use this link to study a fixed-point problem that establishes existence (and uniqueness in a suitable class) of linear mean-variance equilibria. Our results rely on fine properties of dynamic mean-variance hedging in general semimartingale markets.

1 Introduction

The capital asset pricing model (CAPM) of Treynor [32], Sharpe [29], Lintner [19, 20] and Mossin [24] is one of the first general equilibrium models for financial markets. Despite its limitations, it is still one of the cornerstones of modern financial theory and widely used in investment practice; see [18] for a recent overview.

Notwithstanding the enormous influence of the CAPM, a rigorous study of existence and uniqueness of CAPM equilibria was initiated only around 1990 by Nielsen [25, 26, 27] and Allingham [1], with more recent important contributions by Berk [2], Dana [11], Hens et al. [15], Wenzelburger [33] and Koch-Medina/Wenzelburger [17]. With the notable exception of [2], these works focus on preferences described not by expected utility but rather by mean-variance functionals, i.e., functionals of the form $U(\mu, \sigma)$, where U is quasiconcave, increasing in the mean μ and decreasing in the volatility σ . This is because without distributional assumptions on the returns, the only utility functions that are compatible with the CAPM (more precisely, the two-fund separation theorem) are quadratic utility functions; see the discussion in Berk [2, after Corollary 3.2]. For expected quadratic utility, existence and uniqueness of CAPM

^{*}London School of Economics and Political Science, Department of Mathematics, Columbia House, Houghton Street, London WC2A 2AE, UK, email: c.czichowsky@lse.ac.uk. Parts of this research were completed while this author was Visiting Professor at ETH Zürich. He thanks Martin Schweizer and the university for their hospitality.

[†]University of Warwick, Department of Statistics, Coventry, CV4 7AL, UK, email: m.herdegen@warwick.ac.uk.

[‡]ETH Zürich, Department of Mathematics, Rämistrasse 101, 8092 Zürich, Switzerland, email: davidptmartins@gmail.com

equilibria in one period (under suitable assumptions) seem to have been regarded as folklore knowledge from early on.

While generalisations of the CAPM to multi-period and continuous time models have been considered from the late 1970s onwards (see e.g. Stapleton/Subrahmanyam [30] and Breeden [3]), a rigorous study of the existence and uniqueness of CAPM equilibria in multi-period and continuous time models is still missing in the literature — all the papers cited in the previous paragraph study one-period models. The aim of this paper is to close this gap in the literature and to address the important question of existence and uniqueness of CAPM equilibria in multi-period and continuous time models. By exploiting the connection of mean–variance portfolio selection with the classical problem of mean–variance hedging in mathematical finance, see Schweizer [28] for an overview, we provide existence and uniqueness results for CAPM equilibria. Our results apply in both discrete and continuous time and are the first that provide the existence of equilibrium markets for information flows driven by general semimartingales. We also provide examples that illustrate when CAPM equilibria are not unique or fail to exist. Because "mean–variance efficient portfolios approximately maximise expected utility for a wide range of risk-averse (concave) utility functions" (see Markowitz [21, 22]), our results may also provide approximations to equilibrium prices for other utility functions.

In most of the extant literature on the existence and uniqueness of CAPM equilibria, it is assumed that the aggregate random endowment is tradable so that the financial market is complete. The completeness assumption implies the existence of a so-called representative agent, which simplifies the task of showing existence and uniqueness of an equilibrium: Every agent chooses to hedge their idiosyncratic risk and own a fraction of the market portfolio. This structure breaks down in the incomplete case as shown by Koch-Medina/Wenzelburger [17]. They find that in an incomplete market, each agent still hedges their individual endowment as best as possible, even though this cannot be done perfectly. But unlike in the complete case, asset prices are now determined by the so-called extended market portfolio, i.e., the aggregate endowment of all agents, given by the terminal value of the market portfolio together with the unhedgeable parts of the endowments.

Our work extends the work of [17] to multi-period and continuous time. We consider general semimartingale markets and assume that the agents receive endowments at the terminal time T that are partly unhedgeable.

In the case of of quadratic utility, our proof of existence and uniqueness of an equilibrium is based on the construction of a nonstandard type of representative agent, i.e., a fictional agent that aggregates the preferences and endowments of the other agents. We show that the market clears if and only if the representative agent does not trade, and this observation yields a pricing measure for the equilibrium market. To the best of our knowledge, this is the first result that allows for equilibrium prices driven by general semimartingales in an incomplete setting.

In the case of linear mean-variance preferences, we combine the above result with a fixedpoint approach, showing first that under mild assumptions on the model primitives, a linear mean-variance equilibrium corresponds to a quadratic equilibrium for different risk parameters. Due to the linearity assumption in the mean-variance preferences and with the help of the general theory on mean-variance hedging developed in the early 2000s, see Černý/Kallsen [5], the fixed-point problem can be solved explicitly. The case of general mean-variance preferences is covered in a forthcoming paper.

A challenge in moving from the one-period setup of [17] to multi-period and continuous time is that for the latter, the space of attainable payoffs is no longer spanned by the terminal dividends but depends on the equilibrium prices. In addition, one needs to impose integrability conditions on the admissible trading strategies. These conditions can preclude the existence of an equilibrium. Indeed, we exhibit an example where the only candidate equilibrium is such that the buy-and-hold strategy for the risky asset is not admissible in an L^2 -sense. If the asset has positive net supply, by linearity one of the agents must use an inadmissible strategy; therefore, this cannot be an equilibrium market. We give sufficient conditions to ensure that the required integrability conditions are satisfied, so that this issue is prevented and an equilibrium exists.

The remainder of this article is organised as follows. In Section 2, we introduce the market model and introduce the notion of an equilibrium. In Section 3, we study the case of agents with quadratic utility preferences and prove the main existence and uniqueness result in Theorem 3.7. In Section 4, we then consider the case of agents with linear mean–variance preferences and prove existence of equilibria under mild assumptions on the model primitives in Theorem 4.14. Key results on mean–variance hedging that are used throughout the main body of the paper can be found in Appendix A. All proofs and some auxiliary results are delegated to Appendix B.

2 Model and preliminary results

2.1 Financial market

We work on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F} = (\mathcal{F}_t)_{0 \leq t \leq T}, P)$ with a fixed finite time horizon $T \in (0, \infty)$. We assume that the filtration \mathbb{F} satisfies the usual conditions of right-continuity and completeness. Moreover, we assume that \mathcal{F}_0 is *P*-trivial and $\mathcal{F}_T = \mathcal{F}$. We denote the space of locally square-integrable martingales starting from zero by $\mathcal{M}^2_{0,\text{loc}}$.

We consider a financial market consisting of $1 + d = 1 + d_1 + d_2$ assets. The first asset, with price process S^0 , serves as numéraire, and we assume that $(S_t^0)_{0 \le t \le T} \equiv 1$.¹ In addition, we consider d_1 financial assets with price processes $S^{(1)} = (S_t^1, \ldots, S_t^{d_1})_{0 \le t \le T}$ and d_2 productive assets (sometimes also referred to as real assets) with price processes $S^{(2)} = (S_t^{d_1+1}, \ldots, S_t^{d_1+d_2})_{0 \le t \le T}$. The risky assets are collectively expressed as $S := (S^{(1)}, S^{(2)})$. In the following, we likewise use the notation $x = (x^{(1)}, x^{(2)})$ for each $x \in \mathbb{R}^{d_1+d_2}$ with $x^{(i)} \in \mathbb{R}^{d_i}$.

We assume that the price processes $S^{(1)}$ and $S^{(2)}$ are not given a priori but rather determined in equilibrium between K agents trading in the market.

We assume that the initial value and volatility structure of the financial assets are predetermined and known by the market participants, i.e., for $j \in \{1, ..., d_1\}$, we have

$$S_t^j = S_0^j + M_t^j + A_t^j, \quad 0 \le t \le T,$$
(2.1)

where $S_0^j \in \mathbb{R}$ and the local martingale part $M^j \in \mathcal{M}_{0,\text{loc}}^2$ are given a priori. The predictable finite-variation process (which is null at time 0) is to be determined in equilibrium. The financial assets may be regarded as securities constructed by the market participants to enable the trading of short-term risks, determined implicitly by the dynamics of $M^{(1)}$, at appropriate prices set by the market, which are reflected in the dynamics of $A^{(1)}$.

We assume that each productive asset $j \in \{d_1 + 1, \dots, d_1 + d_2\}$ with price process S^j entitles the owner to a random terminal dividend $D^j \in L^2$ at time T so that S^j satisfies the terminal condition

$$S_T^j = D^j; (2.2)$$

the rest of the price process $(S^j)_{0 \le t < T}$ is to be determined by the market in equilibrium.

Finally, we assume that each asset S^j is a local L^2 -semimartingale for $j \in \{1, \ldots, d\}$. This means that there exists a localising sequence of stopping times $(\tau_n)_{n \in \mathbb{N}}$ such that each stopped process $S^{j,\tau_n} = (S^j_{\tau_n \wedge t})_{0 \leq t \leq T}$ is an L^2 -semimartingale, in the sense that

 $\sup \left\{ E[(S^{j,\tau_n}_{\sigma})^2] : \sigma \text{ stopping time} \right\} < \infty;$ (2.3)

¹The dynamics of the numéraire asset cannot be determined in equilibrium as we do not consider intertemporal consumption.

see Delbaen/Schachermayer [12] and Černý/Kallsen [5] for details. We refer to this property by calling (1, S) a *local* L^2 -market. Note that by [5, Lemma A.2], a stochastic process is a local L^2 -semimartingale if and only if it is a special semimartingale whose local martingale part is locally square-integrable. In view of (2.1), this is only a condition on the productive assets as it is automatically satisfied by the financial assets.

2.2 Admissible strategies

In order to describe trading in the market (1, S), we need to specify which strategies are considered *admissible*. To this end, we follow a two-step approach as in Černý/Kallsen [5], to which we also refer for further details.

Definition 2.1. Let (1, S) be a local L^2 -market. A simple integrand for S is a process of the form $\vartheta = \sum_{i=1}^{m-1} \xi_i \mathbb{1}_{[\sigma_i,\sigma_{i+1}]}$, where $m \in \mathbb{N}$, $0 \leq \sigma_1 \leq \cdots \leq \sigma_m$ are [0,T]-valued stopping times, and each ξ_i is a bounded \mathcal{F}_{σ_i} -measurable random vector in \mathbb{R}^d , such that each stopped process $S^{j,\sigma_m} = (S^j_{\sigma_m \wedge t})_{0 \leq t \leq T}$ is an L^2 -semimartingale for $j = 1, \ldots, d$. We denote by $\Theta_{\text{simple}}(S)$ the linear space of all simple integrands for S. We also let L(S) be the set of predictable S-integrable processes on [0, T]; see Jacod/Shiryaev [16, III.6.17].

Definition 2.2. Let (1, S) be a local L^2 -market. Then $\vartheta \in L(S)$ is called L^2 -admissible for S if $\vartheta \cdot S_T \in L^2$ and there exists a sequence $(\vartheta^n)_{n \in \mathbb{N}}$ in $\Theta_{\text{simple}}(S)$ such that

- 1) $\vartheta^n \bullet S_T \xrightarrow{L^2} \vartheta \bullet S_T$,
- **2)** $\vartheta^n \bullet S_\tau \xrightarrow{P} \vartheta \bullet S_\tau$ for all [0, T]-valued stopping times τ .

Here, $\vartheta \cdot S = (\vartheta \cdot S_t)_{0 \le t \le T}$ denotes the stochastic integral $\vartheta \cdot S_t = \int_0^t \vartheta_r dS_r$ for $t \in [0, T]$. The set of all L^2 -admissible trading strategies is denoted by $\overline{\Theta}(S)$.

Remark 2.3. Our definition of L^2 -admissible strategies slightly differs from the definition given in [5], because we stipulate 2) for all stopping times τ and not only for deterministic times $t \in [0, T]$. However, under [5, Assumption 2.1], i.e., if there exists an equivalent local martingale measure (ELMM) Q for S with $\frac{dQ}{dP} \in L^2(P)$, both definitions coincide. The reason for this change is that it allows us to use dynamic programming arguments even if there does not exist an ELMM Q for S with $\frac{dQ}{dP} \in L^2(P)$, as in Czichowsky/Schweizer [10].

We denote by $(e_t^j)_{0 \le t \le T} \equiv (0, \ldots, 0, 1, 0, \ldots, 0) \in \mathbb{R}^d$ the buy-and-hold strategy of the *j*-th risky asset, where 1 is in the *j*-th position. In general, this strategy will not be L^2 -admissible for S because we only assume S^j to be a local L^2 -semimartingale. However, if S^j is an L^2 -semimartingale (i.e., $\sup\{E[(S_{\sigma}^j)^2]: \sigma \text{ stopping time}\} < \infty$ for each $j \in \{1, \ldots, d_1 + d_2\}$), then automatically $e^j \in \Theta_{\text{simple}}(S) \subseteq \overline{\Theta}(S)$ by [5, Corollary 2.9].

Since different strategies may lead to the same stochastic integral, we naturally identify strategies $\vartheta \in \overline{\Theta}(S)$ via the following equivalence relation.

Definition 2.4. Let (1, S) be a local L^2 -market. Then $\vartheta, \vartheta' \in \overline{\Theta}(S)$ are called *S*-equivalent if $\vartheta \cdot S$ and $\vartheta' \cdot S$ are indistinguishable. In this case, we write $\vartheta =_S \vartheta'$; see Czichowsky/Schweizer [9] for more details on how to represent different equivalent classes via the so-called projection onto the predictable range.

We assume that trading is frictionless and that market participants choose self-financing portfolios of the form $(\vartheta_t^0, \vartheta_t)_{0 \le t \le T}$, where ϑ^0 is a predictable process, $\vartheta \in \overline{\Theta}(S)$ and the wealth process $(V_t(\vartheta^0, \vartheta))_{0 \le t \le T}$ satisfies the self-financing condition

$$V_t(\vartheta^0,\vartheta) = \vartheta_t^0 + \vartheta_t^\top S_t = \vartheta_0^0 + \vartheta_0^\top S_0 + \vartheta \bullet S_t, \quad P\text{-a.s. for all } t \in [0,T].$$

Since we shall include the initial wealth of the agents into their endowments, a self-financing portfolio can be specified in terms of $\vartheta \in \overline{\Theta}(S)$ alone. We do not impose any further conditions on the strategies of the agents: for example, they are allowed to borrow or short sell any asset.

2.3 Agents and preferences

We consider $K \ge 1$ agents trading in the financial market (1, S). We assume that each agent $k \in \{1, \ldots, K\}$ owns a *traded endowment* at time 0, consisting of $\eta^{k,j} \in \mathbb{R}$ units of asset $j \in \{0, \ldots, d_1 + d_2\}$, and is also entitled to receive a *non-traded endowment* at time T, which consists of a random income $\Xi^{k,n} \in L^2$.

Because we assume zero interest rates and there are no liquidity constraints on the portfolios of the agents, it does not matter whether a fixed amount of cash is received via the traded or non-traded endowment. Thus by transferring it to the non-traded endowment, we may assume that each agent starts with zero cash. We also make the assumption that the financial assets are in *zero net supply*. This means that these assets are created and traded internally by the market participants, so that any long and short positions in the financial assets must net out, i.e., $\sum_{k=1}^{K} \eta^{k,j} = 0$. Since the initial prices $S_0^1, \ldots, S_0^{d_1}$ are known a priori, each agent is indifferent between receiving an endowment consisting of units of the financial assets or the corresponding cash value via the non-traded endowment. We may thus assume without loss of generality that $\eta^{k,j} = 0$ for $j \in \{0, \ldots, d_1\}$ and $k \in \{1, \ldots, K\}$.

By contrast, the agents may have a nontrivial traded endowment consisting of productive assets. We denote by $\Xi^{k,\text{tr}} := \eta^{k,(2)} D^{(2)} \in L^2$ the value of the traded endowment of agent k at time T. The *total endowment* of agent k at time T is then given by

$$\Xi^k = \Xi^{k, \text{tr}} + \Xi^{k, \text{n}}.$$
(2.4)

Each agent $k \in \{1, \ldots, K\}$ interacts with the market by buying and selling assets according to an L^2 -admissible strategy $\vartheta \in \overline{\Theta}(S)$, which includes the original endowment $\eta^{k,(2)}$ in the productive assets. Since the agent does not own any riskless or financial assets at time 0, their initial wealth is $\eta^{k,(2)} {}^{\top}S_0^{(2)}$, which is the initial value of their traded endowment. Agent k can then generate the wealth process $\eta^{k,(2)} {}^{\top}S_0^{(2)} + \vartheta \cdot S$ by trading with the strategy ϑ in a selffinancing way. Since they additionally receive the non-traded endowment $\Xi^{k,n}$ at time T, their terminal wealth at time T is given by

$$V_T^k(\vartheta) = \eta^{k,(2)} {}^{\mathsf{T}}S_0^{(2)} + \vartheta \bullet S_T + \Xi^{k,\mathbf{n}}.$$
(2.5)

Note that the traded endowment has the terminal value

$$\Xi^{k,\mathrm{tr}} = \eta^{k,(2)} {}^{\top} D^{(2)} = \eta^{k,(2)} {}^{\top} S_T^{(2)} = \eta^{k,(2)} {}^{\top} S_0^{(2)} + \eta^k \bullet S_T,$$

since $\eta^k = (0, \eta^{k,(2)})$ is constant. Thus the terminal wealth of agent k at time T can be equivalently written as

$$V_T^k(\vartheta) = \Xi^{k, \text{tr}} - \eta^k \bullet S_T + \vartheta \bullet S_T + \Xi^{k, \text{n}} = (\vartheta - \eta^k) \bullet S_T + \Xi^k$$
(2.6)

in terms of the total endowment defined in (2.4).

From the right-hand side of (2.6), we see that the total wealth at time T consists of the total endowment as well as any gains or losses generated by the strategy $\vartheta - \eta^k$. This difference may be interpreted as a discretionary strategy that is employed by the agent in addition to the fixed endowment η^k . The right-hand side of (2.5) gives an alternative interpretation. Instead of keeping the traded endowment, agent k may immediately sell it for the price of $\eta^{k,(2)} {}^{\top}S_0^{(2)}$

and then trade with the strategy ϑ ; the non-traded endowment $\Xi^{k,n}$ is then added to the wealth at time T. However, we note that the price $S_0^{(2)}$ is not known a priori, but rather determined in equilibrium. Thus, the right-hand side of (2.6) is more useful for solving the equilibrium problem, since the total endowment Ξ^k is fixed by the primitives, so that only the stochastic integral term $(\vartheta - \eta^k) \cdot S_T$ depends on the (unknown) dynamics of S.

Each agent $k \in \{1, \ldots, K\}$ has preferences over terminal wealth at time T described by a functional $\mathcal{U}_k : L^2 \to \mathbb{R}$. Agent k seeks to maximise utility from terminal wealth at time T, i.e., to solve the problem

$$\mathcal{U}_k((\vartheta - \eta^k) \bullet S_T + \Xi^k) \longrightarrow \max_{\vartheta \in \overline{\Theta}(S)} !$$
(2.7)

We consider two types of functionals \mathcal{U}_k : a quadratic utility functional, which is the subject of Section 3, and a linear mean-variance functional, which is considered in Section 4.

The quadratic utility functional for agent $k \in \{1, ..., K\}$ is given by

$$\mathcal{U}_{k}^{Q}(V) = E[U_{k}^{Q}(V)] = E[2\gamma_{k}V - V^{2}], \quad V \in L^{2},$$
(2.8)

where $U_k^Q(x) := 2\gamma_k x - x^2$ for some risk-tolerance $\gamma_k \in \mathbb{R}$. The parameter γ_k is also sometimes called the *bliss point* of agent k, as it is the optimal wealth that the agent would like to attain at time T in order to maximise their utility.

The linear mean-variance functional is given by

$$\mathcal{U}_k^{\mathrm{MV}}(V) := U_k^{\mathrm{MV}}(E[V], \operatorname{Var}[V]) = E[V] - \frac{\operatorname{Var}[V]}{2\lambda_k}, \quad V \in L^2,$$
(2.9)

where $U_k^{\text{MV}}(\mu, \sigma^2) := \mu - \frac{\sigma^2}{2\lambda_k}$ for some risk tolerance $\lambda_k > 0$.

Remark 2.5. (a) Note that (2.8) differs slightly from the more standard definition of quadratic utility

$$\tilde{\mathcal{U}}_k^{\mathbf{Q}}(V) := E[\tilde{\mathcal{U}}_k^{\mathbf{Q}}(V)] = E\left[V - \frac{V^2}{2\gamma_k}\right], \quad V \in L^2,$$

where $\tilde{U}_k^{\mathbf{Q}}(x) := x - \frac{1}{2\gamma_k}x^2$ for $\gamma_k > 0$. Of course, both formulations are equivalent for $\gamma_k > 0$ (which is the economically relevant case). The reason we use (2.8) is that it is also well defined in the case $\gamma_k \leq 0$, which will be useful for technical reasons when we link quadratic utility to linear mean–variance preferences.

(b) Despite their apparent similarity, the maximisation problems induced by the two preference functionals (2.8) and (2.9) are not equivalent. Indeed, the quadratic utility functional can be written as

$$\mathcal{U}_{k}^{Q}(V) = E[U_{k}^{Q}(V)] = E\left[V - \frac{V^{2}}{2\gamma_{k}}\right] = E[V] - \frac{\operatorname{Var}[V]}{2\gamma_{k}} - \frac{E[V]^{2}}{2\gamma_{k}} = \mathcal{U}_{k}^{Q}(V) - \frac{E[V]^{2}}{2\gamma_{k}},$$

which is not of the form (2.9) due to the presence of the additional term.

2.4 Equilibrium

We can now formulate the key notion of an equilibrium market, which we adapt from the classical concept of a Radner equilibrium. We take the primitives $S_0^{(1)}, M^{(1)}, D^{(2)}, \eta^k, \Xi^{k,n}$ and \mathcal{U}_k defined in Sections 2.1 and 2.3 as given, where \mathcal{U}_k is either $\mathcal{U}_k^{\mathrm{Q}}$ or $\mathcal{U}_k^{\mathrm{MV}}$.

Definition 2.6. A local L^2 -market $(1, S^{(1)}, S^{(2)})$ is called an *equilibrium market* if it satisfies the following conditions:

- 1) For each agent $k \in \{1, ..., K\}$, the maximisation problem (2.7) has a solution $\hat{\vartheta}^k \in \overline{\Theta}(S)$ that is unique up to S-equivalence.
- 2) The market clears, i.e., for $t \in [0, T]$,

$$\sum_{k=1}^{K} \hat{\vartheta}_{t}^{k,j} =_{S} \bar{\eta}^{j} := \begin{cases} 0, & \text{if } j \in \{1, \dots, d_{1}\}, \\ \sum_{k=1}^{K} \eta^{k,j}, & \text{if } j \in \{d_{1}+1, \dots, d_{1}+d_{2}\}. \end{cases}$$
(2.10)

3) $e^j \in \overline{\Theta}(S)$ for $j \in \{d_1 + 1, \dots, d_1 + d_2\}$, i.e., the buy-and-hold strategies of the productive assets are L^2 -admissible.

If $(1, S^{(1)}, S^{(2)})$ is an equilibrium market with respect to the mean-variance functionals defined in (2.8) or (2.9) for some parameters $(\gamma_k)_{k=1}^K$ or $(\lambda_k)_{k=1}^K$, we say that it is a quadratic equilibrium market or mean-variance equilibrium market, respectively.

Remark 2.7. The only slightly non-standard requirement in Definition 2.6 is 3). It ensures that each buy-and-hold strategy η^k is L^2 -admissible, which is a natural requirement since the agents should be allowed to simply hold their respective traded endowments. Mathematically, it also ensures that $\hat{\vartheta}^k \in \overline{\Theta}(S)$ if and only if $\hat{\vartheta}^k - \eta^k \in \overline{\Theta}(S)$, which will allow us to find the optimal strategies $\hat{\vartheta}^k$ by first solving for $\hat{\vartheta}^k - \eta^k$; recall also the discussion below (2.6).

In the remainder of the paper we seek to find equilibrium markets for quadratic and meanvariance preferences. More precisely, we look for conditions on the primitives that ensure the existence and uniqueness of a corresponding equilibrium market, and we seek to characterise that market. We shall study quadratic equilibria in Section 3. This in turn will allow us to obtain results on mean-variance equilibria in Section 4. In both cases, we start by studying the individual optimisation problems of the agents with respect to a (hypothetical) price process S, and then proceed to determining the markets (1, S) that lead to equilibrium.

3 Quadratic utility

In this section, we consider the situation of quadratic utility, i.e., each agent k solves the problem

$$\mathcal{U}_{k}^{Q}(V_{T}^{k}(\vartheta)) = E\left[2\gamma_{k}\left((\vartheta - \eta^{k})\bullet S_{T} + \Xi^{k}\right) - \left((\vartheta - \eta^{k})\bullet S_{T} + \Xi^{k}\right)^{2}\right] \longrightarrow \max_{\vartheta \in \overline{\Theta}(S)}!$$
(3.1)

As customary in the equilibrium literature, we first study the individual problem (3.1) for a given local L^2 -market (1, S) and then solve for equilibrium.

3.1 Individual optimality

Throughout this section, we fix a local L^2 -market (1, S) that satisfies (2.1) and (2.2).

In order to study the maximisation problem (3.1), we note that it is closely linked to the so-called mean-variance hedging problems for a payoff $H \in L^2$:

• The mean-variance hedging (MVH) problem is given by

$$E[(\vartheta \bullet S_T - H)^2] \longrightarrow \min_{\vartheta \in \overline{\Theta}(S)}!$$
(3.2)

• The extended mean-variance hedging (exMVH) problem is given by

$$E[(c+\vartheta \bullet S_T - H)^2] \longrightarrow \min_{(c,\vartheta) \in \mathbb{R} \times \overline{\Theta}(S)}!$$
(3.3)

We say that (3.2) has a unique solution if $\vartheta^1 =_S \vartheta^2$ for any two solutions $\vartheta^1, \vartheta^2 \in \overline{\Theta}(S)$. Similarly, we say that (3.3) has a unique solution if $c_1 = c_2$ and $\vartheta^1 =_S \vartheta^2$ for any two solutions $(c_1, \vartheta^1), (c_2, \vartheta^2) \in \mathbb{R} \times \overline{\Theta}(S)$. A sufficient condition for the existence and uniqueness of solutions to (3.2) and (3.3) is given in Proposition A.7. We refer to Schweizer [28] for a recent overview of mean-variance hedging.

The following result shows that the quadratic utility problem (3.1) is equivalent to a MVH problem (3.2). We recall that all proofs are given in Appendix B.

Lemma 3.1. Let (1, S) be a local L^2 -market and assume that $\eta^k \in \overline{\Theta}(S)$. Then the following are equivalent:

- (a) The quadratic utility problem (3.1) has a unique solution $\hat{\vartheta}^k \in \overline{\Theta}(S)$.
- (b) For $H^k := \gamma_k \Xi^k$, the MVH problem

$$E[(\vartheta \bullet S_T - H^k)^2] \to \min_{\vartheta \in \overline{\Theta}(S)} !$$
(3.4)

has a unique solution $\vartheta^{\text{MVH}}(H^k) \in \overline{\Theta}(S)$.

If either statement holds, then $\hat{\vartheta}^k =_S \eta^k + \vartheta^{\text{MVH}}(H^k)$.

In the above result, $H^k = \gamma_k - \Xi^k$ may be interpreted as the the additional wealth that agent k would like to obtain in order to reach the bliss point γ_k .

We now use part (b) of Lemma 3.1 to deduce a more explicit decomposition for the optimal strategy $\hat{\vartheta}^k$ of agent k into a hedging and pure investment part, where the *hedging problem* of agent k is the extended MVH problem

$$E[(c+\vartheta \bullet S_T - \Xi^k)^2] \to \min_{c \in \mathbb{R}, \vartheta \in \overline{\Theta}(S)}!$$
(3.5)

and the *pure investment problem* is the MVH problem

$$E[(\vartheta \bullet S_T - 1)^2] \to \min_{\vartheta \in \overline{\Theta}(S)} !$$
(3.6)

We assume that $E[(\vartheta^{\text{MVH}}(1) \bullet S_T - 1)^2] > 0$, or equivalently $\vartheta^{\text{MVH}}(1) \bullet S_T \neq 1$, which can be seen as a weak variant of the condition of 'the law of one price' in Černy/Czichowsky [4], that is, a weak no-free-lunch condition on S. Under that assumption, we obtain the following decomposition.

Proposition 3.2. Let (1, S) be a local L^2 -market with $\eta^k \in \overline{\Theta}(S)$. Suppose that there is a unique solution $\vartheta^{\text{MVH}}(1)$ to the pure investment problem (3.6) and $E[(\vartheta^{\text{MVH}}(1) \cdot S_T - 1)^2] > 0$. Then there is a unique solution $\vartheta^{\text{MVH}}(H^k) \in \overline{\Theta}(S)$ to the MVH problem (3.4) if and only if there is a unique solution $(c_k, \vartheta^{\text{ex}}(\Xi^k))$ to the exMVH problem (3.3) with $H = \Xi^k$. In this case ϑ^k can be decomposed as

$$\hat{\vartheta}^k =_S \eta^k - \vartheta^{\text{ex}}(\Xi^k) + (\gamma_k - c_k)\vartheta^{\text{MVH}}(1).$$
(3.7)

The above results shows that $\hat{\vartheta}^k$ can be decomposed into the traded endowment η^k , a hedging component for the total endowment Ξ^k and a an investment component, which is proportional to $\vartheta^{\text{MVH}}(1)$.

3.2 The representative agent

We now return to the question of finding a price process (1, S) that leads to an equilibrium in the sense of Definition 2.6. In order to study quadratic equilibrium markets, we use a standard idea from financial economics to consider a *representative agent* that holds the aggregate endowment of all agents, i.e., the representative agent owns both $\bar{\eta} = \sum_{k=1}^{K} \eta^k$ units of the assets as well as the sum of the non-traded endowments of the agents. Equivalently, the representative agent receives the total endowment $\bar{\Xi} = \sum_{k=1}^{K} \Xi^k$. By the same argument as in (2.6), the representative agent can attain the terminal wealth $(\vartheta - \bar{\eta}) \cdot S_T + \bar{\Xi}$ by trading with a strategy $\vartheta \in \overline{\Theta}(S)$.

The utility function of the representative agent is defined by

$$\bar{U}_{\lambda}^Q(x) = \sup\left\{\sum_{k=1}^K \lambda_k U_k^Q(x_k) : x_1, \dots, x_K \in \mathbb{R}^d, \sum_{k=1}^k x_k = x\right\},\$$

where $\lambda = (\lambda_1, \ldots, \lambda_K) \in \mathbb{R}^K$ is a fixed set of *Negishi weights.*² We denote by $\bar{\gamma} := \sum_{k=1}^K \gamma_k$ the aggregate risk tolerance and make the ansatz $\lambda_k^* := \frac{\bar{\gamma}}{\gamma_k} \mathbf{1}_{\{\gamma_k \neq 0\}} + K \mathbf{1}_{\{\bar{\gamma}=0\}}$. Moreover, we set $\bar{U}^Q := \bar{U}^Q_{\lambda^*}$ for simplicity. Distinguishing between the cases $\bar{\gamma} \neq 0$ and $\bar{\gamma} = 0$, it is not difficult to check that

$$\bar{U}^Q(x) = \sup\left\{\sum_{k=1}^K \lambda_k (2\gamma_k x_k - x_k^2) : x_1, \dots, x_K \in \mathbb{R}^d, \sum_{k=1}^K x_k = x\right\} = 2\bar{\gamma}x - x^2,$$

so that the utility function of the representative agent is of the same form as the utility function of the individual agents. The representative agent then solves the maximisation problem

$$E\left[\bar{U}^{Q}\left(\left(\vartheta-\bar{\eta}\right)\bullet S_{T}+\bar{\Xi}\right)\right]\to\max_{\vartheta\in\overline{\Theta}(S)}!$$
(3.8)

From a mathematical perspective, (3.8) has exactly the same structure as the individual maximisation problem (3.1). Thus we get an analogue of Lemma 3.1 for the representative agent.

Lemma 3.3. Let (1, S) be a local L^2 -market with $\bar{\eta} \in \overline{\Theta}(S)$. Then the following are equivalent:

- (a) The optimisation problem (3.8) has a unique solution $\bar{\vartheta} \in \overline{\Theta}(S)$.
- (b) For $\overline{H} := \overline{\gamma} \overline{\Xi}$, the MVH problem

$$E[(\vartheta \bullet S_T - \bar{H})^2] \to \min_{\vartheta \in \overline{\Theta}(S)} !$$
(3.9)

has a unique solution $\vartheta^{\text{MVH}}(\overline{H}) \in \overline{\Theta}(S)$.

If either statement holds, then $\bar{\vartheta} =_S \bar{\eta} + \vartheta^{\text{MVH}}(\bar{H})$.

In the above result, $\bar{H} = \bar{\gamma} - \bar{\Xi} = \sum_{k=1}^{K} H_k$ may be interpreted as the aggregate additional wealth that the agents would (collectively) like to obtain in order to reach the aggregate bliss point $\bar{\gamma}$.

Similarly to Lemma 3.2, we can also decompose the optimal strategy $\bar{\vartheta}$ of the representative agent into a hedging and a pure investment part, where the *hedging problem of the representative agent* is the extended MVH problem

$$E[(c+\vartheta \bullet S_T - \bar{\Xi})^2] \to \min_{c \in \mathbb{R}, \vartheta \in \overline{\Theta}(S)}!$$
(3.10)

 $^{^{2}}$ Usually, the convention in the literature is that the Negishi weights are positive and sum up to 1. Since we allow for zero and negative risk tolerances, we need to generalise this.

Proposition 3.4. Let (1, S) be a local L^2 -market such that $\bar{\eta} \in \overline{\Theta}(S)$. Suppose that there is a unique solution $\vartheta^{\text{MVH}}(1)$ to the pure investment problem (3.6) and $E[(\vartheta^{\text{MVH}}(1) \bullet S_T - 1)^2] > 0$. Then the optimisation problem (3.8) of the representative agent has a unique solution $\bar{\vartheta} \in \overline{\Theta}(S)$ if and only if the exMVH problem (3.3) for $\bar{\Xi}$ has a unique solution $(c(\bar{\Xi}), \vartheta^{\text{ex}}(\bar{\Xi})) \in \mathbb{R} \times \overline{\Theta}(S)$. In this case $\bar{\vartheta}$ can then be decomposed as

$$\bar{\vartheta} =_{S} \bar{\eta} + \left(\bar{\gamma} - c(\bar{\Xi})\right) \vartheta^{\text{MVH}}(1) - \vartheta^{\text{ex}}(\bar{\Xi}).$$
(3.11)

The following result shows that the optimal strategy for the representative agent is given by the sum of the strategies of the individual agents. This result gives a characterisation of the aggregate demand for the risky assets, which is the key to finding an equilibrium market.

Lemma 3.5. Let (1, S) be a local L^2 -market. Assume that $\eta^1, \ldots, \eta^K \in \overline{\Theta}(S)$ and for each agent $k \in \{1, \ldots, K\}$, the individual optimisation problem (3.1) has a unique solution $\hat{\vartheta}^k$. Then the optimisation problem (3.8) of the representative agent has a unique solution $\bar{\vartheta}$ satisfying

$$\bar{\vartheta} =_S \sum_{k=1}^K \hat{\vartheta}^k. \tag{3.12}$$

3.3 Existence and uniqueness of equilibria

We can now prove our main result on the existence and uniqueness of quadratic equilibrium markets. In a first step, we show that a necessary condition for the existence of a quadratic equilibrium is that the martingale generated by $\bar{H} = \bar{\gamma} - \bar{\Xi}$ yields a (generalised) local martingale measure for (1, S).

Lemma 3.6. Let $(\bar{Z}_t)_{0 \le t \le T}$ be the (square-integrable) *P*-martingale given by $\bar{Z}_t = E[\bar{H} | \mathcal{F}_t]$. If $(1, S) = (1, S^{(1)}, S^{(2)})$ is a quadratic equilibrium market, then for each $j \in \{1, \ldots, d_1 + d_2\}$, the process $(\bar{Z}_t S^j_t)_{0 \le t \le T}$ is a local *P*-martingale.

In order to obtain also a sufficient condition for the existence of a quadratic equilibrium, we need to assume that the process \bar{Z} does not hit 0; this assumption holds in particular in the economically relevant case that $\bar{Z}_T = \bar{H} > 0$ so that \bar{Z} is strictly positive. Then by Lemma 3.6, $\bar{H}/E[\bar{H}]$ is the density of an equivalent local martingale measure for S.

For the following theorem, we need the Galtchouk–Kunita–Watanabe decomposition of \overline{Z} with respect to $M^{(1)}$, the local martingale part $M^{(1)}$ of the financial asset given in (2.1). We have

$$\bar{Z}_t = \bar{Z}_0 + \bar{\xi}^{(1)} \bullet M_t^{(1)} + M_t^{\bar{Z}}, \quad 0 \le t \le T,$$
(3.13)

where $\bar{\xi}^{(1)} \in L^2(M^{(1)})$ and $M^{\bar{Z}}$ is a square-integrable *P*-martingale strongly orthogonal to $M^{(1)}$. By Remark B.1, we may choose $\bar{\xi}^{(1)}$ in (3.13) in such a way that $\bar{\xi}^i \in L^2(M^i)$ for each $i \in \{1, \ldots, d_1\}$.

We can now formulate the main existence and uniqueness theorem for quadratic equilibria. **Theorem 3.7.** Assume that $\bar{Z}_t \neq 0$ and $\bar{Z}_{t-} \neq 0$ for all $t \in [0,T]$ *P-a.s.* If there exists a quadratic equilibrium $(1, S^{(1)}, S^{(2)})$, it is unique and explicitly given by

$$S_{t}^{j} = S_{0}^{j} + M_{t}^{j} - \int_{0}^{t} \frac{\mathrm{d}\langle \bar{Z}, M^{j} \rangle_{s}}{\bar{Z}_{s-}} = S_{0}^{j} + M_{t}^{j} - \sum_{i=1}^{d_{1}} \int_{0}^{t} \frac{\bar{\xi}_{s}^{i}}{\bar{Z}_{s-}} \mathrm{d}\langle M^{i}, M^{j} \rangle_{s}, \qquad j \in \{1, \dots, d_{1}\},$$
(3.14)

$$S_t^j = \frac{E[\bar{H}D^j \mid \mathcal{F}_t]}{\bar{Z}_t} = \frac{E[\bar{Z}_T D^j \mid \mathcal{F}_t]}{\bar{Z}_t}, \qquad j \in \{d_1 + 1, \dots, d_1 + d_2\}.$$
(3.15)

Conversely, the market $(1, S) = (1, S^{(1)}, S^{(2)})$ defined by (3.14) and (3.15) is a quadratic equilibrium if and only if $S^{(2)}$ is a local L^2 -semimartingale and for each $j \in \{d_1 + 1, \ldots, d_1 + d_2\}$, we have $e^j \in \overline{\Theta}(S)$.

The following result gives sufficient conditions on the primitives to ensure the existence of an equilibrium market. We will require that \overline{H} is strictly positive so that \overline{Z} and \overline{Z}_{-} are automatically positive. The remaining assumptions of Theorem 3.7 hold under the stronger assumption that $S^{(2)}$ is an L^2 -semimartingale, which yields for $j \in \{d_1 + 1, \ldots, d_1 + d_2\}$ that $e^j \in \Theta_{\text{simple}}(S) \subseteq \overline{\Theta}(S)$. We now give sufficient (but not necessary) conditions for $S^{(2)}$ to be an L^2 -semimartingale in the case $\overline{H} > 0$ *P*-a.s. This in turn implies that a quadratic equilibrium exists.

Lemma 3.8. Suppose that $\overline{H} > 0$ P-a.s. Then the process $(S_t^{(2)})_{0 \le t \le T}$ defined by (3.15) is an L^2 -semimartingale if any of the following conditions holds:

- (a) $D^j \in L^{\infty}(P)$ for $j \in \{d_1 + 1, \dots, d_1 + d_2\}.$
- (b) $\bar{H}, \bar{H}^{-1} \in L^{\infty}(P).$
- (c) $\bar{H} \in L^{1-p_1}(P) \cap L^{p_2}(P)$ and $D^j \in L^{2q_1q_2}(P)$ for all $j \in \{d_1 + 1, \dots, d_1 + d_2\}$, where $p_1, p_2 \in [1, \infty]$ and $1/p_i + 1/q_i = 1$ for $i \in \{1, 2\}$.³

In the above result, the first two sets of conditions are particularly easy to check; they are special cases of the more elaborate third condition.

We conclude this section by giving an example of a setup where an equilibrium market (in the sense of Definition 2.6) fails to exist due to integrability issues in the candidate for $S^{(2)}$, and not because the process \bar{Z} hits 0. We consider the simplest case of a market with no financial and one productive asset, i.e., $d_1 = 0$ and $d_2 = 1$. The setup is based on the counterexample in Černý/Kallsen [6], which in turn is inspired by the well-known counterexample of Delbaen/Schachermayer [13].

The key point is that the candidate equilibrium price process of the productive asset does not have sufficient integrability for the buy-and-hold strategies to be admissible. In that case, Lemma 3.1 cannot be applied, so that the existence of a solution to the optimisation problem (3.1) is not equivalent to the existence of a solution to the MVH problem (3.4). By part (b) of the proof of Theorem 3.7, there still exists in this case a unique solution to the MVH problem (3.4) for each agent k, but it is unclear whether (3.1) admits a solution.

Example 3.9. After rescaling the time interval $[0, \infty]$ to [0, T], there exists by [6, Lemma 2.2] a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F} = (\mathcal{F}_t)_{0 \le t \le T}, P)$ supporting two probability measures \bar{Q}, Q' and a continuous process $(X_t)_{0 \le t \le T}$ null at 0 with the following properties:

- 1) The measures \bar{Q}, Q' are equivalent to P, with $\frac{\mathrm{d}\bar{Q}}{\mathrm{d}\bar{P}}, \frac{\mathrm{d}Q'}{\mathrm{d}\bar{P}} \in L^2(P)$.
- 2) The process X is a uniformly integrable martingale under \bar{Q} , and a strict local martingale under Q'. Moreover, $X_T \in L^2(P)$.

Fix now some $\bar{\gamma} > 0$, and suppose that $d_1 = 0$, $d_2 = 1$, $D^1 := X_T$ and $\bar{\Xi} := \bar{\gamma} - \frac{d\bar{Q}}{dP}$, so that $\bar{H} = \frac{d\bar{Q}}{dP} > 0$ *P*-a.s. Then it follows from Theorem 3.7 that if a quadratic equilibrium market exists, it must satisfy $S_t^1 = E_{\bar{Q}}[X_T \mid \mathcal{F}_t] = X_t$. However, since $X = 1 \cdot X$ is not a *Q'*-martingale, the strategy $e^1 \equiv 1$ is not admissible by Černý/Kallsen [5, Corollary 2.5]. Therefore, a quadratic equilibrium does not exist in this setup.

³Note that we can always choose $p_2 \ge 2$ as \overline{H} is square-integrable by assumption.

3.4 Existence of equilibria in finite discrete time

Theorem 3.7 provides necessary and sufficient conditions for the existence and uniqueness of a quadratic equilibrium under the assumption that \overline{Z} and \overline{Z}_{-} do not hit 0. If we are only interested in existence (and not uniqueness), this assumption can be relaxed. In this section, we study the special (but important) case of finite discrete time $t \in \{0, \ldots, T\}$ for $T \in \mathbb{N}$. As we shall see, if \overline{Z} is allowed to hit 0, we either end up with nonexistence or nonuniqueness of equilibria.

In the following, we denote by $\Delta X_k = X_k - X_{k-1}$ the increment at time k of a stochastic process X in discrete time.

We start by giving necessary conditions for the existence of an equilibrium that are weaker than the assumption that the process \overline{Z} does not hit 0.

Lemma 3.10. A quadratic equilibrium $(1, (S_t)_{t \in \{0, \dots, T\}})$ can only exist if both of the conditions

$$\{\bar{Z}_{t-1} = 0\} \subseteq \{\bar{\xi}_t^i \Delta \langle M^i \rangle_t = 0\} \quad \text{for } i \in \{1, \dots, d_1\} \text{ and } t \in \{1, \dots, T\},$$
(3.16)

$$\{\bar{Z}_t = 0\} \subseteq \{E[\bar{H}D^j \mid \mathcal{F}_t] = 0\} \quad \text{for } j \in \{d_1 + 1, \dots, d_2\} \text{ and } t \in \{0, \dots, T-1\}$$
(3.17)

hold up to P-null sets.

Lemma 3.10 shows what can go wrong when \overline{Z} is allowed to hit 0. To understand (3.16) and (3.17) more clearly, consider the simple setup of a one-period model with T = 1, and suppose that $d_1 = 0$ and $d_2 = 1$. Thus, there exists a single productive asset S with terminal value $S_1 = D^1 = D$ and unknown initial value $S_0 \in \mathbb{R}$. Suppose that (3.17) is not satisfied, so that $\overline{Z}_0 = 0$ and $E[\overline{H}D] \neq 0$. In this case, there does not exist any value of $S_0 \in \mathbb{R}$ such that $\overline{Z}S$ is a martingale, since $\overline{Z}_0S_0 = 0$ regardless of that choice. On the other hand, if $\overline{Z}_0 = E[\overline{H}D] = 0$, then $\overline{Z}S$ is a martingale for any choice of $S_0 \in \mathbb{R}$, and one can check that (1, S) defines an equilibrium. This also illustrates the issue of non-uniqueness: if $\overline{Z}_t = E[\overline{H}D | \mathcal{F}_t] = 0$ for some $t \in \{0, \ldots, T-1\}$, then the price S_t in equilibrium can be set in an arbitrary way.

The issue is similar for the financial assets. Consider now a one-period model with $d_1 = 1$ and $d_2 = 0$ so that there exists a single financial asset S with $S_1 = S_0 + \Delta A_1 + \Delta M_1$, where $\Delta A_1 \in \mathbb{R}$ is unknown and ΔM_1 is the jump of a martingale. If (3.16) does not hold, then $\overline{Z}S$ is not a martingale for any choice of $A \in \mathbb{R}$, since \overline{Z} is a martingale and hence

$$E[Z_1(S_0 + \Delta A_1 + \Delta M_1)] = E[Z_1 \Delta M_1] = \xi_1 \Delta \langle M \rangle_1 \neq 0 = Z_0 S_0.$$

On the other hand, if $\xi_1 \Delta \langle M \rangle_1 = 0$ then $\overline{Z}S$ is a martingale for any value of ΔA_1 . Thus if $Z_t = 0$ and (3.16) is satisfied, then we would expect that the value ΔA_{t+1} is arbitrary.

As it turns out, (3.16) and (3.17) are the only significant requirements for the existence of an equilibrium (other than integrability conditions, cf. Example 3.9 above). We now show that if these conditions hold, then there exists an equilibrium (but in general will not be unique). In order to construct an explicit equilibrium, we use the fact that $\bar{Z}S$ is a local martingale by Lemma 3.6. However, the construction is more difficult here because we can no longer write \bar{Z} as the stochastic exponential of some (local) martingale \bar{N} . Instead, we define $(\bar{N}_t)_{t \in \{0,...,T\}}$ recursively by $\bar{N}_0 := 0$ and

$$\bar{N}_t := \bar{N}_{t-1} + \frac{\Delta \bar{Z}_t}{\bar{Z}_{t-1}} \mathbb{1}_{\{\bar{Z}_{t-1} \neq 0\}}, \quad t \in \{1, \dots, T\},$$
(3.18)

i.e., we arbitrarily set the increment $\Delta \bar{N}_t$ to 0 whenever $\bar{Z}_{t-1} = 0$. For each $s \in \{0, \ldots, T\}$, we also define the local martingale ${}^s \mathcal{E}(\bar{N}) = ({}^s \mathcal{E}(\bar{N})_t)_{t \in \{s, \ldots, T\}}$ by

$${}^{s}\mathcal{E}(\bar{N})_{t} := \prod_{k=s+1}^{t} (1 + \Delta \bar{N}_{k})$$
(3.19)

In the case where \bar{Z} does not hit 0, we have ${}^{s}\mathcal{E}(\bar{N})_{t} = \bar{Z}_{t}/\bar{Z}_{s}$ for each $s \leq t$. In other words, we may view ${}^{s}\mathcal{E}(\bar{N})$ as "restarting" \bar{Z} at time s (in a multiplicative way) with ${}^{s}\mathcal{E}(\bar{N})_{s} = 1$. The general case is similar, with the important difference that ${}^{s}\mathcal{E}(\bar{N})$ is absorbed at 0 whenever \bar{Z} hits 0 from a nonzero value. Thus, each process ${}^{s}\mathcal{E}(\bar{N})$ reproduces the dynamics of \bar{Z} until the latter hits 0. We note once again that the value of ΔN_{t} may be chosen arbitrarily whenever $\bar{Z}_{t-1} = 0$. Because we set $\Delta N_{t} = 0$ in that case by (3.18), the equilibrium constructed below defaults to behaving as a local martingale whenever \bar{Z} hits 0.

With this construction we obtain the following existence result in discrete time.

Theorem 3.11. Assume that ${}^{s}\mathcal{E}(\bar{N})$ is a square-integrable martingale for each $s \in \{0, \ldots, T\}$ and that (3.16) and (3.17) hold up to P-null sets. Define the process $(S_t)_{t \in \{0, \ldots, T\}}$ by

$$S_t^j := S_0^j + M_t^j + \sum_{k=1}^t \sum_{i=1}^{d_1} \left(-\frac{\bar{\xi}_k^i}{\bar{Z}_{k-1}} \mathbb{1}_{\{\bar{Z}_{k-1} \neq 0\}} \Delta \langle M^i, M^j \rangle_k \right), \quad j \in \{1, \dots, d_1\},$$
(3.20)

$$S_t^j := E[{}^t \mathcal{E}(\bar{N})_T D^j \mid \mathcal{F}_t], \quad j \in \{d_1 + 1, \dots, d_1 + d_2\}.$$
(3.21)

If $(S_t^{(2)})_{t \in \{0,\dots,T\}}$ is square-integrable, then $(1,S) = (1, S^{(1)}, S^{(2)})$ is a quadratic equilibrium.

4 Linear mean–variance preferences

In this section, we consider the situation of linear mean–variance preferences, i.e., each agent k solves the problem

$$\mathcal{U}_{k}^{\mathrm{MV}}(V_{T}^{k}(\vartheta)) = E[(\vartheta - \eta^{k}) \bullet S_{T} + \Xi^{k}] - \frac{\mathrm{Var}[(\vartheta - \eta^{k}) \bullet S_{T} + \Xi^{k}]}{2\lambda_{k}} \longrightarrow \max_{\vartheta \in \overline{\Theta}(S)}!$$
(4.1)

As in Section 3, we first study the individual problem (4.1) for a given local L^2 -market (1, S) and then solve for equilibrium.

4.1 Individual optimality

Throughout this section, we fix a local L^2 -market (1, S) and assume that $e^j \in \overline{\Theta}(S)$ for each $j \in \{d_1 + 1, \dots, d_1 + d_2\}$.

The linear mean-variance optimisation problem (4.1) is connected to the quadratic utility problem (3.1) and the MVH problem (3.2) via the classical notions of mean-variance efficient strategies and the mean-variance efficient frontier.

Definition 4.1. A strategy $\vartheta \in \overline{\Theta}(S)$ is mean-variance efficient with respect to $H \in L^2$ if there does not exist any other strategy $\vartheta' \in \overline{\Theta}(S)$ such that

$$E[\vartheta' \bullet S_T + H] \ge E[\vartheta \bullet S_T + H], \tag{4.2}$$

$$\operatorname{Var}[\vartheta' \bullet S_T + H] \le \operatorname{Var}[\vartheta \bullet S_T + H], \tag{4.3}$$

where one of the inequalities is strict.

We say that ϑ is mean-variance efficient for agent k if $\vartheta - \eta^k$ is mean-variance efficient with respect to $H = \Xi^k$. Recalling the notation $V_T^k(\vartheta) = \vartheta \cdot S_T + \Xi^k$ from (2.6), this is equivalent to the nonexistence of another strategy $\vartheta' \in \overline{\Theta}(S)$ such that both $E[V_T^k(\vartheta')] \ge E[V_T^k(\vartheta)]$ and $\operatorname{Var}[V_T^k(\vartheta')] \le \operatorname{Var}[V_T^k(\vartheta)]$, with one of the inequalities being strict. The mean-variance efficient frontier for agent k is defined as the set

$$\mathcal{E}^{k} := \left\{ \left(E[V_{T}^{k}(\vartheta)], \sqrt{\operatorname{Var}[V_{T}^{k}(\vartheta)]} \right) : \vartheta \in \overline{\Theta}(S) \text{ is mean-variance efficient} \\ \text{for agent } k \right\} \subseteq \mathbb{R} \times \mathbb{R}_{+}.$$

$$(4.4)$$

We begin our analysis by showing that any solution to the linear mean-variance problem (4.1) must be mean-variance efficient.

Lemma 4.2. Any solution $\hat{\vartheta}^k \in \overline{\Theta}(S)$ to (4.1) is mean-variance efficient for agent k.

In light of this result, the next step is to explicitly characterise the set of mean-variance efficient strategies for agent k. To that end, we introduce the following assumption.

Assumption 4.3. There exists an equivalent local martingale measure (ELMM) $Q \approx P$ for S with square-integrable density $dQ/dP \in L^2(P)$.

Assumption 4.3 is a well-known no-free-lunch condition in an L^2 -sense; see Stricker [31, Théorème 2]. If Assumption 4.3 is satisfied, then there exist unique (up to *S*-equivalence) minimisers $\vartheta^{\text{MVH}}(H) \in \overline{\Theta}(S)$ for the MVH problem (3.2) and $(c(H), \vartheta^{\text{ex}}(H)) \in \mathbb{R} \times \overline{\Theta}(S)$ for the exMVH problem (3.3) (see e.g. Černý/Kallsen [5, Lemma 2.11]; this also follows from Proposition A.7).

For future reference, we set

$$\ell := E[(\vartheta^{\text{MVH}}(1) \bullet S_T - 1)^2], \tag{4.5}$$

$$c_k := c(\Xi^k),$$
 $k \in \{1, \dots, K\},$ (4.6)

$$\varepsilon_k^2 := E[(c_k + \vartheta^{\text{ex}}(\Xi^k) \bullet S_T - \Xi^k)^2], \qquad k \in \{1, \dots, K\}.$$
(4.7)

We now find the set of mean-variance efficient strategies for agent k by relating them to the solutions of the quadratic utility problem (3.1) and the MVH problem (3.2) for the payoff $H^k(\gamma_k) := \gamma_k - \Xi^k$. In the subsequent corollary, we obtain an explicit formula for the meanvariance efficient frontier \mathcal{E}^k in terms of the triplet $(\ell, c_k, \varepsilon_k^2) \in (0, 1] \times \mathbb{R} \times \mathbb{R}_+$.

Proposition 4.4. Suppose the market (1, S) satisfies Assumption 4.3. For $\vartheta \in \overline{\Theta}(S)$, the following statements are equivalent:

- (a) ϑ is mean-variance efficient for agent k.
- (b) $\vartheta =_S \vartheta^k(y)$ for some $y \ge 0$, where

$$\vartheta^{k}(y) := y\vartheta^{\mathrm{MVH}}(1) + \eta^{k} - \vartheta^{\mathrm{ex}}(\Xi^{k}).$$
(4.8)

- (c) There exists some $\gamma_k \ge c_k$ such that $\vartheta \eta^k$ is the unique solution to the MVH problem (3.2) for the payoff $H^k(\gamma_k) := \gamma_k \Xi^k$.
- (d) There exists some $\gamma_k \geq c_k$ such that ϑ is the unique solution to the quadratic utility problem (3.1) with risk tolerance γ_k .

Moreover, the constant γ_k can be chosen to be the same in (c) and (d), and $y \ge 0$ can be chosen so that $y + c_k = \gamma_k$.

With the help of Proposition 4.4 and Lemma B.3, it is now straightforward to determine the mean-variance efficient frontier \mathcal{E}^k for agent k.

Corollary 4.5. The mean-variance efficient frontier for agent k is given by

$$\mathcal{E}^{k} = \left\{ \left(\mu_{k}(y), \sigma_{k}(y) \right) = \left(c_{k} + (1 - \ell)y, \sqrt{\varepsilon_{k}^{2} + \ell(1 - \ell)y^{2}} \right) : y \ge 0 \right\}.$$
(4.9)

We recall from Lemma 4.2 that any solution to the mean-variance problem (4.1) is meanvariance efficient for agent k. We can now use Proposition 4.4 and Corollary 4.5 together with the linear form of U_k^{MV} to find the solution to (4.1).

Lemma 4.6. Suppose the market (1, S) satisfies Assumption 4.3. The unique optimal strategy for (4.1) is $\vartheta^k(\lambda_k/\ell)$, where $\vartheta^k(\cdot)$ is given by (4.8).

4.2 Connections to quadratic equilibria

Now that we have characterised the solution to the individual linear mean-variance problem (4.1) in Lemma 4.6, we return to the question of finding a linear mean-variance equilibrium in the sense of Definition 2.6. The first step is to show that any linear mean-variance equilibrium (1, S) satisfying Assumption 4.3 must also be a quadratic equilibrium with respect to some parameters $\gamma_1, \ldots, \gamma_K$. This will allow us to apply the results from Section 3 in order to find linear mean-variance equilibria.

Lemma 4.7. Suppose the market (1, S) is a linear mean-variance equilibrium market and satisfies Assumption 4.3. Then (1, S) is also a quadratic equilibrium market with risk tolerances $\gamma_1, \ldots, \gamma_K$, where $\gamma_k := c_k + \lambda_k/\ell$.

At a first glance, it now seems a straightforward task to find a mean-variance equilibrium by combining Theorem 3.7 with Lemma 4.7. However, things are not so simple because the constants ℓ and c_k that appear in Lemma 4.7 are determined implicitly in terms of the equilibrium price S, and not just the primitives defined in Section 2. Thus, we do not know ℓ and c_k a priori. Moreover, the quadratic equilibrium given by Theorem 3.7 need not satisfy Assumption 4.3 in general.

Nevertheless, Lemma 4.7 suggests that one should look for a linear mean-variance equilibrium within the class of quadratic equilibria given in Theorem 3.7 for some choice of parameters $\gamma_1, \ldots, \gamma_K$. In order to ensure that these quadratic equilibria satisfy Assumption 4.3, we make for the remainder of this section the following assumption on $\overline{\Xi}$.

Assumption 4.8. The aggregate endowment $\overline{\Xi}$ is nonnegative and bounded, so that

$$0 \le \bar{\gamma}_0 := \operatorname{ess\,sup}\bar{\Xi} < \infty. \tag{4.10}$$

Lemma 4.9. Assume that Assumption 4.8 is satisfied. For $\bar{\gamma} > \gamma_0$, set $\bar{H}(\bar{\gamma}) := \bar{\gamma} - \bar{\Xi}$ and define the process $\bar{Z}(\bar{\gamma})$ by $\bar{Z}_t(\bar{\gamma}) := E[\bar{H}(\bar{\gamma}) | \mathcal{F}_t]$. Define the process $S(\bar{\gamma})$ by (3.14) and (3.15) with \bar{H} and \bar{Z} replaced by $\bar{H}(\bar{\gamma})$ and $\bar{Z}(\bar{\gamma})$, respectively. Then $(1, S(\bar{\gamma}))$ is the unique quadratic equilibrium for any choice of parameters $\gamma_1, \ldots, \gamma_K \in \mathbb{R}$ such that $\sum_{k=1}^K \gamma_k = \bar{\gamma}$. Moreover, $S^{(2)}(\bar{\gamma})$ is an L^2 -semimartingale and $S(\bar{\gamma})$ admits an equivalent local martingale measure $Q(\bar{\gamma})$ with bounded density $dQ(\bar{\gamma})/dP := \bar{H}(\bar{\gamma})/E[\bar{H}(\bar{\gamma})]$. In particular, $S(\bar{\gamma})$ satisfies Assumption 4.3.

We now look for a mean-variance equilibrium among the set of "nice" quadratic equilibria $(1, S(\bar{\gamma}))$ given by Lemma 4.9 for $\bar{\gamma} > \bar{\gamma}_0$. In other words, we want to determine which values of $\bar{\gamma} \in (\bar{\gamma}_0, \infty)$ lead to a linear mean-variance equilibrium. We note that, in general, there may exist other mean-variance equilibria with aggregate risk tolerance $\bar{\gamma} \leq \bar{\gamma}_0$, in which case Assumption 4.3 need not be satisfied.

To find a linear mean-variance equilibrium, we study the dependency of $S(\bar{\gamma})$ on $\bar{\gamma}$. To this end, it is important to exclude a special case. Recall from (2.1) that the local martingale parts of the financial assets are denoted by M^j for $j \in \{1, \ldots, d_1\}$ and define the (square-integrable) martingales M^j for $j \in \{d_1 + 1, \ldots, d_1 + d_2\}$ by $M_t^j := E[D^j | \mathcal{F}_t]$. We make the following assumption for the remainder of this section.

Assumption 4.10. There is $j \in \{1, \ldots, d_1 + d_2\}$ such that the local martingales M^j and $Z(\bar{\gamma}_0)$ are not strongly orthogonal.

If Assumption 4.10 fails to hold, we obtain a trivial case where $S = S(\bar{\gamma})$ is (componentwise) a local martingale that does not depend on the choice of $\bar{\gamma}$. In that case, (1, S) is a quadratic equilibrium for any parameters $\gamma_1, \ldots, \gamma_K \in \mathbb{R}$ such that $\sum_{k=1}^K \gamma_k > \bar{\gamma}_0$ and a mean-variance equilibrium for any $\lambda_1, \ldots, \lambda_K > 0$; see Martins [23, Lemma 2.27 and Corollary 2.28] for details. On the other hand, [23, Lemma 2.27] yields under Assumption 4.10 that for any value of $\bar{\gamma} > \bar{\gamma}_0$, $S^j(\bar{\gamma})$ is not a local martingale for at least one $j \in \{1, \ldots, d_1 + d_2\}$. The following result shows that in this case the map $\bar{\gamma} \mapsto S(\bar{\gamma})$ is injective.

Lemma 4.11. Suppose that Assumption 4.10 holds. Then for any $\bar{\gamma} > \bar{\gamma}_0$ and $\gamma_1, \ldots, \gamma_K \in \mathbb{R}$, $(1, S(\bar{\gamma}))$ is a quadratic equilibrium with respect to the risk tolerances $(\gamma_k)_{k=1}^K$ if and only if $\bar{\gamma} = \sum_{k=1}^K \gamma_k$. In particular, for $\bar{\gamma}' > \bar{\gamma}_0$, we have $S(\bar{\gamma}) = S(\bar{\gamma}')$ if and only if $\bar{\gamma} = \bar{\gamma}'$.

4.3 Existence of equilibria

As discussed after Lemma 4.7, we would like to use Theorem 3.7 (or Lemma 4.9) to find a mean-variance equilibrium (1, S). In order to do that, we need to determine $\bar{\gamma}$, which is given via Lemma 4.7 in terms of ℓ and $(c_k)_{k=1}^K$, which themselves depend on S. Although it is not clear a priori how to determine any of these quantities from the primitives, we can express the cyclical dependency (implied by (4.5), (4.6) and Lemmas 4.7 and 4.9) between $(\ell, (c_k)_{k=1}^K), \bar{\gamma}$ and the mean-variance equilibrium (1, S) as a fixed-point condition on $\bar{\gamma}$.

Proposition 4.12. Suppose that Assumptions 4.8 and 4.10 hold. For $\bar{\gamma} > \bar{\gamma}_0$, the market $(1, S(\bar{\gamma}))$ is a mean-variance equilibrium if and only if $\bar{\gamma} = \tilde{\gamma}(\bar{\gamma})$, where

$$\tilde{\gamma}(\bar{\gamma}) := \sum_{k=1}^{K} \left(c_k(\bar{\gamma}) + \frac{\lambda_k}{\ell(\bar{\gamma})} \right).$$
(4.11)

It now remains to solve (4.11) for $\bar{\gamma}$, which is not so easy due to the implicit dependence of ℓ and c_k on $\bar{\gamma}$. However, as it turns out, the probabilistic structure of the quadratic equilibrium leads to a surprisingly simple relationship between $\ell(\bar{\gamma})$ and $\bar{c}(\bar{\gamma}) := \sum_{k=1}^{K} c_k(\bar{\gamma})$. To show this, we use some results on the dynamic theory of mean-variance hedging; see Appendix A.2.

Proposition 4.13. For $\bar{\gamma} > \bar{\gamma}_0$, define $\bar{c}(\bar{\gamma}) := \sum_{k=1}^K c_k(\bar{\gamma})$. Then

$$\bar{\gamma} - E_P[\bar{\Xi}] = (\bar{\gamma} - \bar{c}(\bar{\gamma}))\ell(\bar{\gamma}). \tag{4.12}$$

Equation (4.12) gives us exactly what we need to find an explicit solution $\bar{\gamma}$ to the equation $\bar{\gamma} = \tilde{\gamma}(\bar{\gamma})$, where $\tilde{\gamma}$ is defined by (4.11), and thus to solve the problem of finding mean-variance equilibria of the form $(1, S(\bar{\gamma}))$.

Theorem 4.14. Suppose that Assumptions 4.8 and 4.10 are satisfied. Set

$$\bar{\gamma} := \sum_{k=1}^{K} \lambda_k + E_P[\bar{\Xi}]. \tag{4.13}$$

If $\bar{\gamma} > \bar{\gamma}_0$, then $(1, S(\bar{\gamma}))$ is the unique linear mean-variance equilibrium of the form $(1, S(\bar{\gamma}'))$ for some $\bar{\gamma}' > \bar{\gamma}_0$. If $\bar{\gamma} \leq \bar{\gamma}_0$, then there exists no mean-variance equilibrium of this form.

Note that the condition $\bar{\gamma} > \bar{\gamma}_0$ in Theorem 4.14 is equivalent to

$$\sum_{k=1}^{K} \lambda_k > \operatorname{ess\,sup}\bar{\Xi} - E_P[\bar{\Xi}]. \tag{4.14}$$

Thus there exists an equilibrium of the form given in Lemma 4.9 if and only if the aggregate risk tolerance $\sum_{k=1}^{K} \lambda_k$ is larger than the uncertainty of the aggregate endowment $\bar{\Xi}$, as measured by ess sup $\bar{\Xi} - E_P[\bar{\Xi}] \geq 0$. We note, however, that the condition (4.14) is not necessary for

the existence of a mean-variance equilibrium in the one-period model considered in Koch-Medina/Wenzelburger [17], and so the study of other types of mean-variance equilibria remains an open question for the general case. In particular, there may exist mean-variance equilibria that do not satisfy Assumption 4.3 nor Lemma 4.7, as well as mean-variance equilibria that are quadratic equilibria of the form $(1, S(\bar{\gamma}))$ for $\bar{\gamma} \leq \bar{\gamma}_0$, in which case Lemma 4.9 does not apply. Nevertheless, since the equilibria with $\bar{\gamma} > \bar{\gamma}_0$ are the most meaningful from an economic point of view, Theorem 4.14 still provides a satisfactory existence and uniqueness result.

A Key results on mean–variance hedging

A.1 Static theory

We give here some results on the MVH and exMVH problems (3.2) and (3.3). We omit for conciseness the (technical) proofs of Lemma A.1, Propositions A.3–A.5 and A.7 and Corollary A.6 below; they can be found in Martins [23, Section III.2.5]. On the other hand, the proof of Lemma A.8 is given below.

Lemma A.1. Let $H_1, H_2 \in L^2$ and $\lambda \in \mathbb{R}$.

- 1) Suppose there exist solutions $\vartheta^{\text{MVH}}(H_1), \vartheta^{\text{MVH}}(H_2) \in \overline{\Theta}(S)$ to (3.2) for H_1 and H_2 , respectively. Then $\vartheta^{\text{MVH}}(H_1) + \lambda \vartheta^{\text{MVH}}(H_2)$ is a solution to (3.2) for the payoff $H_1 + \lambda H_2$.
- 2) Suppose there exist solutions $(c(H_1), \vartheta^{\text{ex}}(H_1)), (c(H_2), \vartheta^{\text{ex}}(H_2)) \in \mathbb{R} \times \overline{\Theta}(S)$ to (3.3) for H_1 and H_2 , respectively. Then $(c(H_1) + \lambda c(H_2), \vartheta^{\text{ex}}(H_1) + \lambda \vartheta^{\text{ex}}(H_2))$ is a solution to (3.3) for the payoff $H_1 + \lambda H_2$.

Proof. See [23, Lemma III.1.6].

If S admits an equivalent local martingale measure (ELMM) with square-integrable density, then $\mathcal{G}_T(S)$ and $\mathbb{R} + \mathcal{G}_T(S)$ are closed and the solutions in $\overline{\Theta}(S)$ and $\mathbb{R} \times \overline{\Theta}(S)$ are unique; this follows by Černý/Kallsen [5, Lemma 2.11]. However, without that extra assumption, both closedness of $\mathcal{G}_T(S)$ and $\mathbb{R} + \mathcal{G}_T(S)$ as well as uniqueness of the solutions in $\overline{\Theta}(S)$ and $\mathbb{R} \times \overline{\Theta}(S)$ (if they exist) do not hold in general.

In order to deal with the uniqueness issue, it is useful to introduce the notions of uniqueness of gains processes and uniqueness of value processes associated with a price process S.

Definition A.2. Let (1, S) be a local L^2 -market. It is said to satisfy

- uniqueness of gains processes if for any two trading strategies $\vartheta^1, \vartheta^2 \in \overline{\Theta}(S)$, the equality $\vartheta^1 \cdot S_T = \vartheta^2 \cdot S_T P$ -a.s. implies that $\vartheta^1 =_S \vartheta^2$.
- uniqueness of value processes if for any two trading strategies $\vartheta^1, \vartheta^2 \in \overline{\Theta}(S)$ and initial values $c_1, c_2 \in \mathbb{R}$, the equality $c_1 + \vartheta^1 \cdot S_T = c_2 + \vartheta^2 \cdot S_T$ *P*-a.s. implies that $c_1 = c_2$ and $\vartheta^1 =_S \vartheta^2$.

We have the following two equivalent characterisations of uniqueness of gains and value processes. They follow immediately from the linear structure of the (extended) mean-variance hedging problems given in Lemma A.1, as well as the fact that for H = 0, the problems of MVH (3.2) and exMVH (3.3) admit as solutions $\vartheta = 0$ and $(c, \vartheta) = (0, 0)$, respectively.

Proposition A.3. Let (1, S) be a local L^2 -market. The following are equivalent:

(a) (1, S) satisfies uniqueness of gains processes.

- (b) For some $H \in L^2$, the MVH problem (3.2) admits a unique solution.
- (c) For each $H \in L^2$ for which the MVH problem (3.2) admits a solution, the solution is unique.

Proposition A.4. Let (1, S) be a local L^2 -market. The following are equivalent:

- (a) (1, S) satisfies uniqueness of value processes.
- (b) For some $H \in L^2$, the exMVH problem (3.3) admits a unique solution.
- (c) For each $H \in L^2$ for which the exMVH problem (3.3) admits a solution, the solution is unique.

Proof of Propositions A.3 and A.4. See [23, Propositions III.1.8 and III.1.9].

The following two results show that uniqueness of value processes implies uniqueness of gains processes and link the MVH problem (3.2) and the extended exMVH problem (3.3).

Proposition A.5. Let (1, S) be a local L^2 -market. The following statements hold:

- **1)** If (1, S) satisfies uniqueness of value processes, then (1, S) also satisfies uniqueness of gains processes.
- 2) Suppose that (1, S) satisfies uniqueness of gains processes and that the MVH problem (3.2) for H = 1 admits a solution $\vartheta^{\text{MVH}}(1)$. Then (1, S) satisfies uniqueness of value processes if and only if $E[(\vartheta^{\text{MVH}}(1) \cdot S_T 1)^2] > 0$.

Proof. 1) Since $\vartheta \equiv 0$ is a solution to (3.2) for H = 0 with hedging error 0, any solution to (3.2) for 0 is also a solution to (3.3) for 0 with c = 0. Thus the first assertion follows from Propositions A.3 and A.4, using the fact that if the MVH problem (3.2) for 0 does not have a unique solution, then a fortiori the exMVH problem (3.3) for 0 cannot have a unique solution.

2) We assume first that (1, S) satisfies uniqueness of value processes. Supposing for a contradiction that $E[(\vartheta^{\text{MVH}}(1) \bullet S_T - 1)^2] = 0$, then the exMVH problem (3.3) for 0 has two distinct solutions $(-1, \vartheta^{\text{MVH}}(1))$ and (0, 0), which contradicts Proposition A.4.

To prove the converse statement, suppose that $E[(\vartheta^{\text{MVH}}(1) \cdot S_T - 1)^2] > 0$. We claim that (0,0) is the unique solution to the exMVH problem (3.3) for H = 0. To show the claim, let (c,ϑ) be a solution to the exMVH problem (3.3) for $H \equiv 0$. Then by comparing the MVH (3.2) and exMVH (3.3) problems, we note that ϑ is also a solution to the MVH problem for $H \equiv -c$. The uniqueness of gains processes and Proposition A.3 yield that $\vartheta^{\text{MVH}}(1)$ and ϑ are the unique solutions to the MVH problem (3.2) for $H \equiv 1$ and $H \equiv -c$, respectively. In the case c = 0, we obtain in particular that $\vartheta = 0$ is the unique solution to the MVH problem for $H \equiv 0$, and hence the claim holds in this case since $(c, \vartheta) = (0, 0)$. Suppose now for a contradiction that $c \neq 0$. By the linearity of MVH (see Lemma A.1), we have $\vartheta = -c\vartheta^{\text{MVH}}(1)$. Since both (0,0) and (c, ϑ) are solutions to the exMVH problem for $H \equiv 0$, we deduce that

$$c + \vartheta \bullet S_T = c(1 - \vartheta^{\text{MVH}}(1) \bullet S_T) \equiv 0.$$

However, since $c \neq 0$, this contradicts the assumption so that this case cannot hold. Therefore, the only solution to the exMVH problem for $H \equiv 0$ is (0,0), which shows the claim. The conclusion that (1, S) satisfies uniqueness of value processes then follows by Proposition A.4.

Corollary A.6. Let (1, S) be a local L^2 -market satisfying uniqueness of value processes such that the MVH problem (3.2) for 1 has a unique solution $\vartheta^{\text{MVH}}(1)$. Then for $H \in L^2$, the MVH problem (3.2) for H has a unique solution $\vartheta^{\text{MVH}}(H)$ if and only if the exMVH problem (3.3) for H has a unique solution $(c(H), \vartheta^{\text{ex}}(H))$, in which case we have

$$c(H) = \frac{E[H(1 - \vartheta^{\text{MVH}}(1) \bullet S_T)]}{E[(1 - \vartheta^{\text{MVH}}(1) \bullet S_T)^2]}$$
(A.1)

and $\vartheta^{\text{ex}}(H) =_{S} \vartheta^{\text{MVH}}(H) - c(H)\vartheta^{\text{MVH}}(1).$

Proof. We first show the "if" statement. Since $\vartheta^{\text{ex}}(H)$ solves the MVH problem with payoff H - c(H), it follows by the linearity of MVH (see Lemma A.1) that $\vartheta := \vartheta^{\text{ex}}(H) + c(H)\vartheta^{\text{MVH}}(1)$ is a solution to the MVH problem (3.2) for H. The uniqueness of the solution is ensured by the uniqueness of value processes and Proposition A.3.

Next, we consider the "only if" statement. Suppose for a contradiction that the pair $(c(H), \vartheta^{\text{ex}}(H))$ given in the statement of the corollary is not a solution to the exMVH problem for H. Thus there exists a competitor (c, ϑ) that achieves a lower mean squared error. This competitor may be improved by letting ϑ be the solution to the MVH problem (3.2) for H - c, so that we may assume $\vartheta = \vartheta(H) - c\vartheta(1)$. Moreover, as $E[\vartheta^{\text{MVH}}(H) \cdot S_T(1 - \vartheta^{\text{MVH}}(1) \cdot S_T)] = 0$ by the first-order condition of MVH, the mean squared error attained by (c, ϑ) is given by

$$E\left[\left(c + \left(\vartheta^{\text{MVH}}(H) - c\vartheta^{\text{MVH}}(1)\right) \bullet S_T - H\right)^2\right]$$

= $c^2 E\left[\left(1 - \vartheta^{\text{MVH}}(1) \bullet S_T\right)^2\right] - 2c E\left[H\left(1 - \vartheta^{\text{MVH}}(1) \bullet S_T\right)\right] + E\left[\left(H - \vartheta^{\text{MVH}}(H) \bullet S_T\right)^2\right].$ (A.2)

As a quadratic function of c, the right-hand side of (A.2) has the unique minimiser c(H) given by (A.1), so that the error may be reduced further by setting c = c(H). But then we obtain $(c, \vartheta) = (c(H), \vartheta^{\text{ex}}(H))$, which leads to a contradiction. Therefore, the pair $(c(H), \vartheta^{\text{ex}}(H))$ given in the statement of the corollary is indeed a solution to the exMVH problem for H, and its uniqueness follows by Proposition A.4.

The following result gives simple sufficient conditions for uniqueness of gains and value processes and for the existence of solutions to the MVH and exMVH problems (3.2) and (3.3) in terms of a signed local martingale measure for S. The assumption that such a signed measure exists is not necessary for the existence of solutions to the MVH and exMVH problems, but it is a weaker assumption than the existence of an equivalent local martingale measure for S. By Černý/Czichowsky [4, Theorem 3.2], the conditions in part 2) imply the economic assumption of the so-called law of one price; see [4, Definition 2.3].

Proposition A.7. Let (1, S) be a local L^2 -market and $Z = (Z_t)_{0 \le t \le T}$ a square-integrable martingale such that ZS^j is a local martingale for all $j \in \{1, \ldots, d\}$. Then $Z(\vartheta \cdot S)$ is a *P*-martingale for each $\vartheta \in \overline{\Theta}(S)$. Moreover:

- 1) If $Z_t \neq 0$ P-a.s. for each $t \in [0,T]$, then (1,S) satisfies uniqueness of value processes.
- **2)** If $Z_t \neq 0$ and $Z_{t-} \neq 0$ for all $t \in [0,T]$ *P*-a.s., then the MVH problem (3.2) and exMVH problem (3.3) have unique solutions for each $H \in L^2$.

Proof. See [23, Proposition III.1.12].

We close this appendix by linking the zero solution of an MVH problem to a local martingaletype condition for S, which is used to prove some of the main results in Section 3.2.

Lemma A.8. Let (1, S) be a local L^2 -market and $H \in L^2$. Define the square-integrable martingale $Z = (Z_t)_{0 \le t \le T}$ by $Z_t := E[H | \mathcal{F}_t]$. The following are equivalent:

- (a) $0 \in \overline{\Theta}(S)$ solves the MVH problem (3.2) for H.
- (b) ZS^j is a local *P*-martingale for all $j \in \{1, \ldots, d\}$.

Proof. (a) \Rightarrow (b): As $0 \in \overline{\Theta}(S)$ is a solution to (3.2), we have for all $\vartheta \in \overline{\Theta}(S)$ and $\delta \in \mathbb{R}$ that

$$E[(H - \delta \vartheta \bullet S_T)^2] \ge E[H^2].$$

By taking $\delta \searrow 0$ and $\delta \nearrow 0$, we deduce by examining the first-order term that $E[(\vartheta \cdot S_T)H] = 0$ for any $\vartheta \in \overline{\Theta}(S)$, i.e., H is orthogonal to $\mathcal{G}_T(S) \subseteq L^2$.

Now fix $j \in \{1, \ldots, d\}$ and let σ be a stopping time. Since (1, S) is a local L^2 -market, there exists a localising sequence $(\tau_n)_{n \in \mathbb{N}}$ such that for each $n \in \mathbb{N}$, the stopped process S^{j,τ_n} is an L^2 -semimartingale. Fix $n \in \mathbb{N}$. Then the strategy $e^{j,\sigma,n} := (0, \ldots, 0, \mathbb{1}_{]]0,\sigma\wedge\tau_n], 0, \ldots, 0)$, where the indicator process is at the *j*-th position, belongs to $\Theta_{\text{simple}}(S) \subseteq \overline{\Theta}(S)$. The fact that *Z* is a *P*-martingale with $Z_T = H$ and $E[(e^{j,\sigma,n} \bullet S_T)H] = 0$ yield

$$E[S^j_{\sigma\wedge\tau_n}Z_{\sigma\wedge\tau_n} - S^j_0Z_0] = E[(S^j_{\sigma\wedge\tau_n} - S^j_0)Z_T] = E[(e^{j,\sigma,n} \bullet S_T)H] = 0.$$

Since σ was arbitrary, it follows that $(ZS^j)^{\tau_n}$ is a *P*-martingale. As (τ_n) is a localising sequence, we conclude that ZS^j is a local *P*-martingale.

(b) \Rightarrow (a): Fix $\vartheta \in \overline{\Theta}(S)$. By Proposition A.7, $Z(\vartheta \bullet S)$ is a *P*-martingale, and hence

$$E[(\vartheta \bullet S_T - H)^2] = E[H^2] - 2E[(\vartheta \bullet S_T)Z_T] + E[(\vartheta \bullet S_T)^2]$$

= $E[H^2] + E[(\vartheta \bullet S_T)^2]$
 $\geq E[(H - 0 \bullet S_T)^2].$

Thus $0 \in \overline{\Theta}(S)$ solves the MVH problem (3.2).

A.2 Dynamic theory

We introduce here some key concepts from the dynamic theory of mean-variance hedging, for which we use Černý/Kallsen [5] as a reference, that are used later in the proof of Proposition 4.13. For the remainder of this section, we once again impose Assumption 4.8. Hence [5, Assumption 2.1] is satisfied, since Lemma 4.9 gives an equivalent local martingale measure $Q(\bar{\gamma})$ for $S(\bar{\gamma})$ with bounded density.

As in [5, Definition 3.3], we introduce the *opportunity process* $L(\bar{\gamma})$ as the value process of the pure investment problem, which is given by

$$L_t(\bar{\gamma}) = \operatorname*{ess inf}_{\vartheta \in \overline{\Theta}_{t,T}(S(\bar{\gamma}))} E\left[\left(1 - \left(\vartheta \bullet S(\bar{\gamma}) \right)_T \right)^2 \middle| \mathcal{F}_t \right], \quad 0 \le t \le T,$$
(A.3)

where $\overline{\Theta}_{t,T}(S(\bar{\gamma})) \subseteq \overline{\Theta}(S(\bar{\gamma}))$ is the set of admissible strategies ϑ such that $\vartheta \mathbb{1}_{[0,t]} = 0$. We have by [5, Corollary 3.4 and Lemma 3.10] that $L(\bar{\gamma})$ is an (0,1]-valued submartingale with $L_T(\bar{\gamma}) = 1$. Note that $L_0(\bar{\gamma})$ coincides with $\ell(\bar{\gamma})$ given in (4.5). By [5, Lemma 3.1], there exists for each $t \in [0,T]$ a unique optimal strategy $\vartheta^{(t)}(1; S(\bar{\gamma})) \in \overline{\Theta}_{t,T}(S(\bar{\gamma}))$ to (A.3); we say that it is the optimal pure investment strategy started at time t.

Next, we introduce the mean value process $(\bar{V}_t(\bar{\gamma}))_{0 \leq t \leq T}$ for $\bar{H}(\bar{\gamma}) = \bar{\gamma} - \bar{\Xi}$ in the sense of [5, Definition 4.2]. By [5, Lemmas 3.7 and 4.1 and Proposition 3.13.1], $\bar{V}(\bar{\gamma})$ is the unique semimartingale such that

$$\bar{V}_t(\bar{\gamma}) = \frac{1}{L_t(\bar{\gamma})} E\bigg[\bar{H}(\bar{\gamma}) \bigg(1 - \big(\vartheta^{(t)}\big(1; S(\bar{\gamma})\big) \bullet S(\bar{\gamma})\big)_T\bigg) \bigg| \mathcal{F}_t\bigg], \quad 0 \le t \le T.$$

In particular, $\bar{V}_T(\bar{\gamma}) = \bar{H}(\bar{\gamma})$. By [5, Lemma 4.1], the process $(\bar{V}_s(\bar{\gamma})M_s^{(t)}(\bar{\gamma}))_{t\leq s\leq T}$ is a *P*-martingale on [t,T] for any $t \in [0,T]$, where $(M_s^{(t)}(\bar{\gamma}))_{t\leq s\leq T}$ is the *P*-martingale (see [5, Lemma 3.2]) defined on [t,T] by

$$M_s^{(t)}(\bar{\gamma}) := L_s(\bar{\gamma}) \left(1 - \left(\vartheta^{(t)}(1; S(\bar{\gamma})) \bullet S(\bar{\gamma}) \right)_s \right), \quad 0 \le t \le s \le T.$$
(A.4)

The key property for our purposes is that $\bar{V}_t(\bar{\gamma})$ satisfies the inequality

$$\underset{\vartheta \in \overline{\Theta}_{t,T}(S(\bar{\gamma}))}{\operatorname{ess inf}} E\left[\left(\bar{H}(\bar{\gamma}) - \bar{V}_{t}(\bar{\gamma}) - \left(\vartheta \cdot S(\bar{\gamma})\right)_{T}\right)^{2} \middle| \mathcal{F}_{t}\right] \\ \leq \underset{\vartheta \in \overline{\Theta}_{t,T}(S(\bar{\gamma}))}{\operatorname{ess inf}} E\left[\left(\bar{H}(\bar{\gamma}) - U - \left(\vartheta \cdot S(\bar{\gamma})\right)_{T}\right)^{2} \middle| \mathcal{F}_{t}\right]$$

for any \mathcal{F}_t -measurable random variable U; this follows by [5, Theorem 4.10.2]. In particular, $\bar{V}_0(\bar{\gamma})$ is the first component of the solution to the exMVH problem (3.3) for $\bar{H}(\bar{\gamma})$, and hence $\bar{V}_0(\bar{\gamma}) = c(\bar{H}(\bar{\gamma}); S(\bar{\gamma})) = \bar{\gamma} - c(\bar{\Xi}; S(\bar{\gamma})).$

B Auxiliary results and proofs

Proof of Lemma 3.1. Let $\vartheta \in \overline{\Theta}(S)$ and set $\tilde{\vartheta} := \vartheta - \eta^k \in \overline{\Theta}(S)$. Rewriting $U_k^{\mathbb{Q}}(x) = -(x - \gamma_k)^2 + \gamma_k^2$ yields

$$E[U_k^{\mathbf{Q}}((\vartheta - \eta^k) \bullet S_T + \Xi^k)] = E[U_k^{\mathbf{Q}}(\tilde{\vartheta} \bullet S_T + \Xi^k)] = -E[(\tilde{\vartheta} \bullet S_T - H^k)^2] + \gamma_k^2$$

Because $\overline{\Theta}(S)$ is a vector space, this shows that ϑ is a solution to the maximisation problem (3.1) if and only if $\tilde{\vartheta}$ is a solution to the MVH problem (3.4), and therefore the two problems are equivalent under the assumption that $\eta^k \in \overline{\Theta}(S)$. In particular, (3.1) has a unique solution $\hat{\vartheta}^k$ if and only if (3.4) has a unique solution $\vartheta^{\text{MVH}}(H^k)$, in which case we have the relationship $\hat{\vartheta}^k =_S \eta^k + \vartheta^{\text{MVH}}(H^k)$ between the solutions.

Proof of Proposition 3.2. Since $\vartheta^{\text{MVH}}(1)$ is the unique solution to (3.6), we have by Proposition A.3 that (1, S) satisfies uniqueness of gains processes. Thus the assumption that $E[(\vartheta^{\text{MVH}}(1) \cdot S_T - 1)^2] > 0$ implies by part 2) of Proposition A.5 that (1, S) also satisfies uniqueness of value processes; see Definition A.2. Then by Corollary A.6, there exists a unique solution $\vartheta^{\text{MVH}}(\Xi^k) \in \overline{\Theta}(S)$ to the MVH problem (3.4) for Ξ^k if and only if there exists a unique solution to the exMVH problem for Ξ^k , and we have

$$\vartheta^{\text{MVH}}(\Xi^k) = c_k \vartheta^{\text{MVH}}(1) + \vartheta^{\text{ex}}(\Xi^k).$$
(B.1)

Moreover, by the linearity of MVH (see Lemma A.1), there exists a unique solution to (3.4) for Ξ^k if and only if there exists a unique solution to (3.4) for $H^k = \gamma_k - \Xi^k$, in which case we have

$$\vartheta^{\mathrm{MVH}}(H^k) = \gamma_k \vartheta^{\mathrm{MVH}}(1) - \vartheta^{\mathrm{MVH}}(\Xi^k).$$

Together with (B.1) and Lemma 3.1, this shows the equivalence and gives (3.7).

Proof of Lemma 3.5. By the implication (b) \Rightarrow (c) in Proposition A.3, the map $H \mapsto \vartheta^{\text{MVH}}(H)$ is well defined for all H such that a solution $\vartheta^{\text{MVH}}(H)$ to (3.2) exists, since such a solution is unique up to S-equivalence. We also have by Lemma A.1 that $H \mapsto \vartheta^{\text{MVH}}(H)$ is linear where it is defined. Because $\eta^1, \ldots, \eta^K \in \overline{\Theta}(S)$, we get from Lemma 3.1 that the MVH problem (3.2)

for H^k has the unique solution $\vartheta^{\text{MVH}}(H^k) = \hat{\vartheta}^k - \eta^k$ for each $k \in \{1, \dots, K\}$. Hence there is a unique solution to (3.2) for $\bar{H} = \sum_{k=1}^{K} H^k$, which is given by

$$\vartheta^{\text{MVH}}(\bar{H}) =_S \sum_{k=1}^K \left(\vartheta^{\text{MVH}}(H^k) - \eta^k \right) = \bar{\vartheta} - \bar{\eta}.$$
(B.2)

Thus the claim follows by Lemmas 3.3 and 3.1, as we have

$$\bar{\vartheta} =_{S} \vartheta^{\text{MVH}}(\bar{H}) + \bar{\eta} =_{S} \sum_{k=1}^{K} \left(\vartheta^{\text{MVH}}(H^{k}) + \eta^{k} \right) =_{S} \sum_{k=1}^{K} \hat{\vartheta}^{k},$$

which shows (3.12) and concludes the proof.

Proof of Lemma 3.6. Denote by $\hat{\vartheta}^1, \ldots, \hat{\vartheta}^K \in \overline{\Theta}(S)$ the unique individually optimal strategies. Then by Lemma 3.5, $\bar{\vartheta} := \sum_{k=1}^K \hat{\vartheta}^k \in \overline{\Theta}(S)$ is the unique solution to the optimisation problem (3.8) of the representative agent. Moreover, the market clearing condition (2.10) yields $\bar{\vartheta} =_S \bar{\eta}$, so that by Lemma 3.3, 0 is the unique solution to the MVH problem (3.2) for \bar{H} . Thus Lemma A.8 yields that $(\bar{Z}_t S_t^j)_{0 \le t \le T}$ is a local *P*-martingale for each $j \in \{1, \ldots, d_1 + d_2\}$, as claimed. \Box

Remark B.1. The choice of $\bar{\xi}^{(1)}$ in (3.13) is only unique up to $M^{(1)}$ -equivalence. Because the components of $M^{(1)}$ may be linearly dependent, the components $\bar{\xi}^i \cdot M^i$ need not be well defined in general (see Cherny/Shiryaev [7]), but we choose a particular integrand $\bar{\xi}^{(1)} = (\bar{\xi}^1, \ldots, \bar{\xi}^{d_1})$ with $\bar{\xi}^i \in L^2(M^i)$ for $i \in \{1, \ldots, d_1\}$ as follows: Applying the Gram–Schmidt algorithm to $(M^1, \ldots, M^{d_1}, \bar{Z}) \in \mathcal{M}^2_{0,\text{loc}}$ yields a unique decomposition of the form

$$\bar{Z}_t = \bar{Z}_0 + \sum_{i=1}^{d_1} \bar{\xi}^i \cdot M_t^i + M_t^{\bar{Z}}, \quad 0 \le t \le T,$$

where $\sum_{i=1}^{I} \bar{\xi}^{i} \cdot M^{i}$ is strongly orthogonal to $\sum_{i=I+1}^{d_{1}} \bar{\xi}^{i} \cdot M^{i} + M^{\bar{Z}}$ for each $I \in \{1, \ldots, d_{1}\}$. This orthogonality property and the square-integrability of \bar{Z} yield that $\sum_{i=1}^{I} \bar{\xi}^{i} \cdot M^{i}$ is a square-integrable martingale for each I, and hence so is $\bar{\xi}^{i} \cdot M^{i}$. For this choice of $\bar{\xi}^{(1)} := (\bar{\xi}^{1}, \ldots, \bar{\xi}^{d_{1}})$ and as $M^{(1)}$ is a locally square-integrable martingale by assumption, the predictable quadratic variation

$$\langle \bar{\xi}^i \bullet M^i, M^j \rangle = \bar{\xi}^i {}^{\top} \bullet \langle M^i, M^j \rangle$$

is well defined for each $j \in \{1, \ldots, d_1\}$ and $\overline{\xi^i}^{\top}([M^i, M^j] - \langle M^i, M^j \rangle)$ is a local *P*-martingale. In particular, the process

$$\langle \bar{\xi}^{(1)} \bullet M^{(1)}, M^j \rangle = \sum_{i=1}^{d_1} \bar{\xi}^i \bullet \langle M^i, M^j \rangle$$
(B.3)

is well defined. Although we choose $\bar{\xi}^{(1)}$ as above, note that the martingale $\bar{\xi}^{(1)} \cdot M^{(1)}$ is independent of that choice due to (3.13). Hence the right-hand side of (B.3) is also independent of any choice of $\bar{\xi}^{(1)}$ such that the individual summands are well defined. Likewise, the finitevariation part in (3.14) below also does not depend on the choice of $\bar{\xi}^{(1)}$.

Proof of Theorem 3.7. (a) We show that a quadratic equilibrium (1, S) must be given by (3.14) and (3.15). Lemma 3.6 yields for each $j \in \{1, \ldots, d_1 + d_2\}$ that $\overline{Z}S^j$ is a local *P*-martingale.

We first consider $j \in \{1, \ldots, d_1\}$. Recall the decomposition (3.13) for \overline{Z} and the dynamics (2.1) for S^j . Applying the product formula to $\overline{Z}S^j$ and rearranging terms, we obtain

$$\bar{Z}_t S_t^j - \bar{Z}_- \bullet M_t^j - S_-^j \bullet \bar{Z}_t - \sum_{i=1}^{d_1} \bar{\xi}^i \bullet \left([M^i, M^j] - \langle M^i, M^j \rangle \right)_t$$
$$= \bar{Z}_- \bullet A_t^j + \sum_{i=1}^{d_1} \bar{\xi}^i \bullet \langle M^i, M^j \rangle_t \tag{B.4}$$

for $0 \leq t \leq T$. Note that $\bar{Z}S^j$, M^j , \bar{Z} and $\bar{\xi}^i \cdot ([M^i, M^j] - \langle M^i, M^j \rangle)$ are local *P*-martingales (for the latter, this is shown in Remark B.1), whereas A^j and $\bar{\xi}^i \cdot \langle M^i, M^j \rangle$ are predictable finitevariation processes. Thus both sides of (B.4) must vanish, as they are null at 0. By assumption, we have $\bar{Z}_t \neq 0$ and $\bar{Z}_{t-} \neq 0$ for all $t \in [0, T]$ *P*-a.s. Since \bar{Z} is also càdlàg, this implies that $1/\bar{Z}_-$ is finite-valued and càglàd, thus locally bounded. Integrating $1/\bar{Z}_-$ against the right-hand side of (B.4), which vanishes as we have shown, yields $A^j = -\sum_{i=1}^{d_1} \frac{\bar{\xi}^i}{\bar{Z}_-} \cdot \langle M^i, M^j \rangle$. Plugging into (2.1) then shows (3.14).

Next, consider $j \in \{d_1 + 1, \ldots, d_1 + d_2\}$. By (2.2) and as $\overline{Z}_T = \overline{H}$, we have $\overline{Z}_T S_T^j = \overline{H} D^j$. Since $e^j \in \overline{\Theta}(S)$ and \overline{Z} is a square-integrable martingale, it follows from Proposition A.7 that $\overline{Z}S^j = \overline{Z}S_0^j + \overline{Z}(e^j \cdot S)$ is a *P*-martingale, so that $\overline{Z}_t S_t^j = E[\overline{H}D^j | \mathcal{F}_t]$. Since $\overline{Z}_t \neq 0$ *P*-a.s., this yields (3.15). We have thus shown that any equilibrium (1, S) must satisfy (3.14) and (3.15).

(b) Next, we show the converse statement. Define $(1, S) = (1, S^{(1)}, S^{(2)})$ by (3.14) and (3.15)and assume that $S^{(2)}$ is a local L^2 -semimartingale and $e^j \in \overline{\Theta}(S)$ for $j \in \{d_1 + 1, \ldots, d_1 + d_2\}$. We claim that (1, S) is a quadratic equilibrium. It is clear from (3.14) and (3.15) that $S^{(1)}$ and $S^{(2)}$ satisfy (2.1) and (2.2), respectively. Note that $S^{(1)}$ is a special semimartingale and the local martingale part $M^{(1)}$ is locally square-integrable, by assumption. Thus by Černý/Kallsen [5, Lemma A.2], $S^{(1)}$ is also a local L^2 -semimartingale so that $(1, S^{(1)}, S^{(2)})$ is a local L^2 -market. Next, we want to show that $\overline{Z}S^j$ is a local P-martingale for $j \in \{1, \ldots, d_1 + d_2\}$. This is clear for $j \in \{d_1 + 1, \ldots, d_1 + d_2\}$ by the construction (3.15). For $j \in \{1, \ldots, d_1\}$, we use a result on local \mathcal{E} -martingales by Choulli et al. [8] as in the proof of part 2) of Proposition A.7. Indeed, the assumptions on \overline{Z} yield $\overline{Z} = \overline{Z}_0 \mathcal{E}(\overline{N})$ for some local P-martingale $\overline{N} = (\overline{N}_t)_{0 \le t \le T}$, namely, $\overline{N} = (1/\overline{Z}_-) \cdot \overline{Z}$. Since for $j \in \{1, \ldots, d_1\}$, we have

$$\sum_{i=1}^{d_1} \frac{\bar{\xi}_t^i}{\bar{Z}_{t-}} \mathrm{d}\langle M^i, M^j \rangle_t = \frac{1}{\bar{Z}_{t-}} d\langle \bar{Z}, M^j \rangle_t = d\langle \bar{N}, M^j \rangle_t,$$

we obtain that S^{j} given by (3.14) is a local \mathcal{E} -martingale by [8, Corollary 3.16] (which generalises Girsanov's theorem to local \mathcal{E} -martingales). Thus by [8, Definition 3.11] with n = 0, $\overline{Z}S^{j}$ is a local P-martingale.

Now note that $\overline{Z}_t \neq 0$ and $\overline{Z}_{t-} \neq 0$ for all $t \in [0,T]$ *P*-a.s. by the assumptions on \overline{Z} , and $\overline{Z}S^j$ is a local *P*-martingale for $j \in \{1, \ldots, d_1 + d_2\}$ as shown above. Hence for each agent $k \in \{1, \ldots, K\}$, the MVH problem (3.2) for H^k has a unique solution $\vartheta(H^k)$ by part 2) of Proposition A.7. Since moreover $\eta^k \in \overline{\Theta}(S)$ by the assumption on $S^{(2)}$, it follows by Lemma 3.1 that the individual optimisation problem (3.1) for agent k has a unique solution ϑ^k . This shows condition 1) in Definition 2.6 of an equilibrium market. Moreover, the strategy 0 solves the MVH problem (3.2) for \overline{H} by Lemma A.8. Thus Lemmas 3.3 and 3.5 yield $\sum_{k=1}^{K} \vartheta^k = \overline{\vartheta} = \overline{\eta}$, i.e., the market clears and condition 2) is satisfied. Finally, condition 3) is satisfied by assumption, and thus (1, S) is a quadratic equilibrium.

Proof of Lemma 3.8. (a) This is (c) for $p_1 = 1$ and $p_2 = 2$, so that $q_1 = \infty$ and $q_2 = 2$.

(b) This is (c) for $p_1 = p_2 = \infty$.

(c) We only consider the case that $p_1, p_2 \in (1, \infty)$. The arguments for the other cases are very similar and therefore omitted. Fix $j \in \{d_1 + 1, \ldots, d_1 + d_2\}$ and define $\bar{Q} \approx P$ by

$$\frac{\mathrm{d}\bar{Q}}{\mathrm{d}P} = \frac{\bar{H}}{E[\bar{H}]} =: Z_T^{\bar{Q}} \in L^{1-p_1}(P) \cap L^{p_2}(P).$$

By (3.15) and the Bayes rule, S^j is a (true) \bar{Q} -martingale with $S_T^j = D^j$. Thus the inequalities of Hölder and Doob (with constant C_{q_1}) give

$$E_{P}\left[\sup_{t\in[0,T]}|S_{t}^{j}|^{2}\right] = E_{\bar{Q}}\left[\frac{1}{Z_{T}^{\bar{Q}}}\sup_{t\in[0,T]}|S_{t}^{j}|^{2}\right] \leq E_{\bar{Q}}\left[\left(\frac{1}{Z_{T}^{\bar{Q}}}\right)^{p_{1}}\right]^{1/p_{1}}E_{\bar{Q}}\left[\sup_{t\in[0,T]}|S_{t}^{j}|^{2q_{1}}\right]^{1/q_{1}}$$

$$\leq E_{P}\left[\left(\frac{1}{Z_{T}^{\bar{Q}}}\right)^{p_{1}-1}\right]^{1/p_{1}}C_{q_{1}}E_{\bar{Q}}\left[(D^{j})^{2q_{1}}\right]^{1/q_{1}}$$

$$= C_{q_{1}}E_{P}\left[(Z_{T}^{\bar{Q}})^{1-p_{1}}\right]^{1/p_{1}}E_{P}\left[Z_{T}^{\bar{Q}}(D^{j})^{2q_{1}}\right]^{1/q_{1}}$$

$$\leq C_{q_{1}}E_{P}\left[(Z_{T}^{\bar{Q}})^{1-p_{1}}\right]^{1/p_{1}}E_{P}\left[\left(Z_{T}^{\bar{Q}}\right)^{p_{2}}\right]^{1/(q_{1}p_{2})}E_{P}\left[(D^{j})^{2q_{1}q_{2}}\right]^{1/(q_{1}q_{2})}$$

$$<\infty$$

by the assumptions. This implies that S^j is an L^2 -semimartingale.

Proof of Lemma 3.10. Assume that a quadratic equilibrium (1, S) exists. For a contradiction, suppose that (3.16) is not satisfied, i.e., there exist $i \in \{1, \ldots, d_1\}$ and $t \in \{1, \ldots, T\}$ such that

$$P[\bar{Z}_{t-1} = 0, \,\bar{\xi}_t^i \Delta \langle M^i \rangle_t \neq 0] > 0. \tag{B.5}$$

By Lemma 3.6, $(\bar{Z}_t S_t)_{t=0,...,T}$ is a local *P*-martingale so that $\mathbb{1}_{\{\bar{Z}_{t-1}=0\}}\Delta(\bar{Z}S^i)_t$ is the increment of a local martingale. We decompose

$$\begin{split} \mathbb{1}_{\{\bar{Z}_{t-1}=0\}} \Delta(\bar{Z}S^{i})_{t} &= \mathbb{1}_{\{\bar{Z}_{t-1}=0\}} (\bar{Z}_{t-1} \Delta S^{i}_{t} + S^{i}_{t-1} \Delta \bar{Z}_{t} + \Delta \bar{Z}_{t} \Delta S^{i}_{t}) \\ &= \mathbb{1}_{\{\bar{Z}_{t-1}=0\}} S^{i}_{t-1} \Delta \bar{Z}_{t} + \mathbb{1}_{\{\bar{Z}_{t-1}=0\}} \Delta \bar{Z}_{t} \Delta S^{i}_{t} \\ &= \mathbb{1}_{\{\bar{Z}_{t-1}=0\}} S^{i}_{t-1} \Delta \bar{Z}_{t} + \mathbb{1}_{\{\bar{Z}_{t-1}=0\}} (\Delta [\bar{Z}, S^{i}]_{t} - \Delta \langle \bar{Z}, S^{i} \rangle_{t}) \\ &+ \mathbb{1}_{\{\bar{Z}_{t-1}=0\}} \bar{\xi}^{i}_{t} \Delta \langle M^{i} \rangle_{t}, \end{split}$$

where the last equality follows since $\Delta \langle \bar{Z}, S^i \rangle_t = \bar{\xi}_t^i \Delta \langle M^i \rangle_t$. Like the left-hand side, the first two terms in the last expression of the right-hand side are increments of local martingales, since \bar{Z} is a martingale and $[\bar{Z}, S^i] - \langle \bar{Z}, S^i \rangle$ a local martingale. It follows that the last term $\mathbb{1}_{\{\bar{Z}_{t-1}=0\}} \bar{\xi}_t^i \Delta \langle M^i \rangle_t$ must also be the increment of a local martingale. However, this term is also \mathcal{F}_{t-1} -measurable, and hence null *P*-a.s. This leads to a contradiction with (B.5) so that (3.16) must hold.

Similarly, suppose that (3.17) does not hold, i.e.,

$$P\left[\bar{Z}_t = 0, E[\bar{H}D^j \mid \mathcal{F}_t] \neq 0\right] > 0 \tag{B.6}$$

for some $j \in \{d_1 + 1, \ldots, d_1 + d_2\}$ and $t \in \{0, \ldots, T - 1\}$. Since $\overline{Z}S^j$ is a local *P*-martingale and $1 \in \overline{\Theta}(S^j)$ by condition 3) of Definition 2.6, Proposition A.7 yields that $\overline{Z}S^j$ is a true *P*-martingale. In particular, we have

$$\bar{Z}_t S_t^j = E[\bar{H}D^j \mid \mathcal{F}_t] \quad P ext{-a.s.}$$

This contradicts (B.6), and therefore (3.17) must hold.

24

Proof of Theorem 3.11. We show in the steps (a)–(d) below that the process (1, S) defined by (3.20) and (3.21) is a quadratic equilibrium. In step (a), we check the conditions required by Definition 2.6 except for 1) and 2). In steps (b) and (c), we show that $({}^{s}\mathcal{E}(\bar{N})_{t}S_{t}^{j})_{t \in \{s,...,T\}}$ and $(\bar{Z}_{t}S_{t}^{j})_{t \in \{0,...,T\}}$, respectively, are local *P*-martingales for each $s \in \{0,...,T-1\}$ and $j \in \{1,...,d_{1}+d_{2}\}$. These results are then used in step (d) to check that conditions 1) and 2) of Definition 2.6 are satisfied.

(a) We start by checking (2.1) and (2.2). For $j \in \{1, \ldots, d_1\}$, the process $(A_t^j)_{t \in \{0, \ldots, T\}}$ of S^j given by

$$A_{t}^{j} := \sum_{k=1}^{t} \sum_{i=1}^{d_{1}} \left(-\frac{\bar{\xi}_{k}^{i}}{\bar{Z}_{k-1}} \mathbb{1}_{\{\bar{Z}_{k-1}\neq 0\}} \Delta \langle M^{i}, M^{j} \rangle_{k} \right), \tag{B.7}$$

is predictable, so that by the definition (3.20), $S^{(1)}$ satisfies (2.1). Moreover, plugging t = T into (3.21) yields (2.2) since ${}^{T}\mathcal{E}(\bar{N})_{T} = 1$.

We also have to check that (1, S) is a local L^2 -market. As argued in the proof of Theorem 3.7, $S^{(1)}$ is a local L^2 -semimartingale as $M^{(1)}$ is locally square-integrable and by Černý/Kallsen [5, Lemma A.2]. On the other hand, $S^{(2)}$ is an L^2 -semimartingale as it is square-integrable by assumption and the set $\{0, \ldots, T\}$ of times is finite. The fact that $S^{(2)}$ is an L^2 -semimartingale also implies that condition 3) of Definition 2.6 of an equilibrium market is satisfied.

(b) We first show that $({}^{s}\mathcal{E}(\bar{N})_{t}S_{t}^{j})_{t\in\{s,...,T\}}$ is a local *P*-martingale for each $j \in \{1,...,d_{1}\}$ and $s \in \{0,...,T-1\}$. We use a similar argument as in the proof of part 2) of Proposition A.7. Consider the setup of Choulli et al. [8, Section 3] with the family $\mathcal{E} = ({}^{s}\mathcal{E}(\bar{N}))_{s\in\{0,...,T\}}$. Since for $j \in \{1,...,d_{1}\}$, we have

$$\sum_{i=1}^{d_1} \frac{\bar{\xi}_t^i}{\bar{Z}_{t-1}} \mathbb{1}_{\{\bar{Z}_{t-1}\neq 0\}} \Delta \langle M^i, M^j \rangle_t = \frac{\mathbb{1}_{\{\bar{Z}_{t-1}\neq 0\}}}{\bar{Z}_{t-1}} \Delta \langle \bar{Z}, M^j \rangle_t = \Delta \langle \bar{N}, M^j \rangle_t,$$

we obtain that S^j given by (3.20) is a local \mathcal{E} -martingale by [8, Corollary 3.16], i.e., ${}^s \mathcal{E}(\bar{N})S^j$ is a local P-martingale for each $s \in \{0, \ldots, T\}$.

For $j \in \{d_1 + 1, \ldots, d_1 + d_2\}$, we use the definition (3.21) and the square-integrability of ${}^s \mathcal{E}(\bar{N})$ and D^j to obtain for $t \in \{s, \ldots, T\}$ that

$${}^{s}\mathcal{E}(\bar{N})_{t}S_{t}^{j} = {}^{s}\mathcal{E}(\bar{N})_{t}E[{}^{t}\mathcal{E}(\bar{N})_{T}D^{j} \mid \mathcal{F}_{t}] = E[{}^{s}\mathcal{E}(\bar{N})_{T}D^{j} \mid \mathcal{F}_{t}],$$

and hence ${}^{s}\mathcal{E}(\bar{N})S^{j}$ is a true *P*-martingale for each $s \in \{0, \ldots, T\}$.

(c) Next, we show that $\overline{Z}S^j$ is a local *P*-martingale for $j \in \{1, \ldots, d_1 + d_2\}$. For $j \in \{1, \ldots, d_1\}$ and $t \in \{1, \ldots, T\}$, we have by (3.18) that

$$\begin{aligned} \Delta(\bar{Z}S^{j})_{t} &= \bar{Z}_{t-1}\Delta S_{t}^{j} + S_{t}^{j}\Delta\bar{Z}_{t} \\ &= \bar{Z}_{t-1}\Delta S_{t}^{j} + S_{t}^{j} \left(\mathbb{1}_{\{\bar{Z}_{t-1}\neq 0\}}\bar{Z}_{t-1}\Delta\bar{N}_{t} + \mathbb{1}_{\{\bar{Z}_{t-1}=0\}}\Delta\bar{Z}_{t}\right) \\ &= \mathbb{1}_{\{\bar{Z}_{t-1}\neq 0\}}\bar{Z}_{t-1}(\Delta S_{t}^{j} + S_{t}^{j}\Delta\bar{N}_{t}) + \mathbb{1}_{\{\bar{Z}_{t-1}=0\}}S_{t}^{j}\Delta\bar{Z}_{t}, \end{aligned}$$
(B.8)

where we use $\bar{Z}_{t-1} = \mathbb{1}_{\{\bar{Z}_{t-1}\neq 0\}}\bar{Z}_{t-1}$ for the last equality. Note that we have $t^{t-1}\mathcal{E}(\bar{N})_{t-1} = 1$ and

$$\Delta \left({}^{t-1}\mathcal{E}(\bar{N}) \right)_t = {}^{t-1}\mathcal{E}(\bar{N})_{t-1} \Delta \bar{N}_t = \Delta \bar{N}_t.$$

By plugging in, this yields

$$\Delta \left({}^{t-1}\mathcal{E}(\bar{N})S^j \right)_t = \Delta S^j_t + S^j_{t-1}\Delta \bar{N}_t + \Delta S^j_t \Delta \bar{N}_t = \Delta S^j_t + S^j_t \Delta \bar{N}_t.$$

Since we have already shown in step (b) that ${}^{s}\mathcal{E}(\bar{N})S^{j}$ is a local *P*-martingale for each $s \in \{0, \ldots, T\}$, this implies that $\Delta S_{t}^{j} + S_{t}^{j}\Delta \bar{N}_{t}$ is the increment of a local *P*-martingale, and hence so is the first

term in the right-hand side of (B.8). We now consider the second term. Since $\Delta \bar{Z}_t = \Delta M_t^Z$ on $\{\bar{Z}_{t-1} = 0\}$ by the assumption (3.16), we get

$$\mathbb{1}_{\{\bar{Z}_{t-1}=0\}}S_t^j\Delta\bar{Z}_t = \mathbb{1}_{\{\bar{Z}_{t-1}=0\}}(S_{t-1}^j + \Delta A_t^j + \Delta M_t^j)\Delta M_t^{\bar{Z}}.$$

This is the increment of a local P-martingale, as $M^{\bar{Z}}$ and M^{j} are strongly orthogonal local Pmartingales, whereas $S_{t-1}^{j} + \Delta A_{t}^{j}$ is \mathcal{F}_{t-1} -measurable. Returning to (B.8), we have thus shown that $\bar{Z}S^{j}$ is a local P-martingale for $j \in \{1, \ldots, d_1\}$.

On the other hand, for $j \in \{d_1 + 1, \ldots, d_1 + d_2\}$, we claim that $\overline{Z}S^j$ is even a true *P*martingale. We use backward induction to show this statement, starting with t = T. Since $\overline{Z}_T = \overline{H}$ and $S_T^j = D^j$ are square-integrable, we get that $\overline{Z}_T S_T^j$ is integrable so that $\overline{Z}S^j$ is a martingale on $\{T\}$. For the inductive step, we claim that if $\overline{Z}S^j$ is a martingale on $\{t+1,\ldots,T\}$ for some $t \in \{0,\ldots,T-1\}$, then $E[\overline{Z}_{t+1}S_{t+1}^j | \mathcal{F}_t] = \overline{Z}_t S_t^j P$ -a.s. so that $\overline{Z}S^j$ is a *P*-martingale on $\{t,\ldots,T\}$. To show this claim, note that the definitions (3.18) and (3.19) yield

$$\bar{Z}_{t+1}\mathbb{1}_{\{\bar{Z}_t\neq 0\}} = \bar{Z}_t \left(1 + \frac{\Delta \bar{Z}_{t+1}}{\bar{Z}_t}\right) \mathbb{1}_{\{\bar{Z}_t\neq 0\}} = \bar{Z}_t (1 + \Delta \bar{N}_{t+1}) \mathbb{1}_{\{\bar{Z}_t\neq 0\}} = \bar{Z}_t \, {}^t \mathcal{E}(\bar{N})_{t+1} \mathbb{1}_{\{\bar{Z}_t\neq 0\}}.$$

By plugging in and recalling that ${}^{t}\mathcal{E}(\bar{N})S^{j}$ is a true *P*-martingale and ${}^{t}\mathcal{E}(\bar{N})_{t} = 1$, we get

$$E[\bar{Z}_{t+1}S_{t+1}^{j} \mid \mathcal{F}_{t}] = \bar{Z}_{t}E[{}^{t}\mathcal{E}(\bar{N})_{t+1}S_{t+1}^{j} \mid \mathcal{F}_{t}]\mathbb{1}_{\{\bar{Z}_{t}\neq0\}} + E[\bar{Z}_{t+1}S_{t+1}^{j} \mid \mathcal{F}_{t}]\mathbb{1}_{\{\bar{Z}_{t}=0\}}$$

$$= \bar{Z}_{t}S_{t}^{j}\mathbb{1}_{\{\bar{Z}_{t}\neq0\}} + E[\bar{Z}_{t+1}S_{t+1}^{j} \mid \mathcal{F}_{t}]\mathbb{1}_{\{\bar{Z}_{t}=0\}}.$$
(B.9)

By the inductive hypothesis and the assumption (3.17), we obtain

$$E[\bar{Z}_{t+1}S_{t+1}^{j} \mid \mathcal{F}_{t}]\mathbb{1}_{\{\bar{Z}_{t}=0\}} = E[\bar{Z}_{T}S_{T}^{j} \mid \mathcal{F}_{t}]\mathbb{1}_{\{\bar{Z}_{t}=0\}} = E[\bar{H}D^{j} \mid \mathcal{F}_{t}]\mathbb{1}_{\{\bar{Z}_{t}=0\}} = 0.$$

Plugging into (B.9) yields

$$E[\bar{Z}_{t+1}S_{t+1}^{j} \mid \mathcal{F}_{t}] = \bar{Z}_{t}S_{t}^{j}\mathbb{1}_{\{\bar{Z}_{t}\neq 0\}} = \bar{Z}_{t}S_{t}^{j}$$

It follows by backward induction that $\overline{Z}S^j$ is a true *P*-martingale on $\{0, \ldots, T\}$ for each $j \in \{d_1+1, \ldots, d_1+d_2\}$, as claimed. This also concludes the proof that $\overline{Z}S^j$ is a local *P*-martingale for all $j \in \{1, \ldots, d_1 + d_2\}$.

(d) We are now ready to show that (1, S) satisfies conditions 1) and 2) of Definition 2.6. We first show that (1, S) satisfies uniqueness of value processes. To that end, suppose that $c_1 + \vartheta^1 \cdot S_T = c_2 + \vartheta^2 \cdot S_T$ for some $c_1, c_2 \in \mathbb{R}$ and $\vartheta^1, \vartheta^2 \in \overline{\Theta}(S)$. Recall from (b) that ${}^s \mathcal{E}(\bar{N})S^j$ is a local *P*-martingale on $\{s, \ldots, T\}$ for each $s \in \{0, \ldots, T-1\}$. Since ${}^s \mathcal{E}(\bar{N})$ is square-integrable by assumption, it follows by Proposition A.7 that ${}^s \mathcal{E}(\bar{N})(c_1 + \vartheta^1 \cdot S)$ and ${}^s \mathcal{E}(\bar{N})(c_2 + \vartheta^2 \cdot S)$ are true *P*-martingales on $\{s, \ldots, T\}$. Because ${}^s \mathcal{E}(\bar{N})_s = 1$, this yields

$$c_1 + \vartheta^1 \bullet S_s = E[{}^s \mathcal{E}(N)_T (c_1 + \vartheta^1 \bullet S_T) \mid \mathcal{F}_s] = E[{}^s \mathcal{E}(\bar{N})_T (c_2 + \vartheta^2 \bullet S_T) \mid \mathcal{F}_s] = c_2 + \vartheta^2 \bullet S_s.$$

In particular, taking s = 0 gives $c_1 = c_2$. As $s \in \{0, \ldots, T\}$ is arbitrary, $\vartheta^1 \cdot S$ and $\vartheta^2 \cdot S$ are indistinguishable, so that $\vartheta^1 =_S \vartheta^2$ and (1, S) satisfies uniqueness of value processes.

Next, we show the existence of solutions to the MVH problem (3.2) for each $H \in L^2$. As in step (b), consider once again the family $\mathcal{E} = ({}^s \mathcal{E}(\bar{N}))_{s \in \{0,...,T\}}$, which is square-integrable by assumption. Let τ be a stopping time taking values in $\{0, \ldots, T\}$. Since ${}^s \mathcal{E}(\bar{N})$ is a martingale by assumption for any $s \in \{0, \ldots, T\}$, so is $1 + \mathbb{1}_{\{\tau=s\}}({}^s\mathcal{E}(\bar{N}) - 1)$ as ${}^s\mathcal{E}(\bar{N}) - 1 = 0$ on $\{0, \ldots, s\}$. Thus we obtain that

$${}^{\tau}\mathcal{E}(\bar{N}) = 1 + \sum_{s=0}^{T} \mathbb{1}_{\{\tau=s\}} \left({}^{s}\mathcal{E}(\bar{N}) - 1\right)$$

is a martingale. As this holds for any stopping time τ , the family \mathcal{E} is so-called regular; see [8, Definitions 3.4 and 3.6]. Thus by Czichowsky/Schweizer [10, Theorem 2.16], the set $\mathcal{G}_T(S)$ is closed in L^2 . This implies the existence of a solution to the MVH problem (3.2) for any payoff $H \in L^2$, since it can be seen as a projection problem in L^2 . The uniqueness of value processes (and thus of gains processes) together with Proposition A.3 yields that the solution to (3.2) is unique for each $H \in L^2$. Since $\eta^k \in \overline{\Theta}(S)$ by condition 3) of Definition 2.6, which we already showed in step (a), it follows from Proposition 3.1 that there exists a unique solution $\hat{\vartheta}^k$ to (3.1) for each $k \in \{1, \ldots, K\}$, and thus condition 1) is satisfied.

It remains to check that (1, S) satisfies condition 2) of Definition 2.6, for which we use the same argument as in the proof of Theorem 3.7. By Lemma A.8 and since $\bar{Z}S^j$ is a local *P*-martingale for each $j \in \{1, \ldots, d_1 + d_2\}$, the strategy 0 solves the MVH problem (3.2) for \bar{H} . Thus $\sum_{k=1}^{K} \hat{\vartheta}^k = \bar{\vartheta} = \bar{\eta}$ by Lemmas 3.3 and 3.5, so that the market clears. This concludes the proof that (1, S) is a quadratic equilibrium.

Proof of Lemma 4.2. Suppose by way of contradiction that $\hat{\vartheta}^k$ is not mean-variance efficient for agent k. Then there exists some $\vartheta' \in \overline{\Theta}(S)$ satisfying

$$E[\vartheta' \bullet S_T + \Xi^k] \ge E[(\hat{\vartheta}^k - \eta^k) \bullet S_T + \Xi^k] = E[V_T^k(\hat{\vartheta}^k)],$$

$$Var[\vartheta' \bullet S_T + \Xi^k] \le Var[(\hat{\vartheta}^k - \eta^k) \bullet S_T + \Xi^k] = Var[V_T^k(\hat{\vartheta}^k)],$$

where one of the inequalities is strict. Since $U_k^{\text{MV}}(\mu, \sigma) = \mu - \frac{\sigma^2}{2\lambda_k}$ is strictly increasing in $\mu \in \mathbb{R}$ and strictly decreasing in $\sigma \in \mathbb{R}_+$, we thus have

$$U_k^{\mathrm{MV}}\Big(E[V_T^k(\hat{\vartheta}^k)], \sqrt{\mathrm{Var}[V_T^k(\hat{\vartheta}^k)]}\Big) < U_k^{\mathrm{MV}}\Big(E[V_T^k(\tilde{\vartheta})], \sqrt{\mathrm{Var}[V_T^k(\tilde{\vartheta})]}\Big)$$

where $\tilde{\vartheta} := \vartheta' + \eta^k \in \overline{\Theta}(S)$, and this contradicts the optimality of $\hat{\vartheta}^k$ for (4.1).

Lemma B.2. Suppose the market (1, S) satisfies Assumption 4.3. A strategy $\vartheta \in \overline{\Theta}(S)$ is mean-variance efficient with respect to $H \equiv 0$ if and only if $\vartheta =_S y \vartheta^{\text{MVH}}(1)$ for some $y \geq 0$. In that case, we have

$$E[\vartheta \bullet S_T] = y(1-\ell) \quad and \quad \operatorname{Var}[\vartheta \bullet S_T] = y^2 \ell(1-\ell). \tag{B.10}$$

Proof of Lemma B.2. A strategy is mean-variance efficient with respect to $H \equiv 0$ in the sense of Definition 4.1 if and only if it satisfies the equivalent conditions (a) and (b) of Eberlein/Kallsen [14, Rule 10.43]. Thus the first assertion follows directly from the equivalence with condition (e) in [14, Rule 10.43], and we obtain (B.10) from [14, Rule 10.47].

Lemma B.3. Suppose the market (1, S) satisfies Assumption 4.3. For any strategy $\vartheta \in \overline{\Theta}(S)$, we have

$$E[V_T^k(\vartheta)] = c_k + E[((\vartheta - \eta^k + \vartheta^{\text{ex}}(\Xi^k)) \bullet S_T], \qquad (B.11)$$

$$\operatorname{Var}[V_T^k(\vartheta)] = \operatorname{Var}\left[\left(\vartheta - \eta^k + \vartheta^{\operatorname{ex}}(\Xi^k)\right) \bullet S_T\right] + \varepsilon_k^2, \tag{B.12}$$

where $(c_k, \vartheta^{\text{ex}}(\Xi^k))$ is the unique solution to the exMVH problem (3.3) with $H = \Xi^k$ and ε_k^2 is given by (4.7).

Proof of Lemma B.3. The definition of $(c_k, \vartheta^{\text{ex}}(\Xi^k))$ yields that $c_k + \vartheta^{\text{ex}}(\Xi^k) \cdot S_T$ is the orthogonal projection of Ξ^k onto the set $\{x + \vartheta \cdot S_T : x \in \mathbb{R}, \vartheta \in \Theta(S)\}$, which is closed in L^2 by Černý/Kallsen [5, Lemma 2.9] and Assumption 4.3. Hence we obtain an orthogonal decomposition of the form

$$\Xi^k = c_k + \vartheta^{\text{ex}}(\Xi^k) \bullet S_T + \tilde{\Xi}^k \tag{B.13}$$

where $\tilde{\Xi}^k \in L^2$ is such that $E[\tilde{\Xi}^k] = E[(\tilde{\vartheta} \cdot S_T)\tilde{\Xi}^k] = 0$ for all $\tilde{\vartheta} \in \overline{\Theta}(S)$ by the orthogonality. Moreover, we have by (4.7) that $\operatorname{Var}[\tilde{\Xi}^k] = \varepsilon_k^2$. Therefore, plugging (B.13) into the formula (2.6) for $V_T^k(\vartheta)$ yields

$$E[V_T^k(\vartheta)] = E[(\vartheta - \eta^k) \bullet S_T + c_k + \vartheta^{\text{ex}}(\Xi^k) \bullet S_T + \tilde{\Xi}^k] = c_k + E[((\vartheta - \eta^k + \vartheta^{\text{ex}}(\Xi^k)) \bullet S_T],$$

$$\operatorname{Var}[V_T^k(\vartheta)] = \operatorname{Var}[(\vartheta - \eta^k) \bullet S_T + c_k + \vartheta^{\text{ex}}(\Xi^k) \bullet S_T + \tilde{\Xi}^k] = \operatorname{Var}[(\vartheta - \eta^k + \vartheta^{\text{ex}}(\Xi^k)) \bullet S_T] + \varepsilon_k^2,$$

which shows (B.11) and (B.12).

Proof of Proposition 4.4. (a) \Leftrightarrow (b): Since c_k and ε_k^2 do not depend on the choice of ϑ , it follows by (B.11) and (B.12) together with Definition 4.1 that ϑ is mean-variance efficient for agent k if and only if $\vartheta - \eta^k + \vartheta^{\text{ex}}(\Xi^k)$ is mean-variance efficient with respect to 0. By Lemma B.2, the latter statement is equivalent to

$$\vartheta - \eta^k + \vartheta^{\text{ex}}(\Xi^k) =_S y \vartheta^{\text{MVH}}(1)$$

for some $y \ge 0$. Thus we have (a) \Leftrightarrow (b).

(b) \Leftrightarrow (c): This was already shown in Proposition 3.2, where $y = \gamma_k - c_k$.

(c) \Leftrightarrow (d): By Lemma 3.1, ϑ is a solution to the quadratic utility problem (3.1) with risk tolerance γ_k if and only if $\vartheta - \eta^k$ is a solution to the MVH problem (3.2) for $H^k(\gamma_k) = \gamma_k - \Xi^k$; in particular, the solution to (3.1) is unique by the uniqueness of the solution to (3.2) for $H^k(\gamma_k)$. This shows (c) \Leftrightarrow (d) and concludes the proof.

Proof of Lemma 4.6. Recall the definition (2.9) of $\mathcal{U}_k^{\text{MV}}$ and U_k^{MV} . We claim that $\hat{y}_k := \lambda_k / \ell$ is a maximiser for the problem

$$U_k^{\mathrm{MV}}(\mu_k(y), \sigma_k(y)) \longrightarrow \max_{y \ge 0}!$$
 (B.14)

Indeed, by plugging in we obtain

$$U_k^{\mathrm{MV}}(\mu_k(y), \sigma_k(y)) = c_k + (1-\ell)y - \frac{\varepsilon_k^2 + \ell(1-\ell)y^2}{2\lambda_k},$$

which is a concave quadratic function of y, so that we obtain the minimiser $\hat{y}_k = \lambda_k/\ell$ by differentiating; this minimiser is unique if and only if $\ell \neq 1$. This in turn gives

$$\mathcal{U}_{k}^{\mathrm{MV}}\left(V_{T}^{k}\left(\vartheta^{k}(\hat{y}_{k})\right)\right) = U_{k}^{\mathrm{MV}}\left(\mu_{k}(\hat{y}_{k}), \sigma_{k}(\hat{y}_{k})\right) \ge U_{k}^{\mathrm{MV}}\left(\mu_{k}(y), \sigma_{k}(y)\right) = \mathcal{U}_{k}^{\mathrm{MV}}\left(V_{T}^{k}\left(\vartheta^{k}(y)\right)\right)$$
(B.15)

for all $y \ge 0$, with equality only in the cases $y = \hat{y}_k$ or $\ell = 1$. Thus by Proposition 4.4, the strategy $\vartheta^k(\hat{y}_k)$ is a maximiser of (4.1) within the set of mean-variance efficient strategies.

It remains to show that $\vartheta^k(\hat{y}_k)$ is also the unique solution to (4.1) among all admissible strategies. To that end, let $\vartheta \in \overline{\Theta}(S)$ be any other strategy. Suppose first that $\ell \neq 1$, so that $0 < \ell < 1$. Then by (B.12), the mean-variance efficient strategy $\vartheta' = \vartheta^k(y)$ for agent k with

$$y := \sqrt{\frac{\operatorname{Var}[(\vartheta - \eta^k + \vartheta^{\operatorname{ex}}(\Xi^k)) \bullet S_T]}{\ell(1 - \ell)}}$$

satisfies $\operatorname{Var}[V_T^k(\vartheta')] = \operatorname{Var}[V_T^k(\vartheta)]$. Since ϑ' is mean-variance efficient, we have $E[V_T^k(\vartheta')] \ge E[V_T^k(\vartheta)]$. By (B.15) and as U_k^{MV} is strictly increasing in μ , it follows that

$$\mathcal{U}_{k}^{\mathrm{MV}}\big(V_{T}^{k}(\vartheta)\big) \leq \mathcal{U}_{k}^{\mathrm{MV}}\Big(V_{T}^{k}\big(\vartheta^{k}(y)\big)\Big) \leq \mathcal{U}_{k}^{\mathrm{MV}}\Big(V_{T}^{k}\big(\vartheta(\hat{y}_{k})\big)\Big).$$
(B.16)

Since $\vartheta \in \overline{\Theta}(S)$ is arbitrary, this shows that $\vartheta^k(\hat{y}_k)$ is indeed a solution to (4.1). Moreover, the two inequalities in (B.16) are equalities only if ϑ is mean-variance efficient and $y = \hat{y}_k$, in which case $\vartheta =_S \vartheta^k(y) =_S \vartheta^k(\hat{y}_k)$. Thus $\vartheta^k(\hat{y}_k)$ is the unique solution to (4.1) in the case $\ell \neq 1$.

Finally, in the case $\ell = 1$, i.e., if $0 \in \overline{\Theta}(S)$ is a solution to the pure investment problem, we have by Lemma A.8 that S is a local P-martingale. In this case, there exists a unique meanvariance efficient $\hat{\vartheta} = \vartheta^k(\hat{y}_k) = \vartheta^k(y)$ for all $y \ge 0$. For any strategy $\vartheta \in \overline{\Theta}(S)$, we have by Proposition A.7 that $(\vartheta - \eta^k + \vartheta^{\text{ex}}(\Xi^k)) \cdot S$ is a P-martingale, so that the second term in (B.11) is null. Hence $E[V_T^k(\vartheta)] = E[V_T^k(\hat{\vartheta})]$ and $\operatorname{Var}[V_T^k(\vartheta)] \ge \operatorname{Var}[V_T^k(\hat{\vartheta})]$, with equality if and only if ϑ is mean-variance efficient, i.e., only in the case $\vartheta =_S \hat{\vartheta}$. Since U_k^{MV} is strictly decreasing in σ , it follows that $\hat{\vartheta} = \vartheta(\hat{y}_k)$ is the unique solution to (4.1) also in the case $\ell = 1$.

Proof of Lemma 4.7. Since (1, S) is a mean-variance equilibrium in the sense of Definition 2.6, there exists for each agent $k \in \{1, \ldots, K\}$ a unique solution $\hat{\vartheta}^k$ to (4.1). Lemma 4.6 yields $\hat{\vartheta}^k = \vartheta^k(\lambda_k/\ell)$, where ϑ^k is defined by (4.8). Thus by (b) \Leftrightarrow (d) in Proposition 4.4, $\hat{\vartheta}^k$ is also the unique solution to the quadratic utility problem (3.1) with $\gamma_k := c_k + \lambda_k/\ell$. Therefore, since (1, S) is a mean-variance equilibrium by assumption, it also satisfies all of the conditions in Definition 2.6 for a quadratic equilibrium with individual risk tolerances $\gamma_1, \ldots, \gamma_K$.

Proof of Lemma 4.9. As in Lemma 3.6, define the martingale $(\bar{Z}_t(\bar{\gamma}))_{0 \leq t \leq T}$ by $\bar{Z}_t(\bar{\gamma}) = E[\bar{H}(\bar{\gamma}) | \mathcal{F}_t]$. Note that we have the bounds $\bar{\gamma} - \bar{\gamma}_0 \leq \bar{H}(\bar{\gamma}) \leq \bar{\gamma}$, and hence also

$$0 < \bar{\gamma} - \bar{\gamma}_0 \le \bar{Z}_t(\bar{\gamma}) \le \bar{\gamma}$$

for all $t \in [0, T]$. In particular, the process $\overline{Z}(\bar{\gamma})$ is strictly positive and never hits 0. Moreover, condition (b) in Lemma 3.8 is satisfied, so that $S^{(2)}(\bar{\gamma})$ is an L^2 -semimartingale. We can then apply Theorem 3.7, which yields that $S(\bar{\gamma})$ is the unique quadratic equilibrium with respect to any choice of parameters $\gamma_1, \ldots, \gamma_K$ such that $\sum_{k=1}^K \gamma_k = \bar{\gamma}$. Lemma 3.6 also gives that $\overline{Z}(\bar{\gamma})S(\bar{\gamma})$ is a local *P*-martingale. Thus $Q(\bar{\gamma})$ is a local martingale measure for $S(\bar{\gamma})$ such that $Q(\bar{\gamma}) \approx P$, and $dQ(\bar{\gamma})/dP$ is bounded because $\overline{H}(\bar{\gamma})$ is strictly positive and bounded. This also implies that $S(\bar{\gamma})$ satisfies Assumption 4.3, which concludes the proof.

Proof of Lemma 4.11. By Lemma 4.9, $(1, S(\bar{\gamma}))$ is the unique quadratic equilibrium with aggregate risk tolerance $\bar{\gamma}$ and $\bar{Z}(\bar{\gamma})S(\bar{\gamma})$ is a local martingale. This shows the "if" statement. To prove the converse, suppose for a contradiction that $(1, S(\bar{\gamma}))$ is also a quadratic equilibrium with respect to some risk tolerances $\gamma_1, \ldots, \gamma_K$ such that $\sum_{k=1}^K \gamma_k =: \bar{\gamma}' \neq \bar{\gamma}$. Then by Lemma 3.6, the process $\bar{Z}(\bar{\gamma}')S(\bar{\gamma})$ is also a local martingale. Taking differences yields that

$$\left(\bar{Z}(\bar{\gamma}') - \bar{Z}(\bar{\gamma})\right)S(\bar{\gamma}) = (\bar{\gamma}' - \bar{\gamma})S(\bar{\gamma})$$

is a local martingale as well, and so is $S(\bar{\gamma})$ because $\bar{\gamma}' \neq \bar{\gamma}$. Thus the implication (a) \Rightarrow (d) in [23, Lemma 2.27] leads to a contradiction of Assumption 4.10, so that $\bar{\gamma}' \neq \bar{\gamma}$ cannot hold. This concludes the proof of the equivalence. The second statement follows from the first, since if $S(\bar{\gamma}) = S(\bar{\gamma}')$, then $(1, S(\bar{\gamma}))$ is the unique quadratic equilibrium with aggregate risk tolerance $\bar{\gamma}'$ by Lemma 4.9 so that $\bar{\gamma} = \bar{\gamma}'$. Proof of Proposition 4.12. We start by proving "only if". If $(1, S(\bar{\gamma}))$ is a mean-variance equilibrium, we have by Lemma 4.7 that it is also a quadratic equilibrium with respect to the risk tolerances given by $\bar{\gamma}_k(\bar{\gamma}) := c_k(\bar{\gamma}) + \lambda_k/\ell(\bar{\gamma})$. But then Lemma 4.11 yields $\bar{\gamma} = \tilde{\gamma}(\bar{\gamma})$ as claimed.

To show the "if" statement, suppose that $\bar{\gamma} = \tilde{\gamma}(\bar{\gamma})$. Then by Lemma 4.11, $(1, S(\bar{\gamma}))$ is the unique quadratic equilibrium with risk tolerances $\gamma_k(\bar{\gamma}) := c_k(\bar{\gamma}) + \lambda_k/\ell(\bar{\gamma})$. By Definition 2.6, there exist unique solutions $\hat{\vartheta}^k$ to the individual problems (3.1) with $\gamma_k = \gamma_k(\bar{\gamma})$. By the equivalence (b) \Leftrightarrow (d) in Proposition 4.4, we have $\hat{\vartheta}^k = \vartheta^k(\lambda_k/\ell(\bar{\gamma}))$, and hence by Lemma 4.6, $\hat{\vartheta}^k$ is also the unique solution to the mean–variance problem (4.1). Thus $(1, S(\bar{\gamma}))$ satisfies condition 1) of Definition 2.6 of a mean–variance equilibrium. Since $(1, S(\bar{\gamma}))$ is a quadratic equilibrium, the remaining conditions of Definition 2.6 are also satisfied so that $(1, S(\bar{\gamma}))$ is a mean–variance equilibrium.

Proof of Proposition 4.13. Fix $\bar{\gamma} > \bar{\gamma}_0$ and write $\vartheta^{(0)}(\bar{\gamma})$ as a shorthand for $\vartheta^{(0)}(1; S(\bar{\gamma}))$. Recall from Lemma 4.9 that $\bar{Z}(\bar{\gamma})S(\bar{\gamma})$ is a local *P*-martingale, where $\bar{Z}(\bar{\gamma})$ is a strictly positive *P*-martingale. Since $\vartheta^{(0)}(\bar{\gamma}) \in \overline{\Theta}(S(\bar{\gamma}))$, Proposition A.7 yields that $\bar{Z}(\bar{\gamma})(\vartheta^{(0)}(\bar{\gamma}) \cdot S(\bar{\gamma}))$ is a true *P*-martingale, and hence so is $\bar{Z}(\bar{\gamma})(1 - \vartheta^{(0)}(\bar{\gamma}) \cdot S(\bar{\gamma}))$. As mentioned above (A.4), $(\bar{V}_s(\bar{\gamma})M_s^{(0)}(\bar{\gamma}))_{0\leq s\leq T}$ is also a *P*-martingale. Moreover, by (A.4) and as $L_T(\bar{\gamma}) = 1$, we have

$$\bar{V}_T(\bar{\gamma})M_T^{(0)}(\bar{\gamma}) = \bar{H}(\bar{\gamma})L_T(\bar{\gamma})\Big(1 - \big(\vartheta^{(0)}(\bar{\gamma})\bullet S(\bar{\gamma})\big)_T\Big) = \bar{Z}_T(\bar{\gamma})\Big(1 - \big(\vartheta^{(0)}(\bar{\gamma})\bullet S(\bar{\gamma})\big)_T\Big).$$

Thus by taking expectations under P, we obtain

$$\bar{V}_0(\bar{\gamma})M_0^{(0)}(\bar{\gamma}) = \bar{Z}_0(\bar{\gamma}) \Big(1 - \big(\vartheta^{(0)}(\bar{\gamma}) \cdot S(\bar{\gamma})\big)_0 \Big) = \bar{Z}_0(\bar{\gamma})$$

Recall that $\bar{V}_0(\bar{\gamma}) = \bar{\gamma} - c(\bar{\Xi}; S(\bar{\gamma}))$ and $M_0^{(0)}(\bar{\gamma}) = L_0(\bar{\gamma}) = \ell(\bar{\gamma})$ by (A.4), whereas we have $\bar{Z}_0(\bar{\gamma}) = E_P[\bar{H}(\bar{\gamma})] = \bar{\gamma} - E_P[\bar{\Xi}]$. Plugging in yields

$$\bar{\gamma} - E_P[\bar{\Xi}] = \left(\bar{\gamma} - c(\bar{\Xi}; S(\bar{\gamma}))\right) \ell(\bar{\gamma}).$$

Since the linearity of exMVH (see Lemma A.1) yields

$$c(\bar{\Xi}; S(\bar{\gamma})) = \sum_{k=1}^{K} c(\Xi^{k}; S(\bar{\gamma})) = \sum_{k=1}^{K} c_{k}(\bar{\gamma}) = \bar{c}(\bar{\gamma}),$$

we obtain (4.12), which concludes the proof.

Proof of Theorem 4.14. Let $\gamma' > \bar{\gamma}_0$ and define $\ell(\bar{\gamma}')$ and $c_k(\bar{\gamma}')$ with respect to $S = S(\bar{\gamma}')$ in the same way as below Assumption 4.3. By Proposition 4.12, $(1, S(\bar{\gamma}'))$ is a mean-variance equilibrium if and only if $\bar{\gamma}'$ satisfies the fixed-point condition $\tilde{\gamma}(\bar{\gamma}') = \bar{\gamma}'$, i.e., if and only if

$$\left(\bar{\gamma}' - \bar{c}(\bar{\gamma}')\right)\ell(\bar{\gamma}') = \sum_{k=1}^{K} \lambda_k, \tag{B.17}$$

where we recall $\bar{c}(\bar{\gamma}') := \sum_{k=1}^{K} c_k(\bar{\gamma}')$. Moreover, Proposition 4.13 gives

$$\bar{\gamma}' - E_P[\bar{\Xi}] = \left(\bar{\gamma}' - \bar{c}(\bar{\gamma}')\right) \ell(\bar{\gamma}'). \tag{B.18}$$

By plugging (B.18) into (B.17), we conclude that $\bar{\gamma}' \in (\bar{\gamma}_0, \infty)$ is a fixed point if and only if $\bar{\gamma}' - E_P[\bar{\Xi}] = \sum_{k=1}^K \lambda_k$, i.e., if and only if $\tilde{\gamma} = \bar{\gamma}$. Therefore $\bar{\gamma}$ defined by (4.13) is the only possible solution to the condition $\tilde{\gamma}(\bar{\gamma}') = \bar{\gamma}'$, and it is indeed a solution if $\bar{\gamma} > \bar{\gamma}_0$. In that case, we have by Proposition 4.12 that $(1, S(\bar{\gamma}))$ is a mean-variance equilibrium.

References

- Michael Allingham, Existence theorems in the capital asset pricing model, Econometrica 59 (1991), 1169–1174.
- [2] Jonathan Berk, Necessary conditions for the CAPM, Economic Theory 73 (1997), 245–257.
- [3] D. T. Breeden, An intertemporal asset pricing model with stochastic consumption and investment opportunities, Journal of financial Economics 7 (1979), no. 3, 265–296.
- [4] Aleš Černý and Christoph Czichowsky, The law of one price in quadratic hedging and mean-variance portfolio selection, Preprint, City, University of London, (v2) available at arXiv: https://arxiv.org/abs/2210.15613, 2024.
- [5] Aleš Černý and Jan Kallsen, On the structure of general mean-variance hedging strategies, Annals of Probability 35 (2007), 1479–1531.
- [6] _____, A counterexample concerning the variance-optimal martingale measure, Mathematical Finance 18 (2008), 305–316.
- [7] Alexander Cherny and Albert Shiryaev, Vector stochastic integrals and the fundamental theorem of asset pricing, Tr. Mat. Inst. Steklova (Stokhast. Finans. Mat.) 237 (2002), 12–56.
- [8] Tahir Choulli, Leszek Krawczyk, and Christophe Stricker, *E-martingales and their appli*cations in mathematical finance, Annals of Probability 26 (1998), 853–876.
- [9] Christoph Czichowsky and Martin Schweizer, Closedness in the semimartingale topology for spaces of stochastic integrals with constrained integrands, Séminaire de Probabilités XLIII (Catherine Donati-Martin, Antoine Lejay, and Alain Rouault, eds.), Springer, 2011, pp. 413–436.
- [10] _____, Cone-constrained continuous-time Markowitz problems, Annals of Applied Probability 23 (2013), 764–810.
- [11] Rose-Anne Dana, Existence, uniqueness and determinacy of equilibrium in C.A.P.M. with a riskless asset, Mathematical Economics 32 (1999), 167–175.
- [12] Freddy Delbaen and Walter Schachermayer, Attainable claims with p'th moments, Annales de l'IHP Probabilités et Statistiques 32 (1996), 743–763.
- [13] _____, A simple counterexample to several problems in the theory of asset pricing, Mathematical Finance 8 (1998), 1–11.
- [14] Ernst Eberlein and Jan Kallsen, *Mathematical finance*, Springer, 2019.
- [15] Thorsten Hens, Jörg Laitenberger, and Andreas Löffler, Two remarks on the uniqueness of equilibria in the CAPM, Mathematical Economics 37 (2002), 123–132.
- [16] Jean Jacod and Albert Shiryaev, Limit theorems for stochastic processes, 2nd ed., Springer, 2003.
- [17] Pablo Koch-Medina and Jan Wenzelburger, Equilibria in the CAPM with non-tradeable endowments, Mathematical Economics 75 (2018), 93–107.

- [18] Haim Levy, The capital asset pricing model in the 21st century: Analytical, empirical, and behavioral perspectives, Cambridge University Press, 2011.
- [19] John Lintner, Security prices, risk, and maximal gains from diversification, Finance 20 (1965), 587–615.
- [20] _____, The valuation of risky assets and the selection of risky investments in stock portfolios and capital budgets, Review of Economic Statistics 47 (1965), 346–382.
- [21] Harry Markowitz, Portfolio theory: as i still see it, Annu. Rev. Financ. Econ. 2 (2010), no. 1, 1–23.
- [22] _____, Mean-variance approximations to expected utility, European Journal of Operational Research 234 (2014), no. 2, 346–355.
- [23] David Martins, Aspects of quadratic utility: mean-variance hedging in rough volatility models, and CAPM-type equilibria, Ph.D. thesis, ETH Zürich, 2023.
- [24] Jan Mossin, Equilibrium in a capital asset market, Econometrica **39** (1966), 768–783.
- [25] Lars Nielsen, Uniqueness of equilibrium in the classical capital asset pricing model, Financial and Quantitative Analysis 23 (1988), 329–336.
- [26] _____, Equilibrium in CAPM without a riskless asset, Review of Economic Studies 57 (1990), 315–324.
- [27] _____, Existence of equilibrium in CAPM, Economic Theory 52 (1990), 223–231.
- [28] Martin Schweizer, Mean-variance hedging, Encyclopedia of Quantitative Finance (Rama Cont, ed.), Wiley, 2010, pp. 1177–1181.
- [29] William Sharpe, Capital asset prices: A theory of market equilibrium under conditions of risk, Finance 19 (1964), 425–442.
- [30] R. C. Stapleton and M. G. Subrahmanyam, A multiperiod equilibrium asset pricing model, Econometrica 5 (1978), no. 46, 1077–1096.
- [31] Christophe Stricker, Arbitrage et lois de martingale, Annales de l'I.H.P. Probabilités et statistiques 26 (1990), 451–460 (fre).
- [32] Jack Treynor, Toward a theory of market value of risky assets, Unpublished manuscript, 1962.
- [33] Jan Wenzelburger, The two-fund separation theorem revisited, Annals of Finance 6 (2010), 221–239.