1

Clustering and synchronization analysis of Networks of Bistable Systems

Gianluca Villani and Luca Scardovi

Nonlinear dyn of multiple stab This phenomenor across various r ics, chemistry, a significant role i context of Gene F that control cell multistable dynar

Abstract—This paper studies the dynamics of a network of diffusively-coupled bistable systems. Under mild conditions and without requiring smoothness of the vector field, we analyze the network dynamics and show that the solutions converge globally to the set of equilibria for generic monotone (but not necessarily strictly monotone) regulatory functions. Sufficient conditions for global state synchronization are provided. Finally, by adopting a piecewise linear approximation of the vector field, we determine the existence, location and stability of the equilibria as function of the coupling gain. The theoretical results are illustrated with numerical simulations.

I. INTRODUCTION

Nonlinear dynamical systems can exhibit the coexistence of multiple stable equilibria or, more generally, attractors. This phenomenon, known as multistability [1], is ubiquitous across various natural sciences, including optics, mechanics, chemistry, and biology [2], [3]. Multistability plays a significant role in fundamental biological processes. In the context of Gene Regulatory Networks (GRNs), genetic circuits that control cell fate decisions are traditionally modeled as multistable dynamical systems [4], with different stable steady states representing different cell phenotypes [5]. It has been conjectured [6] and proved [7] that the presence of positive feedback loops is a necessary condition for the existence of multiple equilibria. The synthetic toggle switch is one of the first examples where these theoretical results have been experimentally verified in a biological system [8].

Recently, networks of bistable switches coupled by quorum sensing [9] and diffusive coupling [10]-[13] have been investigated, with their properties of auto-correction, synchronization and convergence to equilibria characterized. In [11], [12] the property of convergence to equilibria has been conjectured without a formal analysis. In [9], the property of (almost) global convergence to equilibria was proved using the theory of strongly monotone systems [14]. However, this property is lost when the graph associated with the signed Jacobian of the vector field is not irreducible everywhere. This occurs, for example, when the interactions between different chemical species are modeled by piece-wise affine functions with intervals where the functions are constant. In [13] this property has been established for a particular class of networks with strictly increasing nonlinearities. Alternative approaches to prove convergence to equilibria approaches include the use of Lyapunov functions [15]–[19], contraction theory [20], [21], and leveraging passivity-like properties [22]-[26].

In this paper, we investigate a fairly general class of networks of bistable systems coupled by diffusion, focusing on convergence to equilibria and synchronization properties. The model encompasses both positive autoregulation loops and mutual repression circuits (toggle switches). Our first result establishes that all trajectories converge to the set of equilibria, regardless of the system's parameters, coupling strength, and network topology, and for generic monotone (not necessarily strictly monotone) regulatory functions. This result builds upon the theory of counterclockwise Input/Output systems [22]–[24] and generalizes results requiring strictly monotone regulation functions [9], [13].

Our second result provides sufficient conditions for asymptotic state synchronization. This result does not require a specific network topology but establishes a condition that involves the degree of connectivity of the network topology and the isolated system's dynamics.

Our last result addresses a biologically relevant networked model by adopting a piece-wise linear approximation of the regulatory functions. In particular, we characterize the local stability of equilibria and, for an all-to-all homogeneous network, we analytically characterize the emergence of stable clustering configurations when the strength of the diffusive coupling is varied. To the best of our knowledge, such an exhaustive study of convergence, synchronization, and clustering properties has not been previously reported in the literature.

The paper is organized as follows. In Section II, we introduce the class of compartmental models under investigation. Section III establishes the property of convergence to equilibria. In Section IV, we provide sufficient conditions for global synchronization. In Section V, we characterize the local stability of equilibria and clustering properties for an all-toall network of piecewise affine bistable systems. Finally, in Section VI, we illustrate the results with numerical simulations and suggest directions for future research. The appendix reports the main technical preliminaries used throughout the paper.

II. A CLASS OF NETWORKS OF BISTABLE SYSTEMS

A. A class of Bistable Motifs

In this section, we present a class of interconnected compartmental models that exemplifies a network of bistable systems. The dynamics of each compartment is described by the ordinary differential equation

$$\dot{x}_1 = -\gamma_1 x_1 + V_1 g_2(x_2)
\dot{x}_2 = -\gamma_2 x_2 + V_2 g_1(x_1),$$
(1)

where the positive scalar values x_1 and x_2 represent the concentration of two chemical species and the positive constants

This research was supported by the National Sciences and Engineering Research Council of Canada.

The authors are with the Department of Electrical and Computer Engineering, University of Toronto, 10 King's College Road, Toronto, ON, M5S 3G4, Canada. gianluca.villani@mail.utoronto.ca, luca.scardovi@utoronto.ca

 V_1 and V_2 determine their synthesis rates. The regulation functions $g_i : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ are assumed to be monotone functions, positive-valued and either both monotonically increasing or monotonically decreasing¹. The class of models in (1)provides a unified framework for studying the dynamics of two important bistable motifs in gene regulatory networks. When both regulation functions are monotonically non-decreasing, (1) includes Positive Auto Regulation (PAR) loops [27], where two transcription factors activate each other. Autoinduction circuits, such as those involved in quorum sensing systems [28]-[30], are typically modeled as PAR loops. Conversely, when the regulation functions are both monotonically decreasing (inhibitory interactions), (1) includes the well-known toggle switch where two proteins mutually inhibit each other's synthesis [8]. Under appropriate assumptions on the regulation functions and the system's parameters, both types of systems exhibit bistability, with two stable steady states. Figure 1 shows an example of a phase portrait of the vector field in (1) where both the regulation functions are piecewise affine functions (4).

One of the most common modelling choices for the regulation functions for an activator is the Hill function

$$g_s(x) = \frac{(x/\theta)^n}{1 + (x/\theta)^n},\tag{2}$$

where $\theta > 0$ defines the concentration of x to significantly activate expression and n is the cooperative degree corresponding to different steepness values of the function g_s . The regulation function for a repressor is instead modelled as

$$g_s^-(x) = \frac{1}{1 + (x/\theta)^n} = 1 - g_s(x), \tag{3}$$

where $\theta > 0$ represents the concentration of x to significantly repress expression. The sigmoidal Hill regulatory functions can be approximated with piecewise affine continuous functions [27], [31]

$$g(x) = \begin{cases} 0, & x < \theta\\ (x - \theta)/\delta, & \theta \le x \le \theta + \delta\\ 1 & x > \theta + \delta \end{cases}$$
(4)

with corresponding inhibitory regulatory functions $g^{-}(x) = 1 - g(x)$. In addition to the activator and repressor functions described above, another important case to consider is when one of the functions is a linear increasing function, i.e., g(x) = x. This case corresponds to a situation where one of the chemical species is a synthase, and the corresponding synthesized molecule is synthesized at a rate proportional to its concentration [29], [30]. In this paper, unless specified, we will not assume specific regulatory functions but will rely on the following assumption.

Assumption 1. 1 The functions g_i , i = 1, 2, in (1) are positive-valued, Lipschitz continuous, and either both monotonically increasing or both monotonically decreasing. Furthermore, at least one of the functions is bounded.

¹In this paper we will use the terminology monotonically increasing (decreasing) to indicate that a function is non-decreasing (non-increasing).



Figure 1: Phase portrait for (1) with regulatory functions g in (4). The grey box with orange edges highlights the forward invariant (and attractive) set \mathcal{B} . The full black circles represent the two stable equilibria and the white circle represents the unstable saddle equilibrium.

This assumption guarantees the boundedness of all forward trajectories of (1), as shown in the following proposition.

Proposition II.1. If Assumption 1 holds, there exist \bar{x}_1 and \bar{x}_2 such that the set

$$\mathcal{B} := [0, \bar{x}_1] \times [0, \bar{x}_2] \tag{5}$$

is forward invariant with respect to the dynamics (1).

Proof. To prove that the set \mathcal{B} is invariant, it is sufficient to prove that on the boundary $\partial \mathcal{B}$, the vector field in (1) satisfies the Nagumo condition [32]. This requires the following conditions to be met on $\partial \mathcal{B}$:

i) $f_1(0, x_2) \ge 0, \forall x_2 \in [0, \bar{x}_2];$ ii) $f_2(x_1, 0) \ge 0, \forall x_1 \in [0, \bar{x}_1];$ iii) $f_1(\bar{x}_1, x_2) \le 0, \forall x_2 \in [0, \bar{x}_2];$ iv) $f_2(x_1, \bar{x}_2) \le 0, \forall x_1 \in [0, \bar{x}_1].$

Since g_1 and g_2 are nonnegative functions, the first two conditions are trivially satisfied. The last two conditions are equivalent to

$$\begin{aligned} \gamma_1 \bar{x}_1 &\geq V_1 g_2(x_2), \forall x_2 \in [0, \bar{x}_2], \\ \gamma_2 \bar{x}_2 &\geq V_2 g_1(x_1), \forall x_1 \in [0, \bar{x}_1]. \end{aligned}$$
(6)

Without loss of generality, assume that g_1 is upper bounded with bound M > 0. From the second equation, we obtain $\bar{x}_2 \ge V_2 M/\gamma_2$. Using this bound in the first inequality we obtain the second bound $\bar{x}_1 \ge g_2(V_2/\gamma_2)V_1/\gamma_1$. We conclude that \mathcal{B} is forward invariant for any $\bar{x}_1 \ge g_2(MV_2/\gamma_2)V_1/\gamma_1$ and $\bar{x}_2 \ge MV_2/\gamma_2$.

B. Interconnected bistable motifs

Simple network motifs as the one described in (1) are often not isolated but interact through a common shared medium (quorum sensing, [33], [9], [34]) or diffusive coupling [35], [10]. We consider here a networked system with N identical compartments, each defined by (1), and coupled via diffusive coupling

$$\dot{x}_{1,i} = -\gamma_1 x_{1,i} + V_1 g_2(x_{2,i}) + \sum_j a_{i,j} \left(x_{1,j} - x_{1,i} \right),$$

$$\dot{x}_{2,i} = -\gamma_2 x_{2,i} + V_2 g_1(x_{1,i}),$$

(7)

i = 1, ..., N. The non-negative coefficients $a_{i,j}$ are coupling coefficients between the first species in different compartments. In the rest of the paper we will assume that the communication between compartments relies solely on the diffusion of the first species over an undirected graph, i.e. $a_{i,j} = a_{j,i}$. This assumption is realistic since it often applies that only one of the regulatory molecules can diffuse through the cell's membrane [29], [30], [34]. By defining $X_1 = [x_{1,1}, ..., x_{1,N}]^T$, $X_2 = [x_{2,1}, ..., x_{2,N}]^T$ and the vectors of activations $G_{\ell}(X_{\ell}) = [g_{\ell}(x_{\ell,1}), ..., g_{\ell}(x_{\ell,N})]$, $\ell = 1, 2$, we can rewrite (7) in the compact form

$$X_1 = -(\Gamma_1 + L)X_1 + V_1G_2(X_2),$$

$$\dot{X}_2 = -\Gamma_2X_2 + V_2G_1(X_1),$$
(8)

where $L = [l_{ij}]$, is the Laplacian matrix defined as

$$l_{ij} = \begin{cases} -a_{ij}^{\ell}, & i \neq j \\ \sum_{i \neq j} a_{ij}^{\ell}, & i = j \end{cases},$$

 $\Gamma_i = \gamma_i I_N$ with I_N being the $N \times N$ identity matrix. The parameters a_{ij} can be associated to a graph where species in different compartments are the nodes and $a_{ij} > 0$ corresponds to a weighted edge in the graph. We will denote the resulting graph as \mathcal{G} . The graph \mathcal{G} is connected if given any two nodes there exists a path that connects them and \mathcal{G} is symmetric (or undirected) if $a_{ij} = a_{ji}$ for every i, j. In the following we will use the concept of algebraic connectivity extended to directed graphs [36].

Definition II.1. [36] For a directed graph with Laplacian matrix L, the algebraic connectivity is the real number defined as

$$\lambda := \min_{z \in \mathcal{P}} z^T L z$$

where $\mathcal{P} = \{z \in \mathbb{R}^n : z \perp \mathbb{1}_n, \|z\|=1\}$ and where $\mathbb{1}_n \triangleq [1, 1, \dots, 1]^T \in \mathbb{R}^n$

From the definition of \mathcal{B} in (5) we can define a forward invariant hyperbox for the networked system in (7)

$$\mathcal{B}^N = \prod_{i=1}^N \mathcal{B}.$$
 (9)

The proof of the invariance of \mathcal{B}^N follows the same lines of the proof for Proposition II.1 and is therefore omitted. For the rest of the paper, we will consider \mathcal{B}^N as the state space.

III. CONVERGENCE ANALYSIS

In the next result, we show that all trajectories of (7) converge to the set of equilibria, independently of the system's parameters, coupling strength, and network topology.

Theorem III.1. Consider system (7) under Assumption 1 and assume that $a_{i,j} = a_{j,i}, i, j = 1, ..., N$. Then, for every initial condition $\xi \in \mathcal{B}^N$ the ω -limit set $\omega(\xi)$, contains only equilibria of (7). If the equilibria of (7) are isolated, for all initial conditions, $\lim_{t\to+\infty} x(t) = \bar{x}$, where \bar{x} is an equilibrium of (7).

In the proof of Theorem III.1, we will make use of the notion of Counterclockwise (CCW) input-output dynamics introduced in [22]–[24]. The main definitions and results used in the paper are reported in the Appendix.

A. Proof of Theorem III.1

As a first step, we rewrite (7) as the positive feedback interconnection

$$\begin{aligned} \dot{x}_{1,i} &= -\gamma_1 x_{1,i} + V_1 g_2(x_{2,i}) + u_i \\ \dot{x}_{2,i} &= -\gamma_2 x_{2,i} + V_2 g_1(x_{1,i}) \\ y_i &= x_{1,i} \\ u &= -Ly. \end{aligned} \tag{10}$$

 $i = 1, ..., N, y = [x_{1,1}, ..., x_{1,N}]^T$ and $u = [u_1, ..., u_N]$. We outline the proof in the following steps. In Proposition III.1 we show that each component of (10) has strictly counterclockwise (CCW) input-output dynamics from u_i to y_i . Then, by exploiting the properties of positive feedback loop interconnections of systems with CCW input-output dynamics, we characterize the omega limit sets of the networked system (7).

The next Lemma is instrumental in the proof of Proposition III.1 and is an adaptation of [23, Lemma III.1]. Note that, unlike the original result in [23, Lemma III.1], we do not require g to be a strictly increasing function of the input.

Lemma III.1. Consider the scalar input-output system

$$\dot{x} = -kx + g(u) = f(x, u), \quad y = h(x)$$
 (11)

where k > 0, $u \in \mathcal{U}$, and \mathcal{U} is the set of Lebesgue measurable, locally essentially bounded functions valued in $U = [0, u_{\text{max}}]$. Let $X = [x_{\min}, x_{\max}], x_{\max} \ge g(u_{\max})/k$ and $x_{\min} \le g(0)/k$, be the state space of (11). Assume that h and g are either both monotonically increasing or both monotonically decreasing Lipschitz continuous functions. Then, (11) has CCW input-output dynamics with respect to arbitrary density functions ρ , i.e.

$$\liminf_{T \to +\infty} \int_{0}^{T} \dot{y}(t) \int_{0}^{u(t)} \rho(\eta, h(x(t))) d\eta \ dt > -\infty$$
(12)

for every $x_0 \in X$, $u \in \mathcal{U}$, and ρ satisfying Definition A.1.

Proof. Let g and h be monotonically increasing functions. For all input values $u(t) \in U$, $f(x_{\min}, u(t)) \geq 0$ and $f(x_{\max}, u(t)) \leq 0$, therefore the state space X is positively invariant under the dynamics defined by (11). We claim that the set $(g(0), g(u_{\max}))$ is also positively invariant. For any

$$x(t) = e^{-kt} x_0 + \int_0^t e^{-k(t-\tau)} g(u(\tau)) d\tau.$$
 (13)

Since $g(0) \leq g(u(t)) \leq g(u_{\max})$, from (13) we obtain

$$x(t) \ge g(0)/k + e^{-kt} (x_0 - g(0)/k) > g(0)/k$$
 (14)

and

$$x(t) \le g(u_{\max})/k + e^{-kt} \left(x_0 - g(u_{\max})/k \right) < g(u_{\max})/k,$$
(15)

thus proving the claim. We can therefore define the partition $X \times U = S_l \cup S_m \cup S_h$, where

$$\begin{split} \mathcal{S}_{l} &= \{(x_{0}, u) : x(t, x_{0}, u(t)) \in [x_{\min}, g(0)/k], \forall t \geq 0\}, \\ \mathcal{S}_{h} &= \{(x_{0}, u) : x(t, x_{0}, u(t)) \in [g(u_{\max})/k, x_{\max}], \forall t \geq 0\}, \\ \mathcal{S}_{m} &= \{(x_{0}, u) : \exists t^{*} \geq 0 : \forall t \geq t^{*}, \\ &x(t, x_{0}, u(t)) \in (g(0)/k, g(u_{\max}/k))\}. \end{split}$$

We now show that (12) holds on each component of the partition. Consider a generic pair $x_0 \times u \in S_1$. By definition of S_1 and since g is non-decreasing, $f(x(t), u(t)) = -kx(t) + g(u(t)) \ge 0$, for every $t \ge 0$. Since h is Lipschitz continuous, its derivative is defined almost everywhere and, by assumption, $Dh(x) := \frac{d}{dx}h(x) \ge 0$, where defined. We conclude that $\dot{y}(t) = Dh(x(t)) f(x(t), u(t)) \ge 0, \forall t \ge 0$ almost everywhere and therefore

$$\liminf_{T \to +\infty} \int_0^T \dot{y}(t) \int_0^{u(t)} \rho(\mu, y(t)) d\mu dt > -\infty, \tag{16}$$

for every ρ satisfying Definition A.1.

We now show that the inequality (12) holds for all pairs $x_0 \times u \in S_m$. Since the function g is non-decreasing, $\forall x \in (g(0)/k, g(u_{\max})/k)$ we can define $\gamma(x) := g^{-1}(kx)$ where g^{-1} is the generalized inverse function of the monotone function g, in the sense of Definition A.5. The function $\gamma(x)$ is strictly increasing by Proposition A.1, and bounded. Since $\dot{y} = Dh(x)f(x,u), \ \dot{y} > 0 \implies f(x,u) = -kx + g(u) > 0 \iff g(u) > kx \implies u > \gamma(x)$ with the final implication following from Proposition A.1. Analogously, $\dot{y} < 0 \implies u < \gamma(x)$ and therefore we obtain the following inequality

$$\dot{y}(t) \int_{0}^{u(t)} \rho(\eta, y(t)) d\eta \ge \dot{y}(t) \int_{0}^{\gamma(x(t))} \rho(\eta, y(t)) d\eta.$$
(17)

The function

$$F(x) := \int_{0}^{x} Dh(\xi) \int_{0}^{\gamma(x)} \rho(\eta, h(\xi)) d\eta d\xi$$
(18)

is Lipschitz in x since it is the integral of a bounded function. Furthermore, F(x(t)) is absolutely continuous, since it is the composition of a Lipschitz function with the absolutely continuous function x(t). We conclude that its time derivative

$$\frac{\mathrm{d}}{\mathrm{d}t}F(x(t)) = \dot{y}(t) \int_{0}^{\gamma(x(t))} \rho(\eta, y(t)) d\eta$$
(19)

is defined almost everywhere. Integrating both sides of (17) we obtain

$$\int_{0}^{T} \dot{y}(t) \int_{0}^{u(t)} \rho(\eta, y(t)) d\eta dt \ge F(x(T)) - F(x(t^*)) + \Delta$$

where

$$\Delta = \int_{0}^{t^*} \dot{y}(t) \int_{0}^{u(t)} \rho(\eta, y(t)) d\eta dt$$

is finite and accounts for the interval of time during which the solution x(t) is outside of $(g(0)/k, g(u_{\max})/k)$. Therefore, by the continuity of F and boundedness of the trajectories, (12) holds. Consider now a generic pair $x_0 \times u \in S_h$. From (11) it holds that $\dot{y}(t) = Dh(x(t))f(x(t), u(t)) \leq 0, \forall t \geq 0$ almost everywhere, and therefore

$$\dot{y}(t)\int_0^{u(t)}\rho(\mu,y(t))d\mu dt \geq \dot{y}(t)\int_0^{u_{\max}}\rho(\mu,y(t))d\mu dt.$$

Integrating both sides we obtain

$$\int_{0}^{T} \dot{y}(t) \int_{0}^{u(t)} \rho(\eta, y(t)) d\eta dt \ge P(x(T)) - P(x(0))$$
(20)

where

$$P(x) = \int_{0}^{x} Dh(\xi) \int_{0}^{u_{\text{max}}} \rho(\eta, h(\xi)) d\eta d\xi$$
(21)

from which (12) follows.

The proof for the case of g and h mononotically decreasing follows the same lines.

Proposition III.1. Consider the following two-dimensional system

$$\dot{x}_1 = -\gamma_1 x_1 + V_1 g_2(x_2) + u$$

$$\dot{x}_2 = -\gamma_2 x_2 + V_2 g_1(x_1)$$

$$y = x_1$$
(22)

where g_i , i = 1, 2, satisfy Assumption 1. Assume that the input signal $u(t) \in [u_{\min}, u_{\max}]$ is bounded. Then, the system has strictly CCW input-output dynamics from u to y.

Proof. System (22) can be represented as a positive loop interconnection of the x_1 and x_2 subsystems

$$\dot{x}_{1} = -\gamma_{1}x_{1} + v + u = f_{1}(x_{1}, v, u)
\dot{x}_{2} = -\gamma_{2}x_{2} + V_{2}g_{1}(x_{1})
v = V_{1}g_{2}(x_{2})
y = x_{1}.$$
(23)

Assumption 1 ensures bounded states and output for bounded input signals with $u(t) \in [u_{\min}, u_{\max}]$. Proposition III.1



Figure 2: Positive feedback interconnection scheme representing the mathematical model in (10).

guarantees that the x_2 subsystem with input x_1 and output v has CCW input-output dynamics with respect to arbitrary density functions. Furthermore, f_1 is C^1 , strictly increasing with respect to u and v and $\partial f_1/\partial u = 1 > 0$. Therefore, by Lemma III.2 in [23] we can conclude that the system in (22) has strictly CCW input-output dynamics with respect to arbitrary density functions from u to y.

By leveraging the results above, we can now conclude the proof of Theorem III.1. System (10) is a bank of SISO scalar nonlinear systems in feedback with the static map $-L: y \rightarrow z$ -Ly where L is the symmetric Laplacian matrix in (8). In Proposition III.1, we showed that each of the subsystems in (10) with input u_i and output y_i has strict CCW input-output dynamics. Furthermore, the map $-L: y \to -Ly$ is a system with CCW input-output dynamics since the Laplacian matrix Lis symmetric [24, Proposition 6.2]. Finally, by [24, Theorem 1] we can conclude that for any initial condition ξ , the ω limit set $\omega(\xi)$ is contained in the largest invariant set in M := $\{x: \dot{y} \equiv 0\}$ where $y = [x_{1,1}, \dots, x_{1,N}]^T$. Furthermore, as the forward orbits are bounded, the omega limit set $\omega(\xi)$ is non-empty, compact and connected [37, Proposition 1.111]. In M, since $\dot{y} = 0$, each component $x_{1,i}$ must be constant, and we write $x_{1,i}(t) = \bar{x}_{1,i}$. Therefore, the dynamics of each compartment is independent with respect to each other and reads

$$\dot{x}_{1,i} = -\gamma_1 \bar{x}_{1,i} + V_1 g_2(x_{2,i}) + \bar{b}_i = 0$$

$$\dot{x}_{2,i} = -\gamma_2 x_{2,i} + V_2 g_1(\bar{x}_{1,i}) = -\gamma_2 x_{2,i} + c_i,$$
(24)

where $\overline{b}_i = \sum_j a_{i,j}^1 (\overline{x}_{1,j} - \overline{x}_{1,i})$ is the constant term depending on the diffusive coupling and $c_i = V_2 g_1(\overline{x}_{1,i})$. Consider an initial condition ξ_0 in $\omega(\xi)$ and assume that ξ_0 is not an equilibrium. The invariance of $\omega(\xi)$ implies that the solution x(t) corresponding to the initial condition ξ_0 is entirely contained in $\omega(\xi)$, that is, $x(t) \in \omega(\xi), \forall t \in \mathbb{R}$. Furthermore, for the dynamics in (24), $\lim_{t \to \pm \infty} |x_{2,i}(t)| = +\infty$ against the fact that $\omega(\xi)$ is bounded. Therefore, all points in $\omega(\xi)$ must be equilibria. Furthermore, if the equilibria of the system are isolated, then for all initial conditions $\xi \in \mathcal{B}^N$, $\lim_{t \to \pm \infty} ||x(t) - \overline{x}|| \to 0$ where \overline{x} is an equilibrium of (7).

IV. SYNCHRONIZATION ANALYSIS

In the following, we derive sufficient conditions on the algebraic connectivity of the coupling graph that ensure that all the equilibria of (8) are synchronized.

Theorem IV.1. Consider a generic equilibrium $\overline{X} = [\overline{X}_1, \overline{X}_2]$ of (8). Assume that Assumption 1 holds and let ℓ_1 and ℓ_2 be the Lipschitz constants of g_1 and g_2 . Let λ_2 be the algebraic connectivity of the Laplacian matrix L. If the following condition is satisfied:

$$\frac{V_1\ell_1 V_2\ell_2}{(\gamma_1 + \lambda_2)\gamma_2} < 1,$$
(25)

then $\overline{X}_1 \in \text{span}\{\mathbb{1}_N\}$ and $\overline{X}_2 \in \text{span}\{\mathbb{1}_N\}$, where $\mathbb{1}_N \in \mathbb{R}^N$ is a vector with all entries equal to one.

Proof. For a generic equilibrium $\overline{X} = [\overline{X}_1, \overline{X}_2]$ of (8), it must hold that

$$(\Gamma_1 + L)\overline{X}_1 = \hat{G}(\overline{X}_1) \tag{26}$$

where $\tilde{G}(\overline{X}_1) := G_2(\frac{V_2}{\gamma_2}G_1(\overline{X}_1))$. If such an equilibrium exists, then

$$Q^{T}Q(\Gamma_{1}+L)\overline{X}_{1} = Q^{T}Q\tilde{G}(\overline{X}_{1}), \qquad (27)$$

where

$$Q = \begin{bmatrix} -1 + (N-1)\nu & 1-\nu & -\nu & \cdots & -\nu \\ -1 + (N-1)\nu & -\nu & 1-\nu & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & -\nu \\ -1 + (N-1)\nu & -\nu & \cdots & -\nu & 1-\nu \end{bmatrix}$$

and

$$\nu = \frac{N - \sqrt{N}}{N(N - 1)} \; .$$

From the definition of Q, it follows that $Q\mathbb{1}_N = 0$, and $Q^T Q = I_N - \frac{1}{N} \mathbb{1}_N \mathbb{1}_N^T$. Therefore, we can write

$$Q^{T}Q(\Gamma_{1}+L)\left(\overline{X}_{1}-\alpha\mathbb{1}_{N}\right)=Q^{T}Q\left(\tilde{G}(\overline{X}_{1})-\tilde{G}(\alpha\mathbb{1}_{N})\right),$$

where
$$\alpha = \frac{1}{N} \mathbb{1}_N^T X_1$$
. Assume that $X_1 \notin \operatorname{span}\{\mathbb{1}_N\}$. Then
 $\left\| Q^T Q(\Gamma_1 + L) \left(\overline{X}_1 - \alpha \mathbb{1}_N \right) \right\|_2 \ge (\gamma_1 + \lambda_2) \left\| \overline{X}_1 - \alpha \mathbb{1}_N \right\|_2$
(28)

where λ_2 is the algebraic connectivity of the Laplacian *L*. \tilde{G} has a Lipschitz constant equal to $\frac{V_1 \ell_1 V_2 \ell_2}{\gamma_2}$, and therefore

$$\left\| Q^T Q \left(\tilde{G}(\overline{X}_1) - \tilde{G}(\alpha \mathbb{1}_N) \right) \right\|_2 \le \frac{V_1 \ell_1 V_2 \ell_2}{\gamma_2} \left\| \overline{X}_1 - \alpha \mathbb{1}_N \right\|_2.$$
(29)

Since

$$\frac{V_1\ell_1V_2\ell_2}{(\gamma_1+\lambda_2)\gamma_2} < 1,$$

the inequalities in (28) and (29) lead to a contradiction, as it must hold that:

$$\left\| Q^T Q(\Gamma_1 + L) \left(\overline{X}_1 - \alpha \mathbb{1}_N \right) \right\|_2 = \left\| Q^T Q \left(\tilde{G}(\overline{X}_1) - \tilde{G}(\alpha \mathbb{1}_N) \right) \right\|_2.$$

Therefore, for all equilibria $\overline{X} = [\overline{X}_1, \overline{X}_2], \overline{X}_1 \in \operatorname{span}\{\mathbb{1}_N\}.$ Since $\overline{X}_2 = V_2/\gamma_2 G_1(\overline{X}_1)$, we conclude that $\overline{X}_2 \in \operatorname{span}\{\mathbb{1}_N\}.$

Remark. In virtue of Theorem IV.1 and Theorem III.1, we can conclude that if the algebraic connectivity of the coupling graph is sufficiently large, all the solutions of (8) asymptotically converge to synchronized equilibria, where the states of each compartment approach each other to a constant value.

V. MULTISTABILITY ANALYSIS

This section focuses on the multistability analysis of a subclass of the network models (7). Specifically, let $g_2(x) = x$ and $g_1(x) = g(x)$, a piecewise linear function defined in (4) and characterized by the positive parameters θ and δ . The resulting system reads

$$\dot{x}_{1,i} = -\gamma_1 x_{1,i} + V_1 x_2 + \sum_j a_{i,j} \left(x_{1,j} - x_{1,i} \right),$$

$$\dot{x}_{2,i} = -\gamma_2 x_{2,i} + V_2 g(x_1).$$
(30)

The dynamics of this piecewise affine model can be seen as an approximation of the dynamics of N bistable switches interconnected by a quorum sensing mechanism. Indeed, in the case of an all-to-all interconnection, it is analogous to those described in [28]–[30], in the limit that autoinducer diffusion is much more rapid than degradation. Under this assumption, intracellular and extracellular concentrations of the autoinducer molecule can be approximated to be equal, as verified in experimental studies, in different biological systems [38].

In the second part of this section, for an all-to-all homogeneous interconnection, we characterize the local stability of the equilibria of (30) and investigate their location as the system's parameters and coupling strength are varied.

The following assumption will be used throughout the remainder of this section.

Assumption 2. The system parameters of (30) satisfy

т 7

$$\frac{V_1 V_2}{\delta \gamma_1 \gamma_2} > 1 + \frac{\theta}{\delta}.$$

Assumption 2 guarantees that the uncoupled system has two stable equilibria $P_{\rm ON} = ((V_1V_2)/(\gamma_2\gamma_1), V_2/\gamma_2), P_{\rm OFF} = (0,0)$ and an unstable equilibrium $P_{\rm s} = (\bar{x}_1, \bar{x}_2)$ with

$$\bar{x}_1 = \frac{V_1}{\gamma_1} \bar{x}_2$$
$$\bar{x}_2 = \frac{V_2 \theta}{\delta \gamma_2} \left(\frac{V_1 V_2}{\delta \gamma_1 \gamma_2} - 1 \right)^{-1}$$

It is useful to partition the phase space \mathcal{B}^N into different *domains* such that the restriction of the vector field is linear-affine in each of them.

Definition V.1. Let $\mathcal{T} \subset \mathbb{R}^N$ denote the set of N dimensional vectors with entries in $\{-1, 0, +1\}$. We can associate each element α in \mathcal{T} with a *domain* Ω_{α} defined as

$$\Omega_{\alpha} := \{ x \in \mathcal{B}^{N} \mid x_{1,j} > \theta + \delta, \text{ if } \alpha_{j} = 1 \\ x_{1,j} < \theta, \text{ if } \alpha_{j} = -1 \\ \theta \le x_{1,j} \le \theta + \delta, \text{ if } \alpha_{j} = 0 \}$$

A domain Ω_{α} is called *saturated* if $\alpha \in \Lambda_e := \{\alpha \in \mathcal{T} : \alpha_j \in \{-1,1\}\}$, *linear* if $\alpha \in \Lambda_0 := \{\alpha \in \mathcal{T} : \alpha_j = 0\}$, and *mixed* if $\alpha \in \Lambda_m := \mathcal{T}/(\Lambda_0 \cup \Lambda_e)$.

Definition V.2. Given a saturated domain Ω_{α} , we say that Ω_{α} has an average level of activation $\overline{q} := m/N, m = 0, \dots, N$ where *m* is the number of entries in α equal to 1.

In the following, we characterize the stability of equilibria in the different types of domains. Similarly to (8), we define the state vector $x = [X_1^T, X_2^T]^T$ such that the restriction of the vector field in (30) to a domain Ω_{α} is described by the linear-affine system

 $M_{\alpha} = \begin{bmatrix} M_{11}(\alpha) & M_{12}(\alpha) \\ M_{21}(\alpha) & M_{22}(\alpha) \end{bmatrix}$

$$\dot{x} = M_{\alpha}x + b_{\alpha},\tag{31}$$

(32)

where

with

$$M_{11}(\alpha) = -(\Gamma_1 + L)$$

$$M_{12}(\alpha) = V_1 I_N$$

$$M_{21}(\alpha) = V_2/\delta \operatorname{diag}(w(\alpha_1), \dots, w(\alpha_N))$$

$$M_{22}(\alpha) = -\Gamma_2$$
(33)

and $w(\alpha_i) = 1$ if $\alpha_i = 0$ and $w(\alpha_i) = 0$, otherwise. We skip here the explicit expression of b_{α} since it is not involved in the subsequent calculations. The following proposition characterizes the local stability of the equilibria of (30).

Proposition V.1. Consider system (30). If an equilibrium exists in a saturated domain, it is unique and locally asymptotically stable. Furthermore, if $\frac{V_1V_2}{\gamma_1\gamma_2\delta} > N$, the equilibria in the interior of non-saturated domains are unstable.

Proof. For all saturated domains, $M_{21}(\alpha) = 0$, therefore the matrix M_{α} is block diagonal, nonsingular and Hurwitz. Since the vector field is differentiable in each saturated domain, if an equilibrium exists in a saturated domain Ω_{α} , it is unique and locally asymptotically stable. We now prove that the equilibria in the interior of mixed and linear domains are unstable if $\frac{V_1V_2}{\gamma_1\gamma_2\delta} > N$. Let Ω_{α} be a mixed or linear domain, we claim that the matrix M_{α} has at least one eigenvalue with strictly positive real part. We proceed by contradiction. Assume that M_{α} has no eigenvalues with strictly positive real part. Then,

$$M_{\alpha}^{\varepsilon} := \begin{bmatrix} M_{11}(\alpha) - \varepsilon I_N & M_{12}(\alpha) \\ M_{21}(\alpha) & M_{22}(\alpha) - \varepsilon I_N \end{bmatrix}$$
(34)

must be Hurwitz, $\forall \varepsilon > 0$. Since M_{α}^{ε} is Metzler, it is Hurwitz if and only if both $M_{22}(\alpha) - \varepsilon I_N$ and the Schur complement $M_{\alpha}^{\varepsilon}/(M_{22}(\alpha) - \varepsilon I_N)$ are Hurwitz [39, Corollary 1]. Since $M_{22}(\alpha) - \varepsilon I_N$ is clearly Hurwitz, M^{ε}_{α} is Hurwitz if and only if

$$M_{\alpha}^{\varepsilon}/(M_{22}(\alpha) - \varepsilon I_N) = M_{11}(\alpha) - \varepsilon I_N + \frac{V_1}{(\gamma_2 + \varepsilon)} M_{21}(\alpha)$$

is Hurwitz. Notice that $M^{\varepsilon}_{\alpha}/(M_{22}(\alpha) - \varepsilon I_N)$ is symmetric and

$$\mathbb{1}_{N}^{T} M_{\alpha}^{\varepsilon} / (M_{22}(\alpha) - \varepsilon I_{N}) \mathbb{1}_{N} = -N(\gamma_{1} + \varepsilon)\gamma_{1} + \frac{V_{2}V_{1}}{(\gamma_{2} + \varepsilon)\delta} n_{\alpha}$$

with $\mathbb{1}_N$ the vector whose entries are all equal to 1 and $n_\alpha \in \{1, ..., N\}$ depends on the domain Ω_α and represents the number of species in the linear regime. By assumption, $\frac{V_1 V_2}{\gamma_1 \gamma_2 \delta} > N$, therefore there exists (small enough) $\varepsilon > 0$ such that $\forall n_\alpha \in \{1, ..., N\}$

$$\mathbb{1}_N^T M_\alpha^\varepsilon / (M_{22}(\alpha) - \varepsilon I_N) \mathbb{1}_N > 0.$$

Therefore, for small enough $\varepsilon > 0$ M^{ε}_{α} is not Hurwitz, thus proving the claim.

In the following, we study the location of (locally asymptotically stable) equilibria in the saturated domains as a function of the coupling strength k for the case of a homogeneous allto-all network, i.e. $L = k (NI_N - \mathbb{1}_N \mathbb{1}_N^T)$.

Notice that in the two saturated domains with average level of activation $\overline{q} = 0$ and $\overline{q} = 1$, the equilibria exist and are unique, each corresponding to all compartments synchronized in the state P_{OFF} and P_{ON} . In the rest of the paper, the following definition will be used to refer to a specific class of equilibria:

Definition V.3. A generic equilibrium $\overline{X} = [\overline{X}_1, \overline{X}_2]$ of (8) is said to be synchronized if $\overline{X}_1 \in \operatorname{span}\{\mathbb{1}_N\}$ and $\overline{X}_2 \in \operatorname{span}\{\mathbb{1}_N\}$.

Theorem V.1. Consider system (30) and assume $L = k \left(NI_N - \mathbb{1}_N \mathbb{1}_N^T \right)$. Let Ω_{α} be a saturated domain with an average level of activation $\overline{q} = m/N$, $m \in \{1, \dots, N-1\}$. There exists a minimum gain $k^{\overline{q}}$ such that $\forall k > k^{\overline{q}}$, no equilibria are contained in the domain Ω_{α} . Additionally, there exists $k^s > 0$ such that for every $k > k^s$, if an equilibrium exists in a saturated domain, it is synchronized.

Proof. A generic equilibrium $\overline{X} = [\overline{X}_1, \overline{X}_2]$ of (8) must satisfy

$$\overline{X}_1 = V_1(\Gamma_1 + L)^{-1}\overline{X}_2$$

$$\overline{X}_2 = V_2/\gamma_2 G_1(\overline{X}_1).$$
(35)

As $\Gamma_1 + L = k (NI_N - \mathbb{1}\mathbb{1}^T) + \gamma_1 I_N$, its inverse (Sherman-Morrison formula [40], [41]) is

$$(\Gamma_1 + L)^{-1} = \frac{1}{Nk + \gamma_1} I_N + \frac{1}{\gamma_1 (Nk + \gamma_1)} k \mathbb{1}_N \mathbb{1}_N^{\mathrm{T}}$$
(36)

and by substituting it in (35), we obtain

$$\overline{X}_{1} = V_{1} \left(\frac{1}{Nk + \gamma_{1}} I_{N} + \frac{1}{\gamma_{1} \left(Nk + \gamma_{1} \right)} k \mathbb{1}_{N} \mathbb{1}_{N}^{\mathrm{T}} \right) \frac{V_{2}}{\gamma_{2}} G_{1}(\overline{X}_{1})$$
(37)

Therefore, for a generic compartment i,

$$\overline{x}_{1,i} = \frac{V_1 V_2 / \gamma_2}{Nk + \gamma_1} g(\overline{x}_{1,i}) + \frac{V_1 V_2 / \gamma_2}{\gamma_1 (Nk + \gamma_1)} Nk \sum_{i=1}^N g(\overline{x}_{1,i}) / N$$

$$\overline{x}_{2,i} = V_2 / \gamma_2 g_1(\overline{x}_{1,i}).$$
(38)

Define $I_{\text{ON}}(\alpha) = \{i : \alpha_i = 1\}$ and $I_{\text{OFF}}(\alpha) = \{i : \alpha_i = -1\}$. As the domain Ω_{α} has an average level of activation $\overline{q} = m/N$, this means that $I_{\text{ON}}(\alpha)$ and $I_{\text{OFF}}(\alpha)$ have cardinality m, and N - m, respectively. Denote with $V := \frac{V_1 V_2}{\gamma_1 \gamma_2}$, the equilibrium $\overline{X} = [\overline{X}_1, \overline{X}_2] \in \Omega_{\alpha}$ if and only if

$$\overline{x}_{1,i} = \gamma_1 \frac{V}{Nk + \gamma_1} + \frac{V}{Nk + \gamma_1} Nk\overline{q} > \theta + \delta, \ i \in I_{\text{ON}}(\alpha)$$
(39)

and

$$\overline{x}_{1,i} = \frac{V}{Nk + \gamma_1} N k \overline{q} < \theta, \ i \in I_{\text{OFF}}(\alpha).$$
(40)

Therefore, $\overline{X} = [\overline{X}_1, \overline{X}_2] \notin \Omega_{\alpha}$ if at least one of (39) and (40) is not verified, i.e. if

$$\gamma_1 \frac{V}{Nk + \gamma_1} + \frac{V}{Nk + \gamma_1} Nk\overline{q} \le \theta + \delta \tag{41}$$

or

$$\frac{V}{Nk + \gamma_1} Nk\overline{q} \ge \theta.$$
(42)

We can now rewrite (41), (42) as

$$kN\left(-V\overline{q} + (\theta + \delta)\right) \ge \gamma_1\left(V - (\theta + \delta)\right) \tag{43}$$

and

$$(V\overline{q} - \theta) Nk \ge \theta \gamma_1. \tag{44}$$

The factor on the right handside of (43) is always strictly positive under Assumption 2. We now show that for any given \overline{q} , it is possible to find a minimum gain $k^{\overline{q}}$ such that for all $k > k^{\overline{q}}$, no equilibria exist in the domain Ω_{α} with an average level of activation \overline{q} . We divide the analysis according to the value of \overline{q} .

1) If \overline{q} is such that $\theta < V\overline{q} < (\theta + \delta)$, by simple manipulation of (43) and (44) we obtain:

$$k \ge k_1^{\overline{q}} := \frac{\gamma_1 \left(\left(\theta + \delta \right) - V \right)}{N \left(V \overline{q} - \left(\theta + \delta \right) \right)} \tag{45}$$

or

$$k \ge k_2^{\overline{q}} := \frac{\theta \gamma_1}{N \left(V \overline{q} - \theta \right)}.$$
(46)

Therefore, $\forall k \geq k^{\overline{q}} = \min\{k_1^{\overline{q}}, k_2^{\overline{q}}\}$, there are no equilibria in the saturated domains corresponding to the *average level of activation* \overline{q} .

2) If \overline{q} is such that $V\overline{q} \ge (\theta + \delta)$, no positive value of k can satisfy (43). From (44) we can instead get the bound

$$k \ge k_2^{\overline{q}} := \frac{\theta \gamma_1}{N \left(V \overline{q} - \theta \right)}.$$
(47)

Therefore, $\forall k \geq k_2^{\overline{q}}$ there are no equilibria in the saturated domains corresponding to the *average level of activation* \overline{q} .

$$k \ge k_1^{\overline{q}} := \frac{\gamma_1 \left((\theta + \delta) - V \right)}{N \left(V \overline{q} - (\theta + \delta) \right)}.$$
(48)

Therefore, $\forall k \geq k_1^{\overline{q}}$ there are no equilibria in the saturated domains corresponding to the *average level of activation* \overline{q} . Finally, for all

$$k > k^{\rm s} := \max_{\bar{q} \in \{m/N \mid m \in \{1, \dots, N-1\}\}} k^{\bar{q}}, \tag{49}$$

the only saturated domain with equilibria are the two saturated domains corresponding to the average level of activation $\overline{q} = 0$ and $\overline{q} = 1$.

VI. NUMERICAL RESULTS

To illustrate the results developed in the previous sections, we consider a network (30) of N = 5 identical bistable compartments with $g_2(x_2) = x_2$, and $g_1(x_1) = g(x_1)$ defined as a piecewise function (4) with parameters $V_1 = V_2 = 1$, $\gamma_1 = \gamma_2 = 1, \ \theta = 0.45, \ \delta = 0.1.$ The chosen parameters and activation functions guarantee that all the solutions converge to the set of equilibria (Theorem III.1) and that all equilibria in the interior of non-saturated domains are unstable (Proposition V.1). Figure 3 (a)-(d) illustrates the number of equilibria (both in the saturated and unsaturated domains) as a function of the homogeneous coupling k for various interconnection topologies. The minimum coupling $k^{\lambda} = \left(\frac{V_1 V_2}{\gamma_2 \delta} - \gamma_1\right)/N$ (red dotted line), computed using Theorem IV.1, represents the minimum coupling gain that ensures global synchronization. This is illustrated in Figure 3, as for $k > k^{\lambda}$, the network has only three synchronized equilibria, corresponding to the three equilibria of the uncoupled bistable system.

In Figure 3 (a) we depict the exact thresholds $k^{\overline{q}}$ (dotted blue lines) computed by using Theorem V.1 in the case where the communication topology is homogeneous and all-to-all. These thresholds represent the coupling strength required to eliminate all the stable equilibria with an average level of activation \overline{q} . In particular, we observe a gradual transition from multistability to bistability of the overall network. Indeed, for all $k > k^{s}$, all stable equilibria are synchronized equilibria (Theorem V.1). The figure also suggests that the bound k^{s} guarantees convergence to the synchronized equilibria almost everywhere. It is worth commenting on the difference between k^{λ} and k^{s} . The bound k^{λ} guarantees that only the three synchronized equilibria exist while k^{s} guarantees that the only locally asymptotically stable equilibria are the synchronized ones.

The gap between the bounds k^{λ} and k^{s} provides a possible explanation on why, in many examples (e.g. the synchronization of Goodwin oscillators in [42]), the minimum coupling gain that guarantees synchronization is higher than the values empirically found in simulation.

This observation is particularly insightful when we consider the limiting case of $\delta \rightarrow 0$, i.e. when the piecewise affine continuous function (4) tends to a discontinuous function. In this case, the minimum coupling gain that guarantees global synchronization computed with (25) grows unbounded with the Lipschitz constant $\ell_1 = 1/\delta$ of the function g_1 . Nevertheless, the minimum coupling gain computed in Theorem V.1 approaches a finite value (Figure 4).

In our final experiments, we compare two different choices for the regulatory functions : piecewise affine approximations (4) and hill-like smooth functions (2). We evaluate how closely these two modelling choices agree in the characterization of clustering and synchronization properties of the networked system. According to [31], these models should produce similar results when Hill coefficients are sufficiently large, with the exact threshold on the cooperativity degree depending on the specific parameters used.

We consider two variants of a network (7) with N = 3 identical bistable compartments. In both variants, $g_2(x_2) = x_2$. For $g_1(x_1)$, we compare:

- A Hill function (2) g₁(x₁) = g_s(x₁) with θ_H = 1.5 and cooperative degree n = 3;
- A piecewise affine activation function (4) g₁(x₁) = g(x₁) with parameters δ = 2 and θ = θ_H − δ/2 = 0.5.

The parameters for the two regulatory functions are chosen such that $g_s(x)|_{x=\theta_H} = g(x)|_{x=\theta_H}$ and $\frac{d}{dx}g_s(x)|_{x=\theta_H} = \frac{d}{dx}g(x)|_{x=\theta_H}$ as shown in Figure 5. The comparison of the number of equilibria as a function of the coupling gain k for the two model variants in Figure 6 indicates that when the Hill function is employed, non-synchronized equilibria are eliminated at slightly smaller values of the coupling gain k. The number of equilibria in the Hill function model is more sensitive to changes in the coupling gain k. Despite this difference, both variants of the models exhibit a similar overall behavior. This suggests that the methodology developed in this paper provides an alternative to the, computationally heavy, numerical computations to locate the equilibria in the smooth system.

VII. CONCLUSION

In this paper, we examined the dynamics of a network of bistable systems coupled by diffusion. Under assumptions, we presented structural conditions (i.e., conditions that do not depend on the values of the system's parameters) that ensure all solutions converge to the set of equilibria. Additionally, under the same technical assumptions, we provided sufficient conditions for the coupling graph to guarantee the bistability of the network and demonstrated that all solutions asymptotically converge to one of the equilibria of the uncoupled system.

For a biologically relevant model of a network of bistable systems, we utilized a piecewise linear approximation of the vector field to determine the location of equilibria as a function of the coupling gain and studied their local stability. This approach provided valuable insights into the impact of diffusive coupling on the stability and synchronization of bistable systems.

Future research directions include extending our results to more general bistable and multistable systems and incorporating other coupling mechanisms such as quorum sensing. Exploring these additional coupling mechanisms could further enhance our understanding of complex network dynamics in biological systems and expand the applicability of our findings to a broader range of real-world scenarios.



(c) Loop topology



Figure 3: In (a), the dotted vertical blue lines correspond to the values $k^{\overline{q}}$ computed analytically. The continuous blue line represents the total number of equilibria in all saturated domains as a function of the coupling parameter k. The continuous orange line represents the total number of equilibria in all domains (including non-saturated) as a function of the coupling parameter k. The dotted red line represents the minimum coupling gain k^{λ} to ensure that only synchronized equilibria exist. In all figures, the minimum values for the blue and orange lines are respectively 2 and 3. The system's parameters used are are $V_1 = V_2 = 1$, $\gamma_1 = \gamma_2 = 1$ and $\theta = 0.45$, $\delta = 0.1$.

APPENDIX

A. Systems with Counterclockwise Input-Output Dynamics

The following is from [23]. Consider the nonlinear differential equation with input u and output y

$$\dot{x} = f(x, u), \ y = h(x),$$
 (50)

defined on the closed state space $X \subset \mathbb{R}^n$, where $f: X \times U \to \mathbb{R}^n$ is a locally Lipschitz function, with input signal $u \in \mathcal{U}$ (the set of $U \subset \mathbb{R}^m$ -valued Lebesgue measurable locally essentially bounded functions) and the output map $h: X \to Y \subset \mathbb{R}^m$ is locally Lipschitz. The input-output transition map $\psi(t,\xi,u)$ can then be defined, for each initial condition $\xi \in X$ and each input signal $u \in \mathcal{U}$, by

$$\psi(t,\xi,u) = h(x(t,\xi,u)). \tag{51}$$

This map relates the input signal u and the initial state ξ to the output of the system at time t. Static input-output maps can be defined as

$$\psi(t, u) = h(u(t)), \tag{52}$$

for some Lipschitz function $h: U \to Y$. The transition map $\psi(t, \xi, u)$ need not be defined for all $t \ge 0$; however, for each $\xi \in X$ and each $u \in \mathcal{U}$ there exists $T_{\xi,u} \in (0, +\infty]$ so that $\psi(t, \xi, u)$ is well defined for all $t \in [0, T_{\xi,u})$. It is useful to define the following subsets of $X \times \mathcal{U}$:

$$\mathcal{S}_{\text{fc}} \doteq \{ (\xi, u) \in X \times \mathcal{U} : T_{\xi, u} = +\infty \}$$
$$\mathcal{S}_{bd} \doteq \{ (\xi, u) \in X \times \mathcal{U} : T_{\xi, u} = +\infty \text{ and } u(\cdot), \psi(\cdot, \xi, u) \text{ are bounded } \}$$



Figure 4: Synchronization bounds k^{λ} (red line) and k^{s} (blue line) for increasing values of the Lipschitz constant $\ell_{2} = 1/\delta$ of the piecewise affine activation function $g_{1}(x_{1}) = g(x_{1})$. Parameters used: N = 5, $V_{1} = V_{2} = 1$, $\gamma_{1} = \gamma_{2} = 1$ and $\theta = 0.45$.



Figure 5: Hill function $g_s(x)$ with cooperativity degree n = 3 (orange line) and piece-wise affine approximation g(x) (blue line). Parameters used: $\theta_H = 1.5$, $\delta = 2$, $\theta = \theta_H - \delta/2$.

We assume, without loss of generality that $0 \in U$, and that $U = U_1 \times U_2 \cdots \times U_m$ and $Y = Y_1 \times Y_2 \cdots \times Y_m$ for some nonempty intervals $U_i, Y_i \subset \mathbb{R}$. We are now ready to report some definitions.

Definition A.1. [23] We say that $\rho : U \times Y \to \mathbb{R}^m$ is a density function if it satisfies the following properties: 1) $\rho(u, y) = [\rho_1(u_1, y_1), \rho_2(u_2, y_2), \dots, \rho_m(u_m, y_m)]$ for scalar functions $\rho_i : U_i \times Y_i \to \mathbb{R}$; 2) $\rho_i(u_i, y_i) > 0$ for almost all $(u_i, y_i) \in U_i \times Y_i$ (according to Lebesgue measure) and all *i* in $\{1 \dots m\}$; 3) ρ is a measurable and locally summable function (jointly in *u* and *y*).

Definition A.2. [23] We say that a system has CCW inputoutput (CCW input-output) dynamics with respect to the density function $\rho(u, y)$ if for any $(\xi, u) \in S_{bd}$ the following inequality holds

$$\liminf_{T \to +\infty} \int_0^T \dot{y}(t)' \int_0^{u(t)} \rho(\mu, y(t)) d\mu dt > -\infty,$$

where $y(t) = \psi(t, \xi, u)$ is assumed to be absolutely continu-



Figure 6: Number of equilibria for different values of k in two variants of a all-to all network of (7) with N = 3bistable compartments with $g_1(x_1) = g_s(x_1)$ (blue line) and $g_1(x_1) = g(x_1)$ (red line). For both lines, the minimum number of equilibria is 3. For both variants $g_2(x_2) = x_2$. Parameters used: $\theta_H = 1.5$, $\delta = 2$, $\theta = \theta_H - \delta/2$, $\gamma_1 = \gamma_2 = 1$, $V_1 = V_2 = 3$.

ous.

Definition A.3. [23] We say that a system has a strict counterclockwise input-output dynamics with respect to the density function $\rho(u, y) : U \times Y \to \mathbb{R}_{\geq 0}$ if the following inequality holds for all pairs $(\xi, u) \in S_{bd}$

$$\liminf_{T \to +\infty} \int_0^T \dot{y}(t)' \int_0^{u(t)} \rho(\mu, y(t)) d\mu - \frac{\tilde{\rho}(|\dot{y}(t)|)}{1 + \gamma(|x(t)|)} dt > -\infty,$$

where $\tilde{\rho}$ is a positive definite function, $\gamma \in \mathcal{K}$ and $y(t) = \psi(t, \xi, u)$ is absolutely continuous.

Remark. The assumption of y(t) being absolutely continuous is a fundamental one as it ensures that the time derivative $\dot{y}(t)$ is defined almost everywhere. Such a property is always guaranteed under the assumption that the output map h is Lipschitz.

B. Generalized inverse

The following definition of generalized inverse of a monotone function is taken from [43].

Definition A.4. [43] Let f be a monotonically increasing function. The generalized inverse is function f^{-1} defined by

$$f^{-1}(y) = \inf\{x \in \overline{\mathbb{R}} : f(x) > y\}, \ \overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\}$$
(53)

In the following we will use the following adapted definition (no need to consider extended $\overline{\mathbb{R}}$).

Definition A.5. Let f be a monotonically increasing function defined on the interval [a, b]. The generalized inverse is function f^{-1} defined by

$$f^{-1}(y) = \inf\{x \in [a,b] : f(x) > y\}, \ y \in (f(a), f(b))$$
(54)

The following proposition holds:

Proposition A.1. [43] The pseudo-inverse f^{-1} of a monotonically increasing function f has following properties:

- 1) f^{-1} is increasing, has left limits and is right continuous.
- 2) The following implications hold for all $x \in \mathbb{R}$ and $y \in \mathbb{R}$:

$$f(x) > y \implies x \ge f^{-1}(y), \tag{55}$$

$$f(x) = y \implies x \le f^{-1}(y), \tag{56}$$

$$f(x) < y \implies x \le f^{-1}(y), \tag{57}$$

$$f^{-1}(y) > x \implies y \ge f(x), \tag{58}$$

$$f^{-1}(y) < x \implies y < f(x).$$
(59)

3) For all $x \in \mathbb{R}$, $f^{-1}(f(x)) \ge x$.

J

4) If f is right continuous at x, then for all $y \in \mathbb{R}$

$$f^{-1}(y) = x \implies y \le f(x), \tag{60}$$

$$f(x) > y \implies x > f^{-1}(y), \tag{61}$$

$$f(f^{-1}(y)) \ge y.$$
 (62)

- 5) f is strictly increasing on \mathbb{R} if, and only if, f^{-1} is continuous on \mathbb{R} .
- 6) f is continuous on \mathbb{R} if, and only if, f^{-1} is strictly increasing on \mathbb{R} .
- 7) f is right-continuous if, and only if, $(f^{-1})^{-1} = f$.
- 8) f is right-continuous if, and only if, $\{x \in \mathbb{R} : f(x) \ge y\}$ is closed for all $y \in \mathbb{R}$.

REFERENCES

- F. Attneave, "Multistability in perception," *Scientific American*, vol. 225, no. 6, pp. 62–71, 1971.
- [2] U. Feudel, "Complex dynamics in multistable systems," *International Journal of Bifurcation and Chaos*, vol. 18, no. 06, pp. 1607–1626, 2008.
- [3] A. N. Pisarchik and U. Feudel, "Control of multistability," *Physics Reports*, vol. 540, no. 4, pp. 167–218, 2014.
- [4] C. H. Waddington, The strategy of the genes. Routledge, 2014.
- [5] R. Shah and D. Del Vecchio, "Reprogramming multistable monotone systems with application to cell fate control," *IEEE transactions on network science and engineering*, vol. 7, no. 4, pp. 2940–2951, 2020.
- [6] R. Thomas, "On the relation between the logical structure of systems and their ability to generate multiple steady states or sustained oscillations," in *Numerical methods in the study of critical phenomena*, pp. 180–193, Springer, 1981.
- [7] E. H. Snoussi, "Necessary conditions for multistationarity and stable periodicity," *Journal of Biological Systems*, vol. 6, no. 01, pp. 3–9, 1998.
- [8] T. S. Gardner, C. R. Cantor, and J. J. Collins, "Construction of a genetic toggle switch in escherichia coli," *Nature*, vol. 403, no. 6767, pp. 339– 342, 2000.
- [9] E. V. Nikolaev and E. D. Sontag, "Quorum-sensing synchronization of synthetic toggle switches: A design based on monotone dynamical systems theory," *PLoS computational biology*, vol. 12, no. 4, p. e1004881, 2016.
- [10] M. Ali Al-Radhawi, D. Del Vecchio, and E. D. Sontag, "Multi-modality in gene regulatory networks with slow promoter kinetics," *PLoS computational biology*, vol. 15, no. 2, p. e1006784, 2019.
- [11] N. Augier, M. Chaves, and J.-L. Gouzé, "Qualitative control strategies for synchronization of bistable gene regulatory networks," *IEEE Transactions on Automatic Control*, 2022.
- [12] M. Chaves, L. Scardovi, and E. Firippi, "Coupling and synchronization of piecewise linear genetic regulatory systems," in 2019 IEEE 58th Conference on Decision and Control (CDC), pp. 2826–2831, IEEE, 2019.
- [13] G. Villani and L. Scardovi, "Global analysis of networks of piecewise affine bistable switches," in 2021 60th IEEE Conference on Decision and Control (CDC), pp. 5544–5549, IEEE, 2021.

- [14] H. L. Smith, Monotone dynamical systems: an introduction to the theory of competitive and cooperative systems: an introduction to the theory of competitive and cooperative systems. No. 41, American Mathematical Soc., 2008.
- [15] L. O. Chua and L. Yang, "Cellular neural networks: Theory," *IEEE Transactions on circuits and systems*, vol. 35, no. 10, pp. 1257–1272, 1988.
- [16] S.-S. Lin and C.-W. Shih, "Complete stability for standard cellular neural networks," *International Journal of Bifurcation and Chaos*, vol. 9, no. 05, pp. 909–918, 1999.
- [17] J. Smillie, "Competitive and cooperative tridiagonal systems of differential equations," *SIAM journal on mathematical analysis*, vol. 15, no. 3, pp. 530–534, 1984.
- [18] M. Pasquini and D. Angeli, "On convergence for piecewise affine models of gene regulatory networks via a lyapunov approach," *IEEE Transactions on Automatic Control*, vol. 65, no. 8, pp. 3333–3348, 2019.
- [19] M. Forti and A. Tesi, "A new method to analyze complete stability of pwl cellular neural networks," *International Journal of Bifurcation and Chaos*, vol. 11, no. 03, pp. 655–676, 2001.
- [20] J. S. Muldowney, "Compound matrices and ordinary differential equations," *The Rocky Mountain Journal of Mathematics*, pp. 857–872, 1990.
- [21] C. Wu, I. Kanevskiy, and M. Margaliot, "k-contraction: Theory and applications," *Automatica*, vol. 136, p. 110048, 2022.
- [22] D. Angeli, "Systems with counterclockwise input-output dynamics," *IEEE Transactions on automatic control*, vol. 51, no. 7, pp. 1130–1143, 2006.
- [23] D. Angeli, "Multistability in systems with counter-clockwise inputoutput dynamics," *IEEE Transactions on automatic control*, vol. 52, no. 4, pp. 596–609, 2007.
- [24] D. Angeli, "Convergence in networks with counterclockwise neural dynamics," *IEEE transactions on neural networks*, vol. 20, no. 5, pp. 794–804, 2009.
- [25] A. G. Ghallab, M. A. Mabrok, and I. R. Petersen, "Extending negative imaginary systems theory to nonlinear systems," in 2018 IEEE Conference on Decision and Control (CDC), pp. 2348–2353, IEEE, 2018.
- [26] A. G. Ghallab and I. R. Petersen, "Negative imaginary systems theory for nonlinear systems: A dissipativity approach," arXiv preprint arXiv:2201.00144, 2022.
- [27] U. Alon, An introduction to systems biology: design principles of biological circuits. Chapman and Hall/CRC, 2006.
- [28] N. Rai, R. Anand, K. Ramkumar, V. Sreenivasan, S. Dabholkar, K. V. Venkatesh, and M. Thattai, "Prediction by promoter logic in bacterial quorum sensing," *PLoS computational biology*, vol. 8, no. 1, p. e1002361, 2012.
- [29] K. Fujimoto and S. Sawai, "A design principle of group-level decision making in cell populations," *PLoS Computational Biology*, vol. 9, no. 6, p. e1003110, 2013.
- [30] M. Schuster, C. Li, P. Smith, and C. Kuttler, "Parameters, architecture and emergent properties of the pseudomonas aeruginosa lasi/lasr quorum-sensing circuit," *Journal of the Royal Society Interface*, vol. 20, no. 200, p. 20220825, 2023.
- [31] A. Polynikis, S. Hogan, and M. di Bernardo, "Comparing different ode modelling approaches for gene regulatory networks," *Journal of Theoretical Biology*, vol. 261, no. 4, pp. 511–530, 2009.
- [32] F. Blanchini, S. Miani, et al., Set-theoretic methods in control, vol. 78. Springer, 2008.
- [33] M. B. Miller and B. L. Bassler, "Quorum sensing in bacteria," Annual Reviews in Microbiology, vol. 55, no. 1, pp. 165–199, 2001.
- [34] A. Boo, R. L. Amaro, and G.-B. Stan, "Quorum sensing in synthetic biology: A review," *Current Opinion in Systems Biology*, vol. 28, p. 100378, 2021.
- [35] R. Silver, J. LeSauter, P. A. Tresco, and M. N. Lehman, "A diffusible coupling signal from the transplanted suprachiasmatic nucleus controlling circadian locomotor rhythms," *Nature*, vol. 382, no. 6594, pp. 810– 813, 1996.
- [36] C. W. Wu, "Algebraic connectivity of directed graphs," *Linear and multilinear algebra*, vol. 53, no. 3, pp. 203–223, 2005.
- [37] C. Chicone, Ordinary differential equations with applications, vol. 34. Springer Science & Business Media, 2006.
- [38] H. B. Kaplan and E. Greenberg, "Diffusion of autoinducer is involved in regulation of the vibrio fischeri luminescence system," *Journal of bacteriology*, vol. 163, no. 3, pp. 1210–1214, 1985.
- [39] M. Souza, F. R. Wirth, and R. N. Shorten, "A note on recursive schur complements, block hurwitz stability of metzler matrices, and related results," *IEEE Transactions on Automatic Control*, vol. 62, no. 8, pp. 4167–4172, 2017.

- [40] J. Sherman and W. J. Morrison, "Adjustment of an inverse matrix corresponding to a change in one element of a given matrix," *The Annals* of *Mathematical Statistics*, vol. 21, no. 1, pp. 124–127, 1950.
- [41] A. W. Max, "Inverting modified matrices," in *Memorandum Rept.* 42, *Statistical Research Group*, p. 4, Princeton Univ., 1950.
- [42] L. Scardovi, M. Arcak, and E. D. Sontag, "Synchronization of interconnected systems with applications to biochemical networks: An inputoutput approach," *IEEE transactions on Automatic Control*, vol. 55, no. 6, pp. 1367–1379, 2010.
- [43] A. de La Fortelle, "A study on generalized inverses and increasing functions part i: generalized inverses," 2015.