

Resummation of threshold double logarithms in hadroproduction of heavy quarkonium

Hee Sok Chung,^{1,*} U-Rae Kim,^{2,†} and Jungil Lee^{1,‡}

¹*Department of Physics, Korea University, Seoul 02841, Korea*

²*Department of Physics, Korea Military Academy, Seoul 01805, Korea*

(Dated: January 20, 2025)

We resum threshold double logarithms that appear in inclusive production of heavy quarkonium. This resolves the catastrophic failure of fixed-order perturbation theory where quarkonium cross sections at large transverse momentum can turn negative due to large radiative corrections. We find that resummation is imperative for describing measured prompt production rates of J/ψ at large transverse momentum.

Introduction.— Recent measurements of charmonium production rates at the LHC at very large transverse momentum [1] posed a serious challenge to the phenomenology of heavy quarkonium production based on the nonrelativistic QCD (NRQCD) factorization formalism [2]. This formalism describes the inclusive cross section of a heavy quarkonium in terms of sums of products of perturbatively calculable short-distance coefficients (SDCs) and nonperturbative long-distance matrix elements (LDMEs). Predictions based on SDCs computed at next-to-leading order (NLO) accuracy in fixed-order (FO) perturbation theory fail to describe the measured J/ψ production rates at the LHC for transverse momentum much larger than 100 GeV. Moreover, predictions based on FO calculations of SDCs can yield unphysical, negative cross sections at large transverse momentum [3, 4]. It has been suggested that this problem arises from threshold logarithms, which originate from singularities in radiative corrections near boundaries of phase space [3, 4]. Hence, the catastrophic failure of FO perturbation theory may be resolved by resumming the threshold logarithms to all orders in perturbation theory, which have not yet been done in the NRQCD factorization formalism for production of heavy quarkonium.

In this Letter, we resum the threshold logarithms in the NRQCD factorization formula for production of J/ψ , $\psi(2S)$, and χ_{cJ} to all orders in perturbation theory. We work at the leading double logarithmic level, which produces the dominant singularities in the SDCs. We focus on the large-transverse momentum region, where the effect of resummation is most significant. By using the Grammer-Yennie approximation [5] we obtain expressions for the singularities in the SDCs in terms of soft functions, which make possible the resummation of threshold logarithms by exponentiation. The resummed SDCs that we obtain are free of singularities, so that we can ensure the positivity of heavy quarkonium cross sections at large p_T . We then compute J/ψ production rates at large p_T from pp collisions at $\sqrt{s} = 13$ TeV by using our resummed SDCs, and find that we obtain substantially improved descriptions of LHC data compared to FO calculations.

NRQCD factorization and fragmentation.—In the

NRQCD factorization formalism, the inclusive cross section of a heavy quarkonium \mathcal{Q} at large transverse momentum p_T is given by [2]

$$\sigma_{\mathcal{Q}} = \sum_{\mathcal{N}} \sigma_{Q\bar{Q}(\mathcal{N})} \langle \mathcal{O}^{\mathcal{Q}}(\mathcal{N}) \rangle, \quad (1)$$

where $\sigma_{Q\bar{Q}(\mathcal{N})}$ is the SDC that corresponds to the production rate of a heavy quark (Q) and a heavy antiquark (\bar{Q}) pair in a specific color and angular momentum state \mathcal{N} , and $\langle \mathcal{O}^{\mathcal{Q}}(\mathcal{N}) \rangle$ is the LDME that represents the probability for a $Q\bar{Q}$ in the state \mathcal{N} to evolve into a quarkonium \mathcal{Q} . The sum over \mathcal{N} can be truncated at a chosen accuracy in the nonrelativistic expansion; for $\mathcal{Q} = J/\psi$ or $\psi(2S)$, dominant contributions come from ${}^3S_1^{[1]}$, ${}^3S_1^{[8]}$, ${}^1S_0^{[8]}$, and ${}^3P_J^{[8]}$ channels, while for $\mathcal{Q} = \chi_{cJ}$ the ${}^3P_J^{[1]}$ and ${}^3S_1^{[8]}$ channels appear at leading order in the nonrelativistic expansion. Because we are interested in the large- p_T region, we focus on the leading large- p_T behaviors of the SDCs which are given by leading-power (LP) fragmentation [6, 7]

$$\sigma_{Q\bar{Q}(\mathcal{N})}^{\text{LP}} = \sum_{i=g,q,\bar{q}} \int_0^1 dz \hat{\sigma}_{i(K)} D_{i \rightarrow Q\bar{Q}(\mathcal{N})}(z), \quad (2)$$

where $\hat{\sigma}_{i(K)}$ is the production rate of a massless parton $i = g, q, \text{ or } \bar{q}$ with momentum K , $D_{i \rightarrow Q\bar{Q}(\mathcal{N})}(z)$ is the fragmentation function (FF) for fragmentation of a parton i into $Q\bar{Q}(\mathcal{N})$, $z = P^+/K^+$ with P the $Q\bar{Q}$ momentum and the $+$ direction defined along P in the lab frame. The correction to Eq. (2) is suppressed by m^2/p_T^2 , with m the heavy quark mass, and can be written in terms of next-to-leading power (NLP) fragmentation [8–10]. In large- p_T hadroproduction, the cross section is dominated by gluon fragmentation ($i = g$). Gluon FFs for $\mathcal{N} = {}^3S_1^{[8]}$, ${}^3P_J^{[8]}$, and ${}^3P_J^{[1]}$ involve distributions that are singular at $z = 1$ [11–13]. It is useful to define the Mellin transform

$$\tilde{D}_{i \rightarrow Q\bar{Q}(\mathcal{N})}(N) = \int_0^1 dz z^{N-1} D_{i \rightarrow Q\bar{Q}(\mathcal{N})}(z). \quad (3)$$

The inverse Mellin transform is singular at $z = 1$ if $\tilde{D}_{i \rightarrow Q\bar{Q}(\mathcal{N})}(N)$ does not vanish as $N \rightarrow \infty$. At leading order in the strong coupling α_s , $\tilde{D}_{g \rightarrow Q\bar{Q}({}^3S_1^{[8]})}^{\text{LO}}(N)$ is

a nonzero constant as $N \rightarrow \infty$, while $\tilde{D}_{g \rightarrow Q\bar{Q}(^3P_J^{[8]})}^{\text{LO}}(N)$ and $\tilde{D}_{g \rightarrow Q\bar{Q}(^3P_J^{[1]})}^{\text{LO}}(N)$ diverge like $\log N$, which imply that these FFs involve singular distributions such as delta functions and plus distributions [11–13]. Moreover, these singularities are exacerbated by radiative corrections; the NLO corrections involve double logarithms in N that are proportional to $\alpha_s \log^2 N$ times the leading-order result, which correspond to $[\log(1-z)/(1-z)]_+$ for $^3S_1^{[8]}$ and $[\log^2(1-z)/(1-z)]_+$ for $^3P_J^{[8]}$ and $^3P_J^{[1]}$ [14–17]. Because these logarithms are associated with singularities at $z = 1$, we refer to them as threshold logarithms. We find that these double logarithmic terms almost completely reproduce the strong logarithmic dependences on p_T of the NLO corrections to $\sigma_{Q\bar{Q}(\mathcal{N})}$. This strongly suggests that resumming threshold logarithms is the key for resolving the negative cross section problem.

Soft approximation.—In order to resum the singularities in the FFs, we obtain an expression for the FFs valid near $z = 1$ by using the Grammer-Yennie approximation for soft gluon attachments to the Q and \bar{Q} produced from the fragmenting gluon [5, 7, 18]. We obtain

$$D_{g \rightarrow Q\bar{Q}}^{\text{soft}}(z) = 2M(-g_{\mu\nu})C_{\text{frag}} \left| \frac{i}{K^2 + i\epsilon} \right|^2 \times \left(g^{\mu\alpha} - \frac{K^\mu n^\alpha}{K^+} \right) \left(g^{\nu\beta} - \frac{K^\nu n^\beta}{K^+} \right) \times \langle 0 | \bar{T} [\mathcal{A}_{\text{soft}}^{\beta,c} \Phi_n^{bc}]^\dagger \delta_z^+ T [\mathcal{A}_{\text{soft}}^{\alpha,a} \Phi_n^{ba}] | 0 \rangle, \quad (4)$$

where $M = \sqrt{P^2}$, $C_{\text{frag}} = z^{d-3} K^+ / [2\pi(N_c^2 - 1)(d-2)]$, $d = 4 - 2\epsilon$ is the number of spacetime dimensions, N_c is the number of colors, n is a lightlike vector defined through $K^+ = n \cdot K$, $|0\rangle$ is the QCD vacuum, $\Phi_k \equiv \Phi_k(\infty, 0) = \mathcal{P} \exp[-ig \int_0^\infty d\lambda k \cdot A^{\text{adj}}(k\lambda)]$ is a Wilson line in the adjoint representation defined along a vector k , with \mathcal{P} the path ordering, g the strong coupling, and A the gauge field, T and \bar{T} are time and anti-time orderings, respectively, and we use the shorthand $\delta_z^+ \equiv 2\pi\delta(n \cdot \hat{p} - (1-z)P^+)$, where \hat{p} is an operator that reads off the momentum of the operator to the right. The operator $\mathcal{A}_{\text{soft}}^{\alpha,a}$ represents an arbitrary number of soft-gluon attachments onto the Q and \bar{Q} lines in the soft approximation, and is given by

$$\mathcal{A}_{\text{soft}}^{\alpha,a} = \bar{u}(p_1) W_{p_1}(\infty, 0) (-ig\gamma^\alpha T^a) W_{p_2}^\dagger(\infty, 0) v(p_2), \quad (5)$$

where p_1 and p_2 are the momenta of the Q and \bar{Q} , respectively, and $W_k(t', t) = \mathcal{P} \exp[-ig \int_t^{t'} d\lambda k \cdot A(k\lambda)]$ is a Wilson line in the fundamental representation defined along a vector k . Note that the operators on the right and left of the δ_z^+ are always time ordered and anti-time ordered, respectively. Hereafter we omit the time and anti-time ordering symbols in expressions involving δ_z^+ .

In order to obtain expressions for specific \mathcal{N} , we first expand in powers of the relative momentum $q \equiv (p_1 -$

$p_2)/2$ and then project onto specific color and angular momentum states. In the $^3S_1^{[8]}$ case, we set $q = 0$ in Eq. (4) to obtain

$$D_{g \rightarrow Q\bar{Q}(^3S_1^{[8]})}^{\text{soft}}(z) = \frac{C_{\text{frag}}(d-2)g^2}{4m^3(d-1)(N_c^2 - 1)} S_{^3S_1^{[8]}}(z), \quad (6)$$

where $S_{^3S_1^{[8]}}(z)$ is the soft function defined by

$$S_{^3S_1^{[8]}}(z) = \langle 0 | [\Phi_p^{ca} \Phi_n^{ba}]^\dagger \delta_z^+ \Phi_p^{cd} \Phi_n^{bd} | 0 \rangle, \quad (7)$$

with $p \equiv (p_1 + p_2)/2 = P/2$. The factor $(d-1)(N_c^2 - 1)$ in the denominator comes from the normalization of the LDME. For the $^3P^{[8]}$ case, we expand Eq. (4) to linear order in q . The expansion of the Wilson line can be carried out by using a straightforward generalization of Polyakov's identity [7, 19]. We obtain

$$D_{g \rightarrow Q\bar{Q}(^3P^{[8]})}^{\text{soft}}(z) = -\frac{C_{\text{frag}}(d-2)g^4 S_{^3P^{[8]}}(z)}{4m^3(d-1)^2(N_c^2 - 1)}, \quad (8)$$

where the soft function $S_{^3P^{[8]}}(z)$ is given by

$$S_{^3P^{[8]}}(z) = \langle 0 | [\mathcal{W}_\alpha^{yx}]^\dagger \delta_z^+ \mathcal{W}_\beta^{yx} | 0 \rangle g^{\alpha\beta}, \quad (9)$$

with

$$\mathcal{W}_\beta^{yx} = \int_0^\infty d\lambda \lambda \Phi_p^{yc}(\infty, \lambda) p^\mu G_{\mu\beta}^b(p\lambda) d^{bcd} \times \Phi_p^{da}(\lambda, 0) \Phi_n^{xa}, \quad (10)$$

where $G_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + ig[A_\mu, A_\nu]$ is the QCD field-strength tensor. Finally, for the $^3P_J^{[1]}$ case, we have

$$D_{g \rightarrow Q\bar{Q}(^3P_J^{[1]})}^{\text{soft}}(z) = -\frac{C_{\text{frag}}(d-2)g^4}{4N_c^2 m^3(d-1)^2} \frac{9}{(2J+1)} \times [c_J S_{^3P^{[1]}}(z) + c_J^{TT} S_{^3P^{[1]}}^{TT}(z)], \quad (11)$$

where $c_0 = (d-1)^{-2}$, $c_1 = (d-2)/[2(d-1)]$, $c_2 = (d-2)(d+1)/[2(d-1)^2]$, $c_0^{TT} = [(d-1)(d-2)]^{-1}$, $c_1^{TT} = -[2(d-2)]^{-1}$, $c_2^{TT} = (d-3)/[2(d-1)(d-2)]$, and the soft functions are given by

$$S_{^3P^{[1]}}(z) = \langle 0 | [\bar{\mathcal{W}}_\alpha^b]^\dagger \delta_z^+ \bar{\mathcal{W}}_\beta^b | 0 \rangle g^{\alpha\beta}, \quad (12a)$$

$$S_{^3P^{[1]}}^{TT}(z) = \langle 0 | [\bar{\mathcal{W}}_\alpha^b]^\dagger \delta_z^+ \bar{\mathcal{W}}_\beta^b | 0 \rangle \times \left[\frac{p^2 n^\alpha n^\beta}{(n \cdot p)^2} + \frac{g^{\alpha\beta}}{d-1} \right], \quad (12b)$$

with

$$\bar{\mathcal{W}}_\beta^b = \int_0^\infty d\lambda \lambda p^\mu G_{\mu\beta}^d(p\lambda) \Phi_p^{da}(\lambda, 0) \Phi_n^{ba}. \quad (13)$$

We use the definitions of the color-singlet LDMEs in Ref. [2], which differ from Refs. [17, 20] by a factor of $2N_c$. The $S_{^3P^{[1]}}^{TT}(z)$ term comes from the anisotropic contribution that arises from projecting onto specific J . As

we will see later, $S_{3P^{[1]}}^{TT}(z)$ will not produce double logarithmic contributions and can be neglected in this work.

The results in Eqs. (6), (8), and (11) reproduce the leading singularities of the FFs at $z = 1$, which are contained in the soft functions. We can verify this at LO by computing the $\mathcal{S}_{\mathcal{N}}$ at leading nonvanishing order:

$$S_{3S_1^{[8]}}^{\text{LO}}(z) = \frac{2\pi(N_c^2 - 1)}{P^+} \delta(1 - z), \quad (14a)$$

$$S_{3P^{[8]}}^{\text{LO}}(z) = -\frac{(d-2)4B_F(N_c^2 - 1)\Gamma(1 + \epsilon)}{2\pi^{1-\epsilon}m^2P^+(1-z)^{1+2\epsilon}}, \quad (14b)$$

$$S_{3P^{[1]}}^{\text{LO}}(z) = -\frac{(d-2)(N_c^2 - 1)\Gamma(1 + \epsilon)}{2\pi^{1-\epsilon}m^2P^+(1-z)^{1+2\epsilon}}, \quad (14c)$$

$$S_{3P^{[1]}}^{TT, \text{LO}}(z) = \frac{(N_c^2 - 1)\epsilon(1 - \epsilon)(1 - 2\epsilon)\Gamma(1 + \epsilon)}{3(3 - 2\epsilon)\pi^{1-\epsilon}m^2P^+(1-z)^{1+2\epsilon}}, \quad (14d)$$

where $B_F = (N_c^2 - 4)/(4N_c)$. Note that $S_{3P^{[8]}}^{\text{LO}}$, $S_{3P^{[1]}}^{\text{LO}}$, and $S_{3P^{[1]}}^{TT, \text{LO}}$ come from diagrams where a single gluon is exchanged between the two field-strength tensors. By using the identity

$$\frac{1}{(1-z)^{1+n\epsilon}} = -\frac{1}{n\epsilon_{\text{IR}}} \delta(1-z) + \left[\frac{1}{(1-z)^{1+n\epsilon}} \right]_+, \quad (15)$$

it is easy to show that the $D_{g \rightarrow Q\bar{Q}(\mathcal{N})}^{\text{soft}}(z)$ reproduce the singular distributions in the LO FFs [11–13]. Note that $S_{3P^{[1]}}^{TT, \text{LO}}$ does not produce singular distributions because it contains an explicit factor of ϵ .

Radiative corrections to soft functions.— We now compute the double logarithms in the soft functions at NLO, which come from contributions involving double poles in ϵ . We first consider $S_{3S_1^{[8]}}(z)$. By explicit calculation we can show that in Feynman gauge, the double logarithms come from NLO diagrams where an additional gluon is exchanged between the Wilson lines Φ_p and Φ_n . The result is

$$\begin{aligned} S_{3S_1^{[8]}}^{\text{NLO}}(z) &= \frac{2\alpha_s C_A (N_c^2 - 1)}{P^+(1-z)^{1+2\epsilon}} \left(\frac{1}{\epsilon_{\text{UV}}} + O(\epsilon^0) \right) \\ &\quad - \frac{\alpha_s C_A (N_c^2 - 1)}{P^+} \delta(1-z) \frac{1}{\epsilon_{\text{UV}}} \\ &\quad \times \left(\frac{1}{\epsilon_{\text{UV}}} - \frac{1}{\epsilon_{\text{IR}}} \right) + \dots, \end{aligned} \quad (16)$$

where $C_A = N_c$, and we neglect any contributions that do not produce double logarithms. The contribution in the first line comes from the real diagram where the gluon crosses the cut, while the remaining terms come from the virtual diagram. By using the identity in Eq. (15), we obtain the expression for the double logarithmic correction to $S_{3S_1^{[8]}}$ at NLO given by

$$\begin{aligned} S_{3S_1^{[8]}}^{\text{NLO}}(z) &= \frac{2\pi(N_c^2 - 1)}{P^+} \frac{\alpha_s C_A}{\pi} \left\{ \frac{-\delta(1-z)}{2\epsilon_{\text{UV}}^2} + \frac{1}{\epsilon_{\text{UV}}(1-z)_+} \right. \\ &\quad \left. + \left[\frac{-2\log(1-z)}{1-z} \right]_+ + \dots \right\}. \end{aligned} \quad (17)$$

This reproduces the double logarithmic term $[\log(1-z)/(1-z)]_+$ in the ${}^3S_1^{[8]}$ FF at NLO [14–16].

Similarly to the ${}^3S_1^{[8]}$ case, double logarithmic corrections to the ${}^3P^{[8]}$ and ${}^3P^{[1]}$ soft functions arise only from planar diagrams where an additional gluon is exchanged between the Wilson lines $\Phi_p(\lambda, 0)$ and Φ_n . The result is

$$S_{3P^{[8]}}^{\text{NLO}}(z) = \frac{\alpha_s C_A}{\pi} \frac{4B_F(N_c^2 - 1)\epsilon_{\text{UV}}^{-2}}{2\pi m^2[P^+(1-z)]^{1+4\epsilon}} + \dots, \quad (18a)$$

$$S_{3P^{[1]}}^{\text{NLO}}(z) = \frac{\alpha_s C_A}{\pi} \frac{(N_c^2 - 1)\epsilon_{\text{UV}}^{-2}}{2\pi m^2[P^+(1-z)]^{1+4\epsilon}} + \dots, \quad (18b)$$

where we neglect any contribution that do not produce any double logarithmic corrections. Again, by using Eq. (15) we can check that the double logarithmic term $[\log^2(1-z)/(1-z)]_+$ agrees with the explicit NLO calculation of the FFs in Ref. [17]. Similarly to the LO case, $S_{3P^{[1]}}^{TT}$ does not produce double logarithmic singularities at NLO because it contains an explicit factor of ϵ .

Resummation.— It is straightforward to resum the double logarithmic corrections to all orders in perturbation theory, by using the fact that they arise from planar diagrams that can be exponentiated in Mellin space [21]:

$$\tilde{S}_{\mathcal{N}}^{\text{resum}}(N) = \exp[J_{\mathcal{N}}^N] \tilde{S}_{\mathcal{N}}^{\text{LO}}(N), \quad (19)$$

where $J_{\mathcal{N}}^N$ is given at leading double logarithmic level by

$$J_{3S_1^{[8]}}^N = \frac{\alpha_s C_A}{\pi} \int_0^1 dz z^{N-1} \left[\frac{-2\log(1-z)}{1-z} \right]_+, \quad (20a)$$

$$J_{3P^{[8]}}^N = J_{3P^{[1]}}^N = \frac{4}{3} J_{3S_1^{[8]}}^N. \quad (20b)$$

By expanding Eq. (19) in powers of α_s we reproduce the double logarithmic corrections at NLO. The resummed expressions for the FFs can be obtained in the same way. At NLO accuracy, the resummed FFs can be written as

$$\begin{aligned} \tilde{D}_{g \rightarrow Q\bar{Q}(\mathcal{N})}^{\text{resum}}(N) &= \exp[J_{\mathcal{N}}^N] \times \left(\tilde{D}_{g \rightarrow Q\bar{Q}(\mathcal{N})}^{\text{FO}}(N) \right. \\ &\quad \left. - J_{\mathcal{N}}^N \tilde{D}_{g \rightarrow Q\bar{Q}(\mathcal{N})}^{\text{LO}}(N) \right), \end{aligned} \quad (21)$$

where the last term in the parenthesis subtracts the double logarithmic correction term in the $\tilde{D}_{g \rightarrow Q\bar{Q}(\mathcal{N})}^{\text{FO}}(N)$ at NLO accuracy to avoid double counting. Note that because the factor $\exp[J_{\mathcal{N}}^N]$ vanishes as $N \rightarrow \infty$ faster than any power of N , the inverse Mellin transform yields regular functions in z that vanish at $z = 1$. The resummed expression for ${}^3S_1^{[8]}$ agrees with the calculation in the soft gluon factorization formalism in Refs. [4, 22] at double logarithmic level. The resummed results for ${}^3P_J^{[8]}$ and ${}^3P_J^{[1]}$ are new.

We note that different expressions for the resummed FFs are possible that are equivalent to Eq. (21) at




FIG. 1. Gluon FFs with resummed threshold double logarithms times z^3 for production of J/ψ (top) and χ_{cJ} (bottom) for $J = 1$ and 2 . Central values of FO results are also shown for comparison. $\text{Br}_J \equiv \text{Br}_{\chi_{cJ} \rightarrow J/\psi + \gamma}$ is the branching fraction for decays of χ_{cJ} into $J/\psi + \gamma$.

the current accuracy. For example, the NLO terms in Eq. (21) may be expanded in powers of α_s , or an expression based on evolution equations may be used [22]. We find that these alternative expressions lead to cross sections that differ by less than 15%, which is not significant compared to other uncertainties in the theory of heavy quarkonium production.

Numerical results.— We now show the numerical results for the resummed FFs. We work with FFs for production of quarkonium \mathcal{Q} written in terms of $D_{i \rightarrow \mathcal{Q} \bar{\mathcal{Q}}(\mathcal{N})}(z)$ as

$$D_{g \rightarrow \mathcal{Q}}(z) = \sum_{\mathcal{N}} D_{g \rightarrow \mathcal{Q} \bar{\mathcal{Q}}(\mathcal{N})}(z) \langle \mathcal{O}^{\mathcal{Q}}(\mathcal{N}) \rangle, \quad (22)$$

where the sum is over $\mathcal{N} = {}^3S_1^{[1]}, {}^3S_1^{[8]}, {}^1S_0^{[8]}$, and ${}^3P_J^{[8]}$ for $\mathcal{Q} = J/\psi$ or $\psi(2S)$, and $\mathcal{N} = {}^3P_J^{[1]}$ and ${}^3S_1^{[8]}$ for $\mathcal{Q} = \chi_{cJ}$. Note that we include the contributions from ${}^3S_1^{[1]}$ and ${}^1S_0^{[8]}$, which do not contain singularities and are not affected by resummation at the current level of accuracy [13, 20, 23, 24]. For consistency, we compute the FO FFs to order α_s^2 , except for the ${}^3S_1^{[1]}$ FF, which we compute at leading nonvanishing order ($\mathcal{O}(\alpha_s^3)$) [20, 23, 24]. Because we are interested in the large p_T region, we evolve the FFs to the $\overline{\text{MS}}$ scale 50 GeV from 3 GeV by using the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) evolution equation [25–28] at leading logarithmic (LL) accuracy. We take the J/ψ LDMEs determined in Ref. [29] in the large p_T re-

gion, and we use the χ_{cJ} LDMEs from Ref. [30]. The LDMEs are renormalized in the $\overline{\text{MS}}$ scheme at the scale m . In order to compensate for the fact that resummation enhances the relative size of the ${}^3P^{[8]}$ SDCs compared to ${}^3S_1^{[8]}$ by about 10%, from which the ${}^3P^{[8]}$ LDME was determined in Ref. [29], we reduce the central value of the ${}^3P^{[8]}$ LDME by 10%. That is, we use $\langle \mathcal{O}^{J/\psi}({}^3S_1^{[1]}) \rangle = 1.18 \pm 0.35 \text{ GeV}^3$, $\langle \mathcal{O}^{J/\psi}({}^3S_1^{[8]}) \rangle = (1.40 \pm 0.42) \times 10^{-2} \text{ GeV}^3$, $\langle \mathcal{O}^{J/\psi}({}^1S_0^{[8]}) \rangle = (-0.63 \pm 3.22) \times 10^{-2} \text{ GeV}^3$, $\langle \mathcal{O}^{J/\psi}({}^3P_0^{[8]}) \rangle = (5.25 \pm 1.86) \times 10^{-2} \text{ GeV}^5$, $\langle \mathcal{O}^{\chi_{c0}}({}^3P_0^{[1]}) \rangle = (8.16 \pm 2.45) \times 10^{-2} \text{ GeV}^5$, and $\langle \mathcal{O}^{\chi_{c0}}({}^3S_1^{[8]}) \rangle = (1.57 \pm 0.47) \times 10^{-3} \text{ GeV}^3$. Note that due to the universality relations in the LDMEs obtained in Refs. [29, 31], the $\psi(2S)$ FF can be obtained by uniformly rescaling the J/ψ FF. We display the gluon FFs for transversely and longitudinally polarized J/ψ in Fig. 1. We show the FFs multiplied by z^3 , because when computing p_T -differential cross sections using Eq. (2), the $\hat{\sigma}_i$ behave approximately like z^3 . For comparison, we also show results for FO FFs, which involve singularities at $z = 1$ that cannot be displayed like regular functions. We see that the resummed J/ψ FFs are smooth functions of z , and they are positive or at least consistent with zero within uncertainties for all $0 < z < 1$, which ensures the positivity of J/ψ production rates. In contrast, the transversely polarized FF in FO perturbation theory rapidly changes sign near $z = 1$. The longitudinal J/ψ FF is unaffected by resummation, because it is free of singularities at the current level of accuracy [14, 32]. The results for the resummed FFs lead to an estimate of the polarization of J/ψ and $\psi(2S)$ at $p_T = 100 \text{ GeV}$ given by $-0.25 \lesssim \lambda_\theta \lesssim +0.15$ in the helicity frame at midrapidity, which is compatible with previous estimates [29, 31] but smaller than recent CMS measurements at large p_T [33]. While in the resummed case we expect λ_θ to stay almost constant as p_T increases, in the FO calculation the parameter may unphysically drop below -1 when the cross section turns negative, because the longitudinal contribution is unaffected by threshold logarithms. Similarly, we show the FFs for χ_{c1} and χ_{c2} in Fig. 1, scaled by the branching fractions into $J/\psi + \gamma$ from PDG [34]. Just like the J/ψ case, the resummed χ_{cJ} FFs are smooth functions of z that are positive for the whole range of z , unlike the FO calculations which change sign rapidly near $z = 1$.

Finally, we compute the prompt J/ψ production rates at the $\sqrt{s} = 13 \text{ TeV}$ LHC to compare with ATLAS measurements in Ref. [1]. We use the method used in Ref. [35] to compute the cross sections in LP fragmentation, except that we use the gluon FFs including threshold double logarithms and DGLAP logarithms resummed to LL accuracy to compute the LP contribution. We also include the contributions from light-quark FFs, which are not affected by resummation at the current level of accuracy. We include feeddown contributions from decays




FIG. 2. Prompt J/ψ production rates from pp collisions at $\sqrt{s} = 13$ TeV computed from resummed SDCs compared to ATLAS data. Central values of FO NLO results are shown for comparison. $B \equiv \text{Br}_{J/\psi \rightarrow \mu^+ \mu^-}$ is the J/ψ dimuon branching fraction.

of χ_{cJ} and $\psi(2S)$, with branching fractions taken from PDG [34]. We include the NLP contributions we obtain from FO SDCs computed at NLO from the FDCHQHP package [36]. We find that the NLP contributions amount to about 10% at $p_T = 60$ GeV, and diminish to less than 1% for p_T larger than 100 GeV. We show the large- p_T cross sections computed from the resummed FFs in Fig. 2 compared to ATLAS data [1]. The resummed results are in fair agreement with data in the large p_T region. In contrast, the results from FO SDCs [37] shown in Fig. 2 fall below measured data and turn negative at large p_T . This shows that resummation of threshold logarithms is absolutely necessary in order to describe heavy quarkonium production rates at very large transverse momentum.

In Fig. 3, we show results for direct J/ψ cross sections computed from other LDME determinations in Refs. [38–40]. Note that the LDMEs in Ref. [38] (B&K) lead to transverse polarization, which is incompatible with measurements [41]. It is known that the LDMEs from Ref. [40] (Bodwin et al.) are incompatible with η_c data [42, 43]. It is important to note that regardless of the choice of LDMEs, in a FO calculation the χ_{cJ} cross sections always turn negative, so that any solid prediction of prompt J/ψ production rate at large p_T requires




FIG. 3. Direct J/ψ production rates from pp collisions at $\sqrt{s} = 13$ TeV for the rapidity range $1.5 \leq |y| \leq 2.0$ computed by using LDMEs adopted in this work, compared to ones from Refs. [38] (B&K), [39] (Chao et al.), and [40] (Bodwin et al.). ATLAS data for prompt J/ψ cross section are shown for comparison.

resummation of threshold logarithms.

Conclusions.—The resummation of threshold double logarithms we computed in this Letter substantially improves the NRQCD description of charmonium production rates at large transverse momentum. This resolves the catastrophic failure of fixed-order perturbation theory where large- p_T cross sections can turn unphysically negative. Resummation of threshold logarithms may also be important in describing measurements involving kinematical cuts near the boundary of phase space such as photoproduction rates [44–46] and quarkonium in jet [47–51]. In the case of bottomonium, we expect that the effect of threshold logarithms will become important if the transverse momentum exceeds 1 TeV, based on the LDME determinations in Refs. [29, 30, 52]. It would be interesting to improve the accuracy of the resummation to single logarithmic level, which requires calculation of the soft functions to single-pole accuracy. The resummation may also be extended to NLP fragmentation contributions, which become important for lower p_T .

We thank Geoffrey Bodwin for fruitful discussions on resummation of threshold logarithms. The work of H. S. C. and J. L. is supported by the National Research Foundation of Korea (NRF) Grant funded by the Korea government (MSIT) under Contract No. NRF2020R1A2C3009918. H. S. C. also acknowledges support by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (Grant No. RS-2023-00248313) and a Korea University grant.

All authors contributed equally to this work.

* neville@korea.ac.kr

† kim87@kma.ac.kr

‡ jungil@korea.ac.kr

- [1] G. Aad *et al.* (ATLAS), Measurement of the production cross-section of J/ψ and $\psi(2S)$ mesons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Eur. Phys. J. C* **84**, 169 (2024), arXiv:2309.17177 [hep-ex].
- [2] G. T. Bodwin, E. Braaten, and G. P. Lepage, Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium, *Phys. Rev. D* **51**, 1125 (1995), [Erratum: *Phys.Rev.D* 55, 5853 (1997)], arXiv:hep-ph/9407339.
- [3] H. S. Chung, Resummation and renormalization of kinematical effects in inclusive P-wave quarkonium production, *J. High Energy Phys.* **07**, 007, arXiv:2303.17240 [hep-ph].
- [4] A.-P. Chen, Y.-Q. Ma, and C. Meng, Resolving negative cross section of quarkonium hadroproduction using soft gluon factorization, *Phys. Rev. D* **108**, 014003 (2023), arXiv:2304.04552 [hep-ph].
- [5] G. Grammer, Jr. and D. R. Yennie, Improved treatment for the infrared divergence problem in quantum electrodynamics, *Phys. Rev. D* **8**, 4332 (1973).
- [6] J. C. Collins and D. E. Soper, Parton Distribution and Decay Functions, *Nucl. Phys. B* **194**, 445 (1982).
- [7] G. C. Nayak, J.-W. Qiu, and G. Sterman, Fragmentation, NRQCD and NNLO factorization analysis in heavy quarkonium production, *Phys. Rev. D* **72**, 114012 (2005), arXiv:hep-ph/0509021.
- [8] Z.-B. Kang, J.-W. Qiu, and G. Sterman, Factorization and quarkonium production, *Nucl. Phys. B Proc. Suppl.* **214**, 39 (2011).
- [9] Z.-B. Kang, Y.-Q. Ma, J.-W. Qiu, and G. Sterman, Heavy Quarkonium Production at Collider Energies: Factorization and Evolution, *Phys. Rev. D* **90**, 034006 (2014), arXiv:1401.0923 [hep-ph].
- [10] Y.-Q. Ma, J.-W. Qiu, G. Sterman, and H. Zhang, Factorized power expansion for high- p_T heavy quarkonium production, *Phys. Rev. Lett.* **113**, 142002 (2014), arXiv:1407.0383 [hep-ph].
- [11] E. Braaten and Y.-Q. Chen, Dimensional regularization in quarkonium calculations, *Phys. Rev. D* **55**, 2693 (1997), arXiv:hep-ph/9610401.
- [12] G. T. Bodwin and J. Lee, Relativistic corrections to gluon fragmentation into spin triplet S wave quarkonium, *Phys. Rev. D* **69**, 054003 (2004), arXiv:hep-ph/0308016.
- [13] G. T. Bodwin, U.-R. Kim, and J. Lee, Higher-order relativistic corrections to gluon fragmentation into spin-triplet S-wave quarkonium, *J. High Energy Phys.* **11**, 020, [Erratum: *J. High Energy Phys.* 07, 170 (2023)], arXiv:1208.5301 [hep-ph].
- [14] E. Braaten and J. Lee, Next-to-leading order calculation of the color octet $3S(1)$ gluon fragmentation function for heavy quarkonium, *Nucl. Phys. B* **586**, 427 (2000), arXiv:hep-ph/0004228.
- [15] J. Lee, Next-to-leading order calculation of a fragmentation function in a light-cone gauge, *Phys. Rev. D* **71**, 094007 (2005), arXiv:hep-ph/0504285.
- [16] Y.-Q. Ma, J.-W. Qiu, and H. Zhang, Heavy quarkonium fragmentation functions from a heavy quark pair. I. S wave, *Phys. Rev. D* **89**, 094029 (2014), arXiv:1311.7078 [hep-ph].
- [17] P. Zhang, C. Meng, Y.-Q. Ma, and K.-T. Chao, Gluon fragmentation into ${}^3P_J^{[1,8]}$ quark pair and test of NRQCD factorization at two-loop level, *J. High Energy Phys.* **08**, 111, arXiv:2011.04905 [hep-ph].
- [18] J. C. Collins and D. E. Soper, Back-To-Back Jets in QCD, *Nucl. Phys. B* **193**, 381 (1981), [Erratum: *Nucl.Phys.B* 213, 545 (1983)].
- [19] A. M. Polyakov, Gauge Fields as Rings of Glue, *Nucl. Phys. B* **164**, 171 (1980).
- [20] P. Zhang, Y.-Q. Ma, Q. Chen, and K.-T. Chao, Analytical calculation for the gluon fragmentation into spin-triplet S-wave quarkonium, *Phys. Rev. D* **96**, 094016 (2017), arXiv:1708.01129 [hep-ph].
- [21] E. Laenen, G. Sterman, and W. Vogelsang, Recoil and threshold corrections in short distance cross-sections, *Phys. Rev. D* **63**, 114018 (2001), arXiv:hep-ph/0010080.
- [22] A.-P. Chen, X.-B. Jin, Y.-Q. Ma, and C. Meng, Fragmentation function of $g \rightarrow Q\bar{Q}({}^3S_1^{[8]})$ in soft gluon factorization and threshold resummation, *JHEP* **06**, 046, arXiv:2103.15121 [hep-ph].
- [23] E. Braaten and T. C. Yuan, Gluon fragmentation into heavy quarkonium, *Phys. Rev. Lett.* **71**, 1673 (1993), arXiv:hep-ph/9303205.
- [24] E. Braaten and T. C. Yuan, Gluon fragmentation into spin triplet S wave quarkonium, *Phys. Rev. D* **52**, 6627 (1995), arXiv:hep-ph/9507398.
- [25] V. N. Gribov and L. N. Lipatov, Deep inelastic e p scattering in perturbation theory, *Sov. J. Nucl. Phys.* **15**, 438 (1972).
- [26] L. N. Lipatov, The parton model and perturbation theory, *Yad. Fiz.* **20**, 181 (1974), *Sov. J. Nucl. Phys.* **20**, 94 (1975).
- [27] Y. L. Dokshitzer, Calculation of the Structure Functions for Deep Inelastic Scattering and e+ e- Annihilation by Perturbation Theory in Quantum Chromodynamics., *Sov. Phys. JETP* **46**, 641 (1977).
- [28] G. Altarelli and G. Parisi, Asymptotic Freedom in Parton Language, *Nucl. Phys. B* **126**, 298 (1977).
- [29] N. Brambilla, H. S. Chung, A. Vairo, and X.-P. Wang, Inclusive production of J/ψ , $\psi(2S)$, and Υ states in pNRQCD, *J. High Energy Phys.* **03**, 242, arXiv:2210.17345 [hep-ph].
- [30] N. Brambilla, H. S. Chung, and A. Vairo, Inclusive production of heavy quarkonia in pNRQCD, *J. High Energy Phys.* **09**, 032, arXiv:2106.09417 [hep-ph].
- [31] N. Brambilla, H. S. Chung, A. Vairo, and X.-P. Wang, Production and polarization of S-wave quarkonia in potential nonrelativistic QCD, *Phys. Rev. D* **105**, L111503 (2022), arXiv:2203.07778 [hep-ph].
- [32] Y.-Q. Ma, J.-W. Qiu, and H. Zhang, Fragmentation functions of polarized heavy quarkonium, *J. High Energy Phys.* **06**, 021, arXiv:1501.04556 [hep-ph].
- [33] A. Hayrapetyan *et al.* (CMS), Measurement of the polarizations of prompt and non-prompt J/ψ and $\psi(2S)$ mesons produced in pp collisions at $\sqrt{s}=13$ TeV, *Phys. Lett. B* **858**, 139044 (2024), arXiv:2406.14409 [hep-ex].
- [34] S. Navas *et al.* (Particle Data Group), Review of Particle Physics, *Phys. Rev. D* **110**, 030001 (2024).
- [35] G. T. Bodwin, K.-T. Chao, H. S. Chung, U.-R. Kim, J. Lee, and Y.-Q. Ma, Fragmentation contributions to hadroproduction of prompt J/ψ , χ_{cJ} , and $\psi(2S)$ states, *Phys. Rev. D* **93**, 034041 (2016), arXiv:1509.07904 [hep-ph].
- [36] L.-P. Wan and J.-X. Wang, FDCHQHP: A Fortran package for heavy quarkonium hadroproduction, *Comput. Phys. Commun.* **185**, 2939 (2014), arXiv:1405.2143 [hep-ph].

- [37] While we use FDCHQHP to compute the FO SDCs, for values of p_T larger than 100 GeV, we compute the FO SDCs from LP fragmentation by using FO FFs. The agreement between LP fragmentation and FO SDCs at large p_T has been verified in Ref. [35].
- [38] M. Butenschoen and B. A. Kniehl, World data of J/ψ production consolidate NRQCD factorization at NLO, Phys. Rev. D **84**, 051501 (2011), arXiv:1105.0820 [hep-ph].
- [39] H. Han, Y.-Q. Ma, C. Meng, H.-S. Shao, and K.-T. Chao, η_c production at LHC and indications on the understanding of J/ψ production, Phys. Rev. Lett. **114**, 092005 (2015), arXiv:1411.7350 [hep-ph].
- [40] G. T. Bodwin, H. S. Chung, U.-R. Kim, and J. Lee, Fragmentation contributions to J/ψ production at the Tevatron and the LHC, Phys. Rev. Lett. **113**, 022001 (2014), arXiv:1403.3612 [hep-ph].
- [41] S. Chatrchyan *et al.* (CMS), Measurement of the Prompt J/ψ and $\psi(2S)$ Polarizations in pp Collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B **727**, 381 (2013), arXiv:1307.6070 [hep-ex].
- [42] R. Aaij *et al.* (LHCb), Measurement of the $\eta_c(1S)$ production cross-section in proton-proton collisions via the decay $\eta_c(1S) \rightarrow p\bar{p}$, Eur. Phys. J. C **75**, 311 (2015), arXiv:1409.3612 [hep-ex].
- [43] R. Aaij *et al.* (LHCb), Measurement of the $\eta_c(1S)$ production cross-section in pp collisions at $\sqrt{s} = 13$ TeV, Eur. Phys. J. C **80**, 191 (2020), arXiv:1911.03326 [hep-ex].
- [44] S. Chekanov *et al.* (ZEUS), Measurements of inelastic J/ψ and ψ -prime photoproduction at HERA, Eur. Phys. J. C **27**, 173 (2003), arXiv:hep-ex/0211011.
- [45] C. Adloff *et al.* (H1), Inelastic photoproduction of J/ψ mesons at HERA, Eur. Phys. J. C **25**, 25 (2002), arXiv:hep-ex/0205064.
- [46] F. D. Aaron *et al.* (H1), Inelastic Production of J/ψ Mesons in Photoproduction and Deep Inelastic Scattering at HERA, Eur. Phys. J. C **68**, 401 (2010), arXiv:1002.0234 [hep-ex].
- [47] M. Baumgart, A. K. Leibovich, T. Mehen, and I. Z. Rothstein, Probing Quarkonium Production Mechanisms with Jet Substructure, J. High Energy Phys. **11**, 003, arXiv:1406.2295 [hep-ph].
- [48] R. Aaij *et al.* (LHCb), Study of J/ψ Production in Jets, Phys. Rev. Lett. **118**, 192001 (2017), arXiv:1701.05116 [hep-ex].
- [49] Z.-B. Kang, J.-W. Qiu, F. Ringer, H. Xing, and H. Zhang, J/ψ production and polarization within a jet, Phys. Rev. Lett. **119**, 032001 (2017), arXiv:1702.03287 [hep-ph].
- [50] R. Bain, L. Dai, A. Leibovich, Y. Makris, and T. Mehen, NRQCD Confronts LHCb Data on Quarkonium Production within Jets, Phys. Rev. Lett. **119**, 032002 (2017), arXiv:1702.05525 [hep-ph].
- [51] A. M. Sirunyan *et al.* (CMS), Study of J/ψ meson production inside jets in pp collisions at $\sqrt{s} = 8$ TeV, Phys. Lett. B **804**, 135409 (2020), arXiv:1910.01686 [hep-ex].
- [52] H. Han, Y.-Q. Ma, C. Meng, H.-S. Shao, Y.-J. Zhang, and K.-T. Chao, $\Upsilon(nS)$ and $\chi_b(nP)$ production at hadron colliders in nonrelativistic QCD, Phys. Rev. D **94**, 014028 (2016), arXiv:1410.8537 [hep-ph].