

Monopole-Fermion Scattering and the Solution to the Semiton/Unitarity Puzzle

Vazha Loladze^{1,*} and Takemichi Okui^{2,3,†}

¹*Rudolf Peierls Centre for Theoretical Physics, University of Oxford, Parks Road, Oxford OX1 3PU, UK*

²*Department of Physics, Florida State University, Tallahassee, FL 32306, USA*

³*Theory Center, High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801, Japan*

We study Polchinski’s “fermion-rotor system” as an accurate description of charged Weyl fermions scattering on a magnetic monopole core in the limit of zero gauge coupling. Traditionally it was thought such scattering could lead to fractional particle numbers (“semitons”). By direct calculations we show those semitonic processes are in fact free propagation, facilitated by composite fermion-rotor operators interpolating the “forbidden” states, effectively “recovering” both ingoing and outgoing states in every lowest partial wave. Non-semitonic Callan-Rubakov processes are unchanged.

The Puzzle: Imagine a spherical monopole M with magnetic charge g sitting at $\vec{r} = 0$ and a left-handed Weyl fermion field f with electric charge q . The equation of motion of f in the background field \vec{A} of M reads $\partial_t f = \vec{\sigma} \cdot \vec{D} f$ with $\vec{D} \equiv \vec{\partial} - iq\vec{A}$. In $f_\alpha(t, r, \theta, \phi) = \frac{\chi(t, r)}{r} \Phi_\alpha(\theta, \phi)$, suppose the spinor $\Phi_\alpha(\theta, \phi)$ is in the lowest- j partial wave, where $\vec{J} = \vec{L} + \frac{\vec{\sigma}}{2}$ with $\vec{L} = \vec{r} \times (-i\vec{D}) - qg\hat{r}$. The equation of motion then becomes a free field equation $\partial_t \chi(t, r) = \text{sgn}(qg) \partial_r \chi(t, r)$, so $\chi(t, r)$ has only radially ingoing (outgoing) waves for $qg > 0$ ($qg < 0$) [1–3][4]. Half the waves seem missing. What does that imply?

Consider, e.g., a minimal ($|qg| = \frac{1}{2}$) SU(5) GUT monopole from the breaking SU(2)_M → U(1)_M embedded in the breaking SU(5) → SU(3) × SU(2) × U(1). Taking only the first generation of SM fermions for simplicity, there are four SU(2)_M doublets,

$$\chi_1, \chi_2, \chi_3, \chi_4 \equiv \begin{pmatrix} e \\ d_3^c \end{pmatrix}_L, \begin{pmatrix} d^3 \\ e^c \end{pmatrix}_L, \begin{pmatrix} u_1^c \\ u^2 \end{pmatrix}_L, \begin{pmatrix} u_2^c \\ u^1 \end{pmatrix}_L, \quad (1)$$

where χ_1 and $\chi_{2,3,4}$ are from the $\bar{\mathbf{5}}$ and $\mathbf{10}$ of SU(5), respectively. Neglecting SM gauge and Yukawa couplings, there is a global SU(4)_F flavor symmetry rotating the four doublets. If $g < 0$, since the lower (upper) fields of the doublets have $q = -\frac{1}{2}$ ($+\frac{1}{2}$) under U(1)_M, the lower (upper) fields contain only ingoing (outgoing) waves.

Here is Rubakov [5] and Callan [6–8]’s 42-year-old puzzle, which has recently become an active area of research [9–17]. Suppose, e.g., an ingoing e^c particle scatters on M with $g < 0$. Since outgoing particles must be interpolated by the upper fields in (1), let A, B, C, D be respectively the numbers of e, d^3, u_1^c, u_2^c particles minus those of their antiparticles in the final state. Since the monopole background conserves four charges (U(1)_M and three diagonal SU(4)_F generators), the four unknowns are fixed by sheer conservation laws, which gives $B = \frac{1}{2}$ and $A = C = D = -\frac{1}{2}$. Thus, $|e^c, M\rangle$ is predicted to become “ $|\frac{1}{2}\bar{e}, \frac{1}{2}d^3, \frac{1}{2}u_1^c, \frac{1}{2}u_2^c, M\rangle$ ” (hence the *semitons*). But there is no such thing as *half* a particle in the Fock

space (hence the *unitarity* puzzle).¹

The Effective Theory: We study the puzzle by employing Polchinski’s *fermion-rotor system* [18], which by construction is an exact effective theory of monopole-fermion scattering in the lowest partial wave in the limit of vanishing U(1)_M gauge coupling (and SM couplings). This limit lets us focus on the exact degrees of freedom and dynamics relevant to the puzzle. It also entails the limits of an infinitesimal core radius and infinite mass of the monopole, or equivalently the low energy limit of the fermions.

Take $|qg| = \frac{1}{2}$ with $g < 0$ and consider N doublets χ_k ($k = 1, \dots, N$ with N even [19]). For each χ_k , let χ_k^- (χ_k^+) be its lower (upper) component. [18] embeds $\chi_k^-(t, r)$ (ingoing) and $\chi_k^+(t, r)$ (outgoing) into a single (1+1)D right-mover $\psi_k(t, x)$ as

$$\psi_k(t, x) \equiv \begin{cases} \chi_k^-(t, r)|_{r=-x} & \text{if } x \leq -r_0, \\ \chi_k^+(t, r)|_{r=x} & \text{if } x \geq r_0, \end{cases} \quad (2)$$

and smoothly varying in-between, where r_0 is the small radius of the core to be taken to zero whenever it is already safe to do so. The U(1)_M charge of $\psi_k(t, x)$ is $q(x) = -\frac{1}{2}$ ($+\frac{1}{2}$) for $x \leq -r_0$ ($x \geq r_0$), varying smoothly in-between. To avoid clutter below we will always give expressions as if $r_0 = 0$ but nonzero r_0 is understood.

Next, [18] introduces $\alpha(t)$, *the rotor*, for U(1)_M invariance. The U(1)_M transformation reads $\psi_k(t, x) \rightarrow e^{iq(x)\beta} \psi_k(t, x)$ with $q(x) = \frac{1}{2} \text{sgn}(x)$. The transformation parameter β is spacetime independent because of the zero U(1)_M gauge coupling. Yet, $q(x)$ being x dependent, the right-mover’s kinetic term $\psi_k^\dagger i(\partial_t + \partial_x) \psi_k$ is not U(1)_M invariant, so an interaction $-q'(x) \alpha(t) \psi_k^\dagger \psi_k(t, x)$ is added with $\alpha(t) \rightarrow \alpha(t) - \beta$ under U(1)_M. This then implies that the only U(1)_M-invariant quadratic term for the rotor is $I[\dot{\alpha}(t)]^2/2$ with a “moment of inertia” I .

¹ SU(4)_F-violating corrections to this result by continuous parameters, which could be imagined as small as we wish, would not fundamentally solve the problem, so by ignoring SM couplings we are focussing on the heart of the puzzle.

The fermion-rotor action \mathcal{S} thus reads

$$\mathcal{S} = \int dt \frac{I}{2} \dot{\alpha}^2 + \int dt dx \sum_{k=1}^N \psi_k^\dagger i \mathcal{D} \psi_k + \mathcal{S}_{ct} \quad (3)$$

with $\mathcal{D} \equiv \partial_t + \partial_x + iq'(x)\alpha(t)$ and \mathcal{S}_{ct} is a counterterm associated with regularization of the singular product $\psi^\dagger(t,x)\psi(t,x)$.² Since $q'(x) \propto \delta(x)$, the fermion-rotor interaction is localized at the monopole core. The conserved $U(1)_M$ charge Q is given by

$$Q = I\dot{\alpha}(t) + \sum_{k=1}^N \int dx q(x) \psi_k^\dagger \psi_k(t, x), \quad (4)$$

so the rotor carries $U(1)_M$ charge $I\dot{\alpha}$. Since $[\alpha(t), I\dot{\alpha}(t)] = i$, any rotor state with a definite $U(1)_M$ charge has indefinite α , so the rotor must be treated fully quantum mechanically.

The following hierarchy of scales is important. If a fermion deposited an $O(1)$ charge on the rotor ($I\dot{\alpha} \sim 1$), the rotor would get excited with energy of $O(I\dot{\alpha}^2) \sim O(1/I)$, which would correspond to a dyon. But we are in the low energy regime with only fermions and no dyon in the spectrum, so

$$\frac{1}{I} \gg E, \quad (5)$$

where E is the energy scale of a given scattering process. The low energy limit (5) is equivalent to a small I limit, but we must be careful with the order of limits. Since the dyon energy ($\sim 1/I$) is of $O(e^2/r_0)$ with the $U(1)_M$ gauge coupling e , the $e \rightarrow 0$ limit is consistent only if

$$I \gg r_0. \quad (6)$$

We thus cannot take $I \rightarrow 0$ until nonzero r_0 becomes unnecessary to regulate singularity. Moreover, we cannot take $r_0 \rightarrow 0$ until the UV cutoff ε_{uv}^{-1} (e.g., for point splitting) becomes unnecessary, because $r_0 \gg \varepsilon_{uv}$.

While $U(1)_M$ and $SU(N)_F$ are exact, a global $U(1)_\psi$ for the ‘‘total fermion number’’ is anomalous due to the rotor. Under a $U(1)_\psi$ transformation:

$$(\psi_1, \dots, \psi_N) \longrightarrow e^{i\gamma}(\psi_1, \dots, \psi_N) \quad (7)$$

with a parameter γ , the path-integral measure changes:

$$[d\psi][d\psi^\dagger] \longrightarrow [d\psi][d\psi^\dagger] \exp\left[-\frac{iN}{2\pi} \int d^2x \gamma q'(x) \dot{\alpha}(t)\right]. \quad (8)$$

² [18] uses point splitting and $\mathcal{S}_{ct} \propto \int dt [\alpha(t)]^2$. This \mathcal{S}_{ct} violates $U(1)_M$ and by design cancels $U(1)_M$ violation from point splitting, like the photon mass counterterm in a QED with a momentum cutoff instead of a gauge invariant regulator.

This leads to an anomalous conservation law for $U(1)_\psi$,

$$(\partial_t + \partial_x) \sum_{k=1}^N J_k(t, x) = \frac{N}{2\pi} q'(x) \dot{\alpha}(t), \quad (9)$$

where $J_k(t, x) \equiv \psi_k^\dagger \psi_k(t, x)$. Thus, the total fermion number is not conserved at the core (as $q'(x) \propto \delta(x)$) whenever nonzero $U(1)_M$ charge ($\dot{\alpha} \neq 0$) is on the rotor.

The Solution:

The semitonic case

Take $N > 2$ and consider an initial state consisting of one χ_i^- particle incident on M .³ This is *semitonic*—the $U(1)_M$ and $SU(N)_F$ conservation would predict fractional fermion numbers in units of $2/N$ in the final state. We find, however, that *the χ_i^- will just propagate freely across the monopole core*. But how could conservation laws be wrong? How can there be an outgoing χ_i^- when the equation of motion forces χ_i^- to be ingoing?

The culprit is the assumption that a k -fermion in the final state must be created by $(\chi_k^+)^\dagger$ for it to be outgoing. This is incorrect. We may multiply χ_k^+ by any functional of the rotor. An especially interesting functional is $e^{i\alpha(t)}$, which transforms under $U(1)_M$ as $e^{i\alpha(t)} \rightarrow e^{-i\beta} e^{i\alpha(t)}$, thereby carrying $U(1)_M$ charge -1 .

This inspires us to consider $\mathcal{O} \equiv e^{i\alpha(\tau)\theta(x)} \psi_k(t, x)$, which has $U(1)_M$ charge $-\frac{1}{2}$ everywhere at all x . When $x < 0$ we have $\mathcal{O} = \chi_k^-$ (see (2)) so \mathcal{O}^\dagger creates an ingoing χ_k^- fermion. At $x > 0$ we have $\mathcal{O} = e^{i\alpha} \chi_k^+$, so \mathcal{O}^\dagger creates some state with $U(1)_M$ charge $-\frac{1}{2}$.

To motivate what value of τ should be used in \mathcal{O} , pretend temporarily that the rotor were a classical background. Then, the fermion equation of motion, $i(\partial_t + \partial_x)\psi(t, x) = q'(x)\alpha(t)\psi(t, x)$, would have a solution of the form $\psi(t, x) = e^{-i\alpha(t-x)\theta(x)} \psi_0(t, x)$, where ψ_0 is a free right-mover, $(\partial_t + \partial_x)\psi_0(t, x) = 0$. This suggests that \mathcal{O} may exactly interpolate a 1-particle state if we choose $\tau = t - x$.

We are thus led to consider the composite operator

$$\Psi_k(t, x) \equiv e^{i\alpha(t-x)\theta(x)} \psi_k(t, x). \quad (10)$$

Our main result is that Ψ_k indeed exactly interpolates a 1-particle state. By a direct calculation to be described in detail shortly, we find

$$\left\langle \Psi_k(t, x) [\Psi_{k'}(t', x')]^\dagger \right\rangle = G_0(t - t', x - x') \delta_{kk'}, \quad (11)$$

where $\langle \dots \rangle \equiv \langle 0 | \hat{T} \{ \dots \} | 0 \rangle$ (with an important clarification about the vacuum to be mentioned later), and

³ We will refer to such a state as a ‘‘1-particle state’’, not counting M (nor the rotor) as a particle. Also remember that our ‘‘particle’’ is only the lowest partial wave.

$G_0(t, x)$ is the free propagator of a massless right-mover:

$$G_0(t, x) = \frac{1}{2\pi i} \frac{t + x}{t^2 - x^2 - i0^+}. \quad (12)$$

Hence Ψ_k exactly interpolates just a massless right-moving particle and no other states (e.g., a left-moving state, a massive state, multi-particle states). To reveal this particle's identity, notice

$$\Psi_k(t, x) = \begin{cases} \chi_k^-(t, r)|_{r=-x} & \text{if } x < 0, \\ e^{i\alpha(t-x)} \chi_k^+(t, r)|_{r=x} & \text{if } x > 0. \end{cases} \quad (13)$$

Thus, at $x < 0$, $[\Psi_k(t, x)]^\dagger$ creates a χ_k^- particle by definition. Then, $\Psi_k(t, x)$ at $x > 0$ must annihilate the same χ_k^- particle as Ψ_k has the exact free 2-point function (11) and no other nonzero 2-point correlators. (This is essentially the definition of “the same particle”.) Hence, $[e^{i\alpha(t-x)} \chi_k^+(t, x)]^\dagger$ creates a χ_k^- at $x > 0$, which, being at $x > 0$, is outgoing!

Similarly, we define $\tilde{\Psi}_k(t, x) \equiv e^{-i\alpha(t-x)\theta(-x)} \psi_k(t, x)$, which is $e^{-i\alpha(t-x)} \chi_k^-(t, -x)$ for $x < 0$ and $\chi_k^+(t, x)$ for $x > 0$. It also exactly has a unique free 2-point function, so the composite operator $[e^{-i\alpha(t-x)} \chi_k^-(t, -x)]^\dagger$ at $x < 0$ creates the same particle to be annihilated by $\chi_k^+(t, x)$ at $x > 0$. That is, $[e^{-i\alpha(t+r)} \chi_k^-(t, r)]^\dagger$ creates an ingoing χ_k^+ !

Therefore, in spite of the equation of motion missing half of the states, we actually have a full 1-particle spectrum after all. That is, for *each* of the lower and upper components of every χ_k doublet, there are *both* ingoing and outgoing states. The states missed by the equation of motion—outgoing χ_k^- and ingoing χ_k^+ —are interpolated by composite operators $e^{i\alpha(t-r)} \chi_k^+(t, r)$ and $e^{-i\alpha(t+r)} \chi_k^-(t, r)$, respectively.

Since the *exact* 2-point correlators are free, the corresponding 1-particle states in the momentum space are eigenstates of the *full* hamiltonian, i.e., stationary states, in (1+1)D. That is, in (3+1)D, *the lowest partial wave of a 1-particle state incident on M propagates freely*. “Free propagation” is not to be confused with “forward scattering”. The latter is the “ $f=i$ ” part of T in $S = \mathbb{1} + iT$, while we have $S = \mathbb{1}$ in the subspace of 1-particle states in the lowest partial wave channel.

We now provide a proof of (11). In [18] Polchinski performs exact path integration over the fermions [Eq. (48) of [18]]. Translating his Euclidean calculation to Lorentzian, his result reads

$$\left\langle \prod_{i=1}^n \psi_{k_i}(t_i, x_i) \prod_{j=1}^{n'} [\psi_{k'_j}(t'_j, x'_j)]^\dagger \right\rangle = \mathcal{Z} W_0, \quad (14)$$

where W_0 is the sum of products of free propagators one

would expect if all the fermions were literally free, and

$$\mathcal{Z} \equiv \int [d\alpha] \exp \left[- \int_\mu^\Lambda \frac{d\omega}{2\pi} \left\{ \frac{N\omega}{4\pi} \alpha(\omega) \alpha(-\omega) + iA\alpha(\omega) - iB\alpha(-\omega) \right\} \right] \quad (15)$$

$$= \exp \left[\frac{2}{N} \int_\mu^\Lambda \frac{d\omega}{\omega} AB \right] \quad (16)$$

$$A = \sum_{i=1}^n \theta(x_i) e^{-i\omega(t_i - x_i)} - \sum_{j=1}^{n'} \theta(x'_j) e^{-i\omega(t'_j - x'_j)}, \quad (17)$$

$$B = \sum_{i=1}^n \theta(-x_i) e^{i\omega(t_i - x_i)} - \sum_{j=1}^{n'} \theta(-x'_j) e^{i\omega(t'_j - x'_j)},$$

where μ is an IR cutoff to be taken to zero (which is subtle when $n \neq n'$ as we will see below), while $\Lambda \sim 1/I$ is taken to infinity as per (5), and $r_0 \rightarrow 0$ has been taken. In (15) the path integration over the rotor is undone so that we can insert our $e^{\pm i\alpha}$.

We are now ready to prove (11). We also want to show that the $e^{i\alpha}$ insertion time must be $t - x$ as in (10). So, let's consider $\langle e^{i\alpha(\tau)\theta(x)} \psi_k(t, x) [e^{i\alpha(\tau')\theta(x')} \psi_{k'}(t', x')]^\dagger \rangle$ with some τ, τ' . This just amounts to adding $i\alpha(\tau)\theta(x) - i\alpha(\tau')\theta(x')$ to the exponent in (15). This then modifies A and B (with $n = n' = 1$ and dropping i, j) as

$$\begin{aligned} A &= A_{\text{from (17)}} - \theta(x) e^{-i\omega\tau} + \theta(x') e^{-i\omega\tau'}, \\ B &= B_{\text{from (17)}} + \theta(x) e^{i\omega\tau} - \theta(x') e^{i\omega\tau'}. \end{aligned} \quad (18)$$

Hence, A vanishes identically if and only if we choose $\tau = t - x$ and $\tau' = t' - x'$, which via (16) implies $\mathcal{Z} = 1$. This proves our result (11) and demonstrates the uniqueness of the time $t - x$ in (10). The same proof works also for $\tilde{\Psi}_k$ except that it is B that vanishes this time.

Naively the proof seems generalizable for any n and n' . That is true for $n = n'$, so $2n$ -point functions of n Ψ 's and n Ψ^\dagger 's are indeed free. For $n \neq n'$, however, the IR cutoff has a subtlety that invalidates (14). As discussed in [18], the IR cutoff removes low frequency modes in α and hence restricts α to having the same value before and after the reaction. But the $U(1)_\psi$ anomaly (9) relates the change of the total fermion number, ΔN_ψ , to the change of α as $\Delta N_\psi = \frac{N}{2\pi} \Delta\alpha$. Therefore, the IR cutoff does not work when $\Delta N_\psi \neq 0$ such as when $n \neq n'$. This subtlety will be especially important below.

The non-semiionic case

We now turn our attention to *non-semiionic* processes. We confirm the so-called Callan-Rubakov (CR) processes and show how that is consistent with free propagation we found above.

We begin with $N = 2$. Let's start with an ingoing χ_1^- incident on M . If we require the final state to be created by $[\chi_{1,2}^+]^\dagger$ only, $U(1)_M$ and $SU(2)_F$ fix the final state to be an outgoing $\overline{\chi_2^+}$ (“anti- χ_2^+ ”). We thus seem to have $\chi_1^- \rightarrow \overline{\chi_2^+}$ (CR). But from (11) we seem to have free $\chi_1^- \rightarrow \chi_1^-$ propagation, with the outgoing χ_1^- created by $[e^{i\alpha}\chi_1^+]^\dagger$.

Free propagation cannot have two different final states, so the outgoing χ_1^- and $\overline{\chi_2^+}$ must be one and the same. That is, we *should* find $\langle \Psi_1(t, x) \psi_2(t', x') \rangle \propto G_0(t - t', x - x')$ for $x, x' > 0$ so the operators $[e^{i\alpha}\chi_1^+]^\dagger (= \Psi_1^\dagger$ at $x > 0$) and χ_2^+ ($= \psi_2$ at $x' > 0$) do create one and the same outgoing particle. To show this, note that this correlator is one of those for which (14) is invalid because $n = 2 \neq 0 = n'$. So, we employ a cluster analysis method similar to one in [18]. Consider the correlator

$$\langle \Psi_i(1) \psi_j(2) \Psi_k^\dagger(3) \psi_\ell^\dagger(4) \rangle, \quad (19)$$

where “1” \equiv “ t_1, x_1 ”, etc., with all $x_{1,2,3,4} > 0$. This correlator is free from the IR subtlety as $n = n' = 2$. So, we start with (14) and insert $e^{i\alpha}$ at “1” and “3” as similarly in (18), then we find the correlator (19) to be

$$(\delta_{ik}\delta_{j\ell} \langle 13 \rangle \langle 24 \rangle - \delta_{i\ell}\delta_{jk} \langle 14 \rangle \langle 23 \rangle) \frac{s_{23}s_{14}}{s_{12}s_{34}}, \quad (20)$$

where $s_{ab} \equiv t_a - t_b - (x_a - x_b)$, $\langle ab \rangle \equiv G_0(t_a - t_b, x_a - x_b) = 1/(2\pi i s_{ab})$. Now, we take $t_1 \sim t_2 \sim t_3 + T \sim t_4 + T$ with large $|T|$. In the $|T| \rightarrow \infty$ limit, the correlator (19) must be clustered as $\langle \Psi_i(1) \psi_j(2) \rangle \langle \Psi_k^\dagger(3) \psi_\ell^\dagger(4) \rangle$. Equating this with the $|T| \rightarrow \infty$ limit of (20), we find

$$\langle \Psi_i(1) \psi_j(2) \rangle = e^{i\vartheta} \epsilon_{ij} \langle 12 \rangle \quad (21)$$

for $x_1, x_2 > 0$, where ϑ is a parameter interpreted as characterizing degenerate vacua. (See [18] for a more discussion on the vacuum degeneracy.) Thus, the outgoing χ_1^- and $\overline{\chi_2^+}$ are indeed one and the same particle.

The identity of this one-and-the-same particle is a $\overline{\chi_2^+}$, not a χ_1^- . The operator χ_2^+ creates a $\overline{\chi_2^+}$ by definition. We do not know what a composite operator $[e^{i\alpha}\psi_1]^\dagger$ creates except that it creates the same particle as $\overline{\chi_2^+}$ does! We thus confirm the known CR result that $\chi_1^- \rightarrow \overline{\chi_2^+}$ with a unit probability.

For $N > 2$, the 2-point functions similar to (21) vanish by $SU(N)_F$ as there are no invariant 2nd-rank antisymmetric tensors like ϵ_{ij} . So, the above situation is unique to $N = 2$ and in particular does not invalidate the particle identification made just below (13). There, we have no potential alternatives like the $\overline{\chi_2^+}$ for the final particle.

But it is now clear that initial states with $N/2$ fermions for $N > 2$ are analogous to the 1-particle states for $N = 2$, because the invariant tensor $\epsilon_{i_1 i_2 \dots i_N}$ permits us to write down an N -point version of (21). For example, consider $N = 4$ with (1) and start with $u^1 + u^2$ incident

on M . We then have either $u^1 + u^2 \rightarrow \bar{e} + \bar{d}^3$ (CR) with the half of the spectrum, or double free propagation $u^1 \rightarrow u^1$ and $u^2 \rightarrow u^2$ with the full spectrum.

When the CR occurs, the two outgoing 2-particle states, $\bar{e} + \bar{d}^3$ and $u^1 + u^2$, must be one and the same. A cluster analysis again shows that is indeed the case. Like (19) we consider

$$\langle \Psi_i(1) \Psi_j(2) \psi_k(3) \psi_\ell(4) \Psi_{i'}^\dagger(5) \Psi_{j'}^\dagger(6) \psi_{k'}^\dagger(7) \psi_{\ell'}^\dagger(8) \rangle \quad (22)$$

with $x_1, \dots, x_8 > 0$, and then pull apart the 1234 cluster from the 5678. Using (14) and inserting $e^{i\alpha}$ like in (18), we find the $N = 4$ version of (21):

$$\langle \Psi_i(1) \Psi_j(2) \psi_k(3) \psi_\ell(4) \rangle = \epsilon_{ijkl} \sqrt{\langle 13 \rangle \langle 23 \rangle \langle 14 \rangle \langle 24 \rangle} \quad (23)$$

up to a phase. When the group of times $t_{1,2}$ are well in the future of $t_{3,4}$, (23) computes the overlap between the state created by an operator-cluster $[\Psi_i(1) \Psi_j(2)]^\dagger$ and that by $\psi_k(3) \psi_\ell(4)$. Since all $x_{1,2,3,4} > 0$, we are studying how similar those two *outgoing* states are to each other. The key is that those two outgoing 2-particle states consist of the same two species when (23) factorizes into free propagators.

Let's choose $(i, j, k, \ell) = (3, 4, 1, 2)$ so the first cluster is $[U^1(1)U^2(2)]^\dagger$ (capitals for $e^{i\alpha}$ insertion), creating outgoing $u^1 + u^2$, while the second is $d^3(3)e(4)$, creating outgoing $\bar{e} + \bar{d}^3$. While (23) does not seem free propagators due to the square root, it *is* when “*intra-cluster separation*” \ll “*inter-cluster separation*”. When 1 and 2 both approach 5 while 3, 4 \rightarrow 6, we have $\sqrt{\langle 13 \rangle \langle 23 \rangle \langle 14 \rangle \langle 24 \rangle} \rightarrow \langle 56 \rangle^2$ so $\langle U^1(1)U^2(2) d^3(3)e(4) \rangle \rightarrow \langle 56 \rangle^2$. This is identical (up to a phase) to $\langle [\psi_a(1)\psi_b(2)]^\dagger \psi_a(3)\psi_b(4) \rangle \rightarrow \langle 56 \rangle^2$ from (14). Therefore, $u^1 + u^2$ and $\bar{e} + \bar{d}^3$ are indeed one and the same outgoing 2-particle state when the two particles are “together”. By the same argument as made for $\overline{\chi_2^+}$, the identity of this outgoing 2-particle state is $\bar{e} + \bar{d}^3$.

Discussions: Directly from the path integral of the rotor-fermion system, we have shown two things: semitonic processes are free propagation, and non-semi-tonic CR processes are confirmed and do not contradict free propagation. Similar pictures were recently suggested in [14, 15] in a very different language of boundary CFT, topological line operators, etc. The fermion-rotor theory would formally be a CFT if we had set $I = 0$ (hence also $r_0 = 0$ as per (5)) but it is essential that the rotor is a dynamical degree of freedom with a kinetic energy. As lucidly discussed in [18], the rotor's kinetic term (or whatever gauge equivalent to it) must be included in order for the limit of zero $U(1)_M$ gauge coupling to be consistent with the semi-classical expansion of path integral of the monopole-fermion system at low energy. Hence, the relation between the two approaches seems not obvious, which is an interesting question worth investigating.

An inevitable next question is to find the (3+1)D counterpart of our $e^{\pm i\alpha}$ insertions. In particular it would be

satisfying to establish a connection between our $e^{\pm i\alpha}$ factors and the abelian vortices with a fractional winding number in [16], which is expected to be also connected to the twisted sectors in [14, 15].

For phenomenology it is imperative to study the implications of the finding that semitonic processes are actually free propagation, especially in the context of monopole-catalyzed nucleon decay, which places by far the strongest bound on the monopole abundance by many orders of magnitude [20, 21]. Traditionally it was thought that, once a monopole enters a nucleon, the nucleon would decay 100% via non-semitonic or (intermediate) semitonic process. Now that the latter is replaced by free propagation, monopole-catalyzed nucleon decay rates should go down. But by how much? What is “free propagation” under confinement? How do $SU(N)_F$ violations from chiral symmetry breaking and quark masses affect our analysis? When we integrate out heavy quarks, what operators do we generate involving the rotor? These are all very interesting questions we save for our future work.

Acknowledgment: TO is supported in part by the US DOE grant DE-SC0010102 and in part by the JSPS Grant-in-Aid for Scientific Research, No. 21H01086. VL is supported by the STFC under Grant No. ST/T000864/1.

* vazha.loladze@physics.ox.ac.uk

† tokui@fsu.edu

- [1] Y. Kazama, C. N. Yang, and A. S. Goldhaber, “Scattering of a Dirac Particle with Charge Ze by a Fixed Magnetic Monopole,” *Phys. Rev. D* **15** (1977) 2287–2299.
- [2] P. Rossi, “Spin 1/2 Particles in the Field of Monopoles,” *Nucl. Phys. B* **127** (1977) 518–536.
- [3] C. J. Callias, “Spectra of Fermions in Monopole Fields: Exactly Soluble Models,” *Phys. Rev. D* **16** (1977) 3068.
- [4] A. S. Goldhaber, “Dirac Particle in a Magnetic Field: Symmetries and their Breaking by Monopole Singularities,” *Phys. Rev. D* **16** (1977) 1815.
- [5] V. A. Rubakov, “Adler-Bell-Jackiw Anomaly and Fermion Number Breaking in the Presence of a Magnetic Monopole,” *Nucl. Phys. B* **203** (1982) 311–348.
- [6] C. G. Callan, Jr., “Disappearing Dyons,” *Phys. Rev. D* **25** (1982) 2141.
- [7] C. G. Callan, Jr., “Dyon-Fermion Dynamics,” *Phys. Rev. D* **26** (1982) 2058–2068.
- [8] C. G. Callan, Jr., “Monopole Catalysis of Baryon Decay,” *Nucl. Phys. B* **212** (1983) 391–400.
- [9] R. Kitano and R. Matsudo, “Missing final state puzzle in the monopole-fermion scattering,” *Phys. Lett. B* **832** (2022) 137271, [arXiv:2103.13639 \[hep-th\]](#).
- [10] C. Csáki, Y. Shirman, O. Telem, and J. Terning, “Pairwise Multiparticle States and the Monopole Unitarity Puzzle,” *Phys. Rev. Lett.* **129** no. 18, (2022) 181601.
- [11] Y. Hamada, T. Kitahara, and Y. Sato, “Monopole-fermion scattering and varying Fock space,” *JHEP* **11** (2022) 116, [arXiv:2208.01052 \[hep-th\]](#).
- [12] V. V. Khoze, “Scattering amplitudes of fermions on monopoles,” *JHEP* **11** (2023) 214, [arXiv:2308.09401 \[hep-th\]](#).
- [13] T. D. Brennan, “A New Solution to the Callan Rubakov Effect,” [arXiv:2309.00680 \[hep-th\]](#).
- [14] M. van Beest, P. Boyle Smith, D. Delmastro, Z. Komargodski, and D. Tong, “Monopoles, Scattering, and Generalized Symmetries,” [arXiv:2306.07318 \[hep-th\]](#).
- [15] M. van Beest, P. Boyle Smith, D. Delmastro, R. Mouland, and D. Tong, “Fermion-Monopole Scattering in the Standard Model,” [arXiv:2312.17746 \[hep-th\]](#).
- [16] C. Csáki, R. Ovadia, O. Telem, J. Terning, and S. Yankielowicz, “Abelian Instantons and Monopole Scattering,” [arXiv:2406.13738 \[hep-th\]](#).
- [17] S. Bolognesi, B. Bucciotti, and A. Luzio, “The Monopole-Fermion Problem in a Chiral Gauge Theory (the $\psi\chi\eta$ Model),” [arXiv:2406.15552 \[hep-th\]](#).
- [18] J. Polchinski, “Monopole Catalysis: The Fermion-Rotor System,” *Nucl. Phys. B* **242** (1984) 345–363.
- [19] E. Witten, “An $SU(2)$ Anomaly,” *Phys. Lett. B* **117** (1982) 324–328.
- [20] **Super-Kamiokande** Collaboration, K. Ueno *et al.*, “Search for GUT monopoles at Super-Kamiokande,” *Astropart. Phys.* **36** (2012) 131–136, [arXiv:1203.0940 \[hep-ex\]](#).
- [21] **IceCube** Collaboration, M. G. Aartsen *et al.*, “Search for non-relativistic Magnetic Monopoles with IceCube,” *Eur. Phys. J. C* **74** no. 7, (2014) 2938, [arXiv:1402.3460 \[astro-ph.CO\]](#). [Erratum: *Eur.Phys.J.C* 79, 124 (2019)].