Hemant K. Mishra^{*}[†]

Abstract. Williason's theorem states that if A is a $2n \times 2n$ real symmetric positive definite matrix then there exists a $2n \times 2n$ real symplectic matrix M such that $M^T AM = D \oplus D$, where D is an $n \times n$ diagonal matrix with positive diagonal entries known as the symplectic eigenvalues of A. The theorem is known to be generalized to $2n \times 2n$ real symmetric positive semidefinite matrices whose kernels are symplectic subspaces of \mathbb{R}^{2n} , in which case, some of the diagonal entries of D are allowed to be zero. In this paper, we further generalize Williamson's theorem to $2n \times 2n$ real symmetric matrices by allowing the diagonal elements of D to be any real numbers, and thus extending the notion of symplectic eigenvalues to real symmetric matrices. Also, we provide an explicit description of symplectic eigenvalues, construct symplectic matrices achieving Williamson's theorem type decomposition, and establish perturbation bounds on symplectic eigenvalues for a class of $2n \times 2n$ real symmetric matrices denoted by EigSpSm(2n). The set EigSpSm(2n) contains $2n \times 2n$ real symmetric positive semidefinite whose kernels are symplectic subspaces of \mathbb{R}^{2n} . Our perturbation bounds on symplectic eigenvalues for EigSpSm(2n) generalize known perturbation bounds on symplectic eigenvalues of positive definite matrices given by Bhatia and Jain [J. Math. Phys. 56, 112201 (2015)].

Contents							
1	Introduction 2						
	1.1	Williamson's theorem	2				
	1.2	Literature review	2				
	1.3	Main contributions	3				
	1.4	Paper organization	4				
2	Review and miscellanea						
	2.1	Matrices	5				
	2.2	Linear algebra on \mathbb{R}^n	5				
	2.3	Symplectic linear algebra on \mathbb{R}^{2n}	5				
3	Williamson's theorem for symmetric matrices						
4	General Williamson's theorem via symplectic orthogonal projec-						
	tion		10				
5	Explicit Williamson's decomposition for a subset of $SpSm(2n)$ 1						
	5.1 Description of symplectic eigenvalues for $EigSpSm(2n)$						

²⁰²⁰ Mathematics Subject Classification: 15B48, 15A18, 15A20, 15A23.

Keywords: Williamson's theorem, symplectic eigenvalue, symplectic matrix, real symmetric matrix, perturbation bound, eigenvalues, symplectic orthogonal projection.

^{*}School of Electrical and Computer Engineering, Cornell University, Ithaca, New York 14850, USA [†]hemant.mishra@cornell.edu

5.2	5.2 Description of symplectic matrices in Williamson's decompo			
	sition for $\operatorname{EigSpSm}(2n)$	15		
5.3	Perturbation bounds on symplectic eigenvalues for $EigSpSm(2n)$	17		
References				

1 Introduction

Williamson's theorem contains germs of modern developments in symplectic topology. It facilitates an immediate proof of Gromov's non-squeezing theorem in the linear case [1], which is one of the most important theorems in symplectic geometry. Also known as Williamson's decomposition, the theorem is fundamental in developing the theory of bosonic Gaussian states in quantum information [2, 3, 4, 5]. In the recent years, Williamson's theorem has attracted much attention of mathematicians and physicists, and it has become a topic of intense study in matrix analysis [6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17], operator theory [18, 19, 20], and quantum physics [21, 22, 23, 4, 24].

1.1 Williamson's theorem

A $2n \times 2n$ real matrix M is called a symplectic matrix if it satisfies $M^T J_{2n} M = J_{2n}$, where $J_{2n} := \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$, I_n being the identity matrix of size n. Williamson's theorem [25] states that for every $2n \times 2n$ real symmetric positive definite matrix A, there exists a symplectic matrix M such that

$$M^{T}AM = \begin{pmatrix} D & 0\\ 0 & D \end{pmatrix}, \tag{1.1}$$

where *D* is an $n \times n$ diagonal matrix with unique positive diagonal entries (up to ordering), called the symplectic eigenvalues of *A*. Several elementary proofs of Williamson's theorem are available in the literature. See [26, 27, 28].

1.2 Literature review

In his original work [25], Williamson showed that for any $2n \times 2n$ real symmetric matrix A there exists a symplectic matrix M such that $M^T AM$ is a (non-diagonal) sparse matrix. In general, $M^T AM$ may not be a diagonal matrix for any symplectic matrix M much less a diagonal matrix of the form $D \oplus D$ for some $n \times n$ diagonal matrix D. See the corollary of Theorem 2 in [25]. Interestingly, if A is positive definite, then it is congruent to a diagonal matrix via a symplectic matrix as stated in (1.1).

Williamson's theorem is known to be generalized to $2n \times 2n$ real symmetric positive semidefinite matrices whose kernels are *symplectic subspaces* of \mathbb{R}^{2n} . More specifically, for a $2n \times 2n$ real symmetric positive semidefinite matrix A there exists a symplectic matrix M such that $M^T AM = D \oplus D$ for some $n \times n$ diagonal matrix D with nonnegative diagonal entries if and only if the kernel of A is a symplectic subspace of \mathbb{R}^{2n} . This was stated in [12, Remark 2.6], and explicitly proved in [29, Theorem 1.3.5]. Also, a constructive proof of this extension was recently given in [17]. Cruz and Faßbender [30] established simple algebraic conditions on $2n \times 2n$ complex matrices that are diagonalizable by symplectic equivalence, similarity, or congruence. In particular, Theorem 21 of [30] states that for a $2n \times 2n$ (complex) matrix A there exists a (complex) symplectic matrix M such that $M^T A M$ is a diagonal matrix if and only if A is symmetric and $AJ^T AJ$ is diagonalizable.

To the best of our knowledge, no precise condition is known for $2n \times 2n$ real symmetric matrices to be diagonalizable in the sense of Williamson's theorem. The main aim of this work is to fill this gap.

1.3 Main contributions

In this paper, we establish explicit necessary and sufficient conditions on $2n \times 2n$ real symmetric matrices to be diagonalizable in the sense of Williamson's theorem, and also investigate several implications of it.

- We show that for a $2n \times 2n$ real symmetric matrix A there exists a symplectic matrix M such that $M^T A M = D \oplus D$ where D is an $n \times n$ real diagonal matrix (unique up to ordering of its diagonal entries) if and only if there exist symplectic subspaces $\mathcal{W}_-, \mathcal{W}_0, \mathcal{W}_+$ of \mathbb{R}^{2n} with dimensions $v(A), \xi(A), \pi(A)$, respectively such that
 - \mathcal{W}_{-} , \mathcal{W}_{0} , \mathcal{W}_{+} are pairwise symplectically orthogonal to each other
 - these subspaces are invariant under $J_{2n}A$,
 - A is negative definite on \mathcal{W}_{-} , the kernel of A is \mathcal{W}_{0} , and A is positive definite on \mathcal{W}_{+} .

Here $v(A), \xi(A), \pi(A)$ denote the number of negative eigenvalues, zero eigenvalues, positive eigenvalues, respectively. See Theorem 3.1.

- We introduce a symplectic analog of orthogonal projection, called *symplectic orthogonal projection*, in Definition 4.1, and discuss some properties of it. Symplectic orthogonal projections can be of independent interest in symplectic geometry. We then re-state the aforementioned result, Theorem 3.1, in terms of symplectic orthogonal projection. See Proposition 4.4. This then leads to a more explicit description of the diagonal form in the generalized Williamson's theorem. See Proposition 4.5.
- We construct explicit Williamson's decomposition and establish perturbation bounds for the diagonal form for a class of $2n \times 2n$ real symmetric matrices. This class, denote by EigSpSm(2n), consists of $2n \times 2n$ real symmetric matrices whose eigenspaces corresponding to negative eigenvalues, zero eigenvalues, and positive eigenvalues form symplectic subspaces of \mathbb{R}^{2n} satisfying the three conditions mentioned above. In particular, EigSpSm(2n) contains the set of $2n \times 2n$ real positive semidefinite matrices with symplectic kernel. The perturbation bounds we obtain generalize known perturbation bounds on symplectic eigenvalues of positive definite matrices given by Bhatia and Jain [6, Theorem 6]. See Section 5.

Symbol	Meaning	Definition
$M(n, k)$ $M(n)$ $Sm(n)$ $Psd(n)$ $Pd(n)$ $I_n \text{ or } I$ $Or(n)$ $ker(A)$ $ran(A)$ $null(A)$ $rank(A)$	set of $n \times k$ real matrices set of $n \times n$ real matrices set of symmetric matrices set of positive semidefinite matrices set of positive definite matrices identity matrix of size n orthogonal group kernel of $A \in M(n, k)$ range of $A \in M(n, k)$ dimension of ker (A) dimension of ran (A)	$M(n, n)$ $\{A \in M(n) : A^{T} = A\}$ $\{A \in Sm(n) : x^{T}Ax \ge 0 \forall x \in \mathbb{R}^{n}\}$ $\{A \in Sm(n) : x^{T}Ax > 0 \forall x \in \mathbb{R}^{n} \setminus \{0\}\}$ $\{U \in M(n) : U^{T}U = I_{n}\}$ $\{x \in \mathbb{R}^{k} : Ax = 0\}$ $\{Ax : x \in \mathbb{R}^{k}\}$
$J_{2n} \text{ or } J$ $Sp(2n, 2k)$ $Sp(2n)$ $OrSp(2n)$ $SpSm(2n)$ $SpPsd(2n)$ $EigSpSm(2n)$	standard symplectic matrix real symplectic group real orthosymplectic group	$ \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} $ $ \{ M \in \mathcal{M}(2n, 2k) : M^T J_{2n} M = J_{2k} \} $ Sp(2n, 2n) $ Or(2n) \cap Sp(2n) $ defined after Remark 3.2 $ \{ A \in Psd(2n) : ker(A) \cap ker(A)^{\perp_s} = \{ 0 \} \} $ defined in Section 5

Table 1: Summary of notations and their mathematical definitions.

1.4 Paper organization

We review some basic theory of matrices, linear algebra, and symplectic linear algebra in Section 2: Subsection 2.1 contains useful concepts from matrix analysis; Subsection 2.2 recalls basic theory of subspaces of the Euclidean space \mathbb{R}^n ; Subsection 2.3 revisits some basic theory of standard symplectic space \mathbb{R}^{2n} , and establishes some symplectic operations that are useful for the development of the paper.

We state and prove the main result in Section 3 (Theorem 3.1) along with an interesting corollary (Corollary 3.3).

In Section 4, we introduce a symplectic analog of the well-known orthogonal projection called symplectic orthogonal projection (Definition 4.1), and re-state the main result in terms of the symplectic orthogonal projection (Proposition 4.4).

Lastly, in Section 5, we study the Williamson's decomposition for EigSpSm(2n): explicit description of the symplectic eigenvalues for matrices in EigSpSm(2n) is given in Subsection 5.1, construction of symplectic matrices achieving the Williamson's decomposition is given in Subsection 5.2, and perturbation bounds on the symplectic eigenvalues are established in Subsection 5.3.

2 Review and miscellanea

In this section, we establish some notations, and briefly recall some basic concepts from matrix analysis, linear algebra, and symplectic linear algebra. We refer the reader to [31, 32] for a comprehensive account of theory of matrices, [33] for linear algebra, and [26,

34] for symplectic linear algebra. A summary of notations with mathematical definitions is provided in Table 1.

2.1 Matrices

Let M(n, k) denote the set of $n \times k$ real matrices. We use the shorthand M(n) for M(n, n). We denote by Sm(n) the subset of M(n) consisting of symmetric matrices. For $A \in Sm(n)$, we shall use the notations v(A), $\xi(A)$, $\pi(A)$ to denote the number of negative eigenvalues, zero eigenvalues, positive eigenvalues of A, respectively. If $K \in M(n)$ is an invertible matrix then the Sylvester's law of inertia states that for any $A \in Sm(n)$, we have $v(A) = v(K^T A K)$, $\xi(A) = \xi(K^T A K)$, and $\pi(A) = \pi(K^T A K)$. See [32, Theorem 4.5.8].

We denote by Psd(n) and Pd(n) the subsets of Sm(n) consisting of positive semidefinite and positive definite matrices, respectively. Let Or(n) denote the set of $n \times n$ real orthogonal matrices, also known as the orthogonal group. For $A \in M(n, k)$, let ker(A)denote the kernel of A, ran(A) denote the column space of A, rank(A) denote the rank of A, and null(A) denote the nullity (dimension of the kernel) of A. A matrix $A \in M(n)$ is called normal if $A^T A = AA^T$. For every $B \in Psd(n)$, there exists a unique $B^{1/2} \in Psd(n)$ such that $(B^{1/2})^2 = B$. The matrices B and $B^{1/2}$ have the same range, and hence the same rank. See [32, Theorem 7.2.6]. Every symmetric matrix $C \in Sm(n)$ can be expressed as a difference of two positive semidefinite matrices $C = C_+ - C_-$, where

$$C_{-} \coloneqq \frac{1}{2}(|C| - C),$$
 (2.1)

$$C_{+} := \frac{1}{2}(|C| + C), \qquad (2.2)$$

and $|C| := (C^2)^{1/2}$ is the absolute value of *C*. We have rank $(C) = \operatorname{rank}(C_+) + \operatorname{rank}(C_-)$ and $C_+C_- = C_-C_+ = 0$. See Proposition 4.1.13 of [32].

2.2 Linear algebra on \mathbb{R}^n

We denote by $\langle \cdot, \cdot \rangle$ the Euclidean inner product given for all $x, y \in \mathbb{R}^n$ by $\langle x, y \rangle := x^T y$. Let \mathcal{W} be a linear subspace of \mathbb{R}^n . \mathcal{W} is said to be an invariant subspace of $A \in M(n)$ if for all $w \in \mathcal{W}$, $Aw \in \mathcal{W}$. We say that A is positive definite on \mathcal{W} if $\langle w, Aw \rangle > 0$ for all non-zero $w \in \mathcal{W}$. We say A is negative definite on \mathcal{W} if -A is positive-definite on \mathcal{W} . The orthogonal complement of \mathcal{W} is defined as

$$\mathscr{W}^{\perp} := \{ u \in \mathbb{R}^n : \langle u, w \rangle = 0, \ \forall w \in \mathscr{W} \}.$$
(2.3)

A matrix $P \in \text{Sm}(n)$ is called an orthogonal projection onto \mathcal{W} if Pw = w and Pw' = 0 for all $w \in \mathcal{W}$ and $w' \in \mathcal{W}^{\perp}$. Any matrix $Q \in \text{Sm}(n)$ that satisfies $Q^2 = Q$ is an orthogonal projection onto ran(Q).

2.3 Symplectic linear algebra on \mathbb{R}^{2n}

Let $J_{2n} := \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$, where I_n is the identity matrix of size $n \times n$. We shall drop the subscript 2n from J_{2n} , and use the notation J instead, when the size of the matrix is clear

from the context. The symplectic orthogonal complement of a subset $\mathcal{X} \subseteq \mathbb{R}^{2n}$ is defined as

$$\mathscr{X}^{\perp_{\mathrm{s}}} \coloneqq \{ u \in \mathbb{R}^{2n} : \forall v \in \mathscr{X}, \langle u, Jv \rangle = 0 \}.$$
(2.4)

A linear subspace \mathcal{W} of \mathbb{R}^{2n} is called a symplectic subspace if for every $u \in \mathcal{W}$ there exists $v \in \mathcal{W}$ such that $\langle u, Jv \rangle \neq 0$. By definition, \mathcal{W} is a symplectic subspace of \mathbb{R}^{2n} if and only if $\mathcal{W} \cap \mathcal{W}^{\perp_s} = \{0\}$. Let \mathcal{W} be a symplectic subspace of \mathbb{R}^{2n} . Then \mathcal{W} has even dimension, say 2k, and it has a *symplectic basis* $\{u_1, \ldots, u_k, v_1, \ldots, v_k\}$ that satisfies for all $1 \leq i, j \leq n$:

$$\langle u_i, Jv_j \rangle = \delta_{ij},\tag{2.5}$$

$$\langle u_i, Ju_j \rangle = 0, \tag{2.6}$$

$$\langle v_i, J v_j \rangle = 0. \tag{2.7}$$

Here $\delta_{ij} = 0$ if $i \neq j$ and $\delta_{ij} = 1$ if i = j. We have $\dim(\mathcal{W}) + \dim(\mathcal{W}^{\perp_s}) = 2n$ and $(\mathcal{W}^{\perp_s})^{\perp_s} = \mathcal{W}$. See [34, Section 1.2]. We say that two symplectic subspaces \mathcal{W} and \mathcal{Z} are said to be symplectically orthogonal to each other if $\mathcal{Z} \subseteq \mathcal{W}^{\perp_s}$.

Let $\operatorname{Sp}(2n, 2k)$ denote the set of $2n \times 2k$ real matrices M that satisfy $M^T J_{2n} M = J_{2k}$. We use the shorthand $\operatorname{Sp}(2n)$ for $\operatorname{Sp}(2n, 2n)$. The set $\operatorname{Sp}(2n)$ consists of $2n \times 2n$ real symplectic matrices, and it is known as the symplectic group. For every $M \in \operatorname{Sp}(2n, 2k)$, ran(M) is a symplectic subspace of \mathbb{R}^{2n} , and the columns of M form a symplectic basis of ran(M). See [34, Section 1.2.1]. We denote by $\operatorname{OrSp}(2n)$ the set of *orthosymplectic* matrices given by $\operatorname{Or}(2n) \cap \operatorname{Sp}(2n)$.

Let n_1, \ldots, n_k be positive integers, and $X_i \in M(n_i)$ for $1 \le i \le k$. Denote by $\oplus X_i$ the usual direct sum of the matrices X_1, \ldots, X_k . Suppose $A_i \in M(2n_i)$ is partitioned into blocks as

$$A_i = \begin{pmatrix} E_i & F_i \\ G_i & H_i \end{pmatrix},\tag{2.8}$$

where $E_i, F_i, G_i, H_i \in M(n_i)$ for all $1 \le i \le k$. The s-direct sum of A_1, \ldots, A_k is defined by

$$\oplus^{s} A_{i} := \begin{pmatrix} \oplus E_{i} & \oplus F_{i} \\ \oplus G_{i} & \oplus H_{i} \end{pmatrix}.$$
(2.9)

Let *M* and *N* be $2n \times 2k$ and $2n \times 2\ell$ matrices whose columns are $u_1, \ldots, u_k, v_1, \ldots, v_k$ and $x_1, \ldots, x_\ell, y_1, \ldots, y_\ell$, respectively. Define the symplectic concatenation of *M* and *N* to be the following $2n \times 2(k + \ell)$ matrix given by

$$M \diamond N \coloneqq [u_1, \dots, u_k, x_1, \dots, x_\ell, v_1, \dots, v_k, y_1, \dots, y_\ell].$$

$$(2.10)$$

3 Williamson's theorem for symmetric matrices

Generalizing Williamson's theorem to symmetric matrices is the main objective of this section. We begin by building some intuition towards generalization of the theorem. Let $A \in \text{Sm}(2n)$ for which there exists $M \in \text{Sp}(2n)$ such that

$$M^{T}AM = \begin{pmatrix} D & 0\\ 0 & D \end{pmatrix}, \tag{3.1}$$

where *D* is an $n \times n$ diagonal matrix. We shall refer to (3.1) as a Williamson's decomposition of *A*. Since the symplectic matrix *M* satisfies $M^{-T} = JMJ^T$, (3.1) gives

$$AM = JMJ^T \begin{pmatrix} D & 0\\ 0 & D \end{pmatrix}.$$
 (3.2)

Let $u_1, \ldots, u_n, v_1, \ldots, v_n$ denote the columns of M and d_1, \ldots, d_n denote the diagonal elements of D. Then (3.2) implies for all $1 \le i \le n$:

$$Au_i = d_i J v_i, \tag{3.3}$$

$$Av_i = -d_i J u_i. \tag{3.4}$$

Define index sets:

$$I_{-} := \{ i : 1 \le i \le n, d_{i} < 0 \}, \tag{3.5}$$

$$I_0 := \{i : 1 \le i \le n, d_i = 0\},\tag{3.6}$$

$$I_{+} := \{i : 1 \le i \le n, d_{i} > 0\},$$
(3.7)

and subspaces:

$$\mathscr{W}_{-} \coloneqq \operatorname{span}\{u_{i}, v_{i} : i \in \mathcal{I}_{-}\},$$
(3.8)

$$\mathcal{W}_0 \coloneqq \operatorname{span}\{u_i, v_i : i \in I_0\},\tag{3.9}$$

$$\mathscr{W}_{+} \coloneqq \operatorname{span}\{u_{i}, v_{i} : i \in \mathcal{I}_{+}\}.$$
(3.10)

By construction, \mathcal{W}_- , \mathcal{W}_0 , \mathcal{W}_+ are symplectic subspaces and are pairwise symplectically orthogonal to each other. Also, by the Sylvester's law of inertia, we have dim(\mathcal{W}_-) = $\nu(A)$, dim(\mathcal{W}_0) = $\xi(A)$, and dim(\mathcal{W}_+) = $\pi(A)$ so that the dimensions of these subspaces add to 2n. The relations (3.3) and (3.4) imply that these subspaces are invariant under *JA*. It is also easy to verify that *A* is negative definite on \mathcal{W}_- . Indeed, let $x \in \mathcal{W}_$ be any non-zero vector given by $x = \sum_{i \in I_-} (a_i u_i + b_i v_i)$, where $a_i, b_i \in \mathbb{R}$ for all $i \in I_-$. We have

$$\langle x, Ax \rangle = \left\langle \sum_{i \in \mathcal{I}_{-}} (a_i u_i + b_i v_i), \sum_{j \in \mathcal{I}_{-}} (a_j A u_j + b_j A v_j) \right\rangle$$
(3.11)

$$= \sum_{i,j\in \mathcal{I}_{-}} \langle a_i u_i + b_i v_i, a_j d_j J v_j - b_j d_j J u_j \rangle$$
(3.12)

$$= \sum_{i,j\in I_{-}} \left(a_i a_j d_j \langle u_i, J v_j \rangle - a_i b_j d_j \langle u_i, J u_j \rangle + b_i a_j d_j \langle v_i, J v_j \rangle - b_i b_j d_j \langle v_i, J u_j \rangle \right)$$
(3.13)

$$+ b_i a_j a_j \langle v_i, J v_j \rangle - b_i b_j a_j \langle v_i, J u_j \rangle$$
(3.13)
$$d_i (a^2 + b^2)$$
(3.14)

$$= \sum_{i \in I_{-}} d_i (a_i^2 + b_i^2)$$
(3.14)

The last inequality follows from the fact that $d_i < 0$ for all $i \in \mathcal{I}_-$. A similar argument shows that *A* is positive definite on \mathcal{W}_+ . Also, we obviously have ker(*A*) = \mathcal{W}_0 .

To summarise everything, the following are necessary conditions on any $A \in Sm(2n)$ that is diagonalizable in the sense of Williamson's theorem:

Condition (i) There exist pairwise symplectically orthogonal symplectic subspaces $\mathcal{W}_{-}, \mathcal{W}_{0}, \mathcal{W}_{+}$ with dimensions $v(A), \xi(A), \pi(A)$, respectively.

Condition (ii) Each of these symplectic subspaces is invariant under *JA*. **Condition** (iii) *A* is negative definite on \mathcal{W}_{-} , the kernel of *A* is \mathcal{W}_{0} , and *A* is positive definite on \mathcal{W}_{+} .

In the following theorem, we show that the above three conditions are sufficient for a symmetric matrix to be diagonalizable in the sense of Williamson's theorem.

Theorem 3.1 For $A \in Sm(2n)$ there exists $M \in Sp(2n)$ and an $n \times n$ diagonal matrix D such that

$$M^{T}AM = \begin{pmatrix} D & 0\\ 0 & D \end{pmatrix}$$
(3.16)

if and only A satisfies Condition (i), Condition (ii), and Condition (iii). The diagonal matrix D so obtained is unique up to ordering of its diagonal entries. Moreover, the diagonal entries of D and -D combined together constitute the 2n eigenvalues of ιJA .

Proof The necessity of the given conditions is already established in the beginning of the section. In what follows, we give an argument for sufficiency of these conditions.

Let $A \in \text{Sm}(2n)$ and suppose \mathcal{W}_{-} , \mathcal{W}_{0} , \mathcal{W}_{+} are symplectic subspaces of \mathbb{R}^{2n} that satisfy **Condition** (i), **Condition** (ii), and **Condition** (iii) for A. Let $k = \frac{1}{2}\nu(A)$, $\ell = \frac{1}{2}\xi(A)$, and $m = \frac{1}{2}\pi(A)$. Choose $M_{-} \in \text{Sp}(2n, 2k)$, $M_{0} \in \text{Sp}(2n, 2\ell)$, and $M_{+} \in \text{Sp}(2n, 2m)$ such that $\operatorname{ran}(M_{-}) = \mathcal{W}_{-}$, $\operatorname{ran}(M_{0}) = \mathcal{W}_{0}$, and $\operatorname{ran}(M_{+}) = \mathcal{W}_{+}$. By **Condition** (i) we have $M_{-} \diamond M_{0} \diamond M_{+} \in \text{Sp}(2n)$. **Condition** (iii) implies $-M_{-}^{T}AM_{-} \in \text{Pd}(2k)$ and $M_{+}^{T}AM_{+} \in \text{Pd}(2m)$. By Williamson's theorem, we thus get $Q_{-} \in \text{Sp}(2k)$ and $Q_{+} \in \text{Sp}(2m)$ such that

$$Q_{-}^{T}M_{-}^{T}AM_{-}Q_{-} = \begin{pmatrix} D_{-} & 0\\ 0 & D_{-} \end{pmatrix}, \qquad (3.17)$$

$$Q_{+}^{T}M_{+}^{T}AM_{+}Q_{+} = \begin{pmatrix} D_{+} & 0\\ 0 & D_{+} \end{pmatrix},$$
(3.18)

where $D_- < 0$ and $D_+ > 0$ are diagonal matrices of size $k \times k$ and $m \times m$, respectively. Set $M := (M_-Q_-) \diamond M_0 \diamond (M_+Q_+)$. It is easy to check that $M \in \text{Sp}(2n)$. In what follows, we show that M diagonalizes A in the sense of Williamson's theorem.

By **Condition** (ii), the columns of JAM_+ lie in the subspace \mathcal{W}_+ . Since \mathcal{W}_- and \mathcal{W}_+ are symplectically orthogonal to each other, we have $M_-^T JJAM_+ = 0$ implying $M_-^T AM_+ = 0$. Also, we have $AM_0 = 0$ which implies that $M_-^T AM_0 = 0$ and $M_+^T AM_0 = 0$. Therefore, we get

$$M^{T}AM = [(M_{-}Q_{-}) \diamond M_{0} \diamond (M_{+}Q_{+})]^{T} A [(M_{-}Q_{-}) \diamond M_{0} \diamond (M_{+}Q_{+})]$$
(3.19)

$$= [(M_{-}Q_{-}) \diamond M_{0} \diamond (M_{+}Q_{+})]^{T} [(AM_{-}Q_{-}) \diamond (AM_{0}) \diamond (AM_{+}Q_{+})] \quad (3.20)$$

$$= \left(Q_{-}^{T} M_{-}^{T} A M_{-} Q_{-} \right) \oplus_{s} \left(M_{0}^{T} A M_{0} \right) \oplus_{s} \left(Q_{+}^{T} M_{+}^{T} A M_{+} Q_{+} \right)$$
(3.21)

$$= \begin{pmatrix} D_{-} & 0\\ 0 & D_{-} \end{pmatrix} \oplus_{s} \begin{pmatrix} 0_{\ell} & 0_{\ell}\\ 0_{\ell} & 0_{\ell} \end{pmatrix} \oplus_{s} \begin{pmatrix} D_{+} & 0\\ 0 & D_{+} \end{pmatrix}$$
(3.22)

$$= \begin{pmatrix} D & 0\\ 0 & D \end{pmatrix}, \tag{3.23}$$

9

where $D \coloneqq D_{-} \oplus 0_{\ell} \oplus D_{+}$ and 0_{ℓ} denotes the zero matrix of size $\ell \times \ell$.

The uniqueness of the diagonal form D and the fact that the combined diagonal entries of D and -D form the eigenvalues of ιJA are established by Pereira et al. [3, Section 5].

Remark 3.2 Pereira et al. [3] provided a method of computing Williamson's decomposition of a $2n \times 2n$ (complex) symmetric matrix A, given that A is guaranteed to admit such a decomposition. Theorem 3.1 complements the work of Pereira et al. [3] in the sense that it provides a characterization of A for existence of its Williamson's decomposition.

Let SpSm(2n) denote the subset of Sm(2n) consisting of matrices satisfying **Condition** (i), **Condition** (ii), and **Condition** (iii). In view of Theorem 3.1, for every $A \in \text{SpSm}(2n)$, there exists $M \in \text{Sp}(2n)$ and a unique $n \times n$ diagonal matrix D with diagonal diagonal entries in ascending order such that $M^T A M = D \oplus D$. We refer to the diagonal elements of D as the symplectic eigenvalues of A. Thus, a matrix in SpSm(2n) can have negative, zero, or positive symplectic eigenvalues.

Let SpPsd(2n) denote the set of $2n \times 2n$ real symmetric positive semidefinite matrices with symplectic kernel. As a corollary of Theorem 3.1, we get the following known result which states that every matrix in SpPsd(2n) exhibits Williamson's decomposition. See [12, Remark 2.6] and [17, Section 2].

Corollary 3.3 *We have* $SpPsd(2n) \subset SpSm(2n)$ *.*

Proof Let $A \in \text{SpPsd}(2n)$. Choose $\mathscr{W}_{-} = 0$, $\mathscr{W}_{0} = \ker(A)$, and $\mathscr{W}_{+} = \mathscr{W}_{0}^{\perp_{s}}$. These symplectic subspaces clearly satisfy **Condition** (i) and **Condition** (iii). It is also straightforward to see that \mathscr{W}_{-} and \mathscr{W}_{0} are invariant under *JA*. It remains to show that \mathscr{W}_{+} is invariant under *JA*. We have $\mathbb{R}^{2n} = \mathscr{W}_{0} \oplus \mathscr{W}_{+}$. Let $y \in \mathscr{W}_{+}$ be arbitrary. For any $x \in \mathscr{W}_{0}$, we have

$$\langle x, J(JA)y \rangle = -\langle x, Ay \rangle$$
 (3.24)

$$= -\langle Ax, y \rangle \tag{3.25}$$

$$= -\langle 0, y \rangle \tag{3.26}$$

$$= 0.$$
 (3.27)

This by definition means $JAy \in \mathcal{W}_0^{\perp_s} = \mathcal{W}_+$, implying that \mathcal{W}_+ is invariant under JA. This shows that **Condition** (ii) is also satisfied by \mathcal{W}_- , \mathcal{W}_0 , \mathcal{W}_+ for A and hence $A \in \text{SpSm}(2n)$.

4 General Williamson's theorem via symplectic orthogonal projection

In this section we introduce a symplectic analog of orthogonal projection, call it symplectic orthogonal projection, and provide an alternate statement for the general Williamson's theorem in terms of symplectic orthogonal projection.

Let \mathcal{W} be a 2k dimensional symplectic subspace of \mathbb{R}^{2n} . Let $M \in \operatorname{Sp}(2n, 2k)$ be any matrix such that $\operatorname{ran}(M) = \mathcal{W}$. The matrix $P_M := JMM^T J^T$ is called the symplectic projection corresponding to M. It is a positive semidefinite matrix with kernel \mathcal{W}^{\perp_s} . See [12, Section 5]. It is known that for $N \in \operatorname{Sp}(2n, 2k)$, the equality $P_M = P_N$ holds if and only there exists $U \in \operatorname{OrSp}(2k)$ such that N = MU [12, Proposition 5.1]. Consequently, we have that $\operatorname{ran}(N) = \mathcal{W}$ is a necessary but not a sufficient condition for the symplectic projection P_N to be equal to P_M (for instance, choose N = MUfor $U \in \operatorname{Sp}(2k) \setminus \operatorname{OrSp}(2k)$). However, it is interesting to observe that the condition $\operatorname{ran}(N) = \mathcal{W}$ is necessary and sufficient for the equality $P_N J P_N = P_M J P_M$. Moreover, the matrix $J^T P_M J P_M$ restricted to \mathcal{W} is the identity operator and its kernel is \mathcal{W}^{\perp_s} as shown in the following proposition.

Proposition 4.1 Let \mathcal{W} be a symplectic subspace of \mathbb{R}^{2n} . Let $M \in \text{Sp}(2n, 2k)$ such that $\operatorname{ran}(M) = \mathcal{W}$, and let P_M be the symplectic projection corresponding to M. Then for all $x \in \mathcal{W}$, we have $J^T P_M J P_M x = x$. Also, $\ker(J^T P_M J P_M) = \mathcal{W}^{\perp_s}$.

Proof Let $M \in \text{Sp}(2n, 2k)$ such that $ran(M) = \mathcal{W}$, and let $u_1, \ldots, u_k, v_1, \ldots, v_k$ be the columns of M. Denote by e_1, \ldots, e_{2n} the standard unit vectors of \mathbb{R}^{2n} . For all $1 \le i \le r$ we have

$$P_M u_i = J M M^T J^T u_i \tag{4.1}$$

$$= JMe_{i+n} \tag{4.2}$$

$$=Jv_i.$$
(4.3)

Similarly, we get $P_M v_i = -Ju_i$. These observations give the following:

$${}^{T}P_{M}JP_{M}u_{i} = J^{T}P_{M}J^{2}v_{i} \tag{4.4}$$

$$= JP_M v_i \tag{4.5}$$

$$= -J^2 u_i \tag{4.6}$$

$$= u_i.$$
 (4.7)

A similar argument gives $J^T P_M J P_M v_i = v_i$. Consequently, for all $x \in \mathcal{W}$, we have $J^T P_M J P_M x = x$.

We have $\ker(J^T P_M J P_M) \supseteq \ker(P_M) = \mathcal{W}^{\perp_s}$, and $\operatorname{ran}(J^T P_M J P_M) \supseteq \mathcal{W}$. The rank-nullity theorem, combined with the fact that $\dim(\mathcal{W}) + \dim(\mathcal{W}^{\perp_s}) = 2n$, implies that $\ker(J^T P_M J P_M) = \mathcal{W}^{\perp_s}$.

Proposition 4.1 states that associated with every symplectic subspace is a *unique* matrix that acts as the identity on the symplectic subspace and its kernel is the symplectic complement of the given symplectic subspace. This leads to the following definition of *symplectic orthogonal projection* onto a symplectic subspace.

Definition 4.1 Let \mathcal{W} be a symplectic subspace of \mathbb{R}^{2n} . We call the $2n \times 2n$ real matrix Π , given by

$$\Pi x = \begin{cases} x & \text{if } x \in \mathcal{W}, \\ 0 & \text{if } x \in \mathcal{W}^{\perp_{s}}, \end{cases}$$
(4.8)

the symplectic orthogonal projection onto $\mathcal W$. It is given by

$$\Pi = J^T P_M J P_M, \tag{4.9}$$

for any $M \in \text{Sp}(2n, 2k)$ such that $ran(M) = \mathcal{W}$.

Remark 4.2 Symplectic orthogonal projections are precisely $2n \times 2n$ real matrices Π that satisfy

- ker(Π) is a symplectic subspace,
- $ran(\Pi) = ker(\Pi)^{\perp_s}$,
- $\Pi^2 = \Pi$.

Proposition 4.3 Let \mathcal{W} be a symplectic subspace of \mathbb{R}^{2n} , and let Π be the associated symplectic orthogonal projection. Then Π^T is the symplectic orthogonal projection onto $J\mathcal{W}$.

Proof Let $M \in \text{Sp}(2n, 2k)$ such that $ran(M) = \mathcal{W}$. We have $P_{JM} = MM^T$. This gives

$$\Pi^{T} = \left(J^{T} P_{M} J P_{M}\right)^{T} \tag{4.10}$$

$$= P_M J^T P_M J \tag{4.11}$$

$$= JMM^{T}J^{T}J^{T}JMM^{T}J^{T}J$$
(4.12)

$$=J^T P_{JM} J P_{JM}. (4.13)$$

We now state Theorem 3.1 in terms of symplectic orthogonal projections as follows.

Proposition 4.4 Let $A \in \text{Sm}(2n)$. We have $A \in \text{SpSm}(2n)$ if and only if there exist symplectic orthogonal projections $\Pi_{-}, \Pi_{0}, \Pi_{+}$ satisfying the following conditions:

(*i*) $\Pi_{-}\Pi_{0} = \Pi_{-}\Pi_{+} = \Pi_{0}\Pi_{+} = 0$, and $\Pi_{-} + \Pi_{0} + \Pi_{+} = I_{2n}$.

- (*ii*) $A = \prod_{-}^{T} A \prod_{-} + \prod_{+}^{T} A \prod_{+}$
- (iii) $\Pi_{-}^{T}A\Pi_{-}$ is negative definite on ran (Π_{-}) and $\Pi_{+}^{T}A\Pi_{+}$ is positive definite on ran (Π_{+}) .

Proof The "if" part is straightforward. Suppose there exist symplectic orthogonal projections Π_-, Π_0, Π_+ satisfying the given conditions. Choose $\mathscr{W}_- = \operatorname{ran}(\Pi_-), \mathscr{W}_0 = \operatorname{ran}(\Pi_0)$, and $\mathscr{W}_+ = \operatorname{ran}(\Pi_+)$. It is easy to see that the symplectic subspaces $\mathscr{W}_-, \mathscr{W}_0$, \mathscr{W}_+ satisfy **Condition** (i), **Condition** (ii), and **Condition** (iii). Therefore, we have $A \in \operatorname{SpSm}(2n)$.

We now prove the "only if" part. Suppose $A \in \text{SpSm}(2n)$. Then there exist symplectic subspaces \mathcal{W}_{-} , \mathcal{W}_{0} , \mathcal{W}_{+} satisfying **Condition** (i), **Condition** (ii), and **Condition** (iii) for A. Let Π_{-} , Π_{0} , and Π_{+} be the symplectic orthogonal projections onto \mathcal{W}_{-} , \mathcal{W}_{0} , and \mathcal{W}_{+} , respectively. **Condition** (i) implies that $\Pi_{-}\Pi_{0} = \Pi_{-}\Pi_{+} = \Pi_{0}\Pi_{+} = 0$ and $\Pi_{-} + \Pi_{0} + \Pi_{+} = I_{2n}$. For any $x \in \mathcal{W}_{-}$, $y \in \mathcal{W}_{0}$, and $z \in \mathcal{W}_{+}$, we get

$$\left(\Pi_{-}^{T}A\Pi_{-} + \Pi_{+}^{T}A\Pi_{+}\right)(x+y+z) = \Pi_{-}^{T}A\Pi_{-}x + \Pi_{+}^{T}A\Pi_{+}z$$
(4.14)

$$=\Pi_{-}^{T}Ax + \Pi_{+}^{T}Az \tag{4.15}$$

$$= Ax + Az \tag{4.16}$$

$$= A(x + y + z).$$
 (4.17)

The equality (4.16) follows from Proposition 4.3 and the fact that \mathcal{W}_{-} and \mathcal{W}_{+} are invariant under *JA*, which is given by **Condition** (ii). We thus have $A = \Pi_{+}^{T}A\Pi_{+} + \Pi_{-}^{T}A\Pi_{-}$. Lastly, $\Pi_{+}^{T}A\Pi_{+}$ being positive definite on \mathcal{W}_{+} and $\Pi_{-}^{T}A\Pi_{-}$ being negative definite on \mathcal{W}_{-} follows directly from **Condition** (iii).

We know that the symplectic eigenvalues of $A \in Pd(2n)$ are the positive eigenvalues of the Hermitian matrix $\iota A^{1/2}JA^{1/2}$. We state an analogous fact for matrices in SpSm(2n) as follows.

Proposition 4.5 Let $A \in \text{SpSm}(2n)$, and let Π_- , Π_0 , and Π_+ be symplectic orthogonal projections given by Proposition 4.4. Then A has $\frac{1}{2} \operatorname{rank}(\Pi_0)$ zero symplectic eigenvalues. The negative symplectic eigenvalues of A are the negative eigenvalues of $\iota \left(-\Pi_-^T A \Pi_-\right)^{1/2} J \left(-\Pi_-^T A \Pi_-\right)^{1/2}$, and the positive symplectic eigenvalues are the positive eigenvalues of $\iota \left(\Pi_+^T A \Pi_+\right)^{1/2} J \left(\Pi_+^T A \Pi_+\right)^{1/2}$.

Proof We know from Proposition 4.4 that $-\Pi_{-}^{T}A\Pi_{-}$ and $\Pi_{+}^{T}A\Pi_{+}$ are positive semidefinite matrices. Also, ker $(-\Pi_{-}^{T}A\Pi_{-})$ = ker (Π_{-}) and ker $(\Pi_{+}^{T}A\Pi_{+})$ = ker (Π_{+}) , which follow from the facts that $\Pi_{-}^{T}A\Pi_{-}$ is negative definite on ran (Π_{-}) and $\Pi_{+}^{T}A\Pi_{+}$ is positive definite on ran (Π_{+}) . Therefore, Williamson's decompositions of $\Pi_{-}^{T}A\Pi_{-}$ and $\Pi_{+}^{T}A\Pi_{+}$ exist. We know from [17, Section 2] that the negative symplectic eigenvalues of $\Pi_{-}^{T}A\Pi_{-}$ are the negative eigenvalues of $\iota (-\Pi_{-}^{T}A\Pi_{-})^{1/2} J (-\Pi_{-}^{T}A\Pi_{-})^{1/2}$, and the positive symplectic eigenvalues of $\Pi_{+}^{T}A\Pi_{+}$ are the positive eigenvalues of

 $\iota (\Pi_{+}^{T}A\Pi_{+})^{1/2} J (\Pi_{+}^{T}A\Pi_{+})^{1/2}$. Therefore, it suffices to show that the non-zero symplectic eigenvalues of $\Pi_{-}^{T}A\Pi_{-}$ and $\Pi_{+}^{T}A\Pi_{+}$ put together.

Suppose the dimensions of $ran(\Pi_{-})$, $ran(\Pi_{0})$, $ran(\Pi_{+})$ are 2k, 2ℓ , 2m, respectively. Thus, $-\Pi_{-}^{T}A\Pi_{-}$ and $\Pi_{+}^{T}A\Pi_{+}$ have ranks 2k and 2m, respectively. Let μ_{1}, \ldots, μ_{k} and $\eta_{1}, \ldots, \eta_{m}$ denote the non-zero symplectic eigenvalues of $\Pi_{-}^{T}A\Pi_{-}$ and $\Pi_{+}^{T}A\Pi_{+}$, respectively. By Theorem 3.1, there exist $M, N \in Sp(2n)$ such that

$$M^{T}\Pi_{-}^{T}A\Pi_{-}M = D_{-} \oplus D_{-}, \qquad (4.18)$$

$$N^{T}\Pi_{+}^{T}A\Pi_{+}N = D_{+} \oplus D_{+}.$$
(4.19)

where D_- and D_+ are the $n \times n$ diagonal matrices given by $D_- = \text{diag}(\mu_1, \ldots, \mu_k, 0, \ldots, 0)$ and $D_+ = \text{diag}(\eta_1, \ldots, \eta_m, 0, \ldots, 0)$. Let $w_1, \ldots, w_n, z_1, \ldots, z_n$ be the columns of M. We have

$$\operatorname{span}\{w_1, \dots, w_k, z_1, \dots, z_k\} = \operatorname{span}\{w_{k+1}, \dots, w_n, z_{k+1}, \dots, z_n\}^{\perp_s}$$
(4.20)

$$= \ker \left(\Pi_{-}^{T} A \Pi_{-} \right)^{\perp_{s}} \tag{4.21}$$

$$= \ker \left(\Pi_{-} \right)^{\perp_{s}} \tag{4.22}$$

$$= \operatorname{ran}\left(\Pi_{-}\right). \tag{4.23}$$

We know from Proposition 4.4 that $\Pi_+\Pi_- = 0$, thus implying for $1 \le i \le k$ that $w_i, z_i \in \ker(\Pi_+)$. Using the fact that $A = \Pi_-^T A \Pi_- + \Pi_+^T A \Pi_+$, we then get $\Pi_-^T A \Pi_- w_i = A w_i$ and $\Pi_-^T A \Pi_- z_i = A z_i$ for all $1 \le i \le k$. The equation (4.18) thus implies for $1 \le i \le k$:

$$Aw_i = \mu_i J z_i \tag{4.24}$$

$$Az_i = -\mu_i J w_i. \tag{4.25}$$

Let $u_1, \ldots, u_n, v_1, \ldots, v_n$ be the columns of *N*. By a similar arguments as given earlier, we get for $j = 1, \ldots, m$:

$$Au_{j} = \eta_{j} Jv_{j} \tag{4.26}$$

$$Av_i = -\eta_i J u_i. \tag{4.27}$$

Let $\{x_1, \ldots, x_\ell, y_1, \ldots, y_\ell\}$ be a symplectic basis of ran (Π_0) . Let us choose $S \coloneqq [w_1, \ldots, w_k, z_1, \ldots, z_k] \diamond [x_1, \ldots, x_\ell, y_1, \ldots, y_\ell] \diamond [u_1, \ldots, u_m, v_1, \ldots, v_m].$ (4.28)

It is easy to verify that $S \in \text{Sp}(2n)$ and $S^T A S = D \oplus D$, where *D* is the $n \times n$ diagonal matrix given by $D = \text{diag}(\mu_1, \dots, \mu_k, 0, \dots, 0, \eta_1, \dots, \eta_m)$. This completes the proof.

5 Explicit Williamson's decomposition for a subset of SpSm(2*n*)

For $A \in \text{Sm}(2n)$, let \mathscr{E}_{-} , \mathscr{E}_{0} , \mathscr{E}_{+} denote the eigen subspaces of A spanned by the eigenvectors corresponding to its negative, zero, and positive eigenvalues, respectively. We define EigSpSm(2n) to be the set of those matrices $A \in \text{Sm}(2n)$ for which \mathscr{E}_{-} , \mathscr{E}_{0} , \mathscr{E}_{+} are pairwise symplectically orthogonal symplectic subspaces, and each of these subspaces is invariant under JA. Observe that $\text{Pd}(2n) \subset \text{SpPsd}(2n) \subset \text{EigSpSm}(2n) \subset \text{SpSm}(2n)$.

In this section, we provide a explicit description of symplectic eigenvalues and diagonalizing symplectic matrices in Williamson's decomposition for matrices in EigSpSm(2n). Furthermore, we establish perturbation bounds on the symplectic eigenvalues of matrices in EigSpSm(2n).

We begin with some preliminary results that will be helpful in the subsequent parts of the section.

Lemma 5.1 Let $A \in \text{Sm}(2n)$ and \mathcal{C} be a symplectic subspace of \mathbb{R}^{2n} of dimension 2k. Suppose \mathcal{C} is an invariant subspace of both JA and A, and that A is positive definite on \mathcal{C} . Then there exist k positive numbers $\gamma_1, \ldots, \gamma_k$ and a symplectic basis $\{u_1, \ldots, u_k, v_1, \ldots, v_k\}$ of \mathcal{C} such that for all $1 \le i \le k$,

$$Au_i = \gamma_i J v_i, \tag{5.1}$$

$$Av_i = -\gamma_i J u_i. \tag{5.2}$$

Proof Let *P* be the orthogonal projection onto the subspace \mathscr{C} . Set $\hat{A} := A + I - P$, where *I* is the identity matrix. Let $x \in \mathscr{C}$ and $x^{\perp} \in \mathscr{C}^{\perp}$ be arbitrary. We have

 $\hat{A}(x + x^{\perp}) = Ax + x - Px + Ax^{\perp} + x^{\perp} - Px^{\perp}$ (5.3)

$$= Ax + x - x + x^{\perp} \tag{5.4}$$

$$= A + x^{\perp}.$$
 (5.5)

If
$$x + x^{\perp} \neq 0$$
, i.e., $x \neq 0$ or $x^{\perp} \neq 0$. This then implies

$$\langle x + x^{\perp}, \hat{A}(x + x^{\perp}) \rangle = \langle x + x^{\perp}, Ax + x^{\perp} \rangle$$
(5.6)

$$= \langle x, Ax \rangle + \langle x, x^{\perp} \rangle + \langle x^{\perp}, Ax \rangle + \langle x^{\perp}, x^{\perp} \rangle$$
(5.7)

$$= \langle x, Ax \rangle + \langle x^{\perp}, x^{\perp} \rangle > 0.$$
(5.8)

This implies that $\hat{A} \in Pd(2n)$, and it is easy to see that \mathscr{C} is invariant under $J\hat{A}$. By Proposition 4.1 of [35], there exists a symplectic basis $\{u_1, \ldots, u_k, v_1, \ldots, v_k\}$ of \mathscr{C} such that

$$\hat{A}u_i = \gamma_i J v_i, \tag{5.9}$$

$$\hat{A}v_i = -\gamma_i J u_i. \tag{5.10}$$

By definition, we have $\hat{A}u_i = Au_i$ and $\hat{A}v_i = Av_i$ for all $1 \le i \le k$. This completes the proof.

The following well-known result on commuting normal matrices plays key role in constructing symplectic matrices in Williamson's decomposition for matrices in EigSpSm(2n). See Theorem 2.5.15 of [32] for a proof.

Lemma 5.2 Let $A, B \in M(n)$ be normal matrices. If A and B commute, then there exists $P \in Or(n)$ and a non-negative integer r such that P^TAP and P^TBP are

block-diagonal matrices of the form:

$$P^{T}AP = \Delta_{1} \oplus \begin{pmatrix} \alpha_{1} & \beta_{1} \\ -\beta_{1} & \alpha_{1} \end{pmatrix} \oplus \cdots \oplus \begin{pmatrix} \alpha_{r} & \beta_{r} \\ -\beta_{r} & \alpha_{r} \end{pmatrix},$$
(5.11)

$$P^{T}BP = \Delta_{2} \oplus \begin{pmatrix} \gamma_{1} & \delta_{1} \\ -\delta_{1} & \gamma_{1} \end{pmatrix} \oplus \cdots \oplus \begin{pmatrix} \gamma_{r} & \delta_{r} \\ -\delta_{r} & \gamma_{r} \end{pmatrix},$$
(5.12)

where $\Delta_1, \Delta_2 \in M(n-2r)$ are diagonal matrices; $\alpha_i, \beta_i, \gamma_i, \delta_i$ are real numbers for all $1 \le i \le r$; and for each $i \in \{1, ..., r\}, \beta_i > 0$ or $\delta_i > 0$.

5.1 Description of symplectic eigenvalues for EigSpSm(2*n*)

The symplectic eigenvalues of a matrix $A \in \text{EigSpSm}(2n)$ are given by a combination of negative and non-negative eigenvalues of the Hermitian matrices $\iota A_{-}^{1/2}JA_{-}^{1/2}$ and $\iota A_{+}^{1/2}JA_{+}^{1/2}$ as stated below.

Theorem 5.3 The symplectic eigenvalues of $A \in \text{EigSpSm}(2n)$ are given by $\frac{1}{2}\xi(A)$ zeros, the negative eigenvalues of $\iota A_{-}^{1/2}JA_{-}^{1/2}$, and the positive eigenvalues of $\iota A_{+}^{1/2}JA_{+}^{1/2}$.

Proof Let $\mathscr{E}_-, \mathscr{E}_0, \mathscr{E}_+$ denote the eigen subspaces of A spanned by the eigenvectors corresponding to its negative, zero, and positive eigenvalues, respectively. Let Π_-, Π_0, Π_+ denote the orthogonal projections onto $\mathscr{E}_-, \mathscr{E}_0, \mathscr{E}_+$, respectively. By definition, Π_-, Π_0, Π_+ are also symplectic orthogonal projections onto the symplectic subspaces $\mathscr{E}_-, \mathscr{E}_0, \mathscr{E}_+$, respectively. Also, we have

$$\Pi_{-}^{T}A\Pi_{-} = -A_{-}, \tag{5.13}$$

$$\Pi_{+}^{T}A\Pi_{+} = A_{+}.$$
(5.14)

By Proposition 4.5, the negative eigenvalues of $\iota A_{-}^{1/2}JA_{-}^{1/2}$ and the positive eigenvalues of $\iota A_{+}^{1/2}JA_{+}^{1/2}$, together with $\frac{1}{2}\xi(A)$ zeros are the symplectic eigenvalues of A.

5.2 Description of symplectic matrices in Williamson's decomposition for EigSpSm(2*n*)

Let $A \in \text{EigSpSm}(2n)$. In what follows, we explicitly construct a symplectic matrix that diagonalizes A in the sense of Williamson's theorem.

We know that the matrices $A_{-}^{1/2}$ and $A_{+}^{1/2}$ commute with each other and satisfy $A_{-}^{1/2}A_{+}^{1/2} = 0$. Therefore, the skew-symmetric matrices $A_{-}^{1/2}JA_{-}^{1/2}$ and $A_{+}^{1/2}JA_{+}^{1/2}$ commute with each other and their product is equal to zero. By Lemma 5.2, there exists $U \in Or(2n)$ and a non-negative integer r such that

$$U^{T}A_{-}^{1/2}JA_{-}^{1/2}U = \Delta_{1} \oplus \begin{pmatrix} \alpha_{1} & \beta_{1} \\ -\beta_{1} & \alpha_{1} \end{pmatrix} \oplus \cdots \oplus \begin{pmatrix} \alpha_{r} & \beta_{r} \\ -\beta_{r} & \alpha_{r} \end{pmatrix},$$
(5.15)

H. K. Mishra

(5.27)

$$U^{T}A_{+}^{1/2}JA_{+}^{1/2}U = \Delta_{2} \oplus \begin{pmatrix} \gamma_{1} & \delta_{1} \\ -\delta_{1} & \gamma_{1} \end{pmatrix} \oplus \cdots \oplus \begin{pmatrix} \gamma_{r} & \delta_{r} \\ -\delta_{r} & \gamma_{r} \end{pmatrix},$$
(5.16)

where Δ_1 , Δ_2 are real diagonal matrices of size $(2n - 2r) \times (2n - 2r)$; the parameters α_i , β_i , γ_i , δ_i are real numbers such that $\beta_i > 0$ or $\delta_i > 0$ for all $1 \le i \le r$. Since both $U^T A_-^{1/2} J A_-^{1/2} U$ and $U^T A_+^{1/2} J A_+^{1/2} U$ are real skew-symmetric matrices, their diagonal elements are zero whence $\Delta_1 = \Delta_2 = 0$ and $\alpha_i = \gamma_i = 0$ for all $1 \le i \le r$. The fact that the product of the matrices in the left-hand sides of (5.15) and (5.16) is zero implies that $\beta_i \delta_i = 0$. This implies that for all $1 \le i \le r$, exactly one of β_i and δ_i is positive.

We know that the kernel of A_- is $\mathscr{C}_0 + \mathscr{C}_+$, which is a symplectic subspace of \mathbb{R}^{2n} of dimension $2(\ell+m)$. It is shown in [17, Section 2] that ker $(A_-^{1/2}JA_-^{1/2}) = \text{ker}(A_-)$, which implies rank $(A_-^{1/2}JA_-^{1/2}) = 2k$. Similarly, we get rank $(A_+^{1/2}JA_+^{1/2}) = 2m$. Therefore, we must have r = k + m, there exist distinct indices $1 \le i_1 < \cdots < i_k \le k + m$ and $1 \le j_1 < \cdots < j_m \le k + m$ such that for $i \in \{i_1, \ldots, i_k\}$, we have $\beta_i > 0, \delta_i = 0$ and for $j \in \{j_1, \ldots, j_m\}$, we have $\beta_j = 0, \delta_j > 0$. Let D_- and D_+ be $2n \times 2n$ diagonal matrices whose *i*th diagonal entries are given by

$$(D_{-})_{i} = \begin{cases} 0 & \text{if } i \in \{1, \dots, \ell\} \cup \{\ell + j_{1}, \dots, \ell + j_{m}\},\\ \beta_{i-\ell} & \text{if } i \in \{\ell + i_{1}, \dots, \ell + i_{k}\}, \end{cases}$$
(5.17)

$$(D_{+})_{i} = \begin{cases} 0 & \text{if } i \in \{1, \dots, \ell\} \cup \{\ell + i_{1}, \dots, \ell + i_{k}\},\\ \delta_{i-\ell} & \text{if } i \in \{\ell + j_{1}, \dots, \ell + j_{m}\}. \end{cases}$$
(5.18)

Let e_1, \ldots, e_{2n} denote the standard unit vectors in \mathbb{R}^{2n} . Let *P* denote the permutation matrix $[e_1, e_3, \ldots, e_{2n-1}, e_2, e_4, \ldots, e_{2n}]$. We then get

$$P^{T}U^{T}A_{-}^{1/2}JA_{-}^{1/2}UP = (D_{-} \oplus D_{-})J, \qquad (5.19)$$

$$P^{T}U^{T}A_{+}^{1/2}JA_{+}^{1/2}UP = (D_{+} \oplus D_{+})J.$$
(5.20)

Let Π_{-} and Π_{+} denote the following isometries

$$\Pi_{-} \coloneqq [e_{\ell+i_1}, \dots, e_{\ell+i_k}, e_{n+\ell+i_1}, \dots, e_{n+\ell+i_k}],$$
(5.21)

$$\Pi_{+} := [e_{\ell+j_{1}}, \dots, e_{\ell+j_{m}}, e_{n+\ell+j_{1}}, \dots, e_{n+\ell+j_{m}}].$$
(5.22)

From (5.19) and (5.20) we thus get

$$\Pi_{-}^{T} P^{T} U^{T} A_{-}^{1/2} J_{2n} A_{-}^{1/2} U P \Pi_{-} = (\widetilde{D}_{-} \oplus \widetilde{D}_{-}) J_{2k},$$
(5.23)

$$\Pi_{+}^{T} P^{T} U^{T} A_{+}^{1/2} J_{2n} A_{+}^{1/2} U P \Pi_{+} = (\widetilde{D}_{+} \oplus \widetilde{D}_{+}) J_{2m},$$
(5.24)

where $\widetilde{D}_{-} := \operatorname{diag}(\beta_{i_1}, \ldots, \beta_{i_k})$ and $\widetilde{D}_{+} := \operatorname{diag}(\delta_{j_1}, \ldots, \delta_{j_m})$. Choose

$$\widetilde{M}_{-} := J_{2n} A_{-}^{1/2} U P \Pi_{-} \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) J_{2k}^{T},$$
(5.25)

$$\widetilde{M}_{+} := J_{2n} A_{+}^{1/2} U P \Pi_{+} \left(\widetilde{D}_{+}^{-1/2} \oplus \widetilde{D}_{+}^{-1/2} \right) J_{2m}^{T}.$$
(5.26)

It is easy to see from (5.23) and (5.24) that $\widetilde{M}_{-} \in \text{Sp}(2n, 2k)$ and $\widetilde{M}_{+} \in \text{Sp}(2n, 2m)$. We observe that $A_{-}^{1/2}JA = -A_{-}^{1/2}JA_{-}$, which follows from the fact that \mathscr{C}_{-} and \mathscr{C}_{+} are invariant under JA. Therefore, we get

$$M_{-}^{I}AM_{-}$$

$$= -J_{2k} \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) \Pi_{-}^{T} P^{T} U^{T} A_{-}^{1/2} J_{2n} A J_{2n} A_{-}^{1/2} U P \Pi_{-} \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) J_{2k}^{T}$$
(5.28)

$$= J_{2k} \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) \Pi_{-}^{T} P^{T} U^{T} A_{-}^{1/2} J_{2n} A_{-} J_{2n} A_{-}^{1/2} U P \Pi_{-} \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) J_{2k}^{T}$$
(5.29)

$$= J_{2k} \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) \Pi_{-}^{T} \left(P^{T} U^{T} A_{-}^{1/2} J_{2n} A_{-}^{1/2} U P \right)^{2} \Pi_{-} \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) J_{2k}^{T}$$
(5.30)

$$= -J_{2k} \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) \Pi_{-}^{T} (D_{-}^{2} \oplus D_{-}^{2}) \Pi_{-} \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) J_{2k}^{T}$$
(5.31)

$$= -J_{2k} \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) (\widetilde{D}_{-}^{2} \oplus \widetilde{D}_{-}^{2}) \left(\widetilde{D}_{-}^{-1/2} \oplus \widetilde{D}_{-}^{-1/2} \right) J_{2k}^{T}$$
(5.32)

$$= -J_{2k}(D_{-} \oplus D_{-})J_{2k}^{I}$$

$$\tag{5.33}$$

$$= -(\widetilde{D}_{-} \oplus \widetilde{D}_{-}). \tag{5.34}$$

By similar arguments, one can show that

$$\widetilde{M}_{+}^{T}A\widetilde{M}_{+} = \widetilde{D}_{+} \oplus \widetilde{D}_{+}.$$
(5.35)

Choose any $\widetilde{M}_0 \in \text{Sp}(2n, 2\ell)$ whose columns form a symplectic basis of \mathcal{E}_0 . Define

$$\widetilde{M} := \widetilde{M}_0 \diamond \widetilde{M}_- \diamond \widetilde{M}_+. \tag{5.36}$$

The matrix \widetilde{M} is symplectic. Indeed, we have $A_+^{1/2}JA_-^{1/2} = 0$ since \mathscr{E}_- and \mathscr{E}_+ are invariant under JA. We thus get from (5.25) and (5.26) that

$$M_{+}^{I}J_{2n}M_{-} = 0_{2m,2k}.$$
(5.37)

Since the subspaces \mathscr{C}_0 and \mathscr{C}_- are perpendicular to each other, and $\operatorname{ran}(A^{1/2}_-) = \operatorname{ran}(A_-) = \mathscr{C}_-$, we get

$$\widetilde{M}_0^T J_{2n} \widetilde{M}_- = 0_{2\ell,2k}.$$
(5.38)

By similar arguments, we also get

$$\tilde{M}_0^T J_{2n} \tilde{M}_+ = 0_{2\ell, 2m}. (5.39)$$

The conditions (5.37), (5.38), (5.39) thus imply that $\widetilde{M} \in \text{Sp}(2n)$. See [14, Subsection 2.3]. By (5.34), (5.35), and the fact that $AM_0 = 0_{2n,2\ell}$, we get

$$M^T A M = D \oplus D, \tag{5.40}$$

where $D := (-\widetilde{D}_{-}) \oplus 0_{\ell,\ell} \oplus \widetilde{D}_{+}$.

5.3 Perturbation bounds on symplectic eigenvalues for EigSpSm(2n)

In this subsection, we provide perturbation bounds on symplectic eigenvalues of matrices in EigSpSm(2n) given by Theorem 5.3. These perturbation bounds generalize the known perturbation bounds on symplectic eigenvalues of positive definite matrices given in [6].

Let $M(n, \mathbb{C})$ denote the set of $n \times n$ complex matrices, and $U(n, \mathbb{C})$ denote the set of $n \times n$ complex unitary matrices. A norm $||| \cdot |||$ on $M(n, \mathbb{C})$ is called unitarily invariant if |||UXV||| = |||X||| for all $X \in M(n, \mathbb{C})$ and $U, V \in U(n, \mathbb{C})$. For $X, Y, Z \in M(n, \mathbb{C})$, every unitarily invariant norm satisfies $|||XYZ||| \le ||X|| \cdot ||Y||| \cdot ||Z||$. Here $|| \cdot ||$ denotes the

matrix operator norm. See Proposition IV.2.4 of [31]. For $A, B \in Psd(n)$, the following inequality holds [31, Theorem X.1.3]:

$$|||A^{1/2} - B^{1/2}||| \le ||||A - B|^{1/2}|||.$$
(5.41)

Given an $n \times n$ complex Hermitian matrix X, let $\lambda(X)$ denote the n-vector consisting of the eigenvalues of X arranged in the *decreasing* order. Let Eig(X) denote the $n \times n$ diagonal matrix whose diagonal elements are given by the entries of $\lambda(X)$. The Lidskii– Wielandt theorem [31, IV.62] gives

$$\|\operatorname{Eig}(X) - \operatorname{Eig}(Y)\| \le \|X - Y\|.$$
(5.42)

For $A \in \text{Sm}(2n)$, let $\widehat{D}(A)$ be the $2n \times 2n$ diagonal matrix

$$\widehat{D}(A) := \operatorname{Eig}(|A_{+}^{1/2}J_{2n}A_{+}^{1/2}|) + \operatorname{Eig}(-|A_{-}^{1/2}J_{2n}A_{-}^{1/2}|).$$
(5.43)

Since the eigenvalues of $\iota A_+^{1/2} J_{2n} A_+^{1/2}$ and $\iota A_-^{1/2} J_{2n} A_-^{1/2}$ occur in pairs of negativepositive, the diagonal elements of $\widehat{D}(A)$ occur in pairs of equal entries, and we denote the diagonal elements of $\widehat{D}(A)$ by $d_1(A), d_1(A), \ldots, d_n(A), d_n(A)$.

The next lemma gives a perturbation bound on D(A). We know from Theorem 5.3 that if $A \in \text{EigSpSm}(2n)$, then the diagonal elements of $\widehat{D}(A)$ are the symplectic eigenvalues of A given by Theorem 5.3, each counted twice.

Proof By definition (5.43) and triangle inequality, we get

$$\|\widehat{D}(A) - \widehat{D}(B)\| \leq \|\operatorname{Eig}(|A_{+}^{1/2}J_{2n}A_{+}^{1/2}|) - \operatorname{Eig}(|B_{+}^{1/2}J_{2n}B_{+}^{1/2}|)\| + \|\operatorname{Eig}(-|A_{-}^{1/2}J_{2n}A_{-}^{1/2}|) - \operatorname{Eig}(-|B_{-}^{1/2}J_{2n}B_{-}^{1/2}|)\|.$$
(5.47)

We know that the eigenvalues of $\iota A_+^{1/2} J_{2n} A_+^{1/2}$ and $\iota B_+^{1/2} J_{2n} B_+^{1/2}$ occur in positive negative pairs. Therefore, using the unitary invariance of the norm, we get

$$\|\operatorname{Eig}(|A_{+}^{1/2}J_{2n}A_{+}^{1/2}|) - \operatorname{Eig}(|B_{+}^{1/2}J_{2n}B_{+}^{1/2}|)\|$$
$$= \|\operatorname{Eig}(\iota A_{+}^{1/2}J_{2n}A_{+}^{1/2}) - \operatorname{Eig}(\iota B_{+}^{1/2}J_{2n}B_{+}^{1/2})\|.$$
(5.48)

Similarly, we also have

$$\|\operatorname{Eig}(-|A_{-}^{1/2}J_{2n}A_{-}^{1/2}|) - \operatorname{Eig}(-|B_{-}^{1/2}J_{2n}B_{-}^{1/2}|)\|$$
$$= \|\operatorname{Eig}(\iota A_{-}^{1/2}J_{2n}A_{-}^{1/2}) - \operatorname{Eig}(\iota B_{-}^{1/2}J_{2n}B_{-}^{1/2})\|.$$
(5.49)

Substituting (5.48) and (5.49) into the right-hand side of (5.47), we get

$$\|\widehat{D}(A) - \widehat{D}(B)\| \leq \|\operatorname{Eig}(\iota A_{+}^{1/2} J_{2n} A_{+}^{1/2}) - \operatorname{Eig}(\iota B_{+}^{1/2} J_{2n} B_{+}^{1/2})\| \\ + \|\operatorname{Eig}(\iota A_{-}^{1/2} J_{2n} A_{-}^{1/2}) - \operatorname{Eig}(\iota B_{-}^{1/2} J_{2n} B_{-}^{1/2})\|.$$
(5.50)

We now apply the same arguments as given in the proof of Theorem 7 of [6] to bound each term in the right-hand side of (5.50).

By the Lidskii–Wielandt theorem (5.42) and the relation (5.41), we get

$$\|\operatorname{Eig}(\iota A_{+}^{1/2} J_{2n} A_{+}^{1/2}) - \operatorname{Eig}(\iota B_{+}^{1/2} J_{2n} B_{+}^{1/2}) \|$$

$$\leq \|A_{+}^{1/2} J_{2n} A_{+}^{1/2} - B_{+}^{1/2} J_{2n} B_{+}^{1/2} \|$$
 (5.51)

$$\leq |||A_{+}^{1/2}J_{2n}A_{+}^{1/2} - A_{+}^{1/2}J_{2n}B_{+}^{1/2}||| + |||A_{+}^{1/2}J_{2n}B_{+}^{1/2} - B_{+}^{1/2}J_{2n}B_{+}^{1/2}|||$$
(5.52)

$$= |||A_{+}^{1/2}J_{2n}(A_{+}^{1/2} - B_{+}^{1/2})||| + |||(A_{+}^{1/2} - B_{+}^{1/2})J_{2n}B_{+}^{1/2}|||$$
(5.53)

$$\leq \|A_{+}^{1/2}J_{2n}\| \cdot \|A_{+}^{1/2} - B_{+}^{1/2}\| + \|A_{+}^{1/2} - B_{+}^{1/2}\| \cdot \|J_{2n}B_{+}^{1/2}\|$$
(5.54)

$$= \left(\|A_{+}^{1/2}\| + \|B_{+}^{1/2}\| \right) \|A_{+}^{1/2} - B_{+}^{1/2}\|$$
(5.55)

$$\leq \left(\|A_{+}^{1/2}\| + \|B_{+}^{1/2}\| \right) \|\|A_{+} - B_{+}\|^{1/2}\|.$$
(5.56)

Similarly,

$$\|\operatorname{Eig}(\iota A_{-}^{1/2}J_{2n}A_{-}^{1/2}) - \operatorname{Eig}(\iota B_{-}^{1/2}J_{2n}B_{-}^{1/2})\|$$

$$\leq \left(\|A_{-}^{1/2}\| + \|B_{-}^{1/2}\| \right) \| \|A_{-} - B_{-}\|^{1/2} \|.$$
(5.57)

Substituting (5.56) and (5.57) into (5.50) gives the desired perturbation bound (5.44). The other perturbation bounds (5.45) and (5.46) follow directly from (5.44).

Remark 5.5 In Proposition 5.4, if the matrices A and B are positive definite, then we have $A_- = B_- = 0$, $A_+ = A$, and $B_+ = B$. The perturbation bound (5.44) in this case reduces to the perturbation bound of symplectic eigenvalues of A and B given in Theorem 7 of [6].

Acknowledgments

The author acknowledges supports from the NSF under grant no. 2304816 and AFRL under agreement no. FA8750-23-2-0031. The author thanks Prof. Tanvi Jain for insightful discussions.

19

References

- Mikhael Gromov. Pseudo holomorphic curves in symplectic manifolds. *Inventiones mathematicae*, 82(2): 307–347, June 1985. http://dx.doi.org/10.1007/BF01388806.
- [2] Alessio Serafini, Fabrizio Illuminati, and Silvio De Siena. Symplectic invariants, entropic measures and correlations of gaussian states. *Journal of Physics B: Atomic, Molecular and Optical Physics*, 37(2):L21, December 2003. arXiv:quant-ph/0307073.
- [3] Jason L. Pereira, Leonardo Banchi, and Stefano Pirandola. Symplectic decomposition from submatrix determinants. Proceedings of the Royal Society A, 477(2255):20210513, November 2021. arXiv:2108.05364.
- [4] F. Nicacio. Williamson theorem in classical, quantum, and statistical physics. *American Journal of Physics*, 89(12):1139–1151, December 2021. http://dx.doi.org/10.1119/10.0005944. arXiv:2106.11965.
- [5] D. Šafránek, A. R. Lee, and I. Fuentes. Quantum parameter estimation using multi-mode Gaussian states. New Journal of Physics, 17(7):073016, 2015. arXiv:1502.07924.
- [6] Rajendra Bhatia and Tanvi Jain. On symplectic eigenvalues of positive definite matrices. Journal of Mathematical Physics, 56(11):112201, November 2015. http://dx.doi.org/10.1063/1.4935852.
 arXiv:1803.04647.
- [7] Fumio Hiai and Yongdo Lim. Log-majorizations for the (symplectic) eigenvalues of the Cartan barycenter. *Linear Algebra and its Applications*, 553:129–144, September 2018. http://dx.doi.org/10.1016/j.laa.2018.04.029. arXiv:1710.00494.
- [8] Hemant K. Mishra. First order sensitivity analysis of symplectic eigenvalues. Linear Algebra and its Applications, 604:324-345, November 2020. http://dx.doi.org/10.1016/j.laa.2020.07.003. arXiv:2007.10572.
- [9] Rajendra Bhatia and Tanvi Jain. A Schur-Horn theorem for symplectic eigenvalues. *Linear Algebra and its Applications*, 599:133–139, August 2020. http://dx.doi.org/10.1016/j.laa.2020.04.005. arXiv:2004.03906.
- [10] Rajendra Bhatia and Tanvi Jain. Variational principles for symplectic eigenvalues. Canadian Mathematical Bulletin, 64(3):553–559, September 2021. http://dx.doi.org/10.4153/S0008439520000648.
- Tanvi Jain. Sums and products of symplectic eigenvalues. *Linear Algebra and its Applications*, 631:67–82, December 2021. http://dx.doi.org/10.1016/j.laa.2021.08.016. arXiv:2108.10741.
- [12] Tanvi Jain and Hemant K. Mishra. Derivatives of symplectic eigenvalues and a Lidskii type theorem. *Canadian Journal of Mathematics*, 74(2):457–485, April 2022. http://dx.doi.org/10.4153/S0008414X2000084X. arXiv:2004.11024.
- [13] Paul-Emile Paradan. The Horn cone associated with symplectic eigenvalues. Comptes Rendus. Mathématique, 360(G10):1163–1168, October 2022. http://dx.doi.org/10.5802/crmath.383. arXiv:2202.10260.
- [14] Gajendra Babu and Hemant K. Mishra. Block perturbation of symplectic matrices in Williamson's theorem. *Canadian Mathematical Bulletin*, 67(1):201–214, March 2024. http://dx.doi.org/10.4153/S0008439523000620. arXiv:2307.01078.
- [15] Nguyen Thanh Son, P.-A. Absil, Bin Gao, and Tatjana Stykel. Computing symplectic eigenpairs of symmetric positive-definite matrices via trace minimization and Riemannian optimization. SIAM Journal on Matrix Analysis and Applications, 42(4):1732–1757, December 2021. http://dx.doi.org/10.1137/21M1390621. arXiv:2101.02618.
- [16] Shaowu Huang. A new version of Schur–Horn type theorem. Linear and Multilinear Algebra, 71(1):41–46, January 2023. http://dx.doi.org/10.1080/03081087.2021.2017834.
- [17] N. T. Son and T. Stykel. Symplectic eigenvalues of positive-semidefinite matrices and the trace minimization theorem. *Electronic Journal of Linear Algebra*, 38:607–616, September 2022. http://dx.doi.org/10.13001/ela.2022.7351. arXiv:2208.05291.
- [18] B. V. Rajarama Bhat and Tiju Cherian John. Real normal operators and Williamson's normal form. Acta Scientiarum Mathematicarum, 85:507–518, December 2019. http://dx.doi.org/10.14232/actasm-018-570-5. arXiv:1804.03921.
- [19] Tiju Cherian John, V. B. Kumar, and Anmary Tonny. An order relation between eigenvalues and symplectic eigenvalues of a class of infinite dimensional operators. June 2023. arXiv:2212.03900.
- [20] V. B. Kumar and Anmary Tonny. On approximating the symplectic spectrum of infinite-dimensional operators. *Journal of Mathematical Physics*, 65(4), April 2024. http://dx.doi.org/10.1063/5.0169600. arXiv:2303.09309.

- [21] Gerardo Adesso, Alessio Serafini, and Fabrizio Illuminati. Extremal entanglement and mixedness in continuous variable systems. *Physical Review A*, 70(2):022318, August 2004. http://dx.doi.org/10.1103/PhysRevA.70.022318, arXiv:quant-ph/0402124.
- [22] Xiao-yu Chen. Gaussian relative entropy of entanglement. *Physical Review A*, 71(6):062320, June 2005. http://dx.doi.org/10.1103/PhysRevA.71.062320, arXiv:quant-ph/0402109.
- [23] Martin Idel, Sebastián Soto Gaona, and Michael M. Wolf. Perturbation bounds for Williamson's symplectic normal form. *Linear Algebra and its Applications*, 525:45–58, July 2017. http://dx.doi.org/10.1016/j.laa.2017.03.013. arXiv:1609.01338.
- [24] Jen-Tsung Hsiang, Onat Arisoy, and Bei-Lok Hu. Entanglement dynamics of coupled quantum oscillators in independent nonMarkovian baths. *Entropy*, 24(12):1814, December 2022. http://dx.doi.org/10.3390/e24121814. arXiv:2211.07124.
- [25] John Williamson. On the algebraic problem concerning the normal forms of linear dynamical systems. American Journal of Mathematics, 58(1):141–163, 1936. http://dx.doi.org/10.2307/2371062.
- [26] Gerald B. Folland. Harmonic Analysis in Phase Space. Number 122. Princeton University Press, 1989. http://dx.doi.org/10.1515/9781400882427.
- [27] R. Simon, S. Chaturvedi, and V. Srinivasan. Congruences and canonical forms for a positive matrix: Application to the Schweinler–Wigner extremum principle. *Journal of Mathematical Physics*, 40(7):3632– 3642, July 1999. arXiv:math-ph/9811003.
- [28] Kh. D. Ikramov. On the symplectic eigenvalues of positive definite matrices. Moscow University Computational Mathematics and Cybernetics, 42:1–4, January 2018.
- [29] Hemant K. Mishra. Differential and subdifferential properties of symplectic eigenvalues. PhD thesis, Indian Statistical Institute, New Delhi, 2021. URL http://hdl.handle.net/10263/7232.
- [30] R. J. de la Cruz and H. Faßbender. On the diagonalizability of a matrix by a symplectic equivalence, similarity or congruence transformation. *Linear Algebra Appl.*, 496:288–306, 2016. http://dx.doi.org/10.1016/j.laa.2016.01.030.
- [31] Rajendra Bhatia. Matrix Analysis. Springer, 1997. http://dx.doi.org/10.1007/978-1-4612-0653-8.
- [32] Roger A. Horn and Charles R. Johnson. Matrix Analysis. Cambridge University Press, 2012. http://dx.doi.org/10.1017/CBO9781139020411.
- [33] Nathaniel Johnston. Advanced Linear and Matrix Algebra. Cambridge University Press, 2012. http://dx.doi.org/10.1017/CBO9781139020411.
- [34] Maurice A. de Gosson. Symplectic Geometry and Quantum Mechanics, volume 166. Springer Science & Business Media, 2006. http://dx.doi.org/10.1007/3-7643-7575-2.
- [35] Hemant K. Mishra. Equality in some symplectic eigenvalue inequalities. arXiv preprint, September 2023. arXiv:2309.04562.