

From new thermodynamics to classical mechanics

Henmei Ni *

School of Chemistry and Chemical Engineering, Southeast University, Southeast University Rd. 2#, Jiangning, Nanjing 211189, China. * henmei_ni@seu.edu.cn

Abstract

The laws of classical mechanics are rebuilt in the frame of new thermodynamics. Heat is the sum of kinetic energy, system work, and system potential of gas, while force is the linear gradients of heat variation. Exporters and importers of force are evident in terms of exotherm and endotherm. Temperature and volume gradients create asymmetric forces driving rotation and spin (self-rotation). It verifies that force transfer doesn't need a medium. As an outstanding achievement, a succinct and general equation is derived to predict the equilibrium distance of molecular interaction: $L_e = \sqrt[3]{\frac{3\pi^{\alpha-1}M_A g}{4N_A kT}}$, without using any assumption, such as van der Waals force and dispersion force. In addition, the origins and attributes of repulsion and attraction are disclosed. Predicting results is applause. For example, at 298 K, L_e for N₂, O₂, and CH₄ are 3.11, 3.11, and 3.68 Å, comparable to the data adopted in MD simulations of the literature. **Furthermore, the relationship between electricity and mass is established. An electron is characterized as a particle with a large α dependent on the interaction distance. Two electrons are mutually attractive from a distance depending on temperature. The electromagnetic phenomena are integrated.**

Keywords: Thermodynamics, Classical Mechanics, Heat, Lennard-Jones potential, Cosmos, Atmosphere, MD simulation

Introduction

The Big Bang is a widely accepted physical theory describing how the universe expanded from an initial high-density and high-temperature state ^[1]. It is proposed that the universe emerged from a "primeval atom," filled homogeneously and isotropically with an extremely high energy density, massive temperatures and pressures, and rapidly expanding and cooling. Under the interactions of the four fundamental forces- the electromagnetic force, the strong nuclear force, the weak nuclear force, and the gravitational force- today's world is created by mass condensation. It is similar to the phase transition of gas-solid and gas-liquid, i.e., evaporation (sublimation) -condensation, which has been well-studied in thermodynamics. Therefore, some common physical laws may exist.

Employing physical concepts to interpret thermodynamic phenomena was common in the past. For example, van der Waals introduced molecular interactions into the state equation of the ideal gas to correct the deviation of real gas ^[2]. Sutherland developed the hard-sphere model (the so-called Sutherland potential) by adding an attractive r^{-3} potential, identical to the van der Waals cohesive force, to improve the kinetic gas theory for atomic and molecular collisions ^[3]. Drude model ^[4] of oscillating dipoles between two atoms provides a straightforward way to introduce the dispersion force ^[5], ^[6].

London gave the famous attractive long-range r^{-6} behavior for the dispersion force [7], [8]. These are the natural responses to Newton's law of universal gravitation that molecular interaction is attractive. However, a repulsive force is needed to study the gas-liquid properties. Mie separated the two-body interaction potential into short-range repulsive and long-range attractive terms [9]. Based on Grüneisen potential within the solid-state and materials physics community [10], Lennard-Jones further developed it and gave a famous equation of potential: [11]-[14]

$$U_{LJ}(r) = 4\varepsilon \left[\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^6 \right] \quad (1)$$

Where r is the distance between the centers of two molecules, ε is the depth of the potential well, and σ the zero-crossing distance of the potential, then the potential U given by Lennard-Jones (LJ). The first term in the square bracket is repulsive, stemming from the electrons' repulsion in an atom, while the second term denotes the attractive. This potential is ideal for testing various algorithms used in atomistic molecular dynamics (MD) and Monte Carlo (MC) simulations [15], [16] to gain deeper insight into the theory of interatomic interactions in extended systems. Also, it has a widely applied interatomic potential for exploring important interactions in atomic and molecular systems and their bulk.

However, applications of classical mechanics cannot integrate the theory of thermodynamics. Quantum concepts are also applied in thermodynamics, but currently, quantum thermodynamics addresses the emergence of thermodynamic laws from quantum mechanics [17], [18]. Therefore, thermodynamics did not benefit from the current knowledge of the particular theory of relativity, quantum physics, and statistical mechanics [19], [20].

Thermodynamics are incoherent. Even a fundamental gas state equation has been argued for decades [21]-[26], let alone to derive a consistent and simple equation to predict the thermodynamic properties of substances. The reasons are summarized as follows. Methodologically, the research depended on intuition and the method of stimulation responses. For example, the gas system has not been detected as a pure gas or gas-liquid coexisting system. The results of system responses are summarized to form the empirical formula. However, the mechanism of system response is rarely a concern. The next is the unclear definition of concepts. Newton proposed the universal gravitation, but even he doubted [27]- "That one body may act upon another at a distance through a vacuum without the mediation of anything else, by and through which their action and force may be conveyed from one another, is to me so great an absurdity that, I believe, no man who has in philosophic matters a competent faculty of thinking could ever fall into it." Heat is another unclear concept in thermodynamics. It is described as "heat is the thermal energy transferred between systems due to a temperature difference" [28]. Thermal energy is the kinetic energy of vibrating and colliding atoms in a substance. However, the mechanism of heat transfer is not clear. It results in the structure defects of thermodynamic theorems. For example, in the definitions of internal energy,

$$dU = \delta Q - \delta W \quad (2)$$

U is the internal energy, and Q and W are the heat and work. In Eq. 1, the exact differential term (dU) is mixed with two inexact differential terms (δQ and δW). In mathematics, it is not compatible.

In our previous papers [29]-[31], a path of heat transfer was postulated based on the assumption that the heat transfers from a gas system to the surroundings through some gas molecules undergoing the inelastic collision with the container walls and then liquefy. It gave a new and concise gas state equation for the saturation system of gas-liquid [29], thereby excluding the classification of the ideal and real gas. Moreover, a new model for the specific heat capacities of gas at the saturation was

established^[30]. Furthermore, it is consistent with the equations of gas solubilities, gas-liquid interfacial tensions, and interfacial pressures independently derived from the analysis of solvation^[31]. Inspired by these achievements and the follow-up thermodynamic equations that will be published, and more importantly, it is independent of class mechanics; therefore, in this paper, I attempt to apply it to mechanics, inverse to the conventional methodology.

Results and Discussion

Definition of Heat

In the new theory of thermodynamics, a critical system energy is assumed above which the gas molecules can undergo inelastic collisions.

$$\varepsilon_{c,exo} = -\frac{3}{2}RT + RT \ln \frac{V_G}{V_L} + \frac{\pi^\alpha mg}{V_G} \quad (3)$$

where R is the gas constant (8.314 J·K⁻¹·mol⁻¹), T is the temperature (K), and V_L and V_G are the molar volumes of liquid and gas (L/mol), respectively. π is the circumference constant (3.14), α is an arbitrary factor, m is the molecule's molar mass, and g is the gravitational constant (9.18 m/s²). In addition, the minus sign denotes “lost,” or in other words, Eq. 3 is an exothermic equation.

On the right side of Eq. 3, the first item refers to the long-distance kinetic energy of molecules, and the second item refers to the spontaneous system work that a system does due to the loss of kinetic energy. The last item is arbitrarily added to correct the discrepancy of the state equation as the temperature approaches the critical temperature, T_c . I term it the system potential. α values have been given in Ref. [29]. Inversely, the endothermal equation of a gas system is,

$$\varepsilon_{c,endo} = \frac{3}{2}RT + RT \ln \frac{V_G}{V_L} - \frac{\pi^\alpha mg}{V_G} \quad (4)$$

Fig. 1 shows the scheme of energy exchange between a gas system and the surroundings. The thermal equilibrium is not Eq. 3+Eq. 4 = 0. The gas system loses or gains the long-term kinetic energy and system potential with spontaneous system work. System work is always positive in the endothermal and exothermic processes and offset by the outer counterparts. The details of how liquid molecules exchange heat with the surroundings via the container walls are unclear. However, since the liquid temperature likely equals gas, the long-term kinetic energy of gas molecules is converted into the vibrational energy of liquid molecules. Therefore, the energy change substantially exchanges system work and system potential. It is quite different from classical mechanics, in which the kinetic energy of particles can be converted into potential through doing work. Still, the work always equals the kinetic energy loss and the potential increase. Analogical to a gas system at T_c , $\frac{3}{2}RT = \frac{\pi^\alpha mg}{V_G}$, the system work is zero due to $V_L=V_G$.^[29] However, this is a case in which the energy exchange is interrupted due to a lack of gas molecules joining the exchange. Consequently, the liquefaction stops no matter how high the pressure is imposed. In other words, the system substantially becomes an adiabat system. At other temperatures of an isolated system, the system work is not equal to the increase of system potential; inversely, it is usually much larger than the kinetic energy loss and less converted into the system potential.^{[29],[30]} That is to say, the system amplifies the kinetic energy loss. However, it is not ridiculous for a system. To some extent, imaginatively like a traffic accident on a highway, a wave of stopping cars forms soon after the kinetic energy of two cars vanishes. The energy

loss of the highway system is much more significant than that of two cars. It indicates that the kinetic energy loss of some particles in a random system is just a triggering factor of an incident where every molecule spontaneously joins to carve up space as needed that the liquefied molecules left, thereby flying for an extra distance along a specific direction to do work. In this way, the system can release energy quickly to attain thermal equilibrium. It is not a paradox to the law of energy conversation but a different system choice. Therefore, classical mechanics suits an adiabat system but not an isolated one.

The molar ratio of liquefying/solidifying gas molecules x_2 at the thermal equilibrium is obtained by applying the Boltzmann distribution.

$$x_2 = \frac{V_L}{V_G} e^{\frac{3}{2} - \frac{\pi^\alpha mg}{RV_G T}} = \frac{V_L}{V_G} e^{\frac{3}{2} - \beta} \quad (5)$$

Eq. 5 works well in calculating gas solubility^[30] and specific heat capacity^[31]. Readers may ask why the molecular interaction is not reflected in the equation of state. As mentioned above, the interaction energy in liquid transforms into the bond energy stored in gas. Since gas has a similar V_G at an equal temperature, an intense interaction gives a small V_L , thus requiring much system work for evaporation. For example, water exhibits a high boiling point and a large latent heat of evaporation.

In summary, different from conventional thermodynamics, the new theory focuses on how gas molecules spontaneously adjust their states in the system when an outer factor stimulates it. Moreover, in classical thermodynamics, heat is defined as the kinetic energy transfer of vibrating and colliding atoms. However, as discussed above, any loss of kinetic energy will trigger the spontaneous actions of system molecules. System work is an indispensable term in the transfer of thermal energy.

Therefore, Eq. 3 and Eq.4 are defined as heat - the bodies of thermal energy exchange.




Figure 1, Scheme of heat exchange and the thermal equilibrium

Definition of Thermal Force

Fig. 2 describes the process in an isolated system where condensation spontaneously undergoes in quest of thermal equilibrium with the surroundings. In an isolated system, once two particles undergo the inelastic collision, the new entity must be stabilized by releasing the thermal energy, i.e., the thermal irradiation. Otherwise, the entity will be degraded. Therefore, the formation of a cool droplet is the result of evaporation (sublimation)- condensation process in which $x'_2 < x_2$. Here, it should be noted that thermal irradiation is also a kind of heat, which is not included in Eq. 4 and Eq. 5. Hence, the inclusion of thermal irradiation is the future work.

Once a droplet forms, the volumetric superiority renders the droplet incorporate more particles than the smaller ones. In addition, the combined effects such as the effect of inverse process, i.e.,

evaporation, the distance-dependent warming effects of thermal irradiation preferably to particles near the droplet surface, and the high probability of collision of high-temperature particles will make the gas phase form a gradient of V_G and T (Fig. 2), which distanced from a droplet, $\frac{\partial T}{\partial(L-r)} < 0$, but

$$\frac{\partial V_G}{\partial(L-r)} > 0.$$

On the other hand, given that the space of the system is large enough and the number of molecules as well, many droplets may form and be randomly distributed. Hence, the fate of gas molecules, including droplets, will be:

1) Fleeing away from the system. It will enlarge V_G . Imagining there is a boundary, according to Eq. 4, a considerable amount of system work will be done for liquefaction, thereby transferring the kinetic energy to the outer. An example is the Joule-Thomson expansion ^[32], in which the gas cools upon expansion. Therefore, if the space of a system is infinitive, it is deduced that those molecules will cool down to 0 K.

2) Circular motion around a droplet. The fleeing probability of a particle inside the system is extremely low by avoiding collisions when moving along the radii directions. Particles must be confined by a droplet acting as partners in the evaporation-condensation process. Some particles circularly move around a droplet to form the boundary layer among them. It does not directly work towards the droplet and the outer surroundings. However, it triggers the periodic fluctuation of V_G along the orbital, producing a ball-type V_G wave spreading to the whole gas phase of a droplet. Indeed, the particle itself will be impacted to produce the orbital wave. These waves do not release the energy at the equilibrium but cause the harmonic vibration of evaporation-condensation equilibrium. It may be the so-called tidal force. Therefore, changing the waves means damaging the equilibrium.

3) Capture by a droplet. The system's work is done towards the droplet, thus warming it. Some irradiation warms the gas phase, while the rest disperses to space. As a result, a gradient of temperature forms high near the droplet and low near the boundary.




Figure 2, Scheme of heat exchange in an isolated system

Based on the above discussion, heat varies during the thermal equilibrium. It can be exactly differentiated over time (t) and dimensions (distance, area and volume). For example, the heat exothermic flow rate is:

$$\frac{d\varepsilon_{c,exo}}{dt} = R \left(\ln \frac{V_G}{V_L} - \frac{3}{2} \right) \cdot \frac{dT}{dt} + (1 - \beta)RT \cdot \frac{d \ln V_G}{dt} - RT \frac{d \ln V_L}{dt} \quad (6)$$

Meanwhile, the linear differentiation, i.e., gradient, of exothermic heat is:

$$\frac{d\varepsilon_{c,exo}}{dL} = R \left(\ln \frac{V_G}{V_L} - \frac{3}{2} \right) \cdot \frac{dT}{dL} + (1 - \beta)RT \cdot \frac{d \ln V_G}{dL} - RT \frac{d \ln V_L}{dr} \quad (7)$$

where L and r are depicted in Fig. 2,

Eq. 7 is defined as the exothermic force.

$$F_{exo} = R \left(\ln \frac{V_G}{V_L} - \frac{3}{2} \right) \cdot \frac{dT}{dL} \vec{L} + (1 - \beta)RT \cdot \frac{d \ln V_G}{dL} \vec{L} - RT \frac{d \ln V_L}{dr} \vec{r} \quad (8)$$

\vec{L} and \vec{r} are the unit of vector indicating the force direction. Meanwhile, the endothermal force is the linear differentiation of Eq. 5,

$$F_{endo} = R \left(\ln \frac{V_G}{V_L} + \frac{3}{2} \right) \cdot \frac{dT}{dL} \vec{L} + (1 + \beta)RT \cdot \frac{d \ln V_G}{dL} \vec{L} - RT \frac{d \ln V_L}{dr} \vec{r} \quad (9)$$

Eq. 8 and Eq. 9 are two sides of a coin due to the heat conversation. If there is a donor, there must be acceptor.

Analysis of Thermal Force

Eq. 8 and Eq. 9 are substantially the statistical equations of a system with many members. Now, they are extended to classical mechanics to deal with the interactions between two isolated objects.

As shown in Fig. 1, the gas and outer systems can be regarded as two particles; the gas system is an exothermic particle (A), i.e., energy donor, while the outer system is an endothermal particle (B), i.e., energy acceptor. B is passive. It records how much A donated. Hence, the exothermic force of Eq. 8 is focused on in the following discussion, using the endothermal force of Eq. 9 as a comparison.

1) Let's discuss the kinetic energy transfer - collision.

Eq. 8 is simplified as,

$$F = -\frac{3}{2}R \cdot \frac{dT}{dL} \vec{L} \quad (10)$$

Back to the original form, $\frac{3}{2}RT = \frac{1}{2}mv^2$, where v is the average speed of particles. Therefore, considering the individual particle, Eq. 10 becomes,

$$F = -\frac{1}{2} \frac{d(mv^2)}{dL} \vec{L} \quad (11)$$

That is, force is the linear gradient of kinetic energy variation.

As shown in Fig. 3, A donates force to B with a velocity v , where B is steady, thus,

$$F = -\frac{1}{2}v^2 \frac{dm_A}{dL} \vec{L} - m_A v \frac{dv}{dL} \vec{L} \quad (12)$$

If $\frac{dm_A}{dL} = 0$, then

$$F = -m_A v \frac{dv}{v dt} = -m_A \vec{a} \quad (13)$$

where \vec{a} is the acceleration. Strictly, force has the "point" attribute. Hence, Eq. 13 deforms into,

$$F = -M_A \vec{a} \quad (14)$$

M_A is the mass of individual particle, i.e., $m_A = N_A \times M_A$, where N_A is Avogadro number.

If \vec{a} is negative, Eq. 13 is positive, meaning that A donates a force with A→B direction to B. Simultaneously B records the accepted force with Eq. 9. However, it should be noted that, standing on B, the linear gradient becomes positive, $\vec{a} > 0$. Thus, B certifies that B accepts a positive force from A. As a result, B is about to move along with the A→B direction. Colloquially, A pushes B.

Since A lost a positive force and slowed down in the A→B direction, we define A's lost force as a force with the inverse direction of A←B. Its quantity equals the positive force that B gains. Force cannot be created out of thin air. Such a relation of forces ←A-B→ is called mutual repulsion.

If \vec{a} is positive, Eq. 13 is negative. It means that A gains a force from B in the A→B direction rather than loses force. Simultaneously standing on B, \vec{a} is negative, Eq. 13 records a negative, too. It indicates that B doesn't accept any force from A; instead, B loses a force in the A→B direction. Similarly, defining B's lost force as a force with the A←B direction, its quantity equals the force that A gains from B with the A→B direction. Colloquially, B pulls A (attraction). Therefore, A→←B is called mutual attraction.

In summary, if A loses/gains a force, B simultaneously must gain/lose the force because of the heat conversation. The sign of Eq. 13 is always the same in exothermic and endothermic processes. Reaction doesn't exist in reality; it is defined to describe force's loss. According to the property of force, once A repulses B, B must repulse A, and inversely, too. This interprets the simultaneity of action and reaction, i.e., Newton's third law.




Figure 3, Interaction of two particles

If $\frac{dm_A}{dL} < 0$, then, Eq. 10 deforms into,

$$F = -\frac{1}{2}v^2 \frac{dM_A}{dL} \vec{L} - M_A \vec{a} \quad (15)$$

It indicates that losing A mass adds a positive force on B, regardless of the initial force's property.

2) Let's consider the temperature variation.

In Eq. 8

$$F = k \ln \frac{V_G}{V_L} \cdot \frac{dT}{dL} \vec{L} \quad (16)$$

k is Boltzmann constant. The positive sign is the same in Eq. 8 and Eq. 9. The term of $kT \ln \frac{V_G}{V_L}$ is the largest value in Eq. 3. Additionally, in most cases, $V_G \gg V_L$, and V_L is almost constant. Hence,

$$F \approx k \ln V_G \cdot \frac{dT}{dL} \vec{L} \quad (17)$$




Figure 4, Scheme of force created by the temperature gradient

As shown in Fig. 4, considering A→B system where B's temperature is zero, since $T = \frac{M_A v^2}{3k}$, and $V_G = \frac{4}{3} N_A \pi L^3$ (N_A is Avogadro number),

$$F = \frac{M_A}{3} \ln \left(\frac{4}{3} N_A \pi L^3 \right) \vec{a} = (18.7289 + \ln L) M_A \vec{a} \quad (18)$$

Here, the SI unit of L is dm . Since the temperature variation gradient is negative, Eq. 18 is negative. It indicates that the high-temperature A donates an attractive force to the low-temperature B. However, B sees a positive \vec{a} . Hence, B records a positive force with Eq. 18. It contrasts with Eq. 13, though $\vec{a} < 0$ in both equations.

The symmetry of action and reaction is destroyed. However, A and B don't lie. A unique explanation is that A rotates around B, and B spins at the request of two forces. There is no exchange of kinetic energy.

3) Let's consider the volumetric term.

As shown in Fig. 5 (left), the V_G variation gradient is created as A approaches B, where B is still.




Figure 5, Scheme of volumetric force

$$\text{If } \frac{dT}{dL} = 0, \text{ and } \frac{dV_L}{dr} = 0,$$

In Eq. 8,

$$F = (1 - \beta) RT \cdot \frac{d \ln V_G}{dL} \vec{L} \quad (19)$$

As shown in Fig. 5 (left), since there is only one particle, A's V_G decreases because L decreases. Hence,

$V_G = \frac{4}{3}N_A\pi L^3$ where N_A is Avogadro constant, 6.022×10^{23} . Here, it should be noted that the unit of L

is dm because the unit of V_G is L/mol . Therefore, remembering that $\frac{dV_G}{dL} < 0$,

$$F = \frac{9\pi^{\alpha-1}M_Ag}{4N_AL^4} - \frac{3kT}{L} \quad (20)$$

or $F = \frac{9\pi^{\alpha-1}M_Ag}{4N_AL^4} - \frac{M_Av^2}{L}$

It says that A exerts two opposite forces on B. One is positive, i.e., repulsive, and inversely proportional to the 4th power of distance, whereas the other is negative, i.e., attractive, but inversely proportional to the distance. At the point of zero force,

$$L_e = \sqrt[3]{\frac{3\pi^{\alpha-1}M_Ag}{4N_AkT}} \quad (21)$$

Provided that A and B are two molecules, L_e is calculated at 298K and listed in Tab 1.

Table 1 Calculated the equilibrium distance of gas molecules at 298 K

Gas	α	m_A (g/mol)	L_e (Å)	Ref. Value	Gas	α	m_A (g/mol)	L_e (Å)	Ref. Value
He	-1.83	4	0.92		CH ₄	0.6	16	3.68	3.74 ^[35] , 3.71 ^[38]
Ne	-1.64	20	1.69	2.80 ^[34]	CH ₃ CH ₃	0.82	30	4.94	3.51 ^[35] , 4.89 ^[36]
Ar	-0.69	40	3.05	3.40 ^[35]	C ₃ H ₈	0.88	44	5.74	5.56 ^[36]
Kr	-0.87	121.8	4.13	3.63 ^[34] , [38]	C ₄ H ₁₀	0.95	58	6.46	
Xe	-0.74	131.2	4.45	3.95 ^[34] , [38]	Ethylene	0.67	28	4.56	3.54 ^[38]
H ₂	0.54	2	1.80		Benzene	0.96	78	7.16	
N ₂	-1/3	28	3.11	3.31 ^[35] , 4.13 ^[36]	Toluene	1.0	92	7.68	
O ₂	-0.45	32	3.11	3.20 ^[37] , 3.95 ^[38]	NF ₃	-1/3	74	4.30	
F ₂	-0.78	40	2.95	2.83 ^[35]	CF ₄	-0.45	92	4.42	5.21 ^[36]
CO	-0.29	28	3.16	3.17 ^[38]	SF ₆	-1/3	152	5.46	5.18 ^[36]
CO ₂	0.07	44	4.21	2.94 ^[35] , 4.38 ^[36]	CH ₃ OH	0.93	32	5.26	
SO ₂	0.25	64	5.11		NH ₃	0.87	17	4.16	3.90 ^[36]
H ₂ O	0.95	18	4.38	3.57 ^[36]	H ₂ S	0.49	34	4.54	

Note: α values have been given in Reference [29]

There is no standard data in the literature, though the van der Waals's radii of elements were calculated based on the equilibrium distance^[33]. However, the data in Tab. 1 sound rational, according to the empirical sense of the equilibrium distance used in the molecule dynamics (MD) simulations^{[34]-[38]}.

Compared with the Lennard-Jones potential (Eq. 1)^{[11]-[14]}, the derivation of Eq. 20 is straightforward and concise without adopting any fictional assumptions, such as the electron repulsion^[3] or the dispersion force^[5]. Moreover, Eq. 20 indicates that L_e decreases as the temperature increases. This is consistent with the anticipation that the faster the particles, the closer the accessible distance, which is vital to the chemical reaction.

Furthermore, the molecules must do extra work due to these forces as the volume expands or shrinks. As shown in Eq 20, the attractive force is contributed by the increase of molar volume in the exothermic process, whereas the repulsive force is stemmed from the system potential. Hence, the

system releases the system potential by doing the negative work.

However, the equation of endothermal force gives a different result. According to Eq. 9, and $\frac{dV_G}{dL} > 0$, Eq. 19 becomes,

$$F = (1 + \beta)kT \cdot \frac{d \ln V_G}{dL} \vec{L} \quad (22)$$

$$F = \frac{9\pi^{\alpha-1} M_A g}{4N_A L^4} + \frac{3kT}{L} \quad (23)$$

It indicates that B accepts two repulsive forces from A. Remembering Eq. 20 that A exports a repulsive ($F_{A \rightarrow B}$) and imports an attractive force from B ($F'_{A \leftarrow B}$). Like Eq. 18, B records that it imported $F_{A \rightarrow B}$ but simultaneously imported $F'_{A \rightarrow B}$. Fig. 6 shows a possible mechanism according to the force's "point" attribute. Considering B's geometry, if A donates a positive force to any point but the central point of B, B immediately creates an inverse force on the opposite end. Therefore, both A and B record getting $F'_{A \rightarrow B}$ from the counterpart. A accepts $F'_{A \rightarrow B}$ to compensate for the loss of $F_{A \rightarrow B}$. Accordingly, A changes the direction of velocity due to attraction.




Figure 6, Possible mechanism of spin created by the asymmetric force

As shown in Fig. 5 (right), B starts to spin while A rotates around B. This is a popular pattern in the cosmos. Nevertheless, equilibrium distance is relevant because, at that position, A attains the force balance, namely that A loses the weight.

On the other hand, as shown in Tab 1, the effects of molecule polarity are not found in the equilibrium distances. Reminiscing the state equation ^[29], where the particularity of polar molecules was not seen yet, it indicates that polar interactions may be shorter-term than molecular interactions.

A general scenario is depicted in Fig. 7, where A approaches B with an angle θ .




Figure 7, A approaches B with an angle

$$d \ln L = -\frac{\cos \theta}{L} \cdot dv + \frac{v \sin \theta}{L} \cdot d\theta \quad (24)$$

$$dF = \frac{Fv}{L} (\cos \theta \cdot d \ln v - \sin \theta \cdot d\theta) \quad (25)$$

If $dF=0$, there are two cases. First, $F=0$. It gives the equilibrium distance, indicating that the mutual attraction turns into mutual repulsion, i.e., a change of force direction. Second, independent of the position of $F=0$, the strength of force turns from strong to weak. It gives the variation of energy.

$$d \ln v = -d \ln \cos \theta \quad (26)$$

Then, conducting the integration from (θ_0, v_0) to (θ_t, v_t) , where $\theta_0 \in [0, \frac{\pi}{2}]$, $\theta_t \in [0, \frac{\pi}{2}]$,

$$v_t \cos \theta_t = v_0 \cos \theta_0 \quad (27)$$

Hence, the variation of kinetic energy or system work is,

$$\Delta E_k = \frac{1}{2} M_A v_0^2 \left(1 - \frac{\cos^2 \theta_0}{\cos^2 \theta_t}\right) \quad (28)$$

If $\theta_t < \theta_0$, $\Delta E_k > 0$, whereas $\theta_t > \theta_0$, $\Delta E_k < 0$. As v_t tangents a cycle, $\theta_t = \frac{\pi}{2}$, the maximum ΔE_k attains. If A is far enough, $\theta_0 \approx \frac{r}{L_0}$, where r is the radius of B and L_0 is the distance. $\theta_t = \frac{\pi}{2}$ is achievable. Furthermore, if $\theta_t = \frac{\pi}{2}$ and $F = 0$, A will cycle B with v_t . Apart from the attractive force, as A approaches B, θ increases, while as A leaves B, θ decreases. Hence, Eq. 25 describes the elliptic motion. In addition, increasing v_0 lifts the frequency of motion and ΔE_k . However, since θ is dependent on L , increasing θ_0 lowers the frequency but increases the long radius of the ellipse. Therefore, for escaping from B, changing θ_0 is more effective. Moreover, Eq. 28 implies that the kinetic energy of a particle will gradually decrease even though it escapes from a planet and travels in a vacuum.

On the other hand, as shown in Fig. 7, provided that the forces exerted on a point of B and turn B to spin. The speed of exerting points obeys,

$$\frac{d\varphi}{dt} = \frac{v \sin \theta}{L} \quad (29)$$

For a homogeneous and rigid sphere B, the net torque is zero as A passes by. However, the net torque will drive B to spin if B is not a sphere or the mass distribution is heterogeneous.

4) Let's consider the liquid or solid term.

$$F = -RT \frac{d \ln v_L}{dr} \vec{r} = -\frac{3kT}{r} \vec{r} \quad (30)$$

This term is negative in both exothermic and endothermal equations. As shown in Fig. 8, in the exothermic process, $\frac{d v_L}{dr} < 0$,^[40] thus, A donates a positive force to B. Inversely, in the endothermal process, B records accepting a negative force from A. It is similar to the asymmetric situation discussed above, but as shown in Fig. 8, they are contract forces in liquid. We cannot imagine that A will rotate around B in a dense liquid and solid. However, resembling the terms in Eq. 20 and Eq. 23, it can be concluded that the property of force is rotational. Since Eq. 19 succeeds in predicting equilibrium

distance, it is expected that Eq. 30 is suitable for predicting the properties of liquid and solid, which most MD simulations work for [34]-[39]. For example, the liquid droplets simultaneously deform into spheres suspended in the atmosphere. Conventionally, it is explained that the droplets simultaneously deform their shapes, which possess the smallest surface area, to decrease the surface/interfacial tension. The unbalanced molecular interactions of surface molecules or cohesive forces are designated as the driving force. However, this explanation is not convincing due to the many exceptions. For example, the interfacial tensions of water-air, benzene-air, and ethanol-air are 72.75, 28.88, and 22.39 mN/m at 293.17 K. [41] It is hard to believe that the molecular interactions in benzene are more potent than the hydrogen bonds in ethanol. Moreover, the interfacial tension is practically zero in the atmosphere of liquid-gas saturation. Hence, Eq. 30 gives the driving force of deformation but weakly relates to the surface tension.




Figure 8, Scheme of volumetric force in liquid

The analysis of heat force has been completed. In summary, the forces created by collision are symmetrical, whereas the forces produced by temperature variation gradient and volumetric variation gradient are asymmetric. The symmetric forces create the linear motion, while the others create the spin or rotation. These disclose the essence of forces and the origins of various motions, such as spin and rotation, which are popular models in the cosmos. All equations solve Newton's "causes hitherto unknown." [27] The origin of force is heat transfer, which doesn't need a medium for the force transfer. These results indicate that the definition of heat is appropriate, and by this definition, the new thermodynamics integrates classical mechanics well.

Newton defined force as the gradient of speed momentum over time. As discussed above, it can describe the collision behavior of objects like Eq. 14. However, it cannot explain the forces stemming from the temperature gradient and V_G gradient. Universal gravitation exists, but it is an integration of various forces. Sun created a temperature gradient in the solar system. Hence, once an object enters the solar system, the Sun donates an attractive force to the object because the Sun sees a decreasing temperature gradient (Eq. 16). However, the object records a repulsive force. Hence, it rotates in the quest of two forces. Linear motion is so tricky in the universe.

At thermal equilibrium, the lucky objects have a common fatality: rotation with an elliptic orbit, which periodically de/accelerates them in the master's temperature gradient. This orbit is eternal, in contrast to the regular circular orbit.

Electricity and Mass

Finally, Eq. 19 is significant in the microscopic world. Inspired by its success in calculating equilibrium distance, I tried to apply it to calculate the force between two electrons in a vacuum ($T=0$).

$$F = \frac{90\pi^{\alpha-1}M_e g}{4N_A L^4} = 3.66 \times 10^{-1} \frac{\pi^{\alpha-1}M_e}{L^4} \quad (31)$$

In Eq. 25, The SI unit of L is m , and the mass of electrons (M_e) is 9.11×10^{-28} g. However, the α value for electrons is unknown. Provided that two electrons are separated with 10^{-10} m, i.e., 1.0 \AA , the repulsive force calculated with Coulomb's law is 2.30×10^{-8} N. However, Yu and Polycarpou issued the force difference of molecular interaction, $\sim 10^{-5}$ N in $1.0 \sim 5.0 \text{ \AA}$ [39]. It is much bigger than the coulomb's force (10^{-8} N/ 1.0 \AA) of two electrons. Hence, their model was not appropriate. In contrast, the calculated repulsive molecular interaction at the equilibrium distance by Eq. 20 is about $\sim 10^{-11}$ N/ 1.0 \AA . Of course, the attractive molecular interaction is also in the same order, depending on temperature. It indicates that Eq. 20 is appulsive in the microscopic world.

Since A and B are exchangeable, if $M_A \neq M_B$, the vector addition gives,

$$F = \frac{3.66 \times 10^{-19}}{L^4} (\pi^{\alpha_A-1} M_A + \pi^{\alpha_B-1} M_B) \quad (32)$$

Assuming that A and B are electrons, and

$$F = \frac{7.32 \times 10^{-19} \pi^{\alpha-1} M_e}{L^4} = \frac{8.99 \times 10^9 Q^2}{L^2} \quad (33)$$

Then,

$$\alpha = 1.7479 \ln L + 36.277 \quad (34)$$

Here, Q is the electric charge of an electron, 1.60×10^{-19} C.

Fig. 9 exhibits the interactions of two electrons. The repulsion calculated with α in Eq. 34 is consistent with Coulomb's law. However, three temperatures are applied because the attractive force depends on the temperature. In addition, the attractive force ($\frac{6kT}{L}$) is doubled, similar to Eq. 33. As shown in Fig. 9, at 4 K, the attractive force is more potent than the repulsive force at a distance longer than about 1 \mu m . It indicates that two electrons mutually attract at a distance $> 1 \text{ \mu m}$. As less than 1 \mu m , the repulsive force is potent. At 298 and 2000 K, the distances are about 10 and 2 nm, respectively. That is to say, as temperature rises, the distance decreases.

Common sense may think it ridiculous that two electrons are mutually attractive. However, this can explain a charged object in which many electrons coexist on the surface rather than homogeneously distributed throughout the body. Moreover, the rotational attribute of attractive force may be regarded by miss as the repulsive force because an electron looks like to avoid another electron.




Figure 9, Mutual interactions between two electrons

The electron is characterized as a particle with a large α , which depends on the interaction distance. It is pretty different from other molecules. Since the mass of an electron is 9.11×10^{-31} kg, the interaction (attraction) distance of two electrons can reach 10^{11} m at 298 K, at which an electron can get an acceleration of about 0.2708 m/s². Obviously, the higher the temperature, the longer the distance. Therefore, the interaction speed is determined by how fast a measuring method is or when an electron appears at the position. That is to say, we can see the present only rather than the past. It discloses the essence of universal gravitation. Moreover, if changing the electron kinetic energy,

$$F = \frac{6k}{L} T \sin(\omega t) = \frac{2M_e}{L} v^2 \sin(\omega t) \quad (35)$$

A distant electron may sense the change and then duplicate it. It resembles the sound in which the kinetic energy of molecules is periodically changed. Of course, electrons' velocity and confined degree determine the duplication speed and deformation degree.

Fig. 9 implies that an atom's temperature is very high, namely that electrons are confined in a tiny space by attraction. This coincides with the thermodynamics of phase transition: an atom is a highly compressed system. Cations and anions are non-equilibrium systems; therefore, they are different from electrons, though they are also “charged.”

Others

On the other hand, if the force definition is correct, the differential of Eq. 4 over V_G should be the gas pressure. For simplification, the gas isotherm is employed.

$$\frac{\varepsilon_{c,endo}}{dV_G} = RT \left(\frac{1}{V_G} - \frac{d \ln V_L}{dV_G} \right) + \frac{\pi^\alpha m g}{V_G^2} \quad (36)$$

At the isotherm, compared with V_G , the variation of V_L is negligible even at high pressure [42].

$$P = RT \frac{1}{V_G} + \frac{\pi^\alpha m g}{V_G^2} = (1 + \beta) \frac{RT}{V_G} \quad (37)$$

Eq. 37 is the state equation of gas at isotherm, on which the paper is underwriting. Eq. 37 is also believed to be able to describe the state of atoms. As shown in Fig. 10, the variation of V_G exhibits the same periodicity as the elements. Since $\beta \rightarrow 0$, the pressure of an atom periodically varies from $\sim 10^4 RT$ to $\sim 10^6 RT$ as the proton number regularly varies. Here, it should be noted that T denotes the temperature in an atom. It may be defined as the electronic temperature.




Figure 10, Periodicity of average V_G of electrons in an atom

Anyway, the purpose of this section is to support the correctness of the definition of force.

Conclusion

Employing principles of classical mechanics to build up thermodynamics has been a hot topic for decades. Inversing this approach, this paper attempts to integrate classical mechanics into the present author's new thermodynamics. First, the concepts of heat and force are defined. Heat is the sum of molecular kinetic energy, system work, and system potential, while force is the linear gradient of heat variation. Since the heat variation can be divided into the exothermic and endothermal processes, the exporter and importer of force are clear. Hence, the forces' exports and imports are checkable. It solves the problem of Newton's "causes hitherto unknown." Second, Newton's three laws are rebuilt based on new definitions. The force equation has four terms of variation gradients: kinetic energy, temperature, gas molar volume, and liquid/solid molar volume. Only the gradient of kinetic energy variation creates the symmetric forces, i.e., mutual repulsion and attraction, though the reaction force doesn't exist. The reaction force is defined to represent the loss of kinetic energy. The other terms create asymmetric forces, namely that the force direction exported is opposite to that imported, which are the deriving forces of rotation and spin (self-rotation). Of most importance, these results relieve Newton's anxiety that the force transfer doesn't need a medium. Therefore, the origins and attributes of forces are disclosed. Third, as an outstanding achievement, a succinct equation is derived to predict the equilibrium distance of molecular interaction.

$$L_e = \sqrt[3]{\frac{3\pi^{\alpha-1}M_A g}{4N_A kT}}$$

Numerous works on calculus, MD simulations, and Mont Carlo simulations have been published based on the Lennard-Jones potential model. Equilibrium distance is a critical factor in those works. Many theories, such as the van der Waals cohesive and dispersion forces, must be proposed to calculate the equilibrium distance. However, the above equation discloses a simple world with the rational results. For example, at 298 K, L_e for N_2 , O_2 , and CH_4 are 3.11, 3.11, and 3.68 Å, comparable to the data adopted in MD simulations of the literature. Moreover, the equation includes the temperature factor.

An electron is characterized as a particle with a large α dependent on the interaction distance. Two electrons are attractive at a distance depending on the temperature. For example, at 4, 298, and 2000 K, the attractive force is more potent than the repulsive force at a distance $>1 \mu\text{m}$, 10, and 2 nm. Since the rotational attribute of attractive force, the attraction is possibly regarded as the repulsion by miss because an electron looks to avoid another one. Interactions of electrons in the cosmos exist independently of time. Hence, the measuring methods determine the interaction speed. The electronic temperature determines the interaction distance of electrons. For example, at 298 K, the distance can reach 10^8 km, at which point an electron can get 0.2708 m/s^2 acceleration. Therefore, we can see the distant stars with high temperatures but not the neighboring planets with low temperatures. Once the motion of electrons is adjusted, all electrons in the cosmos sense it and duplicate it. The status of electrons determines the duplication speed and deformation degree. It is called reflection, refraction, and diffraction. In summary, the above results indicate that classical mechanics can be integrated into the frame of new thermodynamics. Additionally, it disclosed that evaporation (sublimation) –

condensation is a standard method of creating objects in the cosmos. It is believed that all objects, from tiny atoms to giant objects like planets, are thermal equilibrium systems. The new thermodynamics can be applied to study them from the internal structure to the outer behaviors.

Acknowledgments

The paper is completed independently by the author without any supporting funds.

Conflict of Interests


The authors declare no conflict of interest.

References

- [1]. Wikipedia: <https://en.wikipedia.org/wiki/Big-Bang>
- [2]. J. D. Van der Waals, *Over de Continuïteit van den Gasen Vloeistoestand*, Sijthoff, 1873; Vol. 1.
- [3]. W. Sutherland, *London, Edinburgh Dublin Philos. Mag. J. Sci.* 1893, 36, 507–531. LII. The viscosity of gases and molecular force.
- [4]. P. Drude, *Ann. Phys.* 1900, 306, 566–613. Zur Elektronentheorie der Metalle.
- [5]. W. L. Bade, *J. Chem. Phys.* 1957, 27, 1280–1284. Drude-Model Calculation of Dispersion Forces. I. General Theory.
- [6]. J. O. Hirschfelder, C. F. Curtiss, R. B. Bird, *The Molecular Theory of Gases and Liquids*; John Wiley & Sons: New York, 1964.
- [7]. F. London, *Z. Phys.* 1930, 63, 245–279. Zur Theorie und Systematik der Molekularkräfte.
- [8]. F. London, *Trans. Faraday Soc.* 1937, 33, 8b–26. The general theory of molecular forces.
- [9]. G. Mie, *Ann. Phys.* 1903, 316, 657–697. Zur kinetischen Theorie der einatomigen Körper.
- [10]. E. Grüneisen, *Ann. Phys.* 1912, 344, 257–306. Theorie des festen Zustandes einatomiger Elemente.
- [11]. J. E. Jones, *Proc. R. Soc. London, Ser. A* 1924, 106, 441–462. On the Determination of Molecular Fields. I. From the Variation of the Viscosity of a Gas with Temperature.
- [12]. J. E. Jones, *Proc. R. Soc. London, Ser. A* 1924, 106, 463–477. On the Determination of Molecular Fields. II. From the Equation of State of a Gas.
- [13]. J. E. Jones, *Proc. R. Soc. London, Ser. A* 1924, 106, 709–718. On the Determination of Molecular Fields. III. From crystal measurements and kinetic theory data.
- [14]. P. Schwerdtfeger, D.J. Wales, *J. Chem. Theory Comput.* 2024, 20, 3379–3405. 100 Years of the Lennard-Jones Potential.
- [15]. M. P. Allen, D. J. Tildesley, *Computer Simulation of Liquids*; Clarendon Press: New York, 1989.
- [16]. D. Frenkel, B. Smit, *Understanding Molecular Simulation: From Algorithms to Applications*, 2nd ed.; Academic Press: London, 2002.
- [17]. S. Deffner, S. Campbell, *Quantum Thermodynamics: An introduction to the thermodynamics of quantum information*, Morgan & Claypool Publishers, 2019.
- [18]. F. Binder, L.A. Correa, C. Gogolin, J. Anders, G. Adesso, *Thermodynamics in the Quantum Regime. Fundamental Theories of Physics*, Springer, 2018.
- [19]. B. Poirier, *A Conceptual Guide to Thermodynamics*, Wiley, 2014.

- [20]. J. Güemez, J. A. Mier, *Eur. J. Phys.* 2024, 45, 015701. Relativistic mechanics and thermodynamics: IV. Thermodynamic processes.
- [21]. O. Redlich, J. N. S. Kwong, *Chem. Rev.* 1949 44 (1): 233–244. On the Thermodynamics of Solutions. V. An Equation of State. Fugacities of Gaseous Solutions.
- [22]. D. Y. Peng, D. B. Robinson, *Ind. Eng. Chem.: Fundamentals.* 1976 15: 59–64. A New Two-Constant Equation of State.
- [23]. R. Stryjek, J. H. Vera, *Can. J. Chem. Eng.* 1986, 64 (5): 820–826. PRSV2: A cubic equation of state for accurate vapor-liquid equilibria calculations.
- [24]. A. Kaplun, A. Meshalkin, *EPJ Web of Conferences*, 2014, 76, 01026. Unified low-parametrical equation of state for engineering calculations of thermodynamic properties of substances.
- [25]. J. D. Dymond, R. C. Wilhoit, *Virial coefficients of pure gases and mixtures*, Springer, 2003.
- [26]. M. Benedict, G.B. Webb, L. C. Rubin, *J. Chem. Phys.*, 1940, 8 (4): 334–345. An Empirical Equation for Thermodynamic Properties of Light Hydrocarbons and Their Mixtures: I. Methane, Ethane, Propane, and n-Butane.
- [27]. Wikipedia: https://en.wikipedia.org/wiki/Newton%27s_law_of_universal_gravitation.
- [28]. Van W. Gordon, S. Richard, *Fundamentals of Classical Thermodynamics (2nd edition). Chapter 4.7, Definition of Heat*, John Wiley & Sons, 1978.
- [29]. H.-M. Ni, B. Zhu, *ChemRxiv*, 2022, *Preprint*. DOI: 10.26434/chemrxiv-2022-5m718-v4. A general and simple state equation for all gases.
- [30]. H.-M. Ni, *J. Mol. Liquids*, 2022, 368, 120690. On the Hydrophobic Hydration, Solvation and Interface: A Thought Essay (I).
- [31]. X.-Y. Cui, B. Zhu, H.-M. Ni, *ChemistrySelect*, 2024, 9, e202304699 (1 of 10). A New Model for Specific Heat Capacity of Real Gas.
- [32]. W. G. Hoover, C. G. Hoover, K. P. Travis, *Phys. Rev. Lett.* 2014, 112 (14): 144504. Shock-Wave Compression and Joule–Thomson Expansion.
- [33]. S. S. Batsanov, *Inorg. Mat.* 2001, 37 (9): 871–885. Van der Waals Radii of Elements.
- [34]. J. Fischer, M. Wendland, *Fluid Phase Equilibria*, 2023, 573, 113876. On the history of key empirical intermolecular potentials.
- [35]. J. Fischer, R. Lustig, H. Breitenfelder-Manske, W. Lemming, *Mol. Phys.* 1984, 52(2), 485–497. Influence of intermolecular potential parameters on orthobaric properties of fluids consisting of spherical and linear molecules.
- [36]. Y. Song, E. A. Mason, *Phys. Rev. A*, 1990, 42(8), 4749–4755. Analytical equation of state for molecular fluids: Comparison with experimental data.
- [37]. R. Lustig, *Ber. Bunsenges. Phys. Chem.* 1994, 98 (5), 706–711. Thermodynamics of Liquid Oxygen from Molecular Dynamics Simulations.
- [38]. T. Boulik, *J. Chem. Phys.* 1987, 87, 1751–1756. Simple perturbation method for convex-molecule fluids
- [39]. N. Yu, A. A. Polycarpou, *J. Coll. Interf. Sci.* 2004, 278, 428–435. Adhesive contact based on the Lennard–Jones potential: a correction to the value of the equilibrium distance as used in the potential.
- [40]. G. Lamanna, J. van Poppel, M.E.H. van Dongen, *Experiments in Fluids*, 2002, 32, 381–395. Experimental Determination of Droplet Size and Density Field in Condensing Flows.
- [41]. A. W. Adamson, A. P. Gast., *Physical chemistry of surfaces*; 6 Ed, Wiley, 1997.

TOC



Evaporation-Condensation system

$$\text{Force: } F_{exo} = R \left(\ln \frac{V_G}{V_L} - \frac{3}{2} \right) \cdot \frac{dT}{dL} \vec{L} + (1 - \beta) RT \cdot \frac{d \ln V_G}{dL} \vec{L} - RT \frac{d \ln V_L}{dr} \vec{r}$$

$$\text{Equilibrium distance of molecular interaction: } L_e = \sqrt[3]{\frac{3\pi^{\alpha-1} m_A g}{4N_A RT}}$$