A CHARACTERIZATION FOR AN ALMOST MDS CODE TO BE A NEAR MDS CODE AND A PROOF OF THE GENG-YANG-ZHANG-ZHOU CONJECTURE

SHIYUAN QIANG, HUAKAI WEI, AND SHAOFANG HONG*

ABSTRACT. Let \mathbb{F}_q be the finite field of q elements, where $q = p^m$ with p being a prime number and m being a positive integer. Let $C_{(q,n,\delta,h)}$ be a class of BCH codes of length n and designed δ . A linear code C is said to be maximum distance separable (MDS) if the minimum distance d = n - k + 1. If d = n - k, then Cis called an almost MDS (AMDS) code. Moreover, if both of C and its dual code C^{\perp} are AMDS, then C is called a near MDS (NMDS) code. In [A class of almost MDS codes, *Finite Fields Appl.* **79** (2022), #101996], Geng, Yang, Zhang and Zhou proved that the BCH code $C_{(q,q+1,3,4)}$ is an almost MDS code, where $q = 3^m$ and m is an odd integer, and they also showed that its parameters is [q + 1, q - 3, 4]. Furthermore, they proposed a conjecture stating that the dual code $C^{\perp}_{(q,q+1,3,4)}$ is also an AMDS code with parameters [q + 1, 4, q - 3]. In this paper, we first present a characterization for the dual code of an almost MDS code to be an almost MDS code. Then we use this result to show that the Geng-Yang-Zhang-Zhou conjecture is true. Our result together with the Geng-Yang-Zhang-Zhou theorem implies that the BCH code $C_{(q,q+1,3,4)}$ is a near MDS code.

1. INTRODUCTION

Let \mathbb{F}_q be the finite field of q elements, where $q = p^m$ with prime p and positive integer m, \mathbb{F}_q^n is an n-dimensional vector space over \mathbb{F}_q . Let $\mathbb{F}_q^* := \mathbb{F}_q \setminus \{0\}$ be the set of all non-zero elements of \mathbb{F}_q . Then \mathbb{F}_q^* forms a group under the multiplicative operation. For a non-empty set $\mathcal{C} \subseteq \mathbb{F}_q^n$, if \mathcal{C} is a k-dimensional subspace of \mathbb{F}_q^n , then \mathcal{C} is called an [n, k] linear code over \mathbb{F}_q , and n and k are called the *length* and *dimension* of the code \mathcal{C} , respectively. And we call the vector \mathbf{c} in \mathcal{C} codeword. Then the linear code \mathcal{C} holds q^k codewords. The dual code, denoted by \mathcal{C}^{\perp} , of an [n, k] linear code \mathcal{C} over \mathbb{F}_q is defined by $\mathcal{C}^{\perp} := \{\mathbf{c}^{\perp} \in \mathbb{F}_q^n \mid \langle \mathbf{c}, \mathbf{c}^{\perp} \rangle = 0, \mathbf{c} \in \mathcal{C}\}$ with $\langle \mathbf{c}, \mathbf{c}^{\perp} \rangle$ denoting the Euclidean inner product of \mathbf{c} and \mathbf{c}^{\perp} . For any integer $n \geq 1$, we set $\langle n \rangle := \{1, 2, \cdots, n\}$ throughout this paper. As usual, for any finite set A, |A| stands for its cardinality.

In coding theory, the two most common ways to present a linear code are either a generator matrix or a parity-check matrix. A generator matrix for an [n, k] code C is any $k \times n$ matrix G whose rows form a basis of C. An $(n - k) \times n$ matrix H is called a parity-check matrix of an [n, k] code C. Then one has $C = \{\mathbf{x} \in \mathbb{F}_q^n \mid H\mathbf{x}^T = 0\}$. An important invariant of a code is the minimal distance between codewords. Let $\mathbf{a} = (a_1, \dots, a_n)$, $\mathbf{b} = (b_1, \dots, b_n)$ be two codewords in C. Then the Hamming weight $w(\mathbf{a})$ of \mathbf{a} is defined to be the number of nonzero coordinates of \mathbf{a} , that is, $w(\mathbf{a}) = |\{i \in \langle n \rangle \mid a_i \neq 0\}|$. The

²⁰²⁰ Mathematics Subject Classification. 11T71, 94B15, 94B05, 94A60.

Key words and phrases. Cyclic code; Linear code; MDS code; BCH code; AMDS code; NMDS code. *S.F. Hong is the corresponding author and was supported partially by National Science Foundation of China # 12171332.

support set supp(**a**) of $\mathbf{a} \in \mathbb{F}_q^n$ is defined by supp(\mathbf{a}) := $\{i \in \langle n \rangle \mid a_i \neq 0\}$. The Hamming distance $d(\mathbf{a}, \mathbf{b})$ between **a** and **b** is defined to be the number of coordinates in which **a** and **b** differ. In other words, $d(\mathbf{a}, \mathbf{b}) := |\{i \in \langle n \rangle \mid a_i \neq b_i\}|$. Then $d(\mathbf{a}, \mathbf{b}) = w(\mathbf{a} - \mathbf{b})$. The minimum Hamming distance, denoted by $d = d(\mathcal{C})$, of a code $\mathcal{C} \subseteq \mathbb{F}_q^n$ is defined to be the smallest Hamming distance between distinct codewords. Obviously, d is equal to the minimal value of the Hamming weights of all nonzero codewords in \mathcal{C} . Namely, $d = d(\mathcal{C}) := \min_{\mathbf{c}_1, \mathbf{c}_2 \in \mathcal{C}, \mathbf{c}_1 \neq \mathbf{c}_2} \{d(\mathbf{c}_1, \mathbf{c}_2)\}$. In particular, the minimum Hamming distance is important in determining the error-correcting capability of \mathcal{C} . If the minimum Hamming distance d of an [n, k] code is known, then we refer this code to an [n, k, d] code.

For any integer i with $0 \le i \le n$, let A_i stand for the number of codewords in \mathcal{C} with length n and the Hamming weight i. Then the *weight enumerator* of \mathcal{C} is defined to be the polynomial

$$1 + A_1x + A_2x^2 + \dots + A_nx^n.$$

The sequence $(1, A_1, A_2, \dots, A_n)$ is called the *weight distribution* of C. The code C is called a *t*-weight code if $|\{A_i \neq 0 \mid i \in \langle n \rangle\}| = t$. As an important type of linear codes, cyclic codes were first studied by E. Prange in 1957. Not only they have good algebraic structures, but also their encoding and decoding can be easily implemented using the linear shift registers [3, 14, 21, 23]. We say that an [n, k, d] code C over \mathbb{F}_q is cyclic if $(c_0, c_1, \dots, c_{n-1}) \in C$ implies that $(c_{n-1}, c_0, c_1, \dots, c_{n-2}) \in C$.

In the rest of the paper, we impose the restriction gcd(n,q) = 1 and let $(x^n - 1)$ be the ideal $\mathbb{F}_q[x]$ generated by $x^n - 1 \in \mathbb{F}_q[x]$. Then all the elements of the residue class ring $\mathbb{F}_q[x]/(x^n - 1)$ can be represented by polynomials of degree less than n. Clearly, there is an isomorphism φ from \mathbb{F}_q^n as a vector space over \mathbb{F}_q to $\mathbb{F}_q[x]/(x^n - 1)$, defined by

$$\varphi: \mathbb{F}_q^n \to \mathbb{F}_q[x]/(x^n - 1)$$
$$(c_0, c_1, \cdots, c_{n-1}) \mapsto c_0 + c_1 x + \cdots + c_{n-1} x^{n-1}.$$

For convenience, we denote the elements of $\mathbb{F}_q[x]/(x^n-1)$ either as polynomials of degree no more than n-1 modulo x^n-1 or as vectors over \mathbb{F}_q . So we can interpret \mathcal{C} as a subset of $\mathbb{F}_q[x]/(x^n-1)$. A linear code \mathcal{C} is cyclic if and only if \mathcal{C} is an ideal of $\mathbb{F}_q[x]/(x^n-1)$. Note that every ideal of $\mathbb{F}_q[x]/(x^n-1)$ is principal, in particular, every nonzero ideal \mathcal{C} is generated by the monic polynomial q(x) of smallest degree in this ideal.

Let $\mathcal{C} = (g(x))$ be a cyclic code. Then g(x) is called the generator polynomial of \mathcal{C} and $h(x) = (x^n - 1)/g(x)$ is called the parity-check polynomial of \mathcal{C} . Let $m = \operatorname{ord}_n(q)$ be the order of q modulo n, and let α be a generator of $\mathbb{F}_{q^m}^*$. Put $\beta = \alpha \frac{q^{m-1}}{n}$. Then β is a primitive *n*-th root of unity in \mathbb{F}_{q^m} . For any integer i with $0 \le i \le n-1$, let $m_i(x)$ denote the minimal polynomial of β^i over \mathbb{F}_q . Let h and δ be two integers with $0 \le h \le n-1$ and $2 \le \delta \le n$, define

$$g_{(q,n,\delta,h)}(x) := \operatorname{lcm}(m_h(x), m_{h+1}(x), \cdots, m_{h+\delta-2}(x)),$$

where lcm denotes the least common multiple of these minimal polynomials over \mathbb{F}_q , and the addition in the subscript b + i of $m_{b+i}(x)$ means the integer addition modulo n. Let $\mathcal{C}_{(q,n,\delta,h)}$ denote the cyclic code of length n and designed distance δ over \mathbb{F}_q with generator polynomial $g_{(q,n,\delta,h)}(x)$. Then $\mathcal{C}_{(q,n,\delta,h)}$ is called a *BCH code* of length n and designed distance δ . The maximum designed distance of a BCH code is called the *Bose* distance. When h = 1, $\mathcal{C}_{(q,n,\delta,h)}$ is called a narrow-sense BCH code. When $n = q^m - 1$, $\mathcal{C}_{(q,n,\delta,h)}$ is referred to as a primitive BCH code. It is well known that an [n, k, d] code \mathcal{C} is called a maximum distance separable (MDS) code if the minimum distance d reaches

3

the Singleton bound: $d \leq n-k+1$, i.e., if d = n-k+1. The dual of an MDS code is also an MDS code. If d = n - k, then the code is called an *almost MDS code*. For brevity, we write an *AMDS* code for an almost MDS code. Hence AMDS codes have parameters [n, k, n-k]. In general, the dual of an AMDS code may not be an AMDS code. Therefore determining whether the dual of an AMDS code is an AMDS code is an interesting and important problem, see [4], [17], [22] and [25] for some progress about this topic. An AMDS code C is called a *near MDS (NMDS) code* if its dual code is also an AMDS code. MDS codes are widely applied in various fields due to their nice properties, see [14] and [19]. So the study of MDS code, AMDS codes and NMDS code has attracted a lot of attention and made much vital progress, see, for instance, [2], [9], [7], [8], [11], [13] and [15].

BCH codes are a significant class of linear codes that have good algebraic structure, are easy to construct, and can be encoded and decoded relatively easily. Although BCH codes have been studied for decades, their parameters are not easy to determine. In the past 10 years, many mathematicians devoted to this subject and made much vital progress, see [5], [6], [10], [12], [16], [18] and [20]. In 2020, Ding and Tang [6] considered the narrow-sense BCH codes $C_{(q,q+1,3,1)}$ with $q = 3^m$ and its dual code. They showed that the BCH code $C_{(q,q+1,3,1)}$ and the dual code of $C_{(q,q+1,3,1)}$ are NMDS code. In 2022, Geng, Yang, Zhang and Zhou [10] studied a class of AMDS codes from the BCH codes $C_{(q,q+1,3,4)}$ with $q = 3^m$ and determined their parameters.

Theorem 1.1. [10] Let $q = 3^m$ with $m \ge 3$ being odd. Then the BCH code $C_{(q,q+1,3,4)}$ is an AMDS code with parameters [q+1, q-3, 4].

They verified through examples that the code $C_{(q,q+1,3,4)}^{\perp}$ is also an AMDS code, where $q = 3^m$ for m = 3, 5, 7. Based on this observation, they conjectured that if $q = 3^m$ with $m \ge 3$ being odd, then the dual code $C_{(q,q+1,3,4)}^{\perp}$ is an AMDS code with parameters [q+1, 4, q-3].

In the current paper, our main goal is to investigate this conjecture. We will prove that the Geng-Yang-Zhang-Zhou conjecture mentioned above is true as the following theorem shows.

Theorem 1.2. Let $q = 3^m$ with $m \ge 3$ being odd. Then the dual code of the BCH code $C_{(q,q+1,3,4)}$ is an AMDS code with parameters [q+1, 4, q-3].

Combining Theorem 1.1 with Theorem 1.2, we can conclude that the BCH code $C_{(q,q+1,3,4)}$ is a NMDS code. That is, we have the following result.

Theorem 1.3. Let $q = 3^m$ with $m \ge 3$ being odd. Then the BCH code $C_{(q,q+1,3,4)}$ is a NMDS code with parameters [q+1, q-3, 4].

This paper is organized as follows. First of all, we establish in Section 2 a necessary and sufficient condition for the dual code of an almost MDS code to be an almost MDS code. Then in Section 3, we study the BCH code $C_{(q,q+1,3,4)}$ with parameters [q + 1, q - 3, 4] and give the proof of Theorem 1.2.

2. A CHARACTERIZATION FOR THE DUAL CODE OF AN ALMOST MDS CODE TO BE AN ALMOST MDS CODE

In this section, we present a characterization for the dual code of an almost MDS code to be an almost MDS code that is needed in proving Theorem 1.2. We begin with a renowned result due to MacWilliams.

Lemma 2.1. [3, 14, 21, 23] (MacWilliams identity) For a linear [n, k] code C over \mathbb{F}_q , let A_i and A_i^{\perp} be the number of codewords with the Hamming weight i in C and C^{\perp} , respectively. Then for any integer r with $0 \leq r \leq n$, we have

$$\frac{1}{q^k}\sum_{i=0}^{n-r} \binom{n-i}{r} A_i = \frac{1}{q^r}\sum_{i=0}^r \binom{n-i}{n-r} A_i^{\perp}.$$

Lemma 2.2. [24] (Singleton bound) For a linear [n, k, d] code C over \mathbb{F}_q , we have

$$d \le n - k + 1.$$

Lemma 2.3. [23] Let C be a linear [n, k, d] code over \mathbb{F}_q . Then C is an MDS code if and only if the dual code C^{\perp} of C is an MDS code.

Lemma 2.4. Let C be an AMDS code over \mathbb{F}_q with parameters [n, k, n - k]. Then C is a NMDS code if and only if the following identity is true:

$$\frac{1}{q-1}(kA_{n-k} + A_{n-k+1}) = \binom{n}{n-k+1}.$$
(2.1)

Proof. As C is an AMDS code with parameters [n, k, n - k], applying Lemma 2.1 to the case r = k - 1, one derives that

$$\frac{1}{q^k} \sum_{i=0}^{n-k+1} \binom{n-i}{k-1} A_i = \frac{1}{q^{k-1}} \sum_{i=0}^{k-1} \binom{n-i}{n-k+1} A_i^{\perp}.$$

Furthermore, the minimum distance of C equals n - k, and so $A_i = 0$ for any integer i with $1 \le i \le n - k - 1$. It then follows from $A_0 = 1 = A_0^{\perp}$ that

$$\frac{1}{q} \left(\binom{n}{k-1} + \binom{k}{k-1} A_{n-k} + A_{n-k+1} \right) = \binom{n}{n-k+1} + \sum_{i=1}^{k-1} \binom{n-i}{n-k+1} A_i^{\perp}.$$
 (2.2)

First of all, we show the necessity part. To do so, one lets C be a NMDS code. Then C^{\perp} is an AMDS code with $d(C^{\perp}) = k$. This yields that $A_i^{\perp} = 0$ for any integer *i* with $1 \leq i \leq k - 1$. Hence

$$\sum_{i=1}^{k-1} \binom{n-i}{n-k+1} A_i^{\perp} = 0.$$
(2.3)

Then from (2.2), we deduce that

$$\frac{1}{q} \left(\binom{n}{k-1} + kA_{n-k} + A_{n-k+1} \right) = \binom{n}{n-k+1}.$$
(2.4)

Therefore the desired result (2.1) follows immediately. The necessity is proved.

Finally, we show the sufficiency part. Assume that (2.1) is true. Combining (2.2) and (2.4) gives us (2.3). So by (2.3), one can deduce that $A_i^{\perp} = 0$ for any integer *i* with $1 \le i \le k - 1$. This implies that

$$d(\mathcal{C}^{\perp}) \ge k. \tag{2.5}$$

Noticing that C is an AMDS code with parameters [n, k, n - k], so C^{\perp} is a code with parameters $[n, n - k, d(C^{\perp})]$. Since C is not an MDS code, Lemma 2.3 tells us that C^{\perp} is also not an MDS code. Thus from Lemma 2.2, Singleton bound gives us that

$$d(\mathcal{C}^{\perp}) \le n - (n - k) = k. \tag{2.6}$$

It then follows from (2.5) and (2.6), we have $d(\mathcal{C}^{\perp}) = k$. Namely, \mathcal{C}^{\perp} is an AMDS code with parameters [n, n-k, k]. Consequently, \mathcal{C} is a NMDS code as desired. This ends the proof of Lemma 2.4.

Lemma 2.5. Let C be an AMDS code with parameters [n, k, n-k]. Then the Hamming weight of each nonzero codeword \mathbf{c} in C with $\operatorname{supp}(\mathbf{c}) \subseteq \{i_1, \dots, i_{n-k+1}\} \subseteq \langle n \rangle$ is equal to either n-k, or n-k+1.

Proof. Let **c** be a nonzero codeword in C with $\operatorname{supp}(\mathbf{c}) \subseteq \{i_1, \dots, i_{n-k+1}\} \subseteq \langle n \rangle$. On the one hand, since C is an AMDS code with parameters [n, k, n-k], one has $w(\mathbf{c}) \geq d(C) = n-k$. On the other hand, we have $w(\mathbf{c}) = |\operatorname{supp}(\mathbf{c})| \leq |\{i_1, \dots, i_{n-k+1}\}| = n-k+1$. Hence $n-k \leq w(\mathbf{c}) \leq n-k+1$. So the expected result follows immediately.

This completes the proof of Lemma 2.5.

Definition 2.1. Let C be an AMDS code with parameters [n, k, n - k] and let r be an integer with $1 \leq r \leq n$, Then associated with any given subset $\{i_1, \dots, i_r\} \subseteq \langle n \rangle$, we define a subset, denoted by $C(i_1, \dots, i_r)$ or $C(\{i_1, \dots, i_r\})$, of C by

$$\mathcal{C}(i_1,\cdots,i_r):=\{\mathbf{c}\in\mathcal{C}\mid \operatorname{supp}(\mathbf{c})\subseteq\{i_1,\cdots,i_r\}\}.$$

Obviously, for any subset $\{i_1, \dots, i_r\} \subseteq \langle n \rangle$, the set $\mathcal{C}(i_1, \dots, i_r)$ may be empty and forms a vector space under as the usual vector addition and scalar multiplication on \mathbb{F}_q . Now we introduce two subsets of \mathcal{C} as follows:

$$\mathcal{C}_i(i_1,\cdots,i_r) := \{ \mathbf{c} \in \mathcal{C}(i_1,\cdots,i_r) \mid w(\mathbf{c}) = \delta_i \},\$$

where $\delta_i = n - k - 1 + i$ for $i \in \{1, 2\}$.

We have the following result.

Lemma 2.6. Let C be an AMDS code with parameters [n, k, n-k]. Then for any subset $\{i_1, \dots, i_{n-k+1}\} \subseteq \langle n \rangle$, we have the following disjoint union:

 $\mathcal{C}(i_1,\cdots,i_{n-k+1})\setminus\{\mathbf{0}\}=\mathcal{C}_1(i_1,\cdots,i_{n-k+1})\cup\mathcal{C}_2(i_1,\cdots,i_{n-k+1}),$

where $\mathbf{0}$ stands for the zero codeword of \mathcal{C} .

Proof. First of all, it is obvious that $C_1(i_1, \cdots, i_{n-k+1}) \cap C_2(i_1, \cdots, i_{n-k+1}) = \emptyset$.

Now for any $\mathbf{0} \neq \mathbf{c} \in \mathcal{C}(i_1, \dots, i_{n-k+1})$, by Lemma 2.5, one has either $w(\mathbf{c}) = n - k$ or $w(\mathbf{c}) = n - k + 1$. If $w(\mathbf{c}) = n - k$, then $\mathbf{c} \in \mathcal{C}_1(i_1, \dots, i_{n-k+1})$. If $w(\mathbf{c}) = n - k + 1$, then $\mathbf{c} \in \mathcal{C}_2(i_1, \dots, i_{n-k+1})$. By the arbitrariness of \mathbf{c} , one then concludes that

$$\mathcal{C}(i_1,\cdots,i_{n-k+1})\subseteq \mathcal{C}_1(i_1,\cdots,i_{n-k+1})\cup \mathcal{C}_2(i_1,\cdots,i_{n-k+1}).$$

But we have $C_1(i_1, \dots, i_{n-k+1}) \cup C_2(i_1, \dots, i_{n-k+1}) \subseteq C(i_1, \dots, i_{n-k+1})$. So the expected result follows immediately.

This completes the proof of Lemma 2.6.

Lemma 2.7. Let C be an AMDS code over \mathbb{F}_q with parameters [n, k, n-k] and let i be a positive integer no more than n. Then each of the following is true:

- (1) If the *i*-th component of all codewords in C is equal to zero, then $|C(\langle n \rangle \setminus \{i\})| = q^k$ and dim $C(\langle n \rangle \setminus \{i\}) = k$.
- (2) If there exists a codeword in \mathcal{C} with the *i*-th component nonzero, then we have $|\mathcal{C}(\langle n \rangle \setminus \{i\})| = q^{k-1}$ and dim $\mathcal{C}(\langle n \rangle \setminus \{i\}) = k 1$.

Proof. For arbitrary $\alpha \in \mathbb{F}_q$, we define $\mathcal{C}_i(\alpha)$ as a subset of \mathcal{C} by

$$\mathcal{C}_i(\alpha) := \{ \mathbf{c} = (\mathbf{c}_1, \cdots, \mathbf{c}_n) \in \mathcal{C} \mid \mathbf{c}_i = \alpha \}.$$

We claim that $C(\langle n \rangle \setminus \{i\}) = C_i(0)$. In fact, for any $\mathbf{c} \in C(\langle n \rangle \setminus \{i\})$, one has $\mathbf{c}_i = 0$. This implies that $\mathbf{c} \in C_i(0)$. Then $C(\langle n \rangle \setminus \{i\}) \subseteq C_i(0)$. Conversely, for any $\mathbf{c} \in C_i(0)$, we must have $\mathbf{c}_i = 0$. But by Definition 2.1, we have $C(\langle n \rangle \setminus \{i\}) = \{\mathbf{c} \in C \mid \operatorname{supp}(\mathbf{c}) \subseteq \langle n \rangle \setminus \{i\}\}$ which implies that $\operatorname{supp}(\mathbf{c}) \subseteq \langle n \rangle \setminus \{i\}$. Hence $\mathbf{c} \in C(\langle n \rangle \setminus \{i\})$. That is, $C_i(0) \subseteq C(\langle n \rangle \setminus \{i\})$. Thus we have $C(\langle n \rangle \setminus \{i\}) = C_i(0)$ as claimed.

In addition, the subset $C_i(0)$ forms a vector subspace of C under vector addition and scalar multiplication. It then follows that $\dim C(\langle n \rangle \setminus \{i\}) = \dim C_i(0)$.

(1). Let $\mathbf{c}_i = 0$ for any $\mathbf{c} \in \mathcal{C}$. Then $\mathcal{C}_i(0) = \mathcal{C}$. Combining the above claim, one yields that $\mathcal{C}(\langle n \rangle \setminus \{i\}) = \mathcal{C}_i(0) = \mathcal{C}$. Thus $\#\mathcal{C}(\langle n \rangle \setminus \{i\}) = |\mathcal{C}_i(0)| = |\mathcal{C}| = q^k$ and $\dim \mathcal{C}(\langle n \rangle \setminus \{i\}) = \dim \mathcal{C}_i(0) = \dim \mathcal{C} = k$ as required. Part (1) is proved.

(2). Assume that there exists a codeword $\mathbf{c} \in \mathcal{C}$ such that $\mathbf{c}_i \in \mathbb{F}_q^*$. For such \mathbf{c}_i , one writes $\mathbf{c}_i = a$. Namely, $\mathbf{c} \in \mathcal{C}_i(a)$. For any $b \in \mathbb{F}_q^*$, we have $ba^{-1}\mathbf{c} \in \mathcal{C}$ with $(ba^{-1}\mathbf{c})_i = ba^{-1}\mathbf{c}_i = ba^{-1}a = b$. Hence $ba^{-1}\mathbf{c} \in \mathcal{C}_i(b)$. Therefore if we define a map θ from $\mathcal{C}_i(a)$ to $\mathcal{C}_i(b)$ by

$$\theta: \mathcal{C}_i(a) \to \mathcal{C}_i(b),$$
$$\mathbf{c} \mapsto ba^{-1}\mathbf{c},$$

then θ is well defined. First of all, for any $\mathbf{c} \in \mathcal{C}_i(b)$, one has $\mathbf{c}_i = b$. This implies that $(ab^{-1}\mathbf{c})_i = ab^{-1}\mathbf{c}_i = ab^{-1}b = a$. Thus $ab^{-1}\mathbf{c} \in \mathcal{C}_i(a)$. Moreover, we have $\theta(ab^{-1}\mathbf{c}) = ba^{-1} \cdot ab^{-1}\mathbf{c} = \mathbf{c}$. Hence θ is surjective. Consequently, for arbitrary $\mathbf{c}, \mathbf{c}' \in \mathcal{C}_i(b)$ satisfying $\mathbf{c} = \mathbf{c}'$, one can find $\mathbf{\bar{c}}, \mathbf{\bar{c}}' \in \mathcal{C}_i(a)$ such that $\theta(\mathbf{\bar{c}}) = ba^{-1}\mathbf{\bar{c}} = \mathbf{c}$ and $\theta(\mathbf{\bar{c}}') = ba^{-1}\mathbf{\bar{c}}' = \mathbf{c}'$. Since $\mathbf{c} = \mathbf{c}'$, we have $ba^{-1}\mathbf{\bar{c}} = ba^{-1}\mathbf{\bar{c}}'$ which infers that $\mathbf{\bar{c}} = \mathbf{\bar{c}}'$. Hence θ is injective, and so the map θ is bijective. It follows that

$$|\mathcal{C}_i(b)| = |\mathcal{C}_i(a)|, \forall b \in \mathbb{F}_q^*.$$
(2.7)

Consequently, we show that the following is true:

$$|\mathcal{C}_i(0)| = |\mathcal{C}_i(a)|. \tag{2.8}$$

This will be done in what follows. On the one hand, for any $\mathbf{d} = (\mathbf{d}_1, \cdots, \mathbf{d}_n) \in \mathcal{C}_i(a)$, one has $(\mathbf{c} - \mathbf{d})_i = \mathbf{c}_i - \mathbf{d}_i = a - a = 0$ that yields that $\mathbf{c} - \mathbf{d} \in \mathcal{C}_i(0)$. So

$$\{\mathbf{c} - \mathbf{d} \mid \mathbf{d} \in \mathcal{C}_i(a)\} \subseteq \mathcal{C}_i(0).$$

It then follows that

$$\mathcal{C}_{i}(a)| = |\{\mathbf{d} \mid \mathbf{d} \in \mathcal{C}_{i}(a)\}| = |\{\mathbf{c} - \mathbf{d} \mid \mathbf{d} \in \mathcal{C}_{i}(a)\}| \le |\mathcal{C}_{i}(0)|.$$
(2.9)

On the other hand, for any $\mathbf{f} = (\mathbf{f}_1, \cdots, \mathbf{f}_n) \in \mathcal{C}_i(0)$, one has $(\mathbf{f} + \mathbf{c})_i = \mathbf{f}_i + \mathbf{c}_i = 0 + a = a$. Thus $\mathbf{f} + \mathbf{c} \in \mathcal{C}_i(a)$ which implies that

$${\mathbf{f} + \mathbf{c} \mid \mathbf{f} \in \mathcal{C}_i(0)} \subseteq \mathcal{C}_i(a).$$

Therefore

$$|\mathcal{C}_i(0)| = |\{\mathbf{f} \mid \mathbf{f} \in \mathcal{C}_i(0)\}| = |\{\mathbf{f} + \mathbf{c} \mid \mathbf{f} \in \mathcal{C}_i(0)\}| \le |\mathcal{C}_i(a)|.$$
(2.10)

Then the truth of (2.8) follows immediately from (2.9) and (2.10).

Hence, for any two distinct elements b and c in \mathbb{F}_q^* , combining (2.7) and (2.8), we derive that

$$|\mathcal{C}_i(b)| = |\mathcal{C}_i(c)| = |\mathcal{C}_i(0)|$$

7

and

$$\mathcal{C}_i(b) \cap \mathcal{C}_i(c) = \{ \mathbf{c} \in \mathcal{C} \mid \mathbf{c}_i = b \} \cap \{ \mathbf{c} \in \mathcal{C} \mid \mathbf{c}_i = c \} = \emptyset$$

Furthermore, since

$$\mathcal{C} = \mathcal{C}_i(0) \cup (\bigcup_{a \in \mathbb{F}_q^*} \mathcal{C}_i(a)),$$

one deduces that $|\mathcal{C}| = q|\mathcal{C}_i(0)|$. So $|\mathcal{C}_i(0)| = \frac{1}{q}|\mathcal{C}| = q^{k-1}$. Finally, by the claim, one can deduce that $|\mathcal{C}(\langle n \rangle \setminus \{i\})| = |\mathcal{C}_i(0)| = q^{k-1}$ and $\dim \mathcal{C}(\langle n \rangle \setminus \{i\}) = \dim \mathcal{C}_i(0) = k-1$ as expected. Part (2) is proved.

This finishes the proof of Lemma 2.7.

Lemma 2.8. Let C be an AMDS code with parameters [n, k, n - k]. For any subset $\{i_1, \dots, i_{n-k+1}\} \subseteq \langle n \rangle$, we have dim $C(i_1, \dots, i_{n-k+1}) \in \{1, 2\}$.

Proof. For any subset $\{i_1, \cdots, i_{n-k+1}\} \subseteq \langle n \rangle$, we write

$$\{i_{n-k+2},\cdots,i_n\}:=\langle n\rangle\setminus\{i_1,\cdots,i_{n-k+1}\}.$$

Then

$$\{i_1,\cdots,i_{n-k+1}\}=\langle n\rangle\setminus\{i_{n-k+2},\cdots,i_n\}.$$

By Lemma 2.7, we know that

$$\dim \mathcal{C}(i_1, \cdots, i_{n-k+1}) = \dim \mathcal{C}(\langle n \rangle \setminus \{i_{n-k+2}\} \setminus \cdots \setminus \{i_n\}) \ge k - (k-1) = 1.$$

Suppose that there exists subset $\{i_1, \dots, i_{n-k+1}\} \subseteq \langle n \rangle$ such that

$$\dim \mathcal{C}(i_1, \cdots, i_{n-k+1}) > 2.$$

Then by Lemma 2.7, one obtains that

$$\dim \mathcal{C}(i_1, \cdots, i_{n-k-1})$$

= dim($\mathcal{C}(i_1, \cdots, i_{n-k+1}) \setminus \{i_{n-k+1}\}$)\ $\{i_{n-k}\}$)
 \geq dim $\mathcal{C}(i_1, \cdots, i_{n-k+1}) - 2 > 0.$

It follows that there is at least a nonzero codeword $\mathbf{c} \in \mathcal{C}(i_1, \dots, i_{n-k-1})$, we have $\operatorname{supp}(\mathbf{c}) \subseteq \{i_1, \dots, i_{n-k-1}\}$. Hence $w(\mathbf{c}) = |\operatorname{supp}(\mathbf{c})| \leq |\{i_1, \dots, i_{n-k-1}\}| = n - k - 1$. But $w(\mathbf{c}) \geq d(\mathcal{C}) = n - k$. Hence we arrive at

$$n-k-1 \ge w(\mathbf{c}) \ge d(\mathcal{C}) = n-k$$

which is impossible. So $\dim \mathcal{C}(i_1, \cdots, i_{n-k+1}) \leq 2$ for any subset $\{i_1, \cdots, i_{n-k+1}\} \subseteq \langle n \rangle$. It follows that for any subset $\{i_1, \cdots, i_{n-k+1}\} \subseteq \langle n \rangle$, we have $\dim \mathcal{C}(i_1, \cdots, i_{n-k+1}) \in \{1, 2\}$ as desired. Lemma 2.8 is proved. \Box

Lemma 2.9. Let C be an AMDS code with parameters [n, k, n-k]. Let $\delta_i = n-k-1+i$ for $i \in \{1, 2\}$. Let $K_i \subseteq \langle n \rangle$ be an arbitrary subset with $\delta_i \leq |K_i| \leq n$. Then each of the following is true:

(1). We have the disjoint union:

$$\mathcal{C}_i(K_i) = \bigcup_{\substack{I \subseteq K_i \\ |I| = \delta_i}} \mathcal{C}_i(I).$$

(2). We have

$$|\mathcal{C}_i(K_i)| = \sum_{\substack{I \subseteq K_i \\ |I| = \delta_i}} |\mathcal{C}_i(I)|.$$

Proof. Since part (2) is a obvious corollary of part (1), we need just to show part (1). In the following, we show part (1).

Let $i \in \{1, 2\}$ be given. Pick an arbitrary subset $I \subseteq K_i$ with $|I| = \delta_i$ and $C_i(I) \neq \emptyset$. We pick a $\mathbf{c} \in C_i(I)$ with $w(\mathbf{c}) = \delta_i$. Then $\mathbf{c} \in C_i(K_i)$ which implies that $C_i(I) \subseteq C_i(K_i)$. It follows that

$$\bigcup_{\subseteq K_i|I|=\delta_i} \mathcal{C}_i(I) \subseteq \mathcal{C}_i(K_i).$$

Conversely, for any $\mathbf{c} \in \mathcal{C}_i(K_i)$, writing $\operatorname{supp}(\mathbf{c}) = I$, we have

Ι

$$\mathbf{c} \in \mathcal{C}_i(I) \subseteq \bigcup_{\substack{I \subseteq K_i \\ |I| = \delta_i}} \mathcal{C}_i(I).$$

It then follows that

$$\mathcal{C}_i(K_i) \subseteq \bigcup_{I \subseteq K_i |I| = \delta_i} \mathcal{C}_i(I)$$

Thus

$$\mathcal{C}_i(K_i) = \bigcup_{\substack{I \subseteq K_i \\ |I| = \delta_i}} \mathcal{C}_i(I).$$

It remains to show that this union is disjoint that will be done in what follows.

We claim that for $i \in \{1, 2\}$, if $I_1, I_2 \subseteq K$ with $I_1 \neq I_2$ and $\operatorname{supp}(I_1) = \operatorname{supp}(I_2) = \delta_i$, then $\mathcal{C}_i(I_1) \cap \mathcal{C}_i(I_2) = \emptyset$. Picking a $\mathbf{c} \in \mathcal{C}_i(I_1) \cap \mathcal{C}_i(I_2)$ gives that $\operatorname{supp}(\mathbf{c}) \subseteq I_1 \cap I_2$. Since $I_1 \neq I_2$, one deduces that $|\operatorname{supp}(\mathbf{c})| \leq |I_1 \cap I_2| < |I_1| = \delta_i$. This contradicts with the fact $|\operatorname{supp}(\mathbf{c})| = \delta_i$. Hence the proofs of the claim and part (1) are complete. So Lemma 2.9 is proved.

We introduce a concept. We say that a codeword $\mathbf{c} \in \mathcal{C}$ is *monic* if, as a vector, the first nonzero component is the multiplicative identity $1_{\mathbb{F}_q}$ of the finite field \mathbb{F}_q .

Lemma 2.10. Let C be an AMDS code over \mathbb{F}_q with parameters [n, k, n-k]. Then it holds that

$$kA_{n-k} + A_{n-k+1} = (q-1)\binom{n}{n-k+1}$$
(2.11)

if and only if it holds that

$$\dim \mathcal{C}(i_1, \cdots, i_{n-k+1}) = 1, \forall \{i_1, \cdots, i_{n-k+1}\} \subseteq \langle n \rangle.$$
(2.12)

Proof. First, we show the sufficiency part. We assume that (2.12) is true. Now for $i \in \{1, 2\}$, we define three sets F_i , C_i and $E_i(t)$ as follows:

$$F_{i} := \{\{i_{1}, ..., i_{n-k+1}\} \subseteq \langle n \rangle \mid \mathcal{C}_{i}(i_{1}, \cdots, i_{n-k+1}) \neq \emptyset\},\$$
$$C_{1} := \{\mathbf{c} \in \mathcal{C} \mid w(\mathbf{c}) = n-k\}, C_{2} := \{\mathbf{c} \in \mathcal{C} \mid w(\mathbf{c}) = n-k+1\}$$

 $E_i(t) := \{ \mathbf{c} \in C_i \mid \text{ the first nonzero component of codeword } \mathbf{c} \text{ is } t \} \text{ for } t \in \mathbb{F}_q^*.$ Then for $i \in \{1, 2\}, E_i(t) \cap E_i(t') = \emptyset \text{ if } t, t' \in \mathbb{F}_q^* \text{ and } t \neq t'.$ Moreover, $|C_1| = A_{n-k}$ and $|C_2| = A_{n-k+1}$. For brevity, we let $E_i := E_i(1_{\mathbb{F}_q})$ for i = 1, 2.

CLAIM I. We have $|E_i(t)| = |E_i|$ (i = 1, 2) for any $t \in \mathbb{F}_q^*$.

In fact, for any codeword $\mathbf{c_t} \in E_i(t)$, let $(\mathbf{c_t})_{i_1}$ be the first nonzero component of $\mathbf{c_t}$ with $1 \leq i_1 \leq n$. Then $(\mathbf{c_t})_{i_1}^{-1} \mathbf{c_t} \in E_i$, from which one derives $E_i(t) \subseteq E_i$. It then follows that $|E_i(t)| \leq |E_i|$. Conversely, let $\mathbf{c} \in E_i$ be any given element. Since \mathbf{c} is monic, one has $t\mathbf{c} \in E_i(t)$. Thus $E_i \subseteq E_i(t)$, and so $|E_i(t)| \geq |E_i|$ from which one deduces that $|E_i(t)| = |E_i|$ as claimed. So Claim I is proved. For $i \in \{1, 2\}$, since all $E_i(t), t \in \mathbb{F}_q^*$, are disjoint, we have $\bigcup_{t \in \mathbb{F}_q^*} E_i(t) = C_i$. So by Claim I,

$$|C_i| = \sum_{t \in \mathbb{F}_q^*} |E_i(t)| = (q-1)|E_i|.$$

It follows that

$$|E_1| = \frac{1}{q-1}|C_1| = \frac{A_{n-k}}{q-1}$$
(2.13)

and

$$|E_2| = \frac{1}{q-1}|C_2| = \frac{A_{n-k+1}}{q-1}.$$
(2.14)

CLAIM II. We have $|F_1| = k|E_1|$.

In order to prove Claim II, we define the map

$$\vartheta: E_1 \to M_1 := \{ \operatorname{supp}(\mathbf{c}) \mid \mathbf{c} \in E_1 \}$$
$$\mathbf{c} \mapsto \operatorname{supp}(\mathbf{c}).$$

Clearly, it is well defined and is surjective. Now we show it is injective. In fact, let $\operatorname{supp}(\mathbf{c}) = \operatorname{supp}(\mathbf{c}')$, where $\mathbf{c}, \mathbf{c}' \in E_1$. Then $w(\mathbf{c}) = w(\mathbf{c}') = n - k$ and \mathbf{c} and \mathbf{c}' are monic. One may write $\operatorname{supp}(\mathbf{c}) = \operatorname{supp}(\mathbf{c}') = \{j_1, ..., j_{n-k}\} (\subseteq \langle n \rangle)$ with $j_1 < ... < j_{n-k}$. Since \mathbf{c} and \mathbf{c}' are monic, we have $\operatorname{supp}(\mathbf{c} - \mathbf{c}') \subseteq \{j_2, ..., j_{n-k}\}$ and so $w(\mathbf{c} - \mathbf{c}') = |\operatorname{supp}(\mathbf{c} - \mathbf{c}')| \leq n - k - 1$. However, $\mathbf{c} - \mathbf{c}' \in \mathcal{C}$ and \mathcal{C} is an [n, k, n - k] AMDS code. Hence we must have $\mathbf{c} - \mathbf{c}' = \mathbf{0}$, the zero codeword of \mathcal{C} . In other words, $\mathbf{c} = \mathbf{c}'$. Thus ϑ is injective, and so is a bijection. Therefore $|E_1| = |\tilde{M}_1|$.

Now we define a set M_1 by

$$M_1 := \{ \operatorname{supp}(\mathbf{c}) \cup \{ j_{\mathbf{c}} \} \mid \mathbf{c} \in E_1, j_{\mathbf{c}} \in \langle n \rangle \setminus \operatorname{supp}(\mathbf{c}) \}.$$

Then $M_1 = \{\operatorname{supp}(\mathbf{c}) \cup \{j_{\mathbf{c}}\} \mid \mathbf{c} \in \mathcal{C}, w(\mathbf{c}) = n - k, \mathbf{c} \text{ is monic, } j_{\mathbf{c}} \in \langle n \rangle \setminus \operatorname{supp}(\mathbf{c}) \}$. Now let $N := \operatorname{supp}(\mathbf{c}) \cup \{j_{\mathbf{c}}\} = \operatorname{supp}(\mathbf{c}') \cup \{j'_{\mathbf{c}}\}$, where $\mathbf{c}, \mathbf{c}' \in E_1, j_{\mathbf{c}} \in \langle n \rangle \setminus \operatorname{supp}(\mathbf{c})$, $j_{\mathbf{c}'} \in \langle n \rangle \setminus \operatorname{supp}(\mathbf{c}')$. Then #N = n - k + 1, $\operatorname{supp}(\mathbf{c}) \subseteq N$ and $\operatorname{supp}(\mathbf{c}') \subseteq N$. It then follows that $\mathbf{c}, \mathbf{c}' \in \mathcal{C}_1(N) \subseteq \mathcal{C}(N)$. By the condition (2.12), one has $\dim \mathcal{C}(N) = 1$. This implies that $\mathbf{c} = \ell \mathbf{c}'$ with $\ell \in \mathbb{F}_q^*$. Since \mathbf{c} and \mathbf{c}' are monic, one has $\ell = 1$ which implies that $\mathbf{c} = \mathbf{c}'$. Thus $j_{\mathbf{c}} = j_{\mathbf{c}'}$. Hence any two elements in the set M_1 are distinct.

From this and noticing that $\operatorname{supp}(\mathbf{c}) = n - k$ for any $\mathbf{c} \in E_1$, we then derive that

$$|M_1| = |M_1| \cdot |(\langle n \rangle \operatorname{supp}(\mathbf{c}))| = |E_1| \cdot |(\langle n \rangle \operatorname{supp}(\mathbf{c}))| = |E_1| \cdot (n - (n - k)) = k|E_1|.$$
(2.15)

Now we prove that $F_1 = M_1$. On the one hand, for any $\operatorname{supp}(\mathbf{c}) \cup \{j_{\mathbf{c}}\} \in M_1$, we have $\mathbf{c} \in E_1$ and $j_{\mathbf{c}} \in \langle n \rangle \operatorname{supp}(\mathbf{c})$. Thus $|\operatorname{supp}(\mathbf{c})| = n - k$. But $\operatorname{supp}(\mathbf{c}) \subseteq \operatorname{supp}(\mathbf{c}) \cup \{j_{\mathbf{c}}\}$. Hence $\mathbf{c} \in \mathcal{C}_1(\operatorname{supp}(\mathbf{c}) \cup \{j_{\mathbf{c}}\})$, namely, $\mathcal{C}_1(\operatorname{supp}(\mathbf{c}) \cup \{j_{\mathbf{c}}\}) \neq \emptyset$. It the follows from $|\operatorname{supp}(\mathbf{c}) \cup \{j_{\mathbf{c}}\}| = n - k + 1$ that $\operatorname{supp}(\mathbf{c}) \cup \{j_{\mathbf{c}}\} \in F_1$. By the arbitrariness of $\operatorname{supp}(\mathbf{c}) \cup \{j_{\mathbf{c}}\} \in M_1$, one then concludes that $M_1 \subseteq F_1$. It remains to show that $F_1 \subseteq M_1$. Actually, for any $\{i_1, \cdots, i_{n-k+1}\} \in F_1$, we have $\mathcal{C}_1(i_1, \cdots, i_{n-k+1}) \neq \emptyset$. Pick an $\mathbf{e} \in \mathcal{C}_1(i_1, \cdots, i_{n-k+1})$. Then $w(\mathbf{e}) = n - k$ and $\operatorname{supp}(\mathbf{e}) \subseteq \{i_1, \cdots, i_{n-k+1}\}$. Consider the following two cases.

Case 1. $\mathbf{e}_{i_1} = 0$. Then $\mathbf{e}_{i_2} \neq 0$ and $\operatorname{supp}(\mathbf{e}_{i_2}^{-1}\mathbf{e}) = \operatorname{supp}(\mathbf{e}) = \{i_2, \cdots, i_{n-k+1}\}$. So $w(\mathbf{e}_{i_2}^{-1}\mathbf{e}) = |\operatorname{supp}(\mathbf{e}_{i_2}^{-1}\mathbf{e})| = |\operatorname{supp}(\mathbf{e})| = n-k$. It implies that $\mathbf{e}_{i_2}^{-1}\mathbf{e} \in E_1$. At this moment, we have

$$\{i_1, i_2, \cdots, i_{n-k+1}\} = \operatorname{supp}(\mathbf{e}) \cup \{i_1\} = \operatorname{supp}(\mathbf{e}_{i_2}^{-1}\mathbf{e}) \cup \{i_1\} \in M_1$$

So we have $F_1 \subseteq M_1$ as desired.

Case 2. $\mathbf{e}_{i_1} \neq 0$. Likewise, we have $\mathbf{e}_{i_1}^{-1} \mathbf{e} \in E_1$, which infers that $\operatorname{supp}(\mathbf{e}_{i_1}^{-1} \mathbf{e}) = \operatorname{supp}(\mathbf{e}) \subseteq \{i_1, \cdots, i_{n-k+1}\}$. Hence we arrive at

$$\{i_1, \cdots, i_{n-k+1}\} = \operatorname{supp}(\mathbf{e}) \cup (\{i_1, \cdots, i_{n-k+1}\} \setminus \operatorname{supp}(\mathbf{e}))$$
$$= \operatorname{supp}(\mathbf{e}_{i_1}^{-1}\mathbf{e}) \cup (\{i_1, \cdots, i_{n-k+1}\} \setminus \operatorname{supp}(\mathbf{e}_{i_1}^{-1}\mathbf{e})) \in M_1.$$

By the arbitrariness of $\{i_1, \dots, i_{n-k+1}\} \in F_1$, one has $F_1 \subseteq M_1$ as required. One then concludes that $F_1 = M_1$.

Finally, from (2.15) one derives that $|F_1| = |M_1| = k|E_1|$ as claimed. Claim II is proved.

Consequently, combining (2.13) and Claim II, we derive that

$$|F_1| = k|E_1| = \frac{kA_{n-k}}{q-1}.$$
(2.16)

CLAIM III. We have $|F_2| = |E_2|$.

We define a set M_2 by $M_2 := \{ \operatorname{supp}(\mathbf{c}) \mid \mathbf{c} \in E_2 \}$. Then $M_2 = \{ \operatorname{supp}(\mathbf{c}) \mid \mathbf{c} \in \mathcal{C}, w(\mathbf{c}) = n - k + 1, \mathbf{c} \text{ is monic} \}$. Let $N' := \operatorname{supp}(\mathbf{c}) = \operatorname{supp}(\mathbf{c}')$, where $\mathbf{c}, \mathbf{c}' \in E_2$. Then |N'| = n - k + 1. It then follows that $\mathbf{c}, \mathbf{c}' \in \mathcal{C}_2(N') \subseteq \mathcal{C}(N')$. By the condition (2.12), one has dim $\mathcal{C}(N') = 1$. This implies that $\mathbf{c} = \ell \mathbf{c}'$ with $\ell \in \mathbb{F}_q^*$. Since \mathbf{c} and \mathbf{c}' are monic, one has $\ell = 1$ which implies that $\mathbf{c} = \mathbf{c}'$. Thus any two elements in the set M_2 are distinct. It then follows that

$$|M_2| = |E_2|. \tag{2.17}$$

Now we prove that $F_2 = M_2$. On the one hand, take arbitrarily $\operatorname{supp}(\mathbf{e}) \in M_2$. Then we have $\mathbf{e} \in E_2$. It is obvious that $|\operatorname{supp}(\mathbf{e})| = n - k + 1$. Then $\operatorname{supp}(\mathbf{e}) \in F_2$. By the arbitrariness of $\operatorname{supp}(\mathbf{e}) \in M_2$, one concludes that $M_2 \subseteq F_2$. On the other hand, for any $\{i_1, \dots, i_{n-k+1}\} \in F_2$, we have $\mathcal{C}_2(i_1, \dots, i_{n-k+1}) \neq \emptyset$. Then picking an $\mathbf{e}' \in \mathcal{C}_2(i_1, \dots, i_{n-k+1})$ gives us that $\operatorname{supp}(\mathbf{e}') = \{i_1, \dots, i_{n-k+1}\}$ and $w(\mathbf{e}') = n - k + 1$. It follows that $(\mathbf{e}')_{i_1}^{-1} \mathbf{e}' \in E_2$. Therefore

$$\{i_1,\cdots,i_{n-k+1}\} = \operatorname{supp}(\mathbf{e}') = \operatorname{supp}((\mathbf{e}')_{i_1}^{-1}\mathbf{e}') \in M_2.$$

By the arbitrariness of $\{i_1, \dots, i_{n-k+1}\} \in F_2$, one has $F_2 \subseteq M_2$, from which one derives that $F_2 = M_2$ as expected.

We can now deduce from (2.17) that $|F_2| = |M_2| = |E_2|$ as asserted. So Claim III is proved.

Combining (2.14) and Claim III, we can easily deduce that

$$|F_2| = |E_2| = \frac{A_{n-k+1}}{q-1}.$$
(2.18)

Now from (2.16) and (2.18), one yields that

$$|F_1| + |F_2| = \frac{1}{q-1}(kA_{n-k} + A_{n-k+1}).$$
(2.19)

In what follows, we show that $F_1 \cap F_2 = \emptyset$. We assume that $F_1 \cap F_2 \neq \emptyset$. Then we can choose a set $\{i_1, \dots, i_{n-k+1}\} \in F_1 \cap F_2$. So $\{i_1, \dots, i_{n-k+1}\} \in F_1$ and $\{i_1, \dots, i_{n-k+1}\} \in F_2$. This infers that there exist $\mathbf{f}, \mathbf{f}' \in \mathcal{C}$ such that $\mathbf{f} \in \mathcal{C}_1(i_1, \dots, i_{n-k+1})$ and $\mathbf{f}' \in \mathcal{C}_2(i_1, \dots, i_{n-k+1})$. One can deduce that $w(\mathbf{f}) = n - k$ and $w(\mathbf{f}') = n - k + 1$. Moreover, we have $\mathbf{f}, \mathbf{f}' \in \mathcal{C}(i_1, \dots, i_{n-k+1})$. However, the condition (2.12) tells us that $\dim \mathcal{C}(i_1, \dots, i_{n-k+1}) = 1$. Hence we must have $\mathbf{f} = \ell \mathbf{f}'$ with $\ell \in \mathbb{F}_q^*$. This implies that

 $w(\mathbf{f}) = w(\mathbf{f}')$ which is impossible since $w(\mathbf{f}) = n - k$ and $w(\mathbf{f}') = n - k + 1$. Hence we have $F_1 \cap F_2 = \emptyset$ as expected.

Now for any subset $\{i_1, \dots, i_{n-k+1}\} \subseteq \langle n \rangle$, Lemma 2.8 tells us dim $\mathcal{C}(i_1, \dots, i_{n-k+1}) \in \{1, 2\}$ which implies that $\mathcal{C}(i_1, \dots, i_{n-k+1}) \neq \emptyset$. Finally, applying the inclusion-exclusion principle (see, for example, [1]) and Lemma 2.6, we arrive at

$$|F_{1}| + |F_{2}| = |(F_{1} \cup F_{2})| + |(F_{1} \cap F_{2})|$$

$$= |(F_{1} \cup F_{2})|$$

$$= |\{\{i_{1}, \cdots, i_{n-k+1}\} \subseteq \langle n \rangle \mid \mathcal{C}_{1}(i_{1}, \cdots, i_{n-k+1}) \cup \mathcal{C}_{2}(i_{1}, \cdots, i_{n-k+1}) \neq \emptyset\}|$$

$$= |\{\{i_{1}, \cdots, i_{n-k+1}\} \subseteq \langle n \rangle \mid \mathcal{C}(i_{1}, \cdots, i_{n-k+1}) \setminus \{\mathbf{0}\} \neq \emptyset\}|$$

$$= |\{\{i_{1}, \cdots, i_{n-k+1}\} \mid \{i_{1}, \cdots, i_{n-k+1}\} \subseteq \langle n \rangle\}|$$

$$= \binom{n}{n-k+1}.$$
(2.20)

Therefore, by the (2.19) and (2.20), the desired result (2.11) follows immediately. So the sufficiency part is proved.

Now we are in the position to give the proof of the necessity part. Suppose that (2.12) is true. Letting $K_i = \langle n \rangle$ (i = 1, 2) in Lemma 2.9 tells us that

$$kA_{n-k} = k|C_{1}|$$

$$= k|C_{1}(\langle n \rangle)|$$

$$= k \sum_{\substack{I \subseteq \langle n \rangle \\ |I| = \delta_{1} = n-k}} |C_{1}(I)|$$

$$= \sum_{\substack{I \subseteq \langle n \rangle \\ |I| = n-k}} |C_{1}(I)| \sum_{j \in \langle n \rangle \setminus I} 1$$

$$= \sum_{\substack{I \subseteq \langle n \rangle \\ |I| = n-k}} \sum_{j \in \langle n \rangle \setminus I} |C_{1}(I)|$$

$$= \sum_{\substack{I \subseteq \langle n \rangle \\ |J| = n-k+1}} \sum_{j \in J} |C_{1}(J \setminus \{j\})|$$

$$= \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} \sum_{\substack{I \subseteq J \\ I = n-k}} |C_{1}(I)|$$

$$= \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} |C_{1}(J)|$$

$$(2.21)$$

and

$$A_{n-k+1} = |C_2|$$

= $|C_2(\langle n \rangle)|$
= $\sum_{\substack{J \subseteq \langle n \rangle \\ |J| = \delta_2 = n-k+1}} |C_2(J)|.$ (2.22)

It then follows from (2.21) and (2.22) that

$$kA_{n-k} + A_{n-k+1} = \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} |\mathcal{C}_{1}(J)| + \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} |\mathcal{C}_{2}(J)|$$

$$= \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} (|\mathcal{C}_{1}(J)| + |\mathcal{C}_{2}(J)|) + |\mathcal{C}_{1}(J) \cap |\mathcal{C}_{2}(J)|)$$

$$= \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} |\mathcal{C}_{1}(J) \cup \mathcal{C}_{2}(J)| \text{ (since } \mathcal{C}_{1}(J) \cap |\mathcal{C}_{2}(J) = \emptyset)$$

$$= \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} |\mathcal{C}(J) \setminus \{\mathbf{0}\}| \text{ (by Lemma2.6)}$$

$$= \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} (q^{\dim \mathcal{C}(J)} - 1). \quad (2.23)$$

But Lemma 2.8 gives us that for any subset $J \subseteq \langle n \rangle$ with |J| = n - k + 1, one has $\dim \mathcal{C}(J) \in \{1, 2\}$, and hence $q^{\dim \mathcal{C}(J)} - 1 = q - 1$, or $q^2 - 1$. It then follows from (2.23) and (2.11) that

$$kA_{n-k} + A_{n-k+1} = \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} (q^{\dim \mathcal{C}(J)} - 1)$$

$$\geq \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} (q - 1)$$

$$= (q - 1) \sum_{\substack{J \subseteq \langle n \rangle \\ |J| = n-k+1}} 1$$

$$= (q - 1) \binom{n}{n-k+1}$$

$$= kA_{n-k} + A_{n-k+1}.$$

This enables dim $\mathcal{C}(J) = 1$ for any subset $J \subseteq \langle n \rangle$ with |J| = n - k + 1. That is, (2.12) holds as one desires. So the necessity part is proved.

The proof of Lemma 2.10 is complete.

Finally, we can give a characterization for the dual code of an AMDS code to be an AMDS code. In other words, we obtain a characterization for an AMDS code to be a NMDS code. This is the main result of this section.

Theorem 2.1. Let C be an AMDS code over \mathbb{F}_q with parameters [n, k, n-k]. Then the dual code of C is an AMDS code if and only if dim C(I) = 1 holds for any subset $I \subseteq \langle n \rangle$ with |I| = n - k + 1.

Proof. This follows immediately from Lemmas 2.4 and 2.10.

3. On the BCH code $C_{(q,q+1,3,4)}$ with parameters [q+1, q-3, 4] and proof of Theorem 1.2

Now we turn our attention to the BCH code $C_{(q,q+1,3,4)}$. We will explore its properties. Throughout this section, let U_{q+1} denote the cyclic group consisting of all (q+1)-th roots of unity in \mathbb{F}_{q^2} , which is a subgroup of the multiplicative group of \mathbb{F}_{q^2} .

Lemma 3.1. Let $q = p^m$ with p being a prime number and $m \ge 1$ being an integer and let C be the BCH code $C_{(q,q+1,3,4)}$. Then dim $C(i_1, \dots, i_{n-k+1=5}) = 1$ holds for any subset $\{i_1, \dots, i_5\} \subseteq \langle n \rangle$ if and only if for arbitrary two distinct elements $x, y \in U_{q+1} \setminus \{1 := 1_{\mathbb{F}_{q^2}}\} \subseteq \mathbb{F}_{q^2}$, there exists at most one element $z \in U_{q+1} \setminus \{x, y, 1\} \subseteq \mathbb{F}_{q^2}$ such that the following system of homogeneous equations

$$\begin{pmatrix} x^4 & y^4 & z^4 & 1\\ x^5 & y^5 & z^5 & 1 \end{pmatrix} \mathbf{X} = \mathbf{0}$$
(3.1)

has a nonzero solution \mathbf{X} in \mathbb{F}_{q}^{4} .

Proof. At first, we let n = q + 1 and

$$H := \begin{pmatrix} \alpha^4 & (\alpha^4)^2 & \cdots & (\alpha^4)^n = 1\\ \alpha^5 & (\alpha^5)^2 & \cdots & (\alpha^5)^n = 1 \end{pmatrix}$$

with α being a generator of U_{q+1} . Then H is the parity-check matrix of $\mathcal{C}_{(q,q+1,3,4)}$, i.e.,

$$\mathcal{C}_{(q,q+1,3,4)} = \{ \mathbf{c} \in \mathbb{F}_q^{q+1} | H \mathbf{c}^T = \mathbf{0} \}.$$

Writing $h_i = \alpha^i$ for $i \in \langle n \rangle$ gives that

$$H = \begin{pmatrix} h_1^4 & h_2^4 & \cdots & h_n^4 \\ h_1^5 & h_2^5 & \cdots & h_n^5 \end{pmatrix}.$$

We first treat the necessity part. Let $\dim C(i_1, ..., i_{n-k+1=5}) = 1$ hold for all subsets $\{i_1, ..., i_5\} \subseteq \langle n \rangle$. For arbitrary two distinct elements $x, y \in U_{q+1} \setminus \{1 := 1_{\mathbb{F}_{q^2}}\} \subseteq \mathbb{F}_{q^2}$, let $j_1, j_2 \in \langle n \rangle \setminus \{n\}$ be such that $x = h_{j_1}$ and $y = h_{j_2}$. Assume that there are two distinct indexes $i_1, i_2 \in \langle n \rangle \setminus \{j_1, j_2, n\}$ such that the system of homogeneous equations

$$\left(\begin{array}{ccc} h_{j_1}^4 & h_{j_2}^4 & h_{i_1}^4 & 1\\ h_{j_1}^5 & h_{j_2}^5 & h_{i_1}^5 & 1 \end{array}\right) \mathbf{X} = \mathbf{0}$$

has a nonzero solution $\mathbf{X} = (x_1, x_2, x_3, x_4)^T$ in \mathbb{F}_q^4 and the system of homogeneous equations

$$\left(egin{array}{cccc} h_{j_1}^4 & h_{j_2}^4 & h_{i_2}^4 & 1 \ h_{j_1}^5 & h_{j_2}^5 & h_{i_2}^5 & 1 \end{array}
ight) \mathbf{Y} = \mathbf{0}$$

has a nonzero solution $\mathbf{Y} = (y_1, y_2, y_3, y_4)^T$ in \mathbb{F}_q^4 . Then we can pick two nonzero vectors $\mathbf{c}, \mathbf{c}' \in \mathbb{F}_q^n$ such that

$$\mathbf{c}_{j_1} = x_1, \mathbf{c}_{j_2} = x_2, \mathbf{c}_{i_1} = x_3, \mathbf{c}_n = x_4 \text{ and } \mathbf{c}_j = 0 \ \forall j \in \langle n \rangle \setminus \{j_1, j_2, i_1, n\}$$

and

$$\mathbf{c}'_{j_1} = y_1, \mathbf{c}'_{j_2} = y_2, \mathbf{c}'_{i_2} = y_3, \mathbf{c}'_n = y_4 \text{ and } \mathbf{c}_j = 0 \ \forall j \in \langle n \rangle \setminus \{j_1, j_2, i_2, n\},\$$

respectively. So we have $\operatorname{supp}(\mathbf{c}) \subseteq \{j_1, j_2, i_1, n\} \subseteq \{j_1, j_2, i_1, i_2, n\} \subseteq \langle n \rangle$ and $\operatorname{supp}(\mathbf{c}') \subseteq \{j_1, j_2, i_2, n\} \subseteq \langle j_1, j_2, i_1, i_2, n\} \subseteq \langle n \rangle$. Letting $H_i = \begin{pmatrix} h_i^4 \\ h_i^5 \end{pmatrix}$ as the *i*-th column vector

of H for $i \in \langle n \rangle$ then gives us that

$$\begin{aligned} H\mathbf{c}^{T} &= \sum_{i=1}^{n} \mathbf{c}_{i} H_{i} \\ &= \sum_{i \in \langle n \rangle \setminus \{j_{1}, j_{2}, i_{1}, n\}} \mathbf{c}_{i} H_{i} + \mathbf{c}_{j_{1}} H_{j_{1}} + \mathbf{c}_{j_{2}} H_{j_{2}} + \mathbf{c}_{i_{1}} H_{i_{1}} + \mathbf{c}_{n} H_{n} \\ &= \mathbf{0} + \mathbf{c}_{j_{1}} H_{j_{1}} + \mathbf{c}_{j_{2}} H_{j_{2}} + \mathbf{c}_{i_{1}} H_{i_{1}} + \mathbf{c}_{n} H_{n} \\ &= x_{1} H_{j_{1}} + x_{2} H_{j_{2}} + x_{3} H_{i_{1}} + x_{4} H_{n} \\ &= \left(\begin{array}{c} h_{j_{1}}^{4} & h_{j_{2}}^{4} & h_{i_{1}}^{4} & 1 \\ h_{j_{1}}^{5} & h_{j_{2}}^{5} & h_{i_{1}}^{5} & 1 \end{array} \right) (x_{1}, x_{2}, x_{3}, x_{4})^{T} \\ &= \mathbf{0} \end{aligned}$$

and

$$\begin{aligned} H(\mathbf{c}')^T &= \sum_{i=1}^n \mathbf{c}'_i H_i \\ &= \sum_{i \in \langle n \rangle \setminus \{j_1, j_2, i_2, n\}} \mathbf{c}'_i H_i + \mathbf{c}'_{j_1} H_{j_1} + \mathbf{c}'_{j_2} H_{j_2} + \mathbf{c}'_{i_2} H_{i_2} + \mathbf{c}'_n H_n \\ &= \mathbf{0} + \mathbf{c}'_{j_1} H_{j_1} + \mathbf{c}'_{j_2} H_{j_2} + \mathbf{c}'_{i_2} H_{i_2} + \mathbf{c}'_n H_n \\ &= y_1 H_{j_1} + y_2 H_{j_2} + y_3 H_{i_2} + y_4 H_n \\ &= \left(\begin{array}{cc} h_{j_1}^4 & h_{j_2}^4 & h_{i_2}^4 & 1 \\ h_{j_1}^5 & h_{j_2}^5 & h_{i_2}^5 & 1 \end{array} \right) (y_1, y_2, y_3, y_4)^T \\ &= \mathbf{0}. \end{aligned}$$

This implies that $\mathbf{c}, \mathbf{c}' \in \mathcal{C} = \mathcal{C}_{(q,q+1,3,4)}$. Thus $w(\mathbf{c}) = |\operatorname{supp}(\mathbf{c})| \geq d(\mathcal{C}) = 4$ and $w(\mathbf{c}') = |\operatorname{supp}(\mathbf{c}')| \geq d(\mathcal{C}) = 4$. But $|\operatorname{supp}(\mathbf{c})| \leq 4$ and $|\operatorname{supp}(\mathbf{c}')| \leq 4$. Hence we must have $\operatorname{supp}(\mathbf{c}) = \{j_1, j_2, i_1, n\} \subseteq \{j_1, j_2, i_1, i_2, n\}$ and $\operatorname{supp}(\mathbf{c}') = \{j_1, j_2, i_2, n\} \subseteq \{j_1, j_2, i_1, i_2, n\}$. One then deduces that $\mathbf{c}, \mathbf{c}' \in \mathcal{C}(j_1, j_2, i_1, i_2, n)$. It then follows from $\operatorname{supp}(\mathbf{c}) \neq \operatorname{supp}(\mathbf{c}')$ that \mathbf{c} and \mathbf{c}' are linear independent, from which one derives that $\dim \mathcal{C}(j_1, j_2, i_1, i_2, n) \geq 2$. This contradicts with the hypothesis $\dim \mathcal{C}(j_1, j_2, i_1, i_2, n) = 1$. Hence for arbitrary two distinct elements $x, y \in U_{q+1} \setminus \{1 := 1_{\mathbb{F}_{q^2}}\} \subseteq \mathbb{F}_{q^2}$, there exists at most one element $z \in U_{q+1} \setminus \{x, y, 1\} \subseteq \mathbb{F}_{q^2}$ such that the system (3.1) of homogeneous equations has a nonzero solution \mathbf{X} in \mathbb{F}_q^4 . The necessity part is proved.

Consequently, we show the sufficiency part. First of all, Lemma 2.8 tells us that $\dim \mathcal{C}(i_1, ..., i_5) \in \{1, 2\}$ for all subsets $\{i_1, \cdots, i_5\} \subseteq \langle n \rangle$. We assume that there exists a subset $\{j_1, j_2, j_3, i_1, i_2\} \subseteq \langle n \rangle$ such that $\dim \mathcal{C}(j_1, j_2, j_3, i_1, i_2) = 2$. Then there exist two linearly independent nonzero codewords $\mathbf{c}_1, \mathbf{c}_2 \in \mathcal{C}(j_1, j_2, j_3, i_1, i_2)$. Thus $w(\mathbf{c}_1), w(\mathbf{c}_2) \in \{4, 5\}$. We claim that there are two linearly independent nonzero codewords \mathbf{c}_3 and \mathbf{c}_4 in $\mathcal{C}(j_1, j_2, j_3, i_1, i_2)$ with weight 4. If $w(\mathbf{c}_1) = 4$ and $w(\mathbf{c}_2) = 4$, then letting $\mathbf{c}_3 := \mathbf{c}_1 \in \mathcal{C}(j_1, j_2, j_3, i_1, i_2)$ and $\mathbf{c}_4 := \mathbf{c}_2 \in \mathcal{C}(j_1, j_2, j_3, i_1, i_2)$ with $w(\mathbf{c}_3) = w(\mathbf{c}_4) = 4$. When $w(\mathbf{c}_1) = 5$ or $w(\mathbf{c}_2) = 5$, in order to finish the proof of the claim, one needs only to consider the following three cases.

Case 1. $w(\mathbf{c}_1) = w(\mathbf{c}_2) = 5$. Then $\operatorname{supp}(\mathbf{c}_1) = \operatorname{supp}(\mathbf{c}_2) = \{j_1, j_2, j_3, i_1, i_2\}$. Obviously, $(\mathbf{c}_1)_{i_1} \neq 0$ and $(\mathbf{c}_2)_{i_1} \neq 0$. Let $\mathbf{c}_3 = \mathbf{c}_1 - ((\mathbf{c}_1)_{i_1}(\mathbf{c}_2)_{i_1}^{-1})\mathbf{c}_2 \in \mathcal{C}(j_1, j_2, j_3, i_1, i_2)$.

Thus

$$\begin{aligned} (\mathbf{c}_3)_{i_1} &= (\mathbf{c}_1 - ((\mathbf{c}_1)_{i_1} (\mathbf{c}_2)_{i_1}^{-1}) \mathbf{c}_2)_{i_1} \\ &= (\mathbf{c}_1)_{i_1} - (((\mathbf{c}_1)_{i_1} (\mathbf{c}_2)_{i_1}^{-1}) \mathbf{c}_2)_{i_1} \\ &= (\mathbf{c}_1)_{i_1} - (\mathbf{c}_1)_{i_1} (\mathbf{c}_2)_{i_1}^{-1} (\mathbf{c}_2)_{i_1} \\ &= 0. \end{aligned}$$

It implies that $\operatorname{supp}(\mathbf{c}_3) \subseteq \{j_1, j_2, j_3, i_2\}$ and so we have $w(\mathbf{c}_3) = \operatorname{supp}(\mathbf{c}_3) \leq 4$. Since \mathbf{c}_1 and \mathbf{c}_2 are linearly independent nonzero codewords, \mathbf{c}_3 must be a nonzero codeword. This implies that $4 = d(\mathcal{C}) \leq w(\mathbf{c}_3) \leq 4$. Hence we find a codeword $\mathbf{c}_3 \in \mathcal{C}(j_1, j_2, j_3, i_1, i_2)$ with $w(\mathbf{c}_3) = 4$. Likewise, since $(\mathbf{c}_2)_{i_2} \neq 0$, letting $\mathbf{c}_4 := \mathbf{c}_2 - ((\mathbf{c}_2)_{i_2}(\mathbf{c}_1)_{i_2}^{-1})\mathbf{c}_1$ gives us a codeword $\mathbf{c}_4 \in \mathcal{C}(j_1, j_2, j_3, i_1, i_2)$ with $\operatorname{supp}(\mathbf{c}_4) = \{j_1, j_2, j_3, i_1\}$ and $w(\mathbf{c}_4) = 4$. Moreover, $\operatorname{supp}(\mathbf{c}_3) = \{j_1, j_2, j_3, i_2\} \neq \{j_1, j_2, j_3, i_1\} = \operatorname{supp}(\mathbf{c}_4)$ which implies that \mathbf{c}_3 and \mathbf{c}_4 are linearly independent nonzero codewords. Therefore we obtain two linearly independent nonzero codewords $\mathbf{c}_3, \mathbf{c}_4 \in \mathcal{C}(j_1, j_2, j_3, i_1, i_2)$ with $w(\mathbf{c}_3) = w(\mathbf{c}_4) = 4$ as desired. The claim is proved in this case.

Case 2. $w(\mathbf{c}_1) = 5$ and $w(\mathbf{c}_2) = 4$. Then $\operatorname{supp}(\mathbf{c}_1) = \{j_1, j_2, j_3, i_1, i_2\}$ and $\operatorname{supp}(\mathbf{c}_2) \subseteq \{j_1, j_2, j_3, i_1, i_2\}$. Evidently, $(\mathbf{c}_1)_{i_1} \neq 0$ and we may let $(\mathbf{c}_2)_{i_2} \neq 0$. Picking $\mathbf{c}_3 = \mathbf{c}_1 - ((\mathbf{c}_1)_{i_2}(\mathbf{c}_2)_{i_2}^{-1})\mathbf{c}_2$ gives that $\operatorname{supp}(\mathbf{c}_3) \subseteq \{j_1, j_2, j_3, i_1, i_2\}$ and

$$\begin{aligned} (\mathbf{c}_3)_{i_2} = & (\mathbf{c}_1 - ((\mathbf{c}_1)_{i_2}(\mathbf{c}_2)_{i_2}^{-1})\mathbf{c}_2)_{i_2} \\ = & (\mathbf{c}_1)_{i_2} - ((\mathbf{c}_1)_{i_2}(\mathbf{c}_2)_{i_2}^{-1})\mathbf{c}_2)_{i_2} \\ = & (\mathbf{c}_1)_{i_2} - (\mathbf{c}_1)_{i_2}(\mathbf{c}_2)_{i_2}^{-1}(\mathbf{c}_2)_{i_2} \\ - & 0 \end{aligned}$$

Hence supp $(\mathbf{c}_3) \subseteq \{j_1, j_2, j_3, i_1\}$. As \mathbf{c}_1 and \mathbf{c}_2 are linearly independent nonzero codewords, \mathbf{c}_3 must be a nonzero codeword. This implies that $4 = d(\mathcal{C}) \leq w(\mathbf{c}_3) = |\operatorname{supp}(\mathbf{c}_3)| \leq 4$. So one gets a codeword $\mathbf{c}_3 \in \mathcal{C}(j_1, j_2, j_3, i_1, i_2)$ with $w(\mathbf{c}_3) = 4$. Hence we find two linearly independent nonzero codewords \mathbf{c}_3 and $\mathbf{c}_4 = \mathbf{c}_2$ in $\mathcal{C}(j_1, j_2, j_3, i_1, i_2)$ with $w(\mathbf{c}_3) = w(\mathbf{c}_4) = 4$. The claim is true in this case.

Case 3. $w(\mathbf{c}_1) = 4$ and $w(\mathbf{c}_2) = 5$. Then by the symmetry, letting $\mathbf{c}_4 = \mathbf{c}_2 - ((\mathbf{c}_2)_{i_2}(\mathbf{c}_2)_{i_2}^{-1})\mathbf{c}_1$ gives us two linearly independent nonzero codewords $\mathbf{c}_3 := \mathbf{c}_1$ and \mathbf{c}_4 in $\mathcal{C}(j_1, j_2, j_3, i_1, i_2)$ with $w(\mathbf{c}_3) = w(\mathbf{c}_4) = 4$. Hence the claim is proved in this case.

By the above claim, we always have two linearly independent nonzero codewords \mathbf{c}_3 and \mathbf{c}_4 in $\mathcal{C}(j_1, j_2, j_3, i_1, i_2)$ with $w(\mathbf{c}_3) = w(\mathbf{c}_4) = 4$. Suppose that $\operatorname{supp}(\mathbf{c}_3) = \operatorname{supp}(\mathbf{c}_4)$. One may write $\operatorname{supp}(\mathbf{c}_3) = \operatorname{supp}(\mathbf{c}_4) = \{j_1, j_2, j_3, i_1\}$. Let $\mathbf{c}_5 := \mathbf{c}_3 - ((\mathbf{c}_3)_{i_1}(\mathbf{c}_4)_{i_1}^{-1})\mathbf{c}_4$. Then $(\mathbf{c}_5)_{i_1} = 0$ and so $\operatorname{supp}(\mathbf{c}_5) \subseteq \{j_1, j_2, j_3\}$. Since \mathbf{c}_3 and \mathbf{c}_4 are linearly independent in $\mathcal{C}(j_1, j_2, j_3, i_1, i_2)$, it follows that \mathbf{c}_5 is a nonzero codeword. So $w(\mathbf{c}_5) \leq 3$ which contradicts with the fact $w(\mathbf{c}_5) \geq d(\mathcal{C}) = 4$. Thus we must have $\operatorname{supp}(\mathbf{c}_3) \neq \operatorname{supp}(\mathbf{c}_4)$. However, $w(\mathbf{c}_3) = w(\mathbf{c}_4) = 4$. Hence $|(\operatorname{supp}(\mathbf{c}_3) \cap \operatorname{supp}(\mathbf{c}_4)| \leq 3$.

On the other hand, since $\mathbf{c}_3, \mathbf{c}_4 \in \mathcal{C}(j_1, j_2, j_3, i_1, i_2)$, we have $\operatorname{supp}(\mathbf{c}_3) \cup \operatorname{supp}(\mathbf{c}_4) \subseteq \{j_1, j_2, j_3, i_1, i_2\}$. Thus $|\operatorname{supp}(\mathbf{c}_3) \cup \operatorname{supp}(\mathbf{c}_4)| \leq 5$. It then follows from the inclusion-exclusion principle that

$$|\operatorname{supp}(\mathbf{c}_3) \cap \operatorname{supp}(\mathbf{c}_4)| = |\operatorname{supp}(\mathbf{c}_3)| + |\operatorname{supp}(\mathbf{c}_4)| - |\operatorname{supp}(\mathbf{c}_3) \cup \operatorname{supp}(\mathbf{c}_4)|$$
$$\geq 4 + 4 - 5 = 3.$$

This concludes that $|\text{supp}(\mathbf{c}_3) \cap \text{supp}(\mathbf{c}_4)| = 3$. One may write $\text{supp}(\mathbf{c}_3) = \{j_1, j_2, j_3, i_1\}$ and $\text{supp}(\mathbf{c}_4) = \{j_1, j_2, j_3, i_2\}$. Then $\mathbf{X} := ((\mathbf{c}_3)_{j_1}, (\mathbf{c}_3)_{j_2}, (\mathbf{c}_3)_{j_3}, (\mathbf{c}_3)_{i_1})^T$ is a solution of the following system of homogeneous equations

$$\begin{pmatrix} h_{j_1}^4 & h_{j_2}^4 & h_{j_3}^4 & h_{i_1}^4 \\ h_{j_1}^5 & h_{j_2}^5 & h_{j_3}^5 & h_{i_1}^5 \end{pmatrix} \mathbf{X} = \mathbf{0}$$

and $\mathbf{Y} := ((\mathbf{c}_4)_{j_1}, (\mathbf{c}_4)_{j_2}, (\mathbf{c}_4)_{j_3}, (\mathbf{c}_4)_{i_2})^T$ is a solution of the system of homogeneous equations

$$\left(\begin{array}{ccc} h_{j_1}^4 & h_{j_2}^4 & h_{j_3}^4 & h_{i_2}^4 \\ h_{j_1}^5 & h_{j_2}^5 & h_{j_3}^5 & h_{i_2}^5 \end{array}\right) \mathbf{Y} = \mathbf{0}.$$

It then follows that

$$\begin{pmatrix} (h_{j_1}h_{j_3}^{-1})^4 & (h_{j_2}h_{j_3}^{-1})^4 & 1 & (h_{i_1}h_{j_3}^{-1})^4 \\ (h_{j_1}h_{j_3}^{-1})^5 & (h_{j_2}h_{j_3}^{-1})^5 & 1 & (h_{i_1}h_{j_3}^{-1})^5 \end{pmatrix} \mathbf{X} = \begin{pmatrix} 0 \times h_{j_3}^{-4} \\ 0 \times h_{j_3}^{-5} \end{pmatrix} = \mathbf{0}$$

and

$$\begin{pmatrix} (h_{j_1}h_{j_3}^{-1})^4 & (h_{j_2}h_{j_3}^{-1})^4 & 1 & (h_{i_2}h_{j_3}^{-1})^4 \\ (h_{j_1}h_{j_3}^{-1})^5 & (h_{j_2}h_{j_3}^{-1})^5 & 1 & (h_{i_2}h_{j_3}^{-1})^5 \end{pmatrix} \mathbf{Y} = \begin{pmatrix} 0 \times h_{j_3}^{-4} \\ 0 \times h_{j_3}^{-5} \end{pmatrix} = \mathbf{0}.$$

Let $x_0 = h_{j_1} h_{j_3}^{-1}, y_0 = h_{j_2} h_{j_3}^{-1}, z_0 = h_{i_1} h_{j_3}^{-1}$ and $z'_0 = h_{i_2} h_{j_3}^{-1}$. Then $x_0, y_0, z_0, z'_0 \in \mathbb{R}$ U_{q+1} are pairwise distinct, and both of the systems of homogeneous equations

$$\begin{pmatrix} x_0^4 & y_0^4 & z_0^4 & 1\\ x_0^5 & y_0^5 & z_0^5 & 1 \end{pmatrix} \mathbf{X} = \mathbf{0}$$
$$\begin{pmatrix} x_0^4 & y_0^4 & z_0'^4 & 1\\ x^5 & y_0^5 & z'^5 & 1 \end{pmatrix} \mathbf{Y} = \mathbf{0}$$

and

 $\begin{pmatrix} x_0^3 & y_0^3 & z_0'^3 & 1 \end{pmatrix}$ have nonzero solutions in \mathbb{F}_q^4 , which contradicts with the hypothesis. Thus the assumption is not true. Hence we must have $\dim \mathcal{C}(j_1, j_2, j_3, i_1, i_2) = 1$ for any subset $\{j_1, j_2, j_3, i_1, i_2\} \subseteq \langle n \rangle$. The sufficiency part is proved.

This concludes the proof of Lemma 3.1.

By the theory of the system of homogeneous equations over a field, one knows that for arbitrary elements $x, y, z \in U_{q+1} \setminus \{1\}$, the system (3.1) of homogeneous equations always has a nonzero solution X in $\mathbb{F}_{q^2}^4$ when the characteristics of the field is equal to 3. But it does not guarantee that the system (3.1) of homogeneous equations holds a nonzero solution X in $\mathbb{F}_{q^2}^4$.

The following result is known and is given in [10].

Lemma 3.2. [10] Let $q = 3^m$ with m being an odd integer. Let U_{q+1} denote the group consisting of all (q+1)-th roots of unity in \mathbb{F}_{q^2} . If x, y and z are three pairwise distinct elements in U_{q+1} , then

$$\det \begin{pmatrix} x^4 & y^4 & z^4 \\ x^5 & y^5 & z^5 \\ x^{-5} & y^{-5} & z^{-5} \end{pmatrix} \neq 0.$$

Lemma 3.3. Let $q = 3^m$ with m being an odd integer. Let $x, y, z \in U_{q+1} \setminus \{1 := 1_{\mathbb{F}_{q^2}}\} \subseteq$ \mathbb{F}_{q^2} be arbitrary three pairwise distinct elements. Then each of the following is true:

(i). The system (3.1) of homogeneous equations has a nonzero solution in \mathbb{F}_q^4 if and only if the following system of homogeneous equations

$$\begin{pmatrix} x^{4} & y^{4} & z^{4} & 1\\ x^{5} & y^{5} & z^{5} & 1\\ x^{-5} & y^{-5} & z^{-5} & 1\\ x^{-4} & y^{-4} & z^{-4} & 1 \end{pmatrix} \mathbf{X} = \mathbf{0}$$
(3.2)

has a nonzero solution in \mathbb{F}_{q}^{4} .

(ii). The system (3.2) has a nonzero solution in \mathbb{F}_q^4 if and only if the following holds in \mathbb{F}_{q^2} :

$$\det \begin{pmatrix} x^4 & y^4 & z^4 & 1\\ x^5 & y^5 & z^5 & 1\\ x^{-5} & y^{-5} & z^{-5} & 1\\ x^{-4} & y^{-4} & z^{-4} & 1 \end{pmatrix} = 0.$$
(3.3)

Proof. (i). We begin with the proof of the necessity. Let the system (3.1) of the homogeneous equations hold a nonzero solution $(a, b, c, d) \in \mathbb{F}_q^4$. Then for $i \in \{4, 5\}$, we have

$$ax^i + by^i + cz^i + d = 0.$$

Since $x, y, z \in U_{q+1} \setminus \{1\}$ and $a, b, c, d \in \mathbb{F}_q$, we have $x^{q+1} = y^{q+1} = z^{q+1} = 1$ and $a^q = a, b^q = b, c^q = c$ and $d^q = d$. It follows that $x^{-1} = x^q, y^{-1} = y^q, z^{-1} = z^q$. Hence for $i \in \{4, 5\}$, we have

$$ax^{-i} + by^{-i} + cz^{-i} + d = (ax^{i} + by^{i} + cz^{i} + d)^{q} = 0^{q} = 0.$$

It then follows that $(a, b, c, d) \in \mathbb{F}_q^4$ is also a nonzero solution in \mathbb{F}_q^4 of the system (3.2) of homogeneous equations. So the necessity part is proved.

Consequently, we show the sufficiency part. Let the system (3.2) of homogeneous equations have a nonzero solution $(a, b, c, d) \in \mathbb{F}_q^4 \subseteq \mathbb{F}_{q^2}^4$. Then for $i \in \{4, 5\}$, we have $ax^i + by^i + cz^i + d = 0$. In other words, $(a, b, c, d) \in \mathbb{F}_q^4$ is a nonzero solution of the system (3.1) of homogeneous equations. Therefore the sufficiency part is proved. So part (i) is proved.

(ii). First, we show the necessity part. Let the system (3.2) of homogeneous equations have a nonzero solution in $\mathbb{F}_q^4 \subseteq \mathbb{F}_{q^2}^4$. Then by the theory of linear algebra over finite fields, one knows that (3.3) is true over the finite field \mathbb{F}_{q^2} . Hence the necessity is proved.

Conversely, we consider the sufficiency part. Let (3.3) hold over the finite field \mathbb{F}_{q^2} . Then the system (3.2) of homogeneous equations has a nonzero solution $(a, b, c, d) \in \mathbb{F}_{q^2}^4$. So for $i \in \{4, 5\}$, we have

 $ax^i + by^i + cz^i + d = 0$

and

$$ax^{-i} + by^{-i} + cz^{-i} + d = 0.$$

Since $(a, b, c, d) \in \mathbb{F}_{q^2}^4$ is nonzero, at least one of a, b, c and d is a nonzero element in \mathbb{F}_{q^2} . WLOG, one may let $a \neq 0$. Then for $i \in \{4, 5\}$, one has

$$x^i + \frac{b}{a}y^i + \frac{c}{a}z^i + \frac{d}{a} = 0$$

and

$$x^{-i} + \frac{b}{a}y^{-i} + \frac{c}{a}z^{-i} + \frac{d}{a} = 0.$$

Write $\bar{b} = \frac{b}{a}, \bar{c} = \frac{c}{a}$ and $\bar{d} = \frac{d}{a}$. Then $(1, \bar{b}, \bar{c}, \bar{d}) \in \mathbb{F}_{q^2}^4$ is solution of (3.2).

From the condition that $x, y, z \in U_{q+1} \setminus \{1\}$, one derives that $x = x^{-q}, y = y^{-q}$ and $z = z^{-q}$. It then follows that for $i \in \{4, 5\}$, we have

$$x^{i} + \bar{b}^{q}y^{i} + \bar{c}^{q}z^{i} + \bar{d}^{q} = x^{-qi} + \bar{b}^{q}y^{-qi} + \bar{c}^{q}z^{-qi} + \bar{d}^{q} = (x^{-i} + \bar{b}y^{-i} + \bar{c}z^{-i} + \bar{d})^{q} = 0^{q} = 0$$
 and

$$x^{-i} + \bar{b}^q y^{-i} + \bar{c}^q z^{-i} + \bar{d}^q = x^{qi} + \bar{b}^q y^{qi} + \bar{c}^q z^{qi} + \bar{d}^q = (x^i + \bar{b}y^i + \bar{c}z^i + \bar{d})^q = 0^q = 0$$

This implies that $(1, \bar{b}^q, \bar{c}^q, \bar{d}^q) \in \mathbb{F}_{q^2}^4$ is a nonzero solution of (3.2).

On the other hand, since (3.3) is true, we have

$$\operatorname{rank} \begin{pmatrix} x^{4} & y^{4} & z^{4} & 1\\ x^{5} & y^{5} & z^{5} & 1\\ x^{-5} & y^{-5} & z^{-5} & 1\\ x^{-4} & y^{-4} & z^{-4} & 1 \end{pmatrix} \leq 3.$$
(3.4)

But Lemma 3.2 tells us that

$$\operatorname{rank}\begin{pmatrix} x^4 & y^4 & z^4\\ x^5 & y^5 & z^5\\ x^{-5} & y^{-5} & z^{-5} \end{pmatrix} = 3$$

Then

$$\operatorname{rank}\begin{pmatrix} x^{4} & y^{4} & z^{4} & 1\\ x^{5} & y^{5} & z^{5} & 1\\ x^{-5} & y^{-5} & z^{-5} & 1\\ x^{-4} & y^{-4} & z^{-4} & 1 \end{pmatrix} \ge \operatorname{rank}\begin{pmatrix} x^{4} & y^{4} & z^{4}\\ x^{5} & y^{5} & z^{5}\\ x^{-5} & y^{-5} & z^{-5} \end{pmatrix} = 3.$$
(3.5)

Combining (3.4) and (3.5), one can conclude that

$$\operatorname{rank} \begin{pmatrix} x^4 & y^4 & z^4 & 1\\ x^5 & y^5 & z^5 & 1\\ x^{-5} & y^{-5} & z^{-5} & 1\\ x^{-4} & y^{-4} & z^{-4} & 1 \end{pmatrix} = 3.$$

It then follows that the dimension of the solution space over \mathbb{F}_{q^2} of the system (3.2) of homogeneous equations is equal to $4 - \operatorname{rank}(A) = 4 - 3 = 1$. This infers that the two solutions $(1, \bar{b}, \bar{c}, \bar{d})^T$ and $(1, \bar{b}^q, \bar{c}^q, \bar{d}^q)^T$ are linearly dependent which implies that there is a nonzero element $l \in \mathbb{F}_{q^2}^*$ such that $(1, \bar{b}, \bar{c}, \bar{d})^T = l(1, \bar{b}^q, \bar{c}^q, \bar{d}^q)^T$. It follows that l = 1and so we have $\bar{b} = \bar{b}^q, \bar{c} = \bar{c}^q$ and $\bar{d} = \bar{d}^q$. This implies that each of \bar{b}, \bar{c} and \bar{d} belongs to \mathbb{F}_q . Hence $(1, \bar{b}, \bar{c}, \bar{d})$ is a nonzero solution in \mathbb{F}_q^4 of (3.2). The sufficiency part is proved.

This completes the proof of Lemma 3.3.

Lemma 3.4. Let $q = 3^m$ with m being a positive integer. Let $x, y, z \in U_{q+1} \setminus \{1\}$ be three pairwise distinct elements. Then

$$\det \begin{pmatrix} x^4 & y^4 & z^4 & 1\\ x^5 & y^5 & z^5 & 1\\ x^{-5} & y^{-5} & z^{-5} & 1\\ x^{-4} & y^{-4} & z^{-4} & 1 \end{pmatrix} = \frac{XYZ}{((X+1)(Y+1)(Z+1))^5} \det \begin{pmatrix} Y^9 - X^9 & Z^9 - X^9\\ Y^8 - X^8 & Z^8 - X^8 \end{pmatrix},$$

where X := x - 1, Y := y - 1 and Z := z - 1. Moreover, it holds in \mathbb{F}_{q^2} that

$$\det \begin{pmatrix} x^4 & y^4 & z^4 & 1\\ x^5 & y^5 & z^5 & 1\\ x^{-5} & y^{-5} & z^{-5} & 1\\ x^{-4} & y^{-4} & z^{-4} & 1 \end{pmatrix} = 0$$

if and only if it holds in \mathbb{F}_{q^2} that

$$\det \left(\begin{array}{cc} Y^9 - X^9 & Z^9 - X^9 \\ Y^8 - X^8 & Z^8 - X^8 \end{array} \right) = 0.$$

Proof. Let

$$A = \begin{pmatrix} x^4 & y^4 & z^4 & 1\\ x^5 & y^5 & z^5 & 1\\ x^{-5} & y^{-5} & z^{-5} & 1\\ x^{-4} & y^{-4} & z^{-4} & 1 \end{pmatrix}.$$

Then

$$\det(A) = \det \begin{pmatrix} x^4 - x^{-4} & y^4 - y^{-4} & z^4 - z^{-4} \\ x^5 - x^{-4} & y^5 - y^{-4} & z^5 - z^{-4} \\ x^{-5} - x^{-4} & y^{-5} - y^{-4} & z^{-5} - z^{-4} \end{pmatrix}$$
$$= (xyz)^{-4} \det \begin{pmatrix} x^8 - 1 & x^8 - 1 & z^8 - 1 \\ x^9 - 1 & y^9 - 1 & z^9 - 1 \\ x^{-1} - 1 & y^{-1} - 1 & z^{-1} - 1 \end{pmatrix}.$$

Since $q = 3^m$ with $m \ge 1$ being an integer, one has $x^9 - 1 = (x - 1)^9$, $y^9 - 1 = (y - 1)^9$ and $z^9 - 1 = (z - 1)^9$. Moreover, we have

$$\frac{x^8 - 1}{x - 1} = (1 + x + \dots + x^7 + x^8) - x^8 = \frac{(x - 1)^9}{x - 1} - x^8 = (x - 1)^8 - x^8.$$

Likewise, we have

$$\frac{y^8 - 1}{y - 1} = (y - 1)^8 - y^8, \frac{z^8 - 1}{z - 1} = (z - 1)^8 - z^8.$$

It follows that

$$\begin{aligned} \det(A) &= (xyz)^{-4}(x-1)(y-1)(z-1) \\ &\times \det\left(\begin{array}{ccc} (x-1)^8 - x^8 & (y-1)^8 - y^8 & (z-1)^8 - z^8 \\ (x-1)^8 & (y-1)^8 & (z-1)^8 \\ -x^{-1} & -y^{-1} & -z^{-1} \end{array}\right) \\ &= (xyz)^{-4}(x-1)(y-1)(z-1) \det\left(\begin{array}{ccc} x^8 & y^8 & z^8 \\ (x-1)^8 & (y-1)^8 & (z-1)^8 \\ x^{-1} & y^{-1} & z^{-1} \end{array}\right) \\ &= (xyz)^{-4}(x-1)(y-1)(z-1) \\ &\times \det\left(\begin{array}{ccc} x^8 & y^8 - \frac{x}{y}x^8 & z^8 - \frac{x}{z}x^8 \\ (x-1)^8 & (y-1)^8 - \frac{x}{y}(x-1)^8 & (z-1)^8 - \frac{x}{z}(x-1)^8 \\ x^{-1} & 0 & 0 \end{array}\right) \\ &= x^{-5}(yz)^{-4}(x-1)(y-1)(z-1) \\ &\times \det\left(\begin{array}{ccc} y^8 - x^9y^{-1} & z^8 - x^9z^{-1} \\ (y-1)^8 - x(x-1)^8y^{-1} & (z-1)^8 - x(x-1)^8z^{-1} \end{array}\right) \\ &= (xyz)^{-5}(x-1)(y-1)(z-1) \\ &\times \det\left(\begin{array}{ccc} y^9 - x^9 & z^9 - x^9 \\ y(y-1)^8 - x(x-1)^8 & z(z-1)^8 - x(x-1)^8 \end{array}\right). \end{aligned}$$

But

$$y(y-1)^8 = (y-1)^9 + (y-1)^8 = y^9 - 1 + (y-1)^8$$

and

$$z(z-1)^8 = z^9 - 1 + (z-1)^8.$$

Then one arrives at

$$\begin{aligned} \det(A) &= (xyz)^{-5}(x-1)(y-1)(z-1) \\ &\times \det\left(\begin{array}{c} y^9 - x^9 & z^9 - x^9 \\ (y^9 - 1) + (y-1)^8 - (x^9 - 1) - (x-1)^8 & (z^9 - 1) + (z-1)^8 - (x^9 - 1) - (x-1)^8 \end{array}\right) \\ &= (xyz)^{-5}(x-1)(y-1)(z-1) \\ &\times \det\left(\begin{array}{c} y^9 - x^9 & z^9 - x^9 \\ (y^9 - 1 - x^9 + 1) + (y-1)^8 - (x-1)^8 & (z^9 - 1 - x^9 + 1) + (z-1)^8 - (x-1)^8 \end{array}\right) \\ &= (xyz)^{-5}(x-1)(y-1)(z-1) \\ &\times \det\left(\begin{array}{c} y^9 - x^9 & z^9 - x^9 \\ y^9 - x^9 + (y-1)^8 - (x-1)^8 & z^9 - x^9 + (z-1)^8 - (x-1)^8 \end{array}\right) \\ &= (xyz)^{-5}(x-1)(y-1)(z-1) \\ &\times \det\left(\begin{array}{c} y^9 - x^9 & z^9 - x^9 \\ (y-1)^8 - (x-1)^8 & (z-1)^8 - (x-1)^8 \end{array}\right). \end{aligned}$$
 Since the characteristics of the finite field \mathbb{F}_{q^2} is 3, one has

$$Y^9 - X^9 = (y - 1)^9 - (x - 1)^9 = y^9 - x^9$$

and

$$Z^9 - X^9 = (z - 1)^9 - (x - 1)^9 = z^9 - x^9$$

This yields that

$$\det(A) = XYZ((X+1)(Y+1)(Z+1))^{-5} \det \begin{pmatrix} Y^9 - X^9 & Z^9 - X^9 \\ Y^8 - X^8 & Z^8 - X^8 \end{pmatrix}$$

as required.

Since $x,y,z\in U_{q+1}\backslash\{1\}$ are three pairwise distinct elements, we have

$$XYZ((X+1)(Y+1)(Z+1))^{-5} \neq 0.$$

It then follows immediately that det(A) = 0 if and only if

$$\det \left(\begin{array}{cc} Y^9 - X^9 & Z^9 - X^9 \\ Y^8 - X^8 & Z^8 - X^8 \end{array} \right) = 0$$

as desired.

This finishes the proof of Lemma 3.4.

Lemma 3.5. Let $q = 3^m$ with m being an odd integer. Let $x, y, z \in U_{q+1} \setminus \{1\}$ be three pairwise distinct elements. Write X := x - 1, Y := y - 1 and Z := z - 1. Then it holds in \mathbb{F}_{q^2} that

$$\det \begin{pmatrix} Y^9 - X^9 & Z^9 - X^9 \\ Y^8 - X^8 & Z^8 - X^8 \end{pmatrix} = 0$$
(3.6)

if and only if it holds in \mathbb{F}_{q^2} that

$$\frac{(y-1)(z-x)}{(x-1)(z-y)} = -1.$$
(3.7)

Proof. Firstly, we let (3.6) hold in \mathbb{F}_{q^2} . Then

$$Y^{9}(Z^{8} - X^{8}) - Y^{8}(Z^{9} - X^{9}) + X^{8}Z^{9} - X^{9}Z^{8}$$
$$= (Z - X)\left(Y^{9}\frac{Z^{8} - X^{8}}{Z - X} - Y^{8}(Z - X)^{8} + X^{8}Z^{8}\right) = 0.$$

20

Since

$$Z^{8} - X^{8} = \frac{1}{Z}(Z^{9} - X^{9} - X^{8}(Z - X)) = \frac{Z - X}{Z}((Z - X)^{8} - X^{8}),$$

one has

$$(Z-X)\left(\frac{Y^9}{Z}((Z-X)^8-X^8)-Y^8(Z-X)^8+X^8Z^8\right)=0.$$

But $X \neq Z$ since $x \neq z$. Hence

$$\frac{Y^9}{Z}((Z-X)^8 - X^8) - Y^8(Z-X)^8 = -X^8Z^8$$

It follows that

$$\frac{Y^9}{Z}(Z-X)^8 - Y^8(Z-X)^8 = \frac{Y^9X^8}{Z} - X^8Z^8$$

from which one can deduce that

$$Y^{8}(Z-X)^{8}\left(\frac{Y}{Z}-1\right) = \frac{X^{8}}{Z}(Y^{9}-Z^{9}) = \frac{X^{8}}{Z}(Y-Z)^{9}.$$

That is,

$$Y^{8}(Z - X)^{8}(Y - Z) = X^{8}(Y - Z)^{9}.$$

Since $Y - Z \neq 0$, one derives that

$$\left(\frac{Y(Z-X)}{X(Z-Y)}\right)^8 = 1.$$

Now we let (3.7) be true. Let

$$\Delta := \frac{(y-1)(z-x)}{(x-1)(z-y)}.$$
(3.8)

Then $\frac{Y(Z-X)}{X(Z-Y)} = \Delta \in \mathbb{F}_{q^2}$ and $\Delta^8 = 1$. We claim that $\Delta \in \mathbb{F}_q$. Actually, since $x, y, z \in U_{q+1} \setminus \{1\} \subseteq \mathbb{F}_{q^2}$, we have

$$\begin{split} \Delta^q &= \frac{(y-1)^q (z-x)^q}{(x-1)^q (z-y)^q} \\ &= \frac{(y^q-1)(z^q-x^q)}{(x^q-1)(z^q-y^q)} \\ &= \frac{(y^{-1}-1)(z^{-1}-x^{-1})}{(x^{-1}-1)(z^{-1}-y^{-1})} \\ &= \frac{x^{-1}y^{-1}z^{-1}(1-y)(x-z)}{x^{-1}y^{-1}z^{-1}(1-x)(y-z)} \\ &= \frac{(y-1)(z-x)}{(x-1)(z-y)} = \Delta. \end{split}$$

Hence $\Delta \in \mathbb{F}_q$ as claimed. The claim is proved. Evidently, one has $\Delta \neq 0$. Then the claim tells us that $\Delta^{q-1} = 1$. However, since $q = 3^m$ and $m \geq 1$ is an odd integer, we have $q - 1 \equiv 3 - 1 = 2 \pmod{8}$. Then $\gcd(q - 1, 8) = 2$. So from $\Delta^8 = 1$ and $\Delta^{q-1} = 1$ we deduce that $\Delta^2 = 1$. It follows that $\Delta = 1$ or $\Delta = -1$. If $\Delta(z) = 1$, then (y-1)(z-x) = (x-1)(z-y). This implies that x - y = xz - yz. Since $x \neq y$, it then follows that $z = \frac{x-y}{x-y} = 1$ which contradicts with the hypothesis $z \neq 1$. Hence we must have $\Delta = -1$ as (3.7) wanted. The necessity part is proved.

Conversely, let (3.7) hold in \mathbb{F}_{q^2} . Then

$$\left(\frac{Y(Z-X)}{X(Z-Y)}\right)^8 = \left(\frac{(y-1)(z-x)}{(x-1)(z-y)}\right)^8 = 1.$$

Then

$$(Z - Y)Y^8(Z - X)^9 = (Z - X)X^8(Z - Y)^9.$$

Since

$$(Z - X)^9 = Z^9 - X^9$$
 and $(Z - Y)^9 = Z^9 - Y^9$,

it follows that

$$(Z - Y)Y^8(Z^9 - X^9) = (Z - X)X^8(Z^9 - Y^9).$$

It reduces to

$$Y^{8}Z^{10} - Y^{9}Z^{9} - X^{9}Y^{8}Z = X^{8}Z^{10} - X^{9}Z^{9} - X^{8}Y^{9}Z.$$

Then dividing by Z on both sides gives that

$$Y^{9}(Z^{8} - X^{8}) - Y^{8}(Z^{9} - X^{9}) + X^{8}Z^{9} - X^{9}Z^{8} = 0.$$

We then deduce that

$$\det \left(\begin{array}{cc} Y^9 - X^9 & Z^9 - X^9 \\ Y^8 - X^8 & Z^8 - X^8 \end{array} \right) = 0.$$

Namely, (3.6) is true. The sufficiency part is proved.

The proof of Lemma 3.5 is complete.

Lemma 3.6. Let $q = 3^m$ with m being a positive integer. Let $x, y \in U_{q+1}$ be two elements. Then each of the following holds in \mathbb{F}_{q^2} :

(i). We have $x + y + xy = 0 \iff x + y + 1 = 0 \iff x = y = 1$.

(ii). If either $x, y \in U_{q+1} \setminus \{1\}$, or $x, y \in U_{q+1}$ and $x \neq y$, then $x + y + xy \neq 0$ and $x + y + 1 \neq 0$.

Proof. (i). First of all, we show that $x + y + xy = 0 \iff x + y + 1 = 0$.

In fact, since $x, y \in U_{q+1}$, we have $x^q = x^{-1}$ and $y^q = y^{-1}$. Hence

$$x + y + xy = xy(x^{-1} + y^{-1} + 1) = xy(x^{q} + y^{q} + 1) = xy(x + y + 1)^{q}.$$

But $x \neq 0$ and $y \neq 0$. It then follows that x + y + xy = 0 holds if and only if x + y + 1 = 0 holds as required. So the first statement of part (i) is true.

Consequently, we show that $x + y + 1 = 0 \iff x = y = 1$. Let x = y = 1. Since the characteristics of \mathbb{F}_{q^2} is equal to 3, one has x + y + 1 = 0 holds in \mathbb{F}_{q^2} . Conversely, let x + y + 1 = 0. Then we have x + y = -1 = 2. By the first part of (i), one knows that x + y + xy = 0. We then derive that xy = 1 holds in \mathbb{F}_{q^2} . It follows that x and y are the two roots in \mathbb{F}_{q^2} of the quadratic equation

$$u^{2} - 2u + 1 = (u - 1)^{2} = 0.$$

Thus we must have x = y = 1 as desired. Thus part (i) is proved.

(ii). Let either $x, y \in U_{q+1} \setminus \{1\}$, or $x, y \in U_{q+1}$ and $x \neq y$. Then $(x, y) \neq (1, 1)$. So by part (i), we have $x + y + xy \neq 0$ and $x + y + 1 \neq 0$ as desired.

This concludes the proof of Lemma 3.6.

Lemma 3.7. Let $q = 3^m$ with m being a positive integer. Let $x, y \in U_{q+1} \setminus \{1\}$ be two distinct elements. Then we have

$$-\frac{x+y+xy}{x+y+1} \in U_{q+1} \setminus \{x, y, 1\}.$$

Proof. At first, we show that

$$-\frac{x+y+xy}{x+y+1} \in U_{q+1}$$

Since $x, y \in U_{q+1} \setminus \{1\}$, by Lemma 3.6 (ii) we have $x + y + xy \neq 0$ and $x + y + 1 \neq 0$. Thus

$$\left(-\frac{x+y+xy}{x+y+1}\right)^{q} = -\frac{(x+y+xy)^{q}}{(x+y+1)^{q}}$$
$$= -\frac{x^{q}+y^{q}+(xy)^{q}}{x^{q}+y^{q}+1}$$
$$= -\frac{x^{-1}+y^{-1}+(xy)^{-1}}{x^{-1}+y^{-1}+1}$$
$$= -\frac{x+y+1}{x+y+xy}.$$

It then follows that

$$\left(-\frac{x+y+xy}{x+y+1}\right)^{q+1} = 1$$

which implies that $-\frac{x+y+xy}{x+y+1} \in U_{q+1}$ as one wants. It remains to show that

$$-\frac{x+y+xy}{x+y+1} \notin \{x,y,1\}.$$

This will be done in what follows.

First, suppose that $-\frac{x+y+xy}{x+y+1} = x$. Then x + y + xy + x(x + y + 1) = 0. So $2x + 2xy + x^2 + y = 0$ from which one can derive that $-x - xy + x^2 + y = 0$. That is, (y-x)(1-x) = 0. Hence we have x = y or x = 1. This contradicts with the hypothesis $x \neq y$ and $x \neq 1$. Therefore we must have

$$\frac{x+y+xy}{x+y+1} \neq x$$

as required.

Consequently, suppose that $-\frac{x+y+xy}{x+y+1} = y$. Likewise, we can deduce that (x-y)(1-y) = 0. Thus either x = y or y = 1. We arrive at a contradiction with the hypothesis $x \neq y$ and $y \neq 1$. Hence we must have

$$-\frac{x+y+xy}{x+y+1} \neq y$$

as expected.

Finally, suppose that $-\frac{x+y+xy}{x+y+1} = 1$. Then x + y + xy + x + y + 1 = 0. Namely, -x - y + xy + 1 = 0. Equivalently, (1 - x)(1 - y) = 0. This is impossible since the hypothesis that $x \neq 1$ and $y \neq 1$ implies that $(1 - x)(1 - y) \neq 0$. In conclusion, we have

$$-\frac{x+y+xy}{x+y+1} \neq 1$$

as desired.

This completes the proof of Lemma 3.7.

Lemma 3.8. Let $q = 3^m$ with m being an odd integer. Let $x, y \in U_{q+1} \setminus \{1\}$ be two distinct elements. Then there is a unique element $z \in U_{q+1} \setminus \{x, y, 1\}$ such that the system (3.1) of homogeneous equations has a nonzero solution **X** in \mathbb{F}_q^4 .

Proof. This lemma follows immediately from Lemmas 3.3 to 3.7.

We can now give the proof of Theorem 1.2 as the conclusion of this paper.

Proof of Theorem 1.2. Let \mathcal{C} be the BCH code $\mathcal{C}_{(q,q+1,3,4)}$ with parameters [q+1,q-3,4]. On the one hand, Theorem 1.1 tells us that \mathcal{C} is an AMDS code. And Theorem 2.1 infers that the dual code of \mathcal{C} is an AMDS code if and only if dim $\mathcal{C}(I) = 1$ holds for any subset $I \subseteq \langle n \rangle$ with |I| = n - k + 1 = 5. But by Lemma 3.1, we know that dim $\mathcal{C}(I) = 1$ holds for any subset $I \subseteq \langle n \rangle$ with |I| = n - k + 1 = 5 if and only if for arbitrary two distinct elements $x, y \in U_{q+1} \setminus \{1 := 1_{\mathbb{F}_{q^2}}\} \subseteq \mathbb{F}_{q^2}$, there exists at most one element $z \in U_{q+1} \setminus \{x, y, 1\} \subseteq \mathbb{F}_{q^2}$ such that the system (3.1) of homogeneous equations has a nonzero solution \mathbf{X} in \mathbb{F}_q^4 . But the latter one is true because Lemma 3.8 guarantees that for arbitrary two distinct elements $x, y \in U_{q+1} \setminus \{1 := 1_{\mathbb{F}_{q^2}}\} \subseteq \mathbb{F}_{q^2}$, there is a unique element $z \in U_{q+1} \setminus \{x, y, 1\} \subseteq \mathbb{F}_{q^2}$ such that the system (3.1) of homogeneous equations has a nonzero solution \mathbf{X} in \mathbb{F}_q^4 . Then we can conclude that the dual code of \mathcal{C} is an AMDS code.

This finishes the proof of Theorem 1.2.

References

- 1. T.M. Apostol, Introduction to analytic number theory, Springer-Verlag, New York, 1976.
- D. Bartoli, M. Giulietti and I. Platoni, On the covering radius of MDS codes, *IEEE Trans. Inform.* Theory 61 (2015), 801-811.
- 3. L.S. Chen and S.Y. Shen, *Fundamentals of coding theory (Chinese)*, Higher Education Press, Beijing, 2005.
- 4. M.A. de Boer, Almost MDS codes, Des. Codes Cryptogr. 9 (1996), 143-155.
- C.S. Ding, C.L. Fan and Z.C. Zhou, The dimension and minimum distance of two classes of primitive BCH codes, *Finite Fields Appl.* 45 (2017), 237-263.
- C.S. Ding and C.M. Tang, Infinite families of near MDS codes holding t-designs, IEEE Trans. Inform. Theory 66 (2020), 5419-5428.
- 7. S.M. Dodunekov and I.N. Landgev, On near MDS codes, J. Geometry 54 (1995), 30-43.
- S.M. Dodunekov and I.N. Landjev, Near-MDS codes over some small fields, *Discrete Math.* 213 (2000), 55-65.
- R. Dodunekova, S.M. Dodunekov and T. Klove, Almost-MDS and near-MDS codes for error detection, IEEE Trans. Inform. Theory 43 (1997), 285-290.
- X.J. Geng, M. Yang, J. Zhang and Z.C. Zhou, A class of almost MDS codes, *Finite Fields Appl.* 79 (2022), #101996.
- 11. M. Giulietti, On the extendibility of near-MDS elliptic codes, *Appl. Algebra Engrg. Comm. Comput.* **15** (2004), 1-11.
- B.K. Gong, C.S. Ding and C.J. Li, The dual codes of several classes of BCH codes, *IEEE Trans. Inform. Theory* 68 (2022), 953-964.
- 13. Z.L. Heng, C.J. Li and X.R. Wang, Constructions of MDS, near MDS and almost MDS codes from cyclic subgroups of $\mathbb{F}_{q^2}^*$, *IEEE Trans. Inform. Theory* **68** (2022), 7817-7831.
- 14. W.C. Huffman and V. Pless, *Fundamentals of error-correcting codes*, Cambridge University Press, Cambridge, 2003.
- 15. L.F. Jin and H.B. Kan, Self-dual near MDS codes from elliptic curves, *IEEE Trans. Inform. Theory* **65** (2019), 2166-2170.
- 16. S.X. Li, C.S. Ding, M.S. Xiong and G.N. Ge, Narrow-sense BCH codes over GF(q) with length $n = \frac{q^m 1}{q 1}$, *IEEE Trans. Inform. Theory* **63** (2017), 7219-7236.
- 17. X.R. Li and Z.L. Heng, Constructions of near MDS codes which are optimal locally recoverable codes, *Finite Fields Appl.* 88 (2023), #102184.
- C.J. Li, P. Wu and F.M. Liu, On two classes of primitive BCH codes and some related codes, *IEEE Trans. Inform. Theory* 65 (2019), 3830-3840.
- R. Lidl and H. Niederreiter, *Finite fields*, Second edition, Encyclopedia Math. Appl. 20, Cambridge University Press, Cambridge, 1997.
- Y. Liu, R.H. Li, L.B. Guo and H. Song, Dimensions of nonbinary antiprimitive BCH codes and some conjectures, *Discrete Math.* 346 (2023), #113496.

- F.J. MacWilliams and N.J.A. Sloane, *The theory of error-correcting codes*, North-Holland Publishing Company, Amsterdam, 1977.
- 22. A. Meneghetti, M. Pellegrini and M. Sala, A formula on the weight distribution of linear codes with applications to AMDS codes, *Finite Fields Appl.* **77** (2022), #101933.

23. R.M. Roth, Introduction to coding theory, Cambridge University Press, Cambridge, 2006.

- 24. R. Singleton, Maximum distance q-ary codes, IEEE Trans. Inform. Theory 10 (1964), 116-118.
- Q.Y. Wang and Z.L. Heng, Near MDS codes from oval polynomials, *Discrete Math.* 344 (2021), #112277.

MATHEMATICAL COLLEGE, SICHUAN UNIVERSITY, CHENGDU 610064, P.R. CHINA *Email address:* syqiang23@163.com

MATHEMATICAL COLLEGE, SICHUAN UNIVERSITY, CHENGDU 610064, P.R. CHINA *Email address*: 1810097@mail.nankai.edu.cn

MATHEMATICAL COLLEGE, SICHUAN UNIVERSITY, CHENGDU 610064, P.R. CHINA *Email address:* sfhong@scu.edu.cn