Entanglement and the density matrix renormalisation group in the generalised Landau paradigm

LAURENS LOOTENS,^{1,2,*} CLEMENT DELCAMP,³ and FRANK VERSTRAETE^{1,2}

¹Department of Applied Mathematics and Theoretical Physics, University of Cambridge,

Wilberforce Road, Cambridge, CB3 0WA, United Kingdom

²Department of Physics and Astronomy, Ghent University, Krijgslaan 281, 9000 Gent, Belgium

³Institut des Hautes Études Scientifiques, Bures-sur-Yvette, France

We leverage the interplay between gapped phases and dualities of symmetric one-dimensional quantum lattice models to demonstrate that every phase is efficiently characterised by the maximal breaking of the dual (genereralised) symmetry whose structure encodes the quasiparticle excitations. This result has strong implications for the complexity of simulating many-body systems using variational tensor network methods. For every phase in the phase diagram, the dual representation of the ground state that breaks all symmetries minimises both the entanglement entropy and the required number of variational parameters. We demonstrate the applicability of this idea by developing a generalised density matrix renormalisation group algorithm that works on (dual) constrained Hilbert spaces, and quantify the computational gains obtained over traditional DMRG methods in a perturbed Heisenberg model. Our work testifies to the usefulness of generalised non-invertible symmetries and their formal category theoretic description for the nuts and bolts simulation of strongly correlated systems.

SEC. I | Introduction

The concept of symmetry breaking forms one of the cornerstones of many-body physics. This was already recognized by Pierre Curie in 1894 [1, 2]: "Phenomenon may exist in a medium having the same characteristic symmetry or the symmetry of a subgroup of its characteristic symmetry. What is necessary, is that certain elements of symmetry do not exist. Dissymmetry is what creates the phenomenon." Landau [3] formalised this idea, and demonstrated that different gapped phases are characterised by local order parameters which transform nontrivially under the global symmetry. In 1971, Kadanoff and Ceva [4] turned the tables around and realised that non-local disorder parameters can be used to characterise symmetric phases. Their non-local order parameter was obtained by a Kramers-Wannier duality transformation of the local order parameter in the Ising model [5]. Nonlocal order parameters became much more prominent after the discovery of phases of matter that could only be characterized through such non-local order parameters [6, 7]. These topological phases can be constructed through dualities relating theories with global symmetries to dual (gauge) theories [8-11], and their discovery challenged the standard Landau paradigm of symmetry breaking [12].

The modern approach to characterising phases of matter uses the language of entanglement and quantum circuits. Two Hamiltonians with a given symmetry are in the same phase if and only if there exists a symmetrypreserving constant-depth quantum circuit that transforms their ground states into each other [13, 14]. For the case of one-dimensional quantum lattice models, it follows that ground states of local gapped Hamiltonians exhibit an area law for the entanglement entropy [15]. Such ground states can efficiently be represented in terms of variational matrix product states (MPS) [16], and the corresponding density matrix renormalisation group (DMRG) has effectively broken down the exponential complexity wall for simulating ground states of interacting one-dimensional quantum lattice models [17– 19]. The less entanglement in the ground state, the better the algorithm works.

This entanglement-based approach has also unveiled ways of distinguishing different topological phases of matter via the entanglement degrees of freedom of the MPS: besides the subgroup describing the remaining symmetry, phases are also labelled by the possible ways in which the entanglement degrees of freedom transform projectively under this subgroup [20, 21]. Furthermore, it facilitates the definition of symmetry operators that do not act in an on-site manner, but rather encode a correlated action on neighbouring sites, often on a constrained Hilbert space without tensor product structure. Mathematically, such generalised symmetries [22-32] are described by a *fusion* category [33], and they can be represented explicitly as matrix product operators (MPO) [34–38]. The phases of such systems are classified by a choice of module category compatible with those symmetries [27, 39-43], and the generalised Landau paradigm [29, 44–47] entails that the inclusion of these generalised symmetries yields a complete classification of gapped phases. Interestingly, the same classification labels all possible dualities, or all possible ways in which a (generalised) symmetry can be gauged [48–50]. The explicit intertwiners representing those dualities on the lattice are again of the MPO form [38], and they map local order parameters to non-local ones. By incorporating all possible boundary conditions, one can show that these dualities are unitary and hence preserve the full spectrum of the Hamiltonian [51, 52].

^{*} ll708@cam.ac.uk

In this paper, we connect the concepts of symmetry breaking, dualities and entanglement, and come to the remarkable insight that for every Hamiltonian, one can determine an optimal dual Hamiltonian whose ground states exhibit the minimal amount of entanglement. More precisely, consider a Hamiltonian with a (generalised) symmetry and with a ground state in a certain phase. By a (twisted) gauging of the remaining symmetries, the corresponding dual Hamiltonian will spontaneously break its dual symmetry completely, and the corresponding ground states exhibit the minimal amount of entanglement when compared to all other dual models. As all redundancy due to the symmetries is eliminated, those dual ground states have the minimal possible number of variational parameters, and this leads to a large reduction in complexity for their simulation. The ground states in the original theory can be obtained by multiplying the optimal ones with duality intertwiners in the form of MPO's [38, 51], which reintroduces the multiplicities in the entanglement spectra by enlarging the bond dimension. The fact that generalised symmetries emerge in a very natural way, even in cases where the original Hamiltonian only exhibits global on-site symmetries, makes a strong case for the computational usefulness of the generalised Landau paradigm.

The mathematical underpinnings of this result can be summarised in a commutative diagram:



The arrows denote equivalences between the fusion categories that organise the symmetry, the symmetric Hamiltonians and the quasiparticle excitations. More precisely, the category that describes the algebraic properties of the Hamiltonian and the category that describes the symmetry satisfy an equivalence determined by the kinematical degrees of freedom on which both the Hamiltonian and the symmetry act. Similarly, the phase of the Hamiltonian determines an equivalence between the category that describes the symmetry and the one that describes the fusion properties of quasiparticle excitations. Combining these two provides an equivalence between the Hamiltonian and quasiparticle categories, which governs the properties of the entanglement degrees of freedom in the optimal tensor network representation of the ground state. We explain this diagram in more detail in figure 4.

The only price we pay by considering a dual model is that it may not be defined on a tensor product Hilbert space due to kinematical constraints introduced by the gauging procedure. We deal with this aspect by developing a variant of the DMRG algorithm that directly incorporates these constraints, and demonstrate that all the building blocks for state-of-the-art implementations of tensor networks algorithms are still in place. We illustrate our algorithm with a Heisenberg-like model in various phases, before presenting the general mathematical framework.

In retrospect, there is one scenario where this has already been used to great success. Indeed, when applying our method to an ordinary symmetric phase, it is essentially equivalent to using symmetric tensors [53-55] in which the variational degrees of freedom effectively build up the (non-invertible) symmetry-broken ground states of our dual constrained Hilbert space. By defining Hamiltonians directly in the dual space [38, 51], our method yields a much simpler way of implementing symmetrypreserving tensor networks. This strategy was also used in a visionary paper by Sierra and Nishino from 1997 [56], in which the Heisenberg model was studied by means of its dual interaction round a face Hamiltonian. It took the development of a general theory of dualities for the most general kind of symmetry in terms of matrix product operators to turn this is into a practical algorithm for simulating ground states of Hamiltonians with (generalised) symmetries that is optimal even beyond the trivially symmetric phase.

SEC. II | Illustrative example

Consider an open chain of length L, and assign to every site i spin-1 degrees of freedom with spin operators (S_i^x, S_i^y, S_i^z) . These degrees of freedom are governed by the nearest neighbour Hamiltonian [57]

$$\mathbb{H} = \sum_{i=1}^{L-1} \left(\mathbb{h}_{i,0} + J_1 \mathbb{h}_{i,1} + J_2 \mathbb{h}_{i,2} \right), \tag{1}$$

with coupling constants $J_1, J_2 \in \mathbb{R}$ and local operators

$$\begin{split} & \mathbb{h}_{\mathbf{i},0} := S_{\mathbf{i}}^{x} S_{\mathbf{i}+1}^{x} + S_{\mathbf{i}}^{y} S_{\mathbf{i}+1}^{y} + S_{\mathbf{i}}^{z} S_{\mathbf{i}+1}^{z} \,, \\ & \mathbb{h}_{\mathbf{i},1} := (S_{\mathbf{i}}^{x} S_{\mathbf{i}+1}^{x})^{2} + (S_{\mathbf{i}}^{y} S_{\mathbf{i}+1}^{y})^{2} + (S_{\mathbf{i}}^{z} S_{\mathbf{i}+1}^{z})^{2} \,, \\ & \mathbb{h}_{\mathbf{i},2} := \{S_{\mathbf{i}}^{x}, S_{\mathbf{i}}^{y}\} S_{\mathbf{i}+1}^{z} + \{S_{\mathbf{i}}^{z}, S_{\mathbf{i}}^{x}\} S_{\mathbf{i}+1}^{y} + \{S_{\mathbf{i}}^{y}, S_{\mathbf{i}}^{z}\} S_{\mathbf{i}+1}^{x} \,. \end{split}$$

While the term $\mathbb{h}_{i,0}$ defines the spin-1 antiferromagnetic Heisenberg model, which is SO(3) symmetric, the terms $\mathbb{h}_{i,1}$ and $\mathbb{h}_{i,2}$ are perturbations breaking the symmetry down to the finite subgroup $\mathbb{A}_4 \subset SO(3)$ of orientationpreserving symmetries of the tetrahedron. Specifically, \mathbb{A}_4 is isomorphic to the semidirect product $\mathbb{Z}_3 \ltimes \mathbb{D}_2$, where \mathbb{Z}_3 is generated by any cyclic permutation of (S^x, S^y, S^z) , whereas $\mathbb{D}_2 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ is generated by $e^{i\pi S^x}$ and $e^{i\pi S^z}$.

In the presence of a symmetry \mathbb{A}_4 , we distinguish *seven* possible *gapped* phases. These are labelled by a subgroup $H \subseteq \mathbb{A}_4$ characterising the symmetry of the ground states, together with a class $[\psi]$ labeling the projective representations that classify possible symmetry protected topological (SPT) orders [20]. Physically, a non-trivial $[\psi]$ signals the presence of edge modes that transform projectively under the action of the symmetry H. Up to isomorphisms, \mathbb{A}_4 counts five subgroups, namely \mathbb{A}_4 ,

 \mathbb{D}_2 , \mathbb{Z}_3 , \mathbb{Z}_2 as well as the trivial one. Out of these subgroups, only \mathbb{A}_4 and \mathbb{D}_2 have non-trivial second cohomology groups, which are both isomorphic to \mathbb{Z}_2 .

In order to proceed with our analysis, it is crucial to rewrite the local operators entering the definition of the Hamiltonian (1) in such a way that the symmetry \mathbb{A}_4 is manifest. Invoking a finite group version of the Wigner-Eckart theorem, we know that any operator transforming trivially under \mathbb{A}_4 must be expressible as a linear combination of Clebsch-Gordan coefficients. The group \mathbb{A}_4 possesses three one-dimensional irreducible representations $\{\underline{0}, \underline{1}, \underline{1}^*\}$ and a single three-dimensional one $\underline{3}$ satisfying $\underline{3} \otimes \underline{3} \cong \underline{0} \oplus \underline{1} \oplus \underline{1}^* \oplus 2 \cdot \underline{3}$. Given three irreducible representations V_1, V_2 and V_3 such that $V_3 \subset V_1 \otimes V_2$, we interpret the intertwining map $V_3 \to V_1 \otimes V_2$ as the tensor

$$\begin{array}{c} \begin{array}{c} & & \\$$

where the sums are over basis vectors, and *i* enumerates the different ways $V_1 \otimes V_2$ decomposes into V_3 . In this equation, the diagram on the r.h.s. depicts the *Clebsch– Gordan* coefficients valued in \mathbb{C} . In this notation, we can show that the local operators $\mathbb{h}_{i,n}$, $n \in \{0, 1, 2\}$, are all of the form

where $h_n(V, i, j) \in \mathbb{C}$. Notice that in this formulation, the state space of a given spin-1 degree of freedom is spanned by $|\underline{3}, v\rangle$, with $v = 1, \ldots, 3$.

II.A. Dualities

Importantly, we have the following equality of intertwining maps $V_4 \rightarrow V_1 \otimes V_2 \otimes V_3$



where the '*F*-symbols' $(F_{V_4}^{V_1V_2V_3})_{V_5,ij}^{V_6,kl} \in \mathbb{C}$ are provided by the *Racah W-coefficients* of \mathbb{A}_4 . One can use this identity to show that the structure factors of the algebra generated by the local symmetric operators $\{\mathbb{h}_{i,n}\}_{i,n}$ only depends on the *F*-symbols and coefficients $\{h_n\}_n$; the eigenvalues of the full Hamiltonian only depend on those structure factors, and not the specific representation chosen. Interpreting eq. (3) as a tensor network equation, one can ask the following question: Is there another set of tensors, generically depicted as

also indexed by triplets of representations $\{V_1, V_2, V_3 \subset V_1 \otimes V_2\}$, satisfying eq. (3)? Keeping $\{h_n\}_n$ the same, replacing the intertwining maps in eq. (2) by these new tensors would result in an isomorphic algebra of local operators, and thus a dual model with Hamiltonian [38]

$$\mathbb{h}_{\mathbf{i},n}^{\mathrm{dual}} = \sum_{V} \sum_{i,j} h_n(V,i,j) \begin{vmatrix} \frac{1}{3} \\ \frac{3}{1} \\ \frac{1}{2} \\ \frac{1}{3} \end{vmatrix} = \frac{1}{2} \cdot \frac{1}{3} \cdot$$

When dealing with a symmetry \mathbb{A}_4 , one can find collections of tensors of the form (4) satisfying eq. (3) for any $H \subseteq \mathbb{A}_4$ and $[\psi] \in H^2(H, U(1))$ [58]. This is because dualities correspond to (twisted) gauging maps of the symmetry A_4 , and there are as many ways to gauge a symmetry as there are ways to spontaneously break it. Typically, the dual Hamiltonian acts on a distinct microscopic Hilbert space, which is not necessarily a tensor product space as suggested by the graphical notation. This is consistent with the gauging interpretation, leading to theories with gauge degrees of freedom that satisfy local Gauß constraints. Given a pair $(H, [\psi])$, degrees of freedom of the resulting dual model are associated with irreducible ψ -projective representations [38], and as such we label the model by $\operatorname{\mathsf{Rep}}^{\psi}(H)$. Instead of Clebsch-Gordan coefficients, the tensors are found to evaluate to Racah W-coefficients involving linear representations of G and ψ -projective representations of H. Crucially, the initial Hamiltonian can be transmuted into any of its duals via an MPO [51]:



which is true for any V, i and j.

II.B. Numerical simulations

We consider points $(J_1, J_2) = \{(1, 1), (-2, -5), (-5, 1)\}$ in the phase diagram of the model (1) corresponding to the A₄ SPT phase, the A₄ symmetric phase and the D₂ symmetric phase, respectively. For each point, we simulate all the seven dual Hamiltonians labelled by $\operatorname{Rep}^{\psi}(H)$, which result from the ψ -twisted gauging of $H \subseteq G$, the first of which (coined "Original") corresponds to the standard DMRG algorithm without symmetries and hence no constraints. The ground states of these dual models are found using the generalised DMRG algorithm (c.f. app. A), which performs a variational optimisation within the subspace of the constrained Hilbert space that is spanned by MPSs of the form

$$\therefore \underbrace{A_{i-1}}_{A_{i-1}} \underbrace{A_{i}}_{A_{i}} \underbrace{A_{i}}_{A_{i+1}} \underbrace{A_{i}}_{A_{i+1}} \ldots$$
 (7)

Due to the constraints, the MPS matrices carry gauge degrees of freedoms associated with ψ -projective representations of H, which are shared by neighbouring sites. These are represented by the blue lines in eq. (7). For the three phases, we plot the *entanglement spectra* of the dual ground states coloured by the different objects labelling the gauge degrees of freedom, as well as the memory requirements to reach a specific minimal Schmidt value λ_{\min} serving as an error measure. Our findings are displayed on fig. 1, 2 and 3 and analysed below:

• \mathbb{A}_4 SPT, fig. 1: In this phase, entanglement degrees of freedom of the unique ground state transform as pro*jective* representations of A_4 . The three irreducible projective representations of \mathbb{A}_4 are two-dimensional, which explains the two-fold degeneracy for every Schmidt value. Comparing the entanglement spectra of the various dual models, we observe that the entanglement is minimised in the $\operatorname{\mathsf{Rep}}^{\psi}(\mathbb{D}_2)$ and $\operatorname{\mathsf{Rep}}^{\psi}(\mathbb{A}_4)$ models. Additionally, the ground state of the $\operatorname{\mathsf{Rep}}^{\psi}(\mathbb{A}_4)$ model requires the least amount of memory, making this model the most efficient one to simulate. Crucially, this dual model possesses a non-invertible $\mathsf{Rep}(\mathbb{A}_4)$ symmetry, whereby symmetry operators are labelled by representations of \mathbb{A}_4 , which happens to be completely broken in the ground state subspace. An important subtlety with a non-invertible symmetry breaking phase is that the different ground states, being related by non-trivial MPOs, can have different entanglement properties. In this particular example, the ground state labelled by the 3d irreducible representation requires more entanglement and therefore more variational parameters than the other ground states. In practice, it is possible to avoid this ground state by biasing the initial MPS towards the other less entangled ground states.

All the other dual models admit ground states that preserve some symmetry, resulting in these cases in degeneracy in the entanglement spectrum as well excessive memory requirements. This is ostensible in the $\text{Rep}(\mathbb{D}_2)$ model, where every Schmidt value has a four-fold degeneracy due to its ground state preserving a dual \mathbb{A}_4 symmetry, the \mathbb{D}_2 subgroup of which permuting the gauge degrees of freedom that label the Schmidt values, which



Figure 1. Entanglement spectra of the dual models in the middle of the ground state on 60 sites in the \mathbb{A}_4 SPT phase of the initial model $(J_1 = 1, J_2 = 1)$. The colour of a Schmidt value indicates the object that labels the corresponding gauge degree of freedom. Bottom: The memory required to store a ground state MPS tensor in the bulk at double precision for a given truncation error λ_{\min} . The ground state of the Rep^{ψ} (\mathbb{A}_4) model minimizes the entanglement and the number of variational parameters for a fixed truncation error.

happen to be labelled by irreducible representations of \mathbb{D}_2 . A more subtle case is the $\text{Rep}(\mathbb{A}_4)$ model, which has a non-invertible $\text{Rep}(\mathbb{A}_4)$ symmetry that is only partially broken in the ground state subspace. Although the remaining symmetry constrains the entanglement spectrum, the degeneracy is hidden as a fixed $\sqrt{3}$ ratio (the "quantum dimension" of the irrep) of consecutive Schmidt values labelled by the 3d and a 1d irreducible representation. We make this manifest by rescaling the spectrum appropriately.

• \mathbb{A}_4 symmetric, fig. 2: In this phase, entanglement degrees of freedom of the unique ground state transform as *linear* representations of \mathbb{A}_4 , the three-dimensional irreducible representation <u>3</u> explaining occurrences of threefold degeneracy in the entanglement spectrum. Comparing the entanglement spectra of the various dual models, we observe that the entanglement is minimised in the $\operatorname{Rep}(\mathbb{A}_4)$ model. Additionally, it is the model whose ground state requires the least amount of memory, making this model the most efficient one to simulate. As for the previous phase, this optimal dual model has a noninvertible symmetry $\operatorname{Rep}(\mathbb{A}_4)$, which also happens to be completely broken in the ground state subspace.



Figure 2. Top: Entanglement spectra of the dual models in the middle of the ground state on 60 sites in the \mathbb{A}_4 symmetric phase of the original model $(J_1 = -2, J_2 = -5)$. The colour of a Schmidt value indicates the object that labels the corresponding gauge degree of freedom. Bottom: The memory required to store a ground state MPS tensor in the bulk at double precision for a given truncation error λ_{\min} . The ground state of the $\text{Rep}(\mathbb{A}_4)$ model minimises the entanglement and number of variational parameters for a fixed truncation error.

All the other dual models admit ground states sharing the same entanglement spectrum as the initial model, differing only in the labelling of the Schmidt values and the improvement in the memory requirements. These dual models possess either invertible or non-invertible symmetries, and the ground states break various amounts thereof, but never is the whole symmetry broken.

• \mathbb{D}_2 symmetric, fig. 3: In this phase, entanglement degrees of freedom of the ground states transform as linear representations of \mathbb{D}_2 . Since irreducible representations of \mathbb{D}_2 are all one-dimensional, no additional degeneracy in the entanglement spectrum is enforced, the visible twofold degeneracies must originate from a hidden symmetry that might involve time reversal combined with a physical on-site action. Comparing the entanglement spectra of the various dual models, we observe that the entanglement is minimised in the initial model, as well as the $\operatorname{\mathsf{Rep}}^{\psi}(\mathbb{D}_2)$, $\operatorname{\mathsf{Rep}}(\mathbb{D}_2)$ and $\operatorname{\mathsf{Rep}}(\mathbb{Z}_2)$ models. However, the $\mathsf{Rep}(\mathbb{D}_2)$ model stands out as requiring the least amount of memory, making this model the most efficient one to simulate. This optimal dual model has an A_4 symmetry, which happens to be completely broken in the ground state subspace.



Figure 3. Top: Entanglement spectra of the dual models in the middle of the ground state on 60 sites in the \mathbb{D}_2 symmetric phase of the original model $(J_1 = -5, J_2 = 1)$. The colour of a Schmidt value indicates the object that labels the corresponding gauge degree of freedom. Bottom: The memory required to store a ground state MPS tensor in the bulk at double precision for a given truncation error λ_{\min} . The ground state of the $\text{Rep}(\mathbb{D}_2)$ model minimizes the entanglement and the number of variational parameters for a fixed truncation error.

The remaining dual models admit ground states showing more entanglement, as a consequence of Schmidt values in the initial ground state becoming degenerate. This is most easily understood in the $\operatorname{Rep}(\mathbb{Z}_3)$ model that possesses a non-invertible $\operatorname{Rep}(\mathbb{A}_4)$, which is fully preserved by the unique ground state. The one-dimensional irreducible representations act by permuting gauge degrees of freedom labelled by irreducible representations of \mathbb{Z}_3 , enforcing an additional three-fold degeneracy. A similar explanation holds for the degenerate Schmidt values in the $\operatorname{Rep}(\mathbb{A}_4)$ and $\operatorname{Rep}^{\psi}(\mathbb{A}_4)$ models.

What is the main lesson of this analysis? Regardless of the phase, the optimal dual model to simulate is always the single one whose dual symmetry is completely broken in the ground state subspace. This is the model obtained by performing the (possibly ψ -twisted) gauging of the symmetry H that is preserved within the ground state subspace of the initial model. Incidentally, this method may also be employed to identify a gapped phase by comparing entanglement spectra of dual models.

II.C. Symmetry-preserving tensor networks

We close our study of this example with an observation that sheds light on the computational gains that we just observed. In the symmetric phase, the Hamiltonian of the dual model $\operatorname{Rep}(\mathbb{A}_4)$ —resulting from gauging the \mathbb{A}_4 symmetry of the initial model—is transmuted into the initial one via an MPO that takes a particularly simple form. Specifically, the building blocks of this MPO evaluate to Clebsch–Gordan coefficients. By definition, the ground state of the initial model is obtained by acting with this MPO upon the ground state of the dual model $\operatorname{Rep}(\mathbb{A}_4)$. Graphically, it yields an MPS of the form



i.e., a linear combination of A_4 -invariant tensors. But, decomposing an MPS in this way is precisely the starting point of symmetry preserving tensor network algorithms (c.f. app. A) [53–55]. Concisely, in this form, the symmetry of the state can be exploited to yield a computational gain, by targeting for instance a specific charge sector. By directly computing the ground state of the dual $\operatorname{Rep}(\mathbb{A}_4)$ model, our method provides a remarkably simple way to implement symmetry preserving tensor network algorithms, which is applicable for any symmetry group, including generalised symmetries. The projection onto a particular charge sector is enforced locally via the Gauß constraints together with the choice of boundary conditions. Importantly however, our previous analysis revealed that the model $\operatorname{Rep}(\mathbb{A}_4)$ is the optimal one to simulate if and only if the initial model is in the \mathbb{A}_4 symmetric phase. By accessing the other dual models, our approach overcomes this limitation and yields the optimal DMRG simulation in all possible phases.

In practice, symmetric tensor networks have found most of their utility in models with continuous Lie group symmetries such as SU(2). While we restricted ourselves to a finite group, our results readily generalise to these cases, implying for instance that a ground state preserving an SU(2) symmetry is most efficiently parameterised in terms of a dual non-invertible $\operatorname{Rep}(\operatorname{SU}(2))$ symmetry breaking ground state. Irreducible representations with high spins are given weight zero, so we only have to deal with a finite number of labels, just as in standard symmetric DMRG codes. This scenario turns out to be equivalent to simulation a dual interaction-round-a-face model [56]. The different symmetry-broken ground states can be related by acting with MPO symmetry operators. As a consequence, these ground states do not have the same entanglement, showing the importance of properly initialising the DMRG algorithm to favour the ground states with the least amount of entanglement.

SEC. III | General framework

The results presented above hold much more broadly. Consider a *generalised* symmetry in a one-dimensional quantum lattice model, i.e., a symmetry whose operators are not necessarily invertible [24–29, 31, 32, 49]. The symmetry operators take the form of (typically non-local) MPOs [35–38, 51]. Mathematically, a *finite* generalised symmetry can be modelled by a so-called fusion category [59], extending the group theoretical formalism of ordinary symmetries. Similar to ordinary symmetries, generalised symmetries can be spontaneously brokenas we already witnessed in our examples—and may be gauged provided that there is no 't Hooft anomaly. Furthermore, gapped phases with a generalised symmetry are still in one-to-one correspondence with the different ways to gauge a (sub)symmetry thereof [27, 40], and thus dualities. This is the essence of the generalised Landau paradigm [26, 27, 40, 42, 60, 61].

Consider a one-dimensional quantum lattice model with a generalised symmetry. Suppose the symmetry is completely broken in the ground state subspace. By gauging the symmetry, which amounts to identifying the corresponding symmetry operators as well as the corresponding symmetry broken states, we obtain a dual model with a dual symmetry that is fully preserved by the unique ground state. Crucially, gauging this dual symmetry recovers the initial model. More generally, there is always a way to gauge the (sub)symmetry that is preserved in the ground state subspace of a model so as to yield a dual model whose dual generalised symmetry is completely broken (c.f. app. C). Practically, this dual model is obtained by extending the approach followed in our series of examples: We write the Hamiltonian in terms of tensors that make the generalised symmetry of the model manifest; these tensors satisfy equations generalised eq. (3); there is a different set of solutions to these equations, which correspond to the relevant gauging of the preserved subsymmetry, that yield the dual model (c.f. app. C).

We claim that the optimal way of simulating the phase of a given model amounts to simulating the dual phase of this dual model where the dual symmetry is completely broken, after which we recover the original ground state by acting with the MPO that transmutes the corresponding Hamiltonians into each other. Broadly speaking, the reasoning is that any symmetry translates into constraints amongst the variational parameters so they are not all independent. The associated redundancy unequivocally translates into a suboptimal use of computational resources, as we observed in the example above.

In the optimal dual phase, at least one of the ground states has the property that the action of any dual symmetry operator on it yields an orthogonal ground state. For this maximal symmetry breaking state, all order parameters are strictly local. This follows from the fact that the action of the MPOs representing the dual symmetries map such a ground state into a different one, and hence the expectation value of any non-local string order operator vanishes exponentially in the number of sites on which it acts [62]. Conversely, all quasiparticle excitations on top of this maximal symmetry breaking ground state correspond to domain wall excitations. In the case of an infinite one dimensional lattice model, these excitations are created by the action of the symmetry MPOs on one half of the chain [63]. As in the usual ansatz for topological excitations in MPS, additional variational degrees of freedom characterising the precise nature of the excitations emerge at the endpoint of the MPO [64, 65]. The domain wall excitations of the dual symmetry breaking model are mapped to the quasiparticle excitations of the original model, which can be of a very different nature (spinon, holon, etc.). Due to the unitarity of the duality transformation however, they retain the same algebraic properties, and in particular their fusion properties are described by the same categorical data (c.f. figure 4).

SEC. IV | Outlook

In appendix (A) we develop an algorithm that performs the optimal (two-site) DMRG simulation of any gapped phase with an arbitrary generalised symmetry. In symmetric phases, it offers a simpler and more versatile implementation of symmetry-preserving tensor network algorithms, as it bypasses many aspects of traditional implementations such as fusion trees. Our formalism will be particularly interesting for studying second order phase transitions, defects in critical theories, and boundary effects, as dual models exhibit different finite-entanglement scaling regimes [66]. Currently, our algorithm only takes advantage of *internal* symmetries, since it exploits an approach to dualities that has been tailored to this type of symmetry. It will be very interesting to generalise this approach to different types of symmetry, such as spatial symmetries and time reversal, which lead to further refinenements of phase diagrams. Note also that essentially the same ideas can be be used to construct more efficient algorithms for simulating quantum spin systems using neural network states [67].

One of the main merits of our method is that it is systematically extensible to higher dimensional models, and to tensor network network algorithms in terms of projected entangled pair states [68]. In fact, taking advantage of symmetries in higher dimensions is expected to be even more beneficial than in (1+1)d. Although formally more challenging, many aspects of dualities in twodimensional quantum lattice models have been worked out [69]. Specifically, the relevant tensor network operators are already known for a large class of generalised symmetries [70-72]. For instance, a (2+1)d model with an ordinary global symmetry in the symmetric phase can be understood as a dual 1-form symmetry breaking phase obtained by gauging the global symmetry. Our approach reveals that (2+1)d symmetric tensor network methods [54, 55, 73] effectively implement this duality.

Acknowledgments: We would like to thank Jacob Bridgeman, Lukas Devos, José Garre-Rubio, Jutho Haegeman and Sukhwinder Singh for interesting discussions and useful comments. This work has received funding from EOS (grant No. 40007526), IBOF(grant No. IBOF23/064), BOF-GOA (grant BOF23/GOA/021). LL is supported by an No. FWO postdoctoral fellowship (grant No. 12AUN24N) and an EPSRC postdoctoral fellowship (grant No. *EP/Y020456/1*).

References

- P. Curie, Sur la symétrie dans les phénomènes physiques, symétrie d'un champ électrique et d'un champ magnétique, J. Phys. Theor. Appl. 3, 393 (1894).
- [2] E. Castellani and J. Ismael, Which Curie's Principle?, Philosophy of Science 83, 1002 (2016).
- [3] L. D. Landau, On the theory of phase transitions, Zh. Eksp. Teor. Fiz. 7, 19 (1937).
- [4] L. P. Kadanoff and H. Ceva, Determination of an opeator algebra for the two-dimensional Ising model, Phys. Rev. B 3, 3918 (1971).
- [5] H. A. Kramers and G. H. Wannier, Statistics of the twodimensional ferromagnet. Part 1., Phys. Rev. 60, 252 (1941).
- [6] F. J. Wegner, Duality in Generalized Ising Models and Phase Transitions Without Local Order Parameters, J. Math. Phys. 12, 2259 (1971).
- [7] M. den Nijs and K. Rommelse, Preroughening transitions in crystal surfaces and valence-bond phases in quantum spin chains, Phys. Rev. B 40, 4709 (1989).
- [8] X. G. Wen, Topological Order in Rigid States, Int. J. Mod. Phys. B 4, 239 (1990).
- [9] A. Lopez and E. Fradkin, Fractional quantum hall effect and chern-simons gauge theories, Phys. Rev. B 44, 5246 (1991).
- [10] T. Kennedy and H. Tasaki, Hidden $Z_2 \times Z_2$ symmetry breaking in Haldane-gap antiferromagnets, Phys. Rev. B **45**, 304 (1992).
- [11] B. I. Halperin, P. A. Lee, and N. Read, Theory of the half-filled Landau level, Phys. Rev. B 47, 7312 (1993).
- [12] X.-G. Wen, Quantum Field Theory of Many-Body Systems: From the Origin of Sound to an Origin of Light and Electrons (Oxford University Press, 2007).
- [13] M. B. Hastings, Locality in quantum systems (2010), arXiv:1008.5137 [math-ph].
- [14] S. Bravyi, M. B. Hastings, and F. Verstraete, Lieb-Robinson Bounds and the Generation of Correlations and Topological Quantum Order, Phys. Rev. Lett. 97, 050401 (2006), arXiv:quant-ph/0603121.
- [15] M. B. Hastings, An area law for one-dimensional quantum systems, J. Stat. Mech. 0708, P08024 (2007), arXiv:0705.2024 [quant-ph].
- [16] F. Verstraete and J. I. Cirac, Matrix product states represent ground states faithfully, Phys. Rev. B 73, 094423 (2006).
- [17] S. R. White, Density matrix formulation for quantum renormalization groups, Phys. Rev. Lett. 69, 2863 (1992).
- [18] U. Schollwoeck, The density-matrix renormalization group in the age of matrix product states, Annals Phys. 326, 96 (2011), arXiv:1008.3477 [cond-mat.str-el].
- [19] J. I. Cirac, D. Pérez-García, N. Schuch, and F. Ver-

strates, Matrix product states and projected entangled pair states: Concepts, symmetries, theorems, Rev. Mod. Phys. **93**, 045003 (2021).

- [20] X. Chen, Z.-C. Gu, and X.-G. Wen, Complete classification of one-dimensional gapped quantum phases in interacting spin systems, Phys. Rev. B 84, 235128 (2011).
- [21] N. Schuch, D. Pérez-García, and I. Cirac, Classifying quantum phases using matrix product states and projected entangled pair states, Phys. Rev. B 84, 165139 (2011).
- [22] A. Feiguin, S. Trebst, A. W. W. Ludwig, M. Troyer, A. Kitaev, Z. Wang, and M. H. Freedman, Interacting anyons in topological quantum liquids: The golden chain, Phys. Rev. Lett. 98, 160409 (2007).
- [23] N. Bultinck, M. Mariën, D. J. Williamson, M. B. Şahinoğlu, J. Haegeman, and F. Verstraete, Anyons and matrix product operator algebras, Annals Phys. 378, 183 (2017), arXiv:1511.08090 [cond-mat.str-el].
- [24] D. Aasen, R. S. K. Mong, and P. Fendley, Topological defects on the lattice: I. the ising model, Journal of Physics A: Mathematical and Theoretical 49, 354001 (2016).
- [25] M. Buican and A. Gromov, Anyonic Chains, Topological Defects, and Conformal Field Theory, Commun. Math. Phys. 356, 1017 (2017), arXiv:1701.02800 [hep-th].
- [26] C.-M. Chang, Y.-H. Lin, S.-H. Shao, Y. Wang, and X. Yin, Topological Defect Lines and Renormalization Group Flows in Two Dimensions, JHEP 01, 026, arXiv:1802.04445 [hep-th].
- [27] R. Thorngren and Y. Wang, Fusion category symmetry. Part I. Anomaly in-flow and gapped phases, JHEP 04, 132, arXiv:1912.02817 [hep-th].
- [28] D. Aasen, P. Fendley, and R. S. K. Mong, Topological Defects on the Lattice: Dualities and Degeneracies (2020), arXiv:2008.08598 [cond-mat.stat-mech].
- [29] L. Kong, T. Lan, X.-G. Wen, Z.-H. Zhang, and H. Zheng, Algebraic higher symmetry and categorical symmetry: A holographic and entanglement view of symmetry, Phys. Rev. Res. 2, 043086 (2020).
- [30] D. S. Freed, G. W. Moore, and C. Teleman, Topological symmetry in quantum field theory (2022), arXiv:2209.07471 [hep-th].
- [31] S. Schafer-Nameki, ICTP lectures on (non-)invertible generalized symmetries, Phys. Rept. 1063, 1 (2024), arXiv:2305.18296 [hep-th].
- [32] S.-H. Shao, What's Done Cannot Be Undone: TASI Lectures on Non-Invertible Symmetries (2023), arXiv:2308.00747 [hep-th].
- [33] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor categories*, Vol. 205 (American Mathematical Soc., 2016).
- [34] M. B. Şahinoğlu, D. Williamson, N. Bultinck, M. Mariën, J. Haegeman, N. Schuch, and F. Verstraete, Characterizing Topological Order with Matrix Product Operators, Annales Henri Poincare 22, 563 (2021), arXiv:1409.2150 [quant-ph].
- [35] R. Vanhove, M. Bal, D. J. Williamson, N. Bultinck, J. Haegeman, and F. Verstraete, Mapping topological to conformal field theories through strange correlators, Phys. Rev. Lett. **121**, 177203 (2018).
- [36] L. Lootens, J. Fuchs, J. Haegeman, C. Schweigert, and F. Verstraete, Matrix product operator symmetries and intertwiners in string-nets with domain walls, SciPost Phys. 10, 053 (2021).
- [37] J. C. Bridgeman, L. Lootens, and F. Verstraete, Invertible Bimodule Categories and Generalized Schur Or-

thogonality, Commun. Math. Phys. **402**, 2691 (2023), arXiv:2211.01947 [math.QA].

- [38] L. Lootens, C. Delcamp, G. Ortiz, and F. Verstraete, Dualities in one-dimensional quantum lattice models: Symmetric hamiltonians and matrix product operator intertwiners, PRX Quantum 4, 020357 (2023).
- [39] A. Kitaev and L. Kong, Models for Gapped Boundaries and Domain Walls, Commun. Math. Phys. 313, 351 (2012), arXiv:1104.5047 [cond-mat.str-el].
- [40] Z. Komargodski, K. Ohmori, K. Roumpedakis, and S. Seifnashri, Symmetries and strings of adjoint QCD₂, JHEP 03, 103, arXiv:2008.07567 [hep-th].
- [41] R. Thorngren and Y. Wang, Fusion category symmetry. Part II. Categoriosities at c = 1 and beyond, JHEP 07, 051, arXiv:2106.12577 [hep-th].
- [42] K. Inamura, On lattice models of gapped phases with fusion category symmetries, JHEP 03, 036, arXiv:2110.12882 [cond-mat.str-el].
- [43] J. Garre-Rubio, L. Lootens, and A. Molnár, Classifying phases protected by matrix product operator symmetries using matrix product states, Quantum 7, 927 (2023).
- [44] D. Gaiotto and J. Kulp, Orbifold groupoids, JHEP 02, 132, arXiv:2008.05960 [hep-th].
- [45] F. Apruzzi, F. Bonetti, I. n. García Etxebarria, S. S. Hosseini, and S. Schafer-Nameki, Symmetry TFTs from String Theory, Commun. Math. Phys. 402, 895 (2023), arXiv:2112.02092 [hep-th].
- [46] L. Bhardwaj, L. E. Bottini, D. Pajer, and S. Schafer-Nameki, Categorical Landau Paradigm for Gapped Phases (2023), arXiv:2310.03786 [cond-mat.str-el].
- [47] S.-J. Huang and M. Cheng, Topological holography, quantum criticality, and boundary states (2023), arXiv:2310.16878 [cond-mat.str-el].
- [48] J. Frohlich, J. Fuchs, I. Runkel, and C. Schweigert, Duality and defects in rational conformal field theory, Nucl. Phys. B 763, 354 (2007), arXiv:hep-th/0607247.
- [49] L. Bhardwaj and Y. Tachikawa, On finite symmetries and their gauging in two dimensions, JHEP 03, 189, arXiv:1704.02330 [hep-th].
- [50] Y. Tachikawa, On gauging finite subgroups, SciPost Phys. 8, 015 (2020).
- [51] L. Lootens, C. Delcamp, and F. Verstraete, Dualities in one-dimensional quantum lattice models: Topological sectors, PRX Quantum 5, 010338 (2024).
- [52] L. Lootens, C. Delcamp, D. Williamson, and F. Verstraete, Low-depth unitary quantum circuits for dualities in one-dimensional quantum lattice models (2023), arXiv:2311.01439 [quant-ph].
- [53] I. P. McCulloch and M. Gulácsi, The non-abelian density matrix renormalization group algorithm, Europhysics Letters 57, 852 (2002).
- [54] S. Singh, R. N. C. Pfeifer, and G. Vidal, Tensor network decompositions in the presence of a global symmetry, Phys. Rev. A 82, 050301 (2010).
- [55] A. Weichselbaum, Non-abelian symmetries in tensor networks: A quantum symmetry space approach, Annals of Physics **327**, 2972 (2012).
- [56] G. Sierra and T. Nishino, The density matrix renormalization group method applied to interaction round a face hamiltonians, Nuclear Physics B 495, 505 (1997).
- [57] A. Prakash, C. G. West, and T.-C. Wei, Detection of gapped phases of a one-dimensional spin chain with onsite and spatial symmetries, Phys. Rev. B 94, 045136 (2016).

- [58] V. Ostrik, Module categories over the Drinfeld double of a finite group, International Mathematics Research Notices 2003, 1507 (2003).
- [59] P. Etingof, D. Nikshych, and V. Ostrik, On fusion categories (2002), arXiv:math/0203060.
- [60] T.-C. Huang, Y.-H. Lin, and S. Seifnashri, Construction of two-dimensional topological field theories with noninvertible symmetries, JHEP 12, 028, arXiv:2110.02958 [hep-th].
- [61] L. Bhardwaj, L. E. Bottini, D. Pajer, and S. Schäfer-Nameki, Gapped Phases with Non-Invertible Symmetries: (1+1)d (2023), arXiv:2310.03784 [hep-th].
- [62] J. Haegeman, V. Zauner, N. Schuch, and F. Verstraete, Shadows of anyons and the entanglement structure of topological phases, Nature communications 6, 8284 (2015).
- [63] J. Haegeman, B. Pirvu, D. J. Weir, J. I. Cirac, T. J. Osborne, H. Verschelde, and F. Verstraete, Variational matrix product ansatz for dispersion relations, Phys. Rev. B 85, 100408 (2012).
- [64] V. Zauner-Stauber, L. Vanderstraeten, J. Haegeman, I. P. McCulloch, and F. Verstraete, Topological nature of spinons and holons: Elementary excitations from matrix product states with conserved symmetries, Phys. Rev. B 97, 235155 (2018).
- [65] M. Mariën, J. Haegeman, P. Fendley, and F. Verstraete, Condensation-driven phase transitions in perturbed string nets, Phys. Rev. B 96, 155127 (2017).
- [66] R.-Z. Huang, L. Zhang, A. M. Läuchli, J. Haegeman, F. Verstraete, and L. Vanderstraeten, Emergent conformal boundaries from finite-entanglement scaling in matrix product states, Phys. Rev. Lett. **132**, 086503 (2024).
- [67] T. Vieijra, C. Casert, J. Nys, W. De Neve, J. Haegeman, J. Ryckebusch, and F. Verstraete, Restricted boltzmann machines for quantum states with non-abelian or anyonic symmetries, Phys. Rev. Lett. **124**, 097201 (2020).
- [68] F. Verstraete, V. Murg, and J. I. Cirac, Matrix product states, projected entangled pair states, and variational renormalization group methods for quantum spin systems, Adv. Phys. 57, 143 (2008).
- [69] K. Inamura and K. Ohmori, Fusion surface models: 2+1d lattice models from fusion 2-categories, SciPost Phys. 16, 143 (2024).
- [70] J. Haegeman, K. Van Acoleyen, N. Schuch, J. I. Cirac, and F. Verstraete, Gauging quantum states: From global to local symmetries in many-body systems, Phys. Rev. X 5, 011024 (2015).
- [71] C. Delcamp, Tensor network approach to electromagnetic duality in (3+1)d topological gauge models, JHEP 08, 149, arXiv:2112.08324 [cond-mat.str-el].
- [72] C. Delcamp and A. Tiwari, Higher categorical symmetries and gauging in two-dimensional spin systems (2023), arXiv:2301.01259 [hep-th].
- [73] B. Bauer, P. Corboz, R. Orús, and M. Troyer, Implementing global abelian symmetries in projected entangled-pair state algorithms, Phys. Rev. B 83, 125106 (2011).
- [74] S. R. White, Density-matrix algorithms for quantum renormalization groups, Phys. Rev. B 48, 10345 (1993).
- [75] F. Verstraete, D. Porras, and J. I. Cirac, Density matrix renormalization group and periodic boundary conditions: A quantum information perspective, Phys. Rev. Lett. 93, 227205 (2004).
- [76] V. Zauner-Stauber, L. Vanderstraeten, M. T. Fishman, F. Verstraete, and J. Haegeman, Variational optimization

algorithms for uniform matrix product states, Phys. Rev. B ${\bf 97},\,045145$ (2018).

- [77] J. Haegeman and F. Verstraete, Diagonalizing transfer matrices and matrix product operators: A medley of exact and computational methods, Annual Review of Condensed Matter Physics 8, 355 (2017).
- [78] M. Müger, From subfactors to categories and topology i: Frobenius algebras in and morita equivalence of tensor categories, Journal of Pure and Applied Algebra 180, 81 (2003).
- [79] A. Molnar, A. R. de Alarcón, J. Garre-Rubio, N. Schuch, J. I. Cirac, and D. Pérez-García, Matrix product operator algebras I: representations of weak Hopf algebras and projected entangled pair states (2022), arXiv:2204.05940 [quant-ph].
- [80] D. Barter, J. C. Bridgeman, and R. Wolf, Computing associators of endomorphism fusion categories, SciPost Phys. 13, 029 (2022).

SEC. A | Generalised DMRG

In order to compute the ground states of the various models appearing in our examples, we implemented a version of the two-site DMRG algorithm for finite chains with open boundary conditions [74]. The DMRG algorithm is a variational algorithm within the subspace of MPSs, which we recall is a class of wavefunctions that implement the area law of gapped phases at the microscopic level, thereby specifically targeting the physical corner of the total Hilbert space. Briefly, the DMRG algorithm proceeds as follows: The wavefunction being a multilinear function of the variables in all local tensors, the global optimisation problem can iteratively be solved using an alternating least squares approach [75]. The two-site version proceeds by blocking two sites together, before solving the combined least-squares problem, and finally using a singular value decomposition in order to split the two-site tensor into two one-site tensors. This two-site version is typically preferred as it allows for an easier redistribution of Schmidt coefficients in the different tensor blocks.

Typically, MPSs are taken to span a subspace of a tensor product space. But, an important feature of the models we consider is that they are typically not defined on a tensor product space. Rather, states need to satisfy some local kinematical constraints in the form of generalised Gauß laws. Thus, we require MPSs that explicitly enforce these kinematical constraints. For our illustrative example, this is accomplished by considering tensors of the form

$$= A = \sum_{\substack{V_1, V_2 \\ d_1, d_2, i}} (V_1, d_1) = A = (V_2, d_2) (V_2, d_2)$$
(A1)

where V_1, V_2 are summed over ψ -projective irreducible

representations of some subgroup $H \subseteq \mathbb{A}_4$, *i* over basis vectors in the space of intertwining maps $V_1 \otimes \underline{3} \to V_2$, while d_1, d_2 label the remaining variational degrees of freedom. Notice that both entanglement degrees of freedom labelled by (V, d) and physical degrees of freedom labelled by $(V_1 \underline{3} V_2, i)$ carry gauge degrees of freedom represented by blue lines, which are shared by neighbouring physical degrees of freedom, as suggested by our graphical notation. Typically, the dimension of the space of intertwining maps $V_1 \otimes \underline{3} \to V_2$ depends on (V_1, V_2) , which is incompatible with a tensor product Hilbert space. By construction, the action of the Hamiltonian (5) leaves the constrained Hilbert space invariant, and thus explicitly preserves the structure of such an MPS. For the case of an on-site symmetry described by a finite group G, the tensors that build up the dual Hamiltonians are be obtained as Racah W-coefficients of linear representations of Gwith (projective) representations of a subgroup H, which can readily be computed by restricting G-representations to the subgroup.

Our algorithm proceeds like the standard two-site DMRG, but all the basic operations are tailored to preserve the kinematical constraints, which amounts to maintaining the structure displayed in eq. (A1). First of all, by using block-diagonal basis transformations on the entanglement space, any MPS of the form (A1) can be brought into the *left canonical form* defined by the condition:

$$\begin{bmatrix} \bar{A}_i \\ \vdots \\ \vdots \\ A_i \\ A_i \end{bmatrix} \stackrel{!}{=} \begin{bmatrix} \vdots \\ \vdots \\ \vdots \\ \vdots \\ A_i \end{bmatrix} .$$
(A2)

In left canonical form, the Schmidt values λ of the reduced density matrix obtained by tracing out all the sites to the right of the site i are then given by the spectrum of ρ_{i} defined as

$$\rho_{i} \equiv \begin{array}{c} \overline{A_{i}} & \overline{A_{i+1}} & \overline{A_{i+2}} \\ \downarrow \\ \overline{A_{i}} & \overline{A_{i+1}} & \overline{A_{i+2}} \\ \overline{A_{i}} & \overline{A_{i+1}} & \overline{A_{i+2}} \\ \end{array} \right)$$
(A3)

As reviewed above, a crucial step of the two-site DMRG algorithm amounts to decomposing the two-site MPS tensor solving the combined least-squares problem into single-site MPS tensors A. Specifically, consider a tensor whose entries are of the form

Keeping the gauge degree of freedom V_2 fixed, one considers the matrix B^{V_2} with entries

$$[B^{V_2}]_{(V_1,d_1i_1)}^{(V_3,d_3i_2)} := [B^{(V_1\underline{3}V_2,i_1)(V_2\underline{3}V_3,i_2)}]_{(V_1,d_1)}^{(V_3,d_3)}.$$
(A5)

Performing a singular value decomposition yields a factorisation of the form $B^{V_2} = M^{V_2} \Sigma^{V_2} (N^{V_2})^{\dagger}$, where Mand N are unitary matrices, while Σ^{V_2} is a diagonal matrix. The entries of Σ are all positive and are referred to as the singular values of B^{V_2} . Truncating the singular values to the desired precision λ_{\min} yields the low-rank approximation

$$[B^{V_2}]_{(V_1,d_1i_1)}^{(V_3,d_3i_2)} \approx \sum_{k=1}^{\lambda_{\min}} [M^{V_2}]_{(V_1,d_1i_1)}^k [\Sigma^{V_2}]_k^k [N^{V_2}]_{(V_3,d_3i_2)}^k^*.$$

Repeating this operation for all ψ -projective representations V_2 of A, one finally defines MPS tensors A_i and A_{i+1}

$$[A_{i}^{(V_{1}\underline{3}V_{2},i_{1})}]_{(V_{1},d_{1})}^{(V_{2},k)} := [M^{V_{2}}]_{(V_{1},d_{1}i_{1})}^{k}$$

and
$$[A_{i+1}^{(V_{2}\underline{3}V_{3},i_{2})}]_{(V_{2},k)}^{(V_{3},d_{3})} := [\Sigma^{V_{2}}]_{k}^{k} [N^{V_{2}}]_{(V_{3},d_{3}i_{2})}^{k},$$
 (A6)

respectively, so that (A4) is approximated by

$$(V_{1}\underline{3}V_{2},i_{1}) (V_{2}\underline{3}V_{3},i_{2}) \\ | \underbrace{4}_{3} \\ | \underbrace{3}_{1} \\ | \underbrace{4}_{3} \\ | \underbrace{4}_$$

We can then repeat the same steps for the sites i + 1 and i + 2, and keep on sweeping from left to right and then from right to left.

In our simulations, we initialise the bulk of the MPS with random matrices of a given dimension per block V_2 , while on the boundary we restrict ourselves to a single one-dimensional block. This corresponds to a choice of boundary condition for the MPO transmuting the Hamiltonian of the model we are simulating into the initial one. In the very specific cases where our algorithm boils to the standard symmetry-preserving DMRG, this amounts to the customary fixing of the total charge sector of the state. Finally, note that our approach is not specific to the two-site DMRG. In particular, uniform MPS algorithms for infinite chains (including VUMPS and 'pulling-through' algorithms) can also be implemented [76, 77].

SEC. B | Symmetries in tensor networks

Consider a Hamiltonian with an ordinary symmetry encoded into a (finite) group G. Suppose the Hamiltonian is in the G symmetric phase. We claim that the optimal way of simulating this phase via the DMRG algorithm is to simulate the dual model obtained by gauging G, before acting with the MPO transmuting the Hamiltonian of the dual model into the initial one. As commented in the main text, the building blocks of the MPO for this duality evaluate to Clebsch–Gordan coefficients. More precisely, in the case of the model (2), where the local Hilbert space is spanned by $|\underline{3}, v\rangle$, v = 1, ..., 3, we have the following identification:

whereby the MPO tensor acts on the space of intertwining maps $V_1 \otimes \underline{3} \to V_2$. Furthermore, we commented in the main text that in this very specific case, our procedure amounts to directly simulating the initial phase using symmetry-preserving tensor networks. Generally, symmetry-preserving tensor network algorithms exploit a specific expression for the tensors that explicitly enforces the symmetry. Concretely, consider an MPS in the Hilbert space of the model (5). Generically, entanglement degrees of freedom of the MPS tensor live in a vector space U. Let us suppose that the MPS tensors are invariant under the action of \mathbb{A}_4 . As already exploited in the main text, the Wigner-Eckart theorem stipulates that the tensors are expressible as linear combination of Clebsch–Gordan coefficients. More concretely, since the vector space U is equipped with an \mathbb{A}_4 action, it can be decomposed into irreducible representations, i.e. $U \cong \bigoplus_V \langle U, V \rangle V$, where the direct sum is over irreducible representations of \mathbb{A}_4 and $\langle V, U \rangle \in \mathbb{Z}_{\geq 0}$. It follows that we can decompose $u = 1, \ldots, \dim U$ as $(V, v, d) \equiv (V, v) \otimes (V, d)$, where $v = 1, \ldots, \dim V$ and $d = 1, \ldots, \langle U, V \rangle$. Using this notation, the MPS tensors decompose as follows:

$$u_{1} - u_{2} \equiv \sum_{i}^{v} (V_{1}, d_{1}) - \frac{v_{1}}{i} + V_{2} - v_{2}$$
(B2)

revealing in particular the sparse block structure of the tensors. As per our graphical calculus, the matrices on the r.h.s. labelled by i are in fact of the form (A1). From a symmetric tensor network viewpoint, this decomposition is used to target a specific charge sector of the Hilbert space, thereby reducing computational costs, while explicitly enforcing the symmetry [53-55]. Let us now assume that the MPS is the unique ground state of the \mathbb{A}_4 symmetric phase. In this case, eq. (B2) precisely recovers eq. (8) under the identification (B1) such that the entanglement degrees of freedom of the MPS ground state of the dual model are labelled by pairs (V, d); our algorithm then produces results that agree with symmetrypreserving DMRG. However, when comparing our algorithm to current state-of-the-art implementations, our approach turns out to be practically much simpler since it does not require the conventional implementation of the

recoupling theory for symmetric tensors based on fusion trees.

As commented in the main text, this decomposition is tailored to the symmetric phase, for which entanglement degrees of freedom of the unique ground state transform as linear representations of \mathbb{A}_4 . In contrast, this is clearly not suited to the \mathbb{A}_4 SPT phase, for which entanglement degrees of freedom transform as projective representations of \mathbb{A}_4 . Indeed, it is well known that using standard symmetric tensor networks in an SPT phase is more costly because it forces the edge modes to transform according to a linear representation, which requires additional long-range entanglement.

SEC. C | Mathematical formalism

We sketch here the mathematical formalism underpinning the results presented in the main text. We encourage the reader to consult ref. [33] for detailed definitions. Consider any local one-dimensional quantum lattice model with a generalised symmetry encoded into a fusion category \mathcal{C} . In the presence of such a generalised symmetry, gapped phases are characterised by a choice of (indecomposable) *C-module category*, whose objects label the degenerate ground states of the phase [27, 40]. The same module categories also characterise the different ways to gauge (sub)symmetries of the model. After performing the gauging operation associated with a Cmodule category \mathcal{R} , the dual symmetry of the resulting model is encoded into the so-called Morita dual $\mathcal{C}^*_{\mathcal{R}}$ of \mathcal{C} with respect to \mathcal{R} . The fusion category $\mathcal{C}^*_{\mathcal{R}}$ is defined to be the category $\operatorname{Fun}_{\mathcal{C}}(\mathcal{R},\mathcal{R})$ of \mathcal{C} -module endofunctors of \mathcal{R} [33, 78], the fusion structure being provided by the composition of C-module functors. Crucially, \mathcal{R} is also a $\mathcal{C}^*_{\mathcal{R}}$ module category and we have $(\mathcal{C}^*_{\mathcal{R}})^*_{\mathcal{R}} \simeq \mathcal{C}$. In words, there is always a way to gauge a subsymmetry of $\mathcal{C}^*_{\mathcal{R}}$ so as to recover the initial model.

Let us examine this gauging operation in practice. For conciseness, we focus on nearest-neighbour Hamiltonians, but longer range interactions can be accommodated just as easily. As for the example studied in the main text, it is crucial to write the Hamiltonian in such a way that the generalised symmetry C is manifest. Under some mild mathematical assumptions about the symmetry MPOs [36, 79], it follows from a generalised Wigner– Eckart theorem [37] that any local symmetric operator is expressible in terms of generalised Clebsch-Gordan coefficients. Specifically, given a Hamiltonian of the form $\mathbb{H} = \sum_{i=1}^{L-1} \sum_n \mathbb{h}_{i,n}$, the local operators can always be put in the form

$$\mathbb{h}_{i,n} \equiv \sum_{\{Y\}} \sum_{i,j} h_n(\{Y\}, i, j) \begin{vmatrix} \frac{1}{Y_1} & \frac{1}{Y_2} \\ i & \frac{1}{Y_3} & \frac{1}{Y_4} \end{vmatrix}, \quad (C1)$$

in terms of tensors evaluating to the generalised Clebsch-Gordan coefficients. The graphical notation mimics that of eq. (5) and encodes in particular the fact that the Hilbert space of a model with a generalised symmetry is generically not a tensor product space. More concretely, there is a (possibly not unique) choice of C-module category \mathcal{R} such that local operators can be expressed as (C1) where $\{Y\}$ label objects in $\mathcal{C}^*_{\mathcal{R}}$, and the generalised Clesbch-Gordan coefficients are given by the so-called module associator of \mathcal{R} , as a $\mathcal{C}^*_{\mathcal{R}}$ -module category (c.f. [36, 38, 51] for details).

It follows from the defining axioms of the $C_{\mathcal{R}}^*$ -module category \mathcal{R} that the tensors in eq. (C1) satisfy an analogue to eq. (3):



where the '*F*-symbols' $(F_{Y_4}^{Y_1Y_2Y_3})_{Y_5,ij}^{Y_6,kl} \in \mathbb{C}$ enters the definition of the fusion category $\mathcal{C}^*_{\mathcal{R}}$. Together with the complex coefficients (I_{-}) plex coefficients $\{h_n\}_n$, these *F*-symbols govern the algebra generated by local symmetric operators $\{h_{i,n}\}_{i,n}$. Within this framework, performing a gauging operation simply amounts to picking a different $\mathcal{C}^*_{\mathcal{R}}$ -module category \mathcal{R}' . This means replacing the tensors in eq. (C1) by a new set of tensors that now evaluate to generalised Clebsch–Gordan coefficients given by the module associator of \mathcal{R}' . Crucially, this new set of tensors still satisfy eq. (C2), so the algebra of local symmetric operators generated by (C1) is isomorphic to the initial one. The dual symmetry of the resulting model is then encoded into the fusion category $(\mathcal{C}^*_{\mathcal{R}})^*_{\mathcal{R}'}$. Similarly to the examples discussed in the main text, Hamiltonians associated with different choices of $\mathcal{C}^*_{\mathcal{R}}$ -module categories can be transmuted into each other via an MPO. Mathematically, such an operator is described by an object in the category $\operatorname{\mathsf{Fun}}_{\mathcal{C}^*_{\mathcal{R}}}(\mathcal{R},\mathcal{R}')$ of $\mathcal{C}^*_{\mathcal{R}}$ -module functors from \mathcal{R} to \mathcal{R}' , in such a way that the building blocks of the MPO are provided by the module structure of such a functor [51].

Let us now suppose that the initial model, which is defined with respect to the $C_{\mathcal{R}}^*$ -module category \mathcal{R} , is in the phase associated with the \mathcal{C} -module category \mathcal{P} . Our goal is to find a dual model whose dual symmetry is completely broken in the ground state subspace. To achieve this, we must understand how to relate the phase of the dual model to the phase of the initial model, which is not immediate given that the symmetry structures differ. It follows from $(\mathcal{C}_{\mathcal{R}}^*)_{\mathcal{R}}^* \simeq \mathcal{C}$ that every \mathcal{C} module category \mathcal{P} is of the form $\operatorname{Fun}_{\mathcal{C}_{\mathcal{R}}}(\mathcal{R}, \mathcal{Q})$ for some $\mathcal{C}_{\mathcal{R}}^*$ -module category \mathcal{Q} , the module structure being provided by the composition of $\mathcal{C}_{\mathcal{R}}^*$ -module functors. Conversely, the \mathcal{C} -module category \mathcal{P} uniquely specifies a $\mathcal{C}_{\mathcal{R}}^*$ module category, namely $\operatorname{Fun}_{\mathcal{C}}(\mathcal{R}, \mathcal{P})$. The optimal dual model whose dual symmetry is completely broken in the ground state subspace is obtained whenever the module



Figure 4. The MPO representation of the symmetry C is determined by a choice of module category \mathcal{R} , which in turn fixes a fusion category $C_{\mathcal{R}}^*$ that governs the algebra of symmetric Hamiltonians. The phase of this Hamiltonian is given by a module category \mathcal{P} over C, which determines the fusion category $C_{\mathcal{P}}^*$ describing the quasiparticle excitations. Combining these two module categories we obtain $\mathcal{Q} = \operatorname{Fun}_{\mathcal{C}}(\mathcal{R}, \mathcal{P})$, which describes the entanglement degrees of freedom of the optimal tensor network description of the ground state. Indeed, by dualising and replacing \mathcal{R} by \mathcal{Q} , we end up in the maximal symmetry breaking phase $\mathcal{C}_{\mathcal{P}}^*$ of the dual symmetry $\mathcal{C}_{\mathcal{P}}^*$.

category is given by the symmetry fusion category itself. Given $\mathcal{P} \simeq \mathsf{Fun}_{\mathcal{C}^*_{\mathcal{R}}}(\mathcal{R}, \mathcal{Q})$, this is achieved by choosing \mathcal{R}' to be $\mathsf{Fun}_{\mathcal{C}}(\mathcal{R}, \mathcal{P}) \simeq \mathcal{Q}$. The symmetry of this optimal dual model is then given by $\mathcal{C}^*_{\mathcal{P}}$, which also encodes the quasiparticle excitations.

For many fusion categories of interest, all possible module categories — solutions to eq. (C2) —have been obtained. Importantly, the data of the module functors needed to describe the MPO intertwiners relating the different dual models can be obtained as a representation theoretic problem [37, 80], that numerically can be reduced to a linear algebra problem. This MPO intertwiner allows us to relate the ground state of the optimal dual model defined with respect to the module category Q to the ground state of the initial model defined with respect to the module category \mathcal{R} :

$$\mathcal{R} \stackrel{\uparrow}{\models} \mathcal{R} \stackrel{\uparrow}{\models} \mathcal{R} \stackrel{\uparrow}{\models} \mathcal{R} \stackrel{\uparrow}{\models} \mathcal{R} \stackrel{\uparrow}{\models} \mathcal{R}$$

$$\mathcal{P} \stackrel{::}{=} \underbrace{\begin{array}{c} \mathcal{Q} \\ \mathcal{Q} \\ \vdots \end{array}} \stackrel{\uparrow}{=} \underbrace{\mathcal{Q}} \stackrel{\uparrow}{\models} \underbrace{\mathcal{Q}} \stackrel{\downarrow}{\models} \underbrace{\mathcal{Q}} \stackrel{\downarrow}{\downarrow} \underbrace{\mathcal{Q} \stackrel{\downarrow}{\downarrow} \underbrace{\mathcal{Q} \stackrel{\downarrow}{\downarrow} \underbrace{\mathcal{Q}$$

where the individual tensors of which are determined by the data of the category of module functors $\mathcal{P} = \mathsf{Fun}_{\mathcal{C}}(\mathcal{R},\mathcal{Q})$ [51]. In this way, we obtain the optimal tensor network description of the original ground state. The various fusion categories and the Morita equivalences between them are summarised in the commutative diagrams of figure 4.

To conclude, we revisit our example in light of this general formalism. When dealing with an ordinary (invertible) symmetry \mathbb{A}_4 , the corresponding fusion category \mathcal{C} is the category $\mathsf{Vec}_{\mathbb{A}_4}$ of \mathbb{A}_4 -graded vector spaces. The different ways to gauge subsymmetries of \mathbb{A}_4 are provided by $\mathsf{Vec}_{\mathbb{A}_4}$ -module categories, which are known to be classified by pairs $(A, [\psi])$ as defined in the main text [58]. In particular, choosing A = G and $\psi = 1$ amounts to the (untwisted) gauging of A, and the corresponding Vec_{A4}-module category is equivalent to the category Vec of complex vector spaces. The Morita dual $(Vec_{A_4})^*_{Vec}$ of Vec_{A_4} with respect to Vec can be checked to be equivalent to the category $Rep(A_4)$ of representations of A_4 , in agreement with our results. When writing the Hamiltonian as in eq. (1), we are choosing the $Rep(A_4)$ -module category $\mathcal{R} = Vec$ such that the module associator boils down to the ordinary Clebsch–Gordan coefficients of A_4 . We obtain the various dual models by choosing different $Rep(A_4)$ -module categories. Specifically, the dual model resulting for the ψ -twisted gauging of H is obtained by choosing the module category $\mathcal{R}' = Rep^{\psi}(H)$ of ψ -projective representations of H, and the dual symmetry is encoded into $(Rep(A_4))^*_{Rep^{\psi}(H)}$.

We now suppose that the initial model is in the phase associated with the $\operatorname{Vec}_{\mathbb{A}_4}$ -module category $\operatorname{Fun}_{\operatorname{Rep}(\mathbb{A}_4)}(\operatorname{Vec}, \operatorname{Rep}^{\phi}(K))$. The \mathbb{A}_4 SPT, \mathbb{A}_4 symmetric, and \mathbb{D}_2 symmetric phases considered in the main text are obtained by choosing $\operatorname{Rep}^{\phi}(K)$ to be equal to $\operatorname{Rep}^{\psi}(\mathbb{A}_4)$, $\operatorname{Rep}(\mathbb{A}_4)$, and $\operatorname{Rep}(\mathbb{D}_2)$, respectively. The optimal model is always found to be that given by $\mathcal{R}' = \operatorname{Rep}^{\phi}(K) = \mathcal{Q}$, which amounts to performing a ϕ -twisted gauging of K, as predicted. Let us also shed light on some of the suboptimal simulations: For instance, consider the A_4 -symmetric phase and the dual model labelled by $\mathcal{R}' = \mathsf{Rep}^{\psi}(\mathbb{D}_2)$; the symmetry is found to be $(\mathsf{Rep}(\mathbb{A}_4))^*_{\mathsf{Rep}^{\psi}(\mathbb{D}_2)} \simeq \mathsf{Vec}_{\mathbb{A}_4}$, while the dual phase is that associated with the $Vec_{\mathbb{A}_4}$ -module category $\operatorname{Fun}_{\operatorname{Rep}(\mathbb{A}_4)}(\operatorname{Rep}^{\psi}(\mathbb{D}_2), \operatorname{Rep}(\mathbb{A}_4)) \simeq \operatorname{Vec}$; together with the fact that $\operatorname{Rep}^{\psi}(\mathbb{D}_2)$ is equivalent to Vec as category, this explains why the numerical results were the same for this dual model as for the initial one. In a similar vein, in the \mathbb{A}_4 SPT phase, the dual model obtained by choosing $\mathcal{R}' =$ $\operatorname{\mathsf{Rep}}(\mathbb{D}_2)$ has a $(\operatorname{\mathsf{Rep}}(\mathbb{A}_4))^*_{\operatorname{\mathsf{Rep}}(\mathbb{D}_2)} \simeq \operatorname{\mathsf{Vec}}_{\mathbb{A}_4}$ symmetry; the dual phase is associated with the Vec_{A_4} -module category $\operatorname{Fun}_{\operatorname{Rep}(\mathbb{A}_4)}(\operatorname{Rep}(\mathbb{D}_2), \operatorname{Rep}^{\psi}(\mathbb{A}_4))$, which is equivalent to Vec as a category, meaning that the whole symmetry is preserved, in agreement with our numerical results. Finally, let us examine the \mathbb{D}_2 symmetric phase and the dual model obtained by choosing the $Vec_{\mathbb{A}_4}$ -module category $\mathcal{R}' = \mathsf{Rep}(\mathbb{Z}_3)$; the dual symmetry $(\mathsf{Rep}(\mathbb{A}_4))^*_{\mathsf{Rep}(\mathbb{Z}_3)} \simeq$ $\operatorname{Rep}(\mathbb{A}_4)$ is also fully preserved since the module category over it is $\operatorname{Fun}_{\operatorname{Rep}(\mathbb{A}_4)}(\operatorname{Rep}(\mathbb{Z}_3), \operatorname{Rep}(\mathbb{D}_2))$, which happens to be equivalent to Vec, in agreement with our numerical results.