TIME-ASYMPTOTIC STABILITY OF COMPOSITE WAVES OF DEGENERATE OLEINIK SHOCK AND RAREFACTION FOR NON-CONVEX CONSERVATION LAWS

FEIMIN HUANG, YI WANG, AND JIAN ZHANG

Abstract: We are concerned with the large-time behavior of the solution to one-dimensional (1D) cubic non-convex scalar viscous conservation laws. Due to the inflection point of the cubic non-convex flux, the solution to the corresponding inviscid Riemann problem can be the composite wave of a degenerate Oleinik shock and a rarefaction wave and these two nonlinear waves are always attached together. We give a first proof of the timeasymptotic stability of this composite wave, up to a time-dependent shift to the viscous Oleinik shock, for the viscous equation. The Oleinik shock wave strength can be arbitrarily large. The main difficulty is due to the incompatibility of the time-asymptotic stability proof framework of individual viscous shock by the so-called anti-derivative method and the direct L^2 -energy method to rarefaction wave. Here we develop a new type of *a*-contraction method with suitable weight function and the time-dependent shift to the viscous shock, which is motivated by [9, 12]. Another difficulty comes from that the Oleinik shock and rarefaction wave are always attached together and their wave interactions are very subtle. Therefore, the same time-dependent shift needs to be equipped to both Oleinik shock and rarefaction wave such that the wave interactions can be treated in our stability proof. Time-asymptotically, this shift function grows strictly sub-linear with respect to the time and then the shifted rarefaction wave is equivalent to the original self-similar rarefaction wave.

1. INTRODUCTION

We are concerned with the large-time behavior of the solution to the following onedimensional (1D) cubic non-convex scalar conservation laws

$$\begin{cases} u_t + f(u)_x = \mu u_{xx}, & f(u) = u^3, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}, \\ u(0, x) = u_0(x) \to u_{\pm}, & \text{as } x \to \pm \infty, \end{cases}$$
(1.1)

where $u = u(t, x) \in \mathbb{R}$ is the unknown function, the so-called conserved quantity and $\mu > 0$ is the viscosity coefficient, $u_0(x)$ is the given initial data, and $u_{\pm} \in \mathbb{R}$ are the prescribed far field states. Equation (1.1) is closely related to rotationally invariant hyperbolic waves in viscoelasticity and magnetohydrodynamics [1,3] and can also be derived from nonlinear electromagnetic waves [15,23,25].

HUANG, WANG, AND ZHANG

The large-time asymptotic behavior of the solutions to (1.1) with different far fields u_{\pm} is expected to be determined by the corresponding inviscid Riemann problem

$$\begin{cases} u_t + f(u)_x = 0, \\ u(0, x) = u_0(x) = \begin{cases} u_-, & x < 0, \\ u_+, & x > 0. \end{cases}$$
(1.2)

When the flux f(u) is strictly convex (e.g., the classical Burgers equation), the solution to the Riemann problem (1.2) is either a classical shock (for $u_- > u_+$) or a rarefaction wave (for $u_- < u_+$), and the time-asymptotic stability of the corresponding viscous shock wave and rarefaction wave to its viscous conservation laws are extensively studied and well-established since the pioneer work of Il'in-Oleinik [6]. However, when the flux f(u)is non-convex with inflection points, the wave structure of Riemann problem (1.2) can be much more complicated and consist of multiple wave patterns including Oleinik shock, rarefaction waves and even contact discontinuities. Here Oleinik shock means that the shock wave satisfies the Oleinik entropy condition due to the non-convexity of the flux.

There are rich literatures on the time-asymptotic stability of a single viscous Oleinik shock wave with the non-convex flux f(u) for the scalar viscous conservation laws. When the non-convex flux f(u) has only one inflection point, Kawashima-Matsumura [14] first proved the stability of non-degenerate Oleinik shock wave by the anti-derivative and L^2 weighted energy methods, and then Mei [22] proved the stability of its degenerate Oleinik shock wave. For the general non-convex flux f(u), Matsumura-Nishihara [19] proved the stability of both non-degenerate and degenerate Oleinik shock wave by using a suitably choosing unbounded weighted function. See also Liu [17] for the stability of non-degenerate shock wave by a different weight function and Huang-Xu [5] for the decay rate toward the viscous shock profile. Furthermore, Freistühler-Serre [4] established an interesting L^1 stability theorem and Jones-Gardner-Kapitula [7], Osher-Ralston [24] and Weinberger [26] also investigated the time-asymptotic stability of viscous shock wave by different methods.

For a single rarefaction wave, it can only be formed on the convex (or concave) part of the flux, therefore, the stability of the single rarefaction wave is exactly same as the strictly convex (or concave) case by using the direct L^2 -energy method. Therefore, the time-asymptotic stability of a single Oleinik shock (including both non-degenerate and degenerate cases) or a single rarefaction wave to the non-convex scalar viscous conservation laws is rather satisfactory.

However, as we described before, the wave structure of Riemann problem (1.2) for the non-convex flux f(u) can be much more complicated and consist of multiple wave patterns including both Oleinik shock and rarefaction wave. And there is no result on the time-asymptotic stability of these multiple wave patterns including both Oleinik shock and rarefaction as far as we know. In the present paper, we are concerned with the cubic non-convex flux with one inflection point (i. e., $f(u) = u^3$)

and prove the time-asymptotic stability of the composite wave including both degenerate Oleinik shock and rarefaction wave to scalar viscous equation (1.1).

Compared with the single wave pattern case, the main difficulty here is due to the incompatibility of the time-asymptotic stability proof framework of individual viscous shock by the so-called anti-derivative method and the direct L^2 -energy method to rarefaction wave. Here we develop a new type of *a*-contraction method with suitable weight function and the time-dependent shift to the viscous shock, which is motivated by the recent works [9,12]. In [9], Kang-Vasseur first developed the *a*-contraction method for viscous Burgers equation and proved L^2 -contraction of viscous shock wave up to a time-dependent shift function. Then this idea is used in [8] to obtain L^2 -contraction of small viscous shock wave to the 1D barotropic Navier-Stokes equations by choosing the suitable weight function. L^2 -contraction of shock waves were also extensively studied in [10, 11]. Note that the contraction property in [8, 10] holds for small shock essentially due to the choice of weight function. Here we develop a new *a*-contraction with suitable weight function such that the Oleinik shock wave strength can be arbitrarily large. The main idea is explained in Section 2 and the details can be found in Section 4.

Another motivation is the recent work of Kang-Vasseur-Wang [12] for the time-asymptotic stability of the composite wave of viscous shock and rarefaction wave to 1D compressible barotropic Navier-Stokes equations, see also [13] for the time-asymptotic stability of generic Riemann solutions to the compressible Navier-Stokes-Fourier equations. However, the main difference here from [12] is that the Oleinik shock and rarefaction wave in (1.2)are always attached together, while the viscous shock and rarefaction waves for the compressible Navier-Stokes equations in [12] are separate in two different genuinely nonlinear characteristic fields of the corresponding Euler system. Therefore, the wave interactions between degenerate viscous Oleinik shock and rarefaction wave are more subtle here. Since the time-dependent shift function for the viscous Oleinik shock is essential in our stability proof, we need to equip the same time-dependent shift to rarefaction wave such that two shifted waves are still attached together, otherwise, the shifted shock wave will cross through the rarefaction wave and then the wave interactions will cause substantial difficulties. Time-asymptotically, we can prove that the time-dependent shift function grows strictly sub-linear with respect to the time t and then the shifted rarefaction wave is still equivalent to the original self-similar rarefaction wave.

The rest part of this paper is organized as follows. In Section 2, we first list some properties of the viscous Oleinik shock and rarefaction wave, and then present the new ideas for the *a*-contraction method, and finally state our main result on the time-asymptotic stability of composite wave of degenerate Oleinik shock and rarefaction wave. Section 3 is devoted to the proof of our main result. An approximate rarefaction wave is first constructed and then the weight function and time-dependent shift are well established. Then our stability problem is reformulated to the time-asymptotic stability of the shifted Oleinik shock and shifted approximate rarefaction wave. Finally, the main result can be proved by the continuity arguments based on the local existence of the solution and the uniform-in-time a priori estimates. In Section 4, we establish the uniform-in-time a priori estimates by using the modified a-contraction method and the subtle wave interaction estimates between the Oleinik shock and rarefaction wave. Section 5 is an appendix for some detailed calculations.

2. Preliminaries and main result

In this section, we start with the descriptions of the viscous Oleinik shock wave and the rarefaction wave. Then we explain the main idea of our modified *a*-contraction method and state our main result on the time-asymptotic stability of composite wave of degenerate viscous Oleinik shock and rarefaction for the viscous equation (1.1).

Oleinik shock wave. A shock wave with speed σ for the non-convex conservation law (1.2) with the Riemann initial data u_{\pm} is admissible if it satisfies the Rankine-Hugoniot condition

$$\sigma = \frac{f(u_+) - f(u_-)}{u_+ - u_-},\tag{2.1}$$

and the Oleinik (or Liu) entropy condition

$$\sigma \le \frac{f(u) - f(u_-)}{u - u_-}, \text{ for all } u \text{ between } u_- \text{ and } u_+.$$
(2.2)

One can refer to the recent monagraph [16] for details.

In the present paper, we are concerned with non-classical shock wave, that is the shock wave will cross through the inflection point u = 0 for the non-convex flux $f(u) = u^3$. From (2.1) and (2.2), a shock wave for (1.2) with $f(u) = u^3$ is admissible, or called Oleinik shock, if for $u_- < 0$, the right state u_+ satisfies that

$$u_{-} < u_{+} \le -\frac{u_{-}}{2},\tag{2.3}$$

or for $u_- > 0$,

$$-\frac{u_{-}}{2} \le u_{+} < u_{-}. \tag{2.4}$$

The degenerate case $u_{+} = -\frac{u_{-}}{2}$ is calculated through the relation

$$\sigma = f'(u_+)$$

which means that the shock speed σ is equal to the right characteristic speed $f'(u_+)$. In particular, when $u_+ = -\frac{u_-}{2}$ in (2.3) or (2.4), that is, $f'(u_-) > \sigma = f'(u_+)$, Oleinik shock wave is degenerate at u_+ , and we call it degenerate (or sonic) Oleinik shock, and the other case of (2.3) or (2.4) is called non-degenerate (or subsonic) Oleinik shock wave. For definiteness, we only consider the case (2.3), and the case (2.4) can be treated similarly. Correspondingly, the viscous Oleinik shock is the traveling wave $u^{S}(\xi = x - \sigma t; u_{-}, u_{+})$ defined by the following ODE:

$$\begin{cases} -\sigma(u^S)' + (f(u^S))' = \mu(u^S)'', \quad ' = \frac{d}{d\xi}, \\ u^S(\pm\infty) = u_{\pm}. \end{cases}$$
(2.5)

Now we only consider the degenerate viscous shock case, that is, $u_{+} = -\frac{u_{-}}{2}$ in (2.3). Integrating (2.5)₁ from $(-\infty, \xi]$, it holds that

$$\begin{aligned}
\mu u_{\xi}^{S} &= (u^{S} - u_{-})(u^{S} - u_{+})^{2}, \\
u^{S}(\pm \infty) &= u_{\pm}.
\end{aligned}$$
(2.6)

The existence of the viscous Oleinik shock wave $u^{S}(\xi)$ can be found in [14]. The properties of degenerate viscous Oleinik shock can be summarized in the following lemma, whose proof can be found in [18].

Lemma 2.1. For any states u_{\pm} satisfying $u_{-} < 0$ and $u_{+} = -\frac{u_{-}}{2} > 0$ in (2.3) and let $\delta_{S} := |u_{+} - u_{-}|$ be the wave strength of the degenerate Oleinik shock. Then there exists a constant C > 0 such that the following properties hold true:

$$u_{\xi}^S > 0, \qquad \forall \xi \in \mathbb{R},$$

and

$$\begin{split} |u^{S}(\xi) - u_{-}| &\leq C\delta_{S}e^{-C\delta_{S}^{2}|\xi|}, \quad if \ \xi < 0, \\ |u^{S}(\xi) - u_{+}| &\leq \frac{C\delta_{S}}{1 + C\delta_{S}^{2}|\xi|}, \quad if \ \xi > 0, \\ |u^{S}_{\xi}| &\leq C\delta_{S}^{3}e^{-C\delta_{S}^{2}|\xi|}, \quad if \ \xi < 0, \\ |u^{S}_{\xi}| &\leq \frac{C\delta_{S}^{3}}{(1 + C\delta_{S}^{2}|\xi|)^{2}}, \quad if \ \xi > 0, \end{split}$$

$$\begin{aligned} |u_{\xi}| &\leq \frac{1}{(1+C\delta_{S}^{2}|\xi|)^{2}}, \qquad & ij \ \xi > 0, \\ |u_{\xi\xi}^{S}| &\leq C\delta_{S}^{2}|u_{\xi}^{S}|, \qquad & \forall \xi \in \mathbb{R}. \end{aligned}$$

a-contraction method for a single degenerate Oleinik shock. To explain the main ideas of *a*-contraction method applied to the non-convex conservation laws, for simplicity, we fix $u_{-} = -2$ and $u_{+} = 1$ now and the following arguments still hold true for arbitrary u_{\pm} satisfying admissible conditions (2.1) and (2.2). Then its degenerate viscous shock speed is given by $\sigma = f'(u_{+}) = 3u_{+}^2 = 3$ and the viscous shock profile $U(\xi)$ satisfies

$$-3U' + 3U^2U' = \mu U'', \qquad \lim_{\xi \to -\infty} U(\xi) = -2, \qquad \lim_{\xi \to +\infty} U(\xi) = 1,$$

where $\xi = x - 3t$. By (2.6), we have

$$\mu U'(\xi) = (U+2)(U-1)^2 > 0.$$

HUANG, WANG, AND ZHANG

Note that the viscous shock profile $U(\xi)$ itself is strictly monotone, but the derivative of the characteristic field $\partial_{\xi}(f'(U(\xi))) = 6U(\xi)U'(\xi)$ can change the sign at the inflection point U = 0, that is, the strict compressibility of the Oleinik shock profile does not always hold true.

In order to use *a*-contraction method, the following Poincaré type inequality is crucial in our stability proof.

Lemma 2.2. ([8]) For any $f:[0,1] \to \mathbb{R}$ satisfying $\int_0^1 y(1-y)|f'(y)|^2 dy < +\infty$, it holds that

$$\int_{0}^{1} \left| f - \int_{0}^{1} f \, dy \right|^{2} \, dy \le \frac{1}{2} \int_{0}^{1} y(1-y) |f'(y)|^{2} \, dy, \tag{2.7}$$

that is,

$$\int_0^1 f^2(y)dy - \left(\int_0^1 f(y)dy\right)^2 \le \frac{1}{2}\int_0^1 y(1-y)|f'(y)|^2dy.$$
(2.8)

The shifted Oleinik shock $U^{\mathbf{X}}(t,\xi) := U(\xi + \mathbf{X}(t))$ satisfies the equation

$$\partial_t U^{\mathbf{X}} - \dot{\mathbf{X}}(t) U_{\xi}^{\mathbf{X}} - 3U_{\xi}^{\mathbf{X}} + 3(U^{\mathbf{X}})^2 U_{\xi}^{\mathbf{X}} = \mu U_{\xi\xi}^{\mathbf{X}},$$

where the shift function $\mathbf{X}(t)$ is defined in (2.11). Changing variable $(t, x) \to (t, \xi = x - 3t)$, then $u(t, \xi) := u(t, \xi + 3t)$ satisfies

$$u_t - 3u_\xi + 3u^2 u_\xi = \mu u_{\xi\xi}$$

Then the perturbation $\phi(t,\xi) := u(t,\xi) - U(\xi + \mathbf{X}(t))$ satisfies

$$\phi_t - 3\phi_{\xi} + \dot{\mathbf{X}}(t)U_{\xi}^{\mathbf{X}} + [(\phi + U^{\mathbf{X}})^3 - (U^{\mathbf{X}})^3]_{\xi} = \mu\phi_{\xi\xi}.$$
(2.9)

Due to the degeneracy and the partial compressibility of viscous Oleinik shock and the influence of the inflection point of non-convex flux, the *a*-contraction and L^2 relative entropy method for Burgers equation in [9] can not be utilized directly. We need to use suitable weight in the compressibility region of the shock profile, that is, the negative part of the profile for $U(\xi) < 0$.

We denote that $\xi_1 := (U)^{-1}(0)$ and $\xi_* := (U)^{-1}(\frac{1}{2})$, where $(U)^{-1}$ is inverse function of U. The weight function w(U) is defined as

$$w(U(\xi)) := \begin{cases} \frac{5}{2}(1-U), & \xi \in (-\infty,\xi_1) \iff U \in (-2,0), \\ \frac{5}{2}(1-U)(4U^3+1), & \xi \in [\xi_1,\xi_*) \iff U \in [0,\frac{1}{2}), \\ \frac{15}{8}, & \xi \in [\xi_*,+\infty) \iff U \in [\frac{1}{2},1). \end{cases}$$
(2.10)

It is direct to check that the weight function w(U) is C^2 -smooth and satisfies

$$\frac{15}{8} \le w(U) \le \frac{15}{2}.$$

Note that the weight is added essentially on the negative part of the Oleinik shock, that is, on the region $\xi \in (-\infty, \xi_1)$, or equivalently when $U \in (-2, 0)$. In the region $[\xi_*, +\infty)$, the weight $w \equiv \frac{15}{8}$ and in fact, no need for weight, and in the part $[\xi_1, \xi_*)$, the weight wis chosen such that it is C^2 -smooth. We need to point out that this weight w depends on the wave strength such that arbitrarily large shock wave can be handled.

Choose the time-dependent shift $\mathbf{X}(t)$ as

$$\begin{cases} \dot{\mathbf{X}}(t) = \frac{32}{25} \int_{\mathbb{R}} \phi(t,\xi) w^{\mathbf{X}} U_{\xi}^{\mathbf{X}} d\xi = \frac{32}{25} \int_{\mathbb{R}} \phi^{-\mathbf{X}}(t,\xi) w U_{\xi} d\xi, \\ \mathbf{X}(0) = 0, \end{cases}$$
(2.11)

where $w^{\mathbf{X}} := w(U^{\mathbf{X}}(\xi))$ and $\phi^{-\mathbf{X}}(t,\xi) := \phi(t,\xi - \mathbf{X}(t))$. If we do L^2 relative entropy estimate to (2.9) directly, it holds that

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}}\phi^2 d\xi + \int_{\mathbb{R}}\phi_{\xi}^2 d\xi + \int_{\mathbb{R}}3U^{\mathbf{X}}U_{\xi}^{\mathbf{X}}\phi^2 d\xi + (\cdots) = 0.$$
(2.12)

where and in the sequel (\cdots) represents the higher order terms which can be controlled. Since $U_{\xi}^{\mathbf{X}} > 0$, it can be seen from (2.12) that in the positive region of the Oleinik shock profile $U^{\mathbf{X}}$, that is, when $\xi \in [\xi_*, +\infty)$, L^2 relative entropy estimate in (2.12) is good and no need for the weight and Poincaré inequality, while in the negative region of the Oleinik shock profile $U^{\mathbf{X}}$, that is, when $\xi \in (-\infty, \xi_1]$, L^2 relative entropy estimate in (2.12) has bad sign term and therefore, *a*-contraction method should be used with suitable weight *w* in (2.10) and time-dependent shift $\mathbf{X}(t)$ in (2.11).

Precisely, multiplying (2.9) by $w^{\mathbf{X}}\phi$ with the weight function $w^{\mathbf{X}} := w(U^{\mathbf{X}}(\xi))$ defined in (2.10), we can obtain

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}}\phi^{2}w^{\mathbf{X}}d\xi + \int_{\mathbb{R}}\left(3(w')^{\mathbf{X}} - 3(U^{\mathbf{X}})^{2}(w')^{\mathbf{X}} + 3U^{\mathbf{X}}w^{\mathbf{X}} - \frac{(w'')^{\mathbf{X}}}{2}\mu U_{\xi}^{\mathbf{X}}\right)U_{\xi}^{\mathbf{X}}\phi^{2}d\xi + \int_{\mathbb{R}}\mu\phi_{\xi}^{2}w^{\mathbf{X}}d\xi + \dot{\mathbf{X}}(t)\int_{\mathbb{R}}w^{\mathbf{X}}U_{\xi}^{\mathbf{X}}\phi d\xi + (\cdots) = 0.$$
(2.13)

We claim that if the initial perturbation $||u_0 - U||_{H^1(\mathbb{R})}$ is suitably small, then it holds that

$$\begin{aligned} \mathbf{G}(t) &:= \int_{\mathbb{R}} \mu \phi_{\xi}^{2} w^{\mathbf{X}} d\xi + \dot{\mathbf{X}}(t) \int_{\mathbb{R}} w^{\mathbf{X}} U_{\xi}^{\mathbf{X}} \phi d\xi \\ &+ \int_{\mathbb{R}} \left(3(w')^{\mathbf{X}} - 3(U^{\mathbf{X}})^{2}(w')^{\mathbf{X}} + 3U^{\mathbf{X}} w^{\mathbf{X}} - \frac{(w'')^{\mathbf{X}}}{2} \mu U_{\xi}^{\mathbf{X}} \right) U_{\xi}^{\mathbf{X}} \phi^{2} d\xi \end{aligned}$$

$$\geq \frac{5}{16} \int_{\mathbb{R}} \mu \phi_{\xi}^{2} d\xi + \frac{4}{5} \int_{\mathbb{R}} \phi^{2} U_{\xi}^{\mathbf{X}} d\xi + \frac{25}{64} |\dot{\mathbf{X}}(t)|^{2}.$$

$$(2.14)$$

and the following weighted L^2 -contraction holds true,

$$\frac{d}{dt} \int_{\mathbb{R}} \phi^2 w^{\mathbf{X}} d\xi \le 0.$$

If we first change the variable $\xi \to \xi - \mathbf{X}(t)$ in (2.14), then according to the sign of the Oleinik shock profile $U(\xi)$, we can split $\mathbf{G}(t)$ as follows

$$\mathbf{G}(t) = \mathbf{G}_1(t) + \mathbf{G}_2(t) \tag{2.15}$$

where

$$\mathbf{G}_{1}(t) := \int_{-\infty}^{\xi_{*}} \mu w(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} U_{\xi} \Big(3w' - 3U^{2}w' + 3Uw - \frac{w''}{2} \mu U_{\xi} \Big) d\xi + \frac{8}{25} \Big(\int_{-\infty}^{\xi_{*}} \phi^{-\mathbf{X}} w U_{\xi} d\xi \Big)^{2},$$
(2.16)

and

$$\mathbf{G}_{2}(t) := \int_{\xi_{*}}^{+\infty} \mu w(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + 3 \int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} U w U_{\xi} d\xi + \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w U_{\xi} d\xi - \frac{8}{25} \left(\int_{-\infty}^{\xi_{*}} \phi^{-\mathbf{X}} w U_{\xi} d\xi \right)^{2}.$$
(2.17)

The key point is to handle the term $\mathbf{G}_1(t)$. We can use the modified *a*-contraction method with suitably weighted Poincaré inequality to obtain

$$\mathbf{G}_{1}(t) \geq \frac{5}{16} \int_{-\infty}^{\xi_{*}} \mu(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + \frac{4}{5} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} U_{\xi} d\xi.$$
(2.18)

To prove (2.18), we first let

$$y := \frac{U(\xi) - u_{-}}{U(\xi_{*}) - u_{-}} = \frac{2}{5}(U(\xi) + 2),$$
(2.19)

then we have $\xi \in (-\infty, \xi_*] \iff y \in [0, 1]$, which means that we only use weighted Poincaré inequality on the part $\xi \in (-\infty, \xi_*]$, not on the whole line $\xi \in \mathbb{R}$, the main difference from the *a*-contraction method to the classical Burgers equation in [9]. Since $y_{\xi} = \frac{2}{5}U'(\xi) > 0$, we have from the inverse function theorem and (2.19) that there exists a unique inverse

function $\xi = \xi(y)$. For any fixed t > 0, we can write $\psi(t, y) := \phi^{-\mathbf{X}}(t, \xi)w(U(\xi))$. By using the weighted Poincaré inequality in Lemma 2.2 with $\psi(t, y)$, it holds that

$$\frac{8}{25} \left(\int_{-\infty}^{\xi_*} \phi^{-\mathbf{X}} w U_{\xi} d\xi \right)^2 = 2 \left(\int_0^1 \psi(t, y) dy \right)^2
\geq 2 \int_0^1 \psi^2(t, y) dy - \int_0^1 y(1-y) |\psi_y(t, y)|^2 dy
= -\frac{2}{5} \int_{-\infty}^{\xi_*} \mu(\phi_{\xi}^{-\mathbf{X}})^2 w^2 \frac{\frac{1}{2} - U}{(1-U)^2} d\xi
+ \frac{2}{5} \int_{-\infty}^{\xi_*} (\phi^{-\mathbf{X}})^2 U_{\xi} \left[w w''(\frac{1}{2} - U)(U+2) + 2w^2 - w w'(2U + \frac{3}{2}) \right] d\xi.$$
(2.20)

One can refer to Lemma 4.3 for more details. By (2.16) and (2.20), the estimation of $\mathbf{G}_1(t)$ in (2.18) can be proved. Note that the choosing of $\xi_* := (U)^{-1}(\frac{1}{2})$ is not essential, but crucially we need to choose ξ_* such that $U(\xi_*) > 0$ which means that $U(\xi_*)$ should cross the inflection point 0 of the cubic non-convex flux. Since the term $3 \int_{\xi_*}^{+\infty} (\phi^{-\mathbf{X}})^2 U w U_{\xi} d\xi$ has good sign, with the help of the time-dependent shift $\mathbf{X}(t)$ defined in (2.11), we can get

$$\mathbf{G}_{2}(t) \geq \frac{15}{8} \int_{\xi_{*}}^{+\infty} \mu(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + \left(\frac{45}{16} - \frac{9}{8}\ln 2\right) \int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} U_{\xi} d\xi + \frac{25}{64} |\dot{\mathbf{X}}(t)|^{2}.$$
(2.21)

By (2.18) and (2.21), we can prove (2.14), and then weighted L^2 -contraction holds

$$\frac{d}{dt} \int_{\mathbb{R}} \phi^2 w^{\mathbf{X}} d\xi \le 0.$$

Rarefaction wave. We first recall the Riemann problem for the inviscid Burgers equation:

$$\begin{cases} w_t^r + w^r w_x^r = 0, \\ w^r(0, x) = w_0^r(x) = \begin{cases} w_-, x < 0, \\ w_+, x > 0. \end{cases}$$
(2.22)

If $w_- < w_+$, then the Riemann problem (2.22) has a self-similar rarefaction wave fan solution $w^r(t,x) := w^r(\frac{x}{t}; w_-, w_+)$ defined by

$$w^{r}(t,x) = w^{r}(\frac{x}{t};w_{-},w_{+}) := \begin{cases} w_{-}, & x \le w_{-}t, \\ \frac{x}{t}, & w_{-}t \le x \le w_{+}t, \\ w_{+}, & x \ge w_{+}t. \end{cases}$$
(2.23)

For the Riemann problem of general scalar hyperbolic conservation law (1.2), if f''(u) > 0 on $u \in [u_-, u_+]$ and $f'(u_-) < f'(u_+)$, then the self-similar rarefaction wave solution

 $u^{r}(t,x) := u^{r}\left(\frac{x}{t}; u_{-}, u_{+}\right)$ can be defined explicitly by

$$u^{r}\left(\frac{x}{t}; u_{-}, u_{+}\right) = (\lambda)^{-1} \left(w^{r}\left(\frac{x}{t}; \lambda_{-}, \lambda_{+}\right) \right),$$
(2.24)

where $\lambda(u) := f'(u)$ and $\lambda_{\pm} := \lambda(u_{\pm}) = f'(u_{\pm})$.

Composite wave of degenerate Oleinik shock and rarefaction. Given the Riemann initial data u_{\pm} , we consider the cubic Burgers equation, i.e. flux $f(u) = u^3$. If for $u_- < 0$, the state u_+ satisfies that

$$u_{-} < -\frac{u_{-}}{2} < u_{+}, \tag{2.25}$$

or for $u_- > 0$,

$$u_{+} < -\frac{u_{-}}{2} < u_{-}, \tag{2.26}$$

then the Riemann problem (1.2) is solved by composite wave of a degenerate Oleinik shock connecting u_{-} and $u_m(=-\frac{u_{-}}{2})$ and a rarefaction wave connecting u_m and u_{+} . Correspondingly, the time-asymptotic stability of the Cauchy problem (1.1) is conjectured to be determined by composite wave of a degenerate viscous Oleinik shock and a rarefaction wave.

Without loss of generality, in the sequel we consider the far fied states u_{\pm} satisfy (2.25), and we always denote $u_m := -\frac{u_-}{2}$. Then the large-time behavior of the Cauchy problem (1.1) satisfying (2.25) is determined by a superposition wave:

$$\left(u^{S}(x-\sigma t+\mathbf{X}(t);u_{-},u_{m})+u^{r}(\frac{x}{t};u_{m},u_{+})-u_{m}\right),$$
(2.27)

where $u^{S}(x - \sigma t + \mathbf{X}(t); u_{-}, u_{m})$ is defined in (2.5) and $u^{r}(\frac{x}{t}; u_{m}, u_{+})$ is defined in (2.24). **Main result.** Now we can state our main result as follows.

Theorem 2.1. If the far-fields states u_{\pm} in (1.1) satisfy (2.25), then there exist positive constants δ^* , ϵ^* such that if the initial data u_0 satisfies

$$\|u_{0}(\cdot) - u^{S}(\cdot; u_{-}, u_{m})\|_{L^{2}(\mathbb{R}_{-})} + \|u_{0}(\cdot) - (u^{S}(\cdot; u_{-}, u_{m}) + u_{+} - u_{m})\|_{L^{2}(\mathbb{R}_{+})} + \|u_{0x}(\cdot) - u^{S}_{x}(\cdot; u_{-}, u_{m})\|_{L^{2}(\mathbb{R})} < \epsilon^{*},$$

$$(2.28)$$

with $\mathbb{R}_+ := (0, +\infty) =: -\mathbb{R}_-$, and the rarefaction wave strength δ_R satisfies $\delta_R \leq \delta^*$, then the Cauchy problem (1.1) admits a unique global-in-time solution u(t, x) satisfying

$$u(t,x) - \left(u^{S}\left(x - \sigma t + \mathbf{X}(t); u_{-}, u_{m}\right) + u^{r}\left(\frac{x}{t}; u_{m}, u_{+}\right) - u_{m}\right) \in C(0, +\infty; H^{1}(\mathbb{R})),$$

$$u_{xx}(t,x) - u^{S}_{xx}\left(x - \sigma t + \mathbf{X}(t); u_{-}, u_{m}\right) \in L^{2}(0, +\infty; L^{2}(\mathbb{R})),$$

(2.29)

where the shift function $\mathbf{X}(t)$ is absolutely continuous. Moreover, the following timeasymptotic stability of composite wave of degenerate Oleinik shock with shift $\mathbf{X}(t)$ and rarefaction wave holds true,

$$\lim_{t \to +\infty} \sup_{x \in \mathbb{R}} \left| u(t,x) - \left(u^{S} \left(x - \sigma t + \mathbf{X}(t); u_{-}, u_{m} \right) + u^{r} \left(\frac{x}{t}; u_{m}, u_{+} \right) - u_{m} \right) \right| = 0, \quad (2.30)$$

and the shift function $\mathbf{X}(t)$ satisfies

$$\lim_{t \to +\infty} |\dot{\mathbf{X}}(t)| = 0.$$
(2.31)

Remark 2.1. The shift $\mathbf{X}(t)$ is proved to satisfy the time-asymptotic behavior (2.31), which implies

$$\lim_{t \to +\infty} \frac{\mathbf{X}(t)}{t} = 0,$$

that is, the shift $\mathbf{X}(t)$ grows at most sub-linearly with respect to the time t, therefore, the shifted viscous Oleinik shock $u^{S}(x - \sigma t + \mathbf{X}(t); u_{-}, u_{m})$ still keeps the original Oleinik shock profile time-asymptotically.

Remark 2.2. Theorem 2.1 is the first time-asymptotic stability result towards the composite wave of Oleinik shock and rarefaction wave to scalar non-convex conservation laws as far as we know, which is proposed as an open problem in the survey paper of Matsumura [18]. Moreover, in Theorem 2.1, Oleinik shock wave strength can be arbitrarily large, while the rarefaction wave strength should be suitably small due to the subtle wave interactions.

Remark 2.3. Note that both the initial value u_0 and the initial Oleinik shock u^S in the initial perturbation (2.28) belong to $H^1(\mathbb{R})$ up to the constant spatial far-fields, while the initial self-similar rarefaction wave is the discontinuous Riemann data and belongs to piece-wise $H^1(\mathbb{R}_{\pm})$.

3. Reformulation of the problem

In this section, we first construct an approximate rarefaction wave. Then we reformulate the stability problem around the shifted viscous Oleinik shock and shifted approximate rarefaction wave based on the construction of the time-dependent shift $\mathbf{X}(t)$ and suitable weight function $w(u^{S}(\xi))$. Finally, we give the proof of Theorem 2.1 based on the local existence of the solution and the uniform-in-time a priori estimates.

3.1. Construction of approximate rarefaction wave. Note that the self-similar rarefaction wave fan solutions $w^r(t, x)$ and $u^r(t, x)$ in (2.23) and (2.24) are Lipschitz continuous for t > 0 and $x \in \mathbb{R}$. In order to study their time-asymptotic stability for the viscous equation (1.1) with second order derivative, we need to construct their smooth approximations which are equivalent to the corresponding inviscid self-similar rarefaction waves

HUANG, WANG, AND ZHANG

time-asymptotically as in [20]. Precisely, consider the following Cauchy problem of Burgers equation:

$$\begin{cases} w_t^R + w^R w_x^R = 0, \\ w^R(0, x) = w_0^R(x) = \frac{w_+ + w_-}{2} + \frac{w_+ - w_-}{2} \tanh x \to w_{\pm}, \text{ as } x \to \pm \infty. \end{cases}$$
(3.1)

Since $w_- < w_+$ and $(w_0^R)'(x) > 0$ for any $x \in \mathbb{R}$, the Cauchy problem (3.1) has a unique smooth solution $w^R(t, x) := w^R(t, x; w_-, w_+)$ defined by

$$\begin{cases} w^{R}(t,x) = w_{0}^{R}(x_{0}(t,x)) = \frac{w_{+} + w_{-}}{2} + \frac{w_{+} - w_{-}}{2} \tanh(x_{0}(t,x)), \\ x = x_{0}(t,x) + w_{0}^{R}(x_{0}(t,x))t, \end{cases}$$
(3.2)

where $x_0(t, x)$ is the unique intersection point with x-axis of the straight characteristic line of Burgers equation passing through the point (t, x) for t > 0 and $x \in \mathbb{R}$. Correspondingly, a smooth approximate rarefaction wave $u^R(t, x) := u^R(t, x; u_-, u_+)$ of self-similar rarefaction wave $u^r(t, x) := u^r(\frac{x}{t}; u_-, u_+)$ can be constructed by

$$u^{R}(t, x; u_{-}, u_{+}) := (\lambda)^{-1} \big(w^{R}(t, x; \lambda_{-}, \lambda_{+}) \big),$$
(3.3)

with $\lambda(u) := f'(u)$ and $\lambda_{\pm} := \lambda(u_{\pm}) = f'(u_{\pm})$. Then $u^R(t, x)$ in (3.3) satisfies the equation

$$\begin{cases} u_t^R + f(u^R)_x = 0, \\ u^R(0, x) = (\lambda)^{-1} \left(\frac{\lambda_+ + \lambda_-}{2} + \frac{\lambda_+ - \lambda_-}{2} \tanh x \right) \to u_\pm, \text{ as } x \to \pm \infty. \end{cases}$$
(3.4)

Moreover, we have the following lemma.

Lemma 3.1. Assume $u_{-} < u_{+}$ and f''(u) > 0 on $u \in [u_{-}, u_{+}]$ and let $\delta_{R} := |u_{+} - u_{-}|$ be the rarefaction wave strength. Then it holds that

(1). $u_{-} < u^{R}(t, x) < u_{+}, u^{R}_{x}(t, x) > 0$, for $t \ge 0$ and $x \in \mathbb{R}$;

(2). For all $t \ge 1$ and $p \in [1, +\infty]$, there exists a positive constant C_p such that

$$\|u_x^R(t)\|_{L^p(\mathbb{R})} \le C_p \min\left\{\delta_R, \delta_R^{\frac{1}{p}} t^{-1+\frac{1}{p}}\right\} \\ \|u_{xx}^R(t)\|_{L^p(\mathbb{R})} \le C_p \min\left\{\delta_R, t^{-1}\right\};$$

(3). $\lim_{t \to \infty} \sup_{x \in \mathbb{R}} |u^R(t, x) - u^r(\frac{x}{t})| = 0, \text{ that is, the approximate rarefaction wave } u^R(t, x)$

and the inviscid self-similar rarefaction wave fan $u^r(\frac{x}{t})$ are equivalent to each other timeasymptotically for $x \in \mathbb{R}$ uniformly;

(4). For all $t \ge 0$, it holds that

$$|u^{R}(t,x) - u_{+}| \leq C\delta_{R}e^{-2|x-\lambda_{+}t|}, \quad x \geq \lambda_{+}t,$$
$$|u^{R}(t,x) - u_{-}| \leq C\delta_{R}e^{-2|x-\lambda_{-}t|}, \quad x \leq \lambda_{-}t;$$

(5). For all $t \geq 1$ and any $\varepsilon \in (0,1)$, there exists a positive constant C_{ε} such that

$$\begin{aligned} |u^{R}(t,x) - u_{+}| &\leq C_{\varepsilon} \delta_{R}^{\frac{2\varepsilon}{2+\varepsilon}} t^{-1+\varepsilon} e^{-\varepsilon|x-\lambda_{+}t|}, \quad x \geq \lambda_{+}t, \\ |u^{R}(t,x) - u_{-}| &\leq C_{\varepsilon} \delta_{R}^{\frac{2\varepsilon}{2+\varepsilon}} t^{-1+\varepsilon} e^{-\varepsilon|x-\lambda_{-}t|}, \quad x \leq \lambda_{-}t \end{aligned}$$

(6). For all $t \ge 1$ and any $\varepsilon \in (0,1)$, there exists a positive constant C_{ε} such that

$$|u^{R}(t,x) - u^{r}(\frac{x}{t})| \le C_{\varepsilon} \delta_{R}^{\varepsilon} t^{-1+\varepsilon}, \quad \lambda_{-} t \le x \le \lambda_{+} t.$$

Proof. The proof of properties (1)-(3) can be found exactly in [20], and (4) can be proved directly by the solution formula (3.2) and (3.3). Now we prove (5) and (6), whose proofs are motivated by [21], but here we need more detailed information about the rarefaction wave strength δ_R . For simplicity, we only prove (6) and the second inequality in (5), while the first one in (5) can be done similarly.

By (3.3), to prove the second inequality in (5), it is enough to show

$$|w^{R}(t,x) - \lambda_{-}| \le C_{\varepsilon} \delta_{R}^{\frac{2\varepsilon}{2+\varepsilon}} t^{-1+\varepsilon} e^{-\varepsilon|x-\lambda_{-}t|}, \quad \forall t \ge 1 \text{ and } x \le \lambda_{-}t, \tag{3.5}$$

for the solution $w^R(t,x)$ in (3.1) with $w_{\pm} = \lambda_{\pm} := f'(u_{\pm})$. It follows from (3.2) that

$$\frac{\partial x_0}{\partial x} = \frac{1}{1 + (w_0^R)'(x_0)t} > 0, \tag{3.6}$$

which implies that

$$x_0(t,x) \le x_0(t,\lambda_-t), \quad \forall x \le \lambda_-t.$$
 (3.7)

Again by (3.2), $x_0(t, \lambda_- t)$ is given by

$$x_0(t,\lambda_-t) = \lambda_-t - w_0^R \big(x_0(t,\lambda_-t) \big) t < 0, \quad \forall t > 0,$$
(3.8)

which results in

$$\frac{d}{dt}x_0(t,\lambda_-t) = \frac{\lambda_- - w_0^R \left(x_0(t,\lambda_-t)\right)}{1 + (w_0^R)' \left(x_0(t,\lambda_-t)\right)t} < 0, \quad \forall t > 0.$$
(3.9)

Therefore, by (3.8) and (3.9), as $t \to +\infty$,

$$x_0(t,\lambda_-t) \to -\infty. \tag{3.10}$$

By (3.2) and (3.8), we have

$$x_0(t,\lambda_-t) = (\lambda_- - \lambda_+) \frac{e^{2x_0(t,\lambda_-t)}}{e^{2x_0(t,\lambda_-t)} + 1} t < 0, \quad \forall t > 0.$$
(3.11)

Thus, there exist uniform constants $C_1 > C > 0$ such that

$$-C_1 \delta_R t \ e^{2x_0(t,\lambda_-t)} \le x_0(t,\lambda_-t) \le -C \delta_R t \ e^{2x_0(t,\lambda_-t)}, \quad \forall t > 0.$$
(3.12)

Therefore, for any $\varepsilon > 0$, based on the fact $x \leq C_{\varepsilon} e^{\varepsilon x}$, we have

$$\delta_R t \le C_{\varepsilon} e^{-(2+\varepsilon)x_0(t,\lambda_-t)}, \tag{3.13}$$

which implies

$$C_{\varepsilon}(\delta_R t)^{-\frac{2}{2+\varepsilon}} \ge e^{2x_0(t,\lambda_- t)}.$$
(3.14)

Hence by (3.2), (3.8), (3.12) and (3.14), we have $\forall x \leq \lambda_{-}t$,

$$|w^{R}(t,x) - \lambda_{-}| = w_{0}^{R} \left(x_{0}(t,x) \right) - \lambda_{-} \leq w_{0}^{R} \left(x_{0}(t,\lambda_{-}t) \right) - \lambda_{-}$$

$$= -\frac{x_{0}(t,\lambda_{-}t)}{t} \leq C \delta_{R} e^{2x_{0}(t,\lambda_{-}t)} \leq C_{\varepsilon} \delta_{R}^{\frac{\varepsilon}{2+\varepsilon}} t^{-\frac{2}{2+\varepsilon}}.$$
(3.15)

On the other hand, since

$$x = x_0(t, x) + w_0^R(x_0(t, x))t \ge x_0(t, x) + \lambda_- t,$$
(3.16)

we have for $x \leq \lambda_{-}t$,

$$|w^{R}(t,x) - \lambda_{-}| = w_{0}^{R}(x_{0}(t,x)) - \lambda_{-} = (\lambda_{+} - \lambda_{-}) \frac{e^{2x_{0}(t,x)}}{e^{2x_{0}(t,x)} + 1}$$

$$\leq C \delta_{R} e^{2x_{0}(t,x)} \leq C \delta_{R} e^{2(x-\lambda_{-}t)} = C \delta_{R} e^{-2|x-\lambda_{-}t|}.$$
(3.17)

By (3.15) and (3.17), we have for $\forall t \geq 1$ and $\forall x \leq \lambda_{-}t$,

$$w^{R}(t,x) - \lambda_{-}| = |w^{R}(t,x) - \lambda_{-}|^{1-\frac{\varepsilon}{2}}|w^{R}(t,x) - \lambda_{-}|^{\frac{\varepsilon}{2}}$$

$$\leq C_{\varepsilon}\delta_{R}^{\frac{2\varepsilon}{2+\varepsilon}}t^{-\frac{2-\varepsilon}{2+\varepsilon}}e^{-\varepsilon|x-\lambda_{-}t|} \leq C_{\varepsilon}\delta_{R}^{\frac{2\varepsilon}{2+\varepsilon}}t^{-1+\varepsilon}e^{-\varepsilon|x-\lambda_{-}t|}.$$
(3.18)

Now we prove (6). From (3.2), we have

$$\frac{x}{t} - w^R(t, x) = \frac{x_0(t, x)}{t}.$$
(3.19)

Then, by (3.6), we have for $\lambda_{-}t \leq x \leq \lambda_{+}t$,

$$\frac{x_0(t,\lambda_-t)}{t} \le \frac{x}{t} - w^R(t,x) \le \frac{x_0(t,\lambda_+t)}{t}.$$
(3.20)

Thus, we have for $\lambda_{-}t \leq x \leq \lambda_{+}t$,

$$\left|\frac{x}{t} - w^{R}(t,x)\right| \le \max\left\{\left|\frac{x_{0}(t,\lambda_{-}t)}{t}\right|, \left|\frac{x_{0}(t,\lambda_{+}t)}{t}\right|\right\}.$$
(3.21)

By (3.12) and (3.14), we get

$$\left|\frac{x_0(t,\lambda_-t)}{t}\right| \le C\delta_R e^{2x_0(t,\lambda_-t)} \le C_{\varepsilon} \delta_R^{1-\frac{2}{2+\varepsilon}} t^{-\frac{2}{2+\varepsilon}}, \quad \forall \varepsilon > 0, \ \forall t \ge 1.$$
(3.22)

Similarly, we have

$$\left|\frac{x_0(t,\lambda_+t)}{t}\right| \le C_{\varepsilon} \delta_R^{1-\frac{2}{2+\varepsilon}} t^{-\frac{2}{2+\varepsilon}}, \ \forall \varepsilon > 0, \ \forall t \ge 1.$$
(3.23)

Hence, we have $\forall \varepsilon \in (0,1), \forall t \ge 1, \left| u^R(t,x) - u^r(\frac{x}{t}) \right| \le C_{\varepsilon} \delta_R^{\varepsilon} t^{-1+\varepsilon}$ for $\lambda_- t \le x \le \lambda_+ t$. \Box

3.2. Reformulation of the problem. For the Cauchy problem (1.1) with far field states u_{\pm} satisfying (2.25), we want to prove its time-asymptotic ansatz is

$$\bar{u}(t,x) := u^{S} \left(x - \sigma t + \mathbf{X}(t); u_{-}, u_{m} \right) + u^{r} \left(\frac{x}{t}; u_{m}, u_{+} \right) - u_{m},$$
(3.24)

with the time-dependent shift function $\mathbf{X}(t)$ to be determined. Since the self-similar rarefaction wave $u^r(\frac{x}{t}; u_m, u_+)$ is only Lipschitz continuous, in the subsequent stability analysis, the rarefaction wave $u^r(\frac{x}{t}; u_m, u_+)$ is replaced by a shifted approximation rarefaction $u^R(1+t, x+\mathbf{X}(t); u_m, u_+)$ constructed in Section 3.1, with the exactly same time-dependent shift function $\mathbf{X}(t)$ as viscous Oleinik shock, such that the stability ansatz is defined by

$$\tilde{u}(t, x + \mathbf{X}(t)) := u^{S} \left(x - \sigma t + \mathbf{X}(t); u_{-}, u_{m} \right) + u^{R} \left(1 + t, x + \mathbf{X}(t); u_{m}, u_{+} \right) - u_{m}.$$
 (3.25)

For simplification, we denote

$$\tilde{u}^{\mathbf{X}}(t,x) := \tilde{u}(t,x + \mathbf{X}(t)). \tag{3.26}$$

Note that the above defined ansatz is quite different from the case of the time-asymptotic stability of the composite waves including viscous shock wave and rarefaction wave with the time-dependent shift function only for the viscous shock wave to the barotropic compressible Navier-Stokes equations in [12], such that the shifted viscous shock wave and the rarefaction wave are still separate in different characteristic fields and can not pass through each other, while here the Oleinik shock and rarefaction wave are always attached together for scalar non-convex conservation laws. Therefore, the same time-dependent shift function need to be equipped to both Oleinik shock and rarefaction wave such that these two waves can not pass through each other and the wave interactions are still good enough.

For simplification of our analysis, we rewrite the cubic Burgers equation (1.1) through the coordinates transformation $(t, x) \rightarrow (t, \xi = x - \sigma t)$, based on the change of variable associated to the speed of propagation of the shock, $u(t, \xi) := u(t, \xi + \sigma t) = u(t, x)$ satisfies

$$u_t - \sigma u_{\xi} + f(u)_{\xi} = \mu u_{\xi\xi}.$$
 (3.27)

Meanwhile, in the coordinates $(t, \xi = x - \sigma t)$, the ansatz $\tilde{u}(t, x + \mathbf{X}(t)) = \tilde{u}(t, \xi + \sigma t + \mathbf{X}(t)) =: \tilde{u}^{\mathbf{X}}(t, \xi)$ in (3.25) satisfies the equation

$$\tilde{u}_t^{\mathbf{X}} - \sigma \tilde{u}_{\xi}^{\mathbf{X}} + f(\tilde{u}^{\mathbf{X}})_{\xi} - \mu \tilde{u}_{\xi\xi}^{\mathbf{X}} = F^{\mathbf{X}} + \dot{\mathbf{X}}(t)(u^R)_{\xi}^{\mathbf{X}} + \dot{\mathbf{X}}(t) \ (u^S)_{\xi}^{\mathbf{X}}, \tag{3.28}$$

where and in the sequel $(u^R)^{\mathbf{X}} := u^R(1+t, \xi + \sigma t + \mathbf{X}(t)), (u^S)^{\mathbf{X}} := u^S(\xi + \mathbf{X}(t))$ and the error term $F^{\mathbf{X}}$ is given by

$$F^{\mathbf{X}} = \left[f(\tilde{u}^{\mathbf{X}}) - f((u^{R})^{\mathbf{X}}) - f((u^{S})^{\mathbf{X}})\right]_{\xi} - \mu(u^{R})_{\xi\xi}^{\mathbf{X}}$$

$$= \left[\left(f'(\tilde{u}^{\mathbf{X}}) - f'((u^{S})^{\mathbf{X}})\right)(u^{S})_{\xi}^{\mathbf{X}} + \left(f'(\tilde{u}^{\mathbf{X}}) - f'((u^{R})^{\mathbf{X}})\right)(u^{R})_{\xi}^{\mathbf{X}}\right] - \mu(u^{R})_{\xi\xi}^{\mathbf{X}} \qquad (3.29)$$

$$:= F_{1}^{\mathbf{X}} + F_{2}^{\mathbf{X}},$$

with $F_1^{\mathbf{X}}$ being the wave interactions and $F_2^{\mathbf{X}}$ the error due to the inviscid rarefaction wave profile. For simplification, we also denote

$$F := \left[\left(f'(\tilde{u}) - f'(u^S) \right) u_{\xi}^S + \left(f'(\tilde{u}) - f'(u^R) \right) u_{\xi}^R \right] - \mu u_{\xi\xi}^R, \tag{3.30}$$

i.e. $F^{\mathbf{X}}(t,\xi) = F(t,\xi + \mathbf{X}(t)).$

We set the perturbation

$$\phi(t,\xi) := u(t,\xi) - \tilde{u}^{\mathbf{X}}(t,\xi).$$
(3.31)

Then by (3.27) and (3.28), the perturbation ϕ satisfies the equation

$$\begin{cases} \phi_t - \sigma \phi_{\xi} + \left(f(\phi + \tilde{u}^{\mathbf{X}}) - f(\tilde{u}^{\mathbf{X}}) \right)_{\xi} + \dot{\mathbf{X}}(t)((u^S)_{\xi}^{\mathbf{X}} + (u^R)_{\xi}^{\mathbf{X}}) - \mu \phi_{\xi\xi} = -F^{\mathbf{X}}, \\ \phi(0,\xi) = \phi_0(\xi) := u_0(\xi) - \tilde{u}(0,\xi), \end{cases}$$
(3.32)

with $F^{\mathbf{X}}$ defined in (3.29).

3.3. Construction of weight function. By Lemma 2.1: $u_{\xi}^{S} > 0$, then there exists a unique ξ_{1} and a unique ξ_{*} such that $u^{S}(\xi_{1}) = 0$ and $u^{S}(\xi_{*}) = u_{*} := \frac{u_{m}}{2}$, respectively. We define the weight function $w = w(u^{S}(\xi))$ by

$$w(u^{S}(\xi)) := \begin{cases} \frac{5}{2}u_{m}(u_{m} - u^{S}(\xi)), & \xi \in (-\infty, \xi_{1}) \iff u^{S} \in (u_{-}, 0), \\ \frac{5}{2u_{m}^{2}}(u_{m} - u^{S})[4(u^{S})^{3} + u_{m}^{3}], & \xi \in [\xi_{1}, \xi_{*}) \iff u^{S} \in [0, u_{*}), \\ \frac{15}{8}u_{m}^{2}, & \xi \in [\xi_{*}, +\infty) \iff u^{S} \in [u_{*}, u_{m}). \end{cases}$$

$$(3.33)$$

It is easy to check that $w \in C^2(\mathbb{R})$. Notice that

$$\frac{15}{8}u_m^2 \le w < \frac{15}{2}u_m^2,\tag{3.34}$$

$$-\frac{5}{2}u_m \le w' \le 0$$
 and $0 \le w'' \le \frac{15}{2}$, $w' := \frac{dw(u^S)}{du^S}$. (3.35)

This weight function w plays an important role in our stability proof.

3.4. Construction of shift function. Define the shift function $\mathbf{X}(t)$ as a solution to the ODE:

$$\begin{cases} \dot{\mathbf{X}}(t) = \frac{32}{25u_m^2} \int_{\mathbb{R}} \phi(t,\xi) w^{\mathbf{X}}(u^S(\xi)) (u^S)_{\xi}^{\mathbf{X}}(\xi) d\xi, \\ \mathbf{X}(0) = 0, \end{cases}$$
(3.36)

where

$$(u^S)^{\mathbf{X}}(\xi) = u^S(\xi + \mathbf{X}(t))$$
 and $w^{\mathbf{X}}(u^S(\xi)) := w((u^S)^{\mathbf{X}}(\xi)).$

By applying Lemma A.1 in [2] to (3.36), we can prove that (3.36) has a unique absolutely continuous solution defined on any interval in time [0, T]. Moreover, since (1.1) is uniformly

parabolic, the maximum principle [6] implies that $\sup_{t \in [0,T], x \in \mathbb{R}} |u(t,x)| \leq C.$ By (3.31), we have $\sup_{t \in [0,T], \xi \in \mathbb{R}} |\phi(t,\xi)| \leq C.$ Thanks to the facts that $||w||_{C^1(\mathbb{R})} \leq C$ and $||u^S||_{C^1(\mathbb{R})} \leq C$, we have

$$\sup_{\mathbf{X}\in\mathbb{R}} \left| \frac{32}{25u_m^2} \int_{\mathbb{R}} \phi(t,\xi) w^{\mathbf{X}}(u^S(\xi))(u^S)_{\xi}^{\mathbf{X}}(\xi) d\xi \right| \le C,$$
(3.37)

and

$$\sup_{\mathbf{X}\in\mathbb{R}} \left| \partial_{\mathbf{X}} \left(\frac{32}{25u_m^2} \int_{\mathbb{R}} \phi(t,\xi) w^{\mathbf{X}}(u^S(\xi)) (u^S)_{\xi}^{\mathbf{X}}(\xi) d\xi \right) \right| \\
\leq C \int_{\mathbb{R}} \left| \left(w^{\mathbf{X}}(u^S)_{\xi}^{\mathbf{X}} \right)_{\xi} \right| d\xi \leq C ||w||_{C^1(\mathbb{R})} ||u^S||_{C^1(\mathbb{R})} \leq C.$$
(3.38)

Especially, since $|\dot{\mathbf{X}}(t)| \leq C$ by (3.37), we have

$$|\mathbf{X}(t)| \le Ct, \quad \forall t \le T.$$
(3.39)

Proof of Theorem 2.1. In order to prove Theorem 2.1, we shall combine a local existence result together with a priori estimates by continuity arguments.

To state the local existence, we define $\hat{U}(t,x) = u^S(x-\sigma t; u_-, u_m) + u^R(1+t, x; u_m, u_+) - u_m$. In the coordinates transformation $(t,x) \to (t,\xi = x - \sigma t), \ \hat{U}(t,x) = \hat{U}(t,\xi + \sigma t) =: \hat{U}(t,\xi)$ satisfies

$$\hat{U}_t - \sigma \hat{U}_{\xi} + f(\hat{U})_{\xi} - \mu \hat{U}_{\xi\xi} = \hat{F},$$

where

$$\hat{F} = \left(f'(\hat{U}) - f'(u^S)\right)u_{\xi}^S + \left(f'(\hat{U}) - f'(u^R)\right)u_{\xi}^R - \mu u_{\xi\xi}^R.$$

Set perturbation $\hat{\phi}(t,\xi) = u(t,\xi) - \hat{U}(t,\xi)$. By (3.27), we have

$$\hat{\phi}_t - \sigma \hat{\phi}_{\xi} + f(\hat{\phi} + \hat{U})_{\xi} - f(\hat{U})_{\xi} - \mu \hat{\phi}_{\xi\xi} = -\hat{F}.$$

Now, we reformulate the Cauchy problem at general initial time $\tau \ge 0$:

$$\begin{cases} \hat{\phi}_t - \sigma \hat{\phi}_{\xi} + f(\hat{\phi} + \hat{U})_{\xi} - f(\hat{U})_{\xi} - \mu \hat{\phi}_{\xi\xi} = -\hat{F}, \\ \hat{\phi}(\tau, \xi) = \hat{\phi}_{\tau}(\xi), \end{cases} (t,\xi) \in \mathbb{R}_+ \times \mathbb{R}.$$
(3.40)

Proposition 3.1. (Local existence) For any M > 0, there exists a positive constant $t_0 = t_0(M)$ independent of τ such that if $\hat{\phi}_{\tau} \in H^1(\mathbb{R})$ and $\|\hat{\phi}_{\tau}\|_{H^1(\mathbb{R})} \leq M$, then the Cauchy problem (3.40) has a unique solution $\hat{\phi}$ on the time interval $[\tau, \tau + t_0]$ satisfying

$$\begin{cases} \hat{\phi} \in C([\tau, \tau + t_0]; H^1(\mathbb{R})) \cap L^2(\tau, \tau + t_0; H^2(\mathbb{R})), \\ \sup_{t \in [\tau, \tau + t_0]} \| \hat{\phi}(t) \|_{H^1(\mathbb{R})} \le 2M. \end{cases}$$
(3.41)

Proposition 3.2. (A priori estimates) For given $u_{-} < 0$ and the far field states u_{\pm} satisfying (2.25), there exist positive constants $\delta_0, \epsilon_0 > 0$, such that if the Cauchy problem (3.32) has a solution $\phi \in C([0, T]; H^1(\mathbb{R})) \cap L^2(0, T; H^2(\mathbb{R}))$ for some time T > 0 with

$$\mathcal{E}(T) := \sup_{0 \le t \le T} \|\phi\|_{H^1(\mathbb{R})} \le \epsilon_0, \quad \delta_R := |u_+ - u_m| \le \delta_0,$$

then there exists a uniform-in-time positive constant C_0 such that for all $t \in [0, T]$

$$\begin{aligned} \|\phi(t)\|_{H^{1}(\mathbb{R})}^{2} + \int_{0}^{t} \|\phi_{\xi}\|_{H^{1}(\mathbb{R})}^{2} d\tau + \int_{0}^{t} \int_{\mathbb{R}} \phi^{2}((u^{S})_{\xi}^{\mathbf{X}} + (u^{R})_{\xi}^{\mathbf{X}}) d\xi d\tau \\ + \int_{0}^{t} |\dot{\mathbf{X}}(\tau)|^{2} d\tau \leq C_{0}(\|\phi_{0}\|_{H^{1}(\mathbb{R})}^{2} + \delta_{R}^{\frac{8}{33}}). \end{aligned}$$
(3.42)

In addition, by (3.36),

$$|\dot{\mathbf{X}}(t)| \le C \|\phi\|_{L^{\infty}(\mathbb{R})}, \quad \forall t \le T.$$
(3.43)

Proposition 3.1 can be proved in the standard way, and we omit its proof details. The proof of Proposition 3.2 is crucial, and is left in next section.

3.5. The continuity arguments. By the continuity arguments, we can extend the local solution to the global one for all $t \in [0, +\infty)$. Define

$$\epsilon_* := \min\left\{\frac{\epsilon_0}{4}, \sqrt{\frac{\epsilon_0^2}{64C_0} - \delta_R^{\frac{8}{33}}}\right\}, \qquad M := \frac{\epsilon_0}{4},$$

where ϵ_0 and C_0 are given in Proposition 3.2. By the smallness of δ_0 and $\delta_R \leq \delta_0$, we have $\frac{\epsilon_0^2}{64C_0} - \delta_R^{\frac{8}{33}} > 0$. Note that $\|\phi_0\|_{H^1(\mathbb{R})} = \|\hat{\phi}_0\|_{H^1(\mathbb{R})}$. Suppose $\|\hat{\phi}_0\|_{H^1(\mathbb{R})} < \epsilon_* \leq \frac{\epsilon_0}{4} (= M)$, by local existence result in Proposition 3.1, there is a positive constant $T_1 = T_1(M)$ such that a unique solution exists on $[0, T_1]$ and satisfies $\|\hat{\phi}\|_{H^1(\mathbb{R})} \leq 2\|\hat{\phi}_0\|_{H^1(\mathbb{R})} \leq \frac{\epsilon_0}{2}$ for $t \in [0, T_1]$. By Sobolev inequality $\|\hat{\phi}_0\|_{L^{\infty}(\mathbb{R})} \leq \sqrt{2}\|\hat{\phi}_0\|_{H^1(\mathbb{R})}$, we get $\|u_0\|_{L^{\infty}(\mathbb{R})} < C$, by the maximum principle which implies $\|\phi\|_{L^{\infty}(\mathbb{R})} < C$. Hence, (3.37)-(3.39) hold for $t \in [0, T_1]$. Without loss of generality, we can choose T_1 small enough such that for all $t \in [0, T_1]$, we have

$$\begin{aligned} \|u^S(\cdot - \sigma t + \mathbf{X}(t)) + u^R(t, \cdot + \mathbf{X}(t)) - u^S(\cdot - \sigma t) - u^R(t, \cdot)\|_{H^1(\mathbb{R})} \\ &\leq Ct \|u^S_{\xi} + u^R_{\xi}\|_{H^1(\mathbb{R})} \leq \frac{\epsilon_0}{8}. \end{aligned}$$

Therefore, it holds for $t \in [0, T_1]$ that

$$\begin{aligned} \|\phi(t,\cdot)\|_{H^{1}(\mathbb{R})} &\leq \|u^{S}(\cdot - \sigma t + \mathbf{X}(t)) + u^{R}(t,\cdot + \mathbf{X}(t)) - u^{S}(\cdot - \sigma t) - u^{R}(t,\cdot)\|_{H^{1}(\mathbb{R})} \\ &+ \|\hat{\phi}(t,\cdot)\|_{H^{1}(\mathbb{R})} \leq \frac{\epsilon_{0}}{2} + \frac{\epsilon_{0}}{8} < \epsilon_{0}. \end{aligned}$$

Especially, since $\mathbf{X}(t)$ is absolutely continuous, and $\hat{\phi} \in C([0, T_1]; H^1(\mathbb{R}))$, we have $\phi \in C([0, T_1]; H^1(\mathbb{R}))$. We now consider the maximal existence time:

$$T_M := \sup\left\{ t > 0 \Big| \sup_{\tau \in [0,t]} \|\phi(\tau, \cdot)\|_{H^1(\mathbb{R})} \le \epsilon_0 \right\}.$$

If $T_M < \infty$, then the continuity argument implies that $\sup_{\tau \in [0, T_M]} \|\phi(\tau, \cdot)\|_{H^1(\mathbb{R})} = \epsilon_0$. It holds from Theorem 3.2 that

$$\sup_{\in [0,T_M]} \|\phi(\tau,\cdot)\|_{H^1(\mathbb{R})} \le \sqrt{C_0(\|\phi_0\|_{H^1(\mathbb{R})}^2 + \delta_R^{\frac{8}{33}})} \le \frac{\epsilon_0}{8},$$

which contradicts $\sup_{\tau \in [0,T_M]} \|\phi(\tau,\cdot)\|_{H^1(\mathbb{R})} = \epsilon_0$. Thus, $T_M = \infty$, which together with Theorem 3.2 implies

$$\sup_{t>0} \|\phi(t,\cdot)\|_{H^1(\mathbb{R})}^2 + \int_0^\infty \|\phi_{\xi}\|_{H^1(\mathbb{R})}^2 d\tau + \int_0^\infty \int_{\mathbb{R}} \phi^2((u^S)_{\xi}^{\mathbf{X}} + (u^R)_{\xi}^{\mathbf{X}}) d\xi d\tau
+ \int_0^\infty |\dot{\mathbf{X}}(t)|^2 d\tau \le C_0(\|\phi_0\|_{H^1(\mathbb{R})}^2 + \delta_R^{\frac{8}{33}}),$$
(3.44)

and

$$|\dot{\mathbf{X}}(t)| \le C \|\phi(t, \cdot)\|_{L^{\infty}(\mathbb{R})}, \quad \forall t > 0.$$
(3.45)

In addition, since the rarefaction wave u^r is Lipschitz continuous in x for all t > 0 and from Lemma 3.1, we have

$$u(t,x) - \left(u^{S}\left(x - \sigma t + \mathbf{X}(t); u_{-}, u_{m}\right) + u^{r}\left(\frac{x + \mathbf{X}(t)}{t}; u_{m}, u_{+}\right) - u_{m}\right) \in C(0, +\infty; H^{1}(\mathbb{R})).$$
(3.46)

Since $\mathbf{X}(t)$ is absolutely continuous, we have

$$u(t,x) - \left(u^{S}\left(x - \sigma t + \mathbf{X}(t); u_{-}, u_{m}\right) + u^{r}\left(\frac{x}{t}; u_{m}, u_{+}\right) - u_{m}\right) \in C(0, +\infty; H^{1}(\mathbb{R})), \quad (3.47)$$

which is the first line of (2.29).

Since $\phi_{\xi\xi} \in L^2(0, +\infty; L^2(\mathbb{R}))$ by (3.44), and $(u^R)^{\mathbf{X}}_{\xi\xi} \in L^2(0, +\infty; L^2(\mathbb{R}))$ by Lemma 3.1, we have

$$u_{xx}(t,x) - u_{xx}^{S}(x - \sigma t + \mathbf{X}(t); u_{-}, u_{m}) \in L^{2}(0, +\infty; L^{2}(\mathbb{R})),$$

which is the second line of (2.29).

In order to get $\|\phi_0\|_{H^1(\mathbb{R})} \leq \epsilon_*$, we can choose ϵ^* and δ^* which are given in Theorem 2.1 suitably. Using Lemma 3.1, we have

$$\left\| u^{R}(1,x;u_{m},u_{+}) - u_{m} \right\|_{L^{2}(\mathbb{R}_{-})}$$

$$\leq C \delta_{R} \Big(\int_{-\infty}^{0} e^{-4|x-3u_{m}^{2}|} dx \Big)^{\frac{1}{2}} \leq C \delta_{R},$$
(3.48)

$$\begin{aligned} \left\| u^{R}(1,x;u_{m},u_{+}) - u_{+} \right\|_{L^{2}(\mathbb{R}_{+})} \\ &\leq C \Big(\int_{0}^{3u_{+}^{2}} u^{R}(1,x;u_{m},u_{+}) - u_{+})^{2} dx \Big)^{\frac{1}{2}} \\ &+ C \Big(\int_{3u_{+}^{2}}^{+\infty} (u^{R}(1,x;u_{m},u_{+}) - u_{+})^{2} dx \Big)^{\frac{1}{2}} \\ &\leq C \Big(\int_{0}^{3u_{+}^{2}} (u_{m} - u_{+})^{2} dx \Big)^{\frac{1}{2}} + C \delta_{R} \Big(\int_{3u_{+}^{2}}^{+\infty} e^{-4|x-3u_{+}^{2}|} dx \Big)^{\frac{1}{2}} \leq C \delta_{R}, \end{aligned}$$
(3.49)

and

$$\|u_x^R(1,x;u_m,u_+)\|_{L^2(\mathbb{R})} \le C\delta_R.$$
 (3.50)
By (3.48), (3.49) and (3.50), we have

$$\begin{split} \|\phi_0\|_{H^1(\mathbb{R})} &= \left\|u_0 - \left(u^S(x;u_-,u_m) + u^R(1,x;u_m,u_+) - u_m\right)\right\|_{H^1(\mathbb{R})} \\ &= \left\|u_0 - \left(u^S(x;u_-,u_m) + u^R(1,x;u_m,u_+) - u_m\right)\right\|_{L^2(\mathbb{R}_-)} \\ &+ \left\|u_0 - \left(u^S(x;u_-,u_m) + u^R(1,x;u_m,u_+) - u_m\right)\right\|_{L^2(\mathbb{R}_+)} \\ &+ \left\|u_{0x}(\cdot) - u^S_x(\cdot;u_-,u_m) - u^R_x(1,x;u_m,u_+) - u_m\right\|_{L^2(\mathbb{R}_-)} \\ &\leq \left\|u_0 - u^S(x;u_-,u_m)\right\|_{L^2(\mathbb{R}_-)} + \left\|u^R(1,x;u_m,u_+) - u_m\right\|_{L^2(\mathbb{R}_-)} \\ &+ \left\|u_0 - \left(u^S(x;u_-,u_m) + u_+ - u_m\right)\right\|_{L^2(\mathbb{R}_+)} + \left\|u^R(1,x;u_m,u_+) - u_+\right\|_{L^2(\mathbb{R}_+)} \\ &+ \left\|u_{0x}(\cdot) - u^S_x(\cdot;u_-,u_m)\right\|_{L^2(\mathbb{R})} + \left\|u^R_x(1,x;u_m,u_+)\right\|_{L^2(\mathbb{R})} \\ &\leq \left\|u_0 - u^S(x;u_-,u_m)\right\|_{L^2(\mathbb{R}_-)} + \left\|u_0 - \left(u^S(x;u_-,u_m) + u_+ - u_m\right)\right\|_{L^2(\mathbb{R}_+)} \\ &+ \left\|u_{0x}(\cdot) - u^S_x(\cdot;u_-,u_m)\right\|_{L^2(\mathbb{R}_+)} + C_1\delta_R, \end{split}$$

Thus, once we take $\epsilon^* = \frac{\epsilon_*}{2}$ and $\delta^* = \min\left\{\delta_0, \frac{\epsilon_*}{2C_1}\right\}$. Then it holds that $\|\phi_0\|_{H^1(\mathbb{R})} \leq \epsilon_*$.

3.6. Time-asymptotic behaviors. In order to justify the time-asymptotic behaviors (2.30) and (2.31), we set $g(t) := \|\phi_{\xi}(t,\cdot)\|^2$. It is obvious that $g(t) \in L^1(0, +\infty)$. We only need to show that $g'(t) \in L^1(0, +\infty)$. By (3.44), (4.68) and (4.87), we have

$$\begin{split} \int_{0}^{+\infty} |g'(t)| dt &= \int_{0}^{+\infty} \left| \frac{d}{dt} \|\phi_{\xi}\|^{2} \right| dt \\ &\leq C \int_{0}^{+\infty} \left(\|\phi_{\xi}\|^{2}_{H^{1}(\mathbb{R})} + |\dot{\mathbf{X}}(t)|^{2} + \int_{\mathbb{R}} \phi^{2} (u^{S})^{\mathbf{X}}_{\xi} d\xi + \int_{\mathbb{R}} \phi^{2} (u^{R})^{\mathbf{X}}_{\xi} d\xi \right) dt \\ &+ C \int_{0}^{+\infty} \int_{\mathbb{R}} |F^{\mathbf{X}}|^{2} d\xi dt < +\infty. \end{split}$$

Hence

$$\lim_{t \to +\infty} g(t) = \lim_{t \to +\infty} \|\phi_{\xi}(t, \cdot)\|^2 = 0.$$
(3.51)

By Sobolev inequality, we have

$$\lim_{t \to +\infty} \|\phi(t, \cdot)\|_{L^{\infty}(\mathbb{R})} \le \lim_{t \to +\infty} \sqrt{2} \|\phi(t, \cdot)\|^{\frac{1}{2}} \|\phi_{\xi}(t, \cdot)\|^{\frac{1}{2}} = 0.$$
(3.52)

In addition, by (3.43) and (3.52), it holds

 $|\dot{\mathbf{X}}(t)| \le C \|\phi(t, \cdot)\|_{L^{\infty}(\mathbb{R})} \to 0, \quad as \quad t \to +\infty.$ (3.53)

By (2.23), we have $\forall t \geq 1$, there exists a uniform constant C > 0 such that

$$\sup_{x \in \mathbb{R}} |w^r(\frac{x}{1+t}) - w^r(\frac{x}{t})| \le C(1+t)^{-1},$$

which results in

$$\sup_{x \in \mathbb{R}} |u^r(\frac{x}{1+t}) - u^r(\frac{x}{t})| \le C(1+t)^{-1}.$$
(3.54)

From Lemma 3.1 and (3.53)-(3.54), it follows that

$$\sup_{x \in \mathbb{R}} |\bar{u}(t,x) - \tilde{u}^{\mathbf{X}}(t,x)| = \sup_{x \in \mathbb{R}} |u^{r}(\frac{x}{t}; u_{m}, u_{+}) - u^{R}(1 + t, x + \mathbf{X}(t); u_{m}, u_{+})|
\leq \sup_{x \in \mathbb{R}} |u^{r}(\frac{x}{t}; u_{m}, u_{+}) - u^{r}(\frac{x}{1 + t}; u_{m}, u_{+})|
+ \sup_{x \in \mathbb{R}} |u^{r}(\frac{x}{1 + t}; u_{m}, u_{+}) - u^{r}(\frac{x + \mathbf{X}(t)}{1 + t}; u_{m}, u_{+})|
+ \sup_{x \in \mathbb{R}} |u^{r}(\frac{x + \mathbf{X}(t)}{1 + t}; u_{m}, u_{+}) - u^{R}(1 + t, x + \mathbf{X}(t); u_{m}, u_{+})|
\rightarrow 0, (t \rightarrow \infty).$$
(3.55)

By (3.52) and (3.55), we prove (2.30). Hence, the proof of Theorem 2.1 is completed. **Notations.** Throughout this paper, several positive generic constants are denoted by C which is uniform in time. Define

$$u_m := -\frac{u_-}{2}, \quad u_* := \frac{u_m}{2} = -\frac{u_-}{4}.$$

and the far field states u_{\pm} satisfy (2.25) means that $u_{-} < 0 < u_{*} < u_{m} < u_{+}$.

 $L^p(\mathbb{R})(1\leq p\leq +\infty)$ and $H^1(\mathbb{R})$ denote the usual Lebesgue space and Sobolev space in \mathbb{R} with norm

$$\|f\|_{L^{p}(\mathbb{R})} := \left(\int_{\mathbb{R}} |f|^{p} d\xi\right)^{\frac{1}{p}}, \quad \|f\| := \|f\|_{L^{2}(\mathbb{R})}, \quad \|f\|_{H^{1}(\mathbb{R})} := (\|f\|^{2} + \|f_{\xi}\|^{2})^{\frac{1}{2}}.$$

For any function $f: \mathbb{R}^+ \times \mathbb{R} \to \mathbb{R}$ and any time-dependent shift $\mathbf{X}(t)$, we denote

$$f^{\pm \mathbf{X}}(t,\xi) := f(t,\xi \pm \mathbf{X}(t)).$$

HUANG, WANG, AND ZHANG

4. A priori estimates.

In this section, we prove the uniform-in-time a priori estimates in Proposition 3.2. We assume that the Cauchy problem (3.32) has a solution $\phi \in C([0,T]; H^1(\mathbb{R})) \cap L^2(0,T; H^2(\mathbb{R}))$ for some time T > 0. We start from L^2 relative entropy estimate for ϕ .

Proposition 4.1. There exist positive constants $\delta_0, \epsilon_1 > 0$ such that if the rarefaction wave strength $\delta_R := |u_+ - u_m| \leq \delta_0$ and $\mathcal{E}(T) := \sup_{\substack{0 \leq t \leq T \\ 0 \leq t \leq T}} \|\phi(t, \cdot)\|_{H^1(\mathbb{R})} \leq \epsilon_1$, then there exists a uniform constant C > 0 such that for all $t \in [0, T]$

$$\begin{aligned} \|\phi(t,\cdot)\|^{2} + \int_{0}^{t} \|\phi_{\xi}(\tau,\cdot)\|^{2} d\tau + \int_{0}^{t} \int_{\mathbb{R}} \left(\left| (u^{S})_{\xi}^{\mathbf{X}} \right| + \left| (u^{R})_{\xi}^{\mathbf{X}} \right| \right) \phi^{2} d\xi d\tau \\ + \int_{0}^{t} |\dot{\mathbf{X}}(\tau)|^{2} d\tau \leq C(\|\phi_{0}\|^{2} + \delta_{R}^{\frac{8}{33}}). \end{aligned}$$
(4.1)

The proof of Proposition 4.1 can be given by the following Lemmas 4.1 to 4.10.

Lemma 4.1. Let $w(u^{S}(\xi))$ be the weight function defined in (3.33). Then it holds that

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 w d\xi + \dot{\mathbf{X}}(t)\mathbf{Y}(t) + \mathcal{J}^{good}(t) + \mathcal{J}^{bad}(t) = \mathbf{F}(t), \qquad (4.2)$$

where

$$\mathbf{Y}(t) := \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi + \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{R} d\xi - \frac{1}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} w' u_{\xi}^{S} d\xi, \qquad (4.3)$$

$$\mathcal{J}^{good}(t) := \int_{\mathbb{R}} \mu w(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi - 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{S})^{2} w' u_{\xi}^{S} d\xi
+ 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{R} - u_{m}) w u_{\xi}^{S} d\xi + 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u^{R} w u_{\xi}^{R} d\xi
- \frac{3}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{R} - u_{m})^{2} w' u_{\xi}^{S} d\xi - \frac{3}{4} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{4} w' u_{\xi}^{S} d\xi,
\mathcal{J}^{bad}(t) := 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} w u^{S} u_{\xi}^{S} d\xi + 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} w (u^{S} - u_{m}) u_{\xi}^{R} d\xi
- 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{R} - u_{m}) u^{S} w' u_{\xi}^{S} d\xi + \sigma \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} w' u_{\xi}^{S} d\xi
- \frac{1}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} \mu (u_{\xi}^{S})^{2} w'' d\xi + \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} w (u_{\xi}^{S} + u_{\xi}^{R}) d\xi
- 2 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} (u^{S} + u^{R} - u_{m}) w' u_{\xi}^{S} d\xi, \qquad (4.5)$$

and

$$\mathbf{F}(t) := -\int_{\mathbb{R}} F\phi^{-\mathbf{X}} w d\xi.$$
(4.6)

Remark 4.1. Since $w' \leq 0$, $w'' \geq 0$, $u_{\xi}^{S} > 0$, $u_{\xi}^{R} > 0$ and $u^{R} > u_{m}$, $\mathcal{J}^{good}(t)$ consists of good terms, while $\mathcal{J}^{bad}(t)$ consists of bad terms and $\mathbf{F}(t)$ can be controlled by wave interaction estimates in Lemma 4.8 and the decay properties of approximate rarefaction wave.

Proof. Multiplying (3.32) by $w^{\mathbf{X}}\phi$, we can get

$$\left(\frac{1}{2}\phi^{2}w^{\mathbf{X}}\right)_{t} - \frac{1}{2}\dot{\mathbf{X}}(t)\phi^{2}w_{\xi}^{\mathbf{X}} - \sigma\phi_{\xi}\phi w^{\mathbf{X}} + \left(f(\phi + \tilde{u}^{\mathbf{X}}) - f(\tilde{u}^{\mathbf{X}})\right)_{\xi}\phi w^{\mathbf{X}} + \dot{\mathbf{X}}(t)\left[(u^{S})_{\xi}^{\mathbf{X}} + (u^{R})_{\xi}^{\mathbf{X}}\right]\phi w^{\mathbf{X}} - \mu\phi_{\xi\xi}\phi w^{\mathbf{X}} = -F^{\mathbf{X}}\phi w^{\mathbf{X}}.$$
(4.7)

Integrating (4.7) over \mathbb{R} with respect to ξ and changing variable $\xi \to \xi - \mathbf{X}(t)$, we have

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 w d\xi + \dot{\mathbf{X}}(t)\mathbf{Y}(t) + \int_{\mathbb{R}} \left(f(\phi^{-\mathbf{X}} + \tilde{u}) - f(\tilde{u})\right)_{\xi} \phi^{-\mathbf{X}} w d\xi - \sigma \int_{\mathbb{R}} \phi_{\xi}^{-\mathbf{X}} \phi^{-\mathbf{X}} w d\xi - \int_{\mathbb{R}} \mu \phi_{\xi\xi}^{-\mathbf{X}} \phi^{-\mathbf{X}} w d\xi = \mathbf{F}(t),$$

$$(4.8)$$

where $\mathbf{Y}(t)$ and $\mathbf{F}(t)$ are defined in (4.3) and (4.6) respectively. Further calculations yield

$$\int_{\mathbb{R}} -\sigma \phi_{\xi}^{-\mathbf{X}} \phi^{-\mathbf{X}} w d\xi = -\frac{1}{2} \sigma \int_{\mathbb{R}} (\phi^{-\mathbf{X}})_{\xi}^2 w d\xi = \frac{\sigma}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 w' u_{\xi}^S d\xi,$$
(4.9)

and

$$-\int_{\mathbb{R}} \mu \phi_{\xi\xi}^{-\mathbf{X}} \phi^{-\mathbf{X}} w d\xi = \int_{\mathbb{R}} \mu w (\phi_{\xi}^{-\mathbf{X}})^2 d\xi - \frac{1}{2} \int_{\mathbb{R}} \mu (\phi^{-\mathbf{X}})^2 w_{\xi\xi} d\xi$$
$$= \int_{\mathbb{R}} \mu w (\phi_{\xi}^{-\mathbf{X}})^2 d\xi + \frac{\sigma}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 w' u_{\xi}^S d\xi - \frac{3}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 w' (u^S)^2 u_{\xi}^S d\xi \quad (4.10)$$
$$- \frac{1}{2} \int_{\mathbb{R}} \mu (\phi^{-\mathbf{X}})^2 w'' (u_{\xi}^S)^2 d\xi.$$

It holds that

$$\begin{split} &\int_{\mathbb{R}} \left(f(\phi^{-\mathbf{X}} + \tilde{u}) - f(\tilde{u}) \right)_{\xi} \phi^{-\mathbf{X}} w d\xi = -\int_{\mathbb{R}} (\phi^{-\mathbf{X}} w)_{\xi} \left((\phi^{-\mathbf{X}} + \tilde{u})^{3} - \tilde{u}^{3} \right) d\xi \\ &= -\int_{\mathbb{R}} \phi_{\xi}^{-\mathbf{X}} \left((\phi^{-\mathbf{X}})^{3} + 3(\phi^{-\mathbf{X}})^{2} \tilde{u} + 3 \tilde{u}^{2} \phi^{-\mathbf{X}} \right) w d\xi \\ &- \int_{\mathbb{R}} w_{\xi} ((\phi^{-\mathbf{X}})^{4} + 3(\phi^{-\mathbf{X}})^{3} \tilde{u} + 3(\phi^{-\mathbf{X}})^{2} \tilde{u}^{2}) d\xi \\ &= -\int_{\mathbb{R}} \left[\frac{((\phi^{-\mathbf{X}})^{4})_{\xi}}{4} + ((\phi^{-\mathbf{X}})^{3} \tilde{u})_{\xi} - (\phi^{-\mathbf{X}})^{3} \tilde{u}_{\xi} + \left(\frac{3}{2} \tilde{u}^{2} (\phi^{-\mathbf{X}})^{2}\right)_{\xi} - 3 \tilde{u} \tilde{u}_{\xi} (\phi^{-\mathbf{X}})^{2} \right] w d\xi \\ &- \int_{\mathbb{R}} w_{\xi} \left[(\phi^{-\mathbf{X}})^{4} + 3(\phi^{-\mathbf{X}})^{3} \tilde{u} + 3(\phi^{-\mathbf{X}})^{2} \tilde{u}^{2} \right] d\xi \\ &= \int_{\mathbb{R}} 3 \tilde{u} \tilde{u}_{\xi} (\phi^{-\mathbf{X}})^{2} w d\xi - \frac{3}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} \tilde{u} d\xi + \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} \tilde{u}_{\xi} w d\xi \\ &- \int_{\mathbb{R}} \left(\frac{3}{4} (\phi^{-\mathbf{X}})^{4} + 2(\phi^{-\mathbf{X}})^{3} \tilde{u} \right) w_{\xi} d\xi \\ &= 3 \int_{\mathbb{R}} (u^{S} + u^{R} - u_{m}) (u_{\xi}^{S} + u_{\xi}^{R}) (\phi^{-\mathbf{X}})^{2} w d\xi - \frac{3}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{S} + u^{R} - u_{m})^{2} w_{\xi} d\xi \\ &+ \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} (u_{\xi}^{S} + u_{\xi}^{R}) w d\xi - \int_{\mathbb{R}} \left(\frac{3}{4} (\phi^{-\mathbf{X}})^{4} + 2(\phi^{-\mathbf{X}})^{3} (u^{S} + u^{R} - u_{m}) \right) w_{\xi} d\xi \\ &= 3 \int_{\mathbb{R}} (u^{S} + u^{R} - u_{m}) (u_{\xi}^{S} + u_{\xi}^{R}) (\phi^{-\mathbf{X}})^{2} w d\xi - \frac{3}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{R} - u_{m})^{2} w' u_{\xi}^{S} d\xi \\ &- \frac{3}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{S})^{2} w' u_{\xi}^{S} d\xi - 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} (u^{S} + u^{R} - u_{m}) w' u_{\xi}^{S} d\xi \\ &+ \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} w (u_{\xi}^{S} + u_{\xi}^{R}) d\xi - 2 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} (u^{S} + u^{R} - u_{m}) w' u_{\xi}^{S} d\xi \\ &- \frac{3}{4} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{4} w' u_{\xi}^{S} d\xi. \end{split}$$

Hence, by (4.8)-(4.11), we proved Lemma 4.1.

(4.11)

Lemma 4.2. It holds that

$$\dot{\mathbf{X}}(t)\mathbf{Y}(t) + \mathcal{J}^{good}(t) + \mathcal{J}^{bad}(t) = \mathbf{G}^{S}(t) + \mathbf{G}^{R}(t) + \mathbf{G}^{SR}(t) + \mathbf{N}(t) + \mathbf{J}(t), \qquad (4.12)$$

where

$$\mathbf{G}^{S}(t) := \int_{\mathbb{R}} \mu w (\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} \Big(\sigma w' - 3(u^{S})^{2} w' + 3u^{S} w - \frac{w''}{2} \mu u_{\xi}^{S} \Big) d\xi \\
+ \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi - \frac{3}{4} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{4} w' u_{\xi}^{S} d\xi,$$
(4.13)

STABILITY OF DEGENERATE OLEINIK SHOCK AND RAREFACTION WAVES

$$\mathbf{G}^{R}(t) := 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} w u^{R} u_{\xi}^{R} d\xi, \qquad (4.14)$$

$$\mathbf{G}^{SR}(t) := 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 w (u^R - u_m) u_{\xi}^S d\xi - \frac{3}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 (u^R - u_m)^2 w' u_{\xi}^S d\xi, \qquad (4.15)$$

$$\mathbf{N}(t) := \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{R} d\xi - \frac{1}{2} \dot{\mathbf{X}}(t) \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} w' u_{\xi}^{S} d\xi + \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} w (u_{\xi}^{S} + u_{\xi}^{R}) d\xi - 2 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} (u^{S} + u^{R} - u_{m}) w' u_{\xi}^{S} d\xi - 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{R} - u_{m}) u^{S} w' u_{\xi}^{S} d\xi,$$

$$(4.16)$$

and

$$\mathbf{J}(t) := 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 (u^S - u_m) w u_{\xi}^R d\xi.$$
(4.17)

Remark 4.2. First, it is clear that $\mathbf{G}^{R}(t)$ and $\mathbf{G}^{SR}(t)$ are good terms, and $\mathbf{G}^{S}(t)$ are in fact good terms by Lemma 4.5 if we choose the weight function w defined in (3.33) and the time-dependent shift $\mathbf{X}(t)$ defined in (3.36). However, $\mathbf{N}(t)$ are bad terms, which can be controlled by Lemma 4.6 under the assumptions that the perturbations and the rarefaction wave strength are suitably small. While $\mathbf{F}(t)$, $\mathbf{J}(t)$ can be controlled by wave interaction estimates in Lemma 4.8 and the decay properties of the approximate rarefaction wave.

To prove Proposition 4.1, we first estimate $\mathbf{G}^{S}(t)$, $\mathbf{N}(t)$, $\mathbf{J}(t)$ and $\mathbf{F}(t)$ as follows. **Estimation of the** $\mathbf{G}^{S}(t)$. From (3.33), we have $w' = w'' \equiv 0$ for $\xi \in (\xi_{*}, +\infty)$ and then we can split $\mathbf{G}^{\mathbf{S}}(t)$ as follows:

$$\begin{aligned} \mathbf{G}^{\mathbf{S}}(t) &= \int_{-\infty}^{\xi_{*}} \mu w(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} \Big(\sigma w' - 3(u^{S})^{2} w' + 3u^{S} w - \frac{w''}{2} \mu u_{\xi}^{S} \Big) d\xi \\ &+ \int_{\xi_{*}}^{+\infty} \mu w(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + 3 \int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} u^{S} w u_{\xi}^{S} d\xi \\ &+ \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi - \frac{3}{4} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{4} w' u_{\xi}^{S} d\xi \\ &= \int_{-\infty}^{\xi_{*}} \mu w(\phi^{-\mathbf{X}})_{\xi}^{2} d\xi + \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} \Big(\sigma w' - 3(u^{S})^{2} w' + 3u^{S} w - \frac{w''}{2} \mu u_{\xi}^{S} \Big) d\xi \\ &+ \frac{8}{25u_{m}^{2}} \Big(\int_{-\infty}^{\xi_{*}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi \Big)^{2} + \int_{\xi_{*}}^{+\infty} \mu w(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + 3 \int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} u^{S} w u_{\xi}^{S} d\xi \\ &+ \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi - \frac{8}{25u_{m}^{2}} \Big(\int_{-\infty}^{\xi_{*}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi \Big)^{2} - \frac{3}{4} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{4} w' u_{\xi}^{S} d\xi \\ &:= \mathbf{G}_{1}^{S}(t) + \mathbf{G}_{2}^{S}(t), \end{aligned}$$

$$(4.18)$$

where

$$\begin{aligned} \mathbf{G}_{1}^{S}(t) &:= \int_{-\infty}^{\xi_{*}} \mu w (\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} \Big(\sigma w' - 3(u^{S})^{2} w' + 3u^{S} w - \frac{w''}{2} \mu u_{\xi}^{S} \Big) d\xi \\ &+ \frac{8}{25 u_{m}^{2}} \Big(\int_{-\infty}^{\xi_{*}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi \Big)^{2}, \end{aligned}$$

$$(4.19)$$

and

$$\mathbf{G}_{2}^{S}(t) := \int_{\xi_{*}}^{+\infty} \mu w(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + 3 \int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} u^{S} w u_{\xi}^{S} d\xi + \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi - \frac{8}{25u_{m}^{2}} \left(\int_{-\infty}^{\xi_{*}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi \right)^{2} - \frac{3}{4} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{4} w' u_{\xi}^{S} d\xi.$$

$$(4.20)$$

Lemma 4.3. $\mathbf{G}_{1}^{S}(t)$ defined in (4.19) satisfies that

$$\mathbf{G}_{1}^{S}(t) \geq \frac{1}{6} \int_{-\infty}^{\xi_{*}} \mu w(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + \frac{4}{5} u_{m}^{3} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi.$$
(4.21)

Proof. Let $y := \frac{u^S(\xi) - u_-}{u_* - u_-}$, then we have $\xi \in (-\infty, \xi_*] \iff y \in [0, 1]$. Since $y_{\xi} = \frac{2}{5u_m} u_{\xi}^S > 0$, there exists a unique inverse function $\xi = \xi(y)$ by the inverse function theorem. For any fixed t > 0, we can write $\psi(t, y) := \phi^{-\mathbf{X}}(t, \xi) w(u^S(\xi))$. In order to use the weighted Poincaré inequality with $\psi(t, y)$, we first have

$$2\Big(\int_{0}^{1}\psi(t,y)dy\Big)^{2} = 2\Big(\int_{-\infty}^{\xi_{*}}\phi^{-\mathbf{X}}w\frac{u_{\xi}^{S}}{u_{*}-u_{-}}d\xi\Big)^{2} = \frac{8}{25u_{m}^{2}}\Big(\int_{-\infty}^{\xi_{*}}\phi^{-\mathbf{X}}wu_{\xi}^{S}d\xi\Big)^{2}.$$
 (4.22)

Furthermore, we have

$$2\int_{0}^{1}\psi^{2}(t,y)dy = 2\int_{-\infty}^{\xi_{*}}(\phi^{-\mathbf{X}})^{2}w^{2}\frac{u_{\xi}^{S}}{u_{*}-u_{-}}d\xi = \frac{4}{5u_{m}}\int_{-\infty}^{\xi_{*}}(\phi^{-\mathbf{X}})^{2}w^{2}u_{\xi}^{S}d\xi.$$
 (4.23)

and

•

$$\int_{0}^{1} y(1-y) |\psi_{y}(t,y)|^{2} dy = \int_{-\infty}^{\xi_{*}} \left[(\phi^{-\mathbf{X}}w)_{\xi} \right]^{2} \frac{(u^{S}-u_{-})(u_{*}-u^{S})}{u_{\xi}^{S}} d\xi \frac{1}{u_{*}-u_{-}}$$

$$= \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} \left[(\phi^{-\mathbf{X}}w)_{\xi} \right]^{2} \frac{\mu(u_{*}-u^{S})}{(u_{m}-u^{S})^{2}} d\xi$$

$$= \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi_{\xi}^{-\mathbf{X}})^{2} w^{2} \frac{\mu(u_{*}-u^{S})}{(u_{m}-u^{S})^{2}} d\xi + \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} w_{\xi}^{2} \frac{\mu(u_{*}-u^{S})}{(u_{m}-u^{S})^{2}} d\xi$$

$$+ \frac{4}{5u_{m}} \int_{-\infty}^{\xi_{*}} \phi^{-\mathbf{X}} \phi_{\xi}^{-\mathbf{X}} w w_{\xi} \frac{\mu(u_{*}-u^{S})}{(u_{m}-u^{S})^{2}} d\xi := I_{1} + I_{2} + I_{3}.$$

$$(4.24)$$

Noticing that

$$\mu w_{\xi\xi} = \mu [w'(u^S)u^S_{\xi}]_{\xi} = \mu w''(u^S)(u^S_{\xi})^2 + \mu w'(u^S)u^S_{\xi\xi}$$
$$= (u^S - u_-)(u^S - u_m)^2 w''(u^S)u^S_{\xi} + [-3u^2_m + 3(u^S)^2]w'(u^S)u^S_{\xi},$$

we have

$$I_{3} = \frac{4}{5u_{m}} \int_{-\infty}^{\xi_{*}} \phi^{-\mathbf{X}} \phi_{\xi}^{-\mathbf{X}} ww_{\xi} \frac{\mu(u_{*} - u^{S})}{(u_{m} - u^{S})^{2}} d\xi = \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} ((\phi^{-\mathbf{X}})^{2})_{\xi} ww_{\xi} \frac{\mu(u_{*} - u^{S})}{(u_{m} - u^{S})^{2}} d\xi$$

$$= -\frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} \left(ww_{\xi} \frac{\mu(u_{*} - u^{S})}{(u_{m} - u^{S})^{2}} \right)_{\xi} d\xi$$

$$= -I_{2} - \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} ww_{\xi} \left(\frac{\mu(u_{*} - u^{S})}{(u_{m} - u^{S})^{2}} \right)_{\xi} d\xi$$

$$= -I_{2} + \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} ww' u_{\xi}^{S} (2u^{S} + 2u_{m} - u_{*}) d\xi$$

$$- \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} ww' (u_{*} - u^{S}) (u^{S} + 2u_{m}) u_{\xi}^{S} d\xi.$$

$$(4.25)$$

By (4.24) and (4.25), we have

$$\int_{0}^{1} y(1-y) |\psi_{y}(t,y)|^{2} dy = I_{1} + I_{2} + I_{3}$$

$$= \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi_{\xi}^{-\mathbf{X}})^{2} w^{2} \frac{\mu(u_{*}-u^{S})}{(u_{m}-u^{S})^{2}} d\xi$$

$$+ \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} w w' u_{\xi}^{S} (2u^{S} + 2u_{m} - u_{*}) d\xi$$

$$- \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} w w'' (u_{*} - u^{S}) (u^{S} + 2u_{m}) u_{\xi}^{S} d\xi.$$
(4.26)

By weighted Poincaré inequality (2.8), and (4.22), (4.23), (4.26), it holds that

$$\mathbf{G}_{1}^{S}(t) \geq \int_{-\infty}^{\xi_{*}} \mu w(\phi_{\xi}^{-\mathbf{X}})^{2} \left(1 - w \frac{(u_{*} - u^{S})}{(u_{m} - u^{S})^{2}} \frac{2}{5u_{m}}\right) d\xi \\
+ \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} \underbrace{\left(\sigma w' - 3(u^{S})^{2} w' + 3u^{S} w - \frac{w''}{2} \mu u_{\xi}^{S}\right)(u_{*} - u_{-})}_{H_{1}} d\xi \\
+ \frac{2}{5u_{m}} \int_{-\infty}^{\xi_{*}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} \underbrace{w\left(w''(u_{*} - u^{S})(u^{S} + 2u_{m}) + 2w - w'(2u^{S} + 2u_{m} - u_{*})\right)}_{H_{2}} d\xi. \tag{4.27}$$

By (3.33), for $\xi \in (-\infty, \xi_*)$, we have

$$1 - w \frac{(u_* - u^S)}{(u_m - u^S)^2} \frac{2}{5u_m} = \begin{cases} 1 - \frac{\frac{u_m}{2} - u^S}{u_m - u^S} = \frac{\frac{u_m}{2}}{u_m - u^S}, & \xi \in (-\infty, \xi_1), \\ \frac{1}{2u_m^3 (u_m - u^S)} [8(u^S)^4 - 4u_m (u^S)^3 + u_m^4], & \xi \in [\xi_1, \xi_*). \end{cases}$$

$$(4.28)$$

For $\xi \in (-\infty, \xi_1)$, or equivalently, $u^S \in (u_-, 0)$, we have

$$\frac{\frac{u_m}{2}}{u_m - u^S} > \frac{\frac{u_m}{2}}{u_m - u_-} = \frac{\frac{u_m}{2}}{u_m - (-2u_m)} = \frac{1}{6}.$$
(4.29)

For $\xi \in [\xi_1, \xi_*)$, or equivalently, $u^S \in [0, \frac{u_m}{2})$, we have

$$\frac{1}{2u_m^3(u_m - u^S)} [8(u^S)^4 - 4u_m(u^S)^3 + u_m^4] > \frac{1}{4},$$
(4.30)

since if we set $g(u^S) = 8(u^S)^4 - 4u_m(u^S)^3 + u_m^4$, then $g'(u^S) = (u^S)^2(32u^S - 12u_m)$, and it is easy to check that for $u^S \in [0, \frac{u_m}{2})$, $g_{\min}(u^S) = g(\frac{3u_m}{8}) = \frac{485}{512}u_m^4 \approx 0.947u_m^4$ and $\frac{1}{2u_m^3(u_m - u^S)} \ge \frac{1}{2u_m^4}$. By (4.28)-(4.30), we have for $\xi \in (-\infty, \xi_*)$,

$$1 - w \frac{(u_* - u^S)}{(u_m - u^S)^2} \frac{2}{5u_m} > \frac{1}{6}.$$
(4.31)

By the definition of the weight function w defined in (3.33), we can compute

$$H_{1} + H_{2} = \begin{cases} \frac{25}{8}u_{m}^{3}(u_{m} - u^{S}), & \text{for } \xi \in (-\infty, \xi_{1}), \\ -\frac{1800}{u_{m}^{4}}(u^{S})^{8} + \frac{400}{u_{m}^{3}}(u^{S})^{7} + \frac{3950}{u_{m}^{2}}(u^{S})^{6} - \frac{3225}{u_{m}}(u^{S})^{5} + \frac{775}{2}(u^{S})^{4} \\ +325u_{m}(u^{S})^{3} - \frac{75}{2}u_{m}^{2}(u^{S})^{2} - \frac{25}{8}u_{m}^{3}u^{S} + \frac{25}{8}u_{m}^{4}, & \text{for } \xi \in [\xi_{1}, \xi_{*}), \end{cases}$$

$$(4.32)$$

where H_1 and H_2 are defined in (4.27). The detailed calculations can be found in Section 5.

Now we claim that for both $\xi \in (-\infty, \xi_1)$ and $\xi \in [\xi_1, \xi_*)$,

$$H_1 + H_2 > 2u_m^4. (4.33)$$

First, it is obvious that for $\xi \in (-\infty, \xi_1)$, $u^S \in (u_-, 0)$, and then $\frac{25}{8}u_m^3(u_m - u^S) > \frac{25}{8}u_m^4 > 2u_m^4$. For $\xi \in [\xi_1, \xi_*)$, or equivalently, $u^S \in [0, \frac{u_m}{2})$, define

$$h(u^{S}) := h_{1}(u^{S}) - \frac{3225}{u_{m}}(u^{S})^{5} + \frac{775}{2}(u^{S})^{4} + 325u_{m}(u^{S})^{3} - \frac{75}{2}u_{m}^{2}(u^{S})^{2} - \frac{25}{8}u_{m}^{3}u^{S} + \frac{25}{8}u_{m}^{4},$$

where

$$h_1(u^S) := -\frac{1800}{u_m^4} (u^S)^8 + \frac{400}{u_m^3} (u^S)^7 + \frac{3950}{u_m^2} (u^S)^6$$
$$= \frac{(u^S)^6}{u_m^2} \left[-\frac{1800}{u_m^2} (u^S)^2 + \frac{400}{u_m} u^S + 3950 \right] \ge \frac{3700}{u_m^2} (u^S)^6$$

for $u^S \in [0, \frac{u_m}{2})$. Then we have

$$h(u^{S}) \geq \underbrace{\frac{3700}{u_{m}^{2}}(u^{S})^{6} - \frac{3225}{u_{m}}(u^{S})^{5} + 703(u^{S})^{4}}_{h_{2}(u^{S})} - \frac{631}{2}(u^{S})^{4} + 325u_{m}(u^{S})^{3} - \frac{75}{2}u_{m}^{2}(u^{S})^{2} - \frac{25}{8}u_{m}^{3}u^{S} + \frac{25}{8}u_{m}^{4}}_{S}$$

$$\geq -\frac{631}{2}(u^{S})^{4} + 325u_{m}(u^{S})^{3} - \frac{75}{2}u_{m}^{2}(u^{S})^{2} - \frac{25}{8}u_{m}^{3}u^{S} + \frac{25}{8}u_{m}^{4}$$

$$\geq \underbrace{\frac{669}{4}u_{m}(u^{S})^{3} - \frac{75}{2}u_{m}^{2}(u^{S})^{2} - \frac{25}{8}u_{m}^{3}u^{S} + \frac{25}{8}u_{m}^{4}}_{l(u^{S})},$$

where we have used the fact that $h_2(u^S) \ge 0, \forall u^S \in [0, \frac{u_m}{2})$. We can compute that

$$l'(u^S) = \frac{2007}{4}u_m(u^S)^2 - 75u_m^2u^S - \frac{25}{8}u_m^3,$$

which have two roots $u_1 = \frac{150 - 2\sqrt{11896.875}}{2007}u_m < 0$ and $u_2 = \frac{150 + 2\sqrt{11896.875}}{2007}u_m \in (0, u_*)$. Thus for $u^S \in [0, u_2), l'(u^S) < 0$ and for $u^S \in (u_2, u_*), l'(u^S) > 0$. Therefore, we have $l_{\min}(u^S) = l(u_1) > 2u_m^4$, and then the claim in (4.33) is proved. Thus the proof of Lemma 4.3 is completed.

Lemma 4.4. It holds that

$$\mathbf{G}_{2}^{S}(t) \geq \int_{\xi_{*}}^{+\infty} \mu w(\phi_{\xi}^{-\mathbf{X}})^{2} d\xi + (\frac{45}{16} - \frac{9}{8}\ln 2)u_{m}^{3} \int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi + \frac{25}{64}u_{m}^{2}|\dot{\mathbf{X}}(t)|^{2} + \frac{3}{4} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{4} |w'| u_{\xi}^{S} d\xi,$$

$$(4.34)$$

where $\mathbf{G}_{2}^{S}(t)$ is defined by (4.19) and $\dot{\mathbf{X}}(t)$ is defined in (3.36).

Proof. It is sufficient to show that

$$3\int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} u^{S} w u_{\xi}^{S} d\xi + \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi - \frac{8}{25u_{m}^{2}} \Big(\int_{-\infty}^{\xi_{*}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi \Big)^{2} \\ \geq (\frac{45}{16} - \frac{9}{8} \ln 2) u_{m}^{3} \int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi + \frac{25}{64} u_{m}^{2} |\dot{\mathbf{X}}(t)|^{2}.$$

Noticing that $\dot{\mathbf{X}}(t) = \frac{32}{25u_m^2} \int_{\mathbb{R}} \phi w^{\mathbf{X}}(u^S)_{\xi}^{\mathbf{X}} d\xi = \frac{32}{25u_m^2} \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^S d\xi$, we have

$$\frac{1}{2}\dot{\mathbf{X}}(t)\int_{\mathbb{R}}\phi^{-\mathbf{X}}wu_{\xi}^{S}d\xi = \frac{16}{25u_{m}^{2}}\left(\int_{\mathbb{R}}\phi^{-\mathbf{X}}wu_{\xi}^{S}d\xi\right)^{2} \\
= \frac{16}{25u_{m}^{2}}\left(\int_{-\infty}^{\xi_{*}}\phi^{-\mathbf{X}}wu_{\xi}^{S}d\xi\right)^{2} + \frac{16}{25u_{m}^{2}}\left(\int_{\xi_{*}}^{+\infty}\phi^{-\mathbf{X}}wu_{\xi}^{S}d\xi\right)^{2} \\
+ \frac{32}{25u_{m}^{2}}\left(\int_{-\infty}^{\xi_{*}}\phi^{-\mathbf{X}}wu_{\xi}^{S}d\xi\right)\left(\int_{\xi_{*}}^{+\infty}\phi^{-\mathbf{X}}wu_{\xi}^{S}d\xi\right) \\
\geq \frac{8}{25u_{m}^{2}}\left(\int_{-\infty}^{\xi_{*}}\phi^{-\mathbf{X}}wu_{\xi}^{S}d\xi\right)^{2} - \frac{16}{25u_{m}^{2}}\left(\int_{\xi_{*}}^{+\infty}\phi^{-\mathbf{X}}wu_{\xi}^{S}d\xi\right)^{2},$$
(4.35)

where in the last inequality we have used the fact $|ab| \leq \frac{a^2}{4} + b^2$ such that

$$\frac{32}{25u_m^2} \Big(\int_{-\infty}^{\xi_*} \phi^{-\mathbf{X}} w u_{\xi}^S d\xi \Big) \Big(\int_{\xi_*}^{+\infty} \phi^{-\mathbf{X}} w u_{\xi}^S d\xi \Big)$$
$$\geq -\frac{8}{25u_m^2} \Big(\int_{-\infty}^{\xi_*} \phi^{-\mathbf{X}} w u_{\xi}^S d\xi \Big)^2 - \frac{32}{25u_m^2} \Big(\int_{\xi_*}^{+\infty} \phi^{-\mathbf{X}} w u_{\xi}^S d\xi \Big)^2.$$

Hölder inequality yields

$$\left(\int_{\xi_*}^{+\infty} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi \right)^{2} \leq \left(\int_{\xi_*}^{+\infty} 3(\phi^{-\mathbf{X}})^2 u^S w u_{\xi}^{S} d\xi \right) \left(\int_{\xi_*}^{+\infty} \frac{w u_{\xi}^{S}}{3 u^S} d\xi \right)$$

$$= \left(\int_{\xi_*}^{+\infty} 3(\phi^{-\mathbf{X}})^2 u^S w u_{\xi}^{S} d\xi \right) \frac{5 u_m^2}{8} \int_{\xi_*}^{+\infty} \left(\ln |u^S(\xi)| \right)_{\xi} d\xi$$

$$= \left(\int_{\xi_*}^{+\infty} 3(\phi^{-\mathbf{X}})^2 u^S w u_{\xi}^{S} d\xi \right) \frac{5 u_m^2}{8} \left(\ln |u^S(\xi)| \right) \Big|_{\xi_*}^{+\infty}$$

$$= \frac{5 u_m^2 \ln 2}{8} \left(\int_{\xi_*}^{+\infty} 3(\phi^{-\mathbf{X}})^2 u^S w u_{\xi}^{S} d\xi \right).$$

Substituting the above inequality into (4.35) gives that

$$\frac{1}{2}\dot{\mathbf{X}}(t)\int_{\mathbb{R}}\phi^{-\mathbf{X}}wu_{\xi}^{s}d\xi \geq \frac{8}{25u_{m}^{2}} \Big(\int_{-\infty}^{\xi_{*}}\phi^{-\mathbf{X}}wu_{\xi}^{S}d\xi\Big)^{2} - \frac{2}{5}\ln 2\Big(\int_{\xi_{*}}^{+\infty}3(\phi^{-\mathbf{X}})^{2}u^{S}wu_{\xi}^{S}d\xi\Big).$$
(4.36)

From (4.36) and the fact $u^{S}(\xi) \geq \frac{u_{m}}{2}, \forall \xi \geq \xi_{*}$, one has

$$3\int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} u^{S} w u_{\xi}^{S} d\xi + \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi - \frac{8}{25u_{m}^{2}} \Big(\int_{-\infty}^{\xi_{*}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi \Big)^{2}$$

$$\geq 3(1 - \frac{2}{5} \ln 2) \int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} u^{S} w u_{\xi}^{S} d\xi + \frac{1}{2} \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi$$

$$\geq \frac{45}{8} (1 - \frac{2}{5} \ln 2) u_{m}^{2} \int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} u^{S} u_{\xi}^{S} d\xi + \frac{1}{2} \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi$$

$$\geq (\frac{45}{16} - \frac{9}{8} \ln 2) u_{m}^{3} \int_{\xi_{*}}^{+\infty} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi + \frac{25}{64} u_{m}^{2} |\dot{\mathbf{X}}(t)|^{2}.$$
(4.37)
where in the last equality we have used the fact $\frac{1}{2} \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi = \frac{25}{64} u_{m}^{2} |\dot{\mathbf{X}}(t)|^{2}.$
Thus, the proof of Lemma 4.4 is complete.

the proof of Lemma 4.4 is complete.

By Lemma 4.3 and Lemma 4.4, we can get the following lemma.

Lemma 4.5. The following estimation of $\mathbf{G}^{\mathbf{S}}(t)$ holds true.

$$\mathbf{G}^{\mathbf{S}}(t) \geq \frac{5}{16} u_m^2 \int_{\mathbb{R}} \mu(\phi_{\xi}^{-\mathbf{X}})^2 d\xi + \frac{4}{5} u_m^3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 u_{\xi}^S d\xi + \frac{25}{64} u_m^2 |\dot{\mathbf{X}}(t)|^2 + \frac{3}{4} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^4 |w'| u_{\xi}^S d\xi.$$
(4.38)

Estimation of the N(t). The estimation of the N(t) is given in the following lemma.

Lemma 4.6. There exists a positive constant $\epsilon_2 > 0$, such that if $\mathcal{E}(T) := \sup_{0 \le t \le T} \|\phi\|_{H^1(\mathbb{R})} \le \epsilon_2$, then it holds that

$$|\mathbf{N}(t)| \leq (36\sqrt{2} + \frac{35\sqrt{2}}{2})u_m^2 \epsilon_2 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 u_{\xi}^S d\xi + (5\sqrt{2}u_m \delta_R \epsilon_2 + 15u_m^2 \delta_R) \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 u_{\xi}^S d\xi + (\sqrt{2}\epsilon_2 + \frac{48}{5}\delta_R) \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 u_{\xi}^R w d\xi + \frac{25}{128}u_m^2 |\dot{\mathbf{X}}(t)|^2.$$
(4.39)

Proof. We can rewrite $\mathbf{N}(t)$ as:

$$\mathbf{N}(t) = \underbrace{\int_{\mathbb{R}} \left((\phi^{-\mathbf{X}})^{3} u_{\xi}^{S} w - 2(\phi^{-\mathbf{X}})^{3} u^{S} u_{\xi}^{S} w' \right) d\xi}_{\mathbf{N}_{1}(t)} + \underbrace{\int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} u_{\xi}^{R} w d\xi + \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{R} d\xi}_{\mathbf{N}_{3}(t)} - \underbrace{\int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} (u^{R} - u_{m}) w_{\xi} d\xi - 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{R} - u_{m}) u^{S} w_{\xi} d\xi}_{\mathbf{N}_{4}(t)}.$$

$$(4.40)$$

Since $w' \leq 0$ and $u_{\xi}^S > 0$, we have

$$\begin{aligned} |\mathbf{N}_{1}(t)| &= \left| \int_{\mathbb{R}} \left((\phi^{-\mathbf{X}})^{3} u_{\xi}^{S} w - 2(\phi^{-\mathbf{X}})^{3} u^{S} u_{\xi}^{S} w' \right) d\xi \right| \\ &= \left| \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} (w - 2u^{S} w') u_{\xi}^{S} d\xi \right| \\ &\leq \|\phi^{-\mathbf{X}}\|_{L^{\infty}(\mathbb{R})} \left[\|w\|_{L^{\infty}(\mathbb{R})} + 2\|u^{S}\|_{L^{\infty}(\mathbb{R})} \|w'\|_{L^{\infty}(\mathbb{R})} \right] \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi \qquad (4.41) \\ &= \left(\|\phi^{-\mathbf{X}}\|_{L^{\infty}(\mathbb{R})} \frac{35}{2} u_{m}^{2} \right) \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi \\ &\leq \left(\frac{35\sqrt{2}}{2} u_{m}^{2} \|\phi^{-\mathbf{X}}\|_{H^{1}(\mathbb{R})} \right) \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi, \end{aligned}$$

and

$$\begin{aligned} |\mathbf{N}_{2}(t)| &= \left| \dot{\mathbf{X}}(t) \left(-\frac{1}{2} \int_{\mathbb{R}} w' u_{\xi}^{S}(\phi^{-\mathbf{X}})^{2} d\xi \right) \right| \\ &= \left| \frac{32}{25u_{m}^{2}} \left(\int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{S} d\xi \right) \left(-\frac{1}{2} \int_{\mathbb{R}} w' u_{\xi}^{S}(\phi^{-\mathbf{X}})^{2} d\xi \right) \right| \\ &\leq \frac{16}{25u_{m}^{2}} \|\phi^{-\mathbf{X}}\|_{L^{\infty}(\mathbb{R})} \|w\|_{L^{\infty}(\mathbb{R})} \|w'\|_{L^{\infty}(\mathbb{R})} \int_{\mathbb{R}} u_{\xi}^{S} d\xi \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi \end{aligned}$$
(4.42)
$$&\leq \frac{48}{25u_{m}} \|\phi^{-\mathbf{X}}\|_{L^{\infty}(\mathbb{R})} \|w\|_{L^{\infty}(\mathbb{R})} \|w'\|_{L^{\infty}(\mathbb{R})} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi$$
(4.42)
$$&\leq 36\sqrt{2}u_{m}^{2} \|\phi^{-\mathbf{X}}\|_{H^{1}(\mathbb{R})} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi.$$

By ϵ -Cauchy inequality, we have $\forall \epsilon > 0$,

$$\left| \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{R} d\xi \right| \leq \epsilon |\dot{\mathbf{X}}(t)|^{2} + \frac{1}{4\epsilon} \Big(\int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{R} d\xi \Big)^{2}.$$

Taking $\epsilon = \frac{25}{128}u_m^2$ in the above inequality and using Hölder inequality yield

$$\begin{aligned} |\mathbf{N}_{3}(t)| &= \left| \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} w u_{\xi}^{R} d\xi + \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{R} d\xi \right| \\ &\leq \|\phi^{-\mathbf{X}}\|_{L^{\infty}(\mathbb{R})} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} w u_{\xi}^{R} d\xi \\ &+ \frac{25}{128} u_{m}^{2} |\dot{\mathbf{X}}(t)|^{2} + \frac{32}{25u_{m}^{2}} \left(\int_{\mathbb{R}} \phi^{-\mathbf{X}} w u_{\xi}^{R} d\xi \right)^{2} \\ &\leq \|\phi^{-\mathbf{X}}\|_{L^{\infty}(\mathbb{R})} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} w u_{\xi}^{R} d\xi + \frac{25}{128} u_{m}^{2} |\dot{\mathbf{X}}(t)|^{2} \\ &+ \frac{32}{25u_{m}^{2}} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} w u_{\xi}^{R} d\xi \int_{\mathbb{R}} w u_{\xi}^{R} d\xi \\ &\leq \left(\sqrt{2} \|\phi^{-\mathbf{X}}\|_{H^{1}(\mathbb{R})} + \frac{48}{5} \delta_{R} \right) \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{R} w d\xi + \frac{25}{128} u_{m}^{2} |\dot{\mathbf{X}}(t)|^{2}, \end{aligned}$$

$$(4.43)$$

where in the last inequality we have used the fact $\int_{\mathbb{R}} u_{\xi}^R w d\xi \leq \frac{15}{2} u_m^2 \int_{\mathbb{R}} u_{\xi}^R d\xi = \frac{15}{2} u_m^2 \delta_R$. Finally, since $-\frac{5}{2} u_m \leq w' \leq 0$, we have

$$\begin{aligned} |\mathbf{N}_{4}(t)| &= \left| 2 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{3} (u^{R} - u_{m}) w_{\xi} d\xi + 3 \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{R} - u_{m}) u^{S} w_{\xi} d\xi \right| \\ &\leq \left[2\delta_{R} \|\phi^{-\mathbf{X}}\|_{L^{\infty}(\mathbb{R})} \|w'\|_{L^{\infty}(\mathbb{R})} + 3\delta_{R} |u_{-}| \|w'\|_{L^{\infty}(\mathbb{R})} \right] \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi \qquad (4.44) \\ &\leq \left(5\sqrt{2}u_{m}\delta_{R} \|\phi^{-\mathbf{X}}\|_{H^{1}(\mathbb{R})} + 15u_{m}^{2}\delta_{R} \right) \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} u_{\xi}^{S} d\xi. \end{aligned}$$

By (4.40)-(4.44), we proved Lemma 4.6.

HUANG, WANG, AND ZHANG

Integrating (4.2) with respect to t and using Lemmas 4.2-4.6, we have the following lemma.

Lemma 4.7. There exist positive constants $\delta_0 > 0$ and $\epsilon_3 > 0$, such that if $\mathcal{E}(T) := \sup_{0 \le t \le T} \|\phi\|_{H^1(\mathbb{R})} \le \epsilon_3$ and the rarefaction wave strength $\delta_R \le \delta_0$, then it holds that $\forall t \in [0, T]$,

$$\frac{1}{2} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 w d\xi + \frac{5}{16} u_m^2 \int_0^t \int_{\mathbb{R}} \mu(\phi_{\xi}^{-\mathbf{X}})^2 d\xi d\tau + \frac{u_m^3}{2} \int_0^t \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^2 (u_{\xi}^S + u_{\xi}^R) d\xi d\tau + \frac{25}{128} u_m^2 \int_0^t |\dot{\mathbf{X}}(\tau)|^2 d\tau \le \frac{1}{2} \int_{\mathbb{R}} \phi_0^2 w d\xi + \int_0^t \mathbf{F}(\tau) d\tau - \int_0^t \mathbf{J}(\tau) d\tau.$$
(4.45)

It remains to estimate the terms $\int_0^t \mathbf{F}(\tau) d\tau$ and $\int_0^t \mathbf{J}(\tau) d\tau$. First, we handle the wave interaction estimates. Note that the shifted Oleinik shock and the shifted rarefaction wave are always attached together, therefore, their wave interactions are very subtle.

Lemma 4.8. (Wave interaction estimates) There exists generic uniform-in-time constant C > 0, such that

$$\int_{-\infty}^{0} |u^{S} - u_{m}| u_{\xi}^{R} d\xi \le C \delta_{S} \delta_{R}^{\frac{2}{11}} (1+t)^{-\frac{4}{5}}, \qquad (4.46)$$

$$\int_{0}^{+\infty} |u^{S} - u_{m}| u_{\xi}^{R} d\xi \le C \delta_{S}^{-\frac{3}{5}} \delta_{R}^{\frac{1}{5}} (1+t)^{-\frac{4}{5}}, \qquad (4.47)$$

$$\int_{-\infty}^{0} |u^{R} - u_{m}| u_{\xi}^{S} d\xi \le C \delta_{S} \delta_{R}^{\frac{2}{11}} (1+t)^{-\frac{4}{5}}, \qquad (4.48)$$

$$\int_{0}^{[f'(u_{+})-\sigma](t+1)} |u^{R} - u^{r}| u_{\xi}^{S} d\xi \le C \delta_{R}^{\frac{1}{5}} \delta_{S} (1+t)^{-\frac{4}{5}}, \qquad (4.49)$$

$$\int_{0}^{[f'(u_{+})-\sigma](t+1)} |u^{r} - u_{m}| u_{\xi}^{S} d\xi \leq C \delta_{R}^{\frac{1}{5}} \delta_{S}^{-\frac{3}{5}} (1+t)^{-\frac{4}{5}} \ln^{\frac{4}{5}} (1+C \delta_{S}^{2} \delta_{R} t),$$
(4.50)

$$\int_{[f'(u_{+})-\sigma](t+1)}^{+\infty} |u^{R} - u_{m}| u_{\xi}^{S} d\xi \le C \delta_{R} \delta_{S} (1 + C \delta_{R} \delta_{S}^{2} t)^{-1},$$
(4.51)

where $u^{R} = u^{R} (1 + t, \xi + \sigma t; u_{m}, u_{+})$ and $u^{r} = u^{r} (\frac{\xi + \sigma(1+t)}{1+t}; u_{m}, u_{+}).$

Proof. By Lemma 2.1 and Lemma 3.1, $\forall \varepsilon \in (0, 1)$,

$$\int_{-\infty}^{0} |u^{S} - u_{m}| u_{\xi}^{R} d\xi \leq \delta_{S} \int_{-\infty}^{0} u_{\xi}^{R} d\xi$$

$$= \delta_{S} |u^{R} (1 + t, \sigma t) - u_{m}| \leq C_{\varepsilon} \delta_{S} \delta_{R}^{\frac{2\varepsilon}{2+\varepsilon}} (1 + t)^{-1+\varepsilon},$$

$$(4.52)$$

which is exactly (4.46) if we take $\varepsilon = \frac{1}{5}$. By Hölder inequality, $\forall \varepsilon \in (0, 1)$

$$\int_{0}^{+\infty} |u^{S} - u_{m}| u_{\xi}^{R} d\xi \leq \left(\int_{0}^{+\infty} |u^{S} - u_{m}|^{\frac{1}{1-\varepsilon}} d\xi \right)^{1-\varepsilon} \left(\int_{0}^{+\infty} |u_{\xi}^{R}|^{\frac{1}{\varepsilon}} d\xi \right)^{\varepsilon} \\ \leq C \delta_{S} \left(\int_{0}^{+\infty} (1 + C \delta_{S}^{2} \xi)^{-\frac{1}{1-\varepsilon}} d\xi \right)^{1-\varepsilon} \delta_{R}^{\varepsilon} (1 + t)^{-1+\varepsilon} \\ = C_{\varepsilon} \delta_{S}^{2\varepsilon - 1} \delta_{R}^{\varepsilon} (1 + t)^{-1+\varepsilon} \left(\int_{0}^{+\infty} (1 + \xi)^{-\frac{1}{1-\varepsilon}} d\xi \right)^{1-\varepsilon} \\ \leq C \delta_{S}^{-\frac{3}{5}} \delta_{R}^{\frac{1}{5}} (1 + t)^{-\frac{4}{5}}.$$

$$(4.53)$$

If we take $\varepsilon = \frac{1}{5}$ in(4.53), then (4.47) is proved. By Lemma 3.1 (5), we have for $\forall \varepsilon \in (0, 1)$,

$$\int_{-\infty}^{0} |u^R - u_m| u_{\xi}^S d\xi \le C_{\varepsilon} (1+t)^{-1+\varepsilon} \delta_R^{\frac{2\varepsilon}{2+\varepsilon}} \int_{-\infty}^{0} u_{\xi}^S d\xi$$

$$\le C_{\varepsilon} (1+t)^{-1+\varepsilon} \delta_R^{\frac{2\varepsilon}{2+\varepsilon}} \delta_S,$$
(4.54)

which is exactly (4.48) if we take $\varepsilon = \frac{1}{5}$. Now we prove (4.49). It holds that

$$\int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{R} - u^{r}| u_{\xi}^{S} d\xi
\leq \int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{R}(1+t,\xi+\sigma t) - u^{R}(1+t,\xi+\sigma(1+t))| u_{\xi}^{S} d\xi
+ \int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{R}(1+t,\xi+\sigma(1+t)) - u^{r}| u_{\xi}^{S} d\xi
\leq C ||u_{\xi}^{R}||_{L^{\infty}(\mathbb{R})} \int_{0}^{+\infty} u_{\xi}^{S} d\xi + C_{\varepsilon}(1+t)^{-1+\varepsilon} \delta_{R}^{\varepsilon} \int_{0}^{+\infty} u_{\xi}^{S} d\xi
\leq C_{\varepsilon}(1+t)^{-1+\varepsilon} \delta_{R}^{\varepsilon} \delta_{S},$$
(4.55)

which is (4.49) if we take $\varepsilon = \frac{1}{5}$.

On the other hand, noticing that $C^{-1}\delta_R \leq f'(u_+) - \sigma \leq C\delta_R$ for some constant C > 0and the inequality $\ln(1+x) \leq x$ for x > 0, we have

$$\int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{r} - u_{m}|u_{\xi}^{S}d\xi
= \int_{0}^{[f'(u_{+})-\sigma](1+t)} |(f')^{-1}(\frac{\xi + \sigma(1+t)}{1+t}) - (f')^{-1}(\sigma)|u_{\xi}^{S}d\xi
\leq C\delta_{S}^{3} \int_{0}^{[f'(u_{+})-\sigma](1+t)} \frac{\xi}{1+t} \frac{1}{(C\delta_{S}^{2}\xi + 1)^{2}}d\xi
\leq C\delta_{S} \frac{1}{1+t} \int_{0}^{[f'(u_{+})-\sigma](1+t)} \frac{1}{C\delta_{S}^{2}\xi + 1}d\xi
\leq C\delta_{S}^{-1} \frac{\ln[C\delta_{S}^{2}\delta_{R}(1+t) + 1]}{1+t} \leq C\delta_{S}^{-\frac{3}{5}}\delta_{R}^{\frac{1}{5}} \left(\frac{\ln(C\delta_{S}^{2}\delta_{R}t + 1)}{1+t}\right)^{\frac{4}{5}},$$
(4.56)

and

$$\int_{[f'(u_{+})-\sigma](1+t)}^{+\infty} |u^{R} - u_{m}|u_{\xi}^{S}d\xi
\leq \delta_{R} \Big| u_{m} - u^{S} \big([f'(u_{+}) - \sigma](1+t) \big) \Big|
\leq C \delta_{R} \delta_{S} \frac{1}{1 + C \delta_{S}^{2} \big([f'(u_{+}) - \sigma]t \big)} \leq C \delta_{R} \delta_{S} \big(1 + C \delta_{R} \delta_{S}^{2}t \big)^{-1}.$$
(4.57)

Thus, the proof of Lemma 4.8 is complete.

In the following estimation, C may depend on δ_S but is independent of δ_R . Since δ_R is small and we can assume $\delta_R \leq 1$. Thus we have the following lemmas.

Estimation of the $\int_0^t \mathbf{J}(\tau) d\tau$.

Lemma 4.9. $\mathbf{J}(t)$ is defined in (4.17), it holds that

$$\left|\int_{0}^{t} \mathbf{J}(\tau) d\tau\right| \leq \frac{u_{m}^{2}}{16} \int_{0}^{t} \mu \|\phi_{\xi}^{-\mathbf{X}}\|^{2} d\tau + C u_{m}^{2} \sup_{0 \leq t \leq T} \|\phi^{-\mathbf{X}}\|^{2} \delta_{R}^{\frac{4}{11}}, \text{ for all } t \in [0, T].$$
(4.58)

Proof. By Cauchy inequality and Sobolev inequality, using Lemma 4.8, we have $\forall t \in [0, T]$,

$$\begin{aligned} \left| \int_{0}^{t} \mathbf{J}(\tau) d\tau \right| &= \left| 3 \int_{0}^{t} \int_{\mathbb{R}} (\phi^{-\mathbf{X}})^{2} (u^{S} - u_{m}) u_{\xi}^{R} w d\xi d\tau \right| \\ &\leq C u_{m}^{2} \int_{0}^{t} \|\phi^{-\mathbf{X}}\|_{L^{\infty}(\mathbb{R})}^{2} \int_{\mathbb{R}} |u^{S} - u_{m}| u_{\xi}^{R} d\xi d\tau \\ &\leq \frac{u_{m}^{2}}{16} \int_{0}^{t} \mu \|\phi_{\xi}^{-\mathbf{X}}\|^{2} d\tau + C u_{m}^{2} \sup_{0 \leq t \leq T} \|\phi^{-\mathbf{X}}\|^{2} \int_{0}^{t} \left(\int_{\mathbb{R}} |u^{S} - u_{m}| u_{\xi}^{R} d\xi \right)^{2} d\tau \\ &\leq \frac{u_{m}^{2}}{16} \int_{0}^{t} \mu \|\phi_{\xi}^{-\mathbf{X}}\|^{2} d\tau + C u_{m}^{2} \sup_{0 \leq t \leq T} \|\phi^{-\mathbf{X}}\|^{2} \delta_{R}^{\frac{4}{11}}, \text{ for all } t \in [0, T]. \end{aligned}$$

$$(4.59)$$

Estimation of the $\int_0^t \mathbf{F}(\tau) d\tau$.

.

4

Lemma 4.10. $\mathbf{F}(t)$ is defined in (4.16), it holds that $\forall t \in [0, T]$

$$\left|\int_{0}^{t} \mathbf{F}(\tau) d\tau\right| \leq \frac{u_{m}^{2}}{16} \int_{0}^{t} \mu \|\phi_{\xi}^{-\mathbf{X}}\|^{2} d\tau + C u_{m}^{2} \sup_{0 \leq t \leq T} \|\phi^{-\mathbf{X}}\|^{\frac{2}{3}} \delta_{R}^{\frac{8}{33}}.$$
(4.60)

Proof. By Cauchy inequality and Sobolev inequality, we have

$$\begin{aligned} |\mathbf{F}(\tau)| &= \left| \int_{\mathbb{R}} F \phi^{-\mathbf{X}} w d\xi \right| \le C u_m^2 \|\phi^{-\mathbf{X}}\|_{L^{\infty}(\mathbb{R})} \int_{\mathbb{R}} |F| d\xi \le C u_m^2 \|\phi^{-\mathbf{X}}\|^{\frac{1}{2}} \|\phi_{\xi}^{-\mathbf{X}}\|^{\frac{1}{2}} \int_{\mathbb{R}} |F| d\xi \\ &\le \frac{u_m^2}{16} \mu \|\phi_{\xi}^{-\mathbf{X}}\|^2 + C u_m^2 \|\phi^{-\mathbf{X}}\|^{\frac{2}{3}} \Big(\int_{\mathbb{R}} |F| d\xi \Big)^{\frac{4}{3}}, \end{aligned}$$
(4.61)

where

$$\int_{\mathbb{R}} |F|d\xi \le C \int_{\mathbb{R}} |u^S - u_m| u_{\xi}^R d\xi + C \int_{\mathbb{R}} |u^R - u_m| u_{\xi}^S d\xi + \mu \int_{\mathbb{R}} |u_{\xi\xi}^R| d\xi.$$
(4.62)

Using Lemma 4.8, we have

$$\int_{0}^{t} \left(\int_{\mathbb{R}} |u^{S} - u_{m}| u_{\xi}^{R} d\xi \right)^{\frac{4}{3}} d\tau \le C \delta_{R}^{\frac{8}{33}}.$$
(4.63)

Based on the fact

$$\int_{\mathbb{R}} |u^{R} - u_{m}| u_{\xi}^{S} d\xi \leq \int_{-\infty}^{0} |u^{R} - u_{m}| u_{\xi}^{S} d\xi + \int_{0}^{[f'(u_{+}) - \sigma](1+t)} |u^{R} - u^{r}| u_{\xi}^{S} d\xi + \int_{0}^{[f'(u_{+}) - \sigma](1+t)} |u^{R} - u_{m}| u_{\xi}^{S} d\xi + \int_{[f'(u_{+}) - \sigma](1+t)}^{+\infty} |u^{R} - u_{m}| u_{\xi}^{S} d\xi,$$

$$(4.64)$$

and

$$\int_0^t \left(\int_{[f'(u_+)-\sigma](1+t)}^{+\infty} |u^R - u_m| u_{\xi}^S d\xi \right)^{\frac{4}{3}} d\tau \le C \delta_R^{\frac{4}{3}} \int_0^{+\infty} (1 + C \delta_R t)^{-\frac{4}{3}} dt \le C \delta_R^{\frac{1}{3}}.$$

Thus, by Lemma 4.8, we obtain

$$\int_{0}^{t} \left(\int_{\mathbb{R}} |u^{R} - u_{m}| u_{\xi}^{S} d\xi \right)^{\frac{4}{3}} d\tau \le C \delta_{R}^{\frac{8}{33}}.$$
(4.65)

By Lemma 2.2, we have

$$\int_{\mathbb{R}} |u_{\xi\xi}^{R}| d\xi \leq \begin{cases} \delta_{R}, & 1+t \leq \delta_{R}^{-1}, \\ (1+t)^{-1}, & 1+t \geq \delta_{R}^{-1}, \end{cases}$$
(4.66)

thus

$$\int_0^t \left(\int_{\mathbb{R}} |u_{\xi\xi}^R| d\xi \right)^{\frac{4}{3}} d\tau \le C \delta_R^{\frac{1}{3}}.$$

$$(4.67)$$

Thus we complete the proof of Lemma 4.10.

Notice that $\|\phi\|_{H^1(\mathbb{R})} = \|\phi^{-\mathbf{X}}\|_{H^1(\mathbb{R})}$ and $\sup_{0 \le t \le T} \|\phi\|_{H^1(\mathbb{R})} = \sup_{0 \le t \le T} \|\phi^{-\mathbf{X}}\|_{H^1(\mathbb{R})}$. Using Lemma 4.7, 4.9 and 4.10 and changing of variable $\xi \to \xi + \mathbf{X}(t)$, we can prove Proposition 4.1.

To accomplish the a priori estimates in Proposition 3.2, we need to obtain the following L^2 -estimate for ϕ_{ξ} .

Proposition 4.2. There exist positive constants $\delta_0, \epsilon_4 > 0$, if $\mathcal{E}(T) := \sup_{0 \le t \le T} \|\phi\|_{H^1(\mathbb{R})} \le \epsilon_4$ and the rarefaction wave strength $\delta_R = |u_+ - u_m| \le \delta_0$, then there exists a constant Csuch that for all $t \in [0, T]$, it holds that

$$\|\phi_{\xi}(t)\|^{2} + \int_{0}^{t} \|\phi_{\xi\xi}\|^{2} d\tau \leq C(\|\phi_{0}\|_{H^{1}(\mathbb{R})}^{2} + \delta_{R}^{\frac{8}{33}}).$$

Proof. Multiplying (3.32) by $-\phi_{\xi\xi}$ and integrating the resulted equation with respect to ξ , we can obtain

$$\frac{1}{2}\frac{d}{dt}\|\phi_{\xi}\|^{2} + \mu\|\phi_{\xi\xi}\|^{2} = \int_{\mathbb{R}} \left(f(\phi + \tilde{u}^{\mathbf{X}}) - f(\tilde{u}^{\mathbf{X}})\right)_{\xi} \phi_{\xi\xi} d\xi
+ \dot{\mathbf{X}}(t) \int_{\mathbb{R}} \phi_{\xi\xi}((u^{S})^{\mathbf{X}}_{\xi} + (u^{R})^{\mathbf{X}}_{\xi}) d\xi + \int_{\mathbb{R}} \phi_{\xi\xi} F^{\mathbf{X}} d\xi.$$
(4.68)

By Cauchy inequality on the right-hand side of (4.68), we obtain

$$\frac{1}{2}\frac{d}{dt}\|\phi_{\xi}\|^{2} + \frac{\mu}{2}\|\phi_{\xi\xi}\|^{2} \le C \int_{\mathbb{R}} \left| \left(f(\phi + \tilde{u}^{\mathbf{X}}) - f(\tilde{u}^{\mathbf{X}}) \right)_{\xi} \right|^{2} d\xi + C \int_{\mathbb{R}} |F^{\mathbf{X}}|^{2} d\xi + C |\dot{\mathbf{X}}(t)|^{2}.$$
(4.69)

The first term on the right-hand side of (4.69) can be estimated as

$$\int_{\mathbb{R}} \left| \left(f(\phi + \tilde{u}^{\mathbf{X}}) - f(\tilde{u}^{\mathbf{X}}) \right)_{\xi} \right|^{2} d\xi
\leq \int_{\mathbb{R}} \left| \left(f'(\phi + \tilde{u}^{\mathbf{X}}) - f'(\tilde{u}^{\mathbf{X}}) \right) \tilde{u}_{\xi}^{\mathbf{X}} \right|^{2} d\xi + \int_{\mathbb{R}} \left| f'(\phi + \tilde{u}^{\mathbf{X}}) \phi_{\xi} \right|^{2} d\xi
\leq C \int_{\mathbb{R}} \phi^{2} ((u^{S})_{\xi}^{\mathbf{X}})^{2} d\xi + C \int_{\mathbb{R}} \phi^{2} ((u^{R})_{\xi}^{\mathbf{X}})^{2} d\xi + C ||\phi_{\xi}||^{2}
\leq C \int_{\mathbb{R}} \phi^{2} (u^{S})_{\xi}^{\mathbf{X}} d\xi + C \int_{\mathbb{R}} \phi^{2} (u^{R})_{\xi}^{\mathbf{X}} d\xi + C ||\phi_{\xi}||^{2}.$$
(4.70)

By Proposition 4.1, we have

$$\int_{0}^{t} \int_{\mathbb{R}} \left| \left(f(\phi + \tilde{u}^{\mathbf{X}}) - f(\tilde{u}^{\mathbf{X}}) \right)_{\xi} \right|^{2} d\xi d\tau \le C(\|\phi_{0}\|^{2} + \delta_{R}^{\frac{8}{33}}), \quad t \in [0, T].$$
(4.71)

We next estimate the second term on the right-hand side of (4.69). By the definition of $F^{\mathbf{X}}$, (3.29), it holds that

$$\int_{\mathbb{R}} |F^{\mathbf{X}}|^2 d\xi = \int_{\mathbb{R}} |F|^2 d\xi \leq C \int_{\mathbb{R}} |u^S - u_m|^2 (u_{\xi}^R)^2 d\xi + C \int_{\mathbb{R}} |u^R - u_m|^2 (u_{\xi}^S)^2 d\xi + C \int_{\mathbb{R}} |u_{\xi\xi}^R|^2 d\xi.$$
(4.72)

By (4.46), (4.47) and Lemma 3.1, we have

$$\int_{\mathbb{R}} |u^{S} - u_{m}|^{2} (u_{\xi}^{R})^{2} d\xi
\leq C ||u_{\xi}^{R}||_{L^{\infty}(\mathbb{R})} \int_{\mathbb{R}} |u^{S} - u_{m}| u_{\xi}^{R} d\xi
\leq C \delta_{R}^{\frac{3}{5}} (1+t)^{-\frac{2}{5}} \delta_{R}^{\frac{2}{11}} (1+t)^{-\frac{4}{5}} = C \delta_{R}^{\frac{43}{55}} (1+t)^{-\frac{6}{5}},$$
(4.73)

which implies that

$$\int_{0}^{t} \int_{\mathbb{R}} |u^{S} - u_{m}|^{2} (u_{\xi}^{R})^{2} d\xi d\tau \leq C \delta_{R}^{\frac{43}{55}}.$$
(4.74)

We next estimate the second term on the right-hand side of (4.72).

$$\int_{\mathbb{R}} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi$$

$$\leq \int_{-\infty}^{0} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi + \int_{0}^{[f'(u_{+}) - \sigma](1+t)} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi \qquad (4.75)$$

$$+ \int_{[f'(u_{+}) - \sigma](1+t)}^{+\infty} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi.$$

For $\int_{-\infty}^{0} |u^R - u_m|^2 (u_{\xi}^S)^2 d\xi$, by Lemma 3.1, we have

$$\int_{-\infty}^{0} |u^R - u_m|^2 (u_{\xi}^S)^2 d\xi \le C_{\epsilon} \delta_R^{\frac{4\varepsilon}{2+\epsilon}} (1+t)^{-2+2\epsilon}.$$
(4.76)

Taking $\epsilon = \frac{1}{3}$, we have

$$\int_{0}^{t} \int_{-\infty}^{0} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi d\tau \leq C \delta_{R}^{\frac{4}{7}}.$$
(4.77)

For
$$\int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi, \text{ we have}$$
$$\int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi$$
$$\leq \int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{R} - u^{r}|^{2} (u_{\xi}^{S})^{2} d\xi + \int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{r} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi.$$
(4.78)

By Lemma 3.1, we have

$$\int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{R} - u^{r}|^{2} (u_{\xi}^{S})^{2} d\xi
\leq \int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{R}(1+t,\xi+\sigma t) - u^{R}(1+t,\xi+\sigma(1+t))|^{2} (u_{\xi}^{S})^{2} d\xi
+ \int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{R}(1+t,\xi+\sigma(1+t)) - u^{r}|^{2} (u_{\xi}^{S})^{2} d\xi
\leq C ||u^{R}_{\xi}||^{2}_{L^{\infty}(\mathbb{R})} \int_{0}^{+\infty} (u^{S}_{\xi})^{2} d\xi + C_{\varepsilon} \delta^{2\varepsilon}_{R} (1+t)^{-2+2\varepsilon} \int_{0}^{+\infty} (u^{S}_{\xi})^{2} d\xi
\leq C_{\varepsilon} \delta^{2\varepsilon}_{R} (1+t)^{-2+2\varepsilon},$$
(4.79)

$$\begin{split} &\int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{r}-u_{m}|^{2}(u_{\xi}^{S})^{2}d\xi \\ &= \int_{0}^{[f'(u_{+})-\sigma](1+t)} \left| \left(f'\right)^{-1} \left(\frac{\xi+\sigma(1+t)}{1+t}\right) - (f')^{-1}(\sigma) \right|^{2} (u_{\xi}^{S})^{2}d\xi \\ &\leq C \int_{0}^{[f'(u_{+})-\sigma](1+t)} \frac{\xi^{2}}{(1+t)^{2}} \frac{\delta_{S}^{6}}{(C\delta_{S}^{2}\xi+1)^{4}}d\xi \\ &\leq C \frac{1}{(1+t)^{2}} \int_{0}^{[f'(u_{+})-\sigma](1+t)} \frac{1}{C\xi+1}d\xi \\ &\leq C \frac{1}{(1+t)^{2}} \ln(1+C\delta_{R}t) \\ &\leq C \delta_{R}^{\frac{2}{3}} \frac{\ln^{\frac{1}{3}}(1+C\delta_{R}t)}{(1+t)^{\frac{4}{3}}}. \end{split}$$
(4.80)

Taking $\varepsilon = \frac{1}{3}$ in (4.79) and using (4.79), (4.80), we have

$$\int_{0}^{t} \int_{0}^{[f'(u_{+})-\sigma](1+t)} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi d\tau \leq C \delta_{R}^{\frac{2}{3}}.$$
(4.81)

For $\int_{[f'(u_{+})-\sigma](1+t)}^{+\infty} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi$, we have $\int_{[f'(u_{+})-\sigma](1+t)}^{+\infty} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi$

$$\leq \delta_{R}^{2} \|u_{\xi}^{S}\|_{L^{\infty}\left([f'(u_{+})-\sigma](1+t),+\infty\right)} \int_{[f'(u_{+})-\sigma](1+t)}^{f^{*}} u_{\xi}^{S} d\xi \qquad (4.82)$$
$$\leq C \delta_{R}^{2} \frac{1}{(1+C\delta_{R}t)^{2}},$$

which implies

$$\int_{0}^{t} \int_{[f'(u_{+})-\sigma](1+t)}^{+\infty} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi d\tau \leq C\delta_{R}.$$
(4.83)

Substituting (4.77), (4.81), (4.83) into (4.75), we have

$$\int_{0}^{t} \int_{\mathbb{R}} |u^{R} - u_{m}|^{2} (u_{\xi}^{S})^{2} d\xi d\tau \leq C \delta_{R}^{\frac{4}{7}}.$$
(4.84)

We finally estimate the third term on the right-hand side of (4.69).

$$\int_{\mathbb{R}} |u_{\xi\xi}^{R}|^{2} d\xi \leq C ||u_{\xi\xi}^{R}||_{L^{\infty}(\mathbb{R})}^{\frac{2}{3}} ||u_{\xi\xi}^{R}||_{L^{\infty}(\mathbb{R})}^{\frac{1}{3}} (1+t)^{-1} \leq C \delta_{R}^{\frac{2}{3}} (1+t)^{-\frac{4}{3}},$$
(4.85)

which implies that

$$\int_0^t \int_{\mathbb{R}} |u_{\xi\xi}^R|^2 d\xi d\tau \le C \delta_R^{\frac{2}{3}}.$$
(4.86)

Substituting (4.74), (4.84), (4.86) into (4.72), we have

$$\int_0^t \int_{\mathbb{R}} |F^{\mathbf{X}}|^2 d\xi d\tau \le C \delta_R^{\frac{4}{7}}.$$
(4.87)

Integrating (4.68) with respect to t, by (4.71),(4.87) and Proposition 4.1, we can prove Proposition 4.2.

By Proposition 4.1 and 4.2, we finally have the desired uniform-in-time a priori estimates Proposition 3.2. $\hfill \Box$

5. Appendix

In this appendix we show the detailed calculations of $H_1 + H_2$ in (4.32). Notice that H_1 and H_2 are defined in (4.27) with

$$H_1 = \left[(3u_m^2 - 3(u^S)^2)w' + 3u^S w - \frac{1}{2}w'' \mu u_{\xi}^S \right] (u_* - u_-),$$

$$H_2 = ww''(u_* - u^S)(u^S + 2u_m) + 2w^2 + ww'(u_* + u_- - 2u^S).$$

When $\xi \in (-\infty, \xi_1)$, we have $w = \frac{5}{2}u_m(u_m - u^S)$, $w' = -\frac{5}{2}u_m$, w'' = 0. Then

$$H_{1} = \frac{5}{2} u_{m} \left[(3u_{m}^{2} - 3(u^{S})^{2}) \cdot (-\frac{5}{2}u_{m}) + 3u^{S} \cdot \frac{5}{2}u_{m}(u_{m} - u^{S}) \right]$$

$$= \frac{25}{4} u_{m}^{2} \left[3(u^{S})^{2} - 3u_{m}^{2} + 3u_{m}u^{S} - 3(u^{S})^{2} \right]$$

$$= -\frac{75}{4} u_{m}^{4} + \frac{75}{4} u_{m}^{3} u^{S},$$

(5.1)

and

$$H_{2} = 2\left[\frac{5}{2}u_{m}(u_{m} - u^{S})\right]^{2} + \frac{5}{2}u_{m}(u_{m} - u^{S}) \times \left(-\frac{5}{2}u_{m}\right)\left(-\frac{3}{2}u_{m} - 2u^{S}\right)$$

$$= \frac{25}{4}u_{m}^{2}\left[2u_{m}^{2} - 4u_{m}u^{S} + 2(u^{S})^{2} + \frac{3}{2}u_{m}^{2} + \frac{1}{2}u_{m}u^{S} - 2(u^{S})^{2}\right]$$

$$= \frac{25}{4}u_{m}^{2}\left(\frac{7}{2}u_{m}^{2} - \frac{7}{2}u_{m}u^{S}\right)$$

$$= \frac{175}{4}u_{m}^{4} - \frac{175}{2}u_{m}^{3}u^{S}$$
(5.2)

 $= \frac{175}{8}u_m^4 - \frac{175}{8}u_m^3 u^S.$ When $\xi \in [\xi_1, \xi_*)$, that is, $u^S \in [0, \frac{u_m}{2})$, by (3.33), we have

$$H_{1} = \frac{5}{2}u_{m} \Big[\Big(3u_{m}^{2} - 3(u^{S})^{2} \Big) \Big(-\frac{40}{u_{m}^{2}} (u^{S})^{3} + \frac{30}{u_{m}} (u^{S})^{2} - \frac{5}{2}u_{m} \Big) \\ + 3u^{S} \Big(-\frac{10}{u_{m}^{2}} (u^{S})^{4} + \frac{10}{u_{m}} (u^{S})^{3} - \frac{5}{2}u_{m}u^{S} + \frac{5}{2}u_{m}^{2} \Big) \\ - \frac{1}{2} \Big(-\frac{120}{u_{m}^{2}} (u^{S})^{2} + \frac{60}{u_{m}}u^{S} \Big) \Big((u^{S})^{3} - 3u_{m}^{2}u^{S} + 2u_{m}^{3} \Big) \Big] \\ = \frac{375}{u_{m}} (u^{S})^{5} - 225(u^{S})^{4} - 750u_{m} (u^{S})^{3} + 750u_{m}^{2} (u^{S})^{2} - \frac{525}{4}u_{m}^{3}u^{S} - \frac{75}{4}u_{m}^{4},$$

$$(5.3)$$

and

$$H_{2} = w \Big[\Big(-\frac{120}{u_{m}^{2}} (u^{S})^{2} + \frac{60}{u_{m}} u^{S} \Big) \Big(u_{m}^{2} - \frac{3}{2} u_{m} u^{S} - (u^{S})^{2} \Big) - \frac{20}{u_{m}^{2}} (u^{S})^{4} + \frac{20}{u_{m}} (u^{S})^{3} \\ - 5u_{m} u^{S} + 5u_{m}^{2} + \Big(-\frac{40}{u_{m}^{2}} (u^{S})^{3} + \frac{30}{u_{m}} (u^{S})^{2} - \frac{5}{2} u_{m} \Big) \Big(-\frac{3}{2} u_{m} - 2u^{S} \Big) \Big] \\ = \Big[-\frac{10}{u_{m}^{2}} (u^{S})^{4} + \frac{10}{u_{m}} (u^{S})^{3} - \frac{5}{2} u_{m} u^{S} + \frac{5}{2} u_{m}^{2} \Big] \Big[\frac{180}{u_{m}^{2}} (u^{S})^{4} + \frac{140}{u_{m}} (u^{S})^{3} \\ - 255 (u^{S})^{2} + 60u_{m} u^{S} + \frac{35}{4} u_{m}^{2} \Big] \\ = -\frac{1800}{u_{m}^{4}} (u^{S})^{8} + \frac{400}{u_{m}^{3}} (u^{S})^{7} + \frac{3950}{u_{m}^{2}} (u^{S})^{6} - \frac{3600}{u_{m}} (u^{S})^{5} \\ + \frac{1225}{2} (u^{S})^{4} + 1075u_{m} (u^{S})^{3} - \frac{1575}{2} u_{m}^{2} (u^{S})^{2} + \frac{1025}{8} u_{m}^{3} u^{S} + \frac{175}{8} u_{m}^{4}. \end{aligned}$$

$$(5.4)$$

Thus, we get

$$H_1 + H_2 = -\frac{1800}{u_m^4} (u^S)^8 + \frac{400}{u_m^3} (u^S)^7 + \frac{3950}{u_m^2} (u^S)^6 - \frac{3225}{u_m} (u^S)^5 + \frac{775}{2} (u^S)^4 + 325u_m (u^S)^3 - \frac{75}{2} u_m^2 (u^S)^2 - \frac{25}{8} u_m^3 u^S + \frac{25}{8} u_m^4.$$
(5.5)

HUANG, WANG, AND ZHANG

Acknowledgment. The work of Feimin Huang was partially supported by National Key R&D Program of China No. 2021YFA1000800, and National Natural Sciences Foundation of China (NSFC) No. 12288201. The work of Yi Wang was partially supported by NSFC (Grant No. 12171459, 12288201, 12090014) and CAS Project for Young Scientists in Basic Research, Grant No. YSBR-031.

Conflict of Interest: The authors declared that they have no conflicts of interest to this work.

Availability of data and material: Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

References

- M. Brio and J. K. Hunter. Rotationally invariant hyperbolic waves. Comm. Pure Appl. Math., 43 (8): 1037-1053, 1990.
- [2] K. Choi, M.-J. Kang, Y. Kwon, and A. Vasseur. Contraction for large perturbations of traveling waves in a hyperbolic-parabolic system arising from a chemotaxis model. Math. Models Methods Appl. Sci, 30(2):387-437, 2020.
- [3] H. Freistühler and T.-P. Liu. Nonlinear stability of overcompresive shock waves in a rotationally invariant system of viscous conservation laws. Comm. Math. Phys., 153 (1): 147-158, 1993.
- [4] H. Freistühler, D. Serre, L¹ stability of shock waves in scalar viscous conservation laws. Comm. Pure Appl. Math. 51 (1998), no. 3, 291-301.
- [5] F. M. Huang and L. D. Xu, Decay rate toward the traveling wave for scalar viscous conservation law, Commun. Math. Anal. Appl., 1 (2022), no. 3, 395-409.
- [6] A. M. Il'in and O. A. Oleinik. Asymptotic behavior of solution of the Cauchy problem for some quasilinear equations for large values of the time. Mat. Sb. (N.S.), 51 (93):191-216, 1960.
- [7] C. Jones, R. Gardner and T. Kapitula. Stability of traveling waves for non-convex scalar viscous conservation laws. Comm. Pure Appl. Math., 46:505-526, 1993.
- [8] M.-J. Kang and A. Vasseur. Contraction property for large perturbations of shocks of the barotropic Navier-Stokes system. J. Eur. Math. Soc., 23 (2):585-638, 2020.
- M.-J. Kang and A. Vasseur. L²-contraction for shock waves of scalar viscous conservation laws. Ann. l'Institut Henri Poincaré C, Analyse non lineaire, 34 (1):139156, 2017.
- [10] M.-J. Kang. L² -type contraction for shocks of scalar viscous conservation laws with strictly convex flux. J. Math. Pure Appl., 145:1-43, 2021.
- [11] M.-J. Kang, A. Vasseur and Y. Wang. L²-contraction of large planar shock waves for multi-dimensional scalar viscous conservation laws. Journal of Differential Equations, 267(5): 2737-2791, 2019.
- [12] M.-J. Kang, A. Vasseur and Y. Wang. Time-asymptotic stability of composite waves of viscous shock and rarefaction for barotropic Navier-Stokes equations. Adv. Math., 419:108963, 2023.
- [13] M.-J. Kang, A. Vasseur, Y. Wang. Time-asymptotic stability of generic Riemann solutions for compressible Navier-Stokes-Fourier equations. arXiv:2306.05604 (2023).
- [14] S. Kawashima and A. Matsumura. Stability of shock profiles in viscoelasticity with non-convex constitutive relations. Comm. Pure Appl. Math., 47 (12):1547-1569, 1994.

- [15] I. P. Lee-Bapty and D. G. Crighton. Nonlinear Wave Motion Governed by the Modified Burgers Equation. Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences, 323:173-209, 1987.
- [16] T.-P. Liu. Shock Waves. Graduate Studies in Mathematics. American Mathematical Society, 215:106-109, 2021.
- [17] H. Liu. Asymptotic stability of shock profiles for non-convex convection-diffusion equation. Appl. Math. Letters, 10 (1):129-134, 1997.
- [18] A. Matsumura. Waves in compressible fluids: viscous shock, rarefaction, and contact waves. In Handbook of mathematical analysis in mechanics of viscous fluids, pages 2495–2548. Springer, Cham, 2018.
- [19] A. Matsumura and K. Nishihara. Asymptotic stability of traveling waves for scalar viscous conservation laws with non-convex nonlinearity. Comm. Math. Phys., 165:83-96, 1994.
- [20] A. Matsumura and K. Nishihara. Asymptotics toward the rarefaction waves of the solutions of a one-dimensional model system for compressible viscous gas. Japan J. Appl. Math., 3:1-13, 1986.
- [21] A. Matsumura and N. Yoshida. Asymptotic behavior of solutions to the Cauchy problem for the scalar viscous conservation law with partially linearly degenerate flux. SIAM J. Math. Anal., 44 (4):2526-2544, 2012.
- [22] M. Mei. Stability of shock profiles for non-convex scalar viscous conservation laws. Math. Mod. Meth. Appl. Sci., 5 (3):279-296,1995.
- [23] G. A. Nariboli and W. C. Lin. A new type of Burgers' equation. ZAMM-J. Appl. Math. Mech., 53 (8):505-510, 1973.
- [24] S. Osher and J. Ralston. L₁ stability of traveling waves with applications to convective porous media flow. Comm. Pure Appl. Math., 35:737-751, 1982.
- [25] M. Teymur and E. Suhubi. Wave propagation in dissipative or dispersive non-linear media. IMA J. App. Math., 21(1):25-40, 1978.
- [26] H. F. Weinberger. Long-time behavior for a regularized scalar conservation law in the absence of genuine nonlinearity. Ann. Inst. Henri Poincaré, 7:407-425, 1990.

(Feimin Huang)

Institute of Applied Mathematics, AMSS, CAS, Beijing 100190, P. R. China and School of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Email address: fhuang@amt.ac.cn

(Yi Wang)

INSTITUTE OF APPLIED MATHEMATICS, AMSS, CAS, BEIJING 100190, P. R. CHINA AND SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, P. R. CHINA

Email address: wangyi@amss.ac.cn

(Jian Zhang)

INSTITUTE OF APPLIED MATHEMATICS, AMSS, CAS, BEIJING 100190, P. R. CHINA AND SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, P. R. CHINA

Email address: zhangjian2020@amss.ac.cn