

Fading memory and the convolution theorem

Juan-Pablo Ortega* Florian Rossmannek*

Abstract

Several topological and analytical notions of continuity and fading memory for causal and time-invariant filters are introduced, and the relations between them are analysed. A significant generalization of the convolution theorem that establishes the equivalence between the fading memory property and the availability of convolution representations of linear filters is proved. This result extends a previous such characterization to a complete array of weighted norms in the definition of the fading memory property. Additionally, the main theorem shows that the availability of convolution representations can be characterized, at least when the codomain is finite-dimensional, not only by the fading memory property but also by the reunion of two purely topological notions that are called minimal continuity and minimal fading memory property. Finally, when the input space and the codomain of a linear functional are Hilbert spaces, it is shown that minimal continuity and the minimal fading memory property guarantee the existence of interesting embeddings of the associated reproducing kernel Hilbert spaces and approximation results of solutions of kernel regressions in the presence of finite data sets.

Keywords: fading memory property, convolution theorem, linear I/O system, duality, kernel learning, linear kernel, representer theorem, approximation theory.

1. Introduction

Fading memory is a natural modelling assumption that has played a key role in approximation and realization results that are central in control, signal treatment, filtering, and systems theory. Given a causal and time-invariant input/output system, the various notions of fading memory in the literature intend to mathematically encode the idea that the dependence of an output on a past input fades out as this input is further and further into the past. Behavior of this type is expected in the description of many phenomena in the natural and social sciences and is formulated as such in classical works that go back to V. Volterra [35] and N. Wiener [37].

From a mathematical perspective, hypotheses on the memory behavior of certain systems have been central in the formulation of approximation results for systems defined on infinite time intervals. This was done for the first time in the seminal paper [2], which extended to infinite time intervals a classical theorem [27] on the uniform approximation abilities of Volterra series for functions defined on finite time intervals; it is in that paper, also for the first time, that fading memory is formulated as a continuity property with respect to a weighted norm. Similar results have been obtained for recurrent neural networks assuming first finite memory [22, 23, 26, 28, 29, 33] and, later on, fading memory [7, 9, 10]. Other results in connection with the universal approximation properties of recurrent neural systems that also identify fading memory as a key hypothesis are [18–21]. In a nutshell, the main mathematical feature that fading memory brings in its wake is that uniformly bounded semi-infinite input sequences form a compact space with respect to the topology induced by the weighted norms used to define fading memory despite the infinite dimensionality of the corresponding space. This is a key fact needed in classical statements like the Stone-Weierstrass theorem, which are at the core of many universal approximation results. Additionally, fading memory can also be found in the literature in relation to the availability and construction of temporal extensions of filters [1, 4, 30, 31] and the stability of certain differential equations [38].

In this paper, we focus on another important consequence of the fading memory property, namely the availability of convolution representations for linear filters. Realizations of this type are crucial in signal treatment applications since they allow filter estimation using traditional impulse/response analyses. Historically, it was always informally assumed that linear, time-invariant, causal, continuous filters have a convolution representation. Even though this statement is true when the input is defined on finite time intervals, it was

*Division of Mathematical Sciences, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore

pointed out for the first time in [2] that it is actually false in the presence of infinite intervals. Moreover, in this reference, it was proved that a convolution representation exists if and only if the filter has the fading memory property. This statement is called the *convolution theorem* (not to be mistaken with the result on the Fourier transform of the convolution of two functions), and *a significant generalization of its statement is the main subject of the present work*. Recently, linear filters and their convolution representations have regained significant attention with the rise of state-space models for long sequence modelling as an alternative to the Transformer model [34]. Various enhancements of state-space models have enabled them to overcome previous challenges and achieve competitive results in different benchmarks [24]. Among the various models that have been proposed, Mamba [12] and subsequent models based on it such as SiMBA [25] have gained particular interest due to their remarkable balance of performance and efficiency. Beneath the various enhancements and additions to the base model, at the core of these models is a linear state-space system that is typically assumed to admit a convolution representation, to permit both a linear recurrent and a faster kernel-based implementation. Our work makes this latter assumption rigorous and provides the theoretical background.

The contents of this paper are divided into three sections. Section 2 introduces various possible topological and analytical notions of fading memory that generalize the original definition in [2] and spells out the relations between them. Section 3 is the core of the paper and studies the interplay, in the context of linear functionals, between the availability of what are called formal and proper convolution representations and various types of fading memory introduced with different weighted norms. In particular, Theorem 17 contains a significant generalization of the convolution theorem introduced in [2] for linear functionals that have a fading memory property defined using weighted ℓ^p -norms with a fixed $p \in [1, \infty]$. We recall that the original result [2, Theorem 5] was only proved for the case $p = \infty$ and a 1-dimensional domain. Additionally, our theorem shows that the availability of convolution representations can be characterized, at least when the codomain is finite-dimensional, not only by the fading memory property but also by the reunion of two purely topological notions that we shall denominate as minimal continuity and minimal fading memory property. A stronger type of fading memory is shown to be equivalent to finite memory for linear functionals. One of the main messages of Section 3 is that one ought to think carefully how to define fading memory in one's context and that even the domain of the functional under consideration can have a significant effect. Finally, Section 4 considers the case in which the input space \mathcal{Z} and the codomain of linear functionals are defined on Hilbert spaces and studies the natural reproducing kernel Hilbert spaces determined by those functionals. The minimal continuity and minimal fading memory property introduced in the previous section appear here as conditions that allow the embedding of the reproducing kernel Hilbert space in $\ell^2(\mathcal{Z})^*$ and in an approximation result of solutions of kernel regressions in the presence of finite data sets.

2. Notions of fading memory

Suppose $(\mathcal{Z}, \|\cdot\|)$ and $(\mathcal{Y}, \|\cdot\|_{\mathcal{Y}})$ are normed vector spaces over \mathbb{R} . Let $\mathcal{B} = \{z \in \mathcal{Z} : \|z\| \leq 1\}$ be the closed unit ball in \mathcal{Z} . Denote by $\mathbb{N} = \{1, 2, \dots\}$ the set of strictly positive integers and by $\mathbb{Z}_- = \mathbb{Z} \setminus \mathbb{N}$ the set of non-positive integers. Underlined symbols like \underline{z} denote sequences in $\mathcal{Z}^{\mathbb{Z}_-}$, and $\underline{z}_t \in \mathcal{Z}$ the t -th entry, $t \in \mathbb{Z}_-$, of \underline{z} . For any $t \in \mathbb{Z}_-$, let $\delta^t : \mathcal{Z} \rightarrow \mathcal{Z}^{\mathbb{Z}_-}$ be the *inclusion* $\delta^t(z) = (\dots, 0, z, 0, \dots)$ that inserts $z \in \mathcal{Z}$ at the time entry t into the zero vector.

Assumption 1. Throughout, we fix a subset $\underline{\mathcal{Z}} \subseteq \mathcal{Z}^{\mathbb{Z}_-}$ with the following properties.

- (i) The set $\underline{\mathcal{Z}}$ is convex and symmetric; the latter meaning $\underline{\mathcal{Z}} = \{-\underline{z} : \underline{z} \in \underline{\mathcal{Z}}\}$.
- (ii) The set $\underline{\mathcal{Z}}$ contains the inclusions of the unit ball $\bigcup_{t \leq 0} \delta^t(\mathcal{B})$.
- (iii) For any $\underline{z} \in \underline{\mathcal{Z}}$ and any subset $J \subseteq \mathbb{Z}_-$, the set $\underline{\mathcal{Z}}$ contains the sequence $\sum_{t \in J} \delta^t(\underline{z}_t)$, in which \underline{z}_t is replaced by 0 for all $t \in \mathbb{Z}_- \setminus J$.

Historically, the main objects of interest have been time-invariant *filters* $\underline{\mathcal{Z}} \rightarrow \mathcal{Y}^{\mathbb{Z}_-}$ with a shift-invariant domain. We recall that such filters are in a 1:1 correspondence with *functionals* $\underline{\mathcal{Z}} \rightarrow \mathcal{Y}$. This equivalence justifies that we restrict our attention to functionals throughout. Details on this equivalence can be found in Appendix A.

We now study the implications of continuity of a functional $H : \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ when the domain $\underline{\mathcal{Z}}$ is equipped with various interesting topologies. With respect to any of these topologies, the inclusions $\delta^t : \mathcal{B} \rightarrow \underline{\mathcal{Z}}$ of the unit ball become continuous. Thus, the following notion is the common denominator of the various notions of continuity of H that we will encounter.

Definition 2. A functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ is said to be *minimally continuous* if $H \circ \delta^t: \mathcal{B} \rightarrow \mathcal{Y}$ is continuous for any $t \in \mathbb{Z}_-$.

Our first notion of fading memory is designed to capture the basic property that a functional memory should possess to qualify for such a denomination.

Definition 3. A functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ is said to have the *minimal fading memory property (minimal FMP)* if $H(\sum_{t=T}^0 \delta^t(\underline{z}_t))$ converges to $H(\underline{z})$ as $T \rightarrow -\infty$ for any $\underline{z} \in \underline{\mathcal{Z}}$.

The sequence $\sum_{t=T}^0 \delta^t(\underline{z}_t)$ appearing in the definition of the minimal FMP is the one that is constantly zero up to time $T-1$ and agrees with \underline{z} from time T onwards. The minimal FMP is similar to the so-called *input forgetting property* (also known as the *unique steady-state property*) [2, 11, 13]. However, these properties are not the same, and neither one implies the other in general. Compared to other natural notions of fading memory that we introduce below, the minimal FMP is arguably a weak property. However, in the context of linear functionals, the minimal FMP will turn out to be surprisingly strong.

Remark 4. It is customary to define the fading memory of a functional as being continuous with respect to a certain topology. The minimal FMP can alternatively be defined in such a way. Let us say a topology on $\underline{\mathcal{Z}}$ has property (P) if for any $\underline{z} \in \underline{\mathcal{Z}}$ the sequence $\sum_{t=T}^0 \delta^t(\underline{z}_t)$ converges to \underline{z} with respect to this topology as $T \rightarrow -\infty$. Given a functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$, the following are equivalent. (i) H has the minimal FMP. (ii) The initial topology on $\underline{\mathcal{Z}}$ induced by H has property (P). (iii) H is continuous with respect to some topology on $\underline{\mathcal{Z}}$ that has property (P).

The next notion of fading memory has been used in [2, Chapter IX], but only in the context of approximation and not in the context of the convolution theorem, a result at the core of this paper that is later on presented as Theorem 17.


Definition 5. A functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ is said to have the *product fading memory property (product FMP)* if it is continuous with respect to the subspace product topology on $\underline{\mathcal{Z}}$.

The minimal and the product FMP are topological notions. Nevertheless, the definition of fading memory has been historically introduced as continuity with respect to weighted ℓ^p -norms [2, 11], which is an analytical notion, and which we now recall. Given $p \in [1, \infty]$ and a sequence $\underline{w} \in (0, 1]^{\mathbb{Z}_-}$, let $\ell_{\underline{w}}^p(\mathcal{Z}) \subseteq \mathcal{Z}^{\mathbb{Z}_-}$ be the set of all sequences in \mathcal{Z} with $\|\underline{z}\|_{\underline{w}, p} = (\sum_{t \leq 0} \underline{w}_t \|\underline{z}_t\|^p)^{1/p} < \infty$ if $p < \infty$, respectively $\|\underline{z}\|_{\underline{w}, \infty} = \sup_{t \leq 0} \underline{w}_t \|\underline{z}_t\| < \infty$. This makes $(\ell_{\underline{w}}^p(\mathcal{Z}), \|\cdot\|_{\underline{w}, p})$ normed vector spaces. Taking \underline{w} to be constantly 1, we obtain the usual sequence spaces $(\ell^p(\mathcal{Z}), \|\cdot\|_p)$. The closed unit ball in $\ell^p(\mathcal{Z})$ is denoted $\mathbb{B}^p(\mathcal{Z}) = \{\underline{z} \in \ell^p(\mathcal{Z}): \|\underline{z}\|_p \leq 1\}$. The set $c_0(\mathcal{Z}) \subseteq \mathcal{Z}^{\mathbb{Z}_-}$ contains all sequences vanishing at minus infinity. We call $\underline{w} \in (0, 1]^{\mathbb{Z}_-}$ a *weighting sequence* if it is monotone and $\lim_{t \rightarrow -\infty} \underline{w}_t = 0$.

Definition 6. Let $p \in [1, \infty]$. A functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ is said to be *p-continuous* if $\underline{\mathcal{Z}} \subseteq \ell^p(\mathcal{Z})$ and H is continuous with respect to the topology induced by $\|\cdot\|_p$. A functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ is said to have the *p-weighted fading memory property (p-weighted FMP)* if $\underline{\mathcal{Z}} \subseteq \ell^p(\mathcal{Z})$ and there exists a weighting sequence \underline{w} such that H is continuous with respect to the topology induced by $\|\cdot\|_{\underline{w}, p}$.

The fading memory appearing in the pioneering work [2] is the ∞ -weighted FMP. Many works in the literature adopt the same notion of fading memory (see [10, 11, 13, 14, 17] to name just a few). One of the first works to consider the p -weighted FMP for $p \neq \infty$ was [11]. We point out that most of these works assume the underlying space \mathcal{Z} to be compact. In this case, the product FMP and the ∞ -weighted FMP are equivalent [10, 11]. When \mathcal{Z} is not compact, the distinction becomes important. Although \mathcal{Z} is not compact in this work, we have flexibility in choosing the subset $\underline{\mathcal{Z}}$ of $\mathcal{Z}^{\mathbb{Z}_-}$. In particular, given a compact set $K \subseteq \mathcal{Z}$, we could consider $\underline{\mathcal{Z}} \subseteq K^{\mathbb{Z}_-}$, effectively recovering the setups of other works.

Remark 7. Consider a filter $\underline{\mathcal{Z}} \rightarrow \mathcal{Y}$. The product FMP implies both the minimal FMP and minimal continuity. Furthermore, assuming $\underline{\mathcal{Z}} \subseteq \ell^p(\mathcal{Z})$ for a given $p \in [1, \infty]$, the following implications hold.



3. Linear fading memory functionals

We point out that the set $\underline{\mathcal{Z}}$ need not be a linear subspace of $\mathcal{Z}^{\mathbb{Z}_-}$. We call a functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ **linear** if it is the restriction of a (necessarily unique) linear functional defined on the linear span of $\underline{\mathcal{Z}}$. Let $(L(\mathcal{Z}, \mathcal{Y}), \|\cdot\|_{\text{op}})$ be the space of continuous linear functions $\mathcal{Z} \rightarrow \mathcal{Y}$ with the operator norm induced by $\|\cdot\|$ and $\|\cdot\|_{\mathcal{Y}}$. For any $q \in [1, \infty)$ and $\underline{\kappa} \in L(\mathcal{Z}, \mathcal{Y})^{\mathbb{Z}_-}$, denote $\|\underline{\kappa}\|_{\text{op}, q} = (\sum_{t \leq 0} \|\kappa_t\|_{\text{op}}^q)^{1/q}$ and $\|\underline{\kappa}\|_{\text{op}, \infty} = \sup_{t \leq 0} \|\kappa_t\|_{\text{op}}$. We say that a functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ has a **formal convolution representation** if there exists a sequence $\underline{\kappa} \in L(\mathcal{Z}, \mathcal{Y})^{\mathbb{Z}_-}$ such that for all $\underline{z} \in \underline{\mathcal{Z}}$ we have

$$H(\underline{z}) = \lim_{T \rightarrow -\infty} \sum_{t=T}^0 \kappa_t(\underline{z}_t). \quad (1)$$

Note that a functional with a formal convolution representation must be linear and that $\underline{\kappa}$ must be unique. We say that H has a **(proper) convolution representation** if the series in (1) converges absolutely for all $\underline{z} \in \underline{\mathcal{Z}}$. We say that H has **finite memory** if κ_t is zero for all but at most finitely many $t \in \mathbb{Z}_-$. We can immediately relate convolution representations to p -continuity and the p -weighted FMP.

Lemma 8. *Let $p \in [1, \infty]$, and suppose $\mathbb{B}^p(\mathcal{Z}) \subseteq \underline{\mathcal{Z}} \subseteq \ell^p(\mathcal{Z})$ or $c_0(\mathcal{Z}) \cap \mathbb{B}^p(\mathcal{Z}) \subseteq \underline{\mathcal{Z}} \subseteq c_0(\mathcal{Z}) \cap \ell^p(\mathcal{Z})$. Consider a linear functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ with a formal convolution representation $\underline{\kappa}$.*

- (i) *The functional H is p -continuous.*
- (ii) *If $p \neq 1$ and $\|\underline{\kappa}\|_{\text{op}, q} < \infty$ for the Hölder conjugate $q \in [1, \infty)$ of p , then H has the p -weighted FMP.*

Proof. (i) Note that $\text{span}(\underline{\mathcal{Z}}) = \ell^p(\mathcal{Z})$ or $\text{span}(\underline{\mathcal{Z}}) = c_0(\mathcal{Z}) \cap \ell^p(\mathcal{Z})$, which is a Banach space with norm $\|\cdot\|_p$ in either case. For each $n \in \mathbb{N}$, the finite memory linear functional $H_n: \text{span}(\underline{\mathcal{Z}}) \rightarrow \mathcal{Y}$ given by $H_n(\underline{z}) = \sum_{t=-n}^0 \kappa_t(\underline{z}_t)$ is p -continuous. Let \tilde{H} be the unique linear extension of H to $\text{span}(\underline{\mathcal{Z}})$. Since H_n converges point-wise to \tilde{H} , the sequence $(\|H_n(\underline{z})\|_{\mathcal{Y}})_n$ is bounded for all $\underline{z} \in \text{span}(\underline{\mathcal{Z}})$. By the principle of uniform boundedness, the operator norm of H_n with respect to $\|\cdot\|_p$ is uniformly bounded in n . This implies that the point-wise limit of H_n has finite operator norm with respect to $\|\cdot\|_p$, that is, \tilde{H} is p -continuous.

(ii) Let $r = 1/p$ if $p < \infty$ and $r = 1$ if $p = \infty$. Take a weighting sequence \underline{w} such that $\sum_{t \leq 0} \underline{w}_t^{-qr} \|\kappa_t\|_{\text{op}}^q < \infty$. Then, Hölder's inequality shows that H has the p -weighted FMP;

$$\|H(\underline{z})\|_{\mathcal{Y}} \leq \sum_{t \leq 0} \underline{w}_t^{-r} \|\kappa_t\|_{\text{op}} \underline{w}_t^r \|\underline{z}_t\| \leq \left(\sum_{t \leq 0} \underline{w}_t^{-qr} \|\kappa_t\|_{\text{op}}^q \right)^{1/q} \|\underline{z}\|_{\underline{w}, p}.$$

■

We will see later that the case $p = 1$ is special and that a convolution representation satisfying $\|\underline{\kappa}\|_{\text{op}, \infty} < \infty$ is indeed not enough to guarantee the 1-weighted FMP.

Example 9. Let $(\mathcal{X}, \|\cdot\|_{\mathcal{X}})$ be a Banach space, and let $A \in L(\mathcal{X}, \mathcal{X})$ and $C \in L(\mathbb{R}^d, \mathcal{X})$. Suppose the linear state equation $\underline{x}_t = A(\underline{x}_{t-1}) + C(\underline{z}_t)$ admits a unique solution for every input sequence $\underline{z} \in \ell^\infty(\mathbb{R}^d)$. The associated functional $H_{A,C}: \ell^\infty(\mathbb{R}^d) \rightarrow \mathcal{X}$ is minimally continuous since $H_{A,C} \circ \delta^t = A^{-t} \circ C$. If the spectral radius $\rho(A)$ is strictly smaller than 1, then the convolution representation $\underline{\kappa}_t = A^{-t} \circ C$ satisfies $\|\underline{\kappa}\|_{\text{op}, 1} < \infty$, in which case $H_{A,C}$ has the ∞ -weighted FMP. Indeed, $\rho(A) = \lim_{k \rightarrow \infty} \|A^k\|_{\text{op}}^{1/k}$ by Gelfand's formula [5] and, hence, given any $r \in (\rho(A), 1)$ there exists some $t_0 \in \mathbb{Z}_-$ such that $\|A^{-t}\|_{\text{op}} \leq r^{-t}$ for all $t \leq t_0$, which implies that $\|\underline{\kappa}\|_{\text{op}, 1} < \infty$. If $\mathcal{X} \cong \mathbb{R}^n$ is finite-dimensional, then the condition that $\rho(A) < 1$ is implicit in the existence and uniqueness of solutions, and so is the ∞ -weighted FMP. Indeed, if we had $\rho(A) \geq 1$, then there existed an eigenvector $v \in \mathbb{C}^n$ of A for some eigenvalue $\lambda \in \mathbb{C}$ with $|\lambda| \geq 1$. In this case, both the constant zero sequence and the sequence $\underline{x}_t = \text{Re}(\lambda^t v)$ belonged to $\ell^\infty(\mathbb{R}^n)$ and posed a solution to the state equation when the input is the constant zero sequence [17].

The next lemma justifies our introduction of the new notions of minimal continuity and minimal FMP, and it gives us a taste of how strong these seemingly weak notions turn out to be for linear functionals.

Lemma 10. *A linear functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ has a formal convolution representation if and only if it has the minimal FMP and is minimally continuous.¹*

Proof. Suppose H has the minimal FMP and is minimally continuous. Let $\tilde{H}: \text{span}(\underline{\mathcal{Z}}) \rightarrow \mathcal{Y}$ be the unique linear extension of H , and define $\underline{\kappa}_t = \tilde{H} \circ \delta^t$. Restricted to the unit ball, $\underline{\kappa}_t = H \circ \delta^t$ is continuous. Being a linear map, continuity on the unit ball is equivalent to continuity on all of $\underline{\mathcal{Z}}$. Also by linearity, $H(\sum_{t=T}^0 \delta^t(\underline{z}_t)) = \sum_{t=T}^0 \underline{\kappa}_t(\underline{z}_t)$ for any $\underline{z} \in \underline{\mathcal{Z}}$ and $T \in \mathbb{Z}_-$. Taking the limit and applying the minimal FMP, we obtain a formal convolution representation of H . The converse implication is clear. \blacksquare

Example 11. Consider a Banach limit $H: \ell^\infty(\mathbb{R}) \rightarrow \mathbb{R}$, that is, H is a positive, shift-invariant, ∞ -continuous, linear functional such that $H(\underline{z}) = \lim_{t \rightarrow -\infty} \underline{z}_t$ whenever the limit exists [5]. Then, H does not have the minimal FMP and, hence, neither a convolution representation nor the product or the ∞ -weighted FMP. To see this, simply observe that $H(\sum_{t=T}^0 \delta^t(\underline{z}_t)) \equiv 0$, so H would have to be constantly zero.

Next, we fill the gap in Lemma 8.(ii) regarding the case $p = 1$.

Proposition 12. *Suppose $\underline{\mathcal{Z}} \subseteq \ell^1(\mathcal{Z})$. Then, a linear functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ has the 1-weighted FMP if and only if it has a convolution representation satisfying $\lim_{t \rightarrow -\infty} \|\underline{\kappa}_t\|_{\text{op}} = 0$.*

Proof. Suppose H has a convolution representation satisfying $\lim_{t \rightarrow -\infty} \|\underline{\kappa}_t\|_{\text{op}} = 0$. Consider the monotone sequence $\underline{w}_t = \sup_{s \leq t} \min\{1, \|\underline{\kappa}_s\|_{\text{op}}\}$. If \underline{w} is eventually zero, then H has finite memory, which clearly implies the 1-weighted FMP. If \underline{w} is not eventually zero, then it is a weighting sequence. With this weighting sequence, $\|H(\underline{z})\|_{\mathcal{Y}} \leq C \|\underline{z}\|_{\underline{w},1}$, where $C = \sup_{t \leq 0} \max\{1, \|\underline{\kappa}_t\|_{\text{op}}\}$. This shows that H has the 1-weighted FMP. Conversely, if H has the 1-weighted FMP, then there exists a weighting sequence \underline{w} such that H is continuous with respect to $\|\cdot\|_{\underline{w},1}$. Take $(\eta_n)_{n \in \mathbb{N}} \subseteq (0, 1)$ such that $\|H(\underline{z})\|_{\mathcal{Y}} \leq n^{-1}$ for all $\underline{z} \in \underline{\mathcal{Z}}$ with $\|\underline{z}\|_{\underline{w},1} \leq \eta_n$. We know from Remark 7 and Lemma 10 that H has a formal convolution representation with some $\underline{\kappa} \in L(\mathcal{Z}, \mathcal{Y})^{\mathbb{Z}_-}$. Let $\varepsilon > 0$ and $n \geq \varepsilon^{-1}$. Take $T \in \mathbb{Z}_-$ with $\underline{w}_T \leq \eta_n$. For any $t \leq T$ and any $z \in \mathcal{Z}$ with $\|z\| = 1$, we have $\|\delta^t(z)\|_{\underline{w},1} = \underline{w}_t \leq \eta_n$ and, hence, $\|\underline{\kappa}_t(z)\|_{\mathcal{Y}} = \|H(\delta^t(z))\|_{\mathcal{Y}} \leq n^{-1} \leq \varepsilon$. Thus, $\|\underline{\kappa}_t\|_{\text{op}} \rightarrow 0$ as $t \rightarrow -\infty$. In particular, $\|\underline{\kappa}\|_{\text{op},\infty} < \infty$, which implies that the formal convolution representation is a proper one. \blacksquare

Example 13. Let $(\mathcal{V}, \|\cdot\|_{\mathcal{V}})$ and $(\mathcal{W}, \|\cdot\|_{\mathcal{W}})$ be two normed vector spaces over \mathbb{R} . Take $\mathcal{Z} = L(\mathcal{V}, \mathcal{W})$ to be the vector space of continuous linear maps from \mathcal{V} to \mathcal{W} with the operator norm, and $\mathcal{Y} = \ell^1(\mathcal{W})$ with the ℓ^1 -norm. Fix a bounded sequence $\underline{v} \in \ell^\infty(\mathcal{V})$, and consider the sequential evaluation functional

$$H_{\underline{v}}: \ell^1(L(\mathcal{V}, \mathcal{W})) \rightarrow \ell^1(\mathcal{W}), \quad \underline{A} \mapsto (\underline{A}_t(\underline{v}_t))_{t \leq 0}.$$

Then, $H_{\underline{v}}$ is minimally continuous, has the minimal FMP, and has the formal convolution representation $\underline{\kappa}_t(A) = H \circ \delta^t(A) = (\dots, 0, A(\underline{v}_t), 0, \dots, 0)$. In particular, $\|\underline{\kappa}_t\|_{\text{op}} = \|\underline{v}_t\|_{\mathcal{V}}$, and $H_{\underline{v}}$ has the 1-weighted FMP if and only if $\lim_{t \rightarrow -\infty} \|\underline{v}_t\|_{\mathcal{V}} = 0$.

Further above, we introduced the notation $\|\underline{\kappa}\|_{\text{op},q}$ for $\underline{\kappa} \in L(\mathcal{Z}, \mathcal{Y})^{\mathbb{Z}_-}$. We use the same notation for elements in $L(\mathcal{Z}, \mathbb{R})^{\mathbb{Z}_-}$, where $L(\mathcal{Z}, \mathbb{R})$ is equipped with the operator norm induced by $\|\cdot\|$ and the absolute value on \mathbb{R} . Given an $L \in \mathcal{Y}^*$, we denote by $L \circ \underline{\kappa} = (L \circ \underline{\kappa}_t)_{t \leq 0}$ the element-wise composition. The proof of the next result uses the well-known duality of sequence spaces and their duals.

Proposition 14. *Let $p, q \in [1, \infty]$ be Hölder conjugates. Suppose $\{\underline{z} \in \text{span}(\underline{\mathcal{Z}}): \|\underline{z}\|_p \leq 1\} \subseteq \underline{\mathcal{Z}} \subseteq c_0(\mathcal{Z}) \cap \ell^p(\mathcal{Z})$. Then, any p -continuous linear functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ has a convolution representation satisfying $\|L \circ \underline{\kappa}\|_{\text{op},q} < \infty$ for every $L \in \mathcal{Y}^*$.*

Proof. By Remark 7 and Lemma 10, H has a formal convolution representation with some $\underline{\kappa} \in L(\mathcal{Z}, \mathcal{Y})^{\mathbb{Z}_-}$. Since $\underline{\mathcal{Z}}$ contains the unit ball of its span, the unique linear extension \tilde{H} of H to the span of $\underline{\mathcal{Z}}$ is p -continuous. Denote $\mathcal{V} = \ell^p(\underline{\mathcal{Z}})$ if $p < \infty$, respectively $\mathcal{V} = c_0(\mathcal{Z})$ if $p = \infty$. Let $L \in \mathcal{Y}^*$. Since $\text{span}(\underline{\mathcal{Z}})$ is dense in $(\mathcal{V}, \|\cdot\|_p)$ by Assumption 1.(i), $L \circ \tilde{H}$ is the restriction to $\text{span}(\underline{\mathcal{Z}})$ of an element in the dual space of $(\mathcal{V}, \|\cdot\|_p)$.

¹If $\mathcal{Y} = \mathcal{Z} = \mathbb{R}$, then this lemma has an algebraic interpretation. Equip the space of linear functionals $\text{span}(\underline{\mathcal{Z}}) \rightarrow \mathbb{R}$ with the weak* topology, and let \mathcal{V} be the largest linear subspace thereof for which the projections form a Schauder basis. By definition, a linear functional $\underline{\mathcal{Z}} \rightarrow \mathbb{R}$ has a formal convolution representation if and only if it is the restriction to $\underline{\mathcal{Z}}$ of an element in \mathcal{V} . If \mathcal{H} denotes the set of linear functionals $\underline{\mathcal{Z}} \rightarrow \mathbb{R}$ that have the minimal FMP and are minimally continuous, then Lemma 10 states that $\mathcal{H} = \{H|_{\underline{\mathcal{Z}}}: H \in \mathcal{V}\}$. Lemma 16 further below implies that the projections form an unconditional Schauder basis of \mathcal{V} .

It is a classical result that $\underline{\lambda} \mapsto (\underline{z} \mapsto \sum_{t \leq 0} \underline{\lambda}_t(\underline{z}_t))$ is an isomorphism $\ell^q(\mathcal{Z}^*, \|\cdot\|_{\text{op}}) \rightarrow (\mathcal{V}, \|\cdot\|_p)^*$. Thus, there exists an element $\underline{\Delta} \in \ell^q(\mathcal{Z}^*, \|\cdot\|_{\text{op}})$ such that $L \circ \tilde{H}(\underline{z}) = \sum_{t \leq 0} \underline{\Delta}_t(\underline{z}_t)$. This can only be if $L \circ \underline{\kappa}_t = \underline{\Delta}_t$ for all $t \in \mathbb{Z}_-$. In particular, $\|L \circ \underline{\kappa}\|_{\text{op},q} = \|\underline{\Delta}\|_{\text{op},q} < \infty$. \blacksquare

Ideally, we would like to conclude $\|\underline{\kappa}\|_{\text{op},q} < \infty$ without the dual elements $L \in \mathcal{Y}^*$. The full strength of Proposition 14 unfolds when the vector space \mathcal{Y} is finite-dimensional.

Remark 15. Suppose \mathcal{Y} is finite-dimensional. Let $q \in [1, \infty]$. If $\underline{\kappa} \in L(\mathcal{Z}, \mathcal{Y})^{\mathbb{Z}_-}$ satisfies $\|L \circ \underline{\kappa}\|_{\text{op},q} < \infty$ for every $L \in \mathcal{Y}^*$, then $\|\underline{\kappa}\|_{\text{op},q} < \infty$. This follows from the fact that any two norms on a finite-dimensional vector space are equivalent. Explicitly, if $d \in \mathbb{N}$ denotes the dimension of \mathcal{Y} , then there exist $L_1, \dots, L_d \in \mathcal{Y}^*$ and a constant $C > 0$ such that $\|\lambda\|_{\text{op}} \leq C \sum_{j=1}^d \|L_j \circ \lambda\|_{\text{op}}$ for any $\lambda \in L(\mathcal{Z}, \mathcal{Y})$. Therefore, $\|\underline{\kappa}\|_{\text{op},q} \leq Cd \sum_{j=1}^d \|L_j \circ \underline{\kappa}\|_{\text{op},q} < \infty$ by Jensen's inequality.

Sticking to a finite-dimensional codomain, we can upgrade Lemma 10 and prove that the minimal FMP and minimal continuity are equivalent to having a proper convolution representation. The first part of Lemma 16 is a technical result that we will need in some of the proofs to come.

Lemma 16. *Suppose \mathcal{Y} is finite-dimensional with dimension $d \in \mathbb{N}$. Consider a linear functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ that has a formal convolution representation. There exists a constant $c > 0$ independent of H such that for all $\underline{z} \in \underline{\mathcal{Z}}$ there exist disjoint subsets $J_1, \dots, J_{2^d} \subseteq \mathbb{Z}_-$ with*

$$\sum_{t \leq 0} \|\underline{\kappa}_t(\underline{z}_t)\|_{\mathcal{Y}} \leq c \sum_{i=1}^{2^d} \left\| H \left(\sum_{t \in J_i} \delta^t(\underline{z}_t) \right) \right\|_{\mathcal{Y}} < \infty.$$

In particular, the convolution representation of H is proper.

Proof. We make the following claim. There exists a constant $c > 0$ and there exists a partition of \mathcal{Y} into cones C_1, \dots, C_{2^d} intersecting pairwise in the origin, that is, $\bigcup_i C_i = \mathcal{Y}$ and $C_i \cap C_j = \{0\}$ for any $i \neq j$, such that for any $i \in \{1, \dots, 2^d\}$, any $N \in \mathbb{N}$, and any $v_1, \dots, v_N \in C_i$ we have $\sum_{n=1}^N \|v_n\|_{\mathcal{Y}} \leq c \left\| \sum_{n=1}^N v_n \right\|_{\mathcal{Y}}$. Now, fix $\underline{z} \in \underline{\mathcal{Z}}$. Consider $J_i = \{t \in \mathbb{Z}_- : \underline{\kappa}_t(\underline{z}_t) \in C_i \setminus \{0\}\}$ and $\underline{z}^i = \sum_{t \in J_i} \delta^t(\underline{z}_t)$, which belongs to $\underline{\mathcal{Z}}$ by Assumption 1.(ii). Then,

$$\sum_{t \in J_i} \|\underline{\kappa}_t(\underline{z}_t)\|_{\mathcal{Y}} = \liminf_{T \rightarrow -\infty} \sum_{t=T}^0 \|\underline{\kappa}_t(\underline{z}_t^i)\|_{\mathcal{Y}} \leq \liminf_{T \rightarrow -\infty} c \left\| \sum_{t=T}^0 \underline{\kappa}_t(\underline{z}_t^i) \right\|_{\mathcal{Y}} = c \|H(\underline{z}^i)\|_{\mathcal{Y}}.$$

Thus, $\sum_{t \leq 0} \|\underline{\kappa}_t(\underline{z}_t)\|_{\mathcal{Y}} \leq c \sum_{i=1}^{2^d} \|H(\underline{z}^i)\|_{\mathcal{Y}} < \infty$. It remains to prove the claim. It suffices to show it for \mathbb{R}^d with the Euclidean norm since all norms on a finite-dimensional vector space are equivalent. Let C_1, \dots, C_{2^d} be the orthants. The boundaries can be assigned in any way so that $\bigcup_i C_i = \mathbb{R}^d$ and $C_i \cap C_j = \{0\}$. Consider the first orthant C_1 containing the vector $v_0 = (1, \dots, 1)$. Let $\pi: C_1 \rightarrow \{rv_0 : r \geq 0\}$ be the orthogonal projection onto the ray of v_0 . Then, $\|\pi(v)\|_{\mathbb{R}^d} \leq \|v\|_{\mathbb{R}^d}$ for any $v \in C_1$. Also, $\|v\|_{\mathbb{R}^d} \leq \sqrt{2} \|\pi(v)\|_{\mathbb{R}^d}$ because the angle between v and v_0 is at most $\pi/4$. Furthermore, $\|v + w\|_{\mathbb{R}^d} = \|v\|_{\mathbb{R}^d} + \|w\|_{\mathbb{R}^d}$ for any v, w in the ray. Thus,

$$\sum_{n=1}^N \|v_n\|_{\mathbb{R}^d} \leq \sqrt{2} \sum_{n=1}^N \|\pi(v_n)\|_{\mathbb{R}^d} = \sqrt{2} \left\| \sum_{n=1}^N \pi(v_n) \right\|_{\mathbb{R}^d} = \sqrt{2} \left\| \pi \left(\sum_{n=1}^N v_n \right) \right\|_{\mathbb{R}^d} \leq \sqrt{2} \left\| \sum_{n=1}^N v_n \right\|_{\mathbb{R}^d}.$$

By symmetry, the same holds for the other orthants. \blacksquare

Now, we present our main result – a significant generalization of the original convolution theorem [2, Theorem 5]. In the formulation of Theorem 17, the minimal FMP and minimal continuity may be substituted by having a convolution representation, as proved in Lemmas 10 and 16.

Theorem 17 (Convolution Theorem). *Suppose \mathcal{Y} is finite-dimensional. Let $p \in [1, \infty]$, and suppose that $c_0(\mathcal{Z}) \cap \mathbb{B}^p(\mathcal{Z}) \subseteq \underline{\mathcal{Z}} \subseteq \ell^p(\mathcal{Z})$. Consider a linear functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$.²*

²If $\mathcal{Y} = \mathcal{Z} = \mathbb{R}$, then this result admits an algebraic interpretation as did Lemma 10. Suppose $\underline{\mathcal{Z}} = c_0(\mathcal{Z}) \cap \ell^p(\mathbb{R})$. Let \mathcal{V} be the largest linear subspace of the space of linear functionals $\underline{\mathcal{Z}} \rightarrow \mathbb{R}$ for which the projections form a Schauder basis in the weak* topology. It is well-known that $\underline{\mathcal{Z}}^* \subseteq \mathcal{V}$ [15]. From Theorem 17, we deduce the reverse inclusion $\mathcal{V} \subseteq \underline{\mathcal{Z}}^*$.

- (i) Suppose $p = 1$. Then, H has the 1-weighted FMP if and only if it has a convolution representation satisfying $\lim_{t \rightarrow -\infty} \|\underline{\kappa}_t\|_{\text{op}} = 0$. Furthermore, H is 1-continuous if and only if it has a convolution representation satisfying $\|\underline{\kappa}\|_{\text{op},\infty} < \infty$ if and only if it has the minimal FMP and is minimally continuous.
- (ii) Suppose $p \in (1, \infty)$, and let $q \in (1, \infty)$ be the Hölder conjugate of p . Then, H has the p -weighted FMP if and only if it is p -continuous if and only if it has a convolution representation satisfying $\|\underline{\kappa}\|_{\text{op},q} < \infty$ if and only if it has the minimal FMP and is minimally continuous.
- (iii) Suppose $p = \infty$. Then, H has the ∞ -weighted FMP if and only if it has a convolution representation satisfying $\|\underline{\kappa}\|_{\text{op},1} < \infty$ if and only if it has the minimal FMP and is minimally continuous. Furthermore, if $\underline{\mathcal{Z}} \subseteq c_0(\mathcal{Z})$, then H has the ∞ -weighted FMP if and only if it is ∞ -continuous.

Proof. (i) The first part is Proposition 12. The second part follows from Lemma 8, Lemma 10, Proposition 14, and Remark 15.

(ii) This follows from Remark 7, Lemma 8, Lemma 10, Proposition 14, and Remark 15.

(iii) It follows from Remark 7 and Lemma 8 that a convolution representation satisfying $\|\underline{\kappa}\|_{\text{op},1} < \infty$ implies the ∞ -weighted FMP, which in turn implies the minimal FMP and minimal continuity. We know from Lemmas 10 and 16 that the minimal FMP and minimal continuity imply a convolution representation. We show that if H has a convolution representation, then it must satisfy $\|\underline{\kappa}\|_{\text{op},1} < \infty$. Suppose this was not true. Then, there exists a weighting sequence \underline{w} such that $\sum_{t \leq 0} \underline{w}_t \|\underline{\kappa}_t\|_{\text{op}} = \infty$. For any $t \in \mathbb{Z}_-$, take $z_t \in \mathcal{Z}$ with norm one and such that $\|\underline{\kappa}_t\|_{\text{op}} \leq 2 \|\underline{\kappa}_t(z_t)\|_{\mathcal{Y}}$. Let $\underline{z} = \sum_{t \leq 0} \delta^t(\underline{w}_t z_t) \in c_0(\mathcal{Z}) \cap \mathbb{B}^\infty(\mathcal{Z})$. Then, $\sum_{t \leq 0} \underline{w}_t \|\underline{\kappa}_t\|_{\text{op}} \leq 2 \sum_{t \leq 0} \|\underline{\kappa}_t(\underline{z}_t)\|_{\mathcal{Y}} < \infty$ since the convolution representation is proper – a contradiction. Now, assume $\underline{\mathcal{Z}} \subseteq c_0(\mathcal{Z})$. We know from Remark 7 that the ∞ -weighted FMP implies ∞ -continuity and the latter implies the minimal FMP and minimal continuity, which we already showed to be equivalent to the ∞ -weighted FMP. \blacksquare

Remark 18. If $p \in (1, \infty)$ or $p = \infty$ and $\underline{\mathcal{Z}} \subseteq c_0(\mathcal{Z})$, then p -continuity and the p -weighted FMP turn out to be equivalent for linear functionals. The cases $p = 1$ and $p = \infty$ with $\underline{\mathcal{Z}} \not\subseteq c_0(\mathcal{Z})$ are special. We see from Theorem 17.(i) that any convolution representation with $\|\underline{\kappa}\|_{\text{op},\infty} < \infty$ and $\limsup_{t \rightarrow -\infty} \|\underline{\kappa}_t\|_{\text{op}} > 0$ provides an example of a linear functional that is 1-continuous but does not have the 1-weighted FMP. If $\underline{\mathcal{Z}} \not\subseteq c_0(\mathcal{Z})$, then the Banach limit from Example 11 poses an example of a linear functional that is ∞ -continuous but does not have the ∞ -weighted FMP. Theorem 17 also exposes the existence of linear functionals with the p -weighted FMP that are not q -continuous for any $1 \leq p < q \leq \infty$.

Example 19. Consider the linear functional $H^\omega: \ell^\infty(\mathbb{R}) \rightarrow \mathbb{R}$ with the convolution representation $\underline{\kappa}_t^\omega = (1-t)^{1+\omega}$, where $\omega \in (0, \infty)$. Then, $\|\underline{\kappa}^\omega\|_{\text{op},1} < \infty$ and, hence, H^ω has the ∞ -weighted FMP. This example has been studied to illustrate the so-called curse of memory [16]. The weighting sequence \underline{w} for which H^ω is continuous with respect to $\|\cdot\|_{w,\infty}$ captures the rate of decay of the memory of H^ω . It has to satisfy $\sum_{t \leq 0} \underline{w}_t^{-1} |\underline{\kappa}_t^\omega| < \infty$, which excludes any weighting sequence with exponential decay. This is in line with the findings in [16].

If $c_0(\mathcal{Z}) \cap \mathbb{B}^\infty(\mathcal{Z}) \subseteq \underline{\mathcal{Z}} \subseteq \mathbb{B}^\infty(\mathcal{Z})$, then the product FMP admits the same characterization Theorem 17.(iii) as the ∞ -weighted FMP by Remark 7. If the domain $\underline{\mathcal{Z}}$ is larger, then the product FMP is a strictly stronger notion than the ∞ -weighted FMP. In this case, the product FMP turns out to be equivalent to having finite memory. Let $c_{00}(\mathcal{Z}) \subseteq \mathcal{Z}^{\mathbb{Z}_-}$ denote the set of all sequences with at most finitely many non-zero entries.

Proposition 20. Suppose \mathcal{Y} is finite-dimensional and $c_{00}(\mathcal{Z}) \subseteq \underline{\mathcal{Z}}$. Consider a linear functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$.

- (i) The functional H has the product FMP if and only if it has finite memory.
- (ii) If $\underline{\mathcal{Z}} = \mathcal{Z}^{\mathbb{Z}_-}$, then H has the product FMP if and only if it has finite memory if and only if it has the minimal FMP and is minimally continuous.

Proof. (i) It is clear that finite memory implies the product FMP. Conversely, we know from Remark 7 and Lemma 10 that the product FMP implies a formal convolution representation with some $\underline{\kappa} \in L(\mathcal{Z}, \mathcal{Y})^{\mathbb{Z}_-}$. Suppose for contradiction H has infinite memory. Then, there exists a monotone sequence $(t_n)_n \subseteq \mathbb{Z}_-$ such that $\underline{\kappa}_{t_n} \neq 0$ for all $n \in \mathbb{N}$. For any $n \in \mathbb{N}$, take $z_n \in \mathcal{Z}$ such that $\|\underline{\kappa}_{t_n}(z_n)\|_{\mathcal{Y}} = 1$. Consider $\underline{z}^N = \sum_{n \leq N} \delta^{t_n}(z_n)$. By Lemma 16, for any $N \in \mathbb{N}$, there exist $J_1^N, \dots, J_{2^d}^N \subseteq \mathbb{Z}_-$ such that

$$1 = \frac{1}{N} \sum_{n \leq N} \|\underline{\kappa}_{t_n}(z_n)\|_{\mathcal{Y}} = \frac{1}{N} \sum_{t \leq 0} \|\underline{\kappa}_t(\underline{z}^N)\|_{\mathcal{Y}} \leq \frac{c}{N} \sum_{i=1}^{2^d} \left\| H \left(\sum_{t \in J_i^N} \delta^t(\underline{z}^N) \right) \right\|_{\mathcal{Y}}.$$

But $\frac{1}{N} \sum_{t \in J_i^N} \delta^t(\underline{z}_i^N)$ converges to the zero sequence $\underline{0}$ in the product topology. By the product FMP, $H(\frac{1}{N} \sum_{t \in J_i^N} \delta^t(\underline{z}_i^N)) \rightarrow H(\underline{0}) = 0$, a contradiction.

(ii) We need only show that the minimal FMP and minimal continuity imply finite memory. We know from Lemmas 10 and 16 that H has a convolution representation with some $\underline{\kappa} \in L(\mathcal{Z}, \mathcal{Y})^{\mathbb{Z}_-}$. Let J be the set of all $t \in \mathbb{Z}_-$ with $\underline{\kappa}_t \neq 0$. For any $t \in J$, take $z_t \in \mathcal{Z}$ such that $\|\underline{\kappa}_t(z_t)\|_{\mathcal{Y}} = 1$. This time, consider $\underline{z} = \sum_{t \in J} \delta^t(z_t)$. Then, since the convolution representation is proper, $\#J = \sum_{t \in J} \|\underline{\kappa}_t(z_t)\|_{\mathcal{Y}} = \sum_{t \leq 0} \|\underline{\kappa}_t(\underline{z}_t)\|_{\mathcal{Y}} < \infty$. ■

Remark 21. The finite-dimensionality of \mathcal{Y} in Theorem 17 and Proposition 20 is necessary. Indeed, if $\mathcal{Y} = \ell^p(\mathcal{Z})$ is equipped with the topology induced by $\|\cdot\|_p$, then the identity map $\ell^p(\mathcal{Z}) \rightarrow \ell^p(\mathcal{Z})$ is a p -continuous linear functional without the p -weighted FMP. If $\mathcal{Y} = \ell^p(\mathcal{Z})$ is equipped with the subspace product topology, then the identity map has the product FMP and infinite memory.

Example 22. Extend the linear state equation from Example 9 to a state-space system with a readout $y_t = h(\underline{x}_t)$, where $h \in L(\mathcal{X}, \mathbb{R})$. The associated functional becomes $H_{A,C}^h = h \circ H_{A,C}: \ell^\infty(\mathbb{R}^d) \rightarrow \mathbb{R}$. We have seen in Example 9 that a sufficient condition for the ∞ -weighted FMP is A having spectral radius strictly smaller than 1. If A is nilpotent, then $H_{A,C}^h$ has finite memory and, hence, the product FMP. Conversely, as soon as $H_{A,C}^h$ has even the minimal FMP, we must have $\sum_{t \leq 0} \|h \circ A^{-t} \circ C\|_{\text{op}} < \infty$.

Remark 23. Any linear functional $H: \ell^\infty(\mathcal{Z}) \rightarrow \mathcal{Y}$ with the ∞ -weighted FMP can be realized by a linear state-space system [17]. If H has the product FMP and, hence, finite memory, then the state operator of the system can be taken nilpotent. If, in addition, \mathcal{Z} and \mathcal{Y} are finite-dimensional, then the state-space system can be constructed on a finite-dimensional state space.

Example 24. Let $p, q \in [1, \infty]$ be Hölder conjugates. Consider the linear state equation from Example 9 with $\mathcal{X} = \ell^q(\mathbb{R})$, and assume the solutions to the state equation belong to $\ell^\infty(\ell^q(\mathbb{R}))$. Let $T: \ell^\infty(\mathbb{R}^d) \rightarrow \ell^\infty(\mathbb{R}^d)$ denote the shift operator, and let $\sigma: \mathbb{R} \rightarrow \mathbb{R}$ be Lipschitz continuous. Let μ be a Borel probability measure on $\Sigma = \mathbb{R} \times \ell^p(\mathbb{R}) \times \mathbb{R}^d \times \mathbb{R}$ satisfying $\int_\Sigma |w| (\|a\|_p + \|c\|_{\mathbb{R}^d} + |b|) d\mu(w, a, c, b) < \infty$. Consider the functional $H: \ell^\infty(\mathbb{R}^d) \rightarrow \mathbb{R}$ given by

$$H(\underline{z}) = \int_\Sigma w \sigma(a \circ H_{A,C} \circ T(\underline{z}) + c \cdot \underline{z}_0 + b) d\mu(w, a, c, b),$$

where $a \in \ell^p(\mathbb{R})$ is identified with the induced dual element in $\ell^q(\mathbb{R})^*$. Such a functional is a recurrent generalized Barron functional [8]. It is nonlinear in general, but it builds on the linear functional $H_{A,C}$. In particular, if $\sum_{t \leq 0} \|A^{-t} \circ C\|_{\text{op}} < \infty$, then H has the ∞ -weighted FMP.

4. Kernel learning with linear functionals

Suppose that the norms on \mathcal{Y} and \mathcal{Z} are induced by some inner products $\langle \cdot, \cdot \rangle_{\mathcal{Y}}$ and $\langle \cdot, \cdot \rangle_{\mathcal{Z}}$ that turn \mathcal{Y} and \mathcal{Z} into Hilbert spaces. Any (not necessarily linear) functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$ can be used as a feature map to induce a symmetric, positive semi-definite kernel $K_H: \underline{\mathcal{Z}} \times \underline{\mathcal{Z}} \rightarrow \mathbb{R}$ on $\underline{\mathcal{Z}}$ by $K_H(\underline{z}^1, \underline{z}^2) = \langle H(\underline{z}^1), H(\underline{z}^2) \rangle_{\mathcal{Y}}$. Any such kernel has an associated reproducing kernel Hilbert space [3] which, in the particular case of K_H , we specify in Proposition 25 below. This extends a result of [17].

Proposition 25. Consider a functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$. Let \mathcal{Y}_0 be the closure of $\text{span}(H(\underline{\mathcal{Z}}))$ in \mathcal{Y} , and let $(\mathbb{H}, \langle \cdot, \cdot \rangle_{\mathbb{H}})$ be the reproducing kernel Hilbert space associated to the kernel K_H . Then, the map $y \mapsto \langle y, H(\cdot) \rangle_{\mathcal{Y}}$ is an isometric isomorphism of Hilbert spaces $(\mathcal{Y}_0, \langle \cdot, \cdot \rangle_{\mathcal{Y}}) \rightarrow (\mathbb{H}, \langle \cdot, \cdot \rangle_{\mathbb{H}})$.

Proof. The reproducing kernel Hilbert space can be constructed as follows. Let \mathbb{H}_0 be the linear span of the set of maps $\{K_H(\underline{z}^0, \cdot) : \underline{\mathcal{Z}} \rightarrow \mathbb{R}, \underline{z} \mapsto K_H(\underline{z}^0, \underline{z}) : \underline{z}^0 \in \underline{\mathcal{Z}}\}$. Define an inner product $\langle \cdot, \cdot \rangle_{\mathbb{H}_0}$ on \mathbb{H}_0 by linearly extending $\langle K_H(\underline{z}^1, \cdot), K_H(\underline{z}^2, \cdot) \rangle_{\mathbb{H}_0} = K_H(\underline{z}^1, \underline{z}^2)$. Then, $(\mathbb{H}, \langle \cdot, \cdot \rangle_{\mathbb{H}})$ is the completion of $(\mathbb{H}_0, \langle \cdot, \cdot \rangle_{\mathbb{H}_0})$. Consider the map $\phi: (\text{span}(H(\underline{\mathcal{Z}})), \langle \cdot, \cdot \rangle_{\mathcal{Y}}) \rightarrow (\mathbb{H}_0, \langle \cdot, \cdot \rangle_{\mathbb{H}_0})$ given by $y \mapsto \langle y, H(\cdot) \rangle_{\mathcal{Y}}$. This is a well-defined linear bijection, by definition of \mathbb{H}_0 . For any $y_1 = H(\underline{z}^1)$ and $y_2 = H(\underline{z}^2)$, we have

$$\langle \phi(y_1), \phi(y_2) \rangle_{\mathbb{H}_0} = \langle K_H(\underline{z}^1, \cdot), K_H(\underline{z}^2, \cdot) \rangle_{\mathbb{H}_0} = K_H(\underline{z}^1, \underline{z}^2) = \langle y_1, y_2 \rangle_{\mathcal{Y}}.$$

By linear extension, this continues to hold for any $y_1, y_2 \in \text{span}(H(\underline{\mathcal{Z}}))$. Thus, ϕ is an isometric isomorphism of inner product spaces. As such, ϕ extends to the completions $\mathcal{Y}_0 \rightarrow \mathbb{H}$. ■

Next, we show that the reproducing kernel Hilbert space is isometrically isomorphic to $\ell^2(\mathcal{Z})^*$ for orthogonal linear functionals with the minimal FMP. Recall that $\ell^2(\mathcal{Z})^*$ is a Hilbert space with an inner product $\langle \cdot, \cdot \rangle_{\ell^2(\mathcal{Z})^*}$ induced by the inner product $\langle \underline{z}^1, \underline{z}^2 \rangle_{\ell^2(\mathcal{Z})} = \sum_{t \leq 0} \langle \underline{z}_t^1, \underline{z}_t^2 \rangle_{\mathcal{Z}}$ on $\ell^2(\mathcal{Z})$ and the Riesz representation theorem.

Proposition 26. *Suppose $\mathbb{B}^2(\mathcal{Z}) \subseteq \underline{\mathcal{Z}}$. Consider a linear functional $H: \underline{\mathcal{Z}} \rightarrow \mathcal{Y}$. Let \mathcal{Y}_0 be the closure of $\text{span}(H(\underline{\mathcal{Z}}))$ in \mathcal{Y} , and let \mathbb{H} be the reproducing kernel Hilbert space associated to the kernel K_H . Suppose H has the minimal FMP and is minimally continuous. Then, \mathbb{H} embeds linearly in $\ell^2(\mathcal{Z})^*$ by mapping a function $f \in \mathbb{H}$ to the unique linear extension of $f|_{\mathbb{B}^2(\mathcal{Z})}$, and $\text{span}(H(\mathbb{B}^2(\mathcal{Z})))$ is dense in \mathcal{Y}_0 . Furthermore, if the restriction $H: (\mathbb{B}^2(\mathcal{Z}), \langle \cdot, \cdot \rangle_{\ell^2(\mathcal{Z})}) \rightarrow (\mathcal{Y}, \langle \cdot, \cdot \rangle_{\mathcal{Y}})$ is orthogonal, then the embedding is an isometric isomorphism of Hilbert spaces $(\mathbb{H}, \langle \cdot, \cdot \rangle_{\mathbb{H}}) \rightarrow (\ell^2(\mathcal{Z})^*, \langle \cdot, \cdot \rangle_{\ell^2(\mathcal{Z})^*})$, and $\text{span}(H(\mathbb{B}^2(\mathcal{Z}))) = \mathcal{Y}_0$.*

Proof. By Lemmas 8 and 10, the restriction of H to $\mathbb{B}^2(\mathcal{Z})$ is 2-continuous and, hence, admits a unique 2-continuous linear extension $\tilde{H}: \ell^2(\underline{\mathcal{Z}}) \rightarrow \mathcal{Y}$. By Proposition 25, $y \mapsto H_y = \langle y, H(\cdot) \rangle_{\mathcal{Y}}$ is an isomorphism $\mathcal{Y}_0 \rightarrow \mathbb{H}$. We show that the map $\phi: \mathcal{Y}_0 \rightarrow \ell^2(\mathcal{Z})^*$ given by $y \mapsto \langle y, \tilde{H}(\cdot) \rangle_{\mathcal{Y}}$ is injective. Suppose $\langle y, \tilde{H}(\cdot) \rangle_{\mathcal{Y}} = 0$. Then, H_y is constantly zero on $\mathbb{B}^2(\mathcal{Z})$. The minimal FMP and minimal continuity of H transfer to H_y . By Lemma 10, H_y has a formal convolution representation $\underline{\kappa}$. This $\underline{\kappa}$ must be trivial since it is uniquely determined by the restriction of H_y to $\mathbb{B}^2(\mathcal{Z})$. Thus, H_y is constantly zero on all of $\underline{\mathcal{Z}}$, which can only be if $y = 0$. This shows injectivity of ϕ . Note that the orthogonal complement of $\tilde{H}(\ell^2(\mathcal{Z}))$ in \mathcal{Y}_0 belongs to the kernel of ϕ and, hence, is trivial. Having a trivial orthogonal complement is equivalent to density of $\tilde{H}(\ell^2(\mathcal{Z}))$ in \mathcal{Y}_0 .

Now, assume \tilde{H} is orthogonal. Then, $\tilde{H}(\ell^2(\mathcal{Z}))$ is complete. As it is also dense in \mathcal{Y}_0 , we must have $\tilde{H}(\ell^2(\mathcal{Z})) = \mathcal{Y}_0$. Given any $y_1, y_2 \in \mathcal{Y}_0$, take $\underline{z}^{y_1}, \underline{z}^{y_2} \in \ell^2(\mathcal{Z})$ with $\tilde{H}(\underline{z}^{y_i}) = y_i$. By orthogonality of \tilde{H} , $\phi(y_i) = \langle \underline{z}^{y_i}, \cdot \rangle_{\ell^2(\mathcal{Z})}$ and, hence,

$$\langle \phi(y_1), \phi(y_2) \rangle_{\ell^2(\mathcal{Z})^*} = \langle \underline{z}^{y_1}, \underline{z}^{y_2} \rangle_{\ell^2(\mathcal{Z})} = \langle \tilde{H}(\underline{z}^{y_1}), \tilde{H}(\underline{z}^{y_2}) \rangle_{\mathcal{Y}} = \langle y_1, y_2 \rangle_{\mathcal{Y}}.$$

This shows that ϕ is an isometry. To see that ϕ is surjective, let $f \in \ell^2(\mathcal{Z})^*$. By the Riesz representation theorem, there exists some $\underline{z}^0 \in \ell^2(\mathcal{Z})$ such that $f = \langle \underline{z}^0, \cdot \rangle_{\ell^2(\mathcal{Z})}$. Then, $f = \phi(\tilde{H}(\underline{z}^0))$. ■

As a by-product, we can prove the following functional analytic result.

Corollary 27. *Suppose $\underline{\mathcal{Z}}$ is a normed vector space that contains $\ell^2(\mathcal{Z})$ as a non-dense subset. Then, there exists an element $\underline{z} \in \underline{\mathcal{Z}}$ such that the truncated sequence $\sum_{t=T}^0 \delta^t(\underline{z}_t)$ does not converge to \underline{z} as $T \rightarrow -\infty$.*

Proof. The Hahn-Banach theorem guarantees the existence of a non-trivial, continuous, linear functional $H: \underline{\mathcal{Z}} \rightarrow \mathbb{R}$ that vanishes on $\ell^2(\mathcal{Z})$. In particular, $H(\ell^2(\mathcal{Z})) = 0$ is not dense in $H(\underline{\mathcal{Z}}) = \mathbb{R}$. By Proposition 26, H cannot have the minimal FMP. That is, there exists an element $\underline{z} \in \underline{\mathcal{Z}}$ such that $H(\sum_{t=T}^0 \delta^t(\underline{z}_t))$ does not converge to $H(\underline{z})$. Since H is continuous, it follows that $\sum_{t=T}^0 \delta^t(\underline{z}_t)$ cannot converge to \underline{z} . ■

Suppose we are given finitely many points $\underline{z}^i \in \underline{\mathcal{Z}}$, $i = 1, \dots, M$, a strictly increasing function $\Lambda: [0, \infty) \rightarrow [0, \infty)$, playing the role of a regularizer, and a function $\mathcal{E}: \mathbb{R}^M \rightarrow \mathbb{R}$, which is typically an empirical risk. Consider the (regularized empirical risk) minimization problem

$$\inf_{f \in \mathbb{H}} \mathcal{E}(f(\underline{z}^1), \dots, f(\underline{z}^M)) + \Lambda(\|f\|_{\mathbb{H}}). \quad (2)$$

The representer theorem [3] states that any minimizer \hat{f} , if existent, takes the form $\hat{f} = \sum_{i=1}^M \alpha_i \langle H(\underline{z}^i), H(\cdot) \rangle_{\mathcal{Y}}$ for some $\alpha_1, \dots, \alpha_M \in \mathbb{R}$. This is a well-known method to reduce an a priori infinite-dimensional minimization problem to a finite-dimensional one [32]. The coefficients α_i can be computed approximately as the solution of a regression problem, and they can be computed exactly out of the evaluations $\langle H(\underline{z}^i), H(\underline{z}^j) \rangle_{\mathcal{Y}}$ if \mathcal{E} is a mean square error and Λ a Tikhonov regularizer [6, 36]. However, evaluating $\langle H(\underline{z}^i), H(\underline{z}^j) \rangle_{\mathcal{Y}}$ still involves infinite sequences in general and can only be done approximately. To circumvent this, we replace the samples \underline{z}^i by their truncated counterparts $\tau_T(\underline{z}^i) = \sum_{t=T}^0 \delta^t(\underline{z}_t^i)$ for some fixed $T \leq 0$ [6, 17]. Solving for $\alpha_1, \dots, \alpha_M$ using this truncation gives rise to an approximate minimizer \hat{f}_T . Under an orthogonality assumption on H , \hat{f}_T admits another interpretation, as we show next.

Proposition 28. *Consider a linear functional H . Let $(\mathbb{H}, \langle \cdot, \cdot \rangle_{\mathbb{H}})$ be the reproducing kernel Hilbert space associated to the kernel K_H , let $T \in \mathbb{Z}_-$, and let $\mathbb{H}_T \subseteq \mathbb{H}$ be the subspace of all functionals $\underline{z} \mapsto f(\underline{z})$*

that depend only on $\underline{z}_T, \dots, \underline{z}_0$. Suppose H has the minimal FMP and is minimally continuous and that $\text{span}(H \circ \delta^t(\mathcal{B}))$, $t \leq 0$, are orthogonal subspaces of \mathcal{Y} . Then,

$$\arg \min_{f \in \mathbb{H}} \mathcal{E}(f(\tau_T(\underline{z}^1)), \dots, f(\tau_T(\underline{z}^M))) + \Lambda(\|f\|_{\mathbb{H}}) = \arg \min_{f \in \mathbb{H}_T} \mathcal{E}(f(\underline{z}^1), \dots, f(\underline{z}^M)) + \Lambda(\|f\|_{\mathbb{H}}).$$

Proof. Let \mathcal{Y}_0 be the closure of $\text{span}(H(\underline{\mathcal{Z}}))$ in \mathcal{Y} . We know from Proposition 25 that $y \mapsto \langle y, H(\cdot) \rangle_{\mathcal{Y}}$ is an isometric isomorphism $\phi: (\mathcal{Y}_0, \langle \cdot, \cdot \rangle_{\mathcal{Y}}) \rightarrow (\mathbb{H}, \langle \cdot, \cdot \rangle_{\mathbb{H}})$. Denote $\mathcal{Y}_T = \bigoplus_{t=T}^0 \text{span}(H \circ \delta^t(\mathcal{B}))$. Since H has a formal convolution representation by Lemma 10, it follows that $H(\underline{z} - \tau_T(\underline{z}))$ belongs to the orthogonal complement \mathcal{Y}_T^\perp of \mathcal{Y}_T for any $\underline{z} \in \underline{\mathcal{Z}}$. Thus, for any $y = y_T + y^\perp \in \mathcal{Y}_T \oplus \mathcal{Y}_T^\perp$, we can write

$$\langle y, H(\underline{z}) \rangle_{\mathcal{Y}} = \langle y_T, H(\tau_T(\underline{z})) \rangle_{\mathcal{Y}} + \langle y^\perp, H(\underline{z} - \tau_T(\underline{z})) \rangle_{\mathcal{Y}},$$

which shows that $\phi(\mathcal{Y}_T) = \mathbb{H}_T$. Now, suppose $\hat{f}_T \in \arg \min_{f \in \mathbb{H}} \mathcal{E}'(f \circ \tau_T) + \Lambda(\|f\|_{\mathbb{H}})$, where we abbreviate $\mathcal{E}'(f) = \mathcal{E}(f(\underline{z}^1), \dots, f(\underline{z}^M))$. Take $y = y_T + y^\perp \in \mathcal{Y}_T \oplus \mathcal{Y}_T^\perp$ with $\phi(y) = \hat{f}_T$. Then, $\hat{f}_T \circ \tau_T = \phi(y_T) \circ \tau_T$ and, hence,

$$\mathcal{E}'(\phi(y_T) \circ \tau_T) + \Lambda\left(\sqrt{\|y_T\|_{\mathcal{Y}}^2 + \|y^\perp\|_{\mathcal{Y}}^2}\right) = \mathcal{E}'(\hat{f}_T \circ \tau_T) + \Lambda\left(\|\hat{f}_T\|_{\mathbb{H}}\right) \leq \mathcal{E}'(\phi(y_T) \circ \tau_T) + \Lambda(\|y_T\|_{\mathcal{Y}}).$$

Since Λ is strictly increasing, we must have $y^\perp = 0$. This shows that $\hat{f}_T \in \mathbb{H}_T$. That $\mathcal{E}'(f) = \mathcal{E}'(f \circ \tau_T)$ for any $f \in \mathbb{H}_T$ implies $\hat{f}_T \in \arg \min_{f \in \mathbb{H}_T} \mathcal{E}'(f) + \Lambda(\|f\|_{\mathbb{H}})$. Conversely, suppose $\hat{f}_T \in \arg \min_{f \in \mathbb{H}_T} \mathcal{E}'(f) + \Lambda(\|f\|_{\mathbb{H}})$. Then, for any $f = \phi(y_T + y^\perp) \in \mathbb{H}$,

$$\mathcal{E}'(\hat{f}_T) + \Lambda\left(\|\hat{f}_T\|_{\mathbb{H}}\right) \leq \mathcal{E}'(\phi(y_T)) + \Lambda(\|y_T\|_{\mathcal{Y}}) = \mathcal{E}'(f \circ \tau_T) + \Lambda(\|y_T\|_{\mathcal{Y}}) \leq \mathcal{E}'(f \circ \tau_T) + \Lambda(\|f\|_{\mathbb{H}}),$$

which shows that $\hat{f}_T \in \arg \min_{f \in \mathbb{H}} \mathcal{E}'(f \circ \tau_T) + \Lambda(\|f\|_{\mathbb{H}})$. ■

Proposition 28 says that minimizing (2) with truncated samples is equivalent to minimizing (2) over finite memory functionals. In particular, if $c_{00}(\mathcal{Z}) \subseteq \underline{\mathcal{Z}}$, then we minimize (2) over functionals with the product FMP, by Proposition 20. The dimension of \mathbb{H}_T is $\sum_{t=T}^0 \dim(\text{span}(H \circ \delta^t(\mathcal{B}))) \leq (|T| + 1) \times \dim(\mathcal{Z})$. If \mathcal{Z} is finite-dimensional, then minimizing directly over \mathbb{H}_T is itself a finite-dimensional problem. However, even in this case, the kernel trick reduces the dimension of the problem from up to $(|T| + 1) \times \dim(\mathcal{Z})$ down to M . More importantly, the kernel trick turns the optimization problem into a convex one.

Example 29. Suppose $\mathcal{Z} = \mathbb{R}^d$ and $\mathcal{X} = \mathcal{Y} = \ell^2(\mathbb{R}^d)$ with the standard inner products. Let $T^{-1}: \mathcal{X} \rightarrow \mathcal{X}$ denote the right-inverse $T^{-1}(\underline{z}) = (\dots, \underline{z}_{-1}, \underline{z}_0, 0)$ of the shift operator T . Given $\lambda \in (0, 1)$, consider the linear state equation $\underline{x}_t = \lambda T^{-1}(\underline{x}_{t-1}) + \delta^0(\underline{z}_t)$. The associated functional $H: \ell^\infty(\mathbb{R}^d) \rightarrow \ell^2(\mathbb{R}^d)$ is given by $H(\underline{z}) = \sum_{t \leq 0} \lambda^{-t} \delta^t(\underline{z}_t)$, see Example 9. The induced kernel is $K_H(\underline{z}^1, \underline{z}^2) = \sum_{t \leq 0} \lambda^{-2t} \langle \underline{z}_t^1, \underline{z}_t^2 \rangle_{\mathbb{R}^d}$, and it satisfies $K_H(\underline{z}^1, \underline{z}^2) = \langle \underline{z}_0^1, \underline{z}_0^2 \rangle_{\mathbb{R}^d} + \lambda^2 K_H(T(\underline{z}^1), T(\underline{z}^2))$. This recursion makes the kernel and its Gram matrix efficiently computable, which is a desirable property that only few induced kernels are known to have [6].

Let $(\mathbb{H}, \langle \cdot, \cdot \rangle_{\mathbb{H}})$ be the reproducing kernel Hilbert space associated to the kernel K_H . Since the image of H is dense in $\ell^2(\mathbb{R}^d)$, Proposition 25 conveys that $\ell^2(\mathbb{R}^d)$ is isometrically isomorphic to \mathbb{H} via $\underline{y} \mapsto \langle \underline{y}, H(\cdot) \rangle_{\ell^2(\mathbb{R}^d)}$. The functional H has the minimal FMP and is minimally continuous. The linear embedding of \mathbb{H} into $\ell^2(\mathbb{R}^d)^*$ promised by Proposition 26 is given by $\langle \underline{y}, H(\cdot) \rangle_{\ell^2(\mathbb{R}^d)} \mapsto \langle H(\underline{y}), \cdot \rangle_{\ell^2(\mathbb{R}^d)}$. However, this is not an isometry. Indeed, H is not orthogonal as a map. Nonetheless, the subspaces $H \circ \delta^t(\mathbb{R}^d) = \delta^t(\mathbb{R}^d)$, $t \leq 0$, are orthogonal in $\ell^2(\mathbb{R}^d)$, and Proposition 28 is applicable.

If we take $\ell^2(\mathbb{R}^d)$ as the domain of H instead of $\ell^\infty(\mathbb{R}^d)$, then we can permit $\lambda = 1$. In this case, H is the identity on $\ell^2(\mathbb{R}^d)$, which is trivially orthogonal. The isometric isomorphism $\ell^2(\mathbb{R}^d)^d \rightarrow \mathbb{H} \cong \ell^2(\mathbb{R}^d)^*$ from Proposition 25 becomes an instance of the Riesz representation theorem. The linear embedding $\mathbb{H} \hookrightarrow \ell^2(\mathbb{R}^d)^*$ from Proposition 26 is now simply the identity.

5. Conclusion

In this paper, we have conducted a detailed analysis of various types of continuity of linear functionals. The main result linked various so-defined fading memory properties to the availability of convolution representations. This greatly generalizes the classical convolution theorem that first appeared in [2]. The most striking features are, on the one hand, the central role of the two topological notions of minimal continuity and

the minimal FMP; and on the other hand continuity with respect to the product topology turns out to be equivalent to finite memory if the sequences in the input space are not uniformly bounded. This highlights the importance of giving careful consideration to which notion of fading memory to use in one's context. The strength of the two topological notions of minimal continuity and minimal fading memory have been further leveraged to embed reproducing kernel Hilbert spaces associated to such linear functionals defined on Hilbert spaces. Under an additional orthogonality assumption, it is shown that kernel learning with truncated samples is equivalent to kernel learning with finite memory functionals.

An immediate question is whether an analogous convolution theorem holds in a continuous-time setting, in which the inputs are not sequences indexed by integers but functions of the (half-axis of the) real line. Parts of our convolution theorem are based on the duality of the ℓ^p -sequence spaces. We expect these results to transfer easily to the continuous-time setting since analogous duality results hold for the L^p -function spaces. However, the continuous-time and the discrete-time case are not conceptually equivalent in all aspects. For example, the product FMP exhibits fundamentally different behavior in continuous-time. We have seen in Proposition 20 that the product FMP is equivalent to finite memory in discrete-time. In continuous-time, the product FMP and the availability of a convolution representation are incompatible. A proof of this can be found in Appendix B. Morally, the key difference is that any measure on the integers is absolutely continuous with respect to the counting measure. On the real line, on the other hand, functionals that are dual to measures that are not absolutely continuous with respect to the Lebesgue measure cannot admit a convolution representation but can still have fading memory. Thus, although some concepts from the discrete-time case carry over, a careful study of the continuous-time case is desirable.

The idea of having different notions of fading memory as well as using different weighting sequences in the concept of the p -weighted FMP lends itself to the study of the so-called curse of memory. This curse of memory has been approached in continuous-time in [16]. However, as we pointed out in the preceding paragraph, one ought to be careful in distinguishing continuous-time functionals with a convolution representation, which are the objects of interest in [16], from functionals with a stronger concept of fading memory dubbed ‘regularity’ therein. One method of attack to the curse of memory are lower and upper bounds on approximation rates. Another method of attack would be to study continuity of a functional with respect to a weighted ℓ^p -norm and at the same time discontinuity with respect to the ℓ^p -norm weighted by a different weighting sequence.

Finally, it is of great interest whether the ideas from this paper find applications in the study of non-linear functionals. For non-linear functionals, the analog of a convolution representation could be a Volterra series representation. Such representations have been proved to exist in [6, 11]. We hypothesize that some of the assumptions in those results can instead be replaced by careful fading memory assumptions.

Acknowledgments

The authors acknowledge partial financial support from the School of Physical and Mathematical Sciences of the Nanyang Technological University. The second author is funded by an Eric and Wendy Schmidt AI in Science Postdoctoral Fellowship at the Nanyang Technological University. We thank Lyudmila Grigoryeva and Qianxiao Li for helpful discussions and remarks.

Appendix A. Functionals and filters

We recall the equivalence between functionals and time-invariant filters [2, 11]. In this section, \mathcal{Y} and \mathcal{Z} are simply sets. Consider $\underline{\mathcal{Z}}^\pm \subseteq \mathcal{Z}^\mathbb{Z}$ and $\underline{\mathcal{Z}}^- \subseteq \mathcal{Z}^{\mathbb{Z}^-}$. Let $\text{CTI}(\underline{\mathcal{Z}}^\pm)$ and $\text{TI}(\underline{\mathcal{Z}}^-)$ denote the set of all causal and time-invariant filters $\underline{\mathcal{Z}}^\pm \rightarrow \mathcal{Y}^\mathbb{Z}$, respectively the set of all time-invariant filters $\underline{\mathcal{Z}}^- \rightarrow \mathcal{Y}^{\mathbb{Z}^-}$ (which are automatically causal). Let $\mathcal{H}(\underline{\mathcal{Z}}^-)$ denote the set of all functionals $\underline{\mathcal{Z}}^- \rightarrow \mathcal{Y}$. For clarity, we distinguish between the shift $T: \mathcal{Z}^\mathbb{Z} \rightarrow \mathcal{Z}^\mathbb{Z}$ on bi-infinite sequences and the shift $T_-: \mathcal{Z}^{\mathbb{Z}^-} \rightarrow \mathcal{Z}^{\mathbb{Z}^-}$ on semi-infinite sequences. Let $\tau_{\mathcal{Y}}: \mathcal{Y}^\mathbb{Z} \rightarrow \mathcal{Y}^{\mathbb{Z}^-}$ and $\tau_{\mathcal{Z}}: \mathcal{Z}^\mathbb{Z} \rightarrow \mathcal{Z}^{\mathbb{Z}^-}$ be the truncations, and let $p^t(\underline{y}) = \underline{y}_t$ be the projection onto the time entry t .

Lemma A.1. *Suppose $T(\underline{\mathcal{Z}}^\pm) = \underline{\mathcal{Z}}^\pm$ and $\tau_{\mathcal{Z}}(\underline{\mathcal{Z}}^\pm) = \underline{\mathcal{Z}}^-$. Let $\iota: \underline{\mathcal{Z}}^- \rightarrow \underline{\mathcal{Z}}^\pm$ be any right-inverse of $\tau_{\mathcal{Z}}$. Then,*

there is a commutative diagram of bijections

$$\begin{array}{ccc}
 \text{CTI}(\underline{\mathcal{Z}}^\pm) & \xrightarrow{V \mapsto U_V} & \text{TI}(\underline{\mathcal{Z}}^-) \\
 & \searrow V \mapsto H_V & \swarrow H_U \mapsto U \\
 & \mathcal{H}(\underline{\mathcal{Z}}^-) &
 \end{array}
 \quad
 \begin{array}{l}
 U_V = \tau_{\mathcal{Y}} \circ V \circ \iota, \\
 H_V = p^0 \circ V \circ \iota, \\
 H_U = p^0 \circ U.
 \end{array}$$

These bijections are independent of the choice of ι . Their inverses are given component-wise by

$$\begin{array}{ccc}
 \text{CTI}(\underline{\mathcal{Z}}^\pm) & \xleftarrow{V_U \mapsto U} & \text{TI}(\underline{\mathcal{Z}}^-) \\
 & \swarrow V_H \mapsto H & \searrow H \mapsto U_H \\
 & \mathcal{H}(\underline{\mathcal{Z}}^-) &
 \end{array}
 \quad
 \begin{array}{l}
 p^t \circ V_U = p^0 \circ U \circ \tau_{\mathcal{Z}} \circ T^{-t}, \\
 p^t \circ V_H = H \circ \tau_{\mathcal{Z}} \circ T^{-t}, \\
 p^t \circ U_H = H \circ T^{-t}.
 \end{array}$$

Proof. We have to verify that all maps are well-defined. First, observe that $T^{-t}(\underline{\mathcal{Z}}^-) = \underline{\mathcal{Z}}^-$ for any $t \in \mathbb{Z}_-$, which follows from the equality $\tau_{\mathcal{Z}} \circ T^{-t} = T^{-t} \circ \tau_{\mathcal{Z}}$ together with the assumptions that $\underline{\mathcal{Z}}^- = \tau_{\mathcal{Z}}(\underline{\mathcal{Z}}^\pm)$ and $T(\underline{\mathcal{Z}}^\pm) = \underline{\mathcal{Z}}^\pm$. Since $\mathcal{H}(\underline{\mathcal{Z}}^-)$ contains all maps $\underline{\mathcal{Z}}^- \rightarrow \mathcal{Y}$, it is clear that $V \mapsto H_V$ and $U \mapsto H_U$ are well-defined. Note that a filter $U: \underline{\mathcal{Z}}^- \rightarrow \mathcal{Y}^{\mathbb{Z}_-}$ is time-invariant if and only if $p^t \circ U \circ T^{-s} = p^{t+s} \circ U$ for any $t, s \in \mathbb{Z}_-$. From this, it is easy to see that $H \mapsto U_H$ maps into $\text{TI}(\underline{\mathcal{Z}}^-)$. Analogously, a filter $V: \underline{\mathcal{Z}}^\pm \rightarrow \mathcal{Y}^{\mathbb{Z}}$ is time-invariant if and only if $p^t \circ V \circ T^{-s} = p^{t+s} \circ V$ for any $t, s \in \mathbb{Z}$, from which it follows that V_U and V_H are time-invariant for any filter U and any functional H . Next, observe that a filter $V: \underline{\mathcal{Z}}^\pm \rightarrow \mathcal{Y}^{\mathbb{Z}}$ is causal if and only if $p^t \circ V(\underline{z}^1) = p^t \circ V(\underline{z}^2)$ for any $t \in \mathbb{Z}$ and $\underline{z}^1, \underline{z}^2 \in \underline{\mathcal{Z}}^\pm$ that satisfy $\tau_{\mathcal{Z}} \circ T^{-t}(\underline{z}^1) = \tau_{\mathcal{Z}} \circ T^{-t}(\underline{z}^2)$. From this, we find that V_U and V_H are also causal and, hence, that $U \mapsto V_U$ and $H \mapsto V_H$ map into $\text{CTI}(\underline{\mathcal{Z}}^\pm)$. It remains to show that U_V is time-invariant for any $V \in \text{CTI}(\underline{\mathcal{Z}}^\pm)$. The causality and time-invariance of V imply for any $s, t \in \mathbb{Z}_-$ that

$$p^t \circ V \circ \iota \circ T^{-s} = p^t \circ V \circ T^{-s} \circ \iota = p^{t+s} \circ V \circ \iota.$$

Clearly, $p^t \circ \tau_{\mathcal{Y}} = p^t$ for any $t \in \mathbb{Z}_-$. Thus,

$$p^t \circ U_V \circ T^{-s} = p^t \circ \tau_{\mathcal{Y}} \circ V \circ \iota \circ T^{-s} = p^t \circ V \circ \iota \circ T^{-s} = p^{t+s} \circ V \circ \iota = p^{t+s} \circ \tau_{\mathcal{Y}} \circ V \circ \iota = p^{t+s} \circ U_V,$$

which shows that $V \mapsto V_U$ maps into $\text{TI}(\underline{\mathcal{Z}}^-)$. We established that all maps are well-defined. To show that the three maps in the diagram are bijective, we verify that the formulas proposed as inverses are indeed left- and right-inverses. The map ι may not be a left-inverse of $\tau_{\mathcal{Z}}$, but the causality of an element $V \in \text{CTI}(\underline{\mathcal{Z}}^\pm)$ implies that $p^0 \circ V \circ \iota \circ \tau_{\mathcal{Z}} = p^0 \circ V$. Thus, for any $t \in \mathbb{Z}$ and $s \in \mathbb{Z}_-$,

$$\begin{aligned}
 p^t \circ V_{U_V} &= p^0 \circ U_V \circ \tau_{\mathcal{Z}} \circ T^{-t} = p^0 \circ \tau_{\mathcal{Y}} \circ V \circ \iota \circ \tau_{\mathcal{Z}} \circ T^{-t} = p^0 \circ V \circ T^{-t} = p^t \circ V, \\
 p^s \circ U_{V_U} &= p^s \circ \tau_{\mathcal{Y}} \circ V_U \circ \iota = p^s \circ V_U \circ \iota = p^0 \circ U \circ \tau_{\mathcal{Z}} \circ T^{-s} \circ \iota = p^0 \circ U \circ T^{-s} = p^s \circ U.
 \end{aligned}$$

Similarly, for any $t \in \mathbb{Z}$,

$$\begin{aligned}
 p^t \circ V_{H_V} &= H_V \circ \tau_{\mathcal{Z}} \circ T^{-t} = p^0 \circ V \circ \iota \circ \tau_{\mathcal{Z}} \circ T^{-t} = p^0 \circ V \circ T^{-t} = p^t \circ V, \\
 H_{V_H} &= p^0 \circ V_H \circ \iota = H \circ \tau_{\mathcal{Z}} \circ T^0 \circ \iota = H.
 \end{aligned}$$

Last but not least, for any $t \in \mathbb{Z}_-$,


$$\begin{aligned}
 p^t \circ U_{H_U} &= H_U \circ T^{-t} = p^0 \circ U \circ T^{-t} = p^t \circ U, \\
 H_{U_H} &= p^0 \circ U_H = H \circ T^0 = H.
 \end{aligned}$$

This shows that all three maps are bijections with inverses as claimed. Since the inverses do not depend on ι , neither do the original maps. Commutativity of the diagram amounts to the chain of equalities

$$H_{U_V} = p^0 \circ U_V = p^0 \circ \tau_{\mathcal{Y}} \circ V \circ \iota = p^0 \circ V \circ \iota = H_V.$$

■

Remark A.2. Let $\mathcal{F}(\underline{\mathcal{Z}}^\pm)$ and $\mathcal{F}(\underline{\mathcal{Z}}^-)$ be the set of all filters $\underline{\mathcal{Z}}^\pm \rightarrow \mathcal{Y}^\mathbb{Z}$, respectively $\underline{\mathcal{Z}}^- \rightarrow \mathcal{Y}^{\mathbb{Z}-}$. The filter V_U as defined in the previous lemma is causal and time-invariant for any $U \in \mathcal{F}(\underline{\mathcal{Z}}^-)$ even if U is not time-invariant, and the map $\text{TI}(\underline{\mathcal{Z}}^-) \rightarrow \text{CTI}(\underline{\mathcal{Z}}^\pm)$, $U \mapsto V_U$ extends to a map $\mathcal{F}(\underline{\mathcal{Z}}^-) \rightarrow \text{CTI}(\underline{\mathcal{Z}}^\pm)$. It is clear that $\text{CTI}(\underline{\mathcal{Z}}^\pm) \rightarrow \mathcal{H}(\underline{\mathcal{Z}}^-)$, $V \mapsto H_V$ also extends to a map $\mathcal{F}(\underline{\mathcal{Z}}^\pm) \rightarrow \mathcal{H}(\underline{\mathcal{Z}}^-)$, and likewise for $U \mapsto H_U$. Thus, we actually have a larger commutative diagram as follows.



The concatenations $\mathcal{F}(\underline{\mathcal{Z}}^\pm) \rightarrow \mathcal{F}(\underline{\mathcal{Z}}^\pm)$, $V \mapsto V_{H_V}$ and $\mathcal{F}(\underline{\mathcal{Z}}^-) \rightarrow \mathcal{F}(\underline{\mathcal{Z}}^-)$, $U \mapsto U_{H_U}$ are idempotent maps and define natural projections $\mathcal{F}(\underline{\mathcal{Z}}^\pm) \rightarrow \text{CTI}(\underline{\mathcal{Z}}^\pm)$ and $\mathcal{F}(\underline{\mathcal{Z}}^-) \rightarrow \text{TI}(\underline{\mathcal{Z}}^-)$ from the set of all filters onto the set of (causal) time-invariant ones.

Appendix B. Continuous-time linear functionals

In this appendix, we make precise a claim made in the Conclusions Section 5. Let $\mathbb{I} \subseteq \mathbb{R}$ be an interval with non-empty interior, and let $C_c^\infty(\mathbb{I}, \mathbb{R}^d)$ be the set of smooth and compactly supported functions $\mathbb{I} \rightarrow \mathbb{R}^d$. Recall that the topology of point-wise convergence on $C_c^\infty(\mathbb{I}, \mathbb{R}^d)$ is the subspace topology induced by the product topology on $(\mathbb{R}^d)^\mathbb{I}$.

Proposition B.1. *Let $\kappa: \mathbb{I} \rightarrow \mathbb{R}^d$ be locally integrable, and consider the linear functional $H: C_c^\infty(\mathbb{I}, \mathbb{R}^d) \rightarrow \mathbb{R}$,*

$$H(f) = \int_{\mathbb{I}} \kappa(t)^T f(t) dt.$$

Suppose H is continuous with respect to the topology of point-wise convergence. Then, κ vanishes Lebesgue-almost everywhere and, hence, H is constantly zero.

Proof. Suppose for contradiction that κ is non-zero on a set of positive Lebesgue measure. Consider the family of standard symmetric mollifiers $(\phi_\varepsilon)_{\varepsilon \in (0,1)} \subseteq C_c^\infty(\mathbb{R}, \mathbb{R})$, and recall that the support of each ϕ_ε is contained in $[-\varepsilon, +\varepsilon]$. Let $\|\cdot\|$ be the Euclidean norm. Since the convolution $\phi_\varepsilon * \kappa$ converges point-wise Lebesgue-almost everywhere to κ as $\varepsilon \rightarrow 0$, there exists a point t_0 in the interior of \mathbb{I} and constants $\varepsilon_0, \eta > 0$ such that $\|\phi_\varepsilon * \kappa(t_0)\| > \eta$ for all $\varepsilon \in (0, \varepsilon_0)$. Since $\phi_\varepsilon * \kappa$ is continuous, for all $\varepsilon \in (0, \varepsilon_0)$ there exists some $\delta_\varepsilon \in (0, \varepsilon)$ such that $\|\phi_\varepsilon * \kappa(t)\| > \eta$ for all $t \in (t_0 - 3\delta_\varepsilon, t_0)$. Let $\tau_\varepsilon \in C_c^\infty(\mathbb{R}, [0,1])$ be a bump function that is 1 in $(t_0 - 2\delta_\varepsilon, t_0 - \delta_\varepsilon)$ and is zero outside of $(t_0 - 3\delta_\varepsilon, t_0)$. Define $h_\varepsilon \in C_c^\infty(\mathbb{R}, \mathbb{R}^d)$ by $h_\varepsilon(t) = (\delta_\varepsilon)^{-1} \tau_\varepsilon(t) (\phi_\varepsilon * \kappa)(t)$, and set $g_\varepsilon = \phi_\varepsilon * h_\varepsilon \in C_c^\infty(\mathbb{R}, \mathbb{R}^d)$. The support of g_ε is contained in $[t_0 - 4\varepsilon, t_0 + \varepsilon]$. In particular, since t_0 is an interior point of \mathbb{I} , we have $g_\varepsilon \in C_c^\infty(\mathbb{I}, \mathbb{R}^d)$ for sufficiently small ε . Let us compute $H(g_\varepsilon)$. By Fubini's theorem and by symmetry of the mollifiers, that is, $\phi_\varepsilon(t-s) = \phi_\varepsilon(s-t)$, we have

$$H(g_\varepsilon) = \int_{\mathbb{I}} \kappa(t)^T (\phi_\varepsilon * h_\varepsilon)(t) dt = \int_{\mathbb{I}} \int_{\mathbb{R}} \kappa(t)^T \phi_\varepsilon(s-t) h_\varepsilon(s) ds dt = \int_{\mathbb{R}} (\phi_\varepsilon * \kappa)(s)^T h_\varepsilon(s) ds.$$

Unravelling the definitions, we find

$$H(g_\varepsilon) = \int_{\mathbb{R}} (\phi_\varepsilon * \kappa)(s)^T h_\varepsilon(s) ds = \frac{1}{\delta_\varepsilon} \int_{t_0 - 3\delta_\varepsilon}^{t_0} \tau_\varepsilon(s) \|(\phi_\varepsilon * \kappa)(s)\|^2 ds \geq \frac{\eta^2}{\delta_\varepsilon} \int_{t_0 - 3\delta_\varepsilon}^{t_0} \tau_\varepsilon(s) ds \geq \eta^2.$$

Next, for any $\varepsilon \in (0, \varepsilon_0)$, take $r_\varepsilon \in (0, \varepsilon)$ so small that

$$\int_{t_0 - 2r_\varepsilon}^{t_0 + 2r_\varepsilon} |\kappa(t)^T g_\varepsilon(t)| dt \leq \varepsilon.$$

Let $\hat{\tau}_\varepsilon \in C_c^\infty(\mathbb{R}, [0, 1])$ be a bump function that is 1 in $(t_0 - r_\varepsilon, t_0 + r_\varepsilon)$ and is zero outside of $(t_0 - 2r_\varepsilon, t_0 + 2r_\varepsilon)$. Then,

$$|H(\hat{\tau}_\varepsilon g_\varepsilon)| \leq \int_{\mathbb{I}} \hat{\tau}_\varepsilon(t) |\kappa(t)^T g_\varepsilon(t)| dt \leq \int_{t_0 - 2r_\varepsilon}^{t_0 + 2r_\varepsilon} |\kappa(t)^T g_\varepsilon(t)| dt \leq \varepsilon.$$

Finally, set $f_\varepsilon = (1 - \hat{\tau}_\varepsilon)g_\varepsilon \in C_c^\infty(\mathbb{R}, \mathbb{R}^d)$. Clearly, $f_\varepsilon(t_0) = 0$. That the support of f_ε is contained in $[t_0 - 4\varepsilon, t_0 + \varepsilon]$ implies that $(f_\varepsilon(t))_\varepsilon$ is eventually zero for any $t \neq t_0$. Thus, $H(f_\varepsilon) \rightarrow 0$ by continuity of H with respect to the topology of point-wise convergence. This contradicts $H(f_\varepsilon) = H(g_\varepsilon) - H(\hat{\tau}_\varepsilon g_\varepsilon) \geq \eta^2 - \varepsilon$. ■

Let $C_0(\mathbb{I}, \mathbb{R}^d)$ denote the set of continuous functions vanishing at infinity. Consider the set \mathcal{R} of linear functionals $H: C_0(\mathbb{I}, \mathbb{R}^d) \rightarrow \mathbb{R}$ that satisfy $H(f_n) \rightarrow 0$ for any sequence $(f_n)_n \subseteq C_0(\mathbb{I}, \mathbb{R}^d)$ that converges to zero point-wise Lebesgue-almost everywhere and is uniformly bounded in the sense that $\sup_n \sup_{t \in \mathbb{I}} \|f_n(t)\|_{\mathbb{R}^d} < \infty$. Note that any element in \mathcal{R} is continuous with respect to the supremums-norm. Thus, \mathcal{R} is a subset of the dual space $C_0(\mathbb{I}, \mathbb{R}^d)^*$. It can be shown that \mathcal{R} is exactly the set of functionals one obtains by embedding $L^1(\mathbb{I}, \mathbb{R}^d)$ in the dual $C_0(\mathbb{I}, \mathbb{R}^d)^*$ [16]. Proposition B.1 implies that the condition $\sup_n \sup_{t \in \mathbb{I}} \|f_n(t)\|_{\mathbb{R}^d} < \infty$ is crucial in the property above. Indeed, suppose $H: C_0(\mathbb{I}, \mathbb{R}^d) \rightarrow \mathbb{R}$ is a linear functional that satisfies $H(f_n) \rightarrow 0$ for any sequence $(f_n)_n \subseteq C_0(\mathbb{I}, \mathbb{R}^d)$ that converges to zero point-wise Lebesgue-almost everywhere but is permitted to be unbounded. Then, $H \in \mathcal{R} \cong L^1(\mathbb{I}, \mathbb{R}^d)$ admits an integral representation as required in Proposition B.1, and it is continuous with respect to the topology of point-wise convergence. Hence, H is constantly zero. We remark that the property specifying elements in \mathcal{R} cannot be defined as continuity of the functional with respect to a topology.

References

- [1] BORYS, A. Relationships between two definitions of fading memory for discrete-time systems. *International Journal of Electronics and Telecommunications* 61, 4 (2015), 377–380.
- [2] BOYD, S., AND CHUA, L. Fading memory and the problem of approximating nonlinear operators with Volterra series. *IEEE Transactions on Circuits and Systems* 32, 11 (1985), 1150–1161.
- [3] CHRISTMANN, A., AND STEINWART, I. *Support Vector Machines*. Springer, 2008.
- [4] COLEMAN, B. D., AND MIZEL, V. J. On the general theory of fading memory. *Archive for Rational Mechanics and Analysis* 29, 1 (jan 1968), 18–31.
- [5] CONWAY, J. B. *A Course in Functional Analysis*, 2 ed., vol. 96 of *Graduate Texts in Mathematics*. Springer, New York, NY, 2007.
- [6] GONON, L., GRIGORYEVA, L., AND ORTEGA, J.-P. Reservoir kernels and Volterra series. *arXiv:2212.14641v1* (2022).
- [7] GONON, L., GRIGORYEVA, L., AND ORTEGA, J.-P. Approximation error estimates for random neural networks and reservoir systems. *The Annals of Applied Probability* 33, 1 (2023), 28–69.
- [8] GONON, L., GRIGORYEVA, L., AND ORTEGA, J.-P. Infinite-dimensional reservoir computing. *Neural Networks* 179 (2024), 106486.
- [9] GONON, L., AND ORTEGA, J.-P. Fading memory echo state networks are universal. *Neural Networks* 138 (2021), 10–13.
- [10] GRIGORYEVA, L., AND ORTEGA, J.-P. Echo state networks are universal. *Neural Networks* 108 (2018), 495–508.
- [11] GRIGORYEVA, L., AND ORTEGA, J.-P. Differentiable reservoir computing. *Journal of Machine Learning Research* 20, 179 (2019), 1–62.
- [12] GU, A., AND DAO, T. Mamba: Linear-Time Sequence Modeling with Selective State Spaces. *arXiv:2312.00752v2* (2023).
- [13] JAEGER, H. The “echo state” approach to analysing and training recurrent neural networks – with an Erratum note. Tech. Rep. GMD Report 148, German National Research Center for Information Technology, 2010.

- [14] JIANG, H., LI, Q., LI, Z., AND WANG, S. A Brief Survey on the Approximation Theory for Sequence Modelling. *Journal of Machine Learning* 2, 1 (2023), 1–30.
- [15] LI, D., AND QUEFFÉLEC, H. *Introduction to Banach Spaces: Analysis and Probability Volume 1*, vol. 166 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2017. Translated by Danièle Gibbons and Greg Gibbons.
- [16] LI, Z., HAN, J., E, W., AND LI, Q. On the Curse of Memory in Recurrent Neural Networks: Approximation and Optimization Analysis. In *International Conference on Learning Representations* (2021).
- [17] LYUDMILA, G., AND JUAN-PABLO, O. Dimension reduction in recurrent networks by canonicalization. *Journal of Geometric Mechanics* 13, 4 (2021), 647–677.
- [18] MAASS, W., JOSHI, P., AND SONTAG, E. D. Computational aspects of feedback in neural circuits. *PLoS Computational Biology* 3, 1 (2007), e165.
- [19] MAASS, W., NATSCHLÄGER, T., AND MARKRAM, H. Real-time computing without stable states: a new framework for neural computation based on perturbations. *Neural Computation* 14 (2002), 2531–2560.
- [20] MAASS, W., NATSCHLÄGER, T., AND MARKRAM, H. Fading memory and kernel properties of generic cortical microcircuit models. *Journal of Physiology Paris* 98, 4-6 SPEC. ISS. (2004), 315–330.
- [21] MAASS, W., AND SONTAG, E. D. Neural Systems as Nonlinear Filters. *Neural Computation* 12, 8 (aug 2000), 1743–1772.
- [22] MATTHEWS, M. B. *On the Uniform Approximation of Nonlinear Discrete-Time Fading-Memory Systems Using Neural Network Models*. PhD thesis, ETH Zurich, 1992.
- [23] MATTHEWS, M. B. Approximating nonlinear fading-memory operators using neural network models. *Circuits, Systems, and Signal Processing* 12, 2 (jun 1993), 279–307.
- [24] PATRO, B. N., AND AGNEESWARAN, V. S. Mamba-360: Survey of State Space Models as Transformer Alternative for Long Sequence Modelling: Methods, Applications, and Challenges. *arXiv:2404.16112v1* (2024).
- [25] PATRO, B. N., AND AGNEESWARAN, V. S. SiMBA: Simplified Mamba-Based Architecture for Vision and Multivariate Time series. *arXiv:2403.15360v2* (2024).
- [26] PERRYMAN, P. C. *Approximation Theory for Deterministic and Stochastic Nonlinear Systems*. PhD thesis, University of California, Irvine, 1996.
- [27] RUGH, W. J. *Nonlinear System Theory. The Volterra/Wiener Approach*. The Johns Hopkins University Press, 1981.
- [28] SANDBERG, I. W. Approximation theorems for discrete-time systems. *IEEE Transactions on Circuits and Systems* 38, 5 (1991), 564–566.
- [29] SANDBERG, I. W. Structure theorems for nonlinear systems. *Multidimensional Systems and Signal Processing* 2 (1991), 267–286.
- [30] SANDBERG, I. W. Z^+ fading memory and extensions of input-output maps. *International Journal of Circuit Theory and Applications* 29, 11 (2001), 381–388.
- [31] SANDBERG, I. W. Notes of fading-memory conditions. *Circuits, Systems, and Signal Processing* 22, 1 (2003), 43–55.
- [32] SCHÖLKOPF, B., HERBRICH, R., AND SMOLA, A. J. A Generalized Representer Theorem. In *Computational Learning Theory* (2001), D. Helmbold and B. Williamson, Eds., Springer Berlin Heidelberg, pp. 416–426.
- [33] STUBBERUD, A., AND PERRYMAN, P. Current state of system approximation for deterministic and stochastic systems. In *Conference Record of The Thirtieth Asilomar Conference on Signals, Systems and Computers* (1997), vol. 1, IEEE Comput. Soc. Press, pp. 141–145.

- [34] VASWANI, A., SHAZEER, N., PARMAR, N., USZKOREIT, J., JONES, L., GOMEZ, A. N., KAISER, L. U., AND POLOSUKHIN, I. Attention is All you Need. In *Advances in Neural Information Processing Systems* (2017), I. Guyon, U. V. Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, and R. Garnett, Eds., vol. 30, Curran Associates, Inc.
- [35] VOLTERRA, V. *Theory of Functionals and of Integral and Integro-Differential Equations*. Dover, 1959.
- [36] WAHBA, G. *Spline Models for Observational Data*. CBMS-NSF Regional Conference Series in Applied Mathematics. Society for Industrial and Applied Mathematics, 1990.
- [37] WIENER, N. *Nonlinear Problems in Random Theory*. The Technology Press of MIT, 1958.
- [38] ZANG, G., AND IGLESIAS, P. A. Fading memory and stability. *Journal of the Franklin Institute* 340, 6-7 (2004), 489–502.