

ASYMPTOTIC CONVERGENCE FOR A CLASS OF FULLY NONLINEAR INVERSE CURVATURE FLOWS IN A CONE

YA GAO[†], JING MAO^{‡,*}

ABSTRACT. For a given smooth convex cone in the Euclidean $(n + 1)$ -space \mathbb{R}^{n+1} which is centered at the origin, we investigate the evolution of strictly mean convex hypersurfaces, which are star-shaped with respect to the center of the cone and which meet the cone perpendicularly, along an inverse curvature flow with the speed equal to $(f(r)H)^{-1}$, where f is a positive function of the radial distance parameter r and H is the mean curvature of the evolving hypersurfaces. The evolution of those hypersurfaces inside the cone yields a fully nonlinear parabolic Neumann problem. Under suitable constraints on the first and the second derivatives of the radial function f , we can prove the long-time existence of this flow, and moreover the evolving hypersurfaces converge smoothly to a piece of the round sphere.

Keywords: *Inverse curvature flow, Neumann boundary condition, asymptotic convergence.*

MSC 2020: *Primary 53E10, Secondary 35K10.*

1. INTRODUCTION

Let \mathcal{M}_0 be a smooth, closed and convex hypersurface in the Euclidean $(n + 1)$ -space \mathbb{R}^{n+1} which encloses the origin, $n \geq 1$. Li, Sheng and Wang [15] investigated the evolution of \mathcal{M}_0 along the geometric flow

$$(1.1) \quad \begin{cases} \frac{\partial X}{\partial t}(x, t) = -r^\alpha \sigma_k(x, t) \nu, \\ X(x, 0) = X_0(x), \end{cases}$$

where σ_k is the k -th Weingarten curvature (i.e., the k -th elementary symmetric function of principal curvatures) given by

$$\sigma_k(\cdot, t) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} \lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_k},$$

with $\lambda_i = \lambda_i(\cdot, t)$ the principal curvatures of the hypersurface \mathcal{M}_t , parameterized by¹ $X(\cdot, t) : \mathbb{S}^n \rightarrow \mathbb{R}^{n+1}$. Besides, in the system (1.1), $\nu(\cdot, t)$ is the unit outward normal vector of \mathcal{M}_t at $X(\cdot, t)$, $\alpha \in \mathbb{R}^1$ and $r := |X(\cdot, t)|$ denotes the radial distance from the point $X(x, t)$ to the origin. It is not hard to see that: if $\alpha = 0$, $k = 1$, the flow (1.1) degenerates into the well-known mean curvature flow; $\alpha = 0$, $k = n$, the flow (1.1) becomes the Gauss curvature flow. The mean curvature flow and the Gauss curvature flow have been extensively studied in the past six decades, and some interesting, important results have been obtained – see e.g. [1, 3, 5, 11]. BTW, if $\alpha = 0$, then in this setting the evolution equation of (1.1) is simply called σ_k -flow equation and the speed of the σ_k -flow is just the k -th Weingarten curvature (sometimes, k -curvature for abbreviation). If α does not vanish, then the speed of the flow (1.1) depends not only on the curvatures but also on the radial function. As shown by Li, Sheng and Wang in [15], the motivations of their study on the flow (1.1) include not only the close relation with those classical curvature flows just mentioned above but also its background in convex geometry (see

* Corresponding author.

¹ In this paper, \mathbb{S}^n stands for the unit Euclidean n -sphere.

[15, pp. 835-836] for an interesting explanation). For the flow (1.1), Li, Sheng and Wang [15] proved that:

- *If $\alpha \geq k + 1$, the flow exists for all time, preserves the convexity and converges smoothly after normalisation to a sphere centred at the origin. If $\alpha < k + 1$, a counterexample (see [15, Sect. 5]) is given for the above convergence. In the case $k = 1$ and $\alpha \geq 2$, the flow converges to a round point if the initial hypersurface is weakly mean-convex and star-shaped.*

Inspired by the work [15], Mao and his collaborators [2] considered the evolution of closed, star-shaped, strictly mean convex $C^{2,\gamma}$ hypersurfaces in \mathbb{R}^{n+1} along the unit outward normal vector with a speed equal to $(r^\alpha H)^{-1}$, where $H = \sigma_1(\cdot, t)$ is the mean curvature of the hypersurfaces, and proved that if $\alpha \geq 0$, this evolution exists for all time and the evolving hypersurfaces converge smoothly to a round sphere after rescaling. Clearly, when $\alpha = 0$, the expanding flow considered in [2] degenerates into the well-known inverse mean curvature flow (IMCF) and then asymptotic convergence conclusion in [2] certainly covers the convergent result of the classical IMCF firstly obtained by Gerhardt [9] (or Urbas [21]).

The curvature flows mentioned in this paper so far actually correspond to the evolution of closed hypersurfaces along some prescribed curvature flow. What about the situation that the initial hypersurface has boundary? This implies that when considering some prescribed curvature flow, we have not only the initial value condition (IVC) as usual but also the boundary value condition (BVC). This addition of BVC would definitely increase the difficulty of getting a priori estimates for the solutions to the prescribed flow equation.

To the best of our knowledge, the first interesting convergent result concerning the IMCF with boundary is due to T. Marquardt [18, 19]. More precisely, let $M^n \subset \mathbb{S}^n$ ($n \geq 2$) be some piece of \mathbb{S}^n such that $\Sigma^n := \{rx \in \mathbb{R}^{n+1} | r > 0, x \in \partial M^n\}$ be the boundary of a smooth, convex cone that is centered at the origin. Marquardt [18, 19] investigated the evolution of hypersurfaces with boundary (which are star-shaped with respect to the center of the cone Σ^n and which also meet Σ^n perpendicular) along the IMCF, and they can prove that the long-time existence of this flow and moreover the evolving hypersurfaces converge smoothly and exponentially to a piece of the round sphere. Inspired by the previous work [2], together with the success of curvature estimates on the boundary, Mao and Tu [17] generalized Marquardt's above conclusion from the IMCF to the inverse curvature flow with the speed equal to $(r^\alpha H)^{-1}$, $\alpha \geq 0$. Is it possible to consider some inverse curvature flow with boundary in a more general geometric space (not the Euclidean space)? The answer is affirmative and we have obtained several successful cases – see e.g. [6] for the evolution of spacelike (strictly mean convex) graphic hypersurfaces along the IMCF with boundary in a time cone of the Lorentz-Minkowski space \mathbb{R}_1^{n+1} , [7] for the inverse curvature flow with boundary and speed equal to $(r^\alpha H)^{-1}$ in a time cone of \mathbb{R}_1^{n+1} , [8] for the inverse Gauss curvature flow in a time cone of \mathbb{R}_1^{n+1} .

Very recently, inspired by the work [2] of Mao and his collaborators, Gong [12] found that similar convergent result can be obtained if the flow speed $(r^\alpha H)^{-1}$, $\alpha \geq 0$ was replaced by a more general one $(f(r)H)^{-1}$, where f is a positive function of the radial distance parameter $r = |X(\cdot, t)|$ and also satisfies two constraints involving the first and the second derivatives of f (see [12, p. 642] for details).

It is natural to ask whether or not a similar story could be expected in the case that the initial hypersurface also has a boundary when considering the IMCF. The purpose of this paper is trying to give an affirmative answer to this question. In fact, by successfully overcoming difficulties in curvature estimates on the boundary, we can prove the following:

Theorem 1.1. *Let $M^n \subset \mathbb{S}^n$ be some convex piece of the unit sphere $\mathbb{S}^n \subset \mathbb{R}^{n+1}$, and $\Sigma^n = \{rx \in \mathbb{R}^{n+1} | r > 0, x \in \partial M^n\}$ be the boundary of a smooth, convex cone that is centered at the*

origin and has outward unit normal μ . Let $X_0 : M^n \rightarrow \mathbb{R}^{n+1}$ such that $M_0^n := X_0(M^n)$ is a compact, strictly mean convex $C^{2,\gamma}$ -hypersurface ($0 < \gamma < 1$) which is star-shaped with respect to the center of the cone. Assume that

$$M_0^n = \text{graph}_{M^n} u_0$$

is a graph over M^n for a positive map $u_0 : M^n \rightarrow \mathbb{R}$ and

$$\partial M_0^n \subset \Sigma^n, \quad \langle \mu \circ X_0, \nu_0 \circ X_0 \rangle|_{\partial M^n} = 0,$$

where ν_0 is the unit normal vector of M_0^n . Then we have:

(i) There exists a family of strictly mean convex hypersurfaces M_t^n given by the unique embedding

$$X \in C^{2+\gamma, 1+\frac{\gamma}{2}}(M^n \times [0, \infty), \mathbb{R}^{n+1}) \cap C^\infty(M^n \times (0, \infty), \mathbb{R}^{n+1})$$

with $X(\partial M^n, t) \subset \Sigma^n$ for $t \geq 0$, satisfying the following system

$$(1.2) \quad \begin{cases} \frac{\partial}{\partial t} X = \frac{1}{f(|X|)H} \nu & \text{in } M^n \times (0, \infty) \\ \langle \mu \circ X, \nu \circ X \rangle = 0 & \text{on } \partial M^n \times (0, \infty) \\ X(\cdot, 0) = M_0^n & \text{in } M^n \end{cases}$$

where H is the mean curvature of $M_t^n := X(M^n, t) = X_t(M^n)$, ν is the unit outward normal vector of M_t^n , and the function $f(y) \in C^0([0, \infty)) \cap C^2((0, \infty))$ satisfies:

- (1) $f(y) \geq 0$, $f(y) = 0$ if and only if $y = 0$;
- (2) $0 < c_1 \leq yf'(y)f^{-1}(y) \leq c_2$ for some constants c_1 and c_2 , or f is a constant function²;
- (3) If there exist some positive constants c_3, c_4 such that $c_3 \leq f(y) \leq c_4$, then exist some positive constant c_5, c_6 such that $c_5 f(g(t)) \leq f(y \cdot g(t)) \leq c_6 f(g(t))$, where the function $g(t)$ just depends on the time variable t .

Moreover, the Hölder norm on the parabolic space $M^n \times (0, \infty)$ is defined in the usual way (see, e.g., [10, Note 2.5.4]).

(ii) The leaves M_t^n are graphs over M^n , i.e.,

$$M_t^n = \text{graph}_{M^n} u(\cdot, t).$$

(iii) Moreover, the evolving hypersurfaces converge smoothly after rescaling to a piece of a round sphere of radius r_∞ , where r_∞ satisfies

$$\frac{1}{\sup_{M^n} u_0} \left(\frac{\mathcal{H}^n(M_0^n)}{\mathcal{H}^n(M^n)} \right)^{\frac{1}{n}} \leq r_\infty \leq \frac{1}{\inf_{M^n} u_0} \left(\frac{\mathcal{H}^n(M_0^n)}{\mathcal{H}^n(M^n)} \right)^{\frac{1}{n}},$$

where $\mathcal{H}^n(\cdot)$ stands for the n -dimensional Hausdorff measure of a prescribed Riemannian n -manifold.

In Remark 3.1, Remark 3.2 and Remark 4.1 below, we would give the detailed explanations about why the function f needs to satisfy three assumptions (1)-(3) in Theorem 1.1.

Remark 1.1. It is not hard to construct the following possible examples which do satisfy the assumptions (1)-(3) in Theorem 1.1:

- $f(y) = y^\alpha$, $\alpha \geq 0$;
- $f(y) = y + \log(1 + y)$;
- $f(y) = \frac{e^y \cdot y}{1 + e^y}$.

² In $\dot{\varphi}$ estimate, we need the condition that $0 < c_1 \leq yf'(y)f^{-1}(y) \leq c_2$ in order to apply the maximum principle. But when f is a constant function, the third term on the right hand side of the first equation of (3.3) would not appear any more. So here we give the assumption for two different cases of f .

Remark 1.2. It is not hard to find that as long as the function f satisfies three conditions in Theorem 1.1, the evolving hypersurfaces will converge after rescaling to a piece of a round sphere of certain radius r_∞ , which can be controlled by two positive constants that only related to the initial hypersurface.

This paper is organized as follows. In Section 2, we will show that the flow equation (which generally is a system of PDEs) changes into a single scalar second-order parabolic PDE. In Section 3, several estimates, including C^0 , time-derivative and gradient estimates, of solutions to the flow equation will be shown in details. Estimates of higher-order derivatives of solutions to the flow equation, which naturally leads to the long-time existence of the flow, will be investigated in Section 4. We will clearly show the convergence of the rescaled flow in Section 5.

2. THE SCALAR VERSION OF THE FLOW EQUATION

Since the initial $C^{2,\gamma}$ -hypersurface is star-shaped, there exists a function $u_0 \in C^{2,\gamma}(M^n)$ such that $X_0 : M^n \rightarrow \mathbb{R}^{n+1}$ has the form $x \mapsto G_0 := (x, u_0(x))$. The hypersurface M_t^n given by the embedding

$$X(\cdot, t) : M^n \rightarrow \mathbb{R}^{n+1}$$

at time t may be represented as a graph over $M^n \subset \mathbb{S}^n$, and then we can make ansatz

$$X(x, t) = (x, u(x, t))$$

for some function $u : M^n \times [0, T) \rightarrow \mathbb{R}$.

Using techniques as in Ecker [4] (see also [9, 10, 19]), the problem (1.2) is degenerated into solving the following scalar equation with the corresponding initial data and the corresponding NBC

$$(2.1) \quad \begin{cases} \frac{\partial u}{\partial t} = \frac{v}{f(|X|)H} & \text{in } M^n \times (0, \infty) \\ D_\mu u = 0 & \text{on } \partial M^n \times (0, \infty) \\ u(\cdot, 0) = u_0 & \text{in } M^n. \end{cases}$$

By [7, Lemma 2.1], define a new function $\varphi(x, t) = \log u(x, t)$ and then the mean curvature can be rewritten as

$$H = \sum_{i=1}^n h_i^i = \frac{e^{-\varphi}}{v} \left(n - (\sigma^{ij} - \frac{\varphi^i \varphi^j}{v^2}) \varphi_{ij} \right).$$

Hence, the evolution equation in (2.1) can be rewritten as

$$\frac{\partial}{\partial t} \varphi = f^{-1}(e^\varphi) (1 + |D\varphi|^2) \frac{1}{[n - (\sigma^{ij} - \frac{\varphi^i \varphi^j}{v^2}) \varphi_{ij}]} := Q(\varphi, D\varphi, D^2\varphi).$$

In particular,

$$\left(n - (\sigma^{ij} - \frac{\varphi_0^i \varphi_0^j}{v^2}) \varphi_{0,ij} \right)$$

is positive on M^n , since M_0 is strictly mean convex. Thus, the problem (1.2) is again reduced to solve the following scalar equation with the NBC and the initial data

$$(2.2) \quad \begin{cases} \frac{\partial \varphi}{\partial t} = Q(\varphi, D\varphi, D^2\varphi) & \text{in } M^n \times (0, T) \\ D_\mu \varphi = 0 & \text{on } \partial M^n \times (0, T) \\ \varphi(\cdot, 0) = \varphi_0 & \text{in } M^n, \end{cases}$$

where

$$\left(n - (\sigma^{ij} - \frac{\varphi_0^i \varphi_0^j}{v^2}) \varphi_{0,ij} \right)$$

is positive on M^n . Clearly, for the initial graphic hypersurface M_0^n ,

$$\frac{\partial Q}{\partial \varphi_{ij}} \Big|_{\varphi_0} = \frac{1}{e^{2\varphi_0} f(e^{\varphi_0}) H^2} (\sigma^{ij} - \frac{\varphi_0^i \varphi_0^j}{v^2})$$

is positive on M^n , since $f^{-1}(e^{\varphi_0}) > 0$. Based on the above facts, as in [9, 10, 19], we can get the following short-time existence and uniqueness for the parabolic system (1.2).

Lemma 2.1. *Let $X_0(M^n) = M_0^n$ be as in Theorem 1.1. Then there exist some $T > 0$, a unique solution $u \in C^{2+\gamma, 1+\frac{\gamma}{2}}(M^n \times [0, T]) \cap C^\infty(M^n \times (0, T])$, where $\varphi(x, t) = \log u(x, t)$, to the parabolic system (2.2) with the matrix*

$$\left(n - (\sigma^{ij} - \frac{\varphi^i \varphi^j}{v^2}) \varphi_{ij} \right)$$

positive on M^n . Thus there exists a unique map $\psi : M^n \times [0, T] \rightarrow M^n$ such that $\psi(\partial M^n, t) = \partial M^n$ and the map \widehat{X} defined by

$$\widehat{X} : M^n \times [0, T] \rightarrow \mathbb{R}^{n+1} : (x, t) \mapsto X(\psi(x, t), t)$$

has the same regularity as stated in Theorem 1.1 and is the unique solution to the parabolic system (1.2).

Let T^* be the maximal time such that there exists some

$$u \in C^{2+\gamma, 1+\frac{\gamma}{2}}(M^n \times [0, T^*)) \cap C^\infty(M^n \times (0, T^*))$$

which solves (2.2). In the sequel, we shall prove a priori estimates for those admissible solutions on $[0, T]$ where $T < T^*$.

3. C^0 , $\dot{\varphi}$ AND GRADIENT ESTIMATES

Lemma 3.1 (C^0 estimate). *Let φ be a solution of (2.2), we have*

$$c_7 \leq u(x, t) \Theta^{-1}(t, c) \leq c_8, \quad \forall x \in M^n, t \in [0, T]$$

for some positive constants c_7, c_8 , where $\Theta(t, c) := e^{\mathcal{F}(t)+c}$ with

$$\inf_{M^n} \varphi(\cdot, 0) \leq c \leq \sup_{M^n} \varphi(\cdot, 0),$$

where the definition of $\mathcal{F}(t)$ will be given later.

Proof. Let $\varphi(x, t) = \varphi(t)$ (independent of x) be the solution of (2.2) with $\varphi(0) = c$. In this case, the first equation in (2.2) reduces to an ODE

$$\frac{d}{dt} \varphi(t) = \frac{1}{nf(e^{\varphi(t)})} > 0.$$

So we also have

$$\frac{dt}{d\varphi} = nf(e^{\varphi(t)}) > 0.$$

Therefore, using the inverse function theorem, there exists the function $\mathcal{F}(t)$, and $\mathcal{F}'(t) > 0$. One might further request $\mathcal{F}(0) = 0$, and so

$$(3.1) \quad \varphi(t) = \mathcal{F}(t) + c.$$

Using the maximum principle, we can obtain that

$$(3.2) \quad \mathcal{F}(t) + \varphi_1 \leq \varphi(x, t) \leq \mathcal{F}(t) + \varphi_2,$$

where $\varphi_1 := \inf_{M^n} \varphi(\cdot, 0)$ and $\varphi_2 := \sup_{M^n} \varphi(\cdot, 0)$. The estimate is obtained since $\varphi = \log u$. \square

Remark 3.1. Comparing with [7, Lemma 3.1], here since $e^{\varphi(t)} \neq 0$, we have $f(e^{\varphi(t)}) > 0$ (the first assumption in Theorem 1.1). Then using the existence theorem of inverse functions, we know that there exists a function $\mathcal{F}(t)$ such that $\varphi(t) = \mathcal{F}(t) + c$ without the need of explicit expression.

Lemma 3.2 ($\dot{\varphi}$ estimate). *Let φ be a solution of (2.2) and Σ^n be the boundary of a smooth, convex domain defined as in Theorem 1.1, then*

$$\min \left\{ \inf_{M^n} (\dot{\varphi}(\cdot, 0) \cdot f(\Theta(0))), \frac{c_1}{nc_2} \right\} \leq \dot{\varphi}(x, t) f(\Theta(t)) \leq \max \left\{ \sup_{M^n} (\dot{\varphi}(\cdot, 0) \cdot f(\Theta(0))), \frac{c_1}{nc_2} \right\}.$$

Proof. Set

$$\mathcal{M}(x, t) = \dot{\varphi}(x, t) \cdot f(\Theta).$$

Differentiating both sides of the first evolution equation of (2.2), it is easy to get that

$$(3.3) \quad \begin{cases} \frac{\partial \mathcal{M}}{\partial t} = Q^{ij} D_{ij} \mathcal{M} + Q^k D_k \mathcal{M} \\ \quad + f^{-1}(\Theta) \left[\frac{\Theta}{nf(\Theta)} \cdot \frac{\partial f}{\partial \Theta} - \frac{e^\varphi}{f(e^\varphi)} \cdot \frac{\partial f}{\partial e^\varphi} \cdot \mathcal{M} \right] \mathcal{M} & \text{in } M^n \times (0, T) \\ D_\mu \mathcal{M} = 0 & \text{on } \partial M^n \times (0, T) \\ \mathcal{M}(\cdot, 0) = \dot{\varphi}_0 & \text{on } M^n, \end{cases}$$

where $Q^{ij} := \frac{\partial Q}{\partial \varphi_{ij}}$ and $Q^k := \frac{\partial Q}{\partial \varphi_k}$. Then use the second assumption of the function f in the Theorem 1.1, so we have

$$c_1 \leq \frac{e^\varphi}{f(e^\varphi)} \cdot \frac{\partial f}{\partial e^\varphi} \leq c_2.$$

Then the result follows from the maximum principle. \square

Remark 3.2. In Lemma 3.2, since we want to use the maximum principle, we should need the function f to satisfy the assumption (2) in Theorem 1.1. But when f is a constant function, the third term in the right hand side of the equation (3.3) will disappear. Especially, if $f = 1$, the equation (1.2) is the classical inverse mean curvature flow.

Lemma 3.3 (Gradient estimate). *Let φ be a solution of (2.2) and Σ^n be the boundary of a smooth, convex domain described as in Theorem 1.1. Then we have,*

$$(3.4) \quad |D\varphi| \leq \sup_{M^n} |D\varphi(\cdot, 0)|, \quad \forall x \in M^n, t \in [0, T].$$

Proof. Set $\psi = \frac{|D\varphi|^2}{2}$. By differentiating ψ , we have

$$\frac{\partial\psi}{\partial t} = \frac{\partial}{\partial t}\varphi_m\varphi^m = \dot{\varphi}_m\varphi^m = Q_m\varphi^m.$$

Then using the evolution equation of φ in (2.2) yields

$$\frac{\partial\psi}{\partial t} = Q^{ij}\varphi_{ijm}\varphi^m + Q^k\varphi_{km}\varphi^m - e^\varphi f'(e^\varphi)f^{-1}(e^\varphi)Q|D\varphi|^2.$$

Interchanging the covariant derivatives, we have

$$\begin{aligned}\psi_{ij} &= D_j(\varphi_{mi}\varphi^m) \\ &= \varphi_{mij}\varphi^m + \varphi_{mi}\varphi_j^m \\ &= (\varphi_{ijm} + R_{jmi}^l\varphi_l)\varphi^m + \varphi_{mi}\varphi_j^m.\end{aligned}$$

Therefore, we can express $\varphi_{ijm}\varphi^m$ as

$$\varphi_{ijm}\varphi^m = \psi_{ij} - R_{jmi}^l\varphi_l\varphi^m - \varphi_{mi}\varphi_j^m.$$

Then, in view of the fact $R_{ijml} = \sigma_{im}\sigma_{jl} - \sigma_{il}\sigma_{jm}$ on \mathbb{S}^n , we have

$$(3.5) \quad \begin{aligned}\frac{\partial\psi}{\partial t} &= Q^{ij}\psi_{ij} + Q^k\psi_k - Q^{ij}(\sigma_{ij}|D\varphi|^2 - \varphi_i\varphi_j) \\ &\quad - Q^{ij}\varphi_{mi}\varphi_j^m - e^\varphi f'(e^\varphi)f^{-1}(e^\varphi)Q|D\varphi|^2.\end{aligned}$$

Same discussion as in [7, Lemma 3.3], since M^n is convex and using the second assumption in Theorem 1.1, $e^\varphi f'(e^\varphi)f^{-1}(e^\varphi)$ is positive, and so we can get

$$\begin{cases} \frac{\partial\psi}{\partial t} \leq Q^{ij}\psi_{ij} + Q^k\psi_k & \text{in } M^n \times (0, T) \\ D_\mu\psi \leq 0 & \text{on } \partial M^n \times (0, T) \\ \psi(\cdot, 0) = \frac{|D\varphi(\cdot, 0)|^2}{2} & \text{in } M^n. \end{cases}$$

Using the maximum principle, we have

$$|D\varphi| \leq \sup_{M^n} |D\varphi(\cdot, 0)|.$$

Our proof is finished. \square

Remark 3.3. It is worth pointing out that the evolution hypersurfaces M_t^n are always star-shaped under the assumption of Theorem 1.1, since, by Lemma 3.3, we have

$$\left\langle \frac{X}{|X|}, \nu \right\rangle = \frac{1}{v}$$

is bounded from below by a positive constant.

Combing the gradient estimate with $\dot{\varphi}$ estimate, we can obtain

Corollary 3.4. *If φ satisfies (2.2), then we have*

$$(3.6) \quad 0 < c_9 \leq H\Theta \leq c_{10} < +\infty,$$

where c_9 and c_{10} are positive constants independent of φ .

4. HÖLDER ESTIMATES AND CONVERGENCE

In this section, we define the rescaled flow by

$$\tilde{X} = X\Theta^{-1}.$$

Thus,

$$\begin{aligned}\tilde{u} &= u\Theta^{-1}, \\ \tilde{\varphi} &= \varphi - \log \Theta,\end{aligned}$$

and the rescaled mean curvature is given by

$$\tilde{H} = H\Theta.$$

Then, the rescaled scalar curvature equation takes the form

$$\frac{\partial}{\partial t} \tilde{u} = \frac{v}{f(u)\tilde{H}} - \frac{1}{n} \tilde{u} f^{-1}(\Theta).$$

Defining $t = t(s)$ by the relation

$$\frac{dt}{ds} = f(\Theta)$$

such that $t(0) = 0$ and $t(S) = T$. Then \tilde{u} satisfies

$$(4.1) \quad \begin{cases} \frac{\partial}{\partial s} \tilde{u} = \frac{vf(\Theta)}{f(\tilde{u}\Theta)\tilde{H}} - \frac{1}{n} \tilde{u} & \text{in } M^n \times (0, S) \\ D_\mu \tilde{u} = 0 & \text{on } \partial M^n \times (0, S) \\ \tilde{u}(\cdot, 0) = \tilde{u}_0 & \text{in } M^n. \end{cases}$$

Lemma 4.1. *Let X be a solution of (1.2) and $\tilde{X} = X\Theta^{-1}$ be the rescaled solution. Then*

$$\begin{aligned}D\tilde{u} &= Du\Theta^{-1}, \quad D\tilde{\varphi} = D\varphi, \quad \frac{\partial \tilde{u}}{\partial s} = \frac{\partial u}{\partial t} \frac{f(\Theta)}{\Theta} - \frac{1}{n} u\Theta^{-1}, \\ \tilde{g}_{ij} &= \Theta^{-2} g_{ij}, \quad \tilde{g}^{ij} = \Theta^2 g^{ij}, \quad \tilde{h}_{ij} = h_{ij}\Theta^{-1}.\end{aligned}$$

Proof. These relations can be computed directly. \square

Lemma 4.2. *Let u be a solution to the parabolic system (2.2), where $\varphi(x, t) = \log u(x, t)$, and Σ^n be the boundary of a smooth, convex domain described as in Theorem 1.1. Then there exist some $\beta > 0$ and some $C > 0$ such that the rescaled function $\tilde{u}(x, s) := u(x, t(s))\Theta^{-1}(t(s))$ satisfies*

$$(4.2) \quad [D\tilde{u}]_\beta + \left[\frac{\partial \tilde{u}}{\partial s} \right]_\beta + [\tilde{H}]_\beta \leq C(\|u_0\|_{C^{2+\gamma, 1+\frac{\gamma}{2}}(M^n)}, n, \beta, M^n),$$

where $[f]_\beta := [f]_{x, \beta} + [f]_{s, \frac{\beta}{2}}$ is the sum of the Hölder coefficients of f in $M^n \times [0, S]$ with respect to x and s .

Proof. We divide our proof in three steps³.

Step 1: We need to prove that

$$[D\tilde{u}]_{x, \beta} + [D\tilde{u}]_{s, \frac{\beta}{2}} \leq C(\|u_0\|_{C^{2+\gamma, 1+\frac{\gamma}{2}}(M^n)}, n, \beta, M^n).$$

³ In the proof of Lemma 4.2, the constant C may differ from each other. However, we abuse the symbol C for the purpose of convenience.

According to Lemmas 3.1, 3.2 and 3.3, it follows that

$$|D\tilde{u}| + \left| \frac{\partial \tilde{u}}{\partial s} \right| \leq C(\|u_0\|_{C^{2+\gamma, 1+\frac{\gamma}{2}}(M^n)}, M^n).$$

Fix s and the equation (2.2) can be rewritten as an elliptic Neumann problem

$$(4.3) \quad \operatorname{div}_\sigma \left(\frac{D\varphi}{\sqrt{1+|D\varphi|^2}} \right) = \frac{n}{\sqrt{1+|D\varphi|^2}} - f^{-1}(e^\varphi) \cdot \frac{\sqrt{1+|D\varphi|^2}}{\dot{\varphi}}.$$

Other discussions are the same as in [7, Lemma 4.3].

Step 2: The next thing to do is to show that

$$\left[\frac{\partial \tilde{u}}{\partial s} \right]_{x, \beta} + \left[\frac{\partial \tilde{u}}{\partial s} \right]_{s, \frac{\beta}{2}} \leq C(\|u_0\|_{C^{2+\gamma, 1+\frac{\gamma}{2}}(M^n)}, n, \beta, M^n).$$

As $\frac{\partial}{\partial s} \tilde{u} = \tilde{u} \left(\frac{vf(\Theta)}{f(\tilde{u}\Theta)\tilde{u}H} - \frac{1}{n} \right)$, it is enough to bound $\left[\frac{vf(\Theta)}{\tilde{u}Hf(\tilde{u}\Theta)} \right]_\beta$. Set $\Omega(s) := \frac{vf(\Theta)}{\tilde{u}Hf(\tilde{u}\Theta)}$. Let $\tilde{\nabla}$ be the Levi-Civita connection of $\tilde{M}_s := \tilde{X}(M^n, s)$ w.r.t. the metric \tilde{g} . To prove that Ω belongs to the De Giorgi class of functions, we will need the following evolution equation.

Lemma 4.3. *Set $\Phi = \frac{1}{f(e^\varphi)H} = \frac{1}{f(u)H} = \frac{1}{f(\tilde{u}\Theta)H}$, $w = \langle X, \nu \rangle = \frac{u}{v}$ and $\Psi = \frac{\Phi}{w} = \frac{v}{\tilde{u}Hf(\tilde{u}\Theta)}$. Under the assumptions of Theorem 1.1, we have*

$$(4.4) \quad \begin{aligned} \frac{\partial \Psi}{\partial t} &= \operatorname{div}_g(f^{-1}H^{-2}\nabla\Psi) - 2H^{-2}f^{-1}(e^\varphi)\Psi^{-1}|\nabla\Psi|^2 - f'\frac{u}{f}\Psi^2 \\ &\quad - f'f^{-1}\Psi^2\nabla^i u \langle X, X_i \rangle - f'f^{-2}H^{-2}\nabla_i u \nabla^i \Psi. \end{aligned}$$

Proof. Here, for the continuity of the proof of Lemma 4.2, we put the proof of Lemma 4.3 in Appendix at the end of this paper. In fact, the main method used for the proof of Lemma 4.3 is similar to that in [6, 7], but we still give all the details in Appendix to show the differences brought by the usage of the more general function f . \square

Combining (4.4) and $\Omega(s) = \Psi f(\Theta)$, we get

$$(4.5) \quad \begin{aligned} \frac{\partial \Omega}{\partial s} &= \operatorname{div}_{\tilde{g}} \left(f(\Theta)f^{-1}(\tilde{u}\Theta)\tilde{H}^{-2}\tilde{\nabla}\Omega \right) - 2\tilde{H}^{-2}f^{-1}(\tilde{u}\Theta)f(\Theta)\Omega^{-1}|\tilde{\nabla}\Omega|_{\tilde{g}}^2 + \frac{1}{n} \frac{f'(\Theta)\Theta}{f(\Theta)} \cdot \Omega \\ &\quad - \frac{f'(u)u}{f(u)} \cdot \Omega^2 - \frac{f'(u)u}{f(u)} \Omega^2 P - \frac{f'(u)u}{f^2(u)} \frac{f(\Theta)}{\tilde{u}} \tilde{H}^{-2}\tilde{\nabla}_i \tilde{u} \tilde{\nabla}^i \Omega, \end{aligned}$$

where $P := u^{-1}\nabla^i u \langle X, X_i \rangle$. Applying Lemma 3.3, we have

$$|P| \leq |\nabla u|^2 \leq C.$$

The weak formulation of (4.5) is

$$(4.6) \quad \begin{aligned} \int_{s_0}^{s_1} \int_{\tilde{M}_s} \frac{\partial \Omega}{\partial s} \eta d\mu_s ds &= \int_{s_0}^{s_1} \int_{\tilde{M}_s} \operatorname{div}_{\tilde{g}}(f(\Theta)f^{-1}(\tilde{u}\Theta)\tilde{H}^{-2}\tilde{\nabla}\Omega)\eta - 2\tilde{H}^{-2}f^{-1}(\tilde{u}\Theta)f(\Theta)\Omega^{-1}|\tilde{\nabla}\Omega|_{\tilde{g}}^2 \eta d\mu_s ds \\ &\quad + \int_{s_0}^{s_1} \int_{\tilde{M}_s} \left(\frac{1}{n} \frac{f'(\Theta)\Theta}{f(\Theta)} \cdot \Omega - \frac{f'(u)u}{f(u)} \cdot \Omega^2 - \frac{f'(u)u}{f(u)} \Omega^2 P - \frac{f'(u)u}{f^2(u)} \frac{f(\Theta)}{\tilde{u}} \tilde{H}^{-2}\tilde{\nabla}_i \tilde{u} \tilde{\nabla}^i \Omega \right) \eta d\mu_s ds. \end{aligned}$$

Since $\nabla_\mu \tilde{\varphi} = 0$, the boundary integrals all vanish, the interior and boundary estimates are basically the same. We define the test function $\eta := \xi^2 \Omega$, where ξ is a smooth function with

values in $[0, 1]$ and is supported in a small parabolic neighborhood. Then

$$(4.7) \quad \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \frac{\partial \Omega}{\partial s} \xi^2 \Omega d\mu_s ds = \frac{1}{2} \|\Omega \xi\|_{2, \widetilde{M}_s}^2 \Big|_{s_0}^{s_1} - \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \xi \dot{\xi} \Omega^2 d\mu_s ds.$$

where $\dot{\xi} := \frac{\partial \xi}{\partial s}$. Using the divergence theorem and Young's inequality and the second assumption of the function f in the Theorem 1.1, we can obtain

$$(4.8) \quad \begin{aligned} & \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \operatorname{div}_{\widetilde{g}}(f(\Theta) f^{-1}(\widetilde{u}\Theta) \widetilde{H}^{-2} \widetilde{\nabla} \Omega) \xi^2 \Omega d\mu_s ds \\ &= - \int_{s_0}^{s_1} \int_{\widetilde{M}_s} f(\Theta) f^{-1}(\widetilde{u}\Theta) \widetilde{H}^{-2} \xi^2 \widetilde{\nabla}_i \Omega \widetilde{\nabla}^i \Omega d\mu_s ds \\ & \quad - 2 \int_{s_0}^{s_1} \int_{\widetilde{M}_s} f(\Theta) f^{-1}(\widetilde{u}\Theta) \widetilde{H}^{-2} \xi \Omega \widetilde{\nabla}_i \Omega \widetilde{\nabla}^i \xi d\mu_s ds \\ & \leq \int_{s_0}^{s_1} \int_{\widetilde{M}_s} c_{11} \widetilde{H}^{-2} |\widetilde{\nabla} \xi|^2 \Omega^2 d\mu_s ds. \end{aligned}$$

and

$$(4.9) \quad \begin{aligned} & \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \left(\frac{1}{n} \frac{f'(\Theta)\Theta}{f(\Theta)} \cdot \Omega - \frac{f'(u)u}{f(u)} \cdot \Omega^2 - \frac{f'(u)u}{f(u)} \Omega^2 P \right. \\ & \quad \left. - \frac{f'(u)u}{f^2(u)} \frac{f(\Theta)}{\widetilde{u}} \widetilde{H}^{-2} \widetilde{\nabla}_i \widetilde{u} \widetilde{\nabla}^i \Omega \right) \xi^2 \Omega d\mu_s ds \\ & \leq C(c_2, n, P) \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \xi^2 (\Omega^2 + |\Omega|^3) d\mu_s ds \\ & \quad + \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \frac{f'(u)u}{f(u)} \cdot \frac{f(\Theta)}{f(\widetilde{u}\Theta)} \cdot \frac{1}{\widetilde{u}} \widetilde{H}^{-2} |\widetilde{\nabla} \widetilde{u}| |\widetilde{\nabla} \Omega| \xi^2 \Omega d\mu_s ds \\ & \leq C(c_2, n, P) \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \xi^2 (\Omega^2 + |\Omega|^3) d\mu_s ds + \frac{c_2}{2} \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \widetilde{H}^{-2} \frac{f(\Theta)}{f(\widetilde{u}\Theta)} |\widetilde{\nabla} \Omega|^2 \xi^2 \Omega d\mu_s ds \\ & \quad + \frac{c_2}{2} \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \widetilde{H}^{-2} \frac{f(\Theta)}{f(\widetilde{u}\Theta) \cdot \widetilde{u}^2} |\widetilde{\nabla} \widetilde{u}|^2 \xi^2 \Omega^2 d\mu_s ds. \end{aligned}$$

Combing (4.7), (4.8), (4.9) and the third assumption (3) of the function f in Theorem 1.1, we have

$$(4.10) \quad \begin{aligned} & \frac{1}{2} \|\Omega \xi\|_{2, \widetilde{M}_s}^2 \Big|_{s_0}^{s_1} + C(2 - \frac{c_2}{2}) \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \widetilde{H}^{-2} |\widetilde{\nabla} \Omega|^2 \xi^2 \Omega d\mu_s ds \\ & \leq \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \xi |\dot{\xi}| \Omega^2 d\mu_s ds + \int_{s_0}^{s_1} \int_{\widetilde{M}_s} c_{11} \widetilde{H}^{-2} |\widetilde{\nabla} \xi|^2 \Omega^2 d\mu_s ds \\ & \quad + C(c_2, n, P) \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \xi^2 (\Omega^2 + |\Omega|^3) d\mu_s ds \\ & \quad + \frac{c_2 c_5}{2} \int_{s_0}^{s_1} \int_{\widetilde{M}_s} \widetilde{H}^{-2} \widetilde{u}^{-2} |\widetilde{\nabla} \widetilde{u}|^2 \xi^2 \Omega^2 d\mu_s ds, \end{aligned}$$

which implies that

$$\begin{aligned}
(4.11) \quad & \frac{1}{2} \|\Omega \xi\|_{2, \tilde{M}_s}^2 \Big|_{s_0}^{s_1} + \frac{C(c_2)}{\max(\tilde{H}^2)} \int_{s_0}^{s_1} \int_{\tilde{M}_s} |\tilde{\nabla} \Omega|^2 \xi^2 d\mu_s ds \\
& \leq \left(1 + \frac{c_{11}}{\min(\tilde{H}^2)} \right) \int_{s_0}^{s_1} \int_{\tilde{M}_s} \Omega^2 (\xi |\dot{\xi}| + |\tilde{\nabla} \xi|^2) d\mu_s ds \\
& \quad + \left(C(c_2, n, P) + \frac{c_2 c_5 \max |\tilde{\nabla} \tilde{u}|^2}{2 \min(\tilde{H}^2 \tilde{u}^2)} \right) \int_{s_0}^{s_1} \int_{\tilde{M}_s} \xi^2 \Omega^2 + \xi^2 |\Omega|^3 d\mu_s ds.
\end{aligned}$$

This means that Ω belong to the De Giorgi class of functions in $M^n \times [0, S)$. Similar to the arguments in [14, Chap. 5, §1 and §7], there exist constants β and C such that

$$[\Omega]_\beta \leq C \|\Omega\|_{L^\infty(M^n \times [0, S))} \leq C (\|u_0\|_{C^{2+\gamma, 1+\frac{\gamma}{2}}(M^n)}, n, \beta, M^n).$$

Step 3: Finally, we have to show that

$$[\tilde{H}]_{x, \beta} + [\tilde{H}]_{s, \frac{\beta}{2}} \leq C (\|u_0\|_{C^{2+\gamma, 1+\frac{\gamma}{2}}(M^n)}, n, \beta, M^n).$$

This follows from the fact that

$$\tilde{H} = \frac{\sqrt{1 + |D\tilde{\varphi}|^2} \cdot f(\Theta)}{\tilde{u} f(\tilde{u}\Theta)\Omega}$$

together with the estimates for \tilde{u} , Ω , $D\tilde{\varphi}$ and the conditions of f . \square

Remark 4.1. In this paper, the function f is much more general than that in [7, 17] and we don't have the explicit expression. So we only can write $f(\tilde{u}\Theta)$ (related with \tilde{u}) in the rescaled scalar curvature equation rather than $f(\tilde{u})$. Furthermore, by Lemma 3.1 we know $c_7 \leq \tilde{u} \leq c_8$ and $\Theta(t)$ is a function only related to t . Using the third assumption (3) in Theorem 1.1, we can also obtain the conclusion of Lemma 4.2.

Then we can obtain the following higher-order estimates:

Lemma 4.4. *Let u be a solution to the parabolic system (2.2), where $\varphi(x, t) = \log u(x, t)$, and Σ^n be the boundary of a smooth, convex domain described as in Theorem 1.1. Then for any $s_0 \in (0, S)$ there exist some $\beta > 0$ and some $C > 0$ such that*

$$(4.12) \quad \|\tilde{u}\|_{C^{2+\beta, 1+\frac{\beta}{2}}(M^n \times [0, S])} \leq C (\|u_0\|_{C^{2+\gamma, 1+\frac{\gamma}{2}}(M^n)}, n, \beta, M^n)$$

and for all $k \in \mathbb{N}$,

$$(4.13) \quad \|\tilde{u}\|_{C^{2k+\beta, k+\frac{\beta}{2}}(M^n \times [s_0, S])} \leq C (\|u_0(\cdot, s_0)\|_{C^{2k+\beta, k+\frac{\beta}{2}}(M^n)}, n, \beta, M^n).$$

Proof. By [7, Lemma 2.1], we have

$$uvH = n - \left(\sigma^{ij} - \frac{\varphi^i \varphi^j}{v^2} \right) \varphi_{ij} = n - u^2 \Delta_g \varphi.$$

Since

$$u^2 \Delta_g \varphi = \tilde{u}^2 \Delta_{\tilde{g}} \varphi = -|\tilde{\nabla} \tilde{u}|^2 + \tilde{u} \Delta_{\tilde{g}} \tilde{u},$$

then

$$\begin{aligned}\frac{\partial \tilde{u}}{\partial s} &= \frac{\partial u}{\partial t} \Theta^{-1} f(\Theta) - \frac{1}{n} \tilde{u} \\ &= -\frac{uvH}{uH^2 f(e^\varphi)} \Theta^{-1} f(\Theta) + \frac{2v}{H f(e^\varphi)} \Theta^{-1} f(\Theta) - \frac{1}{n} \tilde{u} \\ &= \frac{\Delta_{\tilde{g}} \tilde{u}}{\tilde{H}^2} \cdot \frac{f(\Theta)}{f(e^\varphi)} + \frac{2v}{\tilde{H}} \cdot \frac{f(\Theta)}{f(e^\varphi)} - \frac{1}{n} \tilde{u} - \frac{n + |\tilde{\nabla} \tilde{u}|^2}{\tilde{u} \tilde{H}^2} \cdot \frac{f(\Theta)}{f(e^\varphi)},\end{aligned}$$

which is a uniformly parabolic equation with Hölder continuous coefficients. Therefore, the linear theory (see [16, Chap. 4]) yields the inequality (4.12).

Set $\tilde{\varphi} = \log \tilde{u}$, and then the rescaled version of the evolution equation in (4.1) takes the form

$$\frac{\partial \tilde{\varphi}}{\partial s} = \frac{f(\Theta)}{f(e^\varphi)} \cdot \frac{v^2}{\left[n - \left(\sigma^{ij} - \frac{\tilde{\varphi}^i \tilde{\varphi}^j}{v^2} \right) \tilde{\varphi}_{ij} \right]} - \frac{1}{n},$$

where $v = \sqrt{1 + |D\tilde{\varphi}|^2}$. According to the $C^{2+\beta, 1+\frac{\beta}{2}}$ -estimate of \tilde{u} (see Lemma 4.2), we can treat the equations for $\frac{\partial \tilde{\varphi}}{\partial s}$ and $D_i \tilde{\varphi}$ as second-order linear uniformly parabolic PDEs on $M^n \times [s_0, S]$. At the initial time s_0 , all compatibility conditions are satisfied and the initial function $u(\cdot, t_0)$ is smooth. We can obtain a $C^{3+\beta, \frac{3+\beta}{2}}$ -estimate for $D_i \tilde{\varphi}$ and a $C^{2+\beta, \frac{2+\beta}{2}}$ -estimate for $\frac{\partial \tilde{\varphi}}{\partial s}$ (the estimates are independent of T) by Theorem 4.3 and Exercise 4.5 in [16, Chap. 4]. Higher regularity can be proven by induction over k . \square

Theorem 4.5. *Under the hypothesis of Theorem 1.1, we conclude*

$$T^* = +\infty.$$

Proof. The proof of this result is quite similar to the corresponding argument in [19, Lemma 8] and so is omitted. \square

5. CONVERGENCE OF THE RESCALED FLOW

We know that after the long-time existence of the flow has been obtained (see Theorem 4.5), the rescaled version of the system (2.2) satisfies

$$(5.1) \quad \begin{cases} \frac{\partial}{\partial s} \tilde{\varphi} = \tilde{Q}(\tilde{\varphi}, D\tilde{\varphi}, D^2\tilde{\varphi}) & \text{in } M^n \times (0, \infty) \\ D_\mu \tilde{\varphi} = 0 & \text{on } \partial M^n \times (0, \infty) \\ \tilde{\varphi}(\cdot, 0) = \tilde{\varphi}_0 & \text{in } M^n, \end{cases}$$

where

$$\tilde{Q}(\tilde{\varphi}, D\tilde{\varphi}, D^2\tilde{\varphi}) := \frac{f(\Theta)}{f(e^\varphi \cdot \Theta)} \cdot \frac{v^2}{\left[n - \left(\sigma^{ij} - \frac{\tilde{\varphi}^i \tilde{\varphi}^j}{v^2} \right) \tilde{\varphi}_{ij} \right]} - \frac{1}{n}$$

and $\tilde{\varphi} = \log \tilde{u}$. Similar to what has been done in the C^1 estimate (see Lemma 3.3), we can deduce a decay estimate of $\tilde{u}(\cdot, s)$ as follows.

Lemma 5.1. *Let u be a solution of (2.1), then we have*

$$(5.2) \quad |D\tilde{u}(x, s)| \leq c_{12} \sup_{M^n} |D\tilde{u}(\cdot, 0)|,$$

where c_{12} is a positive constant depending only on c_7, c_8 .

Proof. Set $\tilde{\psi} = \frac{|D\tilde{\varphi}|^2}{2}$. Similar to that in Lemma 3.3, we also can obtain

$$\begin{cases} \frac{\partial \tilde{\psi}}{\partial s} \leq \tilde{Q}^{ij} \tilde{\psi}_{ij} + \tilde{Q}^k \tilde{\psi}_k & \text{in } M^n \times (0, \infty) \\ D_\mu \tilde{\psi} \leq 0 & \text{on } \partial M^n \times (0, \infty) \\ \psi(\cdot, 0) = \frac{|D\tilde{\varphi}(\cdot, 0)|^2}{2} & \text{in } M^n. \end{cases}$$

Using the maximum principle and Hopf's lemma, we can get the gradient estimate of $\tilde{\varphi}$, and then the conclusion (5.2) follows. \square

Lemma 5.2. *Let u be a solution of the flow (2.1). Then,*

$$\tilde{u}(\cdot, s)$$

converges to a real number as $s \rightarrow +\infty$.

Proof. Set $P(t) := \mathcal{H}^n(M_t^n)$, which, as before, represents the n -dimensional Hausdorff measure of M_t^n and is actually the area of M_t^n . According to the first variation of a submanifold, see e.g. [20], and the fact $\operatorname{div}_{M_t^n} \nu = H$, we have

$$\begin{aligned} (5.3) \quad P'(t) &= \int_{M_t^n} \operatorname{div}_{M_t^n} \left(\frac{\nu}{f(|X|)H} \right) d\mathcal{H}^n \\ &= \int_{M_t^n} \sum_{i=1}^n \left\langle \nabla_{e_i} \left(\frac{\nu}{f(|X|)H} \right), e_i \right\rangle d\mathcal{H}^n \\ &= \int_{M_t^n} f^{-1}(u) d\mathcal{H}^n, \end{aligned}$$

where $\{e_i\}_{1 \leq i \leq n}$ is some orthonormal basis of the tangent bundle TM_t^n . By the condition of the function f and (3.2), we have

$$f^{-1}(e^{\mathcal{F}(t)+\varphi_2}) \leq f^{-1}(u) \leq f^{-1}(e^{\mathcal{F}(t)+\varphi_1})$$

Hence, we have

$$f^{-1}(e^{\mathcal{F}(t)+\varphi_2}) P(t) \leq P'(t) \leq f^{-1}(e^{\mathcal{F}(t)+\varphi_1}) P(t).$$

Combining this fact with (5.3), the construction process of function \mathcal{F} , and choosing the constant c in Lemma 3.1 as φ_1 and φ_2 respectively, we know

$$\mathcal{H}^n(M_0^n) \cdot \frac{e^{n(\mathcal{F}(t)+\varphi_2)}}{e^{n\varphi_2}} \leq P(t) \leq \mathcal{H}^n(M_0^n) \cdot \frac{e^{n(\mathcal{F}(t)+\varphi_1)}}{e^{n\varphi_1}}.$$

Therefore, the rescaled hypersurface $\tilde{M}_s = M_t^n \Theta^{-1}$ satisfies the following inequality

$$\frac{\mathcal{H}^n(M_0^n)}{e^{n\varphi_2}} \leq \mathcal{H}^n(\tilde{M}_s) \leq \frac{\mathcal{H}^n(M_0^n)}{e^{n\varphi_1}},$$

which implies that the area of \tilde{M}_s is bounded and the bounds are independent of s . Here $\varphi_1 = \inf_{M^n} \varphi(\cdot, 0)$ and $\varphi_2 = \sup_{M^n} \varphi(\cdot, 0)$. Together with (4.12), Lemma 5.1 and the Arzelà-Ascoli theorem, we conclude that $\tilde{u}(\cdot, s)$ must converge in $C^\infty(M^n)$ to a constant function r_∞ with

$$\frac{1}{e^{\varphi_2}} \left(\frac{\mathcal{H}^n(M_0^n)}{\mathcal{H}^n(M^n)} \right)^{\frac{1}{n}} \leq r_\infty \leq \frac{1}{e^{\varphi_1}} \left(\frac{\mathcal{H}^n(M_0^n)}{\mathcal{H}^n(M^n)} \right)^{\frac{1}{n}},$$

i.e.,

$$(5.4) \quad \frac{1}{\sup_{M^n} u_0} \left(\frac{\mathcal{H}^n(M_0^n)}{\mathcal{H}^n(M^n)} \right)^{\frac{1}{n}} \leq r_\infty \leq \frac{1}{\inf_{M^n} u_0} \left(\frac{\mathcal{H}^n(M_0^n)}{\mathcal{H}^n(M^n)} \right)^{\frac{1}{n}}.$$

This completes the proof. \square

So, we have

Theorem 5.3. *The rescaled flow*

$$\frac{d\tilde{X}}{ds} = \frac{f(\Theta)}{f(|X|)\tilde{H}}\nu - \frac{1}{n}\tilde{X}$$

exists for all time and the leaves converge in C^∞ to a piece of the round sphere of radius r_∞ , where r_∞ satisfies (5.4).

APPENDIX

Proof of Lemma 4.3. Using the Gauss formula, we have

$$\begin{aligned} \partial_t h_{ij} &= \partial_t \langle \partial_i \partial_j X, -\nu \rangle \\ &= -\nabla_{ij}^2 \Phi + \Phi h_{ik} h_j^k. \end{aligned}$$

Direct calculation results in

$$\begin{aligned} \nabla_{ij}^2 \Phi &= \Phi \left(-\frac{1}{H} H_{ij} + \frac{2H_i H_j}{H^2} \right) + 2f' f^{-1} \Phi u_i (\log H)_j \\ &\quad - f' f^{-1} \Phi u_{ij} - (f'' f^{-1} - 2f^{-2} (f')^2) \Phi u_i u_j. \end{aligned}$$

Since we have

$$\Delta h_{ij} = H_{ij} + H h_{ik} h_j^k - h_{ij} |A|^2,$$

so

$$\begin{aligned} \nabla_{ij}^2 \Phi &= -\Phi H^{-1} \Delta h_{ij} + \Phi h_{ik} h_j^k - \Phi H^{-1} |A|^2 h_{ij} + 2\Phi \frac{H_i H_j}{H^2} \\ &\quad + 2f' f^{-1} \Phi u_i (\log H)_j - f' f^{-1} \Phi u_{ij} - (f'' f^{-1} - 2f^{-2} (f')^2) \Phi u_i u_j. \end{aligned}$$

Thus

$$\begin{aligned} \partial_t h_{ij} - \Phi H^{-1} \Delta h_{ij} &= \Phi H^{-1} |A|^2 h_{ij} - \frac{2\Phi}{H^2} H_i H_j - 2f' f^{-1} \Phi u_i (\log H)_j \\ &\quad + f' f^{-1} \Phi u_{ij} + (f'' f^{-1} - 2f^{-2} (f')^2) \Phi u_i u_j. \end{aligned}$$

Then

$$\begin{aligned} \partial_t H &= h_{ij} \partial_t g^{ij} + g^{ij} \partial_t h_{ij} \\ &= \Phi H^{-1} \Delta H - \frac{2\Phi}{H^2} |\nabla H|^2 - \Phi |A|^2 - 2f' f^{-1} \Phi u_i (\log H)^i \\ &\quad + f' f^{-1} \Phi \Delta u + (f'' f^{-1} - 2f^{-2} (f')^2) \Phi |\nabla u|^2 \\ &= f^{-1} H^{-2} \Delta H - 2f^{-1} H^{-3} |\nabla H|^2 - f^{-1} H^{-1} |A|^2 - 2f' f^{-2} H^{-2} u_i H^i \\ &\quad + f' f^{-2} H^{-1} \nabla u + (f'' f^{-1} - 2f^{-2} (f')^2) f^{-1} H^{-1} |\nabla u|^2. \end{aligned}$$

Clearly

$$\partial_t w = \Phi + f' f^{-1} \Phi \nabla^i u \langle X, X_i \rangle + \Phi H^{-1} \nabla^i H \langle X, X_i \rangle,$$

using the Weingarten equation, we have

$$w_i = h_i^k \langle X, X_k \rangle,$$

$$w_{ij} = h_{i,j}^k \langle X, X_k \rangle + h_{ij} - h_i^k h_{kj} \langle X, \nu \rangle = h_{ij,k} \langle X, X^k \rangle + h_{ij} - h_i^k h_{kj} \langle X, \nu \rangle.$$

Thus

$$\Delta w = H + \nabla^i H \langle X, X_i \rangle - |A|^2 \langle X, \nu \rangle,$$

and

$$\partial_t w = f^{-1} H^{-2} \Delta w + f^{-1} H^{-2} w |A|^2 + f' f^{-2} H^{-1} \nabla^i u \langle X, X_i \rangle.$$

Hence

$$\begin{aligned} \frac{\partial \Psi}{\partial t} &= -\frac{f'}{f^2 H w} \partial_t u - \frac{1}{f H^2 w} \partial_t H - \frac{1}{f H w^2} \partial_t w \\ &= -f' f^{-3} H^{-2} w^{-2} u - (f'' f^{-1} - 2f^{-2} (f')^2) f^{-2} H^{-3} w^{-1} |\nabla u|^2 + 2f^{-2} H^{-5} w^{-1} |\nabla H|^2 \\ &\quad + 2f' f^{-3} H^{-4} w^{-1} u_i H^i - f' f^{-3} H^{-3} w^{-1} \Delta u - f^{-2} H^{-4} w^{-1} \Delta H - f^{-2} H^{-3} w^{-2} \Delta w \\ &\quad - f' f^{-3} H^{-2} w^{-2} \nabla^i u \langle X, X_i \rangle. \end{aligned}$$

In order to prove (4.4), we calculate

$$\nabla_i \Psi = -f' f^{-2} H^{-1} w^{-1} \nabla_i u - f^{-1} H^{-2} w^{-1} \nabla_i H - f^{-1} H^{-1} w^{-2} \nabla_i w,$$

and

$$\begin{aligned} \nabla_{ij}^2 \Psi &= -(f'' f^{-1} - 2f^{-2} (f')^2) f^{-1} H^{-1} w^{-1} \nabla_i u \nabla_j u + 2f^{-1} H^{-3} w^{-1} \nabla_i H \nabla_j H + 2f^{-1} H^{-1} w^{-3} \nabla_i w \nabla_j w \\ &\quad + 2f' f^{-2} H^{-2} w^{-1} \nabla_i u \nabla_j H + 2f' f^{-2} H^{-1} w^{-2} \nabla_i u \nabla_j w + 2f^{-1} H^{-2} w^{-2} \nabla_i H \nabla_j w \\ &\quad - f' f^{-2} H^{-1} w^{-1} \nabla_{ij}^2 u - f^{-1} H^{-2} w^{-1} \nabla_{ij}^2 H - f^{-1} H^{-1} w^{-2} \nabla_{ij}^2 w. \end{aligned}$$

Thus

$$\begin{aligned} f^{-1} H^{-2} \Delta \Psi &= -(f'' f^{-1} - 2f^{-2} (f')^2) f^{-2} H^{-3} w^{-1} |\nabla u|^2 + 2f^{-2} H^{-5} w^{-1} |\nabla H|^2 + 2f^{-2} H^{-3} w^{-3} |\nabla w|^2 \\ &\quad + 2f' f^{-3} H^{-4} w^{-1} \nabla_i u \nabla^i H + 2f' f^{-3} H^{-3} w^{-2} \nabla_i u \nabla^i w + 2f^{-2} H^{-4} w^{-2} \nabla_i H \nabla^i w \\ &\quad - f' f^{-3} H^{-3} w^{-1} \Delta u - f^{-2} H^{-4} w^{-1} \Delta H - f^{-2} H^{-3} w^{-2} \Delta w. \end{aligned}$$

So we have

$$\begin{aligned} \operatorname{div}(f^{-1} H^{-2} \nabla \Psi) &= -f' f^{-2} H^{-2} \nabla_i \Psi \nabla^i u - 2f^{-1} H^{-3} \nabla_i \Psi \nabla^i H + f^{-1} H^{-2} \Delta \Psi \\ &= -(f'' f^{-1} - 3f^{-2} (f')^2) f^{-2} H^{-3} w^{-1} |\nabla u|^2 + 4f^{-2} H^{-5} w^{-1} |\nabla H|^2 + 2f^{-2} H^{-3} w^{-3} |\nabla w|^2 \\ &\quad + 5f' f^{-3} H^{-4} w^{-1} \nabla_i u \nabla^i H + 3f' f^{-3} H^{-3} w^{-2} \nabla_i u \nabla^i w + 4f^{-2} H^{-4} w^{-2} \nabla_i H \nabla^i w \\ &\quad - f' f^{-3} H^{-3} w^{-1} \Delta u - f^{-2} H^{-4} w^{-1} \Delta H - f^{-2} H^{-3} w^{-2} \Delta w, \end{aligned}$$

and

$$\begin{aligned} 2H^{-1} w |\nabla \Psi|^2 &= 2(f')^2 f^{-4} H^{-3} w^{-1} |\nabla u|^2 + 2f^{-2} H^{-5} w^{-1} |\nabla H|^2 + 2f^{-2} H^{-3} w^{-3} |\nabla w|^2 \\ &\quad + 4f' f^{-3} H^{-4} w^{-1} \nabla_i u \nabla^i H + 4f' f^{-3} H^{-3} w^{-2} \nabla_i u \nabla^i w + 4f^{-2} H^{-4} w^{-2} \nabla_i H \nabla^i w. \end{aligned}$$

As above, we have

$$\begin{aligned}
& \frac{\partial \Psi}{\partial t} - \operatorname{div}(f^{-1}H^{-2}\nabla\Psi) + 2H^{-1}w|\nabla\Psi|^2 \\
&= -f'f^{-3}H^{-2}w^{-2}u - f'f^{-3}H^{-2}w^{-2}\nabla^i u \langle X, X_i \rangle + (f')^2 f^{-4}H^{-3}w^{-1}|\nabla u|^2 \\
&\quad + f'f^{-3}H^{-4}w^{-1}\nabla_i u \nabla^i H + f'f^{-3}H^{-3}w^{-2}\nabla_i u \nabla^i w \\
&= -f' \frac{u}{f} \Psi^2 - f'f^{-1}\Psi^2 \nabla^i u \langle X, X_i \rangle - f'f^{-2}H^{-2}\nabla_i u \nabla^i \Psi.
\end{aligned}$$

The proof is finished. \square

ACKNOWLEDGMENTS

This research was supported in part by the NSF of China (Grant No. 11926352), the Fok Ying-Tung Education Foundation (China), and Hubei Key Laboratory of Applied Mathematics (Hubei University).

REFERENCES

- [1] B. Andrews, *Gauss curvature flow: the fate of the rolling stones*, Invent. Math. **138**(1) (1999) 151–161.
- [2] L. Chen, J. Mao, Q. Tu, D. Wu, *Asymptotic convergence for a class of inverse mean curvature flows in \mathbb{R}^{n+1}* , Proc. Amer. Math. Soc. **148** (2020) 379–392.
- [3] K. S. Chou, *Deforming a hypersurface by its Gauss-Kronecker curvature*, Commun. Pure Appl. Math. **38**(6) (1985) 867–882.
- [4] K. Ecker, *Regularity Theory for Mean Curvature Flow*, Birkhäuser, Boston, 2004.
- [5] W. J. Firey, *Shapes of worn stones*, Mathematika **21** (1974) 1–11.
- [6] Y. Gao, J. Mao, *Inverse mean curvature flow for spacelike graphic hypersurfaces with boundary in Lorentz-Minkowski space \mathbb{R}_1^{n+1}* , available online at arXiv:2104.10600v5.
- [7] Y. Gao, J. Mao, *An anisotropic inverse mean curvature flow for spacelike graphic hypersurfaces with boundary in Lorentz-Minkowski space \mathbb{R}_1^{n+1}* , Tohoku Math. J. **75** (2023) 347–364.
- [8] Y. Gao, J. Mao, *Inverse Gauss curvature flow in a time cone of Lorentz-Minkowski space \mathbb{R}_1^{n+1}* , available online at arXiv:2108.08686.
- [9] C. Gerhardt, *Flow of nonconvex hypersurfaces into spheres*, J. Differ. Geom. **32** (1990) 299–314.
- [10] C. Gerhardt, *Curvature Problems*, Series in Geom. and Topol., International Press, Somerville, MA. (39) 2006.
- [11] G. Huisken, *Flow by mean curvature of convex surfaces into spheres*, J. Differ. Geom. **20**(1) (1984) 237–266.
- [12] A H. Gong, *A class of inverse mean curvature flow in the Euclid space* (in Chinese), Acta Math. Sinica, Chinese Ser. **67** (2024) 641–651.
- [13] V. A. Ladyženskaja, N. N. Ural'ceva, *Linear and quasilinear elliptic equations*, Academic Press, 1968.
- [14] V. A. Ladyženskaja, V. A. Solonnikov, N. N. Ural'ceva, *Linear and quasilinear equations of parabolic type*, American Mathematical Society, 1968.
- [15] Q. R. Li, W. M. Sheng, X. J. Wang, *Asymptotic convergence for a class of fully nonlinear curvature flows*, J. Geom. Anal. **30** (2020) 834–860.
- [16] G. Lieberman, *Second Order Parabolic Differential Equations*, World Scientific Publishing Co., Singapore, 1996.

- [17] J. Mao, Q. Tu, *A class of inverse curvature flows star-shaped hypersurfaces evolving in a cone*, submitted and available online at arXiv:2104.08884v2.
- [18] T. Marquardt, *The inverse mean curvature flow for hypersurfaces with boundary*, Ph.D. Thesis, Freie Universität Berlin, 2012.
- [19] T. Marquardt, *Inverse mean curvature flow for star-shaped hypersurfaces evolving in a cone*, J. Geom. Anal. **23** (2013) 1303–1313.
- [20] L. Simon, *Lectures on geometric measure theory*, Proceedings of the Center for Mathematical Analysis, Australian National University, Vol. 3, Australian National University, 1983.
- [21] J. Urbas, *On the expansion of starshaped hypersurfaces by symmetric functions of their principal curvatures*, Math. Z. **205** (1990) 355–372.

[†]SCHOOL OF MATHEMATICAL SCIENCE AND ACADEMY FOR MULTIDISCIPLINARY STUDIES, CAPITAL NORMAL UNIVERSITY, BEIJING 100048, CHINA.

Email address: Echo-gaoya@outlook.com

[‡]FACULTY OF MATHEMATICS AND STATISTICS, KEY LABORATORY OF APPLIED MATHEMATICS OF HUBEI PROVINCE, HUBEI UNIVERSITY, WUHAN 430062, CHINA.

Email address: jiner120@163.com