Prescribed duality dynamics in comodule categories

Alexandru Chirvasitu

Abstract

We prove that there exist Hopf algebras with surjective, non-bijective antipode which admit no non-trivial morphisms from Hopf algebras with bijective antipode; in particular, they are not quotients of such. This answers a question left open in prior work, and contrasts with the dual setup whereby a Hopf algebra has injective antipode precisely when it embeds into one with bijective antipode. The examples rely on the broader phenomenon of realizing pre-specified subspace lattices as comodule lattices: for a finite-dimensional vector space V and a sequence $(\mathcal{L}_r)_r$ of successively finer lattices of subspaces thereof, assuming the minimal subquotients of the supremum $\bigvee_r \mathcal{L}_r$ are all at least 2-dimensional, there is a Hopf algebra equipping V with a comodule structure in such a fashion that the lattice of comodules of the r^{th} dual comodule V^{r*} is precisely the given \mathcal{L}_r .

Key words: Diamond Lemma; Tannaka reconstruction; adjoint functor; antipode; comodule; free Hopf algebra; subquotient; triangular

MSC 2020: 16S10; 16T05; 16T15; 16T30; 18A40; 18M05

Introduction

Consider an object in an abelian *left rigid monoidal category* C, i.e. ([16, Definition 3.3.1], [11, §2.10]) on in which each object V has a *left dual* V^{*} [11, Definition 2.10.1]: equipped with morphisms

 $V^* \otimes V \xrightarrow{e \text{ (evaluation)}} \text{monoidal unit } \mathbf{1}, \quad \mathbf{1} \xrightarrow{c \text{ (coevaluation)}} V \otimes V^*$

with

$$V \xrightarrow{id} V \otimes V^* \otimes V \xrightarrow{id \otimes e} V \xrightarrow{id \otimes e} V^* \otimes V \otimes V^* \xrightarrow{e \otimes id} V \xrightarrow{id \otimes e} V^* \xrightarrow{id \otimes e} V^*$$

The (contravariant) functorial nature [11, post Example 2.10.14] of $V \mapsto V^*$ implies that subobjects (quotients) of V produce quotients (respectively subobjects) of V^* . In short, the subobject lattice of V embeds into the dual to that of V^* . That embedding is in general proper (we will recall familiar examples presently), so that iterated duality generates increasingly complicated subobject lattices for the objects $V^{r*} := (V^{(r-1)*})^*$; this is the "duality dynamics" of the paper's title, and one motivating issue is to determine to what extent that "branching" behavior (in the sense that simple objects might acquire non-trivial subobjects after dualization) can be controlled.

The only rigid categories featuring below are those of the form \mathcal{M}_f^H , finite-dimensional (right) comodules over Hopf k-algebras H for a field k. Left rigidity is implemented by means of the antipode S of H [11, Remark 5.3.8], equipping the usual dual vector space V^* of a right H-comodule

 $V \ni v \longmapsto v_0 \otimes v_1 \in V \otimes H$ (Sweedler notation [22, §2.0, pp.32-33])

with the comodule structure defined implicitly by

$$\langle v^*, v_0 \rangle S v_1 = \langle v_0^*, v \rangle v_1^* \in H, \quad \forall v \in V, \ v^* \in V^*, \quad V^* \otimes V \xrightarrow{\text{usual evaluation } \langle -, - \rangle} \Bbbk$$

As noted in the same [11, Remark 5.3.8], this will not, typically, make V^* into a *right* dual [11, Definition 2.10.2] (or *predual*): the usual evaluation is a comodule morphism when defined on $V^* \otimes V$, but not on $V \otimes V^*$; right duals, rather, can be defined analogous using the inverse antipode \overline{S} when it exists (it doesn't always [23, Theorem 11]).

The "control" alluded to two paragraphs up means the degree to which duality can be expected to branch V to just the desired extent and no more. To illustrate the limitations of what can be expected (again, focusing exclusively on categories of comodules), consider an H-comodule V whose dual V^{*} has a 1-dimensional subcomodule $L \leq V^*$. V must then surject onto L^* (as elaborated in Remark 1.15 below; the issue is that left and right duals coincide for 1-dimensional Hopf-algebra comodules), so it could not, for instance, have been irreducible (if at least 2-dimensional).

In part, the goal is to confirm that this is essentially the only difficulty. Extend the notation V^{r*} employed above to negative r, in which case it denotes successive right duals of V. As noted, the distinction does matter once we equip spaces with comodule structures. Corollary 1.11 to Theorem 1.10 yields, via Construction 1.14, the following conclusion.

Theorem A Let V be a finite-dimensional vector space and $(\mathcal{L}_r)_{r\geq d}$ a sequence of increasingly finer lattices of subspaces of V for some $d \in \mathbb{Z}_{\leq 0} \sqcup \{-\infty\}$.

If the minimal subquotients of the supremum lattice $\bigvee_r \mathcal{L}_r$ are all of dimension ≥ 2 there is a Hopf algebra H equipping V^{r*} with comodule structures

$$V^{r*} \xrightarrow{\rho_r} V^{r*} \otimes H, \quad r \in \mathbb{Z}_{>d}$$

so that

• ρ_{r+1} is (left) dual to ρ_r ;

• and for every r the lattice of H-subcomodules of V^{r*} is precisely \mathcal{L}_r^{r*} (meaning \mathcal{L}_r if r is even and its dual if not).

It is not unnatural, at this point, to link back to antipode bijectivity, given the already-mentioned relationship between inverse antipodes and right duality (we refer the reader once more to [11, Remark 5.3.8]). The following observations will serve as a starting point:

• Plainly, Hopf algebras surjected upon (embeddable into) those with bijective antipode have surjective (respectively injective) antipode.

• One of the converses holds: a Hopf algebra with injective antipode embeds into one with bijective antipode [21, Proposition 2.7] (universally, i.e. the embedding is the *unit* [1, Definition 19.3] of an adjunction between Hopf algebras and Hopf algebras with bijective antipode).

The inevitable problem of whether the other converse holds is posed as [6, Question 3]. The first example [21, §3] of a Hopf algebra with surjective, non-injective antipode is explicitly constructed as a quotient of a free Hopf algebra with bijective antipode, so is tautologically not a counterexample to an affirmative answer.

The connection to the preceding material is through the *coefficient coalgebras* C_W attached to *H*-comodules *W* (we recall the notion more extensively in Construction 1.14): the smallest subcoalgebra for which the comodule structure factors as

$$W \xrightarrow{W \otimes C_W} W \otimes H.$$

Given that $C_{W^*} = S(C_W)$, the antipode fails to be injective precisely to the extent to which duality enlarges subspace lattices. This suggests that if the dynamics of a surjective antipode is ill-behaved enough, perhaps the Hopf algebra in question will fail to be surjected upon by one with bijective antipode. Example 1.21 confirms this (in a stronger form than just stated), as a consequence of Corollary 1.20 to Theorem 1.18.

Theorem B There exist Hopf algebras with surjective antipode which do not admit non-trivial morphisms from Hopf algebras with bijective antipode.

1 Free Hopf algebras on coalgebra chains and main results

(Co)algebras are (co)unital, (co)associative and, along with everything else, k-linear for some field k. (Co)op superscripts indicate (co)opposite (co)multiplications [19, Definitions 2.1.1 and 2.1.4].

We take for granted the various universal constructions peppered throughout (free objects and more general types of adjoint functors), as justified by the fact that all categories involved are *locally presentable* in the sense of [2, Definition 1.17]. Sources for this, for the categories featuring most prominently below (algebras ALG, coalgebras COALG, bialgebras BIALG, Hopf algebras HALG or Hopf algebras with bijective antipode $HALG_{\overline{S}}$) and more, include (for instance) [18, Lemmas 1 and 2, Theorem 6 and Proposition 22 4.].

We will need the following variant of the free Hopf algebra H(C) on a coalgebra C [23, Definition 2], i.e. the left adjoint to the forgetful functor HALG \rightarrow COALG.

Definition 1.1 Let $d \in \mathbb{Z}_{\leq 0} \sqcup \{-\infty\}$ be a non-positive extended integer.

(1) A skew chain $(C^r, \theta^r)_{i \geq d}$ of coalgebras is a diagram

$$\cdots \xrightarrow{\theta^{-1}} C^0 \xrightarrow{\theta^0} (C^1)^{cop} \xrightarrow{\theta^1} C^2 \xrightarrow{\theta^2} \cdots \text{ in COALG},$$

with the co-opposites alternating. We write $COALG_{>d}$ for the category of skew coalgebra chains.

(2) The free Hopf algebra $H(\mathbf{C})$ on a skew chain $\mathbf{C} = (C^r, \theta^r)$ is (the image of \mathbf{C} through) the left adjoint to the forgetful functor

$$\mathrm{HALG} \ni H \xrightarrow{\mathrm{FGT}} \left(\cdots \xrightarrow{\mathrm{antipode}\ S} H \xrightarrow{S} H^{op,cop} \xrightarrow{S} H \xrightarrow{S} \cdots \right) \in \mathrm{COALG}_{\geq d}$$

 $H := H(C^r, \theta^r)$ naturally comes equipped with coalgebra morphisms $C^r \xrightarrow{\iota_r} H$ intertwining the antipode on the codomain and the θ^r on the domain(s).

Remark 1.2 Mapping from C^r to $(C^{r+1})^{cop}$ rather than C^{r+1} is only a matter of choice in labeling; the category is of course equivalent to that of functors $\mathbb{Z}_{\geq d} \to \text{COALG}$, with the domain, a poset, regarded as a category as usual [5, Example 1.2.6.b]: exactly one arrow \to for each relation \leq . As

a functor category, it is locally presentable [2, Corollary 1.54] along with its codomain COALG. In particular, the existence of the left adjoint

$$\operatorname{Coalg}_{>d} \longrightarrow \operatorname{HAlg}$$

taken for granted in Definition 1.1(2) is unproblematic, by any number of *adjoint functor theorems* ([1, §18], [2, §0.7], etc.). It is also a simple enough matter to give a (relatively) concrete description of $H(\mathbf{C})$, as we do in the proof of Proposition 1.4.

Instances of the construction of particular interest here:

Examples 1.3 (1) The aforementioned [23, Definition 2] motivating free Hopf algebra H(C) on a coalgebra C can be retrieved as $H(C^r, \theta^r \mid r \ge 0)$ with

$$C^{r} := \begin{cases} C & \text{if } r \text{ is even} \\ C^{cop} & \text{otherwise} \end{cases} \quad \text{and} \quad \theta^{r} = \text{id}, \quad \forall r.$$

$$(1-1)$$

(2) In precisely the same fashion one recovers the free Hopf algebra with bijective antipode on C denoted by $\widehat{H}(C)$ in [21, Lemma 3.1]. The only difference is that this time the chain is bi-infinite, i.e. $\widehat{H}(C) \cong H(C^r, \theta^r \mid r \in \mathbb{Z})$ with C^r and θ^r still as in (1-1).

It will be convenient to abbreviate the description of C^r in (1-1) as $C^r = C^{r \cdot cop}$: applying the *cop* operator an even number of times amounts to doing nothing, and the multiplicative notation $r \cdot cop$ seems less burdensome than the exponential version cop^r .

(3) Even more pertinent to the sorts of phenomena relevant in the present context, consider the quotient

$$H := \widehat{H}(M_4^*)/I := (\text{Hopf ideal generated by } x_{ij}^0, \ i \ge 3, \ j \le 2)$$

where $M_4^* := M_4(\mathbb{k})^*$ is the coalgebra dual to the matrix algebra, $x_{ij}^0 \in M_4^*$ are the basis elements dual to the standard matrix units in M_4 , and the '0' superscript indicates they belong to the initial generating copy of $M_4^* \leq \hat{H}(M_4^*)$ rather than to any antipode iterates thereof. Per the conventions of [21, §3] (also [15, §2]),

$$S^{r}M_{4}^{*} = \operatorname{span} \left\{ x_{ij}^{r} \mid 1 \leq i, j \leq 4 \right\}, \ \forall r \in \mathbb{Z} \text{ and } Sx_{ij}^{r} = x_{ji}^{r+1}.$$

We have $H \cong H(C^r, \theta^r \mid r \in \mathbb{Z})$ for

$$C^r := \begin{cases} (M_4^*)^{r \cdot cop} & \text{if } r < 0\\ (M_4^*/\operatorname{span} \{x_{ij} \mid i \ge 3, \ j \le 2\})^{r \cdot cop} & \text{otherwise,} \end{cases}$$

with the obvious morphisms (identities or surjections). In words: the chain, traveling rightward along identities from $-\infty$, collapses M_4^* to its quotient by the four matrix counits at step 0 and stabilizes afterwards.

As a first general observation, just as the free Hopf algebra construction H(-) (per [23, Corollary 8]), free Hopf algebras on skew chains are compatible with scalar extensions.

Proposition 1.4 For any skew coalgebra chain $\mathbf{C} = (C^r, \theta^r)$ over \Bbbk and field extension $\Bbbk \leq \Bbbk'$ the canonical morphism $H(\mathbf{C}) \otimes \Bbbk' \to H(\mathbf{C} \otimes \Bbbk')$ for

$$\mathbf{C}\otimes \Bbbk':=(C^r\otimes \Bbbk', heta^r\otimes \mathrm{id}_{\Bbbk'})$$

is an isomorphism.

Proof An explicit description of $H(\mathbf{C})$:

• form the usual free coalgebra $H(\bigoplus_r C^r)$, equipped with its injective [23, Corollary 9] structure map

$$\bigoplus C^r \xleftarrow{\iota} H\left(\bigoplus C^r\right),$$

henceforth suppressed, identifying C^r with subcoalgebras of its codomain;

• and quotient out the ideal generated by $\theta^r(x^r) - Sx^r$ for $x^r \in C^r$ and all of their iterates under S.

The (algebra) ideal generated by those differences is already a coideal, so the result is indeed a Hopf algebra; the requisite universality property holds tautologically.

The two bullet points are both invariant under field extensions (the first by the aforementioned [23, Corollary 9]), hence the conclusion.

Remark 1.5 In light of Proposition 1.4 one might occasionally, in analyzing Hopf algebras of the form $H(\mathbf{C})$, assume the ground field algebraically closed. This is sometimes useful for instance, in ensuring that *simple* coalgebras (i.e. [14, 2.4.1(1)] those with no non-obvious subcoalgebras) are matrix coalgebras M_n^* . This is dual to the remark [17, §3.5, Corollary b.] that M_n are the only finite-dimensional simple algebras over algebraically closed fields.

The collapse noted in Example 1.3(3) will replicate in a more general setting, requiring some tooling. We will be concerned, specifically, with skew chains (C^r, θ^r) where each C^r is a dual block-triangular matrix algebra, upper (lower) if r is even (respectively odd), and each θ^r a surjection refining the block-triangular shape. We elaborate.

Definition 1.6 Fix $d \in \mathbb{Z}_{\leq 0} \sqcup \{-\infty\}$, as in Definition 1.1. A *(block-upper-)triangular* $\mathbb{Z}_{\geq d}$ -indexed skew coalgebra chain is one of the following form.

• Each coalgebra C^r , $r \ge d$ has a basis consisting of matrix counits x_{ij}^r in the sense that

$$x_{ij}^r \xrightarrow{\Delta} \sum_{u} x_{iu}^r \otimes x_{uj}^r, \quad \varepsilon(x_{ij}^r) = \delta_{ij} := \begin{cases} 1 & \text{if } i = j \\ 0 & \text{otherwise.} \end{cases}$$

The θ^r operate by

$$x_{ij}^r \xrightarrow{\theta^r} x_{ji}^{r+1}$$
 or 0. (1-2)

We sometimes simply write $x_{ij}^r \mapsto x_{ji}^{r+1}$, understanding that this might vanish if the respective matrix counit is not present in C^{r+1} .

• The subscripts of x_{ij}^r range over a possibly infinite poset (\mathcal{I}, \leq) (common to all r), so that (1-2) does indeed make sense. The connected components of the order \leq are finite, and for each r the subscripts of x_{ij}^r range over a block-triangular pattern with respect to \leq : upper for even r and lower for odd r.

Remarks 1.7 (1) The conditions of Definition 1.6 imply in particular that each C^r is a coproduct (direct sum)

$$C^{r} \cong \bigoplus_{\leq_{r} \text{-connected component } \mathcal{I}_{rs}} C^{rs}, \quad C^{rs} = \operatorname{span} \left\{ x_{ij}^{r} \mid i, j \in \mathcal{I}_{rs} \right\}$$

of finite-dimensional triangular dual matrix coalgebras.

(2) Each triangular pattern imposes an equivalence relation \sim_r on \mathcal{I} :

$$i \sim_r j \iff x_{ij}^r$$
 and x_{ji}^r are both present.

The relation \sim_r become progressively finer with increasing r.

Example 1.8 In Example 1.3(3) (\mathcal{I}, \leq) is simply $\overline{1, 4}$ with the usual ordering. \sim_r is the coarsest relation (one equivalence class) for r < 0 has equivalence classes $\{1, 2\}$ and $\{3, 4\}$ otherwise.

Notation 1.9 In the context of Definition 1.6:

(1) Somewhat abusively, we will also denote by x_{ij}^r the images of those elements through the structure maps $C^r \xrightarrow{\iota_r} H(C^r, \theta^r)$; the latter's antipode thus operates by $Sx_{ij}^r = x_{ji}^{r+1}$.

More generally, following [7, \$1, p.103], for length-t tuples

$$\mathbf{r} = (r_s)_{1 \le s \le t} \quad \mathbf{i} = (i_s)_{1 \le s \le t} \quad \mathbf{j} = (j_s)_{1 \le s \le t}$$

set

$$x_{\mathbf{i}\mathbf{j}}^{\mathbf{r}} := \text{product } x_{i_1j_1}^{r_1} \cdots x_{i_tj_t}^{r_t} \text{ in } H(C^r, \theta^r).$$

(2) In discussing products of matrix counits x_{ij}^r , the two symbols \uparrow and \downarrow indicate the largest (respectively lowest) row/column index for which the factors respective factors are all non-zero in their respective C^r . Some examples follow.

• $x_{i\uparrow}^r$ is the rightmost non-zero matrix counit on row *i* in C^r .

• in $x_{i\downarrow}^r x_{j\downarrow}^s$ the ' \downarrow ' stands for the lowest index that will make both factors non-zero in C^r and C^s respectively. Which specific symbol that is depends only on $\max(r, s)$ (as more of the matrix counits vanish in "later" C^r).

• [15, Theorem 5], describing a k-basis for $H(M_n^*)$, can be phrased in the present language: that basis consists of those words $x_{ij}^{\mathbf{r}}$ (for tuples \mathbf{r} over $\mathbb{Z}_{\geq 0}$ and \mathbf{i} over $\overline{1, n}$) which contain no subwords of the form

$$x_{i\uparrow}^r x_{j\uparrow}^{r+1} \quad x_{\uparrow i}^r x_{\uparrow j}^{r-1} \quad x_{i\uparrow}^r x_{j(\uparrow-1)}^{r+1} x_{k(\uparrow-1)}^{r+2} \quad x_{\uparrow i}^r x_{(\uparrow-1)j}^{r-1} x_{(\uparrow-1)j}^{r-2}$$

• Precisely the same goes for the basis described over the course of (the proof of) [21, Theorem 3.2] for $\hat{H}(M_n^*)$, except this time the superscripts range over all of \mathbb{Z} .

Naturally, in both of these examples $\uparrow = n$ throughout, so there is not much point to the substitution, but in the broader setting below there will be.

• [8, §2] takes a slightly different approach we will find handier here, dispensing with the cubic relations (and the shift in $\uparrow -1$). The bases for $H(M_n^*)$ and $\hat{H}(M_n^*)$ described there are rather those consisting of words containing none of

$$x_{i\uparrow}^r x_{j\uparrow}^{r+1} (r \text{ even}) \qquad x_{i\downarrow}^r x_{j\downarrow}^{r+1} (r \text{ odd}) \qquad x_{\uparrow i}^{r+1} x_{\uparrow j}^r (r \text{ odd}) \qquad x_{\downarrow i}^{r+1} x_{\downarrow j}^r (r \text{ even})$$
(1-3)

as subwords. It is this description that the sequel builds upon.

The usefulness of the $x_{\uparrow\uparrow}^r$ is evident in the statement of the following result, where the book-keeping for which indices are involved and which are not might become cumbersome.

۲

Theorem 1.10 Let $(C^r, \theta^r)_{r \ge d}$ be an upper-triangular skew coalgebra chain in the sense of Definition 1.6. If the classes of the supremum

$$\bigvee_{r} \sim_{r}, \quad \sim_{r} = equivalence \ relation \ of \ Remark \ 1.7(2)$$

all have size ≥ 2 the corresponding k-Hopf algebra $H(C^r, \theta^r)$ has a basis consisting of those words in x_{ij}^r containing no subwords of the form (1-3).

Proof As in all cited sources, this will be an application of the *Diamond Lemma* [3, Theorem 1.2] (which we assume as background, along with its ancillary language and machinery).

As an algebra, $H := H(C^r, \theta^r)$ is the quotient of the free algebra $\Bbbk \langle x_{ij}^r \rangle$ by the relations making $(x_{ij}^r)_{i,j}$ and $(Sx_{ij}^r)_{i,j}$ mutual inverses in an appropriately-sized matrix algebra over H. Said relations translate to the substitutions (or *reductions* [3, p.180])

$$x_{i\uparrow}^r x_{j\uparrow}^{r+1} \longmapsto \delta_{ij} - \sum_{\alpha < \uparrow} x_{i\alpha}^r x_{j\alpha}^{r+1}$$
 (r even) (1-4)

$$x_{i\downarrow}^r x_{j\downarrow}^{r+1} \longmapsto \delta_{ij} - \sum_{\alpha > \downarrow} x_{i\alpha}^r x_{j\alpha}^{r+1} \quad (r \text{ odd})$$
(1-5)

$$x_{\uparrow i}^{r+1} x_{\uparrow j}^r \longmapsto \delta_{ij} - \sum_{\beta < \uparrow} x_{\beta i}^{r+1} x_{\beta j}^r \quad (r \text{ odd})$$
(1-6)

$$x_{\downarrow i}^{r+1} x_{\downarrow j}^r \longmapsto \delta_{ij} - \sum_{\beta > \downarrow} x_{\beta i}^{r+1} x_{\beta j}^r \quad (r \text{ even})$$
(1-7)

These are also [8, equations (2.1) to (2.4)] in that paper's more constrained setting; it is also observed there that the four possible types of *(overlap) ambiguity* [3, post Lemma 1.1] resulting from these reductions are all effectively equivalent.

$$x_{i\uparrow}^r x_{\downarrow\uparrow}^{r+1} x_{\downarrow\downarrow}^r \quad (r \text{ even}) \quad \iff \quad x_{\uparrow i}^{r+1} x_{\uparrow\downarrow}^r x_{j\downarrow}^{r+1} \quad (r \text{ odd}),$$
(1-8)

for instance, in the sense that the computations become symbolically identical, after interchanging the two superscript symbols and the positions of the two subscripts on every $x_{\bullet\bullet}^{\bullet}$. Similar remarks apply to the others, so we fix ideas by focusing on a single one of the four: the right-hand side of (1-8). Note that at this stage we have already made surreptitious use of the hypothesis of having size-(≥ 2) classes: \uparrow and \downarrow are distinct, so there are no "cubical" ambiguities $x_{i\bullet}^r x_{j\bullet}^{r+1} x_{k\bullet}^{r+2}$.

Resolvability was left to the reader on [8, p.86], so we verify it here for some semblance of completeness.

$$\begin{pmatrix} \delta_{i\downarrow} - \sum_{\beta < \uparrow} x_{\beta i}^{r+1} x_{\beta\downarrow}^{r} \end{pmatrix} x_{j\downarrow}^{r+1} \xrightarrow{(1-6)} x_{\uparrow i}^{r+1} x_{\uparrow\downarrow}^{r} x_{j\downarrow}^{r+1} \xrightarrow{(1-5)} x_{\uparrow i}^{r+1} \begin{pmatrix} \delta_{\uparrow j} - \sum_{\alpha > \downarrow} x_{\uparrow \alpha}^{r} x_{j\alpha}^{r+1} \end{pmatrix} x_{j\downarrow}^{r+1} \xrightarrow{(1-5)} x_{\uparrow i}^{r+1} \begin{pmatrix} \delta_{\uparrow j} - \sum_{\alpha > \downarrow} x_{\uparrow \alpha}^{r} x_{j\alpha}^{r+1} \end{pmatrix} x_{\uparrow i}^{r+1} - \sum_{\beta < \uparrow} \delta_{\beta j} x_{\beta i}^{r+1} + \sum_{\substack{\beta < \uparrow \\ \alpha > \downarrow}} x_{\beta i}^{r+1} x_{\beta \alpha}^{r} x_{j\alpha}^{r+1} \xrightarrow{\delta_{\uparrow j}} x_{\uparrow i}^{r+1} - \sum_{\alpha > \downarrow} \delta_{\alpha i} x_{j\alpha}^{r+1} + \sum_{\substack{\alpha > \downarrow \\ \beta < \uparrow}} x_{\beta i}^{r+1} x_{\beta \alpha}^{r} x_{j\alpha}^{r+1}$$

The cubic sums plainly coincide; we thus have

$$\operatorname{left} - \operatorname{right} = \left(\delta_{i\downarrow} x_{j\downarrow}^{r+1} + \sum_{\alpha > \downarrow} \delta_{\alpha i} x_{j\alpha}^{r+1} \right) - \left(\delta_{\uparrow j} x_{\uparrow i}^{r+1} + \sum_{\beta < \uparrow} \delta_{\beta j} x_{\beta i}^{r+1} \right)$$
$$= \sum_{\bullet} \delta_{i\bullet} x_{j\bullet}^{r+1} - \sum_{\bullet} \delta_{\bullet j} x_{\bullet i}^{r+1}$$
$$= x_{ji}^{r+1} - x_{ji}^{r+1} = 0,$$

effecting the resolution.

In applications of the Diamond Lemma one typically also produces a monoid partial order [3, p.181] on the free monoid $\langle X \rangle$ on the set X generators (here $X = \left\{ x_{ij}^r \right\}$). One is given in [8, pp.86-87] which adapts readily here; however, as pointed out in [3, §5.4], over fields (or more generally, in the absence of zero divisors) there is a canonical order that will work provided we know (as we now do, in light of the just-verified resolvability) that all elements are *reduction-finite* (i.e. [3, p.180] all chains of modifications involving (1-4) to (1-7) eventually stabilize):

• a word $w \in \langle X \rangle$ is simply *declared* (strictly) smaller than another, w', if w occurs with non-zero coefficient in some reduction of w';

• the condition on zero divisors will then ensure that the resulting relation < is asymmetric (i.e. [13, Tayble 14.1] w < w' is incompatible with w' < w);

• and by the assumed reduction finiteness it has the descending chain condition (DCC) [4, VIII.1] (admits no infinite chains $w_0 > w_1 > \cdots$).

The validity of the general construction notwithstanding, it certainly functions here: each of the substitutions (1-4) to (1-7) replaces a word with a linear combination of words either strictly shorter or with strictly fewer occurrences of the problematic two-letter left-hand subwords, so that asymmetry and DCC are both self-evident.

In particular, because single-letter words of course have no 2-letter subwords, we have the following counterpart to the injectivity [23, Corollary 9] of the map $C \to H(C)$:

Corollary 1.11 In the setup of Theorem 1.10 the canonical maps $C^s \xrightarrow{\iota_s} H(C^r, \theta^r)$ of Definition 1.1(2) are injective.

Observe also that the only generators entering any of the problematic words (1-3) are those featuring in diagonal blocks of both C^r and C^{r+1} . For that reason, the "off-diagonal generators" simply come along for the ride so to speak, generating an algebra that splits off as a free factor. This is very much analogous to the fact [23, Theorem 32] that for any splitting

$$C = C_0 \oplus V$$
, $C_0 := coradical \text{ of } C \stackrel{[14, \text{ Definition 5.1.5(1)}]}{:=} \sum (\text{simple subcoalgebras})$

we have the algebra-coproduct decomposition $H(C) \cong H(C_0) \coprod TV$ (with TV denoting the tensor algebra of V, i.e. the free algebra on V). Before stating the present analogue, a piece of terminology.

Definition 1.12 The asymptotic coradical $\mathbf{C}_{0\leftarrow}$ of a skew coalgebra chain $\mathbf{C} = (C^r, \theta^r)$ with surjective connecting maps θ^r is the filtered [2, Definition 1.4] colimit

$$\lim_{s \ge d} \mathbf{C}_{0\leftarrow}^{\ge s} := \left(\text{coradical } C_0^s \xrightarrow{\theta^s} \theta^s(C_0^s) \xrightarrow{\theta^{s+1}} (\theta^{s+1} \circ \theta^s)(C_0^s) \xrightarrow{\theta^{s+2}} \cdots \right).$$

The colimit is taken over decreasing s, the connecting maps

$$\mathbf{C}_{0\leftarrow}^{\geq s} \hookrightarrow \mathbf{C}_{0\leftarrow}^{\geq s-1}$$

being the embeddings resulting from the fact [14, Corollary 5.3.5] that the image of the coradical C_0 through a surjection $C \rightarrow D$ contains the coradical D_0 .

Corollary 1.13 For a skew coalgebra chain $\mathbf{C} = (C^r, \theta^r)$ as in Theorem 1.10 we have an algebracoproduct decomposition

$$H(\mathbf{C}) \cong H(\mathbf{C}_{0\leftarrow}) \coprod TV \cong H(\mathbf{C}_{0\leftarrow}) \coprod \left(\coprod_r T(V^r) \right)$$

for

$$V := \bigoplus V^r, \quad C^r = \mathbf{C}^r_{0\leftarrow} \oplus V^r.$$

The passage between the subspace-lattice picture emphasized in the Introduction (Theorem A) and the more coordinate/basis-oriented perspective of Theorem 1.10 is by means of the comodule-coalgebra correspondence familiar from *Tannaka reconstruction* ([20, §2], [16, Chapter 3], etc.). To expand:

Construction 1.14 • A subspace lattice \mathcal{L} on a finite-dimensional vector space V generates a smallest full *exact subcategory* [12, §3.4]

$$\mathcal{C}_{\mathcal{L}} \xrightarrow{\iota_{\mathcal{L}}} \operatorname{VECT}_{f} := \text{finite-dimensional vector spaces.}$$

The inclusion functor $\iota_{\mathcal{L}}$ has a *coendomorphism coalgebra* [20, Definition 2.1.8], equipped with a surjection

$$\operatorname{Coend}(V) \cong M_{\dim V}(\Bbbk)^* \longrightarrow \operatorname{Coend}(\iota_{\mathcal{L}}) =: C_{\mathcal{L}};$$
(1-9)

This is the model for the surjections $C^r \to (C^{r+1})^{cop}$ featuring in a triangular skew chain, coopposites reflecting the dualization in Theorem A.

• The finite-dimensional comodule category $\mathcal{M}^{C_{\mathcal{L}}}$ is precisely $\mathcal{C}_{\mathcal{L}}$ (one version of the aforementioned Tannakian reconstruction [20, Theorem 2.2.8]). Subquotients W of V belonging to $\mathcal{C}_{\mathcal{L}}$, being $C_{\mathcal{L}}$ -comodules, correspond to subcoalgebras of the latter:

 $W \mapsto coefficient \ coalgebra \ C_W$ (associated to the comodule as in [9, Proposition 2.5.3]):

the smallest subcoalgebra of $C_{\mathcal{L}}$ for which the structure map $W \to W \otimes C_{\mathcal{L}}$ takes values in $W \otimes C_W$.

• One can now select a basis for V, appropriately compatible with \mathcal{L} , so as to ensure a block upper-triangular shape for $C_{\mathcal{L}}$. The minimal (non-zero) members of \mathcal{L} sum up to the *socle* [9, Remark 2.4.10(2)] of $V \in \mathcal{M}^{C_{\mathcal{L}}}$: the largest semisimple sup

soc
$$V = \bigoplus_{i} W_i \le V$$
, $W_i \in \mathcal{M}^{C_{\mathcal{L}}}$ simple.

The matrix coalgebras $C_{W_i} \cong \text{Coend}(W_i) \cong M^*_{\dim W_i}$ will be the first diagonal blocks in the (upper) triangular decomposition, with no off-diagonal blocks linking any two:

$$C_{\mathcal{L}} \cong \begin{pmatrix} C_{W_1} & 0 & 0 & \vdots \\ 0 & C_{W_2} & 0 & \vdots \\ 0 & 0 & C_{W_3} & \vdots \\ \vdots & \vdots & \vdots & \vdots \end{pmatrix}$$

• One can then proceed along the *socle filtration*

$$\{0\} = \operatorname{soc}_0 V \le \operatorname{soc}_1 V := \operatorname{soc} V \le \operatorname{soc}_2 V \le \cdots \text{ of } V \text{ in } \mathcal{M}^{C_{\mathcal{L}}},$$

where

$$s^{th}$$
 socle layer $\overline{\operatorname{soc}}_s := \operatorname{soc}_s / \operatorname{soc}_{s-1} := \operatorname{soc}(W / \operatorname{soc}_{s-1})$

The off-diagonal blocks of $C_{\mathcal{L}}$, if any, will link matrix coalgebras C_W and $C_{W'}$ for simple comodules W and W' occurring as summands in distinct socle layers.

Remark 1.15 The hypothesis on subquotients having dimension ≥ 2 is necessary in Corollary 1.11, as even minimal violations thereof will produce counterexamples.

Consider for instance an upper-triangular skew chain $(C^r, \theta^r)_{r>-1}$ with

- C^{-1} a full $n \times n$ matrix algebra for some $n \ge 2$;
- and C^r , $r \ge 0$ triangular, each with two diagonal blocks of sizes 1×1 and $(n-1) \times (n-1)$.

I then claim that $C^{-1} \xrightarrow{\iota_{-1}} H := H(C^r, \theta^r)$ cannot be injective (it factors through a triangular quotient of C^{-1} , again with one 1×1 diagonal block).

This can be checked coordinate-wise (much as the argument driving the related [21, Remark 3.3] does), but is particularly transparent in monoidal-categorical terms (the connection between the two pictures being as in Construction 1.14). Denote by V^r the natural *n*-dimensional C^r -comodule, resulting from the realization of C^r as a quotient of an $n \times n$ matrix coalgebra. Each V^r then also becomes an *H*-comodule by transporting the structure along $C^r \xrightarrow{\iota_r} H$, and the 1×1 block in C^0 means that V^0 has a 1-dimensional C^{0-} (hence also H-)subcomodule *L*.

Now, V^{-1} is the predual $*(V^0)$. Over a Hopf algebra, the coefficient coalgebra of a 1-dimensional comodule L is the span of a grouplike [14, Definition 1.3.4(a)] and the latter's inverse [14, Example 1.5.3] spans a 1-dimensional comodule that is both left and right dual to the original L. In conclusion, the predual V^{-1} of V^0 must surject onto the 1-dimensional $L^* \cong *L$. This will give the image through $C^{-1} \stackrel{\iota_{-1}}{\longrightarrow} H$ a lower-triangular structure with a 1×1 rightmost diagonal block.

We can now return to the statements of the Introduction.

Proof of Theorem A I claim that this is essentially what Corollary 1.11 provides, after producing coalgebras out of subspace lattices as described in Construction 1.14. Set

$$C^{r} := \begin{cases} \text{the coendomorphism coalgebra } C_{\mathcal{L}_{r}} \text{ of } (1-9) & \text{if } r \text{ is even} \\ C_{\mathcal{L}_{r}}^{cop} \cong C_{\mathcal{L}_{r}^{*}} & \text{otherwise,} \end{cases}$$

with \mathcal{L}_r^* denoting the lattice on V^* dual to \mathcal{L}_r . The surjections $C^r \xrightarrow{\theta^r} (C^{r+1})^{cop}$ are those analogous to (1-9), resulting from the refinement $\mathcal{L}_r \rightsquigarrow \mathcal{L}^{r+1}$, and Construction 1.14 also outlines how the coalgebras in question acquire a triangular structure.

This, then, will produce an upper triangular coalgebra chain $\mathbf{C} = (C^r, \theta^r)$ and a free Hopf algebra $H := H(\mathbf{C})$ thereon, with the (dim ≥ 2) hypothesis of Theorem A assuring that of Theorem 1.10. As to the desired conclusion (of Theorem A), it follows from Corollary 1.11 given that [20, Lemma 2.2.12] *embeddings* $C \hookrightarrow D$ of coalgebras induce full, exact inclusions $\mathcal{M}^C \hookrightarrow \mathcal{M}^D$ between the corresponding comodule categories.

A brief remark, prefatory to eventually addressing Theorem B:

Lemma 1.16 Let $\mathbf{C} = (C^r, \theta^r)$ be an upper-triangular skew coalgebra chain meeting the constraints of Theorem 1.10, and $C \leq H(\mathbf{C})$ a simple subcoalgebra.

There is a unique longest tuple $\mathbf{r} = (r_1, \cdots, r_t) \in \langle \mathbb{Z}_{\geq d} \rangle$ (the free monoid on $\mathbb{Z}_{\geq d}$) for which the expansion of every non-zero element of C in the basis of Theorem 1.10 contains some word $x_{ii}^{\mathbf{r}}$.

Proof The Hopf algebra $H := H(\mathbf{C})$ is the sum of its subcoalgebras

$$C^{\mathbf{r}} := C^{r_1} \cdot C^{r_2} \cdots C^{r_t},$$

having identified C^r with a subcoalgebra of H (as Corollary 1.11 allows). C is thus a subcoalgebra of some $C^{\mathbf{r}}$, and the distinguished \mathbf{r} of the statement will be the *minimal* such, length-wise: the relations (1-4) to (1-7) make it clear that $C^{\mathbf{r}}$ consists of linear combinations of $x_{\mathbf{ij}}^{\mathbf{r}'}$ for subwords \mathbf{r}' of \mathbf{r} , hence uniqueness.

Definition 1.17 The tuple $\mathbf{r} \in (\mathbb{Z}_{\geq d})^s$ attached by Lemma 1.16 to a simple subcoalgebra $C \leq H(\mathbf{C})$ is the $(\langle \mathbb{Z}_{\geq d} \rangle$ -valued) rank of C.

The term applies also to the (unique up to isomorphism) simple C- (hence also $H(\mathbf{C})$ -)comodule.

Theorem 1.18 Let $\mathbf{C} = (C^r, \theta^r)$ be a skew coalgebra chain as in Theorem 1.10, $\mathbf{r} = (r_s)_{1 \leq s \leq t} \in \langle \mathbb{Z}_{>d} \rangle$, and

 $D^{r_s} \leq C^{r_s}, \quad 1 \leq s \leq t \quad simple \ subcoalgebras.$

Any simple $H(\mathbf{C})$ -subcoalgebra of $D^{\mathbf{r}} := \prod_{1 \le s \le t} D^{r_s}$ of rank $\mathbf{r} = (r_1, \cdots, r_t)$ has dimension $\ge \max_s \dim D^{r_s}$.

Proof Note that the D^{r_s} are in any case matrix coalgebras, corresponding to square diagonal blocks of C^{r_s} (respectively). There is no harm in assuming the ground field algebraically closed (Remark 1.5), so that the simple rank- $\mathbf{r} \ C \le D^{\mathbf{r}}$ that concerns us is itself a matrix coalgebra. To fix the notation, we prove only that dim $C \ge \dim D^{r_1}$.

The idea behind the proof is that underlying [7, Proposition 2.6]. Let $x \in C$, containing the maximal-length term $x_{ij}^{\mathbf{r}}$ in its expansion in the basis of Theorem 1.10. The s^{th} letter $x_{is,js}^{r_s}$ of $x_{ij}^{\mathbf{r}}$ is one of the basis elements for the matrix coalgebra D^{r_s} , so its subscripts i_s and j_s range independently over a subset

$$\mathcal{I}_s \subseteq \mathcal{I}, \quad |\mathcal{I}_s| = n_s := \sqrt{\dim D^{r_s}}.$$

We construct n_1 length-t tuples $\mathbf{u} = (u_s)_{1 \le s \le t}$ over \mathcal{I} as follows:

• pick the first letter $u_1 \in \mathcal{I}_1$ arbitrarily;

• then pick the second letter $u_2 \in \mathcal{I}_2$ distinct from u_1 (possible by the (≥ 2) assumption transported here from Theorem 1.10);

- then the third letter $u_3 \in \mathcal{I}_3$ distinct from u_2 ;
- and so on.

The choice is arbitrary over \mathcal{I}_1 in first instance, and then ensures that no two consecutive letters in **u** coincide. Now note that

$$x_{\mathbf{ij}}^{\mathbf{r}} \xrightarrow{\Delta} \sum_{\mathbf{u} \text{ chosen in this fashion}} x_{\mathbf{iu}}^{\mathbf{r}} \otimes x_{\mathbf{uj}}^{\mathbf{r}} + \cdots,$$
 (1-10)

where the \cdots signify tensor products of *other* reduced words. The tensorands $x_{iu}^{\mathbf{r}}$ and $x_{uj}^{\mathbf{r}}$ are indeed reduced, by the very shape of the reductions (1-4) to (1-7): the consecutive letters of **u** are distinct so there is no opportunity for reduction there, whereas the **i** and **j** offer no such opportunity by assumption, $x_{ij}^{\mathbf{r}}$ having been assumed reduced to begin with.

The outer \mathbf{i} and \mathbf{j} on the right-hand side of (1-10) identify the word on the left-hand side that produced the tensor product by comultiplication, so no other terms in the expansion of x will produce any of the tensor products $x_{\mathbf{iu}}^{\mathbf{r}} \otimes x_{\mathbf{uj}}^{\mathbf{r}}$. Denoting the '*' superscripts the functionals "dual" to the word basis of Theorem 1.10, as in

$$x_{\mathbf{uv}}^{\mathbf{s}*}\left(x_{\mathbf{u'v'}}^{\mathbf{s'}}\right) = \delta_{\mathbf{r},\mathbf{r'}}\delta_{\mathbf{u},\mathbf{u'}}\delta_{\mathbf{v},\mathbf{v'}},$$

we have

$$(x_{\mathbf{iu}}^{\mathbf{r}*} \otimes \mathrm{id})(\Delta x) = (\mathrm{coefficient}) \cdot x_{\mathbf{uj}}^{\mathbf{r}} + \sum_{\bullet \neq j} (\mathrm{coefficient}) \cdot x_{\mathbf{u\bullet}}^{r}.$$

ranging over the n_1 **u** will thus produce n_1 linearly-independent elements:

dim $(x \triangleleft C^*) \ge n_1$ for the dual action [9, §2.2] $x \triangleleft f := (f \otimes id)(\Delta(x)), \forall f \in C^*.$

The matrix algebra C^* thus has a cyclic module of dimension $\geq n_1$, so must itself have dimension $\geq n_1^2$.

Corollary 1.19 Under the hypotheses of Theorem 1.10, a simple $H(\mathbf{C})$ -comodule V whose k^{th} dual V^{k*} ($k \in \mathbb{Z}_{\geq 0}$) has a simple subquotient whose $\langle \mathbb{Z}_{\geq d} \rangle$ -valued rank contains the letter r + k has dimension

$$\geq \min\left\{\sqrt{\dim C} \mid matrix \ subcoalgebra \ C \leq C^r\right\}.$$

Proof Immediate from Theorem 1.18, given that the $\langle \mathbb{Z}_{\geq d} \rangle$ -valued rank of such a comodule must contain the letter r.

Corollary 1.20 Let $\mathbf{C} = (C^r, \theta^r)$ be a triangular skew coalgebra chain as in Theorem 1.10. If

 $\lim_{r \searrow -\infty} \min \dim (simple \ subcoalgebra \ of \ C^r) = \infty$

(so that in particular the chain is by necessity bi-infinite) then the only Hopf morphisms $H \to H(\mathbf{C})$ for H with bijective antipode are trivial.

Proof Let $H \xrightarrow{\varphi} H(\mathbf{C})$ be a *non*-trivial morphism for bijective-antipode H. Some simple Hcomodule W, regarded as an $H(\mathbf{C})$ -comodule via φ , has a simple subquotient V of non-trivial rank $\mathbf{r} \in \langle \mathbb{Z} \rangle$. The antipode of H being bijective, W has iterated right duals

$$W^{-k*}$$
, $\dim W^{-k*} = \dim W$, $\forall k \in \mathbb{Z}_{\geq 0}$.

Now, if $r_0 \in \mathbb{Z}$ is a letter featuring in \mathbf{r} , $r_0 - k$ must feature in the rank of some simple subquotient of W^{-k*} . As $k \to \infty$ the dimensions of such simples increase indefinitely by Corollary 1.19, contradicting the fact that they can arise as such subquotients.

Example 1.21 Instances of Corollary 1.20 are easily produced. Take, say

- $C^r = M_2^{\oplus \mathbb{Z}}$ and $\theta^r = \text{id for all } r \ge -1;$
- whereas for r < 0 C^r is a sum of countably infinitely many copies of $M_{2^{-r}}$;
- and each θ^r , r < -1 surjects one $2^{-r} \times 2^{-r}$ onto a sum of two diagonal blocks half the size:

$$\begin{pmatrix} * & * \\ * & * \end{pmatrix} \longrightarrow \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}$$

For negative r the simple C^r -comodules are 2^{-r} -dimensional, so that we do fit the framework of Corollary 1.20.

Remark 1.22 Example 1.21 is one particular straightforward way to meet the requirements of Corollary 1.20; it should be clear that the construction admits of much variation. Even restricting attention to cosemisimple coalgebras (as in Example 1.21), one can parametrize the surjections $C^r \rightarrow C^{r+1}$ by the *Bratteli diagrams* [10, Chapter 2] pervasive in the study of *AF* (*approximately finite* [10, Chapter 2, p.11]) operator algebras.

References

- Jiří Adámek, Horst Herrlich, and George E. Strecker. Abstract and concrete categories: the joy of cats. *Repr. Theory Appl. Categ.*, 2006(17):1–507, 2006. 2, 4
- [2] Jiří Adámek and Jiří Rosický. Locally presentable and accessible categories, volume 189 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 1994. 3, 4, 8
- [3] George M. Bergman. The diamond lemma for ring theory. Adv. Math., 29:178–218, 1977. 7, 8
- [4] Garrett Birkhoff. Lattice theory. Corr. repr. of the 1967 3rd ed, volume 25 of Colloq. Publ., Am. Math. Soc. American Mathematical Society (AMS), Providence, RI, 1979. 8
- [5] Francis Borceux. Handbook of categorical algebra. Volume 1: Basic category theory, volume 50 of Encycl. Math. Appl. Cambridge: Cambridge Univ. Press, 1994.
- [6] Alexandru Chirvăsitu. On epimorphisms and monomorphisms of Hopf algebras. J. Algebra, 323(5):1593–1606, 2010.
- [7] Alexandru Chirvăsitu. Subcoalgebras and endomorphisms of free Hopf algebras. J. Pure Appl. Algebra, 215(2):101–107, 2011. 6, 11
- [8] Alexandru Chirvăsitu. Grothendieck rings of universal quantum groups. J. Algebra, 349(1):80– 97, 2012. 6, 7, 8
- [9] Sorin Dăscălescu, Constantin Năstăsescu, and Şerban Raianu. Hopf algebras, volume 235 of Monographs and Textbooks in Pure and Applied Mathematics. Marcel Dekker, Inc., New York, 2001. An introduction. 9, 12

- [10] Edward G. Effros. Dimensions and C*-algebras. (Expository lectures from the CBMS Regional Conference held at Oakland University, June 25-29, 1979), volume 46 of Reg. Conf. Ser. Math. American Mathematical Society (AMS), Providence, RI, 1981. 13
- [11] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik. *Tensor categories*, volume 205 of *Mathe-matical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015. 1, 2
- [12] Peter Freyd. Abelian categories. Repr. Theory Appl. Categ., 2003(3):xxiii + 1–164, 2003. 9
- [13] David Gries and Fred B. Schneider. A logical approach to discrete math. New York, NY: Springer-Verlag, 1993. 8
- [14] S. Montgomery. Hopf algebras and their actions on rings, volume 82 of CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1993. 5, 8, 9, 10
- [15] Warren D. Nichols. Quotients of Hopf algebras. Comm. Algebra, 6(17):1789–1800, 1978. 4, 6
- [16] B. Pareigis. Quantum groups and noncommutative geometry, 2002. available at https://www.mathematik.uni-muenchen.de/~pareigis/Vorlesungen/02SS/QGandNCG.pdf (accessed 2024-06-14). 1, 9
- [17] Richard S. Pierce. Associative algebras, volume 88 of Grad. Texts Math. Springer, Cham, 1982.
 5
- [18] Hans-E. Porst. The formal theory of Hopf algebras. Part II: The case of Hopf algebras. Quaest. Math., 38(5):683-708, 2015. 3
- [19] David E. Radford. Hopf algebras, volume 49 of Series on Knots and Everything. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2012. 3
- [20] Peter Schauenburg. Tannaka duality for arbitrary Hopf algebras, volume 66 of Algebra-Ber. München: R. Fischer, 1992. 9, 11
- [21] Peter Schauenburg. Faithful flatness over Hopf subalgebras: counterexamples. In Interactions between ring theory and representations of algebras (Murcia), volume 210 of Lecture Notes in Pure and Appl. Math., pages 331–344. Dekker, New York, 2000. 2, 4, 6, 10
- [22] Moss E. Sweedler. Hopf algebras. Mathematics Lecture Note Series. W. A. Benjamin, Inc., New York, 1969. 1
- [23] Mitsuhiro Takeuchi. Free Hopf algebras generated by coalgebras. J. Math. Soc. Japan, 23:561– 582, 1971. 2, 3, 4, 5, 8

DEPARTMENT OF MATHEMATICS, UNIVERSITY AT BUFFALO BUFFALO, NY 14260-2900, USA *E-mail address*: achirvas@buffalo.edu