

Exact spectral function and nonequilibrium dynamics of the strongly interacting Hubbard model

Ovidiu I. Păţu,¹ Andreas Klümper,² and Angela Foerster³

¹*Institute for Space Sciences, Bucharest-Măgurele, R 077125, Romania*

²*Fakultät für Mathematik und Naturwissenschaften,*

Bergische Universität Wuppertal, 42097 Wuppertal, Germany

³*Instituto de Física da UFRGS, Av. Bento Gonçalves 9500, Porto Alegre, RS, Brazil*

Analytical results on the correlation functions of strongly correlated many-body systems are rare in the literature and their importance cannot be overstated. We present determinant representations for the space-, time-, and temperature-dependent correlation functions of the strongly interacting one-dimensional Hubbard model in the presence of an external trapping potential. These representations are exact and valid in both equilibrium and nonequilibrium scenarios like the ones initiated by a sudden change of the confinement potential. In addition, they can be implemented numerically very easily significantly outperforming other numerical approaches. As applications of our results we investigate the single particle spectral functions of systems with harmonic trapping and show that dynamical quasicondensation occurs for both fermionic and bosonic spin-1/2 systems released from a Mott insulator state.

Introduction.— One of the most challenging and still unresolved puzzles in physics is the computation of correlation functions in strongly interacting many-body systems. While significant progress has been achieved in understanding integrable single-component systems in one dimension (1D) over the last decades, both at zero [1–3] and finite temperature [3–5], the same level of advancement has not been attained for systems with internal degrees of freedom. The complexity primarily arises from the intricate nature of wavefunctions due to the inclusion of the spin sector, significantly complicating the dynamics. Moreover, multicomponent systems at low temperatures exhibit diverse additional regimes beyond the conventional Luttinger liquid (LL) phase [6, 7]. These additional phases, such as the spin-incoherent Luttinger liquid (SILL) phase [8–14], are absent in their single-component counterparts and have only recently attracted experimental attention [15].

In this Letter, we report on the derivation of determinant representations for the correlation functions of the impenetrable (double occupancy is excluded) 1D Hubbard model in the presence of an external potential. Our results are both exact and remarkably versatile, applicable across a wide range of conditions, including any temperature, particle statistics, and external potential, whether in equilibrium or non-equilibrium scenarios such as those induced by a sudden change in the potential. Furthermore, our determinant representations are extremely easy to implement numerically, allowing for the investigation of a larger number of particles and lattice sites beyond the reach of other numerical techniques like exact diagonalization or time-dependent Density Matrix Renormalization Group. The derivation is based on the factorization of the wavefunctions in the strongly interacting regime [16–25] and the connection with a dual system of spinless fermions subjected to the same trapping potential or quench [26].

As a first application of our results we have investigated the effect of a harmonic trapping potential on the single particle spectral functions of balanced fermionic and bosonic systems at zero temperature. At zero temperature our results describe the SILL regime and in the homogeneous case (no potential) we can highlight the differences between the SILL and LL fermionic spectral functions previously investigated in [27–31]. As a consequence of non-linear LL theory [32, 33] the LL spectral functions exhibit singularities along the dispersion lines of the charge and spin excitations [34–37] feature which is not present in the SILL spectral function due to the spin-incoherence of the system which smears the singularities. In the fermionic case the SILL spectral function does not exhibit the so-called “shadow band” [27] and the dispersion of the charge part is different compared with the LL regime. As a prototypical nonequilibrium scenario we study the release of a system from a Mott insulator (MI) state revealing a dynamical quasicondensation phenomenon [38, 39] characterized by the development of peaks in the momentum distribution at $\pm\pi/(2a_0)$ (a_0 is the lattice constant) as a result of emergence of correlations with the same exponents as those characterizing the spin-incoherent system at zero temperature.

The Hubbard model.— In second quantization the Hamiltonian of the Hubbard model subjected to an external field is $H = H_K + H_U + H_B + H_E$ [40, 41]

$$H_K = -t \sum_{z=1}^{L-1} \sum_{\sigma=\{\uparrow,\downarrow\}} \left(a_{z,\sigma}^\dagger a_{z+1,\sigma} + a_{z+1,\sigma}^\dagger a_{z,\sigma} \right), \quad (1a)$$

$$H_U = U \sum_{z=1}^L n_{z,\uparrow} n_{z,\downarrow}, \quad H_B = B \sum_{z=1}^L (n_{z,\uparrow} - n_{z,\downarrow}) \quad (1b)$$

$$H_E = \sum_{z=1}^L (V(z, t) - \mu) (n_{z,\uparrow} + n_{z,\downarrow}), \quad (1c)$$

where t is the hopping strength, L the number of lat-

tice sites, $U > 0$ the repulsive Coulomb interaction, B the magnetic field, μ the chemical potential and $n_{z,\sigma} = a_{z,\sigma}^\dagger a_{z,\sigma}$. In the following we will be interested in the case of infinite repulsion, $U = \infty$, which means that two particles cannot occupy the same lattice site. We consider the general case of arbitrary statistics with $a_{z,\sigma}^\dagger, a_{z,\sigma}$ anyonic fields [42–44] satisfying $a_{z,\sigma} a_{m,\sigma'}^\dagger = f(z, m) a_{m,\sigma'}^\dagger a_{z,\sigma} + \delta_{\sigma,\sigma'} \delta_{z,m}$, $a_{z,\sigma} a_{m,\sigma'} = \bar{f}(z, m) a_{m,\sigma'} a_{z,\sigma}$, with $f(z, m) = (-1)^{i\pi\kappa \text{sign}(z-m)}$ (bar denotes complex conjugation) where $\kappa \in [0, 1]$ is the statistics parameter and $\text{sign}(k) = |k|/k$, $\text{sign}(0) = 0$. As we vary the statistics parameter κ the commutation relations interpolate continuously between the anticommutation relations for fermions ($\kappa = 0$) and the commutation relations for bosons ($\kappa = 1$). We should remark that the statistics parameter can also be chosen to be $\kappa \in [-1, 0]$ with $\kappa = -1$ ($\kappa = 0$) describing bosons (fermions). At coinciding points the anyonic fields satisfy $(a_{z,\sigma}^\dagger)^2 = (a_{z,\sigma})^2 = 0$ (hard-core condition). The usual Fermi-Hubbard (Bose-Hubbard) model is obtained for $\kappa = 0$ ($\kappa = 1$). For the external potentials we will consider either static trapping potentials like $V(z) = V_0|z - (L+1)/2|^\alpha$ with $\alpha = 1, 2, \dots$ or time dependent potentials of the type $V(z, t=0) = V_I(z)$ and $V(z, t > 0) = V_F(z)$ which correspond to quantum quenches. In the case of quantum quenches we will denote the initial (final) Hamiltonian by H_I (H_F). In equilibrium situations we have $H_I = H_F$. We use units of $\hbar = k_B = 1$ with k_B the Boltzmann constant.

An important role in our results is played by a dual system of spinless fermions subjected to the same external potential like the Hubbard model with the single particle Hamiltonian

$$H^{SP} = -t \sum_{z=1}^{L-1} (|z+1\rangle\langle z| + |z\rangle\langle z+1|) + \sum_{z=1}^L V(z, t) |z\rangle\langle z|,$$

where $|z\rangle = f_z^\dagger |0\rangle$ and f_z^\dagger, f_m are fermionic creation

and annihilation operators satisfying $\{f_z, f_m^\dagger\} = \delta_{z,m}$, $\{f_z, f_m\} = 0$. Similar with the case of the Hubbard model in the case of quenches we will denote the initial (final) single particle Hamiltonian by H_I^{SP} (H_F^{SP}). In equilibrium situations at $t = 0$ the single particle orbitals satisfy $H_I^{SP} \phi_k(z) = \varepsilon(k) \phi_k(z)$ and their time evolution is given by $\phi_k(z, t) = e^{-i\varepsilon(k)t} \phi_k(z)$. In nonequilibrium $\phi_k(z, t)$ is the unique time-dependent solution of the Schrödinger equation $i\hbar \partial \phi_k(z, t) / \partial t = H_F^{SP} \phi_k(z, t)$ with initial boundary condition $\phi_k(z, 0) = \phi_k(z)$ [$\phi_k(z)$ is an eigenfunction of H_I^{SP}] where H_F^{SP} is the final single particle Hamiltonian.

Correlators.— We are interested in deriving determinant representations for the space-, time-, and temperature-dependent correlation functions of the Hubbard model in both equilibrium and nonequilibrium scenarios. The system is initially prepared in a grandcanonical thermal state of the initial Hamiltonian H_I described by the chemical potential μ , magnetic field B and temperature T . The correlators of interest are

$$g_\sigma^{(-)}(x, t; y, t') = \text{Tr} \left[e^{-H_I/T} a_{x,\sigma}^\dagger(t) a_{y,\sigma}(t') \right] / Z, \quad (2a)$$

$$g_\sigma^{(+)}(x, t; y, t') = \text{Tr} \left[e^{-H_I/T} a_{x,\sigma}(t) a_{y,\sigma}^\dagger(t') \right] / Z, \quad (2b)$$

where $Z(\mu, B, T)$ is the partition function of the initial thermal state and the time evolved operators are $a_{x,\sigma}(t) = e^{iH_F t} a_{x,\sigma} e^{-iH_F t}$, $a_{x,\sigma}^\dagger(t) = e^{iH_F t} a_{x,\sigma}^\dagger e^{-iH_F t}$. Due to the symmetry $g_\downarrow^{(\pm)}(x, t; y, t'|B) = g_\uparrow^{(\pm)}(x, t; y, t'|-B)$ it will be sufficient to study only one type of correlators. The densities and momentum distributions are given by $\rho_\sigma(x, t) = g_\sigma^{(-)}(x, t; x, t)$ and $n_\sigma(k, t) = \sum_x \sum_y e^{-ik(x-y)} g_\sigma^{(-)}(x, t; y, t)$.

Determinant representations.— One of the main results of this Letter is the determinant representations for the correlation functions. Introducing the function $F(\gamma, \eta) = 1 + \sum_{p=1}^{\infty} \gamma^{-p} (e^{i\eta p} + e^{-i\eta p})$ where $\gamma = (1 + e^{2B/T})$ the correlators (2a) and (2b) have the representations

$$g_\uparrow^{(-)}(x, t; y, t') = \frac{e^{-i(t-t')\mu_\uparrow}}{2\pi} \int_{-\pi}^{\pi} F(\gamma, \eta) \left[\det \left(1 + \gamma V^{(T,-)}(\eta) + R^{(T,-)} \right) - \det \left(1 + \gamma V^{(T,-)}(\eta) \right) \right] d\eta, \quad (3)$$

$$g_\uparrow^{(+)}(x, t; y, t') = \frac{e^{i(t-t')\mu_\uparrow}}{2\pi} \int_{-\pi}^{\pi} F(\gamma, \eta) \left[\det \left(1 + \gamma V^{(T,+)}(\eta) - \gamma R^{(T,+)}(\eta) \right) + (g-1) \det \left(1 + \gamma V^{(T,+)}(\eta) \right) \right] d\eta, \quad (4)$$

where the chemical potentials of the spin-up (-down) particles are $\mu_\uparrow = \mu - B$ ($\mu_\downarrow = \mu + B$) and $[V^{(T,\pm)}(\eta)]_{ab} = \sqrt{\vartheta(a)} \left(U_{ab}^{(\pm)}(\eta) - \delta_{a,b} \right) \sqrt{\vartheta(b)}$ and

$[R^{(T,\pm)}]_{ab} = \sqrt{\vartheta(a)} R_{ab}^{(\pm)} \sqrt{\vartheta(b)}$ are infinite matrices. $\vartheta(k) = e^{-B/T} / (2 \cosh(B/T) + e^{(\varepsilon(k)-\mu)/T})$ plays the role of the Fermi function for spin up particles. In Eq. (4) $g \equiv g(x, t; y, t') = \sum_{k=1}^{\infty} \phi_k(x, t) \bar{\phi}_k(y, t')$ and in terms of

the function $(\phi_k(z, t))$ are the time evolved orbitals of the dual system of spinless fermions)

$$u(k, q|\eta, x, t) = \delta_{k,q} - \left[1 - e^{i(\pi\kappa - \eta)}\right] \sum_{z=x}^{\infty} \phi_k(z, t) \bar{\phi}_q(z, t),$$

the elements of the relevant matrices are given by $(a, b = 1, 2, \dots)$: $U_{ab}^{(-)}(x, t; y, t'|\eta) = \sum_{q=1}^{\infty} \bar{u}(a, q|\eta, x, t) u(b, q|\eta, y, t')$, $U_{ab}^{(+)}(x, t; y, t'|\eta) = \sum_{k=1}^{\infty} u(k, b|\eta, x, t) \bar{u}(k, a|\eta, y, t')$, and $R_{ab}^{(-)}(x, t; y, t') = \bar{\phi}_a(x, t) \phi_b(y, t')$, $R_{ab}^{(+)}(x, t; y, t'|\eta) = \bar{e}_a(x, t; y, t'|\eta) e_b(x, t; y, t'|\eta)$ with $e_a(x, t; y, t'|\eta) = \sum_{k=1}^{\infty} u(k, a|\eta, x, t) \bar{\phi}_k(y, t')$, $\bar{e}_a(x, t; y, t'|\eta) = \sum_{k=1}^{\infty} \bar{u}(k, a|\eta, y, t') \phi_k(x, t)$. The representations (3), (4) are exact and in practice the numerical computation of the correlators require only the computation of sums, products and determinants of finite size matrices (at zero temperature the matrices have the same dimensions as the number of particles in the system and at finite temperature one can truncate at level N_T when $\theta(N_T)$ is small enough) which can be performed with extreme efficiency, greatly surpassing other numerical schemes like exact diagonalization or the time-dependent Density Matrix Renormalization Group. For fermionic homogeneous systems with periodic boundary conditions determinant representations were derived in [17, 45].

Equilibrium spectral function.— At equilibrium ($H_I = H_F$) our results can be used to investigate the spectral function of the Hubbard model in the presence of a trapping potential. One of the most important quantities in many-body physics the spectral function encodes information about the accessible energy states and their distribution in momentum space and can be computed as the imaginary part of the Fourier transform of the retarded Green's function $A(k, \omega) = -\frac{1}{\pi} \text{Im} G^R(k, \omega)$, with $G^R(k, \omega) = \int e^{i\omega t} dt \sum_x \sum_y e^{-ik(x-y)} G^R(x, t; y, 0)$, where the total retarded Green's function is defined as $G^R(x, t; y, t') = -i\theta(t - t') \sum_{\sigma \in \{\uparrow, \downarrow\}} \left[g_{\sigma}^{(+)}(x, t; y, t') + g_{\sigma}^{(-)}(y, t'; x, t) \right]$. We will focus on the most interesting case: the balanced system ($B = 0$) at zero temperature and investigate the dependence of the spectral function on the strength of the harmonic trapping. It is important to note that using our representations at zero temperature we investigate the spectral function SILL regime [8–14] due to the fact that we take first the limit $U \rightarrow \infty$ and then $T \rightarrow 0$ (for the thermodynamics and quantum criticality in the same regime see [46–49]). The spectral function in the LL regime [6, 7] which is obtained by taking the limit $T \rightarrow 0$ first and then $U \rightarrow \infty$ has been investigated in [27–31].

At zero magnetic field and zero temperature the parameter $\gamma = 1 + e^{2B/T} = 2$ and $F(\gamma = 2, \eta) \equiv F_0(\eta) = 3/(5 - 4 \cos \eta)$. Comparing our results with the ones obtained for single component anyons on the lattice [50, 51]

we obtain the following identities ($g_{LL}^{(\pm)}$ are the equivalent correlation functions of single component anyons)

$$g_{\uparrow}^{(-)}(x, t; y, t'|\kappa) = \frac{1}{4\pi} \int_{-\pi}^{\pi} F_0(\eta) g_{LL}^{(-)}(x, t; y, t'|\kappa - \frac{\eta}{\pi}) d\eta,$$

$$g_{\uparrow}^{(+)}(x, t; y, t'|\kappa) = \frac{1}{2\pi} \int_{-\pi}^{\pi} F_0(\eta) g_{LL}^{(+)}(x, t; y, t'|\kappa - \frac{\eta}{\pi}) d\eta,$$

which reveal the fact that the correlation functions of the Hubbard model with statistical parameter κ can be expressed as integrals over η of single component anyons of statistics parameter $\kappa - \eta/\pi$ with weight $\frac{1}{2\pi} \frac{3}{5 - 4 \cos \eta}$. In the fermionic case similar identities were derived using a different method in [52]. The previous relations also show that the spectral function of the Hubbard model in the SILL regime is closely related to the spectral function of single component anyons. We remind the reader that the singular lines of the spectral function for single component anyons are given by [50, 51, 53]

$$\varepsilon_I(k) = -2t \cos(k + k_F \kappa) - \mu, \quad (5a)$$

$$\varepsilon_{II}(k) = 2t \cos(k + k_F \kappa - 2k_F) + \mu, \quad (5b)$$

$$\varepsilon_{III}(k) = 4t \sin[(k + k_F \kappa - k_F)/2], \quad (5c)$$

where $k_F = \pi N/L$ is the Fermi vector with N the number of particles. In the case of single component anyons decreasing the statistics parameter from $\kappa = 1$ (bosons) the spectral weight from $\varepsilon_{II}(k)$ and $\varepsilon_{III}(k)$ also decreases being transferred to $\varepsilon_I(k)$ which becomes the only singular line at $\kappa = 0$ as it is expected for a free fermionic system.

In Fig. 1 we present results for the spectral function of the Hubbard model (fermionic and bosonic) in the presence of a harmonic potential $V(z) = V_0 |z - (L+1)/2|^2$ with V_0 varying from zero (homogeneous system) to $V_0 = 0.005$ a value for which the system is in the Mott insulator phase (the middle of the trap contains a region in which the density is 1). While we are not aware of any results reported in the literature for the spectral function of the Hubbard model subjected to an external trapping potential in the homogeneous case the spectral function for the Fermi-Hubbard model in the LL regime has been investigated in [27–31]. This gives us the opportunity to highlight the differences between the spectral functions in the two regimes. Both spectral functions present Van Hove singularities at $\pm 2t$ (in the vicinities of $k = \pm\pi$ and $k = 0$) and an additional singularity at $\omega = \mu$ (around $k = 0$). However, the SILL spectral function does not exhibit the so-called “shadow band” [27] and the dispersion of the charge part for $A(k, \omega)$ is given by Eq. (5a) compared with $\varepsilon_{LL}(k) = -2t \cos(|k| + k_F^L)$, ($k_F^L = k_F/2$) for the LL spectral function. Another major difference is that while in the LL regime the spectral function has a power-law behaviour [34–37] $A(k, \omega) \sim |\omega - \varepsilon_j(k)|^{-\mu_j}$ near each singular line $\varepsilon_j(k)$ with exponent μ_j that is momentum dependent this feature is absent in the SILL




FIG. 1. Dependence of the spectral function for the zero temperature balanced Hubbard model on the strength of the potential V_0 ($V(z) = V_0|z - (L+1)/2|^2$). Here $N = 40$, $L = 120$ and $t = 1$. The first row presents results for the fermionic system ($\kappa = 0$) and the second row for the bosonic system ($\kappa = 1$). In the first column the continuous blue, red and yellow lines highlight the $\varepsilon_I(k)$, $\varepsilon_{II}(k)$ and $\varepsilon_{III}(k)$ singular lines [see Eqs. (5a), (5b), (5c)] while the corresponding dashed lines describe $-\varepsilon_I(k)$, $-\varepsilon_{II}(k)$ and $-\varepsilon_{III}(k)$. The dashed white lines are the Van Hove singularities present at $\omega = \pm 2t$ while the dotted white line marks the singularity line present at $\omega = \mu$.

phase. This is due to the fact that $A(k, \omega)$ in the SILL phase can be expressed as an integral over the spectral functions of single component anyonic systems which smoothens the “singular lines” in both fermionic and bosonic cases [Eqs. (5a), (5b)]. The presence of the trapping potential significantly changes the characteristics of the spectral function. For small values of the curvature ($V_0 = 0.0005$) the spectral function presents similar features like in the homogeneous case but with an increased k_F coupled with a broadening of the pseudo-singular lines [Eqs. (5a), (5b) and (5c)]. However, for stronger potentials the homogeneous picture washes out with the majority of the spectral weight migrating along the $\omega = \mu$ singular line and the origin.

Dynamical quasicondensation. — Our determinant representations for the correlators (3) and (4) are also valid in experimentally accessible nonequilibrium scenarios which are initiated by a sudden change of the trapping potential. One such scenario of considerable interest is represented by the free expansion of a Mott insulator (MI) state located initially in the center of the trap (for other nonequilibrium scenarios see [54–58]). In Fig. 2 we present the time evolution of the densities and momentum distributions for a system of $N = 50$ particles on

a lattice with $L = 250$ sites after removal of the harmonic trapping potential with the system initially prepared in a MI state. While the initial quasimomentum distribution is almost flat (for an infinite lattice the distribution is flat) and there are no off-diagonal correlations $\langle a_{j,\sigma}^\dagger a_{j+r,\sigma} \rangle \sim 0$, after release from the trap the momentum distributions develop peaks at $\pm\pi/(2a_0)$ which are more pronounced in the bosonic case [see Fig. 2c] for the bosonic case and Fig. 2e) for the fermionic case]. This rather counterintuitive phenomenon is the multi-component counterpart of the dynamical quasicondensation of hard-core bosons which was theoretically predicted in [38] (see also [59–62]) and validated experimentally in [39]. In the case of single component bosons it was argued [39] that these singularities are due to emerging power-law correlations $\langle a_j^\dagger(t) a_{j+r}(t) \rangle = \mathcal{A}(r) e^{i\Phi(r)}$, $\mathcal{A}(r) \sim r^{-1/2}$, $\Phi(r) = \pm\frac{\pi}{2}r$, with the plus (minus) sign for the right (left) expanding cloud. It is interesting to note that even though we are in a nonequilibrium situation the power law decay of the absolute value is given by the same exponent as in the groundstate (1/2) but with an alternating phase pattern $\pm\pi/2$ between neighbouring sites producing peaks at finite quasimomenta (for each half system). Guided by the single component case in




FIG. 2. Expansion of the zero temperature balanced Hubbard model from a Mott insulator initial state. Here $N = 50$, $L = 250$ and $\tau = \hbar/t$. a) Time evolution of the density $\rho_{\downarrow}(j, t) = \rho_{\uparrow}(j, t)$, which is independent of statistics, after the trapping potential is removed. b) Initial momentum densities $n_{\downarrow}(k, 0) = n_{\uparrow}(k, 0)$ for fermions (green dashed line) and bosons (blue continuous line). Inset: Density profile $\rho_{\downarrow}(j, 0) = \rho_{\uparrow}(j, 0)$ at $t = 0$ which shows that the system is in the MI state in the middle of the trap. c) Time evolution after expansion of the momentum distribution $n_{\downarrow}(k, t)$ for a bosonic system $\kappa = 1$ from which we can see clearly the peaks at $\pm\pi/(2a_0)$. d) $|\langle a_{j,\downarrow}^{\dagger} a_{j+r,\uparrow} \rangle|$ for $j = 26$ at $t = 0$ (black markers, the black continuous line is a guide to the eye) and $t = 0.4N\tau$ (blue markers) with the asymptotic value (continuous blue line) $\mathcal{A}(r)$ given by Eq. (6) with $a = 0.20080$, $k_F^* = 1.48750$. Inset: The phase pattern of the bosonic correlator (blue markers), the blue continuous line is $\frac{\pi}{2}r$. e) Time evolution after expansion of the momentum distribution $n_{\downarrow}(k, t)$ for a fermionic system $\kappa = 0$. f) $|\langle a_{j,\downarrow}^{\dagger} a_{j+r,\uparrow} \rangle|$ for $j = 26$ at $t = 0$ (black markers, the black continuous line is a guide to the eye) and $t = 0.4N\tau$ (blue markers) and the asymptotic value (continuous blue line) $\mathcal{A}(r)$ given by Eq. (7) with $a = 0.22646$, $k_F^* = 1.48228$, $\phi = 0.14982$. Inset: The phase pattern of the fermionic correlator (green markers), the green continuous lines are $\frac{\pi}{2}r$ and $\frac{\pi}{2}r + \pi$.

which the emerging correlation functions have the same asymptotic behavior like in the groundstate with an alternating phase we have discovered that the same phenomenon remains valid in the two-component case. In the bosonic case we have $\langle a_{j,\downarrow}^{\dagger}(t)a_{j+r,\uparrow}(t) \rangle = \mathcal{A}(r)e^{i\Phi(r)}$

with the asymptotic behavior [see Fig. 2d)]

$$\mathcal{A}(r) \sim a e^{-\nu k_F^* r} r^{-\frac{1}{2} + \frac{1}{2}\nu^2}, \quad \Phi(r) = \pm \frac{\pi}{2}r, \quad (6)$$

where a and k_F^* are free parameters [k_F^* is very likely related to the density of the particle cloud at (j, t)] and $\nu = \frac{\ln 2}{\pi}$ which has the same form as the large distance asymptotics of two-component bosons in the SILL regime [63, 64]. It is important to note that because we are in the spin-incoherent regime the asymptotics are exponentially decreasing even at zero temperature with an exponent depending on the number of components which is a general feature of incoherent systems [14]. However, the power exponent is close to the one for single component bosons ($\frac{1}{2}\nu^2 \sim 0.024$) and even though the momentum distribution does not present singularities at $\pm\pi/(2a_0)$ we still have a sizeable number of particles with momenta concentrated around $\pm\pi/(2a_0)$. In the fermionic case the asymptotics are [65] [see Fig. 2f)]

$$\mathcal{A}(r) \sim a e^{-\nu k_F^* r} r^{-1 + \frac{1}{2}\nu^2} |\sin(k_F^* r - \nu \ln r - \phi)|, \quad (7)$$

with a , k_F^* and ϕ free parameters again mimicking the results for two-component fermions at zero temperature in the spin-incoherent regime. In this case the phase is $\Phi(r) = \pm \frac{\pi}{2}r$ plus a contribution of π whenever the sin factor in (7) is negative.

Conclusions.— We have introduced versatile determinant representations for the correlation functions of the strongly interacting Hubbard model, which facilitated the numerical investigation of the spectral function in the presence of a confinement potential and the dynamics of the system after release from the trap. Our results are highly general, easy to implement numerically, and can be used to study the static properties in different trapping geometries, the nonequilibrium dynamics in various scenarios (periodic modulation of the trapping frequency, quenches of the confining potential, etc.), the presence or absence of dynamical localization in periodically kicked systems, and the transport properties at zero and finite temperature. We should also point out that our equal-time representations can be expressed in a Lenard form [26, 66], from which rigorous analytic results regarding the asymptotics can be derived.

- [1] N. Kitanine, K.K. Kozłowski, J.M. Maillet, N.A. Slavnov, and V. Terras, *Algebraic Bethe ansatz approach to the asymptotic behavior of correlation functions*, J. Stat. Mech. P04003 (2009).
- [2] N. Kitanine, K.K. Kozłowski, J.M. Maillet, N.A. Slavnov, and V. Terras, *Form factor approach to dynamical correlation functions in critical models*, J. Stat. Mech. P09001 (2012).
- [3] V.E. Korepin, N.M. Bogoliubov, and A.G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions* (Cambridge University Press, Cambridge, UK, 1993).

- [4] F. Göhmann, A Klümper and A. Seel, *Integral representations for correlation functions of the XXZ chain at finite temperature*, J. Phys. A **37**, 7625 (2004).
- [5] F. Göhmann, M. Karbach, A. Klümper, K.K. Kozłowski, and J. Suzuki, *Thermal form-factor approach to dynamical correlation functions of integrable lattice models*, J. Stat. Mech. 113106 (2017).
- [6] F.D.M. Haldane, *Effective Harmonic-Fluid Approach to Low-Energy Properties of One-Dimensional Quantum Fluids*, Phys. Rev. Lett. **47**, 1840 (1981).
- [7] T. Giamarchi, *Quantum Physics in One Dimension* (Oxford University Press, Oxford, UK, 2003).
- [8] A. Berkovich A and J.H. Lowenstein, *Correlation function of the one-dimensional Fermi gas in the infinite-coupling limit (repulsive case)*, Nucl. Phys. B **285**, 70 (1987).
- [9] A. Berkovich, *Temperature and magnetic field-dependent correlators of the exactly integrable (1+1)-dimensional gas of impenetrable fermions*, J. Phys. A **24** 1543 (1991).
- [10] V.V. Cheianov and M.B. Zvonarev, *Nonunitary Spin-Charge Separation in a One-Dimensional Fermion Gas*, Phys. Rev. Lett. **92**, 176401 (2004).
- [11] V.V. Cheianov and M.B. Zvonarev, *Zero temperature correlation functions for the impenetrable fermion gas*, J. Phys. A **37**, 2261 (2004).
- [12] K. A. Matveev, *Conductance of a Quantum Wire in the Wigner-Crystal Regime*, Phys. Rev. Lett. **92**, 106801 (2004).
- [13] G.A. Fiete and L. Balents, *Green's Function for Magnetically Incoherent Interacting Electrons in One Dimension*, Phys. Rev. Lett. **93**, 226401 (2004).
- [14] G.A. Fiete, *Colloquium: The spin-incoherent Luttinger liquid*, Rev. Mod. Phys. **79**, 801 (2007).
- [15] D. Cavazos-Cavazos, R. Senaratne, A. Kaffle, and R.G. Hulet, *Thermal disruption of a Luttinger liquid*, Nature Communications 14, 3154 (2023).
- [16] M. Ogata and H. Shiba, *Bethe-ansatz wave function, momentum distribution, and spin correlation in the one-dimensional strongly correlated Hubbard model*, Phys. Rev. B **41**, 2326 (1990).
- [17] A.G. Izergin and A.G. Pronko, *Temperature correlators in the two-component one-dimensional gas*, Nucl. Phys. B **520**, 594 (1998).
- [18] F. Deuretzbacher, K. Fredenhagen, D. Becker, K. Bongs, K. Sengstock, and D. Pfannkuche, *Exact Solution of Strongly Interacting Quasi-One-Dimensional Spinor Bose Gases* Phys. Rev. Lett. **100**, 160405 (2008).
- [19] L. Guan, S. Chen, Y. Wang, and Z. Q. Ma, *Exact Solution for Infinitely Strongly Interacting Fermi Gases in Tight Waveguides*, Phys. Rev. Lett. **102**, 160402 (2009).
- [20] A.G. Volosniev, D.V. Fedorov, A.S. Jensen, M. Valiente, and N.T. Zinner, *Strongly interacting confined quantum systems in one dimension*, Nat. Commun. **5**, 5300 (2014).
- [21] J. Levinsen, P. Massignan, G.M. Bruun, and M.M. Parish, *Strong-coupling ansatz for the one-dimensional Fermi gas in a harmonic potential*, Sci. Adv. **1**, e1500197 (2015).
- [22] L. Yang and X. Cui, *Effective spin-chain model for strongly interacting one-dimensional atomic gases with an arbitrary spin*, Phys. Rev. A **93**, 013617 (2016).
- [23] F. Deuretzbacher, D. Becker, J. Bjerlin, S.M. Reimann, and L. Santos, *Spin-chain model for strongly interacting one-dimensional Bose-Fermi mixtures*, Phys. Rev. A **95**, 043630 (2017).
- [24] L. Yang, S.S. Alam, and H. Pu, *Generalized Bose-Fermi mapping and strong coupling ansatz wavefunction for one dimensional strongly interacting spinor quantum gases*, J. Phys. A **55**, 464005 (2022).
- [25] A. Minguzzi and P. Vignolo, *Strongly interacting trapped one-dimensional quantum gases: Exact solution*, AVS Quantum Sci. **4**, 027102 (2022).
- [26] See the Supplemental Material for a sketch of the derivation of the determinant representations.
- [27] K. Penc, K. Hallberg, F. Mila, and H. Shiba, *Shadow Band in the One-Dimensional Infinite-U Hubbard Model*, Phys. Rev. Lett. **77**, 1390 (1996).
- [28] J. Favand, S. Haas, K. Penc, F. Mila, and E. Dagotto, *Spectral functions of one-dimensional models of correlated electrons*, Phys. Rev. B **55**, R4859(R) (1997).
- [29] K. Penc, K. Hallberg, F. Mila, and H. Shiba, *Spectral functions of the one-dimensional Hubbard model in the $U \rightarrow +\infty$ limit: How to use the factorized wave function*, Phys. Rev. B **55**, 15475 (1997).
- [30] D. Bozi, *Deriving the one-electron Spectral Function for the 1D Hubbard Model*, arXiv:2311.16954.
- [31] S. Basak and H. Pu, *Generalized effective spin-chain formalism for strongly interacting spinor gases in optical lattices*, Phys. Rev. A **108**, 063315 (2023).
- [32] A. Imambekov and L.I. Glazman, *Phenomenology of One-Dimensional Quantum Liquids Beyond the Low-Energy Limit*, Phys. Rev. Lett. **102**, 126405 (2009).
- [33] A. Imambekov and L.I. Glazman, *Universal Theory of Nonlinear Luttinger Liquids*, Science **323**, 228 (2009).
- [34] J.M.P. Carmelo, K. Penc, L.M. Martelo, P.D. Sacramento, J.M.B. Lopes Dos Santos, R. Claessen, M. Sing and U. Schwingenschlögl, *One-electron singular branch lines of the Hubbard chain*, Europhys. Lett. **67**, 233 (2004).
- [35] J.M.P. Carmelo, K. Penc, P.D. Sacramento, M. Sing and R. Claessen, *The Hubbard model description of the TCNQ related singular features in photoemission of TTF-TCNQ*, J. Phys. C **18**, 5191 (2006).
- [36] J.M.P. Carmelo, D. Bozi and K. Penc, *Dynamical functions of a 1D correlated quantum liquid*, J. Phys. C. **20**, 415103 (2008).
- [37] Fabian H. L. Essler, *Threshold singularities in the one-dimensional Hubbard model*, Phys. Rev. B **81**, 205120 (2010).
- [38] M. Rigol and A. Muramatsu, *Emergence of quasicondensates of hard-core bosons at finite momentum*, Phys. Rev. Lett. **93**, 230404 (2004).
- [39] L. Vidmar, J.P. Ronzheimer, M. Schreiber, S. Braun, S.S. Hodgman, S. Langer, F. Heidrich-Meisner, I. Bloch, and U. Schneider, *Dynamical Quasicondensation of Hard-Core Bosons at Finite Momenta*, Phys. Rev. Lett. **115**, 175301 (2015).
- [40] J. Hubbard, *Electron correlations in narrow energy bands*, Proc. R. Soc. London, Ser. A **276**, 238 (1963).
- [41] F.H.L. Essler, H. Frahm, F. Göhmann, A. Klümper, and V.E. Korepin, *The One-Dimensional Hubbard Model* (Cambridge University Press, Cambridge, UK, 2005).
- [42] H. J. Schulz and B. Sriram Shastry, *A New Class of Exactly Solvable Interacting Fermion Models in One Dimension*, Phys. Rev. Lett. **80**, 1924 (1998).
- [43] A. Osterloh, L. Amico, and U. Eckern, *Bethe Ansatz solution of a new class of Hubbard-type models*, J. Phys. A **33** L87 (2000).
- [44] L. Cardarelli, S. Greschner and L. Santos, *Engineer-*

- ing interactions and anyon statistics by multicolor lattice depth modulations, *Phys. Rev. A*, **94**, 023615 (2016).
- [45] A. G. Izergin, A. G. Pronko, and N. I. Abarenkova, *Temperature correlators in the one-dimensional Hubbard model in the strong coupling limit*, *Phys. Lett. A* **245**, 537 (1998).
- [46] O.I. Păţu, A Klümper and A. Foerster, *Quantum critical behavior and thermodynamics of the repulsive one-dimensional Hubbard model in a magnetic field*, *Phys. Rev. B* **101**, 035149 (2020).
- [47] O.I. Păţu, A Klümper and A. Foerster, *Universality and quantum criticality of the one-dimensional spinor Bose gas*, *Phys. Rev. Lett.* **120**, 243402 (2018).
- [48] J.J. Luo, H. Pu, and X.W. Guan, *Spin-incoherent liquid and interaction-driven criticality in the one-dimensional Hubbard model*, *Phys. Rev. B* **107**, L201103 (2023).
- [49] J.J. Luo, H. Pu, and X.W. Guan, *Exact results of one-dimensional repulsive Hubbard model*, [arXiv:2307.00890](https://arxiv.org/abs/2307.00890).
- [50] Q.-W. Wang, *Exact dynamical correlations of hard-core anyons in one-dimensional lattices*, *Phys. Rev. B* **105**, 205143 (2022).
- [51] O.I. Păţu, *Exact spectral function of the Tonks-Girardeau gas at finite temperature*, *Phys. Rev. A* **106**, 053306 (2023).
- [52] O. Gamayun, E. Quinn, K. Bidzhiev, and M.B. Zvonarev, *Emergence of anyonic correlations from spin and charge dynamics in one dimension*, *Phys. Rev. A* **109**, 012209 (2024).
- [53] J. Settino, N. Lo Gullo, F. Plastina, and A. Minguzzi, *Exact Spectral Function of a Tonks-Girardeau Gas in a Lattice*, *Phys. Rev. Lett.* **126**, 065301 (2021).
- [54] E. Ilievski, and J. De Nardis, *Ballistic transport in the one-dimensional Hubbard model: The hydrodynamic approach*, *Phys. Rev. B* **96**, 081118 (2017).
- [55] B. Bertini, E. Tartaglia, and P. Calabrese, *Quantum quench in the infinitely repulsive Hubbard model: the stationary state*, *J. Stat. Mech.* 103107 (2017).
- [56] C. Rylands, B. Bertini, and P. Calabrese, *Integrable quenches in the Hubbard model*, *J. Stat. Mech.* 103103 (2022).
- [57] E. Tartaglia, P. Calabrese, and B. Bertini, *Real-time evolution in the Hubbard model with infinite repulsion*, *SciPost Physics* **12**, 028 (2022).
- [58] O. Gamayun, A. Hutsalyuk, B. Pozsgay, and M. Zvonarev, *Finite temperature spin diffusion in the Hubbard model in the strong coupling limit*, *SciPost Physics* **15**, 073 (2023).
- [59] F. Heidrich-Meisner, M. Rigol, A. Muramatsu, A. E. Feiguin, and E. Dagotto, *Ground-state reference systems for expanding correlated fermions in one dimension*, *Phys. Rev. A* **78**, 013620 (2008).
- [60] L. Vidmar, D. Iyer, and Marcos Rigol, *Emergent Eigenstate Solution to Quantum Dynamics Far from Equilibrium*, *Phys. Rev. X* **7**, 021012 (2017).
- [61] A. Takács, S. Scopa, P. Calabrese, L. Vidmar, and J. Dubail, *Quasicondensation and off-diagonal long-range order of hard-core bosons during a free expansion*, [arXiv:2401.16860](https://arxiv.org/abs/2401.16860).
- [62] I. Březinová, M. Stimpfle, S. Donsa, and A. Rubio, *Dynamical quasi-condensation in the weakly interacting Fermi-Hubbard model*, *Phys. Rev. B* **109**, 174308 (2024).
- [63] V.V. Cheianov, H. Smith, and M.B. Zvonarev, *Low-temperature crossover in the momentum distribution of cold atomic gases in one dimension*, *Phys. Rev. A* **71**, 033610 (2005).
- [64] O.I. Păţu, *Correlation functions of one-dimensional strongly interacting two-component gases*, *Phys. Rev. A* **100**, 063635 (2019).
- [65] V.V. Cheianov and M.B. Zvonarev, *One-particle equal time correlation function for the spin-incoherent infinite U Hubbard chain*, *J. Phys. A* **41**, 045002 (2008).
- [66] A. Lenard, *One-dimensional impenetrable bosons in thermal equilibrium*, *J. Math. Phys.* **7**, 1268 (1966).

Supplemental Material: Exact spectral function and nonequilibrium dynamics in the strongly interacting Hubbard model

I. EIGENSTATES AT $t = 0$ AND THEIR TIME EVOLUTION

In the impenetrable limit the doubly occupied states have infinite energy and are effectively excluded from the spectrum. In this limit the dynamics of the system is described by the effective Hamiltonian

$$\tilde{H} = P \left[-t \sum_{z=1}^{L-1} \sum_{\sigma=\{\uparrow,\downarrow\}} \left(a_{z,\sigma}^\dagger a_{z+1,\sigma} + a_{z+1,\sigma}^\dagger a_{z,\sigma} \right) + B \sum_{z=1}^L (n_{z,\uparrow} - n_{z,\downarrow}) + \sum_{z=1}^L (V(z,t) - \mu)(n_{z,\uparrow} + n_{z,\downarrow}) \right] P, \quad (\text{S-1})$$

with

$$P = \prod_{z=1}^L (1 - n_{z,\uparrow} n_{z,\downarrow}), \quad P^\dagger = P, \quad P^2 = P, \quad (\text{S-2})$$

the projector on singly occupied states. In the impenetrable regime the physical Hilbert space denoted by \mathfrak{H} has dimension 3^L while the space \mathfrak{F} in which the canonical Hubbard operators act has dimension 4^L . In the time dependent case we will denote the initial (final) Hamiltonian by \tilde{H}_I (\tilde{H}_F).

At $t = 0$ the eigenstates of the initial Hamiltonian (S-1) in the (N, M) -sector (N particles of which M have spin down) have the form

$$|\psi_{N,M}(\mathbf{k}, \boldsymbol{\lambda})\rangle = \sum_{z_1, \dots, z_N=1}^L \sum_{\alpha_1, \dots, \alpha_N=\{\uparrow,\downarrow\}}^{[N,M]} \chi_{N,M}^\alpha(z_1, \dots, z_N | \mathbf{k}, \boldsymbol{\lambda}) a_{z_N, \alpha_N}^\dagger \cdots a_{z_1, \alpha_1}^\dagger |0\rangle, \quad (\text{S-3})$$

where $[N, M]$ means that the sum is over all combinations of N α 's of which M are equal to \downarrow and $N - M$ are \uparrow , $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_N)$ and $|0\rangle$ is the Fock vacuum satisfying $a_{z,\alpha} |0\rangle = \langle 0 | a_{z,\alpha}^\dagger = 0$ for all values of z and α and $\langle 0 | 0 \rangle = 1$. The eigenstates (S-3) are parameterized by two sets of unequal (in each set) numbers $\mathbf{k} = \{k_j\}_{j=1}^N$ and $\boldsymbol{\lambda} = \{\lambda_j\}_{j=1}^M$ which characterize the charge and spin degrees of freedom. The total wavefunctions are

$$\chi_{N,M}^\alpha(\mathbf{z} | \mathbf{k}, \boldsymbol{\lambda}) = \frac{1}{N!} \left[\sum_{P \in S_N} \xi_{XX}^{(\alpha, P\boldsymbol{\alpha})}(\boldsymbol{\lambda}) e^{i \frac{\pi \kappa}{2} \sum_{1 \leq a < b \leq N} \epsilon(z_a - z_b)} \theta(z_{P(1)} < \cdots < z_{P(N)}) \right] \det_N [\phi_{k_a}(z_b)], \quad (\text{S-4})$$

with $\theta(z_{P(1)} < \cdots < z_{P(N)})$ equal to 1 when $z_{P(1)} < \cdots < z_{P(N)}$ and 0 otherwise and $\epsilon(z) = 1$ for $z \geq 0$ and $\epsilon(z) = -1$ for $z < 0$. The spin sector is described by $\xi_{XX}^{(\alpha, P\boldsymbol{\alpha})}(\boldsymbol{\lambda})$, the wavefunctions of the XX spin chain with periodic boundary conditions, defined by

$$\xi_{XX}^{(\alpha, P\boldsymbol{\alpha})}(\boldsymbol{\lambda}) = \frac{1}{N^{M/2}} \left[\prod_{1 \leq j < l \leq M} \text{sign}(n_l - n_j) \det_M (e^{i\lambda_a n_b}) \right], \quad (\text{S-5})$$

and the BAEs

$$e^{i\lambda_l N} = (-1)^{M+1}, \quad l = 1, \dots, M. \quad (\text{S-6})$$

In (S-5) n_l is the position of the l -th spin down on the auxiliary lattice of the XX spin chain. For example, in the $(N, M) = (4, 2)$ sector and $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \alpha_3, \alpha_4) = (\downarrow\downarrow\uparrow\uparrow)$ for the permutation $P = (1243)$ we have $P\boldsymbol{\alpha} = (\alpha_{P(1)}, \alpha_{P(2)}, \alpha_{P(3)}, \alpha_{P(4)}) = (\downarrow\downarrow\uparrow\uparrow)$ and $(n_1, n_2) = (1, 2)$ while for $P = (3214)$ we have $P\boldsymbol{\alpha} = (\uparrow\downarrow\downarrow\uparrow)$ and $(n_1, n_2) = (2, 3)$. The charge sector is described by a Slater determinant of eigenfunctions of the single particle Hamiltonian with O.B.C. ($\{f_i, f_j^\dagger\} = \delta_{i,j}$, $\{f_i, f_j\} = 0$, are spinless fermions)

$$H_I^{SP} = -t \sum_{z=1}^{L-1} (|z+1\rangle\langle z| + |z\rangle\langle z+1|) + \sum_{z=1}^L V(z, t \leq 0) |z\rangle\langle z|, \quad \text{with } |z\rangle = f_z^\dagger |0\rangle. \quad (\text{S-7})$$

The single particle wavefunctions satisfy $H_I^{SP} \phi_k(z) = \varepsilon(k) \phi_k(z)$ and we have

$$\tilde{H}_I |\psi_{N,M}(\mathbf{k}, \boldsymbol{\lambda})\rangle = E_{N,M}(\mathbf{k}) |\psi_{N,M}(\mathbf{k}, \boldsymbol{\lambda})\rangle, \quad E_{N,M}(\mathbf{k}) = \sum_{j=1}^N \varepsilon(k_j) - \mu N + B(2N - M). \quad (\text{S-8})$$

The wavefunctions (S-4) are the natural generalization of the Bethe ansatz wavefunctions [S1, S2], satisfy the many-body Schrödinger equation with O.B.C., have the appropriate symmetry and they form a complete orthonormal system of dimension 3^L .

The time evolved eigenstates are also described by (S-3) with the time-dependent wavefunctions

$$\chi_{N,M}^{\boldsymbol{\alpha}}(\mathbf{z}, t | \mathbf{k}, \boldsymbol{\lambda}) = \frac{1}{N!} \left[\sum_{P \in S_N} \xi_{XX}^{(\boldsymbol{\alpha}, P \boldsymbol{\alpha})}(\boldsymbol{\lambda}) e^{i \frac{\pi k}{2} \sum_{1 \leq a < b \leq N} \varepsilon(z_a - z_b)} \theta(z_{P(1)} < \dots < z_{P(N)}) \right] \det_N [\phi_{k_a}(z_b, t)], \quad (\text{S-9})$$

where $\phi_k(z, t) = e^{-i\varepsilon(k)t} \phi_k(z)$ in equilibrium situations. In nonequilibrium situations $\phi_k(z, t)$ is the unique time-dependent solution of the Schrödinger equation $i\hbar \partial \phi_k(z, t) / \partial t = H_F^{SP} \phi_k(z, t)$ with initial boundary condition $\phi_k(z, 0) = \phi_k(z)$ [remember $\phi_k(z)$ is an eigenfunction of (S-7)] where H_F^{SP} is the final single particle Hamiltonian

$$H_F^{SP} = -t \sum_{z=1}^{L-1} (|z+1\rangle\langle z| + |z\rangle\langle z+1|) + \sum_{z=1}^L V(z, t > 0) |z\rangle\langle z|, \quad \text{with } |z\rangle = f_z^\dagger |0\rangle. \quad (\text{S-10})$$

II. SKETCH OF THE DERIVATION FOR THE DETERMINANT REPRESENTATIONS

At finite temperature the correlation functions of the system initially prepared in a thermal state of the initial Hamiltonian \tilde{H}_I described by the grandcanonical potential at chemical potential μ , magnetic field B are

$$\begin{aligned} g_{\sigma}^{(-)}(x, t; y, t') &= \langle \tilde{a}_{x,\sigma}^\dagger(t) \tilde{a}_{y,\sigma}(t') \rangle_{\mu, B, T}, \quad \sigma = \{\uparrow, \downarrow\}, \\ &= \text{Tr} \left[e^{-\tilde{H}_I/T} \tilde{a}_{x,\sigma}^\dagger(t) \tilde{a}_{y,\sigma}(t') \right] / \text{Tr} \left[e^{-\tilde{H}_I/T} \right], \\ &= \sum_{N=0}^{\infty} \sum_{M=0}^{N+1} \sum_{\substack{k_1 < \dots < k_{N+1} \\ \lambda_1 < \dots < \lambda_M}} \frac{e^{-E_{N+1,M}(\mathbf{k})/T}}{\mathcal{Z}(\mu, B, T)} \langle \psi_{N+1,M}(\mathbf{k}, \boldsymbol{\lambda}) | \tilde{a}_{x,\sigma}^\dagger(t) \tilde{a}_{y,\sigma}(t') | \psi_{N+1,M}(\mathbf{k}, \boldsymbol{\lambda}) \rangle, \end{aligned} \quad (\text{S-11})$$

$$\begin{aligned} g_{\sigma}^{(+)}(x, t; y, t') &= \langle \tilde{a}_{x,\sigma}(t) \tilde{a}_{y,\sigma}^\dagger(t') \rangle_{\mu, B, T}, \quad \sigma = \{\uparrow, \downarrow\}, \\ &= \text{Tr} \left[e^{-\tilde{H}_I/T} \tilde{a}_{x,\sigma}(t) \tilde{a}_{y,\sigma}^\dagger(t') \right] / \text{Tr} \left[e^{-\tilde{H}_I/T} \right], \\ &= \sum_{N=0}^{\infty} \sum_{M=0}^N \sum_{\substack{q_1 < \dots < q_N \\ \mu_1 < \dots < \mu_M}} \frac{e^{-E_{N,M}(\mathbf{q})/T}}{\mathcal{Z}(\mu, B, T)} \langle \psi_{N,M}(\mathbf{q}, \boldsymbol{\mu}) | \tilde{a}_{x,\sigma}(t) \tilde{a}_{y,\sigma}^\dagger(t') | \psi_{N,M}(\mathbf{q}, \boldsymbol{\mu}) \rangle. \end{aligned} \quad (\text{S-12})$$

where $\mathcal{Z}(\mu, B, T) = \sum_{N=0}^{\infty} \sum_{M=0}^N \sum_{\substack{q_1 < \dots < q_N \\ \mu_1 < \dots < \mu_M}} e^{-E_{N,M}(\mathbf{q})/T} = \prod_{q=1}^{\infty} (1 + 2 \cosh(B/T) e^{-(\varepsilon(q) - \mu)/T})$ is the partition function of the initial thermal state and the time evolved operators are

$$\tilde{a}_{x,\sigma}(t) = e^{i\tilde{H}_F t} \tilde{a}_{x,\sigma} e^{-i\tilde{H}_F t}, \quad \tilde{a}_{x,\sigma}^\dagger(t) = e^{i\tilde{H}_F t} \tilde{a}_{x,\sigma}^\dagger e^{-i\tilde{H}_F t}, \quad (\text{S-13})$$

with

$$\tilde{a}_{x,\sigma} = P a_{x,\sigma} P, \quad \tilde{a}_{x,\sigma}^\dagger = P a_{x,\sigma}^\dagger P. \quad (\text{S-14})$$

It is important to note that the traces in (S-11) and (S-12) are taken in the Hilbert space \mathfrak{H} of dimension 3^L .

The derivation of the determinant representations, Eqs. (3) and (4) of the main text, is based on the summation of the form factors [S1, S2]. Using a resolution of the identity, $\mathbf{1} = \sum_{N=0}^{\infty} \sum_{M=0}^N \sum_{\substack{q_1 < \dots < q_N \\ \mu_1 < \dots < \mu_M}} |\Phi_{N,M}(\mathbf{q}, \boldsymbol{\mu})\rangle \langle \Phi_{N,M}(\mathbf{q}, \boldsymbol{\mu})|$, the mean values of bilocal operators appearing in the right hand-side of (S-11) and (S-12) can be written as

$$\langle \psi_{N+1,M}(\mathbf{k}, \boldsymbol{\lambda}) | \tilde{a}_{x,\sigma}^\dagger(t) \tilde{a}_{y,\sigma}(t') | \psi_{N+1,M}(\mathbf{k}, \boldsymbol{\lambda}) \rangle = \sum_{\substack{q_1 < \dots < q_N \\ \mu_1 < \dots < \mu_M}} \bar{\mathcal{F}}_{N,M}^{(\sigma)}(\mathbf{k}, \boldsymbol{\lambda}; \mathbf{q}, \boldsymbol{\mu} | x, t) \mathcal{F}_{N,M}^{(\sigma)}(\mathbf{k}, \boldsymbol{\lambda}; \mathbf{q}, \boldsymbol{\mu} | y, t'), \quad (\text{S-15})$$

and

$$\langle \psi_{N,\bar{M}}(\mathbf{q}, \boldsymbol{\mu}) | \tilde{a}_{x,\sigma}(t) \tilde{a}_{y,\sigma}^\dagger(t') | \psi_{N,\bar{M}}(\mathbf{q}, \boldsymbol{\mu}) \rangle = \sum_{\substack{k_1 < \dots < k_{N+1} \\ \lambda_1 < \dots < \lambda_M}} \mathcal{F}_{N,M}^{(\sigma)}(\mathbf{k}, \boldsymbol{\lambda}; \mathbf{q}, \boldsymbol{\mu} | x, t) \overline{\mathcal{F}}_{N,M}^{(\sigma)}(\mathbf{k}, \boldsymbol{\lambda}; \mathbf{q}, \boldsymbol{\mu} | y, t'), \quad (\text{S-16})$$

where

$$\mathcal{F}_{N,M}^{(\sigma)}(\mathbf{k}, \boldsymbol{\lambda}; \mathbf{q}, \boldsymbol{\mu} | x, t) = \langle \Psi_{N,\bar{M}}(\mathbf{q}, \boldsymbol{\mu}) | \tilde{a}_{x,\sigma}(t) | \psi_{N+1,M}(\mathbf{k}, \boldsymbol{\lambda}) \rangle, \quad (\text{S-17})$$

is a general form factor of the $\tilde{a}_{x,\sigma}(t)$ operator (the form factor of $\tilde{a}_{x,\sigma}^\dagger(t)$ is obtained by complex conjugation) and

$$\bar{M} = \begin{cases} M & \text{if } \sigma = \uparrow, \\ M - 1 & \text{if } \sigma = \downarrow. \end{cases} \quad (\text{S-18})$$

The determinant representations are obtained performing the following steps [S1, S2]: a) computation of the form factors for a finite size system; b) derivation of determinant formula for the mean values of the bilocal operators using a technique similar with Cauchy-Binet formula; c) taking the thermodynamic limit making use of the von Koch's determinant formula.

III. EQUAL TIME CORRELATORS

The general representations, Eqs. (3) and (4) of the main text, can be used to compute the spectral function at equilibrium, with or without an external trapping potential. In nonequilibrium situations, like the case of release from the trap or evolution from a domain wall boundary scenario, it is sufficient to investigate the $g_\sigma^{(-)}(x, t; y, t')$ at equal time $t = t'$ from which we can extract the dynamics of the real space densities and momentum distributions. For $t = t'$ the integration over η can be performed analytically and the following determinant representation is obtained $g_\uparrow^{(-)}(x, y | t) = g_\uparrow^{(-)}(x, t; y, t)$

$$g^{(-)}(x, y | t) = \det \left(1 + v^{(T,-)} + r^{(T,-)} \right) - \det \left(1 + v^{(T,-)} \right), \quad (\text{S-19})$$

with

$$v_{ab}^{(T,-)} = - \left[\gamma - e^{-i\pi\kappa \text{sign}(y-x)} \right] \sqrt{\vartheta(a)\vartheta(b)} \sum_{z=\min(x,y)}^{\max(x,y)-1} \bar{\phi}_a(z, t) \phi_b(z, t), \quad a, b = 1, 2, \dots, \quad (\text{S-20})$$

$$r_{ab}^{(T,-)} = \sqrt{\vartheta(a)\vartheta(b)} \bar{\phi}_a(x, t) \phi_b(y, t), \quad a, b = 1, 2, \dots. \quad (\text{S-21})$$

It is important to note that $g^{(-)}(x, y | t) = \overline{g^{(-)}(y, x | t)}$. At zero temperature and zero magnetic field (at $T = 0$ even an infinitesimal magnetic field completely polarizes the system making it equivalent to a single component system) we have $\gamma = 2$ and $\vartheta(q) = 1/2$ if $\varepsilon(q) \leq \mu$ and zero otherwise. In this case the relevant matrices are N -dimensional with N the number of energy levels with energy smaller than the chemical potential and

$$v_{ab}^{(0,-)} = -\frac{1}{2} \left[\gamma - e^{-i\pi\kappa \text{sign}(y-x)} \right] \sum_{z=\min(x,y)}^{\max(x,y)-1} \bar{\phi}_a(z, t) \phi_b(z, t), \quad a, b = 1, 2, \dots, N, \quad (\text{S-22})$$

$$r_{ab}^{(0,-)} = \frac{1}{2} \bar{\phi}_a(x, t) \phi_b(y, t), \quad a, b = 1, 2, \dots, N. \quad (\text{S-23})$$

[S1] A. G. Izergin, A. G. Pronko, and N. I. Abarenkova, *Temperature correlators in the one-dimensional Hubbard model in the strong coupling limit*, Phys. Lett. A **245**, 537 (1998).

[S2] A.G. Izergin and A.G. Pronko, *Temperature correlators in the two-component one-dimensional gas*, Nucl. Phys. B **520**, 594 (1998).