

DIRECTIONAL STOCKWELL TRANSFORM OF DISTRIBUTIONS

ASTRIT FERIZI AND KATERINA HADZI-VELKOVA SANEVA

ABSTRACT. We introduce and study the directional Stockwell transform as a hybrid of the directional short-time Fourier transform and the ridgelet transform. We prove an extended Parseval identity and a reconstruction formula for this transform, as well as results for the continuity of both the directional Stockwell transform and its synthesis transform on the appropriate space of test functions. Additionally, we develop a distributional framework for the directional Stockwell transform on the Lizorkin space of distributions $\mathcal{S}'_0(\mathbb{R}^n)$.

1. INTRODUCTION

The one-dimensional wavelet transform has proven to be a very efficient tool for detecting the point singularity of a function. However, it does not perform as well if the singularity is well distributed, such as singularities along curves, along hyperplanes, etc. [1]. One of the developed techniques for dealing with higher-dimensional phenomena is ridgelet analysis. The idea of the ridgelet transform is to use the Radon transform to project a hyperplane singularity into a point singularity and then act with the one-dimensional wavelet transform. It was introduced by Candès in [2, 3] who constructed from a Schwartz function ψ , a family of building blocks, called ridgelets,

$$\mathbf{x} \rightarrow \frac{1}{a} \psi \left(\frac{\mathbf{u} \cdot \mathbf{x} - b}{a} \right), \quad \mathbf{x} \in \mathbb{R}^n,$$

where $\mathbf{u} \in \mathbb{S}^{n-1}$ is the orientation parameter (\mathbb{S}^{n-1} is the unit sphere of \mathbb{R}^n), $b \in \mathbb{R}$ is the location parameter and $a \in \mathbb{R}^+$ is the scale parameter. Then, he used them to define the ridgelet transform of $f \in L^1(\mathbb{R}^n)$,

$$(1.1) \quad \mathcal{R}_\psi f(\mathbf{u}, b, a) = \int_{\mathbb{R}^n} f(\mathbf{x}) \bar{\psi} \left(\frac{\mathbf{u} \cdot \mathbf{x} - b}{a} \right) d\mathbf{x}.$$

The ridgelet transform is extended and studied on the space of Lizorkin distributions $\mathcal{S}'_0(\mathbb{R}^n)$ in [4]. Since the ridgelets are not from the class $\mathcal{S}_0(\mathbb{R}^n)$, the ridgelet transform of Lizorkin distributions can not be evaluated directly as an act of distribution on the ridgelets. Therefore the authors of [4] defined the ridgelet transform on $\mathcal{S}_0(\mathbb{R}^n)$ via a duality approach. They also provided a relationship between the ridgelet, Radon, and wavelet transforms.

Key words and phrases. Stockwell transform, Radon transform, directional Stockwell transform, distributions.

Similarly to the ridgelet analysis, Grafakos and Sansing provided an idea to localize information in time, frequency and direction. They constructed the Gabor ridge functions of a Schwartz function ψ ,

$$\mathbf{x} \rightarrow \psi(\mathbf{u} \cdot \mathbf{x} - b)e^{ia(\mathbf{x} \cdot \mathbf{u})}, \quad \mathbf{x} \in \mathbb{R}^n,$$

where $\mathbf{u} \in \mathbb{S}^{n-1}$ and $a, b \in \mathbb{R}$ [5] and used them to define a directional sensitive variant of the short-time Fourier transform of $f \in L^1(\mathbb{R}^n)$,

$$(1.2) \quad \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}^n} f(\mathbf{x}) \overline{\psi(\mathbf{u} \cdot \mathbf{x} - b)} e^{-ia(\mathbf{x} \cdot \mathbf{u})} d\mathbf{x}.$$

Their results for directionally sensitive time–frequency decompositions in $L^2(\mathbb{R}^n)$ based on Gabor systems in $L^2(\mathbb{R})$ are generalized in [6], by showing similar results for general frames for $L^2(\mathbb{R})$, both in the setting of discrete and continuous frames.

The Stockwell transform was introduced by Stockwell in [7] and combines the best features of the short-time Fourier transform and the wavelet transform. It can be seen as the phase correction of the wavelet transform [8]. A generalized version of the Stockwell transform was given by Du et al. in [9], whereas Riba and Wong introduced and studied the multi-dimensional Stockwell transform [10]. We point out that the Stockwell transform is applied in various fields, including the theory of distributions (see, e.g., [12, 11]), signal and image processing, medicine, mechanical engineering, geoscience (see, e.g., [13, 14] and references therein).

Given the limitations of the Stockwell analysis within the time-frequency plane, we introduce the directional sensitive variant of the Stockwell transform, motivated by the ridgelet transform (1.1) and the transform (1.2) defined by Grafakos and Sansing. This transform, called the directional Stockwell transform, aims to provide a more comprehensive representation of signals with directional features or anisotropic behaviors. We prove an extended Parseval identity, a reconstruction formula, and the continuity of the directional Stockwell transform and its transpose, called here the directional Stockwell synthesis operator, respectively, on the spaces $\mathcal{S}_0(\mathbb{R}^n)$ and $\mathcal{S}(\mathbb{Y}^{n+1})$, where $\mathbb{Y}^{n+1} = \mathbb{S}^{n-1} \times \mathbb{R} \times (\mathbb{R} \setminus \{0\})$.

We then use our results to develop a distributional framework for the directional Stockwell transform. In Section 5, following the duality approach, we define the directional Stockwell transform and its correspondent synthesis operator on $\mathcal{S}'_0(\mathbb{R}^n)$ and $\mathcal{S}'(\mathbb{Y}^{n+1})$, respectively. Here $\mathcal{S}'(\mathbb{Y}^{n+1})$ is a certain space of distributions of slow growth on \mathbb{Y}^{n+1} and $\mathcal{S}'_0(\mathbb{R}^n)$ stands for the Lizorkin distribution space (cf. Subsection 2.1). We emphasize that the Lizorkin spaces play a key role in Holschneider’s approach to the wavelet transform of distributions [15], as well as in the development of the distributional theory for the ridgelet and Stockwell transforms [4, 11]. Many important Schwartz distribution spaces, such as $\mathcal{E}'(\mathbb{R}^n)$, $\mathcal{O}'_C(\mathbb{R}^n)$, $L^p(\mathbb{R}^n)$, or the $\mathcal{D}'_{L^p}(\mathbb{R}^n)$ spaces, are embedded into $\mathcal{S}'_0(\mathbb{R}^n)$.

Unlike the Stockwell transform of a distribution [11], the directional Stockwell transform can not be defined by a direct evaluation of the distribution at a certain family

of test functions. The larger distribution space where the direct approach works is $D'_{L^1}(\mathbb{R}^n)$ (see Section 6 below).

2. PRELIMINARIES

2.1. Notations and spaces. In this section we give some notations and spaces that are used throughout this paper. For $\mathbf{x} = (x_1, x_2, \dots, x_n), \mathbf{y} = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ and $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{N}_0^n$, we use standard notations of n -dimensional calculus, $\mathbf{x}^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_n^{\alpha_n}$, $\partial_{\mathbf{x}}^\alpha = \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \cdots \partial_{x_n}^{\alpha_n} = \frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2} \cdots \partial x_n^{\alpha_n}}$, $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$, $|\mathbf{x}|$ denotes the Euclidean norm and $\mathbf{x} \cdot \mathbf{y} = x_1 y_1 + x_2 y_2 + \cdots + x_n y_n$ the scalar product of \mathbf{x} and \mathbf{y} . We write $A \lesssim B$ when $A \leq C \cdot B$ for some positive constant C . The Fourier transform \mathcal{F} of a function $f \in L^1(\mathbb{R}^n)$ is defined as $\mathcal{F}f(\boldsymbol{\xi}) = \widehat{f}(\boldsymbol{\xi}) = \int_{\mathbb{R}^n} f(\mathbf{x}) e^{-i\mathbf{x} \cdot \boldsymbol{\xi}} d\mathbf{x}$, $\boldsymbol{\xi} \in \mathbb{R}^n$, and it is extended to $L^2(\mathbb{R}^n)$ as usual [16].

We use the notation $(f, \varphi)_{L^2}$ for the L^2 inner product of f and φ and $\langle f, \varphi \rangle$ for the dual pairing between a distribution f and a test function φ . All dual spaces in the paper are equipped with the strong dual topology [17].

The Schwartz space $\mathcal{S}(\mathbb{R}^n)$ consists of all functions $\varphi \in C^\infty(\mathbb{R}^n)$ such that

$$(2.1) \quad \rho_m(\varphi) = \sup_{\mathbf{x} \in \mathbb{R}^n, |\alpha| \leq m} (1 + |\mathbf{x}|)^m |\partial_{\mathbf{x}}^\alpha \varphi(\mathbf{x})| < \infty,$$

for all $m \in \mathbb{N}_0$ [18, 16]. It is topologized by means of seminorms (2.1). Its dual $\mathcal{S}'(\mathbb{R}^n)$ is the space of tempered distributions. The Lizorkin space of test functions $\mathcal{S}_0(\mathbb{R}^n)$ consists of all $\varphi \in \mathcal{S}(\mathbb{R}^n)$ such that $\int_{\mathbb{R}^n} \mathbf{x}^m \varphi(\mathbf{x}) d\mathbf{x} = 0$, for all $m \in \mathbb{N}_0^n$. It is provided with the topology inherited from $\mathcal{S}(\mathbb{R}^n)$ and is of crucial importance for our analysis. Its dual space $\mathcal{S}'_0(\mathbb{R}^n)$ is known as the space of Lizorkin distributions and it is canonically isomorphic to the quotient of $\mathcal{S}'(\mathbb{R}^n)$ by the space of polynomials [15].

The space $D_{L^p}(\mathbb{R}^n)$, $1 \leq p \leq \infty$, consists of all smooth functions φ such that all derivatives belong to $L^p(\mathbb{R}^n)$. Its dual space $D'_{L^p}(\mathbb{R}^n)$ consists of all distributions f which can be represented as

$$(2.2) \quad f = \sum_{j=1}^N \partial^{\alpha_j} f_j, \quad f_j \in L^p(\mathbb{R}^n)$$

for some $N \in \mathbb{N}$, $\alpha_j \in \mathbb{N}_0^n$ ([18], Thm. XXV, page 201).

We use the notation $\mathbb{Y}^{n+1} = \mathbb{S}^{n-1} \times \mathbb{R} \times \mathbb{R}^\times$, where $\mathbb{R}^\times = \mathbb{R} \setminus \{0\}$ and \mathbb{S}^{n-1} stands for the unit sphere of \mathbb{R}^n . We always suppose that $n \geq 2$. We introduce $\mathcal{S}(\mathbb{Y}^{n+1})$ as a space of all functions $\Phi \in C^\infty(\mathbb{Y}^{n+1})$ such that

$$(2.3) \quad \rho_{s,r}^{l,m,k}(\Phi) = \sup_{(\mathbf{u}, b, a) \in \mathbb{Y}^{n+1}} (1 + |b|^2)^{r/2} (|a|^s + |a|^{-s}) |\partial_a^l \partial_b^m \Delta_{\mathbf{u}}^k \Phi(\mathbf{u}, b, a)| < \infty,$$

for all $s, r, l, m, k \in \mathbb{N}_0$, where $\Delta_{\mathbf{u}}$ stands for the Laplace-Beltrami operator on the unit sphere \mathbb{S}^{n-1} . It is topologized in the usually way. Its dual space is denoted by $\mathcal{S}'(\mathbb{Y}^{n+1})$ and will be fundamental in our definition of the directional Stockwell transform of

Lizorkin distributions, as it contains the range of this transform (cf. Section 5). We fix $|a|^{n-2}dbdad\mathbf{u}$ as a standard measure on \mathbb{Y}^{n+1} , where $d\mathbf{u}$ stands for the surface measure on the sphere \mathbb{S}^{n-1} . If F is a locally integrable function of slow growth on \mathbb{Y}^{n+1} , i.e. if there exist $C > 0, s \in \mathbb{N}_0$ such that

$$|F(\mathbf{u}, b, a)| \leq C (|a|^s + |a|^{-s}) (1 + |b|)^s, \quad (\mathbf{u}, b, a) \in \mathbb{Y}^{n+1},$$

then F will be identified with an element of $\mathcal{S}'(\mathbb{Y}^{n+1})$ via the action

$$(2.4) \quad \langle F, \Phi \rangle := \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R} \times \mathbb{R}} \int_{\mathbb{R}} F(\mathbf{u}, b, a) \Phi(\mathbf{u}, b, a) |a|^{n-2} db da d\mathbf{u}, \quad \Phi \in \mathcal{S}(\mathbb{Y}^{n+1}).$$

2.2. One-dimensional Stockwell transform. The Stockwell transform of $f \in L^2(\mathbb{R})$ with respect to a window $\psi \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$ is defined as

$$(2.5) \quad S_\psi f(b, a) = \frac{|a|}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) \bar{\psi}(a(x-b)) e^{-ixa} dx,$$

for $b \in \mathbb{R}$ and $a \in \mathbb{R} \setminus \{0\}$ [7, 8, 9]. The authors of [11] give a complete and rigorous distributional framework for the Stockwell transform based on the duality theory. Moreover, they showed that the distributional Stockwell transform can equivalently be defined by direct evaluation of a distribution on a certain family of test functions. So if $f \in \mathcal{S}'(\mathbb{R})$ and $\psi \in \mathcal{S}(\mathbb{R})$, one can replace (2.5) by

$$S_\psi f(b, a) = \frac{1}{\sqrt{2\pi}} \left\langle f(x), |a| \bar{\psi}(a(x-b)) e^{-ixa} \right\rangle, \quad b \in \mathbb{R}, a \in \mathbb{R} \setminus \{0\}.$$

2.3. The Radon transform. In this subsection we give some preliminaries of the Radon transform that are important for our analysis [19, 20].

The Radon transform of $f \in L^1(\mathbb{R}^n)$ is given by

$$Rf_{\mathbf{u}}(p) = \int_{\mathbf{x} \cdot \mathbf{u} = p} f(\mathbf{x}) d\mathbf{x} = \int_{\mathbb{R}^n} f(\mathbf{x}) \delta(p - \mathbf{x} \cdot \mathbf{u}) d\mathbf{x},$$

where $\mathbf{u} \in \mathbb{S}^{n-1}$, $p \in \mathbb{R}$ and δ is the Dirac-delta function [19]. Using the Fubini's theorem, one can show that $Rf \in L^1(\mathbb{S}^{n-1} \times \mathbb{R})$ for $f \in L^1(\mathbb{R}^n)$. The dual Radon transform $R^*\varrho$ of a function $\varrho \in L^\infty(\mathbb{S}^{n-1} \times \mathbb{R})$ is defined as

$$R^*\varrho(\mathbf{x}) = \int_{\mathbb{S}^{n-1}} \varrho(\mathbf{u}, \mathbf{x} \cdot \mathbf{u}) d\mathbf{u},$$

where $\mathbf{x} \in \mathbb{R}^n$ [19].

A connection between the Fourier transform and the Radon transform, known as the Fourier slice theorem, states that for $f \in L^1(\mathbb{R}^n)$ is true

$$(2.6) \quad \widehat{Rf_{\mathbf{u}}}(p) = \widehat{f}(p\mathbf{u}),$$

where $\mathbf{u} \in \mathbb{S}^{n-1}$ and $p \in \mathbb{R}$ [19].

3. DIRECTIONAL STOCKWELL TRANSFORM OF FUNCTIONS

In this section we define the directional Stockwell transform and prove an extended Parseval identity and a reconstruction formula that suggests the definition of the directional Stockwell synthesis operator.

Definition 3.1. Let $\psi \in \mathcal{S}(\mathbb{R})$. The directional Stockwell transform of an integrable function $f \in L^1(\mathbb{R}^n)$ is defined as

$$(3.1) \quad \begin{aligned} DS_\psi f(\mathbf{u}, b, a) &:= (f(\mathbf{x}), \psi_{\mathbf{u}, b, a}(\mathbf{x}))_{L^2} \\ &= \frac{|a|}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} f(\mathbf{x}) \overline{\psi}(a(\mathbf{x} \cdot \mathbf{u} - b)) e^{-ia(\mathbf{x} \cdot \mathbf{u})} d\mathbf{x}, \end{aligned}$$

where $\psi_{\mathbf{u}, b, a}(\mathbf{x}) = \frac{|a|}{(2\pi)^{n/2}} \psi(a(\mathbf{x} \cdot \mathbf{u} - b)) e^{ia(\mathbf{x} \cdot \mathbf{u})}$, $(\mathbf{u}, b, a) \in \mathbb{Y}^{n+1}$, $\mathbf{x} \in \mathbb{R}^n$.

The following proposition presents a useful relation between the directional Stockwell transform and the Fourier transform.

Proposition 3.2. For $\psi \in \mathcal{S}(\mathbb{R})$ and $f \in L^1(\mathbb{R}^n)$ is true

$$DS_\psi f(\mathbf{u}, b, a) = \frac{e^{-iab}}{(2\pi)^{n/2+1}} \int_{\mathbb{R}} \widehat{f}(\xi \mathbf{u}) \overline{\widehat{\psi}\left(\frac{\xi}{a} - 1\right)} e^{i\xi b} d\xi.$$

Proof. By the Fubini's theorem and the Fourier slice theorem (2.6), we have

$$\begin{aligned} DS_\psi f(\mathbf{u}, b, a) &= \frac{|a|}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} f(\mathbf{x}) \overline{\psi(a(\mathbf{x} \cdot \mathbf{u} - b))} e^{-ia(\mathbf{x} \cdot \mathbf{u})} d\mathbf{x} \\ &= \frac{|a|}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} f(\mathbf{x}) d\mathbf{x} \int_{\mathbb{R}} \overline{\psi(a(p - b))} e^{-iap} \delta(p - \mathbf{x} \cdot \mathbf{u}) dp \\ &= \frac{|a|}{(2\pi)^{n/2}} \int_{\mathbb{R}} \overline{\psi(a(p - b))} e^{-iap} dp \int_{\mathbb{R}^n} f(\mathbf{x}) \delta(p - \mathbf{x} \cdot \mathbf{u}) d\mathbf{x} \\ &= \frac{|a|}{(2\pi)^{n/2}} \int_{\mathbb{R}} Rf_{\mathbf{u}}(p) \overline{\psi(a(p - b))} e^{-iap} dp \\ &= \frac{1}{(2\pi)^{n/2+1}} \int_{\mathbb{R}} \widehat{Rf_{\mathbf{u}}}(\xi) \overline{\widehat{\psi}\left(\frac{\xi}{a} - 1\right)} e^{i(\xi - a)b} d\xi \\ &= \frac{e^{-iab}}{(2\pi)^{n/2+1}} \int_{\mathbb{R}} \widehat{f}(\xi \mathbf{u}) \overline{\widehat{\psi}\left(\frac{\xi}{a} - 1\right)} e^{i\xi b} d\xi. \end{aligned}$$

□

From the previous proof, we obtain the formula

$$(3.2) \quad DS_\psi f(\mathbf{u}, b, a) = \frac{|a|}{(2\pi)^{n/2}} \int_{\mathbb{R}} Rf_{\mathbf{u}}(p) \overline{\psi(a(p - b))} e^{-iap} dp = \frac{1}{(2\pi)^{\frac{n-1}{2}}} S_\psi(Rf_{\mathbf{u}})(b, a).$$

Thus, the directional Stockwell transform is precisely the application of a one-dimensional Stockwell transform to the slices of the Radon transform where \mathbf{u} remains fixed and p varies. The formula (3.2) presents the relation between the Radon, directional Stockwell and Stockwell transforms.

Definition 3.3. Let $\psi \in \mathcal{S}(\mathbb{R})$ be a non-trivial window. A function $\eta \in \mathcal{S}(\mathbb{R})$ is a reconstruction window of ψ if

$$(3.3) \quad C_{\psi,\eta} = \frac{1}{\pi} \int_{\mathbb{R}} \overline{\widehat{\psi}(\xi - 1)} \widehat{\eta}(\xi - 1) \frac{d\xi}{|\xi|^n}$$

is nonzero and finite. If $\eta = \psi$ in (3.3) then ψ is called an admissible window.

One can show that every non-trivial $\psi \in \mathcal{S}(\mathbb{R})$ has a reconstruction window η which may be chosen such that $e^{ix}\eta(x) \in \mathcal{S}_0(\mathbb{R})$.

The next proposition gives an extended Parseval's identity.

Proposition 3.4. (*Extended Parseval's Identity*) Let $\psi \in \mathcal{S}(\mathbb{R})$ be a non-trivial window and let $\eta \in \mathcal{S}(\mathbb{R})$ be a reconstruction window for it. Then

$$\int_{\mathbb{R}^n} f(\mathbf{x}) \overline{h(\mathbf{x})} d\mathbf{x} = \frac{1}{C_{\psi,\eta}} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} DS_\psi f(\mathbf{u}, b, a) \overline{DS_\eta h(\mathbf{u}, b, a)} |a|^{n-2} db da d\mathbf{u},$$

for $f, h \in L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$.

Proof. First we can see that $\widehat{f}(\cdot\mathbf{u}) \overline{\widehat{\psi}\left(\frac{\cdot}{a} - 1\right)}$, $\widehat{h}(\cdot\mathbf{u}) \overline{\widehat{\eta}\left(\frac{\cdot}{a} - 1\right)} \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$. Using Proposition 3.2, the Parseval's identity for the Fourier transform and the Fubini's theorem, we obtain

$$\begin{aligned} & \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} DS_\psi f(\mathbf{u}, b, a) \overline{DS_\eta h(\mathbf{u}, b, a)} |a|^{n-2} db da d\mathbf{u} \\ &= \frac{1}{(2\pi)^{n+2}} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} |a|^{n-2} \left(\int_{\mathbb{R}} \mathcal{F} \left(\widehat{f}(\cdot\mathbf{u}) \overline{\widehat{\psi}\left(\frac{\cdot}{a} - 1\right)} \right) (-b) \mathcal{F} \left(\widehat{h}(\cdot\mathbf{u}) \overline{\widehat{\eta}\left(\frac{\cdot}{a} - 1\right)} \right) (-b) db \right) da d\mathbf{u} \\ &= \frac{1}{(2\pi)^{n+2}} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} |a|^{n-2} \left(\int_{\mathbb{R}} \mathcal{F} \left(\widehat{f}(\cdot\mathbf{u}) \overline{\widehat{\psi}\left(\frac{\cdot}{a} - 1\right)} \right) (b) \mathcal{F} \left(\widehat{h}(\cdot\mathbf{u}) \overline{\widehat{\eta}\left(\frac{\cdot}{a} - 1\right)} \right) (b) db \right) da d\mathbf{u} \\ &= \frac{1}{(2\pi)^{n+1}} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} |a|^{n-2} \left(\int_{\mathbb{R}} \widehat{f}(\xi\mathbf{u}) \overline{\widehat{h}(\xi\mathbf{u}) \widehat{\eta}\left(\frac{\xi}{a} - 1\right)} \widehat{\psi}\left(\frac{\xi}{a} - 1\right) d\xi \right) da d\mathbf{u} \\ &= \frac{1}{(2\pi)^{n+1}} \int_{\mathbb{R}} d\xi \int_{\mathbb{S}^{n-1}} \widehat{f}(\xi\mathbf{u}) \overline{\widehat{h}(\xi\mathbf{u})} d\mathbf{u} \int_{\mathbb{R}^\times} |a|^{n-2} \widehat{\eta}\left(\frac{\xi}{a} - 1\right) \overline{\widehat{\psi}\left(\frac{\xi}{a} - 1\right)} da \\ &= \frac{1}{(2\pi)^{n+1}} \int_{\mathbb{R}} |\xi|^{n-1} d\xi \int_{\mathbb{S}^{n-1}} \widehat{f}(\xi\mathbf{u}) \overline{\widehat{h}(\xi\mathbf{u})} d\mathbf{u} \int_{\mathbb{R}} \widehat{\eta}(t-1) \overline{\widehat{\psi}(t-1)} \frac{dt}{|t|^n} \end{aligned}$$

$$\begin{aligned}
&= \frac{C_{\psi,\eta}}{2} \frac{1}{(2\pi)^n} \int_{\mathbb{R}} |\xi|^{n-1} d\xi \int_{\mathbb{S}^{n-1}} \widehat{f}(\xi \mathbf{u}) \overline{\widehat{h}(\xi \mathbf{u})} d\mathbf{u} \\
&= \frac{C_{\psi,\eta}}{(2\pi)^n} \int_0^\infty |\xi|^{n-1} d\xi \int_{\mathbb{S}^{n-1}} \widehat{f}(\xi \mathbf{u}) \overline{\widehat{h}(\xi \mathbf{u})} d\mathbf{u} \\
&= \frac{C_{\psi,\eta}}{(2\pi)^n} \int_{\mathbb{R}^n} \widehat{f}(\mathbf{w}) \overline{\widehat{h}(\mathbf{w})} d\mathbf{w} = C_{\psi,\eta} \int_{\mathbb{R}^n} f(\mathbf{x}) \overline{h(\mathbf{x})} d\mathbf{x}.
\end{aligned}$$

□

The next proposition gives the reconstruction formula.

Proposition 3.5. (Reconstruction formula) *Let $\psi \in \mathcal{S}(\mathbb{R})$ be a non-trivial window and let $\eta \in \mathcal{S}(\mathbb{R})$ be a reconstruction window for it. If $f \in L^1(\mathbb{R}^n)$ such that $\widehat{f} \in L^1(\mathbb{R}^n)$, then*

$$f(\mathbf{x}) = \frac{1}{C_{\psi,\eta}} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} DS_\psi f(\mathbf{u}, b, a) \eta_{\mathbf{u},b,a}(\mathbf{x}) |a|^{n-2} db da d\mathbf{u}, \quad a.e. \mathbf{x} \in \mathbb{R}^n.$$

Proof. Using Proposition 3.2 and the Fubini's theorem twice, we have

$$\begin{aligned}
&\int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} DS_\psi f(\mathbf{u}, b, a) \eta_{\mathbf{u},b,a}(\mathbf{x}) |a|^{n-2} db da d\mathbf{u} \\
&= \frac{1}{(2\pi)^{n+1}} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} e^{ia(\mathbf{u} \cdot \mathbf{x})} |a|^{n-1} da d\mathbf{u} \int_{\mathbb{R}} \eta(a(\mathbf{u} \cdot \mathbf{x} - b)) e^{-iab} db \int_{\mathbb{R}} \widehat{f}(\xi \mathbf{u}) \overline{\widehat{\psi}\left(\frac{\xi}{a} - 1\right)} e^{i\xi b} d\xi \\
&= \frac{1}{(2\pi)^{n+1}} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} e^{ia(\mathbf{u} \cdot \mathbf{x})} |a|^{n-1} da d\mathbf{u} \int_{\mathbb{R}} \widehat{f}(\xi \mathbf{u}) \overline{\widehat{\psi}\left(\frac{\xi}{a} - 1\right)} d\xi \int_{\mathbb{R}} \eta(a(\mathbf{u} \cdot \mathbf{x} - b)) e^{i(\xi-a)b} db \\
&= \frac{1}{(2\pi)^{n+1}} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} e^{ia(\mathbf{u} \cdot \mathbf{x})} |a|^{n-1} da d\mathbf{u} \int_{\mathbb{R}} \widehat{f}(\xi \mathbf{u}) \overline{\widehat{\psi}\left(\frac{\xi}{a} - 1\right)} d\xi \int_{\mathbb{R}} \frac{1}{|a|} \eta(t) e^{i(\xi-a)\mathbf{u} \cdot \mathbf{x}} e^{-i\left(\frac{\xi}{a}-1\right)t} dt \\
&= \frac{1}{(2\pi)^{n+1}} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} |a|^{n-2} da d\mathbf{u} \int_{\mathbb{R}} e^{i\xi(\mathbf{u} \cdot \mathbf{x})} \widehat{f}(\xi \mathbf{u}) \overline{\widehat{\psi}\left(\frac{\xi}{a} - 1\right)} \widehat{\eta}\left(\frac{\xi}{a} - 1\right) d\xi \\
&= \frac{1}{(2\pi)^{n+1}} \int_{\mathbb{R}} d\xi \int_{\mathbb{S}^{n-1}} e^{i\xi(\mathbf{u} \cdot \mathbf{x})} \widehat{f}(\xi \mathbf{u}) d\mathbf{u} \int_{\mathbb{R}^\times} |a|^{n-2} \overline{\widehat{\psi}\left(\frac{\xi}{a} - 1\right)} \widehat{\eta}\left(\frac{\xi}{a} - 1\right) da \\
&= \frac{1}{(2\pi)^{n+1}} \int_{\mathbb{R}} |\xi|^{n-1} d\xi \int_{\mathbb{S}^{n-1}} \widehat{f}(\xi \mathbf{u}) e^{i(\xi \mathbf{u}) \cdot \mathbf{x}} d\mathbf{u} \int_{\mathbb{R}} \overline{\widehat{\psi}(t-1)} \widehat{\eta}(t-1) \frac{dt}{|t|^n} \\
&= \frac{C_{\psi,\eta}}{2(2\pi)^n} \int_{\mathbb{R}} |\xi|^{n-1} d\xi \int_{\mathbb{S}^{n-1}} \widehat{f}(\xi \mathbf{u}) e^{i(\xi \mathbf{u}) \cdot \mathbf{x}} d\mathbf{u} \\
&= \frac{C_{\psi,\eta}}{(2\pi)^n} \int_0^\infty |\xi|^{n-1} d\xi \int_{\mathbb{S}^{n-1}} \widehat{f}(\xi \mathbf{u}) e^{i(\xi \mathbf{u}) \cdot \mathbf{x}} d\mathbf{u} \\
&= \frac{C_{\psi,\eta}}{(2\pi)^n} \int_{\mathbb{R}^n} \widehat{f}(\mathbf{w}) e^{i\mathbf{w} \cdot \mathbf{x}} d\mathbf{w} = C_{\psi,\eta} f(\mathbf{x}),
\end{aligned}$$

for a.e. $\mathbf{x} \in \mathbb{R}^n$. □

According to our choice of the standard measure on \mathbb{Y}^{n+1} , we denote by $L^2(\mathbb{Y}^{n+1}) := L^2(\mathbb{Y}^{n+1}, |a|^{n-2} dbdad\mathbf{u})$. So, the inner product on this space is

$$(F, G)_{L^2(\mathbb{Y}^{n+1})} := \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} F(\mathbf{u}, b, a) \overline{G(\mathbf{u}, b, a)} |a|^{n-2} dbdad\mathbf{u}.$$

Now, under the conditions of Proposition 3.4, we have

$$(f, h)_{L^2(\mathbb{R}^n)} = \frac{1}{C_{\psi, \eta}} (DS_\psi f, DS_\eta h)_{L^2(\mathbb{Y}^{n+1})}.$$

If ψ is an admissible window and $f \in L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$ then

$$\|DS_\psi f\|_{L^2(\mathbb{Y}^{n+1})} = \sqrt{C_{\psi, \psi}} \|f\|_{L^2(\mathbb{R}^n)}.$$

Since $\mathcal{S}(\mathbb{R}^n)$ is a dense subset of $L^2(\mathbb{R}^n)$ and the last relation holds true for all $f \in \mathcal{S}(\mathbb{R}^n)$, DS_ψ can be extended to a constant multiple of an isometric embedding $L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{Y}^{n+1})$.

The reconstruction formula suggests us to define the directional Stockwell synthesis operator that maps the function on \mathbb{Y}^{n+1} to a function on \mathbb{R}^n . Given $\psi \in \mathcal{S}(\mathbb{R})$, we define the directional Stockwell synthesis operator as

$$DS_\psi^* \Phi(\mathbf{x}) := \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} \Phi(\mathbf{u}, b, a) \psi_{\mathbf{u}, b, a}(\mathbf{x}) |a|^{n-2} dbdad\mathbf{u}, \quad \mathbf{x} \in \mathbb{R}^n.$$

The last integral is absolutely convergent if $\Phi \in \mathcal{S}(\mathbb{Y}^{n+1})$. Now, by using the reconstruction formula, we obtain

$$DS_\eta^*(DS_\psi f)(\mathbf{x}) := \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} DS_\psi f(\mathbf{u}, b, a) \eta_{\mathbf{u}, b, a}(\mathbf{x}) |a|^{n-2} dbdad\mathbf{u} = C_{\psi, \eta} f(\mathbf{x}),$$

for a.e. $\mathbf{x} \in \mathbb{R}^n$.

We also show that the directional Stockwell synthesis operator is in fact the transpose of the directional Stockwell transform in the following sense:

Proposition 3.6. *Let $\psi \in \mathcal{S}(\mathbb{R})$. If $f \in L^1(\mathbb{R}^n)$ and $\Phi \in \mathcal{S}(\mathbb{Y}^{n+1})$, then*

$$\int_{\mathbb{R}^n} f(\mathbf{x}) \overline{DS_\psi^* \Phi(\mathbf{x})} d\mathbf{x} = \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} DS_\psi f(\mathbf{u}, b, a) \Phi(\mathbf{u}, b, a) |a|^{n-2} dbdad\mathbf{u}.$$

Proof. Using the Fubini's theorem, we have

$$\begin{aligned} \int_{\mathbb{R}^n} f(\mathbf{x}) \overline{DS_\psi^* \Phi(\mathbf{x})} d\mathbf{x} &= \int_{\mathbb{R}^n} f(\mathbf{x}) d\mathbf{x} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} \Phi(\mathbf{u}, b, a) \overline{\psi_{\mathbf{u}, b, a}(\mathbf{x})} |a|^{n-2} dbdad\mathbf{u} \\ &= \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} \Phi(\mathbf{u}, b, a) |a|^{n-2} dbdad\mathbf{u} \int_{\mathbb{R}^n} f(\mathbf{x}) \overline{\psi_{\mathbf{u}, b, a}(\mathbf{x})} d\mathbf{x} \\ &= \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} DS_\psi f(\mathbf{u}, b, a) \Phi(\mathbf{u}, b, a) |a|^{n-2} dbdad\mathbf{u}. \end{aligned}$$

□

Under the standard identification (2.4), we write the last relation as

$$\langle f, \overline{DS_\psi^* \Phi} \rangle = \langle DS_\psi f, \Phi \rangle.$$

The last relation will be our model for defining the distributional directional Stockwell transform in Section 5.

4. CONTINUITY OF THE DIRECTIONAL STOCKWELL TRANSFORM ON TEST FUNCTION SPACES

Let $\mathcal{S}_1(\mathbb{R}) = \{\varphi \in \mathcal{S}(\mathbb{R}) : e^{ix}\varphi(x) \in \mathcal{S}_0(\mathbb{R})\}$ and $\psi \in \mathcal{S}_1(\mathbb{R})$. In the next two theorems we show the continuity of $DS_\psi : \mathcal{S}_0(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{Y}^{n+1})$ and $DS_\psi^* : \mathcal{S}(\mathbb{Y}^{n+1}) \rightarrow \mathcal{S}_0(\mathbb{R}^n)$.

Notice that we can extend the definition of the directional Stockwell transform as a sesquilinear mapping $DS : (f, \psi) \rightarrow DS_\psi f$, whereas the directional Stockwell synthesis operator extends to the bilinear form $DS^* : (\Phi, \psi) \rightarrow DS_\psi^* \Phi$.

Theorem 4.1. *The bilinear mapping $DS : \mathcal{S}_0(\mathbb{R}^n) \times \mathcal{S}_1(\mathbb{R}) \rightarrow \mathcal{S}(\mathbb{Y}^{n+1})$ defined as $(f, \psi) \rightarrow DS_\psi f$ is continuous.*

Proof. To prove the theorem is enough to show that for given $l, m, k, r, s \in \mathbb{N}_0$, there exist $\nu, \tau \in \mathbb{N}_0$ and $C > 0$ such that

$$(4.1) \quad \rho_{s,r}^{l,m,k}(DS_\psi f) \leq C \rho_\nu(f) \rho_\tau(\psi), \quad f \in \mathcal{S}_0(\mathbb{R}^n), \quad \psi \in \mathcal{S}_1(\mathbb{R}).$$

We may suppose that r is even and $s \geq 1$.

Step 1. By the definition of the directional Stockwell transform and the Leibniz's formula, we obtain

$$\begin{aligned} & \left| \frac{\partial^l}{\partial a^l} \frac{\partial^m}{\partial b^m} DS_\psi f(\mathbf{u}, b, a) \right| \\ &= \frac{1}{(2\pi)^{n/2}} \left| \frac{\partial^l}{\partial a^l} \frac{\partial^m}{\partial b^m} \int_{\mathbb{R}^n} a f(\mathbf{x}) \overline{\psi(a(\mathbf{x} \cdot \mathbf{u} - b))} e^{-ia(\mathbf{x} \cdot \mathbf{u})} d\mathbf{x} \right| \\ &= \frac{1}{(2\pi)^{n/2}} \left| \frac{\partial^l}{\partial a^l} \int_{\mathbb{R}^n} a^{m+1} f(\mathbf{x}) \overline{\psi^{(m)}(a(\mathbf{x} \cdot \mathbf{u} - b))} e^{-ia(\mathbf{x} \cdot \mathbf{u})} d\mathbf{x} \right| \\ &= \frac{1}{(2\pi)^{n/2}} \left| \int_{\mathbb{R}^n} f(\mathbf{x}) \sum_{k_1+k_2+k_3=l} \binom{l}{k_1, k_2, k_3} \binom{m+1}{k_1} a^{m+1-k_1} \right. \\ & \quad \left. \times e^{-ia(\mathbf{x} \cdot \mathbf{u})} (-i(\mathbf{u} \cdot \mathbf{x}))^{k_2} \overline{\psi^{(m+k_3)}(a(\mathbf{x} \cdot \mathbf{u} - b))} (\mathbf{u} \cdot \mathbf{x} - b)^{k_3} d\mathbf{x} \right| \\ &\lesssim \frac{1}{(2\pi)^{n/2}} \sum_{k_1+k_2+k_3=l} \sum_{|\alpha|=k_2} \sum_{|\beta| \leq k_3} \left| \int_{\mathbb{R}^n} f(\mathbf{x}) a^{m+1-k_1} e^{-ia(\mathbf{x} \cdot \mathbf{u})} \mathbf{x}^\alpha \mathbf{u}^\beta \right| \end{aligned}$$

$$\begin{aligned}
& \times \overline{\psi^{(m+k_3)}(a(\mathbf{x} \cdot \mathbf{u} - b))} \mathbf{x}^\beta \mathbf{u}^\beta b^{k_3 - |\beta|} d\mathbf{x} \Big| \\
& \leq \frac{1}{(2\pi)^{n/2}} \sum_{k_1+k_2+k_3=l} \sum_{|\alpha|=k_2} \sum_{|\beta| \leq k_3} (|a|^m + |a|^{-m}) (1 + |b|^2)^{\frac{k_3 - |\beta|}{2}} \\
& \times \left| \int_{\mathbb{R}^n} a f(\mathbf{x}) e^{-ia(\mathbf{x} \cdot \mathbf{u})} \mathbf{x}^{\alpha+\beta} \overline{\psi^{(m+k_3)}(a(\mathbf{x} \cdot \mathbf{u} - b))} d\mathbf{x} \right| \\
& \leq (|a|^m + |a|^{-m}) (1 + |b|^2)^{l/2} \sum_{k_1+k_2+k_3=l} \sum_{|\alpha|=k_2} \sum_{|\beta| \leq k_3} \left| DS_{\psi_{m+k_3}} f_{\alpha+\beta}(\mathbf{u}, b, a) \right|,
\end{aligned}$$

where $\psi_{m+k_3}(x) = \psi^{(m+k_3)}(x) \in \mathcal{S}_1(\mathbb{R})$, and $f_{\alpha+\beta}(\mathbf{x}) = \mathbf{x}^{\alpha+\beta} f(\mathbf{x}) \in \mathcal{S}_0(\mathbb{R}^n)$. Then

$$\rho_{s,r}^{l,m,k}(DS_\psi f) \lesssim \sum_{k_1+k_2+k_3=l} \sum_{|\alpha|=k_2} \sum_{|\beta| \leq k_3} \rho_{m+s,l+r}^{0,0,k}(DS_{\psi_{m+k_3}} f_{\alpha+\beta}).$$

So we can assume that $m = l = 0$.

Step 2. First, we have

$$\begin{aligned}
& \left| \Delta_{\mathbf{u}}^k DS_\psi f(\mathbf{u}, b, a) \right| = \frac{|a|}{(2\pi)^{n/2}} \left| \Delta_{\mathbf{u}}^k \left(\int_{\mathbb{R}^n} f(\mathbf{x}) \overline{\psi(a(\mathbf{x} \cdot \mathbf{u} - b))} e^{-ia(\mathbf{x} \cdot \mathbf{u})} d\mathbf{x} \right) \right| \\
& = \frac{|a|}{(2\pi)^{n/2}} \left| \sum_{|\alpha|,j,d \leq 2k} P_{\alpha,j,d}(\mathbf{u}) \int_{\mathbb{R}^n} a^d \mathbf{x}^\alpha f(\mathbf{x}) \overline{\psi^{(j)}(a(\mathbf{x} \cdot \mathbf{u} - b))} e^{-ia(\mathbf{x} \cdot \mathbf{u})} d\mathbf{x} \right| \\
& \lesssim (|a|^{2k} + |a|^{-2k}) \sum_{|\alpha|,j \leq 2k} \frac{|a|}{(2\pi)^{n/2}} \left| \int_{\mathbb{R}^n} \mathbf{x}^\alpha f(\mathbf{x}) \overline{\psi^{(j)}(a(\mathbf{x} \cdot \mathbf{u} - b))} e^{-ia(\mathbf{x} \cdot \mathbf{u})} d\mathbf{x} \right|,
\end{aligned}$$

where the $P_{\alpha,j,d}(\mathbf{u})$ are certain polynomials. If we put $f_\alpha(\mathbf{x}) = \mathbf{x}^\alpha f(\mathbf{x})$, $\psi_j(x) = \psi^{(j)}(x)$, we obtain

$$\left| \Delta_{\mathbf{u}}^k DS_\psi f(\mathbf{u}, b, a) \right| \lesssim (|a|^{2k} + |a|^{-2k}) \sum_{|\alpha|,j \leq 2k} \left| DS_{\psi_j} f_\alpha(\mathbf{u}, b, a) \right|,$$

that yields

$$\rho_{s,r}^{0,0,k}(DS_\psi f) \lesssim \sum_{|\alpha|,j \leq 2k} \rho_{2k+s,r}^{0,0,0}(DS_{\psi_j} f_\alpha).$$

Since $\psi_j \in \mathcal{S}_1(\mathbb{R})$, $f_\alpha \in \mathcal{S}_0(\mathbb{R}^n)$, we can suppose that $k = 0$.

Step 3. Using Proposition 3.2, we have

$$\begin{aligned}
& (1 + |b|^2)^{r/2} \left| DS_\psi f(\mathbf{u}, b, a) \right| \\
& = (1 + |b|^2)^{r/2} \left| \frac{1}{(2\pi)^{n/2+1}} \int_{\mathbb{R}} \widehat{f}(\xi \mathbf{u}) \overline{\widehat{\psi}\left(\frac{\xi}{a} - 1\right)} e^{i(\xi-a)b} d\xi \right|
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{(2\pi)^{n/2+1}} \left| \int_{\mathbb{R}} \widehat{f}(\xi \mathbf{u}) \widehat{\psi} \left(\frac{\xi}{a} - 1 \right) \left(1 - \frac{\partial^2}{\partial \xi^2} \right)^{r/2} (e^{i(\xi-a)b}) d\xi \right| \\
&= \frac{1}{(2\pi)^{n/2+1}} \left| \int_{\mathbb{R}} \left(1 - \frac{\partial^2}{\partial \xi^2} \right)^{r/2} \left(\widehat{f}(\xi \mathbf{u}) \widehat{\psi} \left(\frac{\xi}{a} - 1 \right) \right) e^{i(\xi-a)b} d\xi \right| \\
&= \frac{1}{(2\pi)^{n/2+1}} \left| \sum_{|\alpha|, j \leq r} a^{-j} Q_{\alpha, j}(\mathbf{u}) \int_{\mathbb{R}} e^{i(\xi-a)b} \widehat{f}^{(\alpha)}(\xi \mathbf{u}) \widehat{\psi}^{(j)} \left(\frac{\xi}{a} - 1 \right) d\xi \right| \\
&\lesssim (|a|^r + |a|^{-r}) \sum_{|\alpha|, j \leq r} \frac{1}{(2\pi)^{n/2+1}} \left| \int_{\mathbb{R}} e^{i(\xi-a)b} (\mathbf{x}^\alpha \widehat{f}(\mathbf{x})) (\xi \mathbf{u}) (x^j \widehat{\psi}(x)) \left(\frac{\xi}{a} - 1 \right) d\xi \right|,
\end{aligned}$$

for some polynomials $Q_{\alpha, j}(\mathbf{u})$. If we put $f_\alpha(\mathbf{x}) = \mathbf{x}^\alpha f(\mathbf{x})$, $\psi_j(x) = x^j \psi(x)$, then by Proposition 3.2 we obtain

$$\begin{aligned}
&(1 + |b|^2)^{r/2} |DS_\psi f(\mathbf{u}, b, a)| \\
&\lesssim (|a|^r + |a|^{-r}) \sum_{|\alpha|, j \leq r} \frac{1}{(2\pi)^{n/2+1}} \left| \int_{\mathbb{R}} e^{i(\xi-a)b} \widehat{f}_\alpha(\xi \mathbf{u}) \widehat{\psi}_j \left(\frac{\xi}{a} - 1 \right) d\xi \right| \\
&= (|a|^r + |a|^{-r}) \sum_{|\alpha|, j \leq r} |DS_{\psi_j} f_\alpha(\mathbf{u}, b, a)|.
\end{aligned}$$

So we conclude that

$$\rho_{s, r}^{0, 0, 0}(DS_\psi f) \lesssim \sum_{|\alpha|, j \leq r} \rho_{r+s, 0}^{0, 0, 0}(DS_{\psi_j} f_\alpha).$$

Since $\psi_j \in \mathcal{S}_1(\mathbb{R})$, $f_\alpha \in \mathcal{S}_0(\mathbb{R}^n)$ we can suppose that $r = 0$.

Step 4. Since $f \in \mathcal{S}_0(\mathbb{R}^n)$, using the Taylor expansion of \widehat{f} , we have

$$\begin{aligned}
&\frac{1}{|a|^s} |DS_\psi f(\mathbf{u}, b, a)| = \frac{1}{|a|^s (2\pi)^{n/2+1}} \left| \int_{\mathbb{R}} e^{i(\xi-a)b} \widehat{f}(\xi \mathbf{u}) \widehat{\psi} \left(\frac{\xi}{a} - 1 \right) d\xi \right| \\
&= \frac{1}{|a|^{s-1} (2\pi)^{n/2+1}} \left| \int_{\mathbb{R}} e^{ia(\xi-1)b} \widehat{f}(a\xi \mathbf{u}) \widehat{\psi}(\xi - 1) d\xi \right| \\
&\leq \frac{1}{|a|^{s-1}} \sum_{|\alpha|=s-1} \frac{1}{(2\pi)^{n/2+1}} \left| \int_{\mathbb{R}} e^{ia(\xi-1)b} \frac{(a\xi \mathbf{u})^\alpha}{\alpha!} \widehat{f}^{(\alpha)}(a\xi_0 \mathbf{u}) \widehat{\psi}(\xi - 1) d\xi \right| \\
&\lesssim \sum_{|\alpha|=s-1} \left| \int_{\mathbb{R}} e^{ia(\xi-1)b} \xi^{s-1} \widehat{f}^{(\alpha)}(a\xi_0 \mathbf{u}) \widehat{\psi}(\xi - 1) d\xi \right| \\
&\lesssim \sum_{|\alpha|=s-1} \int_{\mathbb{R}^n} |\mathbf{x}^\alpha f(\mathbf{x})| d\mathbf{x} \int_{\mathbb{R}} |\xi|^{s-1} |\widehat{\psi}(\xi - 1)| d\xi \lesssim \rho_{s+n}(f) \rho_{s+1}(\psi).
\end{aligned}$$

Since $\psi \in \mathcal{S}_1(\mathbb{R})$, we have

$$\begin{aligned} |a|^s |DS_\psi f(\mathbf{u}, b, a)| &= \frac{|a|^s}{(2\pi)^{n/2+1}} \left| \int_{\mathbb{R}} e^{i(\xi-a)b} \widehat{f}(\xi \mathbf{u}) \widehat{\psi} \left(\frac{\xi}{a} - 1 \right) d\xi \right| \\ &= \frac{|a|^s}{(2\pi)^{n/2+1}} \left| \int_{\mathbb{R}} e^{i(\xi-a)b} \widehat{f}(\xi \mathbf{u}) \left(\frac{\xi}{a} \right)^s \frac{1}{s!} \overline{\widehat{\psi}^{(s)} \left(\frac{\xi_0}{a} - 1 \right)} d\xi \right| \\ &\lesssim \int_{\mathbb{R}} |\eta^s \psi(\eta)| d\eta \int_{\mathbb{R}} |\xi|^s |\widehat{f}(\xi \mathbf{u})| d\xi \lesssim \rho_{s+n+1}(f) \rho_{s+2}(\psi). \end{aligned}$$

Then, we obtain

$$\rho_{s,0}^{0,0,0}(DS_\psi f) \lesssim \rho_{s+n+1}(f) \rho_{s+2}(\psi).$$

Summing up all the estimates, we find that (4.1) holds true for $\nu = 4k + 4l + 3r + m + s + n + 1$ and $\tau = 4k + 4l + 3r + 2m + s + 2$. \square

We now analyze the directional Stockwell synthesis operator.

Theorem 4.2. *The bilinear mapping $DS^* : \mathcal{S}(\mathbb{Y}^{n+1}) \times \mathcal{S}_1(\mathbb{R}) \rightarrow \mathcal{S}_0(\mathbb{R}^n)$ defined as $(\Phi, \psi) \rightarrow DS_\psi^* \Phi$ is continuous.*

Proof. Since $\Phi \in \mathcal{S}(\mathbb{Y}^{n+1})$ and $\psi \in \mathcal{S}_1(\mathbb{R})$ then $DS_\psi^* \Phi \in \mathbb{C}^\infty(\mathbb{R}^n)$. Using the Parseval's identity and the Fubini's theorem, we can write

$$\begin{aligned} h(\mathbf{x}) &:= DS_\psi^* \Phi(\mathbf{x}) = \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} \Phi(\mathbf{u}, b, a) \psi_{\mathbf{u}, b, a}(\mathbf{x}) |a|^{n-2} db da d\mathbf{u} \\ &= \frac{1}{2\pi} \int_{\mathbb{S}^{n-1}} d\mathbf{u} \int_{\mathbb{R}^\times} |a|^{n-2} da \int_{\mathbb{R}} \widehat{\Phi}(\mathbf{u}, w, a) \overline{\widehat{\psi_{\mathbf{u}, \cdot, a}(\mathbf{x})}}(w) dw \\ &= \frac{1}{(2\pi)^{n/2+1}} \int_{\mathbb{S}^{n-1}} d\mathbf{u} \int_{\mathbb{R}^\times} |a|^{n-2} da \int_{\mathbb{R}} \widehat{\Phi}(\mathbf{u}, w, a) e^{i(a+w)\mathbf{x} \cdot \mathbf{u}} \widehat{\psi} \left(\frac{w}{a} \right) dw \\ &= \frac{1}{(2\pi)^{n/2+1}} \int_{\mathbb{S}^{n-1}} d\mathbf{u} \int_{\mathbb{R}^\times} |a|^{n-2} da \int_{\mathbb{R}} \widehat{\Phi}(\mathbf{u}, w - a, a) e^{iw(\mathbf{x} \cdot \mathbf{u})} \widehat{\psi} \left(\frac{w}{a} - 1 \right) dw \\ &= \frac{1}{(2\pi)^{n/2+1}} \int_{\mathbb{R}} dw \int_{\mathbb{S}^{n-1}} e^{iw(\mathbf{x} \cdot \mathbf{u})} d\mathbf{u} \int_{\mathbb{R}^\times} |a|^{n-2} \widehat{\Phi}(\mathbf{u}, w - a, a) \widehat{\psi} \left(\frac{w}{a} - 1 \right) da \\ &= \frac{1}{(2\pi)^{n/2+1}} \int_0^\infty w^{n-1} dw \int_{\mathbb{S}^{n-1}} e^{iw(\mathbf{x} \cdot \mathbf{u})} d\mathbf{u} \int_{\mathbb{R}} \left(\widehat{\Phi} \left(\mathbf{u}, w - \frac{w}{t}, \frac{w}{t} \right) \right. \\ &\quad \left. + \widehat{\Phi} \left(-\mathbf{u}, -w + \frac{w}{t}, -\frac{w}{t} \right) \right) \widehat{\psi} \left(t - 1 \right) \frac{dt}{|t|^n}, \end{aligned}$$

where $\widehat{\Phi}$ stands for the Fourier transform of $\Phi(\mathbf{u}, b, a)$ with respect to the variable b .

Now we define

$$G(\mathbf{u}, w) := \int_{\mathbb{R}} \left(\widehat{\Phi} \left(\mathbf{u}, w - \frac{w}{t}, \frac{w}{t} \right) + \widehat{\Phi} \left(-\mathbf{u}, -w + \frac{w}{t}, -\frac{w}{t} \right) \right) \widehat{\psi} \left(t - 1 \right) \frac{dt}{|t|^n},$$

$w \in \mathbb{R}, \mathbf{u} \in \mathbb{S}^{n-1}$. Clearly G is a smooth function and

$$(4.2) \quad h(\mathbf{x}) = \frac{1}{(2\pi)^{n/2+1}} \int_0^\infty w^{n-1} dw \int_{\mathbb{S}^{n-1}} G(\mathbf{u}, w) e^{iw(\mathbf{x} \cdot \mathbf{u})} d\mathbf{u}.$$

We will show that $G \in \mathcal{S}(\mathbb{S}^{n-1} \times \mathbb{R})$. For given $N, q, k \in \mathbb{N}_0$, it holds

$$\begin{aligned} & \sup_{(\mathbf{u}, w) \in \mathbb{S}^{n-1} \times \mathbb{R}} \left| w^N \frac{\partial^q}{\partial w^q} \Delta_{\mathbf{u}}^k \left(\int_{\mathbb{R}} \widehat{\Phi}(\mathbf{u}, w - \frac{w}{t}, \frac{w}{t}) \widehat{\psi}(t-1) \frac{dt}{|t|^n} \right) \right| \\ &= \sup_{(\mathbf{u}, w) \in \mathbb{S}^{n-1} \times \mathbb{R}} \left| w^N \int_{\mathbb{R}} \frac{\partial^q}{\partial w^q} \Delta_{\mathbf{u}}^k \widehat{\Phi}(\mathbf{u}, w - \frac{w}{t}, \frac{w}{t}) \widehat{\psi}(t-1) \frac{dt}{|t|^n} \right| \\ &= \sup_{(\mathbf{u}, w) \in \mathbb{S}^{n-1} \times \mathbb{R}} \left| w^N \int_{\mathbb{R}} \widehat{\psi}(t-1) \frac{dt}{|t|^n} \int_{\mathbb{R}} \sum_{l_1+l_2=q} \binom{q}{l_1, l_2} \frac{\partial^{l_1}}{\partial w^{l_1}} \left(\Delta_{\mathbf{u}}^k \Phi(\mathbf{u}, b, \frac{w}{t}) \right) \right. \\ & \quad \left. \times \frac{\partial^{l_2}}{\partial w^{l_2}} \left(e^{-ib(w-\frac{w}{t})} \right) db \right| \\ &= \sup_{(\mathbf{u}, w) \in \mathbb{S}^{n-1} \times \mathbb{R}} \left| w^N \sum_{l_1+l_2=q} \binom{q}{l_1, l_2} \int_{\mathbb{R}} \left(1 - \frac{1}{t} \right)^{l_2} \frac{1}{t^{l_1}} \widehat{\psi}(t-1) \frac{dt}{|t|^n} \right. \\ & \quad \left. \int_{\mathbb{R}} b^{l_2} \frac{\partial^{l_1} \Delta_{\mathbf{u}}^k \Phi(\mathbf{u}, b, \frac{w}{t})}{\partial \left(\frac{w}{t} \right)^{l_1}} e^{-iw(1-\frac{1}{t})b} db \right| \lesssim \rho_{r'}(\widehat{\psi}) \rho_{v'_1, v'_2}^{q'_1, q'_2, k}(\Phi), \end{aligned}$$

for some $q'_1, q'_2, r', v'_1, v'_2 \in \mathbb{N}_0$. Similar, we have

$$\sup_{(\mathbf{u}, w) \in \mathbb{S}^{n-1} \times \mathbb{R}} \left| w^N \frac{\partial^q}{\partial w^q} \Delta_{\mathbf{u}}^k \left(\int_{\mathbb{R}} \widehat{\Phi}(-\mathbf{u}, -w + \frac{w}{t}, -\frac{w}{t}) \widehat{\psi}(t-1) \frac{dt}{|t|^n} \right) \right| \lesssim \rho_{r''}(\widehat{\psi}) \rho_{v''_1, v''_2}^{q''_1, q''_2, k}(\Phi),$$

for some $q''_1, q''_2, r'', v''_1, v''_2 \in \mathbb{N}_0$. So, we have shown that for given $N, q, k \in \mathbb{N}_0$,

$$(4.3) \quad \sup_{(\mathbf{u}, w) \in \mathbb{S}^{n-1} \times \mathbb{R}} \left| w^N \frac{\partial^q}{\partial w^q} \Delta_{\mathbf{u}}^k G(\mathbf{u}, w) \right| \lesssim \rho_r(\widehat{\psi}) \rho_{v_1, v_2}^{q_1, q_2, k}(\Phi),$$

for some $q_1, q_2, r, v_1, v_2 \in \mathbb{N}_0$.

Then, by relation (4.2) and the Fourier inversion formula in polar coordinates, we obtain $h \in \mathcal{S}(\mathbb{R}^n)$ and

$$(4.4) \quad \widehat{h}(w\mathbf{u}) = (2\pi)^{n/2-1} G(\mathbf{u}, w),$$

for $w \in \mathbb{R}^+, \mathbf{u} \in \mathbb{S}^{n-1}$. To show that $h \in \mathcal{S}_0(\mathbb{R}^n)$ is equivalent of showing that

$$\lim_{\mathbf{w} \rightarrow 0} \frac{\widehat{h}(\mathbf{w})}{|\mathbf{w}|^k} = 0, \quad \forall k \in \mathbb{N}_0.$$

Since $\Phi \in \mathcal{S}(\mathbb{Y}^{n+1})$, there exists $C_k > 0$ such that

$$|\widehat{\Phi}(\mathbf{u}, \omega, a)| \leq C_k |a|^{k+1},$$

uniformly for $\omega \in \mathbb{R}$, $\mathbf{u} \in \mathbb{S}^{n-1}$. Then, for $w \in \mathbb{R}^+$ and $\mathbf{u} \in \mathbb{S}^{n-1}$,

$$|\widehat{h}(w\mathbf{u})| \lesssim C_k \int_{\mathbb{R}} \left| \frac{w}{t} \right|^{k+1} |\widehat{\psi}(t-1)| \frac{dt}{|t|^n}.$$

Since $\psi \in \mathcal{S}_1(\mathbb{R})$ we have that $\int_{\mathbb{R}} |\widehat{\psi}(t-1)| \frac{dt}{|t|^{n+k+1}} < \infty$. Then $\lim_{w \rightarrow 0^+} \frac{\widehat{h}(w\mathbf{u})}{|w|^k} = 0$, $\forall k \in \mathbb{N}_0$, i.e.

$$\lim_{\mathbf{w} \rightarrow 0} \frac{\widehat{h}(\mathbf{w})}{|\mathbf{w}|^k} = 0, \quad \forall k \in \mathbb{N}_0.$$

Thus $h = DS_{\psi}^* \Phi \in \mathcal{S}_0(\mathbb{R}^n)$.

We now prove the continuity of the bilinear mapping DS^* . Since $\varphi \rightarrow \widehat{\varphi}$ is an automorphism of the \mathcal{S} spaces, the family

$$\widehat{\rho}_v(\varphi) = \rho_v(\widehat{\varphi}), \quad \varphi \in \mathcal{S}(\mathbb{R}), \quad v \in \mathbb{N}_0,$$

is a base of seminorms for the topology of $\mathcal{S}(\mathbb{R})$. We will define different family of seminorms on $\mathcal{S}_0(\mathbb{R}^n)$. The seminorms

$$\dot{\rho}_{N,q,k}(\varphi) := \sup_{(\mathbf{u},w) \in \mathbb{S}^{n-1} \times \mathbb{R}} \left| w^N \frac{\partial^q}{\partial w^q} \Delta_{\mathbf{u}}^k \widehat{\varphi}(w\mathbf{u}) \right|, \quad N, q, k \in \mathbb{N}_0,$$

are a base of continuous seminorms for the topology of $\mathcal{S}_0(\mathbb{R}^n)$ ([4], p. 10). So, for given $N, q, k \in \mathbb{N}_0$, by equations (4.3) and (4.4) we obtain

$$\begin{aligned} \dot{\rho}_{N,q,k}(h) &= \sup_{(\mathbf{u},w) \in \mathbb{S}^{n-1} \times \mathbb{R}} \left| w^N \frac{\partial^q}{\partial w^q} \Delta_{\mathbf{u}}^k \widehat{h}(w\mathbf{u}) \right| \\ &\lesssim \sup_{(\mathbf{u},w) \in \mathbb{S}^{n-1} \times \mathbb{R}} \left| w^N \frac{\partial^q}{\partial w^q} \Delta_{\mathbf{u}}^k G(\mathbf{u}, w) \right| \lesssim \rho_r(\widehat{\psi}) \rho_{v_1, v_2}^{q_1, q_2, k}(\Phi), \end{aligned}$$

for some $q_1, q_2, r, v_1, v_2 \in \mathbb{N}_0$. □

The next proposition is a direct consequence of Proposition 3.5, Theorem 4.1 and Theorem 4.2.

Proposition 4.3. *Let $\psi \in \mathcal{S}_1(\mathbb{R})$ be a non-trivial window and $\eta \in \mathcal{S}_1(\mathbb{R})$ be a reconstruction window for it. Then*

$$(4.5) \quad \frac{1}{C_{\psi, \eta}} DS_{\eta}^* \circ DS_{\psi} = Id_{\mathcal{S}_0(\mathbb{R}^n)}.$$

5. DIRECTIONAL STOCKWELL TRANSFORM ON $\mathcal{S}'_0(\mathbb{R}^n)$

We are ready to define the directional Stockwell transform of Lizorkin distributions.

Definition 5.1. Let $\psi \in \mathcal{S}_1(\mathbb{R})$. We define the directional Stockwell transform of $f \in \mathcal{S}'_0(\mathbb{R}^n)$ with respect to ψ as the element $DS_\psi f \in \mathcal{S}'(\mathbb{Y}^{n+1})$ whose action on test functions is given by

$$\langle DS_\psi f, \Phi \rangle := \langle f, \overline{DS_\psi^* \Phi} \rangle, \quad \Phi \in \mathcal{S}(\mathbb{Y}^{n+1}).$$

Definition 5.2. Let $\psi \in \mathcal{S}_1(\mathbb{R})$. We define the directional Stockwell synthesis operator $DS_\psi^* : \mathcal{S}'(\mathbb{Y}^{n+1}) \rightarrow \mathcal{S}'_0(\mathbb{R}^n)$ as

$$\langle DS_\psi^* F, \varphi \rangle := \langle F, \overline{DS_\psi \overline{\varphi}} \rangle, \quad F \in \mathcal{S}'(\mathbb{Y}^{n+1}), \quad \varphi \in \mathcal{S}_0(\mathbb{R}^n).$$

The Definitions (5.1) and (5.2) are valid according to Theorem 4.2 and Theorem 4.1, respectively.

The following continuity result is obtained by taking transposes in Theorems 4.1 and Theorem 4.2.

Proposition 5.3. Let $\psi \in \mathcal{S}_1(\mathbb{R})$. The directional Stockwell transform $DS_\psi : \mathcal{S}'_0(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{Y}^{n+1})$ and the directional Stockwell synthesis operator $DS_\psi^* : \mathcal{S}'(\mathbb{Y}^{n+1}) \rightarrow \mathcal{S}'_0(\mathbb{R}^n)$ are continuous linear maps.

We can also generalize the reconstruction formula (4.5) to the space of Lizorkin distributions.

Proposition 5.4. Let $\psi \in \mathcal{S}_1(\mathbb{R})$ be a non-trivial window and $\eta \in \mathcal{S}_1(\mathbb{R})$ be a reconstruction window for it. Then

$$\frac{1}{C_{\psi, \eta}} DS_\eta^* \circ DS_\psi = Id_{\mathcal{S}'_0(\mathbb{R}^n)}.$$

Proof. Let $f \in \mathcal{S}'_0(\mathbb{R}^n)$ and $\varphi \in \mathcal{S}_0(\mathbb{R}^n)$. Using Definitions 5.1 and 5.2 and Proposition 4.3, we obtain

$$\langle DS_\eta^* \circ DS_\psi f, \varphi \rangle = \langle DS_\psi f, \overline{DS_\eta \overline{\varphi}} \rangle = \langle f, \overline{DS_\psi^* (DS_\eta \overline{\varphi})} \rangle = \langle f, \overline{C_{\eta, \psi} \overline{\varphi}} \rangle = C_{\psi, \eta} \langle f, \varphi \rangle.$$

□

6. DIRECTIONAL STOCKWELL TRANSFORM ON $D'_{L^1}(\mathbb{R}^n)$

Since, $\psi_{\mathbf{u}, b, a} \in D_{L^\infty}(\mathbb{R}^n)$, Definition 3.1 also works for $f \in D'_{L^1}(\mathbb{R}^n)$. In this case, by (2.2) and (3.1) we have

(6.1)

$$DS_\psi f(\mathbf{u}, b, a) := \sum_{j=1}^N \langle \partial^{\alpha_j} f_j(\mathbf{x}), \overline{\psi_{\mathbf{u}, b, a}(\mathbf{x})} \rangle_{\mathbf{x}} = \sum_{j=1}^N (-1)^{|\alpha_j|} \int_{\mathbb{R}^n} f_j(\mathbf{x}) \overline{\frac{\partial^{|\alpha_j|}}{\partial \mathbf{x}^{\alpha_j}} \psi_{\mathbf{u}, b, a}(\mathbf{x})} d\mathbf{x},$$

where $f_j \in L^1(\mathbb{R}^n)$.

Since the space $D'_{L^1}(\mathbb{R}^n)$ contains the space of compactly supported distributions $\mathcal{E}'(\mathbb{R}^n)$ and the space of convolutors $\mathcal{O}'_C(\mathbb{R}^n)$, Definition 3.1 also makes sense for $f \in$

$\mathcal{E}'(\mathbb{R}^n)$ or $f \in \mathcal{O}'_C(\mathbb{R}^n)$. On the other hand, Definition 3.1 does not work for $f \in \mathcal{S}'(\mathbb{R}^n)$ since $\psi_{\mathbf{u},b,a} \notin \mathcal{S}(\mathbb{R}^n)$.

In the next Proposition we show that Definition 3.1 coincides with Definition 5.1 under our convention (2.4) for identifying functions with distributions on \mathbb{Y}^{n+1} .

Proposition 6.1. *Let $f \in D'_{L^1}(\mathbb{R}^n)$ and $\psi \in \mathcal{S}_1(\mathbb{R})$. The directional Stockwell transform of f is given by the function (3.1), i.e.*

$$\langle DS_\psi f, \Phi \rangle = \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} DS_\psi f(\mathbf{u}, b, a) \Phi(\mathbf{u}, b, a) |a|^{n-2} db da d\mathbf{u}, \quad \Phi \in \mathcal{S}(\mathbb{Y}^{n+1}).$$

Proof. Since $f \in D'_{L^1}(\mathbb{R}^n)$, by (2.2) we have that $f = \sum_{j=1}^N \partial^{\alpha_j} f_j$, where $f_j \in L^1(\mathbb{R}^n)$ for some $N \in \mathbb{N}$, $\alpha_j \in \mathbb{N}_0^n$. Now for $\Phi \in \mathcal{S}(\mathbb{Y}^{n+1})$, we obtain

$$(6.2) \quad \langle DS_\psi(\partial^{\alpha_j} f_j), \Phi \rangle := \langle \partial^{\alpha_j} f_j, \overline{DS_\psi^* \Phi} \rangle = (-1)^{|\alpha_j|} \left\langle f_j(\mathbf{x}), \frac{\partial^{|\alpha_j|}}{\partial \mathbf{x}^{\alpha_j}} \overline{DS_\psi^* \Phi(\mathbf{x})} \right\rangle.$$

On the other hand,

$$\begin{aligned} \frac{\partial^{|\alpha_j|}}{\partial \mathbf{x}^{\alpha_j}} \overline{DS_\psi^* \Phi(\mathbf{x})} &= \frac{\partial^{|\alpha_j|}}{\partial \mathbf{x}^{\alpha_j}} \overline{\int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} \Phi(\mathbf{u}, b, a) \psi_{\mathbf{u},b,a}(\mathbf{x}) |a|^{n-2} db da d\mathbf{u}} \\ &= \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} \Phi(\mathbf{u}, b, a) \frac{|a|}{(2\pi)^{n/2}} \frac{\partial^{|\alpha_j|}}{\partial \mathbf{x}^{\alpha_j}} \left(\overline{\psi(a(\mathbf{x} \cdot \mathbf{u} - b)) e^{-ia(\mathbf{x} \cdot \mathbf{u})}} \right) |a|^{n-2} db da d\mathbf{u} \\ &= \sum_{k \leq \alpha_j} \binom{\alpha_j}{k} (-i)^{|\alpha_j - k|} \overline{\int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^\times} \int_{\mathbb{R}} (a\mathbf{u})^{\alpha_j} \Phi(\mathbf{u}, b, a) \psi_{\mathbf{u},b,a}^{(|k|)}(\mathbf{x}) |a|^{n-2} db da d\mathbf{u}} \\ (6.3) \quad &= \sum_{k \leq \alpha_j} \binom{\alpha_j}{k} (-i)^{|\alpha_j - k|} \overline{DS_{\psi^{(|k|)}}^* ((a\mathbf{u})^{\alpha_j} \Phi)(\mathbf{x})}. \end{aligned}$$

By relations (6.2) and (6.3), we obtain

$$\begin{aligned} \langle DS_\psi(\partial^{\alpha_j} f_j), \Phi \rangle &= (-1)^{|\alpha_j|} \sum_{k \leq \alpha_j} \binom{\alpha_j}{k} (-i)^{|\alpha_j - k|} \left\langle f_j(\mathbf{x}), \overline{DS_{\psi^{(|k|)}}^* ((a\mathbf{u})^{\alpha_j} \Phi)(\mathbf{x})} \right\rangle \\ &= (-1)^{|\alpha_j|} \sum_{k \leq \alpha_j} \binom{\alpha_j}{k} (-i)^{|\alpha_j - k|} \langle DS_{\psi^{(|k|)}} f_j, (a\mathbf{u})^{\alpha_j} \Phi \rangle. \end{aligned}$$

The last relation implies

$$DS_\psi(\partial^{\alpha_j} f_j) = (-1)^{|\alpha_j|} \sum_{k \leq \alpha_j} \binom{\alpha_j}{k} (-i)^{|\alpha_j - k|} (a\mathbf{u})^{\alpha_j} DS_{\psi^{(|k|)}} f_j.$$

Therefore, we can suppose that $f \in L^1(\mathbb{R}^n)$.

Now, under this assumption, Proposition 3.6 and the standard identification (2.4), we have

$$\begin{aligned} \langle DS_\psi f, \Phi \rangle &= \langle f, \overline{DS_\psi^* \Phi} \rangle = \int_{\mathbb{R}^n} f(\mathbf{x}) \overline{DS_\psi^* \Phi(\mathbf{x})} d\mathbf{x} \\ &= \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R} \times \mathbb{R}} DS_\psi f(\mathbf{u}, b, a) \Phi(\mathbf{u}, b, a) |a|^{n-2} db da d\mathbf{u} = \langle DS_\psi f(\mathbf{u}, b, a), \Phi(\mathbf{u}, b, a) \rangle. \end{aligned}$$

□

REFERENCES

- [1] Candès E.J. Harmonic Analysis of Neural Networks. *Appland Comput Harmon Anal.* 1999; 6: 197-218. <https://doi.org/10.1006/acha.1998.0248>
- [2] Candès E.J. Ridgelet: theory and applications. Ph.D. thesis. Department of Statistics, Stanford University. 1999.
- [3] Candès E.J, Donoho DL. Ridgelets: a key to higher-dimensional intermittency? *Roy Soc Lond Philos Trans Ser A Math Phys Eng Sci.* 1999; 357:2495–2509. <https://doi.org/10.1098/rsta.1999.0444>
- [4] Kostadinova S, Pilipović S, Saneva K, Vindas J. The Ridgelet transform of distributions. *Integral Transforms Spec Funct.* 2014; 25(5), 344-358. <https://doi.org/10.1080/10652469.2013.853057>
- [5] Grafakos L, Sansing C. Gabor frames and directional time–frequency analysis. *Appl Comput Harmon Anal.* 2008; 25, 47-67. <https://doi.org/10.1016/j.acha.2007.09.004>
- [6] Christensen O, Forster B, Massopust P. Directional time frequency analysis via continuous frames. *Bull. Aust. Math. Soc.* 2015; 92(2), 268–281. <https://doi.org/10.1017/S0004972715000337>
- [7] Stockwell R.G, Mansinha L, Lowe R.P. Localization of the Complex Spectrum: The S transform. *IEEE Transactions on Signal Processing.* 1996; 44(4): 998-1001. DOI: 10.1109/78.492555
- [8] Stockwell R.G, Why use the S transform? *Pseudo-Differential Oper Partial Differ Equ Time Frequency Anal.* 2007; 52: 279–309.
- [9] Du J, Wong M.W, Zhu H. Continuous and discrete inversion formulas for the Stockwell transform. *Integral Transforms and Spec Funct.* 2007; 18(8): 537-543.
- [10] Riba L, Wong MW. Continuous inversion formulas for multi-dimensional Stockwell transforms. *Math Model Nat Phenom.* 2013; 8(1), 215-229. <https://doi.org/10.1080/10652460701359032>
- [11] Saneva K, Atanasova S, Buralieva J.V. Tauberian theorems for the Stockwell transform of Lizorkin distributions. *Appl Anal.* 2020;99(4):596-610. <https://doi.org/10.1080/00036811.2018.1506104>
- [12] Atanasova S, Pilipović S, Prangoski B, Saneva K. Characterisation of wave front sets by the Stockwell transform. *J Math Anal Appl.* 2020; 490(2). <https://doi.org/10.1016/j.jmaa.2020.124329>
- [13] Wei D, Zhang Y, Li YM. Linear canonical Stockwell transform: theory and applications. *IEEE Trans Signal Process.* 2022;70:1333–1347. doi: 10.1109/TSP.2022.3152402
- [14] Kumar R, Saxena A, Kumar R, Marwaha S, Singh J, Gyanendra K. A Comprehensive Overview on Modified Versions of Stockwell Transform For Power Quality Monitoring. *IEEE Access.* 2022; 10, 91963-91975. DOI: 10.1109/ACCESS.2022.3202309
- [15] Holschneider M. Wavelets. An Analysis Tool. New York: Clarendon/Oxford University Press; 1995.
- [16] Hörmander L. The analysis of linear partial differential operators I: Distribution theory and Fourier Analysis. Berlin: Springer; 1983.
- [17] Trèves F. Topological vector spaces, distributions and kernels. New York-London: Academic press; 1967.
- [18] Schwartz L. Théorie des Distributions. Paris: Hermann; 1966.

- [19] Hertle A. Continuity of the Radon Transform and its Inverse on Euclidean Space. *Mathematische Zeitschrift*. 1983; 184, 165-192.
- [20] Helgason S. *The Radon transform*. 2nd ed. Boston, MA: Birkhäuser Boston, Inc.; 1999.

FACULTY OF MATHEMATICS AND NATURAL SCIENCES, UNIVERSITY OF PRISHTINA, GEORGE BUSH 31, PRISHTINA, 10000, KOSOVO
Email address: ferizi.astrit@gmail.com

FACULTY OF ELECTRICAL ENGINEERING AND INFORMATION TECHNOLOGIES, Ss. CYRIL AND METHODIUS UNIVERSITY IN SKOPJE, RUGJER BOSHKOVIKJ 18, SKOPJE, 1000, REPUBLIC OF NORTH MACEDONIA
Email address: saneva@feit.ukim.edu.mk