

THE LINEAR TARGETING PROBLEM

KYLE BIERLY, STEPHAN RAMON GARCIA, AND ROGER A. HORN

ABSTRACT. For given real or complex $m \times n$ data matrices X, Y , we investigate when there is a matrix A such that $AX = Y$, and A is invertible, Hermitian, positive (semi)definite, unitary, an orthogonal projection, a reflection, complex symmetric, or normal.

1. INTRODUCTION

The *linear targeting problem* is to construct an $A \in M_m(\mathbb{F})$ such that $AX = Y$, in which $X, Y \in M_{m \times n}(\mathbb{F})$ are given and $\mathbb{F} = \mathbb{R}$ or \mathbb{C} . We say that X is the *source*, Y is the *target*, and A is a *targeting matrix*. If there is an $A \in M_m(\mathbb{F})$ with property \mathcal{P} such that $AX = Y$, we say that A solves the *\mathcal{P} -targeting problem for X and Y* . In this paper, we consider the following properties: invertible, Hermitian, positive (semi)definite, unitary, orthogonal projection, reflection, complex symmetric, and normal.

If $AX = Y$, then $\text{null } X \subseteq \text{null } Y$, so this condition (or something that implies it) is found in all of our results. The invertible linear targeting problem is just the row equivalence problem, which has a solution if and only if $\text{null } X = \text{null } Y$ [1, Theorem 4.4.1].

If $m < n$ and $AX = Y$, then after a simultaneous permutation of the columns of X and Y we may assume that $X = [X_1 \ X_2]$, $Y = [Y_1 \ Y_2]$, $X_1, Y_1 \in M_{m \times k}(\mathbb{F})$, $k \leq m$, and $\text{col } X_2 \subseteq \text{col } X_1$. For example, we could choose $k = \text{rank } X$, but any k between $\text{rank } X$ and m will do. Since there is some $B \in M_{k \times (n-k)}(\mathbb{F})$ such that $X_2 = X_1 B$ [1, Theorem 1.6.21.a], we have $AX_1 = Y_1$, from which it follows that $AX_2 = A(X_1 B) = Y_1 B = Y_2$. Thus, in considering the linear targeting problem $AX = Y$ with $X, Y \in M_{m \times n}(\mathbb{F})$, it suffices to consider only the case $m \geq n \geq 1$.

For any $Z \in M_{m \times n}(\mathbb{F})$ ($m \times n$ matrices over the field \mathbb{R} or \mathbb{C}), Z^* is the *conjugate transpose* of Z and Z^\dagger is its *Moore–Penrose pseudoinverse*; see [1, § 17.4] or [2, 7.3.P7]. The *Euclidean norm* of $\mathbf{x} \in \mathbb{F}^n$ is denoted by $\|\mathbf{x}\| = (\mathbf{x}^* \mathbf{x})^{1/2}$. If $A \in M_n(\mathbb{F})$, we write $A \geq 0$ (respectively, $A > 0$) if $A \in M_n(\mathbb{F})$ is *positive semidefinite* (respectively, *positive definite*), that is, A is Hermitian and $\mathbf{u}^* A \mathbf{u} \geq 0$ (respectively, $\mathbf{u}^* A \mathbf{u} > 0$) for all nonzero $\mathbf{u} \in \mathbb{F}^n$. If $B, S \in M_m(\mathbb{F})$ and S is invertible, then $S^* B S$ is **congruent* to B . If B is, respectively, invertible, Hermitian, or positive semidefinite, then $S^* B S$ has the same respective properties.

2020 *Mathematics Subject Classification.* 15A06, 15A10, 15A24.

Key words and phrases. data mapping, source and target matrices, constrained systems of linear equations.

SRG partially supported by NSF grant DMS-2054002.

2. UNCONSTRAINED LINEAR TARGETING

A linear targeting problem need not have a solution. For example, $AX = Y$ is impossible if

$$X = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad Y = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}.$$

Since $\text{null } X \subseteq \text{null}(AX)$, it is necessary that $\text{null } X \subseteq \text{null } Y$. A computation with pseudoinverses shows that this condition permits us to identify all solutions of an unconstrained linear targeting problem.

Theorem 2.1. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $\text{null } X \subseteq \text{null } Y$.*

- (a) $(YX^\dagger)X = Y$.
- (b) If $Z \in M_m(\mathbb{F})$ and $A = YX^\dagger + Z(I - XX^\dagger)$, then $AX = Y$.
- (c) If $A \in M_m(\mathbb{F})$ and $AX = Y$, then there is a $Z \in M_m(\mathbb{F})$ such that $A = YX^\dagger + Z(I - XX^\dagger)$.

Proof. (a) The hypothesis $\text{null } X \subseteq \text{null } Y$ is equivalent to the condition $\text{col } Y^* \subseteq \text{col } X^*$. Since $X^\dagger X$ is the orthogonal projection (necessarily Hermitian) onto $\text{col } X^*$, we have $X^\dagger X Y^* = Y^*$, and hence $Y = YX^\dagger X = (YX^\dagger)X$.

(b) Compute

$$\begin{aligned} AX &= (YX^\dagger + Z(I - XX^\dagger))X \\ &= YX^\dagger X + Z(X - XX^\dagger X) \\ &= Y + Z(X - X) = Y. \end{aligned}$$

(c) If $AX = Y$, then

$$\begin{aligned} A - YX^\dagger &= (A - YX^\dagger)(I - XX^\dagger + XX^\dagger) \\ &= (A - YX^\dagger)(I - XX^\dagger) + (A - YX^\dagger)XX^\dagger \\ &= (A - YX^\dagger)(I - XX^\dagger) + (AX - (YX^\dagger)X)X^\dagger \\ &= (A - YX^\dagger)(I - XX^\dagger) + (Y - Y)X^\dagger \\ &= (A - YX^\dagger)(I - XX^\dagger) = Z(I - XX^\dagger), \end{aligned}$$

in which $Z = A - YX^\dagger$. □

The following theorem re-examines the unconstrained linear targeting problem in the context of a singular value decomposition of the data matrix X . Its proof introduces notation and relations that we use throughout our discussion. If $X = 0$, the linear targeting problem has a solution if and only if $Y = 0$, so this case is not interesting.

Theorem 2.2. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $X \neq 0$. There is an $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if $\text{null } X \subseteq \text{null } Y$.*

Proof. (\Rightarrow) If $\mathbf{u} \in \text{null } X$, then $Y\mathbf{u} = AX\mathbf{u} = A\mathbf{0} = \mathbf{0}$.

(\Leftarrow) Let $\text{rank } X = r \geq 1$ and let $X = V\Sigma W^*$ be a singular value decomposition, in which $V \in M_m(\mathbb{F})$ and $W \in M_n(\mathbb{F})$ are unitary,

$$\Sigma = \begin{bmatrix} \Sigma_r & 0_{r \times (n-r)} \\ 0_{(m-r) \times r} & 0_{(m-r) \times (n-r)} \end{bmatrix} \in M_{m \times n}(\mathbb{R}), \quad (1)$$

$r \geq 1$, and $\Sigma_r \in M_r(\mathbb{R})$ is diagonal, invertible, and positive definite. Partition $W = [W_1 \ W_2]$ and $V = [V_1 \ V_2]$, in which $W_1 \in M_{n \times r}(\mathbb{F})$ and $V_1 \in M_{m \times r}(\mathbb{F})$. If X has full rank, the block column of zeros in (1) is absent, $W = W_1$, and W_2 is absent. If $r < n$, the columns of W_2 are an orthonormal basis for $\text{null } X$ and the columns of W_1 are an orthonormal basis for $(\text{null } X)^\perp = \text{col } X^*$. Since $\text{null } X \subseteq \text{null } Y$, we have $YW_2 = 0$.

If there is a $B \in M_m$ such that

$$B\Sigma = V^*YW, \quad (2)$$

then $YW = VB\Sigma = (VBV^*)(V\Sigma W^*)W = (VBV^*)XW$ and hence

$$(VBV^*)X = Y.$$

Thus, $A = VBV^*$ solves the unconstrained targeting problem for X and Y . We can partition the right side of (2) in several ways:

$$V^*YW = V^*Y[W_1 \ W_2] = V^*[YW_1 \ YW_2] = [V^*YW_1 \ 0] \quad (3)$$

$$= \begin{bmatrix} V_1^*YW_1 & 0 \\ V_2^*YW_1 & 0 \end{bmatrix} = \begin{bmatrix} Z_1 & 0 \\ Z_2 & 0 \end{bmatrix} = [Z \ 0] \in M_{m \times n}(\mathbb{F}),$$

in which the three matrices $Z = V^*YW_1 \in M_{m \times r}(\mathbb{F})$, $Z_1 = V_1^*YW_1 \in M_r(\mathbb{F})$, and $Z_2 = V_2^*YW_1 \in M_{(m-r) \times r}(\mathbb{F})$ are determined by the data X and Y . Define $B_1 = Z\Sigma_r^{-1} \in M_{m \times r}(\mathbb{F})$ and let

$$B = [B_1 \ B_2] = [Z\Sigma_r^{-1} \ B_2] = \begin{bmatrix} Z_1\Sigma_r^{-1} & R \\ Z_2\Sigma_r^{-1} & S \end{bmatrix}, \quad (4)$$

in which $R \in M_{r \times (m-r)}(\mathbb{F})$ and $S \in M_{m-r}(\mathbb{F})$ are arbitrary; they (and B_2) are absent if X has full rank. Then

$$B\Sigma = [Z\Sigma_r^{-1} \ B_2] \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} = [Z \ 0] = V^*YW,$$

so (2) is satisfied for any choice of B_2 . \square

If a property \mathcal{P} is invariant under unitary similarity, the preceding theorem shows that the \mathcal{P} -targeting problem is equivalent to a matrix completion problem. The source and target matrices determine B_1 ; if we can choose B_2 so that B (and hence also VBV^*) has property \mathcal{P} , then the \mathcal{P} -targeting problem has a solution. For example, if \mathcal{P} is the property that A is invertible, then B must be invertible and hence the data must ensure that B_1 has full rank. The following corollary reveals a familiar criterion for row equivalence of two matrices; see [1, Theorem 4.4.1 and P.8.30].

Corollary 2.3. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $X \neq 0$. There is an invertible $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if $\text{null } X = \text{null } Y$.*

Proof. (\Rightarrow) If $Y = AX$ and A is invertible, then $X = A^{-1}Y$. Consequently, $\text{null } X \subseteq \text{null } Y$ and $\text{null } Y \subseteq \text{null } X$, so $\text{null } X = \text{null } Y$.

(\Leftarrow) Let $r = \text{rank } X \geq 1$. There is a choice of B_2 in (4) that makes B invertible if and only if $\text{rank } B_1 = r$. If $\text{null } X = \text{null } Y$, then $\text{rank } X = \text{rank } Y$, and the relations

$$\text{rank } Y = \text{rank}(V^*YW) = \text{rank}[Z \ 0] = \text{rank } Z = \text{rank}(Z\Sigma_r^{-1}) = \text{rank } B_1 \quad (5)$$

ensure that $\text{rank } B_1 = r$. \square

3. TWO LEMMAS

With the notation in the preceding section, the following lemma memorializes some consequences of the basic assumption $\text{null } X \subseteq \text{null } Y$.

Lemma 3.1. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $X \neq 0$. If $\text{null } X \subseteq \text{null } Y$, then X^*Y is unitarily similar to $\Sigma_r Z_1 \oplus 0_{n-r}$, in which $Z_1 = V_1^* Y W_1$.*

Proof. Compute

$$\begin{aligned} W^*(X^*Y)W &= W^*(W\Sigma V^*)YW = \Sigma(V^*YW) = \Sigma[Z \ 0] \\ &= \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} Z_1 & 0 \\ Z_2 & 0 \end{bmatrix} = \begin{bmatrix} \Sigma_r Z_1 & 0 \\ 0 & 0 \end{bmatrix}. \quad \square \end{aligned} \quad (6)$$

In our computations with block matrices, we require the three types of Schur complements in the next lemma.

Lemma 3.2. *Let $H \in M_r(\mathbb{F})$, $L \in M_{(m-r) \times r}(\mathbb{F})$, and $\lambda \in \mathbb{F}$. Let*

$$B = \begin{bmatrix} H & L^* \\ L & \lambda I_{m-r} \end{bmatrix} \in M_m(\mathbb{F}).$$

- (a) *If $\lambda \in \mathbb{R} \setminus \{0\}$, then B is $*$ -congruent to $(H - \lambda^{-1}L^*L) \oplus (\lambda I_{m-r})$.*
- (b) *If H is Hermitian and invertible, then B is $*$ -congruent to $H \oplus (\lambda I_{m-r} - LH^{-1}L^*)$.*
- (c) *If H is Hermitian and $\text{null } H \subseteq \text{null } L$, then B is $*$ -congruent to $H \oplus (\lambda I_{m-r} - LH^\dagger L^*)$.*

Proof. (a) Compute the $*$ -congruence

$$\begin{bmatrix} I_r & -\lambda^{-1}L^* \\ 0 & I_{m-r} \end{bmatrix} \begin{bmatrix} H & L^* \\ L & \lambda I_{m-r} \end{bmatrix} \begin{bmatrix} I_r & 0 \\ -\lambda^{-1}L & I_{m-r} \end{bmatrix} = \begin{bmatrix} H - \lambda^{-1}L^*L & 0 \\ 0 & \lambda I_{m-r} \end{bmatrix}.$$

(b) The product

$$\begin{bmatrix} I_r & 0 \\ -LH^{-1} & I_{m-r} \end{bmatrix} \begin{bmatrix} H & L^* \\ L & \lambda I_{m-r} \end{bmatrix} \begin{bmatrix} I_r & -H^{-1}L^* \\ 0 & I_{m-r} \end{bmatrix} = \begin{bmatrix} H & 0 \\ 0 & \lambda I_{m-r} - LH^{-1}L^* \end{bmatrix}$$

is a $*$ -congruence since H is Hermitian.

(c) Since $I - H^\dagger H$ is the orthogonal projection onto $\text{null } H$ and $\text{null } H \subseteq \text{null } L$, we have $L(I_{m-r} - H^\dagger H) = 0$ and the product

$$\begin{bmatrix} I_r & 0 \\ -LH^\dagger & I_{m-r} \end{bmatrix} \begin{bmatrix} H & L^* \\ L & \lambda I_{m-r} \end{bmatrix} \begin{bmatrix} I_r & -H^\dagger L^* \\ 0 & I_{m-r} \end{bmatrix} = \begin{bmatrix} H & 0 \\ 0 & \lambda I_{m-r} - LH^\dagger L^* \end{bmatrix}$$

is a $*$ -congruence. □

4. HERMITIAN TARGETING

Theorem 4.1. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $X \neq 0$. There is a Hermitian $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if $\text{null } X \subseteq \text{null } Y$ and X^*Y is Hermitian.*

Proof. (\Rightarrow) If $AX = Y$ and A is Hermitian, then $\text{null } X \subseteq \text{null } Y$ and

$$(X^*Y)^* = (X^*AX)^* = X^*AX = X^*Y.$$

(\Leftarrow) Let $r = \text{rank } X$. It suffices to show that there is a choice of B_2 in (4) such that B is Hermitian. Let

$$B = \begin{bmatrix} Z_1 \Sigma_r^{-1} & (Z_2 \Sigma_r^{-1})^* \\ Z_2 \Sigma_r^{-1} & \lambda I_{m-r} \end{bmatrix} = [B_1 \quad B_2], \quad (7)$$

in which $\lambda \in \mathbb{R}$. Since X^*Y is Hermitian, (6) ensures that $\Sigma_r Z_1$ is Hermitian, which implies that $\Sigma_r^{-*} (\Sigma_r Z_1) \Sigma_r^{-1} = Z_1 \Sigma_r^{-1}$ is Hermitian and hence B is Hermitian. \square

Corollary 4.2. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $X \neq 0$. There is an invertible Hermitian $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if $\text{null } X = \text{null } Y$ and X^*Y is Hermitian.*

Proof. (\Rightarrow) It follows from Theorem 4.1 that X^*Y is Hermitian and $\text{null } X \subseteq \text{null } Y$. Since $X = A^{-1}Y$, it follows that $\text{null } Y \subseteq \text{null } X$, so $\text{null } X = \text{null } Y$.

(\Leftarrow) Let $r = \text{rank } X$. The rank-nullity theorem ensures that $\text{rank } Y = r$ since $\dim \text{null } X = \dim \text{null } Y$. Let

$$B = [B_1 \quad B_2] = \begin{bmatrix} Z_1 \Sigma_r^{-1} & (Z_2 \Sigma_r^{-1})^* \\ Z_2 \Sigma_r^{-1} & \lambda I_{m-r} \end{bmatrix} = \begin{bmatrix} H & L^* \\ L & \lambda I_{m-r} \end{bmatrix}$$

be the Hermitian matrix defined in (7), in which $H = Z_1 \Sigma_r^{-1} \in M_r(\mathbb{F})$ is Hermitian, $L = Z_2 \Sigma_r^{-1} \in M_{(m-r) \times r}(\mathbb{F})$, $B_1 \in M_{m \times r}(\mathbb{F})$, and $\lambda \in \mathbb{R}$ is nonzero. Then (5) ensures that B_1 has full column rank, and hence

$$\text{null } H \cap \text{null}(L^*L) = \text{null } H \cap \text{null } L = \{\mathbf{0}\}. \quad (8)$$

Lemma 3.2.a implies that

$$\det B = (\det(\lambda I_{m-r}))(\det(H - \lambda^{-1}L^*L)) = \lambda^{m-r} \det(H - \lambda^{-1}L^*L).$$

We need to show that there is some real nonzero λ such that $\det(H - \lambda^{-1}L^*L) \neq 0$. Consider the analytic function $f(z) = \det(H - z^{-1}L^*L)$ on $\mathbb{C} \setminus \{0\}$. If $f(\lambda) = 0$ for every real $\lambda \neq 0$, then $f(z) = 0$ for every $z \in \mathbb{C} \setminus \{0\}$. To show that $f(\lambda) \neq 0$ for some real nonzero λ , it suffices to show that $f(i) \neq 0$. If $f(i) = 0$, there is a nonzero $\mathbf{u} \in \mathbb{C}^r$ such that $(H + iL^*L)\mathbf{u} = \mathbf{0}$. However,

$$\begin{aligned} (H + iL^*L)\mathbf{u} = \mathbf{0} &\implies \mathbf{u}^*H\mathbf{u} + i\mathbf{u}^*L^*L\mathbf{u} = \mathbf{u}^*(H + iL^*L)\mathbf{u} = 0 \\ &\implies \text{Im}(\mathbf{u}^*H\mathbf{u} + i\mathbf{u}^*L^*L\mathbf{u}) = 0 \\ &\implies (L\mathbf{u})^*(L\mathbf{u}) = \mathbf{u}^*L^*L\mathbf{u} = 0 \\ &\implies L\mathbf{u} = \mathbf{0} \\ &\implies H\mathbf{u} = (H + iL^*L)\mathbf{u} = \mathbf{0} \\ &\implies \text{null } H \cap \text{null } L \neq \{\mathbf{0}\}, \end{aligned}$$

which contradicts (8). We conclude that B is invertible and Hermitian for some real nonzero λ . \square

If a nonzero $Y \in M_{m \times n}(\mathbb{F})$ is given, what are all the $X \in M_{m \times n}(\mathbb{F})$ such that $Y = AX$ for some Hermitian A ?

Corollary 4.3. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $r = \text{rank } Y \geq 1$. Let $Y = V\Sigma W^*$ be a singular value decomposition, in which*

$$V^*YW = \Sigma = \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} \in M_{m \times n}(\mathbb{R})$$

and $\Sigma_r \in M_r(\mathbb{R})$ is diagonal and positive definite. Let $Z = V^*XW$ and partition

$$Z = \begin{bmatrix} Z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix}$$

conformally to Σ . Let $Z_1 = \begin{bmatrix} Z_{11} \\ Z_{21} \end{bmatrix} \in M_{m \times r}(\mathbb{F})$.

- (a) $X^*Y = Y^*X$ if and only if $Z_{12} = 0$ and $\Sigma_r Z_{11} = Z_{11}^* \Sigma_r$.
- (b) If $Z_{12} = 0$, $\Sigma_r Z_{11} = Z_{11}^* \Sigma_r$, $\text{rank } Z_1 = r$, and $(Z_{21} \text{ null } Z_{11}) \cap \text{col } Z_{22} = \{\mathbf{0}\}$, then there is a Hermitian $A \in M_m(\mathbb{F})$ such that $AX = Y$.
- (c) If $Z_{12} = 0$, $\Sigma_r Z_{11} = Z_{11}^* \Sigma_r$, and Z_{11} is invertible, then for each choice of $Z_{21} \in M_{(m-r) \times r}(\mathbb{F})$ and $Z_{22} \in M_{(m-r) \times (n-r)}(\mathbb{F})$ there is a Hermitian $A \in M_m(\mathbb{F})$ such that $AX = Y$.

Proof. (a) A computation reveals that X^*Y is Hermitian if and only if

$$\begin{aligned} X^*Y = Y^*X &\iff X^*V\Sigma W^* = W\Sigma^T V^*X \\ &\iff (W^*X^*V)\Sigma = \Sigma^T(V^*XW) \\ &\iff \begin{bmatrix} Z_{11}^* & Z_{21}^* \\ Z_{12}^* & Z_{22}^* \end{bmatrix} \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} Z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix} \\ &\iff \begin{bmatrix} Z_{11}^* \Sigma_r & 0 \\ Z_{12}^* \Sigma_r & 0 \end{bmatrix} = \begin{bmatrix} \Sigma_r Z_{11} & \Sigma_r Z_{12} \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

Therefore, X^*Y is Hermitian if and only if $Z_{12} = 0$ and $\Sigma_r Z_{11} = Z_{11}^* \Sigma_r$.

(b) We claim that $\text{null } X \subseteq \text{null } Y$ if and only if $\text{null } Z \subseteq \text{null } \Sigma$. To validate this claim, suppose that $\text{null } X \subseteq \text{null } Y$. Then

$$\begin{aligned} \mathbf{u} \in \text{null } Z &\implies \mathbf{0} = Z\mathbf{u} = (V^*XW)\mathbf{u} = \mathbf{0} \\ &\implies X(W\mathbf{u}) = \mathbf{0} \\ &\implies Y(W\mathbf{u}) = \mathbf{0} \\ &\implies V^*Y(W\mathbf{u}) = \mathbf{0} \\ &\implies \mathbf{0} = (V^*YW)\mathbf{u} = \Sigma\mathbf{u} \\ &\implies \mathbf{u} \in \text{null } \Sigma, \end{aligned}$$

from which we conclude that $\text{null } Z \subseteq \text{null } \Sigma$. The converse follows from a parallel argument. Now assume that $Z_{12} = 0$ and $\Sigma_r Z_{11} = Z_{11}^* \Sigma_r$, which ensures that $X^*Y = Y^*X$. Let $\mathbf{u} = \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} \in \mathbb{F}^n$, in which $\mathbf{u}_1 \in \mathbb{F}^r$. Then $\mathbf{u} \in \text{null } \Sigma$ if and only if $\mathbf{u}_1 = \mathbf{0}$. If $\text{rank } Z_1 = r$ and there is a $\mathbf{u} \in \text{null } Z$ such that $\mathbf{u}_1 \neq \mathbf{0}$, then

$$\mathbf{0} = Z\mathbf{u} = \begin{bmatrix} Z_{11} & 0 \\ Z_{21} & Z_{22} \end{bmatrix} \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} = \begin{bmatrix} Z_{11}\mathbf{u}_1 \\ Z_{21}\mathbf{u}_1 + Z_{22}\mathbf{u}_2 \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ Z_{21}\mathbf{u}_1 + Z_{22}\mathbf{u}_2 \end{bmatrix}.$$

Since $\text{rank } Z_1 = r$, it follows that $Z_{21}\mathbf{u}_1 \neq \mathbf{0}$, which implies that $Z_{22}\mathbf{u}_2 \neq \mathbf{0}$ and hence $(Z_{21} \text{ null } Z_{11}) \cap \text{col } Z_{22} \neq \{\mathbf{0}\}$. We conclude that $\text{null } Z \subseteq \text{null } \Sigma$.

(c) If Z_{11} is invertible, then $\text{rank } Z_1 = r$ and $\text{null } Z_{11} = \{\mathbf{0}\}$, so $(Z_{21} \text{ null } Z_{11}) \cap \text{col } Z_{22} \subseteq Z_{21} \text{ null } Z_{11} = \{\mathbf{0}\}$. \square

5. POSITIVE SEMIDEFINITE TARGETING

We now investigate the positive semidefinite and positive definite targeting problems. An important property of positive semidefinite matrices A is that $A\mathbf{u} = \mathbf{0}$ if and only if $\mathbf{u}^*A\mathbf{u} = 0$; see [1, Corollary 15.1.18] or [2, Observation 7.1.6].

Theorem 5.1. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $X \neq 0$. There is a positive semidefinite $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if $X^*Y \geq 0$ and $\text{null } Y = \text{null}(X^*Y)$.*

Proof. (\Rightarrow) If $AX = Y$ and $A \geq 0$, then $X^*Y = X^*AX$, which is positive semidefinite. To verify the assertion about null spaces, use the fact that $X^*Y \geq 0$ and observe that

$$\begin{aligned} \mathbf{u} \in \text{null}(X^*Y) &\implies (X^*Y)\mathbf{u} = \mathbf{0} \\ &\implies \mathbf{u}^*(X^*Y)\mathbf{u} = 0 \\ &\implies \mathbf{u}^*(X^*AX)\mathbf{u} = 0 \\ &\implies (X\mathbf{u})^*A(X\mathbf{u}) = 0 \\ &\implies A(X\mathbf{u}) = \mathbf{0} \\ &\implies (AX)\mathbf{u} = \mathbf{0} \\ &\implies Y\mathbf{u} = \mathbf{0} \\ &\implies \mathbf{u} \in \text{null } Y. \end{aligned}$$

(\Leftarrow) Let $r = \text{rank } X$. Since X^*Y is Hermitian, we have $\text{null } X \subseteq \text{null}(Y^*X) = \text{null}(X^*Y) = \text{null } Y$, so $\text{null } X \subseteq \text{null } Y$. The hypotheses of Theorem 4.1 are satisfied, and its proof shows that

$$B = \begin{bmatrix} B_1 & B_2 \end{bmatrix} = \begin{bmatrix} Z_1 \Sigma_r^{-1} & (Z_2 \Sigma_r^{-1})^* \\ Z_2 \Sigma_r^{-1} & \lambda I_{m-r} \end{bmatrix} = \begin{bmatrix} H & L^* \\ L & \lambda I_{m-r} \end{bmatrix} \quad (9)$$

is Hermitian for any $\lambda > 0$. The identity (6) and the assumption $X^*Y \geq 0$ imply that $\Sigma_r Z_1 \geq 0$, so $\Sigma_r^{-*}(\Sigma_r Z_1)\Sigma_r^{-1} = Z_1 \Sigma_r^{-1} = H \geq 0$. The hypothesis $\text{null } Y = \text{null}(X^*Y)$, the fact that $H \geq 0$, and (5) ensure that

$$\text{null } H = \text{null } B_1 = \text{null } H \cap \text{null } L = \text{null } H \cap \text{null}(L^*L),$$

which implies that

$$\text{null } H = \text{null } B_1 = \text{null } H \cap \text{null } L \subseteq \text{null } L. \quad (10)$$

Let $\lambda_1 \geq 0$ be the largest eigenvalue of the positive semidefinite matrix $LH^\dagger L^*$. We claim that $B \geq 0$ if $\lambda \geq \lambda_1$. Lemma 3.2.c and (10) ensure that B is * congruent to $H \oplus (\lambda I_{m-r} - LH^\dagger L^*)$, which is positive semidefinite if $\lambda I_{m-r} - LH^\dagger L^* \geq 0$. If $\lambda \geq \lambda_1$ and $\mathbf{x} \in \mathbb{F}^{m-r}$, then

$$\mathbf{x}^*(\lambda I_{m-r} - LH^\dagger L^*)\mathbf{x} = \lambda \mathbf{x}^*\mathbf{x} - \mathbf{x}^*(LH^\dagger L^*)\mathbf{x} \geq (\lambda - \lambda_1)\|\mathbf{x}\|^2 \geq 0,$$

which validates our claim. \square

Corollary 5.2. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $X \neq 0$. There is a positive definite $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if $X^*Y \geq 0$, $\text{null } Y = \text{null}(X^*Y)$, and $\text{rank } Y = \text{rank } X$.*

Proof. (\Rightarrow) If $A \in M_m(\mathbb{F})$ is positive definite and $AX = Y$, then A is invertible and $\text{rank } X = \text{rank } Y$. Moreover, $\text{null } Y = \text{null}(X^*Y)$ and $X^*Y \geq 0$, as in the preceding theorem.

(\Leftarrow) Let $r = \text{rank } X$. Since $\text{rank } Y = \text{rank } X$ and $\text{null } Y = \text{null}(X^*Y)$, it follows that

$$r = \text{rank } Y = \text{rank}(X^*Y) = \text{rank } Z_1 = \text{rank } H.$$

Thus, $H \geq 0$ (as in the preceding theorem), $H \in M_r(\mathbb{F})$ is invertible, and H is positive definite. Lemma 3.2.b ensures that B in (9) is $*$ -congruent to $H \oplus (\lambda I_{n-r} - LH^{-1}L^*)$, which is positive definite if λ is greater than the largest eigenvalue of $LH^{-1}L^*$. \square

Corollary 5.3. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $\text{rank } X = n$. There is a positive definite $A \in M_m$ such that $AX = Y$ if and only if $X^*Y > 0$.*

Proof. (\Rightarrow) If $A \in M_m(\mathbb{F})$ is positive definite and $AX = Y$, then $X^*Y = X^*AX$, which is positive definite since X has full rank.

(\Leftarrow) If $X^*Y > 0$, then $n = \text{rank}(X^*Y) \leq \text{rank } Y$, so $\text{rank } Y = n$ and $\text{null } Y = \{\mathbf{0}\} = \text{null}(X^*Y)$. The preceding corollary ensures that there is a positive definite $A \in M_m$ such that $AX = Y$. \square

6. UNITARY TARGETING

The unitary targeting problem generalizes the familiar fact that, for given $\mathbf{x}, \mathbf{y} \in \mathbb{F}^n$, there is a unitary $U \in M_n(\mathbb{F})$ such that $\mathbf{y} = U\mathbf{x}$ if and only if $\|\mathbf{x}\| = \|\mathbf{y}\|$.

Theorem 6.1. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $X \neq 0$. There is a unitary $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if $X^*X = Y^*Y$.*

Proof. (\Rightarrow) If $Y = AX$ and A is unitary, then $Y^*Y = (AX)^*(AX) = X^*(A^*A)X = X^*IX = X^*X$.

(\Leftarrow) Let $r = \text{rank } X$. If $X^*X = Y^*Y$, then $\text{null } X = \text{null } Y$ [1, Theorem 15.1.9] and (3) ensures that $B = [B_1 \ B_2]$ satisfies (2) with $B_1 = Z\Sigma_r^{-1}$. It suffices to show that B_1 has orthonormal columns, since we may then choose B_2 to make B unitary; see [1, Corollary 6.3.14]. Observe that

$$W^*(Y^*Y)W = (V^*YW)^*(V^*YW) = [Z \ 0]^*[Z \ 0] = \begin{bmatrix} Z^*Z & 0 \\ 0 & 0 \end{bmatrix}.$$

We also have

$$W^*(X^*X)W = \Sigma^*\Sigma = \begin{bmatrix} \Sigma_r^2 & 0 \\ 0 & 0 \end{bmatrix}.$$

Since $X^*X = Y^*Y$, we conclude that $Z^*Z = \Sigma_r^2$ and hence

$$I_r = (Z\Sigma_r^{-1})^*(Z\Sigma_r^{-1}) = B_1^*B_1,$$

that is, B_1 has orthonormal columns. \square

The polar decomposition [1, § 16.3] or [2, § 7.3] provides an alternative approach to the unitary targeting problem. If $X^*X = Y^*Y$ and $Q = (X^*X)^{1/2}$, then $X = U_1Q$ and $Y = V_1Q$, in which $U_1, V_1 \in M_{m \times n}(\mathbb{F})$ have orthonormal columns. If $U = [U_1 \ U_2]$ and $V = [V_1 \ V_2]$ are unitary, then VU^* is unitary and

$$(VU^*)X = (VU^*)U_1Q = V(U^*U_1)Q = [V_1 \ V_2] \begin{bmatrix} I \\ 0 \end{bmatrix} Q = V_1Q = Y.$$

If X and Y do not have the same numbers of columns, there is an interesting story to be told about the identity $X^*X = Y^*Y$. For details, examples, and a historical review, see [3].

7. REFLECTION TARGETING

A matrix $A \in M_m(\mathbb{F})$ is a *reflection* if it is a Hermitian involution ($A = A^*$ and $A^2 = I$), or, equivalently, if it is Hermitian and unitary. We now consider the reflection targeting problem.

Theorem 7.1. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $r = \text{rank } Y \geq 1$. There is a reflection $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if X^*Y is Hermitian and $X^*X = Y^*Y$.*

Proof. (\Rightarrow) If A is a reflection and $AX = Y$, then A is Hermitian and so is $X^*Y = X^*AX$. Since A is unitary, $Y^*Y = (AX)^*AX = X^*A^*AX = X^*IX = X^*X$.

(\Leftarrow) Suppose that $X^*Y = Y^*X$ and $X^*X = Y^*Y$. Let $E = X + Y$ and $F = X - Y$, so that $X = \frac{1}{2}(E + F)$ and $Y = \frac{1}{2}(E - F)$. Let $P = F(F^*F)^\dagger F^*$, which is the (necessarily Hermitian) orthogonal projection onto $\text{col } F$. Define the Hermitian matrix $A = I - 2P$, which is a reflection since

$$A^2 = (I - 2P)^2 = I - 4P + 4P^2 = I - 4P + 4P = I.$$

Observe that $\text{col } E$ is orthogonal to $\text{col } F$ since

$$E^*F = (X + Y)^*(X - Y) = (X^*X - Y^*Y) + (Y^*X - X^*Y) = 0 + 0 = 0.$$

Therefore, $PE = 0$ and $PF = F$. Consequently,

$$\begin{aligned} AX &= \frac{1}{2}(I - 2P)(E + F) = \frac{1}{2}(E + F - 2PE - 2PF) \\ &= \frac{1}{2}(E + F - 0 - 2F) = \frac{1}{2}(E - F) = Y. \quad \square \end{aligned}$$

For a given nonzero $Y \in M_{m \times n}(\mathbb{F})$, the following corollary shows how to construct all the $X \in M_{m \times n}(\mathbb{F})$ such that $Y = AX$ for some reflection A .

Corollary 7.2. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $r = \text{rank } Y \geq 1$. Let $Y = V\Sigma W^*$ be a singular value decomposition, in which*

$$V^*YW = \Sigma = \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} \in M_{m \times n}(\mathbb{R})$$

and $\Sigma_r \in M_r(\mathbb{R})$ is diagonal and positive definite. There is a reflection $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if

$$V^*XW = \begin{bmatrix} U_{11}\Sigma_r & 0 \\ U_{21}\Sigma_r & 0 \end{bmatrix}, \quad (11)$$

in which $U_{11} \in M_r(\mathbb{F})$ is Hermitian and $U_1 = \begin{bmatrix} U_{11} \\ U_{21} \end{bmatrix} \in M_{m \times r}(\mathbb{F})$ has orthonormal columns.

Proof. (\Rightarrow) Partition

$$V^*XW = \begin{bmatrix} Z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix}$$

conformally to Σ . Since X^*Y is Hermitian, Corollary 4.3 ensures that $Z_{12} = 0$ and $\Sigma_r Z_{11} = Z_{11}^* \Sigma_r$. Since $X^*X = Y^*Y$, we have

$$\begin{aligned} W^*X^*XW &= \begin{bmatrix} Z_{11}^* & Z_{21}^* \\ 0 & Z_{22}^* \end{bmatrix} \begin{bmatrix} Z_{11} & 0 \\ Z_{21} & Z_{22} \end{bmatrix} \\ &= \begin{bmatrix} Z_{11}^*Z_{11} + Z_{21}^*Z_{21} & Z_{21}^*Z_{22} \\ Z_{22}^*Z_{21} & Z_{22}^*Z_{22} \end{bmatrix} = \begin{bmatrix} \Sigma_r^2 & 0 \\ 0 & 0 \end{bmatrix} \\ &= W^*Y^*YW. \end{aligned} \tag{12}$$

A comparison of entries in (12) reveals that $Z_{22}^*Z_{22} = 0$, and hence $Z_{22} = 0$. If we partition

$$V^*XW = [Z_1 \ 0],$$

another comparison of entries shows that $Z_1^*Z_1 = \Sigma_r^2$. It follows from the polar decomposition that $Z_1 = U_1\Sigma_r$, in which $U_1 = \begin{bmatrix} U_{11} \\ U_{21} \end{bmatrix} \in M_{m \times r}(\mathbb{F})$ has orthonormal columns. The condition $\Sigma_r Z_{11} = Z_{11}^* \Sigma_r$ ensures that $\Sigma_r U_{11} \Sigma_r = \Sigma_r U_{11}^* \Sigma_r$, so U_{11} is Hermitian.

(\Leftarrow) If V^*XW can be partitioned as in (11), then

$$\begin{aligned} W^*(X^*X)W &= \begin{bmatrix} \Sigma_r U_1^* \\ 0 \end{bmatrix} [U_1 \Sigma_r \ 0] = \begin{bmatrix} \Sigma_r U_1^* U_1 \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} \Sigma_r^2 & 0 \\ 0 & 0 \end{bmatrix} = W^*(Y^*Y)W, \end{aligned}$$

which shows that $X^*X = Y^*Y$. Also,

$$\begin{aligned} W^*(X^*Y)W &= \begin{bmatrix} \Sigma_r U_{11}^* & \Sigma_r U_{21}^* \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} \Sigma_r U_{11}^* \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} \Sigma_r U_{11} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} \end{aligned}$$

is Hermitian, which shows that X^*Y is Hermitian. The preceding theorem now ensures that there is a reflection $A \in M_m(\mathbb{F})$ such that $AX = Y$. \square

One consequence of Theorem 7.1 is that, for given $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$, there is a reflection $A \in M_n(\mathbb{C})$ such that $A\mathbf{x} = \mathbf{y}$ if and only if $\|\mathbf{x}\| = \|\mathbf{y}\|$ and $\mathbf{x}^*\mathbf{y}$ is real; the latter requirement is superfluous if $\mathbb{F} = \mathbb{R}$. In fact, A may be chosen to be a scalar multiple of a Householder matrix in this case. Our proof of Theorem 7.1 constructs a reflection matrix that is a natural analog of a Householder matrix.

8. ORTHOGONAL PROJECTION TARGETING

A computation with the Moore–Penrose pseudoinverse provides a necessary and sufficient condition for there to be a solution to the orthogonal projection targeting problem.

Theorem 8.1. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $Y \neq 0$. There is an orthogonal projection $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if $Y^*X = Y^*Y$.*

Proof. (\Rightarrow) If $Y = AX$, A is Hermitian, and $A^2 = A$, then

$$Y^*Y = (AX)^*(AX) = X^*A^*AX = X^*A^2X = X^*(AX) = X^*Y.$$

(\Leftarrow) If $Y^*X = Y^*Y$, let $A = Y(Y^*Y)^\dagger Y^*$. Then A is Hermitian and

$$A^2 = Y(Y^*Y)^\dagger(Y^*Y)(Y^*Y)^\dagger Y^* = Y(Y^*Y)^\dagger Y^* = A,$$

so A is an orthogonal projection. Since $(Y^*Y)^\dagger(Y^*Y)$ is the orthogonal projection on $\text{col}(Y^*Y)^* = \text{col}(Y^*Y) = \text{col } Y^*$, it follows that $(Y^*Y)^\dagger(Y^*Y)Y^* = Y^*$ and hence $Y = Y(Y^*Y)(Y^*Y)^\dagger$. Therefore,

$$AX = Y(Y^*Y)^\dagger(Y^*X) = Y(Y^*Y)^\dagger(Y^*Y) = Y. \quad \square$$

In the preceding theorem, it is not necessary to assume that $\text{null } X \subseteq \text{null } Y$ because the given hypothesis ensures that

$$\text{null } X \subseteq \text{null}(Y^*X) = \text{null}(Y^*Y) = \text{null } Y.$$

We can use a singular value decomposition to identify all solutions X to the orthogonal projection targeting problem $AX = Y$ for a given Y .

Corollary 8.2. *Let $X, Y \in M_{m \times n}(\mathbb{F})$ with $m \geq n \geq 1$ and $r = \text{rank } Y \geq 1$. Let $Y = V\Sigma W^*$ be a singular value decomposition, in which*

$$V^*YW = \Sigma = \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} \in M_{m \times n}(\mathbb{R})$$

and $\Sigma_r \in M_r(\mathbb{R})$ is diagonal and positive definite. Partition

$$V^*XW = Z = \begin{bmatrix} Z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix}$$

conformally to Σ . There is an orthogonal projection $A \in M_m(\mathbb{F})$ such that $AX = Y$ if and only if

$$V^*XW = \begin{bmatrix} \Sigma_r & 0 \\ Z_{21} & Z_{22} \end{bmatrix},$$

in which $Z_{21} \in M_{(m-r) \times r}(\mathbb{F})$ and $Z_{22} \in M_{(m-r) \times (n-r)}(\mathbb{F})$.

Proof. Compare entries in

$$\begin{aligned} Y^*Y = Y^*X &\iff W\Sigma^\top \Sigma W^* = W\Sigma^\top V^*VZW^* \\ &\iff \Sigma^\top \Sigma = \Sigma^\top Z \\ &\iff \begin{bmatrix} \Sigma_r^2 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} \Sigma_r Z_{11} & \Sigma_r Z_{12} \\ 0 & 0 \end{bmatrix}. \quad \square \end{aligned}$$

9. COMPLEX SYMMETRIC TARGETING

A real symmetric matrix is Hermitian, but a complex symmetric matrix need not even be normal. However, there is an analog of Theorem 4.1 for complex symmetric targeting.

Theorem 9.1. *Let $X, Y \in M_{m \times n}(\mathbb{C})$ with $m \geq n \geq 1$ and $X \neq 0$. There is a complex symmetric $A \in M_m(\mathbb{C})$ such that $AX = Y$ if and only if $\text{null } X \subseteq \text{null } Y$ and $X^\top Y$ is complex symmetric.*

Proof. (\Rightarrow) If $Y = AX$ and A is complex symmetric, then $\text{null } X \subseteq \text{null } Y$, $X^\top AX$ is complex symmetric, and $X^\top Y = X^\top AX = (X^\top Y)^\top$.

(\Leftarrow) Let $\text{rank } X = r \geq 1$ and adopt the notation in the proof of Theorem 2.2. Let $X = V\Sigma W^*$ be a singular value decomposition. If there is a complex symmetric $F \in M_m(\mathbb{C})$ such that $F\Sigma = V^T Y W$, then

$$\overline{V}(F\Sigma)W^* = \overline{V}(V^T Y W)W^* = Y$$

and

$$Y = \overline{V}(F\Sigma)W^* = (\overline{V}FV^*)(V\Sigma W^*) = (\overline{V}FV^*)X.$$

Thus, $A = \overline{V}FV^*$ is complex symmetric and $AX = Y$. If $\text{null } X \subseteq \text{null } Y$, then $YW_2 = 0$ and

$$V^T Y W = [V^T Y W_1 \quad 0].$$

Partition $F = [F_1 \quad F_2]$ with

$$F_1 = \begin{bmatrix} V_1^T Y W_1 \Sigma_r^{-1} \\ V_2^T Y W_1 \Sigma_r^{-1} \end{bmatrix} \in M_{m \times r}(\mathbb{C}).$$

For any choice of $F_2 \in M_{m \times (m-r)}(\mathbb{C})$, we have $F\Sigma = V^T Y W$. Now compute

$$\begin{aligned} W^T (X^T Y) W &= W^T ((V\Sigma W^*)^T Y) W = W^T (\overline{W} \Sigma^T V^T Y) W \\ &= \Sigma^T (V^T Y W) = \Sigma^T [V^T Y W_1 \quad 0] \\ &= \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} V_1^T Y W_1 & 0 \\ V_2^T Y W_1 & 0 \end{bmatrix} = \begin{bmatrix} \Sigma_r V_1^T Y W_1 & 0 \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

If $X^T Y$ is symmetric, then so are $\Sigma_r V_1^T Y W_1$ and $\Sigma_r^{-1} (\Sigma_r V_1^T Y W_1) \Sigma_r^{-1} = V_1^T Y W_1 \Sigma_r^{-1}$. If $G \in M_{m-r}$ is any complex symmetric matrix, then

$$F = \begin{bmatrix} V_1^T Y W_1 \Sigma_r^{-1} & (V_2^T Y W_1 \Sigma_r^{-1})^T \\ V_2^T Y W_1 \Sigma_r^{-1} & G \end{bmatrix}$$

is complex symmetric and $F\Sigma = [V^T Y W_1 \quad 0] = V^T Y W$. Thus, $A = \overline{V}FV^*$ is complex symmetric and $AX = Y$. \square

10. NORMAL TARGETING

What about the normal targeting problem? If $n = 1$ and $\mathbf{x}, \mathbf{y} \in \mathbb{F}^m$ are nonzero, then $\mathbf{x}/\|\mathbf{x}\|$ and $\mathbf{y}/\|\mathbf{y}\|$ have the same norm, so there is a unitary $U \in M_m(\mathbb{F})$ such that $U(\mathbf{x}/\|\mathbf{x}\|) = \mathbf{y}/\|\mathbf{y}\|$. Then $A = (\|\mathbf{y}\|/\|\mathbf{x}\|)U$ is normal and $A\mathbf{x} = \mathbf{y}$.

For $n \geq 2$, however, it is not (yet) clear how to proceed. For a normal $A \in M_m(\mathbb{F})$, we have $A^*A = AA^*$, which is a system of m^2 equations in the entries of A . The equations corresponding to diagonal entries of A^*A and AA^* say that, for each i , the i th row and the i th column of A must have the same Euclidean norm. For example, if $m = 3$, $n = 2$,

$$Y = \begin{bmatrix} 0 & 2 \\ 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad X = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix},$$

and $AX = Y$, then the first two columns of A are determined:

$$A = \begin{bmatrix} 0 & 2 & ? \\ 1 & 0 & ? \\ 0 & 0 & ? \end{bmatrix}.$$

No matter how the third column is chosen, the first column and first row of A cannot have the same norm, so the normal targeting problem for X and Y has no solution. Although the normal targeting problem remains an open problem, the following theorem provides a solution in a special case that provides an alternative solution to the reflection and orthogonal projection targeting problems.

Theorem 10.1. *Let $X, Y \in M_{m \times n}(\mathbb{C})$ and let $\lambda_1, \lambda_2 \in \mathbb{C}$ be distinct.*

- (a) *There is a normal $A \in M_m(\mathbb{C})$ such that the spectrum of A is $\{\lambda, \mu\}$ and $AX = Y$ if and only if $(Y - \lambda X)^*(Y - \mu X) = 0$.*
- (b) *If X, Y, λ , and μ are real and $(Y - \lambda X)^\top(Y - \mu X) = 0$, then there is a real normal $A \in M_m(\mathbb{R})$ such that the spectrum of A is $\{\lambda, \mu\}$ and $AX = Y$.*

Proof. (a) (\Rightarrow) If $AX = Y$, A is normal, and the eigenvalues of A are λ (with multiplicity $p \geq 1$) and μ (with multiplicity $m - p \geq 1$), then there is a unitary $U \in M_m(\mathbb{C})$ and a diagonal $\Lambda = \lambda I_p \oplus \mu I_{m-p}$ such that $A = U\Lambda U^*$. Then

$$\begin{aligned} (Y - \lambda X)^*(Y - \mu X) &= (AX - \lambda X)^*(AX - \mu X) \\ &= X^*(A - \lambda I)^*(A - \mu I)X \\ &= X^*U(\bar{\Lambda} - \bar{\lambda}I)U^*U(\Lambda - \mu I)U^*X \\ &= X^*U \begin{bmatrix} 0_p & 0 \\ 0 & (\bar{\mu} - \bar{\lambda})I_{m-p} \end{bmatrix} \begin{bmatrix} (\lambda - \mu)I_p & 0 \\ 0 & 0_{m-p} \end{bmatrix} U^*X \\ &= X^*U 0_m U^*X = 0. \end{aligned}$$

(\Leftarrow) Suppose that $(Y - \lambda X)^*(Y - \mu X) = 0$. Let $E = Y - \mu X$ and $F = Y - \lambda X$ and observe that $F^*E = 0 = E^*F$, so $\text{col } E$ and $\text{col } F$ are orthogonal subspaces. Moreover,

$$X = (\lambda - \mu)^{-1}(E - F) \quad \text{and} \quad Y = (\lambda - \mu)^{-1}(\lambda E - \mu F).$$

Let $P = E(E^*E)^\dagger E^*$ and $Q = F(F^*F)^\dagger F^*$, which are, respectively, orthogonal projections onto the respective orthogonal subspaces $\text{col } E$ and $\text{col } F$. Let $A = \lambda P + \mu Q$, which is normal with spectrum $\{\lambda, \mu\}$ [1, § 14.9]. Then

$$\begin{aligned} AE &= (\lambda P + \mu Q)E = \lambda PE + \mu QE = \lambda PE = \lambda E, \\ AF &= (\lambda P + \mu Q)F = \lambda PF + \mu QF = \mu QF = \mu F, \end{aligned}$$

and

$$AX = (\lambda - \mu)^{-1}A(E - F) = (\lambda - \mu)^{-1}(AE - AF) = (\lambda - \mu)^{-1}(\lambda E - \mu F) = Y.$$

(b) If X, Y, λ , and μ are real, then the construction in part (a) creates real matrices E, F, P, Q , and A . \square

Let $X, Y \in M_{m \times n}(\mathbb{C})$. An orthogonal projection is a normal matrix with spectrum $\{0, 1\}$. Theorem 10.1 says that there is an orthogonal projection A such that $AX = Y$ if and only if $(Y - X)^*(Y - 0X) = Y^*Y - X^*Y = 0$. This is the condition in Theorem 8.1.

A reflection is a normal matrix with spectrum $\{1, -1\}$. Theorem 10.1 says that there is a reflection A such that $AX = Y$ if and only if $C = (Y + X)^*(Y - X) = 0$. Observe that $C = (Y^*Y - X^*Y) + (X^*Y - Y^*X)$ is the sum of a Hermitian matrix and a skew-Hermitian matrix, and that $C = 0$ if and only if both its Hermitian part and its skew-Hermitian part are zero. This is the condition in Theorem 7.1.

Acknowledgement. We thank Michael A. Dritschel for pointing out that [4, Corollary 1] gives a version of Theorem 5.1 that is valid for bounded linear operators on a Hilbert space.

REFERENCES

- [1] Stephan Ramon Garcia and Roger A. Horn, *Matrix mathematics—a second course in linear algebra*, second ed., Cambridge Mathematical Textbooks, Cambridge University Press, Cambridge, 2023. MR 4574833
- [2] Roger A. Horn and Charles R. Johnson, *Matrix analysis*, second ed., Cambridge University Press, Cambridge, 2013. MR 2978290
- [3] Roger A. Horn and Ingram Olkin, *When does $A^*A = B^*B$ and why does one want to know?*, Amer. Math. Monthly **103** (1996), no. 6, 470–482. MR 1390576
- [4] Zoltán Sebestyén, *Restrictions of positive operators*, Acta Sci. Math. (Szeged) **46** (1983), no. 1-4, 299–301. MR 739047

DEPARTMENT OF MATHEMATICS AND STATISTICS, POMONA COLLEGE, 610 N. COLLEGE AVE., CLAREMONT, CA 91711, USA

Email address: `kebierly@gmail.com`

DEPARTMENT OF MATHEMATICS AND STATISTICS, POMONA COLLEGE, 610 N. COLLEGE AVE., CLAREMONT, CA 91711, USA

Email address: `stephan.garcia@pomona.edu`

URL: <https://stephangarcia.sites.pomona.edu/>

TAMPA, FLORIDA, USA

Email address: `rhorn@math.utah.edu`