

Parafermionic representation of Potts-based cluster chain

Tigran Hakobyan^{1,2,*} and Raffi Varosyan^{1,†}

¹*Yerevan State University, 1 Alex Manoogian Street, Yerevan, 0025, Armenia*

²*Alikhanyan National Laboratory, 2 Alikhanyan br. Street, Yerevan, 0036, Armenia*

The parafermionic representation of cluster chain with $\mathbb{Z}_p \times \mathbb{Z}_p$ symmetry protected topological order is constructed and studied for open and periodic boundary conditions. The resulting Hamiltonian is bilinear in particle operators and splits up into two separate chains composed, respectively, of the standard and time reversal parafermions. It has been demonstrated that each subsystem inherits its own \mathbb{Z}_p component from the symmetry, which corresponds to the parity of all constituent particles. For the open chain, the four invariant parafermions, generating zero-energy modes, are identified. They are located at the edges of each chain. For the closed system with the translational invariance, the boundaries are twisted by the total parafermion parity operator. The structure of edge modes and the space-time symmetry of the system, which depend on the chain length and boundary conditions, are studied in detail.

I. INTRODUCTION

In the last decade, much attention has been paid to the structure and properties of the quantum states of matter at zero temperature. The interest is explained by rapid developments in quantum technologies: quantum computers, quantum teleportation, etc. [1]. The quantum phases have much richer structure compared with the classical ones. According to the Landau theory, the classical phases are characterized by the symmetry of the system, and a change or violation of symmetry leads to a phase transition. Meanwhile, due to the complexity of the wavefunction, the symmetry is not enough in order to describe ultimately the quantum states of matter. The hidden topological properties of the ground state enter into the game. They are characterized by the nonlocal order parameters, might be uncovered also by the appearance of the gapless excitations and anomalous symmetries at the edge.

In more details, consider the quantum matter possessing a unique, gapped ground state (in the bulk) with short-range entanglement. If, in addition, a certain symmetry exists, a single Landau phase splits into a few symmetry-protected topological (SPT) phases [2]. Two states belong to the same SPT phase if they can be transformed to each other without closing the gap and breaking the symmetry on whole path. A ground state which is a product of single-particle states (a product state) belongs to the trivial phase. Important examples exhibiting the SPT phases are the topological insulators with the time reversal symmetry, the spin-Hall state, protected by the time reversal and charge conservation symmetries. Another example is the antiferromagnetic spin-1 Heisenberg chain with the Haldane phase, protected by the $SO(3)$ rotation symmetry, the \mathbb{Z}_2 spin liquids and the chiral spin liquids. In contrast, the quantum states of matter with (pure) topological order exhibit long-range entanglement with specific patterns which are robust against any local perturbation regardless of the symmetry. The topological phases were observed in the conventional superconductors, for the chiral spin and fractional quantum Hall states.

The cluster chain based on the Ising model [3] is the simplest model exhibiting a nontrivial SPT order with the $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry. It has an application in the quantum computers [4] based on the quantum measurements [5]. Recently, the cluster chain is extended to the higher spin case when the trivial phase is described by the p -state Potts model [6–8]. The nontrivial system which possesses the $\mathbb{Z}_p \times \mathbb{Z}_p$ SPT phase. In contrast to the Ising case, the corresponding Hamiltonian is inhomogeneous along the chain but splits into two homogeneous parts composed from the odd and even local terms, which become similar to each other upon the application of the time reversal transformation.

The cluster model is deeply connected with the superconducting Kitaev chain which, in turn, is among the simplest systems demonstrating nontrivial topological order. The system has representation in terms of the lattice spinless Majorana fermions, where the topological order manifests itself as gapless Majorana edge modes [9] which have been recently realised experimentally [10]. The multiple Majorana modes exhibit non-abelian braiding statistics called parastatistics [11]. This property together with a strong gap and robustness of the zero modes against local perturbations makes them a useful tool for the fault-tolerant topological quantum computing. The Jordan-Wigner mapping between the Kitaev chains and cluster models established a deep connection between their topological properties [12]. More precisely, the decoupled Kitaev chains are mapped to cluster-type models with the expanded

* tigran.hakobyan@ysu.am; hakob@yerphi.am

† raffivarosyan88@gmail.com

local stabilisers. In particular, the standard cluster chain corresponds to the two decoupled Kitaev chains which together are equivalent to the Su-Schrieffer-Heeger model [13].

The goal of the current article is to express the Hamiltonian of the Potts-based p -state cluster chain, its useful symmetries, the zero-energy edge modes and other properties in terms of the parafermions. They are specific anyons, which obey the fractional statistics, and might be considered also as an extension of the spinless Majorana fermions. They have larger space of states and can serve as a source for qudits in high-dimensional quantum computers [14] which become popular recently [15]. These quasiparticles can be even more important in quantum computation due to their braiding which allows to perform an entangling quantum gate [16, 17]. In comparison with Majoranas, they subject to a fractional and position-dependent statistics. The lattice parafermions emerge as quasiparticle excitations in the clock models as the Majorana fermions appear in the Ising model [18, 19]. Their protected zero modes appear at the boundaries of certain one-dimensional \mathbb{Z}_p -invariant spin systems [20, 21]. Recent experiments have demonstrated the possibility of the realization of parafermionic modes at the edge of the fractional quantum Hall systems at certain filling factors compatible with the parastatistics [22].

The current article is devoted to the derivation and study of the parafermionic version of the p -state cluster chain with N spins. Similar to the \mathbb{Z}_2 case when it is equivalent by two independent Kitaev chains with Majorana fermions [12], the \mathbb{Z}_p cluster model is also divided into two separate parafermion chains. However, the picture here is more complicated and tangled due to an inhomogeneity between the odd and even local terms in the spin Hamiltonian and complexity of the parafermions. A straightforward application of the spin-to-parafermion map produces a six-particle interaction term in the Hamiltonian which loses its symmetric form. The original symmetric shape is recovered after the inclusion of the time reversal (\mathcal{T}) particle operators. The corresponding Hamiltonian becomes bilinear in particle operators except for the translationally invariant case when an additional factor describing a composite particle appears at the boundary. By composite parafermion we refer to the diagonal generator of the $\mathbb{Z}_p^{\text{odd}} \times \mathbb{Z}_p^{\text{even}}$ symmetry, which protects the SPT phase and is proportional to the total parity of all $2N$ particles. It is a parafermionic analog of the Dirac's chiral gamma matrix and may be considered as a $(2N+1)$ -th particle composed from all parafermions. The composite parafermion appears when the particles are transformed by the lattice reflection (\mathcal{P}) and translation (\mathcal{T}) operations. Being expressed in terms of the parafermions, the cluster model together with the protecting symmetry, zero-energy edge modes and string order operators is divided into two separate parts each described by its own particles. The first (odd) part is encoded by the standard parafermions with the index values $i = 0$ or $i = 3$ modulo 4, and their total parity is related to the $\mathbb{Z}_p^{\text{even}}$ multiplier of the protecting symmetry group. At the same time, the second (even) part is given by the time reversals of the remaining subset of particles with $i = 2, 1 \pmod{4}$, and their total parity describes the $\mathbb{Z}_p^{\text{odd}}$ symmetry. For the open chain, the four independent edge parafermions are discovered which are conserved in time and generate zero-energy modes leading to the p^2 -fold ground-state degeneracy. However, their structure depends on the chain size which is due to difference in symmetries. For odd N , the Hamiltonian is \mathcal{PT} invariant, and the edge particles may be identified with the original parafermions. Meanwhile, for the even N , the Hamiltonian is \mathcal{P} invariant, and at least one of the edge modes must be replaced by the time reversal parafermion. Among the four invariant edge modes, the two might be associated with the odd subsystem, and the remaining two – with even subsystem. The left and right invariants which are responsible for the symmetry fractionalization are expressed in terms of the edge parafermionic modes. The model with periodical boundary conditions, apart from the \mathcal{P} invariance (N is even), possesses an additional $T\mathcal{T}$ (time reversal + lattice translation) symmetry which maps the even and odd subsystems to each other. As a result, the corresponding Hamiltonian becomes twisted by the composite parafermion whose eigenvalues are p -th roots of unity. Being restricted to the invariant sector associated with a specific eigenvalue, it acquires a bilinear form with twisted boundary conditions.

The current article is organized as follows. In Sect. II, the p -state cluster chain with open and periodic boundary conditions is briefly reviewed. In particular, the spin Hamiltonian, its nontrivial SPT phase with the $\mathbb{Z}_p \times \mathbb{Z}_p$ symmetry, the string order parameter, zero-energy boundary modes together with the symmetry fractionalization (for open chains) are described. In Sect. III, using the direct and inverse Fradkin-Kadanoff transformation, the spin representation for the symmetry generators and parafermion bilinears are constructed. Then the parafermionic representation of the Potts-based cluster model is derived for open boundary conditions. The Hamiltonian contains unwanted sixth order terms in particle operators. Sect. IV is devoted to the construction and study of the bilinear representation of the same model and its invariant edge modes based on the space-time symmetries. First, the behavior of parafermions under the \mathcal{P} , \mathcal{T} , and \mathcal{PT} transformations are examined in detail. The joint set of the original and time-reversal particle operators obey an extended exchange relations quite close to the parafermion algebra. Next, the even (unwanted) part of cluster model is formulated in terms of the time reversal parafermions. Then the entire Hamiltonian acquires a bilinear form. The symmetry and structure of the invariant parafermions arising at the chain edges, their dependence on the parity of the chain size, as well as the system factorization into two separate chains has been studied in detail as was described above. In Sect. V, the transformation of the particle operators and their bilinears under the translation is described. Then the parafermionic representations of the closed \mathbb{Z}_p cluster chain and the related string order parameter is constructed, and the space-time symmetries are studied in detail. Finally,

the results are briefly summarised in the last section.

II. CLUSTER CHAIN BASED ON POTTS MODEL

Potts-based cluster chain with periodic boundary conditions. Cluster model is the simplest example which exhibits a nontrivial SPT order [1, 3]. Its ground state is used in the measurement-based quantum computation [4]. The system is unitary equivalent to the noninteracting Ising model. In one dimension, it protects the $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry generated by the spin reflections applied independently on the even and odd sites. In this section, we review an extended cluster chain which corresponds to the SPT version of the p -state Potts model [6–8].

The single-site states of the considered model can be parameterized also by the elements of the cyclic group \mathbb{Z}_p . For $p > 2$, the odd and even local terms in the Hamiltonian are given by different three-spin couplings. So, it is suitable to split up the model into two components:

$$H = H_{\text{odd}} + H_{\text{even}}, \quad (2.1)$$

which describe, respectively, the odd and even subsystems:

$$H_{\text{odd}} = -\frac{1}{2} \sum_{r=1}^L (Z_{2r}^+ X_{2r+1} Z_{2r+2} + Z_{2r} X_{2r+1}^+ Z_{2r+2}^+), \quad (2.2)$$

$$H_{\text{even}} = -\frac{1}{2} \sum_{r=1}^L (Z_{2r-1} X_{2r} Z_{2r+1}^+ + Z_{2r-1}^+ X_{2r}^+ Z_{2r+1}). \quad (2.3)$$

First, we consider the *periodic* boundary conditions with even number of spins: $N = 2L$. Then, the modulo N summation is supposed inside all indexes. Here, p -state extensions ($p = 2, 3, \dots$) of the standard Pauli matrices ($X = \sigma_x, Z = \sigma_z$) are introduced. They obey the following algebra:

$$ZX = \omega XZ, \quad X^p = Z^p = 1, \quad \omega = e^{\frac{2\pi i}{p}}. \quad (2.4)$$

In the standard basis, the matrix X flips the spin cyclically while the Z acts diagonally:

$$\langle n' | Z | n \rangle = \omega^{n-1} \delta_{n'n}, \quad \langle n' | X | n \rangle = \delta_{n'-1, n}, \quad n, n' = 1, \dots, p, \quad (2.5)$$

where the difference in the Kronecker delta is counted modulo p . The extended Pauli operators are unitary but not Hermitian except for $p = 2$ case:

$$X^+ X = Z^+ Z = 1, \quad X^+ = X^{p-1}, \quad Z^+ = Z^{p-1}. \quad (2.6)$$

$\mathbb{Z}_p \times \mathbb{Z}_p$ invariance. From the integrability viewpoint, the cluster Hamiltonian (2.1) has a quite simple structure: it is composed from mutually commuting local stabilizers, which are unitary equivalent to the onsite spin flip operators X_i which produce a trivial, p -state Potts model. Both systems possess a special symmetry, which reveals their topological properties. It is formed by simultaneous cyclic permutations of all spins located separately on the even and odd positions:

$$[H, X_{\text{even}}] = [H, X_{\text{odd}}] = [X_{\text{even}}, X_{\text{odd}}] = 0, \quad (2.7)$$

$$X_{\text{even}} = X_2 X_4 \dots X_{2L}, \quad X_{\text{odd}} = X_1 X_3 \dots X_{2L-1}. \quad (2.8)$$

The elements X_{odd} and X_{even} produce the group $\mathbb{Z}_p \times \mathbb{Z}_p = \mathbb{Z}_p^{\text{odd}} \times \mathbb{Z}_p^{\text{even}}$ (2.6).

Trivial phase: noninteracting Potts model. As was mentioned above, the cluster Hamiltonian (2.1) is obtained from the standard Potts model by the unitary transformation:

$$H = U H_{\text{trivial}} U^+, \quad H_{\text{trivial}} = -\frac{1}{2} \sum_{l=1}^N (X_l + X_l^+). \quad (2.9)$$

The latter can be viewed as an evolution operator generated by the alternating nearest-neighbour Ising (Potts) Hamiltonian in the diagonal basis and respects the $\mathbb{Z}_p \times \mathbb{Z}_p$ symmetry:

$$U |n_1 \dots n_N\rangle = \omega^{\sum_{i=1}^N (-1)^{i-1} n_i n_{i+1}} |n_1 \dots n_N\rangle, \quad [U, X_{\text{even}}] = [U, X_{\text{odd}}] = 0. \quad (2.10)$$

The expressions for the local stabilizers, which constitute the cluster model, depend on the site's parity:

$$UX_{2r}U^+ = Z_{2r-1}X_{2r}Z_{2r+1}^+, \quad UX_{2r+1}U^+ = Z_{2r}^+X_{2r+1}Z_{2r+2}. \quad (2.11)$$

The operator (2.10) has a matrix product structure, rather than represent a disjoint product of local unitaries and, hence, is capable to produce a short-range entanglement when acting on a product state. Note that in quantum computation, this entangler defines a quantum circuit, which is composed from successive product of altering controlled Z quantum gates acting on the adjacent sites (qudits):

$$U = \prod_{\text{odd } l} CZ_{l+1} \prod_{\text{even } l} CZ_{l+1}^+, \quad CZ_{ij}|n_1 \dots n_N\rangle = \omega^{n_i n_j} |n_1 \dots n_N\rangle. \quad (2.12)$$

The trivial system is in disordered phase with a unique, gapped ground state, which is the uniform superposition of all states in the Z -basis:

$$|0\rangle_{\text{trivial}} = p^{-\frac{N}{2}} \sum_{n_1, \dots, n_N} |n_1 \dots n_N\rangle. \quad (2.13)$$

It is the product of the symmetric states $p^{-\frac{1}{2}} \sum_{n=1}^p |n\rangle$. The unitary map (2.10) entangles the trivial state above preserving the symmetry. As a result, the ground state of the shifted Hamiltonian (2.1) acquires additional phase factors in the above superposition giving rise to the decorated domain wall structure:

$$|0\rangle = U|0\rangle_{\text{trivial}} = p^{-\frac{N}{2}} \sum_{n_1, \dots, n_N} \omega^{\sum_{i=1}^N (-1)^{i-1} n_i n_{i+1}} |n_1 \dots n_N\rangle. \quad (2.14)$$

So, this state together with its parent Hamiltonian are in nontrivial topological phase protected by the symmetry $\mathbb{Z}_p \times \mathbb{Z}_p$. Moreover, the powers U^2, U^3, \dots, U^{p-1} produce the Hamiltonians, exhibiting other (nonequivalent) nontrivial SPT phases. This is in agreement with the general classification rule of the SPT phases [23]. In one dimension, they are given by the second cohomology group of the symmetry, which, in fact, is in one-to-one correspondence with the nonequivalent projective representations [24]. In our case, there are exactly $H^2(\mathbb{Z}_p \times \mathbb{Z}_p, U(1)) \cong \mathbb{Z}_p$ different phases.

String order parameter. The trivial ground state is in ferromagnetic order in X -diagonal basis while is disordered in the Z -basis since $X_i|0\rangle_{\text{trivial}} = 1$ while $Z_i|0\rangle_{\text{trivial}} = 0$. Therefore, the vacuum correlations of the products of Pauli X operators on different sites equal unity. Meanwhile, the similar correlations for the Z operators vanish:

$$\begin{aligned} \langle X_i \rangle_{\text{trivial}} &= \langle X_i X_j \rangle_{\text{trivial}} = \langle X_i X_j X_k \rangle_{\text{trivial}} = \dots = 1, \\ \langle Z_i \rangle_{\text{trivial}} &= \langle Z_i Z_j \rangle_{\text{trivial}} = \langle Z_i Z_j Z_k \rangle_{\text{trivial}} = \dots = 0, \end{aligned}$$

where the shortened notation $\langle \dots \rangle := \langle 0 | \dots | 0 \rangle$ is used for the ground-state expectation value.

Since the unitary entangler is diagonal in the Z basis, the second equation above remains valid for the SPT state too. Among the products of X operators, there are special ones whose behaviour under the same entangler is quite simple. They are formed by the strings of m successive operators located on the even or odd sites, $X_i X_{i+2} \dots X_{i+2m-2}$, depending on the parity of i . Any such string acquires two additional Z and Z^+ operators at the ends, which follows from Eq. (2.11). Thus, the SPT ground state is characterized by the two (even and odd) string order parameters defined by the nonzero vacuum expectation value:

$$\langle Z_{2r-1} X_{2r} X_{2r+2} \dots X_{2r+2m-2} Z_{2r+2m-1}^+ \rangle = \langle Z_{2r}^+ X_{2r+1} X_{2r+3} \dots X_{2r+2m-1} Z_{2r+2m} \rangle = 1. \quad (2.15)$$

Here, the presence of the intermediate X operators is essential and distinguishes the string order from the usual ferromagnetic order. They reveal a hidden topological order in the nontrivial SPT phase. The string order parameter of such type was first observed in the spin-1 Heisenberg chain (Haldane chain) [25].

Cluster chain with open boundaries. In contrast to the periodic case, the protecting symmetry does not impose any restriction on the length of the chain with free boundaries. The odd and even components of the cluster Hamiltonian (2.1) with open boundary conditions acquire then the following form:

$$H_{\text{odd}} = -\frac{1}{2} \sum_{r=1}^{L-1} Z_{2r}^+ X_{2r+1} Z_{2r+2} + \text{H. c.}, \quad (2.16)$$

$$H_{\text{even}} = -\frac{1}{2} \sum_{r=1}^{L'} Z_{2r-1} X_{2r} Z_{2r+1}^+ + \text{H. c.} \quad (2.17)$$

Here the spin number of the second Hamiltonian depends on the parity of the total size:

$$L' = \begin{cases} L-1 & \text{for chain with } N = 2L \text{ sites,} \\ L & \text{for chain with } N = 2L + 1 \text{ sites.} \end{cases} \quad (2.18)$$

Respectively, the summation limits both in the noninteracting Potts Hamiltonian (2.9) and unitary entangler (2.10) must be corrected:

$$H_{\text{trivial}} = -\frac{1}{2} \sum_{l=2}^{N-1} (X_l + X_l^+), \quad U|n_1 \dots n_N\rangle = \omega^{\sum_{l=1}^{N-1} (-1)^l n_l n_{l+1}} |n_1 \dots n_N\rangle. \quad (2.19)$$

So, the ground state of the cluster model with free boundaries slightly changes comparing to the closed case (2.14):

$$|0\rangle = p^{-\frac{N}{2}} \sum_{n_1, \dots, n_N} \omega^{\sum_{l=1}^{N-1} (-1)^l n_l n_{l+1}} |n_1 \dots n_N\rangle. \quad (2.20)$$

Note that in contrast to the periodic case (2.14), the boundary spins n_1 and n_N may be excluded from the summation procedure above leaving them as p^2 parameters the above state depends on: $|0\rangle \rightarrow |0\rangle_{n_1 n_N}$. The reason of this degeneracy is the left Z_1, X_1 and right Z_N, X_N boundary operators which do not contribute to the trivial Hamiltonian (2.19). The edge X operators pivot under the unitary entangler but unlike to the bulk case (2.11), they wear only a single nearby Z operator:

$$UX_1U^+ = X_1Z_2, \quad UX_NU^+ = Z_{N-1}^\mp X_N = \begin{cases} Z_{N-1}X_N & \text{for even } N, \\ Z_{N-1}^+X_N & \text{for odd } N. \end{cases} \quad (2.21)$$

Thus, the cluster model has a boundary symmetry generated by the left and right edge operators which can be set, respectively, as follows:

$$L_{\text{even}} = Z_1^+, \quad L_{\text{odd}} = X_1Z_2, \quad \text{and} \quad R_{\text{even}} = Z_{N-1}^\mp X_N, \quad R_{\text{odd}} = Z_N^+. \quad (2.22)$$

They are responsible for the complete spontaneous symmetry breaking for open boundaries. Note that the odd part of this symmetry (2.8) now is generated by the element $X_{\text{odd}} = X_1X_3 \dots X_{2L'-1}$. Moreover, when applied to the ground state, the $\mathbb{Z}_p \times \mathbb{Z}_p$ generators are reduced to the edge symmetries:

$$X_{\text{odd}}|0\rangle = L_{\text{odd}}R_{\text{odd}}|0\rangle, \quad X_{\text{even}}|0\rangle = L_{\text{even}}R_{\text{even}}|0\rangle, \quad (2.23)$$

which can be verified using the stabilizer structure of the cluster model. All states of the p^2 fold degenerate vacuum are obtained from the $X_{\text{odd}}^n X_{\text{even}}^m$ monomial action to the ground state with $0 \leq n, m \leq p-1$.

The left (right) generators (2.22) taken separately do not commute any more. Instead, they obey the generalized Pauli algebra (2.4) which may be regarded as a projective representation of the $\mathbb{Z}_p \times \mathbb{Z}_p$ symmetry which protects topological phases. This phenomenon is known as the symmetry fractionization. Its equivalence class is given by the aforementioned second cohomology coinciding with a single \mathbb{Z}_p group whose elements characterise different SPT phases.

Note that for the two-dimensional cluster model defined on triangular lattice, the gapless excitations live inside the boundary spin chain which has more complex structure than the bulk system. Its local terms do not commute mutually any more [26]. In the Potts case even some numerical studies are needed in order to reveal the low-energy behaviour [27].

III. INHOMOGENEOUS PARAFERMION CLUSTER CHAIN

Lattice parafermions. The parafermions are anyons obeying the fractional exchange statistics (parastatistics), which is intermediate between bosons and fermions. They generalize the Majorana fermions and, like anyons, exist only in one and two dimensions. The lattice parafermions are described by the operators, which obey the following algebra:

$$\chi_i^+ \chi_i = 1, \quad \chi_i^p = 1, \quad \chi_i \chi_j = \omega \chi_j \chi_i \quad \text{with } i < j, \quad (3.1)$$

where the fractional exchange phase factor ω is defined before (2.4) for an integer $p \geq 2$.

For $p = 2$, we use the conventional notation for the particle operators: $\chi = \gamma$. In this case a particle coincides with its own quasiparticle, $\gamma = \gamma^+$, the exchange statistics is fermionic, $\omega = -1$, and the parafermions are reduced to the Majorana (real) fermions. Two such fermions constitute a single Dirac (complex) fermion described by creation-annihilation operators: $c^\pm = \frac{1}{2}(\gamma_1 \pm i\gamma_2)$. A similar description exists in case of more general fractional values of the parameter ω , and the related particles are called Fock parafermions [28, 29]. However, for $p > 2$ the construction becomes polynomial and more complicate. The exchange relation between a particle and antiparticle follows from the standard commutation rules for parafermions (3.1):

$$\chi_i \chi_j^+ = \omega^{-1} \chi_j^+ \chi_i, \quad \chi_i^+ \chi_j = \omega^{-1} \chi_j^+ \chi_i, \quad i < j. \quad (3.2)$$

Fradkin-Kadanoff transformation. The lattice parafermions may be expressed in terms of the onsite spins, represented in terms of the generalized Pauli matrices (2.4), by the Fradkin-Kadanoff transformation which extends the well-known Jordan-Wigner transformation to the particles with fractional statistics [18]:

$$\chi_{2l-1} = Z_l \prod_{k < l} X_k, \quad \chi_{2l} = \epsilon \chi_{2l-1} X_l = Y_l \prod_{k < l} X_k. \quad (3.3)$$

Therefore, there are two different parafermions associated to each lattice site, and the total number of these particles equals $2N$. Here, a p -dimensional analog of the Pauli matrix σ_y is introduced:

$$Y = \epsilon Z X, \quad \epsilon = \omega^{\frac{p-1}{2}} = -\omega^{-\frac{1}{2}}. \quad (3.4)$$

It constitutes the following algebra together with the previously defined X, Y operators (2.4):

$$ZY = \omega YZ, \quad YX = \omega XY, \quad Y^p = 1.$$

Note that the Fradkin-Kadanoff transformation (3.3) preserves the unitarity: all operators presented there are unitary. The complexity increases with the growth of the index value. The first few parafermions have the simplest expressions:

$$\chi_1 = Z_1, \quad \chi_2 = Y_1, \quad \chi_3 = X_1 Z_2, \quad \chi_4 = X_1 Y_2. \quad (3.5)$$

The inverse transformation also can be derived. As a result, the lattice spins are reexpressed in terms of the parafermions as follows:

$$X_l = \epsilon^{-1} \chi_{2l-1}^+ \chi_{2l}, \quad (3.6)$$

$$Z_l = \epsilon^{l-1} \chi_{2l-1}^+ \chi_{2l-2}^+ \chi_{2l-3}^+ \chi_{2l-4}^+ \cdots \chi_2^+ \chi_1, \quad (3.7)$$

$$Y_l = \epsilon^{l-1} \chi_{2l} \chi_{2l-2}^+ \chi_{2l-3}^+ \chi_{2l-4}^+ \cdots \chi_2^+ \chi_1. \quad (3.8)$$

The simplest representation has the element X_l which is proportional to the parity operator of two neighbouring parafermions [16, 30, 31]. The Z_l is related to the total parity of $2l - 1$ successive particles starting from the first.

$\mathbb{Z}_p \times \mathbb{Z}_p$ symmetry generators via parafermions. The first formula in above series immediately provides a parafermionic representation for the symmetry generators of the cluster model (2.8):

$$X_{\text{even}} = \epsilon^{-L} \chi_3^+ \chi_4 \chi_7^+ \chi_8 \cdots \chi_{4L-1}^+ \chi_{4L}, \quad (3.9)$$

$$X_{\text{odd}} = \epsilon^{-L'} \chi_1^+ \chi_2 \chi_5^+ \chi_6 \cdots \chi_{4L'-1}^+ \chi_{4L'},$$

where the L' depends on the length parity of the spin chain (2.18). Their combination generates the diagonal \mathbb{Z}_p subgroup expressed as an altering product over all parafermions:

$$\chi_{2N+1} := X_{\text{odd}} X_{\text{even}} = \epsilon^{-N} \chi_1^+ \chi_2 \chi_3^+ \chi_4 \cdots \chi_{2N-1}^+ \chi_{2N}. \quad (3.10)$$

It obeys a simple commutation relation with parafermions as easy to verify using the algebra (4.12):

$$\chi_i \chi_{2N+1} = \omega \chi_{2N+1} \chi_i, \quad 1 \leq i \leq 2N. \quad (3.11)$$

The element (3.10), which describes the total parafermion parity, resembles a new, composite parafermion added to the system. Therefore, it can be included in the parafermion algebra (3.1) as the last member:

$$\chi_i^p = 1, \quad \chi_i \chi_j = \omega \chi_j \chi_i \quad \text{where} \quad 1 \leq i < j \leq 2N + 1. \quad (3.12)$$

In the Ising case when $p = 2$, the particles are reduced to the Majorana fermions being equivalent to the $2N$ Euclidean gamma matrices. Then the introduced element (3.10) will also be real and obey the same anticommutational relations with all primary gamma matrices, playing the role of the γ^5 in the four-dimensional Minkowski space.

It is worth noticing that the inverse Fradkin-Kadanoff representations for Z_l and Y_l spin operators may be written in terms of the composite parafermion χ_{2N+1}^+ . Then the sting of the particle operators starts from the end but not from the beginning. It is easy to check that the following relation holds:

$$Z_l = \epsilon^{l-N-1} \chi_{2l}^+ \chi_{2l+1}^+ \chi_{2l+2}^+ \cdots \chi_{2N-1}^+ \chi_{2N}^+ \chi_{2N+1}^+. \quad (3.13)$$

This representation is particularly convenient for the spin- Z operators close to the end of the chain. For the last two spins, it gives the simplest expressions:

$$Z_N = \epsilon^{-1} \chi_{2N}^+ \chi_{2N+1}^+, \quad Z_{N-1} = \omega \chi_{2N-2}^+ \chi_{2N-1}^+ \chi_{2N}^+ \chi_{2N+1}^+. \quad (3.14)$$

Spin representations of parafermion bilinears. The Fradkin-Kadanoff transformation (3.3) is essentially nonlocal and may affect on the topological properties of the system. Moreover, the bilinear operators of the type $\chi_i^+ \chi_j$ with $i < j$, being expressed in terms of the spin operators, contain the product of all intermediate spin- X matrices between the two sites equipped by other matrices. In particular, for even i and odd j the following spin representation for the aforementioned bilinear expression is valid:

$$\chi_{2l}^+ \chi_{2(l+m)-1} = \epsilon Z_l^+ X_{l+1} X_{l+2} \cdots X_{l+m-1} Z_{l+m}. \quad (3.15)$$

For the remaining values of i, j indices, the string is bounded by other Pauli matrices:

$$\begin{aligned} \chi_{2l}^+ \chi_{2(l+m)} &= \epsilon Z_l^+ X_{l+1} X_{l+2} \cdots X_{l+m-1} Y_{l+m}, \\ \chi_{2l-1}^+ \chi_{2(l+m)-1} &= \epsilon W_l X_{l+1} X_{l+2} \cdots X_{l+m-1} Z_{l+m}, \\ \chi_{2l-1}^+ \chi_{2(l+m)} &= \epsilon W_l X_{l+1} X_{l+2} \cdots X_{l+m-1} Y_{l+m}. \end{aligned} \quad (3.16)$$

Here we have introduced an onsite spin operator

$$W = \epsilon^{-1} Z^+ X, \quad (3.17)$$

which is a time-reversal version of the second generalized Pauli matrix (see Eq. (4.1) below). It is easy to see that it obeys the following relations:

$$W^p = 1, \quad ZW = \omega WZ, \quad XW = \omega WX, \quad YW = \omega^2 WY = \omega X^2. \quad (3.18)$$

Note that in $p = 2$ case with the usual Pauli matrixes, the new definition does not make sense since $W = -Y$. For $p \geq 3$, the four Pauli matrices can be considered together [32]. With the applied notations in Eqs. (3.16), each lattice site in the string is represented by a single spin operator. In all equations (3.15), (3.16), the restriction $m \geq 1$ is supposed except for the last equation, which holds also for the $m = 0$ case when it reduces to the parafermion representation of the X_l (3.6). Note that the quantities (3.15), (3.16) differ from the string order operators of the Potts-based cluster model, in which the product of the X matrices is taken over the lattice sites with a specific parity (2.15).

Consider now a few simplest cases of above bilinear operators. Note that for $m = 1$, the intermediate matrices are absent in Eqs. (3.16), (3.15), and we obtain:

$$\chi_{2l}^+ \chi_{2l+1} = \epsilon Z_l^+ Z_{l+1}, \quad (3.19)$$

$$\chi_{2l}^+ \chi_{2l+2} = \epsilon Z_l^+ Y_{l+1} = \omega^{-1} Z_l^+ Z_{l+1} X_{l+1}, \quad (3.20)$$

$$\chi_{2l-1}^+ \chi_{2l+1} = \epsilon W_l Z_{l+1} = Z_l^+ Z_{l+1} X_l, \quad (3.21)$$

$$\chi_{2l-1}^+ \chi_{2l+2} = \epsilon W_l Y_{l+1} = \epsilon Z_l^+ Z_{l+1} X_l X_{l+1}. \quad (3.22)$$

Next, for $m = 2$, the spin representations contain a single intermediate X operator, and we arrive at the following expressions:

$$\chi_{2l-2}^+ \chi_{2l+1} = \epsilon Z_{l-1}^+ X_l Z_{l+1}, \quad (3.23)$$

$$\chi_{2l-2}^+ \chi_{2l+2} = \epsilon Z_{l-1}^+ X_l Y_{l+1} = \omega^{-1} Z_{l-1}^+ Z_{l+1} X_l X_{l+1}, \quad (3.24)$$

$$\chi_{2l-3}^+ \chi_{2l+1} = \epsilon W_{l-1} X_l Z_{l+1} = Z_{l-1}^+ Z_{l+1} X_{l-1} X_l, \quad (3.25)$$

$$\chi_{2l-3}^+ \chi_{2l+2} = \epsilon W_{l-1} X_l Y_{l+1} = \epsilon Z_{l-1}^+ Z_{l+1} X_{l-1} X_l X_{l+1} \quad (3.26)$$

where the index substitution $l \rightarrow l - 1$ is applied in order to get more symmetric expressions.

Parafermion representation of Potts-based cluster model with open boundaries. Now let us focus on the parafermionic representation of the the cluster Hamiltonian (2.1). In this section, we consider the free system (2.16), (2.17), (2.18). Other cases will be treated in the below in this article.

First note that Eq. (3.23) encodes the terms from the odd part of the model (2.16) in terms of the parafermions:

$$Z_{l-1}^+ X_l Z_{l+1} = \epsilon^{-1} \chi_{2l-2}^+ \chi_{2l+1}. \quad (3.27)$$

Then, using also Eq. (3.6) and the particle exchange relations (3.1), (3.2), one can write the parafermionic representation for the even stabilisers in the Hamiltonian (2.17):

$$Z_{l-1} X_l Z_{l+1}^+ = (Z_{l-1}^+ X_l Z_{l+1})^+ X_l^2 = \epsilon^{-1} \chi_{2l+1}^+ \chi_{2l-2} (\chi_{2l-1}^+ \chi_{2l})^2 = \epsilon^{-5} \chi_{2l-2} \chi_{2l-1}^{+2} \chi_{2l}^2 \chi_{2l+1}^+. \quad (3.28)$$

As a result, the generalized cluster Hamiltonian (2.1) with open boundary conditions (2.16) and even number of sites ($N = 2L$), being expressed in terms of the parafermions, acquires the following structure:

$$H = -\frac{1}{2\epsilon} \sum_{r=1}^{L-1} (\chi_{4r}^+ \chi_{4r+3} + \omega^2 \chi_{4r-2} \chi_{4r-1}^{+2} \chi_{4r}^2 \chi_{4r+1}^+) + \text{H. c.} \quad (3.29)$$

It consists of two commuting Hamiltonians, described by the first and second local term, respectively. The same model defined on an odd-length chain ($N = 2L + 1$) contains two more terms of the second type, namely the monomial

$$-\epsilon^{-5} \chi_{4L+1} \chi_{4L+3}^{+2} \chi_{4L+4}^2 \chi_{4L+5}^+ \quad (3.30)$$

and its Hermitian conjugate.

A straightforward application of the generalized Jordan-Wigner transform complicates the physical interpretation of the resulting system (3.29). Indeed, the first part of the model is bilinear in parafermions and may be described as free particles. However, in general, the second part drastically differs. It consists of sixth-order monomials which may be regarded as a $(\chi^+ \chi)^3$ -type interaction among particles located at the four neighbouring positions. Recall that in the original spin system both components contain three-spin interactions among the three neighbouring sites, which only slightly differ from each other (2.2), (2.3). Let us consider the two particular values of the exchange phase parameter when the above parafermion system simplifies.

For the $p = 2$ case, the parafermions are reduced to the Majorana fermions γ_i :

$$\epsilon = \iota, \quad \gamma_i = \chi_i : \quad \{\gamma_i, \gamma_j\} = 2\delta_{ij}, \quad (3.31)$$

and the Hamiltonian (3.29) drastically simplifies. The particle operators become self-conjugate, $\gamma^+ = \gamma$, and their squares in the interaction terms disappear. As a result, the even subsystem becomes also a bilinear combination of particle operators, and the entire Hamiltonian reduces to two decoupled Kitaev chains:

$$H = \iota \sum_{l=1}^{N-1} \gamma_{2l} \gamma_{2l+3}. \quad (3.32)$$

The above system coincides with the $\alpha = 2$ chain in classification scheme applied for the fermionic representation of the generalized cluster models [12].

In case of the $p = 3$ state parafermions, the fractional exchange phase takes the value $\epsilon = \omega = e^{\frac{2\pi i}{3}}$, and, due to the relation $\chi^2 = \chi^+$, the members of the H_{even} are reduced to a four-particle interaction of the type $(\chi^+ \chi)^2$. Thus, the Hamiltonian slightly simplifies:

$$H = -\frac{1}{2} \sum_{r=1}^{L-1} (\omega^{-1} \chi_{4r}^+ \chi_{4r+3} + \omega \chi_{4r-2} \chi_{4r-1} \chi_{4r}^+ \chi_{4r+1}^+) + \text{H. c.} \quad (3.33)$$

Note that a four-particle interaction term of similar type arose recently in the extended parafermion chain [33].

In the next section, the time-reversal particle operators will be used in order to present the parafermionic system (3.29) in a bilinear form similar to the Majorana-fermion chain (3.33), which will be more amenable for revealing the symmetries and edge zero-energy modes.

IV. SPLITTING INTO TWO SIMILAR CHAINS WITH STANDARD AND TIME REVERSAL PARAFERMIONS: OPEN BOUNDARIES

In this section, we derive an alternative, bilinear representation of the p -state open cluster model which involves the time reversal parafermions. The symmetries and parafermionic edge modes which elicit the nontrivial SPT phase and are responsible for the ground-state degeneracy are studied. First, let us reveal the behavior of particle operators under the time reversal and lattice parity transformations.

Time reversal. For spinless particles, the time reversal transformation is reduced to the complex conjugation of the wave function ψ . It is the simplest antiunitary involutive operation:

$$\mathcal{T}\psi = \psi^*, \quad \mathcal{T}^2 = 1.$$

Under the time reversal, the extended Pauli matrices (2.4), (3.4), and (3.17) map to each other [34]:

$$\mathcal{T}X\mathcal{T}^{-1} = X, \quad \mathcal{T}Z\mathcal{T}^{-1} = Z^+, \quad \mathcal{T}Y\mathcal{T}^{-1} = W. \quad (4.1)$$

Therefore, the $\mathbb{Z}_p \times \mathbb{Z}_p$ group generated by the elements (2.8) is \mathcal{T} -invariant.

In the following we will use the barred symbols (instead of the asterisk superscript) in order to denote the complex conjugate operators:

$$\bar{\chi} := \chi^* = \mathcal{T}\chi\mathcal{T}^{-1}. \quad (4.2)$$

The barred parafermions obey the inverse statistics (with the fractional phase factor ω replaced by ω^{-1}) compared with the initial one (3.1) [35]:

$$\bar{\chi}_i^p = 1, \quad \bar{\chi}_i\bar{\chi}_j = \omega^{-1}\bar{\chi}_j\bar{\chi}_i, \quad i < j. \quad (4.3)$$

Using Eqs. (4.1), the time reversal version of the Fradkin-Kadanoff transformation (3.3) is easily obtained:

$$\bar{\chi}_{2l-1} = Z_l^+ \prod_{k<l} X_k, \quad \bar{\chi}_{2l} = W_l \prod_{k<l} X_k. \quad (4.4)$$

The corresponding bilinear operators are just the complex conjugations of the corresponding expressions (3.15), (3.16):

$$\begin{aligned} \bar{\chi}_{2l}^+ \bar{\chi}_{2(l+m)-1} &= \epsilon^{-1} Z_l X_{l+1} \dots X_{l+m-1} Z_{l+m}^+, \\ \bar{\chi}_{2l}^+ \bar{\chi}_{2(l+m)} &= \epsilon^{-1} Z_l X_{l+1} \dots X_{l+m-1} W_{l+m}, \\ \bar{\chi}_{2l-1}^+ \bar{\chi}_{2(l+m)-1} &= \epsilon^{-1} Y_l X_{l+1} \dots X_{l+m-1} Z_{l+m}^+, \\ \bar{\chi}_{2l-1}^+ \bar{\chi}_{2(l+m)} &= \epsilon^{-1} Y_l X_{l+1} \dots X_{l+m-1} W_{l+m}, \end{aligned} \quad (4.5)$$

Mixture of the two sets of parafermions, χ and $\bar{\chi}$, obeys a generalized parastatistics with different values of exchange constant among the particle pairs. Namely, in most cases when the particle indexes differ, they obey a simple homogeneous commutation rule:

$$\chi_i \bar{\chi}_j = \omega \bar{\chi}_j \chi_i, \quad i \neq j. \quad (4.6)$$

An exception occurs when we exchange the particle with its own time reversal, and the result depends on the parity of the particle's index:

$$\chi_{2l} \bar{\chi}_{2l} = \omega^2 \bar{\chi}_{2l} \chi_{2l}, \quad \chi_{2l-1} \bar{\chi}_{2l-1} = \bar{\chi}_{2l-1} \chi_{2l-1}. \quad (4.7)$$

In the particular case of the Majorana fermions (3.31), things are simplified drastically. Remember that Majoranas are referred to as real particles coinciding with their antiparticles. Half of them alter the sign under the time reversal while the rests are not changed at all:

$$\bar{\gamma}_i = (-1)^{i-1} \gamma_i. \quad (4.8)$$

Note that the above equation together with Eq. (4.7) are in agreement with the fact that the Majorana operator is involutive: $\gamma_i^2 = 1$.

Clearly, the time reversal particles are not independent but expressed in terms of the original ones. The connection among them can be obtained from the following relations, which are implied by the Fradkin-Kadanoff transformation:

$$\bar{\chi}_{2l-1} = Z_l^{+2} \chi_{2l-1}, \quad \bar{\chi}_{2l} = \omega Z_l^{+2} \chi_{2l}. \quad (4.9)$$

Applying further the inverse transformation for the Pauli Z matrices (3.7), one can get the final expressions for the time-reversal particle operators. In particular, the equations (4.9) result to an equivalence among the respective bilinear terms:

$$\bar{\chi}_{2l-1}^+ \bar{\chi}_{2l} = \omega \chi_{2l-1}^+ \chi_{2l}. \quad (4.10)$$

Finally, we mention that the $\mathbb{Z}_p \times \mathbb{Z}_p$ group, which produces the SPT phases and is generated by the elements (2.8), is \mathcal{T} -invariant because its generators are composed from the \mathcal{T} -invariant operators (4.1). Hence, the composite parafermion (3.10) also remains unchanged:

$$\bar{\chi}_{2N+1} = \chi_{2N+1}. \quad (4.11)$$

Parity transformation. The lattice parity transformation flips the lattice nodes with respect to the middle of the chain. More explicitly, the onsite Pauli matrices transform as follows:

$$\mathcal{P} X_l \mathcal{P} = X_{l'}, \quad \mathcal{P} Z_l \mathcal{P} = Z_{l'}, \quad l' = N + 1 - l, \quad (4.12)$$

where by l' the reflected site is denoted. Clearly, it is involutive and consistent with the time reversal transformation:

$$\mathcal{P}^2 = 1, \quad [\mathcal{T}, \mathcal{P}] = 0. \quad (4.13)$$

From Eq. (4.12) it is easy to see that the $\mathbb{Z}_p \times \mathbb{Z}_p$ generators remain invariant or are mapped to each other depending on the parity of the lattice size:

$$\begin{aligned} N = 2L : \quad & \mathcal{P} X_{\text{even}} \mathcal{P} = X_{\text{odd}}, \\ N = 2L + 1 : \quad & [\mathcal{P}, X_{\text{even}}] = [\mathcal{P}, X_{\text{odd}}] = 0. \end{aligned}$$

In both cases, the combined symmetry operator (3.10) remains invariant with respect to the space parity transformation:

$$\mathcal{P} \chi_{2N+1} \mathcal{P} = \chi_{2N+1}. \quad (4.14)$$

The transformation rules of the basic particle operators can be derived using the Fradkin-Kadanoff mapping (3.3). Indeed, using the reflection rules of the spin operators (4.12), their algebra and transposes: $X^\tau = X^{-1}$, $Z^\tau = Z$, one obtain the following rules:

$$\begin{aligned} \mathcal{P} \chi_{2l} \mathcal{P} &= \varepsilon Z_{l'} \prod_{k=l'}^{2N} X_k = \varepsilon Z_{l'} \left(\prod_{k=1}^{l'-1} X_k^\tau \right) \chi_{2N+1} = \varepsilon \chi_{2l'-1}^\tau \chi_{2N+1}, \\ \mathcal{P} \chi_{2l-1} \mathcal{P} &= Z_{l'} \prod_{k=l'+1}^{2N} X_k = \omega^{-1} \left(\prod_{k=1}^{l'} X_k^\tau \right) Z_{l'} \chi_{2N+1} = \varepsilon \chi_{2l'}^\tau \chi_{2N+1}. \end{aligned} \quad (4.15)$$

Recall that the operator $\chi_{2N+1} = X_1 X_2 \dots X_N$, which generates the diagonal \mathbb{Z}_p symmetry, represents also a composite parafermion given by the altering product of all particle operators (3.10). It appears here due to the nonlocal nature of the Fradkin-Kadanoff transformation (3.3). Note that the second equation in (4.15) follows from the first one and vice versa as is easy to verify using the algebra (4.13). Finally, notice that the above two equations may be combined into a single formula fulfilled for odd and even values of indexes:

$$\mathcal{P} \chi_i \mathcal{P} = \varepsilon \chi_{2N+1-i}^\tau \chi_{2N+1}, \quad 1 \leq i \leq 2N. \quad (4.16)$$

Consider the restriction of the relation (4.16) to the Majorana fermions (3.31). They are described by real operators, so their transposes coincide with the time reversals: $\gamma_i^\tau = \bar{\gamma}_i = (-1)^{i-1} \gamma_i$. Then, the Majorana time reversal rule (4.8) implies:

$$\mathcal{P} \gamma_i \mathcal{P} = \imath (-1)^{i-1} \gamma_{2N+1-i} \gamma_{2N+1}. \quad (4.17)$$

PT transformation. Combining the relations (4.2) and (4.16), one obtains the joint action of the time reversal and lattice parity transformations on the particle operators:

$$\mathcal{PT}\chi_i\mathcal{PT} = \epsilon^{-1}\chi_{2N+1-i}^+\chi_{2N+1}, \quad (4.18)$$

where the reality of the composite parafermion (4.11) is also taken into account. Note that the combined parity-time symmetry holds for wide range of models which are not conserved separately under the time reversal and the space reflection. The \mathcal{PT} invariant systems gain a lot of interest in physics because they have real energy spectra when this symmetry is unbroken, even if the corresponding Hamiltonians are not Hermitian [36].

Finally note that the mapping (4.18) applied to the Majorana fermions has the following form:

$$\mathcal{PT}\gamma_i\mathcal{PT} = -i\gamma_{2N+1-i}\gamma_{2N+1}.$$

Hamiltonian via time-reversal parafermions. As was already discussed, the parafermionic representation of the cluster Hamiltonian obtained in the previous section (3.29), (3.30), (5.17), contains the sixth order interaction terms (3.28). They form the second, even part of the system, H_{even} , and break a visual equivalence with its odd counterpart, H_{odd} . The symmetry between the two subsystems can be recovered if we use a simpler and more elegant version of the parafermionic Hamiltonian which involves the time-reversal particle operators (4.2). Indeed, the complex conjugate local terms of the H_{even} (2.3) are similar to the local terms of the H_{odd} (2.2). Their particle representation is provided by the complex conjugate version of Eq. (3.27):

$$Z_{l-1}X_lZ_{l+1}^+ = \epsilon\bar{\chi}_{2l-2}^+\bar{\chi}_{2l+1}. \quad (4.19)$$

It is bilinear in \mathcal{T} -transformed particle operators. So, the entire system with open boundaries can be rewritten now by is terms of the original and barred particle operators as follows:

$$H = -\frac{1}{2\epsilon}\sum_{r=1}^{L-1}\chi_{4r}^+\chi_{4r+3} - \frac{\epsilon}{2}\sum_{r=1}^{L'}\bar{\chi}_{4r-2}^+\bar{\chi}_{4r+1} + \text{H. c.} \quad (4.20)$$

This is an alternative version of the parafermionic system (3.29). Recall that the upper limit of the second sum depends on the length parity of the spin model (2.18), and the first and second sums describe, respectively, the odd (2.2) and even (2.3) parts of the original spin system.

We conclude that the entire system is separated into two subsystems each described by its own particles. Namely, the subsystem with the Hamiltonian H_{odd} is characterised by the standard parafermions with the indices $i = 0, 3$ modulo 4 while the other with H_{even} is expressed in terms of the time reversal parafermions with the indices $i = 1, 2$ mod 4.

For comparison, below we present also the parafermionic expression of the trivial Hamiltonian (2.19) which immediately follows from the particle representation of the X_l operator (3.6). It contains only the native particles but not their time reversals:

$$H_{\text{trivial}} = -\frac{1}{2}\sum_{l=2}^{N-1}(\epsilon^{-1}\chi_{2l-1}^+\chi_{2l} + \epsilon\chi_{2l}^+\chi_{2l-1}). \quad (4.21)$$

Another interesting simplest system containing only nearest-neighbouring terms describes the potential of the p -state Potts model:

$$H_{\text{ZZ}} = -\frac{1}{2}\sum_{i=1}^{N-1}(Z_i^+Z_{i+1} + Z_{i+1}^+Z_i) = -\frac{1}{2}\sum_{l=1}^{N-1}(\epsilon^{-1}\chi_{2l}^+\chi_{2l+1} + \epsilon\chi_{2l+1}^+\chi_{2l}). \quad (4.22)$$

The combination of two models (4.21), (4.22) with omitted second terms in the bracket, which violates the hermicity of the resulting Hamiltonian, produces the homogeneous version of the free-parafermion model [37]:

$$H_{\text{free}} = \epsilon^{-1}\sum_{i=1}^{2N-1}\chi_i^+\chi_{i+1},$$

which provides the parafermion representation of the Baxter's clock model [38].

Note that the use of the barred particle operators (instead of the usual ones) in the Hamiltonian (4.20) is necessary in order to associate the local terms with stabilisers. In fact, the replacement $\bar{\chi}_i \rightarrow \chi_i$ in the second part will spoil the commutativity among the local terms. Indeed, using the algebra (3.1), it is easy to verify that the "intersecting" local parts do not commute except for the fermionic case when $\omega = -1$:

$$(\chi_i^+\chi_j)(\chi_{i'}^+\chi_{j'}) = \omega^2(\chi_{i'}^+\chi_{j'})(\chi_i^+\chi_j) \quad \text{if} \quad i < i' < j < j'.$$

Splitting into two parafermion chains. It is worth noticing that each of the two parts, constituting the $\mathbb{Z}_p^{\text{odd}} \times \mathbb{Z}_p^{\text{even}}$ symmetry, is composed from the particle operators which form the related subsystem. Indeed, the generator X_{even} from Eq. (3.9) describes the total parafermion parity of the χ -particles which constitute the first subsystem in Eq. (4.20) described by the Hamiltonian H_{odd} . Similarly, the element X_{odd} , which may be presented also as

$$X_{\text{odd}} = \epsilon^{L'} \bar{\chi}_1^+ \bar{\chi}_2 \bar{\chi}_5^+ \bar{\chi}_6 \dots \bar{\chi}_{4L'-1}^+ \bar{\chi}_{4L'}, \quad (4.23)$$

is the total parity of the $\bar{\chi}$ -parafermions forming the second subsystem there corresponding to the H_{even} . So, the p -state cluster chain together with the symmetry, which protects its topological order, *splits* into two separate parafermion chains.

Now, let us investigate the *space-time symmetries* of the parafermionic cluster model (4.20). Obviously, the Hamiltonian (4.20) is not \mathcal{T} -invariant but transforms to an equivalent system given by its conjugate:

$$\bar{H} = \mathcal{T}H\mathcal{T}^{-1} = -\frac{\epsilon}{2} \sum_{r=1}^{L-1} \bar{\chi}_{4r}^+ \bar{\chi}_{4r+3} - \frac{1}{2\epsilon} \sum_{r=1}^{L'} \chi_{4r-2}^+ \chi_{4r+1} + \text{H. c.} \quad (4.24)$$

Obviously, its ground state differs from the ground state of the original system (2.14) except for the Majorana case when both models are identical:

$$|\bar{0}\rangle = \mathcal{T}|0\rangle = p^{-\frac{N}{2}} \sum_{n_1, \dots, n_N} \omega^{\sum_{l=1}^{N-1} (-1)^l n_l n_{l+1}} |n_1 \dots n_N\rangle \quad (4.25)$$

with the omitted term $n_N n_1$ in the exponent due to the free boundaries.

\mathcal{P} or \mathcal{PT} invariance of the Hamiltonian. The behaviour of the cluster model under the lattice reflection depends on the length parity of the spin chain. For *even* N , this transformation maps between local terms of the even and odd Hamiltonians making the entire system reflection-invariant as it follows from the relations (4.12), or (4.16). It is easy to see also that the ground state (2.14) also remains invariant and possesses the even parity quantum number:

$$N = 2L : \quad \mathcal{P}H_{\text{even}}\mathcal{P} = H_{\text{odd}}, \quad \mathcal{P}H\mathcal{P} = H, \quad (4.26)$$

$$\mathcal{P}|0\rangle = p^{-\frac{N}{2}} \sum_{n_1, \dots, n_N} \omega^{\sum_{l=1}^{N-1} (-1)^{l-1} n_{N-l+1} n_N} |n_1 n_2 \dots n_N\rangle = |0\rangle. \quad (4.27)$$

The same equations imply that for the *odd-length* chain, any local term of the odd Hamiltonian is reflected into a complex conjugate term within the same Hamiltonian. The even Hamiltonian also satisfies the same property. Therefore, both subsystems together with the entire system remain invariant under the simultaneous application of the parity and time reversal transformations. Meanwhile, it is easy to see that for odd N , the reflection like the time reversal (4.25) alters the sign in the coefficient's exponent in the ground-state expansion (2.14). Thus, both transformations act identically leaving the ground state \mathcal{PT} invariant with the unit quantum number:

$$N = 2L + 1 : \quad [\mathcal{PT}, H_{\text{even}}] = [\mathcal{PT}, H_{\text{odd}}] = [\mathcal{PT}, H] = 0, \quad (4.28)$$

$$\mathcal{P}|0\rangle = |\bar{0}\rangle, \quad \mathcal{PT}|0\rangle = |0\rangle. \quad (4.29)$$

Boundary parafermions. Consider now the parafermionic modes which appear at the edges of the open chain (4.20). Note that the left χ_1, χ_3 and the right χ_{2N-2}, χ_{2N} boundary operators together with their time reversal counterparts are absent in the Hamiltonian which, in fact, does not imply their conservation.

First consider the *even-length* model which is \mathcal{P} -invariant (4.26). Using the parafermionic algebra (3.1), (4.3), (4.6), it is easy to verify that the following three parafermions are conserved:

$$[H, \chi_1] = [H, \chi_3] = [H, \chi_{2N}] = 0. \quad (4.30)$$

Then, their reflected versions, which are proportional to the bilinear operators $\chi_k^\dagger \chi_{2N+1}$ with $k = 2N, 2N-2, 1$ (4.16) listed in the same order, are also edge invariants. As a result, the \mathcal{T} -counterparts of the edge particles also give rise to zero-energy modes:

$$[H, \bar{\chi}_{2N}] = [H, \bar{\chi}_{2N-2}] = [H, \bar{\chi}_1] = 0. \quad (4.31)$$

Meanwhile, the operators $\bar{\chi}_3$ and χ_{2N-2} are not preserved as is easy to verify. Using the parafermionic representations of the Z_1 (3.5) and Z_N (3.14) spin operators, it is easy to see that the first and last particles are interrelated with their barred versions:

$$\bar{\chi}_1\chi_1 = 1, \quad \bar{\chi}_{2N}\chi_{2N} = w^{-1}\chi_{2N+1}^2. \quad (4.32)$$

Eventually, only four parafermions out of above six ones remain independent. As an independent set of edge operators, one can choose, for instance, the system consisted on the following single particles:

$$\beta_1 = \chi_1, \quad \beta_2 = \chi_3, \quad \beta_3 = \chi_{2N}, \quad \beta_4 = \bar{\chi}_{2N-2}. \quad (4.33)$$

The above list is regrouped in a way to obey the conventional arrangement for the parafermions (3.1) with the commutation rules (4.6):

$$\beta_i^p = 1, \quad \beta_i\beta_j = \omega\beta_j\beta_i, \quad 1 \leq i < j \leq 4. \quad (4.34)$$

They form parafermionic versions of the boundary spins modes appearing due to fractionalization of the $\mathbb{Z}_p \times \mathbb{Z}_p$ symmetry on the left and right parts (2.8). These spin modes also are related to each other by the space parity map as is easy to see from the relations (2.21). They can be reexpressed in terms of the edge parafermions:

$$L_{\text{even}} = \beta_1^+, \quad L_{\text{odd}} = \beta_2, \quad R_{\text{even}} = \mathcal{P}\beta_2\mathcal{P} = \epsilon\beta_4^+\chi_{2N+1}, \quad R_{\text{odd}} = \mathcal{P}\beta_1^+\mathcal{P} = \epsilon^{-1}\beta_3^+\chi_{2N+1}. \quad (4.35)$$

In the derivation of the last equation, the second relation in (4.32) is applied. Before moving to the next case, let us note that the absent edge modes β_2 and β_3 to the first subsystem whereas the β_1, β_4 belong to the second one (4.20). This choice is suggested by their index values in the definition (4.33). [See also below for the translationally invariant case (5.8).] From this point of view, a more natural notation for the edge modes can be chosen by:

$$\beta'_1 = \chi_3, \quad \beta'_2 = \chi_{2N}, \quad \beta'_3 = \bar{\chi}_{2N-2}, \quad \beta'_4 = \bar{\chi}_1, \quad (4.36)$$

where the standard parafermion ordering (4.34) is preserved.

Now turn to the *odd-length* chains: $N = 2L + 1$. Evidently, the left edge parafermions χ_1 and χ_3 remain commuting with the Hamiltonian. Due to the \mathcal{PT} symmetry, their images under this map are also conserved. Therefore, from Eq. (4.18) it is clear that the following boundary particles are invariants of the system:

$$[H, \chi_1] = [H, \chi_3] = [H, \chi_{2N-2}] = [H, \chi_{2N}] = 0. \quad (4.37)$$

In contrast to the even-length cluster model, there is no antiparticle $\bar{\chi}_i$ among the invariant edge modes: all modes are described by particle operators χ_i . Hence, one can number them by retaining the original order:

$$\beta_1 = \chi_1, \quad \beta_2 = \chi_3, \quad \beta_3 = \chi_{2N-2}, \quad \beta_4 = \chi_{2N}. \quad (4.38)$$

The particle representation of the left and right edge symmetries (2.22) are provided, respectively, by the first two equations in (4.35) and the relations:

$$R_{\text{even}} = \mathcal{PT}\beta_2\mathcal{PT} = \epsilon^{-1}\beta_3^+\chi_{2N+1}, \quad R_{\text{odd}} = \mathcal{PT}\beta_1^+\mathcal{PT} = \epsilon^{-1}\beta_4\chi_{2N+1}^+.$$

Notice that in contrast to the even N case (4.36), the edge parafermion $\chi_{2N-2} = \chi_{4L}$ is associated to the odd part of the Hamiltonian while the particle $\chi_{2N} = \chi_{4L+2}$ is related to the even one. In this context, the following designation of the boundary modes might actually make more sense instead of the above one (4.38):

$$\beta'_1 = \chi_3, \quad \beta'_2 = \chi_{2N-2}, \quad \beta'_3 = \bar{\chi}_{2N}, \quad \beta'_4 = \bar{\chi}_1. \quad (4.39)$$

So, one might conclude that the open chain possesses four invariant edge parafermionic modes (4.36) or (4.39), equally distributed among the two parts: two of them are located at the boundaries of the odd subsystem while the remaining two – at the boundaries of the second, even subsystem.

In the Majorana case, the boundary invariants are described by the fermions $\gamma_1, \gamma_3, \gamma_{2N-2}$, and γ_{2N} irrespective on the boundary condition. The arrangement does not matter due to the anticommutation rules.

Note that the trivial model (4.21) does not contain the four edge particles: $\chi_1, \chi_2, \chi_{N-1}$, and χ_N which, as is easy to see, commute with the Hamiltonian. In the nontrivial phase, the second and third particles listed above are replaced by others which manifests a crucial difference between two systems.

V. SPLITTING INTO TWO SIMILAR CHAINS: TRANSLATIONALLY INVARIANT CASE.

Translation of spin lattice. The translation operator cyclically shifts a multispin state to the right on a single lattice step:

$$T|n_1 n_2 \dots n_N\rangle = |n_N n_1 \dots n_{N-1}\rangle, \quad TX_i T^{-1} = X_{i \bmod N+1}. \quad (5.1)$$

It produces \mathbb{Z}_N cyclic group since $T^N = 1$. Clearly, the translation is compatible with the time reversal map and produces the dihedral group \mathbb{D}_N together with the reflection \mathcal{P} .

Clearly, the translation maps to each other the $\mathbb{Z}_p \times \mathbb{Z}_p$ symmetry generators but leaves invariant their combination:

$$TX_{\text{even}} T^{-1} = X_{\text{odd}}, \quad TX_{\text{odd}} T^{-1} = X_{\text{even}}, \quad T\chi_{2N+1} T^{-1} = \chi_{2N+1}. \quad (5.2)$$

Let us describe how the translation act on the primary parafermions. Using their representation via the spin operators (3.3) together with inverse relations (3.5), it is easy to derive the translation rule for $l < N$:

$$T\chi_i T^{-1} = \epsilon^{-1} \chi_1 \chi_2^+ \chi_{i+2}, \quad \text{for } 1 \leq i \leq 2N-2. \quad (5.3)$$

Due to the nonlocal nature of the Fradkin-Kadanoff mapping, the translation rules are more complicate for the last two particles. In that case, the corresponding expressions contain, in addition, the composite parafermion operator (3.10):

$$T\chi_{2N-1} T^{-1} = \epsilon \chi_1 \chi_2^+ \chi_1 \chi_{2N+1}, \quad T\chi_{2N} T^{-1} = \epsilon \chi_1 \chi_{2N+1}. \quad (5.4)$$

More explicitly, the above two equations contain the additional right factor $\omega^{-1} \chi_{2N+1}$ comparing with the general transformation rules (5.3).

It is interesting to consider the conversion rule of the bilinear operators under the cyclic shift on a single spin node. It can derived using Eqs. (5.3), (5.4). Applying the exchange algebra of parafermions, the result can be presented in the compact form:

$$T\chi_i^+ \chi_j T^{-1} = \begin{cases} \chi_{[i+2]}^+ \chi_{[j+2]} & \text{if } i, j \leq 2N-2 \text{ or } i, j \geq 2N-1, \\ \omega^{-1} \chi_{i+2}^+ \chi_{[j+2]} \chi_{2N+1} & \text{if } i \leq 2N-2 \text{ and } j \geq 2N-1, \end{cases} \quad (5.5)$$

where the sum of the indexes in the square bracket must be carried out modulo $2N$. More precisely: $\chi_{[j+2]} := \chi_{1+(j+1) \bmod 2N}$. Hence, we identify: $\chi_{[2N+1]} := \chi_1$, and $\chi_{[2N+2]} := \chi_2$. We see that the translation on a single spin node produces the shift by two steps on the particle lattice. This property once again reflects the fact that a single spin encodes by two parafermions.

Note that the transformation of the time reversal particle operators $\bar{\chi}_i$ are described by the complex conjugate versions of Eqs. (5.3), (5.4), (5.5). The only change is the replacement of the phase factors there by the inverted values: $\omega, \epsilon \rightarrow \omega^{-1}, \epsilon^{-1}$.

The translation rules (5.3), (5.4) for the Majorana fermions acquire the following form:

$$T\gamma_i T^{-1} = -\iota \gamma_1 \gamma_2 \gamma_{i+2}, \quad T\gamma_{2N-1} T^{-1} = -\iota \gamma_2 \gamma_{2N+1}, \quad T\gamma_{2N} T^{-1} = \iota \gamma_1 \gamma_{2N+1} \quad (5.6)$$

where the range of the index i is defined above (5.3).

Finally we remark that the lattice transformations of particles we consider in the current article are inherited from those for the underlying spins (4.12), (5.1). This is a reason of a more complex behavior of particle operators under the chain reflection (4.16) and translation (5.3), (5.4) compared to the case when the parafermions or Majorana fermions [39] are the basic particles.

Hamiltonian and its symmetries. Now we are ready to construct the parafermionic representation of the cluster system with closed boundaries. As was discusses earlier, the chain's size must be even: $N = 2L$. This choice is made in order to guarantee the invariance of the initial spin Hamiltonian (2.1)–(2.3) under the translation on two lattice sites:

$$T^2 H T^{-2} = H. \quad (5.7)$$

One can construct the translationally invariant model starting from the parafermion chain with open boundary conditions. Indeed, from the action of the cyclic shift on the bilinears (5.5), it is easy to see that under the T^2 translation most local terms of the corresponding Hamiltonian (4.20) map to the next term with the index incremented

by four: $\chi_i \rightarrow \chi_{i+4}$. The exceptions are the four terms at the top right edge and their Hermitian conjugates. They, according to Eq. (5.5), are turned into the additional three-body interaction terms

$$T^2 \chi_{2N-4}^+ \chi_{2N-1} T^{-2} = \omega^{-1} \chi_{2N}^+ \chi_3 \chi_{2N+1}, \quad T^2 \bar{\chi}_{2N-6}^+ \bar{\chi}_{2N-3} T^{-2} = \omega \bar{\chi}_{2N-2}^+ \bar{\chi}_1 \chi_{2N+1},$$

which supplement the open system to the closed translationally invariant model. Thus, the parafermionic cluster Hamiltonian with periodical boundary conditions is given as follows:

$$H = -\frac{1}{2} \sum_{r=1}^{L-1} (\epsilon^{-1} \chi_{4r}^+ \chi_{4r+3} + \epsilon \bar{\chi}_{4r-2}^+ \bar{\chi}_{4r+1}) - \frac{1}{2} (\epsilon \chi_{2N}^+ \chi_3 + \epsilon^{-1} \bar{\chi}_{2N-2}^+ \bar{\chi}_1) \chi_{2N+1} + \text{H. c.} \quad (5.8)$$

The above Hamiltonian now contains all primary particles, and none is preserved among them in contrast to the open chain case (4.30), (4.31). This is in good agreement with the uniqueness of the SPT ground state in the closed systems.

Remember now that the composite parafermion operator χ_{2N+1} is the diagonal element of the $\mathbb{Z}_p \times \mathbb{Z}_p$ symmetry of the Hamiltonian (3.10). Therefore, the whole space of states is split into separate q -sectors, each specified by the eigenvalue $\chi_{2N+1} = \omega^q$ where $q = 0, 1, \dots, p-1$. Obviously, due to the parafermion exchange relations, any other particle operator, say, χ_1 together with its integral powers implements a one-to-one map between different q -sectors. Therefore, the sectors are equivalent and have the dimension p^{N-1} . Being restricted to the q -sector, the Hamiltonian (5.8) simplifies taking the bilinear form:

$$H_q = -\frac{1}{2} \sum_{r=1}^{L-1} (\epsilon^{-1} \chi_{4r}^+ \chi_{4r+3} + \epsilon \bar{\chi}_{4r-2}^+ \bar{\chi}_{4r+1}) - \frac{\omega^q}{2} (\epsilon \chi_{2N}^+ \chi_3 + \epsilon^{-1} \bar{\chi}_{2N-2}^+ \bar{\chi}_1) + \text{H. c.} \quad (5.9)$$

Clearly, the translational invariance enhances the symmetry of the Hamiltonian (5.8). Since in the current section we deal with the even N case, the system (5.8) remains invariant under the lattice reflection \mathcal{P} (4.26). Under the translation on a single site, the above Hamiltonian maps to its time reversal:

$$\begin{aligned} THT^{-1} &= \mathcal{T}HT^{-1} = \bar{H} \\ &= -\frac{1}{2} \sum_{r=1}^{L-1} (\epsilon \bar{\chi}_{4r}^+ \bar{\chi}_{4r+3} + \epsilon^{-1} \chi_{4r-2}^+ \chi_{4r+1}) - \frac{1}{2} (\epsilon^{-1} \bar{\chi}_{2N}^+ \bar{\chi}_3 + \epsilon \chi_{2N-2}^+ \chi_1) \chi_{2N+1} + \text{H. c.} \end{aligned} \quad (5.10)$$

This property may be established using Eq. (5.5), or is easier to observe when looking at the spin chain (2.1)–(2.3). Moreover, the joint $T\mathcal{T}$ transformation similar to the space parity \mathcal{P} , maps the even and odd parts of the Hamiltonian to each other. Hence, it is a symmetry of the closed model:

$$[T\mathcal{T}, H] = [\mathcal{P}, H] = 0, \quad T\mathcal{T}H_{\text{even}}(T\mathcal{T})^{-1} = \mathcal{P}H_{\text{even}}\mathcal{P} = H_{\text{odd}}. \quad (5.11)$$

Of course, both transformations mix the stabilisers inside in a different manner.

The two operators $T\mathcal{T}$ and \mathcal{P} , which are subjected to the algebraic relation

$$\mathcal{P}(T\mathcal{T})\mathcal{P} = (T\mathcal{T})^{-1},$$

generate a symmetry group with $2N$ elements which may be regarded as a antiunitary analog of the dihedral group \mathbb{D}_N . Obviously, the even powers of the generator $T\mathcal{T}$ are unitary while its odd powers are antiunitary. In particular, $(T\mathcal{T})^N = (T\mathcal{T})^{2L} = 1$, and $(T\mathcal{T})^2 = T^2$. Therefore, the T^2 (the translation on two spin sites) symmetry (5.7) may be considered as a part of more general $T\mathcal{T}$ symmetry of the Hamiltonian (5.8).

Due to the nondegeneracy of the closed model, the ground state (2.14) must be invariant under both transformation (5.11). Moreover, it form the trial representation of the the symmetry group considered in the current article. Actually, it was already observed that it has an even parity quantum number (4.27) under the reflection. From the other side, both the translation and time reversal (4.25) acting on the vacuum state produce the same result so that their combining action preserves it with the unit quantum number:

$$T\mathcal{T}|0\rangle = |0\rangle : \quad T|0\rangle = p^{-\frac{N}{2}} \sum_{n_1, \dots, n_N} \omega^{\sum_{i=1}^N (-1)^i n_i} |n_N n_1 \dots n_{N-1}\rangle = |\bar{0}\rangle. \quad (5.12)$$

The $\mathbb{Z}_p \times \mathbb{Z}_p$ symmetry, which protects the topological order of the cluster model, also acts trivially on the translationally invariant ground state (2.14):

$$X_{\text{even}}|0\rangle = X_{\text{odd}}|0\rangle = \chi_{2N+1}|0\rangle = |0\rangle. \quad (5.13)$$

The above condition does not hold for the open SPT state Rq. (2.23). The relations (4.27), (5.12), and (5.13) mean that the vacuum state of the closed system form a trivial, or singlet representation with respect to the whole considered symmetry group.

The system consisting on the Majorana fermions is real, and time reversal transformation (4.8) becomes a symmetry of the Hamiltonian regardless on the boundary conditions: $\bar{H} = H$. Therefore, the periodic system remains invariant under the translation on a single spin step (5.6):

$$H = THT^{-1} = \imath \sum_{l=1}^{N-2} \gamma_{2l} \gamma_{2l+3} - \imath (\gamma_{2N} \gamma_3 + \gamma_{2N-2} \gamma_1) \gamma_{2N+1}. \quad (5.14)$$

It is reduced to two decoupled periodic Kitaev chains. There are two sectors with $q = 0, 1$ corresponding to the eigenvalues $(-1)^q = \pm 1$ of the composite Majorana operator γ_{2N+1} . The restriction of the above Hamiltonian on them gives two decomped Majorana chains with antiperiodical and periodical boundary conditions:

$$H_{q=0} = \imath \sum_{l=1}^{N-2} \gamma_{2l} \gamma_{2l+3} - \imath (\gamma_{2N} \gamma_3 + \gamma_{2N-2} \gamma_1), \quad H_{q=1} = \imath \sum_{l=1}^N \gamma_{2l} \gamma_{[2l+3]}. \quad (5.15)$$

Here the square brackets in the last Majorana operator takes into account the periodicity condition alike in Eq. (5.5). It is worth mentioning that the second system with the odd fermion parity is the only case when the spin Hamiltonian retains the (true) translational invariance being expressed in terms of the particle operators. The statement is easily verified using Eq. (5.9) since it requires the condition $\omega^{q-1} = 1$ which is fulfilled only for the fermions in the sector with the odd particle number [39, 40].

Alike the open chain model (3.29), the cluster Hamiltonian also may be expressed also exclusively in terms of the original particle operators. To accomplish this, it remains to write in χ_i operators the last boundary term in Eq. (5.8) which can be done by returning to the spin model and making use of Eqs. (3.5), (3.14) and the exchange relations of particle operators:

$$\epsilon^{-1} \bar{\chi}_{2N-2}^+ \bar{\chi}_1 \chi_{2N+1} = Z_{N-1} X_N Z_1^+ = \epsilon^{-5} \chi_{2N-2} \chi_{2N-1}^2 \chi_{2N}^2 \chi_1^+ \chi_{2N+1}^+. \quad (5.16)$$

As a result, an alternative parafermion representation of the closed model (5.8) is given by the following formula:

$$H = -\frac{1}{2\epsilon} \sum_{r=1}^{L-1} (\chi_{4r}^+ \chi_{4r+3} + \omega^2 \chi_{4r-2} \chi_{4r-1}^2 \chi_{4r}^+ \chi_{4r+1}^+) - \frac{\epsilon}{2} (\chi_{2N}^+ \chi_3 \chi_{2N+1} + \omega^3 \chi_{2N-2} \chi_{2N-1}^2 \chi_{2N}^2 \chi_1^+ \chi_{2N+1}^+) + \text{H. c.} \quad (5.17)$$

String order parameters. Remember that the Hamiltonian (5.8) is built from the local stabilisers, which have to satisfy the condition

$$\epsilon^{-1} \chi_{4r}^+ \chi_{4r+3} |0\rangle = \epsilon \bar{\chi}_{4r-2}^+ \bar{\chi}_{4r+1} |0\rangle = \epsilon^{-1} \chi_3^+ \chi_{2N} |0\rangle = \epsilon \bar{\chi}_1^+ \bar{\chi}_{2N-2} |0\rangle = |0\rangle, \quad r = 1, 2, \dots, \frac{N}{2} - 1, \quad (5.18)$$

where the χ_{2N+1} is excluded for the last two equations (5.13). Thus, any product of above operators also stabilises the SPT state. In particular, the nonlocal string order operator is composed from the product of m sequential even or odd local stabilisers (2.15). Their parafermionic representations are also nonlocal:

$$Z_{2r-1} X_{2r} X_{2r+2} \dots X_{2r+2m-2} Z_{2r+2m-1}^+ = \epsilon^{-m} \prod_{r'=r}^{r+m-1} \bar{\chi}_{4r'-2}^+ \bar{\chi}_{4r'+1}, \quad (5.19)$$

$$Z_{2r}^+ X_{2r+1} X_{2r+3} \dots X_{2r+2m-1} Z_{2r+2m} = \epsilon^m \prod_{r'=r}^{r+m-1} \chi_{4r'+3}^+. \quad (5.20)$$

Note that above operators can be easily modified to the case when the string passes through the edges of the periodic system. Evidently, all these quantities mutually commute, and in case of $m = 1$, they are reduced to the local Hamiltonians.

Alike the Hamiltonian and its $\mathbb{Z}_p \times \mathbb{Z}_p$ symmetry generators, the string parameters are distributed between the two subsystems. Namely, the operator (5.20) is associated with the odd chain which is described in terms of the χ parafermions while the operator (5.19) is associated with the even chain given by the $\bar{\chi}$ particles.

VI. SUMMARY

In summary, we studied the parafermionic representation of the cluster chain, which possesses the $\mathbb{Z}_p^{\text{odd}} \times \mathbb{Z}_p^{\text{even}}$ symmetry protected topological (SPT) order. It corresponds to the p -state extension of the usual spin cluster with $p = 2$ and reduces to the noninteracting Potts model in the trivial phase. The odd and even local Hamiltonians are similar to each other but, in contrast to the Ising case, they are not equal. As a result, the straightforward application of the Fradkin-Kadanoff transformation, which maps the \mathbb{Z}_p -valued spins on one-dimensional lattice to parafermions, introduced more divergence between the odd and even subsystems.

Meanwhile, the use of the time-reversal (\mathcal{T}) particle operators allowed to avoid this discrepancy and recovered a similarity between two subsystems. As a result, we obtained two independent chains bilinear in particle operators. The first chain is composed of the standard parafermions described by the χ_{4r} and χ_{4r+3} parafermions while the second one consists of the time reversal parafermions: $\bar{\chi}_{4r-2}$, $\bar{\chi}_{4r+1}$. They inherit, respectively, the $\mathbb{Z}_p^{\text{odd}}$ and $\mathbb{Z}_p^{\text{even}}$ parts of the symmetry, which describe the total parity of particles contributing to each chain. Under the parity-conservation condition, the topological phases there become pure, without the symmetry-protection requirement. The long-range SPT order parameters turn into the topological order parameters of the χ - and $\bar{\chi}$ -particle chains.

It was shown also that for the open boundary condition, the independent zero-energy edge excitations are equally redistributed along the χ - and $\bar{\chi}$ -parafermionic chains: there is a single invariant (time-reversal) parafermionic mode per edge associated with each chain. At the same time, the structure of these modes depends on the parity of the spin chain's length N . For odd N , the Hamiltonian is \mathcal{PT} invariant, and the edge particles may be identified with the original (χ -type) parafermions. Meanwhile, for the even N , the Hamiltonian is reflection (\mathcal{P}) invariant, and at least one of the edge modes must be replaced by the time reversal ($\bar{\chi}$ -type) parafermion.

In case of the periodical boundary condition when the spin number N is always even, apart from the reflection invariance, the entire system possesses an additional $T\mathcal{T}$ symmetry which maps the χ - and $\bar{\chi}$ -chains to each other. As a result, the corresponding Hamiltonian becomes twisted by the total parafermion parity operator whose eigenvalues are p -th roots of unity. Being restricted to the invariant sector associated with a specific eigenvalue, the Hamiltonian takes a bilinear form with twisted boundary conditions. Note that the transformation rules of particle operators under the lattice reflection and translation (T) are induced by the transformation of the basic spin operators. They were described in detail in the article and have more complicated form than in the case of primary parafermions.

In the Ising case ($p = 2$), the parafermions together with their time reversals turn into the Majorana fermions. Then, the spin cluster splits up into two separate Kitaev chains in the topological phase as has been established recently [12].

ACKNOWLEDGMENTS

The research was supported by the Armenian Science Committee grant No. 21AG-1C047.

-
- [1] B. Zeng, X. Chen, D.-L. Zhou, and X.-G. Wen, "Quantum Information Meets Quantum Matter – From Quantum Entanglement to Topological Phase in Many-Body Systems", Springer New York, NY, 2019, [arXiv:1508.02595](#).
 - [2] Z.-C. Gu and X.-G. Wen, *Tensor-entanglement-filtering renormalization approach and symmetry-protected topological order*, Phys. Rev. B **80**, 155131 (2009), [arXiv:0903.1069](#); F. Pollmann, E. Berg, A.M. Turner, and M. Oshikawa, *Symmetry protection of topological phases in one-dimensional quantum spin systems*, Phys. Rev. B. **85**, 075125 (2012), [arXiv:0909.4059](#).
 - [3] W. Son, L. Amico, R. Fazio, A. Hamma, S. Pascazio, and V. Vedral, *Quantum phase transition between cluster and antiferromagnetic states*, EPL **95**, 50001 (2011), [arXiv:1103.0251](#).
 - [4] R. Raussendorf and H. J. Briegel, *A one-way quantum computer*, Phys. Rev. Lett. **86**, 5188 (2001).
 - [5] H. J. Briegel, D. E. Browne, W. Dür, R. Raussendorf, and M. Van den Nest, *Measurement-based quantum computation*, Nature Physics **5**, 19–26 (2009), [arXiv:0910.1116](#). T.-C. Wei, *Measurement-Based Quantum Computation*, in Oxford Research Encyclopedia of Physics, Ed. Brian Foster. New York: Oxford University Press (2021), [arXiv:2109.10111](#).
 - [6] S. D. Geraedts and O. I. Motrunic, *Exact models for symmetry-protected topological phases in one dimension*, [arXiv:1410.1580](#) (2014).
 - [7] L. H. Santos, *Rokhsar-Kivelson models of bosonic symmetry-protected topological states*, Phys. Rev. B **91**, 155150 (2015), [arXiv:1502.00066](#).
 - [8] J. H. Han, E. Lake, H. T. Lam, R. Verresen, and Y. You, *Topological quantum chains protected by dipolar and other modulated symmetries*, Phys. Rev. B **109**, 125121 (2024), [arXiv:2309.10036](#), 2023.
 - [9] A. Kitaev, *Unpaired Majorana fermions in quantum wires*, Phys.-Usp. **44**, 131 (2001), [arXiv:0010440](#).
 - [10] T. Dvir, *et. al.*, *Realization of a minimal Kitaev chain in coupled quantum dots*, Nature **614**, 445–459 (2023), [arXiv:2206.08045](#).

- [11] D. A. Ivanov, *Non-abelian statistics of half-quantum vortices in p-wave superconductors*, Phys. Rev. Lett. **86**, 268 (2001), [cond-mat/0005069](#).
- [12] R. Verresen, R. Moessner, and F. Pollmann, *One-dimensional symmetry protected topological phases and their transitions*, Phys. Rev. B **96**, 165124 (2017), [arXiv:1707.05787](#).
- [13] W. P. Su, J. R. Schrieffer, and A. J. Heeger, *Solitons in polyacetylene*, Phys. Rev. Lett. **42**, 1698 (1979).
- [14] A. Hutter, J. R. Wootton, and D. Loss, *Parafermions in a Kagome lattice of qubits for topological quantum computation*, Phys. Rev. X **5**, 041040 (2015).
- [15] Y. Wang, Z. Hu, B. C. Sanders, and S. Kais, *Qudits and high-dimensional quantum computing*, Front. Phys. **8**, 479 (2020), [arXiv:2008.00959](#).
- [16] A. Hutter and D. Loss, *Quantum computing with parafermions*, Phys. Rev. B **93**, 125105 (2016), [arXiv:1511.02704](#).
- [17] Zh.-H. Liu, *et. all*, *Topological contextuality and anyonic statistics of photonic-encoded parafermion*, PRX Quantum **2**, 030323 (2021), [arXiv:2011.05008](#).
- [18] E. Fradkin and L. P. Kadanoff, *Disorder variables and para-fermions in two-dimensional statistical mechanics*, Nucl. Phys. B **170** (1970) 1–15.
- [19] F. C. Alcaraz and R. Köberle, *Hidden parafermions in $Z(N)$ theories*, Phys. Rev. D **24**, 1562 (1981).
- [20] P. Fendley, *Parafermionic edge zero modes in Z_n -invariant spin chains*, J. Stat. Mech. (2012), P11020, [arXiv:1209.0472](#).
- [21] J. Alicea and P. Fendley, *Topological phases with parafermions: Theory and blueprints*, Annu. Rev. Condens. Matter Phys. **7**, 119–139 (2016), [arXiv:1504.02476](#); M. T. Batchelor, R. A. Henry, and X. Lu, *A Brief History of Free Parafermions*, AAPS Bull. **33**, 29 (2023), [arXiv:2311.04582](#).
- [22] D. J. Clarke, J. Alicea, and K. Shtengel, *Exotic non-Abelian anyons from conventional fractional quantum Hall states*, Nature Commun. **4**, 1348 (2013), [arXiv:1204.5479](#); N. Schiller, E. Cornfeld, E. Berg, and Y. Oreg, *Predicted signatures of topological superconductivity and parafermion zero modes in fractional quantum Hall edges*, Phys. Rev. Research **2**, 023296 (2020).
- [23] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, *Symmetry protected topological orders and the group cohomology of their symmetry group*, Phys. Rev. B **87**, 155114 (2013), [arXiv:1106.4772](#).
- [24] X. Chen, Z.-C. Gu, and X.-G. Wen, *Classification of gapped symmetric phases in one-dimensional spin systems*, Phys. Rev. B **83**, 035107 (2011), [arXiv:1008.3745](#).
- [25] M. den Nijs and K. Rommelse, *Preroughening transitions in crystal surfaces and valence-bond phases in quantum spin chains*, Phys. Rev. B **40**, 4709 (1989); T. Kennedy and H. Tasaki, *Hidden $Z_2 \times Z_2$ symmetry breaking in Haldane-gap antiferromagnets*, Phys. Rev. B **45**, 304 (1992).
- [26] M. Levin and Z.-C. Gu, *Braiding statistics approach to symmetry-protected topological phases*, Phys. Rev. B **86**, 115109 (2012), [arXiv:1202.3120](#).
- [27] H. Topchyanyan, V. Iugov, M. Mirumyan, Sh. Khachatryan, T. Hakobyan, and T. Sedrakyan, *Z_3 and $(\times Z_3)^3$ symmetry protected topological paramagnets*, JHEP **12**, 199 (2023), [arXiv:2210.01187](#); H. Topchyanyan, V. Iugov, M. Mirumyan, T. Hakobyan, T. Sedrakyan, and A. Sedrakyan, *Two-dimensional topological paramagnets protected by Z_3 symmetry: Properties of the boundary Hamiltonian*, [arXiv:2312.15095](#).
- [28] E. Cobanera and G. Ortiz, *Fock parafermions and self-dual representations of the braid group*, Phys. Rev. A **89**, 012328 (2014), [arXiv:1307.6214](#).
- [29] D. Rossini, M. Carrega, M. C. Strinati, and L. Mazza, *Anyonic tight-binding models of parafermions and of fractionalized fermions*, Phys. Rev. B **99**, 085113 (2019), [arXiv:1811.09422](#).
- [30] J.-S. Hong, S.-Q. Zhang, X. Liu, and X.-J. Liu, *Parafermions with symmetry-protected non-Abelian statistics*, [arXiv:2403.09602](#).
- [31] U. Güngördü, R. Nepal, and A. A. Kovalev, *Parafermion stabilizer codes*, Phys. Rev. A **90**, 042326 (2014), [arXiv:1409.4724](#).
- [32] L. Tsui, Y.-T. Huang, H.-C. Jiang, and D.-H. Lee, *The phase transitions between $Z_n \times Z_n$ bosonic topological phases in 1+1 D, and a constraint on the central charge for the critical points between bosonic symmetry protected topological phases*, Nucl. Phys. B **919**, 470 (2017), [arXiv:1701.00834](#).
- [33] J. Wouters, F. Hassler, H. Katsura, and D. Schuricht, *Phase diagram of an extended parafermion chain*, SciPost Phys. Core **5**, 008 (2022), [arXiv:2106.15823](#).
- [34] R. A. Henry and M. T. Batchelor, *Exceptional points in the Baxter-Fendley free parafermion model*, SciPost Phys. **15** (2023) 016, [arXiv:2301.11031](#).
- [35] W.-T. Xu and G.-M. Zhang, *Enriched classification of parafermionic gapped phases with time reversal symmetry*, Phys. Rev. B **97** (2018) 125135, [arXiv:1709.04657](#).
- [36] C. M. Bender and S. Boettcher, *Real spectra in non-Hermitian Hamiltonians having PT symmetry*, Phys. Rev. Lett. **80**, 5243 (1998), [physics/9712001](#).
- [37] P. Fendley, *Free parafermions*, J. Phys. A **47**, 075001 (2014), [arXiv:1310.6049](#).
- [38] R. J. Baxter, *A simple solvable Z_N Hamiltonian*, Phys. Lett. A **140**, 155 (1989); *Superintegrable chiral Potts model: thermodynamic properties, an “inverse” model, and a simple associated Hamiltonian*, J. Stat. Phys. **57**, 1 (1989).
- [39] N. Seiberg and S.-H. Shao, *Majorana chain and Ising model – (non-invertible) translations, anomalies, and emanant symmetries*, SciPost Phys. **16**, 064 (2024), [arXiv:2307.02534](#).
- [40] E. Lieb, T. Schultz, and D. Mattis, *Two soluble models of an antiferromagnetic chain*, Ann. Phys. **16**, 407–466 (1961).