

PROJECTION THEOREMS IN THE PRESENCE OF EXPANSIONS

K. W. OHM

ABSTRACT. We prove a restricted projection theorem for a certain one dimensional family of projections from \mathbb{R}^n to \mathbb{R}^k .

The family we consider here arises naturally in the study of quantitative equidistribution problems in homogeneous dynamics.

CONTENTS

1.	Introduction	1
2.	Proof of Theorem 1.1	3
3.	Proof of Proposition 1.2	9
	References	11

1. INTRODUCTION

Restricted projection problems are intimately related to central questions in Fourier analysis and incidence geometry, and have been much studied, e.g., by Mattila, Falconer, Bourgain and others. More recently, certain restricted projection theorems have also found striking applications in homogeneous dynamics.

Let us recall the the classical Marstrand projection theorem: Let $K \subset \mathbb{R}^n$ be a compact subset, then for a.e. $v \in \mathbb{S}^{n-1}$

$$(1.1) \quad \dim p_v(K) = \min(1, \dim K),$$

where $p_v(w) = w \cdot v$ is the orthogonal projection in the direction of v and here and in what follows \dim denotes the Hausdorff dimension. Analogous statements hold more generally for orthogonal projection into a.e. m -dimensional subspace, with respect to the Lebesgue measure on $\text{Gr}(m, n)$.

Broadly speaking, restricted projection problems seek to obtain similar results as in (1.1) where v is confined to a proper Borel subset $B \subset \mathbb{S}^{n-1}$. Note, however, that without further restrictions on B , (1.1) fails: e.g., if

$$B = \{(\cos t, \sin t, 0) : 0 \leq t \leq 2\pi\}$$

is the great circle in \mathbb{S}^2 and K is the z -axis, then $p_v(K) = 0$ for every $v \in B$.

To avoid degenerations of this nature, one may, e.g., consider a curve $\gamma : [0, 1] \rightarrow \mathbb{R}^n$ satisfying that

$$\{\gamma'(r), \gamma''(r), \dots, \gamma^{(n)}(r)\} \quad \text{spans } \mathbb{R}^n \text{ for all } r \in [0, 1].$$

It is natural to inquire whether for almost all $r \in [0, 1]$ the following holds

$$(1.2) \quad \dim \text{p}_{\gamma'(r)}(K) = \min(1, \dim K);$$

again $K \subset \mathbb{R}^n$ is a compact subset.

Indeed (1.2) was conjectured by Fässler and Orponen [FO14] in dimension 3. This conjecture was resolved by Käenmäki, Orponen, and Venieri [KOV17] and Pramanik, Yang, and Zahl [PYZ22] — these works rely on the work of Wolff and Schlag on circular Kakeya sets [Wol00, Sch03].

More recently, Gan, Guo, and Wang [GGW22] have established (1.2) in all dimensions using decoupling inequalities for the moment curve due to Bourgain, Demeter, and Guth [BDG16].

In this paper, we study a closely related family of projections. The consideration of these families are mainly motivated by the aforementioned applications to homogeneous dynamics.

Let us fix some notation in order to state the main results of this paper. For every $t \geq 0$ and $r \in [0, 1]$, let $\pi_{t,r} : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be

$$(1.3) \quad \pi_{t,r}(x, y, z) = \left(e^t(x + ry + \frac{r^2}{2}z), y + rz \right)$$

The following is the main theorem in this paper.

1.1. Theorem. *Let $1 \leq \alpha \leq 3/2$, and let $0 < \delta_0 \leq 1$. Let $F \subset B_{\mathbb{R}^3}(0, 1)$ be a finite set satisfying the following:*

$$\#(B_{\mathbb{R}^3}(w, \delta) \cap F) \leq C_0 \cdot \delta^\alpha \cdot (\#F) \quad \text{for all } w \in \mathbb{R}^3 \text{ and all } \delta \geq \delta_0$$

where $C_0 \geq 1$.

Let $0 < \varepsilon < \alpha/100$. For every $\delta \geq e^t \delta_0$, there is a subset $I_\delta \subset [1/2, 1]$ with

$$|[1/2, 1] \setminus I_\delta| \ll_\varepsilon \delta^{\star\varepsilon}$$

so that the following holds. Let $r \in I_\delta$, then there exists $F_{\delta,r} \subset F$ with

$$\#(F \setminus F_{\delta,r}) \ll_\varepsilon \delta^{\star\varepsilon} \cdot (\#F)$$

such that for all $w \in F_{\delta,r}$, we have

$$\#(\{w' \in F_{\delta,r} : \|\pi_{t,r}(w') - \pi_{t,r}(w)\| \leq \delta\}) \ll_\varepsilon C_0 e^{-t/10} \delta^{\alpha-\varepsilon} \cdot (\#F)$$

Remark. Throughout the paper, the notation $a \ll b$ and $a^{\star b}$ mean $a \leq Db$ and a^{Db} , respectively, where D is some positive constant whose dependence is explicated in different statements. Also, for a Borel subset $B \subset \mathbb{R}^d$, we denote the Lebesgue measure of B by $|B|$.

We also state a similar result in all dimensions $n \geq 3$, which relies on the aforementioned work of [GGW22].

Let $2 \leq k \leq n - 1$. For all $t \geq 0$ and all $r \in [0, 1]$, let $\pi_{t,r}^{(k)} : \mathbb{R}^n \rightarrow \mathbb{R}^k$ be

$$\pi_{t,r}^{(k)}(w) = (e^t w \cdot \xi'(r), e^t w \cdot \xi''(r), \dots, e^t w \cdot \xi^{(k-1)}(r), w \cdot \xi^{(k)}(r))$$

where $w \cdot v$ is the usual inner product on \mathbb{R}^n and

$$\xi(r) = \left(\frac{r}{1!}, \frac{r^2}{2!}, \dots, \frac{r^n}{n!} \right) \subset \mathbb{R}^n.$$

Using [GGW22, Thm. 2.1] and some elementary arguments, one can establish the following

1.2. Proposition. *Let $0 < \alpha \leq k$, and let $0 < \delta_0 \leq 1$. Let $F \subset B_{\mathbb{R}^n}(0, 1)$ be a finite set satisfying the following:*

$$\#(B_{\mathbb{R}^n}(w, \delta) \cap F) \leq C_0 \cdot \delta^\alpha \cdot (\#F) \quad \text{for all } w \in F \text{ and all } \delta \geq \delta_0$$

where $C_0 \geq 1$.

Let $0 < \varepsilon < \alpha/100$. For every $\delta \geq e^t \delta_0$, there is a subset $I_\delta \subset [1/2, 1]$ with

$$|[1/2, 1] \setminus I_\delta| \ll_\varepsilon \delta^{*\varepsilon^2}$$

so that the following holds. Let $r \in I_\delta$, then there exists $F_{\delta,r} \subset F$ with

$$\#(F \setminus F_{\delta,r}) \ll_\varepsilon \delta^{*\varepsilon^2} \cdot (\#F)$$

such that for all $w \in F_{\delta,r}$, we have

$$\#(\{w' \in F_{\delta,r} : \|\pi_{t,r}(w') - \pi_{t,r}(w)\| \leq \delta\}) \ll_\varepsilon C_0 e^{-\alpha' t} \delta^{\alpha - \varepsilon} \cdot (\#F)$$

where $\alpha' = \alpha$ if $0 < \alpha \leq k - 1$ and $\alpha' = \alpha - 1$ otherwise.

It is worth noting that Proposition 1.2 does not imply Theorem 1.1; indeed $\alpha' = \alpha - 1 < 1/10$ for $1 \leq \alpha < 11/10$. That said, it seems plausible that the approach developed in [GGW22] can be modified to prove a stronger version of Proposition 1.2 which then would imply (and perhaps improve) Theorem 1.1. However, our proof of Theorem 1.1 relies on much softer arguments and is an adaptation of [OV18].

Also as previously mentioned, our precise formulations here (which at least at first glance may appear somewhat ad hoc) are motivated by recent applications in homogeneous dynamics [LM23, LMW22, LMWY23].

2. PROOF OF THEOREM 1.1

In this section we will prove Theorem 1.1. Let us begin by fixing some notation which will be used throughout this section.

Let μ denote the normalized counting measure on F :

$$\mu(B) = \frac{\#(B \cap F)}{\#F} \quad \text{for any Borel set } B \subset \mathbb{R}^3.$$

Since t is fixed throughout the argument, we will write π_r for $\pi_{t,r}$. For every $w \in F$, all $r \in [0, 1]$, and all $b > 0$, let

$$m^b(\pi_r(w)) = \mu(\{w' \in F : \|\pi_r(w) - \pi_r(w')\| \leq b\}).$$

More generally, given a subset $B \subset \mathbb{R}^3$, let

$$m^b(\pi_r(w)|B) = \mu(\{w' \in B : \|\pi_r(w) - \pi_r(w')\| \leq b\}).$$

For all $w \in F$ and all $b > 0$, let $D_b(w) = \{w' : b \leq \|w - w'\| \leq 2b\}$.

The following lemma is the main step in the proof of Theorem 1.1.

2.1. Lemma. *The following holds for all small enough η and all large enough C . Let $F_{\text{hm}} \subset F$ denote the set of $w \in F$ so that*

$$|\{r \in [\frac{1}{2}, 1] : m^\delta(\pi_r(w)) \geq C_0 e^{-t/10} \delta^{\alpha-18\eta}\}| \geq C\delta^\eta.$$

Then $\mu(F_{\text{hm}}) \leq C\delta^\eta$.

We will use the following elementary lemma in the proof of Lemma 2.1.

2.2. Lemma. *Let σ be a probability measure on $B(0, 10)$ which satisfies*

$$\sigma(B(w, r)) \leq \hat{C}r^\beta \quad \text{for all } w \in \mathbb{R}^3 \text{ and all } r \geq r_0$$

where $1/2 \leq \beta \leq 1$ and $\hat{C} > 0$.

The following holds for all ε_0 small enough. For every $b \geq r_0$, there exists a subset E_b with $\sigma(B(0, 10) \setminus E_b) \leq b^{\varepsilon_0}$, and for every $z \in E_b$, there is a subset $I_z \subset [0, 1]$ with $|[0, 1] \setminus I_z| \leq b^{\varepsilon_0}$ so that for every $r \in I_z$

$$\sigma\{z' : |(1, r, \frac{r^2}{2}) \cdot (z - z')| \leq b\} \ll \hat{C}b^{49\beta/100}$$

Proof. This, rather weak estimate, follows from standard arguments, see e.g. [Ohm23, §3]. It is worth noting that much stronger result holds where $49\beta/100$ is replaced by $\beta - o(1)$, see [LM23, Thm. B.1] and [KOV17]. \square

2.3. Lemma. *Let the notation and assumptions be as in Lemma 2.1. The following holds for all small enough η . There exists some $b \geq \delta^{1-3\eta}$ and a subset $F'_{\text{hm}} \subset F_{\text{hm}}$ with*

$$(2.1) \quad \mu(F'_{\text{hm}}) \geq \delta^{3\eta/2}$$

so that for all $w \in F'_{\text{hm}}$, we have

$$(2.2) \quad |\{r \in [\frac{1}{2}, 1] : m^\delta(\pi_r(w)|D_b(w)) \geq C_0 e^{-t/10} \delta^{\alpha-7\eta}\}| \geq \delta^{3\eta/2}.$$

Proof. Let $b_0 = \delta^{1-3\eta}$ and recall that $1 \leq \alpha \leq 3/2$.

If $e^{-t} \geq \delta^{100\eta}$, then $e^{-t/10} \delta^{\alpha-18\eta} \geq \delta^{\alpha-8\eta}$. Therefore,

$$\begin{aligned} \{r \in [\frac{1}{2}, 1] : m^\delta(\pi_r(w)) \geq C_0 e^{-t/10} \delta^{\alpha-18\eta}\} \subset \\ \{r \in [\frac{1}{2}, 1] : m^\delta(\pi_r(w)) \geq C_0 \delta^{\alpha-8\eta}\}. \end{aligned}$$

Now since $\mu(B(w, b_0)) \leq C_0 b_0^\alpha \leq C_0 \delta^{\alpha-4.5\eta}$, there is some $b = b(w, r) \geq \delta^{1-3\eta}$ so that

$$(2.3) \quad m^\delta(\pi_r(w)|D_b(w)) \geq C_0 e^{-t/10} \delta^{\alpha-7.5\eta}.$$

In view of this we assume $e^{-t} \leq \delta^{100\eta}$ for the rest of the argument. Put

$$\Xi_\delta := \{w : \mu(B(w, b_0)) > C_0 e^{-t/10} \delta^{\alpha-7\eta}\}$$

Also for every $w \in F_{\text{hm}}$, let

$$I(w) = \{r \in [\frac{1}{2}, 1] : m^\delta(\pi_r(w)) \geq C_0 e^{-t/10} \delta^{\alpha-18\eta} \cdot (\#F)\}.$$

We now consider two possibilities:

Case 1: Assume $\mu(F_{\text{hm}} \setminus \Xi_\delta) \geq \delta^\eta/100$.

In this case, let $F''_{\text{hm}} = F_{\text{hm}} \setminus \Xi_\delta$. Then $\mu(F''_{\text{hm}}) \geq \delta^\eta/100$. Moreover, for every $w \in F''_{\text{hm}}$, we have $\mu(B(w, b_0)) \leq C_0 e^{-t/10} \delta^{\alpha-7\eta}$.

Now since for every $r \in I(w)$, we have

$$m^\delta(\pi_r(w)) \geq C_0 e^{-t/10} \delta^{\alpha-18\eta}$$

there exists some $b = b(w, r) \geq \delta^{1-3\eta}$ so that

$$(2.4) \quad m^\delta(\pi_r(w)|D_b(w)) \gg C_0 e^{-t/10} \delta^{\alpha-17\eta}.$$

Case 2: Assume $\mu(F_{\text{hm}} \setminus \Xi_\delta) \leq \delta^\eta/100$.

In this case, we have $\mu(F_{\text{hm}} \cap \Xi_\delta) \geq \frac{C}{2} \delta^\eta$.

Fix a maximal $b_0/2$ separated subset $\{w_1, \dots, w_M\}$ of $F_{\text{hm}} \cap \Xi_\delta$. Discarding a subset of F_{hm} with measure $\leq \mu(F_{\text{hm}})/100$, we will assume that

$$\mu(B(w_i, b_0)) \gg \mu(B(w_i, 4b_0)) \quad \text{for all } 1 \leq i \leq M,$$

where the implied constant is absolute.

Let ν_i be the measure on $B(0, 1)$ which is the image of

$$\frac{1}{\mu(B(w_i, b_0))} \mu|_{B(w_i, b_0)}$$

under the map $w \rightarrow \frac{w_i - w}{b_0}$. Similarly, let $\tilde{\nu}_i$ denote the measure on $B(0, 4)$ is the image of

$$\frac{1}{\mu(B(w_i, 4b_0))} \mu|_{B(w_i, 4b_0)}$$

under the map $w \rightarrow \frac{w_i - w}{b_0}$.

Then for any $1 \leq i \leq M$ and for $\sigma = \nu_i, \tilde{\nu}_i$, we have

$$\sigma(B(z, r)) \leq e^{t/10} \delta^{-7\eta} r^\alpha \quad \text{for all } r \geq \delta_0/b_0.$$

Thus, applying Lemma 2.2 with $\nu_i, \tilde{\nu}_i$ (for any $1 \leq i \leq M$), there exists a subset $E_i \subset B(w_i, b_0)$, with

$$\nu_i(B(w_i, b_0) \setminus E_i) \ll \delta^{\varepsilon_0}$$

and for every $w \in E_i$, there is a subset J_w with $|[0, 1] \setminus J_w| \ll \delta^{\varepsilon_0}$ so that if $r \in J_w$, then for $\sigma = \frac{1}{\mu(B(w_i, b_0))} \mu|_{B(w_i, b_0)}$ and $\sigma = \frac{1}{\mu(B(w_i, 4b_0))} \mu|_{B(w_i, 4b_0)}$,

$$(2.5) \quad \begin{aligned} \sigma(\{w : |(1, r, \frac{r^2}{2}) \cdot \frac{w-w'}{b_0}| \leq e^{-t}\}) &\ll e^{t/10} \delta^{-7\eta} e^{-49t/100} \\ &\ll e^{-0.3t}, \end{aligned}$$

where we used $e^{-t} \leq \delta^{100\eta}$ in the second inequality.

Let $F''_{\text{hm}} = (F_{\text{hm}} \cap \Xi_\delta) \cap (\bigcup_i E_i)$. Then

$$\mu(F''_{\text{hm}}) \gg \delta^\eta$$

For every $w \in F''_{\text{hm}}$, let $\hat{I}(w) = I(w) \cap J_w$. Then $|\hat{I}(w)| \geq \frac{1}{2}\delta^\eta$. Moreover, for every $w \in F''_{\text{hm}}$ and every $r \in \hat{I}(w)$,

$$\{w' \in B(w, b_0) : \|\pi_r(w) - \pi_r(w')\| \leq \delta\} \subset \{w' \in B(w_i, 4b_0) : |(1, r, r^2) \cdot \frac{w-w'}{b_0}| \leq e^{-t}\}$$

where $w \in B(w_i, b_0)$ (see (1.3) for the definition of $\pi_r = \pi_{t,r}$ and recall that $b_0 = \delta^{1-3\eta}$). Applying (2.5) with $\sigma = \frac{1}{\mu(B(w_i, 4b_0))}\mu|_{B(w_i, 4b_0)}$,

$$(2.6) \quad \mu(\{w' \in B(w, b_0) : \|\pi_r(w) - \pi_r(w')\| \leq \delta\}) \ll e^{-0.3t}\mu(B(w_i, 4b_0)) \ll C_0 e^{-0.3t} b_0^\alpha < C_0 e^{-0.2t} \delta^{\alpha-7\eta},$$

where in the last inequality, we used $e^{-t} \leq \delta^{100\eta}$.

In view of (2.6), for every $w \in F''_{\text{hm}}$ and all $r \in \hat{I}(w)$, there exists some $b > b_0 = \delta^{1-3\eta}$ so that (2.4) holds. That is:

$$(2.7) \quad m^\delta(\pi_r(w)|D_b(w)) \gg C_0 e^{-t/10} \delta^{\alpha-17\eta}.$$

Altogether, combining (2.3), (2.4), and (2.7), we have found a subset $F''_{\text{hm}} \subset F_{\text{hm}}$ with $\mu(F''_{\text{hm}}) \gg \delta^\eta$ and for every $w \in F''_{\text{hm}}$ a subset $\hat{I}(w) \subset [\frac{1}{2}, 1]$ with $|\hat{I}(w)| \gg \delta^\eta$ so that the following holds. For every $w \in F''_{\text{hm}}$ and all $r \in \hat{I}(w)$, there exists $b = b(w, r) \geq \delta^{1-3\eta}$ so that

$$m^\delta(\pi_r(w)|D_b(w)) \gg C_0 e^{-t/10} \delta^{\alpha-17\eta}$$

Now applying pigeonhole principle (and Fubini's theorem), there exists $b \geq \delta^{1-3\eta}$ and $F'_{\text{hm}} \subset F''_{\text{hm}}$ with

$$\mu(F'_{\text{hm}}) \gg \delta^{3\eta/2}$$

so that for all $w \in F'_{\text{hm}}$, we have

$$|\{r \in [\frac{1}{2}, 1] : m^\delta(\pi_r(w)|D_b(w)) \geq C_0 e^{-t/10} \delta^{\alpha-7\eta}\}| \geq \delta^{3\eta/2}.$$

The proof is complete. \square

Proof of Lemma 2.1. Assuming C is large enough, we may assume δ is small throughout the proof.

In view of Lemma 2.3, see in particular (2.1), we will replace F_{hm} by F'_{hm} and assume that there is some $b \geq \delta^{1-3\eta}$ so that (2.2) holds for all $w \in F_{\text{hm}}$.

For every $w \in F_{\text{hm}}$, set

$$I'(w) = \{r \in [\frac{1}{2}, 1] : m^\delta(\pi_r(w)|D_b(w)) \geq C_0 e^{-t/10} \delta^{\alpha-7\eta}\}.$$

Then $|I'(w)| \gg \delta^{3/2}$, see (2.2). Choose three subsets $I'_j(w) \subset I'(w)$ for $j = 1, 2, 3$, which satisfy the following properties

$$(2.8) \quad |I'_j(w)| \gg \delta^{2\eta} \quad \text{and} \quad \text{dist}(I'_i(w), I'_j(w)) \gg \delta^{3\eta/2} \quad \text{for } i \neq j$$

For $j = 1, 2, 3$, define

$$E_j(w) = \{w' \in F \cap D_b(w) : \|\pi_r(w) - \pi_r(w')\| \leq \delta \text{ for some } r \in I'_j(w)\}.$$

We claim

$$(2.9) \quad \mu(E_j(w)) \geq C_0 b e^{-t/10} \delta^{\alpha-1-6\eta} \quad \text{for } j = 1, 2, 3.$$

Fix one j and cover $I'_j(w)$ with intervals J_1, \dots, J_N of size $C'\delta/b$ for some C' which will be chosen to be large. Thus

$$(2.10) \quad N \gg \delta^{2\eta}/(C'\delta/b) = b\delta^{2\eta}/(C'\delta) \gg b\delta^{2\eta-1}.$$

Since $b \geq \delta^{1-3\eta}$, (2.10) in particular implies that $N \gg \delta^{-\eta}$.

For each J_i , let $r_i \in J_i \cap I'_j(w)$. Discarding at most half of J_i 's, we will assume $|r_i - r_{i'}| \geq C'\delta/b$, and will continue to denote the collection by J_1, \dots, J_N . For every $1 \leq i \leq N$, let

$$E_{j,i} = \{w' \in E_j(w) : \|\pi_{r_i}(w) - \pi_{r_i}(w')\| \leq \delta\}.$$

Then $\mu(E_{j,i}) \geq C_0 e^{-\kappa t} \delta^{\alpha-7\eta}$. Moreover, $E_{j,i} \cap E_{j,i'} = \emptyset$. Indeed if $w' \in E_{j,i} \cap E_{j,i'}$, then

$$r_i, r_{i'} \in \{r \in [\frac{1}{2}, 1] : \|\pi_r(w) - \pi_r(w')\| \leq \delta\}.$$

Thus the set of $r \in [\frac{1}{2}, 1]$ so that

$$\begin{aligned} |(1, r, \frac{r^2}{2}) \cdot (w - w')| &\leq e^{-t} \delta \quad \text{and} \\ |(0, 1, r) \cdot (w - w')| &\leq \delta \end{aligned}$$

has diameter $\geq C'\delta/b$. Since $b \leq \|w - w'\| \leq 2b$, we get a contradiction so long as C' is large enough.

Using (2.10) and $\mu(E_{j,i}) \geq C_0 e^{-\kappa t} \delta^{\alpha-7\eta}$, thus

$$\mu(E_j(w)) \gg N \cdot \mu(E_{j,i}) \gg C_0 b e^{-t/10} \delta^{\alpha-1-5\eta},$$

for $j = 1, 2, 3$, as we claimed in (2.9).

Using $\mu(F_{\text{hm}}) \gg \delta^{3\eta/2}$ and (2.9), we conclude that

$$(2.11) \quad \mu(\{(w, w_1, w_2, w_3) \in F_{\text{hm}} \times F^3 : w_j \in E_j(w)\}) \gg C_0^3 b^3 e^{-0.3t} \delta^{3\alpha-3-13\eta}$$

We now find an upper bound for the measure of the set on the left side of (2.11). To that end, fix some $(w_1, w_2, w_3) \in F^3$ so that there exists some $w \in F_{\text{hm}}$ with $w_j \in E_j(w)$. In particular, we have $w_j \in D_b(w)$, where $b \geq \delta^{1-3\eta}$, and

$$(2.12) \quad \|w_1 - w_j\| \leq 4b, \quad \text{for } j \in \{1, 2, 3\}.$$

Let $r_j \in I'_j(w)$ be so that $\|\pi_{r_j}(w) - \pi_{r_j}(w_j)\| \leq \delta$. Then $|r_i - r_j| \geq \delta^{3\eta/2}$ for $i \neq j$ and

$$(1, r_j, r_j^2) \cdot w = (1, r_j, r_j^2) \cdot w_j + O(e^{-t} \delta) \quad \text{for } j \in \{1, 2, 3\}.$$

Thus w belongs to a set with diameter $\ll e^{-t} \delta^{1-4.5\eta}$. This and (2.12) imply

$$\mu(\{(w, w_1, w_2, w_3) \in F_{\text{hm}} \times F^3 : w_j \in E_j(w)\}) \ll C_0^3 e^{-\alpha t} \delta^{\alpha-4.5\eta\alpha} b^{2\alpha}$$

Comparing this upper bound and (2.11), we conclude

$$b^3 e^{-3t/10} \delta^{3\alpha-3-13\eta} \ll e^{-\alpha t} \delta^{\alpha-4.5\eta\alpha} b^{2\alpha},$$

which implies

$$\delta^{2\alpha-3-13\eta+4.5\eta\alpha} b^{3-2\alpha} \ll e^{(-\alpha+0.3)t}.$$

Now using $b \geq \delta^{1-3\eta}$, $3-2\alpha > 0$, and the above, we conclude that

$$(2.13) \quad \delta^{2\alpha-3-13\eta+4.5\eta\alpha} \delta^{(3-2\alpha)(1-3\eta)} = \delta^{-13\eta+4.5\eta\alpha-(9-6\alpha)\eta} \ll e^{(-\alpha+0.3)t}$$

However, $-13\eta + 4.5\eta\alpha - (9-6\alpha)\eta \leq -6\eta$, since $3-2\alpha > 0$. Assuming δ is small enough and recalling that $3/10 < \alpha$, (2.13) cannot hold. This contradiction completes the proof. \square

Proof of Theorem 1.1. We now complete the proof of Theorem 1.1, which is based on Lemma 2.1 and Fubini's theorem.

Recall that for every for all $r \in [\frac{1}{2}, 1]$ and all $w \in F$, we put

$$m^\delta(\pi_r(w)) = \mu\{w' : \|\pi_r(w) - \pi_r(w')\| \leq \delta\}$$

Let $\varepsilon > 0$ and let $\eta = \varepsilon/20$. For all $r \in [\frac{1}{2}, 1]$, let

$$F_{\text{bad}}(r) = \{w : m^\delta(\pi_r(w)) \geq C_0 e^{-t/10} \delta^{\alpha-18\eta}\}.$$

The claim in Theorem 1.1 follows if we show that there is a subset $I_\delta \subset [\frac{1}{2}, 1]$ with $|\frac{1}{2}, 1] \setminus I_\delta| \ll \delta^{\eta/2}$ so that for all $r \in I_\delta$, we have

$$\mu(F_{\text{bad}}(r)) \ll \delta^{\eta/2}$$

Let $C > 1$, and assume that there exists a subset $I_{\text{bad}} \subset [\frac{1}{2}, 1]$ with $|I_{\text{bad}}| \geq C\delta^{\eta/2}$ so that for all $r \in I_{\text{bad}}$, we have

$$\mu(F_{\text{bad}}(r)) \geq C\delta^{\eta/2}.$$

We will show this leads to a contradiction provided C is large enough.

Equip $[\frac{1}{2}, 1] \times F$ with the product measure $\text{Leb} \times \mu$. Let

$$\mathbf{E} = \{(r, w) \in [\frac{1}{2}, 1] \times F : m^\delta(\pi_r(w)) \geq C_0 e^{-t/10} \delta^{\alpha-18\eta}\},$$

and for every $w \in F$, let $\mathbf{E}_w = \{r : (r, w) \in \mathbf{E}\}$. The above then implies that

$$\text{Leb} \times \mu(\mathbf{E}) \geq C^2 \delta^\eta.$$

Set $F' = \{w \in F : |\mathbf{E}_w| \geq C\delta^\eta\}$. Then using Fubini's theorem, we conclude

$$\mu(F') \geq \frac{1}{2} C^2 \delta^\eta.$$

Moreover, in view of the definitions, for every $w \in F'$

$$|\{r \in [\frac{1}{2}, 1] : m^\delta(\pi_r(w)) \geq C_0 e^{-t/10} \delta^{\alpha-18\eta}\}| \geq C\delta^\eta$$

This contradicts Lemma 2.1 provided C is large enough.

The proof of complete. \square

3. PROOF OF PROPOSITION 1.2

Let us now turn to the proof of Proposition 1.2. As it was already mentioned, we will prove this theorem using [GGW22, Thm. 2.1]. Let us begin by recalling the notation used in [GGW22, Thm. 2.1] and Proposition 1.2.

Recall that for all $r \in [0, 1]$, we put $\xi(r) = (\frac{r}{1!}, \frac{r^2}{2!}, \dots, \frac{r^n}{n!}) \in \mathbb{R}^n$. For all $1 \leq k \leq n$, let $\mathfrak{p}_r^{(k)} : \mathbb{R}^n \rightarrow \mathbb{R}^k$ be the orthogonal projection onto the space spanned by $\{\xi'(r), \dots, \xi^{(k)}(r)\}$. That is:

$$\mathfrak{p}_r^{(k)}(w) = (w \cdot \xi'(r), w \cdot \xi^{(2)}(r), \dots, w \cdot \xi^{(k)}(r))$$

where $w \cdot v$ is the usual inner product on \mathbb{R}^n .

3.1. Theorem ([GGW22]). *Let $1 \leq k \leq n$ and let $0 < \alpha \leq k$. Let μ be the uniform measure on a finite set $F \subset B_{\mathbb{R}^n}(0, 1)$ satisfying*

$$\mu(B_r(w, \delta) \cap \Theta) \leq C_0 \delta^\alpha \quad \text{for all } w \text{ and all } \delta \geq \delta_0$$

where $C_0 > 0$.

Let $0 < \varepsilon < 10^{-4}\alpha$. For every $\delta \geq \delta_0$, there exists a subset $J_\delta \subset [0, 1]$ with $|[0, 1] \setminus J_\delta| \ll_\varepsilon \delta^{*\varepsilon^2}$ so that the following holds. Let $r \in J_\delta$, then there exists a subset $F_{\delta, r} \subset F$ with

$$\mu(F \setminus F_{\delta, r}) \ll_\varepsilon \delta^{*\varepsilon^2}$$

such that for all $w \in F_{\delta, r}$ we have

$$\mu(\{w' \in F : \|\mathfrak{p}_r^{(k)}(w) - \mathfrak{p}_r^{(k)}(w')\| \leq \delta\}) \ll_\varepsilon C_0 \delta^{\alpha - \varepsilon}$$

Proof. We deduce this from [GGW22, Thm. 2.1]. The argument is more or less standard. Indeed it is similar to the deduction of Theorem 1.1 from Lemma 2.1 and to (a finitary version of) the argument in [GGW22, §2].

Since $1 \leq k \leq n$ is fixed throughout the argument, we will denote $\mathfrak{p}_r^{(k)}$ by \mathfrak{p}_r . Adapting the notation m^δ from the previous section: for every $r \in [\frac{1}{2}, 1]$ and all $w \in F$, put

$$m^\delta(\mathfrak{p}_r(w)) = \mu\{w' : \|\mathfrak{p}_r(w) - \mathfrak{p}_r(w')\| \leq \delta\}.$$

Let D_1, \dots be large constants which will be explicated later. Let $\eta = \varepsilon/D_1$. For all $r \in [\frac{1}{2}, 1]$, let

$$F_{\text{bad}}(r) = \{w : m^\delta(\mathfrak{p}_r(w)) \geq C_0 \delta^{\alpha - D_1 \eta}\}.$$

Assume contrary to the claim in Theorem 3.1 that there exists a subset $I_{\text{bad}} \subset [\frac{1}{2}, 1]$ with $|I_{\text{bad}}| \geq D_2 \delta^{\eta^2/2}$ so that for all $r \in I_{\text{bad}}$, we have

$$\mu(F_{\text{bad}}(r)) \geq D_2 \delta^{\eta^2/2}.$$

We will get a contradiction with [GGW22, Thm. 2.1], provided that D_i 's are large enough.

First note that, for every $r \in I_{\text{bad}}$, the number of δ -boxes $\{\mathbf{B}_{i,r}\}$ required to cover $\mathbf{p}_r(F_{\text{bad}}(r))$ is $\leq D_3 C_0^{-1} \delta^{-\alpha+D_1\eta}$. Following [GGW22], let $\mathcal{T}_r = \{\mathbb{T}_{i,r}\}$ where $\mathbb{T}_{i,r} = \mathbf{p}_r^{-1}(\mathbf{B}_{i,r}) \cap B_{\mathbb{R}^n}(0, 1)$; note that

$$(3.1) \quad \#\mathcal{T}_r \leq D_3 C_0^{-1} \delta^{-\alpha+D_1\eta}.$$

Select a maximal δ -separated subset $\Lambda_\delta \subset I_{\text{bad}}$ and extend this to a maximal δ -separated subset $\hat{\Lambda}_\delta$ of $[\frac{1}{2}, 1]$. The definition of F_{bad}

Let ρ denote the uniform measure on $\hat{\Lambda}_\delta$, and as it was done in the proof of Theorem 1.1, equip $\hat{\Lambda}_\delta \times F$ with the product measure $\rho \times \mu$, and let

$$\begin{aligned} \mathbf{E} &= \{(r, w) \in \Lambda_\delta \times F : m^\delta(\pi_r(w)) \geq C_0 \delta^{\alpha-D_1\eta}\} \\ &= \{(r, w) \in \Lambda_\delta \times F : w \in F_{\text{bad}}(r)\}. \end{aligned}$$

Then the above implies that $\rho \times \mu(\mathbf{E}) \geq D_2^2 \delta^{\eta^2}$.

For every $w \in F$, let $\mathbf{E}_w = \{r \in \Lambda_\delta : (r, w) \in \mathbf{E}\}$, and set

$$F' = \{w \in F : \rho(\mathbf{E}_w) \geq D_2 \delta^{\eta^2}\}.$$

Then using Fubini's theorem, we conclude $\mu(F') \geq \frac{1}{2} D_2^2 \delta^{\eta^2}$.

Recall that ρ is the normalized counting measure on Λ_δ . The above definitions thus imply

$$\sum_{r \in \Lambda_\delta} 1_{\mathbb{T}_r}(w) \geq D_3 \delta^{\eta^2-1} \quad \text{for all } w \in F'$$

where $D_3 = O(D_2)$ and the implied constant is absolute. Let μ' denote the restriction of μ to F' . Now by [GGW22, Thm 2.1], applied with η^2 , μ' and $\{\mathcal{T}_r : r \in \Lambda_\delta\}$, we have

$$\begin{aligned} \sum_{r \in \Lambda_\delta} \#\mathcal{T}_r &\geq D_4(n, \varepsilon, \alpha) C_0^{-1} \mu'(\mathbb{R}^n) \delta^{-1-\alpha+D\eta} \\ &\geq \frac{1}{2} D_2^2 D_4(n, \varepsilon, \alpha) C_0^{-1} \delta^{\eta^2} \delta^{-1-\alpha+D\eta} \end{aligned}$$

where $D = 10^{10n}$ and in the second line we used $\mu'(\mathbb{R}^n) = \mu(F') \geq \frac{1}{2} D_2^2 \delta^{\eta^2}$. Thus there exists some $r \in \Lambda_\delta$ so that

$$(3.2) \quad \#\mathcal{T}_r \geq \frac{1}{2} C_0^{-1} D_4(n, \varepsilon, \alpha) D_2^2 \delta^{-\alpha+(D+1)\eta}.$$

Now comparing (3.2) and (3.1) we get a contradiction so long as D_1 is large enough and δ is small enough. The proof is complete. \square

We now turn to the proof of Proposition 1.2.

Proof of Proposition 1.2. Let δ and k be as in Proposition 1.2. As before, we let μ be the uniform measure on a finite set F .

Apply Theorem 3.1 with $\delta' := e^{-t}\delta \geq \delta_0$ and with k and $k-1$. Then for every $r \in J_\delta$ and $w \in F_{\delta',r}$, we have

$$(3.3) \quad \mu(\{w' \in F : \|\mathbf{p}_r^{(\ell)}(w) - \mathbf{p}_r^{(\ell)}(w')\| \leq e^{-t}\delta\}) \ll_\varepsilon C_0 (e^{-t}\delta)^{\min(\alpha, \ell) - \varepsilon}$$

for $\ell = k, k-1$.

Also note that if $0 < \alpha \leq k - 1$, then

$$\begin{aligned} \{w' : \|\pi_{t,r}(w) - \pi_{t,r}(w')\| \leq \delta\} \subset \\ \{w' \in F : \|\mathfrak{p}_r^{(k-1)}(w) - \mathfrak{p}_r^{(k-1)}(w')\| \leq e^{-t}\delta\}. \end{aligned}$$

Thus (3.3), with $\ell = k - 1$ implies

$$(3.4) \quad \mu(\{w' : \|\pi_{t,r}(w) - \pi_{t,r}(w')\| \leq \delta\}) \ll_{\varepsilon} C_0 e^{-\alpha t} \delta^{\alpha-\varepsilon}$$

as we claimed.

Assume now that $k - 1 < \alpha \leq k$. Let $Q \subset \text{Span}\{\xi'(r), \dots, \xi^{(k)}(r)\} \simeq \mathbb{R}^k$ denote the box $\delta' \times \dots \times \delta' \times \delta$ centered at the origin, where directions correspond directions of $\xi^{(i)}(r)$. That is: Q is box of size δ' in the direction of $\xi^{(i)}(r)$ for all $1 \leq i \leq k - 1$ and of size δ in the direction of $\xi^{(k)}(r)$. We cover

$$Q' := (Q + \pi_{t,r}(w)) \cap \pi_{t,r}(F_{\delta',r})$$

with $\ll e^t$ many boxes B_i of size δ' . Then (3.3), applied with $2\delta'$, implies

$$(3.5) \quad \begin{aligned} \mu(Q') &\ll_{\varepsilon} (C_0(2\delta')^{\alpha-\varepsilon}) \cdot e^t \\ &\ll_{\varepsilon} C_0 e^{(1-\alpha)t} \delta^{\alpha-\varepsilon}. \end{aligned}$$

Finally, note that

$$\{w' : \|\pi_{t,r}(w) - \pi_{t,r}(w')\| \leq \delta\} \subset Q + \pi_{t,r}(w).$$

This and (3.5) complete the proof in this case as well. \square

REFERENCES

- [BDG16] Jean Bourgain, Ciprian Demeter, and Larry Guth. Proof of the main conjecture in vinogradov's mean value theorem for degrees higher than three. *Annals of Mathematics*, 184(2):633–682, 2016.
- [FO14] Katrin Fässler and Tuomas Orponen. On restricted families of projections in \mathbb{R}^3 . *Proc. Lond. Math. Soc. (3)*, 109(2):353–381, 2014.
- [GGW22] Shengwen Gan, Shaoming Guo, and Hong Wang. A restricted projection problem for fractal sets in \mathbb{R}^n , 2022, 2211.09508.
- [KOV17] Antti Käenmäki, Tuomas Orponen, and Laura Venieri. A Marstrand-type restricted projection theorem in \mathbb{R}^3 , 2017, arXiv:1708.04859.
- [LM23] E. Lindenstrauss and A. Mohammadi. Polynomial effective density in quotients of \mathbb{H}^3 and $\mathbb{H}^2 \times \mathbb{H}^2$. *Invent. Math.*, 231(3):1141–1237, 2023.
- [LMW22] Elon Lindenstrauss, Amir Mohammadi, and Zhiren Wang. Effective equidistribution for some one parameter unipotent flows, 2022, 2211.11099.
- [LMWY23] Elon Lindenstrauss, Amir Mohammadi, Zhiren Wang, and Lei Yang. An effective version of the oppenheim conjecture with a polynomial error rate, 2023, 2305.18271.
- [Ohm23] K. W. Ohm. Projection theorems and isometries of hyperbolic spaces, 2023, 2305.12302.
- [OV18] Tuomas Orponen and Laura Venieri. Improved Bounds for Restricted Families of Projections to Planes in \mathbb{R}^3 . *International Mathematics Research Notices*, 2020(19):5797–5813, 08 2018.
- [PYZ22] Malabika Pramanik, Tongou Yang, and Joshua Zahl. A furstenberg-type problem for circles, and a kaufman-type restricted projection theorem in \mathbb{R}^3 , 2022.

- [Sch03] Wilhelm Schlag. On continuum incidence problems related to harmonic analysis. *Journal of Functional Analysis*, 201:480–521, 07 2003.
- [Wol00] T. Wolff. Local smoothing type estimates on L^p for large p . *Geom. Funct. Anal.*, 10(5):1237–1288, 2000.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SAN DIEGO, CA 92093
Email address: `kwohm@ucsd.edu`