The off-equilibrium Kinetic Ising model: The Metric Case

Luca Di Carlo $^{1,\,2}$

¹Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544, USA ²Lewis-Sigler Institute for Integrative Genomics

We investigate the critical behavior of the Kinetic Ising model with non-reciprocal nearest neighbors interactions. A finite-size scaling study suggests that the model belongs to the Ising universality class. We characterize the off-equilibrium behavior of the model by measuring the entropy production rate and by studying the coarsening dynamics, which shows a super-diffusive relaxation. A coarse-grained equation of motion is in compelling agreement with the numerical findings. A one-loop renormalization group calculation shows that the model belongs to the Model A universality class.

Universality is a fundamental concept in the study of critical phenomena and collective behavior. Universality refers to the observation that different systems can exhibit the same collective behavior near their critical points. This means that diverse systems, despite having distinct microscopic interactions and details, share identical critical behavior characterized by the same critical exponents and scaling laws [1]. This allows for a unified understanding and a classification of critical phenomena across various physical contexts, thereby highlighting the underlying simplicity and coherence in the complexity of phase transitions. The renormalization group (RG) theory provides a robust framework to understand this scale invariance and universality at critical points.

Extending the theory of critical phenomena to the realm of off-equilibrium systems presents a hard and yet necessary challenge. Nonequilibrium systems, such as those found in biological contexts, constantly operate far from thermodynamic equilibrium, and are characterized by constant heat dissipation and entropy production. Investigating critical phenomena in this context could provide deeper insights into the fundamental principles of self-organization, pattern formation, and robustness in living systems [2–6].

In recent decades several minimal models of nonequilibrium collective dynamics have been proposed [7–10]. In this work, we study a simple off-equilibrium generalization of the Ising model, the Kinetic Ising model [11]. In constrast to the traditional Ising model which is defined by its steady-state Boltzmann probability distribution

$$P(\boldsymbol{\sigma}) = \frac{1}{Z} e^{\sum_{ij} J_{ij}\sigma_i\sigma_j} \tag{1}$$

the Kinetic Ising model is defined by the update probability,

$$P(\boldsymbol{\sigma}_{t+1}|\boldsymbol{\sigma}_t) = \prod_{i} \frac{e^{h_i^t \boldsymbol{\sigma}_i^{t+1}}}{2\cosh\left(h_i^t\right)}$$
(2)

where the local field is defined as $h_{i}^{t} = \sum_{j} J_{ij}\sigma_{j}^{t}$. It is worth noting the similarity between Eq. (2) and the Markov Chain Monte Carlo update rule used to sample the Ising probability distribution (1). The key difference between (1) and (2) lies in the symmetry properties of the interaction matrix J_{ij} . While in (1) the interactions J_{ij} must be symmetric - or more precisely only the symmetric part of J_{ij} is relevant - in the kinetic Ising model there are no constraints on the symmetry of the interactions J_{ij} . If J_{ij} is symmetric the Markov chain defined by (2) reproduces the Ising equilibrium probability distribution (1). The fully connected Kinetic Ising model has been extensively studied in the literature [11, 12]. In this work we study the asymmetric Ising model with metric interactions; more specifically we focus on a two-dimensional Ising model with non-reciprocal nearest neighbor interactions. We show that, albeit being off-equilibrium, the system belongs to the same static universality class of the equilibrium Ising model. We characterize the off-equilibrium behavior of the system by computing the steady-state entropy production rate and by studying the coarsening dynamics. which is significantly affected by the non-reciprocal interactions. Finally, we study the corresponding coarsegrained field theory, which is in compelling agreement with the numerical results.

THE ASYMMETRIC METRIC ISING MODEL

We consider Ising spins $\sigma_{ij} = \pm$ on a two-dimensional lattice, interacting through a symmetric interaction along the y-axis and an asymmetric interaction along the x-axis:

$$J_{(\mathbf{i}_{\mathbf{x}},\mathbf{i}_{\mathbf{y}}),(\mathbf{i}_{\mathbf{x}},\mathbf{i}_{\mathbf{y}})} = J\delta_{\mathbf{i}_{\mathbf{x}},\mathbf{j}_{\mathbf{x}}} \left[\delta_{\mathbf{i}_{\mathbf{y}},\mathbf{j}_{\mathbf{y}}+1} + \delta_{\mathbf{i}_{\mathbf{y}},\mathbf{j}_{\mathbf{y}}-1} \right] + J\delta_{\mathbf{i}_{\mathbf{x}},\mathbf{j}_{\mathbf{y}}} \left[\delta_{\mathbf{i}_{\mathbf{x}},\mathbf{j}_{\mathbf{x}}-1} + \alpha\delta_{\mathbf{i}_{\mathbf{x}},\mathbf{j}_{\mathbf{x}}+1} \right]$$
(3)

where the parameter α controls the strength of the asymmetry. We call this model the Asymmetric Metric Ising (AMI) model. For $\alpha = 1$ we recover the equilibrium twodimensional Ising model, while for any $\alpha \neq 1$ the system is out of thermodynamic equilibrium. Similarly to what happens in the equilibrium Ising model, we expect a phase transition from disorder to order as we increase the interaction strength J. The level of asymmetry plays



FIG. 1. Finite-size scaling analysis. (a1) The binder cumulant [13] as a function of the coupling J for different values of the off-equilibrium parameter α . Different colors corresponds to different values of α , the color gradient indicates the size of the system ranging from L = 16 to L = 64. The critical value of the coupling J_c depends on the value of α . It shifts from its equilibrium value ($\alpha = 1$) $J_c \simeq 0.442$ to $J_c \simeq 0.7$ for $\alpha = 0$. (a2) Collapse of the binder cumulant with $\nu = 1.0$. (b1, b2) Data collapse of the magnetization with the 2dimensional Ising critical exponents $\beta = 0.125$. (c1, c2) Collapse of the susceptibility with the two-dimensional Ising critical exponent $\eta = 0.25$. While the value of the critical coupling J_c depends on α , the measure critical exponents and the scaling functions seems to be the same as the equilibrium Ising model independently on how far the model is from equilibrium. The scaling of the susceptibility in the ordered phase seems to show some dependence on the value of α .

an important role, in fact if the asymmetry is too strong $(\alpha < 0)$ the system might present dynamical instabilities. For instance, in a system made of just two spins interacting with $J_{12} = -J_{21}$, the spin σ_1 wants to point in the same direction as σ_2 , while σ_2 wants to point in the opposite direction of σ_1 ; this leads to an oscillatory behavior. We expect the phenomenology of the $\alpha < 0$ phase to be very rich. In this work we focus on the ferromagnetic part of the phase diagram, corresponding to $\alpha \ge 0$, and leave the $\alpha < 0$ regime to future studies. In the $\alpha \ge 0$ regime we expect the system to exhibit a ferromagnetic phase transition.

We first address the effect of the asymmetry on the static critical properties of the model - namely we consider only observables measured at the same time. As expected, above a critical coupling J_c the system exhibits a phase transition from a disordered to an ordered phase. As show in Fig. 1 (a1, a2, a3), the critical value of the coupling J_c depends on the strength of asymmetry, going from its equilibrium value ($\alpha = 1$) $J_c \simeq 0.442$ to $J_c \simeq 0.7$ for $\alpha = 0$. A finite-size scaling analysis shows that the static critical exponents and the scaling functions are the same as the equilibrium Ising model, suggesting that the asymmetry does not affect the static universality class of the model. Although the equal time behavior is not affected by the asymmetry, this does not mean that the model is at equilibrium. The system's steady state violates both parity and time-reversal symmetries, as it is evident by inspecting the systems's trajectories [14]. If we interpret the spins as the occupation of a lattice site

by a particle, the asymmetry generates a constant flux of particles in the direction of the asymmetry. We can quantify this flux using the following quantity,

$$\Phi = \frac{1}{L^2} \sum_{\mathbf{x}, \mathbf{y}} \langle \sigma_{\mathbf{x}+1, \mathbf{y}}(t) \sigma_{\mathbf{x}, \mathbf{y}}(t + \Delta t) \rangle - \langle \sigma_{\mathbf{x}-1, \mathbf{y}}(t) \sigma_{\mathbf{x}, \mathbf{y}}(t + \Delta t) \rangle$$
(4)

which measures the average difference in probability between hopping to the right and hopping to the left. Interestingly, Φ coincides, up to a numerical factor, with the entropy production rate of the system [15–17]. The per-particle entropy production rate can be written as [11],

$$\dot{\mathcal{S}} = \frac{1}{L^2} \sum_{ij} J_{ij} \left(\left\langle \sigma_i^t \sigma_j^{t-1} \right\rangle - \left\langle \sigma_j^t \sigma_i^{t-1} \right\rangle \right) = (1 - \alpha) \Phi \quad (5)$$

This equation has a clear interpretation. The energy that is dissipated by the system to maintain the offequilibrium steady-state is the product of the "net force" on the x axis $(1 - \alpha)$ and the 'velocity' Φ along the same direction. We expect the flux Φ to vanish at equilibrium $\alpha = 1$, and at the leading order we have $\Phi \simeq f(J, L)(1 - \alpha)$. Therefore the entropy production is quadratic in the deviation from equilibrium $(1 - \alpha)$,

$$\dot{\mathcal{S}} = f(J,L)(1-\alpha)^2 \tag{6}$$

ensuring that \dot{S} is positive. In Fig. 2 (b,c) we show the entropy production rate for different values of α . In agreement with the previous literature the entropy production



FIG. 2. Entropy production rate. (a) - The entropy production rate of the model peaks around the transition point, unlike what observed in the other moles the entropy production rate is not cusped at the transition point. The color gradient indicates the system's size L = 16, 24, 32, 64. (b) -The entropy production rate seems to be well described by the ansatz (6), suggesting that \dot{S} is a simple quadratic function of the off-equilibrium parameter $(1 - \alpha)$. (c) - Peak of the entropy production for increasing values of the system size L. The curve $\dot{S}(J)$ seems converge to a smooth function.

peaks around the transition point [18, 19]. In particular, \dot{S} is maximum for J slightly below the critical coupling J_c ; this does not seem to be a finite-size effect, as shown in Fig. 2 (c). Fig. 2(c) shows that the ansatz (6) well describes what is observed in the numerical simulations. As opposed to the entropy production rate of other off-equilibrium model, such as the Vicsek model [19] or the Active Ising model [18], which is cusped at the transition point, we observer a smooth entropy production for $J \simeq J_c$, as shown in Fig. 2 (c).

An other macroscopic consequence of the non reciprocity of the interactions regards the coarsening dynamics, which is the process of relaxation to equilibrium through the growth of domains of aligned spins. To study the coarsening dynamics we consider a system in the low temperature phase, initialized in a configuration in which all the spin inside a ball of radius ℓ_0 are pointing up, and all the spins outside the ball are pointing down. We study the coarsening dynamics by measuring how long does it take for the system to relax back to $m \simeq -1$. According to the equilibrium theory of ordering kinetics [20] the positively magnetized region shrinks

linearly in time, $\ell(t) = \sqrt{\ell_0^2 - At}$, implying that the characteristic time scales to relax to the equilibrium state τ grows quadratically with the size of the magnetized region $\tau \sim \ell_0^2$. Essentially, at equilibrium the relaxation process is purely diffusive. In Fig. 3 (a) we show the coarsening dynamics of the system in the ordered phase, $J = \frac{3}{2}J_c$, for $\alpha = 1$. On the contrary, When the system is off-equilibrium $\alpha = 0$, the relaxation dynamics is significantly faster (see coarsening.gif in the supplemental material), and the magnetization does not decrease linearly in time, suggesting a super-diffusive behavior. In Fig. 3 (c) we show the scaling of the relaxation time as a function of the size of the magnetized region ℓ_0 . The relaxation time follows a scaling relation $\tau \sim \ell_0^{z_c}$, with a coarsening exponent z_c that goes from $z_c = 2$ at equilibrium, in agreement with the theory of ordering kinetics, to lower and lower values as we move away from the equilibrium regime, As we move away from $\alpha = 1$, the relaxation dynamics seems to approach a balistic regime $z_{\rm c} = 1.$

COARSE-GRAINED FIELD THEORY

The coarse-grained equation of motion of the AMI can be derived by symmetry arguments. The minimal coarsegrained model compatible with the symmetries of the model takes the following form,

$$\partial_t \varphi = \Gamma_0 \nabla^2 \varphi + v_0 \partial_x \varphi - (m_0 + \lambda_0 \varphi^2) \varphi + \xi \tag{7}$$

where $\xi(x,t)$ is a delta-correlated Gaussian noise, $\langle \xi(x,t)\xi(x',t')\rangle = 2\tilde{\Gamma}_0\delta(x-x')\delta(t-t')$. The term $v_0\partial_x\varphi$ stems for the asymmetric interaction along the x direction. When $v_0 = 0$ the equation of motion (7) coincides with Model A[21], which describes the relaxation dynamics of the equilibrium Ising model. In the Gaussian approximation, the correlation function of the AMI is given by,

$$C_0(k,\omega) = \frac{2\tilde{\Gamma}_0}{(\omega + v_0 k_x)^2 + (\Gamma_0 k^2 + m_0)^2} \qquad , \qquad (8)$$

and the static Gaussian correlation function $C_0(\mathbf{k}, t=0)$ is,

$$C_0(\mathbf{k}, t=0) = \frac{\tilde{\Gamma}_0}{\Gamma_0 k^2 + m_0} \qquad . \tag{9}$$

This suggests that the static properties of the model are independent of the value of the macroscopic parameter v_0 . This results relies only on the fact that the off-equilibrium parameter v_0 appears in the combination $\omega + v_0 k_x$; when computing the Fourier transform of (8) at t = 0 we can redefine the integration variable as $\omega = \omega' - v_0 k_x$ to reabsorb the term proportional to



FIG. 3. Study of the coarsening dynamics (a) - Coarsening dynamics of the system in the ordered phase, $J = \frac{3}{2}J_c$, for $\alpha = 1$. For $\alpha = 1$ the system is at equilibrium and the coarsening dynamics is diffusive. Since we are considering a two-dimensional system, the magnetization relaxes linearly to its steady state value. Results are averaged over 20 different trials, error bars have been obtained as the standard deviation over the trials (b) - Coarsening dynamics in the ordered phase, $J = \frac{3}{2}J_c$, for $\alpha = 0$, the statistics of equal time observable of this model matches the one of the $\alpha = 1.0$ system showed in panel(a). The coarsening dynamics of the off-equilibrium model ($\alpha = 0$) is significantly faster (see the coarsening.gif file in the SM). The magnetization is not decreasing linearly, suggesting a super-diffusive behavior. (c) - Scaling of the time it takes for the system to system to relax back to the steady state when the system is initialized in a configuration in which all the spin inside a ball of radius ℓ_0 are pointing up, and all the spins outside the ball are pointing down. Data are relative to $J = \frac{3}{2}J_c$. As we can see the relaxation time follows a scaling relation $\tau \sim \ell_0^{Z_c}$, where the exponent z_c goes from $z_c = 2$ at equilibrium to lower and lower values as we move away from equilibrium. Data are relative to a system of size L = 256.

 v_0 . Since this results relies only on the form of the Correlation function, we expect it to hold also at the non-Gaussian level. A simple renormalization group calculation shows that the static critical properties are described by the equilibrium model A critical point (see SM) [21].

The presence of the term $v_0 \partial_x \varphi$ in the correlation function (8) suggests a non exponential relaxation; the Gaussian correlation function in the (\mathbf{k}, t) domain reads,

$$C_0(\boldsymbol{k}, t) = \operatorname{Re}\left(\frac{\tilde{\Gamma}_0}{\Gamma_0 \boldsymbol{k}^2 + m_0} e^{+i\omega_{\boldsymbol{k}}t}\right)$$
(10)

where the frequency $\omega_{\mathbf{k}}$ is defined as,

$$\omega_{\boldsymbol{k}} = i\left(m_0 + \Gamma \boldsymbol{k}^2\right) + v_0 \boldsymbol{k}_x \tag{11}$$

In Fig. 4 we show the real and imaginary part of the characteristic frequency measured in numerical simulations close to the critical . In agreement with a renormalization group calculation (see SM), close to the critical point $\omega_{\mathbf{k}}$ is well described by the Gaussian theory $\operatorname{Re}(\omega_{\mathbf{k}}) \sim k^2$ and $\operatorname{Im}(\omega_{\mathbf{k}}) \sim k$.

We finally compute the intensive entropy production rate in the Gaussian approximation. The entropy production rate can be computed by using the following Harada-Sasa relation [22, 23], relating the entropy production rate to the violation of the fluctuation dissipation relations,

$$\mathcal{S}(k,\omega) = \frac{1}{L^d} \frac{L^d \omega}{(2\pi)^{d+1} \tilde{\Gamma}_0} \left[\omega C(\boldsymbol{k},\omega) - 2 \tilde{\Gamma}_0 \operatorname{Im} \left(R(\boldsymbol{k},\omega) \right) \right]$$
(12)

At the Gaussian level this formula leads to,

$$\dot{\mathcal{S}}(\boldsymbol{k}) = \frac{v_0^2 k_x^2}{(2\pi)^d} \frac{1}{\Gamma_0 k^2 + m_0}$$
(13)

This simple calculation is in agreement with the the numerical results shown in Fig. 2. The entropy production rate is a quadratic function of the off-equilibrium parameter $v_0 \simeq (1-\alpha)$, and it peaks around the transition point $m_0 = 0$. The k_x^2 factor in the numerator ensures that the entropy production rate $\dot{S}(\mathbf{k} \to 0)$ remains finite also at the critical point $m_0 = 0$. The L^d factors in Eq. (12) cancel out and the intensive/per-particle entropy production rate is independent of the system size.

DISCUSSION

We have introduced the Asymmetric Metric Ising model, a two-dimensional Ising model with nonreciprocal interactions, and we have studied its critical properties. As its equilibrium counterpart the system exhibits a ferromagnetic phase transition. A numerical finite-size study shows that the model belongs to the



FIG. 4. Real and imaginary part of the characteristic frequency $\omega_{\mathbf{k}}$ measured in the numerical simulations. The frequency is obtained by fitting Eqs. (10) and (11). The scaling of the real and imaginary parts of $\omega_{\mathbf{k}}$ are in agreement with the Gaussian theory, $\operatorname{Re}(\omega_{\mathbf{k}}) \sim k^2$ and $\operatorname{Im}(\omega_{\mathbf{k}}) \sim k$, as predicted by the renormalization group calculation (see SM). Data are relative to a system with $\alpha = 0.5$, $J = 0.99J_c$ of size L = 64. More details on how the characteristic frequency is measured can be found in the SM.

Ising model (Model A) universality class, while being offequilibrium. As observed in similar contexts [24, 25], belonging to an equilibrium static universality class does not necessarily imply that the model is at equilibrium; In fact, the steady state entropy production rate of the model is non zero and it peaks close to the transition point. Unlike observed in other contexts, the entropy production rate is smooth close to the transition point, as opposed to what observed in other off-equilibrium cases. We characterized the coarsening dynamics in the low temperature phase, observing a significant speed up due to the non-reciprocal interactions. In particular, we observe a super-diffusive relaxation to steady state. We finally studied the coarse-grained equation of motion of the model, and showed that at the one-loop level the model belongs to the Model A universality class. The off-equilibrium parameter v_0 does not take any renormalization group corrections, correctly predicting the scaling of the characteristic frequency measured in the numerical simulations. We computed the entropy production rate of the model using a generalized Harada-Sasa relation. The numerical results are in excellent agreement with the theoretical predictions. This worked showed how non trivial behavior can emerge when combining the non-reciprocal interactions o with a metric structure, we expect that the strongly asymmetric regime $\alpha < 0$ to have an even richer phenomenology. We are confident that the study of this simple system might give insights on the behavior of more complicated off-equilibrium phase transition. We believe that the phenomenology of the AMI will find

application in describing real biological system, in particular the super-diffusive relaxation dynamics and the emergent collective transport.

This work was supported by the National Science Foundation, through the Center for the Physics of Biological Function (PHY–1734030). I would like to thank Q. Yu for comments on the manuscript and M. Scandolo and C.W. Lynn for useful discussions.

- M. Kardar, Statistical physics of fields (Cambridge University Press, 2007).
- [2] M. C. Cross and P. C. Hohenberg, Pattern formation outside of equilibrium, Rev. Mod. Phys. 65, 851 (1993).
- [3] M. C. Marchetti, J.-F. Joanny, S. Ramaswamy, T. B. Liverpool, J. Prost, M. Rao, and R. A. Simha, Hydrodynamics of soft active matter, Reviews of modern physics 85, 1143 (2013).
- [4] M. Fruchart, R. Hanai, P. B. Littlewood, and V. Vitelli, Non-reciprocal phase transitions, Nature 592, 363 (2021).
- [5] E. Montbrió and D. Pazó, Kuramoto model for excitation-inhibition-based oscillations, Phys. Rev. Lett. 120, 244101 (2018).
- [6] M. N. Rouches and B. B. Machta, Polymer collapse & liquid-liquid phase-separation are coupled in a generalized prewetting transition, ArXiv (2024).
- [7] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, and O. Shochet, Novel type of phase transition in a system of self-driven particles, Phys. Rev. Lett. **75**, 1226 (1995).
- [8] A. P. Solon and J. Tailleur, Flocking with discrete symmetry: The two-dimensional active ising model, Phys. Rev. E 92, 042119 (2015).
- [9] F. Caballero, C. Nardini, F. van Wijland, and M. E. Cates, Strong coupling in conserved surface roughening: A new universality class?, Physical review letters **121**, 020601 (2018).
- [10] B. VanSaders and V. Vitelli, Informational active matter, arXiv preprint arXiv:2302.07402 (2023).
- [11] M. Aguilera, S. A. Moosavi, and H. Shimazaki, A unifying framework for mean-field theories of asymmetric kinetic ising systems, Nature communications 12, 1197 (2021).
- [12] H. J. Kappen and J. J. Spanjers, Mean field theory for asymmetric neural networks, Phys. Rev. E 61, 5658 (2000).
- [13] K. Binder, Finite size effects at phase transitions, in Computational Methods in Field Theory: Proceedings of the 31. Internationale Universitätswochen für Kernund Teilchenphysik Schladming, Austria, February 1992 (Springer, 2005) pp. 59–125.
- [14] See trajectory.gif in the supplemental material.
- [15] G. E. Crooks, Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences, Phys. Rev. E 60, 2721 (1999).
- [16] U. Seifert, Stochastic thermodynamics, fluctuation theorems and molecular machines, Reports on progress in physics 75, 126001 (2012).
- [17] C. Nardini, E. Fodor, E. Tjhung, F. van Wijland, J. Tailleur, and M. E. Cates, Entropy production in

field theories without time-reversal symmetry: Quantifying the non-equilibrium character of active matter, Phys. Rev. X 7, 021007 (2017).

- [18] Q. Yu and Y. Tu, Energy cost for flocking of active spins: The cusped dissipation maximum at the flocking transition, Phys. Rev. Lett. **129**, 278001 (2022).
- [19] F. Ferretti, S. Grosse-Holz, C. Holmes, J. L. Shivers, I. Giardina, T. Mora, and A. M. Walczak, Signatures of irreversibility in microscopic models of flocking, Phys. Rev. E **106**, 034608 (2022).
- [20] A. J. Bray, Theory of phase-ordering kinetics, Advances in Physics 51, 481 (2002).
- [21] P. C. Hohenberg and B. I. Halperin, Theory of dynamic critical phenomena, Reviews of Modern Physics 49, 435 (1977).
- [22] T. Harada and S.-i. Sasa, Equality connecting energy dissipation with a violation of the fluctuation-response re-

lation, Phys. Rev. Lett. 95, 130602 (2005).

- [23] C. Nardini, E. Fodor, E. Tjhung, F. van Wijland, J. Tailleur, and M. E. Cates, Entropy production in field theories without time-reversal symmetry: Quantifying the non-equilibrium character of active matter, Phys. Rev. X 7, 021007 (2017).
- [24] F. Caballero and M. E. Cates, Stealth entropy production in active field theories near ising critical points, Phys. Rev. Lett. **124**, 240604 (2020).
- [25] F. Caballero, C. Nardini, and M. E. Cates, From bulk to microphase separation in scalar active matter: a perturbative renormalization group analysis, Journal of Statistical Mechanics: Theory and Experiment **2018**, 123208 (2018).