

ALGEBRAIC DEPENDENCE NUMBER AND CARDINALITY OF GENERATING ITERATED FUNCTION SYSTEMS

JUNDA ZHANG

ABSTRACT. For a dust-like self-similar set (generated by IFSs with the strong separation condition), Elekes, Keleti and Máthé found an invariant, called ‘algebraic dependence number’, by considering its generating IFSs and isometry invariant self-similar measures. We find an intrinsic quantitative characterisation of this number: it is the dimension over \mathbb{Q} of the vector space generated by the logarithms of all the common ratios of infinite geometric sequences in the gap length set, minus 1. With this concept, we present a lower bound on the cardinality of generating IFS (with or without separation conditions) in terms of the gap lengths of a dust-like set. We also establish analogous result for dust-like graph-directed attractors on complete metric spaces. This is a new application of the ratio analysis method and the gap sequence.

CONTENTS

1. Introduction	1
2. Preliminaries	3
2.1. Gap lengths	3
2.2. Ratio analysis	4
3. Our Results	4
3.1. Using ratio analysis to obtain logarithmic commensurability	4
3.2. Intrinsic characterisation of algebraic dependence number	5
3.3. Lower bound on the cardinality of generating IFS	6
References	7

1. INTRODUCTION

We recall some basic concepts in fractal geometry. In this paper, a finite set of distinct contractive similarities $\{S_i\}_{i=1}^m$ on a complete metric space (M, d) is called an *iterated function system* (IFS), where for each i , $d(S_i(x), S_i(y)) = r_i d(x, y)$ ($0 < r_i < 1$ is the *contraction ratio* of S_i). According to [27], the *attractor* of the IFS is the unique nonempty compact set $K \subset M$ such that

$$K = \bigcup_{i=1}^m S_i(K), \quad (1.1)$$

which is called a *self-similar set*. We call $\{S_i\}_{i=1}^m$ a *generating IFS* for K . We say that the *strong separation condition* (SSC) holds for the IFS when the union is disjoint in (1.1), and such K is called *dust-like*.

This paper is motivated by the inverse fractal problem in [11, 12, 20, 29]. Given a dust-like set, what can be said about its generating IFSs? This problem is also related to tiling theory and image compression, see [20, Section 1] for details. Our observation is that, the gap lengths of a dust-like

Date: August 22, 2024.

2020 Mathematics Subject Classification. 28A80, 05C20.

Key words and phrases. self-similar set, graph-directed attractor, algebraic dependence number, strong separation condition, generating IFS.

set can determine its *algebraic dependence number* (Theorem 3.6), which gives a lower bound on the cardinality of the generating IFS (Corollary 3.8). In [15], the algebraic dependence number of an IFS is defined as the dimension over \mathbb{Q} of the vector space generated by the logarithms of the contraction ratios of all similarities, minus 1. For a dust-like self-similar set in \mathbb{R}^n , [15, Theorem 5.7] shows that all its generating IFSs with the SSC have the same algebraic dependence number, so this number can be defined for a dust-like self-similar set as its intrinsic characteristic (independent of generating IFSs). This is done by considering certain measures on the dust-like self-similar set, while our characterisation is direct and intrinsic (without considering generating IFSs and measures), and we generalise this to compact metric spaces.

We also consider graph-directed IFSs and graph-directed attractors based on a given directed graph, which generalizes the concept of IFS and self-similar sets. It is introduced in [32], and have been studied a lot including the geometric aspects (see for example [8, 9, 14, 18, 22, 34, 38]) and fractal analysis [4, 6, 24, 33]. A directed graph (V, E) consists of a finite set of vertices V and a finite set of directed edges E with loops and multiple edges allowed. Given a directed graph, we associate each vertex $v \in V$ with a metric space (M^v, d^v) . Let $E_{uv} \subset E$ be the set of edges from vertex u to v . A graph-directed iterated function system (GD-IFS) $\{S_e : e \in E_{uv}\}$ consists of contracting similarities S_e from M_v to M_u , that is,

$$d^u(S_e(x), S_e(y)) = r_e d^v(x, y) \quad (0 < r_e < 1) \quad (1.2)$$

for all $x, y \in M^v$, where r_e is the contraction ratio of S_e . We will assume $d_u \geq 2$ for all $u \in V$ to avoid singletons, where d_u is the number of directed edges leaving u (see related discussion in [14, pp.607]). For a GD-IFS $(V, E, (S_e)_{e \in E})$ based on such a directed graph, there exists a unique list of non-empty compact sets $(F_u \subset M^u)_{u \in V}$ such that, for all $u \in V$,

$$F_u = \bigcup_{v \in V} \bigcup_{e \in E_{uv}} S_e(F_v), \quad (1.3)$$

see [13, Theorem 4.3.5 on p.128]. We call the above $(F_u)_{u \in V}$ the (*list of*) *attractors* of the GD-IFS, and each F_u is called a *GD-attractor*. We say that the *strong separation condition* (SSC) holds for the GD-IFS, if the union is disjoint in (1.3) for each $u \in V$, and the GD-attractors are called *dust-like*.

The graph-directed setting appear frequently in dynamical systems and the study of self-similar sets. Certain complex dynamical systems can be regarded as conformal GD-IFSs using a Markov partition [16, Section 5.5]. The orthogonal projection of certain self-similar sets may be GD-attractors [17, Theorem 1.1]. When exact overlap occurs in an IFS, one can use a corresponding graph-directed system to study its attractor (see for example [10, 26]). Also in fractal analysis, GD-IFSs are used to determine whether the two walk-dimensions coincide for p.c.f. self-similar sets [23]. They are also related to tiling problems [36] and Automata [7].

Our results are useful for *inhomogeneous* (GD-)IFSs and (GD-)attractors, where the contraction ratios are not all equal. The idea is applying the ‘ratio analysis’ method in [28] to the ‘gap length set’ of a dust-like GD-attractor. Separation conditions are often required in fractal geometry to obtain some precise structure information, and relaxing them can be very difficult in many problems. In our paper, the SSC is (only) used for a precise formula of the gap length set [10, Theorem 2] for ratio analysis. We remark that the SSC is a common assumption for inhomogeneous IFSs in related problems (see for example [1, 15, 21]). In Corollary 3.10 we use the ‘full-measure’ condition as in [20, Theorem 4.1] to remove the SSC condition.

This paper is organised as follows. In Section 2, we recall the definition and expression of the gap length set of dust-like GD-attractors, and the required ratio analysis lemmas in previous studies. In Section 3, we present our results. In Section 3.1, we use ratio analysis on the gap length set to obtain some lemmas and the logarithmic commensurability for SSC generating IFSs (Theorem 3.2), which gives an alternative proof of [15, Theorem 5.7]. In Section 3.2, we present

Theorem 3.6 which characterizes the algebraic dependence number of (GD-)IFS in terms of (the geometric sequences in) the gap length set. In Section 3.3, we present the lower bound estimate on the cardinality of generating IFS, with or without separation conditions (Corollaries 3.8 and 3.10).

2. PRELIMINARIES

2.1. Gap lengths. The concept and lemma in this subsection are from [10], which concerns the ‘gap sequence’ (see [35] for details in \mathbb{R}^n), a natural way to define the size and number of holes for disconnected sets. For a compact metric space (K, d) and $\delta > 0$, denote its diameter by $\text{diam}(K)$. We say that $x, y \in K$ are δ -equivalent, if there is a δ -chain connecting x and y , that is, there are points $x_i \in K$ ($i = 1, 2, \dots, t$) such that $x_1 = x, x_t = y, d(x_i, x_{i+1}) \leq \delta$ for all i . According to this equivalence, we can divide K into several δ -equivalence classes (intuitively, these small pieces are separated by a distance δ , while any two points in each piece can be connected by some δ -chain). Let $\kappa(\delta)$ be the number of δ -equivalence classes of K , which is finite due to the compactness of K (since two points in an open $\delta/2$ -metric ball are automatically δ -equivalent, $\kappa(\delta)$ does not exceed the covering number of K using open $\delta/2$ -metric balls).

Definition 2.1 (Gap length set). *The discontinuous points of the function $\kappa(\delta)$ ($0 < \delta < \text{diam}(K)$) are called the gap lengths of K , and the collection of all the gap lengths of K is called the gap length set of K , denoted by $\text{GL}(K)$ (that is, the set of discontinuous points of κ).*

Remark 2.2. *For compact sets on \mathbb{R} , this definition is intuitive and coincides with [28, Definition 2.1], see explanation and an example (middle-third Cantor set) in [35, Section 1].*

Note that the function $\kappa(\delta)$ is non-increasing in $\delta \in (0, \text{diam}(K))$, since a δ -equivalence class must be a δ' -equivalence class when $\delta < \delta'$. Thus the gap lengths are at-most countably many, and arranging them with corresponding multiplicity (according to the value of κ) gives the gap sequence, but we do not concern such multiplicity in our paper.

Definition 2.3 (Contraction ratio set). *The contraction ratio set of a GD-IFS is defined to be the set of the contraction ratios of the similarities, that is, $\{r_e : e \in E\}$ using the notion in (1.2).*

The following Lemma is a direct corollary of [10, Theorem 2]. We define the product of $A, B \subset \mathbb{R}$ to be $AB = \{ab : a \in A, b \in B\}$. We regard a real number as a set when encountering the product with a set in \mathbb{R} , and AB is empty if A is empty. Given a GD-IFS and a directed path $\mathbf{e} = e_1 e_2 \cdots e_k$ with edges e_i , (for which the terminal vertex of e_i is the initial vertex of e_{i+1} when $i = 1, \dots, k-1$), define $r_{\mathbf{e}} := r_{e_1} r_{e_2} \cdots r_{e_k}$.

Lemma 2.4. *Let $G = (V, E)$ be a directed graph satisfying $d_w \geq 2$ for all $w \in V$, and let $(M^u)_{u \in V}$ be a list of complete metric spaces. Suppose that for some $u \in V$, $F_u \subset M^u$ is the GD-attractor of some GD-IFS (based on G) with the SSC. Then for each $u \in V$, there exist finite sets $\Lambda_u, \Gamma_u \subset (0, \infty)$ such that*

$$\text{GL}(F_u) = \Gamma_u \bigcup \Lambda_u \bigcup \left(\bigcup_{v \in V} \Lambda_v R_{uv} \right) \text{ where } R_{uv} := \{r_{\mathbf{e}} : \mathbf{e} \text{ is a directed path from } u \text{ to } v\}. \quad (2.1)$$

Remark 2.5. *When there is no directed path from u to v , then $R_{uv} = \emptyset$. We mention that [10, Theorem 2] gives the construction of Λ_u, Γ_u , although it is not used in our paper. Given a GD-IFS and its attractor list $(F_u \subset M^u)_{u \in V}$, we obtain $\text{GL}(F_u)$ and define $\Gamma_u = \text{GL}(F_u) \cap [\delta, \infty)$ for all $u \in V$, where $\delta := \inf \{d^u(x_e, y_e) : x_e \in S_e(F_v), y_e \in F_u \setminus S_e(F_v), e \in E_{uv}, u, v \in V\} > 0$. Then define $\Lambda_u = \bigcup_{u \in V} \bigcup_{e \in E_{uv}} r_e (\Gamma_v \cap [\delta, r_e^{-1} \delta))$ for each $u \in V$, from which we know $\Lambda_u \subset (0, \delta)$.*

We will use the following notions in [28, Definition 2.5] and a corollary of the above lemma.

Definition 2.6. For a finite set $A = \{a_i\}_{i=1}^n \subset (0, \infty)$, define $A^{\mathbb{Z}^+}$ (resp. $A^{\mathbb{Q}^+}$) to be the union of all products $\prod_{i=1}^n a_i^{m_i}$ where $(m_i)_{i=1}^n$ are non-zero vectors whose entries are nonnegative integers (resp. nonnegative rationals). Let $A^{\mathbb{Z}^+} = \{1\} \cup A^{\mathbb{Z}^+}$.

Corollary 2.7. Let M be a complete metric space, and $K \subset M$ be the attractor of an IFS satisfying the SSC with contraction ratio set X . Then there exist two finite sets $\Lambda, \Gamma \subset (0, \infty)$ such that

$$\text{GL}(K) = \Gamma \bigcup \Lambda X^{\mathbb{Z}^+}. \quad (2.2)$$

2.2. Ratio analysis. In [28], the ‘‘ratio analysis’’ method is used to analyse sets Θ of positive real numbers, in terms of strictly decreasing geometric sequences $\{\theta' r^k\}_{k=0}^\infty$ that are contained in Θ . Whenever we mention ‘geometric sequence’ in this paper, we always assume it is infinitely-long.

Definition 2.8. ([28, Definition 2.4]) Let $\Theta \subset (0, \infty)$. For $\theta \in \Theta$, denote by

$$R_\Theta(\theta) = \{r \in (0, 1) : \text{there exists some } \theta' \in \Theta \text{ such that } \theta \in \{\theta' r^k\}_{k=0}^\infty \subset \Theta\}, \quad (2.3)$$

the set of common ratios of strictly decreasing geometric sequences in Θ that contains θ (which may be empty).

This concept is quite natural since the gap length set of a dust-like attractor contain many geometric sequences. By definition, there is an obvious monotonicity: when $\theta \in \Theta_1 \subset \Theta_2 \subset (0, \infty)$,

$$R_{\Theta_1}(\theta) \subset R_{\Theta_2}(\theta), \quad (2.4)$$

since a geometric sequence in Θ_1 is also in Θ_2 .

We will use the following two lemmas.

Lemma 2.9. ([28, Lemma 2.6 (i)]) Let $A = \{a_i\}_{i=1}^n \subset (0, 1)$ be a finite set, and λ_j ($j = 1, \dots, m$) be positive real numbers (not necessarily distinct). Let $\Theta = \bigcup_{j=1}^m \lambda_j A_j$ where $A_j \subset A^{\mathbb{Z}^+}$ for $1 \leq j \leq m$. Then $R_\Theta(\theta) \subset A^{\mathbb{Q}^+}$ for all $\theta \in \Theta$.

Lemma 2.10. ([28, Corollary 2.7]) Let $X \subset (0, 1)$ and $\Lambda \subset (0, \infty)$ be two finite sets. Then

$$X^{\mathbb{Z}^+} \subset R_{\Lambda X^{\mathbb{Z}^+}}(\theta) \subset X^{\mathbb{Q}^+}$$

for every $\theta \in \Lambda X^{\mathbb{Z}^+}$.

3. OUR RESULTS

3.1. Using ratio analysis to obtain logarithmic commensurability. Logarithmic commensurability is central to the affine-embedding problem [2, 3, 15, 19, 21, 37, 39] and inverse fractal problem (generating IFS), which is stated as follows.

Conjecture 3.1. [19, Conjecture 1.2] Suppose that K, F are two totally disconnected non-singleton self-similar sets in \mathbb{R}^n , which are the attractors of two IFS with contraction ratio sets X, Y respectively. Suppose that there exists an affine map f in \mathbb{R}^n such that $f(F) \subset K$, then the logarithmic commensurability holds: $Y \subset X^{\mathbb{Q}^+}$.

The following theorem can be deduced from [15, Lemma 4.8] in \mathbb{R}^n , which uses geometric measure theoretic arguments. Here we present another proof without using measures.

Theorem 3.2. Let M be a complete metric space. Suppose that $K \subset M$ has two generating IFSs satisfying the SSC with contraction ratio sets A and X , respectively, then $X \subset A^{\mathbb{Q}^+}$, $A \subset X^{\mathbb{Q}^+}$.

This theorem is a direct consequence of Lemma 3.4. To prove Lemma 3.4, we first prove a more general lemma for GD-attractors, which will be used later.

Lemma 3.3. *Let $G = (V, E)$ be a directed graph satisfying $d_w \geq 2$ for all $w \in V$, and let $(M^u)_{u \in V}$ be a list of complete metric spaces. Suppose that for some $u \in V$, $F_u \subset M^u$ is the GD-attractor of some GD-IFS (based on G) satisfying the SSC with contraction ratio set A . Then $R_{\text{GL}(F_u)}(\theta) \subset A^{\mathbb{Q}^+}$ for all $\theta \in \text{GL}(F_u)$.*

Proof. We apply Lemma 2.9 to $\Theta = \text{GL}(F_u)$ given by (2.1), by regarding the numbers in Γ_u , Λ_u and Λ_v ($v \in V$) as λ_j , where the corresponding A_j are $\{1\}$ and R_{uv} respectively. Since A is the contraction ratio set, $r_e \in A^{\mathbb{Z}^+}$, the conditions in Lemma 2.9 are fulfilled, showing the desired. \square

Lemma 3.4. *Let M be a complete metric space and $K \subset M$ be the attractor of an IFS satisfying the SSC with contraction ratio set X . Then for all small enough $\theta \in \text{GL}(K)$,*

$$X \subset X^{\mathbb{Z}^+} \subset R_{\text{GL}(K)}(\theta) \subset X^{\mathbb{Q}^+}. \quad (3.1)$$

Proof. For the left-hand side in (3.1), by Corollary 2.7 and our assumption, there exist two finite sets $\Lambda, \Gamma \subset (0, \infty)$ such that

$$\text{GL}(K) = \Gamma \bigcup \Lambda X^{\mathbb{Z}^+}. \quad (3.2)$$

Thus by the monotonicity (2.4) and Lemma 2.10, we know that for $\theta \in \Lambda X^{\mathbb{Z}^+} \subset \text{GL}(K)$,

$$X^{\mathbb{Z}^+} \subset R_{\Lambda X^{\mathbb{Z}^+}}(\theta) \subset R_{\text{GL}(K)}(\theta),$$

showing the desired for all small enough $\theta \in \text{GL}(K)$ (smaller than the numbers in the finite set Γ in (3.2)).

The right-hand side in (3.1) directly follows from Lemma 3.3. The proof is complete. \square

Proof of Theorem 3.2. By Lemma 3.4, for all small enough $\theta \in \text{GL}(K)$,

$$A \subset R_{\text{GL}(K)}(\theta) \subset A^{\mathbb{Q}^+}, X \subset R_{\text{GL}(K)}(\theta) \subset X^{\mathbb{Q}^+}.$$

Thus

$$A \subset R_{\text{GL}(K)}(\theta) \subset X^{\mathbb{Q}^+}, X \subset R_{\text{GL}(K)}(\theta) \subset A^{\mathbb{Q}^+},$$

which ends the proof. \square

3.2. Intrinsic characterisation of algebraic dependence number. We first generalize the definition of algebraic dependence number to GD-IFSs. Denote by $\text{span} B$ the vector space generated by $B \subset \mathbb{R}$ over \mathbb{Q} , $\log X := \{\log x : x \in X\}$ for a set $X \subset (0, \infty)$.

Definition 3.5 (Algebraic (in)dependence number of GD-IFS). *Given a GD-IFS F with contraction ratio set A , define its algebraic independence number as the dimension of the vector space $\text{span} \log A$, and its algebraic dependence number as its algebraic independence number minus 1.*

Theorem 3.6. *Suppose that F is a GD-IFS satisfying the SSC with attractor list $(F_u \subset M^u)_{u \in V}$ (on complete metric spaces), based on a directed graph (V, E) with $d_w \geq 2$ for all $w \in V$, then its algebraic independence number is no less than the dimension of*

$$\text{span} \left\{ \log r : r \in \bigcup_{u \in V} \bigcup_{\theta \in \text{GL}(F_u)} R_{\text{GL}(F_u)}(\theta) \right\}. \quad (3.3)$$

In particular, for a dust-like self-similar set K (on a complete metric space), its algebraic dependence number plus 1 equals the dimension of

$$\text{span} \left\{ \log r : r \in \bigcup_{\theta \in \text{GL}(K)} R_{\text{GL}(K)}(\theta) \right\} = \text{span} \{ \log r : r \in R_{\text{GL}(K)}(\theta) \} \quad (3.4)$$

for any small enough $\theta \in \text{GL}(K)$ (the vector space generated by the logarithms of all the common ratios of infinite geometric sequences in $\text{GL}(K)$ over \mathbb{Q}).

Remark 3.7. *In this theorem, “infinite geometric sequences in $\text{GL}(K)$ ” has the same effect as “strictly decreasing geometric sequences in $\text{GL}(K)$ ”, since the set $\text{GL}(K)$ has an upper bound $\text{diam}(K)$, which means that the infinite geometric sequences in $\text{GL}(K)$ must have common ratio no greater than 1 (but $\log 1 = 0$ is useless to be a basis for vector spaces).*

In fact we prove a stronger property (3.5) for dust-like self-similar sets: the vector space generated by the logarithms of the contraction ratios of any SSC generating IFS (over \mathbb{Q}) is equal to the vector space generated by the logarithms of all the common ratios of geometric sequences in the gap length set (over \mathbb{Q}).

Proof. For the first inclusion, by using Lemma 3.3 and taking the logarithm,

$$B := \left\{ \log r : r \in \bigcup_{u \in V} \bigcup_{\theta \in \text{GL}(F_u)} R_{\text{GL}(F_u)}(\theta) \right\} \subset \log A^{\mathbb{Q}_+^*},$$

where A is the contraction ratio set of F . It follows by Definition 2.6 that

$$\text{span} B \subset \text{span} \log A^{\mathbb{Q}_+^*} = \text{span} \log A.$$

Then we know the dimension of $\text{span} B$ is no greater than that of $\text{span} \log A$, which is exactly the algebraic independence number of F .

For the second inclusion, by using Lemma 3.4 and taking the logarithm,

$$\log X \subset B' \subset \log X^{\mathbb{Q}_+^*},$$

where X is the contraction ratio set of any SSC generating IFS of K , and B' can be either $\{\log r : r \in \bigcup_{\theta \in \text{GL}(K)} R_{\text{GL}(K)}(\theta)\}$ or $\{\log r : r \in R_{\text{GL}(K)}(\theta)\}$ for any small enough $\theta \in \text{GL}(K)$. Thus

$$\text{span} \log X \subset \text{span} B' \subset \text{span} \log X^{\mathbb{Q}_+^*}.$$

Note that by Definition 2.6, $\text{span} \log X = \text{span} \log X^{\mathbb{Q}_+^*}$, thus

$$\text{span} \log X = \text{span} B' = \text{span} \log X^{\mathbb{Q}_+^*}, \quad (3.5)$$

showing (3.4). Since the algebraic dependence number of K is the algebraic dependence number of any SSC generating IFS, that is, the dimension of $\text{span} \log X$, the proof is finished. \square

3.3. Lower bound on the cardinality of generating IFS. Algebraic independence number provides a natural lower bound for the cardinality of generating IFSs, so we are able to present such lower bound in terms of the fractal itself (using gap lengths).

Corollary 3.8. *Let $G = (V, E)$ be a directed graph satisfying $d_w \geq 2$ for all $w \in V$, and let $(M^u)_{u \in V}$ be a list of complete metric spaces. Suppose that for some $u \in V$, $F_u \subset M^u$ is the GD-attractor of some GD-IFS F satisfying the SSC (based on G). Then the number of edges in G is no less than the dimension of $\text{span}\{\log r : r \in \bigcup_{\theta \in \text{GL}(F_u)} R_{\text{GL}(F_u)}(\theta)\}$.*

In particular, for a dust-like self-similar set $K \subset M$ (where M is a complete metric space), the cardinality of each generating IFS satisfying the SSC is no less than dimension of $\text{span}\{\log r : r \in \bigcup_{\theta \in \text{GL}(K)} R_{\text{GL}(K)}(\theta)\} = \text{span}\{\log r : r \in R_{\text{GL}(K)}(\theta)\}$ for any small enough $\theta \in \text{GL}(K)$.

Proof. For the first claim, just note that $\#E$, the number of edges in G , which is also the cardinality of contracting similarities, is no less than the cardinality of the contraction ratio set of F , and thus no less than the algebraic independence number of F . The first claim then follows by Theorem 3.6 and that $\text{span}\{\log r : r \in \bigcup_{\theta \in \text{GL}(F_u)} R_{\text{GL}(F_u)}(\theta)\} \subset \text{span}\{\log r : r \in \bigcup_{u \in V} \bigcup_{\theta \in \text{GL}(F_u)} R_{\text{GL}(F_u)}(\theta)\}$.

The second claim immediately follows from the first claim and (3.4). \square

Remark 3.9. We delete the union “ $\bigcup_{u \in V}$ ” in (3.3), since we only have the information of one attractor at one vertex u . In addition, we only give a lower bound on the number of edges, not of vertices.

Furthermore, if the dust-like self-similar set is on \mathbb{R} and is full-measure, we can remove the separation condition (SSC) on the generating IFS. We say that a nonempty compact set K is *full-measure* if $\mathcal{H}^{\dim_{\mathbb{H}} K}(K) = (\text{diam}(K))^{\dim_{\mathbb{H}} K}$. ($\dim_{\mathbb{H}}$ is the Hausdorff dimension)

Corollary 3.10. Suppose that a full-measure set $K \subset \mathbb{R}$ is the attractor of an IFS Φ satisfying the SSC, then the cardinality of each generating IFS of K is no less than dimension of $\text{span}\{\log r : r \in \bigcup_{\theta \in \text{GL}(K)} R_{\text{GL}(K)}(\theta)\}$. If further, the logarithm of the contraction ratio of each contracting similarity in Φ is different from each other and linearly independent over \mathbb{Q} , then Φ has the minimal cardinality among all generating IFSs of K .

Proof. For any generating IFS $\{S_i\}_{i=1}^m$ of K , there exists $I \in \{1, 2, \dots, m\}$ such that, the sub-IFS $\Psi := \{S_i\}_{i \in I}$ also has attractor K and satisfies the SSC. Indeed, whenever $S_i(K) \cap S_j(K) \neq \emptyset$, by [20, The claim in the proof of Theorem 4.1], $S_i(K) \subset S_j(K)$ (or $S_i(K) \supset S_j(K)$), so one can remove i (or j) from $\{1, 2, \dots, m\}$, and repeat this process until the SSC is satisfied. The first inclusion then follows by using Corollary 3.8 for the lower bound of $\#\Psi$ (also a lower bound for m).

For the second inclusion, by assumption, the algebraic independence number of Φ is exactly its cardinality $\#\Phi$, thus the algebraic independence number of Ψ is also $\#\Phi$ by [15, Theorem 5.7]. Since the cardinality of Ψ is no less than the algebraic independence number of Ψ , which is $\#\Phi$, we obtain the desired by noting $m \geq \#\Psi$. \square

The full-measure condition is easy to verify by the similarities of an IFS on \mathbb{R} , see [20, Remark 4.1] for a practical sufficient and necessary condition and examples. We end our paper by a brief overview on the full-measure condition. It is firstly discussed in Hausdorff’s paper [25], then extended by Marion (called ‘perfect isotopic’) [30, 31], and independently by Ayer and Strichartz on self-similar sets (several years later): they give the full-measure criterion for dust-like self-similar sets on \mathbb{R} . There is no known analogous criterion in higher dimensional case to the best of our knowledge. The study of full-measure condition for a GD-attractor is more complicated than that for self-similar set (see [5, Theorem 4.6]), and also, there is no known analogous criterion even on \mathbb{R} .

ACKNOWLEDGEMENT. The author was supported by National Natural Science Foundation of China (12271282).

REFERENCES

- [1] A. ALGOM, *Affine embeddings of Cantor sets on the line*, J. Fractal Geom., 5 (2018), pp. 339–350.
- [2] ———, *Affine embeddings of Cantor sets in the plane*, J. Anal. Math., 140 (2020), pp. 695–757.
- [3] A. ALGOM AND M. HOCHMAN, *Self-embeddings of Bedford-McMullen carpets*, Ergodic Theory Dynam. Systems, 39 (2019), pp. 577–603.
- [4] M. BODIN, *Wavelets and Besov spaces on Mauldin-Williams fractals*, Real Anal. Exchange, 32 (2006/07), pp. 119–143.
- [5] G. C. BOORE AND K. J. FALCONER, *Attractors of directed graph IFSs that are not standard IFS attractors and their Hausdorff measure*, Math. Proc. Cambridge Philos. Soc., 154 (2013), pp. 325–349.
- [6] S. CAO, H. QIU, H. TIAN, AND L. YANG, *Spectral decimation for a graph-directed fractal pair*, Sci. China Math., 65 (2022), p. 2503–2520.
- [7] E. CHARLIER, J. LEROY, AND M. RIGO, *An analogue of Cobham’s theorem for graph directed iterated function systems*, Adv. Math., 280 (2015), pp. 86–120.
- [8] M. DAS AND G. A. EDGAR, *Separation properties for graph-directed self-similar fractals*, Topology Appl., 152 (2005), pp. 138–156.
- [9] M. DAS AND S.-M. NGAI, *Graph-directed iterated function systems with overlaps*, Indiana Univ. Math. J., 53 (2004), pp. 109–134.

- [10] J. DENG AND L. XI, *Gap sequences of graph-directed sets*, *Fractals*, 24 (2016), pp. 1650036, 10.
- [11] Q.-R. DENG AND K.-S. LAU, *On the equivalence of homogeneous iterated function systems*, *Nonlinearity*, 26 (2013), pp. 2767–2775.
- [12] ———, *Structure of the class of iterated function systems that generate the same self-similar set*, *J. Fractal Geom.*, 4 (2017), pp. 43–71.
- [13] G. EDGAR, *Measure, topology, and fractal geometry*, Undergraduate Texts in Mathematics, Springer, New York, second ed., 2008.
- [14] G. A. EDGAR AND R. D. MAULDIN, *Multifractal decompositions of digraph recursive fractals*, *Proc. London Math. Soc.* (3), 65 (1992), pp. 604–628.
- [15] M. ELEKES, T. KELETI, AND A. MÁTHÉ, *Self-similar and self-affine sets: measure of the intersection of two copies*, *Ergodic Theory Dynam. Systems*, 30 (2010), pp. 399–440.
- [16] K. FALCONER, *Techniques in fractal geometry*, John Wiley & Sons, Ltd., Chichester, 1997.
- [17] A. FARKAS, *Projections of self-similar sets with no separation condition*, *Israel J. Math.*, 214 (2016), pp. 67–107.
- [18] ———, *Dimension approximation of attractors of graph directed IFSs by self-similar sets*, *Math. Proc. Cambridge Philos. Soc.*, 167 (2019), pp. 193–207.
- [19] D.-J. FENG, W. HUANG, AND H. RAO, *Affine embeddings and intersections of Cantor sets*, *J. Math. Pures Appl.* (9), 102 (2014), pp. 1062–1079.
- [20] D.-J. FENG AND Y. WANG, *On the structures of generating iterated function systems of Cantor sets*, *Adv. Math.*, 222 (2009), pp. 1964–1981.
- [21] D.-J. FENG AND Y. XIONG, *Affine embeddings of Cantor sets and dimension of $\alpha\beta$ -sets*, *Israel J. Math.*, 226 (2018), pp. 805–826.
- [22] A. E. GHENCUI, R. D. MAULDIN, AND M. ROY, *Conformal graph directed Markov systems: beyond finite irreducibility*, *J. Fractal Geom.*, 3 (2016), pp. 217–243.
- [23] Q. GU AND K.-S. LAU, *Dirichlet forms and critical exponents on fractals*, *Trans. Amer. Math. Soc.*, 373 (2020), pp. 1619–1652.
- [24] B. M. HAMBLY AND S. O. G. NYBERG, *Finitely ramified graph-directed fractals, spectral asymptotics and the multidimensional renewal theorem*, *Proc. Edinb. Math. Soc.* (2), 46 (2003), pp. 1–34.
- [25] F. HAUSDORFF, *Dimension und äußeres Maß*, *Math. Ann.*, 79 (1918), pp. 157–179.
- [26] X.-G. HE, K.-S. LAU, AND H. RAO, *Self-affine sets and graph-directed systems*, *Constr. Approx.*, 19 (2003), pp. 373–397.
- [27] J. E. HUTCHINSON, *Fractals and self-similarity*, *Indiana Univ. Math. J.*, 30 (1981), pp. 713–747.
- [28] JIAXIN HU, KENNETH J. FALCONER AND JUNDA ZHANG, *A dichotomy on the self-similarity of graph-directed attractors*, *J. Fractal Geom.*, (2024).
- [29] W. LI AND Y. YAO, *Generating iterated function systems for the vicsek snowflake and the koch curve*, *The American Mathematical Monthly*, (2017).
- [30] J. MARION, *Mesures de Hausdorff et théorie de Perron-Frobenius des matrices non-negatives*, *Ann. Inst. Fourier (Grenoble)*, 35 (1985), pp. 99–125.
- [31] ———, *Mesure de Hausdorff d'un fractal à similitude interne*, *Ann. Sci. Math. Québec*, 10 (1986), pp. 51–84.
- [32] R. D. MAULDIN AND S. C. WILLIAMS, *Hausdorff dimension in graph directed constructions*, *Trans. Amer. Math. Soc.*, 309 (1988), pp. 811–829.
- [33] V. METZ, *“Laplacians” on finitely ramified, graph directed fractals*, *Math. Ann.*, 330 (2004), pp. 809–828.
- [34] T.-J. NI AND Z.-Y. WEN, *Open set condition for graph directed self-similar structure*, *Math. Z.*, 276 (2014), pp. 243–260.
- [35] H. RAO, H.-J. RUAN, AND Y.-M. YANG, *Gap sequence, Lipschitz equivalence and box dimension of fractal sets*, *Nonlinearity*, 21 (2008), pp. 1339–1347.
- [36] H. RAO, Z.-Y. WEN, AND Y.-M. YANG, *Dual systems of algebraic iterated function systems*, *Adv. Math.*, 253 (2014), pp. 63–85.
- [37] P. SHMERKIN, *On Furstenberg’s intersection conjecture, self-similar measures, and the L^q norms of convolutions*, *Ann. of Math.* (2), 189 (2019), pp. 319–391.
- [38] S. TROSCHET, *On the dimensions of attractors of random self-similar graph directed iterated function systems*, *J. Fractal Geom.*, 4 (2017), pp. 257–303.
- [39] M. WU, *A proof of Furstenberg’s conjecture on the intersections of $\times p$ - and $\times q$ -invariant sets*, *Ann. of Math.* (2), 189 (2019), pp. 707–751.