κ -General-Relativity I: a Non-Commutative GR Theory with the κ -Minkowski Spacetime as its Flat Limit

Daniel Rozental¹ and Ofek Birnholtz¹

¹Department of Physics, Bar Ilan University, Ramat Gan 5290002, Israel

We employ a twist deformation on infinitesimal diffeomorphisms to study a modification of General Relativity on a non-commutative spacetime extending the local κ -Minkowski spacetime. This non-commutative spacetime is present in Deformed Special Relativity (DSR) theories, where a fundamental length is relativistically incorporated into Special Relativity as an effective description of Quantum Gravity near Planckian energy scales. To avoid mathematical and physical ambiguities, such as the inability to twist-deform the Poincaré symmetry group such that κ -Minkowski becomes *its* covariant spacetime, we propose a novel approach to DSR and non-commutative theories, twisting the Dilatation-enlarged IGL(3,1) group and constructing deformed General Relativity accordingly. Physically, we interpret this as minimally modifying classical relativistic physics to accommodate quantum effects solely through spacetime uncertainties in a relativistic manner, a possibility new to DSR theories. These tools open the way to developing the dynamic sector of DSR, which would allow for predictions testable by astrophysical observations.

Keywords: Deformed Special Relativity, General Relativity, Quantum Gravity, Non-Commutative Spacetime, κ-Poincaré Group, Twist-deformations.

I. INTRODUCTION

A. Background and Motivation

One of the phenomenological approaches to formulating a quantum gravity (QG) theory involves introducing deformations to the Poincaré symmetries underlying Special Relativity (SR). This deformation is aimed at ensuring that the resulting theory maintains an additional invariant length scale, which is considered fundamental in QG theories [1–5]. These theories, often referred to as Deformed Special Relativity (DSR) or Deformed Relativistic Kinematics (DRK), represent a phenomenological ("bottom-up") approach to addressing the QG problem [1].

Phenomenological theories aim to capture essential properties of various QG theories and incorporate them as deformations into the classical theory. This enables the prediction of deviations from classical gravity in astrophysical systems, which could be probed using technologies within reach [6–8]. In contrast, first-principle theories ("top-down"), such as String Theory and Loop Quantum Gravity, seek to address the problem from a more fundamental level but often yield physical predictions that are difficult or impossible to calculate and are typically applicable only to energies far beyond current technology [1, 9].

Fundamentally, DSR theories involve deforming the symmetries of SR in a manner that preserves the invariance of a QG-length scale, often identified as the Planck length ℓ_p [4, 10–12]:

$$\ell_p = \sqrt{\frac{\hbar G}{c^3}} \sim 10^{-35} \,\mathrm{m} \ . \tag{1}$$

We also define the Planck energy $\kappa = E_p = \hbar c / \ell_p$, so that in Quantum Field Theory (QFT) units ($\hbar = c = 1$),

 $\kappa = \ell_p^{-1}$. It represents the energy scale at which General Relativity (GR), when treated as a QFT, is expected to diverge [13].

Various methods exist for implementing these deformations, each leading to different phenomenological consequences. For example, in the seminal papers [2, 3], one finds two different approaches relying on slightly different basic arguments (whether the starting point is invariant Planck length or invariant Planck energy), resulting in different deformations of how Lorentz boosts act on momenta/energy. These differences affect [14, 15] whether one expects a momenta-dependent speed of light, saturation of energy and momenta, etc. (see also [16, 17]).

Moreover, ensuring length scale invariance in causally connected states requires a deformed composition of momenta. And, from relativistic compatibility, a deformed action of transformations on multi-particle states is needed, an effect also known as backreaction on rapidity for the case of boost transformations [18, 19]. These different possibilities can be understood as different bases of quantum deformations of the Poincaré group [20], establishing connections [7, 8] with related theories such as curved momentum space and Relative Locality [6, 21, 22].

Regardless of the specific model, DSR theories share two key properties. First is the promotion of spacetime variables into *operators* with a non-vanishing commutation relation, known as κ -Minkowski spacetime [23–27]:

$$[\hat{x}^0, \hat{x}^i] = i\ell_p \hat{x}^i, \quad [\hat{x}^i, \hat{x}^j] = 0, \quad \text{for} \quad i, j = 1, 2, 3.$$
 (2)

Second, there are deformations in the Hopf algebra sector of the Poincaré symmetry (ISO(3,1)) [25, 26, 28–31], ensuring the relativistic invariance of relation (2) and the deformed symmetries.

Given a system's symmetry group G, a Hopf algebra is a structure embedded in the Universal Enveloping Algebra $\mathbf{U}(G)$ (the freely generated tensor algebra with the same representations/modules as G's Lie algebra). It determines how observables (symmetry generators) act on multi-particle states represented as the tensor product of irreducible representations. Classically, this structure is trivial and often uncharacterized. For example, the momentum generator (translation) with the one-particle (irreducible) representation D(P) acts on one-particle and two-particle states as follows,

$$D(\mathbf{P})|K\rangle = \mathbf{k}|K\rangle \quad , \quad D(\mathbf{P})|L\rangle = \mathbf{l}|L\rangle \, , \tag{3}$$

$$|KL\rangle \rightarrow (D \otimes D)(\Delta \mathbf{P})(|K\rangle \otimes |L\rangle) = (\mathbf{k} + \mathbf{l}) |KL\rangle, (4)$$

with the trivial co-product in the Hopf algebra defined as

$$\Delta \mathbf{P} \equiv \mathbf{P} \otimes 1 + 1 \otimes \mathbf{P}. \tag{5}$$

In a DSR theory, the form of (5) is generally modified, along with the remaining Hopf algebra structure on U(ISO(3,1)). A symmetry group G with such nontrivial Hopf algebra is called a Quantum Group. Any DSR model can be associated with a specific Quantum Group structure [32] and the spacetime (2) as its invariant space.

Therefore, investigating these Quantum Group and non-commutative space deformations in theories with a dynamical sector, such as the theory of gravity, will impose constraints on any DSR model.

Dynamical theories can often be deformed in their Hopf-algebra sector using deformation quantization tools [33–35], where definitive deformation operations on the classical structures derive the non-commutativity and deformed Hopf algebras. Twisted algebras [36–38] or Seiberg-Witten (SW) maps [39] are such operations schemes, with the former being applied to the Lie algebra of the gauge theory's symmetry. In contrast, the latter directly applies to the gauge variables.

Regarding the non-commutative deformation of GR, foundations of its twist-deformed version were outlined in [40, 41], with consistency proofs provided in [42]. The general view is to formulate an invariant theory under deformed infinitesimal diffeomorphisms (viewed as the Lie algebra of vector fields deformed in its Hopf sector). One should view the non-commutative GR as describing the gravitational interaction on a noncommutative manifold through non-commutative differential geometry. The non-commutative manifold is locally a non-commutative spacetime characterized by the type of deformation describing the *flat* sector of the noncommutative spacetime. In DSR theories, one considered this flat limit to be the κ -Minkowski spacetime, described in (2).

However, definitive results were only obtained for constant non-commutativity [43, 44], a result of a deformed symmetry group that lacks apparent physical meaning [45, 46]. In contrast, no construction has been made explicitly for the physically attractive noncommutative scenario as defined by (2) (only special cases were treated, such as assuming specific symmetries in spacetime, see [42, 47] and references therein). This deficiency can be traced to a profound ambiguity: twisting classical algebras results only in deformations in the Hopf-algebra sector of the symmetry group, leaving the algebra sector classical. However, quantum deformations applied to the Poincaré algebra (κ -Poincaré) that will result in the κ -Minkowski spacetime are only consistent with simultaneous deformation of the algebra and the Hopf-algebra. An illustration of this consistency condition is the attempt to formulate such a quantum deformation of the Poincaré group that deforms only the Hopf sector, known as κ -Poincaré [31]. This formulation suffers from h-acid Topology (one cannot assign the deformation parameter a numerical value), rendering the interpretation of the physical deformation ambiguous. Therefore, one cannot derive κ -Poincaré via a Twist (it was also proven in [48]), an obstacle for the construction of GR (or any other gauge theory) on κ -Minkowski with the minimal-length interpretation.

This is a problem because any reasonable attempt to assign a quantum nature to spacetime should admit a gravitational theory, which should coincide with the GR predictions at scales near the quantum-classical intersection. Thus, for the DSR-type theories, we expect that, near Planckian scales, one should be able to construct a theory of gravity on the non-commutative space of DSR.

B. Overview

To achieve this, we follow a resolution similar to that found in [45, 49]: enlarging the underlying symmetry from the Poincaré group to the IGL(3,1) group (the Inhomogeneous General Linear group, which can be considered as a Dilatation-enlarged Poincaré group) and twisting it in such a way that its covariant spacetime is κ -Minkowski. This way, we can use the known Twist formalism and construct a self-consistent GR on a non-commutative spacetime locally described by the κ -Minkowsky, calling it κ -GR.

However, two pressing questions remain. The first comes from assessing κ -GR on its own right: since twist deformations only alter the Hopf sector and leave the algebra untouched, what is the meaning of the κ -Minkowski spacetime here? Is it relativistically invariant, and does it retain a notion of an invariant length as in other DSR models? The second regards the correspondence principle: since the theory is formulated with deformed IGL(3,1) symmetry, its classical limit will be GR with local symmetry of IGL(3,1), not Poincaré, raising concerns about its physical realism.

We will briefly discuss possible solutions in the discussion section, leaving a more careful analysis to a future paper. Our resolution might be interesting not only for the construction of GR but also for exploring whether spacetime can be related to a quantum nature without imposing modifications in the Poincaré transformations (algebraically), which would expand the scope of DSR theories.

The rest of this paper is organized as follows: In Section Sec.II, we outline the foundations of geometry and the construction of GR elements in a general noncommutative spacetime as it appears in [41]. In Section Sec.III, we re-derive the theory of gravity with a local constant non-commutative spacetime. We utilize the fundamental tools outlined in Section Sec.II to present a proof of concept and compare our results with those found in previous works such as [43, 44]. This effort is essential because the existing derivation has been performed in a somewhat different, simplified manner that only applies to constant non-commutativity. In Sec.IV, we outline the foundational formulas for the construction of GR with a local κ -Minkowski spacetime and IGL(3,1) symmetry and present additional ambiguities that will follow if one attempts to construct it on the Poincaré group. The results are partly based on [45], and some are developed to build the deformed GR. In section Sec.V, we build the theory of twisted GR with a local κ -Minkowski spacetime by following a route similar to that in Sec.III. In section Sec.VI, we discuss the meaning of the derived results, the physical picture of a non-commutative spacetime without algebra deformations, and the future work that aims to solve the ambiguities that arise.

II. BASIC FORMULATION: NON-COMMUTATIVE GEOMETRY AND GRAVITY

A. Syntax and notation conventions

This section gradually outlines the formulae and the mathematical constructions necessary for deformed general relativity extending the κ -Minkowski spacetime. In Sec.IIB, we describe the general formalism of non-commutative spaces by twisting the Lie algebra of vector fields. In particular, it includes the full derivations and definitions of the following operators and operands, enumerated here in brief:

- 1. *-product, a non-commutative product used as an isomorphism between non-commutative spacetime variables and their classical counterparts.
- 2. Φ , the twist operator that generates the *product; it is an invertible rank two tensor composed of vector fields belonging to the symmetry we want to deform. Its inverse is $\overline{\Phi}$.
- 3. \mathcal{R} , an invertible rank two tensor; the universal \mathcal{R} -matrix (as usually defined in Hopf algebras) which, for the current discussion, is used to flip the order of \star -multiplications, demonstrating the non-commutativity. Its inverse is $\overline{\mathcal{R}}$.

- 4. μ , a bidifferential operator, acting as a linear map on products of vector fields (X, Y). We denote its action on some expression of (X, Y) with a " \circ ".
- 5. L^{*}, ⟨·;·⟩, [·;·], ⊗..., the deformed versions of the classical operations (in this case: Lie derivative, the pairing map, vector fields' commutator, and the tensor product) required in a description of differential geometry. Generally, every operator/operand denoted with a *-subscript is the deformed version consistent with non-commutative spacetime variables.

In Sec.II C, we describe the required formalism for a GR theory on non-commutative space, utilizing the formalism established in Sec.II B.

Concerning conventions: we use late Greek alphabet letters for spacetime indices and mid-Latin letters for purely spatial indices (e.g., $\mu, \nu = 0, i$, with i, j = 1, 2, 3); early Latin letters (a, b, c) enumerate generators of algebras; early greek letters (α) enumerate orders of expansion in a parameter λ ; the Gothic \mathfrak{X} is used for geometrical quantities, such as the Riemann tensor, the Ricci tensor, and the Ricci scalar.

B. Non-commutative Spaces & Twisting

Here, we outline how one can deform a commutative space to obtain a non-commutative one in a manner that allows for defining similar concepts as in the classical (commutative) scenario. Definitions such as differentiation, multiplication, and invariance of fields on that space are intrinsic properties of the underlying symmetry; if the symmetry is deformed, the actions on the fields will also be deformed. Understanding the behavior of fields is crucial for constructing any physical theory on non-commutative spaces. Thus, we aim to outline a coherent method for achieving this.

Technically, we first describe how to represent a commutative space with vector fields associated with a Lie algebra, which can be trivially extended to a Hopf algebra. By introducing the twist element Φ , we establish an isomorphism between fields on the non-commutative space and the commutative one via the *-product, defining a deformed algebra of vector fields. From this point, by establishing all other representations of the algebra, we will explore how tensor fields, one-forms, basis vectors, and so on behave in the non-commutative space. Along the way, we introduce other valuable operations, such as the *-tensor-product $\stackrel{*}{\otimes}$ and *-pairing < \cdot, \cdot, \cdot , which lead to new definitions for basis vectors and 1forms on the non-commutative space.

We start (see [25]) by considering the algebra *freely* generated by the non-commutative variables \hat{x}^{μ} , i.e., consider the relation,

$$[\hat{x}^{\mu}, \hat{x}^{\nu}] = C^{\mu\nu}(\hat{x}), \tag{6}$$

where $C^{\mu\nu}(\hat{x})$ is an expansion in \hat{x}^{μ} with constant coefficients,

$$C^{\mu\nu}(\hat{x}) = i\theta^{\mu\nu} + ic^{\mu\nu}_{\rho}\hat{x}^{\rho} + \mathcal{O}(\hat{x}^{\alpha}\hat{x}^{\beta}).$$
(7)

The relation (7) generates the ideal of the algebra, and by considering the formal power series in \hat{x}^{μ} , we obtain the quotient by dividing with the ideal [40]:

$$\hat{\mathcal{A}}_{\hat{x}} = \langle \langle \hat{x}^0 \dots \hat{x}^n \rangle \rangle / ([\hat{x}^\mu, \hat{x}^\nu] - C^{\mu\nu}(\hat{x})), \qquad (8)$$

this algebra represents non-commutative functions, with commuting functions identified as elements in \mathcal{A}_x . Focusing on structures that obey the Poincaré-Birkhoff-Witt property [50], the space of non-commutative polynomials (8) is isomorphic to polynomials in \mathcal{A}_x equipped with a *-product [25, 39] (a *-isomorphism),

$$\hat{\mathcal{A}}_{\hat{x}} \sim (\mathcal{A}_x, \star), \tag{9}$$

implying that polynomials $\hat{P}(\hat{x}) \in \hat{A}_{\hat{x}}$ and $P(x) \in A_x$ are related through the mapping:

$$\hat{P}_1(\hat{x})\hat{P}_2(\hat{x}) \mapsto P_1(x) \star P_2(x).$$
 (10)

The specific form of this mapping arises from deformation in the Hopf algebra sector of the symmetry group on the space under consideration \mathcal{M} (a smooth manifold in general), where spaces with the structures described in equation (7) become the deformed symmetry module algebras.

This Hopf deformation is achieved by first considering the algebra of vector fields Ξ (infinite dimensional, which is the algebra of infinitesimal diffeomorphisms) on \mathcal{M} . This structure possesses a Lie algebra structure, with elements generated by $\{v^1, .., v^N\}$ (N is the group order), and has a Hopf algebra extension on its universal envelope $\mathbf{U}(\Xi)$ (the algebra of tensor fields modulo the ideal of the Lie algebra Ξ). Denoting with Latin indices the generator number, the Lie algebra and the Hopf algebra are,

$$\begin{bmatrix} v^a, v^b \end{bmatrix} = i f^{abc} v^c,$$

$$\Delta(v^a) = v^a \otimes 1 + 1 \otimes v^a \equiv \sum_{i=1}^2 v^a_{(1)i} \otimes v^a_{(2)i}$$

$$(11)$$

$$\equiv v_{(1)}^a \otimes v_{(2)}^a, \tag{12}$$

where we used the Sweedler notation for $v_{(1)}^a, v_{(2)}^a$, with an understood summation, e.g., for (12) we have $v_{(1)2}^a = v_{(2)1}^a = 1$, $v_{(1)1}^a = v_{(2)2}^a = v^a$. A possible deformation of this Hopf structure is through a **Twist Deformation** [36]: a general twist, denoted as the operator element $\Phi \in \mathbf{U}(\Xi) \otimes \mathbf{U}(\Xi)$, is an invertible operator with a deformation nature in its expansion parameter λ , with the order being denoted by the positive integer α ({ $\alpha \in \mathbb{Z} | \alpha \geq 0$ });

$$\Phi = \phi^{\{\alpha\}} \otimes \phi_{\{\alpha\}} = 1 \otimes 1 + \phi^{\{1\}} \otimes \phi_{\{1\}} + \mathcal{O}(\lambda^2), (13)$$

$$\Phi^{-1} \equiv \bar{\Phi} = \bar{\phi}^{\{\alpha\}} \otimes \bar{\phi}_{\{\alpha\}}, \Leftrightarrow \Phi \bar{\Phi} = 1.$$
(14)

When the twist acts on (12), it gives rise to the *twisted* symmetry structure on the co-algebra, leaving the Lie algebra (11) classical,

$$\Delta^{\Phi}(v^a) \coloneqq \Phi \Delta(v^a) \Phi^{-1} = v^a_{(1)\Phi} \otimes v^a_{(2)\Phi} .$$
 (15)

As in (12), we have used the Sweedler notation in $v_{(i)\Phi}^a$, with subscript Φ denoting that this is the twisted quantity. Resulting in the deformed Hopf algebra denoted as $(\mathbf{U}(\Xi)^{\Phi}, \cdot, \Delta^{\Phi})$, which shares the " \cdot "-product when $(\xi, \eta) \in \mathbf{U}(\Xi)$ act on some quantity: $\mathcal{L}_{\xi}\mathcal{L}_{\eta} = \mathcal{L}_{\xi\eta}$, but differs from the classical one in its co-product (15) and therefore in its representations. Thus, a deformation appears in the Leibniz rule of symmetry transformations when they act on field products. This fact allows the derivation of a quantum Hopf algebra from classical structure; the underlying symmetry of the noncommutative space (9) can be derived through a twist. Operationally, consider a set of vector fields $(X, Y, Z) \in$ Ξ with a definitive action of $\mathbf{U}(\Xi)$ on them. If there exists a linear map,

$$\mu: X \times Y \mapsto Z,\tag{16}$$

it can be meaningfully deformed by the twist¹

$$\mu_{\star} \coloneqq \mu \circ \Phi^{-1} \equiv \mu \circ (\bar{\phi}^{\{\alpha\}} \otimes \bar{\phi}_{\{\alpha\}}). \tag{17}$$

Consider, for example, the algebra of commutative functions $Fun(\mathcal{M}) \equiv \mathcal{A}$. The deformation (17) is welldefined since every $(\xi, \eta) \in \mathbf{U}(\Xi)$ has a defined action on $f \in \mathcal{A}$ via an extension of the Lie derivative of vector fields [41],

$$\xi(f) = \mathcal{L}_{\xi}(f)$$
, where $\mathcal{L}_{\xi\eta} = \mathcal{L}_{\xi}\mathcal{L}_{\eta}$. (18)

The deformation of the usual product (16) of $(f,g) \in \mathcal{A}$ will then be

$$fg \mapsto \mu(\Phi^{-1}(f \otimes g))$$

$$\equiv \mu_{\star}(f \otimes g) = f \star g = \bar{\phi}^{\{\alpha\}}(f)\bar{\phi}_{\{\alpha\}}(g).$$
(19)

This deformed multiplication is associative and welldefined for the twists we are considering. The twist deformation consistently defines a non-commutative algebra of functions $Fun_{\star}(\mathcal{M}) \equiv \mathcal{A}_{\star}$. To observe the noncommutativity of this \star -product, one uses the *universal* \mathcal{R} -matrix [41], defined as,

$$\mathcal{R} := \Phi_{21} \Phi^{-1} = \mathbf{R}^{\{\alpha\}} \otimes \mathbf{R}_{\{\alpha\}} = \phi_{\{\alpha\}} \bar{\phi}^{\{\beta\}} \otimes \phi^{\{\alpha\}} \bar{\phi}_{\{\beta\}},$$
$$\bar{\mathcal{R}} := \mathcal{R}^{-1} = \Phi \Phi_{21}^{-1} = \bar{\mathbf{R}}^{\{\alpha\}} \otimes \bar{\mathbf{R}}_{\{\alpha\}},$$
(20)

¹ The validity of this multiplication comes from certain properties the twist must satisfy; for our purposes, this will always hold, see [41].

with the notation,

$$\Phi_{21} \equiv \tau(\Phi) = \phi_{\{\alpha\}} \otimes \phi^{\{\alpha\}}. \tag{21}$$

where we used the flip operator $\tau : U\Xi \otimes U\Xi \mapsto U\Xi \otimes U\Xi$,

$$\tau(\xi \otimes \eta) = \eta \otimes \xi. \tag{22}$$

Using this \mathcal{R} -matrix, we can express the non-cocommutativity of (15),

$$\tau(\Delta^{\Phi}(\xi)) \coloneqq \mathcal{R}\Delta^{\Phi}(\xi)\mathcal{R}^{-1}.$$
 (23)

And the non-commutativity of the product (19),

$$f \star g = \bar{\mathbf{R}}^{\{\alpha\}}(g) \star \bar{\mathbf{R}}_{\{\alpha\}}(f) \neq g \star f.$$
 (24)

The deformed algebra \mathcal{A}_{\star} with the product (19) is a good representation of the deformed Hopf algebra $\mathbf{U}(\Xi)^{\Phi}$. In other words, \mathcal{A}_{\star} is an algebra module of the deformed Hopf algebra; the algebra structure of \mathcal{A}_{\star} is compatible with the action of $\mathbf{U}(\Xi)^{\Phi}$,

$$\xi(a \star b) = \xi_{(1)\Phi}(a) \star \xi_{(2)\Phi}(b), \qquad (25)$$

where we have used Sweedler's notation for the Leibniz rule of ξ , similarly to (12) and (15), defined here by

$$\xi_{(1)\Phi} \otimes \xi_{(2)\Phi} \equiv \Delta^{\Phi}(\xi), \tag{26}$$

for all $\xi \in \mathbf{U}(\Xi) \subset \mathbf{U}(\Xi)^{\Phi}$. To verify (25) explicitly, one computes it as follows,

$$\begin{aligned} \xi(a \star b) &= \xi(\mu \circ \Phi^{-1}(a \otimes b)) = \mu \circ \Phi^{-1} \circ \Delta^{\Phi}(\xi)(a \otimes b) \\ &= \mu \circ \{\Phi^{-1}(\xi_{(1)\Phi}(a) \otimes \xi_{(2)\Phi}(b))\} \\ &= \xi_{(1)\Phi}(a) \star \xi_{(2)\Phi}(b). \end{aligned}$$
(27)

The same holds for the rest of the universal envelope; for any module algebra A of $\mathbf{U}(\Xi)$, there exists a module A_{\star} of $\mathbf{U}(\Xi)^{\Phi}$. Consequently, the algebras of vector fields and tensorfields have a meaningful definition of their \star -product by using the correct form of the Lie derivative (commutator for vector fields and adjoint action for tensor fields). For example, in addition to \mathcal{A}_{\star} the deformed algebra $\mathbf{U}(\Xi_{\star})$ can be defined since any $(\xi, \eta) \in \mathbf{U}(\Xi)$ has a definitive action of $\phi \in \mathbf{U}(\Xi)$ on them,

$$\boldsymbol{\xi} \star \boldsymbol{\eta} \coloneqq \bar{\boldsymbol{\phi}}^{\{\alpha\}}(\boldsymbol{\xi}) \bar{\boldsymbol{\phi}}_{\{\alpha\}}(\boldsymbol{\eta}) = \mathcal{L}_{\bar{\boldsymbol{\phi}}^{\{\alpha\}}}(\boldsymbol{\xi}) \mathcal{L}_{\bar{\boldsymbol{\phi}}_{\{\alpha\}}}(\boldsymbol{\eta}), \qquad (28)$$

and, the algebra is also a $\mathbf{U}(\Xi)^{\Phi}\text{-module},$ as appears from,

$$\zeta(\xi \star \eta) = \zeta_{(1)\Phi}(\xi) \star \zeta_{(2)\Phi}(\eta), \qquad (29)$$

for all $\zeta \in \mathbf{U}(\Xi)$. These properties lead to a deformed

commutator that closes in Ξ ,

$$[u^*, v] = u * v - \bar{R}^{\{\alpha\}}(v) * \bar{R}_{\{\alpha\}}(u)$$

= $[\bar{\phi}^{\{\alpha\}}(u), \bar{\phi}_{\{\alpha\}}(v)].$ (30)

Therefore, also Ξ_{\star} is a deformed Lie algebra (a \star -Lie algebra). However, any product of vector fields $(u, v) \in \Xi$ can be re-written as a product in $\mathbf{U}(\Xi_{\star})$ as $uv = \phi^{\{\alpha\}}(u) \star \phi_{\{\alpha\}}(v)$. Then, the \star -Lie algebra (30) can also be regarded as being generated by \star -vector fields, and consequently, $\mathbf{U}(\Xi_{\star})$ as the universal envelope of Ξ_{\star} .

As a universal envelope structure, $\mathbf{U}(\Xi_{\star})$ is a Hopf algebra ($\mathbf{U}(\Xi_{\star}), \star, \Delta_{\star}$). It has been established [41] that this Hopf algebra is isomorphic to the Hopf algebra on $\mathbf{U}(\Xi)^{\Phi}$: first, define the operator,

$$D(\xi) \coloneqq \bar{\phi}^{\{\alpha\}}(\xi)\bar{\phi}_{\{\alpha\}}, \quad \text{s.t.} \quad D(\xi\star\eta) = D(\xi)D(\eta). \tag{31}$$

With this operator, an isomorphism mapping of the coalgebra is defined,

$$\Delta_{\star} = (D^{-1} \otimes D^{-1}) \circ \Delta^{\Phi} \circ D, \qquad (32)$$

with a corresponding universal \mathcal{R}_{\star} -matrix,

$$\mathcal{R}_{\star} \coloneqq (D^{-1} \otimes D^{-1})(\mathcal{R}), \tag{33}$$

leading to a corresponding relation to that of (20),

$$\tau(\Delta_{\star}(\xi)) = \mathcal{R}_{\star} \star \Delta_{\star}(\xi) \star \mathcal{R}_{\star}^{-1}, \qquad (34)$$

concluding that $\mathbf{U}(\Xi_{\star})$ has a meaningful Hopf algebra structure, isomorphic to that of $\mathbf{U}(\Xi)^{\Phi}$. In principle, one can use the two structures to formulate noncommutative theories; however, $\mathbf{U}(\Xi_{\star})$ exhibits a particular property in its co-product, as derived from (32) [41]:

$$\Delta_{\star}(\Xi_{\star}) \subset \Xi_{\star} \otimes 1 + \mathbf{U}(\Xi_{\star}) \otimes \Xi_{\star}, \qquad (35)$$

which is crucial for a non-commutative algebraic structure to have a well-defined differential calculus [51].

Therefore, in formulating a gravity theory, where a deformed infinitesimal diffeomorphism is central, we shall use the $U(\Xi_*)$ structure.

To use $\mathbf{U}(\Xi_{\star})$, its module algebras must be constructed, which, following the previous analysis, coincide with the modules of $\mathbf{U}(\Xi)^{\Phi}$. Thus, \mathcal{A}_{\star} is a $\mathbf{U}(\Xi_{\star})$ module with the action given by the \star -Lie derivative (Sweedler notation is understood):

$$\mathcal{L}^{\star}_{\xi}(f) \coloneqq \mathcal{L}_{D(\xi)}(f), \tag{36}$$

$$\mathcal{L}^{\star}_{\xi}(f \star g) = \mathcal{L}^{\star}_{\xi(1)\star}(f) \star \mathcal{L}^{\star}_{\xi(2)\star}(g),$$

and similarly for the module $U(\Xi_{\star})$,

$$\mathcal{L}_{\xi}^{\star}(\tau) := \mathcal{L}_{D(\xi)}(\tau), \qquad (37)$$
$$\mathcal{L}_{\xi}^{\star}(\tau \star \tau') = \mathcal{L}_{\xi_{(1)\star}}^{\star}(\tau) \star \mathcal{L}_{\xi_{(2)\star}}^{\star}(\tau').$$

Note that here the product is a "*"-one, hence the product of actions of $(\xi, \eta) \in \mathbf{U}(\Xi_*)$ on some quantity will be $\mathcal{L}_{\xi}^* \mathcal{L}_{\eta}^* = \mathcal{L}_{\xi \star \eta}^*$. Then, the action of the *-Lie derivatives of $u \in \Xi_*$ on another $v \in \Xi_*$ or that of $\xi \in \mathbf{U}(\Xi_*)$ on $\zeta \in \mathbf{U}(\Xi_*)$ is the *-Lie commutator or the *-adjoint action respectively:

$$\mathcal{L}_{u}^{\star}(v) = [u, v], \quad \mathcal{L}_{\xi}^{\star}(\zeta) = adj_{\xi}^{\star}(\zeta) \equiv adj_{D(\xi)}(\zeta). \quad (38)$$

The generalization to the deformed algebra of tensorfields $\mathcal{T}_{\star} \in \Omega_{\star} \otimes ... \otimes \Omega_{\star} \otimes \Xi_{\star} ... \otimes \Xi_{\star}$ (with the *-1-forms denoted as Ω_{\star}) is now achieved through the *-tensorproduct,

$$\forall (\tau, \eta) \in \mathcal{T}_{\star} : \quad \tau \overset{\star}{\otimes} \eta = \bar{\phi}^{\{\alpha\}}(\tau) \otimes \bar{\phi}_{\{\alpha\}}(\eta), \qquad (39)$$

with the following non-commutativity,

$$\tau \star h = \mathcal{L}^{\star}_{\bar{\phi}^{\{\alpha\}}}(\tau) \mathcal{L}^{\star}_{\bar{\phi}_{\{\alpha\}}}(h) = \mathcal{L}^{\star}_{\bar{\phi}_{\{\alpha\}}}(h) \mathcal{L}^{\star}_{\bar{\phi}^{\{\alpha\}}}(\tau)$$
$$= \mathcal{L}^{\star}_{\bar{\mathrm{R}}^{\{\alpha\}}}(h) \star \mathcal{L}^{\star}_{\bar{\mathrm{R}}_{\{\alpha\}}}(\tau) = \bar{\mathrm{R}}^{\{\alpha\}}(h) \star \bar{\mathrm{R}}_{\{\alpha\}}(\tau).(40)$$

The action of $\mathbf{U}(\Xi_{\star})$ on a general tensor is determined by the *-Lie derivative, as in (37). The action on each component is derived using (38). Consequently, the algebra of tensorfields \mathcal{T}_{\star} is also a $\mathbf{U}(\Xi_{\star})$ -module algebra. Specifically, the *-Lie derivative along $u \in \Xi_{\star}$ possesses a deformed Leibniz rule:

$$\mathcal{L}_{u}^{\star}(h \star g) = \mathcal{L}_{u}^{\star}(h) \star g + \bar{\mathrm{R}}^{\{\alpha\}}(h) \star \mathcal{L}_{\bar{\mathrm{R}}_{\{\alpha\}}(u)}^{\star}(g), \quad (41)$$

which agrees with the \star -co-product (32).

With the above definitions of *-vector and *-tensor fields, it is apparent that the notion of commutative *pairing* of *-vector fields and *-1-forms must be deformed (to maintain compatibility with the *-Lie derivative and \mathcal{A}_* -Linearity, see [41]). Thus a *-pairing is introduced,

$$\forall (\xi \in \Xi_{\star}, w \in \Omega_{\star}) : (\xi, w) \mapsto \langle \xi^{\star}, w \rangle \coloneqq \langle \bar{\phi}^{\{\alpha\}}(\xi), \bar{\phi}_{\{\alpha\}}(w) \rangle.$$

$$(42)$$

A proper basis of \star -vector fields and \star -1-forms must satisfy the duality condition dictated from (42). Choosing the \star -1-form basis as in the commutative scenario leads to,

$$\langle \overset{\star}{\partial}_{\mu} \overset{\star}{,} dx^{\nu} \rangle = \delta^{\nu}_{\mu} \Rightarrow \overset{\star}{\partial}_{\mu} = N^{-1_{\star}\nu}_{\mu} \star \partial_{\nu},$$

$$\text{where, } N^{\nu}_{\mu} = \langle \partial_{\mu} \overset{\star}{,} dx^{\nu} \rangle,$$

$$(43)$$

but this product does not generally have to be equal to δ^{ν}_{μ} as if it were classical. This definition concludes the introduction of the theory of non-commutative geometry necessary for the present work.

Let us now summarize the key ideas presented in this subsection. Formulas (6-8) define the algebra of functions on the non-commutative space, and (9, 10)define the form of the isomorphism to the commutative scenario. In (13), the twist element was defined, with the twist-deformed Hopf algebra of vector fields in (15). Using these definitions, we have shown the shape of the \star -product isomorphism map for functions in (19) and the representations of the deformed algebra that are consistent with a quantum differential calculus in (36-38). Additional noteworthy formulas include the deformed Leibniz rule (41), the deformed definition of basis vector fields (43), and each of the non-commutative products (e.g., (39) for the tensor field representation).

C. General Relativity on Non-Commutative Spaces

Here, we demonstrate how the formulation from the previous section can be used to construct a deformed version of Einstein's equation on non-commutative spacetime given its flat limit; we again follow [41]. The underlying philosophy is that GR is a theory with an action invariant under infinitesimal diffeomorphisms, represented by vector fields and their Lie algebra structure. The previous section showed that the structure of infinitesimal diffeomorphism can be adapted to the non-commutative space scenario, with consistent operations defining a differential geometry. Consequently, the construction will follow the classical approach to define the appropriate deformed geometrical quantities that are *-isomorphic to the classical quantities in GR. Thus, the Einstein equation can be composed similarly to the classical scenario, resulting in an equation whose associated action is invariant under *deformed* infinitesimal diffeomorphisms. Note that in contracts to the usual construction of GR, geometrical definitions, like the covariant derivative, curvature tensor, and so on, come before introducing a compatible metric. We begin with the non-commutative version of the covariant derivative.

-Covariant Derivative: The map $\hat{\nabla}_u : \Xi_ \to \Xi_*$ is defined such that the following holds for all $(u, v, z) \in \Xi_*$ and $h \in \mathcal{A}_*$ [41]:

$$\begin{aligned} \overset{\star}{\nabla}_{u+v} z &= \overset{\star}{\nabla}_{u} z + \overset{\star}{\nabla}_{v} z, \quad \overset{\star}{\nabla}_{h \star u} v = h \star \overset{\star}{\nabla}_{u} v, \\ \overset{\star}{\nabla}_{u} (h \star v) &= \mathcal{L}_{u}^{\star}(h) \star v + \bar{\mathrm{R}}^{\{\alpha\}}(h) \star \overset{\star}{\nabla}_{\bar{\mathrm{R}}_{\{\alpha\}}(u)}(v). \end{aligned}$$

$$(44)$$

In a local coordinate neighborhood **U** with coordinate set $\{x^{\mu}\}$ and a suitable, *-dual basis according to (43), the non-commutative connection coefficients are determined by²:

$$\overset{\star}{\nabla}_{\overset{\star}{\partial}_{\mu}}\overset{\star}{\partial}_{\nu} = \Gamma^{\sigma}_{\mu\nu} \star \overset{\star}{\partial}_{\sigma}, \quad \overset{\star}{\nabla}_{\overset{\star}{\partial}_{\mu}}^{\star} dx^{\nu} = -\Gamma^{\nu}_{\mu\sigma} \star dx^{\sigma}, \quad (45)$$

where the *-covariant derivative of a function $h \in \mathcal{A}_{\star}$ is

 $^{^2}$ The classical counterpart of this relation is identical, just drop the *-products and use the classical ∂_{μ} 's.

the *-Lie derivative:

$$\stackrel{\star}{\nabla}_{u}(h) = \mathcal{L}_{u}^{\star}(h) = \mathcal{L}_{D(u)}(h).$$
(46)

The \star -covariant derivative of a general \star -tensorfield (of a \star -bi-vector field, of a \star -1-form, etc.) is defined using the deformed Leibniz rules:

$$\stackrel{\star}{\nabla}_{u}(\tau \overset{\star}{\otimes} \tau') = \stackrel{\star}{\nabla}_{u}(\tau) \overset{\star}{\otimes} \tau' + \bar{\mathrm{R}}^{\{\alpha\}}(\tau) \overset{\star}{\otimes} \stackrel{\star}{\nabla}_{\bar{\mathrm{R}}_{\{\alpha\}}(u)}(\tau').$$
(47)

We Now define a ***-Curvature** for all $(u, v, z) \in \Xi_*$:

$$\mathfrak{X}(u,v,z) \coloneqq \overset{\star}{\nabla}_{u} \overset{\star}{\nabla}_{v} z - \overset{\star}{\nabla}_{\bar{\mathrm{R}}^{\{\alpha\}}(v)} \overset{\star}{\nabla}_{\bar{\mathrm{R}}_{\{\alpha\}}(u)} z - \overset{\star}{\nabla}_{[u,v]} z.$$
(48)

In a local coordinate neighborhood, the expression (48) can be written using its components , and a \star -Riemann tensor and a \star -Ricci tensor are then defined,

$$\mathbf{\mathfrak{X}}^{\sigma}_{\mu\nu\rho} \star \overset{*}{\partial}_{\sigma} = \mathbf{\mathfrak{K}}(\overset{*}{\partial}_{\mu}, \overset{*}{\partial}_{\nu}, \overset{*}{\partial}_{\rho}), \qquad (49)$$

$$\mathbf{\mathfrak{X}}_{\mu\nu} = \mathbf{\mathfrak{K}}(\overset{*}{\partial}_{\mu}, \overset{*}{\partial}_{\nu}) \coloneqq \langle dx^{\sigma}; \mathbf{\mathfrak{K}}(\overset{*}{\partial}_{\sigma}, \overset{*}{\partial}_{\mu}, \overset{*}{\partial}_{\nu}) \rangle'.$$

The "'" in the contraction stresses that the contraction is of a form on the left and a vector field on the right. The ***-Metric tensor** which we call **g** can be constructed by considering a symmetric rank-two ***-**tensor in \mathcal{T}_{\star} ,

$$\mathbf{g} = \mathbf{g}^a \overset{\star}{\otimes} \mathbf{g}_a \in \Omega_\star \overset{\star}{\otimes} \Omega_\star, \tag{50}$$

the left \mathcal{A}_{\star} -linear map is well defined for $(u, v) \in \Xi_{\star}$,

$$\begin{aligned}
\mathbf{g} &: \Xi_{\star} \overset{*}{\otimes} \Xi_{\star} \to \mathcal{A}_{\star}, \\
(u, v) \mapsto \mathbf{g}(u, v) = \langle u \otimes v^{*}; \mathbf{g} \rangle \coloneqq \langle u^{*}_{\star} \langle v^{*}; \mathbf{g} \rangle \rangle.
\end{aligned} \tag{51}$$

In a local neighborhood, with a suitable *-basis, one can write (see [41]) for the *-metric component, and for its associated *-inverse metric³ g^{-1^*} ,

$$\mathbf{g} = dx^{\nu} \overset{\star}{\otimes} dx^{\mu} \star \mathbf{g}_{\mu\nu}, \quad \mathbf{g}^{-1^{\star}} = \overset{\star}{g}^{\mu\nu} \star \overset{\star}{\partial}_{\nu} \overset{\star}{\otimes} \overset{\star}{\partial}_{\mu}. \tag{52}$$

The condition of the metric being compatible as a manifold (assuming a torsion-free metric) is,

$$\nabla_u^* \mathbf{g} = 0. \tag{53}$$

Then, the connection can be determined in terms of the metric, and the ***-Ricci scalar** can be constructed,

$$\mathbf{\mathfrak{X}} = \mathbf{g}^{\star \mu \nu} \star \mathbf{\mathfrak{X}}_{\nu \mu}.$$
 (54)

Establishing these definitions for the geometrical objects, it is now possible to write a deformation of GR

on non-commutative spacetime (see the discussion in [42]) through the deformed Einstein equation that is by definition invariant under deformed infinitesimal diffeomorphisms. Thus, we derive the Einstein Tensor and the Einstein Equation in vacuum,

$$\mathfrak{G}_{\mu\nu} = \mathfrak{R}_{\mu\nu} - \frac{1}{2} \mathbf{g}_{\mu\nu} \star \mathfrak{R} = 0.$$
 (55)

This equation can be perturbatively expanded in the deformation parameter λ , which appears in the twist deformation (13).

In summary, we now have all the formulations necessary for constructing a GR theory on a noncommutative spacetime. The main features to take from this formulation are the deformed actions of the various Lie derivatives \mathcal{L}^* , the *-product appearing in all definitions, and the deformed Leibniz rules (such as (47)). Finally, we are left with equation (55), describing the dynamics of a non-commutative space.

III. TESTING THE METHOD: USING THE NEW GENERAL DESCRIPTION TO RE-DERIVE THE KNOWN CASE OF CONSTANT NON-COMMUTATIVITY

In this section, we re-derive the deformed Einstein equation that extends a constant-non-commutative spacetime. We do this using the prescription outlined in Sec.II, covering the foundations in Sec.III A, and the gravity construction (introducing deformed quantities for the curvature tensor, the Ricci tensor, etc.) in Sec.III B. These results were derived, e.g., in [43, 44] using a simplistic method only applicable in the case of constant non-commutativity; we derive them more generally and compare them to the known results. We thus gain crucial insight into how to derive the non-constant case. To keep the discussion compact, we will present the key steps here and provide the detailed derivations of various expressions in Appendix A.

Note that, until now, we have discussed the general formalism of non-commutative gravity via a twist without specifying the form of the twist. Generally, there are various forms of twists, such as the Jordanian twists, extended Jordanian twists (see [41] and references therein), etc. However, for our present discussion in both the constant and non-constant noncommutativity, we will use the general form of *Abelian Twists*, defined with vector fields $(X_a, X_b) \in \Xi$ [46, 52]:

$$\Phi = \exp\left[-\frac{i}{2}\lambda\Theta^{ab}X_a \otimes X_b\right] = \phi^{\{\alpha\}} \otimes \phi_{\{\alpha\}},$$

$$\Phi^{-1} \equiv \bar{\Phi} = \exp\left[+\frac{i}{2}\lambda\Theta^{ab}X_a \otimes X_b\right] \equiv \bar{\phi}^{\{\alpha\}} \otimes \bar{\phi}_{\{\alpha\}}.(56)$$

The vector fields mutually commute (hence the name), and the matrix Θ^{ab} is anti-symmetric with constant entries. The reason for solely using Abelian twists will become more apparent in section Sec.IV.

³ From the component expression one can understand the " \star " in the metric inverse; we demand that $\mathbf{g}_{\mu\nu}\star\overset{\star}{\mathbf{g}}^{\nu\rho}=\delta^{\rho}_{\mu}$

A. Basic Formulae & Calculations

Here, we consider a non-commutative spacetime locally described by spacetime with constant non-commutativity. Therefore, we only keep the first (constant) term from (7),

$$\left[\hat{x}^{\mu}, \hat{x}^{\nu}\right] = \left[x^{\mu}, x^{\nu}\right] = i\lambda\theta^{\mu\nu} = -i\lambda\theta^{\nu\mu}, \qquad (57)$$

where λ is some small parameter with dimensions of length squared and $\theta^{\mu\nu}$ is a constant anti-symmetric tensor. The twist of the form (56) that will generate this *-isomorphism (see (9)) is [53]:

$$\Phi_{\theta} = \exp\left[-\frac{i}{2}\lambda\theta^{\mu\nu}\partial_{\mu}\otimes\partial_{\nu}\right].$$
 (58)

Here, we have set Θ^{ab} in (56) to $\theta^{\mu\nu}$. The permutation operator (20) will be (see (A1)):

$$\mathcal{R} = \Phi_{21} \Phi^{-1} = \Phi^{-2} = \exp[+i\lambda\theta^{\mu\nu}\partial_{\mu} \otimes \partial_{\nu}].$$
 (59)

To see how the twist (58) generates (57), we first observe that the *-product (19) takes the form

$$f \star g = fg + \frac{i}{2}\lambda\theta^{\mu\nu}(\partial_{\mu}f)(\partial_{\nu}g) + \mathcal{O}(\lambda^2), \qquad (60)$$

then, using the definitions (30) & (13), we directly calculate the spacetime commutator:

$$[x^{\mu} * x^{\nu}] = [\bar{\phi}^{\{\alpha\}}(x^{\mu}), \bar{\phi}_{\{\alpha\}}(x^{\nu})]$$

$$= \bar{\phi}^{\{\alpha\}}(x^{\mu})\bar{\phi}_{\{\alpha\}}(x^{\nu}) - \bar{\phi}_{\{\alpha\}}(x^{\nu})\bar{\phi}^{\{\alpha\}}(x^{\mu})$$

$$= x^{\mu}x^{\nu} + \frac{i}{2}\lambda\theta^{\rho\sigma}(\partial_{\rho}\otimes\partial_{\sigma})(x^{\mu}\otimes x^{\nu}) + \mathcal{O}(\partial^{2}(x^{\mu}))$$

$$-x^{\nu}x^{\mu} - \frac{i}{2}\lambda\theta^{\sigma\rho}(\partial_{\sigma}\otimes\partial_{\rho})(x^{\nu}\otimes x^{\mu}) - \mathcal{O}(\partial^{2}(x^{\mu}))$$

$$= \frac{i}{2}\lambda[\theta^{\mu\nu} - \theta^{\nu\mu}] = i\lambda\theta^{\mu\nu}, \qquad (61)$$

confirming that (58) yields (57).

The first step towards constructing the Einstein Tensor (55) is establishing a set of basis vectors satisfying (43). However, we note that the action of the twist (58) on the classical basis vectors $\{\partial_{\mu}\}$ is through the Lie derivative of ∂_{μ} along the twist' vector fields, which is identically zero (they commute by definition), therefore:

$$N^{\nu}_{\mu} = \langle \partial_{\mu} , dx^{\nu} \rangle = \delta^{\nu}_{\mu} \Rightarrow \partial_{\mu} = \partial^{\star}_{\mu}.$$
 (62)

Then, in any formula from section Sec.II where ∂_{μ} appears, we can replace it with ∂_{μ} . Also, since the action of the twist on ∂_{μ} is trivial, expressions of the form $\partial_{\mu'} \star O^{\mu'}_{\mu}[\lambda, 1]$ will have no contribution from the \star -deformation. As a demonstration, consider the right-

hand side in (45),

$$\Gamma^{\sigma}_{\mu\nu} \star \partial_{\sigma} = \bar{\phi}^{\{\alpha\}} (\Gamma^{\sigma}_{\mu\nu}) \bar{\phi}_{\{\alpha\}} (\partial_{\sigma})$$
$$= \Gamma^{\sigma}_{\mu\nu} \partial_{\sigma} + \frac{i}{2} \theta^{\gamma\delta} \partial_{\gamma} (\Gamma^{\sigma}_{\mu\nu}) \partial_{\delta} (\partial_{\sigma}) + \dots$$
$$= \Gamma^{\sigma}_{\mu\nu} \partial_{\sigma} + 0 = \Gamma^{\sigma}_{\mu\nu} \partial_{\sigma}, \qquad (63)$$

where we have used the fact that after using the twist element, the variables are the commutative ones: the term $\partial_{\gamma}(\partial_{\sigma})$ vanishes as $\mathcal{L}_{\gamma}(\partial_{\sigma}) = (\partial_{\gamma}(1) - \partial_{\gamma}(1))\partial_{\sigma} = 0$. Thus, the *-deformation indeed vanishes. This is also true for functions that are *-multiplied with the 1-form basis vectors.

Thus, the expressions for the connection (45) and the \star -Lie derivative (36) can be simplified (respectively) for vector fields $\xi = \xi^{\mu} \star \overset{\star}{\partial}_{\mu} \in \Xi_{\star}$ such that ξ^{μ} is constant, to

$$\stackrel{\star}{\nabla}_{\partial_{\mu}}\partial_{\nu} = \Gamma^{\sigma}_{\mu\nu}\partial_{\sigma}, \quad \stackrel{\star}{\nabla}_{\partial_{\mu}}dx^{\nu} = -\Gamma^{\nu}_{\mu\sigma}dx^{\sigma}, \quad (64)$$

$$\mathcal{L}_{\xi}^{\star}(h) = \xi^{\mu} \mathcal{L}_{\partial_{\mu}}^{\star}(h) = \xi^{\mu} \mathcal{L}_{\bar{\phi}^{\{\alpha\}}(\partial_{\mu})\bar{\phi}_{\{\alpha\}}}(h) = \mathcal{L}_{\xi}(h).$$
(65)

By the same reasoning, the *-Lie derivative of tensors ((37),(38)) along a constant vector field ξ^{μ} will also reduce to the classical Lie derivative. When the vector field ξ^{μ} is not constant, the expression for \mathcal{L}_{ξ}^{*} gets more complicated. We state here the result, as calculated in (A3) (see also [43])

$$\mathcal{L}_{\xi}^{*}(h) = \sum_{n=0}^{\infty} \frac{1}{n!} (\frac{i\lambda}{2})^{n} \prod_{j=1}^{n} \theta^{\mu_{j}\nu_{j}} \frac{\partial^{n}(\xi^{\gamma})}{\partial^{\mu_{1}} ... \partial^{\mu_{n}}} \frac{\partial^{n}(\partial_{\gamma}h)}{\partial^{\nu_{1}} ... \partial^{\nu_{n}}}.$$
 (66)

B. The Construction of Gravity

We can now turn to construct the deformed Einstein Tensor (55) on the non-commutative spacetime (57) by following Sec.II C and the results outlined in Sec.III A.) Starting with the ***-Covariant Derivative**, we use the general formula (44) to get (see (A4)),

$$\overset{\star}{\nabla}_{z} u = \mathcal{L}_{z}^{\star}(u^{\nu})\partial_{\nu} + (\bar{\phi}^{\{\alpha\}}(z^{\mu})\bar{\phi}_{\{\alpha\}}(u^{\nu})) \star \Gamma^{\sigma}_{\mu\nu}\partial_{\sigma}$$

$$\Rightarrow \overset{\star}{\nabla}_{\mu}u^{\nu} = \partial_{\mu}u^{\nu} + u^{\sigma} \star \Gamma^{\nu}_{\sigma\mu}.$$

$$(67)$$

The ***-Curvature Tensor:** We have the following properties for the constant non-commutative scenario,

$$\mathfrak{K}(\partial_{\mu},\partial_{\nu},\partial_{\rho}) = \mathfrak{K}^{\sigma}_{\mu\nu\rho}\partial_{\sigma} , \quad [\partial_{\mu}{}^{*},\partial_{\nu}] = 0.$$
(68)

Then, by utilizing equation (49), we derive the coordinate-basis expression for the *-curvature tensor (see (A5)),

$$\mathfrak{X}^{\sigma}_{\mu\nu\rho} = \partial_{[\mu}\Gamma^{\sigma}_{\nu]\rho} + \Gamma^{\tau}_{\rho[\nu} \star \Gamma^{\sigma}_{\mu]\tau} \tag{69}$$

This expression is *not* identical to the classical one; a deformation appears in the \star -product between the Christoffel symbols.

The \star -Ricci Tensor: using (49) and (69) we derive (see (A9)),

$$\mathfrak{X}_{\mu\nu} = \langle dx^{\mu} \mathfrak{X}(\partial_{\mu}, \partial_{\nu}, \partial_{\rho}) \rangle = \delta^{\mu}_{\sigma} \mathfrak{X}^{\sigma}_{\mu\nu\rho} \Rightarrow \mathfrak{X}_{\nu\rho} = \mathfrak{X}^{\mu}_{\mu\nu\rho} .$$
(70)

As in classical GR, the *-Ricci tensor is a simple contraction of the *-curvature tensor.

The *-Metric Tensor: Using the *-metric definitions ((50), (51), (52)), we observe that the *-metric can be considered as an undetermined, geometrical, dynamical object that, when solved, provides a structure consistent with the non-commutative spacetime. There is no need to expand it in terms of a classical metric in contracts with the other geometrical quantities. Indeed, Ref. [42] showed that any *-metric is also a commutative metric, in the sense that it is a (non-degenerate) symmetric rank two tensor. Therefore, we can treat the *-metric tensor as a regular metric tensor with the component expression given by (52), keeping in mind that it will be solved via (55), not by the classical Einstein equation. Nevertheless, for the *-metric inverse, we do need to care for the " \star " in it, since the condition $g_{\mu\nu} \star g^{\rho\mu} = \delta_{\nu}^{\rho}$ is not trivial (see eq.(8.18) in [44] for an expansion in the usual inverse $g^{\mu\nu} = (g_{\mu\nu})^{-1}$ and the deformation parameter).

Next, we evaluate the relation between the metric tensor and the Cristoffel symbols, utilizing (53). First, the *-covariant derivative of the metric tensor equals (see (A10)) to,

$$\overset{\star}{\nabla}_{\partial_{\gamma}}(\mathbf{g}) = \overset{\star}{\nabla}_{\gamma}(dx^{\mu} \overset{\star}{\otimes} dx^{\nu} \star g_{\mu\nu})$$

$$= [g_{\mu\nu,\gamma} - \Gamma^{\sigma}_{\gamma(\mu} \star g_{\nu)\sigma}](dx^{\mu} \overset{\star}{\otimes} dx^{\nu}).$$

$$(71)$$

Therefore, the compatibility condition is,

$$\stackrel{\star}{\nabla}_{u}(\mathbf{g}) = 0 \Leftrightarrow g_{\mu\nu,\gamma} = \Gamma^{\sigma}_{\gamma(\mu} \star g_{\nu)\sigma}.$$
(72)

By following the classical procedure, we establish the desired relation to be (see (A14)),

$$\Gamma^{\sigma}_{\nu\gamma} = \overset{*}{g}^{\sigma\mu} \star \frac{1}{2} [g_{\mu(\nu,\gamma)} - g_{\nu\gamma,\mu}].$$
(73)

The \star -Ricci Scalar is derived by use of (54),

$$\mathbf{\mathfrak{K}} = \overset{*}{g}^{\mu\nu} \star \mathbf{\mathfrak{K}}^{\sigma}_{\sigma\mu\nu}. \tag{74}$$

This concludes our analysis of deformed GR defined on a constantly non-commutative spacetime. We computed closed expressions for all of the *-geometrical objects that can be used to construct the Einstein Tensor (55). Regarding solutions to (55), note that $\Re_{\mu\nu}$ is not real nor hermitian. Hence, the resulting equation is not solvable (only trivially). The solution is to force it to be real/hermitian by appropriately symmetrising it, see [41, 42, 47].

All the listed results agree with those found by differ-

ent means in [44] and others. For comparison, consider the following results in [44]; Eq.(7.6) for their version for the *-Lie derivative (66), Eq.(7.13) for the computation 0) of the *-covariant derivative (67), Eq.(7.17) for the Riemann tensor (69), Eq.(8.23) for the deformed relation (73), and finally, Eq.((9.4),(9.5)) for the *-Ricci tensor & scalar ((74),(70)) respectively.

IV. BUILDING BLOCKS OF GEOMETRY IN κ-MINKOWSKI (POSITION-DEPENDENT) NON-COMMUTATIVITY

Here, we shall outline known and new results that will pave the road for our desired construction of a noncommutative gravity theory (of the Einstein Tensor) that extends a local κ -Minkowski spacetime, in which the non-commutativity is position-dependent. For the first time, we give the explicit action of the twist on the basis of vector fields and one-forms. These lead to a crisp characterization of the deformed basis and Leibniz rule, which we verify using our previous results [43]. Again, the detailed derivations of various expressions are found in Appendix B to keep the discussion compact.

The κ -Minkowski spacetime's non-commutativity (see Sec.I), can be written in the *operator* representation as

$$\left[\hat{x}^{0}, \hat{x}^{i}\right] = \frac{i}{\kappa} \hat{x}^{i} , \quad \left[\hat{x}^{i}, \hat{x}^{j}\right] = 0 , \quad i, j = 1, 2, 3.$$
(75)

this is equivalent to the second term in (7) with the anti-symmetric constant tensor $C^{\mu\nu}_{\rho}$, while setting the first term $(\theta^{\mu\nu})$ to zero:

$$C^{\mu\nu}_{\rho} = \frac{1}{\kappa} \left(\delta^{\mu}_{0} \delta^{\nu}_{\rho} - \delta^{\nu}_{0} \delta^{\mu}_{\rho} \right) \coloneqq \lambda \delta^{\mu}_{[0} \delta^{\nu}_{\rho]}.$$
(76)

We would now like to utilize the \star -isomorphism (9) for this non-commutativity. In principle, it is possible to construct a *-product suitable for this noncommutativity that will be a module algebra of the κ -Poincaré algebra (see Sec.I and [25] with references therein) and to derive a deformation quantization procedure from establishing gauge theories on that space. At first sight, this seems a preferable approach; it has a κ -Minkowski spacetime structure, preserves SR symmetries in the classical limit, and adds a fundamental length scale as an additional invariant. However, when one computes the suitable differential calculus [51, 54] (by considering a map of the ideal in (8)) and the associated *-product [55, 56], a profound ambiguity arises: the integral on *-products of fields is not cyclic. Hence, no gauge theory can be built out with this *-product [25, 43]. Several approaches exist to bypass this ambiguity: one is to consider a measure function $\mu(x)$ that will make the integral cyclic [57, 58]; however, such a measure function will spoil the classical limit $\lambda \to 0$ since it is not identity there. Others are to consider different notions of integration [59], etc., see [43, 60, 61] for ongoing research on the subject.

Here, we consider a resolution found in [43]: to approach the spacetime (75) as a result of an abelian twist deformation. This way, we are guaranteed to have a meaningful definition of the integral. However, to obtain (75), we must consider a twist that is not generated by the Poincaré algebra but rather by the algebra of the IGL(3,1) group so that the κ -Poincaré group will no longer be the symmetry group. We will discuss the implications and suggestions to solve this problem of symmetry in Sec.VI. In the following, we follow the twist formalism and construct a deformed GR theory on the spacetime (75).

For that aim, consider first the form of the twist (56) to be [43] (we use $\lambda \coloneqq 1/\kappa = l_p$ for brevity),

$$\Theta^{00} = 0, \ \Theta^{01} = 1, \ \Theta^{10} = -1, \ \Theta^{11} = 0,$$

$$X_0 = \partial_0, \ X_1 = x^j \partial_j \in IGL(3, 1).$$
(77)

Note that these vector fields are not constrained to be killing vector fields as in [42, 47]. Writing the twist explicitly, we have,

$$\Phi = \exp\left[-i\frac{\lambda}{2}\left(\partial_0 \otimes x^j\partial_j - x^j\partial_j \otimes \partial_0\right)\right].$$
(78)

The associated permutation operator (20) is,

$$\mathcal{R} = \Phi_{21}\Phi^{-1} = \exp\left[+i\lambda\left(\partial_0 \otimes x^j\partial_j - x^j\partial_j \otimes \partial_0\right)\right].$$
(79)

As in Sec.III, we first want to show that (78) implies (75). Thus, we calculate the *-product to second order,

$$f \star g = fg + i\frac{\lambda}{2}x^{j}f_{,[0}g_{,j]}$$

$$+ \frac{1}{8}(i\lambda)^{2}x^{j}x^{l}(f_{,0[0} \cdot g_{,l]j} + f_{,j[l} \cdot g_{,0]0}) + \mathcal{O}(\lambda^{3}).$$
(80)

The form of $\bar{\phi}_{\{\alpha\}} \otimes \bar{\phi}^{\{\alpha\}}$ for an Abelian twist is,

$$\bar{\phi}_{\{\alpha\}} \otimes \bar{\phi}^{\{\alpha\}} = \exp\left[i\frac{\lambda}{2}\Theta^{ba}X_b \otimes X_a\right].$$
 (81)

Using (77), we can explicitly calculate (30) to derive,

$$\begin{split} [x^{\mu} * x^{\nu}] &= \bar{\phi}^{\{\alpha\}}(x^{\mu})\bar{\phi}_{\{\alpha\}}(x^{\nu}) - \bar{\phi}_{\{\alpha\}}(x^{\nu})\bar{\phi}^{\{\alpha\}}(x^{\mu}) \\ &= x^{\mu}x^{\nu} + i\frac{\lambda}{2}x^{j}\delta^{\mu}_{[0}\delta^{\nu}_{j]} - x^{\nu}x^{\mu} - i\frac{\lambda}{2}x^{j}\delta^{\nu}_{[0}\delta^{\mu}_{j]} \\ &= i\lambda x^{j}\delta^{\mu}_{[0}\delta^{\nu}_{j]} \qquad (82) \\ \Rightarrow [x^{0} * x^{j}] &= i\lambda x^{j} = -[x^{j} * x^{0}], \end{split}$$

in accordance with (2), confirming that (78) is a suitable twist. The next step is constructing a suitable *-dual basis of vector fields using the condition (43). Thus, we must compute the action of the twist on 1-forms and vectors, i.e., to calculate the Lie derivative (39) along the twists' vector fields (77) of the basis 1-forms and basis vector fields. Note that in [38], the basis was found by different means. However, we calculate with the *- dual basis construction; by comparing the results with those in [38], we can verify our side calculations of the twist's actions on the bases. Then, we can use them for calculations in the next section V.

The vector fields act on basis 1-forms as (see (B1)),

$$X_0(dx^{\mu}) = 0, \quad X_1(dx^{\mu}) = \delta^j_{\mu} dx^{\mu} \neq 0.$$
 (83)

And, on the basis vectors as (see (B2)),

$$X_0(\partial_\mu) = 0, \quad X_1(\partial_\mu) = -\delta^j_\mu \partial_j. \tag{84}$$

Thus, we are not dealing with a *nice* basis (as defined in [47]); the following calculations will not coincide with theirs. The formulae ((83),(84)) dictate the action of the twist on the classical basis of vector fields and of 1-forms, which we now outline. However, we first define a notation for brevity:

$$O^{\mu'}_{\mu}[\lambda,n] \equiv e^{in\frac{\lambda}{2}\partial_0}\delta^j_{\mu}\delta^{\mu'}_j + \delta^0_{\mu}\delta^0_0^{\mu'}.$$
 (85)

Using this notation, we now write the twist's action:

$$\phi^{\{\alpha\}}(\partial_{\mu})\phi_{\{\alpha\}} = O^{\mu'}_{\mu}[\lambda, -1]\partial_{\mu'}$$
(86a)

$$\bar{\phi}^{\{\alpha\}}(\partial_{\mu})\bar{\phi}_{\{\alpha\}} = O^{\mu'}_{\mu}[\lambda, +1]\partial_{\mu'} \tag{86b}$$

$$\phi^{\{\alpha\}}(dx^{\mu})\phi_{\{\alpha\}} = O^{\mu}_{\mu'}[\lambda, +1]dx^{\mu'}$$
(86c)

$$\bar{\phi}^{\{\alpha\}}(dx^{\mu})\bar{\phi}_{\{\alpha\}} = O^{\mu}_{\mu'}[\lambda, -1]dx^{\mu'}$$
 (86d)

By use of (43), we now explicitly derive the \star -pairing of the classical basis,

$$\begin{aligned} \langle \partial_{\mu} ^{*} dx^{\nu} \rangle &= \bar{\phi}^{\{\alpha\}} (\partial_{\mu}) \bar{\phi}_{\{\alpha\}} (x^{\nu}) = O^{\mu'}_{\mu} [\lambda, 1] \partial_{\mu'} (x^{\nu}) \\ &= O^{\mu'}_{\mu} [\lambda, 1] \delta^{\nu}_{\mu'} = O^{\nu}_{\mu} [\lambda, 1]. \end{aligned}$$
(87)

The *-dual basis is therefore (denoted with a " * "),

$$\overset{*}{\partial_0} = \partial_0, \ \overset{*}{\partial_j} = e^{-i\frac{\lambda}{2}\partial_0}\partial_j \iff \langle \overset{*}{\partial_\mu} \overset{*}{,} dx^\nu \rangle = \delta^\nu_\mu.$$
(88)

In [38], various results were derived through a different method from ours. We now recalculate them using our method and results to mutually validate them.

The non-commutativity of basis 1-forms: by definition we have,

$$f \star dx^{\mu} = \bar{\mathbf{R}}^{\{\alpha\}}(dx^{\mu}) \star \bar{\mathbf{R}}_{\{\alpha\}}(f), \qquad (89)$$

then, utilizing (83) and (86), we derive (see (B4)),

$$f \star dx^{\mu} = dx^{\mu'} \star O^{\mu}_{\mu'}[\lambda, 2](f),$$
(90)

in agreement with the known results in [45].

The deformed Leibniz rule of *-basis vectors: in [45] they derived the appropriate Leibniz rule by first considering:

$$\mathbf{d}f = (\partial_{\mu}f)dx^{\mu} = (\partial_{\mu}f) \star dx^{\mu}, \qquad (91)$$
$$\mathbf{d}(f \star g) = \mathbf{d}f \star g + f \star \mathbf{d}g = (\overset{\star}{\partial}_{\mu}(f \star g)) \star dx^{\mu}.$$

Then, by equating between the first and the second equalities in the second line, they derived the following Leibniz rule (see (B8)),

$$\overset{\star}{\partial}_{\mu}(f \star g) = \overset{\star}{\partial}_{\mu'}(f) \star O^{\mu'}_{\mu}[\lambda, -2](g) + f \star (\overset{\star}{\partial}_{\mu}g).$$
(92)

The validity of (86) can now be confirmed by recalculating the Leibniz rule, this time with (36) and by calculating the *-co-product the *-derivative from (35); after calculations, we derive the *-co-product (see (B11)),

$$\Delta_{\star}(\overset{\star}{\partial}_{\mu}) = \overset{\star}{\partial}_{\mu'} \otimes O_{\mu}^{\mu'}[\lambda, -2] + 1 \otimes \overset{\star}{\partial}_{\mu}, \qquad (93)$$

which produces the deformed Leibniz rule (see (B12)),

$$\overset{\star}{\partial}_{\mu}(f \star g) = \overset{\star}{\partial}_{\mu'}(f) \star O^{\mu'}_{\mu}[\lambda, -2](g) + f \star (\overset{\star}{\partial}_{\mu}g), \quad (94)$$

in agreement with (92), supporting the result in (86).

This concludes our analysis of the foundations of geometry in the non-commutative spacetime (75). We note a significant difference compared to the constant non-commutativity outlined in Sec.III; no 'trivial' twist actions appear here; the differential calculus is not classical, etc. Consequently, we anticipate differences in the deformed GR, which we now develop and analyze.

V. GRAVITY EXTENSION OF THE κ -MINKOWSKI SPACETIME

We shall now derive the deformed Einstein equation (deformed Einstein Tensor) on a non-commutative spacetime with a local behavior of the κ -Minkowski spacetime. We follow a similar path as in Sec.III, starting by computing the \star -geometrical objects on the κ -Minkowski spacetime using Sec.II and Sec.IV. Again, the detailed derivations of various expressions are found in Appendix C to keep the discussion compact.

Using (36) and (86b), we calculate the ***-Lie derivative** for a constant vector field ξ ,

$$\mathcal{L}_{\xi^{\mu}\partial_{\mu}}^{*}(h) = \xi^{\mu}\mathcal{L}_{\bar{\phi}^{\{\alpha\}}(\partial_{\mu})\bar{\phi}_{\{\alpha\}}}^{*}(h) \qquad (95)$$
$$= \xi^{\mu}O_{\mu}^{\mu'}[\lambda,1]\partial_{\mu'}^{*}(h) = \xi^{\mu}\partial_{\mu}(h).$$

When the vector field is not constant, the \star -Lie derivative takes a complicated form. We will not directly use such an expression in this work; thus, we leave the calculation and the few first orders to (C1)

and (C3), stating now the general result:

$$\mathcal{L}_{\xi^{\mu}\partial_{\mu}}^{\star} = \sum_{\substack{n=0\\n'=0}}^{\infty} \frac{(i\lambda/2)^{n+n'}}{n!n'!} [(-x^{j}\partial_{j})^{n'} [\partial_{0}^{n}(\xi^{\mu})\partial_{\mu}^{\star}] \partial_{0}^{n'} (x^{j}\partial_{j})^{n}].$$
(96)

The connection coefficients are uniquely determined using (45),

$$\overset{*}{\nabla}_{\overset{*}{\partial}_{\mu}}\overset{*}{\partial}_{\nu} = \Gamma^{\sigma}_{\mu\nu} \star \overset{*}{\partial}_{\sigma} \neq \Gamma^{\sigma}_{\mu\nu}\overset{*}{\partial}_{\sigma}, \qquad (97)$$

holding similarly for the 1-form basis. Note that we will not explicitly expand each \star -product in the construction (it can be done in a future perturbative expansion using (80)).

The ***-Covariant Derivative** is calculated using (44), which, for generic vector fields u and z, simplifies to (see (C5)),

$$\stackrel{\star}{\nabla}_{z} u = \left[\mathcal{L}_{z}^{\star}(u^{\nu}) + \bar{\mathrm{R}}^{\{\alpha\}}(u^{\sigma}) \star \bar{\mathrm{R}}_{\{\alpha\}\mu}^{\mu'}(z) \star \Gamma_{\mu'\sigma}^{\nu} \right] \star \stackrel{\star}{\partial}_{\nu} (98)$$

For the case of $z = \hat{\partial}_{\mu}$ and only considering the $u = u^{\nu}$ part, we derive (C6)

$$\stackrel{\star}{\nabla}_{\stackrel{\star}{\partial}_{\mu}}(u^{\nu}) = \partial_{\mu}(u^{\nu}) + O^{\mu'}_{\mu}[\lambda, 2](u^{\sigma}) \star \Gamma^{\nu}_{\mu'\sigma} \tag{99}$$

Note, in (98), we expressed a *-vector field as $u = u^{\nu} \star \overset{\star}{\partial_{\nu}}$. However, if a vector field $u = u^{\nu} \overset{\star}{\partial_{\nu}}$ is given in the commutative vector space Ξ , its expression in the \star -vector space will be $u = \overset{\star}{u}^{\nu} \star \overset{\star}{\partial_{\nu}}$. The components $\overset{\star}{u}^{\nu}$ are related to the components in the commutative space by equating $u^{\nu} \overset{\star}{\partial_{\nu}}$ to $\overset{\star}{u}^{\nu} \star \overset{\star}{\partial_{\nu}}$ (see [41]). This relation can be solved order by order as $\overset{\star}{u}^{\nu} = \overset{\star}{u}^{\nu}_{(0)} + \lambda \overset{\star}{u}^{\nu}_{(1)} + \dots$ For example: $\overset{\star}{u}^{\nu}_{(0)} = u^{\nu} - \overset{\star}{u}^{\nu}_{(1)} = -\lambda \bar{\phi}^{\{\alpha\}}(u^{\rho}) \bar{\phi}^{\mu}_{\{\alpha\}\rho}$. Here,

 $\bar{\phi}^{\mu}_{\{\alpha\rho\}} \overset{\star}{\partial}_{\mu} = \overset{\star}{\partial}_{\rho}$. Nevertheless, we treat general *-vector fields in our calculations, so we can write $u = u^{\nu} \star \overset{\star}{\partial}_{\nu}$. We must keep in mind that to relate this quantity to its counterpart in the commutative space; we shall need to use the expression for $\overset{\star}{u}^{\nu}$ in terms of u^{ν} and λ .

On top of all that, expression (99) does not require using $\overset{*}{u}^{\nu}$, and the u^{ν} can also be considered as the component in $u \in \Xi$. This is because we can write (98) with $\overset{*}{u}^{\nu}$ and use $\overset{*}{u}^{\nu} \star \overset{*}{\partial}_{\nu} = u^{\nu} \overset{*}{\partial}_{\nu}$.

The \star -Curvature Tensor is calculated using (48) and (49), which, after evaluating, takes the form (see (C7)),

$$\mathfrak{X}^{\sigma}_{\mu\nu\rho} = \partial_{[\mu}\Gamma^{\sigma}_{\nu]\rho} + O^{\gamma}_{[\mu}[\lambda,2](\Gamma^{\tau}_{\nu]\rho}) \star \Gamma^{\sigma}_{\gamma\tau} \qquad (100)$$

As anticipated, there are additional deformations in the \star -curvature tensor compared to (69).

The ***-Ricci Tensor** is calculated using (49) while treating the *****-curvature as some given tensor-coefficients (see (C14))

$$\mathfrak{X}_{\nu\rho} = \langle dx^{\mu} \mathfrak{X}^{\sigma}_{\mu\nu\rho} \star \overset{\star}{\partial}_{\sigma} \rangle' = e^{-i\lambda\partial_0} \mathfrak{X}^j_{j\nu\rho} + \mathfrak{X}^0_{0\nu\rho}.$$
 (101)

To establish the relation between the metric and the Christoffel symbols, we first evaluate the metric compatibility condition (53) as follows (see (C17)),

$$\stackrel{\star}{\nabla}_{\gamma}(g) = \left[-\Gamma^{\sigma}_{\gamma(\mu} \star g_{\nu)\sigma} + g_{\mu\nu,\gamma} \star (dx^{\mu} \stackrel{\star}{\otimes} dx^{\nu}) \right]$$

$$\Rightarrow g_{\mu\nu;\gamma} = g_{\mu\nu,\gamma} - \Gamma^{\sigma}_{\gamma(\mu} \star g_{\nu)\sigma}.$$
(102)

Observe that (102) shares a similar form with its constant non-commutativity counterpart (72); a distinctive property of the metric since the basis 1-forms are \star multiplied from the left. For a general tensor, the covariant derivative will take the form

$$\nabla_{\gamma}(\tau_{\mu\nu} \star dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) = \begin{bmatrix} \tau_{\mu\nu,\gamma} - O_{\gamma}^{\gamma'}[\lambda, 2](\tau_{\sigma\nu}) \star \Gamma_{\gamma'\mu}^{\sigma} & (103) \\ -O_{\gamma}^{\gamma'}[\lambda, 2](\tau_{\mu\sigma}) \star \Gamma_{\gamma'\nu}^{\sigma} \end{bmatrix} \star (dx^{\mu} \overset{\star}{\otimes} dx^{\nu}),$$

differing in *form* from the \star -covariant derivative presented in Sec.III. Using the result (102) and following a similar procedure as in (73) in section Sec.III, we derive the \star -connection in terms of the metric:

$$\Gamma^{\sigma}_{\nu\gamma} = \overset{\star}{g}^{\sigma\mu} \star \Gamma_{\mu\nu\gamma} = \overset{\star}{g}^{\sigma\mu} \star \frac{1}{2} [g_{\mu(\nu,\gamma)} - g_{\nu\gamma,\mu}].$$
(104)

This result does not imply identity between the connection in κ -Minkowski spacetime with (73); the \star -product differs significantly. However, it does signify that the **form** of the deformation is the same.

The \star -Ricci scalar is calculated using (54) and (101),

$$\mathbf{\mathfrak{X}} = \overset{\star^{\mu\nu}}{g} \star \mathbf{\mathfrak{X}}_{\nu\mu} = \overset{\star^{\mu\nu}}{g} \star (O^{\rho}_{\sigma}[\lambda, -2](\mathbf{\mathfrak{X}}^{\sigma}_{\rho\nu\mu})).$$
(105)

Following the same reasoning as in Sec.III, the *metric is treated as a generic dynamical variable, not as a *-quantity with the need to be expanded in the non-commutativity parameter.

Finally, we are now able to write down the *-Einstein Tensor (55) and the resulting Einstein equation in vacuum for a non-commutative spacetime, locally described by the κ -Minkowski spacetime. However, because of the perturbation nature of the twist (78) controlling the *-geometrical quantities, we write the deformed Einstein equation as a perturbation expansion in the deformation

parameter⁴ λ :

$$0 = \mathfrak{X}_{\mu\nu}[\mathbf{g}, \lambda^{1}, \dots, \lambda^{n}, \dots]$$
(107)
$$-\frac{1}{2} \bar{\phi}^{\{\alpha\}}(\mathbf{g}_{\mu\nu}) \bar{\phi}_{\{\alpha\}}(\mathfrak{X}[\mathbf{g}, \lambda^{1}, \dots, \lambda^{n}, \dots]).$$

The brackets in $\mathfrak{X}_{\mu\nu}[\ldots], \mathfrak{X}[\ldots]$ denote some functional dependence in the deformation parameter and the metric. All the terms are uniquely determined and can be expanded to a chosen order in the non-commutativity parameter $\frac{1}{\kappa} = \lambda$. As an example, to first order, one derives an expansion of the form,

$$0 = \mathfrak{X}_{\mu\nu}^{(0)} - \frac{1}{2} \mathbf{g}_{\mu\nu} \mathfrak{X}^{(0)}$$
(108)
+ $[\mathfrak{X}_{\mu\nu}^{(1)} - \frac{1}{2} \bar{\phi}^{\alpha_1}(\mathbf{g}_{\mu\nu}) \bar{\phi}_{\alpha_1}(\mathfrak{X}^{(0)}) - \frac{1}{2} \mathbf{g}_{\mu\nu} \mathfrak{X}^{(1)}] + \mathcal{O}(\lambda^2).$

Denoting the zeroth and first orders of the *-Ricci tensor and scalars as $\mathbf{\mathfrak{X}}_{\mu\nu}^{(0)}, \mathbf{\mathfrak{X}}_{\mu\nu}^{(1)}$ and $\mathbf{\mathfrak{X}}^{(0)}, \mathbf{\mathfrak{X}}^{(1)}$ respectively. Note, the zeroth order is equivalent to the classical expression, and thus (108) exhibits a meaningful classical limit. Equation (108) can theoretically be solved, providing valuable constraints on the perturbation (non-commutativity) parameter. These constraints can serve as a learning tool for various theories involving the κ -Minkowski spacetime (75) (see Sec.I). Note that as in the constant non-commutativity scenario, the Ricci tensor $\mathbf{\mathfrak{X}}_{\mu\nu}$ is not real/hermitian. One must use a symmetrisized version of $\mathbf{\mathfrak{X}}_{\mu\nu}$ before solving, see [42, 47] and references therein.

The ***-Einstein-Hilbert-Action** can also be constructed, for completeness of the discussion, and as an alternative route to the deformed Einstein equation (107). To achieve this, it is crucial to rigorously define the integral over a given *m-form*, with particular emphasis on (graded-)cyclicity. Here, we encounter a distinct advantage of employing a twist of the Abelian kind (56): the integral over a general *-m-form $w \wedge_* w' :=$ $\bar{\phi}^{\{\alpha\}}(w) \wedge \bar{\phi}_{\{\alpha\}}(w')$ is inherently cyclic (see [45, 62]). Furthermore, the "*" in \wedge_* can be omitted during integration since the deformation terms contribute only as total derivatives. Consequently, we can express the Einstein-Hilbert action, which remains invariant under

$$0 = \overset{*}{g}^{\mu\nu} \star \mathfrak{R}_{\mu\nu} - \frac{1}{2} \overset{*}{g}^{\mu\nu} \star g_{\mu\nu} \star (\overset{*}{g}^{\rho\sigma} \star \mathfrak{R}_{\rho\sigma})$$
$$= \overset{*}{g}^{\mu\nu} \star (\mathfrak{R}_{\mu\nu} - \frac{1}{2} \times 4\mathfrak{R}_{\mu\nu}) \qquad (106)$$
$$\Rightarrow \mathfrak{R}_{\mu\nu} = 0.$$

⁴ For empty space (vacuum), equation (55) reduces, as in the classical scenario, to $0 = \Re_{\mu\nu} - \frac{1}{2}g_{\mu\nu} \star \Re$, which implies

deformed infinitesimal diffeomorphisms, as follows:

$$\overset{\star}{\mathbf{S}_{\mathrm{EH}}} = \int \sqrt{-\det[g]} \star \mathbf{\mathfrak{X}} + c.c. = \int \sqrt{-\det[g]} \mathbf{\mathfrak{X}} + c.c.,$$
(109)

where we have utilized the \star -determinant, defined for a general $n \times n$ matrix **A** as

$$\det^{\star}[\mathbf{A}] = \frac{\epsilon^{\mu_1 \dots \mu_n \nu_1 \dots \nu_n}}{n!} \mathbf{A}_{\mu_1 \nu_1} \star \mathbf{A}_{\mu_2 \nu_2} \dots \star \mathbf{A}_{\mu_n \nu_n},$$
(110)

with $(\mu, \nu) \in [0, n]$.

VI. SUMMARY & FUTURE WORK

In this work, we have constructed a gravity theory as a deformed General Relativity theory on a noncommutative spacetime that is locally the κ -Minkowski spacetime.

As discussed in Sec.I, the physical motivation for this endeavor is to develop a theory for the dynamical sector in one of the active approaches in quantum gravity phenomenology: the DSR approach with a noncommutative spacetime characterized by the Planck length scale.

The usual constructions to date (see [43, 44] and Sec.III) of GR on non-commutative spacetime were not suitable for our case because the local non-commutative spacetime we here considered is with non-constant noncommutativity; the non-commutativity associated with theories of minimal length. In contrast, the previous constructions [42–44, 47] were applied to constant noncommutativity and special cases of spacetime symmetries, enabling significant simplifications.

Therefore, we turned to the generalized procedure of geometry on any non-commutative spacetime. We then validated our method by reconstructing the deformed GR for a local constant non-commutative scenario, finding agreement with [43, 44]. Table I summarizes the mathematical quantities, operators, and algebras used, comparing between commutative spacetime (GR), a spacetime of constant non-commutativity, and a κ -non-commutative spacetime.

To ensure that κ -Minkowski spacetime, treated as an algebra of non-commutative functions, is invariant under transformations of a twisted symmetry structure (a module of a quantum group), we first had to enlarge the classical Poincaré symmetry underlying special relativity with a Dilatation generator. As a result, before twisting, the IGL(3,1) Lie group describes the local symmetry of spacetime in contrast with the classical Poincare symmetry group.

By these means, we successfully derived the deformed version of the Einstein equation (107), i.e., an equation that results from the action (109) under variation of the deformed infinitesimal diffeomorphisms.

Moreover, our use of the twisted symmetry scheme implies that while the spacetime is non-commutative, there are no deformations in the algebraic sector of the symmetry's algebra (keeping in mind that one must use (96) for symmetry transformations). This property implies that the energy dispersion relation and other physical quantities (such as bounded particle momenta, see Sec.I) behave classically in the present scheme. By classically, we mean up to the effects originating from the Dilatation enlargement, which we discuss later.

Now, while the spacetime's quantum nature persists in the non-commutativity relations, the traditional interpretation of a DSR theory becomes challenging here. Boosts act classically on momenta, abandoning any classical notion of a fundamental minimal length. Moreover, a significant ambiguity arises when considering the classical limit ($\kappa \to \infty$) of the κ -GR. In this limit, we recover the classical Einstein equation, but with the underlying symmetry of IGL(3,1) instead of the proper Poincaré symmetry of SR, raising concerns on the correspondence principle of the κ -GR. These ambiguities will be addressed in our upcoming paper [63].

In our upcoming paper [63], we present a general procedure to compare predictions from the κ -GR theory to the observations of astrophysical systems that involve highly energetic gravitational events—such as a binary black hole coalescence, among other possibilities. In particular, we discuss the distinction between gravity generation (such as a dynamic deformed metric) and gravity propagation (fuzziness of spacetime originated from (75)) deformations.

ACKNOWLEDGEMENTS

We acknowledge support from the US-Israel Binational Science Fund (BSF) grant No. 2020245 and the Israel Science Fund (ISF) grant No. 1698/22.

Appendix A: Detailed Calculations for Sec.III

Here, we shall detail the calculations beyond the results stated in Sec.III.

The permutation operator (59):

$$\mathcal{R} = \Phi_{21}\Phi^{-1} = (\phi_{\{\alpha\}} \otimes \phi^{\{\alpha\}})(\bar{\phi}^{\{\beta\}} \otimes \bar{\phi}_{\{\beta\}})$$
(A1)
$$= (e^{-i\frac{\lambda}{2}\theta^{\nu\mu}\partial_{\nu}\otimes\partial_{\mu}})(e^{+i\frac{\lambda}{2}\theta^{\mu\nu}\partial_{\mu}\otimes\partial_{\nu}}) = e^{+i\lambda\theta^{\mu\nu}\partial_{\mu}\otimes\partial_{\nu}}.$$

The \star -Lie derivative (66):

Let us first calculate the twist action on a general vector field $\xi \in \Xi$,

$$\begin{split} \bar{\phi}^{\{\alpha\}}(\xi)\bar{\phi}_{\{\alpha\}} &= \mathcal{L}_{\bar{\phi}^{\{\alpha\}}}(\xi)(\mathcal{L}_{\bar{\phi}_{\{\alpha\}}}) \\ &= [\bar{\phi}^{\{\alpha\}},\xi](\mathcal{L}_{\bar{\phi}_{\{\alpha\}}}) = [\bar{\phi}^{\{\alpha\}\mu}\partial_{\mu},\xi^{\nu}\partial_{\nu}] \\ &= (\bar{\phi}^{\{\alpha\}\mu}\partial_{\mu}(\xi^{\nu}) - \xi^{\mu}\partial_{\mu}(\bar{\phi}^{\{\alpha\}\nu}))\partial_{\nu}(\bar{\phi}_{\{\alpha\}}) \\ &= \bar{\phi}^{\{\alpha\}}(\xi^{\nu})\partial_{\nu}(\bar{\phi}_{\{\alpha\}}), \end{split}$$
(A2)

we used the fact that for constant non-commutativity, the $\bar{\phi}^{\{\alpha\}}$ depends on derivatives ∂_{μ} and some constants we denoted by $\bar{\phi}^{\{\alpha\}\mu}$, such that $\bar{\phi}^{\{\alpha\}} = \bar{\phi}^{\{\alpha\}\mu}\partial_{\mu}$. The *-Lie derivative is, therefore,

$$\mathcal{L}_{\xi}^{*} = \mathcal{L}_{\bar{\phi}^{\{\alpha\}}(\xi)\bar{\phi}_{\{\alpha\}}} = \bar{\phi}^{\{\alpha\}}(\xi^{\nu})\partial_{\nu}(\bar{\phi}_{\{\alpha\}})$$

$$= \mu \circ \left(\left\{\exp\left[\frac{i\lambda}{2}\theta^{\mu\nu}\partial_{\mu}\otimes\partial_{\nu}\right]\circ(\xi^{\gamma}\otimes1)\right\}(\partial_{\gamma}\otimes1)\right)\right\}$$

$$= \sum_{n=0}^{\infty}\frac{1}{n!}\left(\frac{i\lambda}{2}\right)^{n}\theta^{\mu_{1}\nu_{1}}...\theta^{\mu_{n}\nu_{n}}\partial_{\mu_{1}}...\partial_{\mu_{n}}(\xi^{\gamma})\partial_{\gamma}\partial_{\nu_{1}}...\partial_{\nu_{n}}(h)$$

$$= \xi + i\lambda/2\theta^{\mu\nu}\partial_{\mu}(\xi^{\gamma})\partial_{\gamma}\partial_{\nu} + \mathcal{O}(\lambda^{2}\partial^{2}(\xi^{\gamma})). \quad (A3)$$

The \star -covariant derivative (67):

$$\begin{split} \bar{\nabla}_{z} u &= \bar{\nabla}_{z} (u^{\nu} \partial_{\nu}) \\ &= \mathcal{L}_{z}^{*} (u^{\nu}) \star \partial_{\nu} + \bar{\mathrm{R}}^{\{\alpha\}} (u^{\nu}) \star \bar{\nabla}_{\bar{\mathrm{R}}_{\{\alpha\}} (z^{\mu} \partial_{\mu})} \star \partial_{\nu} \\ &= \mathcal{L}_{z}^{*} (u^{\nu}) \partial_{\nu} + (\bar{\mathrm{R}}^{\{\alpha\}} (u^{\nu}) \star \bar{\mathrm{R}}_{\{\alpha\}} (z^{\mu})) \star (\Gamma_{\mu\nu}^{\sigma} \partial_{\sigma}) \\ &= [\mathcal{L}_{z}^{*} (u^{\nu}) + (z^{\mu} \star u^{\sigma}) \star \Gamma_{\mu\sigma}^{\nu}] \partial_{\nu}. \end{split}$$
(A4)

For the case when $z^{\mu} = 1$, using only the properties of the *-Lie derivative (65) suffices to get (67).

The *-curvature tensor:

By use of the identities in (68), formula (49) can be simplified to

$$\mathfrak{X}(\partial_{\mu},\partial_{\nu},\partial_{\rho}) = \overset{\star}{\nabla}_{\partial_{\mu}}(\overset{\star}{\nabla}_{\partial_{\nu}}\partial_{\rho}) - \overset{\star}{\nabla}_{\bar{\mathrm{R}}^{\{\alpha\}}(\partial_{\nu})}(\overset{\star}{\nabla}_{\bar{\mathrm{R}}_{\{\alpha\}}(\partial_{\mu})}\partial_{\rho}).$$
(A5)

We calculate the first term,

$$\begin{aligned} \stackrel{\star}{\nabla}_{\partial_{\mu}} (\stackrel{\star}{\nabla}_{\partial_{\nu}} \partial_{\rho}) &= \stackrel{\star}{\nabla}_{\partial_{\mu}} (\Gamma^{\sigma}_{\nu\rho} \star \partial_{\sigma}) \\ &= \mathcal{L}^{\star}_{\partial_{\mu}} (\Gamma^{\sigma}_{\nu\rho}) \star \partial_{\sigma} + \bar{\mathrm{R}}^{\{\alpha\}} (\Gamma^{\sigma}_{\nu\rho}) \star \stackrel{\star}{\nabla}_{\bar{\mathrm{R}}_{\{\alpha\}}(\partial_{\mu})} (\partial_{\sigma}) \\ &= \mathcal{L}^{\star}_{\partial_{\mu}} (\Gamma^{\sigma}_{\nu\rho}) \star \partial_{\sigma} + \bar{\mathrm{R}}^{\{\alpha\}} (\Gamma^{\sigma}_{\nu\rho}) \star \Gamma^{\tau}_{\mu\sigma} \star \partial_{\tau} \\ &= \boxed{\mathcal{L}_{\partial_{\mu}} (\Gamma^{\sigma}_{\nu\rho}) \partial_{\sigma} + (\Gamma^{\sigma}_{\nu\rho} \star \Gamma^{\tau}_{\mu\sigma}) \partial_{\tau}}, \quad (A6) \end{aligned}$$

and then the second term,

$$\overset{*}{\nabla_{\bar{\mathrm{R}}}}_{(\partial_{\nu})} (\overset{*}{\nabla_{\bar{\mathrm{R}}}}_{(\alpha)}(\partial_{\mu}) \partial_{\rho}) = \overset{*}{\nabla_{\bar{\mathrm{R}}}}_{(\alpha)}(\Gamma_{\mu\rho}^{\sigma} \star \partial_{\sigma})$$
(A7)
$$= \mathcal{L}_{\partial_{\nu}}(\Gamma_{\mu\rho}^{\sigma})\partial_{\sigma} + (\Gamma_{\mu\rho}^{\sigma} \star \Gamma_{\nu\sigma}^{\tau})\partial_{\tau}.$$

Subtracting, we get the *-curvature tensor,

$$\begin{aligned} \mathbf{\mathfrak{X}}(\partial_{\mu},\partial_{\nu},\partial_{\rho}) &= \mathcal{L}_{\partial_{\mu}}(\Gamma^{\sigma}_{\nu\rho})\partial_{\sigma} - \mathcal{L}_{\partial_{\nu}}(\Gamma^{\sigma}_{\mu\rho})\partial_{\sigma} + \Gamma^{\sigma}_{\rho[\nu} \star \Gamma^{\tau}_{\mu]\sigma}\partial_{\tau} \\ &= (\partial_{[\mu}\Gamma^{\sigma}_{\nu]\rho} + \Gamma^{\tau}_{\rho[\nu} \star \Gamma^{\sigma}_{\mu]\tau})\partial_{\sigma} \\ \Rightarrow \mathbf{\mathfrak{X}}^{\sigma}_{\mu\nu\rho} &= \overline{\partial_{[\mu}\Gamma^{\sigma}_{\nu]\rho} + \Gamma^{\tau}_{\rho[\nu} \star \Gamma^{\sigma}_{\mu]\tau}} . \end{aligned}$$
(A8)

The \star -Ricci tensor (70):

$$\begin{aligned} \mathbf{\mathfrak{X}}_{\mu\nu} &= \langle dx^{\mu} \mathbf{\mathfrak{X}} (\partial_{\mu}, \partial_{\nu}, \partial_{\rho}) \rangle' = \langle dx^{\mu} \mathbf{\mathfrak{X}} (\partial_{\mu} \Gamma^{\sigma}_{\nu\rho} + ...) \partial_{\sigma} \rangle' \\ &= \langle dx^{\mu} \mathbf{\mathfrak{X}} \partial_{\sigma} (\partial_{\mu} \Gamma^{\sigma}_{\nu\rho} + ...) \rangle' = \langle dx^{\mu} \mathbf{\mathfrak{X}} \partial_{\sigma} \rangle' \cdot (\partial_{\mu} \Gamma^{\sigma}_{\nu\rho} + ...) \\ &= \delta^{\mu}_{\sigma} \mathbf{\mathfrak{X}}^{\sigma}_{\mu\nu\rho} \Rightarrow \mathbf{\mathfrak{X}}_{\mu\nu\rho} = \mathbf{\mathfrak{X}}^{\mu}_{\mu\nu\rho} \end{aligned} .$$
(A9)

The relation between the metric tensor and the Cristoffel symbols:

The \star -covariant derivative of the metric (71) is calculated as,

$$\overset{\star}{\nabla}_{\gamma}(\mathbf{g}) = \overset{\star}{\nabla}_{\gamma}(dx^{\mu} \overset{\star}{\otimes} dx^{\nu} \star g_{\mu\nu})$$

$$= [(\overset{\star}{\nabla}_{\gamma}(dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) \star g_{\mu\nu})$$

$$+ (\bar{\mathbf{R}}^{\{\alpha\}}(dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) \star \overset{\star}{\nabla}_{\bar{\mathbf{R}}_{\{\alpha\}}(\gamma)}(g_{\mu\nu}))]$$

$$:= [(\mathbf{A}) + (\mathbf{B})], \qquad (A10)$$

with

$$\mathbf{A} = \left[\stackrel{\star}{\nabla}_{\gamma} (dx^{\mu}) \stackrel{\star}{\otimes} dx^{\nu} + \bar{\mathbf{R}}^{\{\alpha\}} (dx^{\mu}) \star \bar{\mathbf{R}}^{\mu'}_{\{\alpha\}\gamma} \stackrel{\star}{\otimes} \stackrel{\star}{\nabla}_{\mu'} (dx^{\nu}) \right] \star g_{\mu\nu}$$
$$= \left[\stackrel{\star}{\nabla}_{\gamma} (dx^{\mu}) \stackrel{\star}{\otimes} dx^{\nu} + dx^{\mu} \star \delta^{\mu'}_{\gamma} \stackrel{\star}{\otimes} \stackrel{\star}{\nabla}_{\mu'} dx^{\nu} \right] \star g_{\mu\nu}$$
$$= -\left[\Gamma^{\mu}_{\gamma\sigma} \star (dx^{\sigma} \stackrel{\star}{\otimes} dx^{\nu}) + \Gamma^{\nu}_{\gamma\sigma} \star (dx^{\mu} \stackrel{\star}{\otimes} dx^{\sigma}) \right] \star g_{\mu\nu}$$
$$= \left[-(\Gamma^{\sigma}_{\gamma(\mu} \star g_{\nu)\sigma}) (dx^{\mu} \stackrel{\star}{\otimes} dx^{\nu}) \right], \qquad (A11)$$

where in the first line we used the \star -Leibniz rule (47), and with

$$\mathbf{B} = \bar{\mathbf{R}}^{\{\alpha\}} (dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) \star \bar{\mathbf{R}}^{\mu'}_{\{\alpha\} \gamma} \star \overset{\star}{\nabla}^{\star}_{\mu'}(g_{\mu\nu})$$
(A12)
$$= (dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) \star \mathcal{L}^{\star}_{\gamma}(g_{\mu\nu}) = \boxed{g_{\mu\nu,\gamma}(dx^{\mu} \overset{\star}{\otimes} dx^{\nu})}.$$

The resultant metric compatibility condition is then,

$$\stackrel{\star}{\nabla}_{\gamma}(\mathbf{g}) = [g_{\mu\nu,\gamma} - \Gamma^{\sigma}_{\gamma(\mu} \star g_{\nu)\sigma}](dx^{\mu} \overset{\star}{\otimes} dx^{\nu})$$
$$\Rightarrow \stackrel{\star}{\nabla}_{\gamma}(\mathbf{g}) = 0 \Leftrightarrow g_{\mu\nu,\gamma} = \Gamma^{\sigma}_{\gamma(\mu} \star g_{\nu)\sigma}.$$
(A13)

By combining cyclic permutations of (72), we obtain the relation between the Cristoffel symbols and the metric: first, define,

$$\Gamma^{\sigma}_{\gamma\mu} \star g_{\nu\sigma} = \Gamma^{\sigma}_{\gamma\mu} \star g_{\sigma\nu} \coloneqq \Gamma_{\nu\gamma\mu}, \qquad (A14)$$

$$\Gamma^{\sigma}_{\gamma\nu} \star g_{\mu\sigma} = \Gamma^{\sigma}_{\gamma\nu} \star g_{\sigma\mu} \coloneqq \Gamma_{\mu\gamma\nu}. \tag{A15}$$

Then, by using $g_{\mu\nu,\gamma} = \Gamma_{\nu\gamma\mu} + \Gamma_{\mu\gamma\nu} = \Gamma_{(\nu\mu)\gamma}$, and

$$\frac{1}{2}(g_{\mu(\nu,\gamma)} - g_{\nu\gamma,\mu}) = \frac{1}{2} \left[\Gamma_{(\mu\nu)\gamma} + \Gamma_{(\mu\gamma)\nu} - \Gamma_{(\nu\gamma)\mu} \right]$$
(A16)
$$= 2 \times \frac{1}{2} \Gamma_{\mu\nu\gamma} + \Gamma_{\nu[\mu\gamma]} + \Gamma_{\gamma[\mu\nu]} = \Gamma_{\mu\nu\gamma},$$

we derive,

$$\Gamma^{\sigma}_{\nu\gamma} = g^{\star\sigma\mu} \star \Gamma_{\mu\nu\gamma} = g^{\star\sigma\mu} \star \frac{1}{2} [g_{\mu(\nu,\gamma)} - g_{\nu\gamma,\mu}]. (A17)$$

We used that the metric tensor is symmetric and that the lowered Cristoffell symbol $\Gamma_{\mu\nu\gamma}$ is symmetric in its *last* two indices.

Appendix B: Detailed Calculations for Sec.IV

Here, we shall detail the calculations beyond the results stated in Sec.IV.

The action on basis 1-forms (83):

Using the definition of a Lie derivative of a one-form along a vector field, we derive

$$\begin{aligned} X_0(dx^{\mu}) &= \mathcal{L}_{X_0}(dx^{\mu}) = [X_0^{\mu}\partial_{\mu}(1) + \partial_{\nu}(X_0^{\mu})1] dx^{\nu} \\ &= [0 + \partial_{\nu}(1)] dx^{\nu} = 0, \end{aligned} \tag{B1} \\ X_1(dx^{\mu}) &= \mathcal{L}_{x^j\partial_j}(dx^{\mu}) = (\partial_{\mu}x^j) dx^{\mu} = \delta_{\mu}^j dx^{\mu} \neq 0. \end{aligned}$$

And, for the basis vectors (84), we use the Lie bracket,

$$X_0(\partial_\mu) = [1\partial_\mu(1) - 1\partial_\mu(1)]\partial_\nu = 0,$$

$$X_1(\partial_\mu) = -\partial_\mu(x^j)\partial_j = -\delta^j_\mu\partial_j.$$
 (B2)

The \star -pairing of the classical basis vectors and 1-forms (87):

Explicitly calculating, we get,

$$\begin{aligned} \langle \partial_0^* dx^{\nu} \rangle &= \langle \bar{\phi}^{\{\alpha\}}(\partial_0), \bar{\phi}_{\{\alpha\}}(dx^{\nu}) \rangle \\ &= \langle O_{\nu'}^{\nu}[\lambda, 1] \partial_0, dx^{\nu'} \rangle \\ &= \delta_0^{\nu}, \\ \langle \partial_j^* dx^{\nu} \rangle &= \langle \bar{\phi}^{\{\alpha\}}(\partial_j), \bar{\phi}_{\{\alpha\}}(dx^{\nu}) \rangle \\ &= \langle e^{i\frac{\lambda}{2}\partial_0} \delta_{j'}^{\nu} \partial_j, dx^{j'} \rangle \\ &= e^{i\frac{\lambda}{2}\partial_0} \delta_{j'}^{\nu} \langle \partial_j, dx^{j'} \rangle = e^{i\frac{\lambda}{2}\partial_0} \delta_j^{j'} \delta_{j'}^{\nu}, \\ &\Rightarrow \langle \partial_\mu^* dx^{\nu} \rangle = \boxed{\delta_\mu^0 \delta_0^{\nu} + \delta_\mu^j \delta_j^{\nu} e^{i\frac{\lambda}{2}\partial_0}} . \end{aligned}$$
(B3)

The non-commutativity of \star -1-forms: Since the relevant (non-zero) part of \mathcal{R} here is

$$\bar{\mathcal{R}}_{\text{relevant}} = \exp\left[-i\lambda(-X_1 \otimes X_0)\right], \qquad (B4)$$

we use (86) and (89) to get

$$\bar{\mathbf{R}}^{\{\alpha\}}(dx^{\mu}) \star \bar{\mathbf{R}}_{\{\alpha\}}(f) = dx^{\mu'} \star (e^{+i\lambda\partial_0}\delta^j_{\mu'}\delta^{\mu}_j + \delta^0_{\mu'}\delta^{\mu}_0)(f)$$
$$\coloneqq dx^{\mu'} \star O^{\mu}_{\mu'}[\lambda, 2](f) \quad . \tag{B5}$$

The known calculation of the deformed Leibniz rule of *-basis-vectors (see [38]):

Calculating the left-hand side in the second line of (91),

$$\mathbf{d}(f \star g) = (\overset{\star}{\partial}_{\mu}f) \star dx^{\mu} \star g + f \star (\overset{\star}{\partial}_{\mu}g) \star dx^{\mu}$$
(B6)
$$= ((\overset{\star}{\partial}_{\mu}f) \star O^{\mu}_{\mu'}[\lambda, -2](g)) \star dx^{\mu'} + f \star (\overset{\star}{\partial}_{\mu}g) \star dx^{\mu},$$

and equating with the right-hand side, we derive two equations,

$$(\partial_j^{\star}(f \star g)) \star dx^j = ((\partial_j^{\star}f) \star e^{-i\lambda\partial_0}g + f \star (\partial_j^{\star}g)) \star dx^j, (\partial_0^{\star}(f \star g)) \star dx^0 = f \star (\partial_0^{\star}g) + (\partial_0^{\star}f) \star g.$$
 (B7)

concluding the deformed Leibniz rule for *-derivatives,

$$\partial_{j}^{\star}(f \star g) = (\partial_{j}^{\star}f) \star e^{-i\lambda\partial_{0}}g + f \star \partial_{j}^{\star}g,$$

$$\partial_{0}^{\star}(f \star g) = \overset{\star}{\partial_{0}}f \star g + f \star \partial_{0}^{\star}g.$$
 (B8)

We have used,

$$\bar{\mathbf{R}}^{\{\alpha\}}(h) \star \bar{\mathbf{R}}_{\{\alpha\}}(dx^{\mu}) = O^{\mu}_{\mu'}[\lambda, -2](h) \star dx^{\mu'}.$$
 (B9)

The new calculation of the Leibniz rule of *-basis-vector:

First, we evaluate the twisted co-product of $\hat{\partial}_{\mu}$:

$$\Delta_{\Phi}(\overset{\star}{\partial}_{\mu}) = \Phi \circ (\overset{\star}{\partial}_{\mu} \otimes 1 + 1 \otimes \overset{\star}{\partial}_{\mu}) \circ \Phi^{-1}$$

$$= \phi^{\{\alpha\}}(\overset{\star}{\partial}_{\mu})\bar{\phi}^{\{\beta\}} \otimes \phi_{\{\alpha\}}\bar{\phi}_{\{\beta\}}$$

$$+\phi^{\{\alpha\}}\bar{\phi}^{\{\beta\}} \otimes \phi_{\{\alpha\}}(\overset{\star}{\partial}_{\mu})\bar{\phi}_{\{\beta\}} \qquad (B10)$$

$$= \partial_{\mu'}^{\star} \otimes O_{\mu}^{\mu'}[\lambda, -1] + O_{\mu}^{\mu'}[\lambda, 1] \otimes \partial_{\mu'}^{\star},$$

so that the \star -co-product is then,

$$\Delta_{\star} \begin{pmatrix} \star \\ \partial_{\mu} \end{pmatrix} = (\phi^{\{\alpha\}} \phi_{\{\alpha\}} \otimes \phi^{\{\alpha'\}} \phi_{\{\alpha'\}}) \circ \left[\partial_{\mu'}^{\star} \otimes O_{\mu}^{\mu'} [\lambda, -1] + O_{\mu}^{\mu'} [\lambda, 1] \otimes \partial_{\mu'}^{\star} \right] \circ \bar{\phi}^{\{\beta\}} \bar{\phi}_{\{\beta\}} \\ = \left[\phi^{\{\alpha\}} \phi_{\{\alpha\}} (\partial_{\mu'}^{\star}) \otimes \phi^{\{\alpha'\}} \phi_{\{\alpha'\}} (O_{\mu}^{\mu'} [\lambda, -1]) + \phi^{\{\alpha\}} \phi_{\{\alpha\}} (O_{\mu}^{\mu'} [\lambda, 1]) \otimes \phi^{\{\alpha'\}} \phi_{\{\alpha'\}} (\partial_{\mu'}^{\star}) \right] \circ \bar{\phi}^{\{\beta\}} \bar{\phi}_{\{\beta\}} \\ = \left[\phi_{\{\alpha\}} \partial_{\mu''}^{\star} \otimes \phi^{\{\alpha'\}} O_{\mu}^{\mu''} [\lambda, -2] + \phi_{\{\alpha\}} \otimes \phi^{\{\alpha'\}} \partial_{\mu'}^{\star} \right] \circ \bar{\phi}^{\{\beta\}} \bar{\phi}_{\{\beta\}} = \partial_{\mu'}^{\star} \otimes O_{\mu}^{\mu'} [\lambda, -2] + 1 \otimes \partial_{\mu}^{\star}, \tag{B11}$$

This leads to the deformed Leibniz rule,

$$\hat{\partial}_{\mu}(f \star g) = \mu_{\star} \circ \{\Delta_{\star}(\hat{\partial}_{\mu})(f \otimes g)\}$$
(B12)
$$= \overset{\star}{\partial}_{\mu'}(f) \star O^{\mu'}_{\mu}[\lambda, -2](g) + f \star (\overset{\star}{\partial}_{\mu}g).$$

Appendix C: Detailed Calculations for Sec.V

Here, we shall detail the calculations beyond the results stated in Sec.V.

The ***-Lie derivative:**

For a general vector field $\xi \in \Xi_{\star}$ we have

$$\mathcal{L}_{\xi}^{\star} = \mathcal{L}_{\bar{\phi}^{\{\alpha\}}(\xi)\bar{\phi}_{\{\alpha\}}} \equiv \bar{\phi}^{\{\alpha\}}(\xi)\bar{\phi}_{\{\alpha\}}$$
(C1)
$$= \mu \circ \left(\exp\left[\frac{i\lambda}{2}\left(\partial_{0}\otimes x^{j}\partial_{j} - x^{j}\partial_{j}\otimes\partial_{0}\right)\right]\left(\xi^{\mu}\overset{\star}{\partial}_{\mu}\otimes1\right)\right)$$
$$= \mu \circ \left[\exp\left(\frac{i\lambda}{2}\partial_{0}\otimes x^{j}\partial_{j}\right)\exp\left(-\frac{i\lambda}{2}x^{j}\partial_{j}\otimes\partial_{0}\right)\left(\xi^{\mu}\overset{\star}{\partial}_{\mu}\otimes1\right)\right]$$
$$= \sum_{\substack{n=0\\n'=0}}^{\infty} \frac{(i\lambda/2)^{n+n'}}{n!n'!} \left(-x^{j}\partial_{j}\right)^{n'} \left[\partial_{0}^{n}\left(\xi^{\mu}\right)\overset{\star}{\partial}_{\mu}\right]\partial_{0}^{n'}\left(x^{j}\partial_{j}\right)^{n}.$$

We used that $[x^j \partial_j, \partial_0] = 0$ and that $\partial_0(\partial_\mu) = 0$. To get a taste for the first few orders, first note that

$$\partial_0(\xi) = \mathcal{L}_{\partial_0}(\xi) = (\partial_0(\xi^{\nu}) - \xi^{\mu}\partial_{\mu}(1))\partial_{\nu} = \partial_0(\xi^{\mu})\partial_{\mu},$$

$$x^j\partial_j(\xi) = [x^j\partial_j,\xi] = (x^j\partial_j(\xi^{\mu}) - \xi^{\mu}\delta^{\mu}_j)\partial_{\mu}.(C2)$$

Then, up to the second order, we can write (C1) explicitly;

$$\mathcal{L}_{\xi}^{\star} = \xi^{\mu} \overset{\star}{\partial}_{\mu} + \frac{i\lambda}{2} \Big[x^{j} \left(\partial_{0}(\xi^{\mu}) \partial_{j} - \partial_{j}(\xi^{\mu}) \partial_{0} \right) + \xi^{\mu} \delta^{\mu}_{j} \Big] \overset{\star}{\partial}_{\mu} \partial_{0} \\ + \left(\frac{i\lambda}{2} \right)^{2} x^{j} \Big[\left(\partial_{0}(\xi^{\mu}) - x^{j} \partial_{j} \partial_{0}(\xi^{\mu}) \right) \partial_{j} - \partial_{j}(\xi^{\mu}) \partial_{0} \delta^{\mu}_{j} \\ + 1/2 \left(\partial_{j}(\xi^{\mu}) + x^{j} \partial^{2}_{j}(\xi^{\mu}) \right) \partial_{0} \Big] \overset{\star}{\partial}_{\mu} \partial_{0} \\ + \frac{1}{2} \left(\frac{i\lambda}{2} \right)^{2} \xi^{\mu} \delta^{j}_{\mu} \delta^{\mu'}_{j} \overset{\star}{\partial}_{\mu'} \partial^{2}_{0}.$$
(C3)

Note that if ξ^{μ} is constant, we can calculate the entire sum,

$$\mathcal{L}_{\xi}^{\star} = \xi + \frac{i\lambda}{2}\xi^{\mu}\partial_{\mu}\delta_{j}^{\mu}\partial_{0} + \frac{1}{2}\left[\left(\frac{i\lambda}{2}\right)^{2}\xi^{\mu}\overset{\star}{\partial}_{\mu'}\right]\delta_{j}^{\mu'}\delta_{\mu}^{j}\partial_{0}^{2} + \cdots$$
$$= e^{+(i\lambda/2)\partial_{0}\delta_{j}^{\mu}}\xi. \tag{C4}$$

The *****-Covariant Derivative:

Directly calculating, we get for the general vector fields case (98),

$$\begin{aligned} \stackrel{\star}{\nabla}_{z} u &= \mathcal{L}_{z^{\star} \mu \star}^{\star} \binom{\star^{\nu}}{2} * \stackrel{\star}{\partial}_{\mu} (u^{\star}) * \stackrel{\star}{\partial}_{\nu} + \bar{\mathrm{R}}^{\{\alpha\}} \binom{\star^{\nu}}{u} * \stackrel{\star}{\nabla}_{\bar{\mathrm{R}}_{\{\alpha\}} (z^{\star} \star \stackrel{\star}{\partial}_{\mu})} (\stackrel{\star}{\partial}_{\nu}) \\ &= \left[\mathcal{L}_{z}^{\star} (u^{\star}) + \bar{\mathrm{R}}^{\{\alpha\}} (u^{\star^{\sigma}}) * \bar{\mathrm{R}}_{\{\alpha\}\mu}^{\mu'}(z) * \Gamma_{\mu'\sigma}^{\nu} \right] * \stackrel{\star}{\partial}_{\mu}^{\lambda} (\mathrm{C5}) \end{aligned}$$

This is the farthest way we can simplify the expression. From here one need to use the *-Lie derivative and the composite action of $\bar{\mathbb{R}}^{\{\alpha\}}$ on z. If $z = \partial_{\mu}$ we can get

$$\stackrel{\star}{\nabla}_{\overset{*}{\partial}_{\mu}}(u) = \left[\partial_{\mu}(\overset{\star}{u}^{\nu}) + O^{\mu'}_{\mu}[\lambda, 2](\overset{\star}{u}^{\sigma}) \star \Gamma^{\nu}_{\mu'\sigma}\right] \star \overset{\star}{\partial}_{\nu}$$
$$= \left[\partial_{\mu}(u^{\nu}) + O^{\mu'}_{\mu}[\lambda, 2](u^{\sigma}) \star \Gamma^{\nu}_{\mu'\sigma}\right] \overset{\star}{\partial}_{\nu}.$$
(C6)

Moreover, for the case when we are only interested in the $u = u^{\nu}$ part, it is immediate to derive (99) from (C5).

The *-curvature tensor:

The result (100) is derived through the following calculation,

$$\begin{aligned} \mathbf{1} \mathbf{\hat{x}} \begin{pmatrix} \star & \star & \star & \star \\ \partial_{\mu}, \partial_{\nu}, \partial_{\rho} \end{pmatrix} &= \nabla_{\lambda \mu}^{\star} (\nabla_{\lambda \nu}^{\star} \partial_{\rho}) - \nabla_{[\partial_{\mu}^{\star}; \partial_{\nu}]}^{\star} \begin{pmatrix} \star & \star & \star \\ \partial_{\rho} \end{pmatrix} \\ &- \nabla_{\bar{\mathbf{R}}^{\{\alpha\}}} (\nabla_{\bar{\mathbf{R}}_{\{\alpha\}}} (\partial_{\mu}) \partial_{\rho}) \\ &= \nabla_{\lambda \mu}^{\star} (\Gamma_{\nu \rho}^{\sigma} \star \partial_{\sigma}) - \bar{\mathbf{R}}^{\{\alpha\}\nu'} \star \nabla_{\lambda \nu'}^{\star} (\bar{\mathbf{R}}_{\{\alpha\}\mu}^{\mu'} \star \nabla_{\lambda \mu'}^{\star} (\partial_{\rho})) \\ &= \nabla_{\lambda \mu}^{\star} (\Gamma_{\nu \rho}^{\sigma} \star \partial_{\sigma}) - \bar{\mathbf{R}}^{\{\alpha\}\nu'} \star \nabla_{\lambda \nu'}^{\star} (\bar{\mathbf{R}}_{\{\alpha\}\mu}^{\mu'} \star \Gamma_{\mu'\rho}^{\sigma} \star \partial_{\sigma}) \\ &= (\mathbf{C}) + (\mathbf{D}) , \end{aligned}$$

where we have used $[\overset{\star}{\partial}_{\mu},\overset{\star}{\partial}_{\nu}] = 0$ to go from the first line to the second. We calculate the first term as follows:

$$\mathbf{C} := \nabla_{\partial_{\mu}}^{\star} (\Gamma_{\nu\rho}^{\sigma} \star \overset{\star}{\partial}_{\sigma})$$

$$= \mathcal{L}_{\partial_{\mu}}^{\star} (\Gamma_{\nu\rho}^{\sigma}) \star \overset{\star}{\partial}_{\sigma} + \bar{\mathbf{R}}^{\{\alpha\}} (\Gamma_{\nu\rho}^{\sigma}) \star \overset{\star}{\nabla}_{\bar{\mathbf{R}}_{\{\alpha\}}(\partial_{\mu}^{\star})} (\overset{\star}{\partial}_{\sigma})$$

$$= [\partial_{\mu} (\Gamma_{\nu\rho}^{\sigma}) + \bar{\mathbf{R}}^{\{\alpha\}} (\Gamma_{\nu\rho}^{\tau}) \star \bar{\mathbf{R}}_{\{\alpha\}\mu}^{\mu'} \star \Gamma_{\mu'\tau}^{\sigma}] \star \overset{\star}{\partial}_{\sigma}, (C8)$$

and the second term as

$$\mathbf{D} \coloneqq \bar{\mathbf{R}}_{\nu}^{\{\alpha\}\nu'} \star \stackrel{\star}{\nabla}_{\partial_{\nu'}}^{\star} (\bar{\mathbf{R}}_{\{\alpha\}}^{\mu'} + \Gamma_{\mu'\rho}^{\sigma} \star \stackrel{\star}{\partial}_{\sigma})$$
(C9)
$$= \bar{\mathbf{R}}_{\nu}^{\{\alpha\}\nu'} \star (\bar{\mathbf{R}}^{\{\beta\}}(\bar{\mathbf{R}}_{\{\alpha\}}^{\mu'} + \mu) \star \stackrel{\star}{\nabla}_{\bar{\mathbf{R}}_{\{\beta\}}(\partial_{\nu)}}^{\star} (\Gamma_{\mu'\rho}^{\sigma} \star \stackrel{\star}{\partial}_{\sigma}))$$
$$= \bar{\mathbf{R}}_{\nu}^{\{\alpha\}\nu'} \star [(\bar{\mathbf{R}}^{\{\beta\}}(\bar{\mathbf{R}}_{\{\alpha\}\mu}^{\mu'}) \star \bar{\mathbf{R}}_{\{\beta\}}^{\nu''} \star \stackrel{\star}{\nabla}_{\partial_{\nu''}}^{\star} (\Gamma_{\mu'\rho}^{\sigma} \star \stackrel{\star}{\partial}_{\sigma}))].$$

We also observe that,

$$\bar{\mathbf{R}}^{\{\beta\}}(\bar{\mathbf{R}}^{\mu'}_{\{\alpha\} \ \mu}) \star \bar{\mathbf{R}}^{\nu''}_{\{\beta\} \ \nu'} = \bar{\mathbf{R}}^{\mu'}_{\{\alpha\} \ \mu} \delta^{\nu''}_{\nu'} , \quad (C10)$$
$$\bar{\mathbf{R}}^{\{\alpha\} \ \nu'}_{\ \nu} \star \bar{\mathbf{R}}^{\mu'}_{\{\alpha\} \ \mu} = \delta^{\mu'}_{\mu} \delta^{\nu'}_{\nu} . \quad (C11)$$

thus, we can simplify (C9):

$$\bar{\mathbf{R}}_{\nu}^{\{\alpha\}\nu'} * \nabla_{\partial_{\nu'}}^{\star} (\bar{\mathbf{R}}_{\{\alpha\}\mu}^{\mu'} * \Gamma_{\mu'\rho}^{\sigma} * \overset{\star}{\partial_{\sigma}}) = \nabla_{\lambda}_{\nu} (\Gamma_{\mu\rho}^{\sigma} * \overset{\star}{\partial_{\sigma}})$$
$$= [\partial_{\nu} (\Gamma_{\mu\rho}^{\sigma}) + \bar{\mathbf{R}}^{\{\alpha\}} (\Gamma_{\mu\rho}^{\tau}) * \bar{\mathbf{R}}_{\{\alpha\}\nu}^{\nu'} * \Gamma_{\nu'\tau}^{\sigma}] * \overset{\star}{\partial_{\sigma}} (C12)$$

Finally, (C7) can be now written directly in its components using (49) and the two terms ((C8), (C9)),

$$\begin{aligned} \mathfrak{X}^{\sigma}_{\mu\nu\rho} &= \left[\partial_{\mu}(\Gamma^{\sigma}_{\nu\rho}) + \bar{\mathrm{R}}^{\{\alpha\}}(\Gamma^{\tau}_{\nu\rho}) \star \bar{\mathrm{R}}^{\mu'}_{\{\alpha\} \ \mu} \star \Gamma^{\sigma}_{\mu'\tau}\right] \\ &- \left[\partial_{\nu}(\Gamma^{\sigma}_{\mu\rho}) + \bar{\mathrm{R}}^{\{\alpha\}}(\Gamma^{\tau}_{\mu\rho}) \star \bar{\mathrm{R}}^{\nu'}_{\{\alpha\} \ \nu} \star \Gamma^{\sigma}_{\nu'\tau}\right] \\ &= \left[\partial_{\left[\mu}\Gamma^{\sigma}_{\nu\right]\rho} + O^{\gamma}_{\left[\mu}[\lambda,2](\Gamma^{\tau}_{\nu\right]\rho}) \star \Gamma^{\sigma}_{\gamma\tau}\right], \end{aligned} (C13)$$

where we have used the by-now familiar result,

$$\bar{\mathrm{R}}^{\{\alpha\}}(\Gamma^{\tau}_{\nu\rho}) \star \bar{\mathrm{R}}^{\mu'}_{\{\alpha\} \ \mu} = O^{\mu'}_{\mu}[\lambda, 2](\Gamma^{\tau}_{\nu\rho}).$$

The *****-Ricci tensor:

Equation (101) is derived through,

$$\begin{aligned} \mathbf{\mathfrak{X}}_{\nu\rho} &= \langle dx^{\mu}; \mathbf{\mathfrak{X}}_{\mu\nu\rho}^{\sigma} \star \overleftarrow{\partial}_{\sigma} \rangle' \\ &= \langle dx^{\mu}; \bar{\mathbf{R}}^{\{\alpha\}} (\overleftarrow{\partial}_{\sigma}) \star \bar{\mathbf{R}}_{\{\alpha\}} (\mathbf{\mathfrak{X}}_{\mu\nu\rho}^{\sigma}) \rangle' , \quad (C14) \end{aligned}$$

noting that,

$$\begin{split} \bar{\mathbf{R}}^{\{\alpha\}} \begin{pmatrix} \star \\ \partial_{\sigma} \end{pmatrix} \star \bar{\mathbf{R}}_{\{\alpha\}} (\mathbf{\mathfrak{X}}^{\sigma}_{\mu\nu\rho}) &= \partial_{\sigma'}^{\star} \star (e^{-i\lambda\partial_0} \delta_{\sigma}^j \delta_{j}^{\sigma'} + \delta_{\sigma}^0 \delta_{0}^{\sigma'}) \mathbf{\mathfrak{X}}^{\sigma}_{\mu\nu\rho} \\ &= \partial_{\sigma'}^{\star} \star O_{\sigma}^{\sigma'} [\lambda, -2] (\mathbf{\mathfrak{X}}^{\sigma}_{\mu\nu\rho}), \quad (\text{C15}) \end{split}$$

and using this in (C14), we get

$$\begin{aligned} \mathbf{\mathfrak{X}}_{\nu\rho} &= \langle dx^{\mu} \star \partial_{\sigma'}^{\star} \star O_{\sigma}^{\sigma'} [\lambda, -2] \mathbf{\mathfrak{X}}_{\mu\nu\rho}^{\sigma} \rangle' \\ &= \delta_{\sigma'}^{\mu} \star O_{\sigma}^{\sigma'} [\lambda, -2] (\mathbf{\mathfrak{X}}_{\mu\nu\rho}^{\sigma}) = O_{\sigma}^{\mu} [\lambda, -2] (\mathbf{\mathfrak{X}}_{\mu\nu\rho}^{\sigma}) \\ &= \boxed{e^{-i\lambda\partial_0} \mathbf{\mathfrak{X}}_{j\nu\rho}^{j} + \mathbf{\mathfrak{X}}_{0\nu\rho}^{0}} . \end{aligned}$$
(C16)

The *-metric-compatibility:

Equation (102) is derived as follows,

$$\begin{split} \stackrel{\star}{\nabla}_{\gamma}(g) &= \stackrel{\star}{\nabla}_{\partial_{\gamma}^{*}}((dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) \star g_{\mu\nu}) \\ &= \left[\stackrel{\star}{\nabla}_{\partial_{\gamma}^{*}}(dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) \star g_{\mu\nu} \\ &+ \bar{\mathrm{R}}^{\{\alpha\}}(dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) \star \stackrel{\star}{\nabla}_{\bar{\mathrm{R}}_{\{\alpha\}}(\partial_{\gamma}^{*})}(g_{\mu\nu}) \right] \\ &= \left[(\stackrel{\star}{\nabla}_{\partial_{\gamma}^{*}}(dx^{\mu}) \overset{\star}{\otimes} dx^{\nu}) \star g_{\mu\nu} \\ &+ (\bar{\mathrm{R}}^{\{\alpha\}} \star \bar{\mathrm{R}}_{\{\alpha\}}^{\gamma'} \gamma \overset{\star}{\otimes} \stackrel{\star}{\nabla}_{\partial_{\gamma'}^{*}}(dx^{\nu})) \star g_{\mu\nu} \\ &+ \bar{\mathrm{R}}^{\{\alpha\}}(dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) \star \bar{\mathrm{R}}_{\{\alpha\}}^{\gamma'} \gamma \star \stackrel{\star}{\nabla}_{\partial_{\gamma'}^{*}}(g_{\mu\nu}) \right] \\ &= \left[-\Gamma_{\gamma(\mu}^{\sigma} \star g_{\nu)\sigma} + g_{\mu\nu,\gamma} \star (dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) \right]. (C17) \end{split}$$

Going from the fourth to the fifth line, we used the fact that,

$$\bar{\mathbf{R}}^{\{\alpha\}}(dx^{\mu} \overset{\star}{\otimes} dx^{\nu}) \star \bar{\mathbf{R}}^{\gamma'}_{\{\alpha\} \gamma} = O^{\gamma'}_{\gamma}[\lambda, 2](dx^{\mu} \overset{\star}{\otimes} dx^{\nu})$$
$$= dx^{\mu} \overset{\star}{\otimes} dx^{\nu}, \qquad (C18)$$

and that,

$$\bar{\mathbf{R}}^{\{\alpha\}}(dx^{\mu}) \star \bar{\mathbf{R}}^{\gamma'}_{\{\alpha\} \gamma} = O^{\gamma'}_{\gamma}[\lambda, 2](dx^{\mu}) = dx^{\mu}.$$
(C19)

- a. Addazi, et, Progress in Particle and Nuclear Physics 125, 103948 (2022), arXiv:2111.05659 [hep-ph].
- [2] G. Amelino-Camelia and D. V. Ahluwalia, International Journal of Modern Physics D 11, 35 (2002), arXiv:gr-qc/0012051 [gr-qc].
- [3] J. a. Magueijo and L. Smolin, Phys. Rev. D 67, 044017 (2003).
- [4] S. Hossenfelder, Living Reviews in Relativity 16, 2 (2013), arXiv:1203.6191 [gr-qc].
- [5] G. Amelino-Camelia, M. Matassa, F. Mercati, and G. Rosati, Phys. Rev. Lett. 106, 071301 (2011).
- [6] G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman, and L. Smolin, Phys. Rev. D 84, 084010 (2011).
- [7] J. M. Carmona, J. L. Cortés, and J. J. Relancio, Universe 7, 99 (2021), arXiv:2104.07336 [gr-qc].
- [8] G. Gubitosi and F. Mercati, Classical and Quantum Gravity 30, 145002 (2013), arXiv:1106.5710 [gr-qc].
- C. Rovelli, arXiv e-prints , gr-qc/9803024 (1998), arXiv:gr-qc/9803024 [gr-qc].
- [10] M. Born, Proceedings of the Royal Society of London Series A 165, 291 (1938).
- [11] C. A. Mead, Phys. Rev. 135, B849 (1964).

- [12] L. J. Garay, International Journal of Modern Physics A 10, 145 (1995), arXiv:gr-qc/9403008 [gr-qc].
- [13] J. F. Donoghue, Phys. Rev. D 50, 3874 (1994).
- [14] G. Amelino-Camelia, D. Benedetti, F. D'Andrea, and A. Procaccini, Classical and Quantum Gravity 20, 5353 (2003), arXiv:hep-th/0201245 [hep-th].
- [15] J. M. Carmona, J. L. Cortes, and J. J. Relancio, Class. Quant. Grav. 35, 025014 (2018), arXiv:1702.03669 [hep--th].
- [16] G. Amelino-Camelia and S. Majid, International Journal of Modern Physics A 15, 4301 (2000), arXiv:hep-th/9907110 [hep-th].
- [17] J. Lukierski and A. Nowicki, Acta Physica Polonica B 33, 2537 (2002), arXiv:hep-th/0207022 [hep-th].
- [18] G. Gubitosi and F. Mercati, Classical and Quantum Gravity 30, 145002 (2013).
- [19] G. Gubitosi and S. Heefer, Phys. Rev. D 99, 086019 (2019).
- [20] J. Kowalski-Glikman and S. Nowak, Physics Letters B 539, 126 (2002), arXiv:hep-th/0203040 [hep-th].
- [21] L. Freidel and L. Smolin, arXiv e-prints , arXiv:1103.5626 (2011), arXiv:1103.5626 [hep-th].

- [22] J. R. Oliveira, arXiv e-prints, arXiv:1110.5387 (2011), arXiv:1110.5387 [gr-qc].
- [23] J. Kowalski-Glikman and T. Trześniewski, Physics Letters B 737, 267 (2014).
- [24] T. Trześniewski, Nuclear Physics B 928, 448 (2018).
- [25] M. Arzano and J. Kowalski-Glikman, Springer 986 (2021), 10.1007/978-3-662-63097-6.
- [26] A. Ballesteros, I. Gutierrez-Sagredo, and F. J. Herranz, Physics Letters B 792, 175 (2019).
- [27] P. Maslanka, Journal of Physics A: Mathematical and General 26, L1251 (1993).
- [28] V. Chari and A. N. Pressley, Cambridge university press (1995).
- [29] Y. Kosmann-Schwarzbach, Springer Berlin (1997), 10.1007/BFb0113695.
- [30] S. Zakrzewski, Journal of Physics A: Mathematical and General 27, 2075 (1994).
- [31] A. Borowiec and A. Pachoł, Journal of Physics A Mathematical General 43, 045203 (2010), arXiv:0903.5251 [hep-th].
- [32] J. Kowalski-Glikman and S. Nowak, International Journal of Modern Physics D 12, 299 (2003), arXiv:hep-th/0204245 [hep-th].
- [33] A. Weinstein, Séminaire Bourbaki , 46 (1994).
- [34] M. Kontsevich, Letters in Mathematical Physics 66, 157 (2003), arXiv:q-alg/9709040 [math.QA].
- [35] G. Landi, arXiv e-prints , hep-th/9701078 (1997), arXiv:hep-th/9701078 [hep-th].
- [36] V. G. Drinfel'D, Advanced Series in Mathematical Physics 10, 264 (1990).
- [37] P. Aschieri and L. Castellani, International Journal of Modern Physics A 11, 4513 (1996), arXiv:q-alg/9601006 [math.QA].
- [38] P. P. Kulish and A. I. Mudrov, arXiv Mathematics e-prints , math/9901019 (1999), arXiv:math/9901019 [math.QA].
- [39] J. Madore, S. Schraml, P. Schupp, and J. Wess, European Physical Journal C 16, 161 (2000), arXiv:hep-th/0001203 [hep-th].
- [40] A. Connes, Springer (1994).
- [41] P. Aschieri, M. Dimitrijević, F. Meyer, and J. Wess, Classical and Quantum Gravity 23, 1883–1911 (2006).
- [42] P. Aschieri and L. Castellani, Journal of Geometry and Physics 60, 375 (2010).
- [43] F. Meyer, arXiv e-prints , hep-th/0510188 (2005), arXiv:hep-th/0510188 [hep-th].
- [44] P. Aschieri, C. Blohmann, M. Dimitrijević, F. Meyer, P. Schupp, and J. Wess, Classical and Quantum Gravity 22, 3511 (2005).
- [45] M. Dimitrijević and L. Jonke, Journal of High Energy Physics 2011, 80 (2011), arXiv:1107.3475 [hep-th].
- [46] A. Sykora and C. Jambor, arXiv e-prints , hep--th/0405268 (2004), arXiv:hep-th/0405268 [hep-th].
- [47] N. Herceg, T. Jurić, A. Samsarov, and I. Smolić, Journal of High Energy Physics 2024, 130 (2024), arXiv:2310.06038 [hep-th].
- [48] A. Borowiec, J. Lukierski, and A. Pachoł, Journal of Physics A Mathematical General 47, 405203 (2014), arXiv:1312.7807 [math-ph].
- [49] A. Borowiec and A. Pachol, Phys. Rev. D 79, 045012 (2009), arXiv:0812.0576 [math-ph].
- [50] V. Kharchenko, arXiv Mathematics e-prints , math/0005101 (2000), arXiv:math/0005101 [math.QA].

- [51] S. L. Woronowicz, Communications in Mathematical Physics 122, 125 (1989).
- [52] N. Reshetikhin, Letters in Mathematical Physics 20, 331 (1990).
- [53] R. Oeckl, Nucl. Phys. B 581, 559 (2000), arXiv:hep-th/0003018.
- [54] J. Wess and B. Zumino, Nuclear Physics B Proceedings Supplements 18, 302 (1991).
- [55] S. Meljanac, A. Samsarov, M. Stojić, and K. S. Gupta, European Physical Journal C 53, 295 (2008), arXiv:0705.2471 [hep-th].
- [56] P. Kosiński, P. Maślanka, J. Lukierski, and A. Sitarz, arXiv e-prints, hep-th/0307038 (2003), arXiv:hep-th/0307038 [hep-th].
- [57] G. Felder and B. Shoikhet, arXiv Mathematics e-prints, math/0002057 (2000), arXiv:math/0002057 [math.QA].
- [58] A. Agostini, G. Amelino-Camelia, M. Arzano, and F. D'Andrea, arXiv e-prints , hep-th/0407227 (2004), arXiv:hep-th/0407227 [hep-th].
- [59] L. Möller, Journal of High Energy Physics 2005, 029 (2005), arXiv:hep-th/0409128 [hep-th].
- [60] C. Gonera, P. Kosiński, P. Maślanka, and S. Giller, Physics Letters B 622, 192 (2005), arXiv:hep-th/0504132 [hep-th].
- [61] P. Aschieri, L. Castellani, and M. Dimitrijević, Letters in Mathematical Physics 85, 39 (2008), arXiv:0803.4325 [hep-th].
- [62] P. Aschieri and L. Castellani, Journal of High Energy Physics 2009, 086 (2009), arXiv:0902.3817 [hep-th].
- [63] D. Rozental and O. Birnholtz, In preparation.

Quantity	Commutative Spacetime	Constant Non-Commutativity	$\kappa\text{-Minkowski}$ Non-Commutativity: results the $\kappa\text{-GR}$
The Spacetime Metric	$g_{\mu u}$	$g_{\mu u}[41]$	$g_{\mu u}$
Coor. Commutators	$[x^{\mu}, x^{\nu}] = 0$	$\left[\hat{x}^{\mu}, \hat{x}^{\nu}\right] = i\lambda\theta^{\mu\nu} \ [44]$	$\left[\hat{x}^{\mu}, \hat{x}^{\nu}\right] = i l_p \hat{x}^j \delta^{\mu}_{[0} \delta^{\nu}_{j]}$
The product $f \star g$	fg	$ \begin{aligned} fg + \frac{i}{2} \lambda \theta^{\mu\nu} (\partial_{\mu} f) (\partial_{\nu} g) \\ + \frac{1}{8} (i\lambda)^2 \theta^{\mu\nu} \theta^{\rho\sigma} \partial_{\mu} \partial_{\rho} (f) \partial_{\nu} \partial_{\sigma} (g) \end{aligned} [44] $	$ \begin{aligned} fg &+ i\frac{\lambda}{2} x^j f_{,[0}g_{,j]} \\ &+ \frac{1}{8} (i\lambda)^2 x^j x^l \left(f_{,0[0} \cdot g_{,l]j} + f_{,j[l} \cdot g_{,0]0} \right) \end{aligned} $
Basis Vectors	∂_{μ}	∂_{μ} [43]	$\overset{\star}{\partial}_{\mu} = e^{-i\frac{l_{p}}{2}\partial_{0}\delta_{\mu}^{j}\delta_{j}^{\mu'}}\partial_{\mu'}$
Lie Derivative	$\mathcal{L}_{\xi}\left(\cdot ight)$	$\sum_{n=0}^{\infty} \frac{1}{n!} (\frac{i\lambda}{2})^n \prod_{j=1}^n \theta^{\mu_j \nu_j} \frac{\partial^n (\xi^{\gamma})}{\partial^{\mu_1} \dots \partial^{\mu_n}} \frac{\partial^n (\partial_{\gamma} (\cdot))}{\partial^{\nu_1} \dots \partial^{\nu_n}} $ [43]	$\sum_{\substack{n=0\\n'=0}}^{\infty} \frac{(i\lambda/2)^{n+n'}}{n!n'!} [(-x^j\partial_j)^{n'} [\partial_0^n(\xi^\mu)] \overset{*}{\partial}_{\mu} \partial_0^{n'} (x^j\partial_j)^n] (\cdot)$
Leibniz Rule	$\partial_\mu (fg)$ = $\partial_\mu (f)g + f\partial(g)$	$\partial_{\mu}(f \star g) = \partial_{\mu}(f) \star g + f \star \partial_{\mu}(g) \ [44]$	$\partial_{\mu}(f \star g) = \partial_{\mu'}(f) \star O_{\mu}^{\mu'}[\lambda, -2](g) + f \star (\partial_{\mu}g)$
Metric Inverse	$g^{\mu u}$	$g^{\star\mu\nu} = 2g^{\mu\nu} - g^{\mu\alpha} \star g_{\alpha\beta} \star g^{\beta\nu} \ [44]$	$g^{\star\mu\nu} = 2g^{\mu\nu} - g^{\mu\alpha} \star g_{\alpha\beta} \star g^{\beta\nu} \ [44]$
Metric Determinant	$\frac{1}{4!}\epsilon^{\mu_1\ldots\mu_4\nu_1\ldots\nu_4}g_{\mu_1\nu_1}\ldots g_{\mu_4\nu_4}$	$\frac{1}{4!} \epsilon^{\mu_1 \dots \mu_4 \nu_1 \dots \nu_4} g_{\mu_1 \nu_1} \star g_{\mu_2 \nu_2} \star g_{\mu_3 \nu_3} \star g_{\mu_4 \nu_4}$	$\frac{1}{4!} \epsilon^{\mu_1 \dots \mu_4 \nu_1 \dots \nu_4} g_{\mu_1 \nu_1} \star g_{\mu_2 \nu_2} \star g_{\mu_3 \nu_3} \star g_{\mu_4 \nu_4}$
Cristoffel	$\Gamma^{\sigma}_{\nu\gamma} = g^{\sigma\mu} \frac{1}{2} [g_{\mu(\nu,\gamma)} - g_{\nu\gamma,\mu}]$	$\Gamma^{\sigma}_{\nu\gamma} = g^{\star^{\sigma\mu}} \star \frac{1}{2} [g_{\mu(\nu,\gamma)} - g_{\nu\gamma,\mu}] [44]$	$\Gamma^{\sigma}_{\nu\gamma} = \overset{\star}{g}^{\sigma\mu} \star \frac{1}{2} [g_{\mu(\nu,\gamma)} - g_{\nu\gamma,\mu}]$
Covariant Derivative	$\nabla_{\mu}u^{\nu} = \partial_{\mu}u^{\nu} + u^{\sigma}\Gamma^{\nu}_{\sigma\mu}$	$\stackrel{\star}{\nabla}_{\mu}u^{\nu} = \partial_{\mu}u^{\nu} + u^{\sigma} \star \Gamma^{\nu}_{\sigma\mu} \ [43]$	$\stackrel{\star}{\nabla_{\overset{\star}{\partial_{\mu}}}}(u^{\nu}) = \partial_{\mu}(u^{\nu}) + O^{\mu'}_{\mu}[\lambda, 2](u^{\sigma}) \star \Gamma^{\nu}_{\mu'\sigma}$
Riemann Tensor	$\mathfrak{X}^{\sigma}_{\mu\nu\rho} = \partial_{[\mu}\Gamma^{\sigma}_{\nu]\rho} + \Gamma^{\tau}_{\rho[\nu}\Gamma^{\sigma}_{\mu]\tau}$	$\mathfrak{R}^{\sigma}_{\mu\nu\rho} = \partial_{[\mu}\Gamma^{\sigma}_{\nu]\rho} + \Gamma^{\tau}_{\rho[\nu} \star \Gamma^{\sigma}_{\mu]\tau}[43]$	$\mathfrak{R}^{\sigma}_{\mu\nu\rho} = \partial_{[\mu}\Gamma^{\sigma}_{\nu]\rho} + O^{\gamma}_{[\mu}[\lambda,2](\Gamma^{\tau}_{\nu]\rho}) \star \Gamma^{\sigma}_{\gamma\tau}$
Ricci Tensor	$\mathbf{\mathfrak{X}}_{\nu\rho} = \mathbf{\mathfrak{X}}_{\mu\nu\rho}^{\mu}$	$\mathfrak{R}_{\nu\rho} = \mathfrak{R}^{\mu}_{\mu\nu\rho} \ [43]$	$\mathbf{\mathfrak{X}}_{\nu\rho} = e^{-i\lambda\partial_0}\mathbf{\mathfrak{X}}_{j\nu\rho}^j + \mathbf{\mathfrak{X}}_{0\nu\rho}^0$
Ricci Scalar	$\mathbf{\mathfrak{X}} = g^{\mu\nu} \mathbf{\mathfrak{X}}^{\sigma}_{\sigma\mu\nu}$	$\mathfrak{R} = g^{\star\mu\nu} \star \mathfrak{R}^{\sigma}_{\sigma\mu\nu} \ [43]$	$\mathfrak{X} = \overset{\star}{g}^{\mu\nu} \star \mathfrak{X}_{\nu\mu}$
Einstein Tensor	$\mathbf{\mathfrak{K}}_{\mu u} - \frac{1}{2}g_{\mu u}\mathbf{\mathfrak{K}}$	$\mathfrak{R}_{\mu\nu} - \frac{1}{2}g_{\mu\nu} \star \mathfrak{R} \ [41]$	$\mathbf{\mathfrak{R}}_{\mu u} - \frac{1}{2}g_{\mu u} \star \mathbf{\mathfrak{R}}$
Einstein-Hilbert Lagrangian	$\sqrt{-\det[g]}$ k	$\sqrt{-\det[g]}$ X + c.c	$\sqrt{-\det[g]}$ X + c.c

TABLE I. Differential Geometry & Gravity Theories in Commutative, Constantly Non-Commutative, and κ -Minkowski Non-Commutative Spacetimes