

EVERY FINITE SET OF NATURAL NUMBERS IS REALIZABLE AS ALGEBRAIC PERIODS OF A MORSE–SMALE DIFFEOMORPHISM

GRZEGORZ GRAFF*, WACŁAW MARZANTOWICZ**, ŁUKASZ PATRYK MICHALAK**',***, AND ADRIAN MYSZKOWSKI[◊]

ABSTRACT. A given self-map $f: M \rightarrow M$ of a compact manifold determines the sequence $\{L(f^n)\}_{n=1}^{\infty}$, of the Lefschetz numbers of its iterations. We consider its dual sequence $\{a_n(f)\}_{n=1}^{\infty}$ given by the Möbius inversion formula. The set $\mathcal{AP}(f) = \{n \in \mathbb{N} : a_n(f) \neq 0\}$ is called the set of algebraic periods.

We solve an open problem existing in literature by showing that for every finite subset \mathcal{A} of natural numbers there exist an orientable surface S_g , as well as a non-orientable surface N_g , of genus g , and a Morse–Smale diffeomorphism f of this surface such that $\mathcal{AP}(f) = \mathcal{A}$. For such a map it implies the existence of points of a minimal period n for each odd $n \in \mathcal{A}$. For the orientation-reversing Morse–Smale diffeomorphisms of S_g we identify strong restrictions on $\mathcal{AP}(f)$. Our method also provides an estimate of the number of homotopy classes of Morse–Smale diffeomorphisms, which is exponential in g .

1. INTRODUCTION

Let $f: X \rightarrow X$ be a self-map of a topological space X . One of the classical problems of the topological theory of dynamical systems is to find n -periodic points of f , i.e. $x \in X$ such that $f^n(x) = x$ for some $n \in \mathbb{N}$. If $f^k(x) \neq x$ for $k < n$, then n is called the minimal period of f at x , and x is a point of period n . Then the set $\{x, f(x), f^2(x), \dots, f^{n-1}(x)\}$ is called the orbit of x . We denote by $P^n(f) = \text{Fix}(f^n)$ the set of all n -periodic points, and by $P_n(f)$ the set of all points of period n . We have $P_n(f) = P^n(f) \setminus \bigcup_{k|n, k < n} P^k(f)$.

An important invariant describing the dynamics of f is the set of all minimal periods of f , denoted by $\text{Per}(f) = \{n \in \mathbb{N} : P_n(f) \neq \emptyset\}$.

One of the natural problems in the field of periodic point theory is the following question: *suppose that \mathcal{A} is a subset of natural numbers, is it possible to realize it as the minimal periods of a self-map of some topological space X ?* This problem has been extensively studied (mainly from the arithmetical point of view) by many authors (cf. [EPPW] and the expository paper [BGW]) even in the stronger version, when the number of orbits of each minimal period is also given. In general, it turns out that there are no obstacles for such realizability for very general class of maps and relatively simple spaces. In particular any (also infinite) sequence

2020 *Mathematics Subject Classification*. Primary 37C25 37E30, 57K20; Secondary 55M20, 37D15.

Key words and phrases. Minimal periods, Lefschetz numbers, surfaces, Morse–Smale diffeomorphisms, Nielsen–Thurston classification.

Research supported by the National Science Centre, Poland, within the grant Sheng 1 UMO-2018/30/Q/ST1/00228. The third author was also supported by the Slovenian Research and Innovation Agency program P1-0292 and grant J1-4001.

of natural numbers can be realized for smooth maps on two-dimensional torus \mathbb{T}^2 [Winds]. However, the approach in [Winds] provides the realization in the homotopy class of identity map, so the periodic points are not detectable by topological/homotopical methods. In this paper we address a more subtle version of the problem from the perspective of homotopy invariants and for a more restricted class of maps (see Main Problem below). We also discuss the relations of our problem with the existence of periodic points (refer to Proposition 3.8 and Corollary 3.9).

1.1 (Main Problem). *Can any finite set \mathcal{A} of natural numbers be the set of algebraic periods (see Definition 1.2) of a C^1 Morse–Smale diffeomorphism on a closed surface (orientable or non-orientable) of some genus g ?*

Remind that every Morse–Smale diffeomorphism has only a finite-number of periodic points by the definition (cf. [Smale]).

Below, we sketch the motivation to study the problem in such a formulation. The tools for posing and solving this problem are closely related to the classical concept of algebraic topology, namely the Lefschetz number $L(f)$, which is homotopy invariant (see [Je-Ma] for a comprehensive overview and details).

With the sequence $\{L(f^n)\}_{n=1}^\infty$, we associate the dual sequence given by the Möbius inversion formula

$$(1) \quad a_n(f) = \frac{1}{n} \sum_{k|n} \mu(n/k) L(f^k) = \frac{1}{n} \sum_{k|n} \mu(k) L(f^{n/k}),$$

where $\mu : \mathbb{N} \rightarrow \{-1, 0, 1\}$ is the Möbius function. From the Möbius inversion formula, it follows that $L(f^n) = \sum_{k|n} k a_k(f)$. By the definition, $a_n(f) \in \mathbb{Q}$, but in fact, they are integers due to the Dold’s theorem (cf. [Do]).

Definition 1.2. *The set $\{n \in \mathbb{N} : a_n(f) \neq 0\}$ is called the set of algebraic periods of f and is denoted $\mathcal{AP}(f)$.*

Example 1.3. Let X be a finite set and $f : X \rightarrow X$ a map. Then $L(f^n) = |P^n(f)|$, and consequently $a_n(f) = |P_n(f)|$. Thus, in this example $a_n(f) \neq 0$ implies $P_n(f) \neq \emptyset$ which gives $n \in \text{Per}(f)$.

In general, non-vanishing of $a_n(f)$ does not imply the existence of points of period n . However, for some special classes of smooth self-maps of manifolds, such as the Morse–Smale diffeomorphisms, or transversal C^1 maps, $a_n(f) \neq 0$ implies $P_n(f) \neq \emptyset$ if n is odd, or $P_n(f) \cup P_{\frac{n}{2}}(f) \neq \emptyset$ if n is even (see Proposition 3.8). The above led Llibre, Sirvent and other collaborators to study $\mathcal{AP}(f)$, or more precisely its odd part $\mathcal{AP}_{\text{odd}}(f) = \mathcal{AP}(f) \cap \{2\mathbb{N} - 1\}$. They had been considering situations when it is computable by direct algebraic topology methods (see [Ll-Si3] for an exposition, and the bibliography of the results of this group up to 2013). In fact, instead of algebraic periods, they considered a subset of natural numbers called “the set of minimal Lefschetz periods” denoted by $\text{MPer}_L(f)$ (see Definition 5.1). Later, these authors noted that this set does not contain even numbers (cf. [Ll-Si2]). Recently Graff, Lebedź and Myszowski showed in [GLM] that $\text{MPer}_L(f) = \mathcal{AP}_{\text{odd}}(f)$, which justifies to use only the notion of algebraic periods. The proof of [GLM] uses the k -periodic expansion (cf.

the formula (2)) and is based on the local expression of the fixed point index as a combination of basic periodic sequences. As an appendix, we provide a purely algebraic proof of the fact that $\text{MPer}_L(f) = \mathcal{AP}_{\text{odd}}(f)$ in Theorem 5.2.

A direct inspiration for this work was the papers [Ll-Si1, Ll-Si3] and the following question posed there:

Can any finite set \mathcal{A} of odd positive integers be the minimal set of Lefschetz periods for a C^1 Morse–Smale diffeomorphism on some orientable/non-orientable compact surface without boundary with a convenient genus g .

As stated by the authors in [Ll-Si1], in all their works they do not characterize the sets of homotopy classes of Morse–Smale diffeomorphisms. They only described a set called the minimal sets of Lefschetz periods $\text{MPer}_L(f)$ (Definition 5.1), i.e. $\mathcal{AP}(f) \setminus 2\mathbb{N}$ of quasi-unipotent homeomorphisms (i.e. homeomorphisms for which the induced linear map $H_*(f)$ on the homology of M has spectrum consisting of the roots of unity only). Remind that if $f : M \rightarrow M$ is a Morse–Smale diffeomorphism, then f is quasi-unipotent by the Shub result [Shub], but not conversely. Consequently their results gave only the necessary conditions, i.e. the restrictions on the algebraic periods of Morse–Smale diffeomorphisms never providing a geometrical realization. These restrictions were obtained by purely algebraic computation of the Lefschetz zeta function based on known forms of the homology groups, or cohomology rings, of studied manifolds.

In this work, by a direct geometrical argument we give a complete positive answer to the Problem 1.1, i.e. a stronger version of the stated above question Libre and Sirvent, proving that for any finite set $\mathcal{A} \subset \mathbb{N}$ there exists an orientable surface of genus g and a preserving orientation diffeomorphism $f : S_g \rightarrow S_g$ such that $\mathcal{AP}(f) = \mathcal{A}$ (Theorem 1.4). In the case we demand that the realization is performed in the class of reversing orientation Morse–Smale diffeomorphism we find strong restrictions on $\mathcal{AP}(f)$, namely $\mathcal{AP}(f) \cap \{2\mathbb{N} - 1\} = \emptyset$, i.e. the set of algebraic periods consists only of even numbers if f is a reversing orientation diffeomorphism of an orientable surface, which was first shown by A. Myszowski in his unpublished Ph. D. thesis [Mysz]. Here we provide another proof of this fact. This result corrects the wrong statement regarding S_3 in [Ll-Si1] (Theorem 5.7). Finally, we show that for any finite set $\mathcal{A} \subset \mathbb{N}$ there exists a non-orientable surface of genus g and a Morse–Smale diffeomorphism $f : N_g \rightarrow N_g$ such that $\mathcal{AP}(f) = \mathcal{A}$ (Theorem 1.4).

Theorem 1.4. *Let \mathcal{A} be a finite set of natural numbers. There exist a Morse–Smale diffeomorphism f on a closed surface such that $\mathcal{A} = \mathcal{AP}(f)$. The surface can be chosen both orientable and non-orientable. Moreover, f can be orientation-reversing (in orientable case) if and only if $\mathcal{A} \subset 2\mathbb{N}$.*

One of the consequences of the construction provided in Theorem 1.4 is an exponential growth in g of the number of different algebraically finite type mapping classes, so consequently of homotopy classes of the Morse–Smale diffeomorphisms on a surface of a given genus (Theorem 4.7).

Another important statement that we draw from our main theorem is the fact that for Morse–Smale diffeomorphisms (as well as more for a more general class of transversal maps)

we are able to provide not only the realization of algebraic periods $n \in \mathcal{AP}(f)$ but also a n -periodic point for n odd, see Subsection 3.1.

2. DESCRIPTION OF THE HOMOTOPY CLASSES OF HOMEOMORPHISMS OF SURFACES WHICH CONTAIN THE MORSE–SMALE HOMEOMORPHISM

In this subsection, we give a brief survey of known results that describe the homotopy classes of homeomorphisms, thus diffeomorphisms, of surfaces which contain the Morse–Smale diffeomorphisms. From now on, we assume that our surface is hyperbolic, i.e. it is of genus $g \geq 2$. The case of surfaces with non-negative Euler characteristic was studied separately in [da Rocha].

As the first we remind the Thurston classification theorem [FLP, Farb-Marg], called also the Nielsen–Thurston classification theorem.

Let f be a homeomorphism of a closed orientable surface S_g of genus $g \geq 2$. Denote by $[f]$ the set of all homeomorphisms of the surface S_g that are homotopic to f (the homotopy class of homeomorphisms containing f). According to the Nielsen–Thurston classification (see, [FLP, Thms 11.6, 11.7] and [Gri-Mor-Poch]) the set of all homotopy classes of homeomorphisms on S_g is represented as the union of four disjoint subsets $T1, T2, T3, T4$ distinguished by the conditions described below.

Theorem 2.1 (Nielsen–Thurston). *Let f be a homeomorphism of a closed orientable surface S_g , $g \geq 2$.*

1. *If $[f] \in T1$, then $[f]$ contains a periodic homeomorphism;*
2. *if $[f] \in T2$, then $[f]$ contains a reducible non-periodic homeomorphism of algebraically finite order;*
3. *if $[f] \in T3$, then $[f]$ contains a reducible homeomorphism which is neither periodic nor of algebraically finite order;*
4. *if $[f] \in T4$, then $[f]$ contains a pseudo-Anosov homeomorphism.*

The classes $T1$ and $T2$ are called *algebraically finite type* in the original Nielsen terminology. The Nielsen–Thurston theory was developed for orientable surfaces of genus $g \geq 2$. The set of homotopy classes of Morse–Smale diffeomorphisms of the torus, and also of non-orientable surfaces of genus ≤ 1 , i.e. the projective space and Klein bottle is described in [da Rocha] (see Theorem 2.3).

Roughly speaking, in the reducible case ($T2$ and $T3$), Theorem 2.1 states that one can cut a surface along the invariant collection of closed curves obtaining a “canonical form”. This constitutes a collection of surfaces with boundary, satisfying that for each of them some iteration of the map is its self-homeomorphism. As a consequence, any mapping class can be reduced into either only periodic pieces ($T2$) or periodic and pseudo-Anosov pieces ($T3$).

The analogue of Theorem 2.1 for non-orientable surfaces was shown by Yingqing Wu in [WuYing] (see [Par] for an exhaustive exposition). The statement of the mentioned version has the same formulation and the main idea of its proof is to proceed with $\tilde{f}^+ : S_g \rightarrow S_g$ the

preserving orientation lift of a homeomorphism $f : N_g \rightarrow N_g$, where $p : S_g \rightarrow N_g$ is two-sheets orientable cover of the non-orientable surface N_g .

Before next consideration, let us remind the Baer–Epstein theorem (cf. [Farb-Marg]):

Proposition 2.2. *For homeomorphisms of a surface the homotopy is equivalent to the isotopy.*

Moreover, by the density argument, every homotopy class of homeomorphisms contains a diffeomorphism, and an isotopy between two diffeomorphisms can be replaced by a smooth isotopy. Accordingly, in many formulations of Theorem 2.1, the isotopy classes are used instead of the homotopy classes.

In the forthcoming part of the paper, we will base essentially on the following result of Luis F. da Rocha [da Rocha, THM A]

Theorem 2.3 (L. F. da Rocha, 1985). *If M^2 is a two dimensional compact connected boundaryless manifold (orientable or not) with a negative Euler characteristic, then the condition that f is of algebraically finite type is necessary and sufficient for an isotopy class of f of M^2 to be Morse–Smale i.e. contains a Morse–Smale representant).*

In the case $\chi(M^2) \geq 0$ we have the following situation:

- (a) *if M^2 is the sphere, the projective plane, or the Klein bottle, then every isotopy class is Morse–Smale;*
- (b) *if M^2 is the two dimensional torus, an isotopy class is Morse–Smale if and only if:*

$$H_1(f) : H_1(\mathbb{T}^2; \mathbb{C}) \rightarrow H_1(\mathbb{T}^2; \mathbb{C})$$

has eigenvalues in the set $\{\pm 1, \pm i, 1/2 \pm (\sqrt{3}/2)i, -1/2 \pm (\sqrt{3}/2)i\}$.

Remark 2.4. *Theorem 2.3 was reproved by A.N. Bezdenezhykh and V.Z. Grines [Bez-Grin] for the class T1, and by V. Grines, A. Morozov, and O. Pochinka [Gri-Mor-Poch] for the class T2 of homeomorphisms of orientable surfaces. The authors of the referred articles used subtle analytical methods, and providing constructions of such Morse–Smale diffeomorphisms solving more advanced questions.*

Fact 2.5. *For a homeomorphism of $f : M \rightarrow M$ of a compact surface M the following are equivalent:*

- (1) $\{L(f^n)\}$ *is bounded,*
- (2) $\mathcal{AP}(f)$ *is finite,*
- (3) $H_1(f) : H_1(M; \mathbb{Z}) \rightarrow H_1(M; \mathbb{Z})$ *is quasi-unipotent.*

Proof. The equivalence between the first and the second condition is given in Theorem [Je-Ma, 3.1.46] (cf. [Bab-Bog]). Since the Lefschetz numbers of iterations of a diffeomorphism f of a surface are equal to $1 + \text{tr} H_1(f)^n + \pm 1$ in the orientable case, and respectively $1 + \text{tr} H_1(f)^n$ in the non-orientable case, the first and third conditions are equivalent. \square

However, let us mention here that the spectral radius of the representation $f \mapsto H_1(f)$ is not a strong enough invariant to determine whether $[f] \in T1 \cup T2$ (cf. [FLP] or [Farb-Marg] for the Thurston example of a pseudo-Anosov homeomorphism f with $H_1(f) = \text{id}$).

3. PROOF OF MAIN THEOREM

In the forthcoming considerations we will use the notion of k -periodic expansion of an arithmetic function $\psi : \mathbb{N} \rightarrow \mathbb{C}$ introduced in [Mar-NP], exposed widely in [Je-Ma], and used in [GLM], [Gr-L-NP]. In this language an arithmetic function ψ is represented as a series of elementary periodic functions

$$(2) \quad \psi(n) = \sum_{k|n} a_k(\psi)k = \sum_{k=1}^{\infty} a_k(\psi) \operatorname{reg}_k(n),$$

where the coefficients $a_k(\psi) \in \mathbb{C}$ are given by the Möbius inversion formula as in (1) and

$$(3) \quad \operatorname{reg}_k(n) = \sum_{l=0}^{k-1} \left(e^{2\pi i \frac{l}{k}} \right)^n = \begin{cases} 0 & \text{if } k \nmid n \\ k & \text{if } k | n \end{cases}$$

is the sum of n -powers of all roots of unity of degree k and is called the elementary k -periodic function. Conceptually, the k -periodic expansion is a discrete correspondent of the Fourier expansion, arithmetically it is linked with the Ramanujan sums, and its coefficients are used to study the behaviour of the number sequences which are related to the periodic points (see [EPPW]). In our context, it is just a convenient language to handle the sequence we are studying.

Denote by S_g an orientable, and by N_g a non-orientable closed surface of genus g . Similarly, $S_{g,k}$ and $N_{g,k}$ are compact surfaces of genus g with k boundary components.

Theorem 3.1. *Let $\mathcal{A} \subset \mathbb{N}$ be finite.*

- (1) *There exists a closed orientable surface S_g and an orientation-preserving homeomorphism $f: S_g \rightarrow S_g$ of algebraically finite type such that $\mathcal{AP}(f) = \mathcal{A}$ and the genus g of S_g is equal to*

$$\sum_{n \in \mathcal{A} \setminus \{1\}} n \quad \text{if } 1 \in \mathcal{A} \quad \text{and} \quad 1 + \sum_{n \in \mathcal{A}} n \quad \text{if } 1 \notin \mathcal{A}.$$

- (2) *There exists a closed orientable surface S_g and an orientation-reversing homeomorphism $f: S_g \rightarrow S_g$ of algebraically finite type such that $\mathcal{AP}(f) = \mathcal{A}$ if and only if $\mathcal{A} \subset 2\mathbb{N}$. Moreover, the genus g of S_g is equal to*

$$\sum_{4|n \in \mathcal{A}} 2n + \sum_{4 \nmid n \in \mathcal{A} \setminus \{2\}} n \quad \text{if } 2 \in \mathcal{A} \quad \text{and} \quad 2 + \sum_{4|n \in \mathcal{A}} 2n + \sum_{4 \nmid n \in \mathcal{A}} n \quad \text{if } 2 \notin \mathcal{A}.$$

- (3) *There exists a closed non-orientable surface N_g and a homeomorphism $f: N_g \rightarrow N_g$ of algebraically finite type such that $\mathcal{AP}(f) = \mathcal{A}$ and the genus g of N_g is equal to*

$$\sum_{n \in \mathcal{A} \setminus \{1\}} n \quad \text{if } 1 \in \mathcal{A} \neq \{1\}, \quad 2 + \sum_{n \in \mathcal{A}} n \quad \text{if } 1 \notin \mathcal{A} \quad \text{and} \quad 1 \quad \text{if } \mathcal{A} = \{1\}.$$

Proof. First, let us describe the following construction. Having a finite set $\mathcal{A}' \subset \mathbb{N}$ and an assignment $\tau: \mathcal{A}' \rightarrow \mathbb{N}$, for each $n \in \mathcal{A}'$ consider a surface Σ_n , which is an orientable surface $S_{\tau(n),2}$ in the orientable case or a non-orientable surface $N_{\tau(n),2}$ in the non-orientable case. On each Σ_n we take a periodic homeomorphism f_n of order $\tau(n)$ which cyclically permutes its

1-handles and is a rotation on each boundary component (see Figure 1). More specifically, for an illustration, start with a cylinder $S^1 \times [-1, 1]$ with a rotation by $1/\tau(n)$ of a full angle on each circle $S^1 \times \{t\}$ and remove $\tau(n)$ open disjoint discs $D_1, \dots, D_{\tau(n)}$ in such a way that the rotation cyclically permutes the discs. Now glue $\tau(n)$ copies of $S_{1,1}$ (or $N_{1,1}$) along permuted boundary components and extend the rotation to a periodic homeomorphism of order $\tau(n)$ that cyclically permutes the attached handles.

Next, having surfaces $\Sigma_{n_1}, \dots, \Sigma_{n_k}$ let us join Σ_{n_i} and $\Sigma_{n_{i+1}}$ for $1 \leq i \leq k - 1$ by identifying their boundary components with end circles of a cylinder $C_i = S^1 \times [-1, 1]$. By the construction, we get a surface S with two boundary components and define f to be equal to f_{n_i} on Σ_{n_i} , and on each joining cylinder it is given by a homotopy between two rotations on end circles.




FIGURE 1.

(1) In the case of an orientable surface and an orientation-preserving homeomorphism, take $\mathcal{A}' = \mathcal{A} \cup \{1\}$ if $1 \notin \mathcal{A}$ or $\mathcal{A}' = \mathcal{A} \setminus \{1\}$ if $1 \in \mathcal{A}$, with an assignment $\tau(n) = n$ for $n \in \mathcal{A}'$. The above procedure provides a surface S with an orientation-preserving homeomorphism f . Let us attach two discs to S to get a closed orientable surface \tilde{S} , and f extends on \tilde{S} , being an appropriate rotation on each of two discs. Taking one circle $S^1 \times \{0\}$ for each cylinder C_i we get a system of simple closed curves preserved by f such that f is periodic on the complement of its open tubular neighbourhood $\bigcup \text{Int}C_i$, so f is of algebraically finite type.

Moreover, let $B_n = \{a_1^n, \dots, a_n^n, b_1^n, \dots, b_n^n\}$ consist of circles in S_n such that $\bigcup_{n \in \mathcal{A}'} B_n$ forms the standard (symplectic) basis for \tilde{S} . Then the matrix of $H_1(f)$ on $H_1(\tilde{S})$ is a block diagonal matrix whose blocks correspond to B_n 's and since $f(a_j^n) = f_n(a_j^n) = a_{(j+1) \bmod n}^n$ and similarly $f(b_j^n) = b_{(j+1) \bmod n}^n$, the n th block is the direct sum of two permutation matrices for a cycle of length n :

$$\begin{bmatrix} 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{bmatrix}.$$

Therefore, the characteristic polynomial of $H_1(f)$ is equal to

$$\prod_{n \in \mathcal{A}'} (x^n - 1)^2,$$

and so it is not difficult to observe by the definition of reg_k that: $\text{tr}(H_1(f^l)) = \sum_{n \in \mathcal{A}'} 2 \text{reg}_n(l)$. By the definition of \mathcal{A}' ,

$$(4) \quad L(f^l) = 2 - \text{tr}(H_1(f^l)) = \sum_{n \in \mathcal{A}} a_n(f) \text{reg}_n(l),$$

where $a_n(f) = -2$ for $n \in \mathcal{A} \setminus \{1\}$, $a_1(f) = 2$ if $1 \in \mathcal{A}$ and $a_n(f) = 0$ for $n \notin \mathcal{A}$. Thus $\mathcal{AP}(f) = \mathcal{A}$.

(2) For the orientation-reversing case, we will show later that there are no odd algebraic periods (see Theorem 3.2). Assume that $\mathcal{A} \subset 2\mathbb{N}$ and take $\mathcal{A}' = \mathcal{A} \cup \{2\}$ if $2 \notin \mathcal{A}$ or $\mathcal{A}' = \mathcal{A} \setminus \{2\}$ if $2 \in \mathcal{A}$ with an assignment $\tau(n) = n$ if 4 divides n and $\tau(n) = n/2$ otherwise. Again, the procedure provides an orientable surface S , and we attach a disc to one of its two boundary components (name the resulting surface also S). We take its double $D(S)$, i.e. consider a copy S' of S consisting of surfaces Σ'_n and cylinders C'_i , and join S and S' by a new cylinder $C = S^1 \times [-1, 1]$ along their boundary components. We have defined f on S and S' , and on the new cylinder, let $f|_{S^1 \times \{t\}}$ be the same as $f|_{S^1 \times \{\pm 1\}}$.

Now, let $j: D(S) \rightarrow D(S)$ be an orientation-reversing involution mapping points of S into corresponding points of S' , e.g. it can be seen as the reflection through the plane intersecting $D(S)$ in $S^1 \times \{0\} \subset C$ and separating symmetrically S and S' , see Figure 1 (2). The desired self-homeomorphism of $D(S)$ is $f' = f \circ j$. Indeed, cutting $D(S)$ along circles from connecting cylinders, we get the decomposition into surfaces Σ_n and Σ'_n such that f' maps Σ_n homeomorphically onto Σ'_n and conversely. Thus $(f')^2$ is a self homeomorphism of each piece Σ_n or Σ'_n on which one can check it is $n/2$ -periodic. Hence f' is of algebraically finite type.

Moreover, it is clear that $(f')^2(a_j^n) = a_{(j+2) \bmod \tau(n)}^n$, $(f')^2(b_j^n) = b_{(j+2) \bmod \tau(n)}^n$ on Σ_n and similarly for Σ'_n . If 4 divides n , then $\tau(n) = n$ and so $(f')^2|_{\Sigma_n}$ has order $n/2$, thus the n th block of the matrix of $H_1(f')$ is the direct sum of four permutation matrices for a cycle of length n . However, if n is not divisible by 4, then $\tau(n) = n/2$ is odd and $(f')^2|_{\Sigma_n}$ still has

order $\tau(n) = n/2$. Therefore the characteristic polynomial of f' is equal to

$$\prod_{4|n \in \mathcal{A}'} (x^n - 1)^4 \prod_{4 \nmid n \in \mathcal{A}'} (x^n - 1)^2$$

and so

$$L((f')^l) = 2\text{reg}_2(l) - \text{tr}(H_1((f')^l)) = \sum_{n \in \mathcal{A}} a_n(f') \text{reg}_n(l),$$

where $a_n(f') = -4$ for $n \in \mathcal{A}$ such that $4|n$, $a_n(f') = -2$ for $n \in \mathcal{A} \setminus \{2\}$ such that $4 \nmid n$ and $a_2(f') = 2$ if $2 \in \mathcal{A}$. Obviously, $a_n(f') = 0$ for $n \notin \mathcal{A}$, so $\mathcal{AP}(f') = \mathcal{A}$.

(3) Analogously we construct a desired self-homeomorphism of a non-orientable surface. If $\mathcal{A} = \{1\}$, then just take $\text{id}_{\mathbb{R}P^2}$. Otherwise, consider $\mathcal{A}' = \mathcal{A} \cup \{1\}$ if $1 \notin \mathcal{A}$ or $\mathcal{A}' = \mathcal{A} \setminus \{1\}$ if $1 \in \mathcal{A}$ with an assignment $\tau(n) = n$ for $n \neq 1$ and $\tau(1) = 2$ if $1 \in \mathcal{A}'$. Moreover, in the preliminary construction for the non-orientable case if $1 \in \mathcal{A}'$ we take $f_1 = \text{id}$ on Σ_1 which is of genus $\tau(1) = 2$. We get a non-orientable surface S with two boundary components to which we attach discs as previously. Similarly as in (1) we have a self-homeomorphism f of S of algebraically finite type, which is n -periodic on each Σ_n , except $n = 1$.

Let $B_n = \{a_1^n, \dots, a_{\tau(n)}^n\}$ be a set of circles in Σ_n which are cyclically permuted by f and such that elements of $\bigcup B_n$ represent standard generators of $H_1(S) \cong \mathbb{Z}^{g-1} \oplus \mathbb{Z}/2$ with the relation $2 \sum_{i,n} a_i^n = 0$, where g is the genus of S . The trace of $H_1(f)$ is computed as the trace of $\widetilde{H_1(f)}$ on $H_1(S)/T \cong \mathbb{Z}^{g-1}$, which is induced by $H_1(f)$ after dividing by the torsion part T of $H_1(S)$. Thus $H_1(S)/T \cong \left(\bigoplus_{i,n} \mathbb{Z} a_i^n \right) / \left\langle \sum_{i,n} a_i^n \right\rangle$ has a basis $(\bigcup B_n) \setminus \{a\}$ for some fixed $a \in B_{n_0}$ and a is represented as the vector $(-1, \dots, -1)$ in this basis. Hence the matrix of $\widetilde{H_1(f)}$ is a block matrix if $n_0 = 1 \in \mathcal{A}'$ or almost a block matrix if $n_0 \neq 1$, i.e. the structure of a block matrix is disturbed only in the column which corresponds to a basis element $a_i^{n_0}$ such that $f(a_i^{n_0}) = a$ (if we order the basis in such a way that $a_i^{n_0}$ is the last element, then the matrix is upper block triangular). However, in the second case the characteristic polynomial of $\widetilde{H_1(f)}$ is still the product of characteristic polynomials of permutation matrices for cycles of length n for $n \neq n_0$, the identity matrix of dimension 2 if $1 \in \mathcal{A}'$ and the characteristic polynomial of

$$(5) \quad \begin{bmatrix} 0 & 0 & \cdots & 0 & -1 \\ 1 & 0 & \cdots & 0 & -1 \\ 0 & 1 & \cdots & 0 & -1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & -1 \end{bmatrix},$$

which is the companion matrix of the polynomial $x^{n_0-1} + \dots x + 1 = (x^{n_0} - 1)/(x - 1)$. Since the trace of the companion matrix of a polynomial is equal to the sum of its roots, by the formula (3), the trace of l -power of the matrix (5) is equal to $\text{reg}_{n_0}(l) - \text{reg}_1(l)$. Thus in both the cases we get:

$$\text{tr}(H_1(f^l)) = \sum_{n \in \mathcal{A}' \setminus \{1\}} \text{reg}_n(l) + \sum_{1 \in \mathcal{A}'} 2\text{reg}_1(l) - \text{reg}_1(l),$$

so

$$L(f^l) = 1 - \text{tr}(H_1(f^l)) = \sum_{n \in \mathcal{A}} a_n(f) \text{reg}_n(l),$$

where $a_n(f) = -1$ for $n \in \mathcal{A} \setminus \{1\}$, $a_1(f) = 2$ if $1 \in \mathcal{A}$ and $a_n(f) = 0$ for $n \notin \mathcal{A}$. Hence $\mathcal{AP}(f) = \mathcal{A}$. □

To complete the proof, we are left with showing that $\mathcal{AP}(f) \cap (2\mathbb{N} - 1) = \emptyset$ if f is orientation-reversing, because we have shown only this realization for $\mathcal{AP}(f) \subset 2\mathbb{N}$.

In order to do this, we will need the notion of antisymplectic maps related to the maps induced on homology.

Each orientation-preserving homeomorphism $h: S_g \rightarrow S_g$ induces the linear map $H_1(h)$ that preserves the intersection form on $H_1(S_g; \mathbb{Z}) \cong \mathbb{Z}^{2g} = \langle a_1, \dots, a_g, b_1, \dots, b_g \rangle$:

$$\langle \cdot, \cdot \rangle: \mathbb{Z}^{2g} \times \mathbb{Z}^{2g} \rightarrow \mathbb{Z}$$

given by

$$\langle a_i, b_j \rangle = \delta_{ij} = -\langle b_j, a_i \rangle, \quad \langle a_i, a_j \rangle = 0, \quad \langle b_i, b_j \rangle = 0,$$

whose matrix is

$$\Omega = \begin{bmatrix} 0 & I_g \\ -I_g & 0 \end{bmatrix}.$$

Thus, if A is the matrix of $H_1(h)$ in the basis $\langle a_1, \dots, a_g, b_1, \dots, b_g \rangle$, then

$$(6) \quad \Omega = A^T \Omega A,$$

so A is a symplectic matrix.

Suppose $h: S_g \rightarrow S_g$ is an orientation-reversing homeomorphism. Thus it induces $-\text{id}$ on $H_2(S_g) = \mathbb{Z}$ and so the matrix A of $H_1(h)$ in the standard basis satisfies

$$(7) \quad A^T \Omega A = -\Omega.$$

Such a matrix is called *antisymplectic*.

In particular, one can show that $\det(A) = (-1)^g$. The matrix $M = I_g \oplus (-I_g)$ is clearly antisymplectic and $\det(M) = (-1)^g$. If A is antisymplectic, then AM is symplectic, and so $\det(AM) = 1$. Therefore $\det(A) = (\det(M))^{-1} = (-1)^g$.

It is not difficult to verify that for a square matrix A of dimension n ,

$$\chi_{A^{-1}}(x) = \frac{1}{\det(-A)} x^n \chi_A \left(\frac{1}{x} \right) \quad \text{and} \quad \chi_{-A}(x) = (-1)^n \chi_A(-x).$$

Therefore, if A is antisymplectic of dimension $2g$, then $A^T \Omega = -\Omega A^{-1}$, and so

$$(8) \quad \begin{aligned} \chi_A(x) &= \det(xI - A) = \det(xI - A^T) \det(\Omega) = \det(x\Omega - A^T \Omega) \\ &= \det(x\Omega + \Omega A^{-1}) = \det(xI - (-A^{-1})) = \chi_{-A^{-1}}(x) \\ &= (-1)^{2g} \frac{1}{\det(-A)} (-x)^{2g} \chi_A \left(-\frac{1}{x} \right) = (-1)^g x^{2g} \chi_A \left(-\frac{1}{x} \right). \end{aligned}$$

Theorem 3.2. *Let $h: S_g \rightarrow S_g$ be an orientation-reversing quasi-unipotent homeomorphism (e.g. Morse–Smale diffeomorphism). Then $L(h^{\text{odd}}) = 0$ and so $\mathcal{AP}(h) \subset 2\mathbb{N}$, i.e. h has no odd algebraic periods.*

Proof. For n odd, h^n is also orientation-reversing, so $L(h^n) = \text{tr}(H_1(h^n))$ and $H_1(h^n)$ is antisymplectic. Therefore it suffices to show that $L(h) = 0$ just for h .

The characteristic polynomial $p(x)$ of $H_1(h)$ satisfies (8), so if $p(x) = (x - \xi)^k q(x)$. then

$$p(x) = (x - \xi)^k q(x) = (-1)^g x^{2g} \left(-\frac{1}{x} - \xi \right)^k q \left(-\frac{1}{x} \right).$$

Thus if ξ is a root of $p(x)$ of multiplicity k , then $p(x)$ has a root $-1/\xi$ of multiplicity k . In particular, if ξ is a primitive root of unity of odd order l , then $-1/\xi$ is a primitive root of unity of order $2l$, and conversely, if ξ is primitive of order $2l$, l odd, then $\xi^l = -1$, so $-1/\xi$ is primitive of order l .

Therefore, $p(x)$ has a factor $\varphi_l(x)$, l odd, with multiplicity k if and only if it has a factor $\varphi_{2l}(x)$ with multiplicity k . Since the sum of the roots of $\varphi_m(x)$ is equal to $\mu(m)$, $L(h)$ is a sum of the summands: $\mu(l) + \mu(2l) = \mu(l) - \mu(l) = 0$ for each odd l such that $\varphi_l(x)$ is a factor of $p(x)$ and $\mu(4m) = 0$ for each factor $\varphi_{4m}(x)$ of $p(x)$. Thus $L(h) = 0$.

By the Möbius inversion formula,

$$a_n(h) = \sum_{d|n} \mu(n/d) L(h^d) = 0$$

for odd n since all $d|n$ are also odd. □

Proof of Theorem 1.4. The statement of Theorem 1.4 follows from Theorems 2.3, 3.1, and Theorem 3.2. □

Remark 3.3. *Note that all the statements related to genus g in Theorem 3.1 remain valid for Morse–Smale diffeomorphisms in Theorem 1.4.*

Remark 3.4. *In fact Llibre and coauthors considered orientation-reversing case in [Ll-Si1] (respectively, orientation-preserving case in papers cited in [Ll-Si3]) and they did not take into account the topological restrictions that come from the structure of cohomological ring which are encoded in the formula (7) (or respectively in (6)), which resulted in these papers in listing also non-realizable algebraic periods. In particular, as we showed above, these restrictions force that there are no odd numbers in the set of algebraic periods for orientation-reversing homeomorphisms.*

3.1. Existence of periodic points for Morse–Smale diffeomorphisms and transversal maps. To formulate a consequences of our main result related to the existence of periodic points we remind the notion of a transversal map.

Definition 3.5 ([Do], [Fr]). *Let $f: \mathcal{U} \rightarrow M$ be a C^1 -map of an open subset \mathcal{U} of a manifold M , $\dim M = d$. We say that f is transversal if for any $m \in \mathbb{N}$ and $x \in P^m(f)$, $1 \notin \sigma(Df^m(x))$, where σ denotes the spectrum of the derivative D of $f^m(x)$.*

The set of all transversal maps $\mathcal{U} \rightarrow M$ is denoted by $C_T(\mathcal{U}, M)$ or $C_T(M)$ if $\mathcal{U} = M$. The main property of the class of transversal maps is given in the following theorem (see [Do] if $M = \mathbb{R}^d$, and [Seidel] for the general case, also [Je-Ma] for an exposition).

Theorem 3.6. *The set $C_T^\infty(\mathcal{U}, M)$ is generic in $C^0(\mathcal{U}, M)$, i.e. it is the intersection $C_T^\infty(\mathcal{U}, M) = \bigcap_{n=1}^\infty G_n$ where G_n is open and dense in $C^0(\mathcal{U}, M)$. In particular every map $f: \mathcal{U} \rightarrow M$ is homotopic to a transversal map $h: \mathcal{U} \rightarrow M$.*

A geometric property of a transversal map states the following.

Proposition 3.7. *Let M be a closed manifold. For any $f \in C_T(M)$ and every $m \in \mathbb{N}$ the set $P^m(f)$ consists of isolated points.*

By definition, Morse–Smale diffeomorphisms belong to $C_T(M)$ (cf. [Smale], [Fr]). We denote the set of transversal self-diffeomorphisms of M by $DC_T(M)$ and the set of Morse–Smale diffeomorphisms by $DC_{MS}(M)$.

Finally, all mappings in $C_T(M)$, thus $DC_T(M)$ and $DC_{MS}(M)$, have the following geometric property, which states that non-vanishing $a_n(f)$, n -odd, implies that $P_n(f) \neq \emptyset$, i.e. $n \in \text{Per}(f)$. We have (cf. [Do], and [Fr] for Morse–Smale diffeomorphism).

Proposition 3.8 (see [Je-Ma, Corollary 3.3.10] for an exposition).

Let $f: M \rightarrow M$ be a transversal map. If $a_n(f) \neq 0$ then

$$\begin{cases} P_n(f) \cup P_{\frac{n}{2}}(f) \neq \emptyset & \text{if } n \text{ is even,} \\ P_n(f) \neq \emptyset & \text{if } n \text{ is odd.} \end{cases}$$

The above proposition allows us to formulate the analytical (dynamical) consequence of Theorem 1.4. Put $\mathcal{AP}_{\text{odd}}(f) = \mathcal{AP}(f) \cap \{2\mathbb{N} - 1\}$, and $\mathcal{AP}_{\text{ev}}(f) = \mathcal{AP}(f) \cap \{2\mathbb{N}\}$.

Corollary 3.9. *Let $\mathcal{A} \subset \mathbb{N}$ be a finite subset of natural numbers and $f: S_g \rightarrow S_g$ be a preserving, or reversing orientation Morse–Smale diffeomorphism of orientable surface of genus g given by Theorem 1.4, or correspondingly a Morse–Smale diffeomorphism of a non-orientable surface N_g given by Theorem 1.4, such that $\mathcal{AP}(f) = \mathcal{A}$.*

Then for every $n \in \mathcal{AP}_{\text{odd}}(f)$ and every $h \in C_T(S_g)$, respectively $h \in C_T(N_g)$, homotopic to f , we have $n \in \text{Per}(h)$. For every $n \in \mathcal{AP}_{\text{ev}}(f)$ and every $h \in C_T(S_g)$, respectively $h \in C_T(N_g)$, homotopic to f , we have $(\{n\} \cup \{\frac{n}{2}\}) \cap \text{Per}(h) \neq \emptyset$.

4. FINAL DISCUSSION AND APPLICATIONS

In this section we describe deep relations of the considered concepts with Nielsen periodic point theory. In the next step, we give an estimate from below of the number of algebraically finite type mapping classes of surfaces homeomorphisms.

First, we show that the fact that isotopy classes of Morse–Smale diffeomorphisms are contained in $T1 \cup T2$ follows also from the Nielsen number theory.

We remind that the asymptotic Nielsen number of a self-map $f: M \rightarrow M$ of a compact manifold M , correspondingly the asymptotic generalized Lefschetz number of f , are defined as

$$N^\infty(f) = \limsup \sqrt[n]{N(f^n)}, \quad \text{and respectively} \quad L_\Gamma^\infty(f) = \limsup \sqrt[n]{\|L_\Gamma(f^n)\|},$$

where the generalized Lefschetz number $L_\Gamma(f)$ is defined as an element of the group ring $\mathbb{Z}(\Gamma)$ of $\Gamma = \pi_1(T_f)$ where T_f is the mapping torus f (cf. (1.4) [BoJia]).

For a surface without boundary, [BoJia, Thm 3.7] states the following: *if $f : M \rightarrow M$ is a homeomorphism of a compact connected surface with $\chi(M) < 0$ (orientable or not orientable), then*

$$N^\infty(f) = L_\Gamma^\infty(f) = \lambda,$$

where λ is the largest stretching (expanding) factor of the pseudo-Anosov pieces in the Thurston canonical form of f . Lemma [BoJia, Lem. 3.6] adds that $\log \lambda = \mathbf{h}(f)$ is equal to the topological entropy of the canonical representative of f .

Remark 4.1. *As a matter of fact in [BoJia, Thm 3.7] is shown more. Namely $N^\infty(f) = NI^\infty(f)$, where the latter is equal to $\limsup \sqrt[n]{NP_n(f)}$ and $NP_n(f)$ is so-called n -th Nielsen–Jiang prime periodic number (see [BoJia] for a definition, or [Je-Ma] for a longer exposition). This invariant has the property $NP_n(f) \leq |P_n(f)|$. It shows that in the case of a homeomorphisms of a surface S_g the condition $N^\infty(f) = \lambda > 1$ implies that $P(f) = \bigcup_{n=1}^{\infty} P^n(f)$, and also $\text{Per}(f)$, are infinite. Since $N(f^n)$ and $NP_n(f)$ are homotopy invariants, this yields that a Morse–Smale diffeomorphisms can only occur in the classes $T1$ or $T2$. In other words it provides one direction implication of Theorem 2.3: the set of the isotopy classes of Morse–Smale diffeomorphisms is contained in $T1 \cup T2$.*

4.1. Mapping classes. It is very useful to use the language of mapping classes in the study dynamics of homeomorphisms of surfaces. It is caused by the two factors. First, we are interested in homotopy properties of homeomorphisms of surfaces, second, the theory of mapping classes group is well-developed and several theorems we use are formulated in these terms.

By the definition, *the mapping class group* $\text{Mod}(S_g)$ of S_g is the quotient group

$$\text{Homeo}^+(S_g)/\text{Homeo}_0(S_g),$$

where $\text{Homeo}^+(S_g)$ denotes the group of preserving orientation homeomorphisms of S_g and $\text{Homeo}_0(S_g)$ its subgroup of homeomorphisms isotopic to the identity (cf. [Farb-Marg], [Par]). Thus the assignment $\phi \mapsto [\phi] \in \text{Mod}(S_g)$ is a surjection from $\text{Homeo}^+(S_g)$ onto $\text{Mod}(S_g)$.

This allows us to define an action (a representation) of the group $\text{Mod}(S_g)$ in $H_1(S_g; \mathbb{Z}) \subset H_1(S_g; \mathbb{R})$. More precisely, for a given class $[\phi]$ of $\phi \in \text{Homeo}^+(S_g)$ we define $\Psi([\phi]) := H_1(\phi) \in \text{Aut}(H_1(S_g; \mathbb{Z}))$, where $H_1(\phi)$ is the induced homomorphism of $H_1(S_g; \mathbb{Z})$. It is well defined, i.e. does not depend on a choice of representative of the class, because the subgroup $\text{Homeo}_0(S_g)$ acts trivially on $H_1(S_g; \mathbb{Z})$. In fact, $H_1(\phi)$ belongs to the group $Sp(2g, \mathbb{Z})$ of symplectic matrices of size n according to the property (6).

Now we can formulate a classical fact (that we use below to study transversal maps) about the mapping class group (cf. [Farb-Marg, Proposition 7.3]).

Theorem 4.2 (H. Burkhardt (1889)). *The homomorphism $\Psi : \text{Mod}(S_g) \rightarrow Sp(2g, \mathbb{Z})$ is surjective.*

Another classical fact was shown by J. Nielsen by a direct geometrical consideration and a calculation of the characteristic polynomial (cf. [Nielsen]). Nowadays there are many ways of proving this statement.

Theorem 4.3 (J. Nielsen 1944).

If $[f] \in T1 \cup T2$, then $H_1(f)$ is quasi-unipotent.

Combining the stated above facts we get the following.

Proposition 4.4. *Every $f \in [h]$ of a homeomorphism $h : M \rightarrow M$ of orientable or non-orientable surface M such that $N^\infty(h) = 1$ is isotopic to a Morse–Smale diffeomorphism, thus it is quasi-unipotent and $\mathcal{AP}(f)$ is finite. Conversely, for every homeomorphism f of M which is homotopic to a Morse–Smale diffeomorphism we have $N^\infty(f) = 1$.*

Consequently, a class $[f]$ contains a Morse–Smale diffeomorphism if and only if it contains a homeomorphism f' with the entropy $\mathbf{h}(f') = 0$.

Proof. If $N^\infty(h) = 1$ then by the Boju Jiang theorem there is not a pseudo-Anosov piece in the canonical Nielsen–Thurston form of h . Indeed, this theorem states that if $N^\infty(h) = \lambda$ then λ is the largest stretching factor of the pseudo-Anosov pieces in the canonical form of h . However, for a pseudo-Anosov diffeomorphism we have $\lambda > 1$ which leads to a contradiction if it would be such a piece.

Consequently, $[h] \in T1 \cup T2$. Now, by the Nielsen theorem (Theorem 4.3) it is quasi-unipotent. By da Rocha’s Theorem 2.3 there is a Morse–Smale diffeomorphism $\tilde{h} \in [h]$.

Next, from Fact 2.5 it follows that $\mathcal{AP}(h) = \mathcal{AP}(\tilde{h})$ is finite.

Finally, if $N^\infty(h) > 1$ then h and every $h' \sim h$ has infinitely many periodic orbits which gives a contradiction if h' is Morse–Smale.

The last part, regarding entropy, follows from already quoted [BoJia, Lem. 3.6]. □

Let us remind that the mapping class group $\text{Mod}(S_g)$, can be also defined as the quotient $\text{Diffeo}^+(S_g)/\text{Diffeo}_0(S_g)$, (cf. [Farb-Marg]).

Proposition 4.5. *For every symplectic or antisymplectic matrix $A \in \text{Gl}(2g, \mathbb{Z})$ there exists a transversal diffeomorphism $f \in \text{DC}_T(S_g)$, for which the induced automorphism $H_1(f)$ is equal to A .*

Proof. By Theorem 4.2 for every $A \in \text{Sp}(2g, \mathbb{Z})$ there exists a diffeomorphism h of S_g for which the induced automorphism $H_1(h)$ is equal to A . To get a transversal diffeomorphism $f \in \text{DC}_T(S_g)$ homotopic to h , we use the fact that the set of all transversal self-diffeomorphisms of S_g is dense in $\text{Diffeo}(S_g)$ by Theorem 3.6.

In the case of an antisymplectic matrix A take any orientation-reversing homeomorphism $h : S_g \rightarrow S_g$. Then the product $AH_1(h)$ is symplectic and by the first part of the proof induced by a transversal diffeomorphism f' . Thus $f = f' \circ h^{-1}$ induces $H_1(f) = A$ and again can be approximated by a transversal diffeomorphism. □

4.2. An estimate of the number of algebraically finite type classes. The formulas of Theorem 3.1 lead to an estimate from below of the number of algebraically finite type mapping classes for a fixed genus g . Let us consider two homeomorphisms f', f'' of a surface M . Observe that if they have different k -periodic expansions of Lefschetz numbers of iterations (cf. formula (2)) they could not be homotopic. As result the number of different expansions estimates the number of homotopy classes.

This observation and our construction allow us to find an estimate of the number of non-homotopic diffeomorphisms in $T1 \cup T2$.

Definition 4.6. The number of ways of writing the integer N as a sum of positive integers, where the order of addends is not considered significant, is denoted by $P(N)$ and is called *the number of unrestricted partitions*.

A partition of N can be represented by a sequence (p_1, \dots, p_N) such that $N = \sum_{k=0}^N p_k k$, so $p_k \geq 0$ is the number of integers k in the partition of N .

More information about the functions $P(N)$ one can find in [Wei1]. At this moment let us only mention the asymptotic behavior of it (Hardy-Ramanujan 1918):

$$(9) \quad P(N) \sim \frac{1}{4N\sqrt{3}} e^{\pi\sqrt{2N/3}}.$$

Theorem 4.7. *The number of algebraically finite type mapping classes of an orientable or non-orientable closed surface of genus g is estimated from below by $P(g)$, the number of unrestricted partitions of g . Consequently, there are at least $P(g)$ homotopy classes of Morse–Smale diffeomorphisms on S_g and on N_g .*

Finally, the asymptotic growth in g of this number is greater than or equal to $\frac{1}{4g\sqrt{3}} e^{\pi\sqrt{2g/3}}$.

Proof. Let us first consider the orientable case. We will provide a correspondence between partitions (p_1, \dots, p_g) of g and homeomorphisms $f: S_g \rightarrow S_g$ of algebraically finite type given by

$$(10) \quad a_n(f) = \begin{cases} -2p_n & \text{if } n \neq 1, \\ -2(p_1 - 1) & \text{if } n = 1, \end{cases}$$

where $a_n(f)$ denote the coefficients in the k -periodic expansion of f .

We apply the construction from the proof of Theorem 3.1. Recall that in the orientation-preserving case (1) we realized \mathcal{A} as $\mathcal{AP}(f)$ of a map $f: S_g \rightarrow S_g$ such that $a_k(f) = -2$ for $k \in \mathcal{A} \setminus \{1\}$, $a_1(f) = 2$ if $1 \in \mathcal{A}$ and obviously $a_k(f) = 0$ for $k \notin \mathcal{A}$. From the description of the genus or just from the construction it follows that $2g = 2 - \sum_k a_k(f) \cdot k$ and so one can easily check that $g = \sum_k p_k k$ by the formula (10). Recall that the surface S_g was formed from the pieces Σ_k for $k \in \mathcal{A}'$ and connecting cylinders.

Now, for a given partition (p_1, \dots, p_g) of g and every k we take p_k copies of the surface Σ_k of genus k together with defined k -periodic homeomorphisms and glue them appropriately as in Theorem 3.1. From the construction it is straightforward that $g = \sum p_k k$ is the genus of

the resulted surface and $\text{tr}H_1(f^n) = \sum_k 2p_k \text{reg}_k(n)$. Therefore we obtained an algebraically finite type homeomorphism f of S_g such that

$$L(f^n) = 2 - \text{tr}H_1(f^n) = \sum_k a_k(f) \text{reg}_k(n)$$

is consistent with the formula (10).

Now consider the non-orientable case. The analogous correspondence between partitions (p_1, \dots, p_g) of g and homeomorphisms $f: N_g \rightarrow N_g$ of algebraically finite type is given by

$$(11) \quad a_n(f) = \begin{cases} -p_n & \text{if } n \neq 1, \\ 2 - p_1 & \text{if } n = 1. \end{cases}$$

Recall that in the construction in the proof of Theorem 3.1 in the non-orientable case (3) we realized \mathcal{A} as $\mathcal{A}P(f)$ of $f: N_g \rightarrow N_g$ such that $a_k(f) = -1$ for $k \in \mathcal{A} \setminus \{1\}$. Similarly, for a given partition (p_1, \dots, p_g) of g we repeat the construction with p_k copies of the non-orientable surface Σ_k of genus k on which the obtained function is k -periodic. Therefore the obtained surface is clearly of genus $g = \sum p_k k$ and by the computations during the proof of Theorem 3.1 we get

$$\text{tr}H_1(f^n) = -\text{reg}_1(n) + \sum_k p_k \text{reg}_k(n).$$

Thus f is an algebraically finite type homeomorphism of N_g such that

$$L(f^n) = 1 - \text{tr}H_1(f^n) = \sum_k a_k(f) \text{reg}_k(n),$$

where the coefficients $a_k(f)$ of the k -periodic expansion of f are determined by (11).

In both the cases, by Theorem 2.3 of da Rocha the mapping class of f contains a Morse–Smale representative. The relations (10) and (11) show that diffeomorphisms corresponding in our construction to different unrestricted partitions of g have different k -periodic expansions, and consequently are not homotopic, i.e. they are in different mapping classes. \square

5. APPENDIX - ALGEBRAIC PERIODS AND MINIMAL LEFSCHETZ PERIODS

We remind that the zeta function for the sequence of Lefschetz numbers $L = \{l_n\}_n := \{L(f^n)\}_n$ of iterations is defined as:

$$(12) \quad \zeta_f(z) = \zeta(L; z) := \exp\left(\sum_{n=1}^{\infty} \frac{l_n}{n} z^n\right).$$

In [Ll-Sil] the authors defined *the set of minimal Lefschetz periods* of a diffeomorphism $f: M \rightarrow M$, denoted $\text{MPer}_L(f)$, in the following way.

Definition 5.1.

$$(13) \quad \text{MPer}_L(f) := \bigcap \{r_1, \dots, r_\eta\}$$

where the intersection is taken over all representations of $\zeta(L; z)$ as

$$\zeta(L; z) = \prod_{i=1}^{\eta} (1 + \Delta_i z^{r_i})^{m_i} = \prod_{i=1}^{\eta'} (1 - z^{r_i})^{m'_i} \prod_{j=1}^{\eta''} (1 + z^{r_j})^{m''_j}, \quad m_i, m'_i, m''_j \in \mathbb{Z}, m_i = m'_i + m''_j,$$

where $\Delta_i = \pm 1$, i.e. we take into account rational representations of $\zeta(L; z)$ as products of powers and polynomials $(1 + \Delta_i z^{r_i})$.

Geometrically, it known (see generalized Franks formula being a consequence of (3.3.9) in [Je-Ma]) that $\zeta(L; z)$ has such a representation, with each term related to some periodic orbit in case of transversal maps.

However, the factors and its amount are not canonically determined by the sequence $\{L(f^n)\}_n$ in this formula. Llibre and coauthors derived the set $\text{MPer}_L(f)$ for several examples (see the references of [Ll-Si3]) observing meantime that it does not contain even numbers by a simple algebraic argument included here in the proof of Proposition 5.2 (cf. [Ll-Si2]). Notify also, that formally we have infinitely many of such rational representations potentially with several possible values η , r_i , and m_i , which geometrically reflects the fact that it may appear periodic orbits of high periods whose contributions to $\zeta(L, z)$ annihilate one another.

Recently, in [GLM] the authors showed that for a transversal map $f \in C_T(M)$ of a manifold M we have $\text{MPer}_L(f) = \mathcal{AP}(f) \cap \{2\mathbb{N} - 1\} = \mathcal{AP}_{\text{odd}}(f)$ using the k -periodic expansion of the sequence $\{L(f^n)\}$. Originally this theorem is stated for Morse–Smale diffeomorphisms, but the argument holds for the class of transversal maps. It is worth pointing out that in [GLM] is used the fact that all summands in representation of $\{L(f^n)\}_n$ come from the geometric representation of $\zeta(L, z)$. Below we present another purely algebraic way of showing that $\text{MPer}_L(f) = \mathcal{AP}_{\text{odd}}(f)$ for a larger class of maps.

By the definition, the Lefschetz zeta function $\zeta(L; z) \in \mathbb{Z}[[z]]$ is a power series which is a rational function over \mathbb{Z} , i.e. an element of $\mathbb{Z}(z)$ the field of fractions of the ring of integral polynomials, as follows from the homology formula for the Lefschetz numbers (cf. [Je-Ma, (3.1.27)]). Moreover, $\zeta(L; z)$ has the following multiplicative representation (cf. [Je-Ma, (3.1.22)])

$$(14) \quad \zeta(L; z) = \prod_{n=1}^{\infty} (1 - z^n)^{a_n(f)}.$$

Note that in the case when the sequence $\{L(f^n)\}_n$ is bounded, e.g. if f is a Morse–Smale diffeomorphism, the set $\mathcal{AP}(f) = \{n : a_n(f) \neq 0\}$ is finite, and consequently the product (14) is finite.

Moreover, the formula (14) is a unique rational expression of $\zeta_f(z)$ in terms of polynomials $(1 - z^n)$ since none of the polynomials $(1 - z^n)$ can be expressed as a rational function of polynomials $(1 - z^k)$ for $k \neq n$. Indeed, let $(1 - z^n) = \prod_{k \neq n} (1 - z^k)^{c_k}$ for $c_k \in \mathbb{Z}$ and finitely many $c_k \neq 0$, and let k_0 be the largest number such that $c_{k_0} \neq 0$.

- if $k_0 < n$, then the left-hand side has a zero at a primitive root of unity of degree n , but the right-hand side does not,

- if $k_0 > n$, then the right-hand side has a zero or pole at a primitive root of unity of degree k_0 , but the left-hand side does not.

Proposition 5.2. *Let $f: X \rightarrow X$ be a map of a finite CW-complex X such that the sequence $\{L(f^n)\}_n$ of Lefschetz numbers of iterations is bounded. Then*

$$\text{MPer}_L(f) = \mathcal{AP}_{\text{odd}}(f).$$

Proof. We have just three steps:

- (1) The formula (14) gives $\text{MPer}_L(f) \subset \mathcal{AP}(f)$.
- (2) Since $(1 - z^{2n}) = (1 - z^n)(1 + z^n)$ and $(1 + z^{2n}) = \frac{1 - z^{4n}}{(1 - z^n)(1 + z^n)}$, $\text{MPer}_L(f) \cap 2\mathbb{N} = \emptyset$.
- (3) Finally, if

$$(15) \quad \zeta_f(z) = \prod (1 - z^k)^{c_k} \cdot \prod (1 + z^k)^{d_k},$$

then use $(1 + z^k) = \frac{1 - z^{2k}}{1 - z^k}$ to write

$$\zeta_f(z) = \prod (1 - z^k)^{c_k} \cdot \prod \frac{(1 - z^{2k})^{d_k}}{(1 - z^k)^{d_k}} = \prod (1 - z^k)^{e_k},$$

where $e_k = c_k + d_{k/2} - d_k$ if k is even and $e_k = c_k - d_k$ if k is odd. Since the form (14) is unique, $e_k = a_k(f)$. This means that $c_n - d_n = a_n(f) \neq 0$ for $n \in \mathcal{AP}_{\text{odd}}(f)$. In particular $c_n \neq 0$ or $d_n \neq 0$, so $n \in \text{MPer}_L(f)$ because (15) is arbitrary.

Therefore, $\text{MPer}_L(f) = \mathcal{AP}_{\text{odd}}(f)$. □

REFERENCES

- [Bab-Bog] I. K. Babenko, S. A. Bogatyř, *Behavior of the index of periodic points under iterations of a mapping.* (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 55(1991), no.1, 3–31; translation in Math. USSR-Izv. **38** (1992), no.1, 1–26.
- [Bez-Grin] A. N. Bezdenzhykh, V.Z. Grines, *Realization of gradient-like diffeomorphisms of two-dimensional manifolds*, Sel. Math. Sov. **11**(1) (1992), 19–23.
- [BGW] J. Byszewski, G. Graff and T. Ward, *Dold sequences, periodic points, and dynamics*, Bull. Lond. Math. Soc. **53** (2021), no. 5, 1263–1298.
- [Do] A. Dold, *Fixed point indices of iterated maps*, Inventiones Math., **74** (1985), 419–435.
- [Farb-Marg] B. Farb, D. Margalit, *A primer on mapping class groups*. Princeton Math. Ser., 49 Princeton University Press, Princeton, NJ, 2012. xiv+472 pp.
- [FLP] A. Fathi, F. Laudenbach, V. Ponearu, *Travaux de Thurston sur les surfaces*, Asterisque, **66–67**, Soc. Math. France, 1979), 267–282. English translation by D. Margalit and D. J. Kim: *Thurston’s Work on Surfaces (MN-48)*, Princeton University Press, 2012.
- [Fr] J. Franks, *Some smooth maps with infinitely many hiperbolic point*, Trans. Am. Math. Soc. **226** (1977), 175–179.
- [GLM] G. Graff, M. Lebiedź, A. Myszkowski, *Periodic expansion in determining minimal sets of Lefschetz periods for Morse–Smale diffeomorphisms*, J. Fixed Point Theory Appl. (2019) 21:47.
- [Gr-L-NP] G. Graff, M. Lebiedź, P. Nowak-Przygodzki, *Generating sequences of Lefschetz numbers of iterates*, Monatsh. Math. **188**(3) (2019), 511–525.
- [Gri-Mor-Poch] V. Grines, A. Morozov, O. Pochinka, *Determination of the Homotopy Type of a Morse–Smale Diffeomorphism on an Orientable Surface by a Heteroclinic Intersection*, Qualitative Theory of Dynamical Systems **22:120** (2023), <https://doi.org/10.1007/s12346-023-00809-9>.

- [Gri-Med-Poch] V. Grines, T. Medvedev, O. Pochinka, *Dynamical Systems on 2- and 3-Manifolds*, Springer, 2018.
- [BoJia] B. Jiang, *Estimation of the number of periodic orbits*, Pacific J. Math. **172** (1996), no.1, 151–185.
- [Je-Ma] J. Jezierski, W. Marzantowicz, Homotopy methods in topological fixed and periodic points theory. Topological fixed point theory and its applications, Vol. 3, Springer, Dordrecht (2006)
- [Ll-Si1] J. Llibre, V. F. Sirvent, *Minimal sets of periods for Morse–Smale diffeomorphisms on orientable compact surfaces*, Houston J. Math. **35(3)** (2009), 835–855.
- [Ll-Si2] J. Llibre, V. F. Sirvent, *Erratum: Minimal sets of periods for Morse–Smale diffeomorphisms on orientable compact surfaces*, Houston J. Math. **36** (2010), 335–336.
- [Ll-Si3] J. Llibre, V. F. Sirvent, *A survey on the minimal sets of Lefschetz periods for Morse–Smale diffeomorphisms on some closed manifolds*, (English summary), Publ. Mat. Urug. **14** (2013), 171–186.
- [Mar-NP] W. Marzantowicz, P. Nowak-Przygodzki, *Finding periodic points of a map by use of a k -adic expansion*, Discrete Contin. Dyn. Syst. **5(3)** (1999), 495–514.
- [Mysz] A. Myszkowski, Ph.D. Thesis, Gdańsk University of Technology, 2023, (in Polish).
- [Nielsen] J. Nielsen, *Surface transformation classes of algebraically finite type*, Danske Vid. Selsk. Mat.-Fys. Medd. **21 no.2** (1944), 89 pp.
- [EPPW] G. Everest, A. J. van der Poorten, Y. Puri and T. Ward, *Integer sequences and periodic points*, J. Integer Seq. **5(2)** (2002), Article 02.2.3, 10 pp.
- [Par] L. Paris, *Mapping class groups of non-orientable surfaces for beginners*, (English summary) Winter Braids Lect. Notes1, Winter Braids IV (Dijon, 2014)(2014), Exp. No. 3, 17 pp.
- [da Rocha] L. F. da Rocha, *Characterization of Morse–Smale isotopy classes on surfaces*, Ergod. Th. & Dynam. Sys. (**1985**) **5**, 107–122.
- [Seidel] P. Seidel, *Trasversalität für periodische Punkte differenzierbarer Abbildungen*, Mathematisches Institut, Universität Heidelberg, Forschergruppe: Topologie und nichtkommutative Geometrie, Preprint Nr 97, July 1994.
- [Shub] M. Shub, *Morse–Smale Diffeomorphisms are Unipotent on Homology*, *Dynamical systems*, (Proc. Sympos., Univ. Bahia, Salvador, 1971). Academic Press,
- [Sh-Su] M. Shub, D. Sullivan, *A remark on the Lefschetz fixed point formula for differentiable maps*, Topology **13** (1974), 189–191.
- [Smale] S. Smale, *Differentiable dynamical systems*, Bull. Amer. Math. Soc. **73** (1967), 747–817.
- [Wei1] E. W. Weisstein, Partition Function P , <http://mathworld.wolfram.com/PartitionFunctionP.html>
- [Winds] A. J. Windsor, *Smoothness is not an obstruction to realizability*, Ergodic Theory Dynam. Systems **28** (2008), no.3, 1037–1041.
- [WuYing] Wu Yingqing, *Canonical reducing curves of surface homeomorphism*, Acta Math. Sinica (N.S.) **3 no. 4** (1987), 305–313.

* FACULTY OF APPLIED PHYSICS AND MATHEMATICS, GDAŃSK UNIVERSITY OF TECHNOLOGY, NARUTOWICZA 11/12, 80-233 GDAŃSK, POLAND

Email address: `grzegorz.graff@pg.edu.pl`

** FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, ADAM MICKIEWICZ UNIVERSITY OF POZNAŃ, UL. UNIwersytetu Poznańskiego 4, 61-614 POZNAŃ, POLAND

Email address: `marzan@amu.edu.pl`

*** INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, JADRANSKA 19, SI-1000 LJUBLJANA, SLOVENIA

Email address: `lukasz.michalak@amu.edu.pl`

◇ INDEPENDENT RESEARCHER

Email address: `adrian.myszkowski@pg.edu.pl`