Dynamically syndetic sets and the combinatorics of syndetic, idempotent filters

Daniel Glasscock¹ and Anh N. Le^2

¹Dept. of Mathematics and Statistics, U. of Massachusetts Lowell, Lowell, MA, USA ²Dept. of Mathematics, U. of Denver, Denver, CO, USA

Abstract

A subset of the positive integers is *dynamically central syndetic* if it contains the set of times that a point returns to a neighborhood of itself under a minimal transformation of a compact metric space. These sets are part of the highly-influential link between dynamics and combinatorics forged by Furstenberg and Weiss in the 1970's. Our main result is a characterization of dynamically central syndetic sets as precisely those sets that belong to syndetic, idempotent filters. Idempotent filters are combinatorial objects that abound in ergodic Ramsey theory but have been largely unnoticed and unexplored. We develop the algebra of these objects for the proof of the main theorem and with an eye toward future applications.

The main result is best contextualized as a "global" analogue to Bergelson and Hindman's "local" characterization of Furstenberg's central sets as members of minimal, idempotent ultrafilters. It leads to a dual characterization of sets of topological pointwise recurrence, allowing us to answer a question of Glasner, Tsankov, Weiss, and Zucker. We draw numerous striking contrasts between pointwise recurrence and set recurrence, a topic with a long history in the subject and its applications, and answer four questions posed by Host, Kra, and Maass. We also show that the intersection of a dynamically central syndetic set with a set of pointwise recurrence must be piecewise syndetic, generalizing results of Dong, Glasner, Huang, Shao, Weiss, and Ye.

²⁰²⁰ Mathematics Subject Classification. Primary: 37B20. Secondary: 37B05.

Key words and phrases. return time sets in minimal topological dynamical systems, syndetic sets, thick sets, piecewise syndetic sets, central sets, IP sets, dynamically syndetic sets, dynamically thick sets, sets of pointwise recurrence, topological recurrence, measurable recurrence, idempotent filters, idempotent ultrafilters

Contents

| 1 | Introduction | | 2 |
|----------|--|---|-----------|
| | 1.1 | Part I: Dynamically syndetic sets | 2 |
| | 1.2 | Part II: Dynamically thick sets | 7 |
| | 1.3 | Part III: Dynamically piecewise syndetic sets | 11 |
| | 1.4 | Organization of the paper | 13 |
| | 1.5 | Acknowledgements | 14 |
| 2 | Preliminaries | | |
| | 2.1 | Families of positive integers | 15 |
| | 2.2 | Topology and dynamics | 21 |
| | 2.3 | Families of sets from dynamics | 23 |
| 3 | Part | I: Dynamically syndetic sets | 27 |
| | 3.1 | Definitions and first characterizations | 27 |
| | 3.2 | Central syndetic sets | 35 |
| | 3.3 | Central and dynamically central syndetic sets $\ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots$ | 38 |
| | 3.4 | Combinatorial characterizations: proofs of Theorems A and B $\ldots \ldots \ldots \ldots \ldots \ldots$ | 47 |
| 4 | Part II: Dynamically thick sets and sets of pointwise recurrence | | |
| | 4.1 | Definition and first characterizations | 48 |
| | 4.2 | Combinatorial characterizations: proofs of Theorems C and D $\ \ldots \ $ | 52 |
| | 4.3 | More general examples of dynamically thick sets | 57 |
| | 4.4 | The partition problem and σ -compactness | 61 |
| 5 | Part III: Dynamically piecewise syndetic sets | | 67 |
| | 5.1 | Definitions and first results | 67 |
| | 5.2 | Filter algebra for central and piecewise syndetic family duals | 69 |
| | 5.3 | Polynomial set recurrence and Brauer configurations: a proof of Theorem H \ldots . | 72 |
| 6 | Open questions | | 75 |
| | 6.1 | Dynamically syndetic sets | 76 |
| | 6.2 | Dynamically thick sets | 77 |
| | 6.3 | Dynamically piecewise syndetic sets | 79 |

1. Introduction

1.1. Part I: Dynamically syndetic sets

A subset of the positive integers, \mathbb{N} , is called *dynamically syndetic* if it contains a set of the form

 $R_T(x,U) := \{ n \in \mathbb{N} \mid T^n x \in U \}, \quad x \in X, \quad U \subseteq X \text{ nonempty, open,}$

where (X, d_X) is a compact metric space and the map $T : X \to X$ generates a minimal action of the semigroup $(\mathbb{N}, +)$ on X by continuous maps (see Section 2.2.1). In this case, the pair (X, T) is called a *minimal topological dynamical system*. If $x \in U$, such a set is called *dynamically central syndetic*. Dynamically syndetic and dynamically central syndetic sets are the main objects of study in this paper.

Examples of dynamically syndetic sets abound in combinatorics and number theory. Periodic sets and their almost-periodic and higher-order generalizations, namely Bohr and nil-Bohr sets, are dynamically syndetic. The images of Beatty sequences and the set of those positive integers with even 2-adic valuation are examples of dynamically central syndetic sets. We give further examples in Section 3 and an instructive non-example at the beginning of Section 1.1.2 below.

1.1.1. Motivation: structure in dynamics and combinatorics

Dynamically syndetic sets are structural objects that form part of the bridge between dynamics and combinatorics. We will mention two highly influential sets of results that exemplify this.

Furstenberg and Weiss [19, 22] pioneered the use of ergodic theory and topological dynamics as tools in Ramsey theory and combinatorial number theory. A correspondence principle – a device used to create a dynamical object out of a combinatorial one – makes available a wide range of tools and ideas from dynamics. For example, a topological correspondence principle can be used to reduce van der Waerden's theorem – ostensibly concerning arbitrary finite partitions of \mathbb{N} in its original formulation [50] or arbitrary syndetic subsets of \mathbb{N} in an equivalent formulation [38] – to a statement concerning dynamically syndetic sets. The first step in Furstenberg's highly influential proof of Szemerédi's theorem [19] is also of this nature, transferring the problem from combinatorics to one about measure preserving systems. This set of ideas has had a huge impact on the field and continues to yield dividends [51].

Rotations on compact abelian groups are among the simplest examples of dynamical systems, and the dynamically syndetic sets they generate, *Bohr sets*, are among the simplest examples of dynamically syndetic sets. Nilsystems – algebraic generalizations of compact group rotations – give rise to dynamically syndetic sets called *nil-Bohr sets*. Since ground-breaking work of Host and Kra [33] and Ziegler [53], nilsystems and nil-Bohr sets have been used to describe the "structured" components in powerful decomposition theorems in ergodic theory [7, 16, 45], additive combinatorics and number theory [27, 28, 29, 30] and, recently, in topological dynamics [23]. Thus, dynamically syndetic sets arising from highly structured systems have become essential in defining structure, even in non-dynamical settings.

In their full generality, the families of dynamically (central) syndetic sets have appeared explicitly in the literature in a few places. In various combinations, Dong, Huang, Shao, and Ye [12, 36, 37] have studied these classes (they call them *m*-sets and *sm*-sets) in the context of disjointness between topological dynamical systems. The family of dynamically central syndetic sets was studied in a combinatorial context by Bergelson, Hindman, and Strauss [6] in connection to the quality of sets of return times of points to open sets under polynomial iterates of an irrational rotation. A subfamily of dynamically syndetic sets was considered by Kennedy, Raum, and Salomon [39]; we will discuss some of their results more carefully in the context of our own below.

1.1.2. The main result: a characterization of dynamically syndetic sets

A subset of \mathbb{N} is syndetic if it has bounded gaps, that is, if there exists $N \in \mathbb{N}$ such that it has nonempty intersection with every interval of length N. Syndeticity is a generalization of periodicity that is intimately connected to minimality in topological dynamics. That dynamically syndetic sets are syndetic, for example, is a consequence of a well-known characterization of minimal actions: the system (X, T) is minimal if and only if every set of the form $R_T(x, U)$ is syndetic.

It is not the case, however, that all syndetic sets are dynamically syndetic. Indeed, here

is an instructive example to keep in mind:

$$\left(2\mathbb{N}\cap\bigcup_{\substack{n=0\\n \text{ even}}}^{\infty} \left[2^n, 2^{n+1}\right)\right) \cup \left(\left(2\mathbb{N}+1\right)\cap\bigcup_{\substack{n=0\\n \text{ odd}}}^{\infty} \left[2^n, 2^{n+1}\right)\right).$$
(1.1)

That this set is syndetic is obvious. That this set is not dynamically syndetic is less obvious. We leave the discovery of a short argument to the intrepid reader, opting instead to show this as an application of Theorem A below.

This example raises a question that has appeared explicitly and implicitly in a number of different contexts in the literature: how do we know if a given set is dynamically syndetic? The following theorem – the main result of our paper – answers this question by giving necessary and sufficient conditions for a set to be dynamically syndetic. Theorem A is proved in Section 3.4.

Theorem A. Let $A \subseteq \mathbb{N}$. The following are equivalent.

- (i) The set A is dynamically syndetic.
- (ii) There exists a nonempty subset $B \subseteq A$ that satisfies: for all finite $F \subseteq B$, the set $\bigcap_{f \in F} (B f)$ is syndetic.
- (iii) There exists $n \in \mathbb{N}$ for which the set A n belongs to a syndetic, idempotent filter on \mathbb{N} .

Point (ii) characterizes dynamical syndeticity in terms of a simple combinatorial property. Let us use it to see that the set in (1.1), call it A, is not dynamically syndetic. If $B \subseteq A$ satisfies the condition in (ii), then it must be syndetic. Every syndetic subset of A, in particular, B, must contain both an even integer, b_0 , and an odd integer, b_1 . It is quick to see, however, that the set $(A - b_0) \cap (A - b_1)$ is not syndetic, contradicting what we had supposed about B. In addition to giving a way to check whether a set is dynamically syndetic, the combinatorial condition in (ii) allows us to characterize members of the dual class, the class of dynamically thick sets, as we define and discuss in Section 1.2.

It is worth comparing point (ii) in Theorem A with a recent result of Kennedy, Raum, and Salomon [39, Thm. 8.2]. In the context of the positive integers, it is a consequence of their result that a set $A \subseteq \mathbb{N}$ is dynamically syndetic if and only if there exists $B \subseteq A$ such that for all finite $F_1 \subseteq B$ and $F_2 \subseteq \mathbb{N} \setminus B$, the set

$$\bigcap_{f_1 \in F_1} \left(B - f_1 \right) \cap \bigcap_{f_2 \in F_2} \left((\mathbb{N} \setminus B) - f_2 \right)$$
(1.2)

is syndetic. It is not hard to see that a set $B \subseteq \mathbb{N}$ satisfies the condition in (1.2) if and only if its indicator function 1_B is a uniformly recurrent point in the full shift $(\{0,1\}^{\mathbb{N}},\sigma)$; see Section 2.2.1 for the notation and terminology. That a set is dynamically syndetic if and only if it contains such a subset follows then from, eg., [37, Prop. 2.3] (see also Theorem 3.9). The condition in Theorem A (ii) is substantially weaker than that in (1.2). In particular, it does not imply that the point 1_B is uniformly recurrent. This weaker condition translates to an easier-to-verify sufficient condition for dynamical syndeticity.

The following is an analogous characterization for dynamically central syndetic sets. Theorem B is proved in Section 3.4.

Theorem B. Let $A \subseteq \mathbb{N}$. The following are equivalent.

(i) The set A is dynamically central syndetic.

- (ii) There exists a subset $B \subseteq A$ that satisfies: for all finite $F \subseteq B$, the set $B \cap \bigcap_{f \in F} (B f)$ is syndetic.
- (iii) The set A belongs to a syndetic, idempotent filter on \mathbb{N} .

The novelty and difficulty of Theorems A and B lie in the third conditions, ones that characterize dynamical syndeticity in terms of membership in a combinatorial object called a syndetic, idempotent filter. This connection is, in fact, behind most of the other results in this paper. We turn our attention toward these objects now.

1.1.3. A new tool: the algebra of idempotent families

A family is an upward-closed collection of sets. A filter is a family \mathscr{F} satisfying: for all members A_1 and A_2 of \mathscr{F} , the set $A_1 \cap A_2$ belongs to \mathscr{F} . Filters are combinatorial objects membership in which formalizes what it means for a set to be "large" in some way. The collection of all neighborhoods of a given point in a topological space is the quintessential example of a filter in topology. We will see that an analogue of this filter for minimal topological systems is inextricably linked to the notion of dynamical syndeticity.

Let us briefly narrow the discussion from filters to ultrafilters. Ultrafilters on \mathbb{N} – maximal filters on \mathbb{N} , whose existence is guaranteed by Zorn's lemma – are combinatorial objects with a deep and important history in the study of topological dynamics, especially in applications to combinatorial number theory [2]. The set $\beta \mathbb{N}$ of ultrafilters – identifiable with the Stone-Čech compactification of \mathbb{N} – is a multifaceted topological and algebraic object, the structure of which is behind many of the aforementioned applications. For the purposes of this introduction, it suffices to know that $\beta \mathbb{N}$ is a compact Hausdorff space, and that there is a natural way to lift addition from \mathbb{N} to $\beta \mathbb{N}$, turning $\beta \mathbb{N}$ into a semigroup: if \mathscr{F} and \mathscr{G} are ultrafilters, then

$$\mathscr{F} + \mathscr{G} := \left\{ A \subseteq \mathbb{N} \mid \left\{ n \in \mathbb{N} \mid A - n \in \mathscr{G} \right\} \in \mathscr{F} \right\}$$
(1.3)

is an ultrafilter. The repeated addition of the principal ultrafilter $\{A \subseteq \mathbb{N} \mid 1 \in A\}$ on the left turns $\beta \mathbb{N}$ in to a (non-metrizable) topological dynamical system.

An ultrafilter \mathscr{F} is called *idempotent* if $\mathscr{F} = \mathscr{F} + \mathscr{F}$. Idempotency of ultrafilters is closely related to proximality in topological dynamics [32, Sec. 19.3] and to finite sums sets in additive combinatorics [32, Sec. 5.2]. An ultrafilter is called *minimal* if it is *uniformly recurrent* (see Section 2.2.1) under the dynamics just described. Minimality of ultrafilters is closely related to uniform recurrence [32, Sec. 19.3] and piecewise syndeticity [32, Sec. 4.4]. At the intersection of all of these ideas are Furstenberg's *central sets* [20, Def. 8.3], combinatorially-rich, dynamically-defined subsets of \mathbb{N} introduced to aid in the topological dynamical approach to problems in Ramsey theory. Bergelson and Hindman [4] used ultrafilters to classify the family of central sets: a set is central if and only if it belongs to a minimal, idempotent ultrafilter.

There is no obstruction to extending the familiar definition of sums of ultrafilters in (1.3) to arbitrary families: if \mathscr{F} and \mathscr{G} are families, the sum $\mathscr{F} + \mathscr{G}$ defined in (1.3) is again a family. But while there are books written about the algebra of ultrafilters (we cite, eg., [32] frequently in this work), the algebra of families appears to be largely unexplored. We define a family \mathscr{F} to be *idempotent* if $\mathscr{F} \subseteq \mathscr{F} + \mathscr{F}$. (See Section 2.1.4 for a discussion on containment versus equality.) This definition generalizes the definition of "idempotent filter"

in Krautzberger [42], the only related reference of which we know.

In the setting of topological dynamics, the canonical example of an idempotent filter is one of the form

$$\mathscr{F} := \uparrow \{ R_T(x, U) \mid U \subseteq X \text{ is open and contains } x \}, \tag{1.4}$$

where x is a point in a dynamical system (X, T) and the upward arrow indicates the upward closure of the collection of sets. That \mathscr{F} is a filter is clear. To see that this is an idempotent filter, one needs only to show and interpret the following: for all open $U \subseteq X$ containing x,

$$R_T(x,U) \subseteq \{n \in \mathbb{N} \mid R_T(x,U) - n \in \mathscr{F}\}.$$

If $n \in R_T(x,U)$, then $x \in T^{-n}U$, whereby $R_T(x,U) - n = R_T(x,T^{-n}U)$ is a member of \mathscr{F} . Since $R_T(x,U) \in \mathscr{F}$, we see from the inclusion that $\{n \in \mathbb{N} \mid R_T(x,U) - n \in \mathscr{F}\} \in \mathscr{F}$, demonstrating that $R_T(x,U) \in \mathscr{F} + \mathscr{F}$. Since every member of \mathscr{F} contains a set of the form $R_T(x,U)$, this shows that $\mathscr{F} \subseteq \mathscr{F} + \mathscr{F}$.

A syndetic filter is a filter whose members are all syndetic. If (X, T) is a minimal system, the filter in (1.4) is an example of a syndetic, idempotent filter. Thus, we see that every dynamically central syndetic set belongs to a syndetic, idempotent filter. Theorem B (iii) gives the converse: every member of a syndetic, idempotent filter is dynamically central syndetic. If ultrafilters are "local" combinatorial objects, then syndetic filters are "global" combinatorial objects. One way to contextualize Theorem B, then, is as a global analogue to the local theorem of Bergelson and Hindman [4] characterizing central sets mentioned above.

There are several other "large," idempotent filters that arise naturally in the subject. All three of the following examples play an important role in the results in this paper. See Section 2 for definitions of the families C^*, \mathcal{IP}^* , and \mathcal{PS}^* (thickly syndetic sets).

- (i) By a theorem of Furstenberg [20, Thm. 9.11], if x is a distal point, the filter defined in (1.4) is a C^* , idempotent filter. Such filters feature prominently in Section 5.2.
- (ii) Very general families of sets of return times in topological dynamics and ergodic theory are \mathcal{IP}^* , idempotent filters. Our demonstration of this in a concrete example in Lemma 5.14 is behind the proof of Theorem H below.
- (iii) The family of thickly syndetic sets is a \mathcal{PS}^* , translation-invariant filter (see Lemma 2.11). This filter is behind some of the main applications of our results, such as in Theorem 5.13.

We believe that these examples and the results in this paper demonstrate the usefulness of the algebra of families of large sets and motivate its further development.

1.1.4. A peek into the proofs of Theorems A and B

Let us briefly discuss the proofs of Theorems A and B. The full proofs, finished in Section 3.4, occupy a substantial portion of Part I of this paper. Since any dynamically syndetic set can be translated to become dynamically central syndetic (Lemma 3.4), Theorem A follows relatively easily from Theorem B.

That the combinatorial condition in Theorem B (ii) characterizes membership in a syndetic, idempotent filter, (iii), is proved in Section 3.2.2. The proof is purely combinatorial and highlights some of the family algebra discussed in the previous section. Readers familiar with the characterization of \mathcal{IP} sets (see Section 2.3.2) as members of idempotent ultrafilters will immediately recognize the ideas at play.

That dynamically central syndetic sets belong to a syndetic, idempotent filter (that (i) implies (iii)) is simple and was discussed in the previous section. The challenge lies in the converse, proved in Section 3.3. Since this is the most technical part of the paper, let us briefly describe the main idea. A more detailed outline is given in Section 3.3.1.

We begin with a set $A \subseteq \mathbb{N}$ that satisfies the combinatorial condition in (ii). Were the point $1_{A\cup\{0\}}$ uniformly recurrent in the full symbolic shift $(\{0,1\}^{\mathbb{N}\cup\{0\}}, \sigma)$, it would follow from Theorem 3.8 that A is dynamically central syndetic. It is not true in general, however, that $1_{A\cup\{0\}}$ is uniformly recurrent. The main idea is to modify the symbolic shift by composing it with a "punch" map to form a related system that we call "shift-punch." Under the shift-punch dynamics, the point $1_{A\cup\{0\}}$ is uniformly recurrent. It is this uniform recurrence, then, that allows us to find a subset of A that is dynamically central syndetic.

1.2. Part II: Dynamically thick sets

Dual to the families of dynamically (central) syndetic sets are the families of dynamically thick sets and sets of pointwise recurrence. A set $A \subseteq \mathbb{N}$ is *dynamically thick* if for all minimal systems (X,T) and all points $x \in X$, the set $\{T^n x \mid n \in A\}$ is dense in X. The set A is a set of (minimal topological) pointwise recurrence if for all minimal systems (X,T), all points $x \in X$, and all open $U \subseteq X$ containing x, there exists $a \in A$ such that $T^a x \in U$.

Thick sets – subsets of \mathbb{N} that contain arbitrary long intervals – are the most immediate examples of dynamically thick sets. It is quick to check that a set is dynamically thick if and only if its complement is not dynamically syndetic. Thus, the complement of the set in (1.1) (and, in fact, the set itself) is an example of a dynamically thick set that is, coincidentally, not thick. Further examples of dynamically thick sets and sets of pointwise recurrence are given in Section 4.1.

1.2.1. Motivation: set and point recurrence

The phenomenon of recurrence in dynamical systems has been studied extensively in numerous forms. We will discuss two of those forms most relevant to this work: set recurrence and point recurrence in ergodic theory and topological dynamics.

In a probability measure space (X, μ) with a measure-preserving transformation $T: X \to X$, a set $E \subseteq X$ of positive measure is said to recur at time $n \in \mathbb{N}$ if $\mu(E \cap T^{-n}E) > 0$. A set $A \subseteq \mathbb{N}$ is called a *set of measurable recurrence* if in every probability measure preserving system, every set of positive measure recurs at some time in A. The question of which times a set of positive measure can be guaranteed to recur – that is, the question of which sets are sets of measurable recurrence – is important both conceptually and for applications. Szemerédi's theorem [49], a cornerstone of additive combinatorics, states that every set of positive integers of positive upper density contains arbitrarily long arithmetic progressions. Furstenberg [19] famously reproved this theorem by connecting the existence of arithmetic progressions in sets of positive upper density to the existence of sets of multiple measurable recurrence. Since then, this connection has been explored and exploited to great effect [51].

Set recurrence in topological dynamics is also well-studied. A set $A \subseteq \mathbb{N}$ is a set of topological recurrence if for all minimal systems (X, T) and all nonempty, open sets $U \subseteq X$, there exists $n \in A$ such that the set $U \cap T^{-n}U$ is nonempty. The connection between topological dynamics and partition combinatorics forged by Furstenberg and Weiss [22] parallels the connection between measurable dynamics and density combinatorics described above and has similarly been influential in additive combinatorics and Ramsey theory.

While the dynamical behavior of points in measurable dynamics is necessarily limited to generic points, meaningful statements can be made about the dynamical behavior of all points in topological dynamics. This is manifest in the "for all $x \in X$ " quantifier in the definitions of dynamically thick sets and sets of pointwise recurrence. Despite the fact that these definitions are naturally analogous to those in the topic of set recurrence, far less has been written on them. Many of our results in this paper are best understood as showing that the nature of pointwise recurrence differs substantially from that of set recurrence.

It is a consequence of a result of Pavlov [47] that a set of pointwise recurrence must have positive upper Banach density. Indeed, he showed that if a set $A \subseteq \mathbb{N}$ has zero upper Banach density $(\limsup_{N\to\infty} \max_{n\in\mathbb{N}} |A \cap \{n,\ldots,n+N\}|/N = 0)$ then the set A is not a set of pointwise recurrence: there is a totally minimal, totally uniquely ergodic, and topologically mixing system (X,T) and uncountably many points $x \in X$ for which $x \notin \{T^n x \mid n \in A\}$. That dynamically thick sets and sets of pointwise recurrence are *piecewise syndetic* (the intersection of a syndetic set and a thick set) are the results of Huang and Ye [37, Thm. 2.4] and Dong, Song, Ye [12, Prop. 4.4], respectively. We improve on both of these results in Theorem G, discussed below.

Pointwise recurrence is implicitly studied in [25, 41] in the context of interpolation sets for minimal systems. A set $I \subseteq \mathbb{N}_0$ is *interpolation set for minimal systems* if for every bounded function $f: I \to \mathbb{C}$, there exists a minimal system (X, T), a point $x \in X$, and a continuous function $F: X \to \mathbb{C}$ such that $f(n) = F(T^n x)$ for all $n \in I$. It was proved in [25, 41] that $I \subseteq \mathbb{N}_0$ is an interpolation set for minimal systems if and only if I is not piecewise syndetic. If $A \subseteq \mathbb{N}$ is not piecewise syndetic, then $I := A \cup \{0\}$ is not piecewise syndetic, so there is minimal system (X, T), a point $x \in X$, and a continuous function $F: X \to \mathbb{C}$ such that

$$F(T^n x) = \begin{cases} 1 \text{ if } n = 0\\ 0 \text{ if } n \in A \end{cases}$$

It follows that A is not a set of pointwise recurrence. Contrapositively, a set of pointwise recurrence must be piecewise syndetic, recovering the results in [12, Prop. 4.4] and, hence, in [37, Thm. 2.4] as well.

As far as we know, the term "set of pointwise recurrence" was first introduced by Host, Kra, and Maass [35], who compared and contrasted several different types of recurrence: single and multiple topological recurrence, topological recurrence for nilsystems, pointwise recurrence, etc.... They observed that the family of sets of pointwise recurrence is properly contained in the family of sets of topological recurrence. We answer in this paper several of the questions they raise.

Finally, the families of dynamically thick sets and sets of pointwise recurrence have appeared explicitly in the literature in the context of disjointness. We discuss these appearances in Section 1.2.3 below.

1.2.2. Combinatorial characterizations of dynamically thick sets

There is a useful and well-known combinatorial characterization for sets of topological recurrence (cf. [8, Thm. 2.4]): a set $A \subseteq \mathbb{N}$ is a set of topological recurrence if and only if for all syndetic sets $S' \subseteq S \subseteq \mathbb{N}$, the set $A \cap (S - S')$ is nonempty. Host, Kra, and Maass [35, Question 2.13] asked whether there is a purely combinatorial characterization of sets of pointwise recurrence. Our next result, proved in Section 4.2, gives an answer to this question.

Theorem C. Let $A \subseteq \mathbb{N}$. The following are equivalent.

- (i) A is a set of pointwise recurrence.
- (ii) For all $B \supseteq A$, there exists a finite set $F \subseteq \mathbb{N} \setminus B$ such that the set $B \cup (B F)$ is thick.
- (iii) For all syndetic $S \subseteq \mathbb{N}$, there exists a finite set $F \subseteq A$ such that for all syndetic $S' \subseteq S$,

$$F \cap (S - S') \neq \emptyset.$$

The statement in (ii) is derived from the combinatorial characterization of dynamically central syndetic sets in Theorem B and can be used to produce examples of sets of pointwise recurrence. We demonstrate this in the analogous context of dynamically thick sets below. The statement in (iii) is best understood as a kind of quantitative strengthening of the combinatorial characterization of sets of topological recurrence given at the beginning of this section.

The following is an analogue of Theorem C for dynamically thick sets and is proved in Section 4.2.

Theorem D. Let $A \subseteq \mathbb{N}$. The following are equivalent.

- (i) A is dynamically thick.
- (ii) For all $\mathbb{N} \supseteq B \supseteq A$, there exists a finite set $F \subseteq \mathbb{N} \setminus B$ such that the set B F is thick.
- (iii) For all piecewise syndetic $S \subseteq \mathbb{N}$, there exists a finite set $F \subseteq A$ such that the set S F is thick.

In [24, Question 9.6], Glasner, Tsankov, Weiss and Zucker call for an explicit characterization of dynamically thick sets (they call them *dense orbit sets*) in countable, discrete groups. Theorem D gives an answer to their call for subsets of \mathbb{N} . Kennedy, Raum, and Salomon [39] also provide an answer by translating their characterization of dynamically syndetic sets into one for dynamically thick sets. Because our characterization of dynamically syndetic sets provides a simpler-to-verify sufficient condition (as discussed in Section 1.1.2), Theorem D reveals more about dynamically thick subsets of \mathbb{N} .

The statement in Theorem D (ii) can be used to produce novel examples of dynamically thick sets, such as

$$A := \bigcup_{i=1}^{\infty} \left((p_i \mathbb{N} + 1) \cap H_i \right), \tag{1.5}$$

where $(p_i)_{i \in \mathbb{N}}$ is a sequence of distinct primes and $(H_i)_{i \in \mathbb{N}}$ is a sequence of thick sets. Heuristically, the set A is dynamically thick because it contains sufficiently long pieces of return-time sets from a sufficiently rich collection of systems, in this case, rotations on prime many points. Theorem 4.12 makes this heuristic precise and allows us to generalize the example in (1.5) by showing that no minimal topological dynamical system can be disjoint from an infinite collection of disjoint systems. A collection of further examples of dynamically thick sets is given in Section 4.3.

The dynamically thick set A in (1.5) allows us to answer another question posed by Host, Kra, and Maass [35, Question 3.11]. They asked whether every set of pointwise recurrence for distal systems is an \mathcal{IP} set (see Sections 2.2.1 and 2.3.2 for the requisite definitions). If the thick sets H_i are chosen to be sufficiently spaced (this is clarified in Lemma 4.9), the set A is not an \mathcal{IP} set. It is still, however, dynamically thick – and, hence, a set of pointwise recurrence – giving a negative answer to the question. (Their question was first answered in [41], where an example of a dynamically thick set for distal systems that does not contain the configuration $\{x, y, x + y\}$ is shown.)

1.2.3. Dynamical thickness in the context of disjointness

The notions of pointwise recurrence and dynamical thickness are linked to the topics of product recurrence and disjointness, respectively. We will point out where these topics have appeared together in the literature and what our results have to say about them.

A point x in a dynamical system (X, T) is *recurrent* if for all open $U \subseteq X$ containing x, the set $R_T(x, U)$ is nonempty. The following, then, is immediate from the definitions: for all open $U \subseteq X$ containing x, the set $R_T(x, U)$ is a set of pointwise recurrence if and only if for all minimal system (Y, S) and all $y \in Y$, the point (x, y) is recurrent in the product system $(X \times Y, T \times S)$. Dong, Shao, and Ye [12] call points x with this property \mathcal{F}_s -PR points. Distal points (see Example 5.2) are examples of \mathcal{F}_s -PR points, as are transitive points in the full shift (see below).

Given two dynamically central syndetic sets A and B, there exist minimal systems (X,T)and (Y,S), points $x \in X$ and $y \in Y$, and open sets $U \subseteq X$ and $Y \subseteq Y$ containing x and y, respectively, such that $R_T(x,U) \subseteq A$ and $R_S(y,V) \subseteq B$. If A and B are disjoint, then $R_T(x,U) \cap R_S(y,V) = \emptyset$. In this case, the point (x,y) is not recurrent in the product system $(X \times Y, T \times S)$. We show in the following theorem, proved in Section 3.1.4, that such pairs of sets exist in abundance.

Theorem E. Every dynamically central syndetic set can be partitioned into infinitely many, pairwise disjoint dynamically central syndetic sets.

While Theorem E concerns dynamically central syndetic sets, its consequences for the family of sets of pointwise recurrence are most relevant to this discussion. Host, Kra, and Maass [35, Question 6.5] asked whether or not the family of sets of pointwise recurrence is partition regular. A strong negative answer, given in Theorem 4.3, follows immediately from Theorem E: if $\mathbb{N} = A \cup B$, where A and B are disjoint, dynamically central syndetic sets, then \mathbb{N} has been partitioned into two sets, neither of which is a set of pointwise recurrence. It is well-known that sets of topological and measurable recurrence are partition regular, so Theorem E offers yet another contrast between the nature of set and pointwise recurrence.

A joining of two topological dynamical systems (X, T) and (Y, S) is a nonempty, closed set $Z \subseteq X \times Y$ that satisfies: $(T \times S)Z \subseteq Z$, $\pi_X Z = X$, and $\pi_Y Z = Y$, where π_X, π_Y are the projections from $X \times Y$ to X and Y, respectively. Furstenberg [18, Thm. II.2] defined two topological systems (X, T) and (Y, S) to be *disjoint* if their only joining is $X \times Y$.

To relate the notions of disjointness and dynamical thickness, let (X, T) be a system and $x \in X$ be a point with a dense orbit. The following is simple to verify from the definitions: the system (X,T) is disjoint from every minimal system if and only if for all nonempty, open $U \subseteq X$, the set $R_T(x, U)$ is dynamically thick. It is interesting to compare this to the characterization of recurrent points in the product $X \times Y$ given above.

Furstenberg proved [18, Thm. II.2] that the full shift $(\{0,1\}^{\mathbb{N}},\sigma)$ is disjoint from every

minimal system. This result spurred a number of works in which the notion of dynamical thickness arises either directly or indirectly [12, 24, 36, 37, 39]. We provide a quick explanation of Furstenberg's result in Remark 4.10 by combining the characterization in the previous paragraph with the example in (1.5) above.

This connection between dynamical thickness and disjointness; examples of dynamically thick sets such as the one in (1.5) and its generalizations in Section 4.3; and the call from [24, Question 9.6] led us to wonder whether or not there is a "structure theorem" for dynamically thick sets. Drawing on the results from Part III of this paper, we are able to offer the following description of an arbitrary dynamically thick set, proved in Section 4.3.3. A collection \mathscr{C} of subsets of \mathbb{N} is *robustly syndetic* if for all dynamically syndetic sets $A \subseteq \mathbb{N}$, there exists $B \in \mathscr{C}$ such that the set $A \cap B$ is syndetic.

Theorem F. Let $A \subseteq \mathbb{N}$. The set A is dynamically thick if and only if there exist a robustly syndetic collection \mathscr{C} of subsets of \mathbb{N} and, for each $B \in \mathscr{C}$, a thick set $H_B \subseteq \mathbb{N}$ such that

$$A = \bigcup_{B \in \mathscr{C}} (B \cap H_B).$$
(1.6)

In all of the examples of dynamically thick sets we know, there is much more that can be said about the robustly syndetic collection \mathscr{C} . More needs to be said: we could not even determine using Theorem F whether or not a dynamically thick set can be partitioned into two dynamically thick sets. We speculate more precisely in Section 6.2 on how Theorem F might be improved.

1.3. Part III: Dynamically piecewise syndetic sets

In the last part of this paper, we introduce and study the family of sets formed by intersections of dynamically (central) syndetic and dynamically thick sets (sets of pointwise recurrence). For sets B and C of positive integers, the set $B \cap C$ is called

- dynamically piecewise syndetic if B is dynamically syndetic and C is dynamically thick, and
- dynamically central piecewise syndetic if B is dynamically central syndetic and C is a set of pointwise recurrence.

To our knowledge, the families of dynamically (central) piecewise syndetic sets have not appeared before in the literature. As we discuss next, their study is motivated by the wellestablished notions of piecewise syndeticity and centrality.

1.3.1. Motivation: combinatorial and dynamical piecewise syndeticity

Syndeticity and thickness are *dual*: for every syndetic set A and every thick set B, the intersection $A \cap B$ is nonempty. As mentioned in Section 1.2.1, sets of this form are called piecewise syndetic. Piecewise syndeticity is a "local" analogue of syndeticity that has the advantage of being partition regular: if the union of two sets is piecewise syndetic, one of the sets must be piecewise syndetic. The family of piecewise syndetic sets and the subfamily of central sets (mentioned in Section 1.1.3) are important to the connection between combinatorics and dynamics (via correspondence principles) and to describing algebraic structure in the Stone-Čech compactification of \mathbb{N} .

The families of dynamically syndetic sets and dynamically thick sets are dual in the same

sense, motivating the study of their intersections. By definition, then, dynamically piecewise syndetic sets are nonempty, but what more can be said? The same question applies equally well to the family of dynamically central piecewise syndetic sets. What are these families' relationships to each other and to the families of piecewise syndetic and central sets? In addressing these questions, we generalize some of the aforementioned results on dynamically syndetic and thick sets in the literature and and give some novel combinatorial applications.

1.3.2. Dynamically piecewise syndetic sets are piecewise syndetic

Because the families of dynamically (central) piecewise syndetic sets have not been considered before, our first results concern the relationships between these families and with the families of piecewise syndetic and central sets. Looking ahead, Fig. 2 at the end of Section 2 summarizes the relationships between the families studied in this paper with the established families of subsets of \mathbb{N} in the topic.

There are dynamically syndetic sets that are not dynamically central syndetic, and there are sets of pointwise recurrence that are not dynamically thick. Therefore, from the definitions, there is no apparent relationship between the families of dynamically piecewise syndetic and dynamically central piecewise syndetic sets. As our terminology suggests, however, we prove that dynamically central piecewise syndetic sets are dynamically piecewise syndetic.

We could not find a proof of this fact that does not also elucidate the relationship between the families of dynamically piecewise syndetic and piecewise syndetic sets. We show that, perhaps surprisingly, these families are the same. The following theorem summarizes these relationships and is proved in Section 5.2.1.

Theorem G. Denote by C, \mathcal{PS} , $d\mathcal{PS}$, and $dc\mathcal{PS}$ the families of central, piecewise syndetic, dynamically piecewise syndetic, and dynamically central piecewise syndetic sets, respectively. The following inclusions hold:

$$\mathcal{C} \subsetneq dc \mathcal{PS} \subsetneq d\mathcal{PS} = \mathcal{PS}.$$

In particular, a set is dynamically piecewise syndetic if and only if it is piecewise syndetic.

That dynamically piecewise syndetic sets are piecewise syndetic offers a striking contrast to the analogy with set recurrence given in Section 1.2.1. Indeed, sets of measurable recurrence – and thus, in particular, the intersection of a set of measurable recurrence with the set of times that a particular set of positive measure recurs – are not necessarily even of positive Banach density (and, hence, are not piecewise syndetic). By a result of Furstenberg [20] and Sárközy [48], the set of perfect squares, for example, is such a set of measurable recurrence.

We discussed in Section 1.2.1 how the results in [12, Prop. 4.4], [25], and [37, Thm. 2.4] show that the families of dynamically thick sets and sets of pointwise recurrence are piecewise syndetic. Theorem G generalizes these results by showing that the families of dynamically (central) piecewise syndetic sets are piecewise syndetic. It follows by the abstract family results in Lemma 2.1 that a set of pointwise recurrence modified on a non-piecewise syndetic set is still a set of pointwise recurrence. The same is true for all of the new families described in this paper; see Theorem 5.13.

There are two keys to the proof of Theorem G. The first is our characterization of dynamically central syndetic sets as members of syndetic, idempotent filters in Theorem B, and the second is a collection of abstract results on large, idempotent filters (see, eg., Theorem 5.7).

1.3.3. Sets of polynomial recurrence and combinatorial configurations in dynamically central piecewise syndetic sets

A set $A \subseteq \mathbb{N}$ is called a *set of multiple topological recurrence* if for all $k \in \mathbb{N}$, all minimal systems (X, T), and all nonempty, open sets $U \subseteq X$, there exists $n \in A$ such that

$$U \cap T^{-n}U \cap \dots \cap T^{-kn}U \neq \emptyset.$$

As with the sets of topological recurrence defined in Section 1.2.1, understanding which sets are sets of multiple topological recurrence leads to various applications, for example, generalizations of the classical van der Waerden's theorem [50]. While it is known that a set of topological recurrence is not necessarily a set of multiple topological recurrence [17], Host, Kra, and Maass [35, Question 2.14] asked whether or not a set of pointwise recurrence is a set of multiple topological recurrence.

We show in the next theorem a strong positive answer: dynamically central piecewise syndetic sets (and, hence, sets of pointwise recurrence) are sets of polynomial multiple measurable recurrence for commuting transformations. A set $A \subseteq \mathbb{N}$ is a set of polynomial multiple measurable recurrence for commuting transformations if for all $k, \ell \in \mathbb{N}$, all commuting measure preserving systems $(X, \mu, T_1, \ldots, T_k)$ (see Section 5.3), all $E \subseteq X$ with $\mu(E) > 0$, and, all polynomials $p_{i,j} \in \mathbb{Q}[x]$ with $p_i(0) = 0$ and $p_i(\mathbb{N}_0) \subseteq \mathbb{N}_0$, $1 \leq i \leq k$, $1 \leq j \leq \ell$, there exists $n \in A$ such that

$$\mu\Big(\big(T_1^{p_{1,1}(n)}T_2^{p_{2,1}(n)}\cdots T_k^{p_{k,1}(n)}\big)^{-1}E\cap\cdots\cap\big(T_1^{p_{1,\ell}(n)}T_2^{p_{2,\ell}(n)}\cdots T_k^{p_{k,\ell}(n)}\big)^{-1}E\Big)>0.$$

Because any minimal topological dynamical system admits an invariant measure of full support, sets of polynomial multiple measurable recurrence for commuting transformations are sets of multiple topological recurrence. Therefore, the following theorem answers [35, Question 2.14] positively.

Theorem H. Dynamically central piecewise syndetic sets are sets of polynomial multiple measurable recurrence for commuting transformations.

The proof of Theorem H in Section 5.3 appeals to the \mathcal{IP} polynomial Szemerédi theorem of Bergelson and McCutcheon [9] to show that the family of times of polynomial returns in ergodic theory is an \mathcal{IP}^* , idempotent filter (see Lemma 5.14). Theorem B is then invoked to prove that every member of this \mathcal{IP}^* , idempotent filter intersects all dynamically central piecewise syndetic sets.

Combining Theorem H with the fact that dynamically central piecewise syndetic sets are piecewise syndetic (Theorem G), we show in Theorem 5.17 that every such set contains Brauer-type polynomial configurations. The extent to which dynamically central piecewise syndetic sets are combinatorially rich remains an interesting open question that we speculate on in Section 6.3.

1.4. Organization of the paper

Notation, terminology, and preliminaries are given in Section 2, including new material on the algebra of families of subsets of \mathbb{N} . The main body of work in this paper is then divided into three parts, which comprise Sections 3, 4, and 5, respectively:

- Part I: Dynamically syndetic sets, in which we prove Theorems A, B, and E;

- Part II: Dynamically thick sets, in which we prove Theorems C, D, and F;

- Part III: Dynamically piecewise syndetic sets, in which we prove Theorems G and H. We conclude the paper with a collection of open problems and directions in Section 6. A map of the main results and their logical dependencies is drawn in Fig. 1.



Figure 1: A collection of results in this paper and the logical dependencies between them. Theorems stated in the introduction are in bold. The families of dynamically (central) syndetic, dynamically thick (sets of pointwise recurrence), and dynamically (central) piecewise syndetic sets are denoted by dS (dcS), dT (dcT), and dPS (dcPS), respectively. The acronym "SIF" stands for "syndetic, idempotent filter."

1.5. Acknowledgements

Thanks goes to Andy Zucker and Josh Frisch for helpful discussions at the 2024 Southeastern Logic Symposium Conference and for the references they provided. Thanks also goes to Neil Hindman for correspondence regarding translates of central sets. Last but not least, thanks goes to Andreas Koutsogiannis, Joel Moreira, Florian Richter, and Donald Robertson, with whom the authors were working when a number of the ideas in this paper sharpened.

2. Preliminaries

The set of integers, non-negative integers, and positive integers are denoted by \mathbb{Z} , \mathbb{N}_0 , and \mathbb{N} , respectively. The power set of \mathbb{N} is denoted $\mathcal{P}(\mathbb{N})$. For $N \in \mathbb{N}_0$, we define [N] to be $\{0, 1, \ldots, N-1\}$.

For $A \subseteq \mathbb{N}$ and $n \in \mathbb{N}$, we define

$$\begin{split} A+n &:= \{a+n \mid a \in A\}, \\ A-n &:= \{m \in \mathbb{N} \mid m+n \in A\}, \end{split} \qquad \begin{array}{l} nA &:= \{na \mid a \in A\}, \\ A/n &:= \{m \in \mathbb{N} \mid mn \in A\}. \end{split}$$

For $A, B \subseteq \mathbb{N}$, we define

$$A + B := \bigcup_{b \in B} (A + b), \qquad \qquad A - B := \bigcup_{b \in B} (A - b).$$

We follow the convention that the empty union of subsets of \mathbb{N} is empty while the empty intersection of subsets of \mathbb{N} is equal to \mathbb{N} . Note that all set operations on subsets of \mathbb{N} result in subsets of \mathbb{N} .

2.1. Families of positive integers

Let \mathscr{F} be a collection of subsets of \mathbb{N} . The *upward closure* of \mathscr{F} is

$$\uparrow \mathscr{F} := \left\{ B \subseteq \mathbb{N} \mid \exists A \in \mathscr{F}, \ B \supseteq A \right\}$$

The set \mathscr{F} is upward-closed if $\mathscr{F} = \uparrow \mathscr{F}$. We will call an upward-closed collection of subsets of \mathbb{N} a family. Note that both \emptyset and $\mathcal{P}(\mathbb{N})$ are families. A family is proper if it is neither \emptyset nor $\mathcal{P}(\mathbb{N})$, that is, it is both nonempty and does not contain the empty set.

2.1.1. Dual families and intersections

The *dual* of a family \mathscr{F} of subsets of \mathbb{N} is

$$\mathscr{F}^* := \{ B \subseteq \mathbb{N} \mid \forall A \in \mathscr{F}, \ B \cap A \neq \emptyset \},\$$

the collection of all those sets which have nonempty intersection with all elements of \mathscr{F} . The following facts are quick to check and will be used without mention:

- (i) $(\mathscr{F}^*)^* = \mathscr{F}$, whereby it makes sense to say that \mathscr{F} and \mathscr{F}^* are dual;
- (ii) the empty family Ø and P(N) are dual, while the dual of a proper family is a proper family;
- (iii) $A \in \mathscr{F}$ if and only if $\mathbb{N} \setminus A \notin \mathscr{F}^*$;
- (iv) $\mathscr{F} \subseteq \mathscr{G}$ if and only if $\mathscr{G}^* \subseteq \mathscr{F}^*$.

For nonempty families \mathscr{F} and \mathscr{G} , we define

$$\mathscr{F} \sqcap \mathscr{G} := \{A \cap B \mid A \in \mathscr{F} \text{ and } B \in \mathscr{G}\}.$$

We will adopt the convention that $\emptyset \sqcap \mathscr{F} = \mathscr{F} \sqcap \emptyset = \mathscr{F}$ for all families \mathscr{F} . The following facts are quick to check for all families \mathscr{F} and \mathscr{G} :

(v) the collection $\mathscr{F} \sqcap \mathscr{G}$ is a family that contains both \mathscr{F} and \mathscr{G} (despite what is suggested by the notation);

- (vi) if \mathscr{F} is a proper family, then $\mathscr{F} \sqcap \mathscr{F}^*$ is a proper family;
- (vii) if $\mathscr{F}_1 \subseteq \mathscr{G}_1$ and $\mathscr{F}_2 \subseteq \mathscr{G}_2$, then $\mathscr{F}_1 \sqcap \mathscr{G}_1 \subseteq \mathscr{F}_2 \sqcap \mathscr{G}_2$; in particular, if $\mathscr{F} \subseteq \mathscr{G}$, then $\mathscr{F} \sqcap \mathscr{G} = \mathscr{G}$.

2.1.2. Filters, partition regularity, and ultrafilters

There are two complementary notions of largeness for families that feature prominently in this topic: that of being partition regular and that of being a filter. A family \mathscr{F} is *partition* regular if for all $A \in \mathscr{F}$ and all finite partitions $A = A_1 \cup \cdots \cup A_k$, some piece A_i belongs to \mathscr{F} . The family \mathscr{F} is a *filter* if for all $A_1, A_2 \in \mathscr{F}$, the set $A_1 \cap A_2 \in \mathscr{F}$. These notions are dual in the sense of fact (i) below.

If P is a property of families, we call a family that has property P and that is also a filter a P filter. For example, we will consider proper filters, partition regular filters, translationinvariant filters, syndetic filters, and so on. Note, for example, that \emptyset and $\mathcal{P}(\mathbb{N})$ are both partition regular filters.

The following facts are quick to check:

- (i) a family \mathscr{F} is partition regular if and only if the dual family \mathscr{F}^* is a filter;
- (ii) if \mathscr{F} is a proper filter, then $\mathscr{F} \subseteq \mathscr{F}^*$;
- (iii) if \mathscr{F} and \mathscr{G} are filters, then $\mathscr{F} \sqcap \mathscr{G}$ is the smallest filter containing both \mathscr{F} and \mathscr{G} .

Given a family \mathscr{F} , we will frequently consider the family $\mathscr{F} \sqcap \mathscr{F}^*$. The follow lemma records some useful facts about it.

Lemma 2.1. Let \mathscr{F} be a family and define $\mathscr{G} := \mathscr{F} \sqcap \mathscr{F}^*$. The family \mathscr{G} is partition regular and its dual, \mathscr{G}^* , is a filter. Moreover, for $A \subseteq \mathbb{N}$, the following are all equivalent to the set A belonging to \mathscr{G}^* :

- (i) for all $B \in \mathscr{F}$, the set $A \cap B \in \mathscr{F}$;
- (ii) for all $B \in \mathscr{F}^*$, the set $A \cap B \in \mathscr{F}^*$;
- (iii) for all $B \in \mathscr{G}$, the set $A \cap B \in \mathscr{G}$;
- (iv) for all $B \in \mathscr{G}^*$, the set $A \cap B \in \mathscr{G}^*$.

Proof. If \mathscr{F} is not proper, then $\mathscr{G} = \mathcal{P}(\mathbb{N})$, which is partition regular. Otherwise, the partition regularity of \mathscr{G} is shown in [10, Prop. 2.5 (h)]. That its dual, \mathscr{G}^* , is a filter is mentioned as item (i) before the statement of this lemma.

We will show that (i) is equivalent to A belonging \mathscr{G}^* ; the other statements follow analogously and are left to the reader. Suppose $A \in \mathscr{G}^*$ and $B \in \mathscr{F}$. To see that $A \cap B \in \mathscr{F}$, we will show that for all $C \in \mathscr{F}^*$, the set $A \cap B \cap C$ is nonempty. Let $C \in \mathscr{F}^*$. The set $B \cap C \in \mathscr{G}$, and since $A \in \mathscr{G}^*$, we have that $A \cap B \cap C \neq \emptyset$, as desired.

Conversely, suppose that (i) holds. To see that $A \in \mathscr{G}^*$, we will show that for all $D \in \mathscr{G}$, the set $A \cap D \neq \emptyset$. Let $D \in \mathscr{G}$. There exists $B \in \mathscr{F}$ and $C \in \mathscr{F}^*$ such that $D = B \cap C$. Then $A \cap D = A \cap B \cap C$. By assumption, $A \cap B \in \mathscr{F}$, whereby $A \cap B \cap C \neq \emptyset$, as desired. \Box

A proper family is called an *ultrafilter* if it satisfies one (equivalently, all) of the properties in the following lemma. This lemma is well-known (cf. [32, Thm. 3.6]), but its proof is short, so we provide it here.

Lemma 2.2. Let \mathscr{F} be a proper family of subsets of \mathbb{N} . The following are equivalent:

(i) the family \mathscr{F} is a proper filter and is maximal (by containment) amongst proper filters;

- (ii) the family \mathscr{F} is self-dual, i.e., $\mathscr{F} = \mathscr{F}^*$;
- (iii) the family \mathscr{F} is partition regular and a filter.

Proof. (i) \implies (ii) Suppose \mathscr{F} is a proper filter and is maximal amongst proper filters. Since \mathscr{F} is a proper filter, we have that $\mathscr{F} \subseteq \mathscr{F}^*$. Let $B \in \mathscr{F}^*$. It is quick to check that $\mathscr{G} := \{A \cap B \mid A \in \mathscr{F}\}$ is a proper filter that contains \mathscr{F} . Since \mathscr{F} is maximal, we have that $\mathscr{F} = \mathscr{G}$, whereby $B \in \mathscr{F}$. Since $B \in \mathscr{F}^*$ was arbitrary, we have that $\mathscr{F} = \mathscr{F}^*$.

(ii) \implies (i) Suppose that \mathscr{F} is self-dual, and let \mathscr{G} be a proper filter containing \mathscr{F} . Since \mathscr{G} is a proper filter, we have that $\mathscr{G} \subseteq \mathscr{G}^*$. Since $\mathscr{F} \subseteq \mathscr{G}$, we have that $\mathscr{G}^* \subseteq \mathscr{F}^* = \mathscr{F} \subseteq \mathscr{G} \subseteq \mathscr{G}^*$. Therefore, $\mathscr{F} = \mathscr{G}$, demonstrating that \mathscr{F} is maximal, as desired.

(ii) \implies (iii) Suppose $\mathscr{F} = \mathscr{F}^*$. By Lemma 2.1, the family $\mathscr{F} \sqcap \mathscr{F}^*$ is partition regular, and hence $(\mathscr{F} \sqcap \mathscr{F}^*)^*$ is a filter. But $\mathscr{F} = \mathscr{F}^*$ implies that $\mathscr{F} \sqcap \mathscr{F}^* = \mathscr{F}$ and $(\mathscr{F} \sqcap \mathscr{F}^*)^* = \mathscr{F}^* = \mathscr{F}$, so \mathscr{F} is both partition regular and a filter.

(iii) \implies (ii) Suppose \mathscr{F} is a proper filter that is partition regular. Since \mathscr{F} is proper, we have that $\mathscr{F} \subseteq \mathscr{F}^*$. Since \mathscr{F} is proper and partition regular, \mathscr{F}^* is a proper filter, whereby $\mathscr{F}^* \subseteq (\mathscr{F}^*)^* = \mathscr{F}$. Thus, $\mathscr{F} = \mathscr{F}^*$, as desired.

Ultrafilters exist by a standard application of Zorn's Lemma. The space of ultrafilters on \mathbb{N} is an important object that we will make use of in this paper. We delay further discussion until Section 2.2.3.

2.1.3. Translation algebra

For $A \subseteq \mathbb{N}$ and families \mathscr{F} and \mathscr{G} of subsets of \mathbb{N} , we define

$$A - \mathscr{F} := \{ n \in \mathbb{N} \mid A - n \in \mathscr{F} \}, \\ \mathscr{F} + \mathscr{G} := \{ B \subseteq \mathbb{N} \mid B - \mathscr{G} \in \mathscr{F} \}.$$

Thus, we see that $n \in A - \mathscr{F}$ if and only if $A - n \in \mathscr{F}$ and that $B \in \mathscr{F} + \mathscr{G}$ if and only $B - \mathscr{G} \in \mathscr{F}$. It is simple to check that $\mathscr{F} + \mathscr{G}$ is a family of subsets of \mathbb{N} .

The definition of $\mathscr{F} + \mathscr{G}$ appears for filters in [42] and, when \mathscr{F} and \mathscr{G} are ultrafilters, agrees with the usual definition of sums of ultrafilters (cf. [32, Thm. 4.12]). Family sums will appear throughout the paper. It is useful to note that, loosely speaking, the containment $\mathscr{F} \subseteq \mathscr{G} + \mathscr{H}$ means: for all $A \in \mathscr{F}$, there are \mathscr{G} many positive integers n for which the set A - n belongs to \mathscr{H} .

The following lemma records a number of algebraic facts that will be useful later on.

Lemma 2.3. Let $n \in \mathbb{N}$, $A, B \subseteq \mathbb{N}$, and $\mathscr{F}, \mathscr{G}, \mathscr{F}_1, \mathscr{F}_2, \mathscr{G}_1, \mathscr{G}_2$ be families.

- (i) If \mathscr{F} and \mathscr{G} are proper, then $\mathscr{F} + \mathscr{G}$ is proper.
- (ii) $(A \mathscr{F}) \cap (B \mathscr{G}) \subseteq (A \cap B) (\mathscr{F} \sqcap \mathscr{G})$, with equality if $\mathscr{F} = \mathscr{G}$.
- (iii) If \mathscr{F} is a filter, then $\mathscr{F} + \mathscr{G}$ is a filter.
- (iv) $(A-n) \mathscr{F} = (A \mathscr{F}) n.$
- (v) $A (\mathscr{F} + \mathscr{G}) = (A \mathscr{G}) \mathscr{F}.$
- (vi) If $\mathscr{F} \subseteq \mathscr{G}$, then $A \mathscr{F} \subseteq A \mathscr{G}$.
- (vii) If $\mathscr{F}_1 \subseteq \mathscr{G}_1$ and $\mathscr{F}_2 \subseteq \mathscr{G}_2$, then $\mathscr{F}_1 + \mathscr{F}_2 \subseteq \mathscr{G}_1 + \mathscr{G}_2$.
- $\begin{array}{l} (\text{viii}) \ (\mathscr{F} + \mathscr{G})^* = \mathscr{F}^* + \mathscr{G}^*.\\ (\text{ix}) \ (\mathscr{F}_1 + \mathscr{F}_2) \sqcap (\mathscr{G}_1 + \mathscr{G}_2) \subseteq (\mathscr{F}_1 \sqcap \mathscr{G}_1) + (\mathscr{F}_2 \sqcap \mathscr{G}_2). \end{array}$

Proof. (i) Suppose \mathscr{F} and \mathscr{G} are proper. Since $\emptyset \notin \mathscr{G}$, we have $\emptyset - \mathscr{G} = \emptyset$. Since $\emptyset \notin \mathscr{F}$, we have $\emptyset \notin \mathscr{F} + \mathscr{G}$. Since \mathscr{F} and \mathscr{G} both contain \mathbb{N} , we see that $\mathbb{N} \in \mathscr{F} + \mathscr{G}$. Since $\mathscr{F} + \mathscr{G}$ does not contain \emptyset but contains \mathbb{N} , it is proper.

(ii) Suppose $n \in (A - \mathscr{F}) \cap (B - \mathscr{G})$. Since $A - n \in \mathscr{F}$ and $B - n \in \mathscr{G}$, we have that $(A \cap B) - n = (A - n) \cap (B - n) \in \mathscr{F} \sqcap \mathscr{G}$. Thus, $n \in (A \cap B) - (\mathscr{F} \sqcap \mathscr{G})$, as desired.

If $\mathscr{F} = \mathscr{G}$, then $\mathscr{F} \sqcap \mathscr{G} = \mathscr{F}$. We wish to show in this case that $(A \cap B) - \mathscr{F} \subseteq (A - \mathscr{F}) \cap (B - \mathscr{F})$. Suppose $n \in (A \cap B) - \mathscr{F}$ so that $(A - n) \cap (B - n) \in \mathscr{F}$. Since \mathscr{F} is upward closed, we have that $A - n \in \mathscr{F}$ and $B - n \in \mathscr{F}$. Therefore, $n \in (A - \mathscr{F}) \cap (B - \mathscr{F})$, as desired.

(iii) Suppose \mathscr{F} is a filter. Since $\mathscr{F} + \mathscr{G}$ is a family, we need only to show that for all $A, B \in \mathscr{F} + \mathscr{G}$, the set $A \cap B \in \mathscr{F} + \mathscr{G}$. Let $A, B \in \mathscr{F} + \mathscr{G}$. Thus, $A - \mathscr{G}, B - \mathscr{G} \in \mathscr{F}$, whereby $(A - \mathscr{G}) \cap (B - \mathscr{G}) \in \mathscr{F}$. By (ii), it follows that $(A \cap B) - \mathscr{G} \in \mathscr{F}$, whereby $A \cap B \in \mathscr{F} + \mathscr{G}$, as desired.

(iv) Note that $m \in (A - n) - \mathscr{F}$ if and only if $A - n - m \in \mathscr{F}$, which happens if and only if $A - (m + n) \in \mathscr{F}$. We see that this happens if and only if $m + n \in A - \mathscr{F}$, that is, if and only if $m \in (A - \mathscr{F}) - n$, as desired.

(v) We see that $n \in A - (\mathscr{F} + \mathscr{G})$ if and only if $A - n \in \mathscr{F} + \mathscr{G}$, which happens if and only if $(A - n) - \mathscr{G} \in \mathscr{F}$. By (iv), this happens if and only if $(A - \mathscr{G}) - n \in \mathscr{F}$, which happens if and only if $n \in (A - \mathscr{G}) - \mathscr{F}$, as desired.

(vi) If $n \in A - \mathscr{F}$, then $A - n \in \mathscr{F} \subseteq \mathscr{G}$, whereby $n \in A - \mathscr{G}$.

(vii) Suppose $A \in \mathscr{F}_1 + \mathscr{F}_2$. Then $A - \mathscr{F}_2 \in \mathscr{F}_1 \subseteq \mathscr{G}_1$. Since $\mathscr{F}_2 \subseteq \mathscr{G}_2$, we have that $A - \mathscr{F}_2 \subseteq A - \mathscr{G}_2$. Since \mathscr{G}_1 is upward closed and $A - \mathscr{F}_2 \in \mathscr{G}_1$, we see that $A - \mathscr{G}_2 \in \mathscr{G}_1$. Therefore, $A \in \mathscr{G}_1 + \mathscr{G}_2$, as desired.

(viii) We will show first that $\mathscr{F}^* + \mathscr{G}^* \subseteq (\mathscr{F} + \mathscr{G})^*$. Suppose $A \in \mathscr{F}^* + \mathscr{G}^*$, and let $B \in \mathscr{F} + \mathscr{G}$. Since $A - \mathscr{G}^* \in \mathscr{F}^*$ and $B - \mathscr{G} \in \mathscr{F}$, we have that there exists $n \in (A - \mathscr{G}^*) \cap (B - \mathscr{G})$. We see that $A - n \in \mathscr{G}^*$ and $B - n \in \mathscr{G}$. It follows that $(A - n) \cap (B - n) \neq \emptyset$, whereby $A \cap B \neq \emptyset$, as desired.

Now we can write

$$\mathscr{F}^* + \mathscr{G}^* \subseteq (\mathscr{F} + \mathscr{G})^* = \left((\mathscr{F}^*)^* + (\mathscr{G}^*)^* \right)^* \subseteq \left((\mathscr{F}^* + \mathscr{G}^*)^* \right)^* = \mathscr{F}^* + \mathscr{G}^*,$$

where the first containment follows from the previous paragraph and the second containment follows from the previous paragraph applied to the dual classes. This shows the desired equality.

(ix) Suppose $\mathscr{F}_1 + \mathscr{F}_2 = \emptyset$ or $\mathscr{G}_1 + \mathscr{G}_2 = \emptyset$. If $\mathscr{F}_1 + \mathscr{F}_2 = \emptyset$, then either $\mathscr{F}_1 = \emptyset$ or $\mathscr{F}_2 = \emptyset$. In the first case, the desired inclusion simplifies to $\mathscr{G}_1 + \mathscr{G}_2 \subseteq \mathscr{G}_1 + (\mathscr{F}_2 \sqcap \mathscr{G}_2)$, which follows from (vii). The second case is shown similarly. The case that $\mathscr{G}_1 + \mathscr{G}_2 = \emptyset$ is handled analogously.

Otherwise, we have that $\mathscr{F}_1 + \mathscr{F}_2 \neq \emptyset$ and $\mathscr{G}_1 + \mathscr{G}_2 \neq \emptyset$. To see the desired inclusion, let $C \in (\mathscr{F}_1 + \mathscr{F}_2) \sqcap (\mathscr{G}_1 + \mathscr{G}_2)$. There exists $A \in \mathscr{F}_1 + \mathscr{F}_2$ and $B \in \mathscr{G}_1 + \mathscr{G}_2$ such that $C = A \cap B$. It follows from point (ii) that $(A - \mathscr{F}_2) \cap (B - \mathscr{G}_2) \subseteq (A \cap B) - (\mathscr{F}_2 \sqcap \mathscr{G}_2)$. Since $A - \mathscr{F}_2 \in \mathscr{F}_1$ and $B - \mathscr{G}_2 \in \mathscr{G}_1$, we see that that $(A \cap B) - (\mathscr{F}_2 \sqcap \mathscr{G}_2) \in \mathscr{F}_1 \sqcap \mathscr{G}_1$, whereby $A \cap B \in (\mathscr{F}_1 \sqcap \mathscr{G}_1) + (\mathscr{F}_2 \sqcap \mathscr{G}_2)$, as desired.

2.1.4. Translation-invariant and idempotent families

For $m \in \mathbb{N}$ and a family \mathscr{F} of subsets of \mathbb{N} , we define

$$\mathscr{F} - m := \{B - m \mid B \in \mathscr{F}\}.$$

The family \mathscr{F} is translation invariant if $\mathscr{F} - n \subseteq \mathscr{F}$ for all $n \in \mathbb{N}$. In the algebra of the previous section, this is equivalent to saying that $\mathscr{F} \subseteq \{\mathbb{N}\} + \mathscr{F}$. The family \mathscr{F} is idempotent if $\mathscr{F} \subseteq \mathscr{F} + \mathscr{F}$, equivalently, if for all $A \in \mathscr{F}$, $A - \mathscr{F} \in \mathscr{F}$. (We comment on why the set inclusion in the definition is not an equality in Remark 2.8 below.) Note that all translation-invariant families are idempotent. We will give some less trivial examples of idempotent families in Section 2.3.

Lemma 2.4. Let \mathscr{F} and \mathscr{G} be families.

- (i) If \mathscr{F} and \mathscr{G} are idempotent, then $\mathscr{F} \sqcap \mathscr{G}$ is idempotent.
- (ii) If \mathscr{F} and \mathscr{G} are translation invariant, then $\mathscr{F} \sqcap \mathscr{G}$ is translation invariant.

Proof. (i) We appeal to Lemma 2.3 (ix) with $\mathscr{F}_1 = \mathscr{F}_2 = \mathscr{F}$ and $\mathscr{G}_1 = \mathscr{G}_2 = \mathscr{G}$. Thus,

$$\mathscr{F} \sqcap \mathscr{G} \subseteq (\mathscr{F} + \mathscr{F}) \sqcap (\mathscr{G} + \mathscr{G}) \subseteq (\mathscr{F} \sqcap \mathscr{G}) + (\mathscr{F} \sqcap \mathscr{G}),$$

as desired. Statement (ii) follows similarly and is left to the reader.

Remark 2.5. The following situation arises several times in this paper. A family \mathscr{F} is defined as the upward closure of a collection of special subsets of \mathbb{N} . To show that \mathscr{F} is idempotent, it suffices to show that for all special sets $A \subseteq \mathbb{N}$, the set $A - \mathscr{F}$ belongs to \mathscr{F} . Indeed, if $B \in \mathscr{F}$, then B contains some special set A. If we show that $A - \mathscr{F} \in \mathscr{F}$, then since $A - \mathscr{F} \subseteq B - \mathscr{F}$, we have that $B - \mathscr{F} \in \mathscr{F}$, as desired.

Idempotent ultrafilters are important objects with a deep history in the subject (cf. [2]). It appears that there are two different definitions of "idempotent ultrafilter." In the literature, a family \mathscr{F} is an idempotent ultrafilter if it is an ultrafilter and satisfies $\mathscr{F} = \mathscr{F} + \mathscr{F}$. According to our definitions, an idempotent ultrafilter is an idempotent family (ie. $\mathscr{F} \subseteq \mathscr{F} + \mathscr{F}$) that is also an ultrafilter. We show in the following lemma these two definition are, in fact, the same. We will use facts from Section 2.2.3.

Theorem 2.6. Let \mathscr{F} be a family. The following are equivalent:

- (i) the family \mathscr{F} is a proper, idempotent filter and is maximal (by containment) in the set of proper, idempotent filters;
- (ii) the family \mathscr{F} is an ultrafilter that satisfies $\mathscr{F} \subseteq \mathscr{F} + \mathscr{F}$;
- (iii) the family \mathscr{F} is an ultrafilter that satisfies $\mathscr{F} = \mathscr{F} + \mathscr{F}$.

In particular, every proper, idempotent filter is contained in an idempotent ultrafilter.

Proof. (i) \implies (ii) We will prove first that every proper, idempotent filter is contained in an idempotent family that is an ultrafilter. Supposing this is so, if \mathscr{F} is a maximal proper, idempotent filter, then it is contained in an idempotent family \mathscr{G} that is an ultrafilter. By the maximality of \mathscr{F} as a proper, idempotent filter, we have that $\mathscr{F} = \mathscr{G}$, giving (ii), as desired.

Suppose \mathscr{F} is a proper, idempotent filter and is maximal as such. We claim that the set $\overline{\mathscr{F}} := \bigcap_{A \in \mathscr{F}} \overline{A}$ is a nonempty, compact subsemigroup of $(\beta \mathbb{N}, +)$. Because ultrafilters are maximal filters, there is an ultrafilter containing \mathscr{F} , and so $\overline{\mathscr{F}}$ is nonempty. The set $\overline{\mathscr{F}}$ is

compact because it is closed and $\beta \mathbb{N}$ is compact. Suppose $q_1, q_2 \in \overline{\mathscr{F}}$. Since $\mathscr{F} \subseteq q_1$ and $\mathscr{F} \subseteq q_2$, it follows from Lemma 2.3 (vii) and the idempotency of \mathscr{F} that

$$\mathscr{F} \subseteq \mathscr{F} + \mathscr{F} \subseteq q_1 + q_2.$$

Therefore, we have that $q_1 + q_2 \in \overline{\mathscr{F}}$.

Finally, since $\overline{\mathscr{F}}$ is a nonempty, compact subsemigroup of $(\beta \mathbb{N}, +)$, by [32, Thm. 2.5], there exists an idempotent ultrafilter containing \mathscr{F} .

(ii) \implies (iii) Suppose \mathscr{F} is an ultrafilter that satisfies $\mathscr{F} \subseteq \mathscr{F} + \mathscr{F}$. By Lemma 2.3 (i) and (iii), the family $\mathscr{F} + \mathscr{F}$ is a proper filter. Since \mathscr{F} is a maximal proper filter, we have that $\mathscr{F} = \mathscr{F} + \mathscr{F}$, as desired.

(iii) \implies (i) If \mathscr{F} is an ultrafilter that satisfies $\mathscr{F} = \mathscr{F} + \mathscr{F}$, then it is a proper, idempotent filter. Since it is maximal amongst proper filters, it is maximal amongst proper, idempotent filters.

Remark 2.7. A routine application of Zorn's lemma gives the existence of maximal proper, idempotent filters. It follows from Theorem 2.6 that these are ultrafilters. The proof given here relies on the fact that a compact, Hausdorff, right-topological semigroup contains an idempotent element (see [32, Thm. 2.5] and the end-of-chapter notes). Can one give a topology-free proof? If so, one can arrive at a topology-free proof of Theorem 2.6 and, when combined with Lemma 2.12, of Hindman's theorem (along the lines of that in [32, Cor. 5.9]).

Another routine application of Zorn's lemma gives the existence of maximal proper, translation-invariant filters. It is an easy exercise to show that there are no translation-invariant ultrafilters on N. Therefore, in contrast to maximal proper, idempotent filters, maximal proper, translation-invariant filters are not ultrafilters.

Remark 2.8. The definition of idempotent family we give in this section matches the one for filters in [42]. It may seem more natural to require equality in the definition: $\mathscr{F} = \mathscr{F} + \mathscr{F}$. While certainly some families do satisfy this – in particular, idempotent ultrafilters (Theorem 2.6) – equality appears to be rather rare. Here is an example of a dynamically simple idempotent filter \mathscr{F} which satisfies $\mathscr{F} \subsetneq \mathscr{F} + \mathscr{F}$. See Section 2.2.1 for the notation.

Fix $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, and consider the irrational rotation $(\mathbb{R}/\mathbb{Z}, T_{\alpha} : x \mapsto x + \alpha)$. Put

$$\mathscr{F} := \uparrow \{ R(0, B_{\varepsilon}(0)) \mid \varepsilon > 0 \}$$

The family \mathscr{F} is an idempotent filter. This can be easily checked (see, eg., the proof of the "only if" statement of Theorem 3.18).

Let $(k_i)_{i=1}^{\infty} \subseteq \mathbb{N}$ be such that the only limit point of $i \mapsto k_i \alpha$ is 0. For $n \in \mathbb{N}$, the quantity

$$\varepsilon_n := \min \left\{ \|k_i \alpha - n\alpha\|/2 \mid i \in \mathbb{N}, \ k_i > n \right\}$$

where $\|\cdot\|: \mathbb{R}/\mathbb{Z} \to [0, 1/2)$ denotes the Euclidean distance to \mathbb{Z} , is non-zero. We claim that the set

$$A := \bigcup_{n=1}^{\infty} \left(R(0, B_{\varepsilon_n}(0)) + n \right)$$

belongs to $\mathscr{F} + \mathscr{F}$ but not to \mathscr{F} . Indeed, for all $n \in \mathbb{N}$, the set A - n contains $R(0, B_{\varepsilon_n}(0)) \in \mathscr{F}$, and so $A \in \{\mathbb{N}\} + \mathscr{F} \subseteq \mathscr{F} + \mathscr{F}$.

To see that $A \notin \mathscr{F}$, we will show that for all $\delta > 0$, we have $R(0, B_{\delta}(0)) \not\subseteq A$. Let $\delta > 0$, and let $i \in \mathbb{N}$ such that $k_i \in R(0, B_{\delta}(0))$. Now, for all $n < k_i$, we have that $||k_i \alpha - n\alpha|| \ge 2\varepsilon_n$, whereby $k_i \notin R(0, B_{\varepsilon_n}(0)) + n$. For all $n \ge k_i$, we have that $k_i \notin R(0, B_{\varepsilon_n}(0)) + n$ since $R(0, B_{\varepsilon_n}(0)) + n \subseteq \{n + 1, n + 2, \ldots\}$. Therefore, we have that $k_i \notin A$, as was to be shown.

2.2. Topology and dynamics

For a set U in a topological space X, we denote by \overline{U} , U° , and $\partial U := \overline{U} \setminus U^{\circ}$ the closure, interior, and boundary of U, respectively. In a metric space (X, d), we denote by $B_{\varepsilon}(x)$ the open ball centered at a point $x \in X$ with radius $\varepsilon > 0$.

2.2.1. Topological dynamics

A (topological dynamical) system (X,T) is a nonempty, compact metric space (X, d_X) together with a continuous self-map $T: X \to X$. A subsystem of (X,T) is a system of the form (Y,T), where $Y \subseteq X$ is nonempty, closed (equivalently, compact), and *T*-invariant, meaning $TY \subseteq Y$. A system (X,T) whose only subsystem is itself is called minimal. Given two systems (X,T) and (Y,S), a continuous surjection $\pi: X \to Y$ that satisfies $S \circ \pi = \pi \circ T$ is called a factor map of systems. We write $\pi: (X,T) \to (Y,S)$ to indicate that π is a factor map.

All systems in this paper will be considered as actions of the semigroup $(\mathbb{N}, +)$ by continuous maps on a compact metric spaces. Thus, even in the event that T is a homeomorphism (in which case, the system (X, T) is called *invertible*), the emphasis will be on non-negative iterates of the map T.

We will occasionally have need to consider non-metrizable systems. A not-necessarilymetrizable system is a nonempty, compact Hausdorff space X together with a continuous self-map $T: X \to X$. We will always specify when a system need not be metrizable. All terminology and notation will be shared without confusion between the metric and non-metric cases.

Given a system (X,T), a point $x \in X$, and a set $A \subseteq \mathbb{N}$, we define

$$T^A x := \{ T^a x \mid a \in A \}.$$

The orbit of x is the set $T^{\mathbb{N}}x$ and the orbit closure of x is the set $\overline{T^{\mathbb{N}}x}$. Note that $\overline{T^{\mathbb{N}}x}$ is a nonempty, closed, T-invariant subset of X, whereby $(\overline{T^{\mathbb{N}}x}, T)$ is a subsystem of (X, T). Thus, we see that (X, T) is minimal if and only if every point $x \in X$ has a dense orbit.

Given a system (X,T), a set $U \subseteq X$, and a point $x \in X$, we write

$$R(x,U) := \left\{ n \in \mathbb{N} \mid T^n x \in U \right\}$$

for the times at which the point x visits the set U. It is a simple algebraic fact that we will use without mention that for $n \in \mathbb{N}_0$,

$$R(x, U) - n = R(T^n x, U) = R(x, T^{-n}U).$$

A set $A \subseteq \mathbb{N}$ is syndetic if there exists $N \in \mathbb{N}$ such that $A \cup (A-1) \cup \cdots \cup (A-N) = \mathbb{N}$. Syndetic sets – discussed in more context in Section 2.3.1 – are intimately linked to minimal dynamics. Indeed, let (X, T) be a not-necessarily-metrizable system. Both of the facts below are not hard to show following the results in [20, Ch. 1, Sec. 4].

(i) Let $x \in X$. The subsystem $(\overline{T^{\mathbb{N}_0}x}, T)$ is minimal if and only if for all open $U \subseteq X$ containing x, the set R(x, U) is syndetic. In this case, the point x is said to be uniformly

recurrent.

(ii) The system (X,T) is minimal if and only if for all $x \in X$ and all nonempty, open $U \subseteq X$, the set R(x,U) is syndetic.

We will use these facts several times in this paper.

Lemma 2.9. Let (X,T) be a minimal system. There exists an isolated point in X if and only if X is finite and the system (X,T) is periodic.

Proof. If $x \in X$ is isolated, the set $U := \{x\}$ is open. Since (X, T) is minimal, the set R(x, U) is syndetic. In particular, there exists $n \in \mathbb{N}$ such that $T^n x = x$, whereby x is a periodic point. Since (X, T) is minimal, the set X is equal to the (finite) orbit of x and (X, T) is periodic. The converse is trivial.

The dual notions of distality and proximality arise a few times in this paper. Let (X,T) be a system and $x, y \in X$. The points x and y are *proximal* if $\inf_{n \in \mathbb{N}} d_X(T^n x, T^n y) = 0$. The point x is called *distal* if it is proximal only to itself. The system (X,T) is called *distal* if every point is distal.

2.2.2. Symbolic space and full shift

We denote the space of 0–1 valued functions on \mathbb{N} by $\{0,1\}^{\mathbb{N}}$. Given the product topology, this is a compact Hausdorff space. There are many equivalent metrics that generate the product topology on $\{0,1\}^{\mathbb{N}}$. Instead of specifying one of these metrics explicitly, it suffices for our purposes to fix one and note that two elements of $\{0,1\}^{\mathbb{N}}$ are close if and only if they agree as functions on a long initial interval $\{1,\ldots,N\}$ of \mathbb{N} .

Let $\omega \in \{0,1\}^{\mathbb{N}}$ and $i \in \mathbb{N}$. We write $\omega(i)$ for the value of the function ω at i. Given $k \in \{0,1\}$, the *cylinder set* of all those functions in $\{0,1\}^{\mathbb{N}}$ that evaluate to k at i is denoted by $[k]_i$. This is easily checked to be a clopen subset of $\{0,1\}^{\mathbb{N}}$. The *support* of ω is

$$\operatorname{supp}(\omega) := \left\{ n \in \mathbb{N} \mid \omega(n) = 1 \right\}.$$

We define the (left) shift $\sigma : \{0,1\}^{\mathbb{N}} \to \{0,1\}^{\mathbb{N}}$ at the function ω by the rule $(\sigma\omega)(i) = \omega(i+1), i \in \mathbb{N}$. The shift is easily seen to be a continuous self-map of $\{0,1\}^{\mathbb{N}}$. The *full shift* is the system $(\{0,1\}^{\mathbb{N}}, \sigma)$.

It will frequently be useful to consider functions on \mathbb{N}_0 instead of \mathbb{N} . Everything written above applies equally well to the space $\{0,1\}^{\mathbb{N}_0}$ and the *full shift* $(\{0,1\}^{\mathbb{N}_0},\sigma)$.

2.2.3. Dynamics on the space of ultrafilters

The Stone-Čech compactification of \mathbb{N} , denoted $\beta \mathbb{N}$, is a compact Hausdorff, right-topological semigroup that provides a convenient universal object in the category of minimal systems. We summarize here just what we will need; the interested reader is directed to [32, Ch. 19] and the references therein for more information.

As a set, we realize $\beta \mathbb{N}$ as the set of all ultrafilters on \mathbb{N} (cf. Section 2.1.2). Sets of the form

$$\overline{A} := \{ p \in \beta \mathbb{N} \mid A \in p \}, \quad A \subseteq \mathbb{N},$$

form a base for a non-metrizable, compact Hausdorff topology on $\beta \mathbb{N}$. Since ultrafilters are families, addition of two ultrafilters is as defined in Section 2.1.3. It follows by combining

Lemma 2.2 (ii) and Lemma 2.3 (viii) that the sum of two ultrafilters is an ultrafilter. Thus, $(\beta \mathbb{N}, +)$ is a compact Hausdorff semigroup.

It is not hard to check that for all $n \in \mathbb{N}$ and $q \in \beta\mathbb{N}$, the maps $\beta\mathbb{N} \to \beta\mathbb{N}$ defined by $p \mapsto n+p$ and $p \mapsto p+q$ are continuous. Thus, $(\beta\mathbb{N}, +)$ is called a *right-topological* semigroup. A nonempty subset $L \subseteq \beta\mathbb{N}$ with the property that $p + L \subseteq L$ for all $p \in \beta\mathbb{N}$ is called a *left ideal*. A left ideal that is minimal amongst all left ideals is called a *minimal left ideal*. By a routine application of Zorn's lemma, minimal left ideals exist in $(\beta\mathbb{N}, +)$. If $L \subseteq \beta\mathbb{N}$ is a minimal left ideal, for all $p \in L$, the set $\beta\mathbb{N} + p$ is a left ideal contained in L, and hence is equal to L. Since addition on the right by p is continuous and $\beta\mathbb{N}$ is compact, we see that minimal left ideals are compact.

Denoting addition by 1 on the left by $\lambda_1 : \beta \mathbb{N} \to \beta \mathbb{N}$, we see that $(\beta \mathbb{N}, \lambda_1)$ is a nonmetrizable topological dynamical system. Fix $q \in \beta \mathbb{N}$. Since addition on the right by q is continuous and \mathbb{N} is dense in $\beta \mathbb{N}$, we have that the orbit closure of q under λ_1 is

$$\overline{\lambda_1^{\mathbb{N}}q} = \overline{\mathbb{N}+q} = \overline{\mathbb{N}}+q = \beta\mathbb{N}+q,$$

a left ideal. It is not hard to see that $\beta \mathbb{N} + q$ is a minimal left ideal if and only if the system $(\beta \mathbb{N} + q, \lambda_1)$ is minimal. (Indeed, the system $(\beta \mathbb{N} + q, \lambda_1)$ is minimal if and only if for all $p \in \beta \mathbb{N} + q$, the orbit closure $\beta \mathbb{N} + p$ – a left ideal – is equal to $\beta \mathbb{N} + q$.) In this case, the ultrafilter q is called a *minimal ultrafilter*. We see, then, that the minimal left ideals of the semigroup $(\beta \mathbb{N}, +)$ are precisely the minimal subsystems of $(\beta \mathbb{N}, \lambda_1)$.

Let (X, T) be a not-necessarily-metrizable system, and let $x \in X$. Consider the map $T x : \mathbb{N} \to X$ given by $n \mapsto T^n x$. By the universal property of the Stone-Čech compactification, the map T x lifts to a continuous map $T x : \beta \mathbb{N} \to X$. Concretely, for $p \in \beta \mathbb{N}$, the point $T^p x \in X$ is defined uniquely by the property that for all open $U \subseteq X$ containing $T^p x$, the set R(x, U) is a member of p. The following fact will be useful for us later on: if $U \subseteq X$ is clopen and $p \in \beta \mathbb{N}$, then

$$R(x, U) - p = R(T^{p}x, U).$$
(2.1)

Indeed, for $n \in \mathbb{N}$, we see that $n \in R(x, U) - p$ if and only if $R(T^n x, U) = R(x, U) - n \in p$. If $R(T^n x, U) \in p$, then

$$T^n T^p x = T^{n+p} x = T^{p+n} x = T^p T^n x \in \overline{U} = U,$$

whereby $n \in R(T^p x, U)$. On the other hand, if $n \in R(T^p x, U)$, then $T^p T^n x = T^n T^p x \in U$, whereby $R(T^n x, U) \in p$.

Minimal left ideals of $\beta\mathbb{N}$ under addition by 1 are universal minimal dynamical systems. More precisely, let $L \subseteq \beta\mathbb{N}$ be a minimal left ideal, and let (X,T) be a not-necessarilymetrizable, minimal dynamical system. Fix $x \in X$. The map $T x : \beta\mathbb{N} \to X$ described in the previous paragraph is continuous and satisfies $T^{\lambda_1 p} x = T^{1+p} x = T(T^p x)$ and hence is a factor map $(\beta\mathbb{N}, \lambda_1) \to (X,T)$ of systems. Restricted to L, by the minimality of (X,T), we see that $T x : (L, \lambda_1) \to (X,T)$ is surjective and hence is a factor map of minimal systems.

2.3. Families of sets from dynamics

We outline below those families of subsets of \mathbb{N} that are important in this work. The following nomenclature will be particularly convenient. Let \mathscr{G} be a family. A set $A \subseteq \mathbb{N}$ is called a \mathscr{G} set if it is a member of \mathscr{G} , and a family \mathscr{F} is called a \mathscr{G} family if all members of \mathscr{F} are \mathscr{G}

sets, ie., $\mathscr{F} \subseteq \mathscr{G}$. We will frequently consider in this work, for example, syndetic filters, ie., filters whose members are all syndetic.

2.3.1. Syndetic, thick, and piecewise syndetic sets

A set $A \subseteq \mathbb{N}$ is ...

(i) \ldots syndetic if there exists $N \in \mathbb{N}$ such that

$$A \cup (A-1) \cup \dots \cup (A-N) = \mathbb{N};$$

(ii) ... thick if for all finite $F \subseteq \mathbb{N}$, there exists $n \in \mathbb{N}$ such that $F + n \subseteq A$;

(iii) ... piecewise syndetic if there exists $N \in \mathbb{N}$ such that the set

$$A \cup (A-1) \cup \dots \cup (A-N)$$

is thick.

We denote by S, T, and \mathcal{PS} the families of syndetic, thick, and piecewise syndetic subsets of \mathbb{N} , respectively. All three families are translation invariant. It is well-known and not difficult to show that the families of syndetic and thick sets are dual and that

$$\mathcal{PS} = \mathcal{S} \sqcap \mathcal{T}.$$

By Lemma 2.1, the family \mathcal{PS} is piecewise syndetic and its dual, \mathcal{PS}^* , is a filter. It is not hard to show that a set A belongs to \mathcal{PS}^* if and only if for all finite $F \subseteq \mathbb{N}$, there exists a syndetic set $S \subseteq \mathbb{N}$ such that $F + S \subseteq A$. Thus, members of the filter \mathcal{PS}^* are called *thickly* syndetic sets.

In the following lemmas and in those in the next section, we demonstrate how well-known families of subsets of \mathbb{N} can be characterized by membership in certain types of filters. These results will be useful for us later on.

Lemma 2.10.

- (i) Every proper, translation-invariant filter is a thick, translation-invariant filter.
- (ii) A set $A \subseteq \mathbb{N}$ is thick if and only if it is a member of a proper, translation-invariant filter.

Proof. (i) Let \mathscr{F} be a proper, translation-invariant filter, and let $A \in \mathscr{F}$. For all $N \in \mathbb{N}$, the set $\bigcap_{n=0}^{N} (A-n) \in \mathscr{F}$, and hence is nonempty. It follows that A contains an interval of length N + 1, and hence that A is thick. It follows that \mathscr{F} is thick.

(ii) If $A \subseteq \mathbb{N}$ is a member of a proper, translation-invariant filter, then it is thick, as shown in the proof of (i). Conversely, if $A \subseteq \mathbb{N}$ is thick, then the family

$$\uparrow \left\{ \bigcap_{n=0}^{N} (A-n) \; \middle| \; N \in \mathbb{N} \right\}$$

is quickly checked to be a thick (hence, proper), translation-invariant filter.

The following lemma gives a useful characterization of the filter of thickly syndetic sets as the largest syndetic, translation-invariant filter.

Lemma 2.11. The family \mathcal{PS}^* is a syndetic, translation-invariant filter. If \mathscr{F} is a syndetic, translation-invariant filter, then $\mathscr{F} \subseteq \mathcal{PS}^*$.

Proof. That \mathcal{PS}^* is a syndetic, translation-invariant filter was discussed above.

Let \mathscr{F} be a syndetic, translation-invariant filter, and let $A \in \mathscr{F}$. Since \mathscr{F} is translation invariant, for all $N \in \mathbb{N}$, the sets $A - 1, \ldots, A - N$ all belong to \mathscr{F} . Since \mathscr{F} is a syndetic filter, the set $(A - 1) \cap \cdots \cap (A - N)$ belongs to \mathscr{F} and is syndetic. Since

$$\{1,\ldots,N\} + ((A-1)\cap\cdots\cap(A-N)) \subseteq A,$$

we see that A is thickly syndetic and hence is a member of \mathcal{PS}^* .

2.3.2. Finite sums sets and central sets

The families of infinite, finite sums sets and central sets will also appear in this work. A set $A \subseteq \mathbb{N}$ is called an \mathcal{IP} set (also, an *infinite, finite sums set*) if there exists a sequence $(x_i)_{i=1}^{\infty} \subseteq \mathbb{N}$ such that

$$\operatorname{FS}(x_i)_{i=1}^{\infty} := \left\{ \sum_{f \in F} x_f \; \middle| \; F \subseteq \mathbb{N} \text{ is finite and nonempty} \right\} \subseteq A.$$

Here "FS" stands for "finite sums." It is a consequence of Hindman's theorem [31] that the family \mathcal{IP} is partition regular (see [5, Lemma 2.1]). Thus, the dual family \mathcal{IP}^* is a filter.

It is well known [32, Thm. 5.12] that a set is an \mathcal{IP} set if and only if it is member of an idempotent ultrafilter. The following lemma offers a characterization in terms of idempotent filters, showing that the assumption of maximality (recall, ultrafilters are maximal filters) is superfluous.

Lemma 2.12. A set $A \subseteq \mathbb{N}$ is an \mathcal{IP} set if and only if it is a member of a proper, idempotent filter.

Proof. Suppose $A \subseteq \mathbb{N}$ is an \mathcal{IP} set. There exists $(x_i)_{i=1}^{\infty} \subseteq \mathbb{N}$ such that $FS(x_i)_{i=1}^{\infty} \subseteq A$. We will check that the family

$$\mathscr{F} := \uparrow \left\{ \operatorname{FS}(x_i)_{i=N}^{\infty} \mid N \in \mathbb{N} \right\}$$

is a proper, idempotent filter containing A. It is easy to see from the definition that \mathscr{F} is a proper filter.

To see that \mathscr{F} is idempotent, let $A \in \mathscr{F}$. We must show that $A - \mathscr{F} \in \mathscr{F}$. By definition, there exists $N \in \mathbb{N}$ such that $\mathrm{FS}(x_i)_{i=N}^{\infty} \subseteq A$. We claim that $\mathrm{FS}(x_i)_{i=N}^{\infty} \subseteq A - \mathscr{F}$, from which it will follow that $A - \mathscr{F} \in \mathscr{F}$. Let $x \in \mathrm{FS}(x_i)_{i=N}^{\infty}$. We must show that $A - x \in \mathscr{F}$. There exists $F \subseteq \mathbb{N}$ finite with $\min F \ge N$ such that $x = \sum_{f \in F} x_f$. We see that $A - x \supseteq$ $\mathrm{FS}(x_i)_{i=\max F+1}^{\infty}$, which belongs to \mathscr{F} . Therefore, $A - x \in \mathscr{F}$, as desired.

Conversely, suppose that A is a member of a proper, idempotent filter \mathscr{F} . Since $A \in \mathscr{F}$ and $A - \mathscr{F} \in \mathscr{F}$, there exists $x_1 \in A \cap (A - \mathscr{F})$. Since $A \in \mathscr{F}$ and $A - x_1 \in \mathscr{F}$, we have that $A \cap (A - x_1) \in \mathscr{F}$, and hence that $(A \cap (A - x_1)) - \mathscr{F} \in \mathscr{F}$. There exists

$$x_2 \in A \cap (A - x_1) \cap \left(\left(A \cap (A - x_1) \right) - \mathscr{F} \right).$$

So far, we have found $x_1, x_2 \in \mathbb{N}$ such that $\{x_1, x_2, x_1 + x_2\} \subseteq A$. Iterating, we will take x_3 from the set

$$A \cap (A - x_1) \cap \left(\left(A \cap (A - x_1) \right) - x_2 \right) \cap \left(\left(A \cap (A - x_1) \cap \left(\left(A \cap (A - x_1) \right) - x_2 \right) \right) - \mathscr{F} \right)$$

so that all finite sums from the set $\{x_1, x_2, x_3\}$ belong to A. It is clear how to proceed to find $(x_i)_{i=1}^{\infty} \subseteq \mathbb{N}$ such that $FS(x_i)_{i=1}^{\infty} \subseteq A$, as desired.

We will consider the family of central sets for its importance to the subject and for its connection to the families of dynamically defined sets at the heart of this work. Furstenberg [20, Def. 8.3] defined a set $A \subseteq \mathbb{N}$ to be *central* if there exists a system (X, T), a uniformly recurrent point $x \in X$, a point $y \in X$ proximal to x, and an open set $U \subseteq X$ containing x such that A = R(y, U). We denote the family of central sets by \mathcal{C} .

Bergelson and Hindman [4] showed that a subset of \mathbb{N} is central if and only if it is a member of an idempotent ultrafilter on \mathbb{N} . In analogy to \mathcal{IP} sets and Lemma 2.12, central sets can be characterized in terms of a special class of idempotent filters. Defining this class – "collectionwise piecewise syndetic," idempotent filters – would require too large a diversion, so we leave it to the interested reader. (See [32, Def. 14.19] for a definition of collectionwise piecewise syndetic.)

It was stated by Furstenberg and follows immediately from Bergelson and Hindman's theorem that the family C is partition regular. It is well known that thick sets are central and that central sets are piecewise syndetic and \mathcal{IP} . Dual to the family of central sets is the family C^* , a syndetic filter that will appear prominently in Section 5. Fig. 2 depicts the relationships between the families of subsets of \mathbb{N} introduced so far.



Figure 2: Containment amongst the families appearing in this paper. An arrow $\mathscr{F} \to \mathscr{G}$ indicates that $\mathscr{F} \subsetneq \mathscr{G}$. Those families not prefixed with the letter "d" are defined in Section 2.3. The families dcS / dS, dcT / dT, and dcPS / dPS are the main objects of study in this work and are defined in Sections 3, 4, and 5, respectively. There are eight families and their duals, yielding sixteen families in all, two pairs of which coincide. The diagram's symmetry is explained by the family duality described in Section 2.1.1.

The families \mathcal{IP} and \mathcal{C} are examples of idempotent families that are not translation invariant. It is a short exercise to verify this directly from the definitions. Alternatively, one can note two things: first, by the characterizations given above,

$$\mathcal{IP} = \bigcup_{\substack{p \in \beta \mathbb{N} \\ p \text{ idempotent}}} p \quad \text{and} \quad \mathcal{C} = \bigcup_{\substack{p \in \beta \mathbb{N} \text{ minimal} \\ p \text{ idempotent}}} p,$$

and second, a union of idempotent families is idempotent.

3. Part I: Dynamically syndetic sets

In this section, we define the family of dynamically syndetic sets and give a number of equivalent characterizations of dynamical syndeticity. The main result is Theorem 3.18, a combinatorial characterization of dynamical syndeticity in terms of syndetic, idempotent filters.

3.1. Definitions and first characterizations

A set $A \subseteq \mathbb{N}$ is dynamically syndetic if there exists a minimal topological dynamical system (X, T), a point $x \in X$, and a nonempty, open set $U \subseteq X$ such that $R(x, U) \subseteq A$. If $x \in U$, the set A is called dynamically central syndetic. The families of dynamically syndetic and dynamically central syndetic subsets of \mathbb{N} are denoted dS and dcS, respectively.

It is clear from the definitions and the facts in Section 2.2.1 that

$$dc\mathcal{S} \subseteq d\mathcal{S} \subseteq \mathcal{S}.$$

The set of odd, positive integers and the set in (1.1) from the introduction demonstrate that each of these inclusions is proper. (That $2\mathbb{N} - 1 \notin dcS$ follows from, eg., Lemma 3.6.) Here is a short list of examples of dynamically (central) syndetic sets that may be helpful to have in mind. Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$.

(i) The irrational rotation $(\mathbb{R}/\mathbb{Z}, x \mapsto x + \alpha)$, where α is interpreted as an element of \mathbb{R}/\mathbb{Z} , is a minimal system. For all $x \in \mathbb{R}/\mathbb{Z}$ and all nonempty, open $U \subseteq \mathbb{R}/\mathbb{Z}$, the set

$$R(x,U) = \{ n \in \mathbb{N} \mid x + n\alpha \in U \}$$

is dynamically syndetic and, if $x \in U$, dynamically central syndetic. More generally, supersets of Bohr (Bohr₀) sets – those arising in this way from compact group rotations – are dynamically (central) syndetic.

(ii) When $\alpha > 1$, the sequence $(\lfloor n\alpha \rfloor)_{n \in \mathbb{N}}$ is called a *Beatty sequence*. It is not hard to show that the image of this sequence is equal to the set

$$\{n \in \mathbb{N} \mid \{n/\alpha\} \in ((\alpha - 1)/\alpha, 1)\},\$$

where $\{\cdot\} : \mathbb{R} \to \mathbb{R}/\mathbb{Z}$ is the quotient map. Thus, Beatty sequences yield Bohr, and hence dynamically syndetic, sets. What is not so clear is that, in fact, Beatty sequences yield dynamically central syndetic sets. We will see this as a consequence of Theorem 3.8.

(iii) Using the notation from (i), the set

$$\{n \in \mathbb{N} \mid n^2 \alpha \in U\}$$

is a dynamically syndetic set. It is, in fact, an example of a nil-Bohr set, one that arises from a rotation on a compact homogeneous space of a nilpotent Lie group. Nil-Bohr (nil-Bohr₀) sets are dynamically (central) syndetic [34].

(iv) Denote by $\nu_2 : \mathbb{N} \to \mathbb{N}_0$ the 2-adic valuation. The set

$$A := \{ n \in \mathbb{N} \mid \nu_2(n) \text{ is even} \}$$

is a dynamically syndetic set. To see this, one can show that the point 1_A is uniformly recurrent in the full shift $(\{0,1\}^{\mathbb{N}_0}, \sigma)$, whereby $A = R_{\sigma}(1_A, [1]_0)$ is dynamically syndetic. In fact, the point $1_{A\cup\{0\}} \in [1]_0$ is uniformly recurrent, and so A is dynamically central syndetic. By Lemma 3.6 and the fact that $\mathbb{N}\setminus A = A/2$, the same can be said about the set of positive integers with odd 2-adic valuation.

(v) The Chacon word 00100010100100010001010010010010010... is the unique, nonconstant word in the symbols 0 and 1 that is fixed under the substitutions $0 \mapsto 0010$ and $1 \mapsto 1$. The set of locations of the 1's in the Chacon word is a dynamically syndetic set, for the same reason as in the previous point: the word, interpreted as a point in the full shift ($\{0, 1\}^{\mathbb{N}_0}, \sigma$), is known to be uniformly recurrent [11, Sec. 5.6.6].

Many of the results in this section demonstrate the robustness of the family of dynamically (central) syndetic sets under changes to the combinatorial, topological, or dynamical requirements in its definition. The family of dynamically (central) syndetic subsets of \mathbb{N} remains unchanged if ...

- (i) ... the map T is required to be a homeomorphism, that is, if the system (X,T) is required to be invertible. This is shown in Lemma 3.1.
- (ii) \ldots the space X is allowed to be non-metrizable. This is a consequence of Theorem 3.3.
- (iii) ... the space X is required to be zero dimensional and the set U is required to be clopen. This is a consequence of Theorem 3.8.
- (iv) ... (for dynamically central syndetic sets) the point x is allowed to be in \overline{U} instead of being required to be in U. Sets of this form are called *very strongly central* in [6, Def. 2.10], where a number of equivalent characterizations in terms of ultrafilters are given. That the families of very strongly central sets and dynamically central syndetic sets are the same is shown in Theorem 3.8.
- (v) ... sets of return times R(x, U) are considered up to equivalence on thickly syndetic sets (\mathcal{PS}^* sets). This is a consequence of Theorem 5.13: a set $A \subseteq \mathbb{N}$ is dynamically (central) syndetic if and only if for all thickly syndetic $B \subseteq \mathbb{N}$, the set $A \cap B$ is dynamically (central) syndetic. We will have immediate need for a more basic version of this result for cofinite sets in Lemma 3.2.

There are other directions, however, in which this definitional robustness does not extend.

- (vi) Minimality is a key feature of the definition. For all $A \subseteq \mathbb{N}$, there exists a system (X, T), a point $x \in X$, and a nonempty, open set $U \subseteq X$ containing x such that A = R(x, U). Indeed, note that considering A as a subset of \mathbb{N}_0 , we have that $A = R(1_A, [1]_0)$ in the full shift $(\{0, 1\}^{\mathbb{N}_0}, \sigma)$. Thus, dropping the minimality assumption trivializes the definition of dynamical syndeticity.
- (vii) That we are considering actions of the semigroup $(\mathbb{N}, +)$ instead of the group $(\mathbb{Z}, +)$ matters. For example, if we define dynamically central syndetic subsets of \mathbb{Z} analogously to those in \mathbb{N} , we see that all dynamically central syndetic subsets of \mathbb{Z} contain

0. In this case, if $x \in \overline{U} \setminus U$, then the set $R(x, U) \subseteq \mathbb{Z}$ is, in contrast to Theorem 3.8, not dynamically central syndetic because it does not contain 0. Other results from this paper for $(\mathbb{N}, +)$ -systems fail to generalize under this definition, too, most notably the fact that thickly syndetic (\mathcal{PS}^*) sets (which do not necessarily contain 0) are dynamically central syndetic. Along with other evidence, this suggests that allowing $x \in \overline{U}$ in the definition of the family of dynamically central syndetic sets (as in Theorem 3.8 (ii)) may yield the "right" notion for more general (semi)group actions.

3.1.1. Definitional robustness of dynamically syndetic sets

Here we show that invertibility, changes on finite sets, and metrizability do not affect the definition of the family of dynamically syndetic sets.

Lemma 3.1. A set $A \subseteq \mathbb{N}$ is dynamically (central) syndetic if and only if there exists an invertible system (X,T), a point $x \in X$, and a nonempty, open set $U \subseteq X$ (containing x) such that $R(x,U) \subseteq A$.

Proof. The "if" statement is immediate. To see the "only if" statement, suppose $A \subseteq \mathbb{N}$ is dynamically syndetic. There exists a system (X,T), a point $x \in X$, and a nonempty, open set $U \subseteq X$ such that $R(x,U) \subseteq A$. Let $\pi : (W,T) \to (X,T)$ be the natural extension (see [13, Definition 6.8.10]) of (X,T), and let $w \in W$ be such that $\pi w = x$. Since (X,T) is minimal, so is (W,T) (see, for example, [26, Lemma 2.9]). The system (W,T) is invertible, and $R(w, \pi^{-1}U) = R(x,U)$. Thus, the set A contains $R(w, \pi^{-1}U)$ from the invertible system (W,T). Note that if A is dynamically central syndetic and $x \in U$, then $w \in \pi^{-1}U$, whereby the set $R(x,U) = R(w, \pi^{-1}U)$ is dynamically central syndetic from the invertible system (W,T).

The following lemma shows that dynamically (central) syndetic sets remain such after changes on a finite set. This result is upgraded in Theorem 5.13, which reaches the same conclusion after changes are allowed on a non-piecewise syndetic set.

Lemma 3.2. Let $A \subseteq \mathbb{N}$ be dynamically (central) syndetic. If $B \subseteq \mathbb{N}$ is cofinite, then $A \cap B$ is dynamically (central) syndetic.

Proof. By the definition of dynamically (central) syndetic, it suffices to show that for all minimal systems (X, T), all points $x \in X$, all nonempty, open sets $U \subseteq X$ (containing x), and all $N \in \mathbb{N}$, the set $R(x, U) \setminus \{1, \ldots, N\}$ is dynamically (central) syndetic. So, let (X, T), $x \in X$, $U \subseteq X$ (contain x), and $N \in \mathbb{N}$ be as described.

The set $V := U \setminus \{Tx, T^2x, \ldots, T^Nx\}$ is open and $R(x, U) \setminus \{1, \ldots, N\} \supseteq R(x, V)$. If V is nonempty, then we see that the set $R(x, U) \setminus \{1, \ldots, N\}$ is dynamically syndetic, as desired. Note that if $x \in \{Tx, T^2x, \ldots, T^Nx\}$, then the system is periodic, $X = \{Tx, T^2x, \ldots, T^Nx\}$, and V is empty. Therefore, if V is nonempty and $x \in U$, then $x \in V$, and the set $R(x, U) \setminus \{1, \ldots, N\}$ is dynamically central syndetic, as desired.

If, on the other hand, the set V is empty, then $U \subseteq \{Tx, T^2x, \ldots, T^Nx\}$. Since U is open, we see that some T^ix is an isolated point of X. By Lemma 2.9, the system (X,T) is finite and periodic. In this case, the set $R(x,U) \setminus \{1,\ldots,N\}$ contains an infinite arithmetic progression (of the form $k\mathbb{N}$), which is dynamically (central) syndetic, as desired.

Theorem 3.3. Let (X,T) be a not-necessarily-metrizable, minimal topological dynamical system. For all $x \in X$ and all nonempty, open $U \subseteq X$ (containing x), the set R(x,U) is dynamically (central) syndetic.

Proof. Let $x \in X$ and $U \subseteq X$ be nonempty and open. Fix $L \subseteq \beta \mathbb{N}$ a minimal left ideal. As described in Section 2.2.3, the map $\pi : (L, \lambda_1) \to (X, T)$ given by $\pi(p) = T^p x$ is a factor map of minimal systems. By [32, Thm. 19.23], since x is uniformly recurrent, there exists an idempotent $p \in L$ such that $\pi(p) = x$. It is quick to check that

$$R_T(x,U) = R_{\lambda_1}(p,\pi^{-1}U).$$

We will show that the set $R_{\lambda_1}(p, \pi^{-1}U)$ is dynamically syndetic.

By the definition of the topology on $\beta \mathbb{N}$, since $\pi^{-1}U$ is nonempty and open, there exists $A \subseteq \mathbb{N}$ such that $\emptyset \neq \overline{A} \cap L \subseteq \pi^{-1}U$. In the notation defined in Section 2.1.3, we see that

$$R_{\lambda_1}(p, \pi^{-1}U) \supseteq R_{\lambda_1}(p, \overline{A}) = A - p.$$
(3.1)

Since $\overline{A} \cap L \neq \emptyset$ and (L, λ_1) is minimal, we see that the set A - p is nonempty (indeed, syndetic).

Considering 1_A as an element of $\{0,1\}^{\mathbb{N}_0}$, we see in the full shift $(\{0,1\}^{\mathbb{N}_0}, \sigma)$ that $A = R_{\sigma}(1_A, [1]_0)$. It follows by (2.1) in Section 2.2.3 that

$$A - p = R_{\sigma}(1_A, [1]_0) - p = R_{\sigma}(T^p 1_A, [1]_0).$$

Since p is a minimal idempotent ultrafilter, we have by [32, Thm. 19.23] that the point $T^{p}1_{A}$ is uniformly recurrent under the shift. Since $R_{\sigma}(T^{p}1_{A}, [1]_{0})$ is nonempty, we see that the set $R_{\sigma}(T^{p}1_{A}, [1]_{0})$, and hence the set $A - p = R_{\lambda_{1}}(p, \overline{A})$, is dynamically syndetic. Because these families are upward closed, by (3.1), the same can be said about the set $R_{\lambda_{1}}(p, \pi^{-1}U) = R_{T}(x, U)$, as was to be shown.

In the case that $x \in U$, we wish to show that the set $R_T(x, U)$ is dynamically central syndetic. The argument given above can be followed with some additional information. First, since $x \in U$, the point $p \in \pi^{-1}U$. The set $A \subseteq \mathbb{N}$ can then be assumed to be such that $p \in \overline{A} \cap L$. Considering 1_A as an element of $\{0, 1\}^{\mathbb{N}_0}$, we see

$$R_{\sigma}(1_A, [1]_0) = A \in p,$$

whereby $T^p 1_A \in [1]_0$. Thus, the set $A - p = R_{\sigma}(T^p 1_A, [1]_0)$ is dynamically central syndetic, as desired.

3.1.2. Translates and dilates of dynamically syndetic sets

In this section, we describe what happens to dynamically (central) syndetic sets under translation and dilation. Recall that A + n and A - n are computed as subsets of \mathbb{N} , as described in Section 2.

Lemma 3.4. Let $A \subseteq \mathbb{N}$ be dynamically syndetic.

- (i) For all $n \in \mathbb{N}$, the set A n is dynamically syndetic.
- (ii) The set

$$\{n \in \mathbb{N} \mid A - n \text{ is dynamically central syndetic}\}$$
(3.2)

is dynamically syndetic.

(iii) If A is dynamically central syndetic, then the set in (3.2) is dynamically central syndetic. Moreover, (i), (ii), and (iii) hold with A - n replaced by A + n. Finally,

$$d\mathcal{S} = \bigcup_{n \in \mathbb{N}} \left(dc\mathcal{S} - n \right) = \bigcup_{n \in \mathbb{N}} \left(dc\mathcal{S} + n \right).$$
(3.3)

Proof. Since A is dynamically syndetic, there exists a minimal system (X,T), a point $x \in X$, and a nonempty, open set $U \subseteq X$ such that $R(x,U) \subseteq A$. According to Lemma 3.1, we can assume the system (X,T) is invertible.

For $n \in \mathbb{N}$, we have that

$$R(T^n x, U) = R(x, U) - n \subseteq A - n.$$
(3.4)

Thus, the set A - n is dynamically syndetic, demonstrating (i). It also follows from (3.4) that R(x,U) is a subset of the set in (3.2). Indeed, if $n \in R(x,U)$, then $T^n x \in U$, so the set A - n is dynamically central syndetic. The set R(x,U) is dynamically syndetic, whereby the set in (3.2) is dynamically syndetic, demonstrating (ii). All of this holds in the case that A is dynamically central syndetic, but we have additionally that R(x,U) is dynamically central syndetic, demonstrating (ii).

Now we will show that (i), (ii), and (iii) hold with A - n replaced by A + n.

Let $n \in \mathbb{N}$. Recall that $T: X \to X$ is a homeomorphism. We claim that $R(x,U) + n = R(T^{-n}x,U) \cap \{n+1,n+2,\ldots\}$. Indeed, note that $m \in R(x,U) + n$ if and only if $m - n \in R(x,U)$ if and only if $T^{m-n}x \in U$ and $m \ge n+1$ if and only if $T^mT^{-n}x \in U$ and $m \ge n+1$ if and only if $m \in R(T^{-n}x,U) \cap \{n+1,n+2,\ldots\}$. By Lemma 3.2, the set $R(T^{-n}x,U) \cap \{n+1,n+2,\ldots\}$ is dynamically syndetic. Therefore, the set A + n is dynamically syndetic, demonstrating (i).

Since (X, T) is minimal and invertible, the system (X, T^{-1}) is minimal (see, for example, [26, Lemma 2.7]). We claim that the set $R_{T^{-1}}(x, U)$ is contained in the set in (3.2). This will finish the proof of (ii) and (iii) since $R_{T^{-1}}(x, U)$ is dynamically syndetic and dynamically central syndetic when $x \in U$. Suppose $n \in R_{T^{-1}}(x, U)$ so that $T^{-n}x \in U$. From the previous paragraph, we see that $R(T^{-n}x, U) \cap \{n+1, n+2, \ldots\} = R(x, U) + n \subseteq A+n$. By Lemma 3.2, we have that the set A + n is dynamically central syndetic, whereby n belongs to the set in (3.2), as desired.

Finally, that (3.3) holds follows immediately from (i) and (ii) for A - n and A + n.

It is interesting to formulate the conclusions of Lemma 3.4 in terms of the family algebra developed in Section 2.1. Thus,

- (i) the family dS is translation invariant, that is, $dS \subseteq \{\mathbb{N}\} + dS$;
- (ii) $d\mathcal{S} \subseteq d\mathcal{S} + dc\mathcal{S}$; and
- (iii) the family dcS is idempotent, that is, $dcS \subseteq dcS + dcS$.

The following lemma, well known to experts, describes the families of piecewise syndetic and central sets in dynamical terms. Both equalities can be derived from the result of Auslander [1] and Ellis [14] that any point in any system is proximal to a uniformly recurrent point. We opt for a short derivation of the first from the second, which is shown in [36].

Lemma 3.5. We have that $\mathcal{PS} = dS \sqcap \mathcal{T}$ and $\mathcal{C} = dcS \sqcap \mathcal{T}$.

Proof. The second equality is proven in [36, Thm. 3.7]. We will derive the first from the second. Suppose $A \subseteq \mathbb{N}$ is piecewise syndetic; it is not hard to see from the definition that there exists $N \in \mathbb{N}$ such that $A \cup (A + 1) \cup \cdots \cup (A + N)$ is thick. Since thick sets are central and the family of central sets is partition regular, some A+i is central. By the second equality, there exists $B \in dcS$ and $H \in \mathcal{T}$ such that $A + i \supseteq B \cap H$. Therefore, we have that $A \supseteq (B-i) \cap (H-i)$. By Lemma 3.4, the set B-i is dynamically syndetic and the set H-i is thick, as was to be shown.

For the following lemma, recall the dilation notation set out at the beginning of Section 2.

Lemma 3.6. Let $A \subseteq \mathbb{N}$ and $k \in \mathbb{N}$. If A is dynamically central syndetic, then so are the sets kA and A/k. If A is dynamically syndetic, then so is the set kA.

Proof. Let $A \in dcS$. Let (X,T) be a minimal system, $x \in X$, and $U \subseteq X$ be an open neighborhood of x such that $A \supseteq R(x,U)$. We will show that $kA \in dcS$ by considering a suspension system with base (X,T). More precisely, let $Y = X \times \{0, 1, \ldots, k-1\}$, and define $S: Y \to Y$ by

$$S(x',\eta) = \begin{cases} (x',\eta+1) \text{ if } \eta \leq k-2, \\ (Tx',0) \text{ if } \eta = k-1. \end{cases}$$

It is easy to see that the orbit of every point in Y is dense, whereby (Y, S) is minimal. Also, we see that $kR_T(x, U) = R_S((x, 0), U \times \{0\}) \in dc\mathcal{S}$, as desired.

To see that $A/k \in dc\mathcal{S}$, we appeal to a well-known fact (cf. [52, Cor. 3.2]): if a point in a system is uniformly recurrent under the transformation, then it is uniformly recurrent under every power of the transformation. Since (X, T) is minimal, the point x is uniformly recurrent under T, and hence uniformly recurrent under T^k . Therefore, the system $(\overline{T^{k\mathbb{N}_0}x}, T^k)$ is minimal, contains x, and $A/k \supseteq R_{T^k}(x, U \cap \overline{T^{k\mathbb{N}_0}x}) \in dc\mathcal{S}$, as desired.

Suppose $A \in dS$. By Lemma 3.4 (ii), there exists $n \in \mathbb{N}$ such that $A - n \in dcS$. By the first paragraph of this proof, we have $k(A - n) \in dcS$. Therefore, by Lemma 3.4 (i), we see $kA \supseteq k(A - n) + kn \in dS$, whereby $kA \in dS$, as desired.

3.1.3. Symbolic characterizations of dynamically syndetic sets

We show in this section that the family of dynamically (central) syndetic sets is not affected by requiring return time sets to come from zero-dimensional systems. This is accomplished by considering the indicator functions 1_A of sets $A \subseteq \mathbb{N}$ in the symbolic space $\{0, 1\}^{\mathbb{N}}$.

As a warm-up to the challenges that arise in Theorem 3.8, it is worth noting that when (X,T) is a minimal system, $x \in X$, and $U \subseteq X$ is nonempty and open, the point $1_{R(x,U)} \in \{0,1\}^{\mathbb{N}}$ is not always uniformly recurrent under the shift map σ . Indeed, here is an example.

Example 3.7. Let $\alpha \in (0,1)$ be irrational. Let $X := \mathbb{R}/\mathbb{Z}$ and $T : X \to X$ be defined by $T(x) = x + \alpha$. Let $U := X \setminus \{\alpha\}$ and x := 0. We see that

$$1_{R(0,U)} = (1_U(T^n x))_{n \in \mathbb{N}} = (1_U(\alpha), 1_U(2\alpha), 1_U(3\alpha), \ldots) = (0, 1, 1, \ldots).$$

Since α is irrational, for all $n \ge 2$, $T^n \alpha \ne \alpha$. Therefore, the function $1_{R(0,U)}$ takes the value 0 only at 1. The point $1_{R(0,U)} \in [0]_1$ is not uniformly recurrent because it never returns to the neighborhood $[0]_1$.

In the following theorem, point (ii) says that the set A is very strongly central, according to the terminology in [6, Def. 2.10].

Theorem 3.8. Let $A \subseteq \mathbb{N}$. The following statements are equivalent.

- (i) The set A is dynamically central syndetic.
- (ii) There exists a minimal system (X, T), a nonempty, open set $U \subseteq X$, and a point $x \in \overline{U}$ such that $R(x, U) \subseteq A$.
- (iii) There exists a subset $B \subseteq A$ such that the point $1_{B \cup \{0\}}$ is uniformly recurrent in the full shift $(\{0,1\}^{\mathbb{N}_0}, \sigma)$. In particular, the system $(X := \overline{\sigma^{\mathbb{N}_0} 1_{B \cup \{0\}}}, \sigma)$ is minimal, the set $X \cap [1]_0$ is a clopen neighborhood of $1_{B \cup \{0\}}$, and $B = R(1_{B \cup \{0\}}, X \cap [1]_0) \subseteq A$.

Proof. (i) \Rightarrow (ii) This follows by definition.

(ii) \Rightarrow (iii) Assume that $A \supseteq R(x, U)$, where (X, T) is a minimal system, $U \subseteq X$ is nonempty and open, and $x \in \overline{U}$.

If x is a periodic point of (X,T), let $n_0 \in \mathbb{N}$ be least such that $T^{n_0}x = x$. Since (X,T) is minimal, $X = \{x, Tx, \dots, T^{n_0-1}x\}$. As X is finite and $x \in \overline{U}$, it must be that $x \in U$. Letting $B = n_0 \mathbb{N}$ be the set of positive multiples of n_0 , we have that $A \supseteq R(x,U) \supseteq B$ and that $1_{B \cup \{0\}}$ is a periodic, hence uniformly recurrent, point in $\{0,1\}^{\mathbb{N}_0}$.

Suppose that x is not a periodic point of (X, T). We claim that there exists a nonempty, open set $V \subseteq U$ such that $x \in \overline{V}$ and the boundary ∂V is disjoint from $T^{\mathbb{N}}x := \{T^n x \mid n \in \mathbb{N}\}$.

If $x \in U$, choose $V = B_{\delta}(x)$ for some small $\delta > 0$. Note that the boundary of $B_{\delta}(x)$ is the set $\{z \in X \mid d_X(x, z) = \delta\}$, which may be empty. Since there are uncountably many choices for δ , there exists a choice so that the boundary is disjoint from $T^{\mathbb{N}}x$.

Suppose that $x \in \partial U = \overline{U} \setminus U$. There exists a sequence of pairwise distinct points $(x_k)_{k \in \mathbb{N}}$ in $U \setminus \{x\}$ such that $\lim_{k \to \infty} x_k = x$. Inductively, choose a sequence of disjoint balls $B_{\delta_k}(x_k) \subseteq U$ with $\lim_{k \to \infty} \delta_k = 0$ such that the boundary of each $B_{\delta_k}(x_k)$ has empty intersection with $T^{\mathbb{N}}x$. Let $V = \bigcup_{k=1}^{\infty} B_{\delta_k}(x_k)$. It is clear that V is an open subset of U. Since $\lim_{k \to \infty} x_k = x$, we have that $\partial V = \{x\} \cup \bigcup_{k=1}^{\infty} \partial V_k$, and so the boundary of V is disjoint from $T^{\mathbb{N}}x$. Our claim is proved.

Let B = R(x, V). Since V is nonempty and $V \subseteq U$, we have that B is nonempty and that $B \subseteq A$. It remains to show that $1_{B \cup \{0\}}$ is a uniformly recurrent point in $(\{0, 1\}^{\mathbb{N}_0}, \sigma)$. By the topology on the space $\{0, 1\}^{\mathbb{N}_0}$, we must show that for all $L \in \mathbb{N}$, there exists a syndetic set of positive integers m such that

for all
$$i \in \{0, \dots, L\}, \ 1_{B \cup \{0\}}(m+i) = 1_{B \cup \{0\}}(i).$$
 (3.5)

Let $L \in \mathbb{N}$. As $\partial V \cap T^{\mathbb{N}}x = \emptyset$, for every $n \in \mathbb{N}$, the point $T^n x$ is either in V or in $X \setminus \overline{V}$. For $n \in \{1, \ldots, L\}$, define

$$W_n = \begin{cases} V & \text{if } T^n x \in V \\ X \setminus \overline{V} & \text{if } T^n x \in X \setminus \overline{V} \end{cases}$$

It follows that $x \in \bigcap_{n=1}^{L} T^{-n} W_n$. Since each W_n is open, the set $\bigcap_{n=1}^{L} T^{-n} W_n$ is an open neighborhood of x. Because $x \in \overline{V}$, the set $V \cap \bigcap_{n=1}^{L} T^{-n} W_n$ is nonempty and open. Since (X, T) is minimal, the set

$$R\left(x,V\cap\bigcap_{n=1}^{L}T^{-n}W_{n}\right)$$

is syndetic, and all integers m in this set satisfy the condition in (3.5). This demonstrates that $1_{B\cup\{0\}}$ is uniformly recurrent under the shift, as desired.

(iii) \Rightarrow (i) Suppose A contains a subset B for which $1_{B\cup\{0\}}$ is uniformly recurrent in $(\{0,1\}^{\mathbb{N}_0},\sigma)$. The system $(X := \overline{\sigma^{\mathbb{N}_0} \mathbf{1}_{B\cup\{0\}}},\sigma)$ is minimal, and the cylinder set $X \cap [1]_0$ is open and contains $\mathbf{1}_{B\cup\{0\}}$. Since $A \supseteq B = R(\mathbf{1}_{B\cup\{0\}}, X \cap [1]_0)$, we have that A is dynamically central syndetic, as desired.

A version of the following theorem is given in [37, Prop. 2.3]. The derivation from Theorem 3.8 is short, so we give it here.

Theorem 3.9. Let $A \subseteq \mathbb{N}$. The following statements are equivalent.

- (i) The set A is dynamically syndetic.
- (ii) There exists a nonempty subset $B \subseteq A$ for which the point 1_B is uniformly recurrent in the full shift $(\{0,1\}^{\mathbb{N}}, \sigma)$.

Proof. (i) \Rightarrow (ii) Suppose A is dynamically syndetic. By Lemma 3.4, there exists $k \in \mathbb{N}$ such that A + k is dynamically central syndetic. By Theorem 3.8, there exists a set $B \subseteq A + k$ such that $1_{B\cup\{0\}}$ is uniformly recurrent in $(\{0,1\}^{\mathbb{N}_0},\sigma)$. Note that $B - k \subseteq A$. Since $1_{B\cup\{0\}}$ is uniformly recurrent in $(\{0,1\}^{\mathbb{N}_0},\sigma)$, we have that 1_B is a uniformly recurrent point in $(\{0,1\}^{\mathbb{N}},\sigma)$. Applying σ^k , we have that $\sigma^k 1_B = 1_{B-k}$ is uniformly recurrent, as was claimed.

(ii) \Rightarrow (i) Suppose A contains a nonempty subset B for which 1_B is uniformly recurrent in $(\{0,1\}^{\mathbb{N}}, \sigma)$. Let $k = \min B$. Since 1_B is uniformly recurrent, so is $\sigma^k 1_B = 1_{B-k}$. Since $k \in B$, we have that $1_{(B-k)\cup\{0\}}$ is uniformly recurrent in $\{0,1\}^{\mathbb{N}_0}$. By Theorem 3.8, the set $B - k \in dcS$. Since $A - k \supseteq B - k$, the set A - k is dynamically central syndetic. By Lemma 3.4, the set $A \supseteq (A - k) + k$ is dynamically syndetic, as desired.

3.1.4. Partitioning dynamically central syndetic sets: a proof of Theorem E

Considering some first examples of dynamically central syndetic sets, one may be led to believe that the intersection of two dcS sets is again a dcS set. This is true in some special cases: a result of Furstenberg [20, Thm. 9.11] implies, for example, that the intersection of the set of times of return of a distal point to a neighborhood of itself with any other dcS set is dcS. It is not, however, true in general, as we show in Theorem E. This theorem is best contextualized as a strengthening of [6, Thm. 2.13], which (when combined with Theorem 3.8) says that \mathbb{N} can be partitioned into infinitely many disjoint dcS sets.

Proof of Theorem E. Let $A \in dcS$. We will prove that A can be partitioned into two disjoint dcS sets. Then, by repeatedly partitioning the second set into disjoint dcS sets,

$$A = A_0 \cup A_1 = A_0 \cup A_{10} \cup A_{11} = A_0 \cup A_{10} \cup A_{110} \cup A_{111} = \cdots$$

we see that A can be partitioned into infinitely many disjoint dcS sets. Let (X,T) be a minimal system and $U \subseteq X$ be an open neighborhood of x such that $A \supseteq R(x,U)$. We consider two cases.

Case 1: the point x is an isolated point of X. By Lemma 2.9, the system (X,T) is finite and periodic. Let $(\mathbb{T} := \mathbb{R}/\mathbb{Z}, S)$ be the rotation on the one-dimensional torus by an irrational α . Let $V_1 = (0, 1/2), V_2 = (1/2, 1)$, which are two disjoint, open subsets of \mathbb{T} whose closures contain 0. Then $\{x\} \times V_1$ and $\{x\} \times V_2$ are two disjoint, open subsets of $X \times \mathbb{T}$ whose closures contain (x, 0). Note that the system $(X \times \mathbb{T}, T \times S)$ is minimal, and so by Theorem 3.8, the sets $R((x, 0), \{x\} \times V_1)$ and $R((x, 0), \{x\} \times V_2)$ are dcS. Moreover,

$$A\supseteq R(x,U)\supseteq R(x,\{x\})\supseteq R((x,0),\{x\}\times V_1)\cup R((x,0),\{x\}\times V_2).$$

Case 2: the point x is a limit point of U. Arguing as in the proof of Theorem 3.8, there exists a sequence of disjoint open balls $B_{\delta_k}(x_k)$ in U such that $x_k \to x$ and $\delta_k \to 0$. Take

$$V_0 = \bigcup_{k=1}^{\infty} B_{\delta_{2k}}(x_{2k})$$
 and $V_1 = \bigcup_{k=1}^{\infty} B_{\delta_{2k-1}}(x_{2k-1}).$

Then V_0, V_1 are disjoint open subsets of U satisfying $x \in \overline{V_0} \cap \overline{V_1}$. By Theorem 3.8, the sets $R(x, V_0)$ and $R(x, V_1)$ are dcS sets and, by construction, they are disjoint subsets of A. Put $A_0 := R(x, V_0)$ and $A_1 := A \setminus R(x, V_0)$ to see that $A = A_0 \cup A_1$ is a disjoint partition of A into dcS sets, as desired.

3.2. Central syndetic sets

In this section, we define the family of central syndetic sets and give a combinatorial characterization of them.

Definition 3.10. A set $A \subseteq \mathbb{N}$ is a *central syndetic* if it is a member of a syndetic, idempotent filter on \mathbb{N} .

Canonical examples of central syndetic sets are provided by dynamically central syndetic sets: return time sets R(x,U) where (X,T) is a minimal system, $x \in X$, and $U \subseteq X$ is an open neighborhood of x. That sets of this form are central syndetic is proven in the "only if" direction of Theorem 3.18 below. The "if" direction of that theorem shows that these are essentially the only examples of central syndetic sets.

Let us inject a brief historical remark to motivate the nomenclature. The family of central sets was described first by Furstenberg [20, Def. 8.3]. It was shown by Bergelson and Hindman [4] that a subset of \mathbb{N} is central if and only if it is a member of an idempotent ultrafilter on \mathbb{N} . To discuss these two families separately, it has become customary (see, eg., [32, Sec. 19.3]) to call Furstenberg's sets "dynamically central" and members of idempotent ultrafilters "central." We take a similar approach here: "dynamically central syndetic" sets are defined dynamically as above, while "central syndetic" sets are defined combinatorially as members of syndetic, idempotent filters. We will demonstrate in Theorem 3.18 that these families are the same.

3.2.1. Central syndetic sets are central

In this section, we justify the terminology "central syndetic" by showing that central syndetic sets are, in fact, both central and syndetic. That they are syndetic follows immediately from the definition – they are members of syndetic filters – so we need only to show that they are central. In what follows, for a family \mathscr{F} on \mathbb{N} , we denote by $\overline{\mathscr{F}}$ the set $\bigcap_{A \in \mathscr{F}} \overline{A}$ in $\beta \mathbb{N}$.

Lemma 3.11. For all syndetic filters \mathscr{F} on \mathbb{N} and all minimal left ideals $L \subseteq \beta \mathbb{N}$, we have $L \cap \overline{\mathscr{F}} \neq \emptyset$.

Proof. We will use the following well-known fact ([32, Thm. 4.48]): for all syndetic sets

 $A \subseteq \mathbb{N}$ and all minimal left ideals $L \subseteq \beta \mathbb{N}$, we have $\overline{A} \cap L \neq \emptyset$.

Let \mathscr{F} be a syndetic filter on \mathbb{N} , and let $L \subseteq \beta \mathbb{N}$ be a minimal left ideal. Consider the set $\{\overline{A} \cap L \mid A \in \mathscr{F}\}$. By the fact from the previous paragraph, the set consists of closed, nonempty subsets of L. Since \mathscr{F} is a filter, the set has the finite intersection property. Since L is compact, we have that $\overline{\mathscr{F}} \cap L$ is nonempty, as desired.

Theorem 3.12. For all syndetic, idempotent filters \mathscr{F} on \mathbb{N} and all minimal left ideals $L \subseteq \beta \mathbb{N}$, there exists an idempotent ultrafilter in L containing \mathscr{F} .

Proof. Let \mathscr{F} be a syndetic, idempotent filter on \mathbb{N} . By Lemma 3.11, there exists a minimal ultrafilter $p \in \beta \mathbb{N}$ such that $\mathscr{F} \subseteq p$. Define $L = \beta \mathbb{N} + p$. Since p is minimal, the set L is a minimal left ideal, and hence subsemigroup, of $(\beta \mathbb{N}, +)$. By [32, Thm. 2.6], the set L is compact. Therefore, the set L is a compact subsemigroup of $(\beta \mathbb{N}, +)$ containing p.

It follows from the proof of Theorem 2.6 that the set $\overline{\mathscr{F}}$ is a compact subsemigroup of $(\beta \mathbb{N}, +)$. It clearly contains p.

Define $X = L \cap \overline{\mathscr{F}}$. Since both L and $\overline{\mathscr{F}}$ are compact subsemigroups of $(\beta \mathbb{N}, +)$ containing p, so is X. By [32, Thm. 2.5], there exists an idempotent ultrafilter $q \in X$. Since $X \subseteq L$, we see that q is minimal. Since $X \subseteq \overline{\mathscr{F}}$, we see that $\mathscr{F} \subseteq q$. Thus, we have shown that \mathscr{F} is contained in a minimal, idempotent ultrafilter in L, as desired.

Theorem 3.13. Let $A \subseteq \mathbb{N}$ be central syndetic. For all minimal left ideals $L \subseteq \beta \mathbb{N}$, there exists an idempotent $p \in L$ such that $A \in p$. In particular, for all thick sets $H \subseteq \mathbb{N}$, the set $A \cap H$ is central.

Proof. Since A is central syndetic, there exists a syndetic, idempotent filter \mathscr{F} containing A. Let $L \subseteq \mathbb{N}$ be a minimal left ideal. By Theorem 3.12, there exists an idempotent $p \in L$ such that $\mathscr{F} \subseteq p$. In particular, we have that $A \in p$.

If $H \subseteq \mathbb{N}$ is thick, by [32, Thm. 4.48], there exists a minimal left ideal $L \subseteq \overline{H}$. By the reasoning in the previous paragraph, there exists an idempotent $p \in L$ containing A. Therefore, $A \cap H \in p$, whereby the set $A \cap H$ is central.

Remark 3.14. In [6], a set $A \subseteq \mathbb{N}$ is called *strongly central* if it has the property that for all minimal left ideals $L \subseteq \beta \mathbb{N}$, there exists an idempotent $p \in L$ containing A. Thus, Theorem 3.13 shows that central syndetic sets are strongly central. It will follow, however, from our main result in this section, Theorem 3.18, that strongly central sets need not be central syndetic. Indeed, a corollary of Theorems 3.8 and 3.18 is that the family of central syndetic sets is the same as the family of *very strongly central sets*, in the terminology of [6]. An example of a strongly central, but not very strongly central, set is given in [6, Thm. 2.17].

3.2.2. A combinatorial characterization of central syndetic sets

In this section and the next, we give a combinatorial characterization of central syndetic sets. Consider the following statement for a set $B \subseteq \mathbb{N}$:

for all finite
$$F \subseteq B$$
, the set $B \cap \bigcap_{f \in F} (B - f)$ is syndetic. (3.6)

The following theorem shows that the condition in (3.6) captures membership in a syndetic, idempotent filter.
Theorem 3.15. Let $A \subseteq \mathbb{N}$.

- (i) If A is central syndetic and \mathscr{F} is a syndetic, idempotent filter to which A belongs, then the set $A \cap (A - \mathscr{F})$ satisfies the condition in (3.6).
- (ii) If A satisfies the condition in (3.6), then the family

$$\mathscr{F} := \uparrow \left\{ A \cap \bigcap_{f \in F} (A - f) \mid F \subseteq A \text{ is finite} \right\}$$

is a syndetic, idempotent filter that contains A, whereby A is central syndetic. In particular, the set A is central syndetic if and only if some subset of it satisfies the condition in (3.6).

Proof. (i) Suppose A belongs to a syndetic, idempotent filter \mathscr{F} . Define $A_1 := A \cap (A - \mathscr{F})$. We claim that this set satisfies the condition in (3.6).

First we will show that $A_1 \in \mathscr{F}$ and $A_1 \subseteq A_1 - \mathscr{F}$. That $A_1 \in \mathscr{F}$ follows because $A \in \mathscr{F}$ and $A - \mathscr{F} \in \mathscr{F}$ (since \mathscr{F} is idempotent). Because $\mathscr{F} \subseteq \mathscr{F} + \mathscr{F}$, using the algebra in Lemma 2.3, we see

$$A_{1} = A \cap (A - \mathscr{F}) \subseteq A - \mathscr{F} \subseteq (A - \mathscr{F}) \cap (A - (\mathscr{F} + \mathscr{F}))$$
$$= (A - \mathscr{F}) \cap ((A - \mathscr{F}) - \mathscr{F})$$
$$= (A \cap (A - \mathscr{F})) - \mathscr{F}$$
$$= A_{1} - \mathscr{F}.$$

To see that the condition in (3.6) is satisfied, let $F \subseteq A_1$ be finite. Since $F \subseteq A_1 - \mathscr{F}$, the set $\bigcap_{f \in F} (A_1 - f) \in \mathscr{F}$. Since $A_1 \in \mathscr{F}$, we have that $A_1 \cap \bigcap_{f \in F} (A_1 - f) \in \mathscr{F}$, whereby it is syndetic, as desired.

(ii) That \mathscr{F} is a filter is immediate from its definition. That $A \in \mathscr{F}$ follows immediately from the fact that \mathscr{F} is upward closed.

To see that \mathscr{F} is syndetic, it suffices to show that for all $F \subseteq A$ finite, the set $B := A \cap \bigcap_{f \in F} (A - f)$ is syndetic. But this is immediate from (3.6).

To see that \mathscr{F} is idempotent, we must show that $\mathscr{F} \subseteq \mathscr{F} + \mathscr{F}$. Since $\mathscr{F} + \mathscr{F}$ is a filter, it suffices to show that for all $a \in A \cup \{0\}$, the set $A - a \in \mathscr{F} + \mathscr{F}$.

Let $a \in A \cup \{0\}$. We will show that $(A - a) - \mathscr{F} \in \mathscr{F}$. Since $A - a \in \mathscr{F}$, it suffices to show that $A - a \subseteq (A - a) - \mathscr{F}$. This holds since for any $n \in A - a$, we have that (A - a) - n = A - (n + a), which is a member of \mathscr{F} by the definition of \mathscr{F} since $n + a \in A$. \Box

Remark 3.16. It is a short exercise to show that the condition in (3.6) is equivalent to the following: there exists a chain of syndetic sets $A \supseteq A_1 \supseteq A_2 \supseteq A_3 \supseteq \cdots$ with the property that for all $n \in \mathbb{N}$ and all $a \in A_n$, there exists $m \in \mathbb{N}$ such that $A_n - a \supseteq A_m$. The combinatorial characterization of central sets in [32, Thm. 14.25] says the same thing with with "syndetic" replaced by "collectionwise piecewise syndetic."

We finish this section by showing that sets $B \subseteq \mathbb{N}$ satisfying the condition in (3.6) satisfy the ostensibly stronger condition:

for all finite
$$F \subseteq B$$
 and all $m \in \mathbb{N}$, the set $B \cap \bigcap_{f \in F} (B - f) \cap m\mathbb{N}$ is syndetic. (3.7)

This upgrade will help us with the proof of Theorem 3.18 in the next section.

Lemma 3.17. A set $A \subseteq \mathbb{N}$ satisfies the condition in (3.6) if and only if it satisfies the condition in (3.7).

Proof. If A satisfies the condition in (3.7), then it clearly satisfies the one in (3.6).

Conversely, suppose that $A \subseteq \mathbb{N}$ satisfies the condition in (3.6). To see that it satisfies (3.7), let $F \subseteq A$ be finite and $m \in \mathbb{N}$. To see that the set $A \cap \bigcap_{f \in F} (A - f) \cap m\mathbb{N}$ is syndetic, we will show that it has nonempty intersection with all thick sets.

Let $H \subseteq \mathbb{N}$ be thick. By Theorem 3.15, we see that the set $A \cap \bigcap_{f \in F} (A - f)$ belongs to a syndetic, idempotent filter, and hence is central syndetic. It follows from Theorem 3.13 that the set $A \cap \bigcap_{f \in F} (A - f) \cap H$ is central. The set $m\mathbb{N}$ is \mathcal{IP}^* , hence \mathcal{C}^* , and so the set $A \cap \bigcap_{f \in F} (A - f) \cap m\mathbb{N} \cap H$ is nonempty, as was to be shown. \Box

3.3. Central and dynamically central syndetic sets

We show in this section that the family of dynamically central syndetic sets and central syndetic sets are the same. Combined with Theorem 3.15, we get a combinatorial characterization of the family dcS.

Theorem 3.18. A set $A \subseteq \mathbb{N}$ is dynamically central syndetic if and only if it is central syndetic, that is, a member of a syndetic, idempotent filter on \mathbb{N} .

The proof of Theorem 3.18 is broken into two parts, the first of which we give now.

Proof of the "only if" statement in Theorem 3.18. By assumption, there exists a minimal system (X,T), a point $x \in X$, and an open set $U \subseteq X$ containing x such that $R(x,U) \subseteq A$. Define

$$\mathscr{F} := \uparrow \{ R(x, V) \mid V \subseteq X \text{ is an open set containing } x \}.$$

We claim that \mathscr{F} is a syndetic, idempotent filter containing A.

That \mathscr{F} is a filter, that $A \in \mathscr{F}$, and that \mathscr{F} is syndetic are all immediate. To see that \mathscr{F} is idempotent, it suffices by Remark 2.5 to show that for all open $V \subseteq X$ with $x \in V$, we have that $R(x,V) \subseteq R(x,V) - \mathscr{F}$. Since $R(x,V) \in \mathscr{F}$, this shows that $R(x,V) - \mathscr{F} \in \mathscr{F}$, as desired.

Let $V \subseteq X$ be open and contain x. To see that $R(x, V) \subseteq R(x, V) - \mathscr{F}$, let $n \in R(x, V)$. Thus, $x \in T^{-n}V$. Since $T^{-n}V$ is an open neighborhood of x, we have that $R(x, V) - n = R(x, T^{-n}V) \in \mathscr{F}$, as desired.

The rest of this section is devoted to a proof of the converse, namely that central syndetic sets are dynamically central syndetic. The argument requires a rather lengthy setup.

3.3.1. The main idea

Here is the main idea behind the proof of the "if" statement in Theorem 3.18. By Theorem 3.15, we may assume that the set A satisfies the condition in (3.6).

Our aim is to show that the set A contains a set of the form R(x, U), where the point xbelongs to the open set U in a minimal system. In view of Theorem 3.8, it suffices to find a subset $B \subseteq A$ for which $1_{B \cup \{0\}} \in \{0, 1\}^{\mathbb{N}_0}$ is uniformly recurrent under the shift map σ . Note that the condition in (3.6) is not sufficient for $1_{A \cup \{0\}}$ to be uniformly recurrent under σ . (Indeed, the set $\mathbb{N}_0 \setminus \{n^2 \mid n \in \mathbb{N}\}$ satisfies the condition in (3.6) but has an indicator function that is not uniformly recurrent under σ .) To find the set B, the idea is to modify the shift map σ into a "shift-punch" map κ under which the point 1_A becomes uniformly recurrent.

We append the integer 0 to A and consider it as a subset of \mathbb{N}_0 . The "punch map" $\pi : \{0,1\}^{\mathbb{N}_0} \to \{0,1\}^{\mathbb{N}_0}$, which depends on the set A, is described roughly as follows. At $1_C \in \{0,1\}^{\mathbb{N}_0}$, we find the maximal $N \in \mathbb{N}$ such that $A \cap [N] \subseteq C$. Then, $\pi(1_C) = 1_D$, where $D \cap [N] = A \cap [N]$ and $D \cap (\mathbb{N}_0 \setminus [N]) = C \cap (\mathbb{N}_0 \setminus [N])$. Roughly, the map π "punches out" or "deletes" the elements of C on the interval [N] that are not in $A \cap [N]$. The map π continuously moves points of $\{0,1\}^{\mathbb{N}_0}$ closer to 1_A .

The shift-punch map κ is defined to be $\pi \circ \sigma$. The dynamics of the shift-punch system $(\{0,1\}^{\mathbb{N}_0},\kappa)$ is thus a shift followed by a punch. The idea is to show that in this system, the point 1_A is uniformly recurrent. This happens because the set A satisfies the condition in (3.6) and thus syndetically often contains an initial interval of itself. Keeping track of which elements of A are deleted as the shift-punch map is repeatedly applied to 1_A , we end up with a subset $B \subseteq A$. We will show that the point $1_B \in \{0,1\}^{\mathbb{N}_0}$ is uniformly recurrent under the usual shift map, as desired.

The outline just given is difficult to control in practice because the punches can seemingly overlap in complicated ways. To overcome this, we find it useful to have more control over the length of intervals on which the punches occurs. Thus, the actual shift-punch system is defined as a skew product over a 2-adic odometer. The odometer functions only to control the length of the punch intervals, which are restricted to occur only on dyadic intervals. Dyadic intervals are convenient because a nonempty intersection implies containment. This control on the punch intervals allows us to realize the outline above.

3.3.2. Notation and spaces

In the rest of this section, we will alternate between working in \mathbb{N} and $\mathbb{N}_0 := \{0, 1, 2, ...\}$. Recall that for $N \in \mathbb{N}_0$, we write $[N] := \{0, ..., N - 1\}$. When $A \subseteq \mathbb{N}_0$, we compute the translate A - n as a subset of \mathbb{N}_0 .

It will be convenient to append some elements to \mathbb{N}_0 . Thus, we define $\mathbb{N}_{0,\pm\infty} := \mathbb{N}_0 \cup \{-\infty,\infty\}$. Subspaces, such as $\mathbb{N}_{0,\infty} := \mathbb{N}_0 \cup \{\infty\}$, are defined analogously. We give $\mathbb{N}_{0,\pm\infty}$ the topology that makes the bijection $\mathbb{N}_{0,\pm\infty} \to \{2\} \cup \{1/n \mid n \in \mathbb{N}\} \cup \{0\}$, where $-\infty \mapsto 2$, $x \mapsto 1/(x+1)$, and $\infty \mapsto 0$ a homeomorphism. Thus, every point of $\mathbb{N}_{0,\pm\infty}$, except for ∞ , is isolated. Subspaces of $\mathbb{N}_{0,\pm\infty}$ are given the subspace topology.

3.3.3. Dyadic valuation and intervals

Denote by $\nu_2 : \mathbb{Z} \to \mathbb{N}_{0,\infty}$ the two-adic valuation on \mathbb{Z} , defined by $\nu_2(n) := \sup\{k \in \mathbb{N}_0 \mid 2^k \text{ divides } n\}$. A finite interval $\{a, a + 1, \ldots, a + \ell - 1\}$ in \mathbb{N}_0 is a *dyadic interval* if $\ell = 2^k$ for some $k \in \mathbb{N}_0$ and $\ell \mid a$. Denote by \mathcal{Q}_{2^n} the set of dyadic intervals of cardinality 2^n . We will say that \mathbb{N}_0 is the dyadic interval of infinite cardinality.

The following lemma follows from standard facts about dyadic intervals. We provide a proof for completeness and ease of reference for later.

Lemma 3.19. Let Q and P be finite dyadic intervals in \mathbb{N}_0 .

- (i) $Q \cap 2^{\nu_2(\min Q)} \mathbb{N} = \{\min Q\}.$
- (ii) For all $\ell \in \mathbb{N}_0$, if $(Q \setminus \{\min Q\}) \cap 2^{\ell} \mathbb{N} \neq \emptyset$, then $|Q| \ge 2^{\ell+1}$.
- (iii) If $P \cap Q \neq \emptyset$, then either $P \subseteq Q$ or $Q \subseteq P$.

Proof. (i) There exists $n \in \mathbb{N}$ and $0 \leq k \leq \nu_2(n)$ such that $Q = n + [2^k]$. Since Q is an interval beginning at $n \in 2^{\nu_2(n)} \mathbb{N}$ with $|Q| = 2^k \leq 2^{\nu_2(n)}$, we have that $Q \cap 2^{\nu_2(n)} \mathbb{N} = \{n\}$.

(ii) Write $Q = n + [2^k]$ as in (i). Let $\ell \in \mathbb{N}_0$, and suppose that $(Q \setminus \{n\}) \cap 2^{\ell} \mathbb{N}$ is nonempty. It follows from the previous paragraph that $\ell < \nu_2(n)$. Let $a \in (Q \setminus \{n\}) \cap 2^{\ell} \mathbb{N}$. Since $|Q| = 2^k > a - n$ and $\nu_2(a - n) \ge \ell$, we see that $k \ge \ell + 1$. Therefore, $|Q| \ge 2^{\ell+1}$, as desired.

(iii) Let P and Q be dyadic intervals. Note that if $\min P = \min Q$, then $P \subseteq Q$ or $Q \subseteq P$, and we are done. Otherwise, we will show that if $\min P \in Q \setminus \{\min Q\}$, then $P \subseteq Q$. It will follow that either P and Q are disjoint or that one is contained in the other.

Suppose for a contradiction that $\min P \in Q \setminus \{\min Q\}$ and $P \not\subseteq Q$. Then $\max Q + 1 = \min Q + |Q| \in P$ and is divisible by |Q|. It follows from (ii) that $|P| \ge 2|Q|$. But since $P \not\subseteq Q$ and |P| divides $\min P$, we see by (ii) that $|Q| \ge 2|P|$, a contradiction.

We denote by $\mathcal{Q}_{2^n}^*$ the collection of subsets of \mathbb{N} that have nonempty intersection with all members of \mathcal{Q}_{2^n} . (This notation matches notation for the dual family defined in Section 2.1.1, but note that \mathcal{Q}_{2^n} is not a family since it is not upward closed.)

Lemma 3.20. A set $A \subseteq \mathbb{N}_0$ is syndetic if and only if there exists $k \in \mathbb{N}$ such that $A \in \mathcal{Q}_{2^k}^*$.

Proof. This is left to the reader as an easy exercise.

3.3.4. The 2-adic odometer

We denote by (Z, ρ) the 2-adic odometer, the inverse limit of the family of rotations $(\mathbb{Z}/2^k\mathbb{Z}, +1)$, $k \in \mathbb{N}_0$, as topological dynamical systems. We do not have need to specify a metric on Z explicitly; it suffices for our purposes to know that two elements of Z are near if and only if their projections to $\mathbb{Z}/2^k\mathbb{Z}$ agree for some large value of k.

For $n \in \mathbb{Z}$, we denote by n the element of Z that projects, for all $k \in \mathbb{N}_0$, to $n \in \mathbb{Z}/2^k\mathbb{Z}$. This association gives a dense embedding of \mathbb{Z} into Z. By a convenient abuse of notation, we write $\mathbb{Z} \subseteq Z$, identifying \mathbb{Z} with this copy in Z.

The two-adic valuation $\nu_2 : \mathbb{Z} \to \mathbb{N}_{0,\infty}$ is uniformly continuous in the subspace topology that \mathbb{Z} inherits from the space Z. Indeed, if the images of integers n and m are the same in $\mathbb{Z}/2^k\mathbb{Z}$ for a large value of k, then 2^k divides n-m, so $\nu_2(n)$ is large if and only if $\nu_2(m)$ is large. Therefore, the map $\nu_2 : \mathbb{Z} \to \mathbb{N}_{0,\infty}$ extends uniquely to a continuous map $\nu_2 : \mathbb{Z} \to \mathbb{N}_{0,\infty}$.

3.3.5. The punch maps

Let $X = \{0, 1\}^{\mathbb{N}_0}$, and fix $A \subseteq \mathbb{N}_0$. We will define a family of continuous maps $\pi_n : X \to X$, $n \in \mathbb{N}_{0,\infty}$, depending on A that drive the shift-punch dynamics described in the next section.

First, we define a map $\nu_A: X \to \mathbb{N}_{0,\pm\infty}$ in the following way. For $\omega \in X$,

$$\nu_A(\omega) := \sup \left\{ n \in \mathbb{N}_0 \mid A \cap [2^n] \subseteq \operatorname{supp}(\omega) \right\},\tag{3.8}$$

where the supremum of the empty set is $-\infty$. Note that $\nu_A(\omega) = -\infty$ if and only if $0 \in A$ and $\omega(0) = 0$. Also, note that $\nu_A(\omega) = \infty$ if and only if $A \subseteq \text{supp}(\omega)$.

For $n \in \mathbb{N}_{0,\infty}$ and $\omega \in X$, we define the *punch interval*

$$I_n(\omega) := \left[2^{\min(\nu_A(\omega),n)}\right],$$

where $[2^{-\infty}]$ and $[2^{\infty}]$ are interpreted to be the empty set and \mathbb{N}_0 , respectively. Then, we define the *punch map* $\pi_n : X \to X$ by

$$(\pi_n(\omega))(i) = \begin{cases} 1_A(i) & \text{if } i \in I_n(\omega) \\ \omega(i) & \text{if } i \in \mathbb{N}_0 \backslash I_n(\omega) \end{cases}.$$

Informally, if $A \cap [2^m] \subseteq \operatorname{supp}(\omega)$, where $m \in \{-\infty, 0, 1, \ldots, n\}$ is maximal as such, then $\pi_n(\omega)$ is defined so that $\operatorname{supp}(\pi_n(\omega)) \cap [2^m] = A \cap [2^m]$ and $\operatorname{supp}(\pi_n(\omega)) \cap (\mathbb{N}_0 \setminus [2^m]) = \operatorname{supp}(\omega) \cap (\mathbb{N}_0 \setminus [2^m])$. The word "punch" was chosen since the punch maps "punch out" ones by changing them to zeroes.

Lemma 3.21. Each of the maps $\pi_n : X \to X$, $n \in \mathbb{N}_{0,\infty}$, is continuous, and the map $\mathbb{N}_{0,\infty} \to C(X,X)$ given by $n \mapsto \pi_n$ is continuous, where C(X,X) has the supremum topology.

Proof. Let $n \in \mathbb{N}_{0,\infty}$. To see that $\pi_n : X \to X$ is continuous, let $\omega \in X$ and $N \in \mathbb{N}$. If $\nu_A(\omega) \in \mathbb{N}_{0,-\infty}$, then for all $\omega' \in X$ sufficiently close to ω , the points ω' and ω agree on [N] and $\nu_A(\omega') = \nu_A(\omega)$. If, on the other hand, $\nu_A(\omega) = \infty$, then for all $\omega' \in X$ sufficiently close to ω , the points ω' and ω agree on [N] and $\nu_A(\omega') \ge N$. In both cases, we see that $\pi_n(\omega')$ and $\pi_n(\omega)$ agree on [N]. By the topology of the space X, this shows that π_n is continuous.

To show that the map $\mathbb{N}_{0,\infty} \to C(X,X)$ given by $n \mapsto \pi_n$ is continuous, it suffices by the topology on $\mathbb{N}_{0,\infty}$ to show that $\lim_{n\to\infty} \pi_n = \pi_\infty$, that is, for all $\varepsilon > 0$ and all sufficiently large $n \in \mathbb{N}$,

$$\sup_{\omega \in X} d_X \big(\pi_n(\omega), \pi_\infty(\omega) \big) < \varepsilon.$$

Let $\varepsilon > 0$ and $\omega \in X$. If $\nu_A(\omega) = -\infty$, then for all $n \in \mathbb{N}_0$, $\pi_\infty(\omega) = \pi_n(\omega)$. If $\nu_A(\omega) \in \mathbb{N}_0$, then for all $n \in \mathbb{N}_0$ with $n \ge \nu_A(\omega)$, $\pi_\infty(\omega) = \pi_n(\omega)$. If $\nu_A(\omega) = \infty$, then $\pi_n(\omega)$ and $\pi_\infty(\omega)$ match on $[2^n]$, because they both match 1_A on $[2^n]$. Thus, in all cases, for $n \in \mathbb{N}_0$ sufficiently large, $d_X(\pi_n(\omega), \pi_\infty(\omega)) < \varepsilon$, as was to be shown.

The following lemma gives finer information on the continuity of π_n , $n \in \mathbb{N}_0$.

Lemma 3.22. Let $n, \ell \in \mathbb{N}_0$ with $\ell \ge 2^n$. If $\omega, \xi \in X$ agree on $[\ell]$, then $I_n(\omega) = I_n(\xi)$, and $\pi_n(\omega)$ and $\pi_n(\xi)$ agree on $[\ell]$.

Proof. Suppose that $\omega, \xi \in X$ agree on $[\ell]$. We consider the following cases.

- (i) Case 1: $\nu_A(\omega) = -\infty$ or $\nu_A(\xi) = -\infty$. In this case, since ω and ξ agree on $[\ell]$, we have that $\nu_A(\omega) = \nu_A(\xi) = -\infty$ and $I_n(\omega) = I_n(\xi) = \emptyset$. Since ω and ξ are unchanged by π_n , we have that $\pi_n(\omega)$ and $\pi_n(\xi)$ agree on $[\ell]$.
- (ii) Case 2: $0 \leq \nu_A(\omega) < n$. By the definition of ν_A , we have that $A \cap [2^{\nu_A(\omega)}] \subseteq \text{supp}(\omega)$ but $A \cap [2^{\nu_A(\omega)+1}] \not\subseteq \text{supp}(\omega)$. Since $2^{\nu_A(\omega)} < 2^n \leq \ell$, we have that $[2^{\nu_A(\omega)+1}] \subseteq [\ell]$. Since ω and ξ agree on $[\ell]$, we see that $\nu_A(\omega) = \nu_A(\xi)$, and hence that $I_n(\omega) = I_n(\xi)$. Since π_n changes ω and ξ on the same interval contained in $[\ell]$, we see that $\pi_n(\omega)$ and $\pi_n(\xi)$ agree on $[\ell]$, as desired.
- (iii) Case 3: $0 \le \nu_A(\xi) < n$. We argue just as in Case 2 to arrive at the same conclusion.
- (iv) Case 4: $\nu_A(\omega) \ge n$ and $\nu_A(\xi) \ge n$. In this case, we see that $I_n(\omega) = I_n(\xi) = [2^n]$. Since $2^n \le \ell$ and ω matches ξ on $[\ell]$, we see that $\pi_n(\omega)$ and $\pi_n(\xi)$ agree on $[\ell]$, as desired.

This concludes the casework and the proof of the lemma.

3.3.6. The shift-punch system

The shift-punch system $(Z \times X, \kappa)$ is the topological skew product system defined by the shift-punch map $\kappa : Z \times X \to Z \times X$,

$$\kappa(z,\omega) := \left(\rho z, \pi_{\nu_2(\rho z)}(\sigma \omega)\right)$$

Though not apparent from the notation, the punch maps π_n , the shift-punch map κ , and the shift-punch system $(Z \times X, \kappa)$ all depend highly on the set A.

Lemma 3.23. The shift-punch system $(Z \times X, \kappa)$ is a topological dynamical system.

Proof. The space $Z \times X$ is compact, so we need only to show that the map κ is continuous. Because κ is a skew product, it suffices to check that the map $Z \to C(X, X)$ defined by $z \mapsto \pi_{\nu_2(\rho z)} \circ \sigma$ is continuous. This follows from the fact that the shift $\sigma : X \to X$ is continuous and that the map $z \mapsto \pi_{\nu_2(\rho z)}$ is a composition of three continuous maps: $\rho : Z \to Z$, $\nu_2 : Z \to \mathbb{N}_{0,\infty}$, and, by Lemma 3.21, $\pi : \mathbb{N}_{0,\infty} \to C(X, X)$.

We are interested primarily in the orbit of the point $(0, 1_A)$ under the shift-punch map. In the shift-punch dynamics, it will be important to keep track of where the punches (changes from 1 to 0) occur. The following notation will assist with that. Define $\beta^{(0)} = 1_{\mathbb{N}_0}$ and $\alpha^{(0)} = 1_A$, and for $n \in \mathbb{N}$, define $\beta^{(n)}, \alpha^{(n)} \in X$ to be such that

$$(\rho \times \sigma)\kappa^{n-1}(0, 1_A) = (n, \beta^{(n)}),$$

$$\kappa^n(0, 1_A) = (n, \alpha^{(n)}).$$

By the definition of κ , we see that for all $n \in \mathbb{N}_0$, $\alpha^{(n)} = \pi_{\nu_2(n)}\beta^{(n)}$ and $\beta^{(n+1)} = \sigma\alpha^{(n)}$. Thus, the points $\beta^{(n)}$ and $\alpha^{(n)}$ are the second coordinates of $\kappa^n(0, 1_A)$ before and after, respectively, the punch by $\pi_{\nu_2(n)}$.

The punch window at $n \in \mathbb{N}_0$ is defined to be

$$W(n) := n + I_{\nu_2(n)}(\beta^{(n)}),$$

where W(n) is understood to be the empty set if $I_{\nu_2(n)}(\beta^{(n)})$ is the empty set. Note that $W(0) = \mathbb{N}_0$; that W(n) is an interval beginning at n of length a divisor of $2^{\nu_2(n)}$ (and, hence, is a dyadic interval); and that $\alpha^{(n)}$ and 1_A agree on the interval W(n) - n. Informally, we note that $\sup(\beta^{(n)}) \subseteq A - n$ (as $\beta^{(n)}$ is the result of the first n - 1 shift-punches and a single shift) just before $\pi_{\nu_2(n)}$ is applied to change $\beta^{(n)}$ on the punch interval $I_{\nu_2(n)}(\beta^{(n)})$ to yield $\alpha^{(n)}$, the second coordinate of $\kappa^n(0, 1_A)$. The interval W(n) is the interval in the original set A at which that punch occurs.

Lemma 3.24. For all $n \in \mathbb{N}_0$, (i)

$$\operatorname{supp}\left(\beta^{(n)}\right) = \bigcap_{m=0}^{n-1} \left(\left(A - (n-m)\right) \cup \left(\mathbb{N}_0 \setminus \left(W(m) - n\right)\right) \right),$$
(3.9)

where the empty intersection is interpreted as \mathbb{N}_0 ;

(ii)

$$\operatorname{supp}\left(\alpha^{(n)}\right) = \bigcap_{m=0}^{n} \left(\left(A - (n-m)\right) \cup \left(\mathbb{N}_{0} \setminus \left(W(m) - n\right)\right) \right).$$
(3.10)

Proof. We will prove both (i) and (ii) simultaneously by induction on $n \in \mathbb{N}_0$. The base case n = 0 follows by set algebra, recalling that, by definition, $\beta^{(0)} = 1_{\mathbb{N}_0}$ and $\alpha^{(0)} = 1_A$.

Suppose both (i) and (ii) hold for some $n \in \mathbb{N}_0$. Let $i \in \mathbb{N}_0$. Since $\beta^{(n+1)} = \sigma \alpha^{(n)}$, we see that $(\beta^{(n+1)})(i) = 1$ if and only if $(\alpha^{(n)})(i+1) = 1$. By the inductive hypothesis, we have that $(\alpha^{(n)})(i+1) = 1$ if and only if

$$i+1 \in \bigcap_{m=0}^{n} \left(\left(A - (n-m) \right) \cup \left(\mathbb{N}_0 \setminus \left(W(m) - n \right) \right) \right).$$

The previous line rearranges to

$$i \in \bigcap_{m=0}^{n} \left(\left(A - (n+1-m) \right) \cup \left(\mathbb{N}_0 \setminus \left(W(m) - (n+1) \right) \right) \right), \tag{3.11}$$

which shows (i) for n + 1.

To see (ii) for n + 1, let $i \in \mathbb{N}_0$. Note that by the definition of the map $\pi_{\nu_2(n+1)}$, $(\alpha^{(n+1)})(i) = (\pi_{\nu_2(n+1)}\beta^{(n+1)})(i) = 1$ if and only if $(\beta^{(n+1)})(i) = 1$ and either

- $i \notin I_{\nu_2(n+1)}(\beta^{(n+1)}) = W(n+1) (n+1)$, so that $\pi_{\nu_2(n+1)}$ does not modify $\beta^{(n+1)}$ at *i*; or
- $i \in A$, so that, if $\pi_{\nu_2(n+1)}$ modifies $\beta^{(n+1)}$ at i, it does not change 1 to 0.

By the previous paragraph, we have that $(\beta^{(n+1)})(i) = 1$ if and only if (3.11) holds, and so we see by the previous sentence that $(\alpha^{(n+1)})(i) = 1$ if and only if *i* belongs to the set

$$\bigcap_{m=0}^{n} \left(\left(A - (n+1-m) \right) \cup \left(\mathbb{N}_0 \setminus \left(W(m) - (n+1) \right) \right) \right) \cap \left(A \cup \left(\mathbb{N}_0 \setminus \left(W(n+1) - (n+1) \right) \right) \right).$$

The set on the previous line simplifies to the one in (3.10) for n + 1, demonstrating (ii) for n + 1, as desired.

Lemma 3.25. Let $m \in \mathbb{N}_0$. For all $i \in W(m) - m$, the points $\alpha^{(i)}$ and $\alpha^{(m+i)}$ agree on the interval W(m) - m - i.

Proof. The conclusion is trivial for m = 0, so suppose $m \in \mathbb{N}$. Let $\ell = |W(m)|$. We must show that for all $i \in [\ell]$, the points $\alpha^{(i)}$ and and $\alpha^{(m+i)}$ agree on $W(m) - m - i = [\ell - i]$. We will prove this by induction on i. The base case i = 0 follows from the definition of W(m): the points $\alpha^{(0)} = 1_A$ and $\alpha^{(m)}$ agree on $W(m) - m = [\ell]$.

Let $i \in [\ell - 1]$, and suppose that $\alpha^{(i)}$ and $\alpha^{(m+i)}$ agree on $[\ell - i]$. We will show that $\alpha^{(i+1)}$ and $\alpha^{(m+i+1)}$ agree on $[\ell - i - 1]$. Note that $\alpha^{(i+1)} = \pi_{\nu_2(i+1)}\sigma\alpha^{(i)}$ and $\alpha^{(m+i+1)} = \pi_{\nu_2(m+i+1)}\sigma\alpha^{(m+i)}$.

Since $\alpha^{(i)}$ and $\alpha^{(m+i)}$ agree on $[\ell - i]$, we have that $\sigma \alpha^{(i)}$ and $\sigma \alpha^{(m+i)}$ agree on $[\ell - i - 1]$. If we show that $\nu_2(i+1) = \nu_2(m+i+1)$ and that $2^{\nu_2(m+i+1)} \leq \ell - i - 1$, then it will follow from Lemma 3.22 that $\pi_{\nu_2(i+1)}\sigma\alpha^{(i)}$ and $\pi_{\nu_2(m+i+1)}\sigma\alpha^{(m+i)}$ agree on $[\ell - i - 1]$, concluding the proof of the inductive step. Since W(m) is a dyadic interval, $m + i + 1 \in (W(m) \setminus \{m\}) \cap 2^{\nu_2(m+i+1)} \mathbb{N}$, and $|W(m)| \leq 2^{\nu_2(m)}$, we get from Lemma 3.19 that $\nu_2(m + i + 1) < \nu_2(m)$. By properties of the 2-adic valuation, it follows that $\nu_2(m + i + 1) = \nu_2(i + 1)$. To see that $2^{\nu_2(m+i+1)} \leq \ell - i - 1$, note that $m + i + 1 + [2^{\nu_2(m+i+1)}]$ is a dyadic interval, intersecting, hence contained in, W(m). Therefore, $m + i + 1 + 2^{\nu_2(m+i+1)} \leq m + |W(m)| = m + \ell$, which rearranges to the desired inequality, finishing the proof of the inductive step.

The following lemma shows the "fractal" structure of containment amongst the punch windows.

Lemma 3.26. For all $n, m \in \mathbb{N}_0$, if m < n and $n \in W(m)$, then W(n) - m = W(n - m).

Proof. Let $n, m \in \mathbb{N}_0$ with m < n and $n \in W(m)$. If m = 0, then conclusion is immediate, so suppose $m \ge 1$. Define $\ell = |W(m)|$.

We claim that $\nu_2(n) = \nu_2(n-m)$ and that $2^{\nu_2(n)} \leq \ell - (n-m)$. Indeed, since W(m) is a dyadic interval, $n \in (W(m) \setminus \{m\}) \cap 2^{\nu_2(n)} \mathbb{N}$, and $2^{\nu_2(m)} \geq |W(m)|$, we see from Lemma 3.19 that $\nu_2(n) < \nu_2(m)$. By properties of the 2-adic valuation, this implies that $\nu_2(n) = \nu_2(n-m)$. Since $n + [2^{\nu_2(n)}]$ is a dyadic interval intersecting, and hence contained in, W(m), we see that $n + 2^{\nu_2(n)} \leq m + |W(m)| = m + \ell$. Rearranging, we have that $2^{\nu_2(n)} \leq \ell - (n-m)$.

Next, we claim that $\beta^{(n)}$ and $\beta^{(n-m)}$ agree on $[\ell - n + m]$. Indeed, it follows from Lemma 3.25 with $i = n - m - 1 \in [\ell]$ that the points $\alpha^{(n-m-1)}$ and $\alpha^{(n-1)}$ agree on $[\ell - (n - m - 1)]$. Applying the map σ , we see that $\beta^{(n)} = \sigma \alpha^{(n-1)}$ and $\beta^{(n-m)} = \sigma \alpha^{(n-m-1)}$ agree on $[\ell - n + m]$.

To reach the conclusion of the lemma, since W(n) - m and W(n-m) are both (possibly empty) intervals starting at n-m, it suffices to show that |W(n)| = |W(n-m)|. Recall that |W(n)| is the length of the interval on which $\pi_{\nu_2(n)}$ changes $\beta^{(n)}$, and similarly for |W(n-m)|and $\beta^{(n-m)}$. Since $\beta^{(n)}$ and $\beta^{(n-m)}$ agree on $[\ell - n + m]$, and since $\nu_2(n) = \nu_2(n-m)$ and $2^{\nu_2(n)} \leq \ell - (n-m)$, it follows from Lemma 3.22 that |W(n)| = |W(n-m)|, as desired. \Box

Lemma 3.27. Let $n, \ell \in \mathbb{N}_0$. If

(i) the greatest integer $m \in [n]$ such that $n \in W(m)$ is 0, and

(ii) $n + (A \cap [2^{\ell}]) \subseteq A$,

then $|W(n)| \ge 2^{\min(\ell,\nu_2(n))}$.

Proof. If n = 0, then $W(0) = \mathbb{N}$ and the conclusion holds. Suppose $n \ge 1$. It follows from Lemma 3.24 that $\beta^{(n)}$ is equal to 1_{A-n} on the set $\mathbb{N}_0 \setminus \bigcup_{m=1}^{n-1} (W(m) - n)$. Indeed, if $i \in \mathbb{N}_0 \setminus \bigcup_{m=1}^{n-1} (W(m) - n) = \bigcap_{m=1}^{n-1} (\mathbb{N}_0 \setminus (W(m) - n))$, then *i* belongs to the set in (3.9) if and only if $i \in A - n$. If (i) holds, then the set $\bigcup_{m=1}^{n-1} (W(m) - n)$ contains no non-negative integers. Therefore, the points $\beta^{(n)}$ and 1_{A-n} are equal on the set \mathbb{N}_0 .

Suppose (i) and (ii) hold. Since $\beta^{(n)} = 1_{A-n}$ and $n + (A \cap [2^{\ell}]) \subseteq A$, we see that $\nu_A(\beta^{(n)}) \ge \ell$. It follows from the definition of W(n) that

$$W(n) = n + [2^{\min(\nu_A(\beta^{(n)}), \nu_2(n))}] \supseteq n + [2^{\min(\ell, \nu_2(n))}],$$

whereby $|W(n)| \ge 2^{\min(\ell,\nu_2(n))}$, as desired.

For $\ell \in \mathbb{N}_0$, we define

$$L(\ell) := \left\{ n \in \mathbb{N} \mid |W(n)| \ge 2^{\ell} \right\}$$

to be the set of those positive integers n at which the punch window W(n) is of length at least 2^{ℓ} . The following lemma shows that the collection of return time sets of the point $(0, 1_A)$ to neighborhoods of itself under the shift-punch map is essentially the same as the collection of sets $L(\ell), \ell \in \mathbb{N}_0$.

Lemma 3.28.

(i) For all $\varepsilon > 0$, there exists $\ell \in \mathbb{N}_0$ such that

$$L(\ell) \subseteq R_{\kappa}((0, 1_A), B_{\varepsilon}((0, 1_A)))).$$

(ii) For all $\ell \in \mathbb{N}_0$, there exists $\varepsilon > 0$ such that

$$R_{\kappa}((0,1_A), B_{\varepsilon}((0,1_A))) \subseteq L(\ell).$$

Proof. (i) Let $\varepsilon > 0$. Choose $\ell \in \mathbb{N}_0$ sufficiently large so that for all $n \in \mathbb{N}$ and all $\omega \in X$, if $\nu_2(n) \ge \ell$ and ω and 1_A agree on $[2^{\ell}]$, then $d_{Z \times X}((n, \omega), (0, 1_A)) < \varepsilon$.

Let $n \in L(\ell)$. We see from the definition of W(n) that $\nu_2(n) \ge |W(n)| \ge \ell$ and that the points $\alpha^{(n)}$ and 1_A agree on $W(n) - n \ge [2^{\ell}]$. By the choice of ℓ , we have that $d(\kappa^n(0, 1_A), (0, 1_A)) < \varepsilon$, whereby $n \in R_{\kappa}((0, 1_A), B_{\varepsilon}((0, 1_A)))$, as was to be shown.

(ii) Let $\ell \in \mathbb{N}_0$. Choose $\varepsilon > 0$ so that for all pairs of points $(n, \omega) \in \mathbb{Z} \times X$, if (n, ω) and $(0, 1_A)$ are at a distance of at most ε , then $\nu_2(n) \ge \ell$ and ω and 1_A agree on $[2^\ell]$.

Let $n \in R_{\kappa}((0, 1_A), B_{\varepsilon}((0, 1_A)))$. Since $\kappa^n(0, 1_A) = (n, \alpha^{(n)})$ and $(0, 1_A)$ are at a distance of at most ε , we have that $\nu_2(n) \ge \ell$ and $\alpha^{(n)}$ and 1_A agree on $[2^{\ell}]$. By the definition of $\pi_{\nu_2(n)}$, we have that $\alpha^{(n)} \le \beta^{(n)}$ pointwise. Since $\alpha^{(n)}$ and 1_A agree on $[2^{\ell}]$, we have that $A \cap [2^{\ell}] \subseteq \text{supp}(\beta^{(n)})$. Thus, $\nu_A(\beta^{(n)}) \ge \ell$. It follows that $|W(n)| = 2^{\min(\nu_A(\beta^{(n)}),\nu_2(n))} \ge 2^{\ell}$, whereby $n \in L(\ell)$, as was to be shown.

We are ultimately interested in the result of the punches made on the set A. Thus, we define

$$B := \bigcap_{n=0}^{\infty} \left(\left(A + n \right) \cup \left(\mathbb{N}_0 \backslash W(n) \right) \right).$$
(3.12)

Informally, the set B is the set A after all punches have occurred; this is made precise by the equivalent dynamical characterization of the set B given in Lemma 3.29. Since $W(0) = \mathbb{N}_0$, we see from the definition that $B \subseteq A$. Moreover, we see that if $0 \in A$, then $0 \in B$.

Lemma 3.29. For all $n \in \mathbb{N}_0$, we have that $(\alpha^{(n)})(0) = 1_B(n)$.

Proof. Let $n \in \mathbb{N}_0$. It follows by Lemma 3.24 that $(\alpha^{(n)})(0) = 1$ if and only if

$$0 \in \bigcap_{m=0}^{n} \left(\left(A - (n-m) \right) \cup \left(\mathbb{N}_0 \setminus \left(W(m) - n \right) \right) \right).$$

By set algebra, the previous line holds if and only if

$$n \in \bigcap_{m=0}^{n} \left(\left(A + m \right) \cup \left(\mathbb{N}_0 \backslash W(m) \right) \right).$$

Since for all $m \in \mathbb{N}_0$ greater than $n, W(m) \subseteq \{n+1, n+2, \ldots\}$, we see that the previous line holds if and only if

$$n \in \bigcap_{m=0}^{\infty} \left(\left(A+m\right) \cup \left(\mathbb{N}_0 \setminus W(m)\right) \right) = B,$$

as was to be shown.

The following lemma connects the times of return of the point 1_B to neighborhoods of itself under the usual shift to the family of sets $L(\ell)$ defined above.

Lemma 3.30. For all $\varepsilon > 0$, there exists $\ell \in \mathbb{N}_0$ such that

$$L(\ell) \subseteq R_{\sigma}(1_B, B_{\varepsilon}(1_B))$$

Proof. Let $\varepsilon > 0$. Choose $\ell \in \mathbb{N}_0$ so that if two points in X agree on $[2^{\ell}]$, then they are at a distance of no more than ε .

Let $n \in L(\ell)$. By Lemma 3.25, for all $i \in [2^{\ell}] \subseteq W(n) - n$, the points $\alpha^{(i)}$ and $\alpha^{(n+i)}$ agree on W(n) - n - i. In particular, they agree at 0. It follows from Lemma 3.29 that for all $i \in [2^{\ell}]$, $1_B(i) = 1_B(i+n) = 1_{B-n}(i) = (\sigma^n 1_B)(i)$. Since 1_B and $\sigma^n 1_B$ agree on $[2^{\ell}]$, we see that $n \in R_{\sigma}(1_B, B_{\varepsilon}(1_B))$, as was to be shown.

Remark 3.31. If $0 \notin A$, then $B = \emptyset$. Indeed, since $0 \notin A$, for all $n \in \mathbb{N}_0$, the map $\pi_{\nu_2(n)}$ will punch $\beta^{(n)}$ at least at the zero coordinate (that is, $n \in W(n)$), changing that coordinate to 0 if it is 1. Thus, $\alpha^{(n)}(0) = 0$, and according to Lemma 3.29, the set B is empty. In this case, we see that $R_{\sigma}(1_B, B_{\varepsilon}(1_B)) = \mathbb{N}$. This shows that a result analogous to Lemma 3.28 for the point 1_B under the shift σ does not necessarily hold: it may be that there exists $\ell \in \mathbb{N}_0$ such that for all $\varepsilon > 0$, the set $R_{\sigma}(1_B, B_{\varepsilon}(1_B))$ is not contained in $L(\ell)$.

3.3.7. Central syndetic sets are dynamically central syndetic

To finish the proof of Theorem 3.18, we must show that central syndetic sets are dynamically central syndetic. It suffices by Theorem 3.15 and Lemma 3.17 to show: if $A \subseteq \mathbb{N}$ satisfies the condition in (3.7), then A is dynamically central syndetic. We carry forward all of the notation from the previous sections.

Let $A \subseteq \mathbb{N}$ satisfy the condition in (3.7). We append 0 to A and, by a slight abuse of notation, consider A as a subset of \mathbb{N}_0 . The key step is to show that the condition in (3.7) implies that the sets $L(\ell)$ defined in the previous section are syndetic.

Theorem 3.32. For all $\ell \in \mathbb{N}_0$, the set $L(\ell)$ is syndetic.

Proof. Let $\ell \in \mathbb{N}_0$. We will show first that the set $L(\ell)$ is nonempty. It follows from (3.7) that there exists $n \in 2^{\ell} \mathbb{N}$ such that $n + (A \cap [2^{\ell}]) \subseteq A$. Let $m \in [n]$ be the greatest integer such that $n \in W(m)$. If m = 0, then by Lemma 3.27, we have that $|W(n)| \ge 2^{\ell}$, whereby $n \in L(\ell)$. Otherwise, we have that $1 \le m < n$ and $n \in W(m)$, whereby $(W(m) \setminus \{m\}) \cap 2^{\ell} \mathbb{N} \neq \emptyset$. It follows from Lemma 3.19 that $|W(m)| \ge 2^{\ell+1}$, so $m \in L(\ell)$.

By (3.7), Lemma 3.20, and the previous paragraph, there exists $k \in \mathbb{N}$ so that

$$2^{\ell} \mathbb{N}_0 \cap \bigcap_{a \in A \cap [2^{\ell}]} (A - a) \in \mathcal{Q}_{2^k}^*$$
(3.13)

and so that $L(\ell) \cap [2^k] \neq \emptyset$. We will show that $L(\ell)$ is syndetic by showing that $L(\ell) \in \mathcal{Q}^*_{2^{k+1}}$ and appealing again to Lemma 3.20.

To show that $L(\ell) \in \mathcal{Q}_{2^{k+1}}^*$, we will show that for all $a \in 2^{k+1}\mathbb{N}_0$, the set $L(\ell) \cap (a + ([2^{k+1}] \setminus \{0\}))$ is nonempty. We proceed by induction on a. We have already shown the base case a = 0. (Recall that, by definition, $L(\ell) \subseteq \mathbb{N}$, so that $L(\ell) \cap [2^k] \neq \emptyset$ implies $L(\ell) \cap ([2^k] \setminus \{0\}) \neq \emptyset$.)

Let $a \in 2^{k+1}\mathbb{N}$, and define $Q = a + [2^{k+1}]$. From (3.13), there exists

$$n \in (Q \setminus \{a\}) \cap 2^{\ell} \mathbb{N}_0 \cap \bigcap_{a \in A \cap [2^{\ell}]} (A - a).$$

Let $m \in [n]$ be the greatest integer such that $n \in W(m)$. If m = 0, then by Lemma 3.27, we have that $|W(n)| \ge 2^{\ell}$, whereby $n \in L(\ell) \cap (Q \setminus \{a\})$, as desired. Otherwise, we have that $1 \le m < n$.

If m > a, then $m \in Q \setminus \{a\}$. Since m < n and $n \in W(m)$, we have that $(W(m) \setminus \{m\}) \cap 2^{\ell} \mathbb{N} \neq \emptyset$. It follows from Lemma 3.19 that $|W(m)| \ge 2^{\ell+1}$, so $m \in L(\ell) \cap (Q \setminus \{a\})$, as desired.

If, on the other hand, $m \leq a$, then the fact that W(m) and Q are intersecting dyadic intervals (both contain n) implies that $m \in 2^{k+1}\mathbb{N}$, that $Q \subseteq W(m)$, and that $Q-m \in \mathcal{Q}_{2^{k+1}}$. It follows by the induction hypothesis that there exists $q \in Q \setminus \{a\}$ such that $q-m \in L(\ell) \cap (Q-m)$. Thus, $|W(q-m)| \geq 2^{\ell}$. Since $q \in W(m)$ and $m \leq a < q$, it follows from Lemma 3.26 that W(q-m) = W(q) - m. Therefore, $|W(q)| \geq 2^{\ell}$, whereby $q \in L(\ell) \cap (Q \setminus \{a\})$, as desired. \Box

Let $B \subseteq A$ be the subset of A defined in (3.12) in the previous section. Since $0 \in A$, we see that $0 \in B$. It follows by combining Lemma 3.30 and Theorem 3.32 that the point 1_B is uniformly recurrent in the shift (X, σ) . Since $0 \in B$, it follows from Theorem 3.8 that the set $B \setminus \{0\} \subseteq \mathbb{N}$ is dynamically central syndetic. Therefore, we have that the set $A \setminus \{0\}$ is dynamically central syndetic, concluding the proof of Theorem 3.18.

While we do not have need for it, note that it follows by combining Lemma 3.28 and Theorem 3.32 that the point $(0, 1_A)$ is uniformly recurrent in the shift-punch system $(Z \times X, \kappa)$.

3.4. Combinatorial characterizations: proofs of Theorems A and B

In this section, we bring everything together to prove Theorems A and B from the introduction and give a short application. Logically, it makes more sense to prove Theorem B first.

Proof of Theorem B. This follows immediately by combining Theorems 3.15 and 3.18. \Box

Proof of Theorem A. (i) \iff (iii) If follows by Lemma 3.4 that A is dynamically syndetic if and only if there exists $n \in \mathbb{N}_0$ such that A-n is dynamically central syndetic. By Theorem B, a set is dynamically central syndetic if and only if it belongs to a syndetic, idempotent filter. The desired equivalence is shown by combining these two results.

(iii) \implies (ii) Suppose that there exists $n \in \mathbb{N}_0$ such that A' := A - n belongs to a syndetic, idempotent filter. We have by Theorem B that there exists $B' \subseteq A'$ such that for all finite $F' \subseteq B'$, the set $B' \cap \bigcap_{f' \in F'} (B' - f')$ is syndetic. Put B = B' + n, and note that B is a nonempty subset of A. Let $F \subseteq B$ be finite, and note that $\min F \ge n + 1$. We have

that $F - n \subseteq B'$ is finite, and hence that the set

$$\bigcap_{f'\in F-n} (B'-f')$$

is syndetic. But this set is contained in $\bigcap_{f \in F} (B - f)$, demonstrating syndeticity, as desired.

(ii) \implies (iii) Let $b \in B$. We will show that the set $B - b \subseteq A - b$ satisfies: for all finite $F \subseteq B - b$, the set

$$(B-b)\cap \bigcap_{f\in F}(B-b-f)$$

is syndetic. It will follow by Theorem 3.15 that the set A - b is contained in a syndetic, idempotent filter.

Let $F \subseteq B - b$ be finite. By assumption, since $\{b\} \cup (F + b) \subseteq B$ is finite, we have that the set

$$\bigcap_{f \in \{b\} \cup (F+b)} (B-f) = (B-b) \cap \bigcap_{f \in F} (B-b-f)$$

is syndetic, as was to be shown.

As a short application of Theorem B, we will show that thickly syndetic sets are dynamically central syndetic. This result has appeared in related forms several times in the literature (see [25, Thm. 1], [37, Thm. 2.4], [12, Prop. 4.4]).

Theorem 3.33. Every thickly syndetic set is dynamically central syndetic.

Proof. By Lemma 2.11, the family \mathcal{PS}^* is a syndetic, translation-invariant (hence, idempotent) filter. It follows from Theorem B that every member of \mathcal{PS}^* is dynamically central syndetic, as desired.

Note that we upgrade Theorem 3.33 in Theorem 5.13 to: the intersection of a thickly syndetic set and a dynamically central syndetic set is dynamically central syndetic.

4. Part II: Dynamically thick sets and sets of pointwise recurrence

In this section, we define the families of dynamically thick sets and sets of pointwise recurrence. The main results are the combinatorial characterizations of these families (Theorems C and D) in Section 4.2; a structure result for dynamically thick sets (Theorem F) in Section 4.3.3; and partial progress in Section 4.4 on the problem of partitioning dynamically thick sets.

4.1. Definition and first characterizations

A set $A \subseteq \mathbb{N}$ is dynamically thick if for all minimal systems (X,T) and all points $x \in X$, the set $\{T^n x \mid n \in A\}$ is dense in X. The set A is a set of (minimal topological) pointwise recurrence – also, dynamically central thick – if for all minimal systems (X,T), all points $x \in X$, and all open $U \subseteq X$ containing x, there exists $a \in A$ such that $T^a x \in U$. The

families of dynamically thick sets and sets of pointwise recurrence are denoted $d\mathcal{T}$ and $dc\mathcal{T}$, respectively.

Lemma 4.1. The families of dynamically (central) syndetic and dynamically (central) thick sets are dual, that is,

$$d\mathcal{T} = d\mathcal{S}^*$$
 and $dc\mathcal{T} = dc\mathcal{S}^*$.

Proof. This follows immediately from the definitions.

It can be verified quickly from the definitions or from Lemma 4.1 that

$$\mathcal{T} \subseteq d\mathcal{T} \subseteq dc\mathcal{T}$$

and that each of these inclusions is proper. Here is a short list of examples of dynamically (central) thick sets that may be helpful to have in mind; more elaborate examples are given in Section 4.3.

- (i) For $k \in \mathbb{N}$, the set of positive multiples of k, $k\mathbb{N}$, is a set of pointwise recurrence. In fact, for all thick sets $H \subseteq \mathbb{N}$, the set $k\mathbb{N} \cap H$ is a set of pointwise recurrence. Indeed, this follows from Lemma 3.6 and the fact that H/k is thick: if $A \in dcS$, then $A/k \in dcS$, whereby $A/k \cap H/k \neq \emptyset$, whereby $A \cap k\mathbb{N} \cap H \neq \emptyset$.
- (ii) If A is not dynamically (central) syndetic, then its complement, N\A, is dynamically (central) thick. This follows immediately from Lemma 4.1. The set in (1.1) from the introduction is not dynamically syndetic, hence its complement, a set of the same form, is dynamically thick. A more general class of dynamically thick sets is described in Lemma 4.11.
- (iii) If $x \in X := \{0,1\}^{\mathbb{N}}$ has a dense orbit in the full shift (X, σ) , then for all nonempty, open $U \subseteq X$, the set $R_{\sigma}(x, U)$ is dynamically thick. This is a consequence of a theorem of Furstenberg [18, Thm. II.2]: the full shift is *disjoint* (cf. [18, Def. II.1]) from every minimal system. (This result is generalized in [24, Thm. 6.2] to arbitrary countable, infinite group actions.) Indeed, we will argue that for all minimal systems (Y, S), all $y \in Y$, and all nonempty, open $V \subseteq Y$, the set

$$R_{\sigma \times S}((x,y), U \times V) = R_{\sigma}(x,U) \cap R_S(y,V) \neq \emptyset.$$
(4.1)

By an equivalent definition of the disjointness of (X, σ) and (Y, S) (cf. [18, Lemma II.1]), the system $(X \times Y, \sigma \times S)$ is the only subsystem of the product system that projects to X in the first coordinate and Y in the second. Because x and y have dense orbits in X and Y, respectively, the subsystem $((\sigma \times S)^{\mathbb{N}}(x, y), \sigma \times S)$ projects to X and Y in the first and second coordinates, respectively, and hence is the product system. Since the point (x, y) has a dense orbit in the product system, we see that (4.1) holds. We offer an alternative explanation of this example in Lemma 4.9 and Remark 4.10 below.

We call the reader's attention to the fact that dynamically central thick sets (sets of pointwise recurrence) need not be central, despite the terminology. The example given in Lemma 4.9 demonstrates this.

The robustness of the families of dynamically (central) syndetic sets was discussed in Section 3.1. Because of the duality shown in Lemma 4.1, the families of dynamically (central) thick sets exhibit a similar robustness under changes to the combinatorial, topological, or

dynamical requirements in their definitions. Thus, the families of dynamically (central) thick subsets of \mathbb{N} remain unchanged if ...

- (i) ... "for all minimal systems" is changed to "for all invertible minimal systems," "for all zero dimensional minimal systems," or "for all (not-necessarily-metrizable) minimal systems." By Lemma 4.1, these follow immediately from Lemma 3.1 and Theorems 3.3 and 3.8, respectively.
- (ii) ... (for sets of pointwise recurrence) "all open $U \subseteq X$ with $x \in U$ " is changed to "all open $U \subseteq X$ with $x \in \overline{U}$." This is a consequence of Lemma 4.1 and Theorem 3.8.
- (iii) ... sets are considered up to equivalence on thickly syndetic (\mathcal{PS}^*) sets. This is a consequence of Theorem 5.13: a set $A \subseteq \mathbb{N}$ is dynamically (central) thick if and only if for all thickly syndetic $B \subseteq \mathbb{N}$, the set $A \cap B$ is dynamically (central) thick.

4.1.1. First results

In this section, we collect some first results concerning dynamical thickness. As discussed in Section 1.2.1, that dynamically (central) thick subsets of \mathbb{N} are piecewise syndetic has been shown several times in the literature. We record it here for ease of reference.

Theorem 4.2. Sets of pointwise recurrence (and, hence, dynamically thick sets) are piecewise syndetic.

Proof. This follows immediately from the duality in Lemma 4.1 and Theorem 3.33. \Box

It is easy to see that the family of dynamically thick sets is not partition regular. Indeed, neither the set of even nor the set of odd positive integers is dynamically thick. Host, Kra, and Maass asked in [35, Question 6.5] whether or not the family $dc\mathcal{T}$ is partition regular. The following result gives a strong negative answer to this question.

Theorem 4.3. There exist sets $A, B \subseteq \mathbb{N}$ that are not sets of pointwise recurrence but for which $\mathbb{N} = A \cup B$. In particular, the family of sets of pointwise recurrence is not partition regular.

Proof. Since \mathbb{N} is a dynamically central syndetic set, according to Theorem E, there exists a partition $\mathbb{N} = A \cup B$ into disjoint, dynamically central syndetic sets. Because $A \cap B = \emptyset$, neither A nor B is a set of pointwise recurrence.

We note that the partition guaranteed by Theorem E can be made quite explicit: for $\alpha \in \mathbb{R}\setminus\mathbb{Q}$, define $A := \{n \in \mathbb{N} \mid \{n\alpha\} \in (0, 1/2)\}$ and $B := \{n \in \mathbb{N} \mid \{n\alpha\} \in (1/2, 1)\}$. Neither A nor B is a set of pointwise recurrence, but $\mathbb{N} = A \cup B$.

Lemma 4.4. The intersection of a dynamically (central) syndetic set and a dynamically (central) thick set is infinite.

Proof. Let $A \in d\mathcal{S}$, $B \in d\mathcal{T}$, and $n \in \mathbb{N}$. By Lemma 3.2, we see that $A \cap \{n+1, n+2, \ldots\} \in d\mathcal{S}$. Since $B \in d\mathcal{T}$, we have that $A \cap \{n+1, n+2, \ldots\} \cap B \neq \emptyset$. Therefore, $\max(A \cap B) > n$. Since $n \in \mathbb{N}$ was arbitrary, we see that $A \cap B$ is infinite. The same argument works to show that $A \cap B$ is infinite when $A \in dc\mathcal{S}$ and $B \in dc\mathcal{T}$. Theorem 5.11 in the next section simultaneously strengthens both Theorem 4.2 and Lemma 4.4 by showing that the intersection of a dynamically (central) syndetic set and a dynamically (central) thick set is, in fact, piecewise syndetic.

4.1.2. Translates and dilates of dynamically thick sets

We show in this section what happens to dynamically (central) thick sets under translations and dilations. Most of the statements in Lemma 4.5 can be shown directly from the definitions, but we choose to derive them from Lemma 3.4 using the duality in Lemma 4.1.

Lemma 4.5. Let $A \subseteq \mathbb{N}$.

- (i) If A is dynamically thick, then for all $n \in \mathbb{N}$, the set A n is dynamically thick.
- (ii) If the set

$$\{n \in \mathbb{N} \mid A - n \text{ is a set of pointwise recurrence}\}$$
(4.2)

is dynamically thick, then A is dynamically thick.

(iii) If the set in (4.2) is a set of pointwise recurrence, then A is a set of pointwise recurrence. Moreover, (i), (ii), and (iii) hold with A - n replaced by A + n. Finally,

$$d\mathcal{T} = \bigcap_{n \in \mathbb{N}} \left(dc\mathcal{T} - n \right) = \bigcap_{n \in \mathbb{N}} \left(dc\mathcal{T} + n \right).$$
(4.3)

Proof. (i) Suppose $A \in d\mathcal{T}$ and $n \in \mathbb{N}$. To see that $A - n \in d\mathcal{T}$, by Lemma 4.1, it suffices to show that $A - n \in d\mathcal{S}^*$. Let $B \in d\mathcal{S}$. By Lemma 3.4, the set $B + n \in d\mathcal{S}$. By Lemma 4.4, the set $A \cap (B + n)$ is infinite, so $(A - n) \cap B \supseteq (A - n) \cap (B + n - n) \neq \emptyset$, as desired.

(ii) In the language of the family algebra developed in Section 2.1, this statement is equivalent to $d\mathcal{T} + dc\mathcal{T} \subseteq d\mathcal{T}$. Recall from Lemma 3.4 that $d\mathcal{S} \subseteq d\mathcal{S} + dc\mathcal{S}$. Taking the dual and simplifying using the algebra in Lemma 2.3, we see that $d\mathcal{T} + dc\mathcal{T} \subseteq d\mathcal{T}$.

(iii) This statement is equivalent to $dc\mathcal{T} + dc\mathcal{T} \subseteq dc\mathcal{T}$. Taking the dual of $dc\mathcal{S} \subseteq dc\mathcal{S} + dc\mathcal{S}$ from Lemma 3.4, we see that $dc\mathcal{T} + dc\mathcal{T} \subseteq dc\mathcal{T}$.

That (i) holds with A - n replaced with A + n follows the same argument used in (i). That (ii) and (iii) hold with A - n replaced by A + n requires different reasoning, since the set algebra explanation only treats negative translates.

Suppose that A is such that the set in (4.2) (with A + n instead of A - n) is dynamically thick. We wish to show that A is dynamically thick. It suffices by Lemmas 3.1 and 4.1 to show: for all minimal, invertible (X,T), all $x \in X$, and all nonempty, open $U \subseteq X$, there exists $a \in A$ such that $T^a x \in U$. Thus, let (X,T) be a minimal, invertible system, $x \in X$, and $U \subseteq X$ open and nonempty. Since the system (X,T^{-1}) is minimal (see, for example, [26, Lemma 2.7]) and the set in (4.2) is dynamically thick, there exists $n \in \mathbb{N}$ such that $T^{-n}x \in U$ and $A + n \in dc\mathcal{T}$. By the definition of set of pointwise recurrence, there exists $m \in A + n$ (so m > n) such that $T^m T^{-n} x \in U$. We see that $m - n \in A$ and $T^{m-n} x \in U$, as desired. The argument that (iii) holds with A - n replaced by A + n is very similar and is left to the interested reader.

Finally, that (4.3) holds follows immediately from (i) and (ii) for A - n and A + n.

As demonstrated in the proof of Lemma 4.5, it is interesting and useful to formulate the lemma's conclusions in terms of the family algebra developed in Section 2.1. Thus,

(i) the family $d\mathcal{T}$ is translation invariant, that is, $d\mathcal{T} \subseteq \{\mathbb{N}\} + d\mathcal{T}$;

(ii) $d\mathcal{T} + dc\mathcal{T} \subseteq d\mathcal{T};$ (iii) $dc\mathcal{T} + dc\mathcal{T} \subseteq dc\mathcal{T}.$

Lemma 4.6. Let $A \subseteq \mathbb{N}$ and $k \in \mathbb{N}$. If A is a set of pointwise recurrence, then so are the sets kA and A/k. If A is dynamically thick, then so is the set A/k.

Proof. Suppose $A \in dc\mathcal{T}$. If $B \in dc\mathcal{S}$, then by Lemma 3.6, the sets kB and B/k are $dc\mathcal{S}$ sets, whereby $A \cap (kB) \neq \emptyset$ and $A \cap (B/k) \neq \emptyset$. Since (kB)/k = B and $k(B/k) \subseteq B$, it follows that $A/k \cap B \neq \emptyset$ and $kA \cap B \neq \emptyset$. Since $B \in dc\mathcal{S}$ was arbitrary, we have that A/k and kA are sets of pointwise recurrence.

Suppose $A \in d\mathcal{T}$. To show $A/k \in d\mathcal{T}$ follows the same argument using Lemma 3.6 as in the previous paragraph.

4.2. Combinatorial characterizations: proofs of Theorems C and D

In this section, we give combinatorial characterizations of dynamical thickness and pointwise recurrence. Recall that the empty union and empty intersection are interpreted to be the empty set and \mathbb{N} , respectively.

Theorems C and D will follow immediately from Theorems 4.7 and 4.8, respectively. In each, the second statement comes from interpreting "has nonempty intersection with all dynamically (central) syndetic sets" in combinatorial terms by appealing to the main results on dynamically syndetic sets, while the third, fourth, and fifth statements arise from ultrafilter dynamics, appealing ultimately to Theorem 3.3.

Theorem 4.7. Let $A \subseteq \mathbb{N}$. The following are equivalent.

- (i) The set A is dynamically thick.
- (ii) For all $\mathbb{N} \supseteq B \supseteq A$, there exists a finite set $F \subseteq \mathbb{N} \setminus B$ such that the set B F is thick.
- (iii) For all syndetic $S \subseteq \mathbb{N}$, there exists a finite set $F \subseteq A$ such that for all syndetic $S' \subseteq \mathbb{N}$, the set $F \cap (S S')$ is nonempty.
- (iv) For all piecewise syndetic $S \subseteq \mathbb{N}$, there exists a finite set $F \subseteq A$ such that the set S F is thick.
- (v) For all piecewise syndetic $S \subseteq \mathbb{N}$ and all minimal left ideals $L \subseteq \beta \mathbb{N}$ with $\overline{S} \cap L \neq \emptyset$, there exists a finite set $F \subseteq A$ such that $L \subseteq \overline{S - F}$.

Proof. (i) \iff (ii) Because dS and dT are dual, the set A is dynamically thick if and only if the set $A' := \mathbb{N} \setminus A$ is not dynamically syndetic. By Theorem A, the set A' is not dynamically syndetic if and only if for all nonempty subsets $B' \subseteq A'$, there exists a finite set $F \subseteq B'$ such that the set $\bigcap_{f \in F} (B' - f)$ is not syndetic. Taking the complement of this set in \mathbb{N} , we see that this happens if and only if the set

$$\bigcup_{f \in F} \left((\mathbb{N} \setminus B') - f \right) = \left(\mathbb{N} \setminus B' \right) - F$$

is thick. Set $B := \mathbb{N} \setminus B'$. Note that $B' \subseteq A'$ if and only if $A \subseteq B$ and that $B' \neq \emptyset$ if and only if $B \neq \mathbb{N}$.

Summarizing the previous paragraph, we have shown that the set A is dynamically thick if and only if for all $A \subseteq B \subsetneq \mathbb{N}$, there exists a finite set $F \subseteq \mathbb{N} \setminus B$ such that the set B - Fis thick, as was to be shown. (i) \implies (v) Suppose that $A \in d\mathcal{T}$. Let $S \subseteq \mathbb{N}$ be piecewise syndetic and $L \subseteq \beta \mathbb{N}$ be a minimal left ideal such that $\overline{S} \cap L \neq \emptyset$. Note that $\overline{S} \cap L$ is nonempty and open in L. By the definitional robustness of $d\mathcal{T}$ sets regarding non-metrizable systems (see Section 4.1), for all $p \in L$, there exists $a \in A$ such that $a + p \in \overline{S}$. Therefore,

$$L \subseteq \bigcup_{a \in A} \overline{S - a}.$$

Because L is compact, there exists a finite set $F \subseteq A$ such that

$$L \subseteq \bigcup_{f \in F} \overline{S - f} \subseteq \overline{S - F},$$

as was to be shown.

(v) \implies (iv) Let $S \subseteq \mathbb{N}$ be a piecewise syndetic set. It follows by [32, Thm. 4.40] that there exists a minimal left ideal $L \subseteq \beta \mathbb{N}$ such that $\overline{S} \cap L \neq \emptyset$. Let $F \subseteq A$ be the finite set guaranteed by (v). Since $\overline{S-F}$ contains a minimal left ideal, by [32, Thm. 4.48], the set S-F is thick, as desired.

(iv) \implies (iii) Let $S \subseteq \mathbb{N}$ be syndetic. Let $F \subseteq A$ be the finite set guaranteed by (iv). Since S - F is thick, for all $S' \subseteq \mathbb{N}$ syndetic, the set $S' \cap (S - F)$ is nonempty, whereby $F \cap (S - S') \neq \emptyset$, as desired.

(iii) \implies (i) We will prove the contrapositive: if A is not dynamically thick, then there exists a syndetic set $S \subseteq \mathbb{N}$ such that for all finite $F \subseteq A$, the set S - F is not thick. Taking complements and considering B as $\mathbb{N}\setminus A$ and C as $\mathbb{N}\setminus S$, we must show: if a set $B \subseteq \mathbb{N}$ is dynamically syndetic, then there exists a set $C \subseteq \mathbb{N}$ that is not thick such that for all finite $F \subseteq \mathbb{N}\setminus B$, the set $\bigcap_{f \in F} (C - f)$ is syndetic.

Suppose $B \subseteq \mathbb{N}$ is dynamically syndetic. If $B = \mathbb{N}$, put $C = \emptyset$ and note that the conclusion holds since the empty intersection is \mathbb{N} . Suppose $B \neq \mathbb{N}$. By Theorem 3.8, there exists a minimal system (X,T), a point $x \in X$, and a nonempty, clopen set $U \subseteq X$ such that $R(x,U) \subseteq B$. Define $V := X \setminus U$ and $C := \mathbb{N} \setminus R(x,U) = R(x,V)$. Since $U \neq X$ and the set R(x,U) is syndetic, the set C is nonempty but not thick. Let $F \subseteq \mathbb{N} \setminus B$ be finite. Since $F \subseteq C$, we have that for all $f \in F$, the set $T^{-f}V$ is an open neighborhood of x. We see that

$$\bigcap_{f \in F} (C - f) = R\left(x, \bigcap_{f \in F} T^{-f}V\right),$$

which is syndetic since the set $\bigcap_{f \in F} T^{-f}V$ is an open neighborhood of x, as desired.

Theorem 4.8. Let $A \subseteq \mathbb{N}$. The following are equivalent.

- (i) The set A is a set of pointwise recurrence.
- (ii) For all $B \supseteq A$, there exists a finite set $F \subseteq \mathbb{N} \setminus B$ such that the set $B \cup (B F)$ is thick.
- (iii) For all syndetic $S \subseteq \mathbb{N}$, there exists a finite set $F \subseteq A$ such that for all syndetic $S' \subseteq S$, the set $F \cap (S S')$ is nonempty.
- (iv) For all syndetic $S \subseteq \mathbb{N}$ and all thick $H \subseteq \mathbb{N}$, the piecewise syndetic set $P := S \cap H$ satisfies the following. There exists a finite set $F \subseteq A$ such that for all syndetic $S' \subseteq S$, the piecewise syndetic set $P' := S' \cap H$ is such that the set $P' \cap (P - F)$ is nonempty.
- (v) For all piecewise syndetic $S \subseteq \mathbb{N}$ and all minimal left ideals $L \subseteq \beta \mathbb{N}$ with $\overline{S} \cap L \neq \emptyset$, there exists a finite set $F \subseteq A$ such that $\overline{S} \cap L \subseteq \overline{S - F}$.

Proof. (i) \iff (ii) Because dcS and dcT are dual, the set A is a set of pointwise recurrence

if and only if the set $A' := \mathbb{N} \setminus A$ is not dynamically central syndetic. By Theorem B, the set A' is not dynamically central syndetic if and only if for all $B' \subseteq A'$, there exists a finite set $F \subseteq B'$ such that the set

$$B' \cap \bigcap_{f \in F} (B' - f)$$

is not syndetic. Taking the complement of this set in \mathbb{N} , we see that this happens if and only if the set

$$(\mathbb{N}\backslash B') \cup \bigcup_{f \in F} ((\mathbb{N}\backslash B') - f) = (\mathbb{N}\backslash B') \cup ((\mathbb{N}\backslash B') - F)$$

is thick. Set $B := \mathbb{N} \setminus B'$. Note that $B' \subseteq A'$ if and only if $A \subseteq B$.

Summarizing the previous paragraph, we have shown that the set A is a set of pointwise recurrence if and only if for all $B \supseteq A$, there exists a finite set $F \subseteq \mathbb{N} \setminus B$ such that the set $B \cup (B - F)$ is thick, as was to be shown.

(i) \implies (v) Suppose that $A \in dc\mathcal{T}$. Let $S \subseteq \mathbb{N}$ be piecewise syndetic and $L \subseteq \beta \mathbb{N}$ be a minimal left ideal such that $\overline{S} \cap L \neq \emptyset$. Note that $\overline{S} \cap L$ is nonempty and clopen in L. By the definitional robustness of $dc\mathcal{T}$ sets regarding non-metrizable systems (see Section 4.1), for all $p \in \overline{S} \cap L$, there exists $a \in A$ such that $a + p \in \overline{S}$. Therefore,

$$\overline{S} \cap L \subseteq \bigcup_{a \in A} \overline{S-a}.$$

Because $\overline{S} \cap L$ is compact, there exists a finite set $F \subseteq A$ such that

$$\overline{S} \cap L \subseteq \bigcup_{f \in F} \overline{S - f} \subseteq \overline{S - F},$$

as was to be shown.

(v) \implies (iv) Let $S \subseteq \mathbb{N}$ be syndetic and $H \subseteq \mathbb{N}$ be thick, and put $P := S \cap H$. It follows by [32, Thm. 4.48] that there exists a minimal left ideal $L \subseteq \overline{H}$. Note that since S is syndetic, we have that $\emptyset \neq \overline{S} \cap L = \overline{S} \cap L \cap \overline{H} = \overline{P} \cap L$.

Let $F \subseteq A$ be the finite set guaranteed by (v) for the piecewise syndetic set P. Let $S' \subseteq S$ be syndetic, and put $P' := S' \cap H$. Since S' is syndetic and $S' \subseteq S$, we have that $\emptyset \neq \overline{S'} \cap L = \overline{P'} \cap L$. Since $\emptyset \neq \overline{P'} \cap L \subseteq \overline{P} \cap L \subseteq \overline{P-F}$, we see that $\overline{P'} \cap \overline{P-F} \neq \emptyset$, whereby $P' \cap (P-F) \neq \emptyset$, as desired.

(iv) \implies (iii) Let $S \subseteq \mathbb{N}$ be syndetic. Get $F \subseteq A$ finite from (v) for $H = \mathbb{N}$. Then for all $S' \subseteq S$ syndetic,

$$S' \cap (S - F) \neq \emptyset,$$

whereby $F \cap (S - S') \neq \emptyset$, as desired.

(iii) \implies (i) We will prove the contrapositive: if A is not a set of pointwise recurrence, then there exists a syndetic set $S \subseteq \mathbb{N}$ such that for all finite $F \subseteq A$, there exists a syndetic set $S' \subseteq S$ such that $F \cap (S - S') = \emptyset$. Since $F \cap (S - S') = \emptyset$ if and only if $S' \cap (S - F) = \emptyset$, it is equivalent to show: there exists a syndetic set $S \subseteq \mathbb{N}$ such that for all finite $F \subseteq A$, the set $S \cap (\mathbb{N} \setminus (S - F))$ is syndetic. Taking complements and considering B as $\mathbb{N} \setminus A$ and C as $\mathbb{N} \setminus S$, we must show: if a set $B \subseteq \mathbb{N}$ is dynamically central syndetic, then there exists a set $C \subseteq \mathbb{N}$ that is not thick such that for all finite $F \subseteq \mathbb{N} \setminus B$, the set $(\mathbb{N} \setminus C) \cap \bigcap_{f \in F} (C - f)$ is syndetic.

Suppose $B \subseteq \mathbb{N}$ is dynamically central syndetic. If $B = \mathbb{N}$, put $C = \emptyset$ and note that the conclusion holds since the empty intersection is \mathbb{N} . Suppose $B \neq \mathbb{N}$. By Theorem 3.8, there exists a minimal system (X,T), a point $x \in X$, and a clopen set $U \subseteq X$ with $x \in U$ such that $R(x,U) \subseteq B$. Define $V := X \setminus U$ and $C := \mathbb{N} \setminus R(x,U) = R(x,V)$. Since $U \neq X$ and the set R(x,U) is syndetic, the set C is nonempty but not thick. Let $F \subseteq \mathbb{N} \setminus B$ be finite. Since $F \subseteq C$, we have that for all $f \in F$, the set $T^{-f}V$ is an open neighborhood of x. We see that

$$(\mathbb{N}\setminus C)\cap \bigcap_{f\in F} (C-f) = R\bigg(x, U\cap \bigcap_{f\in F} T^{-f}V\bigg),$$

which is syndetic since the set $U \cap \bigcap_{f \in F} T^{-f}V$ is an open neighborhood of x, as desired. \Box

4.2.1. Examples of dynamically thick sets via the combinatorial characterizations

In this section, we will use the combinatorial characterizations from the previous section to give two concrete examples of dynamically thick sets.

Lemma 4.9. Let $(p_i)_{i \in \mathbb{N}} \subseteq \mathbb{N}$ be a sequence of distinct primes, $(c_i)_{i \in \mathbb{N}} \subseteq \mathbb{Z}$, and $(H_i)_{i \in \mathbb{N}}$ be a sequence of thick sets. The set

$$A := \bigcup_{i=1}^{\infty} \left((p_i \mathbb{N} + c_i) \cap H_i \right)$$

is dynamically thick. Moreover, if for all $n \in \mathbb{N}$, we have that $c_n \not\equiv 0 \mod p_n$ and, for all but finitely many distinct pairs i, j of positive integers, the set $H_i \cap (H_j - n)$ is empty, then the set A is not an \mathcal{IP} set.

Proof. We will apply Theorem 4.7 (ii). Let $\mathbb{N} \supseteq B \supseteq A$. We want to show there exists a finite set $F \subseteq \mathbb{N} \setminus B$ such that B - F is thick. Consider two cases.

Case 1: There exists $i \in \mathbb{N}$ for which the set $\mathbb{N}\setminus B$ contains a complete residue system modulo p_i . Let $F = \{f_0, \ldots, f_{p_i-1}\} \subseteq \mathbb{N}\setminus B$ be a complete system of modulo p_i residues. We see that

$$B - F \supseteq A - F \supseteq \left((p_i \mathbb{N} + c_i) \cap H_i \right) - F \supseteq \bigcap_{\ell=0}^{p_i - 1} (H_i - f_\ell),$$

which is thick, as desired.

Case 2: For all $i \in \mathbb{N}$, the set $\mathbb{N}\setminus B$ avoids some modulo p_i congruence class. In this case, for all $i \in \mathbb{N}$, there exists $a_i \in \mathbb{N}$ such that $B \supseteq p_i \mathbb{N} + a_i$. We will show that B is thick. Indeed, let $k \in \mathbb{N}$. It suffices to show there exists $n \in \mathbb{N}$ such that $n + i \in p_i \mathbb{N} + a_i$ for all $1 \leq i \leq k$. This is equivalent to showing that there exists $n \in \mathbb{N}$ such that $n \equiv a_i - i \mod p_i$ for $1 \leq i \leq k$. This is an immediate consequence of the Chinese Remainder Theorem since the p_i 's are distinct primes.

To see that the set A is not an \mathcal{IP} set under the stipulated conditions, note that for all $n \in \mathbb{N}$, the set $A \cap (A - n)$ is contained in a finite union of sets of the form $(p\mathbb{N} + c) \cap H$, where $c \not\equiv 0 \mod p$ and H is thick. If A was an \mathcal{IP} set, there would exist $n \in \mathbb{N}$ for which $A \cap (A - n)$ is an \mathcal{IP} set. Since the family \mathcal{IP} is partition regular, it would follow that a set

of the form $(p\mathbb{N} + c) \cap H$ is an \mathcal{IP} set, which is clearly false.

Recently, the second author with Koutsogiannis, Moreira, Pavlov, and Richter [41] gave an example of a set of pointwise recurrence for distal systems which is not an \mathcal{IP} set, answering a question of Host, Kra and Maass [35, Question 3.11]. Choosing the thick sets H_i appropriately, the set $A := \bigcup_{i=1}^{\infty} ((p_i \mathbb{N} + 1) \cap H_i)$ is, according to Lemma 4.9, dynamically thick but not an \mathcal{IP} set. This set provides a strong answer to the question of Host, Kra and Maass: the set A is not only a set of pointwise recurrence for distal systems but a dynamically thick set.

Remark 4.10. Lemma 4.9 can be used to explain why the set described in (iii) at the beginning of Section 4.1 is dynamically thick. Indeed, let x be a point with a dense orbit in the full shift $(\{0,1\}^{\mathbb{N}}, \sigma)$, and suppose $U \subseteq X$ contains the cylinder set of all of those words in $\{0,1\}^{\mathbb{N}}$ beginning with some word w of length ℓ . We see that

$$R(x, U) \supseteq \{n \in \mathbb{N} \mid w \text{ appears at the } n^{\text{th}} \text{ position in } x\}.$$

Let $p > \ell$ be prime, and let w_1 be any word of length p starting with w. Let $k \in \mathbb{N}$, and let $w_2 = w_1 w_1 \cdots w_1$ be the word consisting of k copies of w_1 . Since x is transitive, the word w_2 must appear in x: there exists $N_k \in \mathbb{N}$ and $c_k \in \{0, \ldots, p-1\}$ such that $p\{N_k, N_k + 1, \ldots, N_k + k - 1\} + c_k \subseteq R(x, U).$

Let $c \in \{0, \ldots, p-1\}$ be such that for infinitely many $k \in \mathbb{N}$, $c_k = c$. We see that there exists a thick set $H \subseteq \mathbb{N}$ such that $(p\mathbb{N} + c) \cap H \subseteq R(x, U)$. Since p was an arbitrary prime greater than ℓ , we see that the set R(x, U) contains a set of the form in Lemma 4.9 and hence is dynamically thick, as desired.

The class of examples in the following lemma generalizes the dynamically thick set that appears in (1.1) in the introduction.

Lemma 4.11. Let $k \in \mathbb{N}$ and $(H_i)_{i \in \mathbb{N}}$ be a sequence of thick subsets of \mathbb{N} . The set

$$A := \bigcup_{i=0}^{k-1} \left((k\mathbb{N} + i) \cap H_i \right)$$

is dynamically thick.

Proof. We will apply Theorem 4.7 (iv). Let $S \subseteq \mathbb{N}$ be piecewise syndetic. Since piecewise syndeticity is partition regular and $S = \bigcup_{i=0}^{k-1} (S \cap (k\mathbb{N} - i))$, there exists $i \in \{0, \ldots, k-1\}$ such that $S \cap (k\mathbb{N} - i)$ is piecewise syndetic. Since demonstrating (iv) for a subset of S + i suffices to demonstrate it for S, by replacing S with $(S + i) \cap k\mathbb{N}$, we can proceed under the assumption that $S \subseteq k\mathbb{N}$.

Write $S = B \cap C$ where B is syndetic with gap lengths less than $\ell \in \mathbb{N}$ and C is thick. For $i \in \{0, \ldots, k-1\}$, choose an interval I_i of H_i such that $|I_i| > \ell$ and $\max I_i < \min I_{i+1}$. The set

$$F := \bigcup_{i=0}^{k-1} \left((k\mathbb{N} + i) \cap I_i \right)$$

is a finite subset of A. We will show that for all $M \in \mathbb{N}$, there is a finite subset $S' \subseteq S$ for which the set S' - F contains an interval of length greater than M. It will follow that the set S - F is thick. Let $M \in \mathbb{N}$. Choose $S' \subseteq S$ to be the set S intersected with a long interval (whose length will be specified later) on which the distance between consecutive elements of S is less than ℓ . Since $\ell < |I_i|$ and $S' \subseteq k\mathbb{N}$, the set $S' - ((k\mathbb{N} + i) \cap I_i)$ contains the set $(k\mathbb{N} - i) \cap J_i$, where

$$J_i := \{\min S' - \max I_i, \ldots, \max S' - \min I_i\}.$$

It follows that

$$S'-F \supseteq \bigcup_{i=0}^{k-1} \left((k\mathbb{N}-i) \cap J_i \right) \supseteq \bigcap_{i=0}^{k-1} J_i = \left\{ \min S' - \max I_0, \ldots, \max S' - \min I_{k-1} \right\},$$

provided $\min S' - \max I_0 \leq \max S' - \min I_{k-1}$. This interval has length $\max S' - \min S' - (\min I_{k-1} - \max I_0)$ which is greater than M if $\max S' - \min S' > M + \min I_{k-1} - \max I_0$. \Box

4.3. More general examples of dynamically thick sets

In this section, we develop more sophisticated dynamical tools to generalize the examples of dynamically thick sets presented in Lemma 4.9 and Lemma 4.11. We also show that in some sense, every dynamically thick set takes the form exhibited in these examples.

4.3.1. Dynamically thick sets from disjointness

In this section, we generalize the example in Lemma 4.9 to a wider class of dynamically thick sets. We accomplish this by showing that every minimal system must be nearly disjoint from almost all systems in a disjoint collection of minimal systems. This is reminiscent of the fact that in a Hilbert space, according to Bessel's inequality, every vector must be nearly orthogonal to almost all vectors in an orthonormal sequence.

We call a collection of minimal systems (X_i, T_i) , $i \in I$, disjoint if the product system $(\prod_i X_i, \prod_i T_i)$ is minimal. Note that by the definition of the product topology, this is equivalent to having that the product system of any finite subcollection of the systems is minimal. This definition of disjointness is the restriction of the one given in example (iii) in Section 4.1 to minimal systems.

For our purposes below, a *joining* of the systems (X,T) and (Y,S) is a subsystem $(Z \subseteq X \times Y, T \times S)$ of the product system for which $\pi_X Z = X$ and $\pi_Y Z = Y$. Thus, two minimal systems are disjoint if their only joining is the product system. It is easy to see that if (X,T) and (Y,S) are both minimal, then every subsystem of $(X \times Y, T \times S)$ is a joining. In this case, every joining of (X,T) and (Y,S) has a minimal subsystem that is a joining.

Theorem 4.12. Let (Z_i, R_i) , $i \in \mathbb{N}$, be a disjoint collection of minimal systems. For all minimal (X, T), all $\varepsilon > 0$, and all sufficiently large i (depending on the (Z_i, R_i) 's, (X, T), and ε), all joinings $(J, T \times R_i)$ of (X, T) and (Z_i, R_i) satisfy: for all $z \in Z_i$, the set $\pi_1(J \cap (X \times \{z\}))$ is ε -dense in X.

Proof. Let (X,T) be minimal and $\varepsilon > 0$. Suppose for a contradiction that there are infinitely many *i*'s for which there exists a joining $(J_i, T \times R_i)$ of (X,T) and (Z_i, R_i) and a point $y_i \in Z_i$ for which $\pi_1(J_i \cap (X \times \{y_i\}))$ is not ε dense in X. To save on notation, by relabeling, we will ignore those (Z_i, R_i) 's which do not fall into this infinite set. By passing to minimal subsystems, we may assume without loss of generality that the joinings $(J_i, T \times R_i)$ are minimal. Since J_i is closed, there exists an ε -ball $U_i \subseteq X$ and an open set $V_i \subseteq Z_i$ containing y_i such that $J_i \cap (U_i \times V_i) = \emptyset$.

Fix $x_0 \in X$. Since $(J_i, T \times R_i)$ is a joining of (X, T) and (Z_i, R_i) and is minimal, there exists $z_i \in Z_i$ such that $J_i = \overline{(T \times R_i)^{\mathbb{N}}(x_0, z_i)}$. Since $J_i \cap (U_i \times V_i) = \emptyset$, we have that

$$R_{R_i}(z_i, V_i) \subseteq R_T(x_0, X \setminus U_i).$$

$$(4.4)$$

Let $c_1, c_2, \ldots, c_m \in X$ be an $\varepsilon/2$ dense subset of X. For $\ell \in \{1, \ldots, m\}$, define

$$E_{\ell} = \{ i \in \mathbb{N} \mid B_{\varepsilon/2}(c_{\ell}) \subseteq U_i \}.$$

Note that $\mathbb{N} = \bigcup_{\ell=1}^{m} E_{\ell}$. There exists $\ell \in \{1, \ldots, m\}$ for which E_{ℓ} is infinite. Put $U = B_{\varepsilon/2}(c_{\ell})$. It follows now from (4.4) that

$$H \coloneqq \bigcup_{i \in E_{\ell}} R_{R_i}(z_i, V_i) \subseteq \bigcup_{i \in E_{\ell}} R_T(x_0, X \setminus U_i) \subseteq R_T(x_0, X \setminus U).$$
(4.5)

We will show that the set H is thickly syndetic and, hence, thick. This will yield a contradiction since, by the minimality of (X, T), the set $R_T(x_0, X \setminus U)$ cannot be thick.

Let $k \in \mathbb{N}$, and let $e_1, \ldots, e_k \in E_\ell$. We see that

$$\bigcap_{i=1}^{k} (H-i) \supseteq \bigcap_{i=1}^{k} R_{R_{e_i}}(R_{e_i}^i z_{e_i}, V_{e_i}) = R_{R_{e_1} \times \dots \times R_{e_k}} \left((R_{e_1}^1 z_{e_1}, \dots, R_{e_k}^k z_{e_k}), V_{e_1} \times \dots \times V_{e_k} \right),$$

which is syndetic by the minimality of the product system $(Z_{e_1} \times \cdots \times Z_{e_k}, R_{e_1} \times \cdots \times R_{e_k})$. Therefore, intervals of length k appear in H syndetically. Since $k \in \mathbb{N}$ was arbitrary, this shows that the set H is thickly syndetic, as desired.

Theorem 4.13. Let (X_i, T_i) , $i \in \mathbb{N}$, be a disjoint collection of minimal systems. For all minimal systems (Y, S) and all nonempty, open $V \subseteq Y$, there exists $i \in \mathbb{N}$ such that the following holds. For all nonempty, open $U \subseteq X_i$, there exists $s \in \mathbb{N}$ such that for all $(x, y) \in X_i \times Y$, the set $R_{T_i \times S}((x, y), U \times V) = R_{T_i}(x, U) \cap R_S(y, V)$ is syndetic with gap size bounded by s.

Proof. Let (Y, S) be minimal and $V \subseteq Y$ be nonempty and open. Let $\varepsilon > 0$ be such that V contains an ε -ball. According to Theorem 4.12, there exists $i \in \mathbb{N}$ such that all joinings $(J, T_i \times S)$ of (X_i, T_i) and (Y, S) satisfy: for all $x \in X_i$, the set $\pi_2(J \cap (\{x\} \times Y))$ is ε -dense in Y.

Let $U \subseteq X$ be nonempty and open. We claim that for all $(x, y) \in X_i \times Y$, there exists $n \in \mathbb{N}$ such that $(T_i \times S)^n(x, y) \in U \times V$. Indeed, let $(x, y) \in X_i \times Y$. By a theorem of Auslander [1] and Ellis [14], the point (x, y) is proximal to a point $(x_0, y_0) \in X_i \times Y$ that is uniformly recurrent under $T_i \times S$. Since $J := (T_i \times S)^{\mathbb{N}_0}(x_0, y_0)$ is a (minimal) joining of (X_i, T_i) and (Y, S), it is ε -dense in every fiber over X_i . Since V contains an ε -ball, we see that $J \cap (U \times V) \neq \emptyset$. Because (x, y) is proximal to (x_0, y_0) and $R_{T_i \times S}((x_0, y_0), U \times V)$ is syndetic, we see that $R_{T_i \times S}((x, y), U \times V)$ is piecewise syndetic, hence nonempty.

We have shown that for all $(x, y) \in X_i \times Y$, there exists $n \in \mathbb{N}$ such that $(T_i \times S)^n(x, y) \in U \times V$. It follows that $\bigcup_{n=1}^{\infty} (T_i \times S)^{-n}(U \times V) = X_i \times Y$. By compactness, there exists $s \in \mathbb{N}$ such that $\bigcup_{n=1}^{s} (T_i \times S)^{-n}(U \times V) = X_i \times Y$. It follows that for all $(x, y) \in X_i \times Y$, there exists $n \in \{1, \ldots, s\}$ such that $(T_i \times S)^n(x, y) \in U \times V$, whereby for all $(x, y) \in X_i \times Y$, the set $R_{T_i \times S}((x, y), U \times V)$ is syndetic with gap size bounded by s, as was to be shown.

The following is a generalization of Lemma 4.9.

Theorem 4.14. Let (X_i, T_i) , $i \in \mathbb{N}$, be a disjoint collection of minimal systems. For each $i \in \mathbb{N}$, let $x_i \in X_i$, $U_i \subseteq X_i$ be nonempty and open, and $H_i \subseteq \mathbb{N}$ be thick. The set

$$A := \bigcup_{i=1}^{\infty} \left(R_{T_i}(x_i, U_i) \cap H_i \right)$$

is dynamically thick.

Proof. To see that $A \in d\mathcal{T}$, we must show that for all minimal (Y, S), all nonempty, open $V \subseteq Y$, and all $y \in Y$, the set $A \cap R_S(y, V)$ is nonempty. This follows immediately from Theorem 4.13.

4.3.2. Dynamically thick sets from distal points

In this section, we offer a simultaneous generalization of the examples in Lemma 4.9 and Lemma 4.11.

Lemma 4.15. Let (X,T) be a minimal system and $x \in X$. For all piecewise syndetic sets $P \subseteq \mathbb{N}$, the set $T^P x$ is somewhere dense, i.e., $(\overline{T^P x})^\circ \neq \emptyset$.

Proof. Since the set P is piecewise syndetic, there exists $k \in \mathbb{N}$ such that $H := \bigcup_{i=1}^{k} (P-i)$ is thick. Since (X,T) is minimal, the set $T^{H}x$ is dense. We see that

$$X = \overline{\bigcup_{i=1}^{k} T^{-i} T^{P} x} = \bigcup_{i=1}^{k} T^{-i} \overline{T^{P} x}.$$

By the Baire Category Theorem, there exits $i \in \{1, \ldots, k\}$ such that the set $T^{-i}\overline{T^Px}$ has nonempty interior. By semiopenness of minimal maps [40, Thm 2.4], applying *i* many times the map *T*, the set $\overline{T^Px}$ has nonempty interior, as desired.

The following theorem generalizes the examples in both Lemma 4.9 and Lemma 4.11. This can be seen by considering a periodic system in the latter case and an infinite product of periodic systems of prime cardinality in the former case. Recall the definition of a distal point from Section 2.2.1.

Theorem 4.16. Let (X,T) be a minimal system and U_1, U_2, \ldots be nonempty, open subsets of X. Let $x \in X$ be a distal point and $H_1, H_2, \ldots \subseteq \mathbb{N}$ be thick sets. The set

$$A := \bigcup_{i} \left(R(x, U_i) \cap H_i \right)$$

is dynamically thick if and only if the set $\bigcup_{i=1}^{\infty} U_i$ is dense in X.

Proof. If the set $\bigcup_{i=1}^{\infty} U_i$ is not dense in X, then it is disjoint from a nonempty, open set $V \subseteq X$. The set A is clearly disjoint from the set R(x, V), whereby A is not dynamically thick.

Suppose that $\bigcup_{i=1}^{\infty} U_i$ is dense. Let (Y, S) be a minimal system, $y \in Y$, and $V \subseteq Y$ be nonempty and open. Since the set $R_S(y, V)$ is syndetic, by Lemma 4.15, the set $T^{R_S(y,V)}x$ is somewhere dense. Therefore, there exists $i \in \mathbb{N}$ such that $T^{R_S(y,V)}x \cap U_i \neq \emptyset$. It follows that the set $R_{T\times S}((x, y), U_i \times V)$ is nonempty. Since x is a distal point, by [20, Thm. 9.11], the point (x, y) is uniformly recurrent under $T \times S$. Since $U_i \times V$ is open, we have that $R_{T\times S}((x, y), U_i \times V)$ is (dynamically) syndetic. Therefore, the set $R_{T\times S}((x, y), U_i \times V) \cap H_i$ is nonempty, which implies that $A \cap R_S(y, V)$ is nonempty, as desired.

4.3.3. A general form for dynamically thick sets: a proof of Theorem F

In this section, we draw on a future result, Theorem 5.11, to describe the form of any dynamically thick set. (For the astute reader concerned about circular logic: while Theorem F relies on Theorem 5.11, we do not invoke Theorem F anywhere in the paper.)

Definition 4.17. A collection \mathscr{C} of subsets of \mathbb{N} is *robustly syndetic* if for all dynamically syndetic sets $A \subseteq \mathbb{N}$, there exists $B \in \mathscr{C}$ such that $A \cap B$ is syndetic.

The dynamically thick sets described so far all take the form $\bigcup_{B \in \mathscr{C}} (B \cap H_B)$, where \mathscr{C} is a robustly syndetic collection of sets and $(H_B)_{B \in \mathscr{C}}$ is a collection of thick sets. Indeed, in Theorem 4.14, the collection $\{R_{T_i}(x_i, U_i) \mid i \in \mathbb{N}\}$ is robustly syndetic by virtue of the disjointness of the collection of systems $(X_i, T_i), i \in \mathbb{N}$. In Theorem 4.16, the collection $\{R_T(x, U_i) \mid i \in \mathbb{N}\}$ is robustly syndetic by virtue of the fact that x is a distal point and $\bigcup_{i=1}^{\infty} U_i$ is dense in X. Theorem F shows that every dynamically thick set has an underlying robustly syndetic collection.

Proof of Theorem F. Suppose that \mathscr{C} is robustly syndetic and that, for each $B \in \mathscr{C}$, the set $H_B \subseteq \mathbb{N}$ is thick such that (1.6) holds. To see that $A \in d\mathcal{T}$, let $C \in d\mathcal{S}$. There exists $B \in \mathscr{C}$ such that $B \cap C$ is syndetic. Thus, the set $B \cap C \cap H_B$ is nonempty, whereby $A \cap C \neq \emptyset$, as desired.

Conversely, suppose that A is dynamically thick. For all $S \in dS$, we will define a thick set $G_S \subseteq \mathbb{N}$ and a set $B_S \subseteq \mathbb{N}$ in such a way that the collection $\mathscr{C} := \{B_S \mid S \in dS\}$ is robustly syndetic and

$$A \supseteq \bigcup_{S \in d\mathcal{S}} (B_S \cap G_S).$$

This suffices to reach the conclusion: the collection $\mathscr{C} \cup \{A\}$ is robustly syndetic, and setting $H_A := \mathbb{N}$ and, for $B = B_S \in \mathscr{C}$, setting $H_B := G_S$, we see that equality in (1.6) holds.

Let $S \in dS$. The set $A \cap S$ is an intersection of a dynamically thick set and a dynamically syndetic set and, hence, by Theorem 5.11, is piecewise syndetic. There exist $\ell \in \mathbb{N}$ and finite, disjoint intervals I_1, I_2, \ldots of consecutive positive integers satisfying: a) $\lim_{i\to\infty} |I_i| \to \infty$ and, for all $i \in \mathbb{N}$, b) max $I_i + i < \min I_{i+1}$; and c) the set $A \cap S$ has nonempty intersection with all subintervals of I_i of length at least ℓ .

Define $G_S := \bigcup_{i=1}^{\infty} I_i$, and note that it is thick. Define

$$B_S := (A \cap G_S) \cup (\mathbb{N} \backslash G_S). \tag{4.6}$$

Since the set $\mathbb{N}\backslash G_S$ is a disjoint union of finite intervals whose lengths tend to infinity, the set

$$B_S \cap S = (A \cap G_S \cap S) \cup ((\mathbb{N} \setminus G_S) \cap S) \text{ is syndetic.}$$

$$(4.7)$$

Indeed, the set S has nonempty intersection with any interval of length $\ell' \in \mathbb{N}$. Therefore, on G_S , the set $B_S \cap S$ has nonempty intersection with any interval of length at least ℓ , while on $\mathbb{N}\backslash G_S$, the set $B_S \cap S$ has nonempty intersection with any interval of length ℓ' .

Now (4.7) shows that the collection $\mathscr{C} := \{B_S \mid S \in d\mathcal{S}\}$ is robustly syndetic, and (4.6) shows that $A \supseteq \bigcup_{S \in d\mathcal{S}} (B_S \cap G_S)$, as desired.

It would be interesting and useful to improve Theorem F by saying more about the robustly syndetic collection \mathscr{C} . We discuss this further in Question 6.8.

4.4. The partition problem and σ -compactness

It is natural to ask whether any dynamically thick set can be partitioned into two disjoint dynamically thick sets. While we were not able to answer this question – see Question 6.7 for some discussion on how the structure result in Theorem F may help – we were able to show that the established technique for answering this type of question does not work.

The established technique goes as follows. Let \mathscr{F} be a family of subsets of \mathbb{N} , and suppose we wish to show that for all $A \in \mathscr{F}$, there exists a disjoint union $A = A_1 \cup A_2$ such that $A_1, A_2 \in \mathscr{F}$. Let $N_1 < N_2 < \cdots$ be a rapidly increasing sequence of positive integers. Assign the set $A \cap [N_1]$ to A_1 , the set $A \cap ([N_2] \setminus [N_1])$ to A_2 , the set $A \cap ([N_3] \setminus [N_2])$ to A_1 , and so forth. If the family \mathscr{F} is "suitable" and the N_i 's are chosen appropriately, the sets A_1 and A_2 will both belong to \mathscr{F} . This outline is realized in detail in the proof of Theorem 4.25.

There are several instances of this in the literature [15, 43, 44], but to our knowledge, sufficient and necessary conditions for "suitability" have never been explained. In all instances that we know of, "suitability" can be succinctly captured in terms of compactness. We show in Theorem 4.25 below that if the dual family \mathscr{F}^* is σ -compact (Definition 4.18), then all members of \mathscr{F} can be partitioned into two disjoint members of \mathscr{F} . We give several examples of σ -compact families that arise naturally in the subject.

The main result in this section is Theorem 4.30: the families $d\mathcal{T}^* = dS$ and $dc\mathcal{T}^* = dcS$ are not σ -compact. This demonstrates why the usual procedure for splitting dynamically (central) thick sets into two such sets fails. It also indicates that these families are more complex than others usually considered in the subject.

4.4.1. Compact and σ -compact families

The terminology in the following definition is motivated in Theorem 4.19 below.

Definition 4.18. Let \mathscr{F} be a family of subsets of \mathbb{N} .

- (i) The family \mathscr{F} is *compact* if for all $B \in \mathscr{F}^*$, there exists $M \in \mathbb{N}$ such that $B \cap [M] \in \mathscr{F}^*$.
- (ii) The family \mathscr{F} is σ -compact if it is the union of countably many compact families.

Before giving some examples of compact and σ -compact families, we move to explain the terminology. For a family \mathscr{F} of subsets of \mathbb{N} , define

$$X_{\mathscr{F}} := \left\{ 1_A \in \{0,1\}^{\mathbb{N}} \mid A \in \mathscr{F} \right\}.$$

Thus, the set $X_{\mathscr{F}}$ is a subset of the compact metric space $\{0,1\}^{\mathbb{N}}$.

Theorem 4.19. A family \mathscr{F} is compact (resp. σ -compact) if and only if the set $X_{\mathscr{F}}$ is compact (resp. σ -compact).

Proof. Since $X_{\bigcup_{N=1}^{\infty} \mathscr{F}_N} = \bigcup_{N=1}^{\infty} X_{\mathscr{F}_N}$, it suffices to show that \mathscr{F} is compact if and only if $X_{\mathscr{F}}$ is compact.

Suppose $X_{\mathscr{F}}$ is a compact subset of $\{0,1\}^{\mathbb{N}}$. Assume for the sake of a contradiction that \mathscr{F} is not a compact family. Thus, there exists $B \in \mathscr{F}^*$ such that for every $M \in \mathbb{N}$, there exists $A_M \in \mathscr{F}$ such that

$$B \cap [M] \cap A_M = \emptyset. \tag{4.8}$$

Since $X_{\mathscr{F}}$ is compact, the sequence $(1_{A_M})_{M \in \mathbb{N}} \subseteq X_{\mathscr{F}}$ has a subsequence that converges to some point $1_A \in X_{\mathscr{F}}$. By the product topology on $\{0,1\}^{\mathbb{N}}$ and (4.8), we see that $B \cap A = \emptyset$, in contradiction with the fact that $A \in \mathscr{F}$ and $B \in \mathscr{F}^*$.

Conversely, suppose that \mathscr{F} is a compact family. We will show that $X_{\mathscr{F}}$ is compact by showing that it is closed. Let $(x_k)_{k\in\mathbb{N}} \subseteq X_{\mathscr{F}}$ be a convergent sequence with limit point $x \in \{0,1\}^{\mathbb{N}}$. We will show that $x \in X_{\mathscr{F}}$ by showing that for all $B \in \mathscr{F}^*$, the set supp $(x) \cap B$ is nonempty.

Let $B \in \mathscr{F}^*$. Since \mathscr{F} is compact, there exists $M \in \mathbb{N}$ such that for all $k \in \mathbb{N}$,

$$B \cap [M] \cap \operatorname{supp}(x_k) \neq \emptyset.$$

Since $\lim_{k\to\infty} x_k = x$ and $B \cap [M]$ is a finite set, we see that $B \cap [M] \cap \text{supp}(x) \neq \emptyset$. In particular, the set $B \cap \text{supp}(x)$ is nonempty, as was to be shown.

Next we provide some examples of compact and σ -compact families.

Example 4.20. For $N \in \mathbb{N}$, denote by S_N the family of sets that have nonempty intersection with every interval in \mathbb{N} of length N. The family S_N is compact. Indeed, given $B \in S_N^*$, the set B is thick. If $M \in \mathbb{N}$ is large so that $B \cap [M]$ contains an interval of length at least N, then $B \cap [M] \in S_N^*$. We see that the family of syndetic sets, S, is equal to $\bigcup_{N=1}^{\infty} S_N$, whereby it is σ -compact.

Example 4.21. Let $N \in \mathbb{N}$. Denote by \mathcal{IP}_N the upward closure of the collection of sets of the form

$$\operatorname{FS}(x_i)_{i=1}^N := \left\{ \sum_{f \in F} x_f \mid \emptyset \neq F \subseteq \{1, \dots, N\} \right\}, \quad x_1, \dots, x_N \in \mathbb{N}.$$

As in the previous example, the family \mathcal{IP}_N^* is easily seen from the definition to be compact. The dual of the family $\mathcal{IP}_0 := \bigcap_{N=1}^{\infty} \mathcal{IP}_N$ is σ -compact, since $\mathcal{IP}_0^* = \bigcup_{N=1}^{\infty} \mathcal{IP}_N^*$. The family \mathcal{IP}_0^* is important in quantitative strengthenings of recurrence theorems and appears again in this paper in Section 6.3.

Example 4.22. The family of sets of return times in minimal systems is σ -compact. More precisely, let \mathscr{F} be the upward closure of collection of sets of the form

$$\{n \in \mathbb{N} \mid U \cap T^{-n}U \neq \emptyset\},\tag{4.9}$$

where (X,T) is a minimal system and $U \subseteq X$ is a nonempty, open set. The family \mathscr{F} is σ -compact, as we will now show.

It is a fact [8, Thm. 2.4] that $A \in \mathscr{F}$ if and only if there exists a syndetic set $S \subseteq \mathbb{N}$ such that $A \supseteq S - S$. Thus, using the notation from (4.20), if for $N \in \mathbb{N}$ we define

$$\mathscr{F}_N := \uparrow \{ S - S \mid S \in \mathcal{S}_N \},\$$

we have that $\mathscr{F} = \bigcup_{N=1}^{\infty} \mathscr{F}_N$. We have only to show that for all $N \in \mathbb{N}$, the family \mathscr{F}_N is

compact. Let $N \in \mathbb{N}$ and $B \in \mathscr{F}_N^*$. We must show that there exists $M \in \mathbb{N}$ such that for all $A \in \mathscr{F}_N$, the set $A \cap B \cap [M]$ is nonempty.

Assume for the sake of a contradiction that no such M exists. Thus, for all $M \in \mathbb{N}$, there exists $S_M \in \mathcal{S}_N$ such that

$$(S_M - S_M) \cap B \cap [M] = \emptyset. \tag{4.10}$$

Since S_N is compact, passing to a subsequence of $(S_M)_{M\in\mathbb{N}}$, there exists $S \in S_N$ such that $\lim_{M\to\infty} 1_{S_M} = 1_S$. Since $S \in S_N$, the set S - S is a member of \mathscr{F}_N . It follows from (4.10) that $(S - S) \cap B = \emptyset$, contradicting the fact that $B \in \mathscr{F}_N^*$.

Example 4.23. In analogy to Example 4.22, the family of sets of return times in measure preserving systems is σ -compact. More precisely, the upward closure \mathscr{F} of the collection of sets of the form

$$R(E,E) := \{ n \in \mathbb{N} \mid \mu(E \cap T^{-n}E) > 0 \},$$
(4.11)

where (X, μ, T) is a measure preserving system (see Section 5.3) and $E \subseteq X$ is a set of positive measure, is σ -compact. This is most readily seen as a consequence of a result of Forrest [15]: for all sets $B \subseteq \mathbb{N}$ of single measurable recurrence (i.e. for all $B \in \mathscr{F}^*$) and all $\delta > 0$, there exists $M \in \mathbb{N}$ such that for all sets E with measure at least δ , the set $B \cap R(E, E) \cap [M]$ is nonempty. For $N \in \mathbb{N}$, defining

$$\mathscr{F}_N := \uparrow \big\{ R(E,E) \subseteq \mathbb{N} \mid (X,\mu,T) \text{ is a measure preserving system, } E \subseteq X, \, \mu(E) > 1/N \big\},$$

we see that \mathscr{F}_N is compact and $\mathscr{F} = \bigcup_{N=1}^{\infty} \mathscr{F}_N$ is σ -compact.

The previous two examples concerned set recurrence, while the following ones concern pointwise recurrence.

Example 4.24. Bohr₀ sets and Nil_k-Bohr₀ sets are special dynamically central syndetic sets for which the systems in consideration are minimal rotations on compact abelian groups and minimal k-step nilsystems, respectively. (See [34] for a detailed definition of Nil_k-Bohr₀ sets.) It can be shown that these families are σ -compact using the results in [43, Prop. 1.4] and [44, Lemma 3.3], respectively.

4.4.2. The partition problem in dual compact families

The following result offers, under a mild assumption on $\mathscr{F} \sqcap \mathscr{F}^*$, a solution to the partition problem in families whose duals are σ -compact.

Theorem 4.25. Let \mathscr{F} be a family of subsets of \mathbb{N} with the property that all members of $\mathscr{F} \sqcap \mathscr{F}^*$ are infinite. If \mathscr{F}^* is σ -compact, then for all $A \in \mathscr{F}$, there exists a disjoint partition $A = A_1 \cup A_2$ with $A_1, A_2 \in \mathscr{F}$.

Proof. Suppose \mathscr{F}^* is σ -compact. There exists $\mathscr{F}^* = \bigcup_{N=1}^{\infty} \mathscr{G}_N$ where each family \mathscr{G}_N is compact. By taking finite unions, we may assume without loss of generality that $\mathscr{G}_1 \subseteq \mathscr{G}_2 \subseteq \cdots$.

Let $A \in \mathscr{F}$. Since $\mathscr{G}_1 \subseteq \mathscr{F}^*$, the set A belongs to \mathscr{G}_1^* . Since \mathscr{G}_1 is compact, there exists a finite set $A'_1 \subseteq A$ such that $A'_1 \in \mathscr{G}_1^*$. Since all members of $\mathscr{F} \sqcap \mathscr{F}^*$ are infinite, for all $B \in \mathscr{F}^*$, the set $A \cap B$ is infinite, whereby

$$(A \setminus A_1') \cap B \neq \emptyset.$$

Since $B \in \mathscr{F}^*$ was arbitrary, we see that $A \setminus A'_1 \in \mathscr{F}$. Repeating the argument for $A \setminus A'_1$ in place of A, we get a finite set $A'_2 \subseteq A \setminus A'_1$ such that $A'_2 \in \mathscr{G}_2^*$. By the same argument as before, the set $A \setminus (A'_1 \cup A'_2)$ is a member of \mathscr{F} , allowing us to continue on as before.

Repeating ad infinitum, we get a sequence A'_1, A'_2, \ldots of disjoint, finite subsets of A such that for all $N \in \mathbb{N}$, the set $A'_N \in \mathscr{G}^*_N$. Define $A_1 := A'_1 \cup A'_3 \cup \cdots$ and $A_2 := A'_2 \cup A'_4 \cup \cdots$. By appending any elements in $A \setminus (A_1 \cup A_2)$ to A_1 , we may assume that $A = A_1 \cup A_2$.

We need only to show that A_1 and A_2 are members of \mathscr{F} . Indeed, since $\mathscr{F}^* = \bigcup_{N=1}^{\infty} \mathscr{G}_N$, we see that $\mathscr{F} = (\mathscr{F}^*)^* = \bigcap_{N=1}^{\infty} \mathscr{G}_N^*$. Since $\mathscr{G}_1^* \supseteq \mathscr{G}_2^* \supseteq \cdots$, by the construction of the A_i 's, both A_1 and A_2 belong to $\bigcap_{N=1}^{\infty} \mathscr{G}_N^*$.

Remark 4.26. In view of Theorem 4.25 and the examples of σ -compact families presented in Section 4.4.1, every set in the following list of families can be partitioned into two sets in the same family:

- (i) sets of topological recurrence (dual of sets of the form in (4.9)),
- (ii) sets of measurable recurrence (dual of sets of the form in (4.11)),
- (iii) sets of Bohr recurrence (dual of Bohr₀ sets),

(iv) sets of pointwise recurrence for k-step nilsystems (dual of Nil_k -Bohr₀ sets).

These facts are well known to experts, but the framing in terms of compactness is, as far as we know, new.

4.4.3. The family of dynamically syndetic sets is not σ -compact

We show in this section that the families dS and dcS are not σ -compact.

Theorem 4.27. Let \mathscr{G} be a family of subsets of \mathbb{N} . If

there exist
$$B_1, B_2, \ldots \in \mathscr{G}$$
, disjoint, such that for all $B \in \mathscr{G}$
with $B \subseteq \bigcup_{i=1}^{\infty} B_i$, there exists $i \in \mathbb{N}$ such that $|B \cap B_i| = \infty$, (4.12)

then the family \mathscr{G}^* is not σ -compact.

Proof. Suppose property (4.12) holds, and suppose for a contradiction that $\mathscr{G}^* = \bigcup_{i=1}^{\infty} \mathscr{F}_i$, where $\mathscr{F}_1, \mathscr{F}_2, \ldots$ are compact families.

Let $i \in \mathbb{N}$. We claim that there exists a function $\varphi_i : \mathscr{G} \to \mathbb{N}$ such that for all $A \in \mathscr{F}_i$ and for all $B \in \mathscr{G}$,

$$\min(A \cap B) \leqslant \varphi_i(B).$$

Indeed, fix $B \in \mathscr{G}$, and consider the function $X_{\mathscr{F}_i} \to \mathbb{N}$ defined by $1_A \mapsto \min(A \cap B)$. It is locally constant since, more generally, the function $X_{\mathscr{G}} \times X_{\mathscr{G}^*} \to \mathbb{N}$ defined by $(1_A, 1_B) \mapsto \min(A \cap B)$ is locally constant. Since \mathscr{F}_i is compact, the function $X_{\mathscr{F}_i} \to \mathbb{N}$ is bounded from above. We define $\varphi_i(B)$ to be an upper bound. Define the set

$$A := \left(\mathbb{N} \setminus \bigcup_{i=1}^{\infty} B_i\right) \cup \bigcup_{i=1}^{\infty} \left(B_i \setminus \{1, \dots, \varphi_i(B_i)\}\right).$$

We claim that $A \in \mathscr{G}^*$. Indeed, if $B \in \mathscr{G}$, then either $B \cap (\mathbb{N} \setminus \bigcup_{i=1}^{\infty} B_i) \neq \emptyset$, in which case $A \cap B \neq \emptyset$, or $B \subseteq \bigcup_{i=1}^{\infty} B_i$. In the latter case, by (4.12), there exists $i \in \mathbb{N}$ such that $|B \cap B_i| = \infty$. It follows that $B \cap (B_i \setminus \{1, \ldots, \varphi_i(B_i)\}) \neq \emptyset$, whereby $A \cap B \neq \emptyset$.

Since $A \in \mathscr{G}^*$, there exists $j \in \mathbb{N}$ such that $A \in \mathscr{F}_j$. It follows that $\min(A \cap B_j) \leq \varphi_j(B_j)$. But since the B_j 's are disjoint, we see from the definition of A that

$$A \cap B_j = B_j \setminus \{1, \dots, \varphi_j(B_j)\}$$

whereby $\min(A \cap B_j) > \varphi_j(B_j)$, a contradiction.

The following lemma gives a sufficient condition to satisfy the property in (4.12). Recall that all difference sets are computed in \mathbb{N} .

Lemma 4.28. Let \mathscr{G} be a family of subsets of \mathbb{N} . If

for all
$$B \in \mathscr{G}$$
, there exists $n \in \mathbb{N}$ such that $|B \cap (B - n)| = \infty$, (4.13)

and if

there exist
$$B_1, B_2, \dots \in \mathscr{G}$$
, disjoint, such that

$$\lim_{d \to \infty} \min \bigcup_{\substack{i,j=1\\\max(i,j) > d}}^{\infty} (B_i - B_j) = \infty,$$
(4.14)

then the property in (4.12) holds.

Proof. Suppose the properties in (4.13) and (4.14) hold. We will show that the property in (4.12) holds for the same sets B_1, B_2, \ldots

Let $B \in \mathscr{G}$ be such that $B \subseteq \bigcup_{i=1}^{\infty} B_i$. By (4.13), there exists $n \in \mathbb{N}$ such that $|B \cap (B - n)| = \infty$. Writing $B = B \cap \bigcup_{i=1}^{\infty} B_i$, we see that

$$B \cap (B-n) = \bigcup_{i,j=1}^{\infty} (B \cap B_i \cap (B-n) \cap (B_j - n)).$$

By (4.14), there exists $d \in \mathbb{N}$ such that if $\max(i, j) > d$, then $B_i \cap (B_j - n) = \emptyset$. Therefore,

$$B \cap (B-n) = \bigcup_{i,j=1}^{d} \left(B \cap B_i \cap (B-n) \cap (B_j-n) \right) \subseteq \bigcup_{i=1}^{d} \left(B \cap B_i \right).$$

By assumption, the set on the left hand side is infinite, so there must exist an $i \in \{1, \ldots, d\}$ such that $B \cap B_i$ is infinite. This verifies (4.12).

The result in the following lemma is well known. We include a proof for completeness.

Lemma 4.29. If $B \subseteq \mathbb{N}$ is piecewise syndetic, then there exists $n \in \mathbb{N}$ such that the set $B \cap (B - n)$ is piecewise syndetic.

Proof. Suppose $B \subseteq \mathbb{N}$ is piecewise syndetic. There exists a syndetic set $S \subseteq \mathbb{N}$ and a thick set $H \subseteq \mathbb{N}$ such that $B = S \cap H$. There exists $N \in \mathbb{N}$ such that $\bigcup_{i=1}^{N} (S-i) = \mathbb{N}$, and so $\bigcup_{i=1}^{N} (B-i) \supseteq \bigcap_{i=1}^{N} (H-i)$. It follows that the set

$$B \cap \bigcup_{i=1}^{N} (B-i) \supseteq S \cap H \cap \bigcap_{i=1}^{N} (H-i)$$

is piecewise syndetic. Since the family \mathcal{PS} is partition regular, there exists $n \in \{1, \ldots, N\}$ such that the set $B \cap (B - n)$ is piecewise syndetic.

Theorem 4.30. The families $d\mathcal{T}^* = d\mathcal{S}$ and $dc\mathcal{T}^* = dc\mathcal{S}$ are not σ -compact.

Proof. We will show that $d\mathcal{T}$ and $dc\mathcal{T}$ satisfy the conditions in Lemma 4.28 and hence in (4.12). It will follow by Theorem 4.27 that the families $d\mathcal{T}^* = d\mathcal{S}$ and $dc\mathcal{T}^* = dc\mathcal{S}$ are not σ -compact.

To see that (4.13) holds for $d\mathcal{T}$ and $dc\mathcal{T}$, let $B \in d\mathcal{T}$ or $B \in dc\mathcal{T}$. By Theorem 4.2, the set B is piecewise syndetic. Thus, by Lemma 4.29, there exists $n \in \mathbb{N}$ for which the set $B \cap (B-n)$ is piecewise syndetic and hence infinite.

We will show that property (4.14) in Lemma 4.28 holds for the family $d\mathcal{T}$. Since $d\mathcal{T} \subseteq dc\mathcal{T}$, it will follow that $dc\mathcal{T}$ also satisfies (4.14). Let $\{T_{i,j} \mid i, j \in \mathbb{N}\}$ be a doubly-indexed family of thick subsets of \mathbb{N} that are well separated, in the sense that

$$\lim_{d \to \infty} \min \bigcup_{\substack{i,j,k,\ell \in \mathbb{N} \\ (i,j) \neq (k,\ell) \\ \max(i,j,k,\ell) > d}} \min \left(T_{i,j} - T_{k,\ell} \right) = \infty,$$
(4.15)

where the notation (i, j) denotes an element of \mathbb{N}^2 . Let p_1, p_2, \ldots be an increasing enumeration of the primes. For $i \in \mathbb{N}$, define

$$B_i = \bigcup_{j=i}^{\infty} \left(T_{i,j} \cap (p_j \mathbb{N} + 1) \right).$$

The sets B_1, B_2, \ldots are disjoint, and by Lemma 4.9, each is dynamically thick. To verify the limit in (4.14), note that for all $i, k \in \mathbb{N}$,

$$B_i - B_k = \bigcup_{\substack{j=i\\\ell=k}}^{\infty} \left(\left(T_{i,j} \cap (p_j \mathbb{N} + 1) \right) - \left(T_{k,\ell} \cap (p_\ell \mathbb{N} + 1) \right) \right) \subseteq \bigcup_{\substack{j=i\\\ell=k}}^{\infty} \left(T_{i,j} - T_{k,\ell} \right).$$

For all $d \in \mathbb{N}$, considering the cases i = k and $i \neq k$ separately, we see that

$$\min \bigcup_{\substack{i,k=1\\\max(i,k)>d}}^{\infty} \left(B_i - B_k\right) \ge \min\left(\bigcup_{\substack{i,j,k,\ell \in \mathbb{N}\\(i,j)\neq (k,\ell)\\\max(i,j,k,\ell)>d}}\min\left(T_{i,j} - T_{k,\ell}\right), p_d\right).$$

The limit in (4.14) follows now from the limit in (4.15) and the fact that $\lim_{d\to\infty} p_d = \infty$, as desired.

Remark 4.31. Using Lemma 4.28, one can show that the families

$$\mathcal{IP}, \mathcal{IP}_0, \mathcal{C}, \mathcal{PS}, d\mathcal{T}, dc\mathcal{T}, \mathcal{T}, \mathcal{IP}^*, \mathcal{C}^*, \mathcal{PS}^*$$

are not σ -compact. However, members of the families $dS = d\mathcal{T}^*$, $dcS = dc\mathcal{T}^*$, $\mathcal{S} = \mathcal{T}^*$, $\mathcal{IP} = (\mathcal{IP}^*)^*$, $\mathcal{C} = (\mathcal{C}^*)^*$, $\mathcal{PS} = (\mathcal{PS}^*)^*$ can be partitioned into two sets belonging to the same family. This demonstrates that σ -compactness of \mathscr{F}^* is sufficient, but not necessary, for every member of \mathscr{F} to be partitioned into two members of \mathscr{F} .

5. Part III: Dynamically piecewise syndetic sets

In this section, in analogy to the family of piecewise syndetic sets, we introduce and study the families of dynamically (central) piecewise syndetic sets. The main results describe the relationships these families have with other established families (Theorem G) and show that dynamically central piecewise syndetic sets are sets of polynomial multiple measurable recurrence for commuting transformations (Theorem H).

5.1. Definitions and first results

Recall that dS and dcS denote the families of dynamically syndetic and dynamically central syndetic subsets of \mathbb{N} . Their duals, the families of dynamically thick sets and sets of pointwise recurrence, are denoted $d\mathcal{T}$ and $dc\mathcal{T}$.

Definition 5.1. Let $A \subseteq \mathbb{N}$.

(i) The set A is dynamically piecewise syndetic if there exists a dynamically syndetic set $B \subseteq \mathbb{N}$ and a dynamically thick set $C \subseteq \mathbb{N}$ such that $A = B \cap C$. The family of dynamically piecewise syndetic sets is

$$d\mathcal{PS} := d\mathcal{S} \sqcap d\mathcal{T}.$$

(ii) The set A is dynamically central piecewise syndetic if there exists a dynamically central syndetic set $B \subseteq \mathbb{N}$ and a set of pointwise recurrence $C \subseteq \mathbb{N}$ such that $A = B \cap C$. The family of dynamically central piecewise syndetic sets is

$$dc\mathcal{PS} := dc\mathcal{S} \sqcap dc\mathcal{T}.$$

Examples of dynamically (central) piecewise syndetic sets can be gotten by intersecting the examples of dynamically (central) syndetic and dynamically (central) thick sets given in Sections 3.1 and 4.1, respectively. We will show next that return times of distal points to neighborhoods of themselves provide nice dynamical examples of members of the family $dc\mathcal{PS}^*$.

Example 5.2. Let (X,T) be a system, $x \in X$ be a distal point, and $U \subseteq X$ be an open neighborhood of x. We claim that the set $R_T(x,U)$ is a $dc\mathcal{PS}^*$ set. Indeed, by Lemma 2.1, it suffices to show that for all $B \in dcS$, the set $R_T(x,U) \cap B \in dcS$.

Let $B \in dcS$. There exists a minimal system (Y, S), a point $y \in Y$, and an open set $V \subseteq Y$ containing y such that $R_S(y, V) \subseteq B$. By [20, Thm. 9.11], the point $(x, y) \in X \times Y$ is $T \times S$ uniformly recurrent, whereby the set $R_{T \times S}((x, y), U \times V) = R_T(x, U) \cap R_S(y, V)$ is dynamically central syndetic. It follows that $R_T(x, U) \cap B \in dcS$, as desired.

5.1.1. Containment and partition regularity

As $dcS \subseteq dS$ and $dcT \supseteq dT$, the relationship between the families dcPS an dPS is, apriori, not clear. We show in the following lemma that $dcPS \subseteq dPS$, but the proof uses Theorems 5.9 and 5.11, which ultimately rely on the main result in this paper, Theorem 3.18. It would be of interest to find a simpler proof of Theorem 5.3. (A note to the astute reader concerned about circular logic. Prior to Theorems 5.9 and 5.11, on which Theorem 5.3 depends, we use Theorem 5.3 only in the proof of (5.2) in Lemma 5.5. Theorems 5.9 and 5.11 depend only on item (i) in Lemma 5.5, and hence does not rely on Theorem 5.3.)

Theorem 5.3. Dynamically central piecewise syndetic subsets are dynamically piecewise syndetic, that is, $dc\mathcal{PS} \subseteq d\mathcal{PS}$.

Proof. It follows by combining Theorems 5.9 and 5.11 below that $dc\mathcal{PS} \subseteq \mathcal{PS} \subseteq d\mathcal{PS}$, as was to be shown.

Lemma 5.4. The families dcPS and dPS are partition regular, and their duals, $dcPS^*$ and dPS^* , are filters.

Proof. That the families $dc\mathcal{PS}$ and $d\mathcal{PS}$ are partition regular follows immediately from the definition and Lemma 2.1. It follows then from the discussion in Section 2.1.2 that the families $dc\mathcal{PS}^*$ and $d\mathcal{PS}^*$ are filters.

5.1.2. Translates and dilates of dynamically piecewise syndetic sets

The following lemmas clarify how members of the families $dc\mathcal{PS}$ and $d\mathcal{PS}$ behave under translations and dilations. Point (ii) is upgraded in Corollary 5.12 below.

Lemma 5.5. Let $A \subseteq \mathbb{N}$ be dynamically piecewise syndetic.

- (i) For all $n \in \mathbb{N}$, the set A n is dynamically piecewise syndetic.
- (ii) The set

$$\{n \in \mathbb{N} \mid A - n \text{ is dynamically central piecewise syndetic}\}$$
(5.1)

is dynamically syndetic.

Moreover, points (i) and (ii) hold with A - n replaced by A + n. Finally,

$$d\mathcal{PS} = \bigcup_{n \in \mathbb{N}} \left(dc \mathcal{PS} - n \right) = \bigcup_{n \in \mathbb{N}} \left(dc \mathcal{PS} + n \right).$$
(5.2)

Proof. Because the families dS and dT are translation invariant (by Lemmas 3.4 and 4.5), it follows by Lemma 2.4 that the family $d\mathcal{PS} = dS \sqcap dT$ is translation invariant, showing (i).

To see (ii), because A is dynamically piecewise syndetic, there exists a dynamically syndetic set $B \subseteq \mathbb{N}$ and a dynamically thick set $C \subseteq \mathbb{N}$ such that $A = B \cap C$. By Lemma 3.4, there are dynamically syndetically many n's for which B - n is dynamically central syndetic. By Lemma 4.5, for all $n \in \mathbb{N}$, the set C - n is dynamically thick. Therefore, the set in (5.1) is dynamically syndetic, as desired. The same argument can be shown abstractly using prior results about dynamically syndetic sets (from Lemma 3.4) and family algebra (from Section 2.1.3):

$$d\mathcal{PS} = d\mathcal{S} \sqcap d\mathcal{T} \subseteq (d\mathcal{S} + dc\mathcal{S}) \sqcap (d\mathcal{S} + dc\mathcal{T}) \subseteq (d\mathcal{S} \sqcap d\mathcal{S}) + (dc\mathcal{S} \sqcap dc\mathcal{T}) = d\mathcal{S} + dc\mathcal{PS}.$$

The arguments in the previous two paragraphs combine with the positive-translate results in Lemmas 3.4 and 4.5 to show that points (i) and (ii) hold with A - n replaced by A + n. Finally, that (5.2) holds follows immediately and (i) and (ii) for A - n and A + n and from the fact that $dc\mathcal{PS} \subseteq d\mathcal{PS}$ (from Theorem 5.3).

The conclusions of Lemma 5.5 (i) and (ii) formulated in terms of the family algebra developed in Section 2.1 read: $d\mathcal{PS} \subseteq \{\mathbb{N}\} + d\mathcal{PS}$ and $d\mathcal{PS} \subseteq d\mathcal{S} + dc\mathcal{PS}$, respectively. In analogy to Lemma 3.4 (iii), we would like to show that $dc\mathcal{PS} \subseteq dc\mathcal{S} + dc\mathcal{PS}$, that is, that if A is dynamically central piecewise syndetic, then the set in (4.2) is dynamically central syndetic. We were not able to show this. It should be easier to determine whether or not the family $dc\mathcal{PS}$ is idempotent: $dc\mathcal{PS} \subseteq dc\mathcal{PS} + dc\mathcal{PS}$. See Question 6.11 below for more discussion.

Lemma 5.6. Let $A \subseteq \mathbb{N}$ and $k \in \mathbb{N}$. If A is dynamically central piecewise syndetic, then so are the sets kA and A/k. If A is dynamically piecewise syndetic, then so is the set kA.

Proof. Suppose $A \in dc\mathcal{PS}$, and write $A = B \cap C$ where $B \in dc\mathcal{S}$ and $C \in dc\mathcal{T}$. We see that $kA = kB \cap kC$ and $A/k = B/k \cap C/k$, both of which are $dc\mathcal{PS}$ sets in view of Lemmas 3.6 and 4.6.

Suppose $A \in d\mathcal{PS}$, and write $A = B \cap C$ where $B \in d\mathcal{S}$ and $C \in d\mathcal{T}$. To see that $kA \in d\mathcal{PS}$, we will write

$$kA = kB \cap kC = kB \cap \left(kC \cup (\mathbb{N} \setminus k\mathbb{N})\right)$$

and show that $kB \in dS$ and $C' := kC \cup (\mathbb{N} \setminus k\mathbb{N}) \in d\mathcal{T}$.

That $kB \in dS$ follows from Lemma 3.6. To see that $C' \in d\mathcal{T}$, let $D \in dS$. If $D \not\subseteq k\mathbb{N}$, then $D \cap C' \neq \emptyset$. Otherwise, we have that $D \subseteq k\mathbb{N}$. By Lemma 3.4, there exists $n \in \mathbb{N}$ such that $D - n \in dcS$. Since every dcS set contains a multiple of k (by Lemma 3.6), it must be that $n \in k\mathbb{N}$. By Lemma 3.6, the set $(D - n)/k = D/k - n/k \in dcS$. According to Lemma 3.4, the set $D/k \in dS$. It follows that $(D/k) \cap C \neq \emptyset$, whereby $D \cap C' \neq \emptyset$. Since $D \in dS$ was arbitrary, it follows that $C' \in d\mathcal{T}$, as desired.

It is a consequence of Theorem G (proved below independently of Lemma 5.6) that the families of piecewise syndetic sets and dynamically piecewise syndetic sets are the same. It will follow easily from that that dilates of $d\mathcal{PS}$ sets are $d\mathcal{PS}$ sets, offering another proof of the latter statement in Lemma 5.6.

5.2. Filter algebra for central and piecewise syndetic family duals

In this section, we draw on Theorems 3.13 and 3.18 to prove some abstract results concerning special classes of filters. These results ultimately will allow us to relate the families $dc\mathcal{PS}$ and $d\mathcal{PS}$ to themselves and to the families C and \mathcal{PS} in the next section.

Theorem 5.7. If \mathscr{F} is a syndetic, idempotent filter and \mathscr{G} is a \mathcal{C}^* , idempotent filter, then $\mathscr{F} \sqcap \mathscr{G}$ is a syndetic, idempotent filter.

Proof. Since \mathscr{F} and \mathscr{G} are filters, the family $\mathscr{F} \sqcap \mathscr{G}$ is a filter. Thus, to see that it is a syndetic filter, it suffices to show that for all $A \in \mathscr{F}$ and $B \in \mathscr{G}$, the set $A \cap B$ is syndetic. Let $A \in \mathscr{F}$, $B \in \mathscr{G}$, and $H \subseteq \mathbb{N}$ be thick. Since the set A is central syndetic, by Theorem 3.13, the set $A \cap H$ is central. Since $B \in \mathscr{C}^*$, the set $A \cap H \cap B$ is nonempty. Because H was an arbitrary

thick set, we see that the set $A \cap B$ is syndetic, as desired. Finally, that $\mathscr{F} \sqcap \mathscr{G}$ is idempotent follows immediately from Lemma 2.4 and the idempotency of \mathscr{F} and \mathscr{G} .

Theorem 5.8. Let \mathscr{G} be a filter on \mathbb{N} .

(i) If \mathscr{G} is a \mathcal{C}^* , idempotent filter, then $\mathscr{G} \subseteq dc \mathcal{PS}^*$.

(ii) If \mathscr{G} is a syndetic, translation-invariant filter, then $\mathscr{G} \subseteq d\mathcal{PS}^*$.

Proof. (i) Suppose that \mathscr{G} is a \mathcal{C}^* , idempotent filter. To see that $\mathscr{G} \subseteq dc\mathcal{PS}^*$, it suffices by Lemma 2.1 to show that for all $A \in dcS$ and $B \in \mathscr{G}$, the set $A \cap B \in dcS$. Let $A \in dcS$ and $B \in \mathscr{G}$. By Theorem 3.18, the set A belongs to a syndetic, idempotent filter \mathscr{F} . By Theorem 5.7, the family $\mathscr{F} \sqcap \mathscr{G}$ is a syndetic, idempotent filter. Since $A \cap B \in \mathscr{F} \sqcap \mathscr{G}$, we have by Theorem 3.18 that $A \cap B \in dcS$, as desired.

(ii) Suppose that \mathscr{G} is a syndetic, translation-invariant filter. By Lemma 2.11, the family \mathcal{PS}^* contains all syndetic, translation-invariant filters, so we have that $\mathscr{G} \subseteq \mathcal{PS}^* \subseteq \mathcal{C}^*$. Note that \mathscr{G} is idempotent because it is translation invariant.

To see that $\mathscr{G} \subseteq d\mathcal{PS}^*$, it suffices by Lemma 2.1 to show that for all $A \in d\mathcal{S}$ and $B \in \mathscr{G}$, the set $A \cap B \in d\mathcal{S}$. Let $A \in d\mathcal{S}$ and $B \in \mathscr{G}$. By Lemma 3.4, there exists $n \in \mathbb{N}$ such that $A - n \in dc\mathcal{S}$. Since $B - n \in \mathscr{G} - n \subseteq \mathscr{G} \subseteq \mathcal{C}^*$ and \mathscr{G} is idempotent, it follows from (i) and Lemma 2.1 that

$$(A \cap B) - n = (A - n) \cap (B - n) \in dc\mathcal{S} \sqcap \mathscr{G} = dc\mathcal{S}.$$

Adding n, by Lemma 3.4, the set $A \cap B \in dS$, as was to be shown.

It is tempting to read Theorem 5.8 (i) as saying that the intersection of a C^* set with a dcS set is a dcS set. This does not follow from Theorem 5.8 (i) (and is, in fact, false) since not every C^* set belongs to a C^* , idempotent filter. Indeed, the set constructed in Lemma 4.9 is a dynamically thick set that is not \mathcal{IP} . The complement is an example of an \mathcal{IP}^* set, hence C^* set, that is not dS. By Theorem 3.18, this set cannot belong to a C^* , idempotent filter.

It is true, however, that every \mathcal{PS}^* set belongs to a syndetic, translation-invariant filter; indeed, by Lemma 2.11, the family \mathcal{PS}^* is itself such a filter. Therefore, it follows from Theorem 5.8 (ii) and Lemma 2.1 that the intersection of a \mathcal{PS}^* and a dcS set is a dcS set. We expand on this in Theorem 5.13 below.

5.2.1. Combinatorial and dynamical piecewise syndetic families and a proof of Theorem G

In this section, we explain the relationships between the dynamical families $dc\mathcal{PS}$ and $d\mathcal{PS}$ and the combinatorial families C and \mathcal{PS} . In particular, we will show that

$$\mathcal{C} \subsetneq dc \mathcal{PS} \subsetneq d\mathcal{PS} = \mathcal{PS},\tag{5.3}$$

thus completing the proof of Theorem G and the containments stated in Fig. 2.

Theorem 5.9. The family of central sets is properly contained in the family of dynamically central piecewise syndetic sets, i.e., $C \subsetneq dc \mathcal{PS}$, and piecewise syndetic sets are dynamically piecewise syndetic, i.e., $\mathcal{PS} \subseteq d\mathcal{PS}$.

Proof. By Lemma 3.5 and the fact that $\mathcal{T} \subseteq d\mathcal{T} \subseteq dc\mathcal{T}$, we have that

$$\mathcal{C} = dc\mathcal{S} \sqcap \mathcal{T} \subseteq dc\mathcal{S} \sqcap dc\mathcal{T} = dc\mathcal{PS},$$
$$\mathcal{PS} = dS \sqcap \mathcal{T} \subseteq dS \sqcap d\mathcal{T} = d\mathcal{PS}.$$

By Lemma 4.9, there is a set $A \in d\mathcal{T}$ which is not an \mathcal{IP} set. Since every central set is an \mathcal{IP} set, we see that the set $A \in dc\mathcal{PS}$ but $A \notin C$. Thus, the containment $C \subseteq dc\mathcal{PS}$ is proper.

Remark 5.10. Let us offer an alternative, combinatorial proof of the containment $\mathcal{PS} \subseteq d\mathcal{PS}$ in Theorem 5.9. We will show that $d\mathcal{PS}^*$ is a syndetic, translation-invariant filter and appeal to Lemma 2.11 to see that $d\mathcal{PS}^* \subseteq \mathcal{PS}^*$, whereby $\mathcal{PS} \subseteq d\mathcal{PS}$. Translation invariance of $d\mathcal{PS}^*$ follows from the positive translation invariance of $d\mathcal{PS}$ shown in Lemma 5.5 (i). That $d\mathcal{PS}^*$ is a filter is shown in Lemma 5.4. That every member of $d\mathcal{PS}^*$ is syndetic follows from the fact that $\mathcal{T} \subseteq d\mathcal{T} \subseteq d\mathcal{PS}$, whereby the family $d\mathcal{PS}^*$ is syndetic.

The following theorem, when combined with Theorem 5.9, completes the proof of the containments in (5.3) and hence the proof of Theorem G. It is interesting to note that the containment $d\mathcal{PS} \subseteq \mathcal{PS}$ follows immediately from Theorem F. Note, however, that the containment $d\mathcal{PS} \subseteq \mathcal{PS}$ was used to prove Theorem F, so it is best to see Theorem F as an enhancement to the containment $d\mathcal{PS} \subseteq \mathcal{PS}$.

Theorem 5.11. The family of dynamically (central) piecewise syndetic sets is (properly) contained the family of piecewise syndetic sets, i.e., $dcPS \subsetneq PS$ and $dPS \subseteq PS$.

Proof. By Lemma 2.11, the family \mathcal{PS}^* is a syndetic, translation-invariant filter. Since $\mathcal{PS}^* \subseteq \mathcal{C}^*$ and translation-invariance implies idempotency, we have that \mathcal{PS}^* is also a \mathcal{C}^* , idempotent filter. By Theorem 5.8, we see that $\mathcal{PS}^* \subseteq dc\mathcal{PS}^*$ and $\mathcal{PS}^* \subseteq d\mathcal{PS}^*$. It follows that $dc\mathcal{PS} \subseteq \mathcal{PS}$ and $d\mathcal{PS} \subseteq \mathcal{PS}$.

To see that $dc\mathcal{PS} \subsetneq \mathcal{PS}$, note that it follows from Example 5.2 that the set $2\mathbb{N}$ is $dc\mathcal{PS}^*$. Therefore, the set $2\mathbb{N} - 1$, which is \mathcal{PS} , is not $dc\mathcal{PS}$.

Corollary 5.12. For all dynamically piecewise syndetic sets $A \subseteq \mathbb{N}$, the set

$$\left\{ n \in \mathbb{N} \mid A - n \text{ is central} \right\}$$
(5.4)

is dynamically syndetic, that is, $d\mathcal{PS} \subseteq d\mathcal{S} + \mathcal{C}$. The same statement holds with A - n replaced by A + n. Finally,

$$d\mathcal{PS} = \bigcup_{n \in \mathbb{N}} (\mathcal{C} - n) = \bigcup_{n \in \mathbb{N}} (\mathcal{C} + n).$$
(5.5)

Proof. Let $A \in d\mathcal{PS}$. By Theorem 5.11, the set A is piecewise syndetic. By Lemma 3.5, there exists a minimal system (X,T), a point $x \in X$, a nonempty, open set $U \subseteq X$, and a thick set $H \subseteq \mathbb{N}$ such that $A \supseteq R(x,U) \cap H$. For all $n \in R(x,U)$, we see that the set $A-n \supseteq R(T^nx,U) \cap (H-n)$, which, by Lemma 3.5, is central. Thus, the set in (5.4) contains $R(x,U) \in d\mathcal{S}$, as was to be shown.

To arrive at the same result when A - n is replaced by A + n, begin by writing $A \supseteq R(x,U) \cap H$ in the same way, noting that by Lemma 3.1, we may assume that the system (X,T) is invertible. We claim that the set $R_{T^{-1}}(x,U)$ is contained in the set in (5.4) with

A-n replaced by A+n. Since the system (X, T^{-1}) is minimal (see, for example, [26, Lemma 2.7]), this will demonstrate that the set in (5.4) is dynamically syndetic.

Suppose $n \in R_{T^{-1}}(x, U)$ so that $T^{-n}x \in U$. It is quick to check that $R(T^{-n}x, U) \cap \{n + 1, n + 2, \ldots\} \subseteq R(x, U) + n$. Since $R(T^{-n}x, U) \in dcS$, it follows by Lemma 3.2 that the set $R(T^{-n}x, U) \cap \{n + 1, n + 2, \ldots\} \in dcS$. It follows by Lemma 3.5 that the set

$$A+n \supseteq \left(R(x,U)+n\right) \cap \left(H+n\right) \supseteq \left(R(T^{-n}x,U) \cap \{n+1,n+2,\ldots\}\right) \cap \left(H+n\right),$$

is central, as desired.

Finally, that (5.5) holds follows immediately from (5.4) for A - n and A + n.

Theorem 5.13. Let $A \subseteq \mathbb{N}$. The following conditions are equivalent to the set A being thickly syndetic $(A \in \mathcal{PS}^*)$:

- (i) for all $B \in d\mathcal{S}$, the set $A \cap B \in d\mathcal{S}$;
- (ii) for all $B \in d\mathcal{T}$, the set $A \cap B \in d\mathcal{T}$;
- (iii) for all $B \in \mathcal{PS}$, the set $A \cap B \in \mathcal{PS}$;
- (iv) for all $B \in \mathcal{PS}^*$, the set $A \cap B \in \mathcal{PS}^*$.

Thus, the family \mathcal{PS}^* is the largest family of sets that satisfies any one condition (equivalently, all conditions). Moreover, if the set A is thickly syndetic, then

- (v) for all $B \in dcS$, the set $A \cap B \in dcS$;
- (vi) for all $B \in dc\mathcal{T}$, the set $A \cap B \in dc\mathcal{T}$;
- (vii) for all $B \in dc \mathcal{PS}$, the set $A \cap B \in dc \mathcal{PS}$;

(viii) for all $B \in dc \mathcal{PS}^*$, the set $A \cap B \in dc \mathcal{PS}^*$.

Proof. Combining Theorems 5.9 and 5.11, we have that $\mathcal{PS}^* = d\mathcal{PS}^* \subseteq dc\mathcal{PS}^*$. That the first four statements are equivalent to A being a member of \mathcal{PS}^* then follows immediately from Lemma 2.1. If $A \in \mathcal{PS}^*$, then $A \in dc\mathcal{PS}^*$. The final four statements follow since, by Lemma 2.1, they are all equivalent to A being a member of $dc\mathcal{PS}^*$.

5.3. Polynomial set recurrence and Brauer configurations: a proof of Theorem H

5.3.1. Polynomial set recurrence: a proof of Theorem H

In this section, we will show that – in a very general sense made precise by Theorem H – dynamically central piecewise syndetic sets are sets of recurrence in ergodic theory. In what follows, a commuting probability measure preserving system $(X, \mu, T_1, \ldots, T_k)$ is a tuple consisting of a probability measure space (X, μ) together with commuting measure preserving transformations $T_i: X \to X, i = 1, \ldots, k$. The system is *invertible* if the maps $T_i, i = 1, \ldots, k$, are invertible.

Denote by \mathscr{G} the upward closure of the family of all subsets of \mathbb{N} of the form

$$R(\mu, T_i, p_{i,j}, E)_{\substack{i=1,\dots,k\\j=1,\dots,\ell}} := \left\{ n \in \mathbb{N} \mid \mu\left(\bigcap_{j=1}^{\ell} \left(\prod_{i=1}^{k} T_i^{p_{i,j}(n)}\right)^{-1} E\right) > 0 \right\},$$
(5.6)

where $k, \ell \in \mathbb{N}$, the tuple $(X, \mu, T_1, \ldots, T_k)$ is an invertible, commuting probability measure preserving system, the set $E \subseteq X$ satisfies $\mu(E) > 0$, and, for $1 \leq i \leq k, 1 \leq j \leq \ell$, the polynomial $p_{i,j} \in \mathbb{Q}[n]$ satisfies $p_{i,j}(\mathbb{Z}) \subseteq \mathbb{Z}$ and $p_{i,j}(0) = 0$. Thus, the family \mathscr{G} consists of times of returns of any set of positive measure in any probability measure space under polynomial
iterates of a finite collection of invertible, commuting measure preserving transformations. Members of the dual family, \mathscr{G}^* , are called *sets of polynomial multiple measurable recurrence for invertible commuting transformations.*

Lemma 5.14. The family \mathscr{G} is an \mathcal{IP}^* , idempotent filter.

Proof. By a corollary of the IP Polynomial Szemerédi Theorem of Bergelson and McCutcheon [9, Thm. 7.12], every member of \mathscr{G} is an \mathcal{IP}^* set. Therefore, the family \mathscr{G} is \mathcal{IP}^* .

To show that \mathscr{G} is a filter, we must show that the set

$$R(\mu, T_i, p_{i,j}, E)_{\substack{i=1,\dots,k\\j=1,\dots,\ell}} \cap R(\nu, S_i, q_{i,j}, F)_{\substack{i=1,\dots,r\\j=1,\dots,t}}$$
(5.7)

belongs to \mathscr{G} , where $(X, \mu, T_1, \ldots, T_k)$, $p_{i,j} \in \mathbb{Q}[n]$, $E \subseteq X$, $(Y, \nu, S_1, \ldots, S_r)$, $q_{i,j} \in \mathbb{Q}[n]$, and $F \subseteq Y$ are as described above. Denote by I the identity transformation on both X and Y, and consider the commuting probability measure preserving system

$$(X \times Y, \mu \otimes \nu, T_1 \otimes I, \dots, T_k \otimes I, I \otimes S_1, \dots, I \otimes S_r).$$

The set $E \times F \subseteq X \times Y$ satisfies $(\mu \otimes \nu)(E \times F) = \mu(E)\nu(F) > 0$. Moreover,

$$(\mu \otimes \nu) \left(\bigcap_{j_{1}=1}^{\ell} \bigcap_{j_{2}=1}^{t} \left(\prod_{i=1}^{k} (T_{i} \otimes I)^{p_{i,j_{1}}(n)} \cdot \prod_{i=1}^{r} (I \otimes S_{i})^{q_{i,j_{2}}(n)} \right)^{-1} (E \times F) \right) =$$

$$(\mu \otimes \nu) \left(\bigcap_{j=1}^{\ell} \left(\prod_{i=1}^{k} (T_{i} \otimes I)^{p_{i,j}(n)} \right)^{-1} (E \times F) \cap \bigcap_{j=1}^{t} \left(\prod_{i=1}^{r} (I \otimes S_{i})^{q_{i,j}(n)} \right)^{-1} (E \times F) \right) =$$

$$\mu \left(\bigcap_{j=1}^{\ell} \left(\prod_{i=1}^{k} T_{i}^{p_{i,j}(n)} \right)^{-1} E \right) \cdot \nu \left(\bigcap_{j=1}^{t} \left(\prod_{i=1}^{r} S_{i}^{q_{i,j}(n)} \right)^{-1} F \right).$$

It follows that the set in (5.7) is equal to

$$R(\mu \otimes \nu, U_i, m_{i,j}, E \times F)_{\substack{i=1,\dots,k+r,\\j=1,\dots,\ell t}},$$

where

$$U_i := \begin{cases} T_i \otimes I & \text{if } i \leq k \\ I \otimes S_{i-k} & \text{if } i > k \end{cases}, \quad \text{and} \quad m_{i,j} := \begin{cases} p_{i,(j \mod \ell)+1} & \text{if } i \leq k \\ q_{i-k,\lceil j/\ell \rceil} & \text{if } i > k \end{cases}.$$

This shows that the set in (5.7) belongs to \mathscr{G} , as desired.

Next, we will show that \mathscr{G} is idempotent. It suffices by Remark 2.5 to show that if D is a set of the form in (5.6), then $D \subseteq D - \mathscr{G}$, that is, for all $m \in D$, the set $D - m \in \mathscr{G}$.

Let D be a set of the form in (5.6), and let $m \in D$. Define

$$E' := \bigcap_{j=1}^{\ell} \left(\prod_{i=1}^{k} T_i^{p_{i,j}(m)} \right)^{-1} E.$$

Since $m \in D$, we have that $\mu(E') > 0$. For i = 1, ..., k and $j = 1, ..., \ell$, define $q_{i,j} \in \mathbb{Q}[n]$ by

$$q_{i,j}(n) := p_{i,j}(n+m) - p_{i,j}(m).$$

Note that $q_{i,j}(\mathbb{Z}) \subseteq \mathbb{Z}$ and $q_{i,j}(0) = 0$.

By expanding the definition of E', we get

$$\prod_{j=1}^{\ell} \left(\prod_{i=1}^{k} T_{i}^{q_{i,j}(n)} \right)^{-1} E' = \prod_{j=1}^{\ell} \left(\prod_{i=1}^{k} T_{i}^{p_{i,j}(n+m)-p_{i,j}(m)} \right)^{-1} E'$$

$$= \prod_{j=1}^{\ell} \left(\prod_{i=1}^{k} T_{i}^{p_{i,j}(n+m)-p_{i,j}(m)} \right)^{-1} \left(\prod_{j'=1}^{\ell} \left(\prod_{i'=1}^{k} T_{i'}^{p_{i',j'}(m)} \right)^{-1} E \right)$$

$$\subseteq \prod_{j=1}^{\ell} \left(\prod_{i=1}^{k} T_{i}^{p_{i,j}(n+m)} \right)^{-1} E.$$

It follows that

$$R(\mu, T_i, q_{i,j}, E')_{\substack{i=1,\dots,k\\j=1,\dots,\ell}} \subseteq \left\{ n \in \mathbb{N} \; \middle| \; \mu\left(\bigcap_{j=1}^{\ell} \left(\prod_{i=1}^{k} T_i^{p_{i,j}(n+m)}\right)^{-1} E\right) > 0 \right\} = D - m,$$

whereby $D - m \in \mathscr{G}$, as desired.

Theorem 5.15. Every $dc\mathcal{PS}$ set is a \mathscr{G}^* set, that is, dynamically central piecewise syndetic sets are sets of polynomial multiple measurable recurrence for invertible commuting transformations.

Proof. By Lemma 5.14, the family \mathscr{G} is a \mathcal{C}^* , idempotent filter. Therefore, by Theorem 5.8, we have that $\mathscr{G} \subseteq dc\mathcal{PS}^*$. Taking the dual, we see $dc\mathcal{PS} \subseteq \mathscr{G}^*$, as desired.

To prove Theorem H from the introduction, we need only to remove the assumption on invertibility in Theorem 5.15.

Proof of Theorem H. Denote by \mathscr{F} the upward closure of the family of all subsets of \mathbb{N} of the form in (5.6), where $k, \ell \in \mathbb{N}$, the tuple $(X, \mu, T_1, \ldots, T_k)$ is a commuting probability measure preserving system, the set $E \subseteq X$ satisfies $\mu(E) > 0$, and, for $1 \leq i \leq k, 1 \leq j \leq \ell$, the polynomial $p_{i,j} \in \mathbb{Q}[n]$ satisfies $p_{i,j}(\mathbb{N}_0) \subseteq \mathbb{N}_0$ and $p_{i,j}(0) = 0$. To prove Theorem H, we must show that $dc\mathcal{PS} \subseteq \mathscr{F}^*$.

Let $A \in dc\mathcal{PS}$ and let $B \in \mathscr{F}$. The set B contains a set of the form $R(\mu, T_i, p_{i,j}, E)$ as described in the previous paragraph. By the measurable natural extension (cf. [9, Lemma 7.11]), there exists an invertible, commuting probability measure preserving system $(Y, \nu, S_1, \ldots, S_k)$ and a measurable set $F \subseteq Y$ such that for all $n_{i,j} \in \mathbb{N}_0$, $1 \leq i \leq k$, $1 \leq j \leq \ell$,

$$\nu\left(\bigcap_{j=1}^{\ell}\left(\prod_{i=1}^{k}S_{i}^{n_{i,j}}\right)^{-1}F\right) = \mu\left(\bigcap_{j=1}^{\ell}\left(\prod_{i=1}^{k}T_{i}^{n_{i,j}}\right)^{-1}E\right).$$
(5.8)

By Theorem 5.15, there exists $n \in A$ such that

$$\nu\left(\bigcap_{j=1}^{\ell} \left(\prod_{i=1}^{k} S_i^{p_{i,j}(n)}\right)^{-1} F\right) > 0.$$

It follows from (5.8) that $n \in R(\mu, T_i, p_{i,j}, E)$, showing that $A \cap B \neq \emptyset$, as desired.

Remark 5.16. The \mathcal{IP} polynomial Szemerédi theorem [9, Thm 7.12] shows that \mathscr{G} is an \mathcal{IP}^* family, while Theorem 5.15 shows that \mathscr{G} is an $dc\mathcal{PS}^*$ family. Because there is no containment in either direction between the families \mathcal{IP}^* and $dc\mathcal{PS}^*$, Theorem 5.15 is neither an upgrade nor an immediate corollary of the \mathcal{IP} polynomial Szemerédi theorem.

An upgrade to the conclusion of the \mathcal{IP} polynomial Szemerédi theorem is available in the case that the polynomials in (5.6) are required to be linear. In that case, the family \mathscr{G} can be shown to be \mathcal{IP}_0^* (see Section 6.3 for a definition of the family \mathcal{IP}_0^*) by a corollary of a result of Furstenberg and Katznelson [21, Thm. 10.3]. The relationship between the families $dc\mathcal{PS}^*$ and \mathcal{IP}_0^* is not known; it may be the case that $\mathcal{IP}_0^* \subseteq dc\mathcal{PS}^*$, for example. See Question 6.10 below for more discussion in this direction.

5.3.2. Dynamically central piecewise syndetic sets contain polynomial Brauer configurations

Central sets are known to contain an abundance of combinatorial configurations (cf. [32, Ch. 14]). In this section, we demonstrate that dynamically central piecewise syndetic sets also contain some combinatorial configurations. (Recall from Theorem 5.9 that central sets are dynamically central piecewise syndetic.) We speculate in Section 6.3 about the extent to which $dc\mathcal{PS}$ sets exhibit further combinatorial richness.

Theorem 5.17. Let $A \subseteq \mathbb{N}$ be dynamically central piecewise syndetic. For all $k \in \mathbb{N}$ and all $p_1, p_2, \ldots, p_k \in \mathbb{Q}[x]$ with $p_i(\mathbb{Z}) \subseteq \mathbb{Z}$ and $p_i(0) = 0$, there exist $x, y \in \mathbb{N}$ such that

$$x, y, x + p_1(y), \dots, x + p_k(y) \in A.$$
 (5.9)

Proof. By Theorem 5.11, the set A is piecewise syndetic and hence has positive upper Banach density, $d^*(A) > 0$. By a standard application of the Furstenberg Correspondence Principle ([19, Thm. 1.1], see also [46, Thm. 3.2.5]), there exists an invertible probability measure preserving system (X, μ, T) and a set $E \subseteq X$ with $\mu(E) \ge d^*(A) > 0$ such that for all $n_1, \ldots, n_k \in \mathbb{Z}$,

$$\mu(T^{-n_1}E\cap\cdots\cap T^{-n_k}E)\leqslant d^*((A-n_1)\cap\cdots\cap (A-n_k)).$$

Setting $p_0 \equiv 0$, by Theorem 5.15, there exists $y \in A$ such that

$$0 < \mu \left(E \cap T^{-p_1(y)} E \cap \dots \cap T^{-p_k(y)} E \right) \leq d^* \left(A \cap (A - p_1(y)) \cap \dots \cap (A - p_k(y)) \right).$$

Therefore, there exist (many) $x \in A$ for which (5.9) holds, as was to be shown.

6. Open questions

We collect below a number of open questions and directions suggested by the results in this paper. They are roughly categorized as concerning the families of dynamically syndetic, thick, and piecewise syndetic sets, with the exception of the following.

Question 6.1. To which other group and semigroup actions do the main results in this paper generalize? For example, a generalization of Theorem 3.18 to \mathbb{Z} might read: a subset of \mathbb{Z} is dynamically central syndetic if and only if it is a member of a syndetic, idempotent filter on \mathbb{Z} .

Before attempting to answer Question 6.1, it would be necessary to decide how the family of dynamically central syndetic subsets of \mathbb{Z} is defined. We discuss this at more length in point (vii) at the beginning of Section 3.

6.1. Dynamically syndetic sets

By definition, a subset of \mathbb{N} is dynamically central syndetic if it contains a set of the form R(x, U), where (X, T) is a minimal system, $x \in X$, and $U \subseteq X$ is an open neighborhood of x. It can happen, however, that a set of the form R(x, U) where $x \notin \overline{U}$ is dynamically central syndetic. Recall example (iv) in Section 3.1: the set A of positive integers with even 2-adic valuation is dynamically central syndetic because the point $1_{A\cup\{0\}}$ is uniformly recurrent in the full shift $(\{0,1\}^{\mathbb{N}_0}, \sigma)$ and $A = R_{\sigma}(1_{A\cup\{0\}}, [1]_0)$ where $1_{A\cup\{0\}} \in [1]_0$. Since $\{0,1\}^{\mathbb{N}_0}$ is the disjoint union of the cylinders $[0]_0$ and $[1]_0$, by properties of the 2-adic valuation, we see that

$$A/2 = \mathbb{N} \setminus A = R_{\sigma}(1_{A \cup \{0\}}, [0]_0).$$

By Lemma 3.6, this set is dynamically central syndetic despite the fact that $1_{A\cup\{0\}} \notin [0]_0$. This leads us naturally to the following question.

Question 6.2. Let (X,T) be a minimal system, $x \in X$, and $U \subseteq X$ be nonempty and open. Give necessary and sufficient conditions for x and U so that the set R(x,U) is dynamically central syndetic.

An answer to Question 6.2 would be useful in addressing several of the questions posed below. There is precedent for the family of central sets. Indeed, if $U \subseteq X$ is clopen, then the set R(x, U) is central if and only if U contains a point proximal to x (recall the definition of proximal from Section 2.3.2). It is a fact [32, Lem. 19.22, Thms. 19.23, 19.25] that a pair of points $x, y \in X$ is proximal if and only if there exists an idempotent ultrafilter $p \in \beta \mathbb{N}$ such that $T^p x = y$. If R(x, U) is central, then it belongs to a minimal, idempotent ultrafilter p, and so $T^p x \in \overline{U} = U$. On the other hand, if x is proximal to a point $y \in U$, then there is a minimal, idempotent ultrafilter p such that $T^p x = y$, and so $R(x, U) \in p$. It seems plausible that an analogous answer to Question 6.2 could be given in terms of some kind of proximality.

Our main result, Theorem 3.18, gives that if $A \subseteq \mathbb{N}$ belongs to a syndetic, idempotent filter, then there exists a minimal system (X,T), a point $x \in X$, and an open set $U \subseteq X$ containing x such that $R(x,U) \subseteq A$. The proof of Theorem 3.18, however, does not seem to give any indication as to the dynamical nature of the system (X,T). When, for example, can it be guaranteed that (X,T) is totally minimal, weakly mixing, distal, point distal, a nilsystem, or equicontinuous? Here, for example, is a concrete question along these lines. Recall the definition of a distal point from Section 5.1.

Question 6.3. Does every member of an \mathcal{IP}^* , idempotent filter contain a set of the form R(x, U), where (X, T) is a minimal system, $x \in X$ is a distal point, and $U \subseteq X$ is an open neighborhood of x?

Question 6.3 is motivated by a result of Furstenberg [20, Thm 9.11]: a point $x \in X$ is a distal point if and only if for all open neighborhoods $U \subseteq X$ of x, the set R(x, U) belongs to \mathcal{IP}^* if and only if for all open neighborhoods $U \subseteq X$ of x, the set R(x, U) belongs to \mathcal{C}^* . Note that the family of thickly syndetic sets, \mathcal{PS}^* , is a \mathcal{C}^* , idempotent filter. Thus, an

ostensibly easier variant of Question 6.3 is attained by replacing \mathcal{IP}^* with \mathcal{PS}^* .

While \mathcal{IP} sets are sets of pointwise recurrence for distal points, recall from Lemma 4.9 presented in Section 4 that even dynamically thick sets need not be \mathcal{IP} . A positive answer to Question 6.3 would show, via the following lemma, that a set of pointwise recurrence for distal points must be a translate of an \mathcal{IP} set.

Lemma 6.4. Suppose the answer to Question 6.3 is positive. If $A \subseteq \mathbb{N}$ is a set of pointwise recurrence for all distal points (ie., for all systems (X,T), all distal $x \in X$, and all open $U \subseteq X$ containing x, the set $A \cap R(x,U) \neq \emptyset$), then there exists $n \in \mathbb{N}$ such that $A - n \in \mathcal{IP}$.

Proof. Consider the family

$$\mathscr{F} := \{ A \subseteq \mathbb{N} \mid \exists \ \ell \in \mathbb{N}, \ A - \ell \in \mathcal{IP} \}.$$

Using the fact that the family \mathcal{IP} is partition regular, it is not hard to show that

$$\mathscr{F}^* = \{ B \subseteq \mathbb{N} \mid \forall \ \ell \in \mathbb{N}, \ B \cap (B-1) \cap \dots \cap (B-\ell) \in \mathcal{IP}^* \}.$$

The family \mathscr{F}^* is an \mathcal{IP}^* , translation-invariant (hence, idempotent) filter.

Suppose $A \subseteq \mathbb{N}$ is a set of pointwise recurrence for all distal points. We will use the positive answer to Question 6.3 to show that $A \in \mathscr{F}$ and conclude the proof of the lemma.

To show that $A \in \mathscr{F}$, we will show that $A \in (\mathscr{F}^*)^*$. Let $B \in \mathscr{F}^*$. We must show that $A \cap B \neq \emptyset$. By the positive answer to Question 6.3, the set B contains the times of returns of a distal point to a neighborhood of itself. By assumption, the set $A \cap B \neq \emptyset$, as was to be shown.

It was shown in Lemma 3.4 (iii) that the family dcS is idempotent: $dcS \subseteq dcS + dcS$. Is this an equality, and, if not, how close is it to one?

Question 6.5. Is it the case that

$$dc\mathcal{S} = dc\mathcal{S} + dc\mathcal{S}?$$

Because it is known that $dcS \subseteq dcS + dcS$, it is equivalent to ask: If $A \subseteq \mathbb{N}$ is such that the set

$$A - dc\mathcal{S} := \{ n \in \mathbb{N} \mid A - n \in dc\mathcal{S} \}$$

is dynamically central syndetic, is A dynamically central syndetic?

We were not able to show that a set $A \subseteq \mathbb{N}$ is dynamically central syndetic even under the assumption that every shift A - n, $n \in \mathbb{N}$, is dynamically central syndetic. It may be the case that a set with this property is actually thickly syndetic.

The dual form of Question 6.5 is equally appealing. By Lemma 2.3 (viii), the equivalent, dual form asks: Is $dc\mathcal{T} \subseteq dc\mathcal{T} + dc\mathcal{T}$, that is, is the family of sets of pointwise recurrence idempotent? A positive answer to this question would show that the family $dc\mathcal{PS}$ is idempotent, giving a positive answer to Question 6.11 below.

6.2. Dynamically thick sets

It is simple to see from the definitions that if $A \subseteq \mathbb{N}$ is a set of pointwise recurrence, then for all dynamically central syndetic sets $B \subseteq \mathbb{N}$, there exists $n \in A$ such that the set B - n is dynamically central syndetic. Does the converse hold, that is, does this property characterize being a set of pointwise recurrence?

Question 6.6. Is it true that a set $A \subseteq \mathbb{N}$ is a set of pointwise recurrence if and only if for all $B \in dcS$, there exists $n \in A$ such that $B - n \in dcS$?

If we denote by dcS - dcS the family of sets of the form B - dcS where $B \in dcS$, then Question 6.6 asks whether or not $dcT = (dcS - dcS)^*$. The equivalent, dual form of this equality is dcS = dcS - dcS. Thus, an equivalent form of Question 6.6 asks whether or not a set $A \subseteq \mathbb{N}$ is dynamically central syndetic if and only if there exists $B \in dcS$ such that $B - dcS \subseteq A$. An answer to Question 6.2 would likely be very helpful in addressing this question.

There are several other relatives of Question 6.6 that are of interest.

- (i) Is it true that a set $A \in d\mathcal{T}$ if and only if for all $B \in d\mathcal{S}$, there exists $n \in A$ such that $B n \in dc\mathcal{S}$? Equivalently: is it true that $A \in d\mathcal{S}$ if and only if there exists $B \in d\mathcal{S}$ such that $B dc\mathcal{S} \subseteq A$?
- (ii) Is it true that a set $A \in dc\mathcal{T}$ if and only if for all $B \in \mathcal{C}$, there exists $n \in A$ such that $B n \in \mathcal{C}$? Equivalently: is it true that $A \in dc\mathcal{S}$ if and only if there exists $B \in \mathcal{C}$ such that $B \mathcal{C} \subseteq A$?
- (iii) Is it true that a set $A \in d\mathcal{T}$ if and only if for all $B \in \mathcal{PS}$, there exists $n \in A$ such that $B n \in \mathcal{C}$? Equivalently: is it true that $A \in d\mathcal{S}$ if and only if there exists $B \in \mathcal{PS}$ such that $B \mathcal{C} \subseteq A$?

It is a simple exercise to partition a thick set into a disjoint union of two thick sets. The same task for dynamically thick sets or sets of pointwise recurrence does not appear to be quite so simple.

Question 6.7. Can every dynamically thick set (resp. set of pointwise recurrence) be partitioned into two disjoint dynamically thick sets (resp. sets of pointwise recurrence)?

There is a positive answer to the analogue of Question 6.7 for sets of topological recurrence (Example 4.22), sets of measurable recurrence ([15], Example 4.23), sets of Bohr recurrence [43, Prop. 1.4], and sets of pointwise recurrence for nilsystems [44, Lemma 3.3]. Though not explicitly written in these terms, the proofs of all of these results follow by demonstrating the σ -compactness of the dual families, as defined and described in Section 4.4. Since the family of dynamically central syndetic sets is not σ -compact (Theorem 4.30), a new strategy must be devised to answer Question 6.7.

We showed in Theorem F that all dynamically thick sets take the form $\bigcup_{B \in \mathscr{F}} (B \cap H_B)$, where \mathscr{F} is a robustly syndetic collection of sets (recall Definition 4.17) and $(H_B)_{B \in \mathscr{F}}$ is a collection of thick sets. This result seems of limited use, however, since we were not able to say much about the family \mathscr{F} .

Question 6.8. Is it true that for all dynamically thick sets $A \subseteq \mathbb{N}$, there exists a countable, robustly syndetic collection \mathscr{F} of subsets of \mathbb{N} and, for all $B \in \mathscr{F}$, a thick set $H_B \subseteq \mathbb{N}$, such that

$$A = \bigcup_{B \in \mathscr{F}} \left(B \cap H_B \right)? \tag{6.1}$$

A positive answer to Question 6.8 would strengthen Theorem F and would open the door to applications. As an example, we show in the following lemma how a positive answer could be used to give a positive answer to Question 6.7.

Lemma 6.9. If the answer to Question 6.8 is positive, then every dynamically thick set can be partitioned into two disjoint dynamically thick sets.

Proof. Suppose the answer to Question 6.8 is positive. Let $A \in d\mathcal{T}$, and write the set A in the form given in (6.1). Enumerate \mathscr{F} as $\{B_1, B_2, \ldots\}$. For ease of notation, for each $i \in \mathbb{N}$, write H_i instead of H_{B_i} . By passing to subsets of the H_i 's, we may assume without loss of generality that $H_i = \bigcup_{j=1}^{\infty} I_{i,j}$ where $I_{i,j}$ are finite intervals in \mathbb{N} satisfying $\lim_{j\to\infty} |I_{i,j}| = \infty$ and, for all $i, j \in \mathbb{N}$, max $I_{i,j} < \min I_{i,j+1}$.

It is a simple exercise to choose positive integers

$$n_1^{(1)} < n_2^{(1)} < n_1^{(2)} < n_3^{(1)} < n_2^{(2)} < n_1^{(3)} < n_4^{(1)} < n_3^{(2)} < n_2^{(3)} < n_1^{(4)} < \cdots$$

so that the intervals $I_{i,n_i^{(i)}}, i, j \in \mathbb{N}$, are disjoint. For $i \in \mathbb{N}$, define two thick sets

$$H_i^{(1)} = \bigcup_{j=1}^{\infty} I_{i, n_{2j-1}^{(i)}}$$
 and $H_i^{(2)} = \bigcup_{j=1}^{\infty} I_{i, n_{2j}^{(i)}}$

Let

$$A_1 = \bigcup_{i=1}^{\infty} \left(B_i \cap H_i^{(1)} \right) \quad \text{and} \quad A_2 = \bigcup_{i=1}^{\infty} \left(B_i \cap H_i^{(2)} \right).$$

By construction, the sets A_1 and A_2 are disjoint subsets of A. By appending $A \setminus (A_1 \cup A_2)$ to A_1 , we get that $A = A_1 \cup A_2$. Moreover, both A_1 and A_2 have the form in Theorem F, and so they are dynamically thick.

In the event of a negative answer to Question 6.8, one could weaken the requirement on the collection \mathscr{F} while keeping enough to salvage the reasoning in Lemma 6.9. Such a modification to Question 6.8 could be: Is there is a collection \mathscr{F} with the property that $\{1_B \mid B \in \mathscr{F}\}$ is a σ -compact subset of $\{0,1\}^{\mathbb{N}}$ and, for all $B \in \mathscr{F}$, a thick set $H_B \subseteq \mathbb{N}$ such that the map $\mathscr{F} \to \{0,1\}^{\mathbb{N}}$ given by $B \mapsto H_B$ is continuous such that A has the form in (6.1)?

6.3. Dynamically piecewise syndetic sets

It was shown in Section 5.2.1 that family of dynamically central piecewise syndetic sets sits between the families of central and piecewise syndetic sets:

$$\mathcal{C} \subseteq dc \mathcal{PS} \subseteq \mathcal{PS}$$

Central sets are known to have an abundance of combinatorial configurations [32, Ch. 14]. In Theorem 5.17, we show that every $dc\mathcal{PS}$ set contains "Brauer"-type polynomial configurations $x, y, x + p_1(y), \ldots, x + p_k(y)$. On the other hand, Lemma 4.9 shows that $dc\mathcal{PS}$ sets (in fact, dynamically thick sets) need not be \mathcal{IP} sets. This naturally begs the question: to what extent do $dc\mathcal{PS}$ sets contain finite and infinite combinatorial configurations? The following is a specific example of a question along these lines. Recall the definition of an \mathcal{IP}_0 set from Example 4.21.

Question 6.10. Must every dynamically central piecewise syndetic set be an \mathcal{IP}_0 set?

It follows from Theorem 5.17 that $dc\mathcal{PS}$ sets are \mathcal{IP}_2 sets – they contain configurations of the form $\{x, y, x+y\}$ – but we were not able to iterate in order to find higher-order finite sums. A simpler variant of Question 6.10 is: Must dynamically thick sets contain configurations of the form $\{x, y, z, x+y, x+z, y+z, x+y+z\}$?

A positive answer to Question 6.10 would combine with deep results of Furstenberg and Katznelson to give more context to the result in Theorem 5.15, which says that a general family of times of set returns in ergodic theory is $dc\mathcal{PS}^*$. A corollary of a result of Furstenberg and Katznelson [21, Thm. 10.3] (see the discussion and derivation in [3, Sec. 7.1]) gives that a narrower family of times of set returns is \mathcal{IP}_0^* and hence, by a positive answer to Question 6.10, $dc\mathcal{PS}^*$. Thus, Furstenberg and Katznelson's result would go part of the way toward explaining the result in Theorem 5.15. To recover the full result in Theorem 5.15 via this line of reasoning would require an answer to a long-standing open question: Can the results of Bergelson and McCutcheon (eg. [9, Thm. 7.12]) be upgraded from \mathcal{IP}^* to \mathcal{IP}_0^* ?

Finite sums structure is related to idempotency, which is addressed in following question.

Question 6.11. Is the family of dynamically central piecewise syndetic sets idempotent, that is,

$$dc\mathcal{PS} \subseteq dc\mathcal{PS} + dc\mathcal{PS}?$$

Note that the families C and dcS are idempotent. Were the family dcT idempotent (ie., if we had a positive answer to Question 6.5), then by Lemmas 2.4 and 3.4, we would have a positive answer to Question 6.11.

Combining Theorem 5.3 and Lemma 5.5, we have that

$$dc\mathcal{PS} \subseteq d\mathcal{PS} \subseteq d\mathcal{S} + dc\mathcal{PS} \subseteq d\mathcal{PS} + dc\mathcal{PS}.$$

This is superficially close to demonstrating a positive answer to Question 6.11. In fact, it is natural to guess that perhaps $dc\mathcal{PS} \subseteq dc\mathcal{S} + dc\mathcal{PS}$. Note, however, that Theorem 5.3 relies on the shift-punch machinery in Section 3.3. It would be interesting and potentially useful to find a proof of Theorem 5.3 that avoids this heavy machinery.

References

- J. Auslander. On the proximal relation in topological dynamics. Proc. Amer. Math. Soc., 11:890–895, 1960.
- [2] V. Bergelson. Ultrafilters, IP sets, dynamics, and combinatorial number theory. In Ultrafilters across mathematics, volume 530 of Contemp. Math., pages 23–47. Amer. Math. Soc., Providence, RI, 2010.
- [3] V. Bergelson and D. Glasscock. On the interplay between additive and multiplicative largeness and its combinatorial applications. J. Combin. Theory Ser. A, 172:105203, 60, 2020.

- [4] V. Bergelson and N. Hindman. Nonmetrizable topological dynamics and Ramsey theory. Trans. Amer. Math. Soc., 320(1):293–320, 1990.
- [5] V. Bergelson and N. Hindman. Additive and multiplicative Ramsey theorems in N-some elementary results. Combin. Probab. Comput., 2(3):221–241, 1993.
- [6] V. Bergelson, N. Hindman, and D. Strauss. Strongly central sets and sets of polynomial returns mod 1. Proc. Amer. Math. Soc., 140(8):2671–2686, 2012.
- [7] V. Bergelson, B. Host, and B. Kra. Multiple recurrence and nilsequences. *Invent. Math.*, 160(2):261–303, 2005. With an appendix by Imre Ruzsa.
- [8] V. Bergelson and R. McCutcheon. Recurrence for semigroup actions and a noncommutative Schur theorem. In *Topological dynamics and applications (Minneapolis, MN*, 1995), volume 215 of *Contemp. Math.*, pages 205–222. Amer. Math. Soc., Providence, RI, 1998.
- [9] V. Bergelson and R. McCutcheon. An ergodic IP polynomial Szemerédi theorem. Mem. Amer. Math. Soc., 146(695):viii+106, 2000.
- [10] C. Christopherson and J. H. Johnson, Jr. Algebraic characterizations of some relative notions of size. *Semigroup Forum*, 104(1):28–44, 2022.
- [11] J. de Vries. *Topological dynamical systems*, volume 59 of *De Gruyter Studies in Mathematics*. De Gruyter, Berlin, 2014. An introduction to the dynamics of continuous mappings.
- [12] P. Dong, S. Shao, and X. Ye. Product recurrent properties, disjointness and weak disjointness. Israel J. Math., 188:463–507, 2012.
- [13] T. Downarowicz. Entropy in dynamical systems, volume 18 of New Mathematical Monographs. Cambridge University Press, Cambridge, 2011.
- [14] R. Ellis. Locally compact transformation groups. Duke Math. J., 24:119–125, 1957.
- [15] A. Forrest. Recurrence in dynamical systems: A combinatorial approach. ProQuest LLC, Ann Arbor, MI, 1990. Thesis (Ph.D.)–The Ohio State University.
- [16] N. Frantzikinakis. Multiple correlation sequences and nilsequences. Invent. Math., 202(2):875–892, 2015.
- [17] N. Frantzikinakis, E. Lesigne, and M. Wierdl. Sets of k-recurrence but not (k + 1)-recurrence. Ann. Inst. Fourier (Grenoble), 56(4):839–849, 2006.
- [18] H. Furstenberg. Disjointness in ergodic theory, minimal sets, and a problem in Diophantine approximation. *Math. Systems Theory*, 1:1–49, 1967.
- [19] H. Furstenberg. Ergodic behavior of diagonal measures and a theorem of Szemerédi on arithmetic progressions. J. Analyse Math., 31:204–256, 1977.
- [20] H. Furstenberg. Recurrence in ergodic theory and combinatorial number theory. Princeton University Press, Princeton, N.J., 1981.

- [21] H. Furstenberg and Y. Katznelson. An ergodic Szemerédi theorem for IP-systems and combinatorial theory. J. Analyse Math., 45:117–168, 1985.
- [22] H. Furstenberg and B. Weiss. Topological dynamics and combinatorial number theory. J. Analyse Math., 34:61–85, 1978.
- [23] E. Glasner, W. Huang, S. Shao, B. Weiss, and X. Ye. Topological characteristic factors and nilsystems. arXiv:2006.12385, 2020.
- [24] E. Glasner, T. Tsankov, B. Weiss, and A. Zucker. Bernoulli disjointness. Duke Math. J., 170(4):615–651, 2021.
- [25] S. Glasner and B. Weiss. Interpolation sets for subalgebras of $l^{\infty}(\mathbb{Z})$. Israel J. Math., 44(4):345–360, 1983.
- [26] D. Glasscock, A. Koutsogiannis, and F. K. Richter. Multiplicative combinatorial properties of return time sets in minimal dynamical systems. *Discrete Contin. Dyn. Syst.*, 39(10):5891–5921, 2019.
- [27] B. Green and T. Tao. Linear equations in primes. Ann. of Math., 171(3):1753–1850, 2010.
- [28] B. Green and T. Tao. The Möbius function is strongly orthogonal to nilsequences. Ann. of Math., 175(2):541–566, 2012.
- [29] B. Green and T. Tao. The quantitative behaviour of polynomial orbits on nilmanifolds. Ann. of Math. (2), 175(2):465–540, 2012.
- [30] B. Green, T. Tao, and T. Ziegler. An inverse theorem for the Gowers $U^{s+1}[N]$ -norm. Ann. of Math., 176(2):1231–1372, 2012.
- [31] N. Hindman. Finite sums from sequences within cells of a partition of N. J. Combinatorial Theory Ser. A, 17:1–11, 1974.
- [32] N. Hindman and D. Strauss. Algebra in the Stone-Čech compactification. De Gruyter Textbook. Walter de Gruyter & Co., Berlin, 2012. Theory and applications, Second revised and extended edition.
- [33] B. Host and B. Kra. Nonconventional ergodic averages and nilmanifolds. Ann. of Math. (2), 161(1):397–488, 2005.
- [34] B. Host and B. Kra. Nil-Bohr sets of integers. *Ergodic Theory Dynam. Systems*, 31(1):113–142, 2011.
- [35] B. Host, B. Kra, and A. Maass. Variations on topological recurrence. Monatsh. Math., 179(1):57–89, 2016.
- [36] W. Huang, S. Shao, and X. Ye. An answer to Furstenberg's problem on topological disjointness. *Ergodic Theory and Dynamical Systems*, 40(9):2467–2481, 2020.
- [37] W. Huang and X. Ye. Dynamical systems disjoint from any minimal system. Trans. Amer. Math. Soc., 357(2):669–694, 2005.

- [38] S. Kakeya and S. Morimoto. On a Theorem of MM. Bandet and van der Waerden. Japanese journal of mathematics: transactions and abstracts, 7:163–165, 1930.
- [39] M. Kennedy, S. Raum, and G. Salomon. Amenability, proximality and higher-order syndeticity. *Forum Math. Sigma*, 10:Paper No. e22, 28, 2022.
- [40] S. Kolyada, L. Snoha, and S. Trofimchuk. Noninvertible minimal maps. Fund. Math., 168(2):141–163, 2001.
- [41] A. Koutsogiannis, A. N. Le, J. Moreira, R. Pavlov, and F. K. Richter. Interpolation sets for dynamical systems. arXiv:2401.15339, 2024.
- [42] P. Krautzberger. *Idempotent filters and ultrafilters*. Dissertation, Freie Universität Berlin, 2009.
- [43] A. Le. Interpolation sets and nilsequences. Collog. Math., 162(2):181–199, 2020.
- [44] A. Le. Sublacunary sets and interpolation sets for nilsequences. Discrete Contin. Dyn. Syst., 42(4):1855–1871, 2022.
- [45] A. Leibman. Multiple polynomial correlation sequences and nilsequences. Ergodic Theory Dynam. Systems, 30(3):841–854, 2010.
- [46] R. McCutcheon. Elemental methods in ergodic Ramsey theory, volume 1722 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1999.
- [47] R. Pavlov. Some counterexamples in topological dynamics. Ergodic Theory Dynam. Systems, 28(4):1291–1322, 2008.
- [48] A. Sárközy. On difference sets of sequences of integers. III. Acta Math. Acad. Sci. Hungar., 31(3-4):355–386, 1978.
- [49] E. Szemerédi. On the sets of integers containing no k elements in arithmetic progressions. Acta Arith., 27:299–345, 1975.
- [50] B. van der Waerden. Beweis einer Baudetschen Vermutung. Nieuw Arch. Wiskd., II. Ser., 15:212–216, 1927.
- [51] T. Ward. Ergodic theory: interactions with combinatorics and number theory. In *Ergodic theory*, Encycl. Complex. Syst. Sci., pages 577–596. Springer, New York, 2023.
- [52] X. Ye. D-function of a minimal set and an extension of Sharkovskii's theorem to minimal sets. Ergodic Theory Dynam. Systems, 12(2):365–376, 1992.
- [53] T. Ziegler. Universal characteristic factors and Furstenberg averages. J. Amer. Math. Soc., 20(1):53–97, 2007.

Daniel Glasscock UNIVERSITY OF MASSACHUSETTS LOWELL daniel_glasscock@uml.edu

Anh N. Le UNIVERSITY OF DENVER anh.n.le@du.edu