

A conjecture to prove Riemann hypothesis

F.GHABOUSSI

Department of Physics, University of Konstanz

P.O. Box 5560, D 78434 Konstanz, Germany

E-mail: farhad.ghabousi@uni-konstanz.de

In view of the relations between Riemann hypothesis and quantum theory according to the Hilbert-Polya conjecture and further related investigations its proof is important also for the quantum physics. I conjecture to prove Riemann hypothesis according to Riemann's saddle point method which is used by Riemann to derive the integral representation of zeta function.

In view of the relations between Riemann hypothesis and quantum theory according to the Hilbert-Polya conjecture and further related investigations its proof is important also for quantum physics [1]. I conjecture to prove Riemann hypothesis according to the saddle point method which is used by Riemann to derive the integral representation of zeta function [2], [3], [4].

Riemann hypothesis is based on certain essential mathematical tools which are used by Riemann to construct the integral representation of zeta function especially on the critical line [4] and to express the hypothesis about its non-trivial zeros [3]. I distinguish these tools and show that using them one can prove the Riemann hypothesis.

Riemann achieved the integral representation of zeta function $\zeta(s) := \frac{\Gamma(1-s)}{2\pi i} \int_{+\infty}^{+\infty} \frac{(-x)^{s-1}}{e^x-1} dx$ using several substitutions, e. g. by the substitution $x = e^{i\theta}$ [3] and expressed Riemann hypothesis in view of his saddle point technics to approximate the integral on the critical line $s = \frac{1}{2} + it$ [4] [2], [3] with respect to his substitution of power of primes by the integrals $p^{-s} = s \int_p^\infty x^{-s-1} dx, \dots$ [5]. In other words Riemann expressed Riemann hypothesis about such an integral representation under his impression of the approximation of zeta function by the saddle point method on the critical line. Insofar the saddle point approximation is an inherent part of Riemann hypothesis and may be used to prove it.

For the application of saddle point to prove Riemann hypothesis I use the integral representation of zeta function

$$\zeta(s) := 2 \frac{\Gamma(1-s)}{2\pi i} \int_{+\infty}^{+\infty} \frac{(-x)^{2s-1}}{e^{x^2}-1} dx, \quad (1)$$

which is equivalent to the zeta function $\zeta(s) := \frac{\Gamma(1-s)}{2\pi i} \int_{+\infty}^{+\infty} \frac{(-x)^{s-1}}{e^x-1} dx$ according to $x \rightarrow x^2$ substitution.

The analytical detail of saddle point method in this case can be described as follows. First note that Riemann related prime numbers to the variable x and substitutes by $x = e^{i\theta}$ [3]. Therefore considering $x^2 := xx^* = 1$ it is obvious that the denominator of the integral in (1) is constant.

Considering $(-x)^{2s-1} = (-x)^{(2\sigma-1)+it}$ and the fact that according to saddle point method using a new contour the imaginary part of exponent can be considered as constant [6] the desired derivative to determine the saddle point is the derivative of the real exponent, i. e. $\partial_x (-x)^{2\sigma-1} = (2\sigma-1)(-x)^{2\sigma-2}$ and the saddle point is given for $\sigma = \frac{1}{2}$. In other words the application of saddle point method on (1) requires $\sigma = \frac{1}{2}$. Moreover the zeta function is constant for the saddle point, according to the

approximation of zeta function by the saddle point method, in view of the fact that the Taylor expansion of approximated zeta functions contains only the constant term in view of the vanishing of all derivatives $\partial_x^n (-x)^{2\sigma-1} = (2\sigma-1)\dots(2\sigma-n)(-x)^{2\sigma-(n+1)}$ on the saddle point $\sigma = \frac{1}{2}$. Furthermore note that as we discussed above according to the saddle point method the imaginary part of exponent of numerator of the integral in (1) is considered as constant. Hence according to the fact that one can transform constant functions to zero on certain points by suitable coordinate transformations the constant zeta function on the saddle point $\sigma = \frac{1}{2}$ can be transformed to zero. In this manner the Riemann hypothesis can be proved by the application of saddle point method on zeta function.

I show in the following that the same result can be achieved by the application of Riemann's saddle point method on his mentioned substitution of powers of primes by the integrals of powers of variable x , e. g. $p^{-s} = s \int_p^\infty x^{-s-1} dx$ in his paper [5] which also appear in the integral (1) as $(-x)^{2s-1}$.

Riemann substituted powers of prime numbers p by the integrals of powers of variable x in his paper [5].

$$p^{-s} = s \int_p^\infty x^{-s-1} dx, \quad p^{-2s} = s \int_{p^2}^\infty x^{-s-1} dx, \dots \quad (2)$$

Therefore it is reasonable to consider the result of application of Riemann's saddle point method also on these substitutions.

For this note that Riemann's substitution (2) can be extended to

$$p^{2s} = 2s \int_0^p x^{2s-1} dx = 2s \int_0^p x^{(2\sigma-1)+2it} dx \quad (3)$$

In other words the general substitution of power of primes by powers of x includes (3) and the consequences by the application of saddle point method on (3).

Applying the saddle point method to the integral (3) requires again $\sigma = \frac{1}{2}$ as above and leads to the same result.

It is worth mentioning that one may consider the general substitution by Riemann including (2) and (3) as a primary model of zeta function and the application of saddle point method on (3) as a primary model of application of these methods on zeta function as it is performed above.

Footnotes and references

References

- [1] M.V.Berry, "Riemann's zeta function: A model for quantum chaos", in Quantum Chaos and Statistical Nuclear Physics, T.H. Seligman and H. Nishioka, eds., Lecture Notes in Phys. 263, Springer-Verlag, New York, 1986, pp. 1-17. M.V.Berry, "Quantum chaology" (the Bakerian lecture), Proc. Roy. Soc. Lond. Ser. A, 413 (1987) pp. 183-198. M.V. Berry, "Some quantum-to-classical asymptotics", in Chaos and Quantum Physics, Les Houches Lecture Series 52, M.-J. Giannoni, A. Voros, and J. Zinn-Justin, eds., North-Holland, Amsterdam, 1991, pp.251-304. A. Connes, "Trace formula in noncommutative geometry and the zeros of the Riemann zeta function", Sel. Math. 1999, 5 29106. M. Planat (FEMTO-ST), P. Sol, S. Omar, "Riemann hypothesis and Quantum Mechanics", arXiv:1012.4665 [math-ph].
- [2] C. L. Siegel, 1932, "Ueber Riemanns Nachlass zur analytisches Zahlentheorie", Quellen und Studien zur Geschichte der Mathematik, Astronomie, und Physik, 2, 45-80. Translation in "On Riemanns Nachlass for Analytic Number Theory ...", arXiv:1810.05198 [math.HO].
- [3] H. M. Edwards, "Riemann's Zeta Function", New York: Academic Press, (1974). See especially Chapt. 1. 4. for the background of integral and substitution of x by $e^{i\theta}$ and Chapt. 7. for saddle point method. .
- [4] M. V. Berry, "Riemanns Saddle-point method and the Riemann-Siegel formula", The legacy of Bernhard Riemann after One hundred and fifty years, ALM 35, pp. 6978. For application of saddle point method to derive the integral representation of zeta function see here especially the abstract: "Riemanns way to calculate his zeta function on the critical line was based on an application of his saddle-point technique for approximating integrals".
- [5] See Riemann's work, "On the Number of Prime Numbers less than a Given Quantity", in [3], Appenidx, especially p. 302 for substitutions of powers of primes by integrals of powers of x .
- [6] Fedoryuk, M. V. (2001) [1994], "Saddle point method", Encyclopedia of Mathematics, EMS Press.