

A KHINTCHINE INEQUALITY FOR CENTRAL FOURIER SERIES ON NON-KAC COMPACT QUANTUM GROUPS

SANG-GYUN YOUN

ABSTRACT. The study of Khintchin inequalities has a long history in abstract harmonic analysis. While there is almost no possibility of non-trivial Khintchine inequality for central Fourier series on compact connected semisimple Lie groups, we demonstrate a strong contrast within the framework of compact quantum groups. Specifically, we establish a Khintchine inequality with operator coefficients for arbitrary central Fourier series in a large class of non-Kac compact quantum groups. The main examples include the Drinfeld-Jimbo q -deformations G_q , the free orthogonal quantum groups O_F^+ , and the quantum automorphism group $\mathbb{G}_{aut}(B, \psi)$ with a δ -form ψ .

1. INTRODUCTION

The Khintchine inequality has a long tradition in harmonic analysis, and its natural analogue has been studied extensively in noncommutative harmonic analysis [LP86, LPP91, Pis12]. A noncommutative Khintchine inequality with operator coefficients for the cases $2 \leq p < \infty$ is given by

$$\left\| \sum_{i=1}^m \epsilon_i \otimes A_i \right\|_{L^p(\Omega; S_n^p)} \lesssim \max \left\{ \left\| \left(\sum_{i=1}^m A_i^* A_i \right)^{\frac{1}{2}} \right\|_{S_n^p}, \left\| \left(\sum_{i=1}^m A_i A_i^* \right)^{\frac{1}{2}} \right\|_{S_n^p} \right\} \quad (1.1)$$

where $(\epsilon_i)_{i=1}^m$ is a set of independent and identically distributed Rademacher variables with $\epsilon_i : \Omega \rightarrow \{\pm 1\}$, and S_n^p is the matrix algebra M_n with the Schatten p -norm. When comparing two norms $\|\cdot\|$ and $\|\cdot\|'$, let us denote by $\|f\| \lesssim \|f\|'$ if there exists a constant C such that

$$\|f\| \leq C \|f\|' \quad (1.2)$$

where C is independent of the choice of f . Additionally, we write $\|\cdot\| \approx \|\cdot\|'$ if both $\|\cdot\| \lesssim \|\cdot\|'$ and $\|\cdot\|' \lesssim \|\cdot\|$ hold.

Without using independent random variables, one of the most successful approaches has been the study of *lacunary sets*. This subject has a long and rich history for compact abelian groups [GH13, Kah85, LR75, MP81]. Let us denote by $\text{Irr}(G)$ the set of all irreducible unitary representations $\pi : G \rightarrow \mathcal{U}(n_\pi)$ of a compact group G . In the category of compact abelian groups, all irreducible representations are one-dimensional, i.e. $n_\pi \equiv 1$, and

$\{\pi\}_{\pi \in \text{Irr}(G)}$ forms an orthonormal basis of $L^2(G)$. Moreover, there exists an infinite subset $E \subseteq \text{Irr}(G)$ such that

$$\left\| \sum_{\pi \in E} \pi \otimes A_\pi \right\|_{L^p(G; S_n^p)} \lesssim \max \left\{ \left\| \left(\sum_{\pi \in E} A_\pi^* A_\pi \right)^{\frac{1}{2}} \right\|_{S_n^p}, \left\| \left(\sum_{\pi \in E} A_\pi A_\pi^* \right)^{\frac{1}{2}} \right\|_{S_n^p} \right\} \quad (1.3)$$

by [LR75] and [Har99].

There have been two general frameworks to study such infinite subsets $E \subseteq \text{Irr}(G)$ for non-abelian G , as mentioned in [Pis20] for Sidon sets.

(1) One approach is to replace the compact abelian groups G with the duals $\widehat{\Gamma}$ of discrete groups. This approach has led to various astounding positive results. For example, *freely independent* noncommutative random variables can be used to obtain Khintchine inequalities [Buc99, HP93, PP05, Jun05, RX06]. For example, let $2 \leq p \leq \infty$ and $E = \{g_i\}_{i=1}^m$ be a set of canonical generators of \mathbb{F}_m . Then

$$\left\| \sum_{x \in E} \lambda_x \otimes A_x \right\|_{L^p(\widehat{\mathbb{F}}_m; S_n^p)} \lesssim \max \left\{ \left\| \left(\sum_{x \in E} A_x^* A_x \right)^{\frac{1}{2}} \right\|_{S_n^p}, \left\| \left(\sum_{x \in E} A_x A_x^* \right)^{\frac{1}{2}} \right\|_{S_n^p} \right\}. \quad (1.4)$$

Here, $L^p(\widehat{\mathbb{F}}_m; S_n^p)$ is the noncommutative L^p -space from the tensor product von Neumann algebra $M_n \overline{\otimes} \mathcal{L}(\mathbb{F}_m)$ with the noncommutative measure $\text{tr}_n \otimes \tau$, where tr_n is the usual trace on M_n and $\tau = \langle \cdot, \delta_e \rangle_{\ell^2(\mathbb{F}_m)}$ is the vacuum state on the free group von Neumann algebra $\mathcal{L}(\mathbb{F}_m)$.

(2) The other approach is to replace compact abelian groups with compact non-abelian groups G . In this case, the one-dimensional representations π are replaced by matrix coefficient functions $\sqrt{n_\pi} \pi_{ij}$. However, this line of research encountered some disappointing discoveries for compact non-abelian Lie groups. For example, it was proven in [GT80] that compact connected semisimple Lie groups do not admit an infinite $\Lambda(p)$ -set, even in the scalar-valued case. Specifically, for any $1 < p < \infty$, there is no infinite subset $E \subseteq \text{Irr}(G)$ such that

$$\left\| \sum_{\pi \in E} \sum_{i,j=1}^{n_\pi} a_{ij}^\pi \sqrt{n_\pi} \pi_{ij} \right\|_{L^p(G)} \lesssim \left(\sum_{\pi \in E} \sum_{i,j=1}^{n_\pi} |a_{ij}^\pi|^2 \right)^{\frac{1}{2}}. \quad (1.5)$$

One might hope to explore new possibilities by focusing on the *characters* $\chi_\pi = \sum_{i=1}^{n_\pi} \pi_{ii}$. However, it was shown in [GST82] that for any $3 \leq p < \infty$ and for any compact connected semisimple Lie groups G , there is no infinite

subset $E \subseteq \text{Irr}(G)$ satisfying

$$\left\| \sum_{\pi \in E} a_\pi \chi_\pi \right\|_{L^p(G)} \lesssim \left(\sum_{\pi \in E} |a_\pi|^2 \right)^{\frac{1}{2}}. \quad (1.6)$$

Despite the aforementioned negative conclusions in the category of compact non-abelian Lie groups, a positive result was found for a *twisted quantum group* $\mathbb{G} = SU_q(2)$ with $0 < q < 1$. Specifically, [Wan17] constructed an infinite subset $E \subseteq \text{Irr}(SU_q(2))$ such that

$$\left\| \sum_{\pi \in E} a_\pi \chi_\pi \right\|_{L^4(SU_q(2))} \lesssim \left(\sum_{\pi \in E} |a_\pi|^2 \right)^{\frac{1}{2}}. \quad (1.7)$$

This finding suggests potential for further exploration of the Khintchine inequality using characters on the *Drinfeld-Jimbo q -deformations* $\mathbb{G} = G_q$, and more generally on *non-Kac compact quantum groups*.

The main aim of this paper is to establish a very positive result in a large class of non-Kac compact quantum groups. Let $L^p(\mathbb{G}; S_n^p)$ denote the vector-valued noncommutative L^p -space $S_n^p[L^p(\mathbb{G})]$ [Pis98, Pis03]. In Section 3, we prove the following Khintchine inequality with operator coefficients on general compact quantum groups (Theorem 3.2):

$$\left\| \sum_{\alpha \in \text{Irr}(\mathbb{G})} \chi_\alpha \otimes A_\alpha \right\|_{L^p(\mathbb{G}; S_n^p)} \leq K_p \left\| \left(\sum_{\alpha \in \text{Irr}(\mathbb{G})} A_\alpha^* A_\alpha \right)^{\frac{1}{2}} \right\|_{S_n^p} \quad (1.8)$$

for all $p = 2^k$ with natural numbers k , where the constant K_p is given by

$$K_p = \left(\sum_{\alpha \in \text{Irr}(\mathbb{G})} \|\chi_\alpha\|_{L^\infty(\mathbb{G})}^{2-\frac{4}{p}} \left(\frac{n_\alpha}{d_\alpha} \right)^{\frac{2}{p}} \right)^{\frac{1}{2}} \in [1, \infty]. \quad (1.9)$$

Note that the whole set $E = \text{Irr}(\mathbb{G})$ is taken to cover *all central Fourier series* $f \sim \sum_{\alpha \in \text{Irr}(\mathbb{G})} \chi_\alpha \otimes A_\alpha$ with operator coefficients.

In Section 4, we prove that $K_p < \infty$ for all $2 < p < \infty$ in a wide range of non-Kac compact quantum groups, including the Drinfeld-Jimbo q -deformations G_q , non-Kac free orthogonal quantum groups O_F^+ and non-Kac quantum automorphism groups $\mathbb{G}_{\text{aut}}(B, \psi)$ with a δ -form ψ . In these cases, for any $1 \leq r, s < \infty$, we obtain

$$\|f\|_{L^r(\mathbb{G})} \approx \|f\|_{L^s(\mathbb{G})} \quad (1.10)$$

for arbitrary central Fourier series $f \sim \sum_{\alpha \in \text{Irr}(\mathbb{G})} a_\alpha \chi_\alpha$ with scalar coefficients.

2. PRELIMINARIES

2.1. Compact quantum groups and representations. Within the von Neumann algebraic framework [KV00, Vae02, KV03], a *compact quantum group* \mathbb{G} is given by a triple $(L^\infty(\mathbb{G}), \Delta, h)$ where

- (1) $L^\infty(\mathbb{G})$ is a von Neumann algebra,
- (2) $\Delta : L^\infty(\mathbb{G}) \rightarrow L^\infty(\mathbb{G}) \overline{\otimes} L^\infty(\mathbb{G})$ is a normal unital $*$ -homomorphism satisfying

$$(\Delta \otimes \text{id})\Delta = (\text{id} \otimes \Delta)\Delta, \quad (2.1)$$

- (3) $h : L^\infty(\mathbb{G}) \rightarrow \mathbb{C}$ is a normal faithful state satisfying

$$(\text{id} \otimes h)(\Delta(a)) = h(a)1 = (h \otimes \text{id})(\Delta(a)) \quad (2.2)$$

for all $a \in L^\infty(\mathbb{G})$. We call h the *Haar state* on \mathbb{G} .

A (finite dimensional) *unitary representation* of \mathbb{G} is given by a unitary

$$v = \sum_{i,j=1}^{n_v} e_{ij} \otimes v_{ij} \in M_{n_v} \otimes L^\infty(\mathbb{G}) \text{ such that}$$

$$\Delta(v_{ij}) = \sum_{k=1}^{n_v} v_{ik} \otimes v_{kj} \quad (2.3)$$

for all $1 \leq i, j \leq n_v$. Furthermore, v is called *irreducible* if

$$\{T \in M_{n_v} : (T \otimes 1)v = v(T \otimes 1)\} = \{\lambda \cdot \text{Id}_{n_v} : \lambda \in \mathbb{C}\}. \quad (2.4)$$

Two unitary representations $v, w \in M_n \otimes L^\infty(\mathbb{G})$ are called *unitarily equivalent* if there exists a unitary matrix $U \in M_n$ such that

$$(U \otimes 1)v(U^* \otimes 1) = w, \quad (2.5)$$

and we write $v \cong w$ in this case. We denote by $\text{Irr}(\mathbb{G})$ the set of all irreducible unitary representations up to the unitary equivalence. For each equivalence class $\alpha \in \text{Irr}(\mathbb{G})$, let us take a representative element as

$$u^\alpha = \sum_{i,j=1}^{n_\alpha} e_{ij} \otimes u_{ij}^\alpha \in M_{n_\alpha} \otimes L^\infty(\mathbb{G}). \quad (2.6)$$

Here n_α is called the *classical dimension* of u^α . An important distinction from the case of compact groups is that the *conjugate representation*

$$(u^\alpha)^c = \sum_{i,j=1}^{n_\alpha} e_{ij} \otimes (u_{ij}^\alpha)^* \in M_{n_\alpha} \otimes L^\infty(\mathbb{G}) \quad (2.7)$$

is not necessarily a unitary, but there exists an invertible positive matrix $Q_\alpha \in M_{n_\alpha}$ such that

$$u^{\bar{\alpha}} = (Q_\alpha^{\frac{1}{2}} \otimes 1)(u^\alpha)^c(Q_\alpha^{-\frac{1}{2}} \otimes 1) \quad (2.8)$$

is a unitary representation. Here, we may assume that Q_α is diagonal with a suitable choice of an orthonormal basis. Moreover, the matrix Q_α is uniquely determined with the condition $\text{Tr}(Q_\alpha) = \text{Tr}(Q_\alpha^{-1})$. We write

$$d_\alpha = \text{Tr}(Q_\alpha) = \text{Tr}(Q_\alpha^{-1}), \quad (2.9)$$

and call d_α the *quantum dimension* of u^α . A compact quantum group \mathbb{G} is called *Kac type* if one of the following equivalent conditions holds:

- $Q_\alpha = \text{Id}_{n_\alpha}$ for all $\alpha \in \text{Irr}(\mathbb{G})$.
- $d_\alpha = n_\alpha$ for all $\alpha \in \text{Irr}(\mathbb{G})$.
- the Haar state h is tracial, i.e. $h(ab) = h(ba)$ for all $a, b \in L^\infty(\mathbb{G})$.

Let $v = \sum_{i,j=1}^m e_{ij} \otimes v_{ij} \in M_m \otimes L^\infty(\mathbb{G})$ and $w = \sum_{s,t=1}^n e_{st} \otimes w_{st} \in M_n \otimes L^\infty(\mathbb{G})$ be representations of \mathbb{G} . The *direct sum* is defined by

$$v \oplus w = \begin{bmatrix} v_{11} & \cdots & v_{1m} & & & \\ \vdots & \ddots & \vdots & & & \\ v_{m1} & \cdots & v_{mm} & & & \\ & & & w_{11} & \cdots & w_{1n} \\ & & & \vdots & \ddots & \vdots \\ & & & w_{n1} & \cdots & w_{nn} \end{bmatrix} \quad (2.10)$$

as an element in $M_{m+n} \otimes L^\infty(\mathbb{G})$, and the *tensor product* is defined by

$$v \otimes w = \sum_{i,j=1}^m \sum_{s,t=1}^n e_{ij} \otimes e_{st} \otimes v_{ij} w_{st} \quad (2.11)$$

as an element in $M_m \otimes M_n \otimes L^\infty(\mathbb{G})$. Both the direct sum $v \oplus w$ and the tensor product $v \otimes w$ are representations of \mathbb{G} . For any finite-dimensional unitary representation u of \mathbb{G} , there exist $\gamma_1, \gamma_2, \dots, \gamma_m \in \text{Irr}(\mathbb{G})$ such that

$$u \cong u^{\gamma_1} \oplus u^{\gamma_2} \oplus \cdots \oplus u^{\gamma_m}. \quad (2.12)$$

In this case, we call γ_k a component of irreducible decomposition of u and write $\gamma_k \subseteq u$.

2.2. Schur's orthogonality and Fourier series. The space of polynomials

$$\text{Pol}(\mathbb{G}) = \text{span} \{ u_{ij}^\alpha : \alpha \in \text{Irr}(\mathbb{G}), 1 \leq i, j \leq n_\alpha \} \quad (2.13)$$

is a weak*-dense *-subalgebra of $L^\infty(\mathbb{G})$. The Schur's orthogonality relation states that

$$h\left((u_{ij}^\alpha)^* u_{st}^\beta\right) = \frac{\delta_{\alpha,\beta} \delta_{i,s} \delta_{j,t} (Q_\alpha)_{ii}^{-1}}{d_\alpha} \quad (2.14)$$

$$h\left(u_{ij}^\alpha (u_{st}^\beta)^*\right) = \frac{\delta_{\alpha,\beta} \delta_{i,s} \delta_{j,t} (Q_\alpha)_{jj}}{d_\alpha} \quad (2.15)$$

for all $\alpha, \beta \in \text{Irr}(\mathbb{G})$, $1 \leq i, j \leq n_\alpha$ and $1 \leq s, t \leq n_\beta$. We denote by $L^2(\mathbb{G})$ the completion of $\text{Pol}(\mathbb{G})$ with respect to the inner product

$$\langle f, g \rangle = h(g^* f) \quad (2.16)$$

for all $f, g \in \text{Pol}(\mathbb{G})$. Then (2.14) implies that

$$\left\{ \sqrt{d_\alpha (Q_\alpha)_{ii}} u_{ij}^\alpha : \alpha \in \text{Irr}(\mathbb{G}), 1 \leq i, j \leq n_\alpha \right\} \quad (2.17)$$

is an orthonormal basis of $L^2(\mathbb{G})$. In particular, the set of all *characters* $\{\chi_\alpha\}_{\alpha \in \text{Irr}(\mathbb{G})}$ is an orthonormal subset in $L^2(\mathbb{G})$, where $\chi_\alpha = \sum_{i=1}^{n_\alpha} u_{ii}^\alpha$.

While $h(fg) \neq h(gf)$ on non-Kac \mathbb{G} in general, the (non-trivial) *modular automorphism group* $\sigma = (\sigma_z)_{z \in \mathbb{C}}$ of the Haar state h explains how those quantities are related. The automorphism σ_z is determined by

$$\sigma_z(u_{st}^\alpha) = (Q_\alpha)_{ss}^{iz} (Q_\alpha)_{tt}^{iz} u_{st}^\alpha \quad (2.18)$$

for all $\alpha \in \text{Irr}(\mathbb{G})$ and $1 \leq s, t \leq n_\alpha$, and we have

$$h(fg) = h(g\sigma_{-i}(f)) \quad (2.19)$$

for all $f, g \in \text{Pol}(\mathbb{G})$.

We define the *noncommutative L^1 -space* on \mathbb{G} as the predual $L^1(\mathbb{G}) = L^\infty(\mathbb{G})_*$. The *Fourier coefficient* of a normal bounded linear functional $\omega \in L^1(\mathbb{G})$ at $\alpha \in \text{Irr}(\mathbb{G})$ is given by $\widehat{\omega}(\alpha) \in M_{n_\alpha}$ whose entries are

$$\widehat{\omega}(\alpha)_{ij} = \omega\left((u_{ji}^\alpha)^*\right) \quad (2.20)$$

for all $1 \leq i, j \leq n_\alpha$, and we call

$$\omega \sim \sum_{\alpha \in \text{Irr}(\mathbb{G})} \sum_{i,j=1}^{n_\alpha} d_\alpha (\widehat{\omega}(\alpha) Q_\alpha)_{ij} u_{ji}^\alpha \quad (2.21)$$

the *Fourier series* of $\omega \in L^1(\mathbb{G})$. Note that we have a natural contractive embedding $\iota : L^\infty(\mathbb{G}) \hookrightarrow L^1(\mathbb{G})$ given by

$$\iota(a) = \omega_a(\cdot) = h(\cdot a) \quad (2.22)$$

for all $a \in L^\infty(\mathbb{G})$. When we identify $f \in \text{Pol}(\mathbb{G}) \subseteq L^\infty(\mathbb{G})$ with the associated normal bounded functional $\omega_f = h(\cdot f) \in L^1(\mathbb{G})$, we obtain

$$f = \sum_{\alpha \in \text{Irr}(\mathbb{G})} \sum_{i,j=1}^{n_\alpha} d_\alpha(\widehat{f}(\alpha) Q_\alpha)_{ij} u_{ji}^\alpha, \quad (2.23)$$

where the Fourier coefficient $\widehat{f}(\alpha)$ is defined as $\widehat{\omega_f}(\alpha)$.

2.3. Noncommutative L^p -spaces and the crossed product. For a general description, let $\mathcal{M} \subseteq B(H)$ be a von Neumann algebra with a normal faithful positive linear functional $\phi \in \mathcal{M}_*^+$. See [Ter81, PX03, HJX10] for more details of noncommutative L^p -spaces. Since \mathcal{M} is naturally embedded into \mathcal{M}_* similarly as in (2.22), we have a compatible pair $(\mathcal{M}, \mathcal{M}_*)$ of Banach spaces, and the *noncommutative L^p -space* is defined as the complex interpolation space

$$L^p(\mathcal{M}, \phi) = (\mathcal{M}, \mathcal{M}_*)_{\frac{1}{p}} \quad (2.24)$$

for all $1 \leq p \leq \infty$.

Let $1 \leq p_0, p_1 \leq \infty$ and $\theta \in [0, 1]$ such that $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$. Then

$$\|f\|_{L^p(\mathcal{M}, \phi)} \leq \|f\|_{L^{p_0}(\mathcal{M}, \phi)}^{1-\theta} \|f\|_{L^{p_1}(\mathcal{M}, \phi)}^\theta \quad (2.25)$$

for all $f \in \mathcal{M}$.

Although the normal faithful positive functional $\phi \in \mathcal{M}_*^+$ is non-tracial in general, the elements in $L^p(\mathcal{M}, \phi)$ can be realized concretely in a larger noncommutative *tracial* measure space. Using the modular automorphism $\sigma^\phi = (\sigma_t^\phi)_{t \in \mathbb{R}}$, the *crossed product*

$$\mathcal{R} = \mathcal{M} \rtimes_{\sigma^\phi} \mathbb{R} \subseteq B(L^2(\mathbb{R}, H)) \quad (2.26)$$

is defined as the von Neumann algebra generated by two families of operators, $\{\pi(x)\}_{x \in \mathcal{M}}$ and $\{\lambda_s\}_{s \in \mathbb{R}}$ given by

$$(\pi(x)\xi)(t) = \sigma_{-t}^\phi(x)\xi(t) \quad (2.27)$$

$$(\lambda_s\xi)(t) = \xi(t-s) \quad (2.28)$$

for all $x \in \mathcal{M}$ and $s, t \in \mathbb{R}$. Here, $\pi : \mathcal{M} \hookrightarrow \mathcal{R}$ is a normal faithful $*$ -representation, so we can identify \mathcal{M} with $\pi(\mathcal{M})$.

By the Stone's theorem, there exists a densely defined injective positive operator $D = e^A : \mathcal{D}(A) \rightarrow L^2(\mathbb{R}, H)$ such that $\lambda(t) = D^{it}$ for all $t \in \mathbb{R}$, where $\mathcal{D}(A)$ is the set of all $\xi \in L^2(\mathbb{R}, H)$ such that

$$A\xi = \lim_{s \rightarrow 0} \frac{1}{is} (\lambda_s \xi - \xi) \quad (2.29)$$

exists in $L^2(\mathbb{R}, H)$.

For any compactly supported weak*-continuous function $f : \mathbb{R} \rightarrow \mathcal{M}$, let us consider the following operator

$$x_f = \int_{\mathbb{R}} \pi(f(s)) \lambda_s ds. \quad (2.30)$$

The set of all such operators x_f forms a weak*-dense *-subalgebra of \mathcal{R} .

For each normal semifinite weight $\varphi : \mathcal{M}_+ \rightarrow [0, \infty]$, there exists the dual normal semifinite weight $\widehat{\varphi} : \mathcal{R}_+ \rightarrow [0, \infty]$ determined by

$$\widehat{\varphi}(x_f) = \widehat{\varphi} \left(\int_{\mathbb{R}} \pi(f(s)) \lambda_s ds \right) = \varphi(f(0)) \quad (2.31)$$

for any positive x_f . In addition, the dual weight $\widehat{\varphi}$ is faithful if the given weight φ is faithful.

Although $\widehat{\phi}$ of the given faithful $\phi \in \mathcal{M}_*^+$ is non-tracial in general, the following weight

$$\tau(\cdot) = \widehat{\phi}(D^{-\frac{1}{2}} \cdot D^{-\frac{1}{2}}) \quad (2.32)$$

is the unique normal semifinite faithful tracial weight on \mathcal{R} satisfying

$$\sigma_t^{\widehat{\phi}}(x) = \lambda_t \sigma_t^\tau(x) \lambda_t^{-1} = D^{it} x D^{-it}, \quad t \in \mathbb{R}. \quad (2.33)$$

Let $L^0(\mathcal{R}, \tau)$ be the space of all closed densely defined operators T on $L^2(\mathbb{R}, H)$ satisfying the following conditions:

- (1) T is affiliated with \mathcal{R} , i.e. $TV = VT$ for all unitaries V in the commutant algebra \mathcal{R}' .
- (2) T is τ -measurable, i.e. for any $\delta > 0$ there exists a projection $p \in \mathcal{R}$ such that $\text{ran}(p) \subseteq \mathcal{D}(T)$ and $\tau(1 - p) \leq \delta$.

For any normal positive functional $\omega \in \mathcal{M}_*^+$, there exists a unique positive $D_\omega \in L^0(\mathcal{R}, \tau)$ such that

$$\widehat{\omega}(x) = \tau(D_\omega^{\frac{1}{2}} x D_\omega^{\frac{1}{2}}) = \tau(x D_\omega) \quad (2.34)$$

for all $x \in \mathcal{R}_+$, and the modular automorphism of $\widehat{\omega}$ is given by $\sigma_t^{\widehat{\omega}}(x) = D_\omega^{it} x D_\omega^{-it}$. In particular, we have $D_\phi = D = e^A$ for the given normal faithful positive functional $\phi \in \mathcal{M}_*^+$. See Lemma 2.2 and Corollary 2.6 of [Ter81] for more details.

2.4. Dual action and Haagerup L^p -spaces. To define the Haagerup L^p -spaces $L^{p,H}(\mathcal{M}, \phi)$, let us introduce the dual action $\widehat{\sigma}^\phi$ of $\widehat{\mathbb{R}} \cong \mathbb{R}$ on \mathcal{R} using a canonical identification of the dual group $\widehat{\mathbb{R}}$ with \mathbb{R} . Specifically, the dual action $\widehat{\sigma}^\phi$ is given by

$$\widehat{\sigma}_s^\phi(x) = w_s x w_s^* \quad (2.35)$$

for all $s \in \mathbb{R}$ and $x \in \mathcal{R}$, where the unitary representation $w : \widehat{\mathbb{R}} \cong \mathbb{R} \rightarrow \mathcal{U}(L^2(\mathbb{R}, H))$ is given by

$$(w_s \xi)(t) = e^{-ist} \xi(t) \quad (2.36)$$

for all $\xi \in L^2(\mathbb{R}, H)$. In particular, we have

$$\widehat{\sigma}_s^\phi(\pi(a)\lambda_t) = e^{-ist} \pi(a)\lambda_t \quad (2.37)$$

for all $a \in \mathcal{M}$ and $s, t \in \mathbb{R}$. The subalgebra $\pi(\mathcal{M})$ is characterized as the fixed points for the dual action inside \mathcal{R} , i.e.

$$\pi(\mathcal{M}) = \{x \in \mathcal{R} : \widehat{\sigma}_s^\phi(x) = x \forall s \in \mathbb{R}\}. \quad (2.38)$$

The Haagerup L^p -space is defined as

$$L^{p,H}(\mathcal{M}, \phi) = \left\{ x \in L^0(\mathcal{R}, \tau) : \widehat{\sigma}_s^\phi(x) = e^{-\frac{s}{p}} x \forall s \in \mathbb{R} \right\} \quad (2.39)$$

for any $1 \leq p \leq \infty$. In particular, we have $L^{\infty,H}(\mathcal{M}, \phi) = \pi(\mathcal{M})$ for $p = \infty$, and

$$L^{1,H}(\mathcal{M}, \phi) = \text{span} \{ D_\omega \in L^0(\mathcal{R}, \tau) : \omega \in \mathcal{M}_*^+ \} \quad (2.40)$$

for $p = 1$ where D_ω is the unique positive τ -measurable operator associated to $\omega \in \mathcal{M}_*^+$ discussed in Subsection 2.3. Here, (2.40) is thanks to the following identities $\widehat{\omega} = \widehat{\omega} \circ \widehat{\sigma}_t^\phi$ and $e^{-t}\tau = \tau \circ \widehat{\sigma}_t^\phi$.

Note that $\text{Tr}(D_\omega) = \omega(1)$ extends to a contractive positive normal linear functional on $L^{1,H}(\mathcal{M}, \psi)$, which we call *trace*. If $x \in L^{p,H}(\mathcal{M}, \phi)$ and $y \in L^{q,H}(\mathcal{M}, \phi)$ such that $\frac{1}{p} + \frac{1}{q} = 1$ with $p, q \geq 1$, then $xy, yx \in L^{1,H}(\mathcal{M}, \phi)$ and we have

$$\text{Tr}(xy) = \text{Tr}(yx). \quad (2.41)$$

Now we can explain how the complex interpolation space $L^p(\mathcal{M}, \phi)$ is identified with the Haagerup L^p -space $L^{p,H}(\mathcal{M}, \phi)$. Let $x = u|x|$ be the polar decomposition of $x \in L^0(\mathcal{R}, \tau)$. Then $x \in L^{p,H}(\mathcal{M}, \phi)$ is equivalent to that $u \in \mathcal{M}$ and $|x|^p \in L^{1,H}(\mathcal{M}, \phi)$. In this case, the L^p -norm of $x \in L^{p,H}(\mathcal{M}, \phi)$ is defined by

$$\|x\|_{p,H} = \text{Tr}(|x|^p)^{\frac{1}{p}}. \quad (2.42)$$

By [Kos84, Section 9], a linear map $\iota_p : \mathcal{M} \hookrightarrow L^{p,H}(\mathcal{M}, \phi)$ given by

$$\iota_p(x) = \pi(x)D^{\frac{1}{p}} \quad (2.43)$$

extends to an onto isometry $\iota_p : L^p(\mathcal{M}, \phi) \rightarrow L^{p,H}(\mathcal{M}, \phi)$ for all $1 \leq p \leq \infty$.

As in the tracial setting, we have a natural Hölder inequality on the Haagerup L^p -spaces. For any $x \in L^{p,H}(\mathcal{M}, \phi)$ and $y \in L^{q,H}(\mathcal{M}, \phi)$ such that $\frac{1}{p} + \frac{1}{q} = \frac{1}{r}$ with $r \geq 1$, we have $xy \in L^{r,H}(\mathcal{M}, \phi)$ and

$$\|xy\|_{L^{r,H}(\mathcal{M}, \phi)} \leq \|x\|_{L^{p,H}(\mathcal{M}, \phi)} \cdot \|y\|_{L^{q,H}(\mathcal{M}, \phi)}. \quad (2.44)$$

We define the noncommutative vector-valued L^1 -space as $S_n^1[\mathcal{M}_*] = S_n^1 \widehat{\otimes} \mathcal{M}_* \cong (M_n \otimes_{\min} \mathcal{M})_*$ where $\widehat{\otimes}$ and \otimes_{\min} are the projective and the injective tensor product in the category of operator spaces respectively. For the general cases $1 \leq p \leq \infty$, the noncommutative vector-valued L^p -space is realized as the complex interpolation space

$$S_n^p[L^p[\mathcal{M}, \phi]] = (M_n \otimes_{\min} \mathcal{M}, S_n^1 \widehat{\otimes} \mathcal{M}_*)_{\frac{1}{p}} = L^p(\mathcal{M}_n, \phi_n) \quad (2.45)$$

by [Pis98, Corollary 1.4], where $\mathcal{M}_n = M_n \overline{\otimes} \mathcal{M}$, $\phi_n = \text{tr}_n \otimes \phi$ and tr_n is the usual trace on M_n . Thus, we obtain

$$\|f\|_{S_n^p[L^p[\mathcal{M}, \phi]]} = \left[(\text{tr}_n \otimes \text{Tr}) \left(\left| (\text{id} \otimes \pi)(f)(\text{Id}_n \otimes D^{\frac{1}{p}}) \right|^p \right) \right]^{\frac{1}{p}} \quad (2.46)$$

for any $f \in M_n \otimes_{\min} \mathcal{M}$ and $1 \leq p < \infty$. See [Pis98, Pis03] for more details of the noncommutative vector-valued L^p -spaces.

3. A KHINTCHINE INEQUALITY FOR CENTRAL FOURIER SERIES

We will focus on the so-called *central Fourier series with operator coefficients*, which is of the form

$$f \sim \sum_{\alpha \in \text{Irr}(\mathbb{G})} x_\alpha \otimes \chi_\alpha \quad (3.1)$$

with an arbitrary sequence $(x_\alpha)_{\alpha \in \text{Irr}(\mathbb{G})} \subseteq M_n$. A key observation of this paper is the following smoothing effect of the automorphism $\sigma_{-\frac{i}{2p}}$ on the L^p -norm of the characters χ_α .

Lemma 3.1. *For any $p = 2^k$ with a natural number k , we have*

$$\left\| \sigma_{-\frac{i}{2p}}(\chi_\alpha) \right\|_{L^p(\mathbb{G})} \leq \|\chi_\alpha\|_{L^\infty(\mathbb{G})}^{1-\frac{2}{p}} \left(\frac{n_\alpha}{d_\alpha} \right)^{\frac{1}{p}}. \quad (3.2)$$

Proof. For the first step $k = 1$ ($\Leftrightarrow p = 2$), we have

$$\left\| \sigma_{-\frac{i}{4}}(\chi_\alpha) \right\|_{L^2(\mathbb{G})}^2 = \left\| \sum_{j=1}^{n_\alpha} (Q_\alpha)_{jj}^{\frac{1}{2}} u_{jj}^\alpha \right\|_{L^2(\mathbb{G})}^2 \quad (3.3)$$

$$= \sum_{j=1}^{n_\alpha} (Q_\alpha)_{jj} \frac{(Q_\alpha)_{jj}^{-1}}{d_\alpha} = \frac{n_\alpha}{d_\alpha} = \|\chi_\alpha\|_\infty^0 \cdot \left(\frac{n_\alpha}{d_\alpha} \right)^1. \quad (3.4)$$

Now let us assume that (3.2) holds for $p = 2^k$. Then we have

$$\left\| \sigma_{-\frac{i}{4p}}(\chi_\alpha) \right\|_{L^{2p}(\mathbb{G})}^{2p} = \left\| \pi \left(\sigma_{-\frac{i}{4p}}(\chi_\alpha) \right) D^{\frac{1}{2p}} \right\|_{L^{2p,H}(\mathbb{G})}^{2p} \quad (3.5)$$

$$= \text{Tr} \left(\left| D^{\frac{1}{4p}} \pi(\chi_\alpha) D^{-\frac{1}{4p}} D^{\frac{1}{2p}} \right|^{2p} \right) = \text{Tr} \left(\left| D^{\frac{1}{4p}} \pi(\chi_\alpha) D^{\frac{1}{4p}} \right|^{2p} \right) \quad (3.6)$$

$$= \text{Tr} \left(\left(D^{\frac{1}{4p}} \pi(\chi_\alpha)^* D^{\frac{1}{2p}} \pi(\chi_\alpha) D^{\frac{1}{4p}} \right)^p \right) \quad (3.7)$$

$$= \text{Tr} \left(D^{\frac{1}{4p}} \left(\pi(\chi_\alpha)^* D^{\frac{1}{2p}} \pi(\chi_\alpha) D^{\frac{1}{2p}} \right)^{p-1} \pi(\chi_\alpha)^* D^{\frac{1}{2p}} \pi(\chi_\alpha) D^{\frac{1}{4p}} \right) \quad (3.8)$$

$$= \text{Tr} \left(\left(\pi(\chi_\alpha)^* D^{\frac{1}{2p}} \pi(\chi_\alpha) D^{\frac{1}{2p}} \right)^p \right). \quad (3.9)$$

The last equality is thanks to the traciality (2.41). Then, applying the non-commutative Hölder inequality (2.44), we obtain

$$\left\| \sigma_{-\frac{i}{4p}}(\chi_\alpha) \right\|_{L^{2p}(\mathbb{G})}^2 = \left[\text{Tr} \left(\left(\pi(\chi_\alpha)^* D^{\frac{1}{2p}} \pi(\chi_\alpha) D^{\frac{1}{2p}} \right)^p \right) \right]^{\frac{1}{p}} \quad (3.10)$$

$$\leq \left\| \pi(\chi_\alpha)^* \right\|_{L^\infty, H(\mathbb{G})} \cdot \left[\text{Tr} \left(\left| D^{\frac{1}{2p}} \pi(\chi_\alpha) D^{\frac{1}{2p}} \right|^p \right) \right]^{\frac{1}{p}} \quad (3.11)$$

$$= \left\| \pi(\chi_\alpha^*) \right\|_{L^\infty, H(\mathbb{G})} \cdot \left[\text{Tr} \left(\left| D^{\frac{1}{2p}} \pi(\chi_\alpha) D^{-\frac{1}{2p}} D^{\frac{1}{p}} \right|^p \right) \right]^{\frac{1}{p}} \quad (3.12)$$

$$= \left\| \chi_\alpha \right\|_{L^\infty(\mathbb{G})} \cdot \left\| \pi \left(\sigma_{-\frac{i}{2p}}(\chi_\alpha) \right) D^{\frac{1}{p}} \right\|_{L^p, H(\mathbb{G})} \quad (3.13)$$

$$= \left\| \chi_\alpha \right\|_{L^\infty(\mathbb{G})} \cdot \left\| \sigma_{-\frac{i}{2p}}(\chi_\alpha) \right\|_{L^p(\mathbb{G})}. \quad (3.14)$$

Thus, we can conclude that

$$\left\| \sigma_{-\frac{i}{4p}}(\chi_\alpha) \right\|_{L^{2p}(\mathbb{G})} \leq \left\| \chi_\alpha \right\|_{L^\infty(\mathbb{G})}^{\frac{1}{2}} \cdot \left\| \sigma_{-\frac{i}{2p}}(\chi_\alpha) \right\|_{L^p(\mathbb{G})}^{\frac{1}{2}} \quad (3.15)$$

$$\leq \left\| \chi_\alpha \right\|_{L^\infty(\mathbb{G})}^{\frac{1}{2}} \cdot \left\| \chi_\alpha \right\|_{L^\infty(\mathbb{G})}^{\frac{1}{2} - \frac{1}{p}} \left(\frac{n_\alpha}{d_\alpha} \right)^{\frac{1}{2p}} = \left\| \chi_\alpha \right\|_{L^\infty(\mathbb{G})}^{1 - \frac{1}{p}} \left(\frac{n_\alpha}{d_\alpha} \right)^{\frac{1}{2p}} \quad (3.16)$$

where the second inequality comes from the inductive assumption. \square

Now, let us exhibit the main result, an operator-valued Khintchine inequality for an arbitrary central Fourier series with the following constant

$$K_p = \left(\sum_{\alpha \in \text{Irr}(\mathbb{G})} \left\| \chi_\alpha \right\|_{L^\infty(\mathbb{G})}^{2 - \frac{4}{p}} \left(\frac{n_\alpha}{d_\alpha} \right)^{\frac{2}{p}} \right)^{\frac{1}{2}} \in [1, \infty]. \quad (3.17)$$

Theorem 3.2. *Let $p = 2^k$ with a natural number k . Then*

$$\|f\|_{L^p(\mathbb{G}; S_n^p)} \leq K_p \cdot \left\| \left(\sum_{\alpha \in \text{Irr}(\mathbb{G})} x_\alpha^* x_\alpha \right)^{\frac{1}{2}} \right\|_{S_n^p} \quad (3.18)$$

for any central Fourier series $f = \sum_{\alpha \in \text{Irr}(\mathbb{G})} x_\alpha \otimes \chi_\alpha \in M_n \otimes \text{Pol}(\mathbb{G})$ with operator coefficients.

Proof. The case $k = 1$ ($\Leftrightarrow p = 2$) is immediate, so let us suppose that $k \geq 2$. Recall that the vector-valued L^p -norm of $f = \sum_{\alpha \in \text{Irr}(\mathbb{G})} x_\alpha \otimes \chi_\alpha \in M_n \otimes \text{Pol}(\mathbb{G})$ is given by

$$\|f\|_{L^p(\mathbb{G}; S_n^p)} = \left[(\text{tr}_n \otimes \text{Tr}) \left(\left| (\text{id} \otimes \pi)(f)(\text{Id}_n \otimes D^{\frac{1}{p}}) \right|^p \right) \right]^{\frac{1}{p}} \quad (3.19)$$

and note that

$$\left| (\text{id} \otimes \pi)(f)(\text{Id}_n \otimes D^{\frac{1}{p}}) \right|^p \quad (3.20)$$

$$= \left((\text{Id}_n \otimes D^{\frac{1}{p}})(\text{id} \otimes \pi)(f)^*(\text{id} \otimes \pi)(f)(\text{Id}_n \otimes D^{\frac{1}{p}}) \right)^{\frac{p}{2}} \quad (3.21)$$

$$= \left((\text{Id}_n \otimes D^{\frac{1}{p}})(\text{id} \otimes \pi)(f^* f)(\text{Id}_n \otimes D^{-\frac{1}{p}})(\text{Id}_n \otimes D^{\frac{2}{p}}) \right)^{\frac{p}{2}} \quad (3.22)$$

$$= \left((\text{id} \otimes \pi) \left((\text{id} \otimes \sigma_{-\frac{i}{p}})(f^* f) \right) (\text{Id}_n \otimes D^{\frac{2}{p}}) \right)^{\frac{p}{2}}. \quad (3.23)$$

Thus, we have

$$\|f\|_{L^p(\mathbb{G}; S_n^p)}^2 = \left\| (\text{id} \otimes \sigma_{-\frac{i}{p}})(f^* f) \right\|_{S_n^{\frac{p}{2}}[L^{\frac{p}{2}}(\mathbb{G})]} \quad (3.24)$$

Since $\text{span} \{\chi_\alpha\}_{\alpha \in \text{Irr}(\mathbb{G})}$ is a $*$ -subalgebra of $L^\infty(\mathbb{G})$, the element $f^* f = \sum_{\alpha_1, \alpha_2 \in \text{Irr}(\mathbb{G})} x_{\alpha_1}^* x_{\alpha_2} \otimes \chi_{\alpha_1}^* \chi_{\alpha_2}$ can be written as a linear combination of characters $\sum_{\beta \in \text{Irr}(\mathbb{G})} y_\beta \otimes \chi_\beta$, where the operator coefficients are determined by

$$y_\beta = (\text{id} \otimes h) \left((\text{Id}_n \otimes \chi_\beta^*) \cdot f^* f \right) \quad (3.25)$$

$$= \sum_{\alpha_1 \in \text{Irr}(\mathbb{G})} x_{\alpha_1}^* \sum_{\alpha_2 \in \text{Irr}(\mathbb{G})} x_{\alpha_2} \cdot h(\chi_\beta^* \chi_{\alpha_1}^* \chi_{\alpha_2}) \quad (3.26)$$

$$= \sum_{\alpha_1 \in \text{Irr}(\mathbb{G})} x_{\alpha_1}^* \sum_{\alpha_2 \in \text{Irr}(\mathbb{G})} x_{\alpha_2} \cdot h(\chi_{\alpha_1} \chi_\beta \chi_{\alpha_2}^*) = T^* (\text{Id}_n \otimes \chi_\beta) T \quad (3.27)$$

where $T = \sum_{\gamma \in \text{Irr}(\mathbb{G})} x_\gamma \otimes |\chi_\gamma^*\rangle$ with a linear map $|\chi_\gamma^*\rangle : \mathbb{C} \rightarrow L^2(\mathbb{G})$ mapping z to $z \cdot \chi_\gamma^*$. Then we have

$$\|y_\beta\|_{S_n^{\frac{p}{2}}} \leq \|T^*\|_{S^p} \cdot \|\text{Id}_n \otimes \chi_\beta\|_{M_n \overline{\otimes} L^\infty(\mathbb{G})} \cdot \|T\|_{S^p} \quad (3.28)$$

$$= \|\chi_\beta\|_{L^\infty(\mathbb{G})} \cdot \left\| \left(\sum_{\gamma \in \text{Irr}(\mathbb{G})} x_\gamma^* x_\gamma \right)^{\frac{1}{2}} \right\|_{S_n^p}^2. \quad (3.29)$$

Combining the above observations with the triangle inequality and Lemma 3.1, we reach the following conclusion

$$\left\| (\text{id} \otimes \sigma_{-\frac{i}{p}})(f^* f) \right\|_{S_n^{\frac{p}{2}}[L^{\frac{p}{2}}(\mathbb{G})]} \quad (3.30)$$

$$\leq \sum_{\beta \in \text{Irr}(\mathbb{G})} \left\| y_\beta \otimes \sigma_{-\frac{i}{p}}(\chi_\beta) \right\|_{S_n^{\frac{p}{2}}[L^{\frac{p}{2}}(\mathbb{G})]} \quad (3.31)$$

$$\leq \sum_{\beta \in \text{Irr}(\mathbb{G})} \|\chi_\beta\|_{L^\infty(\mathbb{G})} \left\| \left(\sum_{\gamma \in \text{Irr}(\mathbb{G})} x_\gamma^* x_\gamma \right)^{\frac{1}{2}} \right\|_{S_n^p}^2 \cdot \|\chi_\beta\|_{L^\infty(\mathbb{G})}^{1-\frac{4}{p}} \left(\frac{n_\beta}{d_\beta} \right)^{\frac{2}{p}}. \quad (3.32)$$

□

Remark 3.3. *Theorem 3.2 can be rephrased as that $E = \text{Irr}(\mathbb{G})$ is a completely bounded $\Lambda(p)$ -set under the assumption $K_p < \infty$ in a natural sense of [Har99] within the framework of compact quantum groups.*

Then, some standard complex interpolation arguments allow us to show that all L^p -norms of the central Fourier series are equivalent in the scalar-valued case.

Corollary 3.4. *Suppose that $K_p < \infty$ for infinitely many $p = 2^k$ with natural numbers k . Then, for any $1 \leq r, s < \infty$, we have*

$$\|f\|_{L^r(\mathbb{G})} \approx \|f\|_{L^s(\mathbb{G})} \quad (3.33)$$

for arbitrary central Fourier series $f = \sum_{\alpha \in \text{Irr}(\mathbb{G})} a_\alpha \chi_\alpha \in \text{Pol}(\mathbb{G})$.

Proof. It is enough to prove that, for any $1 \leq r < \infty$, the following inequality

$$\|f\|_{L^r(\mathbb{G})} \lesssim \|f\|_{L^1(\mathbb{G})} \quad (3.34)$$

holds for all central Fourier series $f = \sum_{\alpha \in \text{Irr}(\mathbb{G})} a_\alpha \chi_\alpha \in \text{Pol}(\mathbb{G})$.

First of all, suppose that $K_p < \infty$ for $p = 2^k$ with a natural number $k \geq 2$. Then there exists $\theta \in (0, 1)$ such that $\frac{1}{2} = \frac{1-\theta}{1} + \frac{\theta}{p}$ ($\Leftrightarrow \theta = \frac{p}{2(p-1)}$),

and we have

$$\|f\|_{L^2(\mathbb{G})} \leq \|f\|_{L^1(\mathbb{G})}^{1-\theta} \|f\|_{L^p(\mathbb{G})}^\theta \quad (3.35)$$

$$\leq \|f\|_{L^1(\mathbb{G})}^{1-\theta} \cdot K_p^\theta \|f\|_{L^2(\mathbb{G})}^\theta \quad (3.36)$$

by (2.25) and Theorem 3.2. This implies

$$\|f\|_{L^2(\mathbb{G})} \leq K_p^{\frac{\theta}{1-\theta}} \|f\|_{L^1(\mathbb{G})} = K_p^{\frac{p}{p-2}} \|f\|_{L^1(\mathbb{G})}. \quad (3.37)$$

Combining (3.37) with Theorem 3.2, we can conclude that

$$\|f\|_{L^p(\mathbb{G})} \leq K_p \|f\|_{L^2(\mathbb{G})} \leq K_p^{\frac{2p-2}{p-2}} \|f\|_{L^1(\mathbb{G})}. \quad (3.38)$$

For the general cases $1 \leq r < \infty$, there exists $p = 2^k$ with a natural number $k \geq 2$ such that $r < 2^k$ and $K_p < \infty$. Let us write $\frac{1}{r} = \frac{1-\theta'}{1} + \frac{\theta'}{p}$ ($\Leftrightarrow \theta' = \frac{p(r-1)}{r(p-1)}$) with $\theta' \in [0, 1)$. Then, by (2.25) and (3.38), we obtain

$$\|f\|_{L^r(\mathbb{G})} \leq \|f\|_{L^1(\mathbb{G})}^{1-\theta'} \|f\|_{L^p(\mathbb{G})}^{\theta'} = K_p^{\frac{2p(r-1)}{r(p-2)}} \|f\|_{L^1(\mathbb{G})}. \quad (3.39)$$

□

4. MAIN EXAMPLES

In this Section, we demonstrate that the Khintchine inequality with operator coefficients holds for arbitrary central Fourier series in a large class of non-Kac compact quantum groups. More precisely, we will prove that

$$K_p^2 = \sum_{\alpha \in \text{Irr}(\mathbb{G})} \|\chi_\alpha\|_{L^\infty(\mathbb{G})}^{2-\frac{4}{p}} \left(\frac{n_\alpha}{d_\alpha} \right)^{\frac{2}{p}} < \infty \quad (4.1)$$

holds for all $p > 2$ when \mathbb{G} is one of the following compact quantum groups

- Drinfeld-Jimbo q -deformations G_q with $0 < q < 1$,
- Non-Kac free orthogonal quantum groups O_F^+ ,
- Non-Kac quantum automorphism group $\mathbb{G}_{\text{aut}}(B, \psi)$ with a δ -form.

Our unified strategy to cover all the aforementioned examples is as follows. In Subsections 4.1, 4.2 and 4.3, we will prove that there exists $r \in (0, 1)$ such that

$$\frac{n_\alpha}{d_\alpha} \lesssim r^{|\alpha|} \quad (4.2)$$

where $|\cdot| : \text{Irr}(\mathbb{G}) \rightarrow \mathbb{N}$ is a natural length function for each \mathbb{G} in the above list. Assuming the existence of such $r \in (0, 1)$, we can obtain

$$K_p^2 = \sum_{k=0}^{\infty} \sum_{\alpha \in \text{Irr}(\mathbb{G}): |\alpha|=k} \|\chi_\alpha\|_{L^\infty(\mathbb{G})}^{2-\frac{4}{p}} \left(\frac{n_\alpha}{d_\alpha}\right)^{\frac{2}{p}} \quad (4.3)$$

$$\lesssim \sum_{k=0}^{\infty} \left(\sum_{\alpha \in \text{Irr}(\mathbb{G}): |\alpha|=k} \|\chi_\alpha\|_{L^\infty(\mathbb{G})}^2 \right) r^{\frac{2k}{p}}, \quad (4.4)$$

so it is enough for us to see that $\sum_{\alpha \in \text{Irr}(\mathbb{G}): |\alpha|=k} \|\chi_\alpha\|_{L^\infty(\mathbb{G})}^2$ grows polynomially

when k increases.

(1) For the Drinfeld-Jimbo q -deformations $\mathbb{G} = G_q$, we have

$$\sum_{\alpha \in \text{Irr}(\mathbb{G}): |\alpha|=k} \|\chi_\alpha\|_{L^\infty(\mathbb{G})}^2 = \sum_{\alpha \in \text{Irr}(\mathbb{G}): |\alpha|=k} n_\alpha^2, \quad (4.5)$$

thanks to coamenability of G_q , and the right-hand side grows polynomially for k as in the classical situation by [NT13, Theorem 2.4.7 (3)] and [BV09].

(2) For the other cases $\mathbb{G} = O_F^+$ (resp. $\mathbb{G} = \mathbb{G}_{\text{aut}}(B, \psi)$), let us consider $G = SU(2)$ (resp. $G = SO(3)$). Then $\text{Irr}(\mathbb{G}) \cong \text{Irr}(G) \cong \mathbb{N}_0$ with the same fusion rules by [Ban96] (resp. [Bra13]). For each character $\chi_k \in \text{Pol}(\mathbb{G})$, let us denote by $\chi'_k \in \text{Pol}(G)$ the corresponding character. Then we have

$$\|\chi_k\|_{L^\infty(\mathbb{G})} = \|\chi'_k\|_{L^\infty(G)} = k + 1 \text{ (resp. } 2k + 1), \quad (4.6)$$

by [Wan17, Proposition 6.7] and [Ban96] (resp. [Ban99]), so we can conclude that

$$\sum_{\alpha \in \text{Irr}(\mathbb{G}): |\alpha|=k} \|\chi_\alpha\|_{L^\infty(\mathbb{G})}^2 = \|\chi_k\|_{L^\infty(\mathbb{G})}^2 \lesssim (k + 1)^2. \quad (4.7)$$

Hence, the only remaining thing to prove is the existence of $r \in (0, 1)$ satisfying $\frac{n_\alpha}{d_\alpha} \lesssim r^{|\alpha|}$, which will be discussed in the following Subsections 4.1, 4.2, and 4.3 for each class of examples.

4.1. Drinfeld-Jimbo q -deformations. Let G be a simply connected compact semisimple Lie group and let \mathfrak{g} be the associated complexified Lie algebra with a Cartan subalgebra \mathfrak{h} . The *adjoint map* $\text{ad} : \mathfrak{g} \rightarrow \text{End}(\mathfrak{g})$ is given by

$$\text{ad}_X(Y) = [X, Y] \quad (4.8)$$

for all $X, Y \in \mathfrak{g}$. A non-zero linear functional $\alpha \in \mathfrak{h}^*$ is called a *root* if there exists non-zero $X \in \mathfrak{g}$ such that

$$\text{ad}_H(X) = [H, X] = \alpha(H)X \quad (4.9)$$

for all $H \in \mathfrak{h}$. The space of all roots α is denoted by $\Phi \subseteq \mathfrak{h}^* \setminus \{0\}$, and the associated non-trivial eigenspaces

$$\mathfrak{g}_\alpha = \{X \in \mathfrak{g} : \forall H \in \mathfrak{h}, \text{ad}_H(X) = \alpha(H)X\} \quad (4.10)$$

are called *root spaces*. Furthermore, there exists a subset Φ^+ of Φ satisfying the following:

- The subset Φ^+ contains exactly one of α and $-\alpha$ for each $\alpha \in \Phi$.
- If $\alpha, \beta \in \Phi^+$ ($\alpha \neq \beta$) satisfies $\alpha + \beta \in \Phi$, then $\alpha + \beta \in \Phi^+$.

All elements in Φ^+ are called *positive roots*, and a positive root is called *simple* if it cannot be written as the sum of two positive roots. Let us write the simple roots as $\alpha_1, \alpha_2, \dots, \alpha_r$. Then every positive root $\alpha \in \Phi^+$ is a linear combination of simple roots with non-negative integral coefficients.

There exists a non-degenerate symmetric ad-invariant form (\cdot, \cdot) on \mathfrak{g} whose restriction to the real Lie algebra is negative definite. Note that the non-degenerate symmetric form (\cdot, \cdot) provides a natural identification $\mathfrak{h}^* \cong \mathfrak{h}$, so it induces a natural non-degenerate symmetric form on \mathfrak{h}^* . Furthermore, the form (\cdot, \cdot) restricts to an inner product on $\mathfrak{h}_0^* = \text{span}_{\mathbb{R}} \Phi$. Another important basis of \mathfrak{h}_0^* consists of the so-called *fundamental weights* $\omega_1, \omega_2, \dots, \omega_r \in \mathfrak{h}_0^*$ determined by the following relations

$$\frac{2(\omega_i, \alpha_j)}{(\alpha_j, \alpha_j)} = \delta_{ij} \quad (4.11)$$

for all $1 \leq i, j \leq r$. The set of all \mathbb{Z} -linear combinations of the fundamental weights is called the *weight lattice* for \mathfrak{g} , which we denote by P . The subset of all \mathbb{N}_0 -linear combinations of the fundamental weights is denoted by P_+ .

Now, let $0 < q < 1$ and consider the Drinfeld-Jimbo q -deformation G_q . Then $\text{Irr}(G_q)$ is identified with P_+ . See [CP95, Section 10.1] and [NT13, Theorem 2.4.7]) for more details. Let us denote by $\rho = \sum_{i=1}^r \omega_i$, which is equal to $\frac{1}{2} \sum_{\alpha \in \Phi^+} \alpha$ ([Kna02, Proposition 2.69] and [NT13]). The following proposition is important to establish the Khintchine inequality with operator coefficients for arbitrary central Fourier series on G_q .

Proposition 4.1. *Let $t_i = q^{(\omega_i, 2\rho)}$ for all $1 \leq i \leq r$. Then*

$$\|Q_\mu\|_\infty = t_1^{-\mu_1} t_2^{-\mu_2} \dots t_r^{-\mu_r} \quad (4.12)$$

for all $\mu = \sum_{i=1}^r \mu_i \omega_i \in P_+ \cong \text{Irr}(G_q)$. In particular, $\frac{n_\mu}{d_\mu}$ decreases exponentially as $\sum_{i=1}^r \mu_i \rightarrow \infty$.

Proof. For the associated irreducible representation u^μ of G_q , the eigenvalues of the modular matrix Q_μ are of the form $q^{(\nu, -2\rho)}$ where ν is a weight for u^μ ([NT13, Proposition 2.4.10] and [VY20]). Let us prove that $(\nu, 2\rho)$

is maximized when ν is the highest weight μ . Indeed, since μ is the highest weight for u^μ , the difference $\mu - \nu$ can be written as $\sum_{i=1}^r m_i \alpha_i$ with $m_i \geq 0$ and we have

$$(\mu, 2\rho) - (\nu, 2\rho) = \sum_{i,j=1}^r 2m_i(\alpha_i, \omega_j) = \sum_{i=1}^r m_i(\alpha_i, \alpha_i) \geq 0. \quad (4.13)$$

Thus, we can conclude that

$$\|Q_\mu\|_\infty = q^{-(\mu, 2\rho)} = q^{-\sum_{i=1}^r \mu_i(\omega_i, 2\rho)} = \prod_{i=1}^r (q^{(\omega_i, 2\rho)})^{-\mu_i}. \quad (4.14)$$

For the last conclusion, since n_μ is polynomially growing by [NT13, Theorem 2.4.7 (3)] and [BV09, Theorem 2.1], it is enough to see that d_μ is exponentially growing as $\sum_{i=1}^r \mu_i \rightarrow \infty$. Note that

$$t_i = q^{(\omega_i, 2\rho)} = q^{\sum_{\alpha \in \Phi^+} (\omega_i, \alpha)} \leq q^{(\omega_i, \alpha_i)} = q^{\frac{(\alpha_i, \alpha_i)}{2}} < 1 \quad (4.15)$$

for all $1 \leq i \leq r$, implying $t = \max_{1 \leq i \leq r} t_i \in (0, 1)$. Thus, we obtain the following exponentially growing lower bound:

$$d_\mu \geq \|Q_\mu\|_\infty \geq t^{-(\mu_1 + \mu_2 + \dots + \mu_r)}. \quad (4.16)$$

□

4.2. Free orthogonal quantum groups. Let O_F^+ be a non-Kac free orthogonal quantum group with $F \in M_N$ such that $N_q = \text{Tr}(F^*F) > N$. See [Wan95, Ban96, Tim08] for more details. Since Subsection 4.1 covers the case $N = 2$, let us focus on the other general cases $N \geq 3$. Recall that the space $\text{Irr}(O_F^+)$ is identified with \mathbb{N}_0 with the following fusion rule [Ban96]:

$$u^{(k)} \oplus u^{(1)} \cong u^{(k-1)} \oplus u^{(k+1)}, \quad k \in \mathbb{N}. \quad (4.17)$$

Both the quantum dimension d_k and the classical dimension n_k are described by the *Chebyshev polynomials* $(f_k)_{k \in \mathbb{N}_0}$ of the second kind. The polynomial f_k is explicitly given by

$$f_k(t) = \left(\frac{1}{2}\right)^{k+1} \frac{(t + \sqrt{t^2 - 4})^{k+1} - (t - \sqrt{t^2 - 4})^{k+1}}{\sqrt{t^2 - 4}} \quad (4.18)$$

for all $t > 2$, and we have $d_k = f_k(N_q)$ and $n_k = f_k(N)$. Combining these facts with the following identity

$$\lim_{k \rightarrow \infty} f_k(t) \cdot \left(\frac{t + \sqrt{t^2 - 4}}{2}\right)^{-(k+1)} = \frac{1}{\sqrt{t^2 - 4}}, \quad (4.19)$$

we obtain the following estimate

$$\frac{n_k}{d_k} \approx \left(\frac{N + \sqrt{N^2 - 4}}{N_q + \sqrt{N_q^2 - 4}} \right)^{k+1}. \quad (4.20)$$

Since $N < N_q$ is equivalent to $\frac{N + \sqrt{N^2 - 4}}{N_q + \sqrt{N_q^2 - 4}} < 1$, we can conclude that $\frac{n_k}{d_k}$ decreases exponentially as $k \rightarrow \infty$.

4.3. Quantum automorphism group. Let \mathbb{G} be a non-Kac quantum automorphism group $\mathbb{G}_{\text{aut}}(B, \psi)$. We refer the readers to [Wan98, Ban99, Ban02, Bra13] for more details of $\mathbb{G}_{\text{aut}}(B, \psi)$ with a δ -form ψ . By [Ban99, Bra13], the space $\text{Irr}(\mathbb{G}_{\text{aut}}(B, \psi))$ is identified with \mathbb{N}_0 with the following fusion rule:

$$u^{(k)} \oplus u^{(1)} \cong u^{(k-1)} \oplus u^{(k)} \oplus u^{(k+1)}, \quad k \in \mathbb{N}. \quad (4.21)$$

To compute the quantum dimension and the classical dimension, let us consider the following polynomials g_k such that $g_k(4) = 2k+1$ and $g_k(x) = f_{2k}(\sqrt{x})$ for all $x > 4$. Here, f_{2k} is the $2k$ -th Chebyshev polynomial given in (4.18), and the identity (4.19) implies

$$\lim_{k \rightarrow \infty} g_k(x) \cdot \left(\frac{x - 2 + \sqrt{x(x-4)}}{2} \right)^{-k} = \frac{\sqrt{x} + \sqrt{x-4}}{2\sqrt{x-4}} \quad (4.22)$$

for all $x > 4$. Note that $d_k = g_k(d_1) = f_{2k}(\sqrt{d_1})$ and $n_k = g_k(n_1) = f_{2k}(\sqrt{n_1})$ respectively, by [Bra13, Theorem 3.8].

Firstly, if $\dim(B) = 4$, then $n_k = g_k(4) = 2k+1$ grows polynomially and the above identity (4.22) implies

$$d_k \approx \left(\frac{d_1 - 2 + \sqrt{d_1(d_1 - 4)}}{2} \right)^k. \quad (4.23)$$

Then, the non-Kac condition $d_1 > n_1 = 4$ implies $\frac{d_1 - 2 + \sqrt{d_1(d_1 - 4)}}{2} > 1$, so the quantum dimension d_k has an exponential growth.

For the general cases $\dim(B) \geq 5$, we have

$$\frac{n_k}{d_k} \approx \left(\frac{n_1 - 2 + \sqrt{n_1(n_1 - 4)}}{d_1 - 2 + \sqrt{d_1(d_1 - 4)}} \right)^k \quad (4.24)$$

up to constants by (4.22), and the non-Kac condition, i.e., $n_1 < d_1$, implies that the right-hand side decreases exponentially.

Acknowledgements: The author thanks Professor Éric Ricard for his insightful comments, particularly regarding the consideration of operator coefficients. The author was supported by the National Research Foundation of Korea (NRF) grant funded by the Ministry of Science and ICT (MSIT) (No. 2020R1C1C1A01009681), and also by Samsung Science and Technology Foundation under Project Number SSTF-BA2002-01.

REFERENCES

- [Ban96] Teodor Banica. Théorie des représentations du groupe quantique compact libre $O(n)$. *C. R. Acad. Sci. Paris Sér. I Math.*, 322(3):241–244, 1996.
- [Ban99] Teodor Banica. Symmetries of a generic coaction. *Math. Ann.*, 314(4):763–780, 1999.
- [Ban02] Teodor Banica. Quantum groups and Fuss-Catalan algebras. *Comm. Math. Phys.*, 226(1):221–232, 2002.
- [Bra13] Michael Brannan. Reduced operator algebras of trace-perserving quantum automorphism groups. *Doc. Math.*, 18:1349–1402, 2013.
- [Buc99] Artur Buchholz. Norm of convolution by operator-valued functions on free groups. *Proc. Amer. Math. Soc.*, 127(6):1671–1682, 1999.
- [BV09] Teodor Banica and Roland Vergnioux. Growth estimates for discrete quantum groups. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 12(2):321–340, 2009.
- [CP95] Vyjayanthi Chari and Andrew Pressley. *A guide to quantum groups*. Cambridge University Press, Cambridge, 1995. Corrected reprint of the 1994 original.
- [GH13] Colin C. Graham and Kathryn E. Hare. *Interpolation and Sidon sets for compact groups*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC. Springer, New York, 2013.
- [GST82] Saverio Giulini, Paolo M. Soardi, and Giancarlo Travaglini. Norms of characters and Fourier series on compact Lie groups. *J. Functional Analysis*, 46(1):88–101, 1982.
- [GT80] Saverio Giulini and Giancarlo Travaglini. L^p -estimates for matrix coefficients of irreducible representations of compact groups. *Proc. Amer. Math. Soc.*, 80(3):448–450, 1980.
- [Har99] Asma Harcharras. Fourier analysis, Schur multipliers on S^p and non-commutative $\Lambda(p)$ -sets. *Studia Math.*, 137(3):203–260, 1999.
- [HJX10] Uffe Haagerup, Marius Junge, and Quanhua Xu. A reduction method for non-commutative L_p -spaces and applications. *Trans. Amer. Math. Soc.*, 362(4):2125–2165, 2010.
- [HP93] Uffe Haagerup and Gilles Pisier. Bounded linear operators between C^* -algebras. *Duke Math. J.*, 71(3):889–925, 1993.
- [Jun05] Marius Junge. Embedding of the operator space OH and the logarithmic ‘little Grothendieck inequality’. *Invent. Math.*, 161(2):225–286, 2005.
- [Kah85] Jean-Pierre Kahane. *Some random series of functions*, volume 5 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1985.
- [Kna02] Anthony W. Knaapp. *Lie groups beyond an introduction*, volume 140 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, second edition, 2002.
- [Kos84] Hideki Kosaki. Applications of the complex interpolation method to a von Neumann algebra: noncommutative L^p -spaces. *J. Funct. Anal.*, 56(1):29–78, 1984.

- [KV00] Johan Kustermans and Stefaan Vaes. Locally compact quantum groups. *Ann. Sci. École Norm. Sup. (4)*, 33(6):837–934, 2000.
- [KV03] Johan Kustermans and Stefaan Vaes. Locally compact quantum groups in the von Neumann algebraic setting. *Math. Scand.*, 92(1):68–92, 2003.
- [LP86] Françoise Lust-Piquard. Inégalités de Khintchine dans C_p ($1 < p < \infty$). *C. R. Acad. Sci. Paris Sér. I Math.*, 303(7):289–292, 1986.
- [LPP91] Françoise Lust-Piquard and Gilles Pisier. Noncommutative Khintchine and Paley inequalities. *Ark. Mat.*, 29(2):241–260, 1991.
- [LR75] Jorge M. López and Kenneth A. Ross. *Sidon sets*, volume Vol. 13 of *Lecture Notes in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, 1975.
- [MP81] Michael B. Marcus and Gilles Pisier. *Random Fourier series with applications to harmonic analysis*, volume 101 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1981.
- [NT13] Sergey Neshveyev and Lars Tuset. *Compact quantum groups and their representation categories*, volume 20 of *Cours Spécialisés [Specialized Courses]*. Société Mathématique de France, Paris, 2013.
- [Pis98] Gilles Pisier. Non-commutative vector valued L_p -spaces and completely p -summing maps. *Astérisque*, (247):vi+131, 1998.
- [Pis03] Gilles Pisier. *Introduction to operator space theory*, volume 294 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2003.
- [Pis12] Gilles Pisier. Grothendieck’s theorem, past and present. *Bull. Amer. Math. Soc. (N.S.)*, 49(2):237–323, 2012.
- [Pis20] Gilles Pisier. Completely Sidon sets in discrete groups. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 20(3):891–923, 2020.
- [PP05] Javier Parcet and Gilles Pisier. Non-commutative Khintchine type inequalities associated with free groups. *Indiana Univ. Math. J.*, 54(2):531–556, 2005.
- [PX03] Gilles Pisier and Quanhua Xu. Non-commutative L^p -spaces. In *Handbook of the geometry of Banach spaces, Vol. 2*, pages 1459–1517. North-Holland, Amsterdam, 2003.
- [RX06] Éric Ricard and Quanhua Xu. Khintchine type inequalities for reduced free products and applications. *J. Reine Angew. Math.*, 599:27–59, 2006.
- [Ter81] Marianne Terp. L^p spaces associated with von Neumann algebras. *Notes, Math. Institute, Copenhagen Univ*, 3, 1981.
- [Tim08] Thomas Timmermann. *An invitation to quantum groups and duality*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008. From Hopf algebras to multiplicative unitaries and beyond.
- [Vae02] Stefaan Vaes. *Locally compact quantum groups*. PhD thesis, 2002. Ph. D. Thesis, available at: <http://www.wis.kuleuven.ac.be/analyse/>.
- [VY20] Christian Voigt and Robert Yuncken. *Complex semisimple quantum groups and representation theory*, volume 2264 of *Lecture Notes in Mathematics*. Springer, Cham, [2020] ©2020.
- [Wan95] Shuzhou Wang. Free products of compact quantum groups. *Comm. Math. Phys.*, 167(3):671–692, 1995.
- [Wan98] Shuzhou Wang. Quantum symmetry groups of finite spaces. *Comm. Math. Phys.*, 195(1):195–211, 1998.
- [Wan17] Simeng Wang. Lacunary Fourier series for compact quantum groups. *Comm. Math. Phys.*, 349(3):895–945, 2017.

SANG-GYUN YOUN, DEPARTMENT OF MATHEMATICS EDUCATION, SEOUL NATIONAL UNIVERSITY, GWANAK-RO 1, GWANAK-GU, SEOUL 08826, REPUBLIC OF KOREA

Email address: s.youn@snu.ac.kr