

# FOCK PROJECTIONS ON VECTOR-VALUED $L^p$ -SPACES WITH MATRIX WEIGHTS

JIALE CHEN AND MAOFA WANG

**ABSTRACT.** In this paper, we characterize the  $d \times d$  matrix weights  $W$  on  $\mathbb{C}^n$  such that the Fock projection  $P_\alpha$  is bounded on the vector-valued spaces  $L^p_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)$  induced by  $W$ . It is proved that for  $1 \leq p < \infty$ , the Fock projection  $P_\alpha$  is bounded on  $L^p_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)$  if and only if  $W$  satisfies a restricted  $\mathbf{A}_p$ -condition. In particular, when  $p = 1$ , our result establishes a strong (1,1) type estimate for the Fock projections, which is quite different with the case of Calderón–Zygmund operators.

## 1. INTRODUCTION

The purpose of this paper is to establish some weighted norm inequalities for the Fock projections on vector-valued  $L^p$ -spaces with matrix weights. We start from the basic definitions. Fix positive integers  $n$  and  $d$ . A  $d \times d$  matrix-valued function  $W$  on  $\mathbb{C}^n$  is said to be a matrix weight if

- (i)  $W(z)$  is a positive and invertible operator on  $\mathbb{C}^d$  for almost every  $z \in \mathbb{C}^n$ ;
- (ii) both  $W$  and  $W^{-1}$  are locally integrable on  $\mathbb{C}^n$ .

For such a weight  $W$  and  $1 \leq p < \infty$ ,  $\alpha > 0$ , the weighted space  $L^p_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)$  consists of  $\mathbb{C}^d$ -valued measurable functions  $f$  on  $\mathbb{C}^n$  such that

$$\|f\|_{L^p_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)}^p := \int_{\mathbb{C}^n} |W^{1/p}(z)f(z)|^p e^{-\frac{\alpha}{2}|z|^2} dv(z) < \infty,$$

where  $dv$  is the Lebesgue measure on  $\mathbb{C}^n$  and  $|\cdot|$  is the usual Hilbert space norm on  $\mathbb{C}^d$  (or  $\mathbb{C}^n$ ). In the case  $p = \infty$ , the weighted space  $L^\infty_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)$  of  $\mathbb{C}^d$ -valued measurable functions  $f$  is defined by the following norm:

$$\|f\|_{L^\infty_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)} := \operatorname{ess\,sup}_{z \in \mathbb{C}^n} |W(z)f(z)| e^{-\frac{\alpha}{2}|z|^2}.$$

It is easy to see that the subspace of  $L^p_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)$  consisting of  $\mathbb{C}^d$ -valued entire functions is closed. We denote this subspace by  $F^p_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)$  and call it weighted Fock space. If  $W \equiv E_d$ , the  $d \times d$  identity matrix, then these spaces are denoted simply by  $L^p_\alpha(\mathbb{C}^n; \mathbb{C}^d)$  and  $F^p_\alpha(\mathbb{C}^n; \mathbb{C}^d)$ .

---

*Date:* August 24, 2024.

*2020 Mathematics Subject Classification.* Primary 32A50; Secondary 32A25, 42B20, 42B25.

*Key words and phrases.* Fock projection, matrix weight, restricted  $\mathbf{A}_p$ -condition.

Chen was supported by the Fundamental Research Funds for the Central Universities (No. GK202207018) of China. Wang was supported by the National Natural Science Foundation (No. 12171373) of China.

It is well-known that the Fock projection  $P_\alpha$ , i.e. the orthogonal projection from  $L^2_\alpha(\mathbb{C}^n; \mathbb{C})$  onto the Fock space  $F^2_\alpha(\mathbb{C}^n; \mathbb{C})$ , is an integral operator given by

$$P_\alpha f(z) := \int_{\mathbb{C}^n} f(u) \overline{K_z^\alpha(u)} d\lambda_\alpha(u), \quad z \in \mathbb{C}^n,$$

where  $d\lambda_\alpha(u) := \left(\frac{\alpha}{\pi}\right)^n e^{-\alpha|u|^2} dv(u)$  is the Gaussian measure and  $K_z^\alpha(u) := e^{\alpha\langle u, z \rangle}$  denotes the reproducing kernel of  $F^2_\alpha(\mathbb{C}^n; \mathbb{C})$  at  $z$ . Dostanić and Zhu [7] proved that for all  $1 \leq p \leq \infty$ ,  $P_\alpha$  is bounded on  $L^p_\alpha(\mathbb{C}^n; \mathbb{C})$ .

A longstanding theme in harmonic analysis is the weighted theory of singular integral operators, which was initiated by Muckenhoupt [15]. It was proved in [15] that for the scalar-valued setting  $d = 1$ , the Hardy–Littlewood maximal function is bounded on  $L^p(W)$  ( $1 < p < \infty$ ) if and only if the weight  $W$  satisfies what we call the Muckenhoupt  $\mathbf{A}_p$ -condition today. Later, it was shown that the same condition also characterizes the boundedness of the Riesz projection (equivalently, the Hilbert transform) on  $L^p(W)$ ; see [5, 11]. One direction to extend these results is to consider the vector-valued spaces with matrix weights. Treil and Volberg [20] introduced the  $\mathbf{A}_2$ -condition for matrix weights and proved that the Riesz projection is bounded on  $L^2(W)$  if and only if the matrix weight  $W$  satisfies the  $\mathbf{A}_2$ -condition. Nazarov–Treil’ [17] and Volberg [21] introduced the matrix  $\mathbf{A}_p$ -condition and generalized Treil–Volberg’s result to all  $1 < p < \infty$ . Moreover, it was pointed out in [19, 20, 21] that the matrix weighted estimates for the Riesz projection is closely related to the theory of stationary processes, the invertibility of Toeplitz operators with matrix-valued symbols and a two-weight estimate for the Hilbert transform. Therefore, the matrix weighted theory of singular integral operators has drawn widespread attentions since then; see [1, 4, 6, 8, 10, 14, 16, 18] and the references therein. In particular, Frazier and Roudenko [8] introduced the  $\mathbf{A}_1$ -condition for matrix weights, and Cruz-Uribe et al. [6] established some weak (1,1) type estimates for maximal functions and Calderón–Zygmund operators in the case that the matrix weight satisfies the  $\mathbf{A}_1$ -condition.

Similarly to the case of Calderón–Zygmund operators, one can consider the weighted theory of the Fock projections. To determine the bounded and invertible products of Toeplitz operators on Fock spaces  $F^p_\alpha(\mathbb{C}^n; \mathbb{C})$ , Isralowitz [13] proved that the Fock projection  $P_\alpha$  is bounded on the scalar-valued space  $L^p_{\alpha, W}(\mathbb{C}^n; \mathbb{C})$  ( $1 < p < \infty$ ) if and only if  $W$  satisfies the following restricted  $\mathbf{A}_p$ -condition: for some (or any)  $r > 0$ ,

$$\sup_{Q \subset \mathbb{C}^n: l(Q)=r} \left( \frac{1}{v(Q)} \int_Q W dv \right) \left( \frac{1}{v(Q)} \int_Q W^{-p'/p} dv \right)^{p/p'} < \infty.$$

Here,  $Q$  denotes a cube in  $\mathbb{C}^n$  whose side length is denoted by  $l(Q)$ , and  $p' = p/(p-1)$  is the conjugate exponent of  $p$ . Later, Cascante, Fàbrega and Peláez [2] proved that  $P_\alpha$  is bounded on  $L^1_{\alpha, W}(\mathbb{C}; \mathbb{C})$  if and only if the weight  $W$  satisfies the following restricted  $\mathbf{A}_1$ -condition: for some (or any)  $r > 0$ ,

$$\sup_{Q \subset \mathbb{C}: l(Q)=r} \frac{\int_Q W dv}{v(Q) \operatorname{ess\,inf}_{u \in Q} W(u)} < \infty.$$

Motivated by the aforementioned works, the authors [3] recently characterized the matrix weights  $W$  on  $\mathbb{C}$  such that the Fock projection  $P_\alpha$  is bounded on the Hilbert spaces  $L^2_{\alpha,W}(\mathbb{C}; \mathbb{C}^d)$ . It was proved that  $P_\alpha$  is bounded on  $L^2_{\alpha,W}(\mathbb{C}; \mathbb{C}^d)$  if and only if the matrix weight  $W$  satisfies the following  $\mathbf{A}_{2,r}$ -condition for some (or any)  $r > 0$ :

$$\sup_{Q \subset \mathbb{C}: l(Q)=r} \left\| \left( \frac{1}{v(Q)} \int_Q W dv \right)^{1/2} \left( \frac{1}{v(Q)} \int_Q W^{-1} dv \right)^{1/2} \right\| < \infty. \quad (1.1)$$

Here and in the sequel, for a  $d \times d$  matrix  $M$ ,  $\|M\|$  denotes its operator norm acting on  $\mathbb{C}^d$ . In this paper, we are going to extend this result to the full range  $1 \leq p < \infty$  and to the matrix weights defined on  $\mathbb{C}^n$ . Our main result establishes the Fock space analogy of [10, 21].

As was stated in [21], the matrix  $\mathbf{A}_p$ -condition does not have the form as in (1.1) if  $p \neq 2$ . We will follow the line in [21] to define some  $\mathbf{A}_{p,r}$ -conditions for matrix weights. Then it is proved that for  $1 \leq p < \infty$ , the Fock projection  $P_\alpha$  is bounded on  $L^p_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)$  if and only if  $W$  satisfies the  $\mathbf{A}_{p,r}$ -condition for some (or any)  $r > 0$ ; see Theorem 3.1. More precisely, via a class of integral operators induced by the normalized reproducing kernels of  $F^2_\alpha(\mathbb{C}^n; \mathbb{C})$ , we show that if  $P_\alpha$  is bounded on  $L^p_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)$ , then the weight  $W$  satisfies the  $\mathbf{A}_{p,r}$ -condition for any  $r > 0$ . Conversely, based on some estimates for  $\mathbf{A}_{p,r}$ -weights and a duality argument, we prove that if  $W$  satisfies the  $\mathbf{A}_{p,r}$ -condition for some  $r > 0$ , then the “maximal” Fock projection  $P^+_{\alpha,W}$ , defined by

$$P^+_{\alpha,W}(f)(z) := \int_{\mathbb{C}^n} |W^{1/p}(z)W^{-1/p}(u)f(u)| |K^\alpha_z(u)| d\lambda_\alpha(u),$$

is bounded from  $L^p_\alpha(\mathbb{C}^n; \mathbb{C}^d)$  to  $L^p_\alpha(\mathbb{C}^n; \mathbb{C})$ , which implies the boundedness of  $P_\alpha$  on  $L^p_{\alpha,W}(\mathbb{C}^n; \mathbb{C}^d)$ . In particular, when  $p = 1$ , our result establishes a strong (1,1) type estimate for the Fock projections, which is quite different with the case of Calderón–Zygmund operators [6].

The rest part of this paper is organized as follows. In Section 2, we recall some preliminary results and introduce the restricted  $\mathbf{A}_p$ -conditions for matrix weights. Then we state and prove our main result in Section 3.

Throughout the paper, the notation  $\phi_1 \lesssim \phi_2$  (or  $\phi_2 \gtrsim \phi_1$ ) means that there exists a nonessential constant  $c > 0$  such that  $\phi_1 \leq c\phi_2$ . If  $\phi_1 \lesssim \phi_2 \lesssim \phi_1$ , then we write  $\phi_1 \asymp \phi_2$ . For a subset  $E \subset \mathbb{C}^n$ , we use  $\chi_E$  to denote the characteristic function of  $E$ . Given  $z \in \mathbb{C}^n$  and  $r > 0$ ,  $Q_r(z)$  denotes the cube centered at  $z$  with side length  $l(Q) = r$ . Finally, we write  $p'$  for the conjugate exponent of  $p$  for  $1 \leq p < \infty$ , i.e.  $\frac{1}{p} + \frac{1}{p'} = 1$ . In particular,  $1' = \infty$ .

## 2. PRELIMINARIES AND DEFINITIONS

In this section, we recall some preliminaries and introduce the restricted  $\mathbf{A}_p$ -conditions for matrix weights.

Let  $\alpha > 0$  and  $W$  be a  $d \times d$  matrix weight. We use  $\langle \cdot, \cdot \rangle_\alpha$  to denote the pairing defined as follows:

$$\langle f, g \rangle_\alpha := \int_{\mathbb{C}^n} \langle f(z), g(z) \rangle e^{-\alpha|z|^2} dv(z),$$

where  $f$  and  $g$  are  $\mathbb{C}^d$ -valued measurable functions. It is well-known that, with respect to the pairing  $\langle \cdot, \cdot \rangle_\alpha$ , the dual space of  $L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)$  ( $1 < p < \infty$ ) can be represented as

$$(L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d))^* = L_{\alpha, W^{-p'/p}}^{p'}(\mathbb{C}^n; \mathbb{C}^d), \quad (2.1)$$

and the dual space of  $L_{\alpha, W}^1(\mathbb{C}^n; \mathbb{C}^d)$  can be represented as

$$(L_{\alpha, W}^1(\mathbb{C}^n; \mathbb{C}^d))^* = L_{\alpha, W^{-1}}^\infty(\mathbb{C}^n; \mathbb{C}^d). \quad (2.2)$$

We now introduce the restricted  $\mathbf{A}_p$ -conditions for matrix weights. To this end, we consider the more general situation of norm-valued functions. Let  $z \mapsto \rho_z$ ,  $z \in \mathbb{C}^n$  be a function whose values are Banach space norms on  $\mathbb{C}^d$ . We assume this function to be measurable in the sense that for any  $\mathbf{x} \in \mathbb{C}^d$ , the function  $z \mapsto \rho_z(\mathbf{x})$  is measurable. For convenience, we will say the norm-valued function  $z \mapsto \rho_z$  to be a metric and denote it by  $\rho$ .

Let  $\rho$  be a metric. For  $z \in \mathbb{C}^n$ , the dual norm  $(\rho_z)^*$  of  $\rho_z$  is defined by

$$(\rho_z)^*(\mathbf{x}) := \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle \mathbf{x}, \mathbf{y} \rangle|}{\rho_z(\mathbf{y})}, \quad \mathbf{x} \in \mathbb{C}^d.$$

Since  $\mathbb{C}^d$  is reflexive, we have  $(\rho_z)^{**} = \rho_z$ . The dual metric  $\rho^*$  is defined pointwisely by  $\rho_z^* = (\rho_z)^*$ . For  $1 \leq p < \infty$  and a cube  $Q \subset \mathbb{C}^n$ , consider the norms

$$\rho_{p, Q}(\mathbf{x}) := \left( \frac{1}{v(Q)} \int_Q (\rho_z(\mathbf{x}))^p dv(z) \right)^{1/p}, \quad \mathbf{x} \in \mathbb{C}^d$$

and

$$\rho_{\infty, Q}(\mathbf{x}) := \operatorname{ess\,sup}_{z \in Q} \rho_z(\mathbf{x}), \quad \mathbf{x} \in \mathbb{C}^d.$$

The following lemma was proved in [21, Proposition 1.1] for the case  $1 < p < \infty$ . The case  $p = 1$  is similar and we omit the proof here.

**Lemma 2.1.** *Let  $1 \leq p < \infty$  and  $\rho$  be a metric. Then for any  $\mathbf{x} \in \mathbb{C}^d$  and any cube  $Q \subset \mathbb{C}^n$ ,*

$$\rho_{p', Q}^*(\mathbf{x}) \geq (\rho_{p, Q})^*(\mathbf{x}).$$

For  $1 \leq p < \infty$  and  $r > 0$ , a metric  $\rho$  is called an  $\mathbf{A}_{p, r}$ -metric if there exists some constant  $C > 0$  such that the opposite inequality

$$\rho_{p', Q}^*(\mathbf{x}) \leq C (\rho_{p, Q})^*(\mathbf{x})$$

holds for any  $\mathbf{x} \in \mathbb{C}^d$  and any cube  $Q \subset \mathbb{C}^n$  with  $l(Q) = r$ . The  $\mathbf{A}_{p, r}$ -constant of  $\rho$ , denoted by  $\mathcal{C}_{p, r}(\rho)$ , is the least suitable constant  $C$ . Let  $W$  be a  $d \times d$  matrix weight. For  $1 \leq p < \infty$  and  $r > 0$ , we say that  $W$  is an  $\mathbf{A}_{p, r}$ -weight if the metric  $\rho$  defined by

$$\rho_z(\mathbf{x}) = |W^{1/p}(z)\mathbf{x}|, \quad \mathbf{x} \in \mathbb{C}^d, \quad z \in \mathbb{C}^n$$

is an  $\mathbf{A}_{p, r}$ -metric. Moreover, we write  $\mathcal{C}_{p, r}(W) = \mathcal{C}_{p, r}(\rho)$ .

It was pointed out in [21, p. 449] (see also [10, Proposition 1.2]) that for a Banach space norm  $\rho_z$  on  $\mathbb{C}^d$ , there exists a  $d \times d$  self-adjoint matrix  $M_z$ , which is a positive and invertible operator on  $\mathbb{C}^d$ , such that

$$\rho_z(\mathbf{x}) \leq |M_z \mathbf{x}| \leq \sqrt{d} \cdot \rho_z(\mathbf{x}), \quad \mathbf{x} \in \mathbb{C}^d.$$

Then the dual norm  $(\rho_z)^*$  satisfies

$$|M_z^{-1} \mathbf{x}| \leq (\rho_z)^*(\mathbf{x}) \leq \sqrt{d} \cdot |M_z^{-1} \mathbf{x}|, \quad \mathbf{x} \in \mathbb{C}^d.$$

We will call the matrix  $M_z$  the reducing operator of  $\rho_z$ . It is now possible to state the  $\mathbf{A}_{p,r}$ -condition in terms of the reducing operators. Let  $1 \leq p < \infty$  and  $\rho$  be a metric. For any cube  $Q \subset \mathbb{C}^n$ , we use  $\mathcal{R}_Q$  and  $\mathcal{R}_Q^*$  to denote the reducing operators of  $\rho_{p,Q}$  and  $\rho_{p',Q}^*$ , respectively. Consequently,

$$\rho_{p,Q}(\mathbf{x}) \leq |\mathcal{R}_Q \mathbf{x}| \leq \sqrt{d} \cdot \rho_{p,Q}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{C}^d, \quad (2.3)$$

and

$$\rho_{p',Q}^*(\mathbf{x}) \leq |\mathcal{R}_Q^* \mathbf{x}| \leq \sqrt{d} \cdot \rho_{p',Q}^*(\mathbf{x}), \quad \mathbf{x} \in \mathbb{C}^d. \quad (2.4)$$

Combining these inequalities with the definition of the  $\mathbf{A}_{p,r}$ -metric, we obtain that  $\rho$  is an  $\mathbf{A}_{p,r}$ -metric if and only if

$$\sup_{Q \subset \mathbb{C}^n: l(Q)=r} \|\mathcal{R}_Q \mathcal{R}_Q^*\| < \infty.$$

Moreover,

$$\mathcal{C}_{p,r}(\rho) \leq \sup_{Q \subset \mathbb{C}^n: l(Q)=r} \|\mathcal{R}_Q \mathcal{R}_Q^*\| \leq d \mathcal{C}_{p,r}(\rho). \quad (2.5)$$

**Remark 2.2.** Let  $W$  be a matrix weight and let the metric  $\rho$  be defined by  $\rho_z(\mathbf{x}) = |W^{1/2}(z)\mathbf{x}|$ . Then the reducing operators of  $\rho_{2,Q}$  and  $\rho_{2,Q}^*$  can be calculated as follows:

$$\mathcal{R}_Q = \left( \frac{1}{v(Q)} \int_Q W dv \right)^{1/2} \quad \text{and} \quad \mathcal{R}_Q^* = \left( \frac{1}{v(Q)} \int_Q W^{-1} dv \right)^{1/2},$$

which gives the  $\mathbf{A}_{2,r}$ -condition (1.1).

### 3. THE MAIN RESULT

In this section, we state and prove our main result, which characterizes the boundedness of  $P_\alpha$  on  $L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)$  for all  $1 \leq p < \infty$ . Moreover, the corresponding norm estimate for  $P_\alpha$  is established.

**Theorem 3.1.** *Let  $\alpha > 0$ ,  $1 \leq p < \infty$ , and let  $W$  be a  $d \times d$  matrix weight on  $\mathbb{C}^n$ . The following conditions are equivalent:*

- (a)  $P_\alpha$  is bounded on  $L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)$ ;
- (b)  $P_{\alpha,W}^+ : L_\alpha^p(\mathbb{C}^n; \mathbb{C}^d) \rightarrow L_\alpha^p(\mathbb{C}^n; \mathbb{C})$  is bounded;
- (c)  $W$  is an  $\mathbf{A}_{p,r}$ -weight for any  $r > 0$ ;
- (d)  $W$  is an  $\mathbf{A}_{p,r}$ -weight for some  $r > 0$ .

Moreover, for any fixed  $r > 0$ , there exists  $c = c(\alpha, p, r, n) > 0$  such that

$$\left(\frac{\alpha r^2}{\pi}\right)^n e^{-n\alpha r^2} \mathcal{C}_{p,r}(W)^{1/2} \leq \|P_\alpha\| \leq \|P_{\alpha,W}^+\| \leq c d^{c_p} \mathcal{C}_{p,r}(W)^{c(1+\log \mathcal{C}_{p,r}(W))},$$

where  $c_p = 7/2$  for  $1 < p < \infty$ , and  $c_1 = 9/2$ .

As a byproduct of Theorem 3.1, we know that the class of  $\mathbf{A}_{p,r}$ -weight (more generally, the class of  $\mathbf{A}_{p,r}$ -metric) is actually independent of the choice of  $r$ . In fact, we will give a direct and quantitative proof of this fact; see Proposition 3.7.

For the proof of Theorem 3.1, we first note that if  $P_{\alpha,W}^+ : L_\alpha^p(\mathbb{C}^n; \mathbb{C}^d) \rightarrow L_\alpha^p(\mathbb{C}^n; \mathbb{C})$  is bounded, then for  $f \in L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)$ ,

$$\begin{aligned} \|P_\alpha f\|_{L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)}^p &= \int_{\mathbb{C}^n} \left| W^{1/p}(z) \int_{\mathbb{C}^n} f(u) \overline{K_z^\alpha(u)} d\lambda_\alpha(u) \right|^p e^{-\frac{p\alpha}{2}|z|^2} dv(z) \\ &\leq \int_{\mathbb{C}^n} \left( \int_{\mathbb{C}^n} |W^{1/p}(z) f(u)| |K_z^\alpha(u)| d\lambda_\alpha(u) \right)^p e^{-\frac{p\alpha}{2}|z|^2} dv(z) \\ &= \|P_{\alpha,W}^+(W^{1/p} f)\|_{L_\alpha^p(\mathbb{C}^n; \mathbb{C})}^p \\ &\leq \|P_{\alpha,W}^+\|^p \|W^{1/p} f\|_{L_\alpha^p(\mathbb{C}^n; \mathbb{C}^d)}^p \\ &= \|P_{\alpha,W}^+\|^p \|f\|_{L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)}^p. \end{aligned}$$

Therefore, the implication (b) $\implies$ (a) of Theorem 3.1 holds, and

$$\|P_\alpha\| \leq \|P_{\alpha,W}^+\|.$$

The rest part of the paper is devoted to proving the implication (a) $\implies$ (c) and the implication (d) $\implies$ (b).

To establish the implication (a) $\implies$ (c), we consider a class of integral operators induced by the normalized reproducing kernels. Given  $\alpha > 0$  and  $u \in \mathbb{C}^n$ , we use  $k_u^\alpha$  to denote the normalized reproducing kernel of  $F_\alpha^2(\mathbb{C}^n; \mathbb{C})$ , that is,

$$k_u^\alpha(z) = e^{\alpha\langle z, u \rangle - \frac{\alpha}{2}|u|^2}, \quad z \in \mathbb{C}^n.$$

For fixed  $r > 0$ , define an operator  $P_{\alpha,u,r}$  for  $\mathbb{C}^d$ -valued functions  $f$  by

$$P_{\alpha,u,r} f = \chi_{Q_r(u)} k_u^\alpha \int_{Q_r(u)} f \overline{k_u^\alpha} d\lambda_\alpha.$$

The following proposition reveals the relation between the boundedness of  $P_\alpha$  and  $P_{\alpha,u,r}$ .

**Proposition 3.2.** *Let  $\alpha, r > 0$ ,  $1 \leq p < \infty$ ,  $u \in \mathbb{C}^n$ , and let  $W$  be a  $d \times d$  matrix weight on  $\mathbb{C}^n$ . Suppose that  $P_\alpha$  is bounded on  $L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)$ . Then  $P_{\alpha,u,r}$  is bounded on  $L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)$ , and*

$$\|P_{\alpha,u,r}\| \leq e^{n\alpha r^2/2} \|P_\alpha\|.$$

*Proof.* Write  $Q = Q_r(u)$  temporarily to save the notation. Since  $P_\alpha$  is bounded on  $L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)$ , it is clear that for any  $f \in L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)$ ,

$$\|\chi_Q P_\alpha(\chi_Q f)\|_{L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)} \leq \|P_\alpha\| \|f\|_{L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)}. \quad (3.1)$$

We now estimate the norm of  $P_{\alpha,u,r}f - \chi_Q P_\alpha(\chi_Q f)$ . For any  $z \in \mathbb{C}$ ,

$$\begin{aligned} & P_{\alpha,u,r}f(z) - \chi_Q(z)P_\alpha(\chi_Q f)(z) \\ &= \chi_Q(z) \int_Q f(\zeta) e^{\alpha\langle z, \zeta \rangle} \left( \sum_{k=1}^{\infty} \frac{(-1)^k \alpha^k}{k!} \langle u - z, u - \zeta \rangle^k \right) d\lambda_\alpha(\zeta); \end{aligned}$$

see the proof of [3, Proposition 2.2]. Note that for any  $k \geq 1$ ,

$$\begin{aligned} \langle u - z, u - \zeta \rangle^k &= \left( \sum_{j=1}^n (u_j - z_j) \overline{(u_j - \zeta_j)} \right)^k \\ &= \sum_{\substack{k_1, k_2, \dots, k_n \geq 0 \\ k_1 + \dots + k_n = k}} \frac{k!}{k_1! \dots k_n!} \prod_{j=1}^n (u_j - z_j)^{k_j} \overline{(u_j - \zeta_j)^{k_j}}. \end{aligned}$$

For  $u \in \mathbb{C}^n$  and  $k_1, \dots, k_n \geq 0$ , write  $\tau_u^{k_1, \dots, k_n}(z) = \prod_{j=1}^n (u_j - z_j)^{k_j}$ . Then

$$\begin{aligned} & P_{\alpha,u,r}f(z) - \chi_Q(z)P_\alpha(\chi_Q f)(z) \\ &= \chi_Q(z) \int_Q f(\zeta) e^{\alpha\langle z, \zeta \rangle} \left( \sum_{k=1}^{\infty} \frac{(-1)^k \alpha^k}{k!} \sum_{\substack{k_1, k_2, \dots, k_n \geq 0 \\ k_1 + \dots + k_n = k}} \frac{k!}{k_1! \dots k_n!} \tau_u^{k_1, \dots, k_n}(z) \overline{\tau_u^{k_1, \dots, k_n}(\zeta)} \right) d\lambda_\alpha(\zeta) \\ &= \sum_{k=1}^{\infty} \frac{(-1)^k \alpha^k}{k!} \sum_{\substack{k_1, k_2, \dots, k_n \geq 0 \\ k_1 + \dots + k_n = k}} \frac{k!}{k_1! \dots k_n!} \chi_Q(z) \tau_u^{k_1, \dots, k_n}(z) \int_Q \overline{\tau_u^{k_1, \dots, k_n}(\zeta)} f(\zeta) e^{\alpha\langle z, \zeta \rangle} d\lambda_\alpha(\zeta) \\ &= \sum_{k=1}^{\infty} \frac{(-1)^k \alpha^k}{k!} \sum_{\substack{k_1, k_2, \dots, k_n \geq 0 \\ k_1 + \dots + k_n = k}} \frac{k!}{k_1! \dots k_n!} \chi_Q(z) \tau_u^{k_1, \dots, k_n}(z) P_\alpha \left( \chi_Q \overline{\tau_u^{k_1, \dots, k_n}} f \right)(z). \end{aligned}$$

Noting that for any  $k \geq 1$  and any  $k_1, \dots, k_n \geq 0$  with  $k_1 + \dots + k_n = k$ ,

$$\begin{aligned} \|\chi_Q \tau_u^{k_1, \dots, k_n} f\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)}^p &= \int_Q |W^{1/p}(z) f(z)|^p \prod_{j=1}^n |u_j - z_j|^{k_j p} e^{-\frac{p\alpha}{2}|z|^2} dv(z) \\ &\leq 2^{-kp/2} r^{kp} \|f\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)}^p, \end{aligned}$$

we obtain that

$$\begin{aligned} & \|P_{\alpha,u,r}f - \chi_Q P_\alpha(\chi_Q f)\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)} \\ &\leq \sum_{k=1}^{\infty} \frac{\alpha^k}{k!} \sum_{\substack{k_1, k_2, \dots, k_n \geq 0 \\ k_1 + \dots + k_n = k}} \frac{k!}{k_1! \dots k_n!} \left\| \chi_Q \tau_u^{k_1, \dots, k_n} P_\alpha \left( \chi_Q \overline{\tau_u^{k_1, \dots, k_n}} f \right) \right\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)} \\ &\leq \sum_{k=1}^{\infty} \frac{\alpha^k r^k}{2^{k/2} k!} \sum_{\substack{k_1, k_2, \dots, k_n \geq 0 \\ k_1 + \dots + k_n = k}} \frac{k!}{k_1! \dots k_n!} \left\| P_\alpha \left( \chi_Q \overline{\tau_u^{k_1, \dots, k_n}} f \right) \right\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)} \\ &\leq \|P_\alpha\| \sum_{k=1}^{\infty} \frac{\alpha^k r^k}{2^{k/2} k!} \sum_{\substack{k_1, k_2, \dots, k_n \geq 0 \\ k_1 + \dots + k_n = k}} \frac{k!}{k_1! \dots k_n!} \left\| \chi_Q \overline{\tau_u^{k_1, \dots, k_n}} f \right\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)} \end{aligned}$$



$$\begin{aligned}
&\leq \|P_\alpha\| \sum_{k=1}^{\infty} \frac{\alpha^k r^{2k}}{2^k k!} \sum_{\substack{k_1, k_2, \dots, k_n \geq 0 \\ k_1 + \dots + k_n = k}} \frac{k!}{k_1! \dots k_n!} \|f\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)} \\
&= \left( e^{\frac{n\alpha r^2}{2}} - 1 \right) \|P_\alpha\| \|f\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)}.
\end{aligned}$$

Combining this with (3.1) yields the desired result.  $\square$

The following lemma is easy to verify, so we omit the proof.

**Lemma 3.3.** *Let  $\alpha, r > 0$ ,  $1 \leq p < \infty$ ,  $u \in \mathbb{C}^n$ , and let  $W$  be a  $d \times d$  matrix weight on  $\mathbb{C}^n$ . Then the operator  $P_{\alpha, u, r} : L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d) \rightarrow L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)$  is self-adjoint under the pairing  $\langle \cdot, \cdot \rangle_\alpha$ , that is, for  $f \in L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)$  and  $g \in (L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d))^*$ ,*

$$\langle P_{\alpha, u, r} f, g \rangle_\alpha = \langle f, P_{\alpha, u, r} g \rangle_\alpha.$$

**Lemma 3.4.** *Let  $\alpha, r > 0$ ,  $1 \leq p < \infty$ ,  $u \in \mathbb{C}^n$ , and let  $W$  be a  $d \times d$  matrix weight on  $\mathbb{C}^n$  such that  $P_\alpha$  is bounded on  $L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)$ . Let the function  $f$  be defined by*

$$f = c \chi_{Q_r(u)} k_u^\alpha \mathbf{x},$$

where  $c \in \mathbb{C}$  and  $\mathbf{x} \in \mathbb{C}^d$ . Then

(1) for  $1 < p < \infty$ ,

$$\sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle f, \chi_{Q_r(u)} k_u^\alpha \mathbf{y} \rangle_\alpha|}{\|\chi_{Q_r(z)} k_u^\alpha \mathbf{y}\|_{L_{\alpha, W^{-p'/p}}^{p'}(\mathbb{C}^n; \mathbb{C}^d)}} \geq \left( \frac{\alpha r^2}{\pi} \right)^n e^{-n\alpha r^2} \|P_\alpha\|^{-1} \|f\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)}.$$

(2) for  $p = 1$ ,

$$\sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle f, \chi_{Q_r(u)} k_u^\alpha \mathbf{y} \rangle_\alpha|}{\|\chi_{Q_r(z)} k_u^\alpha \mathbf{y}\|_{L_{\alpha, W^{-1}}^\infty(\mathbb{C}^n; \mathbb{C}^d)}} \geq \left( \frac{\alpha r^2}{\pi} \right)^n e^{-n\alpha r^2} \|P_\alpha\|^{-1} \|f\|_{L_{\alpha, W}^1(\mathbb{C}^n; \mathbb{C}^d)}.$$

*Proof.* It is easy to see that  $P_{\alpha, u, r} f = c_{\alpha, u, r} f$ , where  $c_{\alpha, u, r} = \int_{Q_r(u)} |k_u^\alpha|^2 d\lambda_\alpha$ . Suppose first that  $1 < p < \infty$ . Using the duality (2.1) and Lemma 3.3, we obtain that

$$\begin{aligned}
\|f\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)} &= \sup_{\phi} |\langle f, \phi \rangle_\alpha| = \sup_{\phi} \left| \left\langle P_{\alpha, u, r} f, \frac{1}{c_{\alpha, u, r}} \phi \right\rangle_\alpha \right| \\
&= \sup_{\phi} \left| \left\langle f, \frac{1}{c_{\alpha, u, r}} P_{\alpha, u, r} \phi \right\rangle_\alpha \right|,
\end{aligned}$$

where the supremum is taken over  $\phi \in L_{\alpha, W^{-p'/p}}^{p'}(\mathbb{C}^n; \mathbb{C}^d)$  with  $\|\phi\| \leq 1$ . Write  $\mathbf{y}_\phi = \int_{Q_r(u)} \phi \overline{k_u^\alpha} d\lambda_\alpha$ . Then  $P_{\alpha, u, r} \phi = \chi_{Q_r(u)} k_u^\alpha \mathbf{y}_\phi$ . Therefore,

$$\|f\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)} = \sup_{\phi: \mathbf{y}_\phi \neq 0} \left| \left\langle f, \frac{1}{c_{\alpha, u, r}} \chi_{Q_r(u)} k_u^\alpha \mathbf{y}_\phi \right\rangle_\alpha \right|.$$



By Proposition 3.2 and Lemma 3.3, we know that the operator  $P_{\alpha,u,r}$  is bounded on  $L_{\alpha,W^{-p'/p}}^{p'}(\mathbb{C}^n; \mathbb{C}^d)$ , which implies that

$$\begin{aligned} \left\| \frac{1}{c_{\alpha,u,r}} \chi_{Q_r(u)} k_u^\alpha \mathbf{y}_\phi \right\|_{L_{\alpha,W^{-p'/p}}^{p'}(\mathbb{C}^n; \mathbb{C}^d)} &= \frac{1}{c_{\alpha,u,r}} \|P_{\alpha,u,r} \phi\|_{L_{\alpha,W^{-p'/p}}^{p'}(\mathbb{C}^n; \mathbb{C}^d)} \\ &\leq \frac{1}{c_{\alpha,u,r}} \|P_{\alpha,u,r}\| \\ &\leq \frac{1}{c_{\alpha,u,r}} e^{\frac{n\alpha r^2}{2}} \|P_\alpha\|. \end{aligned}$$

It is clear that  $c_{\alpha,u,r} = \left(\frac{\alpha}{\pi}\right)^n \int_{Q_r(u)} e^{-\alpha|z-u|^2} dv(z) \geq \left(\frac{\alpha r^2}{\pi}\right)^n e^{-n\alpha r^2/2}$ . Hence

$$\left\| \frac{1}{c_{\alpha,u,r}} \chi_{Q_r(u)} k_u^\alpha \mathbf{y}_\phi \right\|_{L_{\alpha,W^{-p'/p}}^{p'}(\mathbb{C}^n; \mathbb{C}^d)} \leq \left(\frac{\pi}{\alpha r^2}\right)^n e^{n\alpha r^2} \|P_\alpha\|.$$

Consequently,

$$\begin{aligned} \|f\|_{L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)} &\leq \left(\frac{\pi}{\alpha r^2}\right)^n e^{n\alpha r^2} \|P_\alpha\| \sup_{\phi: \mathbf{y}_\phi \neq 0} \frac{\left| \left\langle f, \chi_{Q_r(u)} k_u^\alpha \frac{\mathbf{y}_\phi}{c_{\alpha,u,r}} \right\rangle_\alpha \right|}{\left\| \chi_{Q_r(u)} k_u^\alpha \frac{\mathbf{y}_\phi}{c_{\alpha,u,r}} \right\|_{L_{\alpha,W^{-p'/p}}^{p'}(\mathbb{C}^n; \mathbb{C}^d)}} \\ &\leq \left(\frac{\pi}{\alpha r^2}\right)^n e^{n\alpha r^2} \|P_\alpha\| \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{\left| \left\langle f, \chi_{Q_r(u)} k_u^\alpha \mathbf{y} \right\rangle_\alpha \right|}{\left\| \chi_{Q_r(u)} k_u^\alpha \mathbf{y} \right\|_{L_{\alpha,W^{-p'/p}}^{p'}(\mathbb{C}^n; \mathbb{C}^d)}}, \end{aligned}$$

which is exactly what we want. In the case  $p = 1$ , we can use the duality (2.2) to obtain the desired result by following the same procedure. The details are omitted.  $\square$

We are now ready to prove the implication (a) $\implies$ (c) of Theorem 3.1.

*Proof of Theorem 3.1.* (a) $\implies$ (c). Fix  $r > 0$ , and let  $\rho$  be the metric defined by

$$\rho_z(\mathbf{x}) = |W^{1/p}(z)\mathbf{x}|, \quad \mathbf{x} \in \mathbb{C}^d, \quad z \in \mathbb{C}^n.$$

It is sufficient to show that  $\rho$  is an  $\mathbf{A}_{p,r}$ -metric. To this end, fix  $u \in \mathbb{C}^n$  and  $\mathbf{x} \in \mathbb{C}^d$ . Define

$$f = \frac{1}{c_{\alpha,u,r}} \chi_{Q_r(u)} k_u^\alpha \mathbf{x}.$$

Then  $P_{\alpha,u,r} f = \chi_{Q_r(u)} k_u^\alpha \mathbf{x}$  and  $\mathbf{x} = \int_{Q_r(u)} f \overline{k_u^\alpha} d\lambda_\alpha$ . Combining the boundedness of  $P_\alpha$  with Proposition 3.2 yields that

$$\begin{aligned} e^{n\alpha r^2/2} \|P_\alpha\| \|f\|_{L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)} &\geq \|P_{\alpha,u,r} f\|_{L_{\alpha,W}^p(\mathbb{C}^n; \mathbb{C}^d)} \\ &= \left( \int_{Q_r(u)} |W^{1/p}(z) k_u^\alpha(z) \mathbf{x}|^p e^{-\frac{p\alpha}{2}|z|^2} dv(z) \right)^{1/p} \\ &= \left( \int_{Q_r(u)} |W^{1/p}(z) \mathbf{x}|^p e^{-\frac{p\alpha}{2}|z-u|^2} dv(z) \right)^{1/p} \end{aligned}$$

$$\begin{aligned}
&\geq e^{-\frac{n\alpha r^2}{4}} \left( \int_{Q_r(u)} |W^{1/p}(z)\mathbf{x}|^p dv(z) \right)^{1/p} \\
&= e^{-\frac{n\alpha r^2}{4}} r^{2n/p} \rho_{p,Q_r(u)}(\mathbf{x}).
\end{aligned} \tag{3.2}$$

Suppose now that  $1 < p < \infty$ . Then we have

$$\begin{aligned}
(\rho_{p',Q_r(u)}^*)^*(\mathbf{x}) &= \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle \mathbf{x}, \mathbf{y} \rangle|}{\rho_{p',Q_r(u)}^*(\mathbf{y})} \\
&= \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{\left| \left\langle \int_{Q_r(u)} f(\zeta) \overline{k_u^\alpha(\zeta)} d\lambda_\alpha(\zeta), \mathbf{y} \right\rangle \right|}{\left( \frac{1}{v(Q_r(u))} \int_{Q_r(u)} |W^{-1/p}(z)\mathbf{y}|^{p'} dv(z) \right)^{1/p'}} \\
&\geq r^{2n/p'} e^{-\frac{n\alpha r^2}{4}} \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{\left| \left( \frac{\alpha}{\pi} \right)^n \int_{Q_r(u)} \left\langle f(\zeta) \overline{k_u^\alpha(\zeta)}, \mathbf{y} \right\rangle e^{-\alpha|\zeta|^2} dv(\zeta) \right|}{\left( \int_{Q_r(u)} |W^{-1/p}(z)\mathbf{y}|^{p'} |k_u^\alpha(z)|^{p'} e^{-\frac{p'\alpha}{2}|z|^2} dv(z) \right)^{1/p'}} \\
&= \left( \frac{\alpha}{\pi} \right)^n r^{2n/p'} e^{-\frac{n\alpha r^2}{4}} \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle f, \chi_{Q_r(u)} k_u^\alpha \mathbf{y} \rangle_\alpha|}{\|\chi_{Q_r(u)} k_u^\alpha \mathbf{y}\|_{L_{\alpha, W^{-p'/p}}^{p'}(\mathbb{C}^n; \mathbb{C}^d)}},
\end{aligned}$$

which, in conjunction with Lemma 3.4, implies that

$$(\rho_{p',Q_r(u)}^*)^*(\mathbf{x}) \geq \left( \frac{\alpha}{\pi} \right)^{2n} r^{2n+2n/p'} e^{-\frac{5n\alpha r^2}{4}} \|P_\alpha\|^{-1} \|f\|_{L_{\alpha, W}^p(\mathbb{C}^n; \mathbb{C}^d)}.$$

Combining the above inequality with (3.2), we deduce that

$$(\rho_{p',Q_r(u)}^*)^*(\mathbf{x}) \geq \left( \frac{\alpha r^2}{\pi} \right)^{2n} e^{-2n\alpha r^2} \|P_\alpha\|^{-2} \rho_{p,Q_r(u)}(\mathbf{x}),$$

which is equivalent to

$$\rho_{p',Q_r(u)}^*(\mathbf{x}) \leq \left( \frac{\pi}{\alpha r^2} \right)^{2n} e^{2n\alpha r^2} \|P_\alpha\|^2 (\rho_{p,Q_r(u)})^*(\mathbf{x}).$$

Since  $u \in \mathbb{C}^n$  and  $\mathbf{x} \in \mathbb{C}^d$  are both arbitrary, we know that  $\rho$  is an  $\mathbf{A}_{p,r}$ -metric. Moreover,

$$\mathcal{C}_{p,r}(\rho) \leq \left( \frac{\pi}{\alpha r^2} \right)^{2n} e^{2n\alpha r^2} \|P_\alpha\|^2.$$

Suppose now that  $p = 1$ . Then similarly,

$$\begin{aligned}
(\rho_{\infty,Q_r(u)}^*)^*(\mathbf{x}) &= \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle \mathbf{x}, \mathbf{y} \rangle|}{\rho_{\infty,Q_r(u)}^*(\mathbf{y})} \\
&= \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{\left| \left\langle \int_{Q_r(u)} f(\zeta) \overline{k_u^\alpha(\zeta)} d\lambda_\alpha(\zeta), \mathbf{y} \right\rangle \right|}{\operatorname{ess\,sup}_{z \in Q_r(u)} |W^{-1}(z)\mathbf{y}|} \\
&\geq \left( \frac{\alpha}{\pi} \right)^n e^{-\frac{n\alpha r^2}{4}} \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle f, \chi_{Q_r(u)} k_u^\alpha \mathbf{y} \rangle_\alpha|}{\|\chi_{Q_r(u)} k_u^\alpha \mathbf{y}\|_{L_{\alpha, W^{-1}}^\infty(\mathbb{C}^n; \mathbb{C}^d)}} \\
&\geq \left( \frac{\alpha r}{\pi} \right)^{2n} e^{-\frac{5n\alpha r^2}{4}} \|P_\alpha\|^{-1} \|f\|_{L_{\alpha, W}^1(\mathbb{C}^n; \mathbb{C}^d)}
\end{aligned}$$

$$\geq \left( \frac{\alpha r^2}{\pi} \right)^{2n} e^{-2n\alpha r^2} \|P_\alpha\|^{-2} \rho_{1, Q_r(u)}(\mathbf{x}),$$

which finishes the proof.  $\square$

We now turn to the implication (d) $\implies$ (b) of Theorem 3.1. Before proceeding, we establish some estimates for  $\mathbf{A}_{p,r}$ -metrics. For a cube  $Q \subset \mathbb{C}^n$ , we use  $3Q$  to denote the cube with the same center but with side length  $3l(Q)$ .

**Lemma 3.5.** *Let  $r > 0$ ,  $1 \leq p < \infty$ , and let  $\rho$  be an  $\mathbf{A}_{p,3r}$ -metric. Then for any cube  $Q \subset \mathbb{C}^n$  with  $l(Q) = r$  and any  $\mathbf{x} \in \mathbb{C}^d$ ,*

$$\rho_{p,3Q}(\mathbf{x}) \leq 3^{2n(1-\frac{1}{p})} \mathcal{C}_{p,3r}(\rho) \rho_{p,Q}(\mathbf{x}).$$

*Proof.* The  $\mathbf{A}_{p,3r}$ -condition implies that

$$\rho_{p,3Q}(\mathbf{x}) = \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle \mathbf{x}, \mathbf{y} \rangle|}{(\rho_{p,3Q})^*(\mathbf{y})} \leq \mathcal{C}_{p,3r}(\rho) \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle \mathbf{x}, \mathbf{y} \rangle|}{\rho_{p',3Q}^*(\mathbf{y})}.$$

It is clear that  $\rho_{p',3Q}^*(\mathbf{y}) \geq 3^{2n(\frac{1}{p}-1)} \rho_{p',Q}^*(\mathbf{y})$ . Hence by Lemma 2.1,

$$\begin{aligned} \rho_{p,3Q}(\mathbf{x}) &\leq 3^{2n(1-\frac{1}{p})} \mathcal{C}_{p,3r}(\rho) \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle \mathbf{x}, \mathbf{y} \rangle|}{\rho_{p',Q}^*(\mathbf{y})} \\ &\leq 3^{2n(1-\frac{1}{p})} \mathcal{C}_{p,3r}(\rho) \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle \mathbf{x}, \mathbf{y} \rangle|}{(\rho_{p,Q})^*(\mathbf{y})} \\ &= 3^{2n(1-\frac{1}{p})} \mathcal{C}_{p,3r}(\rho) \rho_{p,Q}(\mathbf{x}), \end{aligned}$$

which completes the proof.  $\square$

For each  $r > 0$ , we will treat  $r\mathbb{Z}^{2n}$  as a subset of  $\mathbb{C}^n$  via the canonical identification between  $\mathbb{R}^{2n}$  and  $\mathbb{C}^n$ .

**Lemma 3.6.** *Let  $r > 0$ ,  $1 \leq p < \infty$ , and let  $\rho$  be an  $\mathbf{A}_{p,3r}$ -metric. Then for any  $\nu, \nu' \in r\mathbb{Z}^{2n}$ ,*

$$\rho_{p,Q_r(\nu)}(\mathbf{x}) \leq (3^{2n} \mathcal{C}_{p,3r}(\rho))^{\frac{\sqrt{2n}}{r}|\nu-\nu'|} \rho_{p,Q_r(\nu')}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{C}^d.$$

*In particular, the reducing operators satisfy*

$$\left\| \mathcal{R}_{Q_r(\nu)} \mathcal{R}_{Q_r(\nu')}^{-1} \right\| \leq \sqrt{d} (3^{2n} \mathcal{C}_{p,3r}(\rho))^{\frac{\sqrt{2n}}{r}|\nu-\nu'|}, \quad \nu, \nu' \in r\mathbb{Z}^{2n}.$$

*Proof.* Let  $\Gamma(\nu, \nu') = (a_0, a_1, \dots, a_k)$  be the discrete path in  $r\mathbb{Z}^{2n}$  from  $\nu$  to  $\nu'$  defined in [12]. Then  $a_0 = \nu$ ,  $a_k = \nu'$ ,  $k \leq \sqrt{2n}|\nu - \nu'|/r$ , and  $Q_r(a_{j-1}) \subset Q_{3r}(a_j)$  for each  $j \in \{1, \dots, k\}$ . Consequently, for  $\mathbf{x} \in \mathbb{C}^d$ , Lemma 3.5 yields that

$$\frac{\rho_{p,Q_r(\nu)}(\mathbf{x})}{\rho_{p,Q_r(\nu')}(\mathbf{x})} = \prod_{j=1}^k \frac{\rho_{p,Q_r(a_{j-1})}(\mathbf{x})}{\rho_{p,Q_r(a_j)}(\mathbf{x})} \leq \prod_{j=1}^k \frac{3^{2n/p} \rho_{p,Q_{3r}(a_j)}(\mathbf{x})}{\rho_{p,Q_r(a_j)}(\mathbf{x})} \leq (3^{2n} \mathcal{C}_{p,3r}(\rho))^{\frac{\sqrt{2n}}{r}|\nu-\nu'|}.$$

The second assertion is a direct consequence of the first one and (2.3).  $\square$

We now give the proof of the implication (d) $\implies$ (b) of Theorem 3.1.

*Proof of Theorem 3.1.* (d) $\implies$ (b). Suppose  $W \in \mathbf{A}_{p,r_0}$  and write  $r = r_0/3$ . Let  $\rho$  be the metric defined by

$$\rho_z(\mathbf{x}) = |W^{1/p}(z)\mathbf{x}|, \quad \mathbf{x} \in \mathbb{C}^d, \quad z \in \mathbb{C}^n.$$

Then  $\rho$  is an  $\mathbf{A}_{p,3r}$ -metric. The proof will be accomplished by a duality argument. To this end, choose  $f \in L_\alpha^p(\mathbb{C}^n; \mathbb{C}^d)$  and  $g \in L_\alpha^{p'}(\mathbb{C}^n; \mathbb{C})$ . Then

$$\begin{aligned} & \left| \langle P_{\alpha,W}^+(f), g \rangle_\alpha \right| \\ &= \left| \int_{\mathbb{C}^n} \int_{\mathbb{C}^n} |W^{1/p}(z)W^{-1/p}(u)f(u)| |K_z^\alpha(u)| d\lambda_\alpha(u) \overline{g(z)} e^{-\alpha|z|^2} dv(z) \right| \\ &\leq \left( \frac{\alpha}{\pi} \right)^n \int_{\mathbb{C}^n} \int_{\mathbb{C}^n} |W^{1/p}(z)W^{-1/p}(u)f(u)| |g(z)| e^{-\frac{\alpha}{2}|u|^2 - \frac{\alpha}{2}|z|^2} e^{-\frac{\alpha}{2}|z-u|^2} dv(u) dv(z) \\ &\lesssim \sum_{\nu, \nu' \in r\mathbb{Z}^{2n}} \left( e^{-\frac{\alpha}{4}|\nu-\nu'|^2} \times \right. \\ &\quad \left. \int_{Q_r(\nu)} \int_{Q_r(\nu')} |W^{1/p}(z)W^{-1/p}(u)f(u)| |g(z)| e^{-\frac{\alpha}{2}|u|^2 - \frac{\alpha}{2}|z|^2} dv(u) dv(z) \right) \\ &\leq \sum_{\nu, \nu' \in r\mathbb{Z}^{2n}} \left( e^{-\frac{\alpha}{4}|\nu-\nu'|^2} \int_{Q_r(\nu')} |\mathcal{R}_{Q_r(\nu')} W^{-1/p}(u)f(u)| e^{-\frac{\alpha}{2}|u|^2} dv(u) \times \right. \\ &\quad \left. \int_{Q_r(\nu)} \left\| W^{1/p}(z) \mathcal{R}_{Q_r(\nu')}^{-1} \right\| |g(z)| e^{-\frac{\alpha}{2}|z|^2} dv(z) \right), \end{aligned}$$

where  $\mathcal{R}_{Q_r(\nu')}$  is the reducing operator as in (2.3). We now separate into two cases:  $1 < p < \infty$  and  $p = 1$ .

**Case 1:**  $1 < p < \infty$ . Using Hölder's inequality, we obtain that

$$\begin{aligned} & \left| \langle P_{\alpha,W}^+(f), g \rangle_\alpha \right| \\ &\lesssim \left( \sum_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{4}|\nu-\nu'|^2} \left( \int_{Q_r(\nu')} \left\| \mathcal{R}_{Q_r(\nu')} W^{-1/p}(u) \right\| |f(u)| e^{-\frac{\alpha}{2}|u|^2} dv(u) \right)^p \right)^{1/p} \\ &\quad \times \left( \sum_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{4}|\nu-\nu'|^2} \left( \int_{Q_r(\nu)} \left\| W^{1/p}(z) \mathcal{R}_{Q_r(\nu')}^{-1} \right\| |g(z)| e^{-\frac{\alpha}{2}|z|^2} dv(z) \right)^{p'} \right)^{1/p'} \\ &=: \mathcal{S}_1(f)^{1/p} \cdot \mathcal{S}_2(g)^{1/p'}. \end{aligned} \tag{3.3}$$

For the term  $\mathcal{S}_1(f)$ , we have

$$\mathcal{S}_1(f) = \sum_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{4}|\nu-\nu'|^2} \left( \int_{Q_r(\nu')} \left\| \mathcal{R}_{Q_r(\nu')} W^{-1/p}(u) \right\| |f(u)| e^{-\frac{\alpha}{2}|u|^2} dv(u) \right)^p$$

$$\lesssim \sum_{\nu' \in r\mathbb{Z}^{2n}} \|\chi_{Q_r(\nu')} f\|_{L_\alpha^p(\mathbb{C}^n; \mathbb{C}^d)}^p \left( \int_{Q_r(\nu')} \|\mathcal{R}_{Q_r(\nu')} W^{-1/p}(u)\|^{p'} dv(u) \right)^{p/p'}. \quad (3.4)$$

Let  $\{\mathbf{e}_j\}_{1 \leq j \leq d}$  be the standard orthonormal basis of  $\mathbb{C}^d$ . Then it is easy to see that for any  $d \times d$  matrix  $M$ , the operator norm of  $M$  satisfies

$$\|M\| \leq d^{1/2} \max_{1 \leq j \leq d} |M\mathbf{e}_j|.$$

Therefore, for  $\nu' \in r\mathbb{Z}^{2n}$ , we can establish that

$$\begin{aligned} & \int_{Q_r(\nu')} \|\mathcal{R}_{Q_r(\nu')} W^{-1/p}(u)\|^{p'} dv(u) \\ &= \int_{Q_r(\nu')} \|W^{-1/p}(u) \mathcal{R}_{Q_r(\nu')}\|^{p'} dv(u) \\ &\leq \int_{Q_r(\nu')} \left( d^{1/2} \max_{1 \leq j \leq d} |W^{-1/p}(u) \mathcal{R}_{Q_r(\nu')} \mathbf{e}_j| \right)^{p'} dv(u) \\ &\leq d^{p'/2} \sum_{j=1}^d \int_{Q_r(\nu')} |W^{-1/p}(u) \mathcal{R}_{Q_r(\nu')} \mathbf{e}_j|^{p'} dv(u) \\ &\asymp d^{p'/2} \sum_{j=1}^d \left( \rho_{p', Q_r(\nu')}^* (\mathcal{R}_{Q_r(\nu')} \mathbf{e}_j) \right)^{p'}, \end{aligned} \quad (3.5)$$

which, in conjunction with (2.4) and (2.5), implies that

$$\begin{aligned} \int_{Q_r(\nu')} \|\mathcal{R}_{Q_r(\nu')} W^{-1/p}(u)\|^{p'} dv(u) &\lesssim d^{p'/2} \sum_{j=1}^d |\mathcal{R}_{Q_r(\nu')}^* \mathcal{R}_{Q_r(\nu')} \mathbf{e}_j|^{p'} \\ &\leq d^{\frac{p'}{2}+1} \|\mathcal{R}_{Q_r(\nu')}^* \mathcal{R}_{Q_r(\nu')}\|^{p'} \\ &\leq d^{\frac{3p'}{2}+1} \mathcal{C}_{p,r}(W)^{p'} \\ &\lesssim d^{\frac{3p'}{2}+1} \mathcal{C}_{p,r_0}(W)^{p'}. \end{aligned}$$

Combining this with (3.4), we establish that

$$\mathcal{S}_1(f) \lesssim d^{\frac{3p}{2}+\frac{p}{p'}} \mathcal{C}_{p,r_0}(W)^p \|f\|_{L_\alpha^p(\mathbb{C}^n; \mathbb{C}^d)}^p. \quad (3.6)$$

We now turn to the term  $\mathcal{S}_2(g)$ . Hölder's inequality yields that

$$\begin{aligned} \mathcal{S}_2(g) &= \sum_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{4}|\nu-\nu'|^2} \left( \int_{Q_r(\nu)} \|W^{1/p}(z) \mathcal{R}_{Q_r(\nu')}^{-1}\| |g(z)| e^{-\frac{\alpha}{2}|z|^2} dv(z) \right)^{p'} \\ &\leq \sum_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{4}|\nu-\nu'|^2} \|\chi_{Q_r(\nu)} g\|_{L_\alpha^{p'}(\mathbb{C}^n; \mathbb{C})}^{p'} \left( \int_{Q_r(\nu)} \|W^{1/p}(z) \mathcal{R}_{Q_r(\nu')}^{-1}\|^p dv(z) \right)^{\frac{p'}{p}}. \end{aligned} \quad (3.7)$$

For  $\nu, \nu' \in r\mathbb{Z}^{2n}$ , using the same method as in (3.5), we have

$$\int_{Q_r(\nu)} \left\| W^{1/p}(z) \mathcal{R}_{Q_r(\nu')}^{-1} \right\|^p dv(z) \lesssim d^{p/2} \sum_{j=1}^d \left( \rho_{p, Q_r(\nu)} \left( \mathcal{R}_{Q_r(\nu')}^{-1} \mathbf{e}_j \right) \right)^p,$$

which, together with (2.3) and Lemma 3.6, implies that

$$\begin{aligned} \int_{Q_r(\nu)} \left\| W^{1/p}(z) \mathcal{R}_{Q_r(\nu')}^{-1} \right\|^p dv(z) &\lesssim d^{\frac{p}{2}+1} \left\| \mathcal{R}_{Q_r(\nu)} \mathcal{R}_{Q_r(\nu')}^{-1} \right\|^p \\ &\leq d^{p+1} (3^{2n} \mathcal{C}_{p, 3r}(W))^{\frac{p\sqrt{2n}}{r} |\nu - \nu'|}. \end{aligned}$$

Note that for any  $c > 1$ ,

$$\begin{aligned} \sum_{\nu \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{4} |\nu|^2} c^{|\nu|} &\asymp \sum_{\nu \in r\mathbb{Z}^{2n}} \int_{Q_r(\nu)} e^{-\frac{\alpha}{4} |\nu|^2} c^{|\nu|} dv(z) \\ &\lesssim c^{\frac{\sqrt{2n}}{2} r} \int_{\mathbb{C}^n} e^{-\frac{\alpha}{8} |z|^2} c^{|z|} dv(z) \\ &\asymp c^{\frac{\sqrt{2n}}{2} r} \int_0^{+\infty} t^{2n-1} e^{-\frac{\alpha}{8} t^2 + t \log c} dt \\ &= c^{\frac{\sqrt{2n}}{2} r + \frac{2}{\alpha} \log c} \int_{-\frac{4}{\alpha} \log c}^{+\infty} \left( t + \frac{4}{\alpha} \log c \right)^{2n-1} e^{-\frac{\alpha}{8} t^2} dt \\ &\lesssim c^{\frac{\sqrt{2n}}{2} r + \frac{2}{\alpha} \log c} (1 + \log^{2n-1} c), \end{aligned}$$

where the implicit constants depend only on  $\alpha$ ,  $r$  and  $n$ . Consequently, for any  $\nu \in r\mathbb{Z}^{2n}$ ,

$$\begin{aligned} &\sum_{\nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{4} |\nu - \nu'|^2} \left( \int_{Q_r(\nu)} \left\| W^{1/p}(z) \mathcal{R}_{Q_r(\nu')}^{-1} \right\|^p dv(z) \right)^{p'/p} \\ &\lesssim \sum_{\nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{4} |\nu - \nu'|^2} d^{p' + \frac{p'}{p}} (3^{2n} \mathcal{C}_{p, 3r}(W))^{\frac{p'\sqrt{2n}}{r} |\nu - \nu'|} \\ &\lesssim d^{p' + \frac{p'}{p}} \mathcal{C}_{p, r_0}(W)^{p'n + \frac{144p'^2 n^2}{\alpha r_0^2} \log 3 + \frac{36p'^2 n}{\alpha r_0^2} \log \mathcal{C}_{p, r_0}(W)} (1 + \log^{2n-1} \mathcal{C}_{p, r_0}(W)), \end{aligned}$$

which, together with (3.7), yields that

$$\mathcal{S}_2(g) \lesssim d^{p' + \frac{p'}{p}} \mathcal{C}_{p, r_0}(W)^{p'n + \frac{144p'^2 n^2}{\alpha r_0^2} \log 3 + \frac{36p'^2 n}{\alpha r_0^2} \log \mathcal{C}_{p, r_0}(W)} (1 + \log^{2n-1} \mathcal{C}_{p, r_0}(W)) \|g\|_{L_{\alpha}^p(\mathbb{C}^n; \mathbb{C})}^{p'}.$$

Inserting this and (3.6) into (3.3), we conclude that  $P_{\alpha, W}^+ : L_{\alpha}^p(\mathbb{C}^n; \mathbb{C}^d) \rightarrow L_{\alpha}^p(\mathbb{C}^n; \mathbb{C})$  is bounded, and

$$\|P_{\alpha, W}^+\| \lesssim d^{7/2} \mathcal{C}_{p, r_0}(W)^{1+n + \frac{144p'n^2}{\alpha r_0^2} \log 3 + \frac{36p'n}{\alpha r_0^2} \log \mathcal{C}_{p, r_0}(W)} (1 + \log^{2n-1} \mathcal{C}_{p, r_0}(W))^{1/p'},$$

where the implicit constant depends only on  $\alpha$ ,  $p$ ,  $r_0$  and  $n$ .

**Case 2:**  $p = 1$ . We now have

$$\left| \langle P_{\alpha, W}^+(f), g \rangle_{\alpha} \right|$$

$$\begin{aligned}
&\lesssim \sum_{\nu, \nu' \in r\mathbb{Z}^{2n}} \left( e^{-\frac{\alpha}{4}|\nu-\nu'|^2} \int_{Q_r(\nu')} |\mathcal{R}_{Q_r(\nu')} W^{-1}(u) f(u)| e^{-\frac{\alpha}{2}|u|^2} dv(u) \times \right. \\
&\quad \left. \int_{Q_r(\nu)} \|W(z) \mathcal{R}_{Q_r(\nu')}^{-1}\| |g(z)| e^{-\frac{\alpha}{2}|z|^2} dv(z) \right) \\
&\leq \left( \sum_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{8}|\nu-\nu'|^2} \int_{Q_r(\nu')} \|\mathcal{R}_{Q_r(\nu')} W^{-1}(u)\| |f(u)| e^{-\frac{\alpha}{2}|u|^2} dv(u) \right) \times \\
&\quad \left( \sup_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{8}|\nu-\nu'|^2} \int_{Q_r(\nu)} \|W(z) \mathcal{R}_{Q_r(\nu')}^{-1}\| |g(z)| e^{-\frac{\alpha}{2}|z|^2} dv(z) \right) \\
&=: \mathcal{S}_3(f) \cdot \mathcal{S}_4(g).
\end{aligned}$$

For the term  $\mathcal{S}_3(f)$ , similarly as before,

$$\begin{aligned}
\mathcal{S}_3(f) &\lesssim \sum_{\nu' \in r\mathbb{Z}^{2n}} \int_{Q_r(\nu')} \|\mathcal{R}_{Q_r(\nu')} W^{-1}(u)\| |f(u)| e^{-\frac{\alpha}{2}|u|^2} dv(u) \\
&\leq \sum_{\nu' \in r\mathbb{Z}^{2n}} \operatorname{ess\,sup}_{u \in Q_r(\nu')} \|W^{-1}(u) \mathcal{R}_{Q_r(\nu')}\| \int_{Q_r(\nu')} |f(u)| e^{-\frac{\alpha}{2}|u|^2} dv(u) \\
&\leq d^{1/2} \sum_{\nu' \in r\mathbb{Z}^{2n}} \sum_{j=1}^d \rho_{\infty, Q_r(\nu')}^* (\mathcal{R}_{Q_r(\nu')} \mathbf{e}_j) \int_{Q_r(\nu')} |f(u)| e^{-\frac{\alpha}{2}|u|^2} dv(u) \\
&\leq d^{3/2} \sum_{\nu' \in r\mathbb{Z}^{2n}} \|\mathcal{R}_{Q_r(\nu')}^* \mathcal{R}_{Q_r(\nu')}\| \int_{Q_r(\nu')} |f(u)| e^{-\frac{\alpha}{2}|u|^2} dv(u) \\
&\leq d^{5/2} \mathcal{C}_{1,r}(W) \|f\|_{L_\alpha^1(\mathbb{C}^n; \mathbb{C}^d)} \\
&\lesssim d^{5/2} \mathcal{C}_{1,r_0}(W) \|f\|_{L_\alpha^1(\mathbb{C}^n; \mathbb{C}^d)}.
\end{aligned}$$

For the term  $\mathcal{S}_4(g)$ , we have that

$$\mathcal{S}_4(g) \leq \sup_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{8}|\nu-\nu'|^2} \int_{Q_r(\nu)} \|W(z) \mathcal{R}_{Q_r(\nu')}^{-1}\| dv(z) \|g\|_{L_\alpha^\infty(\mathbb{C}^n; \mathbb{C})}.$$

Noting that for  $c > 0$ ,  $\sup_{t \in \mathbb{R}} e^{-\frac{\alpha}{8}t^2} c^t = c^{\frac{2}{\alpha} \log c}$ , we may apply (2.3), Lemma 3.6 and the same method as in (3.5) to obtain that

$$\begin{aligned}
&\sup_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{8}|\nu-\nu'|^2} \int_{Q_r(\nu)} \|W(z) \mathcal{R}_{Q_r(\nu')}^{-1}\| dv(z) \\
&\lesssim d^{1/2} \sup_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{8}|\nu-\nu'|^2} \sum_{j=1}^d \rho_{1, Q_r(\nu)} \left( \mathcal{R}_{Q_r(\nu')}^{-1} \mathbf{e}_j \right) \\
&\leq d^{3/2} \sup_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{8}|\nu-\nu'|^2} \left\| \mathcal{R}_{Q_r(\nu)} \mathcal{R}_{Q_r(\nu')}^{-1} \right\| \\
&\leq d^2 \sup_{\nu, \nu' \in r\mathbb{Z}^{2n}} e^{-\frac{\alpha}{8}|\nu-\nu'|^2} \left( 3^{2n} \mathcal{C}_{1,3r}(W) \right)^{\frac{\sqrt{2n}}{r} |\nu-\nu'|}
\end{aligned}$$



$$\lesssim d^2 \mathcal{C}_{1,r_0}(W)^{\frac{144n^2}{\alpha r_0^2} \log 3 + \frac{36n}{\alpha r_0^2} \log \mathcal{C}_{1,r_0}(W)}.$$

Consequently,

$$\mathcal{S}_4(g) \lesssim d^2 \mathcal{C}_{1,r_0}(W)^{\frac{144n^2}{\alpha r_0^2} \log 3 + \frac{36n}{\alpha r_0^2} \log \mathcal{C}_{1,r_0}(W)} \|g\|_{L_\alpha^\infty(\mathbb{C}^n; \mathbb{C})}.$$

Therefore,  $P_{\alpha,W}^+ : L_\alpha^1(\mathbb{C}^n; \mathbb{C}^d) \rightarrow L_\alpha^1(\mathbb{C}^n; \mathbb{C})$  is bounded, and

$$\|P_{\alpha,W}^+\| \lesssim d^{9/2} \mathcal{C}_{1,r_0}(W)^{1 + \frac{144n^2}{\alpha r_0^2} \log 3 + \frac{36n}{\alpha r_0^2} \log \mathcal{C}_{1,r_0}(W)}$$

with implicit constant depends only on  $\alpha, r_0$  and  $n$ .  $\square$

As stated before, it follows from Theorem 3.1 that for each  $1 \leq p < \infty$ , all the classes of  $\mathbf{A}_{p,r}$ -weights coincide for  $r > 0$ . We end this paper by a quantitative description of this fact, which is independent of Theorem 3.1 and has its own interest.

**Proposition 3.7.** *Let  $1 \leq p < \infty$ ,  $0 < r_1 < r_2 < \infty$ , and let  $\rho : z \mapsto \rho_z$  be a metric on  $\mathbb{C}^n$ . Then  $\rho$  is an  $\mathbf{A}_{p,r_1}$ -metric if and only if it is an  $\mathbf{A}_{p,r_2}$ -metric. Moreover,*

$$\left(\frac{r_1}{r_2}\right)^{2n} \mathcal{C}_{p,r_1}(\rho) \leq \mathcal{C}_{p,r_2}(\rho) \leq 3^{4n^2(1+\frac{3r_2}{r_1})} d^{\frac{5}{2}} \left(2\sqrt{\frac{r_1}{r_2}} + 3\sqrt{\frac{r_2}{r_1}}\right)^{4n} \mathcal{C}_{p,r_1}(\rho)^{1+2n(1+\frac{3r_2}{r_1})}.$$

In particular, for each  $1 \leq p < \infty$ , all the classes of  $\mathbf{A}_{p,r}$ -metrics coincide for  $r > 0$ .

*Proof.* We only consider the case  $1 < p < \infty$ . The case  $p = 1$  follows from the same method. Suppose first that  $\rho$  is an  $\mathbf{A}_{p,r_2}$ -metric. Then for any  $z \in \mathbb{C}^n$  and  $\mathbf{x} \in \mathbb{C}^d$ , noting that

$$\begin{aligned} (\rho_{p,Q_{r_2}(z)})^*(\mathbf{x}) &= \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle \mathbf{x}, \mathbf{y} \rangle|}{\rho_{p,Q_{r_2}(z)}(\mathbf{y})} \\ &\leq \left(\frac{r_2}{r_1}\right)^{2n/p} \sup_{\mathbf{y} \in \mathbb{C}^d \setminus \{0\}} \frac{|\langle \mathbf{x}, \mathbf{y} \rangle|}{\rho_{p,Q_{r_1}(z)}(\mathbf{y})} \\ &= \left(\frac{r_2}{r_1}\right)^{2n/p} (\rho_{p,Q_{r_1}(z)})^*(\mathbf{x}), \end{aligned}$$

we establish that

$$\begin{aligned} \rho_{p',Q_{r_1}(z)}^*(\mathbf{x}) &\leq \left(\frac{r_2}{r_1}\right)^{2n/p'} \rho_{p',Q_{r_2}(z)}^*(\mathbf{x}) \\ &\leq \left(\frac{r_2}{r_1}\right)^{2n/p'} \mathcal{C}_{p,r_2}(\rho) (\rho_{p,Q_{r_2}(z)})^*(\mathbf{x}) \\ &\leq \left(\frac{r_2}{r_1}\right)^{2n} \mathcal{C}_{p,r_2}(\rho) (\rho_{p,Q_{r_1}(z)})^*(\mathbf{x}). \end{aligned}$$

Therefore,  $\rho$  is an  $\mathbf{A}_{p,r_1}$ -metric, and

$$\mathcal{C}_{p,r_1}(\rho) \leq \left(\frac{r_2}{r_1}\right)^{2n} \mathcal{C}_{p,r_2}(\rho). \quad (3.8)$$

Suppose now that  $\rho$  is an  $\mathbf{A}_{p,r_1}$ -metric. Fix  $z \in \mathbb{C}^n$ , and let

$$\Lambda = \left\{ \nu \in z + \frac{r_1}{3}\mathbb{Z}^{2n} : Q_{r_1/3}(\nu) \cap Q_{r_2}(z) \neq \emptyset \right\}.$$

We consider the reducing operators  $\mathcal{R}_Q$  and  $\mathcal{R}_Q^*$  of  $\rho_{p,Q}$  and  $\rho_{p',Q}^*$  respectively. For any  $\mathbf{x} \in \mathbb{C}^d$ , (2.3) implies that

$$\begin{aligned} |\mathcal{R}_{Q_{r_2}(z)}\mathbf{x}| &\leq \sqrt{d}\rho_{p,Q_{r_2}(z)}(\mathbf{x}) \\ &= \sqrt{d} \left( \frac{1}{v(Q_{r_2}(z))} \int_{Q_{r_2}(z)} (\rho_u(\mathbf{x}))^p dv(u) \right)^{1/p} \\ &\leq \sqrt{d} \left( \frac{1}{v(Q_{r_2}(z))} \sum_{\nu \in \Lambda} \int_{Q_{r_1/3}(\nu)} (\rho_u(\mathbf{x}))^p dv(u) \right)^{1/p} \\ &\leq \sqrt{d} \left( \frac{r_1}{3r_2} \right)^{2n/p} \sum_{\nu \in \Lambda} \rho_{p,Q_{r_1/3}(\nu)}(\mathbf{x}) \\ &\leq \sqrt{d} \left( \frac{r_1}{3r_2} \right)^{2n/p} \sum_{\nu \in \Lambda} |\mathcal{R}_{Q_{r_1/3}(\nu)}\mathbf{x}|. \end{aligned}$$

Similarly, (2.4) yields that

$$|\mathcal{R}_{Q_{r_2}(z)}^*\mathbf{x}| \leq \sqrt{d} \left( \frac{r_1}{3r_2} \right)^{2n/p'} \sum_{\nu \in \Lambda} |\mathcal{R}_{Q_{r_1/3}(\nu)}^*\mathbf{x}|.$$

Consequently,

$$\begin{aligned} \left\| \mathcal{R}_{Q_{r_2}(z)} \mathcal{R}_{Q_{r_2}(z)}^* \right\| &= \sup_{\mathbf{x} \in \mathbb{S}_d} |\mathcal{R}_{Q_{r_2}(z)} \mathcal{R}_{Q_{r_2}(z)}^* \mathbf{x}| \\ &\leq \sqrt{d} \left( \frac{r_1}{3r_2} \right)^{2n/p} \sum_{\nu \in \Lambda} \sup_{\mathbf{x} \in \mathbb{S}_d} |\mathcal{R}_{Q_{r_1/3}(\nu)} \mathcal{R}_{Q_{r_2}(z)}^* \mathbf{x}| \\ &= \sqrt{d} \left( \frac{r_1}{3r_2} \right)^{2n/p} \sum_{\nu \in \Lambda} \left\| \mathcal{R}_{Q_{r_1/3}(\nu)} \mathcal{R}_{Q_{r_2}(z)}^* \right\| \\ &= \sqrt{d} \left( \frac{r_1}{3r_2} \right)^{2n/p} \sum_{\nu \in \Lambda} \left\| \mathcal{R}_{Q_{r_2}(z)}^* \mathcal{R}_{Q_{r_1/3}(\nu)} \right\| \\ &= \sqrt{d} \left( \frac{r_1}{3r_2} \right)^{2n/p} \sum_{\nu \in \Lambda} \sup_{\mathbf{x} \in \mathbb{S}_d} |\mathcal{R}_{Q_{r_2}(z)}^* \mathcal{R}_{Q_{r_1/3}(\nu)} \mathbf{x}| \\ &\leq d \left( \frac{r_1}{3r_2} \right)^{2n} \sum_{\nu, \nu' \in \Lambda} \left\| \mathcal{R}_{Q_{r_1/3}(\nu')}^* \mathcal{R}_{Q_{r_1/3}(\nu)} \right\|, \quad (3.9) \end{aligned}$$

where  $\mathbb{S}_d$  is the unit sphere of  $\mathbb{C}^d$ . Since  $\rho$  is an  $\mathbf{A}_{p,r_1}$ -metric, we can use (3.8) to obtain that  $\mathcal{C}_{p,r_1/3}(\rho) \leq 3^{2n} \mathcal{C}_{p,r_1}(\rho)$ . Then it follows from (2.5) that

$$\sup_{u \in \mathbb{C}^n} \left\| \mathcal{R}_{Q_{r_1/3}(u)} \mathcal{R}_{Q_{r_1/3}(u)}^* \right\| \leq 3^{2n} d \mathcal{C}_{p,r_1}(\rho),$$

which implies that for any  $\nu' \in \Lambda$ ,

$$\left( \mathcal{R}_{Q_{r_1/3}(\nu')}^* \right)^2 \leq 3^{4n} d^2 \mathcal{C}_{p,r_1}(\rho)^2 \left( \mathcal{R}_{Q_{r_1/3}(\nu')}^{-1} \right)^2.$$

Combining this with Lemma 3.6 gives that for any  $\nu, \nu' \in \Lambda$ ,

$$\begin{aligned} \left\| \mathcal{R}_{Q_{r_1/3}(\nu')}^* \mathcal{R}_{Q_{r_1/3}(\nu)} \right\| &\leq 3^{2n} d \mathcal{C}_{p,r_1}(\rho) \left\| \mathcal{R}_{Q_{r_1/3}(\nu')}^{-1} \mathcal{R}_{Q_{r_1/3}(\nu)} \right\| \\ &\leq 3^{2n} d^{3/2} \mathcal{C}_{p,r_1}(\rho) \left( 3^{2n} \mathcal{C}_{p,r_1}(\rho) \right)^{\frac{3\sqrt{2n}}{r_1} |\nu - \nu'|}. \end{aligned}$$

Inserting the above estimate into (3.9) yields that

$$\left\| \mathcal{R}_{Q_{r_2}(z)} \mathcal{R}_{Q_{r_2}(z)}^* \right\| \leq 3^{2n} d^{5/2} \left( \frac{r_1}{3r_2} \right)^{2n} \mathcal{C}_{p,r_1}(\rho) \sum_{\nu, \nu' \in \Lambda} \left( 3^{2n} \mathcal{C}_{p,r_1}(\rho) \right)^{\frac{3\sqrt{2n}}{r_1} |\nu - \nu'|}.$$

It is easy to see that for any  $\nu, \nu' \in \Lambda$ ,  $|\nu - \nu'| \leq \sqrt{2n} \left( \frac{r_1}{3} + r_2 \right)$ , and the number  $\#\Lambda$  of elements in the set  $\Lambda$  satisfies  $\#\Lambda \leq \left( 2 + \frac{3r_2}{r_1} \right)^{2n}$ . Hence we can establish that

$$\left\| \mathcal{R}_{Q_{r_2}(z)} \mathcal{R}_{Q_{r_2}(z)}^* \right\| \leq 3^{4n^2 \left( 1 + \frac{3r_2}{r_1} \right)} d^{\frac{5}{2}} \left( 2\sqrt{\frac{r_1}{r_2}} + 3\sqrt{\frac{r_2}{r_1}} \right)^{4n} \mathcal{C}_{p,r_1}(\rho)^{1+2n \left( 1 + \frac{3r_2}{r_1} \right)}.$$

Since  $z \in \mathbb{C}^n$  is arbitrary, in view of (2.5), we conclude that  $\rho$  is an  $\mathbf{A}_{p,r_2}$ -metric, and

$$\mathcal{C}_{p,r_2}(\rho) \leq 3^{4n^2 \left( 1 + \frac{3r_2}{r_1} \right)} d^{\frac{5}{2}} \left( 2\sqrt{\frac{r_1}{r_2}} + 3\sqrt{\frac{r_2}{r_1}} \right)^{4n} \mathcal{C}_{p,r_1}(\rho)^{1+2n \left( 1 + \frac{3r_2}{r_1} \right)}.$$

The proof is complete.  $\square$

## REFERENCES

- [1] K. Bickel, S. Petermichl and B. D. Wick, Bounds for the Hilbert transform with matrix  $A_2$  weights, *J. Funct. Anal.* 270 (2016), no. 5, 1719–1743.
- [2] C. Cascante, J. Fàbrega and J. Á. Peláez, Littlewood–Paley formulas and Carleson measures for weighted Fock spaces induced by  $A_\infty$ -type weights, *Potential Anal.* 50 (2019), no. 2, 221–244.
- [3] J. Chen and M. Wang, Weighted norm inequalities, embedding theorems and integration operators on vector-valued Fock spaces, *Math Z.*, in press.
- [4] M. Christ and M. Goldberg, Vector  $A_2$  weights and a Hardy–Littlewood maximal function, *Trans. Amer. Math. Soc.* 353 (2001), no. 5, 1995–2002.
- [5] R. R. Coifman and C. Fefferman, Weighted norm inequalities for maximal functions and singular integrals, *Studia Math.* 51 (1974), 241–250.
- [6] D. Cruz-Uribe, J. Isralowitz, K. Moen, S. Pott and I. P. Rivera-Ríos, Weak endpoint bounds for matrix weights, *Rev. Mat. Iberoam.* 37 (2021), no. 4, 1513–1538.
- [7] M. Dostanić and K. Zhu, Integral operators induced by the Fock kernel, *Integr. Equ. Oper. Theory* 60 (2008), no. 2, 217–236.

- [8] M. Frazier and S. Roudenko, Matrix-weighted Besov spaces and conditions of  $A_p$  type for  $0 < p \leq 1$ , *Indiana Univ. Math. J.* 53 (2004), no. 5, 1225–1254.
- [9] T. A. Gillespie, S. Pott, S. Treil and A. Volberg, Logarithmic growth for weighted Hilbert transforms and vector Hankel operators, *J. Operator Theory* 52 (2004), no. 1, 103–112.
- [10] M. Goldberg, Matrix  $A_p$  weights via maximal functions, *Pacific J. Math.* 211 (2003), no. 2, 201–220.
- [11] R. Hunt, B. Muckenhoupt and R. Wheeden, Weighted norm inequalities for the conjugate function and Hilbert transform, *Trans. Amer. Math. Soc.* 176 (1973), 227–251.
- [12] J. Isralowitz, Schatten  $p$  class Hankel operators on the Segal–Bargmann space  $H^2(\mathbb{C}^n, d\mu)$  for  $0 < p < 1$ , *J. Operator Theory* 66 (2011), no. 1, 145–160.
- [13] J. Isralowitz, Invertible Toeplitz products, weighted norm inequalities, and  $A_p$  weights, *J. Operator Theory* 71 (2014), no. 2, 381–410.
- [14] M. Lauzon and S. Treil, Scalar and vector Muckenhoupt weights, *Indiana Univ. Math. J.* 56 (2007), no. 4, 1989–2015.
- [15] B. Muckenhoupt, Weighted norm inequalities for the Hardy maximal function, *Trans. Amer. Math. Soc.* 165 (1972), 207–226.
- [16] F. Nazarov, S. Petermichl, S. Treil and A. Volberg, Convex body domination and weighted estimates with matrix weights, *Adv. Math.* 318 (2017), 279–306.
- [17] F. L. Nazarov and S. R. Treil', The hunt for a Bellman function: applications to estimates for singular integral operators and to other classical problems of harmonic analysis, *Algebra i Analiz* 8 (1996), no. 5, 32–162 (Russian); translation in *St. Petersburg Math. J.* 8 (1997), no. 5, 721–824.
- [18] S. Roudenko, Matrix-weighted Besov spaces, *Trans. Amer. Math. Soc.* 355 (2003), no. 1, 273–314.
- [19] I. B. Simonenko, The Riemann boundary-value problem for  $n$  pairs of functions with measurable coefficients and its application to the study of singular integrals in weighted  $L_p$  spaces, *Dokl. Akad. Nauk SSSR* 141 (1961), 36–39.
- [20] S. Treil and A. Volberg, Wavelets and the angle between past and future, *J. Funct. Anal.* 143 (1997), no. 2, 269–308.
- [21] A. Volberg, Matrix  $A_p$  weights via  $S$ -functions, *J. Amer. Math. Soc.* 10 (1997), no. 2, 445–466.

JIALE CHEN, SCHOOL OF MATHEMATICS AND STATISTICS, SHAANXI NORMAL UNIVERSITY, XI'AN 710119, CHINA.

*Email address:* [jialechen@snnu.edu.cn](mailto:jialechen@snnu.edu.cn)

MAOFA WANG, SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN 430072, CHINA.

*Email address:* [mfwang.math@whu.edu.cn](mailto:mfwang.math@whu.edu.cn)