

Smolentsev N. K., Sokolova A. Yu.

Left-invariant Kähler and semi-para-Kähler structures on some six-dimensional unsolvable Lie groups

Abstract

In this article studies questions about the existence of left-invariant Kähler and semi-para-Kähler structures on six-dimensional unsolvable Lie groups whose Lie algebras are semidirect products. According to the classification results, there are four such Lie algebras. It is shown that one of these four Lie algebras admits left-invariant Kähler metrics, and the other three admit left-invariant semi-para-Kähler and semi-Kähler structures. The paper also shows that a six-dimensional symplectic Lie algebra must be solvable except in one case.

1 Introduction

Left-invariant structures on a Lie group are determined by their values on the Lie algebra. Therefore, when studying them, we usually deal with structures on the Lie algebra of a Lie group. In this sense, we will further refer, for example, about symplectic or Kähler Lie algebra, meaning a left-invariant symplectic or Kähler structure on the appropriate Lie group.

It is well known that solvable Lie algebras can admit symplectic structures [3], [5]. In the work of Chu B.-Y. [4] it was shown that a four-dimensional symplectic Lie algebra must be solvable. This is not the case in the six-dimensional case, and Chu gave an example of a six-dimensional symplectic but unsolvable Lie algebra. In the same work it was shown that semisimple Lie algebras cannot be symplectic. If a Lie algebra \mathfrak{g} is unsolvable, then it has a Lévy-Maltsev decomposition $\mathfrak{g} = N \ltimes S$ in the form of a direct sum of the radical N and a solvable subalgebra S . It was shown in [4] if a Lie algebra \mathfrak{g} has a Lévy-Malcev decomposition in the form of a direct product $\mathfrak{g} = N \times S$, then there are no symplectic structures on it. Thus, the question remains about the existence of symplectic structures only on unsolvable six-dimensional Lie algebras, for which the Lévy-Maltsev expansion is a semidirect product $\mathfrak{g} = N \ltimes S$.

According to the classification [2], [8] of six-dimensional unsolvable Lie algebras, which are semidirect products, there are four classes of such Lie algebras. These are the Lie algebras $A_{3.5} \ltimes sl(2, \mathbb{R})$, $A_{3.3} \ltimes sl(2, \mathbb{R})$, $A_{3.1} \ltimes sl(2, \mathbb{R})$ and $A_{3.1} \ltimes so(3)$. Here $A_{3.1}$, $A_{3.3}$, $A_{3.5}$ are three-dimensional solvable Lie algebras with basis $\{e_1, e_2, e_3\}$ and with the following non-zero Lie brackets: $A_{3.1}$ – commutative algebra; $A_{3.3}$: $[e_2, e_3] = e_1$; $A_{3.5}$: $[e_1, e_2] = e_2$, $[e_1, e_3] = e_3$. Thus, $A_{3.1} = \mathbb{R}^3$, $A_{3.3}$ is three-dimensional Heisenberg algebra \mathfrak{h}_3 and $A_{3.5}$ is a three-dimensional solvable and non-unimodular Lie algebra. The Lie algebras $so(3)$ and $sl(2, \mathbb{R})$ are the usual Lie algebras of skew-symmetric matrices of order 3 and matrices of order 2 with zero trace. On these Lie algebras we choose the usual basis $\{e_4, e_5, e_6\}$. Thus, there are [2] four six-dimensional undecidable Lie algebras, which are semidirect products. They are defined by the following Lie brackets:

1. $A_{3.5} \ltimes sl(2, \mathbb{R})$: $[e_1, e_2] = e_2$, $[e_1, e_3] = e_3$, $[e_4, e_5] = 2e_5$, $[e_4, e_6] = -2e_6$, $[e_5, e_6] = e_4$, $[e_2, e_4] = e_2$, $[e_2, e_5] = e_3$, $[e_3, e_4] = -e_3$, $[e_3, e_6] = e_2$.

2. $\mathfrak{h}_3 \ltimes sl(2, \mathbb{R})$: $[e_2, e_3] = e_1$, $[e_4, e_5] = 2e_5$, $[e_4, e_6] = -2e_6$, $[e_5, e_6] = e_4$, $[e_4, e_1] = e_1$, $[e_5, e_2] = e_1$, $[e_6, e_1] = e_2$, $[e_4, e_2] = -e_2$.
3. $\mathbb{R}^3 \ltimes sl(2, \mathbb{R})$: $[e_4, e_5] = 2e_5$, $[e_4, e_6] = -2e_6$, $[e_5, e_6] = e_4$, $[e_4, e_1] = 2e_1$, $[e_5, e_2] = 2e_1$, $[e_6, e_1] = e_2$, $[e_4, e_3] = -2e_3$, $[e_5, e_3] = e_2$, $[e_6, e_2] = 2e_3$.
4. $\mathbb{R}^3 \ltimes so(3)$: $[e_4, e_5] = e_6$, $[e_4, e_6] = -e_5$, $[e_5, e_6] = e_4$, $[e_4, e_2] = e_3$, $[e_5, e_1] = -e_3$, $[e_6, e_1] = e_2$, $[e_4, e_3] = -e_2$, $[e_5, e_3] = e_1$, $[e_6, e_2] = -e_1$.

Let G_1, G_2, G_3 and G_4 be Lie groups with Lie algebras of the list above.

The question of left-invariant symplectic structures on these Lie groups (i.e., symplectic structures of Lie algebras) was considered in the work of the authors [9]. It is shown that only one of the four Lie algebras admits symplectic structures. Thus, Chu's result on six-dimensional unsolvable Lie algebras can be refined as follows: *a six-dimensional symplectic Lie algebra must be solvable except in one case $A_{3,5} \ltimes sl(2, \mathbb{R})$* . Therefore, this exceptional unsolvable symplectic Lie algebra $A_{3,5} \ltimes sl(2, \mathbb{R})$ deserves more detailed study.

This work is an extension of the paper [9]. We will consider questions about the existence of left-invariant Kähler, semi-Kähler and semi-para-Kähler structures on the above four six-dimensional unsolvable Lie groups. It will be shown that only on one of the four Lie groups, namely on G_1 , there exist left-invariant Kähler structures and even with Einstein pseudo-Riemannian metrics. On the other three Lie groups there exist left-invariant semi-para-Kähler and semi-Kähler structures with integrable complex or paracomplex structures.

2 Preliminaries

Let us recall the basic concepts used in this work. An almost complex structure on a $2n$ -dimensional manifold M is a field J of endomorphisms $J : TM \rightarrow TM$ such that $J^2 = -Id$. The almost complex structure J is called integrable if the Nijenhuis tensor, defined by the equality $N_J(X, Y) = [JX, JY] - [X, Y] - J[JX, Y] - J[X, JY]$, vanishes, for any vector fields X, Y on M . In this case, J defines the structure of a complex manifold on M .

We will consider left-invariant almost complex structures on the Lie group G , which are definite by the left-invariant endomorphism field $J : TG \rightarrow TG$ of the tangent bundle TG . Since such a tensor J is determined by a linear operator on the Lie algebra $\mathfrak{g} = T_e G$, we will say that J is an invariant almost complex structure on the Lie algebra \mathfrak{g} . In this case, the integrability condition for J is formulated at the Lie algebra level: $N_J(X, Y) = 0$, for any $X, Y \in \mathfrak{g}$. In this case, we will say that J is a complex structure on the Lie algebra \mathfrak{g} .

A left-invariant symplectic structure ω on a Lie group G is definite by a closed 2-form of maximum rank. The closedness of the form ω is equivalent to the following condition on the Lie algebra \mathfrak{g} : $\omega([X, Y], Z) - \omega([X, Z], Y) + \omega([Y, Z], X) = 0$, $\forall X, Y, Z \in \mathfrak{g}$. In this case, the Lie algebra \mathfrak{g} will be called symplectic.

A left-invariant Kähler structure on a Lie group G is a triple (ω, J, g) consisting of a left-invariant symplectic form ω , a left-invariant complex structure J compatible with the form ω : $\omega(JX, JY) = \omega(X, Y)$, and the (pseudo)Riemannian metric g , defined by the formula $g(X, Y) = \omega(X, JY)$.

An almost paracomplex structure on a $2n$ -dimensional manifold M is the field P of endomorphisms of the tangent bundle TM such that $P^2 = Id$ and the ranks of the eigendistributions $T^\pm M := \ker(Id \mp P)$ are equal. The almost paracomplex structure P is called to be integrable

if the distributions $T^\pm M$ are involutive. In this case, P is called a paracomplex structure. The Nijenhuis tensor N_P of an almost paracomplex structure P is defined by the equality $N_P(X, Y) = [X, Y] + [PX, PY] - P[PX, Y] - P[X, PY]$, for any vector fields X, Y on M . As in the complex case, a para-complex structure P is integrable if and only if $N_P(X, Y) = 0$. A review of the theory of para-complex structures is presented in [1].

In the case of a left-invariant paracomplex structure P on a Lie group G , the eigenspaces subspaces $T_e^\pm G = \mathfrak{g}^\pm$ are subalgebras. Therefore, the paracomplex Lie algebra \mathfrak{g} can be represented as a direct sum of two subalgebras:

$$\mathfrak{g} = \mathfrak{g}^+ \oplus \mathfrak{g}^-.$$

The almost paracomplex structure P on a Lie group G is said to be bi-invariant if it is left-invariant and right-invariant simultaneously. This means that $Ad_g J = J$ for any element $g \in G$. For a connected Lie group, the last condition is equivalent to the following: $[ad_X, J] = 0$, $\forall X \in \mathfrak{g}$, or

$$[X, JY] = J[X, Y].$$

If an paracomplex structure P on a Lie algebra \mathfrak{g} is bi-invariant, then \mathfrak{g}^+ and \mathfrak{g}^- are ideals of \mathfrak{g} . The bi-invariant almost paracomplex structure on a Lie group G is always integrable.

A left-invariant para-Kähler structure on a Lie group G is a triple (ω, P, g) consisting of a left-invariant symplectic form ω , a left-invariant paracomplex structure P compatible with the form ω : $\omega(PX, PY) = -\omega(X, Y)$, and the (pseudo)Riemannian metric g , defined by the formula $g(X, Y) = \omega(X, PY)$.

On the last three Lie algebras of the list presented in the introduction, the condition $d\omega = 0$ leads to degeneracy of the form ω . At the same time, $d(\omega^3) = 3\omega \wedge \omega \wedge d\omega = 0$ for any 2-form ω on a six-dimensional manifold. An intermediate property would be $d(\omega^2) = 2\omega \wedge d\omega = 0$. Therefore, we reduce the closedness property $d\omega = 0$ and require that the following property hold:

$$\omega \wedge d\omega = 0.$$

Note that in the case of a left-invariant almost Hermitian structure on a Lie group G of dimension $2n$, the property $d(\omega^{n-1}) = 0$ of the fundamental form ω defines a class of semi-Kähler manifolds according to the Gray–Harvella classification [6], i.e. such that $\delta\omega = 0$. In our case of pseudo-Hermitian metrics on Lie groups, we will by analogy call such manifolds semi-Kähler.

Definition 2.1. *A left-invariant almost Hermitian (almost para-Hermitian) structure (G, g, ω, J) of dimension $2n$, whose fundamental form ω has the property $d(\omega^{n-1}) = 0$ is called semi-Kähler (semi-para-Kähler).*

The Lie algebras $\mathfrak{h}_3 \ltimes sl(2, \mathbb{R})$, $\mathbb{R}^3 \ltimes sl(2, \mathbb{R})$ and $\mathbb{R}^3 \ltimes so(3)$ do not have symplectic forms. For a non-degenerate 2-form ω its exterior derivative $d\omega$ is not equal to zero. In this case, $d\omega$ can be non-degenerate as a 3-form. The concept of non-degeneracy (stability) for a 3-form Ω on a six-dimensional space was defined in Hitchin's work [10]. For the 3-form Ω , Hitchin constructed a linear operator K_Ω whose square is proportional to the identity operator Id . Let us recall the basic constructions of Hitchin.

Let V be a 6-dimensional real vector space and μ a volume form on V . Let $\Omega \in \Lambda^3 V^*$ and $X \in V$, then $i_X \Omega \in \Lambda^2 V^*$ and $i_X \Omega \wedge \Omega \in \Lambda^5 V^*$. The natural pairing by the exterior product $V^* \otimes \Lambda^5 V^* \rightarrow \Lambda^6 V^* \cong \mathbb{R}\mu$ defines an isomorphism $A : \Lambda^5 V^* \cong V$ and using this, Hitchin defined the linear transformation $K_\Omega : V \rightarrow V$ by the following formula:

$$K_\Omega(X) = A(i_X \Omega \wedge \Omega).$$

In other words, $i_{K_\Omega(X)}\mu = i_X\Omega \wedge \Omega$. The operator K_Ω has the following properties: $\text{trace}(K_\Omega) = 0$ and $K_\Omega^2 = \lambda(\Omega)Id$. Therefore, in the case $\lambda(\Omega) < 0$, we obtain the structure J_Ω of a complex vector space on the space V :

$$J_\Omega = \frac{1}{\sqrt{-\lambda(\Omega)}}K_\Omega,$$

and if $\lambda(\Omega) > 0$, then we obtain the para-complex structure P_Ω , i.e., $P_\Omega^2 = Id$, $P_\Omega \neq 1$ by a similar formula:

$$P_\Omega = \frac{1}{\sqrt{\lambda(\Omega)}}K_\Omega.$$

Thus, the operator $K_{d\omega}$ can define either an almost complex or almost paracomplex structure on six-dimensional Lie algebra \mathfrak{g} when $d\omega$ is non-degenerate and $\mu = \omega^n$.

Let ∇ be the Levi-Civita connection corresponding to the (pseudo)Riemannian metric g . It is determined from a six-term formula [7], which for left-invariant vector fields X, Y, Z on a Lie group takes the form: $2g(\nabla_X Y, Z) = g([X, Y], Z) + g([Z, X], Y) + g(X, [Z, Y])$. The curvature tensor is defined by the formula $R(X, Y) = [\nabla_X, \nabla_Y] - \nabla_{[X, Y]}$. The Ricci tensor $Ric(X, Y)$ is the convolution of the curvature tensor along the first and fourth (upper) indices. The Ricci operator RIC is defined by the formula $Ric(X, Y) = g(RIC(X), Y)$. To calculate the geometric characteristics of left-invariant structures on Lie groups, the Maple system was used.

3 Left-invariant structures on some unsolvable Lie groups

In this section we consider left-invariant geometric structures on each of the four six-dimensional unsolvable Lie groups whose Lie algebras are semidirect products. Recall that left-invariant geometric structures on a Lie group are determined by their values on the Lie algebra; therefore, when studying them, only the Lie algebra of the Lie group is usually used. In this sense, we will refer, for example, about a symplectic or Kähler Lie algebra, meaning a left-invariant symplectic or Kähler structure on the corresponding Lie group.

3.1 Left-invariant Kähler structures on the group G_1 .

The group G_1 has a Lie algebra $A_{3.5} \ltimes sl(2, \mathbb{R})$ with commutation relations: $[e_1, e_2] = e_2$, $[e_1, e_3] = e_3$, $[e_4, e_5] = 2e_5$, $[e_4, e_6] = -2e_6$, $[e_5, e_6] = e_4$, $[e_2, e_4] = e_2$, $[e_2, e_5] = e_3$, $[e_3, e_4] = -e_3$, $[e_3, e_6] = e_2$. This Lie algebra is considered by Chu [4] as an example showing that a six-dimensional symplectic Lie algebra need not be solvable. It consists of matrices of the form:

$$\begin{pmatrix} a_1 & a_2 & a_3 \\ 0 & a_4 & a_5 \\ 0 & a_6 & -a_4 \end{pmatrix}$$

The general form of a symplectic 2-form ω on the Lie algebra $A_{3.5} \ltimes sl(2, \mathbb{R})$ was obtained in [9]:

$$\omega = \begin{pmatrix} 0 & w_{12} & w_{13} & 0 & 0 & 0 \\ -w_{12} & 0 & 0 & w_{12} & w_{13} & 0 \\ -w_{13} & 0 & 0 & -w_{13} & 0 & w_{12} \\ 0 & -w_{12} & w_{13} & 0 & w_{45} & w_{46} \\ 0 & -w_{13} & 0 & -w_{45} & 0 & w_{56} \\ 0 & 0 & -w_{12} & -w_{46} & -w_{56} & 0 \end{pmatrix} \quad (1)$$

with the condition of non-degeneracy, $\det(\omega) = w_{12}^4 w_{45}^2 - 4w_{12}^3 w_{13} w_{45} w_{56} + 2w_{12}^2 w_{13}^2 w_{45} w_{46} + 4w_{12}^2 w_{13}^2 w_{56}^2 - 4w_{12} w_{13}^3 w_{46} w_{56} + w_{13}^4 w_{46}^2 \neq 0$.

It is practically impossible to find a complex structure in a general form compatible with (1). Therefore, we will use special cases where the symplectic 2-form has the simplest form. From the expression $\det(\omega)$ we see that w_{12} and w_{13} cannot be equal to zero at the same time.

Let us first consider the first case, when $w_{12} = 1$ and $w_{13} = 0$. Then $\det(\omega) = w_{12}^4 w_{45}^2$ does not depend on w_{46} and w_{56} . Therefore, we will consider them equal to zero, $w_{46} = 0$ and $w_{56} = 0$. Let, in addition, $w_{45} = -1$.

In the second case, we will make a similar choice: $w_{12} = 0$ and $w_{13} = 1$, $w_{56} = 0$ and $w_{45} = 0$, $w_{46} = 1$.

3.1.1 Case 1.

Let us consider the symplectic structure represented by in Chu's work [4]:

$$\omega_1 = e^1 \wedge e^2 + e^2 \wedge e^4 + e^3 \wedge e^6 - e^4 \wedge e^5. \quad (2)$$

Let us find complex structures J compatible with 2-form (2). To do this, we take a matrix of almost complex structure $J = (J_i^j)$ and require the compatible condition $\omega_1(JX, JY) = \omega_1(X, Y)$, the condition $J^2 = -Id$ and the integrability condition $N_J(X, Y) = [JX, JY] - [X, Y] - J[X, JY] - J[JX, Y] = 0$. The following system of equations is obtained for finding the compatible complex structure J :

$$\begin{cases} \omega_{kj} J_i^k + \omega_{ik} J_j^k = 0, \\ J_k^i J_j^k = -\delta_j^i, \\ J_i^l J_j^m C_{lm}^k - J_i^l J_m^k C_{lj}^m - J_j^l J_m^k C_{il}^m - C_{ij}^k = 0, \end{cases} \quad (3)$$

where δ_j^i is the identity matrix, C_{ij}^k are the structure constants of the Lie algebra, and the indices vary from 1 to 6.

This system of algebraic equations is solved using Maple symbolic calculations. First, we solve the simplest compatible condition, then the integrability condition, and finally we check the condition $J^2 = -Id$.

For the Lie algebra $A_{3.5} \ltimes sl(2, \mathbb{R})$, we obtain six solutions to the system of equations (3). For each case, the associated pseudo-Riemannian metric is defined by the formula $g_{1,J}(X, Y) = \omega_1(X, JY)$. As a result, each solution represents a Kähler structure $(\omega_1, J, g_{1,J})$. Moreover, among all the solutions there is one Kähler structure with an Einstein metric g_1 of scalar curvature $S = 12$. Such a complex structure and the metric tensor are represented by the following matrices:

$$J_1 = \begin{pmatrix} 0 & 1 & 0 & 0 & -\frac{1}{4} & -1 \\ -1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & -\frac{1}{4} & -1 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{4} & 1 & 0 & 0 \end{pmatrix}, \quad g_1 = \begin{pmatrix} -1 & 0 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{4} & 1 & 0 & 0 \\ 1 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\frac{1}{4} & -1 \\ 0 & 0 & 0 & 0 & -1 & 0 \end{pmatrix}. \quad (4)$$

3.1.2 Case 2.

Let us consider the second special case of the symplectic forms (1):

$$\omega_2 = e^1 \wedge e^3 + e^2 \wedge e^5 - e^3 \wedge e^4 + e^4 \wedge e^6. \quad (5)$$

Calculations show that there are five families of complex structures J compatible with form (5). Each solution represents a Kähler structure $(\omega_2, J, g_{2,J})$. Moreover, among all the solutions there is one Kähler structure $(\omega, J_2, g_{2,J})$ with an Einstein metric g_2 of nonzero scalar curvature $S = -12a$, depending on one parameter. This complex structure and metric tensor are represented by the following matrices:

$$J_2 = \begin{pmatrix} 0 & 0 & -a & 0 & \frac{1}{a} & \frac{a}{4} \\ 0 & 0 & 0 & 0 & 0 & -\frac{1}{a} \\ \frac{1}{a} & 0 & 0 & \frac{1}{a} & 0 & 0 \\ 0 & 0 & 0 & 0 & -\frac{1}{a} & -\frac{a}{4} \\ 0 & -\frac{a^3}{4} & 0 & a & 0 & 0 \\ 0 & a & 0 & 0 & 0 & 0 \end{pmatrix}, \quad g_2 = \begin{pmatrix} \frac{1}{a} & 0 & 0 & \frac{1}{a} & 0 & 0 \\ 0 & -\frac{a^3}{4} & 0 & a & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 \\ \frac{1}{a} & a & 0 & \frac{1}{a} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{a} \\ 0 & 0 & 0 & 0 & \frac{1}{a} & \frac{a}{4} \end{pmatrix}. \quad (6)$$

3.1.3 Para-Hermitian structure.

The Lie algebra $A_{3.5} \ltimes sl(2, \mathbb{R})$ has an (integrable) paracomplex structure P_0 corresponding to the semidirect product of subalgebras $\mathfrak{g}^+ = A_{3.5}$ and $\mathfrak{g}^- = sl(2, \mathbb{R})$. It has a diagonal matrix $P_0 = \text{diag}\{1, 1, 1, -1, -1, -1\}$. In addition, on the Lie algebra $A_{3.5} \ltimes sl(2, \mathbb{R})$ there is a 2-form $\Omega_0 = e^1 \wedge e^4 + e^2 \wedge e^5 + e^3 \wedge e^6$, also corresponding to the decomposition into a semidirect product (here e^1, \dots, e^6 is the dual basis). However, the form Ω_0 is not semi-Kähler. Therefore, on the Lie algebra $A_{3.5} \ltimes sl(2, \mathbb{R})$ the para-Hermitian structure (Ω_0, P_0, G_0) is defined, where the pseudo-Riemannian metric g_0 is defined by the formula $G_0(X, Y) = \Omega_0(X, P_0 Y)$. This metric has the following Ricci tensor

$$Ric = -e^1 \odot e^1 - \frac{1}{2}e^5 \odot e^5 - \frac{1}{2}e^6 \odot e^6 - 2e^2 \odot e^5 + 2e^3 \odot e^6 - 2e^4 \odot e^3 - 5e^5 \odot e^6$$

and zero scalar curvature. Recall that $e^i \odot e^j = \frac{1}{2}(e^i \otimes e^j + e^j \otimes e^i)$ is a symmetric product.

Let us formulate the results obtained in the form of a theorem.

Theorem 3.1. *The Lie group G_1 with the Lie algebra $A_{3.5} \ltimes sl(2, \mathbb{R})$ admits a multiparameter family of left-invariant symplectic structures (1). The group G_1 admits left-invariant Kähler structures, including those that have Einstein metrics (4) and (6). The group G_1 also has a natural left-invariant para-Hermitian structure of zero scalar curvature.*

3.2 Left-invariant Kähler structures on the group G_2 .

The group G_2 has a Lie algebra $\mathfrak{h}_3 \ltimes sl(2, \mathbb{R})$ with commutation relations: $[e_2, e_3] = e_1$, $[e_4, e_5] = 2e_5$, $[e_4, e_6] = -2e_6$, $[e_5, e_6] = e_4$, $[e_4, e_1] = e_1$, $[e_5, e_2] = e_1$, $[e_6, e_1] = e_2$, $[e_4, e_2] = -e_2$.

There are no left-invariant symplectic structures on this Lie algebra; the closedness condition $d\omega = 0$ implies the degeneracy of the 2-form ω . It's easy to show. Let $\omega = \omega_{ij}e^i \wedge e^j$ be an arbitrary 2-form. The components of the $d\omega$ have the following expressions through structure constants:

$$d\omega_{ijk} = -C_{ij}^s \omega_{sk} + C_{ik}^s \omega_{sj} - C_{jk}^s \omega_{si}.$$

From the condition $d\omega_{ijk} = 0$ we have, in particular, $d\omega_{134} = -C_{13}^s \omega_{s4} + C_{14}^s \omega_{s3} - C_{34}^s \omega_{s1} = 0$. From the commutation relations we obtain $C_{13}^s = 0$, $C_{34}^s = 0$ and $C_{14}^1 = -1$. Therefore we have: $d\omega_{134} = -\omega_{13} = 0$. The following equalities are obtained in exactly the same way: $\omega_{23} = 0$, $\omega_{34} = 0$, $\omega_{35} = 0$, $\omega_{36} = 0$, from which it follows that the form ω is degenerate.

3.2.1 Semi-para-Kähler structure corresponding to the semidirect product.

On the Lie algebra $\mathfrak{h}_3 \ltimes sl(2, \mathbb{R})$ there is a paracomplex structure P_0 corresponding to the semidirect product of subalgebras: $\mathfrak{g}^+ = \mathfrak{h}_3$ and $\mathfrak{g}^- = sl(2, \mathbb{R})$. It has a diagonal matrix $P_0 = \text{diag}\{1, 1, 1, -1, -1, -1\}$. In addition, the Lie algebra $\mathfrak{h}_3 \ltimes sl(2, \mathbb{R})$ has a natural 2-form $\Omega = ae^1 \wedge e^4 + be^2 \wedge e^5 + ce^3 \wedge e^6$, also corresponding to the semidirect product. It is easy to see that it is compatible with the operator P_0 and is semi-Kähler under the condition $a = -b$. Let us choose the two simplest semi-Kähler 2-forms:

$$\Omega_{01} = -e^1 \wedge e^4 + e^2 \wedge e^5 + e^3 \wedge e^6 \quad \text{and} \quad \Omega_{02} = e^1 \wedge e^4 - e^2 \wedge e^5 + e^3 \wedge e^6. \quad (7)$$

Thus, on the Lie algebra $\mathfrak{h}_3 \ltimes sl(2, \mathbb{R})$ two semi-para-Kähler structures $(\Omega_{0i}, P_0, G_{0i})$, $i = 1, 2$, are defined, whose pseudo-Riemannian metrics $G_{0i}(X, Y) = \Omega_{0i}(X, P_0 Y)$ have scalar curvatures $S_i = -2$ and Ricci tensors:

$$Ric_1 = e^1 \odot e^4 + e^2 \odot e^5 + 2e^3 \odot e^6 - 4e^4 \odot e^4 - 8e^5 \odot e^6,$$

$$Ric_2 = -e^1 \odot e^4 - e^2 \odot e^5 + 2e^3 \odot e^6 - 4e^4 \odot e^4 - 8e^5 \odot e^6.$$

Both 2-forms Ω_{0i} are not only semi-Kähler, but they also have exterior derivatives that are non-degenerate in the Hitchin sense. The operators of almost paracomplex structures corresponding to non-degenerate 3-forms $d\Omega_{01}$ and $d\Omega_{02}$ is:

$$P_{01} = -e_1 \otimes e^1 + e_2 \otimes e^2 + e_3 \otimes e^3 + e_4 \otimes e^4 - e_5 \otimes e^5 - e_6 \otimes e^6 + 4e_1 \otimes e^4 - 4e_2 \otimes e^6 - 4e_3 \otimes e^5,$$

$$P_{02} = -e_1 \otimes e^1 + e_2 \otimes e^2 + e_3 \otimes e^3 + e_4 \otimes e^4 - e_5 \otimes e^5 - e_6 \otimes e^6 - 4e_1 \otimes e^4 + 4e_2 \otimes e^6 - 4e_3 \otimes e^5,$$

Unfortunately, P_{01} and P_{02} are not compatible with the corresponding forms Ω_{01} and Ω_{02} and therefore do not generate semi-para-Kähler structures.

3.2.2 General semi-para-Kähler structures.

There are no closed non-degenerate 2-forms on the Lie algebra $\mathfrak{h}_3 \ltimes sl(2, \mathbb{R})$. Therefore, we will consider semi-Kähler 2-forms, i.e. non-degenerate forms ω and satisfying the condition $\omega \wedge d\omega = 0$.

Let $\omega = \omega_{ij}e^i \wedge e^j$ be an arbitrary 2-form. The condition $\omega \wedge d\omega = 0$ is satisfied for the following parameter values:

$$\omega_{13}\omega_{46} - \omega_{14}\omega_{36} - \omega_{14}\omega_{56} + \omega_{15}\omega_{46} + \omega_{16}\omega_{34} - \omega_{16}\omega_{45} + \omega_{23}\omega_{56} - \omega_{25}\omega_{36} + \omega_{26}\omega_{35} = 0,$$

$$\omega_{12}\omega_{35} - \omega_{13}\omega_{25} + \omega_{15}\omega_{23} = 0, \quad \omega_{12}\omega_{34} - \omega_{13}\omega_{24} + \omega_{14}\omega_{23} = 0,$$

$$-\omega_{12}\omega_{36} + \omega_{13}\omega_{26} - \omega_{16}\omega_{23} = 0,$$

$$-\omega_{13}\omega_{56} + \omega_{15}\omega_{36} - \omega_{16}\omega_{35} - \omega_{23}\omega_{45} + \omega_{24}\omega_{35} - \omega_{25}\omega_{34} = 0.$$

There are many solutions to the system of these equations. Only in eight cases is the 3-form $d\omega$ non-degenerate in the sense of Hitchin. In each of these 8 cases, the 3-form $d\omega$ corresponds to the operator $P_{d\omega}$ of a (nonintegrable) almost paracomplex structure. Note that the expressions for the 2-forms ω and the corresponding operators $P_{d\omega}$ are quite large. Therefore, we consider only those semi-Kähler 2-forms ω with non-degenerate $d\omega$ that are compatible with the paracomplex structure P_0 , i.e. such that $\omega(P_0 X, P_0 Y) = -\omega(X, Y)$. Then we get 5 semi-Kähler 2-forms:

$$\omega_1 = e^1 \wedge \left(\frac{\omega_{16}\omega_{35} - \omega_{24}\omega_{35}}{\omega_{36}} e^5 + \omega_{16} e^6 \right) + e^2 \wedge \omega_{24} e^4 + e^3 \wedge (\omega_{35} e^5 + \omega_{36} e^6), \quad (8)$$

$$\begin{aligned} \omega_2 = e^1 \wedge \left(-\frac{\omega_{25}\omega_{36} - \omega_{26}\omega_{35}}{\omega_{36}} e^4 - \frac{\omega_{24}\omega_{35} - \omega_{25}\omega_{34}}{\omega_{36}} e^5 \right) + e^2 \wedge (\omega_{24}e^4 + \omega_{25}e^5 + \omega_{26}e^6) + \\ + e^3 \wedge (\omega_{34}e^4 + \omega_{35}e^5 + \omega_{36}e^6) + \frac{-\omega_{36}(\omega_{25}\omega_{36} - \omega_{26}\omega_{35}) + \omega_{25}\omega_{36}^2 - \omega_{26}\omega_{35}\omega_{36}}{\omega_{25}\omega_{36} - \omega_{26}\omega_{35}} e^5 \wedge e^6, \end{aligned} \quad (9)$$

$$\omega_3 = \frac{-\omega_{24}\omega_{35} + \omega_{25}\omega_{34}}{\omega_{36}} e^1 \wedge e^5 + e^2 \wedge (\omega_{24}e^4 + \omega_{25}e^5 + \frac{\omega_{25}\omega_{36}}{\omega_{35}} e^6) + e^3 \wedge (\omega_{34}e^4 + \omega_{35}e^5 + \omega_{36}e^6), \quad (10)$$

$$\omega_4 = e^1 \wedge (\omega_{14}e^4 + \omega_{15}e^5 + \omega_{24}e^6) + e^2 \wedge (\omega_{24}e^4 + \omega_{25}e^5) + e^3 \wedge \omega_{35}e^5, \quad (11)$$

$$\omega_5 = e^1 \wedge (\omega_{14}e^4 + \omega_{15}e^5) + e^2 \wedge (\omega_{24}e^4 + \omega_{26}e^6) + e^3 \wedge \omega_{34}e^4. \quad (12)$$

In each case we obtain a semi-para-Kähler structure (ω, P_0, g_ω) with the associated metric $g_\omega(X, Y) = \omega(X, P_0 Y)$. Using the Maple system, the geometric characteristics of the g_ω metric can be easily calculated. In particular, for the last form ω_5 the following expressions for the metric tensor and scalar curvature hold:

$$g_5 = \begin{pmatrix} 0 & 0 & 0 & -\omega_{14} & -\omega_{15} & 0 \\ 0 & 0 & 0 & -\omega_{24} & 0 & -\omega_{26} \\ 0 & 0 & 0 & -\omega_{34} & 0 & 0 \\ -\omega_{14} & -\omega_{24} & -\omega_{34} & 0 & 0 & 0 \\ -\omega_{15} & 0 & 0 & 0 & 0 & 0 \\ 0 & -\omega_{26} & 0 & 0 & 0 & 0 \end{pmatrix}, \quad S_5 = \frac{\omega_{14}^2 + \omega_{15}\omega_{24}}{\omega_{34}\omega_{15}\omega_{26}}.$$

As already noted, each of the above 2-forms has the non-degenerate exterior derivative $d\omega$ in the Hitchin sense and, therefore, has the operator $P_{d\omega}$ of almost paracomplex structure. Let us give the expression for the operator P_5 corresponding to the 3-form $d\omega_5$:

$$\begin{aligned} P_5(e_1) = -e_1, \quad P_5(e_2) = \frac{-2\omega_{14}\omega_{24} - 6\omega_{15}\omega_{26}}{\omega_{14}^2 - \omega_{15}\omega_{24}} e_1 + e_2, \quad P_5(e_3) = \frac{-2\omega_{14}\omega_{34}}{\omega_{14}^2 - \omega_{15}\omega_{24}} e_1 + e_3, \\ P_5(e_4) = \frac{2\omega_{14}\omega_{24} + 18\omega_{15}\omega_{26}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_3 + \frac{\omega_{14}^2 - \omega_{15}\omega_{24}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_4 + \frac{2\omega_{14}\omega_{24}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_5 + \frac{6\omega_{14}\omega_{15}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_6, \\ P_5(e_5) = \frac{2\omega_{14}\omega_{34}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_1 + \frac{6\omega_{15}\omega_{34}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_2 + \frac{-2\omega_{14}^2 - 6\omega_{15}\omega_{24}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_3 + \\ + \frac{2\omega_{14}\omega_{15}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_4 + \frac{-\omega_{14}^2 + \omega_{15}\omega_{24}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_5 + \frac{6\omega_{15}^2}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_6, \\ P_5(e_6) = \frac{6\omega_{26}\omega_{34}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_1 + \frac{-2\omega_{24}\omega_{34}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_2 + \frac{2\omega_{24}^2 - 6\omega_{14}\omega_{26}}{\omega_{14}^2 + \omega_{15}\omega_{24}} e_3 - e_6. \end{aligned}$$

Let us formulate the results obtained in the form of a theorem.

Theorem 3.2. *The group G_2 with Lie algebra $\mathfrak{h}_3 \ltimes sl(2, \mathbb{R})$ does not admit left-invariant symplectic structures. The group G_2 has natural left-invariant semi-para-Kähler structures $(\Omega_{0i}, P_0, G_{0i})$, $i = 1, 2$, of negative scalar curvature, where the semi-Kähler 2-forms Ω_{0i} are represented by formulas (7), $P_0 = \text{diag} 1, 1, 1, -1, -1, -1$, and $G_{0i}(X, Y) = \Omega_{0i}(X, P_0 Y)$. The group G_2 also admits multiparameter families (8) – (12) of left-invariant semi-Kähler 2-forms ω compatible with the paracomplex structure operator P_0 and, therefore, it admits multiparameter families of semi-paraKähler structures (ω, P_0, g_ω) with associated metrics $g_\omega(X, Y) = \omega(X, P_0 Y)$.*

3.3 Left-invariant semi-para-Kähler structures on the group G_3 .

The group G_3 has a Lie algebra $\mathbb{R}^3 \ltimes sl(2, \mathbb{R})$ with commutation relations: $[e_4, e_5] = 2e_5$, $[e_4, e_6] = -2e_6$, $[e_5, e_6] = e_4$, $[e_4, e_1] = 2e_1$, $[e_5, e_2] = 2e_1$, $[e_6, e_1] = e_2$, $[e_4, e_3] = -2e_3$, $[e_5, e_3] = e_2$, $[e_6, e_2] = 2e_3$. There are no left-invariant symplectic structures on this Lie algebra; the closedness condition $d\omega = 0$ implies the degeneracy of the 2-form ω . This is shown just as simply as in the previous section.

3.3.1 Semi-para-Kähler structure corresponding to a semidirect product.

On the considering Lie algebra there is a paracomplex structure P_0 corresponding to the semidirect product of the subalgebras $\mathfrak{g}^+ = \mathbb{R}^3$ and $\mathfrak{g}^- = sl(2, \mathbb{R})$. It has a diagonal matrix $P_0 = \text{diag}\{1, 1, 1, -1, -1, -1\}$. In addition, the Lie algebra $\mathbb{R}^3 \ltimes sl(2, \mathbb{R})$ has a natural 2-form $\Omega = ae^1 \wedge e^4 + be^2 \wedge e^5 + ce^3 \wedge e^6$, corresponding to the semidirect product. It is easy to see that it is compatible with the operator P_0 and is semi-Kähler under the condition $a = -b$. Let us choose the two simplest semi-Kähler 2-forms:

$$\Omega_{01} = -e^1 \wedge e^4 + e^2 \wedge e^5 + e^3 \wedge e^6 \quad \text{and} \quad \Omega_{02} = e^1 \wedge e^4 - e^2 \wedge e^5 + e^3 \wedge e^6,$$

Then two semi-para-Kähler structures $(\Omega_{0i}, P_0, G_{0i})$, $i = 1, 2$, are defined on the Lie algebra $\mathbb{R}^3 \ltimes sl(2, \mathbb{R})$. Their pseudo-Riemannian metrics $G_{0i}(X, Y) = \Omega_{0i}(X, P_0 Y)$ have zero scalar curvature and the following Ricci tensors:

$$Ric_i = -16e^4 \odot e^4 - 16e^5 \odot e^6.$$

Both 2-forms Ω_{0i} have degenerate exterior derivative $d\Omega_{0i}$ in the Hitchin sense.

3.3.2 General semi-para-Kähler structures.

There are no closed non-degenerate 2-forms on this Lie algebra. Let $\omega = \omega_{ij}e^i \wedge e^j$ be an arbitrary 2-form. The condition $\omega \wedge d\omega = 0$ is satisfied for the following parameter values:

$$\begin{aligned} w_{12}w_{35} - w_{13}w_{25} + w_{15}w_{23} &= 0, & w_{12}w_{34} - w_{13}w_{24} + w_{14}w_{23} &= 0, \\ -w_{12}w_{56} - w_{13}w_{45} + w_{14}w_{35} + w_{15}w_{26} - w_{15}w_{34} - w_{16}w_{25} &= 0, \\ w_{13}w_{46} - w_{14}w_{36} + w_{16}w_{34} + w_{23}w_{56} - w_{25}w_{36} + w_{26}w_{35} &= 0, \\ w_{12}w_{46} - w_{14}w_{26} + w_{16}w_{24} - w_{23}w_{45} + w_{24}w_{35} - w_{25}w_{34} &= 0, \\ -w_{12}w_{36} + w_{13}w_{26} - w_{16}w_{23} &= 0. \end{aligned}$$

There are 2 solutions to this system of equations with non-degenerate 2-forms ω :

$$\begin{aligned} \omega_1 = e^1 \wedge (-w_{25}e^4 + w_{15}e^5 - w_{35}e^6) + e^2 \wedge (w_{24}e^4 + w_{25}e^5 + w_{26}e^6) + \\ e^3 \wedge (w_{26}e^4 + w_{35}e^5 - w_{36}e^6) + e^4 \wedge (w_{45}e^5 + w_{46}e^6) + w_{56}e^5 \wedge e^6, \end{aligned}$$

$$\begin{aligned} \omega_2 = e^1 \wedge (w_{14}e^4 + w_{15}e^5 - w_{35}e^6) + e^2 \wedge (w_{24}e^4 - w_{14}e^5 + w_{34}e^6) + \\ e^3 \wedge (w_{34}e^4 + w_{35}e^5) + e^4 \wedge (w_{45}e^5 + w_{46}e^6) + w_{56}e^5 \wedge e^6. \end{aligned}$$

Both of these 2-forms have a degenerate exterior derivative $d\omega_i$, $i = 1, 2$.

The Lie algebra $\mathbb{R}^3 \ltimes sl(2, \mathbb{R})$ has the integrable paracomplex structure $P_0 = \text{diag}\{1, 1, 1, -1, -1, -1\}$, corresponding to the decomposition of the Lie algebra into a

semidirect product. Let us require that the above semi-Kähler 2-forms ω_i be compatible with the operator P_0 : $\omega_i(P_0X, P_0Y) = -\omega_i(X, Y)$. Then,

$$\omega_1 = e^1 \wedge (-\omega_{25}e^4 + \omega_{15}e^5 - \omega_{35}e^6) + e^2 \wedge (\omega_{24}e^4 + \omega_{25}e^5 + \omega_{26}e^6) + e^3 \wedge (\omega_{26}e^4 + \omega_{35}e^5 - \omega_{36}e^6), \quad (13)$$

$$\omega_2 = e^1 \wedge (\omega_{14}e^4 + \omega_{15}e^5 - \omega_{35}e^6) + e^2 \wedge (\omega_{24}e^4 - \omega_{14}e^5 + \omega_{34}e^6) + e^3 \wedge (\omega_{34}e^4 + \omega_{35}e^5). \quad (14)$$

Let us define the associated metric by the formula $g_i(X, Y) = \omega_i(X, P_0Y)$, $i = 1, 2$. Calculations show that every semi-para-Kähler structure (ω_i, P_0, g_i) has zero scalar curvature.

Let us formulate the results obtained in the form of a theorem.

Theorem 3.3. *The group G_3 with Lie algebra $\mathbb{R}^3 \ltimes sl(2, \mathbb{R})$ does not admit left-invariant symplectic structures. The group G_3 has natural left-invariant semi-para-Kähler structures $(\Omega_{0i}, P_0, G_{0i})$, $i = 1, 2$, of zero scalar curvature, where the semi-Kähler 2-forms Ω_{0i} are represented by formulas (7), $P_0 = \text{diag}\{1, 1, 1, -1, -1, -1\}$, and $G_{0i}(X, Y) = \Omega_{0i}(X, P_0Y)$. The group G_3 also admits multiparameter families (13) – (14) of left-invariant semi-Kähler 2-forms ω_i compatible with the paracomplex structure operator P_0 and, therefore, it admits multiparameter families of left-invariant semi-paraKähler structures (ω_i, P_0, g_i) with associated metrics $g_i(X, Y) = \omega_i(X, P_0Y)$ zero scalar curvature.*

3.4 Left-invariant semi-Kähler and semi-para-Kähler structures on the group G_4 .

The group G_4 has a Lie algebra $\mathbb{R}^3 \ltimes so(3)$ with commutation relations: $[e_4, e_5] = e_6$, $[e_4, e_6] = -e_5$, $[e_5, e_6] = e_4$, $[e_4, e_2] = e_3$, $[e_5, e_1] = -e_3$, $[e_6, e_1] = e_2$, $[e_4, e_3] = -e_2$, $[e_5, e_3] = e_1$, $[e_6, e_2] = -e_1$.

This Lie algebra has a paracomplex structure P_0 corresponding to the semidirect product of the subalgebras \mathbb{R}^3 and $so(3)$. It has a diagonal matrix $P_0 = \text{diag}\{1, 1, 1, -1, -1, -1\}$. In addition, on the Lie algebra $\mathbb{R}^3 \ltimes so(3)$ there is a natural 2-form $\Omega = ae^1 \wedge e^4 + be^2 \wedge e^5 + ce^3 \wedge e^6$, also corresponding to the semidirect product. It is easy to see that it is compatible with the operator P_0 and is semi-Kähler for any non-zero values of the parameters a, b, c . Therefore, on the Lie algebra $\mathbb{R}^3 \ltimes so(3)$ the semi-para-Kähler structure (Ω, P_0, g) is defined whose pseudo-Riemannian metric $G(X, Y) = \Omega(X, P_0Y)$ has the Ricci tensor of the form

$$Ric = \frac{a^2 - (b - c)^2}{bc} e^4 \odot e^4 - \frac{b^2 + (a - c)^2}{ab} e^5 \odot e^5 + \frac{c^2 - (a - b)^2}{ac} e^6 \odot e^6$$

and zero scalar curvature.

3.4.1 General semi-para-Kähler structures.

There are no closed non-degenerate 2-forms on this Lie algebra. Let $\omega = \omega_{ij}e^i \wedge e^j$ be an arbitrary 2-form. The condition $\omega \wedge d\omega = 0$ is satisfied for the following parameter values:

$$\begin{aligned} -w_{13}w_{56} + w_{15}w_{36} - w_{16}w_{35} + w_{23}w_{46} - w_{24}w_{36} + w_{26}w_{34} &= 0, \\ w_{12}w_{36} - w_{13}w_{26} + w_{16}w_{23} &= 0, \quad w_{12}w_{34} - w_{13}w_{24} + w_{14}w_{23} = 0, \\ w_{12}w_{56} - w_{15}w_{26} + w_{16}w_{25} - w_{23}w_{45} + w_{24}w_{35} - w_{25}w_{34} &= 0, \quad -w_{12}w_{35} + w_{13}w_{25} - w_{15}w_{23} = 0. \end{aligned}$$

There are 2 solutions to this system of equations with non-degenerate 2-forms ω :

$$\begin{aligned} \omega_1 = e^1 \wedge (w_{14}e^4 + w_{24}e^5 + w_{34}e^6) + e^2 \wedge (w_{24}e^4 + w_{25}e^5 + w_{26}e^6) + \\ e^3 \wedge (w_{34}e^4 + w_{26}e^5 + w_{36}e^6) + e^4 \wedge (w_{45}e^5 + w_{46}e^6) + w_{56}e^5 \wedge e^6, \end{aligned}$$

$$\begin{aligned}\omega_2 = e^1 \wedge (w_{14}e^4 + w_{15}e^5 + w_{34}e^6) + e^2 \wedge (w_{15}e^4 + w_{25}e^5 + w_{35}e^6) + \\ e^3 \wedge (w_{34}e^4 + w_{35}e^5) + e^4 \wedge (w_{45}e^5 + w_{46}e^6) + w_{56}e^5 \wedge e^6.\end{aligned}$$

Let us require that these semi-Kähler 2-forms ω_i be compatible with the operator P_0 : $\omega_i(P_0X, P_0Y) = -\omega_i(X, Y)$. Then they take the form:

$$\omega_1 = e^1 \wedge (\omega_{14}e^4 + \omega_{24}e^5 + \omega_{34}e^6) + e^2 \wedge (\omega_{24}e^4 + \omega_{25}e^5 + \omega_{26}e^6) + e^3 \wedge (\omega_{34}e^4 + \omega_{26}e^5 + \omega_{36}e^6), \quad (15)$$

$$\omega_2 = e^1 \wedge (w_{14}e^4 + w_{15}e^5 + w_{34}e^6) + e^2 \wedge (w_{15}e^4 + w_{25}e^5 + w_{35}e^6) + e^3 \wedge (w_{34}e^4 + w_{35}e^5). \quad (16)$$

Let us define the associated metric by the formula $g_i(X, Y) = \omega_i(X, P_0Y)$, $i = 1, 2$. Calculations show that every semi-para-Kähler structure (ω_i, P_0, g_i) has zero scalar curvature.

3.4.2 Semi-Kähler structures.

The Lie algebra $\mathbb{R}^3 \ltimes so(3)$ admits integrable complex structures compatible with the para-Kähler 2-form:

$$\Omega_{01} = -e^1 \wedge e^4 + e^2 \wedge e^5 + e^3 \wedge e^6. \quad (17)$$

Let us find the matrix $J = (\psi_{ij})$ of this complex structure.

The compatible condition $\Omega_{01}(JX, JY) = \Omega_{01}(X, Y)$ is satisfied for the following values of parameters ψ_{ij} : $\psi_{51} = -\psi_{42}$, $\psi_{61} = -\psi_{43}$, $\psi_{11} = -\psi_{44}$, $\psi_{21} = \psi_{45}$, $\psi_{31} = \psi_{46}$, $\psi_{62} = \psi_{53}$, $\psi_{12} = \psi_{54}$, $\psi_{22} = -\psi_{55}$, $\psi_{32} = -\psi_{56}$, $\psi_{13} = \psi_{64}$, $\psi_{23} = -\psi_{65}$, $\psi_{33} = -\psi_{66}$, $\psi_{24} = -\psi_{15}$, $\psi_{34} = -\psi_{16}$, $\psi_{35} = \psi_{26}$.

Then from the other two equations of system (3) we obtain 4, up to the sign $(\pm J)$ of the solution:

1. $\psi_{15} = 0$, $\psi_{16} = (\psi_{45}^2 + \psi_{46}^2)/(\psi_{41}\psi_{45})$, $\psi_{25} = \psi_{45}^2/\psi_{41}$, $\psi_{26} = (\psi_{45}\psi_{46})/\psi_{41}$, $\psi_{36} = \psi_{46}^2/\psi_{41}$, $\psi_{42} = 0$, $\psi_{43} = 0$, $\psi_{44} = -\psi_{46}/\psi_{45}$, $\psi_{52} = 0$, $\psi_{53} = 0$, $\psi_{54} = 0$, $\psi_{55} = 0$, $\psi_{56} = -1$, $\psi_{64} = 0$, $\psi_{65} = 1$, $\psi_{66} = 0$, $\psi_{14} = -(\psi_{45}^2 + \psi_{46}^2)/(\psi_{41}\psi_{45})$.
2. $\psi_{15} = ((\psi_{45}\psi_{44} + \psi_{46})\psi_{14})/(\psi_{44}^2 + 1)$, $\psi_{16} = (\psi_{15}(\psi_{46}\psi_{44} - \psi_{45}))/(\psi_{45}\psi_{44} + \psi_{46})$, $\psi_{25} = -(\psi_{15}\psi_{45}^2)/(\psi_{45}\psi_{44} + \psi_{46})$, $\psi_{26} = -(\psi_{15}\psi_{45}\psi_{46})/(\psi_{45}\psi_{44} + \psi_{46})$, $\psi_{36} = -(\psi_{15}\psi_{45}^2)/(\psi_{45}\psi_{44} + \psi_{46})$, $\psi_{41} = -(\psi_{45}\psi_{44} + \psi_{46})/\psi_{15}$, $\psi_{42} = 0$, $\psi_{43} = 0$, $\psi_{52} = 0$, $\psi_{53} = 0$, $\psi_{54} = 0$, $\psi_{55} = 0$, $\psi_{56} = -1$, $\psi_{63} = 0$, $\psi_{64} = 0$, $\psi_{65} = 1$, $\psi_{66} = 0$.
3. $\psi_{14} = -(\psi_{16}^2\psi_{41}^2 + \psi_{46}^2)/(\psi_{41}\psi_{46})$, $\psi_{15} = \psi_{46}/\psi_{41}$, $\psi_{25} = 0$, $\psi_{26} = 0$, $\psi_{36} = (\psi_{46}^2)/\psi_{41}$, $\psi_{42} = 0$, $\psi_{43} = 0$, $\psi_{44} = -(\psi_{16}\psi_{41})/\psi_{46}$, $\psi_{45} = 0$, $\psi_{52} = 0$, $\psi_{53} = 0$, $\psi_{54} = 0$, $\psi_{55} = 0$, $\psi_{56} = 1$, $\psi_{63} = 0$, $\psi_{64} = 0$, $\psi_{65} = -1$, $\psi_{66} = 0$.
4. $\psi_{14} = -(\psi_{44}^2 + 1)/\psi_{41}$, $\psi_{15} = 0$, $\psi_{16} = 0$, $\psi_{25} = 0$, $\psi_{26} = 0$, $\psi_{36} = 0$, $\psi_{42} = 0$, $\psi_{43} = 0$, $\psi_{45} = 0$, $\psi_{46} = 0$, $\psi_{52} = 0$, $\psi_{53} = 0$, $\psi_{54} = 0$, $\psi_{55} = 0$, $\psi_{56} = -1$, $\psi_{63} = 0$, $\psi_{64} = 0$, $\psi_{65} = 1$, $\psi_{66} = 0$.

Each of the complex structures J of this list defines a semi-Kähler structure (Ω_{01}, J, g_J) , where $g_J(X, Y) = \Omega_{01}(X, JY)$.

We present explicit expressions for the fourth complex structure, the associated metric, the Ricci tensor and scalar curvature:

$$J_4 = \begin{pmatrix} -\psi_{44} & 0 & 0 & -\frac{\psi_{44}^2+1}{\psi_{41}} & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ \psi_{41} & 0 & 0 & \psi_{44}^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, \quad g_{J_4} = \begin{pmatrix} -\psi_{41} & 0 & 0 & -\psi_{44} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ -\psi_{44} & 0 & 0 & \frac{\psi_{44}^2+1}{\psi_{41}} & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Ric_4 = \begin{pmatrix} -\psi_{41}^2 & 0 & 0 & -\psi_{41}\psi_{44} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \psi_{41}/2 \\ 0 & 0 & 0 & 0 & -\psi_{41}/2 & 0 \\ -\psi_{41}\psi_{44} & 0 & 0 & -\psi_{44}^2 & 0 & 0 \\ 0 & 0 & -\psi_{41}/2 & 0 & 2 & 0 \\ 0 & \psi_{41}/2 & 0 & 0 & 0 & 2 \end{pmatrix}, \quad S_4 = \psi_{41}.$$

Let us formulate the results obtained in the following form.

Theorem 3.4. *The group G_4 with the Lie algebra $\mathbb{R}^3 \ltimes so(3)$ does not admit left-invariant symplectic structures. The group G_4 has natural left-invariant semi-para-Kähler structures (Ω, P_0, G) , with zero scalar curvature, where $\Omega = ae^1 \wedge e^4 + be^2 \wedge e^5 + ce^3 \wedge e^6$ and $P_0 = \text{diag}\{1, 1, 1, -1, -1, -1\}$. The Lie group G_4 also admits multiparameter families (15), (16) of left-invariant semi-Kähler 2-forms ω_i , $i = 1, 2$, compatible with the paracomplex structure P_0 and, therefore, it admits multiparameter families of left-invariant semi-paraKähler structures (ω_i, P_0, g_i) with associated metrics $g_i(X, Y) = \omega_i(X, P_0 Y)$ of zero scalar curvature. The group G_4 admits four multiparameter families of left-invariant complex structures J compatible with the semi-Kähler 2-form (17) and, therefore, it admits multiparameter families of left-invariant semi-Kähler structures (Ω_{01}, J, g_J) with integrable complex structures J , where $g_J(X, Y) = \Omega_{01}(X, JY)$.*

References

- [1] Alekseevsky D.V., Medori C., Tomassini A. Homogeneous para-Kähler Einstein manifolds. Russ. Math. Surv. 2009, Vol. 64, no. 1, pp. 1–43.
- [2] Basarab-Horwath P., Lahno V., Zhdanov R. The structure of Lie algebras and the classification problem for partial differential equations. Acta Appl. Math. 2001. Vol. 69, pp. 43–94.
- [3] Campoamor-Stursberg R. Symplectic forms on six-dimensional real solvable Lie algebras I. Algebra Colloquium. 2009. Vol. 16, no. 2, pp. 253–266.
- [4] Chu Bon-Yao. Symplectic homogeneous spaces. Trans. of the Amer. Math. Soc. 1974. Vol. 197, pp. 154–159.
- [5] Goze M., Khakimdjano Y., Medina A. Symplectic or contact structures on Lie groups. Differential Geom. Appl. 2004, Vol. 21, no. 1, 41–54.

- [6] Gray A., Harvella L.M. The sixteen classes of almost Hermitian manifolds and their linear Invariants. *Ann. Math. Pura Appl.* 1980. Vol. 123, pp. 35–58.
- [7] Kobayashi S. and Nomizu K. *Foundations of Differential Geometry*, Vol. 1 and 2. Interscience Publ. New York, London. 1963.
- [8] Turkowski P. Low-dimensional real Lie algebras. *J. Math. Phys.* 1988. Vol. 29, pp. 2139–2144.
- [9] Smolentsev N.K., Sokolova A.Yu. Parakähler and para-Hermitian structures on six-dimensional unsolvable Lie algebras. *Izvestiya AltGU. Matematika i mekhanika.* 2023. Vol. 4(132), pp. 94–98.
- [10] Hitchin N.J. The geometry of three-forms in six dimensions *J. Diff. Geom.* 2000. Vol. 55, pp. 547–576.