AN ALTERNATIVE FOR CONSTANT MEAN CURVATURE HYPERSURFACES

LIAM MAZUROWSKI AND XIN ZHOU

ABSTRACT. Let M^{n+1} be a closed manifold of dimension $3 \le n+1 \le 7$ equipped with a generic Riemannian metric g. Let c be a positive number. We show that, either there exist infinitely many distinct closed hypersurfaces with constant mean curvature equal to c, or there exist infinitely many distinct closed hypersurfaces with constant mean curvature less than c but enclosing half the volume of M.

1. INTRODUCTION

Let (M, g) be a closed Riemmanian manifold. Let \mathcal{M} denote the moduli space of all smooth, closed, almost-embedded constant mean curvature hypersurfaces in M. Note that we do not fix the topology of elements in \mathcal{M} , and that the value of the mean curvature is allowed to vary between different hypersurfaces in \mathcal{M} . In this short note, we contribute to the study of \mathcal{M} by proving the following alternative for constant mean curvature hypersurfaces.

Theorem 1.1. Let M^{n+1} be a closed manifold of dimension $3 \le n+1 \le 7$. Let g be a generic Riemannian metric on M. Then, for every constant c > 0, at least one of the following two statements is true:

- (i) There exist infinitely many distinct smooth, closed, almost-embedded hypersurfaces in M with constant mean curvature equal to c;
- (ii) There exist infinitely many distinct smooth, closed, almost-embedded hypersurfaces in M with constant mean curvature less than c but enclosing half the volume of M.

The proof of Theorem 1.1 is based on the authors' recently developed min-max theory [12] for finding critical points of functionals of the form $\Omega \mapsto \operatorname{Area}(\partial \Omega) + f(\operatorname{Vol}(\Omega))$. Here $\Omega \subset M$ is an open set with sufficiently nice boundary, and f is an arbitrary smooth function.

1.1. Background and Motivation. In this subsection, we briefly survey some of the known results and open problems concerning the space \mathcal{M} . It is known that for a generic metric g on \mathcal{M} , the moduli space \mathcal{M} of constant mean curvature hypersurfaces has the structure of a 1-dimensional manifold; see for example [13]. This manifold may have both compact and non-compact components.

The finer structure of \mathcal{M} is best understood in the very small mean curvature and very large mean curvature regimes. Regarding surfaces with large mean curvature, Ye [20] proved that there exist foliations by CMC spheres in small neighborhoods of non-degenerate critical points of the scalar curvature. If the scalar curvature function R is Morse, this implies that \mathcal{M} contains at least $\mu(M)$ spheres with constant mean curvature c when c is large enough. Here $\mu(M)$ is the Morse number of M, i.e., the minimum number of critical points of a Morse function on M. Later, Pacard and Xu [15] proved that even when R is not Morse, M still contains at least cat(M) spheres with constant mean curvature c when c is large enough. Here cat(M) is the Lusternik-Schnirelmann category of M. In general, these CMC spheres do not exhaust the high mean curvature portion of \mathcal{M} . For example, Mazzeo and Pacard [13] have constructed CMCs with large mean curvature by perturbing small tubes around geodesics. We still do not have a complete understanding of the space of all CMC hypersurfaces with large mean curvature. In fact, even if one restricts the topology, there is not yet a complete description of the space of CMC spheres with large mean curvature. For instance, Pacard claims in joint work with Malchiodi [14] to be able to construct CMCs which look like a connect sum of two small spheres joined by a Delaunay tube.

In the small mean curvature regime, constant mean curvature surfaces can be found by perturbing minimal surfaces. CMCs with small mean curvature can also be constructed by doubling minimal surfaces; see [10] for a min-max construction of such doublings. Dey [5] proved that the number of CMC hypersurfaces with mean curvature c goes to infinity at a quantitative rate as $c \to 0$. Again a complete description of the space of CMC hypersurfaces with small mean curvature still seems out of reach.

For general values of c > 0, the second named author and J. Zhu [23] proved that M always contains at least one almost-embedded hypersurface with constant mean curvature c. It is conjectured that there should always be a second almost-embedded hypersurfaces with constant mean curvature c.

Conjecture 1.2 (Twin Bubble Conjecture [22]). Let (M^{n+1}, g) be a closed Riemannian manifold of dimension $3 \le n+1 \le 7$. Then, for every constant c > 0, there are at least two distinct closed, almost-embedded hypersurfaces with constant mean curvature c in M.

This conjecture is motivated in part by a famous conjecture of Arnold on the existence of closed magnetic geodesics in Riemannian 2-spheres [3]. Arnold's conjecture states that every Riemannian 2-sphere (S^2, g) should contain at least two closed curves with constant geodesic curvature κ for every constant $\kappa > 0$. Schneider [17] has proved Arnold's conjecture when the metric satisfies certain positive curvature conditions, but the full conjecture remains open.

Remark 1.3. It is not entirely clear that the number two is optimal in the twin bubble conjecture. In fact, to the authors' knowledge, it is not known whether there exist a closed manifold M and a constant c > 0 such that M contains only finitely many closed hypersurfaces with constant mean curvature c.

The present work grew out of the authors' efforts to understand the Twin Bubble Conjecture. Theorem 1.1 shows that, at least for generic metrics, if the Twin Bubble Conjecture fails for a particular constant c > 0, then M must contain infinitely many constant mean curvature hypersurfaces with mean curvature less than c, each enclosing half the volume of M. In particular, this suggests that a thorough understanding of the space of half-volume constant mean curvature hypersurfaces in M may be helpful for constructing hypersurfaces with prescribed constant mean curvature. The work of the authors [11][12] shows that a closed manifold M^{n+1} of dimension $3 \le n+1 \le 5$ equipped with a generic metric g always contains a sequence of closed, almost-embedded half-volume constant mean curvature hypersurfaces Σ_p with $\operatorname{Area}(\Sigma_p) \to \infty$ as $p \to \infty$. In view of Theorem 1.1, it would be very interesting to determine whether the mean curvature of these hypersurfaces goes to 0 as $p \to \infty$.

1.2. Outline of Paper. In Section 2, we first discuss some preliminary material from geometric measure theory. We then recall the authors' min-max theory for functionals of the form $\Omega \mapsto \operatorname{Area}(\partial \Omega) + f(\operatorname{Vol}(\Omega))$. Finally, we briefly review the idea of the suspension of a topological space.

In Section 3, we give the proof of Theorem 1.1. The proof is divided into three steps. In the first step, we apply Dey's lifting construction [5] to find infinitely many relative homotopy classes suitable for applying min-max theory to the functional

$$\Omega \mapsto \operatorname{Area}(\partial \Omega) + c \left| \operatorname{Vol}(\Omega) - \frac{1}{2} \operatorname{Vol}(M) \right|.$$

In the second step, we regularize this functional to

$$F_{\varepsilon,\delta}(\Omega) = \operatorname{Area}(\partial\Omega) - \delta \int_{\Omega} h + c \sqrt{\left(\operatorname{Vol}(\Omega) - \frac{1}{2}\operatorname{Vol}(M)\right)^2} + \varepsilon^2,$$

where $h: M \to \mathbb{R}$ is a suitable Morse function and ε and δ are small positive constants. We then apply min-max theory to find smooth critical points $\Omega_{\varepsilon,\delta}$ of $F_{\varepsilon,\delta}$. Finally, in the third step we let $\delta \to 0$ and then let $\varepsilon \to 0$ and obtain convergence $\Omega_{\varepsilon,\delta} \to \Omega$. We then show that $\Sigma = \partial \Omega$ is either an almost-embedded hypersurface with constant mean curvature c, or an almost-embedded hypersurface with constant mean curvature less than c but enclosing half the volume of M.

1.3. Acknowledgements. L.M. acknowledges the support of an AMS Simons Travel Grant. X.Z. acknowledges the support by NSF grant DMS-1945178, and a grant from the Simons Foundation (1026523, XZ).

2. Preliminaries

In this section we introduce some background material that will be needed for the proof of the main theorem.

2.1. Geometric Measure Theory. In this section, we recall some background from geometric measure theory. For more details we refer to [2], [16, Chapter 2], and [18]. Let M^{n+1} be a closed manifold.

- Let $\mathcal{V}(M)$ denote the space of *n*-dimensional varifolds in *M*.
- Given $V \in \mathcal{V}(M)$, let ||V|| denote the weight measure associated to V.
- Let $\mathcal{Z}_n(M, \mathbb{Z}_2)$ be the space of *n*-dimensional flat chains mod 2 in *M*.
- Given $T \in \mathcal{Z}_n(M, \mathbb{Z}_2)$, let $|T| \in \mathcal{V}(M)$ denote the varifold induced by T.
- Let $\mathcal{C}(M)$ be the space of Caccioppoli sets in M.
- Given $\Omega \in \mathcal{C}(M)$, let $\partial \Omega \in \mathcal{Z}_n(M, \mathbb{Z}_2)$ be the flat chain induced by the boundary of Ω .
- Let $\mathcal{B}(M,\mathbb{Z}_2)$ be the set of all $T \in \mathcal{Z}_n(M,\mathbb{Z}_2)$ such that $T = \partial \Omega$ for some $\Omega \in \mathcal{C}(M)$. This is the connected component of the zero cycle in $\mathcal{Z}_n(M,\mathbb{Z}_2)$.
- Let \mathcal{F} , \mathbf{F} , and \mathbf{M} denote the flat topology, the \mathbf{F} topology, and the mass topology, respectively. By definition, the \mathbf{F} topology on $\mathcal{C}(M)$ is given by

$$\mathbf{F}(\Omega_1, \Omega_2) = \mathcal{F}(\Omega_1, \Omega_2) + \mathbf{F}(|\partial \Omega_1|, |\partial \Omega_2|).$$

• Let $VZ(M, \mathbb{Z}_2)$ and VC(M) denote the VZ and VC spaces. The VZ space was first introduced by Almgren [2], and the related VC space first appeared in [19].

We will describe the VZ and VC spaces in slightly more detail, since this spaces are less wellknown. For more a more thorough introduction to these spaces and their applications in min-max theory see [19] and [12].

Definition 2.1. The space $VZ(M, \mathbb{Z}_2)$ consists of all pairs $(V, T) \in \mathcal{V}(M) \times \mathcal{B}(M, \mathbb{Z}_2)$ such that there exists a sequence $T_i \in \mathcal{B}(M, \mathbb{Z}_2)$ with $|T_i| \to V$ as varifolds and $T_i \to T$ in the flat topology on $\mathcal{B}(M, \mathbb{Z}_2)$. Given $(V,T) \in VZ(M,\mathbb{Z}_2)$ it is not necessarily true that V = |T|. However, it is always true that $||T||| \leq ||V||$ as measures. We equip $VZ(M,\mathbb{Z}_2)$ with the \mathscr{F} topology given by

$$\mathscr{F}((V_1, T_1), (V_2, T_2)) = \mathbf{F}(V_1, V_2) + \mathcal{F}(T_1, T_2).$$

The VC space is defined similarly but using $\mathcal{C}(M)$ in place of $\mathcal{B}(M, \mathbb{Z}_2)$.

Definition 2.2. The space VC(M) consists of all pairs $(V, \Omega) \in \mathcal{V}(M) \times \mathcal{C}(M)$ such that there exists a sequence $\Omega_i \in \mathcal{C}(M)$ with $|\partial \Omega_i| \to V$ as varifolds and $\Omega_i \to \Omega$ as Caccioppoli sets. The \mathscr{F} topology on VC(M) is given by $\mathscr{F}((V_1, \Omega_1), (V_2, \Omega_2)) = \mathbf{F}(V_1, V_2) + \mathcal{F}(\Omega_1, \Omega_2)$.

2.2. Min-Max Notions. In this subsection, we describe the min-max constructions for the E and F functionals developed in [12]. Let M be a closed Riemannian manifold.

First we describe the relevant parameter spaces and homotopy classes. Let I(1,k) denote the cubical complex structure on [0,1] with vertices $[0], [3^{-k}], [2 \cdot 3^{-k}], \ldots, [1]$ and edges $[0,3^{-k}], [3^{-k}, 2 \cdot 3^{-k}], \ldots, [1 - 3^{-k}, 1]$. Then let

$$I(m,k) = \underbrace{I(1,k) \otimes \cdots \otimes I(1,k)}_{m \text{ times}}$$

be the cubical complex structure subdividing I^m into 3^{mk} congruent subcubes.

Let X be a cubical subcomplex of I(m,k) for some $m, k \in \mathbb{N}$. Fix an **F**-continuous map $\Phi: X \to (\mathcal{B}(M, \mathbb{Z}_2), \mathbf{F}).$

Definition 2.3. The X-homotopy class of the map Φ is the set of all sequences of maps $\{\Phi_i \colon X \to (\mathcal{B}(M,\mathbb{Z}_2), \mathbf{F})\}$ such that there exist \mathbf{F} continuous homotopies $H_i \colon X \times [0,1] \to (\mathcal{B}(M,\mathbb{Z}_2), \mathbf{F})$ satisfying $H_i(x, 0) = \Phi(x)$ and $H_i(x, 1) = \Phi_i(x)$.

Now assume that Z is a cubical subcomplex of X. Consider an **F**-continuous map $\Psi: X \to (\mathcal{C}(M), \mathbf{F})$.

Definition 2.4. The (X, Z)-relative homotopy class of the map Ψ is the set of all sequences of maps $\{\Psi_i \colon X \to (\mathcal{C}(M), \mathbf{F})\}$ such that there exist **F**-continuous homotopy maps $H_i \colon X \times [0, 1] \to (\mathcal{C}(M), \mathbf{F})$ satisfying $H_i(x, 0) = \Psi(x)$ and $H_i(x, 1) = \Psi_i(x)$ and

$$\limsup_{i \to \infty} \left[\sup_{(z,t) \in Z \times [0,1]} \mathbf{F}(H_i(z,t), \Psi(z)) \right] = 0.$$

Remark 2.5. In [12], the min-max theory was developed only requiring the homotopies H_i in Definitions 2.3 and 2.4 to be \mathcal{F} continuous. However, inspecting the proofs reveals that it is actually possible to require that H_i is **F** continuous. Indeed, Propositions 1.14 and 1.15 in [21] can be used to ensure all relevant homotopies are **F**-continuous. See the remark above Definition 3.4 in [4], where the same observation was used.

Next, we describe the relevant functionals. Fix a smooth function $f: [0, Vol(M)] \to \mathbb{R}$. Let $\mathfrak{h} = \frac{1}{2} Vol(M)$ and assume f is even in the sense that $f(\mathfrak{h} + t) = f(\mathfrak{h} - t)$ for all $t \in [0, \mathfrak{h}]$.

Definition 2.6. Define $E: \mathcal{B}(M, \mathbb{Z}_2) \to \mathbb{R}$ by

$$E(T) = \mathbf{M}(T) + f(\operatorname{Vol}(\Omega))$$

where $\Omega \in \mathcal{C}(M)$ is any set satisfying $\partial \Omega = T$. Note that E is well-defined since f is assumed to be even. The E functional can also be extended to $VZ(M, \mathbb{Z}_2)$ by setting $E(V, T) = ||V||(M) + f(Vol(\Omega))$ where $\partial \Omega = T$.

Definition 2.7. Let Π be the X-homotopy class of a map $\Phi: X \to (\mathcal{B}(M, \mathbb{Z}_2), \mathbf{F})$. We define the min-max value

$$L^{E}(\Pi) = \inf_{\{\Phi_i\} \in \Pi} \left[\limsup_{i \to \infty} \sup_{x \in X} E(\Phi_i(x)) \right].$$

In [12], the authors proved the following min-max theorem for the E functional.

Theorem 2.8. Let M^{n+1} be a closed manifold of dimension $3 \le n+1 \le 7$. Let Π be the X-homotopy class of a map $\Phi: X \to (\mathcal{B}(M, \mathbb{Z}_2), \mathbf{F})$. Assume that $L^E(\Pi) > 0$. Then there exists $(V,T) \in VZ(M, \mathbb{Z}_2)$ with $E(V,T) = L^E(\Pi)$. Moreover, (V,T) has the following regularity. Choose $\Omega \in \mathcal{C}(M)$ with $\partial\Omega = T$ and then let $H = -f'(\operatorname{Vol}(\Omega))$.

(i) If $H \neq 0$ then there exists a smooth, closed, almost-embedded hypersurface Λ with constant mean curvature H such that $\partial \Omega = \Lambda$. Moreover, there is a (possibly empty) collection of minimal hypersurfaces $\Sigma_1, \ldots, \Sigma_k$ together with multiplicities $m_1, \ldots, m_k \in \mathbb{N}$ such that

$$V = |\Lambda| + \sum_{j=k}^{\ell} m_j |\Sigma_j|.$$

The hypersurfaces $\Lambda, \Sigma_1, \ldots, \Sigma_k$ are all disjoint.

(ii) If H = 0 then there exists a collection of minimal hypersurfaces $\Lambda_1, \ldots, \Lambda_q$ such that $\partial \Omega = \Lambda_1 \cup \ldots \cup \Lambda_q$. Moreover, there is a (possibly empty) collection of minimal hypersurfaces $\Sigma_1, \ldots, \Sigma_k$ and multiplicites $\ell_1, \ldots, \ell_q, m_1, \ldots, m_k \in \mathbb{N}$ such that

$$V = \sum_{i=1}^{q} \ell_i |\Lambda_i| + \sum_{j=1}^{k} m_j |\Sigma_j|.$$

The hypersurfaces $\Lambda_1, \ldots, \Lambda_q, \Sigma_1, \ldots, \Sigma_k$ are all disjoint.

Finally, fix a smooth Morse function $h: M \to \mathbb{R}$. We assume that h satisfies property (T); see [12, Definition 2.10]. The set of Morse functions satisfying property (T) is dense in $C^{\infty}(M)$ by [12, Proposition A.3].

Definition 2.9. Define $F: \mathcal{C}(M) \to \mathbb{R}$ by

$$F(\Omega) = \mathbf{M}(\partial \Omega) - \int_{\Omega} h + f(\operatorname{Vol}(\Omega)).$$

The F functional can also be extended to VC(M) by setting $F(V, \Omega) = ||V||(M) - \int_{\Omega} h + f(Vol(\Omega))$.

Definition 2.10. Let Π be the (X, Z)-homotopy class of a map $\Psi \colon X \to (\mathcal{C}(M), \mathbf{F})$. We define the min-max value

$$L^{F}(\Pi) = \inf_{\{\Psi_{i}\}\in\Pi} \left[\limsup_{i\to\infty} \sup_{x\in X} F(\Psi_{i}(x))\right]$$

In [12], the authors proved the following min-max theorem for the F functional.

Theorem 2.11. Let M^{n+1} be a closed manifold of dimension $3 \le n+1 \le 7$. Let Π be the (X, Z)-homotopy class of a map $\Psi \colon X \to (\mathcal{C}(M), \mathbf{F})$. Assume that

$$L^F(\Pi) > \sup_{z \in Z} F(\Psi(z)).$$

Then there exists $(V, \Omega) \in VC(M)$ satisfying $F(V, \Omega) = L^F(\Pi)$. Moreover, there exists a smooth, almost-embedded hypersurface Λ with mean curvature $H = h|_{\Lambda} - f'(Vol(\Omega))$ such that $\partial \Omega = \Lambda$ and $V = |\Lambda|$. Finally, we give the definition of *p*-sweepouts and the volume spectrum. It is known that the cohomology ring of $\mathcal{B}(M,\mathbb{Z}_2)$ with \mathbb{Z}_2 coefficients is isomorphic to $\mathbb{Z}_2[\lambda]$ with generator λ in degree one. This follows from a more general result of Almgren [1]; also see Marques-Neves [9] for a simpler proof in this special case.

Definition 2.12. Let X be a cubical subcomplex of some I(m,k). A flat continuous map $\Phi: X \to \mathcal{B}(M, \mathbb{Z}_2)$ is called a *p*-sweepout provided $\Phi^* \lambda^p \neq 0$ in $H^p(X, \mathbb{Z}_2)$.

Definition 2.13. Following [8], we say that a *p*-sweepout $\Phi: X \to \mathcal{B}(M, \mathbb{Z}_2)$ has no concentration of mass provided

$$\lim_{r \to 0} \left[\sup \{ \mathbf{M}(\Phi(x) \llcorner B(q, r)) : x \in X, \ q \in M \} \right] = 0.$$

Let \mathcal{P}_p denote the set of all *p*-sweepouts with no concentration of mass.

Definition 2.14. The volume spectrum $\{\omega_p\}_{p\in\mathbb{N}}$ is defined by

$$\omega_p = \inf_{\Phi \in \mathcal{P}_p} \left[\sup_{x \in \operatorname{dom}(\Phi)} \mathbf{M}(\Phi(x)) \right];$$

see [6] and [7].

2.3. The Suspension of a Topological Space. In this short subsection, we briefly recall the idea of the suspension of a topological space. Later we will use suspensions to construct suitable parameter spaces on which to apply min-max theory.

Definition 2.15. Let Y be a topological space. The suspension of Y is the topological space

$$SY = Y \times [-1,1]/\gamma$$

where $(y_1, 1) \sim (y_2, 1)$ and $(y_1, -1) \sim (y_2, -1)$ for all $y_1, y_2 \in Y$.

Note that Y is naturally identified with the subspace $Y \times \{0\} \subset SY$. The suspension SY can also be viewed as the union of two cones on Y, glued along a common copy of Y. More precisely, let $C_+ = Y \times [0,1]/\sim$ where $(y_1,1) \sim (y_2,1)$, and let $C_- = Y \times [-1,0]/\sim$ where $(y_1,-1) \sim (y_2,-1)$. Then C_+ and C_- are both cones on Y, and SY is obtained by gluing C_+ to C_- along $Y \times \{0\}$. Note that if Y is a cubical complex, then SY also has the structure of a cubical complex.

Finally, consider the case where Y admits a free \mathbb{Z}_2 action $y \mapsto -y$. Then this free \mathbb{Z}_2 action extends to a free \mathbb{Z}_2 action on SY given by $(y,t) \mapsto (-y,-t)$. Let $\overline{Y} = Y/\mathbb{Z}_2$ and $\overline{SY} = SY/\mathbb{Z}_2$ denote the quotient spaces. Then \overline{Y} is naturally identified with the subset $(Y \times \{0\})/\mathbb{Z}_2 \subset \overline{SY}$.

3. Proof of the Alternative

The goal of this section is to prove Theorem 1.1. Let M^{n+1} be a closed Riemannian manifold of dimension $3 \le n+1 \le 7$. Let g be a Riemannian metric on M such that no collection of minimal hypersurfaces encloses half the volume of M. Such metrics are generic in the sense of Baire category by [12, Proposition B.1]. Fix a constant c > 0.

Definition 3.1. Given $\epsilon > 0$, let $f_{\varepsilon}(t) = \sqrt{t^2 + \varepsilon^2}$.

Observe that $|f'_{\varepsilon}(t)| \leq 1$ for all $t \in \mathbb{R}$. Moreover, the functions f_{ε} converge uniformly to the absolute value function as $\varepsilon \to 0$.

Definition 3.2. Let $\mathfrak{h} = \frac{1}{2} \operatorname{Vol}(M)$.

Next we define the functionals to which we will apply mix-max theory.

Definition 3.3. For each $\varepsilon > 0$, define $E_{\varepsilon} : \mathcal{B}(M, \mathbb{Z}_2) \to \mathbb{R}$ by

$$E_{\varepsilon}(T) = \mathbf{M}(T) - c \cdot f_{\varepsilon} \big(\operatorname{Vol}(\Omega) - \mathfrak{h} \big),$$

where $\Omega \in \mathcal{C}(M)$ satisfies $\partial \Omega = T$. Also define

$$E_0(T) = \mathbf{M}(T) - c |\operatorname{Vol}(\Omega) - \mathfrak{h}|,$$

where again $\Omega \in \mathcal{C}(M)$ satisfies $\partial \Omega = T$.

Recall that $\{\omega_p\}_{p\in\mathbb{N}}$ denotes the volume spectrum of M and that \mathcal{P}_p is the set of all p-sweepouts with no concentration of mass. In our case, we our interested in the following alternative min-max values where the area functional is replaced by E_0 .

Definition 3.4. Define the min-max values

$$\alpha_p = \inf_{\Phi \in \mathcal{P}_p} \left[\sup_{x \in \operatorname{dom}(\Phi)} E_0(\Phi(x)) \right].$$

Observe that $\omega_p - c\mathfrak{h} \leq \alpha_p \leq \omega_p$ for all $p \in \mathbb{N}$. In particular, we have $\alpha_p \to \infty$ as $p \to \infty$ and so there are infinitely many p's for which there is a gap $\alpha_p < \alpha_{p+1}$. Therefore, the following Theorem 3.5 implies Theorem 1.1 as an immediate corollary. The remainder of this section will be devoted to the proof of Theorem 3.5.

Theorem 3.5. Fix $p \in \mathbb{N}$ for which there is a gap $\alpha_p < \alpha_{p+1}$. Then there exists a closed, almost-embedded hypersurface Λ in M with $\operatorname{Area}(\Lambda) \geq \alpha_{p+1}$ such that either

- (i) Λ has constant mean curvature c, or
- (ii) Λ has constant mean curvature less than c and encloses half the volume of M.

Proof. Fix some $p \in \mathbb{N}$ for which $\alpha_p < \alpha_{p+1}$.

Step 1: We will construct a suitable relative homotopy class. For this step, we will closely follow the arguments of Dey [5]. Choose a *p*-sweepout $\overline{\Phi} \colon \overline{Y} \to \mathcal{B}(M, \mathbb{Z}_2)$ with no concentration of mass for which

$$\sup_{\bar{y}\in\overline{Y}}E_0(\overline{\Phi}(\bar{y})) < \alpha_{p+1}$$

Now let Y be the double cover of \overline{Y} corresponding to the cohomology class $\overline{\Phi}^* \lambda$, where λ is the non-zero element in $H^1(\mathcal{B}(M,\mathbb{Z}_2),\mathbb{Z}_2)$. Let $y \mapsto -y$ denote the deck transformation of Y. Let $\Phi: Y \to \mathcal{C}(M)$ be the lift of $\overline{\Phi}$. Let X = SY be the suspension of Y. Note that Y is naturally identified with the subspace $Y \times \{0\}$ in X. Moreover, X carries a free \mathbb{Z}_2 action $x \mapsto -x$ extending the deck transformation on Y. Let $Z = Y \times [-2^{-1}, 2^{-1}] \subset X$. Also let \overline{X} be the quotient of X by the \mathbb{Z}_2 action, and let $\overline{Z} \subset \overline{X}$ be the quotient of Z by the \mathbb{Z}_2 action.

Next, as in [5, Section 4, Part 2], choose a map $\Lambda \colon [0,1] \to (\mathcal{C}(M),\mathcal{F})$ with no concentration of mass satisfying $\Lambda(0) = M$ and $\Lambda(1) = \emptyset$. Then define $\Psi' \colon X \to (\mathcal{C}(M),\mathcal{F})$ by

$$\Psi'(y,t) = \begin{cases} \Phi(y), & \text{if } |t| \le 2^{-1} \\ \Phi(y) \cap \Lambda(2t-1) & \text{if } t \ge 2^{-1}, \\ M \setminus (\Phi(-y) \cap \Lambda(-1-2t)), & \text{if } t \le -2^{-1}. \end{cases}$$

The map Ψ' is flat continuous by [5, Claim 4.2]. Moreover, Ψ' satisfies

$$\Psi'(x) = M \setminus \Psi'(-x)$$

by construction, and therefore Ψ' induces a map $\overline{\Psi'}: \overline{X} \to (B(M, \mathbb{Z}_2), \mathcal{F})$. The map $\overline{\Psi'}$ has no concentration of mass by [5, Claim 4.3]. Hence, by applying discretization followed by interpolation [21, Section 1.3], we can replace $\overline{\Psi'}$ by an **F**-continuous map $\overline{\Psi}: \overline{X} \to (\mathcal{B}(M, \mathbb{Z}_2), \mathbf{F})$ with the property that

(1)
$$\sup_{\bar{z}\in\overline{Z}}E_0(\overline{\Psi}(\bar{z})) < \alpha_{p+1}.$$

The map $\overline{\Psi}$ is a (p+1)-sweepout by [5, Section 4, Part 3]. Finally let $\Psi: X \to (\mathcal{C}(M), \mathbf{F})$ be the lift of $\overline{\Psi}$. Let Π be the (X, Z) relative homotopy class of the map Ψ .

Step 2: The next step is to apply min-max theory in the homotopy class Π . Fix a smooth Morse function $h: M \to \mathbb{R}$ satisfying property (T). For each $\delta > 0$, define $F_{\varepsilon,\delta}: \mathcal{C}(M) \to \mathbb{R}$ by

$$F_{\varepsilon,\delta}(\Omega) = \mathbf{M}(\partial\Omega) - \delta \int_{\Omega} h - c \cdot f_{\varepsilon}(\operatorname{Vol}(\Omega) - \mathfrak{h}).$$

Also define $\widetilde{E}_0 \colon \mathcal{C}(M) \to \mathbb{R}$ by $\widetilde{E}_0(\Omega) = \mathbf{M}(\partial\Omega) - c |\operatorname{Vol}(\Omega) - \mathfrak{h}|.$

Proposition 3.6. For any sequence $\{\Psi_i \colon X \to (\mathcal{C}(M), \mathbf{F})\}_{i \in \mathbb{N}} \in \Pi$, we have

$$\limsup_{i \to \infty} \left[\sup_{x \in X} \widetilde{E}_0(\Psi_i(x)) \right] \ge \alpha_{p+1}.$$

Proof. The proof is essentially the same as [5, Section 4, Part 4]. Recall that

$$\lim_{i \to \infty} \sup_{z \in Z} \mathbf{F}(\Psi_i(z), \Psi(z)) = 0.$$

Moreover, by definition there are **F**-continuous homotopies $H_i: \mathbb{Z} \times [0,1] \to (\mathcal{C}(M), \mathbf{F})$ satisfying $H_i(z,0) = \Psi(z)$ and $H_i(z,1) = \Psi_i(z)$ and

(2)
$$\lim_{i \to \infty} \sup_{(z,t) \in Z \times [0,1]} \mathbf{F}(H_i(z,t), \Psi(z)) = 0.$$

Let C_+ be the subset of X consisting of all pairs (y, t) with $t \ge 0$. Now define $\Xi_i \colon C_+ \to \mathcal{C}(M)$ by

$$\Xi_i(y,t) = \begin{cases} \Psi_i(y,t), & \text{if } 2^{-1} \le t \le 1\\ H_i((y,t),2t), & \text{if } 0 \le t \le 2^{-1}. \end{cases}$$

Observe that Ξ_i is **F**-continuous and that $\Xi_i(y,0) = \Psi(y,0)$ for all $y \in Y$. Since $\Psi(y,0) = M \setminus \Psi(-y,0)$ for all $y \in Y$, it follows that Ξ_i induces an **F**-continuous map $\overline{\Xi}_i : \overline{X} \to (B(M, \mathbb{Z}_2), \mathbf{F})$. Note that $\overline{\Xi}_i$ is homotopic to $\overline{\Psi}$ and therefore that $\overline{\Xi}_i$ is a (p+1)-sweepout. Moreover, $\overline{\Xi}_i$ is **F**-continuous and hence has no concentration of mass. Therefore, it follows that

$$\sup_{\bar{x}\in\overline{X}} E_0(\overline{\Xi}_i(\bar{x})) \ge \alpha_{p+1}.$$

Next, observe that (1) and (2) imply that

$$\sup_{(z,t)\in Z\times[0,1]}\widetilde{E}_0(H_i(z,t)) < \alpha_{p+1}$$

provided i is sufficiently large. Together with the previous inequality, this implies that

$$\sup_{x \in X} E_0(\Psi_i(x)) \ge \alpha_{p+1}$$

for all sufficiently large i. This proves the proposition.

Consider any sequence $\{\Psi_i \colon X \to (\mathcal{C}(M), \mathbf{F})\} \in \Pi$. It follows from Proposition 3.6 that

$$L^{F_{\varepsilon,\delta}}(\{\Psi_i\}) \ge \alpha_{p+1} - \delta \operatorname{Vol}(M) \cdot \sup_M |h| - c \cdot \sup_{t \in \mathbb{R}} |f_{\varepsilon}(t) - |t||.$$

Likewise, for all $z \in Z$ we have

$$F_{\varepsilon,\delta}(\Psi(z)) < \sup_{\bar{z}\in\overline{Z}} E_0(\overline{\Psi}(\bar{z})) + \delta \operatorname{Vol}(M) \cdot \sup_M |h| + c \cdot \sup_{t\in\mathbb{R}} \left| f_{\varepsilon}(t) - |t| \right|$$

Since f_{ε} converges uniformly to the absolute value function as $\varepsilon \to 0$, it therefore follows that

$$L^{F_{\varepsilon,\delta}}(\Pi) > \sup_{z \in Z} F_{\varepsilon,\delta}(\Psi(z))$$

provided ε and δ are small enough.

Consequently, if ε and δ are sufficiently small, we can apply min-max theory for $F_{\varepsilon,\delta}$ in the homotopy class Π ; see Theorem 1.10 in [12]. It follows that there exists $(V_{\varepsilon,\delta}, \Omega_{\varepsilon,\delta}) \in \mathrm{VC}(M)$ which is stationary for $F_{\varepsilon,\delta}$ and satisfies $F_{\varepsilon,\delta}(V_{\varepsilon,\delta}, \Omega_{\varepsilon,\delta}) = L^{F_{\varepsilon,\delta}}(\Pi)$. Moreover, there is a smooth, almost-embedded hypersurface $\Sigma_{\varepsilon,\delta}$ with $\partial\Omega_{\varepsilon,\delta} = \Sigma_{\varepsilon,\delta}$ and $V_{\varepsilon,\delta} = |\Sigma_{\varepsilon,\delta}|$. The mean curvature of $\Sigma_{\varepsilon,\delta}$ is equal to $\delta h|_{\Sigma_{\epsilon,\delta}} + h_{\varepsilon,\delta}$ where $h_{\varepsilon,\delta} = -c \cdot f'_{\varepsilon}(\mathrm{Vol}(\Omega_{\varepsilon,\delta}) - \mathfrak{h})$ is a constant satisfying $|h_{\varepsilon,\delta}| \leq c$. Finally, $(V_{\varepsilon,\delta}, \Omega_{\varepsilon,\delta})$ satisfies property (R)¹. for $F_{\varepsilon,\delta}$.

Step 3: In the final step, we will pass to the limit $\delta \to 0$ and then the limit $\varepsilon \to 0$ to construct the desired hypersurfaces. First, fix a small $\varepsilon > 0$ and consider a sequence $\delta_j \to 0$. For notational convenience, let $V_{\varepsilon,j} = V_{\varepsilon,\delta_j}$ and let $\Omega_{\varepsilon,j} = \Omega_{\varepsilon,\delta_j}$. Passing to a limit along a subsequence, we can obtain convergence $V_{\varepsilon_j} \to V_{\varepsilon}$ as varifolds and $\partial \Omega_{\varepsilon_j} \to \Omega_{\varepsilon}$ as Caccioppoli sets. The pair $(V_{\varepsilon}, \partial \Omega_{\varepsilon})$ is stationary for E_{ε} . Since the first variation of $(V_{\varepsilon,j}, \Omega_{\varepsilon,j})$ is uniformly bounded, we can apply the compactness theory in [12, Section 5] to deduce that one of the following alternatives holds:

- (i) There is a smooth, closed, almost-embedded hypersurface Λ_{ε} with non-zero constant mean curvature such that $\partial \Omega_{\varepsilon} = \Lambda_{\varepsilon}$ and $V_{\varepsilon} = |\Lambda_{\varepsilon}|$.
- (ii) The varifold V_{ε} is induced by a collection of disjoint, closed minimal hypersurfaces with integer multiplicities. Moreover, some subcollection of the minimal hypersurfaces bounds the region Ω_{ε} .

Finally, note that the fact that $(V_{\varepsilon}, \Omega_{\varepsilon})$ is stationary for E_{ε} implies that the mean curvature of $\sup ||V_{\varepsilon}||$ satisfies $|H_{\varepsilon}| = c |f'_{\varepsilon}(\operatorname{Vol}(\Omega_{\varepsilon}) - \mathfrak{h})|$.

Next, we aim to take a limit of $(V_{\varepsilon}, \Omega_{\varepsilon})$ as $\varepsilon \to 0$. Select a sequence $\varepsilon_k \to 0$. For notational convenience, let $V_k = V_{\varepsilon_k}$ and let $\Omega_k = \Omega_{\varepsilon_k}$. Passing to a limit along a subsequence, we can obtain convergence $V_k \to V$ as varifolds and $\Omega_k \to \Omega$ as Caccioppoli sets. Let H_k be the mean curvature of supp $||V_k||$. Passing to a further subsequence if necessary, we may assume that $H_k \to H$ as $k \to \infty$. Note that $|H| \leq c$. Since H_k is uniformly bounded, we can again appeal to the compactness theory in [12, Section 5] to deduce that one of the following two alternatives holds:

- (i) There is a smooth, closed, almost-embedded hypersurface Λ with non-zero constant mean curvature such that $\partial \Omega = \Lambda$ and $V = |\Lambda|$.
- (ii) The varifold V is induced by a collection of disjoint, closed minimal hypersurfaces with integer multiplicities. Moreover, some subcollection of the minimal hypersurfaces bounds the region Ω .

¹Property (R) asserts that there is a uniform number N such that $(V_{\varepsilon,\delta}, \Omega_{\varepsilon,\delta})$ must be almost minimizing for $F_{\varepsilon,\delta}$ in at least one annulus in any collection of N concentric annuli; c.f. [12, Proposition 3.15]. This property is needed to apply the compactness theory developed in [12, Section 5]

Moreover, the first alternative holds when $H \neq 0$, and in this case Λ has constant mean curvature H. The second alternative holds when H = 0.

We claim that in case (ii) one has $\operatorname{Vol}(\Omega) = \mathfrak{h}$. Indeed, this follows from the relation $|H_k| = c|f'_{\varepsilon_k}(\operatorname{Vol}(\Omega_k) - \mathfrak{h})|$ and the fact that $H_k \to 0$ when alternative (ii) holds. Since the metric g is assumed to be generic, no collection of minimal hypersurfaces in M encloses half the volume of M. Therefore, case (ii) cannot occur.

Next, let us analyze case (i) in more detail. Observe that if $\operatorname{Vol}(\Omega) \neq \mathfrak{h}$, then the relation $|H_k| = c|f'_{\varepsilon_k}(\operatorname{Vol}(\Omega_k) - \mathfrak{h})|$ implies that |H| = c. Therefore, if $\operatorname{Vol}(\Omega) \neq \mathfrak{h}$, the hypersurface Λ has constant mean curvature c. The other possibility is that $\operatorname{Vol}(\Omega) = \mathfrak{h}$, in which case Λ is a hypersurface with constant mean curvature less than c which encloses half the volume of M. Finally, note that

$$E_k(V_k,\partial\Omega_k) \ge \alpha_{p+1} - c \cdot \sup_{t \in \mathbb{R}} \left| f_{\varepsilon_k}(t) - |t| \right|$$

and consequently

$$\|V_k\|(M) \ge \alpha_{p+1} - c \cdot \sup_{t \in \mathbb{R}} \left| f_{\varepsilon_k}(t) - |t| \right|.$$

Letting $k \to \infty$, we obtain $||V||(M) \ge \alpha_{p+1}$.

To summarize, we have now shown that for every $p \in \mathbb{N}$ for which $\alpha_p < \alpha_{p+1}$, there exists a closed, almost-embedded hypersurface Λ in M with $\operatorname{Area}(\Lambda) \ge \alpha_{p+1}$ such that either

- (i) Λ has constant mean curvature c, or
- (ii) Λ has constant mean curvature less than c and encloses half the volume of M.

This completes the proof of Theorem 3.5.

References

- [1] Frederick Almgren. The homotopy groups of the integral cycle groups. Topology, 1(4):257–299, 1962.
- [2] Frederick Almgren. The theory of varifolds. Mimeographed notes, 1965.
- [3] Vladimir I Arnold. Arnold's problems. Springer, 2004.
- [4] Akashdeep Dey. A comparison of the almgren-pitts and the allen-cahn min-max theory. Geometric and Functional Analysis, 32(5):980-1040, 2022.
- [5] Akashdeep Dey. Existence of multiple closed CMC hypersurfaces with small mean curvature. J. Differential Geom., 125(2):379–403, 2023.
- [6] Mikhael Gromov. Dimension, non-linear spectra and width. In Geometric Aspects of Functional Analysis: Israel Seminar (GAFA) 1986-87, pages 132–184. Springer, 2006.
- Yevgeny Liokumovich, Fernando Marques, and André Neves. Weyl law for the volume spectrum. Annals of Mathematics, 187(3):933–961, 2018.
- [8] Fernando C Marques and André Neves. Existence of infinitely many minimal hypersurfaces in positive ricci curvature. *Inventiones mathematicae*, 209(2):577–616, 2017.
- [9] Fernando C Marques and André Neves. Morse index of multiplicity one min-max minimal hypersurfaces. Advances in Mathematics, 378:107527, 2021.
- [10] Liam Mazurowski. CMC doublings of minimal surfaces via min-max. J. Geom. Anal., 32(3):Paper No. 104, 28, 2022.
- [11] Liam Mazurowski and Xin Zhou. The half-volume spectrum of a manifold. arXiv preprint arXiv:2302.07722, 2023.
- [12] Liam Mazurowski and Xin Zhou. Infinitely many half-volume constant mean curvature hypersurfaces via min-max theory. arXiv preprint arXiv:2405.00595, 2024.
- [13] F Pacard and R Mazzeo. Foliations by constant mean curvature tubes. Communications in analysis and geometry, 13(4):633-670, 2005.
- [14] Frank Pacard. Constant mean curvature hypersurfaces in riemannian manifolds. Riv. Mat. Univ. Parma (7), 4:141–162, 2005.
- [15] Frank Pacard and Xingwang Xu. Constant mean curvature spheres in riemannian manifolds. manuscripta mathematica, 128:275–295, 2009.

AN ALTERNATIVE FOR CMCS

- [16] Jon T Pitts. Existence and regularity of minimal surfaces on Riemannian manifolds. (MN-27), volume 27. Princeton University Press, 2014.
- [17] Matthias Schneider. Closed magnetic geodesics on S^2 . Journal of Differential Geometry, 87(2):343–388, 2011.
- [18] Leon Simon. Lectures on geometric measure theory, volume 3 of Proceedings of the Centre for Mathematical Analysis, Australian National University. Australian National University, Centre for Mathematical Analysis, Canberra, 1983.
- [19] Zhichao Wang and Xin Zhou. Existence of four minimal spheres in S^3 with a bumpy metric. arXiv preprint arXiv:2305.08755, 2023.
- [20] Rugang Ye. Foliation by constant mean curvature spheres. Pacific Journal of Mathematics, 147(2):381–396, 1991.
- [21] Xin Zhou. On the multiplicity one conjecture in min-max theory. Annals of Mathematics, 192(3):767–820, 2020.
- [22] Xin Zhou. Mean curvature and variational theory. In ICM—International Congress of Mathematicians. Vol. IV. Sections 5–8, pages 2696–2717. EMS Press, Berlin, [2023] ©2023.
- [23] Xin Zhou and Jonathan J Zhu. Min-max theory for constant mean curvature hypersurfaces. Inventiones mathematicae, 218:441-490, 2019.

LEHIGH UNIVERSITY, DEPARTMENT OF MATHEMATICS, BETHLEHEM, PENNSYLVANIA 18015 *Email address*: lim624@lehigh.edu

CORNELL UNIVERSITY, DEPARTMENT OF MATHEMATICS, ITHACA, NEW YORK 14850 *Email address*: xinzhou@cornell.edu