

A LATTICE APPROACH TO MATRIX WEIGHTS

ZOE NIERAETH

ABSTRACT. In this paper we recontextualize the theory of matrix weights within the setting of Banach lattices. We define an intrinsic notion of directional Banach function spaces, generalizing matrix weighted Lebesgue spaces. Moreover, we prove an extrapolation theorem for these spaces based on the boundedness of the convex-set valued maximal operator. We also provide bounds and equivalences related to the convex body sparse operator. Finally, we introduce a weak-type analogue of directional Banach function spaces. In particular, we show that the weak-type boundedness of the convex-set valued maximal operator on matrix weighted Lebesgue spaces is equivalent to the matrix Muckenhoupt condition, with equivalent constants.

1. INTRODUCTION

1.1. Matrix weighted Lebesgue spaces. Given a Calderón-Zygmund operator T and a space of functions X modeling the initial data of a given partial differential equation, the question whether T maps X to itself is fundamental to the theory of singular integrals. This question has been fully answered when X is a weighted Lebesgue space $L_w^p(\mathbf{R}^d)$. Here, w is an a.e. positive measurable function, and we say that $f \in L_w^p(\mathbf{R}^d)$ if

$$\int_{\mathbf{R}^d} |wf|^p dx < \infty.$$

Then we have $T : L_w^p(\mathbf{R}^d) \rightarrow L_w^p(\mathbf{R}^d)$ precisely when $1 < p < \infty$, and w satisfies the Muckenhoupt A_p condition, i.e.,

$$[w]_p := \sup_Q \left(\frac{1}{|Q|} \int_Q w^p dx \right)^{\frac{1}{p}} \left(\frac{1}{|Q|} \int_Q w^{-p'} dx \right)^{\frac{1}{p'}} < \infty,$$

see, e.g., [Gra14]. Here the supremum is taken over all cubes Q in \mathbf{R}^d , and p' is the Hölder conjugate of p . Note that our normalization of the constant is due to us having introduced our weights using the multiplier approach, see [LN24].

In the '90s, this theory has been extended to the setting of so-called matrix weights. We let \mathbf{F} denote either the field \mathbf{R} or \mathbf{C} . A matrix valued mapping $W : \mathbf{R}^d \rightarrow \mathbf{F}^{n \times n}$ is called a matrix weight if it is Hermitian and positive definite. For an exponent $0 < p \leq \infty$, we define $L_W^p(\mathbf{R}^d; \mathbf{F}^n)$ as the space of measurable functions $f : \mathbf{R}^d \rightarrow \mathbf{F}^n$ for which $|Wf| \in L^p(\mathbf{R}^d)$, i.e., for which

$$\|f\|_{L_W^p(\mathbf{R}^d; \mathbf{F}^n)} := \left(\int_{\mathbf{R}^d} |W(x)f(x)|^p dx \right)^{\frac{1}{p}} < \infty,$$

where the integral is replaced by an essential supremum when $p = \infty$. We have normalized the weight as a natural extension of the multiplier notation in the scalar-valued case;

2020 *Mathematics Subject Classification.* Primary: 42B25; Secondary: 42B35, 46E30.

Key words and phrases. Banach function space, Hardy-Littlewood maximal operator, Muckenhoupt weights, Calderón-Zygmund operators, Convex bodies, Convex-set valued mappings, Matrix weights.

Z. N. is supported by the grant Juan de la Cierva formación 2021 FJC2021-046837-I, the Basque Government through the BERC 2022-2025 program, by the Spanish State Research Agency project PID2020-113156GB-I00/AEI/10.13039/501100011033 and through BCAM Severo Ochoa excellence accreditation SEV-2023-2026.

see also [BC23]. For a linear operator T acting on functions taking values in \mathbf{F} and a measurable function $f = (f_1, \dots, f_n) : \mathbf{R}^d \rightarrow \mathbf{F}^n$, we define

$$\tilde{T}f(x) := (Tf_1(x), \dots, Tf_n(x)).$$

When T is a Calderón-Zygmund operator, finding an alternative to the Muckenhoupt condition for matrix weights was initiated by Nazarov, Treil, and Volberg in the '90s, see [NT96, Tre89, TV97a, TV97b, Vol97]. In these works, they studied the boundedness of \tilde{H} on $L_W^p(\mathbf{R}; \mathbf{R}^n)$, where H is the Hilbert transform. The theory was extended to general Calderón-Zygmund operators by Christ and Goldberg [CG01]. Though usually written in a more involved way using function norms, the matrix A_p condition can be formulated by saying that there is a constant $C > 0$ such that for every cube Q and for every $u \in \mathbf{F}^n$ there is a non-zero $v \in \mathbf{F}^n$ such that

$$(1.1) \quad \left(\frac{1}{|Q|} \int_Q |W(x)u|^p dx \right)^{\frac{1}{p}} \left(\frac{1}{|Q|} \int_Q |W^{-1}(x)v|^{-p'} dx \right)^{\frac{1}{p'}} \leq C|u \cdot v|,$$

where $u \cdot v = \sum_{k=1}^n u_k \overline{v_k}$ is the standard scalar product in \mathbf{F}^n . We denote the optimal constant C by $[W]_p$.

In [NPTV17] it was shown that Calderón-Zygmund operators satisfy a so-called convex body domination. More precisely, if T is a Calderón-Zygmund operator, then there is a constant $C_T > 0$ such that for every bounded function f with bounded support there is a *sparse* collection of cubes \mathcal{S} for which for a.e. $x \in \mathbf{R}^d$ we have

$$\tilde{T}f(x) \in C_T \sum_{Q \in \mathcal{S}} \langle f \rangle_Q \mathbf{1}_Q(x),$$

where

$$(1.2) \quad \langle f \rangle_Q := \left\{ \frac{1}{|Q|} \int_Q hf dx : \|h\|_{L^\infty(\mathbf{R}^d)} \leq 1 \right\},$$

where scalar multiples of a set are considered pointwise, and where the sum of the sets is interpreted as a Minkowski sum. Using this, they showed the bound

$$(1.3) \quad \|\tilde{T}f\|_{L_W^2(\mathbf{R}^d; \mathbf{F}^n)} \lesssim [W]_2^3 \|f\|_{L_W^2(\mathbf{R}^d; \mathbf{F}^n)}.$$

for the Hilbert transform $T = H$, which was later generalized to all Calderón-Zygmund operators by Culiuc, Di Plinio, and Ou in [CDO18]. When $n > 1$, the third power dependence of $[W]_2$ was very recently shown to be sharp for the Hilbert transform by Domelevo, Petermichl, Treil, and Volberg in [DPTV24]. This is in stark contrast with the $n = 1$ case, where sparse domination yields the sharp *square* dependence of $[w]_2$:

$$(1.4) \quad \|Tf\|_{L_w^p(\mathbf{R}^d)} \lesssim [w]_2^2 \|f\|_{L_w^p(\mathbf{R}^d)},$$

see, e.g., [Ler13]. Furthermore, one can apply the sharp version of the Rubio de Francia extrapolation theorem of [DGPP05] to (1.4) to obtain the sharp bound

$$\|Tf\|_{L_w^p(\mathbf{R}^d)} \lesssim [w]_p^{\max\{p, p'\}} \|f\|_{L_w^p(\mathbf{R}^d)},$$

valid for every $1 < p < \infty$. This extrapolation theorem was very recently extended to the matrix weight setting $n > 1$ by Bownik and Cruz-Uribe in [BC23]. Their quantitative bounds match the one of the scalar-valued case $n = 1$. Due to the failure of the bound (1.4) when $n > 2$, applying their result to (1.3) yields

$$\|\tilde{T}f\|_{L_W^p(\mathbf{R}^d; \mathbf{F}^n)} \lesssim [W]_p^{\frac{3}{2} \max\{p, p'\}} \|f\|_{L_W^p(\mathbf{R}^d; \mathbf{F}^n)},$$

which does not recover the known bound

$$\|\tilde{T}f\|_{L_W^p(\mathbf{R}^d; \mathbf{F}^n)} \lesssim [W]_p^{p+p'-1} \|f\|_{L_W^p(\mathbf{R}^d; \mathbf{F}^n)}$$

by Cruz-Uribe, Isralowitz, and Moen in [CIM18, Corollary 1.16]. Thus, finding the sharp bounds outside of $p = 2$ remains open. We refer the reader to [Cru24] for an overview of the history of this problem. Qualitatively, the paper [BC23] marks a significant development in the theory of matrix weights. They fully develop the theory of convex-set valued mappings to define an analogue of the Hardy-Littlewood maximal operator adapted to matrix weights.

Denote the collection of non-empty, closed, convex, and symmetric subsets of \mathbf{F}^n by \mathcal{K} . Given a mapping $F : \mathbf{R}^d \rightarrow \mathcal{K}$, one can define its set of measurable *selections*

$$S^0(\mathbf{R}^d; F) := \{f \in L^0(\mathbf{R}^d; \mathbf{F}^n) : f(x) \in F(x) \text{ a.e.}\},$$

where $L^0(\mathbf{R}^d; \mathbf{F}^n)$ denotes the space of measurable functions $f : \mathbf{R}^d \rightarrow \mathbf{F}^n$. Moreover, we can define the average $\langle F \rangle_Q$ through the so-called Aumann integral as

$$\langle F \rangle_Q := \left\{ \frac{1}{|Q|} \int_Q f \, dx : f \in S^0(\mathbf{R}^d; F) \right\}.$$

This generalizes (1.2) in the sense that if $\mathcal{K}(f)(x)$ denotes the smallest set in \mathcal{K} containing the vector $f(x) \in \mathbf{F}^n$, then

$$\langle \mathcal{K}(f) \rangle_Q = \langle f \rangle_Q.$$

One can now define $M^{\mathcal{K}}F : \mathbf{R}^d \rightarrow \mathcal{K}$ by letting $M^{\mathcal{K}}F(x)$ be the smallest set in \mathcal{K} containing

$$\bigcup_{Q \ni x} \langle F \rangle_Q.$$

Defining $L^p_W(\mathbf{R}^d; \mathcal{K})$ as the space of those $F : \mathbf{R}^d \rightarrow \mathcal{K}$ for which the function

$$h(x) := \sup_{u \in F(x)} |W(x)u|$$

satisfies $h \in L^p(\mathbf{R}^d)$, it was shown in [BC23, Theorem 6.9] that

$$\|M^{\mathcal{K}}F\|_{L^p_W(\mathbf{R}^d; \mathcal{K})} \lesssim [W]_p^{p'} \|F\|_{L^p_W(\mathbf{R}^d; \mathcal{K})}$$

for all $1 < p \leq \infty$. This completely recovering Buckley's sharp bound [Buc93] when $n = 1$. Moreover, defining $T_Q F(x) := \langle F \rangle_Q \mathbf{1}_Q(x)$, they showed in [BC23, Proposition 6.6] that for all $1 \leq p \leq \infty$ we have

$$(1.5) \quad \sup_Q \|T_Q\|_{L^p_W(\mathbf{R}^d; \mathcal{K}) \rightarrow L^p_W(\mathbf{R}^d; \mathcal{K})} \approx [W]_p.$$

Even beyond the setting of weighted Lebesgue spaces, this condition of uniformly bounding the averaging operators over all cubes serves as a useful generalization of the Muckenhoupt condition. We refer the reader to [Nie24] and references therein for an overview.

1.2. Banach function spaces. The theory of singular integrals in more general spaces of functions has received considerable attention in the past years. A space X is called a Banach function space if it is a complete normed subspace of the space $L^0(\mathbf{R}^d)$ of measurable \mathbf{F} -valued functions, and satisfies:

- The ideal property: if $f \in X$ and $|g| \leq |f|$ a.e., then $g \in X$ with $\|g\|_X \leq \|f\|_X$;
- The saturation property: if $E \subseteq \mathbf{R}^d$ satisfies $|E| > 0$, then there is a subset $F \subseteq E$ with $|F| > 0$ and $\mathbf{1}_F \in X$.

The saturation property is equivalent to the existence of a function $\rho \in X$ satisfying $\rho > 0$ a.e., or to the non-degeneracy property that if

$$\int_{\mathbf{R}^d} fg \, dx = 0$$

for all $f \in X$, then $g = 0$. A detailed survey of why these are the right assumptions to make in the definition of a Banach function space can be found in the survey of Lorist and the author in [LN24]. This framework allows one to tackle the theory of singular integrals in

spaces such as weighted Morrey spaces and weighted variable Lebesgue spaces, which are spaces that respectively appear as boundary data of elliptic PDEs, or as non-homogeneous data of PDEs within areas such as image processing and quasi-Newtonian fluids.

In order to extend this framework to the setting of matrix weights, one can define a space X_W as the space of those $f \in L^0(\mathbf{R}^d; \mathbf{F}^n)$ for which $|Wf| \in X$ for a given matrix weight W . Exactly as was done in [BC23] for $X = L^p(\mathbf{R}^d)$, one can study the boundedness properties of singular integrals in these spaces by extending them to spaces of convex-set valued functions. We can define the space $X_W[\mathcal{K}]$ as those $F : \mathbf{R}^d \rightarrow \mathcal{K}$ for which

$$h(x) := \sup_{u \in F(x)} |W(x)u|$$

belongs to X . While this allows one to define spaces such as matrix weight analogues of Morrey spaces and variable Lebesgue spaces, it is unsatisfactory in the sense that the definition of a function space should be intrinsic, i.e., it should not be defined in terms of a given matrix weight. Even in the case $n = 1$, studying properties such as the Muckenhoupt condition and boundedness of singular integrals has a very rich theory in intrinsically defined spaces, see [Nie24] and references therein for an overview.

In the scalar-valued case $n = 1$, the ideal property states that a Banach function space X is a *Banach lattice* in the sense that the norm preserves the a.e. partial ordering of functions $|g(x)| \leq |f(x)|$. When $n > 1$, however, there are many natural partial orderings one can give to \mathbf{F}^n . To create a theory compatible with matrix weights, we need to partially order \mathbf{F}^n in such a way that a vector u dominating a vector v should imply that $|Av| \leq |Au|$ for every matrix $A \in \mathbf{F}^{n \times n}$. As it turns out, this has a very elegant characterization in terms of convex bodies. Given $u \in \mathbf{F}^n$, define

$$\mathcal{K}(u) := \{\lambda u : \lambda \in \mathbf{F}, |\lambda| \leq 1\},$$

which is the smallest set in \mathcal{K} containing u . As a (non-strict) partial ordering, we interpret the inclusion $\mathcal{K}(v) \subseteq \mathcal{K}(u)$ as u dominating v . This can be extended to functions $f, g \in L^0(\mathbf{R}^d; \mathbf{F}^n)$ through writing $\mathcal{K}(f)(x) := \mathcal{K}(f(x))$, and asking that

$$\mathcal{K}(g)(x) \subseteq \mathcal{K}(f)(x)$$

for a.e. $x \in \mathbf{R}^d$. This ordering can be characterized as follows:

Proposition. *Let $f, g \in L^0(\mathbf{R}^d; \mathbf{F}^n)$. The following are equivalent:*

- (i) $\mathcal{K}(g)(x) \subseteq \mathcal{K}(f)(x)$ for a.e. $x \in \mathbf{R}^d$;
- (ii) $g(x) \in \mathcal{K}(f)(x)$ for a.e. $x \in \mathbf{R}^d$;
- (iii) $|g \cdot u| \leq |f \cdot u|$ a.e. for all $u \in \mathbf{F}^n$;
- (iv) $g = hf$ for some $h \in L^\infty(\mathbf{R}^d)$ with $\|h\|_{L^\infty(\mathbf{R}^d)} \leq 1$.

For a proof, see Proposition 2.4 below. In essence, the ordering $\mathcal{K}(g)(x) \subseteq \mathcal{K}(f)(x)$ means that for a.e. $x \in \mathbf{R}^d$, $g(x)$ points in the same direction as $f(x)$, but with a smaller magnitude. We propose the following definition of an \mathbf{F}^n -directional Banach function space:

Definition. We say that \mathbf{X} is an \mathbf{F}^n -directional Banach function space if it is a complete normed subspace of $L^0(\mathbf{R}^d; \mathbf{F}^n)$ and satisfies the following properties:

- *The directional ideal property:* For all $f \in \mathbf{X}$ and $g \in L^0(\mathbf{R}^d; \mathbf{F}^n)$ satisfying $\mathcal{K}(g) \subseteq \mathcal{K}(f)$ a.e., we have $g \in \mathbf{X}$ with $\|g\|_{\mathbf{X}} \leq \|f\|_{\mathbf{X}}$;
- *Non-degeneracy:* If $g \in L^0(\mathbf{R}^d; \mathbf{F}^n)$ satisfies $\int_{\mathbf{R}^d} f \cdot g \, dx = 0$ for all $f \in \mathbf{X}$, then $g = 0$.

The non-degeneracy property allows us to define the Köthe dual \mathbf{X}' of \mathbf{X} through

$$\|g\|_{\mathbf{X}'} := \sup_{\|f\|_{\mathbf{X}}=1} \int_{\mathbf{R}^d} |f \cdot g| \, dx,$$

which again satisfies the directional ideal property. We call \mathbf{X} *Köthe reflexive* if $\mathbf{X}'' = \mathbf{X}$. When $n = 1$, Köthe reflexivity is characterized by the Fatou property through the Lorentz-Luxemburg theorem. A similar assertion is true for \mathbf{F}^n -directional quasi-Banach function spaces under a stronger saturation condition, see Theorem 3.4.

These spaces can be naturally extended to spaces of convex-set valued mappings $F : \mathbf{R}^d \rightarrow \mathcal{K}$. We define $L^0(\mathbf{R}^d; \mathcal{K})$ as the space of $F : \mathbf{R}^d \rightarrow \mathcal{K}$ that are measurable in the sense that for every measurable $E \subseteq \mathbf{F}^n$, the set

$$F^{-1}(E) := \{x \in \mathbf{R}^d : F(x) \cap E \neq \emptyset\}$$

is also measurable. Then we define $\mathbf{X}[\mathcal{K}]$ as the space of those $F \in L^0(\mathbf{R}^d; \mathcal{K})$ for which its space of measurable selections $S^0(\mathbf{R}^d; F)$ is a bounded subset of \mathbf{X} , with

$$\|F\|_{\mathbf{X}[\mathcal{K}]} := \sup_{f \in S^0(\mathbf{R}^d; F)} \|f\|_{\mathbf{X}}.$$

This space naturally contains \mathbf{X} through the isometric embedding $f \mapsto \mathcal{K}(f)$, see Proposition 3.5 below. Working with the general definition of $\mathbf{X}[\mathcal{K}]$ in terms of the selections $S^0(\mathbf{R}^d; F)$ is facilitated by the so-called Filippov selection theorem. It states that if $\phi : \mathbf{R}^d \times \mathbf{F}^n \rightarrow \mathbf{R}$ is measurable in its first coordinate and continuous in its second, then for any $h \in L^0(\mathbf{R}^d)$ for which for a.e. $x \in \mathbf{R}^d$, $h(x)$ is of the form $\phi(x, u)$ for some (x, u) satisfying $u \in F(x)$, then there is a selection $f \in S^0(\mathbf{R}^d; F)$ for which

$$h(x) = \phi(x, f(x)).$$

For example, this can be used to show that if $F(x)$ is a bounded set for a.e. $x \in \mathbf{R}^d$, then, for a given matrix weight $W : \mathbf{R}^d \rightarrow \mathbf{F}^{n \times n}$, there is a selection $f_0 \in S^0(\mathbf{R}^d; F)$ such that

$$|f_0(x)| = \sup_{u \in F(x)} |W(x)u|.$$

As a consequence, when X is a Banach function space with the Fatou property, the space $X_W[\mathcal{K}]$ coincides exactly with our earlier defined space of $F \in L^0(\mathbf{R}^d; \mathcal{K})$ for which the function

$$h(x) := \sup_{u \in F(x)} |W(x)u|$$

belongs to X , see Proposition 3.10 below.

1.3. Main results. Our first main result is a generalization of the extrapolation theorem of Bownik and Cruz-Uribe [BC23] to \mathbf{F}^n -directional Banach function spaces.

Theorem A. *Let $1 \leq p \leq \infty$ and suppose*

$$T : \bigcup_{W \in A_p} L_W^p(\mathbf{R}^d; \mathbf{F}^n) \rightarrow L^0(\mathbf{R}^d; \mathbf{F}^n)$$

is a map for which there is an increasing function $\phi : [0, \infty) \rightarrow [0, \infty)$ such that for all $W \in A_p$ and all $f \in L_W^p(\mathbf{R}^d; \mathbf{F}^n)$ we have

$$\|Tf\|_{L_W^p(\mathbf{R}^d; \mathbf{F}^n)} \leq \phi([W]_p) \|f\|_{L_W^p(\mathbf{R}^d; \mathbf{F}^n)}.$$

Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space for which

$$M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}], \quad M^{\mathcal{K}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}].$$

Then Tf is well-defined for all $f \in \mathbf{X}$, and

$$\|Tf\|_{\mathbf{X}} \lesssim_n \phi(C_n \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}^{\frac{1}{p'}} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]}^{\frac{1}{p}}) \|f\|_{\mathbf{X}}.$$

If $p = \infty$ or $p = 1$, we can omit the bound $M^{\mathcal{K}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]$ or $M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ respectively.

This result is proven in a more general pairs of functions setting in Theorem 6.1 below. Our proof is new, and distinct from the one of Bownik and Cruz-Urbe in how we use our Rubio de Francia extrapolation theorem. In their work, they apply the Rubio de Francia algorithm to an auxiliary operator

$$TF(x) := \sup_{u \in M^{\mathcal{K}F(x)}} |W(x)u|W(x)^{-1}\overline{B},$$

where \overline{B} is the closed unit ball in \mathbf{F}^n . Moreover, they explicitly use the fact that this operator maps the space of scalar function multiples of the ellipsoid $W(x)^{-1}\overline{B}$ to itself. As we have no matrix weight to work with, our proof of Theorem A instead relies on the Filippov selection theorem and a function norm approach, yielding a very elegant intrinsic argument. When $n = 1$, our approach is similar to the one used for Lebesgue spaces in [CMP11]; in particular, it does not recover the *sharp* extrapolation theorem for matrix weighted Lebesgue spaces. We do emphasize that our approach can be adapted to give an intrinsic proof of the extrapolation theorem of [BC23]. We also note that when $n = 1$, our result recovers the sharpest result known in the literature for general Banach function spaces of [CMM22], see also [Nie23, Section 4.7].

If T is a Calderón-Zygmund operator, then we can apply Theorem A for $p = 2$ to (1.3) to conclude that

$$(1.6) \quad \|T\|_{\mathbf{X} \rightarrow \mathbf{X}} \lesssim \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}^{\frac{3}{2}} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]}^{\frac{3}{2}}.$$

This, however, is certainly not sharp. Our second main theorem shows how the power $\frac{3}{2}$ can be reduced to 1 in this estimate using convex body domination.

For a collection of cubes \mathcal{P} and a locally integrably bounded $F : \mathbf{R}^d \rightarrow \mathcal{K}$, we define the averaging operator $T_{\mathcal{P}}$ as

$$T_{\mathcal{P}}F(x) := \sum_{Q \in \mathcal{P}} \langle F \rangle_Q \mathbf{1}_Q(x),$$

where scalar multiples of a set are considered pointwise, and the sum is interpreted as a Minkowski sum. When $F = \mathcal{K}(f)$ for a function f integrable over the cubes in \mathcal{P} , we have

$$T_{\mathcal{P}}(\mathcal{K}(f))(x) = \sum_{Q \in \mathcal{P}} \langle f \rangle_Q \mathbf{1}_Q(x).$$

A collection of cubes \mathcal{S} is called *sparse* if for all $Q \in \mathcal{S}$ we have

$$\left| \bigcup_{\substack{Q' \in \mathcal{S} \\ Q' \subseteq Q}} Q' \right| \leq \frac{1}{2}|Q|.$$

The boundedness properties of $T_{\mathcal{S}}$ in $\mathbf{X}[\mathcal{K}]$ for sparse collections \mathcal{S} dictate precisely on which spaces \mathbf{X} an operator satisfying convex body domination is bounded. Indeed, if

$$\tilde{T}f(x) \in C_T \sum_{Q \in \mathcal{S}} \langle f \rangle_Q \mathbf{1}_Q(x),$$

then, since $\|\mathcal{K}(f)\|_{\mathbf{X}[\mathcal{K}]} = \|f\|_{\mathbf{X}}$,

$$(1.7) \quad \|\tilde{T}f\|_{\mathbf{X}} \lesssim \|T_{\mathcal{S}}(\mathcal{K}(f))\|_{\mathbf{X}[\mathcal{K}]} \leq \|T_{\mathcal{S}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|f\|_{\mathbf{X}}.$$

Thus, to study bounds for T in \mathbf{X} , we need to study the behavior of $T_{\mathcal{S}}$ in $\mathbf{X}[\mathcal{K}]$. We have the following result on the relationship between the boundedness of the sparse operator $T_{\mathcal{S}}$ and of $M^{\mathcal{K}}$:

Theorem B. *Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space. Then the following are equivalent:*

- (i) $T_{\mathcal{S}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ and $T_{\mathcal{S}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]$ uniformly for all sparse collection \mathcal{S} ;
- (ii) $M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ and $M^{\mathcal{K}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]$.

Moreover, in this case we have

$$\begin{aligned} & \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \lesssim_{d,n} \|T_{\mathcal{S}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}; \\ & \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]} \lesssim_{d,n} \|T_{\mathcal{S}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]} \\ & \max\{\|T_{\mathcal{S}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}, \|T_{\mathcal{S}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]}\} \lesssim_n \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]} \end{aligned}$$

The proof of this result can be found in Section 7. Surprisingly, it is not clear if the two bounds in (i) are equivalent. This is in stark contrast with the fact that they *are* equivalent when viewing the operator $T_{\mathcal{S}}$ as a linear operator on \mathbf{X} and \mathbf{X}' instead of their convex-set valued variants. Moreover, when \mathcal{S} is pairwise disjoint, this symmetry *is* true in $\mathbf{X}[\mathcal{K}]$, see Theorem 4.4 below.

To prove Theorem B we need to establish a convex body domination result for $M^{\mathcal{K}}$, which is interesting in its own right. Here, for a collection of cubes \mathcal{P} , we define $M_{\mathcal{P}}^{\mathcal{K}}$ in the same way as $M^{\mathcal{K}}$, but with the union $\bigcup_{Q \in \mathcal{P}} \langle F \rangle_Q \mathbf{1}_Q(x)$ taken only over the $Q \in \mathcal{P}$.

Theorem C. *Let \mathcal{D} be a dyadic grid and let $\mathcal{F} \subseteq \mathcal{D}$ be finite. For each $F \in L^0(\mathbf{R}^d; \mathcal{K})$ satisfying $F \mathbf{1}_Q \in L^1(\mathbf{R}^d; \mathcal{K})$ for all cubes Q , there is a sparse collection $\mathcal{S} \subseteq \mathcal{F}$ for which*

$$M_{\mathcal{F}}^{\mathcal{K}} F(x) \subseteq C_n M_{\mathcal{S}}^{\mathcal{K}} F(x)$$

for a.e. $x \in \mathbf{R}^d$.

This theorem can be found below as Theorem 7.2. As a consequence of Theorem B and (1.7), we obtain the following improvement of (1.6):

Corollary D. *Let T be an operator for which there is a constant $C_T > 0$ such that for every $f \in L_c^\infty(\mathbf{R}^d; \mathbf{F}^n)$ there is a sparse collection \mathcal{S} for which*

$$Tf(x) \in C_T \sum_{Q \in \mathcal{S}} \langle f \rangle_Q \mathbf{1}_Q(x).$$

Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space for which

$$M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}], \quad M^{\mathcal{K}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}].$$

Then, for all $f \in L_c^\infty(\mathbf{R}^d; \mathbf{F}^n) \cap \mathbf{X}$ we have

$$\|Tf\|_{\mathbf{X}} \lesssim C_T \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]} \|f\|_{\mathbf{X}}.$$

Finally, we initiate a study into the boundedness of averaging operators related to the Muckenhoupt condition in \mathbf{F}^n -directional Banach function spaces \mathbf{X} . As we noted in (1.5), the matrix A_p condition can be recovered through the uniform boundedness of the averaging operators

$$T_Q F(x) := \langle F \rangle_Q \mathbf{1}_Q(x)$$

in $L_W^p(\mathbf{R}^d; \mathcal{K})$. Intriguingly, one does not need to bound T_Q on $L_W^p(\mathbf{R}^d; \mathcal{K})$ to recover the matrix A_p condition. Indeed, if we define $T_Q f := \langle f \rangle_Q \mathbf{1}_Q$ for $f \in L^0(\mathbf{R}^d; \mathbf{F}^n)$, then we already have

$$\sup_Q \|T_Q\|_{L_W^p(\mathbf{R}^d; \mathbf{F}^n) \rightarrow L_W^p(\mathbf{R}^d; \mathbf{F}^n)} = [W]_p,$$

this time with a strict equality. In general, given a pairwise disjoint collection of cubes \mathcal{P} and $T_{\mathcal{P}} f := \sum_{Q \in \mathcal{P}} \langle f \rangle_Q \mathbf{1}_Q$, then, by a proof based on the John ellipsoid theorem, we have

$$T_{\mathcal{P}} : \mathbf{X} \rightarrow \mathbf{X} \quad \Leftrightarrow \quad T_{\mathcal{P}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$$

with comparable operator norm, up to a constant depending only on n , see Theorem 4.4 below.

When $n = 1$, the Muckenhoupt condition $X \in A$ is defined as

$$[X]_A := \sup_Q |Q|^{-1} \|\mathbf{1}_Q\|_X \|\mathbf{1}_Q\|_{X'} < \infty,$$

and the quantity $|Q|^{-1} \|\mathbf{1}_Q\|_X \|\mathbf{1}_Q\|_{X'}$ is precisely the operator norm of $T_Q : X \rightarrow X$, see [Nie24, Section 3]. Generalizing this to $n > 1$, we say that an \mathbf{F}^n -directional Banach function space \mathbf{X} satisfies the Muckenhoupt condition $\mathbf{X} \in A$ if

$$[\mathbf{X}]_A := \sup_Q \|T_Q\|_{\mathbf{X} \rightarrow \mathbf{X}} < \infty.$$

As is the case for $n = 1$, we can determine a more or less exact expression for the operator norm $\|T_Q\|_{\mathbf{X} \rightarrow \mathbf{X}}$ in terms of indicator functions. If \mathbf{X} is Köthe reflexive, then we have $X \in A$ precisely if for all cubes Q we have $\mathbf{1}_Q u \in \mathbf{X}, \mathbf{X}'$ for all $u \in \mathbf{F}^n$, and there is a constant $C > 0$ such that for every $u \in \mathbf{F}^n$ and every cube Q there is a non-zero $v \in \mathbf{F}^n$ for which

$$\|\mathbf{1}_Q u\|_{\mathbf{X}} \|\mathbf{1}_Q v\|_{\mathbf{X}'} \leq C|u \cdot v|.$$

Moreover, the optimal C coincides with $[\mathbf{X}]_A$, see Section 4 below. When $\mathbf{X} = L_W^p(\mathbf{R}^d; \mathbf{F}^n)$, then this is precisely the Muckenhoupt condition $W \in A_p$. Several further characterizations of $\mathbf{X} \in A$ are given in Theorem 5.2 below.

To study the relationship between the boundedness of M^K in $\mathbf{X}[\mathcal{K}]$ and the Muckenhoupt condition $\mathbf{X} \in A$, we also define an analogue of the *strong* Muckenhoupt condition. We say that \mathbf{X} satisfies the strong Muckenhoupt condition $\mathbf{X} \in A_{\text{strong}}$ if

$$[\mathbf{X}]_{A_{\text{strong}}} := \sup_{\mathcal{P}} \|T_{\mathcal{P}}\|_{\mathbf{X} \rightarrow \mathbf{X}} < \infty$$

where the supremum is taken over all pairwise disjoint collections of cubes \mathcal{P} . As noted in the above discussion, we have

$$[\mathbf{X}]_A \sim_n \sup_Q \|T_Q\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}, \quad [\mathbf{X}]_{A_{\text{strong}}} \sim_n \sup_{\mathcal{P}} \|T_{\mathcal{P}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}.$$

Thus, it is not surprising that the (strong) Muckenhoupt condition is related to the bounds of M^K in $\mathbf{X}[\mathcal{K}]$. We say that $M^K : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}_{\text{weak}}[\mathcal{K}]$ if there is a constant $C > 0$ such that for all $f \in \mathbf{X}[\mathcal{K}]$ we have

$$\sup_{u \in \mathbf{F}^n} \|\mathbf{1}_{\{x \in \mathbf{R}^d : u \in M^K F(x)\}} u\|_{\mathbf{X}} \leq C \|F\|_{\mathbf{X}[\mathcal{K}]}.$$

Moreover, we denote the optimal constant C by $\|M^K\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}_{\text{weak}}[\mathcal{K}]}$. Weak-type bounds for M^K in $L^p(\mathbf{R}^d; \mathbf{F}^n)$ were also considered in [BC23], but their definition of a weak-type space differs from ours: they considered the condition that the function

$$h(x) := \sup_{u \in M^K F(x)} |u|$$

satisfies $h \in L^{p,\infty}(\mathbf{R}^d)$. It seems that our weak-type bound is better suited for the theory. Indeed, we have the following result extending [Nie24, Theorem B]:

Theorem E. *Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space. Consider the following statements:*

- (a) $M^K : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$;
- (b) $\mathbf{X} \in A_{\text{strong}}$;
- (c) $M^K : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}_{\text{weak}}[\mathcal{K}]$;
- (d) $\mathbf{X} \in A$.

Then (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d) with

$$[\mathbf{X}]_A \leq \|M^K\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}_{\text{weak}}[\mathcal{K}]} \lesssim_{d,n} [\mathbf{X}]_{A_{\text{strong}}} \leq \|M^K\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}.$$

Furthermore, if \mathbf{X} there is a $C \geq 1$ such that for all pairwise disjoint collections of cubes \mathcal{P} and all $f \in \mathbf{X}, g \in \mathbf{X}'$ we have

$$(1.8) \quad \sum_{Q \in \mathcal{P}} \|\mathbf{1}_Q f\|_{\mathbf{X}} \|\mathbf{1}_Q g\|_{\mathbf{X}'} \leq C \|f\|_{\mathbf{X}} \|g\|_{\mathbf{X}'},$$

then (b)-(d) are equivalent, with

$$[\mathbf{X}]_{A_{strong}} \simeq \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}_{weak}[\mathcal{K}]} \simeq [\mathbf{X}]_A.$$

This result can be found as Theorem 5.6 below. The condition (1.8) when $n = 1$ is often denoted as $X \in \mathcal{G}$. Moreover, if $X \in \mathcal{G}$, then for any matrix weight W , the estimate (1.8) is satisfied by X_W . An application of Hölder's inequality shows that it holds for $X = L^p(\mathbf{R}^d)$ with $C = 1$. This yields the following corollary:

Corollary F. *Let $1 \leq p \leq \infty$ and let $W : \mathbf{R}^d \rightarrow \mathbf{F}^{n \times n}$ be a matrix weight. Then the following are equivalent:*

- (i) $W \in A_p$;
- (ii) $M^{\mathcal{K}} : L^p_W(\mathbf{R}^d; \mathcal{K}) \rightarrow L^p_W(\mathbf{R}^d; \mathcal{K})_{weak}$.

Moreover, we have

$$[W]_p \simeq \|M^{\mathcal{K}}\|_{L^p_W(\mathbf{R}^d; \mathcal{K}) \rightarrow L^p_W(\mathbf{R}^d; \mathcal{K})_{weak}}.$$

Further examples of spaces X satisfying $X \in \mathcal{G}$ include variable Lebesgue spaces with global log-Hölder regular exponents. We refer the reader to [Nie24] and references therein for a further overview of this condition.

Organization. This paper is organized as follows:

- In Section 2 we define the preliminary notions of convex bodies and convex-set valued functions that we will require throughout this work.
- In Section 3 we introduce directional quasi-Banach function spaces and their convex-set valued analogues, and prove basic equivalences and results for these spaces. The directional analogue of the Lorentz-Luxemburg theorem is proven in Appendix A.
- In Section 4 we discuss the boundedness of averaging operators in directional Banach function spaces and their convex-set valued analogues in general σ -finite measure spaces.
- In Section 5 we discuss the Muckenhoupt condition and its relations to the bounds of the convex-set valued maximal operator.
- In Section 6 we prove Theorem A.
- In section 7 we prove Theorem B and Theorem C.

Notation. Throughout this work, $d, n \geq 1$ are fixed integers denoting dimension. We let \mathbf{F} denote either the field \mathbf{R} or the field \mathbf{C} . For $u, v \in \mathbf{F}^n$ we write $u = (u_1, \dots, u_n)$, $v = (v_1, \dots, v_n)$, and we define the standard scalar product and norm

$$u \cdot v := \sum_{k=1}^n u_k \overline{v_k}, \quad |u| := |u \cdot u|^{\frac{1}{2}}.$$

We write $A \lesssim B$ to mean that there is some constant C for which $A \leq CB$. If the constant C depends on parameters $\alpha_1, \alpha_2, \dots$, then we write $A \lesssim_{\alpha_1, \alpha_2, \dots} B$. Moreover, $A \gtrsim B$ and $A \gtrsim_{\alpha_1, \alpha_2, \dots} B$ are defined similarly. We write $A \simeq B$ when both $A \lesssim B$ and $A \gtrsim B$, and similarly with parameter subscripts.

2. CONVEX BODIES

Definition 2.1. We let \mathcal{C} denote the collection of non-empty closed sets $K \subseteq \mathbf{F}^n$. Moreover, we denote by \mathcal{K} the subcollection of \mathcal{C} of sets K that satisfy the following additional properties:

- *Convexity:* If $u, v \in K$, then for all $0 \leq t \leq 1$ we have $(1-t)u + tv \in K$;
- *Symmetry:* If $u \in K$, then $\lambda u \in K$ for all $\lambda \in \mathbf{F}$ with $|\lambda| = 1$;

We let \mathcal{K}_b denote the *bounded* sets in \mathcal{K} .

For $K, L \in \mathcal{C}$ and $\lambda \in \mathbf{F}$ we define $K + L$ and λK through

$$K + L := \overline{\{u + v : u \in K, v \in L\}} \in \mathcal{C}, \quad \lambda K := \{\lambda u : u \in K\} \in \mathcal{C}.$$

Taking the closure in the definition of $K + L$ is necessary, unless one of the K or L is compact, in which case the Minkowski sum is closed.

As \mathcal{K} is closed under arbitrary intersections, given a set $B \subseteq \mathbf{F}^n$, we define

$$\mathcal{K}(B) := \bigcap \{K \in \mathcal{K} : B \subseteq K\} \in \mathcal{K},$$

i.e., $\mathcal{K}(B)$ is the smallest set in \mathcal{K} containing B . Note that if B is bounded, then $\mathcal{K}(B) \in \mathcal{K}_b$.

Given a vector $u \in \mathbf{F}^n$, we write $\mathcal{K}(u) := \mathcal{K}(\{u\})$.

Proposition 2.2. *Let $u \in \mathbf{F}^n$. Then $\mathcal{K}(u) = \{\lambda u : \lambda \in \mathbf{F}, |\lambda| \leq 1\}$.*

Proof. Since $K := \{\lambda u : \lambda \in \mathbf{F}, |\lambda| \leq 1\} \in \mathcal{K}$ and $u \in K$, we have $\mathcal{K}(u) \subseteq K$. Conversely, if $v \in K$, then $v = \lambda u$ for some $|\lambda| \leq 1$. Since $u \in \mathcal{K}(u)$, symmetry implies that also $|\lambda|^{-1}v, -|\lambda|^{-1}v \in \mathcal{K}(u)$. Hence, setting $t := \frac{1}{2}(1 - |\lambda|) \in [0, 1]$, by convexity of $\mathcal{K}(u)$ we have

$$v = (1 - t)|\lambda|^{-1}v - t|\lambda|^{-1}v \in \mathcal{K}(u).$$

The result follows. \square

The space \mathcal{C}_b of closed and bounded subsets of \mathbf{F}^n is a complete metric space with respect to the Hausdorff distance

$$d_H(K, L) := \max\{\sup_{u \in K} \inf_{v \in L} |u - v|, \sup_{v \in L} \inf_{u \in K} |u - v|\}.$$

Moreover, \mathcal{K}_b is a closed subset of \mathcal{C}_b with respect to this metric.

2.1. Convex-set valued mappings. Given a σ -finite measure space (Ω, μ) , we define $L^0(\Omega; \mathcal{C})$ as the space of mappings $F : \Omega \rightarrow \mathcal{C}$ (modulo mappings a.e. equal to $\{0\}$) which are measurable in the sense that for every measurable set $E \subseteq \mathbf{F}^n$ the set

$$F^{-1}(E) := \{x \in \Omega : F(x) \cap E \neq \emptyset\}$$

is also measurable. We let $L^0(\Omega; \mathcal{K})$ denote the subset of $L^0(\Omega; \mathcal{C})$ consisting of the mappings $F : \Omega \rightarrow \mathcal{K}$, and similarly for \mathcal{K}_b . Measurability can be characterized as follows:

Proposition 2.3. *Let $F : \Omega \rightarrow \mathcal{C}$. Then the following are equivalent:*

- (i) $F \in L^0(\Omega; \mathcal{C})$;
- (ii) $x \mapsto \inf_{u \in F(x)} |u - v|$ is measurable for all $v \in \mathbf{F}^n$;
- (iii) $F^{-1}(U)$ is measurable for all open $U \subseteq \mathbf{F}^n$;
- (iv) $F(x) = \overline{\{f_k(x) : k \geq 1\}}$ a.e. for a sequence $(f_k)_{k \geq 1}$ in $L^0(\Omega; \mathbf{F}^n)$.

Moreover, if $F : \Omega \rightarrow \mathcal{K}_b$, then the above are also equivalent to

- (v) F is measurable as a function from Ω into \mathcal{K}_b equipped with the Borel σ -algebra induced by the Hausdorff distance.

The equivalences (i)-(iv) can be found in [AF09, Theorem 8.1.4]. For the equivalence with (v), see [CV77, Theorem III.2].

The partial ordering with respect to inclusion of sets of \mathcal{K} can be extended to $L^0(\Omega; \mathcal{K})$ as follows: for $F, G \in L^0(\Omega; \mathcal{K})$ we say that F dominates G if

$$G(x) \subseteq F(x)$$

for a.e. $x \in \Omega$. Moreover, given a function $f \in L^0(\Omega; \mathbf{F}^n)$, we define $\mathcal{K}(f) : \Omega \rightarrow \mathcal{K}$ by

$$\mathcal{K}(f)(x) := \mathcal{K}(f(x)).$$

This is a measurable mapping, since for a countable dense subset $(\lambda_k)_{k \geq 1}$ of the set of $\{\lambda \in \mathbf{F} : |\lambda| \leq 1\}$ we have

$$\mathcal{K}(f)(x) = \overline{\{\lambda_k f(x) : k \geq 1\}}.$$

For $f, g \in L^0(\Omega; \mathbf{F}^n)$ we say that f dominates g if

$$\mathcal{K}(g)(x) \subseteq \mathcal{K}(f)(x)$$

for a.e $x \in \Omega$. Note that this is not a strict partial ordering, since $\mathcal{K}(g) = \mathcal{K}(f)$ a.e. does not imply that $g = f$ a.e. When $n = 1$, this partial ordering is precisely the relation $|g| \leq |f|$ a.e. For general n , f dominates g if and only if $|g \cdot u| \leq |f \cdot u|$ a.e. for all $u \in \mathbf{F}^n$. This follows from the following result:

Proposition 2.4. *Let $f, g \in L^0(\Omega; \mathbf{F}^n)$. The following are equivalent:*

- (i) $\mathcal{K}(g)(x) \subseteq \mathcal{K}(f)(x)$ a.e.;
- (ii) $g(x) \in \mathcal{K}(f)(x)$ a.e.;
- (iii) $|g \cdot u| \leq |f \cdot u|$ a.e. for all $u \in \mathbf{F}^n$;
- (iv) $g = hf$ for some $h \in L^\infty(\Omega)$ with $\|h\|_{L^\infty(\Omega)} \leq 1$.

For the proof we use the following lemma:

Lemma 2.5. *Let $v, w \in \mathbf{F}^n$. Then the following are equivalent:*

- (i) $|v \cdot u| \leq |w \cdot u|$ for all $u \in \mathbf{F}^n$;
- (ii) $\mathcal{K}(v) \subseteq \mathcal{K}(w)$.

Proof. For (ii) \Rightarrow (i), note that by Proposition 2.2 there is a $|\lambda| \leq 1$ for which $v = \lambda w$. Hence, for all $u \in \mathbf{F}^n$ we have

$$|v \cdot u| = |\lambda| |w \cdot u| \leq |w \cdot u|,$$

as desired.

For (i) \Rightarrow (ii), note that the inequality implies that $w^\perp \subseteq v^\perp$, and, hence,

$$\text{span}\{v\} = (v^\perp)^\perp \subseteq (w^\perp)^\perp = \text{span}\{w\}.$$

Thus, there is a $\lambda \in \mathbf{F}$ such that $v = \lambda w$. Then

$$|\lambda| |v|^2 = |\lambda| |v \cdot v| \leq |\lambda| |w \cdot v| = |v|^2,$$

so $|\lambda| \leq 1$, proving the result by Proposition 2.2. \square

Proof of Proposition 2.4. The equivalence (i) \Leftrightarrow (iii) follows from Lemma 2.5. Moreover, for (i) \Leftrightarrow (ii), note that $g(x) \in \mathcal{K}(f)(x)$ implies, per definition of $\mathcal{K}(g)$, that $\mathcal{K}(g)(x) \subseteq \mathcal{K}(f)(x)$.

The implication (iv) \Rightarrow (iii) follows by noting that $|g \cdot u| = |h| |f \cdot u| \leq |f \cdot u|$ a.e. for all $u \in \mathbf{F}^n$. Finally, for (ii) \Rightarrow (iv), note that by Proposition 2.2 for a.e. $x \in \Omega$ there is a $\lambda_x \in \mathbf{F}$, $|\lambda_x| \leq 1$, such that $g(x) = \lambda_x f(x)$, with $\lambda_x = 0$ if $f(x) = 0$. It remains to check that $h(x) := \lambda_x$ is measurable. Writing $f = (f_1, \dots, f_n)$, $g = (g_1, \dots, g_n)$, we have $\text{supp}(f) = \bigcup_{k=1}^n \text{supp}(f_k)$ and $h(x) = \frac{f_k(x)}{g_k(x)}$ whenever $x \in \text{supp}(f_k)$. Define $E_1 := \text{supp}(f_1)$ and iteratively define $E_k := \text{supp}(f_k) \setminus \bigcup_{j=1}^{k-1} E_j$ for $k = 2, \dots, n$. Then

$$h(x) = \sum_{k=1}^n \frac{g_k(x)}{f_k(x)} \mathbf{1}_{E_k},$$

which is measurable. The assertion follows. \square

Remark 2.6. Denoting the standard basis of \mathbf{F}^n by $(e_k)_{k=1}^n$, the map

$$\iota : L^0(\Omega \times \{1, \dots, n\}) \rightarrow L^0(\Omega; \mathbf{F}^n), \quad f \mapsto \sum_{k=1}^n f(\cdot, k) e_k,$$

gives a natural one-to-one correspondence between $L^0(\Omega; \mathbf{F}^n)$ and $L^0(\Omega \times \{1, \dots, n\})$, where $\{1, \dots, n\}$ is equipped with the counting measure, and $\Omega \times \{1, \dots, n\}$ with the

product measure. One could be tempted to define $f \leq g$ on $L^0(\Omega; \mathbf{F}^n)$ by asking for the component-wise ordering

$$(2.1) \quad |f(x) \cdot e_k| \leq |g(x) \cdot e_k|$$

for a.e. $x \in \Omega$. By Proposition 2.4, this partial ordering is weaker than the one we assume. The biggest advantage of ordering vectors the way that we have, is that if $\mathcal{K}(u) \subseteq \mathcal{K}(v)$, then $|Au| \leq |Av|$ for any matrix $A \in \mathbf{F}^{n \times n}$. This fails under (2.1).

We can extend the set operations on \mathcal{K} to $L^0(\Omega; \mathcal{K})$ pointwise. That is, for $F, G \in L^0(\Omega; \mathcal{K})$ and $\lambda \in \mathbf{F}$ we can define $F + G$ and λF by

$$(F + G)(x) := \overline{\{u + v : u \in F(x), v \in G(x)\}}, \quad (\lambda F)(x) := \lambda F(x).$$

Definition 2.7. Let $F \in L^0(\Omega; \mathcal{K})$. We define the space of measurable selections of F by

$$S^0(\Omega; F) := \{f \in L^0(\Omega; \mathbf{F}^n) : f(x) \in F(x) \text{ for a.e. } x \in \Omega\}.$$

Proposition 2.8. Let $F \in L^0(\Omega; \mathcal{K})$. Then the set $S^0(\Omega; F)$ is a non-empty, convex, and symmetric subset of $L^0(\Omega; \mathbf{F}^n)$ that is closed with respect to convergence in measure.

Proof. To see that $S^0(\Omega; F)$ is non-empty, note that it contains the 0 selection $f = 0$. For convexity, if $f, g \in S^0(\Omega; F)$, then for a.e. $x \in \Omega$ and all $0 \leq t \leq 1$ we have $(1-t)f(x) + tg(x) \in F(x)$, since $F(x)$ is convex. Thus, $(1-t)f + tg \in S^0(\Omega; F)$. Symmetry is proven analogously.

To see that $S^0(\Omega; F)$ is closed, let $(f_k)_{k \geq 1}$ be a sequence in $S^0(\Omega; F)$ that converges in measure to a function $f \in L^0(\Omega; \mathbf{F}^n)$. Then there is a subsequence $(f_{k_j})_{j \geq 1}$ that converges pointwise a.e. to f . Hence, for a.e. $x \in \Omega$,

$$f(x) = \lim_{j \rightarrow \infty} f_{k_j}(x) \in \overline{F(x)} = F(x),$$

since $F(x)$ is closed. Thus, $f \in S^0(\Omega; F)$, proving the assertion. \square

Definition 2.9. We define $L^1(\Omega; \mathcal{K})$ as the space of $F \in L^0(\Omega; \mathcal{K})$ for which $S^0(\Omega; F)$ is a bounded subset of $L^1(\Omega; \mathbf{F}^n)$, with

$$\|F\|_{L^1(\Omega; \mathcal{K})} := \sup_{f \in S^0(\Omega; F)} \|f\|_{L^1(\Omega; \mathbf{F}^n)}.$$

For $F \in L^1(\Omega; \mathcal{K})$ we define the *Aumann integral* of F as

$$\int_{\Omega} F \, d\mu := \left\{ \int_{\Omega} f \, d\mu : f \in S^0(\Omega; F) \right\}.$$

If $F \in L^1(\Omega; \mathcal{K})$ then, since the embedding $L^1(\Omega; \mathbf{F}^n) \subseteq L^0(\Omega; \mathbf{F}^n)$ is continuous, it follows from Proposition 2.8 that $S^0(\Omega; F)$ is a non-empty, closed, convex, and symmetric subset of $L^1(\Omega; \mathbf{F}^n)$.

We say that F is integrably bounded if there exists a $0 \leq k \in L^1(\Omega)$ for which for a.e. $x \in \Omega$ we have

$$F(x) \subseteq \{k(x)u \in \mathbf{F}^n : |u| \leq 1\}.$$

It is shown in [BC23, Theorem 3.15] that if F is integrably bounded, then $\int_{\Omega} F \, d\mu \in \mathcal{K}_b$. This is the case for any $F \in L^1(\Omega; \mathcal{K})$:

Proposition 2.10. Let $F \in L^0(\Omega; \mathcal{K})$. Then the following are equivalent:

- (i) $F \in L^1(\Omega; \mathcal{K})$;
- (ii) F is integrably bounded;
- (iii) $h \in L^1(\Omega)$, where $h(x) := \sup_{u \in F(x)} |u|$.

Moreover, in this case we have $\|F\|_{L^1(\Omega; \mathcal{K})} = \|h\|_{L^1(\Omega)}$.

Note that the function h is measurable, as it satisfies $h(x) = \sup_{k \geq 1} |f_k(x)|$ for any sequence $(f_k)_{k \geq 1}$ in $L^0(\Omega; \mathbf{F}^n)$ satisfying

$$F(x) = \overline{\{f_k(x) : k \geq 1\}}.$$

For the proof, we will need [BC23, KNV24, Lemma 3.9]:

Lemma 2.11. *Let $F \in L^0(\Omega; \mathcal{K}_b)$. Then there is a $f_0 \in S^0(\Omega; F)$ for which for a.e. $x \in \Omega$ we have $|f_0(x)| = \sup_{u \in F(x)} |u|$.*

Proof of Proposition 2.10. For (iii) \Rightarrow (ii), note that for any $v \in F(x)$ we have

$$|v| \leq h(x).$$

Hence, for $u := \mathbf{1}_{\text{supp}(h)} h(x)^{-1}v$, we have $|u| \leq 1$ and $v = h(x)u$. We conclude that

$$F(x) \subseteq \{h(x)u \in \mathbf{F}^n : |u| \leq 1\}.$$

As $h \in L^1(\Omega)$, this proves the assertion.

For (ii) \Rightarrow (i), note that for any $f \in S^0(\Omega; F)$ we have $|f(x)| \leq k(x)$. Hence, $f \in L^1(\Omega; \mathbf{F}^n)$ with $\|f\|_{L^1(\Omega; \mathbf{F}^n)} \leq \|k\|_{L^1(\Omega)}$. We conclude that $F \in L^1(\Omega; \mathcal{K})$ with

$$\|F\|_{L^1(\Omega; \mathcal{K})} \leq \|k\|_{L^1(\Omega)},$$

as desired.

To prove (i) \Rightarrow (iii), define

$$F_k(x) := \{u \in F(x) : |u| \leq k\}.$$

Then $F_k \in L^0(\Omega; \mathcal{K}_b)$, so by Lemma 2.11 there is a selection $f_k \in S^0(\Omega; F_k)$ for which

$$h_k(x) := |f_k(x)| = \sup_{u \in F_k(x)} |u|.$$

Note that $h_k(x) \uparrow h(x) := \sup_{u \in F(x)} |u|$ a.e., and that

$$\sup_{k \geq 1} \|h_k\|_{L^1(\Omega)} = \sup_{k \geq 1} \|f_k\|_{L^1(\Omega; \mathbf{F}^n)} \leq \|F\|_{L^1(\Omega; \mathcal{K})} < \infty.$$

Hence, by the monotone convergence theorem, $h \in L^1(\Omega)$ with

$$\|h\|_{L^1(\Omega)} = \sup_{k \geq 1} \|h_k\|_{L^1(\Omega)} \leq \|F\|_{L^1(\Omega; \mathcal{K})}.$$

The result follows. \square

Corollary 2.12. *Let $F \in L^1(\Omega; \mathcal{K})$. Then there is an $f_0 \in S^0(\Omega; F)$ such that*

$$\|F\|_{L^1(\Omega; \mathcal{K})} = \|f_0\|_{L^1(\Omega; \mathbf{F}^n)}.$$

Proof. By Proposition 2.10 the function $h(x) = \sup_{u \in F(x)} |u|$ lies in $L^1(\Omega)$. This implies that $|h(x)| < \infty$ a.e., and, hence $F \in L^0(\Omega; \mathcal{K}_b)$. If we let f_0 be as in Lemma 2.11. Then, by Proposition 2.10,

$$\|F\|_{L^1(\Omega; \mathcal{K})} = \|h\|_{L^1(\Omega)} = \|f_0\|_{L^1(\Omega; \mathbf{F}^n)}.$$

This proves the result. \square

3. DIRECTIONAL QUASI-BANACH FUNCTION SPACES

We say that X is a *quasi-Banach function space over Ω* if it is a complete quasi-normed vector space $X \subseteq L^0(\Omega)$ that satisfies:

- *The ideal property:* for all $f \in X$ and $g \in L^0(\Omega)$ with $|g| \leq |f|$ a.e. we have $g \in X$ with $\|g\|_X \leq \|f\|_X$;
- *The saturation property:* for every $E \subseteq \Omega$ with $\mu(E) > 0$ there is an $F \subseteq E$ with $\mu(F) > 0$ and $\mathbf{1}_F \in X$.

We let K_X denote the optimal constant in the quasi-triangle inequality

$$\|f + g\|_X \leq K_X(\|f\|_X + \|g\|_X).$$

If $K_X = 1$, i.e., the quasi-norm of X is a norm, then we call X a Banach function space over Ω . The saturation property is equivalent to the property that the seminorm

$$\|g\|_{X'} := \sup_{\|f\|_X=1} \int_{\Omega} |fg| dx$$

is a norm, or to the existence of a *weak order unit*, i.e., a function $\rho \in X$ satisfying $\rho(x) > 0$ for a.e. $x \in \Omega$. See [LN24] for an overview.

Definition 3.1. We say that \mathbf{X} is an \mathbf{F}^n -directional quasi-Banach function space over Ω if it is a complete quasi-normed subspace of $L^0(\Omega; \mathbf{F}^n)$ and satisfies the following properties:

- *The directional ideal property:* For all $f \in \mathbf{X}$ and $g \in L^0(\Omega; \mathbf{F}^n)$ satisfying $\mathcal{K}(g)(x) \subseteq \mathcal{K}(f)(x)$ a.e., we have $g \in \mathbf{X}$ with $\|g\|_{\mathbf{X}} \leq \|f\|_{\mathbf{X}}$;
- *Non-degeneracy:* If $g \in L^0(\Omega; \mathbf{F}^n)$ satisfies $\int_{\Omega} f \cdot g d\mu = 0$ for all $f \in \mathbf{X}$, then $g = 0$.

We let $K_{\mathbf{X}}$ denote the optimal constant in the quasi-triangle inequality. If $K_{\mathbf{X}} = 1$, i.e., the quasi-norm of \mathbf{X} is a norm, then we call \mathbf{X} an \mathbf{F}^n -directional Banach function space over Ω . Completeness is equivalent to the Riesz-Fischer property, i.e., for every $(f_k)_{k \geq 1}$ for which $C := \sum_{k=1}^{\infty} K_{\mathbf{X}}^k \|f_k\|_{\mathbf{X}} < \infty$, the partial sums $\sum_{k=1}^K f_k$ have a limit $f \in \mathbf{X}$ with $\|f\|_{\mathbf{X}} \leq K_{\mathbf{X}} C$.

For verifying the directional ideal property, one can use any of the equivalences of Proposition 2.4.

Given an \mathbf{F}^n -directional quasi-Banach function space \mathbf{X} over Ω , we define \mathbf{X}' as the space of $g \in L^0(\Omega; \mathbf{F}^n)$ for which

$$\|g\|_{\mathbf{X}'} := \sup_{\|f\|_{\mathbf{X}}=1} \int_{\Omega} |f \cdot g| d\mu < \infty.$$

The non-degeneracy property is equivalent to the assertion that the seminorm $\|\cdot\|_{\mathbf{X}'}$ is a norm. This follows from the following proposition:

Proposition 3.2. *Let \mathbf{X} be an \mathbf{F}^n -directional quasi-Banach function space over Ω . Then we have $g \in \mathbf{X}'$ if and only if there is a $C \geq 0$ such that for all $f \in \mathbf{X}$ we have*

$$\left| \int_{\Omega} f \cdot g d\mu \right| \leq C \|f\|_{\mathbf{X}}.$$

Moreover, in this case the smallest possible C satisfies $\|g\|_{\mathbf{X}'} = C$.

Proof. If $g \in \mathbf{X}'$, we have

$$\left| \int_{\Omega} f \cdot g d\mu \right| \leq \int_{\Omega} |f \cdot g| d\mu \leq \|g\|_{\mathbf{X}'} \|f\|_{\mathbf{X}},$$

so it remains to prove the converse. Let $f \in \mathbf{X}$ and define

$$\tilde{f} := \mathbf{1}_{\text{supp}(f \cdot g)} \frac{|f \cdot g|}{f \cdot g} f.$$

Then, for all $v \in \mathbf{F}^n$, we have $|\tilde{f}(x) \cdot v| \leq |f(x) \cdot v|$, so that, by the directional ideal property of \mathbf{X} , we have $\tilde{f} \in \mathbf{X}$ with $\|\tilde{f}\|_{\mathbf{X}} \leq \|f\|_{\mathbf{X}}$. Hence,

$$\int_{\Omega} |f \cdot g| d\mu = \left| \int_{\Omega} \tilde{f} \cdot g d\mu \right| \leq C \|\tilde{f}\|_{\mathbf{X}} \leq C \|f\|_{\mathbf{X}}.$$

We conclude that $g \in \mathbf{X}'$, with $\|g\|_{\mathbf{X}'} \leq C$. The assertion follows. \square

We prove several characterizations of non-degeneracy in Proposition A.1 in Appendix A.

Given a complete quasi-normed subspace $\mathbf{X} \subseteq L^0(\Omega; \mathbf{F}^n)$ and $g \in L^0(\Omega; \mathbf{F}^n)$, we define \mathbf{X}_g as the space of functions $h \in L^0(\Omega')$, with $\Omega' := \text{supp}(g)$, for which $hg \in \mathbf{X}$, and set

$$\|h\|_{\mathbf{X}_g} := \|hg\|_{\mathbf{X}}.$$

Saturation of these spaces is equivalent to the directional saturation property:

- *Directional saturation:* For all non-zero $g \in L^0(\Omega; \mathbf{F}^n)$ there is a measurable set $E \subseteq \text{supp}(g)$ with $\mu(E) > 0$ such that $\mathbf{1}_E g \in \mathbf{X}$.

When $n = 1$, directional saturation and non-degeneracy coincide. However, for $n > 1$, the directional saturation property is stronger. We have the following characterizations:

Proposition 3.3. *Let \mathbf{X} be an \mathbf{F}^n -directional quasi-Banach function space over Ω . The following are equivalent:*

- (i) \mathbf{X} satisfies the directional saturation property;
- (ii) \mathbf{X}_g is a quasi-Banach function space over $\text{supp}(g)$ for all non-zero $g \in L^0(\Omega; \mathbf{F}^n)$;
- (iii) For all $g \in L^0(\Omega; \mathbf{F}^n)$ there is a sequence $(f_k)_{k \geq 1}$ in \mathbf{X} for which $|f_k \cdot u| \uparrow |g \cdot u|$ a.e. for all $u \in \mathbf{F}^n$.

Proof. For (i) \Rightarrow (ii), let $g \in L^0(\Omega; \mathbf{F}^n)$ be non-zero and let $E \subseteq \text{supp}(g)$ with $\mu(E) > 0$. Then $\mathbf{1}_E g$ is non-zero, so there is an $F \subseteq E$ with $\mu(F) > 0$ and $\mathbf{1}_F g = \mathbf{1}_F \mathbf{1}_E g \in \mathbf{X}$. Thus, $\mathbf{1}_F \in \mathbf{X}_g$, proving that \mathbf{X}_g is saturated and, hence, a quasi-Banach function space over $\text{supp}(g)$.

To see (ii) \Rightarrow (iii), for $g \in L^0(\Omega; \mathbf{F}^n)$ it follows from [LN24, Proposition 2.5(ii)] that there is an increasing sequence of sets $(E_k)_{k \geq 1}$ with $\bigcup_{k=1}^{\infty} E_k = \text{supp}(g)$ and $\mathbf{1}_{E_k} \in \mathbf{X}_g$ for all $k \geq 1$. The result then follows by setting $f_k := \mathbf{1}_{E_k} g \in \mathbf{X}$ and noting that $|f_k \cdot u| = \mathbf{1}_{E_k} |g \cdot u| \uparrow |g \cdot u|$ a.e. for all $u \in \mathbf{F}^n$.

Finally, for (iii) \Rightarrow (i), let $g \in L^0(\Omega; \mathbf{F}^n)$ be non-zero, and pick $(f_k)_{k \geq 1}$ as in (iii). Then, by Proposition 2.4(iv), there is a sequence $(h_k)_{k \geq 1}$ in $L^\infty(\Omega)$ with $0 \leq |h_k| \uparrow \mathbf{1}_{\text{supp}(g)}$ a.e. and $f_k = h_k g$. This means that for K large enough, the set $E := \{|h_K| > \frac{1}{2}\}$ has positive measure. Since

$$|\mathbf{1}_E g \cdot u| \leq 2|h_K| |g \cdot u| = |2f_K \cdot u|$$

a.e. for all $u \in \mathbf{F}^n$, it follows from the directional ideal property of \mathbf{X} that $\mathbf{1}_E g \in \mathbf{X}$. The assertion follows. \square

We say that an \mathbf{F}^n -directional quasi-Banach function space \mathbf{X} over Ω satisfies the *Fatou property* if:

- For all $(f_k)_{k \geq 1}$ in \mathbf{X} for which there is an $f \in L^0(\Omega; \mathbf{F}^n)$ such that $f_k \rightarrow f$ a.e. and $\liminf_{k \rightarrow \infty} \|f_k\|_{\mathbf{X}} < \infty$, we have $f \in \mathbf{X}$, with

$$\|f\|_{\mathbf{X}} \leq \liminf_{k \rightarrow \infty} \|f_k\|_{\mathbf{X}}.$$

We say that \mathbf{X} satisfies the *monotone convergence property* if:

- For all $(f_k)_{k \geq 1}$ in \mathbf{X} with $\mathcal{K}(f_k) \subseteq \mathcal{K}(f_{k+1})$ a.e. for all $k \geq 1$, $\sup_{k \geq 1} \|f_k\|_{\mathbf{X}} < \infty$, and

$$\bigcup_{k=1}^{\infty} \mathcal{K}(f_k) = \mathcal{K}(f)$$

for $f \in L^0(\Omega; \mathbf{F}^n)$, we have $f \in \mathbf{X}$ with $\|f\|_{\mathbf{X}} = \sup_{k \geq 1} \|f_k\|_{\mathbf{X}}$.

Moreover, we say that \mathbf{X} is *Köthe reflexive* if $\mathbf{X}'' = \mathbf{X}$ with equal norm.

When $n = 1$, the Fatou and monotone convergence properties coincide and, if \mathbf{X} is a Banach function space, then these notions coincide with Köthe reflexivity by the Lorentz-Luxemburg theorem. However, for $n > 1$, the Fatou property is stronger than the monotone convergence property: the Fatou property is defined through pointwise a.e. convergence and, hence, this includes sequences for which f_k and f do not share a direction at any point.

Note that for any \mathbf{F}^n -directional quasi-Banach function space \mathbf{X} over Ω , the space \mathbf{X}' satisfies the Fatou property by Fatou's lemma of integration theory.

The following result is an \mathbf{F}^n -directional analogue of the Lorentz-Luxemburg theorem:

Theorem 3.4 (The directional Lorentz-Luxemburg theorem). *Let \mathbf{X} be an \mathbf{F}^n -directional Banach function space over Ω with the directional saturation property. Then the following are equivalent:*

- (i) \mathbf{X} satisfies the Fatou property;
- (ii) \mathbf{X} is Köthe reflexive.

The proof of this result can be found below as Theorem A.2 in Appendix A.

3.1. Convex-set valued quasi-Banach function spaces. Let \mathbf{X} be an \mathbf{F}^n -directional quasi-Banach function space over Ω . We define $\mathbf{X}[\mathcal{K}]$ as the space of $F \in L^0(\Omega; \mathcal{K})$ for which $S^0(\Omega; F)$ is a bounded subset of \mathbf{X} . Moreover, we set

$$\|F\|_{\mathbf{X}[\mathcal{K}]} := \sup_{f \in S^0(\Omega; F)} \|f\|_{\mathbf{X}}.$$

The space $\mathbf{X}[\mathcal{K}]$ satisfies the ideal property in the sense that if $F \in \mathbf{X}[\mathcal{K}]$, then for any $G \in L^0(\Omega; \mathcal{K})$ for which

$$G(x) \subseteq F(x)$$

for a.e. $x \in \Omega$, we have $G \in \mathbf{X}[\mathcal{K}]$ with $\|G\|_{\mathbf{X}[\mathcal{K}]} \leq \|F\|_{\mathbf{X}[\mathcal{K}]}$. This follows from the observation that $S^0(\Omega; G) \subseteq S^0(\Omega; F)$.

The space $\mathbf{X}[\mathcal{K}]$ naturally contains \mathbf{X} through the embedding $f \mapsto \mathcal{K}(f)$. Indeed, we have the following result:

Proposition 3.5. *Let \mathbf{X} be an \mathbf{F}^n -directional quasi-Banach function space over Ω . Then $f \in \mathbf{X}$ if and only if $\mathcal{K}(f) \in \mathbf{X}[\mathcal{K}]$, with*

$$\|\mathcal{K}(f)\|_{\mathbf{X}[\mathcal{K}]} = \|f\|_{\mathbf{X}}.$$

Proof. If $\mathcal{K}(f) \in \mathbf{X}[\mathcal{K}]$, then, since $f \in S^0(\Omega; \mathcal{K}(f))$, we have $f \in \mathbf{X}$ with

$$\|f\|_{\mathbf{X}} \leq \sup_{g \in S^0(\Omega; \mathcal{K}(f))} \|g\|_{\mathbf{X}} = \|\mathcal{K}(f)\|_{\mathbf{X}[\mathcal{K}]}.$$

Conversely, if $f \in \mathbf{X}$, then for any $g \in S^0(\Omega; \mathcal{K}(f))$ we have $g(x) \in \mathcal{K}(f)(x)$ for a.e. $x \in \Omega$. Thus, by Proposition 2.4 and the directional ideal property of \mathbf{X} , we have $g \in \mathbf{X}$ with $\|g\|_{\mathbf{X}} \leq \|f\|_{\mathbf{X}}$. Taking a supremum over all $g \in S^0(\Omega; \mathcal{K}(f))$, we conclude that $\mathcal{K}(f) \in \mathbf{X}[\mathcal{K}]$ with

$$\|\mathcal{K}(f)\|_{\mathbf{X}[\mathcal{K}]} = \sup_{g \in S^0(\Omega; \mathcal{K}(f))} \|g\|_{\mathbf{X}} \leq \|f\|_{\mathbf{X}}.$$

The assertion follows. □

Given an \mathbf{F}^n -directional quasi-Banach function space \mathbf{X} over Ω , for a sequence $(F_k)_{k \geq 1}$ in $\mathbf{X}[\mathcal{K}]$ we write $F_k \uparrow F$, if $F_k(x) \subseteq F_{k+1}(x)$ a.e. for all $k \geq 1$, and if

$$F(x) = \overline{\bigcup_{k=1}^{\infty} F_k(x)}.$$

Note that $F \in L^0(\Omega; \mathcal{K})$. Indeed, for any open $U \subseteq \mathbf{F}^n$ we have

$$F^{-1}(U) = \bigcup_{k=1}^{\infty} F_k^{-1}(U),$$

which is measurable by measurability of the F_k .

Proposition 3.6. *Let \mathbf{X} be an \mathbf{F}^n -directional quasi-Banach function space \mathbf{X} over Ω with the monotone convergence property. If $(F_k)_{k \geq 1}$ is a bounded sequence in $\mathbf{X}[\mathcal{K}]$ and $F_k \uparrow F$, then $F \in \mathbf{X}[\mathcal{K}]$ with*

$$\|F\|_{\mathbf{X}[\mathcal{K}]} = \sup_{k \geq 1} \|F_k\|_{\mathbf{X}[\mathcal{K}]}.$$

Proof. Let $f \in S^0(\Omega; F)$. Since $\mathcal{K}(f) \cap F_k \in L^0(\Omega; \mathcal{K}_b)$, by Lemma 2.11 there is an $f_k \in S^0(\Omega; F_k)$ for which $|f_k(x)| = \sup_{u \in \mathcal{K}(f)(x) \cap F_k(x)} |u|$. Note that

$$\mathcal{K}(f_k) = \mathcal{K}(f) \cap F_k \uparrow \mathcal{K}(f) \cap F = \mathcal{K}(f).$$

Hence, by the monotone convergence property of \mathbf{X} , $f \in \mathbf{X}$ with

$$\|f\|_{\mathbf{X}} = \sup_{k \geq 1} \|f_k\|_{\mathbf{X}} \leq \sup_{k \geq 1} \|F_k\|_{\mathbf{X}[\mathcal{K}]}.$$

We conclude that $F \in \mathbf{X}[\mathcal{K}]$, with

$$\|F\|_{\mathbf{X}[\mathcal{K}]} \leq \sup_{k \geq 1} \|F_k\|_{\mathbf{X}[\mathcal{K}]}.$$

This proves the assertion. \square

3.2. Matrix weights. A measurable mapping $W : \Omega \rightarrow \mathbf{F}^{n \times n}$ is called a *matrix weight* if for a.e. $x \in \Omega$ the matrix $W(x)$ is Hermitian and positive definite, i.e., for all non-zero $u \in \mathbf{F}^n$ we have $W(x)u \cdot u > 0$.

Given a matrix weight $W : \Omega \rightarrow \mathbf{F}^{n \times n}$ and a quasi-Banach function space X over Ω , we define the \mathbf{F}^n -directional quasi-Banach function space X_W as the space of functions $f \in L^0(\Omega; \mathbf{F}^n)$ for which

$$|Wf| \in X,$$

and set

$$\|f\|_{X_W} := \| |Wf| \|_X.$$

Then $\|\cdot\|_{X_W}$ is indeed a norm, since $\|f\|_{X_W} = 0$ is equivalent to $Wf = 0$ a.e., which is equivalent to $f = W^{-1}Wf = 0$ a.e. To see that X_W satisfies the directional ideal property, suppose that $f \in X_W$ and $g \in L^0(\Omega; \mathbf{F}^n)$ satisfies $\mathcal{K}(g) \subseteq \mathcal{K}(f)$ a.e. Then, by Proposition 2.4, there is a $h \in L^\infty(\Omega)$ with $\|h\|_{L^\infty(\Omega)} \leq 1$ such that $g = hf$. Hence,

$$|Wg| = |h| |Wf| \leq |Wf|$$

a.e., so that by the ideal property of X we have $g \in X_W$, with

$$\|g\|_{X_W} = \| |Wg| \|_X \leq \| |Wf| \|_X = \|f\|_{X_W}.$$

The non-degeneracy property follows from the fact that $(X_W)' = (X')_{W^{-1}}$:

Proposition 3.7. *Let $W : \Omega \rightarrow \mathbf{F}^{n \times n}$ be a matrix weight, and let X be a quasi-Banach function space over Ω . Then $(X_W)' = (X')_{W^{-1}}$.*

In particular, note that if X is a Banach function space with the Fatou property, then $(X_W)'' = (X'')_W = X_W$ by the Lorentz-Luxemburg theorem.

Proof. First, suppose that $g \in (X')_{W^{-1}}$. Then for all $f \in X_W$ we have

$$\begin{aligned} \int_{\Omega} |f \cdot g| \, d\mu &= \int_{\Omega} |Wf \cdot W^{-1}g| \, d\mu \leq \int_{\Omega} |Wf| |W^{-1}g| \, d\mu \\ &\leq \| |Wf| \|_X \| |W^{-1}g| \|_{X'} = \|f\|_{X_W} \|g\|_{(X')_{W^{-1}}}. \end{aligned}$$

Hence, $g \in (X_W)'$ with $\|g\|_{(X_W)'} \leq \|g\|_{(X')_{W^{-1}}}$. Conversely, suppose $g \in (X_W)'$. Let $k \in X$ and define

$$h := \begin{cases} k \frac{W^{-1}g}{|W^{-1}g|} & \text{if } W^{-1}g \neq 0; \\ 0 & \text{if } W^{-1}g = 0. \end{cases}$$

Then $|h| \leq |k|$ a.e., so $|h| \in X$ by the ideal property of X . Setting $f := W^{-1}h \in X_W$, we have

$$\|f\|_{X_W} = \| |h| \|_X \leq \|k\|_X.$$

Hence,

$$\begin{aligned} \int_{\Omega} |k| |W^{-1}g| \, d\mu &= \int_{\Omega} |h \cdot W^{-1}g| \, d\mu = \int_{\Omega} |f \cdot g| \, d\mu \\ &\leq \|f\|_{X_W} \|g\|_{(X_W)'} \leq \|k\|_X \|g\|_{(X_W)'}. \end{aligned}$$

Thus, $g \in (X')_{W^{-1}}$ with $\|g\|_{(X')_{W^{-1}}} \leq \|g\|_{(X_W)'}$. The result follows. \square

As a matter of fact, X_W satisfies the directional saturation property. Indeed, for any non-zero $g \in L^0(\Omega; \mathbf{F}^n)$ we have

$$\|h\|_{(X_W)_g} = \|hg\|_{X_W} = \|h|Wg|\|_X = \|h\|_{X(|Wg|)},$$

showing that $(X_W)_g = X(|Wg|)$. Here, for a function $u \geq 0$, we define $X(u)$ through $\|h\|_{X(u)} := \|hu\|_X$. Since $|Wg| > 0$ a.e. on $\text{supp}(g)$, the space $(X_W)_g$ is saturated by [LN24, Proposition 3.17]. The directional saturation property of X_W now follows from Proposition 3.3.

A very useful result for matrix weighted spaces is Filippov's selection theorem. The following version can be found in [AF09, Theorem 8.2.10]:

Theorem 3.8 (Filippov). *Let $F \in L^0(\Omega; \mathcal{C})$ and $\phi : \Omega \times \mathbf{F}^n \rightarrow \mathbf{R}$ a function for which for all $u \in \mathbf{F}^n$ the function $x \mapsto \phi(x, u)$ is measurable and for a.e. $x \in \Omega$ the function $u \mapsto \phi(x, u)$ is continuous. For each $h \in L^0(\Omega)$ satisfying*

$$h(x) \in \{\phi(x, u) : u \in F(x)\}$$

for a.e. $x \in \Omega$, there is a selection $f \in S^0(\Omega; F)$ for which

$$h(x) = \phi(x, f(x))$$

for a.e. $x \in \Omega$.

As a consequence, we can show that if X satisfies the Fatou property, then any $F \in X_W[\mathcal{K}]$ satisfies $F(x) \in \mathcal{K}_b$ a.e.

Proposition 3.9. *Let X be a quasi-Banach function space over Ω with the Fatou property, and let $W : \Omega \rightarrow \mathbf{F}^{n \times n}$ be a matrix weight. Then $X_W[\mathcal{K}] \subseteq L^0(\Omega; \mathcal{K}_b)$.*

Proof. Define

$$F_k(x) := \{u \in F(x) : |u| \leq k\}.$$

Then, by applying Theorem 3.8 with

$$\phi(x, u) := |W(x)u|, \quad h_k(x) = \sup_{u \in F_k(x)} \phi(x, u),$$

there is a selection $f_k \in S^0(\Omega; F_k)$ such that

$$h_k(x) = |f_k(x)| = \sup_{u \in F_k(x)} |W(x)u|.$$

Since

$$\sup_{k \geq 1} \|h_k\|_X = \sup_{k \geq 1} \|f_k\|_{X_W} \leq \|F\|_{X_W[\mathcal{K}]},$$

it follows from the Fatou property of X that $h(x) = \sup_{k \geq 1} h_k(x) = \sup_{u \in F(x)} |W(x)u|$ satisfies $h \in X$ with

$$\|h\|_X = \sup_{k \geq 1} \|h_k\|_X \leq \|F\|_{X_W[\mathcal{K}]}.$$

This implies that $h(x) < \infty$ a.e. and, hence,

$$\sup_{u \in F(x)} |u| \leq |W(x)^{-1}|h(x) < \infty$$

a.e., as desired. \square

We can also prove the following result:

Proposition 3.10. *Let X be a quasi-Banach function space over Ω , let $W : \Omega \rightarrow \mathbf{F}^{n \times n}$ be a matrix weight, and let $F \in L^0(\Omega; \mathcal{K}_b)$. Then the following are equivalent:*

- (i) $F \in X_W[\mathcal{K}]$;
- (ii) $h \in X$, where $h(x) := \sup_{u \in F(x)} |W(x)u|$.

Moreover, in this case there is an $f_0 \in S^0(\Omega; F)$ such that

$$\|F\|_{X_W[\mathcal{K}]} = \|f_0\|_{X_W} = \|h\|_X.$$

Note that if X satisfies the Fatou property, then any $F \in X_W[\mathcal{K}]$ satisfies $F \in L^0(\Omega; \mathcal{K}_b)$ by Proposition 3.9.

Proof of Proposition 3.10. For (i) \Rightarrow (ii), we apply Theorem 3.8 to

$$\phi(x, u) = |W(x)u|, \quad h(x) = \sup_{u \in F(x)} \phi(x, u),$$

to find an $f_0 \in S^0(\Omega; F)$ for which

$$|f_0(x)| = \sup_{u \in F(x)} |W(x)u|$$

a.e. Then $f_0 \in X_W$, so $h = |Wf_0| \in X$, with

$$\|h\|_X = \|f_0\|_{X_W} \leq \|F\|_{X_W[\mathcal{K}]},$$

as desired. Conversely, for (ii) \Rightarrow (i), if $f \in S^0(\Omega; F)$, then $|Wf| \leq h$ a.e. Hence, by the ideal property of X , we have $f \in X_W$ with $\|f\|_{X_W} = \||Wf|\|_X \leq \|h\|_X$. Thus, $F \in X_W[\mathcal{K}]$ with

$$\|F\|_{X_W[\mathcal{K}]} = \sup_{f \in S^0(\Omega; F)} \|f\|_{X_W} \leq \|h\|_X.$$

The assertion follows. \square

3.3. Infinite sums of convex-set valued mappings. Since we are interested in constructing a version of the Rubio de Francia algorithm in $\mathbf{X}[\mathcal{K}]$, we need to be able to define infinite sums of convex-set valued mappings. We first show that if $F, G \in L^0(\Omega; \mathcal{K})$, then their closed Minkowski sum $F + G$ is again in $L^0(\Omega; \mathcal{K})$.

Proposition 3.11. *Let $F, G \in L^0(\Omega; \mathcal{K})$. Then*

$$(F + G)(x) = \overline{\{u + v : u \in F(x), v \in G(x)\}}$$

satisfies $F + G \in L^0(\Omega; \mathcal{K})$. Moreover, if $h \in S^0(\Omega; F + G)$, then there are sequences $(f_k)_{k \geq 1}$ and $(g_k)_{k \geq 1}$ respectively in $S^0(\Omega; F)$ and $S^0(\Omega; G)$ such that

$$f_k + g_k \rightarrow h$$

a.e. Moreover, if \mathbf{X} is an \mathbf{F}^n -directional quasi-Banach function space with the Fatou property and $F, G \in \mathbf{X}[\mathcal{K}]$, then $F + G \in \mathbf{X}[\mathcal{K}]$ with

$$\|F + G\|_{\mathbf{X}[\mathcal{K}]} \leq K_{\mathbf{X}}(\|F\|_{\mathbf{X}[\mathcal{K}]} + \|G\|_{\mathbf{X}[\mathcal{K}]}).$$

Proof. Let $(f_k)_{k \geq 1}$ and $(g_k)_{k \geq 1}$ be measurable selections of F and G respectively such that

$$F(x) = \overline{\{f_k(x) : k \geq 1\}}, \quad G(x) = \overline{\{g_k(x) : k \geq 1\}}$$

for a.e. $x \in \Omega$. Then the functions $h_{j,k} := f_j + g_k \in S^0(\Omega; F + G)$ satisfy

$$(F + G)(x) = \overline{\{h_{j,k}(x) : j, k \geq 1\}}$$

a.e. As $(h_{j,k})_{j,k \geq 1}$ is countable, we conclude that $F + G \in L^0(\Omega; \mathcal{K})$.

Now, suppose $h \in S^0(\Omega; F + G)$. By applying Theorem 3.8 to

$$\phi(x, (u, v)) = |h(x) - (u + v)|, \quad h_K(x) = \min_{j,k \in \{1, \dots, K\}} |h(x) - (f_k(x) + g_j(x))|,$$

we find a selection $(\tilde{f}_K, \tilde{g}_K) \in S^0(\Omega; F \times G) = S^0(\Omega; F) \times S^0(\Omega; G)$ for which

$$|h(x) - (\tilde{f}_K(x) + \tilde{g}_K(x))| = \min_{j,k \in \{1, \dots, K\}} |h(x) - (f_k(x) + g_j(x))|$$

a.e. Since the right-hand side converges to

$$\inf_{j,k \geq 1} |h(x) - (f_k(x) + g_j(x))| = 0,$$

we conclude that $\tilde{f}_K + \tilde{g}_K \rightarrow h$ a.e., as desired. If \mathbf{X} has the Fatou property and $F, G \in \mathbf{X}[\mathcal{K}]$, then also $h \in \mathbf{X}$ with

$$\begin{aligned} \|h\|_{\mathbf{X}} &\leq \liminf_{k \rightarrow \infty} \|\tilde{f}_K + \tilde{g}_K\|_{\mathbf{X}} \leq \sup_{K \geq 1} K_{\mathbf{X}}(\|\tilde{f}_K\|_{\mathbf{X}} + \|\tilde{g}_K\|_{\mathbf{X}}) \\ &\leq K_{\mathbf{X}}(\|F\|_{\mathbf{X}[\mathcal{K}]} + \|G\|_{\mathbf{X}[\mathcal{K}]}). \end{aligned}$$

Hence, $F + G \in \mathbf{X}[\mathcal{K}]$ with

$$\|F + G\|_{\mathbf{X}[\mathcal{K}]} \leq K_{\mathbf{X}}(\|F\|_{\mathbf{X}[\mathcal{K}]} + \|G\|_{\mathbf{X}[\mathcal{K}]}),$$

proving the result. \square

Given a sequence $(F_k)_{k \geq 1}$ in $\mathbf{X}[\mathcal{K}]$, their partial sums are defined through the closed Minkowski sum

$$S_K(x) := \sum_{k=1}^K F_k(x) = \overline{\left\{ \sum_{k=1}^K u_k : u_k \in F_k(x) \right\}}.$$

If $S_K \uparrow F$, then we write $\sum_{k=1}^{\infty} F_k := F$. Per definition, this means that

$$\sum_{k=1}^{\infty} F_k(x) = \bigcup_{K=1}^{\infty} S_K(x) = \bigcup_{K=1}^{\infty} \overline{\left\{ \sum_{k=1}^K u_k : u_k \in F_k(x) \right\}}.$$

By Proposition 3.11 we have $S_K \in L^0(\Omega; \mathcal{K})$. Hence, we also have $\sum_{k=1}^{\infty} F_k \in L^0(\Omega; \mathcal{K})$.

Theorem 3.12 (Directional Riesz-Fischer). *Let \mathbf{X} be a quasi-Banach function space over Ω with the Fatou property, and let $(F_k)_{k \geq 1}$ be a sequence in $\mathbf{X}[\mathcal{K}]$ satisfying*

$$\sum_{k=1}^{\infty} K_{\mathbf{X}}^k \|F_k\|_{\mathbf{X}[\mathcal{K}]} < \infty.$$

Then $F := \sum_{k=1}^{\infty} F_k \in \mathbf{X}[\mathcal{K}]$ with

$$\|F\|_{\mathbf{X}[\mathcal{K}]} \leq \sum_{k=1}^{\infty} K_{\mathbf{X}}^k \|F_k\|_{\mathbf{X}[\mathcal{K}]}.$$

Proof. By inductively applying Proposition 3.11, we find that $S_K \in \mathbf{X}[\mathcal{K}]$ with

$$\|S_K\|_{\mathbf{X}[\mathcal{K}]} \leq \sum_{k=1}^K K_{\mathbf{X}}^k \|F_k\|_{\mathbf{X}[\mathcal{K}]} \leq \sum_{k=1}^{\infty} K_{\mathbf{X}}^k \|F_k\|_{\mathbf{X}[\mathcal{K}]}.$$

Since $S_K \uparrow F$ and the Fatou property of \mathbf{X} implies the monotone convergence property, it follows from Proposition 3.6 that $F \in \mathbf{X}[\mathcal{K}]$ with

$$\|F\|_{\mathbf{X}[\mathcal{K}]} = \sup_{K \geq 1} \|S_K\|_{\mathbf{X}[\mathcal{K}]} \leq \sum_{k=1}^{\infty} K_{\mathbf{X}}^k \|F_k\|_{\mathbf{X}[\mathcal{K}]}.$$

This proves the assertion. \square

The space $\mathbf{X}[\mathcal{K}]$ is not a quasi-normed vector space, as it lacks an additive structure (there is no additive inverse to Minkowski summation of sets). Nonetheless, it is a quasi-metric space with respect to

$$d_{\mathbf{X}[\mathcal{K}]}(F, G) := \max\left\{ \sup_{f \in S^0(\Omega; F)} \inf_{g \in S^0(\Omega; G)} \|f - g\|_{\mathbf{X}}, \sup_{g \in S^0(\Omega; G)} \inf_{f \in S^0(\Omega; F)} \|f - g\|_{\mathbf{X}} \right\}.$$

Let $\mathcal{K}_b(\mathbf{X})$ denote the collection of closed, bounded, convex, and symmetric subsets of \mathbf{X} . Then $d_{\mathbf{X}[\mathcal{K}]}$ is exactly the Hausdorff quasi-distance inherited from $\mathcal{K}_b[\mathbf{X}]$ through the embedding

$$\mathbf{X}[\mathcal{K}] \hookrightarrow \mathcal{K}_b[\mathbf{X}], \quad F \mapsto \overline{S^0(\Omega; F)}.$$

Note that $d_{\mathbf{X}[\mathcal{K}]}(F, \{0\}) = \|F\|_{\mathbf{X}[\mathcal{K}]}$ and

$$(3.1) \quad d_{\mathbf{X}[\mathcal{K}]}(F + H, G + H) \leq d_{\mathbf{X}[\mathcal{K}]}(F, G), \quad d_{\mathbf{X}[\mathcal{K}]}(\lambda F, \lambda G) = |\lambda| d_{\mathbf{X}[\mathcal{K}]}(F, G)$$

for all $F, G, H \in \mathbf{X}[\mathcal{K}]$ and $\lambda \in \mathbf{F}$.

In [BC23] the metric defined on $L_W^p(\mathbf{R}^d; \mathcal{K})$ is through a pointwise weighted Hausdorff distance. In general, for matrix weighted spaces X_W , this approach is equivalent. Indeed, setting

$$d_W(F, G)(x) := \max\left\{ \sup_{u \in F(x)} \inf_{v \in G(x)} |W(x)(u - v)|, \sup_{v \in G(x)} \inf_{u \in F(x)} |W(x)(u - v)| \right\},$$

we have the following:

Proposition 3.13. *Let X be a quasi-Banach function space over Ω and let $W : \Omega \rightarrow \mathbf{F}^{n \times n}$ be a matrix weight. Then*

$$d_{X_W[\mathcal{K}]}(F, G) \approx \|d_W(F, G)\|_X.$$

Proof of Proposition 3.13. Let $f \in S^0(\Omega; F)$ and apply Theorem 3.8 with

$$\phi(x, v) = |W(x)f(x) - W(x)v|, \quad h(x) = \inf_{v \in G(x)} \phi(x, v)$$

to find a $g \in S^0(\Omega; G)$ such that

$$|W(x)f(x) - W(x)g(x)| = d_W(f, G)(x) := \inf_{v \in G(x)} |W(x)f(x) - W(x)v|$$

a.e. Hence, by the ideal property of X ,

$$\|f - g\|_{X_W} = \| |W(f - g)| \|_X \leq \|d_W(f, G)\|_X.$$

We conclude that

$$\sup_{f \in S^0(\Omega; F)} \inf_{g \in S^0(\Omega; G)} \|f - g\|_{X_W} \leq \|d_W(f, G)\|_X$$

Noting that $d_W(f, G) \leq d_W(F, G)$ a.e. and by repeating the same argument with the roles of F and G reversed, we conclude that

$$d_{X_W[\mathcal{K}]}(F, G) \leq \|d_W(F, G)\|_X.$$

For the converse inequality, apply Theorem 3.8 with

$$\phi(x, u) = \inf_{v \in G(x)} |W(x)u - W(x)v|, \quad h(x) = \sup_{u \in F(x)} \phi(x, u),$$

where measurability of $\phi(\cdot, u)$ follows from Proposition 2.2. This yields an $f_0 \in S^0(\Omega; F)$ for which

$$\inf_{v \in G(x)} |W(x)f_0(x) - W(x)v| = \sup_{u \in F(x)} \inf_{v \in G(x)} |W(x)(u - v)|$$

a.e. Thus, for any $g \in S^0(\Omega; G)$ we have

$$\sup_{u \in F(x)} \inf_{v \in G(x)} |W(x)(u - v)| \leq |W(x)f_0(x) - W(x)g(x)|.$$

a.e. By the ideal property of X this implies

$$\left\| \sup_{u \in F(x)} \inf_{v \in G(x)} |W(x)(u - v)| \right\|_X \leq \|f_0 - g\|_{\mathbf{X}}.$$

Taking an infimum over all $g \in S^0(\Omega; G)$, we conclude that

$$\begin{aligned} \left\| \sup_{u \in F} \inf_{v \in G} |W(u - v)| \right\|_X &\leq \inf_{g \in S^0(\Omega; G)} \|f_0 - g\|_{\mathbf{X}} \\ &\leq \sup_{f \in S^0(\Omega; F)} \inf_{g \in S^0(\Omega; G)} \|f - g\|_{\mathbf{X}}. \end{aligned}$$

Repeating the argument with the roles of F and G reversed, we conclude that

$$\begin{aligned} \|d_W(F, G)\|_X &\leq K_X \left\| \sup_{u \in F} \inf_{v \in G} |W(u - v)| \right\|_X + K_X \left\| \sup_{v \in G} \inf_{u \in F} |W(u - v)| \right\|_X \\ &\leq 2K_X d_{X_W[\mathcal{K}]}(F, G). \end{aligned}$$

This proves the assertion. \square

Completeness of $(X_W, d_{X_W[\mathcal{K}]})$ can be proven analogously to [BC23, Theorem 4.8]. Even without assuming the Fatou property, the space X_W satisfies the directional Riesz-Fischer property of Theorem 3.12:

Proposition 3.14. *Let X be a quasi-Banach function space over Ω and let $W : \Omega \rightarrow \mathbf{F}^{n \times n}$ be a matrix weight. If $(F_k)_{k \geq 1}$ is a sequence in $X_W[\mathcal{K}]$ satisfying*

$$C := \sum_{k=1}^{\infty} K_X^k \|F_k\|_{X_W[\mathcal{K}]} < \infty,$$

then $F := \sum_{k=1}^{\infty} F_k \in X_W[\mathcal{K}]$ with $\|F\|_{X_W[\mathcal{K}]} \leq K_X C$.

Proof. For $f \in S^0(\Omega; F)$ we have

$$(3.2) \quad |W(x)f(x)| \leq \sum_{k=1}^{\infty} \sup_{u_k \in F_k(x)} |W(x)u_k|$$

a.e. Since, by Proposition 3.10, the sequence $h_k(x) := \sup_{u_k \in F_k(x)} |W(x)u_k|$ satisfies $\|h_k\|_X = \|F_k\|_{X_W[\mathcal{K}]}$, we have $\sum_{k=1}^{\infty} K_X^k \|h_k\|_X < \infty$. Thus, by the Riesz-Fischer property of X , the sum on the right-hand side of (3.2) belongs to X with

$$\left\| \sum_{k=1}^{\infty} h_k \right\|_X \leq K_X \sum_{k=1}^{\infty} K_X^k \|h_k\|_X =: C.$$

Hence, by the ideal property of X , $f \in X_W$ with

$$\|f\|_{X_W} = \|Wf\|_X \leq C.$$

Thus, $F \in X_W[\mathcal{K}]$ with $\|F\|_{X_W[\mathcal{K}]} \leq C$, as asserted. \square

We leave it as an open problem whether the Fatou property is necessary for Theorem 3.12 or not.

3.4. Weak-type directional quasi-Banach function spaces. Given an \mathbf{F}^n -directional quasi-Banach function space \mathbf{X} over Ω , we define a weak analogue \mathbf{X}_{weak} as the space of $f \in L^0(\Omega; \mathbf{F}^n)$ for which $\mathbf{1}_{\mathcal{K}(f)^{-1}(\{u\})} u \in \mathbf{X}$ for all $u \in \mathbf{F}^n$, with

$$\|f\|_{\mathbf{X}_{\text{weak}}} := \sup_{u \in \mathbf{F}^n} \|\mathbf{1}_{\mathcal{K}(f)^{-1}(\{u\})} u\|_{\mathbf{X}}.$$

When $n = 1$, this space satisfies the quasi-triangle inequality as a consequence of the fact that for any $K, L \in \mathcal{K}$ we have $K + L \subseteq 2(K \cup L)$. However, this inclusion fails when $n > 2$, and, hence, \mathbf{X}_{weak} might not be a vector space. Moreover, we note that if $\|f\|_{\mathbf{X}_{\text{weak}}} = 0$, then we can conclude that

$$\mu(\{x \in \Omega : u \in \mathcal{K}(f)(x)\}) = 0$$

for all non-zero $u \in \mathbf{F}^n$. However, it is not clear if this implies that $f = 0$ a.e. Despite these difficulties, \mathbf{X}_{weak} does still satisfy the directional ideal and non-degeneracy properties, and inherits the monotone convergence property from \mathbf{X} in case it has it.

Proposition 3.15. *Let \mathbf{X} be an \mathbf{F}^n -directional quasi-Banach function space over Ω . Then \mathbf{X}_{weak} satisfies the directional ideal and non-degeneracy properties. Moreover, we have $\mathbf{X} \subseteq \mathbf{X}_{\text{weak}}$ with*

$$\|f\|_{\mathbf{X}_{\text{weak}}} \leq \|f\|_{\mathbf{X}},$$

and

$$\|F\|_{\mathbf{X}_{\text{weak}}[\mathcal{K}]} = \sup_{u \in \mathbf{F}^n} \|\mathbf{1}_{F^{-1}(\{u\})} u\|_{\mathbf{X}}.$$

Finally, if \mathbf{X} satisfies the monotone convergence property, then so does \mathbf{X}_{weak} .

Proof. Note that if $\mathcal{K}(g) \subseteq \mathcal{K}(f)$ a.e., then $\mathcal{K}(g)^{-1}(\{u\}) \subseteq \mathcal{K}(f)^{-1}(\{u\})$ for all $u \in \mathbf{F}^n$, so the directional ideal property of \mathbf{X}_{weak} follows from the directional ideal property of \mathbf{X} .

For the non-degeneracy property, we first show that $\mathbf{X} \subseteq \mathbf{X}_{\text{weak}}$. Note that since

$$\mathbf{1}_{\mathcal{K}(f)^{-1}(\{u\})}(x)u = \mathbf{1}_{\{x \in \Omega : u \in \mathcal{K}(f)(x)\}}(x)u \in \mathcal{K}(f)(x)$$

a.e. for all $u \in \mathbf{F}^n$, it follows from the directional ideal property that $\mathbf{X} \subseteq \mathbf{X}_{\text{weak}}$ with

$$\|f\|_{\mathbf{X}_{\text{weak}}} \leq \|f\|_{\mathbf{X}},$$

as desired. Now, suppose $g \in L^0(\Omega; \mathbf{F}^n)$ and that $\int_{\Omega} f \cdot g \, d\mu = 0$ for all $f \in \mathbf{X}_{\text{weak}}$. Then, since $\mathbf{X} \subseteq \mathbf{X}_{\text{weak}}$, this also holds for all $f \in \mathbf{X}$. Thus, by the non-degeneracy property of \mathbf{X} , we have $g = 0$ a.e., as desired.

If $F \in \mathbf{X}_{\text{weak}}[\mathcal{K}]$, then for any $f \in S^0(\Omega; F)$ and $u \in \mathbf{F}^n$ we have $\mathcal{K}(f)^{-1}(\{u\}) \subseteq F^{-1}(\{u\})$, so, by the directional ideal property,

$$\|\mathbf{1}_{\mathcal{K}(f)^{-1}(\{u\})} u\|_{\mathbf{X}} \leq \|\mathbf{1}_{F^{-1}(\{u\})} u\|_{\mathbf{X}} \leq \sup_{u \in \mathbf{F}^n} \|\mathbf{1}_{F^{-1}(\{u\})} u\|_{\mathbf{X}}.$$

Taking a supremum over all $u \in \mathbf{F}^n$ then shows $F \in \mathbf{X}_{\text{weak}}[\mathcal{K}]$ with

$$\|F\|_{\mathbf{X}_{\text{weak}}[\mathcal{K}]} \leq \sup_{u \in \mathbf{F}^n} \|\mathbf{1}_{F^{-1}(\{u\})} u\|_{\mathbf{X}}.$$

Conversely, let $u \in \mathbf{F}^n$ and define $f := \mathbf{1}_{F^{-1}(\{u\})} u$. Then $f \in S^0(\Omega; F)$, and

$$\mathcal{K}(f)^{-1}(\{u\}) = F^{-1}(\{u\}).$$

Hence,

$$\|\mathbf{1}_{F^{-1}(\{u\})} u\|_{\mathbf{X}} = \|\mathbf{1}_{\mathcal{K}(f)^{-1}(\{u\})} u\|_{\mathbf{X}} \leq \|f\|_{\mathbf{X}_{\text{weak}}} \leq \|F\|_{\mathbf{X}_{\text{weak}}[\mathcal{K}]}.$$

Taking a supremum over all $u \in \mathbf{F}^n$ now proves the desired identity.

Next, suppose \mathbf{X} has the monotone convergence property. If $(f_k)_{k \geq 1}$ is such that $(\mathcal{K}(f_k))_{k \geq 1}$ is an increasing sequence with union $\mathcal{K}(f)$, then

$$\mathbf{1}_{\mathcal{K}(f_k)^{-1}(\{u\})} \uparrow \mathbf{1}_{\mathcal{K}(f)^{-1}(\{u\})}$$

for all $u \in \mathbf{F}^n$. Hence, $\mathbf{1}_{\mathcal{K}(f)^{-1}(\{u\})} u \in \mathbf{X}$, with

$$\|\mathbf{1}_{\mathcal{K}(f)^{-1}(\{u\})} u\|_{\mathbf{X}} = \sup_{k \geq 1} \|\mathbf{1}_{\mathcal{K}(f_k)^{-1}(\{u\})} u\|_{\mathbf{X}} \leq \sup_{k \geq 1} \|f_k\|_{\mathbf{X}_{\text{weak}}}.$$

Thus, taking a supremum over $u \in \mathbf{F}^n$, proves that $f \in \mathbf{X}_{\text{weak}}$. We conclude that \mathbf{X}_{weak} has the monotone convergence property. \square

Remark 3.16. When X is a quasi-Banach function space over Ω and $W : \Omega \rightarrow \mathbf{F}^n$ is a matrix weight, then we can define the alternative weak-type space as follows. We define X_{weak} through

$$\|f\|_{X_{\text{weak}}} := \sup_{t > 0} \|t \mathbf{1}_{\{|f| > t\}}\|_X.$$

As this is again a quasi-Banach function space over Ω , the space $(X_{\text{weak}})_W$ is a well-defined \mathbf{F}^n -directional quasi-Banach function space over Ω consisting of the $f \in L^0(\Omega; \mathbf{F}^n)$ for which $|Wf| \in X_{\text{weak}}$.

The difference between the two spaces can be more easily illustrated when $n = 1$. If w is a weight, $X = L_w^p(\mathbf{R}^d)$, and $f \in L_w^p(\mathbf{R}^d)$, then for any $u \in \mathbf{F}^n$ we have

$$\{x \in \mathbf{R}^d : u \in \mathcal{K}(f)(x)\} = \{x \in \mathbf{R}^d : |u| \leq |f(x)|\}.$$

Hence, $L_w^p(\mathbf{R}^d)_{\text{weak}}$ as we have defined in this section is given by

$$\|f\|_{L_w^p(\mathbf{R}^d)_{\text{weak}}} = \sup_{u \in \mathbf{F}^n} \|\mathbf{1}_{\mathcal{K}(f)^{-1}(u)} u\|_{L_w^p(\mathbf{R}^d)} = \sup_{t > 0} t w^p(\{|f| > t\})^{\frac{1}{p}}.$$

On the other hand, we have

$$\|f\|_{(L^p(\mathbf{R}^d)_{\text{weak}})_w} = \|wf\|_{L^{p,\infty}(\mathbf{R}^d)} = \sup_{t > 0} t \{|\{wf| > t\}|\}^{\frac{1}{p}}.$$

Thus, the different approaches respectively yield a weak-type space with respect to a change of measure, and a weak-type space with respect to a weight as a multiplier.

4. AVERAGING OPERATORS

4.1. Directional averaging operators. Throughout this section we let $E \subseteq \Omega$ be a measurable set for which $0 < \mu(E) < \infty$. For $f \in L^0(\Omega)$ we set

$$T_E f := \langle f \rangle_E \mathbf{1}_E, \quad \langle f \rangle_E := \frac{1}{\mu(E)} \int_E f \, d\mu.$$

This can be extended component-wise to functions $f \in L^0(\Omega; \mathbf{F}^n)$, which we also denote by $T_E f$.

Given an \mathbf{F}^n -directional quasi-Banach function space Ω , we note that for any $u, v \in \mathbf{F}^n$ with $\mathbf{1}_E u \in \mathbf{X}$, $\mathbf{1}_E v \in \mathbf{X}'$, we have

$$\mu(E)^{-1} \|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} \geq \frac{1}{\mu(E)} \int_{\Omega} |u \mathbf{1}_E \cdot v \mathbf{1}_E| \, d\mu = |u \cdot v|.$$

A converse inequality characterizes the boundedness $T_E : \mathbf{X} \rightarrow \mathbf{X}$.

Proposition 4.1. *Let \mathbf{X} be an \mathbf{F}^n -directional quasi-Banach function space over Ω . Then the following assertions hold:*

- (a) *If $T_E : \mathbf{X} \rightarrow \mathbf{X}$, then for all $v \in \mathbf{F}^n$ we have $\mathbf{1}_E v \in \mathbf{X}'$. Moreover, for all $v \in \mathbf{F}^n$ and all $\varepsilon > 0$ there is a non-zero $u \in \mathbf{F}^n$ for which $\mathbf{1}_E u \in \mathbf{X}$ and*

$$\mu(E)^{-1} \|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} \leq (1 + \varepsilon) \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} |u \cdot v|.$$

(b) If for all $u \in \mathbf{F}^n$ we have $\mathbf{1}_E u \in \mathbf{X}$, and there is a $C > 0$ such that for all $u \in \mathbf{F}^n$ there is a non-zero $v \in \mathbf{F}^n$ for which $\mathbf{1}_E v \in \mathbf{X}'$ and

$$\mu(E)^{-1} \|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} \leq C |u \cdot v|,$$

then $T_E : \mathbf{X} \rightarrow \mathbf{X}$ with

$$\|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} \leq C.$$

Proof. Let $v \in \mathbf{F}^n$, let $f \in \mathbf{X}$ with $\|f\|_{\mathbf{X}} = 1$, and let $u := \langle f \rangle_E$. Then we have

$$\begin{aligned} \left| \int_E f \cdot v \, d\mu \right| &= \mu(E) |\langle f \rangle_E \cdot v| = \mu(E) \frac{|u \cdot v|}{\|\mathbf{1}_E u\|_{\mathbf{X}}} \|\langle f \rangle_E \mathbf{1}_E\|_{\mathbf{X}} \\ &\leq \mu(E) \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} \frac{|u \cdot v|}{\|\mathbf{1}_E u\|_{\mathbf{X}}} \\ &\leq \mu(E) \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} \sup_{\substack{u \in \mathbf{F}^n \setminus \{0\} \\ \mathbf{1}_E u \in \mathbf{X}}} \frac{|u \cdot v|}{\|\mathbf{1}_E u\|_{\mathbf{X}}}. \end{aligned}$$

Hence, by Proposition 3.2 we have $\mathbf{1}_E v \in \mathbf{X}'$. Let $\varepsilon > 0$ and let $f \in \mathbf{X}$ such that

$$\|\mathbf{1}_E v\|_{\mathbf{X}'} \leq (1 + \varepsilon) \left| \int_E f \cdot v \, d\mu \right|.$$

Setting $u := \langle f \rangle_E$, we have $\mathbf{1}_E u = T_E f \in \mathbf{X}$, and, by the same computation as above,

$$\begin{aligned} \|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} &\leq (1 + \varepsilon) \|\mathbf{1}_E u\|_{\mathbf{X}} \left| \int_E f \cdot v \, d\mu \right| \\ &\leq (1 + \varepsilon) \mu(E) \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} |u \cdot v|, \end{aligned}$$

proving (a).

For (b), let $f \in \mathbf{X}$, let $u := \langle f \rangle_E$. Pick $v \in \mathbf{F}^n$ with $\mathbf{1}_E v \in \mathbf{X}'$ for which

$$\mu(E)^{-1} \|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} \leq C |\langle f \rangle_E \cdot v|.$$

Since

$$|\langle f \rangle_E \cdot v| \leq \mu(E)^{-1} \int_{\Omega} |f \cdot \mathbf{1}_E v| \, d\mu \leq \mu(E)^{-1} \|f\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'},$$

we obtain

$$\|T_E f\|_{\mathbf{X}} = \|\mathbf{1}_E u\|_{\mathbf{X}} \leq C \|f\|_{\mathbf{X}},$$

as desired. \square

For the following result, recall that by the directional Lorentz-Luxemburg theorem (Theorem 3.4), if \mathbf{X} has the Fatou property and the directional saturation property, then \mathbf{X} is Köthe reflexive.

Corollary 4.2. *Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space over Ω . For $E \subseteq \Omega$ with $0 < \mu(E) < \infty$, the following are equivalent:*

- (i) $T_E : \mathbf{X} \rightarrow \mathbf{X}$;
- (ii) $T_E : \mathbf{X}' \rightarrow \mathbf{X}'$;
- (iii) For all $u \in \mathbf{F}^n$ we have $\mathbf{1}_E u \in \mathbf{X}$, and there is a $C_1 > 0$ such that for all $u \in \mathbf{F}^n$ there is a $v \in \mathbf{F}^n$ for which $\mathbf{1}_E v \in \mathbf{X}'$ and

$$\mu(E)^{-1} \|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} \leq C_1 |u \cdot v|;$$

- (iv) For all $v \in \mathbf{F}^n$ we have $\mathbf{1}_E v \in \mathbf{X}'$, and there is a $C_2 > 0$ such that for all $v \in \mathbf{F}^n$ there is a $u \in \mathbf{F}^n$ for which $\mathbf{1}_E u \in \mathbf{X}'$ and

$$\mu(E)^{-1} \|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} \leq C_2 |u \cdot v|.$$

Moreover, the optimal constants C_1, C_2 satisfy

$$\|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} = \|T_E\|_{\mathbf{X}' \rightarrow \mathbf{X}'} = C_1 = C_2.$$

Proof. Assuming (i), by Proposition 4.1 we have $\mathbf{1}_E v \in \mathbf{X}'$ for all $v \in \mathbf{F}^n$, and for a given $v \in \mathbf{F}^n$ and $\varepsilon = \frac{1}{k}$, there is a non-zero $u_k \in \mathbf{F}^n$ for which

$$\mu(E)^{-1} \|\mathbf{1}_E u_k\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} \leq (1 + \frac{1}{k}) \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} |u_k \cdot v|.$$

By homogeneity, we may assume that $|u_k| = 1$. Then, by compactness, there is a convergent subsequence $(u_{k_j})_{j \geq 1}$ with limit $u \in \mathbf{F}^n$. Since then $\mathbf{1}_E u_{k_j} \rightarrow \mathbf{1}_E u$ a.e., it follows from the Fatou property that $\mathbf{1}_E u \in \mathbf{X}$ with

$$\begin{aligned} \mu(E)^{-1} \|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} &\leq \liminf_{j \rightarrow \infty} \mu(E)^{-1} \|\mathbf{1}_E u_{k_j}\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} \\ &\leq \liminf_{j \rightarrow \infty} (1 + \frac{1}{k_j}) \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} |u_{k_j} \cdot v| = \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} |u \cdot v|. \end{aligned}$$

This proves (iv) with $C_2 \leq \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}}$. Since $\mathbf{X}'' = \mathbf{X}$, another application of Proposition 4.1 proves that (iv) \Rightarrow (ii) with $\|T_E\|_{\mathbf{X}' \rightarrow \mathbf{X}'} \leq C_2$. Repeating this argument with the roles of \mathbf{X} and \mathbf{X}' reversed, we have proven the chain of implications

$$(i) \Rightarrow (iv) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (i),$$

with

$$\|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} \leq C_1 \leq \|T_E\|_{\mathbf{X}' \rightarrow \mathbf{X}'} \leq C_2 \leq \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}}.$$

The result follows. \square

Note that the above proposition implies that if \mathbf{X} satisfies the directional saturation property and the Fatou property, then $T_E : \mathbf{X} \rightarrow \mathbf{X}$ precisely when $\mathbf{1}_E u \in \mathbf{X}, \mathbf{X}'$ for all $u \in \mathbf{F}^n$, and

$$\|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} = \sup_{|u|=1} \inf_{|v|=1} \frac{\|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'}}{|u \cdot v| \mu(E)} = \sup_{|v|=1} \inf_{|u|=1} \frac{\|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'}}{|u \cdot v| \mu(E)}.$$

If \mathbf{X} satisfies the property that $\mathbf{1}_E u \in \mathbf{X}$ for all $u \in \mathbf{F}^n$, then we can define a norm on \mathbf{F}^n through

$$u \mapsto \|\mathbf{1}_E u\|_{\mathbf{X}}.$$

Using the John ellipsoid theorem, there is a Hermitian positive definite matrix $A_{\mathbf{X},E} \in \mathbf{F}^{n \times n}$ that satisfies

$$|A_{\mathbf{X},E} u| \leq \|\mathbf{1}_E u\|_{\mathbf{X}} \leq n^{\frac{1}{2}} |A_{\mathbf{X},E} u|.$$

We call the matrix $A_{\mathbf{X},E}$ the *reducing matrix in \mathbf{X} of E* . Reducing matrices can be used to give another characterization of the boundedness of T_E :

Proposition 4.3. *Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space over Ω . Then $T_E : \mathbf{X} \rightarrow \mathbf{X}$ if and only if $\mathbf{1}_E u \in \mathbf{X}, \mathbf{X}'$ for all $u \in \mathbf{F}^n$. Moreover, we have*

$$\|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} \sim_n \mu(E)^{-1} \|A_{\mathbf{X},E} A_{\mathbf{X}',E}\|_{\mathbf{F}^{n \times n}}.$$

We note that since $A_{\mathbf{X},E}$ and $A_{\mathbf{X}',E}$ are Hermitian, we have

$$\begin{aligned} \|A_{\mathbf{X},E} A_{\mathbf{X}',E}\|_{\mathbf{F}^{n \times n}} &= \|(A_{\mathbf{X},E} A_{\mathbf{X}',E})^*\|_{\mathbf{F}^{n \times n}} \\ &= \|A_{\mathbf{X}',E}^* A_{\mathbf{X},E}^*\|_{\mathbf{F}^{n \times n}} \\ &= \|A_{\mathbf{X}',E} A_{\mathbf{X},E}\|_{\mathbf{F}^{n \times n}}. \end{aligned}$$

Proof of Proposition 4.3. First assume that $T_E : \mathbf{X} \rightarrow \mathbf{X}$. Then, by Corollary 4.2 we have $\mathbf{1}_E u \in \mathbf{X}, \mathbf{X}'$ for all $u \in \mathbf{F}^n$. Let $w \in \mathbf{F}^n$ and set $u := A_{\mathbf{X}',E} w$. By Corollary 4.2(iii) there is a non-zero $v \in \mathbf{F}^n$ such that

$$\begin{aligned} |A_{\mathbf{X},E} A_{\mathbf{X}',E} w| &\leq \|\mathbf{1}_E u\|_{\mathbf{X}} \leq \mu(E) \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} \frac{|u \cdot v|}{\|\mathbf{1}_E v\|_{\mathbf{X}'}} \\ &\leq \mu(E) \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} \frac{|w \cdot A_{\mathbf{X}',E} v|}{|A_{\mathbf{X}',E} v|} \end{aligned}$$

$$\leq \mu(E) \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} |w|.$$

Hence, $\mu(E)^{-1} \|A_{\mathbf{X},E} A_{\mathbf{X}',E}\|_{\mathbf{F}^{n \times n}} \leq \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}}$.

For the converse, let $f \in \mathbf{X}$ and set $v := \langle A_{\mathbf{X},E} f \rangle_E$. Then

$$\begin{aligned} \mu(E) |\langle A_{\mathbf{X},E} f \rangle_E|^2 &= \int_E A_{\mathbf{X},E} f \cdot v \, d\mu = \int_E f \cdot A_{\mathbf{X},E} v \, d\mu \\ &\leq \|f\|_{\mathbf{X}} \|\mathbf{1}_E A_{\mathbf{X},E} v\|_{\mathbf{X}'} \leq n^{\frac{1}{2}} \|A_{\mathbf{X}',E} A_{\mathbf{X},E} v\| \|f\|_{\mathbf{X}} \\ &\leq n^{\frac{1}{2}} \|A_{\mathbf{X}',E} A_{\mathbf{X},E}\|_{\mathbf{F}^{n \times n}} |\langle A_{\mathbf{X},E} f \rangle_E| \|f\|_{\mathbf{X}}, \end{aligned}$$

so that

$$\|T_E f\|_{\mathbf{X}} \leq n^{\frac{1}{2}} |\langle A_{\mathbf{X},E} f \rangle_E| \leq n \mu(E)^{-1} \|A_{\mathbf{X}',E} A_{\mathbf{X},E}\|_{\mathbf{F}^{n \times n}} \|f\|_{\mathbf{X}}.$$

Since

$$\|A_{\mathbf{X}',E} A_{\mathbf{X},E}\|_{\mathbf{F}^{n \times n}} = \|(A_{\mathbf{X}',E} A_{\mathbf{X},E})^*\|_{\mathbf{F}^{n \times n}} = \|A_{\mathbf{X},E} A_{\mathbf{X}',E}\|_{\mathbf{F}^{n \times n}},$$

this proves the assertion. \square

4.2. Convex-set valued averaging operators. Given a measurable set $E \subseteq \Omega$ with $0 < \mu(E) < \infty$ and $F \in L^0(\Omega; \mathcal{K})$, we define $F \mathbf{1}_E$ pointwise by

$$(F \mathbf{1}_E)(x) := \mathbf{1}_E(x) F(x) = \begin{cases} F(x) & \text{if } x \in E; \\ \{0\} & \text{if } x \notin E. \end{cases}$$

If $F \mathbf{1}_E \in L^1(\Omega; \mathcal{K})$, we use the Aumann integral to define

$$\langle F \rangle_E := \frac{1}{\mu(E)} \int_{\Omega} F \mathbf{1}_E \, d\mu = \{\langle f \rangle_E : f \in S^0(\Omega; F \mathbf{1}_E)\}.$$

The averaging operator T_E of F is defined as

$$T_E F(x) := \langle F \rangle_E \mathbf{1}_E(x) = \{T_E f(x) : f \in S^0(\Omega; F \mathbf{1}_E)\}.$$

We note that if for a function $f \in L^0(\Omega; \mathbf{F}^n)$ with $f \mathbf{1}_E \in L^1(\Omega; \mathbf{F}^n)$ we define

$$\langle\langle f \rangle\rangle_E := \{\langle hf \rangle_E : h \in L^\infty(\Omega), \|h\|_{L^\infty(\Omega)} \leq 1\},$$

then we have

$$T_E(\mathcal{K}(f)) = \langle\langle f \rangle\rangle_E \mathbf{1}_E.$$

This follows from the observation that by Proposition 2.4 the measurable selections of $\mathcal{K}(f)$ are given by

$$S^0(\Omega; \mathcal{K}(f)) = \{hf : \|h\|_{L^\infty(\Omega)} \leq 1\}.$$

Perhaps unexpectedly, it turns out that the boundedness properties of T_E in a space \mathbf{X} are equivalent to its boundedness properties in $\mathbf{X}[\mathcal{K}]$. We will prove this equivalence for more general averaging operators. Given a collection \mathcal{P} measurable sets in Ω satisfying $0 < \mu(E) < \infty$ for all $E \in \mathcal{P}$, we define $T_{\mathcal{P}} := \sum_{E \in \mathcal{P}} T_E$, i.e.,

$$T_{\mathcal{P}} F := \sum_{E \in \mathcal{P}} \langle F \rangle_E \mathbf{1}_E.$$

Then we have the following result.

Theorem 4.4. *Let \mathbf{X} be an \mathbf{F}^n -directional Banach function space over Ω , and let \mathcal{P} be a pairwise disjoint collection of measurable sets in Ω for which $0 < \mu(E) < \infty$ for all $E \in \mathcal{P}$. Then the following are equivalent:*

- (i) $T_{\mathcal{P}} : \mathbf{X} \rightarrow \mathbf{X}$;
- (ii) $T_{\mathcal{P}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$.

Moreover, in this case we have

$$\|T_{\mathcal{P}}\|_{\mathbf{X} \rightarrow \mathbf{X}} \approx_n \|T_{\mathcal{P}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}.$$

Proof. For (i) \Rightarrow (ii), Let $f \in \mathbf{X}$. Since, by Proposition 3.5, we have $\mathcal{K}(f) \in \mathbf{X}[\mathcal{K}]$ with $\|\mathcal{K}(f)\|_{\mathbf{X}[\mathcal{K}]} = \|f\|_{\mathbf{X}}$, noting that $T_{\mathcal{P}}f \in S^0(\Omega; T_{\mathcal{P}}(\mathcal{K}(f)))$, we have

$$\begin{aligned} \|T_{\mathcal{P}}f\|_{\mathbf{X}} &\leq \|T_{\mathcal{P}}(\mathcal{K}(f))\|_{\mathbf{X}} \leq \|T_{\mathcal{P}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|\mathcal{K}(f)\|_{\mathbf{X}[\mathcal{K}]} \\ &= \|T_{\mathcal{P}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|f\|_{\mathbf{X}}. \end{aligned}$$

Thus, $T_{\mathcal{P}} : \mathbf{X} \rightarrow \mathbf{X}$ with $\|T_{\mathcal{P}}\|_{\mathbf{X} \rightarrow \mathbf{X}} \leq \|T_{\mathcal{P}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}$.

For (ii) \Rightarrow (i), let $F \in \mathbf{X}[\mathcal{K}]$. We claim that $\langle F \rangle_E$ is a bounded set in \mathbf{F}^n for all $E \in \mathcal{P}$. Indeed, since $T_E = \mathbf{1}_E T_{\mathcal{P}} : \mathbf{X} \rightarrow \mathbf{X}$, the mapping $u \mapsto \|\mathbf{1}_E u\|_{\mathbf{X}}$ is a norm on \mathbf{F}^n . Thus, there is a constant $c = c_{\mathbf{X}, E} > 0$ for which $\|\mathbf{1}_E u\|_{\mathbf{X}} \geq c|u|$ for all $u \in \mathbf{F}^n$. Let $f \in S^0(\Omega; F \mathbf{1}_E)$ and set $u = \frac{\langle f \rangle_E}{|\langle f \rangle_E|}$. Then

$$c|\langle f \rangle_E| \leq |\langle f \rangle_E| \|\mathbf{1}_E u\|_{\mathbf{X}} = \|T_E f\|_{\mathbf{X}} \leq \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} \|F\|_{\mathbf{X}[\mathcal{K}]},$$

proving the claim. We conclude that $\langle F \rangle_E \in \mathcal{K}_b$. Thus, by the John ellipsoid theorem (see [NPTV17, Section 2.2.1] for a more precise definition), there is a matrix $A_E \in \mathbf{F}^{n \times n}$ for which

$$(4.1) \quad \{A_E u : |u| \leq 1\} \subseteq \langle F \rangle_E \subseteq \{A_E u : |u| \leq n^{\frac{1}{2}}\}.$$

Denoting the columns of A_E by $(v_k^E)_{k=1}^n$, since $v_k^E = A_E e_k \in \langle F \rangle_E$ by the first inclusion in (4.1), we have $v_k^E = \langle f_k^E \rangle_E$ for some $f_k^E \in S^0(\Omega; F \mathbf{1}_E)$. Now, let $g \in S^0(\Omega; T_{\mathcal{P}}F)$. Then, we have $g = \sum_{E \in \mathcal{P}} g_E$ with $g_E = g \mathbf{1}_E \in S^0(\Omega; T_E F)$, and by the second inclusion in (4.1), g_E is of the form

$$g_E(x) = \sum_{k=1}^n h_k^E(x) \langle f_k^E \rangle_E \mathbf{1}_E(x),$$

where $h^E = (h_1^E, \dots, h_n^E)$ satisfies $|h^E(x)| \leq n^{\frac{1}{2}}$. Since \mathcal{P} is pairwise disjoint, setting $h_k(x) := \sum_{E \in \mathcal{P}} h_k^E(x) \mathbf{1}_E(x)$ and $f_k(x) := \sum_{E \in \mathcal{P}} f_k^E(x) \mathbf{1}_E(x)$, we have $|h_k(x)| \leq n^{\frac{1}{2}}$ and

$$g(x) = \sum_{E \in \mathcal{P}} \sum_{k=1}^n h_k^E(x) \langle f_k^E \rangle_E \mathbf{1}_E(x) = \sum_{k=1}^n h_k(x) T_{\mathcal{P}} f_k(x).$$

Hence, by the directional ideal property and the fact that $f_1, \dots, f_n \in S^0(\Omega; F)$,

$$\begin{aligned} \|g\|_{\mathbf{X}} &\leq \sum_{k=1}^n \|h_k T_{\mathcal{P}} f_k\|_{\mathbf{X}} \leq n^{\frac{1}{2}} \sum_{k=1}^n \|T_{\mathcal{P}} f_k\|_{\mathbf{X}} \\ &\leq n^{\frac{1}{2}} \|T_{\mathcal{P}}\|_{\mathbf{X} \rightarrow \mathbf{X}} \sum_{k=1}^n \|f_k\|_{\mathbf{X}} \leq n^{\frac{3}{2}} \|T_E\|_{\mathbf{X} \rightarrow \mathbf{X}} \|F\|_{\mathbf{X}[\mathcal{K}]}. \end{aligned}$$

This proves that $T_{\mathcal{P}}F \in \mathbf{X}[\mathcal{K}]$ with

$$\|T_{\mathcal{P}}F\|_{\mathbf{X}[\mathcal{K}]} \lesssim_n \|T_{\mathcal{P}}\|_{\mathbf{X} \rightarrow \mathbf{X}} \|F\|_{\mathbf{X}[\mathcal{K}]},$$

as desired. \square

As a consequence, we obtain the following characterizations:

Corollary 4.5. *Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space over Ω . Then the following are equivalent:*

- (i) $T_E : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$;
- (ii) $T_E : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]$;
- (iii) $T_E : \mathbf{X} \rightarrow \mathbf{X}$;
- (iv) $T_E : \mathbf{X}' \rightarrow \mathbf{X}'$;
- (v) For all $u \in \mathbf{F}^n$ we have $\mathbf{1}_E u \in \mathbf{X}$, and there is a $C_1 > 0$ such that for all $u \in \mathbf{F}^n$ there is a $v \in \mathbf{F}^n$ for which $\mathbf{1}_E v \in \mathbf{X}'$ and

$$\mu(E)^{-1} \|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} \leq C_1 |u \cdot v|;$$

(vi) or all $v \in \mathbf{F}^n$ we have $\mathbf{1}_E v \in \mathbf{X}'$, and there is a $C_2 > 0$ such that for all $v \in \mathbf{F}^n$ there is a $u \in \mathbf{F}^n$ for which $\mathbf{1}_E u \in \mathbf{X}'$ and

$$\mu(E)^{-1} \|\mathbf{1}_E u\|_{\mathbf{X}} \|\mathbf{1}_E v\|_{\mathbf{X}'} \leq C_2 |u \cdot v|.$$

(vii) $\mathbf{1}_E u \in \mathbf{X}, \mathbf{X}'$ for all $u \in \mathbf{F}^n$.

Moreover, in this case all the associated constants (where for (vii) the associated constant is $\|A_{\mathbf{X},E} A_{\mathbf{X}',E}\|_{\mathbf{F}^{n \times n}}$) are equivalent up to a factor only depending on n .

Proof. The equivalences (i) \Leftrightarrow (iii) and (ii) \Leftrightarrow (iv) follows from Theorem 4.4 with $\mathcal{P} = \{E\}$. The remaining equivalences follows from Corollary 4.2 and Proposition 4.3. \square

5. THE MUCKENHOUPPT CONDITION

We now specify to $\Omega = \mathbf{R}^d$ with the Lebesgue measure. By a cube Q we will mean a cube in \mathbf{R}^d whose sides are parallel to the coordinate axes. We define the Muckenhoupt condition of an \mathbf{F}^n -directional quasi-Banach function space through the averaging operators $T_Q f = \langle f \rangle_Q \mathbf{1}_Q$.

Definition 5.1. We say that an \mathbf{F}^n -directional quasi-Banach function space over \mathbf{R}^d satisfies the *Muckenhoupt condition* if $T_Q : \mathbf{X} \rightarrow \mathbf{X}$ for all cubes Q , and

$$[\mathbf{X}]_A := \sup_Q \|T_Q\|_{\mathbf{X} \rightarrow \mathbf{X}} < \infty.$$

In this case we write $\mathbf{X} \in A$.

By Corollary 4.5 we have the following characterizations of the Muckenhoupt condition:

Theorem 5.2. Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space over \mathbf{R}^d . Then the following are equivalent:

- (i) $\mathbf{X} \in A$;
- (ii) $\mathbf{X}' \in A$;
- (iii) $T_Q : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ for all cubes Q , and

$$[\mathbf{X}]_{A[\mathcal{K}]} := \sup_Q \|T_Q\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} < \infty;$$

- (iv) $T_Q : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]$ for all cubes Q , and

$$[\mathbf{X}']_{A[\mathcal{K}]} := \sup_Q \|T_Q\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]} < \infty;$$

- (v) $\mathbf{1}_Q u \in \mathbf{X}, \mathbf{X}'$ for all cubes Q and $u \in \mathbf{F}^n$, and

$$[\mathbf{X}]_{A_R} := \sup_Q |Q|^{-1} \|A_{\mathbf{X},Q} A_{\mathbf{X}',Q}\|_{\mathbf{F}^{n \times n}} < \infty;$$

- (vi) $\mathbf{1}_Q v \in \mathbf{X}'$ for all cubes Q and $v \in \mathbf{F}^n$, and there is a constant $C_1 > 0$ such that for all cubes Q and all $v \in \mathbf{F}^n$ there is a non-zero $u \in \mathbf{F}^n$ such that $\mathbf{1}_Q u \in \mathbf{X}$, and

$$|Q|^{-1} \|\mathbf{1}_Q u\|_{\mathbf{X}} \|\mathbf{1}_Q v\|_{\mathbf{X}'} \leq C_1 |u \cdot v|;$$

- (vii) $\mathbf{1}_Q u \in \mathbf{X}$ for all cubes Q and $u \in \mathbf{F}^n$, and there is a constant $C_2 > 0$ such that for all cubes Q and all $u \in \mathbf{F}^n$ there is a non-zero $v \in \mathbf{F}^n$ such that $\mathbf{1}_Q v \in \mathbf{X}'$, and

$$|Q|^{-1} \|\mathbf{1}_Q u\|_{\mathbf{X}} \|\mathbf{1}_Q v\|_{\mathbf{X}'} \leq C_2 |u \cdot v|.$$

Moreover, for the optimal C_1, C_2 we have

$$[\mathbf{X}]_A = [\mathbf{X}']_A = C_1 = C_2,$$

and

$$[\mathbf{X}]_A \approx_n [\mathbf{X}]_{A_R} \approx_n [\mathbf{X}]_{A[\mathcal{K}]} \approx_n [\mathbf{X}']_{A[\mathcal{K}]}.$$

Proof. For the equivalence (i) \Leftrightarrow (ii) with equal constant, combining the identity

$$\int_{\Omega} T_Q f \cdot g \, d\mu = |Q| \langle f \rangle_Q \cdot \langle g \rangle_Q = \int_{\Omega} f \cdot T_Q g \, d\mu$$

with Proposition 3.2 and $\mathbf{X}'' = \mathbf{X}$ proves that

$$\|T_Q\|_{\mathbf{X} \rightarrow \mathbf{X}} = \|T_Q\|_{\mathbf{X}' \rightarrow \mathbf{X}'}$$

The equivalence then follows from taking a supremum over all cubes Q . The equivalences of (i) and (iii)-(vii) follow from Corollary 4.5. \square

As is done in [Nie24], we can define the *strong* Muckenhoupt condition $\mathbf{X} \in A_{\text{strong}}$ through

$$[\mathbf{X}]_{A_{\text{strong}}} := \sup_{\mathcal{P}} \|T_{\mathcal{P}}\|_{\mathbf{X} \rightarrow \mathbf{X}},$$

where the supremum is taken over all pairwise disjoint collections of cubes \mathcal{P} , and where $T_{\mathcal{P}} f = \sum_{Q \in \mathcal{P}} \langle f \rangle_Q \mathbf{1}_Q$. Moreover, we also set

$$[\mathbf{X}]_{A_{\text{strong}}[\mathcal{K}]} := \sup_{\mathcal{P}} \|T_{\mathcal{P}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}$$

Analogously to the Muckenhoupt condition, we have the following result:

Proposition 5.3. *Let \mathbf{X} be an \mathbf{F}^n -directional Banach function space over Ω . Then $\mathbf{X} \in A_{\text{strong}}$ if and only if $T_{\mathcal{P}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ uniformly for all pairwise disjoint collections \mathcal{P} , with*

$$[\mathbf{X}]_{A_{\text{strong}}} \sim_n [\mathbf{X}]_{A_{\text{strong}}[\mathcal{K}]}.$$

Proof. This follows directly from Theorem 4.4. \square

Remark 5.4. The proof of Theorem 4.4 fails for *sparse* collections \mathcal{P} . Thus, it is not clear if such an equivalence holds for sparse collections as well.

We define $L_{\text{loc}}^1(\mathbf{R}^d; \mathcal{K})$ to be the space of $F \in L^0(\mathbf{R}^d; \mathcal{K})$ for which $F \mathbf{1}_Q \in L^1(\mathbf{R}^d; \mathcal{K})$ for all cubes Q .

Proposition 5.5. *We have*

$$L_{\text{loc}}^1(\mathbf{R}^d; \mathcal{K}) \subseteq L^0(\mathbf{R}^d; \mathcal{K}_b).$$

Moreover, suppose that \mathbf{X} be an \mathbf{F}^n -directional quasi-Banach function space over \mathbf{R}^d with $\mathbf{X} \in A$. Then

$$\mathbf{X}[\mathcal{K}] \subseteq L_{\text{loc}}^1(\mathbf{R}^d; \mathcal{K}).$$

Proof. Let $F \in L_{\text{loc}}^1(\mathbf{R}^d; \mathcal{K})$. Then for each cube Q we have $F \mathbf{1}_Q \in L^1(\mathbf{R}^d; \mathcal{K})$, so by Proposition 3.9 we have $F(x) \in \mathcal{K}_b$ for a.e. $x \in Q$. Partitioning \mathbf{R}^d into a countable collection of cubes, we conclude that $F(x) \in \mathcal{K}_b$ for a.e. $x \in \mathbf{R}^d$, as desired.

For the second assertion, by Theorem 5.2, we have $T_Q : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ for all cubes Q . In particular, this means that if $F \in \mathbf{X}[\mathcal{K}]$, then $T_Q F$ is well-defined. Hence, $F \mathbf{1}_Q \in L^1(\mathbf{R}^d; \mathcal{K})$. We conclude that $\mathbf{X}[\mathcal{K}] \subseteq L_{\text{loc}}^1(\mathbf{R}^d; \mathcal{K})$. \square

5.1. The convex-set valued maximal operator. Given a collection of cubes \mathcal{P} and $F \in L^0(\Omega; \mathcal{K})$ satisfying $F \in L_{\text{loc}}^1(\mathbf{R}^d; \mathcal{K})$, we define

$$M_{\mathcal{P}}^{\mathcal{K}} F(x) := \sup_{Q \in \mathcal{P}} \langle F \rangle_Q \mathbf{1}_Q(x),$$

where the supremum is taken with respect to the partial ordering given by set inclusion in \mathcal{K} , i.e., $M_{\mathcal{P}}^{\mathcal{K}} F(x)$ is the smallest closed convex set containing $\bigcup_{Q \in \mathcal{P}} T_Q F(x)$. When \mathcal{P} is the collection of all cubes, then we drop the subscript \mathcal{P} and denote the operator by $M^{\mathcal{K}}$.

Just like in the scalar-valued case, to prove the boundedness of $M^{\mathcal{K}}$ it suffices to prove the boundedness $M_{\mathcal{D}}^{\mathcal{K}}$ for 3^d translated dyadic grids \mathcal{D} . For the details, we refer the reader to [BC23, Lemma 5.9].

Note that when \mathcal{P} is finite, the set $\bigcup_{Q \in \mathcal{P}} T_Q F(x)$ is bounded, from which we conclude that $M_{\mathcal{P}}^{\mathcal{K}} F(x) \in \mathcal{K}$. However, if \mathcal{P} is infinite, this need not be the case. Nonetheless, if an \mathbf{F}^n -directional quasi-Banach function space \mathbf{X} over \mathbf{R}^d satisfies the monotone convergence property, then we have

$$M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$$

if and only if we have

$$M_{\mathcal{F}}^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$$

uniformly for all finite collections of cubes \mathcal{F} , and

$$\|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} = \sup_{\mathcal{F}} \|M_{\mathcal{K}}^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}.$$

As an analogue to [Nie24, Theorem B], we prove the following result:

Theorem 5.6. *Let \mathbf{X} be an \mathbf{F}^n -directional Banach function space over \mathbf{R}^d with the Fatou property. Consider the following statements:*

- (a) $M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$;
- (b) $\mathbf{X} \in A_{\text{strong}}$;
- (c) $M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}_{\text{weak}}[\mathcal{K}]$;
- (d) $\mathbf{X} \in A$.

Then (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d) with

$$[\mathbf{X}]_A \leq \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}_{\text{weak}}[\mathcal{K}]} \lesssim_{d,n} [\mathbf{X}]_{A_{\text{strong}}} \leq \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}.$$

Furthermore, if there is a $C \geq 1$ such that for all pairwise disjoint collections of cubes \mathcal{P} and all $f \in \mathbf{X}$, $g \in \mathbf{X}'$ we have

$$\sum_{Q \in \mathcal{P}} \|\mathbf{1}_Q f\|_{\mathbf{X}} \|\mathbf{1}_Q g\|_{\mathbf{X}'} \leq C \|f\|_{\mathbf{X}} \|g\|_{\mathbf{X}'},$$

then (b)-(d) are equivalent, with

$$[\mathbf{X}]_{A_{\text{strong}}} \leq C [\mathbf{X}]_A.$$

Proof. For (a) \Rightarrow (b), let \mathcal{P} be a pairwise disjoint collection of cubes and let $f \in \mathbf{X}$. If $x \in \mathbf{R}^d$ satisfies $x \in Q$ for some $Q \in \mathcal{P}$, then $T_{\mathcal{P}} f(x) = \langle f \rangle_Q \mathbf{1}_Q(x) \in \bigcup_Q \langle f \rangle_Q \mathbf{1}_Q(x)$. Thus, we have

$$T_{\mathcal{P}} f(x) \in M^{\mathcal{K}}(\mathcal{K}(f))(x)$$

a.e. Hence, by the ideal property in $\mathbf{X}[\mathcal{K}]$ and Proposition 3.5, we have

$$\|T_{\mathcal{P}} f\|_{\mathbf{X}} \leq \|M^{\mathcal{K}}(\mathcal{K}(f))\|_{\mathbf{X}[\mathcal{K}]} \leq \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|f\|_{\mathbf{X}}.$$

Thus, $\mathbf{X} \in A_{\text{strong}}$ and $[\mathbf{X}]_{A_{\text{strong}}} \leq \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}$.

For (b) \Rightarrow (c), first note that by the monotone convergence property and a 3^d -lattice reduction, it suffices to prove that $M_{\mathcal{F}}^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}_{\text{weak}}[\mathcal{K}]$ uniformly for finite collections \mathcal{F} contained in a dyadic grid \mathcal{D} . Let $u \in \mathbf{F}^n$ be non-zero and suppose that $x \in M_{\mathcal{F}}^{\mathcal{K}} F^{-1}(\{u\})$. Since the convex hull of a compact set in \mathbf{F}^n is compact – a consequence of Caratheodory's convexity theorem – we have

$$u \in M_{\mathcal{F}}^{\mathcal{K}} F(x) = \text{conv} \left(\bigcup_{Q \in \mathcal{F}} \langle F \rangle_Q \mathbf{1}_Q(x) \right).$$

Thus, there are $\theta_1, \dots, \theta_J \geq 0$ with $\sum_{j=1}^J \theta_j = 1$ and non-zero $u_1, \dots, u_J \in \mathbf{F}^n$ with $u_j \in \bigcup_{Q \in \mathcal{F}} \langle F \rangle_Q$ such that $u = \sum_{j=1}^J \theta_j u_j$. For each j we let \mathcal{P}_j denote the collection of maximal cubes $Q \in \mathcal{F}$ for which $u_j \in \langle F \rangle_Q$. Then

$$\mathbf{1}_{M_{\mathcal{F}}^{\mathcal{K}} F^{-1}(\{u\})} u \in \sum_{j=1}^J \theta_j \sum_{Q \in \mathcal{P}_j} \langle F \rangle_Q \mathbf{1}_Q,$$

so, by the triangle inequality (Proposition 3.11, the directional ideal property of $\mathbf{X}[\mathcal{K}]$, and the fact that each \mathcal{P}_j is pairwise disjoint,

$$\|\mathbf{1}_{M_{\mathcal{F}}^{\mathcal{K}} F^{-1}(\{u\})} u\|_{\mathbf{X}} \leq \sum_{j=1}^J \theta_j \|T_{\mathcal{P}_j} F\|_{\mathbf{X}[\mathcal{K}]} \leq [\mathbf{X}]_{A_{\text{strong}}[\mathcal{K}]} \|F\|_{\mathbf{X}}.$$

Taking a supremum over all $u \in \mathbf{F}^n$ now proves the result.

For (c) \Rightarrow (d), let Q be a cube and let $f \in \mathbf{X}$. If $x \in Q$, then $\langle f \rangle_Q \in M^{\mathcal{K}}(\mathcal{K}(f))(x)$. Hence, for $u = \langle f \rangle_Q$ we have

$$T_Q f(x) = \langle f \rangle_Q \mathbf{1}_Q(x) \in \mathbf{1}_{M^{\mathcal{K}}(\mathcal{K}(f))^{-1}(\{u\})}(x) u$$

a.e. It follows from the ideal property of $\mathbf{X}[\mathcal{K}]$ and Proposition 3.5 that

$$\|T_Q f\|_{\mathbf{X}} \leq \|M^{\mathcal{K}}(\mathcal{K}(f))\|_{\mathbf{X}_{\text{weak}}[\mathcal{K}]} \leq \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}_{\text{weak}}[\mathcal{K}]} \|f\|_{\mathbf{X}},$$

proving that $\mathbf{X} \in A$ with $[\mathbf{X}]_A \leq \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}_{\text{weak}}[\mathcal{K}]}$. The result follows.

Finally, let $\mathbf{X} \in A$ and let \mathcal{P} be a pairwise disjoint collection of cubes. Then

$$\begin{aligned} \left| \int_{\mathbf{R}^d} T_{\mathcal{P}} f \cdot g \, dx \right| &\leq \sum_{Q \in \mathcal{P}} \int_{\mathbf{R}^d} |T_Q f \cdot \mathbf{1}_Q g| \, dx \leq \sum_{Q \in \mathcal{P}} \|T_Q f\|_{\mathbf{X}} \|\mathbf{1}_Q g\|_{\mathbf{X}'} \\ &\leq [\mathbf{X}]_A \sum_{Q \in \mathcal{P}} \|\mathbf{1}_Q f\|_{\mathbf{X}} \|\mathbf{1}_Q g\|_{\mathbf{X}'} \leq C[\mathbf{X}]_A \|f\|_{\mathbf{X}} \|g\|_{\mathbf{X}'}. \end{aligned}$$

Thus, by Proposition 3.2 and Lemma A.3, $\mathbf{X} \in A_{\text{strong}}$ with $[\mathbf{X}]_{A_{\text{strong}}} \leq C[\mathbf{X}]_A$. \square

For matrix weighted spaces, this yields the following corollary:

Corollary 5.7. *Let X be a Banach function space over \mathbf{R}^d for which there is a $C \geq 1$ such that for all pairwise disjoint \mathcal{P} and all $f \in X$, $g \in X'$ we have*

$$(5.1) \quad \sum_{Q \in \mathcal{P}} \|\mathbf{1}_Q f\|_X \|\mathbf{1}_Q g\|_{X'} \leq C \|f\|_X \|g\|_{X'}.$$

If $W : \mathbf{R}^d \rightarrow \mathbf{F}^{n \times n}$ is a matrix weight, then then the following are equivalent:

- (i) $X_W \in A$;
- (ii) $X_W \in A_{\text{strong}}$;
- (iii) $M^{\mathcal{K}} : X_W[\mathcal{K}] \rightarrow (X_W)_{\text{weak}}[\mathcal{K}]$.

Moreover, in this case we have

$$[X_W]_A = [X_W]_{A_{\text{strong}}} \approx_{d,n} \|M^{\mathcal{K}}\|_{X_W[\mathcal{K}] \rightarrow (X_W)_{\text{weak}}[\mathcal{K}]}.$$

Proof. By Proposition 3.7 we have

$$\begin{aligned} \sum_{Q \in \mathcal{P}} \|\mathbf{1}_Q f\|_{X_W} \|\mathbf{1}_Q g\|_{(X_W)'} &= \sum_{Q \in \mathcal{P}} \|\mathbf{1}_Q |Wf|\|_X \|\mathbf{1}_Q |W^{-1}g|\|_{X'} \\ &\leq C \| |Wf| \|_X \| |W^{-1}g| \|_{X'} = C \|f\|_{X_W} \|g\|_{(X_W)'}. \end{aligned}$$

Hence, the result follows from Theorem 5.6. \square

Example 5.8. If $X = L^p(\mathbf{R}^d)$, then (5.1) is satisfied with $C = 1$ by Hölder's inequality. Thus, Corollary 5.7 implies that

$$\|M^{\mathcal{K}}\|_{L^p_W(\mathbf{R}^d; \mathcal{K}) \rightarrow L^p_W(\mathbf{R}^d; \mathcal{K})_{\text{weak}}} \sim_{d,n} [W]_p.$$

In the work [Nie24] the property (5.1) is referred to as property $X \in \mathcal{G}$, and we refer the reader to this work for an overview of further spaces with this property.

Finally, we prove that if $M^{\mathcal{K}}$ is bounded on a space $\mathbf{X}[\mathcal{K}]$, then every $F \in \mathbf{X}[\mathcal{K}]$ satisfies $F(x) \in \mathcal{K}_b$ for a.e. $x \in \mathbf{R}^d$.

Proposition 5.9. *Let \mathbf{X} be an \mathbf{F}^n -directional quasi-Banach function space for which $M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$. Then $\mathbf{X}[\mathcal{K}] \subseteq L^0(\mathbf{R}^d; \mathcal{K}_b)$.*

Proof. Since $M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ implies $\mathbf{X} \in A$ by Theorem 5.6, this follows from Proposition 5.5. \square

6. EXTRAPOLATION

Theorem 6.1. *Let $1 \leq p \leq \infty$, let V be a set, and let $S : V \rightarrow L^0(\Omega; \mathbf{F}^n)$. Suppose*

$$T : \bigcup_{W \in A_p} S^{-1}(L^p_W(\mathbf{R}^d; \mathbf{F}^n)) \rightarrow L^0(\mathbf{R}^d; \mathbf{F}^n)$$

is a map for which there is an increasing function $\phi : [0, \infty) \rightarrow [0, \infty)$ such that for all $W \in A_p$ and all $f \in V$ with $Sf \in L^p_W(\mathbf{R}^d; \mathbf{F}^n)$ we have

$$\|Tf\|_{L^p_W(\mathbf{R}^d; \mathbf{F}^n)} \leq \phi([W]_p) \|Sf\|_{L^p_W(\mathbf{R}^d; \mathbf{F}^n)}.$$

Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space over \mathbf{R}^d for which

$$M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}], \quad M^{\mathcal{K}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}].$$

Then Tf is well-defined for all $f \in V$ with $Sf \in \mathbf{X}$, and

$$\|Tf\|_{\mathbf{X}} \lesssim_n \phi(C_n \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]})^{\frac{1}{p'}} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]}^{\frac{1}{p}} \|Sf\|_{\mathbf{X}}.$$

When $p = \infty$ or $p = 1$, we can omit the bound $M^{\mathcal{K}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]$ or $M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ respectively.

The proof follows from two abstract theorems. The first is the following generalization of the Rubio de Francia algorithm of [BC23, Theorem 7.6]. Recall that we have defined the sum $F + G$ through the closed Minkowski sum

$$(F + G)(x) := \overline{\{u + v : u \in F(x), v \in G(x)\}}.$$

Unlike [BC23, Theorem 7.6], we do not require the operator T to be monotone.

Theorem 6.2 (Rubio de Francia algorithm). *Let \mathbf{X} be an \mathbf{F}^n -directional Banach function space over \mathbf{R}^d with the Fatou property, and let $T : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ be an operator that is sublinear in the sense that*

$$T(F + G)(x) \subseteq TF(x) + TG(x), \quad T(\lambda F)(x) = \lambda TF(x)$$

a.e. for all $F, G \in \mathbf{X}[\mathcal{K}]$ and $\lambda \in \mathbf{F}$. Then there is a measurable mapping $\mathcal{R} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ for which $\|\mathcal{R}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \leq 2$, and for all $F \in \mathbf{X}[\mathcal{K}]$

- (a) $F(x) \subseteq \mathcal{R}F(x)$ a.e.;
- (b) $T(\mathcal{R}F)(x) \subseteq 2\|T\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \mathcal{R}F(x)$ a.e.

For the proof, we require a lemma on nested sets in \mathcal{K} :

Lemma 6.3. *Let $(B_k)_{k \geq 1}$ be a sequence in \mathcal{K} satisfying $B_{k+1} \subseteq B_k$ for all $k \geq 1$. If $\bigcap_{k=1}^{\infty} B_k = \{0\}$, then there is a $K \geq 1$ such that for all $k \geq K$ the set B_k is compact. Moreover, if $(x_k)_{k \geq 1}$ is a sequence for which $x_k \in B_k$ for all $k \geq 1$, then $x_k \rightarrow 0$.*

Proof. Arguing by contrapositive, suppose that B_k is unbounded for all $k \geq 1$. Set $S^{n-1} := \{u \in \mathbf{F}^n : |u| = 1\}$ and, for $k \geq 1$,

$$C_k := S^{n-1} \cap B_k.$$

Then C_k is an intersection of closed sets, and hence, closed. Moreover, as each B_k is unbounded, $C_k \neq \emptyset$. As $C_{k+1} \subseteq C_k$ for all $k \geq 1$ and S^{n-1} is compact, we have

$$S^{n-1} \cap \bigcap_{k=1}^{\infty} B_k = \bigcap_{k=1}^{\infty} C_k \neq \emptyset.$$

Thus, $\bigcap_{k=1}^{\infty} B_k \neq \{0\}$, as asserted.

For the second assertion, let $(x_{j_k})_{k \geq 1}$ be a subsequence. Since for j large enough, B_j is compact, this subsequence has a further convergent subsequence. As this sequence eventually lies in B_j for all $j \geq 1$, the limit is contained in $\bigcap_{k=1}^{\infty} B_k \neq \{0\}$. Hence, every subsequence of $(x_k)_{k \geq 1}$ has a further subsequence converging to 0. This proves that $x_k \rightarrow 0$, as desired. \square

Proof of Theorem 6.2. Let $F \in \mathbf{X}[\mathcal{K}]$ and define

$$F_k(x) := 2^{-k} \|T\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}^{-k} T^k F(x),$$

where T^k is iteratively defined as $T^0 F := F$ and $T^{k+1} F := T(T^k F)$. Then

$$\sum_{k=0}^{\infty} \|F_k\|_{\mathbf{X}[\mathcal{K}]} \leq \sum_{k=0}^{\infty} 2^{-k} \|F\|_{\mathbf{X}[\mathcal{K}]} = 2 \|F\|_{\mathbf{X}[\mathcal{K}]},$$

so, by Theorem 3.12, the series

$$(6.1) \quad \mathcal{R}F(x) := \sum_{k=0}^{\infty} F_k(x) = \overline{\bigcup_{K=0}^{\infty} \left\{ \sum_{k=0}^K u_k : u_k \in F_k(x) \right\}}$$

satisfies $\mathcal{R}F \in \mathbf{X}[\mathcal{K}]$ with

$$\|\mathcal{R}F\|_{\mathbf{X}[\mathcal{K}]} \leq 2 \|F\|_{\mathbf{X}[\mathcal{K}]}.$$

The assertion (a) follows from the fact that the $K = 0$ term of the union in (6.1) is equal to $F(x)$. For (b), sublinearity of T implies that

$$TF_k(x) = 2^{-k} \|T\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}^{-k} T^{k+1} F(x) = 2 \|T\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} F_{k+1}(x)$$

a.e. so that, since for all $J \geq 1$ we have $\mathcal{R}F = \sum_{k=0}^J F_k + S_J$ where $S_J := \sum_{k=J+1}^{\infty} F_k$,

$$(6.2) \quad \begin{aligned} T(\mathcal{R}F)(x) &\subseteq 2 \|T\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \sum_{k=0}^J F_{k+1}(x) + TS_J(x) \\ &\subseteq 2 \|T\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \mathcal{R}F(x) + TS_J(x). \end{aligned}$$

Note that

$$\begin{aligned} \left\| \bigcap_{J=1}^{\infty} TS_J \right\|_{\mathbf{X}[\mathcal{K}]} &\leq \|TS_J\|_{\mathbf{X}[\mathcal{K}]} \leq \|T\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \left\| \sum_{k=J+1}^{\infty} F_k \right\|_{\mathbf{X}[\mathcal{K}]} \\ &\leq \|T\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \sum_{k=J+1}^{\infty} \|F_k\|_{\mathbf{X}[\mathcal{K}]} \rightarrow 0 \end{aligned}$$

as $J \rightarrow \infty$ and, hence, $\bigcap_{J=1}^{\infty} TS_J(x) = \{0\}$.

Since $TS_J(x) \in \mathcal{K}$, by Lemma 6.3 there is a $J_0 \geq 1$ such that for all $J \geq J_0$, the set $TS_J(x)$ is compact. Let $u \in T(\mathcal{R}F)(x)$. Since the Minkowski sum of a closed and a compact set is closed, this means that, by (6.2), for all $J \geq J_0$ we can write $u = v_J + w_J$,

with $v_J \in 2\|T\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \mathcal{R}F(x)$ and $w_J \in TS_J(x)$. Then, since $w_J \rightarrow 0$ as $J \rightarrow \infty$, we have $v_J = u - w_J \rightarrow u$. Hence, $u \in 2\|T\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \mathcal{R}F(x)$. This proves (b), as desired. \square

The second theorem we need for the extrapolation result is the following application of the Rubio de Francia algorithm:

Theorem 6.4. *Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space over \mathbf{R}^d for which*

$$M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}], \quad M^{\mathcal{K}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}].$$

Let $1 \leq p \leq \infty$. Then for all $f \in \mathbf{X}$, $g \in \mathbf{X}'$, there exists a matrix weight $W \in A_p$ with the following properties:

- $[W]_p \lesssim_n \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}^{\frac{1}{p}} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]}^{\frac{1}{p'}}$;
- $\|f\|_{L_W^p(\mathbf{R}^d; \mathbf{F}^n)} \|g\|_{L_{W^{-1}}^{p'}(\mathbf{R}^d; \mathbf{F}^n)} \lesssim_n \|f\|_{\mathbf{X}} \|g\|_{\mathbf{X}'}$.

When $p = \infty$ or $p = 1$, we can omit the bound $M^{\mathcal{K}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]$ or $M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ respectively.

For the proof, we adopt a function norm approach. We call a mapping $\rho : \mathbf{R}^d \times \mathbf{F}^n \rightarrow [0, \infty)$ a function norm if for a.e. $x \in \mathbf{R}^d$ the mapping $u \mapsto \rho(x, u)$ is a norm, and for all $u \in \mathbf{F}^n$ the mapping $x \mapsto \rho(x, u)$ is measurable. We define $L_\rho^p(\mathbf{R}^d)$ as the space of functions $f \in L^0(\mathbf{R}^d; \mathbf{F}^n)$ for which $\rho(x, f(x)) \in L^p(\mathbf{R}^d)$, with

$$\|f\|_{L_\rho^p(\mathbf{R}^d)} := \left(\int_{\mathbf{R}^d} \rho(x, f(x))^p dx \right)^{\frac{1}{p}}.$$

Then, for $1 \leq p \leq \infty$, $L_\rho^p(\mathbf{R}^d)$ is an \mathbf{F}^n -directional Banach function space over \mathbf{R}^d . Moreover, $L_\rho^p(\mathbf{R}^d)' = L_{\rho^*}^{p'}(\mathbf{R}^d)$, where $\rho^* : \mathbf{R}^d \times \mathbf{F}^n \rightarrow [0, \infty)$ is defined as

$$\rho^*(x, v) = \sup_{u \in \mathbf{F}^n \setminus \{0\}} \frac{|u \cdot v|}{\rho(x, u)}.$$

If $W : \mathbf{R}^d \rightarrow \mathbf{F}^{n \times n}$ is a matrix weight, then

$$\rho_W(x, u) := |W(x)u|$$

is a norm function, and $L_{\rho_W}^p(\mathbf{R}^d) = L_W^p(\mathbf{R}^d; \mathbf{F}^n)$. Conversely, for any norm function ρ it follows from the John ellipsoid theorem that there is a matrix weight W for which

$$(6.3) \quad \rho_W(x, u) \leq \rho(x, u) \leq n^{\frac{1}{2}} \rho_W(x, u).$$

We will write $\rho \in A_p$ if there is a constant $C \geq 1$ such that for all cubes Q and all $u \in \mathbf{F}^n$ there is a non-zero $v \in \mathbf{F}^n$ such that

$$\left(\frac{1}{|Q|} \int_Q \rho(x, u)^p dx \right)^{\frac{1}{p}} \left(\frac{1}{|Q|} \int_Q \rho^*(x, v)^{p'} dx \right)^{\frac{1}{p'}} \leq C |u \cdot v|,$$

where for an infinite exponent the integral is interpreted as an essential supremum. The smallest possible C is denoted by $[\rho]_p$. As this condition is precisely the condition $L_\rho^p(\mathbf{R}^d) \in A$, it follows from Theorem 5.2 that $\rho \in A_p$ if and only if $\rho^* \in A_{p'}$, with $[\rho]_p = [\rho^*]_{p'}$.

If W is a matrix weight, then $[\rho_W]_p = [W]_p$. In general, we have $\rho \in A_p$ if and only if $W \in A_p$ for W satisfying (6.3), with $[\rho]_p \approx_n [W]_p$.

Before we get to the proof of Theorem 6.4, we will need an interpolation result. The following lemma is a version of the direct implication in [BC23, Theorem 7.3], as well as a norm function version of [BC23, Proposition 8.7]:

Lemma 6.5. *Let $F_0, F_1 \in L^0(\Omega; \mathcal{K})$ satisfy $F_0 \mathbf{1}_Q, F_1 \mathbf{1}_Q \in L^1(\mathbf{R}^d; \mathcal{K})$ for all cubes Q , and suppose there are $C_0, C_1 > 0$ such that for a.e. $x \in \mathbf{R}^d$ we have*

$$M^{\mathcal{K}} F_0(x) \subseteq C_0 F_0(x), \quad M^{\mathcal{K}} F_1(x) \subseteq C_1 F_1(x)$$

Then

$$\rho_0(x, u) := \sup_{v \in F_0(x)} |u \cdot v|, \quad \rho_1(x, u) := \sup_{v \in F_1(x)} |u \cdot v|$$

satisfy $\rho_0, \rho_1 \in A_1$ with $[\rho_0]_1 \lesssim_n C_0$, $[\rho_1]_1 \lesssim_n C_1$.

Moreover, for all $1 \leq p \leq \infty$, there exists a norm function ρ satisfying

- (a) $\rho \in A_p$ with $[\rho]_p \lesssim_n C_0^{\frac{1}{p'}} C_1^{\frac{1}{p}}$;
- (b) $\rho(x, u) \leq \rho_0^*(x, u)^{\frac{1}{p'}} \rho_1(x, u)^{\frac{1}{p}}$ a.e. for all $u \in \mathbf{F}^n$;
- (c) $\rho^*(x, v) \leq \rho_0(x, v)^{\frac{1}{p'}} \rho_1^*(x, v)^{\frac{1}{p}}$ a.e. for all $v \in \mathbf{F}^n$.

Proof. For the first assertion, by Proposition 5.9 we have $F_0(x), F_1(x) \in \mathcal{K}_b$ for a.e. $x \in \mathbf{R}^d$, proving that ρ_0 and ρ_1 are finite a.e. Thus, we may apply [BC23, Theorem 7.3] to conclude that $\rho_0, \rho_1 \in A_1$ with

$$[\rho_0]_1 \lesssim_n C_0, \quad [\rho_1]_1 \lesssim_n C_1.$$

For the second assertion, set

$$K(x) := \mathcal{K}(\{u \in \mathbf{F}^n : \rho_0^*(x, u)^{\frac{1}{p'}} \rho_1(x, u)^{\frac{1}{p}} \leq 1\}).$$

This set is an absorbing set in \mathcal{K}_b , so we may define $\rho : \mathbf{R}^d \times \mathbf{F}^n \rightarrow [0, \infty)$ as the Minkowski functional

$$\rho(x, u) := \inf\{t > 0 : t^{-1}u \in K(x)\}.$$

Then, since $t^{-1}u \in \{u \in \mathbf{F}^n : \rho_0^*(x, u)^{\frac{1}{p'}} \rho_1(x, u)^{\frac{1}{p}} \leq 1\}$ for $t = \rho_0^*(x, u)^{\frac{1}{p'}} \rho_1(x, u)^{\frac{1}{p}}$, we have

$$\rho(x, u) \leq \rho_0^*(x, u)^{\frac{1}{p'}} \rho_1(x, u)^{\frac{1}{p}},$$

proving (b). For (c), note that

$$|u \cdot v| = |u \cdot v|^{\frac{1}{p'}} |u \cdot v|^{\frac{1}{p}} \leq (\rho_0^*(x, u) \rho_0(x, v))^{\frac{1}{p'}} (\rho_1(x, u) \rho_1^*(x, v))^{\frac{1}{p}}.$$

It follows that if $t > 0$ satisfies $t^{-1}u \in K(x)$, i.e., $t^{-1}u = \sum_{k=1}^K \theta_k u_k$ for $\sum_{k=1}^K \theta_k = 1$, $\rho_0^*(x, u_k)^{\frac{1}{p'}} \rho_1(x, u_k)^{\frac{1}{p}} \leq 1$, we have

$$t^{-1}|u \cdot v| \leq \sum_{k=1}^K \theta_k |u_k \cdot v| \leq \sum_{k=1}^K \theta_k \rho_0^*(x, v)^{\frac{1}{p'}} \rho_1(x, v)^{\frac{1}{p}} = \rho_0^*(x, v)^{\frac{1}{p'}} \rho_1(x, v)^{\frac{1}{p}}.$$

This implies that $|u \cdot v| \leq \rho_0^*(x, v)^{\frac{1}{p'}} \rho_1(x, v)^{\frac{1}{p}} \rho(x, u)$ and, hence,

$$\rho^*(x, v) = \sup_{u \neq 0} \frac{|u \cdot v|}{\rho(x, u)} \leq \rho_0(x, v)^{\frac{1}{p'}} \rho_1^*(x, v)^{\frac{1}{p}},$$

as desired. Finally, for (a), it follows from [BC23, Remark 2.16, Remark 8.10] that $\rho \in A_p$ with the desired bound. The assertion follows. \square

Proof of Theorem 6.4. It suffices to construct a $\rho \in A_p$ instead of $W \in A_p$ for which the desired bounds hold. By applying Theorem 6.2 to $M^{\mathcal{K}}$ in \mathbf{X} and \mathbf{X}' , we obtain the respective operators $\mathcal{R} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ and $\mathcal{R}' : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]$ satisfying (a) and (b). Applying Lemma 6.5 with $F_0 = \mathcal{R}(\mathcal{K}(f))$, $F_1 = \mathcal{R}'(\mathcal{K}(g))$, we obtain a norm function $\rho \in A_p$ for which

$$[\rho]_p \lesssim_n \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}^{\frac{1}{p'}} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]}^{\frac{1}{p}},$$

and (b) and (c) hold for

$$\rho_0(x, u) := \sup_{v \in \mathcal{R}(\mathcal{K}(f))(x)} |u \cdot v|, \quad \rho_1(x, u) := \sup_{v \in \mathcal{R}'(\mathcal{K}(g))(x)} |u \cdot v|.$$

By applying Theorem 3.8 with

$$\phi(x, v) = |f(x) \cdot v|, \quad h(x) = \rho_1(x, f(x)),$$

we can find a $k \in S^0(\mathbf{R}^d; \mathcal{R}'(\mathcal{K}(g)))$ for which

$$|f(x) \cdot k(x)| = \rho_1(x, f(x)).$$

Moreover, since $f(x) \in \mathcal{R}(\mathcal{K}(f))(x)$ a.e., we have $\rho_0(x, u) \geq |f(x) \cdot u|$ a.e., and, hence,

$$\rho_0^*(x, f(x)) \leq 1.$$

Thus, we have

$$\rho(x, f(x))^p \leq \rho_0^*(x, f(x))^{p-1} \rho_1(x, f(x)) \leq |f(x) \cdot k(x)|,$$

or $\rho(x, f(x)) = \rho_0^*(x, f(x)) \leq 1$ if $p = \infty$. We conclude that $f \in L_\rho^p(\mathbf{R}^d)$ with

$$\begin{aligned} \int_{\mathbf{R}^d} \rho(x, f(x))^p dx &\leq \int_{\mathbf{R}^d} |f(x) \cdot k(x)| dx \leq \|f\|_{\mathbf{X}} \|k\|_{\mathbf{X}'} \\ &\leq \|f\|_{\mathbf{X}} \|\mathcal{R}'(\mathcal{K}(g))\|_{\mathbf{X}'[\mathcal{K}]} \leq 2\|f\|_{\mathbf{X}} \|g\|_{\mathbf{X}'}, \end{aligned}$$

or $\|f\|_{L_\rho^\infty(\mathbf{R}^d)} \leq 1$ when $p = \infty$. For the term involving g , an analogous argument as the one for f with the roles of ρ_0 , ρ_1 , and of p , p' reversed, shows that $g \in L_{\rho^*}^{p'}(\mathbf{R}^d)$ with

$$\|g\|_{L_{\rho^*}^{p'}(\mathbf{R}^d)} \leq 2^{\frac{1}{p'}} \|f\|_{\mathbf{X}}^{\frac{1}{p'}} \|g\|_{\mathbf{X}'}^{\frac{1}{p'}}.$$

In conclusion, $f \in L_\rho^p(\mathbf{R}^d)$, $g \in L_{\rho^*}^{p'}(\mathbf{R}^d)$, and

$$\|f\|_{L_\rho^p(\mathbf{R}^d)} \|g\|_{L_{\rho^*}^{p'}(\mathbf{R}^d)} \leq 2\|f\|_{\mathbf{X}} \|g\|_{\mathbf{X}'},$$

as asserted. \square

Proof of Theorem 6.1. Let $f \in V$ for which $Sf \in \mathbf{X}$, and let $g \in \mathbf{X}'$. By Theorem 6.4 we can pick a $W \in A_p$ for which

$$[W]_p \leq C_n \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}^{\frac{1}{p}} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]}^{\frac{1}{p}}$$

and

$$\|Sf\|_{L_W^p(\mathbf{R}^d)} \|g\|_{L_{W^{-1}}^{p'}(\mathbf{R}^d)} \lesssim_n \|Sf\|_{\mathbf{X}} \|g\|_{\mathbf{X}'}.$$

Thus,

$$\begin{aligned} \int_{\mathbf{R}^d} |Tf \cdot g| dx &\leq \|Tf\|_{L_W^p(\mathbf{R}^d)} \|g\|_{L_{W^{-1}}^{p'}(\mathbf{R}^d)} \leq \phi([W]_p) \|Sf\|_{L_W^p(\mathbf{R}^d)} \|g\|_{L_{W^{-1}}^{p'}(\mathbf{R}^d)} \\ &\lesssim_n \phi(C_n \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]}^{\frac{1}{p}} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]}^{\frac{1}{p}}) \|Sf\|_{\mathbf{X}} \|g\|_{\mathbf{X}'}. \end{aligned}$$

As \mathbf{X} is Köthe reflexive, taking a supremum over all $g \in \mathbf{X}'$ with $\|g\|_{\mathbf{X}'} = 1$ proves the result. \square

7. PROOF OF THEOREM B

First, we prove the implication (ii) \Rightarrow (i). This follows from the following result:

Theorem 7.1. *Let \mathbf{X} be a Köthe reflexive \mathbf{F}^n -directional Banach function space over \mathbf{R}^d . Suppose*

$$M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}], \quad M^{\mathcal{K}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}].$$

Then for any sparse collection \mathcal{S} , we have $T_{\mathcal{S}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ with

$$\|T_{\mathcal{S}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \lesssim_n \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]}.$$

Proof. By the monotone convergence property of \mathbf{X} , we may assume that \mathcal{S} is finite. Let $g \in S^0(\mathbf{R}^d; T_S F)$. First, assume that $g = \sum_{Q \in \mathcal{S}} g_Q$ with $g_Q \in S^0(\mathbf{R}^d; \langle F \rangle_Q \mathbf{1}_Q)$. As in the proof of Theorem 4.4, by the John ellipsoid theorem, each g_Q is of the form

$$g_Q(x) = \sum_{k=1}^n h_k^Q(x) \langle f_k^Q \rangle_Q \mathbf{1}_Q(x),$$

where $h^Q = (h_1^Q, \dots, h_n^Q)$ satisfies $|h^Q(x)| \leq n^{\frac{1}{2}}$, and $f_k^Q \in S^0(\mathbf{R}^d; F \mathbf{1}_Q)$. Let $\psi \in \mathbf{X}'$, and set

$$\tilde{h}_k^Q = \frac{\langle f_k^Q \rangle_Q \cdot \langle h_k^Q \psi \rangle_Q}{|\langle f_k^Q \rangle_Q \cdot \langle h_k^Q \psi \rangle_Q|} h_k^Q$$

so that

$$\langle f_k^Q \rangle_Q \cdot \langle \tilde{h}_k^Q \psi \rangle_Q = |\langle f_k^Q \rangle_Q \cdot \langle h_k^Q \psi \rangle_Q|.$$

Then, noting that $\langle \tilde{h}_k^Q \psi \rangle_Q \in C_n \langle \psi \rangle_Q$,

$$\begin{aligned} \left| \int_{\mathbf{R}^d} g \cdot \psi \, dx \right| &\leq \sum_{k=1}^n \left| \sum_{Q \in \mathcal{S}} \int_Q h_k^Q \langle f_k^Q \rangle_Q \cdot \psi \, dx \right| \\ &= \sum_{k=1}^n \left| \sum_{Q \in \mathcal{S}} \langle f_k^Q \rangle_Q \cdot \langle h_k^Q \psi \rangle_Q |Q| \right| \\ &\leq 2 \sum_{k=1}^n \int_{\mathbf{R}^d} \sum_{Q \in \mathcal{S}} \langle f_k^Q \rangle_Q \mathbf{1}_{E_Q} \cdot \sum_{Q \in \mathcal{S}} \langle \tilde{h}_k^Q \psi \rangle_Q \mathbf{1}_{E_Q} \, dx \\ &\leq 2 \sum_{k=1}^n \left\| \sum_{Q \in \mathcal{S}} \langle f_k^Q \rangle_Q \mathbf{1}_{E_Q} \right\|_{\mathbf{X}} \left\| \sum_{Q \in \mathcal{S}} \langle \tilde{h}_k^Q \psi \rangle_Q \mathbf{1}_{E_Q} \right\|_{\mathbf{X}'} \\ &\lesssim_n \|M^{\mathcal{K}} F\|_{\mathbf{X}[\mathcal{K}]} \|M^{\mathcal{K}}(\mathcal{K}(\psi))\|_{\mathbf{X}[\mathcal{K}]} \\ &\leq \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]} \|F\|_{\mathbf{X}[\mathcal{K}]} \|\psi\|_{\mathbf{X}'}. \end{aligned}$$

For a general $g \in S^0(\mathbf{R}^d; T_S F)$, by Proposition 3.11, for each $Q \in \mathbf{S}$ there is a sequence $(g_{k,Q})_{k \geq 1}$ with $g_{k,Q} \in S^0(\mathbf{R}^d; \langle F \rangle_Q \mathbf{1}_Q)$ for which

$$\sum_{Q \in \mathcal{S}} g_{k,Q}(x) \rightarrow g(x)$$

for a.e. $x \in \mathbf{R}^d$. As \mathbf{X} is Köthe reflexive, it satisfies the Fatou property. Thus,

$$\|g\|_{\mathbf{X}} \leq \liminf_{k \rightarrow \infty} \left\| \sum_{Q \in \mathcal{S}} g_{k,Q} \right\|_{\mathbf{X}} \lesssim_n \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]} \|F\|_{\mathbf{X}[\mathcal{K}]}.$$

We conclude that $T_S F \in \mathbf{X}[\mathcal{K}]$, with

$$\|T_S F\|_{\mathbf{X}[\mathcal{K}]} \lesssim_n \|M^{\mathcal{K}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|M^{\mathcal{K}}\|_{\mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]} \|F\|_{\mathbf{X}[\mathcal{K}]}.$$

The assertion follows. \square

For the implication (i) \Rightarrow (ii) we have to prove convex body domination of $M^{\mathcal{K}}$:

Theorem 7.2. *Let \mathcal{D} be a dyadic grid and let $\mathcal{F} \subseteq \mathcal{D}$ be finite. For each $F \in L_{loc}^1(\mathbf{R}^d; \mathcal{K})$ there is a sparse collection $\mathcal{S} \subseteq \mathcal{F}$ for which*

$$M_{\mathcal{F}}^{\mathcal{K}} F(x) \subseteq 2n^{\frac{5}{2}} M_{\mathcal{S}}^{\mathcal{K}} F(x)$$

for a.e. $x \in \mathbf{R}^d$.

Proof. Let \mathcal{F}^* denote the maximal cubes in \mathcal{F} and fix $Q_0 \in \mathcal{F}^*$. By the John ellipsoid theorem there is an orthonormal basis $(e_k)_{k=1}^n$ of \mathbf{F}^n and $\lambda_1, \dots, \lambda_n \geq 0$ such that

$$\mathcal{E}(Q_0) := \left\{ \sum_{k=1}^n u_k \lambda_k e_k : |u| \leq 1 \right\}$$

satisfies

$$\mathcal{E}(Q_0) \subseteq \langle F \rangle_{Q_0} \subseteq n^{\frac{1}{2}} \mathcal{E}(Q_0).$$

For each $k \in \{1, \dots, n\}$ we let $\text{ch}_k(Q_0)$ denote the collection of maximal cubes $Q \in \mathcal{F}$ contained in Q_0 satisfying

$$|\langle F \cdot e_k \rangle_Q| > 2n \langle |F \cdot e_k| \rangle_{Q_0},$$

where

$$|\langle F \cdot e_k \rangle_Q| = \sup_{f \in S^0(\mathbf{R}^d; F)} |\langle f \cdot e_k \rangle_Q|, \quad \langle |F \cdot e_k| \rangle_{Q_0} = \langle x \mapsto \sup_{u \in F(x)} |u \cdot e_k| \rangle_{Q_0}.$$

Moreover, we let $\text{ch}(Q_0)$ denote the maximal cubes in $\bigcup_{k=1}^n \text{ch}_k(Q_0)$. Applying this same procedure to each of the cubes in $\text{ch}(Q_0)$, we iteratively obtain the collections $\mathcal{S}_{j+1}(Q_0) := \bigcup_{Q \in \mathcal{S}_j(Q_0)} \text{ch}(Q)$, where $\mathcal{S}_0(Q_0) := \{Q_0\}$. Setting $\mathcal{S} := \bigcup_{Q_0 \in \mathcal{F}^*} \bigcup_{j=0}^{\infty} \mathcal{S}_j(Q_0)$, we note that for each $Q \in \mathcal{S}$ we have

$$\begin{aligned} \sum_{Q' \in \text{ch}(Q)} |Q'| &\leq \sum_{k=1}^n \sum_{Q' \in \text{ch}_k(Q)} |Q'| \\ &\leq \sum_{k=1}^n \frac{1}{2n \langle |F \cdot e_k| \rangle_Q} \sum_{Q' \in \text{ch}_k(Q)} \int_{Q'} \sup_{u \in F(x)} |u \cdot e_k| dx \\ &\leq \sum_{k=1}^n \frac{1}{2n \langle |F \cdot e_k| \rangle_Q} \int_Q \sup_{u \in F(x)} |u \cdot e_k| dx = \frac{|Q|}{2}. \end{aligned}$$

Hence, \mathcal{S} is sparse.

For each $Q' \in \mathcal{F}$ we let $\pi_{\mathcal{S}}(Q')$ denote the smallest cube $Q \in \mathcal{S}$ containing it. If $Q' \in \mathcal{F}$ satisfies $\pi_{\mathcal{S}}(Q') = Q$ and $Q' \neq Q$, then Q' fails the stopping condition for all $k \in \{1, \dots, n\}$, i.e.,

$$|\langle F \cdot e_k \rangle_{Q'}| \leq 2n \langle |F \cdot e_k| \rangle_Q,$$

where $(e_k)_{k=1}^n$ is the orthonormal basis of associated to $\mathcal{E}(Q)$. Furthermore, this estimate remains true when $Q' = Q$. Thus, we have

$$\langle F \rangle_{Q'} = \sum_{k=1}^n \langle F \cdot e_k \rangle_{Q'} e_k \subseteq 2n \sum_{k=1}^n \langle |F \cdot e_k| \rangle_Q \mathcal{K}(e_k) \subseteq 2n^{\frac{5}{2}} \langle F \rangle_Q.$$

To see this last inclusion, pick an $f \in S^0(\mathbf{R}^d; F)$ for which $\langle |F \cdot e_k| \rangle_Q = \langle |f \cdot e_k| \rangle_Q$. Multiplying f by a unit, we may assume that $f \cdot e_k \geq 0$. Since $\langle f \rangle_Q \in \langle F \rangle_Q \subseteq n^{\frac{1}{2}} \mathcal{E}(Q)$, we can write

$$\langle f \rangle_Q = n^{\frac{1}{2}} \sum_{k=1}^n u_k \lambda_k e_k,$$

where $|u_k| \leq 1$. Hence,

$$\langle |F \cdot e_k| \rangle_Q e_k = (\langle f \rangle_Q \cdot e_k) e_k = n^{\frac{1}{2}} u_k \lambda_k e_k \in n^{\frac{1}{2}} \mathcal{E}(Q) \subseteq n^{\frac{1}{2}} \langle F \rangle_Q,$$

as desired.

In conclusion, we obtain

$$\bigcup_{Q \in \mathcal{F}} \langle F \rangle_Q \mathbf{1}_Q(x) = \bigcup_{Q \in \mathcal{S}} \bigcup_{\substack{Q' \in \mathcal{F} \\ \pi_{\mathcal{S}}(Q')=Q}} \langle F \rangle_{Q'} \mathbf{1}_{Q'}(x) \subseteq 2n^{\frac{5}{2}} \bigcup_{Q \in \mathcal{S}} \langle F \rangle_Q \mathbf{1}_Q(x) \subseteq 2n^{\frac{5}{2}} M_S^K F(x).$$

As $M_{\mathcal{S}}^{\mathcal{K}}F(x) \in \mathcal{K}$, we find that $M_{\mathcal{F}}^{\mathcal{K}}F(x) \subseteq 2n^{\frac{5}{2}}M_{\mathcal{S}}^{\mathcal{K}}F(x)$, as desired. \square

Proof of Theorem B. For the implication (ii) \Rightarrow (i), the bound $T_{\mathcal{S}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ follows from Theorem 7.1. Likewise, as \mathbf{X} is Köthe reflexive, the dual bound $T_{\mathcal{S}} : \mathbf{X}'[\mathcal{K}] \rightarrow \mathbf{X}'[\mathcal{K}]$ follows from Theorem 7.1 applied with \mathbf{X}' instead of \mathbf{X} .

It remains to prove (i) \Rightarrow (ii). By symmetry, we need only prove that $M^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$. By the Fatou property and the 3^d -lattice theorem, it suffices to prove that $M_{\mathcal{F}}^{\mathcal{K}} : \mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]$ for all finite collections \mathcal{F} contained in a dyadic grid \mathcal{D} . Let $F \in \mathbf{X}[\mathcal{K}]$. By Theorem 7.2, there is a sparse collection $\mathcal{S} \subseteq \mathcal{F}$ such that

$$M_{\mathcal{F}}^{\mathcal{K}}F(x) \subseteq C_n M_{\mathcal{S}}^{\mathcal{K}}F(x) \subseteq C_n T_{\mathcal{S}}(x),$$

where the last inclusion follows from the fact that $\bigcup_{Q \in \mathcal{S}} \langle F \rangle_Q \mathbf{1}_Q(x) \subseteq \sum_{Q \in \mathcal{S}} \langle F \rangle_Q \mathbf{1}_Q(x)$. Thus, by the directional ideal property of \mathbf{X} , we have

$$\|M_{\mathcal{F}}^{\mathcal{K}}F\|_{\mathbf{X}[\mathcal{K}]} \lesssim_n \|T_{\mathcal{S}}F\|_{\mathbf{X}[\mathcal{K}]} \leq \|T_{\mathcal{S}}\|_{\mathbf{X}[\mathcal{K}] \rightarrow \mathbf{X}[\mathcal{K}]} \|F\|_{\mathbf{X}[\mathcal{K}]},$$

proving the desired bound. \square

APPENDIX A. BASIC PROPERTIES OF \mathbf{F}^n -DIRECTIONAL QUASI-BANACH FUNCTION SPACES

Proposition A.1. *Let \mathbf{X} be a complete quasi-normed subspace of $L^0(\Omega; \mathbf{F}^n)$ satisfying the directional ideal property. Then the following are equivalent:*

- (i) \mathbf{X} satisfies the non-degeneracy property;
- (ii) $\|\cdot\|_{\mathbf{X}'}$ is a norm;
- (iii) $\mathbf{X} \cap L^2(\Omega; \mathbf{F}^n)$ is dense in $L^2(\Omega; \mathbf{F}^n)$;
- (iv) For all $g \in L^0(\Omega; \mathbf{F}^n)$ there is a sequence $(f_k)_{k \geq 1}$ in \mathbf{X} for which $f_k \rightarrow g$ a.e.

Proof. The equivalence (i) \Leftrightarrow (ii) follows from Proposition 3.2, as it implies that $\int_{\Omega} f \cdot g \, d\mu = 0$ for all $f \in \mathbf{X}$ if and only if $\int_{\Omega} |f \cdot g| \, d\mu = 0$ for all $f \in \mathbf{X}$.

To see (i) \Rightarrow (iii), we note that the statement is equivalent to showing that

$$(\mathbf{X} \cap L^2(\Omega; \mathbf{F}^n))^{\perp} = \{0\},$$

where

$$(\mathbf{X} \cap L^2(\Omega; \mathbf{F}^n))^{\perp} = \left\{ g \in L^2(\Omega; \mathbf{F}^n) : \int_{\Omega} f \cdot g \, d\mu = 0 \text{ for all } f \in \mathbf{X} \cap L^2(\Omega; \mathbf{F}^n) \right\}.$$

Let $f \in \mathbf{X}$, let $(E_k)_{k \geq 1}$ be a sequence of sets for which $\mu(E_k) < \infty$ for all $k \geq 1$ and $\bigcup_{k=1}^{\infty} E_k = \Omega$. Setting

$$f_k := \mathbf{1}_{\{x \in E_k : |f(x)| \leq k\}} f,$$

we have $f_k \in \mathbf{X} \cap L^2(\Omega; \mathbf{F}^n)$, and, by the monotone convergence theorem, for any $g \in L^2(\Omega; \mathbf{F}^n)$ we have

$$\left| \int_{\Omega} f \cdot g \, d\mu \right| \leq \int_{\Omega} |f \cdot g| \, d\mu = \sup_{k \geq 1} \int_{\Omega} |f_k \cdot g| \, d\mu = 0.$$

Hence, $g = 0$ by the non-degeneracy property of \mathbf{X} , as desired.

For (iii) \Rightarrow (iv), let Y denote the closure of $\mathbf{X} \cap L^2(\Omega; \mathbf{F}^n)$ in $L^0(\Omega; \mathbf{F}^n)$ with respect to a.e. pointwise convergence. Since $L^2(\Omega; \mathbf{F}^n)$ convergence implies a.e. pointwise convergence of a subsequence, it follows from (iii) that

$$L^2(\Omega; \mathbf{F}^n) \subseteq Y.$$

For any $g \in L^0(\Omega; \mathbf{F}^n)$, the sequence

$$f_k := \mathbf{1}_{\{x \in E_k : |g(x)| \leq k\}} g$$

satisfies $f_k \in L^2(\Omega; \mathbf{F}^n)$, and $f_k \rightarrow g$ a.e. We conclude that

$$Y = L^0(\Omega; \mathbf{F}^n),$$

as desired.

Finally, for (iv) \Rightarrow (ii), suppose that $\|g\|_{\mathbf{X}'} = 0$, and let $(f_k)_{k \geq 1}$ in \mathbf{X} be such that $f_k \rightarrow g$ a.e. By Fatou's lemma, we have

$$\int_{\Omega} |g|^2 d\mu \leq \liminf_{k \rightarrow \infty} \int_{\Omega} |f_k \cdot g| d\mu = 0.$$

Hence, $g = 0$, as asserted. \square

Next, we prove the \mathbf{F}^n -directional analogue of the Lorentz-Luxemburg theorem. Recall that Köthe reflexivity means that $\mathbf{X}'' = \mathbf{X}$.

Theorem A.2. *Let \mathbf{X} be an \mathbf{F}^n -directional Banach function space over Ω with the directional saturation property. Then the following are equivalent:*

- (i) \mathbf{X} satisfies the Fatou property;
- (ii) \mathbf{X} is Köthe reflexive.

To prove this, we use the following result:

Lemma A.3. *Let \mathbf{X} be an \mathbf{F}^n -directional Banach function space over Ω . If \mathbf{X} satisfies the Fatou property, then*

$$\|f\|_{\mathbf{X}} = \sup_{\|g\|_{\mathbf{X}'}=1} \int_{\Omega} |f \cdot g| d\mu$$

for all $f \in \mathbf{X}$.

Proof. We define the seminorm

$$\|f\|_{\mathbf{X}''} := \sup_{\|g\|_{\mathbf{X}'}=1} \int_{\Omega} |f \cdot g| d\mu,$$

and will show that $\|f\|_{\mathbf{X}''} = \|f\|_{\mathbf{X}}$ for all $f \in \mathbf{X}$ if \mathbf{X} has the Fatou property. Since $\mathbf{X} \subseteq \mathbf{X}''$ with $\|\cdot\|_{\mathbf{X}''} \leq \|\cdot\|_{\mathbf{X}}$, it remains to prove the converse inequality.

Let $B_{\mathbf{X}} := \{f \in \mathbf{X} : \|f\|_{\mathbf{X}} \leq 1\}$ and let

$$Y := B_{\mathbf{X}} \cap L^2(\Omega; \mathbf{F}^n).$$

Then Y is a convex set. We claim that Y is a closed subset of $L^2(\Omega; \mathbf{F}^n)$. Indeed, if $(f_k)_{k \geq 1}$ is a sequence in Y that converges to a function $f \in L^2(\Omega; \mathbf{F}^n)$ in $L^2(\Omega; \mathbf{F}^n)$, then for some subsequence f_{k_j} we have $f_{k_j} \rightarrow f$ a.e so that by the Fatou property of \mathbf{X} we have $f \in \mathbf{X}$ with

$$\|f\|_{\mathbf{X}} \leq \liminf_{j \rightarrow \infty} \|f_{k_j}\|_{\mathbf{X}} \leq 1.$$

Hence, $f \in Y$, as desired.

Now, let $f \in \mathbf{X} \cap L^2(\Omega; \mathbf{F}^n)$ with $\|f\|_{\mathbf{X}} = 1$, let $\varepsilon > 0$, and set $f_0 := (1 + \varepsilon)f$. Then $f_0 \in L^2(\Omega; \mathbf{F}^n)$, but, since $\|f_0\|_{\mathbf{X}} = 1 + \varepsilon > 1$, $f_0 \notin Y$. Since Y is convex, it follows from the Hahn-Banach separation theorem, the Riesz representation theorem, and Proposition 3.2, that there is a $g \in L^2(\Omega; \mathbf{F}^n)$ such that

$$\int_{\Omega} |f_0 \cdot g| d\mu > 1, \quad \int_{\Omega} |h \cdot g| d\mu < 1 \quad \text{for all } h \in Y.$$

We claim that $g \in \mathbf{X}'$ with $\|g\|_{\mathbf{X}'} \leq 1$. Indeed, let $h \in \mathbf{X}$ with $\|h\|_{\mathbf{X}} = 1$. Since Ω is σ -finite we can pick an increasing sequence of sets $(E_k)_{k \geq 1}$ with $\bigcup_{k=1}^{\infty} E_k = \Omega$ and $\mu(E_k) < \infty$ and set

$$h_k := h \mathbf{1}_{\{x \in E_k : |h(x)| \leq k\}} \in Y.$$

Then

$$\int_{\Omega} |h_k \cdot g| d\mu < 1.$$

Since $|h_k \cdot u| \uparrow |h \cdot u|$ a.e. for all $u \in \mathbf{F}^n$, it follows from the monotone convergence theorem that

$$\int_{\Omega} |h \cdot g| \, d\mu = \sup_{k \geq 1} \int_{\Omega} |h_k \cdot g| \, d\mu \leq 1.$$

Thus indeed, $g \in \mathbf{X}'$ with $\|g\|_{\mathbf{X}'} \leq 1$. Noting that

$$1 < (1 + \varepsilon) \int_{\Omega} |f \cdot g| \, d\mu \leq (1 + \varepsilon) \|f\|_{\mathbf{X}''} \|g\|_{\mathbf{X}'} \leq (1 + \varepsilon) \|f\|_{\mathbf{X}''},$$

letting $\varepsilon \downarrow 0$, we conclude that

$$\|f\|_{\mathbf{X}''} \geq 1 = \|f\|_{\mathbf{X}},$$

and, thus, $\|f\|_{\mathbf{X}''} = \|f\|_{\mathbf{X}}$ for all $f \in \mathbf{X} \cap L^2(\Omega; \mathbf{F}^n)$.

Finally, let $f \in \mathbf{X}$ and set

$$f_k := \mathbf{1}_{\{x \in E_k : |f(x)| \leq k\}} f$$

so that $|f_k \cdot u| \uparrow |f \cdot u|$ a.e. for all $u \in \mathbf{F}^n$. Then, by the monotone convergence property of \mathbf{X} and \mathbf{X}'' , $f \in \mathbf{X}$ with

$$\|f\|_{\mathbf{X}} = \sup_{k \geq 1} \|f_k\|_{\mathbf{X}} = \sup_{k \geq 1} \|f_k\|_{\mathbf{X}''} = \|f\|_{\mathbf{X}''}.$$

The assertion follows. \square

Proof of Theorem A.2. If $\mathbf{X} = \mathbf{X}''$, then \mathbf{X} has the Fatou property by Fatou's lemma. Conversely, by Lemma A.3, the Fatou property of \mathbf{X} implies that $\|f\|_{\mathbf{X}} = \|f\|_{\mathbf{X}''}$ for all $f \in \mathbf{X}$. Let $f \in \mathbf{X}''$. By the directional saturation property and Proposition 3.3(iii) there is a sequence $(f_k)_{k \geq 1}$ in \mathbf{X} for which $|f_k \cdot u| \uparrow |f \cdot u|$ a.e. for all $u \in \mathbf{F}^n$. By the monotone convergence property of \mathbf{X} and \mathbf{X}'' we conclude that $f \in \mathbf{X}$, and

$$\|f\|_{\mathbf{X}} = \sup_{k \geq 1} \|f_k\|_{\mathbf{X}} = \sup_{k \geq 1} \|f_k\|_{\mathbf{X}''} = \|f\|_{\mathbf{X}''}.$$

The assertion follows. \square

ACKNOWLEDGEMENTS

I wish to thank Rosemarie Bongers for being better at linear algebra than me, for sharing her knowledge on the Hausdorff distance, and for their insights into the geometry of convex sets. Moreover, I would like to thank her for their feedback on parts of the text, resulting in an overall improvement of the presentation of this work.

REFERENCES

- [AF09] J.-P. Aubin and H. Frankowska. *Set-valued analysis*. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, 2009. Reprint of the 1990 edition [MR1048347].
- [BC23] M. Bownik and D. Cruz-Uribe. Extrapolation and factorization of matrix weights. arXiv:2210.09443, 2023.
- [Buc93] S. Buckley. Estimates for operator norms on weighted spaces and reverse Jensen inequalities. *Trans. Amer. Math. Soc.*, 340(1):253–272, 1993.
- [CDO18] A. Culiuc, F. Di Plinio, and Y. Ou. Uniform sparse domination of singular integrals via dyadic shifts. *Math. Res. Lett.*, 25(1):21–42, 2018.
- [CG01] M. Christ and M. Goldberg. Vector A_2 weights and a Hardy-Littlewood maximal function. *Trans. Amer. Math. Soc.*, 353(5):1995–2002, 2001.
- [CIM18] D. Cruz-Uribe, J. Isralowitz, and K. Moen. Two weight bump conditions for matrix weights. *Integral Equations Operator Theory*, 90(3):Paper No. 36, 31, 2018.
- [CMM22] M. Cao, J.J. Marín, and J.M. Martell. Extrapolation on function and modular spaces, and applications. *Adv. Math.*, 406:Paper No. 108520, 87, 2022.
- [CMP11] D.V. Cruz-Uribe, J.M. Martell, and C. Pérez. *Weights, extrapolation and the theory of Rubio de Francia*, volume 215 of *Operator Theory: Advances and Applications*. Birkhäuser/Springer Basel AG, Basel, 2011.
- [Cru24] D. Cruz-Uribe. Matrix weights, singular integrals, Jones factorization and Rubio de Francia extrapolation. arXiv:2304.03887, 2024.

- [CV77] C. Castaing and M. Valadier. *Convex analysis and measurable multifunctions*, volume Vol. 580 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1977.
- [DGPP05] O. Dragičević, L. Grafakos, M.C. Pereyra, and S. Petermichl. Extrapolation and sharp norm estimates for classical operators on weighted Lebesgue spaces. *Publ. Mat.*, 49(1):73–91, 2005.
- [DPTV24] K. Domelevo, S. Petermichl, S. Treil, and A. Volberg. The matrix A_2 conjecture fails, i.e. $3/2 > 1$. arXiv:2402.06961, 2024.
- [Gra14] L. Grafakos. *Classical Fourier analysis*, volume 249 of *Graduate Texts in Mathematics*. Springer, New York, third edition, 2014.
- [KNV24] S. Kakaroumpas, T.H. Nguyen, and D. Vardakis. Matrix-weighted estimates beyond calderón-zygmund theory. arXiv:2404.02246, 2024.
- [Ler13] A.K. Lerner. On an estimate of Calderón-Zygmund operators by dyadic positive operators. *J. Anal. Math.*, 121:141–161, 2013.
- [LN24] E. Lorist and Z. Nieraeth. Banach function spaces done right. *Indag. Math. (N.S.)*, 35(2):247–268, 2024.
- [Nie23] Z. Nieraeth. Extrapolation in general quasi-Banach function spaces. *J. Funct. Anal.*, 285(10):Paper No. 110130, 2023.
- [Nie24] Z. Nieraeth. The Muckenhoupt condition. arXiv:2405.20907, 2024.
- [NPTV17] F. Nazarov, S. Petermichl, S. Treil, and A. Volberg. Convex body domination and weighted estimates with matrix weights. *Adv. Math.*, 318:279–306, 2017.
- [NT96] F. Nazarov and S. Treil. The hunt for a Bellman function: applications to estimates for singular integral operators and to other classical problems of harmonic analysis. *Algebra i Analiz*, 8(5):32–162, 1996.
- [Tre89] S. Treil. Geometric methods in spectral theory of vector-valued functions: some recent results. In *Toeplitz operators and spectral function theory*, volume 42 of *Oper. Theory Adv. Appl.*, pages 209–280. Birkhäuser, Basel, 1989.
- [TV97a] S. Treil and A. Volberg. Continuous frame decomposition and a vector Hunt-Muckenhoupt-Wheeden theorem. *Ark. Mat.*, 35(2):363–386, 1997.
- [TV97b] S. Treil and A. Volberg. Wavelets and the angle between past and future. *J. Funct. Anal.*, 143(2):269–308, 1997.
- [Vol97] A. Volberg. Matrix A_p weights via S -functions. *J. Amer. Math. Soc.*, 10(2):445–466, 1997.

ZOE NIERAETH (SHE/HER)

BCAM—BASQUE CENTER FOR APPLIED MATHEMATICS, BILBAO, SPAIN

Email address: zoe.nieraeth@gmail.com