A NOTE ABOUT DUAL REPRESENTATIONS OF GROUP ACTIONS ON LIPSCHITZ-FREE SPACES

MICHAEL MEGRELISHVILI

ABSTRACT. Let $\mathcal{F}(M)$ be the Lipschitz-free space of a pointed metric space M. For every isometric continuous group action of G we have an induced continuous dual action on the weak-star compact unit ball $B_{\mathcal{F}(M)^*}$ of the dual space $\operatorname{Lip}_0(M) =$ $\mathcal{F}(M)^*$. We pose the question when a given abstract continuous action of G on a topological space X can be represented through a G-subspace of $B_{\mathcal{F}(M)^*}$. One of such natural examples is the so-called *metric compactification* (of isometric Gspaces) for a pointed metric space. As well as the *Gromov G-compactification* of a bounded metric G-space. Note that there are sufficiently many representations of compact G-spaces on Lipschitz-free spaces.

CONTENTS

1.	Introduction	1
2.	Lipschitz-free spaces	2
3.	Representation of dynamical systems on Lipschitz-free spaces	3
References		7

1. INTRODUCTION

The Lipschitz-free space $\mathcal{F}(M)$ is a Banach space canonically defined for every pointed metric space M which captures several metric properties of M. The theory of Lipschitz-free spaces is a rapidly growing important direction. See, for example, [5, 1, 13] and references therein. Alternative terminology is the *free Banach space* of M as in a work by V. Pestov [11].

In the present note we propose to study dual representations of abstract continuous group actions on Banach spaces $\mathcal{F}(M)$.

First we give necessary definitions. To every Banach space $(V, || \cdot ||)$ one may associate several important structures. Among others: the topological group Is(V) of all linear onto isometries (in its strong operator topology), its canonical dual action on the weak-star compact unit ball B_{V^*} of the dual Banach space V^* . One of the natural ideas is to give a kind of linearization of abstract continuous actions $G \times X \to X$ of a topological group G on a topological space (we say, a G-space) through the dual action on some B_{V^*} .

Date: 2024, August.

²⁰²⁰ Mathematics Subject Classification. 46B04, 46B20, 43A65, 51F30, 53C23, 54H15.

Key words and phrases. Arens–Eells embedding, Lipschitz-free space, group representation, Gromov compactification.

Definition 1.1. [8, 3, 4] Let X be a G-space. A representation of (G, X) on a Banach space V is a pair

$$h: G \to \operatorname{Is}(V), \quad \alpha: X \to V^*$$

where $h: G \to Is(V)$ is a continuous homomorphism and $\alpha: X \to V^*$ is a weak^{*} continuous bounded (e.g., $\alpha(X) \subset B_{V^*}$) G-mapping with respect to the dual action

$$G \times V^* \to V^*, \ (g\varphi)(v) := \varphi(g^{-1}v).$$
$$G \times X \longrightarrow X$$
$$\downarrow^h \qquad \qquad \downarrow^\alpha \qquad \qquad \downarrow^\alpha$$
$$\operatorname{Is}(V) \times V^* \longrightarrow V^*$$

Proper representation will mean that α is a topological embedding. Note that when X is compact then every weak-star continuous $\alpha \colon X \to V^*$ is necessarily bounded.

This definition brings some new tools studying abstract dynamical G-systems using the geometry of Banach spaces. For some applications and more information we refer to [8, 4].

Our aim in the present note is to propose linearizations of actions (in the sense of Definition 1.1) but just on the Lipschitz-free space $V := \mathcal{F}(M)$. See Definition 3.2 as a special case of Definition 1.1. Theorem 3.8 provides a kind of such linearization for the so-called *metric compactification* (of isometric *G*-spaces) for a pointed metric space. As well as the *Gromov G-compactification* of a bounded metric *G*-space.

Note that (see Remark 3.4) there are sufficiently many representations of compact G-spaces on the Lipschitz-free spaces $\mathcal{F}(M)$. Below we give a necessary background and pose some general questions (3.3, 3.5, 3.6).

Acknowledgment. I would like to thank the organizers and participants of the *First Conference on Lipschitz-Free Spaces* (Besancon, September 2023) for their inspiration and stimulating discussions. A part of the present article's ideas was presented on this conference.

2. Lipschitz-free spaces

Here we recall shortly well known facts about Lipschitz-free spaces. Let M be a nonempty set. A *molecule* of M is a formal finite sum

$$m = \sum_{i=1}^{n} c_i (x_i - y_i),$$

where $x_i, y_i \in M, c_i \in \mathbb{R}, n \in \mathbb{N}$. It can be identified with a function $m: M \to \mathbb{R}$ having a finite support such that $\sum_{x \in M} m(x) = 0$. The set Mol(M) of all molecules is a vector space over \mathbb{R} . Now, let d be a pseudometric on M. Define

$$||m||_d := \inf\{\sum_{i=1}^n |c_i| d(x_i, y_i): m = \sum_{i=1}^n c_i(x_i - y_i)\}$$

This is a seminorm on Mol(M). It is well known (and not hard to show) that $|| \cdot ||_d$ is a norm if and only if d is a metric. In this case $(Mol(M), || \cdot ||_d)$ is said to be the *Arens-Eells normed space* of (M, d). Mostly we write simply Mol(M) and $|| \cdot ||$.

Remark 2.1. Note that this norm sometimes is called Kantorovich-Rubinstein norm and it plays a major role in a classical branch of the optimization theory. Namely, in transportation problems.

Denote by $\mathcal{F}(M)$ the completion of $(Mol(M), || \cdot ||)$. This Banach space is said to be the *Lipschitz-free space* of (M, d).

Let $(M, d, \mathbf{0})$ be a *pointed* metric space with a distinguished point $\mathbf{0} \in M$. For every $x \in M$ define the molecule $\chi_x := x - \mathbf{0}$. The set $\{\chi_x : x \in X\}$ is a Hamel base of the vector space Mol(M, d). Define the following natural injection

$$\chi \colon M \to \operatorname{Mol}(M, d), \ x \mapsto \chi_x.$$


Clearly, χ_0 is the zero element **0** of $\mathcal{F}(M)$.

Now recall the description of the dual Banach space $\mathcal{F}(M)^*$ of $\mathcal{F}(M)$ (equivalently, the dual of the normed space $\operatorname{Mol}(M, d)$). For every functional $F: \mathcal{F}(M) \to \mathbb{R}$ we have an induced function $f: M \to \mathbb{R}$, $f(x) := F(\chi_x - \mathbf{0}) = F(\chi_x)$. Conversely, for every real function $f: M \to \mathbb{R}$ with $f(\mathbf{0}) = \mathbf{0}$ define $F: \operatorname{Mol}(M, d) \to \mathbb{R}$ extending flinearly. Formally, $F(m) = \sum_{i=1}^n c_i(f(x_i) - f(y_i))$ for every $m \in \sum_{i=1}^n c_i(x_i - y_i) \in$ $\operatorname{Mol}(M, d)$. Note that for pointed metric spaces every molecule can be represented as $m = \sum_{i=1}^n r_i x_i = \sum_{i=1}^n r_i(\chi_{x_i} - \mathbf{0})$. These observations are useful for example in the verification of Fact 2.2.

By $\operatorname{Lip}_0(X)$ we denote the space of all Lipschitz functions $f: X \to \mathbb{R}$ satisfying $f(\mathbf{0}) = 0$. It is well known that $\operatorname{Lip}_0(X)$ is a Banach space with respect to the natural norm $||f|| := \operatorname{Lip}(f)$. Recall some well known important properties.

Fact 2.2. Let $(M, d, \mathbf{0})$ be a pointed metric space. Then

- (1) $\mathcal{F}(M)^* = \operatorname{Lip}_0(M).$
- (2) Weak-star topology on bounded subsets $\mathcal{F}(M)^* = \operatorname{Lip}_0(X)$ coincides with the topology of pointwise convergence.
- (3) $||m||_d$ is the largest seminorm on $\operatorname{Mol}(M, d)$ such that $||\chi_x \chi_y|| \leq d(x, y)$. Moreover, $||\chi_x - \chi_y||_d = d(x, y)$. That is, $\chi \colon M \to \operatorname{Mol}(M)$ is an isometric embedding.
- (4) (Universal property) Let V be a Banach space and $f \in \operatorname{Lip}_0(M, V)$ there exists a unique bounded linear map $T_f \in L(\mathfrak{F}(M), V)$ such that $f = T_f \circ \chi$ and $||T_f|| = ||f||_{\operatorname{Lip}}$.
- (5) (Canonical linearization) Let $f: (M_1, \mathbf{0}) \to (M_2, \mathbf{0})$ be a Lipschitz map between two pointed metric spaces extends to a unique continuous linear map $\bar{f}: \mathfrak{F}(M_1) \to \mathfrak{F}(M_2)$ such that $\bar{f} \circ \chi_1 = \chi_2 \circ f$.



3. Representation of dynamical systems on Lipschitz-free spaces

As before we denote by $(M, d, \mathbf{0})$ a pointed metric space. Suppose that we have an isometric continuous left action $\pi: G \times M \to M$ of a topological group G on M such

that $g\mathbf{0} = \mathbf{0}$ for every $g \in G$. Naturally extending this action to the normed space Mol(M) of all molecules we get a continuous isometric linear action

$$G \times \operatorname{Mol}(M) \to \operatorname{Mol}(M).$$

Passing to the completion we obtain an unique continuous linear (isometric) extension

$$G \times \mathcal{F}(M) \to \mathcal{F}(M).$$

Let $h: G \to \operatorname{Is}(\mathcal{F}(M))$ be the corresponding canonically defined continuous group homomorphism, where $\operatorname{Is}(\mathcal{F}(M))$ is the topological group of all **linear** isometries (endowed with the strong operator topology). Note that if G is a subgroup of the topological group $\operatorname{Is}(M, d)$ (with the pointwise topology) then h is a topological group embedding. It follows that every pointed metric space with an isometric action (which fixes the distinguished point) admits a natural linearization on $\mathcal{F}(M)$. In Definition 3.2 we deal with a different kind of linearization when the main target is the actions on the dual ball $B_{\mathcal{F}(M)^*}$.

Lemma 3.1. The induced dual action $\pi^* \colon G \times B_{\mathcal{F}(M)^*} \to B_{\mathcal{F}(M)^*}$ is continuous, where $B_{\mathcal{F}(M)^*}$ is the weak-star compact unit ball in the dual space $\mathcal{F}(M)^*$.

Proof. This is a particular case of a general well know fact (see, for example, [8, Fact 2.2]) which is true for all isometric linear actions of G on Banach spaces V. More precisely, the dual action $G \times B_{V^*} \to B_{V^*}$ is continuous for every topological subgroup $G \subseteq \text{Is}(V)$ and every normed space V, where B_{V^*} is the weak-star compact unit ball.

Definition 3.2. Let X be a topological G-space and M be a pointed metric space. A representation of (G, X) on $\mathcal{F}(M)$ is a pair (h, α) where $h: G \to \mathrm{Is}(M)$ is a continuous homomorphism and $\alpha: X \to B_{\mathcal{F}(M)^*}$ is a weak-star continuous G-equivariant map.

Question 3.3. Let \mathcal{M} be a some class of pointed metric spaces. Which dynamical systems (G, X) can be properly represented on $\mathcal{F}(M)$ for some $M \in \mathcal{M}$? Especially, whenever M somehow is related to the original topological space X.

Recall that (in view of Definition 1.1) proper representation simply means that α is a topological embedding. Every properly representable G-space is G-compactifiable.

Remark 3.4. Note that there are sufficiently many representations of compact G-spaces on the Lipschitz-free spaces $\mathcal{F}(M)$. Indeed, one may show that if K is a compact G-space then (G, K) admits a proper representation on $\mathcal{F}(M)$, where $M := B_{C(K)}$ is the norm closed unit ball as the desired metric space (zero of the Banach space C(K) is the distinguished point).

Question 3.5. For every $f \in B_{\mathcal{F}(M)^*}$ (individual 1-Lipschitz function on M) we have a compact G-space $K_f := cl_{w^*}(Gf) \subset B_{\mathcal{F}(M)^*}$. It would be interesting to study dynamical properties of such dynamical systems (G, K_f) .

Question 3.6. For every $f \in B_{\mathcal{F}(M)^*}$ and every $v \in M$ one may consider the corresponding **matrix coefficient** (which is bounded right uniformly continuous)

$$m_{f,v}: G \to \mathbb{R}, \ g \mapsto f(gv)$$

It would be interesting to study whether such functions on G belong to dynamically interesting class of functions. For instance, when $m_{f,v} \in WAP(G)$? When $m_{f,v} \in Asp(G)$? When $m_{f,v} \in Tame(G)$?

For the definitions of the algebras WAP(G), Asp(G), Tame(G) and their roles in Banach representations theory, see for example, [4].

Example 3.7. If d is a stable metric on G (that is, if it satisfies the double limit property of Grothendieck) then (making use the Grothendieck's criterion of WAP) one may show that $m_{\rho_a,v} \in WAP(G)$ for the matrix coefficient defined by any pair $v \in Mol(M)$ and $\rho_a \in B_{\mathcal{F}(M)^*}$, where $\rho_a(x) := d(a, x) - d(a, \mathbf{0})$ (as defined in Theorem 3.8 below).

In the future we are going to study these (and similar) questions. In the present note we discuss the so called metric (hori)compactifications interprating it as a particular case of a representation in the sense of Definition 3.2 and giving a partial response to general Question 3.3.

3.1. Metric equivariant compactifications. Recall that if (X, d) is a bounded metric space (not necessarily pointed), with a continuous *G*-action, then one may consider the following family of elementary (bounded) distance functions

(3.1)
$$\Gamma := \{g_a \colon X \to \mathbb{R}, \ \gamma_a(x) := d(a, x)\}_{a \in X}.$$

Let A_{γ} be a *G*-invariant closed unital subalgebra of $C_b(X)$ (the algebra of all continuous bounded functions with the sup-norm) generated by this family. Denote by $\gamma: X \to \hat{X}^{\gamma}$ the corresponding compactification (maximal ideal space). Then γ is a topological embedding because Γ separates points and closed subsets). In fact, γ is a *G*-compactification as it was mentioned in [9]. That is, there exists a continuous action $G \times \hat{X}^{\gamma} \to \hat{X}^{\gamma}$ which extends the given action on *X*. For some examples and applications (among others, for the Urysohn's sphere) we refer to [7] and [12]. In particular, note that the Gromov compactification of the Urysohn sphere (S_U, d) can be identified with the greatest *G*-compactification of S_U with respect to the action of its Polish isometry group $G := \text{Is}(S_U)$.

Below we deal with a closely related general construction of the so called *metric* compactifications $m: M \to \widehat{M}$. This concept was introduced by M. Gromov but there are several (distinct versions in the literature). Note that m is always continuous injective but not necessarily an embedding. Relevant information about metric (hori)compactifications can be found, for example, in [10, 6, 2].

Let us say that a point x_0 in (M, d) is equidistant if $d(x, x_0) = c_0 > 0$ is constant for every $x \in M \setminus \{x_0\}$. Clearly, then d is bounded, diam $(M, d) \leq 2c_0$. Conversely, for bounded metrics, one may adjoin a new point **0** which is equidistant. This also is very useful in view of actions because in this way any isometric G-action on a bounded metric G-space M can be naturally embedded into an isometric G-action on $M \cup \{0\}$ fixing the new isolated point. In this case assertion (3) of Theorem 3.8 in fact speaks about the "original" non-pointed metric space M and its Gromov compactification.

Below $h: G \to \text{Is}(\mathcal{F}(M))$, as before, be the canonically defined continuous homomorphism induced by the action of G on $\mathcal{F}(M)$. **Theorem 3.8.** Let $(M, d, \mathbf{0})$ be an isometric G-space. Define

$$\rho \colon M \to (\mathcal{F}(M)^*, w^*), \quad \rho(a) = \rho_a,$$

where

$$\rho_a(x) := d(a, x) - d(a, \mathbf{0}).$$

- (1) Then the pair (h, ρ) is a well defined continuous injective representation of the *G*-space *M* on $\mathcal{F}(M)$, with $\rho(\mathbf{0}) = 0_{\mathcal{F}}$ and $||\rho(a)||_{\text{Lip}} = 1$ for every $a \in M \setminus \{\mathbf{0}\}$.
- (2) The induced continuous (injective) G-map

$$\rho \colon M \to \widehat{M} := cl_{w^*}\rho(M) \subset B_{\mathcal{F}(M)^*}$$

is equivalent to the metric (hori)compactification of $(M, d, \mathbf{0})$.

(3) If **0** is equidistant (or, equivalently, if there exists an equidistant point in M) then ρ is a topological embedding and the induced compactification

 $\rho \colon M \setminus \{\mathbf{0}\} \to cl_{w^*}\rho(M \setminus \{\mathbf{0}\})$

is equivalent to the **Gromov compactification** of $(M \setminus \{0\}, d)$.

Proof. (1) First shortly observe that ρ is well defined. We repeatedly use the equality $\mathcal{F}(M)^* = \operatorname{Lip}_0(M)$ (Fact 2.2.1).

Indeed, $\rho_0(x) = 0$ for every $x \in M$. Hence, $\rho_0 = \rho(0) = 0_{\mathcal{F}}$. Also,

$$\rho_a(x) - \rho_a(y)| = |d(a, x) - d(a, \mathbf{0}) - (d(a, \mathbf{0}) - d(a, y))| \le d(x, y).$$

Hence, $||\rho_a||_{\text{Lip}} \leq 1$ for every $a \in M$. Furthermore,

$$|\rho_a(x) - \rho_a(a)| = |d(a, x) - d(a, 0) - d(a, 0) + d(a, a)| = d(a, x).$$

Therefore, $||\rho_a||_{\text{Lip}} = ||\rho(a)||_{\text{Lip}} = 1$ for every $a \in M \setminus \{\mathbf{0}\}$.

 ρ is injective. Indeed, let $a, b \in M$ and $\rho_a(x) = \rho_b(x)$ for every $x \in M$. Then, in particular, $\rho_a(a) = \rho_b(a)$ and $\rho_a(b) = \rho_b(b)$. So, $d(a, \mathbf{0}) - 0 = d(b, \mathbf{0}) - d(a, b)$ and $d(a, \mathbf{0}) - d(b, a) = d(b, \mathbf{0}) - 0$. Then we get 2d(a, b) = 0. Hence, a = b.

For every $x \in M$ define the following function

$$\varphi_x \colon M \to \mathbb{R}, \ \varphi_x(a) := d(a, x) - d(a, \mathbf{0}) = \rho_a(x)$$

Then φ_x is bounded because $|\varphi_x(a)| \leq d(\mathbf{0}, x)$ and continuous by

$$|\varphi_x(a_1) - \varphi_c(a_2)| \le 2d(a_1, a_2)$$

Every function ρ_a can be identified with the following element of \mathbb{R}^M as follows:

$$(\rho_a(x))_{x \in X} = (d(a, x) - d(a, \mathbf{0}))_{x \in M} = (\varphi_x(a))_{x \in X} \in \mathbb{R}^M.$$

Since $\varphi_x(a) := d(a, x) - d(a, \mathbf{0}) = \rho_a(x)$, the function $\rho \colon M \to \mathbb{R}^M$, $a \mapsto \rho_a$ is the diagonal product of the following family of functions $\Phi := \{\varphi_x \colon x \in M\}$. Denote by τ_w the corresponding pointwise (weak) topology on $\rho(M)$ which coincides with the topology of the corresponding precompact uniformity μ_{Φ} on M.

As we already established, $\rho(M) \subset B_{\mathcal{F}(M)^*} \subset \operatorname{Lip}_0(M)$ holds. Since $\rho(M)$ is norm bounded in $\operatorname{Lip}_0(M)$, the weak-star topology inherits on $\rho(M)$ the pointwise topology (Fact 2.2.2). That is exactly the subspace topology τ_w of the product \mathbb{R}^M . Every $\varphi_x \colon M \to \mathbb{R}$ is continuous. Hence, $\tau_w \subseteq top(d)$. This implies that the injection $\rho \colon M \to B_{\mathcal{F}(M)^*}$ is continuous.

$$\rho(ga)(x) = \rho_{ga}(x) = d(ga, x) - d(ga, \mathbf{0}) = d(a, g^{-1}x) - d(a, \mathbf{0}) = \rho_a(g^{-1}x) = (g\rho_a)(x)$$
for every $x \in M$. For a description of the dual action see Definition 1.1.

(2) It is well known that the metric compactification $m: M \to \widehat{M}$ is the completion of the precompact uniformity μ_{Φ} on X generated by the family (of bounded) functions $\Phi := \{\varphi_x : x \in M\}$. See, for example, [2, Section 2.4].

(3) We have to show that ρ is a **topological embedding** of (M, top(d)) into the weak-star compact space $(B_{\mathcal{F}(M)^*}, w^*)$.

Since **0** is equidistant, there exists $c_0 > 0$ such that $d(x, \mathbf{0}) = c_0$ for every $x \in M \setminus \{\mathbf{0}\}$. Therefore, $\varphi_x(a) = d(a, x) - c_0$. We claim that the following family of functions

(3.2)
$$\Gamma_0 := \{\varphi_x \colon X \to \mathbb{R}, \ \varphi_x(a) = d(a, x) - c_0\}_{x \in M}$$

separates points and closed subsets in M. Indeed, for every closed subset $B \subset M$ and a point $x_0 \notin B$ we have $\varphi_{x_0}(x_0) = -c_0$ and $\varphi_{x_0}(b) \ge c_1 - c_0$ for every $b \in B$, where $c_1 := d(x_0, B) > 0$. Hence, $\varphi_{x_0}(x_0) \neq cl(\varphi_{x_0}(B))$.

Clearly, the family $\Phi_X := \{\varphi_x | X : x \in X\}$, with $X := M \setminus \{\mathbf{0}\}$, generates the same unital subalgebra of $C_b(X)$ as the family

$$\Gamma := \{\gamma_a \colon X \to \mathbb{R}, \ \gamma_a(x) := d(a, x)\}_{a \in X}$$

from Equation 3.1.

Remark 3.9.

- (1) One of the corollaries of Theorem 3.8 is that for every isometric pointed metric G-space M the canonically defined representation (h, ρ) on its own Lipschitz-free space $\mathfrak{F}(M)$ is an injective representation of (G, M) which in fact induces the classical metric compactification.
- (2) The metric compactification m: M → M is always continuous and injective for every pointed metric space M but in general it is not a topological embedding. As an important sufficient condition, note that an important fact in metric geometry which asserts that m is a topological embedding for every complete geodesic and proper (the latter means that all closed balls are compact) metric space (M, d).
- (3) The metric compactification is independent (up to the homeomorphism) of the choice of base point. This justifies the first line in assertion (3) of Theorem 3.8.
- (4) The topology of m(M) inherited from M is the weak topology (in terms of [10]) generated by the family Φ (see also [2, Remark 2.6]).
- (5) The continuity of the induced G-action on \hat{X} was verified in [2, Lemma 2.5]. This fact follows directly from the continuity of the dual actions Lemma 3.1.

References

 M. Cuth, M. Doucha and P. Wojtaszczyk, On the structure of Lipschitz-free spaces, Proc. Amer. Math. Soc. 144 (2016), 3833–3846

- B. Duchesne, The Polish topology of the isometry group of the infinite dimensional hyperbolic space, Groups Geom. Dyn. 17 (2023), 633–670
- [3] E. Glasner and M. Megrelishvili, Linear representations of hereditarily non-sensitive dynamical systems, Colloquium Math., 104 (2006), no. 2, 223–283
- [4] E. Glasner and M. Megrelishvili, Representations of dynamical systems on Banach spaces. In: Recent Progress in General Topology III, Springer, Atlantis Press, 2014
- [5] G. Godefroy and N. J. Kalton, *Lipschitz-free Banach spaces*, Studia Math. 159 (2003), no. 1, 121–141. Dedicated to Professor Aleksander Pelczynski on the occasion of his 70th birthday
- [6] A.W. Gutierrez, On the Metric Compactification of Infinite-dimensional l_p Spaces, Canadian Math. Bull. 62 (2019), 491–507
- [7] T. Ibarlucia and M. Megrelishvili, Maximal equivariant compactification of the Urysohn spaces and other metric structures, Advances in Math. v. 380, 2021, 107599. ArXiv:2001.07228
- [8] M. Megrelishvili, Fragmentability and representations of flows, Topology Proc., 27:2 (2003), 497–544. For the updated version see arXiv:math/0411112
- M. Megrelishvili, Topological transformation groups: selected topics. Survey paper in: Open Problems in Topology II (Elliott Pearl, editor), Elsevier Science, 2007, pp. 423–438.
- [10] N. Monod, Superrigidity for irreducible lattices and geometric splitting, J. Amer. Math. Soc., 19(4), 781–814, 2006
- [11] V. Pestov, Free Banach spaces and representations of topological groups, Funct. Anal. Appl. 20 (1986), 70–72
- [12] V. Pestov, A topological transformation group without non-trivial equivariant compactifications, Advances in Math. **311** (2017) 1–17.
- [13] N. Weaver, Lipschitz Algebras, World Scientific Publishing Co., Inc., River Edge, NJ, 1999

(Michael Megrelishvili) Department of Mathematics Bar-Ilan University, 52900 Ramat-Gan Israel

Email address: megereli@math.biu.ac.il