1+1d SPT phases with fusion category symmetry: interface modes and non-abelian Thouless pump

Kansei Inamura^{1,*} and Shuhei Ohyama^{2,[†](#page-0-1)}

¹Institute for Solid State Physics, University of Tokyo, Kashiwa, Chiba 277-8581, Japan ²RIKEN Center for Emergent Matter Science, Wako, Saitama, 351-0198, Japan

(Dated: August 29, 2024)

We consider symmetry protected topological (SPT) phases with finite non-invertible symmetry $\mathcal C$ in 1+1d. In particular, we investigate interfaces and parameterized families of them within the framework of matrix product states. After revealing how to extract the C -SPT invariant, we identify the algebraic structure of symmetry operators acting on the interface of two $\mathcal{C}\text{-}SPT$ phases. By studying the representation theory of this algebra, we show that there must be a degenerate interface mode between different C -SPT phases. This result generalizes the bulk-boundary correspondence for ordinary SPT phases. We then propose the classification of one-parameter families of C-SPT states based on the explicit construction of invariants of such families. Our invariant is identified with a non-abelian generalization of the Thouless charge pump, which is the pump of a local excitation within a C -SPT phase. Finally, by generalizing the results for one-parameter families of SPT phases, we conjecture the classification of general parameterized families of general gapped phases with finite non-invertible symmetries in both 1+1d and higher dimensions.

CONTENTS

[∗] k [inamura@issp.u-tokyo.ac.jp](mailto:k_inamura@issp.u-tokyo.ac.jp)

[†] shuhei.ohyama@riken.jp

2

I. INTRODUCTION AND SUMMARY

Introduction. The theoretical study of topological phases has provided various insights into the physics of many-body systems. One of the most fundamental subjects of this research is a class of quantum states known as invertible states (a.k.a. short-range entangled states). An invertible state is a state that is realized as the unique ground state of a gapped Hamiltonian.^{[1](#page-2-1)} By imposing symmetry, invertible states exhibit various quantum phases, which have been actively studied as Symmetry Protected Topological (SPT) Phases [\[1–](#page-84-1)[8\]](#page-84-2). An interesting physical property of SPT phases is the existence of gapless edge modes. The non-triviality of the SPT phases and these edge modes are related to each other, and their relationship is referred to as bulk-boundary correspondence or anomaly inflow. Furthermore, it is known that topological transport phenomena can occur by adiabatically and periodically driving an invertible state, and this is referred to as generalized Thouless pump phenomena [\[9](#page-84-3)[–19\]](#page-85-0).

Another trend in the study of quantum many-body systems is the generalization of the concept of symmetry [\[20\]](#page-85-1). A typical example of this is the notion known as non-invertible symmetry (a.k.a. categorical symmetry), see, e.g., [\[21–](#page-85-2)[29\]](#page-85-3) for recent reviews. This framework regards invariance under operations on the theory, such as gauging and duality operations, as a symmetry of the theory. While conventional symmetries are described by the mathematical structure of groups, non-invertible symmetries are described by certain types of categories. In particular, the structure of finite non-invertible symmetries in $1+1$ dimensions is described by fusion categories $[30-32]$ $[30-32]$. In this context, it is natural to consider the classification of SPT phases protected by non-invertible symmetry.

Tensor networks are an efficient way to describe highly entangled states. They have been applied not only to numerical computations but also to the classification and description of topological phases [\[2,](#page-84-4) [4–](#page-84-5)[6,](#page-84-6) [8,](#page-84-2) [33–](#page-85-6)[35\]](#page-85-7). Furthermore, it has been pointed out that tensor networks are a useful platform for representing (or microscopically realizing) non-invertible symmetries [\[36–](#page-85-8)[40\]](#page-85-9). Therefore, tensor networks are an excellent tool for studying topological phases that possess non-invertible symmetries.

As pioneering work in the classification of 1+1d SPT phases protected by non-invertible symmetry, a method using effective theories described by TQFT has been proposed [\[32\]](#page-85-5). This paper argues that the classification of fiber functors of the fusion category corresponds to the classification of SPT phases.^{[2](#page-2-2)} However, this paper focuses on continuous systems using field theory and does not discuss lattice systems. Recent studies have also presented several examples of lattice models

¹ Here, we would like to clarify the usage of the term "invertible state." Generally, SPT phases protected by noninvertible symmetry do not possess a group structure. This means that for a given non-invertible symmetric state, there may not necessarily be a non-invertible symmetric state that acts as its inverse. However, we will take the position that the concept of invertible states is independent of symmetry. Therefore, we will refer to a state that is invertible when disregarding symmetry as an invertible state.

 2 See Sec. IIB for a brief review of this claim.

that exhibit SPT phases with non-invertible symmetry [\[41–](#page-85-10)[45\]](#page-86-0). These studies construct generalizations of the cluster model and investigate their phase structures. Additionally, a general theory for classifying gapped systems with symmetry, described by tensor networks known as matrix product operators (MPO), has also been proposed in [\[37\]](#page-85-11). This paper analyzes how matrix product operators (MPO) act on matrix product states (MPS) and points out that phases can be distinguished by the categorical data that appear when MPOs act on an MPS.

For SPT phases protected by non-invertible symmetry, the presence of edge modes and topological pumping phenomena are expected, similar to conventional SPT phases. However, systematic studies of these phenomena have not been carried out in previous research. Furthermore, since gapped lattice models are expected to be described by TQFT in the low-energy limit, it is expected that the data of a fiber functor can be extracted from the ground states of lattice systems. However, it has not been clearly established in previous research how to directly extract the data of the fiber functor from lattice systems.

In this study, we consider SPT phases protected by non-invertible symmetries within the framework of MPS. We first propose a method to extract the fiber functor from invertible states realized in lattice systems. We then investigate the symmetry algebra acting on the interface of SPT phases, generalizing the anomaly inflow to non-invertible symmetries. We also explore the classification of generalized Thouless pumps and their relation to interface modes. A more detailed summary of the paper is provided below.

Outlook and Summary. The structure of the paper is as follows. If the reader is familiar with the contents of Sec. [II,](#page-4-0) each section can be read almost independently. In this paper, we focus only on bosonic systems.

In Sec. [II,](#page-4-0) we provide a brief review of fusion categories and fiber functors. It is believed that SPT phases with fusion category symmetry $\mathcal C$ are classified by fiber functors of $\mathcal C$. We also review the interpretation of this classification from the perspective of Topological Quantum Field Theory (TQFT). In addition, we explain some mathematical facts on fusion categories used in the main part of the paper and provide a brief review of the relation between tensor networks and fusion categories.

In Sec. [III,](#page-17-0) we describe how to extract data of the fiber functor from a C-symmetric MPS. After some preliminary discussions on MPSs and their transfer matrices, we demonstrate that the data of the fiber functor can be extracted as the triple inner product of infinite MPSs. A key to this is the analogy between the partition functions of TQFT with symmetry defect and MPS. Additionally, we introduce the abelianization of the fiber functor. As a result, we can define an invariant that takes values in (abelian) cohomology. As an example, we explicitly compute the data of a fiber functor from the $G \times \text{Rep}(G)$ -symmetric cluster state [\[41,](#page-85-10) [46\]](#page-86-1).

In Sec. [IV,](#page-27-0) we analyze the interface modes between C -symmetric SPT phases. This analysis is a generalization of that of edge modes in conventional SPT phases with group symmetry. We first derive the symmetry operators that act on the interface. These operators form an algebra, which we refer to as the interface algebra. The analysis of interface modes is reduced to classifying the irreducible representations of this algebra. We show that the interface algebra between different SPT phases does not have one-dimensional representations. Consequently, an interface between different SPT phases gives rise to degenerate ground states. This can be regarded as the bulk-boundary correspondence in C-symmetric SPT phases. On the other hand, the self-interface within the same SPT phase always has one-dimensional representations, which are in one-to-one correspondence with the automorphisms of a fiber functor. As an example, we study interfaces of $\text{Rep}(D_8)$ -symmetric

SPT phases.

In Sec. [V,](#page-33-0) we investigate S^1 -parameterized families of C-symmetric invertible states. Based on a careful analysis of gauge redundancy of C-symmetric MPS tensors, we define an invariant that detects the non-triviality of an S^1 -family. We will see that this invariant takes values in the group of automorphisms of a fiber functor. Additionally, we show that the non-triviality of an S^1 -family gives rise to the Thouless pump of non-abelian charges.^{[3](#page-4-2)} This non-abelian Thouless pump is in contrast to conventional Thouless pumps, which provide pumping of abelian charges. As an example, we construct non-trivial S^1 -families of $\text{Rep}(G)$ -symmetric invertible states.

In Sec. [VI,](#page-42-0) we consider parameterized families of general \mathcal{C} -symmetric gapped systems, not limited to invertible states. The central hypothesis is that the moduli space of \mathcal{C} -symmetric gapped systems in a gapped phase labeled by a $\mathcal{C}\text{-module category }\mathcal{M}$ is given by the classifying space $B\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\mathrm{inv}}$ of categorical group $\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\mathrm{inv}}$. Here, $\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\mathrm{inv}}$ is a 2-group consisting of invertible C -module functors and invertible C -module natural transformations. Based on this hypothesis, it is expected that X-parameterized families of a general gapped phase $\mathcal M$ are classified by the non-abelian Čech cohomology $\check{H}^1(X,\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\mathrm{inv}})$. We will present examples supporting this conjecture by considering some special cases. Moreover, we extend this consideration to higher dimensions. Specifically, in the 2+1-dimensional case, it is expected that the moduli space of C-symmetric gapped systems in a non-chiral gapped phase labeled by a C-module 2-category $\mathcal M$ is given by the classifying space $B_{\text{min}}^{\text{Fun}}(M, M)$ ^{inv} of the categorical 2-group $\underline{\underline{\mathrm{Fun}}}_{\mathcal{C}}(M, M)$ ^{inv}. As in the 1+1-dimensional case, we provide several examples supporting this conjecture.

Various technical details are relegated to Appendices. In App. [A,](#page-57-1) we review the basics of the Tambara-Yamagami categories. In App. [B,](#page-58-0) we review the classification of fiber functors of the Tambara-Yamagami categories. In App. [C,](#page-59-0) we compute the data (called L-symbols) of fiber functors of non-anomalous $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami categories, i.e., Rep(D_8), Rep(Q_8), and Rep(H_8). In App. [D,](#page-61-2) we enumerate irreducible representations of the interface algebras for SPT phases with $Rep(D_8)$, $Rep(Q_8)$, and $Rep(H_8)$ symmetries. In App. [E,](#page-71-0) we show that the $G \times Rep(G)$ -symmetric cluster state is obtained as the ground state of a $Rep(G)$ -symmetric model discussed in [\[47\]](#page-86-2). In App. [F,](#page-72-0) we show some identity of tensor network representations of fusion category symmetry. In App. [G,](#page-73-0) we provide a detailed computation of a cohomology group associated with $\text{Rep}(D_8)$. In App. [H,](#page-78-0) we explicitly compute the invariants of S^1 -parameterized families of Rep(D_8)-symmetric invertible states.

II. PRELIMINARIES

In this section, we briefly review mathematical tools that we will use to study SPT phases with fusion category symmetries. Throughout the paper, vector spaces are always finite-dimensional and the base field is always the field $\mathbb C$ of complex numbers. We denote the group cohomology by H_{gp} to distinguish it from other cohomology.

A. Fusion categories

Finite symmetries of 1+1d unitary bosonic systems are generally described by unitary multifusion categories [\[30,](#page-85-4) [31\]](#page-85-12). In what follows, (multi)fusion categories always refer to unitary ones. Each

³ Here, a charge refers to a one-dimensional representation of the self-interface algebra. In general, the charges form a non-abelian group, which is isomorphic to the group of automorphisms of a fiber functor.

object x of a multifusion category C labels a topological line \mathcal{L}_x , and each morphism $\mu \in \text{Hom}_{\mathcal{C}}(x, y)$ labels a topological junction between topological lines \mathcal{L}_x and \mathcal{L}_y . Here, Hom_C(x, y) denotes the finite dimensional C-vector space of all morphisms from x to y. An object x is said to be simple if its endomorphism space $\text{End}_{\mathcal{C}}(x) = \text{Hom}_{\mathcal{C}}(x, x)$ is one-dimensional. Correspondingly, a topological line \mathcal{L}_x labeled by a simple object x is indecomposable. The set of (representatives of isomorphisms classes of) simple objects is denoted by $\text{Simp}(\mathcal{C})$. The number of elements in $\text{Simp}(\mathcal{C})$ is finite.

A multifusion category $\mathcal C$ is equipped with various data that encode the algebraic structures of topological lines and topological junctions between them. Concretely, a multifusion category $\mathcal C$ consists of the data listed below. We refer the reader to [\[48\]](#page-86-3) for a more complete description of multifusion categories.

Fusion coefficients. The fusion of two topological lines \mathcal{L}_x and \mathcal{L}_y defines a tensor product $x \otimes y$ of objects x and y. The object $x \otimes y$ can always be decomposed into a finite direct sum of simple objects as

$$
x \otimes y \cong \bigoplus_{z \in \text{Simp}(\mathcal{C})} N_{xy}^z \ z, \quad N_{xy}^z \in \mathbb{Z}_{\geq 0},\tag{1}
$$

where the sum on the right-hand side is taken over all simple objects of \mathcal{C} . The non-negative integer N_{xy}^z is called a fusion coefficient. The unit of the above fusion rule is denoted by $1 \in \mathcal{C}$ and is called a unit object, which corresponds to the trivial topological line. The unit object 1 is equipped with isomorphisms $l_x : 1 \otimes x \to x$ and $r_x : x \otimes 1 \to x$, referred to as the left and right unitors, that satisfy appropriate coherence conditions. One can always take l_x and r_x to be the identity morphism by identifying $1 \otimes x$ and $x \otimes 1$ with $x \not\to 48$. When the unit object 1 is simple, the multifusion category $\mathcal C$ is called a fusion category. In what follows, we will restrict our attention to the case where $\mathcal C$ is fusion.^{[4](#page-5-0)}

Associator/ F -symbols. The fusion of topological lines is associative only up to isomorphism. Namely, there is a natural isomorphism

$$
\alpha_{x,y,z} : (x \otimes y) \otimes z \to x \otimes (y \otimes z), \tag{2}
$$

which relates two different ways of fusing three topological lines into a single line. The above isomorphism is called an associator and satisfies the pentagon equation represented by the following commutative diagram [\[49\]](#page-86-4):

$$
(x \otimes y) \otimes (z \otimes w)
$$

\n
$$
(x \otimes y) \otimes (z \otimes w)
$$

\n
$$
(x \otimes y) \otimes z) \otimes w
$$

\n
$$
\alpha_{x,y,z \otimes id_w}
$$

\n
$$
(x \otimes (y \otimes z)) \otimes w
$$

\n
$$
(x \otimes (y \otimes z)) \otimes w
$$

\n
$$
(x \otimes (y \otimes z)) \otimes w
$$

\n
$$
\alpha_{x,y,z,w}
$$

\n
$$
\alpha_{x,y,z,w}
$$

\n
$$
x \otimes ((y \otimes z) \otimes w)
$$

\n
$$
(3)
$$

 4 When the symmetry C is not fusion, there are non-trivial topological local operators labeled by endomorphisms of 1, meaning that a bosonic system with multifusion category symmetry $\mathcal C$ must have multiple ground states. Thus, to study SPT phases, it suffices to consider the case where the symmetry $\mathcal C$ is fusion.

More explicitly, the associator [\(2\)](#page-5-1) can be represented by a set of complex numbers $(F_w^{xyz})_{(u;\mu,\nu),(v;\rho,\sigma)}$ defined by

$$
\frac{\frac{x}{\lambda} + \frac{y}{\lambda}}{\frac{w}{\lambda} - \frac{y}{\lambda}} = \sum_{v \in \text{Simp}(\mathcal{C})} \sum_{\rho, \sigma} (F_w^{xyz})_{(u; \mu, \nu), (v; \rho, \sigma)} \left\{ \bigvee_{\rho, \sigma} \int_v^{y} z, (4) \right\}
$$

where the summation on the right-hand side is taken over all simple objects of $\mathcal C$ and all basis vectors of the Hom spaces. The complex numbers $(F_w^{xyz})_{(u;\mu,\nu),(v;\rho,\sigma)}$ are called F-symbols. The F-symbols depend on the choice of basis vectors of the Hom spaces. When $\mathcal C$ is unitary, one can choose these bases so that F_w^{xyz} becomes a unitary matrix for all simple objects $x, y, z, w \in \mathcal{C}$. In the physics literature, the associator in Eq. [\(4\)](#page-6-1) is often chosen to be the identity, which is always possible due to the Mac Lane strictness theorem $[50]$. We emphasize that the F-symbols can be non-trivial even if the associator $\alpha_{x,y,z}$ is the identity.

Quantum dimension. For a unitary fusion category \mathcal{C} , the quantum dimension dim(x) of a simple object x is the unique positive real number that satisfy

$$
\dim(x)\dim(y) = \sum_{z \in \text{Simp}(\mathcal{C})} N_{xy}^z \dim(z),\tag{5}
$$

where N_{xy}^z is the fusion coefficient in Eq. [\(1\)](#page-5-2). We note that $\dim(x)$ is not necessarily an integer. The quantum dimension of a non-simple object $\bigoplus_i x_i \in \mathcal{C}$ is given by the sum of the quantum dimensions $\sum_i \dim(x_i)$. The quantum dimension of x agrees with that of the dual x^* .

The simplest example of a fusion category is the category Vec of finite dimensional C-vector spaces. Objects and morphisms of this category are finite dimensional vector spaces and linear maps between them. The tensor product of objects is defined by the ordinary tensor product of vector spaces. In particular, the unit object of Vec is a one-dimensional vector space C, which is the unique simple object up to isomorphism. The associators and F -symbols are all trivial. The quantum dimension of an object $x \in V_{\text{ec}}$ is given by the dimension of the vector space x, which is a non-negative integer.

B. Fiber functors and SPT phases

In this subsection, we recall the definition of fiber functors and their relation to SPT phases with fusion category symmetries.

A fiber functor of a fusion category C is defined as a tensor functor from C to Vec. Let us unpack this definition. First of all, a fiber functor $F : C \to \text{Vec}$ is a functor, that is, it maps an object $x \in \mathcal{C}$ to a vector space $F(x) \in$ Vec and maps a morphism $\mu \in \text{Hom}_{\mathcal{C}}(x, y)$ to a linear map $F(\mu) \in \text{Hom}_{\text{Vec}}(F(x), F(y))$. The map on morphisms should preserve the composition law and the identity, i.e.,

$$
F(\nu \circ \mu) = F(\nu) \circ F(\mu), \quad F(\mathrm{id}_x) = \mathrm{id}_{F(x)} \tag{6}
$$

for any composable morphisms μ, ν and any object $x \in \mathcal{C}$. The functor $F: \mathcal{C} \to \mathcal{V}$ ec is called a fiber functor if it is equipped with a natural set of isomorphisms

$$
J_{x,y}: F(x) \otimes F(y) \to F(x \otimes y), \quad \varphi: \mathbb{C} \to F(\mathbb{1})
$$
\n⁽⁷⁾

that satisfy the following commutative diagram:^{[5](#page-7-0)}

$$
(F(x) \otimes F(y)) \otimes F(z) \longrightarrow F(x \otimes y) \otimes F(z) \longrightarrow F((x \otimes y) \otimes z) \longrightarrow F(\alpha_{x,y,z}^c)
$$

\n
$$
\xrightarrow[\alpha_{F(x),F(y),F(z)}]{} F(x) \otimes (F(y) \otimes F(z)) \xrightarrow[\mathrm{id}_{F(x)} \otimes J_{y,z}]} F(x) \otimes F(y \otimes z) \longrightarrow F(x \otimes (y \otimes z))
$$

where $\alpha^{\mathcal{C}}$ and α^{Vec} denote the associators of \mathcal{C} and Vec, respectively. The isomorphism $\varphi : \mathbb{C} \to F(1)$ can always be chosen to be the identity. Therefore, a fiber functor is characterized by a set of vector spaces $\{F(x) \mid x \in C\}$ together with isomorphisms $\{J_{x,y} : F(x) \otimes F(y) \to F(x \otimes y) \mid x, y \in C\}$ making the diagram [\(8\)](#page-7-1) commute.

Equivalently, a fiber functor can also be defined as a C -module category whose simple object is unique up to isomorphism. Here, a C-module category $\mathcal M$ is a semisimple category equipped with a tensor functor $F: \mathcal{C} \to \text{End}(\mathcal{M})$, where $\text{End}(\mathcal{M})$ is the category of endofunctors of M. Indeed, when the simple object of $\mathcal M$ is unique, $\mathcal M$ is equivalent to Vec as a semisimple category, and hence the tensor functor $F : \mathcal{C} \to \text{End}(\mathcal{M}) \cong \text{Vec}$ becomes a fiber functor. Conversely, a fiber functor $F: \mathcal{C} \to \mathcal{V}$ ec allows us to endow the target category Vec with a \mathcal{C} -module structure. Thus, a fiber functor of $\mathcal C$ can be identified with a $\mathcal C$ -module category with a single simple object.

Since a fiber functor F preserves the tensor product of objects up to isomorphism J , it follows that

$$
F(x) \otimes F(y) \cong \bigoplus_{z \in \text{Simp}(\mathcal{C})} N_{xy}^z F(z), \tag{9}
$$

where N_{xy}^z is the fusion coefficient of C. The above equation implies

$$
\dim(F(x))\dim(F(y)) = \sum_{z \in \text{Simp}(\mathcal{C})} N_{xy}^z \dim(F(z)).\tag{10}
$$

Recalling that the quantum dimension is the unique positive one-dimensional representation of the fusion rules, we find

$$
\dim(F(x)) = \dim(x),\tag{11}
$$

which means that $F(x) \in \text{Vec}$ is a vector space of dimension dim(x), i.e., $F(x) \cong \mathbb{C}^{\dim(x)}$. In particular, since the dimension of a vector space $F(x)$ must be a positive integer, a fusion category C admits a fiber functor only if the quantum dimension of every object $x \in C$ is a positive integer, namely, dim $(x) \in \mathbb{Z}_{\geq 1}$.

(8)

⁵ We do not specify the coherence conditions on φ here because we will not use them in what follows.

We emphasize that not every fusion category admits a fiber functor. In general, a fusion category C admits a fiber functor if and only if C is equivalent to the category $\text{Rep}(H)$ of representations of a finite dimensional semisimple Hopf algebra H $[48]$.^{[6](#page-8-0)} Here, Rep(H) denotes the category whose objects are finite dimensional representations of H and morphisms are intertwiners between them. A fusion category $\mathcal C$ is said to be non-anomalous if it admits a fiber functor. Otherwise, $\mathcal C$ is said to be anomalous [\[32\]](#page-85-5).

For later convenience, let us introduce a matrix representation of the isomorphism $J_{x,y}$. To this end, we first choose a basis $\{(\phi_x)_i \mid i = 1, 2, \cdots, \dim(x)\}$ of the vector space $F(x)$ for each simple object $x \in \mathcal{C}$. We also fix a basis $\{(\phi_{xy}^z)_{\mu} | \mu = 1, 2, \cdots, N_{xy}^z\}$ of the Hom space $\text{Hom}_{\mathcal{C}}(x \otimes y, z)$. For these bases, one can write down the matrix elements of the isomorphism $J_{x,y}$ as follows:

$$
F((\phi_{xy}^z)_{\mu}) \circ J_{x,y}((\phi_x)_i \otimes (\phi_y)_j) = \sum_k J_{(i,j),(z;k,\mu)}^{xy}(\phi_z)_k, \quad J_{(i,j),(z;k,\mu)}^{xy} \in \mathbb{C}.
$$
 (12)

The set of complex numbers $\{J_{(i)}^{xy}$ $\lim_{(i,j),(z;k,\mu)}\mid z\in \mathrm{Simp}(\mathcal{C}),\ i=1,2,\cdots,\mathrm{dim}(x),\ j=1,2,\cdots,\mathrm{dim}(y),\ k=1,2,\cdots$ $1, 2, \cdots, \dim(z), \mu = 1, 2, \cdots, N_{xy}^z$ form a matrix representation of $J_{x,y}$. That is, each $J_{(i, z)}^{xy}$ $(i,j),(z;k,\mu)$ can be regarded as a component of a square matrix J^{xy} of size $\dim(x)\dim(y)$.

A simple example of a fiber functor is the forgetful functor of the representation category $\text{Rep}(H)$ of a Hopf algebra H. The forgetful functor maps an object $x \in \text{Rep}(H)$ to the underlying vector space of x and maps a morphism to the underlying linear map. The structure isomorphisms J and φ associated with the forgetful functor are trivial.^{[7](#page-8-1)}

Classification of SPT phases. An SPT phase is a symmetric gapped phase that has a unique ground state on any space without boundaries. It is well known that 1+1d bosonic SPT phases with finite internal symmetry group G are classified by the second group cohomology of G [\[3,](#page-84-7) [4,](#page-84-5) [8\]](#page-84-2). More generally, it was proposed in [\[32\]](#page-85-5) that 1+1d bosonic SPT phases with symmetry $\mathcal C$ are classified by fiber functors of $\mathcal C$. In particular, a fusion category symmetry $\mathcal C$ admits an SPT phase if and only if $\mathcal C$ is non-anomalous. We note that the set of SPT phases with symmetry $\mathcal C$ does not have a natural group structure in general.

The data of a fiber functor can be directly interpreted in terms of topological field theory that describes the low energy limit of an SPT phase $[32, 54]$ $[32, 54]$ $[32, 54]$. Concretely, in the context of topological field theory, the vector space $F(x)$ is understood as the state space of the x-tiwsted sector on a circle. Similarly, the isomorphism $J_{x,y} : F(x) \otimes F(y) \to F(x \otimes y)$ can be understood as the transition amplitude on a pants diagram with topological lines x and y inserted on the two legs. The complex number $J_{(i)}^{xy}$ $\lim_{(i,j),(z,k,\mu)}$ represents a three-point function of defect operators, which is related to the transition amplitude on a pair of pants via the state-operator correspondence. See Fig. [1](#page-9-1) for diagrammatic representations of these data.

Classification of general gapped phases. The classification of more general gapped phases with fusion category symmetries was also proposed in [\[32\]](#page-85-5). Specifically, general 1+1d bosonic gapped phases with symmetry $\mathcal C$ are in one-to-one correspondence with module categories over $\mathcal C$ [\[32,](#page-85-5) [55\]](#page-86-7). In particular, SPT phases with symmetry $\mathcal C$ correspond to $\mathcal C$ -module categories with a single simple object, which can be identified with fiber functors of C . As in the case of SPT phases, the data

 6 Any fusion category that may not admit a fiber functor is equivalent to the category of representations of a weak Hopf algebra [\[51,](#page-86-8) [52\]](#page-86-9), which is a generalization of a Hopf algebra [\[53\]](#page-86-10).

⁷ The complex numbers $\{J_{(i,j),(z;k,\mu)}^{xy}\}$ can be non-trivial. This is analogous to the situation where F-symbols of a fusion category C can be non-trivial even if the associator is trivial (i.e., C is strict).

FIG. 1. (a) The vector space $F(x)$ represents the state space on a circle twisted by $x \in C$. (b) The isomorphism $J_{x,y}$ represents the transition amplitude on a pair of pants. (c) The matrix element of $J_{x,y}$ represents the three-point function.

of a module category can be interpreted in terms of topological field theory that describes the low energy limit of a gapped phase with symmetry $[56]$. In this paper, we will focus only on $1+1d$ SPT phases except for Sec. [VI.](#page-42-0)

C. Isomorphisms of fiber functors

In this subsection, we recall the definition of isomorphisms between fiber functors, which play a crucial role in studying interfaces and parameterized families of SPT phases in later sections.

Let (F, J) and (F', J') be fiber functors of a fusion category \mathcal{C}^8 \mathcal{C}^8 . A monoidal natural transformation $\eta: F \Rightarrow F'$ between fiber functors F and F' is a set of linear maps

$$
\eta_x: F(x) \to F'(x), \quad \forall x \in \mathcal{C}
$$
\n(13)

that are natural in $x \in \mathcal{C}$ and satisfy the following commutative diagram:^{[9](#page-9-3)}

$$
F(x) \otimes F(y) \xrightarrow{J_{x,y}} F(x \otimes y)
$$

\n
$$
\eta_x \otimes \eta_y \downarrow \qquad \qquad \downarrow \eta_{x \otimes y}
$$

\n
$$
F'(x) \otimes F'(y) \xrightarrow{J'_{x,y}} F'(x \otimes y)
$$
\n(14)

The monoidal natural transformation η is called a monoidal natural isomorphism if the linear maps η_x are isomorphisms for all $x \in \mathcal{C}$. In what follows, a monoidal natural isomorphism between fiber functors is simply referred to as an isomorphism. In particular, an isomorphism from fiber functor F to itself is called an automorphism of F. Once we choose bases of $F(x)$ and $F'(x)$ to be $\{(\phi_x)_i\}$ and $\{(\phi_x')_i\}$ respectively, we can explicitly write down a monoidal natural transformation $\eta: F \Rightarrow F'$ as

$$
\eta_x((\phi_x)_i) = \sum_{1 \le j \le \dim(x)} (\eta_x)_{ij}(\phi'_x)_j,\tag{15}
$$

where $(\eta_x)_{ij}$ is the (i, j) -component of a $dim(x) \times dim(x)$ matrix η_x . When $\eta_x : F \Rightarrow F'$ is an isomorphism, the matrix η_x is invertible for all $x \in \mathcal{C}$.

⁸ Here and in the subsequent sections, we take the isomorphism φ in Eq. [\(7\)](#page-7-2) to be the identity without loss of generality.

⁹ The linear map η_x is said to be natural in x if it satisfies $\eta_y \circ F(f) = F'(f) \circ \eta_x$ for any morphism $f \in \text{Hom}_{\mathcal{C}}(x, y)$. We note that the structure isomorphisms such as $\alpha_{x,y,z}$ in Eq. [\(2\)](#page-5-1) and $J_{x,y}$ in Eq. [\(7\)](#page-7-2) are required to be natural.

The composition of isomorphisms $\eta : F \Rightarrow F'$ and $\eta' : F' \Rightarrow F''$ defines an isomorphism $\eta' \circ \eta : F \Rightarrow F''$. In particular, automorphisms of a fiber functor F form a finite group $\text{Aut}^{\otimes}(F)$ with the multiplication given by the composition.

For later convenience, let us explicitly write down the commutative diagram [\(14\)](#page-9-4) in terms of matrix representations of $J_{x,y}$ and η_x . To this end, we notice that Eq. [\(14\)](#page-9-4) is equivalent to

$$
\eta_z \circ F((\phi_{xy}^z)_{\mu}) \circ J_{x,y} = F'((\phi_{xy}^z)_{\mu}) \circ J'_{x,y} \circ (\eta_x \otimes \eta_y), \quad \forall (\phi_{xy}^z)_{\mu} \in \text{Hom}_{\mathcal{C}}(x \otimes y, z). \tag{16}
$$

By taking the matrix elements of both sides of the above equation, we obtain

$$
\sum_{k} J_{(i,j),(z;k,\mu)}^{xy}(\eta_z)_{kk'} = \sum_{i',j'} (\eta_x)_{ii'}(\eta_y)_{jj'} (J')_{(i',j'),(z;k',\mu)}^{xy}.
$$
 (17)

Classification of parameterized families of SPT states. We will argue in Sec. [V](#page-33-0) that $Aut^{\otimes}(F)$ classifies S^1 -paramererized families of invertible states in a C-SPT phase labeled by F. The composition of automorphisms of F corresponds to the concatenation of two S^1 -families. In contrast to the classification of SPT phases, the classification of S^1 -parameterized families of SPT states has a natural group structure.

When $\mathcal C$ is an ordinary group symmetry, the set of SPT phases also has a group structure. Accordingly, the classification of parameterized families does not depend on a phase. This is because stacking an SPT state with the whole family gives another family of SPT states in a different phase. Indeed, as we will see below, the group $Aut^{\otimes}(F)$ does not depend on F when C is an ordinary group symmetry. On the other hand, when $\mathcal C$ is a non-invertible symmetry, the set of SPT phases does not necessarily have a group structure. Consequently, the classification of parameterized families may depend on a phase. Indeed, as we will see below, the group $Aut^{\otimes}(F)$ generally depends on F when $\mathcal C$ is non-invertible.

Tannaka-Krein duality. Let us consider the set $\text{End}(F)$ of all (not necessarily monoidal) natural transformations from F to itself. The set $End(F)$ is a vector space because for any natural transformations $\eta, \eta' \in \text{End}(F)$ and complex numbers $\lambda, \lambda' \in \mathbb{C}$, one can define a natural transformation $\lambda \eta + \lambda' \eta' \in \text{End}(F)$ by setting $(\lambda \eta + \lambda' \eta')_x := \lambda \eta_x + \lambda' \eta'_x$ for all $x \in \mathcal{C}$. The vector space $\text{End}(F)$ has the structure of an algebra, whose multiplication is given by the composition of natural transformations. Furthermore, the monoidal structure of F allows us to define a comultiplication of natural transformations, which, together with the antipode, makes $\text{End}(F)$ into a Hopf algebra. The representation category $\text{Rep}(\text{End}(F))$ of this Hopf algebra turns out to be equivalent to the original fusion category C, see, e.g., [\[48,](#page-86-3) Chapter 5]. In particular, when $\mathcal{C} = \text{Rep}(H)$ and $F : \mathcal{C} \to \text{Vec}$ is the forgetful functor, the algebra $\text{End}(F)$ is isomorphic to H as a Hopf algebra, meaning that one can reconstruct a Hopf algebra from its representation category. This phenomenon is known as Tannaka-Krein duality [\[57–](#page-86-12)[61\]](#page-86-13).

We note that monoidal natural automorphisms of a fiber functor $F : C \to \text{Vec}$ are group-like elements of Hopf algebra $\text{End}(F)$. In particular, when C is $\text{Rep}(H)$ and F is the forgetful functor, the group $Aut^{\otimes}(F)$ of automorphisms of F is isomorphic to the group $G(H)$ of group-like elements of H. For example, when F is the forgetful functor of $Rep(G)$ with G a finite group, we have a group isomorphism $Aut^{\otimes}(F) \cong G$. This fact will naturally appear in later sections where we study parameterized families of SPT states with fusion category symmetries.

Examples. Let us see a few simple examples of fiber functors and their automorphisms.

• Vec_G. We first consider fiber functors of Vec_G, the category of G-graded vector spaces. Simple objects of Vec_G are labeled by elements of G, and the fusion rules are given by the group multiplication law:

$$
g \otimes h = gh, \quad \forall g, h \in G. \tag{18}
$$

The associator and F-symbols of Vec_G are all trivial. A fiber functor $F : \text{Vec}_G \to \text{Vec}$ maps every simple object g to a one-dimensional vector space $F(g) \cong \mathbb{C}$ because the quantum dimension of g is one. Therefore, the fiber functor F is completely characterized by a set of isomorphisms $J_{q,h}$: $F(g) \otimes F(h) \to F(gh)$. Since the source and target of $J_{q,h}$ are both one-dimensional, it can be identified with a complex number $J(g, h) \in U(1)$.^{[10](#page-11-0)} With this identification, the commutative diagram [\(8\)](#page-7-1) reduces to the cocycle condition on $J: G \times G \rightarrow$ $U(1)$, i.e.,

$$
J(h,k)J(g,hk) = J(gh,k)J(g,h), \quad \forall g,h,k \in G.
$$
\n
$$
(19)
$$

Similarly, an isomorphism $\eta : F \Rightarrow F'$ between two fiber functors F and F' is characterized by the set of isomorphisms $\eta_g : F(g) \to F'(g)$, which can be identified with complex numbers $\eta(g) \in U(1)$. The commutative diagram [\(14\)](#page-9-4) reduces to

$$
J(g,h) = J'(g,h)\delta\eta(g,h), \quad \delta\eta(g,h) = \eta(g)\eta(h)/\eta(gh), \tag{20}
$$

where J and J' are 2-cocycles associated with F and F' respectively. The above equation implies that two fiber functors are isomorphic to each other if and only if the associated 2-cocycles J and J' are in the same cohomology class. Thus, isomorphism classes of fiber functors of Vec_G are classified by the second group cohomology $H_{gp}^2(G, U(1))$. This agrees with the well-known classification of $1+1d$ bosonic SPT phases with symmetry G [\[3,](#page-84-7) [4,](#page-84-5) [8\]](#page-84-2).

Equation [\(20\)](#page-11-1) also implies that an automorphism of a fiber functor F is given by a function $\eta: G \to U(1)$ that satisfies the cocycle condition

$$
\delta \eta(g, h) = 1, \quad \forall g, h \in G. \tag{21}
$$

Therefore, the group Aut[⊗](F) of automorphisms of F is isomorphic to the first group cohomology $H^1_{gp}(G, U(1))$ regardless of a fiber functor F. We note that $Aut^\otimes(F) = H^1_{gp}(G, U(1))$ classifies \widetilde{S}^1 -parameterized families of G-symmetric invertible states in a 1+1d bosonic SPT phase [\[15,](#page-84-8) [16,](#page-84-9) [62,](#page-86-14) [63\]](#page-86-15).

• $\mathbf{Rep}(G)$. As another example, we consider fiber functors of the representation category $Rep(G)$ of a finite group G. It is known that fiber functors of $Rep(G)$ are classified by (equivalence classes of) pairs (K, ω) , where K is a subgroup of G and $\omega \in \mathbb{Z}_{\text{gp}}^2(K, \mathcal{U}(1))$ is a second cocycle such that the twisted group algebra $\mathbb{C}[K]^\omega$ is simple [\[48,](#page-86-3) Corollary 7.12.20]. In particular, the fiber functor corresponding to the trivial subgroup $K = \{e\}$ is the forgetful functor, which exists for all G.

Due to Tannaka-Krein duality, the group of automorphisms of the forgetful functor of $\text{Rep}(G)$ is isomorphic to G itself. This shows that the group of automorphisms of a fiber functor can

¹⁰ We recall that fusion categories are supposed to be unitary in this paper. Thus, $J(g, h)$ takes values in U(1) rather than \mathbb{C}^{\times} .

be non-abelian when the fusion category has non-invertible objects. This is in contrast to the fact that the group of automorphisms $\text{Aut}^{\otimes}(F) = \text{H}^1_{\text{gp}}(G, \text{U}(1))$ is always abelian for an invertible (i.e., pointed) fusion category Vec_G .

When the fiber functor $F : \text{Rep}(G) \to \text{Vec}$ corresponds to a non-trivial pair (K, ω) , the group of automorphisms of F is not necessarily isomorphic to G . One can indeed construct such a fiber functor if there exists a group G' that is not isomorphic to G but has the equivalent representation category as G. Specifically, the composition of an equivalence $\Phi : \text{Rep}(G) \rightarrow$ $Rep(G')$ and the forgetful functor of $Rep(G')$ gives us a fiber functor $F : Rep(G) \to \text{Vec}$, whose group of automorphisms is isomorphic to G' due to Tannaka-Krein duality. An example of such a pair (G, G') was constructed in $[64]$, where such G and G' are called isocategorical. The fact that $\text{Aut}^{\otimes}(F)$ depends on a fiber functor F is also in contrast to the case of invertible symmetry Vec_G .

D. Matrix product states with fusion category symmetries

The ground states of a gapped phase in $1+1$ dimensions can be efficiently represented by matrix product states (MPSs) [\[33,](#page-85-6) [65–](#page-86-17)[67\]](#page-86-18). Furthermore, the symmetry operators acting on an MPS can often be represented by matrix product operators (MPOs) [\[36–](#page-85-8)[39\]](#page-85-13). Therefore, MPSs symmetric under the action of MPOs are useful tools to study 1+1d gapped phases with fusion category symmetries. In this subsection, following [\[37\]](#page-85-11), we introduce MPSs symmetric under MPOs and discuss the invariants associated with such MPSs. See, e.g., [\[33\]](#page-85-6) for a recent review of MPSs and MPOs.

Matrix product states. An MPS is a tensor network state constructed from a three-leg tensor A as

$$
|A\rangle = \longrightarrow \begin{array}{c}\n\downarrow & \downarrow & \downarrow \\
A & A & A & A & A \\
\end{array}
$$
\n(22)

where each vertical leg represents a physical Hilbert space $\mathcal{H} \cong \mathbb{C}^d$ and each horizontal leg represents a bond Hilbert space $V \cong \mathbb{C}^D$. The arrows on the edges specify the source and target of each tensor $A: V \to V \otimes \mathcal{H}$. These arrows will be omitted when no confusion can arise.^{[11](#page-12-1)} For simplicity, we only consider translationally invariant MPSs in this paper, i.e., the MPS tensor A does not depend on sites. Furthermore, the MPSs discussed in this paper are supposed to obey a periodic boundary condition unless otherwise stated.

Given a basis $\{b_i \mid i = 1, 2, \dots, d\}$ of the physical Hilbert space \mathcal{H} , one can express the MPS [\(22\)](#page-12-2) on a periodic chain as

$$
|A\rangle = \sum_{i_1, i_2, \cdots, i_L} \text{tr}(A^{i_1} A^{i_2} \cdots A^{i_n}) |b_{i_1}, b_{i_2}, \cdots, b_{i_L}\rangle, \qquad (23)
$$

where A^i is a $D \times D$ matrix and L denotes the number of lattice sites. An MPS tensor A is said to be injective if the set $\{A^i \mid i = 1, 2, \cdots, d\}$ generates the full matrix algebra $M_D(\mathbb{C})$. An injective MPS can be realized as the unique ground state of a gapped Hamiltonian belonging to an SPT phase [\[68\]](#page-86-19).

 11 The choice of the orientation of an arrow is just a matter of convention.

Conversely, the ground state of an SPT phase can always be approximated by an injective MPS [\[33,](#page-85-6) [65](#page-86-17)[–67\]](#page-86-18). As such, we can restrict our attention to injective MPSs when studying SPT phases.

Matrix product operators. An MPO is a tensor network operator built from a four-leg tensor O as

^O^b ⁼ O . (24)

We denote the bond Hilbert space of an MPO $\widehat{\mathcal{O}}$ by $V_{\mathcal{O}} \cong \mathbb{C}^{\chi}$, where χ is the bond dimension. The arrows on the edges show that the MPO tensor $\mathcal O$ is a linear map from $M_{\mathcal O}\otimes\mathcal H$ to $\mathcal H\otimes M_{\mathcal O}$. For a given basis $\{b_i \mid i = 1, 2, \dots, d\}$ of the physical Hilbert space \mathcal{H} , the MPO [\(24\)](#page-13-0) on a periodic chain can be expressed in terms of $\chi \times \chi$ matrices $\{\mathcal{O}^{ij} \mid i, j = 1, 2, \cdots, d\}$ as

$$
\widehat{\mathcal{O}} = \sum_{i_1 \cdots, i_L} \sum_{j_1, \cdots, j_L} \text{tr}(\mathcal{O}^{i_1 j_1} \cdots \mathcal{O}^{i_L j_L}) |b_{i_1}, \cdots, b_{i_L}\rangle \langle b_{j_1}, \cdots, b_{j_L}|.
$$
\n(25)

An MPO is said to be injective if the set $\{\mathcal{O}^{ij} \mid i, j = 1, 2, \cdots, d\}$ generates the full matrix algebra $M_{\mathcal{Y}}(\mathbb{C}).$

MPOs can be used to represent fusion category symmetries on the lattice [\[34,](#page-85-14) [36–](#page-85-8)[39\]](#page-85-13). Indeed, for any fusion category C , there exists an MPO representation of C , i.e., a set of injective MPOs $\{\widehat{\mathcal{O}}_x \mid x \in \text{Simp}(\mathcal{C})\}$ together with three-leg tensors $\{(\phi_{xy}^z)_{\mu} \mid \mu = 1, 2, \cdots, N_{xy}^z\}$ and $\{(\overline{\phi}_{xy}^z)_{\mu} \mid \mu = 1, 2, \cdots, N_{xy}^z\}$ $1, 2, \dots, N_{xy}^z$ that allow us to locally fuse two MPOs $\hat{\mathcal{O}}_x$ and $\hat{\mathcal{O}}_y$ as follows:

$$
\mathcal{O}_{x\otimes y} := \mathcal{O}_x \mathcal{O}_y = \overbrace{\rightarrow \bigodot^{\mathcal{O}_x}}^{\mathcal{O}_x} = \sum_{z \in \text{Simp}(\mathcal{C})} \sum_{1 \le \mu \le N_{xy}^z} (\phi_{xy}^z)_{\mu} \overbrace{\rightarrow \bigodot^{\mathcal{O}_z} (\overline{\phi}_{xy}^z)_{\mu}}^{\mathcal{O}_x} .
$$
 (26)

The tensors $\{(\phi_{xy}^z)_\mu\}$ and $\{(\overline{\phi}_{xy}^z)_\mu\}$ are called fusion and splitting tensors respectively. These tensors satisfy the following orthogonality relation:

$$
z \rightarrow \mathbf{e}^{\nu} \xrightarrow{\mu} z' = \delta_{\mu,\nu} \delta_{z,z'} \rightarrow z \tag{27}
$$

On the left-hand side, the fusion and splitting tensors $(\phi_{xy}^z)_{\mu}$ and $(\overline{\phi}_{xy}^z)_{\nu}$ are abbreviated as μ and ν , respectively. Equations [\(26\)](#page-13-1) and [\(27\)](#page-13-2) imply that an MPO representation of C on a periodic chain obeys the same fusion rules as \mathcal{C} , i.e.,

$$
\widehat{\mathcal{O}}_x \widehat{\mathcal{O}}_y = \sum_z N_{xy}^z \widehat{\mathcal{O}}_z. \tag{28}
$$

Using the fusion and splitting tensors in Eq. (26) , one can extract the F-symbols of C as follows:

$$
w \longrightarrow \underbrace{\overbrace{}^{u} \overbrace{}^{w}}_{z} = \sum_{v} \sum_{\rho, \sigma} (F_{w}^{xyz})_{(u; \mu, \nu), (v, \rho, \sigma)} \stackrel{w \longrightarrow \overbrace{}^{w}}{\longrightarrow} \underbrace{\overbrace{}^{x}}_{v} \overbrace{}^{x}}_{z}.
$$
 (29)

The pentagon equation for the F-symbols follows from the associativity of MPOs. Explicit construction of MPOs $\{O_x \mid x \in C\}$ that implement general fusion category symmetry C is discussed in, e.g., [\[36,](#page-85-8) [38,](#page-85-15) [39\]](#page-85-13).

As a simple example, let us consider an MPO representation of $Rep(G)$. For concreteness, we choose the physical Hilbert space $\mathcal H$ to be a G-dimensional vector space faithfully graded by G , i.e.,

$$
\mathcal{H} = \bigoplus_{g \in G} \mathbb{C} \left| g \right\rangle,\tag{30}
$$

where $\{|g\rangle | g \in G\}$ is a basis of H. The bond Hilbert space of an MPO $\widehat{\mathcal{O}}_{\rho}$ is given by the representation space V_ρ of representation $\rho \in \text{Rep}(G)$. One can define the MPO tensor \mathcal{O}_ρ as

$$
\mathcal{O}_{\rho} = \sum_{g \in G} \rho(g) \otimes |g\rangle \langle g| \,,\tag{31}
$$

where $\rho(g)$ is the representation matrix acting on the bond Hilbert space V_{ρ} . We note that the MPO \mathcal{O}_{ρ} labeled by an irreducible representation ρ is injective due to Schur's lemma. For the above MPO representation, the fusion tensors are simply given by projections from the tensor product representation $\rho_1 \otimes \rho_2 \cong \bigoplus_{\rho_3} N_{\rho_1\rho_2\rho_3}^{\rho_3}$ to its fusion channel ρ_3 . Similarly, the splitting tensors are given by inclusion maps from ρ_3 to $\rho_1 \otimes \rho_2$. By construction, the F-symbols defiined by Eq. (29) agree with those of fusion category Rep(G). The MPO representation [\(31\)](#page-14-0) can be generalized to that of $\text{Rep}(H)$ where H is a Hopf algebra [\[47\]](#page-86-2) or a weak Hopf algebra [\[36\]](#page-85-8). Here, we recall that a general fusion category is equivalent to the representation category of a weak Hopf algebra, while a general non-anomalous fusion category is equivalent to the representation category of a Hopf algebra.

MPSs symmetric under MPOs. Let ${\{\widehat{\mathcal{O}}_x \mid x \in \text{Simp}(\mathcal{C})\}}$ be an MPO representation of a fusion category C. An injective MPS A is said to be C-symmetric if there exist three-leg tensors $\{(\phi_x)_i\}$ and $\{(\overline{\phi}_x)_i\}$ called action tensors, which allow us to apply an MPO $\hat{\mathcal{O}}_x$ locally to an injective MPS $|A\rangle$ as follows [\[37\]](#page-85-11):^{[12](#page-14-1)}

$$
\mathcal{O}_x A = \begin{array}{c} \downarrow \mathcal{O}_x \\ \hline \uparrow A \end{array} = \sum_i \begin{array}{c} \downarrow \\ \hline \downarrow A \end{array} \begin{array}{c} \downarrow \\ A \end{array} \begin{array}{c} \downarrow \\ \hline \downarrow A \end{array} \tag{32}
$$

Here, the index i runs from 1 to some positive integer n_x . The action tensors are required to satisfy the following orthogonality relation:

$$
\sum_{\left(\phi_x\right)_i} \overline{\phi_x}_{j}} = \delta_{ij} \longrightarrow . \tag{33}
$$

The sets of action tensors $\{(\phi_x)_i \mid i = 1, 2, \cdots, n_x\}$ and $\{(\phi_x)_i \mid i = 1, 2, \cdots, n_x\}$ are simply written as ϕ and $\overline{\phi}$ when no confusion can arise. Equations [\(32\)](#page-14-2) and [\(33\)](#page-14-3) imply that a C-symmetric injective MPS |A⟩ on a periodic chain is invariant under the action of C up to scalar multiplication:
 $\hat{O}_x |A\rangle = n_x |A\rangle$.

$$
\hat{\mathcal{O}}_x |A\rangle = n_x |A\rangle. \tag{34}
$$

 12 In [\[37\]](#page-85-11), a block-injective MPS symmetric under MPO symmetry is defined by the condition that the set of MPSs with arbitrary boundary conditions is closed under the action of the MPO algebra. The existence of the action tensors follows from this condition.

This equation forces $n_x = \dim(x)$ because the quantum dimension is the unique one-dimensional representation of the fusion rules such that all objects are represented by positive real numbers.

Using the action tensors in Eq. [\(32\)](#page-14-2), one can define a set of complex numbers L_{ij}^{xy} $_{(i,j),(z;k,\mu)}^{xy}$ called L-symbols:

$$
\overbrace{(\phi_x)_i \ (\phi_y)_j}^{\overbrace{\phi_x^2}_b} = \sum_z \sum_{k,\mu} L_{(i,j),(z;k,\mu)}^{xy} \underbrace{\overbrace{(\phi_x^2)_\mu}_{(\phi_z)_k}}^{\overbrace{\phi_x^2}_b}.
$$
 (35)

The associativity of MPO actions implies that the L-symbols satisfy consistency conditions anal-ogous to the pentagon equation [\[37\]](#page-85-11). The inverse of the L-symbol L^{xy} is denoted by \overline{L}^{xy} , which obeys

$$
\overrightarrow{\mathbf{L}_{\mathcal{O}}(z_{xy})} = \sum_{z} \sum_{k,\mu} \overrightarrow{L}_{(z_k,k,\mu),(i,j)}^{\mathcal{I}_{\mathcal{V}}(z_{xy})} \overrightarrow{\mathbf{L}_{(\overrightarrow{\phi}_z)k}}. \tag{36}
$$

Due to the orthogonality relation [\(33\)](#page-14-3), the L-symbols L_{ij}^{xy} $\lim_{(i,j),(z;k,\mu)}$ and $\overline{L}_{(z)}^{xy}$ $\chi_{(z;k,\mu),(i,j)}^{x,y}$ can be expressed in terms of the following closed diagrams:

$$
L_{(i,j),(z;k,\mu)}^{xy} = \frac{1}{D} \underbrace{\left(\overline{\phi}_z\right)_k}_{\left(\phi_x\right)_i \left(\overline{\phi}_x\right)_\mu \left(\phi_y\right)_j}, \quad \overline{L}_{(z;k,\mu),(i,j)}^{xy} = \frac{1}{D} \underbrace{\left(\overline{\phi}_x\right)_i \left(\overline{\phi}_x\right)_\mu}_{\left(\phi_z\right)_k} \left(\overline{\phi}_y\right)_j},\tag{37}
$$

where D is the bond dimension of the MPS.

The L-symbols have gauge ambiguities because they depend on the choice of action tensors. A gauge transformation of the action tensors is defined by

$$
(\phi_x)_i \to {}^U(\phi_x)_i = \sum_j (U_x)_{ij} (\phi_x)_j, \quad (\overline{\phi}_x)_i \to {}^U(\overline{\phi}_x)_i = \sum_j (\overline{\phi}_x)_j (U_x)_{ji}^{-1},
$$
\n(38)

where U_x is a dim(x) \times dim(x) invertible matrix. Correspondingly, a gauge transformation of the L-symbols is given by

$$
L_{(i,j),(z;k,\mu)}^{xy} \to {}^{U}L_{(i,j),(z;k,\mu)}^{xy} = \sum_{i',j',k'} (U_x)_{ii'}(U_y)_{jj'}(U_z)_{k'k}^{-1} L_{(i',j'),(z;k',\mu)}^{xy}, \tag{39}
$$

The L-symbols that are related by the above gauge transformation are said to be equivalent.

Equivalence classes of L-symbols classify C -symmetric SPT phases whose ground states are represented by injective MPSs [\[37\]](#page-85-11).[13](#page-15-0) This is consistent with the classification of SPT phases via

 13 Ref. [\[37\]](#page-85-11) also defined L-symbols for block-injective MPSs symmetric under MPO symmetries. Such L-symbols can be identified with the data of a module category and characterize more general gapped phases.

fiber functors. Indeed, the L-symbols associated with a C -symmetric injective MPS can be identified with the data of a fiber functor as

$$
L_{(i,j),(z;k,\mu)}^{xy} = J_{(i,j),(z;k,\mu)}^{xy},
$$
\n(40)

where the right-hand side is the matrix element of the isomorphism $J_{x,y}$ associated with a fiber functor, see Eq. [\(12\)](#page-8-2). The consistency conditions on the L-symbols agree with the coherence conditions (8) on the isomorphism J. Moreover, the gauge transformation (39) of the L-symbols defines an isomorphism [\(17\)](#page-10-0) of the corresponding fiber functor, meaning that equivalence classes of L-symbols correspond to isomorphism classes of fiber functors. In Sec. [III,](#page-17-0) we will show that one can extract the data of a fiber functor/ L -symbols from an injective MPS by using the triple inner product introduced in [\[69\]](#page-87-0).

Completeness conditions. Before proceeding, we mention that the action tensors $\{(\phi_x)_i, (\phi_x)_i\}$ and the fusion tensors $\{(\phi_{xy}^z)_\mu, (\overline{\phi}_{xy}^z)_\mu\}$ generally do not satisfy the following "completeness" conditions:

$$
\sum_{i} \frac{\partial}{\partial \overline{\phi}_{x}^{i}} \frac{\partial}{\partial \overline{\phi}_{x}^{i}} = \frac{\partial}{\partial \overline{\phi}_{x}^{i}} \qquad \sum_{\mu} (\phi_{xy}^{z})_{\mu} \rightarrow \phi(\overline{\phi}_{xy}^{z})_{\mu} = \frac{\partial}{\partial \overline{\phi}_{x}^{i}} \qquad (41)
$$

Given the orthogonality relations [\(27\)](#page-13-2) and [\(33\)](#page-14-3), one can show that the above completeness conditions are equivalent to the condition that the bond dimension χ_x of the MPO \mathcal{O}_x is equal to dim(x) for all $x \in \text{Simp}(\mathcal{C})$, i.e.,

Eq. (41)
$$
\Leftrightarrow \quad \chi_x = \dim(x) \text{ for all } x \in \text{Simp}(\mathcal{C}).
$$
 (42)

The sketch of a proof goes as follows. First, we notice that the left-hand side of the first equality of Eq. (41) is an idempotent (as a linear map from left to right) due to the orthogonality relation [\(33\)](#page-14-3). The orthogonality further implies that this idempotent is bijective if and only if the bond dimension of \mathcal{O}_x agrees with the quantum dimension of x. Then, the statement follows from the fact that a bijective idempotent is the identity map. A similar argument shows that the second equality of Eq. [\(41\)](#page-16-0) is also satisfied if and only if $\chi_x = \dim(x)$ for all $x \in \text{Simp}(\mathcal{C})$.

Even though the completeness conditions do not hold in general, one can show the following weaker equalities:

$$
\sum_{i} \frac{\psi}{(\phi_x)_j} \frac{\psi}{(\overline{\phi}_x)_i (\phi_x)_i} = \frac{\psi}{(\phi_x)_j},
$$
\n
$$
\sum_{z,\mu} \frac{(\phi_{xy}^z)_{\mu} \longrightarrow \phi(\overline{\phi}_{xy}^z)_{\mu}}{(\phi_x)_i (\phi_y)_j} = \frac{\psi}{(\phi_x)_i (\phi_y)_j}
$$
\n(43)

where $\psi \in Aut(V)$ is an arbitrary unitary operator. The first equality for $\psi = id$ follows from the orthogonality relation [\(33\)](#page-14-3) for the action tensors. The case of a more general ψ will be shown in Appendix [F](#page-72-0) under a mild assumption on the duality structure. The second equality follows from the orthogonality relation (27) for the fusion tensors and the existence of the L-symbols (35) . Equation [\(43\)](#page-16-1) will be referred to as the weak completeness relation.

We note that an MPO representation that satisfies the completeness condition [\(41\)](#page-16-0) exists for any non-anomalous fusion category $\text{Rep}(H)$, where H is a semisimple Hopf algebra. The explicit construction of such MPOs is discussed in [\[36,](#page-85-8) [47\]](#page-86-2). For example, the MPO representation [\(31\)](#page-14-0) of Rep(G) symmetry satisfies the completeness condition because the bond Hilbert space V_{ρ} of an MPO \mathcal{O}_{ρ} is the representation space of $\rho \in \text{Rep}(G)$ and hence we have $\chi_{\rho} = \dim(V_{\rho}) = \dim(\rho)$. For a finite group symmetry Vec_G , the completeness condition (42) means that the symmetry operators are on-site. We do not assume the completeness condition in this paper.

III. FIBER FUNCTORS FROM INJECTIVE MPS

As we reviewed in Sec. [II B,](#page-6-0) SPT phases with fusion category symmetry $\mathcal C$ are classified by fiber functors of $\mathcal C$. In this section, we discuss how to extract these data from the ground state of a lattice system. The key idea is to exploit the correspondence between a TQFT partition function with defects and a generalized overlap of MPSs. In TQFT, a fiber functor $(F, J) : C \to \text{Vec}$ is represented diagrammatically as shown in Fig. [1,](#page-9-1) where $F(x)$ is given by the x-twisted sector on a circle and $J_{x,y}$ is given the transition amplitude on a pants diagram. Correspondingly, in the context of MPS, it is expected that $F(x)$ is given by a vector space spanned by the fixed points of a mixed transfer matrix, and $J_{x,y}$ is given by a triple inner product with three symmetry defects. In the following, we will explain this in detail.

A. Non-abelian factor system and fiber functor

1. Non-abelian factor system

Let us first define a non-abelian generalization of the factor system from an injective MPS with fusion category symmetry. This quantity turns out to be essentially the same as a fiber functor.

Let $\{\mathcal{O}_x \mid x \in \text{Simp}(\mathcal{C})\}$ be an MPO representation of a fusion category symmetry \mathcal{C} , and let A be a C-symmetric injective MPS tensor. The corresponding MPO matrix and MPS matrix are denoted by \mathcal{O}_x^{ij} and A^i , respectively. The bond dimensions of \mathcal{O}_x and A are given by χ_x and D. Since A is C-symmetric, it satisfies the symmetricity condition (32) , which can be rewritten as

$$
\sum_{j} O_{x}^{ij} \otimes A^{j} = \hat{\phi}_{x}^{-1} (1_{x} \otimes A^{i}) \hat{\phi}_{x} \Longleftrightarrow \overbrace{A}^{i} = \overbrace{\hat{\phi}_{x}^{-1} A}^{i} \overbrace{\hat{\phi}_{x}}^{i}.
$$
\n(44)

 $\overline{1}$

Here, 1_x denotes the $\dim(x) \times \dim(x)$ identity matrix, and the dotted line represents an auxiliary bond Hilbert space $F_x = \mathbb{C}^{\dim(x)}$.^{[14](#page-17-3)} On the right-hand side, the crossing of two lines represents the trivial braiding $v \otimes w \mapsto w \otimes v$, and the tensors $\hat{\phi}_x$ and $\hat{\phi}_x^{-1}$ are defined by

$$
\overrightarrow{v_i \bullet \cdots \bullet \bullet}_{\hat{\phi}_x} = \overrightarrow{(\phi_x)_i}, \qquad \overrightarrow{\bullet} \overrightarrow{\phi_x}^1 = \overrightarrow{(\phi_x)_i}, \qquad (45)
$$

¹⁴ When the bond dimension is $\chi_x = \dim(x)$, we can choose F_x to be the bond Hilbert space itself.

where $\{v_i \mid i = 1, 2, \cdots, \dim(x)\}\$ and $\{v_i^i \mid i = 1, 2, \cdots, \dim(x)\}\$ are dual bases of F_x and F_x^* . We note that $\hat{\phi}_x^{-1}$ is the right inverse of $\hat{\phi}_x$ due to the orthogonality relation [\(33\)](#page-14-3).^{[15](#page-18-0)} By abuse of terminology, we will refer to $\hat{\phi}_x$ as the action tensor. The action tensor $\hat{\phi}_x$ is uniquely determined up to the following gauge redundancy:

$$
\hat{\phi}_x \mapsto \hat{\phi}'_x = \frac{\overbrace{f_x}}{\hat{\phi}_x}, \qquad f_x \in GL_{\dim(x)}(\mathbb{C}). \tag{46}
$$

Indeed, if there are two action tensors $\hat{\phi}_x^1$ and $\hat{\phi}_x^2$ that satisfy Eq. [\(44\)](#page-17-4), their composition $\hat{\phi}_x^1(\hat{\phi}_x^2)^{-1}$ commutes with $1_x \otimes A^i$ for all i, and since A is injective, $\hat{\phi}_x^1(\hat{\phi}_x^2)^{-1}$ has to be of the form $f_x \otimes id_D$.

For a pair of two simple objects $x, y \in \text{Simp}(\mathcal{C})$, we can define the action tensor $\hat{\phi}_{x \otimes y}$ for the tensor product object $x \otimes y \in C$ as follows:

$$
\hat{\phi}_{x\otimes y} = \sum_{z,\mu} \underbrace{y \underbrace{\overbrace{\phi_{z}^{x}}^{x \cdots x}}_{\hat{\phi}_{z}} \underbrace{\overbrace{\phi_{xy}^{z}}^{x}}_{\hat{\phi}_{z}}, \qquad \hat{\phi}_{x\otimes y}^{-1} = \sum_{z,\mu} \underbrace{y \underbrace{\overbrace{\phi_{xy}^{x}}^{x}}_{\hat{\phi}_{z}^{-1}} \underbrace{\overbrace{\phi_{xy}^{z}}^{x \cdots x}}_{\hat{\phi}_{z}^{-1}}. \qquad (47)
$$

Here, $(\varphi_{xy}^z)_{\mu}$ and $(\overline{\varphi}_{xy}^z)_{\mu}$ are fusion and splitting tensors for the auxiliary bond Hilbert spaces, which satisfy the orthogonality relation analogous to Eq. [\(27\)](#page-13-2). A straightforward computation shows that the above action tensors satisfy Eq. [\(44\)](#page-17-4) where the MPO \mathcal{O}_x is replaced by $\mathcal{O}_{x\otimes y}$ defined by Eq. [\(26\)](#page-13-1). One can also define the action tensors for the tensor product of more than two simple objects. For $\text{example, for the tensor product of three simple objects } x, y, z \in \text{Simp}(\mathcal{C}), \text{ the action tensors } \hat{\phi}_{(x \otimes y) \otimes z}$ and $\hat{\phi}_{x\otimes(y\otimes z)}$ are given by

$$
\hat{\phi}_{(x\otimes y)\otimes z} = \sum_{u,w,\mu,\nu} \underbrace{z \xrightarrow{u \to u} w}_{\hat{\phi}_w} \underbrace{w \overleftrightarrow{b \nu}}_{\hat{\phi}_w}, \qquad \hat{\phi}_{x\otimes (y\otimes z)} = \sum_{v,w,\rho,\sigma} \underbrace{z \xrightarrow{\sigma} v}_{z \xrightarrow{\sigma} v} \underbrace{w \overleftrightarrow{b \rho}}_{\hat{\phi}_w}, \qquad (48)
$$

where the fusion and splitting tensors are abbreviated as μ , ν , ρ , and σ . The above action tensors may differ from each other by a gauge transformation $f_{x,y,z} \in GL_{dim}(x)$ dim(y) dim(z). Nevertheless, one can always choose φ 's so that $f_{x,y,z}$ becomes the identity because the MPO tensors for $(x\otimes y)\otimes z$ and $x \otimes (y \otimes z)$ are the same, i.e., $\mathcal{O}_{(x \otimes y) \otimes z} = \mathcal{O}_{x \otimes (y \otimes z)}$. More concretely, if we choose φ 's so that they have the same F-symbols as ϕ 's, the two action tensors in Eq. [\(48\)](#page-18-1) agree with each other.^{[16](#page-18-2)} In what follows, we will suppose $\hat{\phi}_{(x\otimes y)\otimes z} = \hat{\phi}_{x\otimes (y\otimes z)}$ without loss of generality.

Since the action tensors $\{\hat{\phi}_x \mid x \in \mathcal{C}\}$ have the gauge ambiguity (46) , $\hat{\phi}_{x \otimes y}$ generally differs from the multiplication of $\hat{\phi}_x$ and $\hat{\phi}_y$ by a gauge transformation. Namely, we have

¹⁵ The tensor $\hat{\phi}_x^{-1}$ is also the left inverse of $\hat{\phi}_x$ if and only if the completeness condition [\(41\)](#page-16-0) holds.

¹⁶ For instance, when $\chi_x = \dim(x)$ for all $x \in \text{Simp}(\mathcal{C})$, we can choose $(\varphi_{xy}^z)_{\mu} = (\varphi_{xy}^z)_{\mu}$.

The linear map $\Omega_{x,y}: F_x \otimes F_y \to F_x \otimes F_y$ will be referred to as a non-abelian factor system. To derive the consistency condition on $\Omega_{x,y}$, let us consider the multiplication of three action tensors $\hat{\phi}_x$, $\hat{\phi}_y$, and $\hat{\phi}_z$ in two different ways:

By comparing Eqs. [\(50\)](#page-19-1) and [\(51\)](#page-19-2), we find that $\Omega_{x,y}$ satisfies a non-abelian cocycle condition:

$$
\Omega_{x\otimes y,z}(\Omega_{x,y}\otimes 1_z)=\Omega_{x,y\otimes z}(1_x\otimes \Omega_{y,z})\Leftrightarrow \frac{\Omega_{x,y}}{\Omega_{x\otimes y,z}}
$$

Here, the diagrams in the above equation are read from left to right, and hence, e.g., the left-hand side represents $\Omega_{x\otimes y,z}(\Omega_{x,y}\otimes 1_z)$ rather than $(\Omega_{x,y}\otimes 1_z)\Omega_{x\otimes y,z}$. We note that Eq. [\(52\)](#page-19-3) agrees with the coherence condition (8) on a fiber functor, provided that the fusion category C is strictified, i.e., $\alpha_{x,y,z}^{\mathcal{C}} = id$. Thus, the isomorphisms $\{\Omega_{x,y} \mid x,y \in \mathcal{C}\}\)$ together with the set of vector spaces ${F_x \mid x \in C}$ can be identified with the data of a fiber functor $(F, \Omega) : C \to \text{Vec}$.

As we mentioned above, the action tensors $\{\hat{\phi}_x \mid x \in \text{Simp}(\mathcal{C})\}$ have a gauge redundancy. Under the gauge transformation [\(46\)](#page-18-3), the non-abelian factor system transforms as

$$
\Omega_{x,y} \mapsto \big(\bigoplus_{z \in x \otimes y} f_z\big)^{-1} \Omega_{x,y} (f_x \otimes f_y).
$$
\n(53)

We note that the gauge transformation of $\hat{\phi}_{x\otimes y}$ is restricted to the one induced by the gauge transformations for simple objects $z \in x \otimes y$. The non-abelian cocycle condition [\(52\)](#page-19-3) is invariant under this gauge transformation.

2. Mixed transfer matrix

In the previous subsection, we argued that the data of a fiber functor is encoded in a \mathcal{C} -symmetric injective MPS as a non-abelian factor system. In the following, we will discuss how to extract this data. To this end, we will use the mixed transfer matrix for a C -symmetric injective MPS and its fixed points.

Given an MPS tensor A , a transfer matrix T_A is defined by

$$
T_A := \sum_i A^{i*} \otimes A^i = \overbrace{A^{\dagger}}^{A^{\dagger}}, \tag{54}
$$

where A^{i*} denotes the complex conjugate of A^i . The left and right eigenvectors of T_A with the largest eigenvalues are called left and right fixed points. When A is injective, the left and right fixed points of T_A are unique and have the same positive eigenvalue $\lambda \in \mathbb{R}$ [\[33,](#page-85-6) [68,](#page-86-19) [70\]](#page-87-1). Without loss of generality, one can choose $\lambda = 1$ by normalizing the MPS tensor A. Thus, for an injective MPS, the left fixed point Λ_A^L and the right fixed point Λ_A^R are the unique (up to scalar) solutions to the following equation:

$$
\Lambda_A^L \bigodot = \Lambda_A^L \bigodot, \qquad \bigodot \Lambda_A^R = \bigodot \Lambda_A^R \ . \tag{55}
$$

Similarly, for a C-symmetric injective MPS tensor A, we define the mixed transfer matrix T_x as

$$
T_x = \sum_{ij} A^{i*} \otimes \mathcal{O}_x^{ij} \otimes A^j = \frac{A^{\dagger}}{\sum_{A} \phi_x} = \sum_i \underbrace{\overbrace{\bigoplus_{\substack{i=1 \ \phi_x \ni_i}}^{A^{\dagger}} A^{\dagger}}_{(\overline{\phi}_x)_i} \qquad (56)
$$

where the last equality follows from Eq. (32) . The fixed point equation for the mixed transfer matrix T_x is given by

$$
\Lambda_x^L \underbrace{\bullet} = \Lambda_x^L \underbrace{\bullet} , \underbrace{\bullet} \underbrace{\bullet} \right) \Lambda_x^R = \underbrace{\bullet} \Lambda_x^R \tag{57}
$$

Combined with the orthogonality relation [\(33\)](#page-14-3), the above equation implies that fixed points Λ_x^L and Λ_x^R have to satisfy

$$
\Lambda_x^L \underbrace{\bigoplus_{(\overline{\phi}_x)_i}}_{(\overline{\phi}_x)_i} = \Lambda_x^L \underbrace{\bigoplus_{(\overline{\phi}_x)_i}}_{(\phi_x)_i}, \underbrace{\bigoplus_{(\phi_x)_i}} \Lambda_x^R = \bigoplus_{(\phi_x)_i} \Lambda_x^R \tag{58}
$$

for all $i = 1, 2, \dots$, $\dim(x)$. This shows that $\Lambda_x^L(\overline{\phi}_x)_i$ and $(\phi_x)_i \Lambda_x^R$ are left and right fixed points of the transfer matrix T_A . Since the fixed point of T_A is unique, they are proportional to Λ_A^L and Λ_A^R , i.e.,

$$
\Lambda_x^L \underbrace{\bigoplus_{(\overline{\phi}_x)_i}}_{\mathfrak{p}} = a_i^L \Lambda_A^L, \qquad \qquad \underset{(\phi_x)_i}{\bigoplus} \Lambda_x^R = a_i^R \Lambda_A^R, \tag{59}
$$

where a_i^L and a_i^R are arbitrary complex numbers. Substituting this back into the fixed point equation [\(57\)](#page-20-0) leads us to

$$
\Lambda_x^L = \sum_i a_i^L \Lambda_x^{L(i)}, \qquad \Lambda_x^R = \sum_i a_i^R \Lambda_x^{R(i)}, \tag{60}
$$

where $\Lambda_x^{L(i)}$ and $\Lambda_x^{R(i)}$ are given by

$$
\Lambda_x^{L(i)} = \Lambda_A^L \underbrace{\left(\begin{matrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot
$$

Here, $\{v_i\}$ and $\{v^i\}$ are dual bases of the auxiliary bond Hilbert space $F_x = \mathbb{C}^{\dim(x)}$ and its dual F_x^* as in Eq. [\(45\)](#page-17-5). Equation [\(60\)](#page-20-1) shows that any fixed point of T_x is given by a linear combination of the fixed points in Eq. [\(61\)](#page-21-1).

In the following, we often write the left fixed point $\Lambda_x^{L(i)}$ as $|\Lambda_x^{L(i)}\rangle$, and define a vector space $F^{MPS}(x)$ assigned to each object $x \in \mathcal{C}$ as

$$
F^{\text{MPS}}(x) = \bigoplus_{i=1}^{\dim(x)} \mathbb{C} \left| \Lambda_x^{L(i)} \right\rangle. \tag{62}
$$

By definition, the dimension of $F^{MPS}(x)$ is dim(x). Similarly, the right fixed point $\Lambda_x^{R(i)}$ will be often denoted by $\langle \Lambda_x^{R(i)} |$. This notation is justified because $\{\Lambda_x^{R(i)}\}$ is the dual basis of $\{\Lambda_x^{L(i)}\}$ due to the orthogonality relation [\(33\)](#page-14-3).

3. Triple inner product

The assignment of the vector space [\(62\)](#page-21-2) defines a functor $F^{\text{MPS}}: \mathcal{C} \to \text{Vec}$, which turns out to be a fiber functor. To see this, let us consider the triple inner product of infinite MPSs [\[69\]](#page-87-0), which is defined by the following diagram:

$$
J_{x,y}^{\text{MPS}} := \hat{\mathcal{O}}_x |A\rangle \begin{matrix} x \otimes y \\ * \\ * \\ * \\ * \end{matrix} \hat{\mathcal{O}}_y |A\rangle : F^{\text{MPS}}(x) \otimes F^{\text{MPS}}(y) \to F^{\text{MPS}}(x \otimes y). \tag{63}
$$

The middle dot in the diagram represents the identity map on the bond Hilbert space $V_x \otimes V_y$. We emphasize that the triple inner product $J_{x,y}^{\text{MPS}}$ is a linear map rather than a single number: terminating each wing of the diagram computes its matrix element. Since each wing is infinitely long, the matrix element of $J_{x,y}^{\text{MPS}}$ vanishes unless the endpoint of each wing is a fixed point of the mixed transfer matrix.^{[17](#page-21-3)} Therefore, $J_{x,y}^{\text{MPS}}$ can be regarded as a linear map from $F^{\text{MPS}}(x) \otimes F^{\text{MPS}}(y)$ to $F^{\text{MPS}}(x\otimes y)$.

Let us explicitly compute the matrix element of $J_{x,y}^{\text{MPS}}$. To this end, we terminate the wings of $J_{x,y}^{\text{MPS}}$ by fixed points $|\Lambda_x^{L(i)}\rangle, |\Lambda_y^{L(j)}\rangle$, and $\langle \Lambda_{x\otimes y}^{R(k)}\rangle$ $\lim_{x \otimes y}$. Here, *i* runs from 1 to dim(*x*), *j* runs from 1

¹⁷ We recall that a fixed point of a (mixed) transfer matrix is an eigenvector with the largest eigenvalue $\lambda = 1$.

to dim(y), and k runs from 1 to dim(x) dim(y). Using the fixed point equation [\(57\)](#page-20-0) successively annihilates the MPO tensors in Eq. [\(63\)](#page-21-4). As a result, the matrix element of $J_{x,y}^{\text{MPS}}$ can be recast into

$$
(J_{x,y}^{\text{MPS}})_{(i,j),k} := \langle \Lambda_{x \otimes y}^{R(k)} | J_{x,y}^{\text{MPS}} | \Lambda_x^{L(i)} \rangle \otimes | \Lambda_y^{L(j)} \rangle = \sum_{\Lambda_x^{L(i)}} \left\{ \bigotimes_{\text{id}}^{R(k)} \Lambda_y^{L(j)} \right\}.
$$
 (64)

To evaluate the right-hand side, we suppose that A is left canonical, meaning that the left fixed point of T_A is the identity, i.e., $\Lambda_A^L = 1_D$. We also suppose that the right fixed point Λ_A^R is normalized so that $tr(\Lambda_A^R) = 1.18$ $tr(\Lambda_A^R) = 1.18$ In this case, by substituting Eq. [\(61\)](#page-21-1) into the right-hand side of Eq. [\(64\)](#page-22-2), we obtain

$$
(J_{x,y}^{\text{MPS}})_{(i,j),k} = \left(\underbrace{v_i \bullet \cdots \bullet}_{\hat{\phi}_x \quad \hat{\phi}_y \quad \hat{\phi}_{x \otimes y}} \right) \Lambda_A^R = \underbrace{v_i \bullet \cdots \bullet}_{v_j \bullet \cdots \bullet} v_y \quad . \tag{65}
$$

This shows that the matrix element of the triple inner product $J_{x,y}^{\text{MPS}}$ agrees with that of the nonabelian factor system $\Omega_{x,y}$. As discussed in Sec. [III A 1,](#page-17-2) $\Omega_{x,y}$ satisfies the coherence condition of a fiber functor. Thus, $J_{x,y}^{\text{MPS}}$ also satisfies the same coherence condition, meaning that $(F^{\text{MPS}}, J^{\text{MPS}})$: $C \rightarrow$ Vec is a fiber functor. It is now clear that one can also compute the L-symbols similarly: by inserting the fusion tensor $(\phi_{xy}^z)_{\mu}$ in the middle of the diagram, we obtain

$$
L_{(i,j),(z;k,\mu)}^{xy} = \frac{1}{D} \underbrace{\left(\overline{\phi}_{z}\right)_{k}}_{(\phi_{x})_{i} \left(\overline{\phi}_{xy}^{z}\right)_{\mu}}\right)_{(\phi_{y})_{j}} = \underbrace{\left(\sum_{\Lambda_{z}^{R(k)}}^{\Lambda_{z}^{R(k)}}\right)_{\Lambda_{y}^{L(j)}}}_{\Lambda_{x}^{L(i)} \left(\overline{\phi}_{xy}^{z}\right)_{\mu}}\right)_{\Lambda_{y}^{L(j)}} = \underbrace{v_{i} \bullet \cdots \bullet}_{v_{j} \bullet \cdots \bullet} \underbrace{\left(\overline{\phi}_{xy}^{z}\right)_{\mu}}_{\Lambda_{x,y}}.
$$
\n
$$
(66)
$$

We recall that $(\varphi_{xy}^z)_\mu$ on the right-hand side is an auxiliary fusion tensor, which has the same *F*-symbols as $(\phi_{xy}^z)_{\mu}$.

4. Torus partition functions

In the preceding subsections, we explained how to compute the fiber functor from the triple inner product of infinite MPSs. From the perspective of TQFT, the data of a fiber functor corresponds to the partition function on a pair of pants with defects as shown in Fig. [1.](#page-9-1) Based on the ideas explained in Sec. [I,](#page-2-0) it is also possible to evaluate more general partition functions in the framework of MPS. In this subsection, we evaluate the partition function on a torus as a specific example.

 18 This is equivalent to the normalization of the infinite MPS.

Let us consider the torus partition function in the presence of symmetry defects of the following configuration:

$$
Z\left[\begin{array}{c}\n(\phi_{xy}^z)_{\mu}x \\
\hline\ny\n\end{array}\right]\n\right]
$$
\n
$$
x\left(\overline{\phi_{yx}^z}\right)_{\nu}\n\left(\overline{\phi_{yx}^z}\right)
$$
\n(67)

As in the computation of a fiber functor, we glue MPSs along these symmetry lines to evaluate the above partition function:

Here, we put a conjugation ∗ on the left side of the defects. The complex number evaluated by the above diagram depends on the number of sites L_1 , L_2 , and L_3 in each interval. However, by taking the large volume limit where L_i goes to infinity for all i, this dependence disappears, and the diagram reduces to the TQFT partition function:

In fact, in this limit, each interval produces a projector to $F^{MPS}(x)$, and hence the right-hand side of the above equation can be computed as

$$
\lim_{L_i \to \infty} \frac{L_2}{\underbrace{\sum_{i=1}^{L_i} \sum_{j=1}^{L_2}}_{L_1 \cup \bigcup_{j=1}^{L_i} L_3}} = \sum_{ijk} L_{(i,j),(z;k,\mu)}^{xy} \overline{L}_{(z;k,\nu),(j,i)}^{yx}.
$$
\n(70)

This is nothing but the torus partition function of an SPT phase with fusion category symmetry [\[32,](#page-85-5) [54,](#page-86-6) [56\]](#page-86-11).

B. Abelianization of the non-abelian factor system

We have seen that the fiber functor or the non-abelian factor system can be used as an invariant that characterizes SPT phases. However, these are classified by non-abelian cohomology theories, and it may not be easy to compute the invariants and classifications in general. In this section, we will abelianize the non-abelian factor system to construct an invariant that takes values in (abelian) cohomology theories. Abelianization results in a loss of information, so these invariants do not provide a complete classification. Nevertheless, if these invariants take different values, it is clear that the two phases are different. Additionally, as it only involves ordinary cohomology theories, one can invoke various techniques for calculating cohomology groups. This makes it relatively easy to identify the structure of the classification.

Before we define the abelianization, let us first define a cohomology theory. For a non-anomalous fusion category \mathcal{C} , we consider the following cochain $C^k(\mathcal{C}; \mathbb{Z})$:

$$
C^k(\mathcal{C}; \mathbb{Z}) := \{ n : \text{Simp}(\mathcal{C})^k \to \mathbb{Z} \}. \tag{71}
$$

A cochain n is extended additively to non-simple objects as $n(\cdots, \rho_i \oplus \rho'_i, \cdots) = n(\cdots, \rho_i, \cdots) +$ $n(\cdots, \rho'_i, \cdots)$. Namely, a cochain n is a For this cochain, we introduce a differential

$$
\delta_k: C^k(\mathcal{C}; \mathbb{Z}) \to C^{k+1}(\mathcal{C}; \mathbb{Z})
$$
\n⁽⁷²⁾

defined by

$$
(\delta_k n)(\rho_1, ..., \rho_{k+1}) := \dim(\rho_1) n(\rho_2, ..., \rho_{k+1}) + \sum_{i=1}^k (-1)^i n(\rho_1, ..., \rho_i \otimes \rho_{i+1}, ..., \rho_{k+1})
$$

+ $(-1)^{k+1} n(\rho_1, ..., \rho_k) \dim(\rho_{k+1}),$ (73)

for $n \in C^k(\mathcal{C}; \mathbb{Z})$. We remark that $\dim(\rho)$ is an integer for all $\rho \in \mathcal{C}$ since \mathcal{C} is non-anomalous. We will drop the subscript k of δ_k when it is clear from the context. Then, we can show that

$$
\delta^2 := \delta_{k+1}\delta_k = 0 : C^k(\mathcal{C}; \mathbb{Z}) \to C^{k+2}(\mathcal{C}; \mathbb{Z}).
$$
\n(74)

See Appendix [G](#page-73-0) for a proof. Therefore, we can define the cohomology theory associated to the complex $(C^k(\mathcal{C}; \mathbb{Z}), \delta)$:

$$
H^{k}(\mathcal{C}, \mathbb{Z}) := \text{Ker}(\delta_{k})/\text{Im}(\delta_{k-1}).
$$
\n(75)

Despite the notation, $H^k(\mathcal{C}, \mathbb{Z})$ depends only on the fusion ring of \mathcal{C} . We note that $H^k(\mathcal{C}, \mathbb{Z})$ is a special case of the Hochschild cohomology.^{[19](#page-24-1)}

One can define an element of $\mathrm{H}^3(\mathcal{C}, \mathbb{Z})$ from the non-abelian factor system $\{\Omega_{x,y}\}\$: we take an R-lift $r_{x,y} \in \mathbb{R}$ of $\det(\Omega_{x,y})$, i.e.,

$$
r_{x,y} := \frac{1}{2\pi} \log(\det(\Omega_{x,y})).
$$
\n(76)

¹⁹ We are grateful to Yosuke Kubota for pointing this out. As a Hochschild cohomology, $H^k(\mathcal{C}, \mathbb{Z})$ should be written as $H^k(\mathrm{Gr}(\mathcal{C}), \mathbb{Z})$, where $\mathrm{Gr}(\mathcal{C})$ is the fusion ring of $\mathcal C$ and $\mathbb Z$ is equipped with a $\mathrm{Gr}(\mathcal{C})$ -bimodule structure defined by $x \cdot m = m \cdot x = \dim(x) m$ for all $x \in \mathcal{C}$ and $m \in \mathbb{Z}$.

$$
n_{x,y,z} := r_{y,z} - \sum_{w_i \in x \otimes y} r_{w_i,z} + \sum_{w_j \in y \otimes z} r_{x,w_j} - r_{x,y}.
$$
 (77)

Due to the non-abelian cocycle condition [\(52\)](#page-19-3), $n_{x,y,z}$ takes values in Z. Therefore, $\{n_{x,y,z}\}$ can be regarded as an element of $C^3(\mathcal{C}; \mathbb{Z})$. Moreover, we can check that $(\delta n)_{x,y,z,w} = 0$. However, under the shift of $r_{x,y}$

$$
r_{x,y} \mapsto r_{x,y} + m_{x,y}, \quad m_{x,y} \in \mathbb{Z}, \tag{78}
$$

the cochain $n_{x,y,z}$ changes by the coboundary of $m_{x,y}$:

$$
n_{x,y,z} \mapsto n_{x,y,z} + (\delta m)_{x,y,z}.\tag{79}
$$

Thus, ${n_{x,y,z}}$ can be regarded as an element of the cohomology:

$$
[n_{x,y,z}] \in \mathrm{H}^3(\mathcal{C}, \mathbb{Z}).\tag{80}
$$

In general, the abelianization $\Omega_{x,y} \mapsto [n_{x,y,z}]$ is neither injective nor surjective.

C. Example: $\mathbf{Rep}(G)$ symmetry

As a simple example, let us compute the L-symbols for the cluster state with $G \times \text{Rep}(G)$ symmetry. We start by recalling an MPS representation of the $G \times \text{Rep}(G)$ cluster state following [\[41\]](#page-85-10), see also [\[46\]](#page-86-1) for an earlier work on this state. The physical and bond Hilbert spaces of the $G \times \text{Rep}(G)$ cluster state are both given by the regular representation of G. In what follows, the basis of the physical Hilbert space is denoted by $\{g\}_{\rm phys} | g \in G\}$, while the basis of the bond Hilbert space is denoted by $\{|g\rangle | g \in G\}$. The MPS tensors on odd sites (black dots) and even sites (white dots) are given by

$$
\overrightarrow{A_o} = \sum_{g \in G} P_g \otimes |g\rangle_{\text{phys}}, \quad \overrightarrow{A_e} = \sum_{g \in G} L_g \otimes |g\rangle_{\text{phys}}, \tag{81}
$$

where P_g and L_g are the projection and the left multiplication acting on the bond Hilbert space, i.e.,

$$
P_g = |g\rangle\langle g| \,, \quad L_g = \sum_{h \in G} |gh\rangle\langle h| \,. \tag{82}
$$

We note that the unit cell of the cluster state consists of two sites. The cluster state given by the above MPS tensors has both G and $\text{Rep}(G)$ symmetries. The G symmetry is implemented by the right multiplication of G on odd sites, while the $\text{Rep}(G)$ symmetry is implemented by the action of Wilson line operators [\(31\)](#page-14-0) on even sites. More specifically, the actions of G and $\text{Rep}(G)$ symmetries are given by

$$
\mathcal{O}_g \cdot A_{\text{cluster}} = R_g^{\text{phys}} \cdot \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = R_g^{-1} \cdot \begin{bmatrix} 1 \\ R_g \\ R_g \end{bmatrix}, \quad \mathcal{O}_\rho \cdot A_{\text{cluster}} = \begin{bmatrix} 2_p^{\text{phys}} \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 2_p \\ 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 2_\rho \\ 1 \\ 0 \end{bmatrix}, \quad (83)
$$

where R_g and Z_ρ are defined by

$$
R_g = \sum_{h \in G} |hg^{-1}\rangle \langle h| \,, \quad Z_\rho = \sum_{g \in G} \rho(g) \otimes |g\rangle \langle g| \,. \tag{84}
$$

We note that $\rho(g)$ in Eq. [\(84\)](#page-26-0) is acting on the MPO's bond Hilbert space V_{ρ} , which is the representation space of ρ . The operators R_g^{phys} and Z_ρ^{phys} in Eq. [\(83\)](#page-25-1) are defined by the same equations except that the bras and kets now have the subscript "phys." Equation [\(83\)](#page-25-1) clearly shows that the cluster state defined above has $G \times \text{Rep}(G)$ symmetry. This cluster state is connected to the trivial product state by a $\text{Rep}(G)$ -symmetric finite depth quantum circuit, although it is not by a $G \times \text{Rep}(G)$ symmetric one [\[41\]](#page-85-10). As such, we say that the $G \times \text{Rep}(G)$ cluster state is in the trivial phase with Rep(G) symmetry.^{[20](#page-26-1)} We note that the cluster state [\(81\)](#page-25-2) is the ground state of the Rep(G) symmetric model discussed in [\[47,](#page-86-2) Section 4.4], see Appendix [E](#page-71-0) for more details.

Now, let us compute the L-symbols. For simplicity, we will focus on the L-symbols associated with Rep(G) symmetry. Computing the L-symbols for the full $G \times \text{Rep}(G)$ symmetry is also straightforward. The MPO tensor \mathcal{O}_{ρ} defined in Eq. [\(83\)](#page-25-1) implies that the action tensors for Rep(G) symmetry are given by

$$
(\phi_{\rho})_i = \frac{\begin{array}{c}\n v_i^{\rho} \\
 \bullet \\
 \hline\n Z_{\rho}\n \end{array}}{(z_{\rho})_i} = \frac{\begin{array}{c}\n v_{\rho}^i \\
 \bullet \\
 \hline\n Z_{\rho}^{-1}\n \end{array}}{(z_{\rho})_i} \tag{85}
$$

where $\{v_i^{\rho}$ $\binom{\rho}{i}$ | $i = 1, 2, \cdots, \dim(\rho)$ } and $\{v^i_\rho = (v^{\rho}_i)\}$ $\binom{\rho}{i}$ [†] | *i* = 1, 2, · · · , dim(ρ)} are dual bases of V_{ρ} and its dual \dot{V}_{ρ}^* . By using these action tensors, we can compute the L-symbols [\(37\)](#page-15-3) as follows:

$$
L_{(i,j),(p_3;k,\mu)}^{\rho_1 \rho_2} = \frac{1}{|G|} \sum_{g \in G} \sum_{i',j',k'} \rho_1(g)_{ii'}^{\dagger} \rho_2(g)_{jj'}^{\dagger} ((\phi_{\rho_1 \rho_2})_{\mu})_{i'j'}^{\kappa'} \rho_3(g)_{k'k},
$$

$$
\overline{L}_{(\rho_3;k,\mu),(i,j)}^{\rho_1 \rho_2} = \frac{1}{|G|} \sum_{g \in G} \sum_{i',j',k'} \rho_3(g)_{kk'}^{\dagger} ((\overline{\phi}_{\rho_1 \rho_2})_{\mu})_{i'j'}^{\kappa'} \rho_1(g)_{i'i} \rho_2(g)_{j'j}.
$$
 (86)

Here, $\rho(g)$ denotes the unitary representation matrix whose (i, j) -component $\rho(g)_{ij}$ is defined by

$$
\rho(g)v_i^{\rho} = \sum_j \rho(g)_{ij} v_j^{\rho}.
$$
\n(87)

Similarly, $((\phi_{\rho_1\rho_2}^{\rho_3})_{\mu})_{ij}^k$ in Eq. [\(86\)](#page-26-2) denotes the matrix element of the fusion tensor $(\phi_{\rho_1\rho_2}^{\rho_3})_{\mu}: V_{\rho_1} \otimes$ $V_{\rho_2} \to V_{\rho_3}$, namely,

$$
(\phi^{\rho_3}_{\rho_1 \rho_2})_{\mu} (v^{\rho_1}_i \otimes v^{\rho_2}_j) = \sum_k ((\phi^{\rho_3}_{\rho_1 \rho_2})_{\mu})_{ij}^k v^{\rho_3}_k. \tag{88}
$$

The corresponding splitting tensor is given by $\left(\overline{\phi}_{\rho_1}^{\rho_3}\right)$ $^{(\rho_3)}_{(\rho_1\rho_2)}|_{\mu} = (\phi_{\rho_1,\rho_2}^{\rho_3})_{\mu}^{\dagger}$. By using the identities^{[21](#page-26-3)}

$$
\rho_1(g)_{ii'} \rho_2(g)_{jj'} = \sum_{\rho_3,\mu} \sum_{k,k'} ((\phi_{\rho_1\rho_2}^{\rho_3})_{\mu})_{ij}^k ((\overline{\phi}_{\rho_1\rho_2}^{\rho_3})_{\mu})_{i'j'}^k \rho_3(g)_{kk'},
$$

$$
\sum_{g \in G} \rho(g)_{ii'}^{\dagger} \lambda(g)_{jj'} = \delta_{\rho,\lambda} \delta_{ij'} \delta_{i'j} |G| / \dim(\rho) \quad \text{for } \rho, \lambda \in \text{Simp}(\text{Rep}(G)),
$$
 (89)

 20 This is just a matter of terminology. In general, defining the notion of a trivial phase with non-invertible symmetry is a subtle problem due to the lack of group structure on the set of SPT phases [\[32,](#page-85-5) [41\]](#page-85-10).

 21 The first equality of eq. [\(89\)](#page-26-4) follows from the fusion rules. The second equality is known as the Schur orthogonality relation for irreducible representations $\rho, \lambda \in \text{Rep}(G)$.

one can rewrite the L-symbols in eq. (86) as

$$
L_{(i,j),(\rho_3;k,\mu)}^{\rho_1\rho_2} = ((\phi_{\rho_1\rho_2}^{\rho_3})_{\mu})_{ij}^k, \quad \overline{L}_{(\rho_3;k,\mu),(i,j)}^{\rho_1\rho_2} = ((\overline{\phi}_{\rho_1\rho_2}^{\rho_3})_{\mu})_{ij}^k. \tag{90}
$$

The above L-symbols correspond to the forgetful functor of $\text{Rep}(G)$.

We note that the L-symbols in eq. (90) are the same as those for the trivial product state with $Rep(G)$ symmetry. This is consistent with the fact the $G \times Rep(G)$ cluster state is in the trivial $Rep(G)$ -symmetric phase.

IV. INTERFACE MODES OF SPT PHASES WITH FUSION CATEGORY SYMMETRIES

In this section, we study interfaces of $1+1d$ SPT phases with fusion category symmetry C. We first derive the symmetry algebra, which we call the interface algebra, acting on the interface of two SPT phases with the same symmetry. The irreducible representations of this algebra can be regarded as localized modes at the interface, i.e. interface modes.^{[22](#page-27-3)} It turns out that the interface algebra does not have one-dimensional representations if the adjacent SPT phases are different. This implies that two different SPT phases must have degenerate interface modes that form a higher dimensional representation of the interface algebra. In other words, the symmetry at the interface of different SPT phases is anomalous. On the other hand, the interface algebra between the same SPT phase has one-dimensional representations, which describe non-degenerate interface modes. These one-dimensional representations are in one-to-one correspondence with automorphisms of a fiber functor F and form a group $Aut^{\otimes}(F)$. As an illustrative example, we will study the interface algebras of SPT phases with $\text{Rep}(G)$ symmetry in detail. Our analysis in this section is similar to that in $[71]$, which studies domain wall excitations in $1+1d$ gapped phases with anomalous finite group symmetries.

A. Symmetry algebra at the interface

Let us consider the interface between two SPT phases SPT_1 and SPT_2 with fusion category symmetry C . The ground states of these SPT phases are represented by C -symmetric injective MPSs A_1 and A_2 . The physical and bond Hilbert spaces of A_i are denoted by \mathcal{H}_i and V_i respectively. Here, \mathcal{H}_1 and V_1 are not necessarily isomorphic to \mathcal{H}_2 and V_2 . The action tensors associated with A_1 and A_2 are written as $\{(\phi_x^1)_i \mid i = 1, 2, \cdots, \dim(x)\}\$ and $\{(\phi_x^2)_i \mid i = 1, 2, \cdots, \dim(x)\}\$. Similarly, the L-symbols for A_1 and A_2 are $\{(L_1)_{(i,\tau)}^{xy}$ $\{ (i,j),(z;k,\mu) \}$ and $\{ (L_2)_{(i,j)}^{xy} \}$ $\{ (i,j),(z;k,\mu) \}$.

To derive the symmetry algebra at the interface, we consider the symmetry action on a periodic MPS in the presence of interfaces. Specifically, we consider the following MPS on a periodic chain

$$
|\psi_{12}, \psi_{21}\rangle = \frac{1}{A_1 A_1} \frac{\psi_{12}}{A_2 A_2} \frac{1}{A_1 A_1} \psi_{21} \frac{1}{A_1 A_1}, \qquad (91)
$$

where a half of the chain is occupied by SPT_1 and the other half is occupied by SPT_2 . The interface degrees of freedom in Eq. [\(91\)](#page-27-4) are represented by arbitrary two-leg tensors $\psi_{12}: V_2 \to V_1$ and

²² This is a similar idea to the correspondence between the irreducible representations of the tube algebra and the anyons in a topologically ordered system.

 $\psi_{21}: V_1 \to V_2$. The action of a symmetry MPO $\widehat{\mathcal{O}}_x$ on MPS [\(91\)](#page-27-4) is computed as

$$
\widehat{\mathcal{O}}_x |\psi_{12}, \psi_{21}\rangle = \frac{\left\{\n\begin{matrix}\n\mathcal{O}_x & \mathcal{O}_x & \mathcal{O}_x \\
\psi_{12} & \psi_{21} & \psi_{21} \\
\vdots & \ddots & \mathcal{O}_x\n\end{matrix}\n\right\}}{A_1 A_1} = \sum_{1 \leq i, j \leq \dim(x)} \left| (\widehat{\mathcal{O}}_x^{12})_{ij} \psi_{12}, (\widehat{\mathcal{O}}_x^{21})_{ji} \psi_{21} \right\rangle, \tag{92}
$$

where the operators $(\widehat{\mathcal{O}}_x^{12})_{ij}$ and $(\widehat{\mathcal{O}}_x^{21})_{ji}$ are defined by

$$
(\widehat{\mathcal{O}}_x^{12})_{ij}\psi_{12} = \frac{\psi_{12}}{(\phi_x^1)_i} \frac{\psi_{12}}{(\overline{\phi}_x^2)_j}, \qquad (\widehat{\mathcal{O}}_x^{21})_{ji}\psi_{21} = \frac{\psi_{21}}{(\phi_x^2)_j} \frac{\psi_{21}}{(\overline{\phi}_x^1)_i}.
$$
 (93)

These operators can be regarded as symmetry operators acting on the interfaces. The symmetry algebra A_{12} acting on the left interface is spanned by operators $\{(\hat{\mathcal{O}}_x^{12})_{ij} \mid x \in \text{Simp}(\mathcal{C}), i, j = 1\}$ $(1, 2, \dots, \text{dim}(x))$. Similarly, the symmetry algebra \mathcal{A}_{21} acting on the right interface is spanned by operators $\{(\widehat{\mathcal{O}}_x^{21})_{ij} \mid x \in \text{Simp}(\mathcal{C}), i,j = 1,2,\cdots, \text{dim}(x)\}\)$. In what follows, we will focus on \mathcal{A}_{12} without loss of generality. The multiplication of symmetry operators in \mathcal{A}_{12} can be computed explicitly as

$$
(\widehat{O}_{x}^{12})_{ii'} (\widehat{O}_{y}^{12})_{jj'} \psi_{12} = \frac{\overbrace{\psi_{12}}^{\psi_{12}}}{(\phi_{x}^{1})_{i} (\phi_{y}^{1})_{j}} \frac{\overbrace{\psi_{12}}^{\psi_{12}}}{(\overline{\phi}_{y}^{2})_{j'} (\overline{\phi}_{x}^{2})_{i'}} = \sum_{z,\mu} \frac{\overbrace{\psi_{12}}^{\psi_{12}}}{(\phi_{x}^{1})_{i} (\phi_{y}^{1})_{j}} \frac{\overbrace{\psi_{12}}^{\psi_{12}}}{(\phi_{x}^{1})_{i} (\phi_{y}^{1})_{j}} \frac{\overbrace{\psi_{12}}^{\psi_{12}}}{(\overline{\phi}_{x}^{2})_{j'} (\overline{\phi}_{x}^{2})_{i'}}
$$
\n
$$
= \sum_{z,\mu} \sum_{k,k'} \frac{\overbrace{\psi_{12}}^{\psi_{12}}}{(\phi_{x}^{1})_{i} (\phi_{y}^{1})_{j} (\overline{\phi}_{z}^{1})_{k}} \frac{\overbrace{\psi_{12}}^{\psi_{12}}}{(\overline{\phi}_{z}^{2})_{k'} (\overline{\phi}_{z}^{2})_{k'} (\overline{\phi}_{y}^{2})_{j'} (\overline{\phi}_{x}^{2})_{i'}}
$$
\n
$$
= \sum_{z,\mu} \sum_{k,k'} (L_{1})_{(i,j),(z;k,\mu)}^{xy} (\overline{L_{2}})_{(z;k',\mu),(i',j')}^{xy} (\widehat{O}_{x}^{12})_{kk'} \psi_{12}.
$$
\n(94)

The second and third equalities follow from the weak completeness relation $(43).^{23}$ $(43).^{23}$ $(43).^{23}$ $(43).^{23}$ The above multiplication law is associative due to the consistency condition on the L-symbols. The unit element of the algebra \mathcal{A}_{12} is the identity operator $\widehat{\mathcal{O}}_1^{12} : \psi_{12} \mapsto \psi_{12}$. Here and in what follows, we omit the indices i and j of an operator $(\widehat{\mathcal{O}}_x^{12})_{ij}$ when $\dim(x) = 1$ because the indices are unique in this case.

Summarizing, we find that the symmetry algebra acting on the interface of \mathcal{C} -symmetric injective MPSs $|A_1\rangle$ and $|A_2\rangle$ is given by

$$
A_{12} = \text{Span}\{(\hat{O}_x^{12})_{ij} \mid x \in \text{Simp}(\mathcal{C}), \ i, j = 1, 2, \cdots, \text{dim}(x)\}\
$$
(95)

that is equipped with the following associative multiplication:

$$
(\widehat{\mathcal{O}}_x^{12})_{ii'} (\widehat{\mathcal{O}}_y^{12})_{jj'} = \sum_{z,\mu} \sum_{k,k'} (L_1)_{(i,j),(z;k,\mu)}^{xy} (\overline{L_2})_{(z;k',\mu),(i',j')}^{xy} (\widehat{\mathcal{O}}_z^{12})_{kk'}.
$$
\n(96)

²³ The last line of Eq. (94) can also be obtained directly from the first line by using the orthogonality relation (27) and the L -move (35) (36) .

We emphasize that the symmetry algebra A_{12} depends only on the L-symbols (or fiber functors), which characterize the SPT phases represented by C-symmetric injective MPSs. In particular, A_{12} does not depend on microscopic details of MPS tensors.

We note that each symmetry operator \mathcal{O}_x acting on the bulk SPT phase is factorized into multiple operators $\{(\hat{O}_x^{12})_{ij} | i, j = 1, 2, \cdots, \dim(x)\}\$ at the interface. Accordingly, the dimension of the symmetry algebra A_{12} is not equal to the number of simple objects of C but coincide with the total dimension of \mathcal{C} , i.e.,

$$
\dim(\mathcal{A}_{12}) = \sum_{x \in \text{Simp}(\mathcal{C})} \dim(x)^2 = \dim(\mathcal{C}).\tag{97}
$$

The interface algebra \mathcal{A}_{12} is a special case of the algebra $\mathcal{A}_{\mathcal{M},\mathcal{N}}$ studied in [\[72\]](#page-87-3), where M and N are general C-module categories. Specifically, $\mathcal{A}_{M,N}$ reduces to \mathcal{A}_{12} when both M and N are module categories associated with fiber functors. When M and N are the same, the algebra $\mathcal{A}_{M,N}$ has the structure of a weak Hopf algebra [\[72\]](#page-87-3), whose representation category is equivalent to the dual fusion category Func(M, M), the category of C-module endofunctors of M [\[72,](#page-87-3) [73\]](#page-87-4). See also [\[74](#page-87-5)[–79\]](#page-87-6) for recent applications of this algebra.

Example. As the simplest example, let us consider the interface algebra of SPT phases with ordinary finite group symmetry G. When the symmetry $C = \text{Vec}_G$ is a finite group, the L-symbols are nothing but group 2-cocycles $\omega \in \mathbb{Z}_{\text{gp}}^2(G, \mathbb{U}(1))$. More specifically, the relation between Lsymbols and 2-cocycles is given by

$$
L^{g,h} = \omega(g,h),\tag{98}
$$

where the indices of the L-symbols are omitted because they are unique. The symmetry algebra \mathcal{A}_{12} at the interface of two SPT phases SPT_{ω_1} and SPT_{ω_2} is spanned by operators $\{\hat{\mathcal{O}}_g^{12} \mid g \in G\}$, which obey the following multiplication law:

$$
\widehat{\mathcal{O}}_g^{12} \widehat{\mathcal{O}}_h^{12} = \omega_1(g, h)\omega_2(g, h)^{-1} \widehat{\mathcal{O}}_{gh}^{12}.
$$
\n(99)

Namely, A_{12} is isomorphic to the twisted group algebra $\mathbb{C}[G]^{\omega_1\omega_2^{-1}}$ with the twist $\omega_1\omega_2^{-1} \in$ $Z_{\text{gp}}^2(G, U(1))$. In particular, if ω_1 and ω_2 belong to different cohomology classes $[\omega_1] \neq [\omega_2] \in$ $\text{H}_{\text{gp}}^2(G, U(1)),$ the symmetry G is realized projectively at the interface. This phenomenon is known as an anomaly inflow [\[80\]](#page-87-7).

As a more interesting example, we will study the interface algebras for the non-invertible $\text{Rep}(D_8)$ symmetry in Sec. [IV C 2.](#page-32-0) The interface algebras for the other non-anomalous $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami symmetries $\text{Rep}(Q_8)$ and $\text{Rep}(H_8)$ will also be discussed in Appendix [D.](#page-61-2)

B. Interface modes

In this subsection, we show that the interface algebra A_{12} has one-dimensional representations if and only if SPT_1 and SPT_2 are the same SPT phase. Consequently, different SPT phases must have degenerate interface modes between them. In other words, the interface algebra between two different SPT phases is anomalous. This can be regarded as an anomaly inflow for fusion category symmetry.

We first show that A_{12} has a one-dimensional representation if SPT_1 and SPT_2 are the same phase. When SPT_1 and SPT_2 are the same, the associated L-symbols L_1 and L_2 are equivalent to each other. Therefore, there exists a set of invertible matrices $\{U_x \mid x \in \text{Simp}(\mathcal{C})\}\$ such that

$$
(L_1)^{xy}_{(i,j),(z;k,\mu)} = \sum_{i',j',k'} (U_x)_{ii'} (U_y)_{jj'} (U_z)^{-1}_{k'k} (L_2)^{xy}_{(i',j'),(z;k',\mu)}.
$$
\n(100)

Since U_x, U_y , and U_z are invertible, one can rewrite the above equation as

$$
(U_x)_{ii'}(U_y)_{jj'} = \sum_{z,\mu} \sum_{k,k'} (L_1)_{(i,j),(z;k,\mu)}^{xy} (\overline{L_2})_{(z;k',\mu),(i',j')}^{xy}(U_z)_{kk'}.
$$
\n(101)

Comparing Eq. [\(101\)](#page-30-0) with Eq. [\(96\)](#page-28-2), we find that $(\widehat{\mathcal{O}}_x^{12})_{ij} = (U_x)_{ij}$ defines a one-dimensional representation of A_{12} . Here, a one-dimensional representation of an algebra A is a representation such that all elements of A are represented by complex numbers.

Conversely, if the interface algebra A_{12} has a one-dimensional representation, one can show that SPT_1 and SPT_2 are the same SPT phase. To see this, we first show that if \mathcal{A}_{12} has a (non-zero) onedimensional representation, so does \mathcal{A}_{21} . To this end, we recall that a fusion category C is unitary as we assume throughout the paper. Since $\mathcal C$ is unitary, there exists a basis such that the L-symbols L_1 and L_2 are unitary matrices.^{[24](#page-30-1)} In this basis, given a one-dimensional representation $(U_x^{12})_{ij}$ of \mathcal{A}_{12} , we can obtain a one-dimensional representation $(U_x^{21})_{ij}$ of \mathcal{A}_{21} by taking the Hermitian conjugate of U_x^{12} , i.e.,

$$
(U_x^{21})_{ij} := (U_x^{12})_{ij}^\dagger = (U_x^{12})_{ji}^*.
$$
\n(102)

Indeed, it immediately follows from the unitarity of the L-symbols that $(U_x)_{ij}^{21}$ defined above is a one-dimesional representation of A_{21} , i.e., it satisfies

$$
(U_x^{21})_{ii'} (U_y^{21})_{jj'} = \sum_{z,\mu} \sum_{k,k'} (L_2)^{xy}_{(i,j),(z;k,\mu)} (\overline{L_1})^{xy}_{(z;k',\mu),(i',j')} (U_z^{21})_{kk'},
$$
\n(103)

as long as $(U_x)_{ij}^{12}$ is a one-dimensional representation of \mathcal{A}_{12} :

$$
(U_x^{12})_{ii'} (U_y^{12})_{jj'} = \sum_{z,\mu} \sum_{k,k'} (L_1)^{xy}_{(i,j),(z;k,\mu)} (\overline{L_2})^{xy}_{(z;k',\mu),(i',j')} (U_z^{12})_{kk'}.
$$
 (104)

Now, we notice that the product $(U_x^{12}U_x^{21})_{ij} := \sum_k (U_x^{12})_{ik} (U_x^{21})_{kj}$ defines a (non-zero) onedimensional representation of the self-interface algebra A_{11} of SPT_1 . Similarly, the product $(U_x^{21}U_x^{12})_{ij} := \sum_k (U_x^{21})_{ik} (U_x^{12})_{kj}$ defines a (non-zero) one-dimensional representation of the selfinterface algebra A_{22} of SPT_2 . The *n*th power $(U_x^{12}U_x^{21})^n$ is also a (non-zero) one-dimensional representation of A_{11} for any positive integer n. Since A_{11} has only finitely many one-dimensional representations, there exists a positive integer n such that the one-dimensional representation given by $(\widehat{O}_x^{11})_{ij} = (U_x^{12}U_x^{21})_{ij}^n$ is isomorphic to the obvious one-dimensional representation $(\widehat{O}_x^{11})_{ij} = \delta_{ij}$. In particular, the determinant of $U_x^{12}U_x^{21}$ is non-zero, which implies that U_x^{12} and U_x^{21} are invertible. Thus, Eq. [\(104\)](#page-30-2) gives an equivalence between the L-symbols L_1 and L_2 , meaning that SPT_1 and SPT_2 are the same SPT phase.

 24 We recall that the L-symbols are matrix elements of the structure isomorphism J of a fiber functor, which is unitary.

In summary, we find that SPT_1 and SPT_2 are the same SPT phase if and only if the interface algebra A_{12} between them has a one-dimensional representation. In particular, the interface algebra between different SPT phases does not have one-dimensional representations. This implies that interface degrees of freedom between different SPT phases always form a higher dimensional representation of A_{12} , which leads to degenerate ground states on a periodic chain in the presence of interfaces. Put differently, the symmetry algebra at the interface of different SPT phases is anoma-lous.^{[25](#page-31-2)} This result generalizes the anomaly inflow on the boundaries of SPT phases with ordinary group symmetries.

From the point of view of symmetry topological field theory, representations of the interface algebra A_{12} (or more generally, $A_{M,N}$) can be understood as boundary excitations of 2+1d topological orders [\[72\]](#page-87-3). We will discuss this perspective in more detail in Sec. [V.](#page-33-0)

Non-degenerate interface modes. One-dimensional representations of the interface algebra \mathcal{A}_{12} describe non-degenerate interface modes between the same SPT phase $SPT_1 = SPT_2$. It is clear from the above discussion that one-dimensional representations of the interface algebra \mathcal{A}_{12} are in one-to-one correspondence with gauge transformations between L-symbols L_1 and L_2 . As we discussed at the end of Sec. IID, gauge transformations between L_1 and L_2 are also in one-to-one correspondence with isomorphisms between the corresponding fiber functors F_1 and F_2 . Therefore, one-dimensional representations of A_{12} are in one-to-one correspondence with isomorphisms of fiber functors F_1 and F_2 . Specifically, given a one-dimensional representation $\{(\widehat{O}_x^{12})_{ij} = (U_x)_{ij}\}$ of \mathcal{A}_{12} , the matrix elements of the corresponding isomorphism $\eta : F_1 \Rightarrow F_2$ are given by

$$
(\eta_x)_{ij} = (U_x)_{ij}.\tag{105}
$$

When the L-symbols L_1 and L_2 are equal to each other, one-dimensional representations of $A_{12} =: A$ can be canonically identified with automorphisms of the corresponding fiber functor F^{26} F^{26} F^{26} These one-dimensional representations form a group under the composition of the interface modes as follows:

$$
(\widehat{\mathcal{O}}_x)_{ij}\psi = (U_x)_{ij}\psi, \quad (\widehat{\mathcal{O}}_x)_{ij}\psi' = (U'_x)_{ij}\psi'
$$

\n
$$
\Rightarrow (\widehat{\mathcal{O}}_x)_{ij}(\psi \circ \psi') = \sum_k (U_x)_{ik}(U'_x)_{kj}\psi \circ \psi' = (U_x U'_x)_{ij}\psi \circ \psi'. \tag{106}
$$

The unit element of this group is given by the obvious one-dimensional representation $\{(\mathcal{O}_x)_{ij} = \delta_{ij}\}.$ Equation [\(105\)](#page-31-4) tells us that the group of one-dimensional representations of A is isomorphic to Aut[⊗] (F) .

C. Examples

1. Non-degenerate interfaces of $Rep(G)$ SPT phases

Let us consider representations of the interface algebra A between two copies of the $G \times \text{Rep}(G)$ cluster state. As in Sec. [III C,](#page-25-0) we will focus on the $Rep(G)$ symmetry of the cluster state rather than the full $G \times \text{Rep}(G)$ symmetry.

²⁵ Symmetry is called anomalous if it does not admit a gapped phase with a unique ground state. This definition of an anomaly was proposed in $1+1d$ in $[32]$ and can be employed in any spacetime dimension, including $0+1d$ as in the current situation.

 26 General representations of A including higher dimensional ones can be identified with monoidal natural transformations of F [\[72,](#page-87-3) [73\]](#page-87-4). These describe local excitations within a single SPT phase.

The interface algebra A is spanned by the operators $\{(\hat{\mathcal{O}}_{\rho})_{ij}\}$ that are labeled by irreducible representations $\rho \in \text{Rep}(G)$ and indices $i, j = 1, 2, \dots$, $\dim(\rho)$. We note that the dimension of A is equal to the order of G. Since the L-symbols for the $Rep(G)$ symmetry of the cluster state are given by Eq. [\(90\)](#page-27-2), the multiplication of operators $\{(\mathcal{O}_{\rho})_{ij}\}$ can be written explicitly as

$$
(\widehat{\mathcal{O}}_{\rho_1})_{ii'} (\widehat{\mathcal{O}}_{\rho_2})_{jj'} = \sum_{\rho_3,\mu} \sum_{k,k'} ((\phi^{\rho_3}_{\rho_1 \rho_2})_{\mu})_{ij}^k ((\overline{\phi}^{\rho_3}_{\rho_1 \rho_2})_{\mu})_{i'j'}^{k'} (\widehat{\mathcal{O}}_{\rho_3})_{kk'}.
$$
\n(107)

From the above equation, we find that $(\widehat{\mathcal{O}}_{\rho})_{ij} = \rho(g)_{ij}$ gives a one-dimensional representation of A for any $g \in G$, cf. the first equality of eq. [\(89\)](#page-26-4). These |G| one-dimensional representations exhaust all the irreducible representations of A because a $|G|$ -dimensional semisimple algebra can have at most |G| irreducible representations. The product of one-dimensional representations $\{(\mathcal{O}_{\rho})_{ij} = \rho(g)_{ij}\}\$ and $\{(\widehat{\mathcal{O}}_{\rho})_{ij} = \rho(h)_{ij}\}$ is given by

$$
(\widehat{\mathcal{O}}_{\rho})_{ij} = \sum_{k} \rho(g)_{ik} \rho(h)_{kj} = \rho(gh)_{ij}.
$$
\n(108)

Thus, we find that the group of one-dimensional representations of A is isomorphic to G . This result can also be derived from the fact that the $G \times \text{Rep}(G)$ cluster state corresponds to the forgetful functor of Rep(G) as an SPT phase with $Rep(G)$ symmetry. Specifically, the discussions at the end of Sec. [IV B](#page-29-0) imply that the group of one-dimensional representations of A is isomorphic to the group of automorphisms of the forgetful functor. This group is isomorphic to G due to Tannaka-Krein duality.

2. Degenerate interfaces of $Rep(D_8)$ SPT phases

As another example, we consider interfaces of different SPT phases with $Rep(D_8)$ symmetry. Here, Rep(D_8) is a fusion category consisting of five simple objects $\{(a, b), D \mid a, b = 0, 1\}$, which obeys the following non-invertible fusion rules:

$$
(a, b) \otimes (c, d) = (a + c, b + d), \quad (a, b) \otimes D = D \otimes (a, b) = D, \quad D \otimes D = \bigoplus_{a, b = 0, 1} (a, b), \quad (109)
$$

where $a + c$ and $b + d$ are defined modulo 2. The Rep(D_8) symmetry admits three SPT phases [\[32\]](#page-85-5), which we denote by SPT_{ν_1} , SPT_{ν_2} , and SPT_{ν_3} . As we derive in Appendix [D,](#page-61-2) the interface algebra between any two of them has two two-dimensional irreducible representations. This means that there are two different interface modes, each of which is two-fold degenerate. The symmetry transformations of these interface modes are determined by the irreducible representations of the interface algebra. See Appendix [D](#page-61-2) for a detailed analysis of the interface algebras and their representations.

In this subsection, using the results in Appendix [D,](#page-61-2) we discuss the action of the $\text{Rep}(D_8)$ symmetry on a periodic chain consisting of two regions occupied by different SPT phases. For concreteness, we suppose that half of the chain is occupied by SPT_{ν_1} and the other half is occupied by SPT_{ν_2} . The other cases can also be studied similarly. We denote the left and right interfaces by $\mathcal{I}_{\nu_1,\nu_2}$ and $\mathcal{I}_{\nu_2,\nu_1}$ respectively. In the presence of these interfaces, the symmetry action on a periodic chain is given by Eq. (92) , that is,

$$
\widehat{\mathcal{O}}_x | \psi_L, \psi_R \rangle = \sum_{1 \le i, j \le \dim(x)} | (\widehat{\mathcal{O}}_x^{\mathcal{I}_{\nu_1, \nu_2}})_{ij} \psi_L, (\widehat{\mathcal{O}}_x^{\mathcal{I}_{\nu_2, \nu_1}})_{ji} \psi_R \rangle , \qquad (110)
$$

where ψ_L and ψ_R are the left and right interface modes. We suppose that ψ_L and ψ_R are in irreducible representations ρ_L and ρ_R of the interface algebras $\mathcal{A}_{\mathcal{I}_{\nu_1,\nu_2}}$ and $\mathcal{A}_{\mathcal{I}_{\nu_2,\nu_1}}$. Specifically, ρ_L and ρ_R are either of the two irreducible representations R_e and R_d listed in Table [V.](#page-67-0) Since ρ_L and ρ_R are two-dimensional, the interface modes ψ_L and ψ_R take values in \mathbb{C}^2 . We denote the Pauli operators acting on them by X, Y , and Z . Based on the data in Table [V,](#page-67-0) one can compute the symmetry action [\(110\)](#page-32-1) for all choices of (ρ_L, ρ_R) .

• When $(\rho_L, \rho_R) = (R_e, R_e)$ or (R_d, R_d) , the symmetry action [\(110\)](#page-32-1) reduces to

$$
\hat{\mathcal{O}}_{(0,0)} |\psi_L, \psi_R\rangle = \hat{\mathcal{O}}_{(1,1)} |\psi_L, \psi_R\rangle = |\psi_L, \psi_R\rangle, \n\hat{\mathcal{O}}_{(0,1)} |\psi_L, \psi_R\rangle = \hat{\mathcal{O}}_{(1,0)} |\psi_L, \psi_R\rangle = Z^L Z^R |\psi_L, \psi_R\rangle, \n\hat{\mathcal{O}}_D |\psi_L, \psi_R\rangle = \frac{1}{2} (S_+^L S_+^R + S_-^L S_-^R) |\psi_L, \psi_R\rangle, \qquad (111)
$$

where the operators with superscript L/R act on the left/right interface and $S_{+} := X \pm iY$. Remarkably, the local factor $Z^{L/R}$ of the invertible operators $\mathcal{O}_{(0,1)}$ and $\mathcal{O}_{(1,0)}$ anti-commutes with the local factors $S_{\pm}^{L/R}$ of the non-invertible operator \mathcal{O}_D . This anti-commutation relation of local factors was originally observed in [\[42\]](#page-85-16) and can be regarded as a manifestation of an anomaly at the interface. We note that the global symmetry operators still obey the correct fusion rules

$$
\widehat{\mathcal{O}}_{(a,b)}\widehat{\mathcal{O}}_{(c,d)} = \widehat{\mathcal{O}}_{(a+c,b+d)}, \quad \widehat{\mathcal{O}}_{(a,b)}\widehat{\mathcal{O}}_{D} = \widehat{\mathcal{O}}_{D}\widehat{\mathcal{O}}_{(a,b)} = \widehat{\mathcal{O}}_{D}, \quad \widehat{\mathcal{O}}_{D}\widehat{\mathcal{O}}_{D} = \sum_{a,b=0,1}\widehat{\mathcal{O}}_{(a,b)}.
$$
 (112)

In particular, $\mathcal{O}_{(a,b)}$ and \mathcal{O}_D commute with each other.

• When $(\rho_L, \rho_R) = (R_e, R_d)$ or (R_d, R_e) , the symmetry action [\(110\)](#page-32-1) reduces to

$$
\begin{aligned}\n\widehat{\mathcal{O}}_{(0,0)} \left| \psi_L, \psi_R \right\rangle &= -\widehat{\mathcal{O}}_{(1,1)} \left| \psi_L, \psi_R \right\rangle = \left| \psi_L, \psi_R \right\rangle, \\
\widehat{\mathcal{O}}_{(0,1)} \left| \psi_L, \psi_R \right\rangle &= -\widehat{\mathcal{O}}_{(1,0)} \left| \psi_L, \psi_R \right\rangle = Z^L Z^R \left| \psi_L, \psi_R \right\rangle, \\
\widehat{\mathcal{O}}_D \left| \psi_L, \psi_R \right\rangle &= 0.\n\end{aligned} \tag{113}
$$

This result is consistent with the fusion rules [\(112\)](#page-33-1).

V. PARAMETERIZED FAMILY AND NON-ABELIAN THOULESS PUMP

In this section, we study families of SPT states parameterized by a circle $S¹$. We first define an invariant of an S^1 -parameterized family of C-symmetric injective MPSs, where C is a fusion category. The invariant turns out to be an automorphism of a fiber functor $F : C \to \text{Vec}$. Physically, this invariant can be understood as a generalized Thouless pump. Indeed, as we will see later, an adiabatic evolution of a C-symmetric injective MPS along a non-trivial family gives rise to a pump of a non-degenerate interface mode.^{[27](#page-33-2)} As an illustrative example, we provide concrete lattice models of S^1 -parameterized families of SPT states with Rep(G) symmetry. In what follows, a family of SPT states parameterized by X will be sometimes abbreviated as an X -family.

 27 We recall that there is a one-to-one correspondence between automorphisms of a fiber functor and non-degenerate interface modes, see Sec. [IV B.](#page-29-0)

A. Invariants of one-parameter families of SPT phases

Let $\{A(\theta) | \theta \in [0, 2\pi]\}$ be a family of C-symmetric injective MPS tensors parameterized by an interval $[0, 2\pi]$. The MPS tensor $A(\theta)$ is supposed to be continuous in θ . In particular, the physical Hilbert space \mathcal{H} and the bond Hilbert space $V = \mathbb{C}^D$ do not depend on θ . We also suppose that the state $|A(\theta)\rangle$ associated with the MPS tensor $A(\theta)$ is 2π -periodic, i.e.,

$$
|A(0)\rangle = |A(2\pi)\rangle \,,\tag{114}
$$

which means that the parameter space is a circle S^1 . We emphasize that $A(\theta)$ is not necessarily 2π -periodic due to the gauge redundancy that we will discuss shortly. Since $A(\theta)$ is C-symmetric, there exists a set of action tensors $\{(\phi(\theta)_x)_i \mid x \in \text{Simp}(\mathcal{C}), i = 1, 2, \cdots, \text{dim}(x)\}\$ that satisfies Eq. [\(32\)](#page-14-2) for each $\theta \in [0, 2\pi]$, that is, we have

$$
\frac{1}{\mathcal{O}_x} \frac{\partial_x}{\partial \theta} = \sum_i \underbrace{\overbrace{\overbrace{\overline{\phi}(\theta)x}_i}_i \underbrace{\overbrace{\overbrace{\phi(\theta)x}_i}_i}_i \cdot \overbrace{\overbrace{\phi(\theta)x}_i}_i \tag{115}
$$

The L-symbols computed from the action tensors $\{(\phi(\theta)_x)_i\}$ are denoted by $\{L(\theta)_{(i,\theta)}^{xy}$ $_{(i,j),(z;k,\mu)}^{xy}$. Since $A(\theta)$ is continuous, one can take $\phi(\theta)$ and $L(\theta)$ to be continuous in $\theta \in [0, 2\pi]$ without loss of generality.

The MPS tensor $A(\theta)$ has a gauge ambiguity because the state $|A(\theta)\rangle$ is invariant under the following gauge transformation:

$$
A(\theta) \to \psi(\theta) A(\theta) = \frac{\psi(\theta)}{A(\theta)} \begin{cases} \psi(\theta)^{-1} \\ \psi(\theta)^{-1} \end{cases},
$$
\n(116)

where $\psi(\theta)$ is an arbitrary automorphism of V, i.e., a $D \times D$ invertible matrix. This gauge transformation induces the following transformation of $\{(\phi(\theta)_x)_i\}$ so that the symmetricity condition [\(115\)](#page-34-1) holds for any $\theta \in [0, 2\pi]$:

$$
(\phi(\theta)_x)_i \to \psi(\theta) (\phi(\theta)_x)_i = \frac{\psi(\theta)}{\begin{pmatrix} \psi(\theta) - 1 \\ (\phi(\theta)_x)_i \end{pmatrix}} , \qquad (\overline{\phi}(\theta)_x)_i \to \psi(\theta) (\overline{\phi}(\theta)_x)_i = \frac{\overline{\psi(\theta)} \psi(\theta)^{-1}}{(\overline{\phi}(\theta)_x)_i} \qquad (117)
$$

The action tensors themselves also have a gauge ambiguity because Eq. [\(115\)](#page-34-1) is invariant under the change of a basis

$$
(\phi(\theta)_x)_i \to {}^{U(\theta)}(\phi(\theta)_x)_i = \sum_j (U(\theta)_x)_{ij} (\phi(\theta)_x)_j,
$$
\n(118)

where $U(\theta)_x$ is a $\dim(x) \times \dim(x)$ invertible matrix. Accordingly, the L-symbols $L(\theta)_{(i,\tau)}^{xy}$ $_{(i,j),(z;k,\mu)}^{xy}$ are also transformed as in Eq. (39) , which we reproduce here:

$$
L(\theta)_{(i,j),(z;k,\mu)}^{xy} \to {}^{U(\theta)}L(\theta)_{(i,j),(z;k,\mu)}^{xy} = \sum_{i',j',k'} (U(\theta)_x)_{ii'} (U(\theta)_y)_{jj'} (U(\theta)_z)_{k'k}^{-1} L(\theta)_{(i',j'),(z;k',\mu)}^{xy}.
$$
 (119)

We note that the transformation [\(117\)](#page-34-2) does not affect the L-symbols because $\psi(\theta)$ and $\psi(\theta)^{-1}$ are canceled out in the definition [\(37\)](#page-15-3) of the L-symbols.

Now, let us define an invariant of an S^1 -parameterized family of SPT states with symmetry C. We provide three different ways to define the invariant for a given S^1 -parameterized family ${A(\theta), \phi(\theta), L(\theta)}.$

• Non-periodicity of $\phi(\theta)$. We first define the invariant of a one-parameter family using the non-periodicity of the action tensors $\phi(\theta)$ in a specific gauge. To this end, we first perform a gauge transformation $\psi(\theta) \in Aut(V)$ so that the MPS tensor $A(\theta)$ becomes 2π -periodic. Such a gauge transformation always exists because $A(0)$ and $A(2\pi)$ are gauge equivalent due to Eq. [\(114\)](#page-34-3). Furthermore, we choose the gauge transformation $U(\theta)$ so that the L-symbols $L(\theta)$ become constant in θ . This choice of a gauge is always possible because the equivalence class of $L(\theta)$ does not depend on θ ^{[28](#page-35-0)}. In the above gauge, the action tensors $\phi(\theta)$ are not necessarily 2π -periodic. Nevertheless, since we have $A(0) = A(2\pi)$, the non-periodicity of $\phi(\theta)$ is at most a gauge transformation

$$
(\phi(2\pi)_x)_i = \sum_j (\eta_x)_{ij} (\phi(0)_x)_j,
$$
\n(120)

where η_x is a dim(x) \times dim(x) invertible matrix. Moreover, since the L-symbols are constant in θ , the gauge transformation η preserves the L-symbols, meaning that η defines an automorphism of the corresponding fiber functor F , cf. equations [\(15\)](#page-9-5) and [\(17\)](#page-10-0). This automorphism η can be regarded as an invariant of an S^1 -parameterized family of injective MPSs.

• Non-periodicity of $A(\theta)$. We can also define an invariant of an S^1 -parameterized family by using the non-periodicity of the MPS tensor $A(\theta)$ in another gauge. To see this, we first choose a gauge so that $A(\theta)$ is 2π -periodic and $L(\theta)$ is constant as above. We then perform a gauge transformation $\psi(\theta) \in \text{Aut}(V)$ so that the action tensors $\phi(\theta)$ become 2π -periodic. A straightforward computation shows that the 2π -periodicity of the action tensor after the gauge transformation is equivalent \rm{to}^{29} \rm{to}^{29} \rm{to}^{29}

$$
\frac{\Psi}{(\phi(0)_x)_i} = \sum_j (\eta_x)_{ij} \xrightarrow{\Psi} \overbrace{(\phi(0)_x)_j}^{\mathbf{U}}, \qquad (121)
$$

where we defined $\Psi := \psi(0)^{-1}\psi(2\pi)$, and η_x is a dim(x) \times dim(x) invertible matrix defined in Eq. (120) . The above equation, together with the orthogonality relation (33) , implies that Ψ has to satisfy

$$
\boxed{\Psi \atop (\phi(0)_x)_i} = (\eta_x)_{ij} \Psi. \tag{122}
$$

That is, Ψ is a one-dimensional representation of the self-interface algebra. Conversely, if Ψ satisfies Eq. [\(122\)](#page-35-3), it also satisfies Eq. [\(121\)](#page-35-4) due to the weak completeness relation [\(43\)](#page-16-1). Therefore, the action tensors become 2π -periodic after the gauge transformation $\psi(\theta)$ if and

²⁸ In other words, we are in the same SPT phase throughout the parameter space $S¹$.

 29 The action tensors in Eq. [\(121\)](#page-35-4) are those before the gauge transformation.
only if Ψ satisfies Eq. [\(122\)](#page-35-0). We denote the solution to this equation by $\Psi = \psi_{\eta}$, which always exists and is unique up to scalar multiplication due to the one-to-one correspondence between one-dimensional representations of the self-interface algebra and automorphisms of a fiber functor. After the above gauge transformation, the MPS tensor $A(\theta)$ is no longer 2π -periodic in general, while the L-symbols remain constant in θ . The non-periodicity of $A(\theta)$ after the gauge transformation is given by $\psi'_{\eta} = \psi(0)\psi_{\eta}\psi(0)^{-1}$:

$$
A(2\pi) = \underbrace{\qquad \psi_{\eta}'}_{A(0)} \underbrace{\qquad (\psi_{\eta}')^{-1}}_{\text{O}} \tag{123}
$$

We note that ψ_{η} and ψ_{η}' are equivalent one-dimensional representations, both of which are associated with the same automorphism η . Any 2π -periodic gauge transformation cannot change this equivalence class because such a gauge transformation ψ' acts on ψ'_{η} by conjugation: $\psi'_\eta \mapsto \psi'(0)\psi'_\eta \psi'(0)^{-1}$. Thus, the equivalence class of the one-dimensional representation ψ_{η} can be regarded as an invariant of an S^1 -parameterized family of injective MPSs. The invariant defined here is related to the one defined by the non-periodicity of $\phi(\theta)$ (cf. Eq. [\(120\)](#page-35-1)) via the one-to-one correspondence between one-dimensional representations of the interface algebra and automorphisms of a fiber functor.

• Automorphism of $L(\theta)$. A yet another way to define the invariant of an S^1 -parameterized family is to look into $L(\theta)$ in a gauge such that both $A(\theta)$ and $\phi(\theta)$ are 2π -periodic. In such a gauge, the L-symbols are not necessarily constant in θ . More specifically, $L(\theta)$ is given by a gauge transformation of $L(0)$ as

$$
L(\theta)_{(i,j),(z;k,\mu)}^{xy} = \sum_{i',j',k'} (U(\theta)_x)_{ii'} (U(\theta)_y)_{jj'} (U(\theta)_z)_{k'k}^{-1} L(0)_{(i',j'),(z;k',\mu)}^{xy},
$$
(124)

where $U(\theta)_x$ is a dim(x) \times dim(x) invertible matrix. Since $L(\theta)$ is 2π -periodic due to the periodicity of $A(\theta)$ and $\phi(\theta)$, the gauge transformation $U(\theta)$ at $\theta = 2\pi$ defines an automorphism of a fiber functor corresponding to $L(0)$. In particular, at $\theta = 2\pi$, we have

$$
U(2\pi)_x = \eta_x,\tag{125}
$$

where η_x on the right-hand side is the same as the one in Eqs. [\(120\)](#page-35-1) and [\(122\)](#page-35-0). Thus, the automorphism η induced at $\theta = 2\pi$ can be identified with the invariant of an S^1 -parameterized family.

From the above discussions, we find that one can associate an automorphism of a fiber functor to a given S^1 -parameterized family of C -symmetric injective MPSs. We call this automorphism a pump invariant due to its relation to a generalized Thouless pump, which will be detailed in Sec. [V B.](#page-38-0) We expect that the pump invariant is a complete invariant of a family, meaning that S^1 -parameterized families of SPT states with symmetry $\mathcal C$ are classified by automorphisms of a fiber functor.

Group of one-parameter families. The set of $S¹$ -parameterized families that share the same MPS at $\theta = 0$ is equipped with a group structure. Specifically, given a pair of families $\{A_1(\theta)\}\$ and ${A_2(\theta)}$ such that $|A_1(0)\rangle = |A_2(0)\rangle$, their product ${A(\theta)}$ is defined by

$$
A(\theta) = \begin{cases} A_1(2\theta) & 0 \le \theta \le \pi, \\ A_2(2\theta - 2\pi) & \pi \le \theta \le 2\pi. \end{cases}
$$
 (126)

Here, we chose a gauge such that $A_1(2\pi) = A_2(0)$, which is possible because $|A_1(2\pi)\rangle$ and $|A_2(0)\rangle$ are the same state. The automorphism (i.e., the pump invariant) associated with the product $\{A(\theta)\}\$ is given by the composition of the automorphisms associated with $\{A_1(\theta)\}\$ and $\{A_2(\theta)\}\$. This implies that the pump invariant gives rise to a group homomorphism from the group of S^1 -parameterized families of injective MPSs to the group $Aut^{\otimes}(F)$ of automorphisms of a fiber functor $F : \mathcal{C} \to \text{Vec}$. We expect that this homomorphism is an isomorphism, meaning that S^1 -parameterized families of SPT states with symmetry C are classified by Aut[⊗](F) as a group. We note that Aut[⊗](F) can be non-abelian in general. In the case of finite group symmetry $\mathcal{C} = \text{Vec}_G$, we have $\text{Aut}^{\otimes}(F) \cong$ $H^1_{gp}(G,\mathcal{U}(1))$ irrespective of F (cf. Sec. IIC). This agrees with the known classification of S^1 parameterized families of SPT phases with G symmetry [\[15,](#page-84-0) [16,](#page-84-1) [62,](#page-86-0) [63\]](#page-86-1).

Abelianization of the pump invariant. As we saw in Sec. [III B,](#page-24-0) we can abelianize the nonabelian factor systems and define a topological invariant that takes values in the third Hochschild cohomology $H^3(\mathcal{C}, \mathbb{Z})$. Similarly, we can abelianize the pump invariant. To define the abelianization, we first choose a gauge where $A(\theta)$ and $\phi(\theta)$ are 2π-periodic. Note that in this gauge, $L(\theta)$ is a 2π -periodic function that is not necessarily constant. As we saw around Eqs. [\(65\)](#page-22-0) and [\(66\)](#page-22-1), the L-symbols can be identified with the matrix elements of the non-abelian factor system. Since the L-symbols are now parameterized by $\theta \in S^1$, the corresponding non-abelian factor system $\Omega_{x,y}(\theta)$ is also parameterized by S^1 . Then, we define the abelianization of the pump invariant by the winding number of det $(\Omega_{x,y}(\theta))$:

$$
n_{x,y} := \frac{1}{2\pi i} \int d\log \det(\Omega_{x,y}(\theta)).
$$
\n(127)

This invariant takes values in $H^2(\mathcal{C}, \mathbb{Z})$. Let us confirm this property. First of all, the winding number $n_{x,y}$ is quantized to integers due to the periodicity of $\Omega_{x,y}(\theta)$. In addition, $n_{x,y}$ satisfies the cocycle condition $(\delta n)_{x,y,z} = 0$, which follows from the cocycle condition for $\Omega_{x,y}$ described in Eq. [\(52\)](#page-19-0). Furthermore, $n_{x,y}$ is defined up to coboundary because gauge transformations can shift it by a coboundary. More concretely, a gauge transformation that preserves the 2π -periodicity of $A(\theta)$ and $\phi(\theta)$ changes the non-abelian factor system as

$$
\Omega_{x,y}(\theta) \mapsto \left(\bigoplus_{z \in x \otimes y} f_z(\theta)\right)^{-1} \Omega_{x,y}(\theta) \left(f_x(\theta) \otimes f_y(\theta)\right),\tag{128}
$$

where $f_x(\theta)$ is a 2π -periodic dim(x) \times dim(x) invertible matrix, cf. Eq. [\(53\)](#page-19-1). Under this gauge transformation, the winding number $n_{x,y}$ changes as

$$
n_{x,y} \mapsto n_{x,y} + \dim(y) \cdot m_x - \sum_{z \in x \otimes y} m_z + \dim(x) \cdot m_y = n_{x,y} + (\delta m)_{x,y},\tag{129}
$$

where $m_x \in \mathbb{Z}$ is the winding number of $\det(f(\theta)_x)$. Thus, the abelianization $[n_{x,y}]$ of the pump invariant takes values in $\mathrm{H}^2(\mathcal{C}, \mathbb{Z})$. In general, the map $\mathrm{Aut}^{\otimes}(F) \to \mathrm{H}^2(\mathcal{C}, \mathbb{Z})$ defined by $\{\Omega_{x,y}(\theta)\} \mapsto$ $[n_{x,y}]$ is neither surjective nor injective.

The term "abelianization" is justified because the composition law of $[n_{x,y}]$ is abelian. Indeed, for the product [\(126\)](#page-36-0) of two families $\{A_1(\theta)\}\$ and $\{A_2(\theta)\}\$, the winding number $n_{x,y}$ can be computed

$$
n_{x,y} = \frac{1}{2\pi i} \int_0^\pi d\theta \frac{d}{d\theta} \log \det(\Omega_{x,y}^1(2\theta)) + \frac{1}{2\pi i} \int_\pi^{2\pi} d\theta \frac{d}{d\theta} \log \det(\Omega_{x,y}^2(2\theta - 2\pi))
$$

=
$$
\frac{1}{2\pi i} \int d\log \det(\Omega_{x,y}^1(\theta)) + \frac{1}{2\pi i} \int d\log \det(\Omega_{x,y}^2(\theta))
$$

=
$$
n_{x,y}^1 + n_{x,y}^2.
$$
 (130)

Here, $\Omega_{x,y}^i(\theta)$ and $n_{x,y}^i$ are the non-abelian factor system for $\{A^i(\theta)\}\$ and its winding number. Put differently, taking the winding number of the non-abelian factor system gives a homomorphism from $\mathrm{Aut}^{\otimes}(F)$ to its abelianization $\mathrm{Aut}^{\otimes}(F)/[\mathrm{Aut}^{\otimes}(F), \mathrm{Aut}^{\otimes}(F)].$

B. Thouless pump of non-degenerate interface modes

1. Adiabatic evolution of C-symmetric injective MPS

The pump invariant defined in the previous subsection can be interpreted as a generalized Thouless pump. To see this, let us consider an injective MPS on a periodic chain:

$$
|A(0;\theta;0)\rangle = \frac{\left|\begin{array}{c}\psi(\theta) & \psi(\theta) \\ A(0) & A(\theta)\end{array}\right|}{A(\theta)} \xrightarrow{f(\theta)} A(0), \qquad (131)
$$

The state in the middle of the chain is adiabatically evolved along an S^1 -parameterized family ${A(\theta)}$. The interface tensor $\psi(\theta) \in Aut(V)$, which we choose to be unitary, is inserted so that the state $|A(0;\theta;0)\rangle$ is C-symmetric for all $\theta \in [0,2\pi]$, i.e.,

$$
\widehat{\mathcal{O}}_x |A(0; \theta; 0)\rangle = \dim(x) |A(0; \theta; 0)\rangle, \quad \forall x \in \mathcal{C}.
$$
 (132)

We choose $\psi(0) = id_V$ so that the initial state is $|A(0)\rangle$. The condition [\(132\)](#page-38-1) can be expressed in terms of diagrams as

$$
\sum_{i,j} \frac{\left| \psi(\theta)_{x}\right\rangle_{ij}}{A(0)} \frac{\left| \langle \overline{\psi}(\theta)_{x}\rangle_{ji} \right|}{A(\theta)} \xrightarrow{A(0)} A(0) = \dim(x) \frac{\left| \psi(\theta) \right|}{A(0)} \frac{\left| \psi(\theta) \right|}{A(\theta)} \frac{\left| \psi(\theta) \right|}{A(0)} \right|}{A(0)} \tag{133}
$$

where $(\psi(\theta)_x)_{ij}$ and $(\overline{\psi}(\theta)_x)_{ji}$ are defined by

$$
(\psi(\theta)_x)_{ij} = \frac{x}{\begin{pmatrix} \psi(\theta) \\ (\phi(0)_x)_i & (\overline{\phi}(\theta)_x)_j \end{pmatrix}}, \quad (\overline{\psi}(\theta)_x)_{ji} = \frac{x}{\begin{pmatrix} \psi(\theta)^{\dagger} \\ (\phi(\theta)_x)_j & (\overline{\phi}(0)_x)_i \end{pmatrix}}.
$$
(134)

In the following, we will show that the tensor $\psi(\theta)$ that satisfies Eq. [\(133\)](#page-38-2) always exists.

To begin with, let us show that the symmetricity condition [\(133\)](#page-38-2) is satisfied if and only if

$$
(\psi(\theta)_x)_{ij} = (U(\theta)_x)_{ij}\psi(\theta), \quad (\overline{\psi}(\theta)_x)_{ij} = (U(\theta)_x)_{ij}^{-1}\psi(\theta)^\dagger,
$$
\n(135)

as

where $U(\theta)_x$ is a dim(x) \times dim(x) invertible matrix.^{[30](#page-39-0)} The "if" part can be readily seen by substituting Eq. [\(135\)](#page-38-3) into the left-hand side of Eq. [\(133\)](#page-38-2). The "only if" part follows from the injectivity of the MPS tensors $A(0)$ and $A(\theta)$. More concretely, since $A(0)$ and $A(\theta)$ are injective, equation [\(133\)](#page-38-2) is satisfied only if

$$
\sum_{i,j} (\psi(\theta)_x)_{ij} A(\overline{\psi}(\theta)_x)_{ji} = \dim(x) \psi(\theta) A \psi(\theta)^\dagger \tag{136}
$$

for any $D \times D$ matrix A with D being the bond dimension of the MPS. In particular, in a gauge where $(\overline{\phi}(\theta)_x)_i = (\phi(\theta)_x)_i^{\dagger}$ $\frac{1}{i}$, the above equation can be written as

$$
\sum_{i,j} \psi(\theta)^{\dagger} (\psi(\theta)_x)_{ij} A [\psi(\theta)^{\dagger} (\psi(\theta)_x)_{ij}]^{\dagger} = \dim(x) A, \qquad (137)
$$

where we used $(\overline{\psi}(\theta)_x)_{ji} = (\psi(\theta)_x)_{ij}^{\dagger}$, which follows from our gauge choice. On the other hand, in the same gauge, the operators of the form $\psi(\theta)^{\dagger}(\psi(\theta))_{ij}$ satisfy the following Kraus condition:

$$
\sum_{i,j} [\psi(\theta)^{\dagger}(\psi(\theta)_x)_{ij}]^{\dagger} \psi(\theta)^{\dagger}(\psi(\theta)_x)_{ij} = \sum_{i,j} (\overline{\psi}(\theta)_x)_{ji} \psi(\theta) \psi(\theta)^{\dagger}(\psi(\theta)_x)_{ij} = \dim(x) \mathrm{id}_V.
$$
 (138)

Here, the second equality follows from the orthogonality [\(33\)](#page-14-0) and weak completeness [\(43\)](#page-16-0) of the ac-tion tensors. Equations [\(137\)](#page-39-1) and [\(138\)](#page-39-2) show that $\left\{\psi(\theta)^{\dagger}(\psi(\theta)_x)_{ij}/\sqrt{\dim(x)} \mid i, j = 1, 2, \cdots, \dim(x)\right\}$ in this gauge are Kraus operators that implement the identity operation on any $D \times D$ matrix A. Such Kraus operators are known to be proportional to the identity.^{[31](#page-39-3)} Therefore, there exits a set of complex numbers $\{(U(\theta)_x)_{ij}\}\)$ that satisfies

$$
(\psi(\theta)_x)_{ij} = (U(\theta)_x)_{ij}\psi(\theta). \tag{139}
$$

The dim(x) \times dim(x) matrix $U(\theta)_x$ has to be unitary (in particular, invertible) so that Eq. [\(133\)](#page-38-2) holds. In a general gauge, $U(\theta)_x$ may not be unitary but is invertible. Thus, we find that the state $|A(0;\theta;0)\rangle$ is C-symmetric if and only if $\psi(\theta)$ satisfies Eq. [\(135\)](#page-38-3).

Due to the orthogonality relation [\(33\)](#page-14-0) and the weak completeness relation [\(43\)](#page-16-0), equation [\(135\)](#page-38-3) can also be written as

$$
(\phi(\theta)_x)_i = \sum_j (U(\theta)_x)_{ij}^{-1} \underbrace{\psi(\theta)^{-1} \downarrow \psi(\theta)}_{(\phi(0)_x)_j} \psi(\theta).
$$
\n(140)

Let us argue that $\psi(\theta)$ that satisfies the above equation always exits. To this end, we first define operators $(\phi(\theta)_x)_i^{\alpha}$ by

$$
(\phi(\theta)_x)_i^{\alpha} = \frac{\int_{0}^{v_x^{\alpha}}}{(\phi(\theta)_x)_i},
$$
\n(141)

where $\{v_x^{\alpha} \mid \alpha = 1, 2, \cdots, \dim V_x\}$ is a basis of the dual of the bond Hilbert space V_x . The operators $\{(\phi(\theta)_x)_i^{\alpha} \in \text{End}(V) \mid x \in \text{Simp}(\mathcal{C}), i = 1, 2, \cdots, \text{dim}(x), \alpha = 1, 2, \cdots, \text{dim}(V_x)\}$ form a

 30 The two equalities in Eq. [\(135\)](#page-38-3) are equivalent to each other due to the weak completeness relation [\(43\)](#page-16-0).

³¹ See, e.g., [https://quantumcomputing.stackexchange.com/questions/25917/what-are-the-possible-kraus-o](https://quantumcomputing.stackexchange.com/questions/25917/what-are-the-possible-kraus-operators-of-the-identity-channel) [perators-of-the-identity-channel](https://quantumcomputing.stackexchange.com/questions/25917/what-are-the-possible-kraus-operators-of-the-identity-channel)

representation of a finite dimensional algebra whose multiplication is given by the composition $(\phi(\theta)_x)_i^{\alpha}(\phi(\theta)_y)_j^{\beta} \in \text{End}(V)$. The structure constant of this algebra is determined by the L-symbols $L(\theta)$. In particular, the algebra structure is independent of θ if we choose a gauge where the Lsymbols are constant in θ . Since everything is supposed to be continuous in θ , the equivalence class of the representation $\{(\phi(\theta)_x)_i^{\alpha}\}\)$ is also continuous in θ . Thus, if we suppose that the above algebra has only finitely many inequivalent finite dimensional representations, the representation $\{\phi(\theta_x)_i^{\alpha}\}\)$ must be in the same equivalence class for all θ . Therefore, there exits an automorphism $\psi(\theta) \in \text{Aut}(V)$ such that

$$
(\phi(\theta)_x)_i = \psi(\theta)(\phi(0)_x)_i \psi(\theta)^{-1}.
$$
\n(142)

The above equation leads to Eq. [\(140\)](#page-39-4) if we perform a gauge transformation $(\psi(\theta)_x)_i \to \sum_j (U(\theta)_x)_{ij} (\psi(\theta)_x)_j$, where $U(\theta)_x$ is an invertible $\dim(x) \times \dim(x)$ matrix such that $U(0) = 1_x$. Thus, there always exists $\psi(\theta)$ that fits into Eq. [\(140\)](#page-39-4), or equivalently, Eq. [\(133\)](#page-38-2). In other words, one can always find the interface tensor $\psi(\theta)$ in Eq. [\(131\)](#page-38-4) that makes the state $|A(0;\theta;0)\rangle$ symmetric under C.

2. Non-abelian Thouless pump

Now, let us show that changing θ from 0 to 2π pumps a non-degenerate interface mode. To this end, we choose a gauge in which the action tensors $\{(\phi(\theta_x)_i)\}\)$ do not depend on θ . This choice of a gauge is possible due to Eq. [\(140\)](#page-39-4). We note that $\psi(2\pi)$ is the identity in this gauge. Since the action tensors $(\phi(\theta)_x)_i$ are 2π -periodic and the L-symbols $L(\theta)$ are constant in θ , the MPS tensor $A(\theta)$ is not necessarily 2π-periodic. Specifically, the MPS tensors at $\theta = 0$ and $\theta = 2\pi$ are related by a gauge transformation

$$
A(2\pi) = \underbrace{\qquad \qquad}_{\blacktriangle} \underbrace{\qquad \qquad}_{A(0)} \underbrace{(\psi_{\eta})^{-1}}_{\blacktriangle} \tag{143}
$$

where ψ_{η} is the pump invariant associated with the S^1 -parameterized family $\{A(\theta)\}\,$ see Eq. [\(123\)](#page-36-1).^{[32](#page-40-0)} Therefore, going around the parameter space S^1 changes the initial state $|A(0)\rangle$ into

$$
|A(0;2\pi;0)\rangle = \frac{1}{A(0)} \qquad \qquad \downarrow \q
$$

This shows that $|A(0;\theta;0)\rangle$ returns to the initial state $|A(0)\rangle$ at $\theta=2\pi$ except that non-degenerate interface modes ψ_{η} and ψ_{η}^{-1} are localized at the interfaces of two regions. This non-degenerate interface mode is regarded as a generalized Thouless pump. We recall that ψ_n is a one-dimensional representation of the self-interface algebra. Since the one-dimensional representations generally form a non-abelian group, they will be referred to as non-abelian Thouless pumps.

One can also see the generalized Thouless pump in a different gauge where $A(\theta)$ is 2π -periodic and $L(\theta)$ is constant in θ . In this gauge, the action tensors $\{(\phi(\theta)_x)_i\}$ are not necessarily 2π -periodic. Specifically, Eq. [\(140\)](#page-39-4) implies that the action tensors at $\theta = 0$ and $\theta = 2\pi$ are related by

$$
(\phi(2\pi)_x)_i = \frac{\psi(2\pi)^{-1} \downarrow \psi(2\pi)}{(\phi(0)_x)_j} \tag{145}
$$

³² The tensor ψ'_η in Eq. [\(123\)](#page-36-1) is now written as ψ_η .

$$
(\phi(2\pi)_x)_i = \sum_j (\eta_x)_{ij} (\phi(0)_x)_i,
$$
\n(146)

where η is the pump invariant associated with the S^1 -parameterized family $\{A(\theta)\}\)$, see Eq. [\(120\)](#page-35-1). Combining Eqs. [\(145\)](#page-40-1) and [\(146\)](#page-41-0), we find that $\psi(2\pi) = \psi_{\eta}$ in this gauge. Therefore, the state $|A(0; 2\pi; 0)\rangle$ is again given by Eq. [\(144\)](#page-40-2). This is consistent with the fact the Thouless pump does not depend on a gauge.

When C is an ordinary finite group symmetry Vec_G, the pumped interface mode ψ_{η} is a onedimensional representation of the group algebra $\mathbb{C}[G]$, i.e., ψ_{η} is a G-charge. Thus, in this case, the generalized Thouless pump reduces to the ordinary one.

C. Example: $\mathbf{Rep}(G)$ symmetry

As an example, let us consider concrete models of S^1 -parameterized families of SPT states with $Rep(G)$ symmetry. We construct these models using the $G \times Rep(G)$ cluster state that we reviewed in Sec. [III C.](#page-25-0) To obtain an S^1 -parameterized family from the cluster state, we first fix an arbitrary group element $g \in G$ and choose any continuous path $\{R_q(\theta) \mid \theta \in [0, 2\pi]\}\$ of unitary operators, acting on the physical Hilbert space $\mathcal{H} = \mathbb{C}^{|G|}$, such that

$$
R_g(0) = \mathrm{id}_{\mathcal{H}}, \quad R_g(2\pi) = R_g. \tag{147}
$$

Namely, the path ${R_g(\theta)}$ interpolates between the identity operator and the right multiplication of g. Such a path always exists because the space of unitary operators on H (i.e., the space of $|G| \times |G|$ unitary matrices) is path-connected. Given a path $\{R_q(\theta)\}\$ of unitary operators, one can construct a family $\{A_q(\theta) | \theta \in [0, 2\pi]\}$ of MPS tensors by applying $R(\theta)$ to odd sites of the cluster state:

$$
A_g(\theta) = \frac{R_g(\theta)}{A_o} \underbrace{\downarrow}_{A_e} \tag{148}
$$

See Eq. [\(81\)](#page-25-1) for the definition of the MPS tensor of the $G \times \text{Rep}(G)$ cluster state. We note that the MPS tensors $A_q(0)$ and $A_q(2\pi)$ are gauge equivalent because G is a symmetry of the cluster state, cf. Eq. [\(83\)](#page-25-2). Therefore, the parameter space is a circle S^1 . The path of Hamiltonians $\{H_g(\theta)\}\$ is explicitly given by

$$
H_g(\theta) = \left(\prod_{i:\text{odd}} R_g(\theta)_i\right) H_{\text{cluster}} \left(\prod_{i:\text{odd}} R_g(\theta)_i\right)^{\dagger},\tag{149}
$$

where $R_q(\theta)_i$ denotes $R_q(\theta)$ acting on site i and H_{cluster} is the Hamiltonian of the $G \times \text{Rep}(G)$ cluster model given in [\[41\]](#page-85-0).

The above family $\{A_g(\theta)\}\$ preserves the Rep(G) symmetry at each $\theta \in [0, 2\pi]$, meaning that ${A_g(\theta)}$ gives rise to an S^1 -parameterized family of SPT states with $Rep(G)$ symmetry. To compute the pump invariant of this family, we notice that the action tensors $(\phi(\theta)_{\rho})_i$ for the Rep(G) symmetry does not depend on θ by construction:

A^o A^e Rg(θ) Z^ρ = A^o A^e Z^ρ Rg(θ) = X i (ϕ^ρ)i (ϕρ)ⁱ Rg(θ) , (150)

where Z_{ρ} , ϕ_{ρ} , and $\overline{\phi}_{\rho}$ are defined by Eqs. [\(84\)](#page-26-0) and [\(85\)](#page-26-1). Since the action tensors do not depend on θ , the L-symbols are constant in θ . Therefore, the pump invariant of the family appears as the non-periodicity of the MPS tensor $A_q(\theta)$. As we can see from Eq. [\(83\)](#page-25-2), the MPS tensor $A_q(\theta)$ is not 2π -periodic when $g \in G$ is not the unit element. Concretely, $A_q(0)$ and $A_q(2\pi)$ differ by a gauge transformation R_q acting on the bond Hilbert space. Accordingly, the family $\{A_q(\theta)\}\$ pumps a non-degenerate interface mode R_g when θ increases from 0 to 2π . Here, we note that R_g is indeed a one-dimensional representation of the interface algebra, i.e.,

$$
(\mathcal{O}_{\rho})_{ij} R_g = \rho(g)_{ij} R_g. \tag{151}
$$

Thus, we find that the pump invariant of the S^1 -parameterized family $\{A_g(\theta)\}\)$ constructed above is a one-dimensional representation $\{(\widehat{\mathcal{O}}_{\rho})_{ij} = \rho(g)_{ij}\}\$. Since $g \in G$ is arbitrary, one can construct as many families as $|G|$ in this way. Remarkably, the above construction exhausts all S^1 -parameterized families of $\text{Rep}(G)$ SPT states in the trivial phase because one-dimensional representations of the interface algebra of the cluster state are classified by G as we discussed in Sec. [IV C 1.](#page-31-0) We note that the generalized Thouless pump in this example can be non-abelian. This is in contrast to the case of an ordinary group symmetry, where the Thouless pump is always abelian.

VI. CLASSIFICATION OF PARAMETERIZED FAMILIES

In this section, we propose conjectures on the classification of various parameterized families of gapped systems, generalizing the classification of S^1 -parameterized families of 1+1d SPT states discussed in the previous section. Our conjectures are supported by general arguments and concrete examples.

A. General gapped phases in 1+1d

1. Conjecture

Let us consider the classification of S^1 -parameterized families of general 1+1d gapped systems with fusion category symmetry C . In general, C -symmetric gapped phases in 1+1d are in one-to-one correspondence with module categories over \mathcal{C} [\[32\]](#page-85-1). In particular, SPT phases with symmetry \mathcal{C} correspond to C-module categories with a single simple object, which can be identified with fiber functors of \mathcal{C} [\[48\]](#page-86-2). In what follows, the gapped phase labeled by a \mathcal{C} -module category \mathcal{M} is denoted by $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$.

Our conjecture on the classification of S^1 -parameterized families of general C-symmetric gapped systems is as follows:

FIG. 2. The symmetry TFT construction of $1+1d$ systems with symmetry C. The left boundary of the slab is topological and supports topological lines described by C . On the other hand, the right boundary is arbitrary and dictates the dynamics of the 1+1d system. When the right boundary is also topological, the whole 1+1d system becomes topological.

Conjecture 1. S^1 -parameterized families of 1+1d C -symmetric gapped systems in gapped phase $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$ are classified by the group $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$ of invertible objects of the category $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})$ of C-module endofunctors of M.

Let us give a motivation for the above conjecture. First of all, since \mathcal{C} -symmetric gapped phases are classified by C-module categories, an S^1 -parameterized family of gapped systems should give rise to an S^1 -parameterized family of C-module categories $\mathcal{M}(\theta)$. When the gapped systems in the family belong to the same phase $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$, the C-module category $\mathcal{M}(\theta)$ is equivalent to M for all $\theta \in S^1 = [0, 2\pi]$. In particular, $\mathcal{M}(0)$ and $\mathcal{M}(2\pi)$ are the same module category, which we choose to be M. Nevertheless, going around S^1 can induce a non-trivial automorphism of M, as we saw in the case of SPT phases (cf. Sec. [V\)](#page-33-0). Thus, an S^1 -parameterized family of gapped systems is associated with an automorphism of M , i.e., an invertible C-module endofunctor of M . We expect that this automorphism classifies S^1 -parameterized families of gapped systems. In the context of MPS, it should be straightforward to construct an invariant of the classification by applying the analysis in Sec. V \AA to \mathcal{C} -symmetric block-injective MPSs.

When M has only one simple object, the group of invertible objects of $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})$ is isomorphic to the group Aut[⊗](F) of the fiber functor F corresponding to M. Thus, as a special case, our conjecture implies that S^1 -parameterized families of SPT states labeled by a fiber functor F are classified by Aut[⊗](F). This is supported by the discussions on pump invariants in Sec. VA.

Relation to symmetry TFT. Conjecture [1](#page-43-0) is also natural from the perspective of symmetry TFT. In the symmetry TFT construction, a \mathcal{C} -symmetric system in 1+1d is realized as a threedimensional TFT on a slab as shown in Fig. [2.](#page-43-1) The 3d TFT in the bulk is the Turaev-Viro-Barrett-Westbury TFT TV (C) [\[81,](#page-87-0) [82\]](#page-87-1), which is the low-energy limit of the 2+1d string-net model [\[83\]](#page-87-2). In this construction, the dynamics of the $1+1d$ system is determined by the choice of a boundary condition of $TV(\mathcal{C})$. In particular, 1+1d gapped systems are obtained by choosing this boundary to be topological. The C-symmetric gapped phase $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$ corresponds to the topological boundary labeled by the same C-module category M . Thus, in this framework, an S^1 -parameterized family of C-symmetric gapped states corresponds to an S^1 -parameterized family of topological boundaries of $TV(\mathcal{C})$.

Given a family $\{\mathcal{M}(\theta) \mid \theta \in S^1\}$ of topological boundaries of $TV(\mathcal{C})$, one can consider a spatially modulated boundary of the symmetry TFT as illustrated in Fig. [3.](#page-44-0) If the whole system is viewed from far away, the modulation looks like a topological line on the boundary. This topological line is invertible because the modulation can be undone by the reversed modulation. Therefore, from the perspective of symmetry TFT, S^1 -parameterized families of C-symmetric gapped states should be classified by invertible topological lines on the topological boundary M . Since topological lines on M are classified by $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})$ [\[72,](#page-87-3) [84,](#page-87-4) [85\]](#page-87-5), we are led to Conjecture [1.](#page-43-0)

FIG. 3. The middle region of the right boundary is modulated gradually from $\mathcal{M}(0)$ to $\mathcal{M}(2\pi)$. This modulated region becomes an invertible topological line when viewed from far away. After squashing the bulk, the invertible line on the boundary becomes an invertible interface in 1+1d. The correspondence between invertible interfaces and S^1 -parameterized families was already observed in Sec. [IV B](#page-29-0) in the case of SPT phases.

Relation to gauging. Provided that the conjecture holds for some gapped phase \mathcal{T} , we can show that the conjecture also holds for other gapped phases obtained by the gauging of $\mathcal T$. To see this, we first recall the gauging of 1+1d gapped phases with fusion category symmetries. Let $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$ be a gapped phase labeled by a C-module category M and let $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}/A$ denote the gauging of $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$ by a separable algebra object $A \in \mathcal{C}$. The symmetry of $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}/A$ is described by the category $A\mathcal{C}_{A}$ of (A, A) -bimodules in C [\[30\]](#page-85-2). Due to Ostrik's theorem [\[51\]](#page-86-3), any C-module category M can be written as the category \mathcal{C}_B of right B-modules in \mathcal{C} , where B is also a separable algebra object in \mathcal{C} . When $\mathcal{M} = \mathcal{C}_B$, the $_A \mathcal{C}_A$ -module category corresponding to the gauged theory $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}/A$ is the category $_A \mathcal{C}_B$ of (A, B) -bimodules in C [\[55,](#page-86-4) Appendix A].

Now, we assume that the conjecture holds for $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$, that is, S^1 -parameterized families of $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$ are classified by the group of invertible objects of $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M}) = \text{Fun}_{\mathcal{C}}(\mathcal{C}_B,\mathcal{C}_B)$. Since the gauging is an invertible operation [\[30\]](#page-85-2),^{[33](#page-44-1)} S¹-parameterized families of the gauged theory $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}/A$ are also classified by Func (C_B, C_B) ^{inv}. On the other hand, Theorem 7.12.16 of [\[48\]](#page-86-2) implies that Func (C_B, C_B) is equivalent to $\text{Fun}_{_A\mathcal{C}_A}(_A\mathcal{C}_B, _A\mathcal{C}_B)$ as a fusion category:

$$
\operatorname{Fun}_{\mathcal{C}}(\mathcal{C}_B, \mathcal{C}_B) \cong \operatorname{Fun}_{A\mathcal{C}_A}(\operatorname{Fun}_{\mathcal{C}}(\mathcal{C}_A, \mathcal{C}_B), \operatorname{Fun}_{\mathcal{C}}(\mathcal{C}_A, \mathcal{C}_B)) \cong \operatorname{Fun}_{A\mathcal{C}_A}(A\mathcal{C}_B, A\mathcal{C}_B). \tag{152}
$$

Therefore, S¹-parameterized families of the gauged theory $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}/A = \mathcal{T}_{A}^{\mathcal{A}\mathcal{C}_{A}}$ $A_{\mathcal{C}_B}^{\mathcal{A} \mathcal{C}_A}$ are classified by the group of invertible objects of $\text{Fun}_{A\mathcal{C}_{A}}(A\mathcal{C}_{B}, A\mathcal{C}_{B})$. This shows that the conjecture holds for the gauged theory $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}/A$ as long as it holds for the original theory $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$.

2. Examples

Example 1: G-SSB phase. As a simple example, we apply our conjecture to the gapped phase that spontaneously breaks a finite group symmetry G down to the trivial group. The fusion category that describes a finite group symmetry G is the category Vec $_G$ of G -graded vector spaces. The Vec_G-module category corresponding to the G-SSB phase is the regular module Vec_G.^{[34](#page-44-2)} Thus, our conjecture implies that S^1 -parameterized families of gapped systems in the G-SSB phase are classified by the group of invertible objects of $\text{Fun}_{\text{Vec}_G}(\text{Vec}_G, \text{Vec}_G) \cong \text{Vec}_G$, which is G. Physically, this group G can be interpreted as the group of invertible domain walls pumped by S^1 -parameterized families.

One can also understand this classification from the point of view of gauging. If we gauge the symmetry G of the G-SSB phase, we obtain a $Rep(G)$ SPT phase that corresponds to the forgetful

³³ Precisely, there exists a separable algebra $A' \in {}_{A}\mathcal{C}_{A}$ such that $T/A/A'$ goes back to the original theory \mathcal{T} .

³⁴ For a fusion category C, a regular C-module is C itself, on which C acts by its tensor product $\otimes : \mathcal{C} \times \mathcal{C} \to \mathcal{C}$.

functor of Rep(G). Our conjecture implies that S^1 -parameterized families of invertible states in this $Rep(G)$ SPT phase are classified by the group of automorphisms of the forgetful functor, which is isomorphic to G due to Tannaka-Krein duality. Thus, S^1 -parameterized families of gapped systems in the G-SSB phase should also be classified by G because the gauging is invertible.

Example 2: C-SSB phase. More generally, we can consider the SSB phase of a general fusion category symmetry C. The C-module category corresponding to the C-SSB phase is the regular module category C. The associated functor category is Func $(\mathcal{C}, \mathcal{C}) \cong \mathcal{C}$, and hence Conjecture [1](#page-43-0) suggests that S^1 -parameterized families of gapped systems in the C-SSB phase are classified by \mathcal{C}^{inv} , the group of invertible objects of \mathcal{C} .

Example 3: $\mathbb{Z}_4/\mathbb{Z}_2$ -SSB phase. One can also consider examples where the symmetry is partially broken. A simple example is the spontaneous symmetry breaking of \mathbb{Z}_4 down to \mathbb{Z}_2 . The Vec_{\mathbb{Z}_4 -} module category corresponding to this gapped phase is Vec_{Z_2} . The action of Vec_{Z_4} on Vec_{Z_2} is defined via the group homomorphism that maps $a \in \mathbb{Z}_4$ to $a \pmod{2} \in \mathbb{Z}_2$. The corresponding functor category is $\text{Fun}_{\text{Vec}_{\mathbb{Z}_4}}(\text{Vec}_{\mathbb{Z}_2}, \text{Vec}_{\mathbb{Z}_2}) \cong \text{Vec}_{\mathbb{Z}_2 \times \mathbb{Z}_2}^{\omega}$ [\[30,](#page-85-2) [86–](#page-87-6)[89\]](#page-87-7), where $[\omega] \neq 0 \in H_{gp}^3(\mathbb{Z}_2 \times$ \mathbb{Z}_2 , U(1)) is given by

$$
\omega((a_1, b_1), (a_2, b_2), (a_3, b_3)) = (-1)^{a_1 a_2 b_3} \tag{153}
$$

for $a_i, b_i \in \mathbb{Z}_2 = \{0, 1\}$ $a_i, b_i \in \mathbb{Z}_2 = \{0, 1\}$ $a_i, b_i \in \mathbb{Z}_2 = \{0, 1\}$. Thus, Conjecture 1 suggests that S^1 -parameterized families of gapped systems in the $\mathbb{Z}_4/\mathbb{Z}_2$ -SSB phase are classified by $\mathbb{Z}_2\times\mathbb{Z}_2$. We note that $[\omega]\in H^3_{\text{gp}}(\mathbb{Z}_2\times\mathbb{Z}_2, U(1))$ does not affect the classification of S^1 -parameterized families. However, it would affect the classification for more general parameter spaces, see Sec. [VI B](#page-45-0) for more details.

B. General parameter space in 1+1d

1. Conjecture

In the discussions so far, the parameter space was supposed to be a circle $S¹$. More generally, one can consider families of $1+1d$ gapped systems parameterized by a general parameter space X, which we assume to be sufficiently nice (e.g., a CW complex). Our conjecture on the classification of such families is as follows^{[35](#page-45-1)}:

Conjecture 2. X-parameterized families of 1+1d gapped systems in a gapped phase $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$ are classified by the set $[X, B_{\text{Fun}_{\mathcal{C}}}(\mathcal{M}, \mathcal{M})^{\text{inv}}]$ of homotopy classes of maps from X to the classifying space of a 2-group $\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$. See below for the definition of $\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$. The set $[X, B\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}]$ is also known as the non-abelian Čech cohomology $\check{H}^1(X, \underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}})$.

The above conjecture is based on the hypothesis that the space of $1+1d$ systems in a C -symmetric gapped phase labeled by M is homotopy equivalent to $B_{\text{Fun}_{\mathcal{C}}}(\mathcal{M},\mathcal{M})^{\text{inv}}$. See [\[90–](#page-87-8)[92\]](#page-87-9) for a similar conjecture for $2+1d$ $2+1d$ $2+1d$ $2+1d$ topologically ordered phases.^{[36](#page-45-2)} Conjecture 2 reduces to Conjecture 1 when the parameter space X is a circle because we have

$$
[S^1, B\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\mathrm{inv}}] = \pi_1(B\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\mathrm{inv}}) \cong \mathrm{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\mathrm{inv}}.
$$
(154)

 35 In the following, we consider the classification of families for a fixed module category M . Therefore, the classification for $X = pt$ is always trivial.

 36 In $[90-92]$ $[90-92]$, it is conjectured that the space of 2+1d gapped Hamiltonians that realize a topological order described by a modular tensor category β (up to invertible topological order) is homotopy equivalent to the classifying space $B\text{Pic}(B)$ of the categorical Picard 2-group $\text{Pic}(B)$. Here, the categorical Picard 2-group $\text{Pic}(B)$ is a monoidal 2-category consisting of invertible β -modules, invertible β -module functors, and invertible β -module natural transformations [\[93\]](#page-87-10). We will return to this conjecture at the end of Sec. [VI D.](#page-54-0)

See Eq. [\(167\)](#page-48-0) for more general homotopy groups of $B\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$. As in the case of Conjecture [1,](#page-43-0) if Conjecture [2](#page-45-3) holds for a C-symmetric gapped phase $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$, it also holds for any other gapped phases obtained by the gauging of $\mathcal{T}_{\mathcal{M}}^{\mathcal{C}}$ because $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})$ is invariant under gauging.

Let us unpack the above conjecture in detail. In Conjecture [2,](#page-45-3) $\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\mathrm{inv}}$ denotes a categorical group consisting of invertible objects and invertible morphisms of $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})$. More concretely, $\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$ is a 2-group $(G,A,\rho,\beta),^{37}$ $(G,A,\rho,\beta),^{37}$ $(G,A,\rho,\beta),^{37}$ where the 0-form group G is $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}},$ the 1-form group A is U(1), the action $\rho: G \to \text{Aut}(A)$ is trivial, and the Postnikov class $[\beta] \in$ $\mathrm{H}^3_{\text{gp}}(\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}},\mathrm{U}(1))$ is given by the F-symbols of the invertible subcategory of $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})$. We suppose that the 1-form group $U(1)$ is equipped with the standard (i.e., continuous) topology of the Lie group U(1), not the discrete topology. On the other hand, the 0-form group $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$ has the discrete topology because it is a finite group. As we will see below, the choice of a topology of $U(1)$ plays an important role in the classification of parameterized families. In what follows, $U(1)$ equipped with the standard topology will be simply written as $U(1)$, whereas the same underlying group equipped with the discrete topology will be denoted by $U(1)_{\delta}$.

The classifying space of $\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$ is denoted by $B\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$. If we use U(1) with the standard topology, the realization of $B\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\mathrm{inv}}$ as a space is given by a Postnikov tower

$$
P_3 \leftarrow K(\mathbb{Z}, 3)
$$
\n
$$
\downarrow
$$
\n
$$
P_2 \leftarrow \qquad * \qquad (155)
$$
\n
$$
\downarrow
$$
\n
$$
B\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}} \longrightarrow P_1 = K(\text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}, 1)
$$

with the Postnikov invariant $\beta_1([\beta]) \in H^4(BFun_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}},\mathbb{Z})$. Here, $\beta_1: H^3(\bullet,U(1)) \to H^4(\bullet,\mathbb{Z})$ is the Bochstein homomorphism associated with $0 \to \mathbb{Z} \to \mathbb{R} \to U(1)$, and surjective arrows represent Serre fibrations. If we use $U(1)_{\delta}$ with the discrete topology, the realization as a space is given by a Postnikov tower

$$
B_{2} \longleftarrow K(U(1)_{\delta}, 2)
$$
\n
$$
\downarrow \qquad K(U(1)_{\delta}, 2)
$$
\n
$$
B_{2} \longleftarrow K(\text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}, 1)
$$
\n
$$
(156)
$$

with the Postnikov invariant $[\beta] \in H^3(BFun_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}},U(1)_{\delta}) \simeq H^3_{\text{gp}}(\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}},U(1))$.^{[38](#page-46-1)} See, e.g., [\[98\]](#page-88-0) for details of the classifying space of a general topological 2-group.

For our purposes, it suffices to know the following fact about the classifying space: for a topological 2-group \mathbb{G} , the homotopy classes of maps from X to $B\mathbb{G}$ are in one-to-one correspondence with elements of the non-abelian Čech cohomology $\check{H}^1(X, \mathbb{G})$ [\[98](#page-88-0)[–102\]](#page-88-1), i.e.,

$$
[X, B\mathbb{G}] = \check{H}^1(X, \mathbb{G}).\tag{157}
$$

³⁷ There are various descriptions of a 2-group [\[94\]](#page-87-11). In this paper, a 2-group G refers to a quadruple (G, A, ρ, β) with G a group, A an abelian group, ρ a group homomorphism from G to Aut(A), and β a group 3-cocycle of G with coefficients in A. We refer the reader to, e.g., [\[95,](#page-88-2) [96\]](#page-88-3) for more detailed expositions of 2-groups for physicists.

³⁸ We note $H^3_{gp}(G, A) \simeq H^i(BG, A)$ if A is discrete [\[97\]](#page-88-4).

The explicit description of the Čech cohomology $\check{H}^1(X,\mathbb{G})$ for a general topological 2-group $\mathbb G$ is given in [\[98,](#page-88-0) Section 4]. The description in [\[98\]](#page-88-0) is based on the crossed module corresponding to $\mathbb{G} = (G, A, \rho, \beta)$. When G is finite and ρ is trivial as in the case of our interest $\mathbb{G} = \underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}$, one can directly describe the Čech cohomology $\check{H}^1(X,\mathbb{G})$ in terms of (G,A,β) [\[103\]](#page-88-5). We will review this description below. See also $[95]$ for the direct description of the Cech cohomology with coefficients in a general discrete 2-group.

Non-abelian Čech cohomology. For concreteness, let us give an explicit description of the Cech cohomology $\check{H}^1(X,\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}})$ following [\[103\]](#page-88-5). We first fix an open cover $\{U_i\}_{i\in I}$ of X, where I is an index set. The overlaps of open subsets are denoted by $U_{ij\cdots k} := U_i \cap U_j \cap \cdots \cap U_k$. Here, any finite overlaps $U_{ij\cdots k}$ are supposed to be contractible, i.e., $\{U_i\}$ is a good open cover. An element of $\check{H}^1(X,\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}})$ is an equivalence class of a pair $(\{g_{ij}\},\{a_{ijk}\})$ of continuous functions

$$
g_{ij}: U_{ij} \to \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}, \qquad a_{ijk}: U_{ijk} \to U(1) \tag{158}
$$

such that

$$
(dg)_{ijk}(x) := g_{jk}(x)g_{ik}(x)^{-1}g_{ij}(x) = 1, \qquad \forall x \in U_{ijk},
$$

\n
$$
(da)_{ijkl}(y) := a_{jkl}(y)a_{ikl}(y)^{-1}a_{ijl}(y)a_{ijk}(y)^{-1} = \beta(g_{ij}(y), g_{jk}(y), g_{kl}(y)), \quad \forall y \in U_{ijkl}.
$$
\n(159)

Here, the 3-cocycle β is normalized so that $\beta(g, h, k) = 1$ if any of g, h, and k is the unit element of $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$. This normalization is always possible without changing the cohomology class of β. Two pairs $({g_{ij}}$, ${a_{ijk}}$) and $({g'_{ij}}$, ${a'_{ijk}}$) are said to be equivalent if and only if there exists a pair $({f_i}, {(\gamma_{ij})})$ of continuous functions

$$
f_i: U_i \to \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}, \qquad \gamma_{ij}: U_{ij} \to U(1) \tag{160}
$$

such that

$$
g'_{ij}(x) = f_i(x)g_{ij}(x)f_j(x)^{-1}, \qquad \forall x \in U_{ij},
$$

\n
$$
a'_{ijk}(y) = a_{ijk}(y)(d\gamma)_{ijk}(y)\zeta_{ijk}(y), \quad \forall y \in U_{ijkl},
$$
\n(161)

where

$$
\zeta_{ijk}(y) := \frac{\beta(g'_{ij}(y), f_j(y), g_{jk}(y))}{\beta(f_i(y), g_{ij}(y), g_{jk}(y))\beta(g'_{ij}(y), g'_{jk}(y), f_k(y))}.
$$
\n(162)

The function ζ is called the first descendant of β [\[95,](#page-88-2) [104\]](#page-88-6) and satisfies

$$
(\delta \zeta)_{ijkl}(y) = \beta(g'_{ij}(y), g'_{jk}(y), g'_{kl}(y)) \beta(g_{ij}(y), g_{jk}(y), g_{kl}(y))^{-1},
$$
\n(163)

which follows from the cocycle condition on β . Summarizing, the Cech cohomology is given by the quotient

$$
\check{H}^{1}(X, \underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}) = \{ (\{g_{ij}\}, \{a_{ijk}\}) \text{ satisfying Eq. (159)} \}/ \sim,
$$
\n(164)

where the equivalence relation $(\{g_{ij}\}, \{a_{ijk}\}) \sim (\{g'_{ij}\}, \{a'_{ijk}\})$ is defined by Eq. [\(161\)](#page-47-1). We note that $g_{ij}(x)$ and $f_i(x)$ in Eqs. [\(159\)](#page-47-0) and [\(161\)](#page-47-1) are constant on U_{ij} and U_i because $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$ is finite and hence has the discrete topology.

It is worth mentioning that the Čech cohomology $\check{H}^1(X,\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\mathrm{inv}})$ depends on the topology of the 1-form group $U(1)$. This is because continuous functions involved in the definition (164) depend on the topology of $U(1)$. Specifically, since $U(1)$ has the standard topology, the values $\alpha_{ijk}(y)$ and $\gamma_{ij}(y)$ in Eqs. [\(159\)](#page-47-0) and [\(161\)](#page-47-1) can vary continuously with respect to y. On the other hand, if we replace U(1) with U(1)_δ, the functions $a_{ijk}(y)$ and $\gamma_{ij}(y)$ have to be constant on U_{ijk} and U_{ij} .

When the Postnikov class $[\beta] \in H^3_{gp}(\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}},U(1))$ is trivial, the functions g_{ij} and a_{ijk} in Eq. (159) are independent of each other. As a result, the Čech cohomology (164) (164) is decomposed into the direct sum

$$
\check{H}^1(X, \underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}) = \check{H}^1(X, \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}) \oplus \check{H}^2(X, U(1))
$$
\n
$$
= \check{H}^1(X, \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}) \oplus \check{H}^3(X, \mathbb{Z}).
$$
\n(165)

Here, the second equality follows from the identity

$$
\check{H}^{2}(X, U(1)) = [X, B^{2}U(1)] = [X, K(\mathbb{Z}, 3)] = \check{H}^{3}(X, \mathbb{Z}),
$$
\n(166)

where $K(\mathbb{Z},3) = B^3\mathbb{Z} = B^2U(1)$ is the third Eilenberg-MacLane space of \mathbb{Z} . We note that the second line of Eq. (165) can also be regarded as the singular cohomology because the Čech cohomology with coefficients in a discrete group agrees with the singular cohomology with the same coefficient.

The Čech cohomology $\check{H}^1(X,\underline{\mathrm{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^\mathrm{inv})$ does not necessarily have a group structure when X is a general parameter space. This is because there is no natural way to define the product of maps from X to $B_{\text{Fun}_{\mathcal{C}}}(\mathcal{M}, \mathcal{M})^{\text{inv}}$ in general.

 S^k -parameterized families. Let us apply Conjecture [2](#page-45-3) to the case where the parameter space X is a k-dimensional sphere S^k . When $k = 1$, the above explicit description of the Čech cohomology shows that $\check{H}^1(S^1, \underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}})$ is given by $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$. To compute the Čech cohomology of $S^{k\geq 2}$, we note that one can always take $\{g_{ij}\}\$ to be trivial when $k\geq 2$ because, for finite group G, any principal G-bundle over $S^{k\geq 2}$ can be trivialized. Therefore, the Čeck cohomology $\check{H}^1(S^k, \underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}})$ for $k \geq 2$ reduces to the second Čech cohomology of S^k with coefficients in $U(1)$. Since $U(1)$ is equipped with the standard topology, the second Čech cohomology is given by

$$
\check{H}^{2}(S^{k}, U(1)) = \check{H}^{3}(S^{k}, \mathbb{Z}) = \delta_{k,3}\mathbb{Z}.
$$
\n(167)

Thus, we find

$$
\check{H}^{1}(S^{k}, \underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}) = \pi_{k}(B\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}) = \begin{cases} \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}} & k = 1, \\ \mathbb{Z} & k = 3, \\ 0 & k \neq 1, 3. \end{cases} \tag{168}
$$

We note that the Čech cohomology of S^3 is given by Z regardless of C and M, meaning that S^3 -parameterized families of any C-symmetric gapped phase $\mathcal M$ are classified by $\mathbb Z$.

We mention that replacing U(1) with U(1)_δ leads us to a different result. Specifically, the Cech cohomology of S^k with coefficients in $\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})_{\delta}^{\text{inv}}$ equipped with the discrete topology is given by

$$
\check{H}^{1}(S^{k}, \underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}_{\delta}) = \pi_{k}(B\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}}_{\delta}) = \begin{cases} \text{Fun}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}} & k = 1, \\ \text{U}(1) & k = 2, \\ 0 & k \ge 3. \end{cases}
$$
(169)

This result is inconsistent with the known classification of invertible states in $1+1d$ [\[11–](#page-84-2)[13\]](#page-84-3). Thus, one should employ the standard topology of $U(1)$ rather than the discrete one. Choosing a suitable topology of U(1) is analogous to choosing a suitable dual (i.e., the Anderson dual) of the bordism group in the classification of SPT phases $[105-107]$ $[105-107]$.^{[39](#page-49-0)}

2. Examples

Example 1: Trivial phase without symmetry. As the simplest example, let us consider parameterized families of trivial states without symmetry. The symmetry category in this case is given by $\mathcal{C} = \text{Vec}$, and the trivial phase corresponds to the Vec-module category $\mathcal{M} = \text{Vec}$. Thus, the functor category $\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})$ is equivalent to Vec. In particular, the 2-group $\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$ consists only of a 1-form group $U(1)$. Consequently, the classification of X-parameterized families of trivial states is given by

$$
\check{H}^1(X, \underline{\mathrm{Vec}}^{\mathrm{inv}}) = \check{H}^3(X, \mathbb{Z}).\tag{170}
$$

This agrees with Kitaev's proposal for the classification of invertible states in $1+1d$ [\[11–](#page-84-2)[13\]](#page-84-3). The invariant of the classification is known as a higher Berry phase [\[17,](#page-84-4) [90,](#page-87-8) [108–](#page-88-9)[117\]](#page-88-10). From our point of view, this classification originates from the fact that symmetry category Vec of the trivial states contains the U(1) 1-form symmetry, which exists in any quantum systems in $1+1d^{40}$ $1+1d^{40}$ $1+1d^{40}$

Example 2: SPT phases with G symmetry. One can also incorporate a finite group symmetry G in the above example. A 1+1d SPT phase with symmetry G is labeled by a fiber functor of Vec_G , or equivalently, a Vec_G -module category with a single simple object. When the simple object of a Vec_G-module category M is unique, the functor category $\text{Fun}_{\text{Vec}_G}(\mathcal{M},\mathcal{M})$ is equivalent to $Rep(G)$ as a fusion category [\[48\]](#page-86-2). Therefore, the group of invertible objects of $Fun_{Vec_G}(\mathcal{M}, \mathcal{M})$ is isomorphic to an abelian group $H^1_{gp}(G, U(1))$, i.e., the group of one-dimensional representations of G. Furthermore, since $\text{Rep}(G)$ is non-anomalous, its invertible subcategory is also non-anomalous, meaning that the F-symbols of this subcategory are trivial. Thus, the Postnikov class β of the 2-group $\underline{\text{Fun}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$ is trivial. Consequently, the Čech cohomology splits into the direct sum

$$
\check{H}^1(X, \underline{\text{Rep}(G)}^{\text{inv}}) = \check{H}^1(X, \mathrm{H}^1_{\text{gp}}(G, \mathrm{U}(1))) \oplus \check{H}^3(X, \mathbb{Z}).\tag{171}
$$

This agrees with the known classification of X-parameterized families of $1+1d$ SPT states with symmetry G [\[14,](#page-84-5) [16,](#page-84-1) [62,](#page-86-0) [90\]](#page-87-8).

Example 3: SPT phases with $\text{Rep}(G)$ **symmetry.** As another example, let us consider parameterized families of Rep(G) SPT states labeled by the forgetful functor of Rep(G). The Rep(G)module category corresponding to this phase is Vec, on which $Rep(G)$ acts via the forgetful functor. The functor category $\text{Fun}_{\text{Rep}(G)}(\text{Vec}, \text{Vec})$ is equivalent to Vec_G [\[48\]](#page-86-2), and hence the group of its invertible objects is isomorphic to G. Furthermore, since Vec_G is non-anomalous, its F-symbols are

 39 The topology of U(1) needs to be set according to the specific objectives. For example, in classifications using effective field theory, the non-triviality of $U(1)$ is detected through the partition function. If one considers the values of the partition function to have physical significance, it is natural to use the discrete topology [\[92\]](#page-87-9). On the other hand, in analyses using lattice systems, U(1) appears as an unphysical phase. Therefore, it is natural to endow U(1) with the standard topology.

⁴⁰ In general, any quantum systems in $(n + 1)d$ have a U(1) *n*-form symmetry, which is generated by scalar multiples of the identity point operator. Usually, a system only with this symmetry is said to have no symmetry.

trivial, meaning that the Postnikov class of $\underline{\text{Fun}}_{\text{Rep}(G)}(\text{Vec},\text{Vec})^{\text{inv}}$ is trivial. Therefore, Conjecture [2](#page-45-3) implies that X-parameterized families are classified by

$$
\check{H}^1(X, \underline{\mathrm{Vec}}_G^{\mathrm{inv}}) = \check{H}^1(X, G) \oplus \check{H}^3(X, \mathbb{Z}).\tag{172}
$$

We note that $\check{H}^1(X, G)$ does not admit a group structure for generic X when G is non-abelian. Equation [\(172\)](#page-50-0) implies that S^1 -parameterized families are classified by G. Concrete models of such families were constructed in Sec. [V C](#page-41-1) by using the $G \times \text{Rep}(G)$ cluster state.

Example 4: SPT phases with $\text{Rep}(H)$ **symmetry.** More generally, for any finite-dimensional semisimple Hopf algebra H, its representation category $\text{Rep}(H)$ has an SPT phase labeled by the forgetful functor of Rep(H). The corresponding Rep(H)-module category is Vec, on which Rep(H) acts via the forgetful functor. The functor category $\text{Fun}_{\text{Rep}(H)}(\text{Vec}, \text{Vec})$ is equivalent to $\text{Rep}(H^*),$ where H^* is the dual of H [\[48,](#page-86-2) Example 7.12.26]. In particular, the group of invertible objects of Fun_{Rep}_(H)(Vec, Vec) is isomorphic to $G(H)$, the group of group-like elements of H. Since Rep^(H*) is also non-anomalous, its group-like subcategory has the trivial F -symbols, meaning that the 2group $\underline{\text{Fun}}_{\text{Rep}(H)}(\text{Vec}, \text{Vec})$ has the trivial Postnikov class. Therefore, Conjecture [2](#page-45-3) suggests that X-parameterized families are classified by

$$
\check{H}^1(X, \underline{\text{Rep}(H^*)}^{\text{inv}}) = \check{H}^1(X, G(H)) \oplus \check{H}^3(X, \mathbb{Z}).\tag{173}
$$

In particular, S^1 -parameterized families are classified by $G(H)$. As in the case of Rep(G), it should be possible to construct concrete models of these families by using the Hopf algebra analogue of the $G \times \text{Rep}(G)$ cluster state [\[43\]](#page-86-5). We leave this construction for the future.

C. Non-chiral SPT phases in 2+1d

1. Conjecture

One can also consider parameterized families of gapped systems in higher dimensions. In this subsection, we focus on S^1 - and S^2 -parameterized families of 2+1d non-chiral SPT states with finite non-invertible symmetries.^{[41](#page-50-1)} More general (non-chiral) gaped systems parameterized by more general parameter spaces will be discussed in Sec. [VI D.](#page-54-0)

Finite non-invertible symmetries in 2+1 dimensions are expected to be described by fusion 2 categories. Physically, a fusion 2-category consists of topological surfaces, topological lines, and topological point operators. The associativity of their fusion product is captured by a higher analog of the F-symbols, known as the 10-j symbols. See $[118, 119]$ $[118, 119]$ $[118, 119]$ for the precise definition of fusion 2-categories.

It is natural to expect that 2+1d non-chiral SPT phases with fusion 2-category symmetry $\mathcal C$ are classified by fiber 2-functors of $\mathcal C$. Here, a fiber 2-functor is a tensor 2-functor from $\mathcal C$ to the 2-category 2Vec of finite semisimple 1-categories. Our conjecture for the classification of $S¹$ - and $S²$ -parameterized families of invertible states in these phases is as follows:

Conjecture 3. S^1 -parameterized families of non-chiral 2+1d SPT states labeled by a fiber 2-functor $F: \mathcal{C} \to 2\text{Vec}$ are classified by the group of isomorphism classes of monoidal natural automorphisms of F. Similarly, S^2 -parameterized families of these SPT states are classified by the group of invertible monoidal modifications of the identity natural transformation of F.

 41 A gapped phase is said to be non-chiral if and only if it admits gapped boundaries.

FIG. 4. An S^2 -family is viewed as an S^1 -family of S^1 -families. Each loop represents an S^1 -family embedded in the S^2 -family. The black dot at the top represents a constant family, which we choose to be the initial and final S^1 -families.

We refer the reader to [\[120\]](#page-89-0) [\[121,](#page-89-1) Chapter 2] for the definitions of monoidal 2-functors, monoidal natural transformations of monoidal 2-functors, and monoidal modifications of monoidal natural transformations.

Our conjecture on S^1 -parameterized families is motivated by a similar argument to the case of 1+1d. Since an SPT phase is labeled by a fiber 2-functor F , an S^1 -parameterized family of SPT states in the same phase gives rise to an S^1 -parameterized family of fiber 2-functors, all of which are isomorphic to F. In particular, going around the parameter space $S¹$ induces a natural automorphism of F. We expect that this automorphism completely characterizes the $S¹$ parameterized family of these SPT states.

To motivate the conjecture on S^2 -parameterized families, we point out that an S^2 -family can be viewed as an S^1 -family of S^1 -families as illustrated in Fig. [4.](#page-51-0) The initial and final S^1 -families are represented by a single dot in the figure, which indicates that these families are constant in $\theta \in S^1$. Since any loops in S^2 are contractible, any S^1 -familes embedded in the S^2 -family are trivial. Consequently, the monoidal natural transformations induced by these $S¹$ -families are isomorphic to the identity natural transformation. In particular, the monoidal natural transformations induced by the initial and final S^1 -families are the identity because these families are constant. Therefore, an S^2 -family, viewed as an S^1 -family of S^1 -families, induces a monoidal modification of the identity natural transformation. We expect that this monoidal modification is an invariant that classifies the S^2 -parameterized family of 2+1d SPT states. See [\[91\]](#page-87-12) for a similar argument in the context of S^2 -families of 2+1d topological orders.

2. Example

As a simple example, let us consider the case of finite group symmetry G. We start by describing in detail the relevant fusion 2-categorical notions such as fiber 2-functors, monoidal natural transformations, and monoidal modifications. We will then apply Conjecture [3](#page-50-2) and find that our conjecture reproduces the correct classification of parameterized families of 2+1d SPT states.

As a fusion 2-category, a finite group symmetry G is described by $2Vec_G$, the 2-category of G-graded 2-vector spaces. Simple objects of $2Vec_G$ are labeled by elements of G and form a group G under the tensor product. The Hom categories consisting of 1-morphisms and 2-morphisms are trivial, that is, $\text{Hom}_{2\text{Vec}_G}(g, h) = \delta_{g,h} \text{Vec}$ as a 1-category. The 10-j symbols of 2Vec_G are also trivial, meaning that the symmetry group G is non-anomalous.^{[42](#page-51-1)}

⁴² An anomalous finite group symmetry G is described by $2Vec_G^{\alpha}$ with the 10-j symbols given by a 4-cocycle $[\alpha] \in$ ${\rm H}^4_{\rm gp}(G,{\rm U}(1)).$

A fiber 2-functor $(F, J, \omega):$ 2Vec $_G \to$ 2Vec consists of 1-isomorphisms $J_{g,h}: F(g) \otimes F(h) \to F(gh)$ and 2-isomorphisms $\omega_{g,h,k}$ fitting into the following diagram:

$$
F(g) \otimes F(h) \otimes F(k) \xrightarrow{J_{g,h}} F(gh) \otimes F(k)
$$

\n
$$
J_{h,k} \downarrow \qquad J_{g,h,k} \downarrow J_{gh,k}
$$

\n
$$
F(g) \otimes F(hk) \xrightarrow{J_{g,h,k}} F(ghk)
$$
\n(174)

The identity 1-morphisms are omitted in the above diagram. The coherence condition on $\omega_{g,h,k}$ is given by the equality of the following two diagrams:

$$
F(gh) \otimes F(k) \otimes F(l) \xrightarrow{J_{gh,k}} F(ghk) \otimes F(l)
$$
\n
$$
F(g) \otimes F(h) \otimes F(l) \xrightarrow{J_{h,k}} F(g) \otimes F(hk) \otimes F(l) \xrightarrow{J_{gh,k,l}} F(ghk) \otimes F(l)
$$
\n
$$
\xrightarrow{J_{h,k}} F(g) \otimes F(h) \otimes F(kl) \xrightarrow{J_{h,k,l}} F(g) \otimes F(hkl)
$$
\n
$$
(175)
$$
\n
$$
\xrightarrow{J_{h,k,l}} F(g) \otimes F(h) \otimes F(kl) \xrightarrow{J_{h,k,l}} F(g) \otimes F(hkl)
$$

$$
F(gh) \otimes F(k) \otimes F(l) \longrightarrow \begin{array}{c} F(gh) \otimes F(k) \otimes F(l) \longrightarrow J_{gh,k} & F(ghk) \otimes F(l) \\ \downarrow J_{gh,k,l} & \downarrow J_{gh,k,l} \\ \downarrow K(g) \otimes F(h) \otimes F(l) & \cong & F(gh) \otimes F(kl) \longrightarrow J_{gh,kl} \\ \downarrow K(g) \otimes F(h) \otimes F(kl) & \downarrow J_{gh,kl} \\ \downarrow K(g) \otimes F(h) \otimes F(kl) & \downarrow J_{gh,kl} \\ \end{array} \tag{176}
$$

The 2-isomorphism $\omega_{g,h,k}$ in Eq. [\(174\)](#page-52-0) can be identified with a complex number $\omega(g, h, k) \in U(1)$ because the source and target 1-morphisms of $\omega_{g,h,k}$ are invertible and hence simple.^{[43](#page-52-1)} With this identification, the equality of two diagrams in Eqs. [\(175\)](#page-52-2) and [\(176\)](#page-52-3) leads to

$$
\omega(h,k,l)\omega(g,hk,l)\omega(g,h,k) = \omega(gh,k,l)\omega(g,h,kl) \Leftrightarrow \delta\omega = 1. \tag{177}
$$

Thus, the 2-isomorphism ω associated with a fiber 2-functor defines a group 3-cocycle $\omega \in$ $Z_{\text{gp}}^3(G, U(1)).$

Given two fiber 2-functors (F, J, ω) and (F', J', ω') , a monoidal natural isomorphism (η, ν) : $(F, J, \omega) \Rightarrow (F', J', \omega')$ consists of 1-isomorphisms $\eta_g : F(g) \to F'(g)$ and 2-isomorphisms $\nu_{g,h}$ fitting into the following diagram:

$$
F(g) \otimes F(h) \xrightarrow{J_{g,h}} F(gh)
$$

\n
$$
\eta_g \otimes \eta_h \downarrow \qquad \qquad \nearrow_{g,h} \qquad \qquad \downarrow \eta_{gh}
$$

\n
$$
F'(g) \otimes F'(h) \xrightarrow{J'_{g,h}} F'(gh)
$$
\n(178)

⁴³ The space of 2-morphisms between simple 1-morphisms is isomorphic to \mathbb{C} [\[118\]](#page-88-11).

The coherence condition on ν is given by the equality of the following two diagrams:

$$
F(g) \otimes F(h) \xrightarrow{J_{n,k}} F(gh) \xrightarrow{J_{g,h,k}} F(ghk)
$$
\n
$$
F(g) \otimes F(h) \otimes F(k) \xrightarrow{J_{g,h}} F(gh) \otimes F(k) \xrightarrow{J_{g,h,k}} F(gh) \otimes F(k) \xrightarrow{J_{g,h,k}} F'(ghk)
$$
\n
$$
\xrightarrow{\eta_g \otimes \eta_h \otimes \eta_k} F'(g) \otimes F'(h) \otimes F'(k) \xrightarrow{\eta_{gh} \otimes \eta_k} F'(gh) \otimes F'(k)
$$
\n
$$
(179)
$$
\n
$$
\xrightarrow{\eta_g,h} F'(g) \otimes F'(h) \otimes F'(k) \xrightarrow{\eta_{g,h}} F'(gh) \otimes F'(k)
$$

$$
F(g) \otimes F(h) \otimes F(h) \longrightarrow F(g) \otimes F(h) \longrightarrow F(g) \otimes F(h) \longrightarrow F(g) \otimes F(h) \longrightarrow F'(g) \otimes F'(h) \longrightarrow F'(g) \otimes F'(k) \longrightarrow F'(g) \otimes F'(k)
$$
\n
$$
(180)
$$

The 2-isomorphism $\nu_{q,h}$ in Eq. [\(178\)](#page-52-4) can be identified with a complex number $\nu(g,h) \in U(1)$ just as in the case of $\omega_{g,h,k}$. With this identification, the equality of the two diagrams in Eqs. [\(179\)](#page-53-0) and [\(180\)](#page-53-1) reduces to

$$
\nu(g,h)\nu(gh,k)\omega(g,h,k)^{-1} = \nu(h,k)\nu(g,hk)\omega'(g,h,k)^{-1} \Leftrightarrow \omega' = \omega\delta\nu.
$$
 (181)

Thus, fiber 2-functors (F, J, ω) and (F', J', ω') are isomorphic to each other if and only if ω and ω' are in the same cohomology class, i.e., $[\omega] = [\omega'] \in H^3_{gp}(G, U(1)).$

Given two monoidal natural isomorphisms $(\eta, \nu), (\eta', \nu') : (F, J, \omega) \Rightarrow (F', J', \omega')$, an invertible monoidal modification $\lambda : (\eta, \nu) \Rightarrow (\eta', \nu')$ consists of 2-isomorphisms $\lambda_g : \eta_g \Rightarrow \eta'_g$ such that

$$
F(g) \otimes F(h) \xrightarrow{J_{g,h}} F(gh) \qquad F(g) \otimes F(h) \xrightarrow{J_{g,h}} F(gh)
$$

\n
$$
\eta_g \otimes \eta_h \downarrow \xrightarrow{\lambda_g \otimes \lambda_h} \eta'_g \otimes \eta'_h \xrightarrow{\gamma'} \eta'_g, \qquad \qquad \eta'_g h \qquad = \eta_g \otimes \eta_h \downarrow \xrightarrow{\nu_{g,h}} \eta_{gh} \underbrace{\xrightarrow{\lambda_{gh}} \eta_{gh}} \eta'_g h
$$

\n
$$
F'(g) \otimes F'(h) \xrightarrow{J'_{g,h}} F'(gh) \qquad F'(g) \otimes F'(h) \xrightarrow{J'_{g,h}} F'(gh)
$$
\n(182)

By identifying the 2-isomorphism λ_g with a complex number $\lambda(g) \in U(1)$, the above equation reduces to

$$
\nu'(g,h)\lambda(g)\lambda(h) = \nu(g,h)\lambda(gh) \Leftrightarrow \nu' = \nu/\delta\lambda.
$$
\n(183)

Now, let us apply Conjecture [3](#page-50-2) to $2+1d$ SPT phases with symmetry G. First of all, Eqs. [\(177\)](#page-52-5) and [\(181\)](#page-53-2) show that isomorphism classes of fiber 2-functors of $2Vec_G$ are classified by the third group cohomology $H_{gp}^3(G, U(1))$. This agrees with the well-known classification of 2+1d SPT phases [\[6,](#page-84-6) [7\]](#page-84-7).

When $\omega = \omega'$, Eq. [\(181\)](#page-53-2) reduces to $\delta \nu = 1$, which implies that a monoidal natural automorphism of a fiber 2-functor (F, J, ω) is associated with a group 2-cocycle ν . Furthermore, Eq. [\(183\)](#page-53-3) implies that natural automorphisms (η, ν) and (η', ν') are isomorphic to each other if and only if the associated 2-cocycles are equivalent. Therefore, isomorphism classes of monoidal natural automorphisms of a fiber 2-functor (F, J, ω) are classified by the second group cohomology $H_{gp}^2(G, U(1))$. This agrees with the known classification of S^1 -parameterized families of G-SPT states in 2+1d [\[15,](#page-84-0) [16,](#page-84-1) [62,](#page-86-0) [90\]](#page-87-8).

When $\nu = \nu'$, Eq. [\(183\)](#page-53-3) reduces to $\delta \lambda = 1$, meaning that an invertible modification of a natural isomorphism (η, ν) is associated with a 1-cocycle λ , i.e., a group homomorphism $\lambda : G \to U(1)$. In particular, the group of invertible modifications of the identity natural transformation of a fiber 2-functor (F, J, ω) is the first group cohomology $H^1_{gp}(G, U(1))$. Thus, Conjecture [3](#page-50-2) suggests that S^2 -parameterized families of G-SPT states in 2+1d are classified by $H_{gp}^1(G, U(1))$. This also agrees with the known classification [\[16,](#page-84-1) [62,](#page-86-0) [90\]](#page-87-8).

More generally, X-parameterized families of G -SPT states in $2+1d$ are known to be classified by [\[16,](#page-84-1) [62,](#page-86-0) [90\]](#page-87-8)

$$
\check{H}^{1}(X, \mathrm{H}_{\mathrm{gp}}^{2}(G, \mathrm{U}(1))) \oplus \check{H}^{2}(X, \mathrm{H}_{\mathrm{gp}}^{1}(G, \mathrm{U}(1))) \oplus \check{H}^{4}(X, \mathbb{Z}). \tag{184}
$$

We will reproduce this classification as a special case of our general conjecture in the next subsection.

D. General non-chiral gapped phases in 2+1d

1. Conjecture

Although we only discussed S^1 - and S^2 -parameterized families of 2+1d non-chiral SPT states above, it is natural to expect that a similar conjecture holds for more general gapped systems parameterized by more general parameter spaces. In this subsection, we consider general parameterized families of general non-chiral gapped systems in 2+1d.

As in the 1+1d case, 2+1d non-chiral gapped phases with fusion 2-category symmetry $\mathcal C$ are expected to be classified by module 2-categories over \mathcal{C} . For these phases, we propose the following conjecture on the classification of parameterized families:

Conjecture 4. X-parameterized families of 2+1d \mathcal{C} -symmetric gapped systems in a non-chiral gapped phase labeled by a C-module 2-category $\mathcal M$ are classified by the set $[X, B\underline{\underline{\mathrm{Fun}}}_{\mathcal C}(\mathcal M, \mathcal M)^\text{inv}]$ of the homotopy classes of maps from X to the classifying space of $\underline{\underline{\mathrm{Fun}}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$. Here, $\underline{\underline{\mathrm{Fun}}}_{\mathcal{C}}(\mathcal{M},\mathcal{M})^{\text{inv}}$ is a 3-group consisting of invertible objects, invertible 1-morphisms, and invertible 2-morphisms of the fusion 2-category $\text{Fun}_{\mathcal{C}}(\mathcal{M},\mathcal{M})$ of C-module 2-endofunctors of M. Equivalently, X-parameterized families of these gapped systems are classified by the non-abelian Cech cohomology $\check{H}^1(X, \underline{\underline{\mathrm{Fun}}}_{\mathcal{C}}(\mathcal{M}, \mathcal{M})^{\text{inv}})$.

We note that the above conjecture is a natural generalization of the conjecture for the $1+1d$ case, cf. Conjecture [2.](#page-45-3) A formal generalization to higher dimensions should be clear. However, we do not state the conjecture in higher dimensions here, partly because the theory of fusion n-categories and related notions for $n \geq 3$ are still under development despite significant advances in recent years [\[122–](#page-89-2)[124\]](#page-89-3).

The conjectural one-to-one correspondence between 2+1d non-chiral gapped phases with symmetry $\mathcal C$ and $\mathcal C$ -module 2-categories is supported by the explicit construction of lattice models in [\[125\]](#page-89-4). As discussed in [\[125\]](#page-89-4), one can construct a C-symmetric commuting projector model from each C-module 2-category. The gapped phase realized in this model should be non-chiral because the Hamiltonian is given by the sum of local commuting projectors. Thus, the lattice models in [\[125\]](#page-89-4) relate C-module 2-categories to C-symmetric non-chiral gapped phases in $2+1d$. For example, when

the symmetry C is trivial, i.e., when $C = 2Vec$, an indecomposable module 2-category over C is given by the 2-category $Mod(\mathcal{D})$ of module categories over a fusion 1-category \mathcal{D} [\[126\]](#page-89-5). The commuting projector model of [\[125\]](#page-89-4) constructed from $Mod(\mathcal{D})$ is the Levin-Wen model [\[83\]](#page-87-2) constructed from \mathcal{D} . This model exhibits a topological order whose anyon contents are described by the Drinfeld center $Z(\mathcal{D})$ of \mathcal{D} . Since the Drinfeld center of a fusion category describes the most general non-chiral topological order in 2+1d [\[84,](#page-87-4) [127\]](#page-89-6), the aforementioned one-to-one correspondence is established at least in the case of the trivial symmetry.^{[44](#page-55-0)}

In what follows, we provide several examples that verify the classification of parameterized families stated in Conjecture [4](#page-54-1)

2. Examples

Example 1: trivial phase without symmetry. The simplest example is the parameterized families of trivial states without symmetry. From the point of view of module 2-categories, the trivial phase corresponds to the regular module $\mathcal{M} = 2$ Vec over $\mathcal{C} = 2$ Vec. The associated functor 2-category Fun_{2Vec}(2Vec, 2Vec) is also equivalent to 2Vec. The 3-group $\underline{\underline{Fun}}_{2Vec}(2Vec, 2Vec)^{inv}$ = 2Vec^{inv} consists only of the 2-form group U(1). We denote this 3-group by U(1)^[2], where the superscript specifies the form degree. Applying Conjecture 4 to this example suggests that Xparameterized families of the trivial states without symmetry are classified by

$$
\check{H}^{1}(X, \underline{\mathbf{2Vec}}^{\text{inv}}) = \check{H}^{1}(X, \mathbf{U}(1)^{[2]}) = [X, K(\mathbb{Z}, 4)] = \check{H}^{4}(X, \mathbb{Z}),
$$
\n(185)

where $K(\mathbb{Z}, 4) = B^3U(1)$ is the fourth Eilenberg-MacLance space of \mathbb{Z} . Here, U(1) is supposed to have the standard topology rather than the discrete one. The above result agrees with the known classification of parameterized families of trivial invertible states $[11-13]$ $[11-13]$ via higher Berry phases [\[90,](#page-87-8) [109,](#page-88-13) [115,](#page-88-14) [129,](#page-89-7) [130\]](#page-89-8). As in 1+1d, this classification originates from the fact that the "trivial" fusion 2-category 2Vec captures a U(1) 2-form symmetry, which exists in any quantum systems in $2+1d.$

Example 2: SPT phases with G symmetry. Another simple example is the parameterized families of SPT states with finite group symmetry G . For simplicity, here we only consider G-symmetric trivial states. 45 From the point of view of module 2-categories, the trivial G-symmetric phase corresponds to a $2Vec$ -module 2-category 2Vec. The associated functor 2category is $\text{Fun}_{2\text{Vec}_G}(2\text{Vec}, 2\text{Vec}) \cong 2\text{Rep}(G)$, the category of 2-representations of G [\[126\]](#page-89-5). Invertible objects and invertible 1-morphisms of $2\text{Rep}(G)$ form $H^2_{gp}(G, U(1))$ and $H^1_{gp}(G, U(1))$ respectively, see e.g. [\[131–](#page-89-9)[133\]](#page-89-10) for a detailed description of $2\text{Rep}(G)$ for physicists. Thus, the 3-group $2\text{Rep}(G)$ ^{inv} consists of a 0-form group $H^2_{gp}(G, U(1))$, a 1-form group $H^1_{gp}(G, U(1))$, and a 2-form group $U(1)$. The 3-group structure on $2\text{Rep}(G)^{inv}$ is trivial, meaning that the actions of the 0-form group on the 1-form and 2-form groups are trivial, and the Postnikov classes are also trivial. 46 Therefore,

 44 General 2+1d topological orders up to invertible ones are believed to be classified by modular tensor categories, which are in one-to-one correspondence with Lagrangian algebras in the Drinfeld center $Z(2\text{Vec})$ of 2Vec [\[128\]](#page-89-11).

⁴⁵ The classification of parameterized families should not depend on the choice of an SPT phase because stacking a G-SPT phase is an invertible operation.

⁴⁶ The invertible objects and invertible 1-morphisms of $2\text{Rep}(G)$ are known as gauged SPT defects in the context of finite group gauge theories [\[134,](#page-89-12) [135\]](#page-89-13). These gauged SPT defects from an anomaly-free 2-group symmetry $H_{\rm gp}^2(G, \mathrm{U}(1))^{[0]} \times H_{\rm gp}^1(G, \mathrm{U}(1))^{[1]}$ with the trivial action of the 0-form group and the trivial Postnikov class [\[134\]](#page-89-12). Put differently, $2\text{Rep}(G)$ ^{inv} has the trivial 3-group structure. The authors thank Ryohei Kobayashi for discussions on this point.

Conjecture [4](#page-54-1) suggests that X-parameterized families of trivial states with G symmetry are classified by

$$
\check{H}^{1}(X, \underbrace{2\mathrm{Rep}(G)}^{\mathrm{inv}}) = \check{H}^{1}(X, \mathrm{H}_{\mathrm{gp}}^{2}(G, \mathrm{U}(1))^{[0]} \times \mathrm{H}_{\mathrm{gp}}^{1}(G, \mathrm{U}(1))^{[1]} \times \mathrm{U}(1)^{[2]})
$$
\n
$$
= \check{H}^{1}(X, \mathrm{H}_{\mathrm{gp}}^{2}(G, \mathrm{U}(1))) \oplus \check{H}^{2}(X, \mathrm{H}_{\mathrm{gp}}^{1}(G, \mathrm{U}(1))) \oplus \check{H}^{4}(X, \mathrm{U}(1)).
$$
\n(186)

This agrees with the known classification of parameterized families of $2+1d$ G-SPT states [\[14,](#page-84-5) [16,](#page-84-1) [62,](#page-86-0) [90\]](#page-87-8).

Example 3: SPT phases with $2\text{Rep}(G)$ **symmetry.** Yet another example is the parameterized families of SPT states with $2\text{Rep}(G)$ symmetry. As opposed to the case of G symmetry, the set of SPT phases with $2Rep(G)$ symmetry does not have a natural group structure in general. As a result, the classification of parameterized families depends on the phases. Here, for simplicity, we consider families of $2\text{Rep}(G)$ -symmetric trivial states, which are obtained by gauging G -symmetry-breaking states.

From the point of view of module 2-categories, the trivial phase with $2\text{Rep}(G)$ symmetry corresponds to a $2\text{Rep}(G)$ -module 2-category 2Vec. The associated functor 2-category is $\text{Fun}_{2\text{Rep}(G)}(2\text{Vec}, 2\text{Vec}) \cong 2\text{Vec}_G$ [\[126\]](#page-89-5). Thus, the 3-group $\underline{\underline{\text{Fun}}}_{2\text{Rep}(G)}(2\text{Vec}, 2\text{Vec})^{\text{inv}}$ consists only of the 0-form group G and the 2-form group $U(1)$. The 3-group structure is again trivial. Therefore, Conjecture [4](#page-54-1) suggests that X-parameterized families of trivial states with $2\text{Rep}(G)$ symmetry are classified by

$$
\check{H}^{1}(X, \underline{2\mathrm{Vec}_{G}}^{\mathrm{inv}}) = \check{H}^{1}(X, G^{[0]} \times U(1)^{[2]}) = \check{H}^{1}(X, G) \oplus \check{H}^{4}(X, \mathbb{Z}). \tag{187}
$$

We note that parameterized families of G -SSB states in $2+1d$ are also classified by the same cohomology group because G -SSB states and $2\text{Rep}(G)$ -symmetric trivial states are related by gauging.

Example 4: topological orders without symmetry. Finally, we consider the parameterized families of topologically ordered states with the trivial symmetry $C = 2Vec$. As already mentioned, an indecomposable module 2-category over 2Vec is given by the 2-category $Mod(\mathcal{D})$ of module categories over a fusion 1-category \mathcal{D} [\[126\]](#page-89-5). The associated functor 2-category is $Fun_{2Vec}(Mod(\mathcal{D}), Mod(\mathcal{D})) \cong Mod(Z(\mathcal{D}))$ [\[126,](#page-89-5) Example 5.3.7], where $Z(\mathcal{D})$ denotes the Drinfeld center of D. The 3-group consisting of invertible objects, invertible 1-morphisms, and invertible 2-morphisms of $Mod(Z(\mathcal{D}))$ is known as the categorical Picard 2-group $Pic(Z(\mathcal{D}))$ of $Z(\mathcal{D})$ [\[93\]](#page-87-10). Thus, we have

$$
\underline{\underline{\text{Fun}}}_{2\text{Vec}}(\text{Mod}(\mathcal{D}), \text{Mod}(\mathcal{D}))^{\text{inv}} = \underline{\underline{\text{Pic}}}(Z(\mathcal{D})).\tag{188}
$$

Therefore, Conjecture [4](#page-54-1) suggests that X-parameterized families of gapped systems in a topologically ordered phase $Z(\mathcal{D})$ are classified by the set $[X, B\underline{\underline{\mathrm{Pic}}} (Z(\mathcal{D}))] = \check{H}^1(X, \underline{\underline{\mathrm{Pic}}} (Z(\mathcal{D})))$. This agrees with the conjecture in [\[91\]](#page-87-12).

VII. FUTURE DIRECTIONS

In this paper, we developed the analysis of interface modes and Thouless pumps in C -SPT phases within the framework of MPS representations. Specifically, by investigating the properties of interface modes, we generalized the bulk-boundary correspondence for usual SPT phases, and by investigating Thouless pumps, we revealed pumping phenomena with respect to non-abelian charges. In addition, through categorical considerations, we proposed several conjectures for the classification of parameterized families of general gapped phases in both 1+1d and higher dimensions. The next task to complete is verifying these conjectures using tensor networks and constructing models that realize their classifications. We leave them for future studies.

ACKNOWLEDGMENTS

We thank Ryohei Kobayashi and Yosuke Kubota for useful discussions. We are grateful to the long-term workshop YITP-T-23-01 held at Yukawa Institute for Theoretical Physics, Kyoto University, where a part of this work was done. S.O. is supported by JSPS KAKENHI Grant Number 23KJ1252 and 24K00522.

Appendix A: Tambara-Yamagami categories

In this appendix, we review the Tambara-Yamagami categories [\[136\]](#page-89-14). A Tambara-Yamagami category TY(A, χ, ϵ) is specified by a triple (A, χ, ϵ) , where A is a finite abelian group, $\chi : A \times A \rightarrow$ U(1) is a symmetric non-degenerate bicharacter of A, and $\epsilon \in \{+1, -1\}$ is a sign. The set of simple objects consists of invertible objects labeled by elements of A and a single non-invertible object D. The fusion rules of these simple objects are given by

$$
a \otimes b = ab, \quad a \otimes D = D \otimes a = D, \quad D \otimes D = \bigoplus_{a \in A} a,
$$
 (A1)

where a and b represent invertible objects labeled by group elements $a, b \in A$. The above fusion rules imply that the quantum dimension of the non-invertible object D is given by $\dim(D) = \sqrt{|A|}$, while the quantum dimensions of invertible objects are all 1 . The F -symbols in a specific gauge can be written as

$$
(F_D^{aDb})_{DD} = (F_b^{DaD})_{DD} = \chi(a, b), \quad (F_D^{DDD})_{a,b} = \frac{\epsilon}{\sqrt{|A|}} \chi(a, b)^{-1}.
$$
 (A2)

The other F-symbols are all trivial.

Simple examples of our interest are $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami categories TY($\mathbb{Z}_2 \times \mathbb{Z}_2, \chi, \epsilon$). When $A = \mathbb{Z}_2 \times \mathbb{Z}_2$, there are two choices for a symmetric non-degenerate bicharacter χ , which we denote by χ_+ and χ_- . These bicharacters are explicitly given by [\[136\]](#page-89-14)

$$
\chi_{\pm}((1,0),(1,0)) = \chi_{\pm}((0,1),(0,1)) = \pm 1, \quad \chi_{\pm}((1,0),(0,1)) = \mp 1,
$$
 (A3)

where $(1,0)$ and $(0,1)$ are generators of $\mathbb{Z}_2 \times \mathbb{Z}_2$. The above equation uniquely determines $\chi_{\pm}(a,b)$ for general $a, b \in \mathbb{Z}_2 \times \mathbb{Z}_2$ because χ_{\pm} is a symmetric bicharacter. Corresponding to different choices for χ and ϵ , there are four inequivalent $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami categories. It is known that $TY(\mathbb{Z}_2\times\mathbb{Z}_2,\chi_+,+1)$ is equivalent to the representation category $\text{Rep}(D_8)$ of the dihedral group D_8 of order 8. Specifically, invertible and non-invertible simple objects of $TY(\mathbb{Z}_2\times\mathbb{Z}_2,\chi_+,+1)$ correspond to one-dimensional and two-dimensional irreducible representations of Rep(D_8). Similarly, TY($\mathbb{Z}_2\times$ $\mathbb{Z}_2, \chi_+, -1$) is equivalent to the representation category Rep(Q_8) of the quaternion group Q_8 , and $TY(\mathbb{Z}_2 \times \mathbb{Z}_2, \chi_-, +1)$ is equivalent to the representation category Rep(H₈) of an eight-dimensional

$\epsilon = 1$	$\epsilon = -1$
	$\chi = \chi_+ \begin{vmatrix} Rep(D_8) \\ Rep(Q_8) \end{vmatrix}$ Rep (Q_8) $\chi = \chi_- \begin{vmatrix} Rep(H_8) \end{vmatrix}$ anomalous

TABLE I. There are four $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami categories depending on the choice of χ and ϵ . Three of them are non-anomalous, while the other one is anomalous. Ref. $[138]$ showed that $\text{Rep}(D_8)$ admits three fiber functors, while $\text{Rep}(Q_8)$ and $\text{Rep}(H_8)$ admit only one fiber functor.

semisimple Hopf algebra H_8 known as the Kac-Paljutkin algebra [\[137\]](#page-89-16). The above three $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami categories are non-anomalous because they are equivalent to representation categories of semisimple Hopf algebras. Thus, these fusion categories admit SPT phases. On the other hand, the remaining one $TY(\mathbb{Z}_2 \times \mathbb{Z}_2, \chi_-, -1)$ is anomalous and hence does not admit SPT phases. See Table [I](#page-58-0) for a summary of four $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami categories.

Appendix B: Fiber functors of Tambara-Yamagami categories

In this appendix, we review fiber functors of Tambara-Yamagami categories following [\[138\]](#page-89-15). The fiber functors and the corresponding L-symbols that we write down below are derived in a specific gauge where the F -symbols of the Tambara-Yamagami categories are given by Eq. $(A2)$.

We first recall the classification of fiber functors of $TY(A, \chi, \epsilon)$ [\[138\]](#page-89-15), see also [\[32\]](#page-85-1). Proposition 3.2 of [\[138\]](#page-89-15) claims that isomorphism classes of fiber functors of $TY(A, \chi, \epsilon)$ are in one-to-one correspondence with equivalence classes of triples (σ, ξ, ν) , where $\sigma : A \to A$ is an involutive automorphism of $A, \xi : A \times A \rightarrow U(1)$ is a 2-cocycle on A, and $\nu : A \rightarrow U(1)$ is a $U(1)$ -valued function on A, such that

$$
\chi(a,b) = \xi(a,\sigma(b))/\xi(\sigma(b),a),
$$

\n
$$
\nu(a)\nu(b)/\nu(ab) = \xi(a,b)/\xi(\sigma(b),\sigma(a)),
$$

\n
$$
\nu(a)\nu(\sigma(a)) = 1,
$$

\n
$$
\sum_{\substack{a \in A \\ \text{s.t. } \sigma(a) = a}} \nu(a) = \epsilon \sqrt{|A|}.
$$
\n(B1)

Here, two triples (σ, ξ, ν) and (σ', ξ', ν') are said to be equivalent if and only if

$$
\sigma' = \sigma, \quad \xi'(a, b) = \xi(a, b)\zeta(a)\zeta(b)/\zeta(ab), \quad \nu'(a) = \nu(a)\zeta(a)/\zeta(\sigma(a))
$$
 (B2)

for some U(1)-valued function $\zeta : A \to U(1)$. The 2-cocycle ξ can always be normalized so that $\xi(q,1) = \xi(1,h) = 1$. In what follows, we suppose that ξ is a normalized 2-cocycle. We note that ξ has to be non-trivial due to the first equality of Eq. [\(B1\)](#page-58-1). In particular, the non-degeneracy of χ implies that the twisted group algebra $\mathbb{C}[A]^\xi$ is simple, i.e., it is isomorphic to a matrix algebra $M_{\sqrt{|A|}}(\mathbb{C}).$

Let us write down a fiber functor in terms of the triple (σ, ξ, ν) . A fiber functor F of a Tambara-Yamagami category $TY(A, \chi, \epsilon)$ is characterized by the natural isomorphism

$$
J_{a,b}: F(a)\otimes F(b)\to F(ab), \quad J_{a,D}: F(a)\otimes F(D)\to F(D),
$$

\n
$$
J_{D,a}: F(D)\otimes F(a)\to F(D), \quad J_{D,D}: F(D)\otimes F(D)\to \bigoplus_{a\in A} F(a).
$$
 (B3)

Here, $F(a) \cong \mathbb{C}$ is a one-dimensional vector space and $F(D) \cong \mathbb{C}$ $\sqrt{|A|}$ is a $\sqrt{|A|}$ -dimensional vector space. Once we choose bases $\{v_a\}$ and $\{v_D^i \mid i = 1, 2, \cdots, \sqrt{|A|}\}$ of $F(a)$ and $F(D)$, the isomorphisms in Eq. [\(B3\)](#page-58-2) can be represented by L-symbols $\{L^{a,b}, L^{a,D}_{i,j}, L^{D,a}_{i,j}, L^{D,D}_{(i,j),a} \mid a,b \in A, i,j = 1\}$ $1, 2, \cdots, \sqrt{|A|}\},\$ that is,

$$
J_{a,b}(v_a \otimes v_b) = L^{a,b}v_{a \otimes b}, \qquad J_{a,D}(v_a \otimes v_D^i) = \sum_j L^{a,D}_{i,j} v_D^j,
$$

$$
J_{D,a}(v_D^i \otimes v_a) = \sum_j L^{D,a}_{i,j} v_D^j, \quad J_{D,D}(v_D^i \otimes v_D^j) = \bigoplus_{a \in A} L^{D,D}_{(i,j),a} v_a.
$$
 (B4)

The L-symbols in the above equation are related to the triple (σ, ξ, ν) as follows [\[138\]](#page-89-15):

$$
L^{a,b} = \xi(a,b), \qquad \sum_{j} L^{a,D}_{i,j} L^{b,D}_{j,k} = \xi(a,b) L^{ab,D}_{i,k},
$$

\n
$$
L^{D,a}_{i,j} = \nu(a) L^{\sigma(a),D}_{i,j}, \qquad L^{D,D}_{(i,j),a} = \xi(a^{-1},a)^{-1} \sum_{k} L^{a^{-1},D}_{i,k} \gamma_{kj}.
$$
\n(B5)

Here, γ_{kj} in the last equation is the (k, j) -component of a non-degenerate $\sqrt{|A|} \times \sqrt{|A|}$ matrix that satisfies

$$
L^{a,D}\gamma = \nu(a)\gamma(L^{\sigma(a),D})^T,\tag{B6}
$$

where $L^{a,D}$ is the $\sqrt{|A|} \times \sqrt{|A|}$ matrix whose (i, j) -component is the L-symbol $L^{a,D}_{i,j}$, and $(L^{a,D})^T$ is its transpose. The matrix γ that satisfies the above equation is unique up to scalar multiplication [\[138\]](#page-89-15).

The second equation on the first line of Eq. [\(B5\)](#page-59-0) implies that $L^{a,D}$ is the representation matrix of a $\sqrt{|A|}$ -dimensional representation of the twisted group algebra $\mathbb{C}[A]^{\xi}$. Such a representation is unique up to unitary equivalence because $\mathbb{C}[A]^\xi$ is a simple algebra as already mentioned. Thus, $L_{i,j}^{a,D}$ is uniquely determined up to the change of the basis of $F(D)$.

Appendix C: L-symbols for fiber functors of $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami categories

In this appendix, we write down the L-symbols for fiber functors of non-anomalous $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami categories, i.e., $\text{Rep}(D_8)$, $\text{Rep}(Q_8)$, and $\text{Rep}(H_8)$.

1. Rep (D_8) symmetry

We first consider the case of $TY(\mathbb{Z}_2 \times \mathbb{Z}_2, \chi_+, +1) \cong \text{Rep}(D_8)$. As we reviewed in Appendix [B,](#page-58-3) fiber functors of TY(A, χ, ϵ) are in one-to-one correspondence with triples (σ, ξ, ν) that satisfies Eq. [\(B1\)](#page-58-1). When $(A, \chi, \epsilon) = (\mathbb{Z}_2 \times \mathbb{Z}_2, \chi_+, +1)$, there are three inequivalent solutions to Eq. (B1) [\[32,](#page-85-1) [138\]](#page-89-15). These solutions share the same ξ and σ , which are given by^{[47](#page-59-1)}

$$
\xi(a,b) = (-1)^{a_2 b_1}, \quad \sigma(a) = a,\tag{C1}
$$

⁴⁷ One can equally choose a different representative ξ of a non-trivial cohomology class $[\xi] \neq 0 \in H_{gp}^2(\mathbb{Z}_2 \times \mathbb{Z}_2, U(1)) \cong$ \mathbb{Z}_2 .

where $a = (a_1, a_2), b = (b_1, b_2) \in \mathbb{Z}_2 \times \mathbb{Z}_2 = \{(0, 0), (0, 1), (1, 0), (1, 1)\}.$ The three solutions are distinguished by different ν 's, which we denote by ν_1 , ν_2 , and ν_3 . Concretely, these ν 's are given by

$$
\nu_1(0,1) = \nu_2(1,0) = \nu_3(1,1) = -1 \tag{C2}
$$

and $\nu_i(a) = +1$ otherwise.

Since all of the three fiber functors have the same ξ and σ , some of the L-symbols are independent of fiber functors. Specifically, $L^{a,b}$, $L^{a,D}$, and their inverses are given regardless of ν as

$$
L^{a,b} = \overline{L}^{a,b} = (-1)^{a_2 b_1}, \qquad L^{a,D}_{i,j} = (-1)^{a_1 a_2} \overline{L}^{a,D}_{i,j} = \begin{cases} \delta_{ij} & a = (0,0), \\ Z_{ij} & a = (0,1), \\ X_{ij} & a = (1,0), \\ -iY_{ij} & a = (1,1), \end{cases}
$$
(C3)

where X, Y, and Z denote Pauli matrices. On the other hand, $L^{D,a}$, $L^{D,D}$ and their inverses depend on the choice of ν .

• When $\nu = \nu_1$, Eqs. [\(B6\)](#page-59-2) and [\(C3\)](#page-60-0) imply $\gamma = X$, and hence we have

$$
L_{i,j}^{D,a} = \begin{cases} \delta_{ij} & a = (0,0), \\ -Z_{ij} & a = (0,1), \\ X_{ij} & a = (1,0), \\ -iY_{ij} & a = (1,1), \end{cases} \qquad L_{(i,j),a}^{D,D} = \begin{cases} X_{ij} & a = (0,0), \\ iY_{ij} & a = (0,1), \\ \delta_{ij} & a = (1,0), \\ Z_{ij} & a = (1,1). \end{cases} \qquad (C4)
$$

• When $\nu = \nu_2$, Eqs. [\(B6\)](#page-59-2) and [\(C3\)](#page-60-0) imply $\gamma = Z$, and hence we have

$$
L_{i,j}^{D,a} = \begin{cases} \delta_{ij} & a = (0,0), \\ Z_{ij} & a = (0,1), \\ -X_{ij} & a = (1,0), \\ -iY_{ij} & a = (1,1), \end{cases} \qquad L_{(i,j),a}^{D,D} = \begin{cases} Z_{ij} & a = (0,0), \\ \delta_{ij} & a = (0,1), \\ -iY_{ij} & a = (1,0), \\ -X_{ij} & a = (1,1). \end{cases} \qquad (C5)
$$

• When $\nu = \nu_3$, Eqs. [\(B6\)](#page-59-2) and [\(C3\)](#page-60-0) imply $\gamma = I$, and hence we have

$$
L_{i,j}^{D,a} = \begin{cases} \delta_{ij} & a = (0,0), \\ Z_{ij} & a = (0,1), \\ X_{ij} & a = (1,0), \\ iY_{ij} & a = (1,1), \end{cases} \qquad L_{(i,j),a}^{D,D} = \begin{cases} \delta_{ij} & a = (0,0), \\ Z_{ij} & a = (0,1), \\ X_{ij} & a = (1,0), \\ iY_{ij} & a = (1,1). \end{cases} \qquad (C6)
$$

In all cases, the inverses of $L^{D,a}$ and $L^{D,D}$ are given by

$$
\overline{L}_{i,j}^{D,a} = (-1)^{a_1 a_2} L_{i,j}^{D,a}, \qquad \overline{L}_{a,(i,j)}^{D,D} = \frac{1}{2} L_{(i,j),a}^{D,D}.
$$
 (C7)

2. Rep (Q_8) symmetry

The $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami category TY($\mathbb{Z}_2 \times \mathbb{Z}_2$, $\chi_+, -1$) \cong Rep(Q_8) has only one fiber functor, which is the forgetful functor. The corresponding solution to Eq. [\(B1\)](#page-58-1) is given by

$$
\xi(a,b) = (-1)^{a_2 b_1}, \quad \sigma(a) = a, \quad \nu(a) = (-1)^{a_1 + a_2 + a_1 a_2}, \tag{C8}
$$

where $a = (a_1, a_2), b = (b_1, b_2) \in \mathbb{Z}_2 \times \mathbb{Z}_2$. The L-symbols $L^{a,b}$, $L^{a,D}_{i,j}$, and their inverses are given by Eq. [\(C3\)](#page-60-0). The other L-symbols $L^{D,a}$ and $L^{D,D}$ are given by

$$
L_{i,j}^{D,a} = \begin{cases} \delta_{ij} & a = (0,0), \\ -Z_{ij} & a = (0,1), \\ -X_{ij} & a = (1,0), \\ iY_{ij} & a = (1,1), \end{cases} \qquad L_{(i,j),a}^{D,D} = \begin{cases} -iY_{ij} & a = (0,0), \\ -X_{ij} & a = (0,1), \\ Z_{ij} & a = (1,0), \\ \delta_{ij} & a = (1,1). \end{cases} \qquad (C9)
$$

Here, we used $\gamma = -iY$, which is the unique solution to Eq. [\(B6\)](#page-59-2) up to scalar. The inverses of $L^{D,a}$ and $L^{D,D}$ are given by

$$
\overline{L}_{i,j}^{D,a} = (-1)^{a_1 a_2} L_{i,j}^{D,a}, \qquad \overline{L}_{a,(i,j)}^{D,D} = \frac{1}{2} L_{(i,j),a}^{D,D}.
$$
\n(C10)

3. Rep (H_8) symmetry

The $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami category TY($\mathbb{Z}_2 \times \mathbb{Z}_2$, $\chi_-, +1) \cong \text{Rep}(H_8)$ has only one fiber functor, which is the forgetful functor. The corresponding solution to Eq. $(B1)$ is given by

$$
\xi(a,b) = (-1)^{a_2 b_1}, \quad \sigma(a) = (a_2, a_1), \quad \nu(a) = 1,
$$
\n(C11)

where $a = (a_1, a_2), b = (b_1, b_2) \in \mathbb{Z}_2 \times \mathbb{Z}_2$. The L-symbols $L^{a,b}$, $L^{a,D}_{i,j}$, and their inverses are given by Eq. [\(C3\)](#page-60-0). The other L-symbols $L^{D,a}$ and $L^{D,D}$ are given by

$$
L_{i,j}^{D,a} = \begin{cases} \delta_{ij} & a = (0,0), \\ X_{ij} & a = (0,1), \\ Z_{ij} & a = (1,0), \\ -iY_{ij} & a = (1,1), \end{cases} \qquad L_{(i,j),a}^{D,D} = \begin{cases} \frac{1}{\sqrt{2}}(X+Z)_{ij} & a = (0,0), \\ \frac{1}{\sqrt{2}}(1+iY)_{ij} & a = (0,1), \\ \frac{1}{\sqrt{2}}(1-iY)_{ij} & a = (1,0), \\ -\frac{1}{\sqrt{2}}(X-Z)_{ij} & a = (1,1). \end{cases} (C12)
$$

Here, we used $\gamma = \frac{1}{\sqrt{2}}$ $\overline{Z}(\overline{X} + \overline{Z})$, which is the unique solution to Eq. [\(B6\)](#page-59-2) up to scalar. The inverses of $L^{D,a}$ and $L^{D,D}$ are given by

$$
\overline{L}_{i,j}^{D,a} = (-1)^{a_1 a_2} L_{i,j}^{D,a}, \qquad \overline{L}_{a,(i,j)}^{D,D} = \frac{1}{2} L_{(i,j),a}^{D,D}.
$$
\n(C13)

Appendix D: Interface algebras for $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami categories

In this appendix, we study representations of symmetry algebras acting on the interface of SPT phases with non-anomalous $\mathbb{Z}_2 \times \mathbb{Z}_2$ Tambara-Yamagami symmetries, i.e, Rep(D_8), Rep(Q_8), and

 $Rep(H_8)$ symmetries. For any of these symmetries, the symmetry algebra A acting on the interface $\mathcal I$ of two SPT phases SPT_1 and SPT_2 is spanned by^{[48](#page-62-0)}

$$
\{\mathcal{O}_a^{\mathcal{I}}, (\mathcal{O}_D^{\mathcal{I}})_{ij} \mid a \in \mathbb{Z}_2 \times \mathbb{Z}_2, \ i, j = 1, 2\}.
$$
 (D1)

In particular, $\mathcal A$ is an eight-dimensional algebra. The multiplication of symmetry operators is given by Eq. [\(96\)](#page-28-0), i.e.,

$$
\mathcal{O}_{a}^{\mathcal{I}} \mathcal{O}_{b}^{\mathcal{I}} = (L_{1})^{a,b} (\overline{L_{2}})^{a,b} \mathcal{O}_{ab}^{\mathcal{I}}, \qquad \mathcal{O}_{a}^{\mathcal{I}} (\mathcal{O}_{D}^{\mathcal{I}})_{ij} = \sum_{k,k'} (L_{1})^{a,D}_{i,k} (\overline{L_{2}})^{a,D}_{k',j} (\mathcal{O}_{D}^{\mathcal{I}})_{kk'},
$$

$$
(\mathcal{O}_{D}^{\mathcal{I}})_{ij} \mathcal{O}_{a}^{\mathcal{I}} = \sum_{k,k'} (L_{1})^{D,a}_{i,k} (\overline{L_{2}})^{D,a}_{k',j} (\mathcal{O}_{D}^{\mathcal{I}})_{kk'}, \quad (\mathcal{O}_{D}^{\mathcal{I}})_{ij} (\mathcal{O}_{D}^{\mathcal{I}})_{kl} = \sum_{a \in \mathbb{Z}_{2} \times \mathbb{Z}_{2}} (L_{1})^{D,D}_{(i,k),a} (\overline{L_{2}})^{D,D}_{a,(j,l)} \mathcal{O}_{a}^{\mathcal{I}},
$$
^(D2)

where L_1 and L_2 are the L-symbols associated with SPT_1 and SPT_2 respectively. Since $L^{a,b}$ and $L^{a,D}$ are given by Eq. [\(C3\)](#page-60-0) regardless of SPT phases, the first line of the above equation reduces to

$$
\mathcal{O}_a^{\mathcal{I}} \mathcal{O}_b^{\mathcal{I}} = \mathcal{O}_{ab}^{\mathcal{I}}, \qquad \mathcal{O}_a^{\mathcal{I}} (\mathcal{O}_D^{\mathcal{I}})_{ij} = \begin{cases}\n(\mathcal{O}_D^{\mathcal{I}})_{ij} & a = (0,0), \\
(Z\mathcal{O}_D^{\mathcal{I}}Z)_{ij} & a = (0,1), \\
(X\mathcal{O}_D^{\mathcal{I}}X)_{ij} & a = (1,0), \\
(Y\mathcal{O}_D^{\mathcal{I}}Y)_{ij} & a = (1,1).\n\end{cases}
$$
\n(D3)

On the other hand, the second line of Eq. [\(D2\)](#page-62-1) depends on SPT phases. Since the interface algebra A should be semisimple, all the irreducible representations of A are obtained by the direct sum decomposition of A.

1. Rep (D_8) symmetry

a. Self-interface

The $\text{Rep}(D_8)$ symmetry has three inequivalent SPT phases, which are distinguished by different values of ν , see Eq. [\(C2\)](#page-60-1). We denote the Rep(D_8) SPT phase associated with triple (σ, ξ, ν) by SPT_{ν} . The interface between SPT_{ν_i} and SPT_{ν_j} is denoted by $\mathcal{I}_{\nu_i,\nu_j}$, and the associated interface algebra is denoted by $\mathcal{A}_{\nu_i,\nu_j}$. In what follows, we write down the symmetry algebra $\mathcal{A}_{\nu,\nu}$ acting on the self-interface $\mathcal{I}_{\nu,\nu}$ and enumerate all irreducible representations of $\mathcal{A}_{\nu,\nu}$. When no confusion can arise, we will omit the subscripts of $\mathcal{I}_{\nu_i,\nu_j}$ and $\mathcal{A}_{\nu_i,\nu_j}$.

• When $SPT_1 = SPT_2 = SPT_{\nu_1}$, the multiplication of symmetry operators on the second line of Eq. $(D2)$ is given by

$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}\mathcal{O}_a^{\mathcal{I}} = \mathcal{O}_a^{\mathcal{I}}(\mathcal{O}_D^{\mathcal{I}})_{ij},
$$

\n
$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}(\mathcal{O}_D^{\mathcal{I}})_{kl} = \frac{1}{2}X_{ik}X_{jl}\mathcal{O}_{(0,0)}^{\mathcal{I}} - \frac{1}{2}Y_{ik}Y_{jl}\mathcal{O}_{(0,1)}^{\mathcal{I}} + \frac{1}{2}\delta_{ik}\delta_{jl}\mathcal{O}_{(1,0)}^{\mathcal{I}} + \frac{1}{2}Z_{ik}Z_{jl}\mathcal{O}_{(1,1)}^{\mathcal{I}}.
$$
 (D4)

⁴⁸ In Sec. [IV,](#page-27-0) a symmetry operator acting on the interface \mathcal{I} was written as $(\hat{\mathcal{O}}_x^{\mathcal{I}})_{ij}$. In this appendix, we omit the hat.

$\mathcal{I} = \mathcal{I}_{\nu_1,\nu_1} \begin{vmatrix} R_1^+ & R_1^- & R_2^+ & R_2^- & R_3^+ & R_3^- & R_4^+ & R_4^- \end{vmatrix}$				
$\begin{array}{c ccccccccc} \hline \mathcal{O}^{\mathcal{I}}_{(0,0)} & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ \mathcal{O}^{\mathcal{I}}_{(0,1)} & 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 \\ \mathcal{O}^{\mathcal{I}}_{(1,0)} & 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 & -1 \\ \mathcal{O}^{\mathcal{I}}_{(1,1)} & 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 \\ \mathcal{O}^{\mathcal{I}}_{(1,1)} & 1 & $				

TABLE II. Irreducible representations of the self-interface algebra $\mathcal A$ for the Rep(D_8) SPT phase SPT_{ν_1}. All irreducible representations are one-dimensional and form a group D_8 under the multiplication.

The multiplications $(D3)$ and $(D4)$ imply that the interface algebra $\mathcal A$ is commutative. Thus, due to the Artin-Wedderburn theorem, the eight-dimensional algebra A can be decomposed into a direct sum of eight one-dimensional subalgebras as

$$
\mathcal{A} = \bigoplus_{s=\pm} \bigoplus_{1 \le i \le 4} \mathbb{C}e_i^s, \tag{D5}
$$

where the direct summands e_i^s are idempotents of A, i.e., they are elements of A such that $e_i^s e_j^t = \delta_{ij} \delta_{st} e_i^s$. The explicit forms of these idempotents are given by

$$
e_1^{\pm} = \frac{1}{2}(e_1 \pm e_1'), \quad e_2^{\pm} = \frac{1}{2}(e_2 \pm ie_2'), \quad e_3^{\pm} = \frac{1}{2}(e_3 \pm e_3'), \quad e_4^{\pm} = \frac{1}{2}(e_4 \pm ie_4'),
$$
 (D6)

where e_i and e'_i for $i = 1, 2, 3, 4$ are defined as follows:

$$
e_1 = \frac{1}{4} \left(\mathcal{O}_{(0,0)}^{\mathcal{I}} + \mathcal{O}_{(0,1)}^{\mathcal{I}} + \mathcal{O}_{(1,0)}^{\mathcal{I}} + \mathcal{O}_{(1,1)}^{\mathcal{I}} \right), \quad e_2 = \frac{1}{4} \left(\mathcal{O}_{(0,0)}^{\mathcal{I}} + \mathcal{O}_{(0,1)}^{\mathcal{I}} - \mathcal{O}_{(1,0)}^{\mathcal{I}} - \mathcal{O}_{(1,1)}^{\mathcal{I}} \right),
$$

\n
$$
e_3 = \frac{1}{4} \left(\mathcal{O}_{(0,0)}^{\mathcal{I}} - \mathcal{O}_{(0,1)}^{\mathcal{I}} + \mathcal{O}_{(1,0)}^{\mathcal{I}} - \mathcal{O}_{(1,1)}^{\mathcal{I}} \right), \quad e_4 = \frac{1}{4} \left(\mathcal{O}_{(0,0)}^{\mathcal{I}} - \mathcal{O}_{(0,1)}^{\mathcal{I}} - \mathcal{O}_{(1,0)}^{\mathcal{I}} + \mathcal{O}_{(1,1)}^{\mathcal{I}} \right),
$$

\n
$$
e_1' = \frac{1}{2} \left((\mathcal{O}_{D}^{\mathcal{I}})_{11} + (\mathcal{O}_{D}^{\mathcal{I}})_{22} \right), \qquad e_2' = \frac{1}{2} \left((\mathcal{O}_{D}^{\mathcal{I}})_{11} - (\mathcal{O}_{D}^{\mathcal{I}})_{22} \right),
$$

\n
$$
e_3' = \frac{1}{2} \left((\mathcal{O}_{D}^{\mathcal{I}})_{12} + (\mathcal{O}_{D}^{\mathcal{I}})_{21} \right), \qquad e_4' = \frac{1}{2} \left((\mathcal{O}_{D}^{\mathcal{I}})_{12} - (\mathcal{O}_{D}^{\mathcal{I}})_{21} \right).
$$
 (D7)

The direct sum decomposition $(D5)$ implies that the interface algebra A has eight onedimensional representations, which we denote by R_i^s where $s = \pm$ and $i = 1, 2, 3, 4$. The (one by one) representation matrix of each idempotent e_j^t in representation R_i^s is given by

$$
R_i^s(e_j^t) = \delta_{ij}\delta_{st}.\tag{D8}
$$

Namely, e_i^s acts as one in R_i^s and zero in the other representations. Going back to the original basis $\{\mathcal{O}_a^{\mathcal{I}},(\mathcal{O}_D^{\mathcal{I}})_{ij} \mid a \in \mathbb{Z}_2 \times \mathbb{Z}_2, i,j = 1,2\}$ of the interface algebra A, one can summarize the action of symmetry operators in each representation R_i^s as in Table [II.](#page-63-1) The group of one-dimensional representations $\{R_i^s | s = \pm, i = 1, 2, 3, 4\}$ is isomorphic to D_8 .

• When $SPT_1 = SPT_2 = SPT_{\nu_2}$, the multiplication of symmetry operators on the second line of Eq. $(D2)$ is given by

$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}\mathcal{O}_a^{\mathcal{I}} = \mathcal{O}_a^{\mathcal{I}}(\mathcal{O}_D^{\mathcal{I}})_{ij},
$$

\n
$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}(\mathcal{O}_D^{\mathcal{I}})_{kl} = \frac{1}{2}Z_{ik}Z_{jl}\mathcal{O}_{(0,0)}^{\mathcal{I}} + \frac{1}{2}\delta_{ik}\delta_{jl}\mathcal{O}_{(0,1)}^{\mathcal{I}} - \frac{1}{2}Y_{ik}Y_{jl}\mathcal{O}_{(1,0)}^{\mathcal{I}} + \frac{1}{2}X_{ik}X_{jl}\mathcal{O}_{(1,1)}^{\mathcal{I}}.
$$
 (D9)

$\begin{array}{c cccccc} \mathcal{I}=\mathcal{I}_{\nu_2,\nu_2} & R_1^+ & R_1^- & R_2^+ & R_2^- & R_3^+ & R_3^- & R_4^+ & R_4^- \\ \hline \mathcal{O}_{(0,0)}^{\mathcal{I}} & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ \mathcal{O}_{(1,1)}^{\mathcal{I}} & 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 \\ \mathcal{O}_{(1,0)}^{\mathcal{I}} & 1 & 1 & -1 & -1 & 1 & 1 & 1 & 1 \\ \mathcal{O}_{(1,1)}^$				

TABLE III. Irreducible representations of the self-interface algebra $\mathcal A$ for the Rep(D_8) SPT phase SPT_{ν_2}. All irreducible representations are one-dimensional and form a group D_8 under the multiplication.

The multiplications $(D3)$ and $(D9)$ imply that the interface algebra A is commutative. Thus, A can be decomposed into a direct sum of eight one-dimensional subalgebras as

$$
\mathcal{A} = \bigoplus_{s=\pm} \bigoplus_{1 \le i \le 4} \mathbb{C}e_i^s, \tag{D10}
$$

where the idempotents e_i^s are given by

$$
e_1^{\pm} = \frac{1}{2}(e_1 \pm e_1'), \quad e_2^{\pm} = \frac{1}{2}(e_2 \pm e_2'), \quad e_3^{\pm} = \frac{1}{2}(e_3 \pm ie_3'), \quad e_4^{\pm} = \frac{1}{2}(e_4 \pm ie_4').
$$
 (D11)

Here, e_i and e'_i for $i = 1, 2, 3, 4$ are defined by Eq. [\(D7\)](#page-63-3). The direct sum decomposition $(D10)$ implies that the interface algebra A has eight one-dimensional representations, which we denote by R_i^s where $s = \pm$ and $i = 1, 2, 3, 4$. The (one by one) representation matrix of each idempotent e_j^t in representation R_i^s is given by

$$
R_i^s(e_j^t) = \delta_{ij}\delta_{st}.\tag{D12}
$$

Going back to the original basis $\{\mathcal{O}_a^{\mathcal{I}},(\mathcal{O}_D^{\mathcal{I}})_{ij} \mid a \in \mathbb{Z}_2 \times \mathbb{Z}_2, i,j = 1,2\}$ of the interface algebra A , one can summarize the action of symmetry operators in each representation R_i^s as in Table [III.](#page-64-1) The group of one-dimensional representations $\{R_i^s \mid s = \pm, i = 1, 2, 3, 4\}$ is isomorphic to D_8 .

• When $SPT_1 = SPT_2 = SPT_{\nu_3}$, the multiplication of symmetry operators on the second line of Eq. $(D2)$ is given by

$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}\mathcal{O}_a^{\mathcal{I}} = \mathcal{O}_a^{\mathcal{I}}(\mathcal{O}_D^{\mathcal{I}})_{ij},
$$

\n
$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}(\mathcal{O}_D^{\mathcal{I}})_{kl} = \frac{1}{2}\delta_{ik}\delta_{jl}\mathcal{O}_{(0,0)}^{\mathcal{I}} + \frac{1}{2}Z_{ik}Z_{jl}\mathcal{O}_{(0,1)}^{\mathcal{I}} + \frac{1}{2}X_{ik}X_{jl}\mathcal{O}_{(1,0)}^{\mathcal{I}} - \frac{1}{2}Y_{ik}Y_{jl}\mathcal{O}_{(1,1)}^{\mathcal{I}}.
$$
\n(D13)

The multiplications [\(D3\)](#page-62-2) and [\(D13\)](#page-64-2) imply that the interface algebra A is commutative. Thus, A can be decomposed into a direct sum of eight one-dimensional subalgebras as

$$
\mathcal{A} = \bigoplus_{s=\pm} \bigoplus_{1 \le i \le 4} \mathbb{C}e_i^s, \tag{D14}
$$

where the idempotents e_i^s are given by

$$
e_1^{\pm} = \frac{1}{2}(e_1 \pm e_1'), \quad e_2^{\pm} = \frac{1}{2}(e_2 \pm e_2'), \quad e_3^{\pm} = \frac{1}{2}(e_3 \pm e_3'), \quad e_4^{\pm} = \frac{1}{2}(e_4 \pm e_4').
$$
 (D15)

$\begin{array}{c cccccc} \mathcal{I}=\mathcal{I}_{\nu_3,\nu_3} & R_1^+ & R_1^- & R_2^+ & R_2^- & R_3^+ & R_3^- & R_4^+ & R_4^- \\ \hline \mathcal{O}_{(0,0)}^{\mathcal{I}} & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ \mathcal{O}_{(0,1)}^{\mathcal{I}} & 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 \\ \mathcal{O}_{(1,0)}^{\mathcal{I}} & 1 & 1 & -1 & -1 & 1 & 1 & 1 & 1 \\ \mathcal{O}_{(1,1)}^$				

TABLE IV. Irreducible representations of the self-interface algebra $\mathcal A$ for the Rep(D_8) SPT phase SPT_{ν_3}. All irreducible representations are one-dimensional and form a group D_8 under the multiplication.

Here, e_i and e'_i for $i = 1, 2, 3, 4$ are defined by Eq. [\(D7\)](#page-63-3). The direct sum decomposition $(D14)$ implies that the interface algebra A has eight one-dimensional representations, which we denote by R_i^s where $s = \pm$ and $i = 1, 2, 3, 4$. The (one by one) representation matrix of each idempotent e_j^t in representation R_i^s is given by

$$
R_i^s(e_j^t) = \delta_{ij}\delta_{st}.\tag{D16}
$$

Going back to the original basis $\{\mathcal{O}_a^{\mathcal{I}},(\mathcal{O}_D^{\mathcal{I}})_{ij} \mid a \in \mathbb{Z}_2 \times \mathbb{Z}_2, i,j = 1,2\}$ of the interface algebra A , one can summarize the action of symmetry operators in each representation R_i^s as in Table [IV.](#page-65-0) The group of one-dimensional representations $\{R_i^s \mid s = \pm, i = 1, 2, 3, 4\}$ is isomorphic to D_8 .

b. Interface between different SPT phases

The interface algebra A between different SPT phases SPT_1 and SPT_2 does not have one-dimensional representations as shown in Sec. [IV B.](#page-29-0) Thus, the direct sum decomposition of A does not contain one-dimensional factors. In addition, since A is an eight-dimensional algebra, the only possible decomposition is

$$
\mathcal{A} = M_2(\mathbb{C}) \oplus M_2(\mathbb{C}). \tag{D17}
$$

The above direct sum decomposition implies that A has two two-dimensional irreducible representations. In what follows, we will explicitly decompose A into a direct sum [\(D17\)](#page-65-1) and write down its irreducible representations. Given a representation R of the interface algebra $\mathcal{A}_{\nu_i,\nu_j}$, one can obtain the corresponding representation of $\mathcal{A}_{\nu_j,\nu_i}$, which we denote by R by abuse of notation, as follows:

$$
R(\mathcal{O}_{a}^{\mathcal{I}_{\nu_{j},\nu_{i}}}) = R(\mathcal{O}_{a}^{\mathcal{I}_{\nu_{i},\nu_{j}}})^{*}, \quad R((\mathcal{O}_{D}^{\mathcal{I}_{\nu_{j},\nu_{i}}})_{kl}) = R(\mathcal{O}_{D}^{\mathcal{I}_{\nu_{i},\nu_{j}}})_{lk})^{*}.
$$
 (D18)

Therefore, we only need to consider three interface algebras $\mathcal{A}_{\nu_1,\nu_2}$, $\mathcal{A}_{\nu_2,\nu_3}$, and $\mathcal{A}_{\nu_3,\nu_1}$.

• When $SPT_1 = SPT_{\nu_1}$ and $SPT_2 = SPT_{\nu_2}$, the multiplication of symmetry operators on the

second line of Eq. $(D2)$ reduces to

$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}\mathcal{O}_a^{\mathcal{I}} = \begin{cases} (\mathcal{O}_D^{\mathcal{I}})_{ij} & a = (0,0), \\ -(Z\mathcal{O}_D^{\mathcal{I}}Z)_{ij} & a = (0,1), \\ -(X\mathcal{O}_D^{\mathcal{I}}X)_{ij} & a = (1,0), \\ (Y\mathcal{O}_D^{\mathcal{I}}Y)_{ij} & a = (1,1), \end{cases}
$$
\n
$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}(\mathcal{O}_D^{\mathcal{I}})_{kl} = \frac{1}{2}X_{ik}Z_{jl}\mathcal{O}_{(0,0)}^{\mathcal{I}} + \frac{1}{2}iY_{ik}\delta_{jl}\mathcal{O}_{(0,1)}^{\mathcal{I}} - \frac{1}{2}\delta_{ik}iY_{jl}\mathcal{O}_{(1,0)}^{\mathcal{I}} - \frac{1}{2}Z_{ik}X_{jl}\mathcal{O}_{(1,1)}^{\mathcal{I}}.
$$
\n(D19)

The multiplications [\(D3\)](#page-62-2) and [\(D19\)](#page-66-0) imply that the interface algebra A is decomposed as Eq. $(D17)$, where the two factors of the direct sum decomposition $(D17)$ are generated by $\{e_i, e'_i \mid i = 1, 4\}$ and $\{e_i, e'_i \mid i = 2, 3\}$ respectively. Here, e_i and e'_i are defined by Eq. [\(D7\)](#page-63-3). Indeed, in a new basis given by

$$
e_{11} = e_1, e_{12} = ie'_1, e_{21} = ie'_4, e_{22} = e_4,
$$

\n $d_{11} = e_2, d_{12} = e'_2, d_{21} = e'_3, d_{22} = e_3,$ (D20)

the multiplication of A can be written as

 $\overline{(\ }$

$$
e_{ij}e_{kl} = \delta_{jk}e_{il}, \quad d_{ij}d_{kl} = \delta_{jk}d_{il}, \quad e_{ij}d_{kl} = d_{ij}e_{kl} = 0,
$$
\n(D21)

which shows that each of $\{e_{ij} | i,j = 1,2\}$ and $\{d_{ij} | i,j = 1,2\}$ spans the full matrix algebra $M_2(\mathbb{C})$. The above direct sum decomposition implies that A has two two-dimensional irreducible representations, which we denote by R_e and R_d . The (two by two) representation matrices of e_{ij} and d_{ij} in these representations are given by

$$
R_e(e_{11}) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad R_e(e_{12}) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad R_e(e_{21}) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad R_e(e_{22}) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad R_e(d_{ij}) = 0,
$$

$$
R_d(d_{11}) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad R_d(d_{12}) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad R_d(d_{21}) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad R_d(d_{22}) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad R_d(e_{ij}) = 0.
$$

(D22)

Going back to the original basis $\{\mathcal{O}_a^{\mathcal{I}},(\mathcal{O}_D^{\mathcal{I}})_{ij} \mid a \in \mathbb{Z}_2 \times \mathbb{Z}_2, i,j = 1,2\}$ of the interface algebra A, one can summarize the action of symmetry operators in each representation as in Table [V.](#page-67-0)

• When $SPT_1 = SPT_{\nu_2}$ and $SPT_2 = SPT_{\nu_3}$, the multiplication of symmetry operators on the second line of Eq. $(D2)$ reduces to

$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}\mathcal{O}_a^{\mathcal{I}} = \begin{cases} (\mathcal{O}_D^{\mathcal{I}})_{ij} & a = (0,0), \\ (Z\mathcal{O}_D^{\mathcal{I}}Z)_{ij} & a = (0,1), \\ -(X\mathcal{O}_D^{\mathcal{I}}X)_{ij} & a = (1,0), \\ -(Y\mathcal{O}_D^{\mathcal{I}}Y)_{ij} & a = (1,1), \end{cases}
$$
(D23)

$$
\mathcal{O}_D^{\mathcal{I}})_{ij}(\mathcal{O}_D^{\mathcal{I}})_{kl} = \frac{1}{2}Z_{ik}\delta_{jl}\mathcal{O}_{(0,0)}^{\mathcal{I}} + \frac{1}{2}\delta_{ik}Z_{jl}\mathcal{O}_{(0,1)}^{\mathcal{I}} - \frac{1}{2}iY_{ik}X_{jl}\mathcal{O}_{(1,0)}^{\mathcal{I}} - \frac{1}{2}X_{ik}iY_{jl}\mathcal{O}_{(1,1)}^{\mathcal{I}}.
$$

The multiplications [\(D3\)](#page-62-2) and [\(D23\)](#page-66-1) imply that the interface algebra A is decomposed as Eq. $(D17)$, where the two factors of the direct sum decomposition $(D17)$ are generated by

$$
\begin{array}{c|c|c|c} \mathcal{I} = \mathcal{I}_{\nu_1, \nu_2} & R_e & R_d & \mathcal{I} = \mathcal{I}_{\nu_2, \nu_1} & R_e & R_d \\ \hline \mathcal{O}_{(0,0)}^{\mathcal{I}} & I_2 & I_2 & \mathcal{O}_{(0,0)}^{\mathcal{I}} & I_2 & I_2 \\ \mathcal{O}_{(1,1)}^{\mathcal{I}} & Z & Z & \mathcal{O}_{(1,0)}^{\mathcal{I}} & Z & Z \\ \mathcal{O}_{(1,0)}^{\mathcal{I}} & I_2 & -Z & \mathcal{O}_{(1,0)}^{\mathcal{I}} & Z & -Z \\ \mathcal{O}_{(1,1)}^{\mathcal{I}} & I_2 & -I_2 & \mathcal{O}_{(1,1)}^{\mathcal{I}} & I_2 & -I_2 \\ \mathcal{O}_{(1,1)}^{\mathcal{I}} & I_2 & -I_2 & \mathcal{O}_{(1,1)}^{\mathcal{I}} & I_2 & -I_2 \\ \mathcal{O}_{(1,1)}^{\mathcal{I}} & I_2 & -I_2 & \mathcal{O}_{(1,1)}^{\mathcal{I}} & I_2 & -I_2 \\ \mathcal{O}_{(1,1)}^{\mathcal{I}} & \left(\frac{-i}{2}S_+ - \frac{i}{2}S_- \right)_{ij} & \left(\frac{1}{2}S_+ - \frac{i}{2}S_+ \right)_{ij} & \left(\mathcal{O}_{D}^{\mathcal{I}}\right)_{ij} & \left(\frac{i}{2}S_- - \frac{i}{2}S_+ \right)_{ij} & \left(\frac{1}{2}S_- - \frac{1}{2}S_+ \right)_{ij} \end{array}
$$

TABLE V. Irreducible representations of the interface algebra $\mathcal A$ between Rep(D_8) SPT phases SPT_{ν_1} and SPT_{ν_2} . Here, I_2 denotes the 2×2 identity matrix and $S_{\pm} := X \pm iY$. The bottom row reads $R_e((\mathcal{O}_D^{\mathcal{I}})_{11}) =$ $-\frac{i}{2}S_+$, etc.

$\mathcal{I}=\mathcal{I}_{\nu_2,\underline{\nu_3}}$			$\mathcal{I}=\mathcal{I}_{\nu_3,\underline{\nu_2}}$ l	R_{e}	
$\mathcal{O}_{(0,0)}^2$	I_2	L,		I_{2}	I ₂
$\mathcal{O}_{(0,1)}^{\mathcal{I}}$	I_2	$\sqrt{2}$		I_2	$-I_2$
$\mathcal{O}_{(1,0)}^{\mathcal{I}}$	Z			Ζ	Z
$\mathcal{O}_{(1,1)}^{\mathcal{I}}$					$-Z$
$(\mathcal{O}_D^{\mathcal{I}})_{ij}$		$\begin{pmatrix} X & 0 \ 0 & iY \end{pmatrix}_{ij} \begin{pmatrix} 0 & X \ iY & 0 \end{pmatrix}_{ij}$	$(\mathcal{O}_D^{\mathcal{I}})_{ij}$		$\begin{pmatrix} X & 0 \ 0 & iY \end{pmatrix}_{ij} \begin{pmatrix} 0 & iY \ X & 0 \end{pmatrix}_{ij}$

TABLE VI. Irreducible representations of the interface algebra A between Rep(D_8) SPT phases SPT_{ν_2} and SPT_{ν_3} . Here, I_2 denotes the 2×2 identity matrix, and the bottom row reads $R_e((\mathcal{O}_D^{\mathcal{I}})_{11}) = X$, etc.

 $\{e_i, e'_i \mid i = 1, 2\}$ and $\{e_i, e'_i \mid i = 3, 4\}$ respectively. Here, e_i and e'_i are defined by Eq. [\(D7\)](#page-63-3). Indeed, in a new basis given by

$$
e_{11} = e_1, e_{12} = e'_1, e_{21} = e'_2, e_{22} = e_2,
$$

\n $d_{11} = e_3, d_{12} = e'_3, d_{21} = e'_4, d_{22} = e_4,$ (D24)

the multiplication of A can be written as

$$
e_{ij}e_{kl} = \delta_{jk}e_{il}, \quad d_{ij}d_{kl} = \delta_{jk}d_{il}, \quad e_{ij}d_{kl} = d_{ij}e_{kl} = 0,
$$
\n(D25)

which shows that each of $\{e_{ij} | i,j = 1,2\}$ and $\{d_{ij} | i,j = 1,2\}$ spans the full matrix algebra $M_2(\mathbb{C})$. The above direct sum decomposition implies that A has two two-dimensional irreducible representations, which we denote by R_e and R_d . The (two by two) representation matrices of e_{ij} and d_{ij} in these representations are given by Eq. [\(D22\)](#page-66-2). Going back to the original basis $\{\mathcal{O}_a^{\mathcal{I}},(\mathcal{O}_D^{\mathcal{I}})_{ij} \mid a \in \mathbb{Z}_2 \times \mathbb{Z}_2, i,j = 1,2\}$ of the interface algebra A, one can summarize the action of symmetry operators in each representation as in Table [VI.](#page-67-1)

• When $SPT_1 = SPT_{\nu_3}$ and $SPT_2 = SPT_{\nu_1}$, the multiplication of symmetry operators on the

$\mathcal{I}=\mathcal{I}_{\nu_3,\nu_1}$			$\mathcal{I}=\mathcal{I}_{\nu_1,\underline{\nu_3}}$		
$\mathcal{O}_{(0,0)}^{\mathcal{I}}$	\mathcal{L}_{2}	I2	$\mathcal{O}_{(0,0)}^{\mathcal{I}}$	Ь	12
$\mathcal{O}_{(0,1)}^{\mathcal{I}}$			$\mathcal{O}_{(0,1)}^{\mathcal{I}}$		
$\mathcal{O}_{(1,0)}^{\mathcal{I}}$	Iо	$-I2$	$\mathcal{O}_{(1,0)}^{\mathcal{I}}$, and $\mathcal{O}_{(1,0)}^{\mathcal{I}}$	I ₂	
$\mathcal{O}_{(1,1)}^{\mathcal{I}}$			$\mathcal{O}_{(1,1)}^{\mathcal{L}}$		
$(\mathcal{O}_D^{\mathcal{I}})_{ij}$		$\begin{pmatrix} \frac{1}{2}S_{+} & \frac{1}{2}S_{-} \\ \frac{1}{2}S_{-} & \frac{1}{2}S_{+} \end{pmatrix}_{ij} \begin{pmatrix} \frac{1}{2}S_{+} & \frac{1}{2}S_{-} \\ -\frac{1}{2}S_{-} & -\frac{1}{2}S_{+} \end{pmatrix}_{ij}$	$(\mathcal{O}_D^{\mathcal{I}})_{ij}$		$\begin{pmatrix} \frac{1}{2}S_{+} & \frac{1}{2}S_{-} \\ \frac{1}{2}S_{-} & \frac{1}{2}S_{+} \end{pmatrix}_{ij} \begin{pmatrix} \frac{1}{2}S_{+} & -\frac{1}{2}S_{-} \\ \frac{1}{2}S_{-} & -\frac{1}{2}S_{+} \end{pmatrix}_{ij}$

TABLE VII. Irreducible representations of the interface algebra A between $\text{Rep}(D_8)$ SPT phases SPT_{ν_3} and SPT_{ν_1} . Here, I_2 denotes the 2×2 identity matrix and $S_{\pm} := X \pm iY$. The bottom row reads $R_e((\mathcal{O}_D^{\mathcal{I}})_{11}) =$ $\frac{1}{2}S_+$, etc.

second line of Eq. $(D2)$ reduces to

$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}\mathcal{O}_a^{\mathcal{I}} = \begin{cases} (\mathcal{O}_D^{\mathcal{I}})_{ij} & a = (0,0), \\ -(Z\mathcal{O}_D^{\mathcal{I}}Z)_{ij} & a = (0,1), \\ (X\mathcal{O}_D^{\mathcal{I}}X)_{ij} & a = (1,0), \\ -(Y\mathcal{O}_D^{\mathcal{I}}Y)_{ij} & a = (1,1), \end{cases}
$$
(D26)

$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}(\mathcal{O}_D^{\mathcal{I}})_{kl} = \frac{1}{2}\delta_{ik}X_{jl}\mathcal{O}_{(0,0)}^{\mathcal{I}} + \frac{1}{2}Z_{ik}iY_{jl}\mathcal{O}_{(0,1)}^{\mathcal{I}} + \frac{1}{2}X_{ik}\delta_{jl}\mathcal{O}_{(1,0)}^{\mathcal{I}} + \frac{1}{2}iY_{ik}Z_{jl}\mathcal{O}_{(1,1)}^{\mathcal{I}}.
$$

The multiplications [\(D3\)](#page-62-2) and [\(D26\)](#page-68-0) imply that the interface algebra A is decomposed as Eq. $(D17)$, where the two factors of the direct sum decomposition $(D17)$ are generated by $\{e_i, e'_i \mid i = 1, 3\}$ and $\{e_i, e'_i \mid i = 2, 4\}$ respectively. Here, e_i and e'_i are defined by Eq. [\(D7\)](#page-63-3). Indeed, in a new basis given by

$$
e_{11} = e_1, e_{12} = e'_1, e_{21} = e'_3, e_{22} = e_3,
$$

\n $d_{11} = e_2, d_{12} = e'_2, d_{21} = e'_4, d_{22} = e_4,$ (D27)

the multiplication of A can be written as

$$
e_{ij}e_{kl} = \delta_{jk}e_{il}, \quad d_{ij}d_{kl} = \delta_{jk}d_{il}, \quad e_{ij}d_{kl} = d_{ij}e_{kl} = 0,
$$
\n(D28)

which shows that each of $\{e_{ij} | i, j = 1, 2\}$ and $\{d_{ij} | i, j = 1, 2\}$ spans the full matrix algebra $M_2(\mathbb{C})$. The above direct sum decomposition implies that A has two two-dimensional irreducible representations, which we denote by R_e and R_d . The (two by two) representation matrices of e_{ij} and d_{ij} in these representations are given by Eq. [\(D22\)](#page-66-2). Going back to the original basis $\{\mathcal{O}_a^{\mathcal{I}},(\mathcal{O}_D^{\mathcal{I}})_{ij} \mid a \in \mathbb{Z}_2 \times \mathbb{Z}_2, i,j = 1,2\}$ of the interface algebra A, one can summarize the action of symmetry operators in each representation as in Table [VII.](#page-68-1)

2. Rep (Q_8) symmetry

The $\text{Rep}(Q_8)$ symmetry has only one SPT phase and hence the only possible interface is the selfinterface. The L-symbols for this SPT phase are given by Eqs. $(C3)$, $(C9)$, and $(C10)$. Substituting

			$\begin{vmatrix} R_1^+ & R_1^- & R_2^+ & R_2^- & R_3^+ & R_3^- & R_4^+ & R_4^- \end{vmatrix}$		
$\begin{array}{c cccccc} \hline \mathcal{O}_{(0,0)}^{\mathcal{I}} & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ \mathcal{O}_{(0,1)}^{\mathcal{I}} & 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 \\ \mathcal{O}_{(1,0)}^{\mathcal{I}} & 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 \\ \hline \mathcal{O}_{(1,1)}^{\mathcal{I}} & 1 & 1 & -1 & -1 & 1 & 1 & 1 & 1 \\ \mathcal{O}_{(1,1)}^{\mathcal{I}} & 1 & 1 & -1 & -1 & -1 & -$					

TABLE VIII. Irreducible representations of the self-interface algebra $\mathcal A$ for the Rep(Q_8) SPT phase. All irreducible representations are one-dimensional and form a group Q_8 under the multiplication.

these L-symbols into the second line of Eq. $(D2)$, one finds that the multiplication of symmetry operators of the self-interface algebra $\mathcal A$ is given by

$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}\mathcal{O}_a^{\mathcal{I}} = \mathcal{O}_a^{\mathcal{I}}(\mathcal{O}_D^{\mathcal{I}})_{ij}
$$

\n
$$
(\mathcal{O}_D^{\mathcal{I}})_{ij}(\mathcal{O}_D^{\mathcal{I}})_{kl} = -\frac{1}{2}Y_{ik}Y_{jl}\mathcal{O}_{(0,0)}^{\mathcal{I}} + \frac{1}{2}X_{ik}X_{jl}\mathcal{O}_{(0,1)}^{\mathcal{I}} + \frac{1}{2}Z_{ik}Z_{jl}\mathcal{O}_{(1,0)}^{\mathcal{I}} + \frac{1}{2}\delta_{ik}\delta_{jl}\mathcal{O}_{(1,1)}^{\mathcal{I}}.
$$
\n(D29)

The multiplications [\(D3\)](#page-62-2) and [\(D29\)](#page-69-0) imply that the interface algebra A is commutative. Thus, A can be decomposed into a direct sum of eight one-dimensional subalgebras as

$$
\mathcal{A} = \bigoplus_{s=\pm} \bigoplus_{1 \le i \le 4} \mathbb{C}e_i^s,\tag{D30}
$$

where the idempotents e_i^s are given by

$$
e_1^{\pm} = \frac{1}{2}(e_1 \pm e_1'), \quad e_2^{\pm} = \frac{1}{2}(e_2 \pm ie_2'), \quad e_3^{\pm} = \frac{1}{2}(e_3 \pm ie_3'), \quad e_4^{\pm} = \frac{1}{2}(e_4 \pm e_4').
$$
 (D31)

Here, e_i and e'_i for $i = 1, 2, 3, 4$ are defined by Eq. [\(D7\)](#page-63-3). The direct sum decomposition [\(D14\)](#page-64-3) implies that the interface algebra A has eight one-dimensional representations, which we denote by R_i^s where $s = \pm$ and $i = 1, 2, 3, 4$. The (one by one) representation matrix of each idempotent e_j^t in representation R_i^s is given by

$$
R_i^s(e_j^t) = \delta_{ij}\delta_{st}.\tag{D32}
$$

Going back to the original basis $\{\mathcal{O}_a^{\mathcal{I}}, (\mathcal{O}_D^{\mathcal{I}})_{ij} \mid a \in \mathbb{Z}_2 \times \mathbb{Z}_2, i, j = 1, 2\}$ of the interface algebra A, one can summarize the action of symmetry operators in each representation R_i^s as in Table [VIII.](#page-69-1) The group of one-dimensional representations $\{R_i^s | s = \pm, i = 1, 2, 3, 4\}$ is isomorphic to Q_8 .

3. Rep (H_8) symmetry

The $\text{Rep}(H_8)$ symmetry has only one SPT phase and hence the only possible interface is the selfinterface. The L-symbols for this SPT phase are given by Eqs. $(C3)$, $(C12)$, and $(C13)$. Substituting these L-symbols into the second line of Eq. [\(D2\)](#page-62-1), one finds that the multiplication of symmetry operators of the self-interface algebra A is given by

$$
(\mathcal{O}_{D}^{\mathcal{I}})_{ij}\mathcal{O}_{a}^{\mathcal{I}} = \begin{cases} (\mathcal{O}_{D}^{\mathcal{I}})_{ij} & a = (0,0), \\ (X\mathcal{O}_{D}^{\mathcal{I}}X)_{ij} & a = (0,1), \\ (Z\mathcal{O}_{D}^{\mathcal{I}}Z)_{ij} & a = (1,0), \\ (Y\mathcal{O}_{D}^{\mathcal{I}}Y)_{ij} & a = (1,1), \end{cases}
$$
\n
$$
(\mathcal{O}_{D}^{\mathcal{I}})_{ij}(\mathcal{O}_{D}^{\mathcal{I}})_{kl} = \frac{1}{4}(X+Z)_{ik}(X+Z)_{jl}\mathcal{O}_{(0,0)}^{\mathcal{I}} + \frac{1}{4}(1+iY)_{ik}(1+iY)_{jl}\mathcal{O}_{(0,1)}^{\mathcal{I}}
$$
\n
$$
+ \frac{1}{4}(1-iY)_{ik}(1-iY)_{jl}\mathcal{O}_{(1,0)}^{\mathcal{I}} + \frac{1}{4}(X-Z)_{ik}(X-Z)_{jl}\mathcal{O}_{(1,1)}^{\mathcal{I}}.
$$
\n(D33)

The multiplications [\(D3\)](#page-62-2) and [\(D33\)](#page-70-0) imply that the interface algebra A is non-commutative. Thus, A cannot be a direct sum of eight one-dimensional subalgebras. On the other hand, since the algebra A is acting on the self-interface, it must have one-dimensional representations as shown in Sec. [IV B.](#page-29-0) In other words, the direct sum decomposition of A has to contain one-dimensional subalgebras. Therefore, it follows that A is decomposed as

$$
\mathcal{A} = \mathbb{C} \oplus \mathbb{C} \oplus \mathbb{C} \oplus \mathbb{C} \oplus M_2(\mathbb{C}).
$$
 (D34)

The basis of each one-dimensional component $\mathbb C$ is given by

$$
e_1^+ = \frac{1}{2}(e_1 + e_1'),
$$
 $e_1^- = \frac{1}{2}(e_1 - e_1'),$ $e_4^+ = \frac{1}{2}(e_4 + ie_4'),$ $e_4^- = \frac{1}{2}(e_4 - ie_4'),$ (D35)

where e_1, e'_1, e_4 , and e'_4 are defined by Eq. [\(D7\)](#page-63-3). On the other hand, a basis $\{e_{ij} | i, j = 1, 2\}$ of the four-dimensional component $M_2(\mathbb{C})$ can be written as

$$
e_{11} = e_2
$$
, $e_{12} = e'_2$, $e_{21} = e'_3$, $e_{22} = e_3$. (D36)

The multiplication of the bases in Eqs. $(D35)$ and $(D36)$ is

$$
e_i^s e_j^t = \delta_{ij} \delta_{st} e_i^s, \qquad e_{ij} e_{kl} = \delta_{jk} e_{il}, \qquad e_i^s e_{jk} = e_{jk} e_i^s = 0.
$$
 (D37)

The direct sum decomposition [\(D34\)](#page-70-3) implies that A has four one-dimensional representations $\{R_i^s\}$ $s = \pm$, $i = 1, 4$ and a single two-dimensional irreducible representation R. The (one by one) representation matrices of e_j^t and e_{jk} in one-dimensional representation R_i^s are given by

$$
R_i^s(e_j^t) = \delta_{ij}\delta_{st}, \qquad R_i^s(e_{jk}) = 0.
$$
 (D38)

Similarly, the (two by two) representation matrices of e_i^s and e_{ij} in two-dimensional representation R are given by

$$
R(e_i^s) = 0, \quad R(e_{11}) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad R(e_{12}) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad R(e_{21}) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad R(e_{22}) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.
$$
\n(D39)

Going back to the original basis $\{\mathcal{O}_a^{\mathcal{I}},(\mathcal{O}_D^{\mathcal{I}})_{ij} \mid a \in \mathbb{Z}_2 \times \mathbb{Z}_2, i,j = 1,2\}$ of the interface algebra A , we can summarize the action of symmetry operators in each representation as in Table [IX.](#page-71-0) The group of one-dimensional representations $\{R_i^s | s = \pm, i = 1, 4\}$ is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$, which is the group of group-like elements of $H_8^* \cong H_8$.

	R_1^+ $R_1^ R_4^+$ R_4^-		
			$\begin{array}{c cccc}\n\hline\n\mathcal{O}^{\mathcal{I}}_{(0,0)} & 1 & 1 & 1 & 1 & I_2\\ \mathcal{O}^{\mathcal{I}}_{(0,1)} & 1 & 1 & 1 & 1 & I_2\\ \mathcal{O}^{\mathcal{I}}_{(1,0)} & 1 & 1 & -1 & -1 & -Z\\ \mathcal{O}^{\mathcal{I}}_{(1,1)} & 1 & 1 & 1 & 1 & -I_2\\ \mathcal{O}^{\mathcal{I}}_{(1,1)} & 1 & 1 & 1 & 1 & -I_2\\ (\mathcal{O}^{\mathcal{I}}_{D})_{ij} &$

TABLE IX. Irreducible representations of the self-interface algebra $\mathcal A$ for the Rep(H_8) SPT phase. Onedimensional representations form a group $\mathbb{Z}_2 \times \mathbb{Z}_2$ under the multiplication. On the rightmost column, I_2 denotes the 2 × 2 identity matrix, and the bottom entry means that $R((\mathcal{O}_D^{\mathcal{I}})_{11}) = \frac{1}{2}(X + iY)$, etc.

Appendix E: The $G \times \text{Rep}(G)$ cluster state as a trivial $\text{Rep}(G)$ SPT state

In this appendix, we show that the $G \times \text{Rep}(G)$ cluster state [\(81\)](#page-25-1) is the exact ground state of the Rep(G) symmetric model discussed in [\[47,](#page-86-6) Section 4.4]. This gives us another piece of evidence for the claim that the $G \times \text{Rep}(G)$ cluster state is in the $\text{Rep}(G)$ -symmetric trivial phase labeled by the forgetful functor of $\text{Rep}(G)$.

To begin with, we review the general construction of commuting projector models and their ground states with $\text{Rep}(H)$ symmetry following [\[47\]](#page-86-6), where H is a general finite semisimple Hopf algebra. The physical Hilbert space of the model is chosen to be a finite semisimple left H-comodule algebra K, i.e., an algebra equipped with a left H-comodule action $\lambda_K : K \to H \otimes K$ that is compatible with the algebra structure on K . Since K is a semisimple algebra, it admits a unique ∆-separable symmetric Frobenius algebra structure [\[139,](#page-89-17) [140\]](#page-89-18). We denote the multiplication and comultiplication of the Frobenius algebra K by $m_K : K \otimes K \to K$ and $\Delta_K : K \to K \otimes K$ respectively. The Hamiltonian of the model is given by the sum of local commuting projectors as follows:

$$
H = -\sum_{i} h_{i,i+1}, \quad h_{i,i+1} = \Delta_K \circ m_K : K \otimes K \to K \otimes K.
$$
 (E1)

Here, $h_{i,i+1}$ is a local term that acts only on sites i and $i+1$. The ground states of this Hamiltonian are in one-to-one correspondence with simple left K-modules. Specifically, the ground state labeled by a simple left K -module M can be written as an MPS

$$
|A_M\rangle = \underbrace{M \underset{A_M}{\overset{K}{\longrightarrow}} \underset{A_M}{\overset{K}{\longrightarrow}} \underset{A_M}{\overset{K}{\longrightarrow}} M},\tag{E2}
$$

where $A_M : M \to K \otimes M$ denotes the left K-comodule action on M defined by

$$
A_M = (\mathrm{id}_K \otimes \rho_M) \circ ((\Delta_K \circ u) \otimes \mathrm{id}_M). \tag{E3}
$$

Here, $\rho_M : K \otimes M \to M$ denotes the left K-action on M and $u : \mathbb{C} \to K$ denotes the unit of K. It turns out that the Hamiltonian $(E1)$ realizes the Rep(H)-symmetric gapped phase that corresponds to a Rep(H)-module category $K\mathcal{M}$, the category of left K-modules [\[47\]](#page-86-6). The above construction produces all gapped phases with $\text{Rep}(H)$ symmetry because any $\text{Rep}(H)$ -module category is equivalent to $_K\mathcal{M}$ for some left H-comodule algebra K [\[141\]](#page-89-19). In particular, the Hamiltonian [\(E1\)](#page-71-1) realizes a Rep(H) SPT phase when K is simple.
Now, let us restrict our attention to the $\text{Rep}(G)$ SPT phase corresponding to the forgetful functor of Rep(G) [\[47,](#page-86-0) Section 4.4]. To obtain this SPT phase, we choose the physical Hilbert space K to be the smash product $K = \mathbb{C}[G]^* \# \mathbb{C}[G]$ of the group algebra $\mathbb{C}[G]$ and its dual $\mathbb{C}[G]^*$. Concretely, K is isomorphic to $\mathbb{C}[G]^* \otimes \mathbb{C}[G]$ as a vector space, and the multiplication and comultiplication of K are given by

$$
(v^{g_1} \# v_{h_1}) \cdot (v^{g_2} \# v_{h_2}) = \delta_{g_1, h_1 g_2} v^{g_1} \# v_{h_1 h_2},
$$

$$
\Delta_K \circ u(1) = \frac{1}{|G|} \sum_{g, h \in G} (v^g \# v_h) \otimes (v^{h^{-1}g} \# v_{h^{-1}}).
$$
 (E4)

Here, $\{v^g \mid g \in G\}$ and $\{v_g \mid g \in G\}$ are dual bases of $\mathbb{C}[G]^*$ and $\mathbb{C}[G]$. The multiplication of K is simply written as \cdot in Eq. [\(E4\)](#page-72-0). The algebra K defined above is simple, ^{[49](#page-72-1)} meaning that an irreducible left K-module M is unique up to isomorphism. Since the dimension of K is $|G|^2$, the dimension of its irreducible module M is $\sqrt{|G|^2} = |G|$. Thus, M is isomorphic to $\mathbb{C}[G]$ as a vector space. If we denote the basis of M as $\{w_q | g \in G\}$, the action of K on M can be written explicitly as

$$
(v_g \# v_h) \cdot w_l = \delta_{l,h^{-1}g} w_{hl},\tag{E5}
$$

where · represents the left K-action $\rho_M : K \otimes M \to M$. Substituting Eqs. [\(E4\)](#page-72-0) and [\(E5\)](#page-72-2) into Eq. [\(E3\)](#page-71-0), we find that the MPS tensor A_M of the ground state is given by

$$
A_M(w_g) = \frac{1}{|G|} \sum_{h \in G} (v^g \# v_h) \otimes w_{h^{-1}g}.
$$
 (E6)

In terms of diagrams, this MPS tensor can be written as^{50} as^{50} as^{50}

$$
\frac{K}{A_M} = \frac{\mathbb{C}[G]^* \mathbb{C}[G]}{A_o A_e},
$$
(E7)

where

$$
\overrightarrow{A_o} = \sum_{g \in G} P_g \otimes |g\rangle_{\text{phys}}, \quad \overrightarrow{A_e} = \frac{1}{|G|} \sum_{g \in G} L_g \otimes |g\rangle_{\text{phys}}.
$$
 (E8)

The operators P_g and L_g are defined by $P_g = |g\rangle \langle g|$ and $L_g = \sum_{h \in G} |gh\rangle \langle h|$ as in Eq. [\(82\)](#page-25-0). The above MPS tensors agree with those of the $G \times \text{Rep}(G)$ cluster state [\(81\)](#page-25-1) up to normalization.

Appendix F: Weak completeness relation

In this appendix, we show the following weak completeness relation, cf. eq. [\(43\)](#page-16-0):

$$
\sum_{i} \frac{\overrightarrow{\psi}}{(\phi_x)_j} \overrightarrow{(\overrightarrow{\phi}_x)_i (\phi_x)_i} = \frac{\overrightarrow{\psi}}{(\phi_x)_j} \tag{F1}
$$

 49 A straightforward computation shows that the center of K is one-dimensional and hence K is simple.

⁵⁰ The mismatch of the orientations of the arrows in Eq. [\(E7\)](#page-72-4) is not a typo: taking the standard basis of $\mathbb{C}[G]$, one can verify that the components of the tensors on both sides agree with each other.

Here, ψ is an arbitrary unitary operator acting on the bond Hilbert space V of the MPS. The bond Hilbert space of the MPO \mathcal{O}_x will be denoted by V_x . In what follows, every diagram is supposed to represent a linear map from the left side to the right side. For instance, the diagrams in eq. [\(F1\)](#page-72-5) represent linear maps from V^* to $V_x \otimes V^*$.

To show eq. $(F1)$, we assume the following dual orthogonality relation:

$$
\overbrace{(\overline{\phi}_x)_i \ (\phi_x)_j}^{\bullet} = \delta_{ij} \longrightarrow . \tag{F2}
$$

This equality should be regarded as the orthogonality relation for the dual object $x^* \in \mathcal{C}$. Equa-tion [\(F2\)](#page-73-0) implies that the action tensor $(\phi_x)_j$ is surjective as a linear map from $V^*_x \otimes V^*$ to V^* . Correspondingly, the image of $(\phi_x)_j$ viewed as a linear map from V^* to $V_x \otimes V^*$ is of the form

$$
\operatorname{Im}((\phi_x)_j) = \operatorname{Span}\{w_\alpha \otimes v_\alpha \mid \alpha = 1, 2, \cdots, \dim V\} \subset V_x \otimes V^*,
$$
 (F3)

where $\{v_\alpha \mid \alpha = 1, 2, \cdots, \dim V\}$ is a basis of V^* and w_α is an element of V_x .

Now, since ψ is unitary, it is diagonalizable by another unitary operator U as follows:

$$
U^{\dagger} \psi U = \text{diag}(\lambda_1, \lambda_2, \cdots, \lambda_{\text{dim } V}), \quad |\lambda_i| = 1. \tag{F4}
$$

If we choose a basis $\{v_\alpha\}$ to be the set of eigenvectors of ψ , the action of $\mathrm{id}_{V_x} \otimes \psi$ on $w_\alpha \otimes v_\alpha$ can be computed as $(id_{V_x} \otimes \psi)(w_\alpha \otimes v_\alpha) = \lambda_\alpha w_\alpha \otimes v_\alpha$. In particular, the action of $id_{V_x} \otimes \psi$ preserves the image of $(\phi_x)_j$, i.e.,

$$
\operatorname{Im}((\operatorname{id}_{V_x} \otimes \psi)(\phi_x)_j) = \operatorname{Im}((\phi_x)_j). \tag{F5}
$$

Recalling that the idempotent $\sum_i (\phi_x)_i (\overline{\phi}_x)_i$ acts as the identity on $\text{Im}((\phi_x)_j)$ due to the orthogo-nality relation [\(33\)](#page-14-0), we find that it also acts as the identity on $\text{Im}((id_{V_{\kappa}} \otimes \psi)(\phi_x)_i)$. Thus, the weak completeness relation [\(F1\)](#page-72-5) holds. We note that the same proof applies to any diagonalizable ψ .

Appendix G: Hochschild cohomology

1. Proof of $\delta^2 = 0$

In Sec. [III B,](#page-24-0) we introduced a cohomology theory $H^k(\mathcal{C}, \mathbb{Z})$. In that section, we utilized the property $\delta^2 = 0$ without proving it. In this section, we show it explicitly:

Proposition 1. $\delta^2 = 0 : C^k(\mathcal{C}, \mathbb{Z}) \to C^{k+2}(\mathcal{C}, \mathbb{Z}).$

Proof.

Let *n* be an aribitrary element in $C^k(\mathcal{C}, \mathbb{Z})$ and $\{\rho_i\}_{i=1}^{k+2}$ be simple objects in \mathcal{C} . We compute

 $(\delta^2 n)(\rho_1, ..., \rho_{k+2})$ directly:

$$
(\delta^{2}n)(\rho_{1},...,\rho_{k+2})
$$
\n
$$
= \dim(\rho_{1})(\delta n)(\rho_{2},...,\rho_{k+1}) - (\delta n)(\rho_{1} \otimes \rho_{2},\rho_{3},...,\rho_{k+1}) + \sum_{i=2}^{k} (-1)^{i}(\delta n)(\rho_{1},...,\rho_{i} \otimes \rho_{i+1},...,\rho_{k+1})
$$
\n
$$
+ (-1)^{k+1}(\delta n)(\rho_{1},...,\rho_{k},\rho_{k+1} \otimes \rho_{k+2}) + (-1)^{k+2}(\delta n)(\rho_{1},...,\rho_{k+1}) \dim(\rho_{k+2}),
$$
\n
$$
= \dim(\rho_{1}) \{\dim(\rho_{2})n(\rho_{3},...,\rho_{k+2}) + \sum_{j=2}^{k+1} (-1)^{j-1}n(\rho_{2},...,\rho_{j} \otimes \rho_{j+1},...,\rho_{k+2})
$$
\n
$$
+ (-1)^{k+1}n(\rho_{2},...,\rho_{k+1}) \dim(\rho_{k+2})\} - \{\dim(\rho_{1} \otimes \rho_{2})n(\rho_{3},...,\rho_{k+2}) - n(\rho_{1} \otimes \rho_{2} \otimes \rho_{3},\rho_{4},...,\rho_{k+2})
$$
\n
$$
+ \sum_{j=3}^{k+1} (-1)^{j+1}n(\rho_{1} \otimes \rho_{2},...,\rho_{j} \otimes \rho_{j+1},...,\rho_{k+2}) + (-1)^{k}n(\rho_{1} \otimes \rho_{2},...,\rho_{k+1}) \dim(\rho_{k+2})\}
$$
\n
$$
+ \sum_{i=2}^{k} (-1)^{i} \{\dim(\rho_{1})n(\rho_{2},...,\rho_{i} \otimes \rho_{i+1},...,\rho_{k+2}) + \sum_{j=1}^{i-2} (-1)^{j}n(\rho_{1},...,\rho_{j} \otimes \rho_{j+1},...,\rho_{i} \otimes \rho_{i+1},...,\rho_{k+2})
$$
\n
$$
+ (-1)^{i+1}n(\rho_{1},...,\rho_{i-1} \otimes \rho_{i} \otimes \rho_{i+1},...,\rho_{k+2}) + (-1)^{i}n(\rho_{1},...,\rho_{i-1},\rho_{i} \otimes \rho_{i+1} \ot
$$

Since the expression is long, let us calculate it term by term. Note that the first line, the second and third lines, the fourth to sixth lines, the seventh and eighth lines, and the ninth line correspond to the first through fifth terms of the first equation, respectively. Let us denote them as $l_1, l_2, ..., l_5$:

$$
l_{1} = \dim(\rho_{1}) \{ \dim(\rho_{2})n(\rho_{3}, ..., \rho_{k+2}) + \sum_{j=2}^{k+1} (-1)^{j-1}n(\rho_{2}, ..., \rho_{j} \otimes \rho_{j+1}, ..., \rho_{k+2})
$$

\n
$$
+ (-1)^{k+1}n(\rho_{2}, ..., \rho_{k+1}) \dim(\rho_{k+2}) \},
$$

\n
$$
l_{2} = - \{ \dim(\rho_{1} \otimes \rho_{2})n(\rho_{3}, ..., \rho_{k+2}) - n(\rho_{1} \otimes \rho_{2} \otimes \rho_{3}, \rho_{4}, ..., \rho_{k+2})
$$

\n
$$
+ \sum_{j=3}^{k+1} (-1)^{j+1}n(\rho_{1} \otimes \rho_{2}, ..., \rho_{j} \otimes \rho_{j+1}, ..., \rho_{k+2}) + (-1)^{k}n(\rho_{1} \otimes \rho_{2}, ..., \rho_{k+1}) \dim(\rho_{k+2}) \},
$$

\n
$$
l_{3} = + \sum_{i=2}^{k} (-1)^{i} \{ \dim(\rho_{1})n(\rho_{2}, ..., \rho_{i} \otimes \rho_{i+1}, ..., \rho_{k+2})
$$

\n
$$
+ \sum_{j=1}^{k-2} (-1)^{j}n(\rho_{1}, ..., \rho_{j} \otimes \rho_{j+1}, ..., \rho_{k} \otimes \rho_{i+1}, ..., \rho_{k+2})
$$

\n
$$
+ (-1)^{i+1}n(\rho_{1}, ..., \rho_{i-1} \otimes \rho_{i} \otimes \rho_{i+1}, ..., \rho_{k+2}) + (-1)^{i}n(\rho_{1}, ..., \rho_{i-1}, \rho_{i} \otimes \rho_{i+1} \otimes \rho_{i+2}, ..., \rho_{k+2})
$$

\n
$$
+ \sum_{j=i+2}^{k+1} (-1)^{j+1}n(\rho_{1}, ..., \rho_{i} \otimes \rho_{i+1}, ..., \rho_{k+2}) \dim(\rho_{k+2}) \},
$$

\n
$$
l_{4} = + (-1)^{k+1} \{ \dim(\rho_{1})n(\rho_{2}, ..., \rho_{k}, \rho_{k+1} \otimes \rho_{k+2}) + \sum_{j=1}^{k-1} (-1)^{j}n(\rho_{1}, ..., \
$$

First, l_1 and l_5 cancel out the first and last terms of l_2 , l_3 , and l_4 . Let us denote the terms obtained by removing the first and last terms from l_2 , l_3 , and l_4 as l'_2 , l'_3 , and l'_4 , respectively:

$$
l'_{2} = -\{-n(\rho_{1} \otimes \rho_{2} \otimes \rho_{3}, \rho_{4}, ..., \rho_{k+2}) + \sum_{j=3}^{k+1} (-1)^{j+1} n(\rho_{1} \otimes \rho_{2}, ..., \rho_{j} \otimes \rho_{j+1}, ..., \rho_{k+2})\},
$$

\n
$$
l'_{3} = +\sum_{i=2}^{k} (-1)^{i} \{\sum_{j=1}^{i-2} (-1)^{j} n(\rho_{1}, ..., \rho_{j} \otimes \rho_{j+1}, ..., \rho_{i} \otimes \rho_{i+1}, ..., \rho_{k+2}) + (-1)^{i+1} n(\rho_{1}, ..., \rho_{i-1} \otimes \rho_{i} \otimes \rho_{i+1}, ..., \rho_{k+2}) + (-1)^{i} n(\rho_{1}, ..., \rho_{i-1}, \rho_{i} \otimes \rho_{i+1} \otimes \rho_{i+2}, ..., \rho_{k+2}) + \sum_{j=i+2}^{k+1} (-1)^{j+1} n(\rho_{1}, ..., \rho_{i} \otimes \rho_{i+1}, ..., \rho_{j} \otimes \rho_{j+1}, ..., \rho_{k+2})\},
$$

\n
$$
l'_{4} = (-1)^{k+1} \{\sum_{j=1}^{k-1} (-1)^{j} n(\rho_{1}, ..., \rho_{j} \otimes \rho_{j+1}, ..., \rho_{k}, \rho_{k+1} \otimes \rho_{k+2}) + (-1)^{k} n(\rho_{1}, ..., \rho_{k-1}, \rho_{k} \otimes \rho_{k+1} \otimes \rho_{k+2})\}.
$$

Let us focus on the second and third terms of l'_3 .

$$
\sum_{i=2}^{k} (-1)^{2i+1} n(\rho_1, ..., \rho_{i-1} \otimes \rho_i \otimes \rho_{i+1}, ..., \rho_{k+2}) + \sum_{i=2}^{k} (-1)^{2i} n(\rho_1, ..., \rho_{i-1}, \rho_i \otimes \rho_{i+1} \otimes \rho_{i+2}, ..., \rho_{k+2})
$$

= $-n(\rho_1 \otimes \rho_2 \otimes \rho_3, \rho_4, ..., \rho_{k+2}) + n(\rho_1, ..., \rho_k \otimes \rho_{k+1} \otimes \rho_{k+2}).$

This term cancels out the first term of l'_1 and the last term of l'_4 . Next, let us focus on the first and fourth terms of l'_3 :

$$
\sum_{i=2}^{k} \sum_{j=1}^{i-2} (-1)^{i+j} n(\rho_1, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_i \otimes \rho_{i+1}, ..., \rho_{k+2}) + \sum_{i=2}^{k} \sum_{j=i+2}^{k+1} (-1)^{i+j+1} n(\rho_1, ..., \rho_i \otimes \rho_{i+1}, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_{k+2})
$$
\n
$$
= \sum_{j=1}^{k-1} \sum_{i=j+2}^{k} (-1)^{i+j} n(\rho_1, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_i \otimes \rho_{i+1}, ..., \rho_{k+2}) - \sum_{i=2}^{k} \sum_{j=i+2}^{k+1} (-1)^{i+j} n(\rho_1, ..., \rho_i \otimes \rho_{i+1}, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_{k+2})
$$
\n
$$
= \sum_{i=1}^{k-1} \sum_{j=i+2}^{k} (-1)^{i+j} n(\rho_1, ..., \rho_i \otimes \rho_{i+1}, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_{k+2}) - \sum_{i=2}^{k} \sum_{j=i+2}^{k+1} (-1)^{i+j} n(\rho_1, ..., \rho_i \otimes \rho_{i+1}, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_{k+2})
$$
\n
$$
= \sum_{j=3}^{k} (-1)^{1+j} n(\rho_1 \otimes \rho_2, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_{k+2}) - \sum_{i=2}^{k-1} (-1)^{i+k+1} n(\rho_1, ..., \rho_i \otimes \rho_{i+1}, ..., \rho_{k+1} \otimes \rho_{k+2}).
$$

Here, we use $\sum_{i=2}^{k} \sum_{j=i+2}^{k+1} = \sum_{j=1}^{k-1} \sum_{i=j+2}^{k}$. On the other hand, the second term of l'_2 and the first term of l'_4 are

$$
-\sum_{j=3}^{k+1}(-1)^{j+1}n(\rho_1 \otimes \rho_2, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_{k+2}) + \sum_{j=1}^{k-1}(-1)^{j+k+1}n(\rho_1, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_k, \rho_{k+1} \otimes \rho_{k+2})
$$

=
$$
-\sum_{j=3}^k(-1)^{j+1}n(\rho_1 \otimes \rho_2, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_{k+2}) + \sum_{j=2}^{k-1}(-1)^{j+k+1}n(\rho_1, ..., \rho_j \otimes \rho_{j+1}, ..., \rho_k, \rho_{k+1} \otimes \rho_{k+2}).
$$

This term cancels out the contribution from the previous one. Consequently, $\delta^2 = 0$.

2. The 2nd Hochschild cohomology for $\text{Rep}(D_8)$

In this section, we show the following proposition:

Proposition 2.

$$
H^{2}(\text{Rep}(D_8), \mathbb{Z}) \simeq (\mathbb{Z}_2 \times \mathbb{Z}_2) \ltimes_{\omega} \mathbb{Z}_4,\tag{G1}
$$

where ω is a nontrivial element in $H^1((\mathbb{Z}_2)^2, \mathbb{Z}_4)^{\otimes 2} \subset H^2((\mathbb{Z}_2)^2, \mathbb{Z}_4)$

Proof. We denote the elements of D_8 as $D_8 = {\sigma^i \tau^j \mid \sigma^4 = 1, \tau^2 = 1, \sigma \tau = \tau \sigma^3}$, and denote the simple objects of $\text{Rep}(D_8)$ as $\text{Simp}(\text{Rep}(D_8)) = \{(0,0), (0,1), (1,0), (1,1), D\}$. Here, (a, b) represents the sign representation with $(\sigma, \tau) = ((-1)^a, (-1)^b)$ and D is the unique 2-dimensional representation of D_8 .

 \Box

Let $n \in \mathbb{Z}^2(\text{Rep}(D_8), \mathbb{Z})$ be a cocycle, i.e.

$$
(\delta n)(\rho_1, \rho_2, \rho_3) = 0,\tag{G2}
$$

for any simple objects ρ_1, ρ_2, ρ_3 . First, we consider the case where ρ_1, ρ_2, ρ_3 are one-dimensional representations. Since the one-dimensional representations follow the multiplication rule of the group $\mathbb{Z}_2 \times \mathbb{Z}_2$, the restrictions on n for one-dimensional representations are the same as that of the cocycles for the group cohomology of $\mathbb{Z}_2 \times \mathbb{Z}_2$. Then, the remaining degrees of freedom are

$$
n((0,0),(0,0)), \quad n((0,1),(0,1)), \quad n((1,0),(1,0)).
$$
 (G3)

The others are determined as

$$
n(\rho_1, \rho_2) = \begin{cases} n((0, 1), (0, 1)) + n((1, 0), (1, 0)) & \text{if } (\rho_1, \rho_2) = ((1, 1), (1, 1)), \\ n((0, 0), (0, 0)) & \text{otherwise,} \end{cases}
$$
(G4)

for one-dimensional representations (ρ_1, ρ_2) . These are the all constraints for the one-dimensional representations.

Similarly, for one-dimensional representation ρ , $n(\rho, D)$ and $n(D, \rho)$ are completely fixed by the cocycle condition as follows: Since

$$
(\delta n)((0,0), (0,0), D) = n((0,0), D) - n((0,0), D) + n((0,0), D) - 2n((0,0), (0,0)) = 0, (G5)
$$

$$
(\delta n)(D, (0,0), (0,0)) = 2n((0,0), (0,0)) - n(D, (0,0)) + n(D, (0,0)) - n(D, (0,0)) = 0, (G6)
$$

we obtain

$$
n((0,0),D) = n(D,(0,0)) = 2n((0,0),(0,0)).
$$
\n(G7)

Also, since

$$
(\delta n)(\rho, \rho, D) = n(\rho, D) - n((0, 0), D) + n(\rho, D) - 2n(\rho, \rho) = 0,
$$
 (G8)

$$
(\delta n)(D, \rho, \rho) = 2n(\rho, \rho) - n(D, \rho) + n(D, (0, 0)) - n(D, \rho) = 0,
$$
 (G9)

we obtain

$$
n(\rho, D) = n(\rho, \rho) - \frac{1}{2}n((0, 0), D) = n(\rho, \rho) - n((0, 0), (0, 0)),
$$
\n(G10)

$$
n(D,\rho) = n(\rho,\rho) - \frac{1}{2}n(D,(0,0)) = n(\rho,\rho) - n((0,0),(0,0)).
$$
\n(G11)

In contrast, there are no restrictions on $n(D, D)$. In summary, $n((0, 0), (0, 0)), n((0, 1), (0, 1)), n((1, 0), (1, 0))$ and $n(D, D)$ remain as undetermined degrees of freedom.

Next, let us consider the coboundary shift by $l \in C^1(\text{Rep}(D_8), \mathbb{Z})$. First, the coboundary action for $n((0, 0), (0, 0)), n((0, 1), (0, 1)), n((1, 0), (1, 0))$ are

$$
(\delta l)((0,0),(0,0)) = l((0,0)), \qquad (G12)
$$

$$
(\delta l)((0,1),(0,1)) = 2l((0,1))-l((0,0)), \qquad (G13)
$$

$$
(\delta l)((1,0),(1,0)) = 2l((1,0)) - l((0,0)). \tag{G14}
$$

The coboundary action for $n(D, D)$ is

$$
(\delta l)(D, D) = 4l(D) - l((0, 0)) - l((0, 1)) - l((1, 0)) - l((1, 1)).
$$
\n(G15)

Since we already fix $l((0,0)), l((0,1)), l((1,0)), l((1,1)),$ we cannot shift $n(D, D)$ by 1. However, by using $(\delta l)(D, D)$, we can shift it by multiples of 4. Thus, in cohomology theory, $n(D, D)$ is regarded as an element of \mathbb{Z}_4 . Consequently, we show that

$$
H^{2}(\text{Rep}(D_8), \mathbb{Z}) = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4,\tag{G16}
$$

as a set, where first and second \mathbb{Z}_2 is a choice of $n((0, 1), (0, 1)) = \pm 1$ and $n((1, 0), (1, 0)) = \pm 1$ respectively, and \mathbb{Z}_4 is a choice of $n(D, D) = 0, 1, 2, 3$.

Finally, we identify the group structure of $H^2(\text{Rep}(D_8), \mathbb{Z})$. Let us take $(1,0,0) \in \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$. This element is represented as

$$
n(\rho, \rho) = \begin{cases} 1, & \text{if } (\rho, \rho) = ((0, 1), (0, 1)), ((1, 1), (1, 1)), \\ 0, & \text{otherwise.} \end{cases}
$$
(G17)

Thus, multiplying this element by two results in

$$
n(\rho, \rho) = \begin{cases} 2, & \text{if } (\rho, \rho) = ((0, 1), (0, 1)), ((1, 1), (1, 1)), \\ 0, & \text{otherwise.} \end{cases}
$$
(G18)

We have decided to normalize $n((0,1),(0,1))$ and $n((1,1),(1,1))$ to 0 or 1 by using $l((0,1))$ and $l((1,1))$. Under this renormalization, $n(D, D)$ shifts by 2. In other words, this means that the element of the first \mathbb{Z}_2 has carried over to become 2 in \mathbb{Z}_4 . Also, the second \mathbb{Z}_2 exhibits the same carry-over structure. Therefore, $H^2(\text{Rep}(D_8), \mathbb{Z})$ is a nontrivial extension of the $\mathbb{Z}_2 \times \mathbb{Z}_2$ by \mathbb{Z}_4 .

Appendix H: Computation of pump invariant for $\text{Rep}(D_8)$

Let us compute the pump invariant for S^1 -parameterized families constructed from the $D_8 \times$ $Rep(D_8)$ cluster state. We note that the pump invariant is already computed in Sec. VC in the case of the general $G \times \text{Rep}(G)$ cluster state. The computations there were performed in a gauge where the action tensors $\hat{\phi}_{\rho}(\theta)$ for $\rho \in \text{Rep}(G)$ are 2π -periodic. In this appendix, we do a similar computation in a gauge where the MPS tensor $A(\theta)$ is 2π -periodic. Furthermore, we not only compute the invariant but also write down $A(\theta)$ and $\hat{\phi}_{\rho}(\theta)$ explicitly for all $\theta \in S^1$.

The dihedral group D_8 of order 8 is defined by

$$
D_8 = \langle \sigma, \tau \mid \sigma^4 = \tau^2 = 1, \sigma\tau = \tau\sigma^3 \rangle.
$$
 (H1)

Under the right regular representation of D_8 , the generators σ and τ are given as follows:

$$
R(\sigma) = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}, \ R(\tau) = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}, \tag{H2}
$$

We denote by $e_{i,j}$ the 8×8 matrix with 1 in the (i, j) entry and 0 elsewhere. By using these matrices, an MPS representation of the $G \times \text{Rep}(G)$ cluster state is given by

$$
A_o^g = e_{g,g} \tag{H3}
$$

for odd sites and

$$
A_e^g = R(g) \tag{H4}
$$

for even sites $[41]$ and the translationally invariant MPS is given by

$$
A_{oe}^{g,h} := A_o^g A_e^h = e_{g,g} R(h) = e_{g,h^{-1}g}.
$$
\n(H5)

We note that this MPS matrix is injective.

Since the parameterization is introduced by a unitary transformation acting on the physical legs (see Eq. [\(148\)](#page-41-1)), the ground state of $H_g(\theta)$ is given by

$$
\left(\prod_{i:\text{odd}} R_g(\theta)_i\right) \left|\left\{A_{oe}^{g,h}\right\}\right\rangle_L,\tag{H6}
$$

where the subscript L denotes the number of unit cells. Therefore, we can easily get an MPS representation of the ground state as follows:

$$
\left(\prod_{i=1}^{L} R_g(\theta)_i\right) \left| \{A_{oe}^{g,h}\}\right\rangle_L = \sum_{\{g_k,h_k\}} \text{tr}\left(A_{oe}^{g_1,h_1} \cdots A_{oe}^{g_L,h_L}\right) \left(R_g(\theta)_1 \left|g_1\right\rangle\right) \left|h_1\right\rangle \cdots \left(R_g(\theta)_L \left|g_L\right\rangle\right) \left|h_L\right\rangle
$$
\n
$$
= \sum_{\{g_k,h_k\}} \text{tr}\left(A_{oe}^{g_1,h_1} \cdots A_{oe}^{g_L,h_L}\right) \left(\sum_{\tilde{g}_1} |\tilde{g}_1\rangle \left[R_g(\theta)_1\right]_{\tilde{g}_1,g_1}\right) \left|h_1\right\rangle \cdots \left(\sum_{\tilde{g}_L} |\tilde{g}_L\rangle \left[R_g(\theta)_L\right]_{\tilde{g}_L,g_L}\right) \left|h_L\right\rangle
$$
\n
$$
= \sum_{\{\tilde{g}_k,h_k\}} \text{tr}\left(A_{oe}^{\tilde{g}_1,h_1}(\theta;g) \cdots A_{oe}^{\tilde{g}_L,h_L}(\theta;g)\right) \left|\tilde{g}_1,h_1,\cdots,\tilde{g}_L,h_L\right\rangle,
$$

where

$$
A_{oe}^{g,h}(\theta;k) = \sum_{\tilde{g}} [R_k(\theta)]_{g,\tilde{g}} A_{oe}^{\tilde{g},h}.
$$
 (H7)

Let us compute the invariant using a gauge where $\phi(\theta)$ is non-periodic. Although $A_{oe}^{g,h}(\theta;k)$ is not 2π -periodic as a matrix, the 2π -periodicity of the MPS guarantees the existence of the transition function \hat{U}_k :

$$
A_{oe}^{g,h}(2\pi;k) = \hat{U}_k A_{oe}^{g,h}(0;k) \hat{U}_k^{\dagger}.
$$
 (H8)

By explicit calculations, we can check that

$$
\hat{U}_{\sigma} = \begin{pmatrix}\n0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 1 & 0 & 0 & 0\n\end{pmatrix}, \quad \hat{U}_{\tau} = \begin{pmatrix}\n0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 1\n\end{pmatrix}, \quad (H9)
$$

and \hat{U}_k is given by a multiplication of them for general $k \in D_8$. We interpolate between \hat{U}_k and the identity matrix as

$$
\hat{U}_k(\theta) := \exp\left(\frac{\theta}{2\pi} \log(\hat{U}_k)\right),\tag{H10}
$$

and define a 2π -periodic MPS matrix as

$$
\tilde{A}_{oe}^{g,h}(\theta;k) = \hat{U}_k(\theta)^\dagger A_{oe}^{g,h}(\theta;k)\hat{U}_k(\theta). \tag{H11}
$$

Note that the L-symbol calculated from this MPS is constant. This is due to the following reasons: First, for the MPS before making it 2π -periodic, the transfer matrix is constant. Thus, we can choose a gauge so that the action tensor is constant. Therefore, the L-symbol calculated in this gauge is constant. Since the L-symbol does not change under the gauge transformation Eq. [\(H11\)](#page-80-0), the L-symbol calculated with $\tilde{A}_{oe}^{g,h}(\theta;k)$ also constant.

Let us compute the fractionalized symmetry operator $\hat{\phi}^k_{\rho}(\theta)$ for $\rho \in \text{Rep}(D_8)$ defined by

$$
\sum_{\tilde{h}} \mathcal{O}_{\rho}^{h\tilde{h}} \tilde{A}_{oe}^{g,\tilde{h}}(\theta; k) = \hat{\phi}^{k}(\theta)_{\rho}^{-1} \tilde{A}_{oe}^{g,h}(\theta; k) \hat{\phi}^{k}(\theta)_{\rho}.
$$
 (H12)

By explicit calculations, we obtain

$$
\hat{\phi}^{\sigma}(\theta)_{(0,0)} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, \quad \hat{\phi}^{\sigma}(\theta)_{(0,1)} = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad (H13)
$$

$$
\hat{\phi}^{\sigma}(\theta)_{(1,0)} = \begin{pmatrix}\n-c\frac{\theta}{2} & 0 & \frac{1+i}{2}s\frac{\theta}{2} & 0 & 0 & 0 & \frac{-1+i}{2}s\frac{\theta}{2} & 0 \\
0 & -c\frac{\theta}{2} & 0 & \frac{-1+i}{2}s\frac{\theta}{2} & 0 & 0 & 0 & \frac{1+i}{2}s\frac{\theta}{2} \\
\frac{1-i}{2}s\frac{\theta}{2} & 0 & c\frac{\theta}{2} & 0 & -\frac{1+i}{2}s\frac{\theta}{2} & 0 & 0 & 0 \\
0 & -\frac{1+i}{2}s\frac{\theta}{2} & 0 & c\frac{\theta}{2} & 0 & \frac{1-i}{2}s\frac{\theta}{2} & 0 & 0 \\
0 & 0 & \frac{-1+i}{2}s\frac{\theta}{2} & 0 & -c\frac{\theta}{2} & 0 & \frac{1+i}{2}s\frac{\theta}{2} & 0 \\
-\frac{1+i}{2}s\frac{\theta}{2} & 0 & 0 & 0 & \frac{1+i}{2}s\frac{\theta}{2} & 0 & -c\frac{\theta}{2} & 0 & \frac{-1+i}{2}s\frac{\theta}{2} \\
0 & \frac{1-i}{2}s\frac{\theta}{2} & 0 & 0 & 0 & \frac{1-i}{2}s\frac{\theta}{2} & 0 & c\frac{\theta}{2} & 0 \\
0 & \frac{1-i}{2}s\frac{\theta}{2} & 0 & 0 & 0 & -\frac{1+i}{2}s\frac{\theta}{2} & 0 & c\frac{\theta}{2}\n\end{pmatrix}, \quad (H14)
$$

$$
\hat{\phi}^{\sigma}(\theta)_{(1,1)} = \begin{pmatrix} c_{\frac{\theta}{2}}^{{\theta}} & 0 & -\frac{1+i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & 0 & 0 & \frac{1-i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 \\ 0 & -c_{\frac{\theta}{2}}^{{\theta}} & 0 & \frac{-1+i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & 0 & 0 & \frac{1+i}{2}s_{\frac{\theta}{2}}^{{\theta}} \\ \frac{-1+i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & -c_{\frac{\theta}{2}}^{{\theta}} & 0 & \frac{1+i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & 0 & 0 \\ 0 & -\frac{1+i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & c_{\frac{\theta}{2}}^{{\theta}} & 0 & \frac{1-i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & 0 \\ 0 & 0 & \frac{1-i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & c_{\frac{\theta}{2}}^{{\theta}} & 0 & -c_{\frac{\theta}{2}}^{{\theta}} & 0 & \frac{-1+i}{2}s_{\frac{\theta}{2}}^{{\theta}} \\ \frac{1+i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & 0 & 0 & \frac{-1+i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & -c_{\frac{\theta}{2}}^{{\theta}} & 0 \\ 0 & \frac{1-i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & 0 & 0 & -\frac{1+i}{2}s_{\frac{\theta}{2}}^{{\theta}} & 0 & c_{\frac{\theta}{2}}^{{\theta}} \end{pmatrix}, \quad (H15)
$$

$$
\hat{\phi}^{\sigma}(\theta)_{D}=\begin{pmatrix} s^{\frac{3}{2}} & c^{\frac{3}{2}}q & 0 & 0 & \frac{\omega_{3}^{2}}{2}s_{\frac{3}{2}}^{2}g^{\frac{3}{2}+\frac{\omega_{4}}{4}}-\frac{\omega_{4}^{3}}{2}s_{\frac{3}{2}}^{2}g^{\frac{2}{2}+\frac{\omega_{4}}{4}} & 0 & 0 & 0\\ 0 & 0 & c^{\frac{3}{2}}q & s^{\frac{3}{2}}q & 0 & 0 & \frac{\omega_{4}^{3}}{2}s_{\frac{3}{2}}^{2}g^{\frac{3}{2}+\frac{\omega_{4}}{4}} & 0 & 0 & \frac{\omega_{4}^{3}}{2}s_{\frac{3}}^{2}g^{\frac{3}{2}+\frac{\omega_{4}}{4}} & 0
$$

,

and

$$
\hat{\phi}^{\tau}(\theta)_{(0,0)} = \begin{pmatrix}\n1 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & 0\n\end{pmatrix}, \quad \hat{\phi}^{\tau}(\theta)_{(0,1)} = \begin{pmatrix}\n-c\frac{\theta}{2} & is\frac{\theta}{2} & 0 & 0 & 0 & 0 & 0 & 0 \\
-is\frac{\theta}{2} & c\frac{\theta}{2} & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & -s\frac{\theta}{2} & is\frac{\theta}{2} & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & -s\frac{\theta}{2} & s\frac{\theta}{2} & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & -s\frac{\theta}{2} & s\frac{\theta}{2}\n\end{pmatrix},
$$
\n
$$
\hat{\phi}^{\tau}(\theta)_{(1,0)} = \begin{pmatrix}\n-1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & 0\n\end{pmatrix}, \quad \hat{\phi}^{\tau}(\theta)_{(1,1)} = \begin{pmatrix}\nc\frac{\theta}{2} & -is\frac{\theta}{2} & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & s\frac{\theta}{2} & -c\frac{\theta}{2} & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0
$$

Here, s and c represent sin and cos, respectively, and ω_8 is the shorthand notation for $e^{2\pi i/8}$, and A and B are defined by

$$
A = \begin{pmatrix} \sin^2 \frac{\theta}{4} & \cos^2 \frac{\theta}{4} & \frac{1}{2} i \sin \frac{\theta}{2} & -\frac{1}{2} i \sin \frac{\theta}{2} \\ -\cos^2 \frac{\theta}{4} & -\sin^2 \frac{\theta}{4} & \frac{1}{2} i \sin \frac{\theta}{2} & -\frac{1}{2} i \sin \frac{\theta}{2} \\ -\frac{1}{2} i \sin \frac{\theta}{2} & \frac{1}{2} i \sin \frac{\theta}{2} & \cos^2 \frac{\theta}{4} & \sin^2 \frac{\theta}{4} \\ -\frac{1}{2} i \sin \frac{\theta}{2} & \frac{1}{2} i \sin \frac{\theta}{2} & -\sin^2 \frac{\theta}{4} & -\cos^2 \frac{\theta}{4} \end{pmatrix}, B = \begin{pmatrix} \cos^2 \frac{\theta}{4} & -\sin^2 \frac{\theta}{4} & -\frac{1}{2} i \sin \frac{\theta}{2} & -\frac{1}{2} i \sin \frac{\theta}{2} \\ -\sin^2 \frac{\theta}{4} & \cos^2 \frac{\theta}{4} & -\frac{1}{2} i \sin \frac{\theta}{2} & -\frac{1}{2} i \sin \frac{\theta}{2} \\ \frac{1}{2} i \sin \frac{\theta}{2} & \frac{1}{2} i \sin \frac{\theta}{2} & \sin^2 \frac{\theta}{4} & -\cos^2 \frac{\theta}{4} \end{pmatrix}.
$$

The action tensor $\hat{\phi}^k_{\rho}(\theta)$ for general $k \in D_8$ is given by a multiplication of the above matrices. The pump invariant is the difference of $\hat{\phi}^k_{\rho}(\theta)$ at $\theta = 0$ and $\theta = 2\pi$:

$$
\hat{\phi}^k(2\pi)_\rho = \eta_\rho^k \hat{\phi}_\rho^k(0). \tag{H16}
$$

By using the above matrices, we can explicitly read off the pump invariant:

$$
\eta_{(0,0)}^{\sigma} = 1, \quad \eta_{(0,1)}^{\sigma} = 1, \quad \eta_{(1,0)}^{\sigma} = -1, \quad \eta_{(1,1)}^{\sigma} = -1, \quad \eta_{D}^{\sigma} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},
$$

$$
\eta_{(0,0)}^{\tau} = 1, \quad \eta_{(0,1)}^{\tau} = -1, \quad \eta_{(1,0)}^{\tau} = 1, \quad \eta_{(1,1)}^{\tau} = -1, \quad \eta_{D}^{\tau} = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix},
$$
(H17)

The pump invariant η_{ρ}^{k} for general $k \in D_{8}$ is given by a multiplication of the above invariants, i.e.,

$$
\eta_{\rho}^{\sigma^i \tau^j} = (\eta_{\rho}^{\sigma})^i (\eta_{\rho}^{\tau})^j. \tag{H18}
$$

According to the Tannaka-Krein duality, it is expected that the pump invariant is classified by D_8 . This is consistent with the fact that the invariants we computed form D_8 as shown in Eq. [\(H18\)](#page-84-0). Our computation demonstrates that all (equivalence classes of) $S¹$ -parameterized families can be constructed in this way.

- [1] Z.-C. Gu and X.-G. Wen, Tensor-entanglement-filtering renormalization approach and symmetryprotected topological order, Physical Review B 80, [10.1103/physrevb.80.155131](https://doi.org/10.1103/physrevb.80.155131) (2009), [arXiv:0903.1069 \[cond-mat.str-el\].](https://arxiv.org/abs/0903.1069)
- [2] F. Pollmann, E. Berg, A. M. Turner, and M. Oshikawa, Symmetry protection of topological phases in one-dimensional quantum spin systems, Phys. Rev. B 85[, 075125 \(2012\),](https://doi.org/10.1103/PhysRevB.85.075125) [arXiv:0909.4059 \[cond](https://arxiv.org/abs/0909.4059)[mat.str-el\].](https://arxiv.org/abs/0909.4059)
- [3] F. Pollmann, A. M. Turner, E. Berg, and M. Oshikawa, Entanglement spectrum of a topological phase in one dimension, Physical Review B 81, [10.1103/physrevb.81.064439](https://doi.org/10.1103/physrevb.81.064439) (2010), [arXiv:0910.1811 \[cond](https://arxiv.org/abs/0910.1811)[mat.str-el\].](https://arxiv.org/abs/0910.1811)
- [4] X. Chen, Z.-C. Gu, and X.-G. Wen, Classification of gapped symmetric phases in one-dimensional spin systems, Phys. Rev. B 83[, 035107 \(2011\),](https://doi.org/10.1103/PhysRevB.83.035107) [arXiv:1008.3745 \[cond-mat.str-el\].](https://arxiv.org/abs/1008.3745)
- [5] X. Chen, Z.-C. Gu, and X.-G. Wen, Complete classification of one-dimensional gapped quantum phases in interacting spin systems, Physical Review B 84, [10.1103/physrevb.84.235128](https://doi.org/10.1103/physrevb.84.235128) (2011), [arXiv:1103.3323](https://arxiv.org/abs/1103.3323) [\[cond-mat.str-el\].](https://arxiv.org/abs/1103.3323)
- [6] X. Chen, Z.-X. Liu, and X.-G. Wen, Two-dimensional symmetry-protected topological orders and their protected gapless edge excitations, Phys. Rev. B 84[, 235141 \(2011\),](https://doi.org/10.1103/PhysRevB.84.235141) [arXiv:1106.4752 \[cond-mat.str-el\].](https://arxiv.org/abs/1106.4752)
- [7] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, Symmetry protected topological orders and the group cohomology of their symmetry group, Phys. Rev. B 87[, 155114 \(2013\),](https://doi.org/10.1103/PhysRevB.87.155114) [arXiv:1106.4772 \[cond-mat.str](https://arxiv.org/abs/1106.4772)[el\].](https://arxiv.org/abs/1106.4772)
- [8] N. Schuch, D. Pérez-García, and I. Cirac, Classifying quantum phases using matrix product states and projected entangled pair states, Physical Review B 84, [10.1103/physrevb.84.165139](https://doi.org/10.1103/physrevb.84.165139) (2011), [arXiv:1010.3732 \[cond-mat.str-el\].](https://arxiv.org/abs/1010.3732)
- [9] D. J. Thouless, Quantization of particle transport, Phys. Rev. B 27[, 6083 \(1983\).](https://doi.org/10.1103/PhysRevB.27.6083)
- [10] J. C. Y. Teo and C. L. Kane, Topological defects and gapless modes in insulators and superconductors, Phys. Rev. B 82[, 115120 \(2010\).](https://doi.org/10.1103/PhysRevB.82.115120)
- [11] A. Kitaev, [Toward a topological classification of many-body quantum states with short-range entan](http://scgp.stonybrook.edu/video_portal/video.php?id=336)[glement](http://scgp.stonybrook.edu/video_portal/video.php?id=336) (2011), talk at Simons Center for Geometry and Physics.
- [12] A. Kitaev, [On the classification of short-range entangled states](http://scgp.stonybrook.edu/video_portal/video.php?id=2010) (2013), talk at Simons Center for Geometry and Physics.
- [13] A. Kitaev, [Homotopy-theoretic approach to spt phases in action:](https://www.ipam.ucla.edu/abstract/?tid=12389) z_{16} classification of three-dimensional [superconductors](https://www.ipam.ucla.edu/abstract/?tid=12389) (2015), talk at Institute for Pure and Applied Mathematics.
- [14] A. Kapustin and L. Spodyneiko, Higher-dimensional generalizations of the Thouless charge pump (2020), [arXiv:2003.09519 \[cond-mat.str-el\].](https://arxiv.org/abs/2003.09519)
- [15] K. Shiozaki, Adiabatic cycles of quantum spin systems, Phys. Rev. B 106[, 125108 \(2022\),](https://doi.org/10.1103/PhysRevB.106.125108) [arXiv:2110.10665 \[cond-mat.str-el\].](https://arxiv.org/abs/2110.10665)
- [16] M. Hermele, [Families of gapped systems and quantum pumps](https://www.youtube.com/watch?v=wtaC0tqXZMU) (2021), talk at Harvard CMSA.
- [17] X. Wen, M. Qi, A. Beaudry, J. Moreno, M. J. Pflaum, D. Spiegel, A. Vishwanath, and M. Hermele, Flow of higher Berry curvature and bulk-boundary correspondence in parametrized quantum systems, Phys. Rev. B 108[, 125147 \(2023\),](https://doi.org/10.1103/PhysRevB.108.125147) [arXiv:2112.07748 \[cond-mat.str-el\].](https://arxiv.org/abs/2112.07748)
- [18] L. Spodyneiko, Hall conductivity pump (2023), [arXiv:2309.14332 \[cond-mat.mes-hall\].](https://arxiv.org/abs/2309.14332)
- [19] S. Ohyama, K. Shiozaki, and M. Sato, Generalized Thouless pumps in (1+1)-dimensional interacting fermionic systems, Phys. Rev. B 106[, 165115 \(2022\),](https://doi.org/10.1103/PhysRevB.106.165115) [arXiv:2206.01110 \[cond-mat.str-el\].](https://arxiv.org/abs/2206.01110)
- [20] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, Generalized Global Symmetries, [JHEP](https://doi.org/10.1007/JHEP02(2015)172) 02, 172, [arXiv:1412.5148 \[hep-th\].](https://arxiv.org/abs/1412.5148)
- [21] C. C´ordova, T. T. Dumitrescu, K. Intriligator, and S.-H. Shao, Snowmass White Paper: Generalized Symmetries in Quantum Field Theory and Beyond, in Snowmass 2021 (2022) [arXiv:2205.09545 \[hep](https://arxiv.org/abs/2205.09545)[th\].](https://arxiv.org/abs/2205.09545)
- [22] J. McGreevy, Generalized Symmetries in Condensed Matter, [Ann. Rev. Condensed Matter Phys.](https://doi.org/10.1146/annurev-conmatphys-040721-021029) 14, [57 \(2023\),](https://doi.org/10.1146/annurev-conmatphys-040721-021029) [arXiv:2204.03045 \[cond-mat.str-el\].](https://arxiv.org/abs/2204.03045)
- [23] S. Schäfer-Nameki, ICTP lectures on (non-)invertible generalized symmetries, [Phys. Rept.](https://doi.org/10.1016/j.physrep.2024.01.007) 1063, 1 [\(2024\),](https://doi.org/10.1016/j.physrep.2024.01.007) [arXiv:2305.18296 \[hep-th\].](https://arxiv.org/abs/2305.18296)
- [24] T. D. Brennan and S. Hong, Introduction to Generalized Global Symmetries in QFT and Particle Physics (2023), [arXiv:2306.00912 \[hep-ph\].](https://arxiv.org/abs/2306.00912)
- [25] L. Bhardwaj, L. E. Bottini, L. Fraser-Taliente, L. Gladden, D. S. W. Gould, A. Platschorre, and H. Tillim, Lectures on generalized symmetries, [Phys. Rept.](https://doi.org/10.1016/j.physrep.2023.11.002) 1051, 1 (2024), [arXiv:2307.07547 \[hep-th\].](https://arxiv.org/abs/2307.07547)
- [26] R. Luo, Q.-R. Wang, and Y.-N. Wang, Lecture notes on generalized symmetries and applications, [Phys.](https://doi.org/10.1016/j.physrep.2024.02.002) Rept. 1065[, 1 \(2024\),](https://doi.org/10.1016/j.physrep.2024.02.002) [arXiv:2307.09215 \[hep-th\].](https://arxiv.org/abs/2307.09215)
- [27] S.-H. Shao, What's Done Cannot Be Undone: TASI Lectures on Non-Invertible Symmetries (2023), [arXiv:2308.00747 \[hep-th\].](https://arxiv.org/abs/2308.00747)
- [28] N. Carqueville, M. Del Zotto, and I. Runkel, Topological defects (2023), [arXiv:2311.02449 \[math-ph\].](https://arxiv.org/abs/2311.02449)
- [29] N. Iqbal, Jena lectures on generalized global symmetries: principles and applications (2024) [arXiv:2407.20815 \[hep-th\].](https://arxiv.org/abs/2407.20815)
- [30] L. Bhardwaj and Y. Tachikawa, On finite symmetries and their gauging in two dimensions, [JHEP](https://doi.org/10.1007/JHEP03(2018)189) 03, [189,](https://doi.org/10.1007/JHEP03(2018)189) [arXiv:1704.02330 \[hep-th\].](https://arxiv.org/abs/1704.02330)
- [31] C.-M. Chang, Y.-H. Lin, S.-H. Shao, Y. Wang, and X. Yin, Topological Defect Lines and Renormalization Group Flows in Two Dimensions, [JHEP](https://doi.org/10.1007/JHEP01(2019)026) 01, 026, [arXiv:1802.04445 \[hep-th\].](https://arxiv.org/abs/1802.04445)
- [32] R. Thorngren and Y. Wang, Fusion category symmetry. Part I. Anomaly in-flow and gapped phases, [JHEP](https://doi.org/10.1007/JHEP04(2024)132) 04, 132, [arXiv:1912.02817 \[hep-th\].](https://arxiv.org/abs/1912.02817)
- [33] J. I. Cirac, D. Pérez-García, N. Schuch, and F. Verstraete, Matrix product states and projected entangled pair states: Concepts, symmetries, theorems, [Rev. Mod. Phys.](https://doi.org/10.1103/RevModPhys.93.045003) 93, 045003 (2021), [arXiv:2011.12127 \[quant-ph\].](https://arxiv.org/abs/2011.12127)
- [34] N. Bultinck, M. Mariën, D. J. Williamson, M. B. Sahinoğlu, J. Haegeman, and F. Verstraete, Anyons and matrix product operator algebras, [Annals Phys.](https://doi.org/10.1016/j.aop.2017.01.004) 378, 183 (2017), [arXiv:1511.08090 \[cond-mat.str](https://arxiv.org/abs/1511.08090)[el\].](https://arxiv.org/abs/1511.08090)
- [35] L. Fidkowski and A. Kitaev, Topological phases of fermions in one dimension, Physical Review B 83, [10.1103/physrevb.83.075103](https://doi.org/10.1103/physrevb.83.075103) (2011), [arXiv:1008.4138 \[cond-mat.str-el\].](https://arxiv.org/abs/1008.4138)
- [36] A. Molnar, A. R. de Alarcón, J. Garre-Rubio, N. Schuch, J. I. Cirac, and D. Pérez-García, Matrix product operator algebras I: representations of weak Hopf algebras and projected entangled pair states (2022), [arXiv:2204.05940 \[quant-ph\].](https://arxiv.org/abs/2204.05940)
- [37] J. Garre-Rubio, L. Lootens, and A. Molnár, Classifying phases protected by matrix product operator symmetries using matrix product states, Quantum 7[, 927 \(2023\),](https://doi.org/10.22331/q-2023-02-21-927) [arXiv:2203.12563 \[cond-mat.str-el\].](https://arxiv.org/abs/2203.12563)
- [38] L. Lootens, C. Delcamp, G. Ortiz, and F. Verstraete, Dualities in One-Dimensional Quantum Lattice Models: Symmetric Hamiltonians and Matrix Product Operator Intertwiners, [PRX Quantum](https://doi.org/10.1103/PRXQuantum.4.020357) 4, 020357 [\(2023\),](https://doi.org/10.1103/PRXQuantum.4.020357) [arXiv:2112.09091 \[quant-ph\].](https://arxiv.org/abs/2112.09091)
- [39] L. Lootens, C. Delcamp, and F. Verstraete, Dualities in One-Dimensional Quantum Lattice Models: Topological Sectors, PRX Quantum 5[, 010338 \(2024\),](https://doi.org/10.1103/PRXQuantum.5.010338) [arXiv:2211.03777 \[quant-ph\].](https://arxiv.org/abs/2211.03777)
- [40] P. Gorantla, S.-H. Shao, and N. Tantivasadakarn, Tensor networks for non-invertible symmetries in 3+1d and beyond (2024), [arXiv:2406.12978 \[quant-ph\].](https://arxiv.org/abs/2406.12978)
- [41] C. Fechisin, N. Tantivasadakarn, and V. V. Albert, Non-invertible symmetry-protected topological order in a group-based cluster state (2023), [arXiv:2312.09272 \[cond-mat.str-el\].](https://arxiv.org/abs/2312.09272)
- [42] S. Seifnashri and S.-H. Shao, Cluster state as a non-invertible symmetry protected topological phase (2024), [arXiv:2404.01369 \[cond-mat.str-el\].](https://arxiv.org/abs/2404.01369)
- [43] Z. Jia, Generalized cluster states from Hopf algebras: non-invertible symmetry and Hopf tensor network representation (2024), [arXiv:2405.09277 \[quant-ph\].](https://arxiv.org/abs/2405.09277)
- [44] Y. Choi, Y. Sanghavi, S.-H. Shao, and Y. Zheng, Non-invertible and higher-form symmetries in 2+1d lattice gauge theories (2024), [arXiv:2405.13105 \[cond-mat.str-el\].](https://arxiv.org/abs/2405.13105)
- [45] Y. Li and M. Litvinov, Non-invertible SPT, gauging and symmetry fractionalization (2024), [arXiv:2405.15951 \[cond-mat.str-el\].](https://arxiv.org/abs/2405.15951)
- [46] C. G. Brell, Generalized cluster states based on finite groups, [New Journal of Physics](https://doi.org/10.1088/1367-2630/17/2/023029) 17, 023029 (2015), [arXiv:1408.6237 \[quant-ph\].](https://arxiv.org/abs/1408.6237)
- [47] K. Inamura, On lattice models of gapped phases with fusion category symmetries, [JHEP](https://doi.org/10.1007/JHEP03(2022)036) 03, 036, [arXiv:2110.12882 \[cond-mat.str-el\].](https://arxiv.org/abs/2110.12882)
- [48] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, [Tensor Categories,](https://doi.org/10.1090/surv/205) Mathematical Surveys and Monographs, Vol. 205 (American Mathematical Society, Providence, RI, 2015).
- [49] G. W. Moore and N. Seiberg, Classical and Quantum Conformal Field Theory, [Commun. Math. Phys.](https://doi.org/10.1007/BF01238857) 123[, 177 \(1989\).](https://doi.org/10.1007/BF01238857)
- [50] S. Mac Lane, [Categories for the Working Mathematician,](https://doi.org/10.1007/978-1-4757-4721-8) 2nd ed., Grad. Texts Math., Vol. 5 (New York, NY: Springer, 1998).
- [51] V. Ostrik, Module categories, weak Hopf algebras and modular invariants, [Transformation Groups](https://doi.org/10.1007/s00031-003-0515-6) 8, [177 \(2003\),](https://doi.org/10.1007/s00031-003-0515-6) [arXiv:math/0111139 \[math.QA\].](https://arxiv.org/abs/math/0111139)
- [52] T. Hayashi, A canonical Tannaka duality for finite seimisimple tensor categories (1999), [arXiv:math/9904073 \[math.QA\].](https://arxiv.org/abs/math/9904073)
- [53] G. Böhm, F. Nill, and K. Szlachányi, Weak Hopf Algebras: I. Integral Theory and C^* -Structure, [Journal of Algebra](https://doi.org/https://doi.org/10.1006/jabr.1999.7984) 221, 385 (1999), [arXiv:math/9805116 \[math.QA\].](https://arxiv.org/abs/math/9805116)
- [54] K. Inamura, Topological field theories and symmetry protected topological phases with fusion category symmetries, [JHEP](https://doi.org/10.1007/JHEP05(2021)204) 05, 204, [arXiv:2103.15588 \[cond-mat.str-el\].](https://arxiv.org/abs/2103.15588)
- [55] Z. Komargodski, K. Ohmori, K. Roumpedakis, and S. Seifnashri, Symmetries and strings of adjoint QCD2, [JHEP](https://doi.org/10.1007/JHEP03(2021)103) 03, 103, [arXiv:2008.07567 \[hep-th\].](https://arxiv.org/abs/2008.07567)
- [56] T.-C. Huang, Y.-H. Lin, and S. Seifnashri, Construction of two-dimensional topological field theories with non-invertible symmetries, [JHEP](https://doi.org/10.1007/JHEP12(2021)028) 12, 028, [arXiv:2110.02958 \[hep-th\].](https://arxiv.org/abs/2110.02958)
- [57] T. Tannaka, Uber den Dualitätssatz der nichtkommutativen topologischen Gruppen, Tohoku Mathematical Journal, First Series 45, 1 (1939).
- [58] M. Krein, A principle of duality for bicompact groups and quadratic block algebras, Doklady AN SSSR 69, 725 (1949).
- [59] K.-H. Ulbrich, On Hopf algebras and rigid monoidal categories, [Israel Journal of Mathematics](https://doi.org/10.1007/BF02764622) 72, 252 [\(1990\).](https://doi.org/10.1007/BF02764622)
- [60] A. Joyal and R. Street, An introduction to Tannaka duality and quantum groups, in [Category Theory,](https://link.springer.com/chapter/10.1007/BFb0084235) edited by A. Carboni, M. C. Pedicchio, and G. Rosolini (Springer Berlin Heidelberg, Berlin, Heidelberg, 1991) pp. 413–492.
- [61] P. Schauenburg, Tannaka duality for arbitrary Hopf algebras, Algebra Berichte 66 (1992).
- [62] R. Thorngren, [Berry phase, diabolical points, and pivot hamiltonians](https://www.ms.u-tokyo.ac.jp/~yasuyuki/yitp2021.htm) (2021), talk at Yukawa Institute for Theoretical Physics.
- [63] S. Bachmann, W. De Roeck, M. Fraas, and T. Jappens, A Classification of G-Charge Thouless Pumps in 1D Invertible States, [Commun. Math. Phys.](https://doi.org/10.1007/s00220-024-05010-w) 405, 157 (2024), [arXiv:2204.03763 \[math-ph\].](https://arxiv.org/abs/2204.03763)
- [64] P. Etingof and S. Gelaki, Isocategorical groups, International Mathematics Research Notices 2001, 59 (2001), [arXiv:math/0007196 \[math.QA\].](https://arxiv.org/abs/math/0007196)
- [65] M. Fannes, B. Nachtergaele, and R. F. Werner, Finitely correlated states on quantum spin chains, Communications in Mathematical Physics 144, 443 (1992).
- [66] F. Verstraete and J. I. Cirac, Matrix product states represent ground states faithfully, [Phys. Rev. B](https://doi.org/10.1103/PhysRevB.73.094423) 73[, 094423 \(2006\),](https://doi.org/10.1103/PhysRevB.73.094423) [arXiv:cond-mat/0505140 \[cond-mat.str-el\].](https://arxiv.org/abs/cond-mat/0505140)
- [67] M. B. Hastings, An area law for one-dimensional quantum systems, J. Stat. Mech. 0708[, P08024 \(2007\),](https://doi.org/10.1088/1742-5468/2007/08/P08024) [arXiv:0705.2024 \[quant-ph\].](https://arxiv.org/abs/0705.2024)
- [68] D. Pérez-García, F. Verstraete, M. M. Wolf, and J. I. Cirac, Matrix product state representations, [Quant. Inf. Comput.](https://doi.org/10.26421/QIC7.5-6-1) 7, 401 (2007), [arXiv:quant-ph/0608197.](https://arxiv.org/abs/quant-ph/0608197)
- [69] S. Ohyama and S. Ryu, Higher structures in matrix product states, Phys. Rev. B 109[, 115152 \(2024\),](https://doi.org/10.1103/PhysRevB.109.115152) [arXiv:2304.05356 \[cond-mat.str-el\].](https://arxiv.org/abs/2304.05356)
- [70] M. Sanz, D. Pérez-García, M. M. Wolf, and J. I. Cirac, A Quantum Version of Wielandt's Inequality, [IEEE Transactions on Information Theory](https://doi.org/10.1109/TIT.2010.2054552) 56, 4668 (2010), [arXiv:0909.5347 \[quant-ph\].](https://arxiv.org/abs/0909.5347)
- [71] J. Garre-Rubio and N. Schuch, Fractional domain wall statistics in spin chains with anomalous symmetries (2024), [arXiv:2405.00439 \[cond-mat.str-el\].](https://arxiv.org/abs/2405.00439)
- [72] A. Kitaev and L. Kong, Models for Gapped Boundaries and Domain Walls, [Communications in Math](https://doi.org/10.1007/s00220-012-1500-5)[ematical Physics](https://doi.org/10.1007/s00220-012-1500-5) 313, 351 (2012), [arXiv:1104.5047 \[cond-mat.str-el\].](https://arxiv.org/abs/1104.5047)
- [73] D. Barter, J. Bridgeman, and R. Wolf, Computing associators of endomorphism fusion categories, SciPost Physics 13, [10.21468/scipostphys.13.2.029](https://doi.org/10.21468/scipostphys.13.2.029) (2022), [arXiv:2110.03644 \[math.QA\].](https://arxiv.org/abs/2110.03644)
- [74] Y. Choi, B. C. Rayhaun, Y. Sanghavi, and S.-H. Shao, Remarks on boundaries, anomalies, and noninvertible symmetries, Phys. Rev. D 108[, 125005 \(2023\),](https://doi.org/10.1103/PhysRevD.108.125005) [arXiv:2305.09713 \[hep-th\].](https://arxiv.org/abs/2305.09713)
- [75] C. Córdova, D. García-Sepúlveda, and N. Holfester, Particle-soliton degeneracies from spontaneously broken non-invertible symmetry, [JHEP](https://doi.org/10.1007/JHEP07(2024)154) 07, 154, [arXiv:2403.08883 \[hep-th\].](https://arxiv.org/abs/2403.08883)
- [76] C. Córdova, N. Holfester, and K. Ohmori, Representation Theory of Solitons (2024), [arXiv:2408.11045](https://arxiv.org/abs/2408.11045) [\[hep-th\].](https://arxiv.org/abs/2408.11045)
- [77] C. Copetti, Defect Charges, Gapped Boundary Conditions, and the Symmetry TFT (2024), [arXiv:2408.01490 \[hep-th\].](https://arxiv.org/abs/2408.01490)
- [78] C. Copetti, L. Córdova, and S. Komatsu, S-Matrix Bootstrap and Non-Invertible Symmetries (2024), [arXiv:2408.13132 \[hep-th\].](https://arxiv.org/abs/2408.13132)
- [79] Y. Zheng, [Symmetry TFTs, boundaries, and non-invertible symmetry resolved Affleck-Ludwig-Cardy](https://www.youtube.com/watch?v=CPoPVG_th-o) [formula](https://www.youtube.com/watch?v=CPoPVG_th-o) (2024), talk at University of Oxford.
- [80] C. G. Callan, Jr. and J. A. Harvey, Anomalies and Fermion Zero Modes on Strings and Domain Walls, [Nucl. Phys. B](https://doi.org/10.1016/0550-3213(85)90489-4) 250, 427 (1985).
- [81] V. G. Turaev and O. Y. Viro, State sum invariants of 3 manifolds and quantum 6j symbols, [Topology](https://doi.org/10.1016/0040-9383(92)90015-A) 31[, 865 \(1992\).](https://doi.org/10.1016/0040-9383(92)90015-A)
- [82] J. W. Barrett and B. W. Westbury, Invariants of piecewise linear three manifolds, [Trans. Am. Math.](https://doi.org/10.1090/S0002-9947-96-01660-1) Soc. 348[, 3997 \(1996\),](https://doi.org/10.1090/S0002-9947-96-01660-1) [arXiv:hep-th/9311155.](https://arxiv.org/abs/hep-th/9311155)
- [83] M. A. Levin and X.-G. Wen, String net condensation: A Physical mechanism for topological phases, Phys. Rev. B 71[, 045110 \(2005\),](https://doi.org/10.1103/PhysRevB.71.045110) [arXiv:cond-mat/0404617.](https://arxiv.org/abs/cond-mat/0404617)
- [84] J. Fuchs, C. Schweigert, and A. Valentino, Bicategories for boundary conditions and for surface defects in 3-d TFT, [Commun. Math. Phys.](https://doi.org/10.1007/s00220-013-1723-0) 321, 543 (2013), [arXiv:1203.4568 \[hep-th\].](https://arxiv.org/abs/1203.4568)
- [85] A. Kapustin and N. Saulina, Topological boundary conditions in abelian Chern-Simons theory, [Nucl.](https://doi.org/10.1016/j.nuclphysb.2010.12.017) Phys. B 845[, 393 \(2011\),](https://doi.org/10.1016/j.nuclphysb.2010.12.017) [arXiv:1008.0654 \[hep-th\].](https://arxiv.org/abs/1008.0654)
- [86] Y. Tachikawa, On gauging finite subgroups, [SciPost Phys.](https://doi.org/10.21468/SciPostPhys.8.1.015) 8, 015 (2020), [arXiv:1712.09542 \[hep-th\].](https://arxiv.org/abs/1712.09542)
- [87] D. Naidu, Categorical Morita equivalence for group-theoretical categories, Communications in Algebra 35, 3544 (2007), [arXiv:math/0605530 \[math.QA\].](https://arxiv.org/abs/math/0605530)
- [88] D. Naidu and D. Nikshych, Lagrangian subcategories and braided tensor equivalences of twisted quantum doubles of finite groups, Communications in mathematical physics 279, 845 (2008), [arXiv:0705.0665 \[math.QA\].](https://arxiv.org/abs/0705.0665)
- [89] B. Uribe Jongbloed, On the classification of pointed fusion categories up to weak Morita equivalence, Pacific Journal of Mathematics 290, 437 (2017), [arXiv:1511.05522 \[math.AT\].](https://arxiv.org/abs/1511.05522)
- [90] P.-S. Hsin, A. Kapustin, and R. Thorngren, Berry Phase in Quantum Field Theory: Diabolical Points and Boundary Phenomena, Phys. Rev. B 102[, 245113 \(2020\),](https://doi.org/10.1103/PhysRevB.102.245113) [arXiv:2004.10758 \[cond-mat.str-el\].](https://arxiv.org/abs/2004.10758)
- [91] D. Aasen, Z. Wang, and M. B. Hastings, Adiabatic paths of Hamiltonians, symmetries of topological order, and automorphism codes, Phys. Rev. B 106[, 085122 \(2022\),](https://doi.org/10.1103/PhysRevB.106.085122) [arXiv:2203.11137 \[quant-ph\].](https://arxiv.org/abs/2203.11137)
- [92] P.-S. Hsin and Z. Wang, On topology of the moduli space of gapped Hamiltonians for topological phases, J. Math. Phys. 64[, 041901 \(2023\),](https://doi.org/10.1063/5.0136906) [arXiv:2211.16535 \[cond-mat.str-el\].](https://arxiv.org/abs/2211.16535)
- [93] P. Etingof, D. Nikshych, V. Ostrik, with an appendix by E. Meir, Fusion categories and homotopy theory, Quantum topology 1, 209 (2010), [arXiv:0909.3140 \[math.QA\].](https://arxiv.org/abs/0909.3140)
- [94] J. C. Baez and A. D. Lauda, Higher-Dimensional Algebra V: 2-Groups, Theory and Applications of Categories 12, 423 (2004), [arXiv:math/0307200 \[math.QA\].](https://arxiv.org/abs/math/0307200)
- [95] A. Kapustin and R. Thorngren, Higher Symmetry and Gapped Phases of Gauge Theories, [Prog. Math.](https://doi.org/10.1007/978-3-319-59939-7_5) 324[, 177 \(2017\),](https://doi.org/10.1007/978-3-319-59939-7_5) [arXiv:1309.4721 \[hep-th\].](https://arxiv.org/abs/1309.4721)
- [96] F. Benini, C. Córdova, and P.-S. Hsin, On 2-Group Global Symmetries and their Anomalies, [JHEP](https://doi.org/10.1007/JHEP03(2019)118) 03[, 118,](https://doi.org/10.1007/JHEP03(2019)118) [arXiv:1803.09336 \[hep-th\].](https://arxiv.org/abs/1803.09336)
- [97] D. Wigner, Algebraic cohomology of topological groups, Bulletin of the American Mathematical Society 76, 825 (1970).
- [98] J. C. Baez and D. Stevenson, The Classifying Space of a Topological 2-Group, in [Algebraic Topology: The Abel Symposium 2007,](https://doi.org/10.1007/978-3-642-01200-6_1) edited by N. Baas, E. M. Friedlander, B. Jahren, and P. A. Østvær (Springer Berlin Heidelberg, Berlin, Heidelberg, 2009) pp. 1–31, [arXiv:0801.3843](https://arxiv.org/abs/0801.3843) [\[math.AT\].](https://arxiv.org/abs/0801.3843)
- [99] C. Wockel, Principal 2-bundles and their gauge 2-groups, Forum Mathematicum 23, [10.1515/form.2011.020](https://doi.org/10.1515/form.2011.020) (2011), [arXiv:0803.3692 \[math.DG\].](https://arxiv.org/abs/0803.3692)
- [100] T. Bartels, [Higher gauge theory I: 2-Bundles](https://arxiv.org/abs/math/0410328) (2006), [arXiv:math/0410328 \[math.CT\].](https://arxiv.org/abs/math/0410328)
- [101] B. Jurčo, Crossed Module Bundle Gerbes; Classification, String Group And Differential Geometry, [International Journal of Geometric Methods in Modern Physics](https://doi.org/10.1142/s0219887811005555) 08, 1079 (2011), [arXiv:math/0510078](https://arxiv.org/abs/math/0510078) [\[math.DG\].](https://arxiv.org/abs/math/0510078)
- [102] N. A. Baas, M. Bökstedt, and T. A. Kro, Two-categorical bundles and their classifying spaces, [Journal](https://doi.org/10.1017/is012001012jkt181) [of K-Theory](https://doi.org/10.1017/is012001012jkt181) 10, 299 (2012), [arXiv:math/0612549 \[math.AT\].](https://arxiv.org/abs/math/0612549)
- [103] D. Berwick-Evans, E. Cliff, L. Murray, A. Nakade, and E. Phillips, [String structures, 2-group bundles,](https://arxiv.org/abs/2110.07571) [and a categorification of the Freed-Quinn line bundle](https://arxiv.org/abs/2110.07571) (2021), [arXiv:2110.07571 \[math.AT\].](https://arxiv.org/abs/2110.07571)
- [104] R. G. Thorngren, Combinatorial Topology and Applications to Quantum Field Theory, [Ph.D. thesis,](https://escholarship.org/uc/item/7r44w49f) UC, Berkeley (main) (2018).
- [105] A. Kapustin, Symmetry Protected Topological Phases, Anomalies, and Cobordisms: Beyond Group Cohomology (2014), [arXiv:1403.1467 \[cond-mat.str-el\].](https://arxiv.org/abs/1403.1467)
- [106] A. Kapustin, R. Thorngren, A. Turzillo, and Z. Wang, Fermionic Symmetry Protected Topological Phases and Cobordisms, [JHEP](https://doi.org/10.1007/JHEP12(2015)052) 12, 052, [arXiv:1406.7329 \[cond-mat.str-el\].](https://arxiv.org/abs/1406.7329)
- [107] D. S. Freed and M. J. Hopkins, Reflection positivity and invertible topological phases, [Geom. Topol.](https://doi.org/10.2140/gt.2021.25.1165) 25[, 1165 \(2021\),](https://doi.org/10.2140/gt.2021.25.1165) [arXiv:1604.06527 \[hep-th\].](https://arxiv.org/abs/1604.06527)
- [108] A. Kitaev, [Differential forms on the space of statistical mechanics models](https://web.ma.utexas.edu/topqft/talkslides/kitaev.pdf) (2019), talk at University of Texas at Austin.
- [109] A. Kapustin and L. Spodyneiko, Higher-dimensional generalizations of Berry curvature, [Phys. Rev. B](https://doi.org/10.1103/PhysRevB.101.235130) 101[, 235130 \(2020\),](https://doi.org/10.1103/PhysRevB.101.235130) [arXiv:2001.03454 \[cond-mat.str-el\].](https://arxiv.org/abs/2001.03454)
- [110] A. Artymowicz, A. Kapustin, and N. Sopenko, Quantization of the Higher Berry Curvature and the Higher Thouless Pump, [Commun. Math. Phys.](https://doi.org/10.1007/s00220-024-05026-2) 405, 191 (2024), [arXiv:2305.06399 \[math-ph\].](https://arxiv.org/abs/2305.06399)
- [111] C. Córdova, D. S. Freed, H. T. Lam, and N. Seiberg, Anomalies in the Space of Coupling Constants and Their Dynamical Applications I, [SciPost Phys.](https://doi.org/10.21468/SciPostPhys.8.1.001) 8, 001 (2020), [arXiv:1905.09315 \[hep-th\].](https://arxiv.org/abs/1905.09315)
- [112] C. Córdova, D. S. Freed, H. T. Lam, and N. Seiberg, Anomalies in the Space of Coupling Constants and Their Dynamical Applications II, [SciPost Phys.](https://doi.org/10.21468/SciPostPhys.8.1.002) 8, 002 (2020), [arXiv:1905.13361 \[hep-th\].](https://arxiv.org/abs/1905.13361)
- [113] Y. Choi and K. Ohmori, Higher Berry phase of fermions and index theorem, [JHEP](https://doi.org/10.1007/JHEP09(2022)022) 09, 022, [arXiv:2205.02188 \[hep-th\].](https://arxiv.org/abs/2205.02188)
- [114] S. Ohyama, Y. Terashima, and K. Shiozaki, Discrete higher Berry phases and matrix product states, Phys. Rev. B 110[, 035114 \(2024\),](https://doi.org/10.1103/PhysRevB.110.035114) [arXiv:2303.04252 \[cond-mat.str-el\].](https://arxiv.org/abs/2303.04252)
- [115] M. Qi, D. T. Stephen, X. Wen, D. Spiegel, M. J. Pflaum, A. Beaudry, and M. Hermele, Charting the space of ground states with tensor networks (2023), [arXiv:2305.07700 \[cond-mat.str-el\].](https://arxiv.org/abs/2305.07700)
- [116] O. E. Sommer, X. Wen, and A. Vishwanath, Higher Berry Curvature from the Wave Function I: Schmidt Decomposition and Matrix Product States (2024), [arXiv:2405.05316 \[cond-mat.str-el\].](https://arxiv.org/abs/2405.05316)
- [117] S. Ohyama and S. Ryu, Higher Berry Connection for Matrix Product States (2024), [arXiv:2405.05327](https://arxiv.org/abs/2405.05327) [\[cond-mat.str-el\].](https://arxiv.org/abs/2405.05327)
- [118] C. L. Douglas and D. J. Reutter, Fusion 2-categories and a state-sum invariant for 4-manifolds (2018), [arXiv:1812.11933 \[math.QA\].](https://arxiv.org/abs/1812.11933)
- [119] J. W. Barrett, C. Meusburger, and G. Schaumann, Gray categories with duals and their diagrams (2024), [arXiv:1211.0529 \[math.QA\].](https://arxiv.org/abs/1211.0529)
- [120] R. Gordon, A. J. Power, and R. Street, [Coherence for tricategories,](https://www.ams.org/books/memo/0558/) Vol. 558 (American Mathematical Soc., 1995).
- [121] C. J. Schommer-Pries, The classification of two-dimensional extended topological field theories (2014), [arXiv:1112.1000 \[math.AT\].](https://arxiv.org/abs/1112.1000)
- [122] D. Gaiotto and T. Johnson-Freyd, Condensations in higher categories (2019), [arXiv:1905.09566](https://arxiv.org/abs/1905.09566) [\[math.CT\].](https://arxiv.org/abs/1905.09566)
- [123] T. Johnson-Freyd, On the Classification of Topological Orders, [Communications in Mathematical](https://doi.org/10.1007/s00220-022-04380-3) Physics 393[, 989 \(2022\),](https://doi.org/10.1007/s00220-022-04380-3) [arXiv:2003.06663 \[math.CT\].](https://arxiv.org/abs/2003.06663)
- [124] L. Kong and H. Zheng, Categories of quantum liquids I, [JHEP](https://doi.org/10.1007/JHEP08(2022)070) 08, 070, [arXiv:2011.02859 \[hep-th\].](https://arxiv.org/abs/2011.02859)
- [125] K. Inamura and K. Ohmori, Fusion Surface Models: 2+1d Lattice Models from Fusion 2-Categories, [SciPost Phys.](https://doi.org/10.21468/SciPostPhys.16.6.143) 16, 143 (2024), [arXiv:2305.05774 \[cond-mat.str-el\].](https://arxiv.org/abs/2305.05774)
- [126] T. D. Décoppet, The Morita Theory of Fusion 2-Categories, [Higher Structures](https://doi.org/10.21136/hs.2023.07) 7, 234 (2023), [arXiv:2208.08722 \[math.CT\].](https://arxiv.org/abs/2208.08722)
- [127] D. S. Freed and C. Teleman, Gapped Boundary Theories in Three Dimensions, [Commun. Math. Phys.](https://doi.org/10.1007/s00220-021-04192-x) 388[, 845 \(2021\),](https://doi.org/10.1007/s00220-021-04192-x) [arXiv:2006.10200 \[math.QA\].](https://arxiv.org/abs/2006.10200)
- [128] T. D. Décoppet and H. Xu, Local modules in braided monoidal 2-categories, [J. Math. Phys.](https://doi.org/10.1063/5.0172042) 65, 061702 [\(2024\),](https://doi.org/10.1063/5.0172042) [arXiv:2307.02843 \[math.CT\].](https://arxiv.org/abs/2307.02843)
- [129] S. Ohyama and S. Ryu, Higher Berry Phase from Projected Entangled Pair States in (2+1) dimensions (2024), [arXiv:2405.05325 \[cond-mat.str-el\].](https://arxiv.org/abs/2405.05325)
- [130] O. E. Sommer, A. Vishwanath, and X. Wen, Higher Berry Curvature from the Wave function II: Locally Parameterized States Beyond One Dimension (2024), [arXiv:2405.05323 \[cond-mat.str-el\].](https://arxiv.org/abs/2405.05323)
- [131] L. Bhardwaj, S. Schäfer-Nameki, and J. Wu, Universal Non-Invertible Symmetries, [Fortsch. Phys.](https://doi.org/10.1002/prop.202200143) 70, [2200143 \(2022\),](https://doi.org/10.1002/prop.202200143) [arXiv:2208.05973 \[hep-th\].](https://arxiv.org/abs/2208.05973)
- [132] T. Bartsch, M. Bullimore, A. E. V. Ferrari, and J. Pearson, Non-invertible Symmetries and Higher Representation Theory I (2022), [arXiv:2208.05993 \[hep-th\].](https://arxiv.org/abs/2208.05993)
- [133] C. Delcamp and A. Tiwari, Higher categorical symmetries and gauging in two-dimensional spin systems, [SciPost Phys.](https://doi.org/10.21468/SciPostPhys.16.4.110) 16, 110 (2024), [arXiv:2301.01259 \[hep-th\].](https://arxiv.org/abs/2301.01259)
- [134] M. Barkeshli, Y.-A. Chen, S.-J. Huang, R. Kobayashi, N. Tantivasadakarn, and G. Zhu, Codimension-2 defects and higher symmetries in (3+1)D topological phases, [SciPost Phys.](https://doi.org/10.21468/SciPostPhys.14.4.065) 14, 065 (2023), [arXiv:2208.07367 \[cond-mat.str-el\].](https://arxiv.org/abs/2208.07367)
- [135] M. Barkeshli, Y.-A. Chen, P.-S. Hsin, and R. Kobayashi, Higher-group symmetry in finite gauge theory and stabilizer codes, [SciPost Phys.](https://doi.org/10.21468/SciPostPhys.16.4.089) 16, 089 (2024), [arXiv:2211.11764 \[cond-mat.str-el\].](https://arxiv.org/abs/2211.11764)
- [136] D. Tambara and S. Yamagami, Tensor Categories with Fusion Rules of Self-Duality for Finite Abelian Groups, [Journal of Algebra](https://doi.org/https://doi.org/10.1006/jabr.1998.7558) 209, 692 (1998).
- [137] G. I. Kac and V. G. Paljutkin, Finite ring groups, [Trans. Mosc. Math. Soc. 15 251-294;](http://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=mmo&paperid=170&option_lang=eng) Translation from Tr. Mosk. Mat. Obs. 15 224-261. (1966).
- [138] D. Tambara, Representations of tensor categories with fusion rules of self-duality for abelian groups, [Israel Journal of Mathematics](https://doi.org/10.1007/BF02803515) 118, 29 (2000).
- [139] J. Fuchs, I. Runkel, and C. Schweigert, TFT construction of RCFT correlators 1. Partition functions, [Nucl. Phys. B](https://doi.org/10.1016/S0550-3213(02)00744-7) 646, 353 (2002), [arXiv:hep-th/0204148.](https://arxiv.org/abs/hep-th/0204148)
- [140] J. Fuchs and C. Stigner, On Frobenius algebras in rigid monoidal categories, Arabian Journal for Science and Engineering **33-2C**, 175 (2009), [arXiv:0901.4886 \[math.CT\].](https://arxiv.org/abs/0901.4886)
- [141] N. Andruskiewitsch and J. M. Mombelli, On module categories over finite-dimensional Hopf algebras, [Journal of Algebra](https://doi.org/https://doi.org/10.1016/j.jalgebra.2007.04.006) 314, 383 (2007), [arXiv:math/0608781 \[math.QA\].](https://arxiv.org/abs/math/0608781)