

# Weak Distributive Laws between Monads of Continuous Valuations and of Non-Deterministic Choice

Jean Goubault-Larrecq

*Université Paris-Saclay, CNRS, ENS Paris-Saclay, Laboratoire Méthodes Formelles,  
91190, Gif-sur-Yvette, France.  
jgl@lml.cnrs.fr*

---

## Abstract

We show that there is weak distributive law of the Smyth hyperspace monad  $\mathcal{Q}_V$  (resp., the Hoare hyperspace monad  $\mathcal{H}_V$ , resp. the monad  $\mathcal{P}\ell_V^q$  of quasi-lenses, resp. the monad  $\mathcal{P}\ell_V$  of lenses) over the continuous valuation monad  $\mathbf{V}$ , as well as over the subprobability valuation monad  $\mathbf{V}_{\leq 1}$  and the probability valuation monad  $\mathbf{V}_1$ , on the whole category  $\mathbf{Top}$  of topological spaces (resp., on certain full subcategories such as the category of locally compact spaces or of stably compact spaces). We show that the resulting weak composite monad is the author's monad of superlinear previsions (resp., sublinear previsions, resp. forks), possibly subnormalized or normalized depending on whether we consider  $\mathbf{V}_{\leq 1}$  or  $\mathbf{V}_1$  instead of  $\mathbf{V}$ . As a special case, we obtain a weak distributive law of the monad  $\mathcal{P}\ell_V^q \cong \mathcal{P}\ell_V$  over the monad of (sub)probability Radon measures  $\mathbf{R}_\bullet$  on the category of stably compact spaces, which specializes further to a weak distributive laws of the Vietoris monad over  $\mathbf{R}_\bullet$ . The associated weak composite monad is the monad of (sub)normalized forks.

*Keywords:* weak distributive law, monad, continuous valuation, Radon measure, hyperspace, Smyth hyperspace, Hoare powerspace, Plotkin hyperspace, quasi-lens, lens

*2020 MSC:* 18C15, 54B20, 28A33, 46E27



For the purpose of Open Access, a CC-BY public copyright licence has been applied by the authors to the present document and will be applied to all subsequent versions up to the Author Accepted Manuscript arising from this submission.

Une montagne en mal d'enfant  
 Jetait une clameur si haute  
 Que chacun, au bruit accourant,  
 Crut qu'elle accoucherait, sans faute,  
 D'une cité plus grosse que Paris;  
 Elle accoucha d'une souris.  
 Quand je songe à cette fable,  
 Dont le récit est menteur  
 Et le sens est véritable,  
 Je me figure un auteur  
 Qui dit: Je chanterai la guerre  
 Que firent les Titans au Maître du tonnerre. »  
 C'est promettre beaucoup: mais qu'en sort-il souvent?  
 Du vent.

*La montagne qui accouche*, Jean de la Fontaine, Fables, livre V, fable 10, 1668.

## 1. Introduction

Jon Beck introduced the notion of distributive laws  $\lambda: TS \rightarrow ST$  of a monad  $S$  over a monad  $T$ , and showed that any such distributive law allowed for the construction of a composite monad  $ST$  [4]. The theory is much richer than what the latter may suggest, and there is a one-to-one correspondence between distributive laws as above, extensions of  $T$  to the Kleisli category of  $S$ , and liftings of  $S$  to the category of  $T$ -algebras.

It was once dreamed that one could perhaps use this construction in order to define new monads, combining various side-effects in the semantics of programming languages. But distributive laws are rare, and, as a case in point, there is no distributive law of the monad of non-deterministic choice over the monad of discrete probability measure on **Set** [40]. The same argument shows that there is no distributive law of any of the monads of non-deterministic choice (the Smyth powerdomain monad  $\mathcal{Q}$ , the Hoare powerdomain monad  $\mathcal{H}$ , the Plotkin powerdomain monad  $\mathcal{P}\ell$ ) over the monad  $\mathbf{V}$  of continuous valuations (or  $\mathbf{V}_1$  [probability valuations], or  $\mathbf{V}_{\leq 1}$  [subprobability valuations], implementing probabilistic choice).

Richard Garner later observed that one could still build a *weak* composite monad of  $S$  and  $T$ —which is not simply the composition  $ST$ —from a *weak* distributive law of  $S$  over  $T$  [9], a notion that he also invented. While distributive laws are natural transformations that must satisfy four equations expressing compatibility with the units and multiplications of each of the two monads, weak distributive laws are only required to satisfy three: compatibility with the unit of the  $T$  monad is ignored. The theory parallels Beck's theory of distributive laws, and notably, weak distributive laws of  $S$  over  $T$

are in one-to-one correspondence with weak extensions of  $T$  to the Kleisli category of  $S$ , and weak liftings of  $S$  to the category of  $T$ -algebras.

It was suggested by Alexandre Goy [20, Chapter 7] that, while there is no distributive law of the Vietoris monad (which sends every space  $X$  to its space of non-empty compact subsets with the Vietoris topology) over the monad  $\mathbf{R}_1$  of Radon probability measures on the category  $\mathbf{KHaus}$  of compact Hausdorff spaces, there should be a weak distributive law relating the two. We will confirm that it exists, as a special case of a weak distributive law between the Plotkin hyperspace monad  $\mathcal{P}l_V$  over  $\mathbf{V}_1 \cong \mathbf{R}_1$ , in the final section of this paper (Section 12). (A toy version of this result was established by Goy and Petrişan [21], on the category  $\mathbf{Set}$ , with the powerset monad instead of the Vietoris monad and the finite distribution monad instead of  $\mathbf{R}_1$ .)

We will show that the resulting weak composite monad is the author’s monad of normalized forks, introduced thirteen years earlier [11] on the category of  $\mathit{dcpo}$ . We take the opportunity to make the relatively obvious generalization to topological spaces of that monad here.

Before we reach that final section on the Vietoris and Radon monads, we will explore essentially all possible combinations of monads of non-deterministic choice on  $\mathbf{Top}$  on the one hand, and of monads of probabilistic choice on the other hand. The latter include the monads  $\mathbf{V}$  of continuous valuations,  $\mathbf{V}_{\leq 1}$  of subprobability valuations,  $\mathbf{V}_1$  of probability valuations. Continuous valuations are a close cousin to measures, and we will soon explain how they relate to each other. The former include the Smyth hyperspace monad  $\mathcal{Q}_V$  (demonic non-determinism) the Hoare hyperspace monad  $\mathcal{H}_V$  (angelic non-determinism), and two variants of the Plotkin hyperspace monad (erratic non-determinism), the monad  $\mathcal{P}l_V$  of lenses and the monad  $\mathcal{P}l_V^q$  of quasi-lenses. We show that all of those have weak distributivity laws over  $\mathbf{V}_\bullet$ , where  $\bullet$  is nothing, “ $\leq 1$ ”, or “ $1$ ”. In the case of  $\mathcal{Q}_V$ , this works on the whole category  $\mathbf{Top}$  of topological spaces. For  $\mathcal{H}_V$ , we have to restrict to certain full subcategories, such as the subcategory of locally compact spaces. For  $\mathcal{P}l_V^q$  and  $\mathcal{P}l_V$ , we need to restrict even further, typically to stably compact spaces—a category that still contains  $\mathbf{KHaus}$  as a proper subcategory.

In each case, we make the resulting weak composite monad explicit. And, in each case, that is a well-known monad: the monad of superlinear (resp., subnormalized or normalized, depending on  $\bullet$ ) previsions in the demonic case, the monad of sublinear (resp., subnormalized or normalized) previsions in the angelic case, and the monad of (resp., subnormalized or normalized) forks in the erratic case. Those monads first appeared in [11] on the category

of *depos*, and we will rely on the (pretty obvious) generalization to **Top** here. “Tout ça pour ça!”, as we say in French; in a more elegant form—and written in La Fontaine’s admirable style—see the quote before this introduction. At least this study should confirm that Garner definitely hit upon a, perhaps the, right way of combining monads.

*Outline.* Section 2 sets some technical preliminaries. We will introduce other notions and results along the way. The rest of the paper is structured in three parts, for each version of non-determinism (demonic, angelic, erratic) we consider. Each part has exactly the same structure, except for the last one, which has an extra, final Section 12 on Radon measures and the Vietoris monad. Explicitly, that structure consists of one section on the monad we will eventually discover as the weak composite monad (superlinear previsions in Section 3, sublinear previsions in Section 6, forks in Section 9), one on the weak distributive law (Section 4, Section 7, Section 10), and one on the resulting weak composite monad (Section 5, Section 8, Section 11).

## 2. Preliminaries

For background on topology, we refer the reader to [13]. We write  $\text{int}(A)$  for the interior of  $A$ ,  $cl(A)$  (or  $cl_X(A)$ ) for the closure of  $A$  (in a space  $X$ ), and  $\mathcal{O}X$  for the lattice of open subsets of  $X$ . The specialization preordering  $\leq$  of a topological space  $X$  is defined on points  $x, y \in X$  by  $x \leq y$  if and only if every open neighborhood of  $x$  contains  $y$ , if and only if  $x$  lies in the closure of  $\{y\}$ . We also write  $x \in A \mapsto f(x)$  for the function that maps every element  $x$  of a set  $A$  to the value  $f(x)$ ; logicians and computer scientists would use the notation  $\lambda x \in A. f(x)$ , but this would interfere with the standard use of the letter  $\lambda$  for (weak) distributive laws.

A *directed* family of elements in a poset is a non-empty family of elements  $D$  such that any two elements of  $D$  have an upper bound in  $D$ . A *filtered* family is defined similarly, with lower bounds replacing upper bounds. We use  $\sup^\uparrow$  for suprema of directed families,  $\bigcup^\uparrow$  for unions of directed families of sets,  $\inf^\downarrow$  for filtered infima,  $\bigcap^\downarrow$  for intersections of filtered families of sets.

An *irreducible* closed subset  $C$  of  $X$  is a non-empty closed subset such that, for any two closed subsets  $C_1$  and  $C_2$  of  $X$  such that  $C \subseteq C_1 \cup C_2$ ,  $C$  is included in  $C_1$  or in  $C_2$  already; equivalently, if  $C$  intersects two open sets, it must intersect their intersection. A space  $X$  is *sober* if and only if it is  $T_0$  and every irreducible closed subset is of the form  $\downarrow x$  for some point  $x \in X$ . Every Hausdorff space, for example, is sober. The notation  $\downarrow x$

stands for the *downward closure* of  $x$  in  $X$ , namely the set of points  $y$  below  $x$ . Symmetrically,  $\uparrow x$  stands for the *upward closure* of  $x$ , namely the set of points  $y$  above  $x$ . This notation extends to  $\uparrow A$ , for any subset  $A$ , denoting  $\bigcup_{x \in A} \uparrow x$ .

A *compact* subset  $A$  of a space  $X$  is one such that one can extract a finite subcover from any of its open covers. No separation property is assumed. A subset  $A$  of  $X$  *saturated* if and only if it is equal to the intersection of its open neighborhoods, or equivalently if and only if it is upwards-closed in the specialization preordering of  $X$ .

A space  $X$  is *locally compact* if and only if every point has a base of compact neighborhoods, or equivalently of compact saturated neighborhoods, since for any compact subset  $K$  of  $X$ , the upward closure  $\uparrow K$  of  $K$  with respect to the specialization preordering of  $X$  is compact saturated. Please beware that, in non-Hausdorff spaces, a compact space may fail to be locally compact.

A space is *coherent* if and only if the intersection of any two compact saturated subsets is compact (and necessarily saturated). That, too, is a property that may fail in non-Hausdorff spaces.

A *stably locally compact* space is a coherent, locally compact, sober space, and a *stably compact* is a compact, stably locally compact space. We note that every compact Hausdorff space is stably compact, as well as any coherent continuous dcpo with a least element. (We will not define dcpos; see [13], [1], or [10].)

Let  $\overline{\mathbb{R}}_+$  be the set of extended non-negative real numbers  $\mathbb{R}_+ \cup \{\infty\}$ , with its usual ordering. When needed, we will consider it with its Scott topology, whose open sets are the intervals  $]t, \infty]$ ,  $t \in \mathbb{R}_+$ , plus  $\emptyset$  and  $\overline{\mathbb{R}}_+$  itself. In general, the *Scott topology* on a poset has as open sets the upward closed subsets  $U$  (i.e.,  $U = \uparrow U$ ) such that every directed family  $D$  whose supremum exists and is in  $U$  intersects  $U$ .

A *continuous valuation* on a space  $X$  is a map  $\nu: \mathcal{O}X \rightarrow \overline{\mathbb{R}}_+$  that is *strict* ( $\nu(\emptyset) = 0$ ), *modular* (for all  $U, V \in \mathcal{O}X$ ,  $\nu(U) + \nu(V) = \nu(U \cup V) + \nu(U \cap V)$ ) and Scott-continuous. We say that  $\nu$  is a *probability valuation* if and only if  $\nu(X) = 1$ , and a *subprobability valuation* if and only if  $\nu(X) \leq 1$ .

Let  $\mathbf{V}X$  denote the space of continuous valuations on a space  $X$ , with the following *weak topology*. That is defined by subbasic open sets  $[U > r] \stackrel{\text{def}}{=} \{\nu \in \mathbf{V}X \mid \nu(U) > r\}$ , where  $U \in \mathcal{O}X$  and  $r \in \mathbb{R}_+$ . We define its subspace  $\mathbf{V}_1 X$  of probability valuations and  $\mathbf{V}_{\leq 1} X$  (subprobability) similarly.

In general, we will write  $\mathbf{V}_\bullet X$ , where  $\bullet$  is nothing, “ $\leq 1$ ”, or “1”.

The specialization ordering of each is the *stochastic ordering*  $\leq$  given by  $\nu \leq \nu'$  if and only if  $\nu(U) \leq \nu'(U)$  for every  $U \in \mathcal{O}X$ ; indeed,  $\nu \leq \nu'$  if and only if for every  $U \in \mathcal{O}X$ , for every  $r \in \mathbb{R}_+$ ,  $\nu \in [U > r]$  implies  $\nu' \in [U > r]$ .

Continuous valuations are an alternative to measures that have become popular in domain theory [26, 25]. The first results that connected continuous valuations and measures are due to Saheb-Djahromi [36] and Lawson [32]. The current state of the art on similar results is the following. In one direction, every measure on the Borel  $\sigma$ -algebra of  $X$  induces a continuous valuation on  $X$  by restriction to the open sets, if  $X$  is hereditarily Lindelöf (namely, if every directed family of open sets contains a cofinal monotone sequence). This is an easy observation, and one half of Adamski’s theorem [2, Theorem 3.1], which states that a space is hereditary Lindelöf if and only if every measure on its Borel  $\sigma$ -algebra restricts to a continuous valuation on its open sets. In the other direction, every continuous valuation on a space  $X$  extends to a measure on the Borel sets provided that  $X$  is an LCS-complete space [7, Theorem 1]. An *LCS-complete* space is a space homeomorphic to a  $G_\delta$  subset of a locally compact sober space— $G_\delta$  meaning a countable intersection of open sets.

There is a monad  $(\mathbf{V}_\bullet, \eta, \mu)$  on **Top**. For every continuous map  $f: X \rightarrow Y$ ,  $\mathbf{V}_\bullet f$  sends every  $\nu \in \mathbf{V}_\bullet X$  to the *pushforward* continuous valuation  $f[\nu]$ , defined by  $f[\nu](V) \stackrel{\text{def}}{=} \nu(f^{-1}(V))$  for every  $V \in \mathcal{O}Y$ . The unit  $\eta$  is defined by  $\eta_X(x) \stackrel{\text{def}}{=} \delta_x$  for every  $x \in X$ , for every space  $X$ ;  $\delta_x$  is the *Dirac valuation* at  $x$ , and maps every open subset  $U$  of  $X$  to 1 if  $x \in U$ , to 0 otherwise. For every continuous map  $f: X \rightarrow \mathbf{V}_\bullet Y$ , its *extension*  $f^\dagger: \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet Y$  is defined by  $f^\dagger(\nu)(V) \stackrel{\text{def}}{=} \int_{x \in X} f(x)(V) d\nu$  for every  $\nu \in \mathbf{V}_\bullet X$  and for every  $V \in \mathcal{O}Y$ , and then  $\mu_X \stackrel{\text{def}}{=} \text{id}_{\mathbf{V}_\bullet X}^\dagger$ . Integration is best defined by a Choquet formula [39, Section 4]. Explicitly, let  $\mathcal{L}X$  be the set of all lower semicontinuous functions  $h$  from  $X$  to  $\overline{\mathbb{R}}_+$ , that is, the set of all continuous maps from  $X$  to  $\overline{\mathbb{R}}_+$  equipped with its Scott topology. For every  $h \in \mathcal{L}X$ ,  $\int_{x \in X} h(x) d\nu$  is defined as the indefinite Riemann integral  $\int_0^\infty \nu(h^{-1}([t, \infty])) dt$ . The following *change of variable* formula follows immediately: for every continuous map  $f: X \rightarrow Y$ , for every  $h \in \mathcal{L}Y$ , for every  $\nu \in \mathbf{V}_\bullet X$ ,  $\int_{y \in Y} h(y) df[\nu] = \int_{x \in X} h(f(x)) d\nu$ . We also note that, for every  $h \in \mathcal{L}X$ , for every  $\xi \in \mathbf{V}_\bullet \mathbf{V}_\bullet X$ ,

$$\int_{x \in X} h(x) d\mu_X(\xi) = \int_{\nu \in \mathbf{V}_\bullet X} \left( \int_{x \in X} h(x) d\nu \right) d\xi$$

See [31, Hilfssatz 6.1] for a proof.

The Choquet formula makes sense when  $\nu$  is not just a continuous valuation, but any monotone map from  $\mathcal{O}X$  to  $\overline{\mathbb{R}}_+$ , and we will use that observation later on.

It will be useful to note that  $\eta_X^{-1}([U > r]) = U$  if  $r < 1$  and is empty otherwise, that  $(\mathbf{V}\bullet f)^{-1}([V > r]) = [f^{-1}(V) > r]$ . The formula for  $(\mu_X)^{-1}([U > r])$  is more involved. For every  $h \in \mathcal{L}X$ , for every  $r \in \mathbb{R}_+$ , let  $[h > r] \stackrel{\text{def}}{=} \{\nu \in \mathbf{V}\bullet X \mid \int_{x \in X} h(x) d\nu > r\}$ . This is also an open subset of  $\mathbf{V}\bullet X$ , and such sets form an alternate base of the weak topology, when  $h$  varies over  $\mathcal{L}X$  and  $r$  over  $\mathbb{R}_+$ , see [27, Theorem 3.3] where this was proved for spaces of probability and subprobability valuations; the proof is similar for arbitrary continuous valuations. We may even restrict  $r$  to be non-zero, since  $[h > 0] = \bigcup_{r>0} [h > r]$ , and in fact to be equal to 1 since  $[h > r] = [h/r > 1]$  when  $r > 0$ . The point is that  $(\mu_X)^{-1}([U > r]) = [(\nu \in \mathbf{V}\bullet X \mapsto \nu(U)) > r]$ , and more generally:

$$(\mu_X)^{-1}([h > r]) = \left[ \left( \nu \in \mathbf{V}\bullet X \mapsto \int_{x \in X} h(x) d\nu \right) > r \right]. \quad (1)$$

We will also need to note that  $(\mathbf{V}f)^{-1}([h > r]) = [h \circ f > r]$ , a consequence of the change of variable formula, and that  $\eta_X^{-1}([h > r]) = h^{-1}([r, \infty])$ .

## Part I

# A weak distributive law between $\mathbf{V}\bullet$ and $\mathcal{Q}_V$

For every topological space  $X$ , let  $\mathcal{Q}X$  be the set of non-empty compact saturated subsets of  $X$ . The *upper Vietoris* topology on that set has basic open subsets  $\square U$  consisting of those non-empty compact saturated subsets of  $X$  that are included in  $U$ , where  $U$  ranges over the open subsets of  $X$ . We write  $\mathcal{Q}_V X$  for the resulting topological space. Its specialization ordering is reverse inclusion  $\supseteq$ .

The  $\mathcal{Q}_V$  constructions was studied by a number of people, starting with Mike Smyth [38], and later by Andrea Schalk [37, Section 7] who studied not only this, but also the variant with the Scott topology, and a localic

counterpart. See also [1, Sections 6.2.2, 6.2.3] or [10, Section IV-8], where the accent is rather on the Scott topology of  $\sqsupseteq$ . In this setting,  $\mathcal{Q}_V$  is a model of *demonic* non-determinism.

There is a monad  $(\mathcal{Q}_V, \eta^\mathcal{Q}, \mu^\mathcal{Q})$  on **Top**. For every continuous map  $f: X \rightarrow Y$ ,  $\mathcal{Q}_V f$  maps every  $Q \in \mathcal{Q}_V X$  to  $\uparrow f[Q]$ , where  $f[Q]$  denotes  $\{f(x) \mid x \in Q\}$ . The unit  $\eta^\mathcal{Q}$  is defined by  $\eta_X^\mathcal{Q}(x) \stackrel{\text{def}}{=} \uparrow x$  for every  $x \in X$ . For every continuous map  $f: X \rightarrow \mathcal{Q}_V Y$ , there is an extension  $f^\sharp: \mathcal{Q}_V X \rightarrow \mathcal{Q}_V Y$ , defined by  $f^\sharp(Q) \stackrel{\text{def}}{=} \bigcup_{x \in Q} f(x)$ . The multiplication  $\mu^\mathcal{Q}$  is defined by  $\mu_X^\mathcal{Q} \stackrel{\text{def}}{=} \text{id}_{\mathcal{Q}_V X}^\sharp$ , namely  $\mu_X^\mathcal{Q}(\mathcal{Q}) \stackrel{\text{def}}{=} \bigcup \mathcal{Q}$  [37, Proposition 7.21].

We note the equalities  $\eta^{\mathcal{Q}^{-1}}(\square U) = U$ ,  $(\mathcal{Q}_V f)^{-1}(\square V) = \square f^{-1}(V)$ , and  $\mu^{\mathcal{Q}^{-1}}(\square U) = \square \square U$ .

### 3. The monad of superlinear previsions

We will show that there is a weak distributive law of  $\mathcal{Q}_V$  of  $\mathbf{V}_\bullet$ , and we will build the resulting weak composite monad. That monad has been well-known for some time, and is the monad of superlinear previsions of [11], except that this paper studied it on the category of dpos and Scott-continuous maps instead of **Top**; the study of superlinear previsions on **Top** was initiated in [14], but it was not stated that they formed a monad there, although, as we will see below, this is pretty elementary.

At any rate, we will need to learn quite a number of facts about superlinear previsions in our endeavor to build the desired weak distributive law and its resulting weak composite monad.

We recall that  $\mathcal{L}X$  is the space of lower semicontinuous maps from  $X$  to  $\overline{\mathbb{R}}_+$ . We equip it with the Scott topology of the pointwise ordering. A *prevision* on  $X$  is a Scott-continuous functional  $F: \mathcal{L}X \rightarrow \overline{\mathbb{R}}_+$  such that  $F(ah) = aF(h)$  for all  $a \in \mathbb{R}_+$  and  $h \in \mathcal{L}X$ . It is *linear* (resp., *superlinear*) if and only if  $F(h + h') = F(h) + F(h')$  (resp.,  $\geq$ ) for all  $h, h' \in \mathcal{L}X$ . It is *subnormalized* (resp., *normalized*) iff  $F(\mathbf{1} + h) \leq \mathbf{1} + F(h)$  (resp.,  $=$ ) for every  $h \in \mathcal{L}X$ , where  $\mathbf{1}$  is the constant function with value 1. We write  $\mathbb{P}_{\text{dp}}X$  for the space of superlinear previsions on  $X$ ,  $\mathbb{P}_{\text{dp}}^{\leq 1}X$  for its subspace of subnormalized superlinear previsions, and  $\mathbb{P}_{\text{dp}}^1X$  for its subspace of normalized superlinear previsions; synthetically, we use the notation  $\mathbb{P}_{\text{dp}}^\bullet X$ . Similarly, we denote by  $\mathbb{P}_p^\bullet X$  the corresponding spaces of linear previsions. The topology on each of those spaces is generated by subbasic open sets  $[h > r] \stackrel{\text{def}}{=} \{F \in \mathbb{P}_{\text{dp}}^\bullet X \text{ (resp., } \mathbb{P}_p^\bullet X) \mid F(h) > r\}$ ,  $h \in \mathcal{L}X$ ,  $r \in \mathbb{R}_+$ .



Its specialization ordering is the pointwise ordering:  $F \leq F'$  if and only if  $F(h) \leq F'(h)$  for every  $h \in \mathcal{L}X$ .

There is a  $\mathbb{P}_{\text{DP}}^\bullet$  endofunctor on **Top**, and its action on morphisms  $f: X \rightarrow Y$  is given by  $\mathbb{P}_{\text{DP}}^\bullet f(F) = (h \in \mathcal{L}Y \mapsto F(h \circ f))$ . The  $\mathbb{P}_{\text{P}}^\bullet$  endofunctor is defined similarly. Those are both functor parts of monads. Those monads, as well as the forthcoming monad of sublinear previsions, were made explicit in [11, Proposition 2], on the category of dpos (resp., pointed dpos) and Scott-continuous maps, except that spaces of previsions were given the Scott topology of the pointwise ordering. The formulae for unit and multiplication is the same below. The proof that those form monads is elementary, and the only change compared to [11] is how we show continuity.

**Proposition 3.1.** *Let  $\bullet$  be nothing, “ $\leq 1$ ”, or “1”. There is a monad  $(\mathbb{P}_{\text{DP}}^\bullet, \eta^{\text{DP}}, \mu^{\text{DP}})$  (resp.,  $(\mathbb{P}_{\text{P}}^\bullet, \eta^{\text{P}}, \mu^{\text{P}})$ ) on **Top**, whose unit and multiplication are defined at every topological space  $X$  by:*

- for every  $x \in X$ ,  $\eta_X^{\text{DP}}(x) \stackrel{\text{def}}{=} (h \in \mathcal{L}X \mapsto h(x))$ ,
- for every  $\mathcal{F} \in \mathbb{P}_{\text{DP}}^\bullet \mathbb{P}_{\text{DP}}^\bullet X$  (resp., in  $\mathbb{P}_{\text{P}}^\bullet \mathbb{P}_{\text{P}}^\bullet X$ ),  $\mu_X^{\text{DP}}(\mathcal{F}) \stackrel{\text{def}}{=} (h \in \mathcal{L}X \mapsto \mathcal{F}(F \mapsto F(h)))$ , where  $F$  ranges over  $\mathbb{P}_{\text{DP}}^\bullet X$ , resp.  $\mathbb{P}_{\text{P}}^\bullet X$ .

PROOF. Unit. The functional  $(h \in \mathcal{L}X \mapsto h(x))$  is a linear prevision, hence also a superlinear prevision, as one checks easily. It is also normalized, hence subnormalized. The map  $\eta_X^{\text{DP}}$  is continuous, since the inverse image of  $[h > r]$  is  $h^{-1}(]r, \infty])$ .

Multiplication. For every  $h \in \mathcal{L}X$ , the map  $F \mapsto F(h)$  is lower semicontinuous, since the inverse image of  $]r, \infty]$  is  $[h > r]$ . Hence  $\mathcal{F}(F \mapsto F(h))$  makes sense. The map  $(h \in \mathcal{L}X \mapsto \mathcal{F}(F \mapsto F(h)))$  is Scott-continuous, because  $\mathcal{F}$  is. Additionally, when  $\mathcal{F}$  is superlinear (resp., linear), and  $F$  ranges over  $\mathbb{P}_{\text{DP}}^\bullet X$ , resp.  $\mathbb{P}_{\text{P}}^\bullet X$ , it is clear that  $(h \in \mathcal{L}X \mapsto \mathcal{F}(F \mapsto F(h)))$  is also superlinear (resp., linear). Hence  $\mu_X^{\text{DP}}$  is a superlinear (resp., linear) prevision. When  $\bullet$  is “ $\leq 1$ ”, namely if  $\mathcal{F}$  is subnormalized and  $F$  ranges over  $\mathbb{P}_{\text{DP}}^{\leq 1} X$ , then for every  $h \in \mathcal{L}X$ ,  $\mu_X^{\text{DP}}(\mathbf{1} + h) = \mathcal{F}(F \mapsto F(\mathbf{1} + h)) \leq \mathcal{F}(F \mapsto \mathbf{1} + F(h)) \leq \mathbf{1} + \mathcal{F}(F \mapsto F(h)) = \mathbf{1} + \mu_X^{\text{DP}}(h)$ , so  $\mu_X^{\text{DP}}(\mathcal{F})$  is subnormalized; similarly, if  $\mathcal{F}$  is normalized, then  $\mu_X^{\text{DP}}(\mathcal{F})$  is normalized. (And yes, the dependency on  $\bullet$  should be made explicit in the notation  $\mu_X^{\text{DP}}$ , but we feel that the notation is heavy enough.) Finally,  $\mu_X^{\text{DP}}$  is continuous since the inverse image of  $[h > r]$  is  $[(F \mapsto F(h)) > r]$ .

The monad laws are easily verified. In fact, they are verified exactly as for the continuation monad, which is defined with the same formulae.  $\square$

There is a homeomorphism between  $\mathbb{P}_p^\bullet X$  and  $\mathbf{V}_\bullet X$ , for every space  $X$ : in one direction,  $\nu \in \mathbf{V}_\bullet X$  is mapped to  $(h \in \mathcal{L}X \mapsto \int_{x \in X} h(x) d\nu)$ , and in the other direction any  $G \in \mathbb{P}_p^\bullet X$  is mapped to  $(U \in \mathcal{O}X \mapsto G(\chi_U))$ , where  $\chi_U$  is the characteristic map of  $U$ . We note in particular that  $\int_{x \in X} \chi_U(x) d\nu = \nu(U)$ . This was first proved by Kirch [31, Satz 8.1], under the additional assumption that  $X$  is core-compact; Tix later observed that this assumption was unnecessary [39, Satz 4.16].

#### 4. The weak distributive law

We will show that there is a weak distributive law of  $\mathcal{Q}_V$  over  $\mathbf{V}_\bullet$  on the whole category of topological spaces. This will be the collection of maps  $\lambda_X^\sharp$  given as follows.

**Proposition 4.1 (Corollary 12.6 of [19]).** *For every topological space  $X$ , there is a continuous map  $\lambda_X^\sharp$  from  $\mathbf{V}_\bullet \mathcal{Q}_V X$  to  $\mathcal{Q}_V \mathbf{V}_\bullet X$ , which maps every  $\mu \in \mathbf{V}_\bullet \mathcal{Q}_V X$  to the collection of continuous valuations  $\nu \in \mathbf{V}_\bullet X$  such that  $\nu(U) \geq \mu(\square U)$  for every  $U \in \mathcal{O}X$ .*

We will also need to know how this is proved.

There is a map  $r_{\text{DP}}: \mathcal{Q}_V(\mathbf{V}_\bullet X) \rightarrow \mathbb{P}_{\text{DP}}^\bullet X$ , defined by  $r_{\text{DP}}(Q)(h) \stackrel{\text{def}}{=} \min_{\nu \in Q} \int_{x \in X} h(x) d\nu$ , and a map  $s_{\text{DP}}^\bullet: \mathbb{P}_{\text{DP}}^\bullet X \rightarrow \mathcal{Q}_V(\mathbf{V}_\bullet X)$  defined by  $s_{\text{DP}}^\bullet(F) \stackrel{\text{def}}{=} \{\nu \in \mathbf{V}_\bullet X \mid \forall h \in \mathcal{L}X, \int_{x \in X} h(x) d\nu \geq F(h)\}$ . Both are continuous, and they form a retraction:  $r_{\text{DP}} \circ s_{\text{DP}}^\bullet = \text{id}_{\mathbb{P}_{\text{DP}}^\bullet X}$ . This is the contents of Proposition 3.22 of [14], only replacing the space  $\mathbb{P}_p^\bullet X$  used there by  $\mathbf{V}_\bullet X$ , since the two are homeomorphic. Additionally,  $r_{\text{DP}}$  and  $s_{\text{DP}}^\bullet$  are natural in  $X$  [19, Lemma 11.2].

This retraction even cuts down to a homeomorphism between  $\mathcal{Q}_V^{cvx}(\mathbf{V}_\bullet X)$  and  $\mathbb{P}_{\text{DP}}^\bullet X$  [14, Theorem 4.15], where the former denotes the subspace of  $\mathcal{Q}_V(\mathbf{V}_\bullet X)$  consisting of convex non-empty compact saturated subsets of  $\mathbf{V}_\bullet X$ . (A subset  $A$  of the latter is *convex* if and only if for all  $G_1, G_2 \in A$ , for every  $r \in [0, 1]$ ,  $rG_1 + (1 - r)G_2 \in A$ .)

Then, there is a continuous map  $\Phi: \mathbf{V}_\bullet \mathcal{Q}_V X \rightarrow \mathbb{P}_{\text{DP}}^\bullet X$  defined by  $\Phi(\mu)(h) \stackrel{\text{def}}{=} \int_{Q \in \mathcal{Q}_V X} \min_{x \in Q} h(x) d\mu$  for every  $h \in \mathcal{L}X$  [19, Lemma 12.5], and the proof of Corollary 12.6 of [19] builds  $\lambda_X^\sharp$  as  $s_{\text{DP}}^\bullet \circ \Phi$ .

**Fact 4.2.**  $\lambda_X^\sharp = s_{\text{DP}}^\bullet \circ \Phi$ .

**Remark 4.3.** In other words, for every  $\mu \in \mathbf{V}\bullet\mathcal{Q}_V X$ ,  $\lambda_X^\sharp(\mu)$  is also the set  $\{\nu \in \mathbf{V}\bullet X \mid \forall h \in \mathcal{L}X, \int_{x \in X} h(x) d\nu \geq \int_{Q \in \mathcal{Q}_V X} \min_{x \in Q} h(x) d\mu\}$ .

In order to illustrate what  $\lambda_X^\sharp$  does, let us look at the special case where  $\mu \stackrel{\text{def}}{=} \sum_{i=1}^n a_i \delta_{Q_i}$ , where  $n \geq 1$ ,  $Q_i \stackrel{\text{def}}{=} \uparrow E_i$ , and where each set  $E_i$  is non-empty and finite. For every  $h \in \mathcal{L}X$ ,  $\int_{Q \in \mathcal{Q}_V X} \min_{x \in Q} h(x) d\mu = \sum_{i=1}^n a_i \min_{x_i \in Q_i} h(x_i) = \sum_{i=1}^n a_i \min_{x_i \in E_i} h(x_i) = \min_{x_1 \in E_1, \dots, x_n \in E_n} \sum_{i=1}^n a_i h(x_i) = \min_{x_1 \in E_1, \dots, x_n \in E_n} \int_{x \in X} h(x) d \sum_{i=1}^n a_i \delta_{x_i}$ . Hence  $\lambda_X^\sharp(\mu) = \uparrow \{\sum_{i=1}^n a_i \delta_{x_i} \mid x_i \in E_i \text{ for every } i \in \{1, \dots, n\}\}$ .

All those maps are continuous. For  $\Phi$ , we have the following.

**Fact 4.4.** For all  $h \in \mathcal{L}X$  and  $r \in \mathbb{R}_+$ ,  $\Phi^{-1}([h > r]) = [h^* > r]$  where  $h^*(Q)$  is defined as  $\min_{x \in Q} h(x)$  for every  $Q \in \mathcal{Q}_V X$ . The map  $h^*$  is in  $\mathcal{L}\mathcal{Q}_V X$ , since for every  $t \in \mathbb{R}_+$ ,  $h^{*-1}([t, \infty]) = \square h^{-1}([t, \infty])$ .

**Fact 4.5.** For every  $h \in \mathcal{L}X$ ,  $(r_{\text{DP}})^{-1}([h > 1]) = \square[h > 1]$ .

The proof that  $s_{\text{DP}}^\bullet$  is continuous is complex [14, Lemma 3.20]. We produce a simpler proof in Lemma 4.6 below, which additionally describes the shape of inverse images of (sub)basic open sets by that map. We note that a base of open subsets of  $\mathcal{Q}_V(\mathbf{V}\bullet X)$  is given by sets of the form  $\square \bigcup_{i \in I} \bigcap_{j=1}^{n_i} [U_{ij} > r_{ij}]$ . Since  $\square$  commutes with directed unions and  $\bigcup_{i \in I} \mathcal{U}_i = \sup_{J \in \mathbb{P}_{\text{fin}}(I)} \bigcup_{i \in J} \mathcal{U}_i$ , since finite unions and finite intersections distribute, and since  $\square$  commutes with finite intersections, a simpler base is given by open subsets of the form  $\square \bigcup_{i=1}^n [U_i > r_i]$ , where each  $U_i$  is open in  $X$  and  $r_i \in \mathbb{R}_+$ . More generally, we consider open subsets of the form  $\square \bigcup_{i=1}^n [h_i > r_i]$  where each  $h_i$  is in  $\mathcal{L}X$  and  $r_i \in \mathbb{R}_+ \setminus \{0\}$ . We may even restrict them to  $n \geq 1$ , since  $\square \emptyset$  is empty. Since  $[h_i > r_i] = [h_i/r_i > 1]$ , we also restrict  $r_i$  to be equal to 1. Let  $\Delta_n \stackrel{\text{def}}{=} \{(t_1, \dots, t_n) \in \mathbb{R}_+^n \mid \sum_{i=1}^n t_i = 1\}$ . We write  $\vec{a}$  for the elements of  $\Delta_n$ ; and  $a_i$  for component number  $i$  of  $\vec{a}$ .

**Lemma 4.6.** For every  $n \geq 1$ , for all  $h_1, \dots, h_n \in \mathcal{L}X$ ,

$$(s_{\text{DP}}^\bullet)^{-1}(\square(\bigcup_{i=1}^n [h_i > 1])) = \bigcup_{\vec{a} \in \Delta_n} \left[ \sum_{i=1}^n a_i h_i > 1 \right].$$

PROOF. Let  $F \in \bigcup_{\vec{a} \in \Delta_n} [\sum_{i=1}^n a_i h_i > 1]$ . Then  $F(\sum_{i=1}^n a_i h_i) > 1$  for some  $\vec{a} \in \Delta_n$ . For every  $\nu \in s_{\text{DP}}^\bullet(F)$ , we then have  $\int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu \geq$

$F(\sum_{i=1}^n a_i h_i) > 1$ . It follows that  $\int_{i=1}^n h_i(x) d\nu > 1$  for some  $i \in \{1, \dots, n\}$ : otherwise  $\int_{i=1}^n h_i(x) d\nu \leq 1$  for every  $i$ , so  $\int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu \leq 1$ . Since  $\nu$  is arbitrary in  $s_{\text{DP}}^\bullet(F)$ ,  $s_{\text{DP}}^\bullet(F) \subseteq \bigcup_{i=1}^n [h_i > 1]$ , so  $F \in s_{\text{DP}}^{-1}(\square(\bigcup_{i=1}^n [h_i > 1]))$ .

Conversely, let  $F \in s_{\text{DP}}^{-1}(\square(\bigcup_{i=1}^n [h_i > r_i]))$ , and let us define  $f: s_{\text{DP}}^\bullet(F) \times \Delta_n \rightarrow \overline{\mathbb{R}}_+$  by  $f(\nu, \vec{a}) \stackrel{\text{def}}{=} \int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu$ . The space  $s_{\text{DP}}^\bullet(F)$ , seen as a subspace of  $\mathbf{V}_\bullet X$ , is non-empty and compact. The map  $f(\_, \vec{a})$  is lower semi-continuous for every  $\vec{a} \in \Delta_n$ , since  $f(\_, \vec{a})^{-1}([t, \infty]) = s_{\text{DP}}^\bullet(F) \cap [\sum_{i=1}^n a_i h_i > t]$  for every  $t \in \mathbb{R}_+$ . The Minimax Theorem 3.3 of [15] then says that: (\*)  $\sup_{\vec{a} \in \Delta_n} \inf_{\nu \in s_{\text{DP}}^\bullet(F)} f(\nu, \vec{a}) = \min_{\nu \in s_{\text{DP}}^\bullet(F)} \sup_{\vec{a} \in \Delta_n} f(\nu, \vec{a})$  (and the min on the right is attained), provided we can show that  $f$  is closely convex in its first argument and closely concave in its second argument. Whatever the latter mean, Remark 3.4 of [15] tells us that maps that preserve pairwise linear combinations of their first arguments (with coefficients  $a$  and  $1-a$ ,  $a \in [0, 1]$ ) are convex hence closely convex in their first argument, and similarly with ‘‘concave’’ instead of ‘‘convex’’; and it is clear that  $f$  is linear in both its arguments.

We recall that  $F = r_{\text{DP}}(s_{\text{DP}}^\bullet(F))$ , so  $F(h) = \min_{\nu \in s_{\text{DP}}^\bullet(F)} \int_{x \in X} h(x) d\nu$  for every  $h \in \mathcal{L}X$ . Since  $F \in s_{\text{DP}}^{-1}(\square(\bigcup_{i=1}^n [h_i > r_i]))$ , for every  $\nu \in s_{\text{DP}}^\bullet(F)$ , there is an  $i \in \{1, \dots, n\}$  such that  $\int_{x \in X} h_i(x) d\nu > 1$ . In particular, there is an  $\vec{a} \in \Delta_n$  such that  $\int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu > 1$ . Therefore  $\min_{\nu \in s_{\text{DP}}^\bullet(F)} \sup_{\vec{a} \in \Delta_n} f(\nu, \vec{a}) > 1$ . By (\*),  $\sup_{\vec{a} \in \Delta_n} \inf_{\nu \in s_{\text{DP}}^\bullet(F)} f(\nu, \vec{a}) > 1$ , so there is an  $\vec{a} \in \Delta_n$  such that  $\inf_{\nu \in s_{\text{DP}}^\bullet(F)} \int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu > 1$ , namely such that  $r_{\text{DP}}(s_{\text{DP}}^\bullet(F))(\sum_{i=1}^n a_i h_i) > 1$ , or equivalently such that  $F(\sum_{i=1}^n a_i h_i) > 1$ . We conclude that  $F \in \bigcup_{\vec{a} \in \Delta_n} [\sum_{i=1}^n a_i h_i > 1]$ .  $\square$

**Lemma 4.7.** *For every  $n \geq 1$ , for all  $h_1, \dots, h_n \in \mathcal{L}X$ ,*

$$\lambda_X^\#^{-1}(\square(\bigcup_{i=1}^n [h_i > 1])) = \bigcup_{\vec{a} \in \Delta_n} \left[ (\sum_{i=1}^n a_i h_i)^* > 1 \right].$$

When  $n = 1$ , there is just one element in  $\Delta_n$ , and that is (1). We obtain the following as a special case.

**Corollary 4.8.** *For every  $h \in \mathcal{L}X$ ,*

$$\begin{aligned} (s_{\text{DP}}^\bullet)^{-1}(\square[h > 1]) &= [h > 1] \\ \lambda_X^\#^{-1}(\square[h > 1]) &= [h^* > 1]. \end{aligned}$$

**Lemma 4.9.**  $\lambda^\sharp$  is a natural transformation.

PROOF. Since  $s_{\text{dp}}^\bullet$  is natural, it suffices to show that  $\Phi$  is natural. Let  $f: X \rightarrow Y$  be any continuous map. We need to show that  $\Phi \circ \mathbf{V}_\bullet \mathcal{Q}_V f = \mathbb{P}_{\text{dp}}^\bullet f \circ \Phi$ . Let  $\mu \in \mathbf{V}_\bullet \mathcal{Q}_V X$  and  $h \in \mathcal{L}Y$ . Then  $(\Phi \circ \mathbf{V}_\bullet \mathcal{Q}_V f)(\mu)(h) = \Phi(\mathcal{Q}_V f[\mu])(h) = \int_{Q \in \mathcal{Q}_V Y} \min_{y \in Q} h(y) d\mathcal{Q}_V f[\mu]$ . By the change of variable formula, this is equal to  $\int_{Q \in \mathcal{Q}_V X} \min_{y \in \mathcal{Q}_V f(Q)} h(y) d\mu$ . Now  $\min_{y \in \mathcal{Q}_V f(Q)} h(y) = \min_{y \in \uparrow f[Q]} h(y) = \min_{x \in Q} h(f(x))$  (because  $h$ , being lower semicontinuous, is monotonic). We therefore obtain that  $(\Phi \circ \mathbf{V}_\bullet \mathcal{Q}_V f)(\mu)(h) = \int_{Q \in \mathcal{Q}_V X} \min_{x \in Q} h(f(x)) d\mu$ . But  $(\mathbb{P}_{\text{dp}}^\bullet f \circ \Phi)(\mu)(h) = \mathbb{P}_{\text{dp}}^\bullet f(\Phi(\mu))(h) = \Phi(\mu)(h \circ f) = \int_{Q \in \mathcal{Q}_V X} \min_{x \in Q} h(f(x)) d\mu$ .  $\square$

**Lemma 4.10.**  $\lambda_X^\sharp \circ \mathbf{V}_\bullet \eta_X^\mathcal{Q} = \eta_{\mathbf{V}_\bullet X}^\mathcal{Q}$ .

PROOF. For every  $\nu \in \mathbf{V}_\bullet X$ , the elements of  $(\lambda_X^\sharp \circ \mathbf{V}_\bullet \eta_X^\mathcal{Q})(\nu)$  are those  $\nu' \in \mathbf{V}_\bullet X$  such that  $\nu'(U) \geq \mathbf{V}_\bullet \eta_X^\mathcal{Q}(\nu)(\square U)$  for every  $U \in \mathcal{O}X$ . Now  $\mathbf{V}_\bullet \eta_X^\mathcal{Q}(\nu)(\square U) = \nu((\eta_X^\mathcal{Q})^{-1}(\square U)) = \nu(U)$ , so  $(\lambda_X^\sharp \circ \mathbf{V}_\bullet \eta_X^\mathcal{Q})(\nu)$  is simply the collection of  $\nu' \in \mathbf{V}_\bullet X$  such that  $\nu' \geq \nu$ , namely  $\eta_{\mathbf{V}_\bullet X}^\mathcal{Q}(\nu)$ .  $\square$

Lemma 4.10 is the first of the weak distributivity laws. The proofs of the others are more complex. We will use the following several times:

**Trick A.** In order to show that two continuous maps  $f, g: Y \rightarrow Z$  are equal, where  $Z$  is a  $T_0$  space, check that  $f^{-1}(V) = g^{-1}(V)$  for every element  $V$  of a subbase of the topology of  $Z$ .

Indeed, if the condition holds, then  $f^{-1}(V) = g^{-1}(V)$  for every open subset  $V$  of  $Z$ . Hence, for every point  $y \in Y$ ,  $f(y)$  and  $g(y)$  will have the same open neighborhoods and therefore be equal, since  $Z$  is  $T_0$ .

We also note that any space of the form  $\mathbf{V}_\bullet Y$ ,  $\mathbb{P}_{\text{dp}}^\bullet Y$ , or  $\mathcal{Q}_V Y$  is  $T_0$ , since their specialization preorderings are  $\leq$ ,  $\leq$ , and  $\subseteq$  respectively, and they are antisymmetric.

In general, the maps  $f$  and  $g$  that are to be shown equal are compositions  $f_1 \circ \dots \circ f_m$  and  $g_1 \circ \dots \circ g_n$ , and we will use Trick A by writing lists of the following kind:  $V \xrightarrow{f_1} V_1 \xrightarrow{f_2} \dots \xrightarrow{f_m} V_m$  and  $V \xrightarrow{g_1} V'_1 \xrightarrow{g_2} \dots \xrightarrow{g_n} V'_n$ , meaning that  $V_1 = f_1^{-1}(V)$ , etc., and finally checking that  $V_m = V'_n$ .

For example, an alternate proof of Lemma 4.10 along these lines is:

$$\begin{aligned}
\sqcap \left( \bigcup_{i=1}^n [h_i > 1] \right) &\xrightarrow{\lambda_X^\sharp} \bigcup_{\vec{a} \in \Delta_n} \left[ \left( \sum_{i=1}^n a_i h_i \right)^* > 1 \right] && \text{by Lemma 4.7} \\
&\xrightarrow{\mathbf{V} \bullet \eta_X^\mathcal{Q}} \bigcup_{\vec{a} \in \Delta_n} \left[ \left( \sum_{i=1}^n a_i h_i \right)^* \circ \eta_X^\mathcal{Q} > 1 \right] \\
&= \bigcup_{\vec{a} \in \Delta_n} \left[ \sum_{i=1}^n a_i h_i > 1 \right],
\end{aligned}$$

since for every  $h \in \mathcal{L}X$ ,  $h^* \circ \eta_X^\mathcal{Q} = h$ , and:

$$\sqcap \left( \bigcup_{i=1}^n [h_i > 1] \right) \xrightarrow{\eta_{\mathbf{V} \bullet X}^\mathcal{Q}} \bigcup_{i=1}^n [h_i > 1].$$

Now  $\bigcup_{\vec{a} \in \Delta_n} [\sum_{i=1}^n a_i h_i > 1] = \bigcup_{i=1}^n [h_i > 1]$ , as open subsets of  $\mathbf{V} \bullet X$ : for every  $\nu \in \mathbf{V} \bullet X$ , if  $\int_{x \in X} h_i(x) d\nu > 1$  for some  $i$ , then  $\int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu > 1$  for the tuple  $\vec{a}$  whose  $i$ th component is 1 and whose other components are 0. Conversely, if  $\int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu > 1$ , then  $\int_{x \in X} h_i(x) d\nu > 1$  for some  $i$ , otherwise  $\int_{x \in X} h_i(x) d\nu \leq 1$  for every  $i$ , so  $\int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu = \sum_{i=1}^n a_i \int_{x \in X} h_i(x) d\nu \leq 1$ .

We will also use the following.

**Trick B.** *In order to show that  $f = g$ , where  $f$  and  $g$  are continuous maps from a space  $Y$  to a space of the form  $\mathcal{Q}_V \mathbf{V} \bullet Z$ , show that  $f(y)$  and  $g(y)$  are convex for every  $y \in Y$ , and then show that  $r_{\text{DP}} \circ f = r_{\text{DP}} \circ g$ .*

Indeed, we recall that  $r_{\text{DP}}$  is a homeomorphism of  $\mathcal{Q}_V^{cvx}(\mathbf{V} \bullet Z)$  onto  $\mathbb{P}_{\text{DP}}^\bullet Z$ .

**Lemma 4.11.**  $\lambda_X^\sharp \circ \mathbf{V} \bullet \mu_X^\mathcal{Q} = \mu_{\mathbf{V} \bullet X}^\mathcal{Q} \circ \mathcal{Q}_V \lambda_X^\sharp \circ \lambda_{\mathcal{Q}_V X}^\sharp$ .

PROOF. Since  $\lambda_X^\sharp = s_{\text{DP}}^\bullet \circ \Phi$  and the image of  $s_{\text{DP}}^\bullet$  consists of convex subsets, the left-hand side takes its values in convex sets. For the right-hand side, for every  $\xi \in \mathbf{V} \bullet \mathcal{Q}_V \mathcal{Q}_V X$ , let  $Q' \stackrel{\text{def}}{=} (\mu_{\mathbf{V} \bullet X}^\mathcal{Q} \circ \mathcal{Q}_V \lambda_X^\sharp \circ \lambda_{\mathcal{Q}_V X}^\sharp)(\xi)$ . We want to show that  $Q'$  is convex. The elements  $\nu$  of  $Q'$  are those of  $\bigcup (\mathcal{Q}_V \lambda_X^\sharp \circ \lambda_{\mathcal{Q}_V X}^\sharp)(\xi)$ , namely those that are in some element  $\mathcal{Q}$  of  $(\mathcal{Q}_V \lambda_X^\sharp \circ \lambda_{\mathcal{Q}_V X}^\sharp)(\xi) = \uparrow \lambda_X^\sharp[\lambda_{\mathcal{Q}_V X}^\sharp(\xi)] = \{\mathcal{Q} \in \mathcal{Q}_V \mathbf{V} \bullet X \mid \exists \mu \in \lambda_{\mathcal{Q}_V X}^\sharp(\xi), \mathcal{Q} \subseteq \lambda_X^\sharp(\mu)\}$ . (Remember that the specialization ordering on  $\mathcal{Q}_V \mathbf{V} \bullet X$  is reverse inclusion  $\supseteq$ .) Simplifying this, and since  $\uparrow \mu$  is always non-empty, compact and saturated, hence

can take the place of  $\mathcal{Q}$ , the elements  $\nu$  of  $\mathcal{Q}'$  are those such that there is a  $\mu \in \lambda_{\mathcal{Q}_V X}^\sharp(\xi)$  such that  $\nu \in \lambda_X^\sharp(\mu)$ .

Let  $a \in [0, 1]$ , and  $\nu_1, \nu_2 \in \mathcal{Q}'$ . Hence there is a  $\mu_1 \in \lambda_{\mathcal{Q}_V X}^\sharp(\xi)$  such that  $\nu_1 \in \lambda_X^\sharp(\mu_1)$  and there is a  $\mu_2 \in \lambda_{\mathcal{Q}_V X}^\sharp(\xi)$  such that  $\nu_2 \in \lambda_X^\sharp(\mu_2)$ . We note that  $\lambda_{\mathcal{Q}_V X}^\sharp(\xi) = (s_{\text{DP}}^\bullet \circ \Phi)(\xi)$  is convex, so  $a\mu_1 + (1-a)\mu_2 \in \lambda_{\mathcal{Q}_V X}^\sharp(\xi)$ . Additionally,  $a\nu_1 + (1-a)\nu_2 \in \lambda_X^\sharp(a\mu_1 + (1-a)\mu_2)$ : for every  $U \in \mathcal{O}X$ ,  $\nu_1(U) \geq \mu_1(\square U)$  and  $\nu_2(U) \geq \mu_2(\square U)$  by definition of  $\lambda_X^\sharp$ , so  $(a\nu_1 + (1-a)\nu_2)(U) \geq (a\mu_1 + (1-a)\mu_2)(\square U)$ . Therefore there is a  $\mu \in \lambda_{\mathcal{Q}_V X}^\sharp(\xi)$ , namely  $\mu \stackrel{\text{def}}{=} a\mu_1 + (1-a)\mu_2$ , such that  $a\nu_1 + (1-a)\nu_2 \in \lambda_X^\sharp(\mu)$ . Equivalently,  $a\nu_1 + (1-a)\nu_2$  is in  $\mathcal{Q}'$ . Hence  $\mathcal{Q}'$  is convex, as promised.

It follows that we can use Trick B, and so we prove the following instead:

$$\Phi \circ \mathbf{V} \bullet \mu_X^\mathcal{Q} = r_{\text{DP}} \circ \mu_{\mathbf{V} \bullet X}^\mathcal{Q} \circ \mathcal{Q}_V \lambda_X^\sharp \circ \lambda_{\mathcal{Q}_V X}^\sharp.$$

Indeed,  $r_{\text{DP}} \circ \lambda_X^\sharp = r_{\text{DP}} \circ s_{\text{DP}}^\bullet \circ \Phi = \Phi$ , which allows us to simplify the left-hand side.

We now use Trick A:

$$\begin{aligned} [h > 1] &\stackrel{\Phi}{\implies} [h^* > 1] \\ &\stackrel{\mathbf{V} \bullet \mu_X^\mathcal{Q}}{\implies} [h^* \circ \mu_X^\mathcal{Q} > 1] = [h^{**} > 1], \end{aligned}$$

where for every  $\mathcal{Q} \in \mathcal{Q}_V \mathcal{Q}_V X$ ,  $h^{**}(\mathcal{Q}) = \min_{Q \in \mathcal{Q}} h^*(Q) = \min_{Q \in \mathcal{Q}} \min_{x \in Q} h(x)$ ; and indeed, the latter is equal to  $\min_{x \in \cup \mathcal{Q}} h(x) = h^*(\mu_X^\mathcal{Q}(\mathcal{Q}))$ .

$$\begin{aligned} [h > 1] &\stackrel{r_{\text{DP}}}{\implies} \square [h > 1] && \text{by Fact 4.5} \\ &\stackrel{\mu_{\mathbf{V} \bullet X}^\mathcal{Q}}{\implies} \square \square [h > 1] \\ &\stackrel{\mathcal{Q}_V \lambda_X^\sharp}{\implies} \square \lambda_X^\sharp{}^{-1}(\square [h > 1]) = \square [h^* > 1] && \text{by Corollary 4.8} \\ &\stackrel{\lambda_{\mathcal{Q}_V X}^\sharp}{\implies} [h^{**} > 1] && \text{by Corollary 4.8.} \end{aligned}$$

□

**Lemma 4.12.**  $\lambda_X^\sharp \circ \mu_{\mathcal{Q}_V X} = \mathcal{Q}_V \mu_X \circ \lambda_{\mathbf{V} \bullet X}^\sharp \circ \mathbf{V} \bullet \lambda_X^\sharp$ .

PROOF. The left-hand side takes its values in convex sets. We claim that the right-hand side does, too. For every  $\xi \in \mathbf{V} \bullet \mathbf{V} \bullet \mathcal{Q}_V X$ , the elements of

$A \stackrel{\text{def}}{=} (\mathcal{Q}_V \mu_X \circ \lambda_{\mathbf{V}\bullet X}^\# \circ \mathbf{V}\bullet \lambda_X^\#)(\xi)$  are the continuous valuations  $\nu \in \mathbf{V}\bullet X$  such that  $\nu \geq \mu_X(\mu)$  for some  $\mu \in \lambda_{\mathbf{V}\bullet X}^\#(\lambda_X^\#[\xi])$ . For all  $a \in [0, 1]$ ,  $\nu_1, \nu_2 \in A$ , there are  $\mu_1, \mu_2 \in \lambda_{\mathbf{V}\bullet X}^\#(\lambda_X^\#[\xi])$  such that  $\nu_1 \geq \mu_X(\mu_1)$  and  $\nu_2 \geq \mu_X(\mu_2)$ . Then  $\mu \stackrel{\text{def}}{=} a\mu_1 + (1-a)\mu_2$  is in  $\lambda_{\mathbf{V}\bullet X}^\#(\lambda_X^\#[\xi])$ , since  $\lambda_{\mathbf{V}\bullet X}^\#$  takes its values in convex sets. In turn,  $a\nu_1 + (1-a)\nu_2 \geq a\mu_X(\mu_1) + (1-a)\mu_X(\mu_2) = a(\mathcal{U} \in \mathcal{O}\mathbf{V}\bullet X \mapsto \int_{\nu \in \mathbf{V}\bullet X} \nu(\mathcal{U}) d\mu_1) + (1-a)(\mathcal{U} \in \mathcal{O}\mathbf{V}\bullet X \mapsto \int_{\nu \in \mathbf{V}\bullet X} \nu(\mathcal{U}) d\mu_2) = \mu_X(\mu)$ , so  $a\nu_1 + (1-a)\nu_2 \in A$ , showing that  $A$  is convex.

We can therefore use Trick B, and since  $r_{\text{DP}} \circ \lambda_X^\# = r_{\text{DP}} \circ s_{\text{DP}}^\bullet \circ \Phi = \Phi$ , it remains to show:

$$\Phi \circ \mu_{\mathcal{Q}_V X} = r_{\text{DP}} \circ \mathcal{Q}_V \mu_X \circ \lambda_{\mathbf{V}\bullet X}^\# \circ \mathbf{V}\bullet \lambda_X^\#.$$

To this end, we use Trick A:

$$\begin{aligned} [h > 1] &\stackrel{\Phi}{\implies} [h^* > 1] && \text{by Fact 4.4} \\ &\stackrel{\mu_{\mathcal{Q}_V X}}{\implies} \left[ \left( \mu \in \mathbf{V}\bullet \mathcal{Q}_V X \mapsto \int_{Q \in \mathcal{Q}_V X} h^*(Q) d\mu \right) > 1 \right] && \text{by (1)} \end{aligned}$$

while:

$$\begin{aligned} [h > 1] &\stackrel{r_{\text{DP}}}{\implies} \square[h > 1] && \text{by Fact 4.5} \\ &\stackrel{\mathcal{Q}_V \mu_X}{\implies} \square \mu_X^{-1}([h > 1]) = \square \left[ \left( \nu \in \mathbf{V}\bullet X \mapsto \int_{x \in X} h(x) d\nu \right) > 1 \right] && \text{by (1)} \\ &\stackrel{\lambda_{\mathbf{V}\bullet X}^\#}{\implies} \left[ \left( \nu \in \mathbf{V}\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)^* > 1 \right] && \text{by Corollary 4.8} \\ &\stackrel{\mathbf{V}\bullet \lambda_X^\#}{\implies} \left[ \left( \nu \in \mathbf{V}\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)^* \circ \lambda_X^\# > 1 \right]. \end{aligned}$$

In order to conclude, it suffices to show that:

$$\left( \nu \in \mathbf{V}\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)^* \circ \lambda_X^\# = \left( \mu \in \mathbf{V}\bullet \mathcal{Q}_V X \mapsto \int_{Q \in \mathcal{Q}_V X} h^*(Q) d\mu \right).$$

The left-hand side maps every  $\mu \in \mathbf{V}\bullet \mathcal{Q}_V X$  to

$$\begin{aligned} \min_{\nu \in s_{\text{DP}}^\bullet(\Phi(\mu))} \int_{x \in X} h(x) d\nu &= r_{\text{DP}}(s_{\text{DP}}^\bullet(\Phi(\mu)))(h) \\ &= \Phi(\mu)(h) = \int_{Q \in \mathcal{Q}_V X} h^*(Q), \end{aligned}$$

and this completes the proof.  $\square$



Putting Lemma 4.9, Lemma 4.10, Lemma 4.11 and Lemma 4.12 together, we obtain the following.

**Theorem 4.13.** *Let  $\bullet$  be nothing, “ $\leq 1$ ”, or “1”. Then  $\lambda^\sharp$  is a weak distributive law of  $\mathcal{Q}_V$  over  $\mathbf{V}_\bullet$  on the category  $\mathbf{Top}$  of topological spaces.*

## 5. The associated weak composite monad

We will show that the associated weak composite monad (see [9, Section 3.2], or [20, page 46]) is isomorphic to the monad  $\mathbb{P}_{\mathbf{DP}}^\bullet$  of superlinear (resp., and subnormalized, resp., and normalized) previsions, or equivalently to the isomorphic monad  $\mathcal{Q}_V^{conv} \mathbf{V}_\bullet$ .

There is an idempotent:

$$\mathcal{Q}_V \mathbf{V}_\bullet X \xrightarrow{\eta_{\mathcal{Q}_V \mathbf{V}_\bullet X}} \mathbf{V}_\bullet \mathcal{Q}_V \mathbf{V}_\bullet X \xrightarrow{\lambda_{\mathbf{V}_\bullet X}^\sharp} \mathcal{Q}_V \mathbf{V}_\bullet \mathbf{V}_\bullet X \xrightarrow{\mathcal{Q}_V \mu_X} \mathcal{Q}_V \mathbf{V}_\bullet X \quad (2)$$

and splitting it defines the action of the (functor part of the) associated weak composite monad on each object  $X$ . *Splitting* an idempotent  $e$  means writing it as  $\iota \circ \pi$ , for a pair of morphisms such that  $\pi \circ \iota = \text{id}$ .

**Lemma 5.1.** *A splitting of (2) is  $\mathcal{Q}_V \mathbf{V}_\bullet X \xrightarrow{r_{\mathbf{DP}}} \mathbb{P}_{\mathbf{DP}}^\bullet X \xrightarrow{s_{\mathbf{DP}}^\bullet} \mathcal{Q}_V \mathbf{V}_\bullet X$ .*

PROOF. Let  $f$  denote the idempotent (2). We recall that  $\lambda_{\mathbf{V}_\bullet X}^\sharp = s_{\mathbf{DP}}^\bullet \circ \Phi$ . In particular, any element in its range is convex. The image of any convex element  $Q$  of a space  $\mathcal{Q}_V \mathbf{V}_\bullet$  under  $\mathcal{Q}_V f$ , where  $f$  is any linear continuous map from  $\mathbf{V}_\bullet Y \rightarrow \mathbf{V}_\bullet Z$ , is convex; and  $\mu_X$  is linear. Therefore  $f$  takes its values in  $\mathcal{Q}_V^{conv} \mathbf{V}_\bullet X$ .

Using Trick B, and simplifying using the fact that  $r_{\mathbf{DP}} \circ s_{\mathbf{DP}}^\bullet$  is the identity, it remains to show that  $r_{\mathbf{DP}}$  is equal to  $r_{\mathbf{DP}} \circ f$ , namely to:

$$\mathcal{Q}_V \mathbf{V}_\bullet X \xrightarrow{\eta_{\mathcal{Q}_V \mathbf{V}_\bullet X}} \mathbf{V}_\bullet \mathcal{Q}_V \mathbf{V}_\bullet X \xrightarrow{\lambda_{\mathbf{V}_\bullet X}^\sharp} \mathcal{Q}_V \mathbf{V}_\bullet \mathbf{V}_\bullet X \xrightarrow{\mathcal{Q}_V \mu_X} \mathcal{Q}_V \mathbf{V}_\bullet X \xrightarrow{r_{\mathbf{DP}}} \mathbb{P}_{\mathbf{DP}}^\bullet X$$

Using Trick A,

$$\begin{aligned}
[h > 1] &\xrightarrow{r_{\text{DP}}} \square[h > 1] && \text{by Fact 4.5} \\
&\xrightarrow{\mathcal{Q}_V \mu_X} \square \mu_X^{-1}([h > 1]) \\
&= \square \left[ \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right) > 1 \right] && \text{by (1)} \\
&\xrightarrow{\lambda_{\mathbf{V}_\bullet X}^\sharp} \left[ \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)^* > 1 \right] && \text{by Corollary 4.8} \\
&\xrightarrow{\eta_{\mathcal{Q}_V \mathbf{V}_\bullet X}} \left\{ Q \in \mathcal{Q}_V \mathbf{V}_\bullet X \mid \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)^* (Q) > 1 \right\} \\
&= \{ Q \in \mathcal{Q}_V \mathbf{V}_\bullet X \mid r_{\text{DP}}(Q)(h) > 1 \} = r_{\text{DP}}^{-1}([h > 1]).
\end{aligned}$$

□

We can always split an idempotent  $e: Z \rightarrow Z$  in **Top** as the composition of its corestriction to  $e[Z]$  with the inclusion map  $e[Z] \rightarrow Z$ . In Lemma 5.1, we might have chosen the homeomorphic space  $\mathcal{Q}_V^{cux} \mathbf{V}_\bullet X$  instead of  $\mathbb{P}_{\text{DP}}^\bullet X$ .

Generalizing (2), there is an idempotent:

$$\mathcal{Q}_V X \xrightarrow{\eta_{\mathcal{Q}_V X}} \mathbf{V}_\bullet \mathcal{Q}_V X \xrightarrow{\lambda_X^\sharp} \mathcal{Q}_V \mathbf{V}_\bullet X \xrightarrow{\mathcal{Q}_V \alpha} \mathcal{Q}_V X \quad (3)$$

for every  $\mathbf{V}_\bullet$ -algebra  $\alpha$ . Finding out what its splitting is is harder in general, especially because we have no complete characterization of  $\mathbf{V}_\bullet$ -algebras in **Top**. We will only need to compute this splitting for one additional  $\mathbf{V}_\bullet$ -algebra.

**Lemma 5.2.** *For every topological space  $X$ , there is a  $\mathbf{V}_\bullet$ -algebra  $\alpha_{\text{DP}}^\bullet: \mathbf{V}_\bullet \mathbb{P}_{\text{DP}}^\bullet X \rightarrow \mathbb{P}_{\text{DP}}^\bullet X$  defined by  $\alpha_{\text{DP}}^\bullet(\xi)(h) \stackrel{\text{def}}{=} \int_{F \in \mathbb{P}_{\text{DP}}^\bullet X} F(h) d\xi$  for all  $\xi \in \mathbf{V}_\bullet \mathbb{P}_{\text{DP}}^\bullet X$  and  $h \in \mathcal{L}X$ .*

*The idempotent (3) with  $\alpha \stackrel{\text{def}}{=} \alpha_{\text{DP}}^\bullet$  splits as:*

$$\mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X \longrightarrow \mathcal{Q}_V^{cux} \mathbb{P}_{\text{DP}}^\bullet X \longrightarrow \mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X,$$

*where the arrow on the right is the inclusion map, and the arrow on the left is the corestriction of (3), and maps every  $Q \in \mathcal{Q}_V^{cux} \mathbb{P}_{\text{DP}}^\bullet X$  to itself.*

PROOF. The map  $F \mapsto F(h)$  is lower semicontinuous, since the inverse image of  $]t, \infty]$  is  $[h > t]$ , for all  $h \in \mathcal{L}X$  and  $t \in \mathbb{R}_+$ . Hence the integral  $\int_{F \in \mathbb{P}_{\text{DP}}^\bullet X} F(h) d\xi$  makes sense. The map  $\alpha_{\text{DP}}^\bullet(\xi)$  is Scott-continuous because each  $F \in \mathbb{P}_{\text{DP}}^\bullet X$  is and because integration with respect to  $\nu$  is Scott-continuous; it is superlinear because  $F$  is and integration with respect to  $\xi$  is linear. Additionally,  $(\alpha_{\text{DP}}^\bullet)^{-1}([h > r]) = [(F \mapsto F(h)) > r]$ , so  $\alpha_{\text{DP}}^\bullet$  is continuous.

We claim that the map  $\alpha_{\text{DP}}^\bullet$  is a  $\mathbf{V}_\bullet$ -algebra. We have  $\alpha_{\text{DP}}^\bullet(\eta_{\mathbb{P}_{\text{DP}}^\bullet X}(F))(h) = \int_{F' \in \mathbb{P}_{\text{DP}}^\bullet X} F'(h) d\delta_F = F(h)$ , so  $\alpha_{\text{DP}}^\bullet \circ \eta_{\mathbb{P}_{\text{DP}}^\bullet X}$  is the identity map. In order to show that  $\alpha_{\text{DP}}^\bullet \circ \mu_{\mathbb{P}_{\text{DP}}^\bullet X} = \alpha_{\text{DP}}^\bullet \circ \mathbf{V}_\bullet \alpha_{\text{DP}}^\bullet$ , we use Trick A:

$$\begin{aligned}
[h > 1] &\xrightarrow{\alpha_{\text{DP}}^\bullet} [(F \in \mathbb{P}_{\text{DP}}^\bullet X \mapsto F(h)) > 1] \\
&\xrightarrow{\mu_{\mathbb{P}_{\text{DP}}^\bullet X}} [(\xi \in \mathbf{V}_\bullet \mathbb{P}_{\text{DP}}^\bullet X \mapsto \int_{F \in \mathbb{P}_{\text{DP}}^\bullet X} F(h) d\xi) > 1] \quad \text{by (1)} \\
[h > 1] &\xrightarrow{\alpha_{\text{DP}}^\bullet} [(F \in \mathbb{P}_{\text{DP}}^\bullet X \mapsto F(h)) > 1] \\
&\xrightarrow{\mathbf{V}_\bullet \alpha_{\text{DP}}^\bullet} [(F \in \mathbb{P}_{\text{DP}}^\bullet X \mapsto F(h)) \circ \alpha_{\text{DP}}^\bullet > 1] \\
&= [(\xi \in \mathbf{V}_\bullet \mathbb{P}_{\text{DP}}^\bullet X \mapsto \alpha_{\text{DP}}^\bullet(\xi)(h)) > 1] \\
&= [(\xi \in \mathbf{V}_\bullet \mathbb{P}_{\text{DP}}^\bullet X \mapsto \int_{F \in \mathbb{P}_{\text{DP}}^\bullet X} F(h) d\xi) > 1].
\end{aligned}$$

In order to justify the announced splitting of (3), we need to show that the image of (3) is exactly  $\mathcal{Q}_V^{cvx} \mathbb{P}_{\text{DP}}^\bullet X$ . For every  $Q \in \mathcal{Q}_V X$ , the image of  $Q$  by (3) is  $\uparrow \alpha_{\text{DP}}^\bullet[Q']$  where  $Q' \stackrel{\text{def}}{=} \lambda_{\mathbb{P}_{\text{DP}}^\bullet X}^\#(\delta_Q)$  is convex, and  $\alpha_{\text{DP}}^\bullet$  is linear (in its  $\xi$  argument), so that image is convex.

In order to see that every element  $Q$  of  $\mathcal{Q}_V^{cvx} \mathbb{P}_{\text{DP}}^\bullet X$  is reached as the image of some element by (3), we simply show that  $Q$  is its own image by (3), or

equivalently, that (3) restricts to the identity on  $\mathcal{Q}_{\mathbb{V}}^{cvx} \mathbb{P}_{\text{DP}}^{\bullet} X$ . We use Trick A:

$$\begin{aligned}
\sqcap \bigcup_{i=1}^n [h_i > 1] &\xrightarrow{\mathcal{Q}_{\mathbb{V}}^{\alpha_{\text{DP}}^{\bullet}}} \sqcap \bigcup_{i=1}^n (\alpha_{\text{DP}}^{\bullet})^{-1}([h_i > 1]) \\
&= \sqcap \bigcup_{i=1}^n [(F \mapsto F(h_i)) > 1] \\
&\xrightarrow{\lambda_{\mathbb{P}_{\text{DP}}^{\bullet} X}^{\#}} \bigcup_{\vec{a} \in \Delta_n} \left[ \left( F \mapsto \sum_{i=1}^n a_i F(h_i) \right)^* > 1 \right] && \text{by Lemma 4.7} \\
&\xrightarrow{\eta_{\mathcal{Q}_{\mathbb{V}}^{\text{DP}}^{\bullet} X}} \bigcup_{\vec{a} \in \Delta_n} \left( \left( F \mapsto \sum_{i=1}^n a_i F(h_i) \right)^* \right)^{-1} ([1, \infty]),
\end{aligned}$$

and we claim that the convex elements  $Q$  of  $\mathcal{Q}_{\mathbb{V}} \mathbb{P}_{\text{DP}}^{\bullet} X$  that are in the latter open set are exactly those of  $\sqcap \bigcup_{i=1}^n [h_i > 1]$ , which will show the claim.

We have  $Q \in \bigcup_{\vec{a} \in \Delta_n} \left( (F \mapsto \sum_{i=1}^n a_i F(h_i))^* \right)^{-1} ([1, \infty])$  if and only there is an  $\vec{a} \in \Delta_n$  such that  $\min_{F \in Q} \sum_{i=1}^n a_i F(h_i) > 1$ . For every  $Q \in \mathcal{Q}_{\mathbb{V}}^{cvx} \mathbb{P}_{\text{DP}}^{\bullet} X$  and for every  $i \in \{1, \dots, n\}$ , let  $f(F, \vec{a}) \stackrel{\text{def}}{=} \sum_{i=1}^n a_i F(h_i)$ . This defines a function from  $Q \times \Delta_n$  to  $\overline{\mathbb{R}}_+$  that is lower semicontinuous in  $Q$  (since  $f$  is the composition of various sums and products with the functions  $F \mapsto F(h_i)$ , which are lower semicontinuous), is such that  $f(aF_1 + (1-a)F_2, i) = af(F_1, i) + (1-a)f(F_2, i)$  for all  $a \in [0, 1]$  and  $F_1, F_2 \in Q$  (we need  $Q$  to be convex simply to make sure that  $aF_1 + (1-a)F_2$  still lies in  $Q$ ) hence is convex (the same property, with  $\geq$  instead of  $=$ ) hence closely convex in its first argument (see Remark 3.4 of [15]). It is also concave hence closely concave in its second argument. By the Minimax Theorem 3.3 of [15], therefore,  $\sup_{\vec{a} \in \Delta_n} \min_{F \in Q} f(F, \vec{a}) = \min_{F \in Q} \sup_{\vec{a} \in \Delta_n} f(F, \vec{a})$ . For every  $Q \in \bigcup_{\vec{a} \in \Delta_n} \left( (F \mapsto \sum_{i=1}^n a_i F(h_i))^* \right)^{-1} ([1, \infty])$ , the left-hand side is strictly larger than 1, so the right-hand side is, too. Hence, for every  $F \in Q$ , there is an  $\vec{a} \in \Delta_n$  such that  $\sum_{i=1}^n a_i F(h_i) > 1$ . It follows easily that  $F(h_i) > 1$  for some  $i \in \{1, \dots, n\}$ , so  $Q \in \sqcap \bigcup_{i=1}^n [h_i > 1]$ . Conversely, if  $Q \in \sqcap \bigcup_{i=1}^n [h_i > 1]$ , then for every  $F \in Q$ , there is an  $i \in \{1, \dots, n\}$  such that  $F(h_i) > 1$ , so there is an  $\vec{a} \in \Delta_n$  such that  $\sum_{i=1}^n a_i F(h_i) > 1$ , and therefore  $\sup_{\vec{a} \in \Delta_n} \min_{F \in Q} f(F, \vec{a}) > 1$ , showing that  $Q$  is in  $\bigcup_{\vec{a} \in \Delta_n} \left( (F \mapsto \sum_{i=1}^n a_i F(h_i))^* \right)^{-1} ([1, \infty])$ .  $\square$

Garner showed that, for all monads  $S$  and  $T$  on a category where idempotent splits, every weak distributive law  $\lambda: TS \rightarrow ST$  yields a weak lifting  $\overline{S}$  of  $S$  to the category of  $T$ -algebras, built as follows.

A  $T$ -semialgebra is a morphism  $\alpha: TX \rightarrow X$  such that  $\alpha \circ \mu_X^T = \alpha \circ T\alpha$ . Given a weak distributive law  $\lambda, S\alpha \circ \lambda_X \circ \eta_{SX}^T: SX \rightarrow SX$  is an idempotent, and also a morphism of  $T$ -semialgebras. We can split it in **Top**, as:

$$SX \xrightarrow{\pi_\alpha} \overline{S}X \xrightarrow{\iota_\alpha} SX$$

since idempotents split in **Top**. We recall that splitting means that the idempotent is equal to  $\iota_\alpha \circ \pi_\alpha$ , and  $\pi_\alpha \circ \iota_\alpha$  is the identity. This actually splits the idempotent in the category of  $T$ -semialgebras, as follows:

$$\begin{array}{ccccc} TSX & \xrightarrow{T\pi_\alpha} & T\overline{S}X & \xrightarrow{T\iota_\alpha} & TSX \\ \lambda_X \downarrow & & \lambda_X \circ T\iota_\alpha \downarrow & & \lambda_X \downarrow \\ STX & & STX & & STX \\ S\alpha \downarrow & & \pi_\alpha \circ S\alpha \downarrow & & S\alpha \downarrow \\ SX & \xrightarrow{\pi_\alpha} & \overline{S}X & \xrightarrow{\iota_\alpha} & SX \end{array}$$

Additionally, the middle vertical arrow  $\overline{S}\alpha \stackrel{\text{def}}{=} \pi_\alpha \circ S\alpha \circ \lambda_X \circ T\iota_\alpha$  is not just a  $T$ -semialgebra, but a  $T$ -algebra.

This defines the object part of an endofunctor  $\overline{S}$  on the category of  $T$ -algebras. On morphisms, the action of  $\overline{S}$  is given by: for all  $T$ -algebras  $\alpha: TX \rightarrow X$  and  $\beta: TY \rightarrow Y$ , for every  $T$ -algebra morphism  $f: \alpha \rightarrow \beta$ ,  $\overline{S}f \stackrel{\text{def}}{=} \pi_\beta \circ Sf \circ \iota_\alpha$ .

$\overline{S}$  is part of a monad, whose unit  $\eta_\alpha^{\overline{S}}$  is  $\pi_\alpha \circ \eta_X^S$ , and whose multiplication is  $\mu_\alpha^{\overline{S}} \stackrel{\text{def}}{=} \pi_\alpha \circ \mu_X^S \circ S\iota_\alpha \circ \iota_{\overline{S}\alpha}$ . All of that was proved by Garner.

**Lemma 5.3.** *For  $S \stackrel{\text{def}}{=} \mathcal{Q}_V$ ,  $T \stackrel{\text{def}}{=} \mathbf{V}_\bullet$ ,  $\lambda \stackrel{\text{def}}{=} \lambda^\sharp$ , the monad  $\overline{S}$  on the category of  $\mathbf{V}_\bullet$ -algebras has the following properties:*

1. *For every free  $\mathbf{V}_\bullet$ -algebra  $\mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$ ,  $\overline{S}\mu_X: \mathbf{V}_\bullet \mathbb{P}_{\text{DP}}^\bullet X \rightarrow \mathbb{P}_{\text{DP}}^\bullet X$  is  $\alpha_{\text{DP}}^\bullet$  (see Lemma 5.2).*
2. *The unit  $\eta_{\mu_X}^{\overline{S}}$  evaluated at the free  $\mathbf{V}_\bullet$ -algebra  $\mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$  is  $r_{\text{DP}} \circ \eta_{\mathbf{V}_\bullet X}^{\mathcal{Q}}$ .*
3. *The multiplication  $\mu_{\mu_X}^{\overline{S}}$  evaluated at the free  $\mathbf{V}_\bullet$ -algebra  $\mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$  is the function that maps every  $Q \in \mathcal{Q}_V^{cvx} \mathbb{P}_{\text{DP}}^\bullet X$  to  $(h \in \mathcal{L}X \mapsto \min_{F \in Q} F(h)) \in \mathbb{P}_{\text{DP}}^\bullet X$ ; the inverse image of  $[h > 1]$  by that function is  $\square[h > 1] \cap \mathcal{Q}_V^{cvx} \mathbb{P}_{\text{DP}}^\bullet X$ , for every  $h \in \mathcal{L}X$ .*

4. The counit  $\epsilon^T$  of the adjunction  $F^T \dashv U^T$  between **Top** and the category of  $T$ -algebras ( $\mathbf{V}_\bullet$ -algebras) is given at each algebra  $\beta: \mathbf{V}_\bullet X \rightarrow X$  as  $\beta$  itself.

PROOF. 1. We know that  $\overline{S}\mu_X$  must be a  $\mathbf{V}_\bullet$ -algebra on  $\overline{S}\mathbf{V}_\bullet X$ , and the latter can be taken as  $\mathbb{P}_{\text{DP}}^\bullet X$ , by Lemma 5.1. (All splittings are up to isomorphism.) We know that  $\overline{S}\mu_X = \pi_{\mu_X} \circ S\mu_X \circ \lambda_X \circ T\iota_{\mu_X} = r_{\text{DP}} \circ \mathcal{Q}_V \mu_X \circ \lambda_X^\# \circ \mathbf{V}_\bullet s_{\text{DP}}^\bullet$ , which we elucidate using Trick A:

$$\begin{aligned}
[h > 1] &\xrightarrow{r_{\text{DP}}} \square[h > 1] && \text{by Fact 4.5} \\
&\xrightarrow{\mathcal{Q}_V \mu_X} \square \mu_X^{-1}([h > 1]) = \square \left[ \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right) > 1 \right] && \text{by (1)} \\
&\xrightarrow{\lambda_{\mathbf{V}_\bullet X}^\#} \left[ \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)^* > 1 \right] && \text{by Corollary 4.8} \\
&\xrightarrow{\mathbf{V}_\bullet s_{\text{DP}}^\bullet} \left[ \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)^* \circ s_{\text{DP}}^\bullet > 1 \right] \\
&= [(F \in \mathbb{P}_{\text{DP}}^\bullet X \mapsto F(h)) > 1].
\end{aligned}$$

The last equality is justified as follows. For every  $F \in \mathbb{P}_{\text{DP}}^\bullet X$ ,

$$\begin{aligned}
\left( \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)^* \circ s_{\text{DP}}^\bullet \right) (F) &= \min_{\nu \in s_{\text{DP}}^\bullet(F)} \int_{x \in X} h(x) d\nu \\
&= r_{\text{DP}}(s_{\text{DP}}^\bullet(F))(h) = F(h).
\end{aligned}$$

Finally,  $[(F \in \mathbb{P}_{\text{DP}}^\bullet X \mapsto F(h)) > 1]$  is the collection of continuous valuations  $\xi \in \mathbf{V}_\bullet \mathbb{P}_{\text{DP}}^\bullet X$  such that  $\int_{F \in \mathbb{P}_{\text{DP}}^\bullet X} F(h) d\xi > 1$ , namely  $(\alpha_{\text{DP}}^\bullet)^{-1}([h > 1])$ .

2.  $\eta_{\mu_X}^{\overline{S}} = \pi_{\mu_X} \circ \eta_{\mathbf{V}_\bullet X}^{\overline{S}} = r_{\text{DP}} \circ \eta_{\mathbf{V}_\bullet X}^{\mathcal{Q}}$ .

3. In general,  $\mu_\alpha^{\overline{S}}$  is a morphism from  $\overline{S}\overline{S}\alpha$  to  $\overline{S}\alpha$  in the category of  $\mathbf{V}_\bullet$ -algebras. When  $\alpha = \mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$ , it is therefore one from  $\mathbb{P}_{\text{DP}}^\bullet \mathbb{P}_{\text{DP}}^\bullet X$  to  $\mathbb{P}_{\text{DP}}^\bullet X$ . We have  $\mu_{\mu_X}^{\overline{S}} \stackrel{\text{def}}{=} \pi_{\mu_X} \circ \mu_{\mathbf{V}_\bullet X}^{\overline{S}} \circ S\iota_{\mu_X} \circ \iota_{\overline{S}\mu_X}$ . We may take  $\iota_{\overline{S}\mu_X} = \iota_{\alpha_{\text{DP}}^\bullet}$  to be the right arrow, from  $\mathcal{Q}_V^{cvx} \mathbb{P}_{\text{DP}}^\bullet X$  to  $\mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X$  in Lemma 5.2—just subspace inclusion. Then  $S\iota_{\mu_X} = \mathcal{Q}_V s_{\text{DP}}^\bullet: \mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X \rightarrow \mathcal{Q}_V \mathcal{Q}_V \mathbf{V}_\bullet X$ , by Lemma 5.1. We compose that with  $\mu_{\mathbf{V}_\bullet X}^{\mathcal{Q}}: \mathcal{Q}_V \mathcal{Q}_V \mathbf{V}_\bullet X \rightarrow \mathcal{Q}_V \mathbf{V}_\bullet X$ , then with  $r_{\text{DP}}: \mathcal{Q}_V \mathbf{V}_\bullet X \rightarrow \mathbb{P}_{\text{DP}}^\bullet X$ . We elucidate what this composition is by relying on

Trick A:

$$\begin{aligned}
[h > 1] &\xrightarrow{r_{\text{DP}}} \square[h > 1] && \text{by Fact 4.5} \\
&\xrightarrow{\mu_{\mathbf{V}\bullet, X}^{\mathcal{Q}}} \square\square[h > 1] \\
&\xrightarrow{\mathcal{Q}_{\mathbf{V}} s_{\text{DP}}^{\bullet}} \square(s_{\text{DP}}^{\bullet})^{-1}(\square[h > 1]) \\
&= \square[h > 1] && \text{by Corollary 4.8} \\
&\xrightarrow{\iota_{\alpha_{\text{DP}}^{\bullet}}} \square[h > 1] \cap \mathcal{Q}_{\mathbf{V}}^{cvx} \mathbb{P}_{\text{DP}}^{\bullet} X.
\end{aligned}$$

Now this looks just like  $r_{\text{DP}}$ , and we imitate it by letting  $r: \mathcal{Q}_{\mathbf{V}}^{cvx} \mathbb{P}_{\text{DP}}^{\bullet} X \rightarrow \mathbb{P}_{\text{DP}}^{\bullet} X$  be defined by  $r(Q)(h) \stackrel{\text{def}}{=} \min_{F \in Q} F(h)$ . This is well-defined and lower semicontinuous, and  $r^{-1}([h > 1]) = \square[h > 1] \cap \mathcal{Q}_{\mathbf{V}}^{cvx} \mathbb{P}_{\text{DP}}^{\bullet} X$ , so  $\mu_{\mu_X}^{\bar{S}} = r$ .

4. Standard category theory, see [34, Chapter VI, Section 2, Theorem 1] for example.  $\square$

Finally, the weak composite monad of  $S$  with  $T$  is  $U^T \bar{S} F^T$ , with unit  $U^T \eta^{\bar{S}} F^T \circ \eta^T$  and multiplication  $U^T \mu^{\bar{S}} F^T \circ U^T \bar{S} \epsilon^T \bar{S} F^T$ , where  $F^T \dashv U^T$  is the adjunction between the base category (**Top** in our case) and the category of  $T$ -algebras, and  $\epsilon^T$  is its counit.

**Theorem 5.4.** *The weak composite monad associated with  $\lambda^{\sharp}: \mathbf{V}\bullet \mathcal{Q}_{\mathbf{V}} \rightarrow \mathcal{Q}_{\mathbf{V}} \mathbf{V}\bullet$  on **Top** is  $(\mathbb{P}_{\text{DP}}^{\bullet}, \eta^{\text{DP}}, \mu^{\text{DP}})$ .*

PROOF. We take  $T \stackrel{\text{def}}{=} \mathbf{V}\bullet$ ,  $S \stackrel{\text{def}}{=} \mathcal{Q}_{\mathbf{V}}$ ,  $\lambda \stackrel{\text{def}}{=} \lambda^{\sharp}$ . The functor  $U^T$  takes every  $\mathbf{V}\bullet$ -algebra  $\alpha: \mathbf{V}\bullet X \rightarrow X$  to  $X$  and every  $\mathbf{V}\bullet$ -algebra morphism  $f: (\alpha: \mathbf{V}\bullet X \rightarrow X) \rightarrow (\beta: \mathbf{V}\bullet Y \rightarrow Y)$  to the underlying continuous map  $f: X \rightarrow Y$ . The functor  $F^T$  takes every space  $X$  to  $\mu_X: \mathbf{V}\bullet \mathbf{V}\bullet X \rightarrow \mathbf{V}\bullet X$ , and every continuous map  $f: X \rightarrow Y$  to  $\mathbf{V}\bullet f$ , seen as a morphism of  $\mathbf{V}\bullet$ -algebras.

Then  $U^T \bar{S} F^T$  maps every space  $X$  to the splitting  $\mathbb{P}_{\text{DP}}^{\bullet} X$  of  $\mathcal{Q}_{\mathbf{V}} \mu_X \circ \lambda_X^{\sharp} \circ \eta_{\mathcal{Q}_{\mathbf{V}} X}$  we have obtained in Lemma 5.1. We also obtain  $\pi_{\mu_X} \stackrel{\text{def}}{=} r_{\text{DP}}$ ,  $\iota_{\mu_X} \stackrel{\text{def}}{=} s_{\text{DP}}^{\bullet}$ .

Given any continuous map  $f: X \rightarrow Y$ ,  $U^T \bar{S} F^T(f)$  is equal to (the map underlying the  $\mathbf{V}\bullet$ -algebra)  $\bar{S}(\mathbf{V}\bullet f)$ , namely  $\pi_{\mu_Y} \circ \mathcal{Q}_{\mathbf{V}} \mathbf{V}\bullet f \circ \iota_{\mu_X} = r_{\text{DP}} \circ \mathcal{Q}_{\mathbf{V}} \mathbf{V}\bullet f \circ s_{\text{DP}}^{\bullet}$ . We claim that this is equal to  $\mathbb{P}_{\text{DP}}^{\bullet}(f)$ , namely that it maps every  $F \in$

$\mathbb{P}_{\text{DP}}^\bullet X$  to  $(h \in \mathcal{L}X \mapsto F(f \circ h))$ . We use Trick A:

$$\begin{aligned}
[h > 1] &\xrightarrow{r_{\text{DP}}} \square[h > 1] && \text{by Fact 4.5} \\
&\xrightarrow{\mathcal{Q}_V \mathbf{V} \bullet f} \square(\mathbf{V} \bullet f)^{-1}([h > 1]) = \square[h \circ f > 1] \\
&\xrightarrow{s_{\text{DP}}^\bullet} [h \circ f > 1] && \text{by Corollary 4.8} \\
&= (\mathbb{P}_{\text{DP}}^\bullet(f))^{-1}([h > 1]).
\end{aligned}$$

From Lemma 5.3, item 2, the unit of  $\bar{S}$  at the algebra  $\mu_X$  is  $\eta_{\mu_X}^{\bar{S}} = r_{\text{DP}} \circ \eta_{\mathbf{V} \bullet X}^{\mathcal{Q}}$ , which is the map  $\nu \in \mathbf{V} \bullet X \mapsto r_{\text{DP}}(\uparrow \nu)$ , namely  $\nu \in \mathbf{V} \bullet X \mapsto (h \in \mathcal{L}X \mapsto \int_{x \in X} h(x) d\nu)$  from  $\mathbf{V} \bullet X$  to  $\mathbb{P}_{\text{DP}}^\bullet X$ .

The unit of the weak composite monad is  $U^T \eta^{\bar{S}} F^T \circ \eta^T$ . At object  $X$ , this maps every  $x \in X$  to  $(h \in \mathcal{L}X \mapsto \int_{x' \in X} h(x') d\delta_x)$ , namely to  $(h \in \mathcal{L}X \mapsto h(x))$ , and that is exactly the unit of the  $\mathbb{P}_{\text{DP}}^\bullet$  monad.

The multiplication is  $U^T \mu^{\bar{S}} F^T \circ U^T \bar{S} \epsilon^T \bar{S} F^T$ , and here is what this means, explicitly. At any object  $X$ ,  $\bar{S} F^T X$  is the  $T$ -algebra  $\bar{S} \mu_X : T \bar{S} T X \rightarrow \bar{S} T X$ . Then  $\epsilon^T$  evaluated at this  $T$ -algebra is the bottom arrow in the following diagram:

$$\begin{array}{ccc}
T T \bar{S} T X & \xrightarrow{T \epsilon_{\bar{S} F^T X}^T} & T \bar{S} T X \\
\mu_{\bar{S} T X} \downarrow & & \downarrow \bar{S} \mu_X \\
T \bar{S} T X & \xrightarrow{\epsilon_{\bar{S} F^T X}^T} & \bar{S} T X
\end{array}$$

from the  $T$ -algebra  $\alpha \stackrel{\text{def}}{=} \mu_{\bar{S} T X}$  that is the leftmost vertical arrow to the  $T$ -algebra  $\beta \stackrel{\text{def}}{=} \bar{S} \mu_X$  on the right. We apply  $\bar{S}$  to the morphism  $\epsilon_{\bar{S} F^T X}^T$  (i.e., we now apply  $\bar{S}$  to a morphism of  $T$ -algebras, not to a  $T$ -algebra, as we did before), and we obtain the composition:

$$\bar{S} T \bar{S} T X \xrightarrow{\iota_\alpha} \bar{S} T \bar{S} T X \xrightarrow{S \epsilon_{\bar{S} F^T X}^T} \bar{S} \bar{S} T X \xrightarrow{\pi_\beta} \bar{S} \bar{S} T X. \quad (4)$$

Applying  $U^T$  to the latter, we obtain the same composition, this time seen as a morphism in the base category, instead of as a morphism in the category of  $T$ -algebras. We finally compose it with  $U^T \mu_{F^T X}^{\bar{S}}$ , which is  $\mu_{\mu_X}^{\bar{S}} : \bar{S} \bar{S} T X \rightarrow \bar{S} T X$ .

By Lemma 5.3, item 4,  $\epsilon_{\bar{S} F^T X}^T : \alpha \rightarrow \beta$  is  $\beta$  itself, where  $\beta = \bar{S} \mu_X = \alpha_{\text{DP}}^\bullet$  (Lemma 5.3, item 1) and  $\alpha = \mu_{\bar{S} T X} = \mu_{\mathbb{P}_{\text{DP}}^\bullet X}$ : we have  $\bar{S} T X = \mathbb{P}_{\text{DP}}^\bullet X$ , using



Lemma 5.1, and  $\iota_\alpha = s_{\text{DP}}^\bullet$ . Using Lemma 5.2,  $\pi_\beta: \mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X \rightarrow \mathcal{Q}_V^{cvx} \mathbb{P}_{\text{DP}}^\bullet X$  is the corestriction of (3) (call it  $f$ ), which maps every convex element of  $\mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X$  to itself.

Hence the multiplication of the weak composite monad, evaluated at  $X$ , is the composition of (4) with  $\mu_{\mu_X}^{\bar{S}}$  (given in Lemma 5.3, item 3), namely:

$$\mathbb{P}_{\text{DP}}^\bullet \mathbb{P}_{\text{DP}}^\bullet X \xrightarrow{s_{\text{DP}}^\bullet} \mathcal{Q}_V \mathbf{V}_\bullet \mathbb{P}_{\text{DP}}^\bullet X \xrightarrow{\mathcal{Q}_V \alpha_{\text{DP}}^\bullet} \mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X \xrightarrow{f} \mathcal{Q}_V^{cvx} \mathbb{P}_{\text{DP}}^\bullet X \xrightarrow{\mu_{\mu_X}^{\bar{S}}} \mathbb{P}_{\text{DP}}^\bullet X \quad (5)$$

For every  $\mathcal{F} \in \mathbb{P}_{\text{DP}}^\bullet \mathbb{P}_{\text{DP}}^\bullet X$ ,  $\mathcal{Q}_V \alpha_{\text{DP}}^\bullet(s_{\text{DP}}^\bullet(\mathcal{F})) = \uparrow\{\alpha_{\text{DP}}^\bullet(\xi) \mid \xi \in s_{\text{DP}}^\bullet(\mathcal{F})\} = \uparrow\{(h \in \mathcal{L}X \mapsto \int_{F \in \mathbb{P}_{\text{DP}}^\bullet X} F(h) d\xi) \mid \xi \in s_{\text{DP}}^\bullet(\mathcal{F})\}$ . It is easy to see that this is a convex set, because  $s_{\text{DP}}^\bullet(\mathcal{F})$  is convex and integration is linear in the continuous valuation. Therefore  $f$  maps it to itself, and, using Lemma 5.3, item 3,  $\mu_{\mu_X}^{\bar{S}}$  maps it to:

$$\begin{aligned} (h \in \mathcal{L}X \mapsto \min_{F \in \mathcal{Q}_V \alpha_{\text{DP}}^\bullet(s_{\text{DP}}^\bullet(\mathcal{F}))} F(h)) &= (h \in \mathcal{L}X \mapsto \min_{\xi \in s_{\text{DP}}^\bullet(\mathcal{F}), F \geq \alpha_{\text{DP}}^\bullet(\xi)} F(h)) \\ &= (h \in \mathcal{L}X \mapsto \min_{\xi \in s_{\text{DP}}^\bullet(\mathcal{F})} \alpha_{\text{DP}}^\bullet(\xi)(h)) \\ &= (h \in \mathcal{L}X \mapsto \min_{\xi \in s_{\text{DP}}^\bullet(\mathcal{F})} \int_{F \in \mathbb{P}_{\text{DP}}^\bullet X} F(h) d\xi) \\ &= (h \in \mathcal{L}X \mapsto r_{\text{DP}}(s_{\text{DP}}^\bullet(\mathcal{F}))(F \in \mathbb{P}_{\text{DP}}^\bullet X \mapsto F(h))) \\ &= (h \in \mathcal{L}X \mapsto \mathcal{F}(F \in \mathbb{P}_{\text{DP}}^\bullet X \mapsto F(h))), \end{aligned}$$

and this is exactly what the multiplication of the  $\mathbb{P}_{\text{DP}}^\bullet$  monad, evaluated at  $X$  and applied to  $\mathcal{F}$ , produces.  $\square$

## Part II

# A weak distributive law between $\mathbf{V}_\bullet$ and $\mathcal{H}_V$

Just as  $\mathcal{Q}_V$  is a model of demonic non-determinism, the Hoare powerdomain is a model of *angelic* non-determinism, and is obtained as follows. For every topological space  $X$ , let  $\mathcal{H}X$  be the set of non-empty closed subsets of  $X$ . The *lower Vietoris* topology on that set has subbasic open subsets  $\diamond U$  consisting

of those non-empty closed subsets that intersect  $U$ , for each open subset  $U$  of  $X$ . We write  $\mathcal{H}_V X$  for the resulting topological space. Its specialization ordering is (ordinary) inclusion  $\subseteq$ .

The  $\mathcal{H}_V$  construction was studied by Schalk [37, Section 6], as well as a localic variant and a variant with the Scott topology. See also [1, Sections 6.2.2, 6.2.3] or [10, Section IV-8].

There is a monad  $(\mathcal{H}_V, \eta^{\mathcal{H}}, \mu^{\mathcal{H}})$  on  $\mathbf{Top}$ . For every continuous map  $f: X \rightarrow Y$ ,  $\mathcal{H}_V f$  maps every  $C \in \mathcal{Q}_V X$  to  $cl(f[C])$ . The unit  $\eta^{\mathcal{H}}$  is defined by  $\eta_X^{\mathcal{H}}(x) \stackrel{\text{def}}{=} \downarrow x$  for every  $x \in X$ . For every continuous map  $f: X \rightarrow \mathcal{Q}_V Y$ , there is an extension  $f^b: \mathcal{H}_V X \rightarrow \mathcal{H}_V Y$ , defined by  $f^b(C) \stackrel{\text{def}}{=} cl(\bigcup_{x \in C} f(x))$ . The multiplication  $\mu^{\mathcal{H}}$  is defined by  $\mu_X^{\mathcal{H}} \stackrel{\text{def}}{=} \text{id}_{\mathcal{H}_V X}^b$ , namely  $\mu_X^{\mathcal{H}}(\mathcal{C}) \stackrel{\text{def}}{=} cl(\bigcup \mathcal{C})$ .

We note the equalities  $\eta^{\mathcal{H}^{-1}}(\diamond U) = U$ ,  $(\mathcal{H}_V f)^{-1}(\diamond V) = \diamond f^{-1}(V)$ , and  $\mu^{\mathcal{H}^{-1}}(\diamond U) = \diamond \diamond U$ .

The remainder of this part is more or less directly copied from Part I, and is therefore pretty boring. For the most part, we replace  $\square$  with  $\diamond$ ,  $\supseteq$  by  $\subseteq$ ,  $\lambda_X^\#$  by  $\lambda_X^b$ , etc. But there are a few differences, if we look closely enough. For example, the sets  $\diamond U$  form a *subbase*, while the sets  $\square U$  formed a *base*. Perhaps more importantly, we will not reason on the whole category  $\mathbf{Top}$ , but on a full subcategory  $\mathbf{Top}^b$  of spaces  $X$  such that  $\mathcal{L}X$  is locally convex. (The latter means that for every  $h \in \mathcal{L}X$ , for every Scott-open neighborhood  $\mathcal{U}$  of  $h$ , there is a convex Scott-open neighborhood  $\mathcal{V}$  of  $h$  included in  $\mathcal{U}$ .) The reason is that there are maps  $r_{\text{AP}}$  and  $s_{\text{AP}}^\bullet$ , similar to the maps  $r_{\text{DP}}$  and  $s_{\text{DP}}^\bullet$ , but they only enjoy similar properties on those spaces  $X$  such that  $\mathcal{L}X$  is locally convex [14, Corollary 3.12]. Also, the formula(e) for inverse images of subbasic open sets by  $s_{\text{AP}}^\bullet$  will be trickier.

The categorical machinery requires the subcategory  $\mathbf{Top}^b$  to be closed under the  $\mathcal{H}_V$  and  $\mathbf{V}_\bullet$  functors, as well as under retracts (so that  $\mathbf{Top}^b$ , just like  $\mathbf{Top}$ , splits idempotents). For example, we may take  $\mathbf{Top}^b$  to be the full subcategory of locally compact spaces (resp. of locally compact, compact spaces if  $\bullet$  is “1”). Indeed, first, every locally compact space  $X$  is LCS-complete, hence  $\odot$ -consonant (whatever that means—we will avoid the definition), hence is such that  $\mathcal{L}X$  is locally convex [7, Figure 1, Lemma 13.6]. Second, for every locally compact space  $X$ ,  $\mathcal{H}_V X$  is locally compact, by [37, Proposition 6.11], and  $\mathbf{V}_\bullet X$  is locally compact if  $\bullet$  is nothing or “ $\leq 1$ ” (see [19, Theorem 12.2], or [18, Theorem 4.1]). When  $\bullet$  is “1”,  $\mathbf{V}_\bullet X$  is locally compact and compact if  $X$  locally compact and compact, and  $\mathcal{H}_V X$  is also locally compact, as we have seen, and compact: the proof of [37, Proposition

6.11] shows in particular that  $\diamond K \stackrel{\text{def}}{=} \{C \in \mathcal{H}_V X \mid C \cap K \neq \emptyset\}$  is compact for every compact subset  $K$  of  $X$ , and the claim follows by taking  $K$  equal to  $X$ . Finally, any retract of a locally compact (resp., compact) space is locally compact (resp., compact); this follows from the arguments developed in the proof of [27, Proposition 2.17]. In brief, we work under the following assumption.

**Definition 5.5.** *Let  $\mathbf{Top}^\bullet$  be any full subcategory of  $\mathbf{Top}$  consisting of spaces  $X$  such that  $\mathcal{L}X$  is locally convex, and closed under  $\mathcal{H}_V$ ,  $\mathbf{V}_\bullet$  and under retracts—for example the category of locally compact spaces if  $\bullet$  is nothing or “ $\leq 1$ ”, or the category of locally compact, compact spaces if  $\bullet$  is “1”.*

## 6. The monad of sublinear previsions

A *sublinear* prevision on a space  $X$  is a prevision  $F$  such that  $F(h+h') \leq F(h) + F(h')$  for all  $h, h' \in \mathcal{L}X$ . We write  $\mathbb{P}_{\text{AP}} X$  for the space of sublinear previsions on  $X$ ,  $\mathbb{P}_{\text{AP}}^{\leq 1} X$  for its subspace of subnormalized sublinear previsions, and  $\mathbb{P}_{\text{AP}}^1 X$  for its subspace of normalized sublinear previsions, and we use the synthetic notation  $\mathbb{P}_{\text{AP}}^\bullet X$  for each. The topology is generated by sets that we continue to write as  $[h > r]$ , with  $h \in \mathcal{L}X$  and  $r \in \mathbb{R}_+$ , and defined as  $\{F \in \mathbb{P}_{\text{AP}}^\bullet X \mid F(h) > r\}$ . Its specialization ordering is the pointwise ordering:  $F \leq F'$  if and only if  $F(h) \leq F'(h)$  for every  $h \in \mathcal{L}X$ .

There is a  $\mathbb{P}_{\text{AP}}^\bullet$  endofunctor on  $\mathbf{Top}$ , which is defined with the exact same formulae as for  $\mathbb{P}_{\text{DP}}^\bullet$  or  $\mathbb{P}_\bullet$ . Its action on morphisms  $f: X \rightarrow Y$  is given by  $\mathbb{P}_{\text{AP}}^\bullet f(F) = (h \in \mathcal{L}Y \mapsto F(h \circ f))$ .

As with superlinear and linear previsions, sublinear previsions form a monad on  $\mathbf{Top}$ , which is a natural counterpart of the monad of sublinear previsions on the category of deposes given in [11, Proposition 2]. The following, which is Proposition 3.1 with “DP” replaced by “AP”, is proved in exactly the same way.

**Proposition 6.1.** *Let  $\bullet$  be nothing, “ $\leq 1$ ”, or “1”. There is a monad  $(\mathbb{P}_{\text{AP}}^\bullet, \eta^{\text{AP}}, \mu^{\text{AP}})$  on  $\mathbf{Top}$ , whose unit and multiplication are defined at every topological space  $X$  by:*

- for every  $x \in X$ ,  $\eta_X^{\text{AP}}(x) \stackrel{\text{def}}{=} (h \in \mathcal{L}X \mapsto h(x))$ ,
- for every  $\mathcal{F} \in \mathbb{P}_{\text{AP}}^\bullet \mathbb{P}_{\text{AP}}^\bullet X$ ,  $\mu_X^{\text{AP}}(\mathcal{F}) \stackrel{\text{def}}{=} (h \in \mathcal{L}X \mapsto \mathcal{F}(F \in \mathbb{P}_{\text{AP}}^\bullet X \mapsto F(h)))$ .

For every space  $X$  such that  $\mathcal{L}X$  is locally convex, there is a function  $r_{\text{AP}}: \mathcal{H}_V(\mathbf{V}_\bullet X) \rightarrow \mathbb{P}_{\text{AP}}^\bullet X$ , defined by  $r_{\text{AP}}(C)(h) \stackrel{\text{def}}{=} \sup_{\nu \in C} \int_{x \in X} h(x) d\nu$ , and a map  $s_{\text{AP}}^\bullet: \mathbb{P}_{\text{AP}}^\bullet X \rightarrow \mathcal{H}_V(\mathbf{V}_\bullet X)$  defined by  $s_{\text{AP}}^\bullet(F) \stackrel{\text{def}}{=} \{\nu \in \mathbf{V}_\bullet X \mid \forall h \in \mathcal{L}X, \int_{x \in X} h(x) d\nu \leq F(h)\}$ . Both are continuous, and  $r_{\text{AP}} \circ s_{\text{AP}}^\bullet = \text{id}_{\mathbb{P}_{\text{AP}}^\bullet X}$ . This is the contents of Corollary 3.12 of [14], only replacing the space  $\mathbb{P}_{\text{P}}^\bullet X$  used there by  $\mathbf{V}_\bullet X$ , since the two are homeomorphic. Additionally,  $r_{\text{AP}}$  and  $s_{\text{AP}}^\bullet$  are natural in  $X$  on the category of spaces  $X$  such that  $\mathcal{L}X$  is locally convex, in particular on the category  $\mathbf{Top}^b$  [19, Lemma 13.1].

**Corollary 6.2.** *The monad  $(\mathbb{P}_{\text{AP}}^\bullet, \eta^{\text{AP}}, \mu^{\text{AP}})$  restricts to a monad on any subcategory  $\mathbf{Top}^b$  as given in Definition 5.5.*

PROOF. It suffices to verify that  $\mathbb{P}_{\text{AP}}^\bullet X$  is an object of  $\mathbf{Top}^b$  for every object  $X$  of  $\mathbf{Top}^b$ , and this follows from the fact that it arises as a retract of  $\mathcal{H}_V \mathbf{V}_\bullet X$  through  $r_{\text{AP}}$  and  $s_{\text{AP}}^\bullet$ .  $\square$

This retraction even cuts down to a homeomorphism between  $\mathcal{H}_V^{cvx}(\mathbf{V}_\bullet X)$  and  $\mathbb{P}_{\text{AP}}^\bullet X$  [14, Theorem 4.11], where the former denotes the subspace of  $\mathcal{H}_V(\mathbf{V}_\bullet X)$  consisting of convex non-empty closed subsets of  $\mathbf{V}_\bullet X$ .

As with  $s_{\text{DP}}^\bullet$ , we propose the following simpler proof that  $s_{\text{AP}}^\bullet$  is continuous, exhibiting the shape of inverse images of subbasic open subsets  $\diamond \bigcap_{i=1}^n [h_i > 1]$  of  $\mathcal{H}_V(\mathbf{V}_\bullet X)$  ( $n \geq 1, h_i \in \mathcal{L}X$ ).

**Lemma 6.3.** *For every  $n \geq 1$ , for every natural number  $N > n$ , let  $\frac{1}{N}\mathbb{N}$  denote the set of non-negative integer multiples of  $\frac{1}{N}$ , and  $\Delta_n^N$  be the finite set  $\{(b_1, \dots, b_n) \in \frac{1}{N}\mathbb{N} \mid 1 - \frac{n}{N} < \sum_{i=1}^n b_i \leq 1\}$ .*

*Let  $X$  be a space such that  $\mathcal{L}X$  is locally convex. For every  $n \geq 1$ , for all  $h_1, \dots, h_n \in \mathcal{L}X$ ,*

$$\begin{aligned} (s_{\text{AP}}^\bullet)^{-1}(\diamond(\bigcap_{i=1}^n [h_i > 1])) &= \bigcap_{\vec{a} \in \Delta_n} \left[ \sum_{i=1}^n a_i h_i > 1 \right] \\ &= \bigcup_{N > n} \bigcap_{\vec{b} \in \Delta_n^N} \left[ \sum_{i=1}^n b_i h_i > 1 \right]. \end{aligned}$$

The first formula produces an intersection of open sets  $[\sum_{i=1}^n a_i h_i > 1]$  over all  $\vec{a} \in \Delta_n$ . That is an infinite intersection in general, and therefore it is not immediately clear that this intersection is open. The next line is an infinite union of finite intersections of open sets, and is therefore open.

PROOF. Letting (1), (2), (3) be the three values above, in order to prove (1) = (2) = (3), we show (1)  $\subseteq$  (3)  $\subseteq$  (2)  $\subseteq$  (1).

(1)  $\subseteq$  (3). Let  $F \in (s_{\text{AP}}^\bullet)^{-1}(\diamond(\bigcap_{i=1}^n [h_i > 1]))$ . There is a  $\nu \in s_{\text{AP}}^\bullet(F)$  such that  $\int_{x \in X} h_i(x) d\nu > 1$  for every  $i \in \{1, \dots, n\}$ . We pick  $N > n$  so large that  $\frac{N}{N-n} < \min_{i=1}^n \int_{x \in X} h_i(x) d\nu$ . Then, for every  $\vec{b} \in \Delta_n^N$ ,  $F(\sum_{i=1}^n b_i h_i) \geq \int_{x \in X} \sum_{i=1}^n b_i h_i(x) d\nu = \sum_{i=1}^n b_i \int_{x \in X} h_i(x) d\nu > \sum_{i=1}^n b_i \frac{N}{N-n} > (1 - \frac{n}{N}) \frac{N}{N-n} = 1$ .

(3)  $\subseteq$  (2). Let  $F \in \mathbb{P}_{\text{AP}}^\bullet X$  be such that there is a natural number  $N > n$  such that, for every  $\vec{b} \in \Delta_n^N$ ,  $F(\sum_{i=1}^n b_i h_i) > 1$ . For every  $\vec{a} \in \Delta_n$ , we define  $\vec{b}$  by  $b_i \stackrel{\text{def}}{=} \frac{1}{N} \lfloor N a_i \rfloor$  for each  $i \in \{1, \dots, n\}$ . Then  $b_i \in \frac{1}{N} \mathbb{N}$ , and  $\sum_{i=1}^n b_i$  lies between  $\sum_{i=1}^n \frac{N a_i - 1}{N} = 1 - \frac{n}{N}$  (strictly) and  $\sum_{i=1}^n a_i = 1$ , so  $\vec{b}$  is in  $\Delta_n^N$ . Also,  $a_i \geq b_i$  for every  $i$ , so, using the monotonicity of  $F$ ,  $F(\sum_{i=1}^n a_i h_i) \geq F(\sum_{i=1}^n b_i h_i) > 1$ ; this holds for every  $\vec{a} \in \Delta_n$ .

(2)  $\subseteq$  (1). Let  $F \in \bigcap_{\vec{a} \in \Delta_n} [\sum_{i=1}^n a_i h_i > 1]$ . Since  $r_{\text{AP}} \circ s_{\text{AP}}^\bullet$  is the identity,  $F(\sum_{i=1}^n a_i h_i) = \sup_{\nu \in s_{\text{AP}}^\bullet(F)} \int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu$ —and  $F(\sum_{i=1}^n a_i h_i) > 1$ —for every  $\vec{a} \in \Delta_n$ .

Let  $f: \Delta_n \times s_{\text{AP}}^\bullet(F) \rightarrow \overline{\mathbb{R}}_+$  map  $(\vec{a}, \nu)$  to  $\int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu$ . We have just shown that  $\sup_{\nu \in s_{\text{AP}}^\bullet(F)} f(\vec{a}, \nu) > 1$  for every  $\vec{a} \in \Delta_n$ .

We equip  $\Delta_n$  with the subspace topology induced by the inclusion in  $(\overline{\mathbb{R}}_+)^n$ , where each copy of  $\overline{\mathbb{R}}_+$  has the Scott topology: this makes  $f$  lower semicontinuous in its first argument, since addition and scalar multiplication are Scott-continuous on  $\overline{\mathbb{R}}_+$ . The topology on  $\Delta_n$  has a base of open subsets  $\Delta_n \cap \prod_{i=1}^n ]b_i, \infty]$ , where each  $b_i \in \mathbb{R}_+$  and  $\sum_{i=1}^n b_i < 1$  (otherwise the intersection with  $\Delta_n$  is empty, and can therefore be disregarded). But  $\Delta_n \cap \prod_{i=1}^n ]b_i, \infty] = \Delta_n \cap \prod_{i=1}^n ]b_i, 1 + \epsilon[$ , where  $\epsilon$  is any strictly positive real number; therefore  $\Delta_n$  also has the subspace topology induced by the inclusion in  $\mathbb{R}^n$  with its usual metric topology. We then know that  $\Delta_n$  is compact (and Hausdorff). The map  $f$  preserves pairwise linear combinations (with coefficients  $a$  and  $1 - a$ ,  $a \in [0, 1]$ ) of its first arguments, and similarly with its second arguments, so by Remark 3.4 and the Minimax Theorem 3.3 of [15],  $\sup_{\nu \in s_{\text{AP}}^\bullet(F)} \inf_{\vec{a} \in \Delta_n} f(\vec{a}, \nu) = \min_{\vec{a} \in \Delta_n} \sup_{\nu \in s_{\text{AP}}^\bullet(F)} f(\vec{a}, \nu)$ .

But  $\sup_{\nu \in s_{\text{AP}}^\bullet(F)} f(\vec{a}, \nu) > 1$  for every  $\vec{a} \in \Delta_n$ , in other words  $\min_{\vec{a} \in \Delta_n} \sup_{\nu \in s_{\text{AP}}^\bullet(F)} f(\vec{a}, \nu) > 1$ , so  $\sup_{\nu \in s_{\text{AP}}^\bullet(F)} \inf_{\vec{a} \in \Delta_n} f(\vec{a}, \nu) > 1$ . Hence there is a continuous valuation  $\nu \in s_{\text{AP}}^\bullet(F)$  such that  $\inf_{\vec{a} \in \Delta_n} f(\vec{a}, \nu) > 1$ , in particular such that  $\int_{x \in X} \sum_{i=1}^n a_i h_i(x) d\nu > 1$  for every  $\vec{a} \in \Delta_n$ , in particular such that  $\int_{x \in X} h_i(x) d\nu > 1$  for every  $i \in \{1, \dots, n\}$ . Therefore  $\nu \in \bigcap_{i=1}^n [h_i > 1]$ , showing that  $s_{\text{AP}}^\bullet(F) \in \diamond \bigcap_{i=1}^n [h_i > 1]$ .  $\square$

The following is immediate.

**Fact 6.4.** *For every topological space  $X$ , for every  $h \in \mathcal{L}X$ ,  $(r_{\text{AP}})^{-1}([h > 1]) = \diamond[h > 1]$ .*

## 7. The weak distributive law

We will show that there is a weak distributive law of  $\mathcal{H}_V$  over  $\mathbf{V}_\bullet$  on  $\mathbf{Top}^b$ . This will be the collection of maps  $\lambda_X^b$  as  $s_{\text{AP}}^\bullet \circ \Psi$ , where  $\Psi$  is given below.

**Lemma 7.1.** *Let  $X$  be a topological space. For every  $h \in \mathcal{L}X$ , the function  $h_*: C \mapsto \sup_{x \in C} h(x)$  is in  $\mathcal{L}\mathcal{H}_V X$ , and for every  $r \in \mathbb{R}_+$ ,  $h_*^{-1}([r, \infty]) = \diamond h^{-1}([r, \infty])$ . There is a continuous map  $\Psi: \mathbf{V}_\bullet \mathcal{H}_V X \rightarrow \mathbb{P}_{\text{AP}}^\bullet X$  such that:*

$$\Psi(\mu)(h) \stackrel{\text{def}}{=} \int_{C \in \mathcal{H}_V X} h_*(C) d\mu$$

for every  $h \in \mathcal{L}X$ . For every subbasic open subset  $[h > r]$  of  $\mathbb{P}_{\text{AP}}^\bullet X$ ,

$$\Psi^{-1}([h > r]) = [h_* > r].$$

PROOF. For every  $r \in \mathbb{R}_+$ ,  $h_*^{-1}([r, \infty]) = \{C \in \mathcal{H}_V X \mid \sup_{x \in C} h(x) > r\} = \{C \in \mathcal{H}_V X \mid \exists x \in C, h(x) > r\} = \diamond h^{-1}([r, \infty])$ . This entails that  $h_*$  is lower semicontinuous.

In particular,  $\Psi(\mu)(h)$  is well-defined for every  $\mu \in \mathbf{V}_\bullet X$  and for every  $h \in \mathcal{L}X$ . It is clear that  $\Psi$  is monotonic in  $h$ . If  $h$  is the (pointwise) supremum of a directed family  $(h_i)_{i \in I}$  in  $\mathcal{L}X$ , then for every  $C \in \mathcal{H}_V X$ ,  $h_*(C) = \sup_{x \in C} \sup_{i \in I}^\uparrow h_i(x) = \sup_{i \in I}^\uparrow h_i^*(C)$ . Since integration with respect to the continuous valuation  $\mu$  is Scott-continuous,  $\Psi(\mu)(h) = \sup_{i \in I}^\uparrow \Psi(\mu)(h_i)$ , showing that  $\Psi(\mu)$  is Scott-continuous. For all  $a \in \mathbb{R}_+$ ,  $h, h' \in \mathcal{L}X$ , we have  $(ah)_* = ah_*$  and  $(h + h')_* \leq h_* + h'_*$ , so  $\Psi(\mu)$  is sublinear. Additionally,  $(\mathbf{1} + h)_* = \mathbf{1} + h_*$ , so  $\Psi(\mu)$  is (sub)normalized if  $\mu$  is a (sub)probability valuation. Hence  $\Psi(\mu) \in \mathbb{P}_{\text{AP}}^\bullet X$ .

Finally,  $\Psi^{-1}([h > r])$  is the collection of all continuous valuations  $\mu \in \mathbf{V}_\bullet X$  such that  $\int_{C \in \mathcal{H}_V X} h_*(C) d\mu > r$ , namely  $[h_* > r]$ . In particular,  $\Psi$  is continuous.  $\square$

**Proposition 7.2.** *For every topological space  $X$  such that  $\mathcal{L}X$  is locally convex, the continuous function  $\lambda_X^b \stackrel{\text{def}}{=} s_{\text{AP}}^\bullet \circ \Psi: \mathbf{V}_\bullet \mathcal{H}_V X \rightarrow \mathcal{H}_V \mathbf{V}_\bullet X$  maps every*

$\mu \in \mathbf{V}_\bullet \mathcal{H}_V X$  to the collection of continuous valuations  $\nu \in \mathbf{V}_\bullet X$  such that  $\nu(U) \leq \mu(\diamond U)$  for every  $U \in \mathcal{O}X$ . On subbasic open sets,

$$\begin{aligned} \lambda_X^b{}^{-1}(\diamond \bigcap_{i=1}^n [h_i > 1]) &= \bigcap_{\vec{a} \in \Delta_n} \left[ \left( \sum_{i=1}^n a_i h_i \right)_* > 1 \right] \\ &= \bigcup_{N > n} \bigcap_{\vec{b} \in \Delta_n^N} \left[ \left( \sum_{i=1}^n b_i h_i \right)_* > 1 \right]. \end{aligned}$$

PROOF. The formula for inverse images is a direct consequence of Lemma 6.3 and of Lemma 7.1.

Let  $\mu \in \mathbf{V}_\bullet \mathcal{H}_V X$ . The elements  $\nu$  of  $\lambda_X^b(\mu)$  are those  $\nu \in \mathbf{V}_\bullet X$  such that  $\int_{x \in X} h(x) d\nu \leq \int_{C \in \mathcal{H}_V X} h_*(C) d\mu$  for every  $h \in \mathcal{L}X$ . In particular, for  $h \stackrel{\text{def}}{=} \chi_U$ , where  $U \in \mathcal{O}X$ , it is easy to check that  $h_* = \chi_{\diamond U}$ , so  $\nu(U) \leq \mu(\diamond U)$ . Conversely, if  $\nu(U) \leq \mu(\diamond U)$  for every  $U \in \mathcal{O}X$ , for every  $h \in \mathcal{L}X$ ,  $\int_{x \in X} h(x) d\nu = \int_0^\infty \nu(h^{-1}([t, \infty])) dt \leq \int_0^\infty \mu(\diamond h^{-1}([t, \infty])) dt = \int_0^\infty \mu(h_*^{-1}([t, \infty])) dt = \int_{x \in X} h_*(x) d\mu$ .  $\square$

When  $n = 1$ ,  $\Delta_n = \{(1)\}$ , and  $\Delta_n^N = \{(1)\}$  for every  $N > n$ . Lemma 6.3 and Proposition 7.2 then specialize to the following.

**Corollary 7.3.** *For every space  $X$  such that  $\mathcal{L}X$  is locally convex, for every  $h \in \mathcal{L}X$ ,*

$$\begin{aligned} (s_{\text{AP}}^\bullet)^{-1}(\diamond [h > 1]) &= [h > 1] \\ \lambda_X^b{}^{-1}(\diamond [h > 1]) &= [h_* > 1]. \end{aligned}$$

**Remark 7.4.** *For every  $\mu \in \mathbf{V}_\bullet \mathcal{Q}_V X$ ,  $\lambda_X^b(\mu)$  is also the set  $\{\nu \in \mathbf{V}_\bullet X \mid \forall h \in \mathcal{L}X, \int_{x \in X} h(x) d\nu \leq \int_{C \in \mathcal{H}_V X} \sup_{x \in C} h(x) d\mu\}$ .*

*In order to illustrate what  $\lambda_X^b$  does, let us look at the special case where  $\mu \stackrel{\text{def}}{=} \sum_{i=1}^n a_i \delta_{C_i}$ , where  $n \geq 1$ ,  $C_i \stackrel{\text{def}}{=} \downarrow E_i$  and where each set  $E_i$  is non-empty and finite. For every  $h \in \mathcal{L}X$ ,  $\int_{C \in \mathcal{H}_V X} \sup_{x \in C} h(x) d\mu = \sum_{i=1}^n a_i \sup_{x_i \in C_i} h(x_i) = \sum_{i=1}^n a_i \max_{x_i \in E_i} h(x_i) = \max_{x_1 \in E_1, \dots, x_n \in E_n} \sum_{i=1}^n a_i h(x_i)$ , which is equal to  $\max_{x_1 \in E_1, \dots, x_n \in E_n} \int_{x \in X} h(x) d \sum_{i=1}^n a_i \delta_{x_i}$ . It follows that  $\lambda_X^b(\mu) = \downarrow \{\sum_{i=1}^n a_i \delta_{x_i} \mid x_i \in E_i \text{ for every } i \in \{1, \dots, n\}\}$ .*

We will use the following easily proved fact.

**Fact 7.5.** *For every topological space  $X$ , for every  $h \in \mathcal{L}X$ , for every subset  $A$  of  $X$ ,  $\sup_{x \in cl(A)} h(x) = \sup_{x \in A} h(x)$ .*

Indeed, we use the fact that an open subset intersects  $cl(A)$  if and only if it intersects  $A$ . Then, for every  $t \in \mathbb{R}_+$ ,  $\sup_{x \in cl(A)} h(x) > t$  if and only if the open set  $h^{-1}(]t, \infty])$  intersects  $cl(A)$ , if and only if it intersects  $A$ , if and only if  $\sup_{x \in A} h(x) > t$ .

**Lemma 7.6.**  $\lambda^b$  is a natural transformation on  $\mathbf{Top}^b$ .

PROOF. Since  $s_{\mathbf{AP}}^\bullet$  is natural, it suffices to show that  $\Psi$  is natural. Let  $f: X \rightarrow Y$  be any continuous map. We need to show that  $\Psi \circ \mathbf{V}\bullet\mathcal{H}_V f = \mathbb{P}_{\mathbf{AP}}^\bullet f \circ \Psi$ . Let  $\mu \in \mathbf{V}\bullet\mathcal{H}_V X$  and  $h \in \mathcal{L}Y$ . Then  $(\Psi \circ \mathbf{V}\bullet\mathcal{H}_V f)(\mu)(h) = \Psi(\mathcal{H}_V f[\mu])(h) = \int_{C \in \mathcal{H}_V Y} \sup_{y \in C} h(y) d\mathcal{H}_V f[\mu]$ . By the change of variable formula, this is equal to  $\int_{C \in \mathcal{H}_V X} \sup_{y \in \mathcal{H}_V f(C)} h(y) d\mu$ . Now  $\sup_{y \in \mathcal{H}_V f(C)} h(y) = \sup_{y \in cl(f[C])} h(y) = \sup_{y \in f[C]} h(y)$  (by Fact 7.5)  $= \sup_{x \in C} h(f(x))$ . We therefore obtain that  $(\Psi \circ \mathbf{V}\bullet\mathcal{H}_V f)(\mu)(h) = \int_{C \in \mathcal{H}_V X} \sup_{x \in C} h(f(x)) d\mu$ . But  $(\mathbb{P}_{\mathbf{AP}}^\bullet f \circ \Psi)(\mu)(h) = \mathbb{P}_{\mathbf{AP}}^\bullet f(\Psi(\mu))(h) = \Psi(\mu)(h \circ f) = \int_{C \in \mathcal{H}_V X} \sup_{x \in C} h(f(x)) d\mu$ .  $\square$

We prove the first of the weak distributivity laws.

**Lemma 7.7.** *For every space  $X$  in  $\mathbf{Top}^b$ ,  $\lambda_X^b \circ \mathbf{V}\bullet\eta_X^{\mathcal{H}} = \eta_{\mathbf{V}\bullet X}^{\mathcal{H}}$ .*

PROOF. For every  $\nu \in \mathbf{V}\bullet X$ , the elements of  $(\lambda_X^b \circ \mathbf{V}\bullet\eta_X^{\mathcal{H}})(\nu)$  are those  $\nu' \in \mathbf{V}\bullet X$  such that  $\nu'(U) \leq \mathbf{V}\bullet\eta_X^{\mathcal{H}}(\nu)(\diamond U)$  for every  $U \in \mathcal{O}X$ . Now  $\mathbf{V}\bullet\eta_X^{\mathcal{H}}(\nu)(\diamond U) = \nu((\eta_X^{\mathcal{H}})^{-1}(\diamond U)) = \nu(U)$ , so  $(\lambda_X^b \circ \mathbf{V}\bullet\eta_X^{\mathcal{H}})(\nu)$  is simply the collection of  $\nu' \in \mathbf{V}\bullet X$  such that  $\nu' \leq \nu$ , namely  $\eta_{\mathbf{V}\bullet X}^{\mathcal{H}}(\nu)$ .  $\square$

For the remaining proofs, we will use Trick A, and the following variant of Trick B.

**Trick C.** *In order to show that  $f = g$ , where  $f$  and  $g$  are continuous map from a space  $Y$  to a space of the form  $\mathcal{H}_V \mathbf{V}\bullet Z$ , where  $Z$  is an object of  $\mathbf{Top}^b$ , show that  $f(y)$  and  $g(y)$  are convex for every  $y \in Y$ , and then show that  $r_{\mathbf{AP}} \circ f = r_{\mathbf{AP}} \circ g$ .*

Indeed, we recall that  $r_{\mathbf{AP}}$  is a homeomorphism of  $\mathcal{H}_V^{cux}(\mathbf{V}\bullet Z)$  onto  $\mathbb{P}_{\mathbf{AP}}^\bullet Z$ .

We also note the following. A cone is a set with a scalar multiplication operation, by scalars from  $\mathbb{R}_+$ , and with an addition operation, satisfying the expected laws. A *semitopological cone* is a cone with a topology that makes



both scalar multiplication and addition separately continuous, where  $\mathbb{R}_+$  is given the Scott topology. For example,  $\mathcal{L}X$ ,  $\mathbf{V}X$ ,  $\mathbb{P}_{\text{DP}}X$ ,  $\mathbb{P}_{\text{AP}}X$  are semi-topological cone, and  $\mathbf{V}_\bullet X$ ,  $\mathbb{P}_{\text{DP}}^\bullet X$ ,  $\mathbb{P}_{\text{AP}}^\bullet X$  are convex subspaces of the latter three. In a semitopological cone, the closure of a convex subsets is convex [28, Lemma 4.10 (a)], and we obtain the following as an easy consequence.

**Fact 7.8.** *Given any convex subspace  $Z$  of a semitopological cone, the closure of any convex subset of  $Z$  in  $Z$  is convex.*

**Lemma 7.9.** *For every object  $X$  of  $\mathbf{Top}^b$ ,  $\lambda_X^b \circ \mathbf{V}_\bullet \mu_X^{\mathcal{H}} = \mu_{\mathbf{V}_\bullet X}^{\mathcal{H}} \circ \mathcal{H}_V \lambda_X^b \circ \lambda_{\mathcal{H}_V X}^b$ .*

The presence of the term  $\lambda_{\mathcal{H}_V X}^b$  is why we require  $\mathbf{Top}^b$  to be closed under  $\mathcal{H}_V$ , see Definition 5.5.

PROOF. Since  $\lambda_X^b = s_{\text{AP}}^\bullet \circ \Psi$  and the image of  $s_{\text{AP}}^\bullet$  consists of convex subsets, the left-hand side takes its values in convex sets. For the right-hand side, for every  $\xi \in \mathbf{V}_\bullet \mathcal{H}_V \mathcal{H}_V X$ ,  $(\mu_{\mathbf{V}_\bullet X}^{\mathcal{Q}} \circ \mathcal{H}_V \lambda_X^b \circ \lambda_{\mathcal{H}_V X}^b)(\xi)$  is the closure of  $\bigcup (\mathcal{H}_V \lambda_X^b \circ \lambda_{\mathcal{H}_V X}^b)(\xi) = \bigcup \text{cl}(\{\lambda_X^b(\mu) \mid \mu \in \lambda_{\mathcal{H}_V X}^b(\xi)\})$ , hence also of  $\bigcup \{\lambda_X^b(\mu) \mid \mu \in \lambda_{\mathcal{H}_V X}^b(\xi)\}$ . In order to show that  $(\mu_{\mathbf{V}_\bullet X}^{\mathcal{Q}} \circ \mathcal{H}_V \lambda_X^b \circ \lambda_{\mathcal{H}_V X}^b)(\xi)$  is convex, it suffices to show that  $A \stackrel{\text{def}}{=} \bigcup \{\lambda_X^b(\mu) \mid \mu \in \lambda_{\mathcal{H}_V X}^b(\xi)\}$  is, by Fact 7.8.

Let  $a \in [0, 1]$ , and  $\nu_1, \nu_2 \in A$ . there is a  $\mu_1 \in \lambda_{\mathcal{H}_V X}^b(\xi)$  such that  $\nu_1 \in \lambda_X^b(\mu_1)$  and there is a  $\mu_2 \in \lambda_{\mathcal{H}_V X}^b(\xi)$  such that  $\nu_2 \in \lambda_X^b(\mu_2)$ . We note that  $\lambda_{\mathcal{H}_V X}^b(\xi) = (s_{\text{AP}}^\bullet \circ \Psi)(\xi)$  is convex, so  $a\mu_1 + (1-a)\mu_2 \in \lambda_{\mathcal{H}_V X}^b(\xi)$ . Additionally,  $a\nu_1 + (1-a)\nu_2 \in \lambda_X^b(a\mu_1 + (1-a)\mu_2)$ : for every  $U \in \mathcal{O}X$ ,  $\nu_1(U) \leq \mu_1(\diamond U)$  and  $\nu_2(U) \leq \mu_2(\diamond U)$  by definition of  $\lambda_X^b$ , so  $(a\nu_1 + (1-a)\nu_2)(U) \leq (a\mu_1 + (1-a)\mu_2)(\diamond U)$ . Therefore there is a  $\mu \in \lambda_{\mathcal{H}_V X}^b(\xi)$ , namely  $\mu \stackrel{\text{def}}{=} a\mu_1 + (1-a)\mu_2$ , such that  $a\nu_1 + (1-a)\nu_2 \in \lambda_X^b(\mu)$ . Equivalently,  $a\nu_1 + (1-a)\nu_2$  is in  $A$ . Hence  $A$  is convex, as promised.

It follows that we can use Trick C, and so we prove the following instead:

$$\Psi \circ \mathbf{V}_\bullet \mu_X^{\mathcal{H}} = r_{\text{AP}} \circ \mu_{\mathbf{V}_\bullet X}^{\mathcal{H}} \circ \mathcal{H}_V \lambda_X^b \circ \lambda_{\mathcal{H}_V X}^b.$$

Indeed,  $r_{\text{AP}} \circ \lambda_X^b = r_{\text{AP}} \circ s_{\text{AP}}^\bullet \circ \Psi = \Psi$ , which allows us to simplify the left-hand side.

We now use Trick A:

$$\begin{aligned} [h > 1] &\xrightarrow{\Psi} [h_* > 1] && \text{by Lemma 7.1} \\ &\xrightarrow{\mathbf{V}_\bullet \mu_X^{\mathcal{H}}} [h_* \circ \mu_X^{\mathcal{H}} > 1] = [h_{**} > 1], \end{aligned}$$

where for every  $\mathcal{C} \in \mathcal{H}_V \mathcal{H}_V X$ ,  $h_{**}(\mathcal{C}) = \sup_{C \in \mathcal{C}} h_*(C) = \sup_{C \in \mathcal{C}} \sup_{x \in C} h(x)$ ; indeed, the latter is equal to  $\sup_{x \in \bigcup \mathcal{C}} h(x) = \sup_{x \in \text{cl}(\bigcup \mathcal{C})} h(x)$  (by Fact 7.5)  $= h_*(\mu_X^{\mathcal{H}}(\mathcal{C}))$ .

$$\begin{aligned}
[h > 1] &\xrightarrow{r_{\text{AP}}} \diamond [h > 1] && \text{by Fact 6.4} \\
&\xrightarrow{\mu_{\mathbf{V}\bullet X}^{\mathcal{H}}} \diamond \diamond [h > 1] \\
&\xrightarrow{\mathcal{H}_V \lambda_X^b} \diamond \lambda_X^b{}^{-1}(\diamond [h > 1]) = \diamond [h_* > 1] && \text{by Corollary 7.3} \\
&\xrightarrow{\lambda_{\mathcal{H}_V X}^b} [h_{**} > 1] && \text{by Corollary 7.3.}
\end{aligned}$$

□

**Lemma 7.10.** *For every object  $X$  of  $\mathbf{Top}^b$ ,  $\lambda_X^b \circ \mu_{\mathcal{H}_V X} = \mathcal{H}_V \mu_X \circ \lambda_{\mathbf{V}\bullet X}^b \circ \mathbf{V}\bullet \lambda_X^b$ .*

The term  $\lambda_{\mathbf{V}\bullet X}^b$  is why we require  $\mathbf{Top}^b$  to be closed under  $\mathbf{V}\bullet$  (see Definition 5.5).

PROOF. The left-hand side takes its values in convex sets. We claim that the right-hand side does, too. For every  $\xi \in \mathbf{V}\bullet \mathbf{V}\bullet \mathcal{H}_V X$ ,  $(\mathcal{H}_V \mu_X \circ \lambda_{\mathbf{V}\bullet X}^b \circ \mathbf{V}\bullet \lambda_X^b)(\xi)$  is the closure of  $A \stackrel{\text{def}}{=} \bigcup \{\mu_X(\mu) \mid \mu \in \lambda_{\mathbf{V}\bullet X}^b(\lambda_X^b[\xi])\}$  so it suffices to show that  $A$  is convex, and to rely on Fact 7.8. For all  $a \in [0, 1]$ ,  $\mu_1, \mu_2 \in \lambda_{\mathbf{V}\bullet X}^b(\lambda_X^b[\xi])$ ,  $\mu \stackrel{\text{def}}{=} a\mu_1 + (1-a)\mu_2$  is in  $\lambda_{\mathbf{V}\bullet X}^b(\lambda_X^b[\xi])$ , since  $\lambda_{\mathbf{V}\bullet X}^b$  takes its values in convex sets. Then  $\mu_X(\mu) = (\mathcal{U} \in \mathcal{O}\mathbf{V}\bullet X \mapsto \int_{\nu \in \mathbf{V}\bullet X} \nu(\mathcal{U}) d\mu) = a\mu_X(\mu_1) + (1-a)\mu_X(\mu_2)$  is in  $A$ , showing that  $A$  is convex.

We can therefore use Trick C, and since  $r_{\text{AP}} \circ \lambda_X^b = r_{\text{AP}} \circ s_{\text{AP}}^\bullet \circ \Psi = \Psi$ , it remains to show:

$$\Psi \circ \mu_{\mathcal{H}_V X} = r_{\text{AP}} \circ \mathcal{H}_V \mu_X \circ \lambda_{\mathbf{V}\bullet X}^b \circ \mathbf{V}\bullet \lambda_X^b.$$

To this end, we use Trick A:

$$\begin{aligned}
[h > 1] &\xrightarrow{\Psi} [h_* > 1] && \text{by Lemma 7.1} \\
&\xrightarrow{\mu_{\mathcal{H}_V X}} \left[ \left( \mu \in \mathbf{V}\bullet \mathcal{H}_V X \mapsto \int_{C \in \mathcal{H}_V X} h_*(C) d\mu \right) > 1 \right] && \text{by (1)}
\end{aligned}$$

while:

$$\begin{aligned}
[h > 1] &\xrightarrow{r_{\text{AP}}} \diamond [h > 1] && \text{by Fact 6.4} \\
&\xrightarrow{\mathcal{H}_V \mu_X} \diamond \mu_X^{-1}([h > 1]) = \diamond \left[ \left( \nu \in \mathbf{V} \bullet X \mapsto \int_{x \in X} h(x) d\nu \right) > 1 \right] && \text{by (1)} \\
&\xrightarrow{\lambda_{\mathbf{V} \bullet X}^b} \left[ \left( \nu \in \mathbf{V} \bullet X \mapsto \int_{x \in X} h(x) d\nu \right)_* > 1 \right] && \text{by Corollary 7.3} \\
&\xrightarrow{\mathbf{V} \bullet \lambda_X^b} \left[ \left( \nu \in \mathbf{V} \bullet X \mapsto \int_{x \in X} h(x) d\nu \right)_* \circ \lambda_X^b > 1 \right].
\end{aligned}$$

In order to conclude, it suffices to show that:

$$\left( \nu \in \mathbf{V} \bullet X \mapsto \int_{x \in X} h(x) d\nu \right)_* \circ \lambda_X^b = (\mu \in \mathbf{V} \bullet \mathcal{H}_V X \mapsto \int_{C \in \mathcal{H}_V X} h_*(C) d\mu).$$

The left-hand side maps every  $\mu \in \mathbf{V} \bullet \mathcal{H}_V X$  to

$$\begin{aligned}
\sup_{\nu \in s_{\text{AP}}^\bullet(\Psi(\mu))} \int_{x \in X} h(x) d\nu &= r_{\text{AP}}(s_{\text{AP}}^\bullet(\Psi(\mu)))(h) \\
&= \Psi(\mu)(h) = \int_{C \in \mathcal{H}_V X} h_*(C),
\end{aligned}$$

and this completes the proof.  $\square$

Putting Lemma 7.6, Lemma 7.7, Lemma 7.9 and Lemma 7.10 together, we obtain the following.

**Theorem 7.11.** *Let  $\bullet$  be nothing, “ $\leq 1$ ”, or “1”. Then  $\lambda^b$  is a weak distributive law of  $\mathcal{H}_V$  over  $\mathbf{V} \bullet$  on any category  $\mathbf{Top}^b$  as given in Definition 5.5.*

## 8. The associated weak composite monad

We proceed as in Section 5. We consider the following idempotent instead of (2):

$$\mathcal{H}_V \mathbf{V} \bullet X \xrightarrow{\eta_{\mathcal{H}_V \mathbf{V} \bullet X}} \mathbf{V} \bullet \mathcal{H}_V \mathbf{V} \bullet X \xrightarrow{\lambda_{\mathbf{V} \bullet X}^b} \mathcal{H}_V \mathbf{V} \bullet \mathbf{V} \bullet X \xrightarrow{\mathcal{H}_V \mu_X} \mathcal{H}_V \mathbf{V} \bullet X \quad (6)$$

**Lemma 8.1.** *For every object  $X$  of  $\mathbf{Top}^b$ , a splitting of the idempotent (6) is  $\mathcal{H}_V \mathbf{V} \bullet X \xrightarrow{r_{\text{AP}}} \mathbb{P}_{\text{AP}}^\bullet X \xrightarrow{s_{\text{AP}}^\bullet} \mathcal{H}_V \mathbf{V} \bullet X$ .*

We note that the middle object  $\mathbb{P}_{\text{AP}}^\bullet X$  occurs as a retract of  $\mathcal{H}_V \mathbf{V}_\bullet X$  in  $\mathbf{Top}$ , through  $r_{\text{AP}}$ . Hence it is in  $\mathbf{Top}^b$ , since  $\mathbf{Top}^b$  is closed under  $\mathcal{H}_V$ ,  $\mathbf{V}_\bullet$  and retracts (Definition 5.5).

PROOF. Let  $f$  denote the idempotent (6). We recall that  $\lambda_{\mathbf{V}_\bullet X}^b = s_{\text{AP}}^\bullet \circ \Psi$ . (And we remember that  $\mathbf{V}_\bullet X$  is in  $\mathbf{Top}^b$ , since  $\mathbf{Top}^b$  is closed under  $\mathbf{V}_\bullet$ .) In particular, any element in its range is convex. The element of any convex element  $C$  of a space  $\mathcal{H}_V \mathbf{V}_\bullet$  by  $\mathcal{H}_V f$ , where  $f$  is any linear continuous map from  $\mathbf{V}_\bullet Y \rightarrow \mathbf{V}_\bullet Z$ , is convex; and  $\mu_X$  is linear. Therefore  $f$  takes its values in  $\mathcal{H}_V^{cvx} \mathbf{V}_\bullet X$ .

Using Trick C, and simplifying using the fact that  $r_{\text{AP}} \circ s_{\text{AP}}^\bullet$  is the identity, it remains to show that  $r_{\text{AP}}$  is equal to  $r_{\text{AP}} \circ f$ , namely to:

$$\mathcal{H}_V \mathbf{V}_\bullet X \xrightarrow{\eta_{\mathcal{H}_V \mathbf{V}_\bullet X}} \mathbf{V}_\bullet \mathcal{H}_V \mathbf{V}_\bullet X \xrightarrow{\lambda_{\mathbf{V}_\bullet X}^b} \mathcal{H}_V \mathbf{V}_\bullet \mathbf{V}_\bullet X \xrightarrow{\mathcal{H}_V \mu_X} \mathcal{H}_V \mathbf{V}_\bullet X \xrightarrow{r_{\text{AP}}} \mathbb{P}_{\text{AP}}^\bullet X$$

Using Trick A,

$$\begin{aligned} [h > 1] &\xrightarrow{r_{\text{AP}}} \diamond [h > 1] && \text{by Fact 6.4} \\ &\xrightarrow{\mathcal{H}_V \mu_X} \diamond \mu_X^{-1}([h > 1]) \\ &= \diamond \left[ \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right) > 1 \right] && \text{by (1)} \\ &\xrightarrow{\lambda_{\mathbf{V}_\bullet X}^b} \diamond \left[ \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)_* > 1 \right] && \text{by Corollary 7.3} \\ &\xrightarrow{\eta_{\mathcal{H}_V \mathbf{V}_\bullet X}} \left\{ C \in \mathcal{H}_V \mathbf{V}_\bullet X \mid \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)_* (C) > 1 \right\} \\ &= \{ C \in \mathcal{H}_V \mathbf{V}_\bullet X \mid r_{\text{AP}}(C)(h) > 1 \} = r_{\text{AP}}^{-1}([h > 1]). \end{aligned}$$

□

Generalizing (6), there is an idempotent:

$$\mathcal{H}_V X \xrightarrow{\eta_{\mathcal{H}_V X}} \mathbf{V}_\bullet \mathcal{H}_V X \xrightarrow{\lambda_X^b} \mathcal{H}_V \mathbf{V}_\bullet X \xrightarrow{\mathcal{H}_V \alpha} \mathcal{H}_V X \quad (7)$$

for every  $\mathbf{V}_\bullet$ -algebra  $\alpha$ . Finding out what its splitting is is harder in general, especially because we have no complete characterization of  $\mathbf{V}_\bullet$ -algebras in  $\mathbf{Top}$ . As before, we will only need to compute this splitting for one additional  $\mathbf{V}_\bullet$ -algebra.

**Lemma 8.2.** *For every topological space  $X$ , there is a  $\mathbf{V}_\bullet$ -algebra  $\alpha_{\mathbf{AP}}^\bullet: \mathbf{V}_\bullet \mathbb{P}_{\mathbf{AP}}^\bullet X \rightarrow \mathbb{P}_{\mathbf{AP}}^\bullet X$  defined by  $\alpha_{\mathbf{AP}}^\bullet(\xi)(h) \stackrel{\text{def}}{=} \int_{F \in \mathbb{P}_{\mathbf{AP}}^\bullet X} F(h) d\xi$  for all  $\xi \in \mathbf{V}_\bullet \mathbb{P}_{\mathbf{AP}}^\bullet X$  and  $h \in \mathcal{L}X$ .*

*If  $X$  is an object of  $\mathbf{Top}^b$ , then a splitting of the idempotent (7) where  $\alpha \stackrel{\text{def}}{=} \alpha_{\mathbf{AP}}^\bullet$  is*

$$\mathcal{H}_V \mathbb{P}_{\mathbf{AP}}^\bullet X \longrightarrow \mathcal{H}_V^{cvx} \mathbb{P}_{\mathbf{AP}}^\bullet X \longrightarrow \mathcal{H}_V \mathbb{P}_{\mathbf{AP}}^\bullet X ,$$

*where the arrow on the right is the inclusion map, and the arrow on the left is the corestriction of (7), and maps every  $C \in \mathcal{H}_V^{cvx} \mathbb{P}_{\mathbf{AP}}^\bullet X$  to itself.*

In the discussion after Lemma 8.1, we have noted that  $\mathbb{P}_{\mathbf{AP}}^\bullet X$  is an object of  $\mathbf{Top}^b$ , for every object  $X$  of  $\mathbf{Top}^b$ . Since  $\mathbf{Top}^b$  is closed under  $\mathcal{H}_V$  and under retracts (see Definition 5.5), the middle object  $\mathcal{H}_V^{cvx} \mathbb{P}_{\mathbf{AP}}^\bullet X$  is in  $\mathbf{Top}^b$ , too.

PROOF. The map  $F \mapsto F(h)$  is lower semicontinuous, since the inverse image of  $]t, \infty]$  is  $[h > t]$ , for all  $h \in \mathcal{L}X$  and  $t \in \mathbb{R}_+$ . Hence the integral  $\int_{F \in \mathbb{P}_{\mathbf{AP}}^\bullet X} F(h) d\xi$  makes sense. The map  $\alpha_{\mathbf{AP}}^\bullet(\xi)$  is Scott-continuous because each  $F \in \mathbb{P}_{\mathbf{AP}}^\bullet X$  is and because integration with respect to  $\nu$  is Scott-continuous; it is sublinear because  $F$  is and integration with respect to  $\xi$  is linear. Additionally,  $(\alpha_{\mathbf{AP}}^\bullet)^{-1}([h > r]) = [(F \mapsto F(h)) > r]$ , so  $\alpha_{\mathbf{AP}}^\bullet$  is continuous.

We claim that the map  $\alpha_{\mathbf{AP}}^\bullet$  is a  $\mathbf{V}_\bullet$ -algebra. We have  $\alpha_{\mathbf{AP}}^\bullet(\eta_{\mathbb{P}_{\mathbf{AP}}^\bullet X}(F))(h) = \int_{F' \in \mathbb{P}_{\mathbf{AP}}^\bullet X} F'(h) d\delta_F = F(h)$ , so  $\alpha_{\mathbf{AP}}^\bullet \circ \eta_{\mathbb{P}_{\mathbf{AP}}^\bullet X}$  is the identity map. In order to show that  $\alpha_{\mathbf{AP}}^\bullet \circ \mu_{\mathbb{P}_{\mathbf{AP}}^\bullet X} = \alpha_{\mathbf{AP}}^\bullet \circ \mathbf{V}_\bullet \alpha_{\mathbf{AP}}^\bullet$ , we use Trick A:

$$\begin{aligned} [h > 1] &\stackrel{\alpha_{\mathbf{AP}}^\bullet}{\implies} [(F \in \mathbb{P}_{\mathbf{AP}}^\bullet X \mapsto F(h)) > 1] \\ &\stackrel{\mu_{\mathbb{P}_{\mathbf{AP}}^\bullet X}}{\implies} [(\xi \in \mathbf{V}_\bullet \mathbb{P}_{\mathbf{AP}}^\bullet X \mapsto \int_{F \in \mathbb{P}_{\mathbf{AP}}^\bullet X} F(h) d\xi) > 1] \quad \text{by (1)} \\ [h > 1] &\stackrel{\alpha_{\mathbf{AP}}^\bullet}{\implies} [(F \in \mathbb{P}_{\mathbf{AP}}^\bullet X \mapsto F(h)) > 1] \\ &\stackrel{\mathbf{V}_\bullet \alpha_{\mathbf{AP}}^\bullet}{\implies} [(F \in \mathbb{P}_{\mathbf{AP}}^\bullet X \mapsto F(h)) \circ \alpha_{\mathbf{AP}}^\bullet > 1] \\ &= [(\xi \in \mathbf{V}_\bullet \mathbb{P}_{\mathbf{AP}}^\bullet X \mapsto \alpha_{\mathbf{AP}}^\bullet(\xi)(h)) > 1] \\ &= [(\xi \in \mathbf{V}_\bullet \mathbb{P}_{\mathbf{AP}}^\bullet X \mapsto \int_{F \in \mathbb{P}_{\mathbf{AP}}^\bullet X} F(h) d\xi) > 1]. \end{aligned}$$

In order to justify the announced splitting of (7), we need to show that the image of (7) is exactly  $\mathcal{H}_V^{cvx} \mathbb{P}_{\mathbf{AP}}^\bullet X$ . For every  $C \in \mathcal{H}_V X$ , the image of  $C$  by

(7) is  $cl(\alpha_{\mathbb{A}\mathbb{P}}^\bullet[C'])$  where  $C' \stackrel{\text{def}}{=} \lambda_{\mathbb{A}\mathbb{P}, X}^\flat(\delta_C)$  is convex, and  $\alpha_{\mathbb{A}\mathbb{P}}^\bullet$  is linear (in its  $\xi$  argument), so that image is convex, using Fact 7.8.

In order to see that every element  $C$  of  $\mathcal{H}_V^{cvx}\mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X$  is reached as the image of some element by (7), we simply show that  $C$  is its own image by (7), or equivalently, that (7) restricts to the identity on  $\mathcal{H}_V^{cvx}\mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X$ . We use Trick A:

$$\begin{aligned}
& \diamond \bigcap_{i=1}^n [h_i > 1] \xrightarrow{\mathcal{H}_V \alpha_{\mathbb{A}\mathbb{P}}^\bullet} \diamond \bigcap_{i=1}^n (\alpha_{\mathbb{A}\mathbb{P}}^\bullet)^{-1}([h_i > 1]) \\
& = \diamond \bigcap_{i=1}^n [(F \mapsto F(h_i)) > 1] \\
& \xrightarrow{\lambda_{\mathbb{A}\mathbb{P}, X}^\flat} \bigcap_{\vec{a} \in \Delta_n} \left[ \left( F \mapsto \sum_{i=1}^n a_i F(h_i) \right)_* > 1 \right] \quad \text{by Proposition 7.2} \\
& \xrightarrow{\eta_{\mathcal{H}_V \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X}} \bigcap_{\vec{a} \in \Delta_n} \left( \left( F \mapsto \sum_{i=1}^n a_i F(h_i) \right)_* \right)^{-1} ([1, \infty]),
\end{aligned}$$

and we claim that the convex elements  $C$  of  $\mathcal{H}_V \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X$  that are in the latter open set are exactly those of  $\diamond \bigcup_{i=1}^n [h_i > 1]$ , which will show the claim.

For every  $C \in \mathcal{H}_V \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X$ ,  $C \in \bigcap_{\vec{a} \in \Delta_n} ((F \mapsto \sum_{i=1}^n a_i F(h_i))_*)^{-1}([1, \infty])$  if and only for every  $\vec{a} \in \Delta_n$ ,  $\sup_{F \in C} \sum_{i=1}^n a_i F(h_i) > 1$ . Let  $f: \Delta_n \times C \rightarrow \overline{\mathbb{R}}_+$  map  $(\vec{a}, F)$  to  $\int_{x \in X} \sum_{i=1}^n a_i F(h_i)$ . This is a lower semicontinuous map in its first argument,  $\Delta_n$  is compact (and Hausdorff), as in the proof of Lemma 6.3, and  $f$  preserves pairwise linear combinations (with coefficients  $a$  and  $1-a$ ,  $a \in [0, 1]$ ) of its first arguments, and similarly with its second arguments (which exist because  $C$  is convex), so by Remark 3.4 and the Minimax Theorem 3.3 of [15],  $\sup_{F \in C} \inf_{\vec{a} \in \Delta_n} f(\vec{a}, F) = \min_{\vec{a} \in \Delta_n} \sup_{F \in C} f(\vec{a}, F)$ . But the elements  $C \in \mathcal{H}_V \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X$  that are in  $\bigcap_{\vec{a} \in \Delta_n} ((F \mapsto \sum_{i=1}^n a_i F(h_i))_*)^{-1}([1, \infty])$  are those such that  $\min_{\vec{a} \in \Delta_n} \sup_{F \in C} f(\vec{a}, F) > 1$ , hence such that  $\sup_{F \in C} \inf_{\vec{a} \in \Delta_n} f(\vec{a}, F) > 1$ , hence such that for some  $F \in C$ ,  $\inf_{\vec{a} \in \Delta_n} f(\vec{a}, F) > 1$ . Since  $f(\vec{a}, F)$  is linear in  $\vec{a}$  (and lower semicontinuous on a compact set), the latter inf is reached at one of the vertices of  $\Delta_n$ , so  $\inf_{\vec{a} \in \Delta_n} f(\vec{a}, F) > 1$  is equivalent to  $F(h_i) > 1$  for every  $i \in \{1, \dots, n\}$ . Therefore, for every  $C \in \mathcal{H}_V \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X$ ,  $C \in \bigcap_{\vec{a} \in \Delta_n} ((F \mapsto \sum_{i=1}^n a_i F(h_i))_*)^{-1}([1, \infty])$  if and only if there is an  $F \in C$  such that  $F(h_i) > 1$  for every  $i \in \{1, \dots, n\}$ , if and only if  $C \in \diamond \bigcap_{i=1}^n [h_i > 1]$ .  $\square$

**Lemma 8.3.** For  $S \stackrel{\text{def}}{=} \mathcal{H}_V$ ,  $T \stackrel{\text{def}}{=} \mathbf{V}_\bullet$ ,  $\lambda \stackrel{\text{def}}{=} \lambda^\flat$ , the monad  $\overline{S}$  on the category of  $\mathbf{V}_\bullet$ -algebras over  $\mathbf{Top}^\flat$  has the following properties:

1. For every free  $\mathbf{V}_\bullet$ -algebra  $\mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$ ,  $\bar{S}\mu_X: \mathbf{V}_\bullet \mathbb{P}_{\mathbf{AP}}^\bullet X \rightarrow \mathbb{P}_{\mathbf{AP}}^\bullet X$  is  $\alpha_{\mathbf{AP}}^\bullet$  (see Lemma 8.2).
2. The unit  $\eta_{\mu_X}^{\bar{S}}$  evaluated at the free  $\mathbf{V}_\bullet$ -algebra  $\mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$  is  $r_{\mathbf{AP}} \circ \eta_{\mathbf{V}_\bullet X}^{\mathcal{H}}$ .
3. The multiplication  $\mu_{\mu_X}^{\bar{S}}$  evaluated at the free  $\mathbf{V}_\bullet$ -algebra  $\mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$  is the function that maps every  $C \in \mathcal{H}_{\mathbf{V}}^{cvx} \mathbb{P}_{\mathbf{AP}}^\bullet X$  to  $(h \in \mathcal{L}X \mapsto \sup_{F \in C} F(h)) \in \mathbb{P}_{\mathbf{AP}}^\bullet X$ ; the inverse image of  $[h > 1]$  by that function is  $\diamond[h > 1] \cap \mathcal{H}_{\mathbf{V}}^{cvx} \mathbb{P}_{\mathbf{AP}}^\bullet X$ , for every  $h \in \mathcal{L}X$ .
4. The counit  $\epsilon^T$  of the adjunction  $F^T \dashv U^T$  between **Top** and the category of  $T$ -algebras ( $\mathbf{V}_\bullet$ -algebras) is given at each algebra  $\beta: \mathbf{V}_\bullet X \rightarrow X$  as  $\beta$  itself.

PROOF. 1. We know that  $\bar{S}\mu_X$  must be a  $\mathbf{V}_\bullet$ -algebra on  $\bar{S}\mathbf{V}_\bullet X$ , and the latter can be taken as  $\mathbb{P}_{\mathbf{AP}}^\bullet X$ , by Lemma 5.1. We know that  $\bar{S}\mu_X = \pi_{\mu_X} \circ S\mu_X \circ \lambda_X \circ T\iota_{\mu_X} = r_{\mathbf{AP}} \circ \mathcal{H}_{\mathbf{V}} \mu_X \circ \lambda_X^b \circ \mathbf{V}_\bullet s_{\mathbf{AP}}^\bullet$ , which we elucidate using Trick A:

$$\begin{aligned}
[h > 1] &\xrightarrow{r_{\mathbf{AP}}} \diamond[h > 1] && \text{by Fact 6.4} \\
&\xrightarrow{\mathcal{H}_{\mathbf{V}} \mu_X} \diamond \mu_X^{-1}([h > 1]) = \diamond \left[ \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right) > 1 \right] && \text{by (1)} \\
&\xrightarrow{\lambda_{\mathbf{V}_\bullet X}^b} \left[ \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)_* > 1 \right] && \text{by Corollary 7.3} \\
&\xrightarrow{\mathbf{V}_\bullet s_{\mathbf{AP}}^\bullet} \left[ \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)_* \circ s_{\mathbf{AP}}^\bullet > 1 \right] \\
&= [(F \in \mathbb{P}_{\mathbf{AP}}^\bullet X \mapsto F(h)) > 1].
\end{aligned}$$

The last equality is justified as follows. For every  $F \in \mathbb{P}_{\mathbf{AP}}^\bullet X$ ,

$$\begin{aligned}
\left( \left( \nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h(x) d\nu \right)_* \circ s_{\mathbf{AP}}^\bullet \right) (F) &= \sup_{\nu \in s_{\mathbf{AP}}^\bullet(F)} \int_{x \in X} h(x) d\nu \\
&= r_{\mathbf{AP}}(s_{\mathbf{AP}}^\bullet(F))(h) = F(h).
\end{aligned}$$

Finally,  $[(F \in \mathbb{P}_{\mathbf{AP}}^\bullet X \mapsto F(h)) > 1]$  is the collection of continuous valuations  $\xi \in \mathbf{V}_\bullet \mathbb{P}_{\mathbf{AP}}^\bullet X$  such that  $\int_{F \in \mathbb{P}_{\mathbf{AP}}^\bullet X} F(h) d\xi > 1$ , namely  $(\alpha_{\mathbf{AP}}^\bullet)^{-1}([h > 1])$ .

$$2. \eta_{\mu_X}^{\bar{S}} = \pi_{\mu_X} \circ \eta_{\mathbf{V}_\bullet X}^S = r_{\mathbf{AP}} \circ \eta_{\mathbf{V}_\bullet X}^{\mathcal{H}}.$$

3. In general,  $\mu_\alpha^{\bar{S}}$  is a morphism from  $\overline{S\bar{S}\alpha}$  to  $\bar{S}\alpha$  in the category of  $\mathbf{V}_\bullet$ -algebras. When  $\alpha = \mu_X: \mathbf{V}_\bullet\mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$ , it is therefore one from  $\mathbb{P}_{\text{AP}}^\bullet\mathbb{P}_{\text{AP}}^\bullet X$  to  $\mathbb{P}_{\text{AP}}^\bullet X$ . We have  $\mu_{\mu_X}^{\bar{S}} \stackrel{\text{def}}{=} \pi_{\mu_X} \circ \mu_{\mathbf{V}_\bullet X}^{\bar{S}} \circ S\iota_{\mu_X} \circ \iota_{\bar{S}\mu_X}$ . We may take  $\iota_{\bar{S}\mu_X} = \iota_{\alpha_{\text{AP}}^\bullet}$  to be the right arrow, from  $\mathcal{H}_{\mathbf{V}}^{cvx}\mathbb{P}_{\text{AP}}^\bullet X$  to  $\mathcal{H}_{\mathbf{V}}\mathbb{P}_{\text{AP}}^\bullet X$  in Lemma 8.2—just subspace inclusion. Then  $S\iota_{\mu_X} = \mathcal{H}_{\mathbf{V}}s_{\text{AP}}^\bullet: \mathcal{H}_{\mathbf{V}}\mathbb{P}_{\text{AP}}^\bullet X \rightarrow \mathcal{H}_{\mathbf{V}}\mathcal{H}_{\mathbf{V}}\mathbf{V}_\bullet X$ , by Lemma 5.1. We compose that with  $\mu_{\mathbf{V}_\bullet X}^{\mathcal{H}}: \mathcal{H}_{\mathbf{V}}\mathcal{H}_{\mathbf{V}}\mathbf{V}_\bullet X \rightarrow \mathcal{H}_{\mathbf{V}}\mathbf{V}_\bullet X$ , then with  $r_{\text{AP}}: \mathcal{H}_{\mathbf{V}}\mathbf{V}_\bullet X \rightarrow \mathbb{P}_{\text{AP}}^\bullet X$ . We elucidate what this composition is by relying on Trick A:

$$\begin{aligned}
[h > 1] &\xrightarrow{r_{\text{AP}}} \diamond[h > 1] && \text{by Fact 6.4} \\
&\xrightarrow{\mu_{\mathbf{V}_\bullet X}^{\mathcal{H}}} \diamond \diamond[h > 1] \\
&\xrightarrow{\mathcal{H}_{\mathbf{V}}s_{\text{AP}}^\bullet} \diamond (s_{\text{AP}}^\bullet)^{-1}(\diamond[h > 1]) \\
&= \diamond[h > 1] && \text{by Corollary 7.3} \\
&\xrightarrow{\iota_{\alpha_{\text{AP}}^\bullet}} \diamond[h > 1] \cap \mathcal{H}_{\mathbf{V}}^{cvx}\mathbb{P}_{\text{AP}}^\bullet X.
\end{aligned}$$

Now this looks just like  $r_{\text{AP}}$ , and we imitate it by letting  $r: \mathcal{H}_{\mathbf{V}}^{cvx}\mathbb{P}_{\text{AP}}^\bullet X \rightarrow \mathbb{P}_{\text{AP}}^\bullet X$  be defined by  $r(C)(h) \stackrel{\text{def}}{=} \sup_{F \in C} F(h)$ . This is well-defined and lower semicontinuous, and  $r^{-1}([h > 1]) = \diamond[h > 1] \cap \mathcal{H}_{\mathbf{V}}^{cvx}\mathbb{P}_{\text{AP}}^\bullet X$ , so  $\mu_{\mu_X}^{\bar{S}} = r$ .

4. Standard category theory, see [34, Chapter VI, Section 2, Theorem 1] for example.  $\square$

Finally, the weak composite monad of  $S$  with  $T$  is  $U^T\bar{S}F^T$ , with unit  $U^T\eta^{\bar{S}F^T} \circ \eta^T$  and multiplication  $U^T\mu^{\bar{S}F^T} \circ U^T\bar{S}\epsilon^T\bar{S}F^T$ , where  $F^T \dashv U^T$  is the adjunction between the base category (**Top** in our case) and the category of  $T$ -algebras, and  $\epsilon^T$  is its counit.

**Theorem 8.4.** *The weak composite monad associated with  $\lambda^{\flat}: \mathbf{V}_\bullet\mathcal{H}_{\mathbf{V}} \rightarrow \mathcal{H}_{\mathbf{V}}\mathbf{V}_\bullet$  on  $\mathbf{Top}^{\flat}$  (see Definition 5.5) is  $(\mathbb{P}_{\text{AP}}^\bullet, \eta^{\text{AP}}, \mu^{\text{AP}})$ .*

PROOF. We take  $T \stackrel{\text{def}}{=} \mathbf{V}_\bullet$ ,  $S \stackrel{\text{def}}{=} \mathcal{H}_{\mathbf{V}}$ ,  $\lambda \stackrel{\text{def}}{=} \lambda^{\flat}$ . The functor  $U^T$  takes every  $\mathbf{V}_\bullet$ -algebra  $\alpha: \mathbf{V}_\bullet X \rightarrow X$  to  $X$  and every  $\mathbf{V}_\bullet$ -algebra morphism  $f: (\alpha: \mathbf{V}_\bullet X \rightarrow X) \rightarrow (\beta: \mathbf{V}_\bullet Y \rightarrow Y)$  to the underlying continuous map  $f: X \rightarrow Y$ . The functor  $F^T$  takes every space  $X$  to  $\mu_X: \mathbf{V}_\bullet\mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$ , and every continuous map  $f: X \rightarrow Y$  to  $\mathbf{V}_\bullet f$ , seen as a morphism of  $\mathbf{V}_\bullet$ -algebras.

Then  $U^T\bar{S}F^T$  maps every space  $X$  to the splitting  $\mathbb{P}_{\text{AP}}^\bullet X$  of  $\mathcal{H}_{\mathbf{V}}\mu_X \circ \lambda_X^{\flat} \circ \eta_{\mathcal{H}_{\mathbf{V}}X}$  we have obtained in Lemma 8.1. We also obtain  $\pi_{\mu_X} \stackrel{\text{def}}{=} r_{\text{AP}}$ ,  $\iota_{\mu_X} \stackrel{\text{def}}{=} s_{\text{AP}}^\bullet$ .



Given any continuous map  $f: X \rightarrow Y$ ,  $U^T \bar{S} F^T(f)$  is equal to (the map underlying the  $\mathbf{V}_\bullet$ -algebra)  $\bar{S}(\mathbf{V}_\bullet f)$ , namely  $\pi_{\mu_Y} \circ \mathcal{H}_V \mathbf{V}_\bullet f \circ \iota_{\mu_X} = r_{\text{AP}} \circ \mathcal{H}_V \mathbf{V}_\bullet f \circ s_{\text{AP}}^\bullet$ . We claim that this is equal to  $\mathbb{P}_{\text{AP}}^\bullet(f)$ , namely that it maps every  $F \in \mathbb{P}_{\text{AP}}^\bullet X$  to  $(h \in \mathcal{L}X \mapsto F(f \circ h))$ . We use Trick A:

$$\begin{aligned}
[h > 1] &\xrightarrow{r_{\text{AP}}} \diamond [h > 1] && \text{by Fact 6.4} \\
&\xrightarrow{\mathcal{H}_V \mathbf{V}_\bullet f} \diamond (\mathbf{V}_\bullet f)^{-1}([h > 1]) = \diamond [h \circ f > 1] \\
&\xrightarrow{s_{\text{AP}}^\bullet} [h \circ f > 1] && \text{by Corollary 7.3} \\
&= (\mathbb{P}_{\text{AP}}^\bullet(f))^{-1}([h > 1]).
\end{aligned}$$

From Lemma 8.3, item 2, the unit of  $\bar{S}$  at the algebra  $\mu_X$  is  $\eta_{\mu_X}^{\bar{S}} = r_{\text{AP}} \circ \eta_{\mathbf{V}_\bullet X}^{\mathcal{H}}$ , which is the map  $\nu \in \mathbf{V}_\bullet X \mapsto r_{\text{AP}}(\downarrow \nu)$ , namely  $\nu \in \mathbf{V}_\bullet X \mapsto (h \in \mathcal{L}X \mapsto \int_{x \in X} h(x) d\nu)$  from  $\mathbf{V}_\bullet X$  to  $\mathbb{P}_{\text{AP}}^\bullet X$ .

The unit of the weak composite monad is  $U^T \eta^{\bar{S}} F^T \circ \eta^T$ . At object  $X$ , this maps every  $x \in X$  to  $(h \in \mathcal{L}X \mapsto \int_{x' \in X} h(x') d\delta_x)$ , namely to  $(h \in \mathcal{L}X \mapsto h(x))$ , and that is exactly the unit of the  $\mathbb{P}_{\text{AP}}^\bullet$  monad.

The multiplication is  $U^T \mu^{\bar{S}} F^T \circ U^T \bar{S} \epsilon^T \bar{S} F^T$ . We recall what that means. At any object  $X$ ,  $\bar{S} F^T X$  is the  $T$ -algebra  $\bar{S} \mu_X: T \bar{S} T X \rightarrow \bar{S} T X$ . Then  $\epsilon^T$  evaluated at this  $T$ -algebra is the bottom arrow in the following diagram:

$$\begin{array}{ccc}
T T \bar{S} T X & \xrightarrow{T \epsilon_{\bar{S} F^T X}^T} & T \bar{S} T X \\
\mu_{\bar{S} T X} \downarrow & & \downarrow \bar{S} \mu_X \\
T \bar{S} T X & \xrightarrow{\epsilon_{\bar{S} F^T X}^T} & \bar{S} T X
\end{array}$$

from the  $T$ -algebra  $\alpha \stackrel{\text{def}}{=} \mu_{\bar{S} T X}$  that is the leftmost vertical arrow to the  $T$ -algebra  $\beta \stackrel{\text{def}}{=} \bar{S} \mu_X$  on the right. We apply  $\bar{S}$  to the morphism  $\epsilon_{\bar{S} F^T X}^T$  (i.e., we now apply  $\bar{S}$  to a morphism of  $T$ -algebras, not to a  $T$ -algebra, as we did before), and we obtain the composition:

$$\bar{S} T \bar{S} T X \xrightarrow{\iota_\alpha} S T \bar{S} T X \xrightarrow{S \epsilon_{\bar{S} F^T X}^T} S \bar{S} T X \xrightarrow{\pi_\beta} \bar{S} S T X. \quad (8)$$

Applying  $U^T$  to the latter, we obtain the same composition, this time seen as a morphism in the base category, instead of as a morphism in the category

of  $T$ -algebras. We finally compose it with  $U^T \mu_{FTX}^{\bar{S}}$ , which is  $\mu_{\mu_X}^{\bar{S}} : \overline{SSTX} \rightarrow \overline{STX}$ .

By Lemma 8.3, item 4,  $\epsilon_{\overline{STX}}^T : \alpha \rightarrow \beta$  is  $\beta$  itself, where  $\beta = \overline{S}\mu_X = \alpha_{\mathbb{A}\mathbb{P}}^\bullet$  (Lemma 8.3, item 1) and  $\alpha = \mu_{\overline{STX}} = \mu_{\mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X}$ : we have  $\overline{STX} = \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X$ , using Lemma 8.1, and  $\iota_\alpha = s_{\mathbb{A}\mathbb{P}}^\bullet$ . Using Lemma 8.2,  $\pi_\beta : \mathcal{H}_V \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X \rightarrow \mathcal{H}_V^{cvx} \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X$  is the corestriction of (7) (call it  $f$ ), which maps every convex element of  $\mathcal{H}_V \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X$  to itself.

Hence the multiplication of the weak composite monad, evaluated at  $X$ , is the composition of (8) with  $\mu_{\mu_X}^{\bar{S}}$  (given in Lemma 8.3, item 3), namely:

$$\mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X \xrightarrow{s_{\mathbb{A}\mathbb{P}}^\bullet} \mathcal{H}_V \mathbf{V}_\bullet \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X \xrightarrow{\mathcal{H}_V \alpha_{\mathbb{A}\mathbb{P}}^\bullet} \mathcal{H}_V \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X \xrightarrow{f} \mathcal{H}_V^{cvx} \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X \xrightarrow{\mu_{\mu_X}^{\bar{S}}} \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X \quad (9)$$

For every  $\mathcal{F} \in \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X$ ,  $\mathcal{H}_V \alpha_{\mathbb{A}\mathbb{P}}^\bullet(s_{\mathbb{A}\mathbb{P}}^\bullet(\mathcal{F})) = cl(\{\alpha_{\mathbb{A}\mathbb{P}}^\bullet(\xi) \mid \xi \in s_{\mathbb{A}\mathbb{P}}^\bullet(\mathcal{F})\}) = cl(\{(h \in \mathcal{L}X \mapsto \int_{F \in \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X} F(h) d\xi) \mid \xi \in s_{\mathbb{A}\mathbb{P}}^\bullet(\mathcal{F})\})$ . It is easy to see that this is a convex set, because  $s_{\mathbb{A}\mathbb{P}}^\bullet(\mathcal{F})$  is convex, integration is linear in the continuous valuation, and closure preserves convexity (Fact 7.8). Therefore  $f$  maps it to itself, and, using Lemma 8.3, item 3,  $\mu_{\mu_X}^{\bar{S}}$  maps it to:

$$\begin{aligned} (h \in \mathcal{L}X \mapsto \sup_{F \in \mathcal{H}_V \alpha_{\mathbb{A}\mathbb{P}}^\bullet(s_{\mathbb{A}\mathbb{P}}^\bullet(\mathcal{F}))} F(h)) &= (h \in \mathcal{L}X \mapsto \sup_{F \in cl(\alpha_{\mathbb{A}\mathbb{P}}^\bullet[s_{\mathbb{A}\mathbb{P}}^\bullet(\mathcal{F})])} F(h)) \\ &= (h \in \mathcal{L}X \mapsto \sup_{F \in \alpha_{\mathbb{A}\mathbb{P}}^\bullet[s_{\mathbb{A}\mathbb{P}}^\bullet(\mathcal{F})]} F(h)) \quad \text{by Fact 7.5} \\ &= (h \in \mathcal{L}X \mapsto \sup_{\xi \in s_{\mathbb{A}\mathbb{P}}^\bullet(\mathcal{F})} \alpha_{\mathbb{A}\mathbb{P}}^\bullet(\xi)(h)) \\ &= (h \in \mathcal{L}X \mapsto \sup_{\xi \in s_{\mathbb{A}\mathbb{P}}^\bullet(\mathcal{F})} \int_{F \in \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X} F(h) d\xi) \\ &= (h \in \mathcal{L}X \mapsto r_{\mathbb{A}\mathbb{P}}(s_{\mathbb{A}\mathbb{P}}^\bullet(\mathcal{F}))(F \in \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X \mapsto F(h))) \\ &= (h \in \mathcal{L}X \mapsto \mathcal{F}(F \in \mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet X \mapsto F(h))), \end{aligned}$$

and this is exactly what the multiplication of the  $\mathbb{P}_{\mathbb{A}\mathbb{P}}^\bullet$  monad, evaluated at  $X$  and applied to  $\mathcal{F}$ , produces.  $\square$

## Part III

# A weak distributive law between $\mathbf{V}_\bullet$ and $\mathcal{P}\ell_{\mathbf{V}}^q$ , resp. $\mathcal{P}\ell_{\mathbf{V}}$

Combining  $\mathcal{Q}_{\mathbf{V}}$  and  $\mathcal{H}_{\mathbf{V}}$ , there is a monad of lenses and a monad of quasi-lenses, which we define below. We start with lenses, which are probably more well-known. Quasi-lenses are easier to work with.

A *lens* is a non-empty set of the form  $Q \cap C$  where  $Q$  is compact saturated and  $C$  is closed in  $X$ . The *Vietoris topology* has subbasic open subsets of the form  $\square U$  (the set of lenses included in  $U$ ) and  $\diamond U$  (the set of lenses that intersect  $U$ ), for each open subset  $U$  of  $X$ . We write  $\mathcal{P}\ell X$  for the set of lenses on  $X$ , and we let  $\mathcal{P}\ell_{\mathbf{V}} X$  denote  $\mathcal{P}\ell X$  with the Vietoris topology. The specialization ordering of  $\mathcal{P}\ell_{\mathbf{V}} X$  is the *topological Egli-Milner ordering*:  $L \sqsubseteq^{\text{TEM}} L'$  if and only if  $\uparrow L \supseteq \uparrow L'$  and  $cl(L) \subseteq cl(L')$  [12, Discussion before Fact 4.1]. This is an ordering, not just a preordering, hence  $\mathcal{P}\ell_{\mathbf{V}} X$  is  $T_0$ .

A *quasi-lens* on a topological space  $X$  is a pair  $(Q, C)$  of a compact saturated subset  $Q$  and a closed subset  $C$  of  $X$  such that:

1.  $Q$  intersects  $C$ ;
2.  $Q \subseteq \uparrow(Q \cap C)$ ;
3. for every open neighborhood  $U$  of  $Q$ ,  $C \subseteq cl(U \cap C)$ .

The notion originates from [23, Theorem 9.6], and was rediscovered in [12, Section 5]. We write  $\mathcal{P}\ell^q X$  for the space of quasi-lenses on  $X$ . The *Vietoris topology* on  $\mathcal{P}\ell^q X$  is generated by the subbasic open sets  $\square^q U \stackrel{\text{def}}{=} \{(Q, C) \in \mathcal{P}\ell^q X \mid Q \subseteq U\}$  and  $\diamond^q U \stackrel{\text{def}}{=} \{(Q, C) \in \mathcal{P}\ell^q X \mid C \cap U \neq \emptyset\}$ . We write  $\mathcal{P}\ell_{\mathbf{V}}^q X$  for  $\mathcal{P}\ell^q X$  with the Vietoris topology. Its specialization preordering is  $\supseteq \times \subseteq$ , which is antisymmetric, so  $\mathcal{P}\ell_{\mathbf{V}}^q X$  is  $T_0$  [19, Lemma 8.2, item 2]; also, the inclusion of  $\mathcal{P}\ell_{\mathbf{V}}^q X$  into  $\mathcal{Q}_{\mathbf{V}} X \times \mathcal{H}_{\mathbf{V}} X$  is a topological embedding [19, Lemma 8.2, item 1], and  $\mathcal{P}\ell_{\mathbf{V}}^q$  extends to an endofunctor on  $\mathbf{Top}$ , whose action on morphisms  $f: X \rightarrow Y$  is defined by  $\mathcal{P}\ell_{\mathbf{V}}^q f(Q, C) \stackrel{\text{def}}{=} (\mathcal{Q}_{\mathbf{V}} f(Q), \mathcal{H}_{\mathbf{V}} f(C)) = (\uparrow f[Q], cl(f[C]))$ , and is such that  $(\mathcal{P}\ell_{\mathbf{V}}^q f)^{-1}(\square^q V) = \square^q f^{-1}(V)$  and  $(\mathcal{P}\ell_{\mathbf{V}}^q f)^{-1}(\diamond^q V) = \diamond^q f^{-1}(V)$  for every  $V \in \mathcal{O}Y$  [19, Lemma 8.3].

The relationship between spaces of lenses and of quasi-lenses is as follows. For every topological space  $X$ , there is a topological embedding  $\iota_X: \mathcal{P}l_V X \rightarrow \mathcal{P}l_V^q X$ , defined by  $\iota_X(L) \stackrel{\text{def}}{=} (\uparrow L, cl(L))$ . There is a map  $\varrho_X: \mathcal{P}l_V^q X \rightarrow \mathcal{P}l_V X$  defined by  $\varrho_X(Q, C) \stackrel{\text{def}}{=} Q \cap C$ , and  $\varrho_X \circ \iota_X = \text{id}_X$ . When  $X$  is weakly Hausdorff [17, Lemma 6.1, Proposition 6.2, Theorem 6.3] or quasi-Polish [19, Theorem 9.6], those define inverse continuous maps, hence a homeomorphism between  $\mathcal{P}l_V X$  and  $\mathcal{P}l_V^q X$ . A space is *weakly Hausdorff* if and only if given any two points  $x$  and  $y$ , every open neighborhood  $W$  of  $\uparrow x \cap \uparrow y$  contains an intersection  $U \cap V$  of an open neighborhood  $U$  of  $x$  and an open neighborhood  $V$  of  $y$  [29, Lemma 6.6]. Every Hausdorff space is weakly Hausdorff, every stably locally compact space is weakly Hausdorff [29, Lemma 8.1]. The *quasi-Polish spaces* are the topological spaces underlying a separable Smyth-complete quasi-metric space [6]. We will not define them any further, except than to notice that every  $\omega$ -continuous dcpo is quasi-Polish in its Scott topology. (See Section 2.2 of [1] for continuous and  $\omega$ -continuous dcpos.)

We will write  $\pi_1: \mathcal{P}l_V^q X \rightarrow \mathcal{Q}_V X$  and  $\pi_2: \mathcal{P}l_V^q X \rightarrow \mathcal{H}_V X$  for the respective projections. For every  $U \in \mathcal{O}X$ ,  $\pi_1^{-1}(\square U) = \square^q U$  and  $\pi_2^{-1}(\diamond U) = \diamond^q U$ , so they are continuous. Those define natural transformations  $\pi_1: \mathcal{P}l_V^q \rightarrow \mathcal{Q}_V$  and  $\pi_2: \mathcal{P}l_V^q \rightarrow \mathcal{H}_V$ , as one can see easily using Trick A.

**Proposition 8.5.** *There is a monad  $(\mathcal{P}l_V^q, \eta^q, \mu^q)$  on  $\mathbf{Top}$ ,*

- *whose unit  $\eta_X^q: X \rightarrow \mathcal{P}l_V^q X$  maps every  $x \in X$  to  $(\eta_X^{\mathcal{Q}}(x), \eta_X^{\mathcal{H}}(x)) = (\uparrow x, \downarrow x)$ , and is such that  $(\eta_X^q)^{-1}(\square^q U) = (\eta_X^q)^{-1}(\diamond^q U) = U$  for every  $U \in \mathcal{O}X$ , and is characterized uniquely by the commutativity of the diagram:*

$$\begin{array}{ccccc}
 X & \xlongequal{\quad} & X & \xlongequal{\quad} & X & (10) \\
 \eta_X^{\mathcal{Q}} \downarrow & & \eta_X^q \downarrow & & \eta_X^{\mathcal{H}} \downarrow \\
 \mathcal{Q}_V X & \xleftarrow{\pi_1} & \mathcal{P}l_V^q X & \xrightarrow{\pi_2} & \mathcal{H}_V X
 \end{array}$$

- *and whose multiplication  $\mu_X^q: \mathcal{P}l_V^q \mathcal{P}l_V^q X \rightarrow \mathcal{P}l_V^q X$  maps every quasi-lens  $(\mathcal{Q}, \mathcal{C})$  on  $\mathcal{P}l_V^q X$  to  $(\bigcup \pi_1[\mathcal{Q}], cl(\bigcup \pi_2[\mathcal{C}]))$ , is such that  $(\mu_X^q)^{-1}(\square^q U) = \square^q \square^q U$  and  $(\mu_X^q)^{-1}(\diamond^q U) = \diamond^q \diamond^q U$  for every  $U \in \mathcal{O}X$ , and is char-*

acterized uniquely by the commutativity of the diagram:

$$\begin{array}{ccccccc}
\mathcal{Q}_V \mathcal{Q}_V X & \xleftarrow{\mathcal{Q}_V \pi_1} & \mathcal{Q}_V \mathcal{P} \ell_V^q X & \xleftarrow{\pi_1} & \mathcal{P} \ell_V^q \mathcal{P} \ell_V^q X & \xrightarrow{\pi_2} & \mathcal{H}_V \mathcal{P} \ell_V^q X & \xrightarrow{\mathcal{H}_V \pi_2} & \mathcal{H}_V \mathcal{H}_V X \\
\downarrow \mu_X^{\mathcal{Q}} & & & & \downarrow \mu_X^q & & & & \downarrow \mu_X^{\mathcal{H}} \\
\mathcal{Q}_V X & \xleftarrow{\pi_1} & \mathcal{P} \ell_V^q X & \xrightarrow{\pi_2} & \mathcal{H}_V X & & & & 
\end{array} \tag{11}$$

PROOF. Unit. For every  $x \in X$ , we claim that  $(Q, C) \stackrel{\text{def}}{=}} (\uparrow x, \downarrow x)$  is a quasi-lens. First,  $\uparrow x \cap \downarrow x$  is non-empty, since it contains  $x$ . Second,  $Q \subseteq \uparrow(Q \cap C)$ , since every element of  $Q$  is larger than or equal to  $x$ , which is in  $Q \cap C$ . Third, since  $x \in Q \cap C$ ,  $C = \downarrow x = cl(\{x\}) \subseteq cl(Q \cap C)$ , from which it follows that for every open neighborhood  $U$  of  $Q$ ,  $C \subseteq (U \cap C)$ . The commutativity of (10) is an equivalent way of stating the definition of  $\eta_X^q$  (modulo the fact that all arrows are indeed continuous maps, which we show next).

For every  $U \in \mathcal{O}X$ , for every  $x \in X$ ,  $x \in (\eta_X^q)^{-1}(\square^q U)$  if and only if  $\eta_X^{\mathcal{Q}}(x) \in \square U$ , if and only if  $x \in U$ , and the equality  $(\eta_X^q)^{-1}(\diamond^q U) = U$  is proved similarly. In particular,  $\eta_X^q$  is continuous.

Finally, the commutativity of (10) characterizes  $\eta_X^q$ , since the pairing  $\langle \pi_1, \pi_2 \rangle: \mathcal{P} \ell_V^q X \rightarrow \mathcal{Q}_V X \times \mathcal{H}_V X$  is the inclusion map, hence is injective.

Multiplication. For every  $(\mathcal{Q}, \mathcal{C}) \in \mathcal{P} \ell_V^q \mathcal{P} \ell_V^q X$ , let  $(Q, C) \stackrel{\text{def}}{=} (\bigcup \pi_1[\mathcal{Q}], cl(\bigcup \pi_2[\mathcal{C}]))$ . We need to check that this is a quasi-lens.

We have  $\mu_X^{\mathcal{Q}}(\mathcal{Q}_V \pi_1(\mathcal{Q})) = \bigcup \uparrow \pi_1[\mathcal{Q}] = \bigcup \pi_1[\mathcal{Q}] = Q$  (because  $\pi_1[\mathcal{Q}]$  is upwards-closed, as one checks easily). In particular,  $Q$  is in  $\mathcal{Q}_V X$ . Also,  $\mu_X^{\mathcal{H}}(\mathcal{H}_V \pi_2(\mathcal{C})) = cl(\bigcup cl(\pi_2[\mathcal{C}]))$ . The outer closure is taken in  $X$ , while the inner closure is taken in  $\mathcal{H}_V X$ . We claim that this is equal to  $C$ . Clearly,  $C \subseteq cl(\bigcup cl(\pi_2[\mathcal{C}]))$ . Conversely, every open set  $U$  that intersects  $cl(\bigcup cl(\pi_2[\mathcal{C}]))$  intersects  $\bigcup cl(\pi_2[\mathcal{C}])$ , so there is an element of  $cl(\pi_2[\mathcal{C}])$  that intersects  $U$ ; in other words,  $cl(\pi_2[\mathcal{C}])$  intersects  $\diamond U$ , so  $\pi_2[\mathcal{C}]$  intersects  $\diamond U$ , equivalently,  $\bigcup \pi_2[\mathcal{C}] = C$  intersects  $U$ . Since  $C = \mu_X^{\mathcal{H}}(\mathcal{H}_V \pi_2(\mathcal{C}))$ , in particular  $C \in \mathcal{H}_V X$ . Also, we have just shown that the diagram (11) commutes (modulo the fact that all arrows are continuous maps, which we will check later).

Since  $(\mathcal{Q}, \mathcal{C})$  is a quasi-lens, there is an element  $(Q', C')$  in  $\mathcal{Q} \cap \mathcal{C}$ . Then there is an element  $x$  in  $Q' \cap C'$ , because  $(Q', C')$  is a quasi-lens. Since  $(Q', C') \in \mathcal{Q}$ ,  $Q'$  is in  $\pi_1[\mathcal{Q}]$ , so  $Q' \subseteq \bigcup \pi_1[\mathcal{Q}] = Q$ ; then  $x \in Q' \subseteq Q$ . Since  $(Q', C') \in \mathcal{C}$ ,  $C'$  is in  $\pi_2[\mathcal{C}]$ , so  $C' \subseteq cl(\bigcup \pi_2[\mathcal{C}]) = C$ ; then  $x \in C' \subseteq C$ . It follows that  $Q \cap C \neq \emptyset$ .

Let us show that  $Q \subseteq \uparrow(Q \cap C)$ . For every  $x \in Q = \bigcup \pi_1[\mathcal{Q}]$ , there is quasi-lens  $(Q', C') \in \mathcal{Q}$  such that  $x \in Q'$ . Since  $(\mathcal{Q}, \mathcal{C})$  is itself a quasi-lens,  $\mathcal{Q} \subseteq \uparrow(\mathcal{Q} \cap \mathcal{C})$ , so there is a quasi-lens  $(Q'', C'') \in \mathcal{Q} \cap \mathcal{C}$  such that  $(Q'', C'') (\supseteq \times \subseteq) (Q', C')$ . We have  $Q'' \supseteq Q'$  and  $x \in Q'$ , so  $x \in Q''$ ; also,  $Q'' \subseteq \uparrow(Q'' \cap C'')$ , so there is a point  $x'' \in Q'' \cap C''$  such that  $x'' \leq x$ . Since  $x'' \in Q''$  and  $(Q'', C'') \in \mathcal{Q}$ ,  $x''$  is in  $\bigcup \pi_1[\mathcal{Q}] = Q$ , and since  $x'' \in C''$  and  $(Q'', C'') \in \mathcal{C}$ ,  $x''$  is in  $\bigcup \pi_2[\mathcal{C}]$ , hence in its closure  $C$ . We have obtained that  $x$  is larger than or equal to a point  $x''$  in  $Q \cap C$ , so  $x \in \uparrow(Q \cap C)$ .

We now show that for every open neighborhood  $U$  of  $Q$ ,  $C \subseteq cl(U \cap C)$ . It suffices to show that for every open neighborhood  $U$  of  $Q$ , every open set  $V$  that intersects  $C$  also intersects  $U \cap C$ . Since  $Q = \bigcup \pi_1[\mathcal{Q}] \subseteq U$ , every quasi-lens  $(Q', C') \in \mathcal{Q}$  is such that  $Q' \subseteq U$ , namely is in  $\square^q U$ . Therefore  $\mathcal{Q} \subseteq \square^q U$ . Since  $(\mathcal{Q}, \mathcal{C})$  is a quasi-lens,  $\mathcal{C} \subseteq cl(\mathcal{C} \cap \square^q U)$ . Hence every open subset  $\mathcal{V}$  of  $\mathcal{P}\ell_V^q X$  that intersects  $\mathcal{C}$  also intersects  $\mathcal{C} \cap \square^q U$ . Since  $C = cl(\bigcup \pi_2[\mathcal{C}])$  and  $V$  intersects  $C$ ,  $V$  intersects  $\bigcup \pi_2[\mathcal{C}]$ , hence  $V$  intersects some closed set  $C'$  such that  $(Q', C') \in \mathcal{C}$ . In other words,  $\diamond^q V$  intersects  $\mathcal{C}$ . Taking  $\mathcal{V} \stackrel{\text{def}}{=} \diamond^q V$ , we obtain that  $\diamond^q V$  intersects  $\mathcal{C} \cap \square^q U$ . In other words, there is a quasi-lens  $(Q', C')$  in  $\mathcal{C}$  such that  $Q' \subseteq U$  and  $C'$  intersects  $V$ . Since  $(Q', C')$  is a quasi-lens,  $C' \subseteq cl(C' \cap U)$ , so  $V$  intersects  $C' \cap U$ . Hence  $C'$  intersects  $U \cap V$ . But, since  $(Q', C') \in \mathcal{C}$ ,  $C'$  is in  $\pi_2[\mathcal{C}]$ , so  $C' \subseteq C = cl(\bigcup \pi_2[\mathcal{C}])$ . Therefore  $C$  also intersects  $U \cap V$ , showing that  $V$  intersects  $U \cap C$ .

It remains to show that  $\mu_X^q$  is continuous. For every open subset  $U$  of  $X$ ,  $(\mu_X^q)^{-1}(\square^q U) = \{(\mathcal{Q}, \mathcal{C}) \in \mathcal{P}\ell_V^q \mathcal{P}\ell_V^q X \mid \bigcup \pi_1[\mathcal{Q}] \subseteq U\} = \{(\mathcal{Q}, \mathcal{C}) \in \mathcal{P}\ell_V^q \mathcal{P}\ell_V^q X \mid \forall (Q, C) \in \mathcal{Q}, Q \subseteq U\} = \square^q \square^q U$ , and  $(\mu_X^q)^{-1}(\diamond^q U) = \{(\mathcal{Q}, \mathcal{C}) \in \mathcal{P}\ell_V^q \mathcal{P}\ell_V^q X \mid cl(\bigcup \pi_2[\mathcal{C}]) \cap U \neq \emptyset\} = \{(\mathcal{Q}, \mathcal{C}) \in \mathcal{P}\ell_V^q \mathcal{P}\ell_V^q X \mid \bigcup \pi_2[\mathcal{C}] \cap U \neq \emptyset\} = \{(\mathcal{Q}, \mathcal{C}) \in \mathcal{P}\ell_V^q \mathcal{P}\ell_V^q X \mid \exists (Q, C) \in \mathcal{C}, C \cap U \neq \emptyset\} = \diamond^q \diamond^q U$ .

There are two easy ways to verify that  $\eta^q$  and  $\mu^q$  are natural, and the monad laws. One is by using Trick A, and the other one is as follows. Given any two maps  $f, g: Y \rightarrow \mathcal{P}\ell_V^q Z$  (for some spaces  $Y$  and  $Z$ ),  $f = g$  if and only if  $\pi_1 \circ f = \pi_1 \circ g$  and  $\pi_2 \circ f = \pi_2 \circ g$ , since the pairing  $\langle \pi_1, \pi_2 \rangle$  is the inclusion map, hence is injective. We use this new trick, as it will be clear that this reduces the question for  $\mathcal{P}\ell_V^q$  to those for  $\mathcal{Q}_V$  and  $\mathcal{H}_V$ .

Naturality. Given any continuous map  $f: X \rightarrow Y$ , we check that  $\eta_Y^q \circ f = \mathcal{P}\ell_V^q f \circ \eta_X^q$ :  $\pi_1 \circ \eta_Y^q \circ f = \eta_Y^q \circ f$  (by (10)) =  $\mathcal{Q}_V f \circ \eta_X^q$  (by naturality of  $\eta^q$ ) =  $\pi_1 \circ \mathcal{P}\ell_V^q f \circ \eta_X^q$ , and similarly  $\pi_2 \circ \eta_Y^q \circ f = \pi_2 \circ \mathcal{P}\ell_V^q f \circ \eta_X^q$ .

We also check that  $\mathcal{P}\ell_V^q f \circ \mu_X^q = \mu_Y^q \circ \mathcal{P}\ell_V^q \mathcal{P}\ell_V^q f$  by:

$$\begin{aligned}
\pi_1 \circ \mathcal{P}\ell_V^q f \circ \mu_X^q &= \mathcal{Q}_V f \circ \pi_1 \circ \mu_X^q && \text{def. of } \mathcal{P}\ell_V^q f \\
&= \mathcal{Q}_V f \circ \mu_X^{\mathcal{Q}} \circ \mathcal{Q}_V \pi_1 \circ \pi_1 && \text{by (11)} \\
&= \mu_Y^{\mathcal{Q}} \circ \mathcal{Q}_V \mathcal{Q}_V f \circ \mathcal{Q}_V \pi_1 \circ \pi_1 && \text{nat. of } \mu^{\mathcal{Q}} \\
&= \mu_Y^{\mathcal{Q}} \circ \mathcal{Q}_V \pi_1 \circ \mathcal{Q}_V \mathcal{P}\ell_V^q f \circ \pi_1 && \text{nat. of } \pi_1 \\
&= \mu_Y^{\mathcal{Q}} \circ \mathcal{Q}_V \pi_1 \circ \pi_1 \circ \mathcal{P}\ell_V^q \mathcal{P}\ell_V^q f && \text{nat. of } \pi_1 \\
&= \pi_1 \circ \mu_Y^q \circ \mathcal{P}\ell_V^q \mathcal{P}\ell_V^q f && \text{by (11),}
\end{aligned}$$

and similarly with  $\pi_2$  instead of  $\pi_1$ ,  $\mathcal{H}$  in lieu of  $\mathcal{Q}$ , and  $\mathcal{H}_V$  instead of  $\mathcal{Q}_V$ .

The monad laws. We proceed similarly. We verify that  $\mu_X^q \circ \eta_{\mathcal{P}\ell_V^q X}^q = \text{id}_{\mathcal{P}\ell_V^q X}$  by:

$$\begin{aligned}
\pi_1 \circ \mu_X^q \circ \eta_{\mathcal{P}\ell_V^q X}^q &= \mu_X^{\mathcal{Q}} \circ \mathcal{Q}_V \pi_1 \circ \pi_1 \circ \eta_{\mathcal{P}\ell_V^q X}^q && \text{by (11)} \\
&= \mu_X^{\mathcal{Q}} \circ \mathcal{Q}_V \pi_1 \circ \eta_{\mathcal{P}\ell_V^q X}^{\mathcal{Q}} && \text{by (10)} \\
&= \mu_X^{\mathcal{Q}} \circ \eta_{\mathcal{Q}_V X} \circ \pi_1 && \text{nat. of } \pi_1 \\
&= \pi_1,
\end{aligned}$$

by the monad law  $\mu_X^{\mathcal{Q}} \circ \eta_{\mathcal{Q}_V X}^{\mathcal{Q}} = \text{id}_{\mathcal{Q}_V X}$  for  $\mathcal{Q}_V$ ; similarly with  $\pi_2$  instead of  $\pi_1$ .

We verify that  $\mu_X^q \circ \mathcal{P}\ell_V^q \eta_X^q = \text{id}_{\mathcal{P}\ell_V^q X}$  by:

$$\begin{aligned}
\pi_1 \circ \mu_X^q \circ \mathcal{P}\ell_V^q \eta_X^q &= \mu_X^{\mathcal{Q}} \circ \mathcal{Q}_V \pi_1 \circ \pi_1 \circ \mathcal{P}\ell_V^q \eta_X^q && \text{by (11)} \\
&= \mu_X^{\mathcal{Q}} \circ \mathcal{Q}_V \pi_1 \circ \mathcal{Q}_V \eta_X^q \circ \pi_1 && \text{def. of } \mathcal{P}\ell_V^q \text{ on morphisms} \\
&= \mu_X^{\mathcal{Q}} \circ \mathcal{Q}_V \eta_X^{\mathcal{Q}} \circ \pi_1 && \text{by (10)} \\
&= \pi_1,
\end{aligned}$$

by the monad law  $\mu_X^{\mathcal{Q}} \circ \mathcal{Q}_V \eta_X^{\mathcal{Q}} = \text{id}_{\mathcal{Q}_V X}$  for  $\mathcal{Q}_V$ ; similarly with  $\pi_2$  instead of  $\pi_1$ .

Finally, we verify that  $\mu_X^q \circ \mu_{\mathcal{P}\ell_V^q X}^q = \mu_X^q \circ \mathcal{P}\ell_V^q \mu_X^q$ . Once again, we do this for  $\pi_1$  only:

$$\begin{aligned}
\pi_1 \circ \mu_X^q \circ \mu_{\mathcal{P}\ell_V^q X}^q &= \mu_X^{\mathcal{Q}} \circ \mathcal{Q}_V \pi_1 \circ \pi_1 \circ \mu_{\mathcal{P}\ell_V^q X}^q && \text{by (11)} \\
&= \mu_X^{\mathcal{Q}} \circ \mathcal{Q}_V \pi_1 \circ \mu_{\mathcal{P}\ell_V^q X}^{\mathcal{Q}} \circ \mathcal{Q}_V \pi_1 \circ \pi_1 && \text{by (11)} \\
&= \mu_X^{\mathcal{Q}} \circ \mu_{\mathcal{P}\ell_V^q X}^{\mathcal{Q}} \circ \mathcal{Q}_V \mathcal{Q}_V \pi_1 \circ \mathcal{Q}_V \pi_1 \circ \pi_1 && \text{nat. of } \mu^{\mathcal{Q}}
\end{aligned}$$

while:

$$\begin{aligned}
\pi_1 \circ \mu_X^q \circ \mathcal{P}\ell_V^q \mu_X^q &= \mu_X^Q \circ \mathcal{Q}_V \pi_1 \circ \pi_1 \circ \mathcal{P}\ell_V^q \mu_X^q && \text{by (11)} \\
&= \mu_X^Q \circ \mathcal{Q}_V \pi_1 \circ \mathcal{Q}_V \mu_X^q \circ \pi_1 && \text{def. of } \mathcal{P}\ell_V^q \text{ on morphisms} \\
&= \mu_X^Q \circ \mathcal{Q}_V \mu_X^Q \circ \mathcal{Q}_V \mathcal{Q}_V \pi_1 \circ \mathcal{Q}_V \pi_1 \circ \pi_1 && \text{by (11),}
\end{aligned}$$

and we conclude since  $\mu_X^Q \circ \mu_{\mathcal{P}\ell_V^q X}^Q = \mu_X^Q \circ \mathcal{Q}_V \mu_X^Q$ , by the final monad law on  $\mathcal{Q}_V$ .  $\square$

**Remark 8.6.** *There is also a  $\mathcal{P}\ell_V$  monad (of lenses) on  $\mathbf{Top}$ , but we will not care about it except on full subcategories of weakly Hausdorff of quasi-Polish spaces, where we can transport the monad structure of  $\mathcal{P}\ell_V^q$  over to  $\mathcal{P}\ell_V$ . We obtain that for every continuous map  $f: X \rightarrow Y$ ,  $\mathcal{P}\ell_V f = \varrho_Y \circ \mathcal{P}\ell_V^q f \circ \iota_X$  maps every lens  $L$  on  $X$  to  $\uparrow f[\uparrow L] \cap cl(f[cl(L)])$  (which we may simplify as  $\uparrow f[L] \cap cl(f[L])$ ), the unit  $\eta_X^{\mathcal{P}\ell}$  maps every  $x \in X$  to  $\varrho_X(\uparrow x, \downarrow x) = \uparrow x \cap \downarrow x$  (which simplifies to  $\{x\}$  if  $X$  is  $T_0$ ), and the multiplication  $\mu_X^{\mathcal{P}\ell} = \varrho_X \circ \mu_X^q \circ \mathcal{P}\ell_V^q \iota_X \circ \iota_{\mathcal{P}\ell X}$  maps every lens  $\mathcal{L} \in \mathcal{P}\ell_V \mathcal{P}\ell_V X$  to  $\uparrow \bigcup \mathcal{L} \cap cl(\bigcup \mathcal{L})$ .*

In order to obtain a distributive law over  $\mathbf{V}_\bullet$ , we need to further restrict the category of topological spaces we can work on. Not only do we need our spaces  $X$  to be such that  $\mathcal{L}X$  is locally convex, but addition on  $\mathcal{L}X$  needs to be *almost open*, meaning that given any two open subsets  $\mathcal{U}$  and  $\mathcal{V}$  of  $\mathcal{L}X$ ,  $\uparrow(\mathcal{U} + \mathcal{V})$  is open. (We will also need  $X$  to be compact when  $\bullet$  is “1”.) This is the case if  $X$  is stably locally compact [14, Lemma 3.24]. (We have already stated that every stably locally compact space  $X$  is weakly Hausdorff [29, Lemma 8.1], so that  $\mathcal{P}\ell_V X \cong \mathcal{P}\ell_V^q X$  in this case.)

We also need our category of spaces to be closed under  $\mathbf{V}_\bullet$ ,  $\mathcal{P}\ell_V^q$ , and under retracts. The category of stably compact spaces is closed under retracts [33, Proposition, bottom of p.153, and subsequent discussion], see also [27, Proposition 2.17].

As far as  $\mathbf{V}_\bullet X$  is concerned, it was shown in [3, Theorem 39] that  $\mathbf{V}_{\leq 1} X$  is stably compact for every stably compact space  $X$ . The same proof shows that  $\mathbf{V}X$  is stably compact under the same assumption. Additionally,  $\mathbf{V}_1 X$  occurs as the subspace  $\langle X \geq 1 \rangle$ , where  $\langle Q \geq r \rangle$  denotes  $\{\nu \in \mathbf{V}_{\leq 1} X \mid \forall U \in \mathcal{O}X, Q \subseteq U \Rightarrow \nu(U) \geq r\}$ , and is compact saturated for every compact saturated subset  $Q$  of the stably compact space  $X$  and for every  $r \in \mathbb{R}_+$  [12, Lemma 6.6]. But any compact saturated subset of a stably compact space is stably compact; more generally, any patch-closed subset of a stably



compact space, namely, any subset that we can obtain as an intersection of finite unions of closed subsets and of compact saturated subsets, is stably compact [27, Proposition 2.16].

As for  $\mathcal{P}\ell_{\mathbf{V}}^{\mathfrak{q}}X$ , it was noted in [12, Fact 5.2] that for every sober space  $X$ ,  $\mathcal{P}\ell_{\mathbf{V}}^{\mathfrak{q}}X$  is homeomorphic to Heckmann’s space of  $\mathbf{A}$ -valuations on  $X$  [24]. If  $X$  is stably compact, then, both spaces are stably compact [12, Proposition 5.13].

**Definition 8.7.** *Let  $\mathbf{Top}^{\mathfrak{q}}$  be any full subcategory of  $\mathbf{Top}$  consisting of spaces  $X$  such that  $\mathcal{L}X$  is locally convex and has an almost open addition map, and which is closed under  $\mathcal{P}\ell_{\mathbf{V}}^{\mathfrak{q}}$ ,  $\mathbf{V}\bullet$  and under retracts—for example the category of stably compact spaces.*

## 9. The monad of forks

A *fork* on a space  $X$  is any pair  $(F^-, F^+)$  of a superlinear prevision  $F^-$  on  $X$  and of a sublinear prevision  $F^+$  on  $X$  satisfying *Walley’s condition*:

$$F^-(h + h') \leq F^-(h) + F^+(h') \leq F^+(h + h')$$

for all  $h, h' \in \mathcal{L}X$  [11, 30]. A fork is *subnormalized*, resp. *normalized* if and only if both  $F^-$  and  $F^+$  are.

We write  $\mathbb{P}_{\text{ADP}}X$  for the set of all forks on  $X$ , and  $\mathbb{P}_{\text{ADP}}^{\leq 1}X$ ,  $\mathbb{P}_{\text{ADP}}^1X$  for their subsets of subnormalized, resp. normalized, forks. The *weak topology* on each is the subspace topology induced by the inclusion into the larger space  $\mathbb{P}_{\text{DP}}X \times \mathbb{P}_{\text{AP}}X$ . A subbase of the weak topology is composed of two kinds of open subsets:  $[h > r]^-$ , defined as  $\{(F^-, F^+) \mid F^-(h) > r\}$ , and  $[h > r]^+$ , defined as  $\{(F^-, F^+) \mid F^+(h) > r\}$ , where  $h \in \mathcal{L}X$ ,  $r \in \mathbb{R}^+$ . The specialization ordering of spaces of forks is the product ordering  $\leq \times \leq$ , where  $\leq$  denotes the pointwise ordering on previsions. In particular, all those spaces of forks are  $T_0$ .

Whether  $\bullet$  is nothing, “ $\leq 1$ ”, or “1”,  $\mathbb{P}_{\text{ADP}}^{\bullet}$  defines an endofunctor on  $\mathbf{Top}$ , whose action on morphisms is given by  $\mathbb{P}_{\text{ADP}}^{\bullet}f \stackrel{\text{def}}{=} (\mathbb{P}f, \mathbb{P}f)$ .

**Proposition 9.1.** *Let  $\bullet$  be nothing, “ $\leq 1$ ”, or “1”. For every topological space  $X$  such that  $\mathcal{L}X$  is locally convex and has an almost open addition map (and such that  $X$  is compact, in case  $\bullet$  is “1”), there are continuous*

maps  $r_{\text{ADP}}: \mathcal{P}\ell_{\vee}^{\text{q}}\mathbf{V}_{\bullet}X \rightarrow \mathbb{P}_{\text{ADP}}^{\bullet}X$  and  $s_{\text{ADP}}^{\bullet}: \mathbb{P}_{\text{ADP}}^{\bullet}X \rightarrow \mathcal{P}\ell_{\vee}^{\text{q}}\mathbf{V}_{\bullet}X$  defined by:

$$\begin{aligned} r_{\text{ADP}}(Q, C) &\stackrel{\text{def}}{=} (r_{\text{DP}}(Q), r_{\text{AP}}(C)) \\ s_{\text{ADP}}^{\bullet}(F^{-}, F^{+}) &\stackrel{\text{def}}{=} (s_{\text{DP}}^{\bullet}(F^{-}), s_{\text{AP}}^{\bullet}(F^{+})). \end{aligned}$$

Additionally,  $r_{\text{ADP}}$  and  $s_{\text{ADP}}^{\bullet}$  are natural in  $X$ .

PROOF. For every quasi-lens  $(Q, C)$  on  $\mathbf{V}_{\bullet}X$ , we check that  $(F^{-}, F^{+}) \stackrel{\text{def}}{=} (r_{\text{DP}}(Q), r_{\text{AP}}(C))$  is a fork. For all  $h, h' \in \mathcal{L}X$ ,

$$\begin{aligned} F^{-}(h + h') &= \min_{\nu \in Q} \int_{x \in X} (h(x) + h'(x)) d\nu \\ &\leq \min_{\nu \in Q \cap C} \int_{x \in X} (h(x) + h'(x)) d\nu \\ &\leq \min_{\nu \in Q \cap C} \left( \int_{x \in X} h(x) d\nu + F^{+}(h') \right) \\ &\quad \text{since } \int_{x \in X} h'(x) d\nu \leq F^{+}(h') \text{ for every } \nu \in Q \cap C \\ &= \min_{\nu \in Q \cap C} \int_{x \in X} h(x) d\nu + F^{+}(h') \\ &= \min_{\nu \in \uparrow(Q \cap C)} \int_{x \in X} h(x) d\nu + F^{+}(h') \\ &\leq \min_{\nu \in Q} \int_{x \in X} h(x) d\nu + F^{+}(h') \quad \text{since } Q \subseteq \uparrow(Q \cap C) \\ &= F^{-}(h) + F^{+}(h'). \end{aligned}$$

For Walley's other inequality, we observe that:

$$\begin{aligned} F^{-}(h) + F^{+}(h') &= \sup_{r \in \mathbb{R}_{+}, r < F^{-}(h)}^{\uparrow} (r + F^{+}(h')) \\ &= \sup_{r \in \mathbb{R}_{+}, r < F^{-}(h), \nu \in C} \left( r + \int_{x \in X} h'(x) d\nu \right). \end{aligned}$$

Then, for every  $r \in \mathbb{R}_{+}$  such that  $r < F^{-}(h)$ , we have  $r < \min_{\nu \in Q} \int_{x \in X} h(x) d\nu$ , so  $Q \subseteq [h > r]$ . Since  $(Q, C)$  is a quasi-lens,  $C \subseteq \text{cl}([h > r] \cap C)$ , so:

$$\begin{aligned} F^{-}(h) + F^{+}(h') &\leq \sup_{r \in \mathbb{R}_{+}, r < F^{-}(h), \nu \in \text{cl}([h > r] \cap C)} \left( r + \int_{x \in X} h'(x) d\nu \right) \\ &= \sup_{r \in \mathbb{R}_{+}, r < F^{-}(h), \nu \in [h > r] \cap C} \left( r + \int_{x \in X} h'(x) d\nu \right) \quad \text{by Fact 7.5.} \end{aligned}$$

Fact 7.5 applies, since  $\nu \in \mathbf{V}_\bullet X \mapsto \int_{x \in X} h'(x) d\nu$  is lower semicontinuous, as the inverse image of any basic open set  $[t, \infty]$  is  $[h' > t]$ ; and therefore  $\nu \in \mathbf{V}_\bullet X \mapsto r + \int_{x \in X} h'(x) d\nu$  is lower semicontinuous, too. Now for every  $r \in \mathbb{R}_+$  such that  $r < F^-(h)$ , for every  $\nu \in [h > r] \cap C$ , we have  $\int_{x \in X} h(x) d\nu > r$  by definition of  $[h > r]$ , so:

$$\begin{aligned} F^-(h) + F^+(h') &\leq \sup_{r \in \mathbb{R}_+, r < F^-(h), \nu \in [h > r] \cap C} \left( \int_{x \in X} h(x) d\nu + \int_{x \in X} h'(x) d\nu \right) \\ &\leq \sup_{\nu \in C} \left( \int_{x \in X} h(x) d\nu + \int_{x \in X} h'(x) d\nu \right) \\ &= F^+(h + h'). \end{aligned}$$

Additionally, if  $\bullet$  is “ $\leq 1$ ” (resp., “1”), then  $F^- = r_{\text{DP}}(Q)$  and  $F^+ = r_{\text{AP}}(C)$  are subnormalized (resp., normalized), so  $r_{\text{ADP}}$  takes its values in  $\mathbb{P}_{\text{ADP}}^\bullet X$ . The fact that  $r_{\text{ADP}}$  is continuous follows from the fact that  $r_{\text{DP}}$  and  $r_{\text{AP}}$  are, and similarly for naturality.

Turning to  $s_{\text{ADP}}^\bullet$ , we must show that for every  $(F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X$ ,  $(Q, C) \stackrel{\text{def}}{=} (s_{\text{DP}}^\bullet(F^-), s_{\text{AP}}^\bullet(F^+))$  is a quasi-lens. That  $s_{\text{ADP}}^\bullet$  is continuous and natural will then follow from the same properties for  $s_{\text{DP}}^\bullet$  and  $s_{\text{AP}}^\bullet$ .

Lemma 3.27 of [14] states that there is a map (called  $s_{\text{ADP}}^\bullet$  there; although this is not the same map as our  $s_{\text{ADP}}^\bullet$ , the two are definitely related) from  $\mathbb{P}_{\text{ADP}}^\bullet X$  to the space of lenses on  $\mathbb{P}_p^\bullet X$ , and which sends  $(F^-, F^+)$  to  $s_{\text{DP}}(F^-) \cap s_{\text{AP}}^\bullet(F^+)$ . In particular,  $Q \cap C = s_{\text{DP}}(F^-) \cap s_{\text{AP}}^\bullet(F^+)$  is a lens, and is therefore non-empty.

Up to the isomorphism  $\mathbb{P}_p^\bullet X \cong \mathbf{V}_\bullet X$ , Lemma 3.28 of [14] states that for every  $\nu' \in \mathbf{V}_\bullet X$ , if  $\nu' \in s_{\text{DP}}^\bullet(F^-)$ , then there is a  $\nu \in \mathbf{V}_\bullet X$  such that  $\nu \leq \nu'$  and for every  $h \in \mathcal{L}X$ ,  $F^-(h) \leq \int_{x \in X} h(x) d\nu \leq F^+(h)$ . In other words,  $Q \subseteq \uparrow(Q \cap C)$ .

Similarly, Lemma 3.29 of [14] states that for every  $\nu' \in \mathbf{V}_\bullet X$  such that  $\nu' \in s_{\text{AP}}^\bullet(F^+)$ , there is a  $\nu \in \mathbf{V}_\bullet X$  such that  $\nu' \leq \nu$  and for every  $h \in \mathcal{L}X$ ,  $F^-(h) \leq \int_{x \in X} h(x) d\nu \leq F^+(h)$ . This requires  $\mathcal{L}X$  to be locally convex and to have an almost open addition map, and also  $X$  to be compact if  $\bullet$  is “1”. In other words,  $C \subseteq \downarrow(Q \cap C)$ . Since the downwards-closure of a set is always included in its closure,  $C \subseteq \text{cl}(Q \cap C)$ . In particular, for every open neighborhood  $U$  of  $Q$ ,  $C \subseteq \text{cl}(U \cap C)$ .  $\square$

**Proposition 9.2.** *Let  $\bullet$  be nothing, “ $\leq 1$ ”, or “1”. For every topological space  $X$  such that  $\mathcal{L}X$  is locally convex and has an almost open addition map (and such that  $X$  is compact, in case  $\bullet$  is “1”),  $r_{\text{ADP}}$  and  $s_{\text{ADP}}^\bullet$  restrict to*

a homeomorphism between the subspace  $\mathcal{P}\ell_V^{q,cvx}\mathbf{V}\bullet X$  of  $\mathcal{P}\ell_V^q\mathbf{V}\bullet X$  consisting of quasi-lenses  $(Q, C)$  such that both  $Q$  and  $C$  are convex, and  $\mathbb{P}_{\text{ADP}}^\bullet X$ .

PROOF. Immediate consequence of Proposition 9.1, of the fact that  $r_{\text{DP}}$  and  $s_{\text{DP}}^\bullet$  form a homeomorphism between  $\mathcal{Q}_V^{cvx}\mathbf{V}\bullet X$  and  $\mathbb{P}_{\text{DP}}^\bullet X$ , and of the fact that  $r_{\text{AP}}$  and  $s_{\text{AP}}^\bullet$  form a homeomorphism between  $\mathcal{H}_V^{cvx}\mathbf{V}\bullet X$  and  $\mathbb{P}_{\text{AP}}^\bullet X$ .  $\square$

Much like there are projection maps  $\pi_1: \mathcal{P}\ell_V^q X \rightarrow \mathcal{Q}_V X$  and  $\pi_2: \mathcal{P}\ell_V^q X \rightarrow \mathcal{H}_V X$ , there are projection maps, which we write with the same notation,  $\pi_1: \mathbb{P}_{\text{ADP}}^\bullet X \rightarrow \mathbb{P}_{\text{DP}}^\bullet X$  and  $\pi_2: \mathbb{P}_{\text{ADP}}^\bullet X \rightarrow \mathbb{P}_{\text{AP}}^\bullet X$ . The pairing  $\langle \pi_1, \pi_2 \rangle$  is then the inclusion map  $\mathbb{P}_{\text{ADP}}^\bullet X \rightarrow \mathbb{P}_{\text{DP}}^\bullet X \times \mathbb{P}_{\text{AP}}^\bullet X$ . It is easy to see that this is a topological embedding.

It was shown in [11, Proposition 3] that there is a monad of forks on the category of dpos (resp., pointed dpos) and Scott-continuous maps, except that spaces of previsions were given the Scott topology of the pointwise ordering. As for previsions, we check that there is a corresponding monad of (subnormalized, normalized) forks on  $\mathbf{Top}$ .

**Proposition 9.3.** *Let  $\bullet$  be nothing, “ $\leq 1$ ”, or “1”. There is a monad  $(\mathbb{P}_{\text{ADP}}^\bullet, \eta^{\text{ADP}}, \mu^{\text{ADP}})$  on  $\mathbf{Top}$  (and, by restriction, on  $\mathbf{Top}^{\text{h}}$ ) whose unit and multiplication are defined by:*

$$\begin{aligned} \eta_X^{\text{ADP}}(x) &\stackrel{\text{def}}{=} (\eta_X^{\text{DP}}(x), \eta_X^{\text{AP}}(x)) \\ \mu_X^{\text{ADP}}(\mathcal{F}^-, \mathcal{F}^+) &\stackrel{\text{def}}{=} (h \in \mathcal{L}X \mapsto \mathcal{F}^-((F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X \mapsto F^-(h)), \\ &\quad h \in \mathcal{L}X \mapsto \mathcal{F}^+((F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X \mapsto F^+(h))) \end{aligned}$$

for every  $x \in X$  in the first case, and for every  $(\mathcal{F}^-, \mathcal{F}^+) \in \mathbb{P}_{\text{ADP}}^\bullet \mathbb{P}_{\text{ADP}}^\bullet X$  in the second case. In other words, unit and multiplication are defined uniquely by the commutativity of the following diagrams:

$$\begin{array}{ccccc} X & \xlongequal{\quad} & X & \xlongequal{\quad} & X \\ \eta_X^{\text{DP}} \downarrow & & \eta_X^{\text{ADP}} \downarrow & & \eta_X^{\text{AP}} \downarrow \\ \mathbb{P}_{\text{DP}}^\bullet X & \xleftarrow{\pi_1} & \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\pi_2} & \mathbb{P}_{\text{AP}}^\bullet X \end{array} \quad (12)$$

and:

$$\begin{array}{ccccc} \mathbb{P}_{\text{DP}}^\bullet \mathbb{P}_{\text{DP}}^\bullet X & \xleftarrow{\mathbb{P}_{\text{DP}}^\bullet \pi_1} & \mathbb{P}_{\text{DP}}^\bullet \mathbb{P}_{\text{ADP}}^\bullet X & \xleftarrow{\pi_1} & \mathbb{P}_{\text{ADP}}^\bullet \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\pi_2} & \mathbb{P}_{\text{AP}}^\bullet \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\mathbb{P}_{\text{AP}}^\bullet \pi_2} & \mathbb{P}_{\text{AP}}^\bullet \mathbb{P}_{\text{AP}}^\bullet X \\ \mu_X^{\text{DP}} \downarrow & & \mu_X^{\text{ADP}} \downarrow & & \mu_X^{\text{ADP}} \downarrow & & \mu_X^{\text{AP}} \downarrow & & \mu_X^{\text{AP}} \downarrow \\ \mathbb{P}_{\text{DP}}^\bullet X & \xleftarrow{\pi_1} & \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\pi_2} & \mathbb{P}_{\text{AP}}^\bullet X & & \mathbb{P}_{\text{AP}}^\bullet X & & \mathbb{P}_{\text{AP}}^\bullet X \end{array} \quad (13)$$

We recall that the functor part acts on morphisms by  $\mathbb{P}_{\text{ADP}}^\bullet f = (\mathbb{P}f, \mathbb{P}f)$ , hence is characterized by the commutativity of the following diagram:

$$\begin{array}{ccccc}
\mathbb{P}_{\text{DP}}^\bullet X & \xleftarrow{\pi_1} & \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\pi_2} & \mathbb{P}_{\text{AP}}^\bullet X \\
\mathbb{P}f \downarrow & & \mathbb{P}_{\text{ADP}}^\bullet f \downarrow & & \downarrow \mathbb{P}f \\
\mathbb{P}_{\text{DP}}^\bullet Y & \xleftarrow{\pi_1} & \mathbb{P}_{\text{ADP}}^\bullet Y & \xrightarrow{\pi_2} & \mathbb{P}_{\text{AP}}^\bullet Y
\end{array} \tag{14}$$

## 10. The weak distributive law

**Proposition 10.1.** *Let  $X$  be a topological space. There is a unique continuous map  $\Theta: \mathbf{V}\bullet\mathcal{P}\ell_{\mathbf{V}}^q X \rightarrow \mathbb{P}_{\text{ADP}}^\bullet X$  such that:*

$$\begin{array}{ccccc}
\mathbf{V}\bullet\mathcal{Q}_{\mathbf{V}} X & \xleftarrow{\mathbf{V}\bullet\pi_1} & \mathbf{V}\bullet\mathcal{P}\ell_{\mathbf{V}}^q X & \xrightarrow{\mathbf{V}\bullet\pi_2} & \mathbf{V}\bullet\mathcal{H}_{\mathbf{V}} X \\
\Phi \downarrow & & \downarrow \Theta & & \downarrow \Psi \\
\mathbb{P}_{\text{DP}}^\bullet X & \xleftarrow{\pi_1} & \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\pi_2} & \mathbb{P}_{\text{AP}}^\bullet X.
\end{array}$$

PROOF. In other words, for every  $\xi \in \mathbf{V}\bullet\mathcal{P}\ell_{\mathbf{V}}^q X$ ,  $\Theta(\xi)$  must be equal to  $(\Phi(\pi_1[\xi]), \Psi(\pi_2[\xi]))$ . It remains to show that the latter is a fork; continuity will follow from the continuity of  $\Phi$  and of  $\Psi$  (for the latter, see Lemma 7.1). Let  $(F^-, F^+) \stackrel{\text{def}}{=} (\Phi(\pi_1[\xi]), \Psi(\pi_2[\xi]))$ . We know that  $F^- \in \mathbb{P}_{\text{DP}}^\bullet X$ , that  $F^+ \in \mathbb{P}_{\text{AP}}^\bullet X$ , and it remains to show Walley's conditions.

For every  $h \in \mathcal{L}X$ , we have:

$$\begin{aligned}
F^-(h) &= \int_{Q \in \mathcal{Q}_{\mathbf{V}} X} \min_{x \in Q} h(x) d\pi_1[\xi] \\
&= \int_{(Q,C) \in \mathcal{P}\ell_{\mathbf{V}}^q X} \min_{x \in Q} h(x) d\xi
\end{aligned} \tag{15}$$

$$\begin{aligned}
F^+(h) &= \int_{C \in \mathcal{H}_{\mathbf{V}} X} \sup_{x \in C} h(x) d\pi_2[\xi] \\
&= \int_{(Q,C) \in \mathcal{P}\ell_{\mathbf{V}}^q X} \sup_{x \in C} h(x) d\xi,
\end{aligned} \tag{16}$$

using the change of variable formula.

For every  $h \in \mathcal{L}X$ , we have:

$$\begin{aligned}
\min_{x \in Q} h(x) &\geq \min_{x \in \uparrow(Q \cap C)} h(x) && \text{since } Q \subseteq \uparrow(Q \cap C) \\
&= \min_{x \in Q \cap C} h(x) \geq \min_{x \in Q} h(x),
\end{aligned}$$

since  $Q \cap C \subseteq C$ . This is a circular list of inequalities, hence all its terms are equal.

For every  $h, h' \in \mathcal{L}X$ , for all  $(Q, C) \in \mathcal{P}\ell_{\mathbf{V}}^{\mathfrak{q}}X$ , we then have:

$$\begin{aligned}
\min_{x \in Q} (h(x) + h'(x)) &= \min_{x \in Q \cap C} (h(x) + h'(x)) \\
&\leq \min_{x \in Q \cap C} (h(x) + \sup_{x \in C} h'(x)) \\
&= \min_{x \in Q \cap C} h(x) + \sup_{x \in C} h'(x) \\
&= \min_{x \in Q} h(x) + \sup_{x \in C} h'(x). \tag{17}
\end{aligned}$$

Using (15) and (16), it follows that  $F^-(h + h') \leq F^-(h) + F^+(h')$ .

We claim that:

$$\min_{x \in Q} h(x) + \sup_{x \in C} h'(x) \leq \sup_{x \in C} (h(x) + h'(x)). \tag{18}$$

It suffices to show that for every  $t \in \mathbb{R}_+$  such that  $t < \min_{x \in Q} h(x)$ ,  $t + \sup_{x \in C} h'(x) \leq \sup_{x \in C} (h(x) + h'(x))$ . Since  $t < \min_{x \in Q} h(x)$ ,  $Q \subseteq h^{-1}(]t, \infty])$ . Since  $(Q, C)$  is a quasi-lens,  $C \subseteq cl(h^{-1}(]t, \infty]) \cap C$ . Hence  $\sup_{x \in C} h'(x) \leq \sup_{x \in cl(h^{-1}(]t, \infty]) \cap C} h'(x) = \sup_{x \in h^{-1}(]t, \infty]) \cap C} h'(x)$  (using Fact 7.5), and since  $h(x) \geq t$  for every  $x \in h^{-1}(]t, \infty]) \cap C$ , we obtain that  $t + \sup_{x \in C} h'(x) \leq \sup_{x \in h^{-1}(]t, \infty]) \cap C} (h(x) + h'(x)) \leq \sup_{x \in C} (h(x) + h'(x))$ .

Using (15) and (16), it follows from (18) that  $F^-(h) + F^+(h') \leq F^+(h + h')$ .  $\square$

**Theorem 10.2.** *Let  $\bullet$  be nothing, “ $\leq 1$ ”, or “1”. For every space  $X$  such that  $\mathcal{L}X$  is locally convex and has an almost open addition map (and such that  $X$  is compact if  $\bullet$  is “1”), there is a unique continuous map  $\lambda_X^{\natural} : \mathbf{V}\bullet\mathcal{P}\ell_{\mathbf{V}}^{\mathfrak{q}}X \rightarrow \mathcal{P}\ell_{\mathbf{V}}^{\mathfrak{q}}\mathbf{V}\bullet X$  such that:*

$$\begin{array}{ccc}
\mathbf{V}\bullet\mathcal{Q}_{\mathbf{V}}X & \xleftarrow{\mathbf{V}\bullet\pi_1} \mathbf{V}\bullet\mathcal{P}\ell_{\mathbf{V}}^{\mathfrak{q}}X & \xrightarrow{\mathbf{V}\bullet\pi_2} \mathbf{V}\bullet\mathcal{H}_{\mathbf{V}}X \\
\lambda_X^{\natural} \downarrow & & \downarrow \lambda_X^{\natural} \\
\mathcal{Q}_{\mathbf{V}}\mathbf{V}\bullet X & \xleftarrow{\pi_1} \mathcal{P}\ell_{\mathbf{V}}^{\mathfrak{q}}\mathbf{V}\bullet X & \xrightarrow{\pi_2} \mathcal{H}_{\mathbf{V}}\mathbf{V}\bullet X
\end{array} \tag{19}$$

and that is  $s_{\text{ADP}}^{\bullet} \circ \Theta$ .

The collection  $\lambda^{\natural}$  of maps  $\lambda_X^{\natural}$  is a weak distributive law of  $\mathcal{P}\ell_{\mathbf{V}}^{\mathfrak{q}}$  over  $\mathbf{V}\bullet$  on  $\mathbf{Top}^{\natural}$ .

PROOF. Uniqueness is because  $\langle \pi_1, \pi_2 \rangle$  is the inclusion map of  $\mathcal{P}\ell_V^q \mathbf{V}\bullet X$ , which is injective. Its existence as  $s_{\text{ADP}}^\bullet \circ \Theta$  follows from Proposition 9.1 and Proposition 10.1. The fact that  $\lambda^\natural$  is natural and satisfies the three equations of weak distributive laws follows from the similar properties of  $\lambda^\sharp$  (Theorem 4.13) and  $\lambda^b$  (Theorem 7.11). Given any morphism  $f: X \rightarrow Y$  in  $\mathbf{Top}^\natural$  (and noticing that the requirements on  $\mathbf{Top}^\natural$  imply those on  $\mathbf{Top}^b$ , which enables us to reuse all our results on  $\lambda^b$ ), we have the following diagram:

$$\begin{array}{ccccc}
\mathbf{V}\bullet Q_V X & \xleftarrow{\mathbf{V}\bullet\pi_1} & \mathbf{V}\bullet \mathcal{P}\ell_V^q X & \xrightarrow{\mathbf{V}\bullet\pi_2} & \mathbf{V}\bullet \mathcal{H}_V X \\
\lambda_X^\sharp \downarrow & & \lambda_X^\natural \downarrow & & \lambda_X^b \downarrow \\
Q_V \mathbf{V}\bullet X & \xleftarrow{\pi_1} & \mathcal{P}\ell_V^q \mathbf{V}\bullet X & \xrightarrow{\pi_2} & \mathcal{H}_V \mathbf{V}\bullet X \\
\downarrow Q_V \mathbf{V}\bullet f & & \downarrow \mathcal{P}\ell_V^q \mathbf{V}\bullet f & & \downarrow \mathcal{H}_V \mathbf{V}\bullet f \\
\mathbf{V}\bullet Q_V Y & \xleftarrow{\mathbf{V}\bullet\pi_1} & \mathbf{V}\bullet \mathcal{P}\ell_V^q Y & \xrightarrow{\mathbf{V}\bullet\pi_2} & \mathbf{V}\bullet \mathcal{H}_V Y \\
\lambda_Y^\sharp \downarrow & & \lambda_Y^\natural \downarrow & & \lambda_Y^b \downarrow \\
Q_V \mathbf{V}\bullet Y & \xleftarrow{\pi_1} & \mathcal{P}\ell_V^q \mathbf{V}\bullet Y & \xrightarrow{\pi_2} & \mathcal{H}_V \mathbf{V}\bullet Y
\end{array}$$

The two squares on the  $X$  (back) face commute by definition of  $\lambda_X^\natural$ , and similarly for the two squares on the  $Y$  (front) face. The vertical, slanted rectangles on the left and on the right commute by naturality of  $\lambda^\sharp$ , resp. of  $\lambda^b$ . The four horizontal (slanted) rectangles commute by naturality of  $\pi_1$  and of  $\pi_2$ . The naturality of  $\lambda^\natural$  is the commutativity of the middle vertical, slanted rectangle, which follows from the fact that all the other rectangles commute, and from the fact that  $\langle \pi_1, \pi_2 \rangle$  is injective.

The first law of weak distributivity laws,  $\lambda_X^\natural \circ \mathbf{V}\bullet\eta_X^q = \eta_{\mathbf{V}\bullet X}^q$ , is proved similarly. The following diagram commutes:

$$\begin{array}{ccccc}
\mathbf{V}\bullet X & \xlongequal{\quad} & \mathbf{V}\bullet X & \xlongequal{\quad} & \mathbf{V}\bullet X \\
\mathbf{v}\bullet\eta_X^q \downarrow & & \mathbf{v}\bullet\eta_X^q \downarrow & & \mathbf{v}\bullet\eta_X^q \downarrow \\
\mathbf{V}\bullet Q_V X & \xleftarrow{\mathbf{V}\bullet\pi_1} & \mathbf{V}\bullet \mathcal{P}\ell_V^q X & \xrightarrow{\mathbf{V}\bullet\pi_2} & \mathbf{V}\bullet \mathcal{H}_V X \\
\lambda_X^\sharp \downarrow & & \lambda_X^\natural \downarrow & & \lambda_X^b \downarrow \\
Q_V \mathbf{V}\bullet X & \xleftarrow{\pi_1} & \mathcal{P}\ell_V^q \mathbf{V}\bullet X & \xrightarrow{\pi_2} & \mathcal{H}_V \mathbf{V}\bullet X
\end{array}$$

Indeed, the top two squares commute by definition of  $\eta^q$  (see (10), Proposition 8.5), and the bottom two squares commute by definition of  $\lambda^\natural$ . The

composition of the leftmost two vertical arrows is  $\eta_{\mathbf{V}\bullet X}^{\mathcal{Q}}$ , and the composition of the rightmost two vertical arrows is  $\eta_{\mathbf{V}\bullet X}^{\mathcal{H}}$ , by the first equation of weak distributive laws for  $\lambda^{\sharp}$  and  $\lambda^{\flat}$  respectively. Hence the composition of the middle two vertical arrows is  $\eta_{\mathbf{V}\bullet X}^{\mathcal{Q}}$ , by the uniqueness of  $\eta^{\mathcal{Q}}$  in (10).

We turn to the second weak distributivity law  $\lambda_X^{\sharp} \circ \mathbf{V}\cdot\mu_X^{\mathcal{Q}} = \mu_{\mathbf{V}\bullet X}^{\mathcal{Q}} \circ \mathcal{P}\ell_V^{\mathcal{Q}}\lambda_X^{\sharp} \circ \lambda_{\mathcal{P}\ell_V^{\mathcal{Q}}X}^{\sharp}$ . The left-hand side is composition of the middle vertical arrows in the following diagram:

$$\begin{array}{ccccc}
\mathbf{V}\cdot\mathcal{Q}_V\mathcal{Q}_V X \xleftarrow{\mathbf{V}\cdot\mathcal{Q}_V\pi_1} \mathbf{V}\cdot\mathcal{Q}_V\mathcal{P}\ell_V^{\mathcal{Q}}X \xleftarrow{\mathbf{V}\cdot\pi_1} \mathbf{V}\cdot\mathcal{P}\ell_V^{\mathcal{Q}}\mathcal{P}\ell_V^{\mathcal{Q}}X \xrightarrow{\mathbf{V}\cdot\pi_2} \mathbf{V}\cdot\mathcal{H}_V\mathcal{P}\ell_V^{\mathcal{Q}}X \xrightarrow{\mathbf{V}\cdot\mathcal{H}_V\pi_2} \mathbf{V}\cdot\mathcal{H}_V\mathcal{H}_V X & & & & \\
\mathbf{V}\cdot\mu_X^{\mathcal{Q}} \downarrow & & \mathbf{V}\cdot\mu_X^{\mathcal{Q}} \downarrow & & \mathbf{V}\cdot\mu_X^{\mathcal{H}} \downarrow \\
\mathbf{V}\cdot\mathcal{Q}_V X \xleftarrow{\mathbf{V}\cdot\pi_1} \mathbf{V}\cdot\mathcal{P}\ell_V^{\mathcal{Q}}X \xrightarrow{\mathbf{V}\cdot\pi_2} \mathbf{V}\cdot\mathcal{H}_V X & & & & \\
\lambda_X^{\sharp} \downarrow & & \lambda_X^{\flat} \downarrow & & \lambda_X^{\flat} \downarrow \\
\mathcal{Q}_V\mathbf{V}\bullet X \xleftarrow{\pi_1} \mathcal{P}\ell_V^{\mathcal{Q}}\mathbf{V}\bullet X \xrightarrow{\pi_2} \mathcal{H}_V\mathbf{V}\bullet X & & & & 
\end{array}$$

where the top two squares are by (11), and the bottom two squares are by definition of  $\lambda^{\sharp}$ . The right-hand side appears as the composition of middle vertical arrows in the following diagram:

$$\begin{array}{ccccc}
\mathbf{V}\cdot\mathcal{Q}_V\mathcal{Q}_V X \xleftarrow{\mathbf{V}\cdot\mathcal{Q}_V\pi_1} \mathbf{V}\cdot\mathcal{Q}_V\mathcal{P}\ell_V^{\mathcal{Q}}X \xleftarrow{\mathbf{V}\cdot\pi_1} \mathbf{V}\cdot\mathcal{P}\ell_V^{\mathcal{Q}}\mathcal{P}\ell_V^{\mathcal{Q}}X \xrightarrow{\mathbf{V}\cdot\pi_2} \mathbf{V}\cdot\mathcal{H}_V\mathcal{P}\ell_V^{\mathcal{Q}}X \xrightarrow{\mathbf{V}\cdot\mathcal{H}_V\pi_2} \mathbf{V}\cdot\mathcal{H}_V\mathcal{H}_V X & & & & \\
\lambda_{\mathcal{Q}_V X}^{\sharp} \downarrow & \text{nat. } \lambda^{\sharp} & \lambda_{\mathcal{P}\ell_V^{\mathcal{Q}} X}^{\sharp} \downarrow & \text{def. } \lambda^{\sharp} & \lambda_{\mathcal{P}\ell_V^{\mathcal{Q}} X}^{\flat} \downarrow & \text{def. } \lambda^{\flat} & \lambda_{\mathcal{P}\ell_V^{\mathcal{Q}} X}^{\flat} \downarrow & \text{nat. } \lambda^{\flat} & \lambda_{\mathcal{H}_V X}^{\flat} \downarrow \\
\mathcal{Q}_V\mathbf{V}\bullet\mathcal{P}\ell_V^{\mathcal{Q}}X \xleftarrow{\pi_1} \mathcal{P}\ell_V^{\mathcal{Q}}\mathbf{V}\bullet\mathcal{P}\ell_V^{\mathcal{Q}}X \xrightarrow{\pi_2} \mathcal{H}_V\mathbf{V}\bullet\mathcal{P}\ell_V^{\mathcal{Q}}X & & & & & & & & \\
\mathcal{Q}_V\mathbf{V}\cdot\pi_1 \swarrow & \text{nat. } \pi_1 & \mathcal{P}\ell_V^{\mathcal{Q}}\mathbf{V}\cdot\pi_1 \swarrow & & \mathcal{P}\ell_V^{\mathcal{Q}}\mathbf{V}\cdot\pi_2 \searrow & \text{nat. } \pi_2 & \mathcal{H}_V\mathbf{V}\cdot\pi_1 \searrow & & \\
\mathcal{Q}_V\mathbf{V}\bullet\mathcal{Q}_V X \xleftarrow{\pi_1} \mathcal{P}\ell_V^{\mathcal{Q}}\mathbf{V}\bullet\mathcal{Q}_V X & & \mathcal{P}\ell_V^{\mathcal{Q}}\lambda_X^{\sharp} \downarrow & & \mathcal{P}\ell_V^{\mathcal{Q}}\mathbf{V}\bullet\mathcal{H}_V X \xrightarrow{\pi_2} \mathcal{H}_V\mathbf{V}\bullet\mathcal{H}_V X & & & & \\
\mathcal{Q}_V\lambda_X^{\sharp} \downarrow & \text{nat. } \pi_1 & \mathcal{P}\ell_V^{\mathcal{Q}}\lambda_X^{\sharp} \downarrow & \text{def. } \lambda^{\sharp} & \text{def. } \lambda^{\flat} & \mathcal{P}\ell_V^{\mathcal{Q}}\lambda_X^{\flat} \downarrow & \text{nat. } \pi_2 & \mathcal{H}_V\lambda_X^{\flat} \downarrow & \\
\mathcal{Q}_V\mathcal{Q}_V\mathbf{V}\bullet X \xleftarrow{\pi_1} \mathcal{P}\ell_V^{\mathcal{Q}}\mathcal{Q}_V\mathbf{V}\bullet X \xleftarrow{\mathcal{P}\ell_V^{\mathcal{Q}}\pi_1} \mathcal{P}\ell_V^{\mathcal{Q}}\mathcal{P}\ell_V^{\mathcal{Q}}\mathbf{V}\bullet X \xrightarrow{\mathcal{P}\ell_V^{\mathcal{Q}}\pi_2} \mathcal{P}\ell_V^{\mathcal{Q}}\mathcal{H}_V\mathbf{V}\bullet X \xrightarrow{\pi_2} \mathcal{H}_V\mathcal{H}_V\mathbf{V}\bullet X & & & & & & & & \\
\mu_X^{\mathcal{Q}} \downarrow & & \mu_{\mathbf{V}\bullet X}^{\mathcal{Q}} \downarrow & & \mu_X^{\mathcal{H}} \downarrow & & & & \\
\mathcal{Q}_V\mathbf{V}\bullet X \xleftarrow{\pi_1} \mathcal{P}\ell_V^{\mathcal{Q}}\mathbf{V}\bullet X \xrightarrow{\pi_2} \mathcal{H}_V\mathbf{V}\bullet X & & & & & & & & 
\end{array}$$

(11) (11)

Those two diagrams have the same (composition of) leftmost vertical arrows, and similarly for the rightmost vertical arrows, because  $\lambda^{\sharp}$  and  $\lambda^{\flat}$  are weak distributivity laws. Since  $\langle \pi_1, \pi_2 \rangle$  is injective, the (composition of) middle vertical arrows are the same, too.



Finally, we address the third weak distributivity law  $\lambda_X^\natural \circ \mu_{\mathcal{P}\ell_V^q X} = \mathcal{P}\ell_V^q \mu_X \circ \lambda_{\mathbf{V}\bullet X}^\natural$ . As above, this consists in checking that the leftmost compositions of vertical arrows in the following two diagrams match, so that the middle compositions of vertical arrows coincide, too.

$$\begin{array}{ccccc}
\mathbf{V}\bullet\mathbf{V}\bullet Q_V X & \xleftarrow{\mathbf{V}\bullet\mathbf{V}\bullet\pi_1} & \mathbf{V}\bullet\mathbf{V}\bullet \mathcal{P}\ell_V^q X & \xrightarrow{\mathbf{V}\bullet\mathbf{V}\bullet\pi_2} & \mathbf{V}\bullet\mathbf{V}\bullet HVX \\
\mu_{Q_V X} \downarrow & \text{nat. } \mu & \mu_{\mathcal{P}\ell_V^q X} \downarrow & \text{nat. } \mu & \downarrow \mu_{HVX} \\
\mathbf{V}\bullet Q_V X & \xleftarrow{\mathbf{V}\bullet\pi_1} & \mathbf{V}\bullet \mathcal{P}\ell_V^q X & \xrightarrow{\mathbf{V}\bullet\pi_2} & \mathbf{V}\bullet \mathcal{H}_V X \\
\lambda_X^\natural \downarrow & \text{def. } \lambda^\natural & \lambda_X^\natural \downarrow & \text{def. } \lambda^\natural & \downarrow \lambda_X^\natural \\
Q_V \mathbf{V}\bullet X & \xleftarrow{\pi_1} & \mathcal{P}\ell_V^q \mathbf{V}\bullet X & \xrightarrow{\pi_2} & \mathcal{H}_V \mathbf{V}\bullet X
\end{array}$$

Here is the second diagram:

$$\begin{array}{ccccc}
\mathbf{V}\bullet\mathbf{V}\bullet Q_V X & \xleftarrow{\mathbf{V}\bullet\mathbf{V}\bullet\pi_1} & \mathbf{V}\bullet\mathbf{V}\bullet \mathcal{P}\ell_V^q X & \xrightarrow{\mathbf{V}\bullet\mathbf{V}\bullet\pi_2} & \mathbf{V}\bullet\mathbf{V}\bullet HVX \\
\mathbf{V}\bullet\lambda_X^\natural \downarrow & \text{def. } \lambda^\natural & \mathbf{V}\bullet\lambda_X^\natural \downarrow & \text{def. } \lambda^\natural & \downarrow \mathbf{V}\bullet\lambda_X^\natural \\
\mathbf{V}\bullet Q_V \mathbf{V}\bullet X & \xleftarrow{\mathbf{V}\bullet\pi_1} & \mathbf{V}\bullet \mathcal{P}\ell_V^q \mathbf{V}\bullet X & \xrightarrow{\mathbf{V}\bullet\pi_2} & \mathbf{V}\bullet \mathcal{H}_V \mathbf{V}\bullet X \\
\lambda_X^\natural \downarrow & \text{def. } \lambda^\natural & \lambda_X^\natural \downarrow & \text{def. } \lambda^\natural & \downarrow \lambda_X^\natural \\
Q_V \mathbf{V}\bullet\mathbf{V}\bullet X & \xleftarrow{\pi_1} & \mathcal{P}\ell_V^q \mathbf{V}\bullet\mathbf{V}\bullet X & \xrightarrow{\pi_2} & \mathcal{H}_V \mathbf{V}\bullet\mathbf{V}\bullet X \\
Q_V \mu_X \downarrow & \text{nat. } \pi_1 & \mathcal{P}\ell_V^q \mu_X \downarrow & \text{nat. } \pi_2 & \downarrow \mathcal{H}_V \mu_X \\
Q_V \mathbf{V}\bullet X & \xleftarrow{\pi_1} & \mathcal{P}\ell_V^q \mathbf{V}\bullet X & \xrightarrow{\pi_2} & \mathcal{H}_V \mathbf{V}\bullet X
\end{array}$$

□

**Remark 10.3.** Using Remark 4.3 and Remark 7.4, in the special case where  $\mu \stackrel{\text{def}}{=} \sum_{i=1}^n a_i \delta_{(Q_i, C_i)}$ , where  $n \geq 1$ , and where  $Q_i \stackrel{\text{def}}{=} \uparrow E_i$ ,  $C_i \stackrel{\text{def}}{=} \downarrow E_i$ , where each set  $E_i$  is finite and non-empty, then  $\lambda_X^\natural(\mu) = (\uparrow \mathcal{E}, \downarrow \mathcal{E})$ , where  $\mathcal{E} \stackrel{\text{def}}{=} \{\sum_{i=1}^n a_i \delta_{x_i} \mid x_1 \in E_1, \dots, x_n \in E_n\}$ .

**Remark 10.4.** We recall that if  $X$  is weakly Hausdorff or quasi-Polish, then  $\mathcal{P}\ell_V^q X \cong \mathcal{P}\ell_V X$ , and that those isomorphisms allow us to transport the  $\mathcal{P}\ell_V^q$  monad structure over to any full subcategory of **Top** consisting of weakly Hausdorff or quasi-Polish spaces. Theorem 10.2 then allows us to obtain a weak distributive law of  $\mathcal{P}\ell_V$  over  $\mathbf{V}\bullet$  on any full subcategory of **Top**<sup>h</sup> consisting of weakly Hausdorff or quasi-Polish spaces—in particular on the category of stably compact spaces, which are all weakly Hausdorff.

**Remark 10.5.** Combining Remark 10.3 and Remark 10.4, in the special case where  $\mu \stackrel{\text{def}}{=} \sum_{i=1}^n a_i \delta_{L_i}$ , where  $n \geq 1$ , and where  $L_i \stackrel{\text{def}}{=} \uparrow E_i \cap \downarrow E_i$ , where each set  $E_i$  is finite and non-empty, then  $\lambda_X^{\natural}(\mu) = \uparrow \mathcal{E} \cap \downarrow \mathcal{E}$ , where  $\mathcal{E} \stackrel{\text{def}}{=} \{\sum_{i=1}^n a_i \delta_{x_i} \mid x_1 \in E_1, \dots, x_n \in E_n\}$ .

## 11. The associated weak composite monad

We proceed as with  $\mathbb{P}_{\text{DP}}^{\bullet}$  and  $\mathbb{P}_{\text{ADP}}^{\bullet}$ , and in fact we reduce to the cases of those monads. We start with the idempotent:

$$\mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\mathbf{V}_{\bullet}X \xrightarrow{\eta_{\mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\mathbf{V}_{\bullet}X}} \mathbf{V}_{\bullet}\mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\mathbf{V}_{\bullet}X \xrightarrow{\lambda_{\mathbf{V}_{\bullet}X}^{\natural}} \mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\mathbf{V}_{\bullet}\mathbf{V}_{\bullet}X \xrightarrow{\mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\mu_X} \mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\mathbf{V}_{\bullet}X \quad (20)$$

whose splitting will define the action of the functor part of the associated weak composite monad on  $X$ .

**Lemma 11.1.** A splitting of (20) is  $\mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\mathbf{V}_{\bullet}X \xrightarrow{r_{\text{ADP}}} \mathbb{P}_{\text{ADP}}^{\bullet}X \xrightarrow{s_{\text{ADP}}^{\bullet}} \mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\mathbf{V}_{\bullet}X$ .

PROOF. Composing (20) with  $\pi_1$ , resp. with  $\pi_2$  on the right (of the diagram—namely computing  $\pi_1 \circ f$  resp.  $\pi_2 \circ f$  where  $f$  is (20)), and simplifying using the naturality of  $\pi_1$  (resp.,  $\pi_2$ ), (19) (see Theorem 10.2) and (12) (see Proposition 9.3), we obtain (2), resp. (6)—the corresponding idempotents in the DP and AP cases—composed with  $\pi_1$ , resp.  $\pi_2$ .

By Lemma 5.1 (resp., Lemma 8.1), this is equal to  $s_{\text{DP}}^{\bullet} \circ r_{\text{DP}} \circ \pi_1$  (resp.,  $s_{\text{AP}}^{\bullet} \circ r_{\text{AP}} \circ \pi_2$ ), and that, in turn, is equal to  $\pi_1 \circ s_{\text{ADP}}^{\bullet} \circ r_{\text{ADP}}$  (resp.,  $\pi_2 \circ s_{\text{ADP}}^{\bullet} \circ r_{\text{ADP}}$ ), by Proposition 9.1. We conclude since  $\langle \pi_1, \pi_2 \rangle$  is an inclusion map, hence is monotonic.  $\square$

Generalizing (20), there is an idempotent:

$$\mathcal{P}\ell_{\mathbf{V}}^{\text{q}}X \xrightarrow{\eta_{\mathcal{P}\ell_{\mathbf{V}}^{\text{q}}X}} \mathbf{V}_{\bullet}\mathcal{P}\ell_{\mathbf{V}}^{\text{q}}X \xrightarrow{\lambda_X^{\natural}} \mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\mathbf{V}_{\bullet}X \xrightarrow{\mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\alpha} \mathcal{P}\ell_{\mathbf{V}}^{\text{q}}X \quad (21)$$

for every  $\mathbf{V}_{\bullet}$ -algebra  $\alpha$ . As before, we will only need to compute this splitting for one additional  $\mathbf{V}_{\bullet}$ -algebra. We write  $\mathcal{P}\ell_{\mathbf{V}}^{\text{q} \text{ convex}}\mathbb{P}_{\text{ADP}}^{\bullet}X$  for the subspace of  $\mathcal{P}\ell_{\mathbf{V}}^{\text{q}}\mathbb{P}_{\text{ADP}}^{\bullet}X$  consisting of *convex* quasi-lenses, namely quasi-lenses  $(\mathcal{Q}, \mathcal{C})$  such that both  $\mathcal{Q}$  and  $\mathcal{C}$  are convex.

**Lemma 11.2.** For every topological space  $X$ , there is a  $\mathbf{V}_\bullet$ -algebra  $\alpha_{\text{ADP}}^\bullet: \mathbf{V}_\bullet \mathbb{P}_{\text{ADP}}^\bullet X \rightarrow \mathbb{P}_{\text{ADP}}^\bullet X$  defined by  $\alpha_{\text{ADP}}^\bullet(\xi) \stackrel{\text{def}}{=} (h \in \mathcal{L}X \mapsto \alpha_{\text{DP}}^\bullet(\pi_1[\xi]), h \in \mathcal{L}X \mapsto \alpha_{\text{AP}}^\bullet(\pi_2[\xi]))$  for every  $\xi \in \mathbf{V}_\bullet \mathbb{P}_{\text{ADP}}^\bullet X$ .

If  $X$  is an object of  $\mathbf{Top}^{\text{h}}$ , then a splitting of the idempotent (21) where  $\alpha \stackrel{\text{def}}{=} \alpha_{\text{ADP}}^\bullet$  is:

$$\mathcal{P}\ell_{\mathbf{V}}^{\text{q}} \mathbb{P}_{\text{ADP}}^\bullet X \longrightarrow \mathcal{P}\ell_{\mathbf{V}}^{\text{q}, \text{cvx}} \mathbb{P}_{\text{ADP}}^\bullet X \longrightarrow \mathcal{P}\ell_{\mathbf{V}}^{\text{q}} \mathbb{P}_{\text{ADP}}^\bullet X,$$

where the arrow on the right is the inclusion map, and the arrow on the left is the corestriction of (21), and maps every  $(Q, C) \in \mathcal{P}\ell_{\mathbf{V}}^{\text{q}, \text{cvx}} \mathbb{P}_{\text{ADP}}^\bullet X$  to itself.

PROOF. For every  $\xi \in \mathbf{V}_\bullet \mathbb{P}_{\text{ADP}}^\bullet X$ , let  $(F_0^-, F_0^+) \stackrel{\text{def}}{=} \alpha_{\text{ADP}}^\bullet(\xi)$ . For every  $h \in \mathcal{L}X$ ,  $F_0^-(h) = \alpha_{\text{DP}}^\bullet(\pi_1[\xi])(h) = \int_{F \in \mathbb{P}_{\text{DP}}^\bullet X} F(h) d\pi_1[\xi] = \int_{(F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X} F^-(h) d\xi$ , by the change of variable formula, and similarly  $F_0^+(h) = \int_{(F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X} F^+(h) d\xi$ . Hence, for all  $h, h' \in \mathcal{L}X$ ,

$$\begin{aligned} F_0^-(h + h') &= \int_{(F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X} F^-(h + h') d\xi \\ &\leq \int_{(F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X} (F^-(h) + F^+(h')) d\xi \\ &= \int_{(F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X} F^-(h) d\xi + \int_{(F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X} F^+(h') d\xi \\ &= F_0^-(h) + F_0^+(h'), \end{aligned}$$

and:

$$\begin{aligned} F_0^-(h) + F_0^+(h') &= \int_{(F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X} (F^-(h) + F^+(h')) d\xi \\ &\leq \int_{(F^-, F^+) \in \mathbb{P}_{\text{ADP}}^\bullet X} F^+(h + h') d\xi \\ &= F_0^+(h + h'), \end{aligned}$$

so  $(F_0^-, F_0^+) \in \mathbb{P}_{\text{ADP}}^\bullet X$ . The map  $\alpha_{\text{ADP}}^\bullet$  is continuous, since the inverse image of the subbasic open set  $[h > r]^-$  is  $[((F^+, F^-) \mapsto F^-(h)) > r]$ , and the inverse image of the subbasic open set  $[h > r]^+$  is  $[((F^+, F^-) \mapsto F^+(h)) > r]$ .

The map  $\alpha_{\text{ADP}}^\bullet$  is defined so that  $\pi_1 \circ \alpha_{\text{ADP}}^\bullet = \alpha_{\text{DP}}^\bullet \circ \mathbf{V}_\bullet \pi_1$  and  $\pi_2 \circ \alpha_{\text{ADP}}^\bullet = \alpha_{\text{AP}}^\bullet \circ \mathbf{V}_\bullet \pi_2$ . The fact that  $\alpha_{\text{ADP}}^\bullet$  is a  $\mathbf{V}_\bullet$ -algebra then follows from the fact that  $\alpha_{\text{DP}}^\bullet$  and  $\alpha_{\text{AP}}^\bullet$  are.

The idempotent (21) with  $\alpha \stackrel{\text{def}}{=} \alpha_{\text{ADP}}^\bullet$  appears as the top row of the following diagram.

$$\begin{array}{ccccccc}
\mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\eta_{\mathcal{P}\ell_V^q X}} & \mathbf{V} \cdot \mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\lambda_{\mathbb{P}_{\text{ADP}}^\bullet X}^\sharp} & \mathcal{P}\ell_V^q \mathbf{V} \cdot \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\mathcal{P}\ell_V^q \alpha_{\text{ADP}}^\bullet} & \mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X \\
\downarrow \pi_1 & \text{nat. } \pi_1 & \downarrow \mathbf{V} \cdot \pi_1 & (19) & \downarrow \pi_1 & \text{nat. } \pi_1 & \downarrow \pi_1 \\
\mathcal{Q}_V \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\eta_{\mathcal{Q}_V \mathbb{P}_{\text{ADP}}^\bullet X}} & \mathbf{V} \cdot \mathcal{Q}_V \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\lambda_{\mathbb{P}_{\text{ADP}}^\bullet X}^\sharp} & \mathcal{Q}_V \mathbf{V} \cdot \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\mathcal{Q}_V \alpha_{\text{ADP}}^\bullet} & \mathcal{Q}_V \mathbb{P}_{\text{ADP}}^\bullet X \\
\downarrow \mathcal{Q}_V \pi_1 & \text{nat. } \eta & \downarrow \mathbf{V} \cdot \mathcal{Q}_V \pi_1 & \text{nat. } \lambda^\sharp & \downarrow \mathcal{Q}_V \mathbf{V} \cdot \pi_1 & \text{def. } \alpha_{\text{ADP}}^\bullet & \downarrow \mathcal{Q}_V \pi_1 \\
\mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X & \xrightarrow{\eta_{\mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X}} & \mathbf{V} \cdot \mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X & \xrightarrow{\lambda_{\mathbb{P}_{\text{DP}}^\bullet X}^\sharp} & \mathcal{Q}_V \mathbf{V} \cdot \mathbb{P}_{\text{DP}}^\bullet X & \xrightarrow{\mathcal{Q}_V \alpha_{\text{DP}}^\bullet} & \mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X
\end{array}$$

The bottom row is the idempotent (3) of Lemma 5.2 (with  $\alpha \stackrel{\text{def}}{=} \alpha_{\text{DP}}^\bullet$ ), which splits as  $\mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X \xrightarrow{r^\sharp} \mathcal{Q}_V^{cvx} \mathbb{P}_{\text{DP}}^\bullet X \xrightarrow{s^\sharp} \mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X$ , where  $s^\sharp$  is the inclusion map and  $r^\sharp$  maps every  $Q \in \mathcal{Q}_V^{cvx} \mathbb{P}_{\text{DP}}^\bullet X$  to itself.

Similarly, (21) also appears as the top row of the following diagram.

$$\begin{array}{ccccccc}
\mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\eta_{\mathcal{P}\ell_V^q X}} & \mathbf{V} \cdot \mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\lambda_{\mathbb{P}_{\text{ADP}}^\bullet X}^\sharp} & \mathcal{P}\ell_V^q \mathbf{V} \cdot \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\mathcal{P}\ell_V^q \alpha_{\text{ADP}}^\bullet} & \mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X \\
\downarrow \pi_2 & \text{nat. } \pi_2 & \downarrow \mathbf{V} \cdot \pi_2 & (19) & \downarrow \pi_2 & \text{nat. } \pi_2 & \downarrow \pi_2 \\
\mathcal{H}_V \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\eta_{\mathcal{H}_V \mathbb{P}_{\text{ADP}}^\bullet X}} & \mathbf{V} \cdot \mathcal{H}_V \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\lambda_{\mathbb{P}_{\text{ADP}}^\bullet X}^\flat} & \mathcal{H}_V \mathbf{V} \cdot \mathbb{P}_{\text{ADP}}^\bullet X & \xrightarrow{\mathcal{H}_V \alpha_{\text{ADP}}^\bullet} & \mathcal{H}_V \mathbb{P}_{\text{ADP}}^\bullet X \\
\downarrow \mathcal{H}_V \pi_2 & \text{nat. } \eta & \downarrow \mathbf{V} \cdot \mathcal{H}_V \pi_2 & \text{nat. } \lambda^\flat & \downarrow \mathcal{H}_V \mathbf{V} \cdot \pi_2 & \text{def. } \alpha_{\text{ADP}}^\bullet & \downarrow \mathcal{H}_V \pi_2 \\
\mathcal{H}_V \mathbb{P}_{\text{AP}}^\bullet X & \xrightarrow{\eta_{\mathcal{H}_V \mathbb{P}_{\text{AP}}^\bullet X}} & \mathbf{V} \cdot \mathcal{H}_V \mathbb{P}_{\text{AP}}^\bullet X & \xrightarrow{\lambda_{\mathbb{P}_{\text{AP}}^\bullet X}^\flat} & \mathcal{H}_V \mathbf{V} \cdot \mathbb{P}_{\text{AP}}^\bullet X & \xrightarrow{\mathcal{H}_V \alpha_{\text{AP}}^\bullet} & \mathcal{H}_V \mathbb{P}_{\text{AP}}^\bullet X
\end{array}$$

The bottom row is the idempotent (7) of Lemma 8.2 (with  $\alpha \stackrel{\text{def}}{=} \alpha_{\text{AP}}^\bullet$ ), which splits as  $\mathcal{H}_V \mathbb{P}_{\text{AP}}^\bullet X \xrightarrow{r^b} \mathcal{H}_V^{cvx} \mathbb{P}_{\text{AP}}^\bullet X \xrightarrow{s^b} \mathcal{H}_V \mathbb{P}_{\text{AP}}^\bullet X$ , where  $s^b$  is the inclusion map and  $r^b$  maps every  $C \in \mathcal{H}_V^{cvx} \mathbb{P}_{\text{AP}}^\bullet X$  to itself.

Let  $r^\sharp$  map every  $(Q, \mathcal{C}) \in \mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X$  to  $(r^\sharp(\mathcal{Q}_V \pi_1(Q)), r^b(\mathcal{H}_V \pi_2(\mathcal{C})))$ . This is a pair in  $\mathcal{Q}_V^{cvx} \mathbb{P}_{\text{ADP}}^\bullet X \times \mathcal{H}_V^{cvx} \mathbb{P}_{\text{ADP}}^\bullet X$ . We need to show that this is also a quasi-lens, and that  $r^\sharp$  is continuous. We observe that  $\pi_1 \circ r^\sharp = r^\sharp \circ \mathcal{Q}_V \pi_1 \circ \pi_1$  is the corestriction of the idempotent (3) to  $\mathcal{Q}_V^{cvx} \mathbb{P}_{\text{ADP}}^\bullet X$ , and  $\pi_2 \circ r^\sharp = r^b \circ \mathcal{H}_V \pi_1 \circ \pi_1$  is the corestriction of the idempotent (7) to  $\mathcal{H}_V^{cvx} \mathbb{P}_{\text{ADP}}^\bullet X$ , so  $r^\sharp$  is the corestriction of the idempotent (21) to  $\mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X$ . In particular,  $r^\sharp$  takes its values in (convex) quasi-lenses and is continuous.

Since  $r^\sharp$  maps convex elements to themselves, and similarly for  $r^\flat$ ,  $r^\sharp$  maps convex quasi-lenses to themselves.

We also define  $s^\sharp: \mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X \rightarrow \mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X$  as the inclusion map. Since  $r^\sharp$  maps convex quasi-lenses to themselves,  $r^\sharp \circ s^\sharp = \text{id}_{\mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X}$ . Also, by definition of  $r^\sharp$  as the corestriction of the idempotent (21) to  $\mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X$ , (21) is equal to  $s^\sharp \circ r^\sharp$ .  $\square$

**Lemma 11.3.** *For  $S \stackrel{\text{def}}{=} \mathcal{P}\ell_V^q$ ,  $T \stackrel{\text{def}}{=} \mathbf{V}_\bullet$ ,  $\lambda \stackrel{\text{def}}{=} \lambda^\sharp$ , the monad  $\bar{S}$  on the category of  $\mathbf{V}_\bullet$ -algebras over  $\mathbf{Top}^\sharp$  has the following properties:*

1. *For every free  $\mathbf{V}_\bullet$ -algebra  $\mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$ ,  $\bar{S}\mu_X: \mathbf{V}_\bullet \mathbb{P}_{\text{ADP}}^\bullet X \rightarrow \mathbb{P}_{\text{ADP}}^\bullet X$  is  $\alpha_{\text{ADP}}^\bullet$  (see Lemma 11.2).*
2. *The unit  $\eta_{\mu_X}^{\bar{S}}$  evaluated at the free  $\mathbf{V}_\bullet$ -algebra  $\mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$  is  $r_{\text{ADP}} \circ \eta_{\mathbf{V}_\bullet X}^q$ .*
3. *The multiplication  $\mu_{\mu_X}^{\bar{S}}$  evaluated at the free  $\mathbf{V}_\bullet$ -algebra  $\mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$  is the function that maps every  $(Q, C) \in \mathcal{P}\ell_V^q \mathbb{P}_{\text{ADP}}^\bullet X$  to  $(h \in \mathcal{L}X \mapsto \min_{(F^-, F^+) \in Q} F^-(h), h \in \mathcal{L}X \mapsto \sup_{(F^+, F^-) \in C} F^+(h)) \in \mathbb{P}_{\text{ADP}}^\bullet X$ .*
4. *The counit  $\epsilon^T$  of the adjunction  $F^T \dashv U^T$  between  $\mathbf{Top}$  and the category of  $T$ -algebras ( $\mathbf{V}_\bullet$ -algebras) is given at each algebra  $\beta: \mathbf{V}_\bullet X \rightarrow X$  as  $\beta$  itself.*

PROOF. 1. We know that  $\bar{S}\mu_X$  must be a  $\mathbf{V}_\bullet$ -algebra on  $\bar{S}\mathbf{V}_\bullet X$ , and the latter can be taken as  $\mathbb{P}_{\text{ADP}}^\bullet X$ , by Lemma 11.1. We know that  $\bar{S}\mu_X = \pi_{\mu_X} \circ S\mu_X \circ \lambda_X \circ T\iota_{\mu_X} = r_{\text{ADP}} \circ \mathcal{P}\ell_V^q \mu_X \circ \lambda_X^\sharp \circ \mathbf{V}_\bullet s_{\text{ADP}}^\bullet$ . Hence  $\pi_1 \circ \bar{S}\mu_X = r_{\text{DP}} \circ Q_V \mu_X \circ \lambda_X^\sharp \circ \mathbf{V}_\bullet s_{\text{DP}}^\bullet \circ \mathbf{V}_\bullet \pi_1$  using the definition of  $r_{\text{ADP}}$  (Proposition 9.1), the naturality of  $\pi_1$ , the definition of  $\lambda^\sharp$  (Theorem 10.2), and the definition of  $s_{\text{ADP}}^\bullet$  (Proposition 9.1). Hence, by Lemma 5.3, item 1,  $\pi_1 \circ \bar{S}\mu_X = \alpha_{\text{DP}}^\bullet \circ \mathbf{V}_\bullet \pi_1$ . Similarly,  $\pi_2 \circ \bar{S}\mu_X = r_{\text{AP}} \circ \mathcal{H}_V \mu_X \circ \lambda_X^\flat \circ \mathbf{V}_\bullet s_{\text{AP}}^\bullet \circ \mathbf{V}_\bullet \pi_2$ , so  $\pi_2 \circ \bar{S}\mu_X = \alpha_{\text{AP}}^\bullet \circ \mathbf{V}_\bullet \pi_2$ , using Lemma 8.3. We also have  $\pi_1 \circ \alpha_{\text{ADP}}^\bullet = \alpha_{\text{DP}}^\bullet \circ \mathbf{V}_\bullet \pi_1$  and  $\pi_2 \circ \alpha_{\text{ADP}}^\bullet = \alpha_{\text{AP}}^\bullet \circ \mathbf{V}_\bullet \pi_2$ , by definition of  $\alpha_{\text{ADP}}^\bullet$  (Lemma 11.2). Since  $\langle \pi_1, \pi_2 \rangle$  is injective,  $\bar{S}\mu_X = \alpha_{\text{ADP}}^\bullet$ .

$$2. \eta_{\mu_X}^{\bar{S}} = \pi_{\mu_X} \circ \eta_{\mathbf{V}_\bullet X}^S = r_{\text{ADP}} \circ \eta_{\mathbf{V}_\bullet X}^q.$$

3. In general,  $\mu_\alpha^{\bar{S}}$  is a morphism from  $\bar{S}\bar{S}\alpha$  to  $\bar{S}\alpha$  in the category of  $\mathbf{V}_\bullet$ -algebras. When  $\alpha = \mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$ , it is therefore one from  $\mathbb{P}_{\text{ADP}}^\bullet \mathbb{P}_{\text{ADP}}^\bullet X$  to  $\mathbb{P}_{\text{ADP}}^\bullet X$ . We have  $\mu_{\mu_X}^{\bar{S}} \stackrel{\text{def}}{=} \pi_{\mu_X} \circ \mu_{\mathbf{V}_\bullet X}^S \circ S\iota_{\mu_X} \circ \iota_{\bar{S}\mu_X}$ . We may take  $\iota_{\bar{S}\mu_X} = \iota_{\alpha_{\text{ADP}}^\bullet}$

to be the right arrow, from  $\mathcal{P}\ell_V^{\text{q} \text{cvx}} \mathbb{P}_{\text{ADP}}^\bullet X$  to  $\mathcal{P}\ell_V^{\text{q}} \mathbb{P}_{\text{ADP}}^\bullet X$  in Lemma 11.2—just subspace inclusion. Let  $m^\sharp$  be that subspace inclusion. Writing  $m^\sharp$  for the subspace inclusion of  $\mathcal{Q}_V^{\text{cvx}} \mathbb{P}_{\text{DP}}^\bullet X$  into  $\mathcal{Q}_V \mathbb{P}_{\text{DP}}^\bullet X$  and  $m^\flat$  for the subspace inclusion of  $\mathcal{H}_V^{\text{cvx}} \mathbb{P}_{\text{AP}}^\bullet X$  into  $\mathcal{H}_V \mathbb{P}_{\text{AP}}^\bullet X$ , we have  $\mathcal{Q}_V \pi_1 \circ \pi_1 \circ m^\sharp = m^\sharp \circ \mathcal{Q}_V \pi_1 \circ \pi_1$  and  $\mathcal{H}_V \pi_2 \circ \pi_2 \circ m^\flat = m^\flat \circ \mathcal{H}_V \pi_2 \circ \pi_2$ .

Then  $S\iota_{\mu_X} = \mathcal{P}\ell_V^{\text{q}} s_{\text{ADP}}^\bullet: \mathcal{P}\ell_V^{\text{q}} \mathbb{P}_{\text{ADP}}^\bullet X \rightarrow \mathcal{P}\ell_V^{\text{q}} \mathcal{P}\ell_V^{\text{q}} \mathbf{V}_\bullet X$ , by Lemma 11.1. We compose that with  $\mu_{\mathbf{V}_\bullet X}^{\text{q}}: \mathcal{P}\ell_V^{\text{q}} \mathcal{P}\ell_V^{\text{q}} \mathbf{V}_\bullet X \rightarrow \mathcal{P}\ell_V^{\text{q}} \mathbf{V}_\bullet X$ , then with  $r_{\text{ADP}}: \mathcal{P}\ell_V^{\text{q}} \mathbf{V}_\bullet X \rightarrow \mathbb{P}_{\text{ADP}}^\bullet X$ . The resulting composition is  $f \stackrel{\text{def}}{=} r_{\text{ADP}} \circ \mu_{\mathbf{V}_\bullet X}^{\text{q}} \circ \mathcal{P}\ell_V^{\text{q}} s_{\text{ADP}}^\bullet \circ m^\sharp$ .

Then  $\pi_1 \circ f = r_{\text{DP}} \circ \mu_{\mathbf{V}_\bullet X}^{\text{Q}} \circ \mathcal{Q}_V \pi_1 \circ \pi_1 \circ \mathcal{P}\ell_V^{\text{q}} s_{\text{ADP}}^\bullet \circ m^\sharp$  by definition of  $r_{\text{ADP}}$  (Proposition 9.1) and by (11) (see Proposition 8.5). Now  $\mathcal{Q}_V \pi_1 \circ \pi_1 \circ \mathcal{P}\ell_V^{\text{q}} s_{\text{ADP}}^\bullet \circ m^\sharp = \mathcal{Q}_V \pi_1 \circ \mathcal{Q}_V s_{\text{ADP}}^\bullet \circ \pi_1 \circ m^\sharp$  (by naturality of  $\pi_1$ ) =  $\mathcal{Q}_V s_{\text{DP}}^\bullet \circ m^\sharp \circ \mathcal{Q}_V \pi_1 \circ \pi_1$  (by definition of  $s_{\text{ADP}}^\bullet$ , see Proposition 9.1) =  $\mathcal{Q}_V s_{\text{DP}}^\bullet \circ m^\sharp \circ \mathcal{Q}_V \pi_1 \circ \pi_1$ . Therefore  $\pi_1 \circ f = r_{\text{DP}} \circ \mu_{\mathbf{V}_\bullet X}^{\text{Q}} \circ \mathcal{Q}_V s_{\text{DP}}^\bullet \circ m^\sharp \circ \mathcal{Q}_V \pi_1 \circ \pi_1$ . The composition  $r_{\text{DP}} \circ \mu_{\mathbf{V}_\bullet X}^{\text{Q}} \circ \mathcal{Q}_V s_{\text{DP}}^\bullet \circ m^\sharp$  was elucidated in Lemma 5.3, item 3, as mapping every  $Q \in \mathcal{Q}_V^{\text{cvx}} \mathbb{P}_{\text{DP}}^\bullet X$  to  $(h \in \mathcal{L}X \mapsto \min_{F \in Q} F(h))$ . Hence  $\pi_1 \circ f$  maps every convex quasi-lens  $(Q, C) \in \mathcal{P}\ell_V^{\text{q} \text{cvx}} \mathbb{P}_{\text{ADP}}^\bullet X$  to  $(h \in \mathcal{L}X \mapsto \min_{F \in \mathcal{Q}_V \pi_1(Q)} F(h)) = (h \in \mathcal{L}X \mapsto \min_{F \in \uparrow \pi_1[Q]} F(h)) = (h \in \mathcal{L}X \mapsto \min_{F \in \pi_1[Q]} F(h)) = (h \in \mathcal{L}X \mapsto \min_{(F^-, F^+) \in Q} F^-(h))$ .

Similarly, and using Lemma 8.3, item 3,  $\pi_2 \circ f$  maps every convex quasi-lens  $(Q, C) \in \mathcal{P}\ell_V^{\text{q} \text{cvx}} \mathbb{P}_{\text{ADP}}^\bullet X$  to  $(h \in \mathcal{L}X \mapsto \sup_{(F^-, F^+) \in Q} F^+(h))$ .

4. Standard category theory [34, Chapter VI, Section 2, Theorem 1].  $\square$

**Theorem 11.4.** *The weak composite monad associated with  $\lambda^\sharp: \mathbf{V}_\bullet \mathcal{P}\ell_V^{\text{q}} \rightarrow \mathcal{P}\ell_V^{\text{q}} \mathbf{V}_\bullet$  on  $\mathbf{Top}^\sharp$  is  $(\mathbb{P}_{\text{ADP}}^\bullet, \eta^{\text{ADP}}, \mu^{\text{ADP}})$ .*

PROOF. We take  $T \stackrel{\text{def}}{=} \mathbf{V}_\bullet$ ,  $S \stackrel{\text{def}}{=} \mathcal{P}\ell_V^{\text{q}}$ ,  $\lambda \stackrel{\text{def}}{=} \lambda^\sharp$ . The functor  $U^T$  takes every  $\mathbf{V}_\bullet$ -algebra  $\alpha: \mathbf{V}_\bullet X \rightarrow X$  to  $X$  and every  $\mathbf{V}_\bullet$ -algebra morphism  $f: (\alpha: \mathbf{V}_\bullet X \rightarrow X) \rightarrow (\beta: \mathbf{V}_\bullet Y \rightarrow Y)$  to the underlying continuous map  $f: X \rightarrow Y$ . The functor  $F^T$  takes every space  $X$  to  $\mu_X: \mathbf{V}_\bullet \mathbf{V}_\bullet X \rightarrow \mathbf{V}_\bullet X$ , and every continuous map  $f: X \rightarrow Y$  to  $\mathbf{V}_\bullet f$ , seen as a morphism of  $\mathbf{V}_\bullet$ -algebras.

Then  $U^T \overline{S} F^T$  maps every space  $X$  to the splitting  $\mathbb{P}_{\text{ADP}}^\bullet X$  of  $\mathcal{P}\ell_V^{\text{q}} \mu_X \circ \lambda_X^\sharp \circ \eta_{\mathcal{H}_V X}$  we have obtained in Lemma 11.1. We also obtain  $\pi_{\mu_X} \stackrel{\text{def}}{=} r_{\text{ADP}}$ ,  $\iota_{\mu_X} \stackrel{\text{def}}{=} s_{\text{ADP}}^\bullet$ .

Given any continuous map  $f: X \rightarrow Y$ ,  $U^T \overline{S} F^T(f)$  is equal to (the map underlying the  $\mathbf{V}_\bullet$ -algebra)  $\overline{S}(\mathbf{V}_\bullet f)$ , namely  $\pi_{\mu_Y} \circ \mathcal{P}\ell_V^{\text{q}} \mathbf{V}_\bullet f \circ \iota_{\mu_X} = r_{\text{ADP}} \circ \mathcal{P}\ell_V^{\text{q}} \mathbf{V}_\bullet f \circ s_{\text{ADP}}^\bullet$ . We claim that this is equal to  $\mathbb{P}_{\text{ADP}}^\bullet(f)$ . It suffices to verify that  $\pi_1 \circ r_{\text{ADP}} \circ \mathcal{P}\ell_V^{\text{q}} \mathbf{V}_\bullet f \circ s_{\text{ADP}}^\bullet = \pi_1 \circ \mathbb{P}_{\text{ADP}}^\bullet(f)$ , and similarly with  $\pi_2$ .

By definition of  $r_{\text{ADP}}$  (Proposition 9.1), naturality of  $\pi_1$  and definition of  $s_{\text{ADP}}^\bullet$  (Proposition 9.1),  $\pi_1 \circ r_{\text{ADP}} \circ \mathcal{P}\ell_{\mathbf{V}}^q \mathbf{V} \bullet f \circ s_{\text{ADP}}^\bullet = r_{\text{DP}} \circ \mathcal{Q}_{\mathbf{V}} \mathbf{V} \bullet f \circ s_{\text{DP}}^\bullet \circ \pi_1$ , namely to  $\mathbb{P}(f) \circ \pi_1 = \pi_1 \circ \mathbb{P}_{\text{ADP}}^\bullet(f)$ . Similarly with  $\pi_2$ .

The unit of the weak composite monad is  $U^T \eta^{\bar{S}} F^T \circ \eta^T$ . At object  $X$ , this maps every  $x \in X$  to  $r_{\text{ADP}} \circ \eta_{\mathbf{V} \bullet X}^q$ , by Lemma 11.3, item 2. In order to show that this is equal to  $\eta_X^{\text{ADP}}$ , it suffices to show that  $\pi_1 \circ r_{\text{ADP}} \circ \eta_{\mathbf{V} \bullet X}^q = \pi_1 \circ \eta_X^{\text{ADP}}$  and similarly with  $\pi_2$ . This follows from the definition of  $r_{\text{ADP}}$  (Proposition 9.1), the definition of  $\eta^q$  (see (10) in Proposition 8.5) and the definition of  $\eta^{\text{ADP}}$  (see (12) in Proposition 9.3).

The multiplication is  $U^T \mu^{\bar{S}} F^T \circ U^T \bar{S} \epsilon^T \bar{S} F^T$ . We recall what that means. At any object  $X$ ,  $\bar{S} F^T X$  is the  $T$ -algebra  $\bar{S} \mu_X: T \bar{S} T X \rightarrow \bar{S} T X$ . Then  $\epsilon^T$  evaluated at this  $T$ -algebra is the bottom arrow in the following diagram:

$$\begin{array}{ccc} T T \bar{S} T X & \xrightarrow{T \epsilon_{\bar{S} F^T X}^T} & T \bar{S} T X \\ \mu_{\bar{S} T X} \downarrow & & \downarrow \bar{S} \mu_X \\ T \bar{S} T X & \xrightarrow{\epsilon_{\bar{S} F^T X}^T} & \bar{S} T X \end{array}$$

from the  $T$ -algebra  $\alpha \stackrel{\text{def}}{=} \mu_{\bar{S} T X}$  that is the leftmost vertical arrow to the  $T$ -algebra  $\beta \stackrel{\text{def}}{=} \bar{S} \mu_X$  on the right. We apply  $\bar{S}$  to the morphism  $\epsilon_{\bar{S} F^T X}^T$  (i.e., we now apply  $\bar{S}$  to a morphism of  $T$ -algebras, not to a  $T$ -algebra, as we did before), and we obtain the composition:

$$\bar{S} T \bar{S} T X \xrightarrow{\iota_\alpha} \bar{S} T \bar{S} T X \xrightarrow{S \epsilon_{\bar{S} F^T X}^T} \bar{S} \bar{S} T X \xrightarrow{\pi_\beta} \bar{S} \bar{S} T X. \quad (22)$$

Applying  $U^T$  to the latter, we obtain the same composition, this time seen as a morphism in the base category, instead of as a morphism in the category of  $T$ -algebras. We finally compose it with  $U^T \mu_{F^T X}^{\bar{S}}$ , which is  $\mu_{\mu_X}^{\bar{S}}: \bar{S} \bar{S} T X \rightarrow \bar{S} T X$ .

By Lemma 11.3, item 4,  $\epsilon_{\bar{S} F^T X}^T: \alpha \rightarrow \beta$  is  $\beta$  itself, where  $\beta = \bar{S} \mu_X = \alpha_{\text{ADP}}^\bullet$  (Lemma 11.3, item 1) and  $\alpha = \mu_{\bar{S} T X} = \mu_{\mathbb{P}_{\text{ADP}}^\bullet X}$ : we have  $\bar{S} T X = \mathbb{P}_{\text{ADP}}^\bullet X$ , using Lemma 11.1), and  $\iota_\alpha = s_{\text{ADP}}^\bullet$ . Using Lemma 11.2,  $\pi_\beta: \mathcal{P}\ell_{\mathbf{V}}^q \mathbb{P}_{\text{ADP}}^\bullet X \rightarrow \mathcal{P}\ell_{\mathbf{V}}^{q \text{cvx}} \mathbb{P}_{\text{ADP}}^\bullet X$  is the corestriction of (21) (call it  $f$ ), which maps every convex element of  $\mathcal{P}\ell_{\mathbf{V}}^q \mathbb{P}_{\text{ADP}}^\bullet X$  to itself.

Hence the multiplication of the weak composite monad, evaluated at  $X$ ,

is the composition of (22) with  $\mu_{\mu_X}^{\bar{S}}$  (given in Lemma 11.3, item 3), namely:

$$\mathbb{P}_{\text{ADP}}^{\bullet} \mathbb{P}_{\text{ADP}}^{\bullet} X \xrightarrow{s_{\text{ADP}}^{\bullet}} \mathcal{P}\ell_{\mathbf{V}}^{\text{q}} \mathbf{V}_{\bullet} \mathbb{P}_{\text{ADP}}^{\bullet} X \xrightarrow{\mathcal{P}\ell_{\mathbf{V}}^{\text{q}} \alpha_{\text{ADP}}^{\bullet}} \mathcal{P}\ell_{\mathbf{V}}^{\text{q}} \mathbb{P}_{\text{ADP}}^{\bullet} X \xrightarrow{f} \mathcal{P}\ell_{\mathbf{V}}^{\text{q} \text{ cvx}} \mathbb{P}_{\text{ADP}}^{\bullet} X \xrightarrow{\mu_{\mu_X}^{\bar{S}}} \mathbb{P}_{\text{ADP}}^{\bullet} X$$

Let us call  $g$  for that composition. Using the explicit expression for  $\mu_{\mu_X}^{\bar{S}}$ , and the appropriate naturality equations, we recognize  $\pi_1 \circ g$  as being equal to the composition (5)  $\circ \mathcal{Q}_{\mathbf{V}} \pi_1 \circ \pi_1$ ; (5) is the expression of the weak composite monad we obtained from  $\lambda^{\sharp}$  in the proof of Theorem 5.4, which we elucidated as mapping every  $\mathcal{F} \in \mathbb{P}_{\text{DP}}^{\bullet} \mathbb{P}_{\text{DP}}^{\bullet} X$  to  $(h \in \mathcal{L}X \mapsto \mathcal{F}(F \in \mathbb{P}_{\text{DP}}^{\bullet} X \mapsto F(h)))$ .

Similarly,  $\pi_2 \circ g$  is equal to (9)  $\circ \mathcal{H}_{\mathbf{V}} \pi_2 \circ \pi_2$ , where (9) was elucidated in the proof of Theorem 8.4 as mapping every  $\mathcal{F} \in \mathbb{P}_{\text{AP}}^{\bullet} \mathbb{P}_{\text{AP}}^{\bullet} X$  to  $(h \in \mathcal{L}X \mapsto \mathcal{F}(F \in \mathbb{P}_{\text{AP}}^{\bullet} X \mapsto F(h)))$ . Hence  $g$  maps every  $(\mathcal{F}^-, \mathcal{F}^+) \in \mathbb{P}_{\text{ADP}}^{\bullet} \mathbb{P}_{\text{ADP}}^{\bullet} X$  to the pair of maps  $(h \in \mathcal{L}X \mapsto \mathcal{H}_{\mathbf{V}} \pi_1(\mathcal{F}^-)(F \in \mathbb{P}_{\text{DP}}^{\bullet} X \mapsto F(h)))$  and  $(h \in \mathcal{L}X \mapsto \mathcal{H}_{\mathbf{V}} \pi_2(\mathcal{F}^+)(F \in \mathbb{P}_{\text{DP}}^{\bullet} X \mapsto F(h)))$ , namely to  $\mu_X^{\text{ADP}}(\mathcal{F}^+, \mathcal{F}^-)$  (see Proposition 9.3).  $\square$

**Remark 11.5.** *The isomorphic weak distributive law of Remark 10.4 on the category of stably compact spaces must induce the same weak composite monad, namely  $(\mathbb{P}_{\text{ADP}}^{\bullet}, \eta^{\text{ADP}}, \mu^{\text{ADP}})$ —or rather, its restriction to the category of stably compact spaces.*

**Remark 11.6.** *A space is  $T_1$  if and only if its specialization preordering is the equality relation. Every Hausdorff space is  $T_1$ . On a  $T_1$  space  $X$ , a lens is the same thing as a non-empty compact subset. Hence  $\mathcal{P}\ell_{\mathbf{V}} X$  is the usual hyperspace of non-empty compact subsets, with the (usual) Vietoris topology, and the  $\mathcal{P}\ell_{\mathbf{V}}$  monad specializes to what has sometimes been called the Vietoris monad on the category  $\mathbf{KHaus}$  of compact Hausdorff spaces (although Vietoris had no knowledge of monads, and while he defined the Vietoris topology on the space of closed subsets of a space [41], spaces of subsets were further explored by Hausdorff [22, §28] then by Michael [35], among others). In other words, relying on Remark 8.6, removing the now useless signs  $\uparrow$  and simplifying,  $\mathcal{P}\ell_{\mathbf{V}} f: \mathcal{P}\ell_{\mathbf{V}} X \rightarrow \mathcal{P}\ell_{\mathbf{V}} Y$  maps every non-empty compact subset  $L$  to its image  $f[L]$  under  $f$ , the unit  $\eta_X^{\mathcal{P}\ell}$  maps every  $x \in X$  to  $\{x\}$ , and the multiplication  $\mu_X^{\mathcal{P}\ell}$  maps every  $\mathcal{L} \in \mathcal{P}\ell_{\mathbf{V}} \mathcal{P}\ell_{\mathbf{V}} X$  to  $\bigcup \mathcal{L}$ .*

*Specializing Remark 11.5 to the appropriate subcategory, we obtain a weak distributive law of the Vietoris monad over  $\mathbf{V}_{\bullet}$  on  $\mathbf{KHaus}$ , and its associated weak composite monad is the appropriate restriction of  $(\mathbb{P}_{\text{ADP}}^{\bullet}, \eta^{\text{ADP}}, \mu^{\text{ADP}})$ , by Theorem 11.4.*



**Remark 11.7.** We refine Remark 10.5 in the case of regular spaces. A space is regular if and only if for every point  $x$ , every open neighborhood  $U$  of  $x$  contains a closed neighborhood  $C$  of  $x$ ; equivalently, if every open set is the union of the directed family of interiors  $\text{int}(C)$  of closed subsets of  $U$ . For every regular space  $X$ ,  $\mathbf{V}_1X$  is Hausdorff: given any two distinct elements  $\mu, \nu \in \mathbf{V}_\bullet$ , there must be an open subset  $U$  of  $X$  such that  $\mu(U) \neq \nu(U)$ , say  $\mu(U) < \nu(U)$ . Let  $r$  be such that  $\mu(U) < r < \nu(U)$ . Since  $r < \nu(U) = \sup_C \nu(\text{int}(C))$ , where  $C$  ranges over the closed subsets of  $U$ , we have  $r < \nu(\text{int}(C))$  for some closed subset  $C$  of  $U$ . Then  $\nu$  is in  $[\text{int}(C) > r]$ ,  $\mu$  is in  $[X \setminus C > 1 - r]$  (since  $1 = \mu(X) = \mu((X \setminus C) \cup U) \leq \mu(X \setminus C) + \mu(U) < \mu(X \setminus C) + r$ , hence  $\mu(X \setminus C) > 1 - r$ ), and  $[\text{int}(C) > r]$  and  $[X \setminus C > 1 - r]$  are disjoint; for the latter, any probability valuation in the intersection must map the disjoint union  $\text{int}(C) \cup (X \setminus C)$  to a number strictly larger than  $r + (1 - r) = 1$ , which is impossible. (One could show that  $\mathbf{V}_1X$  is in fact regular Hausdorff, namely  $T_3$ , but we will not need that.)

Every compact Hausdorff space  $X$  is regular, so the specialization ordering on  $\mathbf{V}_1X$  is equality. Specializing Remark 10.5 to this case, the weak distributive law  $\lambda^\natural$  on  $\mathbf{KHaus}$  maps every  $\mu \stackrel{\text{def}}{=} \sum_{i=1}^n a_i \delta_{E_i}$ , where  $n \geq 1$ , and each set  $E_i$  is finite and non-empty, to  $\lambda_X^\natural(\mu) = \{\sum_{i=1}^n a_i \delta_{x_i} \mid x_1 \in E_1, \dots, x_n \in E_n\}$ . This is the formula given by Goy and Petrişan for their weak distributive law of the powerset monad over the finite distribution monad [21, Lemma 3.1].

## 12. Radon measures

We finish with a discussion of Radon measures. A *Radon measure* on a Hausdorff topological space  $X$  is a Borel measure  $\mu$  such that for every Borel subset  $B$  of  $X$ , for every  $\epsilon > 0$ , there is a compact set  $K_\epsilon$  included in  $B$  such that  $\mu(B \setminus K_\epsilon) \leq \epsilon$  [5, Definition 7.1.1]. One needs to take care to modify this definition slightly on non-Hausdorff spaces [29, Definition 7.2], if only because  $K_\epsilon$  may fail to be Borel: a *Radon measure*  $\mu$  on a coherent sober space  $X$  is a Borel measure on a  $\sigma$ -algebra that contains the patch topology on  $X$ , whose restriction to  $\mathcal{O}(X^{\text{patch}})$  is locally finite (every point has a patch-open neighborhood of finite  $\mu$ -measure) and *inner regular*, meaning that  $\mu(B) = \sup_{K \in \mathcal{C}X, K \subseteq B} \mu(K)$  for every Borel subset  $B$ ; the collection  $\mathcal{C}X$  consists of the intersections of non-empty families of finite unions of lenses. (See Appendix A for Keimel and Lawson's definition, and why we changed it to the current one.) The *patch topology* on  $X$  is the coarsest topology that contains the original topology on  $X$ , as well as the complements of all

compact saturated subsets of  $X$ . The patch topology coincides with the original topology on  $X$  if  $X$  is Hausdorff, since in that case every compact set is closed.

In the following, a *well-filtered* space  $X$  is a topological space such that for every filtered family  $(Q_i)_{i \in I}$  of compact saturated subsets, for every open subset  $U$  of  $Z$ , if  $\bigcap_{i \in I}^\downarrow Q_i \subseteq U$  then  $Q_i \subseteq U$  for some  $i \in I$ . A family  $(Q_i)_{i \in I}$  is *filtered* if and only if it is non-empty, and for any two elements  $Q_i$  and  $Q_{i'}$ , there is an element  $Q_{i''}$  included in both. In a well-filtered space  $X$ , for every filtered family  $(Q_i)_{i \in I}$ ,  $\bigcap_{i \in I}^\downarrow Q_i$  is compact saturated [13, Proposition 8.3.6]. Every sober space is well-filtered [13, Proposition 8.3.5].

**Lemma 12.1.** *For every coherent well-filtered space  $X$ , every element of  $\mathcal{C}X$  is compact in the patch topology, hence in the original topology on  $X$ .*

In the sequel, we would only need to show that every element of  $\mathcal{C}X$  is compact in  $X$ , not in  $X^{\text{patch}}$ . But Lemma 12.1 fixes a flaw. Let me explain.

Lemma 7.1 (5) of [29] states that, in a coherent sober space  $X$ , every element of  $\mathcal{C}X$  is compact in the patch topology, and the proof really only uses well-filteredness, not sobriety. However, the result is obtained from the statement that every patch-closed subset of  $X$  is the union of an element of  $\mathcal{C}X$  with a closed subset of  $X$  [29, Lemma 7.1 (4)], and that is wrong. For a counterexample (to item (4), not (5), of that lemma), consider  $\mathbb{R}$  with its Scott topology for  $X$ . Then  $X^{\text{patch}}$  is  $\mathbb{R}$  with its usual metric topology, while it is an easy exercise to show that the elements of  $\mathcal{C}X$  are the compact subsets of  $\mathbb{R}$  (in accordance with Lemma 12.1; the easiest elementary proof consists in verifying that every element of  $\mathcal{C}X$  is closed and bounded in  $\mathbb{R}$ ). Then any union of an element of  $\mathcal{C}X$  with a closed subset of  $X$  must be bounded from above. Hence, for example,  $\mathbb{Z}$ , which is closed in  $X^{\text{patch}} = \mathbb{R}$ , is not of that form. The fixed proof below, which is in the spirit of the original proof of [29, Lemma 7.1 (5)], was communicated to me by Jimmie Lawson on August 22nd, 2024.

PROOF. Just as in [29, Lemma 7.1], let the *lens topology* be the topology on  $X$  whose closed subsets are the intersections of finite unions of lenses. We will write  $X^\ell$  for  $X$  with the lens topology. The closed subsets of  $X^\ell$  are the elements of  $\mathcal{C}X$ , plus possibly  $X$  itself (the empty intersection) if  $X$  is not compact. It is clear that the lens topology is coarser than the patch topology. It is a consequence of Alexander's subbase lemma that  $X^\ell$  is compact (see [29, Lemma 7.1 (2)]).

We now claim that: (\*) for every  $K \in \mathcal{C}X$ , the lens and the patch topologies on  $X$  induce the same subspace topology on  $K$ . Since the lens topology is coarser than the patch topology on  $X$ , the same inclusion holds for their subspace topologies on  $K$ . Conversely, we must show that given any patch-closed subset  $D$  of  $X$ ,  $D \cap K$  is closed in  $K$  seen as a subspace of  $X^\ell$ . Since patch-closed subsets are intersections of finite unions of closed subsets and of compact saturated subsets of  $X$ , it suffices to show this when  $D$  is either closed or compact saturated in  $X$ .

Let us write  $K$  as  $\bigcap_{i \in I} \bigcup_{j=1}^{m_i} (Q_{ij} \cap C_{ij})$  where  $I \neq \emptyset$ , each  $Q_{ij}$  is compact saturated and each  $C_{ij}$  is closed in  $X$ .

If  $D$  is a closed subset  $C$  of  $X$ , then  $C \cap K = \bigcap_{i \in I} \bigcup_{j=1}^{m_i} (Q_{ij} \cap C \cap C_{ij})$ . Every set  $Q_{ij} \cap C \cap C_{ij}$  is a lens (or empty), so  $C \cap K$  is in  $\mathcal{C}X$ , hence closed in  $X^\ell$ . Then  $C \cap K$  is also equal to the intersection of itself with  $K$ , hence is closed in  $K$  as a subspace of  $X^\ell$ .

If  $D$  is a compact saturated subset  $Q$  of  $X$ , then similarly  $Q \cap K = \bigcap_{i \in I} \bigcup_{j=1}^{m_i} (Q \cap Q_{ij} \cap C_{ij})$ . Since  $X$  is coherent, each set  $Q \cap Q_{ij}$  is compact saturated in  $X$ , so  $Q \cap Q_{ij} \cap C_{ij}$  is a lens (or is empty). Therefore  $Q \cap K$  is in  $\mathcal{C}X$ , hence is closed in  $X^\ell$ . Then  $Q \cap K$  is also equal to the intersection of that set  $Q \cap K$  with  $K$ , hence is closed in  $K$  as a subspace of  $X^\ell$ . This ends the proof of claim (\*).

Using (\*), we can now show that every  $K \in \mathcal{C}X$  is compact in  $X^{\text{patch}}$ .  $K$  is closed in  $X^\ell$ , hence it is also closed in  $K$ , seen as a subspace of  $X^\ell$ . Since  $X^\ell$  is compact,  $K$  is compact in  $X^\ell$ , hence also in  $K$ , seen as a subspace of  $X^\ell$ . But, by (\*), the latter is also equal to  $K$ , seen as a subspace of  $X^{\text{patch}}$ . Any subset of a subspace is compact in the surrounding space, so  $K$  is compact in  $X^{\text{patch}}$ .

Finally, every compact subset of  $X^{\text{patch}}$  is compact in  $X$ : every cover by open subsets from  $X$  is also an open cover by open subsets from  $X^{\text{patch}}$ .  $\square$

Given any topological space  $X$ , the collection  $\blacksquare Q$  of open neighborhoods of an arbitrary compact saturated subset  $Q$  is a Scott-open subset of  $\mathcal{O}X$ . It follows that any union  $\bigcup_{i \in I} \blacksquare Q_i$ , where each  $Q_i$  is compact saturated in  $X$ , is Scott-open in  $\mathcal{O}X$ . A space is *consonant* if and only if all the Scott-open subsets of  $\mathcal{O}X$  are of this form. The notion arises from [8], where it was proved that every regular Čech-complete space is consonant; every locally compact space is consonant, too, as well as every LCS-complete space [7, Proposition 12.1].

**Proposition 12.2.** For  $\bullet$  equal to “ $\leq 1$ ”, resp. “1”, for every coherent sober space  $X$ , let  $\mathbf{R}_\bullet X$  be the set of Radon subprobability (resp. probability) measures on  $X$ , with the weak topology, defined by subbasic open sets  $\{\mu \in \mathbf{R}_\bullet X \mid \mu(U) > r\}$ ,  $U \in \mathcal{O}X$ ,  $r \in \mathbb{R}_+$ .

If  $X$  is consonant, weakly Hausdorff, coherent and sober (e.g., if  $X$  is stably locally compact), then restriction to  $\mathcal{O}X$  defines a homeomorphism between  $\mathbf{R}_\bullet X$  and  $\mathbf{V}_\bullet X$ .

PROOF. Every Radon measure  $\mu$  on the  $\sigma$ -algebra generated by the patch topology on a coherent sober space  $X$  has a restriction  $\nu$  to  $\mathcal{O}X$  that is clearly strict and modular. We claim that it is *tight*, namely that for every open subset  $U$ , for every  $r < \nu(U)$ , there is a compact saturated subset  $Q$  of  $U$  such that  $r < \nu(V)$  for every open neighborhood  $V$  of  $Q$ . Let  $U \in \mathcal{O}X$  and let  $r < \nu(U)$ ; equivalently,  $r < \mu(U)$ . By definition of Radon measures, there is a proper closed subset  $K$  of  $U$  in the lens topology such that  $K \subseteq U$  and  $r < \mu(K)$ . By Lemma 12.1,  $K$  is compact in  $X$ , so  $Q \stackrel{\text{def}}{=} \uparrow K$  is compact saturated in  $X$ . Since  $U$  is upwards-closed,  $Q \subseteq U$ , and we have  $r < \mu(K) \leq \mu(Q) \leq \mu(V) = \nu(V)$  for every open neighborhood  $V$  of  $Q$ . We now use the fact that every tight map from  $\mathcal{O}X$  to  $\overline{\mathbb{R}}_+$  is Scott-continuous [16, Lemma 6.2], so  $\nu$  is a continuous valuation.

In the converse direction, by [29, Theorem 7.3], every locally finite, tight valuation  $\nu$  on a weakly Hausdorff coherent sober space  $X$  extends to a unique Radon measure on the  $\sigma$ -algebra generated by the patch topology. In this sentence, that  $\nu$  is locally finite means that every point  $x$  has an open neighborhood  $U$  such that  $\nu(U) < \infty$ ; this is certainly the case for all (sub)probability valuations. Every continuous valuation  $\nu$  on a consonant space is tight [16, Lemma 6.2]. Hence every (sub)probability valuation on a consonant, weakly Hausdorff, coherent sober space extends to a unique Radon measure on the  $\sigma$ -algebra generated by the patch topology.

In conclusion, restriction to  $\mathcal{O}X$  defines a bijection of  $\mathbf{R}_\bullet X$  onto  $\mathbf{V}_\bullet X$ . It is clear that it is a homeomorphism.  $\square$

It follows that there is a monad of Radon (sub)probability measures  $\mathbf{R}_\bullet$  on the category of stably compact spaces, which is isomorphic to the monad  $\mathbf{V}_\bullet$  of (sub)probability valuations. (We recall that  $\mathbf{V}_\bullet$  preserves stable compactness; otherwise, the previous sentence makes no sense.)

Remark 11.5 then entails the following.

**Proposition 12.3.** *Let  $\bullet$  be “ $\leq 1$ ” or “1”. There is a weak distributive law of  $\mathcal{P}l_{\mathcal{V}}^{\mathbf{q}}$  (equivalently,  $\mathcal{P}l_{\mathcal{V}}$ ) over  $\mathbf{R}_{\bullet}$  (equivalently,  $\mathbf{V}_{\bullet}$ ) on the category of stably compact spaces, and it induces the composite monad  $(\mathbb{P}_{\text{ADP}}^{\bullet}, \eta^{\text{ADP}}, \mu^{\text{ADP}})$ .*

Similarly, Remark 11.6 entails the following.

**Proposition 12.4.** *Let  $\bullet$  be “ $\leq 1$ ” or “1”. There is a weak distributive law of the Vietoris monad over  $\mathbf{R}_{\bullet}$  (equivalently,  $\mathbf{V}_{\bullet}$ ) on  $\mathbf{KHaus}$ , and it induces the composite monad  $(\mathbb{P}_{\text{ADP}}^{\bullet}, \eta^{\text{ADP}}, \mu^{\text{ADP}})$ .*

### Competing interests

The author declares none.

### Acknowledgements

The elegant proof of Lemma 12.1 is due to Jimmie Lawson, and replaces a more complex argument of mine. Thanks, Jimmie!

### References

- [1] Abramsky, S., Jung, A., 1994. Domain theory. In: Abramsky, S., Gabbay, D. M., Maibaum, T. S. E. (Eds.), Handbook of Logic in Computer Science. Vol. 3. Oxford University Press, pp. 1–168.  
URL <https://www.cs.bham.ac.uk/~axj/pub/papers/handy1.pdf>
- [2] Adamski, W., 1977.  $\tau$ -smooth Borel measures on topological spaces. *Mathematische Nachrichten* 78, 97–107.
- [3] Alvarez-Manilla, M., Jung, A., Keimel, K., 2004. The probabilistic powerdomain for stably compact spaces. *Theoretical Computer Science* 328 (3), 221–244.
- [4] Beck, J., 1969. Distributive laws. In: Eckmann, B. (Ed.), Seminar on Triples and Categorical Homology Theory. Vol. 80 of Lecture Notes on in Mathematics. Springer Verlag, pp. 119–140.
- [5] Bogachev, V. I., 2007. Measure Theory. Vol. II. Springer-Verlag.

- [6] de Brecht, M., 2013. Quasi-Polish spaces. *Annals of Pure and Applied Logic* 164 (3), 356–381.  
URL <https://www.sciencedirect.com/science/article/pii/S0168007212001820>
- [7] de Brecht, M., Goubault-Larrecq, J., Jia, X., Lyu, Z., 2019. Domain-complete and LCS-complete spaces. *Electronic Notes in Theoretical Computer Science* Presented at the 8th International Symposium of Domain Theory and its Applications (ISDT'19). Available on arXiv:1902.11142 [math.GN].  
URL <https://www.sciencedirect.com/science/article/pii/S1571066119300337>
- [8] Dolecki, S., Greco, G. H., Lechicki, A., Aug. 1995. When do the upper Kuratowski topology (homeomorphically, Scott topology) and the co-compact topology coincide? *Transactions of the American Mathematical Society* 347 (8), 2869–2884.
- [9] Garner, R., 2020. The Vietoris monad and weak distributive laws. *Applied Categorical Structures* 28 (2).
- [10] Gierz, G., Hofmann, K. H., Keimel, K., Lawson, J. D., Mislove, M., Scott, D. S., 2003. *Continuous Lattices and Domains*. Vol. 93 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press.  
URL <https://www.cambridge.org/core/books/continuous-lattices-and-domains/D27E855CD9C88E6DB3C98196B88F0AF6>
- [11] Goubault-Larrecq, J., Sep. 2007. Continuous previsions. In: Duparc, J., Henzinger, Th. A. (Eds.), *Proceedings of the 16th Annual EACSL Conference on Computer Science Logic (CSL'07)*. Vol. 4646 of *Lecture Notes in Computer Science*. Springer, Lausanne, Switzerland, pp. 542–557.  
URL <http://www.lsv.ens-cachan.fr/Publis/PAPERS/PDF/JGL-cs107.pdf>
- [12] Goubault-Larrecq, J., 2010. De Groot duality and models of choice: Angels, demons and nature. *Mathematical Structures in Computer Science*

20 (2), 169–237.

URL <https://hal.inria.fr/hal-03195342>

- [13] Goubault-Larrecq, J., 2013. Non-Hausdorff Topology and Domain Theory—Selected Topics in Point-Set Topology. Vol. 22 of New Mathematical Monographs. Cambridge University Press.  
URL <https://www.cambridge.org/core/books/nonhausdorff-topology-and-domain-theory/47A93B1951D60717E2E71030CB0A4441>
- [14] Goubault-Larrecq, J., Sep. 2017. Isomorphism theorems between models of mixed choice. *Mathematical Structures in Computer Science* 27 (6), 1032–1067.  
URL <http://www.lsv.fr/Publis/PAPERS/PDF/JGL-mscs16.pdf>
- [15] Goubault-Larrecq, J., 2017. A non-Hausdorff minimax theorem. *Minimax Theory and its Applications* 3 (1), 73–80.
- [16] Goubault-Larrecq, J., 2021. Products and projective limits of continuous valuations on  $T_0$  spaces. *Mathematical Structures in Computer Science* 31 (2), 234–254.  
URL <https://www.cambridge.org/core/journals/mathematical-structures-in-computer-science/article/abs/products-and-projective-limits-of-continuous-valuations-on-t0-spaces/E46326A3597BBD943A9921B525C73EB9w>
- [17] Goubault-Larrecq, J., 2023. On weakly Hausdorff spaces and locally strongly sober spaces. *Topology Proceedings* 62, 117–131, available on arXiv:2211.10400 [math.GN].  
URL <http://topology.nipissingu.ca/tp/reprints/v62/>
- [18] Goubault-Larrecq, J., 2024. Jean. *Topology Proceedings* Submitted, July 2024.
- [19] Goubault-Larrecq, J., 2024. On the preservation of projective limits by functors of non-deterministic, probabilistic, and mixed choice. arXiv, available on arXiv:2407.10235 [math.GN].
- [20] Goy, A., 2021. Compositionnalité des monads par lois de distributivité faibles (on the compositionality of monads via weak distributive laws). Ph.D. thesis, Université Paris-Saclay.

- [21] Goy, A., Petrişan, D., 2020. Combining probabilistic and non-deterministic choice via weak distributive laws. In: Proceedings of the 35th Annual ACM/IEEE Symposium on Logic in Computer Science. LICS '20. Association for Computing Machinery, New York, NY, USA, p. 454–464.  
URL <https://doi.org/10.1145/3373718.3394795>
- [22] Hausdorff, F., 1927. Mengenlehre, 3rd Edition. Springer, Berlin.
- [23] Heckmann, R., 1993. Power domains and second-order predicates. Theoretical Computer Science 111, 59–88.
- [24] Heckmann, R., Dec 1996. Spaces of valuations. In: Andima, S., Flagg, R. C., Itzkowitz, G., Kong, Y., Kopperman, R., Misra, P. (Eds.), Papers on General Topology and its Applications. Vol. 806. Annals of the New York Academy of Science, pp. 174–200.
- [25] Jones, C., 1990. Probabilistic non-determinism. Ph.D. thesis, University of Edinburgh, technical Report ECS-LFCS-90-105.
- [26] Jones, C., Plotkin, G., 1989. A probabilistic powerdomain of evaluations. In: Proceedings of the 4th Annual Symposium on Logic in Computer Science. IEEE Computer Society Press, pp. 186–195.
- [27] Jung, A., 2004. Stably compact spaces and the probabilistic powerspace construction. In: Desharnais, J., Panangaden, P. (Eds.), Domain-theoretic Methods in Probabilistic Processes. Vol. 87 of Electronic Lecture Notes in Computer Science. Elsevier, pp. 5–20, 15pp.
- [28] Keimel, K., 2008. Topological cones: Functional analysis in a  $T_0$ -setting. Semigroup Forum 77 (1), 109–142.
- [29] Keimel, K., Lawson, J., 2005. Measure extension theorems for  $T_0$ -spaces. Topology and its Applications 149 (1–3), 57–83.  
URL <https://www.sciencedirect.com/science/article/pii/S0166864104002755>
- [30] Keimel, K., Plotkin, G., 2009. Predicate transformers for convex powerdomains. Mathematical Structures in Computer Science 19 (3), 501–539.



- [31] Kirch, O., 1993. Bereiche und Bewertungen. Diplomarbeit, Technische Hochschule Darmstadt.
- [32] Lawson, J. D., 1982. Valuations on continuous lattices. In: Hoffmann, R.-E. (Ed.), *Mathematische Arbeitspapiere*. Vol. 27. Universität Bremen, pp. 204–225.
- [33] Lawson, J. D., 1987. The versatile continuous order. In: Main, M. G., Melton, A., Mislove, M. W., Schmidt, D. A. (Eds.), *Proc. 3rd Workshop on Mathematical Foundations of Programming Language Semantics (MFPS'87)*. Springer Verlag LNCS 298, pp. 134–160.
- [34] Mac Lane, S., 1971. *Categories for the Working Mathematician*. Vol. 5 of *Graduate Texts in Mathematics*. Springer-Verlag.
- [35] Michael, E., 1951. Topologies on spaces of subsets. *Transactions of the American Mathematical Society* 71 (1), 152–182.
- [36] Saheb-Djahromi, N., 1980. Cpo's of measures for nondeterminism. *Theoretical Computer Science* 12, 19–37.
- [37] Schalk, A., 1993. Algebras for generalized power constructions. Ph.D. thesis, Technische Hochschule Darmstadt, available from <http://www.cs.man.ac.uk/~schalk/publ/diss.ps.gz>.
- [38] Smyth, M. B., 1983. Powerdomains and predicate transformers: A topological view. In: Diaz, J. (Ed.), *Automata, Languages and Programming*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 662–675.
- [39] Tix, R., 1995. Stetige Bewertungen auf topologischen Räumen. Diplomarbeit, Technische Hochschule Darmstadt.
- [40] Varacca, D., Winskel, G., 2006. Distributing probability over nondeterminism. *Mathematical Structures in Computer Science* 16 (1), 87–113.
- [41] Vietoris, L., 1922. Bereiche zweiter Ordnung. *Monatshefte für Mathematik* 32, 258–280.

## Appendix A. Uniqueness of Radon measures

There is a gap in the proof of [29, Theorem 7.3]. It is claimed that, for a weakly Hausdorff coherent sober space  $X$ , every tight, locally finite valuation  $\nu$  on  $X$  extends to a unique Radon measure on a  $\sigma$ -algebra  $\Sigma$  containing the patch topology of  $X$ . While existence—the difficult part—holds, uniqueness is dealt rather briefly in the final sentence of the proof (“Clearly a Radon measure is uniquely determined by its values on  $\mathcal{C}$ ”; the set  $\mathcal{C}$  is what we call  $\mathcal{C}X$ ). But that only says that the extension is determined uniquely from its values on sets in  $\mathcal{C}X$ , and no reason is given as to why the latter values would be determined uniquely from  $\nu$ .

In trying to find a fix, we needed to change the definition of Radon measures slightly. We define Radon measures on coherent sober spaces as measures on a  $\sigma$ -algebra  $\Sigma$  containing the patch topology, whose restrictions to  $\mathcal{O}(X^{\text{patch}})$  are locally finite, and which are inner regular. Keimel and Lawson’s definition [29, Definition 7.2] is a measure  $\mu$  on  $\Sigma$  that is inner regular and satisfying the following property, instead of local finiteness:

$$\text{(Fin)} \quad \mu(K) < \infty \text{ for every } K \in \mathcal{C}X.$$

Every measure  $\mu$  whose restriction to  $\mathcal{O}(X^{\text{patch}})$  is locally finite satisfies (Fin): for each  $x \in K$ , find a patch-open neighborhood  $O_x$  of  $x$  such that  $\mu(O_x) < \infty$ ; finitely many cover  $K$ , since  $K$  is patch-compact (Lemma 12.1), and their union, which contains  $K$ , has finite  $\mu$ -measure. I do not know whether the converse implication holds, and even less so for inner regular measures.

Now, the extension of  $\nu$  built by Keimel and Lawson—call it  $\mu$ —does not just satisfy property (Fin), but satisfies the stronger property of being locally finite on  $\mathcal{O}(X^{\text{patch}})$ . Indeed, for every  $x \in X$ , since  $\nu$  is locally finite, there is an open neighborhood  $U$  of  $x$  in  $X$  such that  $\nu(U) < \infty$ ; then  $U$  is also patch-open, and  $\nu(U) = \mu(U)$ .

We now claim that there can be only one Radon measure  $\mu$  (in our sense) on the patch space that extends  $\nu$ , provided that  $X$  is weakly Hausdorff, coherent, and well-filtered. As Keimel and Lawson argue, the value of  $\mu$  on Borel sets is entirely determined by its values on elements  $K$  of  $\mathcal{C}X$ , by inner regularity. In proving that local finiteness implies (Fin), we have actually shown a bit more: there is a patch-open neighborhood  $O$  of  $K$  such that  $\mu(O) < \infty$ . Then  $\mu(K) = \mu(O) - \mu(O \setminus K)$ ; the difference makes sense because  $\mu(O \setminus K) \leq \mu(O) < \infty$ . We use inner regularity a second time, and we express  $\mu(O \setminus K)$  as  $\sup_{K'}^{\uparrow} \mu(K')$ , where  $K'$  ranges over the elements of

$\mathcal{C}X$  included in  $O \setminus K$ . Then  $\mu(K) = \inf_{K'}^\downarrow (\mu(O) - \mu(K')) = \inf_{K'}^\downarrow \mu(O \setminus K')$ . Each set  $K'$  is patch-closed, so  $O \setminus K'$  is patch-open. Hence  $\mu(K)$  is entirely determined by the values of  $\mu$  on the patch-open sets.

We argue that the restriction  $\underline{\mu}$  of  $\mu$  to the patch-open subsets of  $X$  is a continuous valuation on  $X^{\text{patch}}$ . It is clearly strict and modular, and Scott-continuity is proved as follows. Let  $(O_i)_{i \in I}$  be a directed family of patch-open subsets of  $X$ , and let  $O$  be its union. Then  $\underline{\mu}(O) = \mu(O) = \sup_{K \subseteq O}^\uparrow \mu(K)$ , where  $K$  (implicitly) ranges over  $\mathcal{C}X$ . Since  $K$  is patch-compact (Lemma 12.1),  $K \subseteq O = \bigcup_{i \in I}^\uparrow O_i$  is equivalent to the existence of an  $i \in I$  such that  $K \subseteq O_i$ . Therefore  $\underline{\mu}(O) = \sup_{i \in I, K \subseteq O_i}^\uparrow \mu(K) = \sup_{i \in I}^\uparrow \mu(O_i) = \sup_{i \in I}^\uparrow \underline{\mu}(O_i)$ . In the previous paragraph, we have argued that  $\mu$  is determined uniquely as a function of  $\underline{\mu}$ , and we will show that  $\underline{\mu}$  is determined uniquely from  $\nu$ .

The family  $\mathcal{F}_\nu$  of open subsets  $U$  (in the original topology on  $X$ ) such that  $\nu(U) < \infty$  is directed, because it contains the empty set and is closed under binary unions, using modularity. The fact that  $\nu$  is locally finite means that  $\bigcup \mathcal{F}_\nu = X$ .

Every patch-open set is a union of sets of the form  $U \setminus Q$  with  $U$  open and  $Q$  compact saturated in  $X$  (with the original topology). Since  $\nu$  is locally finite, we can write  $U \setminus Q$  as the union of the sets  $U \cap U_0 \setminus Q$ ,  $U_0 \in \mathcal{F}_\nu$ . We note that  $U \cap U_0$  is itself in  $\mathcal{F}_\nu$ . Hence every patch-open set is also a union of sets from the collection  $\mathcal{B}$  of sets of the form  $U \setminus Q$  with  $U \in \mathcal{F}_\nu$  (not just  $U \in \mathcal{O}X$ ) and  $Q$  compact saturated in  $X$ . Namely,  $\mathcal{B}$  forms a subbase of the patch topology. Additionally,  $\mathcal{B}$  is closed under finite intersections, and in particular forms a base.

With these observations, we claim that  $\underline{\mu}$  is uniquely determined by its values on elements of  $\mathcal{B}$ . For every patch-open set  $O$ , written as a union  $\bigcup_{i \in I} B_i$  of elements of  $\mathcal{B}$ , hence as a directed union  $\bigcup_{J \in \mathbb{P}_{\text{fin}}(I)}^\uparrow \bigcup_{j \in J} B_j$ ,  $\underline{\mu}(O)$  is equal to  $\sup_{J \in \mathbb{P}_{\text{fin}}(I)}^\uparrow \underline{\mu}(\bigcup_{j \in J} B_j)$ , hence is uniquely determined from the values that  $\underline{\mu}$  takes on finite unions  $\bigcup_{j \in J} B_j$  of elements of  $\mathcal{B}$ . If  $\underline{\mu}(B_j) = \infty$  for some  $j \in J$ , then we must have  $\underline{\mu}(\bigcup_{j \in J} B_j) = \infty$ , since  $\underline{\mu}$  is monotonic. Otherwise, we can rewrite  $\underline{\mu}(\bigcup_{j \in J} B_j)$  as  $\sum_{K \subseteq J, K \neq \emptyset} (-1)^{|K|+1} \underline{\mu}(\bigcap_{j \in K} B_j)$  (this is the inclusion-exclusion formula, a consequence of modularity), and we recall that  $\bigcap_{j \in K} B_j \in \mathcal{B}$ . Hence in both cases, the value of  $\underline{\mu}(\bigcup_{j \in J} B_j)$  is uniquely determined by the values that  $\underline{\mu}$  takes on elements of  $\mathcal{B}$ .

It remains to see that the value  $\underline{\mu}(U \setminus Q)$  is entirely determined by  $\nu$  for

every set  $U \setminus Q$  in  $\mathcal{B}$ . We will show that  $\underline{\mu}(U \setminus Q) = \sup_{V \supseteq Q}^\uparrow (\nu(U) - \nu(U \cap V))$ , where  $V$  ranges over  $\mathcal{O}X$ . We note that the difference  $\nu(U) - \nu(U \cap V)$  makes sense because  $\nu(U \cap V) \leq \nu(U) < \infty$ .

To this end, it is enough to show that for every  $r \in \mathbb{R}_+$  such that  $r < \underline{\mu}(U \setminus Q)$ , there is an open neighborhood  $V$  of  $Q$  such that  $r \leq \nu(U) - \nu(U \cap V)$ . This will show that  $\underline{\mu}(U \setminus Q) \leq \sup_{V \supseteq Q}^\uparrow (\nu(U) - \nu(U \cap V))$ , and the reverse inequality is clear.

Since  $\mu$  is inner regular and  $\underline{\mu}$  is a restriction of  $\mu$  (hence, in particular,  $r < \mu(U \setminus Q)$ ) there is an element  $K \in \mathcal{C}X$  such that  $K \subseteq U \setminus Q$  and  $r \leq \mu(K)$ . Now  $K$  is an intersection  $\bigcap_{i \in I} \bigcup_{j=1}^{m_i} L_{ij}$  where  $I$  is non-empty and each  $L_{ij}$  is a lens, and we can therefore rewrite  $K$  as a filtered intersection  $\bigcap_{J \in \mathbb{P}_{\text{fin}}^*(I)}^\downarrow K_J$ , where  $\mathbb{P}_{\text{fin}}^*(I)$  denotes the collection of non-empty finite subsets of  $I$ , and  $K_J \stackrel{\text{def}}{=} \bigcap_{i \in J} \bigcup_{j=1}^{m_i} L_{ij}$ . By distributing intersections of unions, and realizing that any finite non-empty intersection of lenses is a lens (or the empty set), we see that  $K_J$  is a finite union of lenses. From  $K \subseteq U \setminus Q$ , we deduce that  $\bigcap_{J \in \mathbb{P}_{\text{fin}}^*(I)}^\downarrow K_J \setminus (U \setminus Q)$  is empty. But  $K_J \setminus (U \setminus Q) = (K_J \setminus U) \cup (K_J \cap Q)$  is also a finite union of lenses, because  $L \setminus U$  and  $L \cap Q$  are lenses (or empty) for every lens  $L$ . (The second claim relies on the coherence of  $X$ .) By Lemma 12.1, each set  $K_J \setminus (U \setminus Q)$  is then patch-compact, and is patch-closed. Now any filtered family of compact closed sets whose intersection is empty contains the empty set; this is an easy exercise using the definition of compactness. Therefore  $K_J \setminus (U \setminus Q)$  is empty for some  $J \in \mathbb{P}_{\text{fin}}^*(I)$ . In other words,  $K_J \subseteq U \setminus Q$ . We have  $K \subseteq K_J$ , so  $r \leq \mu(K_J)$ . Therefore, replacing  $K$  by  $K_J$  if necessary, we may assume that  $K$  itself is a finite union of lenses, say  $\bigcup_{i=1}^n (Q_i \cap C_i)$ , where each  $Q_i$  is compact saturated and  $C_i$  is closed in  $X$ , in such a way that  $r \leq \mu(K)$  and  $K \subseteq U \setminus Q$ .

From  $K \subseteq U \setminus Q$ , we obtain that  $Q_i \cap C_i \subseteq U \setminus Q$  for every  $i \in \{1, \dots, n\}$ , namely that  $Q_i \subseteq U \cup U_i$ , where  $U_i$  is the complement of  $C_i$ , and  $Q \cap Q_i \subseteq U$ . Since  $X$  is weakly Hausdorff,  $Q$  has an open neighborhood  $V_i$  and  $Q_i$  has an open neighborhood  $W_i$  such that  $V_i \cap W_i \subseteq U_i$ , for every  $i$ . Then  $Q_i \cap C_i \subseteq W_i \cap C_i \subseteq X \setminus V_i$ , for every  $i \in \{1, \dots, n\}$ . We let  $V \stackrel{\text{def}}{=} \bigcap_{i=1}^n V_i$ . Then  $K \subseteq X \setminus V$ . We also have  $K \subseteq U$ , so  $K \subseteq U \setminus V$ . From  $r \leq \mu(K)$ , we deduce that  $r \leq \mu(U \setminus V)$ . Now  $\mu(U \setminus V) = \mu(U) - \mu(U \cap V)$ , an equality that makes sense because all the sets involved have finite  $\nu$ -measure, equivalently finitely  $\mu$ -measure, since  $\mu$  extends  $\nu$ . Then  $\mu(U \setminus V) = \nu(U) - \nu(U \cap V)$ , and this completes the proof.